

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 November 2009 (12.11.2009)

PCT

(10) International Publication Number
WO 2009/136352 A1

(51) International Patent Classification:
A61K 47/48 (2006.01) **C07K 14/71** (2006.01)

CovX Technologies Ireland Limited 9381, Judicial Drive-Suite 200, San Diego, California 92121 (US).

(21) International Application Number:
PCT/IB2009/051820

(74) Agent: **BENSON, Gregg, C.**; Pfizer Inc., 150 East 42nd Street, New York, NY 10017 (US).

(22) International Filing Date:
4 May 2009 (04.05.2009)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/050,574 5 May 2008 (05.05.2008) US
61/050,569 5 May 2008 (05.05.2008) US
61/050,544 5 May 2008 (05.05.2008) US

(71) Applicant (for all designated States except US): **CovX Technologies Ireland, Ltd.** [IE/IE]; Pottery Road Dub Laoghaire County Dublin, Dublin (IE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DOPPALAPUDI, Venkata Ramana** [US/US]; CovX Technologies Ireland Limited 9381, Judicial Drive-Suite 200, San Diego, California 92121 (US). **LAI, Jing-Yu** [US/US]; CovX Technologies Ireland Limited 9381, Judicial Drive - Suite 200, San Diego, California 92121 (US). **LIU, Bin** [US/US]; CovX Technologies Ireland Limited 9381, Judicial Drive - Suite 200, San Diego, California 92121 (US). **LIU, Dingguo** [CN/US]; CovX Technologies Ireland Limited 9381, Judicial Drive - Suite 200, San Diego, California 92121 (US). **DESHARNAIS, Joel** [CA/US]; CovX Technologies Ireland Limited 9381, Judicial Drive - Suite 200, San Diego, California 92121 (US). **BHAT, Abhijit Suresh** [IN/US]; CovX Technologies Ireland Limited 9381, Judicial Drive - Suite 200, San Diego, California 92121 (US). **FU, Yanwen** [CN/US]; CovX Technologies Ireland Limited 9381, Judicial Drive - Suite 200, San Diego, California 92121 (US). **OATES, Bryan Douglas** [US/US]; CovX Technologies Ireland Limited 9381, Judicial Drive - Suite 200, San Diego, California 92121 (US). **GANG, Chen** [US/US]; CovX Technologies Ireland Limited 9381, Judicial Drive - Suite 200, San Diego, California 92121 (US). **BRADSHAW, Curt William** [US/US];

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to the identity of the inventor (Rule 4.17(i))
- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: ANTI-ANGIOGENIC COMPOUNDS

(57) Abstract: The present invention provides VEGF binding peptides. In addition, the invention provides VEGF peptides conjugated to antibodies alone and in conjunction with other anti-angiogenic molecules. Various uses of the peptides and compounds are provided, including methods to treat disorders associated with abnormal angiogenesis.



WO 2009/136352 A1

ANTI-ANGIOGENIC COMPOUNDS

Field of the disclosure

The present invention relates to novel compounds that possess anti-angiogenic activity and methods of making and using these compounds. In particular, the present invention relates to peptides that bind to the vascular endothelial cell growth factor (VEGF) and macromolecules incorporating these peptides, and methods and uses associated thereof.

Background

Angiogenesis is the fundamental process by which new blood vessels are formed and is essential to a variety of normal body activities such as reproduction, development and wound repair. Although angiogenesis is a highly regulated process under normal conditions, many diseases (characterized as “angiogenic diseases”) are caused or exacerbated by unregulated angiogenesis. For example, ocular neovascularization has been implicated as the most common cause of blindness. In certain existing conditions such as arthritis, newly formed capillary blood vessels invade the joints and destroy cartilage. In diabetes, new capillaries formed in the retina invade the vitreous, bleed, and cause blindness. Growth and metastasis of solid tumors are also angiogenesis-dependent (J. Folkman, Cancer Res., 46:467-473 (1986), J. Folkman, J. Natl. Cancer Inst., 82:4-6 (1989)). It has been shown, for example, that tumors which enlarge to greater than 2 mm obtain their own blood supply by inducing the growth of new capillary blood vessels. Once these new blood vessels become embedded in the tumor, they provide a means for tumor cells to enter the circulation and metastasize to distant sites such as the liver, lungs, and bones (N. Weidner, et. al., N. Engl. J. Med., 324:1-8 (1991)).

Vascular endothelial growth factor (VEGF) has been identified as an extremely powerful angiogenic factor and is required for the growth and metastasis of many human tumours. Much research has focused on attempting to inhibit the VEGF pathway so as to limit or prevent angiogenesis or metastasis. Fairbrother *et al.*: ‘*Novel peptides selected to bind vascular endothelial growth factor target the receptor-binding site*’ (Biochemistry, **1998**, 37, 17754-17764) discloses peptides with varying abilities to bind to VEGF. Various peptides have been identified that bind to the angiogenesis related factor angiopoietin-2 (“Ang-2”) (Oliner, J. *et al.*, Cancer Cell, 204(6), 507-516 (2004)). The Ang-2 binding peptides have been shown to possess anti-angiogenic activity.

It would be desirable to provide compounds that show improved characteristics over known compounds, such as for example, improved binding to VEGF. It would be further desirable to provide compounds showing binding to both VEGF and Ang2.

The reference to any art in this specification is not, and should not be taken as, an acknowledgement of any form or suggestion that the referenced art forms part of the common general knowledge.

Summary of the Invention

[VEGF-Peptides] of the invention

The invention provides peptides, compounds, and pharmaceutical compositions capable of binding to human VEGF (as exemplified by SEQ ID NO:3). Accordingly, in some aspects, the present invention provides for a peptide including a sequence substantially homologous to the sequence:

V¹-E²-P³-N⁴-C⁵-D⁶-I⁷-H⁸-V⁹-M¹⁰-W¹¹-V¹²-W¹³-X¹⁴-C¹⁵-F¹⁶-E¹⁷-R¹⁸-X¹⁹ (SEQ ID NO:122), wherein X¹⁴ is E or V, and X¹⁹ is a natural or unnatural hydrophobic amino acid, or D-isomers thereof.

It has surprisingly been found that this class of peptides, and derivatives thereof (as will be detailed further below) demonstrate improved binding to VEGF over peptides of related structure. In particular mutating position 12 to a neutral hydrophobic residue, especially valine, provides improved binding characteristics over peptides with a negatively charged groups at this position, such as glutamate. In some embodiments, the present invention provides for a peptide including a sequence substantially homologous to SEQ ID NO:122, provided that V¹² is not substituted with E. In some embodiments, X¹⁹ is L. X¹⁹ may be D-Leu. X¹⁹ may be a conservative substitution of L, such as I, or other hydrophobic residue, such as A or V, or D-isomers of any of the preceding.

Further derivatives of the present invention provide for the addition of at least 1-4 residues at the C-terminus. Some embodiments provide for the addition of 2-4 residues at the C-terminus. Some embodiments provide for the addition of 2 residues at the C-terminus. Some embodiments provide for the addition of 3 residues at the C-terminus. Some embodiments provide for the addition of 4 residues at the C-terminus.

Accordingly, the invention also provides for a peptide including a sequence substantially homologous to the sequence:

V¹-E²-P³-N⁴-C⁵-D⁶-I⁷-H⁸-V⁹-M¹⁰-W¹¹-V¹²-W¹³-X¹⁴-C¹⁵-F¹⁶-E¹⁷-R¹⁸-X¹⁹-X²⁰ (SEQ ID NO:123)

wherein X²⁰ is any neutral, hydrophobic or aromatic amino acid. X²⁰ may be any aromatic amino acid, such as Y, F, W, or D-isomers thereof. X²⁰ may be any neutral hydrophobic amino acid, such as M, I, L, Nle, A, or D-isomers thereof.

In some aspects, the invention provides for a peptide including a sequence substantially homologous to the sequence:

V¹-E²-P³-N⁴-C⁵-D⁶-I⁷-H⁸-V⁹-M¹⁰-W¹¹-V¹²-W¹³-X¹⁴-C¹⁵-F¹⁶-E¹⁷-R¹⁸-X¹⁹-X²⁰-X²¹ (SEQ ID NO:124)

wherein X²¹ is any amino acid. In some aspects, X²¹ may be any neutral or positively charged amino acid, such as G, A, I, L, K, R, or K(ac), or D-isomers thereof. For example, X²¹ may be D-Ala.

In some aspects, the invention provides for a peptide including a sequence substantially homologous to the sequence:

V¹-E²-P³-N⁴-C⁵-D⁶-I⁷-H⁸-V⁹-M¹⁰-W¹¹-V¹²-W¹³-X¹⁴-C¹⁵-F¹⁶-E¹⁷-R¹⁸-X¹⁹-X²⁰-X²¹-X²² (SEQ ID NO:125)

wherein X²² is any aliphatic, polar, or negatively charged amino acid, such as V, L, P, E, G, I, S, T,

5 W, F, E, or D-isomers thereof.

In some aspects, the invention provides for a peptide including a sequence substantially homologous to the sequence:

V¹-E²-P³-N⁴-C⁵-D⁶-I⁷-H⁸-V⁹-M¹⁰-W¹¹-V¹²-W¹³-X¹⁴-C¹⁵-F¹⁶-E¹⁷-R¹⁸-X¹⁹-X²⁰-X²¹-X²²-X²³ (SEQ ID

10 NO:126), wherein X²³ is any amino acid. X²³ may be selected from the group consisting of G, A, I, L, Q, E, F, T, W, S, and Y, and D-isomers thereof. X²³ may be selected from the group consisting of G, A, I, L, Q, E, F, T, S, and Y, and D-isomers thereof. In some aspects, X²³ does not comprise a side chain with a double ring. In some aspects, X²³ is not W. In some aspects, X²³ may be E, T, S, L, or F, or D-isomers thereof.

15

In some embodiments, it has been found that it can be preferred for at least one of the 5 residues at the carboxyl terminus to be a D-isomer. In some aspects, at least two of the five C-terminus located residues are D-isomers. In some aspects, no more than three of the of the five C-terminus located residues are D-isomers. In some aspects, between one and three of the five C-terminus located residues are D-isomers. In some aspects, at least one of the four residues at the carboxyl terminus is a D-isomer. In some aspects, at least two of the four C-terminus located residues are D-isomers. In some aspects, no more than three of the four C-terminus located residues are D-isomers. In some aspects, between one and three of the four C-terminus located residues are D-isomers. For each of these, it is preferred that there are at least two and preferably at least three additional residues after X¹⁹. Providing an optimized number of D-isomers at the C-terminus has been found to provide the advantage of increased resistance to enzymatic degradation at the C-terminus, especially enzymatic degradation between residues E¹⁷ and R¹⁸, and enzymatic degradation between residues R¹⁸ and X¹⁹.

25

In some aspects of the invention, X¹⁹ may be L, X²⁰ is present and may be Y, X²¹ is present and may be an aliphatic hydrophobic amino acid (and may be selected from A, L, I, V, G), and X²² is present, and is a hydrophobic amino acid (and may be selected from L, P, V). In certain embodiments, two of X¹⁹, X²¹ and X²² are D-isomers. In some embodiments, X²³ may or may not be present.

30

In certain aspects, specific peptides, and compounds of the invention may include including a peptide sequence substantially homologous to one or more of SEQ ID NOs:36-106

35

In another aspect, the present invention provides for a peptide including a sequence substantially homologous to the sequence:

V¹-E²-P³-N⁴-C⁵-D⁶-I⁷-H⁸-V⁹-M¹⁰-W¹¹-E¹²-W¹³-E¹⁴-C¹⁵-F¹⁶-E¹⁷-R¹⁸-X¹⁹-X²⁰-X²¹-X²²-X²³ (SEQ ID

NO:127), wherein X¹⁹ is a natural or unnatural hydrophobic amino acid, or D-isomers thereof, and

X²⁰ is an aromatic amino acid, neutral amino acid, hydrophobic amino acid or polar amino acid, or D-isomers thereof, and X²¹ is a hydrophobic amino acid or positively charged amino acid, or D-isomers thereof, and X²² is absent, or is a hydrophobic amino acid, aromatic amino acid, negatively charged amino acid, or D-isomers thereof, and X²³ is absent, or is an aromatic amino acid, neutral amino acid, hydrophobic amino acid or D-isomers thereof. X¹⁹ may be L. In alternative embodiments X¹⁹ is

D-Leu. X¹⁹ may be a conservative substitution of L, such as I, or other hydrophobic residue, such as A or V, or D-isomers of any of the preceding. X²⁰ may be selected from the group consisting of A, V, I, L, Y, W, F, M, S, and T or D-isomer thereof. X²⁰ may be selected from the group consisting of A, Y, F, M, and S or D-isomer thereof. X²¹ may be selected from the group consisting of K, R, H, Ornithine, Dap, Dab, G, A, V, I, and L or D-isomers thereof. X²¹ may be selected from the group consisting of K, R, and G or D-isomer thereof. Where X²⁰ is hydrophobic or aromatic, X²¹ may be a positively charged amino acid. Where X²⁰ is one of A, F or Y, X²¹ may be one of K or R or D-isomer thereof. Where X²¹ is hydrophobic, X²⁰ may be polar. X²² if present may be selected from the group consisting of E, D, G, A, L, I, V, M, W, Y, and F or D-isomers thereof. X²² if present may be selected from the group consisting of E, G and W or D-isomers thereof. Where X²¹ is a positively charged residue, X²²

if present may be a negatively charged residue. Where X²¹ is K, X²² may be E. X²³ if present may be selected from the group consisting of W, Y, F, G, A, I, L, V, and M, or D-isomers thereof. X²³ if present may be selected from the group consisting of W, F, G, A, L, and M, or D-isomers thereof.

In certain aspects, specific peptides, compounds and pharmaceutical compositions of the invention

may include including a peptide sequence substantially homologous to one or more of the following:

SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117, SEQ ID NO:118, SEQ ID NO:119, SEQ ID NO:120, SEQ ID NO:121.

Peptides of the invention can be used in a variety of applications, including diagnosis, screening and therapeutics. Peptides and compounds of the invention may be used directly in therapeutic applications, or may find use conjugated covalently or non-covalently to larger molecules which provide additional therapeutic value, such as increased half-life (PK). Examples of molecules which may provide increased half-life include proteins, polypeptides, antibodies, antibody fragments, in particular, the Fc domain of antibodies

The invention provides for a compound of the formula: R¹-[VEGF-Peptide]-R²

wherein [VEGF-Peptide] is a peptide including a sequence substantially homologous to the sequence:

$X^1-X^2-P^3-N^4-C^5-X^6-X^7-X^8-V^9-X^{10}-X^{11}-X^{12}-W^{13}-X^{14}-C^{15}-F^{16}-E^{17}-R^{18}-X^{19}-X^{20}-X^{21}-X^{22}-X^{23}$ (SEQ ID

NO:131) wherein R^1 is absent, CH_3 , $C(O)CH_3$, $C(O)CH_2CH_3$, $C(O)CH_2CH_2CH_3$,

$C(O)CH(CH_3)CH_3$, $C(O)CH_2CH_2CH_2CH_3$, $C(O)CH(CH_3)CH_2CH_3$, $C(O)C_6H_5$,

5 $C(O)CH_2CH_2(CH_2CH_2O)_{1.5}Me$, amido-2-PEG, N-acyl and N-alkyl an amino protecting group, a lipid fatty acid group or a carbohydrate; and R^2 is absent, OH, NH_2 , $NH(CH_3)$, $NHCH_2CH_3$,

$NHCH_2CH_2CH_3$, $NHCH(CH_3)CH_3$, $NHCH_2CH_2CH_2CH_3$, $NHCH(CH_3)CH_2CH_3$, NHC_6H_5 ,

$NHCH_2CH_2OCH_3$, $NHOCH_3$, $NHOCH_2CH_3$, a carboxy protecting group, a lipid fatty acid group or a carbohydrate, and X^1 is a hydrophobic amino acid residue, X^2 is a negatively charged residue, X^6 is a

10 negatively charged residue, X^7 is a hydrophobic amino acid residue, X^8 is a residue comprising a ring structure, X^{10} may be M or any hydrophobic amino acid, X^{11} is an aromatic amino acid, X^{12} is selected

from the group consisting of V, E and Kac, X^{14} is E or V, X^{19} may be any hydrophobic amino acid

residue, or D-isomer thereof, X^{20} may be absent, or may be any neutral, hydrophobic or aromatic

amino acid or D-isomer thereof, X^{21} may be absent, or may be any positively charged residue or any

15 aliphatic non-polar residue or D-isomer thereof, X^{22} may be absent, or may be selected from the group consisting of G, V, L, I, P, S, T, W, F, E, Kac, or D-isomers thereof, X^{23} may be absent, or is selected from the group consisting of G, A, I, L, Q, E, F, T, W, S, Y, and Kac and D-isomers thereof.

X^1 may be V. X^2 may be E. X^6 may be D. X^7 may be I. X^8 may be H. X^{11} may be W. [VEGF-Peptide]

20 may be a peptide including a sequence substantially homologous to the sequence:

$V^1-E^2-P^3-N^4-C^5-D^6-I^7-H^8-V^9-X^{10}-W^{11}-X^{12}-W^{13}-X^{14}-C^{15}-F^{16}-E^{17}-R^{18}-X^{19}-X^{20}-X^{21}-X^{22}-X^{23}$ (SEQ ID

NO:132), where X^{10} may be M or any hydrophobic amino acid, X^{12} is selected from the group

consisting of V, E and Kac, X^{14} is E or V, X^{19} may be any hydrophobic amino acid residue, or D-

isomer thereof, X^{20} may be absent, or may be any neutral, hydrophobic or aromatic amino acid or D-

25 isomer thereof, X^{21} may be absent, or may be any positively charged residue or any aliphatic non-

polar residue or D-isomer thereof, X^{22} may be absent, or may be selected from the group consisting of

G, V, L, I, P, S, T, W, F, E, Kac, or D-isomers thereof, X^{23} may be absent, or is selected from the

group consisting of G, A, I, L, Q, E, F, T, W, S, Y, and Kac and D-isomers thereof.

30 [VEGF-Peptide] includes a sequence substantially homologous to one or more of the following:

V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L (SEQ ID NO:34)

V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-G-W (SEQ ID NO:35)

V-E-P-N-C-D-I-H-V-M-W-V-W-V-C-F-E-R-L-F-R-E-A (SEQ ID NO:36)

V-E-P-N-C-D-I-H-V-M-W-V-W-V-C-F-E-R-L-F-K-E-A (SEQ ID NO:37)

35 V-E-P-N-C-D-I-H-V-M-W-V-W-V-C-F-E-R-L-M-K (SEQ ID NO:38)

V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-R-E-L (SEQ ID NO:39)

V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-I-F (SEQ ID NO:40)

- V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y (SEQ ID NO:41)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-W-G (SEQ ID NO:42)
V-E-P-N-C-D-I-H-V-M-W-V-W-V-C-F-E-R-L-Y-G-G-G (SEQ ID NO:43)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-L-Y (SEQ ID NO:44)
5 V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-S-G-G-G (SEQ ID NO:45)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-M-R-L-T (SEQ ID NO:46)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G (SEQ ID NO:47)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-V-K (SEQ ID NO:48)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-M-R (SEQ ID NO:49)
10 V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-I-L (SEQ ID NO:50)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-(Yome)-G-L-T (SEQ ID NO:51)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-G (SEQ ID NO:52)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-F-K-E-A (SEQ ID NO:53)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-(Nle)-K (SEQ ID NO:54)
15 V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-L-T (SEQ ID NO:55)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-G-F (SEQ ID NO:56)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-I-K (SEQ ID NO:57)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-K (SEQ ID NO:58)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-M-G-L-T (SEQ ID NO:59)
20 V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-(D-Leu)-(Kac) (SEQ ID NO:60)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-G-G (SEQ ID NO:61)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-M-K (SEQ ID NO:62)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-M-R-E-L (SEQ ID NO:63)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L- (SEQ ID NO:64)
25 V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-M-K-E-L (SEQ ID NO:65)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-P-W (SEQ ID NO:66)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-L-K (SEQ ID NO:67)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-L-(Kac) (SEQ ID NO:68)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-L-T (SEQ ID NO:69)
30 V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-E-F (SEQ ID NO:70)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-L-S (SEQ ID NO:71)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-L (SEQ ID NO:72)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-(D-Leu)-T (SEQ ID NO:73)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-V-Q (SEQ ID NO:74)
35 V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-L-E (SEQ ID NO:75)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-P-L (SEQ ID NO:76)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-P-F (SEQ ID NO:77)

- V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-(D-Ala)-(D-Leu) (SEQ ID NO:78)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-R-(D-Leu)-(Kac) (SEQ ID NO:79)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-(D-Tyr)-G-(D-Pro)-L (SEQ ID NO:80)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-(D-Tyr)-G-(D-Pro)-(D-Leu) (SEQ ID NO:81)
5 V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-(D-Tyr)-G-(D-Leu)-(Kac) (SEQ ID NO:82)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-(D-Leu)-Y-(Aib)-L-T (SEQ ID NO:83)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-(D-Leu)-Y-(D-Ala)-V-(D-Gln)(SEQ ID NO:84)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-(D-Leu)-Y-(D-Ala)-L-(D-Thr)(SEQ ID NO:85)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-(Cha)-G-(DLeu)-T (SEQ ID NO:86)
10 V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-(Aib)-L-(D-Thr) (SEQ ID NO:87)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-(Sar)-(D-Leu)-T (SEQ ID NO:88)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-(D-Pro)-L(SEQ ID NO:89)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-(D-Ala)-V-(D-Gln) (SEQ ID NO:90)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-(Aib)-L-T (SEQ ID NO:91)
15 V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-(D-Tyr)-G-(D-Leu)-T (SEQ ID NO:92)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-(D-Tyr)-G-L-T (SEQ ID NO:93)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-(D-Tyr)-D-L-(D-Thr) (SEQ ID NO:94)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-(D-Leu)-(Aib) (SEQ ID NO:95)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-(D-Leu)-(D-Thr) (SEQ ID NO:96)
20 V-E-P-N-C-D-I-H-V-M-W-V-W-V-C-F-E-R-L (SEQ ID NO:97)
V-E-P-N-C-D-I-H-V-M-W-V-W-V-C-F-E-R-L-Y-G-P-F (SEQ ID NO:98)
V-E-P-N-C-D-I-H-V-M-W-V-W-V-C-F-E-R-L-Y-G-P-(SEQ ID NO:99)
V-E-P-N-C-D-I-H-V-M-W-V-W-V-C-F-E-R-L-Y-G-P-E (SEQ ID NO:100)
V-E-P-N-C-D-I-H-V-M-W-V-W-V-C-F-E-R-L-Y-G-P-Q (SEQ ID NO:101)
25 V-E-P-N-C-D-I-H-V-M-W-V-W-V-C-F-E-R-L-Y-G-(D-Leu)-T (SEQ ID NO:102)
V-E-P-N-C-D-I-H-V-M-W-V-W-V-C-F-E-R-L-Y-G-L ((SEQ ID NO:103)
V-E-P-N-C-D-I-H-V-M-W-V-W-V-C-F-E-R-L-(D-Leu)-Kac (SEQ ID NO:104)
V-E-P-N-C-D-I-H-V-M-W-V-W-V-C-F-E-R-L-Y-G-L-T (SEQ ID NO:105)
V-E-P-N-C-D-I-H-V-M-W-V-W-V-C-F-E-R-L-Y-(D-Ala)-(D-Leu) (SEQ ID NO:106)
30 V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-Y-A-K (SEQ ID NO:108)
V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-Y-F-K-E-W (SEQ ID NO:109)
V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-Y-A-R (SEQ ID NO:110)
V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-Y-F-K (SEQ ID NO:111)
V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-Y-A-K-E-F (SEQ ID NO:112)
35 V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-Y-S-G-W-G (SEQ ID NO:113)
V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-Y-S-G-W-F (SEQ ID NO:114)
V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-Y-A-K-E-A (SEQ ID NO:115)

- V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-Y-A-K-E-M (SEQ ID NO:116)
V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-Y-A-K-E-L (SEQ ID NO:117)
V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-Y-F-K-E-L (SEQ ID NO:118)
V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-Y-F-K-E-A (SEQ ID NO:119)
5 V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-Y-Y-G-G-G (SEQ ID NO:120)
V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-Y-M-K (SEQ ID NO:121)
V-E-P-N-C-D-I-H-V-(Nle)-W-V-W-E-C-F-E-R-L-Y-(Aib)-L-T (SEQ ID NO:128)
V-E-P-N-C-D-I-H-V-L-W-V-W-E-C-F-E-R-L-Y-(Aib)-L-T (SEQ ID NO:129)
V-E-P-N-C-D-I-H-V-V-W-V-W-E-C-F-E-R-L-Y-(Aib)-L-T (SEQ ID NO:130)
10 V-E-P-N-C-D-I-H-V-M-W-(Kac)-W-E-C-F-E-R-L-Y-(D-Ala)-V-(D-Gln) (SEQ ID NO:136)
V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-R-L-(D-Leu)-(Kac) (SEQ ID NO:192)
V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-R-L-Y-G-(D-Leu)-T (SEQ ID NO:193)
V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-R-L-Y-G-P-L (SEQ ID NO:194)
V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-R-L-Y-(D-Ala)-(D-Leu) (SEQ ID NO:195)
15 V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-(D-Leu)-K (SEQ ID NO:196)
V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-G-(D-Leu)-K (SEQ ID NO:197)
V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-K-R-(D-Leu) (SEQ ID NO:198)
V-E-P-N-C-D-I-H-V-KW-E-W-E-C-F-(N-methyl glutamate)-R-L (SEQ ID NO:199)
V-E-P-N-C-D-I-H-V-KW-V-E-E-C-F-(D-Glu)-R-L (SEQ ID NO:200)
20 V-E-P-N-C-D-I-H-V-KW-E-W-E-C-F-E-R(N-methyl arginine) (SEQ ID NO:201)
V-E-P-N-C-D-I-H-V-KW-E-W-E-C-F-E-R-(D-Asn) (SEQ ID NO:202)
V-E-P-N-C-D-I-H-V-KW-E-W-E-C-F-E-R-(Aib) (SEQ ID NO:203)
V-E-P-N-C-D-I-H-V-KW-E-W-E-C-F-(Aib)-R-L (SEQ ID NO:204)
V-E-P-N-C-D-I-H-V-KW-E-W-E-C-(Naphthaline)-E-R-L (SEQ ID NO:205)
25 V-E-P-N-C-D-I-H-V-M-W-V-E-C-F-K-R-L-Y-G-L-T (SEQ ID NO:206)
V-E-P-N-C-D-I-H-V-M-W-V-E-C-F-E-R-L-Y-K-L-E (SEQ ID NO:207)
V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-(D-Leu) (SEQ ID NO: 208)
V-E-P-N-C-D-I-H-V-E-W-E-W-E-C-F-K-R-L (SEQ ID NO:209)
V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-(homoarginine)-L-Y-(D-Ala)-(D-Leu) (SEQ ID NO:210)
30 V-E-P-N-C-D-I-H-V-M-W-K-W-E-C-F-E-R-Y-G-(D-Leu)-E (SEQ ID NO:211)
V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-R-L-Y-G-(D-Leu)-E (SEQ ID NO:212)
V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-R-L-Y-(Aib)-L-E (SEQ ID NO:213)
V-E-P-N-C-D-I-H-V-M-W-K-W-E-C-F-E-R-L-Y-(Aib)-L-E (SEQ ID NO:214)
V-E-P-N-C-D-I-H-V-M-W-K-W-E-C-F-E-R-(D-Leu)-K (SEQ ID NO:215)
35 V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-(Kac)-L-Y-(D-Ala)-(D-Leu) (SEQ ID NO:216)
V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-(Kac)-(cyclo-Leu)-Y-(D-Ala)-(D-Leu) (SEQ ID NO:217)
V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-(Kac)-(Tle)-Y-(D-Ala)-(D-Leu) (SEQ ID NO:218)

In some embodiments, the [VEGF-Peptide] of the invention is selected from the group consisting of SEQ ID NOs: 60, 73, 76, 78, 192, 193, 194, or 195.

5 X¹⁰ may be M. [VEGF-Peptide] may be a peptide including a sequence substantially homologous to the sequence:

V¹-E²-P³-N⁴-C⁵-D⁶-I⁷-H⁸-V⁹-M¹⁰-W¹¹-X¹²-W¹³-X¹⁴-C¹⁵-F¹⁶-E¹⁷-R¹⁸-X¹⁹-X²⁰-X²¹-X²²-X²³ (SEQ ID NO:133) where X¹² is selected from the group consisting of V, E and Kac, X¹⁴ is E or V, X¹⁹ may be any hydrophobic amino acid residue, or D-isomer thereof, X²⁰ may be absent, or may be any neutral, hydrophobic or aromatic amino acid or D-isomer thereof, X²¹ may be absent, or may be any positively charged residue or any aliphatic non-polar residue or D-isomer thereof, X²² may be absent, or may be selected from the group consisting of G, V, L, I, P, S, T, W, F, E, Kac, or D-isomers thereof, X²³ may be absent, or is selected from the group consisting of G, A, I, L, Q, E, F, T, W, S, Y, and Kac and D-isomers thereof.

15

X¹² may be E. [VEGF-Peptide] may be a peptide including a sequence substantially homologous to the sequence:

V¹-E²-P³-N⁴-C⁵-D⁶-I⁷-H⁸-V⁹-M¹⁰-W¹¹-E¹²-W¹³-X¹⁴-C¹⁵-F¹⁶-E¹⁷-R¹⁸-X¹⁹-X²⁰-X²¹-X²²-X²³ (SEQ ID NO:134) where X¹⁴ is E or V, X¹⁹ may be any hydrophobic amino acid residue, or D-isomer thereof, X²⁰ may be absent, or may be any neutral, hydrophobic or aromatic amino acid or D-isomer thereof, X²¹ may be absent, or may be any positively charged residue or any aliphatic non-polar residue or D-isomer thereof, X²² may be absent, or may be selected from the group consisting of G, V, L, I, P, S, T, W, F, E, Kac, or D-isomers thereof, X²³ may be absent, or is selected from the group consisting of G, A, I, L, Q, E, F, T, W, S, Y, and Kac and D-isomers thereof.

25

[VEGF-Peptide] may include a sequence substantially homologous to one or more of the following: SEQ ID NO:34, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117, SEQ ID NO:118, SEQ ID NO:119, SEQ ID NO:120, SEQ ID NO:121.

30

X¹² may be V. [VEGF-Peptide] may be a peptide including a sequence substantially homologous to the sequence:

V¹-E²-P³-N⁴-C⁵-D⁶-I⁷-H⁸-V⁹-M¹⁰-W¹¹-V¹²-W¹³-X¹⁴-C¹⁵-F¹⁶-E¹⁷-R¹⁸-X¹⁹-X²⁰-X²¹-X²²-X²³ (SEQ ID NO:135) where X¹⁴ is E or V, X¹⁹ may be any hydrophobic amino acid residue, or D-isomer thereof, X²⁰ may be absent, or may be any neutral, hydrophobic or aromatic amino acid or D-isomer thereof, X²¹ may be absent, or may be any positively charged residue or any aliphatic non-polar residue or D-isomer thereof, X²² may be absent, or may be selected from the group consisting of G, V, L, I, P, S, T,

35

W, F, E, Kac, or D-isomers thereof, X^{23} may be absent, or is selected from the group consisting of G, A, I, L, Q, E, F, T, W, S, Y, and Kac and D-isomers thereof.

[VEGF-Peptide] may include a sequence substantially homologous to one or more of the following:

- 5 SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, ID NO:71, SEQ ID NO:72), SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:128, SEQ ID NO:129, SEQ ID NO:130

- In relation to the various embodiments of [VEGF-Peptide] described herein, in some aspects of the invention, R^1 may be $C(O)CH_3$. In some aspects of the invention, R^2 is NH_2 . R^1 and/ or R^2 may be absent. X^{19} may be selected from the group consisting of L, I, A, V, G or D-isomers thereof. X^{19} may be L. X^{19} may be D-Leu. X^{20} may be Y. X^{20} may be D-Tyr. X^{21} may be selected from the group consisting of D-Ala, G, Aib and Kac. X^{21} may be D-Ala. X^{21} may be G. X^{22} may be selected from the group consisting of L, V, P, D-Leu, D-Pro. X^{22} may be D-Leu. X^{22} may be D-Pro. X^{22} may be L. X^{23} may be absent.

- In some embodiments of the invention, [VEGF-peptide] is (SEQ ID NO:78). In some embodiments, R^1 -[VEGF-peptide]- R^2 is $\{C(O)CH_3\}$ -V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-R-L-Y-(D-Ala)-(D-Leu)- $\{NH_2\}$ (SEQ ID NO:188) and K^{11} is the linking residue.

30

Linking

- The presence of the linking residue (both VEGF-linking residue and Ang2-linking residue respectively) provides the compounds of the invention with great flexibility for linkages to scaffolds, macromolecules and other moieties. In particular, the compounds of the invention may be reliably, securely and efficiently covalently linked to scaffolds, such as antibodies, antibody fragments, PEG molecules, albumin and so on. Surprisingly, it has been found that locating the linking residue at certain key positions in the respective peptide leads to increased stability and/or binding of the

35

peptide. The linking residue can be selected so as to provide a side chain whose chemical characteristics permit specific, reliable, directional and efficient chemical covalent linkages at that location. In some aspects of the invention, the linking residue is covalently linked to the combining site of an antibody directly or via an intermediate linker. The linkage may be irreversible.

5

The compounds of the invention may be covalently linked to a linker moiety, L (or L', as later described) through the linking residue. Numerous linkers are possible; many suitable linkers are disclosed in US2006205670, the contents of which are incorporated herein by reference. In particular, aspects of US2006205670 pertaining to the general formulae describing linkers, specific linker structure, synthesis of linkers and combinations of different elements of X, Y and Z groups as specifically and generally described therein are herein included. The linker may be linear or branched, and optionally includes one or more carbocyclic or heterocyclic groups. Linker length may be viewed in terms of the number of linear atoms, with cyclic moieties such as aromatic rings and the like to be counted by taking the shortest route around the ring. In some embodiments, the linker has a linear stretch of between 5-15 atoms, in other embodiments 15-30 atoms, in still other embodiments 30-50 atoms, in still other embodiments 50-100 atoms, and in still other embodiments 100-200 atoms. Other linker considerations include the effect on physical or pharmacokinetic properties of the resulting compound, such as solubility, lipophilicity, hydrophilicity, hydrophobicity, stability (more or less stable as well as planned degradation), rigidity, flexibility, immunogenicity, modulation of antibody binding, the ability to be incorporated into a micelle or liposome, and the like.


15
20

In some aspects of the invention, the compound comprises a linker (L) or (L') covalently linked to the side chain of the linking residue. The linker may comprise the formula: -[Connector]-X-Y-Z; -[Connector]-X-Y-Z'; -X-Y-Z-; or -X-Y-Z'-; wherein: [Connector] is present where the linker is branched, and where present is covalently linked to the linking residue, and one or more additional Active Molecules, X is a biologically compatible connecting chain including any atom selected from the group consisting of C, H, N, O, P, S, F, CL, Br, and I, and may comprise a polymer or block copolymer, and is covalently linked to the linking residue where the linker is linear, Y is an optionally present recognition group comprising at least a ring structure; and Z is a reactive group that is capable of forming a covalent bond with an amino acid side chain in a combining site of an antibody, and Z' is an attachment moiety comprising a covalent link to an amino acid side chain in a combining site of an antibody.


25

30

When present, Y may have the optionally substituted structure:



or



35

wherein a, b, c, d, and e are independently carbon or nitrogen; f is carbon, nitrogen, oxygen, or sulfur; Y is attached to X and Z independently at any two ring positions of sufficient valence; and no more than four of a, b, c, d, e, or f are simultaneously nitrogen and preferably a, b, c, d, and e in the ring structure are each carbon. In some aspects, Y may be phenyl. Although not wishing to be bound by any theory, it is believed that the Y group can assist in positioning the reactive group into an antibody combining site so that the Z group can react with a reactive amino acid side chain.


The linker may be designed such that it contains a reactive group capable of covalently or non-covalently forming a bond with a macromolecule, such as an antibody, protein, or fragment thereof.

The reactive group is chosen for use with a reactive residue in a particular combining site. For example, a chemical moiety for modification by an aldolase antibody may be a ketone, diketone, beta lactam, active ester haloketone, lactone, anhydride, maleimide, alpha-haloacetamide, cyclohexyl diketone, epoxide, aldehyde, amidine, guanidine, imine, enamine, phosphate, phosphonate, epoxide, aziridine, thioepoxide, masked or protected diketone (ketal for example), lactam, haloketone, aldehyde, and the like. In embodiments of the present invention linking a peptide of the invention with a linker L or L', the moieties Z (or Z' when attached to the macromolecule) is the reactive group.

In some embodiments, Z includes one or more C=O groups arranged to form an azitidinone, diketone, an acyl beta-lactam, an active ester, a haloketone, a cyclohexyl diketone group, an aldehyde, a maleimide, an activated alkene, an activated alkyne or, in general, a molecule comprising a leaving group susceptible to nucleophilic or electrophilic displacement. Other groups may include a lactone, an anhydride, an alpha-haloacetamide, an imine, a hydrazide, or an epoxide. Exemplary linker electrophilic reactive groups that can covalently bond to a reactive nucleophilic group (e.g., a lysine or cysteine side chain) in a combining site of antibody include acyl beta-lactam, simple diketone, succinimide active ester, maleimide, haloacetamide with linker, haloketone, cyclohexyl diketone, aldehyde, amidine, guanidine, imine, enamine, phosphate, phosphonate, epoxide, aziridine, thioepoxide, a masked or protected diketone (a ketal for example), lactam, sulfonate, and the like, masked C=O groups such as imines, ketals, acetals, and any other known electrophilic group. In certain embodiments, the reactive group includes one or more C=O groups arranged to form an acyl beta-lactam, simple diketone, succinimide active ester, maleimide, haloacetamide with linker, haloketone, cyclohexyl diketone, or aldehyde. Z or Z' if present may be a substituted alkyl, substituted cycloalkyl, substituted aryl, substituted arylalkyl, substituted heterocyclyl, or substituted heterocyclylalkyl, wherein at least one substituent is a 1,3-diketone moiety, an acyl beta-lactam, an active ester, an alpha-haloketone, an aldehyde, a maleimide, a lactone, an anhydride, an alpha-haloacetamide, an amine, a hydrazide, or an epoxide. In some aspects, the Z group if present is covalently linked to the combining site of an antibody. In other aspects, the Z group is covalently


linked to a macromolecule scaffold that can provide increased half-life to the peptides of the invention.

In some aspects, Z if present has the structure:



wherein $q=0-5$. q may be 1 or 2. q may be 1. In other aspects, q may be 2.

Z' if present may have the structure:



wherein $q=0-5$ and Antibody-N- is a covalent bond to a side chain in a combining site of an antibody. q may be 1 or 2. q may be 1. In other aspects, q may be 2.

X may be a group comprising three components; $X_p-X_s-X_y$, wherein X_p is a group specifically adapted to be combinable with the side chain of the linking residue, X_s is a spacer region of the X group, and X_y is a group adapted to bind to the Y group. In some aspects, X_y is selected from an amide bond, an enamine bond, or a guaninium bond. X_y may be selected so as to provide a hydrogen molecule adjacent (within two atoms) to the Y group. While not wishing to be bound by theory, it is believed that the H atom can assist the Y group recognition of a hydrophobic pocket through H-bond interaction, particularly in respect of the hydrophobic pocket of the binding cleft of a catalytic antibody, such as h38C2 (e.g. Fig 2A). Thus the amide bond, for example, may be orientated such that the NH group is directly bonded to the Y group, providing the H of the NH group for hydrogen bonding. Alternatively, the C=O group of an amide may be bonded to the Y group, with the H of the NH group may be 2 atoms adjacent to the Y group, but still available for H-bonding. In some aspects, X_s is selected such that X_s does not provide any overly reactive groups. X_s may be selected so as to provide an overall length of the X groups of between 2-15 atoms. X_s may be selected so that the overall length of the X group is between 2 and 10 atoms. X_s may be selected so that the overall length of X group is 4-8 atoms. X_s may be selected so that the overall length of X group is 5 atoms. X_s may be selected so that the overall length of X group is 6 atoms. X_p ideally is selected so as to enable a specific directional covalent linking strategy to the linking residue. For example, where the linking residue comprises a nucleophilic group, X_p may be an electrophilic group and vice versa. For example, if the linking residue side chain comprises an amine group, such as K, H, Y, ornithine, Dap,

or Dab, Xp may be COOH, or other similarly reactive electrophile. If the linking residue is D or E, Xp may comprise a nucleophilic group, such as an amine group. Either of these strategies permits a covalent bond to be formed between the Xp group and the linking residue by amide bond formation strategies. Where the linking group is C, homologs of C, or other thiol-group containing residues, Xp may comprise a maleimide group, permitting a thiol-maleimide addition reaction strategy to covalently link the Xp group to the linking residue. In some aspects, Xp may also comprise a thiol group, allowing a disulphide bridge to be formed between the linking residue and Xp group.


The backbone length of X may be 3-15 atoms.

10 X may be: and in some aspects, X-Y may be:



wherein v and w are selected such that the backbone length of X is 6-12 atoms, or 3-9 atoms, or 4-7 atoms, or 6 atoms, or 7 atoms, or 8 atoms. V and W may each be between 0-8. V may be 1 or 2. W may be 1 or 2. V and W may each be 1. R^b may be hydrogen, substituted or unsubstituted C₁₋₁₀ alkyl, substituted or unsubstituted C₃₋₇ cycloalkyl-C₀₋₆ alkyl, or substituted or unsubstituted aryl-C₀₋₆ alkyl. R^b may be H. In some aspects, V and W are both 1, and R^b is H.

Certain embodiments in accordance of the invention have the structure:



where Peptide is a [VEGF-Peptide] of the invention, and v is 1, 2, 3, 4, or 5; w is 1, 2, 3, 4, or 5; q is 0, 1, 2, 3, 4, or 5; and R^b is hydrogen, substituted or unsubstituted C₁₋₁₀ alkyl, substituted or unsubstituted C₃₋₇ cycloalkyl-C₀₋₆ alkyl, or substituted or unsubstituted aryl-C₀₋₆ alkyl. In certain

embodiments, v is 1, 2, or 3; w is 1, 2, or 3; and q is 0, 1, 2, or 3. In some embodiments, v is 1 or 2; w is 1 or 2; and q is 1 or 2 and R^b is H.

In some aspects, the invention provides for a compound selected from the group consisting of Compounds 2001-2010, Compounds 2014-2016, Compound 2018, Compound 2020, Compound 2022-2024, Compounds 2027-2033, Compound 2036, Compound 2038, Compounds 2041- 2042, Compounds 2045- 2046, Compounds 2048-2050, and Compound 2052-2053. These compounds may be covalently linked to the combining site of a catalytic antibody, for example an aldolase antibody, for example h38C2. In some aspects, the invention provides for a compound of the invention selected from the group consisting of Compounds 2018, 2036, 2045, and 2050. These compounds may be covalently linked to the combining site of a catalytic antibody, for example an aldolase antibody, for example h38C2. In some aspects, the invention provides for Compound 2018.

The present invention also provides for nucleic acid sequences (including DNA sequences, RNA sequences, and DNA-RNA sequences) encoding peptides of the invention, and their precursors.

Bifunctional molecules

In some aspects, the invention provides a compound of the formula:



wherein [Active Molecule-1] is a [VEGF-Peptide] of the invention and [Connector] is a moiety covalently bonded to both [VEGF-Peptide] and [Active Molecule-2]. It will be appreciated that in some aspects of the invention, [Connector] may be covalently attached more than one other [Active Molecule] in addition to the [VEGF-Peptide].

R¹-[VEGF-peptide]-R² may be

{C(O)CH₃}- V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-R-L-Y-(D-Ala)-(D-Leu)- {NH₂} (SEQ ID NO:188) and K¹¹ may be the VEGF-linking residue covalently linked to the [Connector].

An Active Molecule may be any chemical, biochemical, or biological entity capable of being covalently bonded to [Connector] and interacting with biological systems. Examples include therapeutic agents, drugs, pro-drugs, targeting agents, toxins, proteins, peptides, nucleic acid molecules, and lipids. In some aspects of the invention, Active Molecules are Peptides, and may be anti-angiogenic peptides. In some aspects, Active Molecules are [VEGF-Peptides] of the invention and/or [Ang2-Peptides] of the invention as appropriate.


The [VEGF-Peptide] may be covalently linked to [Connector] through a nucleophilic side chain, or N-terminus amino group, or C-terminus carboxyl group of an VEGF-linking residue. Alternatively, where the linker is not branched, the [VEGF-peptide] may be covalently linked to the X-group of L or L' through a nucleophilic side chain or the N-terminus amino group or C-terminus carboxyl group of a VEGF-linking residue. The VEGF-linking residue may be selected from the group comprising K, R, Y, C, T, S, homologs of lysine, homocysteine, homoserine, Dap, Dab, the N-terminus residue and the C-terminus residue. The VEGF-linking residue may be selected from the group consisting of K, Y, T, Dap, and Dab. In some aspects of the invention, the VEGF-linking residue is K. The linking residue may be K¹⁰. The linking residue may be K¹². In some aspects of the invention one of the group consisting of V¹, E², P³, N⁴, V⁹, M¹⁰, V¹², X¹⁴, E¹⁷ and the C-terminus residue of the [VEGF-peptide] is substituted with a (VEGF-)linking residue comprising a nucleophilic side chain or the N-terminus amino group or the C-terminus carboxyl group covalently linked to the combining site of an antibody directly or via an intermediate linker, the linking residue being selected from the group comprising K, R, Y, C, T, S, homologs of lysine, homocysteine, homoserine, Dap, Dab, the N-terminus residue and the C-terminus residue. In some aspects where the linking residue is located at the N-terminus or C-terminus, the linking may occur through the amino group of the N-terminus or carboxyl group of the C-terminus, rather than through the respective amino acid side chain. In some aspects of the invention, one of V¹, N⁴, M¹⁰, V¹², X¹⁴, and E¹⁷ is substituted with the VEGF-linking residue. In some aspects of the invention, one of M¹⁰ and V¹² is substituted with the linking residue. In some aspects of the invention, M¹⁰ is substituted with the VEGF-linking residue. In some aspects of the invention, V¹² is substituted with the VEGF-linking residue.

The [Connector] may comprise the formula: -[AM1-Spacer]-[Branch]-[AM2-Spacer]- wherein [AM-1-Spacer] and [AM-2-Spacer] are each independently a biologically compatible polymer, block copolymer C, H, N, O, P, S, halogen (F, Cl, Br, I), or a salt thereof, alkyl, alkenyl, alkynyl, oxoalkyl, oxoalkenyl, oxoalkynyl, aminoalkyl, aminoalkenyl, aminoalkynyl, sulfoalkyl, sulfoalkenyl, sulfoalkynyl, phosphoalkyl, phosphoalkenyl, or phosphoalkynyl group, covalently bonded to [Branch], and [Branch] is molecule with at least three reactive groups, and [AM-1-Spacer] is covalently linked to [Branch] and to the [Active Molecule-1], and [AM-2-Spacer] is covalently linked to [Branch] and to the [Active Molecule-2]. [AM-1-Spacer] and [AM-2-Spacer] may each independently be a neutral, water soluble molecule able to form covalent bonds with their respective Active Molecule and [Branch]. Where the Active Molecule comprises a peptide, [AM-1-Spacer] and [AM-2-Spacer] may be able to form peptide bonds to the respective Active Molecule.


[AM-1-Spacer] and [AM-2-Spacer] may each independently be selected from the group consisting of: amino polyethylene glycol acids, polyethylene glycol diacids, amino alkanoic acids, amino alkanoic acids, and poly glycine. [AM-1-Spacer] and [AM-2-Spacer] may each independently be selected from

the group consisting of (representative molecules shown below merely to exemplify certain suitable species of Spacer):


Amino polyethylene glycol Acids




Polyethylene glycol diacids



Amino alkanolic acids




Polyglycine







5

Amino alkanolic acids






PEG molecules



10

[AM-1-Spacer] and [AM-2-Spacer] may each independently be selected from the group consisting of: 2-PEG, 4-PEG, and 6-PEG:




15

The [AM-1-Spacer] and [AM-2-Spacer] may each independently be between 6 atoms in length and 15 atoms in length. The [AM-1-Spacer] and [AM-2-Spacer] may each independently be a 4-PEG spacer. The [AM-1-Spacer] and [AM-2-Spacer] may each independently be a 2-PEG spacer. The [AM-1-Spacer] and [AM-2-Spacer] may each independently be a 1-PEG spacer. The [AM-1-Spacer] may be between 6 atoms and 15 atoms in length. [AM-1-Spacer] may be 4-PEG. [AM-2-Spacer] may be between 6 atoms and 15 atoms in length. [AM-2-Spacer] may be 4-PEG.


[Branch] may be a chemical moiety comprising three orthogonal reactive groups. [Branch] may be selected from the group consisting of: Cysteine, Diaminopropionic acid, Diaminobutanoic acid Ornithine, Lysine, Homocysteine, Bismaleimide, and Maleimide-acid, and derivatives and homologs thereof.

[Branch] may be selected from the group consisting of:


Cysteine branch,




Diaminopropionic acid based branch




Diaminobutanoic acid based branch




Ornithine based branch




Lysine based branch




Homocysteine branch



Bismaleimide branch




Maleimide-acid branch



and derivatives and homologs thereof. [Branch] may be cysteine:

Compounds of the invention may comprise the formula: [Branch]-L or [Branch]-L', wherein L is a linker, and L' is a linker covalently attached to an amino acid side chain in a combining site of an antibody, and [Branch] is covalently attached to L or L'. [Branch]-L may be selected from the group consisting of:



[Ang2 peptides]

- 10 In some aspects of the invention, [Active Molecule 2] is an Ang-2 binding peptide [Ang2-peptide]. The [Ang2-Peptide] may be covalently linked to [Connector] through a nucleophilic side chain or N-terminus or C-terminus of an Ang2-linking residue, the Ang2-linking residue being selected from the group comprising K, R, Y, C, T, S, homologs of lysine, homocysteine, homoserine, Dap, Dab, the N-terminus residue and the C-terminus residue]. Where the Ang2-linking residue is located at the C-
- 15 terminus or N-terminus, the linkage may be through the amino group of the N-terminus or the carboxyl group of the C-terminus, rather than the side chain of the particular amino acid at that

position. The Ang2-linking residue may be selected from the group consisting of K, Y, T, Dap, and Dab. The Ang2-linking residue may be K.

In some aspects of the invention, the [Ang2-Peptide] comprises a sequence substantially homologous to:

Q¹ X² Y³ Q⁴ X⁵ L⁶ D⁷ E⁸ X⁹ D¹⁰ X¹¹ X¹² X¹³ X¹⁴ D¹⁵ X¹⁶ F¹⁷ M¹⁸ X¹⁹ Q²⁰ Q²¹ G²² (SEQ ID NO:107)

wherein X² is selected from the group consisting of K, N, R, H, Kac, Nick, and CbcK, and X⁵ is selected from the group consisting of P, hP, dhP, and BnHP, and X⁹ is selected from the group consisting of L, I, ThA, and Kac, and X¹¹ is selected from the group consisting of Q, N, C, K, Kac, Dab, and Dap, and X¹² is selected from the group consisting of L, HL, Nva, I, HchA, HF, and ThA, and X¹³ is selected from the group consisting of L, HL, Nva, I, HchA, HF, and ThA, and X¹⁴ is selected from the group consisting of aromatic residues, and X¹⁶ is selected from the group consisting of Q and N, and X¹⁹ is selected from the group consisting of L, and I, and wherein one of Q¹, E⁸, X⁹, X¹¹, X¹², D¹⁵, X¹⁶, M¹⁸, X¹⁹ or G²² is substituted with the Ang2-linking residue.

In some aspects, X² is selected from the group consisting of K, N, and Kac. X² may be N. X² may be Kac. In some aspects, X⁵ is selected from the group consisting of P, hP, and dhP. X⁵ may be P. In some aspects, X⁹ is L. In some aspects, X⁹ is Kac. In some aspects, X¹¹ is selected from the group consisting of K, Kac. In some aspects, X¹³ is selected from the group consisting of L, HL, Nva, I. X¹³ may be L. In some aspects, X¹⁴ is selected from the group consisting of F, Y, W, BPA, CF, NF. X¹⁴ may be Y. In some aspects, X¹⁶ is Q. The Ang2-linking residue may be substituted for one of the group consisting of X⁹, X¹¹, X¹², D¹⁵, X¹⁶, M¹⁸ and X¹⁹. X¹¹ may be the location of the Ang2-linking residue.

The [Ang2-peptide] may comprise a sequence substantially homologous to one or more compounds selected from the group consisting of: SEQ ID NOs:137-191. The Ang-2 peptide may comprise a compound selected from the group consisting of SEQ ID NOs:137-172 and SEQ ID NO:182. The [Ang2-peptide] may comprise a sequence substantially homologous to one or more compounds selected from the group consisting of: SEQ ID NO:137, SEQ ID NO:138, SEQ ID NO:139, SEQ ID NO:140, SEQ ID NO:153, SEQ ID NO:154, SEQ ID NO:155, SEQ ID NO:157, SEQ ID NO:161, SEQ ID NO:174, and SEQ ID NO:181.

The [Ang2-peptide] may comprises a sequence substantially homologous to

Q(Kac)YQPLDE(Kac)DKTLYDQFMLQQG (SEQ ID NO:153). R¹ may be C(O)CH₃. R² may be NH₂. In some aspects, R¹-[Ang-peptide]-R² may be

{C(O)CH₃}Q(Kac)YQPLDE(Kac)DKTLYDQFMLQQ G-{NH₂} (SEQ ID NO:153) and K¹¹ is the

Ang2-linking residue covalently linked to the [Connector].

In some aspects of the invention, the compound may be selected from the group consisting of: compounds 5001-5028, and 5031-5062. In some aspects, the compound is selected from the group consisting of compounds 5001-5028 and compounds 5031-5074. In some aspects, the compound is selected from the group consisting of 5053, 5060, 5061 and 5062. In some aspects, the compound is compound 5053. In some aspects, the compound is 5037. In some aspects of the invention, the compound selected from the group consisting of: compounds 6001-6028, and 6031-6062. In some aspects, the compound is selected from the group consisting of compounds 6001-6026, 6029, 6031-6069. In some aspects, the compound is selected from the group consisting of compounds 6001-6028 and compounds 6031-6074. In some aspects, the compound is selected from the group consisting of compounds 6053, 6060, 6061 and 6062. In some aspects, the compound is compound 6053. In some aspects, the compound is compound 6037.

Antibodies

The contents of US2006205670 are incorporated herein by reference- in particular paragraphs [0153]-[0233], describing antibodies, useful fragments and variants and modifications thereof, combining sites and CDRs, antibody preparation, expression, humanization, aa modification, glycosylation, ADCC, CDC, increasing serum half life of antibodies, expression vectors, mammalian host systems, and folding, amongst other elements of antibody technology.

As discussed, in certain embodiments, certain antibodies that can be used in conjunction with compounds of the invention may require a reactive side chain in the antibody combining site. A reactive side chain may be present naturally or may be placed in an antibody by mutation. The reactive residue of the antibody combining site may be associated with the antibody, such as when the residue is encoded by nucleic acid present in the lymphoid cell first identified to make the antibody. Alternatively, the amino acid residue may arise by purposely mutating the DNA so as to encode the particular residue (see, e.g., WO 01/22922 to Meares *et al.*). The reactive residue may be a non-natural residue arising, for example, by biosynthetic incorporation using a unique codon, tRNA, and aminoacyl-tRNA as discussed herein. In another approach, the amino acid residue or its reactive functional groups (e.g., a nucleophilic amino group or sulfhydryl group) may be attached to an amino acid residue in the antibody combining site. Thus, covalent linkage with the antibody occurring “through an amino acid residue in a combining site of an antibody” as used herein means that linkage can be directly to an amino acid residue of an antibody combining site or through a chemical moiety that is linked to a side chain of an amino acid residue of an antibody combining site. In some embodiments, the amino acid is cysteine, and the reactive group of the side chain is a sulfhydryl group. In other embodiments, the amino acid residue is lysine, and the reactive group of the side chain is the ϵ -amino group.

Catalytic antibodies are one source of antibodies with suitable combining sites that comprise one or more reactive amino acid side chains. Such antibodies include aldolase antibodies, beta lactamase antibodies, esterase antibodies, amidase antibodies, and the like.

- 5 One embodiment comprises an aldolase antibody such as the mouse monoclonal antibodies mAb 33F12 and mAb 38C2, as well as suitably chimeric and humanized versions of such antibodies (e.g., h38C2, SEQ ID NOs:1 and 2). Mouse mAb 38C2 (andh38C2) has a reactive lysine near to but outside HCDR3, and is the prototype of a new class of catalytic antibodies that were generated by reactive immunization and mechanistically mimic natural aldolase enzymes. See C.F. Barbas 3rd *et al.*,
10 Science 278:2085-2092 (1997)). Other aldolase catalytic antibodies that may be used include the antibodies produced by the hybridoma 85A2, having ATCC accession number PTA-1015; hybridoma 85C7, having ATCC accession number PTA-1014; hybridoma 92F9, having ATCC accession number PTA-1017; hybridoma 93F3, having ATCC accession number PTA-823; hybridoma 84G3, having ATCC accession number PTA-824; hybridoma 84G11, having ATCC accession number PTA-1018;
15 hybridoma 84H9, having ATCC accession number PTA-1019; hybridoma 85H6, having ATCC accession number PTA-825; hybridoma 90G8, having ATCC accession number PTA-1016. Through a reactive lysine, these antibodies catalyze aldol and retro-aldol reactions using the enamine mechanism of natural aldolases. See, e.g., J. Wagner *et al.*, Science 270:1797-1800 (1995); C.F. Barbas 3rd *et al.*, Science 278:2085-2092 (1997); G. Zhong *et al.*, Angew. Chem. Int. Ed. Engl.
20 38:3738-3741 (1999); A. Karlstrom *et al.*, Proc. Natl. Acad. Sci. U.S.A., 97:3878-3883 (2000). Aldolase antibodies and methods of generating aldolase antibodies are disclosed in U.S. Patents Nos. 6,210,938, 6,368,839, 6,326,176, 6,589,766, 5,985,626, and 5,733,75, which are incorporated herein by reference.
- 25 Compounds of the invention may also be formed by linking a compound of the invention to a reactive cysteine, such as those found in the combining sites of thioesterase and esterase catalytic antibodies. Suitable thioesterase catalytic antibodies are described by K.D. Janda *et al.*, Proc. Natl. Acad. Sci. U.S.A. 91:2532-2536 (1994). Suitable esterase antibodies are described by P. Wirsching *et al.*, Science 270:1775-1782 (1995). Reactive amino acid-containing antibodies may be prepared by
30 means well known in the art, including mutating an antibody combining site residue to encode for the reactive amino acid or chemically derivatizing an amino acid side chain in an antibody combining site with a linker that contains the reactive group.

Antibody may be a humanized antibody. Where compounds of the invention are covalently linked to
35 the combining site of an antibody, and such antibodies are humanized, it is important that such antibodies be humanized with retention of high linking affinity for the Z group. Various forms of humanized murine aldolase antibodies are contemplated. One embodiment uses the humanized

aldolase catalytic antibody h38c2 IgG1 or h38c2 Fab with human constant domains C_κ and C_{γ1}1. C. Rader *et al.*, J. Mol. Bio. 332:889-899 (2003) discloses the gene sequences and vectors that may be used to produce h38c2 Fab and h38c2 IgG1. Human germline V_κ gene DPK-9 and human J_κ gene JK4 were used as frameworks for the humanization of the kappa light chain variable domain of m38c2, and human germline gene DP-47 and human J_H gene JH4 were used as frameworks for the humanization of the heavy chain variable domain of m38c2. FIGURE 2A illustrates a sequence alignment between the variable light and heavy chains in m38c2, h38c2, and human germlines. h38c2 may utilize IgG1, IgG2, IgG3, or IgG4 constant domains, including any of the allotypes thereof. FIGURE 2B illustrates one embodiment of h38c2 IgG1 using the G1m(f) allotype, where the light and heavy chain amino acid sequences of this h38c2 IgG1 are set forth in the figure. In certain embodiments of compounds of the invention wherein Antibody is h38c2 IgG1 with the G1m(f) allotype, Z binds to the side chain of the lysine residue at position 99 of the heavy chain. This residue is denoted by bold print in FIGURE 2B. Another embodiment uses a chimeric antibody comprising the variable domains (V_L and V_H) of h38c2 and the constant domains from an IgG1, IgG2, IgG3, or IgG4. Antibody may be a full-length antibody, Fab, Fab', F(ab')₂, F_v, dsF_v, scF_v, V_H, V_L, diabody, or minibody comprising V_H and V_L domains from h38c2. Antibody may be an antibody comprising the V_H and V_L domains from h38c2 and a constant domain selected from the group consisting of IgG1, IgG2, IgG3, and IgG4. Antibody may be h38c2 IgG1. Antibody may be a humanized version of a murine aldolase antibody comprising a constant region from a human IgG, IgA, IgM, IgD, or IgE antibody. In another embodiment, the Antibody is a chimeric antibody comprising the variable region from a murine aldolase antibody and a constant region from a human IgG, IgA, IgM, IgD, or IgE antibody. In further embodiments, the Antibody is a fully human version of a murine aldolase antibody comprising a polypeptide sequence from natural or native human IgG, IgA, IgM, IgD, or IgE antibody.

Various forms of humanized aldolase antibody fragments are also contemplated. One embodiment uses h38c2 F(ab')₂. h38c2 F(ab')₂ may be produced by the proteolytic digestion of h38c2 IgG1. Another embodiment uses an h38c2 scFv comprising the V_L and V_H domains from h38c2 which are optionally connected by the intervening linker (Gly₄Ser)₃. As an alternative to humanization, human antibodies can be generated. For example, it is now possible to produce transgenic animals (e.g., mice) that are capable, upon immunization (or reactive immunization in the case of catalytic antibodies) of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production.

Alternatively, phage display technology (see, e.g., J. McCafferty *et al.*, Nature 348:552-553 (1990); H.J. de Haard *et al.*, J Biol Chem 274, 18218-18230 (1999); and A. Kanppik *et al.*, J Mol Biol, 296, 57-86 (2000)) can be used to produce human antibodies and antibody fragments *in vitro* using

immunoglobulin variable (V) domain gene repertoires from unimmunized donors. As indicated above, human antibodies may also be generated by *in vitro* activated B cells. See, e.g., U.S. Patent Nos. 5,567,610 and 5,229,275; and C.A.K. Borrebaeck *et al.*, Proc. Natl. Acad. Sci. U.S.A. 85:3995-3999 (1988).

Amino acid sequence modification(s) of the antibodies described herein are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody. Amino acid sequence variants of an antibody are prepared by introducing appropriate nucleotide changes into the antibody nucleic acid, or by peptide synthesis. Such modifications include, for example, deletions from, insertions into, and/or substitutions of residues within the amino acid sequences of the antibody. Any combination of deletion, insertion, and substitution is made to arrive at the final construct, provided that the final construct possesses the desired characteristics. The amino acid changes also may alter post-translational processes of the antibody, such as changing the number or position of glycosylation sites. A useful method for identification of certain residues or regions of an antibody that are preferred locations for mutagenesis is called "alanine scanning mutagenesis," as described in B.C. Cunningham and J.A. Wells, Science 244:1081-1085 (1989).

Exemplary uses of compounds and compositions of the invention

The invention provides for the use of compounds of the invention or pharmaceutical compositions of the invention in a method of inhibiting or reducing angiogenesis or for treating or preventing a disease or symptom associated with an angiogenic disorder. The invention provides methods of inhibiting or reducing angiogenesis or treating or preventing a disease or symptom associated with an angiogenic disorder comprising administering to a patient a therapeutically effective dose of compounds and compositions of the invention. Also provided are methods of delivering or administering compounds and compositions of the invention and methods of treatment using compounds and compositions of the invention. As used herein, an angiogenesis-mediated condition is a condition that is caused by abnormal angiogenesis activity or one in which compounds that modulate angiogenesis activity have therapeutic use. One aspect of the invention provides methods for modulating VEGF activity *in vivo* comprising administering an effective amount of a compound or composition as described herein to a subject. Another aspect of the invention includes methods of using compounds and compositions of the invention for diagnostic purposes. Diseases and conditions that may be treated and/or diagnosed with compounds and compositions of the invention include cancer, arthritis, hypertension, kidney disease, psoriasis, angiogenesis of the eye associated with ocular disorder, infection or surgical intervention, macular degeneration, diabetic retinopathy, and the like.

More specifically, examples of "cancer" when used herein in connection with the present invention include cancer selected from lung cancer (NSCLC and SCLC), cancer of the head or neck, ovarian

cancer, colon cancer, rectal cancer, prostate cancer, cancer of the anal region, stomach cancer, breast cancer, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, neoplasms of the central nervous system (CNS), primary CNS lymphoma, non-Hodgkins's lymphoma, spinal axis tumors, carcinomas of the, oropharynx, hypopharynx, esophagus, pancreas, liver, gallbladder and bile ducts, small intestine, urina; or lymphoma or a combination of one or more of the foregoing cancers. Still more specifically, examples of "cancer" when used herein in connection with the present invention include cancer selected from lung cancer (NSCLC and SCLC), breast cancer, ovarian cancer, colon cancer, rectal cancer, prostate cancer, cancer of the anal region, or a combination of one or more of the foregoing cancers.

In some aspects, compounds of the invention may be used in the treatment and/or prevention of ophthalmic diseases such as age-related macular degeneration (both wet and dry), glaucoma, diabetic retinopathy (including diabetic macular edema), choroidal neovascular membrane (CNV), uveitis, myopic degeneration, ocular tumors, central retinal vein occlusion, rubeosis, ocular neovascularization, central serous retinopathy, ocular surface discus such as dry eye, central retinal artery occlusion, cystoid macular edema and other retinal degenerative disease. In some embodiments, it may be advantageous to employ molecules with high binding affinity for the respective target(s), but with a lower PK or half-life, as there will be little enzymatic degradation of the molecule while in the eye, but once the molecule clears the eye, it may be desirable for the molecule to be degraded and undergo renal clearance as soon as possible so as to minimize potential direct effect or side-effects outside the eye. In some aspects of the invention Compounds 6037 is used for the treatment of ophthalmic diseases. In some aspects of the invention Compounds 6044 is used for the treatment of ophthalmic diseases. In some aspects of the invention Compounds 6053 is used for the treatment of ophthalmic diseases.

The compounds according to the invention may be solvated, especially hydrated. Hydration may occur during manufacturing of the compounds or compositions comprising the compounds, or the hydration may occur over time due to the hygroscopic nature of the compounds.

In another aspect, the present invention includes methods of altering at least one physical or biological characteristic of a compounds and compositions. The methods include covalently linking a [VEGF-Peptide] of the invention to a combining site of an antibody, either directly or though a linker. Characteristics of compounds of the invention that may be modified include, but are not limited to, binding affinity, susceptibility to degradation (e.g., by proteases), pharmacokinetics, pharmacodynamics, immunogenicity, solubility, lipophilicity, hydrophilicity, hydrophobicity, stability (either more or less stable, as well as planned degradation), rigidity, flexibility, modulation of antibody binding, and the like. Also, the biological potency of a particular compound of the invention

may be increased by the addition of the effector function(s) provided by the antibody. For example, an antibody provides effector functions such as complement mediated effector functions. Without wishing to be bound by any theory, the antibody portion of a compound of the invention may generally extend the half-life of a smaller sized [VEGF-Peptide] *in vivo*. Thus, in one aspect, the invention provides a method for increasing the effective circulating half-life of a [VEGF-Peptide].

In another aspect, the present invention includes methods of modifying a combining site of an antibody to generate binding specificity for VEGF, or VEGF and one or more other Active Molecules, including Ang2. Such methods include covalently linking a reactive amino acid side chain in a combining site of the antibody to a chemical moiety on a linker of [VEGF-Peptide]-linker compound as described herein. The chemical moiety of the linker is sufficiently distanced from the [VEGF-Peptide] so that the [VEGF-Peptide] can bind its cognate when the [VEGF-Peptide]-linker compound is covalently linked to an antibody combining site (similar constraints will exist for tethering to other half-life increasing macromolecules). Typically, the antibody will not be considered specific for the target molecule. In certain embodiments, an antibody prior to covalent linking would have an affinity for VEGF of less than about 1×10^{-5} moles/liter. However, after the antibody is covalently linked to the [VEGF-Peptide], the modified antibody preferably has an affinity for the target molecule of at least about 1×10^{-6} moles/liter, alternatively, at least about 1×10^{-7} moles/liter, alternatively, at least 1×10^{-8} moles/liter, alternatively at least 1×10^{-9} moles/liter, or alternatively, at least about 1×10^{-10} moles/liter.

The administration of [VEGF-Peptide] of the invention to an immunocompetent individual may result in the production of antibodies against the conjugate. Such antibodies may be directed to the variable region, including the antibody idiotype, as well as to the targeting agent or any linker used to conjugate the targeting agent to the antibody. Reducing the immunogenicity of a [VEGF-Peptide] can be accomplished by methods well known in the art, such as by attaching long chain polyethylene glycol (PEG)-based spacers and the like to the [VEGF-Peptide]. Long chain PEG and other polymers are known for their ability to mask foreign epitopes, resulting in the reduced immunogenicity of therapeutic proteins that display foreign epitopes (N.V. Katre, J. Immunol. 144:209-213 (1990); G.E. Francis *et al.*, Int. J. Hematol. 68:1-18 (1998). Alternatively, or in addition, the individual administered the antibody-[VEGF-Peptide] conjugate may be administered an immunosuppressant such as cyclosporin A, anti-CD3 antibody, and the like. The above also applies *mutatis mutandis* when other Active Molecules, such as Ang2, are covalently linked, via a branched linker, to the compounds of the invention

The invention also provides for stereoisomers, tautomers, solvates, prodrugs, and pharmaceutically acceptable salts of compounds of the invention.

Pharmaceutical Compositions, Methods of Administration and Combination Therapies

The invention also provides pharmaceutical compositions comprising a therapeutically effective amount of one or more compounds of the invention. The invention also provides methods of treatment using such compositions, and methods of preparing pharmaceutical compositions of the invention.

The contents of US2006205670 are incorporated herein by reference- in particular paragraphs [0497]-[0510], describing administration ([0498],[0499]; formulations ([0499],[0500], [0501], [0502]); preparations ([0503],[504]); kits ([0505]); dosage strategies ([0506], [0507], [0508], [0509]); as per the respective targeting agent/ AA argeting agent/ [Peptide] of the invention, and methods of visualizing or localizing the biological target to which the [Peptide] of the invention is directed.

The invention also includes administration of one or more compounds or compositions of the invention in combination with one or more oncology therapeutics, each being administered according to a regimen suitable for that therapeutic. The components of the combination therapy may be administered concurrently or non-concurrently. Components may be administered in the same or in separate compositions, and by the same or different routes of administration. Examples of suitable oncology therapeutics and combinations that may be used in combination with peptides and compounds of the invention are listed in Tables 4-6 of US2006205670 (which is incorporated herein by reference). The pharmaceutical compositions of the invention may further comprise a therapeutically effective amount of one or more chemotherapeutic agent, the chemotherapeutic agent preferably being a compound selected from the group consisting of 5-Flurouracil, Irinotecan, Taxotere, Sunitinib , Axitinib, oxilaplatin, bevacizumab, cetuximab, or chemical equivalents thereof. In some aspects, the chemotherapeutic agent is selected from the group consisting of 5-Flurouracil, Irinotecan, Taxotere , Sunitinib, and Axitinib; or chemical equivalents thereof.

Examples of synthesis of compounds of the invention

Compounds of the invention can be prepared using techniques known in the art. Typically, synthesis of the [VEGF-Peptide] or [Ang2-Peptide] (or other Active Molecule) is the first step and is carried out as described herein. The Active Molecule is then derivatized for linkage to a connecting component (the linker), which is then combined with the antibody. One of skill in the art will readily appreciate that the specific synthetic steps used depend upon the exact nature of the three components. Thus, [VEGF-Peptide] – linker conjugates and compounds of the invention described herein can be readily synthesized. [VEGF-Peptides] and [Ang2-Peptides] may be synthesized by many techniques that are known to those skilled in the art. For solid phase peptide synthesis, a summary of exemplary techniques may be found in *Chemical Approaches to the Synthesis of Peptides and Proteins* (Williams *et al.*, eds.), CRC Press, Boca Raton, FL (1997).

Typically, the desired peptidic Active Molecule is synthesized sequentially on solid phase according to procedures well known in the art. See, e.g., U.S. Patent Application No. 2003/0045477). The linker may be attached to the peptide in part or in full on the solid phase, or may be added using solution phase techniques after the removal of the peptide from the resin (see FIGURES 1A and 1B).

For example, an N-protected amino and carboxylic acid-containing linking moiety may be attached to a resin such as 4-hydroxymethyl-phenoxyethyl-poly(styrene-1% divinylbenzene). The N-protecting group may be removed by the appropriate acid (e.g., TFA for Boc) or base (e.g., piperidine for Fmoc), and the peptide sequence developed in the normal C-terminus to N-terminus fashion (see FIGURE 1A). Alternatively, the peptide sequence may be synthesized first and the linker added to the N-terminal amino acid residue last (see FIGURE 1B). Yet another method entails deprotecting an appropriate side chain during synthesis and derivatizing with a suitably reactive linker. For example, a lysine side chain may be deprotected and reacted with a linker having an active ester. Alternatively, an amino acid derivative with a suitably protected linker moiety already attached to the side chain (see FIGURE 1B) or, in some cases, the alpha-amino nitrogen, may be added as part of the growing peptide sequence.

At the end of the solid phase synthesis, the targeting agent-linker conjugate is removed from the resin and deprotected, either in succession or in a single operation. Removal of the targeting agent-linker conjugate and deprotection can be accomplished in a single operation by treating the resin-bound peptide-linker conjugate with a cleavage reagent, for example, trifluoroacetic acid containing scavengers such as thianisole, water, or ethanedithiol. After deprotection and release of the targeting agent, further derivatization of the targeting agent peptide may be carried out. The fully deprotected peptidyl Active Molecule-linker conjugate is purified by a sequence of chromatographic steps employing any or all of the following types: ion exchange on a weakly basic resin in the acetate form; hydrophobic adsorption chromatography on underivatized polystyrene-divinylbenzene (e.g., AMBERLITE XAD); silica gel adsorption chromatography; ion exchange chromatography on carboxymethylcellulose; partition chromatography, e.g., on SEPHADEX G-25, LH-20 or countercurrent distribution; high performance liquid chromatography (HPLC), especially reverse-phase HPLC on octyl- or octadecylsilyl-silica bonded phase column packing.

Brief Description of the Drawings

FIGURES 1A and FIGURE 1B illustrate the solid phase synthesis of targeting agent-linker conjugates of the present invention.

FIGURE 2A illustrates the amino acid sequence alignment of the variable domains of m38c2, h38c2, and human germlines. Framework regions (FR) and complementarity determining regions (CDR) are defined according to Kabat *et al.* Asterisks mark differences between m38c2 and h38c2 or between

h38c2 and the human germlines. FIGURE 2B illustrates the amino acid sequence of the light and heavy chains (SEQ ID NOs:1 and 2, respectively) of one embodiment of a humanized 38c2 IgG1.

FIGURE 3 shows various structures that may serve as linker reactive groups. Structures A-C form reversible covalent bonds with surface accessible reactive nucleophilic groups (e.g., lysine or cysteine side chain) of a combining site of an antibody. R'₁, R'₂, R'₃, and R₄ in structures A-C represent substituents which include, for example, C, H, N, O, P, S, halogen (F, Cl, Br, I) or a salt thereof. X is N, C, or any other heteroatom. These substituents may also include a group such as an alkyl, alkenyl, alkynyl, oxoalkyl, oxoalkenyl, oxoalkynyl, aminoalkyl, aminoalkenyl, aminoalkynyl, sulfoalkyl, sulfoalkenyl, or sulfoalkynyl group, phosphoalkyl, phosphoalkenyl, phosphoalkynyl group. R'₂ and R'₃ could be cyclic as exemplified in structures B and C while X could be a heteroatom. For example, structure A could form an irreversible covalent bond with a reactive nucleophile if X is N and if R'₁ and R'₃ form part of a cyclic structure. Structures D-G may form nonreversible covalent bonds with reactive nucleophilic groups in a combining site of an antibody. In these structures, R''₁ and R''₂ represent C, O, N, halide or leaving groups such as mesyl or tosyl.

FIGURE 4 shows various electrophiles that are suitable for reactive modification with a reactive amino acid side chain in a combining site of an antibody and thus may serve as linker reactive groups. Key: (A) acyl beta-lactam; (B) simple diketone; (C) succinimide active ester; (D) maleimide; (E) haloacetamide with linker; (F) halo ketone; (G) cyclohexyl diketone; and (H) aldehyde. The squiggle line indicates the point of attachment to the rest of the linker or targeting agent. X refers to a halogen.

FIGURE 5 shows the addition of a nucleophilic ("nu") side chain in an antibody combining site to compounds A-G in FIGURE 3. Antibody- Nu- refers to a covalent bond to an amino acid side chain bearing a nucleophile in a combining site of an antibody.

FIGURE 6 shows the addition of a nucleophilic side chain in an antibody combining to compounds A-H in FIGURE 5. Antibody- Nu- refers to a covalent bond to an amino acid side chain bearing a nucleophile in a combining site of an antibody.

FIGURE 7 shows a synthesis of:




FIGURE 8: Exemplary synthesis of [Ang2-peptide]-[Ang2-Spacer], using SEQ ID NO:153 as example.

FIGURE 9: Exemplary synthesis of [VEGF-Peptide]-[VEGF-Spacer], using SEQ ID NO:188 as example.

5 FIGURE 10: Exemplary synthesis of compound 5053.

FIGURE 11: Tumor volume of Colo205 colon adenocarcinoma xenografts after treatment with Vehicle or Compound 6053 (IP, 1x/wk). Data are depicted as the mean and SE of n=20/group for days 0-35 (n=10 for all groups beyond day 35), vertical arrow indicates first dosing day. * $P < 0.05$,
10 ** $P < 0.01$ versus Vehicle at day 35 (One-way ANOVA with Dunnett's Multiple Comparison Test).

FIGURE 12: Tumor microvessel density of Colo205 colon adenocarcinoma xenografts after treatment with Vehicle or Compound 6053 (IP, 1x/wk). Data are depicted as the mean and SE of n=9-10/group. * $P < 0.05$ versus Vehicle (One-way ANOVA with Dunnett's Multiple Comparison Test).

15

FIGURE 13: Tumor volume of Colo205 colon adenocarcinoma xenografts after treatment with Vehicle or Compound 6053 (IP, 1x/wk). Data are depicted as the mean and SE of n=10/group, vertical arrow indicates first dosing day. ** $P < 0.01$ versus Vehicle (One-way ANOVA with Dunnett's Multiple Comparison Test).

20

FIGURE 14: Viable tumor volume of Colo205 colon adenocarcinoma xenografts after treatment with Vehicle or Compound 6053 (IP, 1x/wk). Data are depicted as the mean and SE of n=9-10/group. ** $P < 0.01$ versus Vehicle (One-way ANOVA with Dunnett's Multiple Comparison Test).

25 FIGURE 15: Ang2 immunoreactivity of Colo205 colon adenocarcinoma xenografts after treatment with Vehicle or Compound 6053 (IP, 1x/wk). Data are depicted as the mean and the SE of n=9-10/group. ** $P < 0.01$ versus Vehicle (One-way ANOVA with Dunnett's Multiple Comparison Test).

FIGURE 16: Phosphorylated VEGFR2 (pVEGFR2) immunoreactivity as a percentage of total
30 VEGFR2 immunoreactivity of Colo205 colon adenocarcinoma xenografts after treatment with Vehicle or Compound 6053 (IP, 1x/wk). Data are depicted as the mean and the SE of n=9-10/group. ** $P < 0.01$ versus Vehicle (One-way ANOVA with Dunnett's Multiple Comparison Test).

FIGURE 17: Tumor volume of MDA-MB-435 breast adenocarcinoma (A) and A431 skin carcinoma
35 (B) after weekly treatment with Vehicle or Compound 6053 (IP, 1x/wk). Data are depicted as the mean and SE of n=9-10/group, vertical arrow indicates first dosing day. * $P < 0.05$ vs Vehicle at day

68; $**P < 0.01$ versus Vehicle at day 35 (One-way ANOVA with Dunnett's Multiple Comparison Test).

FIGURE 18: Tumor volume of Colo205 colon adenocarcinoma xenografts after once weekly treatment with Vehicle, Compound 6053 alone (IP), 5-FU alone (IP), or Compound 6053 in combination with 5-FU (both IP). Data are depicted as the mean and SE of $n=9-10$ /group, vertical arrow indicates first dosing day. $**P < 0.01$ vs Vehicle at day 29 (One-way ANOVA with Bonferroni's Multiple Comparison Test).

FIGURE 19: Tumor volume of Colo205 colon adenocarcinoma xenografts after weekly treatment with Vehicle, Compound 6053 alone (IP), Irinotecan alone (IP), or Compound 6053 in combination with Irinotecan (both IP). Data are depicted as the mean and SE of $n=9-10$ /group, vertical arrow indicates first dosing day. $**P < 0.01$ vs Vehicle at day 29 (One-way ANOVA with Bonferroni's Multiple Comparison Test).

FIGURE 20: Tumor volume of Colo205 colon adenocarcinoma xenografts after once weekly treatment with Vehicle, Compound 6053 alone (IP), Taxotere alone (IP), or Compound 6053 in combination with Taxotere (both IP). Data are depicted as the mean and SE of $n=9-10$ /group, vertical arrow indicates first dosing day. $**P < 0.01$ vs Vehicle, $***P < 0.01$ vs Vehicle and vs Compound 6053 alone at day 29 (One-way ANOVA with Bonferroni's Multiple Comparison Test).

FIGURE 21: Tumor volume of Colo205 colon adenocarcinoma xenografts after once weekly treatment with Vehicle, Compound 6053 alone (IP), daily Sunitinib (PO), or Compound 6053 (IP) in combination with Sunitinib (PO). Data are depicted as the mean and SE of $n=9-10$ /group, vertical arrow indicates first dosing day. $*P < 0.05$, $**P < 0.01$ vs Vehicle at day 28 (One-way ANOVA with Bonferroni's Multiple Comparison Test).

FIGURE 22: Tumor volume of Colo205 colon adenocarcinoma xenografts after once weekly treatment with Vehicle, Compound 6053 alone (IP), daily Axitinib (PO), or Compound 6053 (IP) in combination with Axitinib (PO). Data are depicted as the mean and SE of $n=9-10$ /group, vertical arrow indicates first dosing day. $*P < 0.05$, $**P < 0.01$ vs Vehicle at day 28 (One-way ANOVA with Bonferroni's Multiple Comparison Test).

FIGURE 23: Tumor volume of Colo205 colon adenocarcinoma xenografts after once weekly treatment with Vehicle, Compound 2071, Compound 2049, or Compound 2046 (all IP). Data are depicted as the Mean and SE of $n=10$ /group, vertical arrow indicates first dosing day.

FIGURE 24: Tumor volume of Colo205 colon adenocarcinoma xenografts after once weekly treatment with Vehicle, Compound 2049, Compound 6045, Compound 6042, or Compound 6053 (all IP). Data are depicted as the Mean and SE of n=10/group, vertical arrow indicates first dosing day. ** $P<0.01$ versus Vehicle on day 28 (Compounds 6045, 6042, and 6053).

5

FIGURE 25: Tumor volume of Colo205 colon adenocarcinoma xenografts after once weekly treatment with Vehicle, Compound 4043, Compound 2018, or Compound 6053 (all IP). Data are depicted as the Mean and SE of n=10/group, vertical arrow indicates first dosing day. ** $P<0.01$ versus Vehicle on day 28 (Compounds 4043 and 2018), *** $P<0.001$ versus Vehicle on day 28 (Compound 6053, 10 and 30 mg/kg weekly doses).

10

FIGURE 26: Tumor volume of Colo205 colon adenocarcinoma xenografts after once weekly treatment with Vehicle, Compound 2018, Compound 4043, Compound 6053, Compound 6062, or the combination of Compounds 2018 and 4043 (all IP). Data are depicted as the Mean and SE of n=9/group, vertical arrow indicates first dosing day.

15

FIGURE 27: Tumor volume of Colo205 colon adenocarcinoma xenografts after once weekly treatment with Vehicle, Compound 4043, Compound 2018, Compound 6053, or the combination of Compounds 4043 and 2018 (all IP). Data are depicted as the Mean and SE of n=10/group, vertical arrow indicates first dosing day. ** $P<0.01$ versus Vehicle on day 42 (Compounds 4043 and 6053, and the combination of Compounds 4043 and 2018).

20

FIGURE 28: Tumor volume of Colo205 colon adenocarcinoma xenografts after once weekly treatment with Vehicle, Compound 4043, Compound 2018, Compound 6053, or the combination of Compounds 4043 and 2018 (all IP). Data are depicted as the Mean and SE of n=10/group, vertical arrow indicates first dosing day. * $P<0.05$ versus Vehicle on day 32 (Compounds 2018, and the combination of Compounds 4043 and 2018), ** $P<0.01$ versus Vehicle on day 32 (Compound 6053).

25

FIGURE 29: Fluorescein angiography grading in rabbit VEGF induced retinal leakage 3 days post IVT injection and 48hrs post VEGF dose. Compound 6037 dose response study.

30

FIGURE 30: Fluorescein angiography grading in rabbit VEGF induced retinal leakage 7 days post IVT injection and 48hrs post VEGF dose

35 Definitions


The following abbreviations, terms and phrases are used herein as defined below.

Non-natural amino acid abbreviations


BnHP refers to (2S,4R)-4-Hydroxyproline

Tle refers to tertbutylGlycine


Methyl tyrosine or Tyr-OMe (Yme) refers to:




Norvaline or Nva refers to:




Dehydroproline or DHP refers to




Hydroxyproline or HP refers to:




Homocyclohexyl alanine or HChA refers to:




Homophenyl alanine or HF refers to:




Thiazolyl alanine or ThA refers to:




Homoleucine or HL refers to




Epsilon acyl lysine or Kac (also AcK) refers to:




4-benzoyl phenylalanine or BPA refers to:




Epsilon chloro benzyl carbamate lysine or (ClBnCarbamate) K or CbcK refers to:




Thienyl Alanine or TA refers to



5 Diaminobutyric acid or Dab refers to:




Diaminopropionic acid or Dap refers to:




4-Nitro phenylalanine or NO2F or NF:


10




4-Carboxy phenylalanine or (CO2H)F or CF




15 Amido 2-PEG refers to:




Nicotinyl Lysine (NickK) refers to:




Difluorobenzoyl or DFB refers to:




Dichlorobenzoyl or DCB refers to:



“O” PEG or OP refers to:




Pyridinyl carboxylate or PyC refers to:




5


“4” PEG or 4P (shown attached to a lysine residue) refers to:



Sarcosine or Sar refers to:




Norleucine (Nle) refers to:




10


2-Aminobutyric Acid (Aib) refers to:




O-methyl-L-tyrosine (Yme) refers to:



beta-cyclohexyl-L-alanine (Cha) refers to:



L-Tyrosine methyl ester (YOMe) refers to:



15

Unless indicated otherwise by a "D" prefix, e.g., D-Ala or N-Me-D-Ile, or written in lower case format, e.g., a, i, l, (D versions of Ala, Ile, Leu), the stereochemistry of the alpha-carbon of the amino acids and aminoacyl residues in peptides described in this specification and the appended claims is the natural or "L" configuration. The Cahn-Ingold-Prelog "R" and "S" designations are used to specify the stereochemistry of chiral centers in certain acyl substituents at the N-terminus of the peptides. The designation "R,S" is meant to indicate a racemic mixture of the two enantiomeric forms. This nomenclature follows that described in R. S. Cahn, *et al.*, *Angew. Chem. Int. Ed. Engl.*, 5:385-415 (1966).

"Polypeptide," "peptide," and "protein" are used interchangeably to refer to a polymer of amino acid residues. As used herein, these terms apply to amino acid polymers in which one or more amino acid residues is an artificial chemical analog of a corresponding naturally occurring amino acid. These terms also apply to naturally occurring amino acid polymers. Amino acids can be in the L or D form as long as the binding function of the peptide is maintained. Peptides may be cyclic, having an intramolecular bond between two non-adjacent amino acids within the peptide, e.g., backbone to backbone, side-chain to backbone and side-chain to side-chain cyclization. Cyclic peptides can be prepared by methods well known in the art. See e.g., U.S. Patent No. 6,013,625.

All peptide sequences are written according to the generally accepted convention whereby the alpha-N-terminal amino acid residue is on the left and the alpha-C-terminal amino acid residue is on the right. As used herein, the term "N-terminus" refers to the free alpha-amino group of an amino acid in a peptide, and the term "C-terminus" refers to the free α -carboxylic acid terminus of an amino acid in a peptide. A peptide which is N-terminated with a group refers to a peptide bearing a group on the alpha-amino nitrogen of the N-terminal amino acid residue. An amino acid which is N-terminated with a group refers to an amino acid bearing a group on the alpha-amino nitrogen.

In general, the term "hydrophobic amino acids" or "non-polar amino acids" means amino acid residues that do not contain an ionized group(s) at physiological pH. Examples of hydrophobic amino acid residues include, but are not limited to, Gly, Allylglycine, Cyclohexylglycine (Chg), Dpg, Hydroxyproline, HoLeu, alloIle, Ala, Abu (Aminobutyric acid), Homocycloleucine, Acpc (1-Aminocyclopropane-1-carboxylic acid), Aib, Aic, Val, Leu, Nle, Ile, Met, Phe, Cys, Pro, Gln, Asn, Cha (beta-cyclohexylalanine), Cyclopentylalanine, beta-Cyclopropyl alanine, 3,3-Diphenylalanine, beta-2-Furylalanine, Homocyclohexylalanine (HoCha) 3-(1-Naphthylalanine, 2-Furylalanine, Pyridylalanine, Quinolylalanine, Thiazolylalanine, Theinylalanine, Val, Nva, Ring substituted phenylalanine, Sar, HoSer, Tic, Tic(OH), Ring substituted Tryptophan, Ring substituted Tyrosine, and derivatives thereof.

In general, the term “aliphatic amino acids” and “non-polar aliphatic amino acids” means amino acids with a non-aromatic hydrophobic side chain. Examples include but not limited to Ala, AllylGly, Nva, Chg, Abu, Aib, Aic, Acpc, Homocycloleucine, Cyclopentylalanine, beta-Cyclopropyl alanine, Cha, HoCha, Val, Ile, Leu, and Met and derivatives thereof.

In general, the term “Aromatic amino acids” means amino acids with an aromatic ring side chain. Examples include Phe, Tyr, Trp, His, 2-Furylalanine, Tyr(Me) and derivatives thereof.

In general, the term “Polar amino acids” includes polar and uncharged amino acids, negatively charged amino acids and positively charged amino acids. Examples include Arg, HoArg, Cit, Glu, Asp, Lys, Gln, Asn, Ser, Thr, His, Trp, Tyr, Lys(Ac), and derivatives thereof.

In general, the term “Positively charged amino acids” (or basic amino acids) means amino acids whose side chains are protonated or may be protonated at physiological conditions. Examples include but not are limited to Lys, Ornathine (Orn), Arg, HoArg, Dab, Dap, Trp, His and derivatives thereof.

In general, the term “Negatively charged amino acids” (or acidic amino acids) means amino acids whose side chains may be deprotonated at physiological conditions. Examples include but not limited to Asp, Asu, Glu, Aad, and derivatives thereof.

In general, “Polar, uncharged amino acids” means amino acids with uncharged side chains capable of forming H-bonds with water at physiological conditions. Examples include but are not limited to Gln, Asn, Lys(Ac), Ser, Thr, Cit, and HoCit derivatives thereof.

“Substantially homologous” means at least about 75% (preferably at least about 80%, and more preferably at least about 90% or most preferably at least about 95%, of the amino-acid residues match over the defined length of the peptide sequences. Sequences that are substantially homologous can be identified by comparing the sequences using standard software available in sequence data banks, such as BLAST programs available from the National Cancer Center for Biotechnology Information at ncbi.nlm.nih.gov.

In general, “substituted” refers to a group as defined below in which one or more bonds to a hydrogen atom contained therein are replaced by a bond to non-hydrogen or non-carbon atoms such as, but not limited to, a halogen atom such as F, Cl, Br, and I; an oxygen atom in groups such as hydroxyl groups, alkoxy groups, aryloxy groups, and ester groups; a sulfur atom in groups such as thiol groups, alkyl and aryl sulfide groups, sulfone groups, sulfonyl groups, and sulfoxide groups; a nitrogen atom

in groups such as amines, amides, alkylamines, dialkylamines, arylamines, alkylarylamines, diarylamines, N-oxides, imides, and enamines; a silicon atom in groups such as in trialkylsilyl groups, dialkylarylsilyl groups, alkyldiarylsilyl groups, and triarylsilyl groups; and other heteroatoms in various other groups. Substituted alkyl groups and also substituted cycloalkyl groups and others also include groups in which one or more bonds to a carbon(s) or hydrogen(s) atom is replaced by a bond to a heteroatom such as oxygen in carbonyl, carboxyl, and ester groups; nitrogen in groups such as imines, oximes, hydrazones, and nitriles. As employed herein, a group which is "optionally substituted" may be substituted or unsubstituted. Thus, e.g., "optionally substituted alkyl" refers to both substituted alkyl groups and unsubstituted alkyl groups.

The phrase "unsubstituted alkyl" refers to alkyl groups that do not contain heteroatoms. Thus, the phrase includes straight chain alkyl groups such as methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl and the like. The phrase also includes branched chain isomers of straight chain alkyl groups, including but not limited to, the following which are provided by way of example: $-\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}(\text{CH}_2\text{CH}_3)_2$, $-\text{C}(\text{CH}_3)_3$, $-\text{C}(\text{CH}_2\text{CH}_3)_3$, $-\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)_2$, $-\text{CH}_2\text{C}(\text{CH}_3)_3$, $-\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)_3$, $-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)_2$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_3$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)_3$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, and others. The phrase does not include cycloalkyl groups. Thus, the phrase unsubstituted alkyl groups includes primary alkyl groups, secondary alkyl groups, and tertiary alkyl groups. Unsubstituted alkyl groups may be bonded to one or more carbon atom(s), oxygen atom(s), nitrogen atom(s), and/or sulfur atom(s) in the parent compound. Possible unsubstituted alkyl groups include straight and branched chain alkyl groups having 1 to 20 carbon atoms. Alternatively, such unsubstituted alkyl groups have from 1 to 10 carbon atoms or are lower alkyl groups having from 1 to about 6 carbon atoms. Other unsubstituted alkyl groups include straight and branched chain alkyl groups having from 1 to 3 carbon atoms and include methyl, ethyl, propyl, and $-\text{CH}(\text{CH}_3)_2$.

The phrase "substituted alkyl" refers to an alkyl group in which one or more bonds to a carbon(s) or hydrogen(s) are replaced by a bond to non-hydrogen and non-carbon atoms such as, but not limited to, a halogen atom in halides such as F, Cl, Br, and I; an oxygen atom in groups such as hydroxyl groups, alkoxy groups, aryloxy groups, and ester groups; a sulfur atom in groups such as thiol groups, alkyl and aryl sulfide groups, sulfone groups, sulfonyl groups, and sulfoxide groups; a nitrogen atom in groups such as amines, amides, alkylamines, dialkylamines, arylamines, alkylarylamines, diarylamines, N-oxides, imides, and enamines; a silicon atom in groups such as in trialkylsilyl groups, dialkylarylsilyl groups, alkyldiarylsilyl groups, and triarylsilyl groups; and other heteroatoms in various other groups. Substituted alkyl groups also include groups in which one or more bonds to a

carbon(s) or hydrogen(s) atom is replaced by a bond to a heteroatom such as oxygen in carbonyl, carboxyl, and ester groups; nitrogen in groups such as imines, oximes, hydrazones, and nitriles. Substituted alkyl groups include, among others, alkyl groups in which one or more bonds to a carbon or hydrogen atom is/are replaced by one or more bonds to fluorine atoms. One example of a substituted alkyl group is the trifluoromethyl group and other alkyl groups that contain the trifluoromethyl group. Other alkyl groups include those in which one or more bonds to a carbon or hydrogen atom is replaced by a bond to an oxygen atom such that the substituted alkyl group contains a hydroxyl, alkoxy, aryloxy group, or heterocycloxy group. Still other alkyl groups include alkyl groups that have an amine, alkylamine, dialkylamine, arylamine, (alkyl)(aryl)amine, diarylamine, heterocyclamine, (alkyl)(heterocyclamine), (aryl)(heterocyclamine), or diheterocyclamine group.

The phrase “unsubstituted alkylene” refers to a divalent unsubstituted alkyl group as defined above. Thus methylene, ethylene, and propylene are each examples of unsubstituted alkylenes. The phrase “substituted alkylene” refers to a divalent substituted alkyl group as defined above. Substituted or unsubstituted lower alkylene groups have from 1 to about 6 carbons.

The phrase “unsubstituted cycloalkyl” refers to cyclic alkyl groups such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl and such rings substituted with straight and branched chain alkyl groups as defined above. The phrase also includes polycyclic alkyl groups such as, but not limited to, adamantyl norbornyl, and bicyclo[2.2.2]octyl and the like, as well as such rings substituted with straight and branched chain alkyl groups as defined above. Thus, the phrase would include methylcyclohexyl groups among others. The phrase does not include cyclic alkyl groups containing heteroatoms. Unsubstituted cycloalkyl groups may be bonded to one or more carbon atom(s), oxygen atom(s), nitrogen atom(s), and/or sulfur atom(s) in the parent compound. In some embodiments unsubstituted cycloalkyl groups have from 3 to 20 carbon atoms. In other embodiments, such unsubstituted alkyl groups have from 3 to 8 carbon atoms while in others, such groups have from 3 to 7 carbon atoms.

The phrase “substituted cycloalkyl” has the same meaning with respect to unsubstituted cycloalkyl groups that substituted alkyl groups have with respect to unsubstituted alkyl groups. Thus, the phrase includes, but is not limited to, oxocyclohexyl, chlorocyclohexyl, hydroxycyclopentyl, and chloromethylcyclohexyl groups.

The phrase “unsubstituted aryl” refers to aryl groups that do not contain heteroatoms. Thus the phrase includes, but is not limited to, groups such as phenyl, biphenyl, anthracenyl, and naphthenyl by way of example. Although the phrase “unsubstituted aryl” includes groups containing condensed rings

such as naphthalene, it does not include aryl groups that have other groups such as alkyl or halo groups bonded to one of the ring members, as aryl groups such as tolyl are considered herein to be substituted aryl groups as described below. Typically, an unsubstituted aryl may be a lower aryl, having from 6 to about 10 carbon atoms. One unsubstituted aryl group is phenyl. Unsubstituted aryl groups may be bonded to one or more carbon atom(s), oxygen atom(s), nitrogen atom(s), and/or sulfur atom(s) in the parent compound, however.

The phrase “substituted aryl group” has the same meaning with respect to unsubstituted aryl groups that substituted alkyl groups have with respect to unsubstituted alkyl groups. However, a substituted aryl group also includes aryl groups in which one of the aromatic carbons is bonded to one of the non-carbon or non-hydrogen atoms described above and also includes aryl groups in which one or more aromatic carbons of the aryl group is bonded to a substituted and/or unsubstituted alkyl, alkenyl, or alkynyl group as defined herein. This includes bonding arrangements in which two carbon atoms of an aryl group are bonded to two atoms of an alkyl, alkenyl, or alkynyl group to define a fused ring system (e.g., dihydronaphthyl or tetrahydronaphthyl). Thus, the phrase “substituted aryl” includes, but is not limited to tolyl and hydroxyphenyl among others.

The phrase “unsubstituted alkenyl” refers to straight and branched chain and cyclic groups such as those described with respect to unsubstituted alkyl groups as defined above, except that at least one double bond exists between two carbon atoms. Examples include, but are not limited to vinyl, -CH=C(H)(CH₃), -CH=C(CH₃)₂, -C(CH₃)=C(H)₂, -C(CH₃)=C(H)(CH₃), -C(CH₂CH₃)=CH₂, cyclohexenyl, cyclopentenyl, cyclohexadienyl, butadienyl, pentadienyl, and hexadienyl among others. Lower unsubstituted alkenyl groups have from 1 to about 6 carbons.

The phrase “substituted alkenyl” has the same meaning with respect to unsubstituted alkenyl groups that substituted alkyl groups have with respect to unsubstituted alkyl groups. A substituted alkenyl group includes alkenyl groups in which a non-carbon or non-hydrogen atom is bonded to a carbon double bonded to another carbon and those in which one of the non-carbon or non-hydrogen atoms is bonded to a carbon not involved in a double bond to another carbon. For example, -CH=CH-OCH₃ and -CH=CH-CH₂-OH are both substituted alkenyls. Oxoalkenyls wherein a CH₂ group is replaced by a carbonyl, such as -CH=CH-C(O)-CH₃, are also substituted alkenyls.

The phrase “unsubstituted alkenylene” refers to a divalent unsubstituted alkenyl group as defined above. For example, -CH=CH- is an exemplary unsubstituted alkenylene. The phrase “substituted alkenylene” refers to a divalent substituted alkenyl group as defined above.

The phrase “unsubstituted alkynyl” refers to straight and branched chain groups such as those described with respect to unsubstituted alkyl groups as defined above, except that at least one triple

bond exists between two carbon atoms. Examples include, but are not limited to, $-\text{C}\equiv\text{C}(\text{H})$, $-\text{C}\equiv\text{C}(\text{CH}_3)$, $-\text{C}\equiv\text{C}(\text{CH}_2\text{CH}_3)$, $-\text{C}(\text{H}_2)\text{C}\equiv\text{C}(\text{H})$, $-\text{C}(\text{H})_2\text{C}\equiv\text{C}(\text{CH}_3)$, and $-\text{C}(\text{H})_2\text{C}\equiv\text{C}(\text{CH}_2\text{CH}_3)$ among others. Unsubstituted lower alkynyl groups have from 1 to about 6 carbons.

- 5 The phrase “substituted alkynyl” has the same meaning with respect to unsubstituted alkynyl groups that substituted alkyl groups have with respect to unsubstituted alkyl groups. A substituted alkynyl group includes alkynyl groups in which a non-carbon or non-hydrogen atom is bonded to a carbon triple bonded to another carbon and those in which a non-carbon or non-hydrogen atom is bonded to a carbon not involved in a triple bond to another carbon. Examples include, but are not limited to,
- 10 oxoalkynyls wherein a CH_2 group is replaced by a carbonyl, such as $-\text{C}(\text{O})-\text{CH}\equiv\text{CH}-\text{CH}_3$ and $-\text{C}(\text{O})-\text{CH}_2-\text{CH}\equiv\text{CH}$.

- The phrase “unsubstituted alkynylene” refers to a divalent unsubstituted alkynyl group as defined above. A $-\text{C}\equiv\text{C}-$ is an example of an unsubstituted alkynylene. The phrase “substituted alkynylene”
- 15 refers to a divalent substituted alkynyl group as defined above.

- The phrase “unsubstituted aralkyl” refers to unsubstituted alkyl groups as defined above in which a hydrogen or carbon bond of the unsubstituted alkyl group is replaced with a bond to an aryl group as defined above. For example, methyl ($-\text{CH}_3$) is an unsubstituted alkyl group. If a hydrogen atom of
- 20 the methyl group is replaced by a bond to a phenyl group, such as if the carbon of the methyl were bonded to a carbon of benzene, then the compound is an unsubstituted aralkyl group (*i.e.*, a benzyl group). Thus, the phrase includes, but is not limited to, groups such as benzyl, diphenylmethyl, and 1-phenylethyl ($-\text{CH}(\text{C}_6\text{H}_5)(\text{CH}_3)$).

- 25 The phrase “substituted aralkyl” has the same meaning with respect to unsubstituted aralkyl groups that substituted aryl groups have with respect to unsubstituted aryl groups. However, substituted aralkyls also include groups in which a carbon or hydrogen bond of the alkyl part of the group is replaced by a bond to a non-carbon or a non-hydrogen atom. Examples of substituted aralkyl groups include, but are not limited to, $-\text{CH}_2\text{C}(=\text{O})(\text{C}_6\text{H}_5)$, and $-\text{CH}_2(2\text{-methylphenyl})$.

- 30 The phrase “unsubstituted aralkenyl” refers to unsubstituted alkenyl groups as defined above in which a hydrogen or carbon bond of the unsubstituted alkenyl group is replaced with a bond to an aryl group as defined above. For example, vinyl is an unsubstituted alkenyl group. If a hydrogen atom of the vinyl group is replaced by a bond to a phenyl group, such as if a carbon of the vinyl were bonded to a
- 35 carbon of benzene, then the compound is an unsubstituted aralkenyl group (*i.e.*, a styryl group). Thus,

the phrase includes, but is not limited to, groups such as styryl, diphenylvinyl, and 1-phenylethenyl ($-\text{C}(\text{C}_6\text{H}_5)(\text{CH}_2)-$).

The phrase “substituted aralkenyl” has the same meaning with respect to unsubstituted aralkenyl groups that substituted aryl groups have with respect to unsubstituted aryl groups. A substituted aralkenyl group also includes groups in which a carbon or hydrogen bond of the alkenyl part of the group is replaced by a bond to a non-carbon or a non-hydrogen atom. Examples of substituted aralkenyl groups include, but are not limited to, $-\text{CH}=\text{C}(\text{Cl})(\text{C}_6\text{H}_5)-$, and $-\text{CH}=\text{CH}(2\text{-methylphenyl})-$.

The phrase “unsubstituted aralkynyl” refers to unsubstituted alkynyl groups as defined above in which a hydrogen or carbon bond of the unsubstituted alkynyl group is replaced with a bond to an aryl group as defined above. For example, acetylene is an unsubstituted alkynyl group. If a hydrogen atom of the acetylene group is replaced by a bond to a phenyl group, such as if a carbon of the acetylene were bonded to a carbon of benzene, then the compound is an unsubstituted aralkynyl group. Thus, the phrase includes, but is not limited to, groups such as $-\text{C}\equiv\text{C}\text{-phenyl}$ and $-\text{CH}_2\text{-C}\equiv\text{C}\text{-phenyl}$.

The phrase “substituted aralkynyl” has the same meaning with respect to unsubstituted aralkynyl groups that substituted aryl groups have with respect to unsubstituted aryl groups. However, a substituted aralkynyl group also includes groups in which a carbon or hydrogen bond of the alkynyl part of the group is replaced by a bond to a non-carbon or a non-hydrogen atom. Examples of substituted aralkynyl groups include, but are not limited to, $-\text{C}\equiv\text{C}\text{-C}(\text{Br})(\text{C}_6\text{H}_5)-$ and $-\text{C}\equiv\text{C}(2\text{-methylphenyl})-$.

The phrase “unsubstituted heteroalkyl” refers to unsubstituted alkyl groups as defined above in which the carbon chain is interrupted by one or more heteroatoms chosen from N, O, and S. Unsubstituted heteroalkyls containing N may have NH or N(unsubstituted alkyl) in the carbon chain. For example, unsubstituted heteroalkyls include alkoxy, alkoxyalkyl, alkoxyalkoxy, thioether, alkylaminoalkyl, aminoalkoxy, and other such groups. Typically, unsubstituted heteroalkyl groups contain 1-5 heteroatoms, and particularly 1-3 heteroatoms. In some embodiments unsubstituted heteroalkyls include, for example, alkoxyalkoxyalkoxy groups such as ethyloxyethyloxyethyloxy.

The phrase “substituted heteroalkyl” has the same meaning with respect to unsubstituted heteroalkyl groups that substituted alkyl groups have with respect to unsubstituted alkyl groups.

The phrase “unsubstituted heteroalkylene” refers to a divalent unsubstituted heteroalkyl group as defined above. For example, $-\text{CH}_2\text{-O-CH}_2-$ and $-\text{CH}_2\text{-NH-CH}_2\text{CH}_2-$ are both exemplary unsubstituted heteroalkylenes. The phrase “substituted heteroalkylene” refers to a divalent substituted heteroalkyl group as defined above.

The phrase “unsubstituted heteroalkenyl” refers to unsubstituted alkene groups as defined above in which the carbon chain is interrupted by one or more heteroatoms chosen from N, O, and S.

Unsubstituted heteroalkenyls containing N may have NH or N(unsubstituted alkyl or alkene) in the carbon chain. The phrase “substituted heteroalkenyl” has the same meaning with respect to

5 unsubstituted heteroalkenyl groups that substituted heteroalkyl groups have with respect to unsubstituted heteroalkyl groups.

The phrase “unsubstituted heteroalkenylene” refers to a divalent unsubstituted heteroalkenyl group as defined above. Thus $-\text{CH}_2\text{-O-CH=CH-}$ is an example of an unsubstituted heteroalkenylene. The

10 phrase “substituted heteroalkenylene” refers to a divalent substituted heteroalkenyl group as defined above.

The phrase “unsubstituted heteroalkynyl” refers to unsubstituted alkynyl groups as defined above in which the carbon chain is interrupted by one or more heteroatoms chosen from N, O, and S.

15 Unsubstituted heteroalkynyls containing N may have NH or N(unsubstituted alkyl, alkene, or alkyne) in the carbon chain. The phrase “substituted heteroalkynyl” has the same meaning with respect to unsubstituted heteroalkynyl groups that substituted heteroalkyl groups have with respect to unsubstituted heteroalkyl groups.

20 The phrase “unsubstituted heteroalkynylene” refers to a divalent unsubstituted heteroalkynyl group as defined above. Thus $-\text{CH}_2\text{-O-CH}_2\text{-C}\equiv\text{C-}$ is an example of an unsubstituted heteroalkynylene. The phrase “substituted heteroalkynylene” refers to a divalent substituted heteroalkynyl group as defined above.

25 The phrase “unsubstituted heterocyclyl” refers to both aromatic and nonaromatic ring compounds including monocyclic, bicyclic, and polycyclic ring compounds such as, but not limited to, quinuclidyl, containing 3 or more ring members of which one or more is a heteroatom such as, but not limited to, N, O, and S. Although the phrase “unsubstituted heterocyclyl” includes condensed heterocyclic rings such as benzimidazolyl, it does not include heterocyclyl groups that have other

30 groups such as alkyl or halo groups bonded to one of the ring members as compounds such as 2-methylbenzimidazolyl are substituted heterocyclyl groups. Examples of heterocyclyl groups include, but are not limited to: unsaturated 3 to 8 membered rings containing 1 to 4 nitrogen atoms such as, but not limited to pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, dihydropyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazolyl (e.g., 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl etc.),

35 tetrazolyl, (e.g., 1H-tetrazolyl, 2H tetrazolyl, etc.); saturated 3 to 8 membered rings containing 1 to 4 nitrogen atoms such as, but not limited to, pyrrolidinyl, imidazolidinyl, piperidinyl, piperazinyl; condensed unsaturated heterocyclic groups containing 1 to 4 nitrogen atoms such as, but not limited

to, indolyl, isoindolyl, indolynyl, indolizynyl, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl; unsaturated 3 to 8 membered rings containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms such as, but not limited to, oxazolyl, isoxazolyl, oxadiazolyl (e.g., 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl, etc.); saturated 3 to 8 membered rings containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms such as, but not limited to, morpholynyl; unsaturated condensed heterocyclic groups containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, benzoxazolyl, benzoxadiazolyl, benzoxazinyl (e.g., 2H-1,4-benzoxazinyl, etc.); unsaturated 3 to 8 membered rings containing 1 to 3 sulfur atoms and 1 to 3 nitrogen atoms such as, but not limited to, thiazolyl, isothiazolyl, thiadiazolyl (e.g., 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, etc.); saturated 3 to 8 membered rings containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms such as, but not limited to, thiazolodynyl; saturated and unsaturated 3 to 8 membered rings containing 1 to 2 sulfur atoms such as, but not limited to, thienyl, dihydrodithiynyl, dihydrodithionyl, tetrahydrothiophene, tetrahydrothiopyran; unsaturated condensed heterocyclic rings containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms such as, but not limited to, benzothiazolyl, benzothiadiazolyl, benzothiazinyl (e.g., 2H-1,4-benzothiazinyl, etc.), dihydrobenzothiazinyl (e.g., 2H-3,4-dihydrobenzothiazinyl, etc.), unsaturated 3 to 8 membered rings containing oxygen atoms such as, but not limited to furyl; unsaturated condensed heterocyclic rings containing 1 to 2 oxygen atoms such as benzodioxolyl (e.g., 1,3-benzodioxolyl, etc.); unsaturated 3 to 8 membered rings containing an oxygen atom and 1 to 2 sulfur atoms such as, but not limited to, dihydrooxathiynyl; saturated 3 to 8 membered rings containing 1 to 3 oxygen atoms and 1 to 2 sulfur atoms such as 1,4-oxathiane; unsaturated condensed rings containing 1 to 2 sulfur atoms such as benzothieryl, benzodithieryl; and unsaturated condensed heterocyclic rings containing an oxygen atom and 1 to 3 oxygen atoms such as benzoxathieryl. Heterocyclyl group also include those described above in which one or more S atoms in the ring is double-bonded to one or two oxygen atoms (sulfoxides and sulfones). For example, heterocyclyl groups include tetrahydrothiophene, tetrahydrothiophene oxide, and tetrahydrothiophene 1,1-dioxide. In some embodiments heterocyclyl groups contain 5 or 6 ring members. In other embodiments heterocyclyl groups include morpholine, piperazine, piperidine, pyrrolidine, imidazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, thiomorpholine, thiomorpholine in which the S atom of the thiomorpholine is bonded to one or more O atoms, pyrrole, homopiperazine, oxazolidin-2-one, pyrrolidin-2-one, oxazole, quinuclidine, thiazole, isoxazole, furan, and tetrahydrofuran.

The phrase “substituted heterocyclyl” refers to an unsubstituted heterocyclyl group as defined above in which one of the ring members is bonded to a non-hydrogen atom such as described above with respect to substituted alkyl groups and substituted aryl groups. Examples include, but are not limited to, 2-methylbenzimidazolyl, 5-methylbenzimidazolyl, 5-chlorobenzthiazolyl, 1-methyl piperazinyl, and 2-chloropyridyl.

The phrase “unsubstituted heteroaryl” refers to unsubstituted aromatic heterocyclyl groups as defined above. Thus, unsubstituted heteroaryl groups include but are not limited to furyl, imidazolyl, oxazolyl, isoxazolyl, pyridinyl, benzimidazolyl, and benzothiazolyl. The phrase “substituted heteroaryl” refers to substituted aromatic heterocyclyl groups as defined above.

- 5 The phrase “unsubstituted heterocyclylalkyl” refers to unsubstituted alkyl groups as defined above in which a hydrogen or carbon bond of the unsubstituted alkyl group is replaced with a bond to a heterocyclyl group as defined above. For example, methyl ($-\text{CH}_3$) is an unsubstituted alkyl group. If a hydrogen atom of the methyl group is replaced by a bond to a heterocyclyl group, such as if the carbon of the methyl were bonded to carbon 2 of pyridine (one of the carbons bonded to the N of the
- 10 pyridine) or carbons 3 or 4 of the pyridine, then the compound is an unsubstituted heterocyclylalkyl group.

- The phrase “substituted heterocyclylalkyl” has the same meaning with respect to unsubstituted heterocyclylalkyl groups that substituted aralkyl groups have with respect to unsubstituted aralkyl
- 15 groups. A substituted heterocyclylalkyl group also includes groups in which a non-hydrogen atom is bonded to a heteroatom in the heterocyclyl group of the heterocyclylalkyl group such as, but not limited to, a nitrogen atom in the piperidine ring of a piperidinylalkyl group.

- The phrase “unsubstituted heterocyclylalkenyl” refers to unsubstituted alkenyl groups as defined above in which a hydrogen or carbon bond of the unsubstituted alkenyl group is replaced with a bond
- 20 to a heterocyclyl group as defined above. For example, vinyl is an unsubstituted alkenyl group. If a hydrogen atom of the vinyl group is replaced by a bond to a heterocyclyl group, such as if the carbon of the vinyl were bonded to carbon 2 of pyridine or carbons 3 or 4 of the pyridine, then the compound is an unsubstituted heterocyclylalkenyl group.

- 25 The phrase “substituted heterocyclylalkenyl” has the same meaning with respect to unsubstituted heterocyclylalkenyl groups that substituted aralkenyl groups have with respect to unsubstituted aralkenyl groups. However, a substituted heterocyclylalkenyl group also includes groups in which a non-hydrogen atom is bonded to a heteroatom in the heterocyclyl group of the heterocyclylalkenyl group such as, but not limited to, a nitrogen atom in the piperidine ring of a piperidinylalkenyl group.

- 30 The phrase “unsubstituted heterocyclylalkynyl” refers to unsubstituted alkynyl groups as defined above in which a hydrogen or carbon bond of the unsubstituted alkynyl group is replaced with a bond to a heterocyclyl group as defined above. For example, acetylene is an unsubstituted alkynyl group. If a hydrogen atom of the acetylene group is replaced by a bond to a heterocyclyl group, such as if the carbon of the acetylene were bonded to carbon 2 of pyridine or carbons 3 or 4 of the pyridine, then the
- 35 compound is an unsubstituted heterocyclylalkynyl group.

The phrase "substituted heterocyclalkynyl" has the same meaning with respect to unsubstituted heterocyclalkynyl groups that substituted aralkynyl groups have with respect to unsubstituted aralkynyl groups. A substituted heterocyclalkynyl group also includes groups in which a non-hydrogen atom is bonded to a heteroatom in the heterocycl group of the heterocyclalkynyl group such as, but not limited to, a nitrogen atom in the piperidine ring of a piperidinylalkynyl group.

The phrase "unsubstituted alkoxy" refers to a hydroxyl group (-OH) in which the bond to the hydrogen atom is replaced by a bond to a carbon atom of an otherwise unsubstituted alkyl group as defined above.

The phrase "substituted alkoxy" refers to a hydroxyl group (-OH) in which the bond to the hydrogen atom is replaced by a bond to a carbon atom of an otherwise substituted alkyl group as defined above. A "pharmaceutically acceptable salt" includes a salt with an inorganic base, organic base, inorganic acid, organic acid, or basic or acidic amino acid. Salts of inorganic bases include, for example, alkali metals such as sodium or potassium; alkaline earth metals such as calcium and magnesium or aluminum; and ammonia. Salts of organic bases include, for example, trimethylamine, triethylamine, pyridine, picoline, ethanolamine, diethanolamine, and triethanolamine. Salts of inorganic acids include for example, hydrochloric acid, hydroboric acid, nitric acid, sulfuric acid, and phosphoric acid. Salts of organic acids include for example, formic acid, acetic acid, trifluoroacetic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, malic acid, methanesulfonic acid, benzenesulfonic acid, and p-toluenesulfonic acid. Salts of basic amino acids include, for example, arginine, lysine and ornithine. Acidic amino acids include, for example, aspartic acid and glutamic acid.

"Tautomers" refers to isomeric forms of a compound that are in equilibrium with each other. The concentrations of the isomeric forms will depend on the environment the compound is found in and may be different depending upon, for example, whether the compound is a solid or is in an organic or aqueous solution. For example, in aqueous solution, ketones are typically in equilibrium with their enol forms. Thus, ketones and their enols are referred to as tautomers of each other. As readily understood by one skilled in the art, a wide variety of functional groups and other structures may exhibit tautomerism, and all tautomers of compounds the invention are within the scope of the present invention.

Certain embodiments are derivatives referred to as prodrugs. The expression "prodrug" denotes a derivative of a pharmaceutically or therapeutically active drug, e.g., esters and amides, wherein the derivative has an enhanced characteristic such as, for example, enhanced delivery and therapeutic value as compared to the drug and can be transformed into the drug by an enzymatic or chemical process. See, for example, R.E. Notari, *Methods Enzymol.* 112:309-323 (1985); N. Bodor, *Drugs of*

the Future 6:165-182 (1981); H. Bundgaard, Chapter 1 in *Design of Prodrugs* (H. Bundgaard, ed.), Elsevier, New York (1985); and A. G. Gilman *et al.*, *Goodman And Gilman's The Pharmacological Basis of Therapeutics*, 8th ed., McGraw-Hill (1990). Thus, the prodrug may be designed to alter the metabolic stability or transport characteristics of a drug, mask side effects or toxicity of a drug, improve the flavor of a drug, or to alter other characteristics or properties of a drug. Compounds of the present invention include enriched or resolved optical isomers at any or all asymmetric atoms as are apparent from the depictions. Both racemic and diastereomeric mixtures, as well as the individual optical isomers can be isolated or synthesized so as to be substantially free of their enantiomeric or diastereomeric partners. All such stereoisomers are within the scope of the invention.

The term "carboxy protecting group" as used herein refers to a carboxylic acid protecting ester group employed to block or protect the carboxylic acid functionality while the reactions involving other functional sites of the compound are carried out. Carboxy protecting groups are disclosed in, for example, Greene, *Protective Groups in Organic Synthesis*, pp. 152-186, John Wiley & Sons, New York (1981), which is hereby incorporated herein by reference. In addition, a carboxy protecting group can be used as a prodrug, whereby the carboxy protecting group can be readily cleaved *in vivo* by, for example, enzymatic hydrolysis to release the biologically active parent. T. Higuchi and V. Stella provide a discussion of the prodrug concept in "Pro-drugs as Novel Delivery Systems", Vol. 14 of the A.C.S. Symposium Series, American Chemical Society (1975), which is hereby incorporated herein by reference. Such carboxy protecting groups are well known to those skilled in the art, having been extensively used in the protection of carboxyl groups in the penicillin and cephalosporin fields, as described in U.S. Patent Nos. 3,840,556 and 3,719,667, S. Kukolja, J. Am. Chem. Soc. 93:6267-6269 (1971), and G.E. Gutowski, Tetrahedron Lett. 21:1779-1782 (1970), the disclosures of which are hereby incorporated herein by reference. Examples of esters useful as prodrugs for compounds containing carboxyl groups can be found, for example, at pp. 14-21 in *Bioreversible Carriers in Drug Design: Theory and Application* (E.B. Roche, ed.), Pergamon Press, New York (1987), which is hereby incorporated herein by reference. Representative carboxy protecting groups are C₁ to C₈ alkyl (e.g., methyl, ethyl or tertiary butyl and the like); haloalkyl; alkenyl; cycloalkyl and substituted derivatives thereof such as cyclohexyl, cyclopentyl and the like; cycloalkylalkyl and substituted derivatives thereof such as cyclohexylmethyl, cyclopentylmethyl and the like; arylalkyl, for example, phenethyl or benzyl and substituted derivatives thereof such as alkoxybenzyl or nitrobenzyl groups and the like; arylalkenyl, for example, phenylethenyl and the like; aryl and substituted derivatives thereof, for example, 5-indanyl and the like; dialkylaminoalkyl (e.g., dimethylaminoethyl and the like); alkanoyloxyalkyl groups such as acetoxymethyl, butyryloxymethyl, valerytoxymethyl, isobutyryloxymethyl, isovaleryloxymethyl, 1-(propionyloxy)-1-ethyl, 1-(pivaloyloxy)-1-ethyl, 1-methyl-1-(propionyloxy)-1-ethyl, pivaloyloxymethyl, propionyloxymethyl and the like; cycloalkanoyloxyalkyl groups such as cyclopropylcarbonyloxymethyl, cyclobutylcarbonyloxymethyl,

cyclopentylcarbonyloxymethyl, cyclohexylcarbonyloxymethyl and the like; aroyloxyalkyl, such as benzoyloxymethyl, benzoyloxyethyl and the like; arylalkylcarbonyloxyalkyl, such as benzylcarbonyloxymethyl, 2-benzylcarbonyloxyethyl and the like; alkoxycarbonylalkyl, such as methoxycarbonylmethyl, cyclohexyloxycarbonylmethyl, 1-methoxycarbonyl-1-ethyl, and the like; 5 alkoxycarbonyloxyalkyl, such as methoxycarbonyloxymethyl, t-butyloxycarbonyloxymethyl, 1-ethoxycarbonyloxy-1-ethyl, 1-cyclohexyloxycarbonyloxy-1-ethyl and the like; alkoxycarbonylaminoalkyl, such as t-butyloxycarbonylaminomethyl and the like; alkylaminocarbonylaminoalkyl, such as methylaminocarbonylaminomethyl and the like; alkanoylaminoalkyl, such as acetylaminomethyl and the like; heterocycliccarbonyloxyalkyl, such as 10 4-methylpiperazinylcarbonyloxymethyl and the like; dialkylaminocarbonylalkyl, such as dimethylaminocarbonylmethyl, diethylaminocarbonylmethyl and the like; (5-(alkyl)-2-oxo-1,3-dioxolen-4-yl)alkyl, such as (5-t-butyl-2-oxo-1,3-dioxolen-4-yl)methyl and the like; and (5-phenyl-2-oxo-1,3-dioxolen-4-yl)alkyl, such as (5-phenyl-2-oxo-1,3-dioxolen-4-yl)methyl and the like.

The term "N-protecting group" or "N-protected" as used herein refers to those groups intended to 15 protect the N-terminus of an amino acid or peptide or to protect an amino group against undesirable reactions during synthetic procedures. Commonly used N-protecting groups are disclosed in, for example, Greene, *Protective Groups in Organic Synthesis*, John Wiley & Sons, New York (1981), which is hereby incorporated by reference. For example, N-protecting groups can comprise acyl groups such as formyl, acetyl, propionyl, pivaloyl, t-butylacetyl, 2-chloroacetyl, 2-bromoacetyl, 20 trifluoroacetyl, trichloroacetyl, phthalyl, o-nitrophenoxyacetyl, a-chlorobutyryl, benzoyl, 4-chlorobenzoyl, 4-bromobenzoyl, 4-nitrobenzoyl, and the like; sulfonyl groups such as benzenesulfonyl, p-toluenesulfonyl and the like; carbamate forming groups such as benzyloxycarbonyl, p-chlorobenzyloxycarbonyl, p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, 2-nitrobenzyloxycarbonyl, p-bromobenzyloxycarbonyl, 3,4- 25 dimethoxybenzyloxycarbonyl, 3,5-dimethoxybenzyloxycarbonyl, 2,4-dimethoxybenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 2-nitro-4,5-dimethoxybenzyloxycarbonyl, 3,4,5-trimethoxybenzyloxycarbonyl, 1-(p-biphenyl)-1-methylethoxycarbonyl, α,α -dimethyl-3,5-dimethoxybenzyloxycarbonyl, benzhydryloxycarbonyl, t-butyloxycarbonyl, diisopropylmethoxycarbonyl, isopropylloxycarbonyl, ethoxycarbonyl, methoxycarbonyl, 30 allyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, phenoxycarbonyl, 4-nitrophenoxycarbonyl, fluorenyl-9-methoxycarbonyl, cyclopentylloxycarbonyl, adamantylloxycarbonyl, cyclohexylloxycarbonyl, phenylthiocarbonyl and the like; alkyl groups such as benzyl, triphenylmethyl, benzyloxymethyl and the like; and silyl groups such as trimethylsilyl and the like. In some embodiments N-protecting groups are formyl, acetyl, benzoyl, pivaloyl, t-butylacetyl, 35 phenylsulfonyl, benzyl, 9-fluorenylmethyloxycarbonyl (Fmoc), t-butyloxycarbonyl (Boc), and benzyloxycarbonyl (Cbz).

As used herein, "halo," "halogen" or "halide" refers to F, Cl, Br or I.

As used herein, the abbreviations for any protective groups, amino acids or other compounds, are, unless indicated otherwise, in accord with their common usage, recognized abbreviations, or the IUPAC-IUB Commission on Biochemical Nomenclature, Biochem. 11:942-944 (1972).

As used herein, "substantially pure" means sufficiently homogeneous to appear free of readily detectable impurities as determined by standard methods of analysis, such as thin layer chromatography (TLC), gel electrophoresis, and high performance liquid chromatography (HPLC), used by those of skill in the art to assess such purity, or sufficiently pure such that further purification would not detectably alter the physical and chemical properties, such as enzymatic and biological activities, of the substance. Substantially pure includes compositions in which the compound of the invention forms the major component of the composition, such as constituting about 50%, about 60%, about 70%, about 80%, about 90%, or about 95% or more of the substances in the composition.

Methods for purification of compounds to produce substantially chemically pure compounds are known to those of skill in the art. A substantially chemically pure compound may, however, be a mixture of stereoisomers. In such instances, further purification may increase the specific activity of the compound. However, a compound of the invention need not always be provided in a specific purified state. Partially purified compositions will have utility in certain embodiments and depending on the desired use. For example, purification methods that may yield a greater total recovery of a compound of the invention may produce a lower degree of relative purification.

As used herein, "biological activity" refers to the *in vivo* activities of a compound, composition, or other mixture, or physiological responses that result upon *in vivo* administration of a compound, composition or other mixture. Biological activity thus encompasses therapeutic effects, diagnostic effects and pharmaceutical activity of such compounds, compositions, and mixtures. The term "biologically active" or "functional" when used as a modifier of a composition of the invention containing polypeptides or compositions thereof refers to a polypeptide that exhibits at least one activity that is characteristic of or similar to a compound of the invention.

As used herein, "pharmacokinetics" refers to the concentration of an administered compound in the serum over time. Pharmacodynamics refers to the concentration of an administered compound in target and nontarget tissues over time and the effects on the target tissue (e.g., efficacy) and the nontarget tissue (e.g., toxicity). Improvements in, for example, pharmacokinetics or pharmacodynamics can be designed for a particular targeting agent or biological agent, such as by using labile linkages or by modifying the chemical nature of any linker (e.g., changing solubility, charge, and the like).

As employed herein, the phrases "an effective amount" and "therapeutically effective amount" refer to an amount of a compound of the invention that is useful or able to support an observable change in the level of one or more biological activity characteristic of a composition of the invention, or a dose

sufficient to impart a beneficial effect, e.g., an amelioration of a symptom on the recipient thereof. The specific therapeutically effective dose level for any particular subject will depend upon a variety of factors including the symptom or disorder being treated, the severity of the symptom or disorder, the activity of the specific compound, the route of administration, the rate of clearance of the compound, the duration of treatment, the drugs used in combination or coincident with the compound, the age, body weight, sex, diet, and general health of the subject, and like, as well as other factors well known in the medical arts and sciences. A therapeutically effective amount can be an amount of a compound of the invention sufficient to produce a measurable inhibition of angiogenesis in the tissue being treated, i.e., an angiogenesis-inhibiting amount. Inhibition of angiogenesis can be measured in situ by immunohistochemistry, or by other methods known to one skilled in the art. Various general considerations taken into account in determining the "therapeutically effective amount" are known to those of skill in the art and are described, e.g., in Gilman, A.G., *et al.*, *Goodman And Gilman's The Pharmacological Basis of Therapeutics*, 8th ed., McGraw-Hill (1990); and *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Co., Easton, PA (1990).

As used herein, the terms "concurrently administered" and "concurrent administration" encompass substantially simultaneous administration of one or more compounds of the invention and one other oncology therapeutic.

As used herein, the term, "non-concurrent" administration encompasses administering one or more compounds of the invention at different times, in any order, whether overlapping or not. This includes, but is not limited to, sequential treatment (such as pretreatment, post-treatment, or overlapping treatment) with the components of the combination, as well as regimens in which the drugs are alternated, or wherein one component is administered long-term and the other(s) are administered intermittently.

EXAMPLES

The versatility of the invention is illustrated by the following Examples, which illustrate typical embodiments of the invention and are not limiting of the claims or specification in any way.

Example 1

Synthesis of a compound of the invention

Peptides of the invention can be linked to antibody 38C2 by the following procedure: One mL antibody 38C2 in phosphate buffered saline (10 mg/mL) is added to 12 μ L of a 10 mg/mL stock solution of Peptide and the resulting mixture maintained at room temperature for 2 hours prior to use.


Example 2


C. Rader, *et al.*, J. Mol. Biol. 332:889-899 (2003) details one method of making h38c2. The following details the results, materials and methods in this reference. Humanization Human V κ gene DPK-9 and human J κ gene JK4 were used as frameworks for the humanization of the kappa light chain variable domain, and human V H gene DP-47 and human J H gene J H 4 are used as frameworks for the humanization of the heavy chain variable domain of m38C2. All complementarity determining region (CDR) residues as defined by Kabat *et al.*, as well as defined framework residues in both light chain and heavy chain variable domain, were grafted from m38C2 onto the human framework. The selection of grafted framework residues may be based on the crystal structure of mouse mAb 33F12 Fab (PDB 1AXT). mAb 33F12 Fab shares a 92% sequence homology with m38c2 in the variable domains and identical CDR lengths. Furthermore, both 33F12 and m38C2 have similar catalytic activity. Framework residues consisted of five residues in the light chain and seven residues in the heavy chain (Figure 7A) and encompassed the residues that are likely to participate directly or indirectly in the catalytic activity of m38C2. These include the reactive lysine of m38C2, Lys^{H93}, which is positioned in framework region 3 (FR3) of the heavy chain. Six residues, Ser^{H35}, Val^{H37}, Trp^{H47}, Trp^{H103}, and Phe^{L98}, which are conserved between mouse mAbs 33F12 and 38C2, are within a 5-Å radius of the ϵ amino group of Lys^{H93}. These residues were also conserved in the humanization. Lys^{H93} lies at the bottom of a highly hydrophobic substrate binding sites of mouse mAbs 33F12 and 38C2. In addition to CDR residues, a number of framework residues line this pocket. Among these, Leu^{L37}, Gln^{L42}, Ser^{L43}, Val^{L85}, Phe^{L87}, Val^{H5}, Ser^{H40}, Glu^{H42}, Gly^{H88}, Ile^{H89}, and Thr^{H94} were grafted onto the human framework.

Expression

By fusing the humanized variable domains to human constant domains C κ and C γ 1, h38C2 was initially generated as Fab expressed in *E. coli*. Next, h38c2 IgG was formed from h38c2 Fab using the PIGG vector engineered for human IgG1 expression in mammalian cells. Supernatants from transiently transfected human 293T cells were subjected to affinity chromatography on recombinant protein A, yielding approximately 1 mg/L h38C2 IgG1. Purity was established by SDS-PAGE followed by Coomassie blue staining.

30 β -Diketone Compounds –





3

The enaminone formed by the covalent addition of a β -diketone with m38c2 has a characteristic UV absorbance at $\lambda_{\text{max}} = 318 \text{ nm}$. Like m38C2 IgG, h38C2 IgG showed the characteristic enaminone absorbance after incubation with β -diketone. As a negative control, recombinant human anti-HIV-1 gp120 mAb b12 with the same IgG1 isotype as h38C2 but without reactive lysine, did not reveal enaminone absorbance after incubation with β -diketone **2**. For a quantitative comparison of the binding of β -diketones to m38C2 and h38C2, the authors used a competition ELISA. The antibodies were incubated with increasing concentrations of β -diketones **2** and **3** and assayed against immobilized BSA-conjugated β -diketone **1**. The apparent equilibrium dissociation constants were $38 \mu\text{M}$ (m38C2) and $7.6 \mu\text{M}$ (h38C2) for β -diketone **2** and $0.43 \mu\text{M}$ (m38C2) and $1.0 \mu\text{M}$ (h38C2) for β -diketone **3**, revealing similar β -diketone binding properties for mouse and humanized antibody (Fig. 6).

Molecular modeling – A molecular model of h38C2 Fab was constructed by homology modeling using the crystal structure of a related aldolase antibody, mouse 33F12 Fab (Protein Data Bank ID: 1AXT), as a template. The crystal structure of mouse 33F12 Fab was previously determined at a resolution of $2.15 \text{ \AA}$.⁴ Alignment of mouse 33F12 and 38C2 amino acid sequences using the HOMOLOGY module within INSIGHT II software (Accelrys) confirmed that both sequences are highly homologous. They differ from each other by 19 out of 226 amino acids in the two variable domains, and their CDRs share the same lengths. In addition to the high sequence homology, both structures exhibit considerable structural similarity, as observed by a low-resolution crystal structure of 38C2. Residues in the model were mutated to conform to the h38C2 amino acid sequence and sidechains were placed based on standard rotamers. This model was then minimized with the DISCOVER module in INSIGHT II using 100 steps each of steepest descent minimization followed by conjugate gradient minimization.

Construction of h38C2 Fab –

The sequences of the variable light and heavy chain domains of m38C2 (SEQ ID NOs:4 and 5 respectively) as well as the sequences of human germline sequences DPK-9 (SEQ ID NO:6), JK4 (SEQ ID NO:7), DP-47 (SEQ ID NO:8), and JH4 (SEQ ID NOs:9, 10, and 11) (V BASE; <http://vbase.mrc-cpe.cam.ac.uk/>) were used to design overlapping oligonucleotides for the synthetic assembly of humanized V_{κ} and V_{H} , respectively. N-glycosylation sites with the sequence NXS/T as well as internal restriction sites *HindIII*, *XbaI*, *SacI*, *ApaI*, and *SfiI* were avoided. PCR was carried out by using the Expand High Fidelity PCR System (Roche Molecular Systems). The humanized V_{κ} oligonucleotides were: L flank sense (Rader, C., Ritter, G., Nathan, S., Elia, M., Gout, I., Junbluth, A.A., J. Biol. Chem. 275: 13668-13676 (2000)) (sense 5'-GAGGAGGAGGAGGGCCCCAGGC

GGCCGAGCTCCAGATGACCCAGTCTCTCCA-3' SEQ ID NO:12); h38C2L1 (sense; 5'-GAGCTCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGTGACCGCGTCACCATCAC TTG-3') (SEQ ID NO:13); h38C2L2 (antisense; 5'-ATTCAGATATGGGCTGCCATAAGTGTGCA GGAGGCTCTGACTGGAGCGGCAAGTGATGGTGACGCGGTC-3') (SEQ ID NO:14); h38C2L3
5 (sense; 5'-TATGGCAGCCCATATCTGAATTGGTATCTCCAGAAACCAGGCCAGTCTCCTAAG CTCCTGATCTAT-3') (SEQ ID NO:15); h38C2L4 (antisense; 5'-CTGAAACGTGATGGGACACC ACTGAAACGATTGGACACTTTATAGATCAGGAGCTTAGGAGACTG-3') (SEQ ID NO:16); h38C2L5 (sense; 5'-AGTGGTGTCCCATCACGTTTCAGTGGCAGTGGTTCTGGCACAGATTTCT ACTCTCACCATCAGCAGTCTGCAACCTGAAGATTTTGCAGTG-3') (SEQ ID NO:17); h38C2L6
10 (antisense; 5'-GATCTCCACCTTGGTCCCTCCGCCGAAAGTATAAGGGAGGTGGGTGCCCTGA CTACAGAAGTACACTGCAAAATCTTCAGGTTGCAG-3') (SEQ ID NO:18); L antisense flank (C. Rader *et al.*, J. Biol. Chem. 275:13668-13676 (2000)) (antisense 5'-GACAGATGGTGCAGCCAC AGTTCGTTTGATCTCCACCTTGGTCCTCC-3' SEQ ID NO:19). The humanized V_H oligonucleotides were: H flank sense (C. Rader *et al.*, J. Biol. Chem. 275:13668-13676 (2000))(sense
15 5'-GCTGCCCAACCAGCCATGGCCGAGGTGCAGCTGGTGGAGTCTGGGGGA-3' SEQ ID NO:20); h38C2H1 (sense; 5'-GAGGTGCAGCTGGTGGAGTCTGGCGGTGGCTTGGTACAGCCT GGCGGTTCCCTGCGCCTCTCCTGTGCAGCCTCTGGCT-3') (SEQ ID NO:21); h38C2H2 (antisense; 5'-CTCCAGGCCCTTCTCTGGAGACTGGCGGACCCAGCTCATCCAATAGTTGCTA AAGGTGAAGCCAGAGGCTGCACAGGAGAG-3') (SEQ ID NO:22); h38C2H3 (sense; 5'-TCTCC
20 AGAGAAGGGCCTGGAGTGGGTCTCAGAGATTCGTCTGCGCAGTGACAACTACGCCACGC ACTATGCAGAGTCTGTC-3') (SEQ ID NO:23); h38C2H4 (antisense; 5'-CAGATACAGCGTGTTC TTGGAATTGTCACGGGAGATGGTGAAGCGGCCCTTGACAGACTCTGCATAGTGCGTG-3') (SEQ ID NO:24); h38C2H5 (sense; 5'-CAATTCCAAGAACACGCTGTATCTGCAAATGAACAGC CTGCGCGCCGAGGACACGGGCATTTATTACTGTAAAACG-3') (SEQ ID NO:25); h38C2H6
25 (antisense; 5'-TGAGGAGACGGTGACCAGGGTGCCCTGGCCCCAGTAGCTGAAACTGTAGAA GTACGTTTTACAGTAATAAATGCCCGTG-3') (SEQ ID NO:26); H flank antisense (C. Rader *et al.*, J. Biol. Chem. 275:13668-13676 (2000))(antisense 5'-GACCGATGGGCCCTTGGTGGAGGCT GAGGAGACGGTGACCAGGGTGCC-3' SEQ ID NO:27). Following assembly, humanized V_κ and V_H were fused to human C_κ and C_{γ1}, respectively, and the resulting light chain and heavy chain
30 fragment were fused and *Sfi*I-cloned into phagemid vector pComb3X as described (C. Rader *et al.*, J. Biol. Chem. 275:13668-13676 (2000); C.F. Barbas 3rd *et al.*, *Phage Display: A laboratory manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor N.Y. (2001)). To enrich for clones with the correct h38C2 sequence, Fab were displayed on phage and selected by one round of panning against the immobilized β-diketone 1 (JW) conjugated to BSA. Soluble Fab were produced from single
35 clones and tested for binding to immobilized JW-BSA by ELISA using donkey anti-human F(ab')₂ polyclonal antibodies conjugated to horseradish peroxidase (Jackson ImmunoResearch Laboratories) as secondary antibody. Light chain and heavy chain encoding sequences of positive clones were

analyzed by DNA sequencing using the primers OMPSEQ (5'-AAGACAGCTATCGCGATTGCAG-3' SEQ ID NO:28) and PELSEQ (5'-CTATTGCCTACGGCAGCCGCTG-3' SEQ ID NO:29) (C.F. Barbas 3rd *et al.*, *Phage Display: A laboratory manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor N.Y., (2001)), respectively, to confirm the assembled V_κ and V_H sequences of h38C2.

Construction, production, and purification of h38C2 IgG1 – The recently described vector PIGG (C. Rader *et al.*, FASEB J., 16:2000-2002 (2002)) was used for mammalian expression of h38C2 IgG1. The mammalian expression vector PIGG-h38c2 is illustrated in FIGURE 23. The 9kb vector comprises heavy chain γ 1 and light chain κ expression cassettes driven by a bidirectional CM promoter construct. Using primers PIGG-h38C2H (sense; 5'-GAGGAGGAGGAGGAGGAGCTCAC TCCGAGGTGCAGCTGGTGGAGTCTG-3') (SEQ ID NO:30) and GBACK (5'-GCCCCCTTATTA GCGTTTGCCATC-3' SEQ ID NO:31) (C.F. Barbas 3rd *et al.*, *Phage Display: A laboratory manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor N.Y. (2001)), the VH coding sequence from h38C2 Fab in phagemid vector pComb3X was amplified, digested with *SacI* and *ApaI*, and cloned into the appropriately digested vector PIGG. Using primers PIGG-h38C2L (sense: 5'-GAGGAGGAG GAGGAGAAGCTTGTTGCTCTGGATCTCTGGTGCCTACGGGGAGCTCCAGATGACCCAGTC TCC-3') (SEQ ID NO:32) and LEADB (5'-GCCATGGCTGGTTGGGCAGC-3' SEQ ID NO:33) ((C.F. Barbas 3rd *et al.*, *Phage Display: A laboratory manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor N.Y. (2001)) the light chain coding sequence from h38C2 Fab in phagemid vector pComb3X was amplified, digested with *HindIII* and *XbaI*, and cloned into the appropriately digested vector PIGG that already contained the h38C2 heavy chain. Intermediate and final PIGG vector constructs were amplified in *E. coli* strain SURE (Stratagene) and prepared with the QIAGEN Plasmid Maxi Kit. h38C2 IgG1 were produced from the prepared final PIGG vector construct by transient transfection of human 293T cells using Lipofectamine 2000 (Invitrogen). Transfected cells were maintained in GIBCO 10% ultra-low IgG (<0.1%) FCS (Invitrogen) in RPMI 1640 (Hyclone) for 2 weeks. During this time, the medium was collected and replaced three times. The collected medium was subjected to affinity chromatography on a recombinant Protein A HiTrap column (Amersham Biosciences). This purification step yielded 2.45 mg h38C2 IgG1 from 2,300 mL collected medium as determined by measuring the optical density at 280 nm using an Eppendorf BioPhotometer. Following dialysis against PBS in a Slide-A-Lyzer 10K dialysis cassette (Pierce), the antibody was concentrated to 760 μ g/mL using an Ultrafree-15 Centrifugal Filter Device (UFV2BTK40; Millipore), and sterile filtered through a 0.2- μ m Acrodisc 13MM S-200 Syringe Filter (Pall). The final yield was 2.13 mg (87 %). Purified h38C2 IgG1 was confirmed by nonreducing SDS-PAGE followed by Coomassie Blue staining.

Enaminone formation – Antibody (h38C2 IgG1 or b12 IgG1) was added to β -diketone 2 to a final concentration of 25 μ M antibody binding site and 125 μ M β -diketone. This mixture was incubated at

room temperature for 10 minutes before a UV spectrum was acquired on a SpectraMax Plus 384 UV plate reader (Molecular Devices) using SOFTmax Pro software (version 3.1.2).

Binding assays – Unless noted otherwise, all solutions were phosphate buffered saline (pH 7.4). A 2 x solution of either β -diketone 2 or 3 (50 μ L) was added to 50 μ L of the antibody (either h38C2 or m38C2) and allowed to incubate at 37°C for 1 hr. Solutions were mixed by pipetting. Final concentrations of antibody were 0.4 to 8 nM antibody binding site, and final concentrations of β -diketones 2 and 3 were 10^{-9} to 10^{-2} M and 10^{-10} to 10^{-4} M, respectively. Each well of a Costar 3690 96-well plate (Corning) was coated with 100 ng of the BSA conjugate of β -diketone 1 in TBS. Wells were then blocked with 3% (w/v) BSA in TBS. Then, 50 μ L of the antibody/ β -diketone mixture was added, followed by 50 μ L of a 1:1,000 dilution of either goat anti-human Fc IgG polyclonal antibodies (Pierce) or rabbit anti-mouse Fc IgG polyclonal antibodies (Jackson ImmunoResearch Laboratories) conjugated to horseradish peroxidase. This was followed by 50 μ L ABTS substrate solution.

Between each addition, the plate was covered, incubated at 37°C for 1 hr, and then washed five times with deionized H₂O. The absorbance at 405 nm was monitored as described above until the reaction with no β -diketone reached an appropriate value ($0.5 < A_{405} < 1.0$). For each well, the fractional inhibition of ELISA signal (v_i) was calculated using equation i:

$$v_i = (A_o - A_i) / (A_o) \quad (i)$$

where A_o is the ELISA absorbance obtained in the absence of β -diketone and A_i is the absorbance obtained in the presence of β -diketone. For monovalent binding proteins, the fraction of antibody bound to soluble β -diketone (f) is equal to v_i . However, the IgG antibody is bivalent, and the ELISA signal is inhibited only by the presence of doubly liganded antibody and not by monovalent binding. Therefore, the Stevens correction for a bivalent antibody was used:

$$f_i = (v_i)^{1/2} \quad (ii)$$

The following relationship was used to determine the apparent equilibrium dissociation constant (modified from [ref. 37]):

$$f_i = f_{min} + (f_{max} - f_{min})(1 + K_D/a_0)^{-1} \quad (iii)$$

where a_0 corresponds to the total β -diketone concentration, K_D is the equilibrium dissociation constant, and f_{min} and f_{max} represent the experimentally determined values when the antibody binding sites are unoccupied or saturated, respectively. Because this equation is only valid when the K_D values are at least 10 x higher than the antibody concentration, it was verified that the K_D values determined from equation iii met this criterion. Data were fit using a nonlinear least-squares fitting procedure of KaleidaGraph (version 3.0.5, Abelbeck software) with K_D , f_{max} , and f_{min} as the adjustable parameters and normalized using equation iv:

$$f_{norm} = (f_i - f_{min}) / (f_{max} - f_{min}) \quad (iv)$$

Example 3

Linking Peptides of the invention with Linkers of the invention.

Compounds of the invention may be prepared by several approaches. In one approach, a [Peptide]-linker compound is synthesized with a linker that includes one or more reactive groups designed for covalent reaction with a side chain of an amino acid in a combining site of an antibody. The targeting agent-linker compound and antibody are combined under conditions where the linker reactive group forms a covalent bond with the amino acid side chain.

In another approach, linking can be achieved by synthesizing an antibody-linker compound comprising an antibody and a linker wherein the linker includes one or more reactive groups designed for covalent reaction with an appropriate chemical moiety of the [Peptide]. A [Peptide] may need to be modified to provide the appropriate moiety for reaction with the linker reactive group. The antibody-linker and [Peptide] are combined under conditions where the linker reactive group covalently links to the targeting and/or biological agent.

A further approach for forming an antibody-[Peptide]-conjugate uses a dual linker design. In certain embodiments, a [Peptide]-linker compound is synthesized which comprises a [Peptide]-linker and a linker with a reactive group. An antibody-linker compound is synthesized which comprises an antibody and a linker with a chemical group susceptible to reactivity with the reactive group of the [Peptide]-linker of the first step. These two linker containing compounds are then combined under conditions whereby the linkers covalently link, forming the antibody-[Peptide] compound. Table 9 shows exemplary compounds of the invention comprising [VEGF-Peptides] covalently linked to linkers: D-amino acids are indicated by the use of lower case type script: for example, Compound 2018 (comprising SEQ ID NO:78, substituted with K¹⁰ as linking residue, also equal to SEQ ID NO:195) includes the residues "R-L-Y-(D-Ala)-(D-Leu)", which are written "R-L-Y-a-l". Disulphide bonds are depicted as connecting lines between two cysteine residues.

Z may be a group that forms a reversible or irreversible covalent bond. In some embodiments, reversible covalent bonds may be formed using diketone Z groups such as those shown in FIGURE 3. Thus, structures A-C may form reversible covalent bonds with reactive nucleophilic groups (e.g. lysine or cysteine side chain) in a combining site of an antibody. R'₁, R'₂, R'₃, and R₄ in structures A-C of FIGURE 3 represent substituents which can be C, H, N, O, P, S, halogen (F, Cl, Br, I) or a salt thereof. These substituents also may include a group such as an alkyl, alkenyl, alkynyl, oxoalkyl, oxoalkenyl, oxoalkynyl, aminoalkyl, aminoalkenyl, aminoalkynyl, sulfoalkyl, sulfoalkenyl, or sulfoalkynyl group, phosphoalkyl, phosphoalkenyl, phosphoalkynyl group. R'₂ and R'₃ also could from a ring structure as exemplified in structures B and C. X in FIGURE 3 could be a heteroatom. Other Z groups that form reversible covalent bonds include the amidine, imine, and other reactive groups encompassed by structure G of FIGURE 3. FIGURE 4 includes the structures of other linker

reactive groups that form reversible covalent bonds, e.g., structures B, G, H, and, where X is not a leaving group, E and F.


Z reactive groups that form an irreversible covalent bond with a combining site of an antibody include structures D-G in FIGURE 3 (e.g., when G is an imidate) and structures A, C and D of FIGURE 4.

When X is a leaving group, structures E and F of FIGURE 4 may also form irreversible covalent bonds. Such structures are useful for irreversibly attaching a targeting agent-linker to a reactive nucleophilic group to a combining site of an antibody. As used herein, L' is a linker moiety linking an antibody to the targeting agent and having the formula $-X-Y-Z'-$. FIGURES 5 and 6, respectively,


illustrate the addition mechanism of a reactive, nucleophilic side chain in a combining site of an antibody to the Z moieties illustrated in FIGURES 3 and 4.

This is shown below for the case where the linker has a diketone moiety as the reactive group and linkage occurs with the side chain amino group of a lysine residue in the antibody combining site.

The Antibody is shown schematically as bivalent with a reactive amino acid side chain for each combining site indicated.




Another embodiment shown below is for the case where the linker has a beta lactam moiety as the reactive group and linkage occurs with the side chain amino group of a lysine residue in the antibody combining site. The Antibody is shown schematically as bivalent with a reactive amino acid side chain for each combining site indicated.




Compounds of the invention may also be readily synthesized by covalently linking a [Peptide]-linker compound as described herein to a combining site of a multivalent antibody. For example, a [Peptide]-linker conjugate, where the linker includes a diketone reactive moiety, can be incubated with 0.5 equivalents of an aldolase antibody, such as h38C2 IgG1 to produce a compound of the invention.

A further example of a reaction mechanism combining a [Linker]-[Antibody] to a peptide is shown below, employing 'click chemistry' to combine an alkyne and an azide (where R may be attached to a [Peptide] of the invention).



Example 4

Synthesis of:



is provided in FIGURE 7.

Example 5

The ability of Ang2 binding compounds to interact with Ang2 was measured by competition with Tie-2. These tests were described in US2008166364 (and also in PCT/IE2007/000110), whose contents are herein incorporated by reference in their entirety. In particular, those portions of PCT/IE2007/000110 and US2008166364 detailing the advantages and disadvantages of various Ang2-binding peptides, and the advantages and disadvantages of covalently linking certain Ang2-binding peptides to catalytic antibodies (and those portions specifically relating to linking to h38C2), as well as the advantages and disadvantages of using specific residues of the respective Ang2-specific peptide as the linking residue (in this context, linking residue meaning the residue covalently linked to the linker (for example as used herein in Example 4)). For competitive ELISA, human angiopoietin-2 protein and Tie-2-Fc (R&D Systems) were reconstituted without carrier protein. Mouse anti-human Tie-2 (Pharmingen) was used as the primary antibody and goat anti-mouse-IgG1-HRP (Pierce) was used as the secondary antibody. TMB substrate from Pierce was used.

High-binding half-well plates were coated with Ang-2 (100 ng/well) in 50 μ l PBS and incubated at 4°C overnight. Plates were washed three times with washing buffer (0.1% Tween 20, PBS, pH 7.4) and blocked with Superblock (Scytek), 100 μ l/well at room temperature ("RT") for 1 hour. After removing the blocking solution, 50 μ l of an Ang-2 binding peptide compound (1 μ M and 5x serial dilution) in the presence of 0.25 nM hTie-2-Fc using Superblock as diluent was added and incubated at RT for 2 hours. Plates were washed 3 times with washing buffer. Then, 50 μ l of 0.1 μ g/ml mouse anti-human Tie-2 diluted in Superblock added and incubated at RT for 1 hour. Following incubation, 50 μ l of 1:5,000 dilution of goat anti-mouse IgG-HRP in Superblock was then added and incubated at RT for one hour. After washing 3 times, 50 μ l (25 μ l TMB + 25 μ l H₂O₂) was added, and incubated for 3-5 minutes. Color development was monitored and stopped with 25 μ l of 2 M H₂SO₄. OD450 nm with a correction wavelength of 540 nm was measured. IC₅₀ values (50% inhibition of Ang-2-Tie-2 binding) were calculated using non-linear Sigmoidal dose-response curve fitting function in the Prism 4 software (GraphPad).

For reverse competition ELISA, human Tie-2-Fc, angiopoietin-2 protein, biotinylated anti-human Ang-2 antibody, and streptavidin HRP (R&D Systems) and TMB substrate from Pierce were used.

High-binding half-well plates were coated with Tie-2-Fc (50 ng/well) in 50 μ l PBS and incubated at 4°C overnight. Plates were washed three times with washing buffer (0.1% Tween 20, PBS, pH 7.4) and blocked with Superblock, 150 μ l/well at RT for 1 hour. Plates were washed three times. Following washing, 50 μ l of an Ang-2 binding peptide compound (50 nM, 5x serial dilution) in the presence of 50 ng/ml (0.83 nM) Ang-2 in Superblock were added and incubated at RT for 1 hour. Plates were washed 3 times, 50 μ l of 1 μ g/ml biotinylated anti-Ang-2 detection antibody in Superblock was added and incubated at RT for 2 hours. Plates were washed 3 times, and 50 μ l of

streptavidin HRP (1:200 dilution in Superblock) was added at RT for 20 minutes. Plates were washed 3 times, and 50 μ l (25 μ l TMB + 25 μ l H₂O₂) substrate solution was added and incubated for 20-30 minutes. Color development was stopped with 25 μ l of 2 M H₂SO₄. OD450 nm with a correction wavelength of 540 nm was measured. IC₅₀ values (50% inhibition of Ang-2-Tie-2 binding) were calculated using non-linear Sigmoidal dose-response curve fitting function in the Prism 4 software.

IC₅₀ values for exemplary Ang-2 binding peptide compounds as determined by competitive ELISA are presented in Table 5. IC₅₀ values are provided for the Ang2-binding peptide covalently linked to the Linker as shown in Example 4 via the side chain of the Ang2-linking residue (T) and the Ang2-binding peptide covalently linked to the linker of Example 4 via the side chain of the Ang2-linking residue, wherein the Z group of the linker is covalently linked to the combining site of h38C2 (P).

In Table 5, compounds 4021, 4022 and 4023 are Ang-2 binding peptides alone, not conjugated to either a linker or linker-antibody; compounds 4024 and 4025 are Ang-2 binding peptides conjugated to a linker-antibody, where the linker is 4P ("4" PEG); and compounds 4026-4063 are Ang-2 binding peptides conjugated to a linker-antibody, where the linker is OP ("O" PEG) and has the structure of the linker shown in Example 4. Compounds 4024-4063 were conjugated to humanized aldolase antibody when obtaining the data shown below, except where otherwise indicated. All compounds of the invention shown in the Tables were capped with an acyl group at the N-terminus and an amino group at the C-terminus, except where otherwise indicated (e.g. compounds 4026, 4049, 4050, 4051 and 4052). In Table 5, amino acid sequences of peptide compounds are shown with the position of linker OP or 4P indicated in parentheses following the internal amino acid residue to which the linker is attached. For compound 4026, the N-terminal "OP" linker is indicated at the beginning of the peptide sequence.

For example, compound 4024 in Table 5 has the following sequence: Q(Kac)Y QPL DEL DK(4P)T LYD QFM LQQ G (parent SEQ ID is SEQ ID NO:138). In this example, the second amino acid residue is epsilon acyl lysine, followed by tyrosine, and the linking position (in this case a 4P linker) is the lysine residue 11, followed by threonine. Also, compound 4052 has the following sequence: (Amido 2-PEG)QKacY QPL DEL DK(OP)T LYD QFMLQQ G (SEQ ID NO:162). In this case, the N-terminal glutamine residue is capped by an amido-2-PEG group, the second amino acid residue is epsilon acyl lysine, and the OP linker is attached to lysine residue 11.

Tables 5 and 6 also show half-life ($T_{1/2}$) and "screening" half life (results in parentheses), an alternative method of determining $T_{1/2}$, based on a shorter test period. For "screening" $T_{1/2}$, test compounds were intravenously administered into male Swiss Webster mice. Blood samples were

taken from 4 mice per time point via retroorbital sinus bleed at the following time points: 0.08, 5, and 32 hours. Blood level of test compounds were determined by ELISA. The data were reported as the percentage of test compounds at 32 hours versus 5 min. Normal $T_{1/2}$ was calculated in a similar fashion, after undergoing additional data analysis using WinNonlin version 4.1 (Pharsight

Corporation). The data was fit to a model based upon the shape of the curve (i.e. a bi-exponential decline will be fit to a two compartment model, etc.) The criteria for best fit (i.e. lowest %CV) was based on iterative re-weighted least squares.

IC50 values for exemplary Ang-2 binding peptide compounds as determined by reverse competitive ELISA are presented in Table 5. IC50 values are provided for the targeting peptide plus linker and the targeting peptide linked to an antibody (P) via the linker of Example 4, unless otherwise specified. In Table 5, compounds are conjugated to humanized aldolase antibody h38c2 and the linker structures shown in Example 4 (OP).

Xenograft Studies

Colo205 cells were cultured with 10% FBS RPMI medium and 3×10^6 cells in 0.1 ml Hank's balanced salt solution (HBSS) were injected subcutaneously into the upper right flank of nude mice. After 7-9 days, animals were randomized into appropriate number of groups with average tumor size of 200-300 mm³. Mice were then treated with the requisite amount of compounds of the invention and tumor volumes were measured twice a week. Animals were terminated once their average tumor volume reached 2000 mm³. Upon termination, tumors were weighed and saved for further histological studies. Treatment efficacy was evaluated by measuring the difference in the tumor volumes of treated versus control groups. Results are reported as % T/C, where % T/C was calculated as: $\% T/C = (V_t - V_0)/(C_t - C_0) \times 100$, where, V_0 and V_t were the average tumor volumes of treated groups at the beginning and termination of the group. C_0 and C_t were the average tumor volumes of the control group at the beginning and termination of the group (Table 7).

Example 6

Exemplary [VEGF-Peptide]-[Linker] compounds are shown in Table 9 and exemplary [VEGF-Peptide]-[Linker]-Antibody compounds are shown in Table 10.

Example 7

VEGF Peptide binding assay


High-binding half-well 96-well plates (Costar #3690) were coated with recombinant human VEGFR2/Fc (50 ng/well) in 50 μ l PBS, and incubated at 4°C overnight. After washing the plates three times with 1X washing buffer (KPL Cat#50-65-00) and blocking with Superblock (Scytek #AAA500) at room temperature (RT) for 1 hour, VEGF-peptides of the invention were serially

diluted (concentrations: 0.128 – 10,000 nM) in the presence of 25 ng/ml of recombinant human VEGF165 in 50 µl Superblock, and were then added to the 96-well plates and incubated at RT for 1 hour. Peptides of the invention tested in this fashion were all capped at the amino terminus with a C(O)CH₃ group and at the carboxyl terminus with a NH₂ group. The plates were then washed three times with washing buffer, and a diluted (1:500) biotinylated anti-human VEGF antibody (R&D systems, Cat# BAF293) was added and incubated at RT for 2 hours. After washing the plates, streptavidin HRP conjugate (1:200 in Superblock) (R&D systems, Cat# DY998) was added to detect the bound anti-VEGF antibody, followed by addition of tetramethylbenzidine (TMB) substrate (Pierce, Cat# 34021) for color development. The reaction was stopped with 2 M H₂SO₄. OD450 was measured using SpectraMax (Molecular Device) with a correction wavelength of 540 nM. The data were analyzed using Prism software. OD450 values were plotted as a function of the concentrations of the VEGF peptides. IC₅₀ values, indicative of the potency of the VEGF peptides to inhibit VEGF-VEGFR2 interaction, were obtained using a sigmoid dose-response curve fit in Prism. Results of the VEGF binding assays are shown in Tables 1 and 2 (Table 1 features VEGF Peptides with V¹² (excepting SEQ ID NO:34) and Table 2 features VEGF peptides with E¹²). Tables 1 and 2 indicate the compounds of the invention tested by the parent SEQ ID NO, although the N' and C' termini of the tested compounds were capped as described above. Not all compounds were tested at the same instance: IC₅₀ values given in parantheses have been normalized.

Example 8

Peptide tethering

A solution of linker (as shown in Example 4) (0.1 g, 0.3 mmol), HBTU (O-Benzotriazole-N,N,N',N'-tetramethyl-uronium-hexafluoro-phosphate) (0.11 g, 0.3 mmol) in anhydrous DMF (Dimethylformamide) (2 mL) was cooled to 2-5⁰ C in an ice bath under argon atmosphere. NMM (N-methylmorpholine) (0.12 g, 0.9 mmol) was introduced via syringe and stirred for 5 minutes. Another cold solution of Peptide (SEQ ID NO:64) (0.24 g, 0.1 mmol) in anhydrous DMF (2 ml) was added to the first reaction vessel and stirred for 30 minutes. After the completion of reaction as monitored by LCMS (Liquid Chromatography/ Mass Spectrometry), water (1 ml) was added and the reaction mixture was purified using HPLC (High Performance Liquid Chromatography) to get a pure product (145 mg) with a mass of 2785. In the figure below, full structures of non-natural amino acids and residues involved any bonding other than peptide binding are depicted. Thus the N of both cysteines is shown. Other Compounds of the invention were formed by covalent linkage



Example 9

Covalently linking [Peptide]-[Linker] moiety to [Antibody]

5 Programming and purification of [Peptide]- [Linker]-[Antibody] complexes

Tethered peptides were programmed at a 3:1 ratio with h38C2 at room temperature, overnight. The resultant [Peptide]-[Linker]-[Antibody] macromolecules were purified by PD-10 desalting columns. Protein concentrations were measured by UV A280. [Peptide]-[Linker]-[Antibody] macromolecules were characterized by LCMS analysis for programming efficiency.

10

Example 10

Mouse PK assays (Reverse ELISA format)

PK studies were conducted using male Swiss Webster mice (CFW, Charles Rivers Hollister, CA) weighing approximately 20-22 grams at the start of dosing. [VEGF-Peptide]-[Linker]-[Antibody] complexes were given by IV administration through the tail vein, blood samples were taken from 3 mice per time point via retro orbital sinus bleed at the following time points: 0.08, 0.5, 1, 3, 5, 7 and 24 hours. For time points up to 32 hours mice were anesthetized with isoflurane and bleed volumes did not exceed 0.1 mL/bleed. For the remaining time points (32 - 120 hrs), mice were killed by CO₂ inhalation and terminal cardiac samples were drawn. Protease Inhibitor cocktail was added to all blood tubes prior to sample collection. Blood was allowed to clot on ice for 30 minutes and then centrifuged at 12000 rpm for 5-10 minutes at 4°C to collect serum and immediately stored at -80°C until analysis via ELISA. Dosing solutions were used to establish the standard curves for serum sample analysis by VEGF Reverse ELISA. Aliquots of each serum sample were analyzed by either

20

VEGF Reverse ELISA (also described herein as IgG Coated ELISA) or VEGF ELISA (VEGF Coated ELISA).

VEGF Reverse ELISA

- 5 High-binding half-well 96-well plates (Costar #3690) were coated with 1:100 goat anti-human IgG in coating buffer (Bethyl Laboratories ELISA kit, Cat# E80-104) at 4°C overnight. After washing the plates three times with 1X washing buffer (KPL Cat#50-65-00) and blocking with Superblock (Scytek #AAA500) at room temperature (RT) for 1 hour, prepared dosing solution standards (range: 3.91 - 500 ng/ml) and serum samples were added to the plate and incubated for 1 hour on a plate shaker to
- 10 allow binding of [VEGF-Peptide]-[Linker]-[Antibody] complexes to the plates. A fixed concentration of VEGF (3 nM) was then added and incubated for 1 hour on a plate shaker. The plates were then washed three times with washing buffer, and a diluted (1:500) biotinylated anti-human VEGF antibody (R&D systems, Cat# BAF293) was added and incubated at RT for 2 hours. After washing the plates, streptavidin HRP conjugate (1:200 in Superblock) (R&D systems, Cat# DY998) was added
- 15 to detect the bound anti-VEGF antibody, followed by addition of tetramethylbenzidine (TMB) substrate (Pierce, Cat# 34021) for color development. The reaction was stopped with 2 M H₂SO₄. OD450 was measured using SpectraMax (Molecular Device) with a correction wavelength of 540 nM. Serum concentrations of the [VEGF-Peptide]-[Linker]-[Antibody] complexes were calculated using the standard curves. Serum compound concentrations, as determined by ELISA, were plotted as
- 20 a function of time. Further data analysis was undertaken using WinNonlin version 4.1 (Pharsight Corporation) to determine the β half life ($T_{1/2}$) and the area under the curve (AUC) for [Peptide]-[Linker]-[Antibody] complexes.

VEGF ELISA

- 25 High-binding half-well 96-well plates (Costar #3690) were coated with rhVEGF165 (6.25 ng/well) in 50 μ l PBS/well and incubated at 4°C overnight. After washing the plates three times with 1X washing buffer (KPL Cat#50-65-00) and blocking with Superblock (Scytek #AAA500) at room temperature (RT) for 1 hour, prepared dosing solution standards (range: 3.91 - 500 ng/ml) and serum samples were added to the plate and incubated for 1 hour to allow binding of [VEGF-Peptide]-[Linker]-[Antibody]
- 30 complexes to the coated VEGF on the plates. The plates were then washed three times with washing buffer, diluted goat anti-human IgG-HRP (1:5000) (Bethyl A80-104P-52) was added and incubated at RT for 1 hour, followed by addition of tetramethylbenzidine (TMB) substrate (Pierce, Cat# 34021) for color development. The reaction was stopped with 2 M H₂SO₄. OD450 was measured using SpectraMax (Molecular Device) with a correction wavelength of 540 nM. Serum concentrations of
- 35 the [VEGF-Peptide]-[Linker]-[Antibody] complexes were calculated using the standard curves. Serum [VEGF-Peptide]-[Linker]-[Antibody] complexes concentrations, as determined by ELISA, were plotted as a function of time. Further data analysis was undertaken using WinNonlin version 4.1

(Pharsight Corporation) to determine the β half life ($T_{1/2}$) and the area under the curve (AUC) for [VEGF-Peptide]-[Linker]-[Antibody] complexes.

Table 3 shows the binding of a series of [VEGF-Peptide]-[Linker]-[Antibody] complexes. The [VEGF-Peptide] portion of each complex was based on SEQ ID NO:34, and with one of the residues from X^1 through to X^{19} substituted by K as a linking residue, with the exception of C^5 and C^{15} . In each case, the [VEGF-Peptide] was capped at the amino terminus with a $C(O)CH_3$ group and at the carboxy terminus with a NH_2 group. The [Linker] portion of each complex was that depicted in Example 4. The [Antibody] portion of each complex was h38C2: comprising SEQ ID NO:1 and SEQ ID NO:2. Table 6 also shows half-life data for certain [VEGF-Peptide]-[Linker]-[Antibody] complexes, as well as data representing area under the curve (AUC). The AUC data represents the total area under the curve, taking into account both the α -half life and the β -half life.

Table 4 shows the binding of a series of [VEGF-Peptide]-[Linker]-[Antibody] complexes. The [VEGF-Peptide] portion of each complex was based on the SEQ ID NO indicated, and with the indicated residue substituted by K as a linking residue. In each case, the [VEGF-Peptide] was capped at the amino terminus with a $C(O)CH_3$ group and at the carboxy terminus with a NH_2 group. Residues 1 to 18 of the peptides are not depicted in Table 4, but were as described elsewhere, with the exception of the respective substitution of the linking residue (in each case of these embodiments, the linking residue being K). The [Linker] portion of each complex was that depicted in Example 4. The [Antibody] portion of each complex was h38C2: comprising SEQ ID NO:1 and SEQ ID NO:2. Table 4 shows also shows some half life data for certain [Peptide]-[Linker]-[Antibody] complexes. Results were obtained from both VEGF ELISA and from VEGF Reverse ELISA (results in parentheses).

Example 11


Synthesis of compounds of the invention of the formula: [Ang2-Peptide]-[Connector]-[VEGF-Peptide] is shown in Figures 8-10.

Example 12

Exemplary compounds of the invention of the formula: [Ang2-Peptide]-[Connector]-[VEGF-Peptide] are shown in Table 11.

Example 13

Table 12 shows exemplary compounds of the invention of the formula:



Example 14

Compounds of Example 13 were tested for VEGF binding (IC₅₀), VEGF T¹/₂, Ang2 binding (IC₅₀), and Ang2 T¹/₂. Testing methods were as described above. Results are shown in Table 8. Compounds of the invention exemplified in Example 13 contains a fixed ratio of [Ang-2-Peptide] and [VEGF-Peptide]. The circulating levels of Ang-2 is reported to be about 10 times higher than VEGF circulating levels in prostate (2.5 ng/ml Ang-2 and 0.2 ng/ml VEGF) and breast cancer (2 ng/ml Ang-2 and 0.3 ng/ml VEGF) patients (Ref. *European Journal of Clinical Investigation* (2003) 33, 883–890). It will be apparent that some compounds of the invention (e.g. see Example 14) demonstrate a great variety of binding IC₅₀s and *in vivo* half lives of both Ang2 and VEGF relative to each other within the same molecule, and also relative to other molecules. Thus, for certain applications, it may be desirable to use a Compound whose binding affinity for Ang2 is between about 3 to about 15 times greater than its binding affinity for VEGF, so as to allow for the greater circulating amounts of the respective targets. Naturally, different therapeutic applications may require or permit or favour different Compounds of the invention.

For example, Compound 6053 showed a K_D of 3E-10 for VEGF binding and a K_D of 9E-11 for Ang-2 binding in Biacore assay. The higher affinity for Ang-2 binding may be advantageous as the plasma concentration Ang-2 has been reported as being higher in certain cancer patients. Compound 6053 shows similar VEGF inhibition IC₅₀ (0.7 nM) and Ang-2 inhibition IC₅₀ (0.6 nM). In addition, the stability (T¹/₂) in mice of both the [Ang2-Peptide] and the [VEGF-Peptide] components of Compound 6053 is very similar. A subsequent assay was conducted comparing Compound 2018, Compound 4043 and Compound 6053 (Table 8).

***In vivo* Pharmacology**

The anti-tumor activity of Compound 6053 was evaluated in human xenograft models. Xenografts were induced by subcutaneous (SC) implantation of tumor cells into 5-7 week old female nu/nu mice and allowed to establish to a volume of 200 ~ 400 mm³ prior to initiation of treatment. Once tumors were established, mice were randomized to treatment groups with identical tumor volumes (n=9-10/group), and Compound 6053 treatment was administered once weekly by intraperitoneal (IP) injection. In combination studies, additional anti-cancer agents were administered either weekly by IP injection or daily by oral gavage (PO), with treatments initiated concomitant with Compound 6053. Tumor volumes were measured once or twice weekly, using calipers, and body weights were measured weekly, during the treatment period. In some studies, all mice were killed by CO₂ asphyxiation and tumors were excised once tumor volume in the Vehicle-treated control group reached 2000 mm³. In pseudo-survival studies, mice were killed by CO₂ asphyxiation and tumors were excised, weighed, and processed for further histological and/or immunochemical evaluation once the mean tumor volume of each treatment group exceeded 2000 mm³.

Example 15**Single Agent Studies**

An experiment conducted in the Colo205 (human colon adenocarcinoma) xenograft model is illustrated in Figure 11. Weekly administration of Compound 6053 dose-dependently inhibited Colo205 tumor growth. At day 35, Compound 6053 at 3 mg/kg (IP, 1x/wk) resulted in a significant ~30% reduction in tumor growth, whereas at 10 or 30 mg/kg (IP, 1x/wk) a significant ~50% reduction in tumor growth was seen when compared with the Vehicle-treated controls. Ten or 30 mg/kg Compound 6053 treatment also led to sustained tumor inhibition compared with the control group. Compound 6053 treatment did not affect body weight gain (data not shown) and mice appeared to be in good health throughout the study.

At day 35, Vehicle-treated mice and half of each the Compound 6053-treated groups were killed, and tumors were excised and snap frozen. To assess the anti-angiogenic effect of Compound 6053, tumor microvessel density was assessed immunohistochemically on frozen sections of Colo205 colon adenocarcinoma xenograft tumors treated with Vehicle or Compound 6053. Tumors were stained with a mouse-specific monoclonal antibody to CD31, and immunoreactivity was quantified from 5 areas of 3 sections from each tumor. Tumor microvessel density was significantly reduced ~44% by Compound 6053 treatment (30 mg/kg, 1x/wk) in comparison with the Vehicle-treated group (Figure 12) confirming the anti-angiogenic activity of the molecule.

In an independent Colo205 xenograft study, weekly administration of Compound 6053 again dose-dependently inhibited Colo205 tumor growth. At the 30 mg/kg weekly dose, Compound 6053 resulted in a significant ~70% reduction in tumor growth at day 28 compared with the Vehicle-treated controls (killed by CO₂ asphyxiation (Figure 13). Excised tumors were sectioned and stained with Hematoxylin and Eosin (H&E), and viable tumor area was assessed using standard image analysis computation methods and software. The viable tumor area was determined for 4 sections from each tumor, and the mean value of these measurements was then multiplied by the tumor volume at study termination to yield an estimate of viable tumor volume for each tumor. Viable tumor volume data from the experiment in Colo205 colon adenocarcinoma xenografts are shown in Figure 14. Viable tumor volume significantly decreased ~60% and ~76% in Compound 6053-treated mice at 10 mg/kg and 30 mg/kg dose, respectively, in comparison with the Vehicle-treated group.

To investigate whether Compound 6053 targets both Ang2 and VEGF in vivo, the effects of Compound 6053 on Ang2 expression and phosphorylated VEGFR2 (pVEGFR2) levels were assessed using immunofluorescence in Colo205 xenograft tumors treated with Vehicle or Compound 6053. Frozen sections of these tumors were stained with FITC-labeled anti-Ang2 monoclonal antibody, and

Ang2 immunoreactivity was quantified from 3 images of 1 section from each tumor. Ang2 immunoreactivity was significantly reduced by ~70% in Compound 6053-treated groups (10 and 30 mg/kg) in comparison with the Vehicle-treated group (Figure 15). Frozen sections of these tumors were also double-stained with FITC-labeled VEGFR antibody and a rodamine-labeled pVEGFR2 antibody, and the pVEGFR2 immunoreactivity was quantified from 3 images of 1 section from each tumor and expressed as a percentage of total VEGFR2 immunoreactivity (pVEGFR2/VEGFR2). pVEGFR2/VEGFR2 was significantly reduced by Compound 6053 treatment in a dose-dependent manner in comparison with the Vehicle-treated group, being ~43% and ~70% at 10 and 30 mg/kg, respectively (Figure 16). These data demonstrate that Compound 6053 affects both Ang2 and VEGF pathways in Colo205 xenograft model.

The anti-tumor efficacy of Compound 6053 was also evaluated in an MDA-MB-435 breast carcinoma xenograft model and an A431 skin carcinoma xenograft model. Weekly administration of Compound 6053 (30 mg/kg IP) resulted in a significant 45% reduction (day 68) in tumor growth in MDA-MB-435 model (Figure 17A), and a significant 54% reduction (day 35) in tumor growth of A431 model (Figure 17B). Thus, Compound 6053 demonstrates significant anti-tumor efficacy in three different human xenograft tumor models.

Further single agent studies were conducted with Compounds 2018, 2036, 2049, 2071, 4043, and 6053, and are shown in Figures 23-28.

Example 16

Combination Studies

The anti-tumor efficacy of Compound 6053 was further evaluated in combination therapy studies conducted in the Colo205 xenograft model, comparing the anti-tumor efficacy of Compound 6053 (10 mg/kg IP, weekly) alone and in combination with 5-fluorouracil (5-FU, Figure 18), Irinotecan (Figure 19), Taxotere (Figure 20), Sunitinib (Figure 21), and Axitinib (Figure 22). Combination of Compound 6053 with these chemotherapy agents (5-FU, Irinotecan, Taxotere) or multiple tyrosine kinase inhibitors tested (Sunitinib, Axitinib) resulted in significantly greater inhibition of tumor growth rate than either of the monotherapy alone ($p < 0.05$ by Two-way ANOVA with Bonferroni Multiple Comparison Test). In the Compound 6053 + Taxotere combination study (Figure 17), one tumor was an outlier (attained a tumor volume $> 2000 \text{ mm}^3$ at day 57, requiring euthanasia), and was therefore removed from the analysis. In conclusion, these data provide excellent preclinical rationale for broad application of Compound 6053 in combination with multiple anti-cancer therapeutics targeting both tumor and angiogenic mechanisms of action.

Summary of Preclinical Efficacy

Compound 6053 inhibited human Ang2 and human VEGF binding to their cognate receptors with sub-nanomolar potencies, and demonstrated a prolonged and balanced pharmacokinetic profile across preclinical species. Compound 6053 was efficacious across multiple human xenograft tumor models, and provided additional benefit in combination with multiple standard therapeutic agents, including anti-angiogenics and anti-tumor cell therapeutics, as evidenced by delayed tumor growth and time-to-progression. Taken in total, these data suggest that the high potency of Compound 6053 for Ang2 and VEGF significantly impacts tumor growth and viability alone and in combination with standard chemotherapeutic agents.

Example 17

Anti-angiogenic activity in VEGF induced Rabbit Eye Retinal Leakage Assay

The anti-angiogenic activity of compounds 6054, 6037, and 6044 was evaluated in the VEGF induced rabbit eye retinal leakage assay. This assay was performed in a 3 day or 7 day format. In a 3 day assay format the right eye of each rabbit (3 rabbits/group) was injected (IVT) with compound (1mg/eye) or vehicle on day 0 followed by an injection (IVT) of VEGF (1 µg/eye) into the same eye on the next day or day 1. On day 3, fluorescein was injected by IV and angiography of the eye was taken. In a 7 day assay format compounds (1mg/eye) were injected on day 0, VEGF (1 µg/eye) was injected on day 5 and fluorescein was injected by IV on day 7. In addition, binding and half-life were assayed as described above and are shown in Table 14. As a reference, the VEGF half-life of Compound 6037 is indicated as approximately 36 hours in Table 14, this figure being the VEGF half-life of the analogous molecule, Compound 2053 (see Table 4).

After the fluorescein angiography was taken each eye was given a grading of 0 to 4 based on the fluorescence intensity between retinal vessels compared to background. A grading of 0 indicates fluorescence intensity between retinal vessels equivalent to background not induced by VEGF and represents minimal or no retinal leakage. A grading of 4 indicates fluorescence intensity equivalent to VEGF induced retinal leakage background and indicates heavy retinal leakage of fluorescein. In the 3 day VEGF induced retinal leakage assay Compound 6037 showed efficacy with a retinal leakage grading of 1 at 1 mg/eye dose (Figure 29). The 0.1 mg and 0.01 mg doses showed no efficacy.

Compounds 6037, 6044 and 6054 were evaluated in the 7 day rabbit leakage assay. Compound 6037 showed efficacy at 1 mg dose with a grading of 2 (Figure 30).

The invention thus has been disclosed broadly and illustrated in reference to representative embodiments described above. Those skilled in the art will recognize that various modifications can be made to the present invention without departing from the spirit and scope thereof. All publications, patent applications, and issued patents, are herein incorporated by reference to the same

extent as if each individual publication, patent application or issued patent were specifically and individually indicated to be incorporated by reference in its entirety. Definitions that are contained in text incorporated by reference are excluded to the extent that they contradict definitions in this disclosure.

5

It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination in a single embodiment. Conversely, various features of the invention which are, for brevity, described in the context of a single embodiment, may also be provided separately or in any suitable sub-combination.

10

It is specifically contemplated that any limitation discussed with respect to one embodiment of the invention may apply to any other embodiment of the invention. Furthermore, any composition of the invention may be used in any method of the invention, and any method of the invention may be used to produce or to utilize any composition of the invention. In particular, any aspect of the invention described in the claims, alone or in combination with one or more additional claims and/or aspects of the description, is to be understood as being combinable with other aspects of the invention set out elsewhere in the claims and/or description.

15

20

The use of the term "or" in the claims is used to mean "and/or" unless explicitly indicated to refer to alternatives only or the alternative are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and "and/or."

25

As used herein the specification, "a" or "an" may mean one or more, unless clearly indicated otherwise. As used herein in the claim (s), when used in conjunction with the word "comprising, "the words "a" or "an" may mean one or more than one. As used herein "another" may mean at least a second or more.

30

The words "comprises/comprising" and the words "having/including" when used herein with reference to the present invention are used to specify the presence of stated features, integers, steps or components but does not preclude the presence or addition of one or more other features, integers, steps, components or groups thereof.

TABLE 1: [VEGF-Peptide] Binding

SEQ ID NO:	VEGF Binding IC ₅₀ nM	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23
34	(188)	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L				
35	(325)	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	G	W
36	(172)	V	E	P	N	C	D	I	H	V	M	W	V	W	V	C	F	E	R	L	F	R	E	A
37	(59)	V	E	P	N	C	D	I	H	V	M	W	V	W	V	C	F	E	R	L	F	K	E	A
38	92.6	V	E	P	N	C	D	I	H	V	M	W	V	W	V	C	F	E	R	L	M	K		
39	(70)	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	R	E	L
40	78.3	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	I	F
41	69.9	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y			
42	68.2	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	W	G
43	66	V	E	P	N	C	D	I	H	V	M	W	V	W	V	C	F	E	R	L	Y	G	G	G
44	54.6	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	L	Y
45	51	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	S	G	G	G
46	44.1	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	M	R	L	T
47	43.3	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G		
48	42.4	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	V	K		
49	34.5	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	M	R		
50	34.4	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	I	L
51	32.8	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	YMe	G	L	T
52	32.3	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	G	
53	31.4	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	F	K	E	A
54	28.6	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Nle	K		
55	27	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	dL	Y	G	L	T
56	26.2	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	G	F

TABLE 1 Continued: VEGF-Peptidel Binding

SEQ ID NO:	VEGF Binding IC ₅₀ nM	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23
57	25.4	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	I	K		
58	25.1	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	K		
59	25	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	M	G	L	T
60	25	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	dL	Kac		
61	23.6	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	G	G
62	25	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	M	K		
63	24	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	M	R	E	L
64	24	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L				
65	18	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	M	K	E	L
66	18.7	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	P	W
67	10.1	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	L	K		
68	10	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	L	Kac		
69	9.4	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	L	T
70	9.3	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	E	F
71	5.4	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	L	S
72	4.8	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	L	
73	4.3	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	d-L	T
74	4	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	V	Q
75	3	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	L	E
76	2.3	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	P	L
77	1.8	V	E	P	N	C	D	I	H	V	M	W	V	W	E	C	F	E	R	L	Y	G	P	F

75

TABLE 2: [VEGF-Peptide] Binding

SEQ ID NO #	VEGF Binding IC ₅₀ nM	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23
34	(188)	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L				
108	362	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	A	K		
109	325	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	F	K	E	W
110	(188)	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	A	R		
111	265	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	F	K		
112	115	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	A	K	E	F
113	137	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	S	G	W	G
114	127	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	S	G	G	F
115	122	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	A	K	E	A
116	69	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	A	K	E	M
117	73	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	A	K	E	L
118	79	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	F	K	E	L
119	51	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	F	K	E	A
120	32	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	Y	G	G	G
121	26	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	M	K		

TABLE 3: VEGF-Peptide-Antibody Complex Binding, Half-life and AUC

Compound	VEGF Binding IC ₅₀ nM	Half life (Hrs)	AUC	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	
2077	147			V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	K	
2076	40.5			V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	K	L	
2073	5, 6	83	7x10 ⁵	V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	K	R	L	
2072	>50,000			V	E	P	N	C	D	I	H	V	M	W	E	W	E	C	K	E	R	L	
2075	3.2			V	E	P	N	C	D	I	H	V	M	W	E	W	K	C	F	E	R	L	
2074	625			V	E	P	N	C	D	I	H	V	M	W	E	K	E	C	F	E	R	L	
2071	1	43	2.8x10 ⁶	V	E	P	N	C	D	I	H	V	M	W	W	K	W	E	C	F	E	R	L
2070	1372.5			V	E	P	N	C	D	I	H	V	M	K	E	W	E	C	F	E	R	L	
2069	6	52	2.5x10 ⁶	V	E	P	N	C	D	I	H	V	K	W	E	W	E	C	F	E	R	L	
2068	9.3			V	E	P	N	C	D	I	H	K	M	W	E	W	E	C	F	E	R	L	
2067	166.8			V	E	P	N	C	D	I	K	V	M	W	E	W	E	C	F	E	R	L	
2066	423.5			V	E	P	N	C	D	K	H	V	M	W	E	W	E	C	F	E	R	L	
2065	220			V	E	P	N	C	K	I	H	V	M	W	E	W	E	C	F	E	R	L	
2064	4.5			V	E	P	K	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	
2063	2.5			V	E	K	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	
2062	2			V	K	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	
2028	1,25	32	1.8x10 ⁶	K	E	P	N	C	D	I	H	V	M	W	E	W	E	C	F	E	R	L	

TABLE 4: [VEGF-Peptide]-Antibody Complex:
Binding & Half-life- Linking Residue and X¹⁹-X²³ modifications

Cmpd #	SEQ ID NO:	VEGF Binding IC ₅₀ nM	Half Life(Hrs) VEGF [IgG] Coated	X1-X19 variations from SEQ ID NO:64 *=linking residue	19	20	21	22	23
2043	76			K1*	L	Y	G	P	L
2017	60	2.4	58	K1*	L	D-leu	Kac		
2020	79	0.5	25	K4*	L	R	D-leu	Kac	
2081	60	0.8		K4*	L	D-leu	Kac		
2082	60	2		K9*	L	D-leu	Kac		
2019	97	2.8		K10*	L				
2006	136	2		K10*, (Kac)12	L	Y	D-Ala	V	D-Gln
2013	80	2		K10*	L	D-Tyr	G	D-Pro	L
2025	60	1.65		K10*	L	D-leu	Kac		
2011	82	1.6		K10*	L	D-Tyr	G	D-leu	Kac
2012	81	1.5		K10*	L	D-Tyr	G	D-Pro	D-leu
2026	83	1.5		K10*	D-leu	Y	Aib	L	T
2007	85	1.1		K10*	D-leu	Y	D-Ala	V	D-Gln
2010	85	1		K10*	D-leu	Y	D-Ala	L	D-Thr
2016	86	0.8		K10*	L	Cha	G	D-leu	T
2027	87	0.7		K10*	L	Y	Aib	L	D-Thr
2009	88	0.7		K10*	L	Y	Sar	D-leu	T
2002	74	0.6		K10*	L	Y	G	V	Q
2003	74	0.6		K10*	L	Y	G	V	Q
2048	77	0.5	[23]	K10*	L	Y	G	P	F
2014	89	0.5		K10*	L	Y	G	D-Pro	L
2008	90	0.5		K10*	L	Y	D-Ala	V	D-Gln
2005	74	0.5		K10*	L	Y	G	V	Q
2031	91	0.45	[15]	K10*	L	Y	Aib	L	T
2004	74	0.4		K10*	L	Y	G	V	Q
2001	74	0.4		K10*	L	Y	G	V	Q
2049	73	0.4	[35]	K10*	L	Y	G	D-leu	T
2018	78	0.4	109	K10*	L	Y	D-Ala	D-leu	
2053	75	0.35	36	K10*	L	Y	G	L	E
2046	76	0.2	[19]	K10*	L	Y	G	P	L

TABLE 4 CONTINUED: [VEGF-Peptide]-Antibody Complex:
Binding & Half-life- Linking Residue and X¹⁹-X²³ modifications

Cmpd #	SEQ ID NO:	VEGF Binding IC ₅₀ nM	Half Life(Hrs) VEGF [IgG] Coated	X1-X19 variations from SEQ ID NO:64 *=linking residue	19	20	21	22	23
2022	84	0.7		K12*	D-leu	Y	D-Ala	V	D-Gln
2083	128	0.6		(Nle)10, K12*	L	Y	Aib	L	T
2028	83	0.5		K12*	D-leu	Y	Aib	L	T
2084	129	0.5		L10, K12*	L	Y	Aib	L	T
2036	60	0.45	[134]	K12*	L	D-leu	Kac		
2024	74	0.4		K12*	L	Y	G	V	Q
2085	130	0.4		V10, K12*	L	Y	Aib	L	T
2023	90	0.3		K12*	L	Y	D-Ala	V	D-Gln
2030	91	0.25	[37]	K12*	L	Y	Aib	L	T
2042	75	0.2		K12*	L	Y	G	L	E
2029	87	0.1		K12*	L	Y	Aib	L	D-Thr
2015	73	1.1		K14*	L	Y	G	D-leu	T
2033	92	9.4		K17*	L	D-Tyr	G	D-leu	T
2040	93	8.8		K17*	L	d-Tyr	G	L	T
2037	94	8.1		K17*	L	D-Tyr	D	L	D-Thr
2039	55	7.2		K17*	D-leu	Y	G	L	T
2035	60	6.3		K17*	L	D-leu	Kac		
2038	95	4.2, 4.2		K17*	L	D-leu	Aib		
2047	77	2.6		K17*	L	Y	G	P	F
2038	96	0.9		K17*	L	Y	G	D-leu	D-Thr
2050	73	0.6	[58]	K17*	L	Y	G	D-leu	T
2045	76	0.6	[50]	K17*	L	Y	G	P	L
2052	75	0.4	[38]	K17*	L	Y	G	L	E
2079				K17*	L	Y	G	L	D-Thr
2021		7.2		K21*	L	R	K	K	
2051	88	1.4	33	K21*	L	Y	K	D-leu	T
2041	77			K21*	L	Y	K	P	F
2080				K21*	L	Y	K	L	E
2032	73*	0.25	[<10]	K23*	L	Y	G	D-leu	K
2086	73*		8	K23*	L	Y	G	D-leu	K

TABLE 5:

[Ang2-Peptide] and [Ang2-Peptide]-Antibody-Complex: Binding and Half-life

Compound	Sequence showing position of linker (0P or 4P) and parent SEQ ID NO	Ang-2 T IC50 nM	Ang-2 P IC50 nM	T ^{1/2} (Sc) hours
4021	QKY QPL DEL DKT LYD QFM LQQ G (SEQ ID NO:137)	25.81		
4022	Q(Kac)Y QPL DEL DKT LYD QFM LQQ G (SEQ ID NO:138)	41.9		
4023	QNY QPL DEL DKT LYD QFM LQQ G (SEQ ID NO:139)	27.83		
4024	Q(Kac)Y QPL DEL DK(4P)T LYD QFM LQQ G (SEQ ID NO:138)	206.7	17.4	104 (27)
4025	QNY QPL DEL DK(4P)T LYD QFM LQQ G (SEQ ID NO:139)	300.3	40.456	
4026	(0P)QKY QPL DEL DKT LYD QFM LQQ G (SEQ ID NO:137)	32.3	58.77	(7)
4027	QKY QPL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:137)	29	55.876	117 (32)
4028	QNY QPL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:139)	175	28.63	100 (57)
4029	Q(Kac)Y QPL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:138)	83.3	13.195	77 (33)
4030	Q(Kac)Y QPL DEK(0P) D(Kac)T LYD QFM LQQ G (SEQ ID NO:140)	279	65.4	(17)
4031	Q(Kac)Y QPL DEL DET K(0P)YD QFM LQQ G (SEQ ID NO:141)	>5K	>1000	
4032	Q(Kac)Y QPL DEL D(Kac)T K(0P)YD QFM LQQ G (SEQ ID NO:142)	>5K	>1000	
4033	Q(Kac)Y QPL DEK(0P) DET LYD QFM LQQ G (SEQ ID NO:143)	369.6	88.81	
4034	Q(Kac)Y QD(HP)L DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:144)	224.5	19.75	(31)
4035	Q(Kac)Y Q(HP)L DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:145)	121	16.68	
4036	Q(Kac)Y QPL DEL DK(0P)T IYD QFM LQQ G (SEQ ID NO:146)	2410	152.9	
4037	Q(Kac)Y QPL DEI DK(0P)T LYD QFM LQQ G (SEQ ID NO:147)	1273	80.05	
4038	Q(Kac)Y QPL DEL DK(0P)T(HChA)YD QFM LQQ G (SEQ ID NO:148)	>5K	702.6	
4039	Q(Kac)Y QPL DEL DK(0P)T(HF)YD QFM LQQ G (SEQ ID NO:149)	4894	258.1	
4040	Q(Kac)Y QPL DEL DK(0P)T(ThA)YD QFM LQQ G (SEQ ID NO:150)	>5K	357.8	
4041	Q(Kac)Y QPL DEL DK(0P)T(Nva)YD QFM LQQ G (SEQ ID NO:151)	1339	23.32	(36)
4042	Q(Kac)Y QPL DEL DK(0P)T(HL)YD QFM LQQ G (SEQ ID NO:152)	1342	38.15	(42)

TABLE 5 CONTINUED:

[Ang2-Peptide] and [Ang2-Peptide]-Antibody-Complex: Binding and Half-life

Compound	Sequence showing position of linker (0P or 4P) and parent SEQ ID NO	Ang-2 T IC50	Ang-2 P IC50	T ^{1/2} (Sc) hours
4043	Q(Kac)Y QPL DE(Kac) DK(0P)T LYD QFM LQQ G (SEQ ID NO:153)	240	14.515	110 (38)
4044	Q(Kac)Y QPL DEL DK(0P)T LFD QFM LQQ G (SEQ ID NO:154)	36.9	11.68	58 (31)
4045	Q(Kac)Y Q(HP)L DE(ThA) DK(0P)T L(NO2)FD QFM LQQ G (SEQ ID NO:155)	24.7	12.7	68 (30)
4046	Q(Kac)Y QHPL DE(ThA) DK(0P)T L(BPA)D QFM LQQ G (SEQ ID NO:156)	82.7	25.21	(32)
4047	Q(Kac)Y Q(HP)L DE(ThA) DK(0P)T L(CO2H)FD QFM LQQ G (SEQ ID NO:157)	43.3	20.65	(47)
4048	Q(Kac)Y QPL DE(Kac) DK(0P)T L(NO2)FD QFM LQQ G (SEQ ID NO:158)	82.4	15.75	64 (35)
4049	(DCB)Q(Kac)Y QPL DEL DK(0P)T LYD QFMLQQ G (SEQ ID NO:159)	28.4	12.45	(24)
4050	(DFB)Q(Kac)Y QPL DEL DK(0P)T LYD QFMLQQ G (SEQ ID NO:160)	33	13.56	(23)
4051	(PyC)Q(Kac)Y QPL DEL DK(0P)T LYD QFMLQQ G (SEQ ID NO:161)	14.3	19.38	(18)
4052	(Amido 2-PEG)Q(Kac)Y QPL DEL DK(0P)T LYDQFMLQQ G (SEQ ID NO:162)	133.8	18.14	(31)
4053	Q(CIBnCarbamate)KY QPL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:163)	146.9		
4054	Q(Kac)Y QPL DEL D(Dab)(0P)T LYD QFM LQQ G (SEQ ID NO:164)	291.7	22	
4055	Q(Kac)Y QPL DEL D(Dap)(0P)T LYD QFM LQQ G (SEQ ID NO:165)	598	34.54	
4056	QNY QPL DEL DK(0P)T L(BPA)D QFM LQQ G (SEQ ID NO:166)	92.2	17.2	(29)
4057	QNY QPL DEL DK(0P)T L(CF)D QFM LQQ G (SEQ ID NO:167)	110	15.3	(11)
4058	QNY QPL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:139)	36.9	12.9	(24)
4059	QRY QPL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:168)	176.8	16.1	
4060	QHY QPL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:169)	214	14.4	
4061	Q(Nick)Y QPL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:170)	83.9		
4062	Q(Kac)Y QPL DE(Kac) DK(0P)T L(CF)D QFM LQQ G (SEQ ID NO:171)	499	41.3	
4063	Q(Kac)Y QPL DE(Kac) DK(0P)T LFD QFM LQQ G (SEQ ID NO:153)			

TABLE 6:

[Ang2-Peptide]-Antibody-Complex: Binding and Half-life

Compound No.	Sequence	Ang-2 P IC50 nM	Ang-2 T $\frac{1}{2}$ Hours
4021	QKY QPL DEL DKT LYD QFM LQQ G (SEQ ID NO:137)	36.39	
4022	Q(Kac)Y QPL DEL DKT LYD QFM LQQ G (SEQ ID NO:138)	14.41	
4027	QKY QPL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:137)	0.59	
4028	QNY QPL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:139)	0.15	
4029	Q(Kac)Y QPL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:138)	0.41	
4030	Q(Kac)Y QPL DEK(0P) D(Kac)T LYD QFM LQQ G (SEQ ID NO:140)	1.27	72
4032	Q(Kac)Y QPL DEL D(Kac)T K(0P)YD QFM LQQ G (SEQ ID NO:142)	>100	
4043	Q(Kac)Y QPL DE(Kac) DK(0P)T LYD QFM LQQ G (SEQ ID NO:153)	0.92	
4044	Q(Kac)Y QPL DEL DK(0P)T LFD QFM LQQ G (SEQ ID NO:154)	0.41	
4048	Q(Kac)Y QPL DE(Kac) DK(0P)T L(NO2F)D QFM LQQ G (SEQ ID NO:158)	0.33	
4054	Q(Kac)Y QPL DEL D(Dab)(0P)T LYD QFM LQQ G (SEQ ID NO:164)	0.09	
4055	Q(Kac)Y QPL DEL D(Dap)(0P)T LYD QFM LQQ G (SEQ ID NO:165)	1.61	
4064	K(0P)(Kac)Y QPL DEL D(Kac)T LYD QFM LQQ G (SEQ ID NO:172)	1.92	24
4065	QK(0P)Y QPL DEL D(Kac)T LYD QFM LQQ G (SEQ ID NO:173)	0.31	12
4066	Q(Kac)K(0P) QPL DEL D(Kac)T LYD QFM LQQ G (SEQ ID NO:174)	0.23	17
4067	Q(Kac)Y K(0P)PL DEL D(Kac)T LYD QFM LQQ G (SEQ ID NO:175)	N.I.	
4068	Q(Kac)Y QK(0P)L DEL D(Kac)T LYD QFM LQQ G (SEQ ID NO:176)	44.21	
4069	Q(Kac)Y QPK(0P) DEL D(Kac)T LYD QFM LQQ G (SEQ ID NO:177)	N.I.	
4070	Q(Kac)Y QPL K(0P)EL D(Kac)T LYD QFM LQQ G (SEQ ID NO:178)	N.I.	
4071	Q(Kac)Y QPL DK(0P)L D(Kac)T LYD QFM LQQ G (SEQ ID NO:179)	0.288	35
4072	Q(Kac)Y QPL DEL K(0P)(Kac)T LYD QFM LQQ G (SEQ ID NO:180)	N.I.	
4073	Q(Kac)Y QPL DEL D(Kac)K(0P) LYD QFM LQQ G (SEQ ID NO:181)	0.11	56

TABLE 6 CONTINUED:

[Ang2-Peptide]-Antibody-Complex: Binding and Half-life

4074	Q(Kac)Y QPL DEL D(Kac)T LK(0P)D QFM LQQ G (SEQ ID NO:182)	32.82	
4075	Q(Kac)Y QPL DEL D(Kac)T LYK(0P) QFM LQQ G (SEQ ID NO:183)	0.19	65
4076	Q(Kac)Y QPL DEL D(Kac)T LYD K(0P)FM LQQ G (SEQ ID NO:184)	0.27	94
4077	Q(Kac)Y QPL DEL D(Kac)T LYD QK(0P)M LQQ G (SEQ ID NO:185)	19.53	
4078	Q(Kac)Y QPL DEL D(Kac)T LYD QFK(0P) LQQ G (SEQ ID NO:186)	0.74	72
4079	Q(Kac)Y QPL DEL D(Kac)T LYD QFM K(0P)QQ G (SEQ ID NO:187)	0.077	65
4080	Q(Kac)Y QPL DEL D(Kac)T LYD QFM LK(0P)Q G (SEQ ID NO:188)	0.11	35
4081	Q(Kac)Y QPL DEL D(Kac)T LYD QFM LQK(0P) G (SEQ ID NO:189)	0.27	27
4082	Q(Kac)Y QPL DEL D(Kac)T LYD QFM LQQ K(0P) (SEQ ID NO:190)	0.21	20

TABLE 7:

Compound	Sequence	%TC 10 mg/kg 1x/wk	%TC 3 mg/kg 1x/wk	%TC 1 mg/kg 1x/wk
4027	QKY QPL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:137)	52		
4028	QNY QPL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:139)	38	34	
4029	Q(Kac)Y QPL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:138)	60	28	9
4043	Q(Kac)Y QPL DE(Kac) DK(0P)T LYD QFM LQQ G (SEQ ID NO:153)	53	27	34
4044	Q(Kac)Y QPL DEL DK(0P)T LFD QFM LQQ G (SEQ ID NO:154)	40	22	
4045	Q(Kac)Y Q(HP)L DE(ThA) DK(0P)T L(NO2)FD QFM LQQ G (SEQ ID NO:155)	47		
4048	Q(Kac)Y QPL DE(Kac) DK(0P)T L(NO2)FD QFM LQQ G (SEQ ID NO:158)	45		
4058	QNY QPL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:39)	51		
4092	PL DEL DK(0P)T LYD QFM LQQ G (SEQ ID NO:191)	60		

TABLE 8

	VEGF Part				Ang-2 Part			
Compound No:	Tethering Spot	PEG Length from Branch	VEGF Binding IC50	β Half Life (hrs)	Tethering Spot	PEG Length From Branch	Ang-2 Binding IC50	β Half Life (hrs)
6001	1	4	4		1	0		
6002	1	2			1	0		
6003	1	2			1	0		
6004	1	N/A			C-Ter	N/A	0.5	
6005	1	N/A			C-Ter	N/A	1	
6006	10	0	71		11	2	0.2	
6007	10	0	138		11	0	0.3	
6008	10	2	79		11	0	0.4	
6009	10	2	39		11	2	0.2	
6010	10	0	344		11	0		
6011	12	0	84		11	2	0.2	
6012	10	0	278		11	0	0.6	
6013	17	0	2	67, 91	11	2	0.3	40, 34
6014	10	0	240		11	2	0.1	
6015	1	0	2.8		1	0	0.1	
6016	1	0	6, 8		1	0	0.3	
6017	10	4	26		11	4	0.2	
6018	12	4	24		11	4	0.2	
6019	17	4	1.3	68	11	4	0.13	
6020	1	0	356		11	0	0.8	
6021	10	0	75		11	4	0.3	
6022	12	4	120, 140		18	4	0.5	
6023	10	4	60		18	4	0.3	
6024	10	4	5584		11	4	0.1	
6025	10	4	210, 227		11	4	0.1	
6026	1	4	553, 1129		11	4	0.2	
6027	17	0	2	65	11	0	0.3	32
6028	17	4	4		11	4	0.2, 0.3	
6031	17	0			11	4		
6032	1	0			11	0		
6033	1	0			11	4		
6034	17	0	0.6	42	11	0	1.3	39
6035	1	4			11	4		
6036	17	4	0.5	27	11	4	0.4	41
6037	10	4	0.6		11	4	1.6	
6038	17	0	0.5		11	4	0.8	
6039	17	4	0.6		11	4	1.1	
6040	17	0	0.4		11	0	2.5	








Compound No:	VEGF-Part				Ang2-Part			
	Tethering Spot	PEG Length from Branch	VEGF Binding IC50	β Half Life (hrs)	Tethering Spot	PEG Length From Branch	Ang-2 Binding IC50	β Half Life (hrs)
6041	12	4	2.2		11	4	0.8	
6042	12	4	3.3	56	11	4	0.8	49
6043	10	4	0.6	36	11	4	0.4	72
6044	10	4	0.7	43	11	4	0.9	82
6045	10	0	1.8	52	11	0	1.2	47
6046	12	0	3.3		11	0	1	
6047	10	4	2.5		11	4	0.5	
6048	12	4	4.4		11	4	0.7	
6049	12	4	45		11	0	0.6	
6050	12	4	33		11	4	0.6	
6051	12	4	Not Active		11	4	0.7	
6052	10	4	18		11	4	0.5	
6053	10	4	0.7	68	11	4	0.6	68
6054	1	4	0.3	18	11	4	0.4	
6055	1	4			11	4		
6056	12	4	94, 64		18	4	0.8	
6057	12	0	32, 44		11	12	0.7	
6058	12	12	18		11	0	0.7	
6059	12		15		11		0.6	
6060	10	4	1.2, 1.0		11	0		
6061	10	0	2.9		11	0		
6062	10	0	2.3, 2.2		11	4		
6063	10	4			11	4		
6064	10	4			11	4		
6065	12	4	36		1	0		
6066	4	0	3.1, 7.4		11	4		
6067	1	0	0.6, 0.4		11	4		
6068	9	4	22, 12		11	4		
6070	10	4	1	52(rat)	11	4	0.07	71(rat)
6071	10	4	0.8	66(rat)	11	4	0.05	61(rat)
6072	10	4	1.4	41(rat)	11	4	0.1	74(rat)
6073	10	4	0.6	80(rat)	11	4	0.08	91(rat)

TABLE 9:
[VEGF-Peptide]-Linker Compounds

SEQ ID	Compound #	Structure
74	1001	
74	1002	
74	1003	
74	1004	
74	1005	
90	1006	
84	1007	

90	1008	
88	1009	
85	1010	
82	1011	
81	1012	
80	1013	
89	1014	

73	1015	
86	1016	
60	1017	
78	1018	
97	1019	
79	1020	
196	1021	

84	1022	$\text{Ac-V-E-P-N-C-D-I-H-V-M-W-K-W-E-C-F-E-R-I-Y-a-V-q-CONH}_2$ 
90	1023	$\text{Ac-V-E-P-N-C-D-I-H-V-M-W-K-W-E-C-F-E-R-L-Y-a-V-q-CONH}_2$ 
74	1024	$\text{Ac-V-E-P-N-C-D-I-H-V-M-W-K-W-E-C-F-E-R-L-Y-G-V-Q-CONH}_2$ 
60	1025	$\text{Ac-V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-R-L-I-K-CONH}_2$ 
83	1026	$\text{Ac-V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-R-I-Y-N-CH(CH}_3\text{)-L-T-CONH}_2$ 
87	1027	$\text{Ac-V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-R-L-Y-N-CH(CH}_3\text{)-L-T-CONH}_2$ 
83	1028	$\text{Ac-V-E-P-N-C-D-I-H-V-M-W-K-W-V-W-E-C-F-E-R-I-Y-N-CH(CH}_3\text{)-L-T-CONH}_2$ 

87	1029	
91	1030	
91	1031	
197	1032	
92	1033	
95	1034	
60	1035	

60	1036	
94	1037	
96	1038	
55	1039	
93	1040	
77	1041	
75	1042	

76	1043	
76	1044	
76	1045	
76	1046	
77	1047	
77	1048	
73	1049	
73	1050	

73	1051	
75	1052	
75	1053	
198	1054	
199	1055	
200	1056	
201	1057	

202	1058	
203	1059	
204	1060	
205	1061	
34	1062	
34	1063	

34	1064	
34	1065	
34	1066	
34	1067	
34	1068	
34	1069	
34	1070	
34	1071	





34	1072	
34	1073	
34	1074	
34	1075	
34	1076	
34	1077	
34	1078	
206	1079	

207	1080	Chemical structure diagram showing a peptide sequence: V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-K-L-E-CONH₂. The sequence is flanked by two sulfur atoms (S) connected by a horizontal line. Below the sequence, a chemical structure is shown, featuring a benzene ring substituted with a cyclopropyl group, a carbonyl group, and a side chain containing an amide and an ester linkage.
-----	------	--

TABLE 10:
[VEGF-Peptide]-[Linker]-Antibody compounds






Incorporates SEQ ID #	Compound #	Structure
74	2001	Structure of compound 2001: A VEGF-peptide chain (VEPNCDIHVK) linked via a disulfide bond to a linker chain (WVWECFERLYGVQ). The linker chain is further linked to an antibody via a complex linker structure. The structure is shown in brackets with a subscript '1 or 2'.
74	2002	Structure of compound 2002: A VEGF-peptide chain (VEPNCDIHVK) linked via a disulfide bond to a linker chain (WVWECFERLYGVQ). The linker chain is further linked to an antibody via a complex linker structure. The structure is shown in brackets with a subscript '1 or 2'.
74	2003	Structure of compound 2003: A VEGF-peptide chain (VEPNCDIHVK) linked via a disulfide bond to a linker chain (WVWECFERLYGVQ). The linker chain is further linked to an antibody via a complex linker structure. The structure is shown in brackets with a subscript '1 or 2'.
74	2004	Structure of compound 2004: A VEGF-peptide chain (VEPNCDIHVK) linked via a disulfide bond to a linker chain (WVWECFERLYGVQ). The linker chain is further linked to an antibody via a complex linker structure. The structure is shown in brackets with a subscript '1 or 2'.

74	2005	
90	2006	
84	2007	
90	2008	
88	2009	

85	2010	 Chemical structure of antibody 85. The structure shows a disulfide bridge (S-S) connecting two polypeptide chains. The first chain is VEPNCDIHVK, and the second chain is WVWECFERIY a L q. The structure is shown in brackets, indicating it is a repeating unit. The structure is labeled "Antibody" and "1 or 2".
82	2011	 Chemical structure of antibody 82. The structure shows a disulfide bridge (S-S) connecting two polypeptide chains. The first chain is VEPNCDIHVK, and the second chain is WVWECFERL y G I K. The structure is shown in brackets, indicating it is a repeating unit. The structure is labeled "Antibody" and "1 or 2".
81	2012	 Chemical structure of antibody 81. The structure shows a disulfide bridge (S-S) connecting two polypeptide chains. The first chain is VEPNCDIHVK, and the second chain is WVWECFERL y G I T. The structure is shown in brackets, indicating it is a repeating unit. The structure is labeled "Antibody" and "1 or 2".
80	2013	 Chemical structure of antibody 80. The structure shows a disulfide bridge (S-S) connecting two polypeptide chains. The first chain is VEPNCDIHVK, and the second chain is WVWECFERL y G p L. The structure is shown in brackets, indicating it is a repeating unit. The structure is labeled "Antibody" and "1 or 2".

97	2019	
79	2020	
196	2021	
84	2022	

90	2023	
74	2024	
60	2025	
83	2026	
87	2027	

92	2033	 Antibody 1 or 2
95	2034	 Antibody 1 or 2
60	2035	 Antibody 1 or 2
60 192 (specifying K ¹² as linking residue)	2036	 Antibody 1 or 2
94	2037	 Antibody 1 or 2

96	2038	
55	2039	
93	2040	
77	2041	
75	2042	

76	2043	
76	2044	
76 194 (specifying K ¹⁷ as linking residue)	2045	
76	2046	
77	2047	

202	2058	
203	2059	
204	2060	
205	2061	
34	2062	

34	2063	
34	2064	
34	2065	
34	2066	
34	2067	

34	2068	Antibody 1 or 2
34	2069	Antibody 1 or 2
34	2070	Antibody 1 or 2
34	2071	Antibody 1 or 2
34	2072	Antibody 1 or 2

34	2073	
34	2074	
34	2075	
34	2076	
34	2077	
34	2078	



206	2079	
207	2080	





TABLE 11:**Compounds of the invention of the formula: [Ang2-Peptide]-[Connector]-[VEGF-Peptide]**

Cmpd #	Chemical Structure
5001 VEGF SEQ:34	Chemical structure of compound 5001, showing a VEGF peptide (V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-CONH₂) linked via a disulfide bond to a connector peptide (H₂NOC-G-Q-Q-L-I-F-Q-E-Y-L-R-Q-E-L-D-D-M-P-M-F-N-T-CONH₂).
5002 VEGF SEQ:34	Chemical structure of compound 5002, showing a VEGF peptide (V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-CONH₂) linked via a disulfide bond to a connector peptide (H₂NOC-G-Q-Q-L-I-F-Q-E-Y-L-R-Q-E-L-D-D-M-P-M-F-N-T-CONH₂).
5003 VEGF SEQ:34 P3,N4 are wrongly shown as N3,P4	Chemical structure of compound 5003, showing a VEGF peptide (V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-CONH₂) linked via a disulfide bond to a connector peptide (H₂NOC-G-Q-Q-L-I-F-Q-E-Y-L-R-Q-E-L-D-D-M-P-M-F-N-T-CONH₂).
5004 VEGF SEQ:34	Chemical structure of compound 5004, showing a VEGF peptide (V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-CONH₂) linked via a disulfide bond to a connector peptide (H₂NOC-G-Q-Q-L-I-F-Q-E-Y-L-R-Q-E-L-D-D-M-P-M-F-N-T-CONH₂).
5005 VEGF SEQ:34	Chemical structure of compound 5005, showing a VEGF peptide (V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-CONH₂) linked via a disulfide bond to a connector peptide (H₂NOC-G-Q-Q-L-I-F-Q-E-Y-L-R-Q-E-L-D-D-M-P-M-F-N-T-CONH₂).
5006 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	Chemical structure of compound 5006, showing a VEGF peptide (V-E-P-N-C-D-I-H-V-M-W-E-W-E-C-F-E-R-L-CONH₂) linked via a disulfide bond to a connector peptide (H₂NOC-G-Q-Q-L-I-F-Q-E-Y-L-R-Q-E-L-D-D-M-P-M-F-N-T-CONH₂).

5007 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	
5008 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	
5009 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	
5010 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	
5011 VEGF SEQ:34 E ¹² subs with K as linking residue	
5012 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	

5013 VEGF SEQ:34 E ¹⁷ subs with K as linking residue	
5014 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	
5015 VEGF SEQ:34	
5016 VEGF SEQ:34	
5017 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	
5018 VEGF SEQ:34 E ¹² subs with K as linking residue	





5019 VEGF SEQ:34 E ¹⁷ subs with K as linking residue	
5020 VEGF SEQ:34	
5021 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	
5022 VEGF SEQ:34 E ¹² subs with K as linking residue	
5023 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	

5024 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	
5025 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	
5026 VEGF SEQ:208	
5027 VEGF SEQ:34 E ¹⁷ subs with K as linking residue	
5028 VEGF SEQ:34 E ¹⁷ subs with K as linking residue	

5031 VEGF SEQ:209	
5032 VEGF SEQ:76	
5033 VEGF SEQ:76	
5034 VEGF SEQ:73 E ¹⁷ subs with K as linking residue	
5035 VEGF SEQ:76	

5036 VEGF SEQ:73 E ¹⁷ subs with K as linking residue	
5037 VEGF SEQ:75 M ¹⁰ subs with K as linking residue	
5038 VEGF SEQ:75 E ¹⁷ subs with K as linking residue	
5039 VEGF SEQ:75 E ¹⁷ subs with K as linking residue	
5040 VEGF SEQ:96 E ¹⁷ subs with K as linking residue	

5041 VEGF SEQ:211	
5042 VEGF SEQ:73 V ¹² subs with K as linking residue	
5043 VEGF SEQ:212	
5044 VEGF SEQ:212	

5045 VEGF SEQ:73 M ¹⁰ subs with K as linking residue	
5046 VEGF SEQ:73 V ¹² subs with K as linking residue	
5047 VEGF SEQ:213	
5048 VEGF SEQ:214	

5049 VEGF SEQ:215	
5050 VEGF SEQ:79 V ¹² subs with K as linking residue	
5051 VEGF SEQ:79 V ¹² subs with K as linking residue	
5052 VEGF SEQ:79 V ¹² subs with K as linking residue	

5053 VEGF SEQ:78 M ¹⁰ subs with K as linking residue	
5054 VEGF SEQ:79 V ¹ subs with K as linking residue	
5055 VEGF SEQ:79 V ¹ subs with K as linking residue	
5056 VEGF SEQ:79 V ¹² subs with K as linking residue	
5057 VEGF SEQ:79 V ¹² subs with K as linking residue	

5058 VEGF SEQ:79 V¹² subs with K as linking residue	
5059 VEGF SEQ:79 V¹² subs with K as linking residue	
5060 VEGF SEQ:78 M¹⁰ subs with K as linking residue	
5061 VEGF SEQ:78 M¹⁰ subs with K as linking residue	
5062 VEGF SEQ:78 M¹⁰ subs with K as linking residue	

5063 VEGF SEQ:89 M ¹⁰ subs with K as linking residue	
5064 VEGF SEQ:90 M ¹⁰ subs with K as linking residue	
5070 VEGF SEQ:216	
5071 VEGF SEQ:217	
5072 VEGF SEQ:218	
5073 VEGF SEQ:210	

5074
VEGF
SEQ:73













TABLE 12
{{[Ang2-Peptide]-[Connector]-[VEGF-Peptide]}_{1 or 2}}-[Antibody]

Cmpd #	Chemical Structure
6001 VEGF SEQ:34	Chemical structure of compound 6001: A VEGF peptide (V E P N C D I H V M W E W E C F E R L - CONH₂) linked via a disulfide bond to a connector peptide (H₂NOC-G Q Q L I F Q E Y L R Q E L D D M P M F N T - CO). The connector is further linked to an antibody via a complex linker system including a benzyl group and a cyclic amide. The antibody is labeled "1 or 2ibody".
6002 VEGF SEQ:34	Chemical structure of compound 6002: Similar to 6001, but the connector peptide is H₂NOC-G Q Q L I F Q E Y L R Q E L D D M P M F N t - CO, and the linker system is modified with a different cyclic amide and a benzyl group. The antibody is labeled "1 or 2".
6003 VEGF SEQ:34	Chemical structure of compound 6003: The VEGF peptide and connector peptide are linked via a disulfide bond. The connector is linked to an antibody via a complex linker system including a benzyl group and a cyclic amide. The antibody is labeled "1 or 2".
6004 VEGF SEQ:34	Chemical structure of compound 6004: A VEGF peptide (V E P N C D I H V M W E W E C F E R L - CONH₂) linked via a disulfide bond to a connector peptide (H₂NOC-L R E F C E W E W M V H I D C N P E V - CO). The connector is further linked to an antibody via a complex linker system including a benzyl group and a cyclic amide.
6005 VEGF SEQ:34	Chemical structure of compound 6005: A VEGF peptide (V E P N C D I H V M W E W E C F E R L - CONH₂) linked via a disulfide bond to a connector peptide (H₂NOC-L R E F C E W E W M V H I D C N P E V - CO). The connector is further linked to an antibody via a complex linker system including a benzyl group and a cyclic amide.

6006 VEGF SEQ:34 M¹⁰ subs with K as linking residue	
6007 VEGF SEQ:34 M¹⁰ subs with K as linking residue	
6008 VEGF SEQ:34 M¹⁰ subs with K as linking residue	
6009 VEGF SEQ:34 M¹⁰ subs with K as linking residue	
6010 VEGF SEQ:34 M¹⁰ subs with K as linking residue	

6011 VEGF SEQ:34 E ¹² subs with K as linking residue	1 or 2
6012 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	1 or 2
6013 VEGF SEQ:34 E ¹⁷ subs with K as linking residue	1 or 2
6014 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	1 or 2
6015 VEGF SEQ:34	1 or 2

6016 VEGF SEQ:34	
6017 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	
6018 VEGF SEQ:34 E ¹² subs with K as linking residue	
6019 VEGF SEQ:34 E ¹⁷ subs with K as linking residue	
6020 VEGF SEQ:34	





6021 VEGF SEQ:34 M¹⁰ subs with K as linking residue	 Chemical structure of VEGF SEQ:34 M¹⁰ subunit. The sequence is V E P N C D I H V K W E W E C F E R L-CONH₂. A linker is attached to the Cysteine (C) residue, which is part of a disulfide bridge. The linker consists of a series of ether linkages and amide bonds, ending in a phenyl ring. The phenyl ring is attached to an antibody via a linker. The antibody is shown as a bracketed structure labeled "Antibody". The linker is labeled "1 or 2".
6022 VEGF SEQ:34 E¹² subs with K as linking residue	 Chemical structure of VEGF SEQ:34 E¹² subunit. The sequence is V E P N C D I H V M W K W E C F E R L-CONH₂. A linker is attached to the Cysteine (C) residue, which is part of a disulfide bridge. The linker consists of a series of ether linkages and amide bonds, ending in a phenyl ring. The phenyl ring is attached to an antibody via a linker. The antibody is shown as a bracketed structure labeled "Antibody". The linker is labeled "1 or 2".
6023 VEGF SEQ:34 M¹⁰ subs with K as linking residue	 Chemical structure of VEGF SEQ:34 M¹⁰ subunit. The sequence is V-E-P-N-C-D-I-H-V-K-W-E-W-E-C-F-E-R-L-CONH₂. A linker is attached to the Cysteine (C) residue, which is part of a disulfide bridge. The linker consists of a series of ether linkages and amide bonds, ending in a phenyl ring. The phenyl ring is attached to an antibody via a linker. The antibody is shown as a bracketed structure labeled "Antibody". The linker is labeled "1 or 2".
6024 VEGF SEQ:34 M¹⁰ subs with K as linking residue	 Chemical structure of VEGF SEQ:34 M¹⁰ subunit. The sequence is H₂NOC-L-R-E-F-C-E-W-E-W-K-V-H-I-D-C-N-P-E-V. A linker is attached to the Cysteine (C) residue, which is part of a disulfide bridge. The linker consists of a series of ether linkages and amide bonds, ending in a phenyl ring. The phenyl ring is attached to an antibody via a linker. The antibody is shown as a bracketed structure labeled "Antibody". The linker is labeled "1 or 2".

6025 VEGF SEQ:34 M ¹⁰ subs with K as linking residue	
6026 VEGF SEQ:208	
6029 VEGF SEQ:34 E ¹⁷ subs with K as linking residue- (analogous to 5027)	
6028 VEGF SEQ:34 E ¹⁷ subs with K as linking residue	
6031 VEGF SEQ:209	

6032 VEGF SEQ:76	
6033 VEGF SEQ:76	
6034 VEGF SEQ:73 E ¹⁷ subs with K as linking residue	
6035 VEGF SEQ:76	





6040 SEQ:96 E ¹⁷ subs with K as linking residue	
6041 SEQ:211	
6042 SEQ:73 V ¹² subs with K as linking residue	
6043 SEQ:212	
6044 SEQ:212	





6049 SEQ:215	
6050 SEQ:79 V^{12} subs with K as linking residue	
6051 SEQ:79 V^{12} subs with K as linking residue	
6052 SEQ:79 V^{12} subs with K as linking residue	

6053 SEQ:78 M¹⁰ subs with K as linking residue	
6054 SEQ:79 V¹ subs with K as linking residue	
6055 SEQ:79 V¹ subs with K as linking residue	
6056 SEQ:79 V¹² subs with K as linking residue	

6057 SEQ:79 V¹² subs with K as linking residue	
6058 SEQ:79 V¹² subs with K as linking residue	
6059 SEQ:73 V¹² subs with K as linking residue	
6060 SEQ:78 M¹⁰ subs with K as linking residue	

6061 SEQ:78 M¹⁰ subs with K as linking residue	
6062 SEQ:78 M¹⁰ subs with K as linking residue	
6063 SEQ:89 M¹⁰ subs with K as linking residue	
6064 SEQ:90 M¹⁰ subs with K as linking residue	

6065 SEQ:79 V¹² subs with K as linking residue	
6066 SEQ:79 N⁴ subs with K as linking residue	
6067 SEQ:79 V¹ subs with K as linking residue	
6068 SEQ:79 V⁹ subs with K as linking residue	

6069 SEQ:60 V¹² subs with K as linking residue	
6070 SEQ:216	
6071 SEQ:217	
6072 SEQ:218	



6073
SEQ:2106074
SEQ:73

TABLE 13**Comparison of Compounds 2018, 4043, and 6053**

	IC50 (nM)		Mouse T1/2 (hr)		Rat T1/2 (hr)	
	VEGF	Ang2	VEGF	Ang2	VEGF	Ang2
Cmpd 2018	0.7		109		72	
Cmpd 4043		0.6		~90		95
Cmpd 6053	0.7	0.6	92	92	97	111

5

TABLE 14**Comparison of Compounds 2018, 4043, and 6053**

	IC50 (nM)		Mouse T1/2 (hr)	
	VEGF	Ang2	VEGF	Ang2
Cmpd 6054	0.3	0.4	18	
Cmpd 6037	0.6	1.6	[~ 36]	
Cmpd 6044	0.7	0.9	43	82

Claims

1. A compound of the formula: R^1 -[VEGF-Peptide]- R^2 wherein [VEGF-Peptide] is a peptide having the sequence:

X^1 - X^2 - P^3 - N^4 - C^5 - X^6 - X^7 - X^8 - V^9 - X^{10} - X^{11} - X^{12} - W^{13} - X^{14} - C^{15} - F^{16} - E^{17} - R^{18} - X^{19} - X^{20} - X^{21} - X^{22} - X^{23}


(SEQ ID NO:131) wherein R^1 is absent, CH_3 , $C(O)CH_3$, $C(O)CH_2CH_3$, $C(O)CH_2CH_2CH_3$, $C(O)CH(CH_3)CH_3$, $C(O)CH_2CH_2CH_2CH_3$, $C(O)CH(CH_3)CH_2CH_3$, $C(O)C_6H_5$, $C(O)CH_2CH_2(CH_2CH_2O)_{1-5}Me$, amido-2-PEG, or an N-acyl or N-alkyl amino protecting group, a lipid fatty acid group or a carbohydrate; and R^2 is absent, OH, NH_2 , $NH(CH_3)$, $NHCH_2CH_3$, $NHCH_2CH_2CH_3$, $NHCH(CH_3)CH_3$, $NHCH_2CH_2CH_2CH_3$, $NHCH(CH_3)CH_2CH_3$, NHC_6H_5 , $NHCH_2CH_2OCH_3$, $NHOCH_3$, $NHOCH_2CH_3$, a carboxy protecting group, a lipid fatty acid group or a carbohydrate, and X^1 is a hydrophobic amino acid residue, X^2 is a negatively charged residue, X^6 is a negatively charged residue, X^7 is a hydrophobic amino acid residue, X^8 is a residue comprising a ring structure, X^{10} may be a hydrophobic amino acid, X^{11} is an aromatic amino acid, X^{12} is selected from the group consisting of V, E and Kac, X^{14} is E or V, X^{19} may be any hydrophobic amino acid residue, or D-isomer thereof, X^{20} may be absent, or may be any neutral, hydrophobic or aromatic amino acid or D-isomer thereof, X^{21} may be absent, or may be any positively charged residue or any aliphatic non-polar residue or D-isomer thereof, X^{22} may be absent, or may be selected from the group consisting of G, V, L, I, P, S, T, W, F, E, Kac, or D-isomers thereof, X^{23} may be absent, or is selected from the group consisting of G, A, I, L, Q, E, F, T, W, S, Y, and Kac and D-isomers thereof, and wherein one of the group consisting of, X^1 , X^2 , P^3 , N^4 , V^9 , X^{10} , X^{12} , X^{14} , E^{17} and the C-terminus residue may substituted with a linking residue comprising a nucleophilic side chain or the N-terminus amino group or C-terminus carboxyl group covalently linked to the combining site of an antibody directly or via an intermediate linker, the linking residue being selected from the group consisting of K, R, Y, C, T, S, homologs of lysine, homocysteine, homoserine, Dap, Dab, the N-terminus residue and the C-terminus residue, or a pharmaceutically acceptable salt thereof.

2. A compound as claimed in claim 1, wherein the [VEGF-Peptide] includes a sequence substantially homologous to one or more selected from the group consisting of SEQ ID NOs: 34-136, and SEQ ID NOs:192-195.


3. A compound as claimed in any preceding claim, wherein the [VEGF-Peptide] includes a sequence substantially homologous to:

V-E-P-N-C-D-I-H-V-M-W-V-W-E-C-F-E-R-L-Y-(D-Ala)-(D-Leu) (SEQ ID NO:78).


4. A compound as claimed in claim 3, wherein M¹⁰ is substituted with the linking residue K, comprising the formula (SEQ ID NO:195):
 $\{C(O)CH_3\}-V-E-P-N-C-D-I-H-V-K-W-V-W-E-C-F-E-R-L-Y-(D-Ala)-(D-Leu)-\{NH_2\}$.
5. A compound as claimed in preceding claim, wherein the linking residue is present, and further comprising a linear or branched linker covalently linked to the side chain of the linking residue.
6. A compound as claimed in claim 5, wherein the linker comprises the formula L or L', and L is of the formula: $-[Connector]-X-Y-Z$, or $-X-Y-Z-$, and L' is of the formula: $-[Connector]-X-Y-Z'$, or $-X-Y-Z'-$, wherein: [Connector] is present where the linker is branched, and where present is covalently linked to the linking residue, and one or more additional Active Molecules, X is a biologically compatible connecting chain including any atom selected from the group consisting of C, H, N, O, P, S, F, CL, Br, and I, and may comprise a polymer or block co-polymer, and is covalently linked to the linking residue where the linker is linear, Y is present, and has the optionally substituted structure:



wherein a, b, c, d, and e are independently carbon or nitrogen; f is carbon, nitrogen, oxygen, or sulfur; Y is attached to X and Z independently at any two ring positions of sufficient valence; and no more than four of a, b, c, d, e, or f are simultaneously nitrogen and preferably a, b, c, d, and e in the ring structure are each carbon, and, Z if present has the structure:





and wherein Z' if present has the structure:




wherein q=0-5 and Antibody-N- if present is a covalent bond to a side chain in a combining site of an antibody.

7. A compound as claimed in claims 6, wherein:
 X is:



wherein v and w are selected such that the backbone length of X is 6-12 atoms, and R^b is hydrogen, substituted or unsubstituted C₁₋₁₀ alkyl, substituted or unsubstituted C₃₋₇ cycloalkyl-C₀₋₆ alkyl, or substituted or unsubstituted aryl-C₀₋₆ alkyl, v=1 or 2; w=1 or 2; and R^b is preferably hydrogen.

8. A compound as claimed in any one of claims 6-7, wherein the linker is linear and the [Connector] is absent, selected from the group consisting of Compounds 1001-1080 where Z is present, and selected from the group consisting of Compounds 2001-2080 where Z' is present.
9. A compound as claimed in claim 10, wherein Z' is present, comprising the formula of Compound 2018.
10. A compound comprising the formula:




wherein [Active Molecule-1] is a [VEGF-Peptide] as claimed in any one of claims 5-11, and [Connector] is covalently bonded to both [VEGF-Peptide] and [Active Molecule-2] and [Connector] is also covalently linked to the Linker L or L'.

11. A compound as claimed in claim 10, wherein [Connector] comprises the formula: -[AM1-Spacer]-[Branch]-[AM2-Spacer]-, and wherein [AM-1-Spacer] and [AM-2-Spacer] are each independently a biologically compatible polymer, block copolymer C, H, N, O, P, S, halogen (F, Cl, Br, I), or a salt thereof, alkyl, alkenyl, alkynyl, oxoalkyl, oxoalkenyl, oxoalkynyl, aminoalkyl, aminoalkenyl, aminoalkynyl, sulfoalkyl, sulfoalkenyl, sulfoalkynyl, phosphoalkyl, phosphoalkenyl, or phosphoalkynyl group, covalently bonded to [Branch], and [Branch] is molecule with at least three reactive groups, and [AM-1-Spacer] is covalently


linked to [Branch] and to the [Active Molecule-1], and [AM-2-Spacer] is covalently linked to [Branch] and to the [Active Molecule-2].

12. A compound as claimed in claim 11, wherein the [AM-1-Spacer] and [AM-2-Spacer] are each independently neutral, water soluble molecules able to form covalent bonds with their respective Active Molecule and [Branch], and are selected from the group consisting of: amino polyethylene glycol acids, polyethylene glycol diacids, amino alkanolic acids, amino alkanolic acids, polyglycine, 2-PEG, 4-PEG and 6-PEG, and may be each 4-PEG.


13. A compound as claimed in any one of claims 10-12, wherein [Branch] is a chemical moiety comprising three orthogonal reactive groups, and is selected from the group consisting of: a cysteine branch, diaminopropionic acid based branch, diaminobutanoic acid based branch, ornithine based branch, lysine based branch, homocysteine branch, bismaleimide branch, and maleimide-acid branch, and derivatives and homologs thereof, and the following structures:



14. A compound as claimed in claim 13, wherein [Branch]-L is selected from the group consisting of:



151



15. A compound as claimed in any one of claims 10-14, wherein [Active Molecule 2] is an Ang-2 binding peptide of the formula: R^3 -[Ang2-peptide]- R^4 wherein R^3 is absent, CH_3 , $C(O)CH_3$, $C(O)CH_2CH_3$, $C(O)CH_2CH_2CH_3$, $C(O)CH(CH_3)CH_3$, $C(O)CH_2CH_2CH_2CH_3$, $C(O)CH(CH_3)CH_2CH_3$, $C(O)C_6H_5$, $C(O)CH_2CH_2(CH_2CH_2O)_{1.5}Me$, amido-2-PEG, N-acyl and N-alkyl an amino protecting group, a lipid fatty acid group or a carbohydrate; and R^4 is absent, OH, NH_2 , $NH(CH_3)$, $NHCH_2CH_3$, $NHCH_2CH_2CH_3$, $NHCH(CH_3)CH_3$, $NHCH_2CH_2CH_2CH_3$, $NHCH(CH_3)CH_2CH_3$, NHC_6H_5 , $NHCH_2CH_2OCH_3$, $NHOCH_3$, $NHOCH_2CH_3$, a carboxy protecting group, a lipid fatty acid group or a carbohydrate, and wherein [Ang2-Peptide] comprises a sequence substantially homologous to: $Q^1 X^2 Y^3 Q^4 X^5 L^6 D^7 E^8 X^9 D^{10} X^{11} X^{12} X^{13} X^{14} D^{15} X^{16} F^{17} M^{18} X^{19} Q^{20} Q^{21} G^{22}$ (SEQ ID NO:107), wherein x^2 is selected from the group consisting of K, N, R, H, Kac, Nick, and CbcK, and X^5 is selected from the group consisting of P, hP, dhP, and BnHP, and X^9 is selected from the group consisting of L, I, ThA, and Kac, and X^{11} is selected from the group consisting of Q, N, C, K, Kac, Dab, and Dap, and X^{12} is selected from the group consisting of L, HL, Nva, I, HchA, HF, and ThA, and X^{13} is selected from the group consisting of L, HL, Nva, I, HchA, HF, and ThA, and X^{14} is selected from the group consisting of aromatic residues, and X^{16} is selected from the group consisting of Q and N, and X^{19} is selected from the group consisting of L, and I, and [Ang-2-Peptide] is covalently linked to [Connector] through a nucleophilic side chain or the N-terminus amino group or C-terminus carboxyl

group of an Ang2-linking residue, the Ang2-linking residue being selected from the group comprising K, R, Y, C, T, S, homologs of lysine, homocysteine, homoserine, Dap, Dab, the N-terminus residue and the C-terminus residue, and wherein one of Q¹, E⁸, X⁹, X¹¹, X¹², D¹⁵, X¹⁶, M¹⁸, X¹⁹ or G²² is substituted with the Ang2-linking residue.

5

16. A compound as claimed in claim 15, wherein the [Ang2-peptide] comprises a sequence substantially homologous to one or more compounds selected from the group consisting of: SEQ ID NOs:137-191.

10

17. A compound as claimed in claim 16, wherein R¹ is C(O)CH₃, R² is NH₂ and R¹-[Ang-peptide]-R² is (SEQ ID NO:153):
{C(O)CH₃}-Q-(Kac)-Y-Q-P-L-D-E-(Kac)-D-K-T-L-Y-D-Q-F-M-L-Q-Q-G-{NH₂} and K at position 11 is the Ang2-linking residue covalently linked to the [Connector].

15

18. A compound as claimed in any one of claims 10-17, wherein [VEGF-Peptide] is covalently linked to [Connector] through a nucleophilic side chain or N-terminus amino group or C-terminus carboxyl group of an VEGF-linking residue, the VEGF-linking residue being selected from the group comprising K, R, Y, C, T, S, homologs of lysine, homocysteine, homoserine, Dap, Dab, the N-terminus residue and the C-terminus residue.

20

19. A compound as claimed in any one of claims 10-18, selected from the group consisting of: Compound 5001-Compound 5028, and Compound 5031-Compound 5074, and may be selected from the group consisting of Compounds 5053, 5060, 5061 and 5062.

25

20. A compound as claimed in any one of claims 10-19, wherein the Z group if present is covalently linked to the combining site of an antibody.

30

21. A compound as claimed in any one of claims 10-20, wherein the antibody is present, and the compound is selected from the group consisting of: 6001-6026, 6028, 6029, and 6031-6074, and may be selected from the group consisting of Compounds 6053, 6060, 6061 and 6062.

35

22. A compound as claimed in claim 21, comprising the formula of Compound 6053.

23. A compound as claimed in any one of claims 6-22, wherein Antibody is present and is a catalytic antibody, and may be selected from the group consisting of aldolase antibody, beta lactamase antibody, esterase antibody, and amidase antibody.

24. A compound as claimed in as claimed in any one of claims 6-23, wherein Antibody is present and is a full-length antibody, Fab, Fab', F(ab')₂, F_v, dsF_v, scF_v, V_H, V_L, diabody, minibody comprising V_H and V_L domains from h38c2, or full length antibody comprising the V_H and V_L domains from h38c2 and a constant domain selected from the group consisting of IgG1, IgG2, IgG3, and IgG4, and may be h38C2 IgG1.

25. A pharmaceutical composition comprising a therapeutically effective amount of a compound as claimed in any preceding claim, and optionally further comprising a therapeutically effective amount of one or more chemotherapeutic agent, the chemotherapeutic agent preferably being a compound selected from the group consisting of 5-Flurouracil, Irinotecan, Taxotere, Sunitinib ,Axitinib, oxilaplatin, bevacizumab, and cetuximab.

1/22

FIG. 1





FIG. 1B



3/22

FIG. 2A

VL	FR1	CDR1	FR2	CDR2	FR3	CDR3	FR4			
	1	2	3	4	5	6	7	8	9	0
	12345678901234567890123	45678901abcde234	567890123456789	0123456	7890123456789012345678	901234567	8901234567	8901234567	8901234567	8901234567
m38C2	DVWMTQPLSLPVLRLGDQASISC	RSSQSLHTHTYGSPLYN	WYLQKPGQSPKLLIY	KVSNRFS	GVPDRFSGSGSGTDFTLRISRVEAEDLGVYFC	SGGTHLPYT	FGGGTKLEIK			
	***	*	****	***	*	****	**	*		*
h38C2	ELQMTQSPSSLSASVGDRVTITC	RSSQSLHTHTYGSPLYN	WYLQKPGQSPKLLIY	KVSNRFS	GVPSRFSGSGSGTDFTLTISLSLQPEDFAVYFC	SGGTHLPYT	FGGGTKVEIK			
	**	*	*****	*	**	***	*	*	****	*
DPK-9	DIQMTQSPSSLSASVGDRVTITC	RASQSISS-----YLN	WYQKPGKAPKLLIY	AASSLQS	GVPSRFSGSGSGTDFTLTISLSLQPEDFATYYC	QQSYSTP				
JK4								LT	FGGGTKVEIK	
VH	FR1	CDR1	FR2	CDR2	FR3	CDR3	FR4			
	1	2	3	4	5	6	7	8	9	0
	123456789012345678901234567890	1ab2345	67890123456789	012abc3456789012345	67890123456789012345678901234	56789012	345678901234	56789012	345678901234	5678901234
m38C2	EVKLVEGGGLVQPGGTMKLSCEISGLTFR	N--YWMS	WVRQSPKGLWVA	EIRLRSDNYATHYAESVKG	KFTISRDDSKSRLYLQMN	SLRTEDTGIYYCKT	YFY-SFSY	WGQGT	LVTVSA	*
	*	***	**	*	*	*	*	*	*	*
h38C2	EVOLVESGGGLVQPGGSLRLSCAASGFTFS	N--YWMS	WVRQSPKGLWVS	EIRLRSDNYATHYAESVKG	RFTISRDN	SKNTLYLQMN	SLRAEDTGIYYCKT	YFY-SFSY	WGQGT	LVTVSS
	*	*	*	*	*	*	*	*	*	*
DP-47	EVQLLESGGGLVQPGGSLRLSCAASGFTFS	S--YAMS	WVRQAPKGLWVS	AISG--SGGSTYYADSVKG	RFTISRDN	SKNTLYLQMN	SLRAEDTAVYYCAK			
JH4								YFDY	WGQGT	LVTVSS

4/22

FIG. 2B**Light Chain** (219 amino acids): SEQ ID NO:1

ELQMTQSPSSLSASVGDRVTITCRSSQSLHTYGSPYLNWYLQKPGQSPKLLIYKVS
 N
 RFSGVPSRFSGSGTDFTLTISSLQPEDFAVYFCSCQGTHLPYTFGGGTKVEIKRTVAA
 PSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK
 DSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

Heavy chain (448 amino acids): SEQ ID NO:2

EVQLVESGGGLVQPGGSLRLSCAASGFTFSNYWMSWVRQSPKEGLEWVSEIRLRSD
 NYATHYAESVKGRFTISRDN SKNTLYLQMNSLRAEDTGIYCYCKTYFSFSYWGQGT
 LVTVSSASTKGPSVFPLAPSSKSTSGGTAAALGCLVKDYFPEPTVSWNSGALTSGVHT
 FPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCP
 PCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVVDGVEV
 HNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYCKVSNKALPAPIEKTISKAK
 GQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPPV
 LDSGGSFFLYSKLTVDKSRWQQGNVVFSCSVMEALHNHYTQKSLSLSPGK



5/22

FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 3E




FIG. 3F





FIG. 3G





6/22

FIG. 4A




FIG. 4B




FIG. 4C




FIG. 4D




FIG. 4E




FIG. 4F




FIG. 4G












FIG. 4H



7/22



8/22

FIG. 6A




FIG. 6B




FIG. 6C




FIG. 6D




FIG. 6E




FIG. 6F




FIG. 6G




FIG. 6H






FIG. 7



10/22

FIG. 8

11/22









FIG. 10





13/22

FIG. 11**FIG. 12**

14/22

FIG. 13**FIG. 14**

15/22

FIG.15**FIG.16**

16/22

FIG. 17A





FIG. 17B



17/22

FIG. 18







FIG. 19



18/22

FIG. 20**FIG. 21**

19/22

FIG. 22









FIG. 23



20/22

FIG. 24**FIG. 25**

21/22

FIG. 26**FIG. 27**

22/22

FIG. 28

19/19

FIGURE 28




FIG. 29





FIGURE 30

FIG. 30



INTERNATIONAL SEARCH REPORT

International application No

PCT/IB2009/051820

A. CLASSIFICATION OF SUBJECT MATTER
INV. A61K47/48 C07K14/71

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FAIRBROTHER W J ET AL: "NOVEL PEPTIDES SELECTED TO BIND VASCULAR ENDOTHELIAL GROWTH FACTOR TARGET THE RECEPTOR-BINDING SITE" BIOCHEMISTRY, AMERICAN CHEMICAL SOCIETY, EASTON, PA.; US, vol. 37, no. 51, 22 December 1998 (1998-12-22), pages 17754-17764, XP000876734 ISSN: 0006-2960 compound v114	1-3, 25
X	WO 01/83525 A2 (AMGEN INC [US]) 8 November 2001 (2001-11-08)	1-2, 25
Y	claims; tables 20,21; sequences 1027,1049,1050	1-25
	----- -/-- -----	

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

24 August 2009

Date of mailing of the international search report

28/08/2009

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Vogt, Titus

INTERNATIONAL SEARCH REPORT

International application No

PCT/IB2009/051820

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2006/094269 A2 (COVX TECHNOLOGIES IRELAND LTD [IE]; BRADSHAW CURT W [US]; DOPPAPALAPUDI) 8 September 2006 (2006-09-08) paragraphs [0184] - [0186], [0385] - [0402], [0405]; claims -----	1-25
Y	DOPPAPALAPUDI ET AL: "Chemically programmed antibodies: Endothelin receptor targeting CovX-Bodies(TM)" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, PERGAMON, ELSEVIER SCIENCE, GB, vol. 17, no. 2, 11 January 2007 (2007-01-11), pages 501-506, XP005827262 ISSN: 0960-894X figure 1 -----	1-25
Y	WO 2006/010057 A2 (AMGEN INC [US]; WALKER KENNETH WILLIAM [US]; KINSTLER OLAF B [US]; STI) 26 January 2006 (2006-01-26) claims; sequences 915, 1538 -----	1-25
X,P	WO 2008/081418 A1 (COVX TECHNOLOGIES IRELAND LTD [IE]; BRADSHAW CURT [US]; SAKAMURI SUKUM) 10 July 2008 (2008-07-10) claims -----	1-25
X,P	WO 2008/056346 A2 (COVX TECHNOLOGIES IRELAND LTD [IE]; BRADSHAW CURT [US]; BHAT ABHIJIT []) 15 May 2008 (2008-05-15) claims -----	1-25

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/IB2009/051820

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0183525	A2	08-11-2001	AU 5943201 A	12-11-2001
			AU 2001259432 B2	21-04-2005
			CA 2407956 A1	08-11-2001
			EP 1278778 A2	29-01-2003
			JP 2003533187 T	11-11-2003
			MX PA02010787 A	14-07-2003
			US 2006234307 A1	19-10-2006
WO 2006094269	A2	08-09-2006	AU 2006218437 A1	08-09-2006
			CA 2598833 A1	08-09-2006
			EP 1853294 A2	14-11-2007
			JP 2008531734 T	14-08-2008
			KR 20070108958 A	14-11-2007
			US 2008152660 A1	26-06-2008
WO 2006010057	A2	26-01-2006	CA 2572765 A1	26-01-2006
			EP 1773400 A2	18-04-2007
			JP 2008505928 T	28-02-2008
			US 2008254020 A1	16-10-2008
WO 2008081418	A1	10-07-2008	AU 2008203641 A1	10-07-2008
			US 2009098130 A1	16-04-2009
WO 2008056346	A2	15-05-2008	AU 2007318912 A1	15-05-2008
			CA 2669207 A1	15-05-2008
			EP 2089425 A2	19-08-2009
			US 2008166364 A1	10-07-2008