

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
18 May 2007 (18.05.2007)

PCT

(10) International Publication Number
WO 2007/056614 A1(51) International Patent Classification:
C07K 14/51 (2006.01) *A61K 48/00* (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:
PCT/US2006/044048(22) International Filing Date:
14 November 2006 (14.11.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/736,452 14 November 2005 (14.11.2005) US

(71) Applicant (for all designated States except US): MERIAL LIMITED [US/US]; 3239 SATELLITE BLVD., Duluth, GA 30096 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventor; and

(75) Inventor/Applicant (for US only): FISHER, Laurent, Bernard [FR/FR]; 17, CHEMIN VERT 69110, Sainte Foy Les Lyon (FR).

(74) Agent: JARECKI-BLACK, Judy; MERIAL LIMITED, 3239 SATELLITE BLVD., Duluth, GA 30096 (US).


Published:

— with international search report

[Continued on next page]

(54) Title: GENE THERAPY FOR RENAL FAILURE

050876pPCR-Script PLASMID MAP



(57) Abstract: The present invention relates to recombinant vectors expressing the BMP-7 polypeptide in host cells and to pharmaceutical compositions comprising such recombinant vectors. The invention also encompasses methods for prevention and/or treatment of both acute and chronic renal failure in mammals, advantageously in dogs and cats, by administration of the recombinant vectors and pharmaceutical compositions of the invention.



- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

5 **TITLE OF THE INVENTION***GENE THERAPY FOR RENAL FAILURE***RELATED APPLICATIONS**

This application claims priority to US provisional application 60/736,452, filed November 14, 2005, herein incorporated by reference.

10 **INCORPORATION BY REFERENCE**

All documents cited or referenced herein (“herein cited documents”), and all documents cited or referenced in herein cited documents, together with any manufacturer’s instructions, descriptions, product specifications, and product sheets for any products mentioned herein or in any document incorporated by reference herein, are hereby incorporated herein by reference, and 15 may be employed in the practice of the invention.

FIELD OF THE INVENTION

The present invention relates to recombinant vectors, to pharmaceutical compositions comprising such recombinant vectors, and to methods for prevention and/or treatment of acute and/or chronic renal failure in mammals. The invention also relates to vectors capable of 20 expressing, in a host, a bioactive polypeptide belonging to the Osteogenic Protein-1 / Bone Morphogenetic Protein-7 (OP-1 / BMP-7) family of proteins.

BACKGROUND OF THE INVENTION

The mammalian renal system serves primary roles both in the removal of catabolic waste products from the bloodstream and in the maintenance of fluid and electrolyte balances in the 25 body. Renal failure is, therefore, a life-threatening conditions in which the build-up of catabolites and other toxins, and/or the development of significant imbalances in electrolytes or fluids, may lead to the failure of other major organs systems and death. As a general matter, renal failure is classified as “acute” or “chronic”. As detailed below, acute and chronic renal failure are debilitating and life-threatening diseases for which no adequate treatments exist to delay, and/or 30 reverse kidney structural alterations associated with the disease.

Acute renal failure (ARF) is usually caused by an ischemic or toxic insult that results in an abrupt decline in renal functions. The kidneys are highly susceptible to ischemia and toxicants because of their unique anatomic and physiologic features. The large renal blood flow (approximately 25 % of the cardiac output) results in increased delivery of blood-borne toxicants 35 to the kidney as compared to other organs. The renal cortex is especially susceptible to toxicant

5 exposure because it receives 90 % of renal blood flow and has a large endothelial surface area due to the numerous glomerular capillaries. Within the renal cortex, the proximal tubule (the S3 segment or "pars recta") and the epithelial cells of the thick ascending arm of the loop of Henle, are most frequently affected by ischemic and toxicant-induced injury because of their solute transport functions and high metabolic rates. As water and electrolytes are reabsorbed from the
10 glomerular filtrate, tubular epithelial cells can be exposed to increasingly high concentrations of toxicants. Similarly, in the medulla the counter-current multiplier system may concentrate toxicants. Toxicants that are either secreted or reabsorbed by tubular epithelial cells (such as gentamicin) may accumulate in high concentrations within these cells. Finally, the kidneys also play a role in the biotransformation of many drugs and toxicants. Biotransformation usually
15 results in the formation of metabolites that are less toxic than the parent compound; however, in some cases (such as oxidation of ethylene glycol to glycolate and oxalate) the metabolites are more toxic.

ARF has three distinct phases, which are categorized as initiation, maintenance, and recovery. During the initiation phase, therapeutic measures that reduce the renal insult (e.g.,
20 fluid therapy) can prevent the development of established ARF. The maintenance phase is characterized by tubular lesions and established nephron dysfunction. The recovery phase of ARF occurs when renal function improves subsequent to nephron repair and compensatory hypertrophy. Tubular lesions may be repaired if the tubular basement membrane is intact and viable cells are present. In addition, functional and morphologic hypertrophy of surviving
25 nephrons can, in some cases, adequately compensate for decreased nephron numbers. Even if renal functional recovery is incomplete, adequate function may be re-established in some cases. More commonly, however, tubular damage is severe and irreversible and a large percentage of animals die or are euthanized in the maintenance phase of ARF.

Despite tremendous efforts to decipher the cellular and molecular pathogenesis of ARF
30 during the past decades, no effective treatment is currently available and the incidence of mortality remains very high in veterinary medicine. At least two retrospective studies have documented the poor prognosis associated with ARF in dogs. In a study of hospital acquired ARF, the survival rate was 38 %, whereas in another study of all types of ARF, the survival rate was 24 %. Thus, there is an un-met medical need for improved prevention and/or treatment of
35 ARF.

5 Chronic renal failure (CRF) may be defined as progressive, permanent and significant reduction of glomerular filtration rate (GFR) due to significant and continuing loss of nephrons. CRF typically begins from a point at which a chronic renal insufficiency (i.e., a permanent decrease in renal function of at least 50-60 %) has resulted from some insult to the renal tissues, which has caused a significant loss of nephron functional units. The initial insult may not have
10 been associated with an episode of acute renal failure. Irrespective of the nature of the initial insult, CRF manifests a “final common path” of signs and symptoms as nephrons are progressively lost and GFR progressively declines. This progressive deterioration in renal function is slow and seemingly inevitable, typically spanning several months to years in canine and feline subjects and many decades in human patients.

15 The early stage of CRF typically begins when GFR has been reduced to approximately one-third of the normal level (e.g., 30-40 ml/min for an average human adult). As a result of the significant nephron loss, and in an apparent “attempt” to maintain the overall GFR with fewer nephrons, the average single nephron GFR (SNGFR) is increased by adaptation of the remaining nephrons at both the structural and functional levels. One structural manifestation of this
20 adaptation that is readily detectable by microscopic examination of biopsy samples is a “compensatory hypertrophy” of both the glomeruli and the tubules of the kidney, a process that actually increases the volume of filtrate which can be produced by each remaining nephron by literal enlargement of the glomeruli and tubules.

25 As a result of the hypertrophy or dilatation of the collecting ducts, the urine of subjects with CRF often contains casts which are 2-6 times the normal diameter (referred to herein as “broad casts” or “renal failure casts”). The presence of such broad casts aids in diagnosis of CRF. At the same time, there are functional changes in the remaining nephrons, such as decreased absorption or increased secretion of normally excreted solute, which may be responses to hormonal or paracrine changes elsewhere in the body (e.g., increasing levels of parathyroid
30 hormone (PTH) in response to changes in serum levels of calcium and phosphate).

These adaptations in the early stage CRF are not successful in completely restoring GFR or other parameters of renal function and, in fact, subject the remaining nephrons to increased risk of loss. For example, the increased SNGFR is associated with mechanical stress on the glomerulus due to hypertension and hyperperfusion. The loss of integrity of podocyte juncitures
35 leads to increased permeability of the glomerulus to macromolecules or “leakiness” of the

5 glomerular capsule. Proliferative effects are also observed in mesangial, epithelial and endothelial cells, as well as increases in the deposition of collagen and other matrix proteins. Sclerosis of both the glomeruli and tubules is another common symptom of the hypertrophied nephrons and the risk of coagulation in the glomerulus is increased. In particular, these adaptations of the remaining nephrons, by pushing the SNGFR well beyond its normal level, 10 actually decrease the capacity of the remaining nephrons to respond to acute changes in water, solute, or acid loads, and therefore actually increase the probability of additional nephron loss.

As CRF progresses, and GFR continues to decline to less than 10 % of normal (i.e., around 5-10 ml/min in humans), the subject enters into end-stage renal disease (ESRD). During this phase, the inability of the remaining nephrons to adequately remove waste products 15 and maintain fluid and electrolyte balance, leads to a rapid decline in which many organ systems, and particularly the cardiovascular system, may begin to fail. At this point, renal failure will rapidly progress to death unless the patient receives renal replacement therapy (i.e., chronic hemodialysis, continuous peritoneal dialysis, or kidney transplantation).

The management of CRF must be conducted to ameliorate all identifiable clinical, 20 metabolic, endocrine and biochemical consequences induced by renal failure including, but not limited to, azotemia, nutritional inadequacies, hypoproliferative anaemia, disordered mineral metabolism, electrolyte disturbances, metabolic acidosis, proteinuria, disordered water metabolism, systemic hypertension and the progression of renal injury through interstitial fibrosis that is considered to be the commonly converging outcome of CRF regardless of the 25 specific etiology.

While tremendous progress has been made during the last decade to address several clinical, metabolic, endocrine and biochemical consequences of CRF, the therapy of clinically chronic fibrosis remains extremely challenging and therefore the long-term medical control of renal disease remains an important un-met therapeutic need. Currently, most advanced therapy 30 targeting the reduction of renal disease-associated fibrosis is focused on the reduction of the activity of the renin-angiotensin system (RAS). Although this strategy has been shown to slow the disease evolution, its efficacy remains partial and it does not completely halt the progression of chronic fibrosis in experimental and clinical conditions. This is probably because many factors other than RAS contribute to the pathogenesis of CRF associated fibrosis.

5 The prevalence of CRF in cats and dogs is increasing. For every 1000 cats evaluated in
1980 in the US, four had renal failure regardless of age. By 1990, the number of reported cases
of renal failure has quadrupled with 16 cases identified for every 1000 cats examined. For cats
older than 15 years of age, 153 cases of renal failure were diagnosed in 1990 for every 1000
examinations. The increase in prevalence of renal failure in aging cats may reflect an increase in
10 veterinary care sought by owners as well as greater efforts by veterinarians to detect the disease.
Whatever the reason, these findings emphasize the emerging awareness and importance of CRF
in older animals. The most frequent etiologies of CRF in companion animals include, but are not
limited to, idiopathic chronic interstitial nephritis, irreversible ARF, familial renal dysplasia or
aplasia, congenital polycystic kidney disease, amyloidosis, glomerulonephritis, hypercalcemia,
15 bilateral hydronephrosis, leptospirosis, pyelonephritis, nephrolithiasis bilateral, Falconi-like
syndrome, hypertension, renal lymphosarcoma.

20 In human medicine, approximately 600 patients per million receive chronic dialysis each
year in the USA, at an average cost approaching \$60,000-\$80,000 per patient per year. Of the
new cases of end-stage renal disease each year, approximately 28-33 % are due to diabetic
nephropathy (or diabetic glomerulopathy or diabetic renal hypertrophy), 24-29 % are due to
hypertensive nephrosclerosis (or hypertensive glomerulosclerosis), and 15-22 % are due to
glomerulonephritis. The 5-year survival rate for all chronic human dialysis patients is
approximately 40%, but for patients over 65, the rate drops to approximately 20%. Therefore, a
need remains for treatments to prevent the progressive loss of renal function which has caused
25 almost 200,000 human patients in the USA alone to become dependent upon chronic dialysis,
and which results in the premature deaths of tens of thousands each year.

30 In light of the fact that specific morphogens and/or growth factors that exhibit renotropic
properties and promote tubular repair and recovery of renal function have been recently
identified, it is conceivable that some of these molecules could have the potential to be used as
therapeutic agents for the prevention and/or treatment of ARF and/or CRF. One such agent is
Bone Morphogenetic Protein-7 (BMP-7, or Osteogenic Protein-1, OP-1), which is a member of
the Transforming Growth Factor- β (TGF- β) superfamily. BMP-7 binds to activin receptors
types I and II, but not to TGF- β receptors type I, II and III. Monomeric BMP-7 has a molecular
weight of 17 to 19 kDa and was originally identified by its ability to induce ectopic bone
35 formation. BMP-7 polypeptide is secreted as a homodimer with an apparent molecular weight of

5 approximately 35-36 kDa. Recently, BMP-7 has been shown to be a key morphogen during nephrogenesis. Renal expression of BMP-7 continues in mature kidneys, especially in medullary collecting ducts. Renal tubules also express BMP-7 receptors. In animal models of ARF and CRF, renal expression of BMP-7 is significantly down-regulated and the administration of recombinant BMP-7 protein has been reported to accelerate renal recovery, an effect that was
10 associated with less interstitial inflammation and programmed cell death.

However, because BMP-7 has a short half live *in vivo* (approximately 30 min), maintenance of a sustained level of exogenous protein in the circulation following injection of the purified protein requires multiple short-interval administrations, creating a very significant practical challenge. The cost of such a multi-injection therapy is too high to be applicable in
15 veterinary medicine. Although gene delivery has been successfully promoted as an alternative to protein therapy for various diseases treatment, it's applicability for ARF and/or CRF prevention and/or treatment through BMP-7 polypeptide expression *in vivo* has not been proposed previously, and its potential effectiveness remains uncertain. Indeed, the low molecular weight of the BMP-7 homodimer (i.e., approximately 35 kDa) would theoretically allow for rapid glomerular filtration. Whether or not levels of BMP-7 expressed *in vivo* could reach
20 therapeutically effective plasma concentrations cannot be predicted or determined from the existing literature. To further complicate the evaluation of *in vivo*-expressed BMP proteins, results can be variable depending on the immune status of the treated animal, with significant differences between immune competent and incompetent animals. Thus, when considered
25 collectively as a whole, the literature does not teach whether levels of BMP-7 expressed *in vivo* could reach plasma concentrations that would be therapeutically useful.

Citation or identification of any document in this application does not constitute and admission that such document is available as prior art to the present invention.

SUMMARY OF THE INVENTION

30 The present invention is directed to methods of prevention and treatment of mammalian subjects who are suffering from, or who are at risk of, acute or chronic renal failure, and to recombinant vectors and pharmaceutical compositions for use in such methods. The methods, vectors and compositions of the invention are useful for reducing mortality and/or morbidity rates, and preventing, inhibiting, delaying, or alleviating the progressive loss of renal function
35 which characterizes renal failure. Subjects for which the methods, recombinant vectors, and

5 compositions of the present invention are useful include, but are not limited to, subjects already afflicted with acute or chronic renal failure, subjects who have already received renal replacement therapy, as well as any subject reasonably expected to suffer from an acute or progressive loss of renal function associated with progressive loss of functioning nephron units. Whether a particular subject is at risk of renal disease, and/or whether a subject may benefit from
10 the methods and/or compositions of the present invention, is a determination that can be routinely made by one of ordinary skill in the relevant medical or veterinary art.

In one embodiment the present invention relates to a vector containing and expressing in a host a pre-pro BMP-7 gene, a proBMP-7 gene or a mature BMP-7 gene. The BMP-7 gene encoding the pre-proBMP-7 polypeptide, the proBMP-7 polypeptide or the mature BMP-7 polypeptide may originate from a mammal. In a preferred embodiment, the expression vector may comprise a polynucleotide that encodes a canine pre-proBMP-7, a canine pro-BMP-7 or a canine mature BMP-7 polypeptide. The polynucleotide encoding the BMP-7 polypeptide may be operatively linked to a promoter and optionally an enhancer.
15

In an advantageous embodiment, the invention relates to a vector containing and expressing the canine proBMP-7 polypeptide, wherein the canine proBMP-7 polypeptide is deleted of the "pre" peptide at the N-terminus, and wherein a peptide signal sequence from a different origin is fused to the canine proBMP-7 polypeptide. Advantageously, the peptide signal sequence may be the insulin-like growth factor 1 (IGF-1) or the tissue plasminogen activator (tPA) peptide signal sequence. In another embodiment, the expression vector may comprise a polynucleotide that encodes a canine mature BMP-7 polypeptide wherein said polypeptide is fused with a peptide signal sequence from BMP-7, IGF-1 or tPA.
20
25

In another embodiment the invention relates to a pharmaceutical composition comprising a vector expressing a pre-proBMP-7 polypeptide, a proBMP-7 polypeptide or a mature BMP-7 polypeptide and a pharmaceutically or veterinarily acceptable carrier, excipient or vehicle. In a particular embodiment, the pharmaceutical composition may comprise a substance to improve
30 the efficacy of transfection or transduction of the vector into the host cells.

In yet another embodiment the invention relates to a method for delivering the BMP-7 polypeptide to a mammal which may comprise injecting a vector capable of expressing, *in vivo*, a pre-proBMP-7 polypeptide, a proBMP-7 polypeptide or a mature BMP-7 polypeptide. In an
35 advantageous embodiment, the animal host may be a dog or a cat. The invention relates to the

5 use of such a vector to prevent and/or treat a mammal for chronic or acute renal failure. The pharmaceutical compositions of the invention may be administered by any suitable route of administration including, but not limited to, by the intramuscular or subcutaneous route. In a particular embodiment the vector may be administered to the host using a needle-free injector or using electrotransfer.

10 In a further embodiment the invention relates to the use of pharmaceutical compositions according to the present invention to treat mammals exhibiting an increase of in serum creatinine concentration and/or an increase in serum urea nitrogen concentration. Advantageously a cat may be treated when the plasma creatinine concentration is higher than 1.9 mg/dl and/or when the plasma urea nitrogen concentration is higher than 35 mg/dl. Advantageously a dog may be
15 treated when the plasma creatinine concentration is higher than 1.6 mg/dl, and/or when the plasma urea nitrogen concentration is higher than 30 mg/dl.

20 It is noted that in this disclosure and particularly in the claims and/or paragraphs, terms such as "comprises", "comprised", "comprising" and the like can have the meaning attributed to it in U.S. Patent law; e.g., they can mean "includes", "included", "including", and the like; and that terms such as "consisting essentially of" and "consists essentially of" have the meaning ascribed to them in U.S. Patent law, e.g., they allow for elements not explicitly recited, but exclude elements that are found in the prior art or that affect a basic or novel characteristic of the invention.

25 These and other embodiments are described in, or are obvious from and encompassed by, the following Detailed Description.

BRIEF DESCRIPTION OF THE DRAWINGS

The following detailed description, given by way of example but not intended to limit the invention solely to the specific embodiments described, may best be understood in conjunction with the accompanying drawings, in which:

30 FIG. 1 depicts the 050876pPCR-Script plasmid map and the encoded open reading frame ("ORF") of the canine BMP-7. The nucleotide sequence of the encoded ORF is that of SEQ ID NO: 2 and the amino acid sequence of the encoded ORF is that of SEQ ID NO: 3.

35 FIG. 2 depicts the pNB292 plasmid map and the encoded ORF of the canine BMP-7. The nucleotide sequence of the encoded ORF is that of SEQ ID NO: 2 and the amino acid sequence of the encoded ORF is that of SEQ ID NO: 3.

5 Fig. 3 provides a histogram illustrating the frequency of kidney lesions having certain grades in control rats and in rats treated with a plasmid expressing BMP-7.

Fig. 4 provides a histogram illustrating of frequency of severe lesions in control rats and in rats treated with a plasmid expressing BMP-7.

Also included as part of the present application is a sequence listing in which: SEQ ID
10 N0: 1 is the nucleotide sequence of the canine pre-proBMP-7 polypeptide, SEQ ID N0: 2 is the codon-optimized nucleotide sequence of the canine pre-proBMP-7 polypeptide, SEQ ID N0: 3 is the amino acid sequence of the canine pre-proBMP-7 polypeptide, SEQ ID N0: 4 is the nucleotide sequence of the short signal peptide from tPA (23 amino acids), SEQ ID N0: 5 is the amino acid sequence of the short signal peptide from tPA (23 amino acids), SEQ ID N0: 6 is the
15 nucleotide sequence of the long signal peptide from tPA (28 amino acids), SEQ ID N0: 7 is the amino acid sequence of the long signal peptide from tPA (28 amino acids), SEQ ID N0: 8 is the nucleotide sequence of the equine IGF-1 signal peptide, SEQ ID N0: 9 is the amino acid sequence of the equine IGF-1 signal peptide, SEQ ID N0: 10 is the nucleotide sequence of the pBN292 plasmid, SEQ ID N0: 11 is the nucleotide sequence of the canine IGF-1 signal peptide,
20 and SEQ ID N0: 12 is the amino acid sequence of the canine IGF-1 signal peptide.

DETAILED DESCRIPTION

The methods and compositions of the present invention can be used for preventative treatment of renal failure. The terms "prevention", "prophylaxis", "preventative treatment" and "prophylactic treatment", as they relate to renal failure, and as they are used herein and in the
25 field of human and veterinary medicine, relate to the treatment of either healthy animals or animals suffering from an unrelated disease, but who are considered to be at risk of acute renal failure. The main risk factors for acute renal failure in cats and dogs include, but are not limited to, shock and/or hypovolemia (for example haemorrhage, hypotensive shock, septic shock, prolonged or deep anaesthesia, hypovolemia, heat stroke, trauma, burns, or diuretic abuse),
30 systemic diseases (for example pancreatitis, peritonitis, hepatic failure, disseminated intravascular coagulation, adrenal insufficiency or vasculitis), ischemia (as caused by, for example, thromboembolic occlusion or malignant hypertension), infections (for example leptospirosis, pyelonephritis, feline infectious peritonitis, borreliosis, leishmaniasis, babesiosis, septicaemia or septic emboli), systemic renal disease (for example multiple organ failure, glomerulonephritis,
35 systemic lupus erythematosus, renal vein thrombosis, urinary outflow obstruction, haemolytic

5 uremic syndrome, heme pigmenturia-crush syndrome or polycythemia), advanced age, congenital and/or genetic renal diseases, and other miscellaneous factors such as exposure to nephrotoxins (for example aminoglycosides, amphotericin B, cisplatin, adriamycin, non steroid anti-inflammatory drugs, diuretics, IL-2 or allopurinol), neoplasia (for example lymphoma), hypercalcemia, trauma (for example avulsions), malignant hypertension, oxalate nephrosis, and
10 the like.

15 Treatment for preventative purposes is generally conducted within a few weeks (ideally with 6 to 8 days) before the exposure of a healthy animal to one or more of the aforementioned risk factors for acute renal failure. Alternatively, in diseased animals for which an associated risk factor for acute renal failure has been identified, treatment may be conducted as quickly as possible to limit any negative impact of the primary disease or risk factor on the kidney metabolism and/or the structure and organization of the kidney tissue.

20 In addition to preventative treatments, the methods and compositions of the present invention can also be used for therapeutic treatment of renal failure. The terms "therapy" or "therapeutic treatment", as they relate to renal failure, and as they are used herein and in the field of veterinary medicine, relate to treating, or supporting and/or accelerating treatment of, animals that are already suffering from, or are recovering from (i.e. are in the recovery phase) acute renal failure, or treatments aimed at slowing down and/or reversing lesion evolution in animals diagnosed as having, or at being at risk of, chronic renal failure. A critical objective of therapy is to reduce the risk of an evolution towards CRF subsequent to an ARF event. As used herein, a
25 subject is said to suffer from CRF, or be at risk of developing CRF, if the subject is reasonably expected to suffer a progressive loss of renal function associated with progressive loss of functioning nephron units. Whether a particular subject suffers of CRF, or is at risk of developing CRF, can readily be determination by one with ordinary skill in the relevant veterinary or medical art.

30 The main risks factors for chronic renal failure in dogs include, but are not limited to, idiopathic chronic interstitial nephritis, irreversible ARF, familial renal dysplasia or aplasia (high risk breeds include Norwegian elkhounds, Lhasa apso, Samoyed, Cocker spaniel, Doberman pinsher, Standard poodle, and Golden retriever), congenital polycystic kidney disease (for example in Cairn terriers), amyloidosis, glomerulonephritis, hypercalcemia, bilateral

5 hydronephrosis, leptospirosis, pyelonephritis, nephrolithiasis bilateral, Falconi-like syndrome, and hypertension.

The main risk factors for chronic renal failure in cats include, but are not limited to, idiopathic chronic interstitial nephritis, irreversible ARF, renal lymphosarcoma, polycystic kidney disease (for example in familial in Persian cats), glomerulonephritis, bilateral

10 hydronephrosis, amyloidosis, pyelonephritis, hypercalcemia, and bilateral nephrolithiasis.

Human subjects suffering from CRF, or whom are at risk of developing CRF, or who may be in need of renal replacement therapy, include, but are not limited to, subjects with end-stage renal disease, chronic diabetes nephropathy, hypertensive nephrosclerosis, chronic glomerulonephritis, hereditary nephritis, and/or renal dysplasia, subjects who have had a biopsy 15 indicating glomerular hypertrophy, tubular hypertrophy, chronic glomerulosclerosis, and/or chronic tubulo-interstitial sclerosis, subjects who have had an ultrasound, MRI, CAT scan, or other non-invasive examination indicating the presence of renal fibrosis, subjects having an unusual number of broad casts present in their urinary sediment, subjects having a glomerular filtration rate ("GFR") which is chronically less than 50%, and more particularly less than about 20 40%, 30% or 20%, of the expected GFR for the subject, subjects possessing a number of functional nephron units which is less than about 50%, and more particularly less than about 40%, 30% or 20% of the number of functional nephron units possessed by a healthy but otherwise similar subject, subjects with only a single kidney, and subjects that are kidney transplant recipients.

25 The "glomerular filtration rate" or "GFR" is proportional to the rate of clearance into the urine of "marker" substance which is a plasma-borne substance which is not bound by serum proteins, is freely filtered across glomeruli, and is neither secreted nor reabsorbed by the renal tubules. Thus, as used herein, GFR preferably is defined by the following equation:

$$GFR = \frac{U_{conc} \times V}{P_{conc}}$$

30 where U_{conc} is the urine concentration of the marker substance, P_{conc} is plasma concentration of the marker substance, and V is the urine flow rate in ml/min. Optionally, the GFR can BE corrected for body surface area. Thus, the GFR values may be regarded as being in units of 35 ml/min / 1.73 m^2 . The preferred marker substance for GFR measurements is inulin, however,

5 because of difficulties in measuring the concentration of this substance, creatinine is typically used as the marker for GFR measurements in clinical settings.

An estimate of the "expected GFR" or "GFR_{exp}" may be provided based upon considerations of a subject's age, weight, sex, body surface area, and degree of musculature, and the plasma concentration of some marker compound (e.g., creatinine) as determined by a blood 10 test. Thus, as an example, an expected GFR may be estimated as:

$$GFR_{exp} \approx \frac{(140 - \text{age}) \times \text{weight (kg)}}{72 \times P_{conc} (\text{mg/dL})}$$

This estimate does not take into consideration such factors as surface area, degree of 15 musculature, or percentage of body fat. Nonetheless, using plasma creatinine levels as the marker, this formula has been employed for human males as an inexpensive means of estimating GFR. Because creatinine is produced by striated muscles, the expected GFR of human females subjects is estimated by the same equation multiplied by 0.85 to account for expected difference in muscle mass (*see Lemann et al., 1990 Am. J. Kidney Dis. 16(3); 236-243*).

20 Microscopic examination of urinary sediment for the presence of formed elements is a standard procedure in urine analysis. Amongst the formed elements which may be present in urine, are cylindrical masses of agglutinated materials that typically represent a mold or "cast" of the lumen of a distal convoluted tubule or collecting tube. In healthy human beings, such casts typically have a diameter of 15-25 μm . In subjects with CRF, however, hypertrophy of the 25 tubules may result in the presence of casts which are 2-6 times the diameter of normal casts and often have a homogeneous waxy appearance. These are referred to as "broad casts" or "renal failure casts". As used herein, the term "broad cast" is used to refer to urinary sediment casts having a diameter of 2-6 times normal for the subject, or about 30-150 μm for human casts.

As used herein with respect to clinical indications the term "acute" is used to refer to 30 renal pathologies for which onset occurs rapidly, typically within hours or days of exposure to an insult or risk factor.

As used herein with respect to clinical indications the term "chronic" means persisting for a period of at least three, and more preferably, at least six months. Thus, for example, a subject with a measured GFR chronically below 50% of GFR_{exp} is a subject in which the GFR has been 35 measured and found to be below 50% of GFR_{exp} in at least two measurements separated by at

5 least three, and more preferably, by at least six months, and for which there is no medically sound reason to believe that GFR was substantially (e.g., 10 %) higher during the intervening period. Other indicators of abnormal renal function, such as the presence of broad casts, could similarly be described as chronic if the presence of such indicators persisted in at least two measurements separated by at least three, and more preferably, by at least six months.

10 Table 1 lists some, but not all, of the parameters that may be useful in differentiating between ARF and CRF.

Table 1: Parameters Useful for Differentiating between ARF and CRF

	Acute Renal Failure (ARF)	Chronic Renal Failure (CRF)
History	Ischemic or toxicant exposure	Previous renal disease or renal insufficiency Longstanding polydipsia/polyuria Chronic weight loss, vomiting, diarrhoea
Physical Examination	Good body condition Smooth, swollen, painful kidneys Relatively severe clinical signs for level of dysfunction (azotemia)	Poor body condition Small, irregular kidneys Relatively mild clinical signs for level of dysfunction (azotemia) Osteodystrophy
Clinicopathologic findings	Normal or increased hematocrit Active urine sediment Normal to increased serum potassium More severe metabolic acidosis	Non regenerative anemia Inactive urine sediment Normal to low serum potassium Less severe metabolic acidosis

15 The present invention provides therapies and preventative treatments for renal failure that utilize pharmaceutical compositions comprising vectors capable of expressing the BMP-7 polypeptide *in vivo* and methods and composition for inducing a sustained increase in plasma BMP-7 concentration and thereby reducing the activation of the TGF- β pathway on epithelial cells. TGF- β activation triggers, amongst other things, the phosphohorylation of Smad2 and Smad3 factors and their nuclear import, leading to the promotion of epithelial-mesenchymal

5 transition and to the repression of mesenchymal-epithelial transition, and acting as key trigger for fibrosis. Although BMP-7 is expressed in adult kidneys, its expression is frequently down regulated in the face of renal failure. Therefore, exogenous *in vivo*-produced BMP-7 can help restore levels of BMP-7 to normal physiological levels, leading to the control and regression of the fibrosis associated with tubulo-interstitial nephritis and CRF.

10 As used herein, a pharmaceutical composition according to the invention is said to have “therapeutic efficacy”, or to be “therapeutically effective”, if administration of that amount of the composition is sufficient to cause a significant improvement of the clinical signs or measurable markers of the disease in a mammalian subject suffering from ARF or CRF. As used herein, a pharmaceutical composition according to the invention is said to have “prophylactic efficacy” or
15 to be “prophylactically effective”, if administration of that amount of the composition is sufficient prevent the development of ARF in a subject. The term “therapeutically effective” may also be used herein, in a more general sense, to refer to an amount of a composition that is either sufficient to cause a significant improvement of the clinical signs or measurable markers of disease in a mammalian subject suffering from ARF or CRF, or that is sufficient to prevent the
20 development of ARF in a subject.

Measurable markers of renal function, which are also useful in evaluating the ARF or CRF status of a subject, are well known in the medical and veterinary literature and to those of skill in the art, and include, but are not limited to, blood urea nitrogen or “BUN” levels (both static measurements and measurements of rates of increase or decrease in BUN levels), serum
25 creatinine levels (both static measurements and measurements of rates of increase or decrease in serum creatinine levels), measurements of the BUN/creatinine ratio (static measurements of measurements of the rate of change of the BUN/creatinine ratio), urine/plasma ratios for creatinine, urine/plasma ratios for urea, glomerular filtration rates (GFR), serum concentrations of sodium (Na⁺), urine osmolarity, daily urine output, and the like (see, for example, *Brenner and Lazarus (1994)*, in *Harrison's principles of Internal medicine, 13th edition, Isselbacher et al. eds, McGraw Hill Text, NY*; *Luke and Strom (1994)*, in *Internal Medicine, 4th Edition, J. H. Stein, ed., Mosby-Year Book, Inc. St Louis*). Of the above, measurements of the plasma concentrations of creatinine and/or urea or BUN are particularly important and useful readouts of renal function.

5 Normal values for serum creatinine concentrations range from about 0.5 to about 1.6 mg/decilitre ("dl") in dogs and from about 0.5 to about 1.9 mg/dl in cats. The upper limit of the normal physiological range of serum creatinine levels is slightly higher in cats than in dogs. With the exception of diet, factors influencing physiological values of serum creatinine concentration are poorly understood. It is known that a diet rich in protein has the potential to cause transient
10 hypercreatinemia. For example, an increase of around 25% in serum creatinine concentration can occur over a 6-9 hour period when healthy dogs are fed with commercial food. The relevance of minor variations of creatinemia are difficult to interpret, and the smallest relevant variation between two successive measurements of creatinine levels is considered to be a change in concentration of 35 μ mol/l from normal values.

15 The upper limit of the normal physiological range for BUN levels in fasting dogs and cats ranges from about 8.8 to about 25.9 mg/dl in dogs, and from about 15.4 to about 31.2 mg/dl in cats - the upper limits of the normal range are slightly higher in cats than in dogs. BUN levels, like creatinine levels, are influenced by diet. Other factors that can lead to variation in BUN levels include long-term glucocorticoide treatment and/or hepatocellular failure.

20 Any significant increase of serum creatinine levels and/or BUN levels above their normal physiological ranges is a sign of a reduced ability of the kidneys to eliminate waste and catabolites (i.e., excretory failure).

25 Experimental demonstration of the efficacy of the methods and compositions of the present invention (e.g. the methods and compositions useful for gene therapy with BMP-7 or functional equivalents of BMP-7), can be performed by performed in a variety of ways, for example, by demonstrating that animals treated using the methods and compositions of the present invention exhibit a significantly reduced elevation of plasma creatinine and/or BUN, as compared to placebo-treated animals, when exposed to a trigger or risk factor such as, for example, a toxicant (e.g., $HgCl_2$) or a procedure that induces renal ischemia (e.g., bilateral renal
30 arteries occlusion).

35 Similarly, tissue readouts can be used to demonstrate the efficacy of the methods and compositions of the present invention. Examples of suitable tissular readouts include the quantification of tubulo-interstitial nephritic lesions ("TIN" lesions) within the cortical parenchyma of the kidney, and to a lesser extent, withing the medullary parenchyma of the kidney. It is well documented that renal interstitial fibrosis associated with tubulo-interstitial

5 nephritis (TIN) is a common final pathway of kidney disorders with a wide spectrum of diverse etiologies. Deterioration of renal function is largely determined by the extent of the tubulo-interstitial lesions in many forms of renal diseases, and also in several experimental animal models. Accordingly, method or composition that is able to slow down or reverse the evolution of TIN fibrosis has the potential to benefit all kidney disorders through a disease-modifying mechanism (i.e., by limiting the degradation and disorganization of the structural elements of kidney tissues). Experimental demonstration of the efficacy of the BMP-7 gene therapy methods and compositions of the present invention can be demonstrated from the observation that BMP-7-treated animals have significantly reduced tubulo-interstitial lesions in the kidneys than controls as assessed using the unilateral ureteral obstruction or "UUO" model. The UUO model is a well-established animal model of chronic progression of renal fibrosis associated with progressive tubular atrophy and interstitial collagen accumulation. The UUO model is well known in art (see for example, R. Chevalier et al., *Kidney Int.* 2000, 57, 882-890, the contents of which are hereby incorporated by reference in their entirety), and the unilateral ureteral obstruction procedure can be readily performed by those of ordinary skill in the art. The UUO model is typically associated with very significant tubulo-interstitial pathology and with minimal glomerular lesions, and is a relevant and useful experimental model for demonstrating the efficacy of the methods and compositions of the present invention, for example the demonstrating the efficacy of the gene therapy strategy disclosed herein which is based on the *in vivo* expression of BMP-7 or functional equivalents of BMP-7. Using this model, the evaluation of TIN in the renal cortex can be determined using conventional hematoxylin and eosin (or "H&E") staining and/or collagen-specific Masson Trichrome staining of fixed tissues. Characterization of the lesions is based on the extent of tubular dilatation, epithelial atrophy, and interstitial expansion with myofibroblast activation and matrix deposition. Additional investigations can be based on immunohistochemistry and histomorphometry techniques using, for example, α -smooth muscle actin (" α -SMA") specific antibodies to characterize and quantify the level of epithelial to mesenchyme transition (or "EMT") in the tissue. Complementary immunohistochemical analysis can also be performed with antibodies specific for collagen I or for fibronectin. Quantification of cellular infiltration is an additional readout that can be used to characterize the lesions. Immunohistochemical analysis of the latter can be conducted using, for

5 example, anti ED-1 or anti mac-1 antibodies that are specific for macrophages. Collectively, the results of the above readouts can be used to provide a grade for the lesion.

In addition to the above, any other suitable methods or readouts for studying kidney disease and/or kidney function, including any other suitable animal models, can also be used to demonstrate the efficacy of the methods and compositions of the present invention, and to 10 determine what amount of such compositions, or what modes of administration, will be therapeutically or prophylactically effective.

In one aspect, the present invention related to a vector capable of expressing, *in vivo* in a host, a Bone Morphogenetic Protein-7 (BMP-7) polypeptide, or a variant or a fragment thereof. As used herein “BMP-7 polypeptide” may be used to refer to pre-pro, pro or mature BMP-7 15 polypeptides, wherein the pro and mature BMP-7 polypeptides may be fused to a BMP-7, IGF-1 or tPA signal peptide. The BMP-7 polypeptides of the present invention are preferably of canine origin. In one embodiment the vector contains and expresses in the host a pre-proBMP-7, a proBMP-7 or a mature BMP-7 nucleotide sequence or gene. The nucleotide sequence or gene encoding the pre-proBMP-7 polypeptide, the proBMP-7 polypeptide or the mature BMP-7 20 polypeptide originates from a mammal, for example a cat or a dog. In a preferred embodiment the BMP-7 nucleotide sequence or gene originates from a dog.

BMP-7 is also known as Osteogenic Protein-1 or “OP-1”, and is a member of the transforming growth factor- β or “TGF- β ” superfamily. It is a secreted protein that is processed from the pro-protein to yield the carboxy-terminal mature protein. Within the mature protein 25 there is a conserved pattern of seven cysteine residues defining a domain that extends from amino acid 330 to amino acid 430 of SEQ ID N0: 3. The active form of the protein is a disulfide-bonded homodimer. In its mature, native form, naturally occurring BMP-7 is a glycosylated dimer having an apparent molecular weight of about 30-36 kDa, as determined by SDS- polyacrylamide gel electrophoresis (“SDS-PAGE”). When reduced, the 30 kDa protein 30 gives rise to two glycosylated polypeptide subunits having apparent molecular weights of about 16 kDa and 18 kDa. The unglycosylated protein has an apparent molecular weight of about 27 kDa. When reduced, the 27 kDa unglycosylated protein gives rise to two unglycosylated polypeptide chains, having molecular weights of about 14 kDa and 16 kDa.

Typically, the naturally occurring BMP-7 protein is translated as a precursor, having an 35 N-terminal signal peptide sequence, a “pro” domain, and a “mature” protein domain. The signal

5 peptide is 29 residues long and is cleaved off rapidly upon translation at a cleavage site that can be predicted using the method of *Von Heijne* (1986), *Nucleic Acid Research*, 14; 4683-4691. The “pro” domain has 264 residues in human, canine, swine and bovine BMP-7, and 263 residues in mouse BMP-7. The pro domain is cleaved to yield the “mature” C-terminal domain of 139 residues, which includes the conserved seven-cysteine C-terminal domain of 102 residues.

10 As referred to herein, the “pro form” of the BMP-7 polypeptide refers to a protein comprising a pair of polypeptides, each comprising a pro domain in either covalent or non-covalent association with the mature domain of the BMP-7 polypeptide. The pro form appears to be the primary form secreted from cultured mammalian cells. The “mature form” of the protein refers to the mature C-terminal domain which is not associated, either covalently or non-covalently,

15 with the pro domain.

As used herein the terms “pre-pro BMP-7”, “pro BMP-7”, “mature BMP-7” and “BMP-7” refer not only to the specific polypeptides and sequences illustrated in the specification and in the accompanying sequence listing, but also refer to any and all of the known naturally occurring variants, of these proteins including, but not limited to, derivatives, mutants, homologues, orthologs, allelic variants, allelic polymorphs, polymorphic variants, phylogenetic counterparts, and also any and all non-naturally occurring variants of these proteins, including but not limited to derivatives, mutants, fragments, fusion proteins, and the like. As used herein, the term “variant” encompasses all such naturally occurring and non-naturally occurring variants. In particular, the present invention encompasses all such variants that retain the feature of being useful for the therapeutic or prophylactic treatment of renal diseases including ARF and CRF, and/or that retain BMP-7 activity.

These functionally equivalent variants, derivatives, and fragments, and the like display the ability to retain BMP-7 activity. A functional equivalent, as used herein, refers to any BMP-7 variants, derivatives, fragments, and the like that meet either of the following two criteria (a) they have a significant level of amino acid sequence homology with the protein sequence of BMP-7 as described herein, or is encoded by a nucleotide that has a significant level of nucleotide sequence homology with the protein sequence of BMP-7 as described herein; or (b) they have the ability to provide a statistically different response in the treated group as compared to a placebo treated group in at least one of the following experimental models of renal failure in rodents: (i) a toxicant-induced or ischemia-induced renal failure model, where reduced elevation

5 of plasma creatinine or BUN is expected in the treated as compared to the control/placebo group; (ii) a UUO model of renal failure, where reduced lesion grading is expected in the treated group as compared to the control/placebo group.

By way of illustration of variants, derivatives, and the like that are encompassed by the present invention include, but are not limited to, BMP-7 variants, derivatives, and the like that are encoded by nucleotide sequences that are not exactly the same as the nucleotide sequences disclosed herein, but wherein the changes in the nucleotide sequences do not change the encoded amino acid sequence, or result in conservative substitutions of amino acid residues, deletion of addition of one or a few amino acids, substitution of amino acid residues by amino acid analogs e that do not significantly affect the properties of the encoded polypeptides, and the like.

15 Examples of conservative amino acid substitutions include glycine/alanine substitutions; valine/soleucine/leucine substitutions; asparagine/glutamine substitutions; aspartic acid/glutamic acid substitutions; serine/threonine/methionine substitutions; lysine/arginine substitutions; and phenylalanine/tyrosine/tryptophan substitutions. Other types of substitutions, variations, additions, deletions and derivatives that result in functional BMP-7 derivatives, as described 20 above, are also encompassed by the present invention, and one of skill in the art would readily know how to make, identify, or select such variants or derivatives, and how to test for BMP-7 activity of those variants or derivatives. One of skill in the art may optimize the expression of the BMP-7 polypeptides of the invention by removing cryptic splice sites, by adapting the codon usage by introducing a Kozak consensus sequence before the start codon, by changing the codon 25 usage or combination thereof to improve expression.

In another embodiment, the present invention comprises a canine pre-proBMP-7 polypeptide variant having at least 90 %, at least 91 %, at least 92 %, at least 93 %, at least 94 %, at least 95 %, at least 96 %, at least 97 %, at least 98 % or at least 99 % homology or identity with residues 1 to 431 of SEQ ID NO: 3.

30 In another embodiment the invention comprises a canine mature BMP-7 polypeptide variant having at least 97 %, at least 97.5 %, at least 98 %, at least 98.5 %, or at least 99 % homology or identity with residues 293 to residue 431 of SEQ ID NO: 3.

35 For the purposes of the present invention, sequence identity or homology is determined by comparing the sequences when aligned so as to maximize overlap and identity while minimizing sequence gaps. In particular, sequence identity may be determined using any of a

5 number of mathematical algorithms. A non-limiting example of a mathematical algorithm used for comparison of two sequences is the algorithm of Karlin & Altschul, Proc. Natl. Acad. Sci. USA 1990, 87, 2264-2268, modified as in Karlin & Altschul, Proc. Natl. Acad. Sci. USA 1993, 90, 5873-5877.

10 Another example of a mathematical algorithm used for comparison of sequences is the algorithm of Myers & Miller, CABIOS 1988, 4, 11-17. Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used. Yet another useful algorithm for identifying regions of local sequence similarity and alignment is the FASTA 15 algorithm as described in Pearson & Lipman, Proc. Natl. Acad. Sci. USA 1988, 85, 2444-2448.

19 Advantageous for use according to the present invention is the WU-BLAST (Washington University BLAST) version 2.0 software. WU-BLAST version 2.0 executable programs for several UNIX platforms can be downloaded from <ftp://blast.wustl.edu/blast/executables>. This program is based on WU-BLAST version 1.4, which in turn is based on the public domain 20 NCBI-BLAST version 1.4 (Altschul & Gish, 1996, Local alignment statistics, Doolittle ed., Methods in Enzymology 266, 460-480; Altschul et al., Journal of Molecular Biology 1990, 215, 403-410; Gish & States, Nature Genetics, 1993, 3: 266-272; Karlin & Altschul, 1993, Proc. Natl. Acad. Sci. USA 90, 5873-5877; all of which are incorporated by reference herein).

25 In general, comparison of amino acid sequences is accomplished by aligning an amino acid sequence of a polypeptide of a known structure with the amino acid sequence of a the polypeptide of unknown structure. Amino acids in the sequences are then compared and groups of amino acids that are homologous are grouped together. This method detects conserved regions of the polypeptides and accounts for amino acid insertions and deletions. Homology 30 between amino acid sequences can be determined by using commercially available algorithms (see also the description of homology above). In addition to those otherwise mentioned herein, mention is made too of the programs BLAST, gapped BLAST, BLASTN, BLASTP, and PSI-BLAST, provided by the National Center for Biotechnology Information. These programs are widely used in the art for this purpose and can align homologous regions of two amino acid sequences.

5 In all search programs in the suite, the gapped alignment routines are integral to the database search itself. Gapping can be turned off if desired. The default penalty (Q) for a gap of length one is Q=9 for proteins and BLASTP, and Q=10 for BLASTN, but may be changed to any integer. The default per-residue penalty for extending a gap (R) is R=2 for proteins and BLASTP, and R=10 for BLASTN, but may be changed to any integer. Any combination of
10 values for Q and R can be used in order to align sequences so as to maximize overlap and identity while minimizing sequence gaps. The default amino acid comparison matrix is BLOSUM62, but other amino acid comparison matrices such as PAM can be utilized..

In a preferred embodiment, the present invention provides a vector that contains and expresses a polynucleotide encoding a canine pre-proBMP-7 polypeptide, and more preferably
15 that contains and expresses nucleotides 1 to 1296 of SEQ ID N0: 1. Preferably this vector expresses a polypeptide having the amino acid sequence of SEQ ID N0: 3

In one embodiment, the peptide signal (prepeptide) sequence spans from the Met residue at position (1) to the Ala residue at position (29), with the numbering of the amino acid residues being that of the pre-proBMP-7 sequence identified as SEQ ID N0: 3. Cleavage of the signal
20 peptide may occur after the Ala(29) residue. After cleavage of the preBMP-7 peptide, the proBMP-7 polypeptide is secondarily cleaved after the sequence Arg-X-X-Arg(292) to lead to the mature BMP-7 polypeptide.

The terms "protein", "polypeptide" and "polypeptide fragment" are used interchangeably herein to refer to polymers of amino acid residues of any length.

25 In certain embodiments, the expression vector comprises a polynucleotide that encodes a canine mature BMP-7 polypeptide, wherein the polypeptide is fused to a peptide signal sequence that is, or that comprises or is derived from the canine BMP-7 signal peptide. In other embodiments, the signal peptide sequence may be, or comprise or be derived from, other BMP-7 signal peptides.

30 The present invention further relates to vectors containing and expressing a polynucleotide encoding the proBMP-7 polypeptide, wherein the pre-BMP-7 signal peptide is deleted and wherein a peptide signal sequence from a different origin is fused to the proBMP-7 polypeptide. For example, in certain embodiments, the peptide signal sequence may be the insulin-like growth factor 1 (IGF-1) or the tissue plasminogen activator (tPA) peptide signal
35 sequence. In a preferred embodiment the proBMP-7 encoded by the polynucleotide is a canine

5 proBMP-7 polypeptide. Advantageously the proBMP-7 is encoded by a polynucleotide nucleotide that is, or comprises, or is derived from nucleotides 88 to 1296 of SEQ ID N0: 1, and that encodes, or comprises an amino acid sequence corresponding to amino acid residues 30 to 431 of SEQ ID N0: 3. In another preferred embodiment, the codon-optimized canine nucleotide sequence corresponding to SEQ ID N0: 2 is used.

10 In embodiments where the signal peptide is derived from the IGF-I signal peptides, it is preferred that the peptide signal is, or comprises, or is derived from, the horse IGF-1 peptide signal, and preferably that defined by amino acid residues 1 to 25 of SEQ ID N0: 9, and encoded by nucleotides 1 to 75 of SEQ ID N0: 8. In alternate embodiments, the IGF-1 peptide signal may be, or may comprise, or may be derived from, the canine IGF-1 peptide signal, and preferably is, 15 or comprises, or is derived from, the canine IGF-1 peptide signal defined by amino acid residues 1 to 25 of SEQ ID N0: 12, and that is encoded by nucleotides 1 to 75 of SEQ ID N0: 11.

15 In other embodiments, the peptide signal may be, or may comprise or be derived from, the tPA peptide signal, such as the human tPA signal peptide. In a preferred embodiment, the tPA signal peptide used, is, or comprises or is derived from, amino acid residues 1 to 23 of the 20 human tPA signal peptide sequence of SEQ ID N0: 5, and is encoded by nucleotides 1 to 69 of SEQ ID N0: 4. In an alternative embodiment, a human tPA signal peptide that is, or comprises or is derived from, amino acid residues 1 to 28 of SEQ ID N0: 7 and is encoded by nucleotides 1 to 84 of SEQ ID N0: 6 may be used.

25 According to an advantageous embodiment of the invention, the expression vector comprises the polynucleotides encoding the signal peptide of IGF1 or tPA according to SEQ ID N0: 5, 7, 9 or 12 fused to the pre-proBMP-7 polypeptide deleted of the signal peptide (corresponding to residue 30 to residue 431). According to another embodiment of the invention, the expression vector comprises the polynucleotides encoding the signal peptide of IGF1 or tPA fused to the mature BMP-7 (corresponding to residue 293 to residue 431).

30 Polynucleotides comprising a desired sequence can be inserted into a suitable expression vector, and the vector in turn can be introduced into a suitable host cell, e.g. *E. coli* for replication and amplification.

35 In some embodiments, the present invention encompasses a vector capable of expressing canine pre-proBMP-7, canine proBMP-7, canine mature BMP-7, or a variant or fragment thereof. For the mature BMP-7 or the proBMP-7, it is preferred that the nucleotide sequence

5 encoding the peptide is preceded immediately by a nucleotide sequence in-frame encoding a peptide signal in order to facilitate the secretion of BMP-7 into the extra cellular medium. The signal sequence can be the natural sequence from the pre-proBMP-7 or a peptide signal from a secreted protein e.g. the signal peptide from the tissue plasminogen activator protein (tPA), in particular the human tPA (S. Friezner Degen et al J. Biol. Chem. 1996, 261, 6972-6985; R.
10 Rickles et al J. Biol. Chem. 1988, 263, 1563-1569; D. Berg. et al Biochem. Biophys. Res. Commun. 1991, 179, 1289-1296), or the signal peptide from the Insulin-like growth factor 1 (IGF1), in particular the equine IGF1 (K. Otte et al. Gen. Comp. Endocrinol. 1996, 102(1), 11-15), the canine IGF1 (P. Delafontaine et al. Gene 1993, 130, 305-306), the feline IGF1 (WO-A-03/022886), the bovine IGF1 (S. Lien et al. Mamm. Genome 2000, 11(10), 877-882), the porcine
15 IGF1 (M. Muller et al. Nucleic Acids Res. 1990, 18(2), 364), the chicken IGF1 (Y. Kajimoto et al. Mol. Endocrinol. 1989, 3(12), 1907-1913), the turkey IGF1 (GenBank accession number AF074980). The signal peptide from IGF1 may be natural or optimized, in particular optimized by removing cryptic splice sites and/or by adapting the codon usage.

20 As used herein the term "polynucleotide" is used to refer to a polymeric form of nucleotides of any length, which contain deoxyribonucleotides or ribonucleotides.

The present invention further encompasses a vector containing and expressing a polynucleotide encoding a BMP-7 polypeptide operably linked to a promoter element and optionally also linked to an enhancer. In an advantageous embodiment, the promoter is the promoter of the cytomegalovirus (CMV) immediate early gene, preferably from human- or
25 murine-derived CMV. In other embodiments, the enhancers and/or promoters may be selected from among those promoters that are known in the art, and that are suitable for expression of BMP-7 in the vectors of the present invention. Many such promoters are known in the art, and suitable promoters can readily be selected by those of skill in the art. For example, there are various cell and/or tissue specific promoters (e.g., muscle, endothelial cell, liver, somatic cell, and stem cell specific promoters), and various viral promoters and enhancers, and BMP-7
30 promoters, such as those isogenically specific for each animal species. For example, in one embodiment, if the canine BMP-7 is to be expressed in a canine muscle cell, the enhancers and/or promoters specific to canine muscle cells may be used in order to optimize expression of canine BMP-7 for the desired application. Examples of muscle-specific promoters and
35 enhancers have been described are known to one of skill in the art (see, e.g., Li et al., Gene Ther.

5 1999 Dec, 6(12), 2005-11; Li et al., Nat Biotechnol. 1999 Mar, 17(3), 241-5 and Loirat et al.,
Virology. 1999, Jul 20, 260(1), 74-83; the disclosures of which are incorporated by reference in
their entireties).

Promoters and enhancers that may be employed in the present invention include, but are
not limited to the promoters and enhancers of the LTR of Rous sarcoma virus, the TK gene of
10 HSV-1, the early or late promoters of SV40, the adenovirus major late promoter (MLP),
phosphoglycerate kinase genes, metallothionein genes, α -1 antitrypsin genes, albumin genes,
collagenase genes, elastase I genes, β -actin genes, β -globin genes, γ -globin genes, α -fetoprotein
genes, and muscle creatin kinase genes.

In general, it is advantageous to employ a strong promoter functional in eukaryotic cells. The
15 preferred strong promoter is the immediate early cytomegalovirus promoter (CMV-IE) of human
or murine origin, or optionally having another origin such as the rat or guinea pig. The CMV-IE
promoter can comprise the actual promoter part, which may or may not be associated with the
enhancer part. Reference can be made to EP-A-260 148, EP-A-323 597, U.S. Patents Nos.
5,168,062, 5,385,839, and 4,968,615, as well as to PCT Application No. WO87/03905. The
20 CMV-IE promoter is advantageously a human CMV-IE (Boshart M. et al., Cell., 1985, 41, 521-
530) or murine CMV-IE.

In more general terms, the promoter has either a viral or a cellular origin. A strong viral
promoter other than CMV-IE that may be usefully employed in the practice of the invention is
the early/late promoter of the SV40 virus or the LTR promoter of the Rous sarcoma virus. A
25 strong cellular promoter that may be usefully employed in the practice of the invention is the
promoter of a gene of the cytoskeleton, such as e.g. the desmin promoter (Kwissa M. et al.,
Vaccine, 2000, 18, 2337-2344), or the actin promoter (Miyazaki J. et al., Gene, 1989, 79, 269-
277).

Functional sub fragments of these promoters, i.e., portions of these promoters that
30 maintain an adequate promoting activity, are included within the present invention, e.g. truncated
CMV-IE promoters according to PCT Application No. WO98/00166 or U.S. Patent No.
6,156,567 can be used in the practice of the invention. A promoter in the practice of the
invention consequently includes derivatives and sub fragments of a full-length promoter that
35 maintain an adequate promoting activity and hence function as a promoter, preferably promoting
activity substantially similar to that of the actual or full-length promoter from which the

5 derivative or sub fragment is derived, e.g., akin to the activity of the truncated CMV-IE promoters of U.S. Patent No. 6,156,567 to the activity of full-length CMV-IE promoters. Thus, a CMV-IE promoter in the practice of the invention can comprise or consist essentially of or consist of the promoter portion of the full-length promoter and/or the enhancer portion of the full-length promoter, as well as derivatives and sub fragments.

10 Preferably, the plasmids comprise other expression control elements. It is particularly advantageous to incorporate stabilizing sequence(s), e.g., intron sequence(s), preferably the first intron of the hCMV-IE (PCT Application No. WO89/01036), the intron II of the rabbit β -globin gene (van Ooyen et al., *Science*, 1979, 206, 337-344). As to the polyadenylation signal (polyA) for the plasmids and viral vectors other than poxviruses, use can more be made of the poly(A) 15 signal of the bovine growth hormone (bGH) gene (see U.S. Patent No. 5,122,458), or the poly(A) signal of the rabbit β -globin gene or the poly(A) signal of the SV40 virus.

20 The term "vector", as used herein, refers to a recombinant DNA or RNA plasmid or virus that comprises a heterologous polynucleotide to be delivered to a target cell, such as *in vivo*. The heterologous polynucleotide may comprise a sequence of interest for purposes of therapy, and may optionally be in the form of an expression cassette. As used herein, a "vector" need not be capable of replication in the ultimate target cell or subject.

25 The term "recombinant" as used herein means a polynucleotide semisynthetic, or synthetic origin, which either does not occur in nature or is linked to another polynucleotide in an arrangement not found in nature.

30 The term "heterologous" as used herein derived from a genetically distinct entity from the rest of the entity to which it is being compared. For example, a polynucleotide may be placed by genetic engineering techniques into a plasmid or vector derived from a different source, and is thus a heterologous polynucleotide. A promoter removed from its native coding sequence and operatively linked to a coding sequence other than the native sequence is accordingly a heterologous promoter.

The polynucleotides of the invention may comprise additional sequences, such as additional coding sequences within the same transcription unit, controlling elements such as promoters, ribosome binding sites, transcription terminators, polyadenylation sites, additional transcription units under control of the same or different promoters, sequences that permit

5 cloning, expression, homologous recombination, and transformation of a host cell, and any such construct as may be desirable to provide embodiments of this invention.

Elements for the expression of canine BMP-7 are advantageously present in an inventive vector. In a minimum manner, this comprises, consists essentially of, or consists of an initiation codon (ATG), a stop codon and a promoter, and optionally also a polyadenylation sequence for 10 certain vectors such as plasmid and certain viral vectors, e.g., viral vectors other than poxviruses. When the polynucleotide encodes a polypeptide fragment, e.g. canine BMP-7, advantageously, in the vector, an ATG is placed at 5' of the reading frame and a stop codon is placed at 3'. Other elements for controlling expression may be present, such as enhancer sequences, stabilizing sequences, such as intron and signal sequences permitting the secretion of the protein.

15 Methods for making and/or administering a vector or recombinants or plasmid for expression of gene products of genes of the invention *in vivo* can be any desired method, e.g., a method which is by or analogous to the methods disclosed in, or disclosed in documents cited in: U.S. Patent Nos. 4,603,112; 4,769,330; 4,394,448; 4,722,848; 4,745,051; 4,769,331; 4,945,050; 5,494,807; 5,514,375; 5,744,140; 5,744,141; 5,756,103; 5,762,938; 5,766,599; 5,990,091; 5,174,993; 20 5,505,941; 5,338,683; 5,494,807; 5,591,639; 5,589,466; 5,677,178; 5,591,439; 5,552,143; 5,580,859; 6,130,066; 6,004,777; 6,130,066; 6,497,883; 6,464,984; 6,451,770; 6,391,314; 6,387,376; 6,376,473; 6,368,603; 6,348,196; 6,306,400; 6,228,846; 6,221,362; 6,217,883; 6,207,166; 6,207,165; 6,159,477; 6,153,199; 6,090,393; 6,074,649; 6,045,803; 6,033,670; 6,485,729; 6,103,526; 6,224,882; 6,312,682; 6,348,450 and 6; 312,683; U.S. patent application 25 Serial No. 920,197, filed October 16, 1986; WO 90/01543; WO91/11525; WO 94/16716; WO 96/39491; WO 98/33510; EP 265785; EP 0 370 573; Andreansky et al., Proc. Natl. Acad. Sci. USA 1996, 93:11313-11318; Ballay et al., EMBO J. 1993;4:3861-65; Felgner et al., J. Biol. Chem. 1994;269, 2550-2561; Frolov et al., Proc. Natl. Acad. Sci. USA 1996, 93, 11371-11377; Graham, Tibtech 1990, 8, 85-87; Grunhaus et al., Sem. Virol. 1992, 3, 237-52; Ju et al., 30 Diabetologia 1998, 41, 736-739; Kitson et al., J. Virol. 1991, 65, 3068-3075; McClements et al., Proc. Natl. Acad. Sci. USA 1996, 93, 11414-11420; Moss, Proc. Natl. Acad. Sci. USA 1996, 93, 11341-11348; Paoletti, Proc. Natl. Acad. Sci. USA 1996, 93, 11349-11353; Pennock et al., Mol. Cell. Biol. 1984, 4, 399-406; Richardson (Ed), Methods in Molecular Biology 1995, 39, "Baculovirus Expression Protocols," Humana Press Inc.; Smith et al. (1983) Mol. Cell. Biol. 35 1983, 3, 2156-2165; Robertson et al., Proc. Natl. Acad. Sci. USA 1996, 93, 11334-11340;

5 Robinson et al., *Sem. Immunol.* 1997, 9, 271; and Roizman, *Proc. Natl. Acad. Sci. USA* 1996, 93, 11307-11312. Thus, the vector in the invention can be any suitable recombinant virus or virus vector, such as a poxvirus (e.g., vaccinia virus, avipox virus, canarypox virus, fowlpox virus, raccoonpox virus, swinepox virus, etc.), adenovirus (e.g., human adenovirus, canine adenovirus), herpesvirus (e.g. canine herpesvirus), baculovirus, retrovirus, etc. (as in documents 10 incorporated herein by reference); or the vector can be a plasmid. The cited and incorporated herein by reference documents, in addition to providing examples of vectors useful in the practice of the invention.

According to one embodiment of the invention, the expression vector is a viral vector, in particular an *in vivo* expression vector. In an advantageous embodiment, the expression vector is 15 an adenovirus vector. Advantageously, the adenovirus is a human adenovirus type 5 (hAd5) vector, an E1-deleted and / or an E3-deleted adenovirus.

In one particular embodiment the viral vector is a poxvirus, e.g. a vaccinia virus or an attenuated vaccinia virus, (for instance, MVA, a modified Ankara strain obtained after more than 570 passages of the Ankara vaccine strain on chicken embryo fibroblasts; *see* Stickl & 20 Hochstein-Mintzel, *Munch. Med. Wschr.*, 1971, 113, 1149-1153; Sutter et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1992, 89, 10847-10851; available as ATCC VR-1508; or NYVAC, *see* U.S. Patent No. 5,494,807, for instance, Examples 1 to 6 and *et seq* of U.S. Patent No. 5,494,807 which discuss the construction of NYVAC, as well as variations of NYVAC with additional ORFs deleted from the Copenhagen strain vaccinia virus genome, as well as the insertion of 25 heterologous coding nucleic acid molecules into sites of this recombinant, and also, the use of matched promoters; *see also* WO96/40241), an avipox virus or an attenuated avipox virus (e.g., canarypox, fowlpox, dovepox, pigeonpox, quailpox, ALVAC or TROVAC; *see*, e.g., U.S. Patent No. 5,505,941, 5,494,807), swinepox, raccoonpox, camelpox, or myxomatosis virus.

According to another embodiment of the invention, the poxvirus vector is a canarypox 30 virus or a fowlpox virus vector, advantageously an attenuated canarypox virus or fowlpox virus. In this regard, is made to the canarypox available from the ATCC under access number VR-111. Attenuated canarypox viruses are described in U.S. Patent No. 5,756,103 (ALVAC) and PCT application N° WO01/05934. Numerous fowlpox virus vaccination strains are also available, e.g. the DIFTOSEC CT strain marketed by MERIAL and the NOBILIS VARIOLE vaccine marketed

5 by INTERVET; and, reference is also made to U.S. Patent No. 5,766,599 which pertains to the attenuated fowlpox strain TROVAC.

For information on the method to generate recombinants thereof and how to administer recombinants thereof, the skilled artisan can refer documents cited herein and to PCT application N° WO90/12882, e.g., as to vaccinia virus mention is made of U.S. Patents Nos. 4,769,330, 10 4,722,848, 4,603,112, 5,110,587, 5,494,807, and 5,762,938 *inter alia*; as to fowlpox, mention is made of U.S. Patents Nos. 5,174,993; 5,505,941 and US-5,766,599 *inter alia*; as to canarypox mention is made of U.S. Patent No. 5,756,103 *inter alia*; as to swinepox mention is made of U.S. Patent No. 5,382,425 *inter alia*; and, as to raccoonpox, mention is made of PCT application N° WO00/03030 *inter alia*.

15 When the expression vector is a vaccinia virus, insertion site or sites for the polynucleotide or polynucleotides to be expressed are advantageously at the thymidine kinase (TK) gene or insertion site, the hemagglutinin (HA) gene or insertion site, the region encoding the inclusion body of the A type (ATI); see also documents cited herein, especially those pertaining to vaccinia virus. In the case of canarypox, advantageously the insertion site or sites 20 are ORF(s) C3, C5 and/or C6; see also documents cited herein, especially those pertaining to canarypox virus. In the case of fowlpox, advantageously the insertion site or sites are ORFs F7 and/or F8; see also documents cited herein, especially those pertaining to fowlpox virus. The insertion site or sites for MVA virus are advantageously as in various publications, including, but not limited to, Carroll M. W. et al., Vaccine, 1997, 15 (4), 387-394; Stittelaar K. J. et al., J. 25 Virol. 2000, 74 (9), 4236-4243; Sutter G. et al., 1994, Vaccine, 12 (11), 1032-1040; and, in this regard it is also noted that the complete MVA genome is described in Antoine G., Virology, 1998, 244, 365-396, which enables the skilled artisan to use other insertion sites or other promoters. Advantageously, the polynucleotide to be expressed is inserted under the control of a specific poxvirus promoter, e.g., the vaccinia promoter 7.5 kDa (Cochran et al., J. Virology, 30 1985, 54, 30-35), the vaccinia promoter I3L (Riviere et al., J. Virology, 1992, 66, 3424-3434), the vaccinia promoter HA (Shida, Virology, 1986, 150, 451-457), the cowpox promoter ATI (Funahashi et al., J. Gen. Virol., 1988, 69, 35-47), the vaccinia promoter H6 (Taylor J. et al., Vaccine, 1988, 6, 504-508; Guo P. et al. J. Virol., 1989, 63, 4189-4198; Perkus M. et al., J. Virol., 1989, 63, 3829-3836), *inter alia*.

5 In a particular embodiment the viral vector is an adenovirus, such as a human adenovirus (HAV) or a canine adenovirus (CAV).

In one embodiment the viral vector is a human adenovirus, in particular a serotype 5 adenovirus, rendered incompetent for replication by a deletion in the E1 region of the viral genome, in particular from about nucleotide 459 to about nucleotide 3510 by reference to the 10 sequence of the hAd5 disclosed in Genbank under the accession number M73260 and in the referenced publication J. Chroboczek et al Virol. 1992, 186, 280-285. The deleted adenovirus is propagated in E1-expressing 293 (F. Graham et al J. Gen. Virol. 1977, 36, 59-72) or PER cells, in particular PER.C6 (F. Falloux et al Human Gene Therapy 1998, 9, 1909-1917). The human adenovirus can be deleted in the E3 region, in particular from about nucleotide 28592 to about 15 nucleotide 30470. The deletion in the E1 region can be done in combination with a deletion in the E3 region (see, e.g. J. Shriver et al. Nature, 2002, 415, 331-335, F. Graham et al Methods in Molecular Biology Vol. 7: Gene Transfer and Expression Protocols Edited by E. Murray, The Human Press Inc, 1991, p 109-128; Y. Ilan et al Proc. Natl. Acad. Sci. 1997, 94, 2587-2592; US6,133,028; US6,692,956; S. Tripathy et al Proc. Natl. Acad. Sci. 1994, 91, 11557-11561; B. 20 Tapnell Adv. Drug Deliv. Rev. 1993, 12, 185-199; X. Danthinne et al Gene Therapy 2000, 7, 1707-1714; K. Berkner Bio Techniques 1988, 6, 616-629; K. Berkner et al Nucl. Acid Res. 1983, 11, 6003-6020; C. Chavier et al J. Virol. 1996, 70, 4805-4810). The insertion sites can be the E1 and/or E3 loci (region) eventually after a partial or complete deletion of the E1 and/or E3 25 regions. Advantageously, when the expression vector is an adenovirus, the polynucleotide to be expressed is inserted under the control of a promoter functional in eukaryotic cells, such as a strong promoter, preferably a cytomegalovirus immediate-early gene promoter (CMV-IE promoter), in particular the enhancer/promoter region from about nucleotide -734 to about nucleotide +7 in M. Boshart et al Cell 1985, 41, 521-530 or the enhancer/promoter region from the pCI vector from PROMEGA Corp. The CMV-IE promoter is advantageously of murine or 30 human origin. The promoter of the elongation factor 1 α can also be used. In one particular embodiment a muscle specific promoter can be used (X. Li et al Nat. Biotechnol. 1999, 17, 241-245). Strong promoters are also discussed herein in relation to plasmid vectors. In one embodiment, a splicing sequence can be located downstream of the enhancer/promoter region. For example, the intron 1 isolated from the CMV-IE gene (R. Stenberg et al J. Virol. 1984, 49, 35 190), the intron isolated from the rabbit or human β -globin gene, in particular the intron 2 from

5 the β -globin gene, the intron isolated from the immunoglobulin gene, a splicing sequence from the SV40 early gene or the chimeric intron sequence isolated from the pCI vector from Promega Corp. comprising the human β -globin donor sequence fused to the mouse immunoglobulin acceptor sequence (from about nucleotide 890 to about nucleotide 1022 in Genbank under the accession number CVU47120). A poly(A) sequence and terminator sequence can be inserted downstream the polynucleotide to be expressed, e.g. a bovine growth hormone gene, in particular from about nucleotide 2339 to about nucleotide 2550 in Genbank under the accession number BOVBMP-7, a rabbit β -globin gene or a SV40 late gene polyadenylation signal.

10

In another embodiment the viral vector is a canine adenovirus, in particular a CAV-2 (see, e.g. L. Fischer et al. Vaccine, 2002, 20, 3485-3497; U.S. Patent No. 5,529,780; U.S. Patent No. 5,688,920; PCT Application No. WO95/14102). For CAV, the insertion sites can be in the E3 region and/or in the region located between the E4 region and the right ITR region (see U.S. Patent No. 6,090,393; U.S. Patent No. 6,156,567). In one embodiment the insert is under the control of a promoter, such as a cytomegalovirus immediate-early gene promoter (CMV-IE promoter) or a promoter already described for a human adenovirus vector. A poly(A) sequence and terminator sequence can be inserted downstream the polynucleotide to be expressed, e.g. a bovine growth hormone gene or a rabbit β -globin gene polyadenylation signal.

20

In another particular embodiment the viral vector is a herpesvirus such as a canine herpesvirus (CHV) or a feline herpesvirus (FHV). For CHV, the insertion sites may be in particular in the thymidine kinase gene, in the ORF3, or in the UL43 ORF (see U.S. Patent No. 6,159,477). In one embodiment the polynucleotide to be expressed is inserted under the control of a promoter functional in eukaryotic cells, advantageously a CMV-IE promoter (murine or human). In one particular embodiment a promoter regulated by hypoxia, e.g. the promoter HRE described in K. Boast et al Human Gene Therapy 1999, 13, 2197-2208), can be used. A poly(A) sequence and terminator sequence can be inserted downstream the polynucleotide to be expressed, e.g. bovine growth hormone or a rabbit β -globin gene polyadenylation signal.

30

According to a yet further embodiment of the invention, the expression vector is a plasmid vector or a DNA plasmid vector, in particular an *in vivo* expression vector. In a specific, non-limiting example, the pVR1020 or 1012 plasmid (VICAL Inc.; Luke C. et al., Journal of Infectious Diseases, 1997, 175, 91-97; Hartikka J. et al., Human Gene Therapy, 1996, 7, 1205-35 1217) can be utilized as a vector for the insertion of a polynucleotide sequence. The pVR1020

5 plasmid is derived from pVR1012 and contains the human tPA signal sequence. In one embodiment the human tPA signal comprises from amino acid Met(1) to amino acid Ser(23) or Ala(28) in Genbank under the accession number HUMTPA14. In another specific, non-limiting example, the plasmid utilized as a vector for the insertion of a polynucleotide sequence can contain the signal peptide sequence of equine IGF1 from amino acid Met(24) to amino acid
10 Ala(48) in Genbank under the accession number U28070.

The term plasmid covers any DNA transcription unit comprising a polynucleotide according to the invention and the elements necessary for its *in vivo* expression in a cell or cells of the desired host or target; and, in this regard, it is noted that a supercoiled or non-supercoiled, circular plasmid, as well as a linear form, are intended to be within the scope of the invention.
15 Each plasmid comprises or contains or consists essentially of, in addition to the polynucleotide encoding the pre-proBMP-7, the proBMP-7 or the mature BMP-7 polypeptide, the BMP-7 polypeptide being preferably from canine origin, variant, analog or fragment, operably linked to a promoter or under the control of a promoter or dependent upon a promoter.


The present invention also relates to a pharmaceutical composition comprising a vector
20 expressing *in vivo* under appropriate or suitable conditions or in a suitable host cell. The pharmaceutical compositions can comprise, consist essentially of, or consist of one or more vectors, e.g., expression vectors, such as *in vivo* expression vectors, comprising, consisting essentially or consisting of and expressing one or more polynucleotides encoding a BMP-7 polypeptide, optionally fused with a BMP-7, IGF-1 or tPA signal peptide, in a pharmaceutically
25 or veterinarilly acceptable carrier, excipient or vehicle. Advantageously, the vector contains, consists essentially of, or consists of and expresses at least one polynucleotide encoding a canine BMP-7 polypeptide, optionally fused with a BMP-7, IGF-1 or tPA signal peptide, in a pharmaceutically or veterinarilly acceptable carrier, excipient or vehicle. Thus, according to an embodiment of the invention, the other vector or vectors in the composition comprises a
30 polynucleotide that encodes, and under appropriate circumstances expresses one or more other proteins, polypeptides or peptides than the canine BMP-7 polypeptide.

Compositions containing one or more vectors containing, consisting essentially of or consisting of polynucleotides encoding, and advantageously expressing, advantageously *in vivo*, a canine BMP-7 peptide or fusion protein.

5 In an advantageous embodiment, the invention provides for the administration of a therapeutically effective amount of a formulation for the delivery and expression of a BMP-7 polypeptide in a target cell. Determination of the therapeutically effective amount is routine experimentation for one of ordinary skill in the art. In one embodiment, the formulation comprises an expression vector comprising a polynucleotide that expresses BMP-7 polypeptide
10 and a pharmaceutically or veterinarily acceptable carrier, vehicle or excipient. In an advantageous embodiment, the pharmaceutically or veterinarily acceptable carrier, vehicle or excipient facilitates transfection and/or improves preservation of the vector.

15 The pharmaceutically or veterinarily acceptable carriers or vehicles or excipients are well known to the one skilled in the art. For example, a pharmaceutically or veterinarily acceptable carrier or vehicle or excipient can be water or a 0.9% NaCl (e.g., saline) solution or a phosphate buffer. Other pharmaceutically or veterinarily acceptable carrier or vehicle or excipients that can be used for methods of this invention include, but are not limited to, poly(L-glutamate) or polyvinylpyrrolidone. The pharmaceutically or veterinarily acceptable carrier or vehicle or excipients may be any compound or combination of compounds facilitating the administration of
20 the vector, increasing the level of expression or increasing the duration of expression. Doses and dose volumes are herein discussed in the general description and can also be determined by the skilled artisan from this disclosure read in conjunction with the knowledge in the art, without any undue experimentation.

25 The cationic lipids containing a quaternary ammonium salt which are advantageously but not exclusively suitable for plasmids, are advantageously those having the following formula:



30 in which R_1 is a saturated or unsaturated straight-chain aliphatic radical having 12 to 18 carbon atoms, R_2 is another aliphatic radical containing 2 or 3 carbon atoms and X is an amine or

5 hydroxyl group, e.g. the DMRIE. In another embodiment the cationic lipid can be associated with a neutral lipid, e.g. the DOPE.

Among these cationic lipids, preference is given to DMRIE (N-(2-hydroxyethyl)-N,N-dimethyl-2,3-bis(tetradecyloxy)-1-propane ammonium; PCT Application No. WO96/34109), wherein the cationic lipid can be advantageously associated with a neutral lipid, advantageously 10 DOPE (dioleoyl-phosphatidyl-ethanol amine; Behr J. P., 1994, Bioconjugate Chemistry, 5, 382-389), to form DMRIE-DOPE.

Advantageously, the plasmid mixture with the excipient is formed extemporaneously and advantageously contemporaneously with administration of the preparation or shortly before administration of the preparation; for instance, shortly before or prior to administration, the 15 plasmid- excipient mixture is formed, advantageously so as to give enough time prior to administration for the mixture to form a complex, e.g. between about 10 and about 60 minutes prior to administration, such as approximately 30 minutes prior to administration.

When DOPE is present, the DMRIE:DOPE molar ratio is advantageously about 95: about 5 to about 5:about 95, more advantageously about 1: about 1, e.g., 1:1.

20 The DMRIE or DMRIE-DOPE adjuvant:plasmid weight ratio can be between about 50: about 1 and about 1: about 10, such as about 10: about 1 and about 1:about 5, and advantageously about 1: about 1 and about 1: about 2, e.g., 1:1 and 1:2.

In a specific embodiment, the pharmaceutical composition is directly administered *in vivo*, and the encoded product is expressed by the vector in the host. The methods to deliver *in vivo* 25 a vector encoding a BMP-7 polypeptide, advantageously the canine BMP-7 polypeptide (see, e.g., U.S. Patent No. 6,423,693; EP-A-1 052 286, EP-A-1 205 551, U.S. Patent Application 2004/0057941, PCT Application No. WO9905300 and Draghia-Akli et al., Mol Ther. 2002 Dec, 6(6), 830-6; the disclosures of which are incorporated by reference in their entireties) can be modified to deliver the BMP-7 polypeptide, of the present invention to a dog or a cat. The *in vivo* 30 delivery of a vector encoding and expressing the BMP-7 described herein can be accomplished by one of ordinary skill in the art given the teachings of the above-mentioned references.

Advantageously, the pharmaceutical compositions and/or formulations according to the invention comprise or consist essentially of or consist of an effective quantity of one or more expression vectors to elicit a therapeutic response as discussed herein; and, an effective quantity

5 can be determined from this disclosure, including the documents incorporated herein, and the knowledge in the art, without undue experimentation.

In the case of therapeutic and/or pharmaceutical compositions based on a plasmid vector, a dose can comprise, consist essentially of or consist of, in general terms, about in 1 μ g to about 2000 μ g, advantageously about 50 μ g to about 1000 μ g and more advantageously from about 10 100 μ g to about 800 μ g of plasmid expressing BMP-7 polypeptide. When the pharmaceutical compositions based on a plasmid vector is administered with electrotransfer the dose of plasmid is generally between about 0.1 μ g and 1mg, advantageously between about 1 μ g and 100 μ g, advantageously between about 2 μ g and 50 μ g.

In an advantageous embodiment the pharmaceutical composition comprising a plasmid vector(s) according to the invention is administered preferably by intramuscular route with electrotransfer to improve the uptake of the vector by the host cells. The features of the electrotransfer alternatively or in combination are: (1) a mono or bipolar electric fields, preferably unipolar; (2) an electric field varying from 10 to 250 V / cm, preferably 50 to 200 V / cm; (3) an electric pulse duration of 10 to 50 msec, preferably of 15 to 25 msec; (4) an interval inter pulse varying from 10 to 990 msec, preferably from 50 to 250 msec; (5) a frequency varying from 1 to 50 Hz, preferably from 4 to 20 Hz, most preferably from 6 to 10 Hz; (6) a number of pulses varying from 1 to 15, preferably 4 to 10; (7) a duration of treatment will vary between 0.1 and 5 sec, preferably between 0.5 and 2.5 sec, most preferably between 0.75 and 1.5 sec; (8) the electrodes can be either invasive or non invasive; (9) the number of electrotransfer per treatment will be comprised between 1 and 10, preferably between 1 and 5 and most preferably between 1 and 2, the frequency of treatments will be established based on induced plasma concentrations of BMP-7 polypeptide; (10) the electrotransfer can be applied with or without anaesthesia or sedation. The electrotransfer can be performed also directly on the kidneys.

The pharmaceutical composition comprising plasmid vector(s) or adenovirus vector(s) can be alternatively administer by sonoporation: (1) the conditions are defined in order to avoid shearing induced by ultrasounds exposure; the plasmid(s) can be protected by polymers, preferably by cationic polymers; (2) a commercial contrast agents used in echocardiography (e.g., PESDA perfluorocarbon or Optison) can be used to improve efficacy, based on acoustic cavitation mechanisms (or others); (3) the route of administration is preferably intramuscular or intravascular (intra arterial) that is of interest to target an internal organ like the kidneys; (4) the

5 diagnostic pulsed US is better than continuous wave system; (5) the efficacy is enhanced when plasmid(s) is complexed with cationized gelatine.

Alternatively to enhance *in vivo* gene delivery with minimal tissue damage the pharmaceutical composition can be administered using a femtosecond infrared laser (LBGT technology).

10 The pharmaceutical composition can be also administered by vascular delivery. The pharmacodynamics-based plasmid DNA gene delivery method based on the change of the hydrodynamics of blood circulation in the recipient animals following the injection of a large volume of DNA solution within a short period of time. It has been demonstrated that the delivery of naked DNA through intraportal or intrahepatic vein injection result in high level of gene expression. The specific expression in the mammalian kidney can be achieved following direct 15 injection into the inferior vena cava (IVC). Through this procedure, expression in the kidney was 10- to 1000-fold higher than in other organs.

20 The dose volumes can be between about 0.1 and about 2 ml, advantageously between about 0.2 and about 1 ml. These doses and dose volumes are suitable for the treatment of canines and other mammalian target species such as equines and felines.

25 The therapeutic and/or pharmaceutical composition contains per dose from about 10^4 to about 10^{11} , advantageously from about 10^5 to about 10^{10} and more advantageously from about 10^6 to about 10^9 viral particles of recombinant adenovirus expressing BMP-7 polypeptide. In the case of therapeutic and/or pharmaceutical compositions based on a poxvirus, a dose can be between about 10^2 pfu and about 10^9 pfu. The pharmaceutical composition contains per dose from about 10^5 to 10^9 , advantageously from about 10^6 to 10^8 pfu of poxvirus or herpesvirus recombinant expressing BMP-7 polypeptide.

30 The dose volume of compositions for target species that are mammals, e.g., the dose volume of canine compositions, based on viral vectors, e.g., non-poxvirus-viral-vector-based compositions, is generally between about 0.1 to about 2.0 ml, preferably between about 0.1 to about 1.0 ml, and more preferably between about 0.5 ml to about 1.0 ml.

35 The present invention contemplates at least one administration to an animal of an efficient amount of the therapeutic composition made according to the invention. The animal may be male, female, pregnant female and newborn. This administration may be via various routes including, but not limited to, intramuscular (IM), subcutaneous (SC), intravascular (IV) or

5 intrarenal injection. Alternative routes to reach the kidneys are: renal artery, injection into the renal subcapsular space, retrograde injection from the ureter or parenchymal injection.

The therapeutic composition according to the invention can also be administered by a needle free apparatus (as, for example with a Pigjet, Biojector or Vitajet apparatus (BIOJECT, Oregon, USA)). Another approach to administer plasmid compositions is to use electrotransfer
10 (see, e.g. S. Tollefson et al. Vaccine, 2002, 20, 3370-3378; S. Tollefson et al. Scand. J. Immunol., 2003, 57, 229-238; S. Babiuk et al., Vaccine, 2002, 20, 3399-3408; PCT Application No. WO99/01158). In an advantageous embodiment, the animal is a mammal. In a more advantageous embodiment, the mammal is a dog or a cat.

15 It should be understood by one of skill in the art that the disclosure herein regarding administration of the compositions of the invention is provided by way of example, and that the present invention is not limited to the specific examples described. From the disclosure herein, and from the knowledge in the art, the skilled artisan can determine the number of administrations, the administration route, and the doses to be used for each administration of the compositions of the present invention without any undue experimentation.

20 In a preferred embodiment, the present invention relates to the use of, and to compositions comprising, a viral vector or a plasmid vector encoding and capable of expressing, a canine pre-proBMP-7, a canine proBMP-7, a canine BMP-7 mature polypeptide, or a variant, derivative or fragment thereof, for the treatment and/or prevention of ARF or CRF. However, in other embodiments of the invention, the methods and compositions disclosed herein may be used
25 to treat and/or prevent other diseases and conditions, including, but not limited to, other kidney conditions, disorders and diseases, anorexia, weight loss, dehydratation, depression, vomiting, polyuria and/or polydipsia.

30 In a preferred embodiment the invention relates to the use of the pharmaceutical compositions according to the present invention to treat mammals presenting an increase in their serum creatinine concentration and/or an increase in their BUN concentration, or an increase in their urine specific gravity.

Advantageously a cat is treated when the plasma creatinine concentration is higher than 1.9 mg/dl and/or when the plasma urea nitrogen concentration is higher than 35 mg/dl.
35 Advantageously a dog is treated when the plasma creatinine concentration is higher than 1.6 mg/dl and/or when the plasma urea nitrogen concentration is higher than 30 mg/dl.

5 The invention will now be further described by way of the following non-limiting examples.

Example: 1 construction of the plasmid pNB292

10 The codon-optimized canine BMP7 open reading frame (“ORF”) consists of 1296 bp and encodes a 431 amino acid polypeptide (SEQ ID N0: 2). The codon-optimized cDNA, encoding the polypeptide sequence of SEQ ID N0: 3, and flanked by unique SalI and XbaI restriction sites, was synthesized from overlapping oligonucleotides, assembled by hybridization and cloned into the pCR-Script vector (Invitrogen) to generate plasmid pPCR-Script050876 (Figure 1).

15 The DNA fragment corresponding to the ORF of interest was excised using SalI and XbaI digestions and further cloned into the pVR1012 plasmid (J. Hartikka et al. Human Gene Therapy 1996, 7, 1205-1217) to generate the pNB292 plasmid (Figure 2) in which the expression of the codon-optimized canine BMP-7 is driven by the cytomegalovirus immediate early (CMV IE) promoter/enhancer. The nucleotide sequence of the pNB292 plasmid is that of SEQ ID N0: 10. The pNB292 plasmid was transformed into DH5 α *E. coli* bacteria and 20 subsequently purified using a commercial kit as recommended by the manufacturer (QIAGEN). Final plasmid concentrations were 2 mg/ml in TE buffer.

25 The transient *in vitro* expression of the polypeptide encoded by the pNB292 plasmid was confirmed and observed after transfection of CHO-K1 cells, using Lipofectamin 2000 (INVITROGEN). CHO-K1 cells at 90 % confluence in 6 cm diameter plates were transfected with 5 μ g plasmid and 10 μ l lipofectamine each, according to the manufacturer’s instructions. After transfection, cells were cultivated in MEM-glutamax medium containing 1% foetal calf serum for 24 hours. Cells grown on glass coverslips were washed with PBS, incubated for 10 min in cold acetone for additional fixing and permeabilisation, and again washed in PBS. Recombinant protein production was analysed by indirect immunofluorescence, using an anti-30 human BMP7 polyclonal serum (ABCAM, Cambridge UK). The immunochemical method confirmed that the pre-proBMP-7 polypeptide encoded by pBN292 was expressed in CHO-K1 cells.

Example 2: Therapeutic effect of BMP-7 plasmid-based gene therapy:

5 A study was conducted in rats to demonstrate the ability of BMP-7 gene therapy to reduce the intensity of tubulo-interstitial lesions associated with the evolution of an experimental unilateral ureteral obstruction (UUO) model of chronic renal failure.

20 male Sprague-Dawley rats weighting approximately 200 g at the initiation of the study were purchased from IFFACREDO (L'Arbresle, France). The maximum and minimum of both 10 temperature and hygrometry of the room were recorded daily. The target temperature and hygrometry range were 20-24°C and 20-70 %, respectively. Light was provided using an automatic timer in cycles of 12 hours light and 12 hours dark. Only healthy rats were included in the study. Rats were allocated randomly to 4 groups of 5 animals each (Groups 1 to 4).

15 Unilateral ureteral obstruction (UUO) was performed on rats from groups 2, 3 and 4, using an established procedure (R. Chevalier et al., Kidney Int. 2000, 57, 882-890). Briefly, rats were anaesthetized by intramuscular injection of tiletamine-zolazepam (ZOLETIL® 100 – 20 to 50 mg/kg – VIRBAC, France). The abdomen was clipped free of fur and the ventral skin was scrubbed with providone iodine. A medial incision of the skin and the abdominal lining was performed. The left ureter was exposed and occluded by tightening the tubing with two 5.0 silk 20 sutures approximately 5 mm away from each other. The suture of the abdominal lining and skin was performed using a silk thread (Silk dec. 0, ETHICON, France). The rats in group 2 were sham-operated, i.e. these animals had their ureters surgically exposed and manipulated, but not ligated. The rats in group 1 were kept as a control, with no surgery performed.

25 The plasmid gWIZ-SEAP® expressing the control transgene SEAP was purchased from GTS Inc. (San Diego, USA) and used as a placebo. The pNB292 plasmid was constructed according to example 1. Final plasmid concentrations were 2 mg/ml in TE buffer.

30 Individual animals were treated at two days prior to surgery (D-2) and five days after surgery (D+5), D0 being the day of surgery. An intramuscular pre-treatment with 100 µl of hyaluronidase at 30 U/100 µl was performed on each targeted muscle two hours prior to the injection of plasmids. Rats were subsequently anaesthetized (by intramuscular injection of tiletamine-zolazepam: ZOLETIL® 100 – 20 to 50 mg/kg – VIRBAC, France) and half a dose of plasmid solution (i.e., 200 µL) was administrated by intramuscular (IM) injection into each *tibialis cranialis* muscle region at D-2 and into each *semi-membranosus* muscle region at D+5. Each injection of 200 µl corresponded to half a dose of plasmid, i.e., 400 µg. Each plasmid-

5 injected rat received a total amount of 800 µg of DNA per day of treatment. The following table recapitulates volumes and masses of plasmid injected

Table 2: Plasmid injections

	Group 1		Group 2		Group 3		Group 4	
	D-2	D+5	D-2	D+5	D-2	D+5	D-2	D+5
<i>Tibialis cranialis</i> left	---	---	---	---	200 µl (0.4 mg)	---	200 µl (0.4 mg)	---
<i>Tibialis cranialis</i> right	---	---	---	---	200 µl (0.4 mg)	---	200 µl (0.4 mg)	---
<i>Semi membranous</i> left	---	---	---	---	200 µl (0.4 mg)	---	200 µl (0.4 mg)	200 µl (0.4 mg)
<i>Semi membranous</i> right	---	---	---	---	200 µl (0.4 mg)	---	200 µl (0.4 mg)	200 µl (0.4 mg)

The specific plasmid compositions administered for each group are specified in table 3.

Table 3: Plasmid compositions administered

	Plasmid composition per dose (400 µL)			
	Group 1	Group 2	Group 3	Group 4
pNB292	---	---	400 µg	
gWIZ-SEAP	---	--	400 µg	800 µg

10 Within the five minutes following plasmid intramuscular delivery, electrotransfer (ET) was applied to each injected muscle using non-invasive plaque electrodes (approximately 0.8 cm² each) in the presence of conductive gel between the skin and the electrodes. The inter-electrode distance was measured to be approximately 0.8 cm. A train of 8 electric pulses of 20 msec each was applied at a frequency of 8 Hz over 1.3 sec. The applied voltage was 140 V

15 targeting a field of 175 V/cm.

All rats were euthanized 13 days after surgery (D+13). One half of each left kidney was fixed in 10 % buffered formalin for histopathological analysis. After fixation, each sample was dehydrated in alcohol solutions of increasing concentration, cleared in isoparaffin H and

5 embedded in paraffin. Embedded samples were cut into 5 μm sections using a microtome (MICROM[®], France). Four sections per site were prepared and stained with Hematoxylin-Eosin-Safranin ("HES") and Masson Trichrome. Histological sections were observed using a microscope (ECLIPSE E600) fitted with x2, x4, x10, x25 and x40 objectives. Renal morphological injury, as characterized by tubular dilatation with epithelial atrophy and
10 interstitial expansion with matrix deposition, was scored in a blind fashion based on a scale of 0 (absent), 1 (mild), 2 (moderate), 3 (limited) and 4 (severe). The overall mean scores and the frequency of each grading were calculated based on individual values, which were determined on 10 fields per rat, 6 rats per group.

15 It was found that plasmid-expressed BMP-7 attenuated renal interstitial fibrosis 13 days post unilateral ureteral obstruction (UUO).

Figure 3 provides histograms of the frequency of lesion grades in the control versus treated groups. No alteration of the renal tissue could be observed in any of the 5 rats of the non-obstructed control group 1, all rats maintaining totally normal kidney architecture graded as "0". In contrast, 4 out of 5 rats of the obstructed but non-treated control group 2 had severe lesions at
20 a grade of 4. A single rat in this group scored at a grade of 3, demonstrating the severity of the experimental model. All 4 out of the 4 rats in the SEAP-treated group (group 4) also presented severe lesions scored at grade 4, thus confirming the severity of the challenge in this group treated with a non-relevant transgene. In contrast, 4 out 5 rats in the BMP-7 treated group 3 (group 3) had a lesion score of 3 with only one rat in this group with severe lesions graded 4.
25 Therefore the proportion of severe (grade 4) lesions in the BMP-7-treated group was 20 % as compared to 80 % in the untreated control group (group 2) and 100% in the placebo treated group (group 4) (Fig. 4).

This data clearly demonstrates the therapeutic effect of BMP-7 plasmid-based gene therapy in a very severe experimental model of tubulo-interstitial nephritis.

30 The invention is further described by the following numbered paragraphs:

1. A method of treating a mammalian subject suffering from, or at risk of developing, renal failure, comprising, administering to said mammalian subject a therapeutically effective amount of a plasmid containing a nucleic acid sequence encoding a BMP-7 polypeptide operatively linked to a promoter, wherein the BMP-7 polypeptide is expressed in vivo in the
35 mammalian subject.

5 2. The method according to paragraph 1, wherein the mammalian subject is selected from the group consisting of cats and dogs.

 3. The method according to paragraph 1 wherein the mammalian subject is a dog.

 4. The method according to paragraph 1 wherein the mammalian subject is a cat.

 5. The method according to paragraph 1 wherein the mammalian subject is suffering 10 from, or are at risk from acute renal failure.

 6. The method according to paragraph 1 wherein the mammalian subjects are suffering from, or are at risk from chronic renal failure.

 7. The method according to paragraph 1, wherein the BMP-7 polypeptide is selected from the group consisting of a pre-pro BMP-7 polypeptide, a pro-BMP-7 polypeptide, and a 15 mature BMP-7 polypeptide.

 8. The method according to paragraph 1, wherein the nucleic acid sequence encoding the BMP-7 polypeptide is selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, and fragments, variants, derivatives and homologs thereof that encode polypeptides having BMP-7 activity.

20 9. The method according to paragraph 1, wherein the BMP-7 polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO: 3 and fragments, variants, derivatives and homologs thereof that have BMP-7 activity.

 10. The method according to paragraph 1, wherein the BMP-7 polypeptide comprises a signal peptide.

25 11. The method according to paragraph 10, wherein the signal peptide is selected from the group consisting of the BMP-7 signal sequence, the IGF-1 signal sequence, and the tPA signal sequence.

 12. The method according to paragraph 10, wherein the signal peptide is encoded by a nucleotide sequence selected from the group consisting of SEQ ID NO: 4, SEQ ID NO: 6, SEQ 30 ID NO: 8, SEQ ID NO: 11, and fragments, variants, derivatives and homologs thereof that encode peptides having signal peptide activity.

 13. The method according to paragraph 10, wherein the signal peptide has an amino acid sequence selected from the group consisting of SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 12, and fragments, variants, derivatives and homologs thereof that have signal 35 peptide activity.

5 14. The method according to paragraph 1 wherein the promoter is selected from the group consisting of a CMV IE promoter, a RSV promoter, an HSV-1 TK promoter, a SV40 early promoter, a SV40 late promoter, an adenovirus major late promoter, a phosphoglycerate kinase gene promoter, a metallothionein gene promoter, an α -1 antitrypsin gene promoter, an albumin gene promoter, a collagenase gene promoter, an elastase I gene promoter, a β -actin gene

10 10 promoter, a β -globin gene promoter, a γ -globin gene promoter, an α -fetoprotein gene promoter, and a muscle creatin kinase gene promoter.

15 15. The method according to paragraph 1, wherein the plasmid is pNB292 and has the nucleotide sequence of SEQ ID NO: 10.

16. The method according to paragraph 1, wherein the plasmid comprises the nucleic acid sequence encoding the BMP-7 polypeptide inserted into the VR1012 plasmid.

17. A method of treating a canine suffering from, or at risk of developing, renal failure, comprising, administering to said canine a therapeutically effective amount of a plasmid containing a nucleic acid sequence encoding a BMP-7 polypeptide operatively linked to a promoter.

20 18. The method according to paragraph 17, wherein the canine is suffering from, or are at risk from acute renal failure.

25 19. The method according to paragraph 17, wherein canine is suffering from, or are at risk from chronic renal failure.

20 20. The method according to paragraph 17, wherein the BMP-7 polypeptide is selected from the group consisting of a pre-pro BMP-7 polypeptide, a pro-BMP-7 polypeptide, and a mature BMP-7 polypeptide.

30 21. The method according to paragraph 17, wherein the nucleic acid sequence encoding the BMP-7 polypeptide is selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, and fragments, variants, derivatives and homologs thereof that encode polypeptides having BMP-7 activity.

22. The method according to paragraph 17, wherein the BMP-7 polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO: 3 and fragments, variants, derivatives and homologs thereof that have BMP-7 activity.

35 23. The method according to paragraph 17, wherein the BMP-7 polypeptide comprises a signal peptide.

5 24. The method according to paragraph 23, wherein the signal peptide is selected from the group consisting of the BMP-7 signal sequence, the IGF-1 signal sequence, and the tPA signal sequence.

10 25. The method according to paragraph 23, wherein the signal peptide is encoded by a nucleotide sequence selected from the group consisting of SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 11, and fragments, variants, derivatives and homologs thereof that encode peptides having signal peptide activity.

15 26. The method according to paragraph 23, wherein the signal peptide has an amino acid sequence selected from the group consisting of SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 12, and fragments, variants, derivatives and homologs thereof that have signal peptide activity.

20 27. The method according to paragraph 17 wherein the promoter is selected from the group consisting of a CMV IE promoter, a RSV promoter, an HSV-1 TK promoter, a SV40 early promoter, a SV40 late promoter, an adenovirus major late promoter, a phosphoglycerate kinase gene promoter, a metallothionein gene promoter, an α -1 antitrypsin gene promoter, an albumin gene promoter, a collagenase gene promoter, an elastase I gene promoter, a β -actin gene promoter, a β -globin gene promoter, a γ -globin gene promoter, an α -fetoprotein gene promoter, and a muscle creatin kinase gene promoter.

25 28. The method according to paragraph 17, wherein the plasmid is pNB292 and has the nucleotide sequence of SEQ ID NO: 10.

30 29. The method according to paragraph 17, wherein the plasmid comprises the nucleic acid sequence encoding the BMP-7 polypeptide inserted into the VR1012 plasmid.

30 30. A method of preventing the development of renal failure in a mammalian subject at risk thereof, comprising administering to said mammalian subject a prophylactically effective amount of a plasmid vector containing a nucleic acid sequence encoding a BMP-7 polypeptide operatively linked to a promoter.

30 31. The method according to paragraph 30 wherein the mammalian subject is at risk from acute renal failure.

30 32. The method according to paragraph 30 wherein the mammalian subject is at risk from chronic renal failure.

5 33. The method according to paragraph 30, wherein the BMP-7 polypeptide is selected from the group consisting of a pre-pro BMP-7 polypeptide, a pro-BMP-7 polypeptide, and a mature BMP-7 polypeptide.

10 34. The method according to paragraph 30, wherein the nucleic acid sequence encoding the BMP-7 polypeptide is selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, and fragments, variants, derivatives and homologs thereof that encode polypeptides having BMP-7 activity.

15 35. The method according to paragraph 30, wherein the BMP-7 polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO: 3 and fragments, variants, derivatives and homologs thereof that have BMP-7 activity.

20 36. The method according to paragraph 30, wherein the BMP-7 polypeptide comprises a signal peptide.

25 37. The method according to paragraph 30, wherein the signal peptide is selected from the group consisting of the BMP-7 signal sequence, the IGF-1 signal sequence, and the tPA signal sequence.

30 38. The method according to paragraph 37, wherein the signal peptide is encoded by a nucleotide sequence selected from the group consisting of SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 11, and fragments, variants, derivatives and homologs thereof that encode peptides having signal peptide activity.

35 39. The method according to paragraph 37, wherein the signal peptide has an amino acid sequence selected from the group consisting of SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 12, and fragments, variants, derivatives and homologs thereof that have signal peptide activity.

40 40. The method according to paragraph 30, wherein the promoter is selected from the group consisting of a CMV IE promoter, a RSV promoter, an HSV-1 TK promoter, a SV40 early promoter, a SV40 late promoter, an adenovirus major late promoter, a phosphoglycerate kinase gene promoter, a metallothionein gene promoter, an α -1 antitrypsin gene promoter, an albumin gene promoter, a collagenase gene promoter, an elastase I gene promoter, a β -actin gene promoter, a β -globin gene promoter, a γ -globin gene promoter, an α -fetoprotein gene promoter, and a muscle creatin kinase gene promoter.

5 41. The method according to paragraph 30, wherein the plasmid is pNB292 and has the nucleotide sequence of SEQ ID NO: 10.

10 42. The method according to paragraph 30, wherein the plasmid comprises the nucleic acid sequence encoding the BMP-7 polypeptide inserted into the VR1012 plasmid.

15 43. A recombinant plasmid vector comprising a nucleic acid sequence encoding a BMP-7 polypeptide operatively linked to a promoter.

20 44. The recombinant plasmid vector according to paragraph 43, wherein the BMP-7 polypeptide is selected from the group consisting of a pre-pro BMP-7 polypeptide, a pro-BMP-7 polypeptide, and a mature BMP-7 polypeptide.

25 45. The recombinant plasmid vector according to paragraph 43, wherein the nucleic acid sequence encoding the BMP-7 polypeptide is selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, and fragments, variants, derivatives and homologs thereof that encode polypeptides having BMP-7 activity.

30 46. The recombinant plasmid vector according to paragraph 43, wherein the BMP-7 polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO: 3 and fragments, variants, derivatives and homologs thereof that have BMP-7 activity.

35 47. The recombinant plasmid vector according to paragraph 43, wherein the BMP-7 polypeptide comprises a signal peptide.

40 48. The recombinant plasmid vector according to paragraph 47, wherein the signal peptide is selected from the group consisting of the BMP-7 signal sequence, the IGF-1 signal sequence, and the tPA signal sequence.

45 49. The recombinant plasmid vector according to paragraph 47, wherein the signal peptide is encoded by a nucleotide sequence selected from the group consisting of SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 11, and fragments, variants, derivatives and homologs thereof that encode peptides having signal peptide activity.

50 50. The method according to paragraph 47, wherein the signal peptide has an amino acid sequence selected from the group consisting of SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 12, and fragments, variants, derivatives and homologs thereof that have signal peptide activity.

55 51. The method according to paragraph 43, wherein the promoter is selected from the group consisting of a CMV IE promoter, a RSV promoter, an HSV-1 TK promoter, a SV40 early

5 promoter, a SV40 late promoter, an adenovirus major late promoter, a phosphoglycerate kinase gene promoter, a metallothionein gene promoter, an α -1 antitrypsin gene promoter, an albumin gene promoter, a collagenase gene promoter, an elastase I gene promoter, a β -actin gene promoter, a β -globin gene promoter, a γ -globin gene promoter, an α -fetoprotein gene promoter, and a muscle creatin kinase gene promoter.

10 52. The recombinant plasmid vector according to paragraph 43, wherein the plasmid is pNB292 and has the nucleotide sequence of SEQ ID NO: 10.

53. The recombinant plasmid vector according to paragraph 43, wherein the plasmid comprises the nucleic acid sequence encoding the BMP-7 polypeptide inserted into the VR1012 plasmid.

15 54. A pharmaceutical composition comprising a recombinant plasmid vector according to anyone of paragraphs to 43 to 53, and at least one pharmaceutically or veterinarily acceptable carrier, excipient, or vehicle.

* * *

Having thus described in detail preferred embodiments of the present invention, it is to be
20 understood that the invention defined by the above paragraphs is not to be limited to particular details set forth in the above description as many apparent variations thereof are possible without departing from the spirit or scope of the present invention.

5 **WHAT IS CLAIMED IS:**

1. A recombinant plasmid vector comprising a nucleic acid sequence encoding a BMP-7 polypeptide operatively linked to a promoter, wherein the BMP-7 polypeptide is selected from the group consisting of a pre-pro BMP-7 polypeptide, a pro-BMP-7 polypeptide, and a mature BMP-7 polypeptide; or
10 wherein the BMP-7 polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO: 3 and fragments, variants, derivatives and homologs thereof that have BMP-7 activity.
2. The recombinant plasmid vector according to claim 1, wherein the nucleic acid sequence encoding the BMP-7 polypeptide is selected from the group consisting of SEQ ID NO: 1, SEQ 15 ID NO: 2, and fragments, variants, derivatives and homologs thereof that encode polypeptides having BMP-7 activity.
3. The recombinant plasmid vector according to claim 1, wherein the BMP-7 polypeptide comprises a signal peptide selected from the group consisting of the BMP-7 signal sequence, the 20 IGF-1 signal sequence, and the tPA signal sequence; or
25 wherein the signal peptide is encoded by a nucleotide sequence selected from the group consisting of SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 11, and fragments, variants, derivatives and homologs thereof that encode peptides having signal peptide activity; or
wherein the signal peptide has an amino acid sequence selected from the group consisting of SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 12, and fragments, variants, derivatives and homologs thereof that have signal peptide activity
4. The recombinant plasmid vector according to claim 1, wherein the plasmid is pNB292 and has the nucleotide sequence of SEQ ID NO: 10; or
30 wherein the plasmid comprises the nucleic acid sequence encoding the BMP-7 polypeptide inserted into the VR1012 plasmid.
5. A pharmaceutical composition comprising a recombinant plasmid vector according to claim 1, and optionally at least one pharmaceutically or veterinarily acceptable carrier, excipient, or vehicle.
6. A method of treating a mammalian subject suffering from, or at risk of developing, renal 35 failure, comprising administering to said mammalian subject a therapeutically effective amount

5 of a plasmid containing a nucleic acid sequence encoding a BMP-7 polypeptide operatively linked to a promoter.

7. The method according to claim 6, wherein the BMP-7 polypeptide is selected from the group consisting of a pre-pro BMP-7 polypeptide, a pro-BMP-7 polypeptide, and a mature BMP-7 polypeptide; or

10 wherein the BMP-7 polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO: 3 and fragments, variants, derivatives and homologs thereof that have BMP-7 activity.

8. The method according to claim 6, wherein the nucleic acid sequence encoding the BMP-7 polypeptide is selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, and

15 fragments, variants, derivatives and homologs thereof that encode polypeptides having BMP-7 activity.

9. The method according to claim 6, wherein the BMP-7 polypeptide comprises a signal peptide, wherein the signal peptide is selected from the group consisting of the BMP-7 signal sequence, the IGF-1 signal sequence, and the tPA signal sequence; or

20 wherein the signal peptide has an amino acid sequence selected from the group consisting of SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 12, and fragments, variants, derivatives and homologs thereof that have signal peptide activity; or

wherein signal peptide is encoded by a nucleotide sequence selected from the group consisting of SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 11, and fragments, variants, derivatives and homologs thereof that encode peptides having signal peptide activity.

25 10. The method according to claim 6, wherein the plasmid is pNB292 and has the nucleotide sequence of SEQ ID NO: 10; or

wherein the plasmid comprises the nucleic acid sequence encoding the BMP-7 polypeptide inserted into the VR1012 plasmid.

30 11. The method according to claim 6, wherein the BMP-7 polypeptide is expressed in vivo in the mammalian subject.

12. The method according to claim 6, wherein the mammalian subject is selected from the group consisting of felines and canines.

13. The method according to claim 12, wherein the mammalian subject is a canine.

5 14. A method of preventing the development of renal failure in a mammalian subject at risk thereof, comprising administering to said mammalian subject a prophylactically effective amount of a plasmid vector containing a nucleic acid sequence encoding a BMP-7 polypeptide operatively linked to a promoter.

10 15. The method according to claim 14, wherein the BMP-7 polypeptide is selected from the group consisting of a pre-pro BMP-7 polypeptide, a pro-BMP-7 polypeptide, and a mature BMP-7 polypeptide; or

wherein the BMP-7 polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO: 3 and fragments, variants, derivatives and homologs thereof that have BM16. The method according to claim 14, wherein the nucleic acid sequence

15 encoding the BMP-7 polypeptide is selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, and fragments, variants, derivatives and homologs thereof that encode polypeptides having BMP-7 activity.

20 17. The method according to claim 14, wherein the BMP-7 polypeptide comprises a signal peptide wherein the signal peptide is selected from the group consisting of the BMP-7 signal sequence, the IGF-1 signal sequence, and the tPA signal sequence; or

wherein the signal peptide is encoded by a nucleotide sequence selected from the group consisting of SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 11, and fragments, variants, derivatives and homologs thereof that encode peptides having signal peptide activity; or

25 18. The method according to claim 14, wherein the signal peptide has an amino acid sequence selected from the group consisting of SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 12, and fragments, variants, derivatives and homologs thereof that have signal peptide activity.

18. The method according to claim 14, wherein the plasmid is pNB292 and has the nucleotide sequence of SEQ ID NO: 10; or

30 19. The method according to claim 14, wherein the plasmid comprises the nucleic acid sequence encoding the BMP-7 polypeptide inserted into the VR1012 plasmid.

19. The method according to claim 14, wherein the mammalian subject is selected from the group consisting of felines and canines.

20. The method according to claim 19, wherein the mammalian subject is a canine. P-7 activity.



FIGURE 1: 050876pPCR-Script PLASMID MAP

FIGURE 2: pNB292 PLASMID MAP



Description of the plasmid pNB292:

Optimised canine BMP-7 gene: nucleotides 1651-2949

Kanamycin resistance gene (Tn903): nucleotides 4837-5652

CMV IE promoter: nucleotides 1-683

CMV IE 5' UTR: nucleotides 684-804

Intron A: nucleotides 805-1626

bGH transcriptional terminator / polyA: nucleotides 2979-3525

Origin of replication: nucleotides 3707-4214

FIGURE 3





FIGURE 4



INTERNATIONAL SEARCH REPORT

International application No

PCT/US2006/044048

A. CLASSIFICATION OF SUBJECT MATTER
 INV. C07K14/51
 ADD. A61K48/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, EMBASE, WPI Data, Sequence Search

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>GREGORY KATE E ET AL: "The prodomain of BMP-7 targets the BMP-7 complex to the extracellular matrix." THE JOURNAL OF BIOLOGICAL CHEMISTRY 29 JUL 2005, vol. 280, no. 30, 29 July 2005 (2005-07-29), pages 27970-27980, XP002421582 ISSN: 0021-9258 page 27970, column 2, paragraph 5 – page 27971, column 1; figure 1</p> <p style="text-align: center;">-/-</p>	1,5

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

23 February 2007

Date of mailing of the international search report

07/03/2007

Name and mailing address of the ISA/
 European Patent Office, P.B. 5018 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Deleu, Laurent

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2006/044048

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FRANCESCHI R T ET AL: "Gene therapy for bone formation: In vitro and in vivo osteogenic activity of an adenovirus expressing BMP7" JOURNAL OF CELLULAR BIOCHEMISTRY 2000 UNITED STATES, vol. 78, no. 3, 2000, pages 476-486, XP002421583 ISSN: 0730-2312 page 478, column 1, paragraph 3 page 477, column 1	1,3,5
Y	----- WO 02/47713 A (SULZER BIOLOG INC [US]) 20 June 2002 (2002-06-20) page 12, line 19 - line 22; claims 1-7	6,7,9, 11-15, 17,19,20
Y	----- WO 2004/019876 A (CURIS INC [US]; WASHINGTON UNIVERSITY IN ST LO [US]; CHARETTE MARC F [) 11 March 2004 (2004-03-11) page 57, line 29 - page 58, line 6; claims 1,47-52; figures 18,20,23; table 1	6,7, 9-15, 17-20
Y	----- HARTIKKA J: "AN IMPROVED PLASMID DNA EXPRESSION VECTOR FOR DIRECT INJECTION INTO SKELETAL MUSCLE" HUMAN GENE THERAPY, MARY ANN LIEBERT, NEW YORK, NY, US, vol. 7, no. 10, 20 June 1996 (1996-06-20), pages 1205-1217, XP002050079 ISSN: 1043-0342 page 1206 page 1215	10,18
A	----- VUKICEVIC S ET AL: "Osteogenic protein-1 (bone morphogenetic protein-7) reduces severity of injury after ischemic acute renal failure in rat" JOURNAL OF CLINICAL INVESTIGATION, NEW YORK, NY, US, vol. 102, no. 1, 1 July 1998 (1998-07-01), pages 202-214, XP002297818 ISSN: 0021-9738 page 213; tables 1-4	1-20
P,A	----- BRIGHT CORINNE ET AL: "In vivo evaluation of plasmid DNA encoding OP-1 protein for spine fusion" SPINE, vol. 31, no. 19, September 2006 (2006-09), pages 2163-2172, XP009079493 ISSN: 0362-2436 page 2164	4,10

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box II.1

Although claims 6, 7, 9-15 and 17-20 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box II.2

Claims Nos.: 2, 8, 16 (complete) and 1, 3-5, 7, 9, 10, 15, 17, 18 (partially)

The claims refer to nucleotide sequences selected from the group of SEQ IDs. However the description does not disclose these SEQ IDs but merely refers to the nucleotide sequences of the canine pre-proBMP7 polypeptide, the codon-optimized nucleotide sequence of the canine pre-proBMP-7 polypeptide, the nucleotide sequence of the signal peptide from tPA, from the equine IGF-1 or the canine IGF-1, the nucleotide sequence of the pBN292 plasmid. The search has been restricted to nucleotide sequences encoding the polypeptides designated BMP-7, IGF-1 signal sequence and tPA signal sequence.

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure. If the application proceeds into the regional phase before the EPO, the applicant is reminded that a search may be carried out during examination before the EPO (see EPO Guideline C-VI, 8.5), should the problems which led to the Article 17(2) declaration be overcome.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2006/044048

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 6, 7, 9-15 and 17-20 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

The additional search fees were accompanied by the applicant's protest.
 No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2006/044048

Patent document cited in search report	Publication date	Patent family member(s)			Publication date
WO 0247713	A 20-06-2002	CA 2446832 A1	JP 2004520295 T	20-06-2002	08-07-2004
		US 2002173453 A1			21-11-2002
WO 2004019876	A 11-03-2004	AU 2003268219 A1	CA 2497048 A1	19-03-2004	11-03-2004
		EP 1578360 A2			28-09-2005
		JP 2006516020 T			15-06-2006