

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 November 2009 (12.11.2009)

(10) International Publication Number
WO 2009/137681 A1

(51) International Patent Classification:
C07D 471/04 (2006.01) *A61P 27/00* (2006.01)
C07D 471/06 (2006.01) *A61P 9/00* (2006.01)
C07D 471/16 (2006.01) *A61K 31/4745* (2006.01)

(21) International Application Number:
PCT/US2009/043149

(22) International Filing Date:
7 May 2009 (07.05.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/051,533 8 May 2008 (08.05.2008) US
12/433,978 1 May 2009 (01.05.2009) US

(71) Applicant (for all designated States except US): **ALLEGIAN, INC.** [US/US]; 2525 Dupont Drive, T2-7H, Irvine, CA 92612 (US).

(71) Applicants and
(72) Inventors: **SINHA, Santosh, C.** [IN/US]; 199 Sklar Street, Ladera Ranch, CA 92694 (US). **BHAT, Smita** [US/US]; 62 Barcelona, Irvine, CA 92614 (US). **CHOW, Ken** [US/US]; 20 Tidal Surf, Newport Coast, CA 92657 (US). **BEARD, Richard, L.** [US/US]; 2341 Azure Avenue, Newport Beach, CA 92660 (US). **DONELLO, John, E.** [US/US]; 34041 Pequito Drive, Dana Point, CA 92629 (US). **GARST, Michael, E.** [US/US]; 2627 Raqueta Drive, Newport Beach, CA 92660 (US).

(74) Agents: **WURST, John** et al.; Allergan, Inc., 2525 Dupont Drive, Irvine, CA 92612 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: THERAPEUTICALLY USEFUL SUBSTITUTED HYDROPYRIDO [3,2,1-IJ] QUINOLINE COMPOUNDS



(57) Abstract: Disclosed herein are compounds represented by the structural formula (I): therapeutic methods, compositions, and medicaments related thereto are also disclosed.

WO 2009/137681 A1

18517 PCT (AP)

THERAPEUTICALLY USEFUL SUBSTITUTED HYDROPYRIDO [3,2,1-ij] QUINOLINE COMPOUNDS

5

By Inventors

Santosh C. Sinha, Smita Bhat, Ken Chow, Richard L. Beard, John Donello and Michael E. Garst.

Related Application


This invention claims priority under 35 U.S.C §119 US Patent Application 10 12/433,978, filed May 1, 2009, to which claims priority to US Provisional Application 61/051,533, filed May 8, 2008, the disclosure of which are incorporated by reference herein in their entirety.

Field of the Invention

The present invention provides novel substituted hydropyrido[3,2,1-ij]quinoline compounds, and their uses in medicaments for the treatment of mammals with diseases and conditions that are alleviated by sphingosine-1-phosphate (S1P) receptors modulation.

Background of the Invention

Sphingosine is a compound having the chemical structure shown in the 20 general formula described below, in which Y¹ is hydrogen. It is known that various sphingolipids, having sphingosine as a constituent, are widely distributed in the living body including on the surface of cell membranes of cells in the nervous system.



25 A sphingolipid is one of the lipids having important roles in the living body. A disease called lipidosis is caused by accumulation of a specified sphingolipid in the body. Sphingolipids present on cell membranes function to regulate cell growth; participate in the development and differentiation of cells; function in nerves; are involved in the infection and malignancy of cells; etc. Many of the physiological 30 roles of sphingolipids remain to be solved. Recently the possibility that ceramide, a derivative of sphingosine, has an important role in the mechanism of cell signal

18517 PCT (AP)

transduction has been indicated, and studies about its effect on apoptosis and cell cycle have been reported.

Sphingosine-1-phosphate is an important cellular metabolite, derived from ceramide that is synthesized de novo or as part of the sphingomyeline cycle (in 5 animals cells). It has also been found in insects, yeasts and plants.

The enzyme, ceramidase, acts upon ceramides to release sphingosine, which is phosphorylated by sphingosine kinase, a ubiquitous enzyme in the cytosol and endoplasmic reticulum, to form sphingosine-1-phosphate. The reverse reaction can occur also by the action of sphingosine phosphatases, and the 10 enzymes act in concert to control the cellular concentrations of the metabolite, which concentrations are always low. In plasma, such concentration can reach 0.2 to 0.9 μ M, and the metabolite is found in association with the lipoproteins, especially the HDL. It should also be noted that sphingosine-1-phosphate formation is an essential step in the catabolism of sphingoid bases.

15 Like its precursors, sphingosine-1-phosphate is a potent messenger molecule that perhaps uniquely operates both intra- and inter-cellularly, but with very different functions from ceramides and sphingosine. The balance between these various sphingolipid metabolites may be important for health. For example, within the cell, sphingosine-1-phosphate promotes cellular division (mitosis) as 20 opposed to cell death (apoptosis), which it inhibits. Intracellularly, it also functions to regulate calcium mobilization and cell growth in response to a variety of extracellular stimuli. Current opinion appears to suggest that the balance between sphingosine-1-phosphate and ceramide and/or sphingosine levels in cells is critical for their viability.


25 In common with the lysophospholipids, especially lysophosphatidic acid, with which it has some structural similarities, sphingosine-1-phosphate exerts many of its extra-cellular effects through interaction with five specific G protein-coupled receptors on cell surfaces, known as endothelium differentiation gene receptors ("Edg" or "S1P" receptors).

30 S1P3 receptor is one of the receptors interacting with sphingosine-1-phosphate. S1P3 receptor, alone or together with other S1P receptors, involves in many critical biological processes, such as the growth of new blood vessels, vascular maturation, cardiac development and immunity, as well as for directed cell movement. S1P3 receptor modulators are needed for therapeutic uses.

18517 PCT (AP)

Summary of the Invention

The compounds of the present invention can be represented by the
5 structural formula:



wherein m is an integer of 0, or 1; n is an integer of 0, 1, 2, or 3; each Y is independently carbon (C) or nitrogen (N); Z and X are each independently selected from the group of oxygen (O), sulfur (S), and amine moiety NR^N; B is selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, cyano and X-B together being a heterocyclic ring or ring system; R and R³ are each independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano; each R¹ is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxycarbonyl, aminocarbonyl, aminocarbonxyl, alkylcarboxyl, alkyl amide, amino, alkylamino, and cyano; each R² is independently selected from the group consisting of hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxo, oxycarbonyl, carboxyl, alkyl carboxylate, alkyl amide, aminocarbonyl, amino, alkylamino, and cyano; each R^N is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl, hydroxyl,

18517 PCT (AP)

alkoxyl, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano; including their alternate solid forms, tautomers, stereoisomers, enantiomers, diastereomers, prodrugs, and pharmaceutically acceptable salts, hydrates and solvates; and


5 provided that when Y is carbon, and Z and X are both oxygen, R² is not oxo, or R¹ and R² are not both phenyl or both methyl at the same time.

Applicants have discovered that these compounds modulate sphingosine-1-phosphate (S1P) receptor activity, in particular inhibit S1P3 receptor. These compounds are useful for the treatment of mammals, including human beings, with a range of conditions and diseases that are alleviated by S1P modulation, such as ocular diseases and conditions (glaucoma, elevated intraocular pressure, dry eye, and optical neurodegenerative diseases), cardiovascular diseases and conditions, pulmonary diseases and conditions, skin conditions, angiogenesis, inflammation, sepsis and pain.

15

Detailed Description of the Invention

Disclosed herein are compounds represented by the structural formula:



wherein m is an integer of 0 or 1; n is an integer of 0, 1, 2, or 3; each Y is independently carbon (C) or nitrogen (N); Z and X are each independently selected from the group of oxygen (O), sulfur (S), and amine moiety NR^N ; B is selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, cyano and X-B together being a heterocyclic ring/ring system.

R and R³ are each independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo,

18517 PCT (AP)

halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano; each R¹ is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, 5 halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxycarbonyl, aminocarbonyl, aminocarbonyl, alkylcarboxyl, alkyl amide, amino, alkylamino, and cyano; each R² is independently selected from the group consisting of hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxo, oxycarbonyl, carboxyl, 10 alkyl carboxylate, alkyl amide, aminocarbonyl, amino, alkylamino, and cyano; each R^N is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano; including 15 their alternate solid forms, tautomers, stereoisomers, enantiomers, diastereomers, prodrugs, and pharmaceutically acceptable salts, hydrates and solvates; and optionally, provided that when Y is carbon, and Z and X are both oxygen, R² is not oxo, or R¹ and R² are not both phenyl or both methyl at the same time.

It has been discovered that the compounds of the present invention listed in 20 this patent application modulate sphingosine-1-phosphate (S1P) receptor activity and in particular the S1P3 receptor. These compounds are useful for the treatment of mammals, including humans, with a range of conditions and diseases that are alleviated by S1P modulation: not limited to treating glaucoma, elevated intraocular pressure, ischemic neuropathies, optic neuropathy, pain, visceral pain, corneal 25 pain, headache pain, migraine, cancer pain, back pain, irritable bowel syndrome pain, muscle pain and pain associated with diabetic neuropathy, the treatment of diabetic retinopathy, other retinal degenerative conditions, dry eye, angiogenesis and wounds. Other uses include:

Ocular applications:

30 Retinopathy of prematurity, diabetic retinopathy, optic neuropathy, glaucomatous retinopathy, macular degeneration, choroidal neovascularization, ocular wound healing, and retinal edema;

Cardiovascular applications:

18517 PCT (AP)

Congestive heart failure, cardiac arrhythmia, atherosclerosis, and bradycardia;

Pulmonary applications:

5 Asthma, chronic obstructive pulmonary disease, acute lung injury, acute respiratory distress syndrome, idiopathic pulmonary fibrosis, and ventilation-induced lung injury; and,

Skin applications:

Scar-less wound healing, scar-less skin-wound and cosmetic healing.

For the purposes of this disclosure, "treat," "treating," or "treatment" refer to 10 the diagnosis, cure, mitigation, treatment, or prevention of disease or other undesirable condition.

The compounds of present invention may be identified either by their 15 chemical structures and/or chemical names. If the chemical structure and the chemical name conflict, the chemical structure is determinative of the identity of the compound.

The compounds of the invention may contain one or more chiral centers and/or double bonds and therefore, may exist as stereoisomers, such as double-bond isomers, enantiomers or diastereomers. Accordingly, the chemical structures depicted herein encompass all possible enantiomers and stereoisomers, including 20 the stereoisomerically pure form and enantiomeric and stereoisomeric mixtures. The compounds of the invention may also exist in several tautomeric forms, including but not limiting to, the enol form, the keto form and mixtures thereof. Accordingly, the chemical structures depicted herein encompass all possible 25 tautomeric forms. The compounds of the invention also include isotopically labeled compounds where one or more atoms have an atomic mass different from the atomic mass conventionally found in nature.

Further, the compounds of the invention should be construed broadly to include their pharmaceutically acceptable salts, prodrugs, alternate solid forms, non-covalent complexes, and combinations thereof, unless otherwise indicated.

30 A pharmaceutically acceptable salt is any salt of the parent compound that is suitable for administration to a mammal, including human. A pharmaceutically acceptable salt also refers to any salt which may form *in vivo* as a result of administration of an acid, another salt, or a prodrug which is converted into an acid

18517 PCT (AP)

or salt. A salt comprises one or more ionic forms of the compound, such as a conjugate acid or base, associated with one or more corresponding counter-ions. Salts can form from or incorporate one or more deprotonated acidic groups (e.g. carboxylic acids), one or more protonated basic groups (e.g. amines), or both (e.g. 5 zwitterions).

A prodrug is a compound which is converted to a therapeutically active compound after administration. For example, conversion may occur by hydrolysis of an ester group or some other biologically labile groups. Prodrug preparation is well known in the art. For example, "Prodrugs and Drug Delivery Systems," which 10 is a chapter in Richard B. Silverman, *Organic Chemistry of Drug Design and Drug Action*, 2d Ed., Elsevier Academic Press: Amsterdam, 2004, pp. 496-557, provides further detail on this subject.

Alternate solid forms are different solid forms than those that may result from practicing the procedures described herein. For example, alternate solid 15 forms may be polymorphs, different kinds of amorphous solid forms, glasses, and the like.

Non-covalent complexes are complexes that may form between the compound and one or more additional chemical species that do not involve a covalent bonding interaction between the compound and the additional chemical 20 species. They may or may not have a specific ratio between the compound and the additional chemical species. Examples include solvates, hydrates, charge transfer complexes, and the like.

Hydrocarbyl consists of carbon and hydrogen, wherein each carbon has 4 covalent bonds and each hydrogen has a single bond to a carbon atom. 25 "Hydrocarbyl fragments" has the same meaning as "hydrocarbyl," but is merely used for convenience for counting purposes. For example, one or more hydrocarbyl fragments means, 1, 2, or more distinct parts that each consists of hydrocarbyl, which may be interrupted by another moiety. For example, a functional group may be attached to 2 distinct hydrocarbyl fragments.

30 Hydrocarbyl includes alkyl, alkenyl, alkynyl, aryl containing only hydrogen and carbon, and combinations thereof. Hydrocarbyl may be linear, branched, cyclic (aromatic or non-aromatic), or combinations thereof, which can be further substituted.

18517 PCT (AP)

Alkyl is a hydrocarbyl having no double bonds. Examples include methyl, ethyl, propyl isomers, butyl isomers, pentyl isomers, hexyl isomers, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, etc.

Alkenyl is a hydrocarbyl having one or more double bonds. Examples 5 include ethenyl, propenyl, butenyl isomers, pentenyl isomers, hexenyl isomers, cyclopentenyl, cyclohexenyl, etc.

Alkynyl is a hydrocarbyl having one or more triple bonds. Examples include ethynyl, propynyl, butynyl isomers, pentynyl isomers, hexynyl isomers, cyclopentynyl, cyclohexynyl, etc.

10 Aryl is a substituted or unsubstituted aromatic ring or ring system. It can be hydrocarbon-aryl or heteroaryl. Examples of hydrocarbon-aryl include substituted and unsubstituted phenyl, naphthyl, and biphenyl. Such aryl group can be bonded to other moieties within the molecule at any position.

15 Each hydrogen atom has one covalent bond to carbon (C), nitrogen (N), oxygen (O), or sulfur (S).



















Halo or halo atoms are fluorine (F), chlorine (Cl), bromine (Br), and iodine (I). Each halo atom forms a single bond to a carbon atom. Halohydrocarbyl is a hydrocarbyl having one or more F, Cl, Br, or I as substituents.

20 Heterohydrocarbyl refers to a hydrocarbyl as defined above with at least one non-carbon atom(s) presented at the backbone, including but not limiting to, oxygen (O), sulfur (S), nitrogen (N), phosphor (P), and halo atoms. Heterohydrocarbyl may be linear, branched, cyclic (aromatic or non-aromatic), or combinations thereof, which can be further substituted.

























25 Examples of heterohydrocarbyl include: $-R^{10}-G^1-R^{11}$, $-R^{10}-HI$, $-G^1-R^{10}$, $-G^1-R^{10}-HI$, $G^1-R^{10}-G^2$, and $G^1-R^{10}-G^2-R^{11}$, wherein R^{10} and R^{11} are independently hydrocarbyl or hydrogen (provided that hydrogen is attached to only one C, N, O, or S atom), G^1 and G^2 are independently functional groups, and HI is halo.

30 Additional examples of heterohydrocarbyl are depicted below, wherein R^{10} , R^{11} , R^{12} , and R^{13} are independently hydrocarbyl or hydrogen. Other possibilities exist, but are not depicted here.

18517 PCT (AP)



18517 PCT (AP)




18517 PCT (AP)

Heteroaryl is one type of heterohydrocarbyl, referring to an aromatic ring or ring system containing at least one hetero atom selected from N, O, S, P, and combinations thereof. Examples of heteroaryl include, but not limit to, pyridine, 5 pyrazine, pyrimidine, pyridazine, triazine, furan, pyrrole, thiophene, imidazole, pyrazole, oxazole, isoxazole, thiazole, isothiazole, oxadiazole, thiadiazole, naphthalene, quinoline, quinoxaline, quinazoline, cinnoline, isoquinoline, benzofuran, indole, benzothiophene, benzimidazole, indazole, benzoxazole, benzisoxazole, benzothiazole, isobenzofuran, isoindole, tetraline, chromane, 10 isochromane, thiochromane, chromene, isochromene, thiochromene, indane, indene, coumarine, coumarinone, and the like, which can be further substituted. Such heteroaryl group can be bonded to other moieties within the molecule at any position.

“Substituted” or “a substituent” is hydrogen, one or more hydrocarbyl 15 fragments, one or more heterohydrocarbyl fragments, one or more halo atoms, one or more functional groups, or combinations thereof. Two or more substituents may themselves form an additional ring or ring system.

A functional group comprises of alkyl, aryl, alkenyl, alkynyl, halo, haloalkyl, hydroxyl, alkoxy, hydroxyalkyl, oxo, alkylcarbonyl, formyl, carboxyl, 20 alkylcarboxylate, alkylamide, aminocarbonyl, amino, cyano, diazo, nitro, thio, sulfoxyl, sulfonyl, phosphate, phosphinate, and one of the moieties depicted below.


18517 PCT (AP)



18517 PCT (AP)

If a functional group is asymmetric, it may be oriented in any way possible. For example, the ester functional group is intended to indicate both of the structures below.


5




In a substituent, one or more hydrocarbyl fragments, one or more heterohydrocarbyl fragments, and/or one or more functional groups may be incorporated into one or more rings or ring systems.

The dashed lines on the functional groups indicate that any nitrogen atom 10 on a functional group may form an additional bond with another carbon atom, a hydrogen atom, or may form a double bond with one of the depicted bonds so that an ammonium or a quaternary ammonium type of functional group is formed. Thus, the dashed line functional groups actually represent a group of individual functional groups. For example, the functional group:

15




represents the following possible structures:



20

Similarly, the functional group:




represents the following possible structures:

18517 PCT (AP)




In one embodiment, compounds of the invention are represented by the structural formula



5

and



wherein m is an integer of 0, or 1; n is an integer of 0, 1, 2, or 3;

10 Z and X are each independently selected from the group of oxygen, sulfur, and amine moiety NR^N ;

B is selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, 15 aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, cyano and X-B together being a heterocyclic ring or ring system;

R and R^3 are each independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl,


18517 PCT (AP)

formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano;

each R¹ is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, 5 halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxycarbonyl, aminocarbonyl, aminocarbonxyl, alkylcarboxyl, alkyl alkylcarbonyl, formyl, oxycarbonyl, aminocarbonyl, aminocarbonxyl, alkylcarboxyl, alkyl amide, amino, alkylamino, and cyano;

each R² is independently selected from the group consisting of hydrocarbyl, 10 heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxo, oxycarbonyl, carboxyl, alkyl carboxylate, alkyl amide, aminocarbonyl, amino, alkylamino, and cyano; each R^N is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl, hydroxyl, 15 alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano; including their alternate solid forms, tautomers, stereoisomers, enantiomers, diastereomers, prodrugs, and pharmaceutically acceptable salts, hydrates and solvates.

In another embodiment, compounds of the invention are represented by the 20 structural formula



wherein

n is 0, 1, 2, or 3;

25 Z is O, S, or NR^N;

X is O, S, or NR^N;

18517 PCT (AP)

B is selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, cyano and X-B together being a heterocyclic ring;

5 R and R³ are each independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano;

10 each R¹ is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxycarbonyl, aminocarbonyl, aminocarbonxyl, alkylcarboxyl, alkyl amide, amino, alkylamino, and cyano;


15 each R² is independently selected from the group consisting of hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxo, oxycarbonyl, carboxyl, alkyl carboxylate, alkyl amide, aminocarbonyl, amino, alkylamino, and cyano;

20 each R^N is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano;

25 including its alternate solid forms, tautomers, stereoisomers, enantiomers, diastereomers, prodrugs, and pharmaceutically acceptable salts, hydrates and solvates.

In yet another embodiment, the compounds are represented by

18517 PCT (AP)



wherein o is an integer of 0, 1, 2, or 3;

Z is O or S;

each R^A is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano;

10 R and R³ are each independently selected from the group consisting of hydrogen and a substituent having a formula C₀₋₁₂H₀₋₃₀N₀₋₃O₀₋₅P₀₋₂S₀₋₃F₀₋₆Cl₀₋₃Br₀₋₃I₀₋₃;

each R^N is independently selected from the group consisting of hydrogen and C_{1-12} hydrocarbyl;

B is selected from the group consisting of hydrogen, a substituent having a formula $C_{0-12}H_{0-30}N_{0-3}O_{0-5}P_{0-2}S_{0-3}F_{0-6}Cl_{0-3}Br_{0-3}I_{0-3}$, wherein if X is NR^N , and X-B being a heterocyclic ring/ ring system. The formula $C_{0-12}H_{0-30}N_{0-3}O_{0-5}P_{0-2}S_{0-3}F_{0-6}Cl_{0-3}Br_{0-3}I_{0-3}$ represents a structure having from 0-12 carbon atoms, from 0-30 hydrogen atoms, from 0-3 nitrogen atoms, from 0-5 oxygen atoms, from 0-2 phosphorous atoms, from 0-3 sulfur atoms, from 0-6 fluorine atoms, from 0-3 chlorine atoms, from 0-3 bromine atoms, and from 0-3 iodine atoms.

20 In yet another embodiment, each R^A is independently hydrogen, alkyl, aryl, alkenyl, alkynyl, halo, haloalkyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, carboxyl, alkyl carboxylate, alkyl amide, aminocarbonyl, amino, cyano, diazo, nitro, thio, sulfoxyl, sulfonyl, phosphate, or phosphinate.


In another embodiment, X is O.

25 In another embodiment, Z is O.

In another embodiment, Z is S.

18517 PCT (AP)

In another embodiment, B is C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, C₁₋₆ haloalkyl, hydroxyalkyl, phenyl, benzyl, furylmethyl, or wherein X-B is morpholino. C₁₋₆ means having from one to six (1-6) carbon atoms. C₁₋₆ haloalkyl is C₁₋₆ alkyl having at least one halo atoms of F, Cl, Br, or I as the substituent. Examples of 5 haloalkyl include -CH₂F, -CH₂CHF₂, -C₃H₆F, -C₄H₈F, -C₅H₁₀F, -C₆H₁₂F, fluorocyclopropyl, fluorocyclobutyl, fluorocyclopentyl, fluorocyclohexyl, -CH₂CH₂Cl, -C₃H₆Cl, -C₄H₈Cl, -C₅H₁₀Cl, -C₆H₁₂Cl, chlorocyclopropyl, chlorocyclobutyl, chlorocyclopentyl, chlorocyclohexyl, -CH₂CH₂Br, -C₃H₆Br, -C₄H₈Br, -C₅H₁₀Br, -C₆H₁₂Br, bromocyclopropyl, bromocyclobutyl, bromocyclopentyl, bromocyclohexyl, 10 -CH₂CH₂I, -C₃H₆I, -C₄H₈I, -C₅H₁₀I, -C₆H₁₂I, iodocyclopropyl, iodocyclobutyl, iodocyclopentyl, and iodocyclohexyl. Morpholino is:




In another embodiment, R^A is hydrogen, C₁₋₁₂ alkyl, C₁₋₁₂ alkenyl, C₁₋₁₂ alkynyl, halo, C₁₋₁₂ halohydrocarbyl, C₁₋₁₂ hydroxyalkyl, C₃₋₁₂ cyclic hydrocarbyl, or 15 heteroaryl. C₁₋₁₂ means having from 1-12 carbon atoms.

In yet another embodiment, X is NR^N.

In another embodiment, the compounds can be represented by the structural formula:

20



wherein each R^A is independently hydrogen, C₁₋₁₂ alkyl, C₁₋₁₂ alkenyl, C₁₋₁₂ alkynyl, halo, C₁₋₁₂ halohydrocarbyl, C₁₋₁₂ hydroxyalkyl, C₃₋₁₂ cyclic hydrocarbyl, or heteroaryl.

25 In another embodiment, R^N is hydrogen, methyl, ethyl, propyl, or isopropyl.

18517 PCT (AP)






In one embodiment, R^N is hydrogen, C₁₋₆ alkyl or phenyl.

In another embodiment, B is hydrocarbyl, as described above.

In yet another embodiment, B is substituted or unsubstituted hydrocarbon-aryl or heterohydrocarbyl. Examples of heteroaryl include pyridine, pyrazine, 5 pyrimidine, pyridazine, triazine, furan, pyrrole, thiophene, imidazole, pyrazole, oxazole, isoxazole, thiazole, isothiazole, oxadiazole, thiadiazole, naphthalene, quinoline, quinoxaline, quinazoline, cinnoline, isoquinoline, benzofuran, indole, benzothiophene, benzimidazole, indazole, benzoxazole, benzisoxazole, benzothiazole, isobenzofuran, isoindole, tetraline, chromane, isochromane, 10 thiochromane, chromene, isochromene, thiochromene, indane, indene, coumarine, coumarinone, and the like.




If the aryl or heteroaryl is substituted, the substituents are the same as those defined above. Examples include alkyl, aryl, alkenyl, alkynyl, halo, haloalkyl, hydroxyl, alkoxy, hydroxylalkyl, alkylcarbonyl, formyl, carboxyl, alkyl carboxylate, 15 alkyl amide, aminocarbonyl, amino, cyano, diazo, nitro, thio, sulfoxyl, sulfonyl, phosphate, phosphinate, and the like.

B may also be a combination of one or more of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted hydrocarbon-aryl, or substituted or unsubstituted heteroaryl. For example, B may have one of the structures shown 20 below:




18517 PCT (AP)



Compounds according to the structural formulas below are also contemplated:




5



18517 PCT (AP)



and



5

wherein o is an integer of 0, 1, 2, or 3; Z is O or S;

each R^A is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl,

10 formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano;

R and R^3 are each independently selected from the group consisting of hydrogen and a substituent having a formula $C_{0-12}H_{0-30}N_{0-3}O_{0-5}P_{0-2}S_{0-3}F_{0-6}Cl_{0-3}Br_{0-3}I_{0-3}$;

18517 PCT (AP)

each R^N is independently selected from the group consisting of hydrogen and C_{1-12} hydrocarbyl;

B is selected from the group consisting of hydrogen, a substituent having a formula $C_{0-12}H_{0-30}N_{0-3}O_{0-5}P_{0-2}S_{0-3}F_{0-6}Cl_{0-3}Br_{0-3}I_{0-3}$, wherein if X is NR^N , and X-B being a

5 heterocyclic ring/ ring system. The formula $C_{0-12}H_{0-30}N_{0-3}O_{0-5}P_{0-2}S_{0-3}F_{0-6}Cl_{0-3}Br_{0-3}I_{0-3}$ represents a structure having from 0-12 carbon atoms, from 0-30 hydrogen atoms, from 0-3 nitrogen atoms, from 0-5 oxygen atoms, from 0-2 phosphorous atoms, from 0-3 sulfur atoms, from 0-6 fluorine atoms, from 0-3 chlorine atoms, from 0-3 bromine atoms, and from 0-3 iodine atoms.

10 In yet another embodiment, each R^A is independently hydrogen, alkyl, aryl, alkenyl, alkynyl, halo, haloalkyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, carboxyl, alkyl carboxylate, alkyl amide, aminocarbonyl, amino, cyano, diazo, nitro, thio, sulfoxyl, sulfonyl, phosphate, or phosphinate.

In another embodiment, X is O.

15 In another embodiment, Z is O.

In another embodiment, Z is S.


In another embodiment, B is C_{1-6} alkyl, C_{1-6} alkenyl, C_{1-6} alkynyl, C_{1-6} haloalkyl, hydroxyalkyl, phenyl, benzyl, furylmethyl, or wherein X-B is morpholino.

C_{1-6} means having from one to six (1-6) carbon atoms. C_{1-6} haloalkyl is C_{1-6} alkyl


20 having at least one halo atoms of F, Cl, Br, or I as the substituent. Examples of haloalkyl include $-CH_2F$, $-CH_2CHF_2$, $-C_3H_6F$, $-C_4H_8F$, $-C_5H_{10}F$, $-C_6H_{12}F$, fluorocyclopropyl, fluorocyclobutyl, fluorocyclopentyl, fluorocyclohexyl, $-CH_2CH_2Cl$, $-C_3H_6Cl$, $-C_4H_8Cl$, $-C_5H_{10}Cl$, $-C_6H_{12}Cl$, chlorocyclopropyl, chlorocyclobutyl, chlorocyclopentyl, chlorocyclohexyl, $-CH_2CH_2Br$, $-C_3H_6Br$, $-C_4H_8Br$, $-C_5H_{10}Br$, $-C_6H_{12}Br$, bromocyclopropyl, bromocyclobutyl, bromocyclopentyl, bromocyclohexyl, $-CH_2CH_2I$, $-C_3H_6I$, $-C_4H_8I$, $-C_5H_{10}I$, $-C_6H_{12}I$, iodocyclopropyl, iodocyclobutyl, iodocyclopentyl, iodocyclohexyl

25
















Specific compounds of the present invention include:






18517 PCT (AP)




18517 PCT (AP)




18517 PCT (AP)




18517 PCT (AP)



18517 PCT (AP)



18517 PCT (AP)



The compounds of the present invention can be combined with at least one other therapeutic agent that is already known the art. The compounds of invention and the other therapeutic agent(s) can act additively, or more preferably, synergistically.

The invention is further defined by reference to the following examples, which describe the preparation schemes and methods for obtaining the compounds of the invention, the assays for testing the biological activities of these compounds. It will be apparent to those skilled in the art that many modifications, both to the preparation schemes and assays, may be practiced without departing from the scope of the invention.

15 **EXAMPLES:**


Organic Synthesis:

Reaction Schemes A, B, C , D and E are examples of the preparation methods for obtaining the compounds of the invention.


20

Reaction Scheme A


18517 PCT (AP)



Reaction Scheme B




Reaction Scheme C



10


Reaction Scheme D



15


Reaction Scheme E

18517 PCT (AP)



Example A

5 Method A1: Preparation of methyl 1,7-diphenyl-1,2,3,5,6,7-hexahydronaphthalene-1,4-dicarboxylic acid amide (272)



10 To a solution 4-Nitroaniline (**Intermediate 1**) (1.8 g, 10 mmol) in acetonitrile (8 mL) was added one equivalent of trifluoroacetic acid (1.14 g, 10 mmol). To this suspension was added with stirring a heterogeneous mixture of styrene (**Intermediate 2**), (5.74 mL, 50 mmol) and 37% formaldehyde solution (4.06 mL, 50 mmol) under argon, which gave a yellow precipitate. The precipitate failed to re-dissolve after 30 min. of stirring at room temperature, so the mixture was heated at reflux under argon for further 30 min, during which time the precipitate re-dissolved. The reaction mixture was cooled to room temperature. The precipitate was filtered and wash with acetonitrile gave yellow solid, 9-nitro-1,7-diphenyl-1,2,3,5,6,7-hexahydronaphthalene (**Intermediate 3**), (1.53 g, 41 %).

15

18517 PCT (AP)

A solution of 9-nitro-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinoline (**Intermediate 3**), (1.2 g, 7.06 mmol), in MeOH (100 mL) was subjected to hydrogenation reaction by the action of 10% Pd/C (120 mg) under H₂ balloon at room temperature for 12 h. The mixture was filtered through Celite and freed of solvent under reduced pressure to get 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-amine (**Intermediate 4**) as a solid, (1.08 g, 98%).


To a solution 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-amine (**Intermediate 4**), (207 mg, 0.608 mmol) in dichloromethane (10 mL) was added three equivalent of triethyl amine (0.252 mL, 1.8 mmol), followed by methyl chloroformate (0.071 mL, 0.91 mmol) under argon at 0 °C. The reaction mixture was then stirred at room temperature for overnight. The mixture was quenched with water (30 mL). The residue was isolated in a typical aqueous workup and purified by MPLC (medium pressure liquid chromatography) using silica gel column with 10 to 15 % EtOAc:Hexane to give methyl 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamate (**272**), (181 mg 75%). ¹H NMR (300 MHz, CDCl₃) δ ppm 2.00 - 2.18 (m, 2 H) 2.22 - 2.39 (m, 2 H) 3.03 - 3.22 (m, 4 H) 3.50 - 3.64 (m, 3 H) 3.54 - 3.65 (m, 3 H) 4.05 - 4.23 (m, 2 H) 6.61 (br. s., 2 H) 7.08 - 7.38 (m, 10 H).

20

Method A2: Preparation of methyl (1S,7S)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamate, (**829**), and methyl (1R,7S)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamate (**353**)

25

18517 PCT (AP)



To a solution 4-Nitroaniline (**1**) (4.1 g, 30 mmol) in acetonitrile (30 mL) was added one equivalent of trifluoroacetic acid (2.3 mL, 30 mmol). To this suspension was added with stirring a heterogeneous mixture of styrene (**2**), (19.4 mL, 150 mmol) and 37% formaldehyde solution (12.2 mL, 150 mmol) under argon gave yellow precipitate. The precipitate had failed to re-dissolve after 30min. of stirring at room temperature, so the mixture was heated at reflux under argon for further 30min, during which time the precipitate re-dissolved. The reaction mixture was cooled to room temperature. After general workup afforded mixture of three 5 intermediates **3** (836 mg), **4** (2 g), and **5** (4.2 g), confirmed by Mass Spectra and ¹HNMR (see ref. John M. Mellor; et al; *Tetrahedron*, **1995**, 6115). These intermediates were then converted into the cycloadduct product by heating at reflux with trifluoroacetic acid in acetonitrile yielded 1,7-dimethyl-9-nitro-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinoline (**6**) (7.036 g, 59%) as a solid. This 10 solid product (**6**) was then separated to *trans* and *cis* isomers by washing with ether gave *trans* (**7**), (4.0 g) and hexane:CH₂Cl₂ gave *cis* (**8**), (2.8 g). 15

A mixture of (1*S*,7*S*)-1,7-dimethyl-9-nitro-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinoline (**7**), (2 g, 5 mmol), in THF (60 mL) was subjected

18517 PCT (AP)

to hydrogenation reaction by the action of 10% Pd/C (200 mg) under H₂ balloon at room temperature for 12 h. The mixture was filtered through Celite and freed of solvent under reduced pressure to get (1S,7S)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-amine (**9**) as a solid, (1.5 g, 100%)

5 on the basis of recovered starting material (**7**), (390 mg).

Following a procedure similar to that for (**9**) gained (1*R*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-amine (**10**) as a solid (2.55 g, 100% yield) from (**8**).

To a solution (1*S*,7*S*)-1,7-dimethyl-9-nitro-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinoline (**7**), (115 mg, 0.31 mmol), in dichloromethane (15 mL) was added three equivalent of triethyl amine (0.129 mL, 0.93 mmol), followed by methyl chloroformate (0.031 mL, 0.406 mmol) under argon at 0 °C. The reaction mixture was then stirred at room temperature for overnight. The mixture was quenched with water (30 mL). The residue was isolated in a typical aqueous workup and purified by MPLC (medium pressure liquid chromatography) using silica gel column with 10 to 15 % EtOAc:Hexane to give methyl (1*S*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamate (**829**), (69 mg 52%). 1H NMR (300 MHz, Acetone-d6) δ ppm 1.72 (s, 6H) 1.89 - 2.03 (m, 2 H) 2.17 - 2.30 (m, 2 H) 2.73 - 2.88 (m, 2 H) 2.92 - 3.03 (m, 2 H) 3.56 (s, 3 H) 7.05 - 7.33 (m, 12 H) 8.02 (br. s., 1 H).

Following a procedure similar to that for (**829**), gained (1*R*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamate (**353**), (130g, 62% yield) from (**10**). 1H NMR (300 MHz, Acetone-d6) δ ppm 1.74 (s, 6 H) 1.98 - 2.10 (m, 2 H) 2.13 - 2.24 (m, 2 H) 2.78 - 2.92 (m, 2 H) 2.92 - 3.05 (m, 2 H) 3.55 (s, 3 H) 7.06 (s, 2 H) 7.12 - 7.34 (m, 10 H) 8.02 (br. s., 1 H).

The following compounds were prepared according to the **Reaction Scheme A** and with the steps as shown in **Example A** above.

30 **Ethyl 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamate, 273**

¹H NMR (300 MHz, CDCl₃) δ ppm 1.11 (t, *J*=7.03 Hz, 3 H) 2.02 - 2.17 (m, 2 H) 2.21 - 2.37 (m, 2 H) 3.04 - 3.17 (m, 4 H) 4.04 (q, 2 H) 4.09 - 4.21 (m, 2 H) 6.61 (br. s., 2 H) 7.08 - 7.39 (m, 10 H)

18517 PCT (AP)

Isobutyl 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 274

¹H NMR (300 MHz, CDCl₃) δ ppm 0.94 (d, *J*=6.74 Hz, 6 H) 1.84 - 2.02 (m, 1 H) 5 2.02 - 2.17 (m, 2 H) 2.20 - 2.37 (m, 2 H) 3.31-3.22 (m, 4 H) 4.19-4.14 (m, 2 H) 6.63 (br. s., 2 H) 7.11 - 7.24 (m, 5 H) 7.25 - 7.37 (m, 5 H)

Propyl 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 275

10 ¹H NMR (300 MHz, CDCl₃) δ ppm 0.85 (t, *J*=6.89 Hz, 3 H) 1.46 - 1.64 (m, 2 H) 2.00 - 2.18 (m, 2 H) 2.21 - 2.37 (m, 2 H) 3.03 - 3.19 (m, 4 H) 3.88 - 4.00 (m, 2 H) 4.09 - 4.23 (m, 2 H) 6.61 (br. s., 2 H) 7.07 - 7.37 (m, 10 H)

Butyl 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 276

15 ¹H NMR (300 MHz, CDCl₃) δ ppm 0.87 (t, *J*=7.0 Hz, 3 H) 1.30 (br. s., 2 H) 1.42 - 1.56 (m, 2 H) 2.02 - 2.17 (m, 2 H) 2.19 - 2.39 (m, 2 H) 3.03 - 3.17 (m, 4 H) 3.91 - 4.03 (m, 2 H) 4.10 - 4.22 (m, 2 H) 6.61 (br. s., 2 H) 7.08 - 7.38 (m, 10 H)

20 **Phenyl 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 277**

¹H NMR (300 MHz, CDCl₃) δ ppm 2.05 - 2.19 (m, 2 H) 2.21 - 2.40 (m, 2 H) 3.04 - 3.20 (m, 4 H), 4.11 - 4.24 (m, 2 H) 6.71 (br. s., 2 H) 6.99 - 7.44 (m, 15 H)

25 **Benzyl 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 278**

¹H NMR (300 MHz, CDCl₃) δ ppm 1.91 - 2.09 (m, 2 H) 2.10 - 2.29 (m, 2 H) 2.93 - 3.13 (m, 4 H) 4.01 - 4.15 (m, 2 H) 4.92 (s, 2 H) 6.56 (br. s., 2 H) 7.01 - 7.32 (m, 15 H)

30

Propyl 1,7-bis(3-fluorophenyl)-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 094

18517 PCT (AP)

1H NMR (300 MHz, CD₃OD) δ ppm 0.89 (t, 7.0 Hz, H) 1.47 - 1.67 (m, 2 H) 1.99 - 2.14 (m, 2 H) 2.19 - 2.37 (m, 2 H) 3.08 (t, *J*=5.71 Hz, 4 H) 3.91 (t, *J*=6.74 Hz, 2 H) 4.17 (t, *J*=6.01 Hz, 2 H) 6.67 (br. s., 2 H) 6.81 - 7.04 (m, 6 H) 7.22 - 7.38 (m, 2 H)

5 **Ethyl 1,7-bis(3-fluorophenyl)-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 093**

1H NMR (300 MHz, CD₃OD) δ ppm 1.15 (t, 7.0 Hz, H) 1.98 - 2.16 (m, 2 H) 2.18 - 2.37 (m, 2 H) 2.96 - 3.18 (m, 4 H) 4.00 (q, *J*=7.13 Hz, 2 H) 4.19 (t, *J*=5.86 Hz, 2 H) 6.68 (br. s., 2 H) 6.77 - 7.01 (m, 6 H) 7.20 - 7.35 (m, 2 H)

10 **Butyl 1,7-bis(3-fluorophenyl)-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 202**

1H NMR (300 MHz, Acetone-d₆) δ ppm 0.86 (t, 7.0 Hz, 6 H) 1.20 - 1.39 (m, 2 H) 2.06 - 2.15 (m, 2 H) 2.21 - 2.39 (m, 2 H) 3.02 - 3.17 (m, 4 H) 3.94 (t, *J*=6.59 Hz, 2 H) 4.22 (t, *J*=5.86 Hz, 2 H) 6.83 (br. s., 2 H) 6.88 - 7.09 (m, 6 H) 7.28 - 7.42 (m, 2 H)

Methyl 1,7-bis(3-fluorophenyl)-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 779

20 **1H NMR (300 MHz, CD₃OD) δ ppm 2.01 - 2.15 (m, 2 H) 2.21 - 2.37 (m, 2 H) 3.09 (t, *J*=5.71 Hz, 4 H) 3.57 (s, 3 H) 4.18 (t, *J*=6.15 Hz, 2 H) 6.67 (s, 2 H) 6.81 - 7.03 (m, 6 H) 7.25 - 7.38 (m, 2 H)**

2-fluoroethyl 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 095

1H NMR (300 MHz, CD₃OD) δ ppm 1.99 - 2.16 (m, 2 H) 2.19 - 2.36 (m, 2 H) 3.00 - 3.16 (m, 4 H) 4.09 - 4.27 (m, 4 H) 4.29 - 4.36 (m, 1 H) 4.46 - 4.52 (m, 1 H) 6.66 (br. s., 2 H) 7.07 - 7.21 (m, 6 H) 7.21 - 7.33 (m, 4 H)

30 **But-3-enyl 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 354**

1H NMR (300 MHz, Acetone-d₆) δ ppm 2.04 - 2.15 (m, 2 H) 2.18 - 2.41 (m, 4 H) 3.02 - 3.18 (m, 4 H) 3.89 - 4.01 (m, 2 H) 4.90 - 5.14 (m, 2 H) 5.69 - 5.88 (m, 1 H) 6.78 (br. s., 2 H) 7.08 - 7.25 (m, 6 H) 7.23 - 7.38 (m, 4 H)

18517 PCT (AP)**But-2-ynyl 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 353**

1H NMR (300 MHz, Acetone-d₆) δ ppm 1.74 (s, 3 H) 2.06 - 2.13 (m, 2 H) 2.19 - 2.34 (m, 2 H) 3.02 - 3.20 (m, 4 H) 4.19 (t, *J*=6.01 Hz, 2 H) 4.52 (q, *J*=2.54 Hz, 2 H) 5 6.78 (s, 2 H) 7.12 - 7.24 (m, 6 H) 7.26 - 7.35 (m, 4 H)

Prop-2-ynyl 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 352

1H NMR (300 MHz, Acetone-d₆) δ ppm 2.05 - 2.15 (m, 2 H) 2.20 - 2.36 (m, 2 H) 10 2.90 (t, *J*=2.49 Hz, 1 H) 3.00 - 3.20 (m, 4 H) 4.19 (t, *J*=6.01 Hz, 2 H) 4.59 (d, *J*=2.34 Hz, 2 H) 6.78 (s, 2 H) 7.11 - 7.24 (m, 6 H) 7.25 - 7.37 (m, 4 H)

4-chlorobutyl 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 206

15 1H NMR (300 MHz, Acetone-d₆) δ ppm 1.66 – 1.96 (m, 4 H) 2.06 - 2.11 (m, 2 H) 2.22 – 2.32 (m, 2 H) 3.04 - 3.17 (m, 4 H) 3.58 (t, *J*=6 Hz, 2H) 3.96 (t, *J*=6.3 Hz, 2H) 4.19 (t, *J*=6.3 Hz, 2 H) 6.77 (s, 2 H) 7.16 - 7.20 (m, 6 H) 7.22 - 7.33 (m, 4 H)

3-chloropropyl 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 205

1H NMR (300 MHz, CD₃OD) δ ppm 1.91 - 2.14 (m, 4 H) 2.20 - 2.34 (m, 2 H) 3.02 - 3.15 (m, 4 H) 4.07 (t, *J*=6.01 Hz, 2 H) 4.11 - 4.21 (m, 2 H) 6.64 (br. s., 2 H) 7.09 - 7.21 (m, 6 H) 7.23 - 7.31 (m, 4 H)

25 Ethyl (1*S*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 941

1H NMR (300 MHz, Acetone-d₆) δ ppm 1.16 (t, *J*=7.18 Hz, 3 H) 1.74 (s, 6 H) 1.90 - 2.01 (m, 2 H) 2.16 - 2.32 (m, 2 H) 2.73 - 2.89 (m, 2 H) 2.90 - 3.03 (m, 2 H) 4.02 (q, *J*=7.13 Hz, 2 H) 7.06 - 7.32 (m, 12 H) 8.02 (br. s., 1 H)

30

Propyl (1*S*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-ylcarbamate, 942

18517 PCT (AP)

1H NMR (300 MHz, Acetone-d6) δ ppm 0.88 (t, $J=7.33$ Hz, 3 H) 1.46 - 1.65 (m, 2 H) 1.72 (s, 6 H) 1.88 - 2.03 (m, 2 H) 2.18 - 2.32 (m, 2 H) 2.70 - 2.88 (m, 2 H) 2.89 - 3.03 (m, 2 H) 3.94 (t, $J=6.59$ Hz, 2 H) 7.06 - 7.33 (m, 12 H) 8.02 (br. s., 1 H)

5 **Butyl (1*S*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamate, 943**

1H NMR (300 MHz, Acetone-d6) δ ppm 0.88 (t, $J=7.33$ Hz, 3 H) 1.27 - 1.42 (m, 1 H) 1.45 - 1.61 (m, 1 H) 1.67 - 1.77 (s, 6 H) 1.88 - 2.03 (m, 2 H) 2.18 - 2.31 (m, 2 H) 2.72 - 2.89 (m, 4 H) 2.90 - 3.02 (m, 2 H) 3.93 - 4.05 (m, 2 H) 7.06 - 7.32 (m, 12 H) 10 8.01 (br. s., 1 H)

2-fluoroethyl (1*S*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamate, 944

1H NMR (300 MHz, Acetone-d6) δ ppm 1.73 (s, 6 H) 1.91 - 2.02 (m, 2 H) 2.19 - 2.32 (m, 2 H) 2.73 - 2.85 (m, 2 H) 2.88 - 3.04 (m, 2 H) 4.16 - 4.26 (m, 1 H) 4.25 - 4.36 (m, 1 H) 4.44 - 4.55 (m, 1 H) 4.60 - 4.68 (m, 1 H) 7.09 - 7.31 (m, 12 H) 8.20 (br. s., 1 H)

20 **3-chloropropyl (1*S*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamate, 945**

1H NMR (300 MHz, Acetone-d6) δ ppm 1.73 (s, 6 H) 1.89 - 2.03 (m, 2 H) 2.18 - 2.33 (m, 2 H) 2.18 - 2.31 (m, 2 H) 2.75 - 2.88 (m, 2 H) 2.90 - 3.03 (m, 2 H) 3.58 - 3.70 (m, 2 H) 4.08 - 4.19 (m, 2 H) 7.03 - 7.32 (m, 12 H) 8.11 (br. s., 1 H)

25 **Ethyl (1*R*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamate, 354**

1H NMR (300 MHz, Acetone-d6) δ ppm 1.15 (t, $J=7.03$ Hz, 3 H) 1.74 (s, 6 H) 1.97 - 2.11 (m, 2 H) 2.11 - 2.24 (m, 2 H) 2.73 - 2.91 (m, 2 H) 2.92 - 3.03 (m, 2 H) 3.94 - 4.08 (m, 2 H) 7.08 (br. s., 2 H) 7.12 - 7.35 (m, 10 H) 7.98 (br. s., 1 H)

30

Propyl (1*R*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamate, 355

1H NMR (300 MHz, Acetone-d6) δ ppm 0.87 (t, $J=7.33$ Hz, 3 H) 1.46 - 1.62 (m, 2 H) 1.74 (s, 6 H) 1.98 - 2.09 (m, 2 H) 2.10 - 2.23 (m, 2 H) 2.82 - 2.91 (m, 2 H) 2.91 -

18517 PCT (AP)

3.06 (m, 2 H) 3.87 - 3.98 (m, 2 H) 7.08 (s, 2 H) 7.14 - 7.35 (m, 10 H) 8.01 (br. s., 1 H)

Butyl (1*R*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-***i*j]quinolin-9-ylcarbamate, 356**

1H NMR (300 MHz, Acetone-d6) δ ppm 0.88 (t, *J*=7.33 Hz, 3 H) 1.24 - 1.41 (m, 2 H) 1.47 - 1.60 (m, 2 H) 1.74 (s, 6 H) 1.97 - 2.11 (m, 2 H) 2.08 - 2.24 (m, 2 H) 2.78 - 2.90 (m, 2 H) 2.89 - 3.04 (m, 2 H) 3.91 - 4.01 (m, 2 H) 7.09 (br. s., 2 H) 7.13 - 7.34 (m, 10 H) 8.00 (br. s., 1 H)

10

3-chloropropyl (1*R*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-**hexahdropyrido[3,2,1-*i*j]quinolin-9-ylcarbamate, 357**

1H NMR (300 MHz, Acetone-d6) δ ppm 1.74 (s, 6 H) 1.97 - 2.09 (m, 2 H) 2.11 - 2.25 (m, 2 H) 2.79 - 2.93 (m, 4 H) 2.91 - 3.04 (m, 2 H) 3.56 - 3.72 (m, 2 H) 4.05 - 4.18 (m, 2 H) 7.07 (br. s., 2 H) 7.10 - 7.34 (m, 10 H) 8.08 (s, 1 H)

But-3-enyl (1*R*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-**hexahdropyrido[3,2,1-*i*j]quinolin-9-ylcarbamate, 358**

1H NMR (300 MHz, Acetone-d6) δ ppm 1.74 (s, 6 H) 1.93 - 2.12 (m, 2 H) 2.11 - 2.23 (m, 2 H) 2.23 - 2.37 (m, 2 H) 2.81 - 2.90 (m, 2 H) 2.91 - 3.06 (m, 2 H) 4.02 (t, *J*=6.74 Hz, 2 H) 4.91 - 5.14 (m, 2 H) 5.68 - 5.87 (m, 1 H) 7.09 (br. s., 2 H) 7.12 - 7.35 (m, 10 H) 8.03 (br. s., 1 H)

methyl (1*S*,7*S*)-1,7-bis(4-fluorophenyl)-1,7-dimethyl-1,2,3,5,6,7-**hexahdropyrido[3,2,1-*i*j]quinolin-9-ylcarbamate, 674**

¹H NMR (300 MHz, Acetone-d6): δ = 1.71 (s, 6H) 1.92 - 2.07 (m, 2H) 2.19 - 2.27 (m, 2H) 2.76 - 2.84 (m, 2H) 2.95 - 3.02 (m, 2H) 3.57 (s, 3H) 6.98 - 7.22 (m, 10H)

butyl (1*S*,7*S*)-1,7-bis(4-fluorophenyl)-1,7-dimethyl-1,2,3,5,6,7-**hexahdropyrido[3,2,1-*i*j]quinolin-9-ylcarbamate, 672**

¹H NMR (300 MHz, Acetone-d6): δ = 0.88 (t, *J*=7.5 Hz, 3 H) 1.51 - 1.56 (m, 2H) 1.33 - 1.38 (m, 2H) 1.73 (s, 6H) 1.92 - 2.06 (m, 2H) 2.21 - 2.27 (m, 2H) 2.81 - 2.84 (m, 2H) 2.94 - 3.27 (m, 2H) 3.97 - 4.43 (m, 2 H) 6.98 - 7.22 (m, 10H)

18517 PCT (AP)**S-methyl (1*S*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamothioate, 583**

¹H NMR (300 MHz, Acetone-d6): δ = 1.68 (s, 6H) 2.02 (s, 3H) 2.20 - 1.88 (m, 4H) 2.95 - 3.01 (m, 2H) 2.80 - 2.95 (m, 2H) 6.76 (s, 2H) 7.33 - 7.17 (m, 10H)

5

S-propyl (1*S*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamothioate, 585

¹H NMR (300 MHz, Acetone-d6): δ = 0.91 (t, J =7.2 Hz, 3 H) , 0.97-1.1 (m, 2H) 1.49 - 1.51 (m, 4H) 1.74 (s, 6H) 1.92 - 1.96 (m, 2H) 2.18 - 2.23 (m, 2H) 2.73 - 2.84 (m, 4H) 2.95 - 3.05 (m, 2H) 7.05 (s, 2H) 7.15 - 7.36 (m, 8H)

10

S-methyl (1*R*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamothioate, 582

¹H NMR (300 MHz, Acetone-d6): δ = 1.73 (s, 6H) 2.20 (s, 3H) 1.98 - 2.07 (m, 2H) 2.14 - 2.17 (m, 2H) 2.80 - 2.88(m, 2H) 2.95 - 3.01 (m, 2H) 7.13 - 7.32 (m, 12H)

15

S-propyl (1*R*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamothioate, 584

¹H NMR (300 MHz, Acetone-d6): δ = 1.40 (t, J =9Hz, 3H) 2.05 - 2.12 (m, 2H) 2.28 (s, 6H) 2.66 - 2.72 (m, 6H) 2.51 - 2.64 (m, 2H) 3.47 - 3.54 (m, 2H) 3.31 - 3.39 (m, 2H) 7.59-7.80 (m, 12H)

20

S-methyl (1*S*,7*S*)-1,7-bis(4-fluorophenyl)-1,7-dimethyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-ylcarbamothioate, 671

¹H NMR (300 MHz, Acetone-d6): δ = 1.73 (s, 6H) 2.19 - 2.21 (m, 2H) 2.24-2.27 (m, 2H) 2.81 (s, 3H) 2.76 - 2.86 (m, 2H) 2.96 - 3.03 (m,2H) 6.93 - 7.22 (m, 10H)


25

S-propyl (1*S*,7*S*)-1,7-bis(4-fluorophenyl)-1,7-dimethyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9- ylcaramothioate, 673

¹H NMR (300 MHz, Acetone-d6): δ = 0.92 (t, J =9 Hz, 3 H) 1.08 - 2.11 (m, 2 H) 1.50 - 1.60 (m, 4 H) 1.70 (s, 6 H) 1.90 – 1.98 (m, 1 H) 2.17 – 2.25 (m, 3 H) 2.73-2.78 (m, 1 H) 2.93 – 3.00 (m,1 H) 6.96 – 7.19 (m, 10 H)


30

18517 PCT (AP)

Example BMethod B1: Preparation of 1-(1,7-diphenyl-1,2,3,5,6,7-hexahydropyrido[3,2,1-ij]quinolin-9-yl)-3-ethylurea (484)

5

To a solution 1,7-diphenyl-1,2,3,5,6,7-hexahydropyrido[3,2,1-ij]quinolin-9-amine (**Intermediate 4**), (102 mg, 0.30 mmol) in dichloromethane (15 mL) was added ethyl isocyanate (0.026 mL, 0.33 mmol), mmol) under argon at 0 °C. The reaction mixture was then stirred at room temperature for overnight. The solvent was removed under reduced pressure and purified by MPLC (medium pressure liquid chromatography) using silica gel column with 15 to 20 % EtOAc:Hexane to get 1-(1,7-diphenyl-1,2,3,5,6,7-hexahydropyrido[3,2,1-ij]quinolin-9-yl)-3-ethylurea (**484**) (120 mg 97%). ¹H NMR (300 MHz, CDCl₃) δ ppm 0.91 (t, *J*=7.18 Hz, 3 H) 2.05 - 2.20 (m, 2 H) 2.21 - 2.39 (m, 2 H) 2.96 - 3.28 (m, 6 H) 4.07 - 4.19 (m, 2 H) 4.35 (br. s., 1 H) 5.58 (s, 1 H) 6.43 (s, 2 H) 7.06 - 7.36 (m, 10 H)

Method B2: Preparation of 1-((1*S*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahydropyrido[3,2,1-ij]quinolin-9-yl)-3-ethylurea, **249**:

20

To a solution (1*S*,7*S*)-1,7-dimethyl-9-amino-1,7-diphenyl-1,2,3,5,6,7-hexahydropyrido[3,2,1-ij]quinoline (**9**), (110 mg, 0.30 mmol) in dichloromethane (15 mL) was added ethyl isocyanate (0.026 mL, 0.328 mmol), mmol) under argon at 0 °C. The reaction mixture was then stirred at room temperature for overnight. The solvent was removed under reduced pressure and purified by MPLC) using silica

18517 PCT (AP)

gel column with 15 to 20 % EtOAc:Hexane to get 1-((1*S*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-yl)-3-ethylurea (**249**), (96 mg 73%). 1 H NMR (300 MHz, Acetone-d6) δ ppm 1.00 (t, J =7.18 Hz, 3 H) 1.71 (s, 6 H) 1.91 - 2.02 (m, 2 H) 2.15 - 2.31 (m, 2 H) 2.75 - 2.87 (m, 2 H) 2.90 - 3.03 (m, 2 H) 3.03 - 3.16 (m, 2 H) 6.98 (s, 2 H) 7.09 - 7.31 (m, 10 H)

The following compounds were prepared according to the **Reaction Scheme B** and with the steps as shown in **Example B** above.

10 **1-Butyl-3-(1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-yl)urea, 485**

1 H NMR (300 MHz, CDCl₃) δ ppm 0.85 (t, J =7 Hz, 3 H) 1.04 - 1.38 (m, 4 H) 2.05 - 2.19 (m, 2 H) 2.19 - 2.38 (m, 2 H) 2.89 - 3.29 (m, 6 H) 4.14 (t, J =6.15 Hz, 2 H) 4.38 (br. s., 1 H) 5.61 (s, 1 H) 6.43 (s, 2 H) 7.05 - 7.38 (m, 10 H)

15

1-(1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-yl)-3-pentylurea, 486

1 H NMR (300 MHz, CDCl₃) δ ppm 0.84 (t, J =7.18 Hz, 3 H) 1.03 - 1.33 (m, 6 H) 2.04 - 2.19 (m, 2 H) 2.20 - 2.38 (m, 2 H) 2.90 - 3.28 (m, 6 H) 4.07 - 4.20 (m, 2 H) 4.38 (br. s., 1 H) 5.58 (s, 1 H) 6.43 (s, 5 H) 7.07 - 7.39 (m, 10 H)

1-(1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-yl)-3-hexylurea, 487

1 H NMR (300 MHz, CDCl₃) δ ppm 0.86 (t, J =7.1 Hz, 3 H) 1.06 - 1.33 (m, 8 H) 2.00 - 2.20 (m, 2 H) 2.21 - 2.40 (m, 2 H) 2.93 - 3.29 (m, 6 H) 4.02 - 4.17 (m, 2 H) 4.29 - 4.45 (m, 1 H) 5.57 (br. s., 1 H) 6.43 (br. s., 2 H) 7.03 - 7.40 (m, 10 H)

1-(2-chloroethyl)-3-(1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-yl)urea, 769

30 1 H NMR (300 MHz, CD₃OD) δ ppm 1.99 - 2.15 (m, 2 H) 2.19 - 2.35 (m, 2 H) 3.04 - 3.15 (m, 4 H), 3.49 (m, 2 H) 3.41 - 3.50 (m, 2 H) 4.11 - 4.22 (m, 2 H) 6.55 (s, 2 H) 7.09 - 7.21 (m, 5 H) 7.21 - 7.32 (m, 5 H)

1-allyl-3-(1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-yl)urea,

35 **773**

18517 PCT (AP)

¹H NMR (300 MHz, CD₃OD) δ ppm 2.00 - 2.14 (m, 2 H) 2.20 - 2.36 (m, 2 H) 3.04 - 3.16 (m, 4 H) 3.58 - 3.69 (m, 2 H) 4.11 - 4.23 (m, 2 H) 4.94 - 5.13 (m, 1 H) 5.73 (m, 1 H) 6.56 (s, 2 H) 7.09 - 7.21 (m, 5 H) 7.23 - 7.31 (m, 5 H)

5 **1-tert-butyl-3-(1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-yl)urea, 148**

¹H NMR (300 MHz, CDCl₃)δ ppm 7.12 - 7.33 (m, 10H) 6.41 (s, 2H) 4.14 (t, J=6 Hz, 2H) 3.14 - 3.19 (m, 4H) 2.25 - 2.30 (m, 2H) 2.10 - 2.17 (m, 2H) 2.25 - 2.33 (m, 2H)

10 **1-(1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-yl)-3-(furan-3-ylmethyl)urea, 258**

¹H NMR (300 MHz, CD₃OD) δ ppm 7.12 - 7.33 (m, 11H) 6.56 (s, 2H), 6.25 - 6.27 (m, 1H) 6.08 - 6.10 (m, 1H) 4.19 (s, 2H) 4.16 (t, J=6 Hz, 2H) 3.07 - 3.12 (m, 4H) 2.01 - 2.11 (m, 2H) 2.27 - 2.29 (m, 2H)

15

1-((1S,7S)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-yl)-3-propylurea, 250

¹H NMR (300 MHz, Acetone-d6) δ ppm 0.83 (t, J=6.8 Hz, 3 H) 1.33 - 1.47 (m, 2 H) 1.72 (s, 6H) 1.88 - 2.02 (m, 2 H) 2.17 - 2.30 (m, 2 H) 2.77 - 2.88 (m, 2 H) 2.90 - 2.10 (m, 4 H) 6.98 (s, 2 H) 7.10 - 7.30 (m, 10 H)

1-((1S,7S)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-yl)-3-butylurea, 251

¹H NMR (300 MHz, Acetone-d6) δ ppm 0.86 (t, J=6.8 Hz, 3 H) 1.20 - 1.45 (m, 4 H) 1.70 (s, 6H) 1.90 - 2.02 (m, 2 H) 2.15 - 2.30 (m, 2 H) 2.79 - 2.88 (m, 2 H) 2.90 - 3.02 (m, 2 H) 3.02 - 3.15 (m, 2 H) 6.98 (s, 2 H) 7.09 - 7.33 (m, 10 H)

1-allyl-3-((1S,7S)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-yl)urea, 252

¹H NMR (300 MHz, Acetone-d6) δ ppm 1.73 (s, 6 H) 1.87 - 2.02 (m, 2 H) 2.16 - 2.30 (m, 2 H) 2.74 - 2.87 (m, 2 H) 2.89 - 3.01 (m, 2 H) 3.67 - 3.78 (m, 2 H) 4.91 - 5.16 (m, 2 H) 5.73 - 5.90 (m, 1 H) 7.00 (s, 2 H) 7.06 - 7.35 (m, 10 H)

18517 PCT (AP)**1-t-butyl-3-((1*S*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-yl)urea, 581**

¹H NMR (300 MHz, Acetone-d6): δ = 0.71 (s, 9H) 1.20 (s, 6H) 1.97-2.02 (m, 2H) 2.07 - 2.16 (m, 2H) 2.74 - 2.80 (m, 2H) 2.93 - 2.97 (m, 2H) 6.79 (s, 2H) 7.36 - 7.17 (m, 10H)

1-((1*R*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-yl)-3-ethylurea, 463

¹H NMR (300 MHz, Acetone-d6) δ ppm 1.00 (t, J =7.18 Hz, 3 H) 1.73 (s, 6 H) 1.94 - 2.03 (m, 2 H) 2.10 - 2.25 (m, 2 H) 2.79 - 2.90 (m, 2 H) 2.92 - 3.04 (m, 2 H) 3.10 (t, J =6.45 Hz, 2 H) 5.29 (br. s., 1 H) 6.93 (s, 2 H) 7.07 - 7.34 (m, 10 H)

1-((1*R*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-yl)-3-propylurea, 464

¹H NMR (300 MHz, Acetone-d6) δ ppm 0.82 (t, J =7.00 Hz, 3 H) 1.31 - 1.48 (m, 2 H) 1.75 (s, 6 H) 1.95 - 2.03 (m, 2 H) 2.10 - 2.24 (m, 2 H) 2.75 - 2.89 (m, 4 H) 2.92 - 3.07 (m, 2 H) 5.33 - 5.35 (m, 1 H) 6.95 (s, 2 H) 7.07 - 7.35 (m, 10 H)

1-((1*R*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-yl)-3-butylurea, 465

¹H NMR (300 MHz, Acetone-d6) δ ppm 0.86 (t, J =7.18 Hz, 3 H) 1.19 - 1.44 (m, 4 H) 1.73 (s, 6 H) 1.94 - 2.05 (m, 2 H) 2.11 - 2.23 (m, 2 H) 2.77 - 2.92 (m, 4 H) 2.92 - 3.02 (m, 2 H) 3.02 - 3.12 (m, 2 H) 5.31 (br. s., 1 H) 7.09 - 7.34 (m, 10 H)

1-t-butyl-3-((1*R*,7*S*)-1,7-dimethyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-yl)urea, 580


¹H NMR (300 MHz, Acetone-d6): δ = 1.26 (s, 9H) 1.71 (s, 6H) 1.57 - 1.71 (m, 2H) 1.43 - 1.47 (m, 2H) 2.44 - 2.49 (m, 2H) 2.24 - 2.30 (m, 2H) 6.30 (s, 2H) 6.64 - 6.84 (m, 10H)

30

Example C

Method C: Preparation of 3-(1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinolin-9-yl)-1,1-dimethylurea (405)

18517 PCT (AP)



To a solution 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-amine (**Intermediate 4**), (106 mg, 0.31 mmol) in dichloromethane (10 mL) was added triethyl amine (0.130 mL, 0.933 mmol) followed by dimethylcarbamic chloride (0.043 mL, 0.46 mmol), under argon at 0 °C. The reaction mixture was then stirred at room temperature for overnight. The mixture was quenched with water (30 mL). The residue was isolated in a typical aqueous workup and purified by MPLC (medium pressure liquid chromatography) using silica gel column with 10 to 15 % EtOAc:Hexane to give 3-(1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-yl)-1,1-dimethylurea, (**405**), (63 mg 49%). ¹H NMR (300 MHz, CDCl₃) ppm 2.01 - 2.16 (m, 2 H) 2.20 - 2.37 (m, 2 H) 2.85 (s, 6 H) 3.00 - 3.15 (m, 4 H) 4.12 - 4.25 (m, 2 H) 5.78 (s, 1 H) 6.62 (s, 2 H) 7.10 - 7.22 (m, 5 H) 7.23 - 7.34 (m, 5 H).

15 The following compound was prepared according to the **Reaction Scheme C** and with the steps as shown in **Example C** above.


N-(1,7-diphenyl-1,2,3,5,6,7-hexahydropyrido[3,2,1-ij]quinolin-9-yl)morpholine-4-carboxamide 983

²⁰ ¹H NMR (300 MHz, CDCl₃) δ ppm 2.03 - 2.17 (m, 2 H) 2.20 - 2.38 (m, 2 H) 2.92 - 3.42 (m, 8 H) 3.53 - 3.65 (m, 4 H) 3.98 - 4.20 (m, 2 H) 6.52 - 6.72 (m, 2 H) 7.08 - 7.23 (m, 5 H) 7.24 - 7.37 (m, 5 H)

Example D

25 **Method D: Preparation of 1-(1,7-diphenyl-1,2,3,5,6,7-hexahydropyrido[3,2,1-
ii]quinolin-9-yl)-3-propylthiourea (255)**

18517 PCT (AP)



To a solution 1,7-diphenyl-1,2,3,5,6,7-hexahydropyrido[3,2,1-ij]quinolin-9-amine (**Intermediate 4**), (100 mg, 0.294 mmol) in dichloromethane (15 mL) was added propyl isothiocyanate (0.038 mL, 0.323 mmol) under argon at 0 °C. The reaction mixture was then stirred at room temperature for overnight. The solvent was removed under reduced pressure and purified by MPLC (medium pressure liquid chromatography) using silica gel column with 15 to 20 % EtOAc:Hexane to get 1-(1,7-diphenyl-1,2,3,5,6,7-hexahydropyrido[3,2,1-ij]quinolin-9-yl)-3-propylthiourea (**255**) (64 mg 49%). ¹H NMR (300 MHz, CDCl₃) δ ppm 7.08 - 7.34 (m, 10H) 6.39 (s, 2H) 4.11 (t, J=6 Hz, 2H) 3.32 - 3.46 (m, 2H) 3.16 - 3.23 (m, 4H) 2.25 - 2.30 (m, 2H) 2.13 - 2.17 (m, 2H) 1.32 - 1.39 (m, 2H, 0.73 (t, J=6 Hz, 3H)

The following compound was prepared according to the **Reaction Scheme D** and with the steps as shown in **Example D** above.


15 **1-(1,7-diphenyl-1,2,3,5,6,7-hexahydropyrido[3,2,1-ij]quinolin-9-yl)-3-hexylthiourea, 256**
¹H NMR (300 MHz, CDCl₃)δ ppm 7.07 - 7.34 (m, 10H) 6.39 (s, 2H) 4.11 (t, J=6 Hz, 2H), 3.46 - 3.49 (m, 2H) 3.15 - 3.25 (m, 4H) 2.25 - 2.32 (m, 2H) 2.11 - 2.18 (m, 2H) 1.17 - 1.31 (m, 8H) 0.87 (t, J=6Hz, 3H)

20

Example E

Method E: Preparation of *N*-butyl-1,7-diphenyl-1,2,3,5,6,7-hexahydropyrido[3,2,1-ij]quinoline-9-carboxamide (481)

18517 PCT (AP)



To a solution ethyl 4-aminobenzoate (**11**) (2.47 g, 15 mmol) in acetonitrile (10 mL) was added one equivalent of trifluoroacetic acid (1.71 g, 15 mmol). To this suspension was added with stirring a heterogeneous mixture of styrene (**12**), (8.6 mL, 75 mmol) and 37% formaldehyde solution (6.09 mL, 75 mmol) under argon gave yellow precipitate. The precipitate had failed to redissolve after 30min. of stirring at room temperature, so the mixture was heated at reflux under argon for further 30min, during which time the precipitate redissolved. The reaction mixture was cooled to room temperature. The residue was isolated in a typical aqueous workup and purified by MPLC (medium pressure liquid chromatography) using silica gel column with 10 to 15 % EtOAc:Hexane to give ethyl 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinoline-9-carboxylate (**13**), (4.1 g 68%).

A solution of ethyl 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinoline-9-carboxylate (**13**), (4 g, 10 mmol), in EtOH (40 mL) was subjected to saponification reaction using 2N NaOH (40 mL). The residue was isolated in a typical aqueous workup and purified by MPLC (medium pressure liquid chromatography) using silica gel column with 30 to 40 % EtOAc:Hexane to give 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinoline-9-carboxylic acid (**14**), (2 g 54%).

To a solution of 1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinoline-9-carboxylic acid (**14**), (189 mg, 0.512 mmol) in dichloromethane (10 mL) were added butyl amine (0.027 mL, 0.512 mmol), EDCI (196 mg, 1.02 mmol), followed DMAP (4-(Dimethylamino)pyridine) (70 mg, 1.02 mmol) under argon at 0 °C. The reaction mixture was then stirred at room temperature for overnight. The reaction was quenched with water (30 mL). The residue was isolated in a typical aqueous workup and purified by MPLC (medium pressure liquid chromatography) using

18517 PCT (AP)

silica gel column with 25 to 30 % EtOAc:Hexane to give *N*-butyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinoline-9-carboxamide (**481**), (113 mg 52%). ^1H NMR (300 MHz, CDCl_3) δ ppm 1H NMR (300 MHz, Solvent) δ ppm 0.87 (t, $J=6.0$ Hz, 3 H) 1.17 - 1.38 (m, 2 H) 1.36 - 1.52 (m, 2 H) 2.03 - 2.17 (m, 2 H) 2.18 - 2.36 (m, 2 H) 3.06 - 3.37 (m, 6 H) 4.21 (t, $J=5.27$ Hz, 2 H) 5.61 - 5.75 (m, 1 H) 7.03 - 7.16 (m, 4 H) 7.17 - 7.37 (m, 8 H).

The following compounds were prepared according to the **Reaction Scheme E** and with the steps as shown in **Example E** above.

10

***N*-methyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinoline-9-carboxamide, 775**

^1H NMR (300 MHz, CDCl_3) δ ppm 2.05 - 2.19 (m, 2 H) 2.18 - 2.35 (m, 2 H) 2.79 (d, $J=4.40$ Hz, 3 H) 3.08 - 3.27 (m, 4 H) 4.15 - 4.25 (m, 2 H) 5.78 (bs, 1 H) 7.05 - 7.16 (m, 4 H) 7.16 - 7.26 (m, 2 H) 7.25 - 7.35 (m, 6 H)

1,7-diphenyl-*N*-propyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinoline-9-carboxamide, 409

^1H NMR (300 MHz, CDCl_3) δ ppm 0.86 (t, $J=7.47$ Hz, 3 H) 1.41 - 1.54 (m, 2 H) 2.02 - 2.16 (m, 2 H) 2.15 - 2.34 (m, 2 H) 3.06 - 3.30 (m, 6 H) 4.20 (q, $J=5.66$ Hz, 2 H) 5.71 (br. s., 1 H) 7.03 - 7.16 (m, 4 H) 7.16 - 7.38 (m, 8 H)

***N*-pentyl-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinoline-9-carboxamide, 482**

^1H NMR (300 MHz, CDCl_3) δ ppm 0.85 (t, $J=6.89$ Hz, 3 H) 1.16 - 1.34 (m, 4 H) 1.37 - 1.52 (m, 2 H) 2.04 - 2.17 (m, 2 H) 2.17 - 2.35 (m, 2 H) 3.07 - 3.35 (m, 6 H) 4.20 (q, $J=5.96$ Hz, 2 H) 5.69 (br. s., 1 H) 7.06 - 7.16 (m, 4 H) 7.17 - 7.39 (m, 8 H)

***N*-(2-hydroxyethyl)-1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-*ij*]quinoline-9-carboxamide, 777**

^1H NMR (300 MHz, CD_3OD) δ ppm 2.03 - 2.16 (m, 2 H) 2.18 - 2.33 (m, 2 H) 3.06 - 3.26 (m, 4 H) 3.31 - 3.39 (m, 2 H) 3.50 - 3.60 (m, 2 H) 4.25 (t, $J=5.13$ Hz, 2 H) 7.06 - 7.36 (m, 21 H)

18517 PCT (AP)

N-ethyl-1,7-bis(3-fluorophenyl)-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinoline-9-carboxamide, 611


¹H NMR (300 MHz, Acetone-d6) δ ppm 1.03 (t, *J*=7.18 Hz, 3 H) 2.06 - 2.17 (m, 2 H) 2.20 - 2.35 (m, 2 H) 3.05 - 3.34 (m, 6 H) 4.29 (t, *J*=5.13 Hz, 2 H) 6.84 - 6.92 (m, 2 H) 6.92 - 7.05 (m, 4 H) 7.27 - 7.41 (m, 4 H)

N-butyl-1,7-bis(3-fluorophenyl)-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinoline-9-carboxamide, 593

¹H NMR (300 MHz, Acetone-d6) δ ppm 0.87 (t, *J*=6.9 Hz, 3 H) 1.30 - 1.37 (m, 2 H) 1.44 - 1.51 (m, 2 H) 2.09 - 2.17 (m, 2 H) 2.18 - 2.35 (m, 2 H) 3.20 - 3.27 (m, 6 H) 4.25-4.29 (m, 2 H) 6.97 - 7.02 (m, 4 H) 7.30 - 7.38 (m, 6 H).

Example F

Method F: Preparation of *N*-ethyl-2-(1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-yl)acetamide (371)



To a solution 2-(4-aminophenyl)acetic acid (15) (3.0 g, 19.84 mmol) in acetonitrile (10 mL) was added one equivalent of trifluoroacetic acid (1.53 ml, 19.84 mmol). To this suspension was added with stirring a heterogeneous mixture of styrene (12), (11.3 mL, 99.2 mmol) and 37% formaldehyde solution (9.0 mL, 99.2 mmol) under argon gave yellow precipitate. The precipitate had failed to redissolve after 30 min. of stirring at room temperature, so the mixture was heated at reflux under argon for further 30min, during which time the precipitate redissolved. The reaction mixture was cooled to room temperature. The residue was isolated in a typical aqueous workup and purified by MPLC (medium pressure liquid chromatography) using silica gel column with 10 to 30 % EtOAc:Hexane to give 2-(1,7-diphenyl-1,2,3,5,6,7-hexahdropyrido[3,2,1-ij]quinolin-9-yl)acetic acid (16), (1.9 g, 48%).


18517 PCT (AP)

To a solution of 2-(1,7-diphenyl-1,2,3,5,6,7-hexahydropyrido[3,2,1-ij]quinolin-9-yl)acetic acid (**16**), (85 mg, 0.229 mmol) in dichloromethane (10 mL) were added Ethyl amine (0.015 mL, 0.229 mmol), EDCI (103 mg, 0.538 mmol), followed by HOBT (72 mg, 0.538 mmol) under argon at 0 °C. The reaction mixture was then stirred at room temperature for overnight. The reaction was quenched with water (30 mL). The residue was isolated in a typical aqueous workup and purified by MPLC (medium pressure liquid chromatography) using silica gel column with 5 % MeOH:CH₂Cl₂ to give *N*-ethyl-2-(1,7-diphenyl-1,2,3,5,6,7-hexahydropyrido[3,2,1-*j*]quinolin-9-yl)acetamide (**371**), (27 mg, 29%).

10 ¹H NMR (300 MHz, Acetone-d6) δ ppm 0.87 (t, *J*=6.9 Hz, 3 H) 2.01-2.11 (m, 2H), 2.21 – 2.27 (m, 2 H) 3.0-3.6 (m, 8H), 3.20 - 3.27 (m, 6 H) 4.25-4.29 (m, 2 H) 6.97 - 7.02 (m, 4 H) 7.30 - 7.38 (m, 6 H).

Example G

15 Method G: Procedure for the methyl (4R,10R)-4,10-diphenyl-4,5,6,8,9,10-hexahydropyrido[3,2,1-de][1,5]naphthyridin-2-ylcarbamate (179)



20

To a solution of N-(5-aminopyridin-2-yl)acetamide (**Intermediate 1**) (2.4 g, 15.8 mmol) in acetonitrile (12 mL) was added one equivalent of trifluoroacetic acid (1.2 mL, 15.8 mmol). To this suspension was added with stirring a heterogeneous mixture of styrene (**Intermediate 2**), (7.2 mL, 63.2 mmol) and 37% formaldehyde solution (5.2 mL, 63.2 mmol) under argon, which gave a yellow precipitate. The

18517 PCT (AP)

precipitate failed to redissolve after 30 min. of stirring at room temperature, so the mixture was heated at reflux under argon for further 30 min, during which time the precipitate redissolved. The reaction mixture was cooled to room temperature. The precipitate was filtered and wash with acetonitrile gave yellow solid, *n*-(4R,10R)-

5 4,10-diphenyl-4,5,6,8,9,10-hexahdropyrido[3,2,1-de][1,5]naphthyridin-2-yl)acetamide, (**Intermediate 3**), (2.78 g).

A mixture of *n*-(4R,10R)-4,10-diphenyl-4,5,6,8,9,10-hexahdropyrido[3,2,1-de][1,5]naphthyridin-2-yl)acetamide, (**Intermediate 3**), (0.550 g, 1.43 mmol), in 10 EtOH (12 mL) was Conc. HCl (1.2 mL). The mixture was stirred at 90°C for two hrs. The mixture was concentrated, neutralized with aq. NaOH and extracted in CH₂Cl₂, dried (MgSO₄), filtered and concentrated gave (4R,10R)-4,10-diphenyl-4,5,6,8,9,10-hexahdropyrido[3,2,1-de][1,5]naphthyridin-2-amine (**Intermediate 4**) as a solid, (0.380 g).

15 To a solution (4R,10R)-4,10-diphenyl-4,5,6,8,9,10-hexahdropyrido[3,2,1-de][1,5]naphthyridin-2-amine (**Intermediate 4**), (110 mg, 0.322 mmol) in dichloromethane (10 mL) was added two equivalent of triethyl amine (0.090 mL, 0.644 mmol), followed by methyl chloroformate (0.037 mL, 0.483 mmol) under 20 argon at 0 °C. The reaction mixture was then stirred at room temperature for overnight. The mixture was quenched with water (30 mL). The residue was isolated in a typical aqueous workup and purified by MPLC (medium pressure liquid chromatography) using silica gel column with 10 to 15 % EtOAc:Hexane to give methyl (4R,10R)-4,10-diphenyl-4,5,6,8,9,10-hexahdropyrido[3,2,1-de][1,5]naphthyridin-2-ylcarbamate, (**179**), (20 mg).¹H NMR (300 MHz, CDCl₃) δ ppm 2.00 - 2.18 (m, 2 H) 2.22 - 2.39 (m, 2 H) 3.03 - 3.22 (m, 4 H) 3.50 - 3.64 (m, 3 H) 3.54 - 3.65 (m, 3 H) 4.05 - 4.23 (m, 2 H) 6.61 (br. s., 2 H) 7.08 - 7.38 (m, 10 H)

Biological Data:

30 The compounds of the invention are assessed for their ability to activate or block activation of the human S1P3 receptor in T24 cells stably expressing the human S1P3 receptor using the method described in paragraph [0067] of United States Patent Application Publication No. 20070232682, which published on Oct. 4, 2007, which is hereby incorporated by reference in its entirety.

18517 PCT (AP)

Ten thousands cells/well are plated into 384-well poly-D-lysine coated plates one day prior to use. The growth media for the S1P3 receptor expressing cell line is McCoy's 5A medium supplemented with 10% charcoal-treated fetal bovine serum (FBS), 1% antibiotic-antimycotic and 400 µg/ml geneticin. On the day of the 5 experiment, the cells are washed twice with Hank's Balanced Salt Solution supplemented with 20 mM HEPES (HBSS/Hepes buffer). The cells are then dye loaded with 2 µM Fluo-4 diluted in the HBSS/Hepes buffer with 1.25 mM Probenecid and incubated at 37 °C for 40 minutes. Extracellular dye is removed by washing the cell plates four times prior to placing the plates in the FLIPR 10 (Fluorometric Imaging Plate Reader, Molecular Devices). Ligands are diluted in HBSS/Hepes buffer and prepared in 384-well microplates. The positive control, Sphingosine-1-phosphate (S1P), is diluted in HBSS/Hepes buffer with 4 mg/ml fatty acid free bovine serum albumin. The FLIPR transfers 12.5 µl from the ligand 15 microplate to the cell plate and takes fluorescent measurements for 75 seconds, taking readings every second, and then for 2.5 minutes, taking readings every 10 seconds. Compounds are tested over the concentration range of 0.61 nM to 10,000 nM. Data for calcium ion (Ca^{+2}) responses are obtained in arbitrary fluorescence units and not translated into Ca^{+2} concentrations. IC_{50} values (nM) are determined through a linear regression analysis using the Levenburg Marquardt 20 algorithm.

Table I lists the test results for some of the compounds of the present invention:










18517 PCT (AP)

Table 1 Biological Data: Activity Potency of Compounds against Human S1P3

Receptor nM, (IC₅₀), %Inhibition:

Comp. no.	Structure	S1P3 IC ₅₀	S1P3 % Inhibition
272		8.3	102
273		63	101
274		64	100
275		12.2	100
276		8.7	101
277		901	91
278		66	101
484		77	101










18517 PCT (AP)

485		52	100
486		82	100
487		301	100
405		131	98
983		130	100
769		374	100
773		160	100
094		234	101
093		48	102

18517 PCT (AP)

(+)-enantiomer		3	97
(-)-enantiomer		230	97
202		272	98
779		23	101
268		5	98
(+)-enantiomer		1.6	98
(-)-enantiomer		NA	
095		56	102
067		8	98

18517 PCT (AP)

(+)-enantiomer		3	99
(-)-enantiomer		NA	
354		16	97
353		13	98
352		40	97
206		62	100
205		8	99
(+)-enantiomer		4	99
(-)-enantiomer		1659	

18517 PCT (AP)

699		4	100
(+)-enantiomer		1.6	98
(-)-enantiomer		NA	
700		7	100
(+)-enantiomer		13	98
(-)-enantiomer		NA	
148		38	101
258		59	100
256		78	100

18517 PCT (AP)

255		153	99
829		7	97
941		13	100
942		72	100
943		75	100
944		27	100
945		68	100
354		36	98
355		316	98

18517 PCT (AP)

674		46	96
672		86	89
583		10	98
585		5	98
582		4	99
584		16	98
671		25	97
673		62	97
249		194	100

18517 PCT (AP)








250		523	100
581		349	98
580		62	99
775		619	100
409		874	89
481		210	98

18517 PCT (AP)

TABLE 2: Additional Compounds of the Present Invention:

Compound	Structure
1	
2	
3	
4	
5	
6	
7	

18517 PCT (AP)










8	
9	
10	
11	
12	
13	
14	

18517 PCT (AP)










18517 PCT (AP)

22	
23	
24	
25	
26	
27	
28	
29	
30	

18517 PCT (AP)

31	
32	
33	
34	
35	
36	
37	
38	
39	









18517 PCT (AP)

40	
41	
42	
43	
44	
45	
46	
47	
48	

18517 PCT (AP)

49	
50	
51	
52	
53	
54 (+) Single enantiomer	
55 (-) Single enantiomer	
56	
57	

18517 PCT (AP)

58	
59	
60	
61	
62	
63	
64	
65	

18517 PCT (AP)

18517 PCT (AP)


75	
76	

5

18517 PCT (AP)

Claims:

5 1 A compound and salts and isomers thereof represented by the structural formula:




wherein each Y is independently C or N;
 10 wherein m is an integer of 0, or 1;
 n is an integer of 0, 1, 2, or 3;
 Z and X are each independently selected from the group consisting of oxygen sulfur, and amine moiety NR^N ; B is selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, cyano and X-B together being a heterocyclic ring or ring system;
 15 R and R^3 are each independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano;
 20 each R^1 is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxycarbonyl, aminocarbonyl, aminocarbonxyl, alkylcarboxyl, alkyl
 25


18517 PCT (AP)

alkylcarbonyl, formyl, oxycarbonyl, aminocarbonyl, aminocarbonxyl, alkylcarboxyl, alkyl amide, amino, alkylamino, and cyano; each R² is independently selected from the group consisting of hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxo, oxycarbonyl, carboxyl, alkyl carboxylate, alkyl amide, aminocarbonyl, amino, alkylamino, and cyano; each R^N is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano; including their alternate solid forms, tautomers, stereoisomers, enantiomers, diastereomers, prodrugs, and pharmaceutically acceptable salts, hydrates and solvates.

2. The compound of Claim 1, further represented by the compound selected from the group consisting of:



and



20

wherein m is an integer of 0, or 1;

n is an integer of 0, 1, 2, or 3;

18517 PCT (AP)

Z and X are each independently selected from the group consisting of oxygen, sulfur, and amine moiety NR^N; B is selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkylcarboxyl, alkyl amide, amino, alkylamino, cyano and X-B together being a heterocyclic ring or ring system;


R and R³ are each independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkylcarboxyl, alkyl amide, amino, alkylamino, and cyano;

each R¹ is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxycarbonyl, aminocarbonyl, aminocarbonxyl, alkylcarboxyl, alkylcarbonyl, formyl, oxycarbonyl, aminocarbonyl, aminocarbonxyl, alkylcarboxyl, alkyl amide, amino, alkylamino, and cyano;

each R² is independently selected from the group consisting of hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxo, oxycarbonyl, carboxyl, alkyl carboxylate, alkyl amide, aminocarbonyl, amino, alkylamino, and cyano; each R^N is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkylcarboxyl, alkyl amide, amino, alkylamino, and cyano; including their alternate solid forms, tautomers, stereoisomers, enantiomers, diastereomers, prodrugs, and pharmaceutically acceptable salts, hydrates and solvates.

30 3. The compound of Claim 1, further represented by the structural formula:

18517 PCT (AP)



wherein

m is 0, or 1;

n is 0, 1, 2, or 3;

5 Z is O, S, or NR^N;X is O, S, or NR^N;

B is selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, 10 aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, cyano and X-B together being a heterocyclic ring;

R and R³ are each independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, 15 formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano;

each R¹ is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxycarbonyl, 20 aminocarbonyl, aminocarbonxyl, alkylcarboxyl, alkyl amide, amino, alkylamino, and cyano;

each R² is independently selected from the group consisting of hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxo, oxycarbonyl, carboxyl, 25 alkyl carboxylate, alkyl amide, aminocarbonyl, amino, alkylamino, and cyano;


each R^N is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl,

18517 PCT (AP)

hydroxyl, alkoxyl, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano; including its alternate solid forms, tautomers, stereoisomers, enantiomers, diastereomers, prodrugs, and pharmaceutically acceptable salts, hydrates and solvates.

4. The compound of claim 1, provided that when Z and X are both oxygen, R² is not oxo, or R¹ and R² are not both phenyl or both methyl at the same time.

5. The compound of claim 1, further represented by the structural formula:



10 wherein

n is 0, 1, 2, or 3;

o is 0, 1, 2, or 3;

Z is O or S;

X is O, S, or NR^N;

15 B is selected from the group consisting of hydrogen, a substituent having a formula C₀₋₁₂H₀₋₃₀N₀₋₃O₀₋₅P₀₋₂S₀₋₃F₀₋₆Cl₀₋₃Br₀₋₃I₀₋₃, C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, C₁₋₆ haloalkyl, hydroxyalkyl, phenyl, benzyl, furylmethyl, or wherein X-B is morpholino and X-B being a heterocyclic ring;

20 R and R³ are each independently selected from the group consisting of hydrogen and a substituent having a formula C₀₋₁₂H₀₋₃₀N₀₋₃O₀₋₅P₀₋₂S₀₋₃F₀₋₆Cl₀₋₃Br₀₋₃I₀₋₃;

each R¹ is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxyl, hydroxyalkyl, alkylcarbonyl, formyl, oxycarbonyl, aminocarbonyl, aminocarbonxyl, alkylcarboxyl, alkyl amide, amino, alkylamino, and cyano;

18517 PCT (AP)

each R^N is independently selected from the group consisting of hydrogen and C₁₋₁₂ hydrocarbyl;

each R^A is independently selected from the group consisting of hydrogen, alkyl, hydrocarbyl, C₁₋₁₂ alkyl, C₁₋₁₂ alkenyl, C₁₋₁₂ alkynyl, halo, C₁₋₁₂ halo hydrocarbyl, C₁₋₁₂ hydroxyalkyl, C₃₋₁₂ cyclic hydrocarbyl, heteroaryl, heterohydrocarbyl, substituted or unsubstituted aryl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano;

including its alternate solid forms, tautomers, stereoisomers, enantiomers,

10 diastereomers, prodrugs, and pharmaceutically acceptable salts, hydrates and solvates.

6. The compound of claim 5 wherein X is O.

7. The compound of claim 5 wherein Z is O.

8. The compound of claim 5 wherein B is selected from the group consisting of C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, C₁₋₆ haloalkyl, hydroxyalkyl, phenyl, benzyl, furylmethyl, or wherein X-B is morpholino.


9. The compound of claim 5 wherein each R^A is independently hydrogen, C₁₋₁₂ alkyl, C₁₋₁₂ alkenyl, C₁₋₁₂ alkynyl, halo, C₁₋₁₂ halo hydrocarbyl, C₁₋₁₂ hydroxyalkyl, C₃₋₁₂ cyclic hydrocarbyl, or heteroaryl.

10. The compound of claim 5 wherein X is NR^N.

11. The compound of claim 5, wherein B is selected from the group consisting of C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, C₁₋₆ haloalkyl, hydroxyalkyl, phenyl, benzyl, furylmethyl, or wherein X-B is morpholino.

12. The compound of claim 3, further represented by the structural formula:

18517 PCT (AP)



wherein

n is 0, 1, 2, or 3;

o is 0, 1, 2, or 3;

5 Z is O or S;

X is O, S, or NR^N;

B is selected from the group consisting of hydrogen, a substituent having a formula

C₀₋₁₂H₀₋₃₀N₀₋₃O₀₋₅P₀₋₂S₀₋₃F₀₋₆Cl₀₋₃Br₀₋₃I₀₋₃, and X-B being a heterocyclic ring;

each R³ is independently selected from the group consisting of hydrogen and a

10 substituent having a formula

C₀₋₁₂H₀₋₃₀N₀₋₃O₀₋₅P₀₋₂S₀₋₃F₀₋₆Cl₀₋₃Br₀₋₃I₀₋₃;

each R^N is independently selected from the group consisting of hydrogen and C₁₋₁₂ hydrocarbyl;

each R^A is independently selected from the group consisting of hydrogen, C₁₋₁₂


15 alkyl, C₁₋₁₂ alkenyl, C₁₋₁₂ alkynyl, halo, C₁₋₁₂ halohydrocarbyl, C₁₋₁₂ hydroxyalkyl, C₃₋₁₂ cyclic hydrocarbyl, and substituted or unsubstituted aryl;

including its alternate solid forms, tautomers, stereoisomers, enantiomers,

diastereomers, prodrugs, and pharmaceutically acceptable salts, hydrates and solvates.

20 13. The compound of claim 1, further represented by the structural formula:

18517 PCT (AP)




wherein

- n is 0, 1, 2, or 3;
- Z is O, S, or NR^N;
- 5 X is O, S, or NR^N;
- B is selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, cyano and X-B together being a heterocyclic ring;
- 10 R and R³ are each independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano;
- 15 each R¹ is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxycarbonyl, aminocarbonyl, aminocarbonxyl, alkylcarboxyl, alkyl amide, amino, alkylamino, and cyano;
- 20 each R² is independently selected from the group consisting of hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxo, oxycarbonyl, carbonyl, alkyl carboxylate, alkyl amide, aminocarbonyl, amino, alkylamino, and cyano;
- 25 including its alternate solid forms, tautomers, stereoisomers, enantiomers, diastereomers, prodrugs, and pharmaceutically acceptable salts, hydrates and solvates.

18517 PCT (AP)

14. The compound of claim 1, further represented by the structural formula:



wherein

n is 0, 1, 2, or 3;

5 o is 0, 1, 2, or 3;

X is O, S, or NR^N;

B is selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, C₁₋₆ haloalkyl, C₁₋₆ hydroxyalkyl, phenyl, benzyl, furylmethyl, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halohydrocarbyl, hydroxyl,

10 alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, cyano and X-B together being morpholino or a heterocyclic ring;

R and R³ are each independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, 15 halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl, alkyl amide, amino, alkylamino, and cyano;

each R¹ is independently selected from the group consisting of hydrogen, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, 20 halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl, alkylcarbonyl, formyl, oxycarbonyl, aminocarbonyl, aminocarbonxyl, alkylcarboxyl, alkyl amide, amino, alkylamino, and cyano;

each R^A is independently selected from the group consisting of hydrogen, C₁₋₁₂ alkyl, C₁₋₁₂ alkenyl, C₁₋₁₂ alkynyl, halo, C₁₋₁₂ halohydrocarbyl, C₁₋₁₂ hydroxyalkyl,

25 C₃₋₁₂ cyclic hydrocarbyl, heteroaryl, hydrocarbyl, heterohydrocarbyl, substituted or unsubstituted aryl, halo, halohydrocarbyl, hydroxyl, alkoxy, hydroxyalkyl,

18517 PCT (AP)

alkylcarbonyl, carbonylalkyl, formyl, oxycarbonyl, aminocarbonyl, alkyl carboxyl,

alkyl amide, amino, alkylamino, and cyano;

including its alternate solid forms, tautomers, stereoisomers, enantiomers,

diastereomers, prodrugs, and pharmaceutically acceptable salts, hydrates and

5 solvates.

15. The compound of claim 14 wherein X is O.


16. The compound of claim 14, wherein B is C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, C₁₋₆ haloalkyl, C₁₋₆ hydroxyalkyl, phenyl, benzyl, furylmethyl, or wherein X-B is morpholino.

10 17. The compound of claim 14 wherein X is NR^N.


18. The compound of claim 14, wherein B is C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, C₁₋₆ haloalkyl, C₁₋₆ hydroxyalkyl, phenyl, and benzyl.

19. The compound of claim 14 wherein each R^A is independently hydrogen, C₁₋₁₂ alkyl, C₁₋₁₂ alkenyl, C₁₋₁₂ alkynyl, halo, C₁₋₁₂ halohydrocarbyl, C₁₋₁₂ hydroxyalkyl, 15 C₃₋₁₂ cyclic hydrocarbyl, or heteroaryl.


20. A compound of claim 1 selected from the group consisting of:






18517 PCT (AP)




18517 PCT (AP)




18517 PCT (AP)





18517 PCT (AP)





18517 PCT (AP)





18517 PCT (AP)




21. A compound of claim 1 selected from the group consisting of:




5




18517 PCT (AP)




18517 PCT (AP)



18517 PCT (AP)



18517 PCT (AP)



22. Use of a compound according to claim 1 in the manufacture of a medicament for the treatment of a disease or condition mediated by sphingosine-1-phosphate 3 (S1P3) receptor in a mammal.

23. The use of claim 22, wherein the disease or condition is selected from the group consisting of glaucoma, elevated intraocular pressure, ischemic neuropathies, optic neuropathy, pain, visceral pain, corneal pain, headache pain, migraine, cancer pain, back pain, irritable bowel syndrome pain, muscle pain and pain associated with diabetic neuropathy, diabetic retinopathy, retinal degenerative conditions, dry eye, angiogenesis, retinopathy of prematurity, diabetic retinopathy, optic neuropathy, glaucomatous retinopathy, macular degeneration, choroidal neovascularization, ocular wound healing, retinal edema, congestive heart failure, cardiac arrhythmia, atherosclerosis, bradycardia, asthma, chronic obstructive pulmonary disease, acute lung injury, acute respiratory distress syndrome, idiopathic pulmonary fibrosis, ventilation-induced lung injury, scar-less wound healing, scar-less skin-wound and cosmetic healing.

24. The use of claim 22 wherein the mammal is a human.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2009/043149

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D471/04 C07D471/06 C07D471/16 A61P27/00 A61P9/00
A61K31/4745

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61P A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, EMBASE, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 03/062392 A (CERETEK LLC [US]) 31 July 2003 (2003-07-31) page 136; claim 156; compound 317 page 13 – page 15 page 130 – page 133 -----	1-24
A	LEE S ET AL: "Synthesis and anti-angiogenesis activity of coumarin derivatives" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, PERGAMON, ELSEVIER SCIENCE, GB, vol. 16, no. 17, 1 September 2006 (2006-09-01), pages 4596-4599, XP025107342 ISSN: 0960-894X [retrieved on 2006-09-01] page 4596, left-hand column, paragraph 1; compound 15 ----- -/-	1-24

 Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

- *&* document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

21 July 2009

10/08/2009

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Härtinger, Stefan

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2009/043149

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2004/026864 A (WARNER LAMBERT CO [US]; ANDREANA TONJA LYNN [US]; CHO STEPHEN SUNG YON) 1 April 2004 (2004-04-01) claims 1,7; example 89 -----	1-24
A	WO 2007/056155 A (CHEMBRIDGE RES LAB INC [US]; ANIKIN ALEXEY VYACHESLAVOVICH [US]; GANTL) 18 May 2007 (2007-05-18) page 26, paragraph 391 – page 39 pages 106,279 page 304; claim 5 -----	1-24
X	WO 2004/037213 A (PROCTER & GAMBLE [US]) 6 May 2004 (2004-05-06) page 5, paragraph 6; claims 4,9; figure 1; example 3; table III pages 56,80 -----	1-24
X	HOLT JASON J ET AL: "A microwave-assisted synthesis of julolidine-9-carboxamide derivatives and their conversion to chalcogenoxanthones via directed metalation." THE JOURNAL OF ORGANIC CHEMISTRY 30 MAR 2007, vol. 72, no. 7, 30 March 2007 (2007-03-30), pages 2690-2693, XP002538089 ISSN: 0022-3263 compounds 18,19 -----	1,3,4, 14,19
P,X	DATABASE CHEMCATS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, US; RN 773874-23-0 1 January 2009 (2009-01-01), XP002538090 retrieved from STN order number: AL BF 0592 abstract -----	1
P,X	DATABASE CHEMCATS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, US; RN 311324-17-1 24 March 2009 (2009-03-24), XP002538091 retrieved from STN order number: STK330393 abstract -----	1
		-/-

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2009/043149

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	DATABASE CHEMCATS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, US; RN 773135-85-6 1 January 2009 (2009-01-01), XP002538092 retrieved from STN order number: AL BI 0592 abstract -----	1
P, X	DATABASE CHEMCATS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, US; RN 773129-68-3 1 January 2009 (2009-01-01), XP002538093 retrieved from STN order number: AL BK 0592 abstract -----	1
P, X	DATABASE CHEMCATS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, US; RN 773869-60-6 1 January 2009 (2009-01-01), XP002538094 retrieved from STN order number: AL BE 0592 abstract -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2009/043149

Patent document cited in search report	Publication date	Patent family member(s)			Publication date
WO 03062392	A 31-07-2003	AU 2003214873 A1	CA 2473740 A1	EP 1513522 A2	JP 2005519915 T
					02-09-2003 31-07-2003 16-03-2005 07-07-2005
WO 2004026864	A 01-04-2004	AU 2003263413 A1	BR 0314393 A	CA 2499326 A1	EC SP055676 A
					01-04-2004 06-07-2005 29-06-2005
			EP 1546143 A1	HR 20050248 A2	IS 7710 A
					24-02-2005 31-10-2005
			JP 2006503106 T	KR 20050057404 A	JP 2006503106 T
					26-01-2006 16-06-2005
			MA 27438 A1	MX PA05002007 A	PA 8582601 A1
					01-07-2005 28-04-2005 26-05-2004
			UY 27976 A1		30-04-2004
WO 2007056155	A 18-05-2007	AU 2006311914 A1	EP 1960382 A1	US 2008207635 A1	
					18-05-2007 27-08-2008 28-08-2008
WO 2004037213	A 06-05-2004	AU 2003285042 A1	CA 2500974 A1	CN 1705468 A	EP 1553916 A2
					06-05-2004 07-12-2005 20-07-2005
			JP 2006507287 T	US 2004131648 A1	
					02-03-2006 08-07-2004