

(19)



(11) Publication number:

SG 173642 A1

(43) Publication date:

29.09.2011

(51) Int. Cl:

;

(12)

## Patent Application

(21) Application number: 2011057791

(71) Applicant:

MEDIMMUNE, LLC ONE MEDIMMUNE

(22) Date of filing: 03.02.2010

WAY GAITHERSBURG, MD 20878 ML

(30) Priority: US 61/152,094 12.02.2009

US

THE GOVERNMENT OF THE UNITED

STATES OF AMERICA NATIONAL

INSTITUTES OF HEALTH 6011

EXECUTIVE BOULEVARD, SUITE 325

ROCKVILLE, MD 20852 ML US

(72) Inventor:

JIN, HONG 22385 SANTA PAULA

AVENUE CUPERTINO, CA 95014 US

CHENG, XING 849 NECTARINE AVENUE

SUNNYVALE, CA 94087 US

SUBBARAO, KANTA 1720 WILLARD

STREET, N.W. WASHINGTON, DC 20009

US

(54) Title:

INFLUENZA HEMAGGLUTININ AND NEURAMINIDASE  
VARIANTS

(57) Abstract:

Polypeptides, polynucleotides, reassortant viruses, immunogenic compositions and vaccines comprising influenza hemagglutinin and neuraminidase variants and method using thereof are provided.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
19 August 2010 (19.08.2010)

(10) International Publication Number  
**WO 2010/093537 A3**

(51) International Patent Classification:

*A61K 39/145 (2006.01)*

(21) International Application Number:

PCT/US2010/022970

(22) International Filing Date:

3 February 2010 (03.02.2010)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/152,094 12 February 2009 (12.02.2009) US

(71) Applicants (for all designated States except US): **MED-IMMUNE, LLC** [US/US]; One MedImmune Way, Gaithersburg, MD 20878 (US). **THE GOVERNMENT OF THE UNITED STATES OF AMERICA NATIONAL INSTITUTES OF HEALTH** [US/US]; 6011 Executive Boulevard, Suite 325, Rockville, MD 20852 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **JIN, Hong** [US/US]; 22385 Santa Paula Avenue, Cupertino, CA 95014 (US). **CHENG, Xing** [US/US]; 849 Nectarine Avenue, Sunnyvale, CA 94087 (US). **SUBBARAO, Kanta** [US/US]; 1720 Willard Street, N.W., Washington, DC 20009 (US).

(74) Agents: **HOLMES-SON, Michelle** et al.; MedImmune, LLC, One MedImmune Way, Gaithersburg, MD 20878 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(88) Date of publication of the international search report:

7 October 2010

WO 2010/093537 A3

(54) Title: INFLUENZA HEMAGGLUTININ AND NEURAMINIDASE VARIANTS

(57) Abstract: Polypeptides, polynucleotides, reassortant viruses, immunogenic compositions and vaccines comprising influenza hemagglutinin and neuraminidase variants and method using thereof are provided.

**INFLUENZA HEMAGGLUTININ AND NEURAMINIDASE VARIANTS****BACKGROUND OF THE INVENTION**

[0001] Vaccines against various and evolving strains of influenza are important from a community health stand point, as well as commercially, since each year numerous individuals are infected with different strains and types of influenza virus. Infants, the elderly, those without adequate health care and immuno-compromised persons are at special risk of death from such infections. Compounding the problem of influenza infections is that novel influenza strains evolve readily and can spread amongst various species, thereby necessitating the continuous production of new vaccines.

[0002] Numerous vaccines capable of producing a protective immune response specific for such different and influenza viruses/virus strains have been produced for over 50 years and include whole virus vaccines, split virus vaccines, surface antigen vaccines and live attenuated virus vaccines. However, while appropriate formulations of any of these vaccine types are capable of producing a systemic immune response, live attenuated virus vaccines have the advantage of also being able to stimulate local mucosal immunity in the respiratory tract. Considerable work in the production of influenza viruses, and fragments thereof, for production of vaccines has been done by the present inventors and co-workers; *see, e.g.*, U.S. Application Nos. 60/420,708, filed October 23, 2002; 60/574,117, filed May 24, 2004; 10/423,828, filed April 25, 2003; 60/578,962, filed June 12, 2004; and 10/870,690 filed June 16, 2004, the disclosure of which is incorporated by reference herein.

[0003] Because of the continual emergence (or re-emergence) of different influenza strains, new influenza vaccines are continually desired. Such vaccines typically are created using antigenic moieties of the newly emergent virus strains, thus, polypeptides and polynucleotides of novel, newly emergent, or newly re-emergent virus strains (especially sequences of antigenic genes) are highly desirable.

[0004] The present invention provides new and/or newly isolated influenza hemagglutinin and neuraminidase variants that are capable of use in production of numerous types of vaccines as well as in research, diagnostics, etc. Numerous other benefits will become apparent upon review of the following.

## SUMMARY OF THE INVENTION

[0005] In some aspects herein, the invention comprises an isolated or recombinant polypeptide that is selected from: a polypeptide comprising the amino acid sequence encoded by any one of SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues

5 residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5; a polypeptide comprising the amino acid sequence of any one of SEQ ID

NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2,

residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6; a polypeptide comprising the amino acid sequence encoded by an open reading frame of any one of SEQ

10 ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5; any

alternative (e.g., the mature form without the signal peptide, or the polypeptide as present on the surface of a virus (e.g., influenza)) form of a polypeptide comprising the amino acid sequence of any one of SEQ ID NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues

15 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6; any polypeptide that is encoded by a polynucleotide which hybridizes under

highly stringent conditions over substantially the entire length of a polynucleotide

consisting of the nucleotide sequence selected from SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5

20 and residues 1063-1728 of SEQ ID NO:5; any polypeptide that is encoded by a polynucleotide which hybridizes under highly stringent conditions to a polynucleotide

consisting of the nucleotide sequence selected from SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5; and, a fragment of any of the above wherein the

25 sequence comprises a hemagglutinin or neuraminidase polypeptide, or a fragment of a hemagglutinin or neuraminidase polypeptide. In one embodiment, such polypeptide

fragments generate an antibody that specifically binds a full length polypeptide of the invention. In various embodiments, the isolated or recombinant polypeptides of the invention are substantially identical to about 300 contiguous amino acid residues of any of

30 the above polypeptides. In yet other embodiments, the invention comprises isolated or recombinant polypeptides, that comprise an amino acid sequence that is substantially identical over at least about 350 amino acids; over at least about 400 amino acids; over at

least about 450 amino acids; over at least about 500 amino acids; over at least about 520

amino acids; over at least about 550 amino acids; over at least about 559 amino acids; over at least about 565 amino acids; or over at least about 566 amino acids contiguous of any of the above polypeptides. In some embodiments, the polypeptide sequence (e.g., as listed in "SEQUENCES" herein) comprises less than 565, 559, etc. amino acids. In such

5 embodiments, the shorter listed polypeptides optionally comprise less than 565, 559, etc. amino acids. In yet other embodiments, the polypeptides of the invention (e.g., SEQ ID NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6) optionally comprise fusion proteins, proteins with a leader sequence, a precursor polypeptide, proteins 10 with a secretion signal or a localization signal, or proteins with an epitope tag, an E-tag, or a His epitope tag. In still other embodiments, the invention encompasses a polypeptide comprising an amino acid sequence having at least 85%, at least 90%, at least 93%, at least 95%, at least 98%, at least 98.5%, at least 99%, at least 99.2%, at least 99.4%, at least 99.6%, at least 99.8%, or at least 99.9% sequence identity to a polypeptide comprising the 15 amino acid sequence selected from SEQ ID NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6. In another embodiment, a polypeptide of the invention comprises an amino acid sequence that differs from any one of SEQ ID NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of 20 SEQ ID NO:6, or residues 341-562 of SEQ ID NO:6 at 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, or 50 amino acid residues. The hemagglutinin sequences of the invention can comprise both those sequences with unmodified and modified polybasic cleavage sites (thereby allowing growth of the viruses in eggs). The hemagglutinin polypeptide sequences of SEQ ID NOS:2 and 6 comprise the endogenous amino terminal signal peptide sequences, 25 however, the hemagglutinin polypeptide sequences of the invention also include the mature (amino terminal signal peptide cleaved) form of the hemagglutinin polypeptides. The cleavage sites of any hemagglutinin polypeptide sequence of any influenza strain can be routinely measured or predicted using any number of methods in the art.

**[0006]** In other aspects, the invention comprises a composition with one or more

30 polypeptide listed above, or fragments thereof. The invention also includes polypeptides that are specifically bound by a polyclonal antisera raised against at least 1 antigen that comprises at least one amino acid sequence described above, or a fragment thereof. Such

antibodies specific for the polypeptides described above are also features of the invention. In one embodiment, the polypeptides of the invention are immunogenic.

**[0007]** The invention also encompasses immunogenic compositions comprising an immunologically effective amount of one or more of any of the polypeptides described

5 above (e.g., SEQ ID NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6) as well as methods for stimulating the immune system of an individual to produce a protective immune response against influenza virus by administering to the individual an immunologically effective amount of any of the above polypeptides in a physiologically 10 acceptable carrier.

**[0008]** Additionally, the invention includes a reassortant influenza virus that comprises one or more of the polypeptides or polynucleotides above, in addition to immunogenic compositions comprising an immunologically effective amount of such reassortant influenza virus. Methods for stimulating the immune system of an individual to produce a 15 protective immune response against influenza virus, through administering an immunologically effective amount of such reassortant influenza virus in a physiologically acceptable carrier are also part of the invention.

**[0009]** In other aspects, the invention comprises an isolated or recombinant polynucleotide that is selected from: a polynucleotide comprising any one of the nucleotide sequences of

20 SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5 or complementary sequences thereof, a polynucleotide encoding a polypeptide comprising the amino acid sequence selected from SEQ ID NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 25 341-562 of SEQ ID NO:6 or complementary nucleotide sequences thereof, a polynucleotide which hybridizes under highly stringent conditions over substantially the entire length of any of the above described polynucleotides, and a polynucleotide comprising all or a fragment of any of such nucleotide sequences wherein the sequence encodes a hemagglutinin or neuraminidase polypeptide or a fragment of a hemagglutinin or 30 neuraminidase polypeptide. The invention also includes an isolated or recombinant polynucleotide that encodes an amino acid sequence which is substantially identical over at least about 300 amino acids of any polypeptide encoded by the above polynucleotides, or over at least about 350 amino acids; over at least about 400 amino acids; over at least about

450 amino acids; over at least about 500 amino acids; over at least about 502 amino acids; over at least about 550 amino acids; over at least about 559 amino acids; over at least about 565 amino acids; or over at least about 566 amino acids of any polypeptide encoded by the above polynucleotides. Again, in situations wherein the amino acid is less than, e.g., 566,

5 565, 559, etc. in length (e.g., *see*, “SEQUENCES”) then it should be understood that the length is optionally less than 566, 565, 559, etc. The invention also includes any of the above polynucleotides that comprise a nucleotide sequence encoding a hemagglutinin or neuraminidase polypeptide, or one or more fragments of one or more hemagglutinin or neuraminidase polypeptide. Other aspects of the invention include isolated or recombinant 10 polynucleotides that encode a polypeptide (e.g., a hemagglutinin or neuraminidase polypeptide) whose sequence has at least 98% identity, at least 98.5% identity, at least 99% identity, at least 99.2% identity, at least 99.4% identity, at least 99.6% identity, at least 99.8% identity, or at least 99.9% identity to at least one of the above described 15 polypeptides. The invention also includes isolated or recombinant polynucleotides encoding a polypeptide of hemagglutinin or neuraminidase produced by mutating or recombining one or more above described polynucleotides. In one embodiment, a polynucleotide of the invention may comprise a nucleotide sequence encoding one or more of, e.g., a leader sequence, a precursor sequence, or an epitope tag sequence or the like, and can optionally encode a fusion protein.

20 [0010] In yet other embodiments, the invention comprises a composition of matter having two or more above described polynucleotides (e.g., a library comprising at least about 2, 5, 10, 50 or more polynucleotides). Such compositions can optionally be produced by cleaving one or more above described polynucleotide (e.g., mechanically, chemically, enzymatically with a restriction endonuclease/RNase/DNAse, etc.). Other compositions of 25 the invention include, e.g., compositions produced by incubating one or more above described polynucleotide in the presence of deoxyribonucleotide triphosphates and a thermostable polynucleotide polymerase.

[0011] The invention also encompasses cells comprising at least one of the above described polynucleotides, or a cleaved or amplified fragment or product thereof. Such cells can 30 optionally express a polypeptide encoded by such polynucleotide. Other embodiments of the invention include vectors (e.g., plasmids, cosmids, phage, viruses, virus fragments, etc.) comprising any of above described polynucleotides. Such vectors can optionally comprise an expression vector. Preferred expression vectors of the invention include, but are not

limited to, vectors comprising pol I promoter and terminator sequences or vectors using both the pol I and pol II promoters “the polI/polII promoter system” (e.g., Zobel et al., Nucl. Acids Res. 1993, 21:3607; US20020164770; Neumann *et al.*, Proc. Natl. Acad. Sci. USA 1999, 96:9345; Fodor et al., J. Virol. 1999, 73:9679; and US20030035814). Cells

5 transduced by such vectors are also within the current invention.

[0012] In some embodiments, the invention encompasses a virus (e.g., an influenza virus) comprising one or more above described polynucleotides (e.g., encoding hemagglutinin and/or neuraminidase), or one or more fragments thereof. Immunogenic compositions comprising such virus are also part of the current invention. Such viruses can comprise a 10 reassortant virus such as a 6:2 reassortant virus (e.g., comprising 6 internal genome segments from one or more donor virus and 2 genome segments (e.g., HA or NA genome segments) comprising one or more above described polynucleotide (or one or more fragment thereof). In one embodiment, the genome segment may encode a hemagglutinin and/or neuraminidase polypeptide of the invention. In one embodiment, a reassortant

15 viruses of the invention is a live virus. In another embodiment, a reassortant virus of the invention is a temperature sensitive (ts), cold-adapted (ca), or attenuated (att) virus. In one embodiment, a reassortant virus of the invention comprises at least 1, at least 2, at least 3, at

least 4, at least 5 or 6 internal genome segment of a donor virus (e.g., A/Ann Arbor/6/60, PR8, etc). In another embodiment, a reassortant virus of the invention comprises at least 1, 20 at least 2, at least 3, at least 4, at least 5 or 6 internal genome segment of a donor virus other than A/Ann Arbor/6/60. One preferred embodiment of the invention is a reassortant

influenza virus, wherein the virus is a 6:2 reassortant influenza virus and comprises 6 internal genome segments from A/Ann Arbor/6/60 and 2 genome segments that encode a polypeptide selected from the group consisting of: the polypeptides of SEQ ID NOS:2, 4, 6, 25 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6. In an alternative embodiment, a

reassortant influenza virus of the invention includes a 6:2 reassortant influenza virus, wherein said virus comprises 6 internal genome segments from one or more donor viruses other than A/Ann Arbor/6/60 and 2 genome segments that encode a polypeptide selected 30 from the group consisting of: the polypeptides of SEQ ID NOS:2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6. In another alternative embodiment, a reassortant

influenza virus of the invention includes a 6:2 reassortant influenza virus, wherein said virus

comprises 6 internal genome segments from one or more donor viruses other than A/Ann Arbor/6/60 and 2 genome segments, wherein the 2 genome segments encode HA and/or NA polypeptides from any pandemic influenza strain. Methods of producing a reassortant influenza virus through culturing a host cell harboring an influenza virus in a suitable

5 culture medium under conditions permitting replication of the reassortant influenza virus and, isolating the reassortant influenza virus from one or more of the host cell or the medium are also part of the invention.

[0013] In other embodiments herein, the invention comprises immunogenic compositions having an immunologically effective amount of any of the above described reassortant

10 influenza virus. Other embodiments include methods for stimulating the immune system of an individual to produce a protective immune response against influenza virus by administering to the individual an immunologically effective amount of any of the reassortant influenza virus described above (optionally in a physiologically effective carrier).

15 [0014] Other aspects of the invention include methods of producing an isolated or recombinant polypeptide by culturing any host cell above, in a suitable culture medium under conditions permitting expression of the polypeptide and, isolating the polypeptide from one or more of the host cells or the medium in which the cells are grown.

20 [0015] Immunogenic compositions are also features of the invention. For example, immunogenic compositions comprising one or more of any of the polypeptides and/or polynucleotides described above and, optionally, an excipient such as a pharmaceutically acceptable excipient or one or more pharmaceutically acceptable administration component. Immunogenic compositions of the invention can also comprise any one or more above described virus as well (e.g., along with one or more pharmaceutically acceptable

25 administration component).

30 [0016] Methods of producing immunogenic responses in a subject through administration of an effective amount of any of the above viruses (or immunogenic compositions) to a subject are also within the current invention. Additionally, methods of prophylactic or therapeutic treatment of a viral infection (e.g., viral influenza) in a subject through administration of any one or more above described virus (or immunogenic compositions) in an amount effective to produce an immunogenic response against the viral infection are also part of the current invention. Subjects for such treatment include, but are not limited to, birds (e.g., poultry) and mammals (e.g., humans). Such methods can also comprise *in vivo*

administration to the subject as well as *in vitro* or *ex vivo* administration to one or more cells of the subject. Additionally, such methods can also comprise administration of a composition of the virus and a pharmaceutically acceptable excipient that are administered to the subject in an amount effect to prophylactically or therapeutically treat the viral

5 infection.

[0017] In other aspects the invention includes compositions of matter comprising nucleotide sequences encoding hemagglutinin and/or neuraminidase polypeptides of one or more pandemic influenza strain and nucleotide sequences encoding one or more polypeptide of A/Ann Arbor/6/60. Additionally, the invention includes compositions of matter comprising 10 nucleotide sequences encoding hemagglutinin and/or neuraminidase polypeptides of one or more pandemic influenza strain and nucleotide sequences encoding one or more polypeptide of PR8, A/Leningrad/17 or A/Ann Arbor/6/60. Such sequences can include those listed in the “SEQUENCES” herein. Additionally, preferred embodiments of the invention include compositions of matter comprising sequences encoding hemagglutinin and/or neuraminidase of one or more pandemic influenza strain and nucleotide sequences encoding a selected backbone strain in a 6:2 reassortant. Such compositions preferably include 15 sequences encoding the hemagglutinin and neuraminidase selected from the “SEQUENCES” herein and a backbone strain, wherein the backbone strain is PR8, A/LENINGRAD/17 or A/Ann Arbor/6/60. The invention also includes such compositions 20 as described above wherein the hemagglutinin comprises a modified polybasic cleavage site. The invention also includes live attenuated influenza vaccine comprising such above compositions.

[0018] These and other objects and features of the invention will become more fully apparent when the following detailed description is read in conjunction with the 25 accompanying figures and appendix.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0019] Figure 1: Replication of H2 wt challenge viruses in lungs of ferrets vaccinated with various indicated H2 vaccine viruses. Titer represents average of right and left lungs.

30 [0020] Figure 2: Replication of H2 wt challenge viruses in NT of ferrets vaccinated with various indicated H2 vaccine viruses.

[0021] Figure 3: Replication of H2 wt challenge viruses in lungs of mice vaccinated with various indicated H2 vaccine viruses.

[0022] Figure 4: Replication of H2 wt challenge viruses in NT of mice vaccinated with various indicated H2 vaccine viruses.

#### DETAILED DESCRIPTION

5 [0023] The present invention includes influenza hemagglutinin and neuraminidase polypeptides and polynucleotides as well as vectors, compositions, reassortant influenza viruses and the like comprising such polypeptides and polynucleotides and methods of their use. Additional features of the invention are described in more detail herein.

10 Definitions

[0024] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. The following definitions supplement those in the art and are directed to the current application and are not necessarily to be imputed to any related or unrelated case, e.g., to any commonly owned patent or application. Although any methods and materials similar or equivalent to those described herein can be used in the practice for testing of the present invention, the preferred materials and methods are described herein. Accordingly, the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting.

15 [0025] As used in this specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a virus” includes a plurality of viruses; reference to a “host cell” includes mixtures of host cells, and the like.

20 [0026] The term “reassortant,” when referring to a virus, indicates that the virus includes genetic and/or polypeptide components derived from more than one parental viral strain or source. For example, a 7:1 reassortant comprises 7 viral genome segments (or gene segments) from a first virus, and a single complementary viral genomic segment, e.g., encoding a hemagglutinin or neuraminidase of the invention. A 6:2 reassortant comprises 6 genome segments, e.g., the 6 internal genome segments from a first virus, and two complementary genome segments, i.e., hemagglutinin and neuraminidase encoding genome segments, from a second virus or a second and third virus.

25 [0027] The term “host cell” means a cell that contains a heterologous polynucleotide, such as a vector, and supports the replication and/or expression of the polynucleotide. Host cells

can be prokaryotic cells such as *E. coli*, or eukaryotic cells such as yeast, insect, amphibian, avian or mammalian cells, including human cells. In one embodiment, host cells may be, but are not limited to, Vero (African green monkey kidney) cells, BHK (baby hamster kidney) cells, primary chick kidney (PCK) cells, Madin-Darby Canine Kidney (MDCK) cells, Madin-Darby Bovine Kidney (MDBK) cells, 293 cells (e.g., 293T cells), and COS cells (e.g., COS1, COS7 cells), .

5 [0028] An “immunologically effective amount” of influenza virus is an amount sufficient to enhance an individual's (e.g., a human's) own immune response against a subsequent exposure to influenza virus. Levels of induced immunity can be monitored, e.g., by measuring amounts of neutralizing secretory and/or serum antibodies, e.g., by plaque neutralization, complement fixation, enzyme-linked immunosorbent, or microneutralization assay.

10 [0029] A “protective immune response” against influenza virus refers to an immune response exhibited by an individual (e.g., a human) that is protective against disease when 15 the individual is subsequently exposed to and/or infected with such influenza virus. In some instances, the influenza virus (e.g., naturally circulating) can still cause infection, but it cannot cause a serious infection. Typically, the protective immune response results in detectable levels of host engendered serum and secretory antibodies that are capable of neutralizing virus of the same strain and/or subgroup (and possibly also of a different, non- 20 vaccine strain and/or subgroup) in *vitro* and *in vivo*.

[0030] As used herein, an “antibody” is a protein comprising one or more polypeptides substantially or partially encoded by immunoglobulin genes or fragments of immunoglobulin genes. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon and mu constant region genes, as well as myriad 25 immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively. A typical immunoglobulin (antibody) structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one “light” (about 30 25 kD) and one “heavy” chain (about 50-70 kD). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain (VL) and variable heavy chain (VH) refer to these light and heavy chains respectively. Antibodies exist as intact immunoglobulins or as

a number of well-characterized fragments produced by digestion with various peptidases. Thus, for example, pepsin digests an antibody below the disulfide linkages in the hinge region to produce F(ab)'2, a dimer of Fab which itself is a light chain joined to VH-CH1 by a disulfide bond. The F(ab)'2 may be reduced under mild conditions to break the disulfide linkage in the hinge region thereby converting the (Fab')2 dimer into a Fab' monomer. The 5 Fab' monomer is essentially a Fab with part of the hinge region (see, Fundamental Immunology, W.E. Paul, ed., Raven Press, N.Y. (1999), for a more detailed description of other antibody fragments). While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such Fab' fragments may be 10 synthesized de novo either chemically or by utilizing recombinant DNA methodology. Thus, the term antibody, as used herein, includes antibodies or fragments either produced by the modification of whole antibodies or synthesized de novo using recombinant DNA methodologies. Antibodies include, e.g., polyclonal antibodies, monoclonal antibodies, 15 multiple or single chain antibodies, including single chain Fv (sFv or scFv) antibodies in which a variable heavy and a variable light chain are joined together (directly or through a peptide linker) to form a continuous polypeptide, and humanized or chimeric antibodies.

#### Influenza Virus

**[0031]** The polypeptides and polynucleotides of the invention, e.g., SEQ ID NO: 1-8, are 20 variants of influenza HA and NA sequences. In general, influenza viruses are made up of an internal ribonucleoprotein core containing a segmented single-stranded RNA genome and an outer lipoprotein envelope lined by a matrix protein. The genome of influenza viruses is composed of eight genome segments of linear (-) strand ribonucleic acid (RNA), 25 encoding the immunogenic hemagglutinin (HA) and neuraminidase (NA) proteins, and six internal core polypeptides: the nucleocapsid nucleoprotein (NP); matrix proteins (M); non-structural proteins (NS); and 3 RNA polymerase (PA, PB1, PB2) proteins. During replication, the genomic viral RNA is transcribed into (+) strand messenger RNA and (-) strand genomic cRNA in the nucleus of the host cell. Each of the eight genomic segments is packaged into ribonucleoprotein complexes that contain, in addition to the RNA, NP and 30 a polymerase complex (PB1, PB2, and PA).

**[0032]** Influenza is commonly grouped into influenza A and influenza B categories.

Influenza A and influenza B viruses each contain eight segments of single stranded RNA with negative polarity. The influenza A genome encodes eleven polypeptides. Segments 1-

3 encode three polypeptides, making up a RNA-dependent RNA polymerase. Segment 1 encodes the polymerase complex protein PB2. The remaining polymerase proteins PB1 and PA are encoded by segment 2 and segment 3, respectively. In addition, segment 1 of some influenza strains encodes a small protein, PB1-F2, produced from an alternative reading

5 frame within the PB1 coding region. Segment 4 encodes the hemagglutinin (HA) surface glycoprotein involved in cell attachment and entry during infection. Segment 5 encodes the nucleocapsid nucleoprotein (NP) polypeptide, the major structural component associated with viral RNA. Segment 6 encodes a neuraminidase (NA) envelope glycoprotein.

Segment 7 encodes two matrix proteins, designated M1 and M2, which are translated from

10 differentially spliced mRNAs. Segment 8 encodes NS1 and NS2, two nonstructural

proteins, which are translated from alternatively spliced mRNA variants. The eight genome segments of influenza B encode 11 proteins. The three largest genes code for components of the RNA polymerase, PB1, PB2 and PA. Segment 4 encodes the HA protein. Segment 5 encodes NP. Segment 6 encodes the NA protein and the NB protein. Both proteins, NB and

15 NA, are translated from overlapping reading frames of a bicistronic mRNA. Segment 7 of influenza B also encodes two proteins: M1 and BM2. The smallest segment encodes two products: NS1 is translated from the full length RNA, while NS2 is translated from a spliced mRNA variant.

20 Influenza virus vaccine

[0033] The sequences, compositions and methods herein are primarily, but not solely, concerned with production of influenza viruses for vaccines. Historically, influenza virus vaccines have primarily been produced in embryonated hen eggs using strains of virus selected or based on empirical predictions of relevant strains. More recently, reassortant

25 viruses have been produced that incorporate selected hemagglutinin and/or neuraminidase antigens in the context of an approved attenuated, temperature sensitive master strain.

Following culture of the virus through multiple passages in hen eggs, influenza viruses are recovered and, optionally, inactivated, e.g., using formaldehyde and/or  $\beta$ -propiolactone (or alternatively used in live attenuated vaccines). Thus, it will be appreciated that HA and NA

30 sequences (e.g., SEQ ID NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6) are quite useful in constructing influenza vaccines. The current invention includes viruses/vaccines comprising HA and/or NA polypeptides and polynucleotides of A/Japan/57

and A/swine/MO/2006 (including wherein the HA polypeptides and polynucleotides comprise modified polybasic cleavage sites such as the modifications described herein); and including wherein the viruses/vaccines comprise a backbone (i.e. 6 internal genome segments) such as the backbone of *ca* A/AA/6/60, A/Leningrad/17 or PR8.

5 [0034] Attempts at producing recombinant and reassortant vaccines in cell culture have been hampered by the inability of some of the strains approved for vaccine production to grow efficiently under standard cell culture conditions. However, prior work by the inventors and their coworkers provided a vector system, and methods for producing recombinant and reassortant viruses in culture, thus, making it possible to rapidly produce  
10 vaccines corresponding to one or many selected antigenic strains of virus, e.g., either A or B strains, various subtypes or substrains, etc., e.g., comprising the HA and/or NA sequences herein. See, Multi-Plasmid System for the production of Influenza virus, U.S. Application No. 60/420,708, filed October 23, 2002, U.S. Application No. 10/423,828, filed April 25, 2003 and U.S. Application 60/574,117 filed May 24, 2004. Typically, the cultures are  
15 maintained in a system, such as a cell culture incubator, under controlled humidity and CO<sub>2</sub>, at constant temperature using a temperature regulator, such as a thermostat to insure that the temperature does not exceed 35 °C. Reassortant influenza viruses can be readily obtained by introducing a subset of vectors corresponding to genomic segments of a master influenza virus, in combination with complementary segments derived from strains of interest (e.g.,  
20 HA and/or NA antigenic variants herein). Typically, the master strains are selected on the basis of desirable properties relevant to vaccine administration. For example, for vaccine production, e.g., for production of a live attenuated vaccine, the master donor virus strain may be selected for an attenuated phenotype, cold adaptation and/or temperature sensitivity. In one embodiment, a master donor virus comprises 6 internal genome segments (i.e. a  
25 backbone) that confer one or more of the following properties: temperature sensitive, cold adapted, or attenuated. As explained elsewhere herein and, e.g., in U.S. Patent Application No. 10/423,828, etc., various embodiments of the invention utilize A/Ann Arbor (AA)/6/60 influenza strain as a “backbone” upon which to add HA and/or NA genes (e.g., such as those sequences listed herein, etc.) to create desired reassortant viruses. Thus, for example,  
30 in a 6:2 reassortant, 2 genes (i.e., NA and HA) would be from the influenza strain(s) against which an immunogenic reaction is desired, while the other 6 genes would be from the Ann Arbor strain, or other backbone strain, etc. The Ann Arbor virus is useful for its cold adapted, attenuated, temperature sensitive attributes. Of course, it will be appreciated that

the HA and NA sequences herein are capable of reassortant with a number of other virus genes or virus types (e.g., a number of different “backbones” such as PR8, etc., containing the other influenza genes present in a reassortant, namely, the non-HA and non-NA genes

[0035] Various embodiments herein can comprise live attenuated vaccines, having the HA

5 and/or NA sequences herein, for A/Japan/57 or A/swine/MO/2006. Such vaccines typically comprise, e.g., the HA and/or NA polypeptides of SEQ ID NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6, or their corresponding encoding nucleotides of SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5. One problem arising from growth of vaccine virus strains (e.g., reassortants) in eggs is that certain strains (which can be involved in pandemics) can kill the eggs in which the vaccines are to be produced and are, thus, hard to manipulate, produce, etc. through use of traditional (non-plasmid rescue) reassortant production. Such strains are of interest since evidence indicates

10 they can result in influenza in humans and possible pandemics. Thus, use of plasmid-rescue systems to create/manipulate influenza reassortants with virus strains s (e.g., the HA and NA sequences herein) are quite desirable and are features of the invention. It will be appreciated, however, that the current sequences are also capable of use with non-plasmid or traditional systems.

20 [0036] In various embodiments herein, the antigenic sequences (e.g., the HA and/or NA polypeptides) as well as viruses and vaccines from such viruses comprise modified polybasic cleavage sites. Some highly pathogenic influenza strains comprise multiple basic amino acid cleavage sites within hemagglutinin sequences. *See, e.g., Li et al., J. of Infectious Diseases*, 179:1132-8, 1999. Such cleavage sites, in typical embodiments herein,

25 are, e.g., modified or altered in their sequences in comparison to the wild-type sequences from which the current sequences are derived (e.g., to disable the cleavage or reduce the cleavage there, etc.). Such modifications/alterations can be different in different strains due to the various sequences of the cleavage sites in the wild-type sequences. For example, 4 polybasic residues (arginine-arginine-lysine-lysine) at 326-329 of mature H5 are typically removed in sequences herein (as compared to wt). In various embodiments, the polybasic cleavage sites can be modified in a number of ways (all of which are contained within the invention). For example, the polybasic cleavage site can be removed one amino acid at a time (e.g., one arginine removed, two arginines removed, two arginines and lysine removed,

or two arginines and two lysines removed). Additionally, the amino acid residue directly upstream of the cleavage site can also be removed or altered (e.g., from an R to a T, etc.); also, the nucleotides encoding the amino acid residue directly after the cleavage site can also be modified. In addition, hemagglutinin polypeptide sequences of influenza virus comprise amino terminal signal peptide sequences, thus, the hemagglutinin polypeptide sequences of the invention include both the mature (amino terminal signal peptide cleaved) form of the hemagglutinin polypeptides and the pre-cleaved form of hemagglutinin. The cleavage sites of any hemagglutinin polypeptide sequence of any influenza strain can be routinely measured or predicted using any number of methods in the art.

10 [0037] The terms “temperature sensitive,” “cold adapted” and “attenuated” as applied to viruses (typically used as vaccines or for vaccine production) which optionally encompass the current sequences, are well known in the art. For example, the term “temperature sensitive” (*ts*) indicates, e.g., that the virus exhibits a 100 fold or greater reduction in titer at 39°C relative to 33°C for influenza A strains, or that the virus exhibits a 100 fold or greater reduction in titer at 37°C relative to 33°C for influenza B strains. The term “cold adapted” (*ca*) indicates that the virus exhibits growth at 25°C within 100 fold of its growth at 33°C, while the term “attenuated” (*att*) indicates that the virus replicates in the upper airways of ferrets but is not detectable in their lung tissues, and does not cause influenza-like illness in the animal. It will be understood that viruses with intermediate phenotypes, i.e., viruses exhibiting titer reductions less than 100 fold at 39°C (for A strain viruses) or 37°C (for B strain viruses), or exhibiting growth at 25°C that is more than 100 fold than its growth at 33°C (e.g., within 200 fold, 500 fold, 1000 fold, 10,000 fold less), and/or exhibit reduced growth in the lungs relative to growth in the upper airways of ferrets (i.e., partially attenuated) and/or reduced influenza like illness in the animal, are also useful viruses and can be used in conjunction with the HA and NA sequences herein.

25 [0038] Again, the HA and NA sequences of the current invention are optionally utilized in the production of or in reassortant vaccines (and/or in other *ts*, *cs*, *ca*, and/or *att* viruses and vaccines). However, it should be noted that the HA and NA sequences, etc. of the invention are not limited to specific vaccine compositions or production methods, and can, thus, be utilized in substantially any vaccine type or vaccine production method which utilizes strain specific HA and NA antigens (e.g., any of SEQ ID NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6, or the corresponding nucleotides encoding the specific

HA and NA antigens, e.g., SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5).

5 FluMist™

[0039] As mentioned previously, numerous examples and types of influenza vaccine exist. An example of an influenza vaccine is FluMist™ which is a live, attenuated vaccine that protects children and adults from influenza illness (Belshe et al. (1998) *The efficacy of live attenuated, cold-adapted, trivalent, intranasal influenza virus vaccine in children* N Engl J Med 338:1405-12; Nichol et al. (1999) *Effectiveness of live, attenuated intranasal influenza virus vaccine in healthy, working adults: a randomized controlled trial* JAMA 282:137-44).

10 The methods and compositions of the current invention may be adapted to/used with production of FluMist™ vaccine. However, it will be appreciated by those skilled in the art that the sequences, methods, compositions, etc. herein are also adaptable to production of 15 similar or even different viral vaccines.

[0040] FluMist™ vaccine strains contain, e.g., HA and NA gene segments derived from the strains (e.g., wild-type strains) to which the vaccine is addressed along with six gene segments, PB1, PB2, PA, NP, M and NS, from a common master donor virus (MDV). The HA sequences herein, thus, may be part of various FluMist™ formulations. The MDV for 20 influenza A strains of FluMist™ (MDV-A), was created by serial passage of the wild-type A/Ann Arbor/6/60 (A/AA/6/60) strain in primary chicken kidney tissue culture at successively lower temperatures (Maassab (1967) *Adaptation and growth characteristics of influenza virus at 25 degrees C* Nature 213:612-4). MDV-A replicates efficiently at 25°C (ca, cold adapted), but its growth is restricted at 38 and 39°C (ts, temperature sensitive).

25 Additionally, this virus does not replicate in the lungs of infected ferrets (att, attenuation). The ts phenotype is believed to contribute to the attenuation of the vaccine in humans by restricting its replication in all but the coolest regions of the respiratory tract. The stability of this property has been demonstrated in animal models and clinical studies. In contrast to the ts phenotype of influenza strains created by chemical mutagenesis, the ts property of 30 MDV-A does not revert following passage through infected hamsters or in shed isolates from children (for a recent review, see Murphy & Coelingh (2002) *Principles underlying the development and use of live attenuated cold-adapted influenza A and B virus vaccines* Viral Immunol 15:295-323).

[0041] Clinical studies in over 20,000 adults and children involving 12 separate 6:2 reassortant strains have shown that these vaccines are attenuated, safe and efficacious (Belshe et al. (1998) *The efficacy of live attenuated, cold-adapted, trivalent, intranasal influenza virus vaccine in children* N Engl J Med 338:1405-12; Boyce et al. (2000) *Safety and immunogenicity of adjuvanted and unadjuvanted subunit influenza vaccines administered intranasally to healthy adults* Vaccine 19:217-26; Edwards et al. (1994) *A randomized controlled trial of cold adapted and inactivated vaccines for the prevention of influenza A disease* J Infect Dis 169:68-76; Nichol et al. (1999) *Effectiveness of live, attenuated intranasal influenza virus vaccine in healthy, working adults: a randomized controlled trial* JAMA 282:137-44). Reassortants carrying the six internal genes of MDV-A and the two HA and NA gene segments of a wild-type virus (i.e., a 6:2 reassortant) consistently maintain *ca*, *ts* and *att* phenotypes (Maassab et al. (1982) *Evaluation of a cold-recombinant influenza virus vaccine in ferrets* J. Infect. Dis. 146:780-900).

[0042] Production of such reassorted virus using B strains of influenza is more difficult, however, recent work (see, e.g., Multi-Plasmid System for the Production of Influenza Virus, U.S. Application No. 60/420,708, filed October 23, 2002, U.S. Application No. 10/423,828, filed April 25, 2003, and U.S. Application No. 60/574,117, filed May 24, 2004) has shown an eight plasmid system for the generation of influenza B virus entirely from cloned cDNA. Methods for the production of attenuated live influenza A and B virus suitable for vaccine formulations, such as live virus vaccine formulations useful for intranasal administration were also shown.

[0043] The system and methods described previously are useful for the rapid production in cell culture of recombinant and reassortant influenza A and B viruses, including viruses suitable for use as vaccines, including live attenuated vaccines, such as vaccines suitable for intranasal administration. The sequences (e.g., nucleotide sequences SEQ ID NO: 1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 or residues 1063-1728 of SEQ ID NO:5 and the corresponding amino acids encoded by the nucleotide sequences in SEQ ID NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, or residues 341-562 of SEQ ID NO:6), methods, etc. of the current invention, are optionally used in conjunction with, or in combination with, such previous work involving, e.g., reassorted influenza viruses for vaccine production to produce viruses for vaccines.

[0044] Methods and Compositions for prophylactic administration of vaccines

[0045] As stated above, alternatively, or in addition to, use in production of FluMist<sup>TM</sup> vaccine, the current invention can be used in other vaccine formulations. In general, recombinant and reassortant viruses of the invention (e.g., those comprising polynucleotides of SEQ ID NO: 1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ

5 ID NO:1, residues 88-1062 of SEQ ID NO:5 or residues 1063-1728 of SEQ ID NO:5 or polypeptides of SEQ ID NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, or residues 341-562 of SEQ ID NO:6, or fragments thereof) can be administered prophylactically in an immunologically effective amount and in an appropriate carrier or excipient to stimulate an immune response specific

10 for one or more strains of influenza virus as determined by the HA and/or NA sequence.

Typically, the carrier or excipient is a pharmaceutically acceptable carrier or excipient, such as sterile water, aqueous saline solution, aqueous buffered saline solutions, aqueous dextrose solutions, aqueous glycerol solutions, ethanol, or combinations thereof. The preparation of such solutions insuring sterility, pH, isotonicity, and stability is effected

15 according to protocols established in the art. Generally, a carrier or excipient is selected to minimize allergic and other undesirable effects, and to suit the particular route of administration, e.g., subcutaneous, intramuscular, intranasal, etc.

[0046] A related aspect of the invention provides methods for stimulating the immune system of an individual to produce a protective immune response against influenza virus. In

20 the methods, an immunologically effective amount of a recombinant influenza virus (e.g., comprising an HA and/or NA molecule of the invention), an immunologically effective amount of a polypeptide of the invention, and/or an immunologically effective amount of a nucleic acid of the invention is administered to the individual in a physiologically acceptable carrier.

25 [0047] Generally, the influenza viruses of the invention are administered in a quantity sufficient to stimulate an immune response specific for one or more strains of influenza virus (i.e., against the HA and/or NA strains of the invention). Preferably, administration of the influenza viruses elicits a protective immune response to such strains. Dosages and methods for eliciting a protective immune response against one or more influenza strains are

30 known to those of skill in the art. *See*, e.g., USPN 5,922,326; Wright et al., *Infect. Immun.* 37:397-400 (1982); Kim et al., *Pediatrics* 52:56-63 (1973); and Wright et al., *J. Pediatr.* 88:931-936 (1976). For example, influenza viruses are provided in the range of about 1-1000 HID<sub>50</sub> (human infectious dose), i.e., about 10<sup>5</sup> - 10<sup>8</sup> pfu (plaque forming units) per

dose administered. Typically, the dose will be adjusted within this range based on, e.g., age, physical condition, body weight, sex, diet, time of administration, and other clinical factors. The prophylactic vaccine formulation is systemically administered, e.g., by subcutaneous or intramuscular injection using a needle and syringe, or a needle-less

5 injection device. Alternatively, the vaccine formulation is administered intranasally, either by drops, large particle aerosol (greater than about 10 microns), or spray into the upper respiratory tract. While any of the above routes of delivery results in a protective systemic immune response, intranasal administration confers the added benefit of eliciting mucosal immunity at the site of entry of the influenza virus. For intranasal administration,  
10 attenuated live virus vaccines are often preferred, e.g., an attenuated, cold adapted and/or temperature sensitive recombinant or reassortant influenza virus. *See above.* While stimulation of a protective immune response with a single dose is preferred, additional dosages can be administered, by the same or different route, to achieve the desired prophylactic effect.

15 [0048] Typically, the attenuated recombinant influenza of this invention as used in a vaccine is sufficiently attenuated such that symptoms of infection, or at least symptoms of serious infection, will not occur in most individuals immunized (or otherwise infected) with the attenuated influenza virus. In some instances, the attenuated influenza virus can still be capable of producing symptoms of mild illness (e.g., mild upper respiratory illness) and/or of dissemination to unvaccinated individuals. However, its virulence is sufficiently  
20 abrogated such that severe lower respiratory tract infections do not occur in the vaccinated or incidental host.

25 [0049] Alternatively, an immune response can be stimulated by *ex vivo* or *in vivo* targeting of dendritic cells with influenza viruses comprising the sequences herein. For example, proliferating dendritic cells are exposed to viruses in a sufficient amount and for a sufficient period of time to permit capture of the influenza antigens by the dendritic cells. The cells are then transferred into a subject to be vaccinated by standard intravenous transplantation methods.

30 [0050] While stimulation of a protective immune response may be elicited with a single dose, additional dosages can be administered, by the same or different route, to achieve the desired prophylactic effect. In neonates and infants, for example, multiple administrations may be required to elicit sufficient levels of immunity. Administration can continue at intervals throughout childhood, as necessary to maintain sufficient levels of protection

against wild-type influenza infection. Similarly, adults who are particularly susceptible to repeated or serious influenza infection, such as, for example, health care workers, day care workers, family members of young children, the elderly, and individuals with compromised cardiopulmonary function may require multiple immunizations to establish and/or maintain protective immune responses. Levels of induced immunity can be monitored, for example, by measuring amounts of neutralizing secretory and serum antibodies, and dosages adjusted or vaccinations repeated as necessary to elicit and maintain desired levels of protection.

5 [0051] Optionally, the formulation for prophylactic administration of the influenza viruses also contains one or more adjuvants for enhancing the immune response to the influenza 10 antigens. Suitable adjuvants include: complete Freund's adjuvant, incomplete Freund's adjuvant, saponin, mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil or hydrocarbon emulsions, bacille Calmette-Guerin (BCG), *Corynebacterium parvum*, and the synthetic adjuvant QS-21.

15 [0052] If desired, prophylactic vaccine administration of influenza viruses can be performed in conjunction with administration of one or more immunostimulatory molecules.

Immunostimulatory molecules include various cytokines, lymphokines and chemokines with immunostimulatory, immunopotentiating, and pro-inflammatory activities, such as interleukins (e.g., IL-1, IL-2, IL-3, IL-4, IL-12, IL-13); growth factors (e.g., granulocyte-macrophage (GM)-colony stimulating factor (CSF)); and other immunostimulatory

20 molecules, such as macrophage inflammatory factor, Flt3 ligand, B7.1; B7.2, etc. The immunostimulatory molecules can be administered in the same formulation as the influenza viruses, or can be administered separately. Either the protein (e.g., an HA and/or NA polypeptide of the invention, e.g., any of SEQ ID NO: 2, 4, ,6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and

25 residues 341-562 of SEQ ID NO:6) or an expression vector comprising a polynucleotide (e.g., any of SEQ ID NO: 1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 or residues 1063-1728 of SEQ ID NO:5) encoding the protein can be administered to produce an immunostimulatory effect.

30 [0053] The above described methods are useful for therapeutically and/or prophylactically treating a disease or disorder, typically influenza, by introducing a vector of the invention comprising a heterologous polynucleotide encoding a therapeutically or prophylactically effective HA and/or NA polypeptide (or peptide) or HA and/or NA RNA (e.g., an antisense RNA or ribozyme) into a population of target cells *in vitro*, *ex vivo* or *in vivo*. Typically,

the polynucleotide encoding the polypeptide (or peptide), or RNA, of interest is operably linked to appropriate regulatory sequences as described above in the sections entitled “Expression Vectors” and “Additional Expression Elements.” Optionally, more than one heterologous coding sequence is incorporated into a single vector or virus. For example, in 5 addition to a polynucleotide encoding a therapeutically or prophylactically active HA and/or NA polypeptide or RNA, the vector can also include additional therapeutic or prophylactic polypeptides, e.g., antigens, co-stimulatory molecules, cytokines, antibodies, etc., and/or markers, and the like.

**[0054]** Although vaccination of an individual with an attenuated influenza virus of a

10 particular strain of a particular subgroup can induce cross-protection against influenza virus of different strains and/or subgroups, cross-protection can be enhanced, if desired, by vaccinating the individual with attenuated influenza virus from at least two strains, e.g., each of which represents a different subgroup. Additionally, vaccine combinations can optionally include mixes of pandemic vaccines (e.g., those against pandemic influenza

15 strains such as various avian strains, *see, e.g.*, the sequences herein, or other pandemic strains) and non-pandemic strains. Vaccine mixtures (or multiple vaccinations) can

comprise components from human strains and/or non-human influenza strains (e.g., avian and human, etc.). Similarly, the attenuated influenza virus vaccines of this invention can optionally be combined with vaccines that induce protective immune responses against

20 other infectious agents.

## POLYNUCLEOTIDES OF THE INVENTION

**[0055]** It is well known in the art that the HA and NA polynucleotide segments of influenza viruses comprise both a coding region (encoding the ORF) and noncoding regions (NCRs),

25 both 5' and 3' of the HA and NA coding sequence. It is also known that primers can be made to these NCRs to facilitate amplification of the entire HA and NA segments of influenza virus. (*see, e.g.*, Hoffmann et al. Arch Virol. 2001 Dec;146(12):2275-89).

Further, it is known that the NCRs of the HA and NA of influenza may increase the efficiency of achieving reassortants. Therefore, the nucleotide sequences of these NCRs

30 (including fragments and variants (e.g., at least about 60%, or at least 70%, or at least 80%, or at least 90%, or at least about 91% or at least about 92%, or at least about 93%, or at least about 94%, or at least about 95%, or at least about 96%, or at least about 97%, or at least about 98%, or at least about 98.5%, or at least about 98.7%, or at least about 99%, or at

least about 99.1%, or at least about 99.2%, or at least about 99.3%, or at least about 99.4%, or at least about 99.5%, or at least about 99.6% or at least about 99.7%, or at least about 99.8%, or at least about 99.9% identity) thereof) are within the scope of this invention.

When amplifying the HA and NA segments of any pandemic strain, one could make and  
5 use polynucleotide primers to bind conserved (e.g., among related strains) regions of the HA and NA NCRs for amplification (e.g., by RT-PCR). In one embodiment, HA and NA

polynucleotides of the invention include both the NCR and ORF of the HA and NA sequences (including fragments and variants (e.g., at least about 60%, or at least 70%, or at least 80%, or at least 90%, or at least about 91% or at least about 92%, or at least about 93%, or at least about 94%, or at least about 95%, or at least about 96%, or at least about 97%, or at least about 98%, or at least about 98.5%, or at least about 98.7%, or at least about 99%, or at least about 99.1%, or at least about 99.2%, or at least about 99.3%, or at least about 99.4%, or at least about 99.5%, or at least about 99.6% or at least about 99.7%, or at least about 99.8%, or at least about 99.9%) thereof) of pandemic virus strains. In

10 15 alternative embodiments, the HA and NA polynucleotides of the invention exclude the NCR, but include the ORF (including fragments and variants (e.g., at least about 60%, or at least 70%, or at least 80%, or at least 90%, or at least about 91% or at least about 92%, or at least about 93%, or at least about 94%, or at least about 95%, or at least about 96%, or at least about 97%, or at least about 98%, or at least about 98.5%, or at least about 98.7%, or at least about 99%, or at least about 99.1%, or at least about 99.2%, or at least about 99.3%, or at least about 99.4%, or at least about 99.5%, or at least about 99.6% or at least about 99.7%, or at least about 99.8%, or at least about 99.9%) thereof) of the HA and NA

20 25 sequences of pandemic virus strains (e.g., SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5).

**[0056]** The HA and NA polynucleotides of the invention, e.g., SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5, and fragments thereof, are optionally used in a number of different capacities alternative to, or in addition to, the vaccines described above. Other exemplary uses are described herein for illustrative purpose and not as limitations on the actual range of uses. Different methods of construction, purification, and characterization of the nucleotide sequences of the invention are also described herein. In some embodiments, polynucleotides including one or more

nucleotide sequence of the invention are favorably used as probes for the detection of corresponding or related polynucleotides in a variety of contexts, such as in nucleic hybridization experiments, e.g., to find and/or characterize homologous influenza variants (e.g., homologues to the sequences herein, etc.) infecting other species or in different 5 influenza outbreaks, etc. The probes can be either DNA or RNA molecules, such as restriction fragments of genomic or cloned DNA, cDNAs, PCR amplification products, transcripts, and oligonucleotides, and can vary in length from oligonucleotides as short as about 10 nucleotides in length to full length sequences or cDNAs in excess of 1 kb or more. For example, in some embodiments, a probe of the invention includes a nucleotide sequence 10 or subsequence selected, e.g., from among SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5, or sequences complementary thereto. Alternatively, nucleotide sequences that are variants of one of the above-designated sequences are used as probes. Most typically, such variants include one or a few conservative nucleotide 15 variations. For example, pairs (or sets) of oligonucleotides can be selected, in which the two (or more) nucleotide sequences are conservative variations of each other, wherein one nucleotide sequence corresponds identically to a first variant or and the other(s) corresponds identically to additional variants. Such pairs of oligonucleotide probes are particularly useful, e.g., for specific hybridization experiments to detect polymorphic nucleotides or to, 20 e.g., detect homologous influenza HA and NA variants, e.g., homologous to the current HA and NA sequences, infecting other species or present in different (e.g., either temporally and/or geographically different) influenza outbreaks. In other applications, probes are selected that are more divergent, that is probes that are at least about 91% (or about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 98.5%, about 25 98.7%, about 99%, about 99.1%, about 99.2%, about 99.3%, about 99.4%, about 99.5%, or about 99.6% or more about 99.7%, about 99.8%, about 99.9% or more) identical are selected.

**[0057]** The probes of the invention, e.g., as exemplified by sequences derived from the sequences herein, can also be used to identify additional useful polynucleotide sequences 30 according to procedures routine in the art. In one set of embodiments, one or more probes, as described above, are utilized to screen libraries of expression products or chromosomal segments (e.g., expression libraries or genomic libraries) to identify clones that include sequences identical to, or with significant sequence similarity to, e.g., one or more probe of

the sequences herein, i.e., variants, homologues, etc. It will be understood that in addition to such physical methods as library screening, computer assisted bioinformatic approaches, e.g., BLAST and other sequence homology search algorithms, and the like, can also be used for identifying related polynucleotide sequences. Polynucleotide sequences identified in this manner are also a feature of the invention.

**[0058]** Oligonucleotide probes are optionally produced via a variety of methods well known to those skilled in the art. Most typically, they are produced by well known synthetic methods, such as the solid phase phosphoramidite triester method described by Beaucage and Caruthers (1981) *Tetrahedron Letts* 22(20):1859-1862, e.g., using an automated synthesizer, or as described in Needham-Van Devanter et al. (1984) *Nucl Acids Res*, 12:6159-6168. Oligonucleotides can also be custom made and ordered from a variety of commercial sources known to persons of skill. Purification of oligonucleotides, where necessary, is typically performed by either native acrylamide gel electrophoresis or by anion-exchange HPLC as described in Pearson and Regnier (1983) *J Chrom* 255:137-149.

15 The sequence of the synthetic oligonucleotides can be verified using the chemical degradation method of Maxam and Gilbert (1980) in Grossman and Moldave (eds.) Academic Press, New York, *Methods in Enzymology* 65:499-560. Custom oligos can also easily be ordered from a variety of commercial sources known to persons of skill.

**[0059]** In other circumstances, e.g., relating to attributes of cells or organisms expressing the polynucleotides and polypeptides of the invention (e.g., those harboring virus comprising the sequences of the invention), probes that are polypeptides, peptides or antibodies are favorably utilized. For example, isolated or recombinant polypeptides, polypeptide fragments and peptides derived from any of the amino acid sequences of the invention (e.g., SEQ ID NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6) and/or encoded by polynucleotide sequences of the invention, e.g., selected from SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5 are favorably used to identify and isolate antibodies, e.g., from phage display libraries, combinatorial libraries, polyclonal sera, and the like. Polypeptide fragments of the inventions include a peptide or polypeptide comprising an amino acid sequence of at least 5 contiguous amino acid residues, or at least 10 contiguous amino acid residues, or at least 15 contiguous amino acid residues, or at least 20 contiguous amino acid residues, or at least 25 contiguous amino

acid residues, or at least 40 contiguous amino acid residues, or at least 50 contiguous amino acid residues, or at least 60 contiguous amino residues, or at least 70 contiguous amino acid residues, or at least contiguous 80 amino acid residues, or at least contiguous 90 amino acid residues, or at least contiguous 100 amino acid residues, or at least contiguous 125 amino

5 acid residues, or at least 150 contiguous amino acid residues, or at least contiguous 175 amino acid residues, or at least contiguous 200 amino acid residues, or at least contiguous 250 amino acid residues, or at least contiguous 350, or at least contiguous 400, or at least contiguous 450, or at least contiguous 500, or at least contiguous 550 amino acid residues of the amino acid sequence an HA or NA polypeptide of the invention (e.g., SEQ ID NOS: 2, 10 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16- 340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6). Polynucleotides encoding said polypeptide fragments and antibodies that specifically bind said polypeptides are also preferred embodiments of the invention.

[0060] Antibodies specific for any polypeptide sequence or subsequence, e.g., of SEQ ID NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6, and/or encoded by polynucleotide sequences of the invention, e.g., selected from SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5 are likewise valuable as probes 20 for evaluating expression products, e.g., from cells or tissues. In addition, antibodies are particularly suitable for evaluating expression of proteins comprising amino acid subsequences, e.g., of those given herein, or encoded by polynucleotides sequences of the invention, e.g., selected from those shown herein, *in situ*, in a tissue array, in a cell, tissue or organism, e.g., an organism infected by an unidentified influenza virus or the like.

25 Antibodies can be directly labeled with a detectable reagent, or detected indirectly by labeling of a secondary antibody specific for the heavy chain constant region (i.e., isotype) of the specific antibody. Additional details regarding production of specific antibodies are provided below.

30 Diagnostic Assays

[0061] The polynucleotide sequences of the present invention can be used in diagnostic assays to detect influenza (and/or hemagglutinin and/or neuraminidase) in a sample, to detect hemagglutinin-like and/or neuraminidase-like sequences, and to detect strain

differences in clinical isolates of influenza using either chemically synthesized or recombinant polynucleotide fragments, e.g., selected from the sequences herein. For example, fragments of the hemagglutinin and/or neuraminidase sequences comprising at least between 10 and 20 nucleotides can be used as primers to amplify polynucleotides using polymerase chain reaction (PCR) methods well known in the art (e.g., reverse transcription-PCR) and as probes in polynucleotide hybridization assays to detect target genetic material such as influenza RNA in clinical specimens.

5 [0062] The probes of the invention, e.g., as exemplified by unique subsequences selected from those given herein, can also be used to identify additional useful polynucleotide

10 sequences (such as to characterize additional strains of influenza) according to procedures routine in the art. In one set of preferred embodiments, one or more probes, as described above, are utilized to screen libraries of expression products or cloned viral polynucleotides (i.e., expression libraries or genomic libraries) to identify clones that include sequences identical to, or with significant sequence identity to the sequences herein. In turn, each of 15 these identified sequences can be used to make probes, including pairs or sets of variant probes as described above. It will be understood that in addition to such physical methods as library screening, computer assisted bioinformatic approaches, e.g., BLAST and other sequence homology search algorithms, and the like, can also be used for identifying related polynucleotide sequences.

20 [0063] The probes of the invention are particularly useful for detecting the presence and for determining the identity of influenza polynucleotides in cells, tissues or other biological samples (e.g., a nasal wash or bronchial lavage). For example, the probes of the invention are favorably utilized to determine whether a biological sample, such as a subject (e.g., a human subject) or model system (such as a cultured cell sample) has been exposed to, or 25 become infected with influenza, or particular strain(s) of influenza. Detection of hybridization of the selected probe to polynucleotides originating in (e.g., isolated from) the biological sample or model system is indicative of exposure to or infection with the virus (or a related virus) from which the probe polynucleotide is selected.

[0064] It will be appreciated that probe design is influenced by the intended application.

30 For example, where several allele-specific probe-target interactions are to be detected in a single assay, e.g., on a single DNA chip, it is desirable to have similar melting temperatures for all of the probes. Accordingly, the lengths of the probes are adjusted so that the melting temperatures for all of the probes on the array are closely similar (it will be appreciated that

different lengths for different probes may be needed to achieve a particular  $T_m$  where different probes have different GC contents). Although melting temperature is a primary consideration in probe design, other factors are optionally used to further adjust probe construction, such as selecting against primer self-complementarity and the like.

5 [0065] Vectors, Promoters and Expression Systems

[0066] The present invention includes recombinant constructs incorporating one or more of the polynucleotide sequences described herein. Such constructs optionally include a vector, for example, a plasmid, a cosmid, a phage, a virus, a bacterial artificial chromosome (BAC), a yeast artificial chromosome (YAC), etc., into which one or more of the polynucleotide sequences of the invention, e.g., comprising any of SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5 or a subsequence thereof etc., has been inserted, in a forward or reverse orientation. For example, the inserted polynucleotide can include a viral chromosomal sequence or cDNA including all or part of at least one of the polynucleotide sequences of the invention. In one embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available.

[0067] The polynucleotides of the present invention can be included in any one of a variety of vectors suitable for generating sense or antisense RNA, and optionally, polypeptide (or peptide) expression products (e.g., a hemagglutinin and/or neuraminidase molecule of the invention, or fragments thereof). Such vectors include chromosomal, non-chromosomal and synthetic DNA sequences, e.g., derivatives of SV40; bacterial plasmids; phage DNA; baculovirus; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, pseudorabies, adenovirus, adeno-associated virus, retroviruses and many others (e.g., pCDL). Any vector that is capable of introducing genetic material into a cell, and, if replication is desired, which is replicable in the relevant host can be used.

[0068] In an expression vector, the HA and/or NA polynucleotide sequence of interest is physically arranged in proximity and orientation to an appropriate transcription control sequence (e.g., promoter, and optionally, one or more enhancers) to direct mRNA synthesis. That is, the polynucleotide sequence of interest is operably linked to an appropriate transcription control sequence. Examples of such promoters include: LTR or SV40

promoter, *E. coli* lac or trp promoter, phage lambda P<sub>L</sub> promoter, and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses.

[0069] A variety of promoters are suitable for use in expression vectors for regulating transcription of influenza virus genome segments. In certain embodiments, the

5 cytomegalovirus (CMV) DNA dependent RNA Polymerase II (Pol II) promoter is utilized. If desired, e.g., for regulating conditional expression, other promoters can be substituted which induce RNA transcription under the specified conditions, or in the specified tissues or cells. Numerous viral and mammalian, e.g., human promoters are available, or can be isolated according to the specific application contemplated. For example, alternative  
10 promoters obtained from the genomes of animal and human viruses include such promoters as the adenovirus (such as Adenovirus 2), papilloma virus, hepatitis-B virus, polyoma virus, and Simian Virus 40 (SV40), and various retroviral promoters. Mammalian promoters include, among many others, the actin promoter, immunoglobulin promoters, heat-shock promoters, and the like.

15 [0070] Transcription is optionally increased by including an enhancer sequence. Enhancers are typically short, e.g., 10-500 bp, cis-acting DNA elements that act in concert with a promoter to increase transcription. Many enhancer sequences have been isolated from mammalian genes (hemoglobin, elastase, albumin, alpha-fetoprotein, and insulin), and eukaryotic cell viruses. The enhancer can be spliced into the vector at a position 5' or 3' to  
20 the heterologous coding sequence, but is typically inserted at a site 5' to the promoter.

Typically, the promoter, and if desired, additional transcription enhancing sequences are chosen to optimize expression in the host cell type into which the heterologous DNA is to be introduced (Scharf et al. (1994) *Heat stress promoters and transcription factors* Results Probl Cell Differ 20:125-62; Kriegler et al. (1990) *Assembly of enhancers, promoters, and splice signals to control expression of transferred genes* Methods in Enzymol 185: 512-27).  
25 Optionally, the amplicon can also contain a ribosome binding site or an internal ribosome entry site (IRES) for translation initiation.

[0071] The vectors of the invention also favorably include sequences necessary for the termination of transcription and for stabilizing the mRNA, such as a polyadenylation site or  
30 a terminator sequence. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. In one embodiment, the SV40 polyadenylation signal sequences can provide a bi-directional

polyadenylation site that insulates transcription of (+) strand mRNA molecules from the PolI promoter initiating replication of the (-) strand viral genome.

[0072] In addition, as described above, the expression vectors optionally include one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells, in addition to genes previously listed, markers such as dihydrofolate reductase or neomycin resistance are suitable for selection in eukaryotic cell culture.

[0073] The vector containing the appropriate polynucleotide sequence as described above, as well as an appropriate promoter or control sequence, can be employed to transform a host cell permitting expression of the protein. While the vectors of the invention can be replicated in bacterial cells, most frequently it will be desirable to introduce them into mammalian cells, e.g., Vero cells, BHK cells, MDCK cell, 293 cells, COS cells, or the like, for the purpose of expression.

[0074] As described elsewhere, the HA and NA sequences herein, in various embodiments, can be comprised within plasmids involved in plasmid-rescue reassortant. *See, e.g.*, U.S.

Application Nos. 60/420,708, filed October 23, 2002; 60/574,117, filed May 24, 2004; 10/423,828, filed April 25, 2003; 60/578,962, filed June 12, 2004; and 10/870,690 filed June 16, 2004; and US20020164770, which are incorporated by reference herein. For example, preferred expression vectors of the invention include, but are not limited to, vectors comprising pol I promoter and terminator sequences or vectors using both the pol I and pol II promoters “the polI/polII promoter system” (e.g., Zobel *et al.*, Nucl. Acids Res. 1993, 21:3607; US20020164770; Neumann *et al.*, Proc. Natl. Acad. Sci. USA 1999, 96:9345; Fodor *et al.*, J. Virol. 1999, 73:9679; and US20030035814). The reassortants produced can include the HA and NA genes arranged with the 6 other influenza genes from the A/Ann Arbor/6/60 donor strain (and/or derivatives and modifications thereof), the PR8 donor strain backbone, the A/Leningrad/17 donor strain backbone, etc. Other backbone strains are described, for example, in 20040137013 and 20030147916, which are incorporated by reference herein.

#### [0075] Additional Expression Elements

[0076] Most commonly, the genome segment encoding the influenza virus HA and/or NA protein includes any additional sequences necessary for its expression, including translation into a functional viral protein. In other situations, a minigene, or other artificial construct encoding the viral proteins, e.g., an HA and/or NA protein, can be employed. Again, in such case, it is often desirable to include specific initiation signals that aid in the efficient

translation of the heterologous coding sequence. These signals can include, e.g., the ATG initiation codon and adjacent sequences. To insure translation of the entire insert, the initiation codon is inserted in the correct reading frame relative to the viral protein.

Exogenous transcriptional elements and initiation codons can be of various origins, both

5 natural and synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers appropriate to the cell system in use.

[0077] If desired, polynucleotide sequences encoding additional expressed elements, such as signal sequences, secretion or localization sequences, and the like can be incorporated into the vector, usually, in-frame with the polynucleotide sequence of interest, e.g., to target

10 polypeptide expression to a desired cellular compartment, membrane, or organelle, or to direct polypeptide secretion to the periplasmic space or into the cell culture media. Such sequences are known to those of skill, and include secretion leader peptides, organelle targeting sequences (e.g., nuclear localization sequences, ER retention signals, mitochondrial transit sequences), membrane localization/anchor sequences (e.g., stop 15 transfer sequences, GPI anchor sequences), and the like.

[0078] Where translation of a polypeptide encoded by a polynucleotide sequence of the

invention is desired, additional translation specific initiation signals can improve the

efficiency of translation. These signals can include, e.g., an ATG initiation codon and

adjacent sequences, an IRES region, etc. In some cases, for example, full-length cDNA

20 molecules or chromosomal segments including a coding sequence incorporating, e.g., a polynucleotide sequence of the invention (e.g., as in the sequences herein), a translation initiation codon and associated sequence elements are inserted into the appropriate expression vector simultaneously with the polynucleotide sequence of interest. In such cases, additional translational control signals frequently are not required. However, in cases

25 where only a polypeptide coding sequence, or a portion thereof, is inserted, exogenous translational control signals, including, e.g., an ATG initiation codon is often provided for expression of the relevant sequence. The initiation codon is put in the correct reading frame to ensure transcription of the polynucleotide sequence of interest. Exogenous

transcriptional elements and initiation codons can be of various origins, both natural and

30 synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers appropriate to the cell system in use (see, e.g., Scharf D. et al. (1994) Results Probl Cell Differ 20:125-62; Bittner et al. (1987) Methods in Enzymol 153:516-544).

## Production of recombinant virus

**[0079]** Negative strand RNA viruses can be genetically engineered and recovered using a recombinant reverse genetics approach (see, e.g., USPN 5,166,057 to Palese et al.). Such method was originally applied to engineer influenza viral genomes (Luytjes et al. (1989)

5 Cell 59:1107-1113; Enami et al. (1990) Proc. Natl. Acad. Sci. USA 92:11563-11567), and has been successfully applied to a wide variety of segmented and nonsegmented negative strand RNA viruses, e.g., rabies (Schnell et al. (1994) EMBO J. 13: 4195-4203); VSV (Lawson et al. (1995) Proc. Natl. Acad. Sci. USA 92: 4477-4481); measles virus (Radecke et al. (1995) EMBO J. 14:5773-5784); rinderpest virus (Baron & Barrett (1997) J. Virol. 71: 10 1265-1271); human parainfluenza virus (Hoffman & Banerjee (1997) J. Virol. 71: 3272-3277; Dubin et al. (1997) Virology 235:323-332); SV5 (He et al. (1997) Virology 237:249-260); canine distemper virus (Gassen et al. (2000) J. Virol. 74:10737-44); and Sendai virus (Park et al. (1991) Proc. Natl. Acad. Sci. USA 88: 5537-5541; Kato et al. (1996) Genes to Cells 1:569-579). Those of skill in the art will be familiar with these and similar techniques 15 to produce influenza virus comprising the HA and NA sequences of the invention.

Reassortant influenza viruses produced according to such methods are also a feature of the invention, as are reassortant influenza virus comprising one or more polynucleotides and/or polypeptides of the invention.

20 Cell Culture and Expression Hosts

**[0080]** The present invention also relates to host cells that are introduced (transduced, transformed or transfected) with vectors of the invention, and the production of polypeptides of the invention by recombinant techniques. Host cells are genetically engineered (i.e., transduced, transformed or transfected) with a vector, such as an expression vector, of this invention. As described above, the vector can be in the form of a plasmid, a viral particle, a phage, etc. Examples of appropriate expression hosts include: bacterial cells, such as *E. coli*, *Streptomyces*, and *Salmonella typhimurium*; fungal cells, such as *Saccharomyces cerevisiae*, *Pichia pastoris*, and *Neurospora crassa*; or insect cells such as *Drosophila* and *Spodoptera frugiperda*.

25 **[0081]** Most commonly, mammalian cells are used to culture the HA and NA molecules of the invention. Suitable host cells for the replication of influenza virus include, e.g., Vero cells, BHK cells, MDCK cells, 293 cells and COS cells, including 293T cells, COS7 cells or the like. Commonly, co-cultures including two of the above cell lines, e.g., MDCK cells

and either 293T or COS cells are employed at a ratio, e.g., of 1:1, to improve replication efficiency. Typically, cells are cultured in a standard commercial culture medium, such as Dulbecco's modified Eagle's medium supplemented with serum (e.g., 10% fetal bovine serum), or in serum free medium, under controlled humidity and CO<sub>2</sub> concentration suitable

5 for maintaining neutral buffered pH (e.g., at pH between 7.0 and 7.2). Optionally, the medium contains antibiotics to prevent bacterial growth, e.g., penicillin, streptomycin, etc., and/or additional nutrients, such as L-glutamine, sodium pyruvate, non-essential amino acids, additional supplements to promote favorable growth characteristics, e.g., trypsin, β-mercaptoethanol, and the like.

10 [0082] The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants, or amplifying the inserted polynucleotide sequences. The culture conditions, such as temperature, pH and the like, are typically those previously used with the particular host cell selected for expression, and will be apparent to those skilled in the art and in the references cited herein, including, e.g.,  
15 Freshney (1994) Culture of Animal Cells, a Manual of Basic Technique, 3<sup>rd</sup> edition, Wiley-Liss, New York and the references cited therein. Other helpful references include, e.g., Paul (1975) Cell and Tissue Culture, 5<sup>th</sup> ed., Livingston, Edinburgh; Adams (1980) Laboratory Techniques in Biochemistry and Molecular Biology-Cell Culture for Biochemists, Work and Burdon (eds.) Elsevier, Amsterdam. Additional details regarding tissue culture  
20 procedures of particular interest in the production of influenza virus *in vitro* include, e.g., Merten et al. (1996) *Production of influenza virus in cell cultures for vaccine preparation*. in Cohen and Shafferman (eds.) Novel Strategies in Design and Production of Vaccines, which is incorporated herein in its entirety for all purposes. Additionally, variations in such  
25 procedures adapted to the present invention are readily determined through routine experimentation and will be familiar to those skilled in the art.

[0083] Cells for production of influenza virus (e.g., having the HA and/or NA sequences of the invention) can be cultured in serum-containing or serum free medium. In some cases, e.g., for the preparation of purified viruses, it is typically desirable to grow the host cells in serum free conditions. Cells can be cultured in small scale, e.g., less than 25 ml medium,  
30 culture tubes or flasks or in large flasks with agitation, in rotator bottles, or on microcarrier beads (e.g., DEAE-Dextran microcarrier beads, such as Dormacell, Pfeifer & Langen; Superbead, Flow Laboratories; styrene copolymer-tri-methylamine beads, such as Hillex, SoloHill, Ann Arbor) in flasks, bottles or reactor cultures. Microcarrier beads are small

spheres (in the range of 100-200 microns in diameter) that provide a large surface area for adherent cell growth per volume of cell culture. For example a single liter of medium can include more than 20 million microcarrier beads providing greater than 8000 square centimeters of growth surface. For commercial production of viruses, e.g., for vaccine

5 production, it is often desirable to culture the cells in a bioreactor or fermenter. Bioreactors are available in volumes from under 1 liter to in excess of 100 liters, e.g., Cyto3 Bioreactor (Osmonics, Minnetonka, MN); NBS bioreactors (New Brunswick Scientific, Edison, NJ); laboratory and commercial scale bioreactors from B. Braun Biotech International (B. Braun Biotech, Melsungen, Germany).

10 **[0084]** Regardless of the culture volume, in many desired aspects of the current invention, it is important that the cultures be maintained at an appropriate temperature, to insure efficient recovery of recombinant and/or reassortant influenza virus using temperature dependent multi plasmid systems (*see, e.g.*, Multi-Plasmid System for the Production of Influenza Virus, U.S. Application No. 60/420,708, , filed October 23, 2002, U.S. Application No.

15 10/423,828, filed April 25, 2003, and U.S. Application No. 60/574,117, filed May 24, 2004), heating of virus solutions for filtration, etc. Typically, a regulator, e.g., a thermostat, or other device for sensing and maintaining the temperature of the cell culture system and/or other solution, is employed to insure that the temperature is at the correct level during the appropriate period (e.g., virus replication, etc.).

20 **[0085]** In some embodiments herein (e.g., wherein reassorted viruses are to be produced from segments on vectors) vectors comprising influenza genome segments are introduced (e.g., transfected) into host cells according to methods well known in the art for introducing heterologous polynucleotides into eukaryotic cells, including, e.g., calcium phosphate co-precipitation, electroporation, microinjection, lipofection, and transfection employing

25 polyamine transfection reagents. For example, vectors, e.g., plasmids, can be transfected into host cells, such as COS cells, 293T cells or combinations of COS or 293T cells and MDCK cells, using the polyamine transfection reagent TransIT-LT1 (Mirus) according to the manufacturer's instructions in order to produce reassorted viruses, etc. Thus, in one example, approximately 1  $\mu$ g of each vector is introduced into a population of host cells

30 with approximately 2  $\mu$ l of TransIT-LT1 diluted in 160  $\mu$ l medium, preferably serum-free medium, in a total volume of 200  $\mu$ l. The DNA:transfection reagent mixtures are incubated at room temperature for 45 minutes followed by addition of 800  $\mu$ l of medium. The transfection mixture is added to the host cells, and the cells are cultured as described via

other methods well known to those skilled in the art. Accordingly, for the production of recombinant or reassortant viruses in cell culture, vectors incorporating each of the 8 genome segments, (PB2, PB1, PA, NP, M, NS, HA and NA, e.g., of the invention) are mixed with approximately 20  $\mu$ l TransIT-LT1 and transfected into host cells. Optionally, 5 serum-containing medium is replaced prior to transfection with serum-free medium, e.g., Opti-MEM I, and incubated for 4-6 hours.

**[0086]** Alternatively, electroporation can be employed to introduce such vectors incorporating influenza genome segments into host cells. For example, plasmid vectors incorporating an influenza A or influenza B virus are favorably introduced into Vero cells 10 using electroporation according to the following procedure. In brief, approximately  $5 \times 10^6$  Vero cells, e.g., grown in Modified Eagle's Medium (MEM) supplemented with 10% Fetal Bovine Serum (FBS) are resuspended in 0.4 ml OptiMEM and placed in an electroporation cuvette. Twenty micrograms of DNA in a volume of up to 25  $\mu$ l is added to the cells in the cuvette, which is then mixed gently by tapping. Electroporation is performed according to 15 the manufacturer's instructions (e.g., BioRad Gene Pulser II with Capacitance Extender Plus connected) at 300 volts, 950 microFarads with a time constant of between 28-33 msec. The cells are remixed by gently tapping and approximately 1-2 minutes following 20 electroporation 0.7 ml MEM with 10% FBS is added directly to the cuvette. The cells are then transferred to two wells of a standard 6 well tissue culture dish containing 2 ml MEM, 10% FBS. The cuvette is washed to recover any remaining cells and the wash suspension is divided between the two wells. Final volume is approximately 3.5 mL. The cells are then incubated under conditions permissive for viral growth, e.g., at approximately 33 °C for cold adapted strains.

**[0087]** In mammalian host cells, a number of expression systems, such as viral-based 25 systems, can be utilized. In cases where an adenovirus is used as an expression vector, a coding sequence is optionally ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a nonessential E1 or E3 region of the viral genome will result in a viable virus capable of expressing the polypeptides of interest in infected host cells (Logan and Shenk (1984) Proc Natl Acad Sci 30 81:3655-3659). In addition, transcription enhancers, such as the rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

**[0088]** A host cell strain is optionally chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such

modifications of the protein include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation. Post-translational processing, which cleaves a precursor form into a mature form, of the protein is sometimes important for correct insertion, folding and/or function. Additionally proper location within a host cell (e.g., on the cell surface) is also important. Different host cells such as COS, CHO, BHK, MDCK, 293, 293T, COS7, etc. have specific cellular machinery and characteristic mechanisms for such post-translational activities and can be chosen to ensure the correct modification and processing of the current introduced, foreign protein.

[0089] For long-term, high-yield production of recombinant proteins encoded by, or having

subsequences encoded by, the polynucleotides of the invention, stable expression systems are optionally used. For example, cell lines, stably expressing a polypeptide of the invention, are transfected using expression vectors that contain viral origins of replication or endogenous expression elements and a selectable marker gene. For example, following the introduction of the vector, cells are allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells that successfully express the introduced sequences. Thus, resistant clumps of stably transformed cells, e.g., derived from single cell type, can be proliferated using tissue culture techniques appropriate to the cell type.

[0090] Host cells transformed with a nucleotide sequence encoding a polypeptide of the invention are optionally cultured under conditions suitable for the expression and recovery of the encoded protein from cell culture. The cells expressing said protein can be sorted, isolated and/or purified. The protein or fragment thereof produced by a recombinant cell can be secreted, membrane-bound, or retained intracellularly, depending on the sequence (e.g., depending upon fusion proteins encoding a membrane retention signal or the like) and/or the vector used.

[0091] Expression products corresponding to the polynucleotides of the invention can also be produced in non-animal cells such as plants, yeast, fungi, bacteria and the like. In addition to Sambrook, Berger and Ausubel, all *infra*, details regarding cell culture can be found in Payne et al. (1992) Plant Cell and Tissue Culture in Liquid Systems John Wiley & Sons, Inc. New York, NY; Gamborg and Phillips (eds.) (1995) Plant Cell, Tissue and Organ Culture: Fundamental Methods Springer Lab Manual, Springer-Verlag (Berlin Heidelberg

New York) and Atlas and Parks (eds.) The Handbook of Microbiological Media (1993) CRC Press, Boca Raton, FL.

[0092] In bacterial systems, a number of expression vectors can be selected depending upon the use intended for the expressed product. For example, when large quantities of a

5 polypeptide or fragments thereof are needed for the production of antibodies, vectors that direct high-level expression of fusion proteins that are readily purified are favorably employed. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene), in which the coding sequence of interest, e.g., sequences comprising those found herein, etc., can be ligated into the vector 10 in-frame with sequences for the amino-terminal translation initiating methionine and the subsequent 7 residues of beta-galactosidase producing a catalytically active beta galactosidase fusion protein; pIN vectors (Van Heeke & Schuster (1989) J Biol Chem 264:5503-5509); pET vectors (Novagen, Madison WI); and the like. Similarly, in the yeast 15 *Saccharomyces cerevisiae* a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase and PGH can be used for production of the desired expression products. For reviews, see Ausubel, *infra*, and Grant et al., (1987); Methods in Enzymology 153:516-544.

#### Nucleic Acid Hybridization

20 [0093] Comparative hybridization can be used to identify nucleic acids (e.g., SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5) of the invention, including conservative variations of nucleic acids of the invention. This comparative hybridization method is a preferred method of distinguishing nucleic acids of the invention.

25 In addition, target nucleic acids which hybridize to the nucleic acids represented by, e.g., those shown herein under high, ultra-high and ultra-ultra-high stringency conditions are features of the invention. Examples of such nucleic acids include those with one or a few silent or conservative nucleic acid substitutions as compared to a given nucleic acid sequence.

30 [0094] A test target nucleic acid is said to specifically hybridize to a probe nucleic acid when it hybridizes at least one-half as well to the probe as to the perfectly matched complementary target, i.e., with a signal to noise ratio at least one-half as high as hybridization of the probe and target under conditions in which a perfectly matched probe

binds to a perfectly matched complementary target with a signal to noise ratio that is at least about 5x-10x as high as that observed for hybridization to any of the unmatched target nucleic acids.

[0095] Nucleic acids “hybridize” when they associate, typically in solution. Nucleic acids hybridize due to a variety of well-characterized physico-chemical forces, such as hydrogen bonding, solvent exclusion, base stacking and the like. Numerous protocols for nucleic acid hybridization are well known in the art. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Acid Probes* part I chapter 2, “Overview of principles

10 of hybridization and the strategy of nucleic acid probe assays,” (Elsevier, New York), as well as in Ausubel, Sambrook, and Berger and Kimmel, all *below*. Hames and Higgins (1995) Gene Probes 1 IRL Press at Oxford University Press, Oxford, England, (Hames and Higgins 1) and Hames and Higgins (1995) Gene Probes 2 IRL Press at Oxford University Press, Oxford, England (Hames and Higgins 2) provide details on the synthesis, labeling, 15 detection and quantification of DNA and RNA, including oligonucleotides.

[0096] An example of stringent hybridization conditions for hybridization of complementary nucleic acids which have more than 100 complementary residues on a filter in a Southern or northern blot is 50% formalin with 1 mg of heparin at 42°C, with the hybridization being carried out overnight. An example of stringent wash conditions 20 comprises a 0.2x SSC wash at 65°C for 15 minutes (*see*, Sambrook, *infra* for a description of SSC buffer and other nucleic acid hybridization parameters). Often the high stringency wash is preceded by a low stringency wash to remove background probe signal. An example low stringency wash is 2x SSC at 40°C for 15 minutes. In general, a signal to noise ratio of 5x (or higher) than that observed for an unrelated probe in the particular 25 hybridization assay indicates detection of a specific hybridization.

[0097] After hybridization, unhybridized nucleic acids can be removed by a series of washes, the stringency of which can be adjusted depending upon the desired results. Low stringency washing conditions (e.g., using higher salt and lower temperature) increase sensitivity, but can produce nonspecific hybridization signals and high background signals. 30 Higher stringency conditions (e.g., using lower salt and higher temperature that is closer to the  $T_m$ ) lower the background signal, typically with primarily the specific signal remaining. *See, also*, Rapley, R. and Walker, J.M. eds., Molecular Biotechnology Handbook (Humana Press, Inc. 1998).

[0098] “Stringent hybridization wash conditions” in the context of nucleic acid hybridization experiments such as Southern and northern hybridizations are sequence dependent, and are different under different environmental parameters. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993), *supra*, and in Hames and

5 Higgins, 1 and 2. Stringent hybridization and wash conditions can easily be determined empirically for any test nucleic acid. For example, in determining highly stringent hybridization and wash conditions, the hybridization and wash conditions are gradually increased (e.g., by increasing temperature, decreasing salt concentration, increasing detergent concentration and/or increasing the concentration of organic solvents such as

10 formalin in the hybridization or wash), until a selected set of criteria is met. For example, the hybridization and wash conditions are gradually increased until a probe binds to a perfectly matched complementary target with a signal to noise ratio that is at least 5x as high as that observed for hybridization of the probe to an unmatched target.

[0099] In general, a signal to noise ratio of at least 2x (or higher, e.g., at least 5x, 10x, 20x, 15 50x, 100x, or more) than that observed for an unrelated probe in the particular hybridization assay indicates detection of a specific hybridization. Detection of at least stringent hybridization between two sequences in the context of the present invention indicates relatively strong structural similarity to, e.g., the nucleic acids of the present invention provided in the sequence listings herein.

20 [0100] “Very stringent” conditions are selected to be equal to the thermal melting point ( $T_m$ ) for a particular probe. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the test sequence hybridizes to a perfectly matched probe. For the purposes of the present invention, generally, “highly stringent” hybridization and wash conditions are selected to be about 5°C lower than the  $T_m$  for the specific sequence at a 25 defined ionic strength and pH (as noted below, highly stringent conditions can also be referred to in comparative terms). Target sequences that are closely related or identical to the nucleotide sequence of interest (e.g., “probe”) can be identified under stringent or highly stringent conditions. Lower stringency conditions are appropriate for sequences that are less complementary.

30 [0101] “Ultra high-stringency” hybridization and wash conditions are those in which the stringency of hybridization and wash conditions are increased until the signal to noise ratio for binding of a probe to a perfectly matched complementary target nucleic acid is at least 10x as high as that observed for hybridization to any unmatched target nucleic acids. A

target nucleic acid which hybridizes to a probe under such conditions, with a signal to noise ratio of at least one-half that of the perfectly matched complementary target nucleic acid is said to bind to the probe under ultra-high stringency conditions.

**[0102]** In determining stringent or highly stringent hybridization (or even more stringent

5 hybridization) and wash conditions, the hybridization and wash conditions are gradually increased (e.g., by increasing temperature, decreasing salt concentration, increasing detergent concentration and/or increasing the concentration of organic solvents, such as formamide, in the hybridization or wash), until a selected set of criteria are met. For example, the hybridization and wash conditions are gradually increased until a probe 10 comprising one or more polynucleotide sequences of the invention, e.g., sequences or unique subsequences selected from those given herein (e.g., SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5) and/or complementary polynucleotide sequences, binds to a perfectly matched complementary target (again, a nucleic acid 15 comprising one or more nucleic acid sequences or subsequences selected from those given herein and/or complementary polynucleotide sequences thereof), with a signal to noise ratio that is at least 2x (and optionally 5x, 10x, or 100x or more) as high as that observed for hybridization of the probe to an unmatched target (e.g., a polynucleotide sequence comprising one or more sequences or subsequences selected from known influenza 20 sequences present in public databases such as GenBank at the time of filing, and/or complementary polynucleotide sequences thereof), as desired.

**[0103]** Using the polynucleotides of the invention, or subsequences thereof, novel target nucleic acids can be obtained; such target nucleic acids are also a feature of the invention.

For example, such target nucleic acids include sequences that hybridize under stringent

25 conditions to a unique oligonucleotide probe corresponding to any of the polynucleotides of the invention, e.g., SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5).

**[0104]** Similarly, even higher levels of stringency can be determined by gradually

30 increasing the hybridization and/or wash conditions of the relevant hybridization assay. For example, those in which the stringency of hybridization and wash conditions are increased until the signal to noise ratio for binding of the probe to the perfectly matched complementary target nucleic acid is at least 10X, 20X, 50X, 100X, or 500X or more as

high as that observed for hybridization to any unmatched target nucleic acids. The particular signal will depend on the label used in the relevant assay, e.g., a fluorescent label, a colorimetric label, a radioactive label, or the like. A target nucleic acid which hybridizes to a probe under such conditions, with a signal to noise ratio of at least one-half that of the perfectly matched complementary target nucleic acid is said to bind to the probe under ultra-ultra-high stringency conditions and are also features of the invention.

5 [0105] Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides which they encode are substantially identical.

This occurs, e.g., when a copy of a nucleic acid is created using the maximum codon

10 degeneracy permitted by the genetic code.

[0106] Cloning, Mutagenesis and Expression of Biomolecules of Interest

[0107] General texts which describe molecular biological techniques, which are applicable to the present invention, such as cloning, mutation, cell culture and the like, include Berger and Kimmel, Guide to Molecular Cloning Techniques, Methods in Enzymology volume 152 Academic Press, Inc., San Diego, CA (Berger); Sambrook et al., Molecular Cloning - A Laboratory Manual (3rd Ed.), Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 2000 (“Sambrook”) and Current Protocols in Molecular Biology, F. M.

15 Ausubel et al., eds., Current Protocols, a joint venture between Greene Publishing

20 Associates, Inc. and John Wiley & Sons, Inc., (supplemented through 2002) (“Ausubel”)).

These texts describe mutagenesis, the use of vectors, promoters and many other relevant topics related to, e.g., the generation of HA and/or NA molecules, etc.

[0108] Various types of mutagenesis are optionally used in the present invention, e.g., to produce and/or isolate, e.g., novel or newly isolated HA and/or NA molecules and/or to

25 further modify/mutate the polypeptides (e.g., HA and NA molecules as in SEQ ID NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6) of the invention. They include but are not limited to site-directed, random point mutagenesis, homologous recombination (DNA shuffling), mutagenesis using uracil containing templates, oligonucleotide-directed

30 mutagenesis, phosphorothioate-modified DNA mutagenesis, mutagenesis using gapped duplex DNA or the like. Additional suitable methods include point mismatch repair, mutagenesis using repair-deficient host strains, restriction-selection and restriction-purification, deletion mutagenesis, mutagenesis by total gene synthesis, double-strand break

repair, and the like. Mutagenesis, e.g., involving chimeric constructs, is also included in the present invention. In one embodiment, mutagenesis can be guided by known information of the naturally occurring molecule or altered or mutated naturally occurring molecule, e.g., sequence, sequence comparisons, physical properties, crystal structure or the like.

5 [0109] The above texts and examples found herein describe these procedures as well as the following publications (and references cited within): Sieber, et al., *Nature Biotechnology*, 19:456-460 (2001); Ling et al., *Approaches to DNA mutagenesis: an overview*, *Anal Biochem* 254(2): 157-178 (1997); Dale et al., *Oligonucleotide-directed random mutagenesis using the phosphorothioate method*, *Methods Mol Biol* 57:369-374 (1996); I. A. Lorimer, I.

10 Pastan, *Nucleic Acids Res* 23, 3067-8 (1995); W. P. C. Stemmer, *Nature* 370, 389-91 (1994); Arnold, *Protein engineering for unusual environments*, *Current Opinion in Biotechnology* 4:450-455 (1993); Bass et al., *Mutant Trp repressors with new DNA-binding specificities*, *Science* 242:240-245 (1988); Fritz et al., *Oligonucleotide-directed construction of mutations: a gapped duplex DNA procedure without enzymatic reactions in vitro*, *Nucl Acids Res* 16: 6987-6999 (1988); Kramer et al., *Improved enzymatic in vitro reactions in the gapped duplex DNA approach to oligonucleotide-directed construction of mutations*, *Nucl Acids Res* 16: 7207 (1988); Sakamar and Khorana, *Total synthesis and expression of a gene for the  $\alpha$ -subunit of bovine rod outer segment guanine nucleotide-binding protein (transducin)*, *Nucl Acids Res* 14: 6361-6372 (1988); Sayers et al., *Y-T Exonucleases in phosphorothioate-based oligonucleotide-directed mutagenesis*, *Nucl Acids Res* 16:791-802 (1988); Sayers et al., *Strand specific cleavage of phosphorothioate-containing DNA by reaction with restriction endonucleases in the presence of ethidium bromide*, (1988) *Nucl Acids Res* 16: 803-814; Carter, *Improved oligonucleotide-directed mutagenesis using M13 vectors*, *Methods in Enzymol* 154: 382-403 (1987); Kramer & Fritz *Oligonucleotide-directed construction of mutations via gapped duplex DNA*, *Methods in Enzymol* 154:350-367 (1987); Kunkel, *The efficiency of oligonucleotide directed mutagenesis*, in *Nucleic Acids & Molecular Biology* (Eckstein, F. and Lilley, D.M.J. eds., Springer Verlag, Berlin)) (1987); Kunkel et al., *Rapid and efficient site-specific mutagenesis without phenotypic selection*, *Methods in Enzymol* 154, 367-382 (1987); Zoller & Smith, *Oligonucleotide-directed mutagenesis: a simple method using two oligonucleotide primers and a single-stranded DNA template*, *Methods in Enzymol* 154:329-350 (1987); Carter, *Site-directed mutagenesis*, *Biochem J* 237:1-7 (1986); Eghtedarzadeh & Henikoff, *Use of oligonucleotides to generate large deletions*, *Nucl Acids Res* 14: 5115 (1986); Mandecki,

25

30

*Oligonucleotide-directed double-strand break repair in plasmids of Escherichia coli: a method for site-specific mutagenesis*, Proc Natl Acad Sci USA, 83:7177-7181 (1986);

Nakamaye & Eckstein, *Inhibition of restriction endonuclease Nci I cleavage by phosphorothioate groups and its application to oligonucleotide-directed mutagenesis*, Nucl

5 Acids Res 14: 9679-9698 (1986); Wells et al., *Importance of hydrogen-bond formation in stabilizing the transition state of subtilisin*, Phil Trans R Soc Lond A 317: 415-423 (1986);

Botstein & Shortle, *Strategies and applications of in vitro mutagenesis*, Science 229:1193-1201(1985); Carter et al., *Improved oligonucleotide site-directed mutagenesis using M13 vectors*, Nucl Acids Res 13: 4431-4443 (1985); Grundström et al., *Oligonucleotide-directed*

10 *mutagenesis by microscale 'shot-gun' gene synthesis*, Nucl Acids Res 13: 3305-3316 (1985);

Kunkel, *Rapid and efficient site-specific mutagenesis without phenotypic selection*, Proc Natl Acad Sci USA 82:488-492 (1985); Smith, *In vitro mutagenesis*, Ann Rev Genet

19:423-462(1985); Taylor et al., *The use of phosphorothioate-modified DNA in restriction enzyme reactions to prepare nicked DNA*, Nucl Acids Res 13: 8749-8764 (1985); Taylor et

15 al., *The rapid generation of oligonucleotide-directed mutations at high frequency using phosphorothioate-modified DNA*, Nucl Acids Res 13: 8765-8787 (1985); Wells et al.,

*Cassette mutagenesis: an efficient method for generation of multiple mutations at defined sites*, Gene 34:315-323 (1985); Kramer et al., *The gapped duplex DNA approach to oligonucleotide-directed mutation construction*, Nucl Acids Res 12: 9441-9456 (1984);

20 Kramer et al., *Point Mismatch Repair*, Cell 38:879-887 (1984); Nambiar et al., *Total synthesis and cloning of a gene coding for the ribonuclease S protein*, Science 223: 1299-1301 (1984); Zoller & Smith, *Oligonucleotide-directed mutagenesis of DNA fragments cloned into M13 vectors*, Methods in Enzymol 100:468-500 (1983); and Zoller & Smith, *Oligonucleotide-directed mutagenesis using M13-derived vectors: an efficient and general*

25 *procedure for the production of point mutations in any DNA fragment*, Nucl Acids Res 10:6487-6500 (1982). Additional details on many of the above methods can be found in Methods in Enzymol Volume 154, which also describes useful controls for trouble-shooting problems with various mutagenesis, gene isolation, expression, and other methods.

30 **[0110]** Oligonucleotides, e.g., for use in mutagenesis of the present invention, e.g., mutating libraries of the HA and/or NA molecules of the invention, or altering such, are typically synthesized chemically according to the solid phase phosphoramidite triester method described by Beaucage and Caruthers, Tetrahedron Letts 22(20):1859-1862, (1981) e.g.,

using an automated synthesizer, as described in Needham-VanDevanter et al., Nucleic Acids Res, 12:6159-6168 (1984).

[0111] In addition, essentially any polynucleotide can be custom or standard ordered from any of a variety of commercial sources, such as The Midland Certified Reagent Company

5 (mcrc@oligos(dot)com), The Great American Gene Company (www(dot)genco(dot)com), ExpressGen Inc. (www(dot)expressgen(dot)com), Operon Technologies Inc. (Alameda, CA) and many others. Similarly, peptides and antibodies can be custom ordered from any of a variety of sources, such as PeptidoGenic (available at pkim@ccnet(dot)com), HTI Bio-products, Inc. (www(dot)htibio(dot)com), BMA Biomedicals Ltd. (U.K.), Bio.Synthesis,

10 Inc., and many others.

[0112] The present invention also relates to host cells and organisms comprising a HA and/or NA molecule or other polypeptide and/or polynucleotide of the invention, e.g., SEQ ID NOS: SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID

15 NO:5. Host cells are genetically engineered (e.g., transformed, transduced or transfected) with the vectors of this invention, which can be, for example, a cloning vector or an expression vector. The vector can be, for example, in the form of a plasmid, a bacterium, a

virus, a naked polynucleotide, or a conjugated polynucleotide. The vectors are introduced into cells and/or microorganisms by standard methods including electroporation (*see*, From

20 et al., Proc Natl Acad Sci USA 82, 5824 (1985), infection by viral vectors, high velocity

ballistic penetration by small particles with the nucleic acid either within the matrix of small beads or particles, or on the surface (Klein et al., Nature 327, 70-73 (1987)). Berger, Sambrook, and Ausubel provide a variety of appropriate transformation methods. *See*, above.

25 [0113] Several well-known methods of introducing target nucleic acids into bacterial cells are available, any of which can be used in the present invention. These include: fusion of the recipient cells with bacterial protoplasts containing the DNA, electroporation, projectile bombardment, and infection with viral vectors, etc. Bacterial cells can be used to amplify the number of plasmids containing DNA constructs of this invention. The bacteria are

30 grown to log phase and the plasmids within the bacteria can be isolated by a variety of methods known in the art (*see*, for instance, Sambrook). In addition, a plethora of kits are commercially available for the purification of plasmids from bacteria, (*see*, e.g., EasyPrep<sup>TM</sup>, FlexiPrep<sup>TM</sup>, both from Pharmacia Biotech; StrataClean<sup>TM</sup>, from Stratagene;

and, QIAprep™ from Qiagen). The isolated and purified plasmids are then further manipulated to produce other plasmids, used to transfect cells or incorporated into related vectors to infect organisms. Typical vectors contain transcription and translation terminators, transcription and translation initiation sequences, and promoters useful for regulation of the expression of the particular target nucleic acid. The vectors optionally comprise generic expression cassettes containing at least one independent terminator sequence, sequences permitting replication of the cassette in eukaryotes, or prokaryotes, or both, (e.g., shuttle vectors) and selection markers for both prokaryotic and eukaryotic systems. Vectors are suitable for replication and integration in prokaryotes, eukaryotes, or optionally both. *See*, Giliman & Smith, Gene 8:81 (1979); Roberts, et al., Nature, 328:731 (1987); Schneider, B., et al., Protein Expr Purif 6435:10 (1995); Ausubel, Sambrook, Berger (*all supra*). A catalogue of Bacteria and Bacteriophages useful for cloning is provided, e.g., by the ATCC, e.g., The ATCC Catalogue of Bacteria and Bacteriophage (1992) Gherna et al. (eds.) published by the ATCC. Additional basic procedures for sequencing, cloning and other aspects of molecular biology and underlying theoretical considerations are also found in Watson et al. (1992) Recombinant DNA Second Edition Scientific American Books, NY. *See*, above. Further vectors useful with the sequences herein are illustrated above in the section concerning production of influenza virus for vaccines and the references cited therein.

20

#### Polypeptide Production and Recovery

[0114] Following transduction of a suitable host cell line or strain and growth of the host cells to an appropriate cell density, the selected promoter is induced by appropriate means (e.g., temperature shift or chemical induction) and cells are cultured for an additional period. In some embodiments, a secreted polypeptide product, e.g., a HA and/or NA polypeptide as in a secreted fusion protein form, etc., is then recovered from the culture medium. In other embodiments, a virus particle containing a HA and/or a NA polypeptide of the invention is produced from the cell. Alternatively, cells can be harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification. Eukaryotic or microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents, or other methods, which are well known to those skilled in the art. Additionally, cells expressing a HA and/or a NA

polypeptide product of the invention can be utilized without separating the polypeptide from the cell. In such situations, the polypeptide of the invention is optionally expressed on the cell surface and is examined thus (e.g., by having HA and/or NA molecules (or fragments thereof, e.g., comprising fusion proteins or the like) on the cell surface bind antibodies, etc.

5 Such cells are also features of the invention.

[0115] Expressed polypeptides can be recovered and purified from recombinant cell cultures by any of a number of methods well known in the art, including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography (e.g., using any of the tagging systems known to those skilled in the art), hydroxylapatite chromatography, and lectin chromatography. Protein refolding steps can be used, as desired, in completing configuration of the mature protein. Also, high performance liquid chromatography (HPLC) can be employed in the final purification steps. In addition to the references noted herein, a variety of purification methods are well known in the art, including, e.g., those set forth in Sandana (1997) Bioseparation of Proteins, Academic Press, Inc.; and Bollag et al. (1996) Protein Methods, 2<sup>nd</sup> Edition Wiley-Liss, NY; Walker (1996) The Protein Protocols Handbook Humana Press, NJ, Harris and Angal (1990) Protein Purification Applications: A Practical Approach IRL Press at Oxford, Oxford, England; Harris and Angal Protein Purification Methods: A Practical Approach IRL Press at Oxford, Oxford, England; Scopes (1993) Protein Purification: Principles and Practice 3<sup>rd</sup> Edition Springer Verlag, NY; Janson and Ryden (1998) Protein Purification: Principles, High Resolution Methods and Applications, Second Edition Wiley-VCH, NY; and Walker (1998) Protein Protocols on CD-ROM Humana Press, NJ.

[0116] When the expressed polypeptides of the invention are produced in viruses, the viruses are typically recovered from the culture medium, in which infected (transfected) cells have been grown. Typically, crude medium is clarified prior to concentration of influenza viruses. Common methods include ultrafiltration, adsorption on barium sulfate and elution, and centrifugation. For example, crude medium from infected cultures can first be clarified by centrifugation at, e.g., 1000-2000 x g for a time sufficient to remove cell debris and other large particulate matter, e.g., between 10 and 30 minutes. Optionally, the clarified medium supernatant is then centrifuged to pellet the influenza viruses, e.g., at 15,000 x g, for approximately 3-5 hours. Following resuspension of the virus pellet in an appropriate buffer, such as STE (0.01 M Tris-HCl; 0.15 M NaCl; 0.0001 M EDTA) or

phosphate buffered saline (PBS) at pH 7.4, the virus is concentrated by density gradient centrifugation on sucrose (60%-12%) or potassium tartrate (50%-10%). Either continuous or step gradients, e.g., a sucrose gradient between 12% and 60% in four 12% steps, are suitable. The gradients are centrifuged at a speed, and for a time, sufficient for the viruses

5 to concentrate into a visible band for recovery. Alternatively, and for most large-scale commercial applications, virus is elutriated from density gradients using a zonal-centrifuge rotor operating in continuous mode. Additional details sufficient to guide one of skill through the preparation of influenza viruses from tissue culture are provided, e.g., in Furminger. *Vaccine Production*, in Nicholson et al. (eds.) Textbook of Influenza pp. 324-332; Merten et al. (1996) *Production of influenza virus in cell cultures for vaccine preparation*, in Cohen & Shafferman (eds.) Novel Strategies in Design and Production of Vaccines pp. 141-151, and United States Patent No. 5,690,937. If desired, the recovered viruses can be stored at -80°C in the presence of sucrose-phosphate-glutamate (SPG) as a stabilizer

15 [0117] Alternatively, cell-free transcription/translation systems can be employed to produce polypeptides comprising an amino acid sequence or subsequence of, e.g., the sequences given herein such as SEQ ID NOS: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6, or encoded by the polynucleotide sequences of the invention, e.g., SEQ ID NOS: 20 SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5. A number of suitable *in vitro* transcription and translation systems are commercially available. A general guide to *in vitro* transcription and translation protocols is found in Tymms (1995) In vitro Transcription and Translation Protocols: Methods in Molecular Biology Volume 37, 25 Garland Publishing, NY.

[0118] In addition, the polypeptides, or subsequences thereof, e.g., subsequences comprising antigenic peptides, can be produced manually or by using an automated system, by direct peptide synthesis using solid-phase techniques (see, Stewart et al. (1969) Solid-Phase Peptide Synthesis, WH Freeman Co, San Francisco; Merrifield J (1963) *J Am Chem Soc* 85:2149-2154). Exemplary automated systems include the Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer, Foster City, CA). If desired, subsequences can be chemically synthesized separately, and combined using chemical methods to provide full-length polypeptides.

### Modified Amino Acids

[0119] Expressed polypeptides of the invention can contain one or more modified amino acids. The presence of modified amino acids can be advantageous in, for example, (a)

5 increasing polypeptide serum half-life, (b) reducing/increasing polypeptide antigenicity, (c) increasing polypeptide storage stability, etc. Amino acid(s) are modified, for example, co-translationally or post-translationally during recombinant production (e.g., N-linked glycosylation at N-X-S/T motifs during expression in mammalian cells) or modified by synthetic means (e.g., via PEGylation).

10 [0120] Non-limiting examples of a modified amino acid include a glycosylated amino acid, a sulfated amino acid, a prenylated (e.g., farnesylated, geranylgeranylated) amino acid, an acetylated amino acid, an acylated amino acid, a PEG-ylated amino acid, a biotinylated amino acid, a carboxylated amino acid, a phosphorylated amino acid, and the like, as well as amino acids modified by conjugation to, e.g., lipid moieties or other organic derivatizing agents. References adequate to guide one of skill in the modification of amino acids are replete throughout the literature. Example protocols are found in Walker (1998) Protein Protocols on CD-ROM Human Press, Towata, NJ.

### Fusion Proteins

20 [0121] The present invention also provides fusion proteins comprising fusions of the sequences of the invention (e.g., encoding HA and/or NA polypeptides as exemplified by SEQ ID NOS: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6) or fragments thereof with, e.g., immunoglobulins (or portions thereof), sequences encoding, e.g., GFP 25 (green fluorescent protein), or other similar markers, etc. Nucleotide sequences encoding such fusion proteins are another aspect of the invention. Fusion proteins of the invention are optionally used for, e.g., similar applications (including, e.g., therapeutic, prophylactic, diagnostic, experimental, etc. applications as described herein) as the non-fusion proteins of the invention. In addition to fusion with immunoglobulin sequences and marker sequences, 30 the proteins of the invention are also optionally fused with, e.g., sequences which allow sorting of the fusion proteins and/or targeting of the fusion proteins to specific cell types, regions, etc.

Antibodies

[0122] The polypeptides of the invention can be used to produce antibodies specific for the polypeptides given herein and/or polypeptides encoded by the polynucleotides of the invention, e.g., those shown herein, and conservative variants thereof. Antibodies specific

5 for the above mentioned polypeptides are useful, e.g., for diagnostic and therapeutic purposes, e.g., related to the activity, distribution, and expression of target polypeptides.

[0123] Antibodies specific for the polypeptides of the invention can be generated by methods well known in the art. Such antibodies can include, but are not limited to, polyclonal, monoclonal, chimeric, humanized, single chain, Fab fragments and fragments 10 produced by an Fab expression library.

[0124] Polypeptides do not require biological activity for antibody production (e.g., full length functional hemagglutinin or neuraminidase is not required). However, the polypeptide or oligopeptide must be antigenic. Peptides used to induce specific antibodies typically have an amino acid sequence of at least about 4 amino acids, and often at least 5 or 15 10 amino acids. Short stretches of a polypeptide can be fused with another protein, such as keyhole limpet hemocyanin, and antibody produced against the chimeric molecule.

[0125] Numerous methods for producing polyclonal and monoclonal antibodies are known to those of skill in the art, and can be adapted to produce antibodies specific for the polypeptides of the invention, and/or encoded by the polynucleotide sequences of the

20 invention, etc. See, e.g., Coligan (1991) Current Protocols in Immunology Wiley/Greene, NY; Paul (ed.) (1998) Fundamental Immunology, Fourth Edition, Lippincott-Raven, Lippincott Williams & Wilkins; Harlow and Lane (1989) Antibodies: A Laboratory Manual Cold Spring Harbor Press, NY; Stites et al. (eds.) Basic and Clinical Immunology (4th ed.) Lange Medical Publications, Los Altos, CA, and references cited therein; Goding (1986)

25 Monoclonal Antibodies: Principles and Practice (2d ed.) Academic Press, New York, NY; and Kohler and Milstein (1975) *Nature* 256: 495-497. Other suitable techniques for antibody preparation include selection of libraries of recombinant antibodies in phage or similar vectors. See, Huse et al. (1989) *Science* 246: 1275-1281; and Ward, et al. (1989) *Nature* 341: 544-546. Specific monoclonal and polyclonal antibodies and antisera will 30 usually bind with a  $K_D$  of, e.g., at least about 0.1  $\mu\text{M}$ , at least about 0.01  $\mu\text{M}$  or better, and, typically and at least about 0.001  $\mu\text{M}$  or better.

[0126] For certain therapeutic applications, humanized antibodies are desirable. Detailed methods for preparation of chimeric (humanized) antibodies can be found in U.S. Patent

5,482,856. Additional details on humanization and other antibody production and engineering techniques can be found in Borrebaeck (ed.) (1995) Antibody Engineering, 2<sup>nd</sup> Edition Freeman and Company, NY (Borrebaeck); McCafferty et al. (1996) Antibody Engineering, A Practical Approach IRL at Oxford Press, Oxford, England (McCafferty), and Paul (1995) Antibody Engineering Protocols Humana Press, Towata, NJ (Paul). Additional details regarding specific procedures can be found, e.g., in Ostberg et al. (1983), Hybridoma 2: 361-367, Ostberg, U.S. Pat. No. 4,634,664, and Engelman et al., U.S. Pat. No. 4,634,666.

10 10 Defining Polypeptides by Immunoreactivity

[0127] Because the polypeptides of the invention provide a variety of new polypeptide sequences (e.g., comprising HA and NA molecules), the polypeptides also provide new structural features which can be recognized, e.g., in immunological assays. The generation of antisera which specifically bind the polypeptides of the invention, as well as the polypeptides which are bound by such antisera, are features of the invention.

[0128] For example, the invention includes polypeptides (e.g., HA and NA molecules) that specifically bind to or that are specifically immunoreactive with an antibody or antisera generated against an immunogen comprising an amino acid sequence selected from one or more of the sequences given herein (e.g., SEQ ID NOS: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6), etc. To eliminate cross-reactivity with other homologues, the antibody or antisera is subtracted with the HA and/or NA molecules found in public databases at the time of filing, e.g., the “control” polypeptide(s). Where the other control sequences correspond to a nucleic acid, a polypeptide encoded by the nucleic acid is generated and used for antibody/antisera subtraction purposes.

[0129] In one typical format, the immunoassay uses a polyclonal antiserum which was raised against one or more polypeptide comprising one or more of the sequences corresponding to the sequences herein (e.g., SEQ ID NOS: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6), etc. or a substantial subsequence thereof (i.e., at least about 30% of the full length sequence provided). The set of potential polypeptide immunogens derived from the present sequences are collectively referred to below as “the immunogenic polypeptides.” The resulting antisera is optionally selected to have low cross-

reactivity against the control hemagglutinin and/or neuraminidase homologues and any such cross-reactivity is removed, e.g., by immunoabsorption, with one or more of the control hemagglutinin and neuraminidase homologues, prior to use of the polyclonal antiserum in the immunoassay.

5 [0130] In order to produce antisera for use in an immunoassay, one or more of the immunogenic polypeptides is produced and purified as described herein. For example, recombinant protein can be produced in a recombinant cell. An inbred strain of mice (used in this assay because results are more reproducible due to the virtual genetic identity of the mice) is immunized with the immunogenic protein(s) in combination with a standard 10 adjuvant, such as Freund's adjuvant, and a standard mouse immunization protocol (*see, e.g.,* Harlow and Lane (1988) *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, New York, for a standard description of antibody generation, immunoassay formats and conditions that can be used to determine specific immunoreactivity). Additional references and discussion of antibodies is also found herein and can be applied 15 here to defining polypeptides by immunoreactivity. Alternatively, one or more synthetic or recombinant polypeptide derived from the sequences disclosed herein is conjugated to a carrier protein and used as an immunogen.

[0131] Polyclonal sera are collected and titered against the immunogenic polypeptide in an immunoassay, for example, a solid phase immunoassay with one or more of the 20 immunogenic proteins immobilized on a solid support. Polyclonal antisera with a titer of  $10^6$  or greater are selected, pooled and subtracted with the control hemagglutinin and/or neuraminidase polypeptide(s) to produce subtracted pooled titered polyclonal antisera.

[0132] The subtracted pooled titered polyclonal antisera are tested for cross reactivity 25 against the control homologue(s) in a comparative immunoassay. In this comparative assay, discriminatory binding conditions are determined for the subtracted titered polyclonal antisera which result in at least about a 5-10 fold higher signal to noise ratio for binding of the titered polyclonal antisera to the immunogenic polypeptides as compared to binding to the control homologues. That is, the stringency of the binding reaction is adjusted by the addition of non-specific competitors such as albumin or non-fat dry milk, and/or by 30 adjusting salt conditions, temperature, and/or the like. These binding conditions are used in subsequent assays for determining whether a test polypeptide (a polypeptide being compared to the immunogenic polypeptides and/or the control polypeptides) is specifically bound by the pooled subtracted polyclonal antisera. In particular, test polypeptides which

show at least a 2-5x higher signal to noise ratio than the control receptor homologues under discriminatory binding conditions, and at least about a ½ signal to noise ratio as compared to the immunogenic polypeptide(s), shares substantial structural similarity with the immunogenic polypeptide as compared to the known receptor, etc., and is, therefore a 5 polypeptide of the invention.

[0133] In another example, immunoassays in the competitive binding format are used for detection of a test polypeptide. For example, as noted, cross-reacting antibodies are removed from the pooled antisera mixture by immunoabsorption with the control polypeptides. The immunogenic polypeptide(s) are then immobilized to a solid support 10 which is exposed to the subtracted pooled antisera. Test proteins are added to the assay to compete for binding to the pooled subtracted antisera. The ability of the test protein(s) to compete for binding to the pooled subtracted antisera as compared to the immobilized protein(s) is compared to the ability of the immunogenic polypeptide(s) added to the assay to compete for binding (the immunogenic polypeptides compete effectively with the 15 immobilized immunogenic polypeptides for binding to the pooled antisera). The percent cross-reactivity for the test proteins is calculated, using standard calculations.

[0134] In a parallel assay, the ability of the control protein(s) to compete for binding to the pooled subtracted antisera is optionally determined as compared to the ability of the immunogenic polypeptide(s) to compete for binding to the antisera. Again, the percent 20 cross-reactivity for the control polypeptide(s) is calculated, using standard calculations. Where the percent cross-reactivity is at least 5-10x as high for the test polypeptides as compared to the control polypeptide(s) and or where the binding of the test polypeptides is approximately in the range of the binding of the immunogenic polypeptides, the test polypeptides are said to specifically bind the pooled subtracted antisera.

[0135] In general, the immunoabsorbed and pooled antisera can be used in a competitive binding immunoassay as described herein to compare any test polypeptide to the immunogenic and/or control polypeptide(s). In order to make this comparison, the immunogenic, test and control polypeptides are each assayed at a wide range of concentrations and the amount of each polypeptide required to inhibit 50% of the binding of 25 the subtracted antisera to, e.g., an immobilized control, test or immunogenic protein is determined using standard techniques. If the amount of the test polypeptide required for binding in the competitive assay is less than twice the amount of the immunogenic polypeptide that is required, then the test polypeptide is said to specifically bind to an 30

antibody generated to the immunogenic protein, provided the amount is at least about 5-10x as high as for the control polypeptide.

[0136] As an additional determination of specificity, the pooled antisera is optionally fully immunosorbed with the immunogenic polypeptide(s) (rather than the control

5 polypeptide(s)) until little or no binding of the resulting immunogenic polypeptide subtracted pooled antisera to the immunogenic polypeptide(s) used in the immunosorption is detectable. This fully immunosorbed antisera is then tested for reactivity with the test polypeptide. If little or no reactivity is observed (i.e., no more than 2x the signal to noise ratio observed for binding of the fully immunosorbed antisera to the immunogenic 10 polypeptide), then the test polypeptide is specifically bound by the antisera elicited by the immunogenic protein.

#### Nucleic acid and polypeptide sequence variants

[0137] As described herein, the invention provides for nucleic acid polynucleotide

15 sequences and polypeptide amino acid sequences, e.g., hemagglutinin and neuraminidase sequences, and, e.g., compositions and methods comprising said sequences. Examples of said sequences are disclosed herein (e.g., SEQ ID NOS: 1-8). However, one of skill in the art will appreciate that the invention is not necessarily limited to those sequences disclosed herein and that the present invention also provides many related and unrelated sequences 20 with the functions described herein, e.g., encoding a HA and/or a NA molecule.

[0138] One of skill will also appreciate that many variants of the disclosed sequences are included in the invention. For example, conservative variations of the disclosed sequences that yield a functionally identical sequence are included in the invention. Variants of the nucleic acid polynucleotide sequences, wherein the variants hybridize to at least one

25 disclosed sequence, are considered to be included in the invention. Unique subsequences of the sequences disclosed herein, as determined by, e.g., standard sequence comparison techniques, are also included in the invention.

#### Silent Variations

30 [0139] Due to the degeneracy of the genetic code, any of a variety of nucleic acid sequences encoding polypeptides and/or viruses of the invention are optionally produced, some which can bear lower levels of sequence identity to the HA and NA nucleic acid and polypeptide

sequences herein. The following provides a typical codon table specifying the genetic code, found in many biology and biochemistry texts.

**Table 1** Codon Table

Amino acids			Codon				
Alanine	Ala	A	GCA	GCC	GCG	GCU	
Cysteine	Cys	C	UGC	UGU			
Aspartic acid	Asp	D	GAC	GAU			
Glutamic acid	Glu	E	GAA	GAG			
Phenylalanine	Phe	F	UUC	UUU			
Glycine	Gly	G	GGA	GGC	GGG	GGU	
Histidine	His	H	CAC	CAU			
Isoleucine	Ile	I	AUA	AUC	AUU		
Lysine	Lys	K	AAA	AAG			
Leucine	Leu	L	UUA	UUG	CUA	CUC	CUG
Methionine	Met	M	AUG				
Asparagine	Asn	N	AAC	AAU			
Proline	Pro	P	CCA	CCC	CCG	CCU	
Glutamine	Gln	Q	CAA	CAG			
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG
Serine	Ser	S	AGC	AGU	UCA	UCC	UCG
Threonine	Thr	T	ACA	ACC	ACG	ACU	
Valine	Val	V	GUU	GUC	GUG	GUU	
Tryptophan	Trp	W	UGG				
Tyrosine	Tyr	Y	UAC	UAU			

**[0140]** The codon table shows that many amino acids are encoded by more than one codon.

5 For example, the codons AGA, AGG, CGA, CGC, CGG, and CGU all encode the amino acid arginine. Thus, at every position in the nucleic acids of the invention where an arginine is specified by a codon, the codon can be altered to any of the corresponding codons described above without altering the encoded polypeptide. It is understood that U in an RNA sequence corresponds to T in a DNA sequence.

[0141] Such “silent variations” are one species of “conservatively modified variations,” discussed below. One of skill will recognize that each codon in a nucleic acid (except ATG, which is ordinarily the only codon for methionine, and TTG, which is ordinarily the only codon for tryptophan) can be modified by standard techniques to encode a functionally identical polypeptide. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide is implicit in any described sequence. The invention, therefore, explicitly provides each and every possible variation of a nucleic acid sequence encoding a polypeptide of the invention that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code (e.g., as set forth in **Table 1**, or as is commonly available in the art) as applied to the nucleic acid sequence encoding a hemagglutinin or a neuraminidase polypeptide of the invention. All such variations of every nucleic acid herein are specifically provided and described by consideration of the sequence in combination with the genetic code. One of skill is fully able to make these silent substitutions using the methods herein.

#### Conservative variations

[0142] Owing to the degeneracy of the genetic code, “silent substitutions” (i.e., substitutions in a nucleic acid sequence which do not result in an alteration in an encoded polypeptide) are an implied feature of every nucleic acid sequence of the invention which encodes an amino acid. Similarly, “conservative amino acid substitutions,” in one or a few amino acids in an amino acid sequence are substituted with different amino acids with highly similar properties, are also readily identified as being highly similar to a disclosed construct such as those herein. Such conservative variations of each disclosed sequence are a feature of the present invention.

[0143] “Conservative variations” of a particular nucleic acid sequence refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or, where the nucleic acid does not encode an amino acid sequence, to essentially identical sequences, *see, Table 2 below*. One of skill will recognize that individual substitutions, deletions or additions which alter, add or delete a single amino acid or a small percentage of amino acids (typically less than 5%, more typically less than 4%, 3%, 2% or 1%) in an encoded sequence are “conservatively modified variations” where the alterations result in the deletion of an amino acid, addition of an amino acid, or substitution of an amino acid

with a chemically similar amino acid. Thus, “conservative variations” of a listed polypeptide sequence of the present invention include substitutions of a small percentage, typically less than 5%, more typically less than 4%, 3%, 2% or 1%, of the amino acids of the polypeptide sequence, with a conservatively selected amino acid of the same

5 conservative substitution group. Finally, the addition of sequences which do not alter the encoded activity of a nucleic acid molecule, such as the addition of a non-functional sequence, is a conservative variation of the basic nucleic acid.

Table 2 -- Conservative Substitution Groups

1	Alanine (A)	Serine (S)	Threonine (T)	
2	Aspartic acid (D)	Glutamic acid (E)		
3	Asparagine (N)	Glutamine (Q)		
4	Arginine (R)	Lysine (K)		
5	Isoleucine (I)	Leucine (L)	Methionine (M)	Valine (V)
6	Phenylalanine (F)	Tyrosine (Y)	Tryptophan (W)	

10

#### Unique Polypeptide and Polynucleotide Subsequences

**[0144]** In one aspect, the invention provides a nucleic acid which comprises a unique subsequence in a nucleic acid selected from the sequence of HA and NA molecules disclosed herein, e.g., SEQ ID NOS: SEQ ID NO:1, 3, 5, 7, residues 89-1063 of SEQ ID

15 NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5. The unique subsequence is unique as compared to a nucleic acids corresponding to nucleic acids such as, e.g., those found in GenBank or other similar public databases at the time of filing. Alignment can be performed using, e.g., BLAST set to default parameters. Any unique subsequence is useful, e.g., as a probe to

20 identify the nucleic acids of the invention. *See, above.*

**[0145]** Similarly, the invention includes a polypeptide which comprises a unique subsequence in a polypeptide selected from the sequence of HA and NA molecules disclosed herein, e.g., SEQ ID NOS: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ 25 ID NO:6. Here, the unique subsequence is unique as compared to a polypeptide corresponding to, e.g., the amino acid corresponding to polynucleotide sequences found in, e.g., GenBank or other similar public databases at the time of filing.

[0146] The invention also provides for target nucleic acids which hybridize under stringent conditions to a unique coding oligonucleotide which encodes a unique subsequence in a polypeptide selected from the sequences of HA and NA molecules of the invention wherein the unique subsequence is unique as compared to a polypeptide corresponding to any of the control polypeptides (sequences of, e.g., the nucleic acids corresponding to those found in, e.g., GenBank or other similar public databases at the time of filing). Unique sequences are determined as noted above.

5 Sequence comparison, identity, and homology

10 [0147] The terms “identical” or percent “identity,” in the context of two or more nucleic acid or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same, when compared and aligned for maximum correspondence, as measured using one of the sequence comparison algorithms described below (or other algorithms available to persons

15 of skill) or by visual inspection.

[0148] The phrase “substantially identical,” in the context of two nucleic acids or polypeptides (e.g., DNAs encoding a HA or NA molecule, or the amino acid sequence of a HA or NA molecule) refers to two or more sequences or subsequences that have at least about 90%, preferably 91%, most preferably 92%, 93%, 94%, 95%, 96%, 97%, 98%,

20 98.5%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9% or more nucleotide or amino acid residue identity, when compared and aligned for maximum correspondence, as measured using a sequence comparison algorithm or by visual inspection. Such “substantially identical” sequences are typically considered to be

25 “homologous,” without reference to actual ancestry. Preferably, “substantial identity” exists over a region of the amino acid sequences that is at least about 200 residues in length, at least about 250 residues, at least about 300 residues, 350 residues, 400 residues, 425 residues, 450 residues, 475 residues, 480 residues, 490 residues, 495 residues, 499 residues, 500 residues, 502 residues, 559 residues, 565 residues, or 566 residues, or over the full length of the two sequences to be compared.

30 [0149] For sequence comparison and homology determination, typically one sequence acts as a reference sequence to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are

designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

[0150] Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv Appl Math* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J Mol Biol* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc Natl Acad Sci USA* 85:2444 (1988), by computerized implementations of algorithms such as GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI, or by visual inspection (*see generally*, Ausubel et al., *supra*).

[0151] One example of an algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul et al., *J Mol Biol* 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information

([www\(dot\)ncbi.nlm.nih\(dot\)gov/](http://www.ncbi.nlm.nih.gov/)). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (*see*, Altschul et al., *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always  $> 0$ ) and N (penalty score for mismatching residues; always  $< 0$ ). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, a cutoff of 100, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W)

of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (see, Henikoff & Henikoff (1989) *Proc Natl Acad Sci USA* 89:10915).

[0152] In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul, *Proc Natl Acad Sci USA* 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

[0153] Another example of a useful sequence alignment algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments. It can also plot a tree showing the clustering relationships used to

create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle (1987) *J. Mol. Evol.* 35:351-360. The method used is similar to the

method described by Higgins & Sharp (1989) *CABIOS* 5:151-153. The program can align, e.g., up to 300 sequences of a maximum length of 5,000 letters. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing

a cluster of two aligned sequences. This cluster can then be aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences can be aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program can also be used to plot a dendrogram or tree representation of clustering relationships. The

program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison.

[0154] An additional example of an algorithm that is suitable for multiple DNA, or amino acid, sequence alignments is the CLUSTALW program (Thompson, J. D. et al. (1994) *Nucl. Acids. Res.* 22: 4673-4680). CLUSTALW performs multiple pairwise comparisons

between groups of sequences and assembles them into a multiple alignment based on homology. Gap open and Gap extension penalties can be, e.g., 10 and 0.05 respectively. For amino acid alignments, the BLOSUM algorithm can be used as a protein weight matrix. See, e.g., Henikoff and Henikoff (1992) *Proc. Natl. Acad. Sci. USA* 89: 10915-10919.

## Kits and Reagents

[0155] The present invention is optionally provided to a user as a kit. For example, a kit of the invention contains one or more nucleic acid, polypeptide, antibody, or cell line

5 described herein (e.g., comprising, or with, a HA and/or NA molecule of the invention).

The kit can contain a diagnostic nucleic acid or polypeptide, e.g., antibody, probe set, e.g., as a cDNA micro-array packaged in a suitable container, or other nucleic acid such as one or more expression vector. The kit can also further comprise, one or more additional reagents, e.g., substrates, labels, primers, for labeling expression products, tubes and/or 10 other accessories, reagents for collecting samples, buffers, hybridization chambers, cover slips, etc. The kit optionally further comprises an instruction set or user manual detailing preferred methods of using the kit components for discovery or application of diagnostic sets, etc.

[0156] When used according to the instructions, the kit can be used, e.g., for evaluating a 15 disease state or condition, for evaluating effects of a pharmaceutical agent or other treatment intervention on progression of a disease state or condition in a cell or organism, or for use as a vaccine, etc.

[0157] In an additional aspect, the present invention provides system kits embodying the methods, composition, systems and apparatus herein. System kits of the invention

20 optionally comprise one or more of the following: (1) an apparatus, system, system component or apparatus component; (2) instructions for practicing methods described herein, and/or for operating the apparatus or apparatus components herein and/or for using the compositions herein. In a further aspect, the present invention provides for the use of any apparatus, apparatus component, composition or kit herein, for the practice of any 25 method or assay herein, and/or for the use of any apparatus or kit to practice any assay or method herein.

[0158] Additionally, the kits can include one or more translation system as noted above (e.g., a cell) with appropriate packaging material, containers for holding the components of the kit, instructional materials for practicing the methods herein and/or the like. Similarly, 30 products of the translation systems (e.g., proteins such as HA and/or NA molecules) can be provided in kit form, e.g., with containers for holding the components of the kit, instructional materials for practicing the methods herein and/or the like.

[0159] To facilitate use of the methods and compositions of the invention, any of the vaccine components and/or compositions, e.g., reassorted virus in allantoic fluid, etc., and additional components, such as, buffer, cells, culture medium, useful for packaging and infection of influenza viruses for experimental or therapeutic vaccine purposes, can be packaged in the form of a kit. Typically, the kit contains, in addition to the above components, additional materials which can include, e.g., instructions for performing the methods of the invention, packaging material, and a container.

## EXAMPLES

[0160] H2N2 influenza viruses caused the 1957 pandemic and circulated in humans until 1968 when they were replaced by H3N2 influenza viruses. Having proven capable of causing disease, H2 viruses may have pandemic potential given the lack of H2 specific immunity in persons born after 1968. Fourteen geographically and temporally diverse H2 avian and human influenza viruses were evaluated for their ability to replicate and elicit a broad cross reactive antibody response in ferrets. Sera from ferrets that were inoculated with influenza A/Japan/57 (H2N2), A/mallard/NY/78 (H2N2) and A/swine/MO/2006 (H2N3) viruses elicited a broadly cross-reactive antibody response against heterologous H2 viruses in hemagglutination-inhibition and neutralization assays.

[0161] Using an A/Ann Arbor/6/60 (AA) cold-adapted (*ca*) (H2N2) backbone, three *ca* viruses were generated: *ca* A/Japan/57, *ca* A/mallard/NY/78 and *ca* A/swine/MO/2006. The HA and NA sequences of A/Japan/57 and A/swine/MO/2006 are shown in Table 3. The ability of each *ca* vaccine virus to protect against homologous and heterologous wild type (*wt*) H2 virus challenge was evaluated in ferrets. Efficacy of protection was variable in the upper respiratory tract. The *ca* AA and *ca* A/Japan/57 vaccines provided complete protection against a homologous challenge while the *ca* A/mallard/NY/78 and *ca* A/swine/MO/2006 vaccines provided partial protection from homologous challenge with a significant reduction in virus titers compared with mock-immunized animals. None of the *ca* vaccine viruses conferred complete protection against heterologous challenge in the upper respiratory tract. In the lower respiratory tract, each *ca* vaccine conferred complete protection from the challenge with the homologous *wt* virus. The *ca* AA and *ca* A/swine/MO/2006 vaccines provided complete protection in the lower respiratory tract against all heterologous *wt* challenge viruses.

Table 3. HA/NA sequences of the reassortant vaccine strains.

SEQ ID NO	HA or NA	Strain Name	Amino Acid or Nucleotide
<b>SEQ ID NO: 1</b>	HA (H3)	<i>ca</i> A/Japan/57	Nucleotide
<b>SEQ ID NO: 2</b>	HA (H3)	<i>ca</i> A/Japan/57	Amino Acid
<b>SEQ ID NO: 3</b>	NA (N2)	<i>ca</i> A/Japan/57	Nucleotide
<b>SEQ ID NO: 4</b>	NA (N2)	<i>ca</i> A/Japan/57	Amino Acid
<b>SEQ ID NO: 5</b>	HA (H3)	<i>ca</i> A/swine/MO/2006	Nucleotide
<b>SEQ ID NO: 6</b>	HA (H3)	<i>ca</i> A/swine/MO/2006	Amino Acid
<b>SEQ ID NO: 7</b>	NA (N3)	<i>ca</i> A/swine/MO/2006	Nucleotide
<b>SEQ ID NO: 8</b>	NA (N3)	<i>ca</i> A/swine/MO/2006	Amino Acid

[0162] Figures 1 and 2 show the efficacy of protection conferred by the *ca* AA, *ca* A/Japan/57, *ca* A/mallard/NY/78 and *ca* A/swine/MO/2006 vaccines in ferrets. Ferrets 5 were vaccinated with a single dose of *ca* reassortant virus vaccine. The ferrets were then challenged with wt AA, wt A/Japan/57, wt A/mallard/NY/78 and wt A/swine/MO/2006 influenza virus. Three days post challenge lungs and nasal turbinates of the ferrets were harvested and virus titer in the tissues was determined. Figures 1 and 2 shows efficacy of protection conferred by the recombinant H2 vaccines against homologous and heterologous 10 wild-type H2 viruses in lungs and nasal turbinates, respectively, in ferrets.

[0163] Figures 3 and 4 show the efficacy of protection conferred by the *ca* AA, *ca* A/Japan/57, *ca* A/mallard/NY/78 and *ca* A/swine/MO/2006 vaccines in mice. Mice were vaccinated with a single dose of *ca* reassortant virus vaccine. The mice were then challenged with wt AA, wt A/Japan/57, wt A/mallard/NY/78 and wt A/swine/MO/2006 15 influenza virus. Three days post challenge lungs and nasal turbinates of the mice were harvested and virus titer in the tissues was determined. Figures 3 and 4 show efficacy of protection conferred by the recombinant H2 vaccines against homologous and heterologous wild-type H2 viruses in lungs and nasal turbinates, respectively, in mice.

[0164] While the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be clear to one skilled in the art from a reading of this disclosure that various changes in form and detail can be made without departing from the true scope of the invention. For example, all the techniques and apparatus described above may be used in various combinations. All publications, patents, patent applications, or other 20

documents cited in this application are incorporated by reference in their entirety for all purposes to the same extent as if each individual publication, patent, patent application, or other document were individually indicated to be incorporated by reference for all purposes.

## SEQUENCES

ca A/Japan/57SEQ ID NO:1 Nucleotide Sequence of *ca A/Japan/57 H2*

Entire molecule length: 1773 nt

5	1 agcaaaagca ggggttatac catagacaac caaaagcaaa acaatggcca
	51 tcatttatct cattctcctg ttcacagcag tgagagggga ccagatatgc
101	attggatacc atgccaataa ttccacagag aaggtcgaca caattctaga
151	gcggaacgtc actgtgactc atgccaagga cattcttgag aagaccata
201	acggaaaagtt atgcaaacta aacggaatcc ctccacttga actaggggac
10	251 tgttagcattg ccggatggct ccttgaaat ccagaatgtg ataggcttct
	301 aagtgtgcca gaatggtcct atataatgga gaaagaaaac ccgagagacg
351	gtttgtgtta tccaggcagc ttcaatgatt atgaagaatt gaaacatctc
401	ctcagcagcg taaaacattt cgagaaagta aagattctgc ccaaagatag
451	atggacacag catacaacaa ctggaggtc acgggcctgc gcggtgtctg
15	501 gtaatccatc attcttcagg aacatggct ggctgacaaa gaaaggatca
	551 gattatccgg ttgccaagg atcgtacaac aatacaagcg gagaacaaat
601	gctaataatt tgggggggtgc accatccaa tcatgagaca gaacaaagaa
651	cattgtacca gaatgtggga acctatgtt ccgtaggcac atcaacattg
701	aacaaaaggt caacccaga aatagcaaca aggcctaaag tgaatggaca
20	751 aggaggtaga atggaattct cttggaccct cttggatatg tggacacca
	801 taaattttga gactactggt aatctaattt caccagagta tggattcaaa
851	atatcgaaaa gaggtgttgc agggatcatg aaaacagaag gaacacttga
901	gaactgtgag accaaatgcc aaactcctt gggagcaata aatacaacat
951	tgcctttca caatgtccac ccactgacaa taggtgagtg ccccaaataat
25	1001 gtaaaatccgg agaagttggc cttagcaaca ggactaagga atgttcccc
	1051 gattgaatca agaggattgt ttggggcaat agctggttt atagaaggag
1101	gatggcaagg aatggttgat gttggatgt gataccatca cagcaatgac
1151	cagggatcag ggtatgcagc agacaaagaa tccactcaa aggcatattga
1201	tggaaatcacc aacaaggtaa attctgtgat tggaaagatg aacacccaaat
30	1251 ttgaagctgt tggaaagaa ttcatgact tagagagaag actggagaac
	1301 ttgaacaaaa agatggaga cgggtttcta gatgtgtgga catacaatgc
1351	tgagcttcta gttctgtatgg aaaatgagag gacacttgac tttcatgatt
1401	ctaattgtcaa gaatctgtat gataaaagtca gaatgcagct gagagacaaac
1451	gtcaaagaac taggaaatgg atgtttgaa ttttatcaca aatgtgatga
35	1501 tgaatgcatt aatagtgtga aaaacgggac gtatgattat cccaaatgt
	1551 aagaagagtc taaaactaaat agaaatgaaa tcaaagggtt aaaattgagc
1601	agcatggggg tttatcaaatt ctttgccatt tatgtacag tagcaggttc
1651	tctgtcactg gcaatcatga tggctggat ctcttctgg atgtgctcca
1701	acgggtctct gcagtgcagg atctgcattt gattataagt cattttataaa
40	1751 taaaaaacac cttgtttct act

SEQ ID NO:2 Amino acid sequence of *ca A/Japan/57 H2*

Entire molecule length: 562 aa

45	1 maiylillf tavrgdqici gyhannstek vdtilernvt vthakdilek
	51 thngklickln gipplelgdc siagwllgnp ecdrl11svpe wsyimekenp
101	rdglcypgsf ndyeelkhll ssvkhfekvk ilpkdrwtqh tttggsraca
151	vsgnpsffrn mvwlkkgsd ypvakgsynn tsgeqmliliw gvhhpndete

201 qrtlyqnvgt yvsvgtstln krstpeiatr pkvngqggm efswtlldmw  
 251 dtinfestgn liapeygfki skrgssgimk tegtlencet kcqtplgain  
 301 ttlpfhnvhp ltigecpkv kseklvlatg lrnvpqiesr glfgaiagfi  
 351 eggwqgmvdg wygyhhsndq gsgyaadkes tqkafdgith kvnsviekmn  
 5 401 tqfeavgkef snlerrlenl nkkmedgfld vwtynaellv lmenertldf  
 451 hdsnvknlyd kvrnqlrdnv kelngncfef yhkcddecmn svknngtydyp  
 501 kyeeesklnr neikgvklss mgvyqilaiy atvagslsla immagisfwm  
 551 csngslqcri ci

10 SEQ ID NO:3 Nucleotide Sequence of *ca A/Japan/57 N2*

Entire molecule length: 1466 nt

1 agcaaaagca ggagtgaaaa tgaatccaaa tcaaaaagata ataacaattg  
 51 gctctgtctc ttcaccatt gaaacagtat gcttcctcat gcagattgcc  
 101 atcctggcaa ctactgtgac attgcatttt aagcaacatg agtgcgactc  
 15 151 ccccgcgagc aaccaagtaa tgccatgtga accaataata atagaaagga  
 201 acataacaga gatagtgtat ttgaataaca ccaccataga gaaagagatt  
 251 tgccccgaag tagtggata cagaaattgg tcaaagccgc aatgtcaaat  
 301 tacaggattt gcacctttt ctaaggacaa ttcaatccgg cttctgctg  
 351 gtggggacat ttgggtgacg agagaacctt atgtgtcatg cgatcctggc  
 20 401 aagtgttatac aatttgcact cgggcagggg accacactag acaacaaaca  
 451 ttcaaatggc acaatacatg atagaatccc tcatacgaacc ctattaatga  
 501 atgagttggg tggccattt catttagaa ccaaacaagt gtgtgttagca  
 551 tggccagct caagttgtca cgatggaaaa gcatggttgc atgtttgtgt  
 601 cactggggat gatagaaatg caactgctag cttcatttat gacgggaggc  
 25 651 ttgtggacag tattggtca tggctcaaa atatcctcag gacccaggag  
 701 tcggaatgcg tttgtatcaa tggacttgc acagtagtaa tgactgatgg  
 751 aagtgcata ggaagagccg atactagaat actattcatt aaagagggga  
 801 aaattgtcca tattagccca ttgtcaggaa gtgctcagca tatagaggag  
 851 tggccctgtt accctcgata tcctgacgtc agatgtatct gcagagacaa  
 30 901 ctggaaaggc tctaataaggc ccgttataga cataaatatg gaagattata  
 951 gcattgattc cagttatgt tgctcaggcc ttgttggcga cacacccagg  
 1001 aacgacgaca gctctagcaa tagcaattgc agggatccta acaatgagag  
 1051 agggaatcca ggagtggaaag gctggccctt tgacaatgga gatgatgtat  
 1101 ggatgggaag aacaatcagc aaagattcac gtcaggtt tgaaactttc  
 35 1151 aaagtcatgt gtgggtggc cacaccta atccaaatcgc aggtcaatag  
 1201 acaggtcata gttgacaaca ataattggc tggttactct ggtatttct  
 1251 ctgttgaggc caaaagctgc atcaataggt gctttatgt ggagttgata  
 1301 aggggaaggc cacaggagac tagatgtgg tggacctcaa acagtattgt  
 1351 tgggtttgt ggcacttcag gtacttatgg aacaggctca tggcctgatg  
 40 1401 gggcgaacat caatttcatg cctatataag ctttcgcaat ttttagaaaa  
 1451 actccttgc tctact

SEQ ID NO:4 Amino acid sequence of *ca A/Japan/57 N2*

Entire molecule length: 471 aa

45 1 mnnpnqkitti gsvslietv cflmqiaila ttvtlhfkqh ecdspasnqv  
 51 mpcepiier niteivylnn ttiiekeicpe vveyrnwskp qcqitgfapf  
 101 skdnsirlsa ggdiiwvtrep yvsdpkcy qfalgqggtt dnkhsngtih

151 driphrtllm nelgvpfhlg tkqvcvawss sschdgkawl hvcvtgddrn  
 201 atasfiydgr lvdsgwsq nilrtqesec vcingtctvv mtdgsasgra  
 251 dtrilfikeg kivhisplsg saqhieecsc yprypdvrci crdnwkgsnr  
 301 pvidinmedy sidssyvcsg lvgdptrndd sssnsncrdp nnergnpgvk  
 5 351 gwafdnngddv wmgrtiskds rsgyetfkvi ggwstpnks qvnrvqvivdn  
 401 nnwsgysgif svegkscinr cfyvelirgr pqetrvwwts nsivvfcgts  
 451 gtygtgswpd ganinfmp

ca A/swine/MO/200610 SEQ ID NO:5 Nucleotide Sequence of *ca A/swine/MO/2006 H2*

Entire molecule length: 1772 nt

1 agcaaaagca ggggttatac catagacaac cgaacaaaga caatgaccat  
 51 cactttctc atcctcctgt tcacagtagt gaaaggggac caaatatgca  
 101 tcggatacca tgccaacaat tccacagaaa aagttgacac aatcttggaa  
 151 cggaaacgtca ccgtgactca tgccagaac attcttgaaa agacgcataa  
 201 tggaaagttg tgcagattga gtggaatccc tccattggaa ctgggggatt  
 251 gcagcattgc aggttggctc cttggaaatc cgaatgtga ccggctctt  
 301 agtgtacctg aatggtccta tatagtggaa aaggaaaacc cggtaatgg  
 351 tctgtgctat ccaggcagtt tcaatgatta ttaggaattt aaacatcttc  
 20 401 tcaccagtgt gacacacttt gagaaagttt agattctgcc cagagatcaa  
 451 tggaccgcac acacaacaac tggtggttct cgggcctgtg cagtatctgg  
 501 aaacccgtca ttcttagga acatggtttgc ttacaaag aaagggtcaa  
 551 actactcaat tgctaaaagg tcataacaaca acacaagtgg ggagcaaatg  
 601 ctggtaatat ggggataca tcaccccaat gacgatgcgg aacagaggac  
 25 651 actgtaccag aatgtggaa catatgtttc cgttggaaaca tcaacactaa  
 701 ataagaggtc aatccctgaa atagcaacaa ggcacaaagt caatggacag  
 751 ggaggaagaa tggatttctc ttggactcta ttggagacat gggatgtcat  
 801 aaatttttagt agcactggta atttaatttgc accagaatac ggattcaaaa  
 851 tatcaaagag aggaagctca ggaattatga agacagagaa aatacttggaa  
 30 901 aattgtggaa ccaaattgtca gaccccttg gggcaataa atacaacatt  
 951 gcccac aacattcacc cattgacaat agtgagtgc cccaaatgt  
 1001 taaagtca tagactgatt ttggcgacag gagtaagaaa tggcccccac  
 1051 attgaatcaa gggattttt tggagcaata gctgggttta tagaaggcgg  
 1101 atggcaaggg atggttgtatgg gctggatgg gtaccatcac agcaatgatc  
 35 1151 aaggatcagg atatgcagca gacaaagaat ccactcaaaa ggcaattgtat  
 1201 gggataacta acaaagtaaa ttctgtgatt gaaaagatga acactcagtt  
 1251 tgaggctgtt gggaaagagt tcaacaacctt agagagaagg ctggaaaact  
 1301 taaataaaaaa gatggaaagat ggatttttg atgtatggac atataatgcc  
 1351 gaactcctag ttctaatttggaa aatgagagg acacttgatt tccatgattc  
 40 1401 taatgtgaag aatctgtacg ataaggtcag aatgcaattt agagacaatg  
 1451 ctaagggaaat agggAACGGA tgctttgagt ttatcataa atgtatgtat  
 1501 gaatgcatttga atagtgtcag gaatgggaca tatgattatc ccaaataatgt  
 1551 ggaagagtcc aagctgaaca ggaacgaaat caaaggagtg aaattgagca  
 1601 atatgggggt ttatcaaata cttgctatat acgctacagt tgcaggctct  
 45 1651 ttgtcactgg caatcatgat agctggatt tctttctgga tggttctaa  
 1701 tgggtctctg caatgcagaa tttgcatatg actgtaaatc aatttgtat  
 1751 taaaaacacc cttgtttcttta ct

SEQ ID NO:6 Amino acid sequence of *ca* A/swine/MO/2006 H2

Entire molecule length: 562 aa

1 mtitflillf tvvkqdqici gyhannstek vdtilernvt vthaknilek  
 51 thngklcrls gipplelgdc siagwllgnp ecdrl1svpe wsyivekenp  
 5 101 vnglcypgsf ndyeelkhll tsvthfekvk ilprdqwtqh tttggsraca  
 151 vsgnpsffrn mwltkkgsn ysiakrsynn tsgeqmlviw gihhpnddae  
 201 qrtlyqnvgt yvsvgtstln krsipeiatr pkvngqggm efswtlletw  
 251 dvinfestgn liapeygfki skrgssgimk tekilencet kcqtplgain  
 301 tlpfhnihp ltigecpkvv ksdrllatg vrnvlpqiesr glfgaiagfi  
 10 351 eggwqgmvdg wygyhhsndq gsgyaadkes tqkaidgitn kvnsviekmn  
 401 tqfeavgkef nnlerrlenl nkkmedgfid vwtynaellv lmenertldf  
 451 hdsnvknlyd kvrnqlrdna keingncfef yhkcddcmn svrnngttdyp  
 501 kyeeesklnr neikgvklnsn mgvyqilaiy atvagslsla imiagisfwm  
 551 csngslqcri ci

15

SEQ ID NO: 7 Nucleotide Sequence of *ca* A/swine/MO/2006 N3

Entire molecule length: 1453 nt

1 agcaaaaagca ggtgcgagat gaatccgaat cagaagataa taacaatcg  
 51 ggtagtgaat accactctgt caacaatagc cttcttcatt ggagtggaa  
 20 101 acttaatttt caacacagtc atacatgaga aaataggaga ccatcaaata  
 151 gtgacccatc caacaataac gaccctgca gtaccgaact gcagtgacac  
 201 tataataaca tacaataaca ctgtgataaa caacataaca acaacaataa  
 251 taactgaaga agaaaggcct ttcaagtctc cactaccgct gtgcccctc  
 301 agaggattct tccctttca caaggacaat gcaatacgac tgggtgaaaa  
 25 351 caaagacgtc atagtcacaa gagagcctt ttttagctgc gataatgaca  
 401 actgctggtc ctttgctctc acacaaggag cattgctagg gaccaaacat  
 451 agcaatggga ccattaaaga caggacacca tataaggctc taattcggtt  
 501 cccaatagga acagctccag tactaggaaa ttataaagag atatgcattg  
 551 cttggcgtcag cagcagttgc tttgacgggaa aagagtggat gcatgtgtgc  
 30 601 atgacagggaa acgataatga tgcaagtgc cagataat atggagggag  
 651 aatgacagac tccattaaat catggagaaa ggacatacta agaactcagg  
 701 agtctgaatg ccaatgcatt gacgggactt gtgttgc tgcacagat  
 751 ggcctgctg ctaatagtgc agattacagg gttactgga tacgggaggg  
 801 aaaaataata aagtatgaaa atgttccaa aacaaagata caacacttag  
 35 851 aagaatgttc ctgctatgtg gacattgtat ttactgtat atgttagggac  
 901 aattggaaagg gctctaacag accttggatg agaatcaaca acgagactat  
 951 actggaaaca gggtatgtat gtagtaaatt ccactcagac acccccaggc  
 1001 ccgctgaccc ttcaacaatg tcatgtgact ccccaagcaa tgtcaatgg  
 1051 ggacccggag tgaagggtt tggttcaaa gctggcgatg atgtatgg  
 40 1101 aggtagaaca gtgtcgacta gtggtagatc gggcttggaa attatcaaag  
 1151 ttacagaagg gtggatcaac ttccttaacc atgtcaaatc aattacacaa  
 1201 acactagtgc caaacaatga ctggtcaggc tattccggta gcttcattgt  
 1251 caaagccaag gactgtttc agccctgttt ttatgttgc gttatacgag  
 1301 ggaggcccaa caagaatgat gacgtctctt ggacaagtaa tagtatagtt  
 45 1351 actttctgtg gactagacaa tgaacctgga tcgggaaatt ggccagatgg  
 1401 ttcttaacatt gggttatgc ccaagtaata gaaaaaaagca cttgtttct  
 1451 act

SEQ ID NO: 8 Amino acid sequence of *ca* A/swine/MO/2006 N2

Entire molecule length: 469 aa

1 mnppnqkiiti gvvnttlsti alligvgnli fntvihekig dhqivtypti  
51 ttpavpncsd titynntvi nnitttiite eerpfkspip lcpfrgffpf  
5 101 hkdnairlge nkdvivtrep yvscdndncw sfaltqgall gtkhsngtik  
151 drtpyrlir fpigtapvlg nykeiciaws ssscfdgkew mhvcmtgndn  
201 dasaqiyygg rmtdsikswr kdilrtqese cqcidgtcvv avtdgpaans  
251 adyrvwyire gkiikyenvp ktkiqhleec scyvdidvyc icrdnwkgsn  
301 rpwmrinnet iletgyvcsk fhsdtprpad pstmcdsps nvnggpgvkg  
10 351 fggfkagddvw lgrtvstsgs sgfeiikvte gwinspnhvk sitqtlvpnn  
401 dwsgyssgsfi vkakdcfqpc fyvelirgrp nknddvswts nsivtfcgld  
451 nepgsgnwpd gsnigfmpk

**What is claimed is:**

1. An isolated polypeptide, wherein said polypeptide is selected from the group consisting of:

5           i) a polypeptide encoded by a polynucleotide sequence as shown in any one of SEQ ID NOS: 1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5;

10           ii) a polypeptide comprising an amino acid sequence as shown in any one of SEQ ID NOS: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6;

15           iii) the mature form of the polypeptide comprising the amino acid sequence as shown in any one of SEQ ID NOS: 2, 4, 6, and 8;

20           iv) a polypeptide comprising an amino acid sequence selected from SEQ ID NO: 2, 4, 6, 8, residues 16-340 of SEQ ID NO:2, residues 341-562 of SEQ ID NO:2, residues 16-340 of SEQ ID NO:6, and residues 341-562 of SEQ ID NO:6 in which one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen or fifteen amino acid residues were substituted, inserted or deleted.

2. An immunogenic composition comprising an immunologically effective amount of at least one polypeptide of claim 1.

3. An isolated antibody that specifically binds the polypeptide of claim 1.

4. A method for stimulating the immune system of an individual to produce a

25 protective immune response against influenza virus comprising:

5           i) administering to the individual an immunologically effective amount of the polypeptide of claim 1 in a physiologically acceptable carrier.

5. A reassortant influenza virus comprising the polypeptide of claim 1.

6. An immunogenic composition comprising an immunologically effective amount of 30 the reassortant influenza virus of claim 5.

7. A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus comprising:

i) administering to the individual an immunologically effective amount of the reassortant influenza virus of claim 5 in a physiologically acceptable carrier.

8. An isolated polynucleotide, wherein said polynucleotide is selected from the group

5 consisting of:

i) a polynucleotide comprising a nucleotide sequence as shown in any one of SEQ ID NOS: 1, 3, 5, 7, residues 89-1063 of SEQ ID NO:1, residues 1064-1729 of SEQ ID NO:1, residues 88-1062 of SEQ ID NO:5 and residues 1063-1728 of SEQ ID NO:5, or a complementary sequence thereof;

10 ii) a polynucleotide sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% sequence identity to the polynucleotide sequence of (a);

15 iii) a polynucleotide of (a) in which less than one, less than two, less than three, less than four, less than five, less than six, less than seven, less than eight, less than nine, less than ten, less than eleven, less than twelve, less than thirteen, less than fourteen, less than fifteen, less than twenty, less than twenty five, less than thirty, less than forty, or less than fifty nucleotide residues were substituted, inserted or deleted; and

20 iv) a polynucleotide comprising a nucleotide sequence encoding the polypeptide of claim 1.

9. An immunogenic composition comprising the polynucleotide of claim 8.

10. A reassortant influenza virus comprising the polynucleotide of claim 8.

11. An immunogenic composition comprising an immunologically effective amount of 25 the reassortant influenza virus of claim 10.

12. A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus comprising:

30 i) administering to the individual an immunologically effective amount of the reassortant influenza virus of claim 10 in a physiologically acceptable carrier.

13. A cell comprising the polynucleotide of claim 8.

14. A vector comprising the polynucleotide of claim 8.

15. The vector of claim 14, wherein the vector is a plasmid, a cosmid, a phage, a virus, or a fragment of a virus.

16. The vector of claim 14, wherein the vector is an expression vector.

17. A cell comprising the vector of any one of claims 14-16.

5 18. A method of producing an isolated or recombinant polypeptide, the method comprising:

i) culturing the cell of claim 13 or 17 in a suitable culture medium under conditions permitting expression of said nucleic acid; and,

ii) isolating the polypeptide from the cell or the medium.

10 19. A 6:2 reassortant influenza virus, wherein said virus comprises 6 internal genome segments from one or more donor viruses and at least a first genome segment encoding a first immunogenic influenza surface antigen, wherein the first influenza surface antigen comprises the polypeptide of claim 1.

15 20. The reassortant virus of claim 19, further comprising a second genome segment encoding a second immunogenic influenza surface antigen, wherein the second influenza surface antigen comprises the polypeptide of claim 1.

20 21. A 6:2 reassortant influenza virus, wherein said virus comprises 6 internal genome segments from one or more donor viruses and at least a first genome segment encoding a first immunogenic influenza surface antigen, wherein the first surface antigen encoding genome segment comprises the polynucleotide of claim 8.

22. The reassortant virus of claim 19, further comprising a second genome segment encoding a second immunogenic influenza surface antigen, wherein the second surface antigen encoding genome segment comprises the polynucleotide of claim 8.

23. The reassortant virus of any one of claims 19-22, wherein the one or more donor virus is A/Ann Arbor/6/60.

24. The reassortant virus of any one of claims 19-22, wherein the one or more donor viruses is other than A/Ann Arbor/6/60.

25. The reassortant virus of claim 24, wherein the one or more donor viruses is PR8.

26. The reassortant virus of claim 24, wherein the one or more donor viruses is A/Leningrad/17.

27. The reassortant virus of any one of claims 19-26, wherein the virus comprises a hemagglutinin having a modified polybasic cleavage site.

28. The reassortant virus of any one of claims 19-27, wherein said internal genome segments of the one or more donor viruses confer one or more of the following properties: temperature sensitive, cold adapted, or attenuated.

29. An immunogenic composition comprising an immunologically effective amount of 5 the reassortant influenza virus of any one of claims 19-28.

30. A live attenuated influenza vaccine comprising the reassortant influenza virus of any one of claims 19-28.

31. A split virus or killed virus vaccine comprising the reassortant influenza virus of any one of claims 19-28.

10 32. A live attenuated influenza vaccine comprising the composition of claim 29.

33. A split virus or killed virus vaccine comprising the composition of claim 29.

34. A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus comprising:

i) administering to the individual an immunologically effective amount of 15 the reassortant influenza virus of any one of claims 19-28 in a physiologically effective carrier.

35. A method of prophylactic or therapeutic treatment of a viral infection in a subject, the method comprising:

i) administering to the subject the reassortant influenza virus of any one of 20 claims 19-28 in an amount effective to produce an immunogenic response against the viral infection.

36. A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus comprising:

i) administering to the individual an immunologically effective amount of 25 the immunogenic composition of claim 29.

37. A method of prophylactic or therapeutic treatment of a viral infection in a subject, the method comprising:

i) administering to the subject the immunogenic composition of claim 29 in 30 an amount effective to produce an immunogenic response against the viral infection.

38. A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus comprising:

i) administering to the individual an immunologically effective amount of the vaccine of any one of claims 30-33.

39. A method of prophylactic or therapeutic treatment of a viral infection in a subject, the method comprising:

5 i) administering to the subject the vaccine of any one of claims 30-33 in an amount effective to produce an immunogenic response against the viral infection.

40. The method of any one of claims 34-39, wherein the subject is a human.

41. An isolated antibody that specifically binds the reassortant influenza virus of any 10 one of claims 19-28.

42. A method for producing influenza viruses in cell culture comprising:

i) introducing into a population of host cells a plurality of vectors comprising nucleotide sequences corresponding to at least 6 internal genome segments from one or more donor viruses; and, at least a first genome 15 segment encoding a first immunogenic influenza surface antigen, wherein said first surface antigen comprises the polypeptide of claim 1;

ii) culturing the population of host cells; and,  
iii) recovering a plurality of influenza viruses.

43. The method of claim 42, wherein the plurality of vectors further comprises a

20 nucleotide sequence corresponding to a second genome segment encoding a second immunogenic influenza surface antigen, wherein the second influenza surface antigen comprises the polypeptide of claim 1.

44. A method for producing influenza viruses in cell culture, the method comprising:

i) introducing into a population of host cells a plurality of vectors comprising nucleotide sequences corresponding to at least 6 internal genome segments from one or more donor viruses; and, at least a first genome 25 segment encoding a first immunogenic influenza surface antigen, wherein said first genome segment comprises the polynucleotide of claim 8;

ii) culturing the population of host cells; and,  
iii) recovering a plurality of influenza viruses.

45. The method of claim 44, wherein the plurality of vectors further comprises a nucleotide sequence corresponding to a second genome segment encoding a second immunogenic influenza surface antigen, wherein the second surface antigen encoding genome segment comprises the polynucleotide of claim 8.

5 46. The method of any one of claims 42-45, wherein the one or more donor virus is A/Ann Arbor/6/60.

47. The method of any one of claims 42-45, wherein the one or more donor viruses is other than A/Ann Arbor/6/60.

48. The method of claim 47, wherein the one or more donor viruses is PR8.

10 49. The method of claim 47, wherein the one or more donor viruses is A/Leningrad/17.

50. The method of any one of claims 42-49, wherein the virus comprises a hemagglutinin having a modified polybasic cleavage site.

51. The method of any one of claims 42-50, wherein said one or more donor viruses have one or more of the following properties: temperature sensitive, cold adapted, or

15 attenuated.

52. An immunogenic composition comprising an immunologically effective amount of the influenza virus produced by the method of any one of claims 42-51.

53. A live attenuated influenza vaccine comprising the immunogenic composition of claim 52.

20 54. A split virus or killed virus vaccine comprising the immunogenic composition of claim 52.

55. A live attenuated influenza vaccine comprising the virus produced by the method of any one of claims 42-51.

56. A split virus or killed virus vaccine comprising the virus produced by the method of 25 any one of claims 42-51.

57. A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus comprising:

30 i) administering to the individual an immunologically effective amount of the influenza virus produced by the method of any one of claims 42-51 in a physiologically effective carrier.

58. A method of prophylactic or therapeutic treatment of a viral infection in a subject, the method comprising:

i) administering to the subject the influenza virus produced by the method of any one of claims 42-51 in an amount effective to produce an immunogenic response against the viral infection.

59. A method for stimulating the immune system of an individual to produce a

5 protective immune response against influenza virus comprising:

i) administering to the individual the immunogenic composition of claim 52.

60. A method of prophylactic or therapeutic treatment of a viral infection in a subject,

the method comprising:

10 i) administering to the subject the immunogenic composition of claim 52 in an amount effective to produce an immunogenic response against the viral infection.

61. A method for stimulating the immune system of an individual to produce a

protective immune response against influenza virus comprising:

15 i) administering to the individual the vaccine of any one of claims 53-56.

62. A method of prophylactic or therapeutic treatment of a viral infection in a subject,

the method comprising:

i) administering to the subject the vaccine of any one of claims 53-56 in an amount effective to produce an immunogenic response against the viral infection.

20

63. An isolated antibody that specifically binds the influenza virus produced by the

method of any one of claims 42-51.