



(51) International Patent Classification:

C07K 16/10 (2006.01) A61K 39/00 (2006.01)
A61P 31/14 (2006.01)

(21) International Application Number:

PCT/US2022/077481

(22) International Filing Date:

03 October 2022 (03.10.2022)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

63/251,472 01 October 2021 (01.10.2021) US

(71) Applicant: **ACADEMIA SINICA**; 128, Academia Road, Sec. 2, Nankang, Taipei 115 (TW).

(71) Applicant (for CN only): **LUI, Fu-Tong** [US/US]; P.O. Box 855, West Sacramento, CA 95691 (US).

(72) Inventors: **LIN, Kuo-I**; 17F-2, No. 15, Ln. 61, Sec. 2, Academia Road, Nangang Dist., Taipei 11565 (TW). **MA, Che**; 15F, No. 54 Dongshih Street, Xizhi District, New Taipei City 22141 (TW). **WONG, Chi-Huey**; 6F, No. 25, Ln. 105,

Sec. 1, Hangzhou South Road, Zhongzheng Dist., Taipei City 100 (TW). **WANG, Szu-Wen**; 1F., No. 279-4, Sec. 2, Xiyuan Rd., Wanhua Dist., Taipei City 108035 (TW). **CHANG, Yi-Hsuan**; No. 367, Sec. 4 Ziyou Rd., East Dist, Taichung City 401 (TW). **CHEN, Xiaorui**; 3F., No. 38, Aly. 4 Ln. 69, Sec. 5 Minsheng E. Rd., Songshan Dist., Taipei 105 (TW). **HUANG, Han-Yi**; 12F., No. 185, Zhongshan Rd., Luzhu Dist., Taoyuan City 338 (TW).

(74) Agent: **YANG, Kay** et al.; PROSYLA GROUP PC, 4275 Executive Square, Suite 200, La Jolla, CA 92037 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CV, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IQ, IR, IS, IT, JM, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH,

(54) Title: ANTIBODY SPECIFIC TO SPIKE PROTEIN OF SARS-COV-2 AND USES THEREOF




FIG. 1A

(57) Abstract: The present disclosure relates to an antibody or antigen-binding fragment thereof that specifically binds to a spike protein of SARS-CoV-2. The present disclosure also relates to a pharmaceutical composition, a method for treating and/or preventing diseases and/or disorders caused by a coronavirus in a subject in need thereof, and a method for detecting a coronavirus in a sample.

WO 2023/056482 A1

TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS,
ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, ME, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

ANTIBODY SPECIFIC TO SPIKE PROTEIN OF SARS-COV-2 AND USES THEREOF

PRIORITY INFORMATION

5 [0001] The present application claims priority today and benefit of United States Provisional Patent Application No. 63/251,472, filed October 1, 2021, the disclosure of which is incorporated in its entirety herein.

SEQUENCE LISTING

10 [0002] The instant application contains a Sequence Listing which is submitted electronically in .xml format and is hereby incorporated by reference in its entirety. The .xml copy, created on September 30, 2022, is named "G4590-14200PCT_SeqListing_20220930.xml" and is 10 kilobytes in size.

FIELD OF THE DISCLOSURE

15 [0003] The present disclosure relates to an antibody or antigen-binding fragment thereof, which is specific to a spike protein of SARS-CoV-2, and uses thereof.

BACKGROUND OF THE DISCLOSURE

20 [0004] The COVID-19 pandemic caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has spread globally. The infection causes symptoms of direct cytopathic effects and excessive inflammatory responses in the infected subject. The lack of valid treatment causes high morbidity and mortality. The emergence of these newly identified viruses highlights the need for the development of novel antiviral strategies.

[0005] Thus, there is need for development of an effective treatment for COVID-19.

SUMMARY OF THE DISCLOSURE

25 [0006] The present disclosure provides novel neutralizing therapeutic anti-Coronavirus spike protein (such as anti-SARS-CoV-2-Spike protein) antibodies and their use for treating or preventing viral infection.

30 [0007] Accordingly, the present disclosure provides an antibody or antigen-binding fragment thereof that is specific for an epitope in a spike protein of coronaviruses (CoVs), particularly, SARS-CoV-2. The antibody according to the disclosure is thus useful for treating and/or preventing diseases and/or disorders caused by or related to CoVs, particularly SARS-CoV-2. The antibody of the disclosure is also useful for detecting CoVs (particularly, SARS-CoV-2).

[0008] The present disclosure also provides an epitope comprises a part locating in 419 to 433 amino acid residues or 471 to 482 amino acid residues of SEQ ID NO: 8. In some embodiments, the epitope comprises a part locating in 419 to 433 amino acid residues and 471 to 482 amino

acid residues of SEQ ID NO: 8. Accordingly, the present disclosure provides an antibody binding to the epitope disclosed herein.

[0009] In some embodiments, the disclosure provides an antibody or antigen-binding fragment thereof that specifically binds to an epitope in a spike protein of a CoV; wherein the antibody or antigen-binding fragment thereof comprises complementarity determining regions (CDRs) of a heavy chain variable region and complementarity determining regions of a light chain variable region, wherein the complementarity determining regions of the heavy chain variable region comprise CDRH1, CDRH2, and CDRH3 regions, and the complementarity determining regions of the light chain variable region comprise CDRL1, CDRL2, and CDRL3 regions, and wherein:

the CDRH1 region comprises the amino acid sequence of GYTFTEYT (SEQ ID NO: 1) or a substantially similar sequence thereof; the CDRH2 region comprises the amino acid sequence of INPNIGDT (SEQ ID NO: 2) or a substantially similar sequence thereof; the CDRH3 region comprises the amino acid sequence of AREVYNYSFAY (SEQ ID NO: 3) or a substantially similar sequence thereof; and

the CDRL1 region comprises the amino acid sequence of QSLLYSSNQKNY (SEQ ID NO: 4) or a substantially similar sequence thereof; the CDRL2 region comprises the amino acid sequence of WAS or a substantially similar sequence thereof; the CDRL3 region comprises the amino acid sequence of QQYYRYPLT (SEQ ID NO: 5) or a substantially similar sequence thereof.

[0010] In some embodiments of the disclosure, the spike protein is fully glycosylated. In one embodiment, the spike protein is monoglycosylated.

[0011] In some embodiments of the disclosure, the antibody or antigen-binding fragment thereof comprises a heavy chain variable region comprising the amino acid sequence of MGWSLILLFLVAVATRVEVQLQQSGPEMVKPGASVKISCKTSGYTFTEYTIYWVKQSH GKSLEWLGGINPNIGDTTYNQKFKGKATLTVDTSSSTAYMELRSLTSEDSAVYYCARE VYNYSFAYWGQGLTVTSAASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTV SWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKK AEPKSCDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEV KFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKDYKCKVSNKA LPAPIEKTISKAKGQPREPQVYTLPPSRDELTRNQVSLTCLVKGFYPSDIAVEWESNGQP ENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO: 6) or a substantially similar sequence thereof; and a light chain variable region comprising the amino acid sequence of MRVPAQLLGLLLLWLPGARCDIVMSQSPSSLAVSVGEKVTMCKSSQSLLYSSNQKN YLAWYQQKLGQTPKLLIYWASSRESGVPDRFTGSGSGTDFTLTISSVRAEDLAVYYCQ QQYYRYPLTFGVGKLELKRVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQ

WKVDNALQSGNSQESVTEQDSKDYESTYLSSTLTLSKADYEKHKVYACEVTHQGLSSP
VTKSFNRGEC (SEQ ID NO: 7) or a substantially similar sequence thereof.

[0012] In some embodiments of the disclosure, the antibody is a monoclonal antibody, chimeric antibody, humanized antibody or human antibody.

5 [0013] In some embodiments of the disclosure, the antibody is multispecific.

[0014] The present disclosure also provides a complex comprising the antibody or antigen-binding fragment thereof as disclosed herein bound to a spike protein of a CoV or fragments thereof.

10 [0015] The present disclosure provides a vector encoding the antibody or antigen-binding fragment thereof as disclosed herein.

[0016] The present disclosure provides a genetically engineered cell expressing the antibody or antigen-binding fragment thereof as disclosed herein or containing the vector as disclosed herein.

15 [0017] The present disclosure also provides a method for manufacturing the antibody or antigen-binding fragment thereof as disclosed herein, comprising: (a) introducing into a host cell one or more polynucleotides encoding said antibody or antigen-binding fragment; (b) culturing the host cell under conditions favorable to expression of the one or more polynucleotides; and (c) optionally, isolating the antibody or antigen-binding fragment from the host cell and/or a medium in which the host cell is grown.

20 [0018] The present disclosure provides a pharmaceutical composition comprising the antibody or antigen-binding fragment thereof as disclosed herein and pharmaceutically acceptable carrier and, optionally, a further therapeutic agent.

25 [0019] Examples of the therapeutic agent include but are not limited to an antiviral agent. In some embodiments of the disclosure, the therapeutic agent is an anti-inflammatory agent or an antibody or antigen-binding fragment thereof that specifically binds to a spike protein of a CoV.

[0020] In some embodiments of the disclosure, the subject is vaccinated.

[0021] The present disclosure provides a vessel or injection device comprising the antibody or antigen-binding fragment thereof as disclosed herein.

30 [0022] The present disclosure provides use of the antibody or antigen-binding fragment thereof as described herein in the manufacture of a medicament for treating or preventing infection with a coronavirus in a subject in need thereof. Alternatively, the present disclosure provides a method for treating or preventing infection with a coronavirus in a subject in need thereof, comprising administering a therapeutically effective amount of the antibody or antigen-binding fragment thereof as disclosed herein.

[0023] In some embodiments of the disclosure, the CoV described herein is alpha-CoV, beta-CoV, gamma-CoV, and delta-CoV2. In some embodiments, the CoV described herein includes, but is not limited to, SARS-CoV, MERS-CoV or SARS-CoV-2.

[0024] In some embodiments of the disclosure, the medicament is administered with one or more further therapeutic agents. Alternatively, the subject is administered one or more further therapeutic agents.

[0025] The present disclosure provides a method for neutralizing a coronavirus in a subject in need thereof, comprises administering to the subject the antibody or antigen-binding fragment thereof as disclosed herein.

[0026] The present disclosure provides a method for administering the antibody or antigen-binding fragment thereof as disclosed herein into the body of a subject comprising injecting the antibody or antigen-binding fragment into the body of the subject.

[0027] In some embodiments of the disclosure, the medicament is injected into the body of the subject subcutaneously, intravenously, or intramuscularly. Alternatively, the antibody or antigen-binding fragment is injected into the body of the subject subcutaneously, intravenously, or intramuscularly.

[0028] The present disclosure provides a method for detecting a coronavirus in a sample comprising contacting the sample with the antibody or antigen-binding fragment thereof as disclosed herein.

[0029] The present disclosure provides a kit for detecting a coronavirus in a sample, wherein the kit comprises the antibody or antigen-binding fragment thereof as disclosed herein.

[0030] The present disclosure is described in detail in the following sections. Other characteristics, purposes and advantages of the present disclosure can be found in the detailed description and claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0031] FIGs. 1A to 1C show identification of m31A7 mAb from S_{mg} immunized mice and the comparison of heavy chain of spike specific B cell repertoires elicited by S_{mg} and S_{fg} . FIG. 1A shows overview of the single B cell screening platform. Single spike-specific B cells ($CD3^+ CD19^+ spike^+$) from spleen of immunized mice were sorted into 96-well plates by FACS. The IgH and IgL gene transcripts of each single B cell were amplified by RT-PCR. After sequencing, the cDNAs from the variable regions of IgH or IgL genes were subcloned into the expression vectors containing human Ig heavy chain or light chain constant region, respectively. The chimeric monoclonal antibody was produced by Expi 293, and its binding to spike protein was measured by using spike-expressing 293T cells and FACS. FIG. 1B shows the heavy distribution of the B cell repertoire. Less than 5% usage is shown in white. FIG. 1C shows that analysis of heavy chain IgG repertoires of S_{fg} or S_{mg} immunized mice showed highly-

represented IGHV1-9, IGHV1-18 and IGHV2-3 in the S_{mg} group (*p*-values: 0.048, 1.6x10⁻¹¹ and 0.005, respectively, by using Chi-squared test).

[0032] FIGs. 2A to 2B show the binding of m31A7 with SARS-CoV-2 Spike protein. FIG. 2A shows ELISA binding of m31A7 to S1, S2, RBD or the S protein. FIG. 2B shows FACS analysis of m31A7 binding to HEK293T cells expressing S protein of SARS-CoV-2 WT and variants (including D614G, B.1.1.7 [alpha], B.1.351[beta], and B.1.617.2 [delta]).

[0033] FIGs. 3A and 3B show that m31A7 is able to neutralize various variants of SARS-CoV-2 pseudoviruses. FIG. 3A shows neutralization of m31A7 against SARS-CoV-2 WT and variants pseudoviruses. FIG. 3B shows pseudovirus microneutralization of SARS-CoV-2 WT and variants by m31A7 in comparison with previously reported mAb EY6A (grey). Variant species (including D614G, B.1.1.7 [alpha], B.1.351[beta], and B.1.617.2 [delta]) are labeled on top of each panel.

[0034] FIGs. 4A to 4C exhibit that m31A7 shows prophylactic activity against SARS-CoV-2 challenge in vivo. FIG. 4A shows antibody injection and challenge schedule by using K18hACE2 transgenic mice. FIG. 4B shows weight changes after SARS-CoV-2 infection in PBS or m31A7 treated mice. FIG. 4C shows body temperature changes after SARS-CoV-2 infection in PBS or m31A7 treated mice. **P* < 0.05.

[0035] FIGs. 5A to 5D show biophysical properties of m31A7. FIG. 5A shows dissociation constants of m31A7 IgG and Fab to S protein. FIG. 5B shows dissociation constants of m31A7 IgG and Fab to BA.1 S protein. FIG. 5C shows epitope mapping of RBD by HDX-MS showing potential m31A7-binding peptides, measured at 15 sec. FIG. 5D shows volcano plots of the changes in deuterium uptake in RBD upon addition of m31A7 IgG, with the hits (Δ HDX > 5%, *q* value < 0.01) shown in the top right corner and highlighted in red. Unlabeled hits are redundant peptides with 471-482. **P* < 0.05; ***P* < 0.01; ****P* < 0.001.

DETAILED DESCRIPTION OF THE DISCLOSURE

[0036] It is understood that this invention is not limited to the particular materials and methods described herein. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments and is not intended to limit the scope of the present invention, which will be limited only by the appended claims.

[0037] It must be noted that, as used in this specification and the appended claims, the singular forms "a," "an," and "the" include plural referents unless the content clearly dictates otherwise.

[0038] The term "antibody", as used herein, refers to any antigen-binding molecule or molecular complex comprising at least one complementarity determining region (CDR) that specifically binds to or interacts with a particular antigen (SARS-CoV-2-Spike protein). The term "antibody" includes immunoglobulin molecules comprising four polypeptide chains, two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds, as well as

multimers thereof (e.g., IgM). Each heavy chain comprises a heavy chain variable region (abbreviated herein as HCVR or V_H) and a heavy chain constant region. The heavy chain constant region comprises three domains, C_{H1} , C_{H2} and C_{H3} . Each light chain comprises a light chain variable region (abbreviated herein as LCVR or V_L) and a light chain constant region.

5 The light chain constant region comprises one domain (C_{L1}). The V_H and V_L regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDRs), interspersed with regions that are more conserved, termed framework regions (FR). Each V_H and V_L is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. In
10 different embodiments of the disclosure, the FRs of the anti-SARS-CoV-2-Spike protein antibody (or antigen-binding portion thereof) may be identical to the human germline sequences, or may be naturally or artificially modified. An amino acid consensus sequence may be defined based on a side-by-side analysis of two or more CDRs.

[0039] The terms "antigen-binding portion" of an antibody, "antigen-binding fragment" of an
15 antibody, and the like, as used herein, include any naturally occurring, enzymatically obtainable, synthetic, or genetically engineered polypeptide or glycoprotein that specifically binds an antigen to form a complex.

[0040] As used herein, the term "specifically binds" means that an antibody does not cross react to a significant extent with other epitopes.

20 [0041] As used herein, the term "epitope" refers to the site on the antigen to which an antibody binds.

[0042] As used herein, the term "complementarity determining region" (CDR) refers to the non-contiguous antigen combining sites found within the variable region of both heavy and light chain polypeptides. CDRs have been described by Kabat et al., J. Biol. Chem. 252:6609-6616
25 (1977); Kabat et al., U.S. Dept. of Health and Human Services, "Sequences of proteins of immunological interest" (1991); by Chothia et al., J. Mol. Biol. 196:901-917 (1987); and MacCallum et al., J. Mol. Biol. 262:732-745 (1996), where the definitions include overlapping or subsets of amino acid residues when compared against each other.

[0043] As applied to polypeptides, the term "substantial similarity" or "substantially similar"
30 means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 95% sequence identity, even more preferably at least 98% or 99% sequence identity. Preferably, residue positions which are not identical differ by conservative amino acid substitutions. A "conservative amino acid substitution" is one
35 in which an amino acid residue is substituted by another amino acid residue having a side chain (R group) with similar chemical properties (e.g., charge or hydrophobicity). In general, a conservative amino acid substitution will not substantially change the functional properties of a protein. In cases where two or more amino acid sequences differ from each other by

conservative substitutions, the percent sequence identity or degree of similarity may be adjusted upwards to correct for the conservative nature of the substitution. Means for making this adjustment are well-known to those of skill in the art. Examples of groups of amino acids that have side chains with similar chemical properties include (1) aliphatic side chains: glycine, alanine, valine, leucine and isoleucine; (2) aliphatic-hydroxyl side chains: serine and threonine; (3) amide-containing side chains: asparagine and glutamine; (4) aromatic side chains: phenylalanine, tyrosine, and tryptophan; (5) basic side chains: lysine, arginine, and histidine; (6) acidic side chains: aspartate and glutamate, and (7) sulfur-containing side chains are cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, glutamate-aspartate, and asparagine-glutamine. Alternatively, a conservative replacement is any change having a positive value in the PAM250 log-likelihood matrix disclosed in Gonnet et al. (1992) Science 256: 1443-1445, herein incorporated by reference. A "moderately conservative" replacement is any change having a nonnegative value in the PAM250 log-likelihood matrix.

[0044] The term "monoclonal antibody" as used herein is not limited to antibodies produced through hybridoma technology. A monoclonal antibody is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, by any refers to available or known in the art.

[0045] The term "chimeric" antibody as used herein refers to an antibody having variable sequences derived from a non-human immunoglobulin and human immunoglobulin constant regions, typically chosen from a human immunoglobulin template.

[0046] "Humanized" forms of non-human antibodies are chimeric immunoglobulins that contain minimal sequences derived from non-human immunoglobulin. In general, a humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin sequence.

[0047] As used in the present disclosure, the term "fully glycosylated" refers to a state of glycosylation on a CoV spike protein (particular, SARS-CoV-2-Spike protein) wherein all N-glycan sites within the CoV spike protein (particular, SARS-CoV-2-Spike protein) are glycosylated with at least one sugar moiety.

[0048] As used in the present disclosure, the term "therapeutic agent" refers to any compound, substance, drug, drug or active ingredient having a therapeutic or pharmacological effect that is suitable for administration to a mammal, for example a human.

[0049] As used herein, the term "immunoconjugate" refers to an antigen-binding protein, e.g., an antibody or antigen-binding fragment, which is chemically or biologically linked to a radioactive agent, a cytokine, an interferon, a target or reporter moiety, an enzyme, a peptide or

protein or a therapeutic agent. The antigen-binding protein may be linked to the radioactive agent, cytokine, interferon, target or reporter moiety, enzyme, peptide or therapeutic agent at any location along the molecule so long as it is able to bind its target (CoV-S). Examples of immunoconjugates include antibody-drug conjugates and antibody-toxin fusion proteins. In one embodiment of the invention, the agent may be a second, different antibody that binds specifically to CoV-S. The type of therapeutic moiety that may be conjugated to the anti-CoV-S antigen-binding protein (e.g., antibody or fragment) will take into account the condition to be treated and the desired therapeutic effect to be achieved.

[0050] As used herein, the term "vector" is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked.

[0051] The term "genetically engineered" or "genetic engineering" of cells refers to manipulating genes using genetic materials for the change of gene copies and/or gene expression level in the cell. The genetic materials can be in the form of DNA or RNA. The genetic materials can be transferred into cells by various means including viral transduction and non-viral transfection. After being genetically engineered, the expression level of certain genes in the cells can be altered permanently or temporarily.

[0052] The term "coronavirus" or "CoV" refers to any virus of the coronavirus family, including but not limited to SARS-CoV-2, MERS-CoV, and SARS-CoV. SARS-CoV-2 refers to the newly-emerged coronavirus which is rapidly spreading to other areas of the globe. It binds via the viral spike protein to human host cell receptor angiotensin-converting enzyme 2 (ACE2). The spike protein also binds to and is cleaved by TMPRSS2, which activates the spike protein for membrane fusion of the virus.

[0053] The term "CoV-S", also called "S" or "S protein" refers to the spike protein of a coronavirus, and can refer to specific S proteins such as SARS-CoV-2-S, MERS-CoV S, and SARS-CoV S.

[0054] The term "coronavirus infection" or "CoV infection," as used herein, refers to infection with a coronavirus such as SARS-CoV-2, MERS-CoV, or SARS-CoV. The term includes coronavirus respiratory tract infections, often in the lower respiratory tract. Symptoms can include high fever, dry cough, shortness of breath, pneumonia, gastro-intestinal symptoms such as diarrhea, organ failure (kidney failure and renal dysfunction), septic shock, and death in severe cases.

[0055] As used in the present invention, the term "pharmaceutical composition" refers to a mixture containing a therapeutic agent administered to a mammal, for example a human, for preventing, treating, or eliminating a particular disease or pathological condition that the mammal suffers.

[0056] As used herein, the term "therapeutically effective amount" or "efficacious amount" refers to the amount of an antibody that, when administered to a mammal or other subject for treating a disease, is sufficient to affect such treatment for the disease.

[0057] As used herein, the terms "treatment," "treating," and the like, cover any treatment of a disease in a mammal, particularly in a human, and include: (a) preventing the disease from occurring in a subject which may be predisposed to the disease but has not yet been diagnosed as having it; (b) inhibiting the disease, i.e., arresting its development; and (c) relieving the disease, i.e., causing regression of the disease.

[0058] The term "preventing" or "prevention" is recognized in the art, and when used in relation to a condition, it includes administering, prior to onset of the condition, an agent to reduce the frequency or severity of or to delay the onset of symptoms of a medical condition in a subject, relative to a subject which does not receive the agent.

[0059] As interchangeably used herein, the terms "individual," "subject," "host," and "patient," refer to a mammal, including, but not limited to, murines (rats, mice), non-human primates, humans, canines, felines, ungulates (e.g., equines, bovines, ovines, porcines, caprines), etc. Particularly, the subject is vaccinated.

[0060] As used herein, the term "in need of treatment" refers to a judgment made by a caregiver (e.g. physician, nurse, nurse practitioner, or individual in the case of humans; veterinarian in the case of animals, including non-human mammals) that a subject requires or will benefit from treatment. This judgment is made based on a variety of factors that are in the realm of a care giver's expertise, but that includes the knowledge that the subject is ill, or will be ill, as the result of a condition that is treatable by the compounds of the present disclosure.

[0061] As used herein, the term "sample" encompasses a variety of sample types obtained from an individual, subject or patient and can be used in a diagnostic or monitoring assay. The definition encompasses blood and other liquid samples of biological origin, solid tissue samples such as a biopsy specimen or tissue cultures or cells derived therefrom and the progeny thereof.

[0062] A "neutralizing" refers to a process that a molecule (e.g. antibody) inhibits an activity of a coronavirus to any detectable degree, e.g., inhibits the ability of coronavirus to bind to a receptor, to be cleaved by a protease, or to mediate viral entry into a host cell or viral reproduction in a host cell.

[0063] Coronaviruses (CoVs) infect human and animals and cause varieties of diseases, including respiratory, enteric, renal, and neurological diseases. CoV uses its spike glycoprotein (S), a main target for neutralization antibody, to bind its receptor, and mediate membrane fusion and virus entry. The coronavirus spike protein is highly conserved among all human coronaviruses (CoVs) and is involved in receptor recognition, viral attachment, and entry into host cells. Similarly, SARS-CoV-2 S protein is also highly conserved with that of CoVs. When

the S protein binds to the receptor, TM protease serine 2 (TMPRSS2), a type 2 TM serine protease located on the host cell membrane, promotes virus entry into the cell by activating the S protein. Once the virus enters the cell, the viral RNA is released, polyproteins are translated from the RNA genome, and replication and transcription of the viral RNA genome occur via protein cleavage and assembly of the replicase–transcriptase complex. Viral RNA is replicated, and structural proteins are synthesized, assembled, and packaged in the host cell, after which viral particles are released (*Fehr AR, Perlman S. Coronaviruses: an overview of their replication and pathogenesis. Methods Mol Biol. 2015;1282:1–23*).

[0064] The SARS-CoV-2-Spike protein is a 1273 amino acid type I membrane glycoprotein which assembles into trimers that constitute the spikes or peplomers on the surface of the enveloped coronavirus particle. The protein has two essential functions, host receptor binding and membrane fusion, which are attributed to the N-terminal (S1) and C-terminal (S2) halves of the S protein. CoV-S binds to its cognate receptor via a receptor binding domain (RBD) present in the S1 subunit. The amino acid sequence of full-length SARS-CoV-2 spike protein is exemplified by the amino acid sequence of

QCVNLTTRTQLPPAYTNSFTRGVYYPDKVFRSSVLHSTQDLFLPFFSNVTWFHAIHVS
 GTNGTKRFDNPVLPFNDGVYFASTEKSNIIRGWIFGTTLDLSTQSLIVNNATNVVIKV
 CEFQFCNDPFLGVYYHKNNKSWMESEFRVYSSANNCTFEYVSQPFLMDLEGKQGNF
 KNLREFVFKNIDGYFKIYSKHTPINLVRDLPQGFSALEPLVDLPIGINITRFQTLALHRS
 YLTPGDSSSGWTAGAAAYYVGYLQPRTFLKYNENGTITDAVDCALDPLSETKCTLK
 SFTVEKGIYQTSNFRVQPTESIVRFPNITNLCPFGEVFNATRFASVYAWNRRKRISNCVAD
 YSVLYNSASFSTFKCYGVSPTKLNDLCFTNVYADSFVIRGDEVQRQIAPGQTGKIADYN
 YKLPDDFTGCVIAWNSNNLDSKVGGNYNLYRLFRKSNLKPFERDISTEIQAGSTPC
 NGVEGFNCYFPLQSYGFQPTNGVGYQPYRVVLSFELLHAPATVCGPKKSTNLVKNK
 CVNFFNGLTGTGVLTESNKKFLPFQQFGRDIADTTDAVRDPQTLEILDITPCSFGGVS
 VITPGTNTSNQVAVLYQDVNCTEVPVAIHADQLTPTWRVYSTGNSVVFQTRAGCLIGAE
 HVNNSYECDIPIGAGICASYQTQTNSPGSAGSVASQSIIAYTMSLGAENSVAYSNNNSIAI
 PTNFTISVTTEILPVSMTKTSVDCTMYICGDSTECNLLLQYGSFCTQLNRALTGIAVE
 QDKNTQE VFAQVKQIYKTPPIKDFGGFNFSQILPDPSKPSKRSFIEDLLFNK VTLADAG
 FIKQYGDCLGDIAARDLICAQKFNGLTVLPPLLTDEMIAQYTSALLAGTITSGWTFGA
 GAALQIPFAMQMAYRFNGIGVTQNVLYENQKLIANQFNSAIGKIQDSLSTASALGKL
 QDVVNQNAQALNTLVKQLSSNFGAISSVLNDILSRLDPPEAEVQIDRLITGRLQSLQTY
 VTQQLIRAAEIRASANLAATKMSECVLGQSKRVDFCGKGYHLMSFPQSAPHGVVFLH
 VTYVPAQEKNF TAPAICHDKAHFPREGVVFVSNGTHWFVTQRNFYEPQIITDNTFV
 SGNCDVVIGIVNNTVYDPLQPELDSFKEELDKYFKNHTSPDVLGDISGINASVVNIQ
 KEIDRLNEVAKNLNESLIDLQELGKYEQDIRSLVPRGSPGSGYIPEAPRDGQAYVRKD
 GEWVLLSTFLG (SEQ ID NO: 8). Examples of variants of SARS-CoV-2 spike protein include but are not limited to D614G: D614G; B.1.1.7: 69-70 deletion, 144 deletion, N501Y, A570D,

D614G, P681H, T716I, S982A and D1118H; B.1.351: L18F, D80A, D215G, 242-244 deletion, R246I, K417N, E484K, N501Y, D614G and A701V. The term "CoV-S" includes protein variants of CoV spike protein isolated from different CoV isolates as well as recombinant CoV spike protein or a fragment thereof. The term also encompasses CoV spike protein or a fragment thereof coupled to, for example, a histidine tag, mouse or human Fc, or a signal sequence such as ROR1.

[0065] The present disclosure surprisingly found an epitope comprising a part locating in 419 to 433 amino acid residues or 471 to 482 amino acid residues of SEQ ID NO: 8. Accordingly, the present disclosure provides an antibody binding to the epitope disclosed herein.

[0066] The present disclosure develops an antibody or antigen-binding fragment thereof that is specific for an epitope in a CoV spike protein (particularly, SARS-CoV-2-Spike protein).

[0067] Particularly, the antibody or antigen-binding fragment thereof comprises complementarity determining regions (CDRs) of a heavy chain variable region and complementarity determining regions of a light chain variable region, wherein the complementarity determining regions of the heavy chain variable region comprise CDRH1, CDRH2, and CDRH3 regions, and the complementarity determining regions of the light chain variable region comprise CDRL1, CDRL2, and CDRL3 regions, and wherein:

the CDRH1 region comprises the amino acid sequence of SEQ ID NO: 1 or a substantially similar sequence having at least 90%, at least 95%, at least 98% or at least 99% sequence identity; the CDRH2 region comprises the amino acid sequence of SEQ ID NO: 2 or a substantially similar sequence having at least 90%, at least 95%, at least 98% or at least 99% sequence identity; the CDRH3 region comprises the amino acid sequence of SEQ ID NO: 3 or a substantially similar sequence having at least 90%, at least 95%, at least 98% or at least 99% sequence identity; and the CDRL1 region comprises the amino acid sequence of SEQ ID NO: 4 or a substantially similar sequence having at least 90%, at least 95%, at least 98% or at least 99% sequence identity; the CDRL2 region comprises the amino acid sequence of WAS or a substantially similar sequence having at least 90%, at least 95%, at least 98% or at least 99% sequence identity; the CDRL3 region comprises the amino acid sequence of SEQ ID NO: 5 or a substantially similar sequence having at least 90%, at least 95%, at least 98% or at least 99% sequence identity.

[0068] In some embodiments of the disclosure, the antibody or antigen-binding fragment thereof is m31A7, comprising the amino acid sequence of SEQ ID NO: 6 or a substantially similar sequence having at least 90%, at least 95%, at least 98% or at least 99% sequence identity; and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 7 or a substantially similar sequence having at least 90%, at least 95%, at least 98% or at least 99% sequence identity. the CDRH1 region comprises the amino acid sequence of SEQ ID NO: 1 or a substantially similar sequence having at least 90%, at least 95%, at least 98% or at least

99% sequence identity; the CDRH2 region comprises the amino acid sequence of SEQ ID NO: 4 or a substantially similar sequence having at least 90%, at least 95%, at least 98% or at least 99% sequence identity.

5 [0069] The antibody according to the disclosure can be full-length (for example, an IgG1 or IgG4 antibody) or may comprise only an antigen-binding portion (for example, a Fab, F(ab')₂ or scFv fragment), and may be modified to affect functionality as needed.

10 [0070] In some embodiments of the disclosure, antibody or antigen-binding fragment thereof is conjugated with an anti-CoV-S antigen-binding proteins, e.g., antibodies or antigen-binding fragments, conjugated to another moiety, e.g., a therapeutic moiety (an "immunoconjugate"), such as a toxoid or an antiviral agent to treat coronavirus infection. In an embodiment of the invention, an anti-CoV-S antibody or fragment is conjugated to any of the further therapeutic agents set forth herein.

15 [0071] Various techniques known to persons of ordinary skill in the art can be used to determine whether an antibody "specifically binds to one or more amino acids" within a polypeptide or protein. Exemplary techniques include, e.g., routine cross-blocking assay such as that described Antibodies, Harlow and Lane (Cold Spring Harbor Press, Cold Spring Harb., N.Y.), alanine scanning mutational analysis, peptide blots analysis (Reineke, 2004, Methods Mol Biol 248:443-463), and peptide cleavage analysis. In addition, methods such as epitope excision, epitope extraction and chemical modification of antigens can be employed (Tomer, 2000, Protein Science 9:487-496). Another method that can be used to identify the amino acids within a polypeptide with which an antibody specifically binds is hydrogen/deuterium exchange detected by mass spectrometry. In general terms, the hydrogen/deuterium exchange method involves deuterium-labeling the protein of interest, followed by binding the antibody to the deuterium-labeled protein. Next, the protein/antibody complex is transferred to water to allow hydrogen-deuterium exchange to occur at all residues except for the residues protected by the antibody (which remain deuterium-labeled). After dissociation of the antibody, the target protein is subjected to protease cleavage and mass spectrometry analysis, thereby revealing the deuterium-labeled residues which correspond to the specific amino acids with which the antibody interacts. See, e.g., Ehring (1999) Analytical Biochemistry 267(2):252-259; Engen and Smith (2001) Anal. Chem. 73:256A-265A.

25 [0072] One could easily determine whether an antibody specifically binds to the same epitope as, or competes for binding with, a reference anti-SARS-CoV-2-Spike protein antibody by using routine methods known in the art. For example, to determine if a test antibody binds to the same epitope as a reference anti-SARS-CoV-2-Spike protein antibody of the disclosure, the reference antibody is allowed to bind to a SARS-CoV-2-Spike protein. Next, the ability of a test antibody to bind to the SARS-CoV-2-Spike protein molecule is assessed. If the test antibody is able to bind to SARS-CoV-2-Spike protein following saturation binding with the

reference anti-SARS-CoV-2-Spike protein antibody, it can be concluded that the test antibody binds to a different epitope than the reference anti-SARS-CoV-2-Spike protein antibody. On the other hand, if the test antibody is unable to bind to the SARS-CoV-2-Spike protein molecule following saturation binding with the reference anti-SARS-CoV-2-Spike protein antibody, then the test antibody may bind to the same epitope as the epitope bound by the reference anti-SARS-CoV-2-Spike protein antibody of the disclosure. Additional routine experimentation (e.g., peptide mutation and binding analyses) can then be carried out to confirm whether the observed lack of binding of the test antibody is in fact due to binding to the same epitope as the reference antibody or if steric blocking (or another phenomenon) is responsible for the lack of observed binding. Experiments of this sort can be performed using ELISA, RIA, Biacore, flow cytometry or any other quantitative or qualitative antibody-binding assay available in the art. In accordance with certain embodiments of the present disclosure, two antibodies bind to the same (or overlapping) epitope if, e.g., a 1-, 5-, 10-, 20- or 100-fold excess of one antibody inhibits binding of the other by at least 50% but preferably 75%, 90% or even 99% as measured in a competitive binding assay. Alternatively, two antibodies are deemed to bind to the same epitope if essentially all amino acid mutations in the antigen that reduce or eliminate binding of one antibody reduce or eliminate binding of the other. Two antibodies are deemed to have "overlapping epitopes" if only a subset of the amino acid mutations that reduce or eliminate binding of one antibody reduce or eliminate binding of the other.

[0073] The antibody also includes an antigen-binding fragment of a full antibody molecule. An antigen-binding fragment of an antibody may be derived, e.g., from full antibody molecules using any suitable standard techniques such as proteolytic digestion or recombinant genetic engineering techniques involving the manipulation and expression of DNA encoding antibody variable and optionally constant domains. Such DNA is known and/or is readily available from, e.g., commercial sources, DNA libraries (including, e.g., phage-antibody libraries), or can be synthesized. The DNA may be sequenced and manipulated chemically or by using molecular biology techniques, for example, to arrange one or more variable and/or constant domains into a suitable configuration, or to introduce codons, create cysteine residues, modify, add or delete amino acids, etc.

[0074] Non-limiting examples of an antigen-binding fragment include: (i) Fab fragments; (ii) F(ab')₂ fragments; (iii) Fd fragments; (iv) Fv fragments; (v) single-chain Fv (scFv) molecules; (vi) dAb fragments; and (vii) minimal recognition units consisting of the amino acid residues that mimic the hypervariable region of an antibody (e.g., an isolated complementarity determining region (CDR) such as a CDR3 peptide), or a constrained FR3-CDR3-FR4 peptide. Other engineered molecules, such as domain-specific antibodies, single domain antibodies, domain-deleted antibodies, chimeric antibodies, CDR-grafted antibodies, diabodies, triabodies, tetrabodies, minibodies, nanobodies (e.g. monovalent nanobodies, bivalent nanobodies, etc.),

small modular immunopharmaceuticals (SMIPs), and shark variable IgNAR domains, are also encompassed by the expression "antigen-binding fragment," as used herein.

[0075] An antigen-binding fragment of an antibody typically comprises at least one variable domain. The variable domain may be of any size or amino acid composition and will generally comprise at least one CDR which is adjacent to or in frame with one or more framework sequences. In antigen-binding fragments having a V_H domain associated with a V_L domain, the V_H and V_L domains may be situated relative to one another in any suitable arrangement. For example, the variable region may be dimeric and contain V_H - V_H , V_H - V_L or V_L - V_L dimers. Alternatively, the antigen-binding fragment of an antibody may contain a monomeric V_H or V_L domain.

[0076] In certain embodiments, an antigen-binding fragment of an antibody may contain at least one variable domain covalently linked to at least one constant domain. Non-limiting, exemplary configurations of variable and constant domains that may be found within an antigen-binding fragment of an antibody of the present disclosure include: (i) V_H - C_{H1} ; (ii) V_H - C_{H2} ; (iii) V_H - C_{H3} ; (iv) V_H - C_{H1} - C_{H2} ; (v) V_H - C_{H1} - C_{H2} - C_{H3} ; (vi) V_H - C_{H2} - C_{H3} ; (vii) V_H - C_L ; (viii) V_L - C_{H1} ; (ix) V_L - C_{H2} ; (x) V_L - C_{H3} ; (xi) V_L - C_{H1} - C_{H2} ; (xii) V_L - C_{H1} - C_{H2} - C_{H3} ; (xiii) V_L - C_{H2} - C_{H3} ; and (xiv) V_L - C_L . In any configuration of variable and constant domains, including any of the exemplary configurations listed here, the variable and constant domains may be either directly linked to one another or may be linked by a full or partial hinge or linker region. A hinge region may consist of at least 2 (e.g., 5, 10, 15, 20, 40, 60 or more) amino acids which result in a flexible or semi-flexible linkage between adjacent variable and/or constant domains in a single polypeptide molecule. Moreover, an antigen-binding fragment of an antibody of the present disclosure may comprise a homo-dimer or hetero-dimer (or other multimer) of any of the variable and constant domain configurations listed above in non-covalent association with one another and/or with one or more monomeric V_H or V_L domain (e.g., by disulfide bond(s)).

[0077] The anti-SARS-CoV-2-Spike protein antibody disclosed herein may comprise one or more amino acid substitutions, insertions, and/or deletions in the framework and/or CDR regions of the heavy and light chain variable domains as compared to the corresponding germline sequences from which the antibodies were derived. Such mutations can be readily ascertained by comparing the amino acid sequences disclosed herein to germline sequences available from, for example, public antibody sequence databases. The present disclosure includes an antibody, and an antigen-binding fragment thereof, which are derived from any of the amino acid sequences disclosed herein, wherein one or more amino acids within one or more framework and/or CDR regions are mutated to the corresponding residue(s) of the germline sequence from which the antibody was derived, or to the corresponding residue(s) of another mammalian germline sequence, or to a conservative amino acid substitution of the corresponding germline residue(s) (such sequence changes are referred to herein collectively as "germline mutations"). A person of ordinary skill in the art, starting with the heavy and light

chain variable region sequences disclosed herein, could easily produce numerous antibodies and antigen-binding fragments which comprise one or more individual germline mutations or combinations thereof. In certain embodiments, all of the framework and/or CDR residues within the V_H and/or V_L domains are mutated back to the residues found in the original germline sequence from which the antibody was derived. In other embodiments, only certain residues are mutated back to the original germline sequence, e.g., only the mutated residues found within the first 8 amino acids of FR1 or within the last 8 amino acids of FR4, or only the mutated residues found within CDR1, CDR2 or CDR3. In other embodiments, one or more of the framework and/or CDR residue(s) are mutated to the corresponding residue(s) of a different germline sequence (i.e., a germline sequence that is different from the germline sequence from which the antibody was originally derived). Furthermore, the antibodies of the present disclosure may contain any combination of two or more germline mutations within the framework and/or CDR regions, e.g., wherein certain individual residues are mutated to the corresponding residue of a particular germline sequence while certain other residues that differ from the original germline sequence are maintained or are mutated to the corresponding residue of a different germline sequence. Once obtained, antibodies and antigen-binding fragments that contain one or more germline mutations can be easily tested for one or more desired properties such as, improved binding specificity, increased binding affinity, improved or enhanced antagonistic or agonistic biological properties (as the case may be), reduced immunogenicity, etc. Antibodies and antigen-binding fragments obtained in this general manner are encompassed by the present disclosure.

[0078] The present disclosure also includes an anti-SARS-CoV-2-Spike protein antibody comprising variants of any of the V_H , V_L , and/or CDR amino acid sequences disclosed herein having one or more conservative substitutions. For example, the present disclosure includes an anti-EPHA10 antibody having V_H , V_L , and/or CDR amino acid sequences with, e.g., 10 or fewer, 8 or fewer, 6 or fewer, 4 or fewer, etc. conservative amino acid substitutions relative to any of the V_H , V_L , and/or CDR amino acid sequences disclosed herein.

[0079] In some embodiments of the disclosure, the antibody according to the disclosure is a humanized antibody. In order to improve the binding affinity of the humanized antibody according to the disclosure, some amino acid residues in the human framework region are replaced by the corresponding amino acid residues in the species of CDRs; e.g. a rodent.

[0080] The antibodies of the present disclosure may be monospecific, bi-specific, or multispecific. Multispecific antibodies may be specific for different epitopes of one target polypeptide or may contain antigen-binding domains specific for more than one target polypeptide. The anti-SARS-CoV-2-Spike protein antibodies of the present disclosure can be linked to or co-expressed with another functional molecule, e.g., another peptide or protein. For example, an antibody or fragment thereof can be functionally linked (e.g., by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other molecular entities,

such as another antibody or antibody fragment to produce a bi-specific or a multispecific antibody with a second binding specificity. For example, the present disclosure includes bi-specific antibodies wherein one arm of an immunoglobulin is specific for SARS-CoV-2-Spike protein or a fragment thereof, and the other arm of the immunoglobulin is specific for a second therapeutic target or is conjugated to a therapeutic moiety.

[0081] In another aspect, the present disclosure provides a genetically engineered cell expressing the antibody or antigen-binding fragment thereof or containing the vector. The genetically engineered cell may be an immune cell.

[0082] In one preferred embodiments of the disclosure, the antibody or antigen-binding fragment thereof can be produced using any number of expression systems, including prokaryotic and eukaryotic expression systems. In some embodiments, the expression system is a mammalian cell expression, such as a hybridoma, or a CHO cell expression system. Many such systems are widely available from commercial suppliers. In embodiments in which an antibody comprises both a V_H and V_L region, the V_H and V_L regions may be expressed using a single vector, e.g., in a di-cistronic expression unit, or under the control of different promoters. In other embodiments, the V_H and V_L region may be expressed using separate vectors. A V_H or V_L region as described herein may optionally comprise a methionine at the N-terminus.

[0083] The genes encoding the heavy and light chains of an antibody of interest can be cloned from a cell, e.g., the genes encoding a monoclonal antibody can be cloned from a hybridoma and used to produce a recombinant monoclonal antibody. Gene libraries encoding heavy and light chains of monoclonal antibodies can also be made from hybridoma or plasma cells. Random combinations of the heavy and light chain gene products generate a large pool of antibodies with different antigenic specificity (see, e.g., Kuby, Immunology (3.sup.rd ed. 1997)).

[0084] An example of a method for manufacturing the antibody or antigen-binding fragment comprises: (a) introducing into a host cell one or more polynucleotides encoding said antibody or antigen-binding fragment; (b) culturing the host cell under conditions favorable to expression of the one or more polynucleotides; and (c) optionally, isolating the antibody or antigen-binding fragment from the host cell and/or a medium in which the host cell is grown.

[0085] A vector can be used to introduce a polynucleotides encoding the antibody or antigen-binding fragment of the invention to a host cell. In one embodiment, one type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the

host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "recombinant expression vectors" (or simply, "expression vectors"). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" may be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

[0086] The disclosure provides pharmaceutical compositions comprising the antibody or antigen-binding fragment thereof. The pharmaceutical compositions of the disclosure are formulated with suitable diluents, carriers, excipients, and other agents that provide improved transfer, delivery, tolerance, and the like. The compositions may be formulated for specific uses, such as for veterinary uses or pharmaceutical uses in humans. The form of the composition and the excipients, diluents and/or carriers used will depend upon the intended uses of the antibody and, for therapeutic uses, the mode of administration. A multitude of appropriate formulations can be found in the formulary known to all pharmaceutical chemists: Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa. These formulations include, for example, powders, pastes, ointments, jellies, waxes, oils, lipids, lipid (cationic or anionic) containing vesicles (such as LIPOFECTIN.TM., Life Technologies, Carlsbad, Calif.), DNA conjugates, anhydrous absorption pastes, oil-in-water and water-in-oil emulsions, emulsions carbowax (polyethylene glycols of various molecular weights), semi-solid gels, and semi-solid mixtures containing carbowax. See also Powell et al. "Compendium of excipients for parenteral formulations" PDA (1998) J Pharm Sci Technol 52:238-311.

[0087] In some embodiment of the disclosure, the pharmaceutical composition comprises a further therapeutic agent, such as an antiviral agent. The antiviral agent may be an antibody to an S protein of SARS-CoV-2; anti-inflammatory agent; an antibody to a NTD region of a S protein of SARS-CoV-2; an antibody to a HR1 region of a S protein of SARS-CoV-2; an antibody to a RBD region of a S protein of SARS-CoV-2; a SARS-CoV monoclonal antibody; a MERS-CoV monoclonal antibody; a SARS-CoV-2 monoclonal antibody; a peptide; a protease inhibitor; a PIKfyve inhibitor; a TMPRSS2 inhibitor; and a cathepsin inhibitor; a furin inhibitor; an antiviral peptide; an antiviral protein; an antiviral chemical compound. For example, the antiviral agent may be at least one selected from the group consisting of: 1A9; 201; 311mab-31B5; 311mab-32D4; 47D11; 4A8; 4C2; 80R; Apilimod; B38; camostat mesylate; Casirivimab; CR3014; CR3022; D12; E-64D; EK1; EK1C4; H4; HR2P; IBP02; Imdevimab; m336; MERS-27; MERS-4; MI-701; n3088; n3130; P2B-2F6; P2C-1F11; PI8; S230; S309; SARS-CoV-2 S HR2P fragment (aal 168-1203); Tetrandrine; Viracept (nelfinavir mesylate); YM201636; a-1-PDX; favipiravir; IFN- α ; IFN- α 1b; IFN- α 2a; lopinavir-ritonavir; Q-Griffithsin (Q-GRFT); and Griffithsin; oseltamivir; zanamivir; abacavir; zidovudine;

zalcitabine; didanosine; stavudine; efavirenz; indinavir; ritonavir; nelfinavir; amprenavir; ribavirin; Remdesivir; chloroquine; hydroxychloroquine; rIFN-alpha-2a; rIFN-beta-1b; rIFN-gamma; nIFN-alpha; nIFN-beta; nIFN- gamma; IL-2; PD-L1; Anti-PD-L1; a checkpoint inhibitor; an interferon; interferon mixture; recombinant or natural interferon; Alferon; alpha-
5 interferon species; recombinant or natural interferon alpha; recombinant or natural interferon alpha 2a; recombinant or natural interferon beta; recombinant or natural interferon beta 1b; and recombinant or natural interferon gamma. The alpha-interferon species may be a mixture of at least seven species of alpha-interferon produced by human white blood cells, wherein the seven species are: interferon alpha 2; interferon alpha 4; interferon alpha 7; interferon alpha 8;
10 interferon alpha 10; interferon alpha 16; and interferon alpha 17.

[0088] The dose of antibody administered to a patient may vary depending upon the age and the size of the patient, target disease, conditions, route of administration, and the like. The preferred dose is typically calculated according to body weight or body surface area. When an antibody of the present disclosure is used for treating a condition or disease associated with
15 EPHA10 in an adult patient, it may be advantageous to intravenously administer the antibody of the present disclosure. Depending on the severity of the condition, the frequency and the duration of the treatment can be adjusted. Effective dosages and schedules for administering the antibody may be determined empirically; for example, patient progress can be monitored by periodic assessment, and the dose adjusted accordingly. Moreover, interspecies scaling of
20 dosages can be performed using well-known methods in the art (e.g., Mordenti et al., 1991, Pharmaceut. Res. 8:1351).

[0089] Various delivery systems are known and can be used to administer the pharmaceutical composition of the disclosure, e.g., encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the mutant viruses, receptor mediated endocytosis (see,
25 e.g., Wu et al., 1987, J. Biol. Chem. 262:4429-4432). Methods of introduction include, but are not limited to, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The composition may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together
30 with other biologically active agents. Administration can be systemic or local.

[0090] A pharmaceutical composition of the present disclosure can be delivered subcutaneously or intravenously with a vessel or injection device such as standard needle and syringe. In addition, with respect to subcutaneous delivery, a pen delivery device readily has applications in delivering a pharmaceutical composition of the present disclosure. Such a pen
35 delivery device can be reusable or disposable. A reusable pen delivery device generally utilizes a replaceable cartridge that contains a pharmaceutical composition. Once all of the pharmaceutical composition within the cartridge has been administered and the cartridge is empty, the empty cartridge can readily be discarded and replaced with a new cartridge that

contains the pharmaceutical composition. The pen delivery device can then be reused. In a disposable pen delivery device, there is no replaceable cartridge. Rather, the disposable pen delivery device comes prefilled with the pharmaceutical composition held in a reservoir within the device. Once the reservoir is emptied of the pharmaceutical composition, the entire device is discarded.

[0091] In certain situations, the pharmaceutical composition can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, *supra*; Sefton, 1987, *CRC Crit. Ref. Biomed. Eng.* 14:201). In another embodiment, polymeric materials can be used; see, *Medical Applications of Controlled Release*, Langer and Wise (eds.), 1974, CRC Pres., Boca Raton, Fla. In yet another embodiment, a controlled release system can be placed in proximity of the composition's target, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, 1984, in *Medical Applications of Controlled Release*, *supra*, vol. 2, pp. 115-138). Other controlled release systems are discussed in the review by Langer, 1990, *Science* 249:1527-1533.

[0092] The injectable preparations may include dosage forms for intravenous, subcutaneous, intracutaneous and intramuscular injections, drip infusions, etc. These injectable preparations may be prepared by methods publicly known. For example, the injectable preparations may be prepared, e.g., by dissolving, suspending or emulsifying the antibody or its salt described here in a sterile aqueous medium or an oily medium conventionally used for injections. As the aqueous medium for injections, there are, for example, physiological saline, an isotonic solution containing glucose and other auxiliary agents, etc., which may be used in combination with an appropriate solubilizing agent such as an alcohol (e.g., ethanol), a polyalcohol (e.g., propylene glycol, polyethylene glycol), a nonionic surfactant [e.g., polysorbate 80, HCO-50 (polyoxyethylene (50 mol) adduct of hydrogenated castor oil)], etc. As the oily medium, there are employed, e.g., sesame oil, soybean oil, etc., which may be used in combination with a solubilizing agent such as benzyl benzoate, benzyl alcohol, etc. The injection thus prepared is preferably filled in an appropriate ampoule.

[0093] Advantageously, the pharmaceutical compositions for oral or parenteral use described above are prepared into dosage forms in a unit dose suited to fit a dose of the active ingredients. Such dosage forms in a unit dose include, for example, tablets, pills, capsules, injections (ampoules), suppositories, etc.

[0094] The present disclosure provides a method for detecting a coronavirus in a sample comprising contacting the sample with the antibody or antigen-binding fragment thereof.

[0095] The present disclosure provides a method for neutralizing a coronavirus in a subject in need, comprises administering to the subject the antibody or antigen-binding fragment thereof.

[0096] The present disclosure provides a kit for detecting a coronavirus in a sample, wherein the kit comprises the antibody or antigen-binding fragment thereof.

[0097] The anti-SARS-CoV-2-Spike protein antibody of the present disclosure may also be used to detect and/or measure coronavirus, or SARS-CoV-2-Spike protein-expressing cells in a sample, e.g., for diagnostic purposes. For example, an anti-SARS-CoV-2-Spike protein antibody, or fragment thereof, may be used to diagnose a condition or disease characterized by coronavirus infection. Exemplary diagnostic assays for coronavirus may comprise, e.g., contacting a sample, obtained from a patient, with an anti-SARS-CoV-2-Spike protein antibody of the disclosure, wherein the anti-SARS-CoV-2-Spike protein antibody is labeled with a detectable label or reporter molecule. Alternatively, an unlabeled anti-SARS-CoV-2-Spike protein antibody can be used in diagnostic applications in combination with a secondary antibody which is itself detectably labeled. The detectable label or reporter molecule can be a radioisotope, such as ^3H , ^{14}C , ^{32}P , ^{35}S , or ^{125}I ; a fluorescent or chemiluminescent moiety such as fluorescein isothiocyanate, or rhodamine; or an enzyme such as alkaline phosphatase, beta-galactosidase, horseradish peroxidase, or luciferase. Specific exemplary assays that can be used to detect or measure coronavirus in a sample include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence-activated cell sorting (FACS).

[0098] The following examples are provided to aid those skilled in the art in practicing the present disclosure.

EXAMPLES

[0099] **Material and Method**

[0100] **FACS analysis and sorting of spike-specific B cells.** Splenocytes isolated from S_{mg} immunized mice were incubated with 2 $\mu\text{g}/\text{ml}$ S protein at 4 °C for 1 h, followed by washing and incubation with an antibody cocktail against CD19 (clone: 6D5, PE-Cy7-conjugated, Biolegend), CD3 (clone: 17A2, PE-conjugated, Biolegend), and His (clone: J095G46, APC-conjugated, Biolegend), at 4°C for 15 min. Propidium Iodide (Biolegend) was used to exclude dead cells. Live single spike-specific B cells ($\text{CD3}^- \text{CD19}^+$) were sorted into 96-well PCR plates (Applied Biosystems) containing 10 $\mu\text{l}/\text{well}$ catch buffer (10 mM Tris-HCl, pH 8, and 5 U/ μl RNasin (Promega)) by BD FACSAria II. For repertoire analysis, five spleens from S_{mg} or S_{fg} immunized mice were pooled and stained before sorting.

[0101] **Single-B cell screening.** Primers were designed based on a previous publication. The reaction was then performed at 50 °C for 30 min, 95 °C for 15 min followed by 40 cycles at 94 °C for 30 s, 50°C for 30 s, 72°C for 1 min, and final incubation at 72 °C for 10 min. Semi-nested second round PCR was performed using KOD One PCR master mix (TOYOBO) with 1 μl of unpurified first round PCR product at 98 °C for 2 min followed by 45 cycles of 98 °C for 10 s, 55 °C for 10 s, 68 °C for 10 s, and final incubation at 68 °C for 1 min. PCR products were then analyzed on 1.5% agarose gels and sequencing. The Ig V and L genes were identified by searching on IMGT website (http://imgt.org/IMGT_vquest/input). The genes were then amplified from second round PCR product with single gene-specific V and L gene primers

containing restriction sites for cloning into the vectors containing human IgH or IgL expression backbone. The chimeric IgH and IgL expression constructions were co-transfected into Expi293 for antibody production.

[0102] **Binding of antibody with S protein expressing 293T surface.** 293T cells were transfected with pcDNA6/Spike-P2A-eGFP. Transfected cells were selected under 10 µg/ml of blasticidin for 2-3 weeks. Selected cells were then sorted by FACS Aria II to obtain eGFP⁺ expressing cells. These cells were maintained in DMEM containing 10% FBS and 10 µg/ml of blasticidin. 2-3x10⁵ cells were incubated with serially diluted antibody in FACS buffer on ice for 1 h. Then, cells were washed with FACS buffer for 3 times, followed by staining in BV421 mouse anti-human IgG (BD Biosciences, 562581, 1:100) on ice for 20 min and washed with FACS buffer twice. The percentage of positive cells was quantified using FACS Canto II and the data were analyzed with FlowJo. The S protein variants used here are: WH01 spike: original S protein; D614G: D614G; B.1.1.7: 69-70 deletion, 144 deletion, N501Y, A570D, D614G, P681H, T716I, S982A and D1118H; B.1.351: L18F, D80A, D215G, 242-244 deletion, R246I, K417N, E484K, N501Y, D614G and A701V; BA.1.

[0103] **Affinity and avidity determination using Octet (bio-layer interferometry).** Fab fragment was prepared by using Pierce Fab Micro Preparation Kit (ThermoFisher Scientific) according to manufacturer's instructions. Briefly, m31A7 IgG (250 µg) was digested by incubating with immobilized papain resin at 37°C for 8 h. Fab was then purified by protein A column. Purified m31A7 IgG or Fab fragments was loaded at 10 or 6.7 µg/ml kinetics buffer (0.01% endotoxin-free BSA, 0.002% Tween-20, 0.005% NaN₃ in PBS) onto Protein G or FAB2G biosensors (Molecular Devices, ForteBio), respectively. Association and dissociation of S protein (SARS-CoV-2 WH01) by both IgG or Fab was performed in kinetics buffer at indicated concentrations for 5 min and 15 min, respectively. K_D values were calculated using a 1:1 global fit model (Octet).

[0104] **Results:**

[0105] We here aim to isolate spike specific monoclonal antibodies from mice immunized with the monoglycosylated state (S_{mg}) of SARS-CoV-2. We also compared the spike-specific B cell repertoire between S_{mg} and fully glycosylated Spike (S_{fg}) immunized mice. The sorting of S protein-specific B cells from S_{mg} immunized mice led to the identification of a monoclonal antibody (mAb) m31A7 from IGHV1-18 amplified clones (**FIG. 1A**). Heavy chain repertoire analysis of spike-specific B cells isolated from S_{mg} and S_{fg} immunized mice revealed that several heavy chain loci, including IGHV1-9, IGHV1-18 and IGHV2-3, were selectively expanded in S_{mg} immunized mice (**FIGs. 1B and 1C**). This m31A7 mAb interacts with the full-length S protein, S1 and RBD, but not S2 (**FIG. 2A**) and binds to HEK293T cells that express the S protein from different SARS-CoV-2 variants (**FIG. 2B**). In addition, m31A7 was shown to neutralize various pseudovirus variants (WT, D614G, Alpha, Beta, and Delta) at sub-

picomolar IC_{50} which is up to 1000-fold higher than the reported human mAb EY6A (D. Zhou *et al.*, Structural basis for the neutralization of SARS-CoV-2 by an antibody from a convalescent patient. *Nat Struct Mol Biol* **27**, 950-958 (2020)) (**FIGs. 3A and 3B**). A prophylactic study also demonstrated good *in vivo* efficacy of m31A7 in K18hACE2 transgenic mice (**FIG. 4A**) in
5 maintaining both body weight and temperature after challenge with SARS-CoV-2 virus (**FIGs. 4B and 4C**). Bio-layer interferometry (BLI) analysis measured the dissociation constant of m31A7 and its Fab binding to S protein at 34.9 pM and 0.22 nM, respectively (**FIG. 5A**). Binding avidity of full-length m31A7 IgG1 to full-length Omicron BA.1 S protein was measured by bio-layer interferometry and shown in FIG. 5B. Epitope mapping by hydrogen-
10 deuterium exchange mass spectrometry (HDX-MS) showed the binding regions on RBD (**FIGs. 5C and 5D**). The extremely low usage of IGHV1-18 in the S_{fg} B cell repertoire (**FIGs. 1A and 1B**) suggested that S_{fg} may not elicit m31A7 or related antibodies.

[0106] While the present disclosure has been described in conjunction with the specific embodiments set forth, many alternatives thereto and modifications and variations thereof will
15 be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are regarded as falling within the scope of the present disclosure.

CLAIMS

What is claimed is:

1. An antibody or antigen-binding fragment thereof that is specific for an epitope in a spike protein of a CoV; wherein the epitope comprises a part locating in 419 to 433 amino acid residues or 471 to 482 amino acid residues of SEQ ID NO: 8.
2. The antibody or antigen-binding fragment thereof of claim 1, wherein the antibody or antigen-binding fragment thereof comprises complementarity determining regions (CDRs) of a heavy chain variable region and complementarity determining regions of a light chain variable region, wherein the complementarity determining regions of the heavy chain variable region comprise CDRH1, CDRH2, and CDRH3 regions, and the complementarity determining regions of the light chain variable region comprise CDRL1, CDRL2, and CDRL3 regions, and wherein:
the CDRH1 region comprises the amino acid sequence of SEQ ID NO: 1 or a substantially similar sequence thereof; the CDRH2 region comprises the amino acid sequence of SEQ ID NO: 2 or a substantially similar sequence thereof; the CDRH3 region comprises the amino acid sequence of SEQ ID NO: 3 or a substantially similar sequence thereof; and
the CDRL1 region comprises the amino acid sequence of SEQ ID NO: 4 or a substantially similar sequence thereof; the CDRL2 region comprises the amino acid sequence of WAS or a substantially similar sequence thereof; the CDRL3 region comprises the amino acid sequence of SEQ ID NO: 5 or a substantially similar sequence thereof.
3. The antibody or antigen-binding fragment thereof of claim 1, wherein the spike protein is fully glycosylated.
4. The antibody or antigen-binding fragment thereof of claim 2, wherein the antibody or antigen-binding fragment thereof comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 6 or a substantially similar sequence thereof; and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 7 or a substantially similar sequence thereof.
5. The antibody or antigen-binding fragment thereof of claim 1, wherein the antibody is a monoclonal antibody, chimeric antibody, humanized antibody or human antibody.
6. The antibody or antigen-binding fragment thereof of claim 1, wherein the antibody is multispecific.
7. The antibody or antigen-binding fragment thereof of claim 1, wherein the CoV is alpha-CoV, beta-CoV, gamma-CoV, and delta-CoV2.
8. The antibody or antigen-binding fragment thereof of claim 1, wherein the CoV is SARS-CoV, MERS-CoV or SARS-CoV-2.
9. A complex comprising the antibody or antigen-binding fragment thereof of any of

claims 1 to 8 bound to a spike protein of SARS-CoV-2 or fragments thereof.

10. The complex of claim 9, wherein spike protein is fully glycosylated.

11. A vector encoding the antibody or antigen-binding fragment thereof of any of claims 1 to 8.

5 12. A genetically engineered cell expressing the antibody or antigen-binding fragment thereof of any of claims 1 to 8.

13. A genetically engineered cell containing the vector of claim 11.

10 14. A method for manufacturing the antibody or antigen-binding fragment thereof of any of claims 1 to 8, comprising: (a) introducing into a host cell one or more polynucleotides encoding said antibody or antigen-binding fragment; (b) culturing the host cell under conditions favorable to expression of the one or more polynucleotides; and (c) optionally, isolating the antibody or antigen-binding fragment from the host cell and/or a medium in which the host cell is grown.

15 15. A pharmaceutical composition comprising the antibody or antigen-binding fragment thereof of any of claims 1 to 8 and pharmaceutically acceptable carrier and, optionally, a further therapeutic agent.

16. The pharmaceutical composition of claim 15, wherein the therapeutic agent is an antiviral agent.

20 17. The pharmaceutical composition of claim 15, wherein the therapeutic agent is an anti-inflammatory agent or an antibody or antigen-binding fragment thereof that specifically binds to a spike protein of SARS-CoV-2.

18. A vessel or injection device comprising the antibody or antigen-binding fragment thereof of any of claims 1 to 8.

25 19. Use of the antibody or antigen-binding fragment thereof of any of claims 1 to 7 in the manufacture of a medicament for treating or preventing infection with a coronavirus in a subject in need thereof.

20. The use of claim 19, wherein the coronavirus is selected from the group consisting of SARS-CoV-2, SARS-CoV, and MERS-CoV.

30 21. The use of claim 19, wherein the medicament is administered with one or more further therapeutic agents.

22. The use of claim 21, wherein the one or more further therapeutic agents is an antiviral agent.

23. The use of claim 22, wherein the therapeutic agent is an anti-inflammatory agent or an antibody or antigen-binding fragment thereof that specifically binds to a spike protein of SARS-

CoV-2.

24. The use of claim 19, wherein the subject is vaccinated.

25. The use of claim 19, wherein the medicament is for neutralizing a coronavirus in the subject.

5 26. The use of claim 19, wherein the medicament is injected into the body of the subject subcutaneously, intravenously or intramuscularly.

27. A method for detecting a coronavirus in a sample comprising contacting the sample with the antibody or antigen-binding fragment thereof of any of claims 1 to 8.

10 28. A kit for detecting a coronavirus in a sample, wherein the kit comprises the antibody or antigen-binding fragment thereof of any of claims 1 to 8.

29. An epitope comprising a part locating in 419 to 433 amino acid residues or 471 to 482 amino acid residues of SEQ ID NO: 8.

15 30. A method for manufacturing Spike specific monoclonal antibodies or binding fragments thereof immunized with the monoglycosylated state (S_{mg}) of SARS-CoV-2 wherein the method comprising:

(a) introducing into a host cell one or more polynucleotides encoding said antibody or antigen-binding fragment;

(b) culturing the host cell under conditions favorable to expression of the one or more polynucleotides;

20 (c) optionally, isolating the antibody or antigen-binding fragment from the host cell and/or a medium in which the host cell is grown; and

d) wherein the antibody or binding fragment thereof is specific for an epitope in a spike protein of a CoV; wherein the epitope comprises a part locating in 419 to 433 amino acid residues or 471 to 482 amino acid residues of SEQ ID NO: 8.




FIG. 1A

Heavy Chain




FIG. 1B




FIG. 1C




FIG. 2A




FIG. 2B




FIG. 3A




FIG. 3B




FIG. 4A




FIG. 4B




FIG. 4C





FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5C (Cont.)




FIG. 5D

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2022/077481

A. CLASSIFICATION OF SUBJECT MATTER C07K 16/10(2006.01)i; A61P 31/14(2006.01)i; A61K 39/00(2006.01)i According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C07K 16/10(2006.01); A61K 39/215(2006.01); A61K 39/395(2006.01); A61P 11/00(2006.01); A61P 31/14(2006.01) Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Korean utility models and applications for utility models Japanese utility models and applications for utility models Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) eKOMPASS(KIPO internal) & Keywords: antibody, antigen-binding fragment, epitope, spike protein, SARS-CoV-2, coronavirus, vaccine		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2021-045836 A1 (REGENERON PHARMACEUTICALS, INC.) 11 March 2021 (2021-03-11) claim 1; and paragraphs [0006], [0007], [0013], [0015]-[0022], [0024]-[0028], [0030], [0064], [0066]-[0069], [0153]	1,3,5-28,30
A		2,4,29
X	RAHMAN, M. SHAMINUR et al., "Epitope-based chimeric peptide vaccine design against S, M and E proteins of SARS-CoV-2, the etiologic agent of COVID-19 pandemic: an in silico approach", PeerJ, 27 July 2020 (Publication date), DOI: 10.7717/peerj.9572, Internal pages 1-30, Supplemental Information pages 1, 2 abstract; and supplemental information pages 1, 2	29
Y		1,3,5-28,30
Y	WO 2021-180602 A1 (HARBOUR ANTIBODIES BV et al.) 16 September 2021 (2021-09-16) claims 1-37; and pages 1-6, 96-126	1,3,5-28,30
Y	WO 2021-183195 A1 (ABCELLERA BIOLOGICS INC. et al.) 16 September 2021 (2021-09-16) claims 1-57; and paragraphs [0019]-[0038], [0069]-[0094], [0115]-[0120], [0204]-[0209]	1,3,5-28,30
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "D" document cited by the applicant in the international application "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 27 December 2022		Date of mailing of the international search report 28 December 2022
Name and mailing address of the ISA/KR Korean Intellectual Property Office 189 Cheongsa-ro, Seo-gu, Daejeon 35208, Republic of Korea Facsimile No. +82-42-481-8578		Authorized officer HEO, Joo Hyung Telephone No. +82-42-481-5373

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2022/077481

C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	LIAO, HSIN-YU et al., "Impact of glycosylation on SARS-CoV-2 infection and broadly protective vaccine design", BioRxiv, 25 May 2021 (Posted date), DOI: https://doi.org/10.1101/2021.05.25.445523 , Internal pages 1-48 internal pages 1-44	1-30

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2022/077481

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:
 - a. forming part of the international application as filed.
 - b. furnished subsequent to the international filing date for the purposes of international search (Rule 13ter.1(a)),
 accompanied by a statement to the effect that the sequence listing does not go beyond the disclosure in the international application as filed.
2. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, this report has been established to the extent that a meaningful search could be carried out without a WIPO Standard ST.26 compliant sequence listing.
3. Additional comments:

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.

PCT/US2022/077481

Patent document cited in search report			Publication date (day/month/year)	Patent family member(s)			Publication date (day/month/year)
WO	2021-045836	A1	11 March 2021	AU	2020-340881	A1	21 October 2021
				CA	3115553	A1	11 March 2021
				CN	113766928	A	07 December 2021
				EP	3889177	A1	06 October 2021
				JP	2022-160520	A	19 October 2022
				JP	2022-536429	A	17 August 2022
				JP	7116256	B1	09 August 2022
				KR	10-2021-0134300	A	09 November 2021
				TW	202138004	A	16 October 2021
				US	10787501	B1	29 September 2020
				US	10954289	B1	23 March 2021
				US	10975139	B1	13 April 2021
				WO	2021-180602	A1	16 September 2021
GB	202004340	D0	06 May 2020				
GB	202013024	D0	07 October 2020				
GB	202015240	D0	11 November 2020				
GB	202101720	D0	24 March 2021				
TW	202200612	A	01 January 2022				
US	2021-0340225	A1	04 November 2021				
US	2022-0017606	A1	20 January 2022				
WO	2021-180604	A1	16 September 2021				
WO	2021-183195	A1	16 September 2021	AU	2020-435452	A1	16 September 2021
				AU	2021-234242	A1	16 September 2021
				EP	3878862	A2	15 September 2021
				EP	3878862	A3	06 October 2021
				GB	2593023	A	15 September 2021
				KR	10-2021-0113936	A	17 September 2021
				TW	202142563	A	16 November 2021
				TW	202142564	A	16 November 2021
				US	2021-0292393	A1	23 September 2021
				US	2021-0388066	A1	16 December 2021
				WO	2021-183359	A1	16 September 2021