

(12) UK Patent Application (19) GB (11) 2 280 450 (13) A

(43) Date of A Publication 01.02.1995

(21) Application No **9413040.8**

(22) Date of Filing **29.06.1994**

(30) Priority Data

(31) **08085433**

(32) **30.06.1993**

(33) **US**

(71) Applicant(s)

The Procter & Gamble Company

(Incorporated in USA - Ohio)

**One Procter & Gamble Plaza, Cincinnati, Ohio 45202,
United States of America**

(72) Inventor(s)

Alan Edward Sherry

John Downing Curry

(74) Agent and/or Address for Service

Tony Nicholas Gibson

**Newcastle Technical Centre, P O Box Forest
Hall No 2, Newcastle upon Tyne NE12 9TS,
United Kingdom**

(51) INT CL⁶

C11D 17/00

(52) UK CL (Edition N)

**C5D DHZ D114 D120 D123 D124 D127 D132 D134 D141
D142 D143 D144 D145 D146 D147 D148 D149 D150
D151 D152 D153 D154 D155 D156 D157 D158 D159
D160 D161 D162 D163 D164 D165 D166 D167 D168
D169 D170 D171 D172 D179
U1S S1220 S1699**

(56) Documents Cited

GB 2272449 A GB 2179054 A US 4615819 A

(58) Field of Search

UK CL (Edition M) C5D DDX DHD DHZ

INT CL⁵ C11D 17/00

(54) **Detergent gels containing ethoxylated alkyl sulfates and secondary sulfonates**

(57) A detergent composition comprising a gel wholly or predominantly in hexagonal liquid crystal phase, the gel comprising: (a) from 15% to 50% ethoxylated alkyl sulfate surfactant; (b) from 1% to 20% secondary sulfonate surfactant selected from alkylbenzene sulfonates, alkyltoluene sulfonates, paraffin sulfonates, olefin sulfonates, alpha-sulfonated fatty acid alkyl esters, and mixtures thereof; (c) optionally from 0% to 15% other surfactants; (d) from 40% to 80% water; and (e) from 0.2% to 3% magnesium ions; where the ratio of surfactant (a) to surfactant (b) is between 3:2 and 10:1; and where the total amount of surfactants (a)+(b)+(c) is from 25% to 60% by weight of the gel.

GB 2 280 450 A

DETERGENT GELS CONTAINING ETHOXYLATED ALKYL SULFATES AND
SECONDARY SULFONATES

5

FIELD OF THE INVENTION

10

This invention relates to detergent compositions, in particular detergents in the form of gels in hexagonal liquid crystal phase. A preferred embodiment of the invention is dishcare gels.

Detergents in gel form can present many advantages, for example, ease of use and concentrated cleaning ability. Dishcare gels are preferred for use in washing dishes in some parts of the world. The gel product form best lends itself to the "direct application" habit in which persons apply a wet sponge or other cleaning applicator directly onto the dishcare detergent and then onto the dishes; the dishes are then typically washed and rinsed under running water. Additionally, dishcare gels can be stored in inexpensive tubs or similar packages instead of the more complex plastic bottles required for dishwashing liquids.

It is important for detergent gels to have excellent sudsing properties, and to provide good cleaning and other performance benefits. It is also desirable for detergent gels to have an attractive appearance and to be easy to process. Moreover, it is desirable to make such detergent gels from ingredients that are relatively inexpensive so that the detergent gels are affordable to consumers.

Therefore, it is an object of the present invention to provide detergent compositions in the form of gels.

--2--

It is a particular object of the present invention to make detergent gels that have excellent sudsing characteristics, including a large amount of rich, long-lasting suds.

- 5 It is also an object of the present invention to make detergent gels that provide good cleaning and other performance benefits in addition to the excellent sudsing.

It is another object of the present invention to provide detergent gels that have an attractive appearance and that are easy to
10 process.

It is a further object of the present invention to make such gels from relatively inexpensive ingredients so that the gels are affordable to consumers.

- 15 These and other objects of the invention will be described in further detail herein.

SUMMARY OF THE INVENTION

20 The present invention relates to a detergent composition comprising a gel wholly or predominantly in hexagonal liquid crystal phase, said gel comprising:

- (a) from about 15% to about 50% ethoxylated alkyl sulfate surfactant by weight of the gel, wherein the alkyl group of the ethoxylated alkyl sulfate surfactant has an average from about 8 to about 20 carbon atoms, and wherein the ethoxylated alkyl
25 sulfate surfactant has an average degree of ethoxylation from about 0.5 to about 15;
- (b) from about 1% to about 20%, by weight of the gel, secondary sulfonate surfactant selected from the group consisting of alkylbenzene sulfonates, alkyltoluene sulfonates, paraffin sulfonates, olefin sulfonates, alpha-sulfonated fatty acid
30 alkyl esters, and mixtures thereof;
- (c) optionally from 0% to about 15% other surfactants by weight of the gel;

- (d) from about 40% to about 80% water by weight of the gel; and
- (e) from about 0.2% to about 3% magnesium ions by weight of the gel;

wherein the weight ratio of surfactant (a) to surfactant (b) is
5 between about 3:2 and about 10:1, and wherein the total amount of
surfactants (a)+(b)+(c) is from about 25% to about 60% by weight of
the gel.

DETAILED DESCRIPTION OF THE INVENTION

10 Gels in hexagonal liquid crystal phase are particularly suitable for
use as detergent gels, because they have desirable viscosity and
consistency, are stable, and have a good appearance. Many
surfactants in water undergo phase changes from micellar to
hexagonal to lamellar phase as a function of surfactant
concentration at a given temperature and pressure. Ethoxylated
15 alkyl sulfate surfactants in water, for example, generally display
hexagonal phase regions at from about 25-30% concentration to about
60-65% concentration by weight at ambient temperature and pressure.
Unfortunately, detergent gels made from ethoxylated alkyl sulfate
surfactants alone do not provide very good cleaning or sudsing
properties.

20 Secondary sulfonate surfactants such as alkylbenzene sulfonates,
alkyltoluene sulfonates, paraffin sulfonates, olefin sulfonates, and
alpha-sulfonated fatty acid alkyl esters do not have well-defined
hexagonal phase regions in water at ambient temperature and
pressure. Instead, these surfactants are predominantly in the
25 lamellar phase or in a lamellar phase emulsion even at low
surfactant concentrations. It has now been discovered that by
combining ethoxylated alkyl sulfate surfactants at a concentration
where they are present in a micellar phase near the hexagonal phase
boundary (i.e., about 15-30% surfactant concentration by weight),
with secondary sulfonate surfactants present that display
30 predominantly lamellar phase behavior, exist as lamellar phase
emulsions, or as crystals in water, hexagonal phase gels are formed.
While not intending to be limited by theory, it is believed that

such hexagonal phase gels result from a "phase averaging effect", i.e., a phase averaging of the micellar phase and other phase compositions to form a hexagonal phase composition.

- 5 Importantly, it has also been found that these hexagonal phase detergent gels, containing such secondary sulfonate surfactants in combination with ethoxylated alkyl sulfate surfactants, provide excellent sudsing and good cleaning when the gels also contain small amounts of magnesium ions. Moreover, the secondary sulfonate surfactants are relatively inexpensive so that the detergent gels
10 are readily affordable to consumers. The detergent gel compositions are easy to manufacture because processing of the compositions usually takes place at elevated temperature where the surfactant mixture is liquid, and not in the hexagonal gel phase; upon cooling to room temperature, the compositions enter the hexagonal phase. The detergent gels of this invention do not require additives such
15 as urea to force the compositions into the hexagonal phase. Instead, the hexagonal phase results from mixing of the surfactants in the proportions disclosed herein.

- A detergent gel composition according to the present invention comprises from about 15% to about 50% ethoxylated alkyl sulfate
20 surfactant by weight of the detergent gel, preferably from about 20% to about 45% ethoxylated alkyl sulfate. An ethoxylated alkyl sulfate surfactant, AE_xS, is one having, on average, "x" degree of ethoxylation. The ethoxylated alkyl sulfate surfactant for use in the present invention has an average degree of ethoxylation from about 0.5 to about 15, and preferably from about 1 to about 6.5.
25 The alkyl group of the ethoxylated alkyl sulfate surfactant can have an average from about 8 to about 20 carbon atoms, preferably from about 8 to about 15 carbon atoms, and most preferably from about 12 to about 15 carbon atoms. The alkyl groups are preferably linear, but they can also be branched.

- 30 Blends of different ethoxylated alkyl sulfate surfactants can be used, for example a blend of two surfactants having different degrees of ethoxylation. In general, highly ethoxylated surfactants

(e.g., ethoxylation of 3 or more) provide more mildness to skin, while mono- and di-ethoxylated surfactants contribute more to cleaning ability. As a result, it may be desirable to use a blend of AE₁S and AE₃S or similar blends to provide the optimum
5 combination of cleaning and mildness. Variation in degrees of ethoxylation of the surfactants provides broad formulation flexibility.

The cation of the ethoxylated alkyl sulfate surfactant can be sodium, potassium, lithium, calcium, magnesium, ethylene diamine,
10 ammonium, aluminum, zinc, or lower alkanol ammonium ions, and other cations which are known in the detergent field to be useful in surfactants. As will be discussed below, most preferred are magnesium cations. The preferred magnesium ethoxylated alkyl sulfate surfactant can be either introduced as a raw material, or it can be generated in situ through counterion exchange with Mg⁺⁺ salts
15 (this can also be done for the surfactants with less preferred cations).

Preferred ethoxylated alkyl sulfate surfactants according to the present invention include those where the alkyl group is derived from coconut or palm base, such as mid-cut coconut (C₁₂₋₁₄) or
20 broad-cut coconut (C₁₂₋₁₈). Surfactants of the C₁₂₋₁₄ type are available commercially from Akzo Chemicals, 516 Duren, Germany, under the tradename ELFAN NS 243 S Mg conc. (Mg⁺⁺ cation, alkyl group having an average chain length of C₁₂₋₁₄, average degree of ethoxylation of 3).

25 Synthetic ethoxylated alkyl sulfate surfactants (derived from synthetic alcohols) such as those containing C₁₂₋₁₃ or C₁₂₋₁₅ alkyl groups are also preferred. Such synthetic surfactants are commercially available from South Pearl Corp., Ponce, Puerto Rico, 00731 and other suppliers. Specific examples of preferred surfactants are Mg(C₁₂₋₁₄AE₂S)₂, Mg(C₁₂₋₁₅AE₃S)₂, Mg(C₁₂₋₁₃AE₁S)₂,
30 and their counterparts having other degrees of ethoxylation. Other suitable surfactants include, but are not limited to, ethoxylated

alkyl sulfate surfactants where the alkyl group is lauryl (C₁₂) or myristyl (C₁₄).

5 The ethoxylated alkyl sulfate surfactant used in the detergent gel is preferably a high active surfactant. By "high active" surfactant is meant the surfactant has at least about 60% active content, preferably at least about 65%. At above 60-65% activity, ethoxylated alkyl sulfate surfactants are easily processable lamellar phase materials. Most preferred are high active
10 surfactants that contain little or no alcohols, glycols, inorganic salts or hydrotopes. Ethoxylated alkyl sulfate surfactant stocks which are not "high active" will necessarily contain additives such as alcohols and glycols, or hydrotropes such as toluene, xylene and cumene sulfonates. The ethoxylated alkyl sulfate surfactant stocks of this invention should contain little or no short chain alcohols and glycols (C₆ or lower), preferably less than about 10% by weight,
15 most preferably less than about 5% by weight. Additionally, they should contain little or no hydrotropes, preferably less than about 5% by weight, most preferably less than about 2% by weight. Without limiting the scope of this invention, it has been found that stable hexagonal phase gels can be made in the presence of small quantities of alcohols, glycols and/or hydrotropes, but these gels usually
20 require higher surfactant concentrations. The gels do not contain additives such as urea, thiourea, methyl urea or ethyl urea to force the surfactant system into the hexagonal phase.

The detergent gel of the present invention also comprises from about 1% to about 20%, by weight of the gel, secondary sulfonate
25 surfactants selected from the group consisting of alkylbenzene sulfonates, alkyltoluene sulfonates, paraffin sulfonates, olefin sulfonates, alpha-sulfonated fatty acid alkyl esters, and mixtures thereof. Preferably the detergent gel comprises from about 5% to about 15% secondary sulfonate surfactants by weight of the gel. Preferred secondary sulfonate surfactants are selected from the
30 group consisting of alkylbenzene sulfonates, paraffin sulfonates, and mixtures thereof. Most preferred are alkylbenzene sulfonates.

Suitable alkylbenzene or alkyltoluene sulfonates include the alkali metal (lithium, sodium, and/or potassium), alkaline earth (preferably magnesium), ammonium and/or alkanolammonium salts of straight or branched chain alkylbenzene or alkyltoluene sulfonic acids in which the alkyl group contains from about 9 to about 15 carbon atoms. Alkylbenzene sulfonic acids useful as precursors for these surfactants include decyl benzene sulfonic acid, undecyl benzene sulfonic acid, dodecyl benzene sulfonic acid, tridecyl benzene sulfonic acid, tetrapropylene benzene sulfonic acid and mixtures thereof. Preferred sulfonic acids as precursors of the alkylbenzene sulfonates useful for compositions herein are those in which the alkyl chain is linear and averages about 11 to about 13 carbon atoms in length, most preferably from about 11.3 to about 12.3 carbon atoms in length. Examples of commercially available alkyl benzene sulfonic acids useful in the present invention include Conoco SA 515 and SA 597 marketed by the Continental Oil Company and Calsoft LAS 99 marketed by the Pilot Chemical Company.

Also suitable for use as the secondary sulfonate surfactants are paraffin sulfonates having from about 8 to about 22 carbon atoms, preferably from about 12 to about 16 carbon atoms, in the alkyl moiety. A suitable commercially available paraffin sulfonate useful in the present invention is Hostapur SAS 60 marketed by the Hoechst Celanese Corp.

The secondary sulfonate surfactants can also be olefin sulfonates, which are compounds produced by the sulfonation of alpha-olefin by means of uncomplexed sulfur trioxide followed by neutralization of the acid reaction mixture under conditions such that sulfones formed in the reaction are hydrolyzed to give corresponding hydroxyalkanesulfonates. The alpha-olefins from which the olefin sulfonates are derived are mono-olefins having from about 12 to about 24 carbon atoms, preferably from about 14 to about 16 carbon atoms. Preferably, they are straight chain olefins. Examples of suitable 1-olefins include 1-dodecene, 1-tetradecene, 1-hexadecene, 1-octadecene, 1-eicosene and 1-tetracosene.

The secondary sulfonate surfactants can also be alpha-sulfonated fatty acid alkyl ester surfactants having the general formula:



wherein R¹ is straight or branched alkyl having from about 8 to about 20 carbon atoms; R² is straight or branched alkyl having from about 1 to about 6 carbon atoms; M represents a counter-ion such as sodium, potassium, magnesium, ammonium and alkanolammonium, and i is either 1 or 2 depending on whether the counter-ion is mono- or divalent. Preferred is an ester salt wherein R¹ is C₁₀₋₁₆ alkyl, R² is methyl, and M is Na or K.

The weight ratio of the ethoxylated alkyl sulfate surfactant to the secondary sulfonate surfactant should be between about 3:2 and about 10:1, preferably between about 2:1 and 5:1. Hexagonal phase gels are not formed when the ratio of ethoxylated alkyl sulfate surfactant to secondary sulfonate surfactant is less than 3:2, while cleaning and sudsing properties of the hexagonal phase gels are poor when the ratio is greater than 10:1.

Along with the ethoxylated alkyl sulfate and secondary sulfonate surfactants, the present detergent gel composition can also optionally contain from 0% to about 15% other surfactants by weight of the gel. The other surfactants can be anionic, cationic, nonionic, zwitterionic, ampholytic or amphoteric surfactants known to persons skilled in the art. Preferably not more than about 5% by weight cationic surfactant is used. Preferred optional surfactants are nonionic. Nonionic surfactants useful in the detergent gel of this invention include ethoxylated fatty alcohols, the fatty acyl ethanolamides, alkyl phenols, polypropylene oxides, polyethylene oxides, copolymers of polypropylene oxide and polyethylene oxide, sorbitan esters, and the like.

- Amine oxides and/or betaines can be optional surfactants, each at a level of up to about 0.5% by weight of the gel. Higher levels of amine oxides and/or betaines weaken gel viscosity and complicate processing. Amine oxides are described in U.S. Patent 4,316,824 to Pancheri, which is incorporated herein by reference. The Procter & Gamble Company, Cincinnati, Ohio, manufactures suitable amine oxides such as C₁₀-16 (predominantly C₁₂) alkyl dimethyl amine oxides. The C₁₂, C₁₄, C₁₆, C₁₄-16, and C₁₆-18 alkyl dimethyl amine oxides are available commercially from Stepan Chemical Company under the tradename Ammonyx. Betaines are disclosed in U.S. Patent Nos. 3,950,417; 4,137,191; 4,375,421; and 4,555,360; all of which are incorporated herein by reference. Examples of preferred betaines are cetyl dimethyl betaine, dodecyl dimethyl betaine, coco amido propyl betaine, dodecyl amidopropyldimethyl betaine, and dodecyldimethylammonium hexanoate.
- 15 The total amount of surfactants in the present detergent gel composition should be from about 25% to about 60% by weight of the gel. By "total amount of surfactants" is meant the total amount of ethoxylated alkyl sulfate surfactant, secondary sulfonate surfactant, and any optional surfactant used in the detergent gel. At least about 25% total surfactant is needed to make a suitably thickened gel. Above about 60% total surfactant, processing becomes more difficult and the mixture may not exist in the hexagonal phase. A detergent gel composition according to the present invention preferably contains from about 25% to about 50% total surfactant, and most preferably from about 30% to about 40%.
- 20
- 25 A detergent gel composition according to the invention also contains from about 40% to about 80% water by weight of the gel, preferably from about 45% to about 70%, and most preferably from about 50% to about 65%.
- 30 A detergent gel according to the present invention contains from about 0.2% to about 3% magnesium ions by weight of the gel. At least about 0.2% magnesium ions are needed to ensure satisfactory cleaning performance of the detergent gel, and the magnesium ions

also enhance sudsing performance. If the product contains more than about 3% magnesium ions it is difficult to obtain a gel since additional magnesium ions would need to be introduced as magnesium salts which are electrolytes that weaken the gel structure. The
5 magnesium ions are preferably introduced by use of magnesium neutralized ethoxylated alkyl sulfate surfactants, secondary sulfonate surfactants, and/or optional surfactants. Less preferably for the above-mentioned reason, magnesium ions can also be added in the form of salts, for example, magnesium sulfate, magnesium formate
10 or magnesium chloride; magnesium salts may also be introduced as impurities in the surfactants used (for example, magnesium sulfate impurity is usually present in a magnesium ethoxylated alkyl sulfate surfactant as a result of the synthesis process).

15 The detergent gel compositions of this invention can be clear or opaque as desired. The gels can contain thickening or suspending agents, fillers or abrasives; these materials usually opacify the gel. The detergent gel composition of the invention can contain, if desired, other additives known to persons skilled in the art for use
20 in detergents, for example, bleaching agents, perfumes, builders, amino acids, dyes, antitarnishing agents, antimicrobial agents, suds enhancers, coloring agents, and the like. The amounts of these additives should be limited to avoid interfering with the viscosity of the gel.

25 The detergent compositions of the present invention are in the form of gels which have a viscosity between about 500,000 centipoise and about 6,000,000 centipoise. The viscosity of the gels can be varied within this range depending on the properties desired in the final
30 product. Viscosity measurements of the gels of this invention are taken by means of an Exact Viscometer HAAKE RV20 ROTOVISCO using a cone and plate geometry with Cone PK1, 1°, M=30.2. The viscosity is measured on a 0.5 gram sample of the gel sandwiched between the cone and the instrument's plate, using a shear rate gradient of 0 to 3
35 seconds⁻¹, at a temperature of 23°C (73.4°F). The recorded viscosity corresponds to the highest viscosity reading obtained on the instrument when a sweep time of two minutes is used. Each

sample is measured seven times in this manner and the measurements averaged.

5 A detergent gel according to this invention is wholly or predominantly in hexagonal liquid crystal phase. By "predominantly" is meant greater than about 50%. The liquid crystal phase of the detergent gel can be determined by polarizing light microscope studies, use of X-ray diffraction or other various spectroscopic techniques known to persons skilled in the art. The hexagonal liquid crystal phase is intermediate in rigidity between the
10 lamellar and cubic liquid crystal phases. The hexagonal liquid crystal phase is further described at column 3, lines 12-31, of U.S. Patent 4,615,819 to Leng et al., issued October 7, 1986, this patent being incorporated by reference herein.

15 The detergent gel compositions of this invention are preferably dishwashing detergents for use with the "direct application" habit discussed above, or for use with batch dishwashing typical of liquid detergents; general purpose industrial and household cleaners for use in cleaning hard surfaces such as metal, glass, ceramic, tile and linoleum; concentrated laundry detergents; hand cleaners; shampoos; or other detergent compositions known in the detergent
20 field. Laundry detergents according to the invention will contain surfactant, builder, and typically components such as brighteners, bleach, enzymes, chelating agents, and suds suppressors. General purpose hard surface cleaners will contain surfactants, builder, and sometimes abrasive and solvent.

25 The detergent gels can be prepared in any suitable manner, for instance by simply mixing together the components. The gels are easily processable as fluid mixtures at temperatures of 150°F (65.6°C) to 210°F (98.9°C), preferably about 170°F (76.7°C) to 200°F (93.3°C). The order of mixing of the components is not critical, but a preferred order of addition is to add water first, then
30 ethoxylated alkyl sulfate surfactant and magnesium ions, then nonionic surfactant (if any), and lastly secondary sulfonate surfactant. Abrasives can be added, if desired, at any time during

the processing. Upon cooling, the compositions become viscous and set up as hexagonal phase gels.

The following nonlimiting examples are performed using water at a
5 6-12 grains per gallon hardness. All percentages herein are by weight unless otherwise defined.

EXAMPLE 1

A detergent gel most preferred for its excellent sudsing and good
10 cleaning properties is made as follows. To 20.00 grams of $\text{Mg}(\text{LAS})_2$ (50% active made by Hoechst, Venezuela), 41.02 grams of water and 6.00 grams of silica (100% active Zeodent 119 made by J. M. Huber, USA) are added at about 74°F (23.3°C). The mixture is then well mixed and heated to 180°F (82.2°C). 32.98 grams $\text{Mg}(\text{C}_{12-13}\text{AE}_2\text{S})_2$ (66.7% active made by South Pearl Corp., Puerto Rico, USA) are added
15 and the mixture stirred for 4 hours. The final product is a viscous opaque fluid which quickly forms a gel upon cooling to room temperature. X-ray diffraction identifies the gel phase as predominantly hexagonal. The composition of the gel is 22% $\text{Mg}(\text{AE}_2\text{S})_2$, 10% $\text{Mg}(\text{LAS})_2$, 6% precipitated silica, and 62% water. The gel contains 1.0% magnesium ions. The viscosity of the gel is
20 3,200,000 cps. The IFT (interfacial tension) of the detergent gel is 1.0 dynes/cm.

IFT is measured by use of a Model 500 Spinning Drop Interfacial Tensiometer with a 2mm I.D. x 95mm long capillary tube with cap
(both manufactured by the University of Texas, Austin, Texas). The
25 "soil" used for the IFT measurement is 99.8% Puritan® Canola Oil (Procter & Gamble, Cincinnati, Ohio) and 0.2% oleic acid. The process is as follows:

1. Prepare 100 to 150 grams of a 6% product solution of the detergent gel in distilled water ("soft" water, 3gpg hardness or less).
- 30 2. Let the solution equilibrate to 25°C.
3. Flush the capillary tube three times with product solution using a 10cc disposable syringe.

4. Invert the capillary tube to an approximate 30° angle.
5. Flush a 0.005ml microsyringe with the "soil" described above.
6. Inject the soil:
Add enough soil to elongate it four times its width (see #12).
- 5 7. Put the cap on the capillary tube making sure no air bubbles enter the tube.
8. Dry the tube, especially around the cap area.
9. Making sure the oil droplet is in the center of the tube, insert the tube into the IFT machine and tighten.
10. Turn on the power and then the strobe light.
- 10 11. Locate and center the oil droplet and tighten the eyepiece accordingly.
12. Increase the speed until the oil droplet elongates to four times its width. (The speed cannot go below 7.00 ms./rev.)
13. Start the timer.
14. Measure and record the readings for the top and bottom of the oil droplet as well as the initial speed and the speed after 2, 4, 6, and 10 minutes.
- 15 15. The average IFT is the average of the calculated IFT values at 0, 2, 4, 6 and 10 minutes. IFT is calculated by the equation:
$$IFT = 52,100 \times [(top\ reading - bottom\ reading)^3] \div (speed^2).$$

20

EXAMPLE 2

70.0 grams of NaC₁₂AE₃S solution (28.5% sodium neutralized alkyl ethoxy sulfate with an average of 3 moles of ethoxy groups per mole of surfactant, Steol 4N made by Stepan Chemical Company, USA) and 0.20 grams of MgSO₄ (99% active) are stirred together at 150°F (65.6°C) using a Labmaster 1500 MSV 1500 U mixer. Once all of the MgSO₄ is dissolved, 0.10 grams of a 1% solution of blue dye, 0.50 grams of perfume, and 9.20 grams of water are added and the mixture is vigorously stirred. The temperature is raised to 170°F (76.7°C) and 20.0 grams of Mg(C_{11.8}LAS)₂ (50% active, linear alkylbenzene sulfonate made by Hoechst Corp., Venezuela) are added. The resulting mixture is stirred for 2 hours at 170°F (76.7°C) to achieve homogeneity. The final product is a viscous liquid which forms a gel upon cooling to room temperature. The gel is identified

30

as predominantly hexagonal phase by X-ray diffraction. The composition of the gel is 20% NaAE₃S, 10% Mg(LAS)₂, 0.2% MgSO₄, 69.3% water, 0.5% perfume and 0.001% dye. The gel contains 0.4% magnesium ions. The viscosity of the gel is 1,700,000 cps.

5

EXAMPLE 3

To 49.53 grams of water (8 gpg hardness), 13.33 grams of paraffin sulfonate (60% active, Hostapur SAS 60, sodium neutralized secondary alkane sulfonate made by Hoechst, USA) and 37.14 grams of
10 Mg(C₁₂₋₁₄AE₃S)₂ (70% active magnesium neutralized C₁₂₋₁₄ triethoxylated alkyl sulfate surfactant made by Akzo Chemical Co., Germany) are added and mixed together at 190°F (87.8°C). The product is continuously stirred until a homogeneous solution is obtained. The final product is a clear, water white liquid which forms a gel upon cooling to room temperature. The composition of
15 the gel is 26% Mg(C₁₂₋₁₄AE₃S)₂, 8% paraffin sulfonate, and 64% water. The gel contains 0.75% magnesium ions. The viscosity of the gel is 3,000,000 cps. The gel is identified as predominantly in the hexagonal phase by X-ray diffraction.

EXAMPLE 4

20

To 49.24 grams of water, 13.33 grams of paraffin sulfonate (60% active Hostapur SAS 60 made by Hoechst USA), 31.43 grams of Mg(AE₃S)₂ (70% active, made by Akzo Chemical, Germany), and 6% precipitated silica (100% active Zeodent 119 made by J. M. Huber, USA) are added sequentially and mixed at 190°F (87.8°C). The mixing
25 is continued until the Mg(AE₃S)₂ is completely dissolved and the silica is well dispersed in the mixture. The final product is a white fluid composition that forms an opaque gel upon cooling to room temperature. X-ray diffraction identifies the phase of the gel as predominantly hexagonal. The composition of the gel is 22% Mg(AE₃S)₂, 8% paraffin sulfonate, 6% silica and 64% water. The gel
30 contains 0.6% magnesium ions. The viscosity of the gel is 2,600,000 cps.

The detergent gel products of Examples 1-4 are tested for their sudsing properties by the use of an apparatus consisting of 8 tubes (cylinders) of length 30 cm. and diameter 10 cm. fixed side by side, and rotatable at a speed of 24 rpms about a central axis. Each tube
5 can be charged with 500 ml. of product solution. In short, 0.2% solutions of the products of Examples 1-4 and a control product are inserted into five of the tubes, the tubes are rotated 20 times, and the height of the suds is measured in each of the tubes. One ml. of test soil is injected into each of the tubes containing the product solutions, the tubes are rotated 20 more times, and the height of
10 the suds is again measured. This is repeated with further additions of 1 ml. increments of test soil until the suds are diminished. Following is the procedure in more detail:

1. Prepare test soil: Melt 100 grams of test soil in a water bath using low heat. The test soil is composed of 12.7% Crisco® oil, 27.8% Crisco® shortening, 7.6% lard, 51.7% beef suet,
15 0.14% oleic acid, 0.04% palmitic acid and 0.02% stearic acid.
2. Prepare product solutions: Prepare 500 ml. product solutions at 0.2% concentration using tap water at ambient temperature. The products of Examples 1-4 and one control product are tested. The control product is a commercial detergent gel product containing, as the surfactant, about 36% linear
20 alkylbenzene sulfonate surfactant and about 1% ethoxylated alkyl sulfate surfactant. The control product is known to have very good sudsing properties.
3. Make sure the tubes of the apparatus are clean by rinsing them with distilled water.
4. Pour product solutions into the tubes and replace tube lids.
- 25 5. Secure the tubes in the apparatus.
6. Rotate the tubes 20 times at 24 rpm's.
7. Measure the suds height in each of the tubes.
8. Inject 1 ml of melted test soil into each tube.
9. Repeat 6-8 until the suds are diminished.

**SUDS HEIGHT IN MILLIMETERS AFTER
20 TUBE ROTATIONS PER SOIL LOADING**

5	<u>PRODUCTS</u>	<u>ML. OF SOIL ADDED</u>						
		<u>0</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>6</u>
	Control	49	56	51	55	54	38	24
	Example 1	64	71	64	82	89	59	60
	Example 2	60	65	63	76	77	60	66
	Example 3	59	67	58	67	50	29	21
	Example 4	60	64	58	61	44	22	17

10

		<u>ML. OF SOIL ADDED</u>						<u>TOTAL</u>	<u>SUDS INDEX</u>
		<u>7</u>	<u>8</u>	<u>9</u>	<u>10</u>	<u>11</u>	<u>12</u>		
	Control	11	8	5				351	100.0
	Example 1	39	26	13	5			572	163.0
15	Example 2	45	33	19	11	8	.6	589	167.8
	Example 3	15	10	8	7	7		398	113.4
	Example 4	10	8	6				350	99.7

The "Suds Index" is the total suds of each test product divided by the total suds of the control product. It is seen from the results that the detergent gel products of Examples 1-4 provide excellent sudsing, particularly the detergents of Examples 1 and 2.

EXAMPLE 5

This example illustrates that nonionic surfactants can optionally be used in the present detergent gels. 30.0 grams of $Mg(C_{11.8}LAS)_2$ (50% active, made by Hoechst, Venezuela), 47.12 grams of $NaC_{12-14}AE_2S$ (70% active, Genapol ZRO-V, made by Hoechst, Venezuela), 1.87 grams of C_{10-16} dimethyl amine oxide (32% active, made by Procter & Gamble, USA), and 15.00 grams of alkyl ethoxylated alcohol (C_{12-15} , average of three ethoxylate groups per molecule, Neodol 25-3S made by Shell, USA) are mixed at 185°F (85°C). 122.95 grams of water are added, and the solution stirred continuously at 185°F (85°C). After the solution is made completely homogeneous,

80.76 grams of NaC₁₂₋₁₄AE₁S (26% active Genapol ZRO 1-V made by Hoechst, Venezuela) are added at the same temperature. The final product is a viscous liquid which quickly solidifies into a gel upon cooling to room temperature. The composition of the gel is 7%
 5 NaAE₁S, 11% NaAE₂S, 5% Mg(LAS)₂, 5% Neodol, and 0.2% C₁₀₋₁₆ amine oxide. The gel contains 0.2% magnesium ions.

EXAMPLE 6

The following table demonstrates that the level of amine oxide
 10 should be limited to avoid weakening gel strength of Mg(LAS)₂/AE_xS formulations according to the invention.

	<u>(1)</u>	<u>(2)</u>	<u>(3)</u>	<u>(4)</u>
Mg(C ₁₂₋₁₄ AE ₃ S) ₂	24%	24%	24%	24%
Mg(C _{11.8} LAS) ₂	8%	8%	8%	8%
C ₁₀₋₁₆ Amine Oxide	0%	1%	2%	4%
15 Water	68%	67%	66%	64%
Gel Viscosity	1.0	0.8	0.6	Liquid
	million	million	million	Product
	centipoise	centipoise	centipoise	

EXAMPLE 7

20

The following compositions are additional hexagonal phase gels that further illustrate the invention:

- a) 13.0% NaC₁₂₋₁₃AE₂S, 17% Mg(C_{11.8}LAS)₂, 2.2% NaAE₁S and 67.8% water
- 25 b) 30.0% NaC₁₂₋₁₄AE₃S, 8.0% Mg(C_{11.8}LAS)₂, 30.0% glycerine, 0.72% Mg(OH)₂, 1.53% citric acid and 29.75% water
- c) 23.5% Mg(AE₃S)₂, 8.5% potassium methyl ester sulfonate and 68.0% water

1. A detergent composition comprising a gel wholly or predominantly in hexagonal liquid crystal phase, said gel comprising:
 - (a) from 15% to 50% ethoxylated alkyl sulfate surfactant by weight of the gel, wherein the alkyl group of ethoxylated alkyl sulfate surfactant has an average from 8 to 20 carbon atoms, and wherein the ethoxylated alkyl sulfate surfactant has an average degree of ethoxylation from 0.5 to 15;
 - (b) from 1% to 20%, by weight of the gel, secondary sulfonate surfactant selected from the group consisting of alkylbenzene sulfonates, alkyltoluene sulfonates, paraffin sulfonates, olefin sulfonates, alpha-sulfonated fatty acid alkyl esters, and mixtures thereof;
 - (c) optionally from 0% to 15% other surfactants by weight of the gel;
 - (d) from 40% to 80% water by weight of the gel; and
 - (e) from 0.2% to 3% magnesium ions by weight of the gel;

wherein the weight ratio of surfactant (a) to surfactant (b) is between 3:2 and 10:1, and wherein the total amount of surfactants (a)+(b)+(c) is from 25% to 60% by weight of the gel.
2. A detergent composition according to Claim 1 wherein the secondary sulfonate surfactant is selected from the group consisting of alkylbenzene sulfonates, alkyltoluene sulfonates, paraffin sulfonates, olefin sulfonates, and mixtures thereof.
3. A detergent composition according to Claim 2 wherein the total amount of surfactants (a)+(b)+(c) is from 25% to 50% by weight of the gel.
4. A detergent composition according to Claim 2 which contains from 20% to 45% ethoxylated alkyl sulfate surfactant by weight of the gel.
5. A detergent composition according to Claim 2 wherein the alkyl group of ethoxylated alkyl sulfate surfactant has an average from 8 to 15 carbon atoms, and wherein the ethoxylated alkyl sulfate surfactant has an average degree of ethoxylation from 1 to 6.5.
6. A detergent composition according to Claim 2 wherein the secondary sulfonate surfactant is selected from the group consisting of alkylbenzene sulfonates, paraffin sulfonates, and mixtures thereof.

7. A detergent composition according to Claim 6 wherein the secondary sulfonate surfactant is an alkylbenzene sulfonate.
8. A detergent composition according to Claim 2 wherein the weight ratio of surfactant (a) to surfactant (b) is between 2:1 and 5:1.
9. A detergent composition according to Claim 2 which comprises from 45% to 70% water.
10. A detergent composition according to Claim 2 which is a dishcare gel.

Relevant Technical Fields

(i) UK Cl (Ed.M) C5D DOX DHD DHZ

(ii) Int Cl (Ed.5) C11 D17/00

Search Examiner
M ElliottDate of completion of Search
10 October 1994

Databases (see below)

(i) UK Patent Office collections of GB, EP, WO and US patent specifications.

(ii)

Documents considered relevant
following a search in respect of
Claims :-
1-10

Categories of documents

- X:** Document indicating lack of novelty or of inventive step. **P:** Document published on or after the declared priority date but before the filing date of the present application.
- Y:** Document indicating lack of inventive step if combined with one or more other documents of the same category. **E:** Patent document published on or after, but with priority date earlier than, the filing date of the present application.
- A:** Document indicating technological background and/or state of the art. **&:** Member of the same patent family; corresponding document.

Category	Identity of document and relevant passages		Relevant to claim(s)
P,X	GB 2272449 A	(THE PROCTER & GAMBLE CO) whole document	1 at least
X	GB 2179054 A	(UNILEVER PLC) whole document	1 at least
X	US 4615819 A	(LEVER BROTHERS CO) whole document	1 at least

Databases: The UK Patent Office database comprises classified collections of GB, EP, WO and US patent specifications as outlined periodically in the Official Journal (Patents). The on-line databases considered for search are also listed periodically in the Official Journal (Patents).