



(86) **Date de dépôt PCT/PCT Filing Date:** 2013/05/22
(87) **Date publication PCT/PCT Publication Date:** 2013/11/28
(85) **Entrée phase nationale/National Entry:** 2014/11/20
(86) **N° demande PCT/PCT Application No.:** EP 2013/060543
(87) **N° publication PCT/PCT Publication No.:** 2013/174883
(30) **Priorité/Priority:** 2012/05/22 (EP12168968.1)

(51) **Cl.Int./Int.Cl. A61K 31/55** (2006.01),
A61K 31/5517 (2006.01), **A61K 47/06** (2006.01),
A61K 47/32 (2006.01), **A61K 47/36** (2006.01),
A61K 47/38 (2006.01), **A61K 9/14** (2006.01)

(71) **Demandeur/Applicant:**
PAION UK LIMITED, GB

(72) **Inventeurs/Inventors:**
GRAHAM, JOHN AITKEN, GB;
BAILLIE, ALAN JAMES, GB;
WARD, KEVIN RICHARD, GB;
PEACOCK, THOMAS, GB

(74) **Agent:** NORTON ROSE FULBRIGHT CANADA
LLP/S.E.N.C.R.L., S.R.L.

(54) **Titre : COMPOSITIONS COMPRENANT DES BENZODIAZEPINES A ACTION RAPIDE**
(54) **Title: COMPOSITIONS COMPRISING SHORT-ACTING BENZODIAZEPINES**

(57) **Abrégé/Abstract:**

A composition is provided with a benzodiazepine and at least one hygroscopic excipient, in particular lactose and/or dextran.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)
(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 November 2013 (28.11.2013)

(10) International Publication Number
WO 2013/174883 A1

(51) International Patent Classification:

A61K 31/55 (2006.01) *A61K 47/38* (2006.01)
A61K 47/06 (2006.01) *A61K 31/5517* (2006.01)
A61K 47/32 (2006.01) *A61K 9/14* (2006.01)
A61K 47/36 (2006.01)

(21) International Application Number:

PCT/EP2013/060543

(22) International Filing Date:

22 May 2013 (22.05.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

12168968.1 22 May 2012 (22.05.2012) EP

(71) Applicant: **PAION UK LIMITED** [GB/GB]; Chivers Way, Histon, Cambridge cambridgeshire CB24 9ZR (GB).

(72) Inventors: **GRAHAM, John Aitken**; 43 Fen End, Over, Cambridge Cambridgeshire CB24 5NE (GB). **BAILLIE, Alan James**; 107 Sinclair Street, Helensburgh, Glasgow Lancashire G84 9HY (GB). **WARD, Kevin Richard**; 1 Barn Lane, Oakley, Basingstoke, Hampshire Hampshire RG23 7HT (GB). **PEACOCK, Thomas**; 11 Clarendon Road, Southampton, Hampshire Hampshire S016 4GB (GB).


(74) Agents: **VON RENESSE, Dorothea** et al.; König Szynka Tilmann von Renesse, Mönchenwerther Straße 11, 40545 Düsseldorf (DE).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))



WO 2013/174883 A1

(54) Title: COMPOSITIONS COMPRISING SHORT-ACTING BENZODIAZEPINES

(57) Abstract: A composition is provided with a benzodiazepine and at least one hygroscopic excipient, in particular lactose and/or dextran.

“Compositions comprising short-acting benzodiazepines”

The present invention relates to compositions comprising benzodiazepines or pharmaceutically acceptable salts thereof and their use as pharmaceuticals.

5 Benzodiazepine compounds are known for their capacity to bind to a site on a specific receptor/chloride ion channel complex known as the GABA_A receptor. The binding of the benzodiazepine compound potentiates the binding of the inhibitory neurotransmitter 7-aminobutyric acid (GABA) to the complex, thereby leading to inhibition of normal neuronal function. Therapeutic purposes of the treatment with benzodiazepine compounds are in particular production of sedation or hypnosis, induction of anxiolysis, induction of muscle relaxation, treatment of convulsions or induction and/or maintenance of anesthesia in a mammal. See generally, Goodman and Gilman's The Pharmacological Basis of Therapeutics, Eighth Edition; Gilman, A. G.; Rall, T. W.; Nies, A. S.; Taylor, P., Eds. ; Pergamon Press: New York, 1990; pp. 10 303-304, 346-358.

15 Short-acting benzodiazepines that may provide faster recovery profiles have been the subject of clinical investigations (W. Hering et al., Anesthesiology 1996, 189, 85 (Suppl.); J. Dingemanse et al., Br. J. Anaesth 1997, 79, 567-574). Further compounds of interest are disclosed in WO 96/23790, WO 96/20941 and US 5665 718. Other publications that describe benzodiazepinones include E. Manghisi and A. Salimbemi, Boll. Chim. Farm. 1974, 113, 642-644, W. A. Khan and P. Singh, Org. Prep. Proc. Int.

- 2 -

1978, 10,105-111 and J. B. Hester, Jr, et al., J. Med. Chem. 1980, 23, 643-647. Benzodiazepines such as diazepam, lorazepam, and midazolam all undergo metabolism by hepatic-dependent processes. Active metabolites, which are often much more slowly metabolized than the parent drug, can be generated by these hepatic mechanisms in effect prolonging the duration of action of many benzodiazepines (T. M. Bauer et al, Lancet 1995, 346, 145-7). Inadvertent oversedation has been associated with the use of benzodiazepines (A. Shafer, Crit Care Med 1998, 26, 947-956), particularly in the intensive care unit, where benzodiazepines, such as midazolam, enjoy frequent use.

Short-acting benzodiazepines have been further disclosed in WO 2000/69836 A1. The benzodiazepines as disclosed herein comprise a carboxylic acid ester moiety and are inactivated by non-specific tissue esterases.

WO 2008/007071 A1 discloses a highly crystalline besylate salt of a benzodiazepine with a carboxylic acid ester moiety as disclosed in WO 2000/69836 A1. WO 2008/007081 A1 an esylate salt of a benzodiazepine.

Products which are used as sedatives or anesthetic agents are normally stored at room temperature. Therefore there is a need to provide formulations of short-acting benzodiazepines, which exhibit sufficient stability at room temperature. In addition for products that require to be presented as sterile, for example for injection via various routes, the process used to produce these formulations must be capable of being processed in a manner to ensure sterility assurance e.g. aseptic filtration.

It was in particular an object of the present invention to find pharmaceutically acceptable compositions with sufficient stability at room temperature for short-acting benzodiazepines as disclosed in WO 2000/69836 A1, WO 2008/007071 A1 and WO 2008/007081 A1.

Tests which were carried out with these compounds did show that aqueous solutions of the compounds do not have sufficient stability at room temperature and show strong degradation within a short period of time. Therefore alternative approaches had to be found.

A known technique for stabilizing water-labile compounds is the method of lyophilization. However, lyophilising the benzodiazepine of WO 2008/007071 A1 alone did not result in satisfactory stability of this benzodiazepine.


The inventors have now found that stable lyophilized formulations can be obtained, when mixtures of the benzodiazepine with hygroscopic excipients are formulated

- 3 -

and/or when the lyophilized formulation is at least in part amorphous. In addition it has been demonstrated that an alternative drying process, namely spray -drying, can be used to get the same effect.

A first aspect of the invention is therefore a composition comprising a mixture of at least one benzodiazepine or a pharmaceutically acceptable salt thereof and at least one pharmaceutically acceptable hygroscopic excipient, wherein the benzodiazepine comprises at least one carboxylic acid ester moiety.

According to the invention the benzodiazepine is preferably a compound according to formula (I)



wherein

W is H, a C₁-C₄ branched alkyl, or a straight chained alkyl;

X is CH₂, NH, or NCH₃; n is 1 or 2;

Y is O or CH₂; m is 0 or 1;

Z is O; p is 0 or 1;

R¹ is a C₁-C₇ straight chain alkyl, a C₃-C₇ branched chain alkyl, a C₁-C₄ haloalkyl, a C₃-C₇ cycloalkyl, an aryl, a heteroaryl, an aralkyl, or a heteroaralkyl;

R² is phenyl, 2-halophenyl or 2-pyridyl,

R³ is H, Cl, Br, F, I, CF₃, or NO₂;

(1) R⁴ is H, a C₁-C₄ alkyl, or a dialkylaminoalkyl and R⁵ and R⁶ together represent a single oxygen or S atom which is linked to the diazepine ring by a double bond and p is zero or 1; or (2) R⁴ and R⁵ together form a double bond in the diazepine ring and R⁶


represents the group NHR⁷ wherein R⁷ is H, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, benzyl or benzyl mono or disubstituted independently with halogen substituents, C₁₋₄ alkylpyridyl or C₁₋₄ alkylimidazolyl and p is zero; or (3) R⁴, R⁵ and R⁶ form the group -CR⁸=U-V= wherein R⁸ is hydrogen, C₁₋₄ alkyl or C₁₋₃ hydroxyalkyl, U is N or CR⁹ wherein R⁹ is H, C₁₋₄ alkyl, C₁₋₃ hydroxyalkyl or C₁₋₄ alkoxy-C₁₋₄alkyl, V is N or CH and p is zero.

- 4 -

The term "aryl", alone or in combination, is defined herein as a monocyclic or polycyclic group, preferably a monocyclic or bicyclic group, e.g., phenyl or naphthyl, which can be unsubstituted or substituted, for example, with one or more and, in particular, one to three substituents selected from halogen, C₁₋₄ branched or straight chained alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, hydroxy, nitro, amino, and the like. The term "heteroaryl" is defined herein as a 5-membered or 6-membered heterocyclic aromatic group which can optionally carry a fused benzene ring and wherein said 5-membered or 6-membered heterocyclic aromatic group can be unsubstituted or substituted, for example, with one or more and, in particular, one to three substituents selected from halogen, C₁₋₄ branched or straight chained alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, hydroxy, nitro, amino, and the like. The term "alkoxy", alone or in combination, is defined herein to include an alkyl group, which is attached through an oxygen atom to the parent molecular subunit. Exemplary alkoxy groups include but are not necessarily limited to methoxy, ethoxy and isopropoxy. The term "aralkyl" is defined herein as an alkyl group, in which one of the hydrogen atoms is replaced by an aryl group. The term "heteroaralkyl" is defined herein as an alkyl group, in which one of the hydrogen atoms is replaced by a heteroaryl group.

Exemplary branched or straight chained C₁₋₄ alkyl groups include but are not necessarily limited to methyl, ethyl, propyl, isopropyl, isobutyl and n-butyl. Exemplary C₁₋₇ straight chain alkyl groups include, but are not necessarily limited to, methyl, ethyl, propyl, n-butyl, n-hexyl and n-heptyl. Exemplary C₃₋₇ branched chain alkyl groups include, but are not necessarily limited to, isopropyl, isobutyl, sec-butyl, tert-butyl, isopentyl, neopentyl, tert-pentyl and isohexyl. Exemplary C₃₋₇ cycloalkyl groups include, but are not necessarily limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. Exemplary C₁₋₄ haloalkyl groups include, but are not necessarily limited, to methyl, ethyl, propyl, isopropyl, isobutyl and n-butyl substituted independently with one or more halogens, e.g., fluoro, chloro, bromo and iodo.


The compounds of formula (I) where the groups R⁴ and R⁵ and R⁶ together form the group -CR⁸=U-V= and p is 0 represent a preferred embodiment of the invention and may be conveniently represented by the compound of formula (II):



- 5 -

wherein R^1 , R^2 , R^3 , R^8 , U, V, W, X, Y, n and m have the meanings given for formula (I).

Further preferred are compounds of formula (I)



with

W is H;

X is CH_2 ; n is 1;

Y is CH_2 ; m is 1;

Z is O; p is 0 or 1;

R^1 is CH_3 , CH_2CH_3 , $CH_2CH_2CH_3$, $CH(CH_3)_2$ or $CH_2CH(CH_3)_2$;

R^2 is 2-fluorophenyl, 2-chlorophenyl or 2-pyridyl;

R^3 is Cl or Br;

(1) R^4 is H, a C_1 - C_4 alkyl, or a dialkylaminoalkyl and R^5 and R^6 together represent a single oxygen or S atom which is linked to the diazepine ring by a double bond and p is zero or 1; or (2) R^4 and R^5 together is a double bond in the diazepine ring and R^6 represents the group NHR^7 wherein R^7 is H, C_{1-4} alkyl, C_{1-4} hydroxyalkyl, benzyl or benzyl mono or disubstituted independently with halogen substituents, C_{1-4} alkylpyridyl or C_{1-4} alkylimidazolyl and p is zero; or (3) R^4 , R^5 and R^6 form the group- $CR^8=U-V=$ wherein R^8 is hydrogen, C_{1-4} alkyl or C_{1-3} hydroxyalkyl, U is N or CR^9 wherein R^9 is H, C_{1-4} alkyl, C_{1-3} hydroxyalkyl or C_{1-4} alkoxy, V is N or CH and p is zero.

Preferably, in particular in compounds according to formula (II), W is H, X is CH_2 , n is 1; Y is CH_2 , m is 1; R^1 is CH_3 , CH_2CH_3 , $CH_2CH_2CH_3$, $CH(CH_3)_2$ or $CH_2CH(CH_3)_2$; R^2 is 2-fluorophenyl, 2-chlorophenyl or 2-pyridyl; R^3 is Cl or Br; R^8 is H, CH_3 or CH_2OH ; R^9 is H, CH_3 , CH_2OH or CH_2O -t-butyl; U is CR^9 or N; and V is N or CH.

Particularly preferred amongst these compounds are compounds according to formula (II), wherein in each compound W is H, X is CH_2 , n is 1, Y is CH_2 , m is 1 and wherein R^1 , R^2 , R^3 , R^8 , U and V for each compound are as follows:

R^1	R^2	R^3	R^8	U	V
CH_3	2-fluorophenyl	Cl	H	CH	N

- 6 -

CH ₃	2-fluorophenyl	Cl	CH ₃	CH	N
CH ₃	2-fluorophenyl	Cl	H	C-CH ₃	N
CH ₃	2-fluorophenyl	Cl	H	C-CH ₂ OH	N
CH ₃	2-fluorophenyl	Cl	CH ₂ OH	CH	N
CH ₃	2-pyridyl	Cl	H	CH	N
CH ₃	2-pyridyl	Cl	CH ₃	CH	N
CH ₃	2-pyridyl	Br	CH ₃	CH	N
CH ₃	2-pyridyl	Br	H	C-CH ₃	N
CH ₃	2-pyridyl	Cl	H	C-CH ₃	N
CH ₃	2-pyridyl	Cl	H	C-CH ₂ OH	N
CH ₃	2-pyridyl	Cl	CH ₂ OH	CH	N
CH ₃	2-pyridyl	Cl	CH ₃	C-CH ₃	N
CH ₃	2-chlorophenyl	Cl	CH ₃	N	N
CH ₃	2-fluorophenyl	Cl	CH ₃	N	N
CH ₃	2-fluorophenyl	Cl	CH ₃	N	N
CH ₃	2-fluorophenyl	Cl	H	N	CH
CH ₃	2-fluorophenyl	Cl	CH ₃	N	CH
CH ₃	2-fluorophenyl	Cl	H	C-CH ₂ O-t-butyl	N
CH ₃	2-pyridyl	Cl	CH ₃	C-CH ₂ OH	N

Amongst these compounds the most preferred is remimazolam (INN), wherein W is H, X is CH₂, n is 1, Y is CH₂, m is 1, R¹ is CH₃, R² is 2-pyridyl, R³ is Br, R⁸ is CH₃, U is CH and V is N. According to IUPAC system remimazolam is methyl 3-[(4S)-8-bromo-1-methyl-6-(pyridin-2-yl)-4H-imidazo[1,2-a][1,4]benzodiazepin-4-yl]propanoate. It is clinically developed by PAION AG, Aachen under the internal designation "CNS7056". The besylate form of CNS7056 is also called "CNS7056B" (see the experimental data *infra*).

Compounds according to formula (I) and (II) possess a stereocenter. According to the invention enantiomeric pure forms can be used, which are substantially free of the other enantiomer, but also racemic mixtures can be used.

The composition according to the invention might comprise the free form of the benzodiazepine, but in a preferred embodiment of the invention the benzodiazepine is used in the form of a salt, in particular in the form of an inorganic or organic salt. In a very preferred embodiment the benzodiazepine is used in the salt in a cationic form.

- 7 -

The counter ion of the cationic benzodiazepine is preferably selected from halogenides, in particular fluoride, chloride or bromide, sulfate, organic sulfates, sulfonate, organic sulfonates, nitrate, phosphate, salicylate, tartrate, citrate, maleate, formate, malonate, succinate, isethionate, lactobionate and sulfamate.

The salts of the invention are obtained by reaction of the benzodiazepine with suitable acids, in particular by reaction with the following acids: hydrochloric, hydrobromic, sulfuric, nitric, phosphoric, salicylic, p-toluenesulfonic, tartaric, citric, methanesulfonic, maleic, formic, malonic, succinic, isethionic, lactobionic, naphthalene-2-sulfonic, sulfamic, ethanesulfonic and benzenesulfonic.

In a preferred embodiment the counter ion is selected from organic sulfates and organic sulfonates, in particular from aromatic sulfates and aromatic sulfonates. In a very preferred embodiment an organic sulfonate is used as counter ion, preferably an aromatic sulfonate, in particular p-toluenesulfonic acid (tosylate), naphthalene-2-sulfonic acid, ethanesulfonic acid (esylate) or benzenesulfonic acid, wherein benzenesulfonic acid (besylate) is the most preferred counter ion.

The most preferred salts according to the invention are the besylate salt (as disclosed in WO 2008/007071 A1) or the esylate salt (as disclosed in WO 2008/007081 A1) of remimazolam. The tosylate of remimazolam is also preferred and is subject matter of WO 2013029431 A1.

The compositions according to one aspect of the invention comprise at least one pharmaceutically acceptable hygroscopic excipient. The hygroscopic excipient might be an organic or inorganic substance, but is preferably an organic substance. The hygroscopic excipient does not include water as such but water can be present in addition to a hygroscopic excipient. The hygroscopic excipient is preferably a compound which is able to form stable hydrates.

The hygroscopic excipient is preferably a substance which under normal conditions (25°C, 1013,25 hPa) binds water molecules reversibly and is preferably further able to release the water molecules, when sufficient vacuum and/or heat is applied. Vice versa the hygroscopic excipient as obtained by application of vacuum and/or heat after dehydration is able to bind water molecules again. Under normal temperature conditions (25°C) the water vapour pressure of the hydrated hygroscopic excipients is preferably less than 23 hPa, more preferably less than 20 hPa, preferentially less than 15 hPa, in particular less than 10 hPa. In a particularly preferred embodiment the water vapour pressure of the hydrated hygroscopic excipient is between 2 and 20 hPa, more preferably between 5 and 15 hPa. The dehydrated hygroscopic excipient preferably has a capacity to bind at least 0.01 g, more preferably at least 0.03 g, in

- 8 -

particular at least 0.05 g or at least 1 g water per g of hygroscopic substance. In a preferred embodiment the dehydrated hygroscopic excipient can bind up to 5 g, more preferably up to 10 g, in particular up to 20 g water per g of hygroscopic substance

5 The organic hygroscopic excipients according to the invention preferably possess a molecular weight of less than 400 kD, preferably less than 350 kD, more preferably less than 100 kD., especially preferably less than 20 kD and further preferably less than 1 kD. In a most preferred embodiment the hygroscopic excipient has a molecular weight of less than 0.1 kD.

10 According to the invention the term "excipient" is defined as an ingredient added intentionally to the drug substance which should not have pharmacological properties in the quantity used. Such excipients can provide some other beneficial purpose be this to aid processing, solubility or dissolution, drug delivery via the target route of administration or aid stability.

15 Within the context of the present invention, the definition of "pharmaceutically acceptable" is meant to encompass any substance, which does not interfere with effectiveness of the biological activity of the active ingredient and that is not toxic to the host to which it is administered.

20 In a preferred embodiment the organic hygroscopic excipient is selected from carbohydrates and/or organic polymers.

25 According to the invention a "carbohydrate" is an organic compound with the empirical formula $C_m(H_2O)_n$ (where m could be different from n). Structurally, carbohydrates can be described as polyhydroxy aldehydes and ketones. The term "saccharide" or "sugar" as used hereinafter is a synonym of the term carbohydrate. The carbohydrates are divided into four chemical groups: monosaccharides, disaccharides, oligosaccharides and polysaccharides. The carbohydrates as defined
30 herein encompasses all modifications, derivatives and analogues of carbohydrates such as acidic saccharides containing carboxyl groups, phosphate groups and/or sulfuric ester groups

35 In a very preferred embodiment the organic hygroscopic excipient is a carbohydrate or a mixture of various, at least two types of carbohydrates. Suitable carbohydrates are for example amylose, amylopectin, alginate, dextrans, starches as well as mono-, di- and oligosaccharides.

40 The inventors have found that the use of hygroscopic carbohydrates or mixtures thereof is especially suitable in order to prepare stable solid

formulations/compositions – e.g. lyophilized or spray dried compositions – for benzodiazepines, in particular remimazolam salts, which have a favourable reconstitution time.

5 In a preferred embodiment the carbohydrate possesses a molecular weight of less than 150 kD, preferably less than 100 kD, in particular less than 80 kD, especially preferably less than 20 kD and further preferably less than 1 kD (e.g. less than 0.5 kD). In a preferred embodiment oligo- or polysaccharide chains are non-cyclic, i.e. they are not cyclic hemiacetals or hemiketals.

10 In a particularly preferred embodiment dextrans with a molecular weight of less than 150 kD, preferably less than 100 kD, in particular less than 80 kD are used as hygroscopic excipient. Very preferred dextrans possess a molecular weight of between 5 and 80 kD, in particular of between 10 and 40 kD.

15 In another particularly preferred embodiment mono- or oligosaccharides are used as hygroscopic excipients - either as the sole hygroscopic excipient or in a mixture with at least one further hygroscopic excipient, such as dextran - wherein the oligosaccharides are preferably non-cyclic. In this embodiment the carbohydrate is preferably selected from monosaccharides and C₂₋₆-oligosaccharides, in particular from disaccharides. The disaccharide is preferably selected from lactose, maltose, sucrose and trehalose and is most preferably lactose. In yet another embodiment two or more disaccharides can be used, in particular including lactose. These disaccharides can be combined with further excipients, e.g. dextran.

25 In another very preferred embodiment the organic hygroscopic excipient is a polymer, preferably a polyacrylate or a vinylpolymer, more preferably a polyvinylpyrrolidone.

30 In a preferred embodiment the polyvinylpyrrolidone possesses a molecular weight of less than 150 kD, preferably less than 100 kD, in particular less than 50 kD. Very preferred polyvinylpyrrolidones possess a molecular weight of between 5 and 40 kD, in particular of between 10 and 30 kD.

35 In a further embodiment of the invention the hygroscopic excipient is a mixture of at least two different hygroscopic excipients, in particular exactly two, three, four, five or more excipients. These at least two different excipients can be of the same chemical nature e.g. both are carbohydrates or both are organic polymers. Alternatively they can be of different chemical nature, e.g. one or more are carbohydrates and one or more are organic polymers. In a preferred embodiment of the invention the composition comprises a mixture of at least two carbohydrates.

40

- 10 -

These at least two carbohydrates can be from the same type of carbohydrates, e.g. they all represent monosaccharides, disaccharides, oligosaccharides or polysaccharides, respectively. The carbohydrates can alternatively represent different types of carbohydrates, e.g. one or more monosaccharide combined with one or more disaccharide etc. Preferred is a combination of at least one disaccharide with at least one or more polysaccharide.

Particularly preferred is the combination of one disaccharide and one polysaccharide. The disaccharide preferably is lactose and the polysaccharide is preferably dextran, in particular with a molecular weight of 80 KD or less. The composition preferably contains remimazolam, preferably in besylate, esylate or tosylate salt. Of particular relevance is the besylate salt thereof.

Particular preferred carbohydrate mixtures of the invention comprise or consist of the combinations given in the following table:

Disaccharide 1	Disaccharide 2	Polysaccharide
Lactose	-	Dextran 40
Trehalose	-	Dextran 40
Sucrose	-	Dextran 40
Lactose	-	Dextran 70
Trehalose	-	Dextran 70
Sucrose	-	Dextran 70
Lactose	Trehalose	Dextran 40
Lactose	Sucrose	Dextran 40
Lactose	Trehalose	Dextran 70
Lactose	Sucrose	Dextran 70
Trehalose	Sucrose	Dextran 40
Trehalose	Sucrose	Dextran 70

A composition with the above listed disaccharide/dextran mixture preferably comprise remimazolam, either in its besylate, esylate or tosylate salt. Especially preferred is the besylate salt.

As outlined above the inventors have found that the composition according to the invention, in particular a composition with a mixture of at least one disaccharide such as lactose and at least one dextran can form stable solid, in particular lyophilized or spray dried, formulations with an acceptable lyophilisation (also called "total cycle duration") and/or reconstitution time. According to the invention, a favourable reconstitution time is 5 min or less, preferably 3 minutes or less, more preferably 2 or

- 11 -

even 1 minute. A reconstitution time of 1 min is further preferred and a reconstitution time of less than 1 minute is most preferred.

5 The lyophilisation time for the composition of the invention favourably is less than 120 hours, preferably less than 100 hours, more preferably less than 80 hours and even more preferably less than 70 hours, and specifically 66 hours.

10 This reduction in lyophilisation time in particular applies when the primary drying step is performed at -25 °C and below 100 mTorr (e.g. between 90 and 100 mTorr) or at -15 °C or above and 350 to 750 mTorr.

15 The inventors have found a correlation between the amount of polymer, in particular polysaccharide, more particular dextran, and the time required for the lyophilisation: an increasing amount of polysaccharide in the mixture of carbohydrate excipients increases the collapse temperature of the composition and therewith reduces the time required for lyophilisation. For example a composition with remimazolam salt further comprising lactose and dextran in a weight ratio of 1:1 shows a lyophilisation time of 99 hours, whereas the same composition with a lactose and dextran weight ratio of 1:4 requires a lyophilisation time of only 66 hours or less.

20 Within the mixtures of excipients (excipients of different chemical nature or of different types, *supra*), the wt.-% ratio between the first excipient (e.g. the disaccharide) and the second excipient (e.g. the dextran) can range from 1:1 to 1:10, more preferably from 1:1 to 1:6, even more preferably from 1:1 to 1:5 and particularly preferably from 1:1.0 to 1:4.5. In a specific embodiment said wt.-%ratio is 1:1.5 or 1:4. The first excipient is in particular lactose and the second excipient preferably is dextran, in particular dextran 70 or dextran 40.

25 30 Lactose can be used as a hydrate. However, unless otherwise explicitly mentioned the weight ratios and concentrations provided herein relate to lactose. The same applies to other excipients suitable according to the invention.

35 It is especially preferred when the relative amount of polysaccharide in the mixture exceeds the relative amount of disaccharides therein. Hence 50 wt.-% or more of the mixture of carbohydrates can be a polysaccharide, more preferably 60 wt.-% or more, even more preferred 80 wt.-% or more. In a binary mixture the rest preferably is disaccharide. In these embodiments of the invention it is possible to improve the lyophilisation time, i.e. to obtain higher collapse temperatures. Preferably the polysaccharide is dextran.

40

- 12 -

5 The composition of the invention can comprises the benzodiazepine or a salt thereof, being preferably the besylate or tosylate salt of remimazolam, in a relative amount between 5 and 50 wt.%, more preferably in a relative amount between 8 and 25 wt.-%, even more preferably in a relative amount between 10 and 20 wt.%, and specifically in relative amounts of 10 or 19 wt.%. Notably, all relative amounts, weight ratios etc. of the benzodiazepine, in particular remimazolam, in the compositions of the invention are calculated for the free base; unless otherwise explicitly outlined.

10 The composition of the invention can comprises the total amount of hygroscopic excipients, being preferably a carbohydrate or a mixture of carbohydrates, in a relative amount between 50 and 95 wt.%, more preferably in a relative amount between 75 and 92 wt.%, even more preferably in a relative amount between 80 and 90 wt.-%, and specifically in relative amounts of 81 or 90 wt.%.

15 The wt.% ratio between the total amount of hygroscopic excipients and total amount of benzodiazepines or salts thereof in the composition – calculated for the free base - is preferably at least 1:1, more preferably at least 2:1, 3:1 or 4:1, in particular at least 5:1, 6:1, 7:1 or 9:1. In particularly preferred embodiments the wt.% ratio between the total amount of hygroscopic excipients and the total amount of benzodiazepines or salts thereof, calculated for the free base, in the composition is between 1:1 and 20 100:1, particularly between 3:1 and 50:1, more preferably between 5:1 and 25:1, most preferably between 7:1 and 15:1, and in the most preferred embodiment at 13:1.

25 In one embodiment of the invention the composition contains only hygroscopic excipients.

In one aspect the composition of the invention has a collapse temperature above -20.5°C, preferably above -18° and more preferably above -15.5°C.

30 In a further aspect the composition of the invention the collapse temperature of the composition is increased by the addition of at least one compound with a collapse temperature above -20°C (collapse temperature modifier).

35 In a further aspect the composition of the invention further comprises at least one compound with a collapse temperature above -20°C (hereinafter called “collapse temperature modifier”). This component is added to the composition which is then further dried (in particular by lyophilization) to form a solid composition.

40 The collapse temperature modifier according to the invention can be selected from the group consisting of a sucrose-epichlorhydrin-copolymer (such as Ficoll®), gelatine and hydroxyethyl starch (HES) or dextran. In a preferred embodiment of the invention

- 13 -

the collapse temperature modifier is HES. In yet another preferred embodiment of the invention the collapse temperature modifier is dextran.

5 The collapse temperature modifier can be present within the composition of the invention in a relative amount from 1 to 75 wt.%, more preferably in a relative amount from 5 to 50 wt.%, even more preferably in a relative amount from 10 to 40wt.%. The collapse temperature modifier can be identical with the hygroscopic excipient.

10 The term "collapse temperature" as used in the context of the present invention relates to the temperature at which softening of a solid composition (the "cake") progresses to structural "collapse", a phenomenon that can be observed by Freeze Drying Microscopy (FDM). For a crystallizing system, collapse occurs if the lowest eutectic melting temperature (T_{eu}) is exceeded. For a non-crystallizing system, the collapse temperature is determined by the glass transition temperature (T_g), which
15 can be e.g. measured using differential scanning calorimetry (DSC). The determination of the collapse temperature is common knowledge of the skilled person. Hence, for an amorphous substance the collapse temperature is given by its glass transition temperature.

20 Hence the term "collapse" in particular relates to a loss of the integral structure of the solid composition (cake) and/ or to a reduction of its volume of at least about 10%, 25%, 50%, 75%, 85%, 95% or 100%. The reduction in volume and the loss of structural integrity can be measured using known methodologies, including but not limited to visual inspection or Brunauer–Emmett–Teller (BET) surface area analysis.

25 In one aspect of the invention the counter ion can render the salt hygroscopic. Hence in this embodiment the salt of the benzodiazepine constitutes also the excipient.

30 In another embodiment of the invention the compositions according to the invention might comprise besides the at least one hygroscopic excipient further pharmaceutically acceptable carriers and/or excipients. The further carriers and/or excipients must, if used, of course be acceptable in the sense of being compatible with the other ingredients of the formulation and must not be deleterious to the patient. Accordingly, the present invention provides in a further embodiment a
35 composition as hereinbefore defined and further pharmaceutically acceptable carriers and/or excipients. The further carrier and/or excipient for example might be selected from ascorbic acid, glycine, glycine hydrochloride, sodium chloride, sugar alcohols, and mixtures thereof. In a preferred embodiment the further excipient is selected from sugar alcohols, in particular C_{3-6} sugar alcohols, more preferably C_6 sugar alcohols.

40

- 14 -

In the context of the invention a sugar alcohol (also known as a polyol, polyhydric alcohol or polyalcohol) is defined as a hydrogenated form of carbohydrate, whose carbonyl group (aldehyde or ketone, reducing sugar) has been reduced to a primary or secondary hydroxal group).

5 A second aspect of the invention is a composition comprising at least one benzodiazepine or a pharmaceutically acceptable salt thereof, wherein at least parts of the composition are amorphous and wherein the benzodiazepine comprises at least one carboxylic acid ester moiety. The composition can also contain crystalline
10 parts/compounds.

In yet another aspect of the invention the composition comprises a mixture of said benzodiazepine with at least one hygroscopic excipient, wherein said composition is at least in parts amorphous, but it can also contain crystalline parts.

15 The compositions according to the invention are in a preferred embodiment solid composition, in particular obtained by lyophilization or spray drying. The dried composition contains at least one compound (e.g. the excipient) in amorphous form. In a preferred embodiment the lyophilized composition consists of a mixture of
20 amorphous and crystalline, in particular microcrystalline, parts/compounds. In a preferred embodiment the crystalline part of the lyophilized solids comprises or preferentially substantially consists of the benzodiazepine compounds or salts thereof.

25 In a further embodiment of the invention, at least 50% (w/w), preferably at least 75% (w/w), more preferably at least 90% (w/w) and most preferably at least 95% (w/w) of the benzodiazepine within the composition is in an amorphous state. In preferred embodiment of the invention, at least 96%, 97%, 88% or 99% (w/w) of the benzodiazepine within the composition is in an amorphous state. In a preferred
30 embodiment of the invention, the composition is amorphous for at least 96%, 97%, 88% or 99%.

In one embodiment of the invention the composition contains a mixture of crystalline and amorphous benzodiazepine or the benzodiazepine salt. In one embodiment at
35 least 25 %, 50-75 % or greater than 90% (w/w) of the total benzodiazepine or the benzodiazepine salt of the composition is crystalline.

In the most preferred embodiment of the invention the benzodiazepine salt is remimazolam besylate. When the composition contains crystalline remimazolam
40 besylate, in one embodiment, the crystalline polymorph (herein designated besylate Form 1) exhibits an X-ray powder diffraction (XRPD) pattern which comprises a

- 15 -

characteristic peak at about 7.3, 7.8, 9.4, 12.1, 14.1, 14.4, 14.7, or 15.6 degrees two-theta.

Preferably the besylate Form 1 of remimazolam crystalline polymorph exhibits an XRPD pattern which comprises characteristic peaks at about 7.3, 7.8, 9.4, 12.1, 14.1, 14.4, 14.7, and 15.6 degrees two-theta.

More preferably the besylate Form 1 crystalline polymorph exhibits an XRPD pattern which comprises characteristic peaks at: 7.25 (10.60), 7.84 (72.60), 9.36 (12.10), 12.13 (32.50), 14.06 (48.50), 14.41 (74.30), 14.70 (50.70), 15.60 (26.90) [angle two-theta degrees (percentage relative intensity)].

Preferably the besylate Form 1 crystalline polymorph has a differential scanning calorimetry (DSC) onset melting temperature in the range 187-204°C, preferably about 191-192°C.

The structure of a 2-methoxyethanol:pentyl acetate grown needle habit crystal of Form 1 has been resolved at 190K (R factor of 6.3, example 9 of WO2008/007071 A1). Form I has a stoichiometry of 1:1 compound:besylate. Its crystallographic asymmetric unit contains two independent compound molecules and two besylate molecules. The two independent compound molecules are singly protonated on the imidazole ring. The crystal structure has unit cell dimensions of $a = 7.6868 \text{ \AA}$, $b = 29.2607 \text{ \AA}$, $c = 12.3756 \text{ \AA}$, $\alpha = 90^\circ$, $\beta = 97.7880^\circ$, $\gamma = 90^\circ$, and a space group of $P2_1$. The crystal structure further features the following parameters: system: monoclinic, volume: $2757.86 \text{ \AA}^3$, density: 1.439 g cm^{-3} , absorption: $1.610 \mu [\text{MoK}\alpha] (\text{mm}^{-1})$, $F(000)$: 1224. The Flack "Enantiopole" parameter was determined as 0.03. The crystal structure is also described in more detail in Example 9 of WO2008/007071 A1, and crystallographic coordinates are given in Figure 5A to 5D (corresponding to table 17 of WO2008/007071 A1). Bond lengths and angles for Form 1 are given in Figures 7A-B and 8A-C, respectively (corresponding to Tables 19 and 20 of WO2008/007071 A1).

According to the invention the composition can comprise a besylate salt of remimazolam which is a crystalline polymorph comprising a crystal with unit cell dimensions of $a = 7.6868 \text{ \AA}$, $b = 29.2607 \text{ \AA}$, $c = 12.3756 \text{ \AA}$, $\alpha = 90^\circ$, $\beta = 97.7880^\circ$, $\gamma = 90^\circ$.

There is also provided according to the invention a composition with a besylate salt of remimazolam which is a crystalline polymorph having a crystal structure defined by the structural coordinates as shown in Figure 5A-D. The crystalline form preferably has a bond lengths and angles as shown in Figures 7A-B and 8A-C, respectively.

- 16 -

In a further embodiment the composition according to the invention comprises a polymorph of the besylate salt of remimazolam (herein designated besylate Form 2), that exhibits an XRPD pattern which comprises a characteristic peak at about 8.6, 10.5, 12.0, 13.1, 14.4, or 15.9 degrees two-theta. Preferably the besylate Form 2 crystalline polymorph exhibits an XRPD pattern which comprises characteristic peaks at about 8.6, 10.5, 12.0, 13.1, 14.4, and 15.9 degrees two-theta.

More preferably the besylate Form 2 crystalline polymorph exhibits an XRPD pattern which comprises characteristic peaks at: 8.64 (17.60), 10.46 (21.00), 12.03 (22.80), 13.14 (27.70), 14.42 (11.20), 15.91 (100.00) [angle two-theta degrees (percentage relative intensity)].

Preferably the besylate Form 2 crystalline polymorph has a differential scanning calorimetry (DSC) onset melting temperature in the range 170-200°C, preferably about 180°C.

The structure of an ethanol:ethyl acetate grown plate habit crystal of Form 2 has been resolved at 190K (R factor of 3.8, example 10 of WO2008/007071 A1). Form 2 has stoichiometry of 1:1 compound:besylate. Its crystallographic asymmetric unit contains one compound molecule and one besylate molecule. The compound molecule is singly protonated on the imidazole ring. The crystal structure has unit cell dimensions of $a = 8.92130 \text{ \AA}$, $b = 11.1536 \text{ \AA}$, $c = 25.8345 \text{ \AA}$, $\alpha = 90^\circ$, $\beta = 90^\circ$, $\gamma = 90^\circ$, and a space group of $P2_12_12_1$. The crystal structure further features the following parameters: system: orthorhombic, volume: $2570.65 \text{ \AA}^3$, density: 1.544 g cm^{-3} , absorption: $1.727 \mu [\text{MoK}\alpha] (\text{mm}^{-1})$, $F(000)$: 1224. The Flack "Enantiopole" parameter was determined as 0,011. The crystal structure is described in more detail in Example 10 of WO2008/007071 A1, and crystallographic coordinates are given in Figure 6A-C (corresponding to Table 18 of WO2008/007071 A1). Bond lengths and angles for Form 2 are given in Figures 9 and 10, respectively (corresponding to Tables 21 and 22 of WO2008/007071 A1).

According to the invention there is provided a composition with a besylate salt of remimazolam, which is a crystalline polymorph comprising a crystal with unit cell dimensions of $a = 8.92130 \text{ \AA}$, $b = 11.1536 \text{ \AA}$, $c = 25.8345 \text{ \AA}$, $\alpha = 90^\circ$, $\beta = 90^\circ$, $\gamma = 90^\circ$.

There is also provided according to the invention a composition with a besylate salt of remimazolam which is a crystalline polymorph having a crystal structure defined by the structural coordinates as shown in Figure 6A-C. There is further provided according to the invention a composition with a besylate salt of remimazolam with bond lengths and angles as shown in Figures 9 and 10, respectively.

- 17 -

There is further provided according to the invention a composition with a crystalline polymorph of a besylate salt of remimazolam (herein designated besylate Form 3), that exhibits an X-ray powder diffraction (XRPD) pattern which comprises a characteristic peak at about 7.6, 11.2, 12.4, 14.6, 15.2, 16.4, or 17.7 degrees two-theta. Preferably the besylate Form 3 crystalline polymorph exhibits an XRPD pattern which comprises characteristic peaks at about: 7.6, 11.2, 12.4, 14.6, 15.2, 16.4, and 17.7 degrees two-theta.

More preferably the besylate Form 3 crystalline polymorph exhibits an XRPD pattern which comprises characteristic peaks at: 7.61 (65.70), 11.19 (33.20), 12.38 (48.70), 14.63 (30.60), 15.18 (33.20), 16.40 (29.60), 17.68 (51.30) [angle two-theta degrees (percentage relative intensity)].

Preferably the besylate Form 3 crystalline polymorph has a differential scanning calorimetry (DSC) onset melting temperature in the range 195-205°C, preferably about 200-201°C.

There is further provided according to the invention a composition with a crystalline polymorph of a besylate salt of remimazolam (herein designated besylate Form 4), that exhibits an XRPD pattern which comprises a characteristic peak at about 7.6, 10.8, 15.2, 15.9, or 22.0 degrees two-theta. Preferably the besylate Form 4 crystalline polymorph exhibits an XRPD pattern which comprises characteristic peaks at about: 7.6, 10.8, 15.2, 15.9, and 22.0 degrees two-theta.

Preferably the besylate Form 4 crystalline polymorph exhibits an XRPD pattern which comprises characteristic peaks at: 7.62 (83.50), 10.75 (14.70), 15.17 (37.80), 15.85 (28.70), 22.03 (100) [angle two-theta degrees (percentage relative intensity)].

Preferably the besylate Form 4 crystalline polymorph has a differential scanning calorimetry (DSC) onset melting temperature in the range 180-185°C, preferably about 182°C.

The besylate Forms 1 to 4 may be prepared and crystallised by using the methods and solvents disclosed in WO 2008/007071 A1.

A preferred salt is the besylate Form 1 based on the robustness of formation, yield, purity and chemical and solid form stability.

- 18 -

In one embodiment of the invention the composition comprises a mixture of Forms 1, 2, 3 and 4. However compositions with only one of the Forms 1 to 4 are possible.

5 In another preferred embodiment of the invention the benzodiazepine salt is remimazolam esylate. When the composition contains crystalline remimazolam esylate, in one embodiment, the crystalline polymorph (herein designated esylate Form 1) exhibits an X-ray powder diffraction (XRPD) pattern which comprises a characteristic peak at about 6.2, 9.2, 12.3, 15.0, 17.2, or 20.6 degrees two-theta.

10 Preferably the esylate Form 1 crystalline polymorph exhibits an XRPD pattern which comprises characteristic peaks at about 6.2, 9.2, 12.3, 15.0, 17.2, and 20.6 degrees two-theta.

15 More preferably the esylate Form 1 crystalline polymorph exhibits an XRPD pattern which comprises characteristic peaks at: 6.17 (19.30), 9.21 (20.50), 12.28 (16.40), 14.97 (23.40), 17.18 (52.80), 20.63 (100.00) [angle two-theta degrees (percentage relative intensity)].

20 Preferably the esylate Form 1 crystalline polymorph has a differential scanning calorimetry (DSC) onset melting temperature in the range 195-205°C, preferably about 201-202°C.

25 There is further provided according to the invention a crystalline polymorph of an esylate salt of a compound of formula (I) (herein designated esylate Form 2) that exhibits an X-ray powder diffraction (XRPD) pattern which comprises a characteristic peak at about 3.6, 6.4, 7.1, 12.3, 14.1, or 17.1 degrees two-theta.

30 Preferably the esylate Form 2 crystalline polymorph exhibits an XRPD pattern which comprises characteristic peaks at about 3.6, 6.4, 7.1, 12.3, 14.1, and 17.1 degrees two-theta.

35 More preferably the crystalline polymorph exhibits an XRPD pattern which comprises characteristic peaks at: 3.57 (15.60), 6.42 (21.10), 7.13 (58.30), 12.29 (51.50), 14.10 (58.90), 17.13 (68.00) [angle two-theta degrees (percentage relative intensity)].

Preferably the esylate Form 2 crystalline polymorph has a differential scanning calorimetry (DSC) onset melting temperature in the range 185-195°C, preferably about 190-191°C.

- 19 -

The esylate Forms 1 and 2 may be prepared and crystallised by using the methods and solvents disclosed in WO 2008/007081 A1

5 A preferred salt is the esylate Form 1 based on the robustness of formation, yield, purity and chemical and solid form stability.

In one embodiment of the invention the composition comprises a mixture of Forms 1, and 2. However compositions with only one of the Forms 1 or 2 are possible.

10 The lyophilized form of the composition according to the invention is preferably used for storage of the compositions.

15 The solid form of the compositions, in particular the lyophilized or spray dried solids, preferably show very good storage stability. In a preferred embodiment, they show degradation of the benzodiazepine, in particular hydrolysis of the carboxylic ester moiety, of less than 1 % during storage for 13 weeks, in particular at storage conditions of 40°C/75%RH.

20 The solid, in particular lyophilized or spray dried, compositions according to the invention in a preferred embodiment maintain a room temperature shelf life of at least one year, more preferably of at least two years, in particular of at least three years. They further comprise in a preferred embodiment less than 5 wt.% of water, preferably less than 2 wt.% of water, more preferably less than 1 wt.% of water.

25 In the solid, in particular lyophilized or spray dried, compositions according to the invention the total amount of benzodiazepines or salts thereof and hygroscopic excipients preferably sums up to at least 50 wt.%, more preferably at least 70 wt.%, in particular at least 90 wt.% of the composition.

30 In another preferred embodiment the compositions according to the invention are in liquid form, more preferably aqueous solutions. The liquid form is on the one hand used for the preparation of the lyophilized or spray dried solids, and on the other hand obtained by solubilization of the lyophilized or spray dried solids when transforming the lyophilized composition into a suitable pharmaceutically applicable solution.

35 The liquid contains the reconstituted solid benzodiazepine as free base preferably in an amount of between 0.5 and 30 mg/ml, more preferably in an amount of between 1 and 20 mg/ml, in particular in an amount of between 2 and 10 mg/ml.

40 Further subject of the present invention is a pharmaceutical comprising a composition according to the invention.

- 20 -

Subject of the invention is therefore also a method of manufacturing a composition or pharmaceutical composition according to the invention, wherein the composition or pharmaceutical composition is in the solid state, comprising the following steps:

- 5
- a) providing a solution comprising at least one benzodiazepine with at least one carboxylic acid ester moiety or a pharmaceutically acceptable salt thereof (in particular remimazolam salt) as hereinbefore described and at least one pharmaceutically acceptable hygroscopic excipient or mixtures of at least two
- 10
- hygroscopic excipients as hereinbefore described, wherein the solution is preferably an aqueous solution and wherein the solution preferably possesses a pH of between 2 and 7, preferably 2 and 5 and more preferably 2 and 4;
- b) lyophilizing the solution according to (a).

15

In a preferred embodiment the lyophilisation time of step b) is less than 120 hours, preferably less than 100 hours, more preferably less than 80 hours and even more preferably less than 70 hours, and specifically 66 hours or even lower.

20

Preferably the solid composition resulting from step b) is reconstituted to a liquid pharmaceutical composition in a further step c). Reconstituting the solid composition as of step b) favourably is possible in less than 5 min, less than 3 min, most favourably is less than 1 minute. For reconstitution physiological saline (0.9 wt% sodium chloride) can be used.

25

In yet another embodiment of the invention the lyophilisation of step b) can be replaced by spray-drying.

30

Further subject of the invention is therefore also a method of providing a composition or pharmaceutical according to the invention, wherein the composition or pharmaceutical is in the liquid state, comprising the step of solubilizing a composition according to the invention, wherein the starting composition is in the solid, preferably lyophilized or spray dried, state and wherein the starting composition is preferably at least in part amorphous. Solubilization of the solid, preferably lyophilized or spray dried, composition is preferably carried out with water, an aqueous solution of

35

dextrose or saline solutions.

40

The composition according to the invention, in particular the pharmaceutical, is preferably presented in unit dosage forms such as ampoules or disposable injection devices like syringes. It may also be presented in multi-dose forms such as a bottle or vial, from which the appropriate dose may be withdrawn. All such formulations should be sterile. In a preferred embodiment of the invention the ampoules, injection devices

- 21 -

and multi-dose forms contain the composition according to the invention, in particular the pharmaceutical, in solid, preferably lyophilized or spray dried, form, and the compositions are transformed into ready-to-use pharmaceuticals by solubilization of the compositions only shortly before their use.

5 The formulations according to the invention include those suitable for oral, rectal, topical, buccal (e.g. sub-lingual) and parenteral (e.g. subcutaneous, intramuscular, intradermal or intravenous) administration. It is preferred to present compositions of the present invention in the form of a pharmaceutical formulation for parenteral
10 administration, most preferable for any type of injection, in particular for intravenous, intraarterial, intralumbar, intraperitoneal, intramuscular, intradermal, subcutaneous or intraosseal injection.

Where the pharmaceutical formulation is for parenteral administration, the formulation may be an aqueous or non-aqueous solution or mixture of liquids, which may contain
15 bacteriostatic agents, antioxidants, buffers or other pharmaceutically acceptable additives. The preferred formulation of compositions of the present invention is either an aqueous acidic medium of pH 2-7, preferably 2-5 and more preferably 2-4 or an aqueous solution of a cyclodextrin. Cyclodextrins that can be used for these formulations are either the negatively charged sulfobutylether (SBE) derivatives of β -
20 CD, specifically SBE7- β -CD, marketed under the tradename Captisol by CyDex, Inc. (Critical Reviews in Therapeutic Drug Carrier Systems, 14 (1), 1-104 (1997)), or the hydroxypropyl CD's. The preferred method of formulation (i.e., acid buffer or CD-based) may depend on the physicochemical properties (e.g., aqueous solubility, pKa, etc.) of a particular composition. When the composition is in the solid, in particular
25 lyophilized, state, the solid is correspondingly preferably solubilized before its application in either an aqueous acidic medium preferably resulting in a pH 2-4 of the solution or in an aqueous solution of a cyclodextrin.

According to one aspect of the invention a solid pharmaceutical composition is provided. This composition can comprise 5 to 25% wt.% of remimazolam salt,
30 preferably besylate salt, preferably 8 to 23 wt.%, even more preferred 10 to 19 wt.%.

This composition can further comprise 75 to 95 wt.% of one or more hygroscopic excipients, preferably 77 to 92 wt.% and more preferably 81 to 90 wt.%. The
35 hygroscopic excipients preferably is a mixture of carbohydrates, comprising at up to 40% lactose, 38 wt.%, more preferably up to 33 wt.% disaccharide, preferably lactose. The rest of the mixture can be dextran.

In one embodiment the solid composition as outlined above contains no further excipients. In yet another embodiment the solid composition consists of remimazolam

- 22 -

salt, dextran and a dissaccharide (e.g. lactose) only. In yet another embodiment the sormulation consits only of remimazolam salt and lactose (this might be presented as a hydrate).

5 In another embodiment the composition is a liquid composition consisting of remimazolam, dextran, a disaccharide and a solvent, which preferably is physiological saline (0.9 wt.% sodium chloride). The pH value of such liquid (aqueous) composition, being preferably reconstituted from the solid composition, can range from about 3 to about 4, preferably from about 3.2 to about 3.3 and more preferably from 3.21 to 3.28.

10 Accordingly, the present invention also provides a method for producing sedation or hypnosis in a mammal, which comprises administering to the mammal an effective sedative or hypnotic amount of a pharmaceutical of the present invention as hereinbefore defined. The present invention also provides a method for inducing anxiolysis in a mammal, which comprises administering to the mammal an effective
15 anxiolytic amount of a pharmaceutical of the present invention as hereinbefore defined. The present invention also provides a method for inducing muscle relaxation in a mammal, which comprises administering to the mammal an effective muscle relaxant amount of a pharmaceutical of the present invention as hereinbefore defined. The present invention also provides a method for treating convulsions in a mammal,
20 which comprises administering to the mammal an effective anticonvulsant amount of a pharmaceutical of the present invention as hereinbefore defined. The present invention also provides a method for inducing or maintaining anesthesia in a mammal, which comprises administering to the mammal an effective anesthetic amount of a pharmaceutical of the present invention as hereinbefore defined.

25 The present invention also provides the use of a sedative or hypnotic amount of a composition of the present invention as hereinbefore defined in the manufacture of a medicament for producing sedation or hypnosis in a mammal, including in a human. The present invention also provides the use of an anxiolytic amount of a composition of the present invention as hereinbefore defined in the manufacture of a medicament
30 for producing anxiolysis in a mammal, including in a human. The present invention also provides the use of a muscle relaxant amount of a composition of the present invention as hereinbefore defined in the manufacture of a medicament for producing muscle relaxation in a mammal, including in a human. The present invention also provides the use of an anticonvulsant amount of a composition of the present
35 invention as hereinbefore defined in the manufacture of a medicament for treating convulsions in a mammal, including in a human. The present invention also provides the use of an anesthetic amount of a composition of the present invention as hereinbefore defined in the manufacture of a medicament for inducing or maintaining anesthesia in a mammal, including in a human.

- 23 -

The present invention also provides the use of a pharmaceutical according to the invention for producing sedation or hypnosis and/or inducing anxiolysis and/or inducing muscle relaxation and/or treating convulsions and/or inducing or maintaining anaesthesia in a mammal.

5 Intravenous administration can take the form of bolus injection or, more appropriately, continuous infusion. The dosage for each subject may vary, however, a suitable intravenous amount or dosage of the compounds of the present invention to obtain sedation or hypnosis in mammals would be 0.01 to 5.0 mg/kg of body weight, and more particularly, 0.02 to 0.5 mg/kg of body weight, the above being based on the weight of the compound which is the active ingredient (i.e. the weight of the benzodiazepine). A suitable intravenous amount or dosage of the compounds of the present invention to obtain anxiolysis in mammals would be 0.01 to 5.0 mg/kg of body weight, and more particularly, 0.02 to 0.5 mg/kg of body weight, the above being based on the weight of the compound which is the active ingredient. A suitable intravenous amount or dosage of the compounds of the present invention to obtain muscle relaxation in mammals would be 0.01 to 5.0 mg/kg of body weight, and more particularly, 0.02 to 0.5 mg/kg of body weight, the above being based on the weight of the compound which is the active ingredient. A suitable intravenous amount or dosage of the compounds of the present invention to treat convulsions in mammals would be 0.01 to 5.0 mg/kg of body weight, and more particularly, 0.02 to 0.5 mg/kg of body weight, the above being based on the weight of the compound which is the active ingredient. Thus a suitable pharmaceutical parenteral preparation for administration to humans will preferably contain 0.1 to 20 mg/ml of a compound of the present invention in solution or multiples thereof for multi-dose vials.

25 A yet another aspect of the invention relates to the use of a mixture of at least one disaccharide and at least dextran for preparing a solid composition comprising at least one benzodiazepine comprising at least one carboxylic acid ester moiety or a pharmaceutically acceptable salt thereof, which is preferably a remimazolam salt (particularly its besylate or tosylate salt). Preferably the mixture contains or consists of lactose and dextran, preferably a dextran with 80 kD or less (e.g. dextran 40 or dextran 70). The solid composition has a favourable reconstitution time.

Particularly, the invention relates to the following embodiments:


In the embodiment 1, the invention relates to a composition comprising at least one benzodiazepine comprising at least one carboxylic acid ester moiety or a pharmaceutically acceptable salt thereof, wherein the composition

35 a) comprises at least one pharmaceutically acceptable hygroscopic excipient, and/or

- 24 -

b) the composition is at least in part amorphous.

Embodiment 2 relates to a composition according to embodiment 1, wherein the benzodiazepine is a compound according to formula (I)



wherein

10 W is H, a C₁-C₄ branched or straight chain alkyl;

X is CH₂, NH, or NCH₃; n is 1 or 2;

Y is O or CH₂; m is 0 or 1;

Z is O; p is 0 or 1;

15 R¹ is a C₁-C₇ straight chain alkyl, a C₃-C₇ branched chain alkyl, a C₁-C₄ haloalkyl, a C₃-C₇ cycloalkyl, an aryl, a heteroaryl, an aralkyl, or a heteroaralkyl;

R² is phenyl, 2-halophenyl or 2-pyridyl,

R³ is H, Cl, Br, F, I, CF₃, or NO₂;

20 (1) R⁴ is H, a C₁-C₄ alkyl, or a dialkylaminoalkyl and R⁵ and R⁶ together represent a single oxygen or S atom which is linked to the diazepine ring by a double bond and p is zero or 1; or (2) R⁴ and R⁵ together is a double bond in the diazepine ring and R⁶ represents the group NHR⁷ wherein R⁷ is H, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, benzyl or benzyl mono or disubstituted independently with halogen substituents, C₁₋₄ alkylpyridyl or C₁₋₄ alkylimidazolyl and p is zero; or (3) R⁴, R⁵ and R⁶ form the group-CR⁸=U-V= wherein R⁸ is hydrogen, C₁₋₄ alkyl or C₁₋₃ hydroxyalkyl, U is N or CR⁹ wherein R⁹ is H, C₁₋₄ alkyl, C₁₋₃ hydroxyalkyl or C₁₋₄ alkoxy, V is N or CH and p is zero.

25

Embodiment 3 relates to a composition according to embodiment 2, wherein p is zero and R⁴, R⁵ and R⁶ form the group -CR⁸=U-V= wherein R⁸ is hydrogen, C₁₋₄ alkyl or C₁₋₃ hydroxyalkyl, U is N or CR⁹ wherein R⁹ is H, C₁₋₄ alkyl, C₁₋₃ hydroxyalkyl or C₁₋₄ alkoxy, V is N or CH.

30

Embodiment 4 relates to a composition according to embodiment 2 or 3, wherein W is H; X is CH₂, n is 1; Y is CH₂, m is 1;

R¹ is CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂ or CH₂CH(CH₃)₂;

- 25 -

R² is 2-fluorophenyl, 2-chlorophenyl or 2-pyridyl;
R³ is Cl or Br.

5 Embodiment 5 relates to a composition according to any of embodiments 2 to 4, wherein p is zero and R⁴, R⁵ and R⁶ form the group -CR⁸=U-V= wherein R⁸ is methyl, U is CH₂, V is N;
W is H; X is CH₂, n is 1; Y is CH₂, m is 1;
R₁ is CH₃; R₂ is 2-pyridyl; R₃ is Br.

10 Embodiment 6 relates to a composition according to any of the embodiments 1 to 5, wherein the benzodiazepine is in the form of a pharmaceutically acceptable salt.

15 Embodiment 7 relates to a composition according to any of the embodiments 1 to 6, wherein in the pharmaceutically acceptable salt the benzodiazepine is formulated in cationic form and the counter ion is selected from halogenides, in particular fluoride, chloride or bromide, sulfate, organic sulfates, sulfonate, organic sulfonates, nitrate, phosphate, salicylate, tartrate, citrate, maleate, formate, malonate, succinate, isethionate, lactobionate and sulfamate.

20 Embodiment 8 relates to a composition according to embodiment 7, wherein the counter ion is selected from organic sulfates and sulfonates, in particular aromatic sulfates and sulfonates.

25 Embodiment 9 relates to a composition according to embodiment 8, wherein the counter ion is benzene sulfonate (besylate).

Embodiment 10 relates to a composition according to embodiment 9, wherein the benzodiazepine salt is crystalline remimazolam besylate.

30 Embodiment 11 relates to a composition according to any of the embodiments 1 to 10, wherein the hygroscopic excipient is a compound which is able to form stable hydrates.

35 Embodiment 12 relates to a composition according to any of the embodiments 1 to 11, wherein the hygroscopic excipient is an organic substance, preferably selected from carbohydrates and/or organic polymers.

40 Embodiment 13 relates to a composition according to embodiment 12, wherein the hygroscopic excipient possesses a molecular weight of less than 150 kD.

- 26 -

Embodiment 14 relates to a composition according to embodiment 12 or 13, wherein the carbohydrate is a dextran molecule.

Embodiment 15 relates to a composition according to embodiment 12 or 13, wherein the carbohydrate is selected from monosaccharides and C₂₋₆-oligosaccharides.

Embodiment 16 relates to a composition according to embodiment 15, wherein the carbohydrate is a disaccharide, preferably selected from the group consisting of lactose, maltose, sucrose and trehalose.

Embodiment 17 relates to a composition according to embodiment 12, wherein the organic polymer is a polyvinylpyrrolidone and preferably possesses a molecular weight of between 5 and 40 kD.

Embodiment 18 relates to a composition according to any of the embodiments 1 to 17, wherein the wt.% ratio between the total amount of hygroscopic excipients and the total amount of benzodiazepines or salts thereof in the composition is at least 1:1, preferably at least 2:1, most preferably at least 5:1.

Embodiment 19 relates to a composition according to any of the embodiments 1 to 18, wherein the composition is in the solid state and is preferably a lyophilized solid.

I. STABILITY OF CNS7056; FORMULATIONS WITH SELECTED EXCIPIENTS

1. Formulations

A total of 11 formulations of the besylate salt of remimazolam with a selection of suitable excipients as detailed in Figure 1 and 2 were lyophilized. In addition a formulation containing the besylate salt of remimazolam alone and matching placebos for each formulation were also prepared (see Figure 3). In the following the abbreviation "REM" is used for the besylate salt of remimazolam.

Each formulation was prepared as follows and filled into vials prior to freeze drying: Excipient was dissolved in approximately 50 ml water. REM was added and stirred to dissolve. Once dissolved the pH of the solutions was adjusted to 3.10 ± 0.05 with 0.5 M hydrochloric acid/ 2 M sodium hydroxide. Placebo solutions and the solution containing REM alone were prepared in the same manner. Each solution was made up to 100 ml and 1.2 ml of each solution was aliquoted into 2 ml vials. The formulations were lyophilized using a Virtis Genesis 25 EL freeze dryer according to the following cycle:

- 27 -

Freezing steps

Step	Temperature (°C)	Time (minutes)	Pressure (mTorr)	Hold/Ramp
1	4	10	--	H
2	-45	490	--	R
3	-45	170	--	H

Drying steps

Step	Temperature (°C)	Time (minutes)	Pressure (mTorr)	Hold/Ramp
1	-45	10	100	H
2	-25	200	100	R
3	-25	3640	100	H
4	30	275	70	R
5	30	1300	70	H

- 5 After freeze drying the samples were stored in storage cabinets at 25°C/60%RH and at 40°C/75%RH for 13 weeks, respectively. ("RH" means relative humidity.)

Analysis10 a) Reconstitution time

After 13 weeks of storage the vials were reconstituted in duplicate with 1.2 ml of water for irrigation/injection and swirled gently to mix. The time taken for complete dissolution was recorded.

15 b) HPLC

For HPLC each vial was reconstituted with sample solvent (50/50 % v/v acetonitrile/water) and the contents transferred to a 25 ml volumetric flask (with exception of REM only vial, which was transferred to a 50 ml volumetric flask) with several rinsings. The dextran formulation was insoluble in sample solvent and was diluted in 100% water. For each formulation a placebo was also analysed in the same way. Analyses were performed in duplicate, unless otherwise stated.

Results25 a) Reconstitution time

Reconstitution time was acceptable for all samples.

b) HPLC

- 28 -

5 The investigation of the emergence of the hydrolysis degradant of REM (results are summarized in Figure 4), which is formed by hydrolysis of the ester bond, revealed that the sample with REM only as well as the sample containing mannitol, which is a commonly used excipient for lyophilization of pharmaceuticals, exhibited only poor stability of REM, showing a degradation after 13 weeks at storage conditions of 40 °C/75%RH of more than 8 %.

10 Samples containing glycine showed moderate degradation, whereas all samples containing hygroscopic excipients (carbohydrates or polyvinylpyrrolidone) showed good or excellent stability. In particular the samples containing carbohydrates (disaccharides or dextran) showed excellent stability, i.e., a degradation after 13 weeks at storage conditions of 40 °C/75%RH of less than 1 %.

15 The samples with differing amounts of lactose revealed that the greater the amount of carbohydrate relative to the amount of REM, the better the stability of the REM. Further by incorporating a carbohydrate (e.g. lactose) as a component of a formulation of CNS 7056 that is inherently unstable e.g. mannitol it is possible to improve the stability of this formulation.

20 **2. Stability data of CNS 7056: lactose based formulation batches after storage for up to 36 months**

2.1 Introduction

25 CNS 7056 is presented for clinical use as a sterile lyophilized powder for reconstitution in 20 mL vials with a Bromobutyl stopper, suitable for intravenous injection. Each vial contains 26 mg of CNS 7056. During development further batches with 25, 23 and 26 mg of CNS 7056 have been prepared. On reconstitution with a defined volume of Water for Injection, the concentration of the dose solution is 5mg/mL CNS7056. All these products contain the same CNS 7056 to lactose ratio in the lyophilized product (i.e. 1:13 CNS 7056: lactose monohydrate). Stability data were
30 collected for all intervals through the month shown in bold in the following Table 1:

Table 1: Summary of Stability studies for CNS 7056 batches

Summary of CNS 7056 for Injection on Stability			
Batch	Storage Conditions	Type of Batch	Test Interval (Months)
A01P310	25 °C/60% RH	GMP clinical batch , stability	0, 1, 3, 6, 9, 12, 18, 24, 36 ^c , 48 ^c
	30 °C/65% RH		0, 1, 3, 6, 9, 12
	40 °C/75% RH		0, 1, 3, 6
P310-01 (B)	25 °C/60% RH	Lab development batch, stability	0, 1, 3, 6, 9, 12, 18
	30 °C/65% RH		0, 6, 9, 12
	40 °C/75% RH		0, 1, 3, 6, 9, 12
025CNS27	25 °C/60% RH	Lab development batch, stability	0, 1, 2, 3, 6, 9, 12, 18 ^{ab}
	40 °C/75% RH		0, 1, 2, 3, 6, 12 ^b
026CNS27	25 °C/60% RH	Lab development batch, stability	0, 1, 2, 3, 6, 9, 12, 18 ^{ab}
	40 °C/75% RH		0, 1, 2, 3, 6 ^b
G384	25 °C/60% RH	Lab development batch, stability	0, 1, 2, 3, 6, 9, 12, 18 ^{ab}
	40 °C/75% RH		0, 1, 2, 3, 6 ^b
P02308	25 °C/60% RH	GMP clinical batch, stability	0, 1, 2, 3, 6, 9, 12, 18, 24, 36 ^c , 48 ^c
	40 °C/75% RH		0, 1, 2, 3, 6, 9 ^a , 12
^a = Stability test interval added. Test vials taken from remaining spare vials at specified temperature.			
^b = Stability studies on 025CNS27, 026C0NS27 and G384 are now complete			
^c = Optional time points			

2.1.1 CNS 7056 batch composition

5

Table 2 Composition of the different CNS7056 batches

Substance	Batch #					
	025CNS27	026CNS27	G384	P02308	P310-01(B)	A01P310
CNS7056 base	25 mg	25 mg	25 mg	23 mg	26 mg	26 mg
Lactose monohydrate	433 mg	433 mg	433 mg	398 mg	450.3 mg	450.3 mg
0.12 M NaOH/ 0.12 M HCl	qs to pH 3.1	qs to pH 3.1	qs to pH 3.1	qs to pH 3.1	qs to pH 3.1	qs to pH 3.1

qs = to add sufficiently quantity to

10

2.1.2 Freeze drying conditions

The freeze drying conditions for the batches are given in the following tables 3 to 7:

Table 3: Freeze drying cycle for batch 025CNS27:

- 30 -

Step	Process	Shelf temperature (°C)	Ramp rate (°C/min)	Hold time (min)	Pressure (mTorr)
1	Load	4	0	30	n/a
2	Freezing	-45	0.1	180	n/a
3	Primary drying	-25	0.1	1700	100
4	Secondary drying	30	0.2	1300	75
5	Finish	Vials stoppered under 95% pure nitrogen			

Table 4: Freeze drying cycle for batch 026CNS27:

Step	Process	Shelf temperature (°C)	Ramp rate (°C/min)	Hold time (min)	Pressure (mTorr)
1	Load	4	0	120	n/a
2	Freezing	-45	0.1	300	n/a
3	Primary drying	-30	0.1	2885	100
4	Primary drying	-25	0.2	4100	100
4	Secondary drying	30	0.2	1580	75
5	Finish	Vials stoppered under 95% pure nitrogen			

5

Table 5: Freeze drying cycle for batch G384:

Step	Process	Shelf temperature (°C)	Ramp rate (°C/min)	Hold time (min)	Pressure (mTorr)
1	Load	4	0	10	n/a
2	Freezing	-45	0.1	300	n/a
3	Primary drying	-25	0.1	3640	100
4	Secondary drying	30	0.2	1125	70
5	Finish	Vials stoppered under 95% pure nitrogen			

Table 6: Freeze drying cycle for batch P02308:

Step	Process	Shelf temperature (°C)	Ramp rate (°C/min)	Hold time (min)	Pressure (mTorr)
1	Load	4	0	60	n/a
2	Freezing	-45	0.1	180	n/a
3	Primary drying	-25	0.1	3640	100
4	Secondary drying	30	0.2	1300	75
5	Finish	Vials stoppered under 95% pure nitrogen			

10

Table 7: Freeze drying cycle for batch A01P310:

Step	Process	Shelf temperature (°C)	Ramp rate (°C/min)	Hold time (min)	Pressure (mTorr)
1	Load	4	0	60	n/a
2	Freezing	-45	0.1	210 ^a	n/a
3	Primary drying	-25	0.1	3640	100
4	Secondary drying	30	0.2	1300	75
5	Finish	Vials stoppered under 95% pure nitrogen			
^a = Includes 30 min for condenser preparation					

2.2 Methods of analysis

2.2.1 Appearance of Lyophilized Product

The same CNS 7056 vials (6 at each storage condition) were inspected visually, the appearance recorded and the vials placed back on storage. A comparison was also made to a set of vials that had been stored in the same secondary packaging at 2 to 8°C to assess whether there were any differences (in particular in colour) between these controls and those stored at elevated temperatures.

2.2.2 CNS7056 vial content, concentration on reconstitution and related substances

The CNS7056 assay and related substances determination was determined by HPLC). For this purpose, the appropriate volume of WFI was added to each vial and swirled until complete dissolution was achieved. The seal and the stopper were carefully removed and the stopper was rinsed thoroughly into a 100 ml volumetric flask. The contents of the vial with washings of diluent were transferred to a volumetric flask. The diluent was added to reach a volume of 100 mL (equals a concentration of 0.23, 0.25 or 0.26 mg/mL, respectively). The sample were analysed by HPLC by using the following conditions:

Column: YMC ODS-AQ, 250 x 4.6 mm, 3 µm particle size
Mobile phase: Res A: 0.01% trifluoroacetic acid in water
Res B: 0.01% trifluoroacetic acid in acetonitrile

Gradient:

Time (mins)	%A	%B
0	75	25
20.0	60	40
30.0	20	80
32.0	20	80

32.5	75	25
40.0	75	25

Flow rate: 1.0 ml/min
Column temperature: 40 °C
Autosampler: ambient
Detection: UV at 230 nm
Injection volume: 10µl
Run time: 40 min

The retention time for CNS7056 is about 15 minutes. The CNS 7056 content was assayed by comparison with similarly chromatographed reference solutions. Related substances were determined by normalised area %.

The concentration of the reconstituted solution is calculated by the following equation:

$$\text{CNS7056 base (mg)} = \left(\frac{\text{sample peak area}}{\text{mean peak area std}_1 \text{ throughout run}} \right) \times \frac{\text{Wt std}_1}{50} \times \frac{\text{MWt CNS 7056 base}}{\text{MWt CNS 7056B}} \times \frac{P}{100} \times \text{DF}$$

where

Wt std 1 is the weight of CNS 7056B reference material used to prepared standard 1 (mg)

MWt CNS 7056 base is the molecular weight of the free base of CNS 7056 = 439.3

MWt CNS 7056B is the molecular weight of CNS7056 besylate salt = 597.5

P is the declared assay as per the C of A for the reference standard

DF is the dilution factor

The vial content is calculated according to the following formula:

$$\text{CNS7056 base (mg)} = \left(\frac{\text{sample peak area}}{\text{mean peak area std}_1 \text{ throughout run}} \right) \times \frac{\text{Wt std}_1}{50} \times \frac{\text{MWt CNS 7056 base}}{\text{MWt CNS 7056B}} \times \frac{P}{100} \times \text{DF}$$

where

Wt std 1 is the weight of CNS 7056B reference material used to prepared standard 1 (mg)

MWt CNS 7056 base is the molecular weight of the free base of CNS 7056 = 439.3

MWt CNS 7056B is the molecular weight of CNS7056 besylate salt = 597.5

P is the declared assay as per the C of A for the reference standard

DF is the dilution factor

For determination of related substances CNS7056 is identified by comparison of the retention time to that of CNS7056 in the reference standard chromatograms. The

- 33 -

amount of each individual detected related substance is calculated as area percent for each sample injection according to the following formula:

$$\text{Area \%} = \left(\frac{A}{T} \right) \times 100$$

Where, A = area of the related substance peak
T = total area of all peaks in the chromatogram

5

2.2.3 Chiral purity

The chiral purity of CNS 7056 was determined by HPLC by using the following conditions:

Column: Chiralpak IC, 250 x 4.6 mm, 5µm particle size
Mobile phase: phosphate buffer pH 7.0/water/acetonitrile 10/40/50, v/v/v
Sample solvent: water/acetonitrile 50/50, v/v
Flow rate: 0.7 ml/min
Column temperature: 40°C
Autosampler: ambient
Detection: UV at 250 nm
Injection volume: 10 µl
Run time: 35 minutes

15

20

The retention time for the CNS 7056 S-enantiomer is about 21.3 min and for CNS 7056 R-enantiomer is about 17.8 min (RRT = 0.84). The chiral purity is calculated according to the following formula:

$$\text{Area \%} = \left(\frac{A}{T} \right) \times 100$$

where A = area of R - CNS 7056 peak
T = total area of the CNS 7056 and the R - CNS 7056 peaks

25

2.2.4 Volume of solution in vial following reconstitution (Ph.Eur 2.9.17)

A single vial was reconstituted with 5.0 ml of Water for Injection (WFI), Ph. Eur. using a 5 ml BD syringe fitted with a suitable needle. When fully reconstituted, the contents were removed using a syringe and 21 gauge needle and transferred to a calibrated 10 ml measuring cylinder.

30

2.2.5 Appearance of reconstitution

The appearance of the solution following reconstitution was recorded.

- 34 -

2.2.6 Reconstitution time

Two vials were reconstituted with 5.0 mL of Water for Injection (WFI), Ph. Eur., using a 5 ml BD syringe and appropriate needle, and the time taken to fully dissolve recorded.

2.2.7 pH value

The pH was determined on two reconstituted solutions following addition of 5.0 mL Water for Injection (WFI), Ph. Eur., using a 5 ml BD syringe fitted with a suitable needle. The pH was measured on one aliquot from each of the two vials.

2.2.8 Osmolality

The osmolality was determined on the two reconstituted solutions following addition of 5.0 mL Water for Injection (WFI), Ph. Eur., using a 5 ml BD syringe fitted with a suitable needle. Osmolality was measured on one aliquot from each of the two vials by freezing point depression with reference to a solution of known Osmolality. For this purpose 100µl of the reconstituted CNS 7056 solution is measured in a freezing point depression osmometer.

2.2.9 Water content

The water content was determined by coulometric Karl Fischer titration. The moisture content of vials of CNS 7056 drug product is determined by dissolving the entire contents of a vial of CNS 7056 lyophilised powder in anhydrous dimethylformamide (DMF) and injecting a known volume of the solution into the anolyte of a coulometric Karl Fischer apparatus. In the Karl Fischer reaction, water reacts in a 1:1 ratio with iodine. The amount of water is determined by measuring the number of coulombs of electricity required to oxidise iodide ions to the iodine required for the Karl Fischer reaction. The number of Coulombs is used to calculate the amount of water titrated in µg, which is displayed by the apparatus.

The following equipment and reagents were used:

Karl-Fischer titratus apparatus:	Mitsubishi CA-100
Anolyte:	Hydranal Coulomat AG
Catholyte:	Hydranal Coulomat CG

The water content of CNS7056 lyophilised powder is calculated according the following formula:

- 35 -

$$\text{Moisture per vial (mg)} = \frac{(\text{M}_{\text{sample}} - \text{M}_{\text{solvent}})}{1000} \times \frac{(\text{W}_{\text{solvent}}/\text{D}_{\text{solvent}})}{\text{V}_{\text{titration}}}$$

$$\text{Moisture, \%w/w} = \frac{\text{Moisture per vial} \times 100}{(\text{S}_{\text{vial}})}$$

Where:

- M_{sample} = amount of water in the sample solution added to the titration vessel (μg)
 $\text{M}_{\text{solvent}}$ = mean amount of water in the solvent blanks added to the titration vessel (μg)
 $\text{W}_{\text{solvent}}$ = weight of DMF added to the vial (g)
 $\text{D}_{\text{solvent}}$ = density of the solvent (g/ml)
 For DMF $d=0.944$ g/ml, source CRC handbook 81st edition
 $\text{V}_{\text{titration}}$ = volume of solution added to the titration vessel (ml)
 S_{vial} = Calculated total weight of solid per vial, including water (mg)

2.2.10 ID by UV

Analysis was performed in duplicate on a single vial. The identification by UV was confirmed by comparison of the drug product spectra to reference spectra.

2.2.11 Sub visible particles (EP 2.9.19)

Ten vials were reconstituted with 5 mL WFI using an appropriate sterile syringe and needle. The vials were pooled together under aseptic conditions and analysed according to European Pharmacopeia 2.9.19.

2.2.12 Sub visible particles (EP 2.9.19)

Ten vials were reconstituted with 5 mL WFI using an appropriate sterile syringe and needle. The vials were pooled together under aseptic conditions and analysed according to European Pharmacopeia 2.9.19.

2.2.13 Bacterial Endotoxin

Bacterial Endotoxin was determined by the Limulus amoebocyte lysate (LAL) gel-clot method as a limit test with a limit of < 0.5 EU/mg. For this purpose a LAL with declared sensitivity equal to 0.03 EU/ml is used. The Endotoxins are quantified using the following formula:

$$\text{Endotoxin concentration in the sample (EU/mg)} = \frac{\text{lysate sensitivity } (\lambda) \times \text{test dilution factor}}{\text{sample concentration}}$$

where:

- lysate sensitivity = 0.03 EU/ml
 sample concentration = 5 mg/ml

2.2.14 Sterility

- 36 -

5 Sterility was determined by reconstituting the lyophilised CNS 7056 with 5 ml of sterile peptonate water (0.1%) each and incubating the samples in 100 ml of thioglycollate medium THG) at 30 to 35°C and 100 ml of tryptic soy broth (TSB) at 20 to 25°C. The incubations were performed for not less than 14 days. The media are visually inspected every 2 to 3 days for the presence of microbiological proliferation. If there is no microbial growth, the examined sample meets the test requirements (sterile).

2.3 Results

10 The results of the stability analyses for the above described batches after storage at 25°C/60% relative humidity(RH) or at 40°C/75%RH (so called “accelerated stability” analysis) are summarized in the Figures 11 to 36.

2.4 Summary

15 The tested formulations for CNS 7056 exhibit an excellent long term stability which already supports a shelf life of 36 month for the drug product.

3. Stability data after storage for 36 month

20 The CNS 7056 batch P02308 was subjected to a stability study whereby the vials were stored for 36 months at 25°C/60%RH.

For batch composition and freeze drying conditions see chapter 2.1. For description of the analytical methods see chapter 2.2.

3.1 Results

25 The results of the stability analysis for the batch P02308 after storage at 25°C/60% relative humidity (RH) up to and including 36 months are summarized in figures 27-30.

3.2 Summary

30 All tests performed on batch P02308 after storage at 25°C/60%RH (T=36 months) were within the specified acceptance criteria. Appearance of Lyophilised Product, Completeness of Solution, Time to Reconstitute, pH, and Osmolality of all samples at T=36 months were well within specification.

35 CNS7056B Vial Content at 25°C/60%RH is 23.4 mg/vial, which is in keeping with all previous results. These results are well within specification. The main CNS 7056 hydrolysis product CNS7054X (RRT 0.59) has increased to 0.29% at 25°C/60%RH from 0.07% at initial months.

40 A total impurities result of 0.80% was observed at the T=36 month time point, compared to 0.65% at initial.. These results, together with the supporting data from

- 37 -

storage over 12 months at the accelerated stability storage condition of 40°C/75%RH, reflect only a slight increase in degradation over this significant period of time and demonstrate the stabilising effect of CNS 7056 in combination with lactose. .

5 Moisture content at T=36 months 25°C/60%RH is 0.68% which shows an increase from 0.27% at initial. This increase is thought to be due to water desorption from the stopper, which will occur over time. These results are well within specification.

3.3 Conclusion

10 All parameters are within specification and the only noticeable trends are the expected increase in the hydrolysis product CNS 7054X and moisture content. The rate of increase of CNS 7054X is similar to previous laboratory non GMP development batches of CNS 7056 for injection produced/tested.

15 4. Evaluation of crystalline material in a lyophilised lactose formulation of CNS 7056 by Raman mapping

4.1 Introduction

20 XRPD studies showed that the lyophilised formulation of CNS 7056 is amorphous, however when material is examined under polarised light microscopy there is evidence of crystalline material present in the amorphous matrix. In order to reveal if this crystalline material is due to CNS 7056 or some other component e.g. lactose monohydrate a Raman mapping analysis was performed. This study makes use of a confocal Raman dispersive microscope to study the physical form of CNS 7056 within
25 the lyophilized lactose formulation using Raman mapping. In Raman mapping experiments, once the first Raman spectrum collection is completed from the in focus sample surface, the sample stage is moved at a predefined step (normally a few to few tens micron) and another spectrum is taken. This is continued until the chosen analytical area has been covered and a hyperspectral data set has been constructed.
30 The sample is prepared to ensure its surface is flat as this avoids the need to refocus the microscope during the data collection from one point to another. The hyperspectral data cube is then processed to generate chemical images based on the distinguishable specific Raman peak (fingerprint) of each component of the sample under study. The chemical images thus generated can then establish each
35 component variation over the chosen area of the studied sample. Crystalline (Form I polymorph), amorphous (lyophilized) CNS 7056B and the lyophilized (amorphous) lactose were characterized by Raman, and the characteristic Raman peak of crystalline CNS7056B were used to generate the chemical images of crystalline CNS7056B. The chemical images of the lactose were also generated based on its
40 own characteristic peak. One area of a lyophilized formulation of CNS 7056B was mapped to determine, if present, the content (based on area ratio without calibration)

- 38 -

and distribution of crystalline (Form I) CNS 7056B within the mapping area of the lyophilized lactose formulation. The aim of this study was to establish whether crystalline material within the lactose formulation is due to CNS 7056B or some other component eg. lactose monohydrate.

5

4.2. Material and methods

The following samples were tested in the Raman mapping study:

CNS 7056B (Form I)

- 10 – Item/Lot Number: SOL 12621/5
- Appearance: white powder
- Pharmaterials Ref no: PMO553/08

CNS 7056 for Injection (received from Paion)

- 15 – Item/Lot Number: PO2308
- Appearance: white lyophilised powder
- Pharmaterials Ref no: PM0554/08

Lyophilized CNS 7056 (amorphous)

- 20 – Item/Lot Number: 05/CNS/06
- Appearance: white lyophilised powder
- Pharmaterials Ref no: PM0555/O8

Lyophilized lactose (received from Paion)

- 25 – Item/Lot Number: O28/CNS/27
- Appearance: white lyophilized powder
- Pharmaterials Ref no: PM0548/08

Raman spectra of supplied materials

30 Raman spectroscopy on crystalline (Form I) and amorphous (lyophilized) CNS7056B and amorphous lactose (lyophilised) as supplied was performed using a confocal Nicolet Almega XR dispersive Raman microscope. A distinguishable Raman peak of crystalline (Form I) CNS7056B and amorphous lactose (lyophilised) was respectively selected for generating chemical images and establishing each variation in the
35 examined area of a lyophilized formulation as shown later.

Raman mapping using a confocal dispersive Raman microscope

40 Raman mapping was performed on one area of a lyophilised formulation. For each measurement, Raman mapping was performed on one area (e.g. 300x300 μm^2). The chemical images were then produced, respectively based on the distinguishable Raman peak of crystalline CNS7056B and amorphous lactose (lyophilised). These

- 39 -

operations allowed the identification of crystalline CNS7056B (potentially recrystallized from the lyophilized lactose formulation) and amorphous lactose (lyophilized) in the selected area of the sample. Subsequently, thus produced chemical images were used to indicate the distribution of crystalline CNS7056B and lactose (lyophilised) in the mapping area respectively.

Raman technique

Raman spectra

Samples were analysed by a confocal Nicolet Almega XR Dispersive Raman Microscope for its Raman spectrum using the following conditions:

- Exposure Time: 1.0s
- Exposure Times of each spectrum: 10
- Pinhole Size: 100um
- Spectral range: whole (single grating)
- Laser: He Ne 633nm at 100% power
- Objective: 50x/0.75 (magnifier/numerical aperture number)

Afterwards, the measured Raman spectra were corrected by baseline subtraction (BS) using the software OMNICTM v7.3.

Raman mapping

Each sample was gently pressed by hand so that the mapping area has an approximately flat surface. Raman spectra data for mapping were collected using following conditions:

- Exposure Time: 5.0s
- Exposure Times: 10
- Pinhole Size: 100pm
- Wavelength range: 1700~300 cm^{-1} (multiple gratings)
- Laser: He Ne 633nm (100% power)
- Objective: 50x/0.75
- Area: circa 300 x 300 μm
- Scanning step: 10pm

Then the measured Raman spectra data from mapping were modified by baseline correction and normalization using the software OMNICTM v7.3.

4.3 Results and Discussion

4.3.1 Raman spectra of each component in a lyophilized formulation

- 40 -

The Raman spectra of crystalline (Form I polymorph) and amorphous (lyophilized) CNS7056B and amorphous lactose (lyophilised) as supplied were collected using the procedure described under Material and Methods. As seen in Figure 39, the Raman peak was then selected respectively: circa 1620 cm^{-1} for crystalline CNS7056B and circa 365 cm^{-1} for lyophilized lactose. These peaks are unique to both materials so that the chemical images of crystalline CNS7056B and lactose (lyophilized) can respectively be generated. The whole range of the Raman spectrum for each component contained in a lyophilized formulation is given in Figures 40 and 41.

4.4 Summary

The obtained data demonstrate the presence of crystalline CNS7056B in this lyophilized formulation comprising mostly of amorphous CNS7056B. Furthermore a uniform distribution of CNS7056B and excipient in this lyophilized formulation could be shown.

In the lyophilised product tested approximately 9% of the data points contained a signal corresponding to crystalline CNS 7056B. The actual w/w (%) presence of the crystalline CNS 7056B in the amorphous lactose matrix, however, cannot be concluded from these results as calibration was not performed.

II. PREPARATION AND STABILITY ASSESSMENTS ANALYSIS OF LYOPHILIZED AND SPRAY-DRIED FORMULATIONS

A lyophilised and a spray-dried formulation having the same formulation were prepared and tested for stability.

1. Manufacture of spray-dried CNS7056B formulation (with lactose)

CNS7056B (Form 2 bn 10201126, 5.1g) and Emprove Lactose Monohydrate, (139.2g) were dissolved in 750ml deionized (DI) water with heating to $\sim 50^{\circ}\text{C}$, and then filtered and cooled to room temperature. The pH was checked and not adjusted as it was at 3.1. This solution was spray-dried using the following parameters: Inlet temperature 150°C . Pump rate = 10% (20ml in 7 mins), Fan setting = 50%. Yield 59.5g. The water content was measured via Karl-Fischer and used to calculate the fill weight per vial (997mg). 58 vials were filled with 997mg of spray-dried formulation. 30 vials were placed in the vacuum oven with lids slightly open. These vials were dried under vacuum ($\sim 250\text{psi}$) with a nitrogen bleed at 50°C for 24 hours. The chamber was then flushed with nitrogen, and the vials were then closed quickly under nitrogen. These samples were called 12PM529-8-2. 28 vials were closed without drying. These samples were called 12PM529-8-1. All vials were crimped with aluminium seals.

2. Scale up of spray dried API (CNS7056B)

CNS7056B (Form 2 bn 10201126, 20g) was dissolved in 2900ml DI water. This solution was filtered and then spray-dried using the following parameters: Inlet temperature 130°C, outlet temp. 82-79°C. Pump rate = 10% (20ml in 7 minutes), Fan setting = 30%. Yield not noted. The process was repeated with CNS7056B (Form 2 bn 10201126, 5.6g) was dissolved in 812ml DI water, to give 2.2g overall yield (from both runs) of a white powder. The samples were called 12PM529-9-1.

3. Manufacture of freeze dried (lyophilized) API (CNS7056B)

A solution of CNS7056B in water was prepared (2.2g of bn 10201126, Form 2, PM0232/12 in 230ml water). This was placed in a round bottomed flask (rbf) and 'shell-frozen' in liquid nitrogen and then lyophilised over 5 days. The resulting fluffy white solid was scraped out and broken up (~2g). The samples were designated as 12PM529-10-1.

4. Accelerated stability study on lyophilized and spray dried formulation and spray dried API

The spray-dried CNS7056B formulation, both dried (12PM529-8-2) and non-dried (12PM529-8-1), stored in crimped vials was placed on an accelerated stability study, along with the lyophilized CNS7056B formulation (CNS2501A) as reference, and with the spray-dried amorphous API (12PM529-9-1). Samples were stored at 40°C/75%RH for 4 and 13 weeks, and at 55°C for 4 weeks, and analysed for appearance, assay, related substances, moisture, XRPD, reconstitution time, and appearance following reconstitution.

4.1. Results

The results from the stability study are presented in the Figures 42 to 44 and can be summarized as follows:

The spray-dried formulation (sealed prior to additional drying) had a slightly higher initial total impurity level at t=0 than the lyophilised formulation CNS 2501A i.e. batch (~0.73/0.67% vs 0.48%). This is potentially due to the manufacturing process involving higher temperature, and could be optimised on scale up.

The vacuum dried spray-dried formulation sample (12PM529-8-2) had similar water content to the lyophilised sample supplied (0.24% vs 0.34%). The non-dried spray-dried formulation sample (12PM529-8-1) had significantly higher water content (2.87%), as did the amorphous spray-dried API (CNS7056B, 12PM529-9-1).

The 'dry' spray-dried formulation (12PM529-8-2) showed similar stability to the lyophilized formulation. The total impurities increased ~0.2% for both samples after 4

- 42 -

weeks (slightly more at 55°C than at 40/75), and actually only increased ~0.05% for both samples after 13 weeks at 40/75.

5 The 'wet' spray-dried formulation (12PM529-8-1) had slightly inferior stability than the other formulation samples, but was still within specification for impurities after 13 weeks at 40/75 (total impurities increased from 0.67% at t=0, to 1.33% at t=13 weeks).

10 The spray-dried API (CNS7056B, 12PM529-9-1) showed significant instability, with increase of total impurities to 1.94% (4 weeks at 55°C), 2.56% (4 weeks at 40/75) and 3.35% (13 weeks at 40/75). This confirms that the lactose formulation is stabilising the API significantly during the stability trial, even when there are similar levels of water present in the formulation to the API sample.

15 As expected the major impurity that was observed was the hydrolysis product CNS7054X.

5. Investigation of API distribution and form in the spray-dried and lyophilized formulations using Raman mapping

20 A vial of lyophilized formulated product (CNS7056B in lactose, batch number CNS2501A) was opened and sampled randomly four times. Each sampled portion was then presented on a microscope slide and Raman mapping was carried out over a small area of the surface of the formulation sample (~300x300µm). The data was processed in comparison with reference samples of lyophilized (amorphous) and
25 crystalline API (CNS7056B, forms 1 and 2) and lyophilized (amorphous) and crystalline (monohydrate) lactose. The mapping was analysed to determine the distribution of the API within the formulation, and then if any phase separation (regions of API) was detected, these would be analysed to assess the physical form of the API. A second experiment was carried out where a new vial of lyophilized
30 formulated product (CNS7056B in lactose, batch number CNS2501A) was opened and sampled from top, middle and bottom of the cake. These three samples were again analysed by Raman mapping as above. Also, two regions from the top and bottom samples were mapped over a smaller region (~20x20µm) in more detail. Three more batches of lyophilized formulated product (CNS7056B in lactose) were
35 also analysed by Raman mapping: batches P02308, A01P301 and P301-02N. The size of region mapped in these experiments was ~120x100µm.

5.1 Results for batch CNS2501 A

40 The Raman mapping data was processed to give a 'chemical image' which shows the similarity of the Raman spectra detected at each point on the map with:
a) the excipient main peak at 355cm⁻¹ (i.e. lactose)

- 43 -

- b) the API (CNS7056B) main peak at 1580cm⁻¹
- c) correlation with the whole excipient spectra (lactose).

The data showed that no phase separation and re-crystallization of API was found in batch CNS2501A as supplied after analysis of 7 different grab-samples from 2 different vials. The distribution of API and lactose was uniform and no separate regions or particles of API were found. This suggests that a true molecular dispersion of the API in lactose has been formed in the lyophilised formulation batch CNS2501A.

5.2 Results for batches P02308, A01P301 and P301-02N

The Raman mapping data was processed as described in chapter 5.1.

The data revealed no phase separation and re-crystallization of API in batches P02308, A01P301 and P301-02N as supplied based on one set of mapping data for each batch. The distribution of API and lactose was uniform and no separate regions or particles of API were found. This suggests that a molecular dispersion of the API in lactose has been formed in the lyophilised formulation in batches P02308, A01P301 and P301-02N. (Note: some phase separation was found in previous mapping performed on batch P02308. This suggests that the distribution of separated (crystalline) API in this batch is not uniform.

6. Summary

An equivalent spray dried formulation to the current lyophilised (freeze-dried) product could be successfully developed and tested.

Both the spray dried and lyophilised formulations were shown to be fully amorphous and single phase by XRPD and Raman analysis (i.e. no detectable separated crystalline API). The spray dried formulation had a slightly higher impurity level (~0.7% total impurities vs. ~0.5% for the lyo product). This is presumed to be formed during spray-drying manufacture and could be reduced with process optimisation.

The fully dried spray-dried formulation showed equivalent stability to the lyo product over 13 weeks at 40°C/75%RH and 4 weeks at 55°C. The non-dried spray-dried formulation (3% water) showed slightly worse stability, but stayed within the specification over 13 weeks at 40°C/75%RH.

The spray-dried formulation showed similar colour change to the lyo product in the light stability trial, both turning grey/blue. Physical analysis of the light stressed samples of API and formulation showed some recrystallisation and absorption of water, but no evidence of changes in physical form contributing to the colour changes.

Raman mapping analysis of the current lyo batch (CNS2501A) and previous lyo batches (P02308, A01P301 and P301-02N) of formulated product showed uniform distribution of API and excipients, with no evidence of separation of API and subsequent crystallisation.

III. PREPARATION AND STABILITY ANALYSIS OF DISACCHARIDE BINARY EXCIPIENT CONTAINING FORMULATIONS

1. Purpose and study outline

The purpose of the present study was to evaluate the stability of selected formulations. Several lyophilized formulations of CNS 7056 were prepared containing lactose monohydrate and the pH adjusted to 3.1, the API is present as besylate salt. Two fill concentrations of CNS 7056 were investigated: 5 mg/ml and 10 mg/ml. The formulations were filled in ISO 10R and ISO 6R clear glass vials. Fill volume was reduced to 4 mL/vial (current fill volume is 5.2 mL). The existing formulation was filled in ISO 10R vials that were stoppered with both West 4023/50 art. 1346 stoppers and West S87 J 4416/50 stoppers. The stability of the new formulations manufactured in ISO 10R vials were evaluated together with the existing formulation. In addition to this, the existing formulation lyophilized in the frame of the last clinical batch manufacturing (batch number A01P310, fill volume 5.2 mL, ISO 20R clear glass vial) was tested to generate comparative stability data.

2. Methods

The following tests were performed on the stability samples:

- Appearance of the lyophylisate.
- Reconstitution time.
- Appearance of the reconstituted solution.
- Moisture content by Karl Fischer titration.
- HPLC Assay/Related Substances.
- Osmolality (only at time 0)

3. Batch description

The product composition of the batches submitted to stability is summarised here below.

Formulation (Excipient weight ratio)	CNS 7056 concentration	Vials	Fill volume	Stoppers	Product Reference
Lactose Reference	5 mg/mL	6R	4mL	West 4023/50 art. 1346	L6R5

(Current) Formulation		10R	4mL	West 4023/50 art. 1346	L10R5
				West S87 J 4416/50	L10R5S87
		20R	5.2mL	West 4023/50 art. 1346	L20R5
	10 mg/mL	6R	4mL	West 4023/50 art. 1346	L6R10
		10R	4mL	West 4023/50 art. 1346	L10R10
Lactose : Mannitol (4:1)	5 mg/mL	6R	4mL	West 4023/50 art. 1346	L4M16R5
		10R	4mL	West 4023/50 art. 1346	L4M110R5
	10 mg/mL	6R	4mL	West 4023/50 art. 1346	L4M16R10
		10R	4mL	West 4023/50 art. 1346	L4M110R10
Lactose : Mannitol (2:1)	5 mg/mL	6R	4mL	West 4023/50 art. 1346	L2M16R5
		10R	4mL	West 4023/50 art. 1346	L2M110R5
	10 mg/mL	6R	4mL	West 4023/50 art. 1346	L2M16R10
		10R	4mL	West 4023/50 art. 1346	L2M110R10

4. Stability program

The stability program is summarized in the following table:

Formulation (Excipient weight ratio)	Product Reference	Stability
Lactose Reference (Current) Formulation	L6R5	1 month
	L10R5	3 months
	L10R5S87	3 months
	L20R5	3 months
	L6R10	1 month
	L10R10	1 month
Lactose 4 : Mannitol 1	L4M16R5	1 month

Lactose 2 : Mannitol 1	L4M110R5	1 month
	L4M16R10	1 month
	L4M110R10	1 month
	L2M16R5	1 month
	L2M110R5	1 month
	L2M16R10	1 month
	L2M110R10	1 month

5. Stability schedules

The stability schedules are summarized in the following tables:

Storage conditions 40°C±2 °C / 75%±5% RH and 55°C±5 °C		
Tests for 1 month study	Time 0	1M
Lyo appearance (to be noted on all 5 vials)	√	√
Reconstitution time	√	√
Appearance of reconstituted solution	√	√
Moisture content (KF titration)	√	√
HPLC (Assay/Related Substances)	√	√
Osmolality	√	—

Storage conditions 55°C±5 °C		
Tests for 3 months study	Time 0	1M
Lyo appearance (to be noted on all 5 vials)	√	√
Reconstitution time	√	√
Appearance of reconstituted solution	√	√
Moisture content (KF titration)	√	√
HPLC (Assay/Related Substances)	√	√
Osmolality	√	—

Storage conditions 25°C±2 °C / 60%±5% RH			
Tests for 3 months study	Time 0	1M	3M
Lyo appearance (to be noted on all 5 vials)	√	—	√
Reconstitution time	√	—	√
Appearance of reconstituted solution	√	—	√
Moisture content (KF titration)	√	—	√
HPLC (Assay/Related Substances)	√	—	√
Osmolality	√	—	—

Storage conditions 40°C±2 °C / 75%±5% RH
--

- 47 -

Tests for 3 months study	Time 0	1M	3M
Lyo appearance (to be noted on all 5 vials)	√	√	√
Reconstitution time	√	√	√
Appearance of reconstituted solution	√	√	√
Moisture content (KF titration)	√	√	√
HPLC (Assay/Related Substances)	√	√	√
Osmolality	√	—	—

6. Stability results

The results collected in the frame of the present study are presented in Figures 45 to 51 and can be summarised as follows:

Samples stored at 40°C 75% RH (1 month).

- Some changes in the appearance of the lyo cake in the formulations L2M110R5 and L2M110R10. Some vials of L20R5 (clinical batch vials rejected after visual inspection) showed a different lyo cake appearance
- Expected increase of moisture content (not observed in L20R5; L2M110R10)
- Small increase in total impurities (not observed in L10R10; L10R5S87; L20R5). The HPLC assay kept practically constant.
- Increase of known impurity CNS7054X.

Samples after 3 months storage at 40°C 75% RH (only L10R5; L10R5S87 and L20R5 formulations).

The appearance of both the cake and the reconstituted solutions didn't undergo any variation (some of the cakes of the L20R5 samples were found to be shrunk).

The assay remained unvaried.

L10R5

- Further increase of moisture content (anyway the % H₂O<1.0%).
- Further slight increase of the impurities due to the CNS7054X.

L10R5S87

- Further increase of moisture content (anyway the % H₂O<1.0%).
- Further slight increase of the impurities content due to CNS7054X.

L20R5 (visual inspection)

- Further increase of moisture content (anyway the % H₂O<1.0%).
- Increase of the impurities content mainly due to CNS7054X.

- 48 -

Samples stored at 55°C (1 month).

- The lyo cake of the formulations L2M110R5, L2M110R10, L4M110R5 and L4M110R10 was found to be shrunk and yellowish colored. Some vials of L20R5 (clinical batch vials rejected after visual inspection) showed a charred (insoluble) lyo cake.
- Increase of moisture content (not observed in L20R5)
- Increase in total impurities (total impurities below 1.00% in L10R5; L10R10). Concurrently negligible reduction of the HPLC assay.
- Increase of known impurity CNS7054X.
- Additional impurities exceeding the LOQ in L20R5, L4M110R5; L4M110R10; L2M110R5; L2M110R10
- Slight presence of foam (not persistent) upon reconstitution of.
- L10R5; L10R5S87 and L20R5 formulations after 1 month storage at 25°C/60%RH.
- The appearance of both the cake and the reconstituted solutions didn't undergo any variation.
- The assay remained unvaried.

L10R5

Slight increase of moisture content (anyway the % H₂O<1.0%).
Slight increase of the impurities due to the CNS7054X.

L10R5S87

The moisture content didn't increase.

The impurities content remained practically unvaried.

L20R5 (visual inspection)

Increase of moisture content (anyway the % H₂O<1.0%).

Increase of the impurities content mainly due to CNS7054X.

IV. Preparation and stability analysis of disaccharide/dextran containing formulations, as a means to reduce lyophilization time

1. Purpose

Within this study several CNS7056 lyophilisate formulations containing lactose and dextran were studied. The ratio of the disaccharide to dextran was changed in order to manipulate the glass transition temperature (Tg') and collapse temperature T_c and therefore reduce the lyophilisation time. Compared to the disaccharide lactose the dextran possesses a higher Tg' and therefore can act as a collapse temperature modifier.

Altogether 10 formulations were prepared and tested in different lyophilization protocols.

2. Formulations

2.1. Formulation compositions

Two CNS7056 formulations were prepared containing dextran only (001/PAN/13) or a mixture of lactose and dextran (002/PAN/13) as summarized in the following table:

Name	Formulation
001/PAN/13	50:440:7056:Dextran
002/PAN/13	50:220:220:7056:Lactose:Dextran

2.2. Formulation preparation

2.2.1. Preparation of the solution

The solutions containing 12 mg/mL CNS7056 were prepared according to the following protocol:

- API (CNS 7056 besylate salt) added with magnetic stirring to 85% final volume
- Stirred for 3 hours at ambient, light protected
- Checked pH, nominal pH 3.2 for all formulations, and adjusted to pH 3.0
- Stirred for further 20 minutes: no significant change in appearance
- Made to 90% final volume
- Stirred for further 20 minutes
- Formulations 1-2 appeared light yellow, slightly turbid.
- Checked pH, all nominal pH 3
- Made to final volume and stirred for further 20 minutes
- No change in formulation 1-2 appearance, less undissolved material in the concentrated formulation
- Filtered (0.22 µm PVDF) formulations 1- 2
- Further 25 minutes stirring of concentrated formulationFiltered (0.22 µm PVDF) concentrated formulation
- All filtrates clear, light yellow, free from visible particles
- Filtrates filled in 4.2 mL volumes and lyophilised with protocol as described in 2.1.2

2.2.2. Lyophilisation protocol

The samples were lyophilized according to the following protocol:

Step	Cycle stage	Temperature (°C)	Pressure (mTorr)	Time (min)
1	Load	25	n/a	0
2	Ramp	0	n/a	25
3	Ramp	-45	n/a	225
4	Freezing	-45	n/a	180
5	Hold	-45	93	0
6	Ramp	-25	93	30
7	Primary drying	-25	93	4890
8	Ramp	30	20	120
9	Secondary drying	30	20	480
10	Finish	30	Vials stoppered to 722000 mTorr with (pure) nitrogen.	
Total cycle duration ~ 99 hours (~ 4.1 days)				

2.3 Analysis of the lyophilized samples

The lyophilisate showed a good appearance and a rapid reconstitution time for the carbohydrate:dextran-containing lyophilisates. Both formulations exhibit a purity above 99.72%.

Name	Formulation	Appearance	Recon time in saline	pH	7054X (%)	Purity (%)
001/PAN/13	50: 440, 7056:Dextran	Off white plug	1 m 50 s	3.241	0.11	99.72
002/PAN/13	50:220:220, 7056:Lactose:Dextran	Off white plug	35 s	3.229	0.09	99.74

2.3.1 Appearance

The appearance of the lyophilized samples was determined. The results are listed in the following table:

Sample details			Appearance
001/PAN/13	Initial (T = 0)		Off white plug
	40°C/75%RH	T = 1 m	Off white plug with signs of shrinkage
002/PAN/13	Initial (T = 0)		Off white plug
	40°C/75%RH	T = 1 m	Off white plug with signs of shrinkage

2.3.2 Moisture

The moisture content of the lyophilized samples was determined. The results are listed in the following table:

Sample details			Vial 1	Vial 2	Vial 3	Mean moisture (% w/w)
001/PAN/13	Initial (T = 0)		0.04	0.12	0.19	0.12
	40°C/75%RH	T = 1 m	0.19	0.19	-	0.19
002/PAN/13	Initial (T = 0)		0.16	0.16	0.38	0.23
	40°C/75%RH	T = 1 m	0.36	0.38	-	0.37

2.3.3 Reconstitution time and pH of reconstituted solution

Each vial was reconstituted with 10 mL 0.9% saline. The results regarding reconstitution time and pH are listed in the following table.

5

Sample details			Recon. time (seconds)	pH
001/PAN/13	Initial (T = 0)		110	3.24
	40°C/75%RH	T = 1 m	153	3.21
002/PAN/13	Initial (T = 0)		35	3.23
	40°C/75%RH	T = 1 m	85	3.26

2.3.4 Vial content

The vial content for the samples at T=1m 40°C/75%RH was determined after each vial was reconstituted with 10 mL 0.9% saline. The results are given in the following table:

10

Details	[7056] (mg/vial)		
	Vial 1	Vial 2	Mean
001/PAN/13	49.1136	49.6891	49.401
002/PAN/13	49.0496	49.2652	49.157

2.3.4 Impurities

The impurities for the different formulations at T=1m 40°C/75%RH were determined. The results are given in the following tables:

15

20

001/PAN/13:

- 52 -

			Initial* ² (T=0)	40°C/75%RH T= 1m
	RRT	Name		
Impurity profile (area %)	0.27	n.a.	N.D.	0.03
	0.42	n.a.	N.D.	<LOQ
	0.47	n.a.	<LOQ	<LOQ
	0.51	n.a.	0.07	0.07
	0.57	n.a.	<LOQ	<LOQ
	0.59	7054X	0.11	0.17
	0.64	n.a.	N.D.	<LOQ
	0.68	n.a.	<LOQ	<LOQ
	0.71	n.a.	N.D.	<LOQ
	0.89	n.a.	0.10	N.D.
	0.93	n.a.	N.D.	0.10
	1.00	7056B	99.59	99.45
	1.13	n.a.	N.D.	0.03
	1.31	n.a.	<LOQ	<LOQ
	1.46	n.a.	N.D.	<LOQ
	1.73	n.a.	N.D.	<LOQ
	1.78	n.a.	<LOQ	<LOQ
	1.84	n.a.	<LOQ	<LOQ
	1.91	n.a.	<LOQ	N.D.
	Total imps* ¹ (area%)		0.3	0.4

*¹ Sum of all impurities $\geq 0.03\%$ by area*² Impurities from post lyo samples

Impurities are mean of 2 determinations

n.d. = not detected

002/PAN/13:

			Initial (T=0)	40°C/75%RH T= 1m
	RRT	Name		
Impurity profile (area %)	0.27	n.a.	N.D.	<LOQ
	0.47	n.a.	<LOQ	N.D.
	0.51	n.a.	0.07	0.07
	0.56	n.a.	N.D.	<LOQ
	0.59	7054X	0.09	0.14
	0.64	n.a.	N.D.	<LOQ
	0.68	n.a.	<LOQ	<LOQ
	0.71	n.a.	N.D.	<LOQ
	0.83	n.a.	N.D.	<LOQ
	0.89	n.a.	<LOQ	N.D.
	0.93	n.a.	0.10	0.10
	1.00	7056B	99.62	99.49
	1.10	n.a.	N.D.	0.03
	1.31	n.a.	<LOQ	<LOQ
	1.73	n.a.	N.D.	<LOQ
	1.78	n.a.	<LOQ	<LOQ
	1.84	n.a.	<LOQ	<LOQ
	Total imps* (area%)		0.3	0.3

* Sum of all impurities $\geq 0.03\%$ by area

Impurities are mean of 2 determinations

n.d. = not detected

- 53 -

3. Formulations – Potential Process Improvement with Dextran 40 Based Formulations

3.1. Formulation compositions

Two CNS7056 formulations were prepared containing dextran (007/PAN/13), or a mixture of lactose and dextran 40(009/PAN/13) as summarized in the following table:

Batch	Formulation	7056:excipient(s) (mg)
007/PAN/13	7056:dextran 40	50:440
009/PAN/13	7056:lactose monohydrate:dextran 40	50:88:352

3.2. Formulation preparation

3.2.1. Preparation of the solution

Dissolution of CNS7056B was facilitated by overhead stirring under ambient laboratory conditions, protected from light. A summary of the salient points for the preparation of each formulation are presented below.

007/PAN/13

- API addition (<5 minutes) to ~85% final volume
- Stirring started at 500 rpm and increased to 700 rpm by 120 minutes
- Adjusted to pH 3.0 after 70 minutes
- Increased to ~90% after 120 minutes
- Adjusted to pH 2.8 after 150 minutes
- After 180 minutes, checked pH (2.9), adjusted to pH 3.0, and made to final volume

009/PAN/13

- API addition (<5 minutes) to ~95% final volume
- Following API (500 rpm) stirring immediately increased to 700 rpm and then increased to 800 rpm by 30 minutes
- Adjusted to pH 3.0 after 10 minutes
- After 75 minutes, checked pH (pH 3.1), adjusted to pH 3.0, and made to final volume

Following preparation all formulations were filtered through a 0.22µm PVDF membrane filter

3.2.2. Lyophilisation protocol

Filtrates were filled in 4.2 mL volumes into 20 mL clear Type glass vials and lyophilised, directly from the shelf, with the cycle shown in the following table:

Step	Cycle stage	Temperature (°C)	Pressure (mTorr)	Time (min)
1	Load	25	n/a	0
2	Ramp	0	n/a	25
3	Ramp	-45	n/a	225
4	Freezing	-45	n/a	180
5	Hold	-45	350	30
6	Ramp	-15	350	60
7	Primary drying	-15	350	2861
8	Ramp	30	20	112
9	Secondary drying	30	20	459
10	Finish	30	Vials stoppered to 722000 mTorr with (pure) nitrogen.	
Total cycle duration ~ 66 hours (~ 2.8 days)				

3.3 Sample analysis

There was no significant difference in the appearance of the lyophilised plugs among batches 007 and 009/PAN/13. Lyophilised plugs appeared white/off-white, homogeneous and well formed.

Duplicate vials of each product were used for initial T = 0 testing, a summary of the analytical results is presented below.

3.3.1 Analysis after reconstitution

10 mL normal saline was added to a lyophilised sample, the vial was swirled and observed. The reconstitution time, reconstitution solution appearance, pH and purity (HPLC) were determined. The results are shown in the following table:

Batch	Reconstitution time (s)	Reconstituted solution			
		Appearance	pH	7056 (%area)	7054X (%area)
007/PAN/13	91	Clear, colourless, free from visible particulates	3.3	99.58	0.10
009/PAN/13	81	Clear, colourless, free from visible particulates	3.2	99.60	0.08

3.3.2 Appearance

The appearance of the lyophilized samples was determined. The results are listed in the following table:

Sample details			Appearance
007/PAN/13	Initial (T = 0)		Off white plug
	40°C/75%RH	T = 1 m	Off white plug with signs of shrinkage
	55°C	T = 1 m	Off white plug with signs of shrinkage
009/PAN/13	Initial (T = 0)		Off white plug
	40°C/75%RH	T = 1 m	Off white plug with signs of shrinkage
	55°C	T = 1 m	Off white plug with signs of shrinkage

3.3.3 Moisture

The moisture content of the lyophilized samples was determined. The results are listed in the following table:

Sample details			Vial 1	Vial 2	Vial 3	Mean moisture (% w/w)
007/PAN/13	Initial (T = 0)		0.00	0.00	-	0.00
	40°C/75%RH	T = 1 m	0.14	0.12	-	0.13
	55°C	T = 1 m	0.24	0.32	-	0.28
009/PAN/13	Initial (T = 0)		0.00	0.00	-	0.00
	40°C/75%RH	T = 1 m	0.20	0.21	-	0.21
	55°C	T = 1 m	0.35	0.38	-	0.37

3.3.4 Reconstitution time and pH of reconstituted solution

Each vial was reconstituted with 10 mL 0.9% saline. The results regarding reconstitution time and pH are listed in the following table.

Sample details			Recon. time (seconds)	pH
007/PAN/13	Initial (T = 0)		165*	3.25
	40°C/75%RH	T = 1 m	79	3.21
	55°C	T = 1 m	62	3.28
009/PAN/13	Initial (T = 0)		95*	3.22
	40°C/75%RH	T = 1 m	59	3.21
	55°C	T = 1 m	50	3.21

3.3.5 Vial content

The vial content for the samples at T=1m 40°C/75%RH was determined after each vial was reconstituted with 10 mL 0.9% saline. The results are given in the following table:

Sample details			Vial 1	Vial 2	Mean [7056] (mg/vial)	Recovery ¹ (%)
007/PAN/13	Initial (T = 0)		48.6048	48.5855	48.595	-
	40°C/75%RH	T = 1 m	48.0353	47.9755	48.005	98.8
	55°C	T = 1 m	47.9019	48.6901	48.296	99.4
009/PAN/13	Initial (T = 0)		48.9599	48.9696	48.965	-
	40°C/75%RH	T = 1 m	48.4752	49.4228	48.949	100.0
	55°C	T = 1 m	49.0987	48.3462	48.722	99.5

¹ Recovery is calculated as a percentage of T = 0 result.

3.3.6 Impurities

The impurities for the different formulations at T=1m 40°C/75%RH were determined.
The results are given in the following tables:

5

007/PAN/13:

			Initial (T=0)	40°C/75%RH	55°C
	RRT	Name	T= 1m	T= 1m	T= 1m
Impurity profile (area%)	0.36	n.a.	N.D.	N.D.	<LOQ
	0.41	n.a.	<LOQ	N.D.	N.D.
	0.46	n.a.	<LOQ	<LOQ	<LOQ
	0.51	n.a.	0.07	0.07	0.07
	0.57	n.a.	N.D.	<LOQ	<LOQ
	0.59	7054X	0.10	0.16	0.43
	0.63	n.a.	<LOQ	<LOQ	<LOQ
	0.67	n.a.	<LOQ	<LOQ	<LOQ
	0.70	n.a.	<LOQ	<LOQ	<LOQ
	0.88	n.a.	<LOQ	<LOQ	<LOQ
	0.92	n.a.	0.11	0.11	0.11
	1.00	7056B	99.58	99.54	99.23
	1.31	n.a.	<LOQ	<LOQ	<LOQ
	1.46	ONO	N.D.	N.D.	<LOQ
	1.73	n.a.	<LOQ	<LOQ	<LOQ
	1.77	n.a.	<LOQ	N.D.	N.D.
	1.79	n.a.	<LOQ	<LOQ	<LOQ
	1.84	n.a.	<LOQ	N.D.	N.D.
	1.86	n.a.	N.D.	<LOQ	<LOQ
	Total imps* (area%)		0.3	0.3	0.6

* Sum of all impurities ≥ 0.03% by area
Impurities are mean of 2 determinations
n.d. = not detected

10

009/PAN/13:

			Initial (T=0)	40°C/75%RH T= 1m	55°c T= 1m
	RRT	Name			
Impurity profile (area%)	0.27	n.a.	N.D.	N.D.	<LOQ
	0.36	n.a.	N.D.	N.D.	<LOQ
	0.41	n.a.	<LOQ	N.D.	N.D.
	0.46	n.a.	<LOQ	<LOQ	<LOQ
	0.51	n.a.	0.07	0.07	0.07
	0.57	n.a.	N.D.	<LOQ	<LOQ
	0.59	7054X	0.08	0.12	0.38
	0.63	n.a.	<LOQ	<LOQ	<LOQ
	0.67	n.a.	<LOQ	<LOQ	<LOQ
	0.70	n.a.	<LOQ	<LOQ	<LOQ
	0.88	n.a.	<LOQ	<LOQ	<LOQ
	0.92	n.a.	0.11	0.11	0.10
	1.00	7056B	99.60	99.57	99.27
	1.31	n.a.	<LOQ	<LOQ	<LOQ
	1.47	ONO	N.D.	N.D.	<LOQ
	1.73	n.a.	<LOQ	N.D.	N.D.
	1.75	n.a.	N.D.	<LOQ	<LOQ
	1.77	n.a.	<LOQ	N.D	N.D.
	1.79	n.a.	<LOQ	<LOQ	<LOQ
	1.84	n.a.	<LOQ	N.D.	N.D.
	1.86	n.a.	N.D.	<LOQ	<LOQ
Total imps* (area%)			0.3	0.3	0.6

* Sum of all impurities ≥ 0.03% by area
Impurities are mean of 2 determinations
n.d. = not detected

4. Further formulations

4.1 Parameters:

- Fill concentration of CNS 7056 base = 12 mg/mL
- Vial size = 20R (30mm diameter)
- Fill solution volume 4.2 mL
- Approx 50 vials of each product to be prepared
- Also prepare placebos

4.2 Formulation Compositions (Vial equivalent)

	CNS 7056 (base equivalent)	Total Excipient	Dextran 40	Lactose Monohydrate	Approx Ratio CNS7056: Lactose monohydrate
			80	20	
012/PAN/13	50 mg	440 mg	352 mg	88 mg	1:9
011/PAN/13	50 mg	330 mg	264 mg	66 mg	1:6
010/PAN/13	50 mg	220 mg	176 mg	44 mg	1:4.5

Predicted Tc of formulations from Thermal Assessments = -15 °C

4.3 Parameters

- Freezing to -30 °C @ 0.2 °C/min. Held
- Primary Drying -7 °C @ 700-750mTorr
- Secondary Drying 30 °C

4.4 Test Criteria for analysis of lyo samples

- Appearance
- Recon time - addition of 10 mL saline
- Moisture
- Related substances
- Stability - 1m 55C + 1, 3m 40C/75%RH

4.4.1 Appearance

The appearance of the lyophilized samples was determined. The results are listed in the following table:

Sample details		Appearance
010/PAN/13	Initial (T = 0)	Off white plug
011/PAN/13	Initial (T = 0)	Off white plug with material on vial wall
012/PAN/13	Initial (T = 0)	Off white plug

4.4.2 Moisture

The moisture content of the lyophilized samples was determined. The results are listed in the following table:

Sample details		Vial 1	Vial 2	Vial 3	Mean moisture (% w/w)
010/PAN/13	Initial (T = 0)	0.08	0.02	-	0.05
011/PAN/13	Initial (T = 0)	0.05	0.05	-	0.05
012/PAN/13	Initial (T = 0)	0.11	0.06	-	0.09

4.4.3 Reconstitution time and pH of reconstituted solution

Each vial was reconstituted with 10 mL 0.9% saline. The results regarding reconstitution time and pH are listed in the following table.

Sample details			Recon. time (seconds)	pH
010/PAN/13	Initial (T = 0)		48	3.23
	40°C/75%RH	T = 1 m	-	-
	55°C	T = 1 m	-	-
011/PAN/13	Initial (T = 0)		63	3.20
	40°C/75%RH	T = 1 m	-	-
	55°C	T = 1 m	-	-
012/PAN/13	Initial (T = 0)		64	3.23
	40°C/75%RH	T = 1 m	-	-
	55°C	T = 1 m	-	-

4.4.4 Vial content

5

The vial content for the samples at T=1m 40°C/75%RH was determined after each vial was reconstituted with 10 mL 0.9% saline. The results are given in the following table:

Sample details		Vial 1	Vial 2	Mean [7056] (mg/vial)
010/PAN/13	Initial (T = 0)	49.8316	49.4836	49.658
011/PAN/13	Initial (T = 0)	49.9339	49.1202	49.527
012/PAN/13	Initial (T = 0)	49.0690	47.6608	48.365

4.4.5 Impurities

10

The impurities for the different formulations at T=1m 40°C/75%RH were determined. The results are given in the following tables:

010/PAN/13:

			Initial (T=0)
	RRT	Name	
Impurity profile (area %)	0.31	n.a.	<LOQ
	0.47	n.a.	<LOQ
	0.52	n.a.	0.07
	0.57	n.a.	<LOQ
	0.59	7054X	0.10
	0.65	n.a.	<LOQ
	0.68	n.a.	<LOQ
	0.72	n.a.	<LOQ
	0.89	n.a.	<LOQ
	0.93	n.a.	0.11
	1.00	7056B	99.57
	1.14	n.a.	0.03
	1.31	n.a.	<LOQ
	1.74	n.a.	<LOQ
	1.80	n.a.	<LOQ
	1.86	n.a.	<LOQ
	2.00	n.a.	<LOQ
	Total imps* (area %)		0.3

* Sum of all impurities ≥ 0.03% by area
Impurities are mean of 2 determinations
n.d. = not detected

011/PAN/13:

			Initial (T=0)
	RRT	Name	
Impurity profile (area %)	0.37	n.a.	<LOQ
	0.52	n.a.	0.07
	0.59	7054X	0.10
	0.68	n.a.	<LOQ
	0.89	n.a.	<LOQ
	0.93	n.a.	0.11
	1.00	7056B	99.56
	1.10	n.a.	0.03
	1.74	n.a.	<LOQ
	1.80	n.a.	<LOQ
	1.86	n.a.	<LOQ
	2.00	n.a.	<LOQ
	Total imps* (area %)		0.3

* Sum of all impurities ≥ 0.03% by area
Impurities are mean of 2 determinations
n.d. = not detected

012/PAN/13:

			Initial (T=0)
	RRT	Name	
Impurity profile (area%)	0.31	n.a.	<LOQ
	0.47	n.a.	<LOQ
	0.52	n.a.	0.07
	0.59	7054X	0.08
	0.64	n.a.	<LOQ
	0.68	n.a.	<LOQ
	0.71	n.a.	<LOQ
	0.89	n.a.	<LOQ
	0.93	n.a.	0.11
	1.00	7056B	99.58
	1.14	n.a.	0.03
	1.31	n.a.	<LOQ
	1.74	n.a.	<LOQ
	1.80	n.a.	<LOQ
	1.86	n.a.	<LOQ
Total imp [*] (area%)			0.3

* Sum of all impurities ≥ 0.03% by area
Impurities are mean of 2 determinations
n.d. = not detected

5. Further formulations

5.1 Parameters:

- Fill concentration of CNS 7056 base = 12 mg/mL
- Vial size = 20R (30mm diameter)
- Fill solution volume 4.2 mL
- Approx 50 vials of each product to be prepared
- Also prepare placebos

5.2 Formulation Compositions (Vial equivalent)

	CNS 7056 (base equivalent)	Total Excipient	Dextran 40	Lactose Monohydrate	Approx Ratio CNS7056: Lactose monohydrate
			60	40	
015/PAN/13	50 mg	440 mg	264 mg	176 mg	1:9
014/PAN/13	50 mg	330 mg	198 mg	132 mg	1:6
013/PAN/13	50 mg	220 mg	132 mg	88 mg	1:4.5

Predicted Tc of formulations from Thermal Assessments = -19°C

5.3 Parameters

- 62 -

- Freezing to -30 °C @ 0.2 °C/min
- Primary Drying -15 °C @ 400mTorr
- Secondary Drying 30 °C

5.4 Test Criteria for analysis of lyo samples

- Appearance
- Recon time - addition of 10 mL saline
- Moisture
- Related substances
- Stability - 1m 55C + 1, 3m 40C/75%RH

5.4.1 Appearance

The appearance of the lyophilized samples was determined. The results are listed in the following table:

Sample details		Appearance
013/PAN/13	Initial (T = 0)	Off white plug some material on walls of vial
014/PAN/13	Initial (T = 0)	Off white plug some material on walls of vial
015/PAN/13	Initial (T = 0)	Off white plug some material on walls of vial

5.4.2 Moisture

The moisture content of the lyophilized samples was determined. The results are listed in the following table:

Details		Vial 1	Vial 2	Vial 3	Mean moisture (% w/w)
013/PAN/13	Initial (T = 0)	0.06	0.05	-	0.06
014/PAN/13	Initial (T = 0)	0.00	0.09	-	0.09
015/PAN/13	Initial (T = 0)	0.09	0.00	-	0.09

5.4.3 Reconstitution time and pH of reconstituted solution

Each vial was reconstituted with 10 mL 0.9% saline. The results regarding reconstitution time and pH are listed in the following table.

Sample details		Recon. time (seconds)	pH
013/PAN/13	Initial (T = 0)	39	3.21
014/PAN/13	Initial (T = 0)	35	3.22
015/PAN/13	Initial (T = 0)	43	3.24

5.4.4 Vial content

The vial content for the samples at T=1m 40°C/75%RH was determined after each vial was reconstituted with 10 mL 0.9% saline. The results are given in the following table:

Sample details		Vial 1	Vial 2	Mean [7056] (mg/vial)
013/PAN/13	Initial (T = 0)	48.1356	48.5325	48.334
014/PAN/13	Initial (T = 0)	49.9574	49.7535	49.855
015/PAN/13	Initial (T = 0)	48.3542	47.9459	48.150

5.4.5 Impurities

The impurities for the different formulations at T=1m 40°C/75%RH were determined. The results are given in the following tables:

013/PAN/13:

			Initial (T=0)
	RRT	Name	
Impurity profile (area %)	0.31	n.a.	<LOQ
	0.47	n.a.	<LOQ
	0.52	n.a.	0.07
	0.59	7054X	0.09
	0.64	n.a.	<LOQ
	0.68	n.a.	<LOQ
	0.71	n.a.	<LOQ
	0.89	n.a.	<LOQ
	0.93	n.a.	0.11
	1.00	7056B	99.60
	1.31	n.a.	<LOQ
	1.74	n.a.	<LOQ
	1.77	n.a.	<LOQ
	1.79	n.a.	<LOQ
	1.85	n.a.	<LOQ
Total imps* (area%)			0.3

* Sum of all impurities ≥ 0.03% by area
Impurities are mean of 2 determinations
n.d. = not detected

014/PAN/13:

			Initial (T=0)
	RRT	Name	
Impurity profile (area%)	0.31	n.a.	<LOQ
	0.47	n.a.	<LOQ
	0.51	n.a.	0.07
	0.58	n.a.	<LOQ
	0.59	7054X	0.09
	0.64	n.a.	<LOQ
	0.68	n.a.	<LOQ
	0.71	n.a.	<LOQ
	0.89	n.a.	<LOQ
	0.92	n.a.	0.11
	1.00	7056B	99.61
	1.31	n.a.	<LOQ
	1.74	n.a.	<LOQ
	1.79	n.a.	<LOQ
	1.85	n.a.	<LOQ
	Total imps* (area%)		0.3

* Sum of all impurities $\geq 0.03\%$ by area
Impurities are mean of 2 determinations
n.d. = not detected

015/PAN/13:

			Initial (T=0)
	RRT	Name	
Impurity profile (area%)	0.31	n.a.	<LOQ
	0.47	n.a.	<LOQ
	0.51	n.a.	0.07
	0.56	n.a.	<LOQ
	0.59	7054X	0.08
	0.64	n.a.	<LOQ
	0.68	n.a.	<LOQ
	0.71	n.a.	<LOQ
	0.89	n.a.	<LOQ
	0.92	n.a.	0.11
	1.00	7056B	99.61
	1.31	n.a.	<LOQ
	1.74	n.a.	<LOQ
	1.79	n.a.	<LOQ
	1.85	n.a.	<LOQ
	Total imps* (area%)		0.3

* Sum of all impurities $\geq 0.03\%$ by area
Impurities are mean of 2 determinations
n.d. = not detected

5

V. THERMAL ANALYSIS OF REMIMAZOLAM FORMULATIONS

1. Purpose of the study

In order to increase the lyophilisation temperature, the relative amount of dextran of the lactose:dextran mixture was increased and the critical temperature was determined by differential scanning calorimetry (DSC) and freeze drying microscopy (FDM), for CNS7056B formulations 1 – 6 as shown in the following Table.

10

Formulation	T _g ' by DSC (°C)	T _c by FDM (°C)
001	-13	-11
002	-23	-21
003	-28	-27
004	-24	-20
005	-29	-28
006	-29	-27

Critical temperatures were plotted with respect to dextran, relative to total formulation solute content and shown in Figure. 52.

From the linear equations (Figure 52), for a given critical temperature, the theoretical dextran content of a CNS7056B formulation can be calculated. As shown in the following table, there was a good correlation between data generated for formulations containing lactose.

Target T _c (°C)	2 nd excipient	Technique	Theoretical dextran (%)
-20	Lactose	DSC	48.8
		FDM	56.8
	DSC mean		49
	FDM mean		58
-17.5	Lactose	DSC	60.3
		FDM	69.1
	DSC mean		60
	FDM mean		70
-15	Lactose	DSC	71.9
		FDM	81.3
	DSC mean		71
	FDM mean		82

An alternative presentation of the data, expressing collapse temperature relative to the dextran:lactose ratio in each formulation is shown Figure 53. The Phase I/II sedation formulation was used to represent a formulation containing no dextran (zero on the abscissa). The collapse temperature onset for this formulation has been reported as -31°C.

Similarly, the linear equation from Figure 53 may be used to calculate the theoretical dextran:lactose composition of CNS7056B formulations for given collapse temperatures as shown in the following table:

Target Tc (°C)	Theoretical excipient composition (%)		Example formulation API:lactose:dextran (mg/vial)
	Lactose	Dextran	
-20	45.4	54.6	50:200:240
-17.5	32.8	67.2	50:145:295
-15	20.2	79.8	50:90:350

V.

Figure Legends

Fig. 1: Excipients

Fig. 2: Active formulations

5 Fig. 3: Placebo formulations

Fig. 4: Formulation of hydrolysis degradant of remimazolam (REM) given in % after storage for 13 weeks at 25°C/60% relative humidity (RH) or 40°C/75%RH.

10 Fig. 5A-D: Crystallographic co-ordinates and other relevant data tabulated in the form of a SHELX File for Compound of formula (I) besylate Form 1 of WO2008/007071 A1

Fig. 6A-C: Crystallographic co-ordinates and other relevant data tabulated in the form of a SHELX File for Compound of formula (I) besylate Form 2 of WO2008/007071 A1.

15 Fig. 7A-B: Bond lengths for Compound of formula (I) besylate Form 1 of WO2008/007071 A1

Fig. 8A-C: Bond angles for Compound of formula (I) besylate Form 1 of WO2008/007071 A1

20 Fig. 9: Bond Lengths for Compound of formula (I) besylate Form 2 of WO2008/007071 A1

Fig. 10: Bond angles for Compound of formula (I) besylate Form 2 of WO2008/007071 A1

Fig. 11: Stability data for lot A01P310

Fig. 12: Stability data for lot A01P310, continued

25 Fig. 13: Accelerated stability data for lot A01P310

Fig. 14: Accelerated stability data for lot A01P310, continued

Fig. 15: Long term stability data for lot P310-01

Fig. 16: Long term stability data for lot P310-01, continued

Fig. 17: Accelerated stability data for lot P310-01

30 Fig. 18: Accelerated stability data for lot P310-01, continued

Fig. 19: Long term stability data for lot 026CNS27

Fig. 20: Long term stability data for lot 026CNS27, continued

Fig. 21: Accelerated stability data for lot 026CNS27

Fig. 22: Accelerated stability data for lot 026CNS27, continued

Fig. 23: Long term stability data for lot G384

Fig. 24: Long term stability data for lot G384, continued

Fig. 25: Accelerated stability data for lot G384

5 Fig. 26: Accelerated stability data for lot G384, continued

Fig. 27: Long term stability data for lot P02308

Fig. 28: Long term stability data for lot P02308, continued

Fig. 29: Accelerated stability data for lot P02308

Fig. 30: Accelerated stability data for lot P02308, continued

10 Fig. 31: Long term stability data for lot 25CNS27

Fig. 32: Long term stability data for lot 25CNS27, continued

Fig. 33: Long term stability data for lot 25CNS27, continued

Fig. 34: Accelerated stability data for lot 25CNS27

Fig. 35: Accelerated stability data for lot 25CNS27, continued

15 Fig. 36: Accelerated stability data for lot 25CNS27, continued

Fig. 37: Long term stability data (T=36M) for lot P02308

Fig. 38: Long term stability data (T=36M) for lot P02308, continued

20 Fig. 39: Raman spectra of each component in a lyophilized formulation: each rectangle range showing distinctive peak(s) of crystalline and lyophilized CNS7056B (L) and lyophilized lactose (R).

Fig. 40: Raman spectra of each component in a lyophilized formulation of CNS7056B in lactose: crystalline CNS7056B (top), lyophilized (amorphous) CNS7056B (middle) and lyophilized lactose (amorphous) (bottom).

25 Fig. 41: Representative Raman spectra of crystalline CNS7056B (top 3) selected within the Raman mapping area of the lyophilized formulation and pure crystalline (Form I) CNS7056B (bottom) as a reference.

Fig. 42: Table summarizing the results for the stability study of the Lots 12PM529-8-1, 12PM529-8-2, 12PM529-9-1 and PM0232/12 at initial time point t=0

30 Fig. 43: Table summarizing the results for the stability study of the 12PM529-8-2, 12PM529-9-1 and PM0232/12 at time point t = 4 weeks

Fig. 44: Table summarizing the results for the stability study of the Lots 12PM529-8-1, 12PM529-8-2, 12PM529-9-1 and PM0232/12 at time point t = 13 weeks

- 69 -


- Fig. 45 Table summarizing the results for the stability study of the Lot L10R5
- Fig. 46 Table summarizing the results for the stability study of the Lot L10R10
- Fig. 47 Table summarizing the results for the stability study of the Lot L10R5S87
- Fig. 48 Table summarizing the results for the stability study of the Lot L20R5
- 5 Fig. 49 Table summarizing the results for the stability study of the Lots L4M110R5 and L4M110R10
- Fig. 50 Table summarizing the results for the stability study of the Lot L2M110R5
- Fig. 51 Table summarizing the results for the stability study of the Lot L2M110R10
- 10 Fig. 52 Critical temperature as a function of dextran content for CNS7057B:lactose:dextran formulations
- Fig. 53 Collapse temperature relative to the dextran:lactose ratio for CNS7056B formulations

KÖNIG · SZYNKA · TILMANN · von RENESSE

P A T E N T A N W Ä L T E · P A R T N E R S C H A F T m b B

PCT/EP2013/060543
PAION UK LIMITED54 432 K
July 29, 2014Amended Claims (clean copy)

1. A lyophilized or spray-dried composition comprising at least one benzodiazepine according to formula (I)



with

W is H;

X is CH₂; n is 1;Y is CH₂; m is 1;

Z is O; p is 0 or 1;

R¹ is CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂ or CH₂CH(CH₃)₂;R² is 2-fluorophenyl, 2-chlorophenyl or 2-pyridyl;R³ is Cl or Br;

R⁴, R⁵ and R⁶ form the group-CR⁸=U-V= wherein R⁸ is hydrogen, C₁₋₄ alkyl or C₁₋₃ hydroxyalkyl, U is N or CR⁹ wherein R⁹ is H, C₁₋₄ alkyl, C₁₋₃ hydroxyalkyl or C₁₋₄ alkoxy, V is N or CH and p is zero.

or a pharmaceutically acceptable salt thereof, wherein the composition

comprises at least one pharmaceutically acceptable hygroscopic excipient, wherein the hygroscopic excipient is a carbohydrate selected from the group consisting of disaccharides and dextran and optionally the composition is at least in part amorphous.

2. A composition according to claim 1, wherein p is zero and R⁴, R⁵ and R⁶ form the group -CR⁸=U-V= wherein R⁸ is methyl, U is CH₂, V is N; W is H; X is CH₂, n is 1; Y is CH₂, m is 1; R¹ is CH₃; R² is 2-pyridyl; R³ is Br.
3. A composition according to claim 1 or 2, wherein the benzodiazepine is in the form of a pharmaceutically acceptable salt.
4. A composition according to any of the preceding claims, wherein in the pharmaceutically acceptable salt the benzodiazepine is formulated in cationic form and the counter ion is selected from halogenides, in particular fluoride, chloride or bromide, sulfate, organic sulfates, sulfonate, organic sulfonates, nitrate, phosphate, salicylate, tartrate, citrate, maleate, formate, malonate, succinate, isethionate, lactobionate and sulfamate.
5. A composition according to claim 4, wherein the counter ion is selected from organic sulfates and sulfonates, in particular aromatic sulfates and sulfonates.
6. A composition according to claim 5, wherein the counter ion is benzene sulfonate (besylate).
7. A composition according to claim 5, wherein the benzodiazepine salt is remimazolam besylate.
8. A composition according to claim 5, wherein the benzodiazepine salt is a tosylate, in particular remimazolam tosylate.
9. A composition according to claim 5, wherein the benzodiazepine salt is an esylate, in particular remimazolam esylate.
10. A composition according to any of the preceding claims, wherein the carbohydrate possesses a molecular weight of less than 150 kD.
11. A composition according to claim 10, wherein the carbohydrate is a dextran molecule.
12. A composition according to claim 10, wherein the carbohydrate is selected from monosaccharides and C₂₋₆-oligosaccharides.

13. A composition according to claim 12, wherein the carbohydrate is a disaccharide, preferably selected from the group consisting of lactose, maltose, sucrose and trehalose.
14. A composition according to any of the preceding claims, wherein the composition comprises a mixture of a first hygroscopic excipient and a second hygroscopic excipient.
15. A composition according to claim 14, wherein the first excipient is a disaccharide.
16. A composition according to claim 15, wherein the disaccharide is selected from the list consisting of lactose, maltose, sucrose and trehalose.
17. A composition according to claim 14 to 16, wherein the second excipient is polymer, preferably a polysaccharide or organic polymer.
18. A composition according to claim 17, wherein the polysaccharide is dextran, preferably with a molecular weight of 80 KD or less.
19. A composition according to any of the claims 14 to 18, wherein the wt% ratio of the first excipient and the second excipient is from 20:1 to 1:10, preferably from 5:1 to 1:5 and more preferably 1:1.5 or 1:4.
20. A composition according to any of the preceding claims, wherein the composition is solid, preferably lyophilized and comprises a mixture of lactose and dextran.
21. A composition according to claim 20, wherein the wt% ratio between lactose and dextran is from 1:1.0 to 1:10, preferably from 1:1.2 to 1:5, and more preferably is 1:1.5 or 1:4.
22. A composition according to any of the preceding claims wherein the wt% ratio between the total amount of excipients and the total amount of benzodiazepines or salts, calculated for the base, in the composition is at least 1:1, preferably between 9:1 and 4.5 :1, most preferably 6:1.
23. A composition according to claim 22, wherein the excipients are carbohydrates and the benzodiazepine is remimazolam.
24. A composition according to any of the preceding claims wherein the wt% ratio between the total amount of hygroscopic excipients and the total amount of

benzodiazepines or salts thereof, calculate for the base, in the composition is from 20:1 to 1:1, preferably from 12:1 to 3:1, most preferably from 9:1 to 3:1.

25. A composition according to any of the preceding claims, wherein the composition is in the solid state and is preferably a lyophilized solid.

26. A composition according to any of the preceding claims, wherein said composition comprises a compound with a collapse temperature above -20°C (collapse temperature modifier).

27. A composition according to claim 26, wherein the collapse temperature modifier is selected from the group consisting of a sucrose-epichlorhydrin-copolymer, being preferably Ficoll®, gelatine and hydroxyethyl starch (HES), preferred is HES.


28. A composition according to any of the above claims, wherein the benzodiazepine or the salt thereof is amorphous for at least 50%, preferably for at least 75%, more preferably for at least 90% and most preferably for at least 95%.

29. A composition of any of the above claims which is a pharmaceutical formulation.

30. Pharmaceutical composition according claim 29 comprising a benzodiazepine or a salt thereof according to any of the above claims in an amount from 5 to 50% and at least one hygroscopic carbohydrate in an amount from 50% to 95%.

31. The composition according to any of the preceding claims, wherein the benzodiazepine is remimazolam, and the hygroscopic excipient is selected from the group consisting of lactose, maltose, sucrose, trehalose and dextran.

32. A stable lyophilized or stable freeze-dried composition comprising at least one benzodiazepine according to formula (I)



with

W is H;

X is CH₂; n is 1;

Y is CH₂; m is 1;

Z is O; p is 0 or 1;

R¹ is CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂ or CH₂CH(CH₃)₂;


R² is 2-fluorophenyl, 2-chlorophenyl or 2-pyridyl;

R³ is Cl or Br;

R⁴, R⁵ and R⁶ form the group-CR⁸=U-V= wherein R⁸ is hydrogen, C₁₋₄ alkyl or C₁₋₃ hydroxyalkyl, U is N or CR⁹ wherein R⁹ is H, C₁₋₄ alkyl, C₁₋₃ hydroxyalkyl or C₁₋₄ alkoxy, V is N or CH and p is zero.

or a pharmaceutically acceptable salt thereof, wherein the composition comprises at least one pharmaceutically acceptable hygroscopic excipient, wherein the hygroscopic excipient is a carbohydrate and optionally the composition is at least in part amorphous.

33. A composition comprising at least one benzodiazepine according to formula (I)



with

W is H;

X is CH₂; n is 1;

Y is CH₂; m is 1;

Z is O; p is 0 or 1;

R¹ is CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂ or CH₂CH(CH₃)₂;

R² is 2-fluorophenyl, 2-chlorophenyl or 2-pyridyl;

R³ is Cl or Br;

R^4 , R^5 and R^6 form the group- $CR^8=U-V=$ wherein R^8 is hydrogen, C_{1-4} alkyl or C_{1-3} hydroxyalkyl, U is N or CR^9 wherein R^9 is H , C_{1-4} alkyl, C_{1-3} hydroxyalkyl or C_{1-4} alkoxy, V is N or CH and p is zero.

or a pharmaceutically acceptable salt thereof, wherein the composition comprises at least one pharmaceutically acceptable hygroscopic excipient, wherein the hygroscopic excipient is a carbohydrate and optionally the composition is at least in part amorphous and upon lyophilization or freeze-drying yields a stable lyophilized or stable freeze-dried composition.

34. The composition according to any of claims 32 or 33, wherein the stable composition is a composition in which less than 1% of the carboxylic ester moiety of the benzodiazepine is hydrolysed during storage for 13 weeks at storage conditions of 40°C and 75% relative humidity.

35. The composition according to any of claims 32 to 34, wherein the hygroscopic excipient is a substance which at 25°C and 1013.25 hPa reversibly binds water molecules.

36. Use of a mixture of at least one disaccharide and at least one dextran for preparing a pharmaceutical composition comprising a benzodiazepine with a carboxylic acid ester moiety as referred to in claim 1 or a pharmaceutically acceptable salt thereof, in particular remimazolam.

37. Method of preparing a pharmaceutical composition comprising the steps of

- a) providing a solution comprising a benzodiazepine as referred to in claim 1;
- b) adding at least one hygroscopic excipient, which is a carbohydrate selected from the group consisting of disaccharides and dextran or a mixture thereof with at least another hygroscopic excipient;
- c) drying the solution from step b) by lyophilization or spray-drying.

Fig. 1

Material	Supplier
Lactose monohydrate PhEur	DMV-Frontera Excipients
Mannitol PhEur	S. Black Ltd
Trehalose	Sigma-Aldrich
Sucrose PhEur	Fisher Scientific Ltd
Maltose monohydrate PhEur	S. Black Ltd
Dextran 70kD	Sigma-Aldrich
Povidone K17	IMCD Ltd
Glycine	Fisher Scientific Ltd
Hydrochloric acid SG 1.18	BDH Ltd
2M Sodium hydroxide	Fisher Scientific Ltd
Water for injection	Baxter Healthcare

Fig. 2

Formulation number	Formulation	REM mg/vial	excipient 1 mg/vial	excipient 2 mg/vial
01	REM only	16.32	--	--
02	REM/lactose (1:9)	8.16	73.44	--
03	REM/lactose (1:4.5)	8.16	36.72	--
04	REM/lactose (1:1)	8.16	8.16	--
05	REM/mannitol (1:9)	8.16	73.44	--
06	REM/trehalose (1:9)	8.16	73.44	--
07	REM/sucrose (1:9)	8.16	73.44	--
08	REM/maltose (1:9)	8.16	73.44	--
09	REM/dextran (1:9)	8.16	73.44	--
10	REM/lactose/mannitol (1:4.5:4.5)	8.16	36.72	36.72
11	REM/povidone (1:9)	8.16	73.44	--
12	REM/glycine (1:9)	8.16	73.44	--

Fig. 3

Formulation number	Formulation	excipient 1 mg/vial	excipient 2 mg/vial
02P	lactose	73.44	--
05P	mannitol	73.44	--
06P	trehalose	73.44	--
07P	sucrose	73.44	--
08P	maltose	73.44	--
09P	dextran	73.44	--
10P	lactose/mannitol	36.72	36.72
11P	povidone	73.44	--
12P	glycine	73.44	--

Fig. 4

Formulation number	Formulation	25°C/60%RH	40°C/75%RH
01	REM only	1.06	8.56
02	REM/lactose (1:9)	0.13	0.45
03	REM/lactose (1:4.5)	0.20	0.90
04	REM/lactose (1:1)	0.56	3.54
05	REM/mannitol (1:9)	1.62	8.23
06	REM/trehalose (1:9)	0.14	0.51
07	REM/sucrose (1:9)	0.10	0.22
08	REM/maltose (1:9)	0.14	0.45
09	REM/dextran (1:9)	0.16	0.57
10	REM/lactose/mannitol (1:4.5:4.5)	0.19	0.89
11	REM/povidone (1:9)	0.53	2.99
12	REM/glycine (1:9)	1.30	1.54

Fig. 5A

```

TITL 12161316      Compound CNS7056 Form 1
CELL 0.71073  7.687 29.261 12.376  90.000  97.788  90.000
ZERR 2 0.0001 0.0005 0.0003 0.0000 0.0008 0.0000
LATT -1
SYMM -X,Y+0.500,-Z
SFAC C      2.3100  20.8439  1.0200  10.2075  1.5886  0.5687  0.8650 =
          51.6512  0.2156  0.0033  0.0016  1.15  0.7700 12.0110
SFAC H      0.4930  10.5109  0.3229  26.1257  0.1402  3.1424  0.0408 =
          57.7998  0.0030  0.0000  0.0000  0.06  0.3200  1.0079
SFAC O      3.0485  13.2771  2.2868  5.7011  1.5463  0.3239  0.8670 =
          32.9089  0.2508  0.0106  0.0060  3.25  0.7700 15.9994
SFAC BR     17.1789  2.1723  5.2358  16.5796  5.6377  0.2609  3.9851 =
          41.4328  2.9557 -0.2901  2.4595 1000.00  1.1000 79.9040
SFAC N     12.2126  0.0057  3.1322  9.8933  2.0125 28.9975  1.1663 =
          0.5826 -11.5290  0.0061  0.0033  1.96  0.7700 14.0067
SFAC S      6.9053  1.4679  5.2034  22.2151  1.4379  0.2536  1.5863 =
          56.1720  0.8669  0.1246  0.1234  53.20  1.1100 32.0660
UNIT 108. 100. 20.  4. 16.  4.

S80  6  0.23964  0.43139  0.09908 11.00000  0.04634  0.03299 =
      0.04052  0.00002  0.01880 -0.00340
O81  3  0.16028  0.39374  0.15143 11.00000  0.06864  0.04111 =
      0.05255 -0.00210  0.02801  0.00002
O82  3  0.14598  0.47435  0.11207 11.00000  0.08099  0.03603 =
      0.04614  0.00545  0.03373 -0.00236
O83  3  0.42589  0.43401  0.12925 11.00000  0.05754  0.08564 =
      0.05198 -0.01536  0.01792 -0.00644
C84  1  0.20581  0.41866 -0.04324 11.00000  0.05949  0.04444 =
      0.02903  0.00359  0.01728  0.00704
C85  1  0.03624  0.41100 -0.09142 11.00000  0.06649  0.10092 =
      0.05586  0.01088  0.01751  0.00507
C86  1  0.00323  0.39810 -0.20187 11.00000  0.08670  0.14765 =
      0.05902 -0.02096 -0.03160 -0.00004
C87  1  0.14311  0.39209 -0.25693 11.00000  0.07916  0.11651 =
      0.06238 -0.01696  0.00195  0.02481
C88  1  0.30473  0.39806 -0.20987 11.00000  0.09246  0.09710 =
      0.04155  0.00157  0.01795  0.02685
C89  1  0.33456  0.41126 -0.10133 11.00000  0.05999  0.09817 =
      0.07178 -0.01451  0.00886  0.02173
S90  6  0.68868  0.81145  0.51625 11.00000  0.04072  0.02869 =
      0.05437  0.00158  0.00214  0.00223
O91  3  0.79129  0.77464  0.57315 11.00000  0.08025  0.03751 =
      0.04867 -0.00213 -0.00954  0.01626
O92  3  0.52601  0.81933  0.56122 11.00000  0.04778  0.05360 =
      0.06934 -0.00642  0.01702  0.00039
O93  3  0.78935  0.85213  0.50763 11.00000  0.07515  0.04369 =
      0.05025 -0.01354  0.01764 -0.01547
C94  1  0.62446  0.78970  0.38130 11.00000  0.04232  0.04028 =
      0.05049  0.00898  0.00929  0.00525
C95  1  0.74659  0.76959  0.32396 11.00000  0.06194  0.06998 =
      0.03238  0.00341 -0.00103  0.00990
C96  1  0.69911  0.75023  0.22476 11.00000  0.12417  0.10337 =
      0.03441  0.01537  0.02421  0.03314
C97  1  0.51941  0.75295  0.17732 11.00000  0.11897  0.11939 =
      0.02308 -0.01324 -0.00963 -0.00586
C98  1  0.40301  0.77268  0.23169 11.00000  0.06106  0.10242 =
      0.05463  0.00570 -0.01263 -0.00283
C99  1  0.45446  0.79193  0.33547 11.00000  0.05307  0.07089 =
      0.04982  0.00728 -0.00426 -0.01944
BR1  4  0.06011  0.52462  0.55140 11.00000  0.04153  0.05204 =
      0.07369 -0.00524  0.02434  0.00670
C2   1  0.25757  0.50395  0.49005 11.00000  0.02832  0.04536 =
      0.03350 -0.00752  0.01511  0.00763
C3   1  0.28921  0.45781  0.47911 11.00000  0.03135  0.03107 =
      0.04579  0.00145  0.00221 -0.00479

```

Fig. 5B

C4	1	0.42954	0.44393	0.43174	11.00000	0.03767	0.03461 =
		0.02980	-0.00320	-0.00151	-0.00125		
C5	1	0.54674	0.47556	0.39943	11.00000	0.03535	0.02939 =
		0.03479	-0.00390	0.00647	0.00183		
C6	1	0.51907	0.52242	0.41134	11.00000	0.04226	0.03479 =
		0.04333	-0.00172	0.00236	0.00188		
C7	1	0.37213	0.53602	0.45794	11.00000	0.03598	0.02793 =
		0.04586	-0.00044	0.01652	0.00336		
C8	1	0.64321	0.55824	0.38118	11.00000	0.03964	0.02453 =
		0.02719	0.00516	0.00457	0.00373		
C9	1	0.68998	0.59645	0.46059	11.00000	0.03743	0.03694 =
		0.04454	-0.00375	0.01588	0.00649		
N10	5	0.69097	0.58514	0.56581	11.00000	0.06070	0.03116 =
		0.04918	-0.00640	0.02020	-0.00054		
C11	1	0.74090	0.61847	0.63822	11.00000	0.06804	0.05787 =
		0.04752	-0.00600	0.01695	-0.00669		
C12	1	0.78515	0.66221	0.61053	11.00000	0.05480	0.04458 =
		0.05526	-0.02125	0.01554	-0.00787		
C13	1	0.77550	0.67229	0.50132	11.00000	0.04463	0.03102 =
		0.05452	0.00407	0.01432	-0.00038		
C14	1	0.73186	0.63955	0.42553	11.00000	0.04272	0.03021 =
		0.04282	-0.00243	0.01499	0.00270		
N15	5	0.71451	0.55972	0.29408	11.00000	0.04979	0.02502 =
		0.03692	0.00975	0.01748	0.00775		
C16	1	0.67500	0.52204	0.21324	11.00000	0.04463	0.02346 =
		0.04948	-0.00464	0.01738	0.00561		
C17	1	0.75857	0.47996	0.26673	11.00000	0.04549	0.02673 =
		0.01954	-0.00693	0.00506	-0.00121		
N18	5	0.70009	0.45973	0.35317	11.00000	0.03293	0.02806 =
		0.02597	-0.00088	0.00321	0.00207		
C19	1	0.81334	0.42409	0.39181	11.00000	0.03678	0.02848 =
		0.03351	-0.00426	0.00585	0.00488		
C20	1	0.93968	0.42402	0.32661	11.00000	0.03371	0.02802 =
		0.03711	0.00202	0.00106	0.00680		
N21	5	0.90585	0.45925	0.25315	11.00000	0.04775	0.03416 =
		0.02231	-0.01051	0.01052	-0.00308		
C22	1	0.79597	0.39511	0.48941	11.00000	0.03997	0.03711 =
		0.04548	0.01039	0.00508	0.00197		
C23	1	0.74788	0.53407	0.10940	11.00000	0.05650	0.04712 =
		0.03514	0.00836	0.00449	0.00605		
C24	1	0.68780	0.50047	0.01647	11.00000	0.08242	0.04077 =
		0.03001	-0.00046	0.01385	0.00523		
C25	1	0.71419	0.51690	-0.09234	11.00000	0.06429	0.06543 =
		0.03392	0.00018	0.00559	-0.00499		
O26	3	0.76261	0.55440	-0.11450	11.00000	0.12347	0.08282 =
		0.04188	0.01501	0.01658	-0.04001		
O27	3	0.65910	0.48459	-0.16756	11.00000	0.10340	0.06919 =
		0.03191	0.00253	0.01824	-0.00449		
C28	1	0.66642	0.49760	-0.27953	11.00000	0.19131	0.12699 =
		0.01390	-0.01417	0.02134	-0.05279		
BR51	4	1.06737	0.71057	0.98743	11.00000	0.03812	0.08781 =
		0.06774	0.00566	-0.00531	0.00447		
C52	1	0.84276	0.73306	0.93243	11.00000	0.03132	0.05952 =
		0.03819	0.00358	0.00226	-0.00263		
C53	1	0.81293	0.77906	0.93249	11.00000	0.04627	0.06820 =
		0.03723	-0.00581	0.00481	-0.00474		
C54	1	0.65043	0.79579	0.88269	11.00000	0.04551	0.03939 =
		0.04858	-0.00084	0.00376	-0.01071		
C55	1	0.51946	0.76552	0.84226	11.00000	0.04294	0.03573 =
		0.03413	0.00062	0.00952	-0.00208		
C56	1	0.54512	0.71765	0.84581	11.00000	0.02688	0.03659 =
		0.04586	-0.00025	0.00561	0.00047		
C57	1	0.71139	0.70186	0.88914	11.00000	0.03105	0.04840 =
		0.04447	-0.00668	-0.00429	0.00504		
C58	1	0.40956	0.68443	0.79765	11.00000	0.03348	0.02893 =
		0.04334	0.00070	0.00351	0.00421		

Fig. 5C

C59	1	0.38048	0.64253	0.86694	11.00000	0.03165	0.03488 =
		0.04951	0.00002	0.00425	0.00528		
N60	5	0.42879	0.64650	0.97247	11.00000	0.03542	0.05694 =
		0.03178	0.00872	0.00154	0.00467		
C61	1	0.38962	0.61026	1.03529	11.00000	0.04457	0.06338 =
		0.05765	0.01416	0.00707	0.00171		
C62	1	0.30187	0.57202	0.98967	11.00000	0.06548	0.04957 =
		0.11303	0.03456	0.03582	0.00696		
C63	1	0.25733	0.56863	0.88018	11.00000	0.07395	0.04664 =
		0.09803	0.00115	0.01240	-0.01007		
C64	1	0.29561	0.60475	0.81590	11.00000	0.08355	0.04152 =
		0.05459	-0.00010	0.00128	-0.02308		
N65	5	0.31344	0.68797	0.70771	11.00000	0.03846	0.03072 =
		0.04952	-0.00160	0.00032	0.00597		
C66	1	0.33129	0.72953	0.64125	11.00000	0.03574	0.02676 =
		0.05519	0.00406	0.00580	0.00330		
C67	1	0.26347	0.76733	0.70231	11.00000	0.03803	0.03316 =
		0.04166	0.01528	0.00868	0.00029		
N68	5	0.35122	0.78274	0.79764	11.00000	0.03387	0.03259 =
		0.05055	0.00549	0.00427	0.00218		
C69	1	0.24763	0.81583	0.84108	11.00000	0.05345	0.03305 =
		0.04570	0.00005	0.02067	-0.00546		
C70	1	0.09873	0.81841	0.77077	11.00000	0.04465	0.03799 =
		0.06107	0.00794	0.01464	0.00936		
N71	5	0.10819	0.78841	0.68720	11.00000	0.03892	0.03266 =
		0.05306	0.00974	0.01063	0.00803		
C72	1	0.30218	0.84064	0.94469	11.00000	0.08091	0.04934 =
		0.08052	-0.01505	0.02392	-0.00661		
C73	1	0.22541	0.72388	0.52948	11.00000	0.04039	0.05583 =
		0.03295	0.00047	0.00724	-0.00165		
C74	1	0.30154	0.68566	0.46508	11.00000	0.05896	0.05343 =
		0.05504	-0.00576	0.00667	0.02016		
C75	1	0.18003	0.67204	0.36587	11.00000	0.05296	0.05447 =
		0.04241	0.00546	0.01355	0.00171		
O76	3	0.06782	0.69497	0.31818	11.00000	0.05552	0.07543 =
		0.05719	-0.00702	-0.00194	0.02108		
O77	3	0.22119	0.62976	0.33149	11.00000	0.08466	0.04267 =
		0.04376	-0.00714	0.00726	0.00488		
C78	1	0.10717	0.61220	0.23887	11.00000	0.06302	0.09312 =
		0.07465	-0.02449	0.02418	-0.00980		
H611	2	10.42342	10.61111	11.10933	11.00000	0.06582	
H621	2	10.27371	10.54835	11.03412	11.00000	0.09086	
H631	2	10.20282	10.54235	10.84949	11.00000	0.08585	
H641	2	10.26600	10.60396	10.74163	11.00000	0.07058	
H661	2	10.45616	10.73494	10.63683	11.00000	0.04658	
H701	2	10.00528	10.83765	10.77749	11.00000	0.05724	
H721	2	10.20390	10.85662	10.96784	11.00000	0.10482	
H722	2	10.39143	10.86250	10.93477	11.00000	0.10500	
H723	2	10.34863	10.81975	11.00178	11.00000	0.10479	
H731	2	10.22647	10.75279	10.49048	11.00000	0.05050	
H732	2	10.10462	10.71635	10.53573	11.00000	0.05107	
H741	2	10.41143	10.69632	10.44327	11.00000	0.06599	
H742	2	10.32279	10.65905	10.51273	11.00000	0.06616	
H571	2	10.73613	10.67093	10.88928	11.00000	0.04893	
H531	2	10.89874	10.79871	10.96543	11.00000	0.05990	
H541	2	10.63029	10.82681	10.87790	11.00000	0.05285	
H161	2	10.54702	10.51731	10.19609	11.00000	0.04687	
H201	2	11.03302	10.40374	10.33036	11.00000	0.03977	
H221	2	10.90306	10.37871	10.51025	11.00000	0.06107	
H222	2	10.77354	10.41394	10.54853	11.00000	0.06102	
H223	2	10.70245	10.37370	10.47387	11.00000	0.06087	
H231	2	10.71028	10.56434	10.08666	11.00000	0.05487	
H232	2	10.87494	10.53365	10.12431	11.00000	0.05471	
H241	2	10.56546	10.49241	10.01723	11.00000	0.06095	
H242	2	10.75795	10.47323	10.02815	11.00000	0.06099	
H111	2	10.74728	10.61186	10.71244	11.00000	0.06882	
H121	2	10.81997	10.68398	10.66349	11.00000	0.06182	

Fig. 5D

H131	2	10.79812	10.70154	10.48020	11.00000	0.05215
H141	2	10.72939	10.64544	10.35226	11.00000	0.04595
H71	2	10.35042	10.56684	10.46668	11.00000	0.04408
H31	2	10.21444	10.43638	10.50355	11.00000	0.04223
H41	2	10.44931	10.41280	10.42055	11.00000	0.04056
H891	2	10.44977	10.41481	9.93226	11.00000	0.09285
H881	2	10.39917	10.39332	9.75106	11.00000	0.09266
H871	2	10.12372	10.38356	9.66972	11.00000	0.10194
H861	2	9.88808	10.39388	9.76390	11.00000	0.11607
H851	2	9.94416	10.41466	9.94909	11.00000	0.08904
H951	2	10.86472	10.76918	10.35546	11.00000	0.06580
H961	2	10.78321	10.73544	10.18942	11.00000	0.10497
H971	2	10.48493	10.74055	10.10914	11.00000	0.10604
H981	2	10.28646	10.77378	10.20054	11.00000	0.08719
H991	2	10.37377	10.80653	10.37249	11.00000	0.07037
H781	2	10.14480	10.58182	10.22240	11.00000	0.11588
H782	2	10.11102	10.63197	10.17669	11.00000	0.11581
H783	2	9.98883	10.61082	10.25546	11.00000	0.11600
H711	2	10.01359	10.78308	10.62464	11.00000	0.05205
H211	2	10.98261	10.46785	10.19729	11.00000	0.04161
H281	2	10.62358	10.47180	9.67092	11.00000	0.11566
H282	2	10.59036	10.52501	9.70225	11.00000	0.11566
H283	2	10.79029	10.50514	9.71088	11.00000	0.11566

Fig. 6A

```

TITL 1142055      Compound CNS7056 form 2
CELL  0.71073   8.921 11.154 25.834  90.000  90.000  90.000
ZERR   4 0.0001 0.0002 0.0004 0.0000 0.0000 0.0000
LATT  -1
SYMM  X+0.500,-Y+0.500,-Z
SYMM  -X,Y+0.500,-Z+0.500
SYMM  -X+0.500,-Y,Z+0.500
SFAC C      2.3100  20.8439  1.0200  10.2075  1.5886  0.5687  0.8650 =
          . 51.6512  0.2156  0.0033  0.0016  1.15  0.7700 12.0110
SFAC H      0.4930  10.5109  0.3229  26.1257  0.1402  3.1424  0.0408 =
          57.7998  0.0030  0.0000  0.0000  0.06  0.3200  1.0079
SFAC BR     17.1789  2.1723  5.2358  16.5796  5.6377  0.2609  3.9851 =
          41.4328  2.9557 -0.2901  2.4595 1000.00  1.1000 79.9040
SFAC N     12.2126  0.0057  3.1322  9.8933  2.0125 28.9975  1.1663 =
          0.5826 -11.5290  0.0061  0.0033  1.96  0.7700 14.0067
SFAC O      3.0485  13.2771  2.2868  5.7011  1.5463  0.3239  0.8670 =
          32.9089  0.2508  0.0106  0.0060  3.25  0.7700 15.9994
SFAC S      6.9053  1.4679  5.2034  22.2151  1.4379  0.2536  1.5863 =
          56.1720  0.8669  0.1246  0.1234  53.20  1.1100 32.0660
UNIT  108. 100.  4. 16. 20.  4.

BR1      3 -0.04819 -0.10880 -0.27710 11.00000  0.07032  0.03277 =
          0.03090  0.00144 -0.01238 -0.02224
C2      1 -0.15018 -0.21830 -0.32054 11.00000  0.02777  0.02177 =
          0.02345 -0.00009 -0.00209 -0.00471
C3      1 -0.17401 -0.18875 -0.37205 11.00000  0.02963  0.01861 =
          0.02702  0.00623  0.00188 -0.00107
C4      1 -0.24491 -0.26965 -0.40362 11.00000  0.02825  0.02442 =
          0.01718  0.00327  0.00106 -0.00145
C5      1 -0.29275 -0.37943 -0.38401 11.00000  0.02223  0.01822 =
          0.01875 -0.00067  0.00141  0.00066
C6      1 -0.27139 -0.40894 -0.33163 11.00000  0.02028  0.01967 =
          0.01926  0.00182  0.00105 -0.00153
C7      1 -0.20042 -0.32532 -0.29979 11.00000  0.02809  0.02763 =
          0.01685  0.00206  0.00190 -0.00055
C8      1 -0.32197 -0.52600 -0.30927 11.00000  0.01670  0.02233 =
          0.01945  0.00135 -0.00476 -0.00144
C9      1 -0.39853 -0.52353 -0.25770 11.00000  0.01623  0.02317 =
          0.01584  0.00259 -0.00384 -0.00281
N10     4 -0.46099 -0.41943 -0.24363 11.00000  0.02251  0.02613 =
          0.02353 -0.00189  0.00408  0.00155
C11     1 -0.52777 -0.41652 -0.19697 11.00000  0.02617  0.03441 =
          0.02357 -0.00451  0.00365  0.00346
C12     1 -0.53610 -0.51390 -0.16425 11.00000  0.02740  0.04329 =
          0.02040 -0.00335  0.00652 -0.00779
C13     1 -0.47518 -0.62062 -0.17997 11.00000  0.03584  0.03200 =
          0.02405  0.00767  0.00645 -0.00687
C14     1 -0.40334 -0.62685 -0.22730 11.00000  0.02879  0.02223 =
          0.02565  0.00090  0.00272 -0.00057
N15     4 -0.30040 -0.62781 -0.33049 11.00000  0.02151  0.02416 =
          0.01713  0.00287 -0.00002  0.00182
C16     1 -0.21928 -0.62991 -0.38036 11.00000  0.02330  0.02286 =
          0.01602  0.00057  0.00417  0.00450

```

Fig. 6B

C17	1	-0.32510	-0.57975	-0.41920	11.00000	0.02824	0.02308 =
		0.01704	-0.00121	0.00336	-0.00285		
N18	4	-0.36294	-0.46298	-0.41818	11.00000	0.02482	0.02037 =
		0.01483	0.00150	-0.00070	0.00079		
C19	1	-0.46920	-0.44117	-0.45641	11.00000	0.03022	0.02725 =
		0.01634	0.00325	0.00039	-0.00224		
C20	1	-0.49445	-0.54753	-0.47911	11.00000	0.03071	0.03401 =
		0.01669	0.00110	-0.00174	-0.00215		
N21	4	-0.40440	-0.63226	-0.45591	11.00000	0.03619	0.02354 =
		0.02146	-0.00463	0.00147	-0.00154		
C22	1	-0.54310	-0.32298	-0.46595	11.00000	0.03636	0.03429 =
		0.03074	0.00778	-0.00982	-0.00011		
C23	1	-0.15995	-0.75547	-0.39193	11.00000	0.03430	0.02640 =
		0.01793	-0.00359	0.00177	0.00554		
C24	1	-0.06166	-0.79435	-0.34621	11.00000	0.04707	0.03881 =
		0.02350	0.00041	0.00034	0.01530		
C25	1	0.06625	-0.87542	-0.35603	11.00000	0.03182	0.02650 =
		0.01948	0.00340	-0.00125	-0.00016		
O26	5	0.17233	-0.88334	-0.32760	11.00000	0.03778	0.06570 =
		0.03313	-0.01160	-0.01173	0.00417		
O27	5	0.05245	-0.94265	-0.39885	11.00000	0.03130	0.03874 =
		0.02467	-0.00799	-0.00330	0.01418		
C28	1	0.17574	-1.02443	-0.40865	11.00000	0.05622	0.08123 =
		0.03697	-0.01153	-0.00496	0.04396		
S80	6	-0.94275	-0.52899	-0.49624	11.00000	0.03340	0.02679 =
		0.02442	0.00000	0.00210	-0.00075		
O81	5	-0.83867	-0.47114	-0.53020	11.00000	0.05118	0.08336 =
		0.03575	0.02297	-0.00622	-0.02476		
O82	5	-1.08156	-0.46260	-0.49186	11.00000	0.04015	0.07788 =
		0.05503	-0.01022	-0.00539	0.01721		
O83	5	-0.97025	-0.65272	-0.50726	11.00000	0.13945	0.03230 =
		0.06071	-0.01467	0.01447	-0.00725		
C84	1	-0.86288	-0.52210	-0.43343	11.00000	0.02735	0.05893 =
		0.02832	0.01509	0.00686	-0.00534		
C85	1	-0.87781	-0.41462	-0.40588	11.00000	0.03763	0.08695 =
		0.03855	-0.01799	0.00427	-0.00754		
C86	1	-0.81420	-0.39965	-0.35764	11.00000	0.05438	0.16315 =
		0.04455	-0.02905	0.00147	-0.02905		
C87	1	-0.73766	-0.49241	-0.33773	11.00000	0.06202	0.20226 =
		0.06481	0.03510	-0.02105	-0.05062		
C88	1	-0.71885	-0.60444	-0.36221	11.00000	0.04217	0.17120 =
		0.11388	0.10762	-0.01320	-0.03729		
C89	1	-0.78500	-0.61610	-0.41251	11.00000	0.03725	0.08786 =
		0.07642	0.05538	-0.00772	-0.01074		
H891	2	9.22557	9.31210	9.56883	11.00000	0.08027	
H881	2	9.33331	9.33306	9.65289	11.00000	0.13097	
H851	2	9.06867	9.64846	9.57936	11.00000	0.06577	
H861	2	9.17563	9.67239	9.66111	11.00000	0.10509	
H161	2	9.86530	9.42517	9.62245	11.00000	0.02469	
H111	2	9.42959	9.65626	9.81326	11.00000	0.03383	
H121	2	9.41618	9.49292	9.86839	11.00000	0.03606	
H131	2	9.51614	9.31066	9.84059	11.00000	0.03697	
H141	2	9.64103	9.30191	9.76144	11.00000	0.03108	
H231	2	9.89972	9.24922	9.57680	11.00000	0.03066	
H232	2	9.75764	9.18723	9.60372	11.00000	0.03099	
H241	2	9.87585	9.16237	9.67759	11.00000	0.04434	
H242	2	9.97980	9.27746	9.67100	11.00000	0.04489	

Fig. 6C

H281	2	10.15353	8.92912	9.56085	11.00000	0.08666
H282	2	10.18989	8.92278	9.62053	11.00000	0.08723
H283	2	10.26566	9.02166	9.58620	11.00000	0.08710
H201	2	9.44027	9.43682	9.49457	11.00000	0.03327
H221	2	9.36727	9.66624	9.51370	11.00000	0.05146
H222	2	9.52479	9.72860	9.51527	11.00000	0.05104
H223	2	9.43193	9.71611	9.56601	11.00000	0.05131
H41	2	9.73983	9.74902	9.56204	11.00000	0.02807
H31	2	9.85823	9.88568	9.61518	11.00000	0.03001
H71	2	9.81367	9.65791	9.73490	11.00000	0.02870
H871	2	9.30621	9.51762	9.69480	11.00000	0.13226
H211	2	9.59801	9.29339	9.53630	11.00000	0.03270

Fig. 7A

S80	O81	1.454(5)Å
S80	O83	1.432(6)Å
C84	C85	1.376(12)Å
C85	C86	1.408(14)Å
C86	C87	1.360(16)Å
C87	C88	1.310(15)Å
C88	C89	1.386(14)Å
C89	H891	0.932Å
S90	O92	1.454(6)Å
S90	C94	1.793(8)Å
C94	C99	1.354(11)Å
C95	H951	0.938Å
C96	H961	0.934Å
C97	H971	0.924Å
C98	H981	0.927Å
Br1	C2	1.886(6)Å
C2	C7	1.381(9)Å
C3	H31	0.928Å
C4	H41	0.937Å
C5	N18	1.454(8)Å
C6	C8	1.498(9)Å
C8	C9	1.500(9)Å
C9	N10	1.343(9)Å
N10	C11	1.345(10)Å
C11	H111	0.933Å
C12	H121	0.927Å
C13	H131	0.918Å
N15	C16	1.492(9)Å
C16	C23	1.511(9)Å
C17	N18	1.352(8)Å
N18	C19	1.400(8)Å
C19	C22	1.496(9)Å
C20	H201	0.927Å
C22	H221	0.958Å
C22	H223	0.953Å
C23	H231	0.962Å
C24	C25	1.470(11)Å
C24	H242	0.962Å
C25	O27	1.354(10)Å
C28	H281	1.000Å
C28	H283	1.000Å
C52	C53	1.366(11)Å
C53	C54	1.404(11)Å
C54	C55	1.383(10)Å
C55	C56	1.414(9)Å
C56	C57	1.396(9)Å
C57	H571	0.925Å
C58	N65	1.254(8)Å
C59	C64	1.391(10)Å
C61	C62	1.386(14)Å
C62	C63	1.355(15)Å

S80	O82	1.468(5)Å
S80	C84	1.784(7)Å
C84	C89	1.318(12)Å
C85	H851	0.927Å
C86	H861	0.936Å
C87	H871	0.934Å
C88	H881	0.935Å
S90	O91	1.459(5)Å
S90	O93	1.431(5)Å
C94	C95	1.383(11)Å
C95	C96	1.356(13)Å
C96	C97	1.428(17)Å
C97	C98	1.323(15)Å
C98	C99	1.409(13)Å
C99	H991	0.924Å
C2	C3	1.382(9)Å
C3	C4	1.358(10)Å
C4	C5	1.388(9)Å
C5	C6	1.398(9)Å
C6	C7	1.394(9)Å
C7	H71	0.926Å
C8	N15	1.274(8)Å
C9	C14	1.386(9)Å
C11	C12	1.379(11)Å
C12	C13	1.375(11)Å
C13	C14	1.351(10)Å
C14	H141	0.921Å
C16	C17	1.500(9)Å
C16	H161	0.988Å
C17	N21	1.315(8)Å
C19	C20	1.344(9)Å
C20	N21	1.376(8)Å
N21	H211	1.000Å
C22	H222	0.950Å
C23	C24	1.536(11)Å
C23	H232	0.969Å
C24	H241	0.971Å
C25	O26	1.202(10)Å
O27	C28	1.445(10)Å
C28	H282	1.000Å
Br51	C52	1.886(7)Å
C52	C57	1.412(10)Å
C53	H531	0.927Å
C54	H541	0.921Å
C55	N68	1.427(9)Å
C56	C58	1.489(9)Å
C58	C59	1.530(10)Å
C59	N60	1.314(9)Å
N60	C61	1.372(10)Å
C61	H611	0.918Å
C62	H621	0.928Å

Fig. 7B

C63	C64	1.378(13)Å	C63	H631	0.932Å
C64	H641	0.917Å	N65	C66	1.485(8)Å
C66	C67	1.474(9)Å	C66	C73	1.516(10)Å
C66	H661	0.982Å	C67	N68	1.354(9)Å
C67	N71	1.334(8)Å	N68	C69	1.406(9)Å
C69	C70	1.343(11)Å	C69	C72	1.484(12)Å
C70	N71	1.366(10)Å	C70	H701	0.925Å
N71	H711	1.000Å	C72	H721	0.964Å
C72	H722	0.958Å	C72	H723	0.965Å
C73	C74	1.535(10)Å	C73	H731	0.975Å
C73	H732	0.967Å	C74	C75	1.493(12)Å
C74	H741	0.972Å	C74	H742	0.977Å
C75	O76	1.185(9)Å	C75	O77	1.360(9)Å
O77	C78	1.440(11)Å	C78	H781	0.965Å
C78	H782	0.966Å	C78	H783	0.960Å

Fig. 8A

O81	S80	O82	111.0(3)°
O82	S80	O83	114.4(4)°
O82	S80	C84	106.2(3)°
S80	C84	C85	117.7(6)°
C85	C84	C89	118.3(8)°
C84	C85	H851	119.626°
C85	C86	C87	118.1(10)°
C87	C86	H861	121.303°
C86	C87	H871	119.251°
C87	C88	C89	119.3(10)°
C89	C88	H881	120.264°
C84	C89	H891	118.485°
O91	S90	O92	111.7(3)°
O92	S90	O93	113.5(3)°
O92	S90	C94	105.7(3)°
S90	C94	C95	120.6(6)°
C95	C94	C99	119.3(8)°
C94	C95	H951	118.566°
C95	C96	C97	118.4(10)°
C97	C96	H961	121.695°
C96	C97	H971	119.699°
C97	C98	C99	120.8(9)°
C99	C98	H981	120.094°
C94	C99	H991	119.276°
Br1	C2	C3	121.0(5)°
C3	C2	C7	120.5(5)°
C2	C3	H31	120.203°
C3	C4	C5	120.6(6)°
C5	C4	H41	118.766°
C4	C5	N18	119.6(5)°
C5	C6	C7	117.8(6)°
C7	C6	C8	118.8(6)°
C2	C7	H71	119.721°
C6	C8	C9	117.5(5)°
C9	C8	N15	115.9(6)°
C8	C9	C14	121.2(6)°
C9	N10	C11	115.5(6)°
N10	C11	H111	118.526°
C11	C12	C13	117.4(7)°
C13	C12	H121	121.289°
C12	C13	H131	119.499°
C9	C14	C13	118.3(6)°
C13	C14	H141	121.419°
N15	C16	C17	105.9(5)°
C17	C16	C23	112.4(5)°
C17	C16	H161	109.539°
C16	C17	N18	122.7(6)°
N18	C17	N21	106.5(5)°
C5	N18	C19	127.0(5)°
N18	C19	C20	105.2(5)°
C20	C19	C22	129.4(6)°

O81	S80	O83	112.9(4)°
O81	S80	C84	105.5(3)°
O83	S80	C84	106.0(4)°
S80	C84	C89	123.6(7)°
C84	C85	C86	120.0(9)°
C86	C85	H851	120.377°
C85	C86	H861	120.636°
C86	C87	C88	121.8(10)°
C88	C87	H871	118.984°
C87	C88	H881	120.392°
C84	C89	C88	122.5(10)°
C88	C89	H891	119.061°
O91	S90	O93	112.8(4)°
O91	S90	C94	104.5(3)°
O93	S90	C94	108.0(3)°
S90	C94	C99	120.1(6)°
C94	C95	C96	121.6(9)°
C96	C95	H951	119.820°
C95	C96	H961	119.911°
C96	C97	C98	119.9(8)°
C98	C97	H971	120.397°
C97	C98	H981	119.080°
C94	C99	C98	119.9(9)°
C98	C99	H991	120.819°
Br1	C2	C7	118.5(5)°
C2	C3	C4	119.7(6)°
C4	C3	H31	120.109°
C3	C4	H41	120.600°
C4	C5	C6	120.6(6)°
C6	C5	N18	119.8(6)°
C5	C6	C8	123.3(6)°
C2	C7	C6	120.6(6)°
C6	C7	H71	119.679°
C6	C8	N15	126.6(6)°
C8	C9	N10	114.9(6)°
N10	C9	C14	123.9(6)°
N10	C11	C12	124.4(7)°
C12	C11	H111	117.061°
C11	C12	H121	121.279°
C12	C13	C14	120.4(6)°
C14	C13	H131	120.125°
C9	C14	H141	120.274°
C8	N15	C16	118.0(5)°
N15	C16	C23	109.4(5)°
N15	C16	H161	110.723°
C23	C16	H161	108.851°
C16	C17	N21	130.3(6)°
C5	N18	C17	123.1(5)°
C17	N18	C19	109.8(5)°
N18	C19	C22	125.3(6)°
C19	C20	N21	108.0(5)°

Fig. 8B

C19	C20	H201	126.017°	N21	C20	H201	126.026°
C17	N21	C20	110.5(5)°	C17	N21	H211	124.840°
C20	N21	H211	124.681°	C19	C22	H221	109.508°
C19	C22	H222	109.778°	H221	C22	H222	108.808°
C19	C22	H223	110.905°	H221	C22	H223	108.786°
H222	C22	H223	109.018°	C16	C23	C24	112.3(6)°
C16	C23	H231	109.392°	C24	C23	H231	108.812°
C16	C23	H232	108.378°	C24	C23	H232	109.105°
H231	C23	H232	108.825°	C23	C24	C25	114.3(7)°
C23	C24	H241	109.968°	C25	C24	H241	110.030°
C23	C24	H242	108.195°	C25	C24	H242	105.346°
H241	C24	H242	108.752°	C24	C25	O26	126.4(7)°
C24	C25	O27	109.4(7)°	O26	C25	O27	123.9(7)°
C25	O27	C28	115.2(7)°	O27	C28	H281	109.674°
O27	C28	H282	109.261°	H281	C28	H282	109.475°
O27	C28	H283	109.465°	H281	C28	H283	109.476°
H282	C28	H283	109.476°	Br51	C52	C53	119.3(6)°
Br51	C52	C57	119.0(5)°	C53	C52	C57	121.7(7)°
C52	C53	C54	118.9(7)°	C52	C53	H531	120.141°
C54	C53	H531	120.985°	C53	C54	C55	119.8(7)°
C53	C54	H541	120.227°	C55	C54	H541	120.000°
C54	C55	C56	122.1(6)°	C54	C55	N68	119.4(6)°
C56	C55	N68	118.5(6)°	C55	C56	C57	117.2(6)°
C55	C56	C58	123.2(6)°	C57	C56	C58	119.5(6)°
C52	C57	C56	120.2(7)°	C52	C57	H571	119.709°
C56	C57	H571	120.138°	C56	C58	C59	116.5(6)°
C56	C58	N65	126.7(6)°	C59	C58	N65	116.8(6)°
C58	C59	N60	116.3(6)°	C58	C59	C64	118.5(7)°
N60	C59	C64	125.0(7)°	C59	N60	C61	116.1(7)°
N60	C61	C62	121.7(8)°	N60	C61	H611	119.342°
C62	C61	H611	118.993°	C61	C62	C63	120.6(8)°
C61	C62	H621	120.029°	C63	C62	H621	119.353°
C62	C63	C64	118.4(9)°	C62	C63	H631	120.452°
C64	C63	H631	121.124°	C59	C64	C63	118.1(8)°
C59	C64	H641	120.844°	C63	C64	H641	121.057°
C58	N65	C66	118.2(6)°	N65	C66	C67	105.4(5)°
N65	C66	C73	109.7(5)°	C67	C66	C73	111.5(6)°
N65	C66	H661	109.122°	C67	C66	H661	108.890°
C73	C66	H661	112.017°	C66	C67	N68	121.8(6)°
C66	C67	N71	130.3(7)°	N68	C67	N71	107.4(6)°
C55	N68	C67	122.5(6)°	C55	N68	C69	128.7(6)°
C67	N68	C69	108.7(6)°	N68	C69	C70	105.5(6)°
N68	C69	C72	124.0(7)°	C70	C69	C72	130.5(7)°
C69	C70	N71	109.1(6)°	C69	C70	H701	125.444°
N71	C70	H701	125.502°	C67	N71	C70	109.2(6)°
C67	N71	H711	125.400°	C70	N71	H711	125.366°
C69	C72	H721	110.667°	C69	C72	H722	109.838°
H721	C72	H722	108.539°	C69	C72	H723	110.831°
H721	C72	H723	108.455°	H722	C72	H723	108.445°
C66	C73	C74	111.0(6)°	C66	C73	H731	108.535°

Fig. 8C

C74	C73	H731	110.248°		C66	C73	H732	110.751°
C74	C73	H732	108.249°		H731	C73	H732	108.042°
C73	C74	C75	112.4(6)°		C73	C74	H741	108.496°
C75	C74	H741	109.125°		C73	C74	H742	108.155°
C75	C74	H742	108.578°		H741	C74	H742	110.035°
C74	C75	O76	126.2(7)°		C74	C75	O77	110.7(7)°
O76	C75	O77	123.0(7)°		C75	O77	C78	115.6(7)°
O77	C78	H781	109.214°		O77	C78	H782	109.848°
H781	C78	H782	109.923°		O77	C78	H783	109.687°
H781	C78	H783	109.026°		H782	C78	H783	109.127°

Fig. 9

Br1	C2	1.892(3)Å
C2	C7	1.383(5)Å
C3	H31	0.938Å
C4	H41	0.921Å
C5	N18	1.428(4)Å
C6	C8	1.497(4)Å
C8	C9	1.497(4)Å
C9	N10	1.338(4)Å
N10	C11	1.345(4)Å
C11	H111	0.935Å
C12	H121	0.948Å
C13	H131	0.936Å
N15	C16	1.478(4)Å
C16	C23	1.527(5)Å
C17	N18	1.346(4)Å
N18	C19	1.391(4)Å
C19	C22	1.494(5)Å
C20	H201	0.912Å
C22	H221	0.965Å
C22	H223	0.960Å
C23	H231	0.969Å
C24	C25	1.478(5)Å
C24	H242	0.988Å
C25	O27	1.342(4)Å
C28	H281	0.964Å
C28	H283	0.962Å
S80	O82	1.447(3)Å
S80	C84	1.774(4)Å
C84	C89	1.369(7)Å
C85	H851	0.932Å
C86	H861	0.943Å
C87	H871	0.934Å
C88	H881	0.925Å

C2	C3	1.387(5)Å
C3	C4	1.371(5)Å
C4	C5	1.392(5)Å
C5	C6	1.406(4)Å
C6	C7	1.395(5)Å
C7	H71	0.924Å
C8	N15	1.276(4)Å
C9	C14	1.395(5)Å
C11	C12	1.378(5)Å
C12	C13	1.370(5)Å
C13	C14	1.382(5)Å
C14	H141	0.934Å
C16	C17	1.487(5)Å
C16	H161	0.976Å
C17	N21	1.320(4)Å
C19	C20	1.342(5)Å
C20	N21	1.378(5)Å
N21	H211	0.854Å
C22	H222	0.966Å
C23	C24	1.534(5)Å
C23	H232	0.981Å
C24	H241	0.960Å
C25	O26	1.201(4)Å
O27	C28	1.451(5)Å
C28	H282	0.965Å
S80	O81	1.431(3)Å
S80	O83	1.430(3)Å
C84	C85	1.400(7)Å
C85	C86	1.380(7)Å
C86	C87	1.342(13)Å
C87	C88	1.410(13)Å
C88	C89	1.433(10)Å
C89	H891	0.940Å

Fig. 10

Br1	C2	C3	119.3(3)°	Br1	C2	C7	118.9(3)°
C3	C2	C7	121.8(3)°	C2	C3	C4	119.0(3)°
C2	C3	H31	120.033°	C4	C3	H31	120.959°
C3	C4	C5	120.3(3)°	C3	C4	H41	119.485°
C5	C4	H41	120.261°	C4	C5	C6	121.0(3)°
C4	C5	N18	118.9(3)°	C6	C5	N18	120.1(3)°
C5	C6	C7	118.2(3)°	C5	C6	C8	122.3(3)°
C7	C6	C8	119.5(3)°	C2	C7	C6	119.7(3)°
C2	C7	H71	120.432°	C6	C7	H71	119.874°
C6	C8	C9	117.7(3)°	C6	C8	N15	124.4(3)°
C9	C8	N15	117.9(3)°	C8	C9	N10	116.6(3)°
C8	C9	C14	120.0(3)°	N10	C9	C14	123.4(3)°
C9	N10	C11	116.7(3)°	N10	C11	C12	123.7(3)°
N10	C11	H111	117.041°	C12	C11	H111	119.278°
C11	C12	C13	118.8(3)°	C11	C12	H121	120.443°
C13	C12	H121	120.783°	C12	C13	C14	119.3(3)°
C12	C13	H131	120.694°	C14	C13	H131	119.952°
C9	C14	C13	118.1(3)°	C9	C14	H141	120.942°
C13	C14	H141	120.983°	C8	N15	C16	117.6(3)°
N15	C16	C17	105.7(3)°	N15	C16	C23	110.8(3)°
C17	C16	C23	115.7(3)°	N15	C16	H161	107.681°
C17	C16	H161	107.726°	C23	C16	H161	108.910°
C16	C17	N18	120.7(3)°	C16	C17	N21	131.2(3)°
N18	C17	N21	108.0(3)°	C5	N18	C17	122.3(3)°
C5	N18	C19	128.6(3)°	C17	N18	C19	109.0(3)°
N18	C19	C20	105.7(3)°	N18	C19	C22	124.9(3)°
C20	C19	C22	129.3(3)°	C19	C20	N21	108.6(3)°
C19	C20	H201	127.007°	N21	C20	H201	124.433°
C17	N21	C20	108.7(3)°	C17	N21	H211	125.926°
C20	N21	H211	125.351°	C19	C22	H221	110.223°
C19	C22	H222	109.368°	H221	C22	H222	108.664°
C19	C22	H223	111.184°	H221	C22	H223	109.452°
H222	C22	H223	107.885°	C16	C23	C24	107.9(3)°
C16	C23	H231	107.712°	C24	C23	H231	110.073°
C16	C23	H232	111.123°	C24	C23	H232	109.430°
H231	C23	H232	110.583°	C23	C24	C25	118.8(3)°
C23	C24	H241	107.661°	C25	C24	H241	104.516°
C23	C24	H242	109.365°	C25	C24	H242	106.503°
H241	C24	H242	109.671°	C24	C25	O26	123.3(3)°
C24	C25	O27	114.4(3)°	O26	C25	O27	122.4(3)°
C25	O27	C28	115.2(3)°	O27	C28	H281	108.952°
O27	C28	H282	110.269°	H281	C28	H282	109.738°
O27	C28	H283	108.681°	H281	C28	H283	110.225°
H282	C28	H283	108.963°	O81	S80	O82	111.9(2)°
O81	S80	O83	115.1(2)°	O82	S80	O83	111.2(3)°
O81	S80	C84	106.30(18)°	O82	S80	C84	104.5(2)°
O83	S80	C84	107.0(2)°	S80	C84	C85	117.6(4)°
S80	C84	C89	122.1(4)°	C85	C84	C89	120.2(5)°
C84	C85	C86	121.6(6)°	C84	C85	H851	119.148°
C86	C85	H851	119.275°	C85	C86	C87	117.5(8)°
C85	C86	H861	121.859°	C87	C86	H861	120.606°
C86	C87	C88	124.9(7)°	C86	C87	H871	117.763°
C88	C87	H871	117.376°	C87	C88	C89	116.0(7)°
C87	C88	H881	122.592°	C89	C88	H881	121.435°
C84	C89	C88	119.8(8)°	C84	C89	H891	120.080°
C88	C89	H891	120.078°				

Fig. 11

Stability Data, CNS 7056 for Injection, 26 mg (Lot A01P310)

Product Name: CNS 7056 for Injection		Date Manufactured: July 2009		Container Closure: Type I, Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt				
Lot: A01P310								
				Storage Condition: 25 °C/60% RH				

Product Name: CNS 7056 for Injection		Date Manufactured: July 2009		Container Closure: Type I, Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt				
Lot: A01P310				Drug Substance Lot: 11084				
				Storage Condition: 25 °C/60% RH				
Tests^a	Acceptance Criteria^a	Time (Months)						
		0	1	3	6	9	12	18
Related Substances (Area %):								
CNS 7054X	NMT 1.5%	0.06	0.10	0.09	0.14	0.13	0.14	0.17
RRT 0.67	NMT 0.4%	0.03	0.04	0.03	0.03	0.03	0.03	0.03
RRT 0.70	NMT 0.4%	0.06	0.05	0.06	0.05	0.06	0.06	0.05
RRT 0.92	NMT 0.4%	0.11	0.12	0.11	0.11	0.11	0.10	0.11
RRT 1.31	NMT 0.4%	0.23	0.24	0.24	0.24	0.25	0.24	0.24
RRT 1.82	NMT 0.4%	0.03	<LOQ	<LOQ	0.03	0.03	0.03	<LOQ
Total	NMT 2.0%	0.52	0.55	0.53	0.60	0.61	0.60	0.60
Chiral (%R)	NMT 0.5%	0.27	—	0.28	0.28	—	0.25	—
Endotoxins USP <85>	NMT 0.5 EU/mg	<0.2 EU/mg	—	—	—	—	<0.2 EU/mg	—
Sterility USP <71>	Complies with test	Complies	—	—	—	—	Complies	—
These tests and acceptance criteria are based on a previous specification.								
Not performed.								

Fig. 13

Accelerated Stability Data, CNS 7056 for Injection, 26 mg (Lot A01P310)

Product Name: CNS 7056 for Injection		Date Manufactured: July 2009		Container Closure: Type I, Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt		
Lot: A01P310				Drug Substance Lot: 11084		
				Storage Condition: 40 °C/75% RH		
Tests ^a	Acceptance Criteria ^a	Time (Months)				
		0	1	3	6	
Appearance of Lyophilized Product	White to off white lyophilized powder.	Complies	Complies	Complies	Complies	
Appearance of Reconstituted Product	A clear, colorless to pale yellow solution, free from visible particulates.	Clear, colorless solution, essentially free from particulates.	Clear, very pale yellow solution, essentially free from particulates.	Clear, very pale yellow solution, essentially free from particulates.	Clear, very pale yellow solution, 1 vial contained small fiber	
Time to Reconstitute (min)	NMT 2 min	1-2 secs	1-2 secs	1-2 secs	1-2 secs	
Osmolality	290 ± 30 mOsmol/kg	281	-	279	295	
pH on Reconstitution	2.9 to 3.5	3.3	3.2	3.2	3.2	
Moisture Karl Fischer (% w/w)	NMT 1.5	0.16	0.32	0.37	0.59	
CNS 7056 Vial Content (mg/vial)	24.1 to 28.0 mg/vial	26.5	26.3	26.3	26.0	
	92.5 to 107.5% nominal amount	102.0	101.0	101.3	99.8	
Concentration on Reconstitution (mg/mL)	4.5 to 5.5 (Upright)	4.9	4.9	5.0	5.2	
	4.5 to 5.5 (Inverted)	4.8	4.8	4.8	5.1	
Subvisible particles	≥10 µm report per vial	≥10 µm:79	—	—	—	
	≥25 µm report per vial	≥25 µm:1	—	—	—	

Fig. 14

Accelerated Stability Data, CNS 7056 for Injection, 26 mg (Lot A01P310)						
Product Name: CNS 7056 for Injection		Container Closure: Type I, Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt				
Lot: A01P310						
		Storage Condition: 40 °C/75% RH				
Tests ²	Acceptance Criteria ³	Time (Months)				
		0	1	3	6	
Related Substances (Area %):						
CNS 7054X	NMT 1.5%	0.06	0.14	0.18	0.33	
RRT 0.67	NMT 0.4%	0.03	0.03	0.03	0.03	
RRT 0.70	NMT 0.4%	0.06	0.05	0.05	0.05	
RRT 0.92	NMT 0.4%	0.11	0.11	0.11	0.11	
RRT 1.31	NMT 0.4%	0.23	0.24	0.25	0.25	
RRT 1.82	NMT 0.4%	0.03	<LOQ	<LOQ	<LOQ	
Total	NMT 2.0%	0.52	0.57	0.62	0.77	
Chiral (%R)	NMT 0.5% R-enantiomer	0.27	—	0.28	0.28	
Endotoxins USP <8>	NMT 0.5 EU/mg	<0.2 EU/mg	—	—	—	
Sterility USP <71>	Complies with test	Complies	—	—	—	
⁴ =	These tests and acceptance criteria are based on a previous specification.					
— =	Not performed.					

Fig. 15

Long-Term Stability Data, CNS 7056 for Injection, 26 mg (Lot P310-01 (B))									
Product Name: CNS 7056 for Injection		Date Manufactured: May 2009			Container Closure: Type I, Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt				
Lot: P310-01 (B)					Drug Substance Lot: I1084				
					Storage Condition: 25 °C/60% RH				
Tests ^a	Acceptance Criteria ^a	Time (Months)							
		0	1	3	6	9	12	18	
Appearance of Lyophilized Product	White to off white lyophilized powder.	Complies	Complies	Complies	Complies	Complies	Complies	Complies	
Appearance of Reconstituted Product	A clear, colorless to pale yellow solution, free from visible particulates.	Very pale yellow, clear solution essentially free from particulates with several fibers not thought to be product related.	Clear, virtually colorless to very pale yellow solution, free from visible particulates. Several fibers visible not thought to be product related.	Very pale yellow, clear solution essentially free from particulates with several fibers not thought to be product related.	Very pale yellow, clear solution essentially free from particulates with several fibers not thought to be product related.	Very pale yellow, clear solution essentially free from particulates with several fibers not thought to be product related.	Very pale yellow, clear solution essentially free from particulates with several fibers not thought to be product related.	Very pale yellow, clear solution essentially free from particulates with several fibers not thought to be product related.	
Time to Reconstitute (Min)	NMT 2 min	1-2 secs	1-2 secs	1-2 secs	1-2 secs	1-2 secs	1-2 secs	1-2 secs	
Osmolality	290 ± 30 mOsmol/kg	289	280	277	281	282	286	278	
pH on Reconstitution	2.9 to 3.5	3.1	3.2	3.2	3.2	3.1	3.2	3.1	
Moisture Karl Fischer (% w/w)	NMT 1.5	0.39	0.60	0.62	0.64	0.55	0.76	0.86	
CNS 7056 Vial Content (mg/vial)	24.1 to 28.0 mg/vial	26.4	26.2	26.1	26.1	26.4	26.6	26.4	

Long-Term Stability Data, CNS 7056 for Injection, 26 mg (Lot P310-01 (B))									
Product Name:		Date Manufactured: May 2009							
CNS 7056 for Injection		Container Closure: Type I, Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt							
Lot: P310-01 (B)		Drug Substance Lot: 11084							
		Storage Condition: 25 °C/60% RH							
Tests ^a	Acceptance Criteria ^a	Time (Months)							
		0	1	3	6	9	12	18	
Concentration on Reconstitution (mg/mL)	92.5 to 107.5% nominal amount	101.7	100.6	100.5	100.5	101.5	102.2	101.4	
	4.5 to 5.5 (Upright)	5.0	5.0	4.9	4.9	5.1	4.9	5.1	
	4.5 to 5.5 (Inverted)	5.0	4.9	4.9	4.9	5.1	4.9	5.0	
Related Substances (Area %):									
CNS 7054X	NMT 1.5%	0.06	0.07	0.13	0.14	0.16	0.18	0.19	
RRT 0.68	NMT 0.40%	0.03	0.03	0.03	0.03	0.03	0.03	0.04	
RRT 0.70	NMT 0.40%	0.06	0.06	0.06	0.06	0.06	0.06	0.06	
RRT 0.92	NMT 0.40%	0.11	0.12	0.12	0.12	0.10	0.11	0.11	
RRT 0.94	NMT 0.40%	ND	0.03	ND	ND	ND	ND	ND	
RRT 1.31	NMT 0.40%	0.24	0.24	0.24	0.25	0.24	0.25	0.24	
RRT 1.85	NMT 0.40%	<LOQ	0.03	<LOQ	<LOQ	<LOQ	0.03	0.03	
Total	NMT 2.0%	0.50%	0.58%	0.58	0.60	0.59	0.66	0.67	
Chiral (%R)	NMT 0.5%	0.25%	—	—	0.30	—	0.3	—	
^a	=	tests and acceptance criteria are based on a previous specification.							
—	=	Not performed.							

Fig. 17

Accelerated Stability Data, CNS 7056 for Injection, 26 mg (Lot P310-01 (B))									
Product Name: CNS 7056 for Injection		Container Closure: Type I, Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt							
Lot: P310-01 (B)		Drug Substance Lot: 11084							
		Storage Condition: 40 °C/75% RH							
Tests ^a	Acceptance Criteria ^a	Time (Months)							
		0	1	3	6	9	12		
Appearance of Lyophilized Product	White to off white lyophilized powder.	Complies	Complies	Complies	Complies	Complies	Complies		
Appearance of Reconstituted Product	A clear, colorless to pale yellow solution, free from visible particulates.	Very pale yellow, clear solution essentially free from particulates with several fibers not thought to be product related.	Clear, virtually colorless to very pale yellow solution free from visible particulates. Several fibers visible not thought to be product related.	Very pale yellow, clear solution essentially free from particulates with several fibers not thought to be product related.	Very pale yellow, clear solution essentially free from particulates with several fibers not thought to be product related.	Very pale yellow, clear solution essentially free from particulates with several fibers not thought to be product related.	Very pale yellow, clear solution essentially free from particulates with several fibers not thought to be product related.		
Time to Reconstitute (min)	NMT 2 min	1-2 secs	1-2 secs	1-2 secs	1-2 secs	1-2 secs	1-2 secs		
Osmolality	290 ± 30 mOsmol/kg	289	281	274	281	284	281		
pH on Reconstitution	2.9 to 3.5	3.1	3.3	3.2	3.1	3.1	3.2		
Moisture Karl Fischer (% w/w)	NMT 1.5	0.39	0.51	0.58	0.84	0.82	0.82		
CNS 7056 Vial Content (mg/vial)	24.1 to 28.0 mg/vial	26.4	26.0	26.3	26.3	26.2	26.5		
	92.5 to 107.5% nominal amount	101.7	100.0	101.3	101.0	100.6	102.0		

Accelerated Stability Data, CNS 7056 for Injection, 26 mg (Lot P310-01 (B))									
Product Name: CNS 7056 for Injection		Container Closure: Type I, Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt							
Lot: P310-01 (B)		Drug Substance Lot: 11084							
		Storage Condition: 40 °C/75% RH							
Tests ^a	Acceptance Criteria ^a	Time (Months)							
		0	1	3	6	9	12		
Concentration on Reconstitution (mg/mL)	4.5 to 5.5 (Upright)	5.0	4.9	4.9	4.8	5.0			
	4.5 to 5.5 (Inverted)	5.0	4.9	4.9	4.9	5.0			4.9
Related Substances (Area %):									
RRT 0.51	NMT 0.4%	<LOQ	<LOQ	0.03	<LOQ	<LOQ			<LOQ
CNS 7054X	NMT 1.5%	0.06	0.11	0.22	0.36	0.45			0.68
RRT 0.63	NMT 0.4%	<LOQ	<LOQ	0.03	0.03	<LOQ			<LOQ
RRT 0.68	NMT 0.4%	0.03	0.03	0.03	0.03	0.03			0.03
RRT 0.70	NMT 0.4%	0.06	0.06	0.06	0.06	0.06			0.06
RRT 0.92	NMT 0.4%	0.11	0.12	0.12	0.12	0.11			0.11
RRT 0.94	NMT 0.4%	ND	0.03	ND	ND	ND			ND
RRT 1.31	NMT 0.4%	0.24	0.25	0.24	0.25	0.25			0.25
RRT 1.85	NMT 0.4%	<LOQ	0.03	<LOQ	0.03	<LOQ			0.03
Total	NMT 2.0%	0.50	0.63	0.73	0.88	0.90			1.16
Chiral (%R)	NMT 0.5%	0.25	—	0.3	0.3	—			0.3
^a	=	These tests and acceptance criteria are based on a previous specification.							
—	=	Not performed							

Fig. 19

Long-Term Stability Data, CNS 7056 for Injection, 25 mg (026CNS27)										
Product Name: CNS 7056		Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt								
Lot: 026CNS27		Drug Substance Lot: SOL12621/5								
		Storage Condition: 25 °C/60% RH								
Test	Acceptance Criteria	Time (Months)								
		0	1	2	3	6	9	12	18	
Appearance of Lyophilized Product	Record	White lyophilized plug with shrinkage. Although no layering evident, the bottom of the plug (2-3 mm) appeared to be slightly more shrunken	As initial	As initial	As initial	As initial	As initial	As initial	As initial	As initial
Completeness of Solution	A clear, colorless solution, free from visible particulates	A clear, colorless solution, free from visible particulates	As initial	—	As initial	As initial	As initial	As initial	As initial	As initial
Time to Reconstitute (min)	NMT 2 min	<1	<1	—	<1	<1	<1	<1	<1	<1
pH on Reconstitution	2.9 to 3.5	3.1	3.1	—	3.1	3.1	3.1	3.2	3.1	3.1
Moisture Karl Fischer (% w/w)	NMT 1.5	0.12	0.20	0.30	0.34	0.42	0.48	0.40	0.59	0.59
CNS 7056 Vial Content (mg/vial)	23.1 to 26.9 mg/vial	25.4	25.2	24.8	25.8	25.9	25.5	25.4	25.2	25.2
	92.5 to 107.5% nominal amount	101.6	100.8	99.2	103.2	103.6	102.0	101.6	100.8	100.8
Concentration on Reconstitution (mg/mL)	4.5 to 5.5	4.6	—	—	4.7	4.7	4.6	4.8	4.9	4.9

Fig. 21

Accelerated Stability Data, CNS 7056 for Injection, 25 mg (Lot 026CNS27)									
Product Name: CNS 7056					Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt				
Lot: 026CNS27								Drug Substance Lot: SOL12621/5	
Bulk Batch Size: 800 mL					Storage Condition: 40 °C/75% RH				
Test	Acceptance Criteria	Time (Months)							
			0	1	2	3	6		
Appearance of Lyophilized Product	Record	White lyophilized plug with shrinkage. Although no layering evident, the bottom of the plug (2-3 mm) appeared to be slightly more shrunken	As initial	As initial	As initial	As initial	As initial	As initial	
Completeness of Solution	A clear, colorless solution, free from visible particulates	A clear, colorless solution, free from visible particulates.	As initial	—	—	As initial	As initial	As initial	
Time to Reconstitute (min)	NMT 2 min	<1	<1	—	—	<1	<1	<1	
pH on Reconstitution	2.9 to 3.5	3.1	3.1	—	—	3.1	3.1	3.1	
Moisture Karl Fischer (% w/w)	NMT 1.5	0.12	0.28	0.46	0.54	0.54	0.64	0.64	
CNS 7056 Vial Content (mg/vial)	23.1 to 26.9 mg/vial	25.4	25.4	24.8	24.8	24.8	25.1	25.1	
	92.5 to 107.5% nominal amount	101.6	101.6	99.2	99.2	99.2	100.4	100.4	
Concentration on Reconstitution (mg/mL)	4.5 to 5.5	4.6	—	—	—	4.7	4.9	4.9	

Fig. 22

Accelerated Stability Data, CNS 7056 for Injection, 25 mg (Lot 026CNS27)									
Product Name: CNS 7056		Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt							
Lot: 026CNS27		Drug Substance Lot: SOL12621/5							
Bulk Batch Size: 800 mL		Storage Condition: 40 °C/75% RH							
Test	Acceptance Criteria	Time (Months)							
		0	1	2	3	6			
Related Substances (Area %):									
CNS 7054X	NMT 1.5%	1.12	1.16	1.18	1.21	1.32			
RRT 0.68	NMT 0.4%	0.09	0.09	0.09	0.08	0.08			
RRT 0.93	NMT 0.4%	0.13	0.13	0.12	0.14	0.13			
RRT 0.94	NMT 0.4%	0.07	0.09	0.09	0.11	0.07			
RRT 1.31	NMT 0.4%	0.24	0.23	0.24	0.23	0.22			
Total	NMT 2.0%	1.65	1.70	1.72	1.77	1.82			
Chiral (% R)	NMT 0.5% R-enantiomer	0.16	0.17	—	0.16	0.16			
— = Not performed.									

Fig. 23

Long-Term Stability Data, CNS 7056 for Injection, 25 mg (Lot G384)										
Product Name:		Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt								
CNS 7056										
Lot: G384		Drug Substance Lot: SOL12621/5								
		Storage Condition: 25 °C/60% RH								
Test	Acceptance Criteria	Time (Months)								
		0	1	2	3	6	9	12	18	
Appearance of Lyophilized Product	Record	White lyophilized plug with shrinkage. Although no splitting evident, the bottom of the plug (2-3 mm) appeared to be slightly more shrunken.	As initial	As initial	As initial	As initial	As initial	As initial	As initial	As initial
Completeness of Solution	A clear, colorless solution, free from visible particulates	A clear, colorless solution, free from visible particulates.	As initial	As initial	As initial	As initial	As initial	As initial	As initial	As initial
Time to Reconstitute (min)	NMT 2 min	<1	<1	<1	<1	<1	<1	<1	<1	<1
pH on Reconstitution	2.9 to 3.5	3.3	3.3	3.4	3.3	3.3	3.3	3.3	3.3	3.3
Moisture Karl Fischer (% w/w)	NMT 1.5	0.10	0.22	0.30	0.31	0.43	0.50	0.45	0.60	0.60

Long-Term Stability Data, CNS 7056 for Injection, 25 mg (Lot G384)												
Product Name: CNS 7056		Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt										
Lot: G384		Drug Substance Lot: SOL12621/5										
		Storage Condition: 25 °C/60% RH										
		Time (Months)										
Test	Acceptance Criteria	0	1	2	3	6	9	12	18			
CNS 7056 Vial Content (mg/vial)	23.1 to 26.9 mg/vial	23.4	23.7	23.8	23.7	23.7	23.6	23.8	23.8			
	92.5 to 107.5% nominal amount	93.6	94.8	95.2	94.8	94.8	94.4	95.2	95.2			
Concentration on Reconstitution (mg/mL)	4.5 to 5.5	4.4 ^a	—	4.5	—	4.8	—	4.6	4.5			
Related Substances (Area %):												
CNS 7054X	NMT 1.5%	1.08	1.10	1.10	1.11	1.11	1.14	1.13	1.16			
RRT 0.68	NMT 0.4%	0.09	0.09	0.09	0.09	0.10	0.10	0.08	0.08			
RRT 0.93	NMT 0.4%	0.12	0.12	0.14	0.14	0.12	0.12	0.13	0.18			
RRT 0.94	NMT 0.4%	0.07	0.09	0.10	0.10	^b	^b	<LOQ	^c			
RRT 1.31	NMT 0.4%	0.23	0.24	0.24	0.24	0.24	0.24	0.22	0.23			
Total	NMT 2.0%	1.59	1.64	1.67	1.68	1.57	1.60	1.56	1.65			
Chiral (% R)	NMT 0.5% R-enantiomer	0.16	0.16	—	0.16	0.17	0.15	0.14	0.16			
^a	= See Section 7.2.13.1 for discussion.											
^b	= Peak at RRT 0.94 not resolved from CNS 7056.											
^c	= Peak not detected. Appeared to be unresolved from previous peak.											
—	= Not performed.											

Fig. 25

Accelerated Stability Data, CNS 7056 for Injection, 25 mg (Lot G384)									
Product Name: CNS 7056				Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt					
Lot: G384				Drug Substance Lot: SOL12621/5					
				Storage Condition: 40 °C/75% RH					
Test	Acceptance Criteria	Time (Months)							
		0	1	2	3	6			
Appearance of Lyophilized Product	Record	White lyophilized plug with shrinkage. Although no splitting evident, the bottom of the plug (2-3 mm) appeared to be slightly more shrunken	As initial	As initial	As initial	As initial	As initial		
Completeness of Solution	A clear, colorless solution, free from visible particulates	A clear, colorless solution, free from visible particulates.	As initial	As initial	As initial	As initial			
Time to Reconstitute (min)	NMT 2 min	<1	<1	<1	<1	<1			
pH on Reconstitution	2.9 to 3.5	3.3	3.3	3.4	3.4	3.3			
Moisture Karl Fischer (% w/w)	NMT 1.5	0.10	0.48	0.44	0.50	0.69			
CNS 7056 Vial Content (mg/vial)	23.1 to 26.9 mg/vial	23.4	23.6	23.8	23.7	23.7			
	92.5 to 107.5% nominal amount	93.6	94.4	95.2	94.8	94.8			
Concentration on Reconstitution (mg/mL)	4.5 to 5.5	4.4 ^a	—	4.5	—	4.7			

Accelerated Stability Data, CNS 7056 for Injection, 25 mg (Lot G384)									
Product Name: CNS 7056		Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt							
Lot: G384		Drug Substance Lot: SOL12621/5							
		Storage Condition: 40 °C/75% RH							
Test	Acceptance Criteria	Time (Months)							
		0	1	2	3	6			
Related Substances (Area %):									
CNS 7054X	NMT 1.5%	1.08	1.14	1.14	1.18	1.28			
RRT 0.68	NMT 0.4%	0.09	0.08	0.08	0.08	0.10			
RRT 0.93	NMT 0.4%	0.12	0.12	0.14	0.14	0.12			
RRT 0.94	NMT 0.4%	0.07	0.09	0.12	0.10	0.12			
RRT 1.31	NMT 0.4%	0.23	0.24	0.23	0.24	0.23			
Total	NMT 2.0%	1.59	1.67	1.71	1.74	1.73			
Chiral (% R)	NMT 0.5% R-enantiomer	0.16	0.14	—	0.16	0.16			
^a	=	See Section 7.2.13.1 for discussion.							
^b	=	Peak at RRT 0.94 not resolved from CNS 7056.							
—	=	Not performed.							

Fig. 27

Long-Term Stability Data, CNS 7056 for Injection, 23 mg (Lot P02308)											
Product Name: CNS 7056		Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt									
Lot: P02308		Drug Substance Lot: 11084									
		Storage Condition: 25 °C/60% RH									
Test	Acceptance Criteria	Time (Months)									
		0	1	3	6	9	12	18	24	36	
Appearance of Lyophilized Product	White to off-white lyophilized plug	Off-white lyophilized plug	As initial	As initial	As initial	As initial	As initial	As initial	As initial	As initial	
Completeness of Solution	A clear, colorless to pale yellow solution, free from visible particulates	All vials - a clear, very pale yellow solution, free from visible particulates	As initial	As initial	As initial	As initial	As initial	As initial	As initial	As initial	
Time to Reconstitute (min)	NMT 2 min	<1	<1	<1	<1	<1	<1	<1	<1	<1	
Subvisible Particulates	≥10 µm: report	66	65	—	93	—	24	16	13	40	
	≥25 µm: report	1	0	—	<1	—	<1	<1	0	<1	
Osmolality (mOsm/kg)	290 ± 30	280	283	292	288	289	287	282	284	288	
pH on Reconstitution	2.9 to 3.5	3.1	3.1	3.1	3.2	3.2	3.1	3.2	3.1	3.1	
Moisture Karl Fischer (%) w/w	NMT 1.5	0.27	0.27	0.36	0.50	0.47	0.46	0.62	0.59	0.68	
CNS 7056 Vial Content (mg/vial)	21.3 to 24.7 mg/vial	23.6	23.6	23.4	23.6	23.6	23.5	23.5	23.4	23.4	
	92.5 to 107.5% nominal amount	102.6	102.6	101.7	102.6	102.3	102.2	102.1	101.5	101.6	

Long-Term Stability Data, CNS 7056 for Injection, 23 mg (Lot P02308)														
Product Name: CNS 7056			Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt											
Lot: P02308			Drug Substance Lot: 11084											
			Storage Condition: 25 °C/60% RH											
Test	Acceptance Criteria	Time (Months)												
		0	1	3	6	9	12	18	24	36				
Test	Acceptance Criteria	Time (Months)												
Concentration on Reconstitution (mg/mL)	4.5 to 5.5	0	1	3	6	9	12	18	24	36				
		4.9	5.1	5.0	5.0	5.1	5.0	5.1	5.1	5.0				
Related Substances (Area %):														
CNS 7054X	NMT 1.5%	0.07	0.10	0.11	0.13	0.16	0.16	0.20	0.23	0.29				
RRT 0.67	NMT 0.4%	0.03	0.03	0.04	0.03	0.03	0.03	0.03	0.03	0.03				
RRT 0.71	NMT 0.4%	0.06	0.06	0.06	0.06	0.05	0.05	0.06	0.06	0.06				
RRT 0.92	NMT 0.4%	0.12	0.12	0.12	0.11	0.12	0.12	0.11	0.11	0.11				
RRT 0.96	NMT 0.4%	0.12	<LOQ	0.10	0.04	0.09	0.10	0.04	0.03	0.03				
RRT 1.31	NMT 0.4%	0.25	0.27	0.25	0.25	0.25	0.26	0.26	0.25	0.25				
RRT 1.80	NMT 0.4%	<LOQ	<LOQ	<LOQ	0.03	<LOQ	<LOQ	<LOQ	0.03	0.03				
Total	NMT 2.0%	0.65	0.58	0.68	0.65	0.70	0.72	0.70	0.74	0.80				
Chiral (% R)	NMT 0.5% R-enantiomer	0.26	0.27	—	0.27	—	0.27	0.27	0.26	0.28				
Sterility	Complies	Complies	—	—	—	—	Complies	—	Complies	Complies				
— =		Not performed.												

Fig. 29

Accelerated Stability Data, CNS 7056 for Injection, 23 mg (Lot P02308)									
Product Name: CNS 7056		Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt			Drug Substance Lot: 11084				
Lot: P02308					Storage Condition: 40 °C/75% RH				
Test	Acceptance Criteria	Time (Months)							
		0	1	3	6	9	12		
Appearance of Lyophilized Product	White to off-white lyophilized plug	Off-white lyophilized plug.	As initial	As initial	As initial	As initial	As initial	As initial	As initial
Completeness of Solution	A clear, colorless to pale yellow solution, free from visible particulates	A clear, very pale yellow solution free from visible particulates.	As initial	As initial	As initial	As initial	As initial	As initial	As initial
Time to Reconstitute (min)	NMT 2 min	<1	<1	<1	<1	<1	<1	<1	<1
Subvisible Particulates	≥10 µm: report	66	76	12	74	—	20	—	20
	≥25 µm: report	1	<1	0	<1	—	0	—	0
pH on Reconstitution	2.9 to 3.5	3.1	3.1	3.1	3.2	3.2	3.1	3.2	3.1
Osmolality (mOsm/kg)	290 ± 30	280	289	291	286	290	288	286	288
Moisture Karl Fischer (% w/w)	NMT 1.5	0.27	0.40	0.58	0.67	0.78	0.86	0.67	0.86
CNS 7056 Vial Content (mg/vial)	21.3 to 24.7 mg/vial	23.6	23.6	23.4	23.5	23.4	23.2	23.5	23.2
	92.5 to 107.5% nominal amount	102.6	102.6	101.7	102.2	101.7	101.0	102.2	101.0

Accelerated Stability Data, CNS 7056 for Injection, 23 mg (Lot P02308)									
Product Name: CNS 7056		Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt							
Lot: P02308		Drug Substance Lot: 11084							
		Storage Condition: 40 °C/75% RH							
Test	Acceptance Criteria	Time (Months)							
		0	1	3	6	9	12		
Concentration on Reconstitution (mg/mL)	4.5 to 5.5	4.9	5.2	5.0	5.0	5.1	5.1		
Related Substances (Area %):									
CNS 7054X	NMT 1.5%	0.07	0.12	0.20	0.32	0.41	0.54		
RRT 0.67	NMT 0.4%	0.03	0.03	0.04	0.03	0.03	0.03		
RRT 0.71	NMT 0.4%	0.06	0.07	0.05	0.06	0.05	0.05		
RRT 0.92	NMT 0.4%	0.12	0.12	0.12	0.11	0.12	0.12		
RRT 0.96	NMT 0.4%	0.12	0.03	0.10	0.05	0.10	0.09		
RRT 1.31	NMT 0.4%	0.25	0.27	0.27	0.28	0.29	0.29		
RRT 1.80	NMT 0.4%	<LOQ	<LOQ	<LOQ	0.03	<LOQ	<LOQ		
Total	NMT 2.0%	0.65	0.64	0.78	0.88	1.00	1.12		
Chiral (% R)	NMT 0.5% R-enantiomer	0.26	0.27	0.27	0.27	—	0.27		
Sterility	Complies	Complies	—	—	—	—	—		
— = Not performed									—

Fig. 31

Long-Term Stability Data, CNS 7056 for Injection, 25 mg (Lot 025CNS27)												
Product Name: CNS 7056		Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt										
Lot: 025CNS27		Drug Substance Lot: SOL12621/5										
		Storage Condition: 25 °C/60% RH										
Test	Acceptance Criteria	Plug	Time (Months)									
			0	1	2	3	6	9	12	18		
Appearance of Lyophilized Product	Record	Unsplit	White lyophilized plug with shrinkage.	As initial	—	—	—	—	—	—	—	—
		Split	White lyophilized plug with shrinkage. 2 distinct layers evident.	As initial	As initial	As initial	As initial	As initial	As initial	As initial	White lyophilized plug with shrinkage.	
Completeness of Solution	A clear, colorless solution, free from visible particulates		The bulk of the plug dissolved rapidly. The resultant solution was clear, colorless, and free from visible particulates.	As initial	As initial	As initial	As initial	As initial	As initial	As initial	As initial	As initial
		Time to Reconstitute (Min)	NMT 2 min	Unsplit	<1	—	—	—	<1 ^b	<1	<1 ^b	<1 ^b
pH on Reconstitution	2.9 to 3.5	Split	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
		Unsplit	3.1	—	—	—	—	—	3.1 ^b	3.1	3.1 ^b	3.2 ^b
		Split	3.2	3.1	3.2	3.1	3.2	3.2	3.1	3.1	3.1	3.1

Fig. 32

Long-Term Stability Data, CNS 7056 for Injection, 25 mg (Lot 025CNS27)												
Product Name: CNS 7056		Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt										
Lot: 025CNS27		Drug Substance Lot: SOL12621/5										
		Storage Condition: 25 °C/60% RH										
Test	Acceptance Criteria	Plug	Time (Months)									
			0	1	2	3	6	9	12	18		
Moisture Karl Fischer (% w/w)	NMT 1.5	Unsplit	0.36	0.31	—	—	—	0.79 ^a	0.76	0.62 ^b		
		Split	0.83	0.78	0.86	0.87	0.87	—	—	—		
CNS 7056 Vial Content (mg/vial)	23.1 to 26.9 mg/vial	Unsplit	24.6	25.0	24.7	24.4	24.7	24.8	25.0	25.0		
			98.5	100.0	98.7	97.6	98.7	99.2	100.0	100.0		
Concentration on Reconstitution (mg/mL)	4.5 to 5.5	Unsplit	4.6	—	—	4.6	4.7	4.8	4.8	4.7		
Related Substances (Area %): CNS 7054X RRT 0.68 RRT 0.93 RRT 0.94 RRT 1.31 Total												
	NMT 1.5%	Unsplit	1.08	1.09	—	—	—	—	—	—		
	NMT 0.4%		0.09	0.09	—	—	—	—	—	—		
	NMT 0.4%		0.12	0.13	—	—	—	—	—	—		
	NMT 0.4%		0.07	0.08	—	—	—	—	—	—		
	NMT 0.4%		0.22	0.24	—	—	—	—	—	—		
Total			1.58	1.628	—	—	—	—	—	—		

Long-Term Stability Data, CNS 7056 for Injection, 25 mg (Lot 025CNS27)												
Product Name: CNS 7056		Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt										
Lot: 025CNS27		Drug Substance Lot: SOL12621/S										
		Storage Condition: 25 °C/60% RH										
Test	Acceptance Criteria	Plug	Time (Months)									
			0	1	2	3	6	9	12	18		
Related Substances (Area %):												
CNS 7054X	NMT 1.5%	Split	1.08	1.09	1.12	1.14	1.15	1.20			1.20	1.16
RRT 0.68	NMT 0.4%		0.09	0.09	0.11	0.08	0.09	0.08			0.10	0.10
RRT 0.93	NMT 0.4%		0.12	0.13	0.20	0.12	0.14	0.12			0.12	0.13
RRT 0.94	NMT 0.4%		0.07	0.08	^c	0.09	0.09	^d			^d	<0.05
RRT 1.31	NMT 0.4%		0.22	0.23	0.24	0.24	0.24	0.23			0.24	0.24
Total	NMT 2.0%		1.58	1.62	1.67	1.67	1.71	1.63			1.66	1.63
Chiral (% R)	NMT 0.5% R-enantiomer	Unsplit	0.16	0.16	—	0.12	0.13				0.15	0.16
^a	= Two distinct layers evident in only 9 of 11 vials.											
^b	= At 39 and 78 weeks, it is not known whether the vials used for the analysis were unsplit or split.											
^c	= Peaks at RRT 0.93 and 0.94 are not resolved; total area remains the same.											
^d	= Peak at RRT 0.94 not resolved from CNS 7056 peak.											
—	= Not performed.											

Fig. 34

Accelerated Stability Data, CNS for Injection, 25 mg (Lot 025CNS27)									
Product Name: CNS 7056		Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt							
Lot: 025CNS27		Drug Substance Lot: SOL12621/5							
		Storage Condition: 40 °C/75% RH							
Test	Acceptance Criteria	Plug	Time (Months)						
			0	1	2	3	6	9	12
Appearance of Lyophilized Product	Record	Unsplit	White lyophilized plug with shrinkage.	As initial	As initial	As initial	As initial	As initial	As initial
		Split	White lyophilized plug with shrinkage. 2 distinct layers evident.	As initial	As initial	As initial	As initial	As initial	White lyophilized plug with shrinkage.
Completeness of Solution	A clear, colorless solution, free from visible particulates	Unsplit	The bulk of the plug dissolved rapidly. The resultant solution was clear, colorless and free from visible particulates.	As initial	As initial	As initial	As initial	As initial	As initial
Time to Reconstitute (min)	NMT 2 min	Unsplit	<1	<1	—	—	—	<1 ^a	<1 ^a
		Split	<1	<1	<1	<1	<1		
pH on Reconstitution	2.9 to 3.5	Unsplit	3.1	3.2	—	—	—	3.1 ^a	3.1 ^a
		Split	3.2	3.1	3.3	3.1	3.2		

Fig. 35

Accelerated Stability Data, CNS for Injection, 25 mg (Lot 025CNS27)											
Product Name: CNS 7056		Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt									
Lot: 025CNS27		Drug Substance Lot: SOL12621/5									
		Storage Condition: 40 °C/75% RH									
		Time (Months)									
Test	Acceptance Criteria	Plug	Time (Months)								
			0	1	2	3	6	9	12		
Moisture Karl Fischer (% ww)	NMT 1.5	Unsplit	0.36	0.47	—	—	—	1.24 ^a	0.82		
		Split	0.83	1.18	1.00	0.78	1.11				
CNS 7056 Vial Content (mg/vial)	23.1 to 26.9 mg/vial	Unsplit	24.6	24.6	24.4	24.1	24.6	24.3	24.6		
			98.5	98.4	97.5	96.4	98.3	97.0	98.4		
	92.5 to 107.5% nominal amount										

Accelerated Stability Data, CNS for Injection, 25 mg (Lot 025CNS27)											
Product Name: CNS 7056		Container Closure: Type I Glass Vials, 20 mL; Bromobutyl Rubber Stopper; and Aluminum Skirt									
Lot: 025CNS27		Storage Condition: 40 °C/75% RH Drug Substance Lot: SOL12621/5									
Test	Acceptance Criteria	Plug	Time (Months)								
			0	1	2	3	6	9	12		
Conc. on Reconst. (mg/mL)	4.5 to 5.5	Unsplit	4.6	4.6	4.5	4.6	4.6	4.9			
Rel. subst. (Area %)											
CNS 7054X	NMT 1.5%	Unsplit	1.08	1.15	—	—	—	—	—	—	
RRT 0.68	NMT 0.4%		0.09	0.09	—	—	—	—	—	—	
RRT 0.93	NMT 0.4%		0.12	0.13	—	—	—	—	—	—	
RRT 0.94	NMT 0.4%		0.07	0.08	—	—	—	—	—	—	
RRT 1.31	NMT 0.4%		0.22	0.24	—	—	—	—	—	—	
Total	NMT 2.0%		1.58	1.67	—	—	—	—	—	—	
Rel. Sub. (Area %):											
CNS 7054X	NMT 1.5%	Split	1.08	1.15	1.24	1.25	1.36	1.40		1.51	
RRT 0.68	NMT 0.4%		0.09	0.09	0.11	0.08	0.09	0.08		0.10	
RRT 0.93	NMT 0.4%		0.12	0.13	0.20	0.12	0.13	0.11		0.12	
RRT 0.94	NMT 0.4%		0.07	0.08	^d	0.09	0.10	^d		^d	
RRT 1.31	NMT 0.4%		0.22	0.24	0.24	0.23	0.24	0.23		0.23	
Total	NMT 2.0%		1.58	1.69	1.79	1.77	1.92	1.82		1.96	
Chiral (% R)	NMT 0.5% R-enantiomer	Unsplit	0.16	0.16	—	0.12	0.16	0.15		0.15	
^a	= It is not known whether the vials used for the analysis were unsplit or split.										
^b	= Not done due to shortage of vials stored at 40 °C/75% RH.										
^c	= It is not known whether the vials used for the analysis were unsplit or split; aligned with split data for ease of comparison.										
^d	= Peaks at RRT 0.93 and 0.94 were not resolved.										
—	= Not performed.										

Fig. 37

Test (Release Specification and Shelf Life Specification)	Method	Results
Appearance of Lyophilised Product (White to off white lyophilised plug)	C/TE/8257	25°C/60% RH All vials off white lyophilised plug
Completeness of Solution (on reconstitution) (A clear, colourless to pale yellow solution free from visible particulates)	USP <641>	All vials - a clear, colourless to pale yellow solution, free from visible particulates
Time to reconstitute (NMT 2 minutes)	C/TE/8256	Vial 1: 38 secs Vial 2: 30 secs Mean : 34 secs
Identity CNS7056B UV (Absorbance maxima at 216±1nm and 230.5±1 nm)	QC058	NR
HPLC retention achiral method Identity CNS7056B (Retention time practically identical to reference standard)	ADP104 achiral HPLC	Retention time practically identical to reference standard
HPLC retention chiral method Identity CNS7056B (Retention time practically identical to reference standard)	ADP104 chiral HPLC	Retention time practically identical to reference standard
Sterility (Complies)	USP <71>	Sterile
Sub-visible particulates (≥10µm : Record ≥25µm : Record)	USP <788>	≥10µm : 40 particles ≥25µm : <1 particle

NR: Test not required as part of stability programme.

Fig. 38

Test (Release Specification)	Method	Results
		25°C/60%RH
pH on reconstitution (2.9 – 3.5)	USP<791>	Vial 1: pH 3.13 Vial 2: pH 3.15 Mean: pH 3.14
Osmolality (290 ± 30 mOsm/Kg)	USP<785>	Vial 1: 289mOsm/kg Vial 2: 287mOsm/kg Mean: 288mOsm/kg
CNS 7056B Vial content/Recovery (Release 21.9 to 24.2 mg per vial, 95.0-105.0% of stated amount) (Shelf Life 21.3 to 24.7 mg per vial, 92.5-107.5% of stated amount)	ADP104 achiral HPLC	Vial 1: 23.2 mg per vial (101.0%) Vial 2: 23.5 mg per vial (102.2%) Mean: 23.4 mg per vial (101.6 %)
Chiral Purity (% area) (Release and Shelf Life NMT 0.5% R enantiomer)	ADP104 chiral HPLC	0.28%
Moisture by Karl Fischer (% w/w) (Release NMT1.0%) (Shelf Life NMT1.5%)	ADP104 KF	Vial 1: 0.71% Vial 2: 0.67% Vial 3: 0.67% Mean: 0.68%

NR: Test not required as part of stability programme.

Fig. 39




Fig. 40




Fig. 41




Figure 42

Initial

Test Parameters	Specification	12PM529-8-2	12PM529-8-1	PM0237/12	12PM529-9-1
Appearance	White to off white powder	White powder	White powder	White powder	White powder
Assay (mg/vial)	24.94 – 27.66	25.97	26.11	26.25	N/A
% Assay	95%-105%	98.73%	99.29%	99.79%	98.75%
Total Impurities	2%	0.73%	0.67%	0.48%	0.97%
Impurities Above 0.045%	N/A	0.43%	0.39%	0.32%	0.76%
Moisture	1.5%w/w	0.24%	2.87%	0.34%	2.98%
Reconstitution Time	60 seconds	38.5s	30.5s	34.5s	-
Appearance after reconstitution time	A clear colourless to pale yellow solution free from visible particulates	clear very pale yellow solution	clear pale yellow solution	Clear colourless solution	-
XRPD	Assess whether crystalline material is present	No crystalline material detected	No crystalline material detected	No crystalline material detected	No crystalline material detected

Figure 43

4 Weeks

Test Parameters	Specification	12PM529-8-2 (40 °C/75%RH)	12PM529-8-2 (55 °C)	PM0237/12 (40 °C/75%RH)	PM0237/12 (55 °C)	12PM529-9-1 (40 °C/75%RH)	12PM529-9-1 (55 °C)
Appearance	White to off white powder	White Colour free flowing powder	White Colour free flowing powder	White Colour not free flowing powder	White Colour not free flowing powder	Off white colour free flowing powder	Light grey colour free flowing powder
Assay (mg/vial)	24.94 – 27.66	25.64mg/vial	25.72mg/vial	25.65mg/vial	25.72mg/vial	N/A	N/A
% Assay	95%-105%	102.56%	102.88%	102.61%	102.88%	95.51%	96.25%
Total Impurities	2%	0.92%	0.99%	0.61%	0.71%	2.56%	1.94%
Impurities Above 0.045%	N/A	0.89%	0.65%	0.40%	0.53%	2.35%	1.71%
Moisture	1.5%w/w	0.25%	0.36%	0.25%	0.23%	-	-
Reconstitution Time	60 seconds	42.5s	35.0s	35.0s	37.5s	-	-
Appearance after reconstitution time	A clear colourless to pale yellow solution free from visible particulates	Clear very pale yellow solution, free from visual particulates	Clear very pale yellow solution, free from visual particulates	Clear colourless solution free from visual particulates	Clear colourless solution free from visual particulates	-	-
XRPD	Assess whether crystalline material is present	No crystalline material detected	No crystalline material detected	No crystalline material detected	No crystalline material detected	-	-

Figure 44

13 Weeks

Test Parameters	Specification	12PM529-8-2 (40 °C/75%RH)	12PM529-8-1 (40 °C/75%RH)	PM0237/12 (40 °C/75%RH)	12PM529-9-1 (40 °C/75%RH)
Appearance	White to off white powder	White Colour free flowing powder	White Colour free flowing powder	White Colour not free flowing powder	Light beige colour free flowing powder
Assay (mg/vial)	24.94 – 27.66	25.69mg/vial	23.92mg/vial	25.48mg/vial	N/A
% Assay	95%-105%	102.76%	95.67%	101.92%	95.35%
Total Impurities	2%	0.78%	1.33%	0.57%	3.35%
Impurities Above 0.045%	N/A	0.49%	0.98%	0.41%	3.08%
Moisture	1.5%w/w	0.28%	2.96%	0.28%	-
Reconstitution Time	60 seconds	32.5s	37.5s	37.5s	-
Appearance after reconstitution time	A clear colourless to pale yellow solution free from visible particulates	Clear very pale yellow solution, free from visual particulates	Clear colourless solution free from visual particulates	Clear colourless solution free from visual particulates	-
XRPD	Assess whether crystalline material is present	No crystalline material detected	No crystalline material detected	No crystalline material detected	-

Figure 45

L10R5					
Test	Time 0	1 month 40°C±2°C / 75%±5% RH	3 months 40°C±2°C / 75%±5% RH	3 months 25°C±2°C / 60%±5% RH	1 month 55°C
Appearance of the lyophilisate	Off white cohesive cake, bright surface, some vials show cracks	Off white cohesive cake, bright surface, some vials show cracks	Off white cohesive cake, bright surface, some vials show cracks	Off white cohesive cake, bright surface, some vials show cracks	Off white cohesive cake, bright surface, some vials show cracks
Reconstitution time (V=4 mL)	30 s	35 s	72 s	63 s	38 s
Appearance of reconstituted solution	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles
Osmolality	303 mOsm/Kg	-	-	-	-
Moisture content	0.29% (0.30%; 0.28%; 0.29%)	0.69% (0.69%; 0.64%; 0.75%)	0.89% (0.93%; 0.86%; 0.89%)	0.56% (0.62%; 0.51%; 0.53%)	1.02% (1.12%; 0.98%; 0.95%)
HPLC content/assay	20.4 mg/vial 102%	20.2 mg/vial 101%	20.1 mg/vial 101%	20.3 mg/vial 102%	20.2 mg/vial 101%
HPLC sum of RS & RS details	Total RS = 0.52% CNS7054X = 0.06% RRT 0.66 = 0.04% RRT 0.69 = 0.05% RRT 0.92 = 0.11% RRT 1.32 = 0.22% RRT 1.92 = 0.03%	Total RS = 0.60% CNS7054X = 0.13% RRT 0.66 = 0.03% RRT 0.68 = 0.05% RRT 0.92 = 0.11% RRT 1.31 = 0.25% RRT 1.91 = 0.03%	Total RS = 0.74% CNS7054X = 0.28% RRT 0.66 = 0.03% RRT 0.69 = 0.05% RRT 0.92 = 0.11% RRT 1.31 = 0.24% RRT 1.91 = 0.03%	Total RS = 0.57% CNS7054X = 0.12% RRT 0.66 = 0.04% RRT 0.69 = 0.05% RRT 0.92 = 0.10% RRT 1.31 = 0.23% RRT 1.91 = 0.03%	Total RS = 0.93% CNS7054X = 0.53% RRT 0.66 = 0.03% RRT 0.67 = 0.04% RRT 0.92 = 0.10% RRT 1.31 = 0.23% RRT 1.93 < LOQ

Figure 46

L10R10			
Test	Time 0	1 month 40°C±2°C / 75%±5% RH	1 month 55°C
Appearance of the lyophilisate	Off-white cohesive cake, bright surface, irregular edge all around the cake	Off-white cohesive cake, bright surface, irregular edge all around the cake	Off-white cohesive cake, bright surface, irregular edge all around the cake
Reconstitution time (V=8 mL)	50 s	45 s	38 s
Appearance of reconstituted solution	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles
Osmolality	616 mOsm/kg	-	-
Moisture content	1.22% (1.29%; 1.16%)	0.71% (0.70%; 0.73%)	1.23% (1.71%*; 0.74%)
HPLC content/assay	39.7 mg/vial 99%	39.1 mg/vial 98%	39.4 mg/vial 99%
HPLC sum of RS & RS details	Total RS = 0.57% CNS7054X = 0.05% RRT 0.66 = 0.05% RRT 0.69 = 0.07% RRT 0.92 = 0.13% RRT 1.32 = 0.25% RRT 1.92=0.03%	Total RS = 0.61% CNS7054X = 0.13% RRT 0.66 = 0.04% RRT 0.68 = 0.05% RRT 0.92 = 0.11% RRT 1.31 = 0.25% RRT 1.92=0.03%	Total RS = 0.84% CNS7054X = 0.42% RRT 0.67 = 0.03% RRT 0.71 = 0.05% RRT 0.92 = 0.10% RRT 1.32 = 0.24% RRT 1.97<LOQ (**)

(*) Sample partially melt

(**) Slight shift of the RRT, this RS has to be compared with RRT 1.92

Figure 47

L10R5S87					
Test	Time 0	1 month 40°C±2°C / 75%±5% RH	3 months 40°C±2°C / 75%±5% RH	3 months 25°C±2°C / 60%±5% RH	1 month 55°C
Appearance of the lyophilisate	Off white cohesive cake, bright surface, some vials show irregular edge only on a small part of the cake	Off white cohesive cake, bright surface, some vials show irregular edge only on a small part of the cake	Off white cohesive cake, bright surface, some vials show irregular edge only on a small part of the cake	Off white cohesive cake, bright surface, some vials show irregular edge only on a small part of the cake	Off white cohesive cake, bright surface, some vials show irregular edge only on a small part of the cake
Reconstitution time (V=4 mL)	30 s	38 s	68 s	41 s	40 s
Appearance of reconstituted solution	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles
Osmolality	303 mOsm/Kg	-	-	-	-
Moisture content	0.56% (0.76%; 0.48%; 0.46%)	0.75% (0.78%; 0.73%; 0.76%)	0.92% (0.87%; 1.07%; 0.80%)	0.64% (0.74%; 0.57%; 0.61%)	1.03% (1.01%; 1.08%; 1.00%)
HPLC content/assay	20.7 mg/vial 104%	20.2 mg/vial 101%	20.2 mg/vial 101%	20.2 mg/vial 101%	20.2 mg/vial 101%
HPLC sum of RS & RS details	Total RS = 0.55% CNS7054X = 0.06% RRT 0.66 = 0.04% RRT 0.69 = 0.06% RRT 0.92 = 0.13% RRT 1.32 = 0.23% RRT 1.92 = 0.03%	Total RS = 0.61% CNS7054X = 0.16% RRT 0.66 = 0.03% RRT 0.68 = 0.05% RRT 0.92 = 0.11% RRT 1.31 = 0.25% RRT 1.92 = 0.03%	Total RS = 0.74% CNS7054X = 0.28% RRT 0.66 = 0.03% RRT 0.69 = 0.05% RRT 0.92 = 0.11% RRT 1.31 = 0.24% RRT 1.91 = 0.04%	Total RS = 0.57% CNS7054X = 0.12% RRT 0.66 = 0.03% RRT 0.69 = 0.05% RRT 0.92 = 0.10% RRT 1.31 = 0.24% RRT 1.91 = 0.03%	Total RS = 1.10% CNS7054X = 0.69% RRT 0.67 = 0.03% RRT 0.71 = 0.05% RRT 0.92 = 0.10% RRT 1.32 = 0.23% RRT 1.97 < LOQ(*)

(*) Slight shift of the RRT, this RS has to be compared with RRT 1.92

Figure 48

L20R5 (*)					
Test	Time 0	1 month 40°C±2°C / 75%±5% RH	3 months 40°C±2°C / 75%±5% RH	3 months 25°C±2°C / 60%±5% RH	1 month 55°C
Appearance of the lyophilisate	Off white cohesive cake, bright surface	Off white cohesive cake, bright surface (some cakes were found to be shrunk)	Off white cohesive cake, bright surface (some cakes were found to be shrunk)	Off white cohesive cake, bright surface	Off white cohesive cake, bright surface (some vials were found to be charred)
Reconstitution time (V=5 mL)	30 s	34 s	68 s	59 s	32 s
Appearance of reconstituted solution	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles (the charred cake was insoluble)
Osmolality	-	-	-	-	-
Moisture content	0.22% (0.19%; 0.23%; 0.22%)	0.37% (0.36%; 0.37%; 5.40% ^{**})	0.76% (0.44%; 1.07% (■))	0.63% (1.01%; 4.85% (■); 0.26%)	0.53% (0.52%; 0.52%; 0.55%)
HPLC content/assay	25.5 mg/vial 98%	26.9 mg/vial 103%	25.8 mg/vial 99%	25.6 mg/vial 98%	25.6 mg/vial 98%
HPLC sum of RS & RS details	Total RS = 0.58% CNS7054X = 0.07% RRT 0.66 = 0.05% RRT 0.69 = 0.06% RRT 0.92 = 0.12% RRT 1.32 = 0.24% RRT 1.92 = 0.03%	Total RS = 0.69% CNS7054X = 0.25% RRT 0.66 = 0.03% RRT 0.68 = 0.05% RRT 0.92 = 0.11% RRT 1.32 = 0.24% RRT 1.92 = 0.03%	Total RS = 1.02% CNS7054X = 0.51% RRT 0.61 = 0.03% RRT 0.66 = 0.03% RRT 0.69 = 0.05% RRT 0.92 = 0.11% RRT 1.31 = 0.23% RRT 1.91 = 0.03% RRT 2.07 = 0.03%	Total RS = 1.60% CNS7054X = 1.11% RRT 0.61 = 0.03% RRT 0.66 = 0.04% RRT 0.69 = 0.05% RRT 0.92 = 0.11% RRT 1.31 = 0.23% RRT 1.91 = 0.03% RRT 2.07 < LOQ	Total RS = 1.47% CNS7054X = 1.03% RRT 0.64 = 0.03% RRT 0.67 = 0.03% RRT 0.71 = 0.05% RRT 0.92 = 0.10% RRT 1.32 = 0.23% RRT 1.97 < LOQ (●)

(*) vials from clinical batch A01P310. Vials rejected after visual inspection; (**) Moisture content of a shrunk cake. This result wasn't considered in the average one as the other two results refer to cakes whose appearance didn't change from time 0. (***) Sample with a shrunk cake. (●) Slight shift of the RRT, this RS has to be compared with RRT 1.92; (■) result not included in the average; (■) Only two vials available for the analysis, the other vials were found to be shrunk.

Figure 49

L4M110R5			
Test	Time 0	1 month 40°C±2°C / 75%±5% RH	1 month 55°C
Appearance of the lyophilisate	Off white cohesive cake, bright surface, some vials show cracks	Off white cohesive cake, bright surface, some vials show cracks	Yellowish shrunk cake
Reconstitution time (V=4 mL)	40 s	25 s	31 s
Appearance of reconstituted solution	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles
Osmolality	292 mOsm/Kg	-	-
Moisture content	0.33% (0.34%; 0.33%)	0.69% (0.73%; 0.65%)	1.05% (1.10%; 1.01%)
HPLC content/assay	20.6 mg/vial 103%	20.3 mg/vial 102%	20.2 mg/vial 101%
HPLC sum of RS & RS details	Total RS = 0.56% CNS7054X = 0.07% RRT 0.66 = 0.04% RRT 0.69 = 0.06% RRT 0.92 = 0.13% RRT 1.32 = 0.23% RRT 1.92 = 0.03%	Total RS = 0.66% CNS7054X = 0.20% RRT 0.66 = 0.03% RRT 0.68 = 0.04% RRT 0.92 = 0.11% RRT 1.32 = 0.25% RRT 1.92 = 0.03%	Total RS = 1.33% RRT 0.36 = 0.03% CNS7054X = 0.90% RRT 0.67 = 0.03% RRT 0.71 = 0.05% RRT 0.92 = 0.10% RRT 1.32 = 0.24% RRT 1.97 < LOQ (*)
L4M110R10			
Test	Time 0	1 month 40°C±2°C / 75%±5% RH	1 month 55°C
Appearance of the lyophilisate	Off-white cohesive cake, irregular edge all around the cake	Off-white cohesive cake, irregular edge all around the cake	Yellowish shrunk cake
Reconstitution time (V=8 mL)	50 s	81 s	65 s
Appearance of reconstituted solution	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles
Osmolality	596 mOsm/Kg	-	-
Moisture content	0.53% (0.54%; 0.52%)	0.95% (0.79%; 1.12%)	0.98% (0.95%; 1.01%)
HPLC content/assay	41.0 mg/vial 103%	40.1 mg/vial 100%	39.7 mg/vial 99%
HPLC sum of RS & RS details	Total RS = 0.55% CNS7054X = 0.06% RRT 0.66 = 0.04% RRT 0.69 = 0.06% RRT 0.92 = 0.12% RRT 1.32 = 0.24% RRT 1.92 = 0.03%	Total RS = 0.61% CNS7054X = 0.16% RRT 0.66 = 0.03% RRT 0.68 = 0.06% RRT 0.92 = 0.11% RRT 1.32 = 0.25% RRT 1.92 < LOQ	Total RS = 1.34% RRT 0.34 = 0.03% RRT 0.36 = 0.03% CNS7054X = 0.88% RRT 0.67 = 0.03% RRT 0.71 = 0.06% RRT 0.92 = 0.10% RRT 1.32 = 0.24% RRT 1.97 < LOQ (*)

Figure 50

L2M110R5			
Test	Time 0	1 month 40°C±2°C / 75%±5% RH	1 month 55°C
Appearance of the lyophilisate	Off white cohesive cake, surface less brighter than the other formulations, some vials show irregular edge all around the cake	Off-white cohesive cake further shrunk , irregular edge all around the cake, surface less brighter than the other formulations	Yellowish shrunk cake
Reconstitution time (V=4 mL)	50 s	43 s	45 s
Appearance of reconstituted solution	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles
Osmolality	329 mOsm/Kg	-	-
Moisture content	0.34% (0.31%; 0.36%)	0.78% (0.82%; 0.73%)	0.83% (0.73%; 0.92%)
HPLC content/assay	20.5 mg/vial 103%	20.2 mg/vial 101%	19.9 mg/vial 100%
HPLC sum of RS & RS details	<u>Total RS = 0.58%</u> CNS7054X = 0.08% RRT 0.66 = 0.05% RRT 0.69 = 0.06% RRT 0.92 = 0.13% RRT 1.32 = 0.24% RRT 1.92 = 0.03%	<u>Total RS = 0.68%</u> CNS7054X = 0.22% RRT 0.61 = 0.03% RRT 0.66 = 0.03% RRT 0.68 = 0.05% RRT 0.92 = 0.11% RRT 1.32 = 0.23% RRT 1.92 = 0.03%	<u>Total RS = 1.76%</u> RRT 0.34 = 0.05% RRT 0.36 = 0.05% CNS7054X = 1.21% RRT 0.64 <LOQ RRT 0.67 = 0.03% RRT 0.71 = 0.06% RRT 0.92 = 0.10% RRT 1.32 = 0.24% RRT 1.46 = 0.03% RRT 1.97 <LOQ (*)

Figure 51

L2M110R10			
Test	Time 0	1 month 40°C±2°C / 75%±5% RH	1 month 55°C
Appearance of the lyophilisate	Off-white cohesive cake, bright surface, irregular edge all around the cake, shrunk cake, surface less brighter than the other formulations	Off-white cohesive cake further shrunk , irregular edge all around the cake, surface less brighter than the other formulations	Yellowish shrunk cake
Recon. time (V=8 mL)	62 s	81 s	54 s
Appearance of recon solution	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles	Clear, pale yellow solution free of visible particles
Osmolality	648 mOsm/Kg	-	-
Moisture content	1.03% (1.16%; 0.91%)	1.00% (0.92%; 1.08%)	0.95% (0.96%; 0.94%)
HPLC content/assay	40.6 mg/vial 102%	40.5 mg/vial 101%	39.8 mg/vial 100%
HPLC sum of RS & RS details	<u>Total RS = 0.58%</u> CNS7054X = 0.07% RRT 0.66 = 0.04% RRT 0.69 = 0.07% RRT 0.92 = 0.12% RRT 1.32 = 0.25% RRT 1.92 = 0.03%	<u>Total RS = 0.72%</u> CNS7054X = 0,25% RRT 0,61 = 0,03% RRT 0.66 = 0,03% RRT 0.68 = 0,05% RRT 0.92 = 0,11% RRT 1.32 = 0,25% RRT 1.92 < LOQ	<u>Total RS = 2.23%</u> RRT 0,34 = 0.04% RRT 0,36 = 0.05% CNS7054X = 1.65% RRT 0.64 = 0.03% RRT 0.67 = 0.03% RRT 0.71 = 0.07% RRT 0.92 = 0.10% RRT 1.32 = 0.24% RRT 1.97 < LOQ (*)

(*)Slight shift of the RRT, this RS has to be compared with RRT 1.92

Figure 52




Figure 53

