




(86) Date de dépôt PCT/PCT Filing Date: 2011/06/03
(87) Date publication PCT/PCT Publication Date: 2011/12/08
(45) Date de délivrance/Issue Date: 2017/04/25
(85) Entrée phase nationale/National Entry: 2012/12/03
(86) N° demande PCT/PCT Application No.: US 2011/039135
(87) N° publication PCT/PCT Publication No.: 2011/153477
(30) Priorités/Priorities: 2010/06/03 (US61/351,125);
2011/01/07 (US61/430,645); 2011/04/13 (US61/474,764)

(51) Cl.Int./Int.Cl. *A61K 39/395* (2006.01),
A61P 17/00 (2006.01), *C07K 16/24* (2006.01)
(72) Inventeurs/Inventors:
OKUN, MARTIN M., US;
HARRIS, THOMAS C., US
(73) Propriétaire/Owner:
ABBVIE BIOTECHNOLOGY LTD., BM
(74) Agent: TORYS LLP

(54) Titre : UTILISATIONS ET COMPOSITIONS POUR LE TRAITEMENT DE L'HIDROSADENITE (HS)
(54) Title: USES AND COMPOSITIONS FOR TREATMENT OF HIDRADENITIS SUPPURATIVA (HS)



(57) **Abrégé/Abstract:**

The disclosure provides methods and compositions for the treatment of hidradenitis suppurativa. The disclosure describes the methods and uses for treating hidradenitis suppurativa in a subject using a TNF α inhibitor, such as a human TNF α antibody, or an antigen-binding portion thereof. Also described are methods for determining the efficacy of a TNF α inhibitor for treating hidradenitis suppurativa in a subject.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
8 December 2011 (08.12.2011)(10) International Publication Number
WO 2011/153477 A3

(51) International Patent Classification:

A61K 39/395 (2006.01) *C07K 16/00* (2006.01)
A61K 39/00 (2006.01)

(21) International Application Number:

PCT/US2011/039135

(22) International Filing Date:

3 June 2011 (03.06.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/351,125	3 June 2010 (03.06.2010)	US
61/430,645	7 January 2011 (07.01.2011)	US
61/474,764	13 April 2011 (13.04.2011)	US

(71) Applicant (for all designated States except US): **AB-BOTT BIOTECHNOLOGY LTD.**; Clarendon House, 2 Church Street, Hamilton, HM 11 (BM).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **OKUN, Martin, M.** [US/US]; 207 W. Park Avenue, Libertyville, IL 60048(US). **HARRIS, Thomas, C.** [US/US]; 432 Saint Andrews Lane, Gurnee, IL 60031 (US).(74) Agents: **COWLES, Cristin, Howley** et al.; McCarter & English, LLP, 265 Franklin Street, Boston, MA 02110 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,

[Continued on next page]

(54) Title: USES AND COMPOSITIONS FOR TREATMENT OF HIDRADENITIS SUPPURATIVA (HS)

Figure 1



(57) Abstract: The disclosure provides methods and compositions for the treatment of hidradenitis suppurativa. The disclosure describes the methods and uses for treating hidradenitis suppurativa in a subject using a TNFa inhibitor, such as a human TNFa antibody, or an antigen-binding portion thereof. Also described are methods for determining the efficacy of a TNPa inhibitor for treating hidradenitis suppurativa in a subject.

WO 2011/153477 A3

LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:**Declarations under Rule 4.17:**

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
- *with sequence listing part of description (Rule 5.2(a))*

(88) Date of publication of the international search report:

15 March 2012

USES AND COMPOSITIONS FOR
TREATMENT OF HIDRADENITIS SUPPURATIVA (HS)

RELATED APPLICATIONS

This application claims the benefit of the filing date under 35 U.S.C. §119(e) of
5 U.S. Provisional Application No. 61/351,125, filed June 3, 2010; U.S. Provisional
Application No. 61/430,645, filed January 7, 2011; and U.S. Provisional Applicaion No.
61/474,764, filed April 13, 2011.

BACKGROUND OF THE INVENTION

10 Hidradenitis suppurativa (HS) refers to a skin disorder of the apocrine glands
(sweat glands found on certain parts of the body) and hair follicles in which swollen,
painful, chronically inflamed lesions or lumps develop in the groin and sometimes under
the arms and under the breasts. Hidradenitis suppurativa is characterized by recurrent
inflamed nodules, abscesses, and fistulas, and it occurs when apocrine gland outlets
15 become blocked by perspiration or are unable to drain normally because of incomplete
gland development. Secretions trapped in the glands force perspiration and bacteria into
surrounding tissue, causing subcutaneous induration, inflammation, and infection.
Hidradenitis suppurativa is confined to areas of the body that contain apocrine glands.
These areas are the axillae, areola of the nipple, groin, perineum, circumanal, and
20 periumbilical regions.

HS is a chronic inflammatory skin disease, particularly of young adults, and
affects approximately 1% of the general population in the West, with women affected 2
to 5 times more commonly than men (Naldi L. Epidemiology. *In*: Hidradenitis
Suppurativa (Jemec G, Revuz J, Leyden J, eds). Heidelberg: Springer. 2006), and with
25 an average age of onset of 23 years. The poorly understood disease is believed to be
under-reported by those who suffer from it. HS is also associated with obesity and
smoking.

The disease is associated with significant morbidity. Given the pain and
consequent physical impairment associated with the tender lesions of this disease, it has

been reported that health-related quality of life is lower for patients with hidradenitis suppurativa than other dermatological diseases. In addition, a recent study estimated that up to 20% of patients with HS report the co-existence of depression, and HS patients report a high level of stigmatization. Use of emergency services and antibiotics for
5 incision and drainage of painful abscesses are high when lesions flare.

It is speculated that immunological abnormalities of the hair follicle play a role in the etiology of this disease, and that the underlying mechanisms may be pathogenetically related to those of Crohn's disease (Kurzen *et al.*, *Exp Dermatol.* 17: 455-456, 2008).

Treatment options have been largely disappointing. Some researchers also
10 suggest that changes in diet, warm compresses and baths, and zinc gluconate supplements can relieve symptoms of HS, bring about remission, and/or minimize recurrence.

To date, no systemic therapy has been demonstrated to be effective for HS in a randomized, double-blind, placebo-controlled trial. Non-evidence-based approaches for
15 moderate to severe HS include long-term antibiotic therapy to control inflammation; case reports or series have described corticosteroids, cyclosporine, or methotrexate as occasionally effective. Surgical intervention is utilized for more advanced HS cases (Alikhan *et al.*, *J Am Acad Dermatol.* 60: 539-561, 2009). That is, the evidentiary basis behind current therapies for moderate-severe disease, including short- or long-term oral
20 or topical antibiotics, retinoids, intralesional steroids, oral steroids, immunosuppressive agents, radiation, laser therapy, or disfiguring surgical removal of involved areas in more severe cases, is largely limited to anecdotal experience or open-label studies. Surgery is the preferred treatment option in Europe. Currently, there are no approved therapies for this disease in the US.

25 While some recent case reports have described successful use of tumor necrosis factor- α (TNF- α) antagonists in HS (Haslund *et al.*, *Acta Derm Venereol.* 89: 595-600, 2009), TNF α antagonists have met with limited success in treating HS. For example, a study examining treatment of HS with etanercept failed to show improvement of HS over a 24-week treatment period (Adams *et al.*, *Arch Dermatol.* 146(5): 501-504, 2010).

SUMMARY OF THE INVENTION

Given the limited success of treatments for hidradenitis suppurativa (HS), there remains a need for an effective treatment, especially in view of the debilitating nature of this disease.

5 The invention described herein provides a safe and effective treatment of hidradenitis suppurativa (HS) using TNF α inhibitors, particularly human anti-TNF α antibodies, such as adalimumab / D2E7.

Adalimumab is a monoclonal IgG antibody that contains only human peptide sequences. It binds with high specificity and affinity to soluble and membrane-bound
10 TNF α , thereby neutralizing the biological activities of TNF α . Thus, the instant invention provides improved methods and compositions for treating hidradenitis suppurativa.

In one embodiment, the invention provides a means for treating patients suffering from moderate to severe chronic hidradenitis suppurativa.

15 In one embodiment, the invention provides a means for treating patients suffering from moderate to severe hidradenitis suppurativa.

The invention provides improved methods of treatment, including methods of improving disease reduction in patients having hidradenitis suppurativa and improvements in quality of life for the hidradenitis suppurativa patients.

20 In one embodiment, the invention provides a method for treating certain subpopulations of patients, including, for example, those who have failed prior therapy or have had a sub-therapeutic response, including, for example, a subject who has an inadequate response to or is intolerant to, or has a contraindication to, oral antibiotics. In certain embodiments, the invention is used to treat HS in a subject who was
25 unresponsive or intolerant to oral antibiotics for treatment for their hidradenitis suppurativa.

In one embodiment, the invention includes the treatment of a subject who has an AN count of greater than or equal to 3 at baseline. In another embodiment, the subject is a female. In a further embodiment, the subject who is over 40 years old. In yet another
30 embodiment, the subject is a smoker.

In yet a further embodiment, the subject may have any combination of the specific features recited herein. For example, the subject may be a female smoker who is over 40 years old and who may have a history of intolerance to antibiotics.

In one embodiment, the TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, is administered weekly. In one embodiment, the TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, is administered every other week (eow or biweekly).

In one embodiment, the invention provides a method of treating hidradenitis suppurativa (HS) in a subject having HS, comprising administering human TNF α antibody, or an antigen-binding portion thereof, such as adalimumab, to the subject as a fixed dose, *e.g.*, about 40 mg, according to a weekly dosing regimen. In one embodiment, the dosing regimen includes an induction dose or doses, followed by weekly dosing of the antibody to treat HS.

In one embodiment, the subject being treated is first selected as having HS and is subsequently treated with a human TNF α antibody, or an antigen-binding portion thereof, according to the methods described herein, *e.g.*, weekly.

In one aspect, the invention provides a method of achieving a clinical response in a subject suffering from hidradenitis suppurativa, comprising administering an effective amount of a human TNF α antibody, or antigen-binding portion thereof, to the subject such that the clinical response in hidradenitis suppurativa is achieved.

In one embodiment, the TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, is administered to the subject on a multiple variable dosing regimen.

In one embodiment, the invention provides a method for treating a subject having hidradenitis suppurativa (HS), the method comprising administering an isolated human anti-TNF α antibody, or an antigen binding portion thereof, to the subject according to a multiple variable dose regimen, such that HS is treated, wherein the multiple variable dose regimen comprises administering a first loading dose, administering a second loading dose which is less than the first loading dose, and administering a treatment dose

which is less than the second loading dose, wherein the treatment dose is administered to the subject weekly.

In one embodiment, the invention provides a method for treating a subject having hidradenitis suppurativa (HS), the method comprising administering an isolated human
5 anti-TNF α antibody, or an antigen binding portion thereof, to the subject according to a multiple variable dose regimen, such that HS is treated, wherein the multiple variable dose regimen comprises administering a first loading dose, administering a second loading dose which is less than the first loading dose, and administering a treatment dose which is less than the second loading dose, wherein the treatment dose is administered to
10 the subject biweekly.

In one embodiment, the maintenance dose is administered on a weekly dosing regimen, followed by a biweekly dosing regimen.

In one embodiment, the second loading dose is about 40-60% of the first loading dose. In one embodiment, the treatment dose is about 40-60% of the second loading
15 dose.

In one embodiment, the first loading dose is about 140-180 mg, *e.g.*, about 160 mg.

In one embodiment, the second loading dose is about 60-100 mg, *e.g.*, about 80 mg.

20 In one embodiment, the treatment dose is about 30-50 mg, *e.g.*, about 40 mg.

In a further embodiment, the invention provides a method for decreasing the number of inflammatory lesions (AN count) in a subject having HS, said method comprising systemically administering an isolated human anti-TNF α antibody, or an antigen binding portion thereof, to the subject, such that the AN count is decreased.

25 In one embodiment, the AN count is reduced by at least a 50% reduction in the subject relative to baseline AN count.

In one embodiment, the subject has no increase in an abscess count and/or no increase in a draining fistula count following administration with the anti-TNF α antibody, or an antigen binding portion thereof.

In one embodiment, the anti-TNF α antibody, or antigen binding portion thereof, is administered to the subject on a weekly basis.

In yet a further embodiment of the invention, the subject has HS lesions in at least two distinct anatomic areas prior to treatment.

5 In one embodiment, the TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, is administered to the subject on a weekly or a biweekly dosing regimen, including at a dose of about 40 mg administered weekly or biweekly, respectively.

10 In yet another embodiment, the TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, is administered to the subject via subcutaneous administration.

In certain embodiments, the clinical response is measured by reduction in pain.

15 In certain embodiments, the reduction in pain is greater than 30% or greater than 10 mm reduction from a baseline measured at the beginning of the treatment, after 2, 4, 8, 12, or 16 weeks of treatment.

In another aspect, the invention further provides a method of treating hidradenitis suppurativa in a subject comprising administering an initial loading dose of a human TNF α antibody or antigen-binding portion thereof, to the subject, then administering subsequent doses, *e.g.*, maintenance or treatment doses, of the human TNF α antibody or antigen-binding portion thereof, to the subject, wherein the maintenance doses are about one-half to one-fourth of the dose amount of the loading dose. In certain embodiments, the loading dose may comprise one or more doses. In certain embodiments, the loading dose comprises a first dose of about 160 mg and a second dose of about 80 mg, optionally administered 2 weeks apart. In certain other embodiments, the loading dose 25 comprises a single dose of about 80 mg.

In one embodiment, the initial dose is given in its entirety on Day 1 (of week 0), or is given twice, once each at week 0 and week 2. In one embodiment, the second dose, *e.g.*, maintenance or treatment dose, is administered to the subject about one week or two weeks after the last loading dose, and is given on a weekly or biweekly dosing 30 regimen.

In certain embodiments, the loading dose comprises about 160 mg of adalimumab administered at week 0 and about 80 mg of adalimumab administered at week 2, and the maintenance dose comprises weekly administration of about 40 mg of adalimumab starting from week 4 .

- 5 In certain embodiments, the loading dose comprises about 80 mg of adalimumab administered at week 0, and the maintenance dose comprises biweekly / every-other-week (eow) administration of about 40 mg of adalimumab starting from week 1.

 In one embodiment, the invention provides a method of improving (*i.e.*, reducing) a Hidradenitis Suppurativa-Physician's Global Assessment (HS-PGA) score of
10 a subject having hidradenitis suppurativa from a high score (*e.g.*, an HS-PGA score of 3 or more) to a no or small impact score (*e.g.*, an HS-PGA score of 0-2), comprising administering a TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, to the subject, such that the HS-PGA score improves from the high score to the no or small impact score. The invention also provides a method of decreasing an HS-
15 PGA score of a subject having hidradenitis suppurativa by at least about 2 grades, comprising administering a TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, to the subject, such that the HS-PGA score is decreased by at least about 2 grades.

 In one aspect, the invention provides a method of decreasing an HS-PGA score
20 of a subject having hidradenitis suppurativa from a high score (*e.g.*, a score of 3 or more) to a no or small impact score (*e.g.*, a score of 0-2), comprising administering a human TNF α antibody to the subject, such that the HS-PGA score is decreased from the high score to the no or small impact score.

 In one aspect, the invention provides a method of decreasing an HS-PGA score
25 of a subject having hidradenitis suppurativa by at least about 2 grades, comprising administering a human TNF α antibody to the subject, such that the HS-PGA score is decreased by at least about 2 grades.

 The invention also includes administering an effective TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, to a subject or patient
30 population having hidradenitis suppurativa.

In one embodiment, the TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, is administered to the patient population or subject on a weekly or biweekly dosing regimen.

In still another embodiment, the TNF α inhibitor is administered in a multiple
5 variable dose regimen. In one embodiment, the multiple variable dose regimen comprises one or more induction or loading doses, which are at least double or quadruple the treatment or maintenance doses. In certain embodiments, the TNF α inhibitor is administered weekly or biweekly to the patient population or subject. In one embodiment, the induction dose comprises about 160 or 80 mg. In one
10 embodiment, the treatment dose comprises about 40 mg.

In one embodiment, the TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, is administered to the patient population or subject at a dose of about 40 mg on a weekly or biweekly dosing regimen.

In another embodiment, the TNF α inhibitor, *e.g.*, human TNF α antibody, or
15 antigen-binding portion thereof, is administered to the patient population or subject via subcutaneous administration.

The invention provides an article of manufacture comprising: a human TNF α antibody, or antigen-binding portion thereof, and a label or package insert contained within the packaging material indicating that an adverse event which has been reported
20 in the use of the human TNF α antibody is super infection of hidradenitis suppurativa lesions and/or pilonidal cyst flare.

The invention includes a package comprising a TNF α inhibitor and instructions for administering the TNF α inhibitor to a human subject for the treatment of adults with hidradenitis suppurativa, *e.g.*, moderate to severe chronic hidradenitis suppurativa, who
25 have been unresponsive or intolerant to oral antibiotics for treatment for their hidradenitis suppurativa. The invention also includes a package comprising a TNF α inhibitor, wherein the package contains, on the label and in a position which is visible to a subject, including a prospective purchaser, a printed statement which informs a subject, including a prospective purchaser, that the TNF α inhibitor is indicated for the
30 treatment of adults with moderate to severe chronic hidradenitis suppurativa who have

been unresponsive or intolerant to oral antibiotics for treatment for their hidradenitis suppurativa.

The invention further provides a package comprising a TNF α inhibitor, wherein the package contains, on the label and in a position which is visible to a subject,
5 including a prospective purchaser, a printed statement which informs a subject, including a prospective purchaser, that the recommended dose of the TNF α inhibitor for patients with hidradenitis suppurativa is 40 mg TNF α inhibitor administered every week or every other week, as a single dose via subcutaneous injection.

In one aspect, the invention provides an article of manufacture comprising a
10 human TNF α antibody and a package insert, wherein the package insert indicates the recommended human TNF α antibody dose regimen for adult patients with hidradenitis suppurativa is about 160 mg at week 0, about 80 mg at week 2, followed by 40 mg every week beginning at week 4.

In one aspect, the invention provides an article of manufacture comprising a
15 human TNF α antibody and a package insert, wherein the package insert indicates the recommended human TNF α antibody dose regimen for adult patients with hidradenitis suppurativa is about 80 mg at week 0, followed by about 40 mg every other week beginning at week 1.

In one aspect, the invention provides an article of manufacture which comprising
20 adalimumab and a package insert, wherein the package insert indicates that the adalimumab may be used to treat hidradenitis suppurativa in patients who have been unresponsive or intolerant to oral antibiotics for treatment for their hidradenitis suppurativa.

The invention also provides a means for determining the efficacy of a TNF α
25 inhibitor for the treatment of hidradenitis suppurativa.

The invention further provides a method of determining the efficacy of a TNF α inhibitor, *e.g.*, a human TNF α antibody, or an antigen-binding portion thereof, for treating hidradenitis suppurativa in a subject, comprising determining a proportion of treated subjects achieving Hidradenitis Suppurativa Clinical Response (HiSCR) within a
30 patient population having hidradenitis suppurativa who was administered the human TNF α antibody, or antigen-binding portion thereof, wherein a statistically significant

(*e.g.*, $p < 0.05$, or $p < 0.01$, or $p < 0.005$, or $p < 0.001$) increase in the proportion achieving clinical response in the treatment group compared to placebo indicates that the TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, is an effective TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, for the treatment of hidradenitis suppurativa in the subject.

In one embodiment, at least about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25% or more of the patient population achieve statistically significant Hidradenitis Suppurativa Clinical Response (HiSCR) indicates that the TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, is an effective human TNF α antibody, or antigen-binding portion thereof, for the treatment of hidradenitis suppurativa in the subject.

In one embodiment, the invention includes a method for determining the efficacy of a TNF α inhibitor, *e.g.*, human TNF α antibody, or an antigen-binding portion thereof, for improving the functional limitations of a subject having hidradenitis suppurativa comprising determining an improvement in an HS-PGA score from a patient population administered with the TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, wherein a statistically significant difference in reduction of the HS-PGA score (*e.g.*, by at least 2 grades, and/or to an HS-PGA score of 0-2) for the patient population indicates that the human TNF α antibody, or antigen-binding portion thereof, is an effective TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, for improving the functional limitations of a subject having hidradenitis suppurativa.

The invention further provides a method of determining the efficacy of a TNF α inhibitor, *e.g.*, a human TNF α antibody, or an antigen-binding portion thereof, for treating hidradenitis suppurativa in a subject, comprising determining a proportion of treated subject achieving clinical response (*e.g.*, as defined by HS-PGA score reduction) within a patient population having hidradenitis suppurativa who was administered the human TNF α antibody, or antigen-binding portion thereof, wherein a statistically significant (*e.g.*, $p < 0.05$, or $p < 0.01$, or $p < 0.005$, or $p < 0.001$) increase in the proportion achieving clinical response in the treatment group compared to placebo indicates that the TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding

portion thereof, is an effective TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, for the treatment of hidradenitis suppurativa in the subject.

In one embodiment, at least about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25% or more of the patient population achieve statistically
5 significant clinical response (*e.g.*, as measured by reduction of HS-PGA score as defined herein) indicates that the TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, is an effective human TNF α antibody, or antigen-binding portion thereof, for the treatment of hidradenitis suppurativa in the subject.

The invention also provides a method of treating hidradenitis suppurativa in a
10 subject comprising administering an effective amount of a TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, to the subject, wherein the effective amount of the TNF α inhibitor, *e.g.*, human TNF α antibody, or antigen-binding portion thereof, has previously been identified as achieving a statistically significant clinical response in at least about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22,
15 23, 24, 25% or more of a patient population having hidradenitis suppurativa. Ranges of values using a combination of any of the above recited values as upper and/or lower limits are intended to be included in the scope of the invention.

In one embodiment, the efficacy of a TNF α inhibitor for treating hidradenitis suppurativa in a patient population may be evaluated by determining the percentage of
20 the patient population achieving Hidradenitis Suppurativa Clinical Response (HiSCR). HiSCR is defined as at least a 50% reduction in the total inflammatory lesion (abscess and nodule) count (AN count) relative to baseline with no increase in abscess count and no increase in draining fistula count.

In one embodiment, the efficacy of a TNF α inhibitor for treating hidradenitis
25 suppurativa in a patient population may be evaluated by determining the percentage of the patient population achieving clinical response (as defined by HS-PGA score reduction) for whom the TNF α inhibitor has been effective for treating hidradenitis suppurativa.

In one embodiment, the invention provides a kit for the treatment of HS in a
30 subject, said kit comprising containers providing the loading dose(s) and/or treatment dose(s), *e.g.*, at least seven containers, of an isolated human anti-TNF α antibody, or an

antigen binding portion thereof. The kit may also further provide instructions for administration of the anti-TNF α antibody, or antigen binding portion thereof, to a subject having HS. In one embodiment, the container is a preloaded syringe. In one embodiment, the container is an autoinjector. In one embodiment, each container in the
5 kit contains about 40 mg of the anti-TNF α antibody, or antigen binding portion thereof.

In one embodiment, the TNF α inhibitor is selected from the group consisting of an anti-TNF α antibody, or an antigen-binding portion thereof, a TNF fusion protein, or a recombinant TNF binding protein.

In one embodiment, the TNF fusion protein is etanercept.

10 In one embodiment, the TNF α antibody, or antigen-binding portion thereof, is selected from the group consisting of a chimeric antibody, a humanized antibody, and a multivalent antibody.

In one embodiment of the invention, the TNF α antibody, or antigen-binding portion thereof, is a human antibody.

15 In another embodiment, the TNF α antibody, or antigen-binding portion thereof, is an isolated human antibody that dissociates from human TNF α with a K_d of 1×10^{-8} M or less and a k_{off} rate constant of $1 \times 10^{-3} \text{ s}^{-1}$ or less, both determined by surface plasmon resonance, and neutralizes human TNF α cytotoxicity in a standard *in vitro* L929 assay with an IC_{50} of 1×10^{-7} M or less.

20 In certain embodiments, the TNF α antibody is an isolated human antibody, or antigen-binding portion thereof, with the following characteristics: a) dissociates from human TNF α with a k_{off} rate constant of $1 \times 10^{-3} \text{ s}^{-1}$ or less, as determined by surface plasmon resonance; (b) has a light chain CDR3 domain comprising the amino acid sequence of SEQ ID NO: 3, or modified from SEQ ID NO: 3 by a single alanine
25 substitution at position 1, 4, 5, 7 or 8 or by one to five conservative amino acid substitutions at positions 1, 3, 4, 6, 7, 8 and/or 9; (c) has a heavy chain CDR3 domain comprising the amino acid sequence of SEQ ID NO: 4, or modified from SEQ ID NO: 4 by a single alanine substitution at position 2, 3, 4, 5, 6, 8, 9, 10 or 11 or by one to five conservative amino acid substitutions at positions 2, 3, 4, 5, 6, 8, 9, 10, 11 and/or 12.

In one embodiment, the TNF α antibody is an isolated human antibody, or an antigen binding portion thereof, comprising a light chain variable region comprising CDRs having the amino acid sequences described in SEQ ID NOs: 3, 5, and 7, and a heavy chain variable region comprising CDRs having the amino acid sequences
5 described in SEQ ID NOs: 4, 6, and 8.

In one embodiment, the TNF α antibody is an isolated human antibody, or an antigen binding portion thereof, with a light chain variable region (LCVR) comprising the amino acid sequence of SEQ ID NO: 1 and a heavy chain variable region (HCVR) comprising the amino acid sequence of SEQ ID NO: 2.

10 In one embodiment, the anti-TNF α antibody, or antigen-binding portion thereof, is adalimumab.

In one embodiment, the anti-TNF α antibody, or antigen-binding portion thereof, is an adalimumab biosimilar antibody or an adalimumab interchangeable antibody.

In one embodiment, the TNF α antibody, or antigen-binding portion thereof, is a
15 40 mg dose.

In another embodiment, the TNF α antibody, or antigen-binding portion thereof, is administered subcutaneously.

In certain embodiments, the TNF α antibody, or antigen-binding portion thereof, is infliximab or golimumab. In yet another embodiment, the TNF α antibody, or antigen-binding portion thereof, is certolizumab.
20

In one embodiment, the TNF α inhibitor is administered weekly to the patient population or subject having hidradenitis suppurativa. In another embodiment, TNF α inhibitor is administered biweekly to the patient population or subject having hidradenitis suppurativa.

25 In is contemplated that, all embodiments described herein, including those described under different aspects of the invention, can be combined with one another where not specifically prohibited.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the study design for the Phase II clinical trial described in Examples 1-5. a: Dose escalation to ew dosing for patients with PGA ≥ 3 at Weeks 28 or 31. b: From Week 4, after 160 mg dose at Week 0, 80 mg at Week 2. c: From Week 1, after 80 mg dose at Week 0. d: From Week 17, after 80 mg dose at Week 16.

5 Figure 2 shows disposition of patients screened for the study. ADA: adalimumab; eow: every other week; ew: every week.

Figure 3 shows efficacy of adalimumab over 52 weeks. A significantly greater proportion of patients in the ew group achieved the primary endpoint, a clinical response at Week 16, compared with patients allocated to the placebo group; *p < 0.05, ew vs. placebo, ITT analysis.

10

DETAILED DESCRIPTION OF THE INVENTION

I. Definitions

The term “human TNF α ” (abbreviated herein as hTNF α , or simply hTNF), as used herein, is intended to refer to a human cytokine that exists as a 17 kDa secreted form and a 26 kDa membrane associated form, the biologically active form of which is composed of a trimer of non-covalently bound 17 kDa molecules. The structure of hTNF α is described further in, for example, Pennica, D., *et al.* (1984) *Nature* 312:724-729; Davis, J.M., *et al.* (1987) *Biochemistry* 26:1322-1326; and Jones, E.Y., *et al.* (1989) *Nature* 338:225-228. The term human TNF α is intended to include recombinant human TNF α (rhTNF α), which can be prepared by standard recombinant expression methods or purchased commercially (R & D Systems, Catalog No. 210-TA, Minneapolis, MN). TNF α is also referred to as TNF. In addition, the terms “TNF α antibody” and “anti-TNF α antibody” are used interchangeably throughout.

15

20

25 The term “TNF α inhibitor” includes agents which interfere with TNF α activity. The term also includes each of the anti-TNF α human antibodies and antibody portions described herein as well as those described in U.S. Patent Nos. 6,090,382; 6,258,562; 6,509,015; 7,223,394, and in U.S. Publication No. US20030219438. In one embodiment, the TNF α inhibitor used in the invention is an anti-TNF α antibody, or a

fragment thereof, including infliximab (REMICADE[®], Centocor; described in U.S. Patent No. 5,656,272), CDP571 (a humanized monoclonal anti-TNF-alpha IgG4 antibody), CIMZIA[®] (certolizumab (or CDP870) a humanized monoclonal anti-TNF-alpha antibody fragment; UCB), an anti-TNF dAb (Peptech), SIMPONI[®] (golimumab; also referred to as CNTO 148; Centocor Ortho Biotech, see WO 02/12502), and adalimumab (HUMIRA[®], Abbott Laboratories, a human anti-TNF α mAb, described in US 6,090,382 as D2E7). Additional TNF α antibodies which may be used in the invention are described in U.S. Patent Nos. 6,593,458; 6,498,237; 6,451,983; and 6,448,380.

10 In another embodiment, the TNF α inhibitor is a TNF-R fusion protein, *e.g.*, etanercept (ENBREL[®], Amgen; described in WO 91/03553 and WO 09/406476).

In another embodiment, the TNF α inhibitor is a recombinant TNF binding protein (r-TBP-I) (Serono).

The term “antibody,” as used herein, is intended to refer to immunoglobulin molecules comprised of four polypeptide chains, two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as HCVR or VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as LCVR or VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The antibodies of the invention are described in further detail in U.S. Patent Nos. 6,090,382; 6,258,562; and 6,509,015 .

30 The term “antigen-binding portion” or “antigen-binding fragment” of an antibody (or simply “antibody portion”), as used herein, refers to one or more fragments of an antibody that retain the ability to specifically bind to an antigen (*e.g.*, hTNF α). It has been shown that the antigen-binding function of an antibody can be performed by

fragments of a full-length antibody. Binding fragments include Fab, Fab', F(ab')₂, Fabc, Fv, single chains, and single-chain antibodies. Examples of binding fragments encompassed within the term "antigen-binding portion" of an antibody include (i) a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (v) a dAb fragment (Ward *et al.* (1989) *Nature* 341:544-546), which consists of a VH domain; and (vi) an isolated complementarity determining region (CDR).

Furthermore, although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the VL and VH regions pair to form monovalent molecules (known as single chain Fv (scFv); see *e.g.*, Bird *et al.* (1988) *Science* 242:423-426; and Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883). Such single chain antibodies are also intended to be encompassed within the term "antigen-binding portion" of an antibody. Other forms of single chain antibodies, such as diabodies are also encompassed. Diabodies are bivalent, bispecific antibodies in which VH and VL domains are expressed on a single polypeptide chain, but using a linker that is too short to allow for pairing between the two domains on the same chain, thereby forcing the domains to pair with complementary domains of another chain and creating two antigen binding sites (see *e.g.*, Holliger *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:6444-6448; Poljak *et al.* (1994) *Structure* 2:1121-1123). The antibody portions of the invention are described in further detail in U.S. Patent Nos. 6,090,382, 6,258,562, 6,509,015.

25

Still further, an antibody or antigen-binding portion thereof may be part of a larger immunoadhesion molecules, formed by covalent or non-covalent association of the antibody or antibody portion with one or more other proteins or peptides. Examples of such immunoadhesion molecules include use of the streptavidin core region to make a tetrameric scFv molecule (Kipriyanov, S.M., *et al.* (1995) *Human Antibodies and Hybridomas* 6:93-101) and use of a cysteine residue, a marker peptide and a C-terminal polyhistidine tag to make bivalent and biotinylated scFv molecules (Kipriyanov, S.M., *et*

30

al. (1994) *Mol. Immunol.* 31:1047-1058). Antibody portions, such as Fab and F(ab')₂ fragments, can be prepared from whole antibodies using conventional techniques, such as papain or pepsin digestion, respectively, of whole antibodies. Moreover, antibodies, antibody portions and immunoadhesion molecules can be obtained using standard
5 recombinant DNA techniques, as described herein.

A “clinical response” as used herein is refers to an indicator of therapeutic effectiveness of an agent. In one embodiment, a clinical response is defined by whether or not a subject achieves an HiSCR. In another embodiment, a clinical response is defined as achieving an Hidradenitis Suppurativa Physician’s Global Assessment (HS-
10 PGA) score, or HS-PGA score, as defined below in Table 1, of clear (0), minimal (1), or mild (2), with an improvement (*i.e.*, reduction) from baseline HS-PGA score of at least 2 grades. The baseline HS-PGA score is the HS-PGA score measured just prior to the commence of treatment, to which the HS-PGA score obtained after a period of treatment is compared. Both the baseline HS-PGA score and the HS-PGA score obtained after a
15 treatment period are assessed based on the system and criteria in Table 1.

The term “Hidradenitis Suppurativa Clinical Response” or “HiSCR” as used herein is defined as at least a 50% reduction in the total inflammatory lesion (abscess and nodule) count (AN count) relative to baseline with no increase in abscess count and no increase in draining fistula count.

20 A “conservative amino acid substitution,” as used herein, is one in which one amino acid residue is replaced with another amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art, including basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine,
25 glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine).

“Chimeric antibodies” refers to antibodies wherein one portion of each of the
30 amino acid sequences of heavy and light chains is homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular

class, while the remaining segment of the chains is homologous to corresponding sequences from another species. In one embodiment, the invention features a chimeric antibody or antigen-binding fragment, in which the variable regions of both light and heavy chains mimics the variable regions of antibodies derived from one species of mammals, while the constant portions are homologous to the sequences in antibodies derived from another species. In one embodiment of the invention, chimeric antibodies are made by grafting CDRs from a mouse antibody onto the framework regions of a human antibody.

“Humanized antibodies” refer to antibodies which comprise at least one chain comprising variable region framework residues substantially from a human antibody chain (referred to as the acceptor immunoglobulin or antibody) and at least one complementarity determining region (CDR) substantially from a non-human-antibody (*e.g.*, mouse). In addition to the grafting of the CDRs, humanized antibodies typically undergo further alterations in order to improve affinity and/or immunogenicity.

The term “multivalent antibody” refers to an antibody comprising more than one antigen recognition site. For example, a “bivalent” antibody has two antigen recognition sites, whereas a “tetravalent” antibody has four antigen recognition sites. The terms “monospecific,” “bispecific,” “trispecific,” “tetraspecific,” *etc.* refer to the number of different antigen recognition site specificities (as opposed to the number of antigen recognition sites) present in a multivalent antibody. For example, a “monospecific” antibody’s antigen recognition sites all bind the same epitope. A “bispecific” or “dual specific” antibody has at least one antigen recognition site that binds a first epitope and at least one antigen recognition site that binds a second epitope that is different from the first epitope. A “multivalent monospecific” antibody has multiple antigen recognition sites that all bind the same epitope. A “multivalent bispecific” antibody has multiple antigen recognition sites, some number of which bind a first epitope and some number of which bind a second epitope that is different from the first epitope.

The term “human antibody,” as used herein, is intended to include antibodies having variable and constant regions derived from human germline immunoglobulin sequences. The human antibodies of the invention may include amino acid residues not encoded by human germline immunoglobulin sequences (*e.g.*, mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*), for example

in the CDRs and in particular CDR3. However, the term “human antibody,” as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

5 The term “recombinant human antibody,” as used herein, is intended to include all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies expressed using a recombinant expression vector transfected into a host cell (described further below), antibodies isolated from a recombinant, combinatorial human antibody library (described further below), antibodies isolated
10 from an animal (*e.g.*, a mouse) that is transgenic for human immunoglobulin genes (see *e.g.*, Taylor *et al.* (1992) *Nucl. Acids Res.* 20:6287) or antibodies prepared, expressed, created or isolated by any other means that involves splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies have variable and constant regions derived from human germline immunoglobulin sequences.
15 In certain embodiments, however, such recombinant human antibodies are subjected to *in vitro* mutagenesis (or, when an animal transgenic for human Ig sequences is used, *in vivo* somatic mutagenesis) and thus the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human germline VH and VL sequences, may not naturally exist within the human
20 antibody germline repertoire *in vivo*.

Such chimeric, humanized, human, and dual specific antibodies can be produced by recombinant DNA techniques known in the art, for example using methods described in PCT International Application No. PCT/US86/02269; European Patent Application No. 184,187; European Patent Application No. 171,496; European Patent Application
25 No. 173,494; PCT International Publication No. WO 86/01533; U.S. Pat. No. 4,816,567; European Patent Application No. 125,023; Better *et al.* (1988) *Science* 240:1041-1043; Liu *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:3439-3443; Liu *et al.* (1987) *J. Immunol.* 139:3521-3526; Sun *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:214-218; Nishimura *et al.* (1987) *Cancer Res.* 47:999-1005; Wood *et al.* (1985) *Nature* 314:446-449; Shaw *et al.* (1988) *J. Natl. Cancer Inst.* 80:1553-1559; Morrison (1985) *Science* 229:1202-
30 1207; Oi *et al.* (1986) *BioTechniques* 4:214; U.S. Pat. No. 5,225,539; Jones *et al.* (1986) *Nature* 321:552-525; Verhoeyan *et al.* (1988) *Science* 239:1534; and Beidler *et al.*

(1988) *J. Immunol.* 141:4053-4060, Queen *et al.*, *Proc. Natl. Acad. Sci. USA* 86:10029-10033 (1989), US 5,530,101, US 5,585,089, US 5,693,761, US 5,693,762, Selick *et al.*, WO 90/07861, and Winter, US 5,225,539.

An “isolated antibody,” as used herein, is intended to refer to an antibody that is substantially free of other antibodies having different antigenic specificities (*e.g.*, an isolated antibody that specifically binds hTNF α is substantially free of antibodies that specifically bind antigens other than hTNF α). An isolated antibody that specifically binds hTNF α may, however, have cross-reactivity to other antigens, such as TNF α molecules from other species. Moreover, an isolated antibody may be substantially free of other cellular material and/or chemicals.

A “neutralizing antibody,” as used herein (or an “antibody that neutralized hTNF α activity”), is intended to refer to an antibody whose binding to hTNF α results in inhibition of the biological activity of hTNF α . This inhibition of the biological activity of hTNF α can be assessed by measuring one or more indicators of hTNF α biological activity, such as hTNF α -induced cytotoxicity (either *in vitro* or *in vivo*), hTNF α -induced cellular activation and hTNF α binding to hTNF α receptors. These indicators of hTNF α biological activity can be assessed by one or more of several standard *in vitro* or *in vivo* assays known in the art (see U.S. Patent No. 6,090,382). Preferably, the ability of an antibody to neutralize hTNF α activity is assessed by inhibition of hTNF α -induced cytotoxicity of L929 cells. As an additional or alternative parameter of hTNF α activity, the ability of an antibody to inhibit hTNF α -induced expression of ELAM-1 on HUVEC, as a measure of hTNF α -induced cellular activation, can be assessed.

The term “surface plasmon resonance,” as used herein, refers to an optical phenomenon that allows for the analysis of real-time biospecific interactions by detection of alterations in protein concentrations within a biosensor matrix, for example using the BIAcore™ system (Pharmacia Biosensor AB, Uppsala, Sweden and Piscataway, NJ). For further descriptions, see Example 1 of U.S. Patent 6,258,562 and Jönsson *et al.* (1993) *Ann. Biol. Clin.* 51:19; Jönsson *et al.* (1991) *Biotechniques* 11:620-627; Johnson *et al.* (1995) *J. Mol. Recognit.* 8:125; and Johnson *et al.* (1991) *Anal. Biochem.* 198: 268.

The term “ k_{off} ,” as used herein, is intended to refer to the off rate constant for dissociation of an antibody from the antibody/antigen complex.

The term “ K_d ,” as used herein, is intended to refer to the dissociation constant of a particular antibody-antigen interaction.

5 The term “ IC_{50} ” as used herein, is intended to refer to the concentration of the inhibitor required to inhibit the biological endpoint of interest, *e.g.*, neutralize cytotoxicity activity.

The term “dose,” as used herein, refers to an amount of TNF α inhibitor (*e.g.*, an anti-TNF α antibody) which is administered to a subject.

10 The term “dosing,” as used herein, refers to the administration of a TNF α inhibitor (*e.g.*, an anti-TNF α antibody) to achieve a therapeutic objective (*e.g.*, treatment of hidradenitis suppurativa).

A “dosing regimen” describes a treatment schedule for a TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof), *e.g.*, a treatment schedule
15 over a prolonged period of time or throughout the course of treatment, *e.g.* administering a first dose of a TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) at week 0 followed by a second dose of a TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) on a weekly or biweekly dosing regimen.

20 The term “multiple-variable dose” includes different doses of a TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) which are administered to a subject for therapeutic treatment. “Multiple-variable dose regimen” or “multiple-variable dose therapy” describes a treatment schedule which is based on administering different amounts of TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an
25 antigen-binding portion thereof) at various time points throughout the course of treatment. Exemplary multiple-variable dose regimens are described in PCT application no. PCT/US05/12007 and US 20060009385 .

The term “induction dose” or “loading dose,” used interchangeably herein, refers to the first dose(s) of TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding
30 portion thereof) which is initially used to treat hidradenitis suppurativa. The loading

dose may be larger in comparison to the subsequent maintenance or treatment dose. The induction dose can be a single dose or, alternatively, a set of doses. For example, a 160 mg dose may be administered as a single 160 mg dose, as two doses of 80 mg each, or four doses of 40 mg each. In one embodiment, an induction dose is subsequently
5 followed by administration of smaller doses of TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof), *e.g.*, the treatment or maintenance dose(s). The induction dose is administered during the induction or loading phase of therapy. In one embodiment of the invention, the induction dose is at least twice the given amount of the treatment dose. In one embodiment of the invention, the induction
10 dose is 80 mg. In one embodiment of the invention, the induction dose comprise a 160 mg dose followed by an 80 mg dose, wherein the two induction doses are administered 2 weeks apart.

The term “maintenance therapy” or “maintenance dosing regimen” refers to a treatment schedule for a subject or patient diagnosed with a disorder/disease, *e.g.*,
15 hidradenitis suppurativa, to enable them to maintain their health in a given state, *e.g.*, reduced number of inflammatory lesions or achieving a clinical response. In one embodiment, a maintenance therapy of the invention is used for a subject or patient diagnosed with a disorder/disease, *e.g.*, hidradenitis suppurativa to enable them to maintain their health in a state which is completely free of symptoms or a reduction in
20 symptoms associated with the disease. In one embodiment, a maintenance therapy of the invention is used for a subject or patient diagnosed with a disorder/disease, *e.g.*, hidradenitis suppurativa, to enable them to maintain their health in a state which is substantially free of symptoms associated with the disease. In one embodiment, a maintenance therapy of the invention is used for a subject or patient diagnosed with a
25 disorder/disease, *e.g.*, hidradenitis suppurativa, to enable them to maintain their health in a state where there is a significant reduction in symptoms associated with the disease.

The term “treatment phase” or “maintenance phase,” as used herein, refers to a period of treatment comprising administration of a TNF α inhibitor (*e.g.*, an anti-TNF α antibody) to a subject in order to maintain a desired therapeutic effect, *e.g.*, improved
30 symptoms associated with hidradenitis suppurativa.

The term “maintenance dose” or “treatment dose” is the amount of TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) taken by a

subject to maintain or continue a desired therapeutic effect. A maintenance dose can be a single dose or, alternatively, a set of doses. A maintenance dose is administered during the treatment or maintenance phase of therapy. In one embodiment, a maintenance dose(s) is smaller than the induction dose(s) and may be equal to each other when
5 administered in succession. In one embodiment, the invention provides a maintenance dose of 40 mg of adalimumab administered subcutaneously to a subject weekly or biweekly. In one embodiment, the maintenance dose is administered every week or every other week beginning 1 or 2 weeks after the last loading dose. In one embodiment, a maintenance dose is administered about 4 weeks following the initial loading dose.

10 The terms “biweekly dosing regimen,” “biweekly dosing,” and “biweekly administration,” as used herein, refer to the time course of administering a substance (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) to a subject to achieve a therapeutic objective, *e.g.*, throughout the course of treatment. The biweekly dosing regimen is not intended to include a weekly dosing regimen. Preferably, the
15 substance is administered every 9-19 days, or 10-18 days, more preferably, every 11-17 days, or 12-16 days, even more preferably, every 13-15 days, and most preferably, every 14 days. In one embodiment, the biweekly dosing regimen is initiated in a subject at week 0 of treatment. In another embodiment, a maintenance dose is administered on a biweekly dosing regimen. In one embodiment, both the loading and maintenance doses
20 are administered according to a biweekly dosing regimen. In one embodiment, biweekly dosing includes a dosing regimen where doses of a TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) are administered to a subject every other week consecutively for a given time period, *e.g.*, 4 weeks, 8 weeks, 16 weeks, 24 weeks, 26 weeks, 32 weeks, 36 weeks, 42 weeks, 48 weeks, 52 weeks, 56 weeks, *etc.* Biweekly
25 dosing methods are also described in US 20030235585.

Biweekly administration is also referred to as “eow” or “every other week”.

The terms “qwk,” “qw,” or “ew,” as used interchangeably herein, refer to a weekly dosing regimen, where a substance (*e.g.*, a human anti-TNF α antibody, or an antigen-binding portion thereof) is administered to a subject once a week (or every
30 week) to achieve a therapeutic objective, *e.g.*, treating HS. A “weekly dosing regimen” as used herein, refers to the time course of administering a substance (*e.g.*, an anti-TNF α antibody) to a subject to achieve a therapeutic objective, *e.g.*, throughout the course of

treatment. Weekly administration is more frequent than biweekly, *e.g.*, every 6-8 days, every 5-8 days, or every 7 days.

The term “fixed dose” or “total body dose” refers to a dose which is a constant amount delivered with each administration and is not dependent on the weight of the subject being treated. The term “fixed dose” does not include weight-based dosing, *i.e.*, mg/kg dosing determinations. In one embodiment, a human TNF α antibody, or antigen-binding portion thereof, is administered to the subject at a fixed dose ranging from 10-180 mg. In one embodiment, a human TNF α antibody, or antigen-binding portion thereof, is administered to the subject in a fixed dose of 30 mg, 35 mg, 40 mg, 45 mg, 50 mg, 55 mg, 60 mg, 65 mg, 70 mg, 75 mg, 80 mg, 85 mg, 90 mg, 95 mg, 100 mg, 105 mg, 110 mg, 115 mg, 120 mg, 125 mg, 130 mg, 135 mg, 140 mg, 145 mg, 150 mg, 155 mg, 160 mg, 165 mg, 170 mg, 175 mg, 180 mg *etc.* Ranges of values between any of the aforementioned recited values are also intended to be included in the scope of the invention, *e.g.*, 31 mg, 32 mg, 33 mg, 34 mg, 35 mg, 36 mg, 37 mg, 38 mg, 39 mg, 40 mg, 41 mg, 42 mg, 43 mg, 44 mg, 45 mg, 46 mg, 47 mg, 48 mg, 49 mg, 85 mg, 95 mg, as are ranges based on the aforementioned doses, *e.g.*, 30-50mg, 20-80 mg, 20-70 mg, 20-60 mg, and 20-50 mg.

The term “combination” as in the phrase “a first agent in combination with a second agent” includes co-administration of a first agent and a second agent, which for example may be dissolved or intermixed in the same pharmaceutically acceptable carrier, or administration of a first agent, followed by the second agent, or administration of the second agent, followed by the first agent. The present invention, therefore, includes methods of combination therapeutic treatment and combination pharmaceutical compositions.

The term “concomitant” as in the phrase “concomitant therapeutic treatment” includes administering an agent in the presence of a second agent. A concomitant therapeutic treatment method includes methods in which the first, second, third, or additional agents are co-administered. A concomitant therapeutic treatment method also includes methods in which the first or additional agents are administered in the presence of a second or additional agents, wherein the second or additional agents, for example, may have been previously administered. A concomitant therapeutic treatment method may be executed step-wise by different actors. For example, one actor may administer to

a subject a first agent and a second actor may administer to the subject a second agent, and the administering steps may be executed at the same time, or nearly the same time, or at distant times, so long as the first agent (and additional agents) are after administration in the presence of the second agent (and additional agents). The actor and
5 the subject may be the same entity (*e.g.*, human).

The term “combination therapy,” as used herein, refers to the administration of two or more therapeutic substances, *e.g.*, an anti-TNF α antibody and another drug. The other drug(s) may be administered concomitant with, prior to, or following the administration of an anti-TNF α antibody.

10 The term “treatment,” as used within the context of the present invention, is meant to include therapeutic treatment, as well as prophylactic or suppressive measures, for the treatment of hidradenitis suppurativa. For example, the term treatment may include administration of a TNF α inhibitor prior to or following the onset of hidradenitis suppurativa thereby preventing or removing signs of the disease or disorder. As another
15 example, administration of a TNF α inhibitor after clinical manifestation of hidradenitis suppurativa to combat the symptoms and/or complications and disorders associated with hidradenitis suppurativa comprises “treatment” of the disease. Further, administration of the agent after onset and after clinical symptoms and/or complications have developed where administration affects clinical parameters of the disease or disorder and perhaps
20 amelioration of the disease, comprises “treatment” of the hidradenitis suppurativa.

Those “in need of treatment” include mammals, such as humans, already having hidradenitis suppurativa, including those in which the disease or disorder is to be prevented.

The terms “subject” and “patient”, as used herein, are used interchangeably. In
25 one embodiment, a subject refers to an individual who may be treated therapeutically with a TNF α inhibitor, *e.g.*, a human TNF α antibody, or an antigen-binding portion thereof.

Various aspects of the invention are described in further detail herein.

The invention provides improved uses and compositions for treating hidradenitis
30 suppurativa disease with a TNF α inhibitor, *e.g.*, a human TNF α antibody, or an antigen-binding portion thereof. Compositions and articles of manufacture, including kits,

relating to the methods and uses for treating hidradenitis suppurativa are also contemplated as part of the invention.

II. TNF Inhibitors

5 A TNF α inhibitor which is used in the methods and compositions of the invention includes any agent which interferes with TNF α activity. In a preferred embodiment, the TNF α inhibitor can neutralize TNF α activity, particularly detrimental TNF α activity which is associated with hidradenitis suppurativa, and related complications and symptoms.

10 In one embodiment, the TNF α inhibitor used in the invention is an TNF α antibody (also referred to herein as a TNF α antibody), or an antigen-binding fragment thereof, including chimeric, humanized, and human antibodies. Examples of TNF α antibodies which may be used in the invention include, but not limited to, infliximab (REMICADE[®], Johnson and Johnson; described in U.S. Patent No. 5,656,272),
15 CDP571 (a humanized monoclonal anti-TNF-alpha IgG4 antibody), CDP 870 (a humanized monoclonal anti-TNF-alpha antibody fragment), an anti-TNF dAb (Peptech), SIMPONI (golimumab) (also referred to as CNTO 148; Centocor Ortho Biotech, see WO 02/12502), CIMZIA (certolizumab) (UCB), and adalimumab (HUMIRA[®], Abbott Laboratories, a human anti-TNF mAb, described in US
20 6,090,382 as D2E7). Additional TNF antibodies which may be used in the invention are described in U.S. Patent Nos. 6,593,458; 6,498,237; 6,451,983; and 6,448,380 .

Other examples of TNF α inhibitors which may be used in the methods and compositions of the invention include etanercept (ENBREL[™], described in WO
25 91/03553 and WO 09/406476), soluble TNF receptor Type I, a PEGylated soluble TNF receptor Type I (PEGs TNF-R1), p55 TNFRIgG (Lenercept), and recombinant TNF binding protein (r-TBP-I) (Serono).

In one embodiment, the term "TNF α inhibitor" excludes infliximab. In one embodiment, the term "TNF α inhibitor" excludes adalimumab. In another embodiment,
30 the term "TNF α inhibitor" excludes adalimumab and infliximab.

In one embodiment, the term “TNF α inhibitor” excludes etanercept, and, optionally, adalimumab, infliximab, and adalimumab and infliximab.

In one embodiment, the term “TNF α antibody” excludes infliximab. In one embodiment, the term “TNF α antibody” excludes adalimumab. In another embodiment,
5 the term “TNF α antibody” excludes adalimumab and infliximab.

In one embodiment, the invention features uses and composition for treating or determining the efficacy of a TNF α antibody, or antigen binding portion thereof, for the treatment of hidradenitis suppurativa, wherein the TNF α antibody, or antigen binding portion thereof, is an isolated human antibody, or antigen-binding portion thereof, that
10 binds to human TNF α with high affinity and a low off rate, and also has a high neutralizing capacity. Preferably, the human antibodies used in the invention are recombinant, neutralizing human anti-hTNF α antibodies. The most preferred recombinant, neutralizing antibody of the invention is HUMIRA[®] (adalimumab; also referred to as D2E7 (the amino acid sequence of the adalimumab VL region is shown in
15 SEQ ID NO: 1; the amino acid sequence of the adalimumab VH region is shown in SEQ ID NO: 2). The properties and sequences of adalimumab / HUMIRA[®] (also referred to as D2E7) have been described in Salfeld *et al.*, U.S. Patent Nos. 6,090,382, 6,258,562, and 6,509,015.

The methods of the invention may also be performed using chimeric and
20 humanized murine anti-hTNF α antibodies which have undergone clinical testing for treatment of rheumatoid arthritis (see *e.g.*, Elliott, M.J., *et al.* (1994) *Lancet* 344:1125-1127; Elliot, M.J., *et al.* (1994) *Lancet* 344:1105-1110; Rankin, E.C., *et al.* (1995) *Br. J. Rheumatol.* 34:334-342).

In one embodiment, the method of the invention includes determining the
25 efficacy of adalimumab and antibody portions, adalimumab-related antibodies and antibody portions, or other human antibodies and antibody portions with equivalent properties to adalimumab, such as high affinity binding to hTNF α with low dissociation kinetics and high neutralizing capacity, for the treatment of hidradenitis suppurativa. In one embodiment, the invention provides treatment with an isolated human antibody, or
30 an antigen-binding portion thereof, that dissociates from human TNF α with a K_d of 1×10^{-8} M or less and a k_{off} rate constant of $1 \times 10^{-3} \text{ s}^{-1}$ or less, both determined by surface

plasmon resonance, and neutralizes human TNF α cytotoxicity in a standard *in vitro* L929 assay with an IC₅₀ of 1×10^{-7} M or less. More preferably, the isolated human antibody, or antigen-binding portion thereof, dissociates from human TNF α with a k_{off} of $5 \times 10^{-4} \text{ s}^{-1}$ or less, or even more preferably, with a k_{off} of $1 \times 10^{-4} \text{ s}^{-1}$ or less. More
5 preferably, the isolated human antibody, or antigen-binding portion thereof, neutralizes human TNF α cytotoxicity in a standard *in vitro* L929 assay with an IC₅₀ of 1×10^{-8} M or less, even more preferably with an IC₅₀ of 1×10^{-9} M or less and still more preferably with an IC₅₀ of 1×10^{-10} M or less. Notably, the aforementioned characteristics are all properties of adalimumab. In a preferred embodiment, the antibody is an isolated human
10 recombinant antibody, or an antigen-binding portion thereof.

It is well known in the art that antibody heavy and light chain CDR3 domains play an important role in the binding specificity/affinity of an antibody for an antigen. Accordingly, in another aspect, the invention pertains to treating hidradenitis suppurativa by administering human antibodies that have slow dissociation kinetics for association
15 with hTNF α and that have light and heavy chain CDR3 domains that structurally are identical to or related to those of adalimumab. Position 9 of the adalimumab VL CDR3 can be occupied by Ala or Thr without substantially affecting the K_{off} . Accordingly, a consensus motif for the adalimumab VL CDR3 comprises the amino acid sequence: Q-R-Y-N-R-A-P-Y-(T/A) (SEQ ID NO: 3). Additionally, position 12 of the adalimumab
20 VH CDR3 can be occupied by Tyr or Asn, without substantially affecting the K_{off} . Accordingly, a consensus motif for the adalimumab VH CDR3 comprises the amino acid sequence: V-S-Y-L-S-T-A-S-S-L-D-(Y/N) (SEQ ID NO: 4). Moreover, as demonstrated in Example 2 of U.S. Patent No. 6,090,382, the CDR3 domain of the adalimumab heavy and light chains is amenable to substitution with a single alanine
25 residue (at position 1, 4, 5, 7 or 8 within the VL CDR3 or at position 2, 3, 4, 5, 6, 8, 9, 10 or 11 within the VH CDR3) without substantially affecting the k_{off} . Still further, the skilled artisan will appreciate that, given the amenability of the adalimumab VL and VH CDR3 domains to substitutions by alanine, substitution of other amino acids within the CDR3 domains may be possible while still retaining the low off rate constant of the
30 antibody, in particular substitutions with conservative amino acids. Preferably, no more than one to five conservative amino acid substitutions are made within the adalimumab VL and/or VH CDR3 domains. More preferably, no more than one to three conservative

amino acid substitutions are made within the adalimumab VL and/or VH CDR3 domains. Additionally, conservative amino acid substitutions should not be made at amino acid positions critical for binding to hTNF α . Positions 2 and 5 of the adalimumab VL CDR3 and positions 1 and 7 of the adalimumab VH CDR3 appear to be critical for
5 interaction with hTNF α and thus, conservative amino acid substitutions preferably are not made at these positions (although an alanine substitution at position 5 of the adalimumab VL CDR3 is acceptable, as described above) (*see* U.S. Patent No. 6,090,382).

Accordingly, in another embodiment, the antibody or antigen-binding portion
10 thereof preferably contains the following characteristics:

- a) dissociates from human TNF α with a k_{off} rate constant of $1 \times 10^{-3} \text{ s}^{-1}$ or less, as determined by surface plasmon resonance;
- b) has a light chain CDR3 domain comprising the amino acid sequence of SEQ ID NO: 3, or modified from SEQ ID NO: 3 by a single alanine substitution at position 1,
15 4, 5, 7 or 8 or by one to five conservative amino acid substitutions at positions 1, 3, 4, 6, 7, 8 and/or 9;
- c) has a heavy chain CDR3 domain comprising the amino acid sequence of SEQ ID NO: 4, or modified from SEQ ID NO: 4 by a single alanine substitution at position 2, 3, 4, 5, 6, 8, 9, 10 or 11 or by one to five conservative amino acid substitutions at
20 positions 2, 3, 4, 5, 6, 8, 9, 10, 11 and/or 12.

More preferably, the antibody, or antigen-binding portion thereof, dissociates from human TNF α with a k_{off} of $5 \times 10^{-4} \text{ s}^{-1}$ or less. Even more preferably, the antibody, or antigen-binding portion thereof, dissociates from human TNF α with a k_{off} of $1 \times 10^{-4} \text{ s}^{-1}$ or less.

25 In yet another embodiment, the antibody or antigen-binding portion thereof preferably contains a light chain variable region (LCVR) having a CDR3 domain comprising the amino acid sequence of SEQ ID NO: 3, or modified from SEQ ID NO: 3 by a single alanine substitution at position 1, 4, 5, 7 or 8, and with a heavy chain variable region (HCVR) having a CDR3 domain comprising the amino acid sequence of SEQ ID
30 NO: 4, or modified from SEQ ID NO: 4 by a single alanine substitution at position 2, 3, 4, 5, 6, 8, 9, 10 or 11. In one embodiment, the LCVR further has a CDR2 domain

comprising the amino acid sequence of SEQ ID NO: 5 (*i.e.*, the adalimumab VL CDR2) and the HCVR further has a CDR2 domain comprising the amino acid sequence of SEQ ID NO: 6 (*i.e.*, the adalimumab VH CDR2). In one embodiment, the LCVR further has CDR1 domain comprising the amino acid sequence of SEQ ID NO: 7 (*i.e.*, the
5 adalimumab VL CDR1) and the HCVR has a CDR1 domain comprising the amino acid sequence of SEQ ID NO: 8 (*i.e.*, the adalimumab VH CDR1). The framework regions for VL preferably are from the V_κI human germline family, more preferably from the A20 human germline V_κ gene and most preferably from the D2E7 VL framework sequences shown in Figures 1A and 1B of U.S. Patent No. 6,090,382. In one
10 embodiment, the framework regions for VH are from the VH3 human germline family, from the DP-31 human germline VH gene, or from the adalimumab VH framework sequences shown in Figures 2A and 2B of U.S. Patent No. 6,090,382.

In one embodiment, the anti-TNF α antibody, or antigen binding portion thereof, used in the methods and compositions of the invention comprises the six CDRs of
15 adalimumab (*i.e.*, SEQ ID NOs: 3 to 8).

Accordingly, in another embodiment, the antibody or antigen-binding portion thereof preferably contains a light chain variable region (LCVR) comprising the amino acid sequence of SEQ ID NO: 1 (*i.e.*, the adalimumab VL) and a heavy chain variable region (HCVR) comprising the amino acid sequence of SEQ ID NO: 2 (*i.e.*, the
20 adalimumab VH). In certain embodiments, the antibody comprises a heavy chain constant region, such as an IgG1, IgG2, IgG3, IgG4, IgA, IgE, IgM or IgD constant region. Preferably, the heavy chain constant region is an IgG1 heavy chain constant region or an IgG4 heavy chain constant region. Furthermore, the antibody can comprise a light chain constant region, either a kappa light chain constant region or a lambda light
25 chain constant region. Preferably, the antibody comprises a kappa light chain constant region. Alternatively, the antibody portion can be, for example, a Fab fragment or a single chain Fv fragment.

In one embodiment, the anti-TNF α antibody, or antigen binding portion thereof, used in the methods and compositions of the invention comprises the variable light
30 and/or heavy chain as described in SEQ ID NOs: 9 and 10, respectively.

Also included in the methods and compositions of the invention are antibodies that are bioequivalent to adalimumab. In one embodiment, the methods and compositions of the invention include an antibody that is a biosimilar antibody to adalimumab, also referred to as an “adalimumab biosimilar antibody”. An adalimumab biosimilar antibody is highly similar to adalimumab notwithstanding minor differences in clinically inactive components, and for which there are no clinically meaningful differences between the adalimumab biosimilar antibody and adalimumab in terms of the safety, purity and potency. In another embodiment, the methods and compositions of the invention include an antibody that is an interchangeable antibody to adalimumab, also referred to as an “adalimumab interchangeable antibody”. An adalimumab interchangeable antibody is an adalimumab biosimilar antibody (as defined above) that is also expected to produce the same clinical result as adalimumab in any given patient. Furthermore, with respect to an adalimumab interchangeable antibody, the risk in terms of safety or diminished efficacy in alternating or switching between the use of the adalimumab interchangeable antibody and adalimumab must not be greater than the risk of using adalimumab without such alteration or switch.

In one embodiment, the invention is directed to an antibody having a light and/or heavy chain variable region having an amino acid sequence which is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to SEQ ID NO: 1 or SEQ ID NO: 2 (adalimumab light and heavy chain sequences, respectively). Preferably, the antibody is an IgG, such as an IgG1 or IgG4 antibody.

In still other embodiments, the invention includes uses of an isolated human antibody, or an antigen-binding portions thereof, containing adalimumab-related VL and VH CDR3 domains. For example, antibodies, or antigen-binding portions thereof, with a light chain variable region (LCVR) having a CDR3 domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 3, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20, SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25 and SEQ ID NO: 26 or with a heavy chain variable region (HCVR) having a CDR3 domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 4, SEQ ID NO: 27,

SEQ ID NO: 28, SEQ ID NO: 29, SEQ ID NO: 30, SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34 and SEQ ID NO: 35.

The TNF α antibody used in the methods and compositions of the invention may be modified for improved treatment of hidradenitis suppurativa. In some embodiments, the TNF α antibody or antigen binding fragments thereof, is chemically modified to provide a desired effect. For example, PEGylation of antibodies and antibody fragments of the invention may be carried out by any of the PEGylation reactions known in the art, as described, for example, in the following references: *Focus on Growth Factors* 3:4-10 (1992); EP 0 154 316; and EP 0 401 384 .

Preferably, the PEGylation is carried out via an acylation reaction or an alkylation reaction with a reactive polyethylene glycol molecule (or an analogous reactive water-soluble polymer). A preferred water-soluble polymer for PEGylation of the antibodies and antibody fragments of the invention is polyethylene glycol (PEG). As used herein, "polyethylene glycol" is meant to encompass any of the forms of PEG that have been used to derivatize other proteins, such as mono (C1-C10) alkoxy- or aryloxy-polyethylene glycol.

Methods for preparing PEGylated antibodies and antibody fragments of the invention will generally comprise the steps of (a) reacting the antibody or antibody fragment with polyethylene glycol, such as a reactive ester or aldehyde derivative of PEG, under conditions whereby the antibody or antibody fragment becomes attached to one or more PEG groups, and (b) obtaining the reaction products. It will be apparent to one of ordinary skill in the art to select the optimal reaction conditions or the acylation reactions based on known parameters and the desired result.

PEGylated antibodies and antibody fragments may generally be used to treat hidradenitis suppurativa by administration of the TNF α antibodies and antibody fragments described herein. Generally the PEGylated antibodies and antibody fragments have increased half-life, as compared to the nonpegylated antibodies and antibody fragments. The PEGylated antibodies and antibody fragments may be employed alone, together, or in combination with other pharmaceutical compositions.

In yet another embodiment of the invention, TNF α antibodies or fragments thereof can be altered wherein the constant region of the antibody is modified to reduce

at least one constant region-mediated biological effector function relative to an unmodified antibody. To modify an antibody of the invention such that it exhibits reduced binding to the Fc receptor, the immunoglobulin constant region segment of the antibody can be mutated at particular regions necessary for Fc receptor (FcR)

5 interactions (see *e.g.*, Canfield, S.M. and S.L. Morrison (1991) *J. Exp. Med.* 173:1483-1491; and Lund, J. *et al.* (1991) *J. of Immunol.* 147:2657-2662). Reduction in FcR binding ability of the antibody may also reduce other effector functions which rely on FcR interactions, such as opsonization and phagocytosis and antigen-dependent cellular cytotoxicity.

10 An antibody or antibody portion used in the methods of the invention can be derivatized or linked to another functional molecule (*e.g.*, another peptide or protein). Accordingly, the antibodies and antibody portions of the invention are intended to include derivatized and otherwise modified forms of the human anti-hTNF α antibodies described herein, including immunoadhesion molecules. For example, an antibody or
15 antibody portion of the invention can be functionally linked (by chemical coupling, genetic fusion, non-covalent association or otherwise) to one or more other molecular entities, such as another antibody (*e.g.*, a bispecific antibody or a diabody), a detectable agent, a cytotoxic agent, a pharmaceutical agent, and/or a protein or peptide that can mediate associate of the antibody or antibody portion with another molecule (such as a
20 streptavidin core region or a polyhistidine tag).

One type of derivatized antibody is produced by cross-linking two or more antibodies (of the same type or of different types, *e.g.*, to create bispecific antibodies). Suitable cross-linkers include those that are heterobifunctional, having two distinctly reactive groups separated by an appropriate spacer (*e.g.*, m-maleimidobenzoyl-N-
25 hydroxysuccinimide ester) or homobifunctional (*e.g.*, disuccinimidyl suberate). Such linkers are available from Pierce Chemical Company, Rockford, IL.

Useful detectable agents with which an antibody or antibody portion of the invention may be derivatized include fluorescent compounds. Exemplary fluorescent detectable agents include fluorescein, fluorescein isothiocyanate, rhodamine, 5-
30 dimethylamine-1-naphthalenesulfonyl chloride, phycoerythrin and the like. An antibody may also be derivatized with detectable enzymes, such as alkaline phosphatase, horseradish peroxidase, glucose oxidase and the like. When an antibody is derivatized

with a detectable enzyme, it is detected by adding additional reagents that the enzyme uses to produce a detectable reaction product. For example, when the detectable agent horseradish peroxidase is present, the addition of hydrogen peroxide and diaminobenzidine leads to a colored reaction product, which is detectable. An antibody
5 may also be derivatized with biotin, and detected through indirect measurement of avidin or streptavidin binding.

An antibody, or antibody portion, used in the methods and compositions of the invention, can be prepared by recombinant expression of immunoglobulin light and heavy chain genes in a host cell. To express an antibody recombinantly, a host cell is
10 transfected with one or more recombinant expression vectors carrying DNA fragments encoding the immunoglobulin light and heavy chains of the antibody such that the light and heavy chains are expressed in the host cell and, preferably, secreted into the medium in which the host cells are cultured, from which medium the antibodies can be recovered. Standard recombinant DNA methodologies are used to obtain antibody heavy
15 and light chain genes, incorporate these genes into recombinant expression vectors and introduce the vectors into host cells, such as those described in Sambrook, Fritsch and Maniatis (eds), *Molecular Cloning; A Laboratory Manual, Second Edition*, Cold Spring Harbor, N.Y., (1989), Ausubel, F.M. *et al.* (eds.) *Current Protocols in Molecular Biology*, Greene Publishing Associates, (1989) and in U.S. Patent No. 4,816,397 by Boss
20 *et al.*

To express adalimumab (D2E7) or an adalimumab (D2E7)-related antibody, DNA fragments encoding the light and heavy chain variable regions are first obtained. These DNAs can be obtained by amplification and modification of germline light and heavy chain variable sequences using the polymerase chain reaction (PCR). Germline
25 DNA sequences for human heavy and light chain variable region genes are known in the art (see *e.g.*, the "Vbase" human germline sequence database; see also Kabat, E.A., *et al.* (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242; Tomlinson, I.M., *et al.* (1992) "The Repertoire of Human Germline V_H Sequences Reveals about Fifty Groups
30 of V_H Segments with Different Hypervariable Loops" *J. Mol. Biol.* 227:776-798; and Cox, J.P.L. *et al.* (1994) "A Directory of Human Germ-line V₇₈ Segments Reveals a Strong Bias in their Usage" *Eur. J. Immunol.* 24:827-836 .

To obtain a DNA fragment encoding the heavy chain variable region of D2E7, or a D2E7-related antibody, a member of the V_H3 family of human germline VH genes is amplified by standard PCR. Most preferably, the DP-31 VH germline sequence is amplified. To obtain a DNA fragment
5 encoding the light chain variable region of adalimumab, or a adalimumab -related antibody, a member of the V_KI family of human germline VL genes is amplified by standard PCR. Most preferably, the A20 VL germline sequence is amplified. PCR primers suitable for use in amplifying the DP-31 germline VH and A20 germline VL sequences can be designed based on the nucleotide sequences disclosed in the references
10 cited *supra*, using standard methods. The variable light and heavy chain nucleic acid sequences of adalimumab are described in SEQ ID NOs: 36 and 37, respectively.

Once the germline VH and VL fragments are obtained, these sequences can be mutated to encode the adalimumab or adalimumab -related amino acid sequences disclosed herein. The amino acid sequences encoded by the germline VH and VL DNA
15 sequences are first compared to the adalimumab or adalimumab -related VH and VL amino acid sequences to identify amino acid residues in the adalimumab or adalimumab -related sequence that differ from germline. Then, the appropriate nucleotides of the germline DNA sequences are mutated such that the mutated germline sequence encodes the adalimumab or adalimumab -related amino acid sequence, using the genetic code to
20 determine which nucleotide changes should be made. Mutagenesis of the germline sequences is carried out by standard methods, such as PCR-mediated mutagenesis (in which the mutated nucleotides are incorporated into the PCR primers such that the PCR product contains the mutations) or site-directed mutagenesis.

Moreover, it should be noted that if the “germline” sequences obtained by PCR
25 amplification encode amino acid differences in the framework regions from the true germline configuration (*i.e.*, differences in the amplified sequence as compared to the true germline sequence, for example as a result of somatic mutation), it may be desirable to change these amino acid differences back to the true germline sequences (*i.e.*, “backmutation” of framework residues to the germline configuration).

30 Once DNA fragments encoding adalimumab or adalimumab related VH and VL segments are obtained (by amplification and mutagenesis of germline VH and VL genes, as described above), these DNA fragments can be further manipulated by standard

recombinant DNA techniques, for example to convert the variable region genes to full-length antibody chain genes, to Fab fragment genes or to a scFv gene. In these manipulations, a VL- or VH-encoding DNA fragment is operatively linked to another DNA fragment encoding another protein, such as an antibody constant region or a flexible linker. The term “operatively linked,” as used in this context, is intended to mean that the two DNA fragments are joined such that the amino acid sequences encoded by the two DNA fragments remain in-frame.

The isolated DNA encoding the VH region can be converted to a full-length heavy chain gene by operatively linking the VH-encoding DNA to another DNA molecule encoding heavy chain constant regions (CH1, CH2 and CH3). The sequences of human heavy chain constant region genes are known in the art (see *e.g.*, Kabat, E.A., *et al.* (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The heavy chain constant region can be an IgG1, IgG2, IgG3, IgG4, IgA, IgE, IgM or IgD constant region, but most preferably is an IgG1 or IgG4 constant region. For a Fab fragment heavy chain gene, the VH-encoding DNA can be operatively linked to another DNA molecule encoding only the heavy chain CH1 constant region.

The isolated DNA encoding the VL region can be converted to a full-length light chain gene (as well as a Fab light chain gene) by operatively linking the VL-encoding DNA to another DNA molecule encoding the light chain constant region, CL. The sequences of human light chain constant region genes are known in the art (see *e.g.*, Kabat, E.A., *et al.* (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The light chain constant region can be a kappa or lambda constant region, but most preferably is a kappa constant region.

To create a scFv gene, the VH- and VL-encoding DNA fragments are operatively linked to another fragment encoding a flexible linker, *e.g.*, encoding the amino acid sequence (Gly₄-Ser)₃, such that the VH and VL sequences can be expressed as a contiguous single-chain protein, with the VL and VH regions joined by the flexible

linker (see *e.g.*, Bird *et al.* (1988) *Science* 242:423-426; Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883; McCafferty *et al.*, *Nature* (1990) 348:552-554).

To express the antibodies, or antibody portions used in the invention, DNAs encoding partial or full-length light and heavy chains, obtained as described above, are
5 inserted into expression vectors such that the genes are operatively linked to transcriptional and translational control sequences. In this context, the term “operatively linked” is intended to mean that an antibody gene is ligated into a vector such that transcriptional and translational control sequences within the vector serve their intended function of regulating the transcription and translation of the antibody gene. The
10 expression vector and expression control sequences are chosen to be compatible with the expression host cell used. The antibody light chain gene and the antibody heavy chain gene can be inserted into separate vector or, more typically, both genes are inserted into the same expression vector. The antibody genes are inserted into the expression vector by standard methods (*e.g.*, ligation of complementary restriction sites on the antibody
15 gene fragment and vector, or blunt end ligation if no restriction sites are present). Prior to insertion of the adalimumab or adalimumab -related light or heavy chain sequences, the expression vector may already carry antibody constant region sequences. For example, one approach to converting the adalimumab or adalimumab -related VH and VL sequences to full-length antibody genes is to insert them into expression vectors
20 already encoding heavy chain constant and light chain constant regions, respectively, such that the VH segment is operatively linked to the CH segment(s) within the vector and the VL segment is operatively linked to the CL segment within the vector. Additionally or alternatively, the recombinant expression vector can encode a signal peptide that facilitates secretion of the antibody chain from a host cell. The antibody
25 chain gene can be cloned into the vector such that the signal peptide is linked in-frame to the amino terminus of the antibody chain gene. The signal peptide can be an immunoglobulin signal peptide or a heterologous signal peptide (*i.e.*, a signal peptide from a non-immunoglobulin protein).

In addition to the antibody chain genes, the recombinant expression vectors of
30 the invention carry regulatory sequences that control the expression of the antibody chain genes in a host cell. The term “regulatory sequence” is intended to include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals) that

control the transcription or translation of the antibody chain genes. Such regulatory sequences are described, for example, in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). It will be appreciated by those skilled in the art that the design of the expression vector, including
5 the selection of regulatory sequences may depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, *etc.* Preferred regulatory sequences for mammalian host cell expression include viral elements that direct high levels of protein expression in mammalian cells, such as promoters and/or enhancers derived from cytomegalovirus (CMV) (such as the CMV promoter/enhancer),
10 Simian Virus 40 (SV40) (such as the SV40 promoter/enhancer), adenovirus, (*e.g.*, the adenovirus major late promoter (AdMLP)) and polyoma. For further description of viral regulatory elements, and sequences thereof, see *e.g.*, U.S. Patent No. 5,168,062 by Stinski, U.S. Patent No. 4,510,245 by Bell *et al.* and U.S. Patent No. 4,968,615 by Schaffner *et al.*

15 In addition to the antibody chain genes and regulatory sequences, the recombinant expression vectors used in the invention may carry additional sequences, such as sequences that regulate replication of the vector in host cells (*e.g.*, origins of replication) and selectable marker genes. The selectable marker gene facilitates selection of host cells into which the vector has been introduced (see *e.g.*, U.S. Patents
20 Nos. 4,399,216, 4,634,665 and 5,179,017, all by Axel *et al.*). For example, typically the selectable marker gene confers resistance to drugs, such as G418, hygromycin or methotrexate, on a host cell into which the vector has been introduced. Preferred selectable marker genes include the dihydrofolate reductase (DHFR) gene (for use in DHFR⁻ host cells with methotrexate selection/amplification) and the *neo* gene (for G418
25 selection).

For expression of the light and heavy chains, the expression vector(s) encoding the heavy and light chains is transfected into a host cell by standard techniques. The various forms of the term “transfection” are intended to encompass a wide variety of techniques commonly used for the introduction of exogenous DNA into a prokaryotic or
30 eukaryotic host cell, *e.g.*, electroporation, calcium-phosphate precipitation, DEAE-dextran transfection and the like. Although it is theoretically possible to express the antibodies of the invention in either prokaryotic or eukaryotic host cells, expression of

antibodies in eukaryotic cells, and most preferably mammalian host cells, is the most preferred because such eukaryotic cells, and in particular mammalian cells, are more likely than prokaryotic cells to assemble and secrete a properly folded and immunologically active antibody. Prokaryotic expression of antibody genes has been
5 reported to be ineffective for production of high yields of active antibody (Boss, M.A. and Wood, C. R. (1985) *Immunology Today* 6:12-13).

Preferred mammalian host cells for expressing the recombinant antibodies of the invention include Chinese Hamster Ovary (CHO cells) (including dhfr- CHO cells, described in Urlaub and Chasin, (1980) *Proc. Natl. Acad. Sci. USA* 77:4216-4220, used
10 with a DHFR selectable marker, *e.g.*, as described in R.J. Kaufman and P.A. Sharp (1982) *Mol. Biol.* 159:601-621), NS0 myeloma cells, COS cells and SP2 cells. When recombinant expression vectors encoding antibody genes are introduced into mammalian host cells, the antibodies are produced by culturing the host cells for a period of time sufficient to allow for expression of the antibody in the host cells or, more preferably,
15 secretion of the antibody into the culture medium in which the host cells are grown. Antibodies can be recovered from the culture medium using standard protein purification methods.

Host cells can also be used to produce portions of intact antibodies, such as Fab fragments or scFv molecules. It is understood that variations on the above procedure are
20 within the scope of the present invention. For example, it may be desirable to transfect a host cell with DNA encoding either the light chain or the heavy chain (but not both) of an antibody of this invention. Recombinant DNA technology may also be used to remove some or all of the DNA encoding either or both of the light and heavy chains that is not necessary for binding to hTNF α . The molecules expressed from such truncated
25 DNA molecules are also encompassed by the antibodies of the invention. In addition, bifunctional antibodies may be produced in which one heavy and one light chain are an antibody of the invention and the other heavy and light chain are specific for an antigen other than hTNF α by cross-linking an antibody of the invention to a second antibody by standard chemical cross-linking methods.

30 In a preferred system for recombinant expression of an antibody, or antigen-binding portion thereof, of the invention, a recombinant expression vector encoding both the antibody heavy chain and the antibody light chain is introduced into dhfr-CHO cells

by calcium phosphate-mediated transfection. Within the recombinant expression vector, the antibody heavy and light chain genes are each operatively linked to CMV enhancer/AdMLP promoter regulatory elements to drive high levels of transcription of the genes. The recombinant expression vector also carries a DHFR gene, which allows
5 for selection of CHO cells that have been transfected with the vector using methotrexate selection/amplification. The selected transformant host cells are culture to allow for expression of the antibody heavy and light chains and intact antibody is recovered from the culture medium. Standard molecular biology techniques are used to prepare the recombinant expression vector, transfect the host cells, select for transformants, culture
10 the host cells and recover the antibody from the culture medium.

In view of the foregoing, nucleic acid, vector and host cell compositions that can be used for recombinant expression of the antibodies and antibody portions used in the invention include nucleic acids, and vectors comprising said nucleic acids, comprising the human TNF α antibody adalimumab (D2E7). The nucleotide sequence encoding the
15 adalimumab light chain variable region is shown in SEQ ID NO: 36. The CDR1 domain of the LCVR encompasses nucleotides 70-102, the CDR2 domain encompasses nucleotides 148-168 and the CDR3 domain encompasses nucleotides 265-291. The nucleotide sequence encoding the adalimumab heavy chain variable region is shown in SEQ ID NO: 37. The CDR1 domain of the HCVR encompasses nucleotides 91-105, the
20 CDR2 domain encompasses nucleotides 148-198 and the CDR3 domain encompasses nucleotides 295-330. It will be appreciated by the skilled artisan that nucleotide sequences encoding D2E7-related antibodies, or portions thereof (*e.g.*, a CDR domain, such as a CDR3 domain), can be derived from the nucleotide sequences encoding the adalimumab LCVR and HCVR using the genetic code and standard molecular biology
25 techniques.

Recombinant human antibodies of the invention in addition to adalimumab or an antigen binding portion thereof, or adalimumab -related antibodies disclosed herein can be isolated by screening of a recombinant combinatorial antibody library, preferably a scFv phage display library, prepared using human VL and VH cDNAs prepared from
30 mRNA derived from human lymphocytes. Methodologies for preparing and screening such libraries are known in the art. In addition to commercially available kits for generating phage display libraries (*e.g.*, the Pharmacia *Recombinant Phage Antibody*

System, catalog no. 27-9400-01; and the Stratagene *SurfZAP*TM phage display kit, catalog no. 240612), examples of methods and reagents particularly amenable for use in generating and screening antibody display libraries can be found in, for example, Ladner *et al.* U.S. Patent No. 5,223,409; Kang *et al.* PCT Publication No. WO 92/18619; Dower *et al.* PCT Publication No. WO 91/17271; Winter *et al.* PCT Publication No. WO 92/20791; Markland *et al.* PCT Publication No. WO 92/15679; Breitling *et al.* PCT Publication No. WO 93/01288; McCafferty *et al.* PCT Publication No. WO 92/01047; Garrard *et al.* PCT Publication No. WO 92/09690; Fuchs *et al.* (1991) *Bio/Technology* 9:1370-1372; Hay *et al.* (1992) *Hum Antibod Hybridomas* 3:81-65; Huse *et al.* (1989) *Science* 246:1275-1281; McCafferty *et al.*, *Nature* (1990) 348:552-554; Griffiths *et al.* (1993) *EMBO J* 12:725-734; Hawkins *et al.* (1992) *J Mol Biol* 226:889-896; Clackson *et al.* (1991) *Nature* 352:624-628; Gram *et al.* (1992) *PNAS* 89:3576-3580; Garrard *et al.* (1991) *Bio/Technology* 9:1373-1377; Hoogenboom *et al.* (1991) *Nuc Acid Res* 19:4133-4137; and Barbas *et al.* (1991) *PNAS* 88:7978-7982.

15 In a preferred embodiment, to isolate human antibodies with high affinity and a low off rate constant for hTNF α , a murine anti-hTNF α antibody having high affinity and a low off rate constant for hTNF α (*e.g.*, MAK 195, the hybridoma for which has deposit number ECACC 87 050801) is first used to select human heavy & light chain sequences having similar binding activity toward hTNF α , using the epitope imprinting methods
20 described in Hoogenboom *et al.*, WO 93/06213. The antibody libraries used in this method are preferably scFv libraries prepared and screened as described in McCafferty *et al.*, PCT Publication No. WO 92/01047, McCafferty *et al.*, *Nature* (1990) 348:552-554; and Griffiths *et al.*, (1993) *EMBO J* 12:725-734. The scFv antibody libraries preferably are screened using recombinant human TNF α as the antigen.

25 Once initial human VL and VH segments are selected, “mix and match” experiments, in which different pairs of the initially selected VL and VH segments are screened for hTNF α binding, are performed to select preferred VL/VH pair combinations. Additionally, to further improve the affinity and/or lower the off rate constant for hTNF α binding, the VL and VH segments of the preferred VL/VH pair(s)
30 can be randomly mutated, preferably within the CDR3 region of VH and/or VL, in a process analogous to the *in vivo* somatic mutation process responsible for affinity maturation of antibodies during a natural immune response. This *in vitro* affinity

maturation can be accomplished by amplifying VH and VL regions using PCR primers complimentary to the VH CDR3 or VL CDR3, respectively, which primers have been “spiked” with a random mixture of the four nucleotide bases at certain positions such that the resultant PCR products encode VH and VL segments into which random
5 mutations have been introduced into the VH and/or VL CDR3 regions. These randomly mutated VH and VL segments can be re-screened for binding to hTNF α and sequences that exhibit high affinity and a low off rate for hTNF α binding can be selected.

Following screening and isolation of an anti-hTNF α antibody of the invention from a recombinant immunoglobulin display library, nucleic acid encoding the selected
10 antibody can be recovered from the display package (*e.g.*, from the phage genome) and subcloned into other expression vectors by standard recombinant DNA techniques. If desired, the nucleic acid can be further manipulated to create other antibody forms of the invention (*e.g.*, linked to nucleic acid encoding additional immunoglobulin domains, such as additional constant regions). To express a recombinant human antibody isolated
15 by screening of a combinatorial library, the DNA encoding the antibody is cloned into a recombinant expression vector and introduced into a mammalian host cells, as described in further detail in above.

Methods of isolating human neutralizing antibodies with high affinity and a low off rate constant for hTNF α are described in U.S. Patent Nos. 6,090,382, 6,258,562, and
20 6,509,015 .

Antibodies, antibody-portions, and other TNF α inhibitors for use in the methods of the invention, can be incorporated into pharmaceutical compositions suitable for administration to a subject. Typically, the pharmaceutical composition comprises an antibody, antibody portion, or other TNF α inhibitor, and a pharmaceutically acceptable
25 carrier. As used herein, “pharmaceutically acceptable carrier” includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. Examples of pharmaceutically acceptable carriers include one or more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol and the like, as well as combinations thereof.
30 In many cases, it is preferable to include isotonic agents, *e.g.*, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Pharmaceutically

acceptable carriers may further comprise minor amounts of auxiliary substances such as wetting or emulsifying agents, preservatives or buffers, which enhance the shelf life or effectiveness of the antibody, antibody portion, or other TNF α inhibitor.

The compositions for use in the methods and compositions of the invention may
5 be in a variety of forms. These include, for example, liquid, semi-solid and solid dosage forms, such as liquid solutions (*e.g.*, injectable and infusible solutions), dispersions or suspensions, tablets, pills, powders, liposomes and suppositories. The preferred form depends on the intended mode of administration and therapeutic application. Typical preferred compositions are in the form of injectable or infusible solutions, such as
10 compositions similar to those used for passive immunization of humans with other antibodies or other TNF α inhibitors. The preferred mode of administration is parenteral (*e.g.*, intravenous, subcutaneous, intraperitoneal, intramuscular). In one embodiment, the antibody or other TNF α inhibitor is administered by intravenous infusion or injection. In another embodiment, the antibody or other TNF α inhibitor is administered
15 by intramuscular or subcutaneous injection.

Therapeutic compositions typically must be sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, microemulsion, dispersion, liposome, or other ordered structure suitable to high drug concentration. Sterile injectable solutions can be prepared by incorporating the
20 active compound (*i.e.*, antibody, antibody portion, or other TNF α inhibitor) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated
25 above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. The proper fluidity of a solution can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required
30 particle size in the case of dispersion and by the use of surfactants. Prolonged absorption of injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.

In one embodiment, the invention includes pharmaceutical compositions comprising an effective TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) and a pharmaceutically acceptable carrier, wherein the effective TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) may
5 be used to treat hidradenitis suppurativa.

In one embodiment, the antibody or antibody portion for use in the methods of the invention is incorporated into a pharmaceutical formulation as described in PCT/IB03/04502 and U.S. Appln. No. 20040033228 .
This formulation includes a concentration 50 mg/ml of adalimumab, wherein one pre-
10 filled syringe contains 40 mg of antibody for subcutaneous injection. Alternative low-ionic formulations of adalimumab are described in US20090291062 .

The antibodies, antibody-portions, and other TNF α inhibitors of the present invention can be administered by a variety of methods known in the art, although for
15 many therapeutic applications, the preferred route/mode of administration is parenteral, *e.g.*, subcutaneous injection. In another embodiment, administration is via intravenous injection or infusion.

As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. In certain embodiments, the
20 active compound may be prepared with a carrier that will protect the compound against rapid release, such as a controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation of such
25 formulations are patented or generally known to those skilled in the art. See, *e.g.*, *Sustained and Controlled Release Drug Delivery Systems*, Robinson, ed., Dekker, Inc., New York, 1978.

In one embodiment, the TNF α antibodies and inhibitors used in the invention are delivered to a subject subcutaneously. In one embodiment, the subject administers the
30 TNF α inhibitor, including, but not limited to, a TNF α antibody, or antigen-binding portion thereof, to himself/herself.

The TNF α antibodies and inhibitors used in the invention may also be administered in the form of protein crystal formulations which include a combination of protein crystals encapsulated within a polymeric carrier to form coated particles. The coated particles of the protein crystal formulation may have a spherical morphology and
5 be microspheres of up to 500 micro meters in diameter or they may have some other morphology and be microparticulates. The enhanced concentration of protein crystals allows the antibody of the invention to be delivered subcutaneously. In one embodiment, the TNF α antibodies of the invention are delivered via a protein delivery system, wherein one or more of a protein crystal formulation or composition, is
10 administered to a subject with a TNF α -related disorder. Compositions and methods of preparing stabilized formulations of whole antibody crystals or antibody fragment crystals are also described in WO 02/072636 .

In one embodiment, a formulation comprising the crystallized antibody fragments described in PCT/IB03/04502 and U.S. Appln. No. 20040033228,
15 are used to treat rheumatoid arthritis using the treatment methods of the invention.

In certain embodiments, an antibody, antibody portion, or other TNF α inhibitor of the invention may be orally administered, for example, with an inert diluent or an assimilable edible carrier. The compound (and other ingredients, if desired) may also be
20 enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the subject's diet. For oral therapeutic administration, the compounds may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. To administer a compound of the invention by other than parenteral administration, it may be necessary
25 to coat the compound with, or co-administer the compound with, a material to prevent its inactivation.

Supplementary active compounds can also be incorporated into the compositions. In certain embodiments, an antibody or antibody portion for use in the methods of the invention is co-formulated with and/or co-administered with one or more additional
30 therapeutic agents, including an hidradenitis suppurativa inhibitor or antagonist. For example, an anti-hTNF α antibody or antibody portion of the invention may be co-formulated and/or co-administered with one or more additional antibodies that bind

other targets associated with TNF α related disorders (*e.g.*, antibodies that bind other cytokines or that bind cell surface molecules), one or more cytokines, soluble TNF α receptor (see *e.g.*, PCT Publication No. WO 94/06476) and/or one or more chemical agents that inhibit hTNF α production or activity (such as cyclohexane-ylidene
5 derivatives as described in PCT Publication No. WO 93/19751) or any combination thereof. Furthermore, one or more antibodies of the invention may be used in combination with two or more of the foregoing therapeutic agents. Such combination therapies may advantageously utilize lower dosages of the administered therapeutic agents, thus avoiding possible side effects, complications or low level of response by the
10 patient associated with the various monotherapies.

The pharmaceutical compositions of the invention may include a “therapeutically effective amount” or a “prophylactically effective amount” of an antibody or antibody portion of the invention. A “therapeutically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic
15 result. A therapeutically effective amount of the antibody, antibody portion, or other TNF α inhibitor may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the antibody, antibody portion, other TNF α inhibitor to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the antibody, antibody
20 portion, or other TNF α inhibitor are outweighed by the therapeutically beneficial effects. A “prophylactically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result. Typically, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount will be less than the therapeutically effective amount.

25 Additional description regarding methods and uses of the invention comprising administration of a TNF α inhibitor are described in Part III of this specification.

The invention also pertains to packaged pharmaceutical compositions or kits for administering the anti-TNF antibodies of the invention for the treatment of hidradenitis suppurativa. In one embodiment of the invention, the kit comprises a TNF α inhibitor,
30 such as an antibody, and instructions for administration of the TNF α inhibitor for treatment of hidradenitis suppurativa. The instructions may describe how, *e.g.*,

subcutaneously, and when, *e.g.*, at week 0, week 2, week 4, *etc.*, the different doses of TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) shall be administered to a subject for treatment.

Another aspect of the invention pertains to kits containing a pharmaceutical composition comprising a TNF α inhibitor, such as an antibody, and a pharmaceutically acceptable carrier and one or more pharmaceutical compositions each comprising an additional therapeutic agent useful for treating hidradenitis suppurativa, and a pharmaceutically acceptable carrier. Alternatively, the kit comprises a single pharmaceutical composition comprising an anti-TNF α antibody, one or more drugs useful for treating hidradenitis suppurativa, and a pharmaceutically acceptable carrier. The instructions may describe how, *e.g.*, subcutaneously, and when, *e.g.*, at week 0, week 2, week 4, *etc.*, the different doses of TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) and/or the additional therapeutic agent shall be administered to a subject for treatment.

The kit may contain instructions for dosing of the pharmaceutical compositions for the treatment of hidradenitis suppurativa. Additional description regarding articles of manufacture of the invention are described in subsection III.

The package or kit alternatively can contain the TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) and it can be promoted for use, either within the package or through accompanying information, for the uses or treatment of the disorders described herein. The packaged pharmaceuticals or kits further can include a second agent (as described herein) packaged with or co-promoted with instructions for using the second agent with a first agent (as described herein).

III. Uses and Compositions for Treating Hidradenitis Suppurativa with a TNF α Inhibitor

Hidradenitis suppurativa (HS) is a skin disorder of the apocrine glands (sweat glands found on certain parts of the body) and hair follicles in which swollen, painful, inflamed lesions or lumps develop in the groin and sometimes under the arms and under the breasts. Hidradenitis suppurativa occurs when apocrine gland outlets become

blocked by perspiration or are unable to drain normally because of incomplete gland development. Secretions trapped in the glands force perspiration and bacteria into surrounding tissue, causing subcutaneous induration, inflammation, and infection. Hidradenitis suppurativa is confined to areas of the body that contain apocrine glands.

- 5 These areas are the axillae, areola of the nipple, groin, perineum, circumanal, and periumbilical regions.

TNF α is an important cytokine in the pathogenesis of hidradenitis suppurativa, with elevated concentrations of TNF α playing a role in the hidradenitis suppurativa pathologic condition. The methods and uses described herein provide a means of
10 determining the efficacy of a TNF α inhibitor for treating hidradenitis suppurativa, and the use of such TNF α inhibitor for treating hidradenitis suppurativa. Thus in one embodiment, the invention provides a method for treating hidradenitis suppurativa in a subject having hidradenitis suppurativa.

Certain subtypes of hidradenitis suppurativa may be treated in accordance with
15 the invention. In one embodiment, moderate to severe hidradenitis suppurativa, is treated by administering a TNF α inhibitor, *e.g.*, antibody, or antigen-binding portion thereof, to a subject suffering therefrom. In one embodiment, chronic HS, *e.g.*, moderate to severe chronic HS, is treated by administering a TNF α inhibitor, *e.g.*, antibody, or antigen-binding portion thereof, to a subject suffering therefrom.

20 The invention also provides a method for treating certain subpopulations of hidradenitis suppurativa patients who may be especially difficult to treat (described in more detail below). For example, in one embodiment, the invention provides a method for treating patients who have a subtherapeutic response to a therapy, such as those who have been unresponsive or intolerant to oral antibiotics for treatment for their
25 hidradenitis suppurativa.

The invention also provides methods for improving hidradenitis suppurativa symptoms in a subject based on indices used to measure the disease state.

Treatment of HS using a TNF α inhibitor, such as a human anti-TNF α antibody or antigen binding portion thereof, may also be determined using measures known in the
30 art. Treatment of HS may be determined using any of the measures described herein,

e.g., improvement in Hurley Staging or the Sartorius scale, or any measure known to those in the art.

For example, in one embodiment, an improvement in the Hurley stage of the subject having HS, or any of the measures described herein, is evidence of effective HS treatment. In one embodiment, the severity of HS is determined according to the Hurley staging system. Hurley staging is based on assigning the subject having HS one of three different “Stages” depending on the disease level. More specifically, Stage I refers to abscess formation, single or multiple, without sinus tracts and cicatrization; Stage II refers to recurrent abscesses with tract formation and cicatrization, as well as single or multiple, widely separated lesions; and Stage III, which refers to diffuse or near-diffuse involvement, or multiple interconnected tracts and abscesses across the entire area. Hurley Stage III is the most severe form. In one embodiment, the subject having HS has HS lesions that are present in at least two distinct anatomic areas (*e.g.* left and right axilla; or left axilla and left inguinal-crural fold), one of which is at least Hurley Stage II. In another embodiment, the subject being treated has at least one lesion that is at least a Hurley Stage II.

In one embodiment, treatment of HS with a TNF α inhibitor, such as a human anti-TNF α antibody or antigen binding portion thereof, is determined by an improved Hurley score relative to a given baseline, *e.g.*, the Hurley stage of the subject prior to treatment with the TNF α inhibitor. In one embodiment, improvement in a Hurley score indicates that the Hurley score of the subject has either improved or been maintained following treatment with a TNF α inhibitor, such as a human anti-TNF α antibody or antigen binding portion thereof. Severity of hidradenitis suppurativa may be determined according to standard clinical definitions. See, for example, Hurley staging {III vs. (I or II)} for hidradenitis suppurativa (Poli F, Jemec GBE, Revuz J., Clinical Presentation. In: Jemec GBE, Revuz J, Leyden JJ, editors. Hidradenitis Suppurativa. Springer, New York, 2006, pp 11-24).

Hurley stage III disease is the most severe stage of hidradenitis suppurativa, reflecting diffuse or near-diffuse involvement of affected areas.

In one embodiment, the Sartorius scale may be used as an index for measuring efficacy of a TNF α inhibitor, *e.g.*, a human anti-TNF α antibody, for treating hidradenitis suppurativa. The Sartorius scale is described by Sartorius *et al.* in British Journal of

Dermatology, 149: 211-213 . Briefly, the following outcome variables are explicitly mentioned in reports based on the Sartorius scale: (1) anatomical region involved (axilla, groin, gluteal or other region or inframammary region left and/or right: 3 points per region involved); (2) number and scores of lesions (abscesses, nodules, fistulas, scars: points per lesion of all regions involved: nodules 2; fistulas 4; scars 1; others 1); (3) the longest distance between two relevant lesions, *i.e.*, nodules and fistulas, in each region, or size if only one lesion (< 5 cm, 2; < 10 cm, 4; > 10 cm, 8); and (4) are all lesions clearly separated by normal skin? In each region (yes 0/no 6). By assigning numerical scores to these variables, disease intensity can be quantified in a more clinically meaningful way on an open-ended scale. A total score as well as scores of selected regions chosen for surgical or other intervention can be calculated and followed over time.

In one embodiment, treatment of HS with a TNF α inhibitor, such as a human anti-TNF α antibody or antigen binding portion thereof, is determined according to an achieving an HiSCR (Hidradenitis Suppurativa Clinical Response) of the subject being treated. The HiSCR is defined as at least a 50% reduction in the total inflammatory lesion (abscess and inflammatory nodule) count (AN count) in a subject relative to baseline, with no increase in abscess count and no increase in draining fistula count. In one embodiment, treatment of HS in a subject is defined as an at least 50% reduction in the inflammatory lesion (abscess and nodule) count.

The HiSCR scoring system was designed to assess hidradenitis suppurativa activity in an affected subject before and after a treatment. It is described in more detail in Example 5 below.

In another embodiment, treatment of HS with a TNF α inhibitor, such as a human anti-TNF α antibody or antigen binding portion thereof, is defined as achieving an Hidradenitis Suppurativa Physician's Global Assessment (HS-PGA) score, or HS-PGA score, as defined below, of clear (0), minimal (1), or mild (2), with an improvement (*i.e.*, reduction) from baseline HS-PGA score of at least 2 grades, optionally, at the end of a treatment period (such as week 16). The baseline HS-PGA score is the HS-PGA score measured just prior to the commencement of treatment, to which the HS-PGA score obtained after a period of treatment is compared. Both the baseline HS-PGA score and

the HS-PGA score obtained after a treatment period are assessed based on the following system and criteria:

HS-PGA Scoring System

Score	Rating	Description
0	Clear	No abscesses, no draining fistulas, no nodules
1	Minimal	No abscesses, no draining fistulas, no inflammatory nodules, presence of non-inflammatory nodules
2	Mild	No abscesses or draining fistulas, and less than 5 inflammatory nodules, or Single abscess or draining fistula, and no inflammatory nodules
3	Moderate	No abscesses or draining fistulas, and at least 5 inflammatory nodules, or Single abscess or draining fistula in the presence of inflammatory nodules, or Between 2 and 5 abscesses or draining fistulas with or without inflammatory nodules, up to 10
4	Severe	Between 2 and 5 abscesses and draining fistulas with or without inflammatory nodules that are greater than 10
5	Very severe	More than 5 abscesses or draining fistulas

5 The HS-PGA scoring system was designed for use in assessing hidradenitis suppurativa activity before and after a treatment. It is a six-point score that partly depends on the presence / absence of abscesses, draining fistulas, and/or nodules (inflammatory or non-inflammatory), and, if present, the extent of such presence.

10 The invention also includes a method of decreasing an HS-PGA score of a subject comprising administering a human TNF α antibody, or antigen-binding portion thereof, to the subject, such that partial remission of hidradenitis suppurativa is induced. In one embodiment, the invention provides an improvement of at least about 2 grades in the HS-PGA score of a subject having hidradenitis suppurativa.

15 In one embodiment, the invention provides a method for improving the DLQI score of a subject. In one embodiment, the improvement in the DLQI score is determined by achieving a score, *e.g.*, a statistically significant score, correlating with a “no” or “small impact” of the disease state on the subject. In one embodiment, the

improvement in the DLQI score is determined by achieving an improvement in the DLQI score of the subject. Examples of such improvements are provided in the examples described herein.

Methods of treatment described herein may include administration of a

5 TNF α inhibitor, *e.g.*, antibody, or antigen-binding portion thereof, to a subject to achieve a therapeutic goal, *e.g.*, treatment of hidradenitis suppurativa, or achievement of a clinical response as defined herein. Also included in the scope of the invention are uses of a TNF α inhibitor, *e.g.*, antibody, or antigen-binding portion thereof, in the manufacture of a medicament to achieve a therapeutic goal, *e.g.*, treatment, of

10 hidradenitis suppurativa, increase in clinical response as defined herein, and/or maintenance or improvement (reduction) of a HiSCR. Thus, where methods are described herein, it is also intended to be part of this invention that the use of the TNF α inhibitor in the manufacture of a medicament for the purpose of the method is also considered within the scope of the invention. Likewise, where a use of a

15 TNF α inhibitor, *e.g.*, antibody, or antigen-binding portion thereof, in the manufacture of a medicament for the purpose of achieving a therapeutic goal is described, methods of treatment resulting in the therapeutic goal are also intended to be part of the invention.

In one embodiment, the invention provides a method for treating a subject having hidradenitis suppurativa (HS) comprising administering an isolated human anti-TNF α

20 antibody, or an antigen binding portion thereof, to the subject according to a multiple variable dose regimen, such that HS is treated, wherein the multiple variable dose regimen comprises administering a first loading dose, administering a second loading dose which is less than the first loading dose, and administering a treatment dose which is less than the second loading dose, wherein the treatment dose is administered to the

25 subject weekly.

In one embodiment, the invention provides a method for treating a subject having hidradenitis suppurativa (HS) comprising administering an isolated human anti-TNF α antibody, or an antigen binding portion thereof, to the subject according to a multiple variable dose regimen, such that HS is treated, wherein the multiple variable dose

30 regimen comprises administering a first loading dose, administering a second loading dose which is less than the first loading dose, and administering a treatment dose which

is less than the second loading dose, wherein the treatment dose is administered to the subject biweekly.

In one embodiment, the invention provides a method for decreasing the number of inflammatory lesions (AN count) in a subject having HS, said method comprising
5 systemically administering an isolated human anti-TNF α antibody, or an antigen binding portion thereof, to the subject, such that the AN count is decreased. The decrease in AN count may be anything greater than 10%, *e.g.*, the AN count may be reduced by at least a 50% reduction in the subject relative to baseline AN count. The subject may also exhibit other improvements in HS following treatment with a TNF α inhibitor, *e.g.*, anti-TNF α
10 antibody. For example the subject may have no increase in an abscess count and/or no increase in a draining fistula count following administration with the anti-TNF α antibody, or an antigen binding portion thereof.

In one embodiment, treatment of hidradenitis suppurativa is achieved by administering a human TNF α antibody, or an antigen-binding portion thereof, to a
15 subject having hidradenitis suppurativa, wherein the human TNF α antibody, or an antigen-binding portion thereof, is administered on a weekly or biweekly dosing regimen, or any combination thereof. Biweekly dosing regimens can be used to treat disorders in which TNF α activity is detrimental, and are further described in US Appln. No. 10/163657 (US 20030235585).

In one
20 embodiment, biweekly dosing includes a dosing regimen wherein doses of a TNF α inhibitor are administered to a subject every other week, *e.g.*, beginning at week 1. In one embodiment, biweekly dosing includes a dosing regimen where doses of a TNF α inhibitor are administered to a subject every other week consecutively for a given time period, *e.g.*, 4 weeks, 8 weeks, 16, weeks, 24 weeks, 26 weeks, 32 weeks, 36
25 weeks, 42 weeks, 48 weeks, 52 weeks, 56 weeks or more, *etc.* Biweekly or weekly dosing is preferably administered parenterally, including subcutaneously. In one embodiment, the human TNF α antibody, or an antigen-binding portion thereof, is administered in a dose of about 40 mg for each weekly or biweekly dosing. In one embodiment, the human TNF α antibody, or an antigen-binding portion thereof, is
30 adalimumab. Additional examples of dosing regimens within the scope of the invention are provided herein in the Examples.

In one embodiment, treatment of hidradenitis suppurativa is achieved using multiple variable dosing methods of treatment. Examples of such multiple variable dosing regimens are described in PCT appln. no. PCT/US05/12007 .

For example, a loading dose of about 160 and/or 80 mg of a
5 TNF α inhibitor, *e.g.*, TNF α antibody, or an antigen-binding portion thereof, may first be administered to a subject having hidradenitis suppurativa, followed by a maintenance or treatment dose of about 40 mg.

In one embodiment, the invention provides a method of treating hidradenitis suppurativa in a subject comprising administering a loading dose(s) of a TNF α inhibitor,
10 *e.g.*, TNF α antibody, or an antigen-binding portion thereof, to the subject at week 0, optionally another loading dose at week 2. In one embodiment, the loading dose(s) is given in its entirety on one day or is divided over multiple days (*e.g.*, divided over two days). In one embodiment, the loading dose(s) is administered subcutaneously. Following administration of the loading dose(s), one or more maintenance or treatment
15 dose(s) of the TNF α inhibitor, *e.g.*, TNF α antibody, or an antigen-binding portion thereof, may be administered to the subject, wherein the maintenance or treatment dose is about half or 1/4 of the dose amount of the loading dose(s). In one embodiment, the maintenance or treatment dose is administered to the subject about one or two weeks after the last loading dose(s). In one embodiment, the maintenance or treatment dose is
20 administered subcutaneously. Subsequent doses may be administered following the same or different maintenance or treatment dosing regimen.

In one embodiment, two loading doses are administered to the subject. In one embodiment, the second loading dose is about 40-60% of the first or initial loading dose. In one embodiment, the treatment dose is about 40-60% of the second loading
25 dose.

In one embodiment, the first loading dose is about 140-180 mg. Numbers intermediate to the stated range are also included in the invention, *e.g.*, 145-175 mg, 150-170 mg, and 155-165 mg. In one embodiment, the first loading dose is about 160 mg.

In one embodiment, the second loading dose is about 60-100 mg. Numbers intermediate to the stated range are also included in the invention, *e.g.*, 70-90 mg, 75-85 mg, and 65-95 mg. In one embodiment, the second loading dose is about 80 mg.

In one embodiment, the treatment dose is about 30-50 mg. Numbers
5 intermediate to the stated range are also included in the invention, *e.g.*, 35-45 mg. In one embodiment, the treatment dose is about 40 mg.

In another embodiment, the loading dose(s) of the human TNF α antibody, or antigen-binding portion thereof, comprises about 80 mg, and may be given at week 0, followed by at least one maintenance dose of the human TNF α antibody, or antigen-
10 binding portion thereof, comprising about 40 mg, administered on a biweekly dosing regimen, optionally from week 1. Alternatively, in another embodiment, the loading dose(s) of the human TNF α antibody, or antigen-binding portion thereof, comprises a first dose of about 160 mg administered on week 0, and a second loading dose of about 80 mg administered on week 2, followed by at least one maintenance dose of the human
15 TNF α antibody, or antigen-binding portion thereof, comprising about 40 mg, administered weekly thereafter. In one embodiment, the maintenance dose is administered to the subject starting at about week 4 (wherein week 0 is the initial loading dose).

In one embodiment, the subject is first selected for having HS and is then
20 administered a TNF α inhibitor, *e.g.*, TNF α antibody, or an antigen-binding portion thereof, in accordance with the methods described herein.

Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic
25 effect in association with the required pharmaceutical carrier. Such dosage unit forms may be a tablet or pill with a pre-determined amount of therapeutic agents, or a vial with therapeutic agents to be reconstituted by a solution to produce a drug product of a pre-determined final concentration. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the
30 active compound and the particular therapeutic or prophylactic effect to be achieved, and

(b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

Dosage regimens described herein may be adjusted (*e.g.*, in individual patients) to provide the optimum desired response, *e.g.*, maintaining remission of hidradenitis
5 suppurativa, in consideration of the teachings herein.

It is to be noted that dosage values can vary with the type and severity of hidradenitis suppurativa. It is to be further understood that for any particular subject, specific dosage regimens may be adjusted over time according to the teachings of the specification and the individual need and the professional judgment of the person
10 administering or supervising the administration of the compositions, and that dosage amounts and ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed invention.

Subpopulations

15 The invention provides uses and methods for treating certain subpopulations of hidradenitis suppurativa patients with a TNF α inhibitor, *e.g.*, TNF α antibody, or an antigen-binding portion thereof.

In one embodiment, the subject has HS lesions in at least two distinct anatomic areas prior to treatment.

20 In one embodiment, the subject had an inadequate response to or was intolerant to oral antibiotics for treatment of their HS.

In one embodiment, the subject has an AN count of greater than or equal to 3 at baseline, a female, a subject who is over 40 years old, a subject who is a smoker, or any combination thereof.

25 In one embodiment, the invention provides a method of treating moderate to severe hidradenitis suppurativa in a subject comprising administering to the subject a TNF α inhibitor, *e.g.*, TNF α antibody, or an antigen-binding portion thereof, such that moderate to severe hidradenitis suppurativa is treated. Subjects having moderate to severe hidradenitis suppurativa may be administered a TNF α inhibitor, *e.g.*, TNF α
30 antibody, or an antigen-binding portion thereof, such that moderate to severe

hidradenitis suppurativa is treated and advancement of the disease is prevented. The invention also provides use of a TNF α inhibitor, *e.g.*, TNF α antibody, or an antigen-binding portion thereof, in the manufacture of a medicament for the treatment of moderate to severe hidradenitis suppurativa in a subject who has moderate to severe

5 hidradenitis suppurativa. In one embodiment, a patient having moderate to severe hidradenitis suppurativa is defined as a patient having a HS-PGA score no less than 3. In one embodiment, such patients have been unresponsive or intolerant to oral antibiotics for treatment for their hidradenitis suppurativa. In one embodiment, such patients have had a diagnosis of moderate to severe hidradenitis suppurativa for at least 6 months prior

10 to Baseline HS-PGA measurement, and involve at least two distinct anatomic areas (*e.g.* left and right axilla; or left axilla and left inguinal-crural fold).

In one embodiment, the invention provides an article of manufacture comprising adalimumab and a package insert, wherein the package insert indicates that adalimumab may be used to treat hidradenitis suppurativa in patients who have a HS-PGA score no

15 less than 3, who have been unresponsive or intolerant to oral antibiotics for treatment for their hidradenitis suppurativa, and/or who have had a diagnosis of moderate to severe hidradenitis suppurativa for at least 6 months prior to Baseline HS-PGA measurement, and involve at least two distinct anatomic areas (*e.g.* left and right axilla and left inguinal-crural fold).

20

Articles of Manufacture

The invention also provides a packaged pharmaceutical composition wherein the TNF α inhibitor, *e.g.*, human TNF α antibody, is packaged within a kit or an article of manufacture. The kit or article of manufacture of the invention contains materials useful

25 for the treatment, including induction and/or remission, prevention and/or diagnosis of hidradenitis suppurativa. The kit or article of manufacture comprises a container and a label or package insert or printed material on or associated with the container which provides information regarding use of the TNF α inhibitor, *e.g.*, a TNF α antibody, for the treatment of hidradenitis suppurativa.

30 A kit or an article of manufacture refers to a packaged product comprising components with which to administer a TNF α inhibitor for treatment of a hidradenitis

suppurativa. The kit preferably comprises a box or container that holds the components of the kit. The box or container is affixed with a label or a Food and Drug Administration approved label, including a protocol for administering the TNF α inhibitor. The box or container holds components of the invention which are preferably
5 contained within plastic, polyethylene, polypropylene, ethylene, or propylene vessels. The vessels can be capped-tubes or bottles. The kit can also include instructions for administering the TNF α antibody of the invention. In one embodiment the kit of the invention includes the formulation comprising the human antibody adalimumab (or D2E7), as described in PCT/IB03/04502 and U.S. Appln. No. 10/222140.

10

The term “package insert” is used to refer to instructions customarily included in commercial packages of therapeutic products, that contain information about the indications, usage, dosage, administration, contraindications and/or warnings concerning the use of such therapeutic products.

15 In one embodiment, the article of manufacture of the invention comprises (a) a first container with a composition contained therein, wherein the composition comprises a TNF α antibody; and (b) a package insert indicating that the TNF α antibody may be used for reducing signs and symptoms and inducing and maintaining remission of hidradenitis suppurativa. In a preferred embodiment, the label or package insert
20 indicates that the TNF α inhibitor, *e.g.*, a TNF α antibody, is used for treating hidradenitis suppurativa.

In one embodiment, the invention features a kit comprising a sufficient number of containers to provide both loading and treatment doses of the TNF α inhibitor, *e.g.*, anti-TNF α antibody. For example, the kit may contain at least seven containers
25 containing about 40 mg of an isolated human anti-TNF α antibody, or an antigen binding portion thereof. Seven containers each containing 40 mg of an anti-TNF α antibody could, for example, provide enough antibody for a loading dose of about 160 mg (4 x 40 mg), a loading dose of about 80 mg (2 x 40 mg), and one treatment dose of about 40 mg.

Suitable containers for the TNF α inhibitor, *e.g.*, a TNF α antibody, include, for
30 example, bottles, vials, syringes, including preloaded syringes, pens, including autoinjector pens, *etc.* The containers may be formed from a variety of materials such as

glass or plastic. The container holds a composition which is by itself or when combined with another composition effective for treating, preventing and/or diagnosing the condition and may have a sterile access port.

In one embodiment, the article of manufacture comprises a TNF α inhibitor, *e.g.*,
5 a human TNF α antibody, and a label or package insert which indicates to a subject who will be administering the TNF α inhibitor about using the TNF α inhibitor for the treatment of hidradenitis suppurativa. The label may be anywhere within or on the article of manufacture. In one embodiment, the article of manufacture comprises a container, such as a box, which comprises the TNF α inhibitor and a package insert or
10 label providing information pertaining to use of the TNF α inhibitor for the treatment of hidradenitis suppurativa. In another embodiment, the information is printed on a label which is on the outside of the article of manufacture, in a position which is visible to prospective purchasers.

In one embodiment, the label or package insert of the invention informs a reader,
15 including a subject, *e.g.*, a purchaser, who will be administering the TNF α inhibitor for treatment, that the TNF α inhibitor, *e.g.*, a TNF α antibody such as adalimumab, is an indicated treatment of hidradenitis suppurativa, including of moderately to severely active disease in adult patients.

In one embodiment, the label or package insert describes certain patient
20 populations who may respond favorably to the TNF α inhibitor within the article of manufacture. For example, the label or package insert may indicate that the TNF α antibody, *e.g.*, adalimumab, may be used to treat hidradenitis suppurativa in patients who have been unresponsive or intolerant to oral antibiotics for treatment for their hidradenitis suppurativa.

25 In one embodiment, the label or package insert of the invention describes certain therapeutic benefits of the TNF α antibody, *e.g.*, adalimumab, including specific symptoms of hidradenitis suppurativa which may be reduced by using the TNF α antibody, *e.g.*, adalimumab. It should be noted that the package insert may also contain information pertaining to other disorders which are treatable using the
30 TNF α antibody, *e.g.*, adalimumab. Information described herein which is provided in a label or package insert and pertains to other disorders, *i.e.*, diseases other than

hidradenitis suppurativa, is also included within the scope of the invention. The package insert of the invention may indicate that extra TNF α in the body can attack normal healthy body tissues and cause inflammation especially in the tissues in your bones, cartilage, joints and digestive tract. The package insert of the invention may also
5 indicate that adalimumab helps reduce the signs and symptoms of immune diseases, including rheumatoid and psoriatic arthritis (pain and swollen joints), ankylosing spondylitis (morning stiffness and back pain), and psoriasis (abdominal pain and diarrhea).

In another embodiment, the package insert of the invention describes the dose
10 and administration of adalimumab, for the treatment of hidradenitis suppurativa. The label may indicate that the initiation of therapy includes a loading dose of about 160 mg and 80 mg administered at weeks 0 and 2, respectively. The label may also indicate that the maintenance dosing for the treatment of hidradenitis suppurativa with adalimumab is about 40 mg every week thereafter, such as starting from week 4. Alternatively, the
15 label may indicate that the initiation of therapy includes a loading dose of about 80 mg administered at week 0, followed by maintenance doses of about 40 mg every other week thereafter, such as starting from week 1. Regardless of the initial treatment dosing regimen, the label may also indicate that additional maintenance doses are administered at about 40 mg every other week. In another embodiment, the label or package insert of
20 the invention indicates that the TNF α inhibitor (*e.g.*, adalimumab) is administered by subcutaneous injection.

The label or the package insert of the invention may also provide information to subjects who will be receiving adalimumab regarding combination uses with other hidradenitis suppurativa therapeutic agents for both safety and efficacy purposes.

25 The label or the package insert of the invention may contain warnings and precautions regarding the use of the TNF α inhibitor, *e.g.*, a TNF α antibody such as adalimumab. In one embodiment, the information provided in the label or the package insert describes super infection of hidradenitis suppurativa lesions and/or pilonidal cyst flare.

30 The label or the package insert of the invention may contain information regarding the use of the TNF α inhibitor, *e.g.*, a TNF α antibody such as adalimumab, in

clinical studies for hidradenitis suppurativa. In one embodiment, the label of the invention describes the studies described herein as Examples 1 to 2, either as a whole or in portion. The label of the invention may also indicate that the safety profile for patients with hidradenitis suppurativa treated with HUMIRA[®] was similar to the safety
5 profile seen in patients with other indications treatable by HUMIRA[®], such as rheumatoid arthritis.

The label of the invention may contain information regarding the pharmacodynamics of the TNF α inhibitor, *e.g.*, a TNF α antibody such as adalimumab.

In one embodiment of the invention, the kit comprises a TNF α inhibitor, such as
10 an antibody, an second pharmaceutical composition comprising an additional therapeutic agent, and instructions for administration of both agents for the treatment of hidradenitis suppurativa. The instructions may describe how, *e.g.*, subcutaneously, and when, *e.g.*, at week 0, week 2, and weekly or biweekly thereafter, doses of TNF α antibody and/or the additional therapeutic agent shall be administered to a subject for treatment.

15 Another aspect of the invention pertains to kits containing a pharmaceutical composition comprising an anti-TNF α antibody and a pharmaceutically acceptable carrier and one or more additional pharmaceutical compositions each comprising a drug useful for treating a TNF α related disorder (such as hidradenitis suppurativa) and a pharmaceutically acceptable carrier. Alternatively, the kit comprises a single
20 pharmaceutical composition comprising an anti-TNF α antibody, one or more drugs useful for treating a TNF α related disorder (such as hidradenitis suppurativa) and a pharmaceutically acceptable carrier. The kits further contain instructions for dosing of the pharmaceutical compositions for the treatment of a TNF α related disorder (such as hidradenitis suppurativa).

25 The package or kit alternatively may contain the TNF α inhibitor and it may be promoted for use, either within the package or through accompanying information, for the uses or treatment of hidradenitis suppurativa. The packaged pharmaceuticals or kits further can include a second agent (as described herein) packaged with or co-promoted with instructions for using the second agent with a first agent (as described herein).

Additional therapeutic agents

TNF α inhibitors, including TNF α antibodies, or antigen binding portions thereof, may be used in the methods, uses, and compositions of the invention either alone or in combination with an additional therapeutic agent. It should be understood that the

5 TNF α inhibitors can be used alone or in combination with an additional agent, *e.g.*, a therapeutic agent, said additional agent being selected by the skilled artisan for its intended purpose. For example, the additional agent can be a therapeutic agent art-recognized as being useful to treat the disease or condition being treated by the TNF α inhibitors. The additional agent also can be an agent that imparts a beneficial attribute to

10 the therapeutic composition, *e.g.*, an agent which effects the viscosity of the composition.

It should further be understood that the combinations which are to be included within this invention are those combinations useful for their intended purpose. The agents set forth below are illustrative for purposes and not intended to be limited. The

15 combinations, which are part of this invention, can be the TNF α inhibitors of the present invention and at least one additional agent selected from the lists below. The combination can also include more than one additional agent, *e.g.*, two or three additional agents if the combination is such that the formed composition can perform its intended function.

20 TNF α inhibitors described herein may be used in combination with additional therapeutic agents such as a Disease Modifying Anti-Rheumatic Drug (DMARD) or a Non-steroidal Anti-inflammatory Drug (NSAID) or a steroid or any combination thereof. Preferred examples of a DARED are hydroxychloroquine, leflunomide, methotrexate, parenteral gold, oral gold and sulfasalazine. Preferred examples of non-steroidal anti-

25 inflammatory drug(s) also referred to as NSAIDS include drugs like ibuprofen. Other preferred combinations are corticosteroids including prednisolone; the well known side effects of steroid use can be reduced or even eliminated by tapering the steroid dose required when treating patients in combination with TNF α inhibitors of this invention.

Preferred combinations of therapeutic agents may interfere at different points in

30 the autoimmune and subsequent inflammatory cascade; preferred examples include TNF antagonists such as soluble p55 or p75 TNF receptors, derivatives, thereof, (p75

TNFRlgG (ENBRELTM) or p55 TNFRlgG (Lenercept), chimeric, humanized or human TNF antibodies, or a fragment thereof, including infliximab (REMICADE[®], Johnson and Johnson; described in U.S. Patent No. 5,656,272),
 PSORIASIS P571 (a humanized monoclonal anti-TNF-alpha IgG4 antibody),
 5 PSORIASIS P 870 (a humanized monoclonal anti-TNF-alpha antibody fragment), an anti-TNF dAb (Peptech), SIMPONI[®] (golimumab; also referred to as CNTO 148; Centocor Ortho Biotech, see WO 02/12502), and adalimumab (HUMIRA[®], Abbott Laboratories, a human anti-TNF mAb, described in US 6,090,382 as D2E7). Additional TNF antibodies which can be used in the invention are described in U.S. Patent Nos.
 10 6,593,458; 6,498,237; 6,451,983; and 6,448,380 .

Other combinations including TNF α converting enzyme (TACE) inhibitors; IL-1 inhibitors (Interleukin-1-converting enzyme inhibitors, IL-1RA *etc.*) may be effective for the same reason. Other preferred combinations include Interleukin 11. Yet another preferred combination are other key players of the autoimmune response
 15 which may act parallel to, dependent on or in concert with TNF α inhibitors function; especially preferred are IL-18 antagonists including IL-18 antibodies or soluble IL-18 receptors, or IL-18 binding proteins. Yet another preferred combination are non-depleting anti-PSORIASIS 4 inhibitors. Yet other preferred combinations include antagonists of the co-stimulatory pathway PSORIASIS 80 (B7.1) or PSORIASIS 86
 20 (B7.2) including antibodies, soluble receptors or antagonistic ligands.

The TNF α inhibitors used in the invention may also be combined with agents, such as methotrexate, 6-MP, azathioprine sulphasalazine, mesalazine, olsalazine chloroquine/hydroxychloroquine, pencillamine, aurothiomalate (intramuscular and oral), azathioprine, cochlincine, corticosteroids (oral, inhaled and local injection), beta-2
 25 adrenoreceptor agonists (salbutamol, terbutaline, salmeteral), xanthines (theophylline, aminophylline), cromoglycate, nedocromil, ketotifen, ipratropium and oxitropium, cyclosporin, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs, for example, ibuprofen, corticosteroids such as prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement inhibitors, adrenergic agents,
 30 agents which interfere with signaling by proinflammatory cytokines such as TNF α or IL-1 (*e.g.* IRAK, NIK, IKK , p38 or MAP kinase inhibitors), IL-1 β converting enzyme inhibitors, TNF α converting enzyme (TACE) inhibitors, T-cell signaling inhibitors such

as kinase inhibitors, metalloproteinase inhibitors, sulfasalazine, azathioprine, 6-mercaptapurines, angiotensin converting enzyme inhibitors, soluble cytokine receptors and derivatives thereof (*e.g.* soluble p55 or p75 TNF receptors and the derivatives p75 TNFRIgG (EnbrelTM and p55 TNFRIgG (Lenercept)), sIL-1RI, sIL-1RII, sIL-6R), anti-
 5 inflammatory cytokines (*e.g.* IL-4, IL-10, IL-12, IL-13 and TGF β), celecoxib, folic acid, hydroxychloroquine sulfate, rofecoxib, etanercept, infliximab, naproxen, valdecoxib, sulfasalazine, methylprednisolone, meloxicam, methylprednisolone acetate, gold sodium thiomalate, aspirin, triamcinolone acetonide, propoxyphene napsylate/apap, folate, nabumetone, diclofenac, piroxicam, etodolac, diclofenac sodium, oxaprozin, oxycodone
 10 hcl, hydrocodone bitartrate/apap, diclofenac sodium/misoprostol, fentanyl, anakinra, human recombinant, tramadol hcl, salsalate, sulindac, cyanocobalamin/fa/pyridoxine, acetaminophen, alendronate sodium, prednisolone, morphine sulfate, lidocaine hydrochloride, indomethacin, glucosamine sulf/chondroitin, amitriptyline hcl, sulfadiazine, oxycodone hcl/acetaminophen, olopatadine hcl, misoprostol, naproxen
 15 sodium, omeprazole, cyclophosphamide, rituximab, IL-1 TRAP, MRA, CTLA4-IG, IL-18 BP, anti-IL-18, Anti-IL15, BIRB-796, SCIO-469, VX-702, AMG-548, VX-740, Roflumilast, IC-485, PSORIASIS C-801, and Mesopram.

Non-limiting examples of therapeutic agents for hidradenitis suppurativa with which TNF α inhibitor of the invention can be combined include the following: antiseptic
 20 and antiperspirant agents (*e.g.*, 6.25% aluminum chloride hexahydrate in absolute ethanol), anti-inflammatory or anti-androgen therapy such as tetracycline, intralesional triamcinolone, or finasteride.

The antibodies of the invention, or antigen binding portions thereof, may also be combined with agents, such as methotrexate, cyclosporin, FK506, rapamycin,
 25 mycophenolate mofetil, leflunomide, NSAIDs, for example, ibuprofen, corticosteroids such as prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement inhibitors, adrenergic agents, agents which interfere with signaling by proinflammatory cytokines such as TNF α or IL-1 (*e.g.*, IRAK, NIK, IKK, p38 or MAP kinase inhibitors), IL-1 β converting enzyme inhibitors, TNF α converting enzyme
 30 inhibitors, T-cell signaling inhibitors such as kinase inhibitors, metalloproteinase inhibitors, sulfasalazine, azathioprine, 6-mercaptapurines, angiotensin converting enzyme inhibitors, soluble cytokine receptors and derivatives thereof (*e.g.*, soluble p55

or p75 TNF receptors, sIL-1RI, sIL-1RII, sIL-6R) and anti-inflammatory cytokines (*e.g.*, IL-4, IL-10, IL-12, IL-13 and TGF β).

Additional examples of therapeutic agents for hidradenitis suppurativa in which a TNF α inhibitor can be combined include the following: combinations of TNF antagonists, for example, anti-TNF antibodies, D2E7 (PCT Publication No. WO 97/29131; HUMIRA[®]), Ca2 (REMICADE[®]), PSORIASIS P 571, TNFR-Ig constructs, (p75 TNFRIgG (ENBREL[™]) and p55 TNFRIgG (LENERCEPT) inhibitors and PDE4 inhibitors. TNF α inhibitors of the invention can be combined with corticosteroids, for example, budenoside and dexamethasone. TNF α inhibitors of the invention may also be combined with agents such as sulfasalazine, 5-aminosalicylic acid and olsalazine, and agents which interfere with synthesis or action of proinflammatory cytokines such as IL-1, for example, IL-1 β converting enzyme inhibitors and IL-1ra. TNF α inhibitors may also be used with T cell signaling inhibitors, for example, tyrosine kinase inhibitors 6-mercaptopurines. TNF α inhibitors can be combined with IL-12. TNF α inhibitors can be combined with mesalamine, prednisone, azathioprine, mercaptopurine, infliximab, methylprednisolone sodium succinate, diphenoxylate/atrop sulfate, loperamide hydrochloride, methotrexate, omeprazole, folate, ciprofloxacin/dextrose-water, hydrocodone bitartrate/apap, tetracycline hydrochloride, fluocinonide, metronidazole, thimerosal/boric acid, cholestyramine/sucrose, ciprofloxacin hydrochloride, hyoscyamine sulfate, meperidine hydrochloride, midazolam hydrochloride, oxycodone hcl/acetaminophen, promethazine hydrochloride, sodium phosphate, sulfamethoxazole / trimethoprim, celecoxib, polycarbophil, propoxyphene napsylate, hydrocortisone, multivitamins, balsalazide disodium, codeine phosphate/apap, colesevelam hcl, cyanocobalamin, folic acid, levofloxacin, methylprednisolone, natalizumab and interferon-gamma.

The TNF α inhibitors may also be combined with agents, such as alemtuzumab, dronabinol, Unimed, daclizumab, mitoxantrone, xaliproden hydrochloride, fampridine, glatiramer acetate, natalizumab, sinnabidol, a-immunokine NNSO3, ABR-215062, AnergiX.MS, chemokine receptor antagonists, BBR-2778, calagualine, CPI-1189, LEM (liposome encapsulated mitoxantrone), THC.CBD (cannabinoid agonist) MBP-8298, mesopram (PDE4 inhibitor), MNA-715, anti-IL-6 receptor antibody, neurovax, pirfenidone allotrap 1258 (RDP-1258), sTNF-R1, talampanel, teriflunomide, TGF-beta2,

tiplimotide, VLA-4 antagonists (for example, TR-14035, VLA4 Ultrahaler, Antegran-ELAN/Biogen), interferon gamma antagonists, IL-4 agonists, and the humanized IL-6 antibody tocilizumab.

In yet another embodiment, the invention includes an article of manufacture or a
5 method comprising the combination of a TNF α inhibitor and an antibiotic or anti-infective agent. Anti-infective agents include those agents known in the art to treat viral, fungal, parasitic or bacterial infections. The term, "antibiotic," as used herein, refers to a chemical substance that inhibits the growth of, or kills, microorganisms. Encompassed
10 by this term are antibiotic produced by a microorganism, as well as synthetic antibiotics (*e.g.*, analogs) known in the art. Antibiotics include, but are not limited to, clarithromycin (BIAXIN[®]), ciprofloxacin (CIPRO[®]), and metronidazole (FLAGYL[®]).

Any one of the above-mentioned therapeutic agents, alone or in combination therewith, can be administered to a subject suffering from hidradenitis suppurativa, in combination with the TNF α antibody using a multiple variable dose treatment regimen.
15 In one embodiment, any one of the above-mentioned therapeutic agents, alone or in combination therewith, can be administered to a subject suffering from hidradenitis suppurativa in addition to a TNF α antibody to treat another TNF α -related disorder, such as rheumatoid arthritis. It should be understood that the additional therapeutic agents can be used in combination therapy as described above, but also may be used in other
20 indications described herein wherein a beneficial effect is desired.

The combination of agents used within the methods and pharmaceutical compositions described herein may have a therapeutic additive or synergistic effect on the condition(s) or disease(s) targeted for treatment. The combination of agents used within the methods or pharmaceutical compositions described herein also may reduce a
25 detrimental effect associated with at least one of the agents when administered alone or without the other agent(s) of the particular pharmaceutical composition. For example, the toxicity of side effects of one agent may be attenuated by another agent of the composition, thus allowing a higher dosage, improving patient compliance, and improving therapeutic outcome. The additive or synergistic effects, benefits, and
30 advantages of the compositions apply to classes of therapeutic agents, either structural or functional classes, or to individual compounds themselves.

IV. *Efficacy of TNF α Inhibitor*

The invention also provides methods for determining whether a TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) is effective at
5 treating hidradenitis suppurativa in a subject. Such methods may be used to determine the efficacy of a TNF α inhibitor, including those which are unknown or unconfirmed to have such efficacy. Using the methods described herein, effective TNF α inhibitors (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) may be determined or confirmed, and, subsequently, used in the method of treating hidradenitis suppurativa.

10 The efficacy of a TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) for treatment of hidradenitis suppurativa in a patient or patient population who has hidradenitis suppurativa, may be evaluated by determining the percentage of the patient population in whom a clinical response has been achieved following administration of the TNF α inhibitor (*e.g.*, a test or candidate TNF α
15 inhibitor).

In one embodiment, the invention provides a method for determining the efficacy of a TNF α inhibitor, including a human TNF α antibody, for treating hidradenitis suppurativa in a subject, using the HS-PGA score to assess the proportion of treated subjects who have achieved clinical response defined above. Alternatively, or in
20 addition, the HiSCR scoring system, the DLQI, the Sartorius scale may also be used to determine efficacy.

In certain embodiments, a candidate TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) is efficacious for treating hidradenitis suppurativa if at least about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22,
25 23, 24, 25% or more treated patients achieve statistically significant (as compared to placebo treatment) clinical response, as defined herein based on HS-PGA score. Ranges of values using a combination of any of the above recited values as upper and/or lower limits are intended to be included in the scope of the invention.

The DLQI is an additional validated instrument used to assess dermatologic-
30 related functional limitations. Characteristics of the DLQI include: (1) ten items on an

overall scoring range of 0-30; higher scores represent greater quality of life impairment and lower scores represent lower quality of life impairment; (2) well-established properties of reliability and validity for the DLQI total score in a dermatology setting (see Badia *et al.* (1999) *Br J Dermatol* 141:698; Finlay *et al.* (1994) *Clin Exp Dermatol* 5 19:210; and Shikiar *et al.* (2003) *Health and Quality of Life Outcomes* 1:53); (3) six subcategories: symptoms and feelings; daily activities; leisure; work/school; personal relationships; and treatment; and, (4) all data are observed values. Patients who discontinued before the time point were not included in this analysis.

Ranges of DLQI scores can be evaluated for their correspondence to categories of
10 disease impact.

In certain embodiments, the DLQI score may also be used as an index for measuring efficacy of a TNF α inhibitor in a patient or patient population having hidradenitis suppurativa, where mean improvement within a population of treated patients in their DLQI scores that is statistically significant (as compared to placebo) indicates that the
15 TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) is effective for treating hidradenitis suppurativa. In one embodiment, the invention provides a method for determining whether a human TNF α antibody is effective for treating hidradenitis suppurativa.

In certain embodiments, the Pain VAS score may also be used as an index for
20 measuring efficacy of a TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) in a patient or patient population having hidradenitis suppurativa, where mean statistically significant (as compared to placebo) improvement within a population of treated patients in their Pain VAS scores that is at least 30% indicates that the TNF α inhibitor is effective for treating hidradenitis suppurativa. In
25 one embodiment, the invention provides a method for determining whether a human TNF α antibody is effective for treating hidradenitis suppurativa based on improvement in Pain VAS score.

In one embodiment, the invention provides a method of treating hidradenitis suppurativa in a subject, comprising administering an effective amount of a
30 TNF α inhibitor, *e.g.*, a human TNF α antibody, to the subject such that hidradenitis suppurativa is treated, wherein the effective TNF α inhibitor, *e.g.*, human

TNF α antibody, is previously identified as achieving a statistically significant clinical response within a patient or patient population.

In one embodiment, at least about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25% or more of the treated patients achieve a clinical response as
5 defined herein.

Time points for determining efficacy will be understood by those of skill in the art to depend on the type of efficacy being determined, *e.g.*, treatment of hidradenitis suppurativa. In one embodiment, measurements in scores, *e.g.*, the HS-PGA score of a subject, may be measured against a subject's baseline score. Generally, a baseline refers
10 to a measurement or score of a patient before treatment, *i.e.* week 0. In certain embodiments, however, other time points may also be included as a starting point in determining efficacy.

Patients or patient populations described in the methods of the invention are generally selected based on common characteristics, such as, but not limited to, subjects
15 diagnosed with hidradenitis suppurativa. Such a patient or patient population would be appropriate for determining the efficacy of the TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) for treating hidradenitis suppurativa in the given patient population. In one embodiment, the patient or patient population is an adult population, *e.g.*, older than 17 years of age or older than 18 years of age. In certain
20 embodiments, the patient or patient population has a diagnosis of moderate to severe hidradenitis suppurativa for at least 6 months prior to baseline measurement of HS-PGA, and involved at least two distinct anatomic areas (*e.g.*, left and right axilla; or left axilla and left inguinal-crural fold). In one embodiment, subjects have been unresponsive or intolerant to oral antibiotics for treatment for their hidradenitis suppurativa. In one
25 embodiment, subjects have an HS-PGA score of 3 or greater.

In one embodiment, the methods of the invention is used to determine whether a TNF α inhibitor is an effective TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) with respect to a patient population who has already been administered the TNF α inhibitor. Such a patient population may be pre-selected
30 according to common characteristics, *e.g.*, HS-PGA score, and may have already been given the TNF α inhibitor. Administration of the TNF α inhibitor (*e.g.*, an anti-TNF α

antibody, or an antigen-binding portion thereof) may or may not be performed by the same person of ordinary skill who is determining the efficacy of the TNF α inhibitor in accordance with the teachings of the specification.

In one embodiment, the methods of the invention comprise administering the
5 TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) to the subjects of a patient population and determining the efficacy of the TNF α inhibitor (*e.g.*, an anti-TNF α antibody, or an antigen-binding portion thereof) by determining changes, improvements, measurements, *etc.*, using HS-PGA scores of the patient population in comparison to the Examples set forth below.

10 In addition, while some methods are described in terms of patient populations, methods of efficacy described herein may also be applied to individual subjects. For example, a method for determining efficacy may comprise determining whether a subject who has hidradenitis suppurativa, and who is on a dosage regimen comprising a human TNF α antibody, is able to achieve a clinical response as defined herein, in order to
15 determine if the human TNF α antibody is an effective human TNF α antibody. In one embodiment, if the subject is able to achieve a clinical response as defined herein for at least about 8, 12, 16, 20, 24, 30, 36, 42, 48, 52, 56 weeks or more, then the human TNF α antibody is effective at treating hidradenitis suppurativa.

The Examples and discoveries described herein are representative of a
20 TNF α inhibitor, *i.e.*, adalimumab, which is effective for treating hidradenitis suppurativa. As such, the studies and results described in the Examples section herein may be used as a guideline for determining the efficacy of a TNF α inhibitor, *i.e.*, whether a TNF α inhibitor is an effective TNF α inhibitor for the treatment of hidradenitis suppurativa. In one embodiment, methods of determining efficacy described herein may
25 be used to determine whether a TNF α inhibitor is bioequivalent to another TNF α inhibitor.

In one embodiment, the article of manufacture of the invention comprises instructions regarding how to determine the efficacy of the TNF inhibitor for the treatment of hidradenitis suppurativa.

30 The present invention is further illustrated by the following examples which should not be construed as limiting in any way.

EXAMPLES

Example 1 Safety and Efficacy of Adalimumab in Subjects with Moderate to Severe Chronic Hidradenitis Suppurativa (HS)

5 This example demonstrates that anti-TNF α antibodies, such as the fully human anti-TNF α antibody adalimumab, are efficacious and safe for treating human hidradenitis suppurativa (HS) patients, especially human patients with moderate to severe chronic hidradenitis suppurativa.

10 In this Phase II clinical trial study, qualified hidradenitis suppurativa patients were first randomized into three treatment groups in a 1:1:1 ratio: (1) weekly subcutaneous (*s.c.*) injections of about 40 mg adalimumab (40 mg qwk); (2) biweekly *s.c.* injection of about 40 mg adalimumab (40 mg eow); and (3) matching placebo with no adalimumab injection. More details regarding the three treatment arms are described below. Randomization was stratified by Hurley staging (III vs. I or II) for HS.

15 This period of the study (Period 1) was conducted over 16 weeks, as a double-blind, placebo-controlled treatment period to evaluate efficacy and safety. Patients were then invited to participate in a 36-week open-label follow-up period (Period 2), in which all patients receive open-label adalimumab (40 mg eow) *s.c.* injection for an evaluation of long-term safety and efficacy (see Example 2 below). Patients who had received
20 placebo in Period 1 received an initial blinded 80 mg dose of adalimumab at Week 16, and patients who had received active therapy in Period 1 received blinded placebo at Week 16. Patients with HS-PGA scores ≥ 3 (moderate or worse) at Weeks 28 or 31 also had the option to dose escalate to 40 mg adalimumab weekly. Those whose dose escalated remained on ew dosing for the remainder of the study.

25

Study Treatment Arms

 Arm A (Adalimumab 40 mg ew): subjects randomized to Arm A received a loading dose of adalimumab 160 mg at Week 0 and adalimumab 80 mg at Week 2, followed by adalimumab 40 mg weekly starting at Week 4 through Week 15.

Arm B (Adalimumab 40 mg eow): subjects randomized to Arm B received a loading dose of adalimumab 80 mg at Week 0, followed by adalimumab 40 mg eow starting at Week 1 through Week 15.

Arm C (Placebo): subjects randomized to Arm C received matching placebo
5 administered weekly, starting at Week 0 through Week 15.

In the event that a patient had an acutely painful lesion, investigators had the option to intervene with either an injection of intralesional triamcinolone acetonide suspension or performance of incision and drainage. Two protocol-allowed interventions were permitted during Period 1. Patients requiring more than two
10 interventions during Period 1 were to be discontinued from the study.

Population

Qualified subjects were males and females ≥ 18 years old, had a diagnosis of moderate to severe hidradenitis suppurativa (HS-PGA score of moderate or worse; see
15 Table 1 for HS-PGA Scoring System) for at least 6 months prior to Baseline that involved at least two distinct anatomic areas (*e.g.*, left and right axilla; or left axilla and left inguinal-crural fold). Subjects must have been unresponsive or intolerant to oral antibiotics for treatment for their hidradenitis suppurativa. In addition, qualified subjects were required to have a hidradenitis suppurativa-PGA score of 3 or greater. Patients
20 who had prior treatment with any anti-TNF therapy, including adalimumab, infliximab or etanercept, were excluded from the study. Also excluded are patients who had used systemic non-biologic therapies (other than certain permitted oral antibiotics) for HS within 4 weeks prior to Baseline visit.

Use of the following oral and/or topical antibiotic therapy for HS was permitted
25 if patients had received a stable dose for ≥ 4 weeks prior to Baseline visit and the dose remained stable during study: 1% topical clindamycin bid; Tetracycline (up to 500 mg po bid); Doxycycline (up to 100 mg po bid); or Minocycline (up to 100 mg po bid).

More specifically, patient inclusion criteria included adults with stable, moderate to severe hidradenitis suppurativa. Patients also had to have had a negative Chest X-ray
30 and PPD test at Screening. If a participant had a past ulcerative reaction to PPD placement and/or chest X-ray consistent with prior tuberculosis exposure, the participant

had to initiate, or have documented completion of, a course of anti-tuberculosis therapy. Participants had to have the ability to administer subcutaneous injections, and be in general good health otherwise.

Patient exclusion criteria included patients who had prior anti-TNF therapy and unstable antibiotic therapy for HS. The exclusion criteria also required medication washouts for other HS treatments. Patient exclusion criteria further included prior exposure to Tysabri® (natalizumab), recent infection requiring treatment, any significant medical events or conditions that may put patients at risk for participation, female subjects who were pregnant or breast-feeding or considering becoming pregnant during the study, a history of cancer, except successfully treated skin cancer, and any recent history of drug or alcohol abuse.

Participating sites were predominantly in the United States (Alabama, California, Florida, Georgia, Illinois, Indiana, Massachusetts, Missouri, Nebraska, New York, North Carolina, Pennsylvania, Texas, Virginia), with some sites located in the Netherlands, Denmark, and Germany.

One hundred ninety five patients were screened for entry in the study: 41 were screen failures, and 154 patients from 26 centers in four countries (Denmark, Germany, Netherlands, and the United States) were enrolled (Figure 1). The most common reason for patients failing screening was not meeting screening criteria.

Efficacy Endpoint(s)

The assessed efficacy endpoints include: Hidradenitis Suppurativa Physician's Global Assessment (HS-PGA) response (scale below, definition of HS-PGA response described above); change in abscess, draining fistula, and inflammatory nodules; and QoL assessment, change in pain score. Specifically, at screening and study visits, physicians assessed counts of nodules (inflammatory and non-inflammatory), abscesses, and fistulas (draining and non-draining), and used these counts to assign patients to one of six ordinal categories (clear, minimal, mild, moderate, severe, very severe) of the HS-PGA scale according to Table 1 below.

Table 1 HS-PGA Scoring System

Score	Rating	Description
0	Clear	No abscesses, no draining fistulas, no nodules
1	Minimal	No abscesses, no draining fistulas, no inflammatory nodules, presence of non-inflammatory nodules
2	Mild	No abscesses or draining fistulas, and less than 5 inflammatory nodules, or Single abscess or draining fistula, and no inflammatory nodules
3	Moderate	No abscesses or draining fistulas, and at least 5 inflammatory nodules, or Single abscess or draining fistula in the presence of inflammatory nodules, or Between 2 and 5 abscesses or draining fistulas with or without inflammatory nodules, up to 10
4	Severe	Between 2 and 5 abscesses and draining fistulas with or without inflammatory nodules that are greater than 10
5	Very severe	More than 5 abscesses or draining fistulas

A clinical response to treatment was defined as an HS-PGA score of clear, minimal, or mild with at least a two-grade improvement relative to baseline score. Pain was assessed with a visual analogue score (VAS), ranging from 0 mm (no pain) to 100 mm (maximal pain). The following patient reported instruments were included in the study: the Dermatology Life Quality Index (DLQI) (measures dermatology-specific health-related quality of life ranging from 0-30 with 0 being no impairment), Work Productivity and Activity Impairment-Specific Health Problem: Psoriasis (WPAI-SHP) (ranges from 0-100 with 0 being no impairment), Patient Health Questionnaire (PHQ-9) (self-assessment for depression ranging 0-27 with 0 being no depressive symptoms). Blood samples were collected to measure levels of C-reactive protein (CRP) and other hematologic and biochemical markers and immunogenicity.

The primary endpoint of the study was the proportion of patients achieving clinical response at Week 16 (see Table 1 for HS-PGA Scale). Secondary efficacy measures included the proportion of patients achieving clinical response at Weeks 2, 4, 8, and 12, and all study visits during Period 2; the proportion of patients achieving at least a two-grade improvement in PGA scale at Week 16; the proportion of patients achieving a PGA score of clear, minimal or mild at Week 16; the proportion of patients achieving complete clearance of abscesses, draining fistulas, or inflammatory nodules at Week 16 and mean improvement in these counts from Baseline to Week 16; the

proportion of patients achieving $\geq 30\%$ reduction and 10 point absolute reduction in VAS score (among those patients with at least 10 mm VAS score at baseline); mean change in CRP levels from Baseline to Week 16; and, mean change in DLQI, total work productivity impairment (TWPI) outcome from the WPAI-SHP, and PHQ-9 scores from

5 Baseline to Week 16.

More specifically, main secondary endpoints assessed included the following:

Percent Change From Baseline in Number of All Inflammatory Nodules and Plaques at Week 16. Specifically, the number of all inflammatory nodules and plaques includes inflammatory nodules that were tender, erythematous, and had diameters less

10 than 5 cm and included plaques that had diameters greater than or equal to 5 cm. Range for percent change was negative infinity to infinity. Negative percent changes from Baseline indicated improvement;

Percentage of Participants Achieving Clinical Response at Week 2 from Baseline. Specifically, clinical response was defined as Hidradenitis Suppurativa-

15 Physician's Global Assessment (HS-PGA) of clear, minimal, or mild (scores of 0, 1, or 2) with a minimum of 2 grades improvement (reduction) from baseline. PGA is a physician's assessment of the severity of hidradenitis suppurativa based on a 6-point scale (score of 0 = clear and 5 = very severe);

Percentage of Participants Achieving Clinical Response at Week 4 from

20 Baseline. Specifically, clinical response was defined as a Hidradenitis Suppurativa-Physician's Global Assessment (HS-PGA) score of clear, minimal, or mild (scores of 0, 1, or 2) with a minimum of 2 grades improvement (reduction) from baseline. PGA is a physician's assessment of the severity of hidradenitis suppurativa based on a 6-point scale (score of 0 = clear and 5 = very severe);

Percentage of Participants Achieving Clinical Response at Week 8 from

25 Baseline. Specifically, clinical response was defined as a Hidradenitis Suppurativa-Physician's Global Assessment (HS-PGA) of clear, minimal, or mild (scores of 0, 1, or 2) with a minimum of 2 grades improvement (reduction) from baseline. HS-PGA is a physician's assessment of the severity of disease based on a 6-point scale (score of 0 =

30 clear and 5 = very severe);

Percentage of Participants Achieving Clinical Response at Week 12 from Baseline. Specifically, clinical response was defined as a Hidradenitis Suppurativa-Physician's Global Assessment (HS-PGA) of clear, minimal, or mild (scores of 0, 1, or 2) with a minimum of 2 grades improvement (reduction) from baseline. HS-PGA is a physician's assessment of the severity of hidradenitis suppurativa based on a 6-point scale (score of 0 = clear and 5 = very severe);

Change From Baseline in Modified Sartorius Scale at Week 16. Specifically, the Modified Sartorius Scale reflects changes in hidradenitis suppurative symptoms, namely the number of lesions (abscesses, nodules, and fistulas) and the longest distance between lesions. A total score was derived based on assessments at up to 8 distinct anatomical regions and ranges from 5 to indefinite. Smaller numbers are better scores and indicate less lesion involvement, thus decreases (negative changes) from baseline indicate improvement in severity of disease;

Change From Baseline in Modified Sartorius Scale at Week 52. Specifically, the Modified Sartorius Scale reflects changes in hidradenitis suppurative symptoms, namely the number of lesions (abscesses, nodules, and fistulas) and the longest distance between lesions. A total score was derived based on assessments at up to 8 distinct anatomical regions and ranges from 5 to indefinite. Smaller numbers are better scores and indicate less lesion involvement, thus decreases (negative changes) from baseline indicate improvement in severity of disease;

Percent Change From Baseline in Number of All Inflammatory Nodules and Plaques at Week 52. Specifically, the number of all inflammatory nodules and plaques includes inflammatory nodules that were tender, erythematous, and had diameters less than 5 cm and included plaques that had diameters greater than or equal to 5 cm. Range for percent change was negative infinity to infinity. Negative percent changes from Baseline indicate improvement;

The proportion of patients achieving clinical success at Weeks 2, 4, 8, and 12;

The proportion of patients achieving an HS-PGA score of clear, minimal, or mild at Week 16; and

The proportion of patients achieving $\geq 30\%$ reduction and ≥ 10 point absolute

reduction in VAS (Visual Analog Scale) (among patients with VAS pain score ≥ 10 at Baseline) at Weeks 2, 4, 8, 12, and 16.

Safety assessments included incidences of adverse events (AE) assessed throughout the study, and up to 70 days after last dose of study drug, or up to the date
5 prior to the first dose in Period 2 for those who entered Period 2.

Statistical Methods

The study was planned to enroll 150 patients to have 80% power to detect a clinically relevant treatment difference, assuming a 10% clinical response rate for
10 placebo-treated patients and a 35% response rate for adalimumab-treated patients.

Efficacy analyses were conducted on the intent-to-treat (ITT) populations in each period: for Period 1, the ITT population consisted of all patients randomized at Week 0; for Period 2, the integrated ITT population consisted of all patients randomized to adalimumab eow and adalimumab ew arms at Week 0, and patients randomized to the
15 placebo arm at Week 0 who entered Period 2. Primary efficacy analysis was performed by Cochran-Mantel-Haenszel (CMH) test adjusting for baseline Hurley stage with non-responder imputation (NRI) as the primary approach and LOCF as sensitivity approach to impute missing data. To control for multiplicity, an initial overall test with all three treatment groups was performed and pair-wise comparison of each adalimumab dose
20 group vs. placebo group were conducted only when the overall test was significant.

CMH and Analysis of Covariance (ANCOVA) with factors of treatment and Hurley stage were used for categorical and continuous secondary efficacy variables, respectively; NRI, last-observation-carried forward (LOCF), and as-observed approaches were used as appropriate. All statistical tests were two-sided with the significance level
25 of 0.05.

All statistical tests were 2-sided with the significance level of 0.05. Pair-wise comparisons of adalimumab group vs. placebo group were conducted if overall comparison was significant.

The safety analyses were conducted in the safety population (patients in the ITT
30 population receiving ≥ 1 dose of study drug); safety variables were summarized by

treatment group.

Demographics and Clinical Characteristics

A total of 154 subjects were randomly assigned to one of the three treatment arms at Week 0: 51 to placebo (pbo), 52 to every other week (eow), and 51 to weekly (qw) therapy. Randomization was stratified by Hurley staging {III vs. (I or II)} for hidradenitis suppurativa (*see* Poli F, Jemec GBE, Revuz J., Clinical Presentation. In: Jemec GBE, Revuz J, Leyden JJ, editors. Hidradenitis Suppurativa. Springer, New York, 2006, pp 11-24).

Hurley staging is a severity scale which assesses both current activity and past scarring, ranging from isolated abscesses in the primary stage to coalescing lesions with scarring and sinus tracts in the tertiary stage. Hurley stage III disease is the most severe stage of hidradenitis suppurativa, reflecting diffuse or near-diffuse involvement of affected areas. The percentage of enrolled subjects with Hurley stage III was not to exceed 50%.

Table 2 Baseline Demographics and Clinical Characteristics

	Placebo (n=51)	ADA eow (n=52)	ADA ew (n=51)
Age (yrs), mean (SD)	37.8 (12.10)	36.1 (12.50)	35.1 (10.69)
Female, n (%)	36 (70.6)	38 (73.1)	36 (70.6)
White, n (%)	37 (72.5)	36 (69.2)	37 (72.5)
Black	8 (15.7)	12 (23.1)	9 (17.6)
Nicotine users, n (%)	29 (56.9)	26 (50.0)	30 (58.8)
Body weight (kg), mean (SD)	96.5 (24.80)	99.8 (26.75)	95.4 (22.94)
BMI, n (%)			
<25	9 (17.6)	6 (11.5)	9 (17.6)
≥25-<30	6 (11.8)	11 (21.2)	12 (23.5)
≥30	36 (70.6)	35 (67.3)	30 (58.8)
Disease duration (yrs), mean (SD)	13.4 (10.4)	10.9 (9.0)	11.3 (9.1)
CRP (mg/L), mean (SD) ^a	13.3 (15.0)	17.8 (26.9)	21.5 (33.1)
Hurley stage I or II, n (%)	36 (70.6)	37 (71.2)	36 (70.6)
Hurley stage III, n (%)	15 (29.4)	15 (28.8)	15 (29.4)
HS-PGA moderate, n (%)	33 (64.7)	35 (67.3)	35 (68.6)
HS-PGA severe/very severe, n (%)	17 (33.3)	16 (30.8)	16 (31.4)
Patients receiving p.o. doxycycline or minocycline, n (%)	4 (7.8)	6 (11.5)	8 (15.7)
Prior topical therapy, n (%)	27 (52.9)	26 (50.0)	23 (45.1)
Prior systemic therapy, n (%)	49 (96.1)	52 (100.0)	50 (98.0)

Prior pain medication use, n (%)	23 (45.1)	17 (32.7)	17 (33.3)
Prior opioid use, n (%)	7 (13.7)	7 (13.5)	7 (13.7)
VAS skin pain, mean (SD)	57.8 (28.51)	53.0 (26.35)	52.0 (24.51)
DLQI, mean (SD)	15.4 (7.71)	13.5 (7.65)	16.4 (7.48)
PHQ-9, mean (SD)	9.1 (6.8)	8.1 (6.1)	11.1 (7.0)
TWPI, mean (SD)	31.4 (34.7)	35.1 (29.5)	45.5 (32.8)
HS-PGA Components			
Abscesses, median (IQR)	1 (2.0)	1 (2.0)	1 (2.0)
Patients with abscess, n (%)	29 (56.9)	29 (55.8)	20 (39.2)
Draining fistulas, mean (median)	1 (3.0)	1 (3.5)	1 (3.0)
Patients with draining fistulas, n (%)	33 (64.7)	28 (53.8)	30 (58.8)
Inflammatory nodules, mean (median)	6 (10.0)	7 (14.5)	7 (10.0)
Patients with inflammatory nodules, n (%)	49 (96.1)	48 (92.3)	50 (98.0)

a: Based on patients with non-missing values; placebo, n = 39; eow, n = 40; ew, n = 38. ADA, adalimumab; BMI, body mass index; HS-PGA, Hidradenitis suppurativa physician's global assessment; VAS, visual analog scale; DLQI, dermatology life quality index; TWPI, total work productivity impairment; IQR, interquartile range.

5

It is apparent that baseline demographics were generally well-balanced across the treatment arms (see Table 2). The majority of all enrolled subjects were female (71.4%), white (71.4%), less than 40 years old (63.6% were less than 40 years old, with a mean age of 36.3 years), smokers (55.2%), and had a Hurley Stage II (55.2%). Mean weight for all enrolled subjects was 97.2 kg. Baseline characteristics were similar across treatment groups. Mean pain score was 54.3/100 and the percentages of patients with HS-PGA scores of moderate, severe, or very severe were 66.9%, 9.7%, and 22.1%, respectively (Table 3).

15

Table 3 Distribution of HS-PGA at Baseline (%)

PGA Category	Placebo	Every Other Week	Weekly
Clear (0)	0	0	0
Minimal (1)	2.0	0	0
Mild (2)	0	1.9	0
Moderate (3)	64.7	67.3	68.6
Severe (4)	9.8	9.6	9.8
Very Severe (5)	23.5	21.2	21.6

Mean DLQI scores at baseline indicated that HS had a large deleterious effect on patients' dermatology-specific quality of life. The mean baseline DLQI score of 15.1 was worse than the mean baseline DLQI score of 11.4 for patients enrolled in an adalimumab Phase III psoriasis trial (REVEAL, see Revicki *et al.*, *Dermatology* 216: 260-270, 2008). Overall baseline demographics and clinical characteristics were similar across the three treatment arms.

Of the 154 subjects who enrolled, 11 discontinued during Period 1: 5 from the placebo group and 6 from the qw group. One of the discontinued subjects in the qw group withdrew for the primary reason of adverse event.

Overall, 90.2% of placebo patients, 100% of adalimumab (ADA) eow patients, and 88.2% of ADA ew patients completed Period 1. See Table 4 below. Of patients entering Period 2, 73.9% of placebo/eow patients, 74.5% of eow/eow patients, and 68.9% of ew/eow patients completed Period 2. Baseline demographics and clinical characteristics are summarized in Table 2. Of note, these patients were markedly obese, with average weights over 90 kg. Over 20% of the enrollees were African-American, suggesting that the epidemiology of the disease may be linked to the African-American population. They also had high entry levels of pain and substantial use of narcotics for this problem, with around 13% reporting opioid use. Prior systemic antibiotic usage for treatment of HS was reported by 144 patients (94%); of these, 104 (72%) reported no satisfactory response. Other systemic therapies that had been utilized included corticosteroids (29 patients, 19%) and retinoids (28 patients, 18%).

Table 4 Patient Disposition

	Placebo (n=51)	ADA eow (n=52)	ADA ew (n=51)
Completed Period 1	46 (90.2%)	52 (100%)	45 (88.2%)
Primary reason for discontinuation			
Adverse event	0	0	1 (2.0%)
Withdraw consent	2 (3.9%)	0	1 (2.0%)
Lack of efficacy	0	0	1 (2.0%)
Lost to follow-up	2 (3.9%)	0	1 (2.0%)
Exceeded protocol specified number of interventions ^a	1 (2.0%)	0	0

other ^b	0	0	2 (3.9%)
--------------------	---	---	----------

a. A maximum of 2 interventions were permitted during Period 1.

b. One discontinued due to non-compliance and 1 per investigator discretion.

Efficacy Results

5 The primary endpoint for this study was the proportion of subjects achieving a clinical response, defined as achieving an HS-PGA of clear (0), minimal (1), or mild (2), with an improvement (*i.e.*, reduction) from Baseline of at least 2 grades at Week 16.

All responders were in the Hurley Stage I/II stratum with the exception of one qw subject that was a responder in the Hurley Stage III stratum.

10 When evaluating the proportion of subjects in the eow arm and in the qw arm who improved at least two HS-PGA grades from Baseline to Week 16, both eow and qw treatment groups had response rates significantly higher than that of placebo: 3.9% for placebo subjects, 21.2% for eow subjects ($p=0.009$ vs. placebo), and 21.6% for qw subjects ($p=0.008$ vs. placebo).

15 Table 5 describes the proportion of patients achieving clinical response (defined as achieving an HS-PGA of clear, minimal, or mild, and at least 2 grade improvement relative to baseline) in each group. Specifically, a significantly greater proportion of patients allocated to the ew group achieved the primary endpoint, a clinical response at Week 16, compared with patients allocated to the placebo group (17.6% vs. 3.9%, $p =$
20 0.03; Figures 1 and 7). For the primary endpoint, point estimates for number of patients needed to treat (NNT) were 18 for the eow group and 8 for the ew group. The clinical response rate at Week 16 with ew dosing was 22.2% (8/36) for patients with Hurley Stage I or II at baseline, compared with 6.7% (1/15) for patients with Hurley Stage III at baseline.

25 Table 5. Proportion (Percentage) of Patients Achieving Clinical Response

Patient Group	Week 2	Week 4	Week 8	Week 12	Week 16
Placebo	2	2	7.8	5.9	3.9
EOW	9.6	5.8	5.8	7.7	9.6
QWK	2	11.8	7.8	21.6*	17.6*

* $p=0.005$ at Week 12, or $p=0.006$ at Week 16, qwk versus placebo.

Post-analysis also revealed that a higher proportion of current smokers receiving weekly therapy achieved a Clinical Response at Week 16 compared with current non-smokers receiving weekly therapy (23.3% vs. 9.5%). The percentage of subjects receiving concomitant oral antibiotics was too small to permit meaningful inferences about the effect of antibiotic usage on clinical response rates. Among patients receiving weekly therapy, a higher proportion of patients with BMI greater than or equal to the median BMI achieved a clinical response at Week 16 compared with patients below the median BMI (23% vs. 14%).

At Week 16, an HS-PGA score of clear, minimal, or mild was achieved by 49.0% of ew patients, 21.2% of eow patients, and 23.5% of placebo patients ($p < 0.01$, ew vs. pbo).

Mean reduction (improvement) from baseline to Week 16 in the modified Sartorius scale was 33.0 for ew patients, 32.0 for eow patients, and 16.7 for placebo patients. Clinical photographs of the perineum of a patient receiving ew therapy showed marked improvement (data not shown).

Significant improvement was also seen when evaluating the proportion of subjects who improved at least one HS-PGA grade from Baseline to Week 16: 28.0% for placebo subjects, 40.4% for eow subjects, and 66.0% for ew subjects ($p < 0.001$ vs. placebo). Upon further analysis, the proportion of subjects who improved at least one HS-PGA grade from Baseline to Week 16 is: 27.5% for placebo subjects, 40.4% for eow subjects, and 56.9% for ew subjects ($p = 0.002$ vs. placebo).

Overall, at Week 16, a statistically significantly greater proportion of ADA ew patients (49.0%) achieved an HS-PGA of clear, minimal, or mild as compared with placebo patients (23.5%) ($p < 0.01$, Table 6). This data also suggests that the tested adalimumab weekly dosing (ew) may be more effective in achieving an HS-PGA of clear, minimal or mild as compared to once-every-other week (eow or biweekly) dosing (21.2%) at Week 16.

Table 6. Proportion (Percentage) of Patients Achieving an HS-PGA of Clear, Minimal or mild at Week 16 of Treatment

Patient Group	Week 16
Placebo (n=51)	23.5
EOW (n=52)	21.2
EW (n=51)	49.0*

*p<0.01, placebo vs. ADA ew at Week 16

In fact, each of the individual components of the HS-PGA score improved at Week 16 for patients receiving ew therapy compared with placebo-treated patients; patients receiving eow therapy generally experienced less improvement than ew treated patients on each of these components (see below).

Adalimumab therapy, particularly ew dosing, was associated with significant improvement in various other patient-reported outcomes. Improvement in Pain VAS scores of $\geq 30\%$ was considered to constitute a clinically relevant improvement in pain, and was thus a predefined secondary analysis. The proportion of subjects whose pain scores improved at least 30% from baseline to Week 16 was significantly higher than placebo in the qw arm only: 27.1% for placebo subjects, 36.2% for eow subjects, and 47.9% for qw subjects (p=0.037 vs. placebo). See Table 7.

Moreover, improvement in pain was rapid, with significantly more patients in both active treatment arms achieving at least 30% and at least 10 mm reduction in pain by Week 2. Specifically, by Week 2, 44.7% (p < 0.01 vs. placebo) of subjects receiving adalimumab every other week (ADA eow) and 41.7% (p < 0.05 vs. placebo) of subjects receiving adalimumab every week (ADA ew) experienced at least a 30% and at least a 10 mm reduction in pain when compared to placebo's 18.8%. By week 4, 58.3% (p < 0.001 vs. placebo) of ADA ew subjects and 46.8% (p < 0.05 vs. placebo) of ADA eow subjects showed a greater than 30 % and greater than 10 mm reduction from baseline pain, compared to the 22.9% of the placebo group. At week 12, 60.4% (p < 0.01 vs. placebo) of ADA ew subjects maintained statistically significant pain reduction over the placebo group (29.2%). See Table 7.

Table 7. Proportion (Percentage) of Patients Achieving a $\geq 30\%$ and ≥ 10 mm Reduction from Baseline in Pain at Weeks 2, 4, 8, 12 and 16 of Treatment Among Patients with ≥ 10 mm VAS Pain Score at Baseline

Patient Group	Week 2	Week 4	Week 8	Week 12	Week 16
Placebo (n = 48)	18.8	22.9	29.2	29.2	27.1

EOW (n = 47)	44.7 [†]	46.8*	40.4	42.6	36.2
EW (n = 48)	41.7**	58.3 [§]	43.8	60.4 [‡]	47.9**

*p < 0.05, ADA eow vs. placebo; **p < 0.05, ADA ew vs. placebo;

[†]p < 0.01, ADA eow vs. placebo; [‡]p < 0.01, ADA ew vs. placebo;

[§]p < 0.001, ADA ew vs. placebo

5 Among patients with ≥ 10 mm VAS pain scores at baseline, the proportion with a clinically significant improvement in pain (at least 30% reduction and 10 mm reduction in pain) at Week 16 was significantly higher for patients in the ew group compared with the placebo group (47.9% vs. 27.1%, $P < 0.05$); more than 40% of patients receiving ew or eow therapy crossed this threshold of pain reduction at Week 2.

10 Mean reduction (improvement) in DLQI scores was significantly greater for subjects in the qw arm only when compared to placebo: 1.9 for placebo subjects, 2.8 for eow subjects, and 6.0 for qw subjects ($p < 0.001$ ew vs. placebo). Work productivity was significantly improved among patients in the ew group as compared with the placebo group: mean reduction (improvement) in TWPI scores between baseline and Week 16
15 were 17.4 for the ew group and 0.94 for the eow group; placebo patients experienced a 2.93 increase (deterioration) in TWPI score ($p < 0.001$, ew vs. placebo). Mean reduction (improvement) in the PHQ-9 depression measure between baseline and Week 16 was 3.8 for the ew group, 1.4 for the eow group, and 1.2 for the placebo group ($p < 0.05$, ew vs. placebo).

20 Raw counts of the abscesses, draining fistulas, and inflammatory nodules were the key clinical components of the PGA score. Table 8 below depicts the mean decrease in counts (positive value represents improvement) for these lesion types, from baseline to Week 16, for subjects in the three treatment arms:

25 Table 8. Mean Decrease in Counts for Abscesses, Draining Fistulas and Inflammatory Nodules

Patient Group	Abscesses	Draining Fistulas	Inflammatory Nodules
Placebo	0.42	-1.03	1.93
EOW	1.43	0.04	6.18*
QW	1.85	-4.93	5.66*

*p < 0.05, qw vs. placebo

For inflammatory nodules, abscesses, and draining fistulas, Table 9 below provides the mean/median absolute change from baseline to Week 16, the mean/median percentage change from baseline to Week 16, and the proportion of patients achieving complete clearance at Week 16. The qwk dosing arm notably outperformed the eow dosing arm and placebo arm in mean/median percentage decrease in inflammatory nodules, mean/median percentage decrease in draining fistulas, and in the proportion of patients with complete clearance of draining fistulas. While the eow dosing arm outperformed the qwk dosing arm in mean absolute decrease in draining fistulas, the treatment effect for eow dosing relative to placebo was small with this endpoint (difference in mean change of less than 1 draining fistula per patient), and the mean increase in the qwk group was impacted by a single outlier.

Table 9. Percent Reduction, Absolute Reduction, Proportion of Patients with Complete Clearance (%) Changes in Primary Lesions at Week 16

	Placebo	Eow	Qwk
Percentage Reduction ¹			
Inflammatory Nodules (mean)	13.7	30.4	50.7 (p<0.05 vs. placebo)
Inflammatory Nodules (median)	17.86	48.39	66.67
Abscesses (mean)	25.0	46.2	51.8
Abscesses (median)	100.00	75.00	92.86
Draining Fistulas (mean)	7.5	7.7	44.4 (p<0.05 vs. placebo)
Draining Fistulas (median)	11.11	12.5	96.15
Absolute Reduction ¹			
Inflammatory Nodules (mean)	1.93	6.18 (p<.01 vs. placebo)	5.66 (p<0.05 vs. placebo)
Inflammatory Nodules (median)	1.0	3.5	4.0
Abscesses (mean)	0.42	1.43	1.85
Abscesses (median)	1.0	1.0	1.0

Draining Fistulas (mean)	-1.03	0.04	-4.93*
Draining Fistulas (median)	1.0	0.5	1.5
Proportion of patients with complete clearance ¹ (%)			
Inflammatory Nodules	8.2	20.8	20.0
Abscesses	44.8	48.3	50.0
Draining Fistulas	27.3	25.0	43.3
¹ Among patients with any lesion at baseline. Positive value connotes patient improvement (reduction in counts). * The mean increase in qwk group was heavily impacted by a single outlier: a patient for whom was reported an increase of 210 draining fistula (from 100 at baseline to 310 at Week 16), while the maximum increase in the other two groups was 14.			

Table 10 below describes the proportion of patients achieving complete clearance of abscesses at Weeks 2, 4, 8, 12 and 16 among those with at least one lesion at baseline. The qwk dosing arm outperformed the eow dosing arm at every study visit except Week 5 2.

Table 10. Proportion (Percentage) of Patients Achieving Complete Clearance of Abscesses at Weeks , 4, 8, 12 and 16

Patient Group	Week 2	Week 4	Week 8	Week 12	Week 16
Placebo	37.9	37.9	44.8	44.8	44.8
EOW	41.4	48.3	51.7	58.6	48.3
QWK	35.0	50.0	55.0	60.0	50.0

Among patients with any abscesses at baseline. PBO = 29; EOW = 29; QWK = 20.

10 For all of these components (except for complete clearance of abscesses), patients with Hurley Stage III disease at baseline who received weekly adalimumab dosing experienced better improvement relative to those receiving placebo (data not shown). For all patients receiving ew dosing, at least half of the mean percentage improvement noted at Week 16 was observed by Week 4: mean percentage improvement
15 was 26% at Week 4 for inflammatory nodule count; 67% at Week 4 for abscess count; and 46% at Week 4 for draining fistula count. Between baseline and Week 16, mean serum CRP levels declined by 7.9 mg/L and 3.1 mg/L for the ew and eow groups, respectively, compared with an increase of 4.4 mg/L for the placebo group ($p < 0.05$, ew vs. placebo).

In one particular example, at baseline, a 45 year old subject with hidradenitis suppurativa duration of 10 years, 8 months, had an HS-PGA and VAS pain score of 47. Following 16 weeks of adalimumab 40 mg administered every other week (eow), the subject achieved an HS-PGA of 3 and a VAS pain score of 33. Reduction in HS lesions in this patient following adalimumab 40 mg administered every other week (eow) for 16 weeks was a clinically relevant improvement (reduction in pain of 34%).

Among patients receiving weekly therapy, a higher proportion of patients with BMI greater than or equal to the median BMI achieved a clinical response at Week 16 compared with patients below the median BMI (22.7% vs. 13.8); among placebo-treated patients, the analogous Clinical Response ratio for those $\geq$ median BMI: those $<$ median BMI was 0% vs. 7.7%.

Safety Results

For safety assessment, the 3 treatment groups in Period 1 (Adalimumab 40 mg Qwk, Adalimumab 40 mg Eow, and Placebo) were assessed from Week 0 up to Week 16. For Open-label adalimumab (Period 2), patients were assessed from Week 16 through Week 52, plus 70 days after the last dose. Assessment and recording of adverse events (AE) and serious adverse events (SAE) was performed by the investigator at each study visit. Information on events that occurred 70-days after the last dose of study drug (Week 52 or early termination) was collected during a follow-up phone call.

The proportion of subjects with any adverse event (AE) was 54.9% for placebo arm, 63.5% for eow arm, and 70.6% for qw arm. Generally, the types and frequency of adverse events were consistent with results observed in the adalimumab safety database for other indications. Three subjects (all Hurley Stage III) developed serious infections: an eow subject with super infection of scrotal hidradenitis, an eow subject with a pilonidal cyst flare, and a qw subject with polymicrobial infection of the penis and scrotum. One subject in the qw arm developed a vocal cord benign neoplasm (granular cell tumor).

No fatalities, tuberculosis, opportunistic infections, demyelinating disease, congestive heart failure, lupus-like syndrome, lymphoma, or nonmelanoma skin cancer were reported during Period 1.

Nine patients experienced serious adverse events (SAE) were reported during Period 1: small intestinal obstruction and suicide attempt (1 each in the placebo group); hidradenitis, interstitial lung disease, and pilonidal cyst (1 each in the ADA eow group); anemia, non-cardiac chest pain, genital bacterial infection and Escherichia infection [same patient], and vocal cord neoplasm in ew group. See Tables 11 and 12 below.

Table 11. Treatment-Emergent Adverse Events During Period 1

	Placebo (n=51)	ADA eow (n=52)	ADA ew (n=51)
Any AE	30 (58.8%)	33 (63.5%)	36 (70.6%)
Infectious AE	18 (35.3%)	22 (42.3%)	17 (33.3%)
Serious AE	2 (3.9%)	3 (5.8%)	4 (7.8%)
Serious Infectious AE	0	1 (1.9%)	1 (2.0%)
Malignancies	0	0	0
AE leading to withdrawal	0	2 (3.8%)	1 (2.0%)

AE: adverse event.

10 A detailed break-down of different categories of AE is presented in Table 12:

Table 12. Treatment-emergent Adverse Events During Period 1

	Adalimumab 40 mg Qwk DB	Adalimumab 40 mg Eow DB	Placebo DB
Total # participants affected / at risk	36/51 (70.59%)	33/52 (63.46%)	30/51 (58.82%)
Gastrointestinal disorders			
Gastrooesophageal reflux disease [†] _A			
# participants affected/at risk	3/51 (5.88%)	0/52 (0%)	0/51 (0%)
Nausea [†] _A			
# participants affected/at risk	4/51 (7.84%)	2/52 (3.85%)	1/51 (1.96%)
Vomiting [†] _A			
# participants affected/at risk	1/51 (1.96%)	2/52 (3.85%)	3/51 (5.88%)
General disorders			
Fatigue [†] _A			

# participants affected/at risk	3/51 (5.88%)	2/52 (3.85%)	2/51 (3.92%)
Infections and infestations			
Folliculitis † A			
# participants affected/at risk	0/51 (0%)	0/52 (0%)	3/51 (5.88%)
Nasopharyngitis † A			
# participants affected/at risk	6/51 (11.76%)	7/52 (13.46%)	6/51 (11.76%)
Upper respiratory tract infection † A			
# participants affected/at risk	4/51 (7.84%)	4/52 (7.69%)	2/51 (3.92%)
Musculoskeletal and connective tissue disorders			
Arthralgia † A			
# participants affected/at risk	3/51 (5.88%)	0/52 (0%)	1/51 (1.96%)
Nervous system disorders			
Headache † A			
# participants affected/at risk	8/51 (15.69%)	7/52 (13.46%)	2/51 (3.92%)
Respiratory, thoracic and mediastinal disorders			
Cough † A			
# participants affected/at risk	3/51 (5.88%)	1/52 (1.92%)	0/51 (0%)
Oropharyngeal pain † A			
# participants affected/at risk	1/51 (1.96%)	3/52 (5.77%)	1/51 (1.96%)
Skin and subcutaneous tissue disorders			
Hidradenitis † A			
# participants affected/at risk	4/51 (7.84%)	6/52 (11.54%)	6/51 (11.76%)
Pruritus † A			
# participants affected/at risk	1/51 (1.96%)	3/52 (5.77%)	0/51 (0%)
Gastrointestinal disorders			
Abdominal pain upper † A			
# participants affected/at risk	1/51 (1.96%)	2/52 (3.85%)	1/51 (1.96%)
Diarrhoea † A			
# participants affected/at risk	0/51 (0%)	2/52 (3.85%)	2/51 (3.92%)
General disorders			
Injection site pruritus † A			
# participants affected/at risk	2/51 (3.92%)	0/52 (0%)	0/51 (0%)

Oedema peripheral † A			
# participants affected/at risk	0/51 (0%)	1/52 (1.92%)	2/51 (3.92%)
Pain † A			
# participants affected/at risk	0/51 (0%)	2/52 (3.85%)	0/51 (0%)
Pyrexia † A			
# participants affected/at risk	0/51 (0%)	2/52 (3.85%)	1/51 (1.96%)
Infections and infestations			
Bronchitis † A			
# participants affected/at risk	1/51 (1.96%)	0/52 (0%)	2/51 (3.92%)
Ear infection † A			
# participants affected/at risk	0/51 (0%)	0/52 (0%)	2/51 (3.92%)
Herpes simplex † A			
# participants affected/at risk	0/51 (0%)	2/52 (3.85%)	0/51 (0%)
Influenza † A			
# participants affected/at risk	2/51 (3.92%)	1/52 (1.92%)	0/51 (0%)
Sinusitis † A			
# participants affected/at risk	2/51 (3.92%)	0/52 (0%)	1/51 (1.96%)
Tonsillitis † A			
# participants affected/at risk	0/51 (0%)	0/52 (0%)	2/51 (3.92%)
Investigations			
Blood cholesterol increased † A			
# participants affected/at risk	0/51 (0%)	0/52 (0%)	2/51 (3.92%)
Metabolism and nutrition disorders			
Hypercholesterolaemia † A			
# participants affected/at risk	1/51 (1.96%)	2/52 (3.85%)	0/51 (0%)
Musculoskeletal and connective tissue disorders			
Myalgia † A			
# participants affected/at risk	1/51 (1.96%)	1/52 (1.92%)	2/51 (3.92%)
Nervous system disorders			
Dizziness † A			
# participants affected/at risk	0/51 (0%)	1/52 (1.92%)	2/51 (3.92%)
Respiratory, thoracic and mediastinal disorders			

Dyspnoea † A			
# participants affected/at risk	0/51 (0%)	2/52 (3.85%)	0/51 (0%)
Sinus congestion † A			
# participants affected/at risk	2/51 (3.92%)	0/52 (0%)	0/51 (0%)
Skin and subcutaneous tissue disorders			
Dermatitis allergic † A			
# participants affected/at risk	2/51 (3.92%)	0/52 (0%)	0/51 (0%)
Pityriasis rosea † A			
# participants affected/at risk	2/51 (3.92%)	0/52 (0%)	0/51 (0%)
Psoriasis † A			
# participants affected/at risk	0/51 (0%)	0/52 (0%)	2/51 (3.92%)
Rash † A			
# participants affected/at risk	2/51 (3.92%)	0/52 (0%)	0/51 (0%)
Urticaria † A			
# participants affected/at risk	2/51 (3.92%)	0/52 (0%)	0/51 (0%)

† Indicates events were collected by systematic assessment.

A Term from vocabulary, MedDRA 13.1

Frequency Threshold Above Which Other Adverse Events are Reported: 3%

5 In summary, no deaths, malignancies, cases of tuberculosis or opportunistic infections occurred. During Period 1, the proportion of patients in each treatment arm experiencing adverse events (Tables 11 and 12) was higher for the eow and ew groups compared with the placebo group. Headaches, typically described as mild or moderate in severity, accounted for much of the numerical imbalance in adverse events among the treatment groups. A greater proportion of patients who started with weekly dosing in Period 1 and switched to eow dosing in Period 2 experienced any adverse event, any infectious adverse event, or any serious adverse event compared with patients who received eow dosing in Periods 1 and/or 2 (Tables 11 and 12 also see Table 13 below). Patients who underwent dose escalation in Period 2 had similar adverse event frequencies compared with patients who received eow dosing. Fifteen patients receiving adalimumab experienced one or more serious adverse events during the trial, with the most common types of serious adverse events being hidradenitis suppurativa (5 patients), infectious complications of hidradenitis suppurativa (4 patients), and anemia (2 patients;

one patient had a history of intermittent GI bleeding from ulcerative colitis and one patient had low hemoglobin concentration at screening).

The proportion of patients experiencing treatment-emergent common toxicity criteria (CTC) grade 3 adverse events in clinical chemistry or hematology values was low and similar across the treatment groups in Period 1, and remained low during Period 2. The prevalence of anti-adalimumab antibodies for the 52 week study was 10.4% (16 of 154 subjects).

The types and frequencies of adverse events were generally consistent with what has been observed in clinical trials of adalimumab in other indications. However, super infection of hidradenitis suppurativa lesions and/or pilonidal cyst flare, though infrequent, was observed with active treatment but not with placebo treatment. While not wishing to be bound by any particular theory, it is possible that this is a safety signal associated with adalimumab-mediated immunosuppression of contaminated skin and soft tissue, although it cannot be ruled out at this stage that it is a spurious signal. That said, the types and frequency of adverse events were consistent with results observed in other studies for treatment of other indications with adalimumab.

In conclusion, the above study showed that human anti-TNF α antibodies, such as adalimumab, are efficacious and safe for treating subjects having human hidradenitis suppurativa (HS), including patients with moderate to severe forms of the disease.

Example 2 Safety and Efficacy of Adalimumab in Subjects with Moderate to Severe Chronic Hidradenitis Suppurativa (HS)

The following example is a continuation of the study described in Example 1. This example shows the partial maintenance of efficacy and continued safety of adalimumab 40 mg every other week (eow) administration for hidradenitis suppurativa patients over a period of at least several weeks (at least 8 weeks) after the initial 16-week, double-blind, placebo-controlled treatment period. The pharmacokinetics and immunogenicity of adalimumab following subcutaneous (*s.c.*) injection were also assessed.

Subjects who have received 16 weeks of either weekly or biweekly injections of adalimumab (or placebo) in the randomized, double-blind, placebo-controlled treatment period, as described in Example 1 above, receive open-label adalimumab (40 mg eow) *s.c.* injection for an additional 36 weeks for evaluating long-term safety and efficacy of adalimumab administration in this patient population (study design is described in Figure 1). Patients having a sub-optimal response at Weeks 28 or 31 had the option of dose-escalating to ew dosing.

As described above in Example 1, 51 placebo (pbo), 52 every other week (eow), and 51 weekly (ew) patients were enrolled in Period 1. Week 16 Clinical Response rates were 3.9% for pbo, 9.6% for eow, 17.6% for ew patients ($P=0.025$, ew vs. pbo ($n=46/51/45$)).

Week 52 Clinical Response rates were 17.4% for pbo->eow ($n=46$), 5.9% for eow->eow ($n=51$), and 8.9% for ew->eow ($n=45$) patients. 89 patients had sub-optimal responses at Weeks 28 or 31, and were dose-escalated to ew dosing; of these, 13 (15%) had a clinical response at Week 52. The percentage of patients with serious adverse events was 3.9% for pbo, 5.8% for eow, and 7.8% ew patients during Period 1; 2.2% for pbo->eow, 3.9% for eow->eow, and 4.4% for ew->eow patients during Period 2; and 5.6% for dose escalation patients.

Among patients who had received ew dosing in Period 1, the proportion of patients achieving clinical response was maintained for 8 weeks after switching to eow dosing in Period 2 (Figure 3). However, with continued eow dosing, the proportion of patients achieving clinical response declined, and eventually became similar to the response rate for patients who had never received weekly dosing (but have received eow dosing since week 16). Of 24 ew-treated patients who had HS-PGA of mild or better response at the entry of Period 2, 12 (50%) retained this response at Week 28, 12 weeks following step-down to eow dosing. Eighty-nine patients (63% of those who entered Period 2) had a sub-optimal response at Weeks 28 or 31 and were dose-escalated to ew dosing; of these, 13 (15%) had a clinical response at Week 52. Thus it appears that ew dosing (for 40 mg ADA) was more effective than eow dosing, although eow dosing was sufficient for a proportion of the patients.

The safety profile was similar for patients receiving adalimumab weekly therapy compared to those receiving every other week therapy. Table 13 below compares AE between Period 1 and Period 2.

Table 13. Adverse Events (AE) During Period 1 and Period 2

	Period 1			Periods 1 and 2 ^a		
	Placebo n=51	ADA eow n=52	ADA ew n=51	PBO/eow+eow/eow N=98	ew/eow N=51	Dose Escalation N=89
n (%)						
Any adverse event	30 (58.8)	33 (63.5)	36 (70.6)	60 (61.2)	44 (86.3)	50 (56.2)
Any infectious adverse event	18 (35.3)	22 (42.3)	17 (33.3)	41 (41.8)	30 (58.8)	24 (27.0)
Serious adverse event	2 (3.9)	3 (5.8)	4 (7.8)	5 (5.1)	6 (11.8)	5 (5.6)
Serious infectious adverse event	0	1 (1.9)	1 (2.0)	1 (1.0)	3 (5.9)	3 (3.4)
Any adverse event leading to study drug discontinuation	0	2 (3.8)	2 (3.9)	4 (4.1)	5 (9.8)	5 (5.6)
Malignancies	0	0	0	0	0	0
Adverse Events in >7% of patients						
Nausea	1 (2.0)	2 (3.8)	4 (7.8)	2 (2.0)	6 (11.8)	1 (1.1)
Fatigue ^b	2 (3.9)	2 (3.8)	3 (5.9)	5 (5.1)	5 (9.8)	1 (1.1)
Influenza ^b	0	1 (1.9)	2 (3.9)	2 (2.0)	4 (7.8)	1 (1.1)
Nasopharyngitis	6 (11.8)	7 (13.5)	6 (11.8)	13 (13.3)	9 (17.6)	5 (5.6)
Upper Respiratory Tract Infection	2 (3.9)	4 (7.7)	4 (7.8)	7 (7.1)	6 (11.8)	0 (0)
Arthralgia ^b	1 (2.0)	0	3 (5.9)	5 (5.1)	4 (7.8)	0 (0)
Headache	2 (3.9)	7 (13.5)	8 (15.7)	9 (9.2)	10 (19.6)	5 (5.6)
Cough ^b	0	1 (1.9)	3 (5.9)	1 (1.0)	5 (9.8)	2 (2.2)
Hidradenitis	6 (11.8)	7 (13.5)	4 (7.8)	17 (17.3)	12 (23.5)	6 (6.7)
Vomiting	3 (5.9)	2 (3.8)	1 (2.0)			

a: AEs after dose escalation were not included in the eow groups.

b: Occurred in <7% of patients in all treatment groups during Period 1.

This Phase 2 study demonstrated the efficacy and tolerability of adalimumab in treating the signs of inflammation, pain, and health-related quality of life impairment and work productivity in patients with hidradenitis suppurativa, and confirmed that hidradenitis suppurativa is mediated, directly or indirectly, in part by pathogenically elevated levels of TNF α .

Significant, dose-dependent improvement in the global assessment of these patients was achieved at the Week 16 primary endpoint, with evidence of improvement in important secondary endpoints at earlier time points. Patients across the spectrum of severity benefited, and patients with less severe and more reversible (*e.g.*, less scarring) disease at baseline (Hurley Stage I or II) were more likely to benefit. Results from the placebo-controlled portion of the study showed that benefit is more consistent and stable with weekly than every other week dosing; this finding was corroborated by the outcomes in the open-label portion of the study, in which the majority of patients escalated their dosing frequency to weekly because of sub-optimal response to every other week dosing and in which dose escalation improved treatment outcomes. A larger treatment effect was noted for improvement of inflammatory nodules and draining fistulas than for improvement of abscesses.

In addition to evaluating therapeutic effects, this study corroborated and expanded the knowledge of several important epidemiologic characteristics of the HS population. Consistent with previous epidemiologic studies of risk factors associated with HS, the majority of the study patient population was female, their average weight exceeded 90 kg, and the majority were smokers. The observation that approximately 20% of the study patients, who were mostly from the United States, were African American is important, as it supports the impression of an increased prevalence of this disease in this population that is subject to multiple health disparities. Moreover, the proportion of non-white patients in the study population may be an underestimate of the proportion in the general population, as numerous studies have demonstrated a reduced willingness by non-white patients to participate in clinical trials.

Additionally, the substantial baseline level of pain medication used (particularly of opioid use by 10% of the enrollees), as well as the health-related quality of life and work productivity impairment for these patients, was striking, underscoring the debilitating nature of hidradenitis suppurativa and its significant unmet medical need.

The mean baseline DLQI score of 15 in this study was 3.6 points higher (worse) than the mean baseline DLQI of moderate-severe psoriasis patients enrolled in an adalimumab Phase III psoriasis trial; the mean baseline total work productivity impairment was twice as great as that of moderate-severe psoriasis patients enrolled in an adalimumab Phase III trial; and the mean baseline PHQ-9 score of 9.5 indicated moderate depression.

Conversely, treatment with weekly adalimumab resulted in significant and clinically relevant improvement in patient-reported outcomes: the 6-point mean improvement in DLQI between Weeks 0 and 16 exceeded the minimum clinically important difference (MCID) for DLQI (2.3-5.7 points); a significantly higher proportion of patients receiving weekly dosing achieved the clinically meaningful reduction in pain; the mean 3.8-point reduction (improvement) in PHQ-9 scores from Week 0 to 16 exceeded the MCID ($\frac{1}{2}$ of the baseline standard deviation for PHQ-9); and adalimumab ew patients achieved statistically significantly ($p < 0.001$) greater improvement compared to placebo by 20.34 units in TWPI, exceeding the MCID ($\frac{1}{2}$ of the baseline standard deviation).

Adalimumab was well tolerated in this study. The proportion of patients with infectious adverse events were similar in both active-treatment groups and the placebo group during the 16-week double blind period, and the proportion of patients who experienced adverse events leading to study discontinuation or serious adverse events was low. The adverse event pattern did not change through 52 weeks of therapy. The safety profile was similar for patients receiving adalimumab weekly therapy compared to those receiving every other week therapy. Given the apparent increase in efficacy achieved with the higher dose, the risk-benefit balance appears to favor weekly dosing.

In conclusion, weekly adalimumab therapy was effective for improvement of moderate to severe HS for up to 52 weeks. A decline in response rate was seen following switch from ew to eow dosing during Period 2. Dose escalation to ew dosing resulted in improved efficacy. The proportion of patients experiencing serious adverse events was low through 52 weeks of treatment.

In summary, results described above show that adalimumab is the first systemic therapy to have demonstrated significant efficacy in patients with moderate to severe hidradenitis suppurativa in achieving control of objective signs of disease and in reducing pain. Serious adverse event rates associated with the treatment were low, and were similar across all treatment groups.

Example 3 Adalimumab Is Effective at Treating Subpopulations with Hidradenitis Suppurativa (HS)

This example provides a subanalysis of the 52-week Phase II study described above in Examples 1 and 2. Specifically, the following example examines the
5 adalimumab response across subgroups of patients with moderate to severe hidradenitis suppurativa.

The goal of this subanalysis was to assess the relationship between age, sex, race, median weight, and current smoking status, and clinical efficacy in a clinical trial of adalimumab (ADA) in hidradenitis suppurativa (HS).

10 The methods of the study are described above in Examples 1 and 2, and are reiterated here. Moderate to severe HS patients (patients with HS-Physician's Global Assessment [HS-PGA] $\geq$ moderate) were randomized 1:1:1 to ADA 40 mg weekly (ew) (after 160 mg at Week 0, 80 mg at Week 2), ADA 40 mg every other week (eow) (after 80 mg at Week 0), or placebo (pbo) in a 52-week, Phase II clinical trial. The primary
15 endpoint was the proportion of patients at Week 16 with a Clinical Response (defined as achieving an HS-PGA score of clear, minimal, or mild, with at least a two grade improvement relative to baseline) at Week 16.

For this analysis, data from subsets of the Intention-to-Treat population were grouped based on patient age, sex, race, median weight, and current smoking status, for a
20 post hoc analysis of the effect of these factors on clinical efficacy. Non-responder imputation was used for missing data.

Overall Week 16 Clinical Response rates for pbo, eow, and ew patients were 3.9% (n=2/51), 9.6% (n=5/52), and 17.6% (n=9/51) (P=0.025, ew vs. pbo).

Age: At Week 16, Clinical Responses were achieved as follows in patients <40
25 years: pbo 5.9% (2/34), eow 6.5% (2/31) and ew 12.1% (4/33); for patients $\geq$ 40 years: pbo 0% (0/17), eow 15.0% (3/21) and ew 29.4% (5/18; P=0.047, ew vs. pbo).

Sex: At Week 16, Clinical Responses were achieved as follows: for males: pbo 0% (0/15), eow 14.3% (2/14), and ew 13.3% (2/15); for females: pbo 5.6% (2/36), eow 7.9% (3/38), and ew 19.4% (7/36).

Race: At Week 16, Clinical Responses were achieved as follows: for whites: pbo 2.6% (1/38), eow 7.9% (3/38), and ew 21.6% (8/37; P=0.016, ew vs. pbo); for non-whites: pbo 7.7% (1/13), eow 14.3% (2/14), and ew 7.1% (1/14).

Body weight: At Week 16, Clinical Responses were achieved as follows: for
5 patients <96.5 kg: pbo 7.7% (2/26), eow 4.5% (1/30), and ew 17.2% (5/29); for patients >96.5 kg: pbo 0% (0/25), eow 13.3% (4/30; P=0.048, eow vs. pbo), and ew 18.2% (4/22; P=0.013, ew vs. pbo).

Current smoking status: At Week 16, Clinical Responses were achieved as follows: for current smokers: pbo 3.4% (1/29), eow 11.5% (3/26), and ew 23.3% (7/30; P=0.041, ew vs. pbo); and for current non-smokers: pbo 4.5% (1/22), eow 7.7% (2/26),
10 and ew 9.5% (2/21).

The results show that weekly adalimumab therapy was effective for the treatment of moderate to severe HS at Week 16. In this post-hoc analysis, the greatest Clinical Response rates were seen in patients greater than 40 years old, females, whites, and
15 current smokers. No apparent differences were seen between body weight subgroups in this analysis.

Example 4 Impact of Weight and Body Mass Index on High-Sensitivity C-Reactive Protein Response to Adalimumab in Hidradenitis Suppurativa Patients
20

This example provides a subanalysis of the 52-week Phase II study described above in Examples 1 and 2. The goal of the subanalysis was to determine the impact of adalimumab treatment on high-sensitivity C-reactive protein (hs-CRP) in moderate to severe hidradenitis suppurativa (HS) patients, and to determine whether weight or BMI
25 had an effect on hsCRP response to adalimumab.

The following example is based on data from a Phase II, 52-week (wk) trial, where the initial 16 wk portion was double-blind and placebo-controlled (see Examples 1 and 2). Patients with moderate to severe HS (HS-Physician's Global Assessment [HS-PGA] $\geq$ moderate) were randomized 1:1:1 to adalimumab 40 mg weekly (ew) (after 160
30 mg at Wk 0 and 80 mg at Wk 2), adalimumab 40 mg every other wk (eow) (after 80 mg

at Wk 0), or placebo (pbo). The primary efficacy variable was the proportion of patients who achieved Clinical Response, defined as an HS-PGA score of clear, minimal, or mild, with at least a two grade improvement relative to baseline, at Wk 16.

This post-hoc analysis assessed mean change in hs-CRP concentration from baseline to Wk 16 by treatment group; patients were further grouped by weight ≤ 100 kg and >100 kg and BMI <25 , ≥ 25 - <30 , and ≥ 30 . Missing data were imputed as non-response.

154 patients were enrolled: 51 pbo/52 eow/51 ew. At Week 16, the proportion of patients achieving Clinical Response was 3.9%/9.6%/17.6% for pbo/eow/ew groups ($P=.025$, ew vs. pbo). At baseline, mean hs-CRP concentrations were: 13.8/17.8/22.2 mg/L for pbo/eow/ew groups ($n=36/40/35$). At Week 16, mean change from baseline in hs-CRP was 5.5/-1.6/-3.2 mg/L for pbo/eow/ew patients ($P=0.034$, ew vs. pbo). For pbo/ew/eow patients ≤ 100 kg and >100 kg, mean change from baseline to Wk 16 was 8.3/-4.0/-4.4 mg/L ($n=20/19/24$; $P=0.020$, eow vs. pbo; $P=0.011$, ew vs. pbo) and 2.0/0.5/-0.7 mg/L ($n=16/21/11$). For pbo/eow/ew patients with BMI $<25/\geq 25$ - $<30/\geq 30$, mean change from baseline to Wk 16 was 4.7/-1.0/-6.0 ($n=6/4/6$); 14.2/-11.6/4.8 ($n=3/8/9$; $P=0.018$, eow vs. pbo); 4.7/1.2/-6.0 ($n=27/28/20$).

The conclusion of the subanalysis was that weekly adalimumab therapy was effective in reducing hs-CRP concentrations in moderate to severe HS patients regardless of weight or BMI. For weight, the greatest hs-CRP reductions occurred in patients ≤ 100 kg receiving weekly dosing; for BMI, comparable hs-CRP reductions were seen in normal and obese patients receiving weekly dosing.

Example 5. HiSCR Scoring System for Treatment of Hidradenitis Suppurativa (HS)

The following example describes a new scoring system for assessing improvements in a subject having HS. The new scoring system, called Hidradenitis Suppurativa Clinical Response (HiSCR), was developed based on clinical trial data (described above) obtained using the anti-TNF α antibody adalimumab for the treatment of HS.

HiSCR is defined as at least a 50% reduction in the total inflammatory lesion (abscess and inflammatory nodule) count (AN count) relative to baseline, with no increase in abscess count and no increase in draining fistula count. Treated subjects are declared clinical responders only if they experience at least 50% reduction in total inflammatory lesion (abscess + inflammatory nodule) count, and also no increase in abscess count and no increase in draining fistula count.

HiSCR was applied to the raw clinical data from the Phase II clinical trial described in Examples 1 - 4 to assess the efficacy of adalimumab using HiSCR. The results from this analysis are shown in Table 14.

Table 14 shows the proportion of patients achieving HiSCR, *i.e.*, at least 50% reduction in inflammatory lesion count with no increase in abscess count and no increase in draining fistula count, at Weeks 2, 4, 8, 12 and 16, among those patients who were Hurley Stage II or III, and who had more than two abscesses or inflammatory nodules, but less than 20 draining fistulas at baseline. The response rates observed at week 12 for HiSCR were about 61% and about 16% in the adalimumab qwk group and placebo group, respectively.

Table 14. Proportion (Percentage) of Patients Achieving HiSCR
at Weeks 2, 4, 8, 12 and 16

Patient Group	Week 2	Week 4	Week 8	Week 12	Week 16
Placebo	8.1	13.5	32.4	16.2	21.6
EOW	21.1	31.6 ^{\$}	31.6	28.9	31.6
QWK	41.7 [#]	47.2 [#]	50.0	61.1 ^{#, &}	55.6 ^{#, &}

Excluding subjects with ≤ 2 abscesses/inflammatory nodules and >20 draining fistulas at baseline; N=37 for PBO, N=38 for EOW, N=36 for QWK, ITT, NRI.

#: $p < 0.05$, qwk vs. pbo; \$: $p < 0.05$, eow vs. pbo; &: $p < 0.05$, qwk vs. eow.

HiSCR may be used to assess clinical efficacy, as the results described in Table 14 showed that improved efficacy would be seen at all study visits for patients treated with qwk dosing compared with those treated with eow dosing. At Week 12, the number of treatments needed to achieve clinical success is projected to be 2.2 with qwk dosing, compared with 7.9 with eow dosing.

REFERENCES

1. Esmann S, Jemec GB. Psychosocial Impact of Hidradenitis Suppurativa: A Qualitative Study. *Acta Derm Venereol* 2011.
2. Jemec GB, Wendelboe P. Topical clindamycin versus systemic tetracycline in the treatment of hidradenitis suppurativa. *J Am Acad Dermatol* 1998;39:971-4.
3. Kurzen H, Kurokawa I, Jemec GB, et al. What causes hidradenitis suppurativa? *Exp Dermatol* 2008;17:455-6; discussion 7-72.
4. Shah N. Hidradenitis suppurativa: a treatment challenge. *Am Fam Physician* 2005;72:1547-52.
5. Fimmel S, Zouboulis CC. Comorbidities of hidradenitis suppurativa (acne inversa). *Dermato-endocrinology* 2010;2:9-16.
6. van der Zee HH, van der Woude CJ, Florencia EF, Prens EP. Hidradenitis suppurativa and inflammatory bowel disease: are they associated? Results of a pilot study. *Br J Dermatol* 2010;162:195-7.
7. Revuz JE, Canoui-Poitaine F, Wolkenstein P, et al. Prevalence and factors associated with hidradenitis suppurativa: results from two case-control studies. *J Am Acad Dermatol* 2008;59:596-601.
8. Jemec GB. Hidradenitis suppurativa. *J Cutan Med Surg* 2003;7:47-56.
9. Zouboulis CC. Disorders of the Apocrine Sweat Glands. In: Wolff K, Goldsmith LA, Katz SI, Gilchrest BA, Paller AS, Leffell DJ, eds. *Fitzpatrick's Dermatology in General Medicine*, New York City: McGraw Hill; 2010.
10. König A, Lehmann C, Rompel R, Happle R. Cigarette smoking as a triggering factor of hidradenitis suppurativa. *Dermatology* 1999;198:261-4.
11. Sartorius K, Emtestam L, Jemec GB, Lapins J. Objective scoring of hidradenitis suppurativa reflecting the role of tobacco smoking and obesity. *Br J Dermatol* 2009;161:831-9.
12. von der Werth JM, Jemec GB. Morbidity in patients with hidradenitis suppurativa. *Br J Dermatol* 2001;144:809-13.

13. Wolkenstein P, Loundou A, Barrau K, Auquier P, Revuz J. Quality of life impairment in hidradenitis suppurativa: a study of 61 cases. *J Am Acad Dermatol* 2007;56:621-3.
14. Using PHQ-9 Diagnosis and Score for Initial Treatment Selection. (Accessed
5 February 22, 2010, at
15. Matusiak L, Bieniek A, Szepietowski JC. Psychophysical aspects of hidradenitis suppurativa. *Acta Derm Venereol* 2010;90:264-8.
16. Alikhan A, Lynch PJ, Eisen DB. Hidradenitis suppurativa: a comprehensive
10 review. *J Am Acad Dermatol* 2009;60:539-61; quiz 62-3.
17. Plewig G. Acne and Rosacea. In: Burgdorf WHC, Plewig G, Wolff HH, Landthaler M, eds. *Braun-Falco's Dermatology*, 3rd ed. Heidelberg: Springer Medlzin Verlag 2009:1002-4.
18. Clemmensen OJ. Topical treatment of hidradenitis suppurativa with clindamycin.
15 *Int J Dermatol* 1983;22:325-8.
19. Mortimer PS, Dawber RP, Gales MA, Moore RA. A double-blind controlled cross-over trial of cyproterone acetate in females with hidradenitis suppurativa. *Br J Dermatol* 1986;115:263-8.
20. van der Zee HH, de Ruitter L, van den Broecke DG, Dik WA, Laman JD, Prens
20 EP. Elevated levels of TNF-alpha, IL-1beta and IL-10 in hidradenitis suppurativa skin; a rationale for targeting TNF-alpha and IL-1beta. *Br J Dermatol* 2011.
21. Lebwohl B, Sapadin AN. Infliximab for the treatment of hidradenitis suppurativa. *J Am Acad Dermatol* 2003;49:S275-6.
22. Rosi YL, Lowe L, Kang S. Treatment of hidradenitis suppurativa with infliximab
25 in a patient with Crohn's disease. *J Dermatolog Treat* 2005;16:58-61.
23. Haslund P, Lee RA, Jemec GB. Treatment of hidradenitis suppurativa with tumour necrosis factor-alpha inhibitors. *Acta Derm Venereol* 2009;89:595-600.
24. Harde V, Mrowietz U. Treatment of severe recalcitrant hidradenitis suppurativa with adalimumab. *J Dtsch Dermatol Ges* 2009;7:139-41.

25. Adams DR, Yankura JA, Fogelberg AC, Anderson BE. Treatment of hidradenitis suppurativa with etanercept injection. *Arch Dermatol* 2010;146:501-4.
26. Grant A, Gonzalez T, Montgomery MO, Cardenas V, Kerdel FA. Infliximab therapy for patients with moderate to severe hidradenitis suppurativa: a randomized, double-blind, placebo-controlled crossover trial. *J Am Acad Dermatol* 2010;62:205-17.
27. Miller I, Lynggaard CD, Lophaven S, Zachariae C, Dufour DN, Jemec GB. A double blind Placebo Controlled randomised Trial of Adalimumab in the treatment of Hidradenitis Suppurativa. *Br J Dermatol* 2011.
- 10 28. Hurley HJ. Axillary hyperhidrosis, apocrine bromhidrosis, hidradenitis suppurativa, and familial benign pemphigus: surgical approach. In: Roenigk RK, Roenigk HH, eds. *Dermatologic Surgery*. New York: Dekker; 1989:729.
29. Manton KG, Patrick CH, Johnson KW. Health differentials between blacks and whites: recent trends in mortality and morbidity. *Milbank Q* 1987;65 Suppl 1:129-99.
- 15 30. Corbie-Smith G, Thomas SB, Williams MV, Moody-Ayers S. Attitudes and beliefs of African Americans toward participation in medical research. *J Gen Intern Med* 1999;14:537-46.
31. Shavers VL, Lynch CF, Burmeister LF. Racial differences in factors that influence the willingness to participate in medical research studies. *Ann Epidemiol* 2002;12:248-56.
- 20 32. Revicki DA, Willian MK, Menter A, Saurat JH, Harnam N, Kaul M. Relationship between clinical response to therapy and health-related quality of life outcomes in patients with moderate to severe plaque psoriasis. *Dermatology* 2008;216:260-70.
- 25 33. Kimball AB, Yu AP, Signorovitch J, et al. The effects of adalimumab treatment and psoriasis severity on self-reported work productivity and activity impairment for patients with moderate to severe psoriasis,. accepted.

34. Shiklar R, Willian MK, Okun MM, Thompson CS, Revicki DA. The validity and responsiveness of three quality of life measures in the assessment of psoriasis patients: results of a phase II study. *Health Qual Life Outcomes* 2006;4:71.
35. Farrar JT, Berlin JA, Strom BL. Clinically important changes in acute pain
5 outcome measures: a validation study. *J Pain Symptom Manage* 2003;25:406-11.
36. Norman GR, Sloan JA, Wyrwich KW. Interpretation of changes in health-related quality of life: the remarkable universality of half a standard deviation. *Med Care* 2003;41:582-92.

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention
5 described herein. Such equivalents are intended to be encompassed by the following claims.

The invention claimed is:

1. Use of adalimumab, in multiple doses for treating moderate to severe hidradenitis suppurativa (HS) in an adult, wherein the multiple doses comprise:
 - a first loading dose of 160 mg of adalimumab for subcutaneous administration to the subject at week 0;
 - a second loading dose of 80 mg of adalimumab for subcutaneous administration to the subject at week 2; and
 - a weekly maintenance dose of 40 mg of adalimumab for subcutaneous administration to the adult starting at week 4, wherein said multiple doses are not subject to any discretionary adjustment by a physician or medical practitioner.
2. The use of claim 1, further comprising use of a biweekly maintenance dose of 40 mg of adalimumab for subcutaneous administration to the adult starting at week 16, wherein said weekly administration of the maintenance dose stops after week 15.
3. The use of any one of claims 1 to 2 wherein said treating decreases the number of inflammatory lesions in the adult.
4. The use of any one of claims 1 to 3 wherein said treating prevents worsening of abscesses in the adult.
5. The use of any one of claims 1 to 4 wherein said treating prevents worsening of draining fistulas in the adult.
6. The use of any one of claims 1 to 5, further comprising use of antibiotics in the adult.
7. The use of any one of claims 1 to 6, wherein the adult had an inadequate response to oral antibiotics prior to said treating.

Figure 1




Figure 2






Figure 3





Up to 28 days 16 weeks 36 weeks

