

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2014/001836 A1

(43) International Publication Date
3 January 2014 (03.01.2014)

(51) International Patent Classification:

A01N 37/02 (2006.01) *A01N 25/34* (2006.01)
A01N 37/06 (2006.01) *A01P 19/00* (2006.01)
A01N 37/36 (2006.01)

(21) International Application Number:

PCT/IB2012/001341

(22) International Filing Date:

25 June 2012 (25.06.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(71) Applicant (for all designated States except US): **INSTITUT DE RECHERCHE EN SEMIOCHIMIE ET ETHOLOGIE APPLIQUEE** [FR/FR]; Le Rieu Neuf, F-84490 Saint Saturnin d'Apt (FR).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **PAGEAT, Patrick** [FR/FR]; Le Fénier du Plan, Route de Saint-Saturnin, F-84400 Apt (FR).

(74) Agents: **GUTMANN, Ernest** et al.; 3, rue Auber, F-75009 Paris (FR).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: FELINE SCRATCH MARKING SEMIOCHEMICALS

(57) Abstract: Semiochemical compositions comprising a feline scratch marking pheromone, analogues of these semiochemical compositions and a method to prevent cats from scratching in particular places and on multiple objects are described.



WO 2014/001836 A1

FELINE SCRATCH MARKING SEMIOCHEMICALS

1. Field of the Invention

The present invention relates to semiochemical compositions comprising a feline scratch marking semiochemical analogues of these semiochemical compositions and a method to prevent cats from scratching in particular places and on multiple objects.

2. Background and Prior Art

Cats are known for their scratching behavior, which is seen by most cat owners as one of the most unacceptable nuisance in this species. Most cat owners misunderstand this behavior interpreting it as a sole means for controlling the length of their claws. However, cats usually scratch for three reasons: to hone or sharpen their claws, which are the cat's chief offensive and defensive weapons, to stretch and tone their muscles and to mark their territory.

Felines actually have sweat glands in their paw pads and when they scratch against furniture, curtains, wallpaper, corners of upholstered furniture and other objects they are actually depositing their own scent onto those objects. Vertical scratches also leave a visible mark for other cats to see as well as to smell. Thus, there are actually two different marks that cats make when they scratch various objects; a visible marking and an odorous marking.

Currently to rid felines of scratching various objects it is suggested that the feline be trained to scratch in particular areas by providing scratching posts, scratching pads or cat trees. Other alternatives are herbal sprays that repel the felines from objects due to their scent or double-sided tape. Herbal sprays replace the odorous territorial markers left behind by the scratching to discourage repeat scratching in the same place. Since cats' paws are extremely sensitive to touch, using sticky surfaces, such as double-sided tape, is exceptionally annoying to cats so that they avoid touching those surfaces.

Soft plastic nail caps called Softpaws[®] are also available which should be applied to the claws every 4 to 6 weeks and keep the nails blunt. However, the

application of these plastic nail caps is difficult, since they are applied with glue. Usually they are applied by an expert or veterinarian.

Electric carpets are yet another solution. These carpets are placed by the objects that are being scratched and discharge a strong electric volts but weak
5 amperage in the footpads of the cats that walk on them.

Yet another alternative is to dull the cat's claws by trimming their nails as a part of a general routine or declawing the cat. However these solutions are not really recommended by veterinarian's since it may lead to other behavioral problems in cats.

10 U.S. Patent No. 5,709,863 discloses compositions containing an emulsion and a mixture of fatty acids or derivatives thereof and a compound of vegetal origin that has an attractive effect on cats for preventing cats from urinating in a marked spot or to prevent anxiety in cats. These compositions comprise an emulsion of oleic acid, azelaic acid, palmitic acid, pimelic acid and a compound of vegetal origin that has an
15 attractive effect on cats. However, this patent does not solve the problem of preventing cats from scratching on multiple objects.

Thus, there is a need in the art to address the problem of feline's scratching on multiple household and other objects.

It is object of the present invention to provide an alternative treatment for
20 preventing cats to scratch on multiple objects.

It is another object to provide compositions that attract cats to a particular object which the cat can scratch.

It is another object to provide a scratching post that attracts cats and thus prevents them from scratching other multiple objects.

25 It is yet another object to provide a method of attracting cats to a particular object so that they can scratch that object, while avoiding other objects.

These and other objects are achieved by the present invention as evidenced by the summary of the invention, description of the preferred embodiments and the claims.

SUMMARY OF THE PRESENT INVENTION

In one of the composition aspects, the present invention provides a
5 semiochemical composition comprising a secretion obtained from the interdigital
paws of a cat.

In another aspect a semiochemical composition comprising valeric acid, lactic
acid and linoleic acid and/or derivatives thereof and/or isomers thereof is provided.

10 A semiochemical composition comprising between about 3.5% to 10.5%
(w%/w%) of valeric acid, between about 5.5% to 15.5% (w%/w%) of lactic acid and
between 68.5% to 83.5 (w%/w%) of linoleic acid and/or derivatives thereof and/or
isomers thereof is yet another aspect of the invention.

15 In another aspect a semiochemical composition comprising between about
3.5% to 10.5% (w%/w%) of valeric acid, between about 5.5% to 15.5% (w%/w%) of
lactic acid, between about 48.4 % to 58.9 % (w%/w%) of linoleic acid and between
about 20.1% to 24.5 % (w%/w%) of oleic acid and/or derivatives thereof and/or
20 isomers thereof is provided.

The present invention provides a semiochemical composition analogue
comprising 10% to 20% (w%/w%) of valeric acid, lactic acid and linoleic acid and/or
derivatives thereof and/or isomers thereof and 60% to 98% (w%/w%) of linoleic acid.

25 A semiochemical composition analogue comprising 10% to 20% (w%/w%) of a
composition comprising between about 3.5% to 10.5% (w%/w%) of valeric acid,
between about 5.5% to 15.5% (w%/w%) of lactic acid and between 68.5% to 83.5
(w%/w%) of linoleic acid and/or derivatives thereof and/or isomers thereof and
between 60% to 98% (w%/w%) of linoleic acid is yet another embodiment of the
30 present invention.

In another embodiment a semiochemical composition comprising 20% of a composition comprising between about 3.5% to 10.5% (w%/w%) of valeric acid, between about 5.5% to 15.5% (w%/w%) of lactic acid and between 68.5% to 83.5 (w%/w%) of linoleic acid and/or derivatives thereof and/or isomers thereof and 75%
5 linoleic acid.

Solutions of the above described semiochemical compositions and analogues thereof are also one aspect of the present invention.

The semiochemical compositions can further comprise a solvent and optionally at least one cat attractant and colorants.

10 A method for preventing a cat from scratching on multiple objects, said method comprising placing at least one semiochemical composition and/or semiochemical composition analogues and/or derivatives thereof and/or isomers thereof on an object that a cat uses for scratching.

A cat scratching post, board, carpet, trees and toys comprising the
15 semiochemical compositions or semiochemical composition analogues, and/or derivatives thereof and/or isomers thereof as described herein, are also an aspect of the present invention.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a X bar graph with joined points showing the average of the NPL
20 results using the cat scratching semiochemical analogues. The limits of control are 3 Sigma.

Fig. 2 is a X bar graph with joined points showing the average of the DBFL results using the cat scratching semiochemical analogues. The limits of control are 3 Sigma.

25 Fig. 3 is a X bar graph with joined points showing the average of the SOA results using the cat scratching semiochemical analogues. The limits of control are 3 Sigma.

Fig. 4 is a X bar graph with joined points showing the average of the NPL results using the cat scratching semiochemical analogues. The limits of control are 3

Sigma. OL stands for the object that the cat habitually lacerated. The mean value is 9.583, the inferior value from the mean value is 7.576, while the superior value from the mean value is 11.591.

Fig. 5 is a X bar graph with joined points showing the average of the DBFL results using the cat scratching semiochemical analogues. The limits of control are 3 Sigma. OL stands for the object that the cat habitually lacerated. The mean value is 2.665, the inferior value from the mean value is 1.194, while the superior value from the mean value is 4.137.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

As used herein "semiochemical" means a chemical emitted by a plant or an animal that evokes a behavioral or physiological response in another organism. When the semiochemical affects an individual of the same species, it is called a pheromone. When the semiochemical affects an individual of a different species, it is called an allelochemical.

By "enhancer composition" is meant an active semiochemical composition or analogues thereof that is species-specific in cats and which can be used to enhance or act synergistically with the basic semiochemical composition or analogues thereof and/or derivatives thereof and/or isomers thereof to increase the effectiveness in felines of the basic semiochemical composition or analogues thereof.

The terms cats and felines are used interchangeably herein. "Feline" means of or belonging to the family Felidae, which includes lions, tigers, jaguars and wild and domestic cats.

The term "object or objects" means any physical thing that can be seen and touched. Examples of objects includes any object that a feline may scratch including, but limited to, rugs, furniture, carpets, drapes, upholstered corners of couches and chairs, wall papered surfaces, cardboard storage boxes, wood trim on doors and walls. These are only a very few examples defined to encompass the term objects that is used herein.

As used herein the term "analogue" means a group of chemical compounds similar in structure but different in respect to an elemental composition.

"Derivatives of fatty acids" as used herein includes fatty acid esters, salts, alcohols, ketones, ethers and amides.

By "isomers" is meant herein a chemical species with the same number and types of atoms as another chemical species and include conjugated fatty acids, which are positional and geometric isomers. Due to the configuration of the double bond in some fatty acids, they can take two forms. The cis form, in which both of the hydrogen atoms are on the same side of the chain and the trans form in which the hydrogen atoms are on opposite sides. For example, conjugated linoleic acid is a mixture of positional and geometric isomers of linoleic acid involving double bonds at positions 8 and 10, 9 and 11, 10 and 12 or 11 and 13. Each of these positional diene isomers can occur in cis-trans, trans-cis, cis-cis or trans-trans geometrical configurations.

The term "consisting essentially of" means that the semiochemical compositions or analogues thereof and/or derivatives thereof and/or isomers thereof can contain other ingredients that do not affect the semiochemical properties of the compositions or analogues thereof.

More specifically, the present invention relates to semiochemical compositions derived from the interdigital area of the cat's paws that is used to attract the cat's to a specific area for scratching.

Analogues, derivatives and isomers of the semiochemical compositions, described herein, are also encompassed by the present invention.

The compositions and analogues of the present invention are based on semiochemical compositions and made up of volatile molecules, the essential components of these molecules being fatty acids and methyl esters of the fatty acids.

In one aspect, the composition of the present invention comprises a mixture of three fatty acids; namely valeric acid, lactic acid and linoleic acid and/or derivatives thereof and/or isomers thereof which represent between about 3.5% to 83.5% (w%/w%) of the total composition. The remaining ingredients being compounds having an attractive affect on cats, colorants and solvents.

In another aspect the semiochemical composition comprises a mixture of propionic acid, cyclohexylacetic acid and cyclopentylpropionic acid and/or derivatives thereof and/or isomers thereof, which represents between 11.5% to 45.5% (w%/w%) of the total composition. This composition can also contain compounds that have an attractive affect on cats, colorants and solvents.

In yet another aspect the semiochemical composition comprises a mixture of valeric acid, lactic acid, linoleic acid and oleic acid and/or derivatives thereof and/or isomers thereof, which represents between 5.5% to 58.9% (w%/w%) of the total composition.

The semiochemical analogues of the present invention comprise 10% to 20% (w%/w%) of a mixture of three fatty acids; namely valeric acid, lactic acid and linoleic acid and/or derivatives thereof and/or isomers thereof and from 60% to 98% (w%/w%) of linoleic acid.

The semiochemical analogues or semiochemical compositions or derivatives thereof and/or isomers thereof can be attached to a chemical carrier provided that the bioactive structure of the fatty acids is preserved. Such carrier molecules include resins, liposomes, crown compounds, proteins, polymers and the like.

The fatty acids can be in their pure form, i.e., as a free fatty acid as well as their derivatives forms such as esters of fatty acids, salts of fatty acids, alcohols of fatty acids, ketones of fatty acids, ethers of fatty acids, and amides of fatty acids.

Isomers of the fatty acids in the semiochemical compositions and/or derivatives and/or analogues thereof, described herein, can also be used in the formulations of the semiochemical compositions.

More specifically, the semiochemical compositions of the present invention comprise between about 3.5% to 10.5% (w%/w%) valeric acid, between about 5.5% to 15.5% (w%/w%) lactic acid and between 68.5% to 83.5% (w%/w%) linoleic acid and/or derivatives thereof and/or isomers thereof.

A solution of a semiochemical compositions comprising between about 3.5% to 10.5% (w%/w%) valeric acid, between about 5.5% to 15.5% (w%/w%) lactic acid

and between 68.5% to 83.5%(w%/w%) linoleic acid and/or derivatives thereof and/or isomers thereof is also encompassed by the present invention.

In one aspect the semiochemical compositions of the present invention comprise between about 3.5% to 10.5% (w%/w%) valeric acid, between about 5.5%
5 to 15.5% (w%/w%) lactic acid and between 68.5% to 83.5%(w%/w%) linoleic acid and/or isomers thereof and/or derivatives thereof that are esters, salts, alcohols, ketones , ethers and amides of valeric acid, lactic acid and linoleic acid. These derivatives are in the same percentages as their fatty acid compositions.

A solution of the semiochemical compositions of the present invention
10 comprise between about 3.5% to 10.5% (w%/w%) valeric acid, between about 5.5% to 15.5% (w%/w%) lactic acid and between 68.5% to 83.5%(w%/w%) linoleic acid and/or isomers thereof and/or derivatives thereof that are esters, salts, alcohols, ketones , ethers and amides of valeric acid, lactic acid and linoleic acid. These derivatives are in the same percentages as their fatty acid compositions.

15 Another embodiment of the present invention is a semiochemical composition comprising between about 3.5% to 10.5% (w%/w%)of valeric acid, between about 5.5% to 15.5% (w%/w%)of lactic acid, between about 48.4 % to 58.9 % (w%/w%) of linoleic acid and between about 20.1% to 24.5 % (w%/w%) of oleic acid and/or derivatives thereof and/or isomers thereof.

20 A solution of the semiochemical composition comprising between about 3.5% to 10.5% (w%/w%)of valeric acid, between about 5.5% to 15.5% (w%/w%)of lactic acid, between about 48.4 % to 58.9 % (w%/w%) of linoleic acid and between about 20.1% to 24.5 % (w%/w%) of oleic acid and/or derivatives thereof and/or isomers thereof is part of the present invention.

25 Another embodiment of the present invention is a semiochemical composition comprising between about 3.5% to 10.5% (w%/w%)of valeric acid, between about 5.5% to 15.5% (w%/w%)of lactic acid, between about 48.4 % to 58.9 % (w%/w%) of linoleic acid and between about 20.1% to 24.5 % (w%/w%) of oleic acid and/or isomers thereof and derivatives thereof that are esters, salts, alcohols, ketones,
30 ethers and amides of valeric acid, lactic acid, linoleic acid and oleic acid. These derivatives are in the same percentage as their fatty acid counterparts.

In yet another aspect of the present invention a solution of the semiochemical composition comprising between about 3.5% to 10.5% (w%/w%) of valeric acid, between about 5.5% to 15.5% (w%/w%) of lactic acid, between about 48.4 % to 58.9 % (w%/w%) of linoleic acid and between about 20.1% to 24.5 % (w%/w%) of oleic acid and/or isomers thereof and/or derivatives thereof that are esters, salts, alcohols, ketones, ethers and amides of valeric acid, lactic acid, linoleic acid and oleic acid. These derivatives are in the same percentage as their fatty acid counterparts.

In another aspect the semiochemical compositions of the present invention comprises a composition comprising about 36.5% to 43.5% (w%/w%) propionic acid, about 11.5% to 26.5% (w%/w%) cyclohexylacetic acid and about 33.5% to 45.5% (w%/w%) cyclopentylpropionic acid and/or derivatives thereof and/or isomers thereof. These derivatives are in the same percentage as their fatty acid counterparts.

A solution of the semiochemical composition comprising about 36.5% to 43.5% (w%/w%) propionic acid, about 11.5% to 26.5% (w%/w%) cyclohexylacetic acid and about 33.5% to 45.5% (w%/w%) cyclopentylpropionic acid and/or derivatives thereof and/or isomers thereof is another aspect of the invention. These derivatives are in the same percentage as their fatty acid counterparts.

In yet another aspect the semiochemical compositions of the present invention comprise about 36.5% to 43.5% (w%/w%) propionic acid, about 11.5% to 26.5% (w%/w%) cyclohexylacetic acid and about 33.5% to 45.5% (w%/w%) cyclopentylpropionic acid and/or isomers thereof and/or derivatives thereof that are esters, salts, alcohols, ketones, ethers and amides of propionic acid, cyclohexylacetic acid and cyclopentylpropionic acid. These derivatives are in the same percentage as their fatty acid counterparts.

A solution of semiochemical compositions of the present invention comprise about 36.5% to 43.5% (w%/w%) propionic acid, about 11.5% to 26.5% (w%/w%) cyclohexylacetic acid and about 33.5% to 45.5% (w%/w%) cyclopentylpropionic acid and/or isomers thereof and/or derivatives thereof that are esters, salts, alcohols, ketones, ethers and amides of propionic acid, cyclohexylacetic acid and cyclopentylpropionic acid is yet another aspect of the invention. These derivatives are in the same percentage as their fatty acid counterparts.

The present invention provides a semiochemical composition analogue comprising 10% to 20% (w%/w%) of valeric acid, lactic acid and linoleic acid and/or derivatives thereof and/or isomers thereof and 60% to 98% (w%/w%) of linoleic acid.

In yet another aspect analogues of the semiochemical composition comprises
5 between about 3.5% to 10.5% (w%/w%) valeric acid, between about 5.5% to 15.5% (w%/w%) lactic acid and between 68.5% to 83.5%(w%/w%) linoleic acid and/or derivatives thereof and/or isomers thereof in concentrations of 10% to 20% (w%/w%) and between 60% to 98% (w%/w%) of linoleic acid. These derivatives are in the same percentage as their fatty acid counterparts.

10 A solution of the semiochemical composition comprising between about 3.5% to 10.5% (w%/w%) valeric acid, between about 5.5% to 15.5% (w%/w%) lactic acid and between 68.5% to 83.5%(w%/w%) linoleic acid and/or derivatives thereof and/or isomers thereof in concentrations of 10% to 20% (w%/w%) and between 60% to 98% (w%/w%) of linoleic acid is part of the present invention. These derivatives are in the
15 same percentage as their fatty acid counterparts.

In yet another aspect analogues of the semiochemical composition comprises between about 3.5% to 10.5% (w%/w%) valeric acid, between about 5.5% to 15.5% (w%/w%) lactic acid (w%/w%) and between 68.5% to 83.5%(w%/w%) of linoleic acid in concentrations of 10% to 20% and between 60% to 98% (w%/w%) of linoleic acid
20 and/or isomers thereof and/or derivatives thereof that are esters, salts, alcohols, ketones, ethers and amides of valeric acid, lactic acid and linoleic acid. These derivatives are in the same percentage as their fatty acid counterparts.

In another aspect a solution of the semiochemical composition comprises between about 3.5% to 10.5% (w%/w%) valeric acid, between about 5.5% to 15.5% (w%/w%) lactic acid and between 68.5% to 83.5%(w%/w%) linoleic acid in
25 concentrations of 10% to 20% (w%/w%) and between 60% to 98% (w%/w%) of linoleic acid and/or isomers thereof and/or derivatives thereof that are esters, salts, alcohols, ketones, ethers and amides of valeric acid, lactic acid and linoleic acid forms part of the invention.

30 In one embodiment the semiochemical analogues such as SC10-98, which has 10% of the fraction Sc1 (See, Table III below) and 98% linoleic acid; SC10-75,

which has 10%, of the fraction Sc1 (See, Table III below) and 75% linoleic acid; SC10-60, which has 10% of the fraction Sc1 (See, Table III below) and 60% linoleic acid; SC20-98, which has 20% of the fraction Sc1 (See, Table III below) and 98% linoleic acid; SC20-75, which has 20% of the fraction Sc1 (See, Table III below) and 75% linoleic acid; and SC20-60, which has 20% of the fraction Sc1 (See, Table III below) and 60% linoleic acid form another aspect of the semiochemical compositions of the invention.

Solutions of the semiochemical analogues such as SC10-98, which has 10% of the fraction Sc1 (See, Table III below) and 98% linoleic acid; SC10-75, which has 10%, of the fraction Sc1 (See, Table III below) and 75% linoleic acid; SC10-60, which has 10% of the fraction Sc1 (See, Table III below) and 60% linoleic acid; SC20-98, which has 20% of the fraction Sc1 (See, Table III below) and 98% linoleic acid; SC20-75, which has 20% of the fraction Sc1 (See, Table III below) and 75% linoleic acid; and SC20-60, which has 20% of the fraction Sc1 (See, Table III below) and 60% linoleic acid form another aspect of the invention.

The semiochemical compositions described herein can be diluted in various solvents and can be used in various forms. The compositions may be in the form of a solution, aerosol spray, gel, slow release matrix, microencapsulated product, liposomes, nanoparticles and the like.

In a preferred embodiment the fatty acids can be microencapsulated and put into a suspension in alcohol.

An enhancer composition containing between 5% to 35% (w%/w%) can also be added to the semiochemical composition, if desired. This enhancer composition comprises volatile organic compounds and mixtures thereof and includes, but not limited to amines and fatty acids from indolic derivatives, esters of these amines and fatty acids, ketones such as acetone, alcohols, sterols and the like.

Cat attractants can also be optionally added to the composition. These include catnip (*Nepeta Cataria*), essential oil of catnip, tatarian honeysuckle (*Lonicera tartarica*), Valerian (*Valeriana officinalis*), silver vine/ matatabi (*Actinidia polygama*), cat thyme (*Teucrium marum*), Buckbean (*Menyanthes trifoliata*), actinidine, actinidiolide, boschnialactone, boschniakine, dihydroactinidiolide,

dihydronepetalactone, epinepetalactone, iridomyrmecin, isodihydronepetalactone, missugashiwalactone, neonepetalactone, onikulactone and mixtures thereof.

The cat attractants are present in the semiochemical compositions described herein in an amount of 0.05% to 0.15% (w%/w%). The cat attractants can also be present in the amount of 0.02% to 0.20% (w%/w%). They also can be present in the amount of 0.01% to 0.10% (w%/w%).

The concentration of the above-mentioned fatty acids may vary depending on the final form of use. However, the concentrations of the specific fatty acids that may be utilized and their concentration may be ascertained and tested according to the methods set forth herein.

The fatty acids, which are generally solid in nature, can be diluted in any nonaqueous solvent to form a solution of the present invention. More particularly, solvents such as ethyl alcohol, propylene glycol, alcohol, ether, chloroform, ethanol, benzene, acetone, propyl alcohol, isopropanol, 2-propanol fixed and volatile oils and the like. It is preferable to use ethyl alcohol in the formulation or acetone and propylene glycol. The ethyl alcohol can be 95% ethyl alcohol.

Fatty acids are commercially available from various chemical companies in solid form. However, since it is difficult to solubilize fatty acids, the fatty acid is generally added to the solvent under constant agitation and at a temperature of between about 37°C to about 38°C, more preferably 37.5°C.

The semiochemical compositions and analogues, as well as their derivatives and isomers, described herein, can be applied to scratching posts or cat trees or any object in which the cat has a tendency to scratch. For example, cat platforms, cat carpets, cat boards, cat toys and the like. Thus, the cat is attracted by the semiochemical compositions, analogues, derivatives and isomers and using that semiochemical coated object to scratch and tends not to scratch other objects that the cat had a habit to do prior to the semiochemical usage.

The above-described compositions were discovered after the detailed analysis of the interdigital area of cat's paws. This procedure involved recuperating from surfaces where the cats scratched secretions and analyzing the chemical

composition of these secretions via mass spectroscopy or gas chromatography/mass spectroscopy.

A first secretion called fraction Sc1 and a second secretion called Sc2 were obtained after analysis. Several analogues based on the Sc1 fraction and linoleic acid were also tested for their effectiveness to prevent cat's from scratching multiple objects. These analogues contained various amounts of the Sc1 fraction and differing amounts of linoleic acid.

The present invention also relates to a method of preventing cats from scratching on multiple objects, said method comprising placing one of the semiochemical compositions, analogues, derivatives and/or isomers of the present invention, as described herein, on an object such that the cat is attracted to the object and scratches on this objection and not other objects.

Cat posts, cat platforms, cat carpets, cat boards, cat toys and the like, having the semiochemical compositions, analogues, derivatives and isomers, as described herein, placed on these objects also forms a part of the invention.

In order to illustrate the present invention and the advantages thereof, the following specific examples are given, it being understood that the same are intended only as illustrative and in nowise limitative.

Example 1-A Population Study

The cats that were chosen for this study based on the major genetic branching and ethnic classification of cats. The three major types of cats that were chosen for this study were European cats, Siamese cats and Persian cats. The sex of the cats that were chosen for this study included females (F), castrated females (CF), males(M) and castrated males (CM).

The cats constituted twelve groups with five cats per group. This population was considered as an initial survey. The results obtained showed a large heterogeneity in the secretion compositions based either on the individual cats or their race. Therefore the sampling had to be repeated with a larger cat population.

Besides the criteria of race and sex, other criteria were established to be included in the dozen groups which were: (1) registered in a studbook kept by one of

- the recognized breeding clubs; (2) have an age between 8 months and 5 years; (3) did not receive any medical treatment for the last 3 months; (4) did not have any commercial cat scratching post; (5) did not use any litter that was chemical or perfumed; the cat litter had to be constituted of pure clay; (6) did not present any symptom of anxiety; (7) the scratch marking had to be executed every day on the same posts in proximity of a territorial zone in which the cat badly tolerated intrusions that is verified by a territorial aggression during an intrusion; and (8) the cat had a behavior of scratching that was observed at least one time per day by its owners and for at least one month.

The following Table I shows the cats that were selected for this study.

Table I

NUMBER & TYPE OF CATS	SEX	AGE IN MONTHS	DURATION OF SCRATCHING IN MONTHS
EUR1	M	12	3
EUR2	FC	9.5	4
EUR3	FC	11	2
EUR4	MC	25	5
EUR5	MC	9	3.5
EUR6	M	12.5	2
EUR7	F	11	1.5
EUR8	FC	10	2
EUR9	MC	9.5	3.5
EUR10	F	11.5	3
EUR11	M	10.5	3
EUR12	M	11	2.5
EUR13	F	12	4
EUR14	MC	15	2
EUR15	FC	18	1.5
EUR16	F	11.5	6
EUR17	MC	9.5	2

EUR18	F	18.5	3
EUR19	M	21	5
EUR20	FC	14	3
SIA1	FC	9.5	9
SIA2	MC	11	4
SIA3	MC	13.5	4
SIA4	FC	12	2.5
SIA5	F	14	2
SIA6	M	17	5
SIA7	F	9	3
SIA8	F	10.5	2
SIA9	M	12	2
SIA10	MC	18	2
SIA11	FC	14.5	7
SIA12	FC	9	6.5
SIA13	M	11	3
SIA14	MC	16.5	2
SIA15	FC	12	4
SIA16	F	10.5	6
SIA17	M	13	2.5
SIA18	F	18.5	3
SIA19	MC	20	2
SIA20	M	14	8
PER1	MC	9.5	6
PER2	F	10	4
PER3	M	11	3.5
PER4	MC	10.5	2
PER5	FC	14	3
PER6	FC	16	2.5
PER7	F	18	4
PER8	M	22	3
PER9	F	17	7
PER10	M	9.5	2.5

PER11	F	10	2
PER12	MC	12.5	5
PER13	MC	16	6
PER14	M	15.5	4
PER15	M	18	3
PER16	MC	9	2
PER17	FC	20	8
PER18	FC	21.5	4
PER19	F	12.5	1.5
PER20	FC	11	3

The homogeneity of the groups were confirmed by the student T test which gave the following results for age and race and for duration of scratching and race.

Unpaired T-test for AGE with the variable group of RACE with a theoretical

5 gap of 0

cat origin	average difference	Degrees of Freedom DDL	t	p
European, Siamese	-0.176	38	-0.145	0.8855
European, Persian	-1.075	38	-0.795	0.4313
Siamese, Persian	-0.899	38	-0.753	0.4562

Information of the group by AGE variable group of RACE

cats origin	number	average	variance	standard deviation	standard error
European	20	13.100	18.674	4.321	0.966
Siamese	20	13.276	10.693	3.270	0.731

Persian	20	14.175	17.849	4.225	0.945
---------	----	--------	--------	-------	-------

Unpaired t-test for DURATION OF SCRATCHING with a variable group of RACE with a theoretical gap of 0

origin of cats	average difference	Degrees of Freedom DDL	t	p
European, Siamese	-0.900	38	-1.589	0.1203
European, Persian	-0.725	38	-1.495	0.1431
Siamese, Persian	0.175	38	0.276	0.7842

- 5 Information of the DURATION OF SCRATCHING group with a variable group of RACE

origin of cats	number	average	variance	standard deviation	standard error
European	20	3.075	1.534	1.238	0.277
Siamese	20	3.975	4.881	2.209	0.494
Persian	20	3.800	3.168	1.780	0.398

Example 2-Isolation and Analysis to Identify the Semiochemical associated with Cat's Scratching

10 Isolation

To assure that the secretions were taken which were spontaneously deposited by the cats a device was used that permitted to obtain the scratches on a support that was compatible with the requirements of chemical analysis.

- 15 The samples were taken during a period of 48 hours by fixing a surgical compress using latex examining gloves on part of the surface lacerated by the cat.

The lacerated surface was divided in two in the sense of the height. One of the two halves was recovered with the aid of a sterile compress. To ensure that there was not any dissymmetry in the deposit of the secretion, the sample was collected alternatively half left and half right at each site for two days to collect the
5 pheromones.

The sterile compresses that were collected were replaced every 6 hours during the day at the same hour that the owners went to sleep and woke up. This represented 9 compresses per day per cat, thus 18 compresses on average per cat during two days of collection. During the collection, the sterile compresses were
10 manipulated with gloved hands and immediately placed in a flask that contained 10 ml of acetonitrile. This solvent was the one that was utilized during the analysis of the samples.

The flasks distributed to the cat's owners carried a coded number associated with their animal at the moment of recruitment. Each owner returned 5 flasks that
15 contained 3 to 4 sterile compresses.

During this collection phase, if the compresses came torn off from their support in the course of the execution of scratching and were found on the ground during collecting, they were eliminated from the study to not risk the contamination of the samples with other material. In the course of this study 9 sterile compresses had
20 to be eliminated.

Analysis

The solvent used throughout this study was HPLC grade acetonitrile (CH_3CN) that is commercialized by SIGMA (catalogue number 43,913-4). The samples were treated by differentiating the groups to which the individual cats belonged. For each
25 cat, the samples which were taken over a 48 hour period were mixed.

The sterile compresses were first mixed with the aid of a glass stirrer that was washed and rinsed with acetonitrile prior to inserting the samples from another cat. 30 ml of solution was obtained for each cat. 10 ml of each solution was evaporated under a nitrogen stream prior to being diluted in 1 ml of acetonitrile to realize a 10-
30 fold concentration.

The samples were then subjected to gas chromatography/mass spectroscopy using a Turbo Mass spectrometer made by Perkin Elmer. The detection was effectuated on impact using (EI+) at an energy of 70 eV at 180°C. A JW column type DB1 having a length of 60 m (id=0.25 mm film = 0.25 µm) at a split of 1/20 and a split/splitless of 3 seconds was used. 1.0 µl from the sample was injected.

The initial results were analyzed using a data base to obtain the most probable spectrums. Data bases containing such data are well known in the art.

The initial results obtained with the 60 chromatographic profiles presented little differences between the cats. These chromatographs contained 11 chemical substances, which were uniquely typical fatty acids or fatty acids in the form of methyl esters. Four of the seven fatty acids were in the form of an ester, which leads to the number of 11 components. These 11 components are set forth in Table II below

Table II

Compounds	Formula
propionic acid	$C_2H_5CO_2H$
lactic acid	$CH_3CH(OH)CO_2H$
valeric acid	$CH_3(CH_2)_3CO_2H$
methyl valerate	$CH_3(CH_2)_3CO_2CH_3$
isovaleric acid	$(CH_3)_2CHCH_2CO_2H$
cyclohexylacetic acid	$C_6H_{11}CH_2CO_2H$
methyl cyclohexylacetate	$C_6H_{11}CH_2CO_2CH_3$
3-cyclopentylpropionic acid	$C_5H_9CH_2CH_2CO_2H$
methyl 3-cyclopentylpropionate	$C_5H_9CH_2CH_2CO_2CH_3$
linoleic acid	$CH_3(CH_2)_4CH=CHCH_2CH=CH(CH_2)_7CO_2H$
methyl linoleate	$CH_3(CH_2)_4CH=CHCH_2CH=CH(CH_2)_7CO_2CH_3$

Since isovaleric acid was only present in 34% of cats it was excluded in the specific secretions. Also excluded were the esters since they could be considered as degradation products of the semiochemical secretion due to cutaneous bacteria. The secretions can also undergo a biotransformation due to symbiotes or saprophytes

before acquiring their active structure. In this the transformation of the pro-semiochemical is important and goes in the direction of simplifying the structure. This is what happens in the anal sacks of carnivores. The glands excrete proteins that are transformed into amines by the local saprophytes. In this case, the passage of the fatty acid to its methyl ester does not affect very much the spatial structure of the fatty acid, which is merely anecdotal and therefore permits to increase the portion of the fatty acids due to the ester.

Two fractions were identified having six different fatty acids. One fraction Sc1 was present in 69% to 83% of the total secretions. This fraction was called Sc1 and its compounds are present in Table III. The second fraction Sc2 was very rare and never appeared alone in the population of cats that was studied. Sc2 was present in the total secretions between 17% and 31 %. Sc2 was present in cats that were more stressed during the study. The compounds of Sc2 are presented in Table IV.

Table III

Compounds	Amounts in wt%/wt%
valeric acid	3.5 to 10.5%
lactic acid	5.5 to 15.5%
linoleic acid	68.5 to 83.5%

Table IV

Compounds	Amounts in wt%/wt%
propionic acid	36.5% to 43.5%
cyclohexylacetic acid	11.5% to 26.5%
cyclopentylpropionic acid	33.5% to 45.5%

Example 3-Testing of the Structural Analogues of the Cat Scratching Semiochemical

70 cats were tested in this study using the solution SC10-98, SC10-75, SC10-60, SC20-98, SC20-75 and SC20-60, which are structural analogues of the feline scratch marking pheromone. SC10-98 had 10% of the fraction Sc1 (See, Table III

above) and 98% linoleic acid; SC10-75 had 10% of the fraction Sc1 (See, Table III above) and 75% linoleic acid; SC10-60 had 10% of the fraction Sc1 (See, Table III above) and 60% linoleic acid; SC20-98 had 20% of the fraction Sc1 (See, Table III above) and 98% linoleic acid; SC20-75 had 20% of the fraction Sc1 (See, Table III above) and 75% linoleic acid; and SC20-60 had 20% of the fraction Sc1 (See, Table III above) and 60% linoleic acid.

Ethanol was used as a placebo, which was the same solvent used in the solutions. This study was not conducted as a double blind study since although the six analogues could not be distinguished by the human nose, the ethanol could in fact be distinguished. However the treatment was conducted randomly.

For this study the analogues of the feline scratch marking semiochemical were applied to a new cat scratching post for cats that were habituated to scratch another object.

The cats for this study were pets from the Paris area. To be included in the study, besides their approval from their owners, the cats must have satisfied the following conditions: (1) cats of all different races; (2) castrated or non-castrated males and females; (3) the minimum age was 9 months and the maximum age was 8 years; (4) have no signs of an ailment and have not been treated; and (5) the cat scratched at a specific site constantly for a period of at least 2 months.

The criteria for non-inclusion were (1) that the cats had onychetomies (declawed); (2) the cats were in any bad health; (3) the cats were undergoing an intervention during the 7 days of the test; (4) the cats were carriers of torn nail sheaths; and (5) the cats underwent behavioral therapy for scratching.

The criteria for exclusion included (1) occurrence of any organic disease; (2) the cats underwent anesthesia; (3) the cats were administered drugs with the exception of intern and extern antiparasites; (4) any modification of the cats environment including moving; and (5) introducing a new animal into the household.

70 cats were used in this study; 10 cats for each treatment. E stands for European S stands for Siamese and P stands for Persian. The treatment and background of each cat that was tested is set forth in Table V below.

Table V

CATS	RACE	SEX	AGE in months	DURATION OF SCRATCHING IN MONTHS	TREATMENT GIVEN
C1	E	M	9	2	SC10-60
C2	S	MC	12	3	SC20-98
C3	E	MC	11	4	SC20-60
C4	E	FC	28	11	SC10-75
C5	E	FC	26	15	P
C6	S	MC	33	24	SC20-75
C7	S,E	F	21	15	SC20-98
C8	SB	FC	14	6	P
C9	E	MC	9.5	2	SC20-60
C10	S	MC	12.5	3	SC10-98
C11	P	FC	11.5	2	SC10-75
C12	E	MC	14.5	5	SC20-75
C13	P	M	17	3	SC20-75
C14	P,E	FC	14	2	SC10-60
C15	E	FC	15.5	3	SC20-60
C16	P	F	16	6	SC20-98
C17	P,E	MC	10	2	P
C18	E	MC	10.5	4	SC10-60
C19	S	MC	15	5	P
C20	P,E	M	16.5	6	SC10-98
C21	SB	FC	14.5	8	SC10-75
C22	SB,E	FC	46	38	SC20-98
C23	P	FC	36.5	28	SC10-60
C24	P	F	25	11	SC10-98
C25	E	MC	33	25	SC20-60
C26	P	MC	32.5	24	SC20-75
C27	P,E	MC	18	11	SC10-98

C28	E	FC	16.5	8	SC20-98
C29	E	FC	62	54	SC20-75
C30	E	F	65.5	58	SC10-75
C31	E	FC	23.5	16	SC10-60
C32	S,E	MC	36.5	29	SC10-75
C33	SB	MC	42.5	37	P
C34	P	M	41	34	SC10-98
C35	E	MC	25	16	SC20-75
C36	P,E	MC	33.5	25	SC10-60
C37	E	MC	36	28	SC20-60
C38	P	FC	14	6	SC10-75
C39	SB	M	12.5	5	SC20-98
C40	P	FC	9.5	2	SC10-60
C41	E	MC	28	20	SC20-60
C42	E	MC	29.5	22	P
C43	E	MC	21	14	SC10-98
C44	E	FC	22.5	13	SC10-75
C45	P	FC	24	15	SC20-98
C46	P,E	F	28.5	16	P
C47	S	FC	26.5	19	SC20-60
C48	E	MC	39	30	SC10-60
C49	E	MC	44.5	35	SC10-98
C50	S	MC	42.5	36	SC20-75
C51	P	FC	68	60	SC20-98
C52	S,P	MC	86	78	P
C53	E	MC	26	17	SC10-75
C54	E	M	44	36	SC10-98
C55	E	FC	43.5	34	SC20-98
C56	P	F	28.5	20	SC20-75
C57	S	MC	29.5	21	SC10-60
C58	E	FC	41.5	32	SC20-60
C59	E	FC	42.5	31	SC10-98
C60	S,E	FC	44	36	P

C61	P	FC	46.5	39	SC20-60
C62	S	MC	57.5	48	SC10-60
C63	P,E	MC	58	52	SC10-75
C64	E	M	62.5	51	SC20-98
C65	E	MC	96	56	SC10-98
C66	S	FC	56	47	SC20-75
C67	P	MC	9	3	P
C68	SB	MC	10.5	3	SC20-60
C69	P	MC	13	5	SC20-75
C70	E	MC	12.5	4	SC10-75

P stands for placebo in the above Table. SB stands for the Birmin cat (sacred cat of Burma.)

Example 4-Testing the Analogue Semiochemical

5 This study utilized two types of material. A commercial scratching post composed of a board having the dimension 25 x 60 cm covered with a hemp string arranged in the direction of the width to the entire length. The scratching post was covered on part of the surface by yellow peeling paper in the direction of the length that permits the validation that the scratching post was utilized. The paper was
10 changed each 7 to 8 hours that permitted to have 3 measures taken by day. When the paper was replaced it was placed in a different area on the scratching post that was not used during the 8 hours precedent to obtain an alternative dimension. Lacerated or not the papers were replaced and conserved for testing.

15 A brown glass bottle containing 50 ml of solution of the analogue of the semiochemical in an alcoholic solution without an adjuvant or a placebo was prepared. This product was applied to the scratching post with the aid of a sterile complex and applied using latex gloves. The product was applied using four longitudinal bands occupying the entire length of the scratching post. This application was renewed every day and once a day during the study.

20 The cats selected for this study were those that already had a habit of using a scratching post. Their scratching post habitual was removed and replaced with the

scratching post described above. The study was performed over a period of seven days.

The following parameters were studied:

NPL: the numbers of papers that were lacerated in 7 days

5 DBFL: the delay before the first laceration

SOA: the number of scratches on another object

The NPL permitted to evaluate visually the frequency of new scratches, The DBFL permitted to estimate the number of lacerations with the new scratching post. The SOA permitted to identify the failure leading to the lacerations and then
10 dispensed to other objects besides the scratching post.

The method of evaluation of this study was the following. The evaluation of NPL was done using an evaluation of 1 for all of the papers that had lacerations thereon. A paper that simply became unhooked from the support was given a score of 0, lacking an intact non-lacerated paper. The DBFL was not evaluated until the
15 replacement paper was added. It was taken into account the number of first order paper that was scratched. If no paper was scratched a note of 22 was given. The value of DBFL was between 1 and 22. SOA was evaluated for every scratch that was noted. If no scratches were present then a note of zero was assigned. SOA was thus the sum of the scores during the seven days. A scratch was scored if it was directly
20 observed or if traces of a scratch could be found.

The results are set forth in Table VI below.

TABLE VI

CATS	TREATMENT	NPL	DBFL	SOA
C1	SC10-60	3	6	2
C2	SC20-98	18	1	0
C3	SC10-60	7	3	1
C4	SC10-75	9	5	0
C5	P	1	5	5
C6	SC20-75	20	1	0

C7	SC20-98	19	2	0
C8	P	0	22	6
C9	SC20-60	7	6	2
C10	SC10-98	12	3	3
C11	SC10-75	4	4	5
C12	SC20-75	17	1	0
C13	SC20-75	18	1	1
C14	SC10-60	3	8	6
C15	SC20-60	9	4	2
C16	SC20-98	20	1	0
C17	P	4	17	4
C18	SC10-60	2	7	6
C19	P	0	22	7
C20	SC10-98	11	6	2
C21	SC10-75	3	7	5
C22	SC20-98	21	1	0
C23	SC10-60	4	9	4
C24	SC10-98	6	5	1
C25	SC20-60	8	7	3
C26	SC20-75	18	2	0
C27	SC10-98	6	9	7
C28	SC20-98	20	1	0
C29	SC20-75	19	1	0
C30	SC10-75	6	11	3
C31	SC10-60	3	9	6
C32	SC10-75	7	8	4
C33	P	5	14	7
C34	SC10-98	14	3	0
C35	SC20-75	19	1	0
C36	SC10-60	2	14	6
C37	SC20-60	7	6	2
C38	SC10-75	6	12	5
C39	SC20-98	20	2	0

C40	SC10-60	2	18	7
C41	SC20-60	6	9	0
C42	P	1	15	3
C43	SC10-98	17	3	1
C44	SC10-75	3	11	3
C45	SC20-98	17	3	0
C46	P	3	17	7
C47	SC20-60	11	8	6
C48	SC10-60	5	14	6
C49	SC10-98	18	2	1
C50	SC20-75	21	1	0
C51	SC20-98	20	1	0
C52	P	4	16	4
C53	SC10-75	9	8	3
C54	SC10-98	14	6	3
C55	SC20-98	21	1	0
C56	SC20-75	17	2	0
C57	SC10-60	3	16	7
C58	SC20-60	12	6	2
C59	SC10-98	15	3	1
C60	P	0	22	6
C61	SC20-60	9	11	0
C62	SC10-60	2	14	1
C63	SC10-75	6	9	5
C64	SC20-98	18	2	0
C65	SC10-98	11	7	5
C66	SC20-75	19	1	1
C67	P	2	17	7
C68	SC20-60	4	14	3
C69	SC20-75	15	3	0
C70	SC10-75	3	13	7

P is the placebo.

The results are set forth in Figures 1 to 3. The results of this analysis permit to conclude that solutions SC20-98 and SC20-75 are the semiochemical that were more efficient.

The results of the t tests for SC20-75 and SC20-98 are set forth below.

- 5 Unpaired t-test for NPL variable group:treatment Theoretical gap=0 for SC20-75, SC20-98

average gap	DDL	t	p
-1.100	18	-1.601	0.1268

Information of the group for NPL variable group:treatment

treatment	number	average	variance	standard deviation	standard error
SC20-75	10	18.300	2.900	1.703	0.539
SC20-98	10	19.400	1.822	1.350	0.427

- 10 Unpaired t-test for DBFL variable group:treatment Theoretical gap=0 for SC20-75, SC20-98

average gap	DDL	t	p
-1.00	18	-0.318	0.741

Information of the group for DBFL variable group:treatment

treatment	number	average	variance	standard deviation	standard error
SC20-75	10	1.400	0.489	0.699	0.221
SC20-98	10	1.500	0.500	0.707	0.224

- 15 Unpaired t-test for SOA variable group:treatment Theoretical gap=0 for SC20-75, SC20-98

average gap	DDL	t	p
0.200	18	1.500	0.1510

Information of the group for SOA variable group:treatment

treatment	number	average	variance	standard deviation	standard error
SC20-75	10	0.200	0.178	0.422	0.133
SC20-98	10	0.000	0.000	0.000	0.000

This analysis highlighted the absence of any statistically significant difference between the two treatments for all of the parameters that were studied. This confirms the efficacy of the analogs of the scratching pheromone and permitting to discriminate between the different solutions and to choose the solution which is more efficient.

Example 5- Optimization of SC20-75

SC20-75 was chosen for this study since it was easier to fabricate at a lower cost than SC20-98. This study was performed to optimize SC20-75 either by using an olfactory adjuvant or by the association with visual signals that reproduce an attractive effect to the cats associated with vertical lacerations and are perfectly mimicked by the presence of vertical stripes such as wallpaper, fabrics etc.

This study was performed on cats that had consultations with two veterinarians at the same clinic and thus was a monocentric study. The cats were described as destructors since they had a behavior of marking by scratching and destroying objects.

To be included in this study, the cats (1) had a behavior of marking by scratching for at least 2 months; (2) were in good health and behavior other than the scratching; (3) were between the ages of 9 months and 8 years; and (4) sex and race were indifferent.

The cats that were not included in this study were those that (1) had onychectomies (declawed) (2) were subjected to a therapy to remove the claws (3)

presented somatic or behavior disorders (4) had treatments for psychotropic medications or hormones for at least 6 months (5) cats whose scratching was lessened by semiochemical therapy.

In the course of the study the cats would be excluded for the following reasons: (1) exacerbation of scratching; (2) default of compliance with the study such as paper with the scratch marks not returned; (3) appearance of urinary marking or aggression; (4) were given anesthesia; (5) had a surgical intervention; (6) caught an organic disease; and (7) administration of a drug other than an antiparasite internally or externally.

Sixteen cats were selected for this study that lasted 15 days. E stands for Eurasian, S for Siamese and P for Persian in the table below. The cats had the following characteristics:

Table VII

CATS	RACE	SEX	AGE IN MONTHS	DURATION OF SCRATCHING IN MONTHS
C1	E	MC	15	3
C2	E	MC	18	3
C3	E	MC	26	2
C4	S	FC	24	6
C5	E	MC	28	4
C6	P	FC	27	2
C7	S,E	F	32	6
C8	P	FC	11	3
C9	P,E	MC	9	2
C10	E	FC	15	8
C11	E	FC	17	5
C12	E	M	9	2
C13	P	FC	10	4
C14	E	MC	26	18

C15	SB	M	23	14
C16	E	MC	9	3

SB stands for the Birmin cat (sacred cat of Burma.)

The statistics on the group of cats chosen for this study that had the appropriate duration of scratching are as follows:

5

Average	5.312
Standard Deviation	4.571
Standard Error	1.143
Number	16
Minimum	2.000
Maximum	18.000
Number missing	0

The scratching posts were the same as used in Example 4, but two scratching posts were contributed to each cat to respect the methods used in each test. One of the scratching posts was labeled S hidden from the face of the cat. It was coated with the adjuvanted analogue scratching semiochemical from a flask also marked S. The other scratching post was marked A and was coated with an adjuvanted alcohol from a flask marked A.

The 50 ml solutions of analogue scratching semiochemical were contained in a brown glass flask and contained SC20-75 adjuvanted with a 10% of tincture of catnip. The control was a solution of 95% alcohol adjuvanted with tincture of catnip at 10% of the total volume. The scratching posts were coated as in Example 4. The paper that was utilized on the scratching post was the same as in Example 4.

To compare the efficacy of the two scratching posts, they were placed 60 centimeters apart and on either side of a surface that was previously scratched. The scratching post (S) had the semiochemical analogue coated thereon, while the post (A) was coated with the control. Like in Example 4 a part of the scratching post was

recovered with peeling paper to detect the scratches. In this case the papers were also placed according to the same modalities on the object that was already scratched by the cat. One sole object (if there were many) was left accessible to the cat, which is the object that should be the most scratched. The evaluation was

5 based on this object. The study was performed for a period of 7 days.

The evaluation of this study was done with the same parameters as that set forth in Example 4 with the exception of SOA that was replaced by the measure of the object that was habitually scratched by the cat (OL). This object was considered as the place of scratching in competition with the other scratching posts of the cat.

10 The results are set forth in Table VIII below for the study done with the analogue of cat scratch semiochemical (S).

Table VIII

CATS	NPL	DBFL
C1	17	1
C2	18	2
C3	15	1
C4	19	2
C5	20	1
C6	18	3
C7	19	1
C8	21	1
C9	18	2
C10	15	4
C11	18	2
C12	19	1
C13	19	1
C14	19	1
C15	20	1
C16	17	3

The results are set forth below in Table IX for the control (A).

Table IX

CATS	NPL	DBFL
C1	2	9
C2	2	15
C3	1	18
C4	0	22
C5	3	7
C6	1	17
C7	1	18
C8	2	14
C9	1	12
C10	3	16
C11	1	17
C12	3	14
C13	0	22
C14	0	22
C15	1	16
C16	0	22

5

The results are set forth below in Table X for the object that was habitually scratched by the cat (OL).

Table X

CATS	NPL	DBFL
C1	14	1
C2	11	1
C3	15	1

C4	9	3
C5	10	2
C6	6	1
C7	13	1
C8	8	1
C9	14	2
C10	15	1
C11	3	2
C12	6	2
C13	4	1
C14	8	1
C15	8	1
C16	3	2

The results are shown in Figures 4 and 5.

Statistical analysis was then performed using the above results.

Analogue Scratch Semiochemical SC20-75

Test	Average	Standard Deviation	Standard Error	Number	Minimum	Maximum	Number missing
NPL	18.250	1.653	0.4313	16	15.000	21.000	0
DBFL	1.688	0.946	0.237	16	1.000	4.000	0

5

Control (A)

Test	Average	Standard Deviation	Standard Error	Number	Minimum	Maximum	Number missing
NPL	1.312	1.078	0.270	16	000	3.000	0
DBFL	16.312	4.527	1.132	16	7.000	22.000	0

The object that was habitually scratched by the cat

Test	Average	Standard Deviation	Standard Error	Number	Minimum	Maximum	Number missing
NPL	9.188	4.167	1.042	16	3.000	15.000	0
DBFL	1.438	0.629	0.157	16	1.000	3.000	0

The SC20-75 analogue of feline interdigital semiochemicals was tested in sixteen (16) cats. Those cats were described as scratching intensively at home by their owners. For each cat, the owners were asked to define a “target area” which was a place where the cat had intensive scratching activity. Two cardboard scratching posts were fixed 60 cm on each side of the target area. One of those scratching posts was coated with a placebo and the other the semiochemical solution. The assessment parameter was the total number of scratches during 7 days of the study, counted by putting a piece of paper on half of the scratching surfaces. Comparison was made using the Mann-Whitney test.

The scratching post sprayed with the semiochemical was used more by the cats in a significant way when compared to the target area ($p < 0.05$) and in a very highly significant way when compared to the target area ($p < 0.001$).

While the invention has been described in terms of various preferred embodiments, the skilled artisan will appreciate that various modifications, substitutions, omissions and changes may be made without departing from the scope thereof. Accordingly, it is intended that the scope of the present invention be limited by the scope of the following claims, including equivalents.

What is claimed is:


1. A semiochemical composition comprising a secretion obtained from the interdigital paws of a cat.
2. A semiochemical composition comprising valeric acid, lactic acid and linoleic acid or derivatives thereof.
3. The semiochemical composition according to Claim 2, comprising between about 3.5% to 10.5% (w%/w%) of valeric acid, between about 5.5% to 15.5% (w%/w%) of lactic acid and between 68.5% to 83.5 (w%/w%) of linoleic acid and/or derivatives thereof and/or isomers thereof.
4. A semiochemical composition comprising between about 3.5% to 10.5% (w%/w%) of valeric acid, between about 5.5% to 15.5% (w%/w%) of lactic acid, between about 48.4 % to 58.9 % (w%/w%) of linoleic acid and between about 20.1% to 24.5 % (w%/w%) of oleic acid and/or derivatives thereof and/or isomers thereof.
5. A semiochemical composition analogue comprising 10% to 20% of a composition comprising valeric acid, lactic acid and f linoleic acid and between 60% to 90% (w%/w%) of oleic acid and/or derivatives thereof and/or isomers thereof.
6. A semiochemical composition analogue comprising 10% to 20% of a composition comprising between about 3.5% to 10.5% (w%/w%) of valeric acid, between about 5.5% to 15.5% (w%/w%) of lactic acid and between 68.5% to 83.5 (w%/w%) of linoleic acid and between 60% to 90% (w%/w%) of oleic acid and/or derivatives thereof and/or isomers thereof.
7. A semiochemical composition comprising 20% of a composition comprising between about 3.5% to 10.5% (w%/w%) of valeric acid, between about 5.5% to 15.5% (w%/w%) of lactic acid and between 68.5% to 83.5

(w%/w%) of linoleic acid and/or derivatives thereof and/or isomers thereof and 75% linoleic acid.


- 5 8. A semiochemical composition comprising 20% of a composition comprising between about 3.5% to 10.5% (w%/w%) of valeric acid, between about 5.5% to 15.5% (w%/w%) of lactic acid and between 68.5% to 83.5 (w%/w%) of linoleic acid or derivatives thereof and/or isomers thereof and 98% linoleic acid
- 10 9. The semiochemical composition or semiochemical composition analogue according to any one of Claims 1 to 8, wherein the derivatives are esters, salts, alcohols, ketones, ethers and amides of valeric acid, lactic acid and linoleic acid.
- 15 10. The semiochemical composition or semiochemical composition analogue according to any one of Claims 1 to 8, wherein the isomers are positional isomers and/or geometrical isomers of cis- trans, trans-cis, cis-cis or trans-trans configurations.
- 20 11. The semiochemical composition according to any one of Claims 1 to 10, further comprising a cat attractant.
- 25 12. The semiochemical composition according to Claim 11, wherein the cat attractant is catnip.
- 30 13. A solution comprising the semiochemical compositions of the present invention according to any one of Claims 1 to 12 and a solvent.
- 35 14. A method for preventing a cat from scratching on multiple objects, said method comprising placing at least one semiochemical composition according to any one of Claims 1 to 13 on an object that a cat uses for scratching.
15. An object for cats covered with the semiochemical composition and/or derivatives thereof and/or isomers thereof or semiochemical composition

analogue and/or derivatives thereof and/or isomers thereof according to any one of Claims 1 to 13.


1/5

**FIGURE 1**

2/5

**Figure 2**

3/5

**FIGURE 3**





FIGURE 4

5/5

**FIGURE 5**

INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2012/001341

A. CLASSIFICATION OF SUBJECT MATTER INV. A01N37/02 A01N37/06 A01N37/36 A01N25/34 A01P19/00 ADD.		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) A01N A01P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EP0-Internal, WPI Data, CHEM ABS Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	A. COZZI ET AL: "The interest of the use of feline interdigital semiochemical (ScratchyLicious) to induce scratching behaviour in cat", 2011 ACVB/AVSAB VETERINARY BEHAVIOR SYMPOSIUM, ST. LOUIS, MISSOURI, 15 July 2011 (2011-07-15), pages 4-5, XP055056716, pages 4-5 ----- -/--	1-15
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search	Date of mailing of the international search report	
18 March 2013	02/04/2013	
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Hateley, Martin	

INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2012/001341

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	COZZI A ET AL: "The maternal cat appeasing pheromone: Exploratory study of the effects on aggressive and affiliative interactions in cats", JOURNAL OF VETERINARY BEHAVIOR: CLINICAL APPLICATIONS AND RESEARCH, ELSEVIER, AMSTERDAM, NL, vol. 5, no. 1, 1 January 2010 (2010-01-01) , pages 37-38, XP026855547, ISSN: 1558-7878 [retrieved on 2010-01-14] the whole document	1-15
A	----- GB 2 345 635 A (DYNAVET S A R L [FR]) 19 July 2000 (2000-07-19)	1-15
A	----- EP 0 724 832 A1 (PAGEAT PATRICK [FR] FIDELINE [FR]) 7 August 1996 (1996-08-07) cited in the application the whole document	1-15
A	----- US 2002/189549 A1 (WRITER TYA B [US] ET AL) 19 December 2002 (2002-12-19) the whole document -----	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/IB2012/001341

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 2345635	A	19-07-2000	NONE
EP 0724832	A1	07-08-1996	AT 185046 T 15-10-1999
		AU 692675 B2 11-06-1998	
		AU 4716996 A 21-08-1996	
		CA 2210992 A1 08-08-1996	
		DE 69512509 D1 04-11-1999	
		DE 69512509 T2 31-05-2000	
		DK 724832 T3 10-04-2000	
		EP 0724832 A1 07-08-1996	
		ES 2139853 T3 16-02-2000	
		FI 963902 A 27-09-1996	
		GR 3032140 T3 27-04-2000	
		HK 1014629 A1 07-09-2001	
		JP 3886146 B2 28-02-2007	
		JP H10513164 A 15-12-1998	
		NO 963781 A 21-11-1996	
		NZ 301681 A 29-11-1999	
		US 5709863 A 20-01-1998	
		WO 9623414 A1 08-08-1996	
US 2002189549	A1	19-12-2002	NONE