



(51) International Patent Classification:

C07C 235/34 (2006.01) C07D 249/18 (2006.01)  
C07C 233/11 (2006.01) C07D 263/54 (2006.01)  
C07C 233/07 (2006.01) C07D 277/62 (2006.01)  
C07C 237/20 (2006.01) A61K 31/165 (2006.01)  
C07C 255/00 (2006.01) A61K 31/17 (2006.01)  
C07C 271/28 (2006.01) A61K 31/18 (2006.01)  
C07C 275/28 (2006.01) A61K 31/277 (2006.01)  
C07C 307/10 (2006.01) A61K 31/404 (2006.01)  
C07C 311/03 (2006.01) A61K 31/416 (2006.01)  
C07D 209/18 (2006.01) A61K 31/4184 (2006.01)  
C07D 215/02 (2006.01) A61K 31/428 (2006.01)  
C07D 231/56 (2006.01) A61K 31/47 (2006.01)  
C07D 213/74 (2006.01) A61P 9/12 (2006.01)  
C07D 235/04 (2006.01) A61P 13/12 (2006.01)

(21) International Application Number:

PCT/AU2014/000922

(22) International Filing Date:

17 September 2014 (17.09.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

2013903571 17 September 2013 (17.09.2013) AU  
2013903572 17 September 2013 (17.09.2013) AU

(71) Applicant: VECTUS BIOSYSTEMS PTY LTD  
[AU/AU]; Level 3, 11 Julius Avenue, North Ryde, New  
South Wales 2113 (AU).

(72) Inventor: DUGGAN, Karen Annette; 9 Surfside Avenue,  
Clovelly, New South Wales 2031 (AU).

(74) Agent: SHELSTON IP; Level 21, 60 Margaret Street,  
Sydney, New South Wales 2000 (AU).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,  
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,  
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,  
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,  
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,  
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,  
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,  
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,  
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,  
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,  
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,  
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,  
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,  
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

(54) Title: COMPOSITIONS FOR THE TREATMENT OF HYPERTENSION AND/OR FIBROSIS

(57) Abstract: The present invention relates to novel compounds and their use in the prophylactic and/or therapeutic treatment of hypertension and/or fibrosis.



WO 2015/039172 A1

## COMPOSITIONS FOR THE TREATMENT OF HYPERTENSION AND/OR FIBROSIS

### *FIELD OF THE INVENTION*

5 The present invention relates to novel compounds and their use in the prophylactic and/or therapeutic treatment of cardiovascular disease, and in particular the treatment of prehypertension, hypertension and/or fibrotic conditions.

The invention has been developed primarily for the prophylactic and/or therapeutic treatment of cardiovascular disease and will be described hereinafter with reference to  
10 this application. However, it will be appreciated that the invention is not limited to this particular field of use.

### *BACKGROUND OF THE INVENTION*

Any discussion of the prior art throughout the specification should in no way be considered as an admission that such prior art is widely known or forms part of the  
15 common general knowledge in the field.

Hypertension (high blood pressure) affects 26% of the adult population worldwide with an incidence of 30-33% in western countries. The world wide incidence of hypertension is expected to reach 29% by 2025 as a consequence of the westernisation of India and China. Current studies indicate that fewer than 20% of patients with hypertension attain  
20 their recommended blood pressure (BP) target and that to achieve these targets >75% of patients require therapy with multiple antihypertensive agents. Prehypertension (slightly elevated blood pressure) affects 31% of adults in the US and may develop into hypertension if not treated.

All currently available therapies have side effects:

- 25 • Angiotensin Converting Enzyme Inhibitors (ACEI) – cough, angioneurotic oedema, hyperkalaemia;
- Angiotensin Receptor Blockers (ARB's) - angioneurotic oedema, hyperkalaemia;
- Calcium Channel Blockers (CCB) – flushing, leg / ankle oedema, constipation;
- Thiazide diuretics – new onset diabetes, gout, hyponatraemia;
- 30 • Beta ( $\beta$ ) Blockers – new onset diabetes, inability to exercise, bradycardia, masking hypoglycaemia in diabetics; and

- 2 -

- Aldosterone Antagonists – gynaecomastia, menorrhagia, hyperkalaemia.

The need to use combination therapy increases the likelihood that patients will experience side effects and as a consequence not attain their BP target.


Hypertension and prehypertension are a major factor in the development of heart, kidney  
5 and blood vessel damage, resulting in the replacement of normal functional tissue by scar  
tissue or fibrosis. Some of the current antihypertensive agents – ACE inhibitors, ARB's  
renin inhibitors and aldosterone antagonists are able to slow the progression of the  
replacement of functional tissue by fibrosis, none have been shown to reverse existing  
fibrosis and restore normal tissue architecture. There is thus a need for agents which have  
10 the efficacy to reduce BP significantly and thus enable a larger proportion of patients to  
attain BP target with single agent therapy and/or to reverse existing fibrosis and/or restore  
normal tissue architecture.

It is an object of the present invention to overcome or ameliorate at least one of the  
disadvantages of the prior art, or to provide a useful alternative.

#### 15 **SUMMARY OF THE INVENTION**

Surprisingly, the present inventors have found that certain novel terphenyl compounds  
have blood pressure lowering and/or anti-fibrotic effects. These effects may be seen in  
intravenous and/or oral dosing studies.


According to one aspect, the present invention provides a compound of the formula



20

wherein:

A is selected from the group consisting of:



Q is independently selected from halo, alkyl, hydroxy, amino and substituted amino;  
n is 0, 1, 2, 3, 4 or 5;


- 5 R<sub>1</sub>, R<sub>3</sub> and R<sub>4</sub> are independently C, CH, CH<sub>2</sub>, O, N, NH or S, and  
R<sub>2</sub> is C, CH, CH<sub>2</sub>, N, NH, C-CF<sub>3</sub>, CH-CF<sub>3</sub> or C=O,  
or a stereoisomer or pharmaceutically acceptable salt thereof,  
wherein when n is 1, Q cannot be hydroxy.

In one embodiment, Q is halo selected from the group consisting of F, Cl, Br and I.

- 10 In one embodiment, Q is substituted amino of the formula -NHW and wherein:  
W is selected from -CN, -SO<sub>2</sub>(X)<sub>a</sub>Y and -CO(X)<sub>a</sub>Y,  
a is 0 or 1,  
X is selected from -NH- and -O-, and  
Y is selected from -H, -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>OH and -CH<sub>2</sub>CH<sub>2</sub>OH.
- 15 In one embodiment, Q is substituted amino selected from the group consisting of  
-NHSO<sub>2</sub>CH<sub>3</sub>, -NHCOH, -NHCONHCH<sub>3</sub>, -NHCONHCH<sub>2</sub>CH<sub>3</sub>, -NHSO<sub>2</sub>NHCH<sub>3</sub>,  
-NHSO<sub>2</sub>NHCH<sub>2</sub>CH<sub>3</sub>, -NHCOCH<sub>3</sub>, -NHCOOCH<sub>3</sub>, -NHCOOCH<sub>2</sub>CH<sub>2</sub>OH, -NHCONH<sub>2</sub> and  
-NHCN.


- In one embodiment, Q is alkyl selected from the group consisting of methyl, ethyl,  
20 propyl, butyl and pentyl.

In one embodiment, A is selected from:



and

5 In one embodiment, A is




wherein:

X, Y or Z are C or N, wherein only one of X, Y or Z can be N,

R<sub>5</sub> to R<sub>9</sub> are independently selected from hydrogen, halo, alkyl, hydroxy, amino and  
 10 substituted amino, with the proviso that when one of R<sub>5</sub> to R<sub>9</sub> is halo, then at least one of  
 the remaining R<sub>5</sub> to R<sub>9</sub> cannot be hydrogen

In one embodiment, A is

- 5 -




wherein:

X, Y or Z are C or N, wherein only one of X, Y or Z can be N,

R<sub>5</sub> to R<sub>9</sub> are independently selected from hydrogen, halo, alkyl, hydroxy, amino and substituted amino, with the proviso that when one of R<sub>5</sub> to R<sub>9</sub> is halo, then at least one of the remaining R<sub>5</sub> to R<sub>9</sub> must be halo, alkyl, hydroxy, amino or substituted amino.

In one embodiment, A is




wherein:

10 X, Y or Z are C or N, wherein only one of X, Y or Z can be N,

R<sub>5</sub> to R<sub>9</sub> are independently selected from hydrogen, halo, alkyl, hydroxy, amino and substituted amino, with the proviso that when one of R<sub>5</sub> to R<sub>9</sub> is halo, then at least one of the remaining R<sub>5</sub> to R<sub>9</sub> must be alkyl, hydroxy, amino or substituted amino.

In one embodiment, A is



- 6 -


wherein:

X, Y or Z are C or N, wherein only one of X, Y or Z can be N,

R<sub>5</sub> to R<sub>9</sub> are independently selected from hydrogen, halo, alkyl, hydroxy, amino and substituted amino, with the proviso that when one of R<sub>5</sub> to R<sub>9</sub> is alkyl, then at least one of

5 the remaining R<sub>5</sub> to R<sub>9</sub> cannot be hydrogen

In one embodiment, A is




wherein:

X, Y or Z are C or N, wherein only one of X, Y or Z can be N,

10 R<sub>5</sub> to R<sub>9</sub> are independently selected from hydrogen, halo, alkyl, hydroxy, amino and

substituted amino, with the proviso that when one of R<sub>5</sub> to R<sub>9</sub> is alkyl, then at least one of the remaining R<sub>5</sub> to R<sub>9</sub> must be halo, alkyl, hydroxy, amino or substituted amino.

In one embodiment, A is



15 wherein:


X, Y or Z are C or N, wherein only one of X, Y or Z can be N,

R<sub>5</sub> to R<sub>9</sub> are independently selected from hydrogen, halo, alkyl, hydroxy, amino and substituted amino, with the proviso that when one of R<sub>5</sub> to R<sub>9</sub> is alkyl, then at least one of

the remaining R<sub>5</sub> to R<sub>9</sub> must be halo, hydroxy, amino or substituted amino.

20 In one embodiment, A is

- 7 -




wherein:

X, Y or Z are C or N, wherein only one of X, Y or Z can be N,

- R<sub>5</sub> to R<sub>9</sub> are independently selected from hydrogen, halo, alkyl, hydroxy, amino and substituted amino, with the proviso that when one of R<sub>5</sub> to R<sub>9</sub> is hydroxy, then at least one of the remaining R<sub>5</sub> to R<sub>9</sub> cannot be hydrogen

In one embodiment, A is




wherein:

- X, Y or Z are C or N, wherein only one of X, Y or Z can be N,

R<sub>5</sub> to R<sub>9</sub> are independently selected from hydrogen, halo, alkyl, hydroxy, amino and substituted amino, with the proviso that when one of R<sub>5</sub> to R<sub>9</sub> is hydroxy, then at least one of the remaining R<sub>5</sub> to R<sub>9</sub> must be halo, hydroxy, amino or substituted amino.


In one embodiment, A is





wherein:


X, Y or Z are C or N, wherein only one of X, Y or Z can be N,


R<sub>5</sub> to R<sub>9</sub> are independently selected from hydrogen, halo, alkyl, hydroxy, amino and substituted amino, with the proviso that when one of R<sub>5</sub> to R<sub>9</sub> is hydroxy, then at least one  
 5 of the remaining R<sub>5</sub> to R<sub>9</sub> must be halo, amino or substituted amino.





In one embodiment, A is , Q is substituted amino and n is 1.





In one embodiment, A is , Q is substituted amino and n is 2.





In one embodiment, A is , Q is amino and n is 1.




In one embodiment, A is , Q is amino and n is 2.




10 In one embodiment, A is , Q is hydroxy and n is 2.




In one embodiment, A is




In one embodiment, A is




In one embodiment, A is




In one embodiment, A is



5 In one embodiment, A is




In one embodiment, A is




In one embodiment, A is  
is amino.


, X and Y are C, Z is N and one of R<sub>5</sub> to R<sub>9</sub>




In one embodiment, A is  $\text{C}_6\text{H}_4$ , X, Y and Z are all C, one of R<sub>5</sub> to R<sub>9</sub> is hydroxy and at least one of the remaining R<sub>5</sub> to R<sub>9</sub> is halo.




In one embodiment, A is  $\text{C}_6\text{H}_4$ , Z, Y and X are all C, two of R<sub>5</sub> to R<sub>9</sub> are hydroxy and at least one of the remaining R<sub>5</sub> to R<sub>9</sub> is halo.




5 In one embodiment, A is  $\text{C}_6\text{H}_4$ , Z, Y and X are all C, two of R<sub>5</sub> to R<sub>9</sub> are hydroxy and at least one of the remaining R<sub>5</sub> to R<sub>9</sub> is alkyl.



In one embodiment, A is  $\text{C}_6\text{H}_4$ , Z, Y and X are all C, one of R<sub>5</sub> to R<sub>9</sub> is alkyl and at least one of the remaining R<sub>5</sub> to R<sub>9</sub> is hydroxy.




In one embodiment, A is  $\text{C}_6\text{H}_4$ , Z, Y and X are all C, one of R<sub>5</sub> to R<sub>9</sub> is halo and at least one of the remaining R<sub>5</sub> to R<sub>9</sub> is hydroxy.




In one embodiment, A is , Z, Y and X are all C, two of R<sub>5</sub> to R<sub>9</sub> is halo and at least one of the remaining R<sub>5</sub> to R<sub>9</sub> is hydroxy.


In one embodiment, the compound is selected from the group consisting of:




(T1),




(T2),




(T3),



(T4),




(T5),




(T6),


5















(T10),




(T11),




(T12),







(T31),




(T32),




(T33),




(T35),




(T37),




(T38),




(T39),




(T48),




(T58),



(T63),




(T64),



(T65),

- 14 -



According to another aspect, the present invention relates to a pharmaceutical composition comprising a compound of the present invention and a pharmaceutically acceptable excipient.

According to another aspect, the present invention relates to a method for the therapeutic treatment of hypertension or prehypertension in a subject comprising administering to the subject a compound according to the present invention.

According to another aspect, the present invention relates to a method for the therapeutic treatment of fibrosis in a subject comprising administering to the subject a compound according to the present invention.

According to another aspect, the present invention relates to a method for the prophylactic treatment of fibrosis in a subject comprising administering to the subject a compound according to the present invention.

According to another aspect, the present invention relates to a method for the therapeutic treatment of hypertension and fibrosis in a subject comprising administering to the subject a compound according to the present invention.

According to another aspect, the present invention relates to a method for the therapeutic treatment of prehypertension and fibrosis in a subject comprising administering to the subject a compound according to the present invention.

In one embodiment, the fibrosis is myocardial fibrosis or kidney fibrosis.

- 5 In another embodiment, the fibrosis is myocardial fibrosis and kidney fibrosis.

According to another aspect, the present invention relates to a compound of the present invention for use in the therapeutic treatment of hypertension or prehypertension.

According to another aspect, the present invention relates to a compound of the present invention for use in the therapeutic treatment of fibrosis.

- 10 According to another aspect, the present invention relates to a compound of the present invention for use in the prophylactic treatment of fibrosis.

According to another aspect, the present invention relates to a compound of the present invention for use in the therapeutic treatment of hypertension and fibrosis.

- 15 According to another aspect, the present invention relates to a compound of the present invention for use in the therapeutic treatment of prehypertension and fibrosis.

According to another aspect, the present invention relates to use of a compound of the present invention for the manufacture of a medicament for the therapeutic treatment of hypertension or prehypertension.

- 20 According to another aspect, the present invention relates to use of a compound of the present invention for the manufacture of a medicament for the therapeutic treatment of fibrosis.

According to another aspect, the present invention relates to use of a compound of the present invention for the manufacture of a medicament for the prophylactic treatment of fibrosis.

- 25 According to another aspect, the present invention relates to use of a compound of the present invention for the manufacture of a medicament for the therapeutic treatment of hypertension and fibrosis.

- 30 According to another aspect, the present invention relates to use of a compound of the present invention for the manufacture of a medicament for the therapeutic treatment of prehypertension and fibrosis.

Unless the context clearly requires otherwise, throughout the description and the claims, the words “comprise”, “comprising”, and the like are to be construed in an inclusive sense as opposed to an exclusive or exhaustive sense; that is to say, in the sense of “including, but not limited to”.

## 5 **BRIEF DESCRIPTION OF THE FIGURES**

Figure 1: Synthesis of 3-formylbiphenyl-4-yl trifluoromethanesulfonate.

Figure 2: Synthesis of T1, T2, T10 and T18.

Figure 3: Synthesis of diethyl(carbamoylmethyl)phosphonate.

Figure 4: Synthesis of T20.

10 Figure 5: Synthesis of T70.

Figure 6: Synthesis of T48.

Figure 7: Synthesis of 3-(3-amino-3-oxopropyl)biphenyl-4-yltrifluoromethanesulfonate.

Figure 8: Synthesis of T25.

Figure 9: Synthesis of indolone pinacol boronic ester.

15 Figure 10: Synthesis of T31.

Figure 11: Baseline normalised cell index for various compounds at three concentrations, 62.5µM (open bars), 125µM (hatched bars) and 250µM (solid bars), on rat A10 vascular smooth muscle cells as determined using the xCELLigence RTCA instrument.

20 Figure 12: Baseline normalised cell index for various compounds at three concentrations, 62.5µM (open bars), 125µM (hatched bars) and 250µM (solid bars), on bovine aortic endothelial cells as determined using the xCELLigence RTCA instrument.

Figure 13: Systolic (hatched bars) and diastolic (open bars) blood pressures in controls and treated spontaneous hypertensive rats (SHR) on a 2.2% salt diet after 4 weeks therapy. T1, T2, T20, T31 and T48 were administered at 500pmol/kg/min in the drinking solution (5% ethanol) for 4 weeks, T70 was administered at 100 pmol/kg/min in the  
25 drinking solution. \* p<0.05, \*\* p<0.01, \*\*\* p<0.005 and \*\*\*\* p<0.0005 treated systolic vs control systolic; # p<0.05, ## p<0.025 and ### p<0.005 diastolic treated vs diastolic control.

Figure 14: Relationship between baseline normalised cell index for A10 vascular smooth muscle cells and systolic blood pressure for various compounds.

Figure 15: Relationship between baseline normalised cell index for bovine aortic endothelial cells and systolic blood pressure for various compounds.

5 Figure 16: Myocardial fibrosis quantitated by computerised histomorphometry on Masson's trichrome stained histological sections in SHR on a 2.2% salt diet at 14 weeks and after 4 weeks treatment with drug in the drinking solution or vehicle control. \* p<0.005, \*\* p<0.001 and \*\*\* p<0.0005 vs 18 week vehicle treated control. # p<0.05, ## p<0.01, ### p<0.005 and ##### p<0.0005 vs 14 week control. The latter comparison  
10 indicates the ability to reverse existing pathology.

Figure 17: Interstitial fibrosis in the kidney quantitated by computerised histomorphometry on Masson's trichrome stained histological sections in SHR on a 2.2% salt diet at 14 weeks and after 4 weeks treatment with drug in the drinking solution or vehicle control. \* p<0.005, \*\* p<0.001 and \*\*\* p<0.0005 vs 18 week vehicle treated  
15 control. # p<0.05 vs 14 week control. The latter comparison indicates the ability to reverse existing pathology.

Figure 18: Relationship between baseline normalised cell index for bovine aortic endothelial cells and myocardial fibrosis for various compounds.

Figure 19: Relationship between baseline normalised cell index for bovine aortic  
20 endothelial cells and renal interstitial fibrosis for various compounds.

Figure 20: Micrographs of heart for control rats (A) and rats treated for four weeks with 500pmol/kg/min of T1 (B), T2 (C), T20 (D) or T31 (E).


Figure 21: Micrographs of kidney for control rats (A) and rats treated for four weeks with 500pmol/kg/min of T1 (B), T2 (C), T20 (D) or T31 (E).

## 25 ***DETAILED DESCRIPTION OF THE INVENTION***

The present invention relates to certain novel terphenyl compounds that show blood pressure lowering and anti-fibrotic effects in oral dosing studies in an experimental animal model. With respect to anti-fibrotic activity, the compounds of the present invention are effective in preventing fibrosis, slowing down progression of established  
30 fibrosis and/or reducing the degree (reversal) of established fibrosis. These are important


findings with respect to the range and severity of conditions which can be treated with the compounds of the present invention.

The compounds of the present invention are represented by the formula:



5 wherein:










A is selected from the group consisting of:






Q is independently selected from halo, alkyl, hydroxy, amino and substituted amino;  
 10 n is 0, 1, 2, 3, 4 or 5;


R<sub>1</sub>, R<sub>3</sub> and R<sub>4</sub> are independently C, CH, CH<sub>2</sub>, O, N, NH or S, and  
 R<sub>2</sub> is C, CH, CH<sub>2</sub>, N, NH, C-CF<sub>3</sub>, CH-CF<sub>3</sub> or C=O,  
 or a stereoisomer or pharmaceutically acceptable salt thereof,  
 wherein when n is 1, Q cannot be hydroxy.













The following compounds are specific, but non-limiting, examples of the compounds of the present invention:




5









As used herein, the term "halo" designates -F, -Cl, -Br or -I; the term "hydroxy" means -OH; the term "amino" means -NH<sub>2</sub>; and the term "substituted amino" includes -NHW, wherein W is selected from -CN, -SO<sub>2</sub>(X)<sub>a</sub>Y and -CO(X)<sub>a</sub>Y, a is 0 or 1, X is selected from -NH- and -O-, and Y is selected from -H, -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>OH and -CH<sub>2</sub>CH<sub>2</sub>OH.

As used herein, the abbreviations Me, Et, Ph, Ms represent methyl, ethyl, phenyl, and methanesulfonyl, respectively. A more comprehensive list of the abbreviations utilized by organic chemists of ordinary skill in the art appears in the first issue of each volume of the Journal of Organic Chemistry; this list is typically presented in a table entitled Standard List of Abbreviations. The abbreviations contained in said list, and all abbreviations utilized by organic chemists of ordinary skill in the art are hereby incorporated by reference.

Compounds of the present invention may exist in particular geometric or stereoisomeric forms. The present invention contemplates all such compounds, including cis- and trans-isomers, (R)- and (S)-enantiomers, diastereomers, (d)-isomers, (l)-isomers, the racemic mixtures thereof, and other mixtures thereof, as falling within the scope of the invention. All such isomers, as well as mixtures thereof, are intended to be included in this invention.

If, for instance, a particular enantiomer of a compound of the present invention is desired, it may be prepared by asymmetric synthesis, or by derivatization with a chiral auxiliary, where the resulting diastereomeric mixture is separated and the auxiliary group cleaved to provide the pure desired enantiomers. Alternatively, diastereomeric salts may be formed with an appropriate optically active acid or base, followed by resolution of the diastereomers thus formed by fractional crystallization or chromatographic means well known in the art, and subsequent recovery of the pure enantiomers.

In general, the compounds of the present invention may be prepared by the methods illustrated in the general reaction schemes as, for example, described below, or by modifications thereof, using readily available starting materials, reagents and conventional synthesis procedures. In these reactions, it is also possible to make use of  
5 variants which are in themselves known, but are not mentioned here.

The present invention also contemplates pharmaceutically acceptable salts of the compounds. The term "pharmaceutically acceptable salt" includes both acid and base addition salts and refers to salts which retain the biological effectiveness and properties of the free bases or acids, and which are not biologically or otherwise undesirable. The  
10 pharmaceutically acceptable salts are formed with inorganic or organic acids or bases, and can be prepared in situ during the final isolation and purification of the compounds, or by separately reacting a purified compound in its free base or acid form with a suitable organic or inorganic acid or base, and isolating the salt thus formed.

The term "fibrosis" as used in the context of the present invention includes, but is not  
15 limited to, myocardial fibrosis and/or kidney fibrosis.

In addition to treatment of established fibrosis, the compounds of the present invention may be used prophylactically in subjects at risk of developing fibrosis. As an example of subjects in the risk category for developing fibrosis are those having hypertension, diabetes, myocarditis, ischaemic heart disease, Conn's Syndrome, pheochromocytoma,  
20 genetic predisposition high salt diet and/or receiving drugs used in cancer chemotherapy (such as daunorubicin). The term "prophylactic" as used in the context of the present invention is intended inter alia to encompass treatments used to prevent or slow down the development of fibrosis in the at risk group. Subjects who may be given prophylactic treatment may already have signs of early heart failure on echocardiography.

25 The term "hypertension" as used in the context of the present invention indicates an adult blood pressure of above about 139 mmHg systolic and/or above about 89 mmHg diastolic.

The term "prehypertension" as used in the context of the present invention indicates an adult blood pressure in the range about 120-139 mmHg systolic and/or about 80-89  
30 mmHg diastolic.

The present invention also contemplates pharmaceutical compositions which include the compounds of the present invention, in conjunction with acceptable pharmaceutical excipients. The term “pharmaceutically acceptable excipient” as used in the context of the present invention means any pharmaceutically acceptable inactive component of the composition. As is well known in the art, excipients include diluents, buffers, binders, lubricants, disintegrants, colorants, antioxidants/preservatives, pH-adjusters, etc. The excipients are selected based on the desired physical aspects of the final form: e.g. obtaining a tablet with desired hardness and friability being rapidly dispersible and easily swallowed etc. The desired release rate of the active substance from the composition after its ingestion also plays a role in the choice of excipients. Pharmaceutical compositions may include any type of dosage form such as tablets, capsules, powders, liquid formulations, delayed or sustained release, patches, snuffs, nasal sprays and the like. The physical form and content of the pharmaceutical compositions contemplated are conventional preparations that can be formulated by those skilled in the pharmaceutical formulation field and are based on well established principles and compositions described in, for example, Remington: The Science and Practice of Pharmacy, 19th Edition, 1995; British Pharmacopoeia 2000 and similar formulation texts and manuals.

For example, where the compounds or compositions are to be administered orally, they may be formulated as tablets, capsules, granules, powders or syrups; or for parenteral administration, they may be formulated as injections (intravenous, intramuscular or subcutaneous), drop infusion preparations or suppositories. For application by the ophthalmic mucous membrane route, they may be formulated as eyedrops or eye ointments. These formulations can be prepared by conventional means, and, if desired, the active ingredient may be mixed with any conventional additive, such as an excipient, a binder, a disintegrating agent, a lubricant, a corrigent, a solubilizing agent, a suspension aid, an emulsifying agent or a coating agent.

When the compound(s) of the present invention are administered as pharmaceuticals, to humans and animals, they can be given per se or as a pharmaceutical composition containing, for example, 0.1 to 99.5% (more preferably, 0.5 to 90%) of active ingredient in combination with a pharmaceutically acceptable carrier.

The dosage of a compound and frequency of administration that should be used can also be easily determined by the practicing physician in order to produce the desired response.

Although the dosage will vary depending on the symptoms, age and body weight of the patient, the nature and severity of the disorder to be treated or prevented, the route of administration and the form of the drug, in general, a daily dosage of from 0.0001 mg to 200 mg of the compound of the present invention may be a suitable effective amount for an adult human patient, and this may be administered in a single dose or in divided doses. A “patient” or “subject” to be treated by the subject method can mean either a human or non-human subject.

An “effective amount” of a subject compound, with respect to a method of treatment, refers to an amount of the therapeutic in a preparation which, when applied as part of a desired dosage regimen provides a benefit according to clinically acceptable standards for the treatment or prophylaxis of a particular disorder.

The present invention will now be described in more detail with reference to specific but non-limiting examples describing specific compositions and methods of use. It is to be understood, however, that the detailed description of specific procedures, compositions and methods is included solely for the purpose of exemplifying the present invention. It should not be understood in any way as a restriction on the broad description of the inventive concept as set out above.

### ***EXAMPLES***

#### Example 1 – Synthesis of 3-formylbiphenyl-4-yl trifluoromethanesulfonate

The synthetic route used to prepare 3-formylbiphenyl-4-yl trifluoromethanesulfonate (14) is shown in Figure 1. Briefly, a Suzuki cross-coupling reaction between 5-bromo-2-hydroxybenzaldehyde and phenylboronic acid was used to generate 2-hydroxy-5-phenylbenzaldehyde (13), which was subsequently reacted with N-phenyltriflamide to produce 3-formylbiphenyl-4-yl trifluoromethanesulfonate (14).

#### *Production of 2-Hydroxy-5-phenylbenzaldehyde (13)*

5-Bromosalicylaldehyde (2.49 g, 12.4 mmol), phenyl boronic acid (1.51 g, 12.4 mmol), palladium(II) acetate (14 mg, 0.5 mol%) and potassium carbonate (5.14 g, 37.2 mmol) were stirred in degassed water (75 mL) at ambient temperature for 2 h, under an argon atmosphere. The reaction was monitored by TLC (1:1 dichloromethane/pentane). Water (75 mL) was added and the reaction mixture acidified (pH 6) with 10% HCl, then extracted with ethyl acetate (3x). The combined organic extracts were washed with brine,

- 26 -

then dried and concentrated. The crude material was passed through a short column of silica, eluting with 1:1 dichloromethane/pentane, then recrystallised from ethyl acetate/pentane to afford 2-hydroxy-5-phenylbenzaldehyde (1.89 g, 77%) as dark yellow crystals (can be triturated with pentane instead recrystallised if desired); mp 100-101 °C.

5 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 10.99 (s, 1H); 9.97 (s, 1H); 7.78-7.73 (m, 2H); 7.56-7.52 (m, 2H); 7.47-7.41 (m, 2H); 7.37-7.32 (m, 1H); 7.09-7.04 (m, 1H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 196.9, 161.2, 139.6, 136.0, 133.6, 132.1, 129.2, 127.6, 126.8, 121.0, 118.4. EIMS: m/z 198 [M]<sup>+</sup>. HRMS calcd for C<sub>13</sub>H<sub>10</sub>O<sub>2</sub> 198.0675, found 198.0677.

*Production of 3-formylbiphenyl-4-yl trifluoromethanesulfonate (14)*

10 2-Hydroxy-5-phenylbenzaldehyde (13) (100 mg, 0.50 mmol), *N*-phenyltriflimide (180.0 mg, 0.51 mmol) and potassium carbonate (209 mg, 1.51 mmol) were stirred in dry THF in a sealed tube, and heated at 120 °C for 6 min, using microwave irradiation. The solvent was removed under reduced pressure; water and dichloromethane were added and the layers separated. The aqueous layer was extracted further with dichloromethane (2x). The  
15 combined organic extracts were washed with brine (1x), then dried and concentrated. Purified by radial chromatography, eluting with 1: 1 dichloromethane/pentane, to afford 3-formylbiphenyl-4-yl-trifluoromethanesulfonate (143 mg, 86%) as a clear, colourless oil.  
<sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 10.32 (s, 1H); 8.17 (d, 1H, *J*=2.4 Hz); 7.89 (dd, 1H, *J*=8.6, 2.5 Hz); 7.63-7.36 (m, 6H). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 186.5, 149.1, 142.3, 138.0,  
20 134.1, 129.2, 129.1, 128.8, 128.6, 127.2, 122.9, 118.7 (q, *J*<sub>CF</sub>=320.9 Hz). <sup>19</sup>F NMR (188 MHz, CDCl<sub>3</sub>) δ -73.2. EIMS: m/z 330 [M]<sup>+</sup>. HRMS calcd for C<sub>14</sub>H<sub>9</sub>F<sub>3</sub>O<sub>2</sub>S 330.0168, found 330.0163.

Example 2 - Synthesis of T1, T2, T10 and T18

The synthetic route used to prepare T1, T2, T10 and T18 is shown in Figure 2. Briefly, 3-  
25 formylbiphenyl-4-yl trifluoromethanesulfonate (14) was cross-coupled with 3-nitrophenylboronic acid to produce nitro terphenyl (17), which then underwent a Horner-Wadsworth-Emmons reaction with diethyl(carbamoylmethyl)phosphonate (18) to produce terphenyl acrylamide (19). Hydrogenolysis of compound 19 simultaneously reduced the olefin and nitro groups to produce 3-(3-Amino-[1,1':4',1''-terphenyl]-2'-  
30 yl)propanamide (T18), which was then used to prepare 3-(3-(Methylsulfonamido)-[1,1':4',1''-terphenyl]-2'-yl)propanamide (T1) through reaction with methanesulfonyl chloride, 3-(3-Formamido-[1,1':4',1''-terphenyl]-2'-yl)propanamide (T2) through reaction

with formic acid, and 3-(3-Acetamido-[1,1':4',1''-terphenyl]-2'-yl)propanamide (T10) through reaction with acetic anhydride.

Diethyl(carbamoylmethyl)phosphonate (18) was generated from an Arbuzov reaction between 2-chloroacetamide and triethyl phosphite (prepared as shown in Figure 3)

5 *Production of 3-Nitro-[1,1':4',1''-terphenyl]-2'-carbaldehyde (17)*

3-formyl-[1,1'-biphenyl]-4-yl trifluoromethanesulfonate (14) (4.15 g, 12.60 mmol), 3-nitrophenylboronic acid (2.52 g, 15.10 mmol), potassium phosphate (4.01 g, 18.90 mmol) and tetrakis(triphenylphosphine)palladium(0) (0.33 g, 0.28 mmol) in dioxane (50 mL) were placed in a Schlenk flask, under an argon atmosphere. Degassed 1,4-dioxane (2 mL)  
10 was added and the mixture purged with argon. The reaction mixture was heated at 85 °C until complete conversion was observed (monitored by GCMS); generally required overnight reaction time. The crude material was purified by chromatography (DCVC) eluting with a gradient of ethyl acetate in heptane (0 - 25% ethyl acetate) to give 3-nitro-[1,1':4',1''-terphenyl]-2-carbaldehyde (17) as a pale tan solid (2.05 g, 67%) after recovery  
15 of un-reacted triflate (0.83 g); mp 113.6 – 116.3°C (NB: Product was contaminated with ~25% by <sup>1</sup>H NMR of 3,3'-dinitro-1,1'-biphenyl). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 10.02 (s, 1H), 8.29 (m, 3H), 7.92 (dd, 1H, *J* 8.0, 2.1 Hz), 7.72 (m, 1H), 7.66 (m, 3H), 7.50 (m, 3H), 7.42 (m, 1H). <sup>13</sup>C NMR (100 MHz, DMSO-*d*<sub>6</sub>) δ 191.7, 147.8, 140.8, 140.5, 139.1, 138.4, 136.4, 133.8, 132.0, 131.9, 130.6, 129.9, 128.3, 127.0, 126.8, 124.2, 122.8. EIMS:  
20 *m/z* Found: M<sup>+</sup> 303.0880, C<sub>19</sub>H<sub>13</sub>NO<sub>3</sub> requires 303.0890. EIMS: *m/z* 303 (M<sup>+</sup>, 100%), 256 (52).

*Production of (E)-3-(3-Nitro-[1,1':4',1''-terphenyl]-2'-yl)acrylamide (19)*

3-Nitro-[1,1':4',1''-terphenyl]-2-carbaldehyde (17) (2.35 g, 7.77 mmol) and diethyl(carbamoylmethyl)phosphonate (18) (1.51 g, 7.75 mmol) were dissolved in dry  
25 THF (100 mL), and added slowly to a vigorously stirred suspension of powdered potassium hydroxide (0.86 g, 15.40 mmol). Following stirring for 1 h at rt, the material was precipitated from the reaction mixture by addition of water and diethyl ether to give (E)-3-(3-nitro-[1,1':4',1''-terphenyl]-2'-yl)acrylamide (19) (1.8 g, 82%) as a pale lemon solid. A small portion was purified by chromatography (DCVC) eluting with a gradient  
30 of ethyl acetate in DCM (0 – 20% ethyl acetate) for characterisation to give (E)-3-(3-nitro-[1,1':4',1''-terphenyl]-2'-yl)acrylamide (19) as a colourless solid; mp 206 - 210°C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 8.35 – 8.28 (m, 1H), 8.18 – 8.15 (m, 1H), 8.02 – 7.98

- 28 -

(m, 1H), 7.85 – 7.76 (m, 5H), 7.56 – 7.41 (m, 4H), 7.49 (br s, 1H), 7.33 (d, 1H,  $J$  15.7 Hz), 7.15 (br s, 1H), 6.78 (d, 1H,  $J$  15.7 Hz).  $^{13}\text{C}$  NMR (100 MHz, DMSO- $d_6$ )  $\delta$  166.2, 147.8, 140.9, 140.6, 139.1, 138.3, 136.5, 136.2, 133.4, 131.1, 130.0, 129.0, 128.0, 127.8, 126.8, 125.0, 124.8, 123.8, 122.5. EIMS:  $m/z$  Found:  $M^{++}$  344.1153,  $\text{C}_{21}\text{H}_{16}\text{N}_2\text{O}_3$  requires  
5 344.1155. EIMS:  $m/z$  344 ( $M^{++}$ , 37%), 326 (50), 252 (100).

*Production of 3-(3-Amino-[1,1':4',1''-terphenyl]-2'-yl)propanamide (T18)*

To a solution of (*E*)-3-(3-nitro-[1,1':4',1''-terphenyl]-2'-yl)acrylamide (19) (1.70 g, 4.94 mmol) in methanol (50 mL) and ethyl acetate (25 mL) was added 10% palladium on carbon (50% wt water). The reaction mixture was stirred vigorously in an autoclave  
10 under hydrogen at 140psi for 2 hours. The reaction mixture was filtered through Celite washing well with methanol and ethyl acetate. The filtrate was concentrated then pre-absorbed onto Celite and chromatographed (DCVC) eluting with a gradient of methanol in DCM (0 – 3% methanol). Fractions containing a single spot on TLC were combined to give 3-(3-amino-[1,1':4',1''-terphenyl]-2'-yl)propanamide (T18) as a colourless solid (0.92  
15 g, 59%); mp 157.3 – 157.9 °C.  $^1\text{H}$  NMR (400 MHz, DMSO- $d_6$ )  $\delta$  7.69 (d, 2H,  $J$  7.4 Hz), 7.59 (s, 1H), 7.51 – 7.46 (m, 3H), 7.37 (m, 1H), 7.23 (br s, 1H), 7.19 (d, 1H,  $J$  7.9 Hz), 7.08 (m, 1H), 6.74 (br s, 1H), 6.57 (d, 1H,  $J$  8.4 Hz), 6.52 (s, 1H), 6.46 (d, 1H,  $J$  7.5 Hz), 5.13 (br s, 2H), 2.84 (m, 2H), 2.31 (m, 2H).  $^{13}\text{C}$  NMR (100 MHz, DMSO- $d_6$ )  $\delta$  173.5, 148.5, 141.5, 141.3, 140.1, 139.1, 128.8, 130.1, 128.9, 128.7, 127.3, 127.1, 126.6, 124.0,  
20 116.5, 114.4, 112.6, 36.4, 28.3. EIMS:  $m/z$  Found:  $M^{++}$  316.1566,  $\text{C}_{21}\text{H}_{20}\text{N}_2\text{O}$  requires 316.1570. EIMS:  $m/z$  316 ( $M^{++}$ , 100%). HPLC purity (40% ACN /  $\text{H}_2\text{O}$ , 258 nm): 100.0%.

*Production of 3-(3-(Methylsulfonamido)-[1,1':4',1''-terphenyl]-2'-yl)propanamide (T1)*

To a suspension of 3-(3-amino-[1,1':4',1''-terphenyl]-2'-yl)propanamide (T18) (0.50 g,  
25 1.57 mmol) in DCM (7 mL), cooled to -5 °C, was added triethylamine (0.33 mL, 2.36 mmol), followed by dropwise addition of methanesulfonyl chloride (0.21 g, 1.83 mmol) at such a rate as to maintain the temperature below 0 °C (~20 minutes). The reaction mixture was partitioned between 2M hydrochloric acid and ethyl acetate and the layers separated. The organic phase was washed again with 2M hydrochloric acid, saturated  
30 bicarbonate solution and brine. The crude material was pre-absorbed onto Celite and chromatographed (DCVC) eluting with a gradient of methanol in DCM (0 – 3% methanol). Like fractions were combined to give 3-(3-(methylsulfonamido)-[1,1':4',1''-

terphenyl]-2'-yl)propanamide (T1) as colourless fine needles (0.25 g, 41%); mp 166.7 – 168.4 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.12 (br s, 1H), 7.58 (m, 2H), 7.53 (m, 1H), 7.47 – 7.31 (m, 5H), 7.27 – 7.24 (m, 2H), 7.19 (m, 1H), 7.12 (m, 1H), 5.87 (br s, 1H), 5.78 (br s, 1H), 2.99, (s, 3H), 2.94 (m, 2H), 2.43 (m, 2H). <sup>13</sup>C NMR (100 MHz, DMSO-*d*<sub>6</sub>) δ 173.3, 141.8, 139.9 (two signals coincident), 139.5, 139.1, 138.4, 130.3, 129.3, 128.9, 127.5, 127.4, 126.7, 124.4, 124.3, 119.9, 118.2, 39.3, 36.2, 28.3. EIMS: *m/z* Found: M<sup>+</sup> 394.1341, C<sub>22</sub>H<sub>22</sub>N<sub>2</sub>O<sub>3</sub>S requires 394.1346. EIMS: *m/z* 394 (M<sup>+</sup>, 12%), 376 (22), 256 (100). HPLC purity (40% ACN / H<sub>2</sub>O, 256 nm): 99.84%.

*Production of 3-(3-Formamido-[1,1':4',1''-terphenyl]-2'-yl)propanamide (T2)*

10 A solution of 3-(3-amino-[1,1':4,1''-terphenyl]-2'-yl)propanamide (T18) (0.41 g, 1.30 mmol) in formic acid (5 mL) was heated at reflux for 5 hours, then concentrated to dryness. The crude material was pre-absorbed onto Celite then chromatographed (DCVC) eluting with a gradient of methanol in DCM (0 – 5% methanol). Like fractions were combined to give 3-(3-formamido-[1,1':4,1''-terphenyl]-2'-yl)propanamide (T2) as  
15 colourless solid (0.21 g, 47%); mp 213 °C. Existed as a mixture of *E* and *Z* amide isomers. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 10.30 (s) and 10.22 (d, *J* 11.0 Hz; 1H), 8.88 (d, *J* 11.0 Hz) and 8.31 (d, *J* 1.8 Hz; 1H), 7.70 (m, 2H), 7.63 – 7.19 (m, 10H), 7.08 (m, 1H), 6.76 (br s, 1H), 2.83 (m, 2H), 2.32 (m, 2H). <sup>13</sup>C NMR (50 MHz, DMSO-*d*<sub>6</sub>) δ 173.4, 162.7, 159.7, 142.0, 141.4, 140.2, 140.0, 139.9, 139.4, 139.3, 139.1, 138.3, 138.2, 130.2,  
20 129.3, 128.9, 128.8, 127.5, 127.3, 126.7, 124.3, 124.2, 119.6, 117.8, 117.7, 116.0, 36.2, 28.2 (a number of signals were coincident). EIMS: *m/z* Found: M<sup>+</sup> 344.1518, C<sub>22</sub>H<sub>20</sub>N<sub>2</sub>O<sub>2</sub> requires 344.1519. EIMS: *m/z* 344 (M<sup>+</sup>, 20%), 299 (34), 254 (100). HPLC purity (50% ACN / H<sub>2</sub>O, 255 nm): 99.53%.

*Production of 3-(3-Acetamido-[1,1':4',1''-terphenyl]-2'-yl)propanamide (T10)*

25 A solution of 3-(3-amino-[1,1':4,1''-terphenyl]-2'-yl)propanamide (T18) (0.42 g, 1.33 mmol) and *N,N*-dimethylaminopyridine (0.04 g, cat.) in acetic anhydride (7 mL) was stirred at ambient temperature for 20 hours. The reaction mixture was partitioned between water and ethyl acetate. The layers were separated and the organic phase washed with water (2x) and brine and filtered through a plug of silica gel 60, washing well with  
30 ethyl acetate. The filtrate was concentrated to dryness and recrystallised from methanol and 1,2-dichloroethane to give 3-(3-acetamido-[1,1':4,1''-terphenyl]-2'-yl)propanamide (T10) as a beige solid (0.36 g, 75%); mp 208 – 209 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ

- 30 -

10.03 (br s, 1H), 7.70 (m, 2H), 7.62 (m, 2H), 7.58 – 7.47 (m, 4H), 7.40 – 7.35 (m, 2H),  
7.23 (m, 2H), 7.02 (d, 1H, 7.7 Hz), 6.75 (br s, 1H), 2.83 (m, 2H), 2.31 (m, 2H), 2.06 (s,  
3H). <sup>13</sup>C NMR (100 MHz, DMSO-*d*<sub>6</sub>) δ 173.4, 168.4, 141.2, 140.4, 139.9, 139.3, 139.2,  
139.1, 130.2, 128.9, 128.6, 127.4, 127.3, 126.7, 124.2, 123.6, 119.4, 117.5, 36.2, 28.2,  
5 24.1. EIMS: *m/z* Found: M<sup>+</sup> 358.1666, C<sub>23</sub>H<sub>22</sub>N<sub>2</sub>O<sub>2</sub> requires 358.1676. EIMS: *m/z* 358  
(M<sup>+</sup>, 8%), 299 (33), 254 (100). HPLC purity (50% ACN / H<sub>2</sub>O, 255 nm): 99.53%.

### Example 3 - Synthesis of T20

The synthetic route used to prepare T20 is shown in Figure 4. Briefly, 3-formylbiphenyl-  
4-yl trifluoromethanesulfonate (14) was cross-coupled with 3,5-dimethoxyphenylboronic  
10 acid to produce dimethoxy terphenyl (20), which then underwent a Horner-Wadsworth-  
Emmons reaction with diethyl(carbamoylmethyl)phosphonate (18) to afford terphenyl  
acrylamide (21). Hydrogenolysis of compound 21 yielded propanamide (22), which,  
subsequently, was demethylated using boron tribromide to afford T20.

#### *Production of 3,5-Dimethoxy-[1,1':4',1''-terphenyl]-2'-carbaldehyde (20)*

15 To a solution of 3,5-dimethoxyphenylboronic acid (4.0 g, 22.0 mmol), 3-formylbiphenyl-  
4-yl trifluoromethanesulfonate (14) (6.6 g, 20.0 mmol) and sodium carbonate (47.2 g,  
40.0 mmol) in degassed dioxane/ethanol/H<sub>2</sub>O (5:1:1, 165 mL) was added  
tetrakis(triphenylphosphine)palladium(0) (1.16 g, 1.0 mmol). The reaction was heated at  
110 °C for 2 hours in a sealed tube. Analysis by TLC (1:2 DCM/PE) indicated the triflate  
20 had been consumed. The reaction was concentrated, then taken up in water and extracted  
with ethyl acetate (3x). The combined organic extracts were washed with water and brine  
then dried (MgSO<sub>4</sub>) and concentrated. The crude material was filtered through a short  
column of silica, eluting with 1:1 DCM:PE to afford 3,5-dimethoxy-[1,1':4',1''-terphenyl]-  
2'-carbaldehyde (20) (6.1 g, 96%) as a pale yellow solid. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ  
25 10.09 (s, 1H), 8.26 (d, 1H, *J* 1.8 Hz), 7.87 (dd, 1H, *J* 2.1, 8.0 Hz), 7.68 (m, 2H), 7.58 -  
7.35 (m, 4H), 6.56 (s, coincident, 3H), 3.84 (s, 6H). <sup>13</sup>C NMR (50 MHz, CDCl<sub>3</sub>) δ 192.4,  
160.9, 144.8, 140.9, 139.6, 134.2, 132.0, 131.1, 129.1, 128.1, 127.2, 125.8, 108.6, 100.2,  
55.6 (one signal not observed). EIMS: *m/z* Found: M<sup>+</sup> 318.1255, C<sub>21</sub>H<sub>18</sub>O<sub>3</sub> requires  
318.1250. EIMS: *m/z* 318 (M<sup>+</sup>, 55%).

- 31 -

*Production of (E/Z)-3-(3,5-Dimethoxy-[1,1':4',1''-terphenyl]-2'-yl)acrylamide (21)*

3,5-Dimethoxy-[1,1':4',1''-terphenyl]-2'-carbaldehyde (20) (6.1 g, 19.1 mmol) and diethyl (carbamoylmethyl)phosphonate (18) (3.7 g, 19.1 mmol) were dissolved in dry THF (180 mL), and added slowly to a vigorously stirred suspension of powdered KOH (2.1 g, 38.2 mmol) in THF (70 mL). The reaction was stirred at rt for 1 h under an argon atmosphere. Analysis by TLC (1:2 DCM:PE) indicated the carbaldehyde had been consumed. The THF was removed under reduced pressure, and the residue taken up in water and extracted with DCM (x3). The combined organic extracts were washed with brine (x1) then dried (MgSO<sub>4</sub>) and concentrated to ~50 mL. The solution was filtered through a short column of silica eluting with DCM to afford (E/Z)-3-(3,5-dimethoxy-[1,1':4',1''-terphenyl]-2'-yl)acrylamide (21) (2.5 g, 36%) as an orange foam. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 7.85 (d, 1H, *J* 1.7 Hz); 7.78 – 7.56 (m, 4H); 7.53 – 7.32 (m, 4H); 6.54 – 6.38 (m, 4H); 5.70 (brs, 2H); 3.81 (s, 6H). <sup>13</sup>C NMR (50 MHz, CDCl<sub>3</sub>) δ 167.9, 160.8, 141.8, 141.7, 141.4, 140.9, 140.5, 133.4, 130.9, 129.1, 128.4, 127.9, 127.3, 125.7, 121.5, 108.3, 100.1, 55.7. EIMS: *m/z* Found: M<sup>+</sup> 359.1504, C<sub>23</sub>H<sub>21</sub>O<sub>3</sub>N requires 359.1516. EIMS: *m/z* 359 (M<sup>+</sup>, 3%).

*Production of 3-(3,5-Dimethoxy-[1,1':4',1''-terphenyl]-2'-yl)propanamide (22)*

(E/Z)-3-(3,5-Dimethoxy-[1,1':4',1''-terphenyl]-2'-yl)acrylamide (21) (2.5 g, 6.9 mmol) and 10% palladium on carbon (50% wt in H<sub>2</sub>O, 1.0 g) in methanol (100 mL) were stirred at rt under a hydrogen atmosphere at 50 psi for 2h. The reaction mixture was gravity filtered through GF paper washing thoroughly with methanol, then concentrated. The residue was then taken up in DCM and gravity filtered through GF paper, washing thoroughly with DCM, then concentrated. The crude material was then filtered through a short column of silica, washing thoroughly with DCM, then eluting the desired compound with 1:49 methanol:DCM to afford 3-(3,5-dimethoxy-[1,1':4',1''-terphenyl]-2'-yl)propanamide (22) (2.2 g, 90%) as a white solid. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 7.70 – 7.24 (m, 8H); 6.50 (m, 3H); 5.78 (br s, 1H); 5.34 (br s, 1H); 3.82 (s, 6H); 3.05 (m, 2H); 2.39 (m, 2H). <sup>13</sup>C NMR (50 MHz, CDCl<sub>3</sub>) δ 174.8, 160.8, 143.3, 141.0, 140.8, 140.8, 138.6, 130.6, 129.0, 128.1, 127.6, 127.2, 125.1, 107.6, 99.2, 55.5, 37.2, 29.2. EIMS: *m/z* Found: M<sup>+</sup> 361.1672, C<sub>23</sub>H<sub>23</sub>O<sub>3</sub>N requires 361.1672. EIMS: *m/z* 361 (M<sup>+</sup>, 100%).

*Production of 3-(3,5-Dihydroxy-[1,1':4',1''-terphenyl]-2'-yl)propanamide (T20)*

A solution of 3-(3,5-dimethoxy-[1,1':4',1''-terphenyl]-2'-yl)propanamide (22) (500 mg, 1.4 mmol) was dissolved in dry DCM (5 mL) and cooled to -78 °C, under an argon atmosphere. Boron tribromide (2.9 mL, 2.9 mmol, 1.0 M solution in hexanes) was added and the reaction allowed to warm to rt overnight. The solution was cooled (ice/water bath) and water (5 mL) and methanol (2 mL) slowly added. The layers were separated and the aqueous phase extracted further with DCM (x2). The combined organic extracts were washed with 1.0 M sodium thiosulfate (x1), water (x1) and brine (x1), then dried (MgSO<sub>4</sub>) and concentrated. Purified by radial chromatography using a gradient elution (DCM → 4:96 methanol:DCM → 6:94 methanol:DCM → 8:92 methanol:DCM to afford 3-(3,5-dihydroxy-[1,1':4',1''-terphenyl]-2'-yl)propanamide (T20) (122 mg, 26%) as a white solid; mp 232 – 233 °C. <sup>1</sup>H NMR (200 MHz, DMSO-*d*<sub>6</sub>) δ 9.32 (brs, 2H); 7.72 – 7.64 (m, 2H); 7.58 (d, 1H, *J* 1.8 Hz); 7.53 – 7.33 (m, 4H); 7.24 (overlap, brs, 1H); 7.18 (overlap, d, 1H, *J* 7.9 Hz); 6.75 (brs, 1H); 6.23 (t, 1H, *J* 2.1); 6.15 (d, 2H, *J* 2.1 Hz); 2.84 (m, 2H); 2.31 (m, 2H). <sup>13</sup>C NMR (100 MHz, DMSO-*d*<sub>6</sub>) δ 173.5, 158.1, 142.6, 140.9, 140.0, 139.0, 138.9, 129.9, 128.9, 127.3, 127.1, 126.6, 124.1, 107.1, 101.2, 36.3, 28.2. EIMS: *m/z* Found: M<sup>+</sup> 333.1344, C<sub>21</sub>H<sub>19</sub>O<sub>3</sub>N requires 333.1359. EIMS: *m/z* 333 (M<sup>+</sup>, 94%). HPLC purity (40% ACN / H<sub>2</sub>O, 264 nm): 95.97%.

Example 4 - Synthesis of T70

The synthetic route used to prepare T70 is shown in Figure 5. Briefly, 3-formyl-[1,1'-biphenyl]-4-yl trifluoromethanesulfonate (14) was cross-coupled with 3,5-dinitrophenyl pinacol boronic ester (34) [prepared from a reaction between 1-iodo-3,5-dinitrobenzene and bis(pinacolato)diborane] to provide 3,5-dinitroterphenyl (35). A subsequent Horner-Wadsworth-Emmons reaction with diethyl(carbamoylmethyl)phosphonate (18) afforded 3,5-dinitroterphenyl acrylamide (36). Compound 36 was then hydrogenated to give propanamide (37), which was reacted with methanesulfonyl chloride to produce T70.

*Production of 2-(3,5-Dinitrophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (34).*

1-Iodo-3,5-dinitrobenzene (5.00 g, 17.00 mmol), bispinacolatodiboron (4.75 g, 18.7 mmol), potassium acetate (5.00 g, 51.00 mmol) and dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (0.35 g, 0.48 mmol) in DMSO (80 mL) was stirred at 70 °C for 17 hours. The reaction mixture was cooled to ambient temperature, diluted with ethyl acetate then washed with saturated

sodium bicarbonate solution and brine. The crude material was pre-absorbed onto Celite, then chromatographed (DCVC) eluting with a gradient of ethyl acetate in heptane (0 – 100% ethyl acetate). Like fractions were combined to give 2-(3,5-dinitrophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (34) as a pale yellow solid (2.10 g, 40%); mp 144.0 – 148.0°C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.08 (t, 1H, *J* 2.2 Hz), 8.90 (d, 2H, *J* 2.2 Hz), 1.37, (s, 12H).

*Production of 3,5-Dinitro-[1,1':4',1''-terphenyl]-2'-carbaldehyde (35).*

Prepared according to the method of P5; from 3-formyl-[1,1'-biphenyl]-4-yl trifluoromethanesulfonate (14) (1.77 g, 5.36 mmol), 2-(3,5-dinitrophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (34) (1.81 g, 6.16 mmol), tetrakis(triphenylphosphine)palladium(0) (0.44 g, 0.38mmol) and aqueous sodium carbonate (1M) (11.0 mL, 11.0 mmol) in toluene (36 mL) and ethanol (7 mL). A solid was filtered from the interface during extraction which was found to be the desired product (0.86 g, 46%). The ethyl acetate extract was purified by chromatography (DCVC) eluting with a gradient of dichloromethane in heptane (10 – 50% DCM) to give a further amount of 3,5-dinitro-[1,1':4',1''-terphenyl]-2'-carbaldehyde (35) as a pale tan solid (0.62 g, 33%) (Total yield: 79%); mp 209 – 212 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 10.04 (s, 1H), 8.90 (s, 1H), 8.74 (s, 2H), 8.33 (s, 1H), 8.14 (d, 1H, *J* 8.0 Hz), 7.83 (d, 2H, *J* 7.3 Hz), 7.72 (d, 1H, *J* 8.0 Hz), 7.56 (m, 2H), 7.47 (m, 1H). <sup>13</sup>C NMR (100 MHz, DMSO-*d*<sub>6</sub>) δ 192.0, 147.9, 141.2, 138.2, 138.1, 134.0, 132.3, 131.8, 130.0, 129.3, 128.5 (two signals coincident), 126.9, 117.7 (one signal not observed). EIMS: *m/z* Found: M<sup>+</sup> 348.0731, C<sub>19</sub>H<sub>12</sub>N<sub>2</sub>O<sub>5</sub> requires 348.0741. EIMS: *m/z* 348 (M<sup>+</sup>, 100%).

*Production of (E)-3-(3,5-Dinitro-[1,1':4',1''-terphenyl]-2'-yl)acrylamide (36).*

Prepared according to the method used to generate compound 19; from 3,5-dinitro-[1,1':4',1''-terphenyl]-2'-carbaldehyde (35) (1.75 g, 5.03 mmol), diethyl(carbamoylmethyl)phosphonate (18) (1.09 g, 5.59 mmol) and sodium hydroxide (0.50 g, 12.50 mmol) in THF (70 mL). The crude solid was recrystallised from acetone to give (E)-3-(3,5-dinitro-[1,1':4',1''-terphenyl]-2'-yl)acrylamide (36) as a pale tan solid (1.40 g, 72%); mp 221 – 223 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 8.90 (s, 1H), 8.58 (s, 2H), 8.02 (s, 1H), 7.85 (d, 1H, *J* 8.0 Hz), 7.80 (d, 2H, *J* 7.6 Hz), 7.64 (d, 1H, *J* 8.0 Hz), 7.56 – 7.43 (m, 4H), 7.32 (d, 1H, <sup>3</sup>*J*<sub>trans</sub> 15.7 Hz), 7.16 (br s, 1H), 6.76 (d, 1H, <sup>3</sup>*J*<sub>trans</sub> 15.7 Hz). <sup>13</sup>C NMR (100 MHz, DMSO-*d*<sub>6</sub>) δ 166.1, 148.0, 142.4, 141.3, 138.9, 136.2, 135.9,

- 34 -

133.8, 131.3, 129.7, 129.1, 128.2, 127.8, 126.9, 126.1, 125.3, 117.6. EIMS:  $m/z$  Found:  $M^{++}$  389.1000,  $C_{21}H_{15}N_3O_5$  requires 389.1006. EIMS:  $m/z$  389 ( $M^{++}$ , 42%), 252 (100).

*Production of 3-(3,5-Diamino-[1,1':4',1''-terphenyl]-2'-yl)propanamide (37).*

Prepared according to the method used to generate T18; from (*E*)-3-(3,5-dinitro-  
5 [1,1':4',1''-terphenyl]-2'-yl)acrylamide (36) (1.40 g, 3.60 mmol) and 10% palladium on carbon (50% wt water) (0.28 g) in methanol (40 mL). Catalyst was removed by filtration and the filtrate concentrated to dryness to give 3-(3,5-diamino-[1,1':4',1''-terphenyl]-2'-yl)propanamide (37) as a tan solid (1.07 g, 90%); mp 87.4 – 90.6°C.  $^1H$  NMR (400 MHz, DMSO- $d_6$ )  $\delta$  7.68 (m, 2H), 7.65 (m, 1H), 7.49 – 7.45 (m, 3H), 7.36 (m, 1H), 7.22 (br s, 1H), 7.15 (d, 1H,  $J$  7.9 Hz), 6.76 (br s, 1H), 5.83 (m, 1H), 5.75 (m, 2H), 4.79 (br s, 4H),  
10 2.86 (m, 2H), 2.31 (m, 2H).  $^{13}C$  NMR (100 MHz, DMSO- $d_6$ )  $\delta$  173.7, 148.9, 142.2, 141.9, 140.2, 139.0, 138.4, 129.9, 128.9, 127.2, 127.0, 126.6, 123.8, 104.0, 98.7, 36.6, 28.4. EIMS:  $m/z$  Found:  $M^{++}$  331.1678,  $C_{21}H_{21}N_3O$  requires 331.1679. EIMS:  $m/z$  331 ( $M^{++}$ , 67%), 287 (100), 273 (72).

15 *Production of 3-(3,5-Di(methylsulfonamido)-[1,1':4',1''-terphenyl]-2'-yl)propanamide (T70).*

Prepared according to the method used to generate T1; from 3-(3,5-diamino-[1,1':4',1''-terphenyl]-2'-yl)propanamide (37) (0.46 g, 1.38 mmol), methanesulfonyl chloride (2.56 mL, 3.30 mmol) and triethylamine (0.58 mL, 4.14 mmol) in DCM (15 mL). The crude  
20 material was purified by chromatography (DCVC) eluting with a gradient of methanol in DCM (0 – 5% methanol) and then radial chromatography eluting with 3% methanol in DCM to give 3-(3,5-dimethylsulfonamido)-[1,1':4',1''-terphenyl]propanamide (T70) as a beige solid (0.15 g, 22%); mp 227 – 230 °C.  $^1H$  NMR (400 MHz, DMSO- $d_6$ )  $\delta$  9.96 (s, 2H), 7.73 – 7.66 (m, 2H), 7.63 (d, 1H,  $J$  1.9 Hz), 7.55 (dd, 1H,  $J$  1.9, 7.9 Hz), 7.52 – 7.45  
25 (m, 2H), 7.42 – 7.36 (m, 1H), 7.26 (d, 1H,  $J$  7.9 Hz), 7.24 (br s, 1H), 7.19 -7.15 (m, 1H), 6.91 (d, 2H,  $J$  1.9 Hz), 6.77 (br s, 1H), 3.06 (s, 6H), 2.83 (t, 2H,  $J$  8.0 Hz), 2.32 (t, 2H,  $J$  8.0 Hz).  $^{13}C$  NMR (50 MHz, DMSO- $d_6$ )  $\delta$  173.3, 142.6, 139.8, 139.6, 139.5, 139.3, 139.1, 130.1, 128.9, 127.5 (two signals coincident), 126.7, 124.4, 114.8, 108.8, 39.3, 36.3, 28.3. EIMS:  $m/z$  Found:  $M^{++}$  487.1226,  $C_{23}H_{25}N_3O_5^{32}S_2$  requires 487.1230. EIMS:  $m/z$   
30 487 ( $M^{++}$ , 4%), 408 (75), 349 (100), 271 (78). HPLC purity (40% ACN /  $H_2O$ , 264 nm): 94.72%.

Example 5 - Synthesis of T48

The synthetic route used to prepare T48 is shown in Figure 6. Briefly, 3-formyl-[1,1'-biphenyl]-4-yl trifluoromethanesulfonate (14) was cross-coupled with pyridyl pinacol boronic ester (30 - Ihle, N. C.; Krause, A. E. *J. Org. Chem.* 1996, 61, 4810) to produce  
5 teraryl (31), which then underwent a Horner-Wadsworth-Emmons reaction with diethyl(carbamoylmethyl)phosphonate (18) to afford teraryl acrylamide (32). Hydrogenation of compound 32 yielded propanamide (33), which was subsequently deprotected to afford T48.

*Production of tert-Butyl (4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridin-2-yl)carbamate (30).*

2-Aminopyridine-4-boronic acid pinacol ester (2.0 g, 9.1 mmol) was stirred as a suspension in *tert*-butanol (30 mL), under an argon atmosphere. The Boc anhydride (2.20 g, 10.0 mmol) in *tert*-butanol (20 mL) was added slowly, and the reaction stirred at 35 °C for 18 hours. Analysis by <sup>1</sup>H NMR showed the pinacol ester starting material had been  
15 consumed. The reaction mixture was concentrated under reduced pressure, and the crude material stirred in water for 5 minutes. The solid was collected by filtration and dried *in vacuo* at 50 °C, to afford *tert*-butyl (4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridin-2-yl)carbamate (31) as a white solid (2.9 g, 98%); mp 172 – 178.0°C. (Lit. 188 – 193 °C). <sup>1</sup>H NMR (200 MHz, DMSO-*d*<sub>6</sub>) δ 9.75 (br s, 1H), 8.26 (dd, 1H, *J* 0.9, 4.8 Hz),  
20 8.08 (m, 1H), 7.18 (dd, 1H, *J* 0.7, 4.8 Hz), 1.47 (s, 9H), 1.31 (s, 12H).

*Production of tert-Butyl (4-(3-formyl-[1,1'-biphenyl]-4-yl)pyridin-2-yl)carbamate (31).*

To a solution of *tert*-butyl (4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridin-2-yl)carbamate (30) (2.9 g, 8.9 mmol), 3-formylbiphenyl-4-yl trifluoromethanesulfonate (14) (2.7 g, 8.1 mmol) and sodium carbonate (1.7 g, 16.2 mmol) in degassed  
25 dioxane/ethanol/H<sub>2</sub>O mixture (5:1:1, 75 mL) was added tetrakis(triphenylphosphine)palladium(0) (467 mg, 0.40 mmol). The reaction was heated at 110 °C for 2 hours in a sealed tube. Analysis by <sup>1</sup>H NMR indicated the triflate had been consumed. The reaction was concentrated, then taken up in DCM and poured into water. The layers were separated and the aqueous phase extracted further with DCM  
30 (2x). The combined organic extracts were washed with water (x1) and brine then dried and concentrated to approx. 20 – 30 mL volume. The solution was filtered through a short column of silica eluting with DCM to afford *tert*-butyl (4-(3-formyl-[1,1'-biphenyl]-

- 36 -

4-yl)pyridin-2-yl)carbamate (31) as a yellow solid (1.5 g, 48%); mp 168.8 – 171.5 °C. <sup>1</sup>H NMR (200 MHz, DMSO-*d*<sub>6</sub>) δ 10.00 (overlap s, 1H), 9.98 (overlap, br s, 1H), 8.35 (dd, 1H, *J* 0.7, 5.1 Hz), 8.20 (d, 1H, *J* 1.9 Hz), 8.10 (dd, 1H, *J* 2.1, 8.0 Hz), 7.88 (m, 1H), 7.79 (m, 2H), 7.63 (d, 1H, *J* 8.0 Hz), 7.59 - 7.40 (m, 3H), 7.16 (dd, 1H, *J* 1.6, 5.1 Hz), 1.47 (s, 9H). <sup>13</sup>C NMR (50 MHz, DMSO-*d*<sub>6</sub>) δ 191.4, 152.8, 152.6, 147.8, 146.8, 141.2, 140.7, 138.4, 133.6, 132.0, 131.2, 129.2, 128.3, 126.8, 126.1, 119.4, 112.8, 79.7, 28.0. EIMS: *m/z* Found: M<sup>+</sup> 374.1611, C<sub>23</sub>H<sub>22</sub>O<sub>3</sub>N<sub>2</sub> requires 374.1625. EIMS: *m/z* 374 (M<sup>+</sup>, 7%), 57 (100).

*Production of (E)-tert-Butyl (4-(3-(3-amino-3-oxoprop-1-en-1-yl)-[1,1'-biphenyl]-4-yl)pyridin-2-yl)carbamate (32).*

Prepared according to the method used to generate compound 19; from *tert*-butyl (4-(3-formyl-[1,1'-biphenyl]-4-yl)pyridin-2-yl)carbamate (31) (1.44 g, 3.85 mmole), diethyl (carbamoylmethyl)phosphonate (18) (0.75 g, 3.85 mmole) and sodium hydroxide (0.31 g, 7.70 mmol) in THF (40 mL). (*E*)-*tert*-Butyl (4-(3-(3-amino-3-oxoprop-1-en-1-yl)-[1,1'-biphenyl]-4-yl)pyridin-2-yl)carbamate (32) was precipitated from the reaction mixture upon addition of water and diethyl ether as a colourless solid (1.32 g, 83%); mp 179.5 – 182.2 °C. <sup>1</sup>H NMR (200 MHz, DMSO-*d*<sub>6</sub>) δ 9.95 (s, 1H); 8.33 (m, 1H); 7.99 (d, 1H, *J* 1.6 Hz); 7.82-7.73 (m, 4H); 7.58-7.42 (m, 5H); 7.34 (d, 1H, *J* 15.8 Hz); 7.14 (br s, 1H); 6.99 (dd, 1H, *J* 1.5, 5.1 Hz); 6.77 (d, 1H, *J* 15.7 Hz); 1.46 (s, 9H). <sup>13</sup>C NMR (50 MHz, DMSO-*d*<sub>6</sub>) δ 166.3, 152.7, 149.0, 147.7, 140.6, 139.2, 138.7, 136.4, 133.1, 130.4, 129.0, 128.0, 127.7, 126.8, 124.6, 124.6, 119.2, 112.5, 79.7, 28.0. EIMS: *m/z* Found: M<sup>+</sup> 415.1873, C<sub>22</sub>H<sub>25</sub>O<sub>3</sub>N<sub>3</sub> requires 415.1890. EIMS: *m/z* 415 (M<sup>+</sup>, 5%), 315 (58), 297 (64), 271 (100).

*Production of tert-Butyl (4-(3-(3-amino-3-oxopropyl)-[1,1'-biphenyl]-4-yl)pyridin-2-yl)carbamate (33).*

Prepared according to the method used to prepare T18; from (*E*)-*tert*-butyl (4-(3-(3-amino-3-oxoprop-1-en-1-yl)-[1,1'-biphenyl]-4-yl)pyridin-2-yl)carbamate (32) (1.17 g, 2.80 mmol) and 10% palladium on carbon (50% wt water) (0.50 g) in methanol (75 mL). The filtrate was concentrated to give *tert*-butyl (4-(3-(3-amino-3-oxopropyl)-[1,1'-biphenyl]-4-yl)pyridin-2-yl)carbamate (33) as a colourless solid (1.05 g, 89%); mp 161.5 – 164.5 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 9.90 (s, 1H), 8.32 - 8.29 (m, 1H), 7.78 (s, 1H), 7.73 - 7.69 (m, 2H), 7.66 (s, 1H), 7.61 - 7.56 (m, 1H), 7.53 - 7.47 (m, 2H), 7.43 -

- 37 -

7.37 (m, 1H), 7.30 - 7.26 (m, 1H), 7.25 (br s, 1H), 7.09 - 7.05 (m, 1H), 6.76 (br s, 1H),  
2.88-2.81 (m, 2H), 2.36-2.29 (m, 2H), 1.47 (s, 9H). <sup>13</sup>C NMR (50 MHz, DMSO-*d*<sub>6</sub>) δ  
173.2, 152.8, 152.6, 150.3, 147.6, 140.1, 139.7, 138.9, 138.2, 129.8, 128.9, 127.6, 127.5,  
126.7, 124.5, 118.8, 112.4, 79.6, 36.1, 28.04, 28.00. EIMS: *m/z* Found: M<sup>+</sup> 417.2028,  
5 C<sub>25</sub>H<sub>27</sub>N<sub>3</sub>O<sub>3</sub> requires 417.2047. EIMS: *m/z* 417 (M<sup>+</sup>, 5%), 317 (15), 284 (89), 258 (100).

*Production of 3-(4-(2-Aminopyridin-4-yl)-[1,1'-biphenyl]-3-yl)propanamide (T48).*

A mixture of *tert*-butyl (4-(3-(3-amino-3-oxopropyl)-[1,1'-biphenyl]-4-yl)pyridin-2-  
yl)carbamate (33) (0.94 g, 2.26 mmol) and TFA (7.0 mL) in DCM (10mL) was stirred at  
10 ambient temperature for 3 hours. The reaction mixture was partitioned with iced water  
and ethyl acetate, then neutralised with sodium hydroxide (~pH 6), then basified to pH 10  
with 1M sodium carbonate solution. The crude material was collected by filtration then  
recrystallised from methanol to give 3-(4-(2-aminopyridin-4-yl)-[1,1'-biphenyl]-3-  
yl)propanamide (T48) as a colourless solid (0.48 g, 67 %); mp 248 – 249 °C. <sup>1</sup>H NMR  
(400 MHz, DMSO-*d*<sub>6</sub>) δ 7.95 (d, 1H, *J* 5.2 Hz), 7.73 – 7.66 (m, 2H), 7.62 (s, 1H), 7.57 –  
15 7.52 (m, 1H), 7.52 -7.45 (m, 2H), 7.42 -7.35 (m, 1H), 7.27 (br s, 1H), 7.24 – 7.20 (m,  
1H), 6.78 (br s, 1H), 6.48 (d, 1H, *J* 5.2 Hz), 6.39 (s, 1H), 5.99 (s, 2H), 2.84 (t, 2H, *J* 7.9  
Hz), 2.33 (t, 2H, *J* 7.9 Hz). <sup>13</sup>C NMR (100 MHz, DMSO-*d*<sub>6</sub>) δ 173.3, 159.9, 149.4, 147.7,  
139.8, 139.7, 139.0, 138.9, 129.6, 128.9, 127.5, 127.4, 126.7, 124.4, 112.7, 107.8, 36.2,  
28.1. EIMS: *m/z* Found: M<sup>+</sup> 317.1516, C<sub>20</sub>H<sub>19</sub>N<sub>3</sub>O requires 317.1523. EIMS: *m/z* 317  
20 (M<sup>+</sup>, 12%), 273 (53), 258 (100). HPLC purity (35% ACN / 0.1% TFA, 291 nm):  
98.76%.


Example 6 - Synthesis of T3, T11, T12 and T15 from T18

A solution of 3-(3-amino-1,1':4',1''-terphenyl-2'-yl)propanamide (T18 - produced in  
Example 2) (1 equiv.) in dichloromethane (12.5 mL/mmol) was added to a solution of  
25 triphosgene (0.3 equiv) in dichloromethane (6.25 mL/mmol). Triethylamine (0.3  
mL/mmol) was added and the mixture stirred at room temperature under nitrogen for 30  
min. An amine or alcohol (2-5 equiv.) was added and the mixture stirred at room  
temperature under nitrogen. The neat reaction mixture was purified by flash  
chromatography (methanol/dichloromethane). The product was suspended in 1:1  
30 dichloromethane/hexanes and isolated by filtration.

The following compounds were prepared by this procedure:


- 38 -

3-{3-[(Methylcarbamoyl)amino]-1,1':4',1''-terphenyl-2'-yl}propanamide (T3)




White powder (82 mg, 27%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 8.59 (s, 1H), 7.70 (d, *J*=7.2 Hz, 2H), 7.61 (d, *J*=1.6 Hz, 1H), 7.43 - 7.56 (m, 4H), 7.26 - 7.42 (m, 3H), 7.23 (d, *J*=7.8 Hz, 2H), 6.87 (d, *J*=7.2 Hz, 1H), 6.72 (br. s., 1H), 6.06 (br. q, *J*=4.5 Hz, 1H), 2.79 - 2.87 (m, 2H), 2.64 (d, *J*=4.7 Hz, 3H), 2.27 - 2.35 (m, 2H); LCMS [M+H]<sup>+</sup> = 374.2; HPLC (water/ACN + 0.1% TFA gradient) 100% at 220nm.

Methyl [2'-(3-amino-3-oxopropyl)-1,1':4',1''-terphenyl-3-yl]carbamate (T11)



White powder (131 mg, 55%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 9.74 (s, 1H), 7.70 (d, *J*=7.4 Hz, 2H), 7.62 (d, *J*=1.4 Hz, 1H), 7.42 - 7.58 (m, 5H), 7.32 - 7.41 (m, 2H), 7.23 (s, 2H), 6.99 (d, *J*=7.4 Hz, 1H), 6.73 (br. s., 1H), 3.68 (s, 3H), 2.78 - 2.87 (m, 2H), 2.26 - 2.35 (m, 2H); LCMS [M+H]<sup>+</sup> = 375.3; HPLC (water/ACN + 0.1% TFA gradient) 99.4% at 220nm.

2-Hydroxyethyl [2'-(3-amino-3-oxopropyl)-1,1':4',1''-terphenyl-2'-yl]carbamate (T12)




White powder (129 mg, 50%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 9.79 (s, 1H), 7.70 (d, *J*=7.2 Hz, 2H), 7.62 (d, *J*=1.6 Hz, 1H), 7.42 - 7.58 (m, 5H), 7.31 - 7.41 (m, 2H), 7.15 - 7.27 (m, 2H), 6.98 (d, *J*=7.6 Hz, 1H), 6.73 (br. s., 1H), 4.81 (t, *J*=5.3 Hz, 1H), 4.11 (t,

- 39 -

$J=5.1$  Hz, 2H), 3.63 (q,  $J=5.3$  Hz, 2H), 2.76 - 2.88 (m, 2H), 2.25 - 2.37 (m, 2H); LCMS  $[M+H]^+ = 405.1$ ; HPLC (water/ACN + 0.1% TFA gradient) 99.4% at 220nm.


*3-[3-(Carbamoylamino)-1,1':4',1''-terphenyl-2'-yl]propanamide (T15)*



5 White powder (74 mg, 32%).  $^1\text{H NMR}$  (400MHz, DMSO- $d_6$ ) 8.62 (s, 1H), 7.70 (d,  $J=7.4$  Hz, 2H), 7.61 (d,  $J=1.6$  Hz, 1H), 7.43 - 7.56 (m, 4H), 7.26 - 7.42 (m, 3H), 7.18 - 7.26 (m, 2H), 6.88 (d,  $J=7.0$  Hz, 1H), 6.72 (br. s., 1H), 5.87 (s, 2H), 2.78 - 2.87 (m, 2H), 2.27 - 2.36 (m, 2H); LCMS  $[M+H]^+ = 360.3$ ; HPLC (water/ACN + 0.1% TFA gradient) 97.2% at 220nm.

10 Example 7 - Synthesis of T4 from T18

Ethyl isocyanate (50  $\mu\text{L}$ , 0.63 mmol) was added to a solution of 3-(3-amino-1,1':4',1''-terphenyl-2'-yl)propanamide (155 mg, 0.49 mmol) (T18 - produced in Example 2) in dichloromethane (10 mL). The mixture was stirred at room temperature under nitrogen for 3 days. The reaction mixture was evaporated to dryness. The residue was dissolved in  
 15 a mixture of dichloromethane (10 mL) and methanol (2 mL), adsorbed onto silica gel 60, and purified by flash chromatography (methanol/dichloromethane). The product was suspended in 1:1 dichloromethane/hexanes and isolated by filtration to give 3-{3-  
 [(Ethylcarbamoyl) amino]-1,1':4',1''-terphenyl-2'-yl} propanamide (T4):



20 White powder (115 mg, 60%).  $^1\text{H NMR}$  (400MHz, DMSO- $d_6$ ) 8.50 (s, 1H), 7.70 (d,  $J=7.4$  Hz, 2H), 7.61 (d,  $J=1.6$  Hz, 1H), 7.42 - 7.57 (m, 4H), 7.25 - 7.42 (m, 3H), 7.18 - 7.26 (m, 2H), 6.87 (d,  $J=7.2$  Hz, 1H), 6.72 (br. s., 1H), 6.14 (t,  $J=5.5$  Hz, 1H), 3.03 - 3.18

- 40 -


(m, 2H), 2.77 - 2.90 (m, 2H), 2.24 - 2.38 (m, 2H), 1.05 (t,  $J=7.1$  Hz, 3H); LCMS  $[M+H]^+$  = 388.3; HPLC (water/ACN + 0.1% TFA gradient) 98.5% at 220nm.

#### Example 8 - Synthesis of T5 and T6 from T18

3-(3-Amino-1,1':4',1''-terphenyl-2'-yl)propanamide (T18 - produced in Example 2) (1  
 5 equiv.) and triethylamine (1.3-2.0 equiv.) were dissolved in dichloromethane (18  
 mL/mmol). A solution of an alkylsulfamoyl chloride (1.3- 2.0 equiv.) in dichloromethane  
 (4 mL/mmol) was added drop wise. The mixture was stirred at room temperature under  
 nitrogen for 1 h. The neat reaction mixture was purified by flash chromatography  
 (methanol/dichloromethane). The product was suspended in 1:1 dichloromethane/hexanes  
 10 and isolated by filtration.


The following compounds were prepared by this procedure:

#### 3-{3-[(Methylsulfamoyl)amino]-1,1':4',1''-terphenyl-2'-yl}propanamide (T5)



White powder (60 mg, 30%).  $^1\text{H NMR}$  (400MHz, DMSO- $d_6$ ) 9.77 (br. s., 1H), 7.70 (d,  
 15  $J=7.2$  Hz, 2H), 7.62 (d,  $J=1.6$  Hz, 1H), 7.54 (dd,  $J=7.9, 1.7$  Hz, 1H), 7.49 (t,  $J=7.6$  Hz,  
 2H), 7.31 - 7.42 (m, 3H), 7.22 - 7.29 (m, 2H), 7.13 - 7.21 (m, 2H), 6.99 (d,  $J=7.6$  Hz, 1H),  
 6.78 (br. s., 1H), 2.75 - 2.88 (m, 2H), 2.48 (obscured by DMSO- $d_6$ ), 2.27 - 2.36 (m, 2H);  
 LCMS  $[M+H]^+$  = 410.2; HPLC (water/ACN + 0.1% TFA gradient) 97.4% at 220nm.

#### 3-{3-[(Ethylsulfamoyl)amino]-1,1':4',1''-terphenyl-2'-yl}propanamide (T6)



20


White powder (69 mg, 34%).  $^1\text{H NMR}$  (400MHz, DMSO- $d_6$ ) 9.72 (br. s., 1H), 7.70 (d,  
 $J=7.2$  Hz, 2H), 7.62 (d,  $J=1.6$  Hz, 1H), 7.43 - 7.58 (m, 4H), 7.31 - 7.42 (m, 2H), 7.20 -

- 41 -

7.29 (m, 2H), 7.11 - 7.20 (m, 2H), 6.97 (d,  $J=7.6$  Hz, 1H), 6.77 (br. s., 1H), 2.76 - 2.97 (m, 4H), 2.26 - 2.37 (m, 2H), 0.98 (t,  $J=7.2$  Hz, 3H); LCMS  $[M+H]^+ = 424.3$ ; HPLC (water/ACN + 0.1% TFA gradient) 99.6% at 220nm.

Example 9 - Synthesis of T16 from T18

5 3-(3-Amino-1,1':4',1''-terphenyl-2'-yl)propanamide (T18 - produced in Example 2) (181 mg, 0.57 mmol) was dissolved in methanol (3.8 mL) with gentle heating. Potassium acetate (170 mg, 1.73 mmol) was added and the mixture cooled in an ice-water bath. A solution of cyanogen bromide (61 mg, 0.58 mmol) in methanol (1.1 mL) was added drop wise. The mixture was stirred in the ice-water bath under nitrogen for 1 h, then at room  
10 temperature under nitrogen overnight. The reaction mixture was evaporated to dryness. The residue was dissolved in 10% methanol/dichloromethane (60 mL). The organic phase washed with water (3×20 mL) and brine (20 mL), dried over anhydrous sodium sulfate, and filtered. The filtrate was evaporated to dryness. The residue was purified by flash chromatography (methanol/dichloromethane). The product was dissolved in ethyl acetate  
15 (20 mL) and the organic phase washed with hydrochloric acid (1 M, 3×20 mL) and brine (20 mL), dried over anhydrous sodium sulfate, and filtered. The filtrate was evaporated to dryness. The residue was suspended in 1:1 dichloromethane/hexanes and isolated by filtration. The resultant product was purified by flash chromatography (methanol/dichloromethane). The product was suspended in 1:1 dichloromethane/hexanes  
20 and isolated by filtration to give 3-[3-(Cyanoamino)-1,1':4',1''-terphenyl-2'-yl]propanamide (T16):



White powder (68 mg, 35%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 10.30 (br. s., 1H), 7.70 (d,  $J=7.2$  Hz, 2H), 7.63 (d,  $J=1.6$  Hz, 1H), 7.55 (dd,  $J=7.8, 1.8$  Hz, 1H), 7.35 - 7.52 (m, 4H),  
25 7.20 - 7.30 (m, 2H), 7.04 (d,  $J=7.6$  Hz, 1H), 6.99 (dd,  $J=8.0, 1.8$  Hz, 1H), 6.88 (s, 1H), 6.75 (br. s., 1H), 2.82 (t,  $J=7.8$  Hz, 2H), 2.25 - 2.35 (m, 2H); LCMS  $[M+H]^+ = 342.3$ ; HPLC (water/ACN + 0.1% TFA gradient) 97.6% at 220nm.

Example 10 - Synthesis of 3-(3-Amino-3-oxopropyl)biphenyl-4-yl trifluoromethanesulfonate

The synthesis of 3-(3-Amino-3-oxopropyl)biphenyl-4-yl trifluoromethanesulfonate (7) is shown in Figure 7.

5 *Production of 4-Hydroxybiphenyl-3-carbaldehyde (2)*

5-Bromosalicylaldehyde (**1**) (50.00 g, 0.249 mol), K<sub>2</sub>CO<sub>3</sub> (103.13 g, 0.746 mol), phenylboronic acid (30.33 g, 0.249 mmol), and Pd(OAc)<sub>2</sub> (0.28 g, 1.2 mmol) were added to a magnetically stirred 2L round bottom flask containing freshly degassed H<sub>2</sub>O (1.5 L, degassed by purging with N<sub>2</sub> (4 x 2.5L balloons)). The reaction mixture was stirred under  
10 N<sub>2</sub> overnight and analysed by TLC (**2** observed but **1** was still present). The reaction mixture was stirred for an additional 24 h before pouring carefully into HCl (aq., 0.2M, 3 L) over several hours maintaining the pH of the mixture (~pH 2) by addition of small amounts of HCl (aq., 33%). The mixture was then stirred for 1 h with EtOAc (500 mL) and filtered through Celite into a separatory funnel. The organic layer was collected and  
15 the aqueous layer extracted with EtOAc (500 mL, washing through the Celite filtrate) and the two organic layers combined, dried over MgSO<sub>4</sub> and concentrated to give a yellow solid residue (64 g). The residue was taken up in hot EtOH (200 mL) and H<sub>2</sub>O (200 mL) added slowly with vigorous stirring and allowed to cool to room temperature over 48 h. The resulting precipitate was collected by vacuum filtration and washed with H<sub>2</sub>O/EtOH  
20 (1:1, 200 mL) and air dried to give the crude biphenyl **2** (41.97 g, containing 25 mol% **1** as an impurity) as a pale yellow solid. The crude biphenyl **2** (39.8 g, containing approx. 0.050 mol of **1**), phenylboronic acid (6.02 g, 0.050 mol) and K<sub>2</sub>CO<sub>3</sub> (40.76 g, 0.295 mol) were added to H<sub>2</sub>O (1.0 L) in a 2L round bottom flask with magnetic stirring. The reaction mixture was purged with N<sub>2</sub> (2 x 2.5 L balloons, over 15 min) before adding  
25 Pd(OAc)<sub>2</sub> (223 mg, 1.0 mmol) and slowly heating to reflux for 3 h under N<sub>2</sub>. Additional phenylboronic acid (1.2 g, 9.84 mmol) was added and stirring continued at reflux for 4 h, then cooled to room temperature and left to stand over the weekend. The mixture was poured into HCl (aq., 3.3 M, 1.5 L) over 1 min and stirred well for 10 min before collecting the solid by vacuum filtration and sucking to dry for 30 min. The solid was  
30 transferred to a vacuum desiccator and dried overnight to give 39.7 g of a 12:1 mixture of biphenyl **2** (mass equiv. 36.6 g, 74%) and **1** (equiv. to an 8 mol% impurity). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 7.08 (d, *J*=8.61 Hz, 1 H), 7.32 - 7.39 (m, 1 H), 7.45 (t, *J*=7.43 Hz, 2

- 43 -

H), 7.55 (d,  $J=7.43$  Hz, 2 H), 7.72 - 7.80 (m, 2 H), 9.93 - 10.00 (m, 1 H), 11.01 (s, 1 H).

*Production of 4-(Benzyloxy)biphenyl-3-carbaldehyde (3)*

A magnetically stirred mixture of phenol **2** (38.10 g, 0.192 mol), K<sub>2</sub>CO<sub>3</sub> (33.78 g, 0.250 mol) and benzylbromide (29.7 mL, 0.250 mol) in CH<sub>3</sub>CN (370 mL) in a 500 mL round  
5 bottom flask was heated slowly to 70°C for 3 h and analysed by TLC (silica, 10% EtOAc/hexane, visualised by UV). The TLC showed the reaction was progressing but some phenol **2** remained. The reaction mixture was heated to reflux for 2 h and then analysed by TLC (reaction complete, no phenol **2** observed). The reaction mixture was cooled to room temperature and transferred to a 1L conical flask and carefully acidified  
10 with HCl (aq., 2M, 200 mL, some effervescence observed, continued until pH < 2). Water was added (200 mL) and extracted with EtOAc (3 x 500 mL). The extracts were dried over MgSO<sub>4</sub> and concentrated to give a light brown solid. The solid was suspended in hexane (150 mL) and stirred vigorously for 10 min before collecting the product by vacuum filtration and washing with hexane (2 x 60 mL) to give compound **3** as a light  
15 brown powder (44.50 g, 80%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 5.25 (s, 2 H), 7.13 (d,  $J=9.00$  Hz, 1 H), 7.29 - 7.39 (m, 2 H), 7.39 - 7.49 (m, 6 H), 7.57 (d,  $J=7.43$  Hz, 2 H), 7.77 (dd,  $J=8.61, 2.35$  Hz, 1 H), 8.10 (d,  $J=2.35$  Hz, 1 H), 10.60 (s, 1 H).

*Production of (2E)-3-[4-(Benzyloxy)biphenyl-3-yl]prop-2-enoic acid (4)*

Piperidine (2.2 mL, 0.022 mol) was added to a magnetically stirred mixture of aldehyde  
20 (**3**) (44.5 g, 0.154 mol) and malonic acid (19.25 g, 0.185 mol) in pyridine (250 mL) and slowly heated to a gentle reflux for 5 h. Effervescence was noticed as the reaction temperature approached 90°C. TLC of the reaction mixture (silica, 10% EtOAc/hexane, visualised by UV) showed only a faint spot corresponding to the starting aldehyde **3** and an intense spot of fluorescent material on the baseline, corresponding to the product (**4**).  
25 The reaction was cooled to room temperature and concentrated on the rotovap (60°C). EtOAc (200 mL) and HCl (aq., 2M, 200 mL) was added to give a thick slurry of white paste. The solid (compound **4**) was collected by vacuum filtration and the biphasic filtrate transferred to a separatory funnel. The organic phase was collected, washed with HCl (aq., 2M, 1 x 100 mL), H<sub>2</sub>O (2 x 200 mL) and brine (1 x 75 mL), dried over MgSO<sub>4</sub> and  
30 concentrated to provide additional compound **4** as a pale yellow-brown solid. The two crops of compound **4** were combined and dried in a vacuum desiccator to give compound **4** (48.5 g, 95%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 5.23 (s, 2 H), 6.64 (d,  $J=16.04$  Hz, 1 H),

- 44 -

7.03 (d,  $J=8.61$  Hz, 1 H), 7.34 (d,  $J=5.48$  Hz, 2 H), 7.38 - 7.49 (m, 6 H), 7.55 (d,  $J=7.43$  Hz, 3 H), 7.68 - 7.85 (m, 1 H), 8.22 (d,  $J=16.04$  Hz, 1 H).

*Production of (2E)-3-[4-(Benzyloxy)biphenyl-3-yl]prop-2-enamide (5)*

Oxalyl chloride (25 mL, 0.29 mol) was added slowly via a dropping funnel over 30 min  
5 to a magnetically stirred mixture of carboxylic acid **4** (48.2 g, 0.146 mol) and DMF (0.8 mL) in CH<sub>2</sub>Cl<sub>2</sub> (500 mL) in a 1L 3 necked round bottom flask equipped with a dropping funnel, a stopper and an oil bubbler. The temperature of the reaction was maintained by placing the vessel in a water bath for the duration of the addition. Upon addition of approximately 2/3 of the oxalyl chloride, the reaction mixture became homogenous with  
10 the disappearance of the suspended solid. The reaction mixture was allowed to stir for an additional 1 h before concentrating the reaction mixture on the rotary evaporator (60°C) to give the intermediate acid chloride as a yellow solid. The yellow intermediate acid chloride was suspended in a magnetically stirred solution of 1,4-dioxane (200 mL) and a solution of NH<sub>3</sub> (31 mL, 28% in H<sub>2</sub>O, 0.438 mol) in 1,4-dioxane (200 mL) was added  
15 over 15 min. The temperature of the reaction was maintained by placing the vessel in a water bath for the duration of the addition. A thick slurry resulted. The slurry was stirred at room temperature for an additional 30 min before pouring the mixture into a 1L conical flask and then adding H<sub>2</sub>O to give a final volume of 1 L. The slurry was stirred for 5 min and the solid collected by vacuum filtration, washing the solid with H<sub>2</sub>O (2 x 300 mL).  
20 The solid was dried in a vacuum desiccator overnight and then on the rotary evaporator (60°C, approx. 1-5 mmHg) to give the first crop of compound **5** (41.6 g, 86%) as an off white powder. The aqueous filtrates were concentrated to dryness, H<sub>2</sub>O (200 mL) added and the solid collected by vacuum filtration to give a second crop of compound **5** (8.12 g) as an off white powder. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 5.23 (s, 2 H), 6.61 - 6.67 (m, 1 H),  
25 7.04 (d,  $J=8.61$  Hz, 1 H), 7.31 - 7.38 (m, 3 H), 7.44 (m, 7 H), 7.55 (m, 2 H), 7.78 (d,  $J=1.57$  Hz, 1 H), 8.22 (d,  $J=16.04$  Hz, 1 H).

*Production of 3-(4-Hydroxybiphenyl-3-yl)propanamide (6)*

Compound **5** (41.55 g, 0.126 mol) was suspended in a magnetically stirred mixture of EtOAc (1 L) and NEt<sub>3</sub> (1.5 mL). N<sub>2</sub> (3 x 1L balloons) was bubbled through the mixture  
30 before adding Pd/C (10% w/w, 4.15 g) and placing the flask under vacuum briefly before back filling the atmosphere with H<sub>2</sub> from a balloon. The balloon was recharged with fresh H<sub>2</sub> and opened to the reaction mixture and stirred for 6 h, recharging the balloon with H<sub>2</sub>

approximately every 1-2 h (3 times) and then stirred overnight. The balloon was again recharged with H<sub>2</sub> and the reaction mixture slowly heated to reflux for 3 h and then cooled to room temperature and stirred under H<sub>2</sub> for 4 days. The balloon was again recharged with H<sub>2</sub> and the reaction mixture slowly heated to reflux for 3 h before cooling and then bubbling N<sub>2</sub> (2 x 1L balloons) through the reaction mixture. The reaction mixture was filtered through Celite, washing the Celite pad with EtOAc (2 x 150 mL) and the filtrate concentrated on the rotary evaporator (60°C) to give a yellow oil. Et<sub>2</sub>O was added to the yellow oil and then removed on the rotary evaporator to give a pale yellow powder (crude **6**). The TLC (silica, 70% EtOAc/hexane) of the pale yellow powder showed several products. The yellow powder was stirred vigorously in hexane (150 mL) for 30 min and the solid collected by vacuum filtration, washing with hexane (2 x 30 mL) to give compound **6** (30.22 g, containing approx. 15 mol% unknown impurity) as pale yellow powder. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 2.69 - 2.75 (m, 2 H), 2.94 - 3.01 (m, 2 H), 5.57 (br. s., 2 H), 6.99 (d, *J*=8.22 Hz, 1 H), 7.29 (d, *J*=2.35 Hz, 2 H), 7.33 - 7.43 (m, 4 H), 7.53 (d, *J*=7.43 Hz, 2 H).

*Production of 3-(3-Amino-3-oxopropyl)bi(phenyl)-4-yl trifluoromethanesulfonate (7)*

1,1,1-Trifluoro-*N*-phenyl-*N*-[(trifluoromethyl)sulfonyl]methanesulfonamide (PhNTf<sub>2</sub>) (42.5 g, 0.119 mol) was added to a solution of **6** (24.0 g, 0.0995 mol) and NEt<sub>3</sub> (15.3 mL, 0.109 mol) in CH<sub>3</sub>CN (480 mL) and the reaction mixture was stirred for 1.5 h. Additional PhN(Tf)<sub>2</sub> (2.25 g, 6.30 mmol) and NEt<sub>3</sub> (1.5 mL, 10.7 mmol) was added and the reaction mixture stirred for a further 30 min. The reaction showed only a slight trace of remaining **6**. The reaction mixture was concentrated on the rotary evaporator (60°C) to give an orange oil. A small aliquot of the oil was taken up in EtOAc (15 mL) and washed with Na<sub>2</sub>CO<sub>3</sub> (aq., 2M, 2 x 20 mL) and NaOH (aq., 0.5 M, 2 x 20 mL), dried over MgSO<sub>4</sub> and concentrated to give a crop of crude **7** (422 mg). Subsequent HPLC analysis indicated the desired product was in the organic phase. This first crop of crude **7** was recombined with the orange oil and taken up in EtOAc (300 mL), washed with Na<sub>2</sub>CO<sub>3</sub> (aq., 2M, 2 x 250 mL), dried over MgSO<sub>4</sub> and concentrated on the rotary evaporator (60°C) to give an orange oil (65 g, <sup>1</sup>H NMR showed significant impurities including NEt<sub>3</sub>). This oil was redissolved in EtOAc (300 mL) and washed with citric acid (aq., 10% w/w, 2 x 250 mL) and water (2 x 350 mL), dried over MgSO<sub>4</sub> and concentrated on the rotary evaporator (60°C) to give an orange oil (59 g). This oil was again taken up in EtOAc (300 mL) and

- 46 -


washed with NaOH (aq., 0.5 M, 3 x 200 mL), HCl (2M, 2 x 200 mL) and H<sub>2</sub>O (1 x 300 mL), dried over MgSO<sub>4</sub> and concentrated on the rotary evaporator (60°C) to give an orange oil that solidified upon standing. This solid was suspended in Et<sub>2</sub>O (150 mL) and stirred vigorously for 30 min, collected by vacuum filtration and washed with Et<sub>2</sub>O (2 x 30 mL) to give compound **7** as a white powder (12.3 g, 37%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 2.60 (t, *J*=7.63 Hz, 2 H), 3.13 (t, *J*=7.83 Hz, 2 H), 5.48 (br. s., 2 H), 7.32 (d, *J*=8.61 Hz, 1 H), 7.36 - 7.42 (m, 1 H), 7.45 (t, *J*=7.43 Hz, 2 H), 7.50 (dd, *J*=8.61, 1.96 Hz, 1 H), 7.54 (d, *J*=7.43 Hz, 2 H), 7.59 (d, *J*=1.96 Hz, 1 H).

#### Example 11 - Synthesis of T22 and T23

T22 and T23 were prepared from 3-(3-Amino-3-oxopropyl)biphenyl-4-yl trifluoromethanesulfonate (**7** - produced in Example 10). A mixture of **7** (1 equiv.), a heterocyclic boronic acid (1.2 equiv.) and potassium carbonate (2 equiv.) was suspended in 1,4-dioxane (4 mL/mmol) and water (5 drops/mmol). Nitrogen was bubbled through the mixture for 15 min. Tetrakis(triphenylphosphine)palladium(0) (0.1 equiv) was added and the mixture heated at 85°C under nitrogen for 20 h. The mixture was diluted with ethyl acetate and filtered. The residue was washed with ethyl acetate (2×). The combined filtrates were evaporated to dryness and purified by flash chromatography (methanol/dichloromethane). The product was suspended in hexanes (4 mL) and isolated by filtration.

The following compounds were prepared by this procedure:

#### *3-[4-(1H-Indol-4-yl)biphenyl-3-yl]propanamide (T22)*




Pale beige powder (58 mg, 32%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>) 8.27 (br. s., 1H), 7.64 - 7.71 (m, 2H), 7.58 - 7.63 (m, 1H), 7.54 (dd, *J*=7.8, 1.8 Hz, 1H), 7.40 - 7.50 (m, 4H), 7.36 (br. t, *J*=7.5 Hz, 1H), 7.28 (d, *J*=8.2 Hz, 1H), 7.22 (t, *J*=2.7 Hz, 1H), 7.03 - 7.09 (m, 1H),

- 47 -

6.27 - 6.33 (m, 1H), 4.96 (br. s., 1H), 4.88 (br. s., 1H), 2.99 (br. s., 2H), 2.23 (t,  $J=7.9$  Hz, 2H); LCMS  $[M+H]^+ = 341.2$ ; HPLC (water/ACN + 0.1% TFA gradient) 97.1% at 220nm

*3-[4-(1H-Indol-6-yl)biphenyl-3-yl]propanamide (T23)*




- 5 Pale brown powder (13 mg, 7%).  $^1\text{H NMR}$  (400MHz,  $\text{CDCl}_3$ ) 8.27 (br. s., 1H), 7.69 (d,  $J=8.0$  Hz, 1H), 7.64 (d,  $J=7.2$  Hz, 2H), 7.56 (d,  $J=1.6$  Hz, 1H), 7.47 - 7.53 (m, 1H), 7.42 - 7.46 (m, 2H), 7.32 - 7.41 (m, 3H), 7.12 (dd,  $J=8.1, 1.3$  Hz, 1H), 6.60 (br. s., 1H), 5.06 (br. s., 2H), 3.04 - 3.16 (m, 2H), 2.29 - 2.40 (m, 2H); LCMS  $[M+H]^+ = 341.3$ ; HPLC (water/ACN + 0.1% TFA gradient) 99.5% at 220nm.

10 Example 12 - Synthesis of T29, T38, T63, T64, T65 and T66


T29, T38, T63, T64, T65 and T66 were prepared from 3-(3-Amino-3-oxopropyl) biphenyl-4-yl trifluoromethanesulfonate (**7** - produced in Example 10). A mixture of **7** (1 equiv.), an aromatic boronic acid or heterocyclic boronic acid pinacol ester (1.1 equiv.), and potassium carbonate (2-3 equiv.) was dissolved in a mixture of 1,4-dioxane (3.1 mL/mmol), ethanol (0.65 mL/mmol) and water (0.65 mL/mmol). Nitrogen was bubbled through the mixture for 10 min. Tetrakis(triphenylphosphine)palladium(0) (0.1 equiv) was added and the mixture heated at 85°C under nitrogen for 20 h. The mixture was partitioned between ethyl acetate and water. The aqueous phase was extracted with ethyl acetate. The combined ethyl acetate extracts were washed with water and brine, dried over anhydrous sodium sulfate, and filtered. The filtrate was evaporated to dryness and purified by flash chromatography (methanol/dichloromethane). The product was suspended in 1:1 dichloromethane/hexanes and isolated by filtration

The following compounds were prepared by this procedure:


- 48 -

*3-[4-(2-Oxo-2,3-dihydro-1,3-benzothiazol-5-yl)biphenyl-3-yl]propanamide (T29)*

Light brown powder (39 mg, 11%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 11.96 (br. s., 1H), 7.70 (d, *J*=7.2 Hz, 2H), 7.60 - 7.67 (m, 2H), 7.55 (dd, *J*=7.9, 1.7 Hz, 1H), 7.49 (t, *J*=7.6 Hz, 2H), 7.35 - 7.43 (m, 1H), 7.20 - 7.31 (m, 2H), 7.09 - 7.16 (m, 1H), 7.05 (d, *J*=1.2 Hz, 1H), 6.75 (br. s., 1H), 2.84 (t, *J*=7.8 Hz, 2H), 2.25 - 2.37 (m, 2H); LCMS [M+H]<sup>+</sup> = 375.1; HPLC (water/ACN + 0.1% TFA gradient) 98.6% at 220nm.

*3-[4-(1H-Indazol-6-yl)biphenyl-3-yl]propanamide (T38)*


Pale yellow powder (72 mg, 44%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 13.09 (s, 1H), 8.12 (s, 1H), 7.82 (d, *J*=8.2 Hz, 1H), 7.71 (d, *J*=7.4 Hz, 2H), 7.65 (d, *J*=1.4 Hz, 1H), 7.56 (dd, *J*=8.0, 1.8 Hz, 1H), 7.43 - 7.53 (m, 3H), 7.35 - 7.43 (m, 1H), 7.32 (d, *J*=7.8 Hz, 1H), 7.21 (br. s., 1H), 7.11 (dd, *J*=8.3, 0.9 Hz, 1H), 6.70 (br. s., 1H), 2.77 - 2.93 (m, 2H), 2.25 - 2.38 (m, 2H); LCMS [M+H]<sup>+</sup> = 342.1; HPLC (water/ACN + 0.1% TFA gradient) 98.5% at 220nm.

*3-(2-Fluoro-3-hydroxy-1,1':4',1''-terphenyl-2'-yl)propanamide (T63).*

- 49 -


White powder (87 mg, 53%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 9.92 (br. s., 1H), 7.70 (d, *J*=7.2 Hz, 2H), 7.64 (d, *J*=1.6 Hz, 1H), 7.54 (dd, *J*=7.9, 1.7 Hz, 1H), 7.49 (t, *J*=7.6 Hz, 2H), 7.35 - 7.42 (m, 1H), 7.16 - 7.29 (m, 2H), 7.03 - 7.10 (m, 1H), 6.94 - 7.02 (m, 1H), 6.65 - 6.78 (m, 2H), 2.72 (t, *J*=7.8 Hz, 2H), 2.28 (t, *J*=7.9 Hz, 2H); LCMS [M+H]<sup>+</sup> =  
5 336.2; HPLC (water/ACN + 0.1% TFA gradient) 99.3% at 220nm.

*3-(4-Fluoro-3-hydroxy-1,1':4',1''-terphenyl-2'-yl)propanamide (T64)*




White powder (98 mg, 61%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 9.95 (br. s., 1H), 7.69 (d, *J*=7.2 Hz, 2H), 7.60 (d, *J*=1.6 Hz, 1H), 7.43 - 7.55 (m, 3H), 7.33 - 7.42 (m, 1H), 7.12 -  
10 7.28 (m, 3H), 6.90 (dd, *J*=8.5, 2.1 Hz, 1H), 6.65 - 6.80 (m, 2H), 2.76 - 2.87 (m, 2H), 2.25 - 2.36 (m, 2H); LCMS [M+H]<sup>+</sup> = 336.2; HPLC (water/ACN + 0.1% TFA gradient) 99.6% at 220nm.

*3-(3-Fluoro-5-hydroxy-1,1':4',1''-terphenyl-2'-yl)propanamide (T65)*




15 White powder (89 mg, 55%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 10.05 (br. s., 1H), 7.69 (d, *J*=7.4 Hz, 2H), 7.61 (d, *J*=1.6 Hz, 1H), 7.44 - 7.56 (m, 3H), 7.34 - 7.42 (m, 1H), 7.20 - 7.31 (m, 2H), 6.76 (br. s., 1H), 6.54 - 6.64 (m, 3H), 2.83 (t, *J*=7.8 Hz, 2H), 2.32 (t, *J*=7.8 Hz, 2H); LCMS [M+H]<sup>+</sup> = 336.2; HPLC (water/ACN + 0.1% TFA gradient) 99.6% at 220nm.

*3-(2-Fluoro-5-hydroxy-1,1':4',1''-terphenyl-2'-yl)propanamide (T66)*

White powder (75 mg, 45%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 9.51 (s, 1H), 7.70 (d, *J*=7.2 Hz, 2H), 7.63 (d, *J*=1.4 Hz, 1H), 7.54 (dd, *J*=7.9, 1.7 Hz, 1H), 7.49 (t, *J*=7.5 Hz, 2H),  
 5 7.35 - 7.42 (m, 1H), 7.19 - 7.29 (m, 2H), 7.10 (t, *J*=9.1 Hz, 1H), 6.79 (dt, *J*=8.6, 3.6 Hz, 1H), 6.74 (br. s., 1H), 6.66 (dd, *J*=6.3, 2.9 Hz, 1H), 2.68 - 2.78 (m, 2H), 2.24 - 2.34 (m, 2H); LCMS [M+H]<sup>+</sup> = 336.2; HPLC (water/ACN + 0.1% TFA gradient) 97.6% at 220nm.

10 Example 13 - Synthesis of 3-[4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)biphenyl-3-yl]propanamide

3-[4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)biphenyl-3-yl]propanamide (**8**) was prepared from 3-(3-Amino-3-oxopropyl) biphenyl-4-yl trifluoromethanesulfonate (**7** - produced in Example 10). A mixture of **7** (1.81 g, 4.84 mmol), bis(pinacolato)diboron (1.35 g, 5.31 mmol), 1,1'-bis(diphenylphosphino)ferrocenepalladium(II) dichloride  
 15 dichloromethane complex (790 mg, 0.97 mmol) and potassium acetate (1.43 g, 14.5 mmol) were suspended in anhydrous dimethylsulfoxide (31 mL) under nitrogen. The mixture was heated at 85°C under nitrogen for 4 h. The mixture was diluted with ethyl acetate (90 mL) and eluted through a silica gel column with ethyl acetate. The fractions that contained the major band were washed with water (2 × 200 mL) and brine (200 mL),  
 20 dried over anhydrous sodium sulfate, and filtered. The filtrate was evaporated to dryness. The residue was purified by flash chromatography (ethyl acetate/dichloromethane) to give compound **8**:




- 51 -

A light brown oil that solidified on standing (1.02 g, 59%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>) 7.91 (d, *J*=8.4 Hz, 1H), 7.60 (d, *J*=7.2 Hz, 2H), 7.40 - 7.50 (m, 4H), 7.32 - 7.39 (m, 1H), 5.82 (br. s., 1H), 5.33 (br. s., 1H), 3.22 - 3.31 (m, 2H), 2.51 - 2.59 (m, 2H), 1.38 (s, 12H).

Example 14 - Synthesis of T24, T26, T27, T30, T32, T33, T35, T37, T39 and T58


- 5 T24, T26, T27, T30, T32, T33, T35, T37, T39 and T58 were prepared from 3-[4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)biphenyl-3-yl]propanamide (**8** - produced in Example 13). A mixture of **8** (1equiv.), a bromo-aromatic or bromo-heterocycle (1.1 equiv.) and 1,1'-bis(diphenylphosphino)-ferrocenepalladium(II) dichloride dichloromethane complex (0.1 equiv.) was dissolved in anhydrous *N,N*-
- 10 dimethylformamide (10.6 mL/mmol) under nitrogen. A degassed sodium carbonate solution (2 M, 5.3 mL/mmol) was added. The mixture was heated at 80°C under nitrogen. The mixture was partitioned between ethyl acetate and water. The aqueous phase was extracted with ethyl acetate. The combined ethyl acetate extracts were washed with water (3×) and brine, dried over anhydrous sodium sulfate, and filtered. The filtrate was
- 15 evaporated to dryness and purified by flash chromatography (methanol/dichloromethane). The product was suspended in 1:1 dichloromethane/hexanes and isolated by filtration
- The following compounds were prepared by this procedure:

*3-[4-(2-Oxo-2,3-dihydro-1H-indol-4-yl)biphenyl-3-yl]propanamide (T24)*




- 20 Pale yellow powder (72 mg, 58%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 10.46 (s, 1H), 7.70 (d, *J*=7.2 Hz, 2H), 7.63 (d, *J*=1.4 Hz, 1H), 7.43 - 7.56 (m, 3H), 7.34 - 7.42 (m, 1H), 7.16 - 7.30 (m, 3H), 6.83 (t, *J*=7.3 Hz, 2H), 6.67 (br. s., 1H), 3.26 (br. s., 2H), 2.72 (br. t, *J*=7.0 Hz, 2H), 2.25 (t, *J*=7.7 Hz, 2H); LCMS [M+H]<sup>+</sup> = 357.2; HPLC (water/ACN + 0.1% TFA gradient) 100.0% at 220nm.

## 3-[4-(2-Oxo-2,3-dihydro-1,3-benzoxazol-7-yl)biphenyl-3-yl]propanamide (T26)




Light orange powder (30 mg, 23%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 11.72 (br. s., 1H), 7.70 (d, *J*=7.4 Hz, 2H), 7.65 (s, 1H), 7.56 (dd, *J*=7.8, 1.2 Hz, 1H), 7.48 (t, *J*=7.6 Hz, 2H),  
 5 7.34 - 7.42 (m, 1H), 7.31 (d, *J*=7.8 Hz, 1H), 7.15 - 7.26 (m, 2H), 7.10 (d, *J*=7.0 Hz, 1H), 7.02 (d, *J*=7.4 Hz, 1H), 6.69 (br. s., 1H), 2.75 (t, *J*=8.0 Hz, 2H), 2.26 (t, *J*=8.0 Hz, 2H);  
 LCMS [M+H]<sup>+</sup> = 359.1; HPLC (water/ACN + 0.1% TFA gradient) 98.9% at 220nm.

## 3-[4-(2-Oxo-2,3-dihydro-1,3-benzoxazol-5-yl)biphenyl-3-yl]propanamide (T27)



10 Pale orange powder (49 mg, 28%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 11.69 (br. s., 1H), 7.70 (d, *J*=7.4 Hz, 2H), 7.62 (d, *J*=1.6 Hz, 1H), 7.45 - 7.57 (m, 3H), 7.31 - 7.43 (m, 2H), 7.17 - 7.30 (m, 2H), 7.00 - 7.09 (m, 2H), 6.72 (br. s., 1H), 2.83 (t, *J*=7.8 Hz, 2H), 2.25 - 2.35 (m, 2H); LCMS [M+H]<sup>+</sup> = 359.2; HPLC (water/ACN + 0.1% TFA gradient) 97.0%  
 at 220nm.

## 15 3-[4-(2-Oxo-2,3-dihydro-1H-benzimidazol-4-yl)biphenyl-3-yl]propanamide (T30)




- 53 -

Light beige powder (110 mg, 63%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 10.69 (s, 1H), 10.48 (s, 1H), 7.71 (d, *J*=7.4 Hz, 2H), 7.63 (d, *J*=1.6 Hz, 1H), 7.55 (dd, *J*=7.8, 1.8 Hz, 1H), 7.50 (t, *J*=7.6 Hz, 2H), 7.35 - 7.43 (m, 1H), 7.19 - 7.30 (m, 2H), 6.97 - 7.05 (m, 1H), 6.91 - 6.97 (m, 1H), 6.78 - 6.82 (m, 1H), 6.75 (br. s., 1H), 2.60 - 2.86 (m, 2H), 2.29 (br. s., 2H);

5 LCMS [M+H]<sup>+</sup> = 358.2; HPLC (water/ACN + 0.1% TFA gradient) 96.4% at 220nm.


*3-[4-(2-Oxo-1,2,3,4-tetrahydroquinolin-7-yl)biphenyl-3-yl]propanamide (T32)*



Pale beige powder (118 mg, 66% yield). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 10.13 (s, 1H), 7.69 (d, *J*=7.4 Hz, 2H), 7.61 (d, *J*=1.6 Hz, 1H), 7.44 - 7.56 (m, 3H), 7.34 - 7.41 (m, 1H),

10 7.18 - 7.28 (m, 3H), 6.90 (dd, *J*=7.5, 1.5 Hz, 1H), 6.82 (d, *J*=1.2 Hz, 1H), 6.74 (br. s., 1H), 2.94 (t, *J*=7.5 Hz, 2H), 2.77 - 2.87 (m, 2H), 2.53 (obscured by DMSO-d<sub>6</sub>), 2.28 - 2.37 (m, 2H); LCMS [M+H]<sup>+</sup> = 371.2; HPLC (water/ACN + 0.1% TFA gradient) 97.9% at 220nm.

*3-[4-[2-(Trifluoromethyl)-1H-benzimidazol-4-yl]biphenyl-3-yl]propanamide (T33)*



15


White powder (107 mg, 53%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 13.81 - 14.10 (m, 1H), 7.08 - 7.88 (m, 12H), 6.59 - 6.81 (m, 1H), 2.71 (br. t, *J*=7.3 Hz, 2H), 2.28 (m, 2H),

Spectrum was split into two species due to hydrogen exchange on the benzimidazole

- 54 -


moiety; LCMS  $[M+H]^+ = 410.2$ ; HPLC (water/ACN + 0.1% TFA gradient) 100% at 220nm.

3-[4-(1H-Benzimidazol-4-yl)biphenyl-3-yl]propanamide (T35)




- 5 Light brown powder (54 mg, 32%).  $^1\text{H NMR}$  (400MHz, DMSO- $d_6$ ) 12.32 - 12.57 (m, 1H), 8.16 (d,  $J=13.3$  Hz, 1H), 7.03 - 7.79 (m, 12H), 6.54 - 6.75 (m, 1H), 2.67 - 2.81 (m, 2H), 2.20 - 2.33 (m, 2H), Spectrum was split into two species due to hydrogen exchange on the benzimidazole moiety; LCMS  $[M+H]^+ = 342.2$ ; HPLC (water/ACN + 0.1% TFA gradient) 96.9% at 220nm.


10 3-[4-(1H-Indazol-4-yl)biphenyl-3-yl]propanamide (T37)



- Pale brown powder (69 mg, 42%).  $^1\text{H NMR}$  (400MHz, DMSO- $d_6$ ) 13.20 (s, 1H), 7.68 - 7.79 (m, 4H), 7.54 - 7.63 (m, 2H), 7.50 (t,  $J=7.6$  Hz, 2H), 7.32 - 7.47 (m, 3H), 7.17 (br. s., 1H), 7.03 (d,  $J=6.8$  Hz, 1H), 6.67 (br. s., 1H), 2.77 (t,  $J=7.8$  Hz, 2H), 2.21 - 2.29 (m, 2H);
- 15 LCMS  $[M+H]^+ = 342.2$ ; HPLC (water/ACN + 0.1% TFA gradient) 99.7% at 220nm.

*3-[4-(1H-Benzotriazol-4-yl)biphenyl-3-yl]propanamide (T39)*

Pale yellow powder (10 mg, 6%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 15.79 (br. s., 1H), 7.81 - 8.05 (m, 1H), 7.68 - 7.79 (m, 3H), 7.62 (d, *J*=7.8 Hz, 1H), 7.52 (t, *J*=7.7 Hz, 3H), 7.29 - 7.45 (m, 3H), 7.21 (br. s., 1H), 6.71 (br. s., 1H), 2.68 - 2.80 (m, 2H), 2.27 (t, *J*=7.8 Hz, 2H); LCMS [M+H]<sup>+</sup> = 343.2; HPLC (water/ACN + 0.1% TFA gradient) 98.2% at 220nm.

*3-(2,4-Difluoro-3-hydroxy-1,1':4',1''-terphenyl-2'-yl)propanamide (T58)*

After the first ethyl acetate/water extraction, the aqueous layer was adjusted to pH 6 by the addition of 1M hydrochloric acid and workup continued per the general procedure. Obtained as a pale brown powder (106 mg, 62%). <sup>1</sup>H NMR (400MHz, DMSO-d<sub>6</sub>) 10.24 (br. s., 1H), 7.70 (d, *J*=7.4 Hz, 2H), 7.64 (d, *J*=1.4 Hz, 1H), 7.55 (dd, *J*=7.9, 1.7 Hz, 1H), 7.49 (t, *J*=7.6 Hz, 2H), 7.35 - 7.43 (m, 1H), 7.17 - 7.27 (m, 2H), 7.11 (t, *J*=9.2 Hz, 1H), 6.65 - 6.81 (m, 2H), 2.71 (t, *J*=7.7 Hz, 2H), 2.28 (t, *J*=7.9 Hz, 2H); LCMS [M+H]<sup>+</sup> = 354.3; HPLC (water/ACN + 0.1% TFA gradient) 99.5% at 220nm.

Example 15 - Synthesis of 5-Iodo-2-methylbenzene-1,3-diol for use in the production of T67

A mixture of 4-chloro-3,5-dimethoxyaniline (3.0 g, 16.0 mmol), palladium(II)acetate (180 mg, 0.80 mmol), 2-dicyclohexylphosphino-2,4,6-triisopropylbiphenyl (XPhos)

- 56 -

(381 mg, 0.80 mmol), potassium carbonate (6.73 g, 48.7 mmol) and methylboronic acid (1.15 g, 19.2 mmol) in water (100 mL) and dioxane (100 mL) was heated to 100°C (oil bath temperature) under nitrogen for 18 h. The reaction was not complete and heated to reflux for an additional 3 h, cooled to room temperature, diluted with water (200 mL) and  
5 extracted with ethyl acetate (3 x 150 mL), dried over magnesium sulfate, concentrated and purified by flash chromatography (ethyl acetate/hexanes) to give 3,5-dimethoxy-4-methylaniline (720 mg, 27%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 5.93 (s, 2H), 3.77 (s, 6H), 3.58 (bs, 2H), 1.98 (s, 3H). Sodium nitrite (340 mg, 4.93 mmol) was added to a mixture of 3,5-dimethoxy-4-methylaniline (720 mg, 4.31 mmol) in sulfuric acid (1.1 mL) and  
10 water (13 mL) at 0°C and stirred for 30 min. The ensuing mixture was added to a preheated mixture of sodium iodide (2.58 g, 17.2 mmol) and iodine (555 mg, 2.19 mmol) in sulfuric acid (1.1 mL) and water (13 mL) at 80°C, and the mixture heated to reflux for 30 min. The reaction mixture was cooled to room temperature and diluted with a solution of sodium sulfite (20% w/w, 100 mL) and water (100 mL), and extracted with ethyl  
15 acetate (3 x 100 mL). The organics were combined, dried over magnesium sulfate, concentrated and purified by flash chromatography (ethyl acetate/hexanes) to give 5-iodo-1,3-dimethoxy-2-methylbenzene as a white powder (388 mg, 32%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 6.84 (s, 2H), 3.79 (s, 6H), 2.02 (s, 3H). 5-Iodo-1,3-dimethoxy-2-methylbenzene (388 mg, 1.39 mmol) in dichloromethane (8 mL) was cooled to 0°C before adding neat  
20 boron tribromide (0.8 mL, 8 mmol) slowly over 1 min under nitrogen. The reaction mixture was allowed to warm to room temperature slowly over 3 h and stirred for 18 h. The reaction mixture was slowly and cautiously poured onto ice water (100 mL) and extracted with ethyl acetate (3 x 60 mL) and the combined organics washed with brine (1 x 50 mL), dried over magnesium sulfate, concentrated and purified by flash  
25 chromatography (ethyl acetate/hexanes) to give 5-iodo-2-methylbenzene-1,3-diol as a white powder (260 mg, 74%). <sup>1</sup>H NMR (400 MHz, d<sub>6</sub>-DMSO) 9.44 (s, 2H), 6.63 (s, 2H), 1.87 (s, 3H).

Example 16 - Synthesis of 5-Bromo-2-fluorobenzene-1,3-diol for use in the production of T68

30 A mixture of Oxone (1.44 g) in water (2 mL) was added to a solution of 5-bromo-2-fluoro-1,3-phenylenediboronic acid, pinacol ester (500 mg, 1.17 mmol) in acetone (1.5 mL) over 1 min and stirred at room temperature for 15 min. Additional Oxone (0.512 g)

and acetone (1 mL) was added and stirred for a further 20 min. Sodium bisulfite solution (10 % w/w, 10 mL) was added followed by water (10 mL) and extracted with dichloromethane (3 x 20 mL), dried over magnesium sulphate and purified by flash chromatography (ethyl acetate/hexanes) to give 5-bromo-2-fluorobenzene-1,3-diol as a white powder (159 mg, 66%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 6.73 (d, *J* = 6.8 Hz, 2H), 5.29 (bs, 2H).

Example 17 - Synthesis of 2-Chloro-5-iodobenzene-1,3-diol for use in the production of T69


2-Chloro-5-iodobenzene-1,3-diol was synthesised in 3 steps from 3,5-dimethoxyaniline according to WO 2011/027106 A1, with a modification to the first step to prepare 4-chloro-3,5-dimethoxyaniline in which the reagent *N*-chlorosuccinimide was added portion wise as a suspension in acetic acid (50 mL) to a solution of 3,5-dimethoxyaniline (10.01 g, 65.35 mmol) in acetic acid (50 mL) at 0°C and warmed to room temperature after 30 min.

Example 18 -Synthesis of T67, T68 and T69

T67, T68 and T69 were prepared from 3-[4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)biphenyl-3-yl]propanamide (**8** - produced in Example 13). A mixture of **8** (1 equiv.), a substituted benzene-1,3-diol (produced in examples 15-17) (1.1 equiv.), Pd(dppf)Cl<sub>2</sub>.CH<sub>2</sub>Cl<sub>2</sub> (0.1 equiv.) and sodium carbonate (2M, 5 mL/mmol) in *N,N*-dimethylformamide (10 mL/mmol) was bubbled through with nitrogen for 5 min before being heated to 80°C under nitrogen for 18 h. The reaction mixture was cooled to room temperature and partitioned between 1M hydrochloric acid and ethyl acetate. The combined organics were washed with brine, dried over magnesium sulfate, concentrated and purified by flash chromatography (ethyl acetate/hexanes).


The following compounds were prepared by this procedure:

3-(3,5-Dihydroxy-4-methyl-1,1':4',1''-terphenyl-2'-yl)propanamide (T67)




White powder (121 mg, 59%). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) 9.17 (s, 2H), 7.68 (d, *J*=7.43 Hz, 2H), 7.58 (br. s, 1H), 7.48 (t, *J*=7.80 Hz, 3H), 7.37 (s, 1H), 7.23 (br. s., 1H),  
 5 7.17 (d, *J*=7.83 Hz, 1H), 6.76 (br. s, 1H), 6.25 (s, 2H), 3.33 (s, 3H), 2.85 (t, *J*=7.83 Hz, 2H), 2.30 (t, *J*=7.83 Hz, 2H). LCMS [M+H]<sup>+</sup> = 348. HPLC (water/ACN + 0.1% TFA gradient) 100% at 220 nm.

3-(4-Fluoro-3,5-dihydroxy-1,1':4',1''-terphenyl-2'-yl)propanamide (T68)



10 Pale yellow solid (82 mg, 41%). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 9.70 (br. s, 2H), 7.68 (d, *J*=7.43 Hz, 2H), 7.58 (s, 1H), 7.43 - 7.53 (m, 3H), 7.32 - 7.42 (m, 1H), 7.25 (br. s, 1H), 7.18 (d, *J*=7.83 Hz, 1H), 6.75 (br. s, 1H), 6.33 (d, *J*=7.43 Hz, 2H), 2.83 (t, *J*=7.80 Hz, 2H), 2.30 (t, *J*=8.20 Hz, 2H). LCMS [M+H]<sup>+</sup> = 352. HPLC (water/ACN + 0.1% TFA gradient) 96.1% at 220 nm.

15 3-(4-Chloro-3,5-dihydroxy-1,1':4',1''-terphenyl-2'-yl)propanamide (T69)



White powder (26 mg, 12%). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) 9.93 - 10.12 (m, 2H), 7.69 (d, *J*=7.43 Hz, 2H), 7.59 (s, 1H), 7.43 - 7.54 (m, 3H), 7.34 - 7.42 (m, 1H), 7.25 (br. s., 1H), 7.19 (d, *J*=7.83 Hz, 1H), 6.76 (br. s., 1H), 6.38 (s, 2H), 2.84 (t, *J*=7.40 Hz, 2H), 2.30 (t, *J*=8.20 Hz, 2H). LCMS [M+H]<sup>+</sup> = 368. HPLC (water/ACN + 0.1% TFA gradient)  
5 100% at 254 nm.

#### Example 19 - Synthesis of T25

The synthetic route used to prepare T25 is shown in Figure 8. Briefly, 3-formylbiphenyl-4-yl trifluoromethanesulfonate (14) was subjected to a Horner-Wadsworth-Emmons reaction with diethyl(carbamoylmethyl)phosphonate (18) to afford biphenyl acrylamide (23), which cross-coupled with indolone pinacol boronic ester (24) to produce indolone acrylamide (25). Subsequent hydrogenation of compound 25 yielded T25.  
10

To synthesise T25, the requisite indolone pinacol boronic ester (24) was needed. Thus, 1,4-dibromo-2-nitrobenzene was alkylated with dimethyl malonate to give aryl malonate (26), which was decarboxylated and cyclised to form bromoindolone (27); this was in turn reacted with bis(pinacolato)diborane to form indolone pinacol boronic ester (24) (Figure 9).  
15

#### *Production of Dimethyl 2-(4-bromo-2-nitrophenyl)malonate (26)*

To a mixture of potassium *tert*-butoxide (21.6 g, 193.00 mmol) in DMF (75 mL) was added dimethyl malonate (22.40 mL, 196.00 mmol). The reaction was exothermic and a solid precipitated out. The reaction mixture was heated to 90 °C for 10 minutes, then cooled to ambient temperature. 2,5-Dibromonitrobenzene (25.50 g, 91 mmol) was added as a solid. The reaction mixture turned purple and was stirred at 90 °C for 2 hours. After cooling to ambient temperature, it was poured onto ice cold 5% hydrochloric acid solution and transferred to a separating funnel. The crude material was extracted out with ethyl acetate (2x). The combined ethyl acetate extracts were washed with water and brine to give a bright yellow oil. The crude oil was pre-absorbed onto Celite, then chromatographed (DCVC) eluting with a gradient of ethyl acetate in heptane (0 -10% ethyl acetate). Like fractions were combined and were recrystallised from DCM and heptane to give dimethyl 2-(4-bromo-2-nitrophenyl)malonate (26) as pale yellow needles  
25  
30 (26.78 g, 87%); mp 85.8 – 87.1°C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.18 (d, 1H, *J* 2.1 Hz), 7.75 (dd, 1H, *J* 2.1, 8.4 Hz), 7.40 (d, 1H, *J* 8.4 Hz), 5.26 (s, 1H), 3.78 (s, 6H).

- 60 -

*Production of 6-Bromoindolin-2-one (27)*

Lithium chloride (6.36 g, 156.0 mmol) was added to a solution of dimethyl 2-(4-bromo-2-nitrophenyl)malonate (26) (26.0 g, 78.30 mmol) in dimethylsulfoxide (100 mL) and was heated at 100 °C for 20 hours. Once cooled to ambient temperature the reaction mixture  
5 was partitioned between ethyl acetate and brine. The layers were separated, then washed with brine again and concentrated. The dark tan oil was dissolved in acetic acid (100 mL) and iron powder (17.50 g, 313.0 mmol) was added (exotherm). The reaction was then heated at 110 °C for 1 hour. The acetic acid was removed by rotary evaporation and the residue dissolved in ethyl acetate and the iron powder was removed by filtering through  
10 Celite. The filtrate was washed with 1M hydrochloric acid and water, then filtered through phase separation paper (IPS). The crude material was pre-absorbed onto Celite, then chromatographed (DCVC) eluting with chloroform. Fractions containing the desired material were combined, pre-absorbed onto Celite and chromatographed (DCVC) again eluting with a gradient of ethyl acetate in heptane (20 -80% ethyl acetate). Clean  
15 fractions were combined and recrystallised from DCM and methanol to give 6-bromoindolin-2-one (27) as yellow needles (4.32 g, 26%); mp 208 - 214°C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 10.47 (br s, 1H), 7.14 (d, 1H, *J* 7.9 Hz), 7.09 (dd, 1H, *J* 1.8, 7.9 Hz), 6.94 (d, 1H, *J* 1.8 Hz), 3.44 (s, 2H).

*Production of 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)indolin-2-one (24)*

20 6-Bromoindolin-2-one (27) (2.00 g, 9.40 mmol), bispinacolatodiboron (6.00 g, 23.60 mmol), potassium acetate (2.76 g, 28.2 mmol) and dichloro[1,1'-bis(diphenylphosphino)ferrocene] palladium (II) dichloromethane adduct (0.40 g, 0.55 mmol) in DMSO (30 mL) were stirred at 90 °C for 18 hours. The reaction mixture was cooled to ambient temperature, then partitioned between water and ethyl acetate. The  
25 layers were separated and the aqueous layer extracted again with ethyl acetate (2x). The combined organic layers were washed with water and brine and concentrated to give a purple solid. The crude material was pre-absorbed onto Celite and chromatographed (DCVC) eluting with a gradient of ethyl acetate in heptane (0 - 50% ethyl acetate). Like fractions were combined and recrystallised from DCM and PE to give 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)indolin-2-one (24) as a colourless solid in 2 crops  
30 (1.33 g, 55%); mp 178.5 - 181.4 °C. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 8.61 (br s, 1H), 7.46 (d, 1H, *J* 7.4 Hz), 7.30 (s, 1H), 7.21 (d, 1H, *J* 7.4 Hz), 3.53 (s, 2H), 1.32 (s, 12H).

*Production of (E)-3-(3-Amino-3-oxoprop-1-en-1-yl)-[1,1'-biphenyl]-4-yl trifluoromethanesulfonate (23)*

3-Formylbiphenyl-4-yltrifluoromethanesulfonate (14) (3.80 g, 11.50 mmol) and diethyl (2-amino-2-oxoethyl)phosphonate (18) (2.25 g, 11.50 mmol) were dissolved in dry THF (100 mL), and added slowly to a vigorously stirred suspension of powdered sodium hydroxide (0.92 g, 23.00 mmol). Following stirring for 1 h at rt, the reaction mixture was partitioned between brine and ethyl acetate. A yellow by-product was removed by filtration and the layers separated. The organic layer was concentrated and then purified by chromatography (DCVC) eluting with a gradient of ethyl acetate in heptane (0 – 20% ethyl acetate), and then recrystallised from DCM and PE to give (E)-3-(3-amino-3-oxoprop-1-en-1-yl)-[1,1'-biphenyl]-4-yl trifluoromethanesulfonate (23) as a beige solid (0.82 g, 19%); mp 130.6 – 132.3 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 8.09 – 8.04 (m, 1H), 7.88 – 7.82 (m, 1H), 7.79 – 7.73 (m, 2H), 7.65 – 7.41 (m, 6H), 7.33 (br s, 1H), 6.93 (d, 1H, <sup>3</sup>J<sub>trans</sub> 16 Hz). <sup>13</sup>C NMR (100 MHz, DMSO-*d*<sub>6</sub>) δ 165.6, 146.4, 141.1, 138.0, 130.0, 129.5, 129.1, 128.6, 128.4, 127.7, 127.1, 126.4, 122.8, 118.1 (q, *J* 321 Hz). EIMS: *m/z* Found: M<sup>+</sup> 371.0420, C<sub>16</sub>H<sub>12</sub>F<sub>3</sub>NO<sub>4</sub><sup>32</sup>S requires 371.0434. EIMS: *m/z* 371 (M<sup>+</sup>, 62%), 195 (100), 167 (100).

*Production of (E)-3-(4-(2-Oxindolin-6-yl)-[1,1'-biphenyl]-3-yl)acrylamide (25)*

Prepared according to the method used to generate P5; from (E)-3-(3-amino-3-oxoprop-1-en-1-yl)-[1,1'-biphenyl]-4-yl trifluoromethanesulfonate (23) (0.50 g, 1.35 mmol), 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)indolin-2-one (24) (0.43 g, 1.68 mmol), tetrakis(triphenylphosphine)palladium(0) (0.100 g, 0.09 mmol) and aqueous sodium carbonate (1M) (3.0 mL, 3.00 mmol) in toluene (10 mL) and ethanol (2 mL). The crude material was collected by filtration from the aqueous work-up, then purified by trituration in DCM and methanol to give (E)-3-(4-(2-oxindolin-6-yl)-[1,1'-biphenyl]-3-yl)acrylamide (25) as pale lemon needles (0.36 g, 75%); mp 263 - 267°C (Dec.). <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 10.47 (s, 1H), 7.95 (s, 1H), 7.80 – 7.70 (m, 3H), 7.57 – 7.37 (m, 5H), 7.46 (br s, 1H), 7.32 (d, 1H, *J* 7.6 Hz), 7.12 (br s, 1H), 6.89 (d, 1H, *J* 7.6 Hz), 6.80 – 6.72 (m, 2H), 3.55 (s, 2H). <sup>13</sup>C NMR (100 MHz, DMSO-*d*<sub>6</sub>) δ 177.4, 166.5, 143.9, 141.1, 139.6, 139.4, 138.8, 137.4, 133.1, 130.9, 129.0, 127.8, 127.6, 126.8, 125.3, 124.4, 124.3, 123.7, 122.6, 110.1, 35.6. EIMS: *m/z* Found: M<sup>+</sup> 354.1356, C<sub>23</sub>H<sub>18</sub>N<sub>2</sub>O<sub>2</sub> requires 354.1363. EIMS: *m/z* 354 (M<sup>+</sup>, 13%), 310 (100), 309 (43).

*Production of 3-(4-(2-Oxoindolin-6-yl)-[1,1'-biphenyl]-3-yl)propanamide (T25).*

Prepared according to the method used to generate T18; from (*E*)-3-(4-(2-oxindolin-6-yl)-[1,1'-biphenyl]-3-yl)acrylamide (25) (0.11 g, 0.30 mmol) and 10% palladium on carbon (50% wt water) in methanol (30mL). The filtrate was concentrated to give 3-(4-(2-oxoindolin-6-yl)-[1,1'-biphenyl]-3-yl)propanamide (T25) as a pale yellow solid (0.96 g, 89%); mp 219 - 222°C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 10.44 (s, 1H), 7.74 – 7.65 (m, 2H), 7.65 – 7.58 (m, 1H), 7.56 – 7.43 (m, 3H), 7.42 – 7.34 (m, 1H), 7.31 – 7.19 (m, 3H), 6.94 – 6.87 (m, 1H), 6.80 – 6.71 (m, 2H), 3.53 (s, 2H), 2.84 (t, 2H, *J* 7.9 Hz), 2.31 (t, 2H, *J* 7.9 Hz). <sup>13</sup>C NMR (50 MHz, DMSO-*d*<sub>6</sub>) δ 176.5, 173.4, 143.8, 140.6, 140.2, 140.0, 139.2, 139.1, 130.3, 128.9, 127.4, 127.3, 126.7, 124.6, 124.2 (two signals coincident), 121.9, 109.7, 36.2, 35.6, 28.2. EIMS: *m/z* Found: M<sup>+</sup> 356.1531, C<sub>23</sub>H<sub>20</sub>N<sub>2</sub>O<sub>2</sub>N requires 356.1531. EIMS: *m/z* 356 (M<sup>+</sup>, 100%), 297 (70). HPLC purity (35% ACN / 0.1% TFA, 256 nm): 97.57%.

Example 20 - Synthesis of T31

The synthetic route used to prepare T31 is shown in Figure 10. Briefly, 3-formylbiphenyl-4-yl trifluoromethanesulfonate (14) was cross-coupled with benzimidazolone pinacol boronic ester (24) to produce benzimidazolone (28), which then underwent a Horner-Wadsworth-Emmons reaction with diethyl(carbamoylmethyl)phosphonate (18) to afford benzimidazolone acrylamide (29). Subsequent hydrogenation of compound 29 yielded T31.

*Production of 4-(2-Oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-[1,1'-biphenyl]-3-carbaldehyde (28)*

To a suspension of 2-oxo-2,3-dihydro-1H-benzoimidazole-5-boronic acid pinacol ester (24) (574 mg, 2.2 mmol), 3-formylbiphenyl-4-yl trifluoromethanesulfonate (14) (663 mg, 2.0 mmol) and sodium carbonate (426 mg, 4.0 mmol) in degassed dioxane/ethanol/H<sub>2</sub>O (5:1:1, 20 mL) was added tetrakis(triphenylphosphine)palladium(0) (116 mg, 0.1 mmol). The reaction was heated at 110 °C for 2 h in a sealed tube. Analysis by TLC (1:2 DCM;PE) indicated the triflate had been consumed. The reaction was concentrated to dryness, then taken up in equal volumes of DCM and water and stirred vigorously for 20 minutes, ensuring all lumps were broken up and a fine precipitate was achieved. The solid was collected by filtration through hardened ashless paper (540) on a Buchner funnel, and washed thoroughly with DCM and water. The solid was dried *in vacuo* at

- 63 -

40 °C to afford 4-(2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-[1,1'-biphenyl]-3-carbaldehyde (28) (365 mg, 58%) as a pale yellow solid. <sup>1</sup>H NMR (200 MHz, DMSO-*d*<sub>6</sub>) δ 10.82 (brs, 2H), 9.95 (s, 1H), 8.15 – 7.96 (m, 2H), 7.76 (m, 2H), 7.68 – 7.36 (m, 4H), 7.12 – 6.95 (m, 3H). <sup>13</sup>C NMR (50 MHz, DMSO-*d*<sub>6</sub>) δ 192.1, 155.4, 144.6, 139.1, 138.7, 133.7, 131.8, 131.7, 130.1, 130.0, 129.2, 128.0, 126.7, 125.1, 122.9, 109.7, 108.4 (one signal not observed). EIMS: *m/z* Found: M<sup>+</sup> 314.1050, C<sub>20</sub>H<sub>14</sub>O<sub>2</sub>N<sub>2</sub> requires 314.1055. EIMS: *m/z* 314 (M<sup>+</sup>, 100%).

*Production of (E/Z)-3-(4-(2-Oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-[1,1'-biphenyl]-3-yl)acrylamide (29)*

4-(2-Oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-[1,1'-biphenyl]-3-carbaldehyde (28) (350 mg, 1.1 mmol) and diethyl (carbamoylemethyl)phosphonate (18) (217 mg, 1.1 mmol) were dissolved in dry THF (15 mL), and added slowly to a vigorously stirred suspension of powdered KOH (125 mg, 2.2 mmol) in THF (10 mL). The reaction was stirred at rt for 1 h under an argon atmosphere. Analysis by TLC (1:99 methanol:DCM) indicated the carbaldehyde had been consumed. The THF was removed under reduced pressure, and the residue taken up in equal volumes of DCM and water and stirred vigorously for 30 minutes; ensuring all lumps were broken up and a fine precipitate achieved. The solid was collected by filtration through hardened ashless paper (540) on a Buchner funnel, and washed thoroughly with DCM and water. The solid was dried *in vacuo* at 40 °C to afford (E/Z)-3-(4-(2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-[1,1'-biphenyl]-3-yl)acrylamide (29) (247 mg, 56%) as a yellow/brown solid. <sup>1</sup>H NMR (200 MHz, DMSO-*d*<sub>6</sub>) δ 10.74 (brs, 1H), 10.70 (brs, 1H), 7.93 (d, 1H, *J* 1.8), 7.80 – 7.67 (m, 3H), 7.58 – 7.34 (m, 6H), 7.14 – 6.98 (m, 2H), 6.94 – 6.83 (m, 2H), 6.74 (d, 1H, *J* 15.8 Hz). <sup>13</sup>C NMR (100 MHz, DMSO-*d*<sub>6</sub>) δ 166.6, 155.4, 141.5, 139.5, 139.1, 137.8, 133.1, 131.7, 131.2, 129.8, 129.3, 129.0, 127.7, 127.5, 126.7, 124.4, 123.4, 122.4, 109.5, 108.3. EIMS: *m/z* Found: M<sup>+</sup> 355.1315, C<sub>22</sub>H<sub>17</sub>O<sub>2</sub>F<sub>3</sub> requires 355.1315. EIMS: *m/z* 355 (M<sup>+</sup>, 31%).

*Production of 3-(4-(2-Oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-[1,1'-biphenyl]-3-yl)propanamide (T31)*

(E/Z)-3-(4-(2-Oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-[1,1'-biphenyl]-3-yl)acrylamide (29) (240 mg, 0.7 mmol) and 10% palladium on carbon (50% wt in H<sub>2</sub>O, 100 mg) in methanol (20 mL) were stirred at rt under a hydrogen atmosphere at 50 psi for 2h. The reaction mixture was gravity filtered through GF paper washing thoroughly with

- 64 -

methanol, then concentrated. Purified by preparative HPLC (55% methanol/H<sub>2</sub>O, 70 mL/min, 280 nm, 300 x 40 mm Deltaprep C<sub>18</sub> column) to afford 3-(4-(2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-[1,1'-biphenyl]-3-yl)propanamide (T31) (177 mg, 73%) as a pink solid; mp 250 – 251 °C. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 10.67 (d, 2H, *J* 7.8 Hz), 7.69 (m, 2H), 7.60 (d, 1H, *J* 1.9 Hz), 7.53 – 7.45 (m, 3H), 7.37 (m, 1H), 7.28 – 7.21 (m, 2H), 6.99 (d, 1H, *J* 7.8 Hz), 6.90 (dd, 1H, *J* 1.6, 7.9 Hz), 6.89 – 6.86 (m, 1H), 6.74 (brs, 1H), 2.84 (m, 2H), 2.30 (m, 2H). <sup>13</sup>C NMR (100 MHz, DMSO-*d*<sub>6</sub>) δ 173.4, 155.4, 140.9, 140.0, 139.3, 138.8, 133.1, 130.6, 129.7, 128.9, 128.8, 127.3, 127.2, 126.6, 124.1, 121.3, 109.0, 108.1, 36.1, 28.3. EIMS: *m/z* Found: M<sup>+</sup> 357.1469, C<sub>22</sub>H<sub>19</sub>N<sub>3</sub>O<sub>2</sub> requires 357.1472. EIMS: *m/z* 357 (M<sup>+</sup>, 30%). HPLC purity (40% ACN / H<sub>2</sub>O, 282 nm): 94.51%.

#### Example 21 - *in vitro* screening

The xCELLigence SP system (Roche) was used to measure changes in cellular impedance (cell index) following the treatment of A10 embryonic vascular smooth muscle cells (ATCC, CRL-1476) with test compound. This *in vitro* assay was correlated with blood pressure data obtained in the animal model described below in Example 22, so that it can be used for faster screening of larger number of compounds. In this *in vitro* cell based experimental system a negative impedance profile correlates with blood pressure reduction in rats - a decrease in impedance is associated with vasodilatation and an increase in impedance is associated with vasoconstriction (Stallaert W, Dorn JF, van der Westhuizen E, Audet M & Bouvier M. Impedance responses reveal β-adrenergic signaling pluridensitometry and allow classification of ligands with distinct signalling profiles PLoS ONE 2012; 7(1):e29420, doi:10.1371/journal.pone.0029420).

Briefly, 50 µl of cell culture medium (DMEM low glucose supplemented with 10% fetal bovine serum at 37°C) was added to each well of an E-Plate 96 (Roche), and the background impedance in each well was measured. 50 µl of A-10 cell suspension (10,000 cells/well) was then added to the appropriate wells of the E-Plate 96. Cell index was monitored for each well of the E-Plate 96 in RTCA SP Station within the cell culture incubator. After overnight incubation for 16-20 hours at 5% CO<sub>2</sub> and 95% humidity, 100 µl of test compound solution (test compounds were prepared in DMSO and diluted with cell culture medium to a final DMSO concentration of 0.25%) was added to the appropriate wells of the E-Plate 96 and cell index values were measured immediately following compound treatment every 20 seconds for 3 hours. Cell index value is baseline-

corrected by subtracting the cell index of vehicle-treated cells and normalized by dividing by the cell index at the time point immediately before compound addition. Baseline normalized cell index as a function of time is plotted using Roche RTCA software.

Compounds may achieve reductions in blood pressure by interaction with vascular smooth muscle cells causing these cells to relax resulting in vasodilatation and a reduction in blood pressure. These are termed direct vasodilators. A negative impedance response for A10 vascular smooth muscle cells indicates that a test compound is a direct vasodilator (Figure 11).

The xCELLigence SP system (Roche) was also used to measure changes in cellular impedance (cell index) following the treatment of bovine aortic endothelial cells (European Collection of Cell Cultures) with test compound. The method employed is the same for the A10 embryonic vascular smooth muscle cells described above but with the cell culture medium supplemented with 15% fetal bovine serum instead of 10%.

Compounds may interact with vascular endothelial cells causing the release of substances such as nitric oxide and endothelium-derived hyperpolarising factor, which in turn act on the vascular smooth muscle cells causing vasodilatation and lowering blood pressure. Such compounds are termed indirect vasodilators. A negative impedance response for bovine aortic endothelial cells indicates that a test compound is an indirect vasodilator (Figure 12).

#### 20 Example 22 – *in vivo* screening

##### *Oral Studies*

Fourteen week old SHR (on a 2.2% salt diet; Glen Forrest Stockfeeders) were randomly assigned to zero time control, test compound treatment (100 or 500 pmol/kg/min) in the drinking solution or control drinking solution (5% ethanol in deionised distilled water (n=5 each group). The rats assigned to zero time control group were anaesthetised and had their heart and kidneys harvested while rats assigned to control and test compound treatment were weighed twice weekly and had their drinking solution intake monitored to allow adjustment of the test compound concentration in the drinking solution to maintain a constant dose over the 4-week study period. Blood pressure was measured twice weekly by tail cuff plethysmography (PowerLab, ADInstruments, Castle Hill, NSW, Australia).

After 4 weeks rats were anaesthetised, and their hearts and kidneys harvested for quantitation of fibrosis.

#### *Fibrosis quantitation*

To quantitate tissue fibrosis tissue slices  $\leq 3$ mm thick were fixed in 10% buffered formalin for 24 hours, processed and embedded in paraffin. Three micron transverse sections were stained using Masson's trichrome stain. A minimum of 20 random fields at magnification x20 from transverse sections (5 at each of 2 levels) were digitized and the degree of fibrosis determined as a percent of field area of each digitized image using Image-Pro Plus V.7 (Media Cybernetics, Bethesda, MD, USA) then averaged to determine the level of fibrosis for tissue for each rat.

#### *Results*

Mean systolic blood pressure results observed for animals on 2.2% salt diet after 4 weeks treatment with 100 or 500pmol/kg/min T1, T2, T20, T31, T48 or T70 orally showed decreased blood pressure compared to controls (Figure 13). Mean diastolic blood pressure was also decreased compared to controls for T1, T2, T31 and T70.

The mean systolic blood pressure results for T1, T2, T20, T31, T48 and T70 were compared to the baseline normalised cell indexes of the compounds on A10 vascular smooth muscle cells (Figure 14) and the bovine aortic endothelial cells (Figure 15) and showed a correlation between the *in vivo* and *in vitro* results.

Fibrosis in the heart after 4 weeks treatment with 500 pmol/kg/min T1, T2, T20, T31, T48 or T70 orally in 18 week old SHR on 2.2% salt diet is decreased compared to controls (Figure 16).

Fibrosis in the kidney after 4 weeks treatment with 500 pmol/kg/min T1, T2, T20, T31, T48 or T70 orally in 18 week old SHR on 2.2% salt diet is decreased compared to controls (Figure 17).

The baseline normalised cell indexes of T1, T2, T20, T31, T48 and T70 on bovine aortic endothelial cells were compared with the myocardial fibrosis results (Figure 18) and kidney fibrosis results (Figure 19) for the compounds and showed a correlation between the *in vivo* and *in vitro* results.


- 67 -

Histological sections from the hearts (Figure 20) of control rats (A) or rats treated for four weeks with 500 pmol/kg/min T1 (B), T2 (C), T20 (D) or T31 (e) on 2.2% salt diet showed that the control has extensive fibrosis (see arrows) appearing as a band of light grey in the right lower quarter extending diagonally up and out surrounding the large  
5 blood vessel as well as numerous muscle fibres with smaller amounts present throughout the micrograph (muscle fibres appear as discrete darker grey areas). In the sections from T1, T2, T20 and T31 treated rats no discrete areas of fibrosis are present, muscle fibres appear in cross section as varying shades of dark grey.

Histological sections from the kidneys (Figure 21) of control rats (A) or rats treated for  
10 four weeks with 500 pmol/kg/min T1 (B), T2 (C), T20 (D) or T31 (e) on 2.2% salt diet showed that the control has extensive fibrosis appearing as thick lighter grey bands completely surrounding all tubules while in the centre 2 tubules have been obliterated (arrows). In the sections from T1, T2, T20 and T31 treated rats the fibrosis has been reduced to thin bands incompletely surrounding some but not all tubules.


THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:-

1. A compound of the formula



5 wherein:

A is selected from the group consisting of:



Q is independently selected from halo, alkyl, hydroxy, amino and substituted amino;

10 n is 0, 1, 2, 3, 4 or 5;


R<sub>1</sub>, R<sub>3</sub> and R<sub>4</sub> are independently C, CH, CH<sub>2</sub>, O, N, NH or S, and


R<sub>2</sub> is C, CH, CH<sub>2</sub>, N, NH, C-CF<sub>3</sub>, CH-CF<sub>3</sub> or C=O,

or a stereoisomer or pharmaceutically acceptable salt thereof,

wherein when n is 1, Q cannot be hydroxy.




2. The compound, or a stereoisomer or pharmaceutically acceptable salt thereof, according to claim 1 wherein Q is halo selected from the group consisting of F, Cl, Br and I.
3. The compound, or a stereoisomer or pharmaceutically acceptable salt thereof, according to claim 1, wherein Q is substituted amino of the formula -NHW and wherein:  
 W is selected from -CN, -SO<sub>2</sub>(X)<sub>a</sub>Y and -CO(X)<sub>a</sub>Y,  
 a is 0 or 1,  
 X is selected from -NH- and -O-, and  
 Y is selected from -H, -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>OH and -CH<sub>2</sub>CH<sub>2</sub>OH.
4. The compound, or a stereoisomer or pharmaceutically acceptable salt thereof, according to claim 1 or claim 3, wherein Q is substituted amino selected from the group consisting of -NHSO<sub>2</sub>CH<sub>3</sub>, -NHCOH, -NHCONHCH<sub>3</sub>, -NHCONHCH<sub>2</sub>CH<sub>3</sub>,  
 -NHSO<sub>2</sub>NHCH<sub>3</sub>, -NHSO<sub>2</sub>NHCH<sub>2</sub>CH<sub>3</sub>, -NHCOCH<sub>3</sub>, -NHCOOCH<sub>3</sub>, -NHCOOCH<sub>2</sub>CH<sub>2</sub>OH,  
 -NHCONH<sub>2</sub> and -NHCN.
5. The compound, or a stereoisomer or pharmaceutically acceptable salt thereof, according to claim 1 wherein Q is alkyl selected from the group consisting of methyl, ethyl, propyl, butyl and pentyl.
6. The compound, or a stereoisomer or pharmaceutically acceptable salt thereof, according to claim 1 wherein A is selected from:


















and


7. The compound, or a stereoisomer or pharmaceutically acceptable salt thereof,  
 5 according to claim 1, wherein the compound is selected from the group consisting of:









(T31),




(T32),




(T33),




(T35),




(T37),




(T38),




(T39),




(T48),




(T58),



(T63),








(T64),



(T65),

- 73 -



8. A pharmaceutical composition comprising a compound, or a stereoisomer or  
 5 pharmaceutically acceptable salt thereof, according to any one of claims 1 to 7 and a  
 pharmaceutically-acceptable excipient.

9. A method for the therapeutic treatment of hypertension or prehypertension in a subject  
 comprising administering to the subject a compound, or a stereoisomer or  
 10 pharmaceutically acceptable salt thereof, according to any one of claims 1 to 7 or a  
 composition according to claim 8.

10. A method for the prophylactic treatment of fibrosis in a subject comprising  
 administering to the subject a compound, or a stereoisomer or pharmaceutically  
 15 acceptable salt thereof, according to any one of claims 1 to 7 or a composition according  
 to claim 8.

11. A method for the therapeutic treatment of fibrosis in a subject comprising  
 administering to the subject a compound, or a stereoisomer or pharmaceutically

- 74 -

acceptable salt thereof, according to any one of claims 1 to 7 or a composition according to claim 8.

12. A method for the therapeutic treatment of hypertension and fibrosis in a subject  
5 comprising administering to the subject a compound, or a stereoisomer or pharmaceutically acceptable salt thereof, according to any one of claims 1 to 7 or a composition according to claim 8.

13. A method for the treatment of prehypertension and fibrosis in a subject comprising  
10 administering to the subject a compound, or a stereoisomer or pharmaceutically acceptable salt thereof, according to any one of claims 1 to 7 or a composition according to claim 8.

14. The method according to any one of claims 10 to 13 wherein the fibrosis is  
15 myocardial fibrosis or kidney fibrosis.

15. The method according to any one of claims 10 to 13 wherein the fibrosis is kidney fibrosis and kidney fibrosis.

20 16. Use of a compound, or a stereoisomer or pharmaceutically acceptable salt thereof, according to any one of claims 1 to 7 for the manufacture of a medicament for the therapeutic treatment of hypertension or prehypertension.

17. Use of a compound, or a stereoisomer or pharmaceutically acceptable salt thereof,  
25 according to any one of claims 1 to 7 for the manufacture of a medicament for the prophylactic treatment of fibrosis.

18. Use of a compound, or a stereoisomer or pharmaceutically acceptable salt thereof,  
30 according to any one of claims 1 to 7 for the manufacture of a medicament for the therapeutic treatment of fibrosis.

- 75 -

19. Use of a compound, or a stereoisomer or pharmaceutically acceptable salt thereof, according to any one of claims 1 to 7 for the manufacture of a medicament for the therapeutic treatment of hypertension and fibrosis or prehypertension and fibrosis.

1/16




Figure 1




Figure 2

2/16



Figure 3




Figure 4

3/16




Figure 5




Figure 6

4/16




Figure 7




Figure 8

5/16




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15




Figure 16




Figure 17




Figure 18




Figure 19




Figure 20




Figure 21

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/AU2014/000922

## A. CLASSIFICATION OF SUBJECT MATTER

[See Supplemental Sheet]

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN: Chemical Abstracts Registry and CAPLUS. Substructure search of general formula in claim 1.

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	Documents are listed in the continuation of Box C	



Further documents are listed in the continuation of Box C



See patent family annex

* Special categories of cited documents:		
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family	
"P" document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search  
29 October 2014Date of mailing of the international search report  
29 October 2014

## Name and mailing address of the ISA/AU

AUSTRALIAN PATENT OFFICE  
PO BOX 200, WODEN ACT 2606, AUSTRALIA  
Email address: pct@ipaaustralia.gov.au

## Authorised officer

Scott Makin  
AUSTRALIAN PATENT OFFICE  
(ISO 9001 Quality Certified Service)  
Telephone No. +61 2 6283 7971

INTERNATIONAL SEARCH REPORT		International application No.
C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		PCT/AU2014/000922
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PETERS, M. et al, "A Modular Synthesis of Teraryl-Based $\alpha$ -Helix Mimetics, Part 1: Synthesis of Core Fragments with Two Electronically Differentiated Leaving Groups", Chemistry – A European Journal, 2013, vol. 19, pp. 2442-2449 Abstract; Introduction; Compounds 1a-c, scheme 5, p. 2445	1-19
A	HUANG, W.-J. et al, "Synthesis and Biological Evaluation of ortho-Aryl N-Hydroxycinnamides as Potent Histone Deacetylase (HDAC) $\alpha$ Isoform-Selective Inhibitors", ChemMedChem, 2012, vol. 7, pp. 1815-1824 Abstract; Fig. 2, p. 1818; Table 4, p. 1820; Compound 22d, table 3, p. 1819	1-19
A	US 2011/0082109 A1 (MIYANAGA et al) 07 April 2011 Abstract; Claims; Paragraph [0118]; Examples 40-41 (p. 24), 48 (p. 26), 72 (p. 35), 75-78 (pp. 36-37) and 80 (p. 38)	1-19
A	WO 2006/018325 A1 (GALDERMA RESEARCH & DEVELOPMENT, S.N.C.) 23 February 2006 Abstract; Claims; p. 14, lines 25-26; Compounds 85-87, 95, pp. 9-10)	1-19
A	WO 2005/120545 A1 (DUGGAN) 22 December 2005 Abstract; Claims; Examples, pp. 10-14	1-19

## Supplemental Box – IPC Marks

*C07C 235/34 (2006.01)*

*C07C 233/11 (2006.01)*

*C07C 233/07 (2006.01)*

*C07C 237/20 (2006.01)*

*C07C 255/00 (2006.01)*

*C07C 271/28 (2006.01)*

*C07C 275/28 (2006.01)*

*C07C 307/10 (2006.01)*

*C07C 311/03 (2006.01)*

*C07D 209/18 (2006.01)*

*C07D 215/02 (2006.01)*

*C07D 231/56 (2006.01)*

*C07D 213/74 (2006.01)*

*C07D 235/04 (2006.01)*

*C07D 249/18 (2006.01)*

*C07D 263/54 (2006.01)*

*C07D 277/62 (2006.01)*

*A61K 31/165 (2006.01)*

*A61K 31/17 (2006.01)*

*A61K 31/18 (2006.01)*

*A61K 31/277 (2006.01)*

*A61K 31/404 (2006.01)*

*A61K 31/416 (2006.01)*

*A61K 31/4184 (2006.01)*

*A61K 31/428 (2006.01)*

*A61K 31/47 (2006.01)*

*A61P 9/12 (2006.01)*

*A61P 13/12 (2006.01)*

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/AU2014/000922

This Annex lists known patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document/s Cited in Search Report		Patent Family Member/s	
Publication Number	Publication Date	Publication Number	Publication Date
US 2011/0082109 A1	07 April 2011	US 2012088737 A2	12 Apr 2012
		JP 2012012376 A	19 Jan 2012
WO 2006/018325 A1	23 February 2006	AR 050695 A1	15 Nov 2006
		AU 2005274277 A1	23 Feb 2006
		BR PI0515209 A	08 Jul 2008
		CA 2575029 A1	23 Feb 2006
		CN 101014328 A	08 Aug 2007
		EP 1781274 A1	09 May 2007
		EP 1781274 B1	05 Nov 2008
		FR 2874378 A1	24 Feb 2006
		FR 2874378 B1	27 Oct 2006
		JP 2008509965 A	03 Apr 2008
		KR 20070042553 A	23 Apr 2007
		MX 2007001943 A	09 May 2007
		RU 2007109778 A	27 Sep 2008
		US 2007207175 A1	06 Sep 2007
WO 2005/120545 A1	22 December 2005	AU 2005251386 A1	22 Dec 2005
		AU 2005251386 B2	13 May 2010
		BR PI0511367 A	04 Dec 2007
		CA 2573439 A1	22 Dec 2005
		CN 1993137 A	04 Jul 2007
		EP 1768689 A1	04 Apr 2007
		EP 1768689 B1	20 Aug 2014
		IL 179931 A	29 Aug 2013
		JP 2008501723 A	24 Jan 2008
		JP 4824682 B2	30 Nov 2011
		KR 20140057411 A	12 May 2014
		MX PA06014462 A	23 May 2007
		NZ 552130 A	30 Oct 2009
		RU 2007101164 A	20 Jul 2008
		US 2008108573 A1	08 May 2008

Due to data integration issues this family listing may not include 10 digit Australian applications filed since May 2001.

Form PCT/ISA/210 (Family Annex)(July 2009)

**INTERNATIONAL SEARCH REPORT**

Information on patent family members

International application No.

**PCT/AU2014/000922**

This Annex lists known patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

<b>Patent Document/s Cited in Search Report</b>		<b>Patent Family Member/s</b>	
<b>Publication Number</b>	<b>Publication Date</b>	<b>Publication Number</b>	<b>Publication Date</b>
		US 7951777 B2	31 May 2011
		ZA 200700070 A	28 May 2008

**End of Annex**

Due to data integration issues this family listing may not include 10 digit Australian applications filed since May 2001.

Form PCT/ISA/210 (Family Annex)(July 2009)