




(72) SHARPE, DAVID, US

(72) NATRAJAN, ANAND, US

(72) JIANG, QINGPING, US

(72) PARSONS, GEORGE, US

(72) LAW, SAY-JONG, US


(71) BAYER CORPORATION, US

(51) Int.Cl.⁷ G01N 33/58, C07D 215/50, C12Q 1/32, C12Q 1/00,
C07D 221/12, C07D 219/04

(30) 1998/11/25 (60/109,823) US


(54) **MESURE D'HYDRURE PAR UTILISATION DE COMPOSES
CHIMIO Luminescents D'ACRIDINIUM**

(54) **MEASUREMENT OF HYDRIDE USING CHEMILUMINESCENT
ACRIDINIUM COMPOUNDS**



(57) La présente invention concerne une méthode permettant de mesurer un hydrure par utilisation d'un composé chimioluminescent, tel qu'un composé d'acridinium. La source de l'hydrure pour la réduction du composé d'acridinium peut être d'origine chimique ou biochimique, ou le résultat d'une catalyse enzymatique. La source chimique de l'hydrure pourrait être NaBH₄. La source biochimique de l'hydrure

(57) The present invention discloses a method for the measurement of hydride using a chemiluminescent compound, such as an acridinium compound. The source of hydride for the reduction of acridinium compound may be of chemical or biochemical origin, or the result of enzymatic catalysis. The chemical source of hydride, might be NaBH₄. A biochemical source of hydride might be that derived from NADH, or NADPH, while an



(21) (A1) **2,319,187**
(86) 1999/11/24
(87) 2000/06/02

pourrait dériver de NADH, ou de NADPH, tandis qu'une source enzymatique serait de la classe des oxydoréductases appelées déshydrogénases qui convertissent NADH ou NADPH à partir de NAD ou NADP. Parmi les applications des composés d'acridinium comme indicateurs chimioluminescents d'hydruure, citons des essais diagnostiques qui utilisent les déshydrogénases comme réactifs, indicateurs, marqueurs diagnostiques ou encore comme étiquettes. L'éthanol, par exemple, pourrait être détecté avec la chimioluminescence d'un ester d'acridinium par réaction de déshydrogénase d'alcool sur l'éthanol, ladite réaction produisant NADH. Dans son rôle d'étiquette, la déshydrogénase peut être utilisée dans une ELISA pour la détection d'un analyte spécifique, auquel cas l'ester d'acridinium fournit le signal de réponse.

enzymatic source would be the class of oxidoreductases termed dehydrogenases which convert NADH or NADPH from NAD or NADP. Among applications for acridinium compounds as chemiluminescent indicators of hydride are diagnostic assays using dehydrogenases as reagents, indicators, diagnostic markers or as labels. Ethanol, for example, might be detected with acridinium ester chemiluminescence through the reaction of alcohol dehydrogenase on ethanol, said reaction producing NADH. As a label, dehydrogenase might be used in an ELISA for the detection of a specific analyte with acridinium ester providing the signaling response.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : G01N 33/58, C12Q 1/00, 1/32, C07D 219/04, 221/12, 215/50	A1	(11) International Publication Number: WO 00/31543 (43) International Publication Date: 2 June 2000 (02.06.00)
(21) International Application Number: PCT/IB99/01894 (22) International Filing Date: 24 November 1999 (24.11.99) (30) Priority Data: 60/109,823 25 November 1998 (25.11.98) US (71) Applicant (for all designated States except US): BAYER CORPORATION [US/US]; 333 Coney Street, East Walpole, MA 02032 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): SHARPE, David [US/US]; 61 Cross Street, Foxborough, MA 02035 (US). NATRAJAN, Anand [US/US]; 77 Madeline Road, Manchester, NH 03104 (US). JIANG, Qingping [US/US]; 14 Stratton Way, Northborough, MA 01532 (US). PARSONS, George [US/US]; 23 Brewster Road, Arlington, MA 02746 (US). LAW, Say-Jong [US/US]; 15 Forbes Road, Westwood, MA 02090 (US). (74) Agents: ROESLER, Judith, A.; Law & Patent Department, Bayer Corporation, 63 North Street, Medfield, MA 02052 (US) et al.		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>

(54) Title: MEASUREMENT OF HYDRIDE USING CHEMILUMINESCENT ACRIDINIUM COMPOUNDS**(57) Abstract**

The present invention discloses a method for the measurement of hydride using a chemiluminescent compound, such as an acridinium compound. The source of hydride for the reduction of acridinium compound may be of chemical or biochemical origin, or the result of enzymatic catalysis. The chemical source of hydride, might be NaBH₄. A biochemical source of hydride might be that derived from NADH, or NADPH, while an enzymatic source would be the class of oxidoreductases termed dehydrogenases which convert NADH or NADPH from NAD or NADP.

Among applications for acridinium compounds as chemiluminescent indicators of hydride are diagnostic assays using dehydrogenases as reagents, indicators, diagnostic markers or as labels. Ethanol, for example, might be detected with acridinium ester chemiluminescence through the reaction of alcohol dehydrogenase on ethanol, said reaction producing NADH. As a label, dehydrogenase might be used in an ELISA for the detection of a specific analyte with acridinium ester providing the signaling response.



WO 00/31543

PCT/IB99/01894

TITLE OF THE INVENTION

Measurement of Hydride Using Chemiluminescent Acridinium Compounds

5

CROSS REFERENCE TO RELATED APPLICATIONS

U.S. Provisional Application No. 60/109,823 filed November 25, 1998 entitled: Modulation of Chemiluminescent Acridinium Compounds by Hydrides or Hydride Generating Systems and Applications Thereof.

10

BACKGROUND OF THE INVENTION

Acridinium esters (AE) have provided for an extremely sensitive method of detection and have been used extensively as chemiluminescent labels in both immuno- and nucleic acid assays. Hydrolytically stable, Polysubstituted Aryl Acridinium Esters (PAAE) have proven useful as analytical labels (U.S. Patent Nos. 4,745,181; 4,918,192; and 5,110,932) with a variety of linkages (U.S. Patent Nos. 5,241,070; 5,538,901; and 5,663,074) and were the first chemiluminescent, acridinium compounds to satisfy the stringent requirements of commercial ligand binding assays. The utility of PAAE was further enhanced with the advent of Functionalized Hydrophilic PAAE (U.S. Patent No. 5,656,426) which increased the quantum yield of PAAE and enhanced the performance of PAAE-labeled binding partners in terms of the observed signal to noise ratios and the sensitivities of various binding assays. Additionally, introduction of ionizable groups at the phenoxy moiety produced another sub-class of hydrophilic PAAE (U.S. Patent Nos. 5,227,489; 5,449,556; and 5,595,875).

15

20

25

30

WO 00/31543

PCT/IB99/01894

-2-

5 M. Kawaguichi, et al. (Bioluminescence and Chemiluminescence, Proceedings of 9th International Symposium 1996, Ed. Hastings, Kricka and Stanley, John Wiley & Sons, 1997, pp. 480-484) have described
10 stabilized phenyl acridinium esters for chemiluminescent immunoassays. AE derivatives with additional methyl substitutions at C-1, which are optional at C-3 of the acridinium nucleus with matching mono- or di-methyl
15 substitutions at the ortho-positions of the phenoxy moiety, were shown to have excellent stability in aqueous solution.

EP 0324,202 A1 and subsequently EP 0609,885 A1 both describe acridinium esters with functional groups substituted at the nitrogen atom of the acridinium
20 nucleus. The latter application further describes alternate substituents such as the biphenyl or naphthyl moieties as possible replacements for the phenyl group.

Mattingly, et al. (U.S. Patent Nos. 5,468,646 and 5,543,524) describe chemiluminescent acridinium salts,
25 and their applications in immunoassays. These acridinium salts belong to another class of compounds termed acridinium sulfonylamides (or N-sulfonylacridinium carboxamides). The acridinium sulfonylamides (AS) have aqueous stabilities which are
30 comparable with PAAE. Mattingly, et al. further describe and claim the analogous chemiluminescent phenanthridinium salts, and their applications in immunoassays, in U.S. Patent Nos. 5,545,739; 5,565,570, and 5,669,819. Additionally, in these patents a general structure of acridinium sulfonylamides is described showing possible substitutents of a Markush group at the acridinium nucleus.

WO 00/31543

PCT/IB99/01894

-3-

Conventional acridinium compounds, such as those described in the aforementioned patents and literature, emit light with maxima at about 428 nm upon reaction with hydrogen peroxide in strong alkaline solution.

5 Acridinium compounds which emit light of wavelength maxima >500 nm have also been described in the prior arts. U.S. Patent Nos. 5,395,752; 5,702,887 and 5,879,894 describe novel, long-emission acridinium esters (LEAE), where a fused, benzacridinium system is employed

10 to extend the wavelength of emission of the acridinium ester. In the copending PCT application PCT/IB98/00831 Jiang et al. have further extended the PAAE emission maxima well into the region of 600-700 nm by utilizing the principle of energy transfer. This entailed the

15 covalent coupling of luminophores to acridinium ester. When the chemiluminescent reactions of these conjugates were initiated by treatment with alkaline peroxide, light emission was observed at long wavelengths where the wavelength maxima depended upon the structure of the

20 luminophore. In the more recent copending PCT application PCT/US99/18076, Natrajan et al. describe novel acridinium compounds that have intrinsic emission maxima close to or in the near infrared region (>590 nm). The structural requirements for such long wavelength-

25 emitting acridinium compounds are disclosed.

N-Alkylacridan esters obtained from the reduction of acridinium esters have been used as enzyme substrate indicators for the determination of phosphatases and oxidases and their substrates or products. N-alkylacridan

30 phosphate esters have been engineered as substrates for the direct detection of minute concentrations of alkaline

WO 00/31543

PCT/IB99/01894

-4-

phosphatase (Akhavan-Tafti, H et al. "Lumagen™ APS: New Substrates for the Chemiluminescent Detection of Phosphatase Enzymes"; Proc. 9th. Internat'l. Symp. Bioluminescence and Chemiluminescence; (1996); Hastings, J. W.; Kricka, L. J.; Stanley; P. E., (Eds.); John Wiley & Sons, Inc., New York, NY; pp. 311-314). Similarly, N-alkylacridancarboxylate esters have been applied as oxidizable indicators for horseradish peroxidase, where the chemiluminescence from the oxidized acridinium ester product was used to quantify either horseradish peroxidase or oxidases and their substrates in coupled enzymatic reactions (Akhavan-Tafti, H et al.; Chemiluminescent Detection of Oxidase Enzymes by Peroxidase-mediated Oxidation of Acridan Compounds; Proc. 9th. Internat'l. Symp. Bioluminescence and Chemiluminescence; (1996); Hastings, J. W.; Kricka, L. J.; Stanley; P. E., (Eds.); John Wiley & Sons, Inc., New York, NY; pp. 501-504).

Luminols along with peroxidase have been used as chemiluminescent detectors of hydrogen peroxide generated from dehydrogenase and its cofactors (WO 95/29255).

Various classes of chromogenic, hydride-reducible indicators have been described (Ottaway, J. M.; Oxidation-Reduction Indicators; Internat'l Ser. Monographs Anal. Chem.; (1968); Belcher, R.; Frieser, H., (Eds.); Plenum; pp. 469-529) (Bird, C. L.; Kuhn, A. T.; Electrochemistry of the Viologens; Chem. Soc. Rev.; (1981), 10, pp. 49-82). Several of these, including the oxidized salts of phenazines, phenoxazines and phenothiazines, have been used as chromogenic indicators of hydride from the reduced nicotinamide cofactors

WO 00/31543

PCT/IB99/01894

-5-

(dihydronicotinamide adenine dinucleotide, NADH, and dihydronicotinamide adenine dinucleotide phosphate, NADPH) and the reduced flavin cofactors (dihydroflavin mononucleotide, FMNH₂ and dihydroflavin adenine dinucleotide, FADH₂) generated by the enzymatic activity of dehydrogenases (Czerlinski, G. H., et al., "Coupling of Redox Indicator Dyes into an Enzymatic Reaction Cycle", J. Biochem. Biophys. Methods, (1988) 15, pp. 241-248) (Nakamura, S., et al., "Use of 1-Methoxy-5-methylphenaziniummethylsulfate in the Assay of Some Enzymes of Diagnostic Importance", Clin. Chim. Acta, (1980), 101, p. 321).

BRIEF SUMMARY OF THE INVENTION

The present invention discloses a method for the measurement of hydride using a chemiluminescent compound. The preferred chemiluminescent molecule is an acridinium compound. The source of hydride for the reduction of acridinium compound may be of chemical or biochemical origin, or the result of enzymatic catalysis. The chemical source of hydride, for example, might be metal hydrides, such as NaBH₄. A biochemical source of hydride might be that derived from NADH, NADPH, FMNH₂ or FADH₂, while an enzymatic source would be the class of oxidoreductases termed dehydrogenases which convert in redox reactions NADH, NADPH, FMNH₂ or FADH₂ from NAD, NADP, FMN or FAD.

There are numerous potential applications for acridinium compounds as chemiluminescent indicators of hydride. Any applied tests or diagnostic assays, in which hydride is either present at the onset of or generated through the course of a reaction, would benefit

WO 00/31543

PCT/IB99/01894

-6-

from the present invention. Such tests, which could encompass many different formats as discussed below in detail, may involve the quantitation or detection of metal hydrides, or enzyme cofactors such as NADH, NADPH, FMNH₂, or FADH₂. Of particular importance, are those diagnostic assays which might use dehydrogenases as reagents, indicators, diagnostic markers or as labels. Ethanol, for example, might be detected with acridinium ester chemiluminescence through the reaction of alcohol dehydrogenase on ethanol, said reaction producing NADH. As a label, dehydrogenase might be used in an ELISA for the detection of a specific analyte with acridinium ester providing the signaling response. Nucleic acid assays using dehydrogenase as a label are also envisioned. Assays for the detection of clinically relevant dehydrogenases such as elevated glutamate dehydrogenase as an indicator of hepatocellular damage might also be developed.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is the reaction showing the addition of hydride to C-9 of the acridinium nucleus.

Figure 2 shows reduction of 2',6'-Dimethylphenyl-10-Methyl Acridinium-9-Carboxylate with NADH in DMF/0.1 M phosphate pH 7.4 at room temperature for 10 minutes.

Figure 3 shows the general structure of the main acridinium compounds useful as the chemiluminescent indicators of the present invention.

Figure 4 shows the additional structures that can be linked to R₂ and R₃ to form an additional ring.

Figure 5 shows the polysubstituted aryl moiety identified as Y.

WO 00/31543

PCT/IB99/01894

-7-

Figure 6 shows examples of the leaving group R_{10} .

Figure 7 shows the general structure of phenanthridinium and quinolinium compounds useful as the chemiluminescent indicators of the present invention.

5 Figure 8 shows the addition of a hydride to an electron-deficient ester moiety to form an electron-rich moiety.

10 Figure 9 shows the addition of hydride to a long-wavelength emission compound to form a short-wavelength emission compound.

Figure 10 shows an acridinium nucleus wherein the reduction of the side chain quencher is preferentially achieved.

15 Figure 11 shows an acridinium nucleus where reduction of the side chain fluorophore is preferentially achieved.

Figure 12 shows an analytical procedure wherein NADH formation can be quantified.

20 Figure 13 shows a homogeneous immunoassay using an acridinium compound as a hydride indicator.

Figure 14 shows an assay in which interfering substances are eliminated via the use of a solid phase in the assay.

25 Figure 15 illustrates heterogeneous assays using the instant invention.

Figure 16 shows the chemiluminometric Emit® theophylline assay using NSP-DMAE₄-BSA as a hydride indicator.

30 Figure 17 shows the chemiluminometric Emit® valproate assay using NSP-DMAE₄-BSA as a hydride indicator.

WO 00/31543

PCT/IB99/01894

-8-

Figure 18 shows the chemiluminometric Emit® quinidine assay using NSP-DMAE₄-BSA as a hydride indicator.

Figure 19 shows the chemiluminometric Emit® theophylline assay using DMAE- ϕ as a hydride indicator.

Figure 20 shows the chemiluminometric Emit® valproate assay using DMAE- ϕ as a hydride indicator.

Figure 21 shows referenced vs. determined analyte concentration for controls.

Figure 22 shows a chemiluminometric ethanol assay.

DETAILED DESCRIPTION OF THE INVENTION

Central to the present invention is our discovery that the chemiluminescent activity of an acridinium compound can be modulated (increased or decreased) by hydride. Although the examples discussed herein result in a decrease in chemiluminescent activity, several mechanisms are contemplated wherein one will see an increase in chemiluminescent activity. For example, in Section B2 and part of Section B3, below, chemiluminescence can be increased when the detector is sensitive to a selective range of emission wavelength. If the reaction results in a shift of the emission to a wavelength where a detector is more sensitive, increased chemiluminescence will be observed. In the other part of Section B3, below, an increase is observed because the moiety causing quenching is reduced and inactivated by the hydride.

A. Novel chemiluminescent indicators of hydride

We have discovered that reduction of acridinium ester with hydride suppresses the chemiluminescent

WO 00/31543

PCT/IB99/01894

-9-

activity of the acridinium ester. The reduction in chemiluminescent activity is the result of the addition of hydride to C-9 of the acridinium nucleus (see Figure 1). The product, which is the acridan, is unable to participate in the light-generating reaction with alkaline hydrogen peroxide. Other chemiluminescent acridinium compounds or analogs (e.g. benzacridinium, phenanthridinium, quinolinium compounds or isomers) and their derivatives, as well as chemiluminescent compounds such as lucigenin and its derivatives, share the same or similar mechanism of hydride attack to the electron-deficient acridinium nucleus to form the reduced acridans.

Our initial studies focused on the reduction of 2',6'-dimethylphenyl-10-methyl acridinium-9-carboxylate (DMAE- Φ) with the chemical reductant sodium borohydride in methanol solvent. Chromatographic analysis (HPLC) of this reduction reaction revealed clean conversion to the corresponding N-methylacridan, correlating with a drop in chemiluminescent activity when the reaction mixture was treated with alkaline peroxide. A similar result was also observed when the hydride donor NADH was employed to reduce the acridinium ester. Chromatographic analysis of this reaction too revealed clean formation of the acridan with a concomitant loss of chemiluminescent activity. The extent of this decrease was found to depend upon the concentration of NADH. As shown in the table in Figure 2, increasing quantities of NADH were accompanied by a corresponding decrease in the chemiluminescent activity of the reaction mixture. The reduction reaction was complete in less than 10 minutes.

WO 00/31543

PCT/IB99/01894

-10-

Measurement of hydride generated in enzymatic systems has been used in the prior art to devise sensitive assays. For example, Cook et al. in Clin. Chem. 1993, 39/6, 965-971 describe an enzyme amplification system to detect human proinsulin in human plasma. This assay involves using alkaline phosphatase as the primary label which is used to dephosphorylate NADP⁺ to NAD⁺. The latter compound is then employed as a cofactor in a redox cycle using the enzymes alcohol dehydrogenase and diaphorase. While the former enzyme utilizes NAD⁺ to oxidize ethanol to acetaldehyde with generation of NADH, the latter enzyme diaphorase utilizes the NADH for the reduction of a tetrazolium dye to produce a colored formazan dye. Using this system, Cook et al. were able to detect 1 zeptomole of alkaline phosphatase and 0.017 pmol/L of human proinsulin. The same system was employed by Johannsson et al. (J. Immunol. Meth. 1986, 87, 7-11) to devise a highly sensitive assay for TSH (sensitivity of 0.0013 μ IU/L). Both the above assays use a colored dye as an indicator. Since chemiluminescent indicators such as acridinium esters are much more sensitive, measuring hydride using these compounds is likely to produce more sensitive assays. We envisage other interesting and useful applications of our finding which we will elaborate in greater detail later.

The reduction of 2',6'-dimethylphenyl-10-methyl acridinium ester with NADH involves addition of hydride to C-9 to generate the non-chemiluminescent acridan. It is evident that acridinium compounds and analogs with widely different structural modifications, but containing the same core acridinium nucleus or the analogous

WO 00/31543

PCT/IB99/01894

-11-

benzacridinium, phenanthridinium, or quinolinium nucleus will be amenable to this transformation. Such compounds can be generalized with the structure(s), which have been described in all the prior arts references incorporated herein in the background section at the beginning of this disclosure. These structures include not only blue-emitting (about 420-490 nm), but also green (about 490-570 nm), yellow (about 570-580 nm), orange (about 580-595 nm) and red (about 595-780 nm)-emitting acridinium compounds and said analogs.

For further illustration, the general structure of the main acridinium compounds useful as the chemiluminescent indicators of the present invention can be schematically represented as shown in Figure 3, wherein

R_1 , is an alkyl, alkenyl, alkynyl or aralkyl containing optionally up to 20 heteroatoms; preferably R_1 is methyl or sulfoalkyl group.

R_2 , R_2' , and R_3 , are identical or different, selected from hydrogen, -R, substituted or unsubstituted aryl (ArR or Ar), halide, amino, hydroxyl, nitro, sulfonate, -CN, -COOH, -SCN, -OR, -SR, -SSR, -C(O)R, -C(O)OR, -C(O)NHR, or -NHC(O)R;

Throughout this application, R is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, and aralkyl containing optionally up to 20 heteroatoms;

Alternatively, R_2 and R_3 can be linked as shown in Figure 4, so as to form an additional ring fused to the attached acridinium nucleus. The C_1 , C_2 , C_3 , C_4 , C_5 , and C_8 , peri-positions of the acridinium nucleus are optionally substituted as represented by R_2' .

WO 00/31543

PCT/IB99/01894

-12-

A⁻ is a counter ion which is introduced to pair with the quaternary nitrogen of the acridinium nucleus either as a result of quaternarizing the acridine ring nitrogen by the use of alkylating agents during the synthesis, modification of the R₁, or subsequent exchange that occurs during the work-up of reaction mixtures and purification of desired compounds in a solution or fluid containing excess amount of other anions. Examples of the counter ions include CH₃SO₄⁻, FSO₃⁻, CF₃SO₃⁻, C₄F₉SO₃⁻, CH₃C₆H₄SO₃⁻, halide, CF₃COO⁻, CH₃COO⁻, and NO₃⁻,

X is nitrogen, oxygen or sulfur;

When X is oxygen or sulfur, Z is omitted and Y is a polysubstituted aryl moiety shown in Figure 5, where R₄ and R₈ can be (1) hydrogen or (2) alkyl, alkenyl, alkynyl, alkoxyl (-OR), alkylthiol (-SR), or substituted amino groups that serve to stabilize the -COX- linkage between the acridinium nucleus and the Y moiety, through steric and/or electronic effects. Preferably, one of them is as defined below while the other is a hydrogen, if the C₁ or C₈ position of the acridinium nucleus is substituted with a lower alkyl group, preferably methyl. More preferably, R₄ and R₈ are alkyl, alkenyl, alkynyl, alkoxyl (-OR), alkylthiol (-SR), or substituted amino groups that serve to stabilize the -COX- linkage between the acridinium nucleus and the Y moiety, through steric and/or electronic effect. Most preferably R₄ and R₈ are lower alkyl (e.g., a methyl group).

R₅ and R₇ are any of R₂, R₂', and R₃ defined above;

R₆ is also any of R₂, R₂', and R₃ defined above, when the acridinium compound is used as a free chemiluminescent indicator.

WO 00/31543

PCT/IB99/01894

-13-

Alternatively, the acridinium compounds can be covalently attached to a more water-soluble polymer (natural or synthetic) or biopolymers (e.g. proteins, polysaccharides, glycoproteins, and nucleic acids) in a conjugate form to enhance its water solubility for practical, commercial utility as chemiluminescent indicator. It will then be necessary to impart a reactive functional group, preferably, onto R_6 to facilitate covalent linkage formation between the acridinium compound and the water soluble polymer or biopolymer of choice.

Thus, R_6 can also be $-R_9-R_{10}$, where R_9 is not required but optionally can be branched or straight-chained alkyl, substituted or unsubstituted aryl or aralkyl containing optionally up to 20 heteroatoms, and R_{10} is a leaving group or an electrophilic functional group attached with a leaving group including but not limited to those shown in Figure 6. R_{10} can also be $-Q-R-Nu$, $-Q-R-(I)nNu-$, $-Q-Nu$, $-R-Nu$, or $-Nu$, where n is a number of at least 1, Nu is a nucleophilic group, Q is a functional linkage, I is an ionic or ionizable group; detailed definitions of Nu , Q , and I can be found in the US Patent #5,241,070, column 3, line 45 to column 3, line 16. The reactions contemplated for Nu were also described in the same patent, column 3, line 48 to column 4, line 18.

R_5 and R_6 , and R_6 and R_7 are interchangeable; and

When X is nitrogen, then Z is $-SO_2-Y'$, Y' has the same definition of Y as described above, and both can be the same or different. Additionally, Y itself can be a branched or straight-chained alkyl containing optionally up to 20 carbon atoms, halogenated or

WO 00/31543

PCT/IB99/01894

-14-

unhalogenated, or a substituted aryl, or heterocyclic ring system.

Similarly, the general structure of the phenanthridinium and quinolinium compounds useful as the chemiluminescent indicators of the present invention can be schematically represented as shown in Figure 7. All the definitions of the substituents and counter-ion of the phenanthridinium and quinolinium nucleus discussed above are the same as those described for the acridinium compounds earlier, except for the possible peri-positions for R₂' substituent have to be re-designated at the C₁, C₂, C₃, C₄, C₆, and C₉ positions of the phenanthridinium nucleus and C₂, C₃, C₅, and C₈ positions of the quinolinium nucleus.

B. Alternative modes of hydride modulation of chemiluminescent acridinium compounds

In addition to reduction of acridinium esters to acridans, hydride such as NADH can also be employed to modulate the chemiluminescent activity of acridinium compounds in other novel and interesting ways which can be potentially useful. Several such approaches are summarized below all of which require chemical selectivity (regioselectivity) in the reduction reaction.

1. Modulation of the Acridinium Ester or Sulfonylamide cleavage lability

In one approach, hydride can be directed to the ester or sulfonylamide moiety (instead of the acridinium nucleus) of the acridinium compound to effect a change in the emission characteristics of the compound. It is well

WO 00/31543

PCT/IB99/01894

-15-

known that acridinium esters with poor leaving groups are poorly chemiluminescent. Poor leaving groups are typically electron-rich and can be generated by hydride addition to electron-poor functional groups. By
5 employing this procedure, an acridinium ester or sulfonylamide that contains an electron-deficient ester or sulfonylamide moiety can be converted to one which is electron-rich, thereby converting a strongly chemiluminescent compound to a weak one. Figure 8
10 illustrates how this is achieved. While carrying some of the necessary basic substituents and the counter-ion as defined in section A above, the acridinium nucleus in the illustration is deliberately engineered to be electron-rich by the attachment of one or more desired electron-donating groups (relative to hydrogen and designated with
15 letter D), which are as defined in "Advanced Organic Chemistry" Jerry March Ed., 4th edition, page 18-19, at one or more optimal positions of the acridinium nucleus, so that reduction of the N-methylpyridinium moiety is
20 achieved preferentially. If more than one electron-donating groups (D's) are substituted at the acridinium nucleus, the D's can be the same or different. To impart other important or desirable structural characteristics (e.g. hydrophilicity, functional group, etc.), other
25 additional substituents are permissible at the peripositions of the acridinium nucleus, as long as the combined electronic effect of all substituents is that of electron donating. Obviously, the same teaching applies to the analogs of the acridinium ester and sulfonylamide
30 as described earlier in section A.

WO 00/31543

PCT/IB99/01894

-16-

2. Modulation of emission wavelength maxima of acridinium compounds:

Reduction can also be used to alter the emission-wavelength of an acridinium compound. The emission-wavelength of blue-emitting acridinium compounds can be shifted to long wavelengths by creating an extended electronic conjugation system at the acridinium nucleus as discussed in PCT application PCT/US99/18076. By directing reduction to disrupt the extended conjugated system in an acridinium compound, emission wavelength of the acridinium compound can be shifted from long to short wavelength, hence varying the chemiluminescent intensity of the selected wavelength range under monitoring. Figure 9 illustrates the above principle. Again, while carrying some of the necessary basic substituents and the counter-ion as defined in section A above, the acridinium nucleus in the illustration is also deliberately engineered to be electron-rich by the attachment of one or more desired electron-donating groups (relative to hydrogen and designated with letter D), at one or more optimal positions of the acridinium nucleus, so that reduction of the side chain extended conjugation system is achieved preferentially. If more than one electron-donating groups (D's) are substituted at the acridinium nucleus, the D's can be the same or different. To impart other important or desirable structural characteristics (e.g. hydrophilicity, functional group, etc.), other additional substituents are permissible at the peri-positions of the acridinium nucleus, as long as the combined electronic effect of all substituents is that of electron donating. Obviously, the same teaching applies to the analogs of

WO 00/31543

PCT/IB99/01894

-17-

the acridinium ester and sulfonylamide as described earlier in section A.

3. Modulation of quencher/fluorophore moiety of
5 acridinium compounds:

Other modes of modulating the chemiluminescent activity are illustrated in the scheme in Figure 9 where the hydride resulting from an enzymatic reaction alters the structure of the hydride-reducible quencher that is
10 covalently linked to the acridinium moiety having group D, the purpose of which was described above, and thus alters the quenching effect. The prerequisite of the quencher here is its ability of quenching the chemiluminescent activity of the acridinium compound via
15 dipole-dipole resonance energy transfer or other mechanisms, and its ability to be reduced by hydride preferentially over the reduction of the acridinium moiety. The reduction in quantum yield due to the quenching effect of the quencher can be reversed by the
20 interaction of hydride with the quencher. Thus, in the presence of hydride, the quantum yield of the acridinium moiety of the conjugate will increase. One obvious advantage of modulating the chemiluminescent activity by hydride using acridinium-quencher conjugate is the
25 positive correlation between the concentration of the hydride and chemiluminescent activity of the conjugate.

Again, while carrying some of the necessary basic substituents and the counter-ion as defined in section A above, the acridinium nucleus in Figure 10 can be
30 attached with one or more desired electron-donating groups, D's at one or more optimal positions of the acridinium nucleus, so that reduction of the side chain

WO 00/31543

PCT/IB99/01894

-18-

quencher is achieved preferentially. If more than one electron-donating groups (D's) are substituted at the acridinium nucleus, the D's can be the same or different. Similarly, to impart other important or desirable structural characteristics (e.g. hydrophilicity, functional group, etc.), other additional substituents are permissible at the peripositions of the acridinium nucleus, as long as the combined electronic effect of all substituents is that of electron donating. Obviously, the same teaching also applies to the analogs of the acridinium ester and sulfonylamide as described earlier in section A.

An alternative mode similar to the use of the acridinium compound-quencher conjugate is to replace the quencher with a fluorophore to form the AE-fluorophore conjugate. Acridinium-fluorophore conjugates, where the fluorophore is linked to the acridinium nucleus, in their chemiluminescent reactions, emit light in a region of the spectrum characteristic of the fluorophore. This occurs via resonance energy transfer from the electronically excited, acridone moiety to the fluorophore. The principle and examples of the AE-fluorophore conjugates are disclosed in more detail in PCT Application PCT/IB98/00831, which is incorporated herein by reference. One of the examples of this novel class of compounds is the conjugate of acridinium ester with rhodamine as shown below. The conjugate emits light at 628 nm due to efficient energy transfer from the AE moiety to rhodamine after it is treated with hydrogen peroxide in strong alkaline solution. The acridinium moiety can be modified by varying the D group so that the rhodamine moiety is selectively or

WO 00/31543

PCT/IB99/01894

-19-

preferentially reduced by hydride. As the result of the reduction, the light emission intensity at long wavelength due to the rhodamine moiety can decrease and the emission intensity at short wavelength due the
5 acridinium moiety can increase. Thus, the quantitation of hydride is made possible either by detecting the decrease of the long emission signal or by detecting the increase of the short emission signal.

Again, while carrying some of the necessary basic
10 substituents and counter-ions as defined in section A above, the acridinium nucleus in Figure 11 can be attached with one or more desired electron-donating groups, D's at one or more optimal positions of the acridinium nucleus, so that reduction of the side chain
15 fluorophore moiety is achieved preferentially. If more than one electron-donating groups (D's) are substituted at the acridinium nucleus, the D's can be the same or different. Similarly, to impart other important or desirable structural characteristics (e.g.
20 hydrophilicity, functional group, etc.), other additional substituents are permissible at the peripositions of the acridinium nucleus, as long as the combined electronic effect of all substituents is that of electron donating. Obviously, the same teaching also
25 applies to the analogs of the acridinium ester and sulfonylamide as described earlier in section A.

C. Applications of the chemiluminescent indicators of hydride

1. Conversion of a colorimetric to
30 chemiluminescent Emit[®] assay:

Modulating the chemiluminescent activity of

WO 00/31543

PCT/IB99/01894

-20-

acridinium compounds with hydride has numerous applications in analytical measurements. As mentioned earlier, colored indicator molecules to monitor NADH formation have been used to devise sensitive immunoassays (see above). We have discovered that NADH (and by analogy any hydride) formation in a system can be quantified using acridinium ester (and by analogy any acridinium-type compound), the present invention discloses an analytical procedure as shown in Figure 12. Here, A and AH represent oxidized and reduced forms of an acridinium compound whose chemiluminescent-emission properties are distinguishable. The hydride source can be chemical or biological. In addition, the hydride can be transferred either directly to the acridinium compound or indirectly through the intervention of an enzyme such as diaphorase.

One additional benefit of using the instant invention is that it allows a laboratory to measure hydride generation via the use of an instrument that measures chemiluminescence, thus eliminating the need to use a UV-visible spectrophotometer, which was needed in the past to measure hydride generation.

A widely used, commercial diagnostic test called Emit[®] (Emit[®] is a registered trademark of Behring Diagnostics GmbH and is defined below, Schneider, R. S. "Recent Advances in Enzyme Immunoassay" in Ligand Assay; Langan, J. and Clapp, J., Editors: Masson Publishing USA, Inc. 1981) employs a homogeneous immunoassay format and monitors and quantifies the formation of NADH (directly by UV spectrophotometry) whose concentration is correlated ultimately to the concentration of analyte.

WO 00/31543

PCT/IB99/01894

-21-

To demonstrate the utility of acridinium compounds as an excellent, quantitative indicator of NADH, we utilized this commercial test for three analytes (theophylline, quinidine and valproate). Similarly, an enzymatic assay for the determination of ethanol is also given in example 8 to demonstrate the utility of this approach. We also chose to employ the direct reduction of the acridinium compound with the NADH generated in the assay although it is obvious that this reduction can also be performed through the intervention of diaphorase.

The model chemiluminescent hydride indicators used for this study were 2',6'-dimethylphenyl-10-methyl-acridinium-9-carboxylate (DMAE- ϕ) free label and 2',6'-dimethylphenyl-10-(3'-sulfopropyl)-acridinium-9-carboxylate (NSP-DMAE) conjugated via a carboxamide linkage to bovine serum albumin (BSA). A specific demonstration of the utility of acridinium esters as chemiluminescent redox indicators of hydride is described below where chemiluminescent signal from NSP-DMAE or DMAE- ϕ is used in an enzyme-immunoassay for the quantitation of theophylline, quinidine and valproate in serum and also in an enzymatic assay for the quantitation of ethanol in serum. For this purpose a prototype assay has been developed on the Bayer Diagnostics Corp. ACS:180[®] (ACS:180[®] is a registered trademark of Bayer Corporation for the Automated Chemiluminescence Analyzer: a fully automated immunodiagnostic assay system) using Behring Diagnostics Emit[®] 2000 Theophylline and Model 1 Valproate Assay reagents and also Bayer Corporation Immuno 1[™] Emit[®] 2000 Quinidine Assay reagents as well as an enzymatic assay for the determination of ethanol in serum

WO 00/31543

PCT/IB99/01894

-22-

with the inclusion of an additional Chemiluminescent Redox Indicator reagent. DMAE- ϕ or NSP-DMAE-BSA conjugates are useful for applications in both organic and aqueous media. Since a broad variety of acridinium compounds which include the prior art hydrophilic AE's as described above have been well documented, the users of the present invention can select a highly water soluble acridinium compound to eliminate the use of organic solvents which are potentially detrimental to assay. Therefore, the user is provided with a number of options for selection of the chemiluminescent hydride indicator with appropriate solubility for application in an assay matrix with a given polarity.

Alternatively the acridinium compound selected as the chemiluminescent indicator for hydride might be conjugated to any number of small molecules, macromolecules or particulates, covalently or otherwise, to improve or impart properties such as, but not limited to, solubility, quantum yield, stability and resonance energy transfer.

Enzyme-multiplied immunoassay technique (Emit[®]) was originally developed by Syva Co., Inc. as a bioanalytical technology and is currently marketed in kits by Dade-Behring, Inc., chiefly for therapeutic drug monitoring (TDM) with some minor application for the detection of drugs of abuse. The automated, homogeneous Behring Diagnostics, Inc. Emit[®] assay series is currently one of several preferred diagnostics systems for TDM analytes. In these assays, a drug present in a patient sample will compete with a drug-enzyme conjugate for a limited quantity of antibody, directed against the drug. The

WO 00/31543

-23-

PCT/IB99/01894

conjugate's enzymatic activity, which in this case is that of the glucose-6-phosphate dehydrogenase (G6PDH) from Leuconostoc mesenteroides, is partially inhibited when bound to the drug-specific antibody. G6PDH catalyzes the reduction of oxidized nicotinamide adenine dinucleotide (NAD) to the reduced form, NADH, by hydride transfer from glucose-6-phosphate (G6P) and is monitored by the increased UV absorbance at 340 nm due to NADH formation. Endogenous sample G6PDH requires NADP, not NAD, for catalytic activity and will, therefore, not interfere with NAD reduction. Consequently, the G6PDH catalytic activity of the conjugate is correlated to the drug concentration in the patient sample through the rate of NADH formation, for example $\Delta A_{340 \text{ nm}}/\text{minute}$. A Behring Diagnostics, Inc. Emit[®] assay kit holds two reagents: Reagent A contains a buffered solution of G6P (substrate), NAD (enzyme cofactor), and murine monoclonal antibodies reactive to a particular drug analyte, while Reagent B consists of a buffered solution of drug-G6PDH conjugate.

Since the Bayer Diagnostics ACS:180[®] readout system employs a photomultiplier tube to quantitate chemiluminescent signal, adaptation of a Behring Diagnostics, Inc. Emit[®] assay to the Bayer Diagnostics ACS:180[®] requires an additional reagent for transduction of the colorimetric signal to chemiluminescence. This additional reagent is termed a chemiluminescent redox indicator of hydride (CHI), and the CHI is an acridinium compound. The following experiments have indicated that several acridinium ester derivatives are reduced by hydride transfer (most likely to the C-9 position of the

WO 00/31543

PCT/IB99/01894

-24-

N-methylacridinium nucleus) from an equimolar concentration of NADH in the milli- and micromolar range. The rate of reduction is slow, most likely because the predominant pseudobase form is not susceptible to hydride reduction, ensuring that the differential decrease in chemiluminescence is observed even in the presence of a great excess of available hydride. For example, reduced DMAE- ϕ (N-alkylacridan) is weakly chemiluminescent, when compared to DMAE- ϕ , and yields little detectable signal when flashed; therefore, although the comparative concentration of NADH introduced into a mixture may be far greater than that of DMAE- ϕ , the relative NADH concentration is correlated to the decrease in total chemiluminescence. This correlation has been established. DMAE- ϕ , as well as NSP-DMAE, should, therefore, function as a chemiluminescent redox indicator of hydride in an Emit[®] type assay, where increased analyte concentrations would increase the rate of NADH formation, thereby decreasing the chemiluminescence of a fixed quantity of acridinium compound included within the assay reaction.

The ability to measure changes in NADH concentrations which are in the micro to millimolar range, with acridinium compounds or conjugates in the nanomolar range, has permitted us to measure changes in analyte concentrations (as reflected in changes in NADH concentrations) directly, without having to either dilute NADH containing samples or adding a secondary sink reagent to scavenge the vast excess of NADH in undiluted samples.

A schematic representation of the above assay is

shown in Figure 13 and examples of three analytes (theophylline, quinidine and valproate) measured using the aforementioned technique are described in detail in example 5-8 as well as for an enzymatic assay for the quantitation of ethanol. The data clearly indicate that acridinium compounds are indeed excellent, quantitative indicators of hydride.

3. Solving the assay interference from a whole blood sample with the insertion of a solid phase capture step:

The homogenous assay described above measures the residual chemiluminescent activity in the assay mixture i.e. increasing concentrations of analyte lead to decreasing levels of chemiluminescent activity remaining in the assay mixture. Since the assay is homogenous, extraneous substances in the sample e.g. from a whole blood sample, could potentially interfere or inhibit the chemiluminescent reaction of the acridinium compound with hydrogen peroxide. In the event this is observed, it is possible to manipulate the assay so that this interference/inhibition is removed. This can be accomplished in a most direct way as illustrated in Figure 14. In this assay, a solid phase such as paramagnetic particles are introduced at the end of the homogenous assay. The particles are coated with an antibody which is specific for the acridinium compounds (Ref: U. Piran, et al., U.S. Patent No. 5,445,936, "Method for Non-Competitive Binding Assays"). Following completion of the homogenous assay, the particles are introduced to capture the acridinium compound. The particles can then be washed to remove interfering substances prior to triggering the

chemiluminescent reaction with hydrogen peroxide.

4. Heterogeneous chemiluminescent Emit[®] assays

Assays that employ a heterogeneous format can also be coupled to the present invention. Figure 15 illustrate how this may be performed. For a sandwich assay utilizing two antibodies, one of them can be coupled to the hydride generating system such as G6PDH, alcohol dehydrogenase etc. The concentration of the enzyme label in the assay is then measured by using the NADH generated by the enzyme to reduce the acridinium compound indicator molecule. Alternatively, in a competitive assay for a small analyte, the hydride generating system such as G6PDH can be coupled to the analyte for use in the assay as a tracer. Signal readout in the assay is accomplished in a similar manner as the sandwich assay.

It is also obvious that utilizing the hydride generating system such as the enzyme G6PDH as a label for nucleic acids will permit nucleic acid assays to be devised with acridinium compounds as the indicator of hydride.

Those with relevant expertise will recognize further variations which are consistent with the invention disclosed. This invention is illustrated, but not limited, by the following examples.

25

Example 1

2',6'-Dimethylcarbonylphenyl 10-methylacridinium-9-carboxylate trifluoromethanesulfonate (DMAE- ϕ).

Synthesis of 2',6'-Dimethylphenyl acridine-9-carboxylate

30 Acridine-9-carboxylic acid (5 g) was refluxed with thionyl chloride (~25 mL) under a nitrogen atmosphere in

WO 00/31543

PCT/IB99/01894

-27-

an oil-bath. Reflux was maintained till the reaction turned clear. The solution was then cooled to room temperature and poured into benzene (200 mL, anhydrous). The benzene suspension was chilled in the refrigerator overnight to complete the precipitation of the acid chloride which was isolated by filtration and rinsed with anhydrous ether. Yield = 5.3 g (quant.)

The acid chloride (5.3 g) was mixed with 2,6-dimethylphenol, dimethylaminopyridine (0.5 g) in anhydrous pyridine (40 mL). The reaction was heated in an oil-bath at 100°C for 3 hours. The reaction was then cooled to room temperature and the crude product was purified by flash chromatography on silica gel using 1:4, ethyl acetate in hexanes as eluent. Concentration of the flash fractions afforded the product as a bright yellow powder.

Synthesis of 2',6'-Dimethylphenyl 10-methylacridinium-9-carboxylate trifluoromethanesulfonate

A solution of 2',6'-dimethylphenyl acridine-9-carboxylate (20 mg, 0.061 mmol) in anhydrous dichloromethane (~2 mL) was treated with methyl trifluoromethanesulfonate (0.175 mL, 25 eq.). The reaction was stirred at room temperature. A yellow precipitate started forming in ~1 hour. The reaction was stirred for ~16 hours and then anhydrous ether (~50 mL) was added to precipitate the product which was collected by filtration and rinsed with ether and then air-dried. Yield = 29 mg, MALDI-TOF MS 342.9 obs. (342.4 calc).

Example 2

2',6'-Dimethylphenyl 10-methylacridan-9-carboxylate.

Synthesis: A solution of 2',6'-dimethylphenyl-10-methylacridinium-9-carboxylate trifluoromethanesulfonate (50 mg, 0.105 mmol) in methanol (20 mL, partial solution) was cooled in an ice-bath and treated with sodium borohydride (20 mg, 0.525 mmol). The yellow color of the solution was bleached instantly. After 1 hour additional sodium borohydride (20 mg) was added and the reaction was warmed to room temperature and stirred for ~16 hours. The reaction was then quenched with acetic acid (1 mL) and evaporated to dryness. The residue was dissolved in acetonitrile. HPLC analysis on a 3.9 x 300 mm C18 column, with a 30-min. gradient of 10-->100% MeCN/ water (0.05% TFA in each) at a flow rate of 1 mL/min. and UV-detection at 260 nm showed Rt = 27 min (acridan), Rt = 18 min (acridinium ester). The product was purified by preparative HPLC and the HPLC fractions were lyophilized to dryness to afford a white powder. Yield = 30 mg (60%).

Example 3

2',6'-Dimethylphenyl 10-(3'-sulfopropyl)-acridinium-9-carboxyamidyl-BSA Conjugate.

Synthesis: Bovine serum albumin (1.65 mg, 25 nmol) was dissolved in 475 μ L of 0.20 M NaHCO₂, pH 9.0. NSP-DMAE-NHS (US Patent 5,656,426) (0.61 mg, 1.03 μ mol) was dissolved in 103 μ L of N,N-dimethylformamide DMF to make a 10 mM solution. Twenty-five microliters of the 10 mM NSP-DMAE-NHS was mixed with the 475 μ L of BSA solution and incubated at 4°C for 16 hours. The conjugate was

isolated in water by SEC. NSP-DMAE incorporation onto BSA was approximately four labels per protein molecule as calculated from the known chemiluminescent specific activity of the label and protein determination using the
5 Bradford protein assay (6.4 μ M).

Example 4

Preparation of Chemiluminescent Hydride Indicator.

a. Nine milligrams (32 μ moles) of N(10)-methylacridinium
10 tetrafluoroborate (NMA, Sigma-Aldrich) was dissolved in 320 μ LS of N,N-dimethylformamide (DMF) to make a 0.10 M solution. A mass of 6.25 mgs (13 μ moles) of DMAE- ϕ was dissolved in 2.607 mLs of methanol, and was serially diluted in methanol to a concentration of 0.50 μ M. Three
15 hundred microliters of the 0.10 M NMA was mixed with 750 μ LS of the 0.50 μ M DMAE- ϕ into 1.95 mLs of water. Alternatively, NMA and two other hydride sinks, cetylpyridium chloride and crotonic acid, were dissolved in 0.20 M glycine buffer, 0.1 % BgG, pH 7.4, when the
20 chemiluminescent indicator of hydride used was a covalent conjugate of acridinium ester and IgG (Bayer Diagnostics). The acridinium ester-IgG conjugate was diluted to 12 nM in the same buffer.

b. Alternatively, NSP-DMAE₄-BSA conjugate was diluted to a
25 concentration of 12 nM in a buffer of 0.20 M glycine, 1.0 % (w/v) BSA, pH 7.4.

Example 5

Theophylline Assay using Acridinium Ester as a
Chemiluminescent Indicator of Hydride.

In this automated, homogeneous enzyme-immunoassay, theophylline present in a patient sample will compete with a theophylline-G6PDH conjugate for binding to a limited quantity of anti-theophylline, murine, monoclonal antibody. Binding of the anti-theophylline antibody to the theophylline-G6PDH conjugate partially inhibits enzymatic activity. G6PDH catalyzes the reduction of oxidized nicotinamide adenine dinucleotide (NAD) to the reduced form NADH by hydride transfer from glucose-6-phosphate (G6P). Hydride is then chemically transferred from the NADH to DMAE- ϕ or NSP-DMAE conjugated to BSA. Consequently, the G6PDH catalytic activity of the conjugate is correlated to the theophylline concentration in the patient sample and inversely correlated to the residual chemiluminescence of the assay reaction.

The following assay was run in entirety by the Bayer Diagnostics ACS:180[®] (Bayer Diagnostics Corp., Walpole, MA). Cuvettes were loaded into the track, then 10 μ Ls of Bayer Diagnostics ACS Theophylline Assay standards or controls were added to their respective cuvettes in replicates of three. The standards contained theophylline in concentrations of 0.00, 13.9, 27.8, 55.5, 111 and 222 μ M. The Bayer Diagnostics Ligand Controls 1, 2 and 3 contained theophylline in concentrations specified in the following table. To each cuvette, 300 μ Ls of Behring Diagnostics Emit[®] 2000 Theophylline Assay Reagent 1, containing a buffered solution of G6P, NAD, and anti-theophylline antibody, was added and incubated

WO 00/31543

PCT/IB99/01894

-31-

at 37°C for 2 minutes and 40 seconds. To this was added 150 µLs Behring Diagnostics Emit® 2000 Theophylline Assay Reagent 2, consisting of a buffered solution of theophylline-G6PDH conjugate. The cuvettes were incubated at 37°C for another 2 minutes and 40 seconds, then 20 µLs of the Chemiluminescent Indicator, containing either 10 mM NMA and 125 nM DMAE- ϕ in aqueous 10% (v/v) DMF and 25% (v/v) methanol or 12 nM NSP-DMAE₄-BSA conjugate, 0.20 M glycine, 1.0 % (w/v) BSA, pH 7.4, was added for a final incubation at 37°C for 5 minutes. Since the assay is homogeneous Washes 1 and 2, as well as, Vacuums 1, 2 and 3 were switched off. The reaction mixtures were sequentially mixed with Bayer Diagnostics ACS Reagents 1 (0.1 N HNO₃, 0.5 % (w/v) H₂O₂) and Reagent 2 (0.25 N NaOH, 0.5 % (w/v) N,N,N,N-hexadecyltrimethylammonium chloride surfactant) to initiate the chemiluminescent reaction.

Example 6

Valproate Assay using Acridinium Ester as a Chemiluminescent Indicator of Hydride.

In this automated, homogeneous enzyme-immunoassay, valproate present in a patient sample will compete with a valproate-G6PDH conjugate for binding to a limited quantity of anti-valproate, murine, monoclonal antibody. Binding of the anti-valproate antibody to the valproate-G6PDH conjugate partially inhibits enzymatic activity. G6PDH catalyzes the reduction of oxidized nicotinamide adenine dinucleotide (NAD) to the reduced form NADH by hydride transfer from glucose-6-phosphate (G6P). Hydride is then chemically transferred from the NADH to DMAE- ϕ or

WO 00/31543

PCT/IB99/01894

-32-

NSP-DMAE conjugated to BSA. Consequently, the G6PDH catalytic activity of the conjugate is correlated to the valproate concentration in the patient sample and inversely correlated to the residual chemiluminescence of the assay reaction.

The following assay was run in entirety by the Bayer Diagnostics ACS:180[®] (Bayer Diagnostics Corp., Walpole, MA). Cuvettes were loaded into the track, then 10 μ Ls of Bayer Diagnostics in-house ACS Valproate Assay standards or controls were added to their respective cuvettes in replicates of three. The standards contained valproate in concentrations of 0.000, 86.7, 173, 347, 693 and 1387 μ M. The Bayer Diagnostics in-house TDM Controls A, B and C contained valproate in concentrations specified in the following table. To each cuvette, 225 μ Ls of Behring Diagnostics Emit[®] Model 1 Valproate Assay Reagent A, containing a buffered solution of G6P, NAD, and anti-valproate antibody, was added and incubated at 37°C for 2 minutes and 40 seconds. To this was added 225 μ Ls Behring Diagnostics Emit[®] Model 1 Valproate Assay Reagent B, consisting of a buffered solution of valproate-G6PDH conjugate. The cuvettes were incubated at 37°C for another 2 minutes and 40 seconds, then 20 μ Ls of the Chemiluminescent Indicator, containing either 10 mM NMA and 125 nM DMAE- ϕ in aqueous 10% (v/v) DMF and 25% (v/v) methanol or 12 nM NSP-DMAE₄-BSA conjugate, 0.20 M glycine, 1.0 % (w/v) BSA, pH 7.4, was added for a final incubation at 37°C for 5 minutes. Since the assay is homogeneous Washes 1 and 2, as well as, Vacuums 1, 2 and 3 were switched off. The reaction mixtures were sequentially

WO 00/31543

-33-

PCT/IB99/01894

mixed with Bayer Diagnostics ACS Reagents 1 (0.1 N HNO₃, 0.5 % (w/v) H₂O₂) and Reagent 2 (0.25 N NaOH, 0.5 % (w/v) N,N,N,N-hexadecyltrimethylammonium chloride surfactant) to initiate the chemiluminescent reaction.

5 Chemiluminescence data were collected for five seconds as photons detected by ACS luminometer and expressed in relative light units (RLUs).

Example 7

10 Quinidine Assay using Acridinium Ester as a Chemiluminescent Indicator of Hydride.

In this automated, homogeneous enzyme-immunoassay, quinidine present in a patient sample will compete with a quinidine-G6PDH conjugate for binding to a limited

15 quantity of anti-quinidine, murine, monoclonal antibody. Binding of the anti-quinidine antibody to the quinidine-G6PDH conjugate partially inhibits enzymatic activity. G6PDH catalyzes the reduction of oxidized nicotinamide adenine dinucleotide (NAD) to the reduced form NADH by

20 hydride transfer from glucose-6-phosphate (G6P). Hydride is then chemically transferred from the NADH to NSP-DMAE conjugated to BSA. Consequently, the G6PDH catalytic activity of the conjugate is correlated to the quinidine concentration in the patient sample and inversely

25 correlated to the residual chemiluminescence of the assay reaction.

The following assay was run in entirety by the Bayer Diagnostics ACS:180[®] (Bayer Diagnostics Corp., Walpole, MA). Cuvettes were loaded into the track, then 10μLs of

30 Bayer Diagnostics in-house ACS Quinidine Assay standards were added to their respective cuvettes in replicates of

WO 00/31543

PCT/IB99/01894

-34-

three. The standards contained quinidine in concentrations of 0.00, 0.77, 1.5, 3.1, 6.2, 12, 18 and 25 μM . To each cuvette, 160 μLs of Bayer Corporation Emit[®] 2000 Quinidine Assay Reagent A, containing a buffered solution of G6P, NAD, and anti-quinidine antibody, was added and incubated at 37°C for 2 minutes and 40 seconds. To this was added 80 μLs Bayer Corporation Emit[®] 2000 Quinidine Assay Reagent B, consisting of a buffered solution of quinidine-G6PDH conjugate. The cuvettes were incubated at 37°C for another 2 minutes and 40 seconds, then 20 μLs of the Chemiluminescent Indicator, containing 12 nM NSP-DMAE₄-BSA conjugate, 0.20 M glycine, 1.0 % (w/v) BSA, pH 7.4, was added for a final incubation at 37°C for 5 minutes. Since the assay is homogeneous Washes 1 and 2, as well as, Vacuums 1, 2 and 3 were switched off. The reaction mixtures were sequentially mixed with Bayer Diagnostics ACS Reagents 1 (0.1 N HNO₃, 0.5 % (w/v) H₂O₂) and Reagent 2 (0.25 N NaOH, 0.5 % (w/v) N,N,N,N-hexadecyltrimethylammonium chloride surfactant) to initiate the chemiluminescent reaction. Chemiluminescence data were collected for five seconds as photons detected by ACS luminometer and expressed in relative light units (RLUs).

25

Example 8

Homogeneous, Chemiluminometric, Enzyme Assay for Ethanol using Acridinium Ester as a Chemiluminescent Indicator of Hydride.

30

In this automated, homogeneous enzyme assay, ethanol from the patient sample is oxidized to acetaldehyde through

WO 00/31543

PCT/IB99/01894

-35-

the specific catalytic action of a fixed quantity of yeast alcohol dehydrogenase (ADH) with concurrent hydride reduction of the cofactor nicotinamide adenine dinucleotide (NAD) to NADH. Equilibrium in favor of oxidation of ethanol to acetaldehyde is further promoted with the addition of a hydrazide aldehyde scavenger. Hydride is chemically transferred from the NADH to NSP-DMAE conjugated to BSA. Consequently, the ethanol concentration is correlated to NADH formation and inversely related to the residual chemiluminescence of the assay reaction.

The following assay was run in entirety by the Bayer Diagnostics ACS:180[®] (Bayer Diagnostics Corp., Walpole, MA). Cuvettes were loaded into the track, then 10 μ Ls of standards prepared by serial dilution of ethanol in serum were added to their respective cuvettes in replicates of three. The standards contained ethanol in concentrations of 0.00, 0.025, 0.050, 0.10, 0.20, and 0.40 % (w/v). To each cuvette, 150 μ Ls of Ethanol Assay Reagent A, containing 0.15 M sodium phosphate, 0.15 M hydrazinecarboxamide hydrochloride, pH 9.0, was added and incubated at 37°C for 2 minutes and 40 seconds. To this was added 150 μ Ls Ethanol Assay Reagent B, consisting of 22 mM glycine, 12 mM NAD, 14 μ M ADH, 1.0 mg/mL BSA, pH 7.0. The cuvettes were incubated at 37°C for another 2 minutes and 40 seconds, then 20 μ Ls of the Chemiluminescent Indicator, containing 12 nM NSP-DMAE₄-BSA conjugate, 0.20 M glycine, 1.0 % (w/v) BSA, pH 7.4, was added for a final incubation at 37°C for 5 minutes. Since the assay is homogeneous Washes 1 and 2, as well as, Vacuums 1, 2 and 3 were switched off. The reaction

WO 00/31543

PCT/IB99/01894

-36-

mixtures were sequentially mixed with Bayer Diagnostics ACS Reagents 1 (0.1 N HNO₃, 0.5 % (w/v) H₂O₂) and Reagent 2 (0.25 N NaOH, 0.5 % (w/v) N,N,N,N-hexadecyltrimethylammonium chloride surfactant) to initiate the chemiluminescent reaction. Chemiluminescence data were collected for five seconds as photons detected by ACS luminometer and expressed in relative light units (RLUs).

Method of Calculation for Assay Parameters. Arithmetic means for RLUs resulting from a specific analyte concentration, represented here as μ , were calculated from three replicates. An inverse, non-linear relationship exists between the analyte concentration present in the standard and the detected RLUs,

$$y = y_0 + \frac{(y_{\infty} - y_0)}{(1 + 10^{-m \log x - b})}$$

where x is the analyte concentration, and y is the observed signal generated as RLUs (Rodbard, David; Ligand Analysis; (1981); Langon, J.; Clapp, J. (Eds.); Masson Publishing, Inc., New York; pp. 45 - 101) (Nix, Barry; The Immunoassay Handbook; (1994); Wild, David (Ed.); Stockton Press, Inc., New York; pp. 117 - 123) (Van Lente, Frederick, Galen, Robert S.; Enzyme-immunoassay; (1980); Maggio, Edward T. (Ed.); CRC Press, Inc., Boca Raton; pp. 135 - 153).

Additionally, there are four parameters, namely the regression constant, b , the regression coefficient, m , the asymptotic limit at infinite dose, y_{∞} , and the asymptotic limit for the zero dose, y_0 . The latter three of these parameters were calculated directly using the

WO 00/31543

PCT/IB99/01894

-37-

iterative, standard, four-parameter logistic (4PL-STD) analysis function of the DOSECALC.EXE Rev.1.73 program (Bayer Diagnostics Corp., Walpole, MA). The arithmetic mean of the regression constant b was determined over the
 5 entire range of analyte concentrations as calculated from the dose response expression re-written as

$$b = -\log \frac{y_{\infty} - y}{y - y_0} - m \log x.$$

10 Analyte concentrations of unknowns were subsequently calculated using the dose response equation arranged as

$$x = 10^{\frac{\log[(y_{\infty} - y)/(y - y_0)] + b}{-m}}.$$

Theophylline, Quinidine and Valproate Assay Standard
 15 Curves Using Acridinium Ester as a Chemiluminescent Indicator of Hydride. A plot of the RLUs versus analyte concentration illustrates the inverse, non-linear relationship between chemiluminescence and analyte concentration. Precision was good, being under 5 % C.V.
 20 for all points.

Clearly, acridinium compounds will function as chemiluminescent indicators of hydride either as free indicators dissolved in organic solvent or as hydrophilic
 25 conjugates when covalently linked to protein.

Assay Accuracy in Determination of Theophylline and Valproate Concentrations. Analyte concentrations were determined for the Ligand Controls using the standard 4PL

WO 00/31543

PCT/IB99/01894

-38-

function. The determined values were compared to the established ranges stated in the associated product literature, as shown in Figure 21.

5 Variation in specified analyte ranges between different assay constructs is considerable for a number of multi-level ligand controls because of the different susceptibility to the matrix effect. Since acceptable ranges of analyte concentrations have not yet been
10 established for the new assay format described in the present invention, some difference is expected. Determined values are similar enough to referenced control limits that we can demonstrate the utility of an acridinium compound as a chemiluminescent indicator of
15 hydride for the quantitation of theophylline and valproate in a homogeneous, enzyme-immunoassay.

Example 9

Ethanol Assay Standard Curves Using Acridinium Ester as a
20 Chemiluminescent Indicator of Hydride.

A plot of the RLUs versus analyte concentration (see Figure 22) illustrates the inverse, non-linear relationship between chemiluminescence and analyte concentration. As with the chemiluminometric Emit[®] assays
25 precision in the ethanol assay was good being under 5 % C.V. for all points.

WO 00/31543

PCT/IB99/01894

-39-

CLAIMS

We claim:

- 5 1. A chemiluminescent assay method to detect or
quantitate hydride, said assay comprising measuring the
amount of chemiluminescence generated from a
chemiluminescent compound in a reaction in which said
chemiluminescent compound reacts with a hydride
10 generated from a chemical, biochemical or enzyme-
catalyzed reaction.
2. The method of claim 1 in which said chemiluminescent
compound is an acridinium, benzacridinium,
15 phenanthridinium, quinolinium or lucigenin compound, or
the conjugate of said acridinium, benzacridinium,
phenanthridinium, quinolinium or lucigenin compound.
3. The method of claim 1 in which said hydride reacts
20 with the nucleus of said chemiluminescent compound.
4. The method of claim 1 in which said hydride reacts
with the conjugate of said chemiluminescent compound.
- 25 5. The method of claim 1 in which said hydride reacts
with the ester or sulfonylamide derivative of said
chemiluminescent compound.
6. The method of claim 1 in which said hydride reacts
30 with said chemiluminescent compound containing (a) an
extended electronic conjugation system, (b) a hydride-

WO 00/31543

PCT/IB99/01894

-40-

reducible quencher, (c) one or more electron donating groups, or (d) a fluorescent resonance energy acceptor.

5 7. A method for determining the amount of hydride produced in a reaction comprising (a) adding a known amount of a chemiluminescent compound to said reaction mixture, (b) measuring the amount of chemiluminescence given off from said reaction mixture after the addition of reagents which release said chemiluminescence, (c)
10 comparing the amount of chemiluminescence produced with the expected chemiluminescence in the absence of hydride, and (d) calculating the amount of said hydride present by comparison with a standard curve.

15 8. The method of claim 7 wherein said hydride is generated in an assay method for an analyte.

9. The method of claim 7 in which said chemiluminescent compound is an acridinium, benzacridinium,
20 phenanthridinium, quinolinium or lucigenin compound, or the conjugate of said acridinium, benzacridinium, phenanthridinium, quinolinium or lucigenin compound.

10. The method of claim 9 in which said acridinium or
25 benzacridinium compound is an acridinium ester or benzacridinium ester, acridinium sulfonylamide or benzacridinium sulfonylamide.

11. The method of claim 7 in which said hydride is
30 produced in a biochemical redox reaction of an enzyme cofactor.

WO 00/31543

PCT/IB99/01894

-41-

12. The method of claim 11 in which said cofactor is NAD⁺, NADP⁺, FMN or FAD.

13. A method for determining the amount of an analyte in a sample by determining the amount of hydride produced in a reaction comprising (a) adding a known amount of a chemiluminescent compound to said reaction mixture, (b) measuring the amount of chemiluminescence given off from said reaction mixture after the addition of reagents which release said chemiluminescence, (c) comparing the amount of chemiluminescence produced with the expected chemiluminescence in the absence of hydride, and (d) calculating the amount of said analyte present by comparison with a standard curve.

15

14. The method of claim 13 in which said analyte is theophylline, valproate, quinidine or ethanol and said sample is serum.

20

15. A method for chemiluminescent determination of analyte in a sample comprising (a) adding sequentially to said sample the reagents from a colorimetric assay for hydride and a chemiluminescent indicator of hydride, (b) measuring the chemiluminescence generated and (c) comparing said luminescence generated with a calibration curve to determine the amount of analyte in said sample.

25

16. A kit comprising (a) the reagents for a colorimetric assay for hydride and (b) a chemiluminescent indicator of hydride.

30


WO 00/31543

PCT/IB99/01894

-42-

17. The method of claim 15 in which said chemiluminescence generated is measured on an automated chemiluminescence analyzer or diagnostic system.

18. A chemiluminescent hydride indicator having the structure



wherein

R_1 , is an alkyl, alkenyl, alkynyl or aralkyl containing optionally up to 20 heteroatoms; preferably R_1 is methyl or sulfoalkyl group;

R_2 , R_2' , and R_3 , are identical or different, selected from hydrogen, -R, substituted or unsubstituted aryl (ArR or Ar), halide, amino, hydroxyl, nitro, sulfonate, -CN, -COOH, -SCN, -OR, -SR, -SSR, -C(O)R, -C(O)OR, -C(O)NHR, or -NHC(O)R;

R is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, and aralkyl containing optionally up to 20 heteroatoms;

WO 00/31543

PCT/IB99/01894

-43-

alternatively, R_2 and R_3 can be linked so as to form an additional ring fused to the attached acridinium nucleus;

5 C_1 , C_2 , C_3 , C_4 , C_5 , and C_8 , peri-positions of the acridinium nucleus are optionally substituted as represented by R_2' ;

10 A^- is a counter ion which is introduced to pair with the quaternary nitrogen of the acridinium nucleus either as a result of quaternizing the acridine ring nitrogen by the use of alkylating agents during the synthesis, modification of the R_1 , or subsequent exchange that occurs during the work-up of reaction mixtures and purification of desired compounds in a solution or fluid containing excess amount of other anions;

15 X is nitrogen, oxygen or sulfur; such that, when X is oxygen or sulfur, Z is omitted and Y is a polysubstituted aryl moiety;

R_5 and R_7 are any of R_2 , R_2' , and R_3 defined above;

20 R_6 is also any of R_2 , R_2' , and R_3 defined above, when the acridinium compound is used as a free chemiluminescent indicator.

25 said acridinium compounds being optionally covalently attached to a more water-soluble polymer (natural or synthetic) or biopolymers (e.g. proteins, polysaccharides, glycoproteins, and nucleic acids) in a conjugate form for enhanced water solubility;

30 R_5 and R_6 , and R_6 and R_7 are interchangeable; such that when X is nitrogen, then Z is $-SO_2-Y'$, Y' being defined as Y above, and both can be the same or different, with Y optionally being a branched or straight-chained alkyl

WO 00/31543


PCT/IB99/01894

-44-

containing up to 20 carbon atoms, or a substituted aryl, or heterocyclic ring system.

19. The chemiluminescent hydride indicator of claim 18 in which said counter ion is selected from the group consisting of CH_3SO_4^- , FSO_3^- , CF_3SO_3^- , $\text{C}_4\text{F}_9\text{SO}_3^-$, $\text{CH}_3\text{C}_6\text{H}_4\text{SO}_3^-$, halide, CF_3COO^- , CH_3COO^- , and NO_3^- ;

said polysubstituted aryl moiety is



10


where R_4 and R_8 can be (1) hydrogen or (2) alkyl, alkenyl, alkynyl, alkoxyl ($-\text{OR}$), alkylthiol ($-\text{SR}$), or substituted amino groups that serve to stabilize the $-\text{COX}-$ linkage between the acridinium nucleus and the Y moiety, through steric and/or electronic effects, preferably, one of them is a hydrogen, if the C_1 or C_8 position of the acridinium nucleus is substituted with a lower alkyl group, preferably methyl; more preferably, R_4 and R_8 are alkyl, alkenyl, alkynyl, alkoxyl ($-\text{OR}$), alkylthiol ($-\text{SR}$), or substituted amino groups that serve to stabilize the $-\text{COX}-$ linkage between the acridinium nucleus and the Y moiety, through steric and/or electronic effect; most preferably R_4 and R_8 are lower alkyl (e.g., a methyl group);

WO 00/31543

PCT/IB99/01894

-45-

5 R_6 can also be $-R_9-R_{10}$, where R_9 is not required but optionally can be branched or straight-chained alkyl, substituted or unsubstituted aryl or aralkyl containing optionally up to 20 heteroatoms, and R_{10} is a leaving group or an electrophilic functional group attached with a leaving group having the structure



a halide, $-COOH$

10 R_{10} can also be $-Q-R-Nu$, $-Q-R-(I)_nNu-$, $-Q-Nu$, $-R-Nu$, or $-Nu$, where n is a number of at least 1, Nu is a nucleophilic group, Q is a functional linkage, I is an ionic or ionizable group.


WO 00/31543

PCT/IB99/01894

-46-

20. The chemiluminescent hydride indicator of claim 18, in which said indicator is a phenanthridium or quinolinium compound having the structure

5



with substituents being the same as in claim 18, except that the peri-positions for R₂' substituent are re-designated at the C₁, C₂, C₃, C₄, C₆, and C₉ positions of said phenanthridinium nucleus and C₂, C₃, C₅, and C₈ positions of said quinolinium nucleus.

21. The method of claim 13 for elimination of interference resulting from the presence of whole blood comprising, before the measurement of chemiluminescence, (a) the addition of a solid phase coated with an antibody specific for an acridinium compound, (b) incubation of the reaction mixture to allow capture of said acridinium compound, (c) separating said solid phase and washing said separated solid phase to remove said interfering substance.

WO 00/31543

PCT/IB99/01894

-47-

22. The method of claim 15 for elimination of interference resulting from the presence of whole blood comprising, before the measurement of chemiluminescence, (a) the addition of a solid phase coated with an antibody specific for an acridinium compound, (b) incubation of the reaction mixture to allow capture of said acridinium compound, (c) separating said solid phase and washing said separated solid phase to remove said interfering substance.

10

WO 00/31543

1/11

PCT/IB99/01894




Figure 1

NADH eq.	% Residual Chemiluminescent Activity
0	100
0.1	94
0.2	85
0.4	46
0.6	38
0.8	30
1.0	17

Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

WO 00/31543

4/11

PCT/IB99/01894




Figure 7




Figure 8

WO 00/31543

5/11

PCT/IB99/01894




Figure 9




Figure 10

WO 00/31543

6/11

PCT/IB99/01894




Figure 11




Figure 12

Homogeneous Immunoassay Using Acridinium Compound as Hydride Indicator





Figure 13



AC = acridinium compound; ACH = reduced acridinium compound

Figure 14




Figure 15

Chemiluminometric *Emit*[®] Theophylline Assay using NSP-DMAE₄-BSA as a Hydride Indicator




Figure 16

9/11

WO 00/31543

PCT/IB99/01894

Chemiluminometric *Emit*[®] Valproate Assay using NSP-DMAE₄-BSA as a Hydride Indicator




Figure 17

Chemiluminometric *Emit*[®] Quinidine Assay using NSP-DMAE₄-BSA as a Hydride Indicator




Figure 18

**Chemiluminometric *Emit*[®] Theophylline Assay
Using DMAE- ϕ as a Hydride Indicator**




Figure 19

**Chemiluminometric *Emit*[®] Valproate Assay
using DMAE- ϕ as a Hydride Indicator**




Figure 20

WO 00/31543

11/11

PCT/IB99/01894

Referenced vs. Determined Analyte Concentration for Controls						
	Theophylline Concentration (μM)			Valproate Concentration (μM)		
	Controls					
Referenced	1	2	3	A	B	C
Ciba-Corning ACS:180®	21-33	60-86	119-189	104-173	381-451	589-659
Syva Emit 2000®	27-41	75-111	159-237	-	-	-
Determined	15.4	42.3	133	116	337	567

Figure 21

Chemiluminometric Ethanol Assay





Figure 22



Chemiluminescent Acridinium Ester

Nonchemiluminescent Acridan

