



Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2732320 C 2017/08/29

(11)(21) **2 732 320**

(12) **BREVET CANADIEN**  
**CANADIAN PATENT**

(13) **C**

(86) **Date de dépôt PCT/PCT Filing Date:** 2009/07/31  
(87) **Date publication PCT/PCT Publication Date:** 2010/02/04  
(45) **Date de délivrance/Issue Date:** 2017/08/29  
(85) **Entrée phase nationale/National Entry:** 2011/01/27  
(86) **N° demande PCT/PCT Application No.:** US 2009/052439  
(87) **N° publication PCT/PCT Publication No.:** 2010/014921  
(30) **Priorité/Priority:** 2008/08/01 (US61/085,588)


(51) **Cl.Int./Int.Cl. C07D 473/34 (2006.01),**  
**A61K 31/52 (2006.01)**

(72) **Inventeurs/Inventors:**  
JACOBSON, KENNETH A., US;  
MELMAN, ARTEM, US;  
WANG, BEN BIH-REN, US

(73) **Propriétaire/Owner:**  
THE UNITED STATES OF AMERICA, AS  
REPRESENTED BY THE SECRETARY,  
DEPARTMENT OF HEALTH AND HUMAN  
SERVICES, US

(74) **Agent:** RIDOUT & MAYBEE LLP

(54) **Titre : ANTAGONISTES ET AGONISTES PARTIELS DES RECEPTEURS DE L'ADENOSINE A<sub>3</sub> ET AGONISTES PARTIELS**  
(54) **Title: A<sub>3</sub> ADENOSINE RECEPTOR ANTAGONISTS AND PARTIAL AGONISTS**



(I)

(57) **Abrégé/Abstract:**

Disclosed are A<sub>3</sub> adenosine receptor antagonists and/or partial agonists of formula (I): wherein R<sup>1</sup> to R<sup>5</sup> are as described herein, as well as pharmaceutical compositions thereof and methods of use thereof. The antagonists or partial agonists find use in treating a

**(57) Abrégé(suite)/Abstract(continued):**

number of diseases including cancer, glaucoma, inflammatory diseases, asthma, stroke, myocardial infarction, allergic reactions, rhinitis, poison ivy induced responses, urticaria, scleroderma, arthritis, brain arteriole diameter constriction, bronchoconstriction, and myocardial ischemia, as well as in preventing cardiac ischemia. Also disclosed are radiolabeled compounds of formula (I) and the use thereof in diagnostic imaging of tissues and organs.

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
4 February 2010 (04.02.2010)(10) International Publication Number  
WO 2010/014921 A3(51) International Patent Classification:  
*C07D 473/34* (2006.01)    *A61K 31/52* (2006.01)(74) Agents: **PILLAI, Xavier** et al.; Two Prudential Plaza, Suite 4900, 180 N. Stetson Avenue, Chicago, IL 60601 (US).

(21) International Application Number:

PCT/US2009/052439

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:

31 July 2009 (31.07.2009)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/085,588    1 August 2008 (01.08.2008)    US

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(71) Applicant (for all designated States except US): **THE UNITED STATES OF AMERICA, AS REPRESENTED BY THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]**; Office Of Technology Transfer, National Institutes Of Health, 6011 Executive Boulevard, Suite 325, MSC 7660, Bethesda, MD 20892-7660 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(72) Inventors; and


(75) Inventors/Applicants (for US only): **JACOBSON, Kenneth, A.** [US/US]; 11218 Watermill Lane, Silver Spring, MD 20902-3438 (US). **MELMAN, Artem** [IL/US]; 469 County Route 59, Potsdam, NY 13676 (US). **WANG, Ben, Bih-Ren** [US/US]; 14630 Bauer Drive, #3, Rockville, MD 20853 (US).

(81) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

[Continued on next page]

(54) Title: A<sub>3</sub> ADENOSINE RECEPTOR ANTAGONISTS AND PARTIAL AGONISTS

(57) Abstract: Disclosed are A<sub>3</sub> adenosine receptor antagonists and/or partial agonists of formula (I): wherein R<sup>1</sup> to R<sup>5</sup> are as described herein, as well as pharmaceutical compositions thereof and methods of use thereof. The antagonists or partial agonists find use in treating a number of diseases including cancer, glaucoma, inflammatory diseases, asthma, stroke, myocardial infarction, allergic reactions, rhinitis, poison ivy induced responses, urticaria, scleroderma, arthritis, brain arteriole diameter constriction, bronchoconstriction, and myocardial ischemia, as well as in preventing cardiac ischemia. Also disclosed are radiolabeled compounds of formula (I) and the use thereof in diagnostic imaging of tissues and organs.

**WO 2010/014921 A3**

**(88) Date of publication of the international search report:**

16 September 2010

- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

## A<sub>3</sub> ADENOSINE RECEPTOR ANTAGONISTS AND PARTIAL AGONISTS

### CROSS-REFERENCE TO A RELATED APPLICATION

[0001] This application claims the benefit of United States Provisional Patent Application No. 61/085,588, filed August 1, 2008.

### BACKGROUND OF THE INVENTION

[0002] There are four subtypes of receptors for adenosine, designated A<sub>1</sub>, A<sub>2A</sub>, A<sub>2B</sub>, and A<sub>3</sub>. The A<sub>3</sub> adenosine receptor is found primarily in the central nervous system, brain, testes, and the immune system, where it appears to be involved in the modulation of release from mast cells of mediators of the immediate hypersensitivity reaction (Ramkumar et al., *J. Biol. Chem.*, 268, 16887-16890 (1993)).

[0003] It is believed that A<sub>3</sub> adenosine receptor selective antagonists should serve as cerebroprotective, antiasthmatic, or anti-inflammatory agents. It is also believed that A<sub>3</sub> adenosine receptor selective antagonists should serve in the treatment of glaucoma, for example, in reducing intraocular pressure. Research activity is evident in the area of A<sub>3</sub> adenosine receptor antagonists; see, for example, U.S. Patents 6,066,642 and 6,528,516 and WO 2008/055711. Accordingly, there is a desire to find new A<sub>3</sub> adenosine receptor antagonists.

[0004] Further, A<sub>3</sub> adenosine receptor partial agonists, are advantageous in cardioprotection and produce anti-ischemic effects. Partial agonists also tend to have less side effects than full agonists. In addition, partial agonists are less likely to produce desensitization of the receptor as compared to full agonists. Accordingly, partial agonists can activate the receptor for a longer duration and achieve longer lasting response. Accordingly, there is a desire to find new A<sub>3</sub> adenosine receptor partial agonists.

### BRIEF SUMMARY OF THE INVENTION

[0005] The invention provides compounds, pharmaceutical compositions, and methods of use of the compounds. The compounds of the invention are antagonists, or partial agonists, of the A<sub>3</sub> adenosine receptor and are purine analogs having substituents at the N<sup>6</sup>-, 2-, and 9-, and optionally at the 8-position, of the purine core. The compounds have a constrained ring or a rigid bicyclo[3.1.0] hexane ring at the 9-position of the purine core, which provides high potency and selectivity to the A<sub>3</sub> adenosine receptor and at the same time lack a substituent on the 4'-position of the bicyclo hexane ring. The absence of a 4'-substituent in many of the

compounds leads to lack of activation of the A<sub>3</sub> adenosine receptor. Many of the compounds act as partial agonists.

#### BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

[0006] Figure 1 depicts a reaction scheme to prepare compounds **7b-13b** in accordance with an embodiment of the invention. a) 7 steps (see Joshi et al. *Nucleosides, Nucleotides, and Nucleic Acids* **2008**, 27, 279 and Joshi et al. *J. Org. Chem.* **2005**, 70, 439); b) TBDPS-Cl, imidazole, DMF; c) NaOH, H<sub>2</sub>O, MeOH, reflux; d) 2-mercaptopurine N-oxide, DCC, toluene; e) (Me<sub>3</sub>Si)<sub>3</sub>SiH, AIBN, toluene; f) Bu<sub>4</sub>NF, THF; g) 2,6-dichloropurine, PPh<sub>3</sub>, DIAD, THF; h) RNH<sub>2</sub>, EtOH; i) TFA/H<sub>2</sub>O/MeOH.

[0007] Figure 2 depicts a reaction scheme to prepare compounds **19a-19g** in accordance with an embodiment of the invention.

[0008] Figure 3 depicts compounds **22a-22g** in accordance with an embodiment of the invention and reaction schemes to prepare compounds **20a-20g** and **21a-21g**.

[0009] Figure 4 depicts functional antagonism by the compound **7b** of the invention in the guanine nucleotide binding assay ([<sup>35</sup>S]GTPγS) in membranes of CHO cells expressing human A<sub>3</sub>AR.

[0010] Figure 5 depicts functional agonism of compounds **7b** and **9b** in accordance with an embodiment of the invention in an assay of adenylate cyclase membranes of CHO cells expressing hA<sub>3</sub>AR. The full agonist NECA (5'-N-ethylcarboxamidoadenosine), representing 100% efficiency, is shown comparison. Figure 6 depicts a (radio)iodination of compound **7b** on its N<sup>6</sup>-3-iodobenzyl substituent by iododestannylation of a 3-(trimethylstannyl)benzyl precursor.


[0011] Figure 7A depicts the non-specific, specific, and total binding of [<sup>125</sup>I] **7b** on mouse A<sub>3</sub> adenosine receptor. Figure 7B depicts the extent of specific binding as a function of the concentration of the compound.

[0012] Figure 8 depicts the biodistribution of Br-76 labeled compound **9b** at 15, 60, and 120 min post injection in rats. The Y-axis represents %Initial Dose per gram and X-axis shows various organs.

#### DETAILED DESCRIPTION OF THE INVENTION

[0013] The present invention is predicated on the concept that compounds having a ring constrained substituent or a rigid bicyclo[3.1.0] hexane ring at the 9-position which provides high potency as an antagonist and selectivity to the A<sub>3</sub> adenosine receptor, or as a partial agonist of the A<sub>3</sub> adenosine receptor, and at the same time lack a substituent on the 4'-position of the bicyclic hexane ring.

[0014] Accordingly, the present invention provides a compound of Formula I:



wherein

[0015]  $R^1$  is selected from the group consisting of hydrogen,  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkoxy, hydroxyl,  $C_3$ - $C_8$  cycloalkyl,  $C_3$ - $C_8$  cycloalkyl  $C_1$ - $C_6$  alkyl,  $C_3$ - $C_8$  dicycloalkyl  $C_1$ - $C_6$  alkyl,  $C_7$ - $C_{12}$  bicycloalkyl  $C_1$ - $C_6$  alkyl,  $C_7$ - $C_{14}$  tricycloalkyl  $C_1$ - $C_6$  alkyl,  $C_6$ - $C_{14}$  aryl,  $C_6$ - $C_{14}$  aryl  $C_1$ - $C_6$  alkyl,  $C_6$ - $C_{14}$  diaryl  $C_1$ - $C_6$  alkyl,  $C_6$ - $C_{14}$  aryl  $C_1$ - $C_6$  alkoxy,  $C_1$ - $C_6$  alkyl carbonyl, sulfonyl,  $C_1$ - $C_6$  alkyl sulfonyl,  $C_6$ - $C_{14}$  aryl sulfonyl, heterocyclyl  $C_1$ - $C_6$  alkyl, heterocyclyl, heteroaryl  $C_1$ - $C_6$  alkyl, 4-[[4-[[[(2-amino  $C_1$ - $C_6$  alkyl) amino]-carbonyl]-  $C_1$ - $C_6$  alkyl] aniline] carbonyl]  $C_1$ - $C_6$  alkyl]  $C_6$ - $C_{14}$  aryl, and  $C_6$ - $C_{14}$  aryl  $C_3$ - $C_8$  cycloalkyl, wherein the aryl or heterocyclyl portion of  $R^1$  is optionally substituted with one or more substituents selected from the group consisting of halo, amino, hydroxyl, carboxy,  $C_1$ - $C_6$  alkoxy carbonyl, aminocarbonyl,  $C_1$ - $C_6$  alkylaminocarbonyl,  $C_1$ - $C_6$  dialkyl aminocarbonyl,  $C_1$ - $C_6$  alkyl,  $C_2$ - $C_6$  alkenyl,  $C_2$ - $C_6$  alkynyl,  $C_1$ - $C_6$  alkoxy,  $C_6$ - $C_{14}$  aryloxy, hydroxy  $C_1$ - $C_6$  alkyl, hydroxy  $C_2$ - $C_6$  alkenyl, hydroxy  $C_2$ - $C_6$  alkynyl, carboxy  $C_1$ - $C_6$  alkyl, carboxy  $C_2$ - $C_6$  alkenyl, carboxy  $C_2$ - $C_6$  alkynyl, aminocarbonyl  $C_1$ - $C_6$  alkyl, aminocarbonyl  $C_2$ - $C_6$  alkenyl, aminocarbonyl  $C_2$ - $C_6$  alkynyl, and  $C\equiv C$ -( $CH_2$ )<sub>n</sub>-COR<sup>7</sup> wherein R<sup>7</sup> is selected from the group consisting of OH, OR<sup>8</sup>, and NR<sup>9</sup>R<sup>10</sup>, wherein R<sup>8</sup> is selected from the group consisting of  $C_1$ - $C_6$  alkyl,  $C_3$ - $C_8$  cycloalkyl,  $C_3$ - $C_8$  cycloalkyl  $C_1$ - $C_6$  alkyl,  $C_3$ - $C_8$  dicycloalkyl  $C_1$ - $C_6$  alkyl,  $C_7$ - $C_{12}$  bicycloalkyl  $C_1$ - $C_6$  alkyl,  $C_7$ - $C_{14}$  tricycloalkyl  $C_1$ - $C_6$  alkyl,  $C_6$ - $C_{14}$  aryl,  $C_6$ - $C_{14}$  aryl  $C_1$ - $C_6$  alkyl,  $C_6$ - $C_{14}$  and diaryl  $C_1$ - $C_6$  alkyl; and R<sup>9</sup> and R<sup>10</sup> are independently selected from the group consisting of hydrogen,  $C_1$ - $C_6$  alkyl, and ( $CH_2$ )<sub>n</sub>R<sup>11</sup> wherein R<sup>11</sup> is NR<sup>12</sup>R<sup>13</sup>, wherein R<sup>12</sup> and R<sup>13</sup> are independently selected from the group consisting of hydrogen,  $C_1$ - $C_6$  alkyl, and COR<sup>14</sup> wherein R<sup>14</sup> is hydrogen or  $C_1$ - $C_6$  alkyl; wherein n is an integer from 1 to 10; and the alkyl or

cycloalkyl portion of R<sup>1</sup> is optionally substituted with one or more substituents selected from the group consisting of halo, amino, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>6</sub>-C<sub>14</sub> aryloxy, C<sub>1</sub>-C<sub>6</sub> hydroxyalkyl, C<sub>2</sub>-C<sub>6</sub> hydroxyalkenyl, C<sub>2</sub>-C<sub>6</sub> hydroxy alkynyl, aminocarbonyl C<sub>1</sub>-C<sub>6</sub> alkoxy, and C<sub>6</sub>-C<sub>14</sub> aryl C<sub>1</sub>-C<sub>6</sub> alkoxy;

[0016] R<sup>2</sup> is selected from the group consisting of hydrogen, halo, amino, hydrazido, mercapto, C<sub>1</sub>-C<sub>20</sub> alkylamino, C<sub>6</sub>-C<sub>14</sub> aryl amino, C<sub>6</sub>-C<sub>14</sub> aryloxy, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>1</sub>-C<sub>20</sub> alkoxy, C<sub>1</sub>-C<sub>20</sub> thioalkoxy, pyridylthio, C<sub>7</sub>-C<sub>12</sub> cycloalkyl C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkyl C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkenyl C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>6</sub>-C<sub>14</sub> aryl C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>7</sub>-C<sub>12</sub> cycloalkyl C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkyl C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>6</sub>-C<sub>14</sub> aryl C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>2</sub>-C<sub>20</sub> alkynyl, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-C(=O)-O-C<sub>1</sub>-C<sub>6</sub> alkyl, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-C(=O)-NH-(CH<sub>2</sub>)<sub>n</sub>-NH<sub>2</sub>, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-C<sub>1</sub>-C<sub>6</sub> alkyl, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-aryl, wherein m and n are independently 1 to 10, C<sub>7</sub>-C<sub>12</sub> cycloalkyl C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkyl C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkenyl C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>6</sub>-C<sub>14</sub> aryl C<sub>2</sub>-C<sub>20</sub> alkynyl, and the alkyl, cycloalkyl, or aryl portion of R<sup>2</sup> is optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, amino, alkylamino, dialkylamino, sulfur, carboxy, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkyl aminocarbonyl, aminoalkyl aminocarbonyl, and trialkylsilyl;

[0017] R<sup>3</sup> and R<sup>4</sup> are independently selected from the group consisting of hydroxyl, amino, thiol, ureido, C<sub>1</sub>-C<sub>6</sub> alkyl carbonylamino, hydroxy C<sub>1</sub>-C<sub>6</sub> alkyl, and hydrazinyl; and

[0018] R<sup>5</sup> is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, heteroaryl, and C<sub>1</sub>-C<sub>6</sub> aminoalkyl;

[0019] or a pharmaceutically acceptable salt thereof.

[0020] The term "aryl" refers to aromatic moieties such as phenyl, naphthyl, anthracenyl, and biphenyl. The term "heterocyclyl" refers to 3-7 membered rings which can be saturated or unsaturated or heteroaromatic, comprising carbon and one or more heteroatoms such as O, N, and S, and optionally hydrogen; optionally in combination with one or more aromatic rings. Examples of heterocyclyl groups include pyridyl, piperidinyl, piperazinyl, pyrazinyl, pyrrolyl, pyranyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrrolidinyl, furanyl, tetrahydrofuranyl, thienyl, furyl, thiophenyl, tetrahydrothiophenyl, purinyl, pyrimidinyl, thiazolyl, thiazolidinyl, thiazolinyl, oxazolyl, tetrazolyl, tetrazinyl, benzoxazolyl, morpholinyl, thiomorpholinyl, quinolinyl, and isoquinolinyl. Examples of heteroaryl alkyl include heteroaryl methyl such as 2- or 3- methyl substituted groups, e.g., thienylmethyl, pyridylmethyl, and furylmethyl.

[0021] The alkyl, alkoxy, and alkylamino groups can be linear or branched. When an aryl group is substituted with a substituent, e.g., halo, amino, alkyl, hydroxyl, alkoxy, and others, the aromatic ring hydrogen is replaced with the substituent and this can take place in any of the available hydrogens, e.g., 2, 3, 4, 5, and/or 6-position wherein the 1-position is the point of attachment of the aryl group in the compound of the present invention.

[0022] The term "halo" refers to fluorine, chlorine, bromine, and iodine.

[0023] Examples of bicycloalkyls include norbornyl, s-endornorbornyl, carbamethylcyclopentyl, and bicyclohexyl. An example of a tricycloalkyl is adamantyl.

[0024] The phrase "salt" or "pharmaceutically acceptable salt" is intended to include nontoxic salts synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two. Generally, nonaqueous media such as ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing Company, Easton, PA, 1990, p. 1445, and *Journal of Pharmaceutical Science*, 66, 2-19 (1977). For example, they can be a salt of an alkali metal (e.g., sodium or potassium), alkaline earth metal (e.g., calcium), or ammonium of salt.

[0025] Examples of pharmaceutically acceptable salts for use in the present inventive pharmaceutical composition include those derived from mineral acids, such as hydrochloric, hydrobromic, phosphoric, metaphosphoric, nitric and sulphuric acids, and organic acids, such as tartaric, acetic, citric, malic, lactic, fumaric, benzoic, glycolic, gluconic, succinic, maleic and arylsulfonic, for example, benzenesulfonic and p-toluenesulfonic, acids.

[0026] In accordance with an embodiment of the invention, R<sup>1</sup> is selected from the group consisting of C<sub>6</sub>-C<sub>14</sub> aryl C<sub>1</sub>-C<sub>6</sub> alkyl and C<sub>6</sub>-C<sub>14</sub> aryl C<sub>3</sub>-C<sub>8</sub> cycloalkyl, wherein the aryl portion of R<sup>1</sup> is optionally substituted with one or more substituents selected from the group consisting of halo, amino, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>6</sub>-C<sub>14</sub> aryloxy, hydroxy C<sub>1</sub>-C<sub>6</sub> alkyl, hydroxy C<sub>2</sub>-C<sub>6</sub> alkenyl, hydroxy C<sub>2</sub>-C<sub>6</sub> alkynyl, aminocarbonyl C<sub>1</sub>-C<sub>6</sub> alkoxy, and C<sub>6</sub>-C<sub>14</sub> aryl C<sub>1</sub>-C<sub>6</sub> alkoxy; and in a particular embodiment, R<sup>1</sup> is selected from the group consisting of benzyl, phenyl cyclopropyl, or 1-naphthyl methyl, wherein the phenyl or naphthyl portion of R<sup>1</sup> is optionally substituted with one or more substituents selected from the group consisting of halo, amino, hydroxyl, carboxy, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl,

dialkyl aminocarbonyl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, phenoxy, hydroxy C<sub>1</sub>-C<sub>6</sub> alkyl, hydroxy C<sub>2</sub>-C<sub>6</sub> alkenyl, and hydroxy C<sub>2</sub>-C<sub>6</sub> alkynyl.

[0027] In a specific embodiment of the invention, R<sup>1</sup> is benzyl, phenyl cyclopropyl, or 1-naphthyl methyl, wherein the phenyl or naphthyl portion of R<sup>1</sup> is optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, and alkoxy. Examples of R<sup>1</sup> are benzyl and benzyl substituted with one or more substituents selected from the group consisting of halo and C<sub>1</sub>-C<sub>6</sub> alkoxy.

[0028] In any of the embodiments above, R<sup>1</sup> is selected from the group consisting of 3-chlorobenzyl, 3-bromobenzyl, 3-iodobenzyl, 2-hydroxy-5-methoxy-benzyl, and 2,5-dimethoxybenzyl. In an embodiment, the phenyl cyclopropyl is *trans*-2-phenyl-1-cyclopropyl.

[0029] In any of the embodiments above, R<sup>2</sup> is halo, specifically chloro, bromo, or iodo, or R<sup>2</sup> is -C≡C-(CH<sub>2</sub>)<sub>m</sub>-CH<sub>3</sub>, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-aryl, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-C(=O)-O-CH<sub>3</sub>, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-C(=O)-NH-(CH<sub>2</sub>)<sub>n</sub>-NH<sub>2</sub>, wherein m and n are independently 1 to 10, where in certain embodiments m and n are 2 to 6, and in certain other embodiments m and n are 3 to 5, and wherein the CH<sub>3</sub> or aryl group is optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, amino, alkylamino, dialkylamino, sulfur, carboxy, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkyl aminocarbonyl, aminoalkyl aminocarbonyl, and trialkylsilyl; or a pharmaceutically acceptable salt thereof.

[0030] In any of the embodiments above, R<sup>3</sup> and R<sup>4</sup> are particularly hydroxyl.

[0031] In any of the embodiments above, R<sup>5</sup> is particularly hydrogen.

[0032] The term "one or more substituents" in any of the embodiments of the invention refers to 1, 2, 3, 4, or more substituents.

[0033] Particular examples of compounds of the invention are those wherein R<sup>2</sup> is chloro, R<sup>1</sup> is 3-chlorobenzyl, 3-iodobenzyl, 3-bromobenzyl, 1-naphthylmethyl, 2,5-dimethoxybenzyl, 2-hydroxy-5-methoxybenzyl, or *trans*-2-phenyl-cyclopropyl, R<sup>3</sup> and R<sup>4</sup> are hydroxyl, and R<sup>5</sup> is hydrogen.

[0034] Many of the compounds described above have antagonistic as well as partial agonistic properties at the A<sub>3</sub> adenosine receptor, depending upon the parameter studied. The definition of antagonist or agonist is highly dependent upon the cell system and the parameter studied, receptor density, species, and the like.

[0035] The compounds of the invention can be prepared by any suitable method. For example, Fig. 1 illustrates a method of preparing compounds 7b-13b. Figure 2 illustrates a

method of preparing compounds **19a-19g**. Fig. 3 illustrates a method of preparing compounds **20a-20g** and **21a-21g**.

**[0036]** The present invention further provides a pharmaceutical composition comprising a compound as described above and a pharmaceutically acceptable carrier. The present invention provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier and an effective amount, e.g., a therapeutically effective amount, including a prophylactically effective amount, of one or more of the aforesaid compounds, or salts thereof, of the present invention.

**[0037]** The pharmaceutically acceptable carrier can be any of those conventionally used and is limited only by chemico-physical considerations, such as solubility and lack of reactivity with the compound, and by the route of administration. It will be appreciated by one of skill in the art that, in addition to the following described pharmaceutical compositions; the compounds of the present invention can be formulated as inclusion complexes, such as cyclodextrin inclusion complexes, or liposomes.

**[0038]** The pharmaceutically acceptable carriers described herein, for example, vehicles, adjuvants, excipients, or diluents, are well known to those who are skilled in the art and are readily available to the public. It is preferred that the pharmaceutically acceptable carrier be one which is chemically inert to the active compounds and one which has no detrimental side effects or toxicity under the conditions of use.

**[0039]** The choice of carrier will be determined in part by the particular active agent, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of the pharmaceutical composition of the present invention. The following formulations for oral, aerosol, parenteral, subcutaneous, intravenous, intraarterial, intramuscular, interperitoneal, intrathecal, rectal, and vaginal administration are merely exemplary and are in no way limiting.

**[0040]** Formulations suitable for oral administration can consist of (a) liquid solutions, such as an effective amount of the compound dissolved in diluents, such as water, saline, or orange juice; (b) capsules, sachets, tablets, lozenges, and troches, each containing a predetermined amount of the active ingredient, as solids or granules; (c) powders; (d) suspensions in an appropriate liquid; and (e) suitable emulsions. Liquid formulations may include diluents, such as water and alcohols, for example, ethanol, benzyl alcohol, and the polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant, suspending agent, or emulsifying agent. Capsule forms can be of the ordinary

hard- or soft-shelled gelatin type containing, for example, surfactants, lubricants, and inert fillers, such as lactose, sucrose, calcium phosphate, and cornstarch. Tablet forms can include one or more of lactose, sucrose, mannitol, corn starch, potato starch, alginic acid, microcrystalline cellulose, acacia, gelatin, guar gum, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium stearate, calcium stearate, zinc stearate, stearic acid, and other excipients, colorants, diluents, buffering agents, disintegrating agents, moistening agents, preservatives, flavoring agents, and pharmacologically compatible carriers. Lozenge forms can comprise the active ingredient in a flavor, usually sucrose and acacia or tragacanth, as well as pastilles comprising the active ingredient in an inert base, such as gelatin and glycerin, or sucrose and acacia, emulsions, gels, and the like containing, in addition to the active ingredient, such carriers as are known in the art.

**[0041]** The compounds of the present invention, alone or in combination with other suitable components, can be made into aerosol formulations to be administered via inhalation. These aerosol formulations can be placed into pressurized acceptable propellants, such as dichlorodifluoromethane, propane, nitrogen, and the like. They also may be formulated as pharmaceuticals for non-pressured preparations, such as in a nebulizer or an atomizer.

**[0042]** Formulations suitable for parenteral administration include aqueous and non-aqueous, isotonic sterile injection solutions, which can contain anti-oxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. The compound can be administered in a physiologically acceptable diluent in a pharmaceutical carrier, such as a sterile liquid or mixture of liquids, including water, saline, aqueous dextrose and related sugar solutions, an alcohol, such as ethanol, isopropanol, or hexadecyl alcohol, glycols, such as propylene glycol or polyethylene glycol, glycerol ketals, such as 2,2-dimethyl-1,3-dioxolane-4-methanol, ethers, such as poly(ethyleneglycol) 400, an oil, a fatty acid, a fatty acid ester or glyceride, or an acetylated fatty acid glyceride with or without the addition of a pharmaceutically acceptable surfactant, such as a soap or a detergent, suspending agent, such as pectin, carboomers, methylcellulose, hydroxypropylmethylcellulose, or carboxymethylcellulose, or emulsifying agents and other pharmaceutical adjuvants.

**[0043]** Oils, which can be used in parenteral formulations include petroleum, animal, vegetable, or synthetic oils. Specific examples of oils include peanut, soybean, sesame, cottonseed, corn, olive, petrolatum, and mineral. Suitable fatty acids for use in parenteral

formulations include oleic acid, stearic acid, and isostearic acid. Ethyl oleate and isopropyl myristate are examples of suitable fatty acid esters. Suitable soaps for use in parenteral formulations include fatty alkali metal, ammonium, and triethanolamine salts, and suitable detergents include (a) cationic detergents such as, for example, dimethyl dialkyl ammonium halides, and alkyl pyridinium halides, (b) anionic detergents such as, for example, alkyl, aryl, and olefin sulfonates, alkyl, olefin, ether, and monoglyceride sulfates, and sulfosuccinates, (c) nonionic detergents such as, for example, fatty amine oxides, fatty acid alkanolamides, and polyoxyethylene-polypropylene copolymers, (d) amphoteric detergents such as, for example, alkyl-beta-aminopropionates, and 2-alkyl-imidazoline quaternary ammonium salts, and (3) mixtures thereof.

**[0044]** The parenteral formulations will typically contain from about 0.5 to about 25% by weight of the active ingredient in solution. Suitable preservatives and buffers can be used in such formulations. In order to minimize or eliminate irritation at the site of injection, such compositions may contain one or more nonionic surfactants having a hydrophile-lipophile balance (HLB) of from about 12 to about 17. The quantity of surfactant in such formulations ranges from about 5 to about 15% by weight. Suitable surfactants include polyethylene sorbitan fatty acid esters, such as sorbitan monooleate and the high molecular weight adducts of ethylene oxide with a hydrophobic base, formed by the condensation of propylene oxide with propylene glycol. The parenteral formulations can be presented in unit-dose or multi-dose sealed containers, such as ampoules and vials, and can be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, water, for injections, immediately prior to use. Extemporaneous injection solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described.

**[0045]** The compounds of the present invention may be made into injectable formulations. The requirements for effective pharmaceutical carriers for injectable compositions are well known to those of ordinary skill in the art. See *Pharmaceutics and Pharmacy Practice*, J. B. Lippincott Co., Philadelphia, Pa., Banker and Chalmers, eds., pages 238-250 (1982), and *ASHP Handbook on Injectable Drugs*, Toissel, 4th ed., pages 622-630 (1986).

**[0046]** Additionally, the compounds of the present invention may be made into suppositories by mixing with a variety of bases, such as emulsifying bases or water-soluble bases. Formulations suitable for vaginal administration may be presented as pessaries,

tampons, creams, gels, pastes, foams, or spray formulas containing, in addition to the active ingredient, such carriers as are known in the art to be appropriate.

**[0047]** The present invention also provides a method of treating a disease in an animal, e.g., a mammal, comprising administering to the animal an effective amount of a compound or a pharmaceutically acceptable salt of the invention, wherein the disease is selected from the group consisting of cancer, glaucoma, inflammatory diseases, asthma, stroke, myocardial infarction, allergic reactions, rhinitis, poison ivy induced responses, urticaria, scleroderma, arthritis, brain arteriole diameter constriction, bronchoconstriction, and myocardial ischemia. The invention also provides a method for selectively inactivating an A<sub>3</sub> adenosine receptor, or partially activating an A<sub>3</sub> adenosine receptor, in an animal in need thereof, comprising administering to the mammal an effective amount of a compound or pharmaceutically acceptable salt of the invention. The methods of the invention can be applied to any suitable mammal, particularly human.

**[0048]** The term “animal” refers to any member of the animal kingdom. In embodiments, “animal” refers to a human at any stage of development. In embodiments, “animal” includes mammals, birds, reptiles, amphibians, fish, and worms. In certain embodiments, the non-human animal is a mammal, e.g., a rodent, a mouse, a rat, a rabbit, a monkey, a dog, a cat, a sheep, cattle, a primate, or a pig. The animal may also be a transgenic animal, genetically engineered animal, or a clone.


**[0049]** The present invention further provides a method for inactivating A<sub>3</sub> adenosine receptors, or partially activating such a receptor, in a cell comprising contacting the cell with an effective amount of one or more of the inventive compounds or a pharmaceutically acceptable salt thereof. The contacting can be *in vitro* or *in vivo*. When the contacting is done *in vitro*, the contacting can be done by any suitable method, many of which are known in the art. For example, the cell can be provided in a culture medium and the inventive compound introduced into the culture medium *per se*, or as a solution of the compound in an appropriate solvent.

**[0050]** The present invention further provides a method of cardioprotection for preventing or reducing ischemic damage to the heart in an animal in need thereof comprising administering to the animal a compound or salt as described above, particularly, a compound or salt of formula I, wherein R<sup>1</sup> is 3-bromobenzyl or 3-iodobenzyl, R<sup>2</sup> is halo, R<sup>3</sup> and R<sup>4</sup> are hydroxyl, and R<sup>5</sup> is hydrogen.

**[0051]** The compounds or salts thereof can be used in any suitable dose. Suitable doses and dosage regimens can be determined by conventional range finding techniques. Generally treatment is initiated with smaller dosages, which are less than the optimum dose. Thereafter, the dosage is increased by small increments until optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired. In proper doses and with suitable administration of certain compounds, the present invention provides for a wide range of responses. Typically the dosages range from about 0.001 to about 1000 mg/kg body weight of the animal being treated/day. For example, in embodiments, the compounds or salts may be administered from about 100 mg/kg to about 300 mg/kg, from about 120 mg/kg to about 280 mg/kg, from about 140 mg/kg to about 260 mg/kg, from about 150 mg/kg to about 250 mg/kg, from about 160 mg/kg to about 240 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect.

**[0052]** In accordance with another embodiment, the invention provides isotopically labeled compounds described above, for example, compounds labeled with a radioactive or non-radioactive isotope, for use in the determination of drug/tissue distribution assays, in the manipulation of oxidative metabolism via the primary kinetic isotope effect, in identifying potential therapeutic agents for the treatment of diseases or conditions associated with target-receptor mediation. The compounds of the invention can be prepared with a radioactive isotope. Any suitable atom can be replaced with a radioactive isotope, for example, a carbon atom, hydrogen atom, a halogen atom, a sulfur atom, nitrogen atom, or an oxygen atom can be replaced with a corresponding isotope. Thus, for example, a halogen atom can be replaced with  $^{18}\text{F}$ ,  $^{36}\text{Cl}$ ,  $^{75}\text{Br}$ ,  $^{76}\text{Br}$ ,  $^{77}\text{Br}$ ,  $^{82}\text{Br}$ ,  $^{122}\text{I}$ ,  $^{123}\text{I}$ ,  $^{125}\text{I}$ , or  $^{131}\text{I}$ . The use of radiolabeled compounds that may be detected using imaging techniques, such as the Single Photon Emission Computerized Tomography (SPECT), Magnetic Resonance Spectroscopy (MRS), or the Positron Emission Tomography (PET), are known in the art. See, for example, U.S. Pat. Nos. 6,395,742 and 6,472,667.

**[0053]** In accordance with a further embodiment, the invention provides a radiolabeled compound of Formula I:



(I),

wherein

$R^1$  is selected from the group consisting of  $C_6$ - $C_{14}$  aryl,  $C_6$ - $C_{14}$  aryl  $C_1$ - $C_6$  alkyl,  $C_6$ - $C_{14}$  diaryl  $C_1$ - $C_6$  alkyl,  $C_6$ - $C_{14}$  aryl  $C_1$ - $C_6$  alkoxy,  $C_6$ - $C_{14}$  aryl sulfonyl, heterocyclyl  $C_1$ - $C_6$  alkyl, heterocyclyl, heteroaryl  $C_1$ - $C_6$  alkyl, 4-[[[4-[[[(2-amino  $C_1$ - $C_6$  alkyl) amino]-carbonyl]- $C_1$ - $C_6$  alkyl] aniline] carbonyl]  $C_1$ - $C_6$  alkyl]  $C_6$ - $C_{14}$  aryl, and  $C_6$ - $C_{14}$  aryl  $C_3$ - $C_8$  cycloalkyl, wherein the aryl or heterocyclyl portion of  $R^1$  is substituted with one or more halogen atoms that are radioactive;

$R^2$  is selected from the group consisting of hydrogen, halo, amino, hydrazido, mercapto,  $C_1$ - $C_{20}$  alkylamino,  $C_6$ - $C_{14}$  aryl amino,  $C_6$ - $C_{14}$  aryloxy,  $C_1$ - $C_{20}$  alkyl,  $C_1$ - $C_{20}$  alkoxy,  $C_1$ - $C_{20}$  thioalkoxy, pyridylthio,  $C_7$ - $C_{12}$  cycloalkyl  $C_1$ - $C_{20}$  alkyl,  $C_7$ - $C_{12}$  bicycloalkyl  $C_1$ - $C_{20}$  alkyl,  $C_7$ - $C_{12}$  bicycloalkenyl  $C_1$ - $C_{20}$  alkyl,  $C_6$ - $C_{14}$  aryl  $C_1$ - $C_{20}$  alkyl,  $C_2$ - $C_{20}$  alkenyl,  $C_7$ - $C_{12}$  cycloalkyl  $C_2$ - $C_{20}$  alkenyl,  $C_7$ - $C_{12}$  bicycloalkyl  $C_2$ - $C_{20}$  alkenyl,  $C_7$ - $C_{12}$  bicycloalkenyl  $C_2$ - $C_{20}$  alkenyl,  $C_6$ - $C_{14}$  aryl  $C_2$ - $C_{20}$  alkenyl,  $C_2$ - $C_{20}$  alkynyl, carboxy alkyl  $C_2$ - $C_{20}$  alkynyl,  $-C\equiv C-$  ( $CH_2$ )<sub>m</sub>- $C(=O)$ - $O$ - $C_1$ - $C_6$  alkyl,  $-C\equiv C-$  ( $CH_2$ )<sub>m</sub>- $C(=O)$ - $NH$ - $(CH_2)_n$ - $NH_2$ ,  $-C\equiv C-$  ( $CH_2$ )<sub>m</sub>- $C_1$ - $C_6$  alkyl,  $-C\equiv C-$  ( $CH_2$ )<sub>m</sub>-aryl, wherein m and n are independently 1 to 10,  $C_7$ - $C_{12}$  cycloalkyl  $C_2$ - $C_{20}$  alkynyl,  $C_7$ - $C_{12}$  bicycloalkyl  $C_2$ - $C_{20}$  alkynyl,  $C_7$ - $C_{12}$  bicycloalkenyl  $C_2$ - $C_{20}$  alkynyl,  $C_6$ - $C_{14}$  aryl  $C_2$ - $C_{20}$  alkynyl, and the alkyl, cycloalkyl, or aryl portion of  $R^2$  is optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, amino, alkylamino, dialkylamino, sulfur, carboxy, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkyl aminocarbonyl, aminoalkyl aminocarbonyl, and trialkylsilyl;

$R^3$  and  $R^4$  are independently selected from the group consisting of hydroxyl, amino, thiol, ureido, C<sub>1</sub>-C<sub>6</sub> alkyl carbonylamino, hydroxy C<sub>1</sub>-C<sub>6</sub> alkyl, and hydrazinyl; and

$R^5$  is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, heteroaryl, and C<sub>1</sub>-C<sub>6</sub> aminoalkyl;

or a pharmaceutically acceptable salt thereof.

[0054] The halogen atom of the radiolabeled compound or salt in  $R^1$  of the invention can be any suitable isotope, for example, <sup>18</sup>F, <sup>76</sup>Br, or <sup>125</sup>I, preferably <sup>76</sup>Br or <sup>125</sup>I.

[0055] In a particular embodiment, the invention provides radiolabeled compounds or salts wherein  $R^1$  is 3-bromobenzyl or 3-iodobenzyl,  $R^2$  is halo,  $R^3$  and  $R^4$  are hydroxyl, and  $R^5$  is hydrogen.

[0056] Accordingly, the present invention further provides a method of diagnostic imaging of an A<sub>3</sub> adenosine receptor in a tissue or organ of an animal comprising administering an effective amount of a radiolabeled compound or salt as described above to the animal and obtaining an image of the organ or tissue of the animal. The image can be obtained by any suitable imaging technique, for example, SPECT, MRS, and/or PET.

[0057] The present invention also provides a diagnostic method for determining a treatment of a patient for a possible agonist or antagonist of the A<sub>3</sub> adenosine receptors, the treatment comprising:

- (a) administering a radiolabeled compound or salt as described above;
- (b) obtaining a biological sample from the patient;
- (c) determining the level of expression of the A<sub>3</sub> adenosine receptor;
- (d) comparing the level of expression of the receptor to that of a normal population;

and

(e) if the patient's level of expression is higher than that of the normal population, determining a treatment regimen comprising administering an agonist or antagonist of the adenosine receptor whose expression was higher in the patient than that of the normal population.

[0058] The following examples further illustrate the invention but, of course, should not be construed as in any way limiting its scope.

#### EXAMPLE 1

[0059] This example demonstrates a method of preparing compounds in accordance with an embodiment of the invention. D-ribose was protected with TBDPS-Cl followed by alkaline hydrolysis, thus providing acid **2**. Reductive decarboxylation of acid **2** was carried

out using non-toxic tris(trimethylsilyl)silane as a hydrogen donor and produced the silyl ether **3** in 40% yield. The silyl ether **3** was deprotected with TBAF. The resultant alcohol **4** was converted into a key dichloropurine derivative **6** through a Mitsonobu reaction (Fig. 1). Derivative **6** reacted with an excess of the corresponding primary amine to give the  $N^6$  substituted and 2',3'-isopropylidene protected derivatives compounds **7a – 13a**, followed by acid catalyzed deprotection to give the  $N^6$ -3-halobenzyl and related arylmethyl derivatives **7b – 13b**.

**[0060] (1R, 2S, 3R, 4R, 5R)-3,4-O-(Isopropylidene)-2-O-(tert-butyldiphenylsilyl)-2,3,4-trihydroxybicyclo[3.1.0]hexane-1-carboxylic acid (2).** *tert*-Butyldiphenylsilyl chloride (2.70 g, 10 mmol) and triethylamine (2.0 g, 20 mmol) were added to a solution of alcohol **1** (prepared from D-ribose following the standard procedure (Joshi et al. *supra*) 1.22 g, 5 mmol) and imidazole (140 mg, 2 mmol) in DMF (3 mL) while stirring at room temperature. The solution was stirred at 60 °C for 16 h. The reaction mixture was cooled to room temperature and diluted with a 4:1 ethyl acetate – hexane mixture (50 mL), washed with water, dried, and solvent was evaporated. The residue was purified by flash chromatography (0 to 10% ethyl-acetate – hexane) to give ethyl (1*R*, 2*S*, 3*R*, 4*R*, 5*R*)-2,3,4-O-(isopropylidene)-4-O-(*tert*-butyldiphenylsilyl)-2,3,4-trihydroxybicyclo[3.1.0]hexane-1-carboxylate. The compound was dissolved in MeOH (5 mL), 2N aq. NaOH (5 mL) was added, and the reaction mixture was refluxed for 2 h. The reaction mixture was neutralized with NaH<sub>2</sub>PO<sub>4</sub>, and extracted with DCM. The combined DCM solutions were dried and evaporated, and the residue was purified by flash chromatography to give title compound **2** (1.65 g, 73%). <sup>1</sup>H NMR (CDCl<sub>3</sub>),  $\delta$ : 7.72 (d, 4H, J=7.8 Hz), 7.39 (m, 6H), 5.05 (d, 1H, J=6.3 Hz), 4.43 (t, 1H, J=6.0 Hz), 4.08 (t, 1H, J=6.6 Hz), 2.26 (m, 1H), 1.97 (s, 3H), 1.56 (s, 3H), 1.52 (m, 1H), 1.21 (s, 3H), 1.08 (s, 9H).

**[0061] (1S, 2S, 3R, 4R, 5R)-3,4-O-(Isopropylidene)-2-O-(tert-butyldiphenylsilyl)-2,3,4-trihydroxybicyclo[3.1.0]hexane (3).** A 1M solution of DCC in oxygen-free toluene (0.96 mL) was added to a solution of acid **2** (363 mg, 0.80 mmol), 2-mercaptopyridine *N*-oxide (112 mg, 0.88 mmol), and AIBN (40 mg, 0.24 mmol) in dry oxygen-free toluene (4 mL). The reaction mixture was stirred for 4 h at 25 °C, tris(trimethylsilyl)silane (0.50 mL,

1.6 mmol) was added, and the reaction mixture was heated at 85 °C for 4 h. The reaction mixture was evaporated, and the residue was separated by flash chromatography (0 to 10%

ethyl acetate – hexane mixture) to afford the title compound **3** (121 mg, 40%). <sup>1</sup>H NMR (CDCl<sub>3</sub>), δ: 7.76 (d, 4H, J=7.8 Hz), 7.39 (m, 6H), 4.66 (t, 1H, J=6.0 Hz), 4.44 (t, 1H, J=6.6 Hz), 4.03 (t, 1H, J=6.6 Hz), 1.6 (m, 1H), 1.57 (s, 3H), 1.45 (m, 1H), 1.33 (s, 1H), 1.20 (s, 3H), 1.09 (s, 9H), 0.58 (m, 1H).

**[0062] (1*R*, 2*R*, 3*S*, 4*S*, 5*S*)-2,3,4-trihydroxybicyclo[3.1.0]hexane (4), Method B.** A 1M solution of *tert*-butylammonium fluoride in THF (1 mL) was added to a solution of silylether **3** (102 mg, 0.25 mmol) in THF (1 mL). The reaction mixture was left at 20 °C for 16 h and evaporated. The residue was diluted with ethyl acetate (20 mL) and washed with a small amount of brine. The ethyl acetate solution was dried and evaporated, and the residue was purified by flash chromatography to afford the title compound **4** (33 mg, 84%). <sup>1</sup>H NMR and MS are provided under Method A.

**[0063] General procedure for preparation of compounds 7b – 13b.** An amine (RNH<sub>2</sub> in Scheme 3, 0.5 mmol) was added to a solution of **6** (20 mg, 0.06 mmol) in DCM (0.1 mL). The reaction mixture was stirred at room temperature for 16 h. The solvent was removed under vacuum, and the residue was separated by flash chromatography (30 to 100% ethyl acetate-hexane) to afford the corresponding 6-alkylaminopurine derivative that was dissolved in a mixture of MeOH (4 mL), TFA (0.2 mL) and water (2 mL). The reaction mixture was stirred at 70 °C for 16 h, and then evaporated. The residue was evaporated twice with water, and the residue was purified by flash chromatography (50 to 100% ethyl acetate).

**[0064] (1'R, 2'R, 3'S, 4'R, 5'S)-4'-(2-Chloro-6-(3-iodobenzylamino)purine)-2',3'-O-dihydroxybicyclo-[3.1.0]hexane (7b).** Yield 15 mg (51%) <sup>1</sup>H NMR (CD<sub>3</sub>OD), δ: 8.16 (s, 1H), 7.49 (s, 1H), 7.60 (d, 1H, 8.5 Hz), 7.40 (d, 1H, 8.5 Hz), 7.10 (t, 1H, 8.5 Hz), 4.71 (s, 2H), 3.90 (d, 3.3Hz, 1H), 3.65 (s, 1H), 2.05-1.95 (m, 1H), 1.67-1.63 (m, 1H), 1.36 (s, 1H), 1.31-1.27 (m, 1H), 0.95-0.87 (m, 1H), 0.77-0.75 (m, 1H). HRMS calculated for C<sub>18</sub>H<sub>18</sub>ClIN<sub>5</sub>O<sub>2</sub><sup>+</sup> (M+H)<sup>+</sup>: 498.0194; found, 498.0194. HPLC: RT 21.6 min (98%) in solvent system A, 17.0 min (98%) in system B.

**[0065] (1'R, 2'R, 3'S, 4'R, 5'S)-4'-(2-Chloro-6-(3-chlorobenzylamino)purine)-2',3'-O-dihydroxybicyclo-[3.1.0]hexane (8b).** Yield 58%. <sup>1</sup>H NMR (CD<sub>3</sub>OD), δ: 8.16 (br.s., 1H), 7.41 (s, 1H), 7.29 (m, 3H), 4.79 (s, 1H), 4.75 (br. s, 2H), 4.70 (br. t., 1H, J=5.4 Hz), 3.86 (d, 1H, J=6.6 Hz), 1.97 (m, 1H), 1.65 (m, 1H), 1.30 (m, 1H), 0.75 (m, 1H). HRMS (ESI MS m/z): calculated for C<sub>18</sub>H<sub>18</sub>Cl<sub>2</sub>N<sub>5</sub>O<sub>2</sub><sup>+</sup> (M+H)<sup>+</sup>, 406.0832; found, 406.0825. HPLC RT 20.3 min (98%) in solvent system A, 15.6 min (98%) in system B.

**[0066] (1'R, 2'R, 3'S, 4'R, 5'S)-4'-(2-Chloro-6-(3-bromobenzylamino)purine]-2',3'-O-dihydroxybicyclo-[3.1.0]hexane (9b)** Yield 65%.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ): 8.03 (s, 1H), 7.45 (s, 1H), 7.29 (m, 2H), 7.12 (t, 1H,  $J=7.8$  Hz), 4.68 (s, 1H), 4.63 (br. s, 2H), 4.59 (br. t., 1H,  $J=5.4$  Hz), 3.79 (d, 1H,  $J=6.6$  Hz), 1.86 (m, 1H), 1.55 (m, 1H), 1.20 (m, 1H), 0.64 (m, 1H). HRMS (ESI MS m/z) calculated for  $\text{C}_{18}\text{H}_{18}\text{BrClN}_5\text{O}_2^+$  ( $\text{M}+\text{H}$ ) $^+$ , 450.0327; found 450.0315. HPLC RT 20.74 min (98%) in solvent system A, 16.1 min (99%) in system B.

**[0067] (1'R, 2'R, 3'S, 4'R, 5'S)-4'-(2-Chloro-6-(1-naphthylamino)purine]-2',3'-O-dihydroxybicyclo[3.1.0]hexane (10b)** Yield 48%.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ): 8.13 (br.d., 2H,  $J=7.8$  Hz), 7.84 (m, 2H), 7.49 (m, 4H), 5.21 (s, 1H), 4.79 (br. s, 1H), 4.78 (br. s, 2H), 4.67 (br. t., 1H,  $J=5.1$  Hz), 3.88 (d, 1H,  $J=6.6$  Hz), 1.93 (m, 1H), 1.62 (m, 1H), 1.25 (m, 1H), 0.73 (m, 1H). HRMS (ESI MS m/z) calculated for  $\text{C}_{22}\text{H}_{21}\text{ClN}_5\text{O}_2^+$  ( $\text{M}+\text{H}$ ) $^+$ , 422.1378; found 422.1385. HPLC RT 21.5 min (97%) in solvent system A, 17.0 min (98%) in system B.

**[0068] (1'R, 2'R, 3'S, 4'R, 5'S)-4'-(2-Chloro-6-(2,5-dimethoxybenzylamino)purine]-2',3'-O-dihydroxybicyclo-[3.1.0]hexane (11b)** Yield 44%.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ): 8.4 (very br. s, 1H), 6.95 (s, 1H,  $J=2.7$  Hz), 6.89 (d, 1H,  $J=9.3$  Hz), 6.78 (dd, 1H,  $J=2.7, 9.0$  Hz), 4.80 (s, 1H), 4.75 (br. m, 3H), 3.87 (d, 1H,  $J=6.3$  Hz), 3.83 (s, 3H), 3.71 (s, 3H), 1.95 (m, 1H), 1.64 (m, 1H), 1.29 (m, 1H), 0.74 (m, 1H). HRMS (ESI MS m/z) calculated for  $\text{C}_{20}\text{H}_{23}\text{ClN}_5\text{O}_4^+$  ( $\text{M}+\text{H}$ ) $^+$ , 432.1433; found 432.1439. HPLC RT 18.7 min (98%) in solvent system A, 16.6 min (98%) in system B.

**[0069] (1'R, 2'R, 3'S, 4'R, 5'S)-4'-(2-Chloro-6-(2-hydroxy-5-methoxybenzylamino)purine]-2',3'-O-dihydroxybicyclo-[3.1.0]hexane (12b)** Yield 39%.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ): 8.07 (s, 1H), 6.60-6.82 (m, 3H), 4.69 (s, 1H), 4.59 (br. t., 1H,  $J=6.0$  Hz), 4.56 (br. s, 2H), 3.79 (d, 1H,  $J=6.6$  Hz), 3.61 (s, 3H) 1.86 (m, 1H), 1.55 (m, 1H), 1.20 (m, 1H), 0.65 (m, 1H). HRMS (ESI MS m/z) calculated for  $\text{C}_{19}\text{H}_{21}\text{ClN}_5\text{O}_4^+$  ( $\text{M}+\text{H}$ ) $^+$ , 418.1277; found, 418.1277. HPLC RT 16.0 min (100%) in solvent system A, 11.0 min (98%) in system B.

**[0070] (1'R, 2'R, 3'S, 4'R, 5'S)-4'-(2-Chloro-6-(*trans*-2-phenylcyclopropylamino)purine]-2',3'-O-dihydroxybicyclo-[3.1.0]hexane (13b)** Yield 52%.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ): 8.16 (very br.s., 1H), 7.0-7.48 (m, 5H), 4.79 (s, 1H), 4.68 (br. s, 2H), 3.88 (d, 1H,  $J=5.7$  Hz), 2.17 (m, 1H) 1.97 (m, 1H), 1.65 (m, 1H), 1.29 (m, 2H), 0.74 (m, 1H). HRMS (ESI MS m/z) calculated for  $\text{C}_{20}\text{H}_{21}\text{ClN}_5\text{O}_2^+$  ( $\text{M}+\text{H}$ ) $^+$ , 398.1378; found, 398.1372. HPLC RT 20.3 min (99%) in solvent system A, 15.6 min (98%) in system B.

## EXAMPLE 2

[0071] This Example illustrates the ability of the compounds in accordance with an embodiment of the invention to bind to A<sub>3</sub> adenosine receptors. The binding affinity values are set forth in Table 1.

**Receptor binding and functional assays**

[0072] [<sup>125</sup>I]N<sup>6</sup>-(4-Amino-3-iodobenzyl)adenosine-5'-N-methyluronamide (I-AB-MECA; 2000 Ci/mmol), [<sup>3</sup>H]cyclic AMP (40 Ci/mmol), and other radioligands were purchased from Perkin-Elmer Life and Analytical Science (Boston, MA). [<sup>3</sup>H]CCPA (2-chloro-N<sup>6</sup>-cyclopentyladenosine) was a custom synthesis product (Perkin Elmer). Test compounds were prepared as 5 mM stock solutions in DMSO and stored frozen.

[0073] Cell culture and membrane preparation: CHO (Chinese hamster ovary) cells expressing the recombinant human A<sub>3</sub>AR were cultured in DMEM (Dulbecco's modified Eagle's medium) supplemented with 10% fetal bovine serum, 100 units/mL penicillin, 100 µg/mL streptomycin, 2 µmol/mL glutamine and 800 µg/mL geneticin. The CHO cells expressing rat A<sub>3</sub>ARs were cultured in DMEM and F12 (1:1). Cells were harvested by trypsinization. After homogenization and suspension, cell membranes were centrifuged at 500 g for 10 min, and the pellet was re-suspended in 50 mM Tris-HCl buffer (pH 8.0) containing 10 mM MgCl<sub>2</sub>, 1 mM EDTA and 0.1 mg/mL CHAPS (3[(3-cholamidopropyl)dimethylammonio]-propanesulfonic acid). The suspension was homogenized with an electric homogenizer for 10 sec, and was then re-centrifuged at 20,000 g for 20 min at 4°C. The resultant pellets were resuspended in buffer in the presence of adenosine deaminase (3 Units/mL), and the suspension was stored at -80°C until the binding experiments. The protein concentration was measured using the Bradford assay. Bradford, M.M. *Anal. Biochem.* **1976**, 72, 248.

[0074] Binding assays at the A<sub>1</sub> and A<sub>2A</sub> receptors: For binding to human A<sub>1</sub> receptors, see (a) Schwabe, U.; Trost, T. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **1980**, 313, 179. (b) Perreira, M.; Jiang, J.K.; Klutz, A.M.; Gao, Z.G.; Shainberg, A.; Lu, C.; Thomas, C.J.; Jacobson, K.A. *J. Med. Chem.* **2005**, 48, 4910.

[0075] [<sup>3</sup>H]R-PIA (N<sup>6</sup>-[(R)-phenylisopropyl]adenosine, 2 nM) or [<sup>3</sup>H]CCPA (0.5 nM) was incubated with membranes (40 µg/tube) from CHO cells stably expressing human A<sub>1</sub> receptors at 25°C for 60 min in 50 mM Tris-HCl buffer (pH 7.4; MgCl<sub>2</sub>, 10 mM) and increasing concentrations of the test ligand in a total assay volume of 200 µl. Nonspecific binding was determined using 10 µM of CPA (N<sup>6</sup>-cyclopentyladenosine). For human A<sub>2A</sub>

receptor binding (Jarvis, M. F.; Schutz, R.; Hutchison, A. J.; Do, E.; Sills, M. A.; Williams, M. J. *Pharmacol. Exp. Ther.* **1989**, *251*, 888-893) membranes (20 µg/tube) from HEK-293 cells stably expressing human A<sub>2A</sub> receptors were incubated with [<sup>3</sup>H]CGS21680 (2-[p-(2-carboxyethyl)phenyl-ethylamino]-5'-N-ethylcarboxamido-adenosine, 15 nM) and increasing concentrations of the test ligand at 25°C for 60 min in 200 µl 50 mM Tris-HCl, pH 7.4, containing 10 mM MgCl<sub>2</sub>. NECA (10 µM) was used to define nonspecific binding. The reaction was terminated by filtration with GF/B filters.

[0076] Binding assay at the human A<sub>3</sub> receptor: For the competitive binding assay, each tube contained 50 µL membrane suspension (20 µg protein), 25 µL of [<sup>125</sup>I]I-AB-MECA (1.0 nM), Olah, M.E., Gallo-Rodriguez, C., Jacobson, K.A., Stiles, G.L. *Mol. Pharmacol.* **1994**, *45*, 978, and 25 µL of increasing concentrations of the test ligands in Tris-HCl buffer (50 mM, pH 8.0) containing 10 mM MgCl<sub>2</sub>, 1 mM EDTA. Nonspecific binding was determined using 10 µM of Cl-IB-MECA in the buffer. The mixtures were incubated at 37°C for 60 min. Binding reactions were terminated by filtration through Whatman GF/B filters under reduced pressure using a MT-24 cell harvester (Brandell, Gaithersburgh, MD, USA). Filters were washed three times with 9 mL ice-cold buffer. Radioactivity was determined in a Beckman 5500B γ-counter. IC<sub>50</sub> values were converted to K<sub>i</sub> values as described in Cheng, Y.; Prusoff, W. H. *Biochem. Pharmacol.* **1973**, *22*, 3099.

[0077] Cyclic AMP accumulation assay: Intracellular cyclic AMP levels were measured with a competitive protein binding method. Nordstedt, C.; Fredholm, B. B. *Anal. Biochem.* **1990**, *189*, 231; Post, S. R.; Ostrom, R. S.; Insel, P. A. *Methods Mol. Biol.* **2000**, *126*, 363. CHO cells that expressed the recombinant human or rat A<sub>3</sub>AR or the human A<sub>1</sub> or A<sub>2B</sub>AR were harvested by trypsinization. After centrifugation and resuspended in medium, cells were planted in 24-well plates in 1.0 mL medium. After 24 h, the medium was removed and cells were washed three times with 1 mL DMEM, containing 50 mM HEPES, pH 7.4. Cells were then treated with the agonist NECA and/or test compound (e.g. 7b) in the presence of rolipram (10 µM) and adenosine deaminase (3 units/mL). After 45 min forskolin (10 µM) was added to the medium, and incubation was continued for an additional 15 min. The reaction was terminated by removing the supernatant, and cells were lysed upon the addition of 200 µL of 0.1 M ice-cold HCl. The cell lysate was resuspended and stored at -20°C. For determination of cyclic AMP production, protein kinase A (PKA) was incubated with [<sup>3</sup>H]cyclic AMP (2 nM) in K<sub>2</sub>HPO<sub>4</sub>/EDTA buffer (K<sub>2</sub>HPO<sub>4</sub>, 150 mM; EDTA, 10 mM), 20

$\mu$ L of the cell lysate, and 30  $\mu$ L 0.1 M HCl or 50  $\mu$ L of cyclic AMP solution (0-16 pmol/200  $\mu$ L for standard curve). Bound radioactivity was separated by rapid filtration through Whatman GF/C filters and washed once with cold buffer. Bound radioactivity was measured by liquid scintillation spectrometry.

[0078]  $[^{35}\text{S}]$ GTP $\gamma$ S binding assay:  $[^{35}\text{S}]$ GTP $\gamma$ S binding was measured by a variation of the method described. (a) Lorenzen, A.; Lang H.; Schwabe U. *Biochem. Pharmacol.* **1998**, *56*, 1287. (b) Jacobson, K.A.; Ji, X.-d.; Li, A.H.; Melman, N.; Siddiqui, M.A.; Shin, K.J.; Marquez, V.E.; Ravi, R.G. *J. Med. Chem.* **2000**, *43*, 2196. Each assay tube consisted of 200  $\mu$ L buffer containing 50 mM Tris HCl (pH 7.4), 1 mM EDTA, 1 mM MgCl<sub>2</sub>, 1  $\mu$ M GDP, 1 mM dithiothreitol, 100 mM NaCl, 3 U/ml ADA, 0.2 nM  $[^{35}\text{S}]$ GTP $\gamma$ S, 0.004% 3-[(3-cholamidopropyl) dimethylammonio]propanesulfonate (CHAPS), and 0.5% bovine serum albumin. Incubations were started upon addition of the membrane suspension (CHO cells stably expressing either the native human A<sub>1</sub>AR or A<sub>3</sub>AR, 5  $\mu$ g protein/tube) to the test tubes, and they were carried out in duplicate for 30 min at 25 °C. The reaction was stopped by rapid filtration through Whatman GF/B filters, pre-soaked in 50 mM Tris HCl, 5 mM MgCl<sub>2</sub> (pH 7.4) containing 0.02% CHAPS. The filters were washed twice with 3 mL of the same buffer, and retained radioactivity was measured using liquid scintillation counting. Non-specific binding of  $[^{35}\text{S}]$ GTP $\gamma$ S was measured in the presence of 10  $\mu$ M unlabelled GTP $\gamma$ S. None of the compounds >10% stimulation; thus, they are antagonists of the A<sub>3</sub> adenosine receptor

Table 1. Affinity data for compounds in accordance with an embodiment of the invention.

Compound	R <sup>1</sup> In Formula I, R <sup>2</sup> = Cl, R <sup>3</sup> and R <sup>4</sup> = OH and R <sup>5</sup> = H	Affinity (K <sub>i</sub> , nM) or % inhibition <sup>a</sup>				% Efficac y <sup>b</sup>
		A <sub>1</sub>	A <sub>2A</sub>	A <sub>3</sub>	A <sub>3</sub>	
<b>7b</b>	3-I-Phenyl-CH <sub>2</sub>	3040 ± 610	1080 ± 310	1.44 ± 0.60	1.0 ± 3.2	
<b>8b</b>	3-Cl-Phenyl-CH <sub>2</sub>	3070 ± 1500	4510 ± 910	1.06 ± 0.36	2.9 ± 3.7	
<b>9b</b>	3-Br-Phenyl-CH <sub>2</sub>	1760 ± 1010	1600 ± 480	0.73 ± 0.30	5.8 ± 0.8	
<b>10b</b>	1-Naphthyl-CH <sub>2</sub>	1120 ± 640	1530 ± 350	1.42 ± 0.12	3.1 ± 0.3	
<b>11b</b>	2,5-diMeO-Ph-CH <sub>2</sub>	3000 ± 1260	2620 ± 730	1.58 ± 0.56	4.6 ± 3.8	
<b>12b</b>	2-OH-5-MeO-Ph-CH <sub>2</sub>	1110 ± 300	6870 ± 1440	4.06 ± 0.35	0.4 ± 1.3	
<b>13b</b>	<i>trans</i> -2-Ph-cyclopropyl	1790 ± 1430	2010 ± 890	1.30 ± 0.39	9.7 ± 4.1	

<sup>a</sup> All experiments were done on CHO or HEK (A<sub>2A</sub> only) cells stably expressing one of four subtypes of human ARs. The binding affinity for A<sub>1</sub>, A<sub>2A</sub> and A<sub>3</sub>ARs was expressed as K<sub>i</sub> values (n = 3-5) and was determined by using agonist radioligands ([<sup>3</sup>H]CCPA or ([<sup>3</sup>H]R-PIA), (<sup>3</sup>H]CGS21680), [<sup>125</sup>I]I-AB-MECA, respectively. The potency at the A<sub>2B</sub>AR was expressed as EC<sub>50</sub> values and was determined by stimulation of cyclic AMP production in AR-transfected CHO cells. A percent in parentheses refers to inhibition of radioligand binding at 10 μM.

<sup>b</sup> measured by [<sup>35</sup>S]GTPγS binding assay.

[0079] In accordance with one method of biological assay, compounds **7b** – **9b** (3-halobenzyl) in the (N)-methanocarba series were potent A<sub>3</sub> AR antagonists with binding K<sub>i</sub> values of 0.7 – 1.4 nM. Compound **9b** (3-bromobenzyl analogue) proved to be the most potent A<sub>3</sub>AR antagonist of this series in binding with a K<sub>i</sub> value of 0.73 nM, and it displayed high selectivity (2400-fold and 2190-fold in comparison to the A<sub>1</sub> and A<sub>2A</sub>AR, respectively). The most A<sub>3</sub>AR selective compound was the 3-chloro analogue **8b** with 2900-fold and 4250-fold selectivity in comparison to the A<sub>1</sub> and A<sub>2A</sub>AR, respectively. The SAR of substitution of the N<sup>6</sup>-benzyl group further showed that dimethoxy substitution (**11b**), fusion of the phenyl ring to a second ring (**10b**), and extension by one carbon (i.e., in the rotationally constrained 2-phenylcyclopropyl analogue, **13b**) were all tolerated with nanomolar binding affinity at the A<sub>3</sub>AR. Compound **12b**, a demethylated analogue of **11b**, was slightly less potent in binding to the A<sub>3</sub>AR.

[0080] In a functional assay of [<sup>35</sup>S]GTPγS binding induced by A<sub>3</sub>AR activation, **7b** completely inhibited stimulation by 1 μM NECA (5'-N-ethylcarboxamidoadenosine) with an IC<sub>50</sub> of 29.8 nM (Figure 1). Schild analysis of the right shifts by **7b** of the response curves in the inhibition of adenylate cyclase by NECA provided a K<sub>B</sub> value of 8.9 nM.

[0081] When compared in the ability to stimulate the A<sub>3</sub>AR using multiple functional criteria, different results were obtained. In the cAMP assays, compounds **7b** and **9b** exhibited partial agonism at A<sub>3</sub>AR with percent relative efficacies of 44±6 and 46±4, respectively, and the EC<sub>50</sub> values were respectively, 12±1 and 4.2±0.6 nM.

### EXAMPLE 3

[0082] This example illustrates a method of preparing a radioiodinated compound in accordance with an embodiment of the invention. Compound **7b** having <sup>125</sup>I was prepared as follows. The (radio)iodination of compound **7b** on its N<sup>6</sup>-3-iodobenzyl substituent was accomplished in high yield by iododestannylation of a 3-(trimethylstannyl)benzyl precursor through a “cold” iodination reaction as shown in Figure 6.

[0083] **Materials and instrumentation.** Hexamethyltin and other reagents, including pharmacological agents, were purchased from Sigma-Aldrich Chemical Company, except where noted. Sodium [<sup>125</sup>I]iodide (17.4 Ci/mg) in NaOH (1.0x10<sup>-5</sup> M) was supplied by Perkin–Elmer Life and Analytical Science. <sup>1</sup>H NMR spectra were obtained with a Varian Gemini 300 spectrometer using CDCl<sub>3</sub> and CD<sub>3</sub>OD as solvents. Chemical shifts are expressed in δ values (ppm) with tetramethylsilane (δ 0.00) for CDCl<sub>3</sub> and water (δ 3.30) for

CD<sub>3</sub>OD. TLC analysis was carried out on aluminum sheets precoated with silica gel F<sub>254</sub> (0.2 mm) from Aldrich. HPLC mobile phases consisted of CH<sub>3</sub>CN/tetrabutyl ammonium phosphate (5 mM) from 20/80 to 60/40 in 20 min, flow rate 1.0 ml/min. High-resolution mass measurements were performed on Micromass/Waters LCT Premier Electrospray Time of Flight (TOF) mass spectrometer coupled with a Waters HPLC system.

[0084] **Preparation of 23: (1'R, 2'R, 3'S, 4'R, 5'S)-4'-(2-Chloro-6-(3-trimethylstannylbenzylamino)purine]-2',3'-O-dihydroxybicyclo-[3.1.0]hexane (1).** 7b (8.95 mg, 0.018 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (2.7 mg), and hexamethyltin (11 µL, 0.054 mmol) were mixed together in anhydrous dioxane (2 ml), and the resulting reaction mixture was stirred at 70 °C for 2 h. The mixture was concentrated under reduced pressure. The product was purified by flash chromatography by using CHCl<sub>3</sub>: MeOH (10:1) as the eluant to afford the stannyl derivative 23 (9.3 mg, 90%) as an oil. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>), 7.81 (s, 1H), 7.53 (s, 1H), 7.34 (m, 2H), 7.33 (m, 1H), 6.49 (br s, 1H), 4.88 (br s, 2H), 4.00 (m, 2H), 3.71 (s, 1H), 3.65 (m, 1H), 3.47 (m, 1H), 2.02 (m, 1H), 1.96 (s, 1H), 1.64 (m, 1H), 1.28 (m, 2H), 0.81 (m, 1H), 0.29 (s, 9H). HRMS (M + 1)<sup>+</sup>: calculated for C<sub>21</sub>H<sub>27</sub>ClIN<sub>5</sub>O<sub>2</sub>Sn<sup>+</sup> (M+H)<sup>+</sup> 535.6338, found 536.0823 HPLC: R<sub>t</sub> = 22.1 min. HPLC system: 5 mM TBAP/CH<sub>3</sub>CN from 80/20 to 60/40 in 25 min, then isocratic for 2 min; flow rate of 1 ml/min.

[0085] The trimethylstannyl intermediate 23 (0.1 mg) was reacted sodium [<sup>125</sup>I] iodide in NaOH (1.0 x 10<sup>-5</sup> M) to obtain [<sup>125</sup>I] 7b, following the procedure disclosed in Vaidyanathan G., et al., *Nat. Protocols* 1: 707-713 (2006).

[0086] Figure 7A depicts the non-specific, specific, and total binding of [<sup>125</sup>I] 7b on mouse A<sub>3</sub> adenosine receptor. Figure 7B depicts the extent of specific binding as a function of the concentration of the compound. The compound was an agonist of the mouse A<sub>3</sub> adenosine receptor.

#### EXAMPLE 4

[0087] This example illustrates a method of preparing a radiolabeled ligand, that is <sup>76</sup>Br-labeled compound 9b in accordance with an embodiment of the invention. Bromine-76 was prepared from an arsenic metal target using the <sup>75</sup>As (<sup>3</sup>He, 2n) yielding <sup>76</sup>Br nuclear reaction. The <sup>76</sup>Br was processed after allowing for the decay of the simultaneously produced Br-75 (t<sub>1/2</sub> = 1.6 h).

[0088] An aliquot of the aqueous solution of Br-76 (about 10-20 µl, 18.5-37.0 MBq) is added to a 1-mL reaction vial and the solvent evaporated with argon flow. Trimethylstannyl intermediate 23 in acetonitrile is added to the vial containing the Br-76 radioactivity and

followed by adding 37% peracetic acid in acetonitrile. The vial is sealed and placed on an 80 °C heating block and heated for 30 min. At the end of the reaction, the reaction mixture is loaded onto a Phenomenex Luna C18 (2) column (250 x 4.6 mm) and eluted with 100 mM ammonium acetate/acetonitrile (60/40) at a flow rate of 1.2 mL/min. The radioactivity peak containing the desired product ( $t_R=10$  min) is collected and analyzed on a separate HPLC system for determination of purity and specific activity.

[0089] In vivo biodistribution of compound Br-76 labeled compound **9b** was carried out in rats. All studies in live animals were conducted under protocol approved by the NIH Animal Care and Use Committee. The biodistribution was evaluated after intravenous administration to adult Sprague-Dawley rats. The animals were sacrificed at 15, 30, 60, and 120 min and various tissues were harvested for gamma counting. The data are reported in units of percentage of injected dose per gram in Figure 8. The compound exhibited antagonistic properties to the A<sub>3</sub> adenosine receptor albeit at a low magnitude of uptake. The low uptake may be due to the lower age of the animals. The uptake in the A<sub>3</sub>AR-containing testes continued to increase with time after injection (0.09% ID/g at 15 min to 0.18% ID/g at 2 h). Blood continued to provide an input function over 2h. In spite of a potential testes-blood barrier, uptake of the antagonist increased with time, which indicates that the compound may be a viable molecular imaging probe for pathological conditions with elevated A<sub>3</sub>AR.

[0090] [BLANK]


[0091] The use of the terms “a” and “an” and “the” and similar referents in the context of describing the invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. The terms “comprising,” “having,” “including,” and “containing” are to be construed as open-ended terms (i.e., meaning “including, but not limited to,”) unless otherwise noted. Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or

otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., “such as”) provided herein, is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention unless otherwise claimed. No language in the specification should be construed as indicating any non-claimed element as essential to the practice of the invention.

**[0092]** Preferred embodiments of this invention are described herein, including the best mode known to the inventors for carrying out the invention. Variations of those preferred embodiments may become apparent to those of ordinary skill in the art upon reading the foregoing description. The inventors expect skilled artisans to employ such variations as appropriate, and the inventors intend for the invention to be practiced otherwise than as specifically described herein. Accordingly, this invention includes all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in all possible variations thereof is encompassed by the invention unless otherwise indicated herein or otherwise clearly contradicted by context.

## WE CLAIM:

1. A compound of Formula I:



wherein

$R^1$  is selected from the group consisting of hydrogen,  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkoxy, hydroxyl,  $C_3$ - $C_8$  cycloalkyl,  $C_3$ - $C_8$  cycloalkyl  $C_1$ - $C_6$  alkyl,  $C_3$ - $C_8$  dicycloalkyl  $C_1$ - $C_6$  alkyl,  $C_7$ - $C_{12}$  bicycloalkyl  $C_1$ - $C_6$  alkyl,  $C_7$ - $C_{14}$  tricycloalkyl  $C_1$ - $C_6$  alkyl,  $C_6$ - $C_{14}$  aryl,  $C_6$ - $C_{14}$  aryl  $C_1$ - $C_6$  alkyl,  $C_6$ - $C_{14}$  diaryl  $C_1$ - $C_6$  alkyl,  $C_6$ - $C_{14}$  aryl  $C_1$ - $C_6$  alkoxy,  $C_1$ - $C_6$  alkyl carbonyl, sulfonyl,  $C_1$ - $C_6$  alkyl sulfonyl,  $C_6$ - $C_{14}$  aryl sulfonyl, heterocyclyl  $C_1$ - $C_6$  alkyl, heterocyclyl, heteroaryl  $C_1$ - $C_6$  alkyl, 4-[[[4-[[[(2-amino  $C_1$ - $C_6$  alkyl) amino]-carbonyl]- $C_1$ - $C_6$  alkyl] anilino] carbonyl]  $C_1$ - $C_6$  alkyl]  $C_6$ - $C_{14}$  aryl, and  $C_6$ - $C_{14}$  aryl  $C_3$ - $C_8$  cycloalkyl, wherein the aryl or heterocyclyl portion of  $R^1$  is optionally substituted with one or more substituents selected from the group consisting of halo, amino, hydroxyl, carboxy,  $C_1$ - $C_6$  alkoxy carbonyl, aminocarbonyl,  $C_1$ - $C_6$  alkylaminocarbonyl,  $C_1$ - $C_6$  dialkyl aminocarbonyl,  $C_1$ - $C_6$  alkyl,  $C_2$ - $C_6$  alkenyl,  $C_2$ - $C_6$  alkynyl,  $C_1$ - $C_6$  alkoxy,  $C_6$ - $C_{14}$  aryloxy, hydroxy  $C_1$ - $C_6$  alkyl, hydroxy  $C_2$ - $C_6$  alkenyl, hydroxy  $C_2$ - $C_6$  alkynyl, carboxy  $C_1$ - $C_6$  alkyl, carboxy  $C_2$ - $C_6$  alkenyl, carboxy  $C_2$ - $C_6$  alkynyl, aminocarbonyl  $C_1$ - $C_6$  alkyl, aminocarbonyl  $C_2$ - $C_6$  alkenyl, aminocarbonyl  $C_2$ - $C_6$  alkynyl, and  $C\equiv C-(CH_2)_n-COR^7$  wherein  $R^7$  is selected from the group consisting of  $OH$ ,  $OR^8$ , and  $NR^9R^{10}$ , wherein  $R^8$  is selected from the group consisting of  $C_1$ - $C_6$  alkyl,  $C_3$ - $C_8$  cycloalkyl,  $C_3$ - $C_8$  cycloalkyl  $C_1$ - $C_6$  alkyl,  $C_3$ - $C_8$  dicycloalkyl  $C_1$ - $C_6$  alkyl,  $C_7$ - $C_{12}$  bicycloalkyl  $C_1$ - $C_6$  alkyl,  $C_7$ - $C_{14}$  tricycloalkyl  $C_1$ - $C_6$  alkyl,  $C_6$ - $C_{14}$  aryl,  $C_6$ - $C_{14}$  aryl  $C_1$ - $C_6$  alkyl,  $C_6$ - $C_{14}$  diaryl  $C_1$ - $C_6$  alkyl; and  $R^9$  and  $R^{10}$  are independently selected from

the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, and (CH<sub>2</sub>)<sub>n</sub>R<sup>11</sup> wherein R<sup>11</sup> is NR<sup>12</sup>R<sup>13</sup>, wherein R<sup>12</sup> and R<sup>13</sup> are independently selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, and COR<sup>14</sup> wherein R<sup>14</sup> is hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl; wherein n is an integer from 1 to 10; and the alkyl or cycloalkyl portion of R<sup>1</sup> is optionally substituted with one or more substituents selected from the group consisting of halo, amino, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>6</sub>-C<sub>14</sub> aryloxy, C<sub>1</sub>-C<sub>6</sub> hydroxyalkyl, C<sub>2</sub>-C<sub>6</sub> hydroxyalkenyl, C<sub>2</sub>-C<sub>6</sub> hydroxy alkynyl, aminocarbonyl C<sub>1</sub>-C<sub>6</sub> alkoxy, and C<sub>6</sub>-C<sub>14</sub> aryl C<sub>1</sub>-C<sub>6</sub> alkoxy;

R<sup>2</sup> is selected from the group consisting of hydrogen, halo, amino, hydrazido, mercapto, C<sub>1</sub>-C<sub>20</sub> alkylamino, C<sub>6</sub>-C<sub>14</sub> aryl amino, C<sub>6</sub>-C<sub>14</sub> aryloxy, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>1</sub>-C<sub>20</sub> alkoxy, C<sub>1</sub>-C<sub>20</sub> thioalkoxy, pyridylthio, C<sub>7</sub>-C<sub>12</sub> cycloalkyl C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkyl C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkenyl C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>6</sub>-C<sub>14</sub> aryl C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>7</sub>-C<sub>12</sub> cycloalkyl C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkyl C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>6</sub>-C<sub>14</sub> aryl C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>2</sub>-C<sub>20</sub> alkynyl, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-C(=O)-O-C<sub>1</sub>-C<sub>6</sub> alkyl, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-C(=O)-NH-(CH<sub>2</sub>)<sub>n</sub>-NH<sub>2</sub>, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-C<sub>1</sub>-C<sub>6</sub> alkyl, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-aryl, wherein m and n are independently 1 to 10, C<sub>7</sub>-C<sub>12</sub> cycloalkyl C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkyl C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkenyl C<sub>2</sub>-C<sub>20</sub> alkynyl, and C<sub>6</sub>-C<sub>14</sub> aryl C<sub>2</sub>-C<sub>20</sub> alkynyl, and the alkyl, cycloalkyl, or aryl portion of R<sup>2</sup> is optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, amino, alkylamino, dialkylamino, thiol, carboxy, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkyl aminocarbonyl, aminoalkyl aminocarbonyl, and trialkylsilyl;

R<sup>3</sup> and R<sup>4</sup> are independently selected from the group consisting of hydroxyl, amino, thiol, ureido, C<sub>1</sub>-C<sub>6</sub> alkyl carbonylamino, hydroxy C<sub>1</sub>-C<sub>6</sub> alkyl, and hydrazinyl; and

R<sup>5</sup> is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, heteroaryl, and C<sub>1</sub>-C<sub>6</sub> aminoalkyl;

or a pharmaceutically acceptable salt thereof.

2. The compound of claim 1, wherein R<sup>1</sup> is selected from the group consisting of C<sub>6</sub>-C<sub>14</sub> aryl C<sub>1</sub>-C<sub>6</sub> alkyl, heteroaryl, and C<sub>6</sub>-C<sub>14</sub> aryl C<sub>3</sub>-C<sub>8</sub> cycloalkyl, wherein the aryl portion of R<sup>1</sup>

is optionally substituted with one or more substituents selected from the group consisting of halo, amino, hydroxyl, carboxy, C<sub>1</sub>-C<sub>6</sub> alkoxycarbonyl, aminocarbonyl, C<sub>1</sub>-C<sub>6</sub> alkylaminocarbonyl, C<sub>1</sub>-C<sub>6</sub> dialkyl aminocarbonyl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>6</sub>-C<sub>14</sub> aryloxy, hydroxy C<sub>1</sub>-C<sub>6</sub> alkyl, hydroxy C<sub>2</sub>-C<sub>6</sub> alkenyl, hydroxy C<sub>2</sub>-C<sub>6</sub> alkynyl, carboxy C<sub>1</sub>-C<sub>6</sub> alkyl, carboxy C<sub>2</sub>-C<sub>6</sub> alkenyl, carboxy C<sub>2</sub>-C<sub>6</sub> alkynyl, aminocarbonyl C<sub>1</sub>-C<sub>6</sub> alkyl, aminocarbonyl C<sub>2</sub>-C<sub>6</sub> alkenyl, aminocarbonyl C<sub>2</sub>-C<sub>6</sub> alkynyl, and C≡C-(CH<sub>2</sub>)<sub>n</sub>-COR<sup>7</sup> wherein R<sup>7</sup> is selected from the group consisting of OH, OR<sup>8</sup>, and NR<sup>9</sup>R<sup>10</sup>, wherein R<sup>8</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>8</sub> dicycloalkyl C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkyl C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>7</sub>-C<sub>14</sub> tricycloalkyl C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>6</sub>-C<sub>14</sub> aryl, C<sub>6</sub>-C<sub>14</sub> aryl C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>6</sub>-C<sub>14</sub> and diaryl C<sub>1</sub>-C<sub>6</sub> alkyl; and R<sup>9</sup> and R<sup>10</sup> are independently selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, and (CH<sub>2</sub>)<sub>n</sub>R<sup>11</sup> wherein R<sup>11</sup> is NR<sup>12</sup>R<sup>13</sup>, wherein R<sup>12</sup> and R<sup>13</sup> are independently selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, and COR<sup>14</sup> wherein R<sup>14</sup> is hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl; wherein n is an integer from 1 to 10; and the alkyl or cycloalkyl portion of R<sup>1</sup> is optionally substituted with one or more substituents selected from the group consisting of halo, amino, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>6</sub>-C<sub>14</sub> aryloxy, hydroxy C<sub>1</sub>-C<sub>6</sub> alkyl, hydroxy C<sub>2</sub>-C<sub>6</sub> alkenyl, hydroxy C<sub>2</sub>-C<sub>6</sub> alkynyl, aminocarbonyl C<sub>1</sub>-C<sub>6</sub> alkoxy, and C<sub>6</sub>-C<sub>14</sub> aryl C<sub>1</sub>-C<sub>6</sub> alkoxy, or a pharmaceutically acceptable salt thereof.

3. The compound of claim 1 or 2, wherein R<sup>1</sup> is selected from the group consisting of benzyl, phenyl cyclopropyl, or 1-naphthyl methyl, wherein the phenyl or naphthyl portion of R<sup>1</sup> is optionally substituted with one or more substituents selected from the group consisting of halo, amino, hydroxyl, carboxy, C<sub>1</sub>-C<sub>6</sub> alkoxycarbonyl, aminocarbonyl, C<sub>1</sub>-C<sub>6</sub> alkylaminocarbonyl, C<sub>1</sub>-C<sub>6</sub> dialkyl aminocarbonyl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, phenoxy, hydroxy C<sub>1</sub>-C<sub>6</sub> alkyl, hydroxy C<sub>2</sub>-C<sub>6</sub> alkenyl, and hydroxy C<sub>2</sub>-C<sub>6</sub> alkynyl, or a pharmaceutically acceptable salt thereof.

4. The compound of any one of claims 1-3, wherein R<sup>1</sup> is benzyl, phenyl cyclopropyl, or 1-naphthyl methyl, wherein the phenyl or naphthyl portion of R<sup>1</sup> is optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, and C<sub>1</sub>-C<sub>6</sub> alkoxy, or a pharmaceutically acceptable salt thereof.

5. The compound of any one of claims 1-4, wherein R<sup>1</sup> is benzyl, or a pharmaceutically acceptable salt thereof.

6. The compound of any one of claims 1-4, wherein R<sup>1</sup> is benzyl substituted with one or more substituents selected from the group consisting of halo and C<sub>1</sub>-C<sub>6</sub> alkoxy, or a pharmaceutically acceptable salt thereof.

7. The compound of claim 6, wherein the halo is chloro, bromo, or iodo, or a pharmaceutically acceptable salt thereof.

8. The compound of any one of claims 1-4 and 6-7, wherein R<sup>1</sup> is selected from the group consisting of 3-chlorobenzyl, 3-bromobenzyl, 3-iodobenzyl, 2-hydroxy-5-methoxy-benzyl, and 2,5-dimethoxybenzyl, or a pharmaceutically acceptable salt thereof.

9. The compound of claim 3 or 4, wherein the phenyl cyclopropyl is *trans*-2-phenyl-1-cyclopropyl, or a pharmaceutically acceptable salt thereof.

10. The compound of any one of claims 1-9, wherein R<sup>2</sup> is halo, or a pharmaceutically acceptable salt thereof.

11. The compound of any one of claims 1-10, wherein R<sup>2</sup> is chloro, bromo, or iodo, or a pharmaceutically acceptable salt thereof.

12. The compound of any one of claims 1-9, wherein R<sup>2</sup> is -C≡C-(CH<sub>2</sub>)<sub>m</sub>-CH<sub>3</sub>, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-aryl, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-C(=O)-O-CH<sub>3</sub>, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-C(=O)-NH-(CH<sub>2</sub>)<sub>n</sub>-NH<sub>2</sub>, wherein m and n are independently 1 to 10, wherein the CH<sub>3</sub> or aryl is optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, amino, alkylamino, dialkylamino, thiol, carboxy, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, aminoalkyl aminocarbonyl, and trialkylsilyl; or a pharmaceutically acceptable salt thereof.

13. The compound of any one of claims 1-12, wherein R<sup>3</sup> and R<sup>4</sup> are hydroxyl, or a pharmaceutically acceptable salt thereof.

14. The compound of any one of claims 1-13, wherein R<sup>5</sup> is hydrogen, or a pharmaceutically acceptable salt thereof.

15. The compound of claim 1, wherein R<sup>2</sup> is chloro, R<sup>1</sup> is 3-chlorobenzyl, 3-iodobenzyl, 3-bromobenzyl, 1-naphthylmethyl, 2,5-dimethoxybenzyl, 2-hydroxy-5-methoxybenzyl, or *trans*-2-phenyl-cyclopropyl, R<sup>3</sup> and R<sup>4</sup> are hydroxyl, and R<sup>5</sup> is hydrogen, or a pharmaceutically acceptable salt thereof.

16. A pharmaceutical composition comprising a compound of any one of claims 1-15 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

17. A compound of any one of claims 1-15 or a pharmaceutically acceptable salt thereof for use in treating a disease or condition selected from the group consisting of cancer, glaucoma, inflammatory diseases, asthma, stroke, myocardial infarction, allergic reactions, rhinitis, poison ivy induced responses, urticaria, scleroderma, arthritis, brain arteriole diameter constriction, bronchoconstriction, and myocardial ischemia.


18. Use of a compound of any one of claims 1-15 or a pharmaceutically acceptable salt thereof as a partial activator or selective inactivator of A<sub>3</sub> adenosine receptor.

19. The use of claim 18, wherein, in the compound of formula (I), R<sup>1</sup> is 3-bromobenzyl or 3-iodobenzyl, R<sup>2</sup> is halo, R<sup>3</sup> and R<sup>4</sup> are hydroxyl, and R<sup>5</sup> is hydrogen.

20. The use of claim 19, wherein, in the compound of formula (I), R<sup>2</sup> is chloro.

21. A compound or pharmaceutically acceptable salt of claim 1, wherein R<sup>1</sup> is 3-bromobenzyl or 3-iodobenzyl, R<sup>2</sup> is halo, R<sup>3</sup> and R<sup>4</sup> are hydroxyl, and R<sup>5</sup> is hydrogen for use in preventing or reducing ischemic damage to the heart in an animal in need thereof.

22. A radiolabeled compound of Formula I:



wherein

R<sup>1</sup> is selected from the group consisting of C<sub>6</sub>-C<sub>14</sub> aryl, C<sub>6</sub>-C<sub>14</sub> aryl C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>6</sub>-C<sub>14</sub> diaryl C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>6</sub>-C<sub>14</sub> aryl C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>6</sub>-C<sub>14</sub> aryl sulfonyl, heterocyclyl C<sub>1</sub>-C<sub>6</sub> alkyl, heterocyclyl, heteroaryl C<sub>1</sub>-C<sub>6</sub> alkyl, 4-[[[4-[[[(2-amino C<sub>1</sub>-C<sub>6</sub> alkyl) amino]-carbonyl]-C<sub>1</sub>-C<sub>6</sub> alkyl] anilino] carbonyl] C<sub>1</sub>-C<sub>6</sub> alkyl] C<sub>6</sub>-C<sub>14</sub> aryl, and C<sub>6</sub>-C<sub>14</sub> aryl C<sub>3</sub>-C<sub>8</sub> cycloalkyl, wherein the aryl or heterocyclyl portion of R<sup>1</sup> is substituted with one or more halogen atoms that are radioactive;

R<sup>2</sup> is selected from the group consisting of hydrogen, halo, amino, hydrazido, mercapto, C<sub>1</sub>-C<sub>20</sub> alkylamino, C<sub>6</sub>-C<sub>14</sub> aryl amino, C<sub>6</sub>-C<sub>14</sub> aryloxy, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>1</sub>-C<sub>20</sub> alkoxy, C<sub>1</sub>-C<sub>20</sub> thioalkoxy, pyridylthio, C<sub>7</sub>-C<sub>12</sub> cycloalkyl C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkyl C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkenyl C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>6</sub>-C<sub>14</sub> aryl C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>7</sub>-C<sub>12</sub> cycloalkyl C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkyl C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkenyl C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>6</sub>-C<sub>14</sub> aryl C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>2</sub>-C<sub>20</sub> alkynyl, carboxy alkyl C<sub>2</sub>-C<sub>20</sub> alkynyl, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-C(=O)-O-C<sub>1</sub>-C<sub>6</sub> alkyl, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-C(=O)-NH-(CH<sub>2</sub>)<sub>n</sub>-NH<sub>2</sub>, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-C<sub>1</sub>-C<sub>6</sub> alkyl, -C≡C-(CH<sub>2</sub>)<sub>m</sub>-aryl, wherein m and n are independently 1 to 10, C<sub>7</sub>-C<sub>12</sub> cycloalkyl C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>7</sub>-C<sub>12</sub>

bicycloalkyl C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>7</sub>-C<sub>12</sub> bicycloalkenyl C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>6</sub>-C<sub>14</sub> aryl C<sub>2</sub>-C<sub>20</sub> alkynyl, and the alkyl, cycloalkyl, or aryl portion of R<sup>2</sup> is optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, amino, alkylamino, dialkylamino, thiol, carboxy, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkyl aminocarbonyl, aminoalkyl aminocarbonyl, and trialkylsilyl;

R<sup>3</sup> and R<sup>4</sup> are independently selected from the group consisting of hydroxyl, amino, thiol, ureido, C<sub>1</sub>-C<sub>6</sub> alkyl carbonylamino, hydroxy C<sub>1</sub>-C<sub>6</sub> alkyl, and hydrazinyl; and

R<sup>5</sup> is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, heteroaryl, and C<sub>1</sub>-C<sub>6</sub> aminoalkyl;

or a pharmaceutically acceptable salt thereof.

23. The radiolabeled compound or salt of claim 22, wherein the halogen atom of R<sup>1</sup> is <sup>18</sup>F, <sup>76</sup>Br, or <sup>125</sup>I.

24. The radiolabeled compound or salt of claim 22 or 23, wherein R<sup>1</sup> is 3-bromobenzyl or 3-iodobenzyl, R<sup>2</sup> is halo, R<sup>3</sup> and R<sup>4</sup> are hydroxyl, and R<sup>5</sup> is hydrogen.

25. A method of imaging an A<sub>3</sub> adenosine receptor in a tissue or organ of an animal comprising administering an effective amount of a radiolabeled compound or salt of any one of claims 22-24 to the animal and obtaining an image of the organ or tissue of the animal.

26. The method of claim 25, wherein the image is obtained by a technique selected from the group consisting of Single Photon Emission Computerized Tomography (SPECT), Magnetic Resonance Spectroscopy (MRS), and Positron Emission Tomography (PET) and combinations thereof.

1/8




FIG. 1

2/8




FIG. 2




FIG. 3

4/8




FIG. 4

5/8




Figure 5

6/8




Figure 6

7/8

Figure 7a





Figure 7b



8/8





Figure 8



(I)