



(72) CHATTERJEE, DEB K., US

(72) SOLUS, JOSEPH, US

(72) YANG, SHUWEI, US

(71) LIFE TECHNOLOGIES, INC., US

(51) Int.Cl.⁶ C12Q 1/68, C12N 15/85, C12N 15/63, C12N 15/00, C12P 19/34,
C12N 9/12

(30) 1997/02/07 (60/037,393) US

(30) 1998/01/06 (60/070,562) US

(54) **POLYMERASES POUR L'ANALYSE ET LE TYPAGE DE
FRAGMENTS D'ACIDE NUCLEIQUE POLYMORPHES ET
LEURS UTILISATIONS**

(54) **POLYMERASES FOR ANALYZING OR TYPING
POLYMORPHIC NUCLEIC ACID FRAGMENTS AND USES
THEREOF**

(57) La présente invention concerne des procédés d'identification, d'analyse et de typage de fragments d'ADN polymorphes, notamment des fragments minisatellites, microsatellites ou d'ADN STR. L'invention concerne plus particulièrement des procédés utilisant des ADN polymérase, plus spécifiquement des ADN polymérase thermostables et encore plus spécifiquement des polymérase Thermotoga, ou des mutants ou des dérivés de celles-ci. Les molécules minisatellites, microsatellites ou d'ADN STR peuvent être amplifiées et analysées en vue de détecter des polymorphismes. L'invention concerne également des polymérase ayant réduit, sensiblement réduit ou éliminé leur capacité à ajouter des nucléotides 3' non matriciels à une molécule d'acide nucléique synthétisée. Selon l'invention, une telle réduction ou élimination peut être effectuée par la modification ou la mutation de la polymérase souhaitée.

(57) The present invention provides methods for use in identifying, analyzing and typing polymorphic DNA fragments, particularly minisatellite, microsatellite or STR DNA fragments. In particular, the invention provides methods using DNA polymerases, more particularly thermostable DNA polymerases, and most particularly Thermotoga polymerases or mutants or derivatives thereof, whereby minisatellite, microsatellite or STR DNA molecules may be amplified and analyzed for polymorphisms. The invention also relates to polymerases having reduced, substantially reduced or eliminated ability to add non-template 3' nucleotides to a synthesized nucleic acid molecule. In accordance with the invention, such reduction or elimination may be accomplished by modifying or mutating the desired polymerase.

CORRECTED
VERSION*CORRECTED
VERSION**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68, C12P 19/34, C12N 9/12, 15/00, 15/63, 15/85	A1	(11) International Publication Number: WO 98/35060 (43) International Publication Date: 13 August 1998 (13.08.98)
<p>(21) International Application Number: PCT/US98/02791</p> <p>(22) International Filing Date: 9 February 1998 (09.02.98)</p> <p>(30) Priority Data: 60/037,393 7 February 1997 (07.02.97) US 60/070,562 6 January 1998 (06.01.98) US</p> <p>(71) Applicant: LIFE TECHNOLOGIES, INC. [US/US]; 9800 Medical Center Drive, Rockville, MD 20850 (US).</p> <p>(72) Inventors: CHATTERJEE, Deb, K.; 6 Forest Ridge Court, North Potomac, MD 20878 (US). SOLUS, Joseph; 4 Ivyberry Court, Gaithersburg, MD 20879 (US). YANG, Shuwei; 15509 Moravia Court, Rockville, MD 20855 (US).</p> <p>(74) Agents: ESMOND, Robert, W. et al.; Sterne, Kessler, Goldstein & Fox P.L.L.C., Suite 600, 1100 New York Avenue, N.W., Washington, DC 20005-3934 (US).</p>	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i> <i>With an indication in relation to a deposited microorganism furnished under Rule 13bis separately from the description.</i> <i>Date of receipt by the International Bureau:</i> 27 March 1998 (27.03.1998)</p>	
(54) Title: POLYMERASES FOR ANALYZING OR TYPING POLYMORPHIC NUCLEIC ACID FRAGMENTS AND USES THEREOF		
(57) Abstract		
<p>The present invention provides methods for use in identifying, analyzing and typing polymorphic DNA fragments, particularly minisatellite, microsatellite or STR DNA fragments. In particular, the invention provides methods using DNA polymerases, more particularly thermostable DNA polymerases, and most particularly <i>Thermotoga</i> polymerases or mutants or derivatives thereof, whereby minisatellite, microsatellite or STR DNA molecules may be amplified and analyzed for polymorphisms. The invention also relates to polymerases having reduced, substantially reduced or eliminated ability to add non-template 3' nucleotides to a synthesized nucleic acid molecule. In accordance with the invention, such reduction or elimination may be accomplished by modifying or mutating the desired polymerase.</p>		

Polymerases for Analyzing or Typing Polymorphic Nucleic Acid Fragments and Uses Thereof

FIELD OF THE INVENTION

The present invention is in the field of molecular and cellular biology. The invention relates to compositions and methods for use in analyzing and typing polymorphic regions of DNA. More particularly, the invention is directed to compositions of polymerases (preferably DNA polymerases and most preferably thermostable DNA polymerases), and methods using these compositions, whereby polymorphic, minisatellite, microsatellite or STR DNA fragments may be amplified and analyzed. The compositions and methods of the present invention are useful in a variety of techniques employing DNA amplification and polymorphism analysis, including medical genetic, forensic, and plant breeding applications.

The present invention also relates to polymerases having reduced, substantially reduced or eliminated ability to add one or more non-templated nucleotides to the 3' terminus of a synthesized nucleic acid molecule. Preferably, the polymerases of the invention are thermostable or mesophilic polymerases. Specifically, the polymerases of the present invention (e.g., DNA or RNA polymerases) have been mutated or modified to reduce, substantially reduce or eliminate such activity (compared to the unmodified, unmutated, or wild type polymerase), thereby providing a polymerase which synthesizes nucleic acid molecules having little or no non-templated 3' terminal nucleotides. Such polymerases thus have enhanced or greater ability to produce a double stranded nucleic acid molecule having blunt ended termini which may facilitate cloning of such molecules. The present invention also relates to cloning and expression of the polymerases of the invention, to nucleic acid molecules containing the cloned genes, and to host cells which express said genes. The polymerases of the present invention may be used in DNA sequencing, amplification, nucleic acid synthesis, and polymorphism analysis.

The invention also relates to polymerases of the invention which have one or more additional mutations or modifications. Such mutations or modifications

-2-

include those which (1) substantially reduce 3'→5' exonuclease activity; and/or (2) substantially reduce 5'→3' exonuclease activity. The polymerases of the invention can have one or more of these properties. These polymerases may also be used in nucleic acid analysis including but not limited to DNA sequencing, amplification, nucleic acid synthesis, and polymorphism analysis.

BACKGROUND OF THE INVENTION

DNA Structure

The genetic framework (*i.e.*, the genome) of an organism is encoded in the double-stranded sequence of nucleotide bases in the deoxyribonucleic acid (DNA) which is contained in the somatic and germ cells of the organism. The genetic content of a particular segment of DNA, or gene, is only manifested upon production of the protein which the gene ultimately encodes. There are additional sequences in the genome that do not encode a protein (*i.e.*, "noncoding" regions) which may serve a structural, regulatory, or unknown function. Thus, the genome of an organism or cell is the complete collection of protein-encoding genes together with intervening noncoding DNA sequences. Importantly, each somatic cell of a multicellular organism contains the full complement of genomic DNA of the organism, except in cases of focal infections or cancers, where one or more xenogeneic DNA sequences may be inserted into the genomic DNA of specific cells and not into other, non-infected, cells in the organism.

Minisatellite and Microsatellite DNA

Interspersed throughout the genomic DNA of most eukaryotic organisms are short stretches of polymorphic repetitive nucleotide sequences known as "minisatellite DNA" sequences or fragments (Jeffreys, A.J., *et al.*, *Nature* 314:67-73 (1985)). These repeating sequences often appear in tandem and in variable numbers within the genome, and they are thus sometimes referred to as "short tandem repeats" ("STRs") or "variable numbers of tandem repeats" ("VNTRs")

-3-

(see U.S. Patent No. 5,075,217; Nakamura *et al.*, *Science* 235:1616-1622 (1987)). Typically, however, minisatellite repeat units are about 9 to 60 bases in length (Nakamura *et al.*, *Science* 235:1616-1622 (1987); Weber and May, *Am. J. Hum. Genet.* 44:388-396 (1989)) which are repeated in tandem about 20-50 times (Watson, J.D., *et al.*, eds., *Recombinant DNA*, 2nd ed., New York: Scientific American Books, p. 146 (1992)). Other short, simple sequences which are analogous to minisatellite DNAs, termed "microsatellite DNAs" (Litt, M., and Luty, J.A., *Am. J. Hum. Genet.* 44:397-401 (1989); Weber and May, *Am. J. Hum. Genet.* 44:388-396 (1989)), are usually about 1-6 bases in repeat unit length and thus give rise to monomeric (Economou, E.T., *et al.*, *Proc. Natl. Acad. Sci. USA* 87:2951-2954 (1990)), dimeric, trimeric, quatrameric, pentameric or hexameric repeat units (Litt, M., and Luty, J.A., *Am. J. Hum. Genet.* 44:397-401 (1989); Weber and May, *Am. J. Hum. Genet.* 44:388-396 (1989)). The most prevalent of these highly polymorphic microsatellite sequences in the human genome is the dinucleotide repeat (dC-dA)_n•(dG-dT)_n (where n is the number of repetitions in a given stretch of nucleotides), which is present in a copy number of about 50,000-100,000 (Tautz, D., and Renz, M., *Nucl. Acids Res.* 12:4127-4138 (1984); Dib, C., *et al.*, *Nature* 360:152-154 (1996)), although the existence of a variety of analogous repeat sequences in the genomes of evolutionarily diverse eukaryotes has been reported (Hamada, H., *et al.*, *Proc. Natl. Acad. Sci. USA* 79:6465-6469 (1982)).

The actual *in vivo* function of minisatellite and microsatellite sequences is unknown. However, because these tandemly repeated sequences are dispersed throughout the genome of most eukaryotes, exhibit size polymorphism, and are often heterozygous (Weber, J.L., *Genomics* 7:524-530 (1990)), they have been explored as potential genetic markers in assays attempting to distinguish closely related individuals, and in forensic and paternity testing (see, *e.g.*, U.S. Patent No. 5,075,217; Jeffreys, A.J., *et al.*, *Nature* 332:278-281 (1988)). The finding that mutations often are observed in microsatellite DNA regions in cancer cells (Loeb, L.A., *Cancer Res.* 54:5059-5063 (1994)), potentially linking genomic instability

-4-

to the carcinogenic process and providing useful genetic markers of cancer, lends additional significance to methods facilitating the rapid analysis and genotyping of polymorphisms in these genomic DNA regions.

Methods of Genotyping Minisatellite or STR DNA Sequences

5 To analyze minisatellite, microsatellite or STR DNA sequence polymorphisms, a variety of molecular biological techniques have been employed. These techniques include restriction fragment length polymorphism (RFLP) or "DNA fingerprinting" analysis (Wong, Z., et al., *Nucl. Acids Res.* 14:4605-4616 (1986); Wong, Z., et al., *Ann. Hum. Genet* 51:269-288 (1987); Jeffreys, A.J., et
10 *al.*, *Nature* 332:278-281 (1988); U.S. Patent Nos. 5,175,082; 5,413,908; 5,459,039; and 5,556,955). Far more commonly employed for STR genotyping than RFLP and hybridization, however, are amplification-based methods, such as those relying on the polymerase chain reaction (PCR) method invented by Mullis and colleagues (*see* U.S. Patent Nos. 4,683,195; 4,683,202; and 4,800,159).
15 These methods use "primer" sequences which are complementary to opposing regions flanking the polymorphic DNA sequence to be amplified from the sample of genomic DNA to be analyzed. These primers are added to the DNA target sample, along with excess deoxynucleotides and a DNA polymerase (*e.g.*, *Taq* polymerase; *see* below), and the primers bind to their target via base-specific
20 binding interactions (*i.e.*, adenine binds to thymine, cytosine to guanine). By repeatedly passing the reaction mixture through cycles of increasing and decreasing temperatures (to allow dissociation of the two DNA strands on the target sequence, synthesis of complementary copies of each strand by the polymerase, and re-annealing of the new complementary strands), the copy
25 number of the minisatellite or STR sequence of DNA may be rapidly increased, and detected by size separation methods such as gel electrophoresis.

PCR and related amplification approaches have been used in attempts to develop methods for typing and analyzing STRs or minisatellite regions. For example, PCR has been employed to analyze polymorphisms in microsatellite

-5-

sequences from different individuals, including (dC-dA)_n·(dG-dT)_n (Weber, J.L, and May, P.E., *Am. J. Hum. Genet.* 44:388-396 (1989); Weber, J. L., *Genomics* 7:524-530 (1990); U.S. Patent Nos. 5,075,217; 5,369,004; and 5,468,613). Similar methods have been applied to a variety of medical and forensic samples to perform DNA typing and to detect polymorphisms between individual samples (U.S. Patent Nos. 5,306,616; 5,364,759; 5,378,602; and 5,468,610).

In Vitro Use of DNA Polymerases

The above-described amplification-based techniques require the use of DNA polymerases, which catalyze the addition of deoxynucleoside triphosphate (dNTP) bases into the newly forming DNA strands. Together with other enzymes (*e.g.*, helicases, ligases and ATPases), the DNA polymerases ensure rapid and relatively faithful replication of DNA in preparation for proliferation *in vivo* in prokaryotes, eukaryotes and viruses.

DNA polymerases synthesize the formation of DNA molecules which are complementary to a DNA template. Upon hybridization of a primer to the single-stranded DNA template, polymerases synthesize DNA in the 5' to 3' direction, successively adding nucleotides to the 3'-hydroxyl group of the growing strand. Thus, in the presence of deoxyribonucleoside triphosphates (dNTPs) and a primer, a new DNA molecule, complementary to the single stranded DNA template, can be synthesized.

In addition to an activity which adds dNTPs to DNA in the 5' to 3' direction (*i.e.*, "polymerase" activity), many DNA polymerases also possess activities which remove dNTPs in the 5' to 3' and/or the 3' to 5' direction (*i.e.*, "exonuclease" activity). This dual activity of certain DNA polymerases is, however, a drawback for some *in vitro* applications. For example, the *in vitro* synthesis of an intact copy of a DNA fragment by the polymerase activity, an elongation process which proceeds in a 5' to 3' direction along the template DNA strand, is jeopardized by the exonuclease activities which may simultaneously or subsequently degrade the newly formed DNA.

Limitations of PCR-based Genotyping of Minisatellite, Microsatellite and STR DNA Sequences

Application of PCR-based methods to analysis of minisatellite or STR DNA sequences has a number of significant limitations. It has been shown, for example, that use of *Taq* and other thermostable DNA polymerases commonly employed in PCR and related automated amplification methods causes the accumulation of amplification products containing non-templated 3' terminal nucleotides (Clark, J.M., *et al.*, *J. Molec. Biol.* 198:123-127 (1987); Clark, J.M., *Nucl. Acids Res.* 16:9677-9686 (1988); Hu, G., *DNA Cell Biol.* 12:763-770 (1993)). That is, some of the newly synthesized DNA strands produced in each round of amplification have had an extra nucleotide added to their 3' termini, such that the newly synthesized strands may be longer by one base.

Non-templated nucleotide addition is a slow process compared to template-directed synthesis (Clark, J.M., *Nucl. Acids Res.* 16:9677-9686 (1988)), and its extent is sequence-dependent (Hu, G., *DNA Cell Biol.* 12:763-770 (1993); Brownstein, M.J., *et al.*, *BioTechniques* 20:1004-1010 (1996)). Consequently, the PCR product is often heterogeneous in regard to extra nucleotide addition depending upon the primers and the reaction conditions used by the investigator (Magnuson, V.L., *et al.*, *BioTechniques* 21:700-709 (1996)). Extra nucleotide addition, in combination with "stutter" due to slippage during PCR amplification (Levinson, G., and Gutman, G.A., *Molec. Biol. Evol.* 4:203-221 (1987); Schlotterer, C., and Tautz, D., *Nucl. Acids Res.* 20:211-215 (1992)), often results in complex DNA fragment patterns which are difficult to interpret, especially by automated methods. This can result in improper genotyping analysis, particularly if the percentage of non-templated nucleotide addition is between 30-70% of the PCR product (Smith, J.R., *et al.*, *Genome Res.* 5:312-317 (1995)).

Thus, a need currently exists for a rapid, automated method for identifying, analyzing and typing polymorphic DNA fragments, particularly minisatellite, microsatellite or STR DNA fragments, that will not result in the problematic results described above. The present invention provides such a method.

-7-

BRIEF SUMMARY OF THE INVENTION

The present invention satisfies these needs in the art by providing methods useful in the identification, analysis or typing of polymorphic DNA fragments, particularly minisatellite, microsatellite or STR DNA fragments, in samples of DNA from a cell, particularly a eukaryotic cell. Specifically, the invention provides a method of producing a population of amplified DNA molecules, for use in analyzing or typing a DNA molecule in a DNA sample isolated from a cell, preferably a eukaryotic cell. The method of the present invention comprises contacting a DNA sample with a DNA polymerase (preferably a thermostable DNA polymerases) reduced, substantially reduced or eliminated in the ability to add one or more non-templated nucleotides to the 3' terminus of a DNA molecule, amplifying a polymorphic DNA fragment, preferably a minisatellite, microsatellite or STR DNA fragment, within the DNA sample and analyzing the amplified polymorphic DNA fragment. In the method of the invention, the analysis step may comprise, for example, sizing or sequencing the amplified DNA molecule and optionally comparing the size and/or sequence of the amplified DNA molecule to a different DNA sample which has been amplified according to the invention. In preferred embodiments of the present invention, the thermostable DNA polymerase is a *Thermotoga* DNA polymerase, preferably a *Thermotoga* DNA polymerase substantially reduced in 3'-5' exonuclease activity, more preferably a *Tne* polymerase, a *Tma* polymerase, or a mutant or derivative thereof, and most preferably a mutant of *Tne* polymerase selected from the group consisting of *Tne* N' Δ 219, D323A; *Tne* N' Δ 283, D323A; *Tne* N' Δ 284, D323A; *Tne* N' Δ 193, D323A; *Tne* D137A, D323A; *Tne* D8A, D323A; *Tne* G195D, D323A; *Tne* G37D, D323A; *Tne* N' Δ 283; *Tne* D137A, D323A, R722K; *Tne* D137A, D323A, R722Y; *Tne* D137A, D323A, R722L; *Tne* D137A, D323A, R722H; *Tne* D137A, D323A, R722Q; *Tne* D137A, D323A, F730Y; *Tne* D137A, D323A, K726R; *Tne* D137A, D323A, K726H; *Tne* D137A, D323A, R722K, F730Y; *Tne* D137A, D323A, R722K, K726R; *Tne* D137A, D323A, R722K, K726H; *Tne* D137A, D323A,

-8-

R722H, F730Y; *Tne* D137A, D323A, R722H, K726R; *Tne* D137A, D323A, R722H, K726H; *Tne* D137A, D323A, R722Q, F730Y; *Tne* D137A, D323A, R722Q, K726R; *Tne* D137A, D323A, R722Q, K726H; *Tne* D137A, D323A, R722N, F730Y; *Tne* D137A, D323A, R722N, K726R; *Tne* D137A, D323A, R722N, K726H; *Tne* D137A, D323A, F730S; *Tne* N'Δ283, D323A, R722K/H/Q/N/Y/L; *Tne* N'Δ219, D323A, R722K; *Tne* N'Δ219, D323A, F730Y; *Tne* N'Δ219, D323A, K726R; *Tne* N'Δ219, D323A, K726H; *Tne* D137A, D323A, F730S, R722K/Y/Q/N/H/L, K726R/H; *Tne* D137A, D323A, F730T, R722K/Y/Q/N/H/L, K726R/H; *Tne* D137A, D323A, F730T; *Tne* F730S; *Tne* F730A; *Tne* K726R; *Tne* K726H; and *Tne* D137A, D323A, R722N. The present invention is particularly directed to the above methods wherein the eukaryotic cell is an plant cell or an animal cell, preferably a mammalian cell, more preferably a normal, diseased, cancerous, fetal or embryonic mammalian cell, and most preferably a human cell. The invention is also directed to the above methods, further comprising isolating the polymorphic, minisatellite, microsatellite or STR DNA fragment and inserting it into a vector, preferably an expression vector. By the present methods, the polymorphic or microsatellite DNA fragment may be amplified prior to being inserted into the vector.

The present invention also provides a method of determining the relationship between a first individual and a second individual, comprising contacting a DNA sample from the first and second individuals with a DNA polymerase (*e.g.* a thermostable DNA polymerase) reduced, substantially reduced or eliminated in the ability to add one or more non-templated nucleotides to the 3' terminus of a DNA molecule, amplifying one or more DNA molecules in the DNA sample to generate a collection of amplified polymorphic DNA fragments, separating the amplified DNA fragments by length, and comparing the pattern of amplified DNA fragments from the first individual to that of the second individual. This method also allows the identification of one or more unique polymorphic DNA fragments, particularly a minisatellite, microsatellite or STR DNA fragment, that is specifically present in only one of the two individuals. This method may

-9-

further comprise determining the sequence of the unique polymorphic, minisatellite, microsatellite or STR DNA fragment. In this embodiment of the present invention, the thermostable DNA polymerase may be a *Thermotoga* DNA polymerase, preferably a *Thermotoga* DNA polymerase substantially reduced in 3'-5' exonuclease activity, more preferably a *Tne* polymerase, a *Tma* polymerase, or a mutant or derivative thereof, and most preferably a mutant of *Tne* polymerase selected from the group consisting of *Tne* N'Δ219, D323A; *Tne* N'Δ283, D323A; *Tne* N'Δ284, D323A; *Tne* N'Δ193, D323A; *Tne* D137A, D323A; *Tne* D8A, D323A; *Tne* G195D, D323A; *Tne* G37D, D323A; *Tne* N'Δ283; *Tne* D137A, D323A, R722K; *Tne* D137A, D323A, R722Y; *Tne* D137A, D323A, R722L; *Tne* D137A, D323A, R722H; *Tne* D137A, D323A, R722Q; *Tne* D137A, D323A, F730Y; *Tne* D137A, D323A, K726R; *Tne* D137A, D323A, K726H; *Tne* D137A, D323A, R722K, F730Y; *Tne* D137A, D323A, R722K, K726R; *Tne* D137A, D323A, R722K, K726H; *Tne* D137A, D323A, R722H, F730Y; *Tne* D137A, D323A, R722H, K726R; *Tne* D137A, D323A, R722H, K726H; *Tne* D137A, D323A, R722Q, F730Y; *Tne* D137A, D323A, R722Q, K726R; *Tne* D137A, D323A, R722Q, K726H; *Tne* D137A, D323A, R722N, F730Y; *Tne* D137A, D323A, R722N, K726R; *Tne* D137A, D323A, R722N, K726H; *Tne* D137A, D323A, F730S; *Tne* N'Δ283, D323A, R722K/H/Q/N/Y/L; *Tne* N'Δ219, D323A, R722K; *Tne* N'Δ219, D323A, F730Y; *Tne* N'Δ219, D323A, K726R; *Tne* N'Δ219, D323A, K726H; *Tne* D137A, D323A, F730S, R722K/Y/Q/N/H/L, K726R/H; *Tne* D137A, D323A, F730T, R722K/Y/Q/N/H/L, K726R/H; *Tne* D137A, D323A, F730T; *Tne* F730S; *Tne* F730A; *Tne* K726R; *Tne* K726H; and *Tne* D137A, D323A, R722N. The present invention is particularly directed to the above methods wherein the first or second individual is an animal or a plant, and most preferably wherein the first or second individual is a human.

The present invention also provides isolated nucleic acid molecules encoding mutant *Tne* DNA polymerase proteins, wherein the mutant *Tne* DNA polymerase proteins have an amino acid sequence as set forth in any one of SEQ ID NOs: 4-10. The invention also provides mutant *Tne* DNA polymerase proteins

-10-

having an amino acid sequence as set forth in any one of SEQ ID NOs:4-10, most preferably a mutant *Tne* polymerase protein selected from the group consisting of *Tne* N'Δ283, D323A (SEQ ID NO:4); *Tne* N'Δ193, D323A (SEQ ID NO:5); *Tne* D137A, D323A (SEQ ID NO:6); *Tne* D8A, D323A (SEQ ID NO:7); *Tne* G195D, D323A (SEQ ID NO:8); *Tne* G37D, D323A (SEQ ID NO:9); and *Tne* N'Δ283 (SEQ ID NO:10). The invention also relates to nucleic acid molecules and the proteins encoded by such nucleic acid molecules for mutant *Tne* polymerases selected from the group consisting of *Tne* n'Δ283; *Tne* D137A, D323A, R722K; *Tne* D137A, D323A, R722Y; *Tne* D137A, D323A, R722L; *Tne* D137A, D323A, R722H; *Tne* D137A, D323A, R722Q; *Tne* D137A, D323A, F730Y; *Tne* D137A, D323A, K726R; *Tne* D137A, D323A, K726H; *Tne* D137A, D323A, R722K, F730Y; *Tne* D137A, D323A, R722K, K726R; *Tne* D137A, D323A, R722K, K726H; *Tne* D137A, D323A, R722H, F730Y; *Tne* D137A, D323A, R722H, K726R; *Tne* D137A, D323A, R722H, K726H; *Tne* D137A, D323A, R722Q, F730Y; *Tne* D137A, D323A, R722Q, K726R; *Tne* D137A, D323A, R722Q, K726H; *Tne* D137A, D323A, R722N, F730Y; *Tne* D137A, D323A, R722N, K726R; *Tne* D137A, D323A, R722N, K726H; *Tne* D137A, D323A, F730S; *Tne* N'Δ283, D323A, R722K/H/Q/N/Y/L; *Tne* N'Δ219, D323A, R722K; *Tne* N'Δ219, D323A, F730Y; *Tne* N'Δ219, D323A, K726R; *Tne* N'Δ219, D323A, K726H; *Tne* D137A, D323A, F730S, R722K/Y/Q/N/H/L, K726R/H; *Tne* D137A, D323A, F730T, R722K/Y/Q/N/H/L, K726R/H; *Tne* D137A, D323A, F730T; *Tne* F730S; *Tne* F730A; *Tne* K726R; *Tne* K726H; and *Tne* D137A, D323A, R722N. These mutations may be made to sequence ID NO:2 to produce the mutant polymerases having the indicated amino acid mutations (where, for example, "D137A" indicates that the Asp (D) residue at position 137 in SEQ ID NO:2 has been mutated to an Ala (A) residue, and, for example, "R722K/Y/Q/N/H/L" indicates that the Arg (R) residue at position 722 in SEQ ID NO:2 has been mutated to a Lys (K), Tyr (Y), Gln (Q), Asn (N), His (H) or Leu (L) residue).

-11-

The present invention also provides kits for the identification, analysis or typing of a polymorphic DNA fragment, particularly a minisatellite, microsatellite or STR DNA fragment, comprising a first container containing one or more DNA polymerases reduced, substantially reduced or eliminated in the ability to add non-templated 3' terminal nucleotides. Kits according to the invention may contain additional containers selected from the group consisting of a container containing one or more DNA primer molecules, a container containing one or more deoxynucleoside triphosphates needed to synthesize a DNA molecule complementary to the DNA template, and a container containing a buffer suitable for identifying, analyzing or typing a polymorphic DNA fragment by the methods of the invention. Any number of these components of the kit may be combined in a single or multiple containers to provide the kit of the invention. According to the invention, the DNA polymerase of the kit is preferably a *Thermotoga* DNA polymerase, more preferably a *Thermotoga* DNA polymerase substantially reduced in 3'-5' exonuclease activity, still more preferably a *Tne* polymerase, a *Tma* polymerase, or a mutant or derivative thereof, and most preferably a mutant of *Tne* polymerase selected from the group consisting of *Tne* N'Δ283; *Tne* D137A, D323A, R722K; *Tne* D137A, D323A, R722Y; *Tne* D137A, D323A, R722L; *Tne* D137A, D323A, R722H; *Tne* D137A, D323A, R722Q; *Tne* D137A, D323A, F730Y; *Tne* D137A, D323A, K726R; *Tne* D137A, D323A, K726H; *Tne* D137A, D323A, R722K, F730Y; *Tne* D137A, D323A, R722K, K726R; *Tne* D137A, D323A, R722K, K726H; *Tne* D137A, D323A, R722H, F730Y; *Tne* D137A, D323A, R722H, K726R; *Tne* D137A, D323A, R722H, K726H; *Tne* D137A, D323A, R722Q, F730Y; *Tne* D137A, D323A, R722Q, K726R; *Tne* D137A, D323A, R722Q, K726H; *Tne* D137A, D323A, R722N, F730Y; *Tne* D137A, D323A, R722N, K726R; *Tne* D137A, D323A, R722N, K726H; *Tne* D137A, D323A, F730S; *Tne* N'Δ283, D323A, R722K/H/Q/N/Y/L; *Tne* N'Δ219, D323A, R722K; *Tne* N'Δ219, D323A, F730Y; *Tne* N'Δ219, D323A, K726R; *Tne* N'Δ219, D323A, K726H; *Tne* D137A, D323A, F730S, R722K/Y/Q/N/H/L, K726R/H; *Tne* D137A, D323A, F730T, R722K/Y/Q/N/H/L, K726R/H; *Tne* D137A, D323A,

-12-

F730T; *Tne* F730S; *Tne* F730A; *Tne* K726R; *Tne* K726H; and *Tne* D137A, D323A, R722N.

5 The present invention also relates generally to mutated or modified polymerases (DNA or RNA polymerases) which have reduced, substantially reduced or eliminated ability to add one or more non-templated nucleotides to the 3' terminus of a synthesized nucleic acid molecule (compared to the corresponding wildtype, unmutated or unmodified polymerase). Preferably, such mutant or modified polymerases have substantially reduced ability to add one or more non-templated nucleotides to the 3' terminus of a synthesized nucleic acid molecule.

10 Such polymerases of the invention may be thermostable or mesophilic polymerases. Thus, the present invention relates to such mutated or modified polymerases and to kits containing such polymerases. The invention also relates to the use of such mutant or modified polymerases in a number of procedures including DNA sequencing, amplification reactions, nucleic acid synthesis, and

15 polymorphism analysis.

Mutant or modified polymerases of particular interest in the invention include *Taq* DNA polymerase, *Tne* DNA polymerase, *Tma* DNA polymerase, *Pfu* DNA polymerase, *Tfl* DNA polymerase, *Tth* DNA polymerase, *Tbr* DNA polymerase, *Pwo* DNA polymerase, *Bst* DNA polymerase, *Bca* DNA polymerase, VENT™ DNA polymerase, DEEP VENT™ DNA polymerase, T7 DNA polymerase, T5 DNA polymerase, DNA polymerase III, Klenow fragment DNA polymerase, Stoffel fragment DNA polymerase, and mutants, fragments or derivatives thereof. RNA polymerases of interest include T7, SP6, and T3 RNA polymerases and mutants, variants and derivatives thereof.

20

25 The present invention relates in particular to mutant PolII type DNA polymerases (preferably thermostable DNA polymerases) wherein one or more mutations have been made in the O-helix which reduces, substantially reduces or eliminates the ability of the enzyme to add one or more non-templated nucleotides to the 3' terminus of a synthesized nucleic acid molecule. The O-helix is defined as RXXXXKXXXXFXXXXYX (SEQ ID NO:11), wherein X may be any amino acid.

30

-13-

The preferred sites for mutation or modification to produce the polymerases of the invention are the R and/or F and/or K and/or Y positions in the O-helix, although other changes (or combinations thereof) within the O-helix may be made to make the desired polymerase. In this preferred aspect of the invention, R and/or F
5 and/or K and/or Y may be replaced with any other amino acid including Ala, Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr and Val.

In accordance with the invention, other functional changes (or combinations thereof) may be made to the polymerases having reduced ability to
10 add non-templated nucleotides to the 3' terminus of a synthesized nucleic acid molecule. For example, the polymerase may also be modified to reduce, substantially reduce or eliminate 5' exonuclease activity, and/or 3' exonuclease activity. Thus, the invention relates to mutant or modified DNA polymerases having reduced ability to add non-templated nucleotides which are modified in at
15 least one way selected from the group consisting of

(a) to reduce or eliminate the 3'-5' exonuclease activity of the polymerase;

(b) to reduce or eliminate the 5'-3' exonuclease activity of the polymerase;

20 Any one or a number of these mutations or modifications (or combinations thereof) may be made to provide the polymerases of the invention. Preferred polymerases of the invention, in addition to having reduced ability to add non-templated 3' nucleotides, also have reduced, substantially reduced or eliminated 3' exonuclease activity.

25 The present invention is also directed to nucleic acid molecules (preferably vectors) containing a gene encoding the mutant or modified polymerases of the present invention and to host cells containing such molecules. Any number of hosts may be used to express the gene of interest, including prokaryotic and eukaryotic cells. Preferably, prokaryotic cells are used to express the polymerases

-14-

of the invention. The preferred prokaryotic host according to the present invention is *E. coli*.

The invention also relates to a method of producing the polymerases of the invention, said method comprising:

- 5 (a) culturing the host cell comprising a gene encoding a polymerase of the invention;
- (b) expressing said gene; and
- (c) isolating said polymerase from said host cell.

10 The invention also relates to a method of synthesizing a nucleic acid molecule comprising:

- (a) mixing one or more nucleic acid templates (e.g. RNA or DNA) with one or more polymerases of the invention; and
- (b) incubating said mixture under conditions sufficient to synthesize nucleic acid molecules complementary to all or a portion of said templates. Such condition may include incubation with one or more deoxy- and/or dideoxyribonucleoside triphosphates. Such deoxy- and dideoxyribonucleoside triphosphates include dATP, dCTP, dGTP, dTTP, dITP, 7-deaza-dGTP, 7-deaza-dATP, dUTP, ddATP, ddCTP, ddGTP, ddITP, ddTTP, [α -S]dATP, [α -S]dTTP, [α -S]dGTP, and [α -S]dCTP. The synthesized nucleic acid molecules may in accordance with the invention be cloned into one or more vectors.
- 15
- 20

The invention also relates to a method of sequencing a DNA molecule, comprising:

- (a) hybridizing a primer to a first DNA molecule;
- (b) contacting said molecule of step (a) with deoxyribonucleoside triphosphates, one or more DNA polymerases of the invention, and one or more terminator nucleotides;
- 25
- (c) incubating the mixture of step (b) under conditions sufficient to synthesize a random population of DNA molecules complementary to said first DNA molecule, wherein said synthesized DNA molecules are shorter in length

-15-

than said first DNA molecule and wherein said synthesized DNA molecules comprise a terminator nucleotide at their 3' termini; and

(d) separating said synthesized DNA molecules by size so that at least a part of the nucleotide sequence of said first DNA molecule can be determined.

5 Such terminator nucleotides include but are not limited to dideoxynucleoside triphosphates such as ddTTP, ddATP, ddGTP, ddITP or ddCTP.

The invention also relates to a method for amplifying a double stranded DNA molecule, comprising:

10 (a) providing a first and second primer, wherein said first primer is complementary to a sequence at or near the 3'-termini of the first strand of said DNA molecule and said second primer is complementary to a sequence at or near the 3'-termini of the second strand of said DNA molecule;

15 (b) hybridizing said first primer to said first strand and said second primer to said second strand in the presence of one or more polymerases of the invention, under conditions such that a third DNA molecule complementary to said first strand and a fourth DNA molecule complementary to said second strand are synthesized;

(c) denaturing said first and third strands, and said second and fourth strands; and

20 (d) repeating steps (a) to (c) one or more times. The amplified double-stranded nucleic acid molecules produced by the method of the invention may be cloned into one or more vectors. Thus, the invention relates also to a method of cloning an amplified DNA molecule comprising:

25 (a) amplifying one or more DNA molecules with one or more polymerases of the invention; and

(b) ligating said amplified DNA molecules in one or more vectors.

The invention further relates to a method of cloning a nucleic acid molecule comprising:

30 (a) mixing a nucleic acid template (or one or more templates) with one or more polymerases of the invention;

-16-

(b) incubating said mixture under conditions sufficient to synthesize a nucleic acid molecule complementary to all or a portion of said template, thereby producing a double-stranded nucleic acid molecule (preferably a double-stranded DNA molecule); and

5 (c) ligating said double-stranded nucleic acid molecule into one or more vectors.

Preferably, the vectors used for ligating the amplified or synthesized double-stranded nucleic acid molecules have blunt ended termini and may be prepared by digesting a vector with any one or a number of restriction enzymes
10 known in the art which provide blunt end cleavage. Such restriction enzymes include *ScaI*, *SmaI*, *HpaI*, *HincII*, *HaeIII*, *AluI*, and the like.

The invention also relates to kits for sequencing, amplifying, synthesizing or cloning of nucleic acid molecules comprising one or more polymerases of the invention and one or more other components (or combinations thereof) selected
15 from the group consisting of

- (a) one or more dideoxynucleoside triphosphates;
- (b) one or more deoxynucleoside triphosphates;
- (c) one or more primers;
- (d) one or more suitable buffers; and
- 20 (e) one or more ligases.

Other preferred embodiments of the present invention will be apparent to one of ordinary skill in light of the following drawings and description of the invention, and of the claims.

BRIEF DESCRIPTION OF THE FIGURES

25 **FIGURE 1** shows the restriction map of the approximate DNA fragment which contains the *Tne* DNA polymerase gene in pSport 1 and pUC19. This figure also shows the region containing the O-helix homologous sequences.

-17-

FIGURE 2A schematically depicts the construction of plasmids pUC-Tne (3'→5') and pUC-Tne FY.

FIGURE 2B schematically depicts the construction of plasmids pTrcTne35 and pTrcTne FY.

5 **FIGURE 3** schematically depicts the construction of plasmid pTrcTne35 FY.

FIGURE 4 schematically depicts the construction of plasmids pTTQTne5FY and pTTQTne535FY.

FIGURE 5 depicts a plasmid containing the *Taq* DNA polymerase gene.

10 **FIGURE 6** depicts an autoradiogram showing of the ability of polymerase mutants to add non-templated 3' nucleotides.

FIGURE 7 is an autoradiogram of the product of PCR amplification of the upper and lower alleles of the CD4 locus, using primers corresponding to these alleles, demonstrating nontemplated nucleotide addition (n+1) by *Taq* DNA polymerase but not by *Tne* DNA polymerase.

15

FIGURE 8 is an autoradiogram of the product of PCR amplification of the upper and lower alleles of the D20S27 locus, using primers corresponding to these alleles, demonstrating nontemplated nucleotide addition (n+1) by *Taq* DNA polymerase but not by *Tne* DNA polymerase.

20 **FIGURE 9** is a composite of electropherogram gel scans of PCR amplifications at the D15S153 (Figures 9A and 9B) and D15S127 loci (Figures 9C and 9D), demonstrating nontemplated nucleotide addition (n+1) by *Taq* DNA

-18-

polymerase (Figures 9A and 9C) but not by *Tne* DNA polymerase (Figures 9B and 9D).

FIGURE 10A and B are composites of a electropherogram gel scan of PCR amplifications at D16S405 and D16S401 loci.

5 **FIGURE 11** is a composite of a electropherogram gel scan of PCR amplifications at D16S401 locus.

FIGURE 12A and B are composites of a electropherogram gel scan of PCR amplifications at D15S127 and D15S153 loci.

10 **FIGURE 13** is a composite of a electropherogram gel scan of PCR amplifications at D16S401 locus.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

15 In the description that follows, a number of terms used in recombinant DNA technology are extensively utilized. In order to provide a clearer and consistent understanding of the specification and claims, including the scope to be given such terms, the following definitions are provided.

20 **Polymorphic.** As is understood by one of ordinary skill in the art, a nucleic acid molecule is said to be "polymorphic" if it may exist in more than one form. For example, a nucleic acid molecule is said to be polymorphic if it may have more than one specific nucleotide sequence (such as degenerate nucleic acid molecules or genes that may each encode the same protein). More commonly, a nucleic acid molecule is said to be polymorphic if it displays size differences (*i.e.*, differences in length), particularly when comparisons of nucleic acid molecules

-19-

from different individuals are made. Of course, other definitions of the term "polymorphic" will be apparent to one of ordinary skill and are also encompassed within this definition.

Cloning vector. A plasmid, cosmid or phage DNA or other DNA molecule which is able to replicate autonomously in a host cell, and which is characterized by one or a small number of restriction endonuclease recognition sites at which such DNA sequences may be cut in a determinable fashion without loss of an essential biological function of the vector, and into which DNA may be spliced in order to bring about its replication and cloning. The cloning vector may further contain a marker suitable for use in the identification of cells transformed with the cloning vector. Markers, for example, are tetracycline resistance or ampicillin resistance.

Recombinant host. Any prokaryotic or eukaryotic microorganism which contains the desired cloned genes in an expression vector, cloning vector or any DNA molecule. The term "recombinant host" is also meant to include those host cells which have been genetically engineered to contain the desired gene on the host chromosome or genome.

Host. Any prokaryotic or eukaryotic microorganism that is the recipient of a replicable expression vector, cloning vector or any DNA molecule. The DNA molecule may contain, but is not limited to, a structural gene, a promoter and/or an origin of replication.

Promoter. A DNA sequence generally described as the 5' region of a gene, located proximal to the start codon. At the promoter region, transcription of an adjacent gene(s) is initiated.

Gene. A DNA sequence that contains information necessary for expression of a polypeptide or protein. It includes the promoter and the structural gene as well as other sequences involved in expression of the protein.

Structural gene. A DNA sequence that is transcribed into messenger RNA that is then translated into a sequence of amino acids characteristic of a specific polypeptide.

-20-

Operably linked. As used herein "operably linked" means that the promoter is positioned to control the initiation of expression of the polypeptide encoded by the structural gene.

5 **Expression.** Expression is the process by which a gene produces a polypeptide. It includes transcription of the gene into messenger RNA (mRNA) and the translation of such mRNA into polypeptide(s).

10 **Substantially Pure.** As used herein "substantially pure" means that the desired purified protein is essentially free from contaminating cellular contaminants which are associated with the desired protein in nature. Contaminating cellular components may include, but are not limited to, phosphatases, exonucleases, endonucleases or undesirable DNA polymerase enzymes.

15 **Primer.** As used herein "primer" refers to a single-stranded oligonucleotide that is extended by covalent bonding of nucleotide monomers during amplification or polymerization of a nucleic acid molecule. Minisatellite primers used for the amplification of minisatellite dimer, trimer, tetramer, etc., sequences are well-known in the art.

20 **Template.** The term "template" as used herein refers to a double-stranded or single-stranded nucleic acid molecule which is to be amplified, synthesized or sequenced. In the case of a double-stranded DNA molecule, denaturation of its strands to form a first and a second strand is performed before these molecules may be amplified, synthesized or sequenced. A primer, complementary to a portion of a template is hybridized under appropriate conditions and the polymerase of the invention may then synthesize a molecule complementary to said template or a portion thereof. The newly synthesized molecule, according to
25 the invention, may be equal or shorter in length than the original template. Mismatch incorporation or strand slippage during the synthesis or extension of the newly synthesized molecule may result in one or a number of mismatched base pairs. Thus, the synthesized molecule need not be exactly complementary to the template.

-21-

Incorporating. The term "incorporating" as used herein means becoming a part of a nucleic acid (*e.g.*, DNA) molecule or primer.

Amplification. As used herein "amplification" refers to any *in vitro* method for increasing the number of copies of a nucleotide sequence with the use of a DNA polymerase. Nucleic acid amplification results in the incorporation of nucleotides into a DNA molecule or primer thereby forming a new DNA molecule complementary to a DNA template. The formed DNA molecule and its template can be used as templates to synthesize additional DNA molecules. As used herein, one amplification reaction may consist of many rounds of DNA replication. DNA amplification reactions include, for example, polymerase chain reactions (PCR). One PCR reaction may consist of 5 to 100 "cycles" of denaturation and synthesis of a DNA molecule.

Oligonucleotide. "Oligonucleotide" refers to a synthetic or natural molecule comprising a covalently linked sequence of nucleotides which are joined by a phosphodiester bond between the 3' position of the pentose of one nucleotide and the 5' position of the pentose of the adjacent nucleotide.

Nucleotide. As used herein "nucleotide" refers to a base-sugar-phosphate combination. Nucleotides are monomeric units of a nucleic acid sequence (DNA and RNA). The term nucleotide includes deoxyribonucleoside triphosphates such as dATP, dCTP, dITP, dUTP, dGTP, dTTP, or derivatives thereof. Such derivatives include, for example, [α S]dATP, 7-deaza-dGTP and 7-deaza-dATP. The term nucleotide as used herein also refers to dideoxyribonucleoside triphosphates (ddNTPs) and their derivatives. Illustrated examples of dideoxyribonucleoside triphosphates include, but are not limited to, ddATP, ddCTP, ddGTP, ddITP, and ddTTP. According to the present invention, a "nucleotide" may be unlabeled or detectably labeled by well known techniques. Detectable labels include, for example, radioactive isotopes, fluorescent labels, chemiluminescent labels, bioluminescent labels and enzyme labels.

Thermostable. As used herein "thermostable" refers to a polymerase which is resistant to inactivation by heat. DNA polymerases synthesize the

-22-

formation of a DNA molecule complementary to a single-stranded DNA template by extending a primer in the 5'-to-3' direction. This activity for mesophilic DNA polymerases may be inactivated by heat treatment. For example, T5 DNA polymerase activity is totally inactivated by exposing the enzyme to a temperature of 90°C for 30 seconds. As used herein, a thermostable polymerase activity is more resistant to heat inactivation than a mesophilic polymerase. However, a thermostable polymerase does not mean to refer to an enzyme which is totally resistant to heat inactivation and thus heat treatment may reduce the polymerase activity to some extent. A thermostable polymerase typically will also have a higher optimum temperature than mesophilic polymerases.

Hybridization. The terms "hybridization" and "hybridizing" refers to the pairing of two complementary single-stranded nucleic acid molecules (RNA and/or DNA) to give a double-stranded molecule. As used herein, two nucleic acid molecules may be hybridized, although the base pairing is not completely complementary. Accordingly, mismatched bases do not prevent hybridization of two nucleic acid molecules provided that appropriate conditions, well known in the art, are used. In the present invention, the term "hybridization" refers particularly to hybridization of an oligonucleotide to a template molecule.

3'-5' Exonuclease Activity. "3'-5' exonuclease activity" is an enzymatic activity well known to the art. This activity is often associated with DNA polymerases, and is thought to be involved in a DNA replication "editing" or correction mechanism.

A "DNA polymerase substantially reduced in 3'-5' exonuclease activity" (which may also be represented as "3'exo-") is defined herein as either (1) a mutated DNA polymerase that has about or less than 10%, or preferably about or less than 1%, of the 3'-5' exonuclease activity of the corresponding unmutated, wildtype enzyme, or (2) a DNA polymerase having a 3'-5' exonuclease specific activity which is less than about 1 unit/mg protein, or preferably about or less than 0.1 units/mg protein. A unit of activity of 3'-5' exonuclease is defined as the amount of activity that solubilizes 10 nmoles of substrate ends in 60 min. at 37°C,

-23-

assayed as described in the "BRL 1989 Catalogue & Reference Guide", page 5, with *Hha*I fragments of *lambda* DNA 3'-end labeled with [³H]dTTP by terminal deoxynucleotidyl transferase (TdT). Protein is measured by the method of Bradford, *Anal. Biochem.* 72:248 (1976). As a means of comparison, natural, wildtype T5-DNA polymerase (DNAP) or T5-DNAP encoded by pTTQ19-T5-2 has a specific activity of about 10 units/mg protein while the DNA polymerase encoded by pTTQ19-T5-2(exo⁻) (U.S. Patent No. 5,270,179) has a specific activity of about 0.0001 units/mg protein, or 0.001% of the specific activity of the unmodified enzyme, a 10⁵-fold reduction.

5
10 **5'-3' Exonuclease Activity.** "5'-3' exonuclease activity" is also an enzymatic activity well known in the art. This activity is often associated with DNA polymerases, such as *E. coli* PolI and PolIII.

15 A "DNA polymerase substantially reduced in 5'-3' exonuclease activity" (which may also be represented as "5'exo-") is defined herein as either (1) a mutated DNA polymerase that has about or less than 10%, or preferably about or less than 1%, of the 5'-3' exonuclease activity of the corresponding unmutated, wildtype enzyme, or (2) a DNA polymerase having 5'-3' exonuclease specific activity which is less than about 1 unit/mg protein, or preferably about or less than 0.1 units/mg protein.

20 Both of the 3'-5' and 5'-3' exonuclease activities can be observed on sequencing gels. Active 5'-3' exonuclease activity will produce nonspecific ladders in a sequencing gel by removing nucleotides from the 5'-end of the growing primers. 3'-5' exonuclease activity can be measured by following the degradation of radiolabeled primers in a sequencing gel. Thus, the relative amounts of these activities, e.g. by comparing wildtype and mutant polymerases, can be determined with no more than routine experimentation.

25 **Minisatellite DNA.** As used herein, the term "minisatellite DNA" refers to a DNA fragment comprising a short stretch of tandemly repetitive nucleotide sequence. *In vivo*, minisatellite DNA fragments are found interspersed throughout the genomes of most eukaryotic organisms thus far examined. These repeating

30

-24-

sequences appear in tandem and often in variable numbers within the genome; thus, the terms "short tandem repeats" ("STRs") or "variable numbers of tandem repeats" ("VNTRs") may be used synonymously when referring to these regions. Minisatellite DNA fragments are typically about 9 bases to about 60 bases in length and are repeated about 20-50 times at a typical locus in a eukaryotic genome.

5

10

Microsatellite DNA. As used herein, the term "microsatellite DNA" refers to DNA fragments which are typically of a repeat unit size of about 1-6 bases in length. The most prevalent of these microsatellite DNA fragments in the human genome is the dinucleotide repeat $(dC-dA)_n \bullet (dG-dT)_n$ (where n is the number of repetitions in a given stretch of nucleotides). The terms "STRs" and "VNTRs" may also be used synonymously to denote these structures.

15

20

25

30

Non-templated 3' Terminal Nucleotide Addition. As used herein, the term "non-templated 3' terminal nucleotide addition" or "extranucleotide addition" means the propensity of an enzyme such as a DNA polymerase to incorporate one or more additional nucleotides, which are not found in the template strand at the 3' terminus of a newly synthesized nucleic acid molecule in a synthesis or amplification reaction, such as PCR. As a result of non-templated 3' terminal nucleotide addition, the synthesized or amplification products (*i.e.*, the newly synthesized DNA strand) will be longer by one or more nucleotides than is the template, in such a fashion that if the template is " n " nucleotides in length, the synthesis or amplification products will be " $n+1$," " $n+2$," " $n+3$," etc., nucleotides in length. A "polymerase substantially reduced in the ability to add one or more non-templated nucleotides to the 3' terminus of a nucleic acid molecule" is defined herein as a DNA polymerase, which when it has no 3' exonuclease activity or has substantially reduced 3' exonuclease activity, it will produce a collection of amplification products in which less than about 50%, preferably less than about 30%, more preferably less than about 20%, still more preferably less than about 10%, still more preferably less than about 5%, and most preferably less than about 1% of the amplification products contain one or more non-templated nucleotides

-25-

at their 3' termini compared to amplification products produced by *Taq* DNA polymerase assayed under the same conditions. Preferably, the conditions used for assaying 3' non-templated nucleotide addition is performed such that less than 100% of the amplification products of *Taq* DNA polymerase exhibits 3' non-templated nucleotide addition. Included in this definition are those polymerases that satisfy this definition for any primer set used. Thus, if the use of any primer set provides the indicated reduction of 3' non-templated nucleotide addition, the polymerase is said to be substantially reduced in the ability to add one or more non-templated nucleotides to the 3' terminus of a nucleic acid molecule.

When referring to polymerases which have been mutated or modified to reduce or eliminate 3' non-templated nucleotide addition, the mutated or modified polymerase is said to be "reduced in the ability to add one or more non-templated nucleotides to the 3' terminus of a nucleic acid molecule" when the polymerase has a lower or reduced or eliminated ability to add non-templated 3' nucleotides compared to the corresponding unmutated, unmodified or wildtype polymerase. For example, when testing the affect of a point mutation in the O-helix of a polymerase on non-templated nucleotide addition, the polymerase unmodified in the same position of the O-helix is preferably used for comparison purposes. Such mutated or modified polymerases are said to "substantially reduced in the ability to add one or more non-templated nucleotides to the 3' terminus of a nucleic acid molecule" if the mutated or modified polymerase has less than about 50%, preferably less than about 30%, more preferably less than about 20%, still more preferably less than about 10%, still more preferably less than about 5%, and most preferably less than about 1% of the activity for adding non-templated 3' terminal nucleotides compared to the corresponding unmutated, unmodified or wildtype polymerase. Preferably, the conditions used for assaying 3' non-templated nucleotide addition is performed such that less than 100% of the amplification products produced by the unmutated, unmodified or wildtype polymerase control exhibits 3' non-templated nucleotide addition. Included in this definition are those

-26-

mutant or modified polymerases that satisfy this definition for any primer set tested.

5 The ability of a polymerase to add a non-templated 3' terminal nucleotide to the growing strand may be assessed by a variety of techniques, most preferably by gel electrophoresis of the synthesized or amplification products for a direct size comparison and by comparison to markers of known size (see Figures 6-13).

Other terms used in the fields of recombinant DNA technology and molecular and cell biology as used herein will be generally understood by one of ordinary skill in the applicable arts.

10 Sources of Polymerases

The methods of the present invention rely on the use of polymerases (thermostable or mesophilic DNA or RNA polymerases) reduced, substantially reduced or eliminated in the ability to add one or more non-templated 3' terminal nucleotide to a growing nucleic acid strand. These thermostable DNA polymerases may be obtained from any strain of any thermophilic microorganism, including but not limited to strains of *Thermus aquaticus* (*Taq* polymerase; see U.S. Patent Nos. 4,889,818 and 4,965,188), *Thermus thermophilus* (*Tth* polymerase), *Thermococcus litoralis* (*Tli* or VENT™ polymerase), *Pyrococcus furiosus* (*Pfu* or DEEPVENT™ polymerase), *Pyrococcus woosii* (*Pwo* polymerase) and other *Pyrococcus* species, *Bacillus sterothermophilus* (*Bst* polymerase), *Sulfolobus acidocaldarius* (*Sac* polymerase), *Thermoplasma acidophilum* (*Tac* polymerase), *Bacillus caldophilus* (*Bca* polymerase), *Thermus flavus* (*Tfl/Tub* polymerase), *Thermus ruber* (*Tru* polymerase), *Thermus brockianus* (DYNAZYME™ polymerase), *Thermotoga neapolitana* (*Tne* polymerase; see WO 96/10640 and WO96/41014), *Thermotoga maritima* (*Tma* polymerase; see U. S. Patent No. 5,374,553) and other species of the *Thermotoga* genus (*Tsp* polymerase) and *Methanobacterium thermoautotrophicum* (*Mth* polymerase). Mesophilic DNA polymerases of interest in the invention include but are not limited to T7 DNA polymerases, T5 DNA polymerase, DNA polymerase

-27-

III, Klenow fragment DNA polymerase and mutants, fragments or derivatives thereof. RNA polymerases such as T3, T5, SP6 and mutants, variants and derivatives thereof may also be used in accordance with the invention. Polymerases having reduced or substantially reduced ability to add a non-templated 3' nucleotide to a growing nucleic acid strand may be wildtype polymerases, or may be made by mutating such wildtype polymerases by standard techniques (for example, by generating point mutations, insertions, deletions, etc., in the wildtype gene or protein). Polymerases that are reduced or substantially reduced in the ability to add a non-templated 3' nucleotide to a growing strand may be identified by assaying the synthesized products (e.g. PCR products) formed by such enzymes, as is well-known in the art and as generally described below in the Examples.

The nucleic acid polymerases used in the present invention may be mesophilic or thermophilic, and are preferably thermophilic. Preferred mesophilic DNA polymerases include T7 DNA polymerase, T5 DNA polymerase, Klenow fragment DNA polymerase, DNA polymerase III and the like. Preferred thermostable DNA polymerases that may be used in the methods of the invention include *Taq*, *Tne*, *Tma*, *Pfu*, *Tfl*, *Tth*, Stoffel fragment, VENT™ and DEEPVENT™ DNA polymerases, and mutants, variants and derivatives thereof (U.S. Patent No. 5,436,149; U.S. Patent 4,889,818; U.S. Patent 4,965,188; U.S. Patent 5,079,352; U.S. Patent 5,614,365; U.S. Patent 5,374,553; U.S. Patent 5,270,179; U.S. Patent 5,047,342; U.S. Patent No. 5,512,462; WO 92/06188; WO 92/06200; WO 96/10640; Barnes, W.M., *Gene* 112:29-35 (1992); Lawyer, F.C., *et al.*, *PCR Meth. Appl.* 2:275-287 (1993); Flaman, J.-M., *et al.*, *Nucl. Acids Res.* 22(15):3259-3260 (1994)). For amplification of long nucleic acid molecules (e.g., nucleic acid molecules longer than about 3-5 Kb in length), at least two DNA polymerases (one substantially lacking 3' exonuclease activity and the other having 3' exonuclease activity) are typically used. See U.S. Patent No. 5,436,149; U.S. Patent No. 5,512,462; Farnes, W.M., *Gene* 112:29-35 (1992); and copending U.S. Patent Application No. 08/689,814, filed February 14, 1997, the disclosures

-28-

of which are incorporated herein in their entireties. Examples of DNA polymerases substantially lacking in 3' exonuclease activity include, but are not limited to, *Taq*, *Tne(exo⁻)*, *Tma(exo⁻)*, *Pfu (exo⁻)*, *Pwo(exo⁻)* and *Tth* DNA polymerases, and mutants, variants and derivatives thereof.

5 Polypeptides having nucleic acid polymerase activity are preferably used in the present methods at a final concentration in solution of about 0.1-200 units per milliliter, about 0.1-50 units per milliliter, about 0.1-40 units per milliliter, about 0.1-3.6 units per milliliter, about 0.1-34 units per milliliter, about 0.1-32 units per milliliter, about 0.1-30 units per milliliter, or about 0.1-20 units per
10 milliliter, and most preferably at a concentration of about 20-40 units per milliliter. Of course, other suitable concentrations of nucleic acid polymerases suitable for use in the invention will be apparent to one of ordinary skill in the art.

In a preferred aspect of the invention, polymerases of the invention and preferably the mutant or modified polymerases of the invention are made by
15 recombinant techniques. A number of cloned polymerase genes are available or may be obtained using standard recombinant techniques.

To clone a gene encoding a polymerase, which may be modified in accordance with the invention, isolated DNA which contains the polymerase gene is used to construct a recombinant library in a vector. Any vector, well known in
20 the art, can be used to clone the DNA polymerase of interest. However, the vector used must be compatible with the host in which the recombinant DNA library will be transformed.

Prokaryotic vectors for constructing the plasmid library include plasmids such as those capable of replication in *E. coli* such as, for example, pBR322, ColE1, pSC101, pUC-vectors (pUC18, pUC19, etc.: In: *Molecular Cloning, A
25 Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1982); and Sambrook *et al.*, In: *Molecular Cloning A Laboratory Manual* (2d ed.) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989)). Bacillus plasmids include pC194, pC221, pC217, etc. Such
30 plasmids are disclosed by Glyczan, T. In: *The Molecular Biology Bacilli*,

-29-

Academic Press, York (1982), 307-329. Suitable *Streptomyces* plasmids include pIJ101 (Kendall *et al.*, *J. Bacteriol* 169:4177-4183 (1987)). *Pseudomonas* plasmids are reviewed by John *et al.*, (*Rad. Insec. Dis.* 8:693-704 (1986)), and Igaki, (*Jpn. J. Bacteriol.* 33:729-742 (1978)). Broad-host range plasmids or cosmids, such as pCP13 (Darzins and Chakrababary, *J. Bacteriol.* 159:9-18, 1984) can also be used for the present invention. The preferred vectors for cloning the genes of the present invention are prokaryotic vectors. Preferably, pCP13 and pUC vectors are used to clone the genes of the present invention.

The preferred host for cloning the polymerase genes of interest is a prokaryotic host. The most preferred prokaryotic host is *E. coli*. However, the desired polymerase genes of the present invention may be cloned in other prokaryotic hosts including, but not limited to, *Escherichia*, *Bacillus*, *Streptomyces*, *Pseudomonas*, *Salmonella*, *Serratia*, and *Proteus*. Bacterial hosts of particular interest include *E. coli* DH10B, which may be obtained from Life Technologies, Inc. (LTI) (Rockville, MD).

Eukaryotic hosts for cloning and expression of the polymerases of interest include yeast, fungi, and mammalian cells. Expression of the desired polymerase in such eukaryotic cells may require the use of eukaryotic regulatory regions which include eukaryotic promoters. Cloning and expressing the polymerase gene in eukaryotic cells may be accomplished by well known techniques using well known eukaryotic vector systems.

Once a DNA library has been constructed in a particular vector, an appropriate host is transformed by well known techniques. Transformed colonies are preferably plated at a density of approximately 200-300 colonies per petri dish. For thermostable polymerase selection, colonies are then screened for the expression of a heat stable DNA polymerase by transferring transformed *E. coli* colonies to nitrocellulose membranes. After the transferred cells are grown on nitrocellulose (approximately 12 hours), the cells are lysed by standard techniques, and the membranes are then treated at 95°C for 5 minutes to inactivate the endogenous *E. coli* enzyme. Other temperatures may be used to inactivate the

-30-

host polymerases depending on the host used and the temperature stability of the polymerase to be cloned. Stable polymerase activity is then detected by assaying for the presence of polymerase activity using well known techniques (*see, e.g., Sagner et al., Gene 97:119-123 (1991)*, which is hereby incorporated by reference in its entirety). The gene encoding a polymerase of the present invention can be cloned using the procedure described by Sanger *et al., supra*. Other techniques for selecting cloned polymerases in accordance with the present invention will be well-known to those of ordinary skill in the art.

Modifications or Mutations of Polymerases

In accordance with the invention, the nucleotide binding domain of the polymerase of interest is modified or mutated in such a way as to produce a mutated or modified polymerase having reduced, substantially reduced or eliminated activity for adding non-templated 3' nucleotides. The O-helix region typically defines the nucleotide binding domain of DNA polymerases. The O-helix may be defined as RXXXKXXXFXXXYYX (SEQ ID NO:11), wherein X may be any amino acid. One or more mutations or combinations of mutations may be made in the O-helix of any polymerase in order to reduce or eliminate non-templated 3' nucleotide addition in accordance with the invention. Such mutations include point mutation, frame-shift mutations, deletions and insertions. Preferably, one or more point mutations, resulting in one or more amino acid substitutions, are used to produce polymerases having such activity. Such mutations may be made by a number of methods that will be familiar to one of ordinary skill, including but not limited to site-directed mutagenesis. In a preferred aspect of the invention, one or more mutations at positions R, K, F, and/or Y in the polymerase O-helix may be made to produce a polymerase having the desired activity. Most preferably, one or more mutations at position R and/or F and/or K and/or Y within the O-helix results in polymerases having reduced, substantially reduced or eliminated activity for adding non-templated 3' nucleotides. In the preferred aspect, amino acid substitutions are made at position R and/or F and/or K and/or

-31-

Y (or combinations thereof). Thus, R (Arg) and/or F (Phe) and/or K (Lys) may be substituted with any other amino acid including Ala, Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr, and Val. Preferably, R (Arg) is substituted with amino acids Lys, Tyr, Leu, His, Gln, Met, or Asn. F (Phe) is preferably substituted with amino acids Tyr, Ala, Leu, Thr, and Ser. K (Lys) is preferably substituted with amino acids Arg, Tyr, Leu, His, Gln, Met or Asn, and more preferably with Arg or His. Y (Tyr) is preferably substituted with amino acids Lys, Arg, Ala, Thr, Phe, Leu, His, Gln, Met, or Asn. Positions corresponding to R, K, F and Y for RNA polymerases may also be determined by comparing nucleotide and/or amino acid sequences with those of DNA polymerases, to determine homologies therebetween. Corresponding mutations or modification may then be made to produce the desired result in any RNA polymerase.

The O-helix has been identified and defined for a number of polymerases and may be readily identified for other polymerases by one with skill in the art. Thus, given the defined O-helix region and the methods and assays described herein, one with skill in the art can make one or a number of modifications which would result in polymerases having reduced, substantially reduced or eliminated activity for adding non-templated 3' nucleotides. Accordingly, the invention relates to methods for producing such polymerases having modifications in the O-helix domain resulting in reduction, substantial reduction or elimination of activity for adding non-templated 3' nucleotides, methods for producing nucleic acid molecules encoding such polymerases, and polymerases and nucleic acid molecules produced by such methods.

The following table illustrates identified O-helix regions for known polymerases.

-32-

Polymerase	O-Helix Region	SEO ID NO.
Poll	754 RRS AKAINFG LIYG	12
<i>Taq</i>	659 RRA AKTINFG VLYG	13
T7	518 RDN AKTFIYG FLYG	14
<i>Tne</i>	722 RRV GK MVNFS IY G	15
T5	588 RQA AKAITFG ILYG	16
<i>Tma</i>	722 RRAGK MVNFS IY G	17

Thus, in accordance with a preferred aspect of the invention, corresponding mutations in the R and/or F and/or K positions of the O-helix can be made for the following enzymes based on the tables below.

Polymerase	Mutation Position
Poll	Arg ⁷⁵⁴
T5	Arg ⁵⁸⁸
T7	Arg ⁵¹⁸
<i>Taq</i>	Arg ⁶⁵⁹
<i>Tne</i>	Arg ⁷²²
<i>Tma</i>	Arg ⁷²²
<i>Bca</i>	Arg ⁷⁰⁵
<i>Bst</i>	Arg ⁷⁰²
<i>Tth</i>	Arg ⁶⁶¹

Polymerase	Mutation Position
Poll	Phe ⁷⁶²
T5	Phe ⁵⁹⁶
T7	Phe ⁵²⁸
<i>Taq</i>	Phe ⁶⁶⁷
<i>Tne</i>	Phe ⁷³⁰
<i>Tma</i>	Phe ⁷³⁰
<i>Bca</i>	Phe ⁷¹³
<i>Bst</i>	Phe ⁷¹⁰
<i>Tth</i>	Phe ⁶⁶⁹

-33-

Polymerase	Mutation Position
PoiI	Lys ⁷⁵⁸
T5	Lys ⁵⁹²
T7	Lys ⁵²²
5 <i>Taq</i>	Lys ⁶⁶³
<i>Tne</i>	Lys ⁷²⁶
<i>Tma</i>	Lys ⁷²⁶
<i>Bca</i>	Lys ⁷⁰⁷
10 <i>Bst</i>	Lys ⁷⁰⁶
<i>Tth</i>	Lys ⁶⁶⁵

The mutation position of Arg⁷⁰⁵ for *Bca* is based on the sequence information in GenBank. It should be noted, however, that according to the sequence described by Vemori *et al. J. Biochem. (Japan) 113*:401-410 (1993), the position of Arg in *Bca* is 703.

15 **Additional Modifications or Mutations of Polymerases**

In accordance with the invention, in addition to the mutations or modifications described above, one or more additional mutations or modifications (or combinations thereof) may be made to the polymerases of interest. Mutations or modifications of particular interest include those modifications of mutations which (1) reduce or eliminate 3' to 5' exonuclease activity; and (2) reduce or eliminate 5' to 3' exonuclease activity.

If the DNA polymerase has 3'-to-5' exonuclease activity, this activity may be reduced, substantially reduced, or eliminated by mutating the polymerase gene. Such mutations include point mutations, frame shift mutations, deletions and insertions. Preferably, the region of the gene encoding the 3'-to-5' exonuclease activity is mutated or deleted using techniques well known in the art (Sambrook *et al.*, (1989) in: *Molecular Cloning, A Laboratory Manual (2nd Ed.)*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

-34-

The 3'-to-5' exonuclease activity can be reduced or impaired by creating site specific mutants within the 3'→5' exonuclease domain. *See infra*. In a specific embodiment of the invention Asp³²³ of *Tne* DNA polymerase is changed to any amino acid, preferably to Ala³²³ to substantially reduce 3'→5' exonuclease activity.

5 In another specific embodiment of the invention, Asp³²³ of *Tma* may be changed to any other amino acid, preferably to Ala to substantially reduce 3'→5' exonuclease activity. The following represents a domain of interest for a number of polymerases for preparing 3'→5' exonuclease mutants.

	<i>Tne</i>	318	PSFALDLETSS	328	(SEQ ID NO:18)
10	Pol I	350	PVFAFDTETDS	360	(SEQ ID NO:19)
	T5	159	GPVAFDSETSA	169	(SEQ ID NO:20)
	T7	1	MIVSDIEANA	10	(SEQ ID NO:21)

Mutations, such as insertions, deletions and substitutions within the above domain can result in substantially reduced 3'→5' exonuclease activity. By way of

15 example, Asp³⁵⁵ (PolI), Asp¹⁶⁴ (T5), and Asp⁵ (T7) may be substituted with any amino acid to substantially reduce 3'→5' exonuclease activity. For example, Asp at these positions may be substituted with Ala.

The 5'→3' exonuclease activity of the polymerases can be reduced, substantially reduced or eliminated by mutating the polymerase gene or by deleting

20 the 5' to 3' exonuclease domain. Such mutations include point mutations, frame shift mutations, deletions, and insertions. Preferably, the region of the gene encoding the 5'→3' exonuclease activity is deleted using techniques well known in the art. In embodiments of this invention, any one of six conserved amino acids that are associated with the 5'→3' exonuclease activity can be mutated. Examples

25 of these conserved amino acids with respect to *Tne* DNA polymerase include Asp⁸, Glu¹¹², Asp¹¹⁴, Asp¹¹⁵, Asp¹³⁷, and Asp¹³⁹. Other possible sites for mutation are Gly¹⁰², Gly¹⁸⁷ and Gly¹⁹⁵.

-35-

Corresponding amino acid to target for other polymerases to reduce or eliminate 5'→3' exonuclease activity as follows:

5 *E. coli* polI: Asp¹³, Glu¹¹³, Asp¹¹⁵, Asp¹¹⁶, Asp¹³⁸, and Asp¹⁴⁰.
Taq pol: Asp¹⁸, Glu¹¹⁷, Asp¹¹⁹, Asp¹²⁰, Asp¹⁴², and Asp¹⁴⁴.
Tma pol: Asp⁸, Glu¹¹², Asp¹¹⁴, Asp¹¹⁵, Asp¹³⁷, and Asp¹³⁹.

Amino acid residues of *Taq* DNA polymerase are as numbered in U.S. 5,079,352. Amino acid residues of *Thermotoga maritima* (*Tma*) DNA polymerase are numbered as in U.S. Patent No. 5,374,553.

10 Examples of other amino acids which may be targeted for other polymerases to reduce 5'→3' exonuclease activity include:

Enzyme or source	Mutation positions
<i>Streptococcus pneumoniae</i>	Asp ¹⁰ , Glu ¹¹⁴ , Asp ¹¹⁶ , Asp ¹¹⁷ , Asp ¹³⁹ , Asp ¹⁴¹
<i>Thermus flavus</i>	Asp ¹⁷ , Glu ¹¹⁶ , Asp ¹¹⁸ , Asp ¹¹⁹ , Asp ¹⁴¹ , Asp ¹⁴³
<i>Thermus thermophilus</i>	Asp ¹⁸ , Glu ¹¹⁸ , Asp ¹²⁰ , Asp ¹²¹ , Asp ¹⁴³ , Asp ¹⁴⁵
15 <i>Deinococcus radiodurans</i>	Asp ¹⁸ , Glu ¹¹⁷ , Asp ¹¹⁹ , Asp ¹²⁰ , Asp ¹⁴² , Asp ¹⁴⁴
<i>Bacillus caldotenax</i>	Asp ⁹ , Glu ¹⁰⁹ , Asp ¹¹¹ , Asp ¹¹² , Asp ¹³⁴ , Asp ¹³⁶

Coordinates of *S. pneumoniae*, *T. flavus*, *D. radiodurans*, *B. caldotenax* were obtained from Gutman and Minton. Coordinates of *T. thermophilus* were obtained from International Patent No. WO 92/06200.

20 Typically, the mutant polymerases of the invention can be affected by substitution of amino acids typically which have different properties. For example, an acidic amino acid such as Asp may be changed to a basic, neutral or polar but uncharged amino acid such as Lys, Arg, His (basic); Ala, Val, Leu, Ile, Pro, Met, Phe, Trp (neutral); or Gly, Ser, Thr, Cys, Tyr, Asn or Gln (polar but uncharged).
 25 Glu may be changed to Asp, Ala, Val Leu, Ile, Pro, Met, Phe, Trp, Gly, Ser, Thr, Cys, Tyr, Asn or Gln.

-36-

5 Preferably, oligonucleotide directed mutagenesis is used to create the mutant polymerases which allows for all possible classes of base pair changes at any determined site along the encoding DNA molecule. In general, this technique involves annealing a oligonucleotide complementary (except for one or more mismatches) to a single stranded nucleotide sequence coding for the DNA polymerase of interest. The mismatched oligonucleotide is then extended by DNA polymerase, generating a double stranded DNA molecule which contains the desired change in the sequence on one strand. The changes in sequence can of course result in the deletion, substitution, or insertion of an amino acid. The double stranded polynucleotide can then be inserted into an appropriate expression vector, and a mutant polypeptide can thus be produced. The above-described oligonucleotide directed mutagenesis can of course be carried out via PCR.

Enhancing Expression of Polymerases

15 To optimize expression of the polymerases of the present invention, inducible or constitutive promoters are well known and may be used to express high levels of a polymerase structural gene in a recombinant host. Similarly, high copy number vectors, well known in the art, may be used to achieve high levels of expression. Vectors having an inducible high copy number may also be useful to enhance expression of the polymerases of the invention in a recombinant host.

20 To express the desired structural gene in a prokaryotic cell (such as, *E. coli*, *B. subtilis*, *Pseudomonas*, etc.), it is necessary to operably link the desired structural gene to a functional prokaryotic promoter. However, the natural promoter of the polymerase gene may function in prokaryotic hosts allowing expression of the polymerase gene. Thus, the natural promoter or other promoters may be used to express the polymerase gene. Such other promoters may be used to enhance expression and may either be constitutive or regulatable (i.e., inducible or derepressible) promoters. Examples of constitutive promoters include the *int* promoter of bacteriophage λ , and the *bla* promoter of the β -lactamase gene of pBR322. Examples of inducible prokaryotic promoters include the major right and left promoters of bacteriophage λ (P_R and P_L), *trp*,

25
30

-37-

recA, *lacZ*, *lacI*, *tet*, *gal*, *trc*, and *tac* promoters of *E. coli*. The *B. subtilis* promoters include α -amylase (Ulmanen *et al.*, *J. Bacteriol* 162:176-182 (1985)) and *Bacillus* bacteriophage promoters (Gryczan, T., In: *The Molecular Biology Of Bacilli*, Academic Press, New York (1982)). *Streptomyces* promoters are described by Ward *et al.*, *Mol. Gen. Genet.* 203:468478 (1986)). Prokaryotic promoters are also reviewed by Glick, *J. Ind. Microbiol.* 1:277-282 (1987); Cenatiemto, Y., *Biochimie* 68:505-516 (1986); and Gottesman, *Ann. Rev. Genet.* 18:415-442 (1984). Expression in a prokaryotic cell also requires the presence of a ribosomal binding site upstream of the gene-encoding sequence. Such ribosomal binding sites are disclosed, for example, by Gold *et al.*, *Ann. Rev. Microbiol.* 35:365404 (1981).

To enhance the expression of polymerases of the invention in a eukaryotic cell, well known eukaryotic promoters and hosts may be used. Preferably, however, enhanced expression of the polymerases is accomplished in a prokaryotic host. The preferred prokaryotic host for overexpressing the polymerases of the invention is *E. coli*.

Isolation and Purification of Polymerases

The enzyme(s) of the present invention is preferably produced by fermentation of the recombinant host containing and expressing the desired polymerase gene. However, the polymerases of the present invention may be isolated from any strain which produces the polymerase of the present invention. Fragments of the polymerase are also included in the present invention. Such fragments include proteolytic fragments and fragments having polymerase activity.

Any nutrient that can be assimilated by a host containing the polymerase gene may be added to the culture medium. Optimal culture conditions should be selected case by case according to the strain used and the composition of the culture medium. Antibiotics may also be added to the growth media to insure maintenance of vector DNA containing the desired gene to be expressed. Media formulations have been described in DSM or ATCC Catalogs and Sambrook *et*

-38-

al., In: *Molecular Cloning, a Laboratory Manual* (2nd ed.), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989).

5 Host cells producing the polymerases of this invention can be separated from liquid culture, for example, by centrifugation. In general, the collected microbial cells are dispersed in a suitable buffer, and then broken down by ultrasonic treatment or by other well known procedures to allow extraction of the enzymes by the buffer solution. After removal of cell debris by ultracentrifugation or centrifugation, the polymerase can be purified by standard protein purification techniques such as extraction, precipitation, chromatography, affinity chromatography, electrophoresis or the like. Assays to detect the presence of the polymerase during purification are well known in the art and can be used during conventional biochemical purification methods to determine the presence of these enzymes.

10

***Thermotoga* Polymerases**

15 *Thermotoga* polymerases for use in the present invention are obtained from any strain of *Thermotoga* species, more preferably from a strain of *Thermotoga neapolitana* (WO 96/10640 or WO96/41014) or *Thermotoga maritima* (U.S. Patent No. 5,374,553). Enzymes suitable for use in the present invention from these more preferred sources are the wildtype DNA polymerases (*Tne* from *T. neapolitana*; *Tma* from *T. maritima*), or mutants or derivatives thereof.

20

The present invention provides isolated nucleic acid molecules encoding preferred mutant *Tne* DNA polymerases, mutant *Tne* DNA polymerases encoded by such isolated nucleic acid molecules, and specific mutant *Tne* DNA polymerase proteins. Most preferred are the wildtype *Tne* DNA polymerase (SEQ ID NOs:1,2), the wildtype *Tma* DNA polymerase (U.S. Patent No. 5,374,553), and the following mutants of *Tne* DNA polymerase: *Tne* N'Δ219, D323A (SEQ ID NO:3); *Tne* N'Δ283, D323A (SEQ ID NO:4); *Tne* N'Δ192, D323A (SEQ ID NO:5); *Tne* D137A, D323A (SEQ ID NO:6); *Tne* D8A, D323A (SEQ ID NO:7); *Tne* G195D, D323A (SEQ ID NO:8); *Tne* G37D, D323A (SEQ ID NO:9);

25

-39-

Tne N'Δ283 (SEQ ID NO:10); *Tne* D137A, D323A, R722K; *Tne* D137A, D323A, R722Y; *Tne* D137A, D323A, R722L; *Tne* D137A, D323A, R722H; *Tne* D137A, D323A, R722Q; *Tne* D137A, D323A, F730Y; *Tne* D137A, D323A, K726R; *Tne* D137A, D323A, K726H; *Tne* D137A, D323A, R722K, F730Y; *Tne* D137A, D323A, R722K, K726R; *Tne* D137A, D323A, R722K, K726H; *Tne* D137A, D323A, R722H, F730Y; *Tne* D137A, D323A, R722H, K726R; *Tne* D137A, D323A, R722H, K726H; *Tne* D137A, D323A, R722Q, F730Y; *Tne* D137A, D323A, R722Q, K726R; *Tne* D137A, D323A, R722Q, K726H; *Tne* D137A, D323A, R722N, F730Y; *Tne* D137A, D323A, R722N, K726R; *Tne* D137A, D323A, R722N, K726H; *Tne* D137A, D323A, F730S; *Tne* N'Δ283, D323A, R722K/H/Q/N/Y/L; *Tne* N'Δ219, D323A, R722K; *Tne* N'Δ219, D323A, F730Y; *Tne* N'Δ219, D323A, K726R; *Tne* N'Δ219, D323A, K726H; *Tne* D137A, D323A, F730S, R722K/Y/Q/N/H/L, K726R/H; *Tne* D137A, D323A, F730T, R722K/Y/Q/N/H/L, K726R/H; *Tne* D137A, D323A, F730T; *Tne* F730S; *Tne* F730A; *Tne* K726R; *Tne* K726H; and *Tne* D137A, D323A, R722N. It will of course be understood by the skilled artisan that the designations of the above-described mutant polymerases indicate the position of the amino acid residue in the wildtype amino acid sequence (SEQ ID NO:2) that is being mutated, as well as to what residue the amino acid is being mutated. Thus, for example, "D137A" indicates that the Asp (D) residue at position 137 in SEQ ID NO:2 has been mutated to an Ala (A) residue, and, for example, "R722K/Y/Q/N/H/L" indicates that the Arg (R) residue at position 722 in SEQ ID NO:2 has been mutated to a Lys (K), Tyr (Y), Gln (Q), Asn (N), His (H) or Leu (L) residue. Mutant polymerases having one or more mutations or modifications corresponding to the *Tne* mutants of the invention are also contemplated by the invention.

The following chart indicates the nucleic acid sequences of the nucleic acid molecules encoding the above-described mutant *Tne* DNA polymerases (SEQ ID NOs:3-10), each with reference to the wildtype *Tne* DNA polymerase (SEQ ID NO:1):

-40-

SEQ ID NO:	Deletion of SEQ ID NO: 1	Insertion	Substitution to SEQ ID NO: 1
3	Deletion of positions 1-657 from the 5'-end	ATG AGC TTC at the 5'-end	A replaces G at position 966; C replaces A at 968 and G replaces C at 969
4	Deletion of positions 1-849 from the 5'-end	None	A replaces G at 966; C replaces A at 968 and G replaces C at 969
5	Deletion of positions 1-576 from the 5'-end	ATG AAT TCG AGC TCG GTA CCC at the 5'-end	A replaces G at 966; C replaces A at 968 and G replaces C at 969; A replaces G at 584
6	None	None	A replaces G at 966; C replaces A at 968 and G replaces C at 969; C replaces T at 408 and C replaces A at 410
7	None	None	A replaces G at 966; C replaces A at 968 and G replaces C at 969; C replaces A at 23 and C replaces T at 24
8	None	None	A replaces G at 966; C replaces A at 968 and G replaces C at 969; T replaces C at 576 and A replaces G at 584
9	None	None	A replaces G at 966; C replaces A at 968 and G replaces C at 969; A replaces G at 110
10	Deletion of positions 1-849 from the 5'-end	None	None

Using these same approaches, the sequence guidance provided herein, and knowledge of appropriate nucleotide substitutions to be made to SEQ ID NO:1, one of ordinary skill can readily produce other nucleic acid molecules encoding mutant polymerases, such as those described in detail above, having the desired activity. In addition, other nucleic acid molecules which comprise a sequence

-41-

substantially different from those described above but which, due to the degeneracy of the genetic code, still encode a mutant *Tne* DNA polymerase having an amino acid sequence set forth above, are also encompassed by the present invention. Since the genetic code is well known in the art, it is routine for one of ordinary skill in the art to produce such mutants and degenerate variants without undue experimentation.

Each of these mutant *Tne* DNA polymerases are reduced or substantially reduced in the ability to add a non-templated 3' terminal nucleotide to the growing strand. These mutant *Tne* DNA polymerase proteins may be prepared by recombinant DNA techniques routine to one of ordinary skill. Preferably, such mutant *Tne* polymerases are prepared by inserting an isolated DNA molecule having a nucleotide sequence as described above for each individual mutant into a recombinant vector, inserting the vector into a host cell, preferably an *Escherichia coli* cell, and culturing the host cell under conditions favoring the production of the mutant *Tne* DNA polymerase. The mutant *Tne* polymerase is then isolated from the host cell according to standard protein purification techniques. Further guidance for the preparation and isolation of mutant DNA polymerases from thermostable microorganisms can be found, for example, in U.S. Patent No. 5,374,553, in co-pending U.S. Patent Application No. 08/689,818 of Deb K. Chatterjee and A. John Hughes, entitled "Cloned DNA Polymerases from *Thermotoga* and Mutants Therof," filed September 6, 1996, and in co-pending U.S. Patent Application No. 08/689,807 of Deb K. Chatterjee, entitled "Cloned DNA Polymerases from *Thermotoga* and Mutants Therof," filed September 6, 1996, the disclosures of all of which are incorporated herein in their entirety.

In the methods of the present invention, *Thermotoga* DNA polymerases substantially reduced in 3'-5' exonuclease activity (such as a *Tne* mutant having an amino acid sequence as set forth in any one of SEQ ID NOs:3-9), or *Thermotoga* DNA polymerases not substantially reduced in 3'-5' exonuclease activity (such as *Tne* DNA polymerase (SEQ ID NOs:1,2), *Tma* DNA polymerase (U.S. Patent No. 5,374,553), or the *Tne* mutant *Tne*N Δ 283 (SEQ ID NO:10)), may be used with

-42-

similar results, since both types of *Thermotoga* DNA polymerase are substantially reduced in the ability to add a nontemplated 3' terminal nucleotide to a DNA template. Other thermostable DNA polymerases substantially reduced in 3'-5' exonuclease activity, such as *Taq*, VENT™(exo-), DEEPVENT™(exo-), Dtok(exo-) and THERMOLASE™ *Tbr*, are not preferred for use in the present methods as they will add non-templated nucleotides to the 3' termini of the amplification products as described below. However, such thermostable polymerase can be made which have reduced, substantially reduced or eliminated activity to add 3' non-template nucleotides by mutating or modifying the polymerase in accordance with the invention. The preferred *Thermotoga* polymerases of the invention contain such mutations or modifications in their O-helix.

The recombinant host comprising the gene encoding *Tne* DNA polymerase, *E. coli* DH10B(pUC-Tne), was deposited on September 30, 1994, with the Collection, Agricultural Research Culture Collection (NRRL), 1815 North University Street, Peoria, Illinois 61604 USA, as Deposit No. NRRL B-21238. The gene encoding *Tma* DNA polymerase has also been cloned and sequenced (U.S. Patent No. 5,374,553, which is expressly incorporated by reference herein in its entirety). Methods for preparing mutants and derivatives of these *Tne* and *Tma* polymerases are well-known in the art, and are specifically described in co-pending U.S. Patent Application No. 08/689,818 of Deb K. Chatterjee and A. John Hughes, entitled "Cloned DNA Polymerases from *Thermotoga* and Mutants Therof," filed September 6, 1996, and co-pending U.S. Patent Application No. 08/689,807 of Deb K. Chatterjee, entitled "Cloned DNA Polymerases from *Thermotoga* and Mutants Therof," filed September 6, 1996, the disclosures of which are incorporated herein in their entirety.

Advantages of Thermostable Polymerases

The use of thermostable polymerases (e.g. *Thermotoga* polymerases) or mutants or derivatives thereof in the methods of the present invention provide

-43-

several distinct advantages. These advantages are particularly apparent in the application of the present methods to analysis and typing of minisatellite, microsatellite and STR DNA regions.

5 With respect to traditional thermolabile DNA polymerases used in DNA amplification and sequencing, such as T4, T7 or *E. coli* Klenow fragment polymerases, thermostable polymerases such as *Thermotoga* DNA polymerases maintain their enzymatic activity in the multiple high-temperature cycles used in PCR and analogous automated amplification methodologies. It is therefore unnecessary to add fresh enzyme at the beginning of each amplification cycle when
10 using thermostable polymerases, as must be done when thermolabile enzymes are used.

With respect to other thermostable enzymes, it has been unexpectedly discovered in the present invention (as described in more detail in the Examples below) that the use of *Tne* or *Tma* DNA polymerase mutants or derivatives
15 thereof, does not result in the incorporation of non-templated 3' nucleotides into the newly synthesized DNA strands during DNA amplification reactions. This non-templated incorporation is a common problem when using certain other commonly employed thermostable enzymes, such as *Taq*, VENT™(exo-), DEEPVENT™(exo-), Dtok(exo-) and THERMOLASE™ *Tbr*. It has also been
20 unexpectedly discovered that mutants of these polymerases can be made to reduce or eliminate addition of non-templated 3' nucleotides. In particular, such mutations are preferably made within the O-helix of such polymerases.

Thus, the use of *Tne* or *Tma* DNA polymerases or mutants or derivatives thereof (or other mutant polymerases produced according to the invention) in
25 amplifying and typing DNA sequences, particularly hypervariable DNA sequences such as minisatellite, microsatellite or STR regions, will allow a faithful amplification and resolution of polymorphisms in these regions. This faithful resolution is not possible using other thermostable polymerases due to their propensity for non-templated incorporation. Thus, these enzymes are suitable for
30 use in automated amplification systems such as PCR.

-44-

Sources of DNA

Suitable sources of DNA, including a variety of cells, tissues, organs or organisms, may be obtained through any number of commercial sources (including American Type Culture Collection (ATCC), Rockville, Maryland; Jackson Laboratories, Bar Harbor, Maine; Cell Systems, Inc., Kirkland, Washington; Advanced Tissue Sciences, La Jolla, California). Cells that may be used as starting materials for genomic DNA preparation are preferably eukaryotic (including fungi or yeasts, plants, protozoans and other parasites, and animals including humans and other mammals). Although any mammalian cell may be used for preparation of DNA, preferred are blood cells (erythrocytes and leukocytes), endothelial cells, epithelial cells, neuronal cells (from the central or peripheral nervous systems), muscle cells (including myocytes and myoblasts from skeletal, smooth or cardiac muscle), connective tissue cells (including fibroblasts, adipocytes, chondrocytes, chondroblasts, osteocytes and osteoblasts) and other stromal cells (*e.g.*, macrophages, dendritic cells, Schwann cells), although other cells, including the progenitors, precursors and stem cells that give rise to the above-described somatic cells, are equally suitable. Also suitable for use in the preparation of DNA are mammalian tissues or organs such as those derived from brain, kidney, liver, pancreas, blood, bone marrow, muscle, nervous, skin, genitourinary, circulatory, lymphoid, gastrointestinal and connective tissue sources, as well as those derived from a mammalian (including human) embryo or fetus. These cells, tissues and organs may be normal, or they may be pathological such as those involved in infectious diseases (caused by bacteria, fungi or yeast, viruses (including AIDS) or parasites), in genetic or biochemical pathologies (*e.g.*, cystic fibrosis, hemophilia, Alzheimer's disease, schizophrenia, muscular dystrophy or multiple sclerosis), or in cancerous processes.

More specifically, in one aspect of the invention, the relationship between a first individual and a second individual may be determined by analyzing and typing a particular polymorphic DNA fragment, such as a minisatellite or microsatellite DNA sequence. In such a method, the amplified fragments for each

-45-

individual are compared to determine similarities or dissimilarities. Such an analysis is accomplished, for example, by comparing the size of the amplified fragments from each individual, or by comparing the sequence of the amplified fragments from each individual. In another aspect of the invention, genetic identity can be determined. Such identity testing is important, for example, in paternity testing, forensic analysis, etc. In this aspect of the invention, a sample containing DNA (*e.g.*, a crime scene sample or a sample from an individual) is analyzed and compared to a sample from one or more individuals. In one such aspect of the invention, one sample of DNA may be derived from a first individual and another sample may be derived from a second individual whose relationship to the first individual is unknown; comparison of these samples from the first and second individuals by the methods of the invention may then facilitate a determination of the genetic identity or relationship between the first and second individual. In a particularly preferred such aspect, the first DNA sample may be a known sample derived from a known individual and the second DNA sample may be an unknown sample derived, for example, from crime scene material. In an additional aspect of the invention, one sample of DNA may be derived from a first individual and another sample may be derived from a second individual who is related to the first individual; comparison of these samples from the first and second individuals by the methods of the invention may then facilitate a determination of the genetic kinship of the first and second individuals by allowing examination of the Mendelian inheritance, for example, of a polymorphic, minisatellite, microsatellite or STR DNA fragment. In another aspect of the invention, DNA fragments important as genetic markers for encoding a gene of interest can be identified and isolated. For example, by comparing samples from different sources, DNA fragments which may be important in causing diseases such as infectious diseases (of bacterial, fungal, parasitic or viral etiology), cancers or genetic diseases, can be identified and characterized. In this aspect of the invention a DNA sample from normal cells or tissue is compared to a DNA sample from diseased cells or tissue. Upon comparison according to the invention, one

-46-

or more unique polymorphic fragments present in one DNA sample and not present in the other DNA sample can be identified and isolated. Identification of such unique polymorphic fragments allows for identification of sequences associated with, or involved in, causing the diseased state.

5 Once the starting cells, tissues, organs or other samples are obtained, DNA may be prepared therefrom by methods that are well-known in the art (*See, e.g.,* Maniatis, T., *et al., Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press, pp. 9.16-9.23 (1989); Kaufman, P.B., *et al., Handbook of Molecular and Cellular Methods in Biology*
10 *and Medicine*, Boca Raton, Florida: CRC Press, pp. 1-26 (1995)). The DNA samples thus prepared may then be used to identify, analyze and type polymorphic DNA fragments, including minisatellite, microsatellite and STR DNA fragments, by amplification, preferably by PCR amplification, as modified by the methods of the present invention.

15 General methods for amplification and analysis of DNA fragments are well-known to one of ordinary skill in the art (*see, e.g.,* U.S. Pat. Nos. 4,683,195; 4,683,202; and 4,800,159; Innis, M.A., *et al., eds., PCR Protocols: A Guide to Methods and Applications*, San Diego, California: Academic Press, Inc. (1990); Griffin, H.G., and Griffin, A.M., *eds., PCR Technology: Current Innovations*,
20 Boca Raton, Florida: CRC Press (1994)). Typically, these methods comprise contacting the DNA sample with a thermostable DNA polymerase in the presence of one or more primer sequences, amplifying the DNA sample to generate a collection of amplified polymorphic, minisatellite, microsatellite or STR DNA fragments, preferably by PCR or equivalent automated amplification technique,
25 separating the amplified DNA fragments by size, preferably by gel electrophoresis, and analyzing the gels for the presence of polymorphic, minisatellite, microsatellite or STR DNA fragments by direct comparison of the pattern of fragments generated from a first sample of DNA to those from a second sample of DNA, or by a more indirect comparison using known size markers.

-47-

As noted above, amplification protocols used heretofore for analyzing and typing polymorphic DNA fragments, particularly minisatellite, microsatellite or STR DNA sequences, use certain thermostable DNA polymerases such as *Taq* (U.S. Patent Nos. 4,683,195; 4,683,202; and 4,800,159). However, as discussed in detail above, these approaches yield amplification products in which one or more non-templated nucleotides is added to the 3' termini of the products by the polymerases, thus leading to heterogeneity in the amplification products, and ambiguity concerning the correct size of the amplification products.

This problem is overcome in the present invention by contacting the DNA sample in the amplification reaction mixtures with one or more DNA polymerases of the invention which are reduced, substantially reduced or eliminated in the ability to add a nontemplated 3' terminal nucleotide to the growing strand. Preferably, such DNA polymerases are *Thermotoga* DNA polymerases, more preferably a *Thermotoga* DNA polymerase substantially reduced in 3'-5' exonuclease activity, still more preferably a *Tne* polymerase (SEQ ID NOs:1,2), a *Tma* polymerase (U.S. Patent No. 5,374,553), or a mutant or derivative thereof, and most preferably one of the following mutants of *Tne* polymerase: *Tne* N Δ 219, D323A (SEQ ID NO:3); *Tne* N Δ 283, D323A (SEQ ID NO:4); *Tne* N Δ 192, D323A (SEQ ID NO:5); *Tne* D137A, D323A (SEQ ID NO:6); *Tne* D8A, D323A (SEQ ID NO:7); *Tne* G195D, D323A (SEQ ID NO:8); *Tne* G37D, D323A (SEQ ID NO:9); *Tne* N Δ 283 (SEQ ID NO:10); *Tne* D137A, D323A, R722K; *Tne* D137A, D323A, R722Y; *Tne* D137A, D323A, R722L; *Tne* D137A, D323A, R722H; *Tne* D137A, D323A, R722Q; *Tne* D137A, D323A, F730Y; *Tne* D137A, D323A, K726R; *Tne* D137A, D323A, K726H; *Tne* D137A, D323A, R722K, F730Y; *Tne* D137A, D323A, R722K, K726R; *Tne* D137A, D323A, R722K, K726H; *Tne* D137A, D323A, R722H, F730Y; *Tne* D137A, D323A, R722H, K726R; *Tne* D137A, D323A, R722H, K726H; *Tne* D137A, D323A, R722Q, F730Y; *Tne* D137A, D323A, R722Q, K726R; *Tne* D137A, D323A, R722Q, K726H; *Tne* D137A, D323A, R722N, F730Y; *Tne* D137A, D323A, R722N, K726R; *Tne* D137A, D323A, R722N, K726H; *Tne* D137A,

-48-

D323A, F730S; *Tne* N'Δ283, D323A, R722K/H/Q/N/Y/L; *Tne* N'Δ219, D323A, R722K; *Tne* N'Δ219, D323A, F730Y; *Tne* N'Δ219, D323A, K726R; *Tne* N'Δ219, D323A, K726H; *Tne* D137A, D323A, F730S, R722K/Y/Q/N/H/L, K726R/H; *Tne* D137A, D323A, F730T, R722K/Y/Q/N/H/L, K726R/H; *Tne* D137A, D323A, F730T; *Tne* F730S; *Tne* F730A; *Tne* K726R; *Tne* K726H; and *Tne* D137A, D323A, R722N.

It will be understood, however, that other thermostable DNA polymerases or mutants thereof, any of which are reduced, substantially reduced, or eliminated in the ability to add a non-templated 3' terminal nucleotide to the growing strand, may be used in the methods of the present invention equivalently. The DNA polymerases are used in the methods of the present invention at a concentration of about 0.0001 units/ml to about 10 units/ml, preferably at a concentration of about 0.001 units/ml to about 5 units/ml, more preferably at a concentration of about 0.004 units/ml to about 1 unit/ml, and most preferably at a concentration of about 0.04 units/ml. Thus, the methods of the present invention produce a population of amplified DNA fragments, most preferably of polymorphic or microsatellite DNA fragments, which comprise substantially no non-templated 3' terminal nucleotides. By "substantially no non-templated 3' terminal nucleotides" is meant that the population of amplified DNA fragments demonstrates about 0-50%, about 0-30%, about 0-20%, preferably about 0-10%, more preferably about 0-5%, still more preferably about 0-1% and most preferably about 0%, of DNA molecules containing non-templated 3' nucleotides compared to amplified DNA fragments produced by the polymerase control. When testing the ability of a DNA polymerase to add 3' non-templated nucleotides, the polymerase, when it has substantially reduced or eliminated 3' exonuclease activity, is compared to *Taq* DNA polymerase (see above). When testing polymerases which have been modified or mutated to reduce or eliminate 3' non-templated nucleotide addition, the mutated or modified polymerase is compared to the corresponding wildtype, unmodified or unmutated polymerase (see above).

-49-

Following amplification by the methods of the present invention, the amplified DNA fragments may be analyzed to identify or type a polymorphic, minisatellite, microsatellite or STR DNA fragment. This step is usually accomplished by separation of the amplified DNA fragments by size, a procedure which permits the determination of the presence of unique polymorphic fragments in one or more of the DNA samples. The fragments may be separated by any physical or biochemical means including gel electrophoresis, capillary electrophoresis, chromatography (including sizing, affinity and immunochromatography), density gradient centrifugation and immunoadsorption. For carrying out the present invention, separation of DNA fragments by gel electrophoresis is particularly preferred, as it provides a rapid and highly reproducible means of sensitive separation of a multitude of DNA fragments, and permits direct, simultaneous comparison of the fragments in several samples of DNA, or samples of DNA from a first and a second individual.

Gel electrophoresis is typically performed on agarose or polyacrylamide sequencing gels according to standard protocols, preferably using gels containing polyacrylamide at concentrations of 3-12% and most preferably at about 8%, and containing urea at a concentration of about 4-12M, most preferably about 8M. Samples are loaded onto the gels, usually with samples containing amplified DNA fragments prepared from different sources of genomic DNA being loaded into adjacent lanes of the gel to facilitate subsequent comparison. Reference markers of known sizes may be used to facilitate the comparison of samples. Following electrophoretic separation, DNA fragments may be visualized and identified by a variety of techniques that are routine to those of ordinary skill in the art, such as autoradiography. One can then examine the autoradiographic films either for differences in polymorphic fragment patterns ("typing") or for the presence of one or more unique bands in one lane of the gel ("identifying"); the presence of a band in one lane (corresponding to a single sample, cell or tissue type) that is not observed in other lanes indicates that the DNA fragment comprising that unique band is source-specific and thus a potential polymorphic DNA fragment.

-50-

5 A variety of DNA fragments comprising polymorphic, minisatellite, microsatellite or STR DNA fragments can thus be identified using the methods of the present invention by comparing the pattern of bands on the films depicting various samples. Importantly, using the present methods the amplification products of the polymorphic DNA fragments will be faithful copies of the template (allele) material -- *i.e.*, they will not exhibit undesired additional nucleotides at their 3' termini via non-templated addition of nucleotides by the polymerases. One can extend this approach, in another preferred embodiment, to isolate and characterize these fragments or any DNA fragment amplified without the non-templated addition of a 3' terminal nucleotide. In this embodiment, one or more of the unique DNA fragments are removed from the gel which was used for identification (see above), according to standard techniques such as electroelution or physical excision.

15 The isolated unique DNA fragments may then be inserted into standard nucleotide vectors, including expression vectors, suitable for transfection or transformation of a variety of prokaryotic (bacterial) or eukaryotic (yeast, plant or animal including human and other mammalian) cells. In particular, the present invention provides methods of cloning such isolated unique DNA fragments, or any PCR-amplified DNA fragment, by blunt-end cloning. As described above, 20 *Taq* DNA polymerase adds a non-templated nucleotide, typically a deoxyadenosine ("A"), to the 3' terminus of the amplified DNA fragment. Thus, *Taq*-catalyzed PCR generates a collection of DNA fragments with 3' A overhangs. To clone such *Taq*-amplified fragments, two approaches are commonly used: either the 3' A overhang is removed by treating the amplified fragment with, for example, T4 DNA polymerase (a technique sometimes called "3' polishing"), or 25 a special cloning vector with a 3' T overhang ("TA cloning vector") is used. Of course, such approaches are more time-consuming and expensive than if direct insertion of the amplified fragment were done. Such a direct approach is possible using the methods of the invention, which generates little or no 3' A overhangs 30 (and thus, blunt ends) on the amplified DNA fragments. The DNA fragments,

-51-

amplified according to the methods of the invention, may thus be directly inserted into corresponding blunt-ended vectors according to standard techniques (for example, using T4 DNA ligase). Thus, the present invention provides a method of blunt-end cloning of a DNA fragment that obviates the use of TA cloning vectors or 3' polishing.

5

To identify the presence of minisatellite DNA fragments, the polymorphic DNA fragments that are identified and isolated by the methods of the present invention may be further characterized, for example by sequencing (*i.e.*, determining the nucleotide sequence of the polymorphic fragments), by methods described above and others that are standard in the art (*see, e.g.*, U.S. Patent Nos. 4,962,022 and 5,498,523, which are directed to methods of DNA sequencing).

10

Kits

The invention also provides kits for use in the identification, analysis and typing of a polymorphic DNA fragment, particularly a minisatellite or STR DNA fragment, according to the present methods. Kits according to the present invention may comprise a carrying means being compartmentalized to receive in close confinement therein one or more containers such as vials, tubes, bottles and the like. Each of such containers may comprise components or a mixture of components needed to perform DNA amplification or analysis.

15

Such kits may comprise of one or more thermostable DNA polymerases reduced, substantially reduced or eliminated in the ability to add a non-templated 3' nucleotide to a growing DNA strand. Preferably the container contains a *Thermotoga* DNA polymerase or a mutant or a derivative thereof, particularly those described in full detail above. The kit may also contain one or more DNA primer molecules, one or more deoxyribonucleoside triphosphates needed to synthesize a DNA molecule complementary to a DNA template, and/or a buffer suitable for amplification of a nucleic acid molecule (or combinations thereof).

20

25

A kit for DNA analysis may include one or more of the above components, and may further include containers which contain reagents necessary for

-52-

separation and analysis of DNA fragments, such as polyacrylamide, agarose, urea, detergents and the like.

Of course, it is also possible to combine one or more of these reagents in a single tube. A detailed description of such formulations at working concentrations is described in co-pending U.S. Application No. 08/689,815 of
5 Ayoub Rashtchian and Joseph Solus, entitled "Stable Compositions for Nucleic Acid Amplification and Sequencing" filed on August 14, 1996, the disclosure of which is incorporated by reference herein in its entirety.

The invention also relates to kits for detectably labeling molecules, sequencing, amplifying and synthesizing molecules by well known techniques. See
10 U.S. Patent Nos. 4,962,020, 5,173,411, 4,795,699, 5,498,523, 5,405,776 and 5,244,797. Such kits may comprise a carrying means being compartmentalized to receive in close confinement one or more container means such as vials, test tubes and the like. Each of such container means comprises components or a mixture
15 of components needed to perform nucleic acid synthesis, sequencing, labeling, or amplification.

A kit for sequencing DNA may comprise a number of container means. Such a kit may comprise one or more of the polymerases of the invention, one or a number of types of nucleotides needed to synthesize a DNA molecule
20 complementary to DNA template, one or a number of different types of terminators (such as dideoxynucleoside triphosphates), a pyrophosphatase, one or a number of primers and/or a suitable sequencing buffer (or combinations of such components).

A kit used for amplifying or synthesizing of nucleic acids will comprise,
25 one or more polymerases of the invention, and one or a number of nucleotides or mixtures of nucleotides. Various primers may be included in a kit as well as a suitable amplification or synthesis buffers.

When desired, the kit of the present invention may also include container means which comprise detectably labeled nucleotides which may be used during
30 the synthesis or sequencing of a nucleic acid molecule. One of a number of labels

-53-

may be used to detect such nucleotides. Illustrative labels include, but are not limited to, radioactive isotopes, fluorescent labels, chemiluminescent labels, bioluminescent labels and enzyme labels.

Use of the Methods and Kits

5 The polymerases, methods and kits embodied in the present invention will have general utility in any application utilizing nucleic acid amplification methodologies, particularly those directed to the analysis and typing of polymorphic or minisatellite DNA fragments, and most particularly those directed to the analysis and typing of minisatellite, microsatellite and STR DNA fragments.

10 Amplification techniques in which the present methods may be used include PCR (U.S. Patent Nos. 4,683,195 and 4,683,202), Strand Displacement Amplification (SDA; U.S. Patent No. 5,455,166; EP 0 684 315), and Nucleic Acid Sequence-Based Amplification (NASBA; U.S. Patent No. 5,409,818; EP 0 329 822). Nucleic acid analysis and typing techniques which may employ the present

15 compositions include nucleic acid sequencing methods such as those disclosed in U.S. Patent Nos. 4,962,022 and 5,498,523, as well as more complex PCR-based nucleic acid fingerprinting techniques such as Random Amplified Polymorphic DNA (RAPD) analysis (Williams, J.G.K., *et al.*, *Nucl. Acids Res.* 18(22):6531-6535, 1990), Arbitrarily Primed PCR (AP-PCR; Welsh, J., and McClelland, M., *Nucl. Acids Res.* 18(24):7213-7218, 1990), DNA Amplification Fingerprinting (DAF; Caetano-Anollés *et al.*, *Bio/Technology* 9:553-557, 1991), and microsatellite PCR or Directed Amplification of Minisatellite-region DNA (DAMD; Heath, D.D., *et al.*, *Nucl. Acids Res.* 21(24): 5782-5785, 1993). In particular, the polymerases, methods and kits of the present invention will be

20 useful in the fields of medical genetics, therapeutics and diagnostics, forensics (particularly identity and paternity testing), and agricultural (*e.g.*, plant breeding) and other biological sciences, in any procedure utilizing DNA polymerases for analysis and typing of polymorphic, minisatellite, microsatellite or STR DNA fragments. Particularly suitable for diagnosis by the methods of the present

25

-54-

invention are genetic diseases such as cystic fibrosis, hemophilia, Alzheimer's disease, schizophrenia, muscular dystrophy or multiple sclerosis. Together, these abilities will assist medical professionals and patients in diagnostic and prognostic determinations as well as in the development of treatment and prevention regimens for these and other disorders.

5

It will also be apparent to one of ordinary skill in the art that the present methods may be used to screen animal tissues to be subsequently used in medical procedures such as tissue or organ transplants, blood transfusions, zygote implantations and artificial inseminations. In such procedures, pre-screening of the subject tissues for the presence of particular polymorphic DNA fragments may improve the success of tissue or organ transplants (by decreasing the likelihood of rejection due to donor-recipient genetic incompatibility) and of zygote implantations (by eliminating the use of genetically defective zygotes). Similarly, use of these methods will reduce the chances of transmission of infectious diseases (e.g., hepatitis and AIDS) in medical procedures that are often prone to such transmission, such as blood transfusions and artificial insemination. Finally, use of the present invention for identification of unique polymorphic, minisatellite, microsatellite and STR DNA fragments will assist in forensic science in such applications as crime-scene analysis of blood, tissue and body secretions containing small amounts of DNA, as well as in paternity testing.

10

15

20

It will be readily apparent to those skilled in the relevant arts that other suitable modifications and adaptations to the methods and applications described herein are obvious and may be made without departing from the scope of the invention or any embodiment thereof. Having now described the present invention in detail, the same will be more clearly understood by reference to the following examples, which are included herewith for purposes of illustration only and are not intended to be limiting of the invention.

25

Example 1: Bacterial Strains And Growth Conditions

Thermotoga neapolitana DSM No. 5068 was grown under anaerobic conditions as described in the DSM catalog (addition of resazurin, Na₂S, and sulfur granules while sparging the media with nitrogen) at 85°C in an oil bath from 12 to 24 hours. The cells were harvested by filtering the broth through Whatman #1 filter paper. The supernatant was collected in an ice bath and then centrifuged in a refrigerated centrifuge at 8,000 rpms for twenty minutes. The cell paste was stored at -70°C prior to total genomic DNA isolation.

E. coli strains were grown in 2X LB broth base (Lennox L broth base: GIBCO/BRL) medium. Transformed cells were incubated in SOC (2% tryptone, 0.5% yeast extract, yeast 10 mM NaCl, 2.5 mM KCl, 20mM glucose, 10mM MgCl₂, and 10mM MgSO₄ per liter) before plating. When appropriate antibiotic supplements were 20 mg/l tetracycline and 100 mg/l ampicillin. *E. coli* strain DH10B (Lorow *et al.*, *Focus* 12:19-20 (1990)) was used as host strain. Competent DH10B may be obtained from Life Technologies, Inc. (LTI) (Rockville, MD).

Example 2: DNA Isolation

Thermotoga neapolitana chromosomal DNA was isolated from 1.1g of cells by suspending the cells in 2.5 ml TNE (50mM Tris-HCl, pH 8.0, 50mM NaCl, 10mM EDTA) and treated with 1% SDS for 10 minutes at 37°C. DNA was extracted with phenol by gently rocking the lysed cells overnight at 4°C. The next day, the lysed cells were extracted with chloroform:isoamyl alcohol. The resulting chromosomal DNA was further purified by centrifugation in a CsCl density gradient. Chromosomal DNA isolated from the density gradient was extracted three times with isopropanol and dialyzed overnight against a buffer containing 10 mM Tris-HCl (pH 8.0) and 1 mM EDTA (TE).

-56-

Example 3: Construction of Genomic Libraries

The chromosomal DNA isolated in Example 2 was used to construct a genomic library in the plasmid pCP13. Briefly, 10 tubes each containing 10 μ g of *Thermotoga neapolitana* chromosomal DNA was digested with 0.01 to 10 units of *Sau*IIIAl for 1 hour at 37°C. A portion of the digested DNA was tested in an agarose (1.2%) gel to determine the extent of digestion. Samples with less than 50% digestion were pooled, ethanol precipitated and dissolved in TE. 6.5 μ g of partially digested chromosomal DNA was ligated into 1.5 μ g of pCP13 cosmid which had been digested with *Bam*HI restriction endonuclease and dephosphorylated with calf intestinal alkaline phosphatase. Ligation of the partially digested *Thermotoga* DNA and *Bam*HI cleaved pCP13 was carried out with T4 DNA ligase at 22°C for 16 hours. After ligation, about 1 μ g of ligated DNA was packaged using λ -packaging extract (obtained from Life Technologies, Inc., Rockville, MD). DH10B cells (Life Tech. Inc.) were then infected with 100 μ l of the packaged material. The infected cells were plated on tetracycline containing plates. Serial dilutions were made so that approximately 200 to 300 tetracycline resistant colonies were obtained per plate.

Example 4: Screening for Clones Expressing Thermotoga neapolitana DNA Polymerase

Identification of the *Thermotoga neapolitana* DNA polymerase gene of the invention was cloned using the method of Sagner *et al.*, *Gene* 97:119-123 (1991) which reference is herein incorporated in its entirety. Briefly, the *E. coli* tetracycline resistant colonies from Example 3 were transferred to nitrocellulose membranes and allowed to grow for 12 hours. The cells were then lysed with the fumes of chloroform:toluene (1:1) for 20 minutes and dried for 10 minutes at room temperature. The membranes were then treated at 95°C for 5 minutes to inactivate the endogenous *E. coli* enzymes. Surviving DNA polymerase activity

-57-

was detected by submerging the membranes in 15 ml of polymerase reaction mix (50 mM Tris-HCl (pH 8.8), 1 mM MgCl₂, 3 mM β-mercaptoethanol, 10 μM dCTP, dGTP, dTTP, and 15 μCi of 3,000 Ci/mmol [α^{32} P]dATP) for 30 minutes at 65°C.

5 Using autoradiography, three colonies were identified that expressed a *Thermotoga neapolitana* DNA polymerase. The cells were grown in liquid culture and the protein extract was made by sonication. The presence of the cloned thermostable polymerase was confirmed by treatment at 90°C followed by measurement of DNA polymerase activity at 72°C by incorporation of radioactive
10 deoxyribonucleoside triphosphates into acid insoluble DNA. One of the clones, expressing *Tne* DNA polymerase, contained a plasmid designated pCP13-32 and was used for further study.

Example 5: Subcloning of Tne DNA polymerase

15 Since the pCP13-32 clone expressing the *Tne* DNA polymerase gene contains about 25 kb of *T. neapolitana* DNA, subcloning a smaller fragment of the *Tne* polymerase gene was attempted. The molecular weight of the *Tne* DNA polymerase purified from *E. coli*/pCP13-32 was about 100 kd. Therefore, a 2.5-3.0 kb DNA fragment will be sufficient to code for full-length polymerase. A second round of *Sau*3A partial digestion similar to Example 3 was done using
20 pCP13-32 DNA. In this case, a 3.5 kb region was cut out from the agarose gel, purified by Gene Clean (BIO 101, La Jolla, CA) and ligated into plasmid pSport 1 (Life Technologies, Inc.) which had been linearized with *Bam*HI and dephosphorylated with calf intestinal alkaline phosphatase. After ligation, DH10B was transformed and colonies were tested for DNA polymerase activity as
25 described in Example 1. Several clones were identified that expressed *Tne* DNA polymerase. One of the clones (pSport-*Tne*) containing about 3 kb insert was further characterized. A restriction map of the DNA fragment is shown in Fig. 1. Further, a 2.7 Kb *Hind*III-*Sst*I fragment was subcloned into pUC19 to generate

-58-

pUC19-*Tne*. *E. coli*/pUC19-*Tne* also produced *Tne* DNA polymerase. *E. coli* DH10B (pUC19-*Tne*) was deposited on September 30, 1994 with the Collection, Agricultural Research Culture Collection (NRRL), 1815 Peoria, IL 61604 as Deposit No. NRRL B-21338. The nucleotide and amino acid sequence of *Tne* polymerase is described in U.S. application serial nos. 08/706,702 and 08/706,706 filed September 9, 1996, both of which are incorporated by reference herein.

Example 6: Purification of *Thermotoga neapolitana* DNA Polymerase from *E. coli*

Twelve grams of *E. coli* cells expressing cloned *Tne* DNA polymerase (DH10B/pSport-*Tne*) were lysed by sonication (four thirty-second bursts with a medium tip at the setting of nine with a Heat Systems Ultrasonics Inc., model 375 sonicator) in 20 ml of ice cold extraction buffer (50 mM Tris HCl (pH 7.4), 8% glycerol, 5 mM mercaptoethanol, 10 mM NaCl, 1 mM EDTA, 0.5 mM PMSF). The sonicated extract was heated at 80°C for 15 min. and then cooled in ice for 5 min. 50 mM KCl and PEI (0.4%) was added to remove nucleic acids. The extract was centrifuged for clarification. Ammonium sulfate was added to 60%, the pellet was collected by centrifugation and resuspended in 10 ml of column buffer (25 mM Tris-HCl (pH 7.4), 8% glycerol, 0.5% EDTA, 5mM 2-mercaptoethanol, 10 mM KCl). A Blue-Sepharose (Pharmacia) column, or preferably a Toso heparin (Tosohaas) column, was washed with 7 column volumes of column buffer and eluted with a 15 column volume gradient of buffer from 10mM to 2 M KCl. Fractions containing polymerase activity were pooled. The fractions were dialyzed against 20 volumes of column buffer. The pooled fractions were applied to a Toso650Q column (Tosohaas). The column was washed to baseline OD₂₈₀ and elution effected with a linear 10 column volume gradient of 25 mM Tris (pH 7.4), 8% glycerol, 0.5 mM EDTA, 10 mM KCl, 5 mM β-mercaptoethanol to the same buffer plus 650 mM KCl. Active fractions were pooled.

-59-

Example 7: Construction of *Thermotoga neapolitana* 3'-to-5' Exonuclease Mutant

The amino acid sequence of portions of the *Tne* DNA polymerase was compared with other known DNA polymerases such as *E. coli* DNA polymerase 1, *Taq* DNA polymerase, T5 DNA polymerase, and T7 DNA polymerase to localize the regions of 3'-to-5' exonuclease activity, and the dNTP binding domains within the DNA polymerase. One of the 3'-to-5' exonuclease domains was determined based on the comparison of the amino acid sequences of various DNA polymerases (Blanco, L., et al. *Gene* 112: 139-144 (1992); Braithwaite and Ito, *Nucleic Acids Res.* 21: 787-802 (1993)) is as follows:

<i>Tne</i>	318	PSFALDLETSS	328	(SEQ ID NO:18)
Pol I	350	PVFAFDTETDS	360	(SEQ ID NO:19)
T5	159	GPVAFDSETSA	169	(SEQ ID NO:20)
T7	1	MIVSDIEANA	10	(SEQ ID NO:21)

As a first step to make the *Tne* DNA polymerase devoid of 3'→5' exonuclease activity, a 2kb *Sph* fragment from pSport-Tne was cloned into M13mp19 (LTI, Rockville, MD). The recombinant clone was selected in *E. coli* DH5αF'IQ (LTI, Rockville, MD). One of the clones with the proper insert was used to isolate uracilated single-stranded DNA by infecting *E. coli* CJ236 (Biorad, California) with the phage particle obtained from *E. coli* DH5αF'IQ. An oligonucleotide, GA CGT TTC AAG CGC TAG GGC AAA AGA (SEQ ID NO:22) was used to perform site directed mutagenesis. This site-directed mutagenesis converted Asp³²³ (indicated as * above) to Ala³²³. An *Eco*47III restriction site was created as part of this mutagenesis to facilitate screening of the mutant following mutagenesis. The mutagenesis was performed using a protocol as described in the Biorad manual (1987) except T7 DNA polymerase was used instead of T4 DNA polymerase (USB, Cleveland, OH). The mutant clones were

-60-

screened for the *Eco47III* restriction site that was created in the mutagenic oligonucleotide. One of the mutants having the created *Eco47III* restriction site was used for further study. The mutation Asp³²³ to Ala³²³ was confirmed by DNA sequencing.

5 To incorporate the 3'→5' exonuclease mutation in an expression vector, the mutant phage was digested with *SphI* and *HindIII*. A 2 kb fragment containing the mutation was isolated. This fragment was cloned in pUC-Tne to replace the wild type fragment. See Figure 2A. The desired clone, pUC-Tne (3'→5'), was isolated. The presence of the mutant sequence was confirmed by the presence of
10 the unique *Eco47III* site. The plasmid was then digested with *SstI* and *HindIII*. The entire mutant polymerase gene (2.6 kb) was purified and cloned into *SstI* and *HindIII* digested pTrc99 expression vector (Pharmacia, Sweden). The clones were selected in DH10B (LTI, Rockville, MD). The resulting plasmid was designated pTrcTne35. See Figure 2B. This clone produced active heat stable
15 DNA polymerase.

Example 8: Phenylalanine to Tyrosine Mutant

The polymerase active site including the dNTP binding domain is usually present at the carboxyl terminal region of the polymerase. The sequence of the *Tne* polymerase gene suggests that the amino acids that presumably contact and
20 interact with the dNTPs are present within the 694 bases starting at the internal *BamHI* site. See Figure 1. This conclusion is based on homology with a prototype polymerase *E. coli* DNA polymerase I. See Polisky et al., *J. Biol. Chem.* 265:14579-14591 (1990). A comparison was made of the O-helix for various polymerases:

-61-

	<i>Tne</i>	722	RRVGKMVNFSIYG	735	(SEQ ID NO:12)
	Pol I	754	RRSAKAINFGLIYG	767	(SEQ ID NO:13)
	T5	562	RQAAKAITFGILYG	575	(SEQ ID NO:14)
	T7	518	RDNAKTFIYGFLYG	531	(SEQ ID NO:15)
5	<i>Taq</i>	659	RRAAKTINFGVLYG	672	(SEQ ID NO:16)

10 In order to change Phe⁷³⁰ of the *Tne* polymerase to a Tyr⁷³⁰ site directed mutagenesis was performed using the oligonucleotide GTA TAT TAT AGA GTA GTT AAC CAT CTT TCC A (SEQ ID NO:23). As part of this oligonucleotide directed mutagenesis, a *HpaI* restriction site was created in order to screen mutants easily. The same uracilated single-stranded DNA and mutagenesis procedure described in Example 7 were used for this mutagenesis. Following mutagenesis, the mutants were screened for the *HpaI* site. Mutants with the desired *HpaI* site were used for further study. The mutation was confirmed by DNA sequencing.

15 The Phe⁷³⁰ to Tyr⁷³⁰ mutation was incorporated into pUC-*Tne* by replacing the wild type *SphI* -*HindIII* fragment with the mutant fragment obtained from the mutant phage DNA. The presence of the desired clone, pUC-*TneFY*, was confirmed by the presence of the unique *HpaI* site, see Figure 2A. The entire mutant polymerase gene was subcloned into pTrc99 as an *SstI*-*HindIII* fragment as described above in DH10B. The resulting plasmid was designated pTrc*TneFY*. (Figure 2B). The clone produced active heat stable polymerase.

Example 9: 3'→5' Exonuclease and Phe⁷³⁰→Tyr⁷³⁰ Double Mutants

25 In order to introduce the 3'→5' exonuclease mutation and the Phe⁷³⁰→Tyr⁷³⁰ mutation in the same expression vector, pTrc99, it was necessary to first reconstitute both mutations in the pUC-*Tne* clone. See Figure 3. Both the pUC-*Tne* (3'→5') and the pUC-*TneFY* were digested with *Bam*HI. The digested pUC-*Tne* (3'→5') was dephosphorylated to avoid recirculation in the following

-62-

ligations. The resulting fragments were purified on a 1% agarose gel. The largest *Bam*HI fragment (4.4 kb) was purified from pUC-Tne (3'→5') digested DNA and the smallest *Bam*HI fragment (0.8 kb) containing the Phe⁷³⁰→Tyr⁷³⁰ mutation was purified and ligated to generate pUC-Tne35FY. The proper orientation and the presence of both mutations in the same plasmid was confirmed by *Eco*47III, *Hpa*I, and *Sph*I-*Hind*III restriction digests. See Figure 3.

The entire polymerase containing both mutations was subcloned as a *Sst*I-*Hind*III fragment in pTrc99 to generate pTrcTne35FY in DH10B. The clone produced active heat stable polymerase.

Example 10: 3'→5' Exonuclease, 5'→3' Exonuclease, and Phe⁷³⁰→Tyr⁷³⁰ Triple Mutants

In most of the known polymerases, the 5'-to-3' exonuclease activity is present at the amino terminal region of the polymerase (Ollis, D.L., *et al.*, *Nature* 313, 762-766, 1985; Freemont, P.S., *et al.*, *Proteins* 1, 66-73, 1986; Joyce, C.M., *Curr. Opin. Struct. Biol.* 1: 123-129 (1991). There are some conserved amino acids that are implicated to be responsible for 5'-to-3' exonuclease activity (Gutman and Minton, *Nucl. Acids Res.* 21, 4406-4407, 1993). See *supra*. It is known that 5'-to-3' exonuclease domain is dispensable. The best known example is the Klenow fragment of *E. coli* Pol I. The Klenow fragment is a natural proteolytic fragment devoid of 5'-to-3' exonuclease activity (Joyce, C.M., *et al.*, *J. Biol. Chem.* 257, 1958-1964, 1990). In order to generate an equivalent mutant for *Tne* DNA polymerase devoid of 5'-to-3' exonuclease activity, the presence of a unique *Sph*I site present 680 bases from the *Sst*I site was exploited. pUC-Tne35FY was digested with *Hind*III, filled-in with Klenow fragment to generate a blunt-end, and digested with *Sph*I. The 1.9 kb fragment was cloned into an expression vector pTTQ19 (Stark, M.J.R., *Gene* 51, 255-267, 1987) at the *Sph*I-*Sma*I sites and was introduced into DH10B. This cloning strategy generated an in-frame polymerase clone with an initiation codon for methionine from the

-63-

vector. The resulting clone is devoid of 219 amino terminal amino acids of *Tne* DNA polymerase. This clone is designated as pTTQTne535FY (Fig. 4). The clone produced active heat stable polymerase. No exonuclease activity could be detected in the mutant polymerase as evidenced by lack of presence of unusual sequence ladders in the sequencing reaction. This particular mutant polymerase is highly suitable for DNA sequencing.

Example 11: 5'→3' Exonuclease Deletion and Phe⁷³⁰→Tyr⁷³⁰ Substitution Mutant

In order to generate the 5'-to-3' exonuclease deletion mutant of the *Tne* DNA polymerase Phe⁷³⁰→Tyr⁷³⁰ mutant, the 1.8 kb *SphI-SpeI* fragment of pTTQTne535FY was replaced with the identical fragment of pUC-Tne FY. See Fig. 4. A resulting clone, pTTQTne5FY, produced active heat stable DNA polymerase. As measured by the rate of degradation of a labeled primer, this mutant has a modulated, low but detectable, 5'-to-3' exonuclease activity compared to wild type *Tne* DNA polymerase. M13/pUC Forward 23-Base Sequencing PrimerTM, obtainable from LTI, Rockville, MD, was labeled at the 5' end with [P³²] ATP and T4 kinase, also obtainable from LTI, Rockville, MD, as described by the manufacturer. The reaction mixtures contained 20 units of either wildtype or mutant *Tne* DNA polymerase, 0.25 pmol of labeled primer, 20 mM tricine, pH 8.7, 85 mM potassium acetate, 1.2 mM magnesium acetate, and 8% glycerol. Incubation was carried out at 70°C. At various time points, 10 µl aliquots were removed to 5 µl cycle sequencing stop solution and were resolved in a 6 % polyacrylamide sequencing gel followed by autoradiography. While the wildtype polymerase degraded the primer in 5 to 15 minutes, it took the mutant polymerase more than 60 minutes for the same amount of degradation of the primer.

Example 12: Purification of the Mutant Polymerases

The purification of the mutant polymerases was done essentially as described Example 6, *supra*, with minor modifications. Specifically, 5 to 10 grams of cells expressing cloned mutant *Tne* DNA polymerase were lysed by sonication with a Heat Systems Ultrasonic, Inc. Model 375 machine in a sonication buffer comprising 50 mM Tris-HCl (pH 7.4); 8% glycerol; 5 mM 2-mercaptoethanol, 10 mM NaCl, 1 mM EDTA, and 0.5 mM PMSF. The sonication sample was heated at 75°C for 15 minutes. Following heat treatment, 200 mM NaCl and 0.4% PEI was added to remove nucleic acids. The extract was centrifuged for clarification. Ammonium sulfate was added to 48%, the pellet was resuspended in a column buffer consisting of 25 mM Tris-HCl (pH 7.4); 8% glycerol; 0.5% EDTA; 5 mM 2-mercaptoethanol; 10 mM KCl and loaded on a heparin agarose (LTI) column. The column was washed with 10 column volumes using the loading buffer and eluted with a 10 column volume buffer gradient from 10 mM to 1 M KCl. Fractions containing polymerase activity were pooled and dialyzed in column buffer as above with the pH adjusted to 7.8. The dialyzed pool of fractions were loaded onto a MonoQ (Pharmacia) column. The column was washed and eluted as described above for the heparin column. The active fractions are pooled and a unit assay was performed.

The unit assay reaction mixture contained 25 mM TAPS (pH 9.3), 2 mM MgCl₂, 50 mM KCl, 1 mM DTT, 0.2 mM dNTPs, 500 µg/ml DNase I treated salmon sperm DNA, 21 mCi/ml [α P³²] dCTP and various amounts of polymerase in a final volume of 50 µl. After 10 minutes incubation at 70°C, 10 µl of 0.5 M EDTA was added to the tube. TCA perceptible counts were measured in GF/C filters using 40 µl of the reaction mixture.

-65-

Example 13: Generation of 5'-to-3' exonuclease mutant of full length *Tne* DNA polymerase

1. Identification of Two Amino Acids Responsible for 5'-to-3' Exonuclease Activity

5 *Tne* DNA polymerase contains three enzymatic activities similar to *E. coli* DNA polymerase I: 5'-to-3' DNA polymerase activity, 3'-to-5' exonuclease activity and 5'-to-3' exonuclease activity. This example is directed to the elimination of the 5'-to-3' exonuclease activity in full length *Tne* DNA polymerase. Gutman and Minton (*Nucleic Acids Res.* 1993, 21, 4406-4407) identified six (A-F) conserved 5'-to-3' exonuclease domains containing a total of 10 carboxylates in various DNA polymerases in the polI family. Seven out of 10 carboxylates (in domains A, D and E) have been implicated to be involved in divalent metal ions binding as judged from the crystal structure (Kim et al. *Nature*, 1995, 376, 612-616) of *Taq* DNA polymerase. However, there was no clear demonstration that these carboxylates are actually involved 5'-to-3' exonuclease activity. In order to find out the biochemical characteristics of some of these carboxylates, two of the aspartic acids in domains A and E were chosen for mutagenesis. The following aspartic acids in these two domains were identified:

20 *Tne* DNA polymerase: 5 FLFD⁸GT 10 (domain A) (SEQ ID NO:24)
 Taq DNA polymerase: 15 LLVD¹⁸GH 20 (SEQ ID NO:25)
 and
 Tne DNA polymerase: 132 SLITGD¹³⁷KDML141 (domain E) (SEQ ID NO:26)
 Taq DNA polymerase: 137 RILTAD¹⁴²KDLY146 (SEQ ID NO:27)

25 **2. Isolation of Single Stranded DNA for Mutagenesis**

Single stranded DNA was isolated from pSportTne (see *infra*). pSportTne was introduced into DH5 α F'IQ (LTI, Rockville, MD) by transformation. A single

-66-

colony was grown in 2 ml Circle Grow (Bio 101, CA) medium with ampicillin at 37°C for 16 hrs. A 10 ml fresh media was inoculated with 0.1 ml of the culture and grown at 37°C until the A590 reached approximately 0.5. At that time, 0.1 ml of M13KO7 helper phage (1×10^{11} pfu/ml, LTI) was added to the culture. The infected culture was grown for 75 min. Kanamycin was then added at 50 µg/ml, and the culture was grown overnight (16 hrs.). The culture was spun down. 9 ml of the supernatant was treated with 50 µg each of RNaseA and DNaseI in the presence of 10 mM MgCl₂ for 30 min. at room temperature. To this mixture, 0.25 volume of a cocktail of 3M ammonium acetate plus 20% polyethylene glycol was added and incubated for 20 min. on ice to precipitate phage. The phage was recovered by centrifugation. The phage pellet was dissolved in 200 µl of TE (10 mM Tris-HCl (pH 8) and 1 mM EDTA). The phage solution was extracted twice with equal volume of buffer saturated phenol (LTI, Rockville, MD), twice with equal volume of phenol:chloroform:isoamyl alcohol mixture (25:24:1, LTI, Rockville, MD) and finally, twice with chloroform: isoamyl alcohol (24:1). To the aqueous layer, 0.1 volume of 7.5 M ammonium acetate and 2.5 volume of ethanol were added and incubated for 15 min. at room temperature to precipitate single stranded DNA. The DNA was recovered by centrifugation and suspended in 200 µl TE.

3. Mutagenesis of D⁸ and D¹³⁷

Two oligos were designed to mutagenize D⁸ and D¹³⁷ to alanine. The oligos are: 5' GTAGGCCAGGGCTGTGCCGGCAAAGAGAAATAGTC 3' (D8A) (SEQ ID NO:28) and 5' GAAGCATATCCTTGGCGCCGGTTAT TATGAAAATC 3' (D137A) (SEQ ID NO:29). In the D8A oligo a *Ngo*AIV (bold underlined) and in the oligo D137A a *Kas*I (bold underlined) site was created for easy identification of clones following mutagenesis. 200 pmol of each oligo was kinased according to the Muta-gene protocol (Bio-Rad, CA) using 5 units of T4 Kinase (LTI, Rockville, MD). 200 ng of single stranded DNA was

-67-

annealed with 2 pmol of oligo according to the Muta-gene protocol. The reaction volume was 10 μ l. Following the annealing step, complementary DNA synthesis and ligation was carried out using 5 units of wildtype T7 DNA polymerase (USB, Ohio) and 0.5 unit T4 ligase (LTI). 1 μ l of the reaction was used to transform a MutS *E. coli* (obtainable from Dr. Paul Modrich at the Duke University, NC) and selected in agar plates containing ampicillin. A control annealing and synthesis reaction was carried out without addition of any oligo to determine the background. There were 50-60 fold more colonies in the transformation plates with the oligos than without any oligo. Six colonies from each mutagenic oligo directed synthesis were grown and checked for respective restriction site (*NgoAIV* or *KasI*). For D8A (*NgoAIV*), 4 out of 6 generated two fragments (3 kb and 4.1 kb). Since pSportTne has an *NgoAIV* site near the *f1* intergenic region, the new *NgoAIV* site within the *Tne* DNA polymerase produced the expected fragments. The plasmid was designated as pSportTne*NgoAIV*. For D137A (*KasI*), 5 out of 6 clones produced two expected fragments of 1.1 kb and 6 kb in size. Since pSportTne has another *KasI* site, the newly created *KasI* site generated these two expected fragments. The plasmid was designated as pSportTne*KasI*. Both D8A and D137A mutations were confirmed by DNA sequencing.

4. Reconstruction of the Mutant Polymerase into Expression Vector

During the course of expression of *Tne* DNA polymerase or mutant *Tne* DNA polymerase, a variety of clones were constructed. One such clone was designated as pTTQ Tne SeqS1. This plasmid was constructed as follows: first, similar to above mutagenesis technique glycine 195 was changed to an aspartic acid in pSportTne. A mutation in the corresponding amino acid in *E. coli* DNA polymerase I (polA214, domain F) was found to have lost the 5'-to-3' exonuclease activity (Gutman and Minton, see above). An *SspI* site was created in the mutant polymerase. Second, a 650 bp *SstI-SphI* fragment containing the G195D mutation was subcloned in pUCTne35FY (see *infra*) to replace the wild type fragment.

-68-

This plasmid was called pUCTne3022. Finally, the entire mutant *Tne* DNA polymerase was subcloned from pUCTne3022 into pTTQ18 as *SstI-HindIII* fragment to generate pTTQTneSeqS1. To introduce the mutation D8A or D137A in this expression vector, the 650 bp *SstI-SphI* was replaced with the same *SstI-SphI* fragment from pSportTneNgoAIV or pSportTneKasI. The plasmids were designated as pTTQTneNgo(D8A) and pTTQTneKas(D137A), respectively.

5. Confirmation of the Mutations by DNA Sequencing

DNA sequencing of both mutant polymerases confirmed the presence of the restriction site *NgoAIV* as well as the mutation D8A; and *KasI* site as well as the mutation D137A. Also confirmed by DNA sequencing was the presence of the mutation D323A and the *Eco47III* restriction site in the 3'-to-5' exonuclease region. In addition, confirmed by DNA sequencing was the F730Y mutation and the *HpaI* restriction site in the O-helix region of the mutant *Tne* DNA polymerase.

6. 5'-to-3' exonuclease Activity of the Mutant *Tne* DNA Polymerases

The full length mutant DNA polymerase was purified as described above. The 5'-to-3' exonuclease activity was determined as described in the LTI catalog. Briefly, 1 pmol of labeled (³²P) *HaeIII* digested λ DNA (LTI) was used for the assay. The buffer composition is: 25 mM Tris-HCl (pH 8.3), 5 mM MgCl₂, 50 mM NaCl, 0.01% gelatin. The reaction was initiated by the addition of 0, 2, 4, 6 and 10 units of either wild type or mutant *Tne* DNA polymerase in a 50 μ l reaction. The reaction mix was incubated for 1 hr at 72°C. A 10 μ l aliquot was subjected to PEI-cellulose thin layer chromatography and the label released was quantitated by liquid scintillation. In this assay, both D8A and D137A mutants showed less than 0.01% label release compared to the wild type *Tne* DNA polymerase. The result demonstrates that in both D8A and D137A mutants the 5'-to-3' exonuclease activity has been considerably diminished. Thus, it has been

-69-

confirmed that these two aspartates are involved with the 5'-to-3' exonuclease activity.

Example 14: Generation of double mutants, R722K/F730Y, R722Q/F730Y, R722H/F730Y and R722N/F730Y of Tne DNA polymerase

5 For all mutations, the PCR method was used. A common 5'-oligo, CAC CAG ACG **GGT ACC** GCC ACT GGC AGG TTG (SEQ ID NO:30), was used. This oligo contains a *KpnI* site (shown above in bold italics). The template used for PCR was pTTQTneSeqS1 (Example 13) which already contains the F730Y mutation in the Tne polymerase gene. For the R722K/F730Y mutation, the oligo used was TAT AGA GTA **GTT AAC** CAT CTT TCC AAC CCG TTT CAT TTC TTC GAA CAC (SEQ ID NO:31). For the R722Q/F730Y mutation, the oligo used was TAT AGA GTA **GTT AAC** CAT CTT TCC AAC CCG TTG CAT TTC TTC GAA CAC (SEQ ID NO:32). For the R722N/F730Y mutation, the oligo used was TAT AGA GTA **GTT AAC** CAT CTT TCC AAC CCG GTT CAT TTC TTC GAA CAC (SEQ ID NO:33) and for the R722H/F730Y the oligo used was TAT AGA GTA **GTT AAC** CAT CTT TCC AAC CCG ATG CAT TTC TTC GAA CAC (SEQ ID NO:34). Each of these oligos contains a *HpaI* site (bold italics). The underlined codons were the mutated codons for arginine at the position 722 for respective amino acids. The PCR generated a 318 bp product containing a *KpnI* and a *HpaI* site. The PCR products were digested with *KpnI* and *HpaI* and cloned into pUC-TneFY digested with *KpnI* and *HpaI* to replace the original fragment to generate pUC19TneFY-R722K, pUC19TneFY-R722Q, pUC19TneFY-R722H and pUC19TneFY-R722N. Finally, the *KpnI*-*HindIII* fragment (~800bp) of pTTQTneKasI(D137A) was replaced by the ~800 bp *KpnI*-*HindIII* fragment from these plasmids to generate pTne11 (R722K/F730Y), pTne10 (R722Q/F730Y), pTne13 (R722H/F730Y) and pTne9 (R722N/F739Y), respectively. The mutations were confirmed by DNA sequencing.

-70-

Example 15: Generation of *Tne* DNA Polymerase mutants F730A and F730S

F730A was constructed using PCR. The forward oligo was AAG ATG ***GTT AAC GCG TCT*** ATA ATA TAC GG (SEQ ID NO:35) which contains a *HpaI* site and a *MluI* site (bold italics). The reverse oligo was CAA GAG GCA CAG AGA GTT TCA CC (SEQ ID NO:36) which anneals downstream of *SpeI* present in the *Tne* polymerase gene. The template used for PCR was pTTQ*Tne* KasI (D137A). The 482bp PCR product was digested with *HpaI* and *SpeI* and cloned into pUC-*TneFY* thereby replacing the amino acid tyrosine at position 730 with alanine. This construct was called pUC-*Tne FA*.

10 F730S was constructed by site directed mutagenesis. The oligo was GTA TAT TAT AGA GGA ***GTT AAC*** CAT CTT TCC (SEQ ID NO:37) where a *HpaI* site was created (bold italics). The single stranded DNA used was isolated from pSport-*Tne* that contains the double mutation D137A and D323A. This construct was designated p*Tne* 47. The *Tne* polymerase gene was then cloned as an *SstI* and *HindIII* fragment into the plasmid pUC19 and the resulting clone was designated p*Tne*101.

Example 16: Generation of *Tne* DNA polymerase with a *HpaI* site in front of the amino acid phenylalanine at position 730.

20 A construct of *Tne* polymerase was made using PCR where a *HpaI* restriction enzyme site was introduced into the gene in front of the amino acid phenylalanine at position 730. The forward oligonucleotide was AAG ATG ***GTT AACTTC*** TCT ATA ATA TAC GG (SEQ ID NO:38) which contains a *HpaI* site (shown above in bold italics) and the reverse oligo was the same as in Example 15 above. The template used for PCR was p*Tne*33 which contains the *Tne* polymerase gene with D137A and D323A mutations cloned in pUC19. The 482bp PCR product was digested with *HpaI* and *SpeI* and was used to replace the corresponding fragment in p*Tne*101 (see example 15). The construct was

-71-

sequenced to verify that the amino acid at position 730 was indeed phenylalanine and the plasmid was numbered pTne106.

Example 17: Generation of double mutants R722Y/F730A and R722L/F730A of the Tne DNA polymerase.

5 For both the mutations PCR method was used. The common 5' oligo was the same as in Example 14. For R722Y/F730A mutation the oligo used was TAT AGA GTA ***GTT AAC*** CAT CTT TCC AAC CCG ***GTA*** ***CAT GTC*** TTC GTT CAC (SEQ ID NO:39). For R722L/F730A mutation the oligo used was TAT AGA GTA ***GTT AAC*** CAT CTT TCC AAC CCG ***CAA*** ***CAT GT C*** TTC GTT CAC (SEQ ID NO:40). Each of these oligos contain a *HpaI* site (shown above in bold italics). The underlined codons were the mutated codons for arginine at the position 722 for respective amino acids. An *AflIII* site was also created (shown above in bold italics next to the underlined codon) in order to confirm the mutation. The PCR generated a 318 bp product containing a *KpnI* and a *HpaI* site. 10 The PCR products were digested with *KpnI* and *HpaI* and cloned into pUC-TneFA (see example 15). The constructs were named as pUCTneYA and pUCTneLA. 15

Example 18: Generation of Tne DNA Polymerase mutants R722Y and R722L.

20 The plasmid pTne 106 (see example 16) was digested with *HpaI* and *KpnI* and the 318 bp fragment was replaced with the corresponding fragment from pUCTneYA or pUCTneLA (see Example 17) to generate the mutants R722Y or R722L. In these constructs the amino acid at position 730 is the same as wild type *Tne* (phenylalanine). The constructs were sequenced to confirm the R722Y and 25 the R722L mutations. The *Tne* DNA polymerase gene was then cloned as a *SstI/HindIII* fragment into the plasmid pSport1.

-72-

Example 19: Generation of *Tne* DNA Polymerase mutants R722K, R722Q and R722H.

5 The construct pTne 106 (see example 16) was digested with *Hpa*I and *Kpn*I and the 318 bp fragment was replaced with the corresponding fragment from the construct pUC19TneFY-R722K, pUC19TneFY-R722H or pTne10 (see Example 14), to generate the mutants R722K, R722H and R722Q. The constructs were sequenced to confirm the mutations. The *Tne* DNA polymerase gene was then subcloned into the vector pSport1 as a *Sst*I/*Hind*III fragment.

Example 20: Purification of the mutant *Tne* DNA Polymerases

10 The purification of the mutants of *Tne* DNA polymerase was carried out based on the method described above with minor modifications. Two to three grams of cells expressing cloned mutant *Tne* DNA polymerase were resuspended in 15-20 ml of sonication buffer (50 mM Tris-HCl, pH 8.0, 10% glycerol, 5mM 2-mercaptoethanol, 50 mM NaCl, 1 mM EDTA, and 0.5 mM PMSF and
15 sonicated with a 550 Sonic Dismembrator (Fisher Scientific). The sonicated sample was heated at 82°C for 20 min and then cooled in ice-water for 5 min. In the sample, 20 mM NaCl and 0.2% PEI were added and centrifuged at 13,000 rpm for 10 min. Ammonium sulfate (305g/L) was added to the supernatant. The pellet was collected by centrifugation and resuspended in 4 ml of MonoQ column
20 buffer (50 mM Tris-HCl, pH 8.0, 10% glycerol, 5mM 2-mercaptoethanol, 50 mM NaCl, and 1 mM EDTA). The sample was dialyzed against one liter of MonoQ buffer overnight. Following the centrifugation at 13,000 rpm to remove any insoluble materials, the sample was loaded onto a MonoQ column (HR5/5, Pharmacia). The column was washed with MonoQ column buffer to baseline of
25 OD₂₈₀ and then eluted with a linear gradient of 50-300 mM NaCl in 20 ml MonoQ column buffer. The fractions were analyzed by 8% SDS-PAGE and the *Tne* DNA

-73-

polymerase activity determined as described earlier. The fractions containing active and pure *Tne* DNA polymerase were pooled.

Example 21: Generation of *Taq* DNA Polymerase Mutants R659K, R659H and R659Y

5 A 2.5 kb portion of the gene encoding *Taq* DNA polymerase (Figure 5) was cloned as a *Hind* III-*Xba* I fragment into M13mp19. Site directed mutagenesis was performed using the BioRad mutagene kit (BioRad California) using the following oligonucleotides:

10 CTTGGCCGCCCGATGCATCAGGGGGGTC (SEQ ID NO:41) for the R659H mutation where an *Nsi*I site was created (see bold italics);

CTTGGCCGCCCGCTTCATGAGGGGGTCCAC (SEQ ID NO:42) for the R659K mutation where a *Bsp*HI site was created (see bold italics); and

CTTGGCCGCCCTGTACATCAGGGGGGTC (SEQ ID NO:43) for the R659Y mutation where a *Bsr*GI site was created (see bold italics).

15 For each mutation, six clones were screened by analyzing the M13RF DNA for the expected restriction sites. Mutations were confirmed by DNA sequencing. DNA shown to contain the mutation by the presence of the expected restriction site was digested with *Ngo*AIV and *Xba* I and the approximately 1600 base pair fragment was used to replace corresponding fragment in the wildtype
20 *Taq* DNA polymerase gene. These constructs were made in a plasmid containing *Taq* polymerase gene under the control of Tac promoter (pTTQ *Taq*) to generate pTTQ *Taq* (R659K), pTTQ *Taq* (R659H) and pTTQ *Taq* (R659Y). These plasmids were transformed into *E. coli* DH10B (LTI).

-74-

Example 22: Construction of Tne polymerase mutants containing F730S and F730T

Single stranded DNA was isolated from pSportTne (Tne35) containing D137A and D323A mutations as described in the section 2 of example 13. These D137 and D323A mutations rendered Tne DNA polymerase devoid of 5'-exonuclease and 3'-to-5'-exonuclease activities, respectively. Thus, Tne 35 is devoid of both exonuclease activities. The site-directed mutagenesis was done following the protocol described in section 3 of Example 13. The oligos used were 5' GTA TAT TAT AGA GGA GTT AAC CAT CTT TCC 3' (SEQ ID NO:37) for F730S and 5' GTA TAT TAT AGA GGT GTT AAC CAT CTT TCC 3' (SEQ ID NO:44) for F730T. Each of these two oligos contain a diagnostic *HpaI* site for screening of mutants in the MutS strain. The mutant plasmids were transferred to DH10B strains. The mutations were finally confirmed by DNA sequencing. The mutant polymerases were purified by the procedure as described in Example 20.

Example 23: Determination of the Activity of Non-templated One Base Addition for Tne and Taq DNA Polymerase by Primer Extension Assay

The following 34-mer primer was ³²P labeled at the 5' end with [γ -³²P] ATP and T4 polynucleotide kinase by standard protocol (Molecular Cloning, A Laboratory Manual, Cold Spring Harbor, NY):

5'-GGGAGACCGGAATTCTCCTTCATTAATTCCTATA-3' (SEQ ID NO:45)

The unincorporated ATP was removed by a BioRad P6 column(1.0 ml). The labeled primer was annealed to the following homogenous (purified) 48-mer template:

-75-

5'-TGGAGACCCTGGA ACTATAGGAATTAATGAAGGAGAATTCCGGT
CTCCC-3' (SEQ ID NO:46).

Wildtype or mutant DNA polymerases (0.125-1.0 unit) were incubated at
72° C for 2 min in 20 mM Tris-HCl (pH8.3), 1.5 mM MgCl₂, 50 mM KCl, 1.0
5 mM DTT, 200 uM of dCTP, dGTP, TTP, dATP, and 0.02 pmol of the annealed
primer-template. After addition of sequencing stop buffer and heated at 90°C for
2 min, the mixture was loaded onto 10% polyacrylamide-7 M urea. Following the
electrophoresis, the gel was dried and the reaction products were analyzed by
autoradiography. The non-templated one base addition products shown in Figure
10 6 were quantified by a PhosphorImager (Molecular Dynamics).

	The DNA polymerases	% of N+1
	1 D137A	18.5
	2 D137A D323A	78.5
15	3 D137A D323A R722K	0.7
	4 D137A D323A R722Y	0.7
	5 D137A D323A R722L	5.7
	6 D137A D323A R722H	1.2
20	7 D137A D323A R722Q	1.4
	8 D137A D323A F730Y	61.3
	9 D137A D323A R722 K F730Y	6.8
	10 D137A D323A R722H F730Y	2.1
	11 D137A D323A R722Q F730Y	6.1
25	12 D137A D323A R722N F730Y	15.9
	13 D137A D323A F730S	8.3
	14 D137A D323A F730T	24.2

-76-

	Taq DNA Polymerases	% of N+1
	1 W.T.	37
	2 R659K	1.4
	3 R659Y	0.9
5	4 R659H	0.5
	5 F667Y	39.1

Example 24: Comparison of DNA Synthesis by Taq and Tne

To examine its propensity to add a nontemplated nucleotides to the 3' termini of PCR products, *Tne* DNA polymerase (5'exo⁻, 3'exo⁻) was compared side-by-side with *Taq* DNA polymerase in amplifications of short tandem repeats at 23 different marker loci (see Table 1). Reactions comprising 20 mM TRIS-HCl, pH 8.4, 50 mM KCl, 1.5 mM MgCl₂, 200 mM each dNTP, 200 nM [³²P] α-dATP, 200 nM each of the upper and lower primers, 25 ng of human DNA, 0.1% nonionic detergent and 1 unit of DNA polymerase (in a volume of 25 ml) were assembled on ice. Published sequences for upper and lower primers for each locus, as shown in Table 1, were used for all amplifications.

Reactions were loaded into a Perkin Elmer model 9600 thermocycler preheated to 94°C and PCR was done using standard cycling conditions (1 minute pre-denaturation at 94°C; 30 cycles of 30 seconds at 94°C, 30 seconds at 55°C, and 1 minute at 72°C; 1 minute post-extension at 72°C; overnight soak at 4°C). A portion of each reaction was mixed with an equal volume of 95% formamide containing dyes to indicate the progress of electrophoresis. Samples were heated to 90°C for 2 min, and 5 ml of each was loaded on a 6% denaturing polyacrylamide gel. Sequencing ladders were loaded to provide size markers, and electrophoresis was performed at 70 watts. After electrophoresis the gel was transferred to filter paper and dried. Autoradiography and phosphoimage analysis was performed to visualize the PCR products and estimate the percentage of

-77-

product which contained the added nucleotide by direct comparison of bands produced by each enzyme.

5 Examples of the side-by-side comparisons of amplification products produced by *Taq* DNA polymerase and *Tne* DNA polymerase are shown in Figure 7 for the CD4 locus and in Figure 8 for the D20S27 locus. At both of these loci, a significant portion of the *Taq* PCR product contained an extra non-templated nucleotide (n+1), while *Tne* polymerase demonstrated no apparent non-templated nucleotide incorporation for either the CD4 locus (Figure 7) or the D20S27 locus (Figure 8). Complete results for the 23 marker loci examined are summarized in Table 2. In PCRs using *Taq* DNA polymerase, a portion of the amplification product contained an extra non-templated nucleotide (n+1) at every locus examined. In PCRs using *Tne* DNA polymerase, however, no detectable portion of the product at any of the loci examined contained an additional non-templated nucleotide. These results indicate that *Tne* DNA polymerase, in contrast to *Taq* DNA polymerase, is substantially reduced in the ability to add a nontemplated 3' terminal nucleotide to the growing strand. Since the *Tne* DNA polymerase used in these amplifications was a 3'exo- mutant (*i.e.*, it was substantially reduced in 3' exonuclease activity), these results are consistent with the notion that the *Tne* polymerase was unable to add the extra nucleotide to the product rather than adding the nucleotide and then removing it via a 3' exonuclease activity.

10

15

20

Table 1. Primers Used in Example 24.

Locus	Upper Primer (SEQ ID NO:)	Lower Primer (SEQ ID NO:)	Reference
D13S71	GTATTTTGGTATGCTTGTGC (47)	CTATTTTGGAAATATATGTGCCT (48)	Nucl. Acids Res. 18:4638 (1990)
D1S103	ACGAACATTTCTACAAGTTAC (49)	TTTCAGAGAAACTGACCCTGT (50)	Nucl. Acids Res. 18:2199 (1990)
D1S87	GATAAATGCCAAACAIGTTGT (51)	TGCTCTCAGGATTTCCCTCCA (52)	Nucl. Acids Res. 18:4640 (1990)
D2S136	AGCTTGAGACCTCTGTGTCC (53)	ATTCAGAAGAAACAGTGATGGT (54)	Nature Genet. 7:246-339 (1994)
CD4	TTGGAGTCGCAAGCTGAACTAGC (55)	GCCTGAGTGACAGAGTGAGAACC (56)	Nucl. Acids Res. 19:4791 (1991)
PLA2A	CCCACTAGGTTGTAAAGCTCCATGA (57)	TACTATGTGCCAGGCTCTGTCTTA (58)	Nucl. Acids Res. 18:7468 (1990)
D19S49	ACTCATGAAGGTGACAGTTC (59)	GTGTTGTGACCTATTGCAAT (60)	Nucl. Acids Res. 18:1927 (1990)
D4S175	ATCTCTGTCCCTCCCTGTT (61)	CTTATTTGGCCCTTGAAGGTAG (62)	Genomics 14:209-219 (1992)
APOC2AC	AGCCCGTGTGGAAACCATGACTG (63)	TACATAGCGAGACTCCATCTCCC (64)	Hum. Genet. 83:245-251 (1989)
D20S27	TTTATGCGAGCGTATGGATA (65)	CACCACCATTGATCTGGAAG (66)	Nucl. Acids Res. 18:2202 (1990)
D15S127	CCAACCACACTGGGAA (67)	AACAGTTGCCACCGGT (68)	Nature Genet. 7:246-339 (1994)
D4S398	CATGAAATGCTGACTGGGTA (69)	TCAATTTATGTGCAGCCAAT (70)	Nature Genet. 7:246-339 (1994)
APOC2	CATAGCGAGACTCCATCTCC (71)	GGGAGAGGGCAAAGATCGAT (72)	Am. J. Hum. Genet. 44:388-396 (1989)
D10S89	AACACTAGTGACATTTTICA (73)	AGCTAGGCCCTGAAGGCTTCT (74)	Nucl. Acids Res. 18:4637 (1990)
VWA	CCCTAGTGGATGATAAGAATAATC (75)	GGACAGATGATAAATACATAGGATGGATGG (76)	Hum. Molec. Genet. 1:287 (1992)
D16S401	TTCCTTACAAACACTGCCCC (77)	ATTTGGATGGCTTGACAGAG (78)	Nature Genet. 7:246-339 (1994)
D7S440	ACATTTCTAAGACTTTCCCAAT (79)	AGAGCATGCACCCCTGAAATG (80)	Nucl. Acids Res. 18:4039 (1990)
D4S174	AAGAACCATGCGATACGACT (81)	CATTCCTAGATGGTAAAGC (82)	Nucl. Acids Res. 18:4636 (1990)
D16S520	GCCTAGTCAATACGAGCGG (83)	TCCACAGCCCATGTAAACC (84)	Nature Genet. 7:246-339 (1994)
D16S511	CCCCGGAGCAAGTICA (85)	CAGCCCAAAGCCAGATTA (86)	Nature Genet. 7:246-339 (1994)
D21S11	ATATGTGAGTCAATTCCTCCAAG (87)	TGTATTAGTCAATGTTCTCCAG (88)	Hum. Molec. Genet. 1:67 (1992)
THO1	CAGCTGCCCTAGTCAGCAC (89)	GCTTCCGAGTGCAGGTCACA (90)	Nucl. Acids Res. 19:3753 (1991)
ACTBP2	ATTTCTGGCGGCACAAAGAGTGA (91)	ACATCTCCCTACCCGCTATA (92)	Nucl. Acids Res. 20:1432 (1992)

Table 2. Non-templated 3' Terminal Nucleotide Addition by *Taq* and *Tne* DNA Polymerases at 23 Microsatellite DNA Loci.

Locus	Repeat Type	<i>Taq</i> (% n+1)	<i>Tne</i> (% n+1)
D13S71	dinucleotide	100	0
5 D1S103	dinucleotide	75-100	0
D15S87	dinucleotide	30-50	0
D2S136	dinucleotide	90-100	0
HUMCD4	pentanucleotide	50	0
HUMPLA2A	trinucleotide	25	0
10 D19S49	dinucleotide	75	0
D4S175	dinucleotide	75	0
APOC2AC	dinucleotide	50	0
D20S27	dinucleotide	50	0
D15S127	dinucleotide	100	0
15 D4S398	dinucleotide	50	0
APOC2	dinucleotide	50	0
D10S89	dinucleotide	75-100	0
HUMVWA	tetranucleotide	90	0
D16S401	dinucleotide	100	0
20 D7S440	dinucleotide	90	0
D4S174	dinucleotide	75	0
D16S520	dinucleotide	100	0
D16S511	dinucleotide	100	0
HUMD21S11	tetranucleotide	100	0
25 HUMTHO1	tetranucleotide	75	0
HUMACTBP2	tetranucleotide	25	0

-80-

Example 25: Comparison of DNA Synthesis by *Tne* and Other Thermostable Enzymes

To further evaluate the differences in the propensities of *Tne* and other thermostable DNA polymerases to add non-templated 3' terminal nucleotides to PCR products, side-by-side amplifications were performed using a single marker locus D1S103 and a variety of thermostable enzymes, including 3' exonuclease deficient (3'exo-) enzymes, and 3' exonuclease competent (3'exo+) enzymes. PCR amplifications, electrophoresis and analysis were performed as described for Example 24, using 200 nM of D1S103-specific upper and lower primers.

Results for the amplifications using 3'exo- DNA polymerases are shown in Table 3. With the exception of *Tne*(3'exo-), all of the 3'exo- DNA polymerases examined exhibited a propensity to add a non-templated 3' terminal nucleotide (n+1) to the PCR product. For *Taq* and *Tbr* DNA polymerases, up to 100% of the PCR products contained an additional non-templated 3' terminal nucleotide, while *Vent*, *Deep Vent*, and *Dtok* 3'exo- mutants polymerases added this non-templated nucleotide to 25-100% of the PCR products. In contrast, the 3'exo- mutant of *Tne* DNA polymerase was substantially reduced in the ability to add a nontemplated 3' terminal nucleotide to the DNA molecule; none of the PCR products from reactions using *Tne*(3'exo-) had an additional non-templated nucleotide at their 3' termini.

Results from amplifications using 3'exo+ DNA polymerases are shown in Table 4. Five polymerases were examined as well as two commercially available enzyme mixes (mixtures of a primary 3'exo- polymerase and a secondary 3'exo+ polymerase). At this locus, the 3'exo+ DNA polymerases (*Tne*, *Tma*, *Pfu*, *Pwo* and 9°North) yielded product which did not contain an extra non-templated nucleotide. The enzyme mixtures (*Elongase* and *Expand HiFi*) yielded a mixture of products with and without an additional non-templated nucleotide. Together, these results indicate that *Tne* polymerases, whether 3'exo- or 3'exo+, are substantially reduced in the ability to add a nontemplated 3' terminal nucleotide to the DNA molecule. Moreover, of the preferred 3'exo- polymerases, only *Tne*(3'exo-) was substantially reduced in this activity, indicating its favorableness

-81-

in PCR applications where non-templated nucleotide addition to the amplification product is undesirable.

Table 3. Non-templated 3' Terminal Nucleotide Addition by 3'exo- DNA Polymerases.

5

Enzyme	n Sized Fragment	n+1 Sized Fragment
<i>Tne(3'exo-)</i>	+	-
<i>Taq</i>	-	+
<i>Vent(3'exo-)</i>	+	+
<i>Deep Vent(3'exo-)</i>	+	+
<i>Dtok(3'exo-)</i>	+	+
<i>ThermolaseTbr</i>	-	+

10

Table 4. Non-templated 3' Terminal Nucleotide Addition by 3'exo+ DNA Polymerases.

15

Enzyme	n Sized Fragment	n+1 Sized Fragment
<i>Tne(3'exo+)</i>	+	-
<i>U1Tma</i>	+	-
<i>Pfu</i>	+	-
<i>Pwo</i>	+	-
<i>9°North</i>	+	-
<i>Elongase</i>	+	+
<i>Expand HiFi</i>	+	+

20

Example 26: Comparison of DNA Synthesis by *Tne* Mutants

25

To examine the utility of *Tne* DNA polymerase and various mutants thereof in amplification of microsatellite DNA sequences, the experiments described in Example 25 were repeated with 11 different *Tne* DNA polymerase

-82-

mutants. Of these mutants, 3 were 5'exo+, while the remainder were 5'exo- either due to N-terminal deletions of the protein, or to point mutations in the 5' exonuclease domain of the polymerase.

5 As shown in Table 5, use of the 5'exo- *Tne* mutants resulted in productive amplifications, yielding PCR products with no non-templated 3' terminal nucleotide additions. Results were identical for all seven *Tne*(3'exo-/5'exo-) polymerase mutants, as well as for the single *Tne*(3'exo+/5'exo-) mutant tested. Results with 5'exo+ *Tne* mutants were inconclusive under the conditions tested.

10 These results indicate that the mutants of *Tne* DNA polymerase tested in the present studies are substantially reduced in the ability to add nontemplated 3' terminal nucleotides to the growing strand, particularly a DNA template comprising a microsatellite DNA sequence or an STR.

Table 5. Non-templated 3' Terminal Nucleotide Addition by *Tne* DNA Polymerase Mutants

15	Enzyme	5'exo Activity	3'exo Activity	n Sized Fragment	n+1 Sized Fragment
	<i>Tne</i> N' Δ 219, D323A	-	-	+	-
	<i>Tne</i> N' Δ 283, D323A	-	-	+	-
	<i>Tne</i> N' Δ 192, D323A	-	-	+	-
	<i>Tne</i> D137A, D323A	-	-	+	-
20	<i>Tne</i> D8A, D323A	-	-	+	-
	<i>Tne</i> G195D, D323A	-	-	+	-
	<i>Tne</i> G37D, D323A	-	-	+	-
	<i>Tne</i> N' Δ 283	-	+	+	-

-83-

Example 27: Fluorescent Analysis of DNA Synthesis by *Tne* and *Taq* DNA Polymerases

In an alternative analysis approach, the propensities of *Taq* DNA polymerase and *Tne* DNA polymerase to add non-templated nucleotides to the PCR products were compared using fluorescent detection. The polymerases were compared in side-by-side amplifications utilizing a commonly used commercially available marker panel (ABI Prism Linkage Mapping Set Panel 21), examining ten different loci. Reaction mixtures (15 ml) containing 1.5 mM MgCl₂, 250 mM of each deoxynucleoside triphosphate, 333 nM of each primer, 50 ng of human DNA and 0.6 units of *Taq* or *Tne* DNA polymerase were assembled on ice. Reactions were loaded into a Perkin Elmer model 9600 thermocycler preheated to 95°C, and PCR was performed using recommended cycling conditions (5 minutes pre-denaturation at 95°C; 10 cycles of 15 seconds at 95°C, 15 seconds at 55°C, and 60 seconds at 72°C; and 20 cycles of 15 seconds at 89°C, 15 seconds at 55°C, and 60 seconds at 72°C). Two sets of extension reactions were conducted for each locus, one with a 10 minute post-extension incubation at 72°C followed by an overnight soak and storage at 4°C (conditions which favor nontemplated 3' nucleotide addition), the other with no post-extension incubation followed by immediate storage at -20°C (conditions which inhibit nontemplated 3' nucleotide addition). A portion of each reaction was diluted, mixed with loading cocktail, heat denatured and loaded on an 8% polyacrylamide sequencing gel. The ABI 373 Stretch Automated Sequencer was run for 5-6 hours at 15W in order to obtain single base resolution, and data were analyzed using GeneScan software. Areas of the peaks recognized by the software were used to estimate the percentage of nontemplated 3' nucleotide addition ("n+1") for each locus by the two polymerases under the two different extension conditions. The total area under the allelic peaks was used to compare the yields of specific PCR product obtained in *Tne* and *Taq* amplifications, and yields produced by *Tne* polymerase were expressed for each locus as a percentage of those produced by *Taq* polymerase. Table 6 summarizes the results obtained.

-84-

Table 6. Comparison of DNA Amplification by Taq and Tne DNA Polymerases by Fluorescent Detection

	locus	color	expected size	cycling conditions	Taq pattern	Tne pattern	Tne yield (% Taq)
5	D16S40 5	blue	107-145	no final ext	0% n+1	100% n	89%
				10' final	94% n+1	100% n	178%
				ext			
	D15S12 7	green	114-148	no final ext	53% n+1	100% n	133%
				10' final	100%	100% n	142%
				ext	n+1		
	D16S52 0	yellow	144-160	no final ext	40% n+1	98% n	275%
				10' final	100%	100% n	252%
				ext	n+1		
10	D16S51 1	green	182-222	no final ext	62% n+1	100% n	51%
				10' final	100%	100% n	160%
				ext	n+1		
	D16S41 1	blue	215-235	no final ext	0% n+1	100% n	218%
				10' final	64% n+1	100% n	257%
				ext	67% n+1	95% n	305%
				10' final ext			
15	D15S13 1	yellow	237-275	no final ext	0% n+1	100% n	48%
				10' final	69% n+1	95% n	231%
				ext			
	D15S13 0	blue	280-294	no final ext	0% n+1	100% n	101%
	D16S50 3	yellow	294-310	no final ext	0% n+1	100% n	102%
				10' final	73% n+1	100% n	166%
				ext			
20	D15S11 7	green	316-334	no final ext	17% n+1	100% n	130%
				10' final	77% n+1	100% n	326%
				ext			
	D16S51 5	blue	320-350	no final ext	32% n+1	100% n	486%
				10' final	100%	100% n	298%
				ext	n+1		

-85-

The results shown in Table 6 confirm that under conditions favoring ("10' final ext") or inhibiting ("no final ext") 3' nontemplated nucleotide addition, *Tne* DNA polymerase produced PCR products that were 95-100% free from non-templated nucleotide addition ("n") for each locus examined. *Taq* DNA polymerase, however, demonstrated significant addition of nontemplated nucleotides under inhibiting conditions in most loci tested, while under permissive conditions well over half, and in some cases all, of the PCR product produced by *Taq* DNA polymerase demonstrated an additional nontemplated 3' nucleotide. Furthermore, under most conditions the amount of PCR product yielded by *Tne* DNA polymerase was at least as high as that of *Taq* DNA polymerase, and for some loci was 3- to 4-fold higher.

Figure 9 shows two examples of electropherogram gel scans, aligned by PCR product size, comparing the PCR products obtained with *Taq* and *Tne* polymerases with a 10-minute final extension. For the D15S153 locus, *Taq* exhibited non-templated nucleotide addition to 40% of the PCR product (Figure 39), while *Tne* exhibited no such addition of non-templated nucleotides (Figure 9B). Similar results were obtained with the D15S127 locus: 53% of the *Taq* PCR products demonstrated non-templated nucleotide addition (Figure 9C), while none of the *Tne* PCR products demonstrated non-templated nucleotide addition (Figure 9D). These results demonstrate the difficulty in identifying alleles in a heterogeneous pattern as generated by *Taq* amplification, compared to the more homogeneous, simple pattern generated by amplification with *Tne*.

Together with Examples 24-26, these results indicate that *Tne* DNA polymerase and the mutants thereof tested in the present studies are substantially reduced in the ability to add a nontemplated 3' terminal nucleotide to DNA templates, particularly DNA templates comprising microsatellite DNA sequences or STRs. Conversely, *Taq* DNA polymerase demonstrates significant addition of nontemplated 3' nucleotides to PCR products.

-86-

Example 28: Comparison of *Taq* and *Tne*

To examine the ability of a truncated form of *Tne* DNA polymerase (N Δ 283, 5'exo-, 10% 3'exo activity) to add a nucleotide to the end of the PCR product, the enzyme was compared side-by-side with wild type *Taq* DNA polymerase in amplifications of short tandem repeats at 5 different marker loci. A portion of ABI Prism Linkage Mapping Set Panel 21 was used for the primer sets for the loci. 15 ul reactions (20 mM Tris-HCl, pH 8.4, 50 mM KCl, 1.5 mM MgCl₂, 200 uM each dNTP, 333 nM each primer, 60 ng human DNA, 0.1% nonionic detergent, 0.6 U DNA polymerase) were assembled on ice.

Reactions were loaded into a Perkin Elmer model 9600 thermocycler preheated to 95 °C and PCR was done using recommended cycling conditions (5 min. pre-denaturation at 95°C; 10 cycles of 15 sec at 95°C, 15 sec at 55 °C, and 60 sec at 72°C; 20 cycles of 15 sec at 89 °C, 15 sec at 55 °C, and 60 sec at 72°C; 10min final extension at 72°C). A portion of each reaction was diluted, mixed with loading cocktail, heat denatured and loaded on a 8% sequencing gel. The ABI 373 Stretch Automated Sequencer was run for 5-6hr at 15W in order to obtain 1base resolution. Data was analyzed using GeneScan software. Areas of the peaks recognized by the software were used to estimate the percent of extranucleotide addition. Table 7 summarizes the results obtained. Examples of the electropherogram data is shown in Figure 10.

Table 7: Percent extranucleotide addition exhibited by *Taq* and *Tne* DNA polymerases at specific loci.

Locus	<i>Taq</i> (% n+1)	<i>Tne</i> (% n+1)
D16S405	46	0
D16S401	100	45
D16S520	63	0
D15S131	51	0
D16S411	53	0

-87-

Example 29: Comparison of Tne Mutants

In order to evaluate the effect of amino acid substitutions in *Tne* DNA polymerase in regard to extra nucleotide addition, different mutations at position F730 in the untruncated polymerase were compared in side-by-side amplifications with *Taq*(wild type) and a truncated *Tne*(N Δ 219, D323A, F730Y) utilizing a portion of ABI Prism Linkage Mapping Set Panel 21. Six loci were examined. 15 ul reactions (20 mM Tris-HCl, pH 8.4, 50 mM KCl, 1.5 mM MgCl₂, 200 uM each dNTP, 333 nM each primer, 50-60 ng human DNA, 0.1% nonionic detergent, 0.15-0.6 U DNA polymerase) were assembled on ice.

Reactions were loaded into a Perkin Elmer model 9600 thermocycler preheated to 95 °C and PCR was done using recommended cycling conditions (5 min. pre-denaturation at 95°C; 10 cycles of 15 sec at 95°C, 15 sec at 55 °C, and 60 sec at 72 °C; 20 cycles of 15 sec at 89°C, 15 sec at 55 °C, and 60 sec at 72°C; 10min final extension at 72°C). A portion of each reaction was diluted, mixed with loading cocktail, heat denatured and loaded on a 8% sequencing gel. The ABI 373 Stretch Automated Sequencer was run for 5-6hr at 15W in order to obtain 1base resolution. Data was analyzed using GeneScan software. Areas of the peaks recognized by the software were used to estimate the percent of extranucleotide addition. Table 8 summarizes the results obtained. An example of the electropherogram data is shown in Figure 11.

Table 8: Percent extranucleotide addition exhibited by mutant *Tne* DNA polymerases at specific loci.

mutant:	locus:	D16S405	D16S401	D15S131	D15S127	D16S511	D15S153
<i>Taq</i> (wild type)		46%	100%	51%	100%	100%	100%
Tne-1 (N'Δ219, D323A, F730Y)		0%	0%	0%	0%	0%	0%
5 Tne-35 (D137A, D323A)		0%	52%	0%	0%	0%	0%
Tne-18 (D137A, D323A, F730Y)		0%	2%	0%	0%	0%	0%
Tne-13 (D137A, D323A, R722H, F730Y)		0%	0%	0%	0%	0%	0%
Tne-14 (D137A, D323A, F730A)	nd	nd	0%	0%	nd	nd	nd
10 Tne-47 (D137A, D323A, F730S)	0%	0%	0%	0%	0%	0%	0%
Tne-48 (D137A, D323A, F730T)	0%	0%	0%	0%	0%	0%	0%

Example 30: Comparison of Tne and Taq Mutants

In order to evaluate the effect of amino acid substitution at position F667 in *Taq* DNA polymerase (equivalent to F730 in *Tne* DNA polymerase) in regard to extra nucleotide addition, a commercially available mutant of *Taq* DNA polymerase (*Taq* FS) (N^Δ3, G46D, F667Y) was compared in side-by-side amplifications with *Taq* DNA polymerase (wild type) and *Tne*-1 DNA polymerase (N^Δ219, D323A, F730Y). Three loci were examined (a portion of ABI Prism Linkage Mapping Set Panel 21). 15 ul reactions (20 mM Tris-HCl, pH 8.4, 50 mM KCl, 1.5 mM MgCl₂, 200 uM each dNTP, 333 nM each primer, 60 ng human DNA, 0.1% nonionic detergent, 0.6 U DNA polymerase) were assembled on ice.

Reactions were loaded into a Perkin Elmer model 9600 thermocycler preheated to 95 °C and PCR was done using recommended cycling conditions (5 min. pre-denaturation at 95 °C; 10 cycles of 15 sec at 95 °C, 15 sec at 55 °C, and 60 sec at 72 °C; 20 cycles of 15 sec at 89 °C, 15 sec at 55 °C, and 60 sec at 72 °C; 10min final extension at 72 °C). A portion of each reaction was diluted, mixed with loading cocktail, heat denatured and loaded on a 8% sequencing gel. The ABI 373 Stretch Automated Sequencer was run for 5-6hr at 15W in order to obtain 1base resolution. Data was analyzed using GeneScan software. Areas of the peaks recognized by the software were used to estimate the percent of extranucleotide addition. Table 9 summarizes the results obtained. Examples of the electropherogram data are shown in Figure 12.

-90-

Table 9: Percent extranucleotide addition exhibited by Taq and Tne DNA polymerases at specific loci.

locus	Taq(% n+1)	TaqFS(% n+1)	Tne-1(% n+1)
D16S411	48	0	0
D15S127	100	31	5
D15S153	100	29	0

Example 31: Comparison of Tne Mutants

In order to evaluate the effect of amino acid substitutions at position R722 in *Tne* DNA polymerase in regard to extranucleotide addition, different mutations in the polymerase were compared in side-by-side amplifications utilizing a portion of ABI Prism Linkage Mapping Set Panel 21. Six loci were examined. 15 ul reactions (20 mM Tris-HCl, pH 8.4, 50 mM KCl, 1.5 mM MgCl₂, 200 uM each dNTP, 333 nM each primer, 50-60 ng human DNA, 0.1% nonionic detergent, 0.2-0.6 U DNA polymerase) were assembled on ice.

Reactions were loaded into a Perkin Elmer model 9600 thermocycler preheated to 95 °C and PCR was done using recommended cycling conditions (5 min. pre-denaturation at 95 °C; 10 cycles of 15 sec at 95 °C, 15 sec at 55 °C, and 60 sec at 72 °C; 20 cycles of 15 sec at 89 °C, 15 sec at 55 °C, and 60 sec at 72 °C; 10min final extension at 72 °C). A portion of each reaction was diluted, mixed with loading cocktail, heat denatured and loaded on a 8% sequencing gel. The ABI 373 Stretch Automated Sequencer was run for 5-6hr at 15W in order to obtain 1base resolution. Data was analyzed using GeneScan software. Heights of the n and n+1 peaks recognized by the software were used to estimate the percent of extranucleotide addition. Table 10 summarizes the results obtained. An example of the electropherogram data is shown in Figure 13.

-91-

Table 10: Percent extranucleotide addition exhibited by mutant Tne DNA polymerases at specific loci.

mutant:	locus:	D16S405	D16S401	D15S131	D15S127	D16S511	D15S153
Tne-35 (D137A, D323A)		0%	54%	0%	0%	0%	0%
Tne-109 (D137A, D323A, R722Y)		0%	0%	0%	0%	0%	0%
Tne-110 (D137A, D323A, R722L)		0%	0%	0%	0%	0%	0%
Tne-114 (D137A, D323A, R722K)		0%	0%	0%	0%	0%	0%
Tne-115 (D137A, D323A, R722Q)		0%	0%	0%	0%	0%	0%
Tne-116 (D137A, D323A, R722H)		0%	0%	0%	0%	0%	0%

-92-

Example 32: Generation of *Tne* DNA Polymerase Mutant K726R

The mutation of the *Tne* polymerase was done by essentially the same procedure as described above in Example 13. The single-stranded DNA was isolated from pSport-*Tne* containing D137A and D323A mutations. The oligonucleotide used for the mutagenesis was 5'-GAA GTT CAC CAT CCG GCC GAC CCG TCG CAT TTC 3' (SEQ ID NO:93). An *Xma*III site (bold italics in the above sequence) was introduced into the oligonucleotide for easy screening of the mutants. The mutation was confirmed by DNA sequencing. The clone was named pTne129 (D137A, D323A, K726R).

Example 33: Determination of the Activity of Non-templated One Base Addition for *Tne* DNA Polymerase and its Mutant D137A, D323A, K726R, by Primer Extension Assay

The mutant *Tne* DNA polymerase (*Tne* D137A, D323A, K726R) prepared in Example 32 was purified as described in Example 20. The assay for non-templated one base addition was conducted as described in Example 23. The results were as follows:

<i>Tne</i> DNA Polymerase	% of Product With N+1
D137A, D323A	78.4
D137A, D323A, R722H	1.7
D137A, D323A, K726R	0.9

These results demonstrate that mutation of the lysine residue at position 726 of *Tne*, particularly to arginine, substantially reduces the activity of the polymerase in adding non-templated bases.

Having now fully described the present invention in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious

-93-

to one of ordinary skill in the art that the same can be performed by modifying or changing the invention within a wide and equivalent range of conditions, formulations and other parameters without affecting the scope of the invention or any specific embodiment thereof, and that such modifications or changes are intended to be encompassed within the scope of the appended claims.

All publications, patents and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains, and are herein incorporated by reference to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- 5 (A) NAME: Life Technologies, Inc.
 (B) STREET: 9800 Medical Center Drive
 (C) CITY: Rockville
 (D) STATE: Maryland
 (E) COUNTRY: USA
 (F) POSTAL CODE (ZIP): 20850

- 10 (ii) TITLE OF INVENTION: Polymerases for Analyzing or Typing Polymorphic
 Nucleic Acid Fragments and Uses Thereof

(iii) NUMBER OF SEQUENCES: 93

(iv) COMPUTER READABLE FORM:

- 15 (A) MEDIUM TYPE: Floppy disk
 (B) COMPUTER: IBM PC compatible
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(v) CURRENT APPLICATION DATA:

- 20 (A) APPLICATION NUMBER: (To be assigned)
 (B) FILING DATE: (Herewith)

(vi) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: US (To be assigned)
 (B) FILING DATE: 06-JAN-1998

25 (vi) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: US 60/037,393
 (B) FILING DATE: 07-FEB-1997

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- 30 (A) LENGTH: 2682 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATGGCGAGAC TATTTCTCTT TGATGGCACA GCCCTGGCCT ACAGGGCATA TTACGCCCTC 60
 GACAGATCCC TTTCCACATC CACAGGAATT CCAACGAACG CCGTCTATGG CGTTGCCAGG 120

	ATGCTCGTTA AATTCATTAA GGAACACATT ATACCCGAAA AGGACTACGC GGCTGTGGCC	180
	TTCGACAAGA AGGCAGCGAC GTTCAGACAC AAAC TGCTCG TAAGCGACAA GGCGCAAAGG	240
	CCAAAGACTC CGGCTCTTCT AGTTCAGCAG CTACCTTACA TCAAGCGGCT GATAGAAGCT	300
	CTTGGTTTCA AAGTGCTGGA GCTGGAGGGA TACGAAGCAG ACGATATCAT CGCCACGCTT	360
5	GCAGTCAGGG CTGCACGTTT TTTGATGAGA TTTTCATTAA TAACCGGTGA CAAGGATATG	420
	CTTCAACTTG TAAACGAGAA GATAAAGGTC TGGAGAATCG TCAAGGGGAT ATCGGATCTT	480
	GAGCTTTACG ATTCGAAAAA GGTGAAAGAA AGATACGGTG TGAACCACA TCAGATACCG	540
	GATCTTCTAG CACTGACGGG AGACGACATA GACAACATTC CCGGTGTAAC GGAATAGGT	600
	GAAAAGACCG CTGTACAGCT TCTCGGCAAG TATAGAAATC TTGAATACAT TCTGGAGCAT	660
10	GCCCGTGAAC TCCCCAGAG AGTGAGAAAG GCTCTCTTGA GAGACAGGGA AGTTGCCATC	720
	CTCAGTAAAA AACTTGCAAC TCTGGTGACG AACGCACCTG TTGAAGTGGA CTGGGAAGAG	780
	ATGAAATACA GAGGATACGA CAAGAGAAAA CTACTTCCGA TATTGAAAGA ACTGGAGTTT	840
	GCTTCCATCA TGAAGGAACT TCAACTGTAC GAAGAAGCAG AACCCACCGG ATACGAAATC	900
	GTGAAGGATC ATAAGACCTT CGAAGATCTC ATCGAAAAGC TGAAGGAGGT TCCATCTTTT	960
15	GCCCTGGACC TTGAAACGTC CTCCCTTGAC CCGTTCAACT GTGAGATAGT CGGCATCTCC	1020
	GTGTCGTTCA AACCGAAAAC AGCTTATTAC ATTCCACTTC ATCACAGAAA CGCCCAGAAT	1080
	CTTGATGAAA CACTGGTGCT GTCGAAGTTG AAAGAGATCC TCGAAGACCC GTCTTCGAAG	1140
	ATTGTGGGTC AGAACCTGAA GTACGACTAC AAGGTTCTTA TGGTAAAGGG TATATCGCCA	1200
	GTTTATCCGC ATTTTGACAC GATGATAGCT GCATATTTGC TGGAGCCAAA CGAGAAAAAA	1260
20	TTCAATCTCG AAGATCTGTC TTTGAAATTT CTCGGATACA AAATGACGTC TTATCAGGAA	1320
	CTGATGTCGT TTTCTCACC ACTTTTTGGT TTCAGCTTTG CGGATGTTCC GGTAGACAAG	1380
	GCTGCGAACT ACTCCTGCGA GGATGCAGAC ATCACTTATA GGCTCTACAA GATACTCAGC	1440
	ATGAAGCTCC ATGAAGCGGA ACTTGAGAAC GTCTTCTACA GGATAGAGAT GCCGTTGGTG	1500
	AACGTTCTTG CACGCATGGA ATTGAACGGG GTGTATGTGG ACACAGAATT CCTGAAAAAG	1560
25	CTCTCGGAGG AGTACGGCAA AAAGCTCGAG GAACTGGCCG AAAAAATCTA CCAGATAGCA	1620
	GGTGAGCCCT TCAACATCAA TTCTCCAAAA CAGGTTTCAA AGATCCTTTT TGAGAAGCTG	1680
	GGAATAAAAC CCCGTGGAAA AACGACAAAA ACAGGAGAGT ACTCTACCAG GATAGAGGTG	1740
	TTGGAAGAGA TAGCGAATGA GCACGAGATA GTACCCCTCA TTCTCGAGTA CAGAAAGATC	1800

WO 98/35060

PCT/US98/02791

96

CAGAAACTGA AATCGACCTA CATAGACACC CTTCCGAAAC TTGTGAACCC GAAAACCGGA 1860
 AGAATTCATG CATCTTTCCA CCAGACGGGT ACCGCCACTG GCAGGTTGAG TAGCAGTGAT 1920
 CCAAATCTTC AGAATCTTCC GACAAAGAGC GAAGAGGGAA AAGAAATTAG AAAAGCGATT 1980
 GTGCCCCAGG ATCCAGACTG GTGGATCGTC AGTGCGGATT ATTCCCAAAT AGAACTCAGA 2040
 5 ATCCTCGCTC ATCTCAGTGG TGATGAGAAC CTTGTGAAGG CCTTCGAGGA GGGCATCGAT 2100
 GTGCACACCT TGACTGCCTC CAGGATCTAC AACGTAAAGC CAGAAGAAGT GAACGAAGAA 2160
 ATGCGACGGG TTGGAAAGAT GGTGAACTTC TCTATAATAT ACGGTGTCAC ACCGTACGGT 2220
 CTTTCTGTGA GACTTGGAAT ACCGGTTAAA GAAGCAGAAA AGATGATTAT CAGCTATTTTC 2280
 ACACTGTATC CAAAGGTGCG AAGCTACATC CAGCAGGTTG TTGCAGAGGC AAAAGAGAAG 2340
 10 GGCTACGTCA GGACTCTCTT TGGAAGAAAA AGAGATATTC CCCAGCTCAT GGCAAGGGAC 2400
 AAGAACACCC AGTCCGAAGG CGAAAGAATC GCAATAAACA CCCCATTCA GGAACGGCG 2460
 GCAGATATAA TAAAATTGGC TATGATAGAT ATAGACGAGG AGCTGAGAAA AAGAAACATG 2520
 AAATCCAGAA TGATCATTCA GGTTCATGAC GAACTGGTCT TCGAGGTTCC CGATGAGGAA 2580
 AAAGAAGAAC TAGTTGATCT GGTGAAGAAC AAAATGACAA ATGTGGTGAA ACTCTCTGTG 2640
 15 CCTCTTGAGG TTGACATAAG CATCGGAAAA AGCTGGTCTT GA 2682

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 893 amino acids
 (B) TYPE: amino acid
 20 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

25 Met Ala Arg Leu Phe Leu Phe Asp Gly Thr Ala Leu Ala Tyr Arg Ala
 1 5 10 15
 Tyr Tyr Ala Leu Asp Arg Ser Leu Ser Thr Ser Thr Gly Ile Pro Thr
 20 25 30
 Asn Ala Val Tyr Gly Val Ala Arg Met Leu Val Lys Phe Ile Lys Glu
 35 40 45

97

His Ile Ile Pro Glu Lys Asp Tyr Ala Ala Val Ala Phe Asp Lys Lys
 50 55 60

Ala Ala Thr Phe Arg His Lys Leu Leu Val Ser Asp Lys Ala Gln Arg
 65 70 75 80

5 Pro Lys Thr Pro Ala Leu Leu Val Gln Gln Leu Pro Tyr Ile Lys Arg
 85 90 95

Leu Ile Glu Ala Leu Gly Phe Lys Val Leu Glu Leu Glu Gly Tyr Glu
 100 105 110

10 Ala Asp Asp Ile Ile Ala Thr Leu Ala Val Arg Ala Ala Arg Phe Leu
 115 120 125

Met Arg Phe Ser Leu Ile Thr Gly Asp Lys Asp Met Leu Gln Leu Val
 130 135 140

Asn Glu Lys Ile Lys Val Trp Arg Ile Val Lys Gly Ile Ser Asp Leu
 145 150 155 160

15 Glu Leu Tyr Asp Ser Lys Lys Val Lys Glu Arg Tyr Gly Val Glu Pro
 165 170 175

His Gln Ile Pro Asp Leu Leu Ala Leu Thr Gly Asp Asp Ile Asp Asn
 180 185 190

20 Ile Pro Gly Val Thr Gly Ile Gly Glu Lys Thr Ala Val Gln Leu Leu
 195 200 205

Gly Lys Tyr Arg Asn Leu Glu Tyr Ile Leu Glu His Ala Arg Glu Leu
 210 215 220

Pro Gln Arg Val Arg Lys Ala Leu Leu Arg Asp Arg Glu Val Ala Ile
 225 230 235 240

25 Leu Ser Lys Lys Leu Ala Thr Leu Val Thr Asn Ala Pro Val Glu Val
 245 250 255

Asp Trp Glu Glu Met Lys Tyr Arg Gly Tyr Asp Lys Arg Lys Leu Leu
 260 265 270

30 Pro Ile Leu Lys Glu Leu Glu Phe Ala Ser Ile Met Lys Glu Leu Gln
 275 280 285

Leu Tyr Glu Glu Ala Glu Pro Thr Gly Tyr Glu Ile Val Lys Asp His
 290 295 300

Lys Thr Phe Glu Asp Leu Ile Glu Lys Leu Lys Glu Val Pro Ser Phe
 305 310 315 320

35 Ala Leu Asp Leu Glu Thr Ser Ser Leu Asp Pro Phe Asn Cys Glu Ile
 325 330 335

WO 98/35060

PCT/US98/02791

98

	Val	Gly	Ile	Ser	Val	Ser	Phe	Lys	Pro	Lys	Thr	Ala	Tyr	Tyr	Ile	Pro
				340					345						350	
	Leu	His	His	Arg	Asn	Ala	Gln	Asn	Leu	Asp	Glu	Thr	Leu	Val	Leu	Ser
			355					360					365			
5	Lys	Leu	Lys	Glu	Ile	Leu	Glu	Asp	Pro	Ser	Ser	Lys	Ile	Val	Gly	Gln
		370					375					380				
	Asn	Leu	Lys	Tyr	Asp	Tyr	Lys	Val	Leu	Met	Val	Lys	Gly	Ile	Ser	Pro
	385					390					395					400
10	Val	Tyr	Pro	His	Phe	Asp	Thr	Met	Ile	Ala	Ala	Tyr	Leu	Leu	Glu	Pro
				405						410					415	
	Asn	Glu	Lys	Lys	Phe	Asn	Leu	Glu	Asp	Leu	Ser	Leu	Lys	Phe	Leu	Gly
				420					425					430		
	Tyr	Lys	Met	Thr	Ser	Tyr	Gln	Glu	Leu	Met	Ser	Phe	Ser	Ser	Pro	Leu
			435					440					445			
15	Phe	Gly	Phe	Ser	Phe	Ala	Asp	Val	Pro	Val	Asp	Lys	Ala	Ala	Asn	Tyr
	450						455					460				
	Ser	Cys	Glu	Asp	Ala	Asp	Ile	Thr	Tyr	Arg	Leu	Tyr	Lys	Ile	Leu	Ser
	465					470					475					480
20	Met	Lys	Leu	His	Glu	Ala	Glu	Leu	Glu	Asn	Val	Phe	Tyr	Arg	Ile	Glu
				485						490					495	
	Met	Pro	Leu	Val	Asn	Val	Leu	Ala	Arg	Met	Glu	Leu	Asn	Gly	Val	Tyr
				500					505					510		
	Val	Asp	Thr	Glu	Phe	Leu	Lys	Lys	Leu	Ser	Glu	Glu	Tyr	Gly	Lys	Lys
			515					520					525			
25	Leu	Glu	Glu	Leu	Ala	Glu	Lys	Ile	Tyr	Gln	Ile	Ala	Gly	Glu	Pro	Phe
		530					535					540				
	Asn	Ile	Asn	Ser	Pro	Lys	Gln	Val	Ser	Lys	Ile	Leu	Phe	Glu	Lys	Leu
	545					550					555					560
30	Gly	Ile	Lys	Pro	Arg	Gly	Lys	Thr	Thr	Lys	Thr	Gly	Glu	Tyr	Ser	Thr
					565						570				575	
	Arg	Ile	Glu	Val	Leu	Glu	Glu	Ile	Ala	Asn	Glu	His	Glu	Ile	Val	Pro
				580					585					590		
	Leu	Ile	Leu	Glu	Tyr	Arg	Lys	Ile	Gln	Lys	Leu	Lys	Ser	Thr	Tyr	Ile
			595					600					605			
35	Asp	Thr	Leu	Pro	Lys	Leu	Val	Asn	Pro	Lys	Thr	Gly	Arg	Ile	His	Ala
	610						615					620				
	Ser	Phe	His	Gln	Thr	Gly	Thr	Ala	Thr	Gly	Arg	Leu	Ser	Ser	Ser	Asp

	625				630					635					640	
	Pro	Asn	Leu	Gln	Asn	Leu	Pro	Thr	Lys	Ser	Glu	Glu	Gly	Lys	Glu	Ile
					645					650					655	
5	Arg	Lys	Ala	Ile	Val	Pro	Gln	Asp	Pro	Asp	Trp	Trp	Ile	Val	Ser	Ala
				660					665						670	
	Asp	Tyr	Ser	Gln	Ile	Glu	Leu	Arg	Ile	Leu	Ala	His	Leu	Ser	Gly	Asp
			675					680					685			
	Glu	Asn	Leu	Val	Lys	Ala	Phe	Glu	Glu	Gly	Ile	Asp	Val	His	Thr	Leu
		690					695					700				
10	Thr	Ala	Ser	Arg	Ile	Tyr	Asn	Val	Lys	Pro	Glu	Glu	Val	Asn	Glu	Glu
	705					710					715					720
	Met	Arg	Arg	Val	Gly	Lys	Met	Val	Asn	Phe	Ser	Ile	Ile	Tyr	Gly	Val
					725					730					735	
15	Thr	Pro	Tyr	Gly	Leu	Ser	Val	Arg	Leu	Gly	Ile	Pro	Val	Lys	Glu	Ala
				740					745					750		
	Glu	Lys	Met	Ile	Ile	Ser	Tyr	Phe	Thr	Leu	Tyr	Pro	Lys	Val	Arg	Ser
			755					760					765			
	Tyr	Ile	Gln	Gln	Val	Val	Ala	Glu	Ala	Lys	Glu	Lys	Gly	Tyr	Val	Arg
		770					775						780			
20	Thr	Leu	Phe	Gly	Arg	Lys	Arg	Asp	Ile	Pro	Gln	Leu	Met	Ala	Arg	Asp
	785					790					795					800
	Lys	Asn	Thr	Gln	Ser	Glu	Gly	Glu	Arg	Ile	Ala	Ile	Asn	Thr	Pro	Ile
					805					810					815	
25	Gln	Gly	Thr	Ala	Ala	Asp	Ile	Ile	Lys	Leu	Ala	Met	Ile	Asp	Ile	Asp
				820					825					830		
	Glu	Glu	Leu	Arg	Lys	Arg	Asn	Met	Lys	Ser	Arg	Met	Ile	Ile	Gln	Val
			835					840					845			
	His	Asp	Glu	Leu	Val	Phe	Glu	Val	Pro	Asp	Glu	Glu	Lys	Glu	Glu	Leu
		850					855					860				
30	Val	Asp	Leu	Val	Lys	Asn	Lys	Met	Thr	Asn	Val	Val	Lys	Leu	Ser	Val
	865					870					875					880
	Pro	Leu	Glu	Val	Asp	Ile	Ser	Ile	Gly	Lys	Ser	Trp	Ser			
					885						890					

(2) INFORMATION FOR SEQ ID NO:3:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 677 amino acids

(B) TYPE: amino acid

100

(C) STRANDEDNESS: not relevant

(D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

5	Met	Ser	Leu	His	Ala	Arg	Glu	Leu	Pro	Gln	Arg	Val	Arg	Lys	Ala	Leu	1	5	10	15
	Leu	Arg	Asp	Arg	Glu	Val	Ala	Ile	Leu	Ser	Lys	Lys	Leu	Ala	Thr	Leu	20	25	30	
10	Val	Thr	Asn	Ala	Pro	Val	Glu	Val	Asp	Trp	Glu	Glu	Met	Lys	Tyr	Arg	35	40	45	
	Gly	Tyr	Asp	Lys	Arg	Lys	Leu	Leu	Pro	Ile	Leu	Lys	Glu	Leu	Glu	Phe	50	55	60	
	Ala	Ser	Ile	Met	Lys	Glu	Leu	Gln	Leu	Tyr	Glu	Glu	Ala	Glu	Pro	Thr	65	70	75	80
15	Gly	Tyr	Glu	Ile	Val	Lys	Asp	His	Lys	Thr	Phe	Glu	Asp	Leu	Ile	Glu	85	90	95	
	Lys	Leu	Lys	Glu	Val	Pro	Ser	Phe	Ala	Leu	Ala	Leu	Glu	Thr	Ser	Ser	100	105	110	
20	Leu	Asp	Pro	Phe	Asn	Cys	Glu	Ile	Val	Gly	Ile	Ser	Val	Ser	Phe	Lys	115	120	125	
	Pro	Lys	Thr	Ala	Tyr	Tyr	Ile	Pro	Leu	His	His	Arg	Asn	Ala	Gln	Asn	130	135	140	
	Leu	Asp	Glu	Thr	Leu	Val	Leu	Ser	Lys	Leu	Lys	Glu	Ile	Leu	Glu	Asp	145	150	155	160
25	Pro	Ser	Ser	Lys	Ile	Val	Gly	Gln	Asn	Leu	Lys	Tyr	Asp	Tyr	Lys	Val	165	170	175	
	Leu	Met	Val	Lys	Gly	Ile	Ser	Pro	Val	Tyr	Pro	His	Phe	Asp	Thr	Met	180	185	190	
30	Ile	Ala	Ala	Tyr	Leu	Leu	Glu	Pro	Asn	Glu	Lys	Lys	Phe	Asn	Leu	Glu	195	200	205	
	Asp	Leu	Ser	Leu	Lys	Phe	Leu	Gly	Tyr	Lys	Met	Thr	Ser	Tyr	Gln	Glu	210	215	220	
	Leu	Met	Ser	Phe	Ser	Ser	Pro	Leu	Phe	Gly	Phe	Ser	Phe	Ala	Asp	Val	225	230	235	240

101

Pro Val Asp Lys Ala Ala Asn Tyr Ser Cys Glu Asp Ala Asp Ile Thr
 245 250 255
 Tyr Arg Leu Tyr Lys Ile Leu Ser Met Lys Leu His Glu Ala Glu Leu
 260 265 270
 5 Glu Asn Val Phe Tyr Arg Ile Glu Met Pro Leu Val Asn Val Leu Ala
 275 280 285
 Arg Met Glu Leu Asn Gly Val Tyr Val Asp Thr Glu Phe Leu Lys Lys
 290 295 300
 10 Leu Ser Glu Glu Tyr Gly Lys Lys Leu Glu Glu Leu Ala Glu Lys Ile
 305 310 315 320
 Tyr Gln Ile Ala Gly Glu Pro Phe Asn Ile Asn Ser Pro Lys Gln Val
 325 330 335
 Ser Lys Ile Leu Phe Glu Lys Leu Gly Ile Lys Pro Arg Gly Lys Thr
 340 345 350
 15 Thr Lys Thr Gly Glu Tyr Ser Thr Arg Ile Glu Val Leu Glu Glu Ile
 355 360 365
 Ala Asn Glu His Glu Ile Val Pro Leu Ile Leu Glu Tyr Arg Lys Ile
 370 375 380
 20 Gln Lys Leu Lys Ser Thr Tyr Ile Asp Thr Leu Pro Lys Leu Val Asn
 385 390 395 400
 Pro Lys Thr Gly Arg Ile His Ala Ser Phe His Gln Thr Gly Thr Ala
 405 410 415
 Thr Gly Arg Leu Ser Ser Ser Asp Pro Asn Leu Gln Asn Leu Pro Thr
 420 425 430
 25 Lys Ser Glu Glu Gly Lys Glu Ile Arg Lys Ala Ile Val Pro Gln Asp
 435 440 445
 Pro Asp Trp Trp Ile Val Ser Ala Asp Tyr Ser Gln Ile Glu Leu Arg
 450 455 460
 30 Ile Leu Ala His Leu Ser Gly Asp Glu Asn Leu Val Lys Ala Phe Glu
 465 470 475 480
 Glu Gly Ile Asp Val His Thr Leu Thr Ala Ser Arg Ile Tyr Asn Val
 485 490 495
 Lys Pro Glu Glu Val Asn Glu Glu Met Arg Arg Val Gly Lys Met Val
 500 505 510
 35 Asn Phe Ser Ile Ile Tyr Gly Val Thr Pro Tyr Gly Leu Ser Val Arg
 515 520 525
 Leu Gly Ile Pro Val Lys Glu Ala Glu Lys Met Ile Ile Ser Tyr Phe

WO 98/35060

PCT/US98/02791

102

	530		535			540										
	Thr	Leu	Tyr	Pro	Lys	Val	Arg	Ser	Tyr	Ile	Gln	Gln	Val	Val	Ala	Glu
	545					550					555					560
5	Ala	Lys	Glu	Lys	Gly	Tyr	Val	Arg	Thr	Leu	Phe	Gly	Arg	Lys	Arg	Asp
					565					570					575	
	Ile	Pro	Gln	Leu	Met	Ala	Arg	Asp	Lys	Asn	Thr	Gln	Ser	Glu	Gly	Glu
				580					585					590		
	Arg	Ile	Ala	Ile	Asn	Thr	Pro	Ile	Gln	Gly	Thr	Ala	Ala	Asp	Ile	Ile
			595					600					605			
10	Lys	Leu	Ala	Met	Ile	Asp	Ile	Asp	Glu	Glu	Leu	Arg	Lys	Arg	Asn	Met
	610						615					620				
	Lys	Ser	Arg	Met	Ile	Ile	Gln	Val	His	Asp	Glu	Leu	Val	Phe	Glu	Val
	625					630					635					640
15	Pro	Asp	Glu	Glu	Lys	Glu	Glu	Leu	Val	Asp	Leu	Val	Lys	Asn	Lys	Met
					645					650					655	
	Thr	Asn	Val	Val	Lys	Leu	Ser	Val	Pro	Leu	Glu	Val	Asp	Ile	Ser	Ile
				660					665					670		
	Gly	Lys	Ser	Trp	Ser											
				675												

20 (2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 610 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: not relevant

25

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

	Met	Lys	Glu	Leu	Gln	Leu	Tyr	Glu	Glu	Ala	Glu	Pro	Thr	Gly	Tyr	Glu
	1				5					10					15	
30	Ile	Val	Lys	Asp	His	Lys	Thr	Phe	Glu	Asp	Leu	Ile	Glu	Lys	Leu	Lys
				20					25					30		
	Glu	Val	Pro	Ser	Phe	Ala	Leu	Ala	Leu	Glu	Thr	Ser	Ser	Leu	Asp	Pro
			35					40					45			
35	Phe	Asn	Cys	Glu	Ile	Val	Gly	Ile	Ser	Val	Ser	Phe	Lys	Pro	Lys	Thr
	50						55					60				

WO 98/35060

PCT/US98/02791

103

Ala Tyr Tyr Ile Pro Leu His His Arg Asn Ala Gln Asn Leu Asp Glu
 65 70 75 80
 Thr Leu Val Leu Ser Lys Leu Lys Glu Ile Leu Glu Asp Pro Ser Ser
 85 90 95
 5 Lys Ile Val Gly Gln Asn Leu Lys Tyr Asp Tyr Lys Val Leu Met Val
 100 105 110
 Lys Gly Ile Ser Pro Val Tyr Pro His Phe Asp Thr Met Ile Ala Ala
 115 120 125
 10 Tyr Leu Leu Glu Pro Asn Glu Lys Lys Phe Asn Leu Glu Asp Leu Ser
 130 135 140
 Leu Lys Phe Leu Gly Tyr Lys Met Thr Ser Tyr Gln Glu Leu Met Ser
 145 150 155 160
 Phe Ser Ser Pro Leu Phe Gly Phe Ser Phe Ala Asp Val Pro Val Asp
 165 170 175
 15 Lys Ala Ala Asn Tyr Ser Cys Glu Asp Ala Asp Ile Thr Tyr Arg Leu
 180 185 190
 Tyr Lys Ile Leu Ser Met Lys Leu His Glu Ala Glu Leu Glu Asn Val
 195 200 205
 20 Phe Tyr Arg Ile Glu Met Pro Leu Val Asn Val Leu Ala Arg Met Glu
 210 215 220
 Leu Asn Gly Val Tyr Val Asp Thr Glu Phe Leu Lys Lys Leu Ser Glu
 225 230 235 240
 Glu Tyr Gly Lys Lys Leu Glu Glu Leu Ala Glu Lys Ile Tyr Gln Ile
 245 250 255
 25 Ala Gly Glu Pro Phe Asn Ile Asn Ser Pro Lys Gln Val Ser Lys Ile
 260 265 270
 Leu Phe Glu Lys Leu Gly Ile Lys Pro Arg Gly Lys Thr Thr Lys Thr
 275 280 285
 30 Gly Glu Tyr Ser Thr Arg Ile Glu Val Leu Glu Glu Ile Ala Asn Glu
 290 295 300
 His Glu Ile Val Pro Leu Ile Leu Glu Tyr Arg Lys Ile Gln Lys Leu
 305 310 315 320
 Lys Ser Thr Tyr Ile Asp Thr Leu Pro Lys Leu Val Asn Pro Lys Thr
 325 330 335
 35 Gly Arg Ile His Ala Ser Phe His Gln Thr Gly Thr Ala Thr Gly Arg
 340 345 350
 Leu Ser Ser Ser Asp Pro Asn Leu Gln Asn Leu Pro Thr Lys Ser Glu

104

			355				360				365							
			Glu	Gly	Lys	Glu	Ile	Arg	Lys	Ala	Ile	Val	Pro	Gln	Asp	Pro	Asp	Trp
			370						375					380				
5			Trp	Ile	Val	Ser	Ala	Asp	Tyr	Ser	Gln	Ile	Glu	Leu	Arg	Ile	Leu	Ala
			385					390					395					400
			His	Leu	Ser	Gly	Asp	Glu	Asn	Leu	Val	Lys	Ala	Phe	Glu	Glu	Gly	Ile
						405						410					415	
			Asp	Val	His	Thr	Leu	Thr	Ala	Ser	Arg	Ile	Tyr	Asn	Val	Lys	Pro	Glu
						420					425					430		
10			Glu	Val	Asn	Glu	Glu	Met	Arg	Arg	Val	Gly	Lys	Met	Val	Asn	Phe	Ser
					435					440					445			
			Ile	Ile	Tyr	Gly	Val	Thr	Pro	Tyr	Gly	Leu	Ser	Val	Arg	Leu	Gly	Ile
			450						455					460				
15			Pro	Val	Lys	Glu	Ala	Glu	Lys	Met	Ile	Ile	Ser	Tyr	Phe	Thr	Leu	Tyr
			465					470					475					480
			Pro	Lys	Val	Arg	Ser	Tyr	Ile	Gln	Gln	Val	Val	Ala	Glu	Ala	Lys	Glu
						485						490					495	
20			Lys	Gly	Tyr	Val	Arg	Thr	Leu	Phe	Gly	Arg	Lys	Arg	Asp	Ile	Pro	Gln
						500					505					510		
			Leu	Met	Ala	Arg	Asp	Lys	Asn	Thr	Gln	Ser	Glu	Gly	Glu	Arg	Ile	Ala
					515					520					525			
			Ile	Asn	Thr	Pro	Ile	Gln	Gly	Thr	Ala	Ala	Asp	Ile	Ile	Lys	Leu	Ala
			530						535					540				
25			Met	Ile	Asp	Ile	Asp	Glu	Glu	Leu	Arg	Lys	Arg	Asn	Met	Lys	Ser	Arg
			545				550						555					560
			Met	Ile	Ile	Gln	Val	His	Asp	Glu	Leu	Val	Phe	Glu	Val	Pro	Asp	Glu
						565						570					575	
30			Glu	Lys	Glu	Glu	Leu	Val	Asp	Leu	Val	Lys	Asn	Lys	Met	Thr	Asn	Val
					580						585					590		
			Val	Lys	Leu	Ser	Val	Pro	Leu	Glu	Val	Asp	Ile	Ser	Ile	Gly	Lys	Ser
					595					600					605			
			Trp	Ser														
			610															

35 (2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 708 amino acids

105

- (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met Asn Ser Ser Ser Val Pro Ile Pro Gly Val Thr Gly Ile Gly Glu
 1 5 10 15

Lys Thr Ala Val Gln Leu Leu Gly Lys Tyr Arg Asn Leu Glu Tyr Ile
 20 25 30

10 Leu Glu His Ala Arg Glu Leu Pro Gln Arg Val Arg Lys Ala Leu Leu
 35 40 45

Arg Asp Arg Glu Val Ala Ile Leu Ser Lys Lys Leu Ala Thr Leu Val
 50 55 60

15 Thr Asn Ala Pro Val Glu Val Asp Trp Glu Glu Met Lys Tyr Arg Gly
 65 70 75 80

Tyr Asp Lys Arg Lys Leu Leu Pro Ile Leu Lys Glu Leu Glu Phe Ala
 85 90 95

Ser Ile Met Lys Glu Leu Gln Leu Tyr Glu Glu Ala Glu Pro Thr Gly
 100 105 110

20 Tyr Glu Ile Val Lys Asp His Lys Thr Phe Glu Asp Leu Ile Glu Lys
 115 120 125

Leu Lys Glu Val Pro Ser Phe Ala Leu Ala Leu Glu Thr Ser Ser Leu
 130 135 140

25 Asp Pro Phe Asn Cys Glu Ile Val Gly Ile Ser Val Ser Phe Lys Pro
 145 150 155 160

Lys Thr Ala Tyr Tyr Ile Pro Leu His His Arg Asn Ala Gln Asn Leu
 165 170 175

Asp Glu Thr Leu Val Leu Ser Lys Leu Lys Glu Ile Leu Glu Asp Pro
 180 185 190

30 Ser Ser Lys Ile Val Gly Gln Asn Leu Lys Tyr Asp Tyr Lys Val Leu
 195 200 205

Met Val Lys Gly Ile Ser Pro Val Tyr Pro His Phe Asp Thr Met Ile
 210 215 220

Ala Ala Tyr Leu Leu Glu Pro Asn Glu Lys Lys Phe Asn Leu Glu Asp

WO 98/35060

PCT/US98/02791

106

	225				230					235				240		
	Leu	Ser	Leu	Lys	Phe	Leu	Gly	Tyr	Lys	Met	Thr	Ser	Tyr	Gln	Glu	Leu
					245					250					255	
5	Met	Ser	Phe	Ser	Ser	Pro	Leu	Phe	Gly	Phe	Ser	Phe	Ala	Asp	Val	Pro
				260					265					270		
	Val	Asp	Lys	Ala	Ala	Asn	Tyr	Ser	Cys	Glu	Asp	Ala	Asp	Ile	Thr	Tyr
			275					280					285			
	Arg	Leu	Tyr	Lys	Ile	Leu	Ser	Met	Lys	Leu	His	Glu	Ala	Glu	Leu	Glu
		290					295					300				
10	Asn	Val	Phe	Tyr	Arg	Ile	Glu	Met	Pro	Leu	Val	Asn	Val	Leu	Ala	Arg
	305					310					315					320
	Met	Glu	Leu	Asn	Gly	Val	Tyr	Val	Asp	Thr	Glu	Phe	Leu	Lys	Lys	Leu
				325						330					335	
15	Ser	Glu	Glu	Tyr	Gly	Lys	Lys	Leu	Glu	Glu	Leu	Ala	Glu	Lys	Ile	Tyr
				340					345					350		
	Gln	Ile	Ala	Gly	Glu	Pro	Phe	Asn	Ile	Asn	Ser	Pro	Lys	Gln	Val	Ser
			355					360					365			
	Lys	Ile	Leu	Phe	Glu	Lys	Leu	Gly	Ile	Lys	Pro	Arg	Gly	Lys	Thr	Thr
		370					375					380				
20	Lys	Thr	Gly	Glu	Tyr	Ser	Thr	Arg	Ile	Glu	Val	Leu	Glu	Glu	Ile	Ala
	385					390					395					400
	Asn	Glu	His	Glu	Ile	Val	Pro	Leu	Ile	Leu	Glu	Tyr	Arg	Lys	Ile	Gln
				405						410					415	
25	Lys	Leu	Lys	Ser	Thr	Tyr	Ile	Asp	Thr	Leu	Pro	Lys	Leu	Val	Asn	Pro
			420					425						430		
	Lys	Thr	Gly	Arg	Ile	His	Ala	Ser	Phe	His	Gln	Thr	Gly	Thr	Ala	Thr
			435					440					445			
	Gly	Arg	Leu	Ser	Ser	Ser	Asp	Pro	Asn	Leu	Gln	Asn	Leu	Pro	Thr	Lys
		450					455					460				
30	Ser	Glu	Glu	Gly	Lys	Glu	Ile	Arg	Lys	Ala	Ile	Val	Pro	Gln	Asp	Pro
	465					470					475					480
	Asp	Trp	Trp	Ile	Val	Ser	Ala	Asp	Tyr	Ser	Gln	Ile	Glu	Leu	Arg	Ile
				485						490					495	
35	Leu	Ala	His	Leu	Ser	Gly	Asp	Glu	Asn	Leu	Val	Lys	Ala	Phe	Glu	Glu
			500						505					510		
	Gly	Ile	Asp	Val	His	Thr	Leu	Thr	Ala	Ser	Arg	Ile	Tyr	Asn	Val	Lys

107

	515	520	525
	Pro Glu Glu Val Asn Glu Glu Met Arg Arg Val Gly Lys Met Val Asn 530 535 540		
5	Phe Ser Ile Ile Tyr Gly Val Thr Pro Tyr Gly Leu Ser Val Arg Leu 545 550 555 560		
	Gly Ile Pro Val Lys Glu Ala Glu Lys Met Ile Ile Ser Tyr Phe Thr 565 570 575		
	Leu Tyr Pro Lys Val Arg Ser Tyr Ile Gln Gln Val Val Ala Glu Ala 580 585 590		
10	Lys Glu Lys Gly Tyr Val Arg Thr Leu Phe Gly Arg Lys Arg Asp Ile 595 600 605		
	Pro Gln Leu Met Ala Arg Asp Lys Asn Thr Gln Ser Glu Gly Glu Arg 610 615 620		
15	Ile Ala Ile Asn Thr Pro Ile Gln Gly Thr Ala Ala Asp Ile Ile Lys 625 630 635 640		
	Leu Ala Met Ile Asp Ile Asp Glu Glu Leu Arg Lys Arg Asn Met Lys 645 650 655		
	Ser Arg Met Ile Ile Gln Val His Asp Glu Leu Val Phe Glu Val Pro 660 665 670		
20	Asp Glu Glu Lys Glu Glu Leu Val Asp Leu Val Lys Asn Lys Met Thr 675 680 685		
	Asn Val Val Lys Leu Ser Val Pro Leu Glu Val Asp Ile Ser Ile Gly 690 695 700		
25	Lys Ser Trp Ser 705		

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 893 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: not relevant

30

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Ala Arg Leu Phe Leu Phe Asp Gly Thr Ala Leu Ala Tyr Arg Ala

WO 98/35060

PCT/US98/02791

108

	1			5					10					15		
	Tyr	Tyr	Ala	Leu	Asp	Arg	Ser	Leu	Ser	Thr	Ser	Thr	Gly	Ile	Pro	Thr
				20					25					30		
5	Asn	Ala	Val	Tyr	Gly	Val	Ala	Arg	Met	Leu	Val	Lys	Phe	Ile	Lys	Glu
			35					40					45			
	His	Ile	Ile	Pro	Glu	Lys	Asp	Tyr	Ala	Ala	Val	Ala	Phe	Asp	Lys	Lys
		50					55					60				
	Ala	Ala	Thr	Phe	Arg	His	Lys	Leu	Leu	Val	Ser	Asp	Lys	Ala	Gln	Arg
	65					70					75					80
10	Pro	Lys	Thr	Pro	Ala	Leu	Leu	Val	Gln	Gln	Leu	Pro	Tyr	Ile	Lys	Arg
					85					90					95	
	Leu	Ile	Glu	Ala	Leu	Gly	Phe	Lys	Val	Leu	Glu	Leu	Glu	Gly	Tyr	Glu
				100					105					110		
15	Ala	Asp	Asp	Ile	Ile	Ala	Thr	Leu	Ala	Val	Arg	Ala	Ala	Arg	Phe	Leu
			115					120					125			
	Met	Arg	Phe	Ser	Leu	Ile	Thr	Gly	Ala	Lys	Asp	Met	Leu	Gln	Leu	Val
		130					135					140				
	Asn	Glu	Lys	Ile	Lys	Val	Trp	Arg	Ile	Val	Lys	Gly	Ile	Ser	Asp	Leu
	145					150					155					160
20	Glu	Leu	Tyr	Asp	Ser	Lys	Lys	Val	Lys	Glu	Arg	Tyr	Gly	Val	Glu	Pro
					165					170					175	
	His	Gln	Ile	Pro	Asp	Leu	Leu	Ala	Leu	Thr	Gly	Asp	Asp	Ile	Asp	Asn
				180					185					190		
25	Ile	Pro	Gly	Val	Thr	Gly	Ile	Gly	Glu	Lys	Thr	Ala	Val	Gln	Leu	Leu
			195					200					205			
	Gly	Lys	Tyr	Arg	Asn	Leu	Glu	Tyr	Ile	Leu	Glu	His	Ala	Arg	Glu	Leu
		210					215					220				
	Pro	Gln	Arg	Val	Arg	Lys	Ala	Leu	Leu	Arg	Asp	Arg	Glu	Val	Ala	Ile
	225					230					235					240
30	Leu	Ser	Lys	Lys	Leu	Ala	Thr	Leu	Val	Thr	Asn	Ala	Pro	Val	Glu	Val
					245					250					255	
	Asp	Trp	Glu	Glu	Met	Lys	Tyr	Arg	Gly	Tyr	Asp	Lys	Arg	Lys	Leu	Leu
			260						265					270		
	Pro	Ile	Leu	Lys	Glu	Leu	Glu	Phe	Ala	Ser	Ile	Met	Lys	Glu	Leu	Gln
35			275					280					285			
	Leu	Tyr	Glu	Glu	Ala	Glu	Pro	Thr	Gly	Tyr	Glu	Ile	Val	Lys	Asp	His

	290		295		300											
	Lys	Thr	Phe	Glu	Asp	Leu	Ile	Glu	Lys	Leu	Lys	Glu	Val	Pro	Ser	Phe
	305					310					315					320
5	Ala	Leu	Ala	Leu	Glu	Thr	Ser	Ser	Leu	Asp	Pro	Phe	Asn	Cys	Glu	Ile
					325					330					335	
	Val	Gly	Ile	Ser	Val	Ser	Phe	Lys	Pro	Lys	Thr	Ala	Tyr	Tyr	Ile	Pro
				340					345					350		
	Leu	His	His	Arg	Asn	Ala	Gln	Asn	Leu	Asp	Glu	Thr	Leu	Val	Leu	Ser
			355					360					365			
10	Lys	Leu	Lys	Glu	Ile	Leu	Glu	Asp	Pro	Ser	Ser	Lys	Ile	Val	Gly	Gln
		370					375					380				
	Asn	Leu	Lys	Tyr	Asp	Tyr	Lys	Val	Leu	Met	Val	Lys	Gly	Ile	Ser	Pro
	385					390					395					400
15	Val	Tyr	Pro	His	Phe	Asp	Thr	Met	Ile	Ala	Ala	Tyr	Leu	Leu	Glu	Pro
					405					410					415	
	Asn	Glu	Lys	Lys	Phe	Asn	Leu	Glu	Asp	Leu	Ser	Leu	Lys	Phe	Leu	Gly
				420					425					430		
	Tyr	Lys	Met	Thr	Ser	Tyr	Gln	Glu	Leu	Met	Ser	Phe	Ser	Ser	Pro	Leu
			435					440					445			
20	Phe	Gly	Phe	Ser	Phe	Ala	Asp	Val	Pro	Val	Asp	Lys	Ala	Ala	Asn	Tyr
		450					455					460				
	Ser	Cys	Glu	Asp	Ala	Asp	Ile	Thr	Tyr	Arg	Leu	Tyr	Lys	Ile	Leu	Ser
	465					470					475					480
25	Met	Lys	Leu	His	Glu	Ala	Glu	Leu	Glu	Asn	Val	Phe	Tyr	Arg	Ile	Glu
					485					490					495	
	Met	Pro	Leu	Val	Asn	Val	Leu	Ala	Arg	Met	Glu	Leu	Asn	Gly	Val	Tyr
				500					505					510		
	Val	Asp	Thr	Glu	Phe	Leu	Lys	Lys	Leu	Ser	Glu	Glu	Tyr	Gly	Lys	Lys
			515					520					525			
30	Leu	Glu	Glu	Leu	Ala	Glu	Lys	Ile	Tyr	Gln	Ile	Ala	Gly	Glu	Pro	Phe
		530					535					540				
	Asn	Ile	Asn	Ser	Pro	Lys	Gln	Val	Ser	Lys	Ile	Leu	Phe	Glu	Lys	Leu
	545					550					555					560
35	Gly	Ile	Lys	Pro	Arg	Gly	Lys	Thr	Thr	Lys	Thr	Gly	Glu	Tyr	Ser	Thr
					565					570					575	
	Arg	Ile	Glu	Val	Leu	Glu	Glu	Ile	Ala	Asn	Glu	His	Glu	Ile	Val	Pro

WO 98/35060

PCT/US98/02791

110

	580					585					590					
	Leu	Ile	Leu	Glu	Tyr	Arg	Lys	Ile	Gln	Lys	Leu	Lys	Ser	Thr	Tyr	Ile
			595					600					605			
5	Asp	Thr	Leu	Pro	Lys	Leu	Val	Asn	Pro	Lys	Thr	Gly	Arg	Ile	His	Ala
	610					615					620					
	Ser	Phe	His	Gln	Thr	Gly	Thr	Ala	Thr	Gly	Arg	Leu	Ser	Ser	Ser	Asp
	625					630					635					640
	Pro	Asn	Leu	Gln	Asn	Leu	Pro	Thr	Lys	Ser	Glu	Glu	Gly	Lys	Glu	Ile
					645					650					655	
10	Arg	Lys	Ala	Ile	Val	Pro	Gln	Asp	Pro	Asp	Trp	Trp	Ile	Val	Ser	Ala
				660					665					670		
	Asp	Tyr	Ser	Gln	Ile	Glu	Leu	Arg	Ile	Leu	Ala	His	Leu	Ser	Gly	Asp
			675					680					685			
15	Glu	Asn	Leu	Val	Lys	Ala	Phe	Glu	Glu	Gly	Ile	Asp	Val	His	Thr	Leu
	690					695						700				
	Thr	Ala	Ser	Arg	Ile	Tyr	Asn	Val	Lys	Pro	Glu	Glu	Val	Asn	Glu	Glu
	705					710					715					720
	Met	Arg	Arg	Val	Gly	Lys	Met	Val	Asn	Phe	Ser	Ile	Ile	Tyr	Gly	Val
					725					730					735	
20	Thr	Pro	Tyr	Gly	Leu	Ser	Val	Arg	Leu	Gly	Ile	Pro	Val	Lys	Glu	Ala
				740					745					750		
	Glu	Lys	Met	Ile	Ile	Ser	Tyr	Phe	Thr	Leu	Tyr	Pro	Lys	Val	Arg	Ser
			755					760					765			
25	Tyr	Ile	Gln	Gln	Val	Val	Ala	Glu	Ala	Lys	Glu	Lys	Gly	Tyr	Val	Arg
	770					775							780			
	Thr	Leu	Phe	Gly	Arg	Lys	Arg	Asp	Ile	Pro	Gln	Leu	Met	Ala	Arg	Asp
	785					790					795					800
	Lys	Asn	Thr	Gln	Ser	Glu	Gly	Glu	Arg	Ile	Ala	Ile	Asn	Thr	Pro	Ile
					805					810					815	
30	Gln	Gly	Thr	Ala	Ala	Asp	Ile	Ile	Lys	Leu	Ala	Met	Ile	Asp	Ile	Asp
				820					825					830		
	Glu	Glu	Leu	Arg	Lys	Arg	Asn	Met	Lys	Ser	Arg	Met	Ile	Ile	Gln	Val
			835					840					845			
35	His	Asp	Glu	Leu	Val	Phe	Glu	Val	Pro	Asp	Glu	Glu	Lys	Glu	Glu	Leu
	850					855					860					
	Val	Asp	Leu	Val	Lys	Asn	Lys	Met	Thr	Asn	Val	Val	Lys	Leu	Ser	Val

112

	Ile	Pro	Gly	Val	Thr	Gly	Ile	Gly	Glu	Lys	Thr	Ala	Val	Gln	Leu	Leu
			195					200					205			
	Gly	Lys	Tyr	Arg	Asn	Leu	Glu	Tyr	Ile	Leu	Glu	His	Ala	Arg	Glu	Leu
		210					215					220				
5	Pro	Gln	Arg	Val	Arg	Lys	Ala	Leu	Leu	Arg	Asp	Arg	Glu	Val	Ala	Ile
	225					230					235					240
	Leu	Ser	Lys	Lys	Leu	Ala	Thr	Leu	Val	Thr	Asn	Ala	Pro	Val	Glu	Val
					245					250					255	
10	Asp	Trp	Glu	Glu	Met	Lys	Tyr	Arg	Gly	Tyr	Asp	Lys	Arg	Lys	Leu	Leu
			260						265						270	
	Pro	Ile	Leu	Lys	Glu	Leu	Glu	Phe	Ala	Ser	Ile	Met	Lys	Glu	Leu	Gln
			275					280					285			
	Leu	Tyr	Glu	Glu	Ala	Glu	Pro	Thr	Gly	Tyr	Glu	Ile	Val	Lys	Asp	His
		290					295					300				
15	Lys	Thr	Phe	Glu	Asp	Leu	Ile	Glu	Lys	Leu	Lys	Glu	Val	Pro	Ser	Phe
	305					310					315					320
	Ala	Leu	Ala	Leu	Glu	Thr	Ser	Ser	Leu	Asp	Pro	Phe	Asn	Cys	Glu	Ile
					325					330					335	
20	Val	Gly	Ile	Ser	Val	Ser	Phe	Lys	Pro	Lys	Thr	Ala	Tyr	Tyr	Ile	Pro
				340					345					350		
	Leu	His	His	Arg	Asn	Ala	Gln	Asn	Leu	Asp	Glu	Thr	Leu	Val	Leu	Ser
			355					360					365			
	Lys	Leu	Lys	Glu	Ile	Leu	Glu	Asp	Pro	Ser	Ser	Lys	Ile	Val	Gly	Gln
		370					375					380				
25	Asn	Leu	Lys	Tyr	Asp	Tyr	Lys	Val	Leu	Met	Val	Lys	Gly	Ile	Ser	Pro
	385					390					395					400
	Val	Tyr	Pro	His	Phe	Asp	Thr	Met	Ile	Ala	Ala	Tyr	Leu	Leu	Glu	Pro
					405					410					415	
30	Asn	Glu	Lys	Lys	Phe	Asn	Leu	Glu	Asp	Leu	Ser	Leu	Lys	Phe	Leu	Gly
				420					425					430		
	Tyr	Lys	Met	Thr	Ser	Tyr	Gln	Glu	Leu	Met	Ser	Phe	Ser	Ser	Pro	Leu
			435					440					445			
	Phe	Gly	Phe	Ser	Phe	Ala	Asp	Val	Pro	Val	Asp	Lys	Ala	Ala	Asn	Tyr
		450					455					460				
35	Ser	Cys	Glu	Asp	Ala	Asp	Ile	Thr	Tyr	Arg	Leu	Tyr	Lys	Ile	Leu	Ser
	465					470					475					480
	Met	Lys	Leu	His	Glu	Ala	Glu	Leu	Glu	Asn	Val	Phe	Tyr	Arg	Ile	Glu

WO 98/35060

PCT/US98/02791

113

	485	490	495
	Met Pro Leu Val Asn Val Leu Ala Arg Met Glu Leu Asn Gly Val Tyr 500	505	510
5	Val Asp Thr Glu Phe Leu Lys Lys Leu Ser Glu Glu Tyr Gly Lys Lys 515	520	525
	Leu Glu Glu Leu Ala Glu Lys Ile Tyr Gln Ile Ala Gly Glu Pro Phe 530	535	540
	Asn Ile Asn Ser Pro Lys Gln Val Ser Lys Ile Leu Phe Glu Lys Leu 545	550	555
10	Gly Ile Lys Pro Arg Gly Lys Thr Thr Lys Thr Gly Glu Tyr Ser Thr 565	570	575
	Arg Ile Glu Val Leu Glu Glu Ile Ala Asn Glu His Glu Ile Val Pro 580	585	590
15	Leu Ile Leu Glu Tyr Arg Lys Ile Gln Lys Leu Lys Ser Thr Tyr Ile 595	600	605
	Asp Thr Leu Pro Lys Leu Val Asn Pro Lys Thr Gly Arg Ile His Ala 610	615	620
	Ser Phe His Gln Thr Gly Thr Ala Thr Gly Arg Leu Ser Ser Ser Asp 625	630	635
20	Pro Asn Leu Gln Asn Leu Pro Thr Lys Ser Glu Glu Gly Lys Glu Ile 645	650	655
	Arg Lys Ala Ile Val Pro Gln Asp Pro Asp Trp Trp Ile Val Ser Ala 660	665	670
25	Asp Tyr Ser Gln Ile Glu Leu Arg Ile Leu Ala His Leu Ser Gly Asp 675	680	685
	Glu Asn Leu Val Lys Ala Phe Glu Glu Gly Ile Asp Val His Thr Leu 690	695	700
	Thr Ala Ser Arg Ile Tyr Asn Val Lys Pro Glu Glu Val Asn Glu Glu 705	710	715
30	Met Arg Arg Val Gly Lys Met Val Asn Phe Ser Ile Ile Tyr Gly Val 725	730	735
	Thr Pro Tyr Gly Leu Ser Val Arg Leu Gly Ile Pro Val Lys Glu Ala 740	745	750
35	Glu Lys Met Ile Ile Ser Tyr Phe Thr Leu Tyr Pro Lys Val Arg Ser 755	760	765
	Tyr Ile Gln Gln Val Val Ala Glu Ala Lys Glu Lys Gly Tyr Val Arg		

114

	770		775		780														
	Thr	Leu	Phe	Gly	Arg	Lys	Arg	Asp	Ile	Pro	Gln	Leu	Met	Ala	Arg	Asp			
	785					790					795					800			
5	Lys	Asn	Thr	Gln	Ser	Glu	Gly	Glu	Arg	Ile	Ala	Ile	Asn	Thr	Pro	Ile			
					805					810					815				
	Gln	Gly	Thr	Ala	Ala	Asp	Ile	Ile	Lys	Leu	Ala	Met	Ile	Asp	Ile	Asp			
				820					825					830					
	Glu	Glu	Leu	Arg	Lys	Arg	Asn	Met	Lys	Ser	Arg	Met	Ile	Ile	Gln	Val			
			835					840					845						
10	His	Asp	Glu	Leu	Val	Phe	Glu	Val	Pro	Asp	Glu	Glu	Lys	Glu	Glu	Leu			
	850						855					860							
	Val	Asp	Leu	Val	Lys	Asn	Lys	Met	Thr	Asn	Val	Val	Lys	Leu	Ser	Val			
	865					870					875					880			
15	Pro	Leu	Glu	Val	Asp	Ile	Ser	Ile	Gly	Lys	Ser	Trp	Ser						
					885					890									

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 893 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

25	Met	Ala	Arg	Leu	Phe	Leu	Phe	Asp	Gly	Thr	Ala	Leu	Ala	Tyr	Arg	Ala			
	1				5					10					15				
	Tyr	Tyr	Ala	Leu	Asp	Arg	Ser	Leu	Ser	Thr	Ser	Thr	Gly	Ile	Pro	Thr			
				20					25					30					
	Asn	Ala	Val	Tyr	Gly	Val	Ala	Arg	Met	Leu	Val	Lys	Phe	Ile	Lys	Glu			
			35					40					45						
30	His	Ile	Ile	Pro	Glu	Lys	Asp	Tyr	Ala	Ala	Val	Ala	Phe	Asp	Lys	Lys			
	50						55					60							
	Ala	Ala	Thr	Phe	Arg	His	Lys	Leu	Leu	Val	Ser	Asp	Lys	Ala	Gln	Arg			
	65					70					75					80			
	Pro	Lys	Thr	Pro	Ala	Leu	Leu	Val	Gln	Gln	Leu	Pro	Tyr	Ile	Lys	Arg			

115

	85					90					95					
	Leu	Ile	Glu	Ala	Leu	Gly	Phe	Lys	Val	Leu	Glu	Leu	Glu	Gly	Tyr	Glu
				100					105					110		
5	Ala	Asp	Asp	Ile	Ile	Ala	Thr	Leu	Ala	Val	Arg	Ala	Ala	Arg	Phe	Leu
			115					120					125			
	Met	Arg	Phe	Ser	Leu	Ile	Thr	Gly	Asp	Lys	Asp	Met	Leu	Gln	Leu	Val
		130					135					140				
	Asn	Glu	Lys	Ile	Lys	Val	Trp	Arg	Ile	Val	Lys	Gly	Ile	Ser	Asp	Leu
	145				150						155				160	
10	Glu	Leu	Tyr	Asp	Ser	Lys	Lys	Val	Lys	Glu	Arg	Tyr	Gly	Val	Glu	Pro
					165					170					175	
	His	Gln	Ile	Pro	Asp	Leu	Leu	Ala	Leu	Thr	Gly	Asp	Asp	Ile	Asp	Asn
				180					185					190		
15	Ile	Pro	Asp	Val	Thr	Gly	Ile	Gly	Glu	Lys	Thr	Ala	Val	Gln	Leu	Leu
			195					200					205			
	Gly	Lys	Tyr	Arg	Asn	Leu	Glu	Tyr	Ile	Leu	Glu	His	Ala	Arg	Glu	Leu
		210					215					220				
	Pro	Gln	Arg	Val	Arg	Lys	Ala	Leu	Leu	Arg	Asp	Arg	Glu	Val	Ala	Ile
	225					230					235				240	
20	Leu	Ser	Lys	Lys	Leu	Ala	Thr	Leu	Val	Thr	Asn	Ala	Pro	Val	Glu	Val
					245					250					255	
	Asp	Trp	Glu	Glu	Met	Lys	Tyr	Arg	Gly	Tyr	Asp	Lys	Arg	Lys	Leu	Leu
				260					265					270		
25	Pro	Ile	Leu	Lys	Glu	Leu	Glu	Phe	Ala	Ser	Ile	Met	Lys	Glu	Leu	Gln
			275					280					285			
	Leu	Tyr	Glu	Glu	Ala	Glu	Pro	Thr	Gly	Tyr	Glu	Ile	Val	Lys	Asp	His
		290					295					300				
	Lys	Thr	Phe	Glu	Asp	Leu	Ile	Glu	Lys	Leu	Lys	Glu	Val	Pro	Ser	Phe
	305				310						315				320	
30	Ala	Leu	Ala	Leu	Glu	Thr	Ser	Ser	Leu	Asp	Pro	Phe	Asn	Cys	Glu	Ile
					325					330					335	
	Val	Gly	Ile	Ser	Val	Ser	Phe	Lys	Pro	Lys	Thr	Ala	Tyr	Tyr	Ile	Pro
				340					345					350		
35	Leu	His	His	Arg	Asn	Ala	Gln	Asn	Leu	Asp	Glu	Thr	Leu	Val	Leu	Ser
			355				360						365			
	Lys	Leu	Lys	Glu	Ile	Leu	Glu	Asp	Pro	Ser	Ser	Lys	Ile	Val	Gly	Gln

WO 98/35060

PCT/US98/02791

116

	370		375		380												
	Asn	Leu	Lys	Tyr	Asp	Tyr	Lys	Val	Leu	Met	Val	Lys	Gly	Ile	Ser	Pro	
	385					390					395					400	
5	Val	Tyr	Pro	His	Phe	Asp	Thr	Met	Ile	Ala	Ala	Tyr	Leu	Leu	Glu	Pro	
				405						410					415		
	Asn	Glu	Lys	Lys	Phe	Asn	Leu	Glu	Asp	Leu	Ser	Leu	Lys	Phe	Leu	Gly	
				420					425					430			
	Tyr	Lys	Met	Thr	Ser	Tyr	Gln	Glu	Leu	Met	Ser	Phe	Ser	Ser	Pro	Leu	
			435					440					445				
10	Phe	Gly	Phe	Ser	Phe	Ala	Asp	Val	Pro	Val	Asp	Lys	Ala	Ala	Asn	Tyr	
	450						455					460					
	Ser	Cys	Glu	Asp	Ala	Asp	Ile	Thr	Tyr	Arg	Leu	Tyr	Lys	Ile	Leu	Ser	
	465					470					475					480	
15	Met	Lys	Leu	His	Glu	Ala	Glu	Leu	Glu	Asn	Val	Phe	Tyr	Arg	Ile	Glu	
				485						490					495		
	Met	Pro	Leu	Val	Asn	Val	Leu	Ala	Arg	Met	Glu	Leu	Asn	Gly	Val	Tyr	
				500					505					510			
	Val	Asp	Thr	Glu	Phe	Leu	Lys	Lys	Leu	Ser	Glu	Glu	Tyr	Gly	Lys	Lys	
			515					520					525				
20	Leu	Glu	Glu	Leu	Ala	Glu	Lys	Ile	Tyr	Gln	Ile	Ala	Gly	Glu	Pro	Phe	
	530						535					540					
	Asn	Ile	Asn	Ser	Pro	Lys	Gln	Val	Ser	Lys	Ile	Leu	Phe	Glu	Lys	Leu	
	545					550					555					560	
25	Gly	Ile	Lys	Pro	Arg	Gly	Lys	Thr	Thr	Lys	Thr	Gly	Glu	Tyr	Ser	Thr	
				565						570					575		
	Arg	Ile	Glu	Val	Leu	Glu	Glu	Ile	Ala	Asn	Glu	His	Glu	Ile	Val	Pro	
				580					585					590			
	Leu	Ile	Leu	Glu	Tyr	Arg	Lys	Ile	Gln	Lys	Leu	Lys	Ser	Thr	Tyr	Ile	
			595					600					605				
30	Asp	Thr	Leu	Pro	Lys	Leu	Val	Asn	Pro	Lys	Thr	Gly	Arg	Ile	His	Ala	
	610						615					620					
	Ser	Phe	His	Gln	Thr	Gly	Thr	Ala	Thr	Gly	Arg	Leu	Ser	Ser	Ser	Asp	
	625					630					635					640	
35	Pro	Asn	Leu	Gln	Asn	Leu	Pro	Thr	Lys	Ser	Glu	Glu	Gly	Lys	Glu	Ile	
				645						650					655		
	Arg	Lys	Ala	Ile	Val	Pro	Gln	Asp	Pro	Asp	Trp	Trp	Ile	Val	Ser	Ala	

WO 98/35060

PCT/US98/02791

117

	660	665	670
	Asp Tyr Ser Gln Ile Glu Leu Arg Ile Leu Ala His Leu Ser Gly Asp 675	680	685
5	Glu Asn Leu Val Lys Ala Phe Glu Glu Gly Ile Asp Val His Thr Leu 690	695	700
	Thr Ala Ser Arg Ile Tyr Asn Val Lys Pro Glu Glu Val Asn Glu Glu 705	710	715
	Met Arg Arg Val Gly Lys Met Val Asn Phe Ser Ile Ile Tyr Gly Val 725	730	735
10	Thr Pro Tyr Gly Leu Ser Val Arg Leu Gly Ile Pro Val Lys Glu Ala 740	745	750
	Glu Lys Met Ile Ile Ser Tyr Phe Thr Leu Tyr Pro Lys Val Arg Ser 755	760	765
15	Tyr Ile Gln Gln Val Val Ala Glu Ala Lys Glu Lys Gly Tyr Val Arg 770	775	780
	Thr Leu Phe Gly Arg Lys Arg Asp Ile Pro Gln Leu Met Ala Arg Asp 785	790	800
	Lys Asn Thr Gln Ser Glu Gly Glu Arg Ile Ala Ile Asn Thr Pro Ile 805	810	815
20	Gln Gly Thr Ala Ala Asp Ile Ile Lys Leu Ala Met Ile Asp Ile Asp 820	825	830
	Glu Glu Leu Arg Lys Arg Asn Met Lys Ser Arg Met Ile Ile Gln Val 835	840	845
25	His Asp Glu Leu Val Phe Glu Val Pro Asp Glu Glu Lys Glu Glu Leu 850	855	860
	Val Asp Leu Val Lys Asn Lys Met Thr Asn Val Val Lys Leu Ser Val 865	870	875
	Pro Leu Glu Val Asp Ile Ser Ile Gly Lys Ser Trp Ser 885	890	

30 (2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 893 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: not relevant

(D) TOPOLOGY: not relevant

35

(ii) MOLECULE TYPE: protein

WO 98/35060

PCT/US98/02791

118

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

	Met	Ala	Arg	Leu	Phe	Leu	Phe	Asp	Gly	Thr	Ala	Leu	Ala	Tyr	Arg	Ala	
	1				5					10					15		
5	Tyr	Tyr	Ala	Leu	Asp	Arg	Ser	Leu	Ser	Thr	Ser	Thr	Gly	Ile	Pro	Thr	
				20					25					30			
	Asn	Ala	Val	Tyr	Asp	Val	Ala	Arg	Met	Leu	Val	Lys	Phe	Ile	Lys	Glu	
			35					40					45				
	His	Ile	Ile	Pro	Glu	Lys	Asp	Tyr	Ala	Ala	Val	Ala	Phe	Asp	Lys	Lys	
		50					55					60					
10	Ala	Ala	Thr	Phe	Arg	His	Lys	Leu	Leu	Val	Ser	Asp	Lys	Ala	Gln	Arg	
	65					70					75					80	
	Pro	Lys	Thr	Pro	Ala	Leu	Leu	Val	Gln	Gln	Leu	Pro	Tyr	Ile	Lys	Arg	
					85					90					95		
15	Leu	Ile	Glu	Ala	Leu	Gly	Phe	Lys	Val	Leu	Glu	Leu	Glu	Gly	Tyr	Glu	
				100					105					110			
	Ala	Asp	Asp	Ile	Ile	Ala	Thr	Leu	Ala	Val	Arg	Ala	Ala	Arg	Phe	Leu	
			115					120					125				
	Met	Arg	Phe	Ser	Leu	Ile	Thr	Gly	Asp	Lys	Asp	Met	Leu	Gln	Leu	Val	
		130					135					140					
20	Asn	Glu	Lys	Ile	Lys	Val	Trp	Arg	Ile	Val	Lys	Gly	Ile	Ser	Asp	Leu	
	145					150					155					160	
	Glu	Leu	Tyr	Asp	Ser	Lys	Lys	Val	Lys	Glu	Arg	Tyr	Gly	Val	Glu	Pro	
				165						170					175		
25	His	Gln	Ile	Pro	Asp	Leu	Leu	Ala	Leu	Thr	Gly	Asp	Asp	Ile	Asp	Asn	
				180					185					190			
	Ile	Pro	Gly	Val	Thr	Gly	Ile	Gly	Glu	Lys	Thr	Ala	Val	Gln	Leu	Leu	
			195					200					205				
	Gly	Lys	Tyr	Arg	Asn	Leu	Glu	Tyr	Ile	Leu	Glu	His	Ala	Arg	Glu	Leu	
		210					215					220					
30	Pro	Gln	Arg	Val	Arg	Lys	Ala	Leu	Leu	Arg	Asp	Arg	Glu	Val	Ala	Ile	
	225					230					235					240	
	Leu	Ser	Lys	Lys	Leu	Ala	Thr	Leu	Val	Thr	Asn	Ala	Pro	Val	Glu	Val	
					245					250					255		
35	Asp	Trp	Glu	Glu	Met	Lys	Tyr	Arg	Gly	Tyr	Asp	Lys	Arg	Lys	Leu	Leu	
			260						265					270			
	Pro	Ile	Leu	Lys	Glu	Leu	Glu	Phe	Ala	Ser	Ile	Met	Lys	Glu	Leu	Gln	
			275					280					285				

	Leu	Tyr	Glu	Glu	Ala	Glu	Pro	Thr	Gly	Tyr	Glu	Ile	Val	Lys	Asp	His	
	290						295					300					
	Lys	Thr	Phe	Glu	Asp	Leu	Ile	Glu	Lys	Leu	Lys	Glu	Val	Pro	Ser	Phe	
	305					310					315					320	
5	Ala	Leu	Ala	Leu	Glu	Thr	Ser	Ser	Leu	Asp	Pro	Phe	Asn	Cys	Glu	Ile	
					325					330					335		
	Val	Gly	Ile	Ser	Val	Ser	Phe	Lys	Pro	Lys	Thr	Ala	Tyr	Tyr	Ile	Pro	
				340					345						350		
10	Leu	His	His	Arg	Asn	Ala	Gln	Asn	Leu	Asp	Glu	Thr	Leu	Val	Leu	Ser	
			355					360						365			
	Lys	Leu	Lys	Glu	Ile	Leu	Glu	Asp	Pro	Ser	Ser	Lys	Ile	Val	Gly	Gln	
	370						375					380					
	Asn	Leu	Lys	Tyr	Asp	Tyr	Lys	Val	Leu	Met	Val	Lys	Gly	Ile	Ser	Pro	
	385					390					395					400	
15	Val	Tyr	Pro	His	Phe	Asp	Thr	Met	Ile	Ala	Ala	Tyr	Leu	Leu	Glu	Pro	
					405					410					415		
	Asn	Glu	Lys	Lys	Phe	Asn	Leu	Glu	Asp	Leu	Ser	Leu	Lys	Phe	Leu	Gly	
				420					425					430			
20	Tyr	Lys	Met	Thr	Ser	Tyr	Gln	Glu	Leu	Met	Ser	Phe	Ser	Ser	Pro	Leu	
			435					440					445				
	Phe	Gly	Phe	Ser	Phe	Ala	Asp	Val	Pro	Val	Asp	Lys	Ala	Ala	Asn	Tyr	
	450						455					460					
	Ser	Cys	Glu	Asp	Ala	Asp	Ile	Thr	Tyr	Arg	Leu	Tyr	Lys	Ile	Leu	Ser	
	465					470					475					480	
25	Met	Lys	Leu	His	Glu	Ala	Glu	Leu	Glu	Asn	Val	Phe	Tyr	Arg	Ile	Glu	
					485					490					495		
	Met	Pro	Leu	Val	Asn	Val	Leu	Ala	Arg	Met	Glu	Leu	Asn	Gly	Val	Tyr	
				500					505					510			
30	Val	Asp	Thr	Glu	Phe	Leu	Lys	Lys	Leu	Ser	Glu	Glu	Tyr	Gly	Lys	Lys	
			515					520					525				
	Leu	Glu	Glu	Leu	Ala	Glu	Lys	Ile	Tyr	Gln	Ile	Ala	Gly	Glu	Pro	Phe	
	530						535					540					
	Asn	Ile	Asn	Ser	Pro	Lys	Gln	Val	Ser	Lys	Ile	Leu	Phe	Glu	Lys	Leu	
	545					550					555					560	
35	Gly	Ile	Lys	Pro	Arg	Gly	Lys	Thr	Thr	Lys	Thr	Gly	Glu	Tyr	Ser	Thr	
					565					570					575		

WO 98/35060

PCT/US98/02791

120

	Arg	Ile	Glu	Val	Leu	Glu	Glu	Ile	Ala	Asn	Glu	His	Glu	Ile	Val	Pro
				580					585					590		
	Leu	Ile	Leu	Glu	Tyr	Arg	Lys	Ile	Gln	Lys	Leu	Lys	Ser	Thr	Tyr	Ile
			595					600					605			
5	Asp	Thr	Leu	Pro	Lys	Leu	Val	Asn	Pro	Lys	Thr	Gly	Arg	Ile	His	Ala
	610						615					620				
	Ser	Phe	His	Gln	Thr	Gly	Thr	Ala	Thr	Gly	Arg	Leu	Ser	Ser	Ser	Asp
	625					630					635					640
10	Pro	Asn	Leu	Gln	Asn	Leu	Pro	Thr	Lys	Ser	Glu	Glu	Gly	Lys	Glu	Ile
					645					650					655	
	Arg	Lys	Ala	Ile	Val	Pro	Gln	Asp	Pro	Asp	Trp	Trp	Ile	Val	Ser	Ala
				660					665					670		
	Asp	Tyr	Ser	Gln	Ile	Glu	Leu	Arg	Ile	Leu	Ala	His	Leu	Ser	Gly	Asp
			675					680					685			
15	Glu	Asn	Leu	Val	Lys	Ala	Phe	Glu	Glu	Gly	Ile	Asp	Val	His	Thr	Leu
	690						695					700				
	Thr	Ala	Ser	Arg	Ile	Tyr	Asn	Val	Lys	Pro	Glu	Glu	Val	Asn	Glu	Glu
	705				710						715					720
20	Met	Arg	Arg	Val	Gly	Lys	Met	Val	Asn	Phe	Ser	Ile	Ile	Tyr	Gly	Val
					725					730					735	
	Thr	Pro	Tyr	Gly	Leu	Ser	Val	Arg	Leu	Gly	Ile	Pro	Val	Lys	Glu	Ala
				740					745					750		
	Glu	Lys	Met	Ile	Ile	Ser	Tyr	Phe	Thr	Leu	Tyr	Pro	Lys	Val	Arg	Ser
			755					760					765			
25	Tyr	Ile	Gln	Gln	Val	Val	Ala	Glu	Ala	Lys	Glu	Lys	Gly	Tyr	Val	Arg
	770						775					780				
	Thr	Leu	Phe	Gly	Arg	Lys	Arg	Asp	Ile	Pro	Gln	Leu	Met	Ala	Arg	Asp
	785					790					795					800
30	Lys	Asn	Thr	Gln	Ser	Glu	Gly	Glu	Arg	Ile	Ala	Ile	Asn	Thr	Pro	Ile
				805						810					815	
	Gln	Gly	Thr	Ala	Ala	Asp	Ile	Ile	Lys	Leu	Ala	Met	Ile	Asp	Ile	Asp
				820					825					830		
	Glu	Glu	Leu	Arg	Lys	Arg	Asn	Met	Lys	Ser	Arg	Met	Ile	Ile	Gln	Val
			835					840					845			
35	His	Asp	Glu	Leu	Val	Phe	Glu	Val	Pro	Asp	Glu	Glu	Lys	Glu	Glu	Leu
	850						855						860			

121

Val Asp Leu Val Lys Asn Lys Met Thr Asn Val Val Lys Leu Ser Val
865 870 875 880

Pro Leu Glu Val Asp Ile Ser Ile Gly Lys Ser Trp Ser
885 890

5 (2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 610 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Lys Glu Leu Gln Leu Tyr Glu Glu Ala Glu Pro Thr Gly Tyr Glu
1 5 10 15
Ile Val Lys Asp His Lys Thr Phe Glu Asp Leu Ile Glu Lys Leu Lys
20 25 30
Glu Val Pro Ser Phe Ala Leu Asp Leu Glu Thr Ser Ser Leu Asp Pro
35 40 45
Phe Asn Cys Glu Ile Val Gly Ile Ser Val Ser Phe Lys Pro Lys Thr
50 55 60
Ala Tyr Tyr Ile Pro Leu His His Arg Asn Ala Gln Asn Leu Asp Glu
65 70 75 80
Thr Leu Val Leu Ser Lys Leu Lys Glu Ile Leu Glu Asp Pro Ser Ser
85 90 95
Lys Ile Val Gly Gln Asn Leu Lys Tyr Asp Tyr Lys Val Leu Met Val
100 105 110
Lys Gly Ile Ser Pro Val Tyr Pro His Phe Asp Thr Met Ile Ala Ala
115 120 125
Tyr Leu Leu Glu Pro Asn Glu Lys Lys Phe Asn Leu Glu Asp Leu Ser
130 135 140
Leu Lys Phe Leu Gly Tyr Lys Met Thr Ser Tyr Gln Glu Leu Met Ser
145 150 155 160
Phe Ser Ser Pro Leu Phe Gly Phe Ser Phe Ala Asp Val Pro Val Asp
165 170 175
Lys Ala Ala Asn Tyr Ser Cys Glu Asp Ala Asp Ile Thr Tyr Arg Leu
35

WO 98/35060

PCT/US98/02791

122

	180				185				190							
	Tyr	Lys	Ile	Leu	Ser	Met	Lys	Leu	His	Glu	Ala	Glu	Leu	Glu	Asn	Val
			195				200						205			
5	Phe	Tyr	Arg	Ile	Glu	Met	Pro	Leu	Val	Asn	Val	Leu	Ala	Arg	Met	Glu
		210					215					220				
	Leu	Asn	Gly	Val	Tyr	Val	Asp	Thr	Glu	Phe	Leu	Lys	Lys	Leu	Ser	Glu
	225					230					235					240
	Glu	Tyr	Gly	Lys	Lys	Leu	Glu	Glu	Leu	Ala	Glu	Lys	Ile	Tyr	Gln	Ile
				245						250					255	
10	Ala	Gly	Glu	Pro	Phe	Asn	Ile	Asn	Ser	Pro	Lys	Gln	Val	Ser	Lys	Ile
				260							265				270	
	Leu	Phe	Glu	Lys	Leu	Gly	Ile	Lys	Pro	Arg	Gly	Lys	Thr	Thr	Lys	Thr
			275					280							285	
15	Gly	Glu	Tyr	Ser	Thr	Arg	Ile	Glu	Val	Leu	Glu	Glu	Ile	Ala	Asn	Glu
		290					295					300				
	His	Glu	Ile	Val	Pro	Leu	Ile	Leu	Glu	Tyr	Arg	Lys	Ile	Gln	Lys	Leu
	305					310					315					320
	Lys	Ser	Thr	Tyr	Ile	Asp	Thr	Leu	Pro	Lys	Leu	Val	Asn	Pro	Lys	Thr
						325					330				335	
20	Gly	Arg	Ile	His	Ala	Ser	Phe	His	Gln	Thr	Gly	Thr	Ala	Thr	Gly	Arg
				340							345				350	
	Leu	Ser	Ser	Ser	Asp	Pro	Asn	Leu	Gln	Asn	Leu	Pro	Thr	Lys	Ser	Glu
			355					360					365			
25	Glu	Gly	Lys	Glu	Ile	Arg	Lys	Ala	Ile	Val	Pro	Gln	Asp	Pro	Asp	Trp
		370					375					380				
	Trp	Ile	Val	Ser	Ala	Asp	Tyr	Ser	Gln	Ile	Glu	Leu	Arg	Ile	Leu	Ala
	385					390					395					400
	His	Leu	Ser	Gly	Asp	Glu	Asn	Leu	Val	Lys	Ala	Phe	Glu	Glu	Gly	Ile
				405							410				415	
30	Asp	Val	His	Thr	Leu	Thr	Ala	Ser	Arg	Ile	Tyr	Asn	Val	Lys	Pro	Glu
				420						425				430		
	Glu	Val	Asn	Glu	Glu	Met	Arg	Arg	Val	Gly	Lys	Met	Val	Asn	Phe	Ser
			435					440					445			
35	Ile	Ile	Tyr	Gly	Val	Thr	Pro	Tyr	Gly	Leu	Ser	Val	Arg	Leu	Asn	Ile
		450					455					460				
	Pro	Val	Lys	Glu	Ala	Glu	Lys	Met	Ile	Ile	Ser	Tyr	Phe	Thr	Leu	Tyr
	465					470					475					480

124

- (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

5 Arg Arg Ser Ala Lys Ala Ile Asn Phe Gly Leu Ile Tyr Gly
 1 5 10

(2) INFORMATION FOR SEQ ID NO:13:

- 10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 14 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

15 Arg Arg Ala Ala Lys Thr Ile Asn Phe Gly Val Leu Tyr Gly
 1 5 10

(2) INFORMATION FOR SEQ ID NO:14:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 14 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

25 Arg Asp Asn Ala Lys Thr Phe Ile Tyr Gly Phe Leu Tyr Gly
 1 5 10

(2) INFORMATION FOR SEQ ID NO:15:

- 30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 14 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant

125

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

5 Arg Arg Val Gly Lys Met Val Asn Phe Ser Ile Ile Tyr Gly
 1 5 10

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 14 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

15 Arg Gln Ala Ala Lys Ala Ile Thr Phe Gly Ile Leu Tyr Gly
 1 5 10

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 14 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

25 Arg Arg Ala Gly Lys Met Val Asn Phe Ser Ile Ile Tyr Gly
 1 5 10

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 11 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

127

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Met	Ile	Val	Ser	Asp	Ile	Glu	Ala	Asn	Ala
1			5						10

(2) INFORMATION FOR SEQ ID NO:22:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 26 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

10 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GACGTTTCAA GCGCTAGGGC AAAAGA

26

(2) INFORMATION FOR SEQ ID NO:23:

- 15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 31 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GTATATTATA GAGTAGTTAA CCATCTTTCC A

31

(2) INFORMATION FOR SEQ ID NO:24:

- 25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

WO 98/35060

PCT/US98/02791

128

Phe Leu Phe Asp Gly Thr
1 5

(2) INFORMATION FOR SEQ ID NO:25:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Leu Leu Val Asp Gly His
1 5

(2) INFORMATION FOR SEQ ID NO:26:

- 15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 10 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Ser Leu Ile Thr Gly Asp Lys Asp Met Leu
1 5 10

(2) INFORMATION FOR SEQ ID NO:27:

- 25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 10 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Arg Ile Leu Thr Ala Asp Lys Asp Leu Tyr

WO 98/35060

PCT/US98/02791

129

1

5

10

(2) INFORMATION FOR SEQ ID NO:28:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

10 GTAGGCCAGG GCTGTGCCGG CAAAGAGAAA TAGTC

35

(2) INFORMATION FOR SEQ ID NO:29:

- 15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

GAAGCATATC CTTGGCGCCG GTTATTATGA AAATC

35

20 (2) INFORMATION FOR SEQ ID NO:30:

- 25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

CACCAGACGG GTACCGCCAC TGGCAGGTTG

30

(2) INFORMATION FOR SEQ ID NO:31:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 48 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

TATAGAGTAG TTAACCATCT TTCCAACCCG TTTCATTTCT TCGAACAC

48

(2) INFORMATION FOR SEQ ID NO:32:

- 10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 48 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

15 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

TATAGAGTAG TTAACCATCT TTCCAACCCG TTGCATTTCT TCGAACAC

48

(2) INFORMATION FOR SEQ ID NO:33:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 48 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

TATAGAGTAG TTAACCATCT TTCCAACCCG GTTCATTTCT TCGAACAC

48

(2) INFORMATION FOR SEQ ID NO:34:

- 30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 48 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both

(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

TATAGAGTAG TTAACCATCT TTCCAACCCG ATGCATTTCT TCGAACAC

48

5 (2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: both

(D) TOPOLOGY: both

10

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

AAGATGGTTA ACGCGTCTAT AATATACGG

29

(2) INFORMATION FOR SEQ ID NO:36:

15 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: both

(D) TOPOLOGY: both

20

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

CAAGAGGCAC AGAGAGTTTC ACC

23

(2) INFORMATION FOR SEQ ID NO:37:

25

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: both

(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

GTATATTATA GAGGAGTTAA CCATCTTTCC

30

(2) INFORMATION FOR SEQ ID NO:38:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 29 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

AAGATGGTTA ACTTCTCTAT AATATACGG

29

(2) INFORMATION FOR SEQ ID NO:39:

- 15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 48 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

20 TATAGAGTAG TTAACCATCT TTCCAACCCG GTACATGTCT TCGTTCAC

48

(2) INFORMATION FOR SEQ ID NO:40:

- 25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 48 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

WO 98/35060

PCT/US98/02791

133

TATAGAGTAG TTAACCATCT TTCCAACCCG CAACATGTCT TCGTTCAC

48

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

5

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

10 CTTGGCCGCC CGATGCATCA GGGGGTC

27

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

15

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

CTTGGCCGCC CGCTTCATGA GGGGGTCCAC

30

20 (2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:

25

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

CTTGGCCGCC CTGTACATCA GGGGGTC

27

(2) INFORMATION FOR SEQ ID NO:44:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

GTATATTATA GAGGTGTTAA CCATCTTTCC

30

(2) INFORMATION FOR SEQ ID NO:45:

- 10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 34 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

15 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

GGGAGACCGG AATTCTCCTT CATTAATTCC TATA

34

(2) INFORMATION FOR SEQ ID NO:46:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 49 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

TGGAGACCCT GGA ACTATAG GAATTAATGA AGGAGAATTC CGGTCTCCC

49

(2) INFORMATION FOR SEQ ID NO:47:

- 30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both

135

(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

GTATTTTGGT ATGCTTGTGC

20

5 (2) INFORMATION FOR SEQ ID NO:48:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: both

10 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

CTATTTTGGA ATATATGTGC CT

22

(2) INFORMATION FOR SEQ ID NO:49:

15 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: both

(D) TOPOLOGY: both

20 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

ACGAACATTC TACAAGTTAC

20

(2) INFORMATION FOR SEQ ID NO:50:

25 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: both

(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

WO 98/35060

PCT/US98/02791

136

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

TTTCAGAGAA ACTGACCTGT

20

(2) INFORMATION FOR SEQ ID NO:51:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 21 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

GATAAATGCC AACATGTTG T

21

(2) INFORMATION FOR SEQ ID NO:52:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

20 TGCTCTCAGG ATTCCTCCA

20

(2) INFORMATION FOR SEQ ID NO:53:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

AGCTTGAGAC CTCTGTGTCC

20

(2) INFORMATION FOR SEQ ID NO:54:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

5

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

10 ATTCAGAAGA AACAGTGATG GT

22

(2) INFORMATION FOR SEQ ID NO:55:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

15

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

TTGGAGTCGC AAGCTGAACT AGC

23

20 (2) INFORMATION FOR SEQ ID NO:56:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

25

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

GCCTGAGTGA CAGAGTGAGA ACC

23

(2) INFORMATION FOR SEQ ID NO:57:

WO 98/35060

PCT/US98/02791

138

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

CCCCTAGGT TGTAAGCTCC ATGA

24

(2) INFORMATION FOR SEQ ID NO:58:

- 10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

15 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

TACTATGTGC CAGGCTCTGT CCTA

24

(2) INFORMATION FOR SEQ ID NO:59:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

ACTCATGAAG GTGACAGTTC

20

(2) INFORMATION FOR SEQ ID NO:60:

- 30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both

WO 98/35060

PCT/US98/02791

139

(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

GTGTTGTIGA CCTATTGCAT

20

5 (2) INFORMATION FOR SEQ ID NO:61:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

10

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

ATCTCTGTTT CCTCCCTGTT

20

(2) INFORMATION FOR SEQ ID NO:62:

- 15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

20

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

CTTATTGGCC TTGAAGGTAG

20

(2) INFORMATION FOR SEQ ID NO:63:

- 25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

AGCCCGTGTT GGAACCATGA CTG

23

(2) INFORMATION FOR SEQ ID NO:64:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 23 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

TACATAGCGA GACTCCATCT CCC

23

(2) INFORMATION FOR SEQ ID NO:65:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

20 TTTATGCGAG CGTATGGATA

20

(2) INFORMATION FOR SEQ ID NO:66:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

WO 98/35060

PCT/US98/02791

141

CACCACCATT GATCTGGAAG

20

(2) INFORMATION FOR SEQ ID NO:67:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 16 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: both
 - (D) TOPOLOGY: both

5

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

10 CCAACCACAC TGGGAA

16

(2) INFORMATION FOR SEQ ID NO:68:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 16 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: both
 - (D) TOPOLOGY: both

15

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

AACAGTTGCC CACGGT

16

20 (2) INFORMATION FOR SEQ ID NO:69:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: both
 - (D) TOPOLOGY: both

25

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

CATGAAATGC TGACTGGGTA

20

(2) INFORMATION FOR SEQ ID NO:70:

142

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both
- (ii) MOLECULE TYPE: cDNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:
 TCAATTTATG TGCAGCCAAT 20
- (2) INFORMATION FOR SEQ ID NO:71:
- 10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both
- 15 (ii) MOLECULE TYPE: cDNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:
 CATAGCGAGA CTCCATCTCC 20
- (2) INFORMATION FOR SEQ ID NO:72:
- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both
- (ii) MOLECULE TYPE: cDNA
- 25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:
 GGGAGAGGGC AAAGATCGAT 20
- (2) INFORMATION FOR SEQ ID NO:73:
- 30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both

143

(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

AACACTAGTG ACATTATTTT CA

22

5 (2) INFORMATION FOR SEQ ID NO:74:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: both

10 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

AGCTAGGCCT GAAGGCTTCT

20

(2) INFORMATION FOR SEQ ID NO:75:

15 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: both

(D) TOPOLOGY: both

20 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

CCCTAGTGGA TGATAAGAAT AATC

24

(2) INFORMATION FOR SEQ ID NO:76:

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: both

(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

GGACAGATGA TAAATACATA GGATGGATGG

30

(2) INFORMATION FOR SEQ ID NO:77:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: both
(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

TTCTCTTACA AACTGCCCC

20

(2) INFORMATION FOR SEQ ID NO:78:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: both
(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

20 ATTTGGATGG CTTGACAGAG

20

(2) INFORMATION FOR SEQ ID NO:79:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: both
(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

WO 98/35060

PCT/US98/02791

145

ACATTCTAAG ACTTTCCCAA T

21

(2) INFORMATION FOR SEQ ID NO:80:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: both
 - (D) TOPOLOGY: both

5

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

10 AGAGCATGCA CCCTGAATTG

20

(2) INFORMATION FOR SEQ ID NO:81:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: both
 - (D) TOPOLOGY: both

15

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

AAGAACCATG CGATACGACT

20

20 (2) INFORMATION FOR SEQ ID NO:82:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: both
 - (D) TOPOLOGY: both

25

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

CATTCTAGA TGGGTAAAGC

20

(2) INFORMATION FOR SEQ ID NO:83:

146

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

GCTTAGTCAT ACGAGCGG

18

(2) INFORMATION FOR SEQ ID NO:84:

- 10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

15 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

TCCACAGCCA TGTAACC

18

(2) INFORMATION FOR SEQ ID NO:85:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 16 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

CCCCGGAGCA AGTTCA

16

(2) INFORMATION FOR SEQ ID NO:86:

- 30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both

WO 98/35060

PCT/US98/02791

147

(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

CAGCCCAAAG CCAGATTA

18

5 (2) INFORMATION FOR SEQ ID NO:87:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: both

10 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

ATATGTGAGT CAATCCCCA AG

22

(2) INFORMATION FOR SEQ ID NO:88:

15 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: both

(D) TOPOLOGY: both

20 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

TGTATTAGTC AATGTTCTCC AG

22

(2) INFORMATION FOR SEQ ID NO:89:

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 19 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: both

(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

WO 98/35060

PCT/US98/02791

148

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

CAGCTGCCCT AGTCAGCAC

19

(2) INFORMATION FOR SEQ ID NO:90:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

GCTTCCGAGT GCAGGTCACA

20

(2) INFORMATION FOR SEQ ID NO:91:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 21 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

20 ATTCTGGGCG CACAAGAGTG A

21

(2) INFORMATION FOR SEQ ID NO:92:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:92:

WO 98/35060

PCT/US98/02791

149

ACATCTCCCC TACCGCTATA

20

(2) INFORMATION FOR SEQ ID NO:93:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 33 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: both
(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:93:

10 GAAGTTCACC ATCCGGCCGA CCCGTCGCAT TTC

33

WO 98/35060

- 149.1 -

PCT/US98/02791

Applicant's or agent's file reference number: 0942.425PC02	International application No. TO BE ASSIGNED
---------------------------------------------------------------	----------------------------------------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM (JAPAN)
(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>42</u> , line <u>14</u> .	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection (NRRL)	
Address of depositary institution (including postal code and country) 1815 North University Street Peoria, Illinois 61604 United States of America	
Date of deposit 30 September 1994	Accession Number: NRRL B-21238
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
<i>E. coli</i> DH10B(pUC-Tne)	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application <i>Hal Jander</i>	<input type="checkbox"/> This sheet was received by the International Bureau on:

WO 98/35060

- 149.2 -

PCT/US98/02791

Applicant's or agent's file reference number: 0942.425PC02	International application No. TO BE ASSIGNED
---------------------------------------------------------------	----------------------------------------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM (JAPAN)
(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>58</u> , line <u>2</u> .	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection (NRRL)	
Address of depositary institution (including postal code and country) 1815 North University Street Peoria, Illinois 61604 United States of America	
Date of deposit 30 September 1994	Accession Number: NRRL B-21338
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
<i>E.coli</i> DH10B(pUC19-Tne)	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application <i>Hal Saunders</i>	<input type="checkbox"/> This sheet was received by the International Bureau on:

WHAT IS CLAIMED IS:

1. A method of identifying, analyzing or typing a polymorphic DNA fragment in a sample of DNA, said method comprising contacting said sample of DNA with one or more DNA polymerases substantially reduced in the ability to add one or more non-templated nucleotides to the 3' terminus of a DNA molecule, amplifying said polymorphic DNA fragment within said sample and analyzing said amplified polymorphic DNA fragment.
5
2. A method of producing amplified copies of a polymorphic DNA fragment which comprise substantially no non-templated 3' terminal nucleotides, said method comprising contacting a DNA sample with one or more DNA polymerases substantially reduced in the ability to add one or more non-templated nucleotides to the 3' terminus of a DNA molecule and amplifying said polymorphic DNA fragment within said DNA sample.
10
3. A method of cloning a DNA molecule comprising contacting said DNA molecule with one or more DNA polymerases substantially reduced in the ability to add one or more non-templated nucleotides to the 3' terminus of a DNA molecule, amplifying said DNA molecule and inserting said DNA molecule into a vector.
15
4. The method of claim 3, wherein said vector is blunt-ended.
5. The method of claim 1, wherein said polymorphic DNA fragment is selected from the group of polymorphic DNA fragments comprising a minisatellite DNA fragment, a microsatellite DNA fragment and a STR DNA fragment.
20
6. The method of claim 1, wherein said polymerases are thermostable DNA polymerases.
25

7. The method of claim 6, wherein said thermostable DNA polymerases are *Thermotoga* DNA polymerases and mutants or derivatives thereof.

5 8. The method of claim 7, wherein said DNA polymerase is a *Tne* or *Tma* DNA polymerase.

9. The method of claim 1, wherein said DNA polymerases are substantially reduced in 3'-5' exonuclease activity.

10. The method of claim 1, wherein said DNA polymerases are substantially reduced in 5'-3' exonuclease activity.

10 11. The method of claim 9, wherein said DNA polymerases are substantially reduced in 5'-3' exonuclease activity.

15 12. The method of claim 1, wherein said DNA polymerases contain one or more modifications or mutations which reduce the ability of the polymerase to add one or more non-templated 3' nucleotides to a synthesized nucleic acid molecule.

13. The method of claim 12, wherein said DNA polymerases are substantially reduced in at least one activity selected from the group consisting of:

- (a) 3'-5' exonuclease activity; and
- (b) 5'-3' exonuclease activity.

20 14. The method of claim 13, wherein said polymerases have substantially reduced 3'-5' exonuclease and 5'-3' exonuclease activity.

15. The method of claim 13, wherein said polymerase is substantially

reduced in 3'-5' exonuclease activity.

16. The method of claim 12, wherein said polymerases comprise one or more mutations or modifications in the O-helix of said polymerase.

5 17. The method of claim 16, wherein said O-helix is defined as RXXXKXXXFXXXYYX (SEQ ID NO:11), wherein X is any amino acid.

18. The method of claim 17, wherein said mutation or modification is at position R (Arg) and/or F (Phe) and/or K (Lys) of said O-helix or combinations thereof.

10 19. The method of claim 16, wherein said mutation or modification is an amino acid substitution at position R and/or F and/or K of said O-helix or combinations thereof.

20. The method of claim 1, wherein said polymerase is selected from the group consisting of:

- 15 *Tne* N'Δ219, D323A;
Tne N'Δ283, D323A;
Tne N'Δ284, D323A;
Tne N'Δ193, D323A;
Tne D137A, D323A;
Tne D8A, D323A;
20 *Tne* G195D, D323A;
Tne G37D, D323A;
Tne N'Δ283;
Tne D137A, D323A, R722K;
Tne D137A, D323A, R722Y;
25 *Tne* D137A, D323A, R722L;
Tne D137A, D323A, R722H;

- Tne* D137A, D323A, R722Q;
Tne D137A, D323A, F730Y;
Tne D137A, D323A, K726R;
Tne D137A, D323A, K726H;
5 *Tne* D137A, D323A, R722K, F730Y;
Tne D137A, D323A, R722K, K726R;
Tne D137A, D323A, R722K, K726H;
Tne D137A, D323A, R722H, F730Y;
Tne D137A, D323A, R722H, K726R;
10 *Tne* D137A, D323A, R722H, K726H;
Tne D137A, D323A, R722Q, F730Y;
Tne D137A, D323A, R722Q, K726R;
Tne D137A, D323A, R722Q, K726H;
Tne D137A, D323A, R722N, F730Y;
15 *Tne* D137A, D323A, R722N, K726R;
Tne D137A, D323A, R722N, K726H;
Tne D137A, D323A, F730S;
Tne N' Δ 283, D323A, R722K/H/Q/N/Y/L;
Tne N' Δ 219, D323A, R722K;
20 *Tne* N' Δ 219, D323A, F730Y;
Tne N' Δ 219, D323A, K726R;
Tne N' Δ 219, D323A, K726H;
Tne D137A, D323A, F730S, R722K/Y/Q/N/H/L, K726R/H;
Tne D137A, D323A, F730T, R722K/Y/Q/N/H/L, K726R/H;
25 *Tne* D137A, D323A, F730T;
Tne F730S;
Tne F730A;
Tne K726R;
Tne K726H; and
30 *Tne* D137A, D323A, R722N.

21. A method of determining the relationship between a first individual and a second individual, said method comprising comparing a population of amplified DNA molecules in a sample of DNA from said first individual to that of said second individual prepared according to the method of claim 1.

5 22. The method of claim 21, wherein said sample of DNA from said first individual is a known sample and said sample of DNA from said second individual is an unknown sample.

10 23. A kit for the identification, analysis or typing of a polymorphic DNA fragment, said kit comprising one or more DNA polymerases substantially reduced in the ability to add one or more non-templated nucleotides to the 3' terminus of a DNA molecule.

15 24. The kit of claim 23, said kit further comprising one or more components selected from the group consisting of one or more DNA primers, one or more deoxynucleoside triphosphates, and a buffer suitable for use in the identification, analysis or typing of a polymorphic DNA fragment.

 25. The kit of claim 23, wherein said polymerases are thermostable DNA polymerases.

 26. The kit of claim 25, wherein thermostable DNA polymerases are *Thermotoga* DNA polymerases.

20 27. The kit of claim 23, wherein said DNA polymerase is substantially reduced in 3'-5' exonuclease activity.

 28. The kit of claim 23, wherein said DNA polymerase is substantially reduced in 5'-3' exonuclease activity.

29. The kit of claim 23, wherein said DNA polymerases comprise one or more modifications or mutations which reduce the ability of the polymerase to add one or more non-templated 3' nucleotides to a synthesized nucleic acid molecule.

5 30. The kit of claim 29, wherein said polymerases comprise one or more mutations in the O-helix of said polymerase.

31. The kit of claim 30, wherein said O-helix is defined as RXXXKXXXFXXXYX (SEQ ID NO:11), wherein X is any amino acid.

10 32. The kit of claim 31, wherein said mutation or modification is at position R (Arg) and/or F (Phe) and/or K (Lys) of said O-helix or combinations thereof.

33. The method of claim 31, wherein said mutation or modification is an amino acid substitution at position R and/or F and/or K of said O-helix or combinations thereof.

15 34. A polymerase which has been modified or mutated to reduce, substantially reduce or eliminate the ability of the polymerase to add non-templated 3' nucleotides to a synthesized nucleic acid molecule.

35. The polymerase of claim 34, wherein said polymerase is a DNA or RNA polymerase.

20 36. The polymerase of claim 34, wherein said polymerase is substantially pure.

37. The polymerase of claim 34, wherein said polymerase is mesophilic or thermostable.

38. The polymerase of claim 34, wherein said polymerase is selected from the group consisting of *Tne* DNA polymerase, *Taq* DNA polymerase, *Tma* DNA polymerase, *Tth* DNA polymerase, *Tli* DNA polymerase, VENT™ DNA polymerase, *Pfu* DNA polymerase, DEEPVENT™ DNA polymerase, *Pwo* DNA polymerase, *Bst* DNA polymerase, *Bca* DNA polymerase, *Tfl* DNA polymerase, and mutants, variants and derivatives thereof.

39. The polymerase of claim 34, wherein said polymerase is substantially reduced in at least one activity selected from the group consisting of:

(a) 3'→5' exonuclease activity; and

(b) 5'→3' exonuclease activity.

40. The polymerase of claim 39, wherein said polymerase is substantially reduced in 3'-5' exonuclease activity.

41. The polymerase of claim 39, wherein said polymerase is substantially reduced in 5'-3' exonuclease activity.

42. The polymerase of claim 41, which is modified or mutated to reduce or eliminate 3'-5' exonuclease activity.

43. The polymerase of claim 34, which comprises one or more modifications or mutations in the O-helix of said polymerase.

44. The polymerase of claim 43, wherein said O-helix is defined as RXXXKXXXFXXXYX (SEQ ID NO:11), wherein X is any amino acid.

45. The polymerase of claim 44, wherein said mutation or modification is at position R (Arg) and/or F (Phe) and/or K (Lys) of said O-helix or combinations thereof.

46. The polymerase of claim 44, wherein said mutation or modification is an amino acid substitution at position R and/or F and/or K of said O-helix or combinations thereof.

5 47. The polymerase of claim 46, wherein R (Arg) is substituted with an amino acid selected from the group consisting of Ala, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Try and Val.

48. The polymerase of claim 46, wherein R (Arg) is substituted with Lys or His.

10 49. The polymerase of claim 46, wherein F (Phe) is substituted with an amino acid selected from the group consisting of Ala, Asn, Arg, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Pro, Ser, Thr, Trp, Try and Val.

50. The polymerase of claim 46, wherein K (Lys) is substituted with an amino acid selected from the group consisting of Ala, Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Met, Phe, Pro, Ser, Thr, Trp, Try and Val.

15 51. The polymerase of claim 46, wherein K (Lys) is substituted with Arg or His.

52. A mutant *Tne* DNA polymerase protein selected from the group consisting of:

Tne N'Δ219, D323A;

20 *Tne* N'Δ283, D323A;

Tne N'Δ284, D323A;

Tne N'Δ193, D323A;

Tne D137A, D323A;

Tne D8A, D323A;

25 *Tne* G195D, D323A;

- Tne G37D, D323A;*
- Tne N'Δ283;*
- Tne D137A, D323A, R722K;*
- Tne D137A, D323A, R722Y;*
- 5 *Tne D137A, D323A, R722L;*
- Tne D137A, D323A, R722H;*
- Tne D137A, D323A, R722Q;*
- Tne D137A, D323A, F730Y;*
- Tne D137A, D323A, K726R;*
- 10 *Tne D137A, D323A, K726H;*
- Tne D137A, D323A, R722K, F730Y;*
- Tne D137A, D323A, R722K, K726R;*
- Tne D137A, D323A, R722K, K726H;*
- Tne D137A, D323A, R722H, F730Y;*
- 15 *Tne D137A, D323A, R722H, K726R;*
- Tne D137A, D323A, R722H, K726H;*
- Tne D137A, D323A, R722Q, F730Y;*
- Tne D137A, D323A, R722Q, K726R;*
- Tne D137A, D323A, R722Q, K726H;*
- 20 *Tne D137A, D323A, R722N, F730Y;*
- Tne D137A, D323A, R722N, K726R;*
- Tne D137A, D323A, R722N, K726H;*
- Tne D137A, D323A, F730S;*
- Tne N'Δ283, D323A, R722K/H/Q/N/Y/L;*
- 25 *Tne N'Δ219, D323A, R722K;*
- Tne N'Δ219, D323A, F730Y;*
- Tne N'Δ219, D323A, K726R;*
- Tne N'Δ219, D323A, K726H;*
- Tne D137A, D323A, F730S, R722K/Y/Q/N/H/L, K726R/H;*
- 30 *Tne D137A, D323A, F730T, R722K/Y/Q/N/H/L, K726R/H;*
- Tne D137A, D323A, F730T;*

The F730S;
The F730A;
The K726R;
The K726H; and
The D137A, D323A, R722N.

5

53. A vector comprising a gene encoding the polymerase of claim 34.

54. The vector of claim 53, wherein said gene is operably linked to a promoter.

10 55. The vector of claim 54, wherein said promoter is selected from the group consisting of a λ -P_L promoter, a *tac* promoter, a *trp* promoter, and a *trc* promoter.

56. A host cell comprising the vector of claim 53.

15 57. A method of producing a polymerase, said method comprising:
(a) culturing the host cell of claim 56;
(b) expressing said gene; and
(c) isolating said polymerase from said host cell.

20 58. A method of synthesizing a nucleic acid molecule comprising:
(a) mixing a nucleic acid template with one or more polymerases of claim 34; and
(b) incubating said mixture under conditions sufficient to make a nucleic acid molecule complementary to all or a portion of said template.

59. The method of claim 58, wherein said mixture further comprises one or more nucleotides selected from the group consisting of dATP, dCTP, dGTP, dTTP, dITP, 7-deaza-dGTP, dUTP, ddATP, ddCTP, ddGTP, ddITP,

ddTTP, [α -S]dATP, [α -S]dTTP, [α -S]dGTP, and [α -S]dCTP.

60. The method of claim 59, wherein one or more of said nucleotides are detectably labeled.

61. A method of sequencing a DNA molecule, comprising:

5

(a) hybridizing a primer to a first DNA molecule;

(b) contacting said DNA molecule of step (a) with deoxyribonucleoside triphosphates, one or more DNA polymerases of claim 34, and a terminator nucleotide;

10

(c) incubating the mixture of step (b) under conditions sufficient to synthesize a random population of DNA molecules complementary to said first DNA molecule, wherein said synthesized DNA molecules are shorter in length than said first DNA molecule and wherein said synthesized DNA molecules comprise a terminator nucleotide at their 3' termini; and

15

(d) separating said synthesized DNA molecules by size so that at least a part of the nucleotide sequence of said first DNA molecule can be determined.

62. The method of claim 61, wherein said deoxyribonucleoside triphosphates are selected from the group consisting of dATP, dCTP, dGTP, dTTP, dITP, 7-deaza-dGTP, dUTP, [α -S]dATP, [α -S]dTTP, [α -S]dGTP, and [α -S]dCTP.

20

63. The method of claim 61, wherein said terminator nucleotide is ddTTP, ddATP, ddGTP, ddITP or ddCTP.

64. The method of claim 61, wherein one or more of said deoxyribonucleoside triphosphates is detectably labeled.

25

65. The method of claim 61, wherein one or more of said terminator

nucleotides is detectably labeled.

66. A method for amplifying a double stranded DNA molecule, comprising:

5 (a) providing a first and second primer, wherein said first primer is complementary to a sequence at or near the 3'-termini of the first strand of said DNA molecule and said second primer is complementary to a sequence at or near the 3'-termini of the second strand of said DNA molecule;

10 (b) hybridizing said first primer to said first strand and said second primer to said second strand in the presence of the one or more DNA polymerases of claims 34, under conditions such that a third DNA molecule complementary to said first strand and a fourth DNA molecule complementary to said second strand are synthesized;

(c) denaturing said first and third strand, and said second and fourth strands; and

15 (d) repeating steps (a) to (c) one or more times.

67. A kit for sequencing, amplifying or sequencing a DNA molecule comprising one or more polymerases of claim 34.

20 68. The kit of claim 67, further comprising one or more dideoxyribonucleoside triphosphates and/or one or more deoxyribonucleoside triphosphates.




FIG.1

2/20




FIG.2A

3/20




FIG.2B

4/20




FIG.3




FIG.4

6/20




FIG.5

7/20




FIG. 6

8 / 20




FIG.7

9 / 20




FIG.8

10/20




FIG.9A




FIG.9B

11/20




FIG. 9C




FIG. 9D

12/20




FIG.10A




FIG.10B




FIG.10C






FIG.10D






FIG.12A




FIG.12B

16/20




FIG.12C




FIG.12D




FIG.12E




FIG.12F




FIG.13A




FIG.13B

20/20




FIG.13C