



(86) Date de dépôt PCT/PCT Filing Date: 1993/10/08
 (87) Date publication PCT/PCT Publication Date: 1994/04/28
 (45) Date de délivrance/Issue Date: 2002/02/12
 (85) Entrée phase nationale/National Entry: 1994/06/08
 (86) N° demande PCT/PCT Application No.: CA 1993/000425
 (87) N° publication PCT/PCT Publication No.: 1994/008564
 (30) Priorité/Priority: 1992/10/08 (07/957,998) US

(51) Cl.Int.⁵/Int.Cl.⁵ A61K 9/107
 (72) Inventeurs/Inventors:
 Alakhov, Vallery Yu, CA;
 Sveshnikov, Peter G., RU;
 Severin, Eugenie S., RU;
 KABANOV, ALEXANDER V., CA
 (73) Propriétaires/Owners:
 Alakhov, Vallery Yu, CA;
 Sveshnikov, Peter G., RU;
 Severin, Eugenie S., RU;
 KABANOV, ALEXANDER V., CA
 (74) Agent: DEETH WILLIAMS WALL LLP

(54) Titre : COMPOSITION D'AGENTS ANTINEOPLASIQUES, INCORPOREE DANS DES MICELLES
 (54) Title: COMPOSITION OF ANTINEOPLASTIC AGENTS INCORPORATED IN MICELLES

(57) **Abrégé/Abstract:**

Compositions of anti-neoplastic agents incorporate the agent in micelles of at least one block copolymer of poly(oxyethylene)-poly(oxypropylene) in which the ratio of (oxypropylene) blocks to the (oxyethylene) blocks is from about 0.25 to about 1-5 and the micelles have an average diameter of from about 10 to about 25 nm.

2 125 500

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : A61K 9/107, 47/10</p>	<p>A1</p>	<p>(11) International Publication Number: WO 94/08564 (43) International Publication Date: 28 April 1994 (28.04.94)</p>
<p style="text-align: center;">2125500</p> <p>(21) International Application Number: PCT/CA93/00425 (22) International Filing Date: 8 October 1993 (08.10.93) (30) Priority data: 07/957,998 8 October 1992 (08.10.92) US (71)(72) Applicants and Inventors: ALAKHOV, Valery, Yu [RU/CA]; 20726 Gay Cedars, Baie d'Urfe, Quebec H9X 2T4 (CA). KABANOV, Alexander, V. [RU/CA]; 920 Place Sims, #58, Dorval, Quebec H9S 2A1 (CA). SVESHNIKOV, Peter G. [RU/RU]; Teply Stan str., 2-6, Moscow, 117133 (RU). SEVERIN, Eugenii S. [RU/RU]; Novoslobodskaya str., 57/65-65, Moscow, 103055 (RU).</p>		<p>(74) Agent: BLAKE, CASSELS & GRAYDON; P.O. Box 25, Commerce Court West, Suite 2800, Toronto, Ontario M5L 1A9 (CA). (81) Designated States: AU, BB, BG, BR, BY, CA, CZ, FI, HU, JP, KP, KR, KZ, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.</p>

(54) Title: COMPOSITION OF ANTINEOPLASTIC AGENTS INCORPORATED IN MICELLES

(57) Abstract

Compositions of anti-neoplastic agents incorporate the agent in micelles of at least one block copolymer of poly(oxyethylene)-poly(oxypropylene) in which the ratio of (oxypropylene) blocks to the (oxyethylene) blocks is from about 0.25 to about 1.5 and the micelles have an average diameter of from about 10 to about 25 nm.

2125500

COMPOSITION OF ANTINEOPLASTIC AGENTS INCORPORATED IN MICELLES

The present invention pertains to improvements in pharmaceutical compositions and in particular improvements in pharmaceutical compositions used in chemotherapy.

5 A number of anti-neoplastic agents currently are in use in chemotherapy (see generally "Cutting's Handbook of Pharmacology, 7th Ed., Chapter 13, Csáky and Barnes) and many additional agents are under investigation.

10 Because of their often complex structure, these agents can exhibit low stability in the blood stream. Many are extremely insoluble and possess poor transport properties with respect to cell membranes. In addition, binding of the anti-neoplastic agent with plasma proteins, as well as other nonspecific interactions in the blood stream prior to its
15 reaching its target, can greatly reduce the effective amount actually available to combat the neoplastic cells. Moreover, multidrug resistance often is observed with such agents; i.e., the sensitivity of the neoplastic cells to the agent is observed to decrease, often by a factor of 10^3 ,
20 over the course of treatment and this resistance thereafter may manifest itself even with respect to structurally different anti-neoplastic agents.

In accordance with the present invention, the anti-neoplastic agent of choice (which may include a mixture of several distinct anti-neoplastic agents) is incorporated into a
25 micelle of a block copolymer of poly(oxyethylene)-poly(oxypropylene) in an aqueous dispersion as hereinafter described.

The use of the block copolymer micelle in administrat-
30 ing the anti-neoplastic agent provides non-covalent solubilization which reduces water-instability and increases the

2125500

solubility of the anti-neoplastic agent.

Moreover, while block copolymers of poly(oxyethylene)-poly(oxypropylene) have been used as nonionic surfactants, the effects observed here clearly extend beyond mere solubilization. For example, undesired pretarget protein binding of the anti-neoplastic agent is reduced; i.e., the anti-neoplastic agent appears to be "shielded" from proteins which otherwise would bind to it. Increased sensitivity with respect to the target anti-neoplastic cells also is observed. Finally a reversion in multidrug resistance is observed. While the multidrug resistance (MDR) phenomenon is not fully understood, it is accompanied by an overexpression of a transmembrane P-glycoprotein of M_r about 170 kD (P-170) which mediates the ATP-dependent efflux of numerous drugs from such cells (although drug efflux may involve other membrane components of MDR cells as well). The present compositions appear to possess increased cytotoxic activity with respect to P-170 dependent and P-170 independent MDR cancer cells as compared with sensitive cells, thereby reducing the multidrug resistance effect.


A variety of anti-neoplastic agents are suitable for use in the present composition. These include alkaloids such as vinblastine, colchicine, and demecoline; antibiotics such as those of the rhodomycin group as for example as daunorubicin and doxorubicin, those of the mitomycin group as for example mitomycin C and N-methyl mitomycin C, and those of the bleomycin group such as bleomycin A₂; and antifolates such as methotrexate, aminopterin, and dideazatetrahydrofolic acid. It will be appreciated that this improvement extends to mixtures of several such agents.

The present invention is not directed to the underlying anti-neoplastic activity of these agents but rather to an improvement in the manifestation of this activity through formulation.

WO 94/08564

PCT/CA93/00425

The block copolymers of poly(oxyethylene)-poly(oxypropylene) generally are characterized by the structural formula:



I

in which each of x and z , independently of the other, has a value of from about 5 to about 100 and y has a value of from about 20 to about 80. Such block copolymers are known (see Stanton, *Am. Perfumer. Cosmet.* 72(4), 54-58 (1958); Schmolka, *Loc. cit.* 82(7), 25-30 (1967); and *Nonionic Surfactants*, Schick, Ed., (Dekker, NY, 1967) 300-371). A number of these copolymers are commercially available under the generic names of "poloxamers"[™] and "pluronic"[™].

The hydrophobic/hydrophilic properties of a given block copolymer depends upon the ratio of the number of oxypropylene groups to the number of oxyethylene groups. For a composition containing a single block copolymer of poly(oxyethylene)-poly(oxypropylene), for example, this relationship, taking into account the molecular masses of the central hydrophobic block and the terminal hydrophilic blocks, can be expressed as follows:

$$n = \frac{y}{x + z} * 1.32$$

in which y is the number of oxypropylene units and x and z are number of oxyethylene units.

Selecting a block copolymer with the appropriate n value depends upon the hydrophobic/hydrophilic properties of the specific anti-neoplastic agent, or the composite hydrophobic/hydrophilic properties of a mixture of anti-neo-

plastic agents, to be formulated. Typically n will range in value from about 0.25 to about 1.5. This range should be viewed not as numerically critical but as expressing the optimum hydrophobic-hydrophilic balance between the predominantly hydrophilic poly(oxyethylene) blocks and the predominantly hydrophobic poly(oxypropylene) blocks.

An important aspect of the present invention involves utilizing mixture of different block copolymers of poly(oxyethylene)-poly(oxypropylene) to achieve a more specific hydrophobic-hydrophilic balance suitable for a given anti-neoplastic agent or mixture of several anti-neoplastic agents, preserving the optimal size of particles. For example, a first block copolymer may have an n of 1.00 whereas a second may have a value of 1.5. If material having an n of 1.3 is desired, a mixture of one weight portion of the first block copolymer and 1.5 weight portion of the second block copolymer can be employed.

A more generalized relationship therefore for such mixtures can be expressed as follows:

$$N = \frac{y_1 * a}{(x_1 + z_1) * (a + b)} + \frac{y_2 * b}{(x_2 + z_2) * (a + b)} * 1.32$$

in which:

y_1 and y_2 are the number of oxypropylene units in the first and second block copolymers, respectively;
 x_1 and z_1 are number of oxyethylene units in the first block copolymer;

x_2 and z_2 are number of oxyethylene units in the second block copolymer;

a is the weight proportion in the first block copolymer; and

b is the weight proportion in the second block copolymer.

If only one block copolymer of poly(oxyethylene)-poly(oxypropylene) is utilized, N will equal n . An analogous relationship will apply to compositions employing more than two block copolymers of poly(oxyethylene)-
5 poly(oxypropylene).

Using the above parameters, one or more block copolymers of poly(oxyethylene)-poly(oxypropylene) are combined so as to have a value for N of from about 0.25 to about 1.5. The combined copolymers form micelles, the value of N
10 affecting in part the size of the micelles thus produced. Typically the micelles will have an average diameter of from about 10 to about 25 nm., although this range can vary widely. The average diameter of any given preparation can be readily determined by quasielastic light scattering tech-
15 niques.

The anti-neoplastic compound or compounds in the copolymer micelle are administered parenterally in aqueous formulations, alone or in combination with other therapeutic agents including other anti-neoplastic agents, steroids,
20 etc., to a mammal suffering from neoplasm and in need of treatment. Parenteral routes of administration include intramuscular, intrathecal, intraperitoneal, intravenous and intra-arterial. Isotonic micellar solution of one or more block copolymers of poly(oxyethylene)-poly(oxypropylene)
25 incorporating one or more anti-neoplastic agents are used for parenteral administration. Dosages typically are those associated with the specific anti-neoplastic agent, although as in every case the regimen must be titrated to the particular neoplasm, the condition of the patient, and the
30 response. For example, an isotonic micellar solution of daunorubicin in the block copolymer micelles is administered so as to provide about 1 mg of daunorubicin per kg of body weight. Vinblastine on the other hand is administered in the same fashion but in accordance with conventional usage
35 at lower doses of from about 0.1 to about 0.2 mg/kg. Often

the amount required can be reduced.

The following examples will serve to further typify the nature of the invention but should not be construed as a limitation on the scope thereof which is defined solely by the appended claims.

Example 1

5 A. A block copolymer of poly(oxyethylene)-poly(oxypropylene) in which $N = 0.25$ (Pluronic F-68) is diluted with RPMI 1640 medium to a final concentration of 2.0% at 4°C. 10 The mixture is incubated for 30 minutes at 37°C and then sterilely filtered through a 0.22 μm filter. An equal volume of a solution of 200 μg daunorubicin in RPMI 1640 medium is added and this mixture is incubated for 30 minutes at 37°C.

15 B. Human ovarian carcinoma cells (CRL157) are precultured in 1% solution of the same block copolymer but without daunorubicin in RPMI 1640 medium supplemented with 10% calf fetal serum. The preparation of part A is added and the mixture is incubated for 60 minutes at 37°C and the cells 20 then washed three times with RPMI 1640 and cultured in RPMI 1640 supplemented with 10% calf fetal serum for 3 days (Prep. A). Cytotoxicity is measured, both for this preparation and a parallel preparation of free daunorubicin (Prep. B), using the method of Alley et al., *Cancer Res.*, 25 48, 589-601 (1988). The results are as follows:

conc. (ng/mL)	50000	10000	2000	400	80	16
---------------	-------	-------	------	-----	----	----

% Inhibition

5

Prep. A	100	100	92	24	6	2
Prep. B	100	81	53	38	20	1

10

Following the same procedure, cytotoxicity is determined against human T-lymphoma (Jurkat) cells:

conc. (ng/mL)	50000	10000	2000	400	80	16	3.2
---------------	-------	-------	------	-----	----	----	-----

15

% Inhibition

20

Prep. A	100	100	100	100	92	33	3
Prep. B	100	100	100	84	51	44	22

Following the same procedure, cytotoxicity is determined against human small cell carcinoma of lung (H-69):

25

conc. (ng/mL)	50000	10000	2000	400	80	16	3.2
---------------	-------	-------	------	-----	----	----	-----

% Inhibition

30

Prep. A	100	100	100	100	100	42	12
Prep. B	100	100	100	91	69	42	20

Example 2

35

Block copolymers of poly(oxyethylene)-poly(oxypropylene) having the ratios of poly(oxypropylene) to poly(oxyethylene) indicated below are dispersed in RPMI 1640 medium at the concentration indicated below. The mixtures are incubated for 40 minutes at 30°C. The average micelle diameter is measured by quasielastic light scattering and the value of N calculated as previously indicated. The results are as follows:

40

	copolymer	conc.	Avg. Diameter	N
5	F-68 ¹	1.0 %	726.0 nm	0.25
	P-85 ²	1.0 %	18.0 nm	1.00
	L-64 ³	1.0 %	20.4 nm	1.50
10	1:1.5 P-85:L-64	0.01 %	17.0 nm	1.30
	1:2.5 F-68:L-64	1.0 %	33.5 nm	1.38

15
 Note 1: $x = 80$, $y = 30$, and $z = 80$
 Note 2: $x = 75/2$, $y = 55$, and $z = 75/2$
 Note 3: $x = 27/2$, $y = 30$, and $z = 27/2$

Example 3

20 A. A 1:1.5 mixture of block copolymers of poly(oxyethylene)-poly(oxypropylene) (Pluronic P-85 and L-64) having individual ratios (n) of (oxypropylene) to (oxyethylene) blocks of 1.00 and 1.50, respectively, and a combined value (N) of 1.30, is diluted with RPMI 1640 medium to a final concentration of 2.0% at 4°C. The mixture is
 25 incubated for 30 minutes at 37°C and then sterilely filtered through a 0.22 μm filter. An equal volume of a solution of 200 μg daunorubicin in RPMI 1640 medium is added and this mixture is incubated for 30 minutes at 37°C.

30 B. Cytotoxicity to human ovarian cancer cells (CRL157) is measured, both for this preparation (Prep. A) and a parallel preparation of free daunorubicin (Prep. B) as described in Example 1B. The results are as follows:

conc. (ng/mL)	50000	10000	2000	400	80	16	3.2
---------------	-------	-------	------	-----	----	----	-----

% Inhibition

5

Prep. A	100	100	100	100	94	53	8
Prep. B	100	100	81	50	29	10	2

10

Example 4

Daunorubicin in the composition of Example 3 is evaluated for cytotoxicity in (i) human T-lymphoma (Jurkat) cells as described in Example 1 and (ii) normal human mononuclear cells. The results are as follows:

15

conc. (ng/mL)	50000	10000	2000	400	80	16	3.2
---------------	-------	-------	------	-----	----	----	-----

Cell

% Inhibition

20

Prep. A	Jur.	100	100	100	100	100	74	28
Prep. B	Jur.	100	100	100	83	59	36	21
Prep. A	Norm.	100	100	91	60	21	5	2
Prep. B	Norm.	100	100	80	58	23	18	1

25

Example 5

30

IC₅₀ values for (i) human T-lymphoma (Jurkat) cells and (ii) normal human mononuclear cells are determined for the daunorubicin composition of Example 3 and compared to those for free daunorubicin. Measurements are made at the indicated intervals of the drug contact with the cells from 15 minutes to 12 hours. The results are as follows:

2125500

time (hours)	0.25	0.50	0.75	1.0	2.0	4.0	8.0	12
--------------	------	------	------	-----	-----	-----	-----	----

Cell

IC₅₀ (ng/mL)

5	Prep. A	Jur.	150	46	25	17	9.0	0.80	0.50	0.30
	Prep. B	Jur.	120	68	35	25	19	16	3.0	5.2
10	Prep. A	Norm.	3570	950	620	450	250	220	160	140
	Prep. B	Norm.	4900	980	405	310	290	275	280	240

Example 6

15 The antineoplastic agent vinblastine is incorporated into the block copolymer mixture described in Example 3. The IC₅₀ of this preparation against SKVO₃ cells, a drug-sensitive human ovarian carcinoma line, is determined to be 0.121 $\mu\text{g/mL}$; the IC₅₀ against SKVLB cells, an MDR subline

20 expressing high levels of P-170 obtained through long term cultivation of SKVO₃ in the presence of vinblastine, is 0.0012 $\mu\text{g/mL}$. The IC₅₀ of free vinblastine against SKVO₃ cells is determined to be 0.095 $\mu\text{g/mL}$; the IC₅₀ against SKVLB cells is 0.615 $\mu\text{g/mL}$.

25

Example 7

The antineoplastic agent mitomycin C is incorporated into the block copolymer mixture described in Example 3. The IC₅₀ of this preparation against SKVO₃ cells is determined to be 0.265 $\mu\text{g/mL}$; the IC₅₀ against SKVLB cells is

30 0.005 $\mu\text{g/mL}$. The IC₅₀ of free mitomycin against SKVO₃ cells is determined to be 0.320 $\mu\text{g/mL}$; the IC₅₀ against SKVLB cells is 1.120 $\mu\text{g/mL}$.

Example 8

The antineoplastic agent methotrexate is incorporated into the block copolymer mixture described in Example 3. The IC₅₀ of this preparation against SKVO₃ cells is determined to be 0.880 µg/mL; the IC₅₀ against SKVLB cells is 0.0175 µg/mL. The IC₅₀ of free methotrexate against SKVO₃ cells is determined to be 1.090 µg/mL; the IC₅₀ against SKVLB cells is 1.340 µg/mL.

Example 9

The antineoplastic agent colchicine is incorporated into the block copolymer mixture described in Example 3. The IC₅₀ of this preparation against SKVO₃ cells is determined to be 0.720 µg/mL; the IC₅₀ against SKVLB cells is 0.045 µg/mL. The IC₅₀ of free colchicine against SKVO₃ cells is determined to be 0.950 µg/mL; the IC₅₀ against SKVLB cells is 7.450 µg/mL.

Example 10

The antineoplastic agent daunorubicin is incorporated into the block copolymer mixture described in Example 3. The IC₅₀ of this preparation against SKVO₃ cells is determined to be 0.600 µg/mL; the IC₅₀ against SKVLB cells is 0.0068 µg/mL. The IC₅₀ of free daunorubicin against SKVO₃ cells is determined to be 0.620 µg/mL; the IC₅₀ against SKVLB cells is 5.850 µg/mL.

25

Example 11

To 30 µL of a 20 mg/mL solution of bovine serum albumin in phosphate buffered saline are added 30 µL of daunorubicin

solution in the block copolymer mixture described in Example 3 (Prep. A). A second formulation (Prep. B) is prepared in parallel fashion using free daunorubicin.

5 The preparations are incubated for 10 minutes at 25°C, and then analyzed by HPLC on a TSK-3000 SW gel-filtration column in PBS containing 0.3 M sodium chloride and 5% acetonitrile. Detection is performed at 280 nm and 470 nm. The portion of the drug bound with BSA is determined as:

$$D_b = S_b/S_f$$

10 in which:

S_b is relative area of the 470 nm peak (corresponding to daunorubicin) which coincides in retention time for the 280 nm peak (corresponding to BSA); and

15 S_f is relative area of the peak (or peaks) corresponding to daunorubicin which does not coincide in retention time of the BSA peak.

The results are as follows:

20

Composition	D_b
Prep. A	0.01
Prep. B	0.39

25

Example 12

Micellar daunorubicin obtained as described in Example 3 (Prep. A) and free daunorubicin (Prep. B) are incubated in the dark at 37°C and cytotoxicity to CRL157 cells is then determined in the manner discussed in Part B of Example 1.

30

The results are as follows:

time (hours)	2	4	12	24	48	96
--------------	---	---	----	----	----	----

5

IC₅₀, µg/mL

Prep. A	9.1	10.05	9.8	10.4	10.7	11.3
Prep. B	400	475	1120	6300	10180	48900

10

Example 13

The daunorubicin composition of Example 3 (Prep. A) is evaluated against daunorubicin-sensitive human breast cancer (MCF-7) and two cell lines demonstrating resistance: daunorubicin/verapamil-resistant (MCF-7AU) not expressing P-170, and daunorubicin-resistant, verapamil-sensitive (Dox-MCF-7), expressing P-170, in each case in comparison to free daunorubicin (Prep. B). The results are as follows:

20

conc. (ng/mL)	50000	10000	2000	400	80	16
---------------	-------	-------	------	-----	----	----

Cell

% Inhibition

25

Prep. A	MCF-7	100	100	84	65	42	12
	MCF-7AU	100	100	100	96	69	39
	Dox-MCF-7	100	100	100	89	73	45
Prep. B	MCF-7	100	100	91	69	43	15
	MCF-7AU	100	89	65	37	9	3
	Dox-MCF-7	100	86	62	39	7	2

30

35

Free daunorubicin (Prep. B) exhibits higher IC₅₀'s (is less toxic) against both resistant lines. Daunorubicin incorporated in the block copolymers (Prep. A) exhibited lower IC₅₀'s (is more toxic) against both resistant lines.

40

Example 14

5 Groups (6 animals/dose point) of C57B1/6 7-week-old female mice are inoculated i.p. with free or micellar (N = 1.3) daunorubicin obtained as described in Example 3 (Prep. B and Prep. A, respectively), and are observed for 14 days. Drug concentrations are adjusted so that a maximum volume of 0.5 mL is injected in each mouse.

10 The MTD is defined as a dose which leads to no daunorubicin-deaths (any higher dose leads to the daunorubicin-related death of at least 1 animal per group). The experiment is repeated twice. The results are reproducible with less than 10% variation.

15 The MTD of free and micellar (N = 1.3) daunorubicin is determined to be 2.0 and 1.0 $\mu\text{g}/\text{kg}$ body weight, respectively.

Example 15

20 Daunorubicin possesses high specificity with respect to bone marrow, manifesting itself as reversible leukopenia, i.e., a decrease in the number of WBS (leukocyte count) during drug administration. Bone marrow suppression, as well as anticancer effects of daunorubicin, are conditioned by DNA-intercollating activity, whereas the most harmful side effect of anthracyclines, cardiotoxicity, results mainly from membrane interactions with metabolites (which have low anticancer activity and do not produce significant effects on bone marrow). Therefore, the leukocyte count during in vivo administration of MTD daunorubicin allows the assessment of the ratio between specific (DNA-intercollation) activity of the drug and non-specific toxicity.

30 Groups (6 animals/group) of C57B1/6 7-week-old female mice are inoculated i.p. with free or micellar (N = 1.3)

daunorubicin obtained as described in Example 3 (Prep. B and Prep. A, respectively). Drug concentrations (MTD) are adjusted so that a maximum volume of 0.5 mL is injected in each mouse. Blood samples are collected and viable leukocytes are counted as described in Michisch *et al.*, *Proc. Natl. Acad. Sci. USA* 88, 547-551 (1991). The number of WBC after administration of 0.1 mL PBS, 15-16 mln cells/mL, is used as the control. The experiment is repeated twice. The results are reproducible with less than 10% variation.

10 The results obtained are as follows:

Days	0	3	7	10	14
WBS, % of control					
Prep. A	100	20	46.6	86.6	100
Prep. B	100	40	60	93.8	100

20 Example 16

The effects of free and micellar daunorubicin obtained as described in Example 3 (Prep. B and Prep. A, respectively) on leukocyte count are determined three days after administration as described in Example 15.

25 The results obtained are as follows:

Dose of daunorubicin % of MTD	25	50	75	100
WBS, % of control				
Prep. A	85	73	45	21
Prep. B	78	61	36	39

35 The data shown in Examples 14 through 16 indicate that solubilization of daunorubicin in the block copolymer

micelles does not essentially affect the drug's overall toxicity (MTD of 2 mg/kg and 1 mg/kg for free and micellar drug, respectively), whereas an increase in reversible bone marrow suppression is observed which does not markedly influence the animal survivability.

Example 17

Anti-neoplastic activity is determined by evaluation of the cytotoxic activity of plasma of mammals inoculated with the test composition (see de Valeriola et al., *Cancer Chemother. Pharmacol.* 29, 133-140, 1991).

Groups (6 animals/group) of C57B1/6 7-week-old female mice are inoculated i.v. (via the tail vein) with free or micellar (N = 1.3) daunorubicin obtained as described in Example 3 (Prep. B and Prep. A, respectively). Drug concentrations (MTD) are adjusted so that a maximum volume of 0.1 mL is injected in each mouse. The experiment is repeated twice. The results are reproducible with less than 10 % variation.

To obtain plasma samples, blood (10 μ l) is collected from the tail artery one hour after drug administration, diluted 1:10 with sterile RPMI 1640 medium, and centrifuged at 400 g for 15 minutes. The supernatants obtained are diluted as shown in the table with plasma analogously obtained from mice not inoculated with the drug (the plasma of mice not inoculated with the drug does not produce any significant cytotoxic effect on H-69 cells) and mixed with an equal volume of H-69 cell suspension on RPMI 1640 medium supplemented with 10% fetal calf serum. The cells are incubated for two hours at 37°C and 5% CO₂, and then washed three times with RPMI 1640. The pretreated cells are incubated in RPMI 1640 supplemented with 10% fetal calf serum at 37°C and 5% CO₂ for three days, after which cytotoxicity is determined as described in Example 1.

The results obtained are as follows:

Dilution of plasma	1:20	1:200	1:2000	1:20000
	Inhibition, %			
Prep. A	100	58	8	0
Prep. B	42	5	0	0

Thus cytotoxic titers, the dilution at which the plasma of mice inoculated with preparations B or A produced 50% inhibition of H-69 cell growth, of plasma of mice inoculated with preparations B and A with respect to H-69 cells are determined to be 1:228 and 1:48, respectively.

Example 18

The procedure of Example 16 is repeated utilizing SKVLB and SRVO₃ cells. The results are as follows:

a) when MTD of daunorubicin is introduced

5

Plasma Dilution	1:20	1:200	1:2000
-----------------	------	-------	--------

Inhibition, %

10

Prep. A	SKVLB	32	61	18
Prep. B	SKVLB	0	0	0
Prep. A	SKVO ₃	11	0	0
Prep. B	SRVO ₃	9	0	0

15

b) when 10 mg/kg daunorubicin are introduced

20

Plasma Dilution	1:20	1:200	1:2000
-----------------	------	-------	--------

Inhibition, %

25

Prep. A	SKVLB	100	94	69
Prep. B	SKVLB	8	0	0
Prep. A	SKVO ₃	62	31	0
Prep. B	SRVO ₃	22	6	0

30

Example 19

35

A composition suitable for parental administration is prepared by dissolving 400 mg of Pluronic P-85 and 600 mg of Pluronic L-64 in 50 mL of RPMI 1640 at 4°C. The mixture is incubated for 30 minutes at 37°C and then sterilely filtered through a 0.22 μ m filter. This is mixed with a solution of 10 mg of sterile lyophilized daunorubicin powder dissolved in 50 mL of RPMI and incubated for 30 minutes at 37°C.

40

The composition can be stored in the dark at room temperature for 7 days without any essential loss of activity or can be lyophilized and stored for at least 1 year in the dark at room temperature.

Example 20

5 A further composition suitable for parental administration is prepared by dissolving 400 mg of Pluronic P-85 and 600 mg of Pluronic L-64 in 50 mL of PBS at 4°C. The mixture is incubated for 30 minutes at 37°C and then sterilely filtered through a 0.22 μ m filter. This is mixed with a solution of 1 mg of sterile lyophilized daunorubicin powder and 5 mg of glucose dissolved in 50 mL of PBS and the mixture is incubated for 30 minutes at 37°C.

10 The composition can be stored in the dark at room temperature for 7 days without any essential loss of activity or can be lyophilized and stored for at least 1 year in the dark at room temperature.


Example 21

15 A further composition suitable for parental administration is prepared by dissolving 100 mg of sodium ascorbate in a 9% aqueous solution of sodium chloride. To one-half of this solution are added at 4°C 400 mg of Pluronic P-85 and 600 mg of Pluronic L-64. The mixture is incubated for 30
20 minutes at 37°C and then sterilely filtered through a 0.22 μ m filter. Separately 10 mg of sterile lyophilized daunorubicin powder and 50 mg of glucose are dissolved in the remaining sodium ascorbate-sodium chloride solution and the two solutions are mixed and incubated for 30 minutes at
25 37°C.

This composition can be stored for 30 days in the dark at room temperature without any essential loss of activity or can be lyophilized and stored for at least 1 year in the dark at room temperature.

Claims:

1. A pharmaceutical composition comprising an anti-neoplastic agent incorporated into micelles of at least one block copolymer of poly(oxyethylene)-poly(oxypropylene) in which the ratio of poly(oxypropylene) blocks to the poly(oxyethylene) blocks is from 0.25 to 1.5 and the micelles have an average diameter of from 10 to 25 nm.
2. A composition according to claim 1 wherein each block copolymer of poly(oxyethylene)-poly(oxypropylene) is independently represented by the formula:



in which each of x and z, independently of the other, has a value of from 5 to 100 and y has a value of from 20 to 80.

3. A composition according to claim 1 in which the micelles are formed from a plurality of block copolymers of poly(oxyethylene)-poly(oxypropylene) having different hydrophobic/hydrophilic properties.
4. A composition according to claim 3 in which the composite of the block copolymers satisfies the equation:

$$N = \frac{y_1^{*a}}{(x_1 + z_1)^{*}(a + b)} + \frac{y_2^{*b}}{(x_2 + z_2)^{*}(a + b)} * 1.32$$

in which:

y_1 and y_2 are the number of oxypropylene units in the first and second block copolymers, respectively;

x_1 and z_1 are number of oxyethylene units in the first block copolymer;


x_2 and z_2 are number of oxyethylene units in the second block copolymer;

a is the weight proportion of the first block copolymer; and

b is the weight proportion of the second block copolymer

such that the value of N is from 0.25 to 1.5.

5. A composition according to claim 1 wherein the anti-neoplastic agent is an alkaloid, antibiotic, or antifolate.
6. A composition according to claim 5 in which the anti-neoplastic agent is selected from the group consisting of vinblastine, colchicine, demecolone, daunorubicin, doxorubicin, mitomycin C, N-methyl mitomycin C, bleomycin A₂, methotrexate, aminopterin, and dideazatetrahydrofolic acid.
7. A composition according to claim 6 in which the anti-neoplastic agent is daunorubicin.
8. An aqueous dispersion of a quantity of a composition according to claim 1 at least sufficient to provide an effective dose of said anti-neoplastic agent upon parenteral administration.
9. The use of an anti-neoplastic agent incorporated into micelles of at least one block copolymer of poly(oxyethylene)-poly(oxypropylene) in which the ratio of poly(oxypropylene) blocks to the poly(oxyethylene) blocks is from 0.25 to 1.5 and the micelles have an average diameter of from 10 to 25 nm for combatting neoplasms.
10. The use of a rhodamycin antibiotic selected from the group consisting of daunorubicin and doxorubicin leading to a reduction in multidrug resistance which comprises administering to said mammal an effective amount of said rhodamycin antibiotic in micelles comprising at least one block copolymer, each of said block copolymer having of the formula:



in which x and z have values of from 5 to 100 and y has a value of from 20 to 80, such that in the expression:

$$N = \left[\frac{y_1^{*a}}{(x_1 + z_1)^*(a+b)} + \frac{y_2^{*b}}{(x_2 + z_2)^*(a+b)} \right] * 1.32$$

where

x_1 , z_1 , and y_1 correspond to x , z , and y , respectively, of a first block copolymer,

a is the weight proportion of said first block copolymer,

and

if an additional block copolymer is present, x_2 , z_2 , and y_2 correspond to x , z , and y , respectively, of said additional block copolymer, and

b is the weight proportion of said additional block copolymer,

the variable N has a value of from 0.25 to 1.5 for combating the growth of cancer cells in a mammal.

11. The use according to claim 10 wherein said micelles comprise a single block copolymer.
12. The use according to claim 10 wherein N has a value of 1.
13. The use according to claim 10 wherein said rhodamycin antibiotic is daunorubicin.
14. The use according to claim 10 wherein said rhodamycin antibiotic is doxorubicin.