

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 2015293633 B2

(54) Title
Coronavirus

(51) International Patent Classification(s)
C12N 7/04 (2006.01) **A61K 39/215** (2006.01)
A61K 39/00 (2006.01) **C07K 14/165** (2006.01)

(21) Application No: **2015293633** (22) Date of Filing: **2015.07.23**

(87) WIPO No: **WO16/012793**

(30) Priority Data

(31) Number
1413020.7 (32) Date
2014.07.23 (33) Country
GB

(43) Publication Date: **2016.01.28**
(44) Accepted Journal Date: **2021.04.08**

(71) Applicant(s)
The Pirbright Institute

(72) Inventor(s)
Bickerton, Erica;Keep, Sarah;Britton, Paul

(74) Agent / Attorney
Davies Collison Cave Pty Ltd, GPO Box 3876, Sydney, NSW, 2001, AU

(56) Related Art
V. D. MENACHERY ET AL, "Attenuation and Restoration of Severe Acute Respiratory Syndrome Coronavirus Mutant Lacking 2'-O-Methyltransferase Activity", JOURNAL OF VIROLOGY, (2014-01-29), vol. 88, no. 8, pages 4251 - 4264
WO 2005049814 A2
WO 2004092360 A2
PAUL BRITTON ET AL, "Modification of the avian coronavirus infectious bronchitis virus for vaccine development", BIOENGINEERED, (2012-03-01), vol. 3, no. 2, doi:10.4161/bbug.18983, ISSN 2165-5979, pages 114 - 119

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



WIPO | PCT



(10) International Publication Number

WO 2016/012793 A1

(43) International Publication Date

28 January 2016 (28.01.2016)

(51) International Patent Classification:

C12N 7/04 (2006.01) A61K 39/00 (2006.01)
C07K 14/165 (2006.01) A61K 39/215 (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:

PCT/GB2015/052124

(22) International Filing Date:

23 July 2015 (23.07.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

1413020.7 23 July 2014 (23.07.2014) GB

(71) Applicant: THE PIRBRIGHT INSTITUTE [GB/GB];
Ash Road, Pirbright, Woking, Surrey GU24 0NF (GB).

(72) Inventors: BICKERTON, Erica; c/o The Pirbright Institute, Ash Road, Pirbright, Woking, Surrey GU24 0NF (GB). KEEP, Sarah; c/o The Pirbright Institute, Ash Road, Pirbright, Woking, Surrey GU24 0NF (GB). BRITTON, Paul; c/o The Pirbright Institute, Ash Road, Pirbright, Woking, Surrey GU24 0NF (GB).

(74) Agent: ELMHIRST, Elizabeth; 120 Holborn, London EC1N 2DY (GB).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))



WO 2016/012793 A1

(54) Title: CORONAVIRUS

(57) Abstract: The present invention provides a live, attenuated coronavirus comprising a variant replicase gene encoding polyproteins comprising a mutation in one or more of non-structural protein(s) (nsp)-10, nsp-14, nsp-15 or nsp-16. The coronavirus may be used as a vaccine for treating and/or preventing a disease, such as infectious bronchitis, in a subject.

CORONAVIRUS

FIELD OF THE INVENTION

5 The present invention relates to an attenuated coronavirus comprising a variant replicase gene, which causes the virus to have reduced pathogenicity. The present invention also relates to the use of such a coronavirus in a vaccine to prevent and/or treat a disease.

BACKGROUND TO THE INVENTION

10 Avian infectious bronchitis virus (IBV), the aetiological agent of infectious bronchitis (IB), is a highly infectious and contagious pathogen of domestic fowl that replicates primarily in the respiratory tract but also in epithelial cells of the gut, kidney and oviduct. IBV is a member of the Order *Nidovirales*, Family *Coronaviridae*, Subfamily *Coronavirinae* and
15 Genus *Gammacoronavirus*; genetically very similar coronaviruses cause disease in turkeys, guinea fowl and pheasants.

Clinical signs of IB include sneezing, tracheal rales, nasal discharge and wheezing. Meat-type birds have reduced weight gain, whilst egg-laying birds lay fewer eggs and
20 produce poor quality eggs. The respiratory infection predisposes chickens to secondary bacterial infections which can be fatal in chicks. The virus can also cause permanent damage to the oviduct, especially in chicks, leading to reduced egg production and quality; and kidney, sometimes leading to kidney disease which can be fatal.

25 IBV has been reported to be responsible for more economic loss to the poultry industry than any other infectious disease. Although live attenuated vaccines and inactivated vaccines are universally used in the control of IBV, the protection gained by use of vaccination can be lost either due to vaccine breakdown or the introduction of a new IBV serotype that is not related to the vaccine used, posing a risk to the poultry industry.

30 Further, there is a need in the industry to develop vaccines which are suitable for use *in ovo*, in order to improve the efficiency and cost-effectiveness of vaccination programmes. A major challenge associated with *in ovo* vaccination is that the virus must be capable of replicating in the presence of maternally-derived antibodies against the virus, without
35 being pathogenic to the embryo. Current IBV vaccines are derived following multiple passage in embryonated eggs, this results in viruses with reduced pathogenicity for chickens, so that they can be used as live attenuated vaccines. However such viruses almost always show an *increased* virulence to embryos and therefore cannot be used for

in ovo vaccination as they cause reduced hatchability. A 70% reduction in hatchability is seen in some cases.

5 Attenuation following multiple passage in embryonated eggs also suffers from other disadvantages. It is an empirical method, as attenuation of the viruses is random and will differ every time the virus is passaged, so passage of the same virus through a different series of eggs for attenuation purposes will lead to a different set of mutations leading to attenuation. There are also efficacy problems associated with the process: some mutations will affect the replication of the virus and some of the mutations may make the 10 virus too attenuated. Mutations can also occur in the S gene which may also affect immunogenicity so that the desired immune response is affected and the potential vaccine may not protect against the required serotype. In addition there are problems associated with reversion to virulence and stability of vaccines.

15 It is important that new and safer vaccines are developed for the control of IBV. Thus there is a need for IBV vaccines which are not associated with these issues, in particular vaccines which may be used for *in ovo* vaccination.

SUMMARY OF ASPECTS OF THE INVENTION

20 The present inventors have used a reverse genetics approach in order to rationally attenuate IBV. This approach is much more controllable than random attenuation following multiple passages in embryonated eggs because the position of each mutation is known and its effect on the virus, i.e. the reason for attenuation, can be derived.

25 Using their reverse genetics approach, the present inventors have identified various mutations which cause the virus to have reduced levels of pathogenicity. The levels of pathogenicity may be reduced such that when the virus is administered to an embryonated egg, it is capable of replicating without being pathogenic to the embryo. 30 Such viruses may be suitable for *in ovo* vaccination, which is a significant advantage and has improvement over attenuated IBV vaccines produced following multiple passage in embryonated eggs.

35 Thus in a first aspect, the present invention provides a live, attenuated coronavirus comprising a variant replicase gene encoding polyproteins comprising a mutation in one or more of non-structural protein(s) (nsp)-10, nsp-14, nsp-15 or nsp-16.

The variant replicase gene may encode a protein comprising one or more amino acid mutations selected from the list of:

Pro to Leu at position 85 of SEQ ID NO: 6,
Val to Leu at position 393 of SEQ ID NO: 7;
Leu to Ile at position 183 of SEQ ID NO: 8;
Val to Ile at position 209 of SEQ ID NO: 9.

In a particular embodiment of the first aspect the coronavirus is an infectious bronchitis virus (IBV).

In a particular embodiment of the first aspect there is provided a live, attenuated IBV comprising a variant replicase gene encoding polyproteins comprising a mutation in one or more of the non-structural protein(s) (nsp)-10 and nsp-14; wherein the variant replicase gene encodes a protein comprising an amino acid mutation of Pro to Leu at the position corresponding to position 85 of SEQ ID NO: 6, and/or wherein the variant replicase gene encodes a protein comprising an amino acid mutation of Val to Leu at the position corresponding to position 393 of SEQ ID NO: 7.

The IBV may further comprise a mutation in one or more of nsp-15 corresponding to SEQ ID NO: 8 and nsp-16 corresponding to SEQ ID NO: 9.

The replicase gene may encode a protein comprising the amino acid mutation Pro to Leu at position 85 of SEQ ID NO: 6.

The replicase gene may encode a protein comprising the amino acid mutations Val to Leu at position 393 of SEQ ID NO: 7; Leu to Ile at position 183 of SEQ ID NO: 8; and Val to Ile at position 209 of SEQ ID NO: 9.

The replicase gene may encode a protein comprising the amino acid mutations Pro to Leu at position 85 of SEQ ID NO: 6; Val to Leu at position 393 of SEQ ID NO: 7; Leu to Ile at position 183 of SEQ ID NO: 8; and Val to Ile at position 209 of SEQ ID NO: 9.

The replicase gene may comprise one or more nucleotide substitutions selected from the list of:

2015293633 10 Mar 2021

3a

C to T at nucleotide position 12137;
G to C at nucleotide position 18114;
T to A at nucleotide position 19047; and
G to A at nucleotide position 20139;
compared to the sequence shown as SEQ ID NO: 1.

The coronavirus may be an infectious bronchitis virus (IBV).

The coronavirus may be IBV M41.

The coronavirus may comprise an S protein at least part of which is from an IBV serotype other than M41.

For example, the S1 subunit or the entire S protein may be from an IBV serotype other than M41.

5 The coronavirus according to the first aspect of the invention has reduced pathogenicity compared to a coronavirus expressing a corresponding wild-type replicase, such that when the virus is administered to an embryonated egg, it is capable of replicating without being pathogenic to the embryo.

In a second aspect, the present invention provides a variant replicase gene as defined in connection with the first aspect of the invention.

10 In a third aspect, the present invention provides a protein encoded by a variant coronavirus replicase gene according to the second aspect of the invention.

In a fourth aspect, the present invention provides a plasmid comprising a replicase gene according to the second aspect of the invention.

15 In a fifth aspect, the present invention provides a method for making the coronavirus according to the first aspect of the invention which comprises the following steps:

(i) transfecting a plasmid according to the fourth aspect of the invention into a host cell;

20 (ii) infecting the host cell with a recombining virus comprising the genome of a coronavirus strain with a replicase gene;

(iii) allowing homologous recombination to occur between the replicase gene sequences in the plasmid and the corresponding sequences in the recombining virus genome to produce a modified replicase gene; and

25 (iv) selecting for recombining virus comprising the modified replicase gene.

The recombining virus may be a vaccinia virus.

The method may also include the step:

30 (v) recovering recombinant coronavirus comprising the modified replicase gene from the DNA from the recombining virus from step (iv).

In a sixth aspect, the present invention provides a cell capable of producing a coronavirus according to the first aspect of the invention.

35 In a seventh aspect, the present invention provides a vaccine comprising a coronavirus according to the first aspect of the invention and a pharmaceutically acceptable carrier.

In an eighth aspect, the present invention provides a method for treating and/or preventing a disease in a subject which comprises the step of administering a vaccine according to the seventh aspect of the invention to the subject.

5

Further aspects of the invention provide:

- the vaccine according to the seventh aspect of the invention for use in treating and/or preventing a disease in a subject.
- use of a coronavirus according to the first aspect of the invention in the manufacture of a vaccine for treating and/or preventing a disease in a subject.

10 The disease may be infectious bronchitis (IB).

15 The method of administration of the vaccine may be selected from the group consisting of; eye drop administration, intranasal administration, drinking water administration, post-hatch injection and *in ovo* injection.

Vaccination may be by *in ovo* vaccination.

20 The present invention also provides a method for producing a vaccine according to the seventh aspect of the invention, which comprises the step of infecting a cell according to the sixth aspect of the invention with a coronavirus according to the first aspect of the invention.

25 DESCRIPTION OF THE FIGURES

Figure 1 - Growth kinetics of M41-R-6 and M41-R-12 compared to M41-CK (M41 EP4) on CK cells

30 **Figure 2** – Clinical signs, snicking and wheezing, associated with M41-R-6 and M41-R-12 compared to M41-CK (M41 EP4) and Beau-R (Bars show mock, Beau-R, M41-R 6, M41-R 12, M41-CK EP4 from left to right of each timepoint).

35 **Figure 3** - Ciliary activity of the viruses in tracheal rings isolated from tracheas taken from infected chicks. 100% ciliary activity indicates no effect by the virus; apathogenic, 0% activity indicates complete loss of ciliary activity, complete ciliostasis, indicating the virus is pathogenic (Bars show mock, Beau-R, M41-R 6, M41-R 12, M41-CK EP4 from left to right of each timepoint).

5 **Figure 4** - Clinical signs, snicking, associated with M41R-nsp10rep and M41R-nsp14,15,16rep compared to M41-R-12 and M41-CK (M41 EP5) (Bars show mock, M41-R12; M41R-nsp10rep; M41R-nsp14,15,16rep and M41-CK EP5 from left to right of each timepoint).

10 **Figure 5** - Ciliary activity of M41R-nsp10rep and M41R-nsp14,15,16rep compared to M41-R-12 and M41-CK in tracheal rings isolated from tracheas taken from infected chicks (Bars show mock; M41-R12; M41R-nsp10rep; M41R-nsp14,15,16rep and M41-CK EP5 from left to right of each timepoint).

15 **Figure 6** - Clinical signs, snicking, associated with M41R-nsp10, 15rep, M41R-nsp10, 14, 15rep, M41R-nsp10, 14, 16rep, M41R-nsp10, 15, 16rep and M41-K compared to M41-CK (Bars show mock, M41R-nsp10,15rep1; M41R-nsp10,14,16rep4; M41R-nsp10,15,16rep8; M41R-nsp10,14,15rep10; M41-K6 and M41-CK EP4 from left to right of each timepoint).

20 **Figure 7** - Clinical signs, wheezing, associated with M41R-nsp10, 15rep, M41R-nsp10, 14, 15rep, M41R-nsp10, 14, 16rep, M41R-nsp10, 15, 16rep and M41-K compared to M41-CK (Bars show mock, M41R-nsp10,15rep1; M41R-nsp10,14,16rep4; M41R-nsp10,15,16rep8; M41R-nsp10,14,15rep10; M41-K6 and M41-CK EP4 from left to right of each timepoint).

25 **Figure 8** - Ciliary activity of M41R-nsp10, 15rep, M41R-nsp10, 14, 15rep, M41R-nsp10, 14, 16rep, M41R-nsp10, 15, 16rep and M41-K compared to M41-CK in tracheal rings isolated from tracheas taken from infected chicks (Bars show mock, M41R-nsp10,15rep1; M41R-nsp10,14,16rep4; M41R-nsp10,15,16rep8; M41R-nsp10,14,15rep10; M41-K6 and M41-CK EP4 from left to right of each timepoint).

30 **Figure 9** - Growth kinetics of rIBVs compared to M41-CK on CK cells. Fig 9A shows the results for M41-R and M41-K. Fig 9B shows the results for M41-nsp10 rep; M41R-nsp14, 15, 16 rep; M41R-nsp10, 15 rep; M41R-nsp10, 15, 16 rep; M41R-nsp10, 14, 15 rep; and M41R-nsp10, 14, 16.

35 **Figure 10** – Position of amino acid mutations in mutated nsp10, nsp14, nsp15 and nsp16 sequences.

Figure 11 – A) Snicking; B) Respiratory symptoms (wheezing and rales combined) and C) Ciliary activity of rIBV M41R-nsp 10,14 rep and rIBV M41R-nsp 10,16 rep compared to M41-CK (Bars show mock, M41R-nsp10,14rep; M41R-nsp10,16rep and M41-K from left to right of each timepoint).

5

DETAILED DESCRIPTION

The present invention provides a coronavirus comprising a variant replicase gene which, when expressed in the coronavirus, causes the virus to have reduced pathogenicity 10 compared to a corresponding coronavirus which comprises the wild-type replicase gene.

CORONAVIRUS

15 *Gammacoronavirus* is a genus of animal virus belonging to the family *Coronaviridae*. Coronaviruses are enveloped viruses with a positive-sense single-stranded RNA genome and a helical symmetry.

The genomic size of coronaviruses ranges from approximately 27 to 32 kilobases, which is the longest size for any known RNA virus.

20 Coronaviruses primarily infect the upper respiratory or gastrointestinal tract of mammals and birds. Five to six different currently known strains of coronaviruses infect humans. The most publicized human coronavirus, SARS-CoV which causes severe acute respiratory syndrome (SARS), has a unique pathogenesis because it causes both upper 25 and lower respiratory tract infections and can also cause gastroenteritis. Middle East respiratory syndrome coronavirus (MERS-CoV) also causes a lower respiratory tract infection in humans. Coronaviruses are believed to cause a significant percentage of all common colds in human adults.

30 Coronaviruses also cause a range of diseases in livestock animals and domesticated pets, some of which can be serious and are a threat to the farming industry. Economically significant coronaviruses of livestock animals include infectious bronchitis virus (IBV) which mainly causes respiratory disease in chickens and seriously affects the poultry industry worldwide; porcine coronavirus (transmissible gastroenteritis, TGE) and 35 bovine coronavirus, which both result in diarrhoea in young animals. Feline coronavirus has two forms, feline enteric coronavirus is a pathogen of minor clinical significance, but spontaneous mutation of this virus can result in feline infectious peritonitis (FIP), a disease associated with high mortality.

There are also two types of canine coronavirus (CCoV), one that causes mild gastrointestinal disease and one that has been found to cause respiratory disease. Mouse hepatitis virus (MHV) is a coronavirus that causes an epidemic murine illness with 5 high mortality, especially among colonies of laboratory mice.

Coronaviruses are divided into four groups, as shown below:

Alpha

10

- Canine coronavirus (CCoV)
- Feline coronavirus (FeCoV)
- Human coronavirus 229E (HCoV-229E)
- Porcine epidemic diarrhoea virus (PEDV)
- Transmissible gastroenteritis virus (TGEV)

15

- Human Coronavirus NL63 (NL or New Haven)

Beta

20

- Bovine coronavirus (BCoV)
- Canine respiratory coronavirus (CRCoV) - Common in SE Asia and Micronesia
- Human coronavirus OC43 (HCoV-OC43)
- Mouse hepatitis virus (MHV)
- Porcine haemagglutinating encephalomyelitis virus (HEV)
- Rat coronavirus (RCV). Rat Coronavirus is quite prevalent in Eastern 25 Australia where, as of March/April 2008, it has been found among native and feral rodent colonies.
- (No common name as of yet) (HCoV-HKU1)
- Severe acute respiratory syndrome coronavirus (SARS-CoV)
- Middle East respiratory syndrome coronavirus (MERS-CoV)

30

Gamma

- Infectious bronchitis virus (IBV)
- Turkey coronavirus (Bluecomb disease virus)
- Pheasant coronavirus

35

- Guinea fowl coronavirus

Delta

- Bulbul coronavirus (BuCoV)
- Thrush coronavirus (ThCoV)
- Munia coronavirus (MuCoV)
- Porcine coronavirus (PorCov) HKU15

5

The variant replicase gene of the coronavirus of the present invention may be derived from an alphacoronavirus such as TGEV; a betacoronavirus such as MHV; or a gammacoronavirus such as IBV.

10

As used herein the term "derived from" means that the replicase gene comprises substantially the same nucleotide sequence as the wild-type replicase gene of the relevant coronavirus. For example, the variant replicase gene of the present invention may have up to 80%, 85%, 90%, 95%, 98% or 99% identity with the wild type replicase sequence. The variant coronavirus replicase gene encodes a protein comprising a mutation in one or more of non-structural protein (nsp)-10, nsp-14, nsp-15 or nsp-16 when compared to the wild-type sequence of the non-structural protein.

IBV

20

Avian infectious bronchitis (IB) is an acute and highly contagious respiratory disease of chickens which causes significant economic losses. The disease is characterized by respiratory signs including gasping, coughing, sneezing, tracheal rales, and nasal discharge. In young chickens, severe respiratory distress may occur. In layers, respiratory distress, nephritis, decrease in egg production, and loss of internal egg quality and egg shell quality are common.

In broilers, coughing and rattling are common clinical signs, rapidly spreading in all the birds of the premises. Morbidity is 100% in non-vaccinated flocks. Mortality varies depending on age, virus strain, and secondary infections but may be up to 60% in non-vaccinated flocks.

The first IBV serotype to be identified was Massachusetts, but in the United States several serotypes, including Arkansas and Delaware, are currently circulating, in addition to the originally identified Massachusetts type.

The IBV strain Beaudette was derived following at least 150 passages in chick embryos. IBV Beaudette is no longer pathogenic for hatched chickens but rapidly kills embryos.

H120 is a commercial live attenuated IBV Massachusetts serotype vaccine strain, attenuated by approximately 120 passages in embryonated chicken eggs. H52 is another Massachusetts vaccine, and represents an earlier and slightly more pathogenic passage virus (passage 52) during the development of H120. Vaccines based on H120 are commonly used.

IB QX is a virulent field isolate of IBV. It is sometimes known as "Chinese QX" as it was originally isolated following outbreaks of disease in the Qingdao region in China in the mid 1990s. Since that time the virus has crept towards Europe. From 2004, severe egg production issues have been identified with a very similar virus in parts of Western Europe, predominantly in the Netherlands, but also reported from Germany, France, Belgium, Denmark and in the UK.

The virus isolated from the Dutch cases was identified by the Dutch Research Institute at Deventer as a new strain that they called D388. The Chinese connection came from further tests which showed that the virus was 99% similar to the Chinese QX viruses. A live attenuated QX-like IBV vaccine strain has now been developed.

IBV is an enveloped virus that replicates in the cell cytoplasm and contains a non-segmented, single-stranded, positive sense RNA genome. IBV has a 27.6 kb RNA genome and like all coronaviruses contains the four structural proteins; spike glycoprotein (S), small membrane protein (E), integral membrane protein (M) and nucleocapsid protein (N) which interacts with the genomic RNA.

The genome is organised in the following manner: 5'UTR - polymerase (replicase) gene – structural protein genes (S-E-M-N) – UTR 3'; where the UTR are untranslated regions (each ~ 500 nucleotides in IBV).

The lipid envelope contains three membrane proteins: S, M and E. The IBV S protein is a type I glycoprotein which oligomerizes in the endoplasmic reticulum and is assembled into homotrimer inserted in the virion membrane via the transmembrane domain and is associated through non-covalent interactions with the M protein. Following incorporation into coronavirus particles, the S protein is responsible for binding to the target cell receptor and fusion of the viral and cellular membranes. The S glycoprotein consists of

four domains: a signal sequence that is cleaved during synthesis; the ectodomain, which is present on the outside of the virion particle; the transmembrane region responsible for anchoring the S protein into the lipid bilayer of the virion particle; and the cytoplasmic tail.

- 5 All coronaviruses also encode a set of accessory protein genes of unknown function that are not required for replication *in vitro*, but may play a role in pathogenesis. IBV encodes two accessory genes, genes 3 and 5, which both express two accessory proteins 3a, 3b and 5a, 5b, respectively.
- 10 The variant replicase gene of the coronavirus of the present invention may be derived from an IBV. For example the IBV may be IBV Beaudette, H120, H52, IB QX, D388 or M41.

15 The IBV may be IBV M41. M41 is a prototypic Massachusetts serotype that was isolated in the USA in 1941. It is an isolate used in many labs throughout the world as a pathogenic lab stain and can be obtained from ATCC (VR-21TM). Attenuated variants are also used by several vaccine producers as IBV vaccines against Massachusetts serotypes causing problems in the field. The present inventors chose to use this strain as they had worked for many years on this virus, and because the sequence of the complete 20 virus genome is available. The M41 isolate, M41-CK, used by the present inventors was adapted to grow in primary chick kidney (CK) cells and was therefore deemed amenable for recovery as an infectious virus from a cDNA of the complete genome. It is representative of a pathogenic IBV and therefore can be analysed for mutations that cause either loss or reduction in pathogenicity.

25

The genome sequence of IBV M41-CK is provided as SEQ ID NO: 1.

SEQ ID NO: 1 IBV M41-CK Sequence

30 AACTAAGATAGATATTAATATATATCTATCACACTAGCCTTGCCTAGATTTCCAACCTTA
ACAAAACGGACTTAAATAACCTACAGCTGGCCTCATAGGTGTTCCATTGCAGTGCACCTT
AGTGCCTGGATGGCACCTGGCCACCTGTCAGGTTTTGTTATTAAAATCTTATTGTTGC
TGGTATCACTGCTTGTGTTGCCGTGTCACTTATACATCCGTTGCTGGGCTACCTAG
TATCCAGCGTCCCTACGGGCGCCGTGGCTGGTCAGTGCGAAGAACCTCTGGTTCATCTA
GCGGTAGGCGGGTGTGGAAGTAGCACCTCAGACGTACCGGTTCTGTTGTGAAATAC
GGGGTCACCTCCCCCACATACCTCTAACGGCTTTGAGCCTAGCGTGGCTACGTTCT
35 CGCATAAGGTGGCTATACGACGTTGTAGGGGGTAGTGCCAAACAACCCCTGAGGTGAC
AGGTTCTGGTGGTGTAGTGAGCAGACATAACATAGACAGTGACAACATGGCTCAAGC
CTAAAACAGGGAGTATCTCCCAAACTAAGGGATGTCATTCTGTATCAAAGACATTCT
GAACAACTTGTGACGCTTGTGTTCTATACGTCACACAACCCCTAAGGATTACGCTGAT
40 GCTTTGCAGTTAGGCAGAAGTTGATGTAATCTGCAGACTGGAAACAGTTCAAATT
GAAACTGTGTGGCTCTTCCCTCTGAAGGGAGTTGACAAAATAACACCTGGCGTCCCA
GCAAAAGTCTAAAAGCCACTTCTAAGTTGGCAGATTAGAAGACATCTTGGTGTCTCT

CCCTTGCAAGAAAATATCGTGAACCTTTGAAGACAGCAGGCCAGTGGTCTCTTACTGTA
 GAAACACTGGATGCTCGTCACAAACTCTGATGAAATTGGCTACTGAAATACTT
 TGGCTTCAGGTGGCAGCAAAATCCAAGTTCGGCTATGGCGATGCGCAGGCTTGG
 GAAGTAACTGCAAAAGTCATGGATGCTTGGCTCAAATATGAGTGCTCTTCCAGATT
 5 TTTAAACAAACAAATAGTCAGAATTTCAGGCTGGCTATTTGAGAATGTGAGT
 GAATTACCACAGCGTATTGCAGCACTTAAGATGGCTTGCTAAGTGTGCCAAGTCCATT
 ACTGTTGTGGTTATGGAGAGGACTCTAGTTGTTAGAGAGTTCGCAGGAACCTGCTTGCA
 AGCATTAAATGGTGCCTTGCAAAATTCTTGAAGAACCTCCAAATGGTTATGGGTGCT
 AAAATTTCACTACACTGCCTCTTGGGAGGCTGCAGTGAAAATGTGGATAACATA
 10 CCAAATGCACCGAGAGGCATAAAGGGTTGAAGTCGTTGGTAATGCCAAGGTACACAA
 GTTGGTGTGGTGCATGCGAAATGACTTAACACTGCTGACCAAAAGCTGAAATTCCCT
 GTGGAGTCAGAAGGTTGGTCTGCAATTGGGTGACATCTTGCTATGTCTTAAGAGT
 GGTGATGCTTTACGGGCACCTCTTCAGGAAATTTCGATTGCATGATGTGCATTGT
 TGTGAGCGTGTGCTGTCTTGATGGTGTAAACACGGAGATAATGATGGACTTATT
 15 CTTGCAGCAATCTACTCTTCTTAGTGTGGCAGAACATTGGCAGGCATTAAGGGGT
 GAACCATTAAAGTTCTGGGTCAATAATTGTGTATGCCAAGGGATGCAGCAGTTCTTT
 ACATTAGCGAAGGCTGCTACTATTGCAGATGTTGAAGCTGTTCAATCAGCGCTGTG
 AAAGTAGAAGATGTTGGTCTTCACTTACTGAAAGTCTTGAATTCTGGAGGCTTGCA
 TATGGAAAAGTGCATACTCGAAGAATTGTTAGACTTGGTTGAGGCTCAAATG
 20 GCGATTGTGATTAGCGACAGTGCTGGAGAGGGCATTGGCATCTGTTGCAAGTC
 ATCTATAAAGTAGGTGGTCTTTACTAAAGTTGACTTTGTGAAAATATTGGAAA
 GGTTTTGTGCACAGTTGAAAAGAGCTAACGCTCATTGTCACTGAAACCTCTGTGTTTG
 AAAGGAGTTGCACAGCATTGTTCAACTATTGCTGGATGCAATACAGTTATGTATAAA
 AGTTTAAGAAGTGTGCACTTGGTAGAATCCATGGAGACTTGCTCTGGAAAGGAGGT
 25 GTGCACAAATTATTCAAGAGGGCGATGAAATTGGTTGACGCCATTGATAGTATTGAT
 GTTGAAGATCTGGGTGTTCAAGAAAATTGATTGATTGATGTTGTGATAATGTG
 ACACCTCCAGAGAACCAACCGGTCAATGGTCAAATCGAGGATGACGGAAAGAACTAC
 ATGTTCTCCGCTTCAAAAGGATGAGAACATTATTACACCAATGTCACAGCTGGT
 GCTATTAAATGTGGTTGCAAAGCAGGCCGTAAGACTGTCACCTTGGAGAAACTACTGTG
 30 CAAGAAAATACCACCACTGATGTTGTTATTAAAGTTAGCATTGAGTGTGGTGAA
 CCATGGAATACAATCTCAAAAGGTTATAAGGAGCCATTGAAGTAGAGACAGACCTC
 ACAGTTGAACAATTGCTCTGTGGTCTATGAGAAAATGTGTGATGATCTCAAGCTGTT
 CCGGAGGCTCCAGAACCAACCACTTGAGAACATGTCACACTGTTGATAAGAATGGTAAA
 GATTGGATTGCATAAAATCATGCCATCTGATCTCGTATTGAGAGCGATGATGAC
 35 ATCGAGGAAGAAGATGCAGAAGAATGTGACACGGATTAGGTGATGCTGAGGAGTGTGAC
 ACTAATTCAAGAATGTGAGAACAGATGAGGAACTAAAGTGTGGCTTTATACAAGAC
 CCGGCAAGTAACAAATATCCTCGCCTTGATGATGATTAGCGCTACAATGGATGT
 ATTGTTCATAAAGGACGCTCTCGATGTTGAATTACCATCTGGTGAAGAACCTTGT
 GTCAATAACTGCTTGAAGGGCTGTTAAAGCTTCCGAGAAAGTATTGATGTTCTA
 40 GGTGACTGGGTGAGGCTGTTGATGCGCAAGAACATTGTGTCACAAAGAACACTCGG
 GTCATATCTGAGAAATCAGTTGAGGTTTACTGGTAGTTGATGCAATGGCTGAACAA
 GCTATTGTTGAGAGCAGGAAATAGTACCTGTTGAACAAAGTCAGGATGTTGTT
 TTTACACCTGCAGACCTAGAAGTTGTTAAAGAACAGCAGAACAGGTTGATGAGTTATT
 CTCATTCTGCTGTCCTAAAGAACAGTTGTCAGGAGAACAGGAGGCCACAGGTT
 45 GAGCAAGAGCCTACCCAGTGTGTTAAAGCACAAACGTGAGAACAGGCTAAAAGTTCAA
 GTTAAACCAGCTACATGTGAAAACCCAAATTGGAGTACAAACATGTGTGGTGAT
 TTGGCTGTTGTAATTGCCAACAGCATTGGATGAGTTAAAGAGTTCTGCATTGTAACGCT
 GCAAATGAGCACATGTCGATGGTGGTGCAGGCAATTGCAAGACTTTGTGGA
 CCGGACTTTGTTGAATTGCGCGACTATGTTAAAGAACATGGCCACAGCAAAACTT
 50 GTCACACCTTCATTGTTAAAGCATTCAATGTGATAATGTTGAGGACCTGCCAT
 GGAGACAGCAACTTGCCTGAGAACAGCTTGGTGTGCTTACAAGAGTGTCTTGAGGTGGA
 GTGGTTAACTATGTTGTCAGTTCTCATCAGGGATTGGTGTAGATTAAAATA
 TCAATAGATGCTATGCGCAAGCTTAAAGGTTGCCATACCGCTCTTTATTTCT
 CTGAGTCAGAACACATCGATTATTCGATGCAACTGTAAGCAGAACAGAACATTATCTT
 55 ACGGAGGATGGTGTAAATACCGCTCTGGTTAAAACCTGGTATTCTTGGTCAA
 TTTGGACAGGTTTGCAAGAAATAAGGAGTCTTCCGGCTGATGATGTTGAGGATAAA
 GAAATCCTCTTATACCCACAACGACAAGACTATTCTGAATATTGTTAGATGCG

CAAAAGTATGTAACATATTGCAAACGCTTGCAGAATGGGATGTCATATAGAGAC
 AATTTGTTATATTAGAGTGGCGTGACGAAATTGCTGGATTAGTCAGCAATAGTC
 CTTCAAGCTGCTAAAATTAGATTAAAGGTTCTGCAGAAGCATGGCTAAACTGTTG
 GGGGAGATCCTACAGACTTGGTGCCTGGTGTATGCAAGTGCAATGCTAAAGTAGGT
 5 GATTTTCAGATGCTAATTGGCTTGGCAATTAGCAGAACATTGACCGAGATTAC
 ACAATGCACTCTTAAGAAGTGTGTCGTGCAATTGTTAGGTTAAGAGTTATGAACTT
 AGGGGCTTGAAGCCTGTATTCAAGCCAGTCGAGCACCTAATCTACATTAAACG
 CAATATTCAAATTGCCAACCTGTGGTCAAGTAGTACGGATGAAGTAATAGAACGTTCA
 TTACCGTACTTATTGCTTTGCTACTGATGGTCCTGCTACAGTTGATTGATGAAAAT
 10 GCTGTAGGGACTGTTGCTTCAATTGGCTACTAATAGTGGCATTGTTATACACAAGCC
 GATGGTAAGGCTTTGACAATTGCTAAGGATAGAAAATTGGAAGGAAGTCCCTAC
 ATTACAGCAATGTATACACGTTCTTAGGAGTAAAAGAGATGTTCAACCTGCTACTAGTTCT
 CATACTAAGGGTAAAGCTAAAGTAGTAAAAGAAGATGTTCAACCTGCTACTAGTTCT
 AAAGCCAGTTGACCGATCTTACTGACTTGAACAGTGGTATGATAGAACATCTATGAG
 15 AGTCTTAAAGTCAGGAGACACTGATAATCTGATGAATATGTCATTACGACAAAG
 GAAGATTCTAAGTTGCCACTGACACTTAAAGTTAGAGGTATCAAATCAGTTGACTTT
 AGGTCTAAGGATGGTTTACTTATAAGTTAACACCTGATACTGATGAAAATTCAAAACA
 CCAGTCTACTACCCAGTCTGGATTCTATTAGTCTTAGGGCAATATGGGTTGAAGGCAGT
 GCTAATTGTTGTTGGCATCCAAATTATTATAGTAAGTCCTCCGAATTCCACGTTT
 20 TGGGAAATGCCAGAGCTTGTAAAATGGTTATAAAATTGATGGTAACTATGGGC
 CTTGGCGTGCAGAACACCTTAATAAAACCTAATTGGAGAGAATTAAACATTGCTAAG
 AAAGCTATTGTTGGATCTAGTGTGTTACTACGCAGTGTGGTAAAATACTAGTTAAAGCA
 GCTACATACGTTGCCATAAAGTAGGTGATGGTGTAGTCGCAATATTACAGATAGAATT
 AAGGGTCTTGTGGATTACACGTGGCATTGAAAAGAAAATGTCCTACAATTCTA
 25 AAGACACTTGTGTTCTTTATTCTTAAAGGCTAGTGCTAAGAGTTAGTTCT
 AGCTATAAGATTGTTATGTAAGGTGGTTGCTACCTTACTTATAGTGTGGTTATA
 TACACAAGTAATCCAGTAGTGTGTTACTGGAATACGTGTCAGACTCCTATTGAAAGGT
 TCTTATGTTGCTTATAATGACTACGTAAAGATTCTTGTATGTTACGCTATTG
 GCAGGTGATTACTTGTGTTGTTACATGATAGAGATTCACTCATCTGTACAAA
 30 CATGCTTATAGCGTAGAACAAATTATAAGGATGCAGCTCTGGCATTAACCTTAATTGG
 AATTGGCTTATTGGTCTTCTAATATTATTGTTAAGCCAGTGGCAGGTTGTTATT
 ATTGTTATTGTTAAGTATTGGTATTGAGTTCAACTGTGTTGCAAACACTGGTGTAGGT
 TTTCTAGATTGGTTGTAACAGTTTACCCATTAAATTATGGGAGCAGGGATT
 TATTCTGGCTCTTACAAGATATACGTACAAGTCATCATATTGTAAGGAT
 35 GAAACATGTGAAGTGTGCAAGAGAGTTGCACGCAGCAACAGGAAGAGGTTAGCGTTGTA
 GTTGGTGGACGCAAGCAAATAGTCATGTTACACTAATTCTGGCTATAACTTTGTAAG
 AGACATAATTGGTATTGTAAGAATTGTGATGATTGGTCACCAAAATACATTATGTC
 CCTGAAGTTGCTGGCGAGCTTCTGAAAGCTTAAGGCCATGTTAAACCTACAGCATAT
 GCTTACACGTTGTTGAGGATAATGCATCTCTGCTGTTAAGTGTGTTAGTGTACAGAT
 40 GCTGCAATTCTGGTAAGGATAATGCATCTCTGCTGTTAAGTGTGTTAGTGTACAGAT
 TTTTAAAGAAAGCTGTTCTTAAGGAGGATTGAAATGTAACAAATATCTAATGAT
 GGTTTATAGTGTGTAATACACAGACTGCCATGCACTAGAGGAAGCAAAGAATGCAGCC
 GTCTATTATGCGCAATATCTGTTGTAAGCCAATACTTATGTTGACCAGGCACCTTATGAG
 CAATTAAATAGTAGAGCCTGTTCAAGAGTGTATAGATAAAAGTGTGAGCATTTGTC
 45 AATATAATATCTGTAAGACTGCAGCTTAAATTATAAGGCAGGCACACTCGTGATGCT
 CTGCTTTCTATTACTAAAGACGAAGAAGCGTAGATATGGCTATCTCTGCCACAATCAT
 GAAGTGGAAATACACTGGTGACGGTTTACTAATGTGATACCGTCATATGGTATGGACACT
 GATAAGTTGACACCTCGTGATAGAGGGTTTGATAATGCAGATGCTCTATTGCTAAT
 TTAAGAGTCAAAATGCTCCTCCGGTAGTATGGAAGTTCTGATCTATTAAATTGTCT
 50 GACAGTTGCCTTAAATTATACTCAGTACTGTCAGTCAGTCAGGAGGTGTTCTTATA
 ACAAAAGTCTGGTCTAAACAAAGTTATTCTGTCATACCCAGAAACTGTTGAGGAA
 AAGGCAGGTGGTGTATTAAACACTTTAAATGGTTATGAGTTGTTAAATGGCTT
 TTTGTTCTTATATACTTTACAGCATGTTGTTGGTTACTACTATATGGAGATGAAT
 AAAAGTTGTTACCCATGTATGATGAAACTCCACACTGCACTGTTGAAGGGTTCAAA
 55 GTTATAGACAAAGGTGTATTAGAGAGATTGTCAGAACAGATAATTGTTCTCTAATAAG
 TTTGTTAATTGACGCCCTTGGGTAATCATGAAAATAAAACTGTCCAATT
 GTTACAGTTGTTAGATGGTGACGGGACAGTAGCTGTTGGTGTCCCTGGTTGTATCA

TGGGTTATGGATGGTGTATGTTGTGCATATGACACAGACTGATCGTAGACCTGGTAC
 ATTCCCTACCTGGTTAATAGAGAAATTGGTGGTTACACTCAGGATTCAATTATCACTGAG
 GGTAGTTTATACATCTATAGCATTATTTCTGCTAGATGTTATATTAAACAGCCAGC
 AATACACCTCAATTGTATTGTTAATGGCGACAATGATGCACCTGGAGCCTACCATT
 5 GGTAGTATTATTCTCATAGAGTATACTCCAACCTAATGGTGTAGGCTTATAGTCCA
 CAACAAATACTGCATACACCCATACATAGTGAAGTTGTTCAGACAGCTATTGTAGAGGT
 AGTGTATGTGAGTATACTAAACCAGGTTACTGTGTCACTAGACTCCAATGGGTTTG
 TTTAATGATGAATACTAGTAAACCTGGCGTTCTGTGGTTACTGTTAGAGAACTT
 ATGTTAATATGGTTAGTACATTCTTACTGGTGTCAACCCATAATATTATTCAGCTA
 10 GCAACTATGTTTAATACTAGTGTATTGTGTTAATTGCAATGGTTAAAGTT
 CAAGGTGTTTAAAGCTTATGCGACCATTGTGTTACAATAATGTTAGTTGGGTTATT
 AATGCATTGTTTGTGTACATAGTTATAATAGTGTAGCTGTATATTATTAGTA
 CTCTATTGCTATGCATCATTGGTTACAAGTCGAATACTGCTATAATAATGCATTGTTGG
 CTTGTTTACCTTGGTTAATAGTACCCACATGGTGGCTTGTGCTATCTGGGATT
 15 ATTCTTATATGTACACACCGTGGTTCTGGTGTACGGTACTACTAAAAACTCGT
 AAGTTGTATGATGGCAACGAGTTGTTGTAATTATGACCTGCTGCGAAGAGCACTTT
 GTTATTGTTGACTGAATTGTTAAGCTTACGAATGAGATAGGTGATAATTGAAAGCC
 TATCTTCTCGTATGCTAGACTAAACTATTAGGCACTGGTAGTGAGCAAGATTAC
 TTGCAAGCTTGTGCGATGGTAGCTTATGCTTGGACCAATATAGAAATAGTGGTGT
 20 GAGGTTGTTATACCCCACCGCGTTACTCTATTGGTGTAGTAGACTACACGCTGGTTT
 AAAAAGTCTCTCTAGTAGTGCTGTTGAGAAGTGCAATTGTTAGTGCTCTTATAGA
 GGCAATAATCTTAAATGGACTGTGGCTGGGTTAGCTATTACTGCCACGCCATGTGTTA
 GGTAAGTTAGTGGTAGGCAAGTGGGTTAGCTAAACCTGCTAATAATCATGAGTT
 GAAGTTGTAACTCAAATGGTGTACTTGAATGTTGTCAGCAGCGGCTTAAAGGAGCA
 25 GTTTAATTACAAACTGCAGTTGCCAATGCTGAAACTCTAAGTATAAGTTGTTAAA
 GCTAATTGTTGATAGTTCACTATAGCTTCTTATGGTGTACAGTTATAGGACTT
 TACCCGTCACTATGCGTTCTAATGGTACTATTAGAGCATCTTCTAGCAGGAGCCTGT
 GGCTCAGTTGGTTTAATATAGAAAAGGGTAGTTAATTCTTATATGCACCATCTT
 GAGTTACCTAATGCATTACACACTGGAACGTACCTAATGGTGAGTTATGGTGGTTAT
 30 GTAGATGAAGAGGTTGCGCAAAGAGTGCCACCAAGATAATCTAGTTACTAACATATTGTA
 GCATGGCTCATGGCAATTATTAGTGTAAAGAAAAGTAGTTTCAACACCTAAATGG
 TTGGAGAGTACTACTGTTCTATTGAAGATTACAATAGGTGGCTAGTGATAATGGTT
 ACTCCATTTCACACTGCTATTACTAAATTAGTGCTATAACTGGGTTGATGTT
 TGTAAACTCCTCGCACTATTATGGTAAAAGTGCTCAATGGGTTAGTGATCCCATT
 35 GGACAATATAATTGAAAGACGAATTGACACCAGAACTGTATTAACTCAAGTTGGTGGT
 GTTAGGTTACAGTCTTGTAAAGAAAAGCTACATCTTGGTTTGAGTAGATGTGTA
 TTAGCTGCTTGTGTGTGCTATTGCTTATTACGGCAGTGCCACTTAAG
 TTTATGTACATGCAGCTGTTATTGTTGATGGCTGTGCTTTATTCTTACTGTT
 AACATGTTATGGCATACATGGACACTTCCATTGCCTACATTGATTACAGTTATT
 40 GGAGTTGTGCTGAAGTCCCTTCATATACAATACTCTAATTAGTCAGTTGTTATTTC
 TTAAGCCAATGGTATGATCCTGTAGTCTTGTGATACTATGGTACCATGGATGTTATTGCCA
 TTAGTGTGACTACTGCTTTAAGTGTACAAGGCTGCTATATGAATTCTTCAAACT
 TCTTGTAAATGCTGATCAGTTATGAAGTTAGGTTGTTATTACACCTCTTCAAAC
 ACTCTTACTGCATATACAGAAGGTAATTGGGAGTTATTCTTGAGTTGGTACACTATT
 45 GTGGTGGCTAATGTTAGTAGTAATTCTTAATTGGTTAATTGTTAAGTGTGCTAAG
 TGGATTTATATTATTGCAATGCAACATACTTAAATAATTATGTTAATGGCAGTCATG
 GTTAATGGCATAGGCTGGCTTGCACCTGTTACTTGGATTGTATTGGGGTTAATAAA
 GTTTTGGTTAACCTTAGGTAATAACATAATTAAAGTTCAAGTAGATCAATATAGGTAT
 ATGTGTTGCATAAGGTAATCCACCTAAACTGTGTGGGAGGTCTTACTACAAATATA
 50 CTTATACAAGGAATTGGAGGCATCGTGTGTTGCCATAGCTACAGTGCATCTAAATTG
 AGTGTGTAAGGTACAACACTGTTGTTTAATGCAGCTTGTGACTAACGTTAATGTTGAA
 GCAAATTCAAAATGCATGCTTATCTTGTGAGTTACACAATAAAATCCTCGCATCTGAT
 GATGTTGGAGAGTGCATGGATAATTATTGGGTATGCTTATAACACTATTGTATAGAT
 TCTACTATTGATTGGGTGAGTATTGTGATGATATACTTAAGAGGGTCAACTGTATTACAA
 55 TCGGTTACTCAAGAGTTTCGCACATACCCCTGCTATGCTGAATATGAAAGAGCTAAGAGT
 ATTATGAAAAGGTTAGCCATTCTAAACATGGGGTTAACACAGCAAGAGCTTGCT
 GCATATCGTAAAGCTGCCAATATTGCAAAGTCAGTTTGATAGAGACTTGCTGTTCAA

AAGAAGTTAGATAGCATGGCAGAACGTGCTATGACAACAATGTATAAAGAGGCGCGTGTAA
ACTGATAGAAGAGCAAAATTAGTTCATCATTACATGCACTACTTTTCAATGCTTAAG
AAAATAGATTCTGAGAAGCTTAATGTCTTATTGACCAGGCGAATAGTGGTGTGTACCC
CTAGCAACTGTTCAATTGTTGTAGTAATAAGCTTACCCCTGTTATACCAGACCCAGAG
5 ACCTGGGTCAAGTGTGGAGGGTGTGCATGTTACATATTCAACAGTGTGTTGGAATATA
GACTGTGTTACTGATGCCGATGGCACAGAGTTACACCCACTCTACAGGTAGTGGATTG
ACTTACTGTATAAGTGGTGATAATATAGCATGGCCTTAAAGGTTAACTGACTAGGAAT
GGGCATAATAAGGTTGATGTTGCCCTGCAAAATAATGAGCTTATGCCACGGGTGTAAAG
ACAAAGGCTTGCCTAGCAGGTGAGATCAAGCACATTGAGCGTTGAGTCTAAATGTTAT
10 TATACAAGTATTAGTGGCAGTCAGTTGAGCTGCTATTACCTCTCAAATCTAATCTG
AAAGTAGCCTCTTTGAATGAGGCAGGTAATCAGATTATGTAGACTTAGACCCACCA
TGTAATTTGGTATGAAAGTGGTGATAAGGTTGAGTTACCTGTATTGTTATAAAA
AATACGAGGTCTATTGTAAGAGGTATGGTACTTGGTGCTATATCTAATGTTGTTGTTA
CAATCTAAAGGTATGAGACAGAGGAAGTGGATGCTGAGGCATTCTCTCACTTTGTTCT
15 TTTGCAGTAGATCCTGCGGATACATATTGTAATATGAGCGAGGTAATCAACCTTA
GGTAACTGTGTTAAATGTTGACAGTACATAATGTTGAGTGGTTGCAATAACATCAAAG
CCAAGTCCAACCTCCGGATCAGGATTCTATGGAGGAGCTCTGTGTCTTATTGAGA
GCACATATAGCACACCCCTGGCGGAGCAGGAAATTAGATGGACGCTGCAATTAAAGGT
TCTTTGTGCAAATACCTACTACGGAGAAAGATCCTGTTGAGTCTGTCTACGTAACAAG
20 GTTTGCACTGTTGTCAGTGTGGATTGTTATGGATGTCAGTGTGATTCACTTAGACAA
CCTAAACCTCTGTTCACTGAGTTGCTGATCTGGTTGATAAGAATTATTAAAC
GGGTACGGGGTAGCAGTGAGGCTCGGCTGATAACCCCTAGCTAATGGATGTGACCCCGATG
TTGTAAGCGAGCCTTGATGTTGTAATAAGGAATCAGCCGGTATGTTCAAATTTGA
AGCGTAACTGTGCAAGCATTCCAAGAAGTACGTGATACTGAAGATGGAAATCTGAGTATT
25 GTGATTCTATTGTTGTTGAAACAAACACTCCTAGTAATTATGAACATGAGAAAGCTT
GTTATGAAGACTTAAAGTCAGAAGTAACAGCTGATCATGATTCTTGTTGTTCAATAAGA
ACATTATATAATTAGTAGGCAGAGGCTACTAAGTACTATGATGGATTGTTGCTATG
CTTGCCTGCACTTTGACCCAAAGGATTGCGAAGTCTTAAAGAAATACTTGTCACTTATG
GTTGTATAGAAGATTATCACCCTAAGTGGTTGAAGAGAAATAAGGATGGTACGACCCAA
30 TAGAAAACCTAAATATTATGCATGTTGGCTAAAATGGGACCTATTGTAACGACGTGCTT
TATTGAATGCTATTGAGTCGGAAACCTCATGGTTGAAAAAGGTTATGTTGGTTATTAA
CACTTGATAACCAAGATCTTAATGGCAAATTGATTGATTGGTGTATTTCAGAAGACAG
CGCCTGGTGTGGTGTCTGTTGATACGTATTCTTACATGATGCCCATCATAG
CCATGACTGATGCGTGGCACCTGAGAGGTATTTGAATATGATGTGCATAAGGGTTATA
35 AATCTTATGATCTCCTCAAGTATGATTACTGAGGAGAAACAAGATTGTTCAGAAGT
ACTTTAAGTATTGGGATCAAGAGTATCACCCCTAAGTGTGCGACTGTAGTGTGACAGGT
GTTTGATACATTGTGCAAACATCTCAACATCTGTTCTACACTGTACCGCAGACTCTT
TCGGTAATTGTTGAGAAAGGTTTGTGATGGTGTACCATTTAGCTACTTGTGGCT
ATCATTCTAAGGAACCTGGTGTATTATGAATCAAGATAACACCATGTCATTTCAAAAAA
40 TGGGTTGAGTCACACTGCAAGTGTGGAGATCCTGCCCTGTTAGTGGGACATCCA
ATAAATTAGTGGATCTTAGAACGCTTGTGTTAGTGTGCTTAGCGTCTGGTATTAA
CTCATCAAACGGTAAACCAAGGTCACTTAACAAGGATTCTACGATTGTCAGAGAAGG
CTGGTATGTTAAGGAAGGTTCTCTATACCACTTAAACATTTCTTACCCACAGACTG
GTAATGCTGCTATAAACGATTATGATTATTATCGTTATAACAGGCCCTACCATGTTGATA
45 TACGTCAACCTTATTGTTAGAAGTACCTCTAAATATTGAAATGTTATGAAGGCG
GCTGTATACCAAGCAAGCCAAGTGTAGTTAACAAATTAGATAAGAGTGCAGGTTATCCGT
TCAATAAGTTGGAAAGGCCGCTCTATTATGAATGAGTCTAGAGGAGCAGGACCAAC
TCTTGAGAGTACAAAGAAGAACGCTCTGCCTACTATAACTCAGATGAATTAAATATG
CCATATCCCGAAAAATAGAGCGCGTACAGTGGCAGGTGTCTATCCTTCTACTATGA
50 CTAATAGGCAGTTCATCAGAACGATTCTTAAGTCTATAGTCACACTAGAACGCTCCTG
TAGTTATTGGAACAAACCAAGTTTATGCCGGTGGATAACATGTTGAGAAACCTTATTG
AGGGTGTGAGACCCGATTCTTATGGGGTGGATTATCCAAGTGTGATAGAGCAATGC
CTAATTGTTGCGTATAGCAGCATCTTAGTACTCGCTCGTAAACACACTAATTGTTGTA
CTTGGTCTGAACGCGTTATAGGTTGATAATGAATGCGCTCAGGTTTATCTGAAACTG
55 TCTTAGCTACAGGTGGTATATGTGAAACCTGGTGGTACTAGCAGTGGAGATGCTACTA
CTGCTTATGCAAACAGTGTGTTCAACATAATAAGCCACATCTGCTAATGTTGCGCGTC
TTTGAGTGTATAACGCGTGTATTGTATATGATGACATTAAGAGCTTGCAGTATGAAT

TGTACCAGCAGGTTATAGGCAGTCATTGACCCAGCATTGTTGAAAAGTTTATT
 CTTATTGTGTAAGAATTCTCATTGATGATCTGTCTGACGACGGTGTGTTATA
 ACAACACATTAGCCAAACAAGGTCTGTAGCAGATATTCTGGTTTAGAGAAGTCT
 ACTATCAGAACATGTTTATGGCTGATTCTAAATGTTGGTTAACAGATTAGAAA
 5 AAGGCCACATGAATTGTTACAGCACACAATGTTAGTGGAGGTTGATGGTGAGCCTA
 GATACTTGCCATATCCAGACCCATCACGTATTTGTGTCATGTGTTTGAGATGATT
 TGGATAAGACAGAACATCTGTCGTTATGGAGCCTATATCGCTCTGCCATAGATGCCT
 ACCCACTAGTACATCATGAAAATGAGGAGTACAAGAACGTTACTCTTGCTCTTCAT
 ACATCAGAAAACCTATCAAGAGCTTCTCAGAATATGCTTATGGACTACTCTTTGAA
 10 TGGATATAGATAAGGGTAGTAAATTGGAAACAGGAGTCTATGAAAATATGTATAGAG
 CCCCTACAACATTACAGTCTTGTGGCGTTGTAGTGTGTAATAGTCAAACATATATTGC
 GCTGTGGTAATTGATTGCAAAACCATTGTGTTGTAAGTGTGCTATGACCATGTCA
 TGCACACAGACCACAAAATGTTGTCTATAAATCCTACATTGCTCACAGCCAGGTT
 GTGGTGAAGCAGATGTTACTAAATTGTACCTCGGAGGTATGTCATACCTCTGCCGTAATC
 15 ATAACCAAAGTTATCAATACCGTTAGTATCTAATGGTACAGTGTGGAAATTACAGGG
 CTAATTGTGAGGTAGCGAAAATGTTGATGATTTAATCAACTAGCTACTAATTGGT
 CTACTGTGGAACCTTATATTGGCAAATCGTTGTAGATCGTTGAGACGCTTGCTG
 CAGAGACAGTAAAAGCTACAGAAGAATTACATAAGCAACAAATTGCTAGTGCAGAAGTGA
 GAGAAGTACTCTCAGATCGTAATTGATTGATTGCTTGGAGGCCAGGTTAAACCAGGCCTC
 20 CATTGAATAGAAATTATGTTTCACTGGCTTCACTTACTAGAACTAGTAAAGTTGCTCAGC
 TCGGTGATTTACATTGAAAAAGGTGAAGGTAAGGACGTTGTCTATTATCGAGCGACGT
 CTACTGCTAAATTGTCGTGAGACATTGTTGTTAACCTCACACAATGTTGTTCTC
 TTATAGCGCCAACGTTGTGTCCTCAGCAAACCTTTCTAGGTTGTGAATTAAAGACCTA
 ATGTGATGGTACCTCGCTGTTTGTAAATAACATTCCATTGTCACATTAGTAGGCAAGC
 25 AGAAGCGTACTACAGTACAAGGCCCTCCTGGCAGTGGTAAATCCCATTGCTATAGGAT
 TGGCGGCTTACTTTAGTAACGCCGTGTCGTTTACTGCATGCTCATGCAGCTGTTG
 ATGCTTATGTGAAAAAGCTTTAAGTTCTAAAGTAGATGATTGCACTCGTATAGTAC
 CTCAAAGGACTACTATCGATTGCTCTCAAGTTAAAGCTAATGACACAGGAAAAGT
 ACATTTTAGTACTATTAAATGCCTGCCAGAAGTTAGTTGACATTCTTGTTGACG
 30 AGGTTAGTATGTTGACCAATTACGAATTGTCCTTATTAAATGGTAAGATAAACTATCAAT
 ATGTTGTTGATGTAGGGTATGCTCAATTACCGGCGCTCGTACGTTGCTAACGGTT
 CACTCTCTCAAAGGATTATAATGTTGTCACAAACCTTATGGTTGTGTTAACCTGACA
 TTTTCCTGCAAAGTGTACCGTTGTCCTAAAGAAATTGTAGATACTGTTCTACTCTG
 TATATGATGAAAGTTATTGCAAATAACCCGAATCACGTCAGTGTGTTCAAGGTTATAG
 35 TTAATAATGTAATTCTGATGTAGGACATGAAAGTGGCTAGCCTACAAACATAACTCAAT
 TAGAATTGTTGAAAGATTGTCGCAATAAGGAATGGCGGGAAAGCAACATTCAATT
 CACCTTATAATGCTATGAAACCAGAGAGCCTACCGTATGCTTGGACTTAATGTTGACAG
 TAGACTCGTCTCAAGGTTCGGAGTATGATTGTTATCTTGTGTTACTGCAGATTGCG
 AGCATGCACTGAATATTAAACAGATTCAATGTTAGCGCTTACAAGAGCCAAGCGTGGTATAC
 40 TAGTTGTCATGCGTCAGCGTGTGAACTATATTCAAGCTCTTAAGTTATAGAGCTTGATA
 GTGTAGCAAGTCTGCAAGGTACAGGTTGTTAAAATTGCAACAAAGAGTTAGTGGTG
 TTCAACCCAGCTTATGCACTGTTAACGTTGGAAGCTGGTTAGAAATAACATATAACATCTTATT
 CTTTGTGTTAGGGTTAAAGATGAGTGTGTTATGTTGAAAGGCTGCCAACATGTTATAACAC
 45 GTGATGAGGCTATCCGCAACGTAAGAGGTTGGTAGGTTGATGTTAGAAGCAACACATG
 CTTGCGGTACTAACATTGGTACTAACCTGCCTTCCAAGTAGGTTCTACTGGTGCAG
 ACTTTGTAGTTACGCCGTGAGGGACTGTAGATACTCAATAGGCAATAATTGAGCCTG
 TGAATTCTAAAGCACCTCCAGGTGAACAATTAAATCACTTGAGAGCCTATTCAAAGTG
 CTAACACCTGGCATGTTGTAAGGCCAAGGATTGTCAAATGTTAGCGATAACCTGTGCA
 50 ACGTTTCAGATTGTTAGTGTGTTGTCAGTGGTGTATGGCCTAGAAACTAACCACTTTGC
 GCTATTGTTAAAATAGGCAAGGACCAAGTTGTTCTGGCTTCTAGAGCAACAAACTT
 TTAATTCTCATACTCAGGCTTATGCTTGGAGCATTGCTTGGGTTTGTGTTATTGTT
 ATAATCCACTCTTAGGATATTCAACAGTGGGTTATTCTGGTAACCTACAATTAAACC
 ATGATTGCAATTGTAATGTCATGGACACGACATGTAGCTCTGCGGATGCTATTATGA
 55 CGCGTTGCTTGCAATTAAATGCAATTGTTAGGAGTCAAGATGTCAGACTGGGATTAACTTACC
 CTCATATGCAAATGAGGATGAAAGTCATTCTAGCTGTAGATATTACAACGCACTGATTC
 TTAATGCAATTGTTGATGCTCTTAAAGTTAACGTTGCTATGATATTAGGCAACCCCTAAAG

GTATAAAATGTGTTAGACGTGGAGACTTAAATTAGATTCTATGATAAGAACATCCAATAG
TACCCAATGTCAAGCAGTTGAGTATGACTATAATCAGCACAAAGATAAGTTGCTGATG
GTCTTTGTATGTTTGGAAATTGTAATGTGGATTGTTATCCCGACAATTCCCTAGTTGTA
GGTACGACACACGAAATTGAGTGTGTTAACCTACCTGGTGTAAATGGTGGTAGCTTGT
5 ATGTTAACAAAGCATGCATTCCACACACCTAAATTGATCGCACTAGCTTCGTAAATTGA
AAGCTATGCCATTCTTTCTATGACTCATCGCCTGCGAGACCATTCAATTGGATGGAG
TTGCGCAAGACCTGTCATTAGCTACGAAAGATTGATCACAAATGCAACATAGGCG
GTGCTGTTGTAAGAACGACGACAAATGTATGCAGATTGACTCTTATAATGCAG
CTGTTACTGCTGGTTACTTTGGGTTACTAATAATTAAACCCATATAATTGTTGGA
10 AAAGTTTCAGCTCTCCAGTCTATCGACAATTGCTTATAATATGATAAGGGTGGTC
ATTATGATGCTATTGCAAGGAGAAATGCCACTATCGTAACTGGAGATAAAAGTTTGTAA
TAGATCAAGCGTAGAAAAGCAGTTTTTAATCAAACAATTCTGCCTACATCTGTAG
CGTTGAGCTGTATGCGAAGAGAAATATTGCAACTGCCAACACCGTATTGAAAG
GTTGGGTGTAGATGTACTAATGGATTGTAATTGGGATTACACGAACCAAACACCAC
15 TATAACCGTAATACTGTTAACGATGTGCAATTACAGACATAGAACCAATGCCCTAATAG
TGCTGTATGATGATAGATATGGTATTACAGCTTTCTAGCTGCTGATAATGCTGTT
TAGTTCTACACAGTGTACAAGCGGTATTGCTATGTAGAAATACCGTCAAACCTGCTTG
TTCAGAACGGTATTCCGTTAAAGATGGAGCGAACCTGTATGTTATAAGCGTGTAAATG
GTGCGTTGTTACGCTACCTAACACATTAAACACACAGGGTCGCAGTTGAAACTTTG
20 AACCTCGTAGTGTGAGCGTGATTTCCTCGACATGTCTGAGGAGAGTTGAGAAA
AGTATGGTAAAGAATTAGGTCTACAGCACATACTGTATGGTGAAGTTGATAAGCCCCAAT
TAGGTGGTTACACACTGTTAGGTATGTCAGACTTTACGTGCGAATAAGTTGAACG
CAAAGTCTGTTACTAATTCTGATTCTGATGTATGCAAAATTATTGTTATTGGCAGACA
ATGGTTCTACAAGCAAGTGTACTGTTGGATTGCTGCTGTTGATGATTCTTAGAAC
25 TTCTTAGGAACATACTGAAAGAGTATGGTACTAATAAGTCTAAAGTTGAAACAGTGTCAA
TTGATTACCATAGCATAAATTATGACTTGGTTGAAGATGGCATTATTAAACATGTT
ATCCACAGCTTCAATCAGCATGGACGTGTTATAATATGCCACTTATAAAAGTTC
AGAATTGTTATGGAACCTTGCACATTCTAATTATGGTGTGGAATAGCGTTGCCAA
GTGGTATTATGATGAATGTGGCAAAGTATACACAACCTGTCAATTACCTTCGAAAACAA
30 CAATGTGTGACCGCATAATATGCGAGTAATGCATTGGAGCTGGAAAGTGACAAAGGAG
TGGCTCCAGGTAGTACTGTTCTAAACAATGGCTCCAGAAGGGACACTCCTGTCGATA
ATGATATTGTAGACTATGTCGATGCACATGTTCTGCTTCAAGATTGCAATAAAT
ATAAGACAGAGCACAAGTTGATCTGATATCTGATATGTATAACAGACAATGATTCAA
AAAGAAAGCATGAAGGCAGTATGCCAATAATGGCAATGATGACGTTTCAATTCT
35 CAAGTTTCTCGTAATAATTGGCTCTAGGTGGTAGTTGCTGTAAGTTGACAGAGA
CAAGTTGGCACGAAGTTTATATGACATTGCACAGGATTGTCATGGGGACAATGTTT
GTACAGCAGTGAATGCCCTTCTCAGAACGATTCTGGTTGGTTAATTATTGGGTG
CAAGTGAAGGGTTAAGGTTAGTGGAAAACGCTGCACGCAAATTATATATTGGAGGA
ATTGTAATTATTACAAACCTCTGCTTATAGTATATTGACGTTGCTAAGTTGATTG
40 GATTGAAAGCAACACCAAGTTGTTAATTGAAAAGTGAACAAAAGACAGACACTTGTCTTAA
ATTAAATTAAAGTGTGGTAAGTACTGGTAAGAGAGTGTGGTAACACCTCTTTACTAGTG
ACTCTTTGTTGTAAGTACTATGCTGCTGTTGATGACAGTAGTTCTACGTTACTAC
TACCAAAAGTGCCTTACGACACTAATGGTGGCATTACACGGGGTGCTTATGCCGTA
GTAAATTATTCTAGCGAATCTAATAATGCAGGCTCTCACCTGGGTATTGTTGGTACT
45 ATTCACTGGGGCGTGTGTTAATGCTTCTATAGCTATGACGGCACCGTCATCAGGT
ATGGCTTGGCTAGCAGTCTGACTGCACACTGTAACCTTCAGATACTACAGTG
TTGTTACACATTGTTAAATATGATGGTGTCTATAACTGGCATGCTTCAAAAGAAT
TTTTACGTGTTCTGCTATGAAAGGTTAGGCCAGCTTCTATAATTAAACAGTTAGTGTA
GCTAAAGTACCCACTTTAAATCATTCACTGTTAATAATTAAACATCCGTATATTAA
50 AATGGTGTACTTGTGTTACACCTCTAATGAGACCACAGATGTACATCTGCAGGTGTTAT
TTTAAAGCTGGGGACCTATAACCTAAAGTTAGGAGAGAAGTTAAAGCCCTGGCTTAT
TTGTTAATGGTACTGCACAAGATGTTATTGTTGTTGATGGATCACCTAGAGGGCTTGTAA
GCATGCCAGTATAACTGGCAATTTCAGATGGCTTATCCTTATTAAATAGTAGT
TTAGTTAACGAGAAGTTATTGCTATCGTGAAGGTTATTATAATTAAACACTTACGTTA
55 CACAATTTCACCTTCAATGAGACTGGCGCAACCCATAACCTAGTGGTGTCAA
ATTCAAACCTACCAACACAAACAGCTCAGAGTGGTTATTATAATTAAACCTTACGTTA
CTGAGTAGTTGTTATAAGGAGTCTAATTGTTGATGGATCTTATCACCCAAGTTGT

AATTTAGACTAGAAACTATTAATAATGGCTTGTGGTTAACACTTCAGTTCAATT
GCTTACGGTCCTCTCAAGGTGGTGCAGCAAGCAATCTGCTTTAGTGGTAGAGCAACTTGT
TGTATGCTTATTCAATGGAGGTCTCGCTGTAAAGGTGTTATTCAGGTGAGTTA
GATCTTAATTGAATGTGGACTGTTAGTTATGTTACTAAGAGCGGGCTCGTATA
5 CAAACAGCCACTGAACCGCCAGTTATAACTCGACACAATTATAATAATTACTTAAAT
ACTTGTGTTGATTATAATATATGGCAGAACTGGCCAAGGTTTATTACTAATGTAACC
GACTCAGCTGTTAGTTATAATTATCTAGCAGACGCAGGTTGGCTATTAGATACATCT
GGTCCATAGACATCTTGTGACAAGGTGAATATGGTCTTACTTATTATAAGGTTAAC
CCTTGCAGATGTCAACCAGCAGTTGTAGTTCTGGTGGTAAATTAGTAGGTATTCTT
10 ACTTCACGTAATGAGACTGGTCTCAGCTCTGAGAACAGTTACATAAAATCACT
AATGGAACACGTCGTTAGACGTTCTATTACTGAAAATGTCGAAATTGCCCTTATGTT
AGTATGGTAAGTTGTATAAAACCTGATGGTCAATTGCCACAATAGTACCAAAACAA
TTGGAACAGTTGTGGCACCTTACTTAATGTTACTGAAAATGTCGTCATACCTAACAGT
TTAATTAACTGTTACAGATGAGTACATACAAACGCGTATGGATAAGGTCCAATTAAAT
15 TGTCTGCAGTATGTTGTGGCAATTCTCTGGATTGTAGAGATTGTTCAACAAATATGGG
CCTGTTGTGACAACATATTGTCGTAGTAAATAGTATTGGTCAAAAAGAAGATATGGAA
CTTTGAATTCTATTCTACTAAACCGGCTGGTTAATACACCATTCTTAGTAAT
GTTAGCACTGGTGAGTTAATATTCTCTGTTAACAACTCCTAGTAGTCCTAGAAGG
CGTTCTTTATTGAAGACCTCTATTACAAGCGTTAACATGTTGGATTACCAACAGAT
20 GACGCATACAAAAATTGCACTGCAGGACCTTGTAGGTTCTAACGGACCTTGCCTGCT
CGTGAATATAATGGTTGCTGTGCTCCATTATAACAGCAGAAATGCAAATTGG
TATACTAGTCTCTAGTAGCTCTATGGCTTTGGTGTATTACTGCAGCTGGTGCTATA
CCTTTGCCACACAACCTGCAGGCTAGAATTAACTCACTGGGTATTACCCAGTCATT
TTGAGAATCAAGAAAAATTGCTGCTCCTTAAATAAGGCCATTGGTGTATGCAGGAA
25 GGTTTGAAGTACATCTAGCATTACAACAAATTCAAGATGTTGTTAACAGCAGAGT
GCTATTCTACTGAGACTATGGCATCACTAAATAAAATTGGTGTATTCTCTATG
ATTCAAGAAATCTACCAAGCAACTTGACGCCATACAAGCAAATGCTCAAGTGGATCGTCTT
ATAACTGGTAGATTGTCATCACTTCTGTTTAGCATCTGCTAACAGCAGGCGGAGCATATT
AGAGTGTACAACAGCGTGAGTTAGCTACTCAGAAAATTAAATGAGTGTGTTAACAG
30 TCTATTAGGTACTCCTTGTGGTAATGGACGACATGTTCTAACCATACCGCAAATGCA
CCTAATGGTATAGTGTAAACACTTTCTTACTCCAGATAGTTGTTAACATGTTACT
GCAATAGTGGTTTTGTGTAAGCCAGCTAATGCTAGTCAGTATGCAATAGTACCCGCT
AATGGTAGGGTATTAAACAAAGTTAATGGTAGTTACTACATCACAGCAGGAGATATG
TATATGCCAAGAGCTATTACTGCAGGAGATAGTTACGCTACTCTTGTCAAGCAAAT
35 TATGTAAGTGTAAATAAGACCGTCATTACTACATTGCTAGACAATGATGATTGATT
AATGACGAATTGTCAAAATGGTGAATGACACTAACGATGAGCTACCAAGACTTGACAAA
TTCAATTACACAGTACCTATACTTGACATTGATAGTGAATTGATCGTATTCAAGGCGTT
ATACAGGGCTTAATGACTCTTAATAGACCTTGAAAACATTCAAAACTTAACTCAAAC
ATTAAAGTGGCTTGGTATGTTGTTAGCCATAGCTTGCCTACTATTATCTCATCTTA
40 ATACTAGGATGGTTCTTCATGACTGGATGTTGGTGTGTTGATGCTTGGC
ATTATGCCTCTAAATGAGTAAGTGTGGTAAGAAATCTTCTTATTACACGACTTTGATAAC
GATGTGGTAACTTAACAATACAGACCTAAAAGTCTGTTAATGATTCAAAGTCCCACGT
CCTCCTAATAGTATTAAATTCTTGTGTAACATTGACTAACAGTGTGTTAGAGAGT
TTATTATAGCGCTCCAACAACAAACTAACAGTTTACTCCAAATTATCAATAGTAAC
45 AGCCTAGACTGACCCTTGTACAGTCTAGACTAACAGTAAACTAGAACGAAATTATTGA
AACTGGTGAGCAAGTGATTCAAAAAACTCAGTTCAATTACAGCATAATTCAAGTGTATT
AAACACAGAAGTATTGACCCCTTGACTATTGTTATTACAGAGGAGTAATTGGGA
AATAGAGTCAGCTGAAGATTGTCAGGTGATGATGAATTATTGAATAAGTCGTAGAGG
AAAATGGAAGTTTCTAACAGCGCTTATATATTGTTAGGATTGTTAGCACTTATCTC
50 TAGGTAGAGCACTTCAAGCATTGTCAGGCTGCTGATGCTGTTGTTATTGGTATA
CATGGTAGTAATTCCAGGAGCTAAGGGTACAGCCTTGTATATAAGTATAACATATGGTA
GAAAACCTAACATCCGGATTAGAACAGCTATTGTCAACGAGTTCTAAGAACGGTT
GGAATAATAAAATCCAGCAAATTCAAGATGTCACAGGAGAACATTGACTCTTGAC
TTGAAACAGTCAGTTGAGCTTTAAAGAGTATAATTATTATAACTGCATTCTGTTG
55 TTCTTAACCATAACTTCAGTATGGCTATGCAACAAGAACAGTAAAGTTATTATATACTG
AAAATGATAGTGTATGGTCTTGGCCCTAACATTGCAAGTAGGTGTAATTGATG
ATATACCCACCAACACAGGAGGTCTTGTGCGAGCGATAATACTTACAGTGTGCGTGT

CTGTCTTTGTAGGTTATTGGATCCAGAGTATTAGACTCTTAAGCGGTGTAGGTCATGG
 TGGTCATTAAACCCAGAACATCTAATGCCGTAGGTTCAATACTCTAACATAATGGTCAACAA
 TGTAATTTGCTATAGAGAGTGTGCCAATGGTGCTTCTCCAATTATAAAGAACATGGTGT
 CTTTATTGTGAGGGTCAGTGGCTGCTAAGTGTGAACCAGACCACCTGCCTAAAGATATA
 5 TTTGTTGTACACCGGATAGACGTAATATCTACCGTATGGTCAGAAATAACTGGTGAC
 CAAAGCGGAAATAAGAAACGGTTGCTACGTTGTCTATGCAAAGCAGTCAGTAGATACT
 GGCGAGCTAGAAAGTGTAGCAACAGGAGGGAGTAGTCTTACACCTAAATGTGTGTGT
 AGAGAGTATTAAAATTATTCTTAATAGTGCCTCTATTAAAGAGGCATAATAGTATT
 ATTTTGAGGATATTAATATAAATCCTCTGTGTTATACTCTCTTCAAGAGCTATTAA
 10 TTTAAAAAACAGTTTCCACTCTTGCCAAAAACTATTGTTGTAATGGTGTAAACC
 TTTCAAGTAGATAATGGAAAGTCTACTACGAAGGAAACCAATTTCAGAAAGGTTGT
 TGAGGTTGCGGTAGTTATAAAAAGATTAAACTACCTACTACACCTATTTTATAAG
 AGGCCTTTATCTTACAAGCGCTTAATAAACGACGATGAAATGGCTGACTAGTTTG
 TAAGGGCAGTTATTCATGTTATAAACCCCTATTATAACTCAATTAAGAGTATTAGATA
 15 GGTAAATCTTAGATCATGGACCAAAACACATCTAACGTGTGTTAGGTGCGTGTATTG
 TTCAATTAGATTAGTTATAGGTTGGCGTATACCCCTACTCAATCGCTGGTATGAATAA
 TAGTAAAGATAATCCTTTTGCGGAGCAATAGCAAGAAAAGCGCGAATTATCTGAGAGA
 AGGATTAGATTGTGTTACTTTCTTAACAAAGCAGGACAAGCAGACTTGTCCCCTG
 TACCTCTCTAGTATTCCAGGGAAAACCTGTGAGGAACACAAATATAAATAATCTTT
 20 GTCATGGCAAGCGTAAGGCAACTGGAAAGACAGATGCCAGCTCCAGTCATCAAACTA
 GGAGGACCAAGCCACCTAAAGTTGGTTCTCTGGAAATGTATCTTGGTTCAAGCAATA
 AAAGCCAAGAAGTTAAATTCAACCTCCGCTTAAGTTGAAGGTAGCGGTGTTCTGATAAT
 GAAAATCTAAAACCAAGTCAGCAGCATGGATATTGGAGACGCCAAGCTAGGTTAAGCCA
 GGTAAAGGTGGAAGAAAACCAGTCCCAGATGCTGGTATTTTACTATACTGGAACAGGA
 25 CCAGCCGCTAACCTGAATTGGGGTGTAGCCAAGATGGTATAGTGTGGGTGCTGGTAAG
 GGTGCTGATACTAAATTAGATCTAACCTGGTACTCGTACTGACAAGTTGACCAA
 TATCCGCTACGGTTTCAGACGGAGGACCTGATGGTAATTCCGTTGGGATTTCATTCT
 CTGAATCGTGGCAGGAGTGGGAGATCAACAGCAGCTTACAGCAGCATCTAGTAGAGCA
 CCATCACGTGAAGTTCGCGTGGCAGGAGTGGTCTGAAGATGATCTTATTGCTCGT
 30 GCAGCAAGGATAATTCAAGGATCAGCAGAAGAAGGGTCTCGCATTACAAAGGCTAACGGCT
 GATGAAATGGCTCACCGCCGGTATTGCAAGCGCACTATTCCACCTAATTATAAGGTTGAT
 CAAGTGTGTTGGTCCCCGTACTAAAGGTAAGGAGGGAAATTGGTGTGACAAGGATGAAT
 GAGGAAGGTATTAAGGATGGGCGCGTTACAGCAATGCTAACCTAGTCCCTAGCAGCCAT
 GCTTGTCTTTCGGAAGTAGAGTGACGCCAGACTCAACCAGATGGCTGCACTTGAAA
 35 TTGGAATTACTACTGTGGTCCACGTGATGATCCGAGTTGATAATTATGTTAAAGGTT
 TGTGATCAGTGTGTTGATGGTAGGAACACGTCCAAAAGATGATGAACCAAGACCAAG
 TCACGCTCAAGTTCAAGACCTGCAACAAGAGGAAATTCTCAGCGCCAAGACAGCAGCG
 CCTAAGAAGGAGAAAAGCCAAGAAGCAGGATGATGAAGTGGATAAAGCATTGACCTCA
 GATGAGGAGAGGAACAATGCACAGCTGGAATTGATGATGAACCCAAGGTAATTAACTGG
 40 GGGGATTCCAGCCCTAGGAGAGAATGAACCTTGAGTAAAATTCAATAGTAAGAGTTAAGGA
 AGATAGGCATGTAGCTTGATTACCTACATGTCTATGCCAGGGAAATGTCTAATTGTCT
 ACTTAGTAGCCTGGAAACGAACCGTAGACCCCTAGATTAAATTAGTTAATTGTTAATT
 TTAGTTAAGTTAGTTAGAGTAGGTATAAAGATGCCAGTGCCGGGGCACGCCAGTAC
 GACCGAGGGTACAGCACTAGGACGCCATTAGGGGAAGAGCTAAATTAGTTAAGTTA
 45 AGTTAATTGGCTATGTATAGTAAAATTATAGGCTAGTATAGAGTTAGAGCAAAAAAA
 AAAAAAAAAAAAAAA

REPLICASE

50 In addition to the structural and accessory genes, two-thirds of a coronavirus genome
 comprises the replicase gene (at the 5' end of the genome), which is expressed as two
 polyproteins, pp1a and pp1ab, in which pp1ab is an extension product of pp1a as a result
 of a -1 ribosomal shift mechanism. The two polyproteins are cleaved by two types of

virus-encoded proteinases usually resulting in 16 non-structural proteins (Nsp1-16); IBV lacks Nsp1 thereby encoding Nsp2-16.

Thus Gene 1 in IBV encodes 15 (16 in other coronaviruses) non-structural proteins (nsp2-16), which are associated with RNA replication and transcription.

The term 'replicase protein' is used herein to refer to the pp1a and pp1ab polyproteins or individual nsp subunits.

10 The term 'replicase gene' is used herein to refer to a nucleic acid sequence which encodes for replicase proteins.

A summary of the functions of coronavirus nsp proteins is provided in Table 1.

Table 1

Nsp Protein	Key features
1	Conserved within but not between coronavirus genetic groups; potential regulatory functions in the host cell.
2	Disposable for MHV and SARS-CoV replication in tissue culture
3	Acidic domain; macro domain with ADRP and poly(ADP-ribose)-binding activities; one or two ZBD-containing papain-like proteases; Y domain
4	Transmembrane domain
5	3C-like main protease, homodimer
6	Transmembrane domain
7	Interacts with nsp8 to form a hexadecamer complex
8	Noncanonical RNA polymerase; interacts with nsp7 to form a hexadecameric complex
9	ssRNA-binding protein, dimer
10	RNA-binding protein, homododecamer, zinc-binding domain, known to interact with nsp14 and nsp16
11	Unknown
12	RNA-dependent RNA polymerase
13	Zinc-binding domain, NTPase, dNTPase, 5'-to-3' RNA and DNA helicase, RNA 5'-triphosphate
14	3'-to 5' exoribonuclease, zinc-binding domain and N7-methyltransferase
15	Uridylate-specific endoribonuclease, homohexamer
16	Putative ribose-2'-O-methyltransferase

The variant replicase gene encoded by the coronavirus of the present invention
5 comprises a mutation in one or more of the sections of sequence encoding nsp-10, nsp-14, nsp-15 or nsp-16.

Nsp10 has RNA-binding activity and appears to be involved in homo and/or heterotypic interactions within other nsps from the pp1a/pp1ab region. It adopts an α/β fold comprised of five α -helices, one 3_{10} -helix and three β -strands. Two zinc-binding sites have been identified that are formed by conserved cysteine residues and one histidine residue (Cys-74/Cys-77/His-83/Cys-90; Cys-117/Cys-120/Cys-128/Cys-130). The protein has been confirmed to bind single-stranded and double-stranded RNA and DNA without obvious specificity. Nsp-10 can be cross-linked with nsp-9, suggesting the existing of a complex network of protein-protein interactions involving nsp-7, -8, -9 and -10. In addition, nsp-10 is known to interact with nsp-14 and nsp-16.

Nsp-14 comprises a 3'-to-5' exoribonuclease (ExoN) active domain in the amino-terminal region. SARS-CoV ExoN has been demonstrated to have metal ion-dependent 3'-to-5' exoribonuclease activity that acts on both single-stranded and double-stranded RNA, but not on DNA. Nsp-14 has been shown to have proof-reading activity. This nsp has also been shown to have N7-methyltransferase (MT) activity in the carboxyl-terminal region.

Nsp-15 associated NendoU (nidoviral endoribonuclease, specific for U) RNase activity has been reported for a number of coronaviruses, including SARS-CoV, MHV and IBV. The activities were consistently reported to be significantly enhanced by Mn²⁺ ions and there was little activity in the presence of Mg²⁺ and Ca²⁺. NendoU cleaves at the 3' side of uridylate residues in both single-stranded and double-stranded RNA. The biologically relevant substrate(s) of coronavirus NendoUs remains to be identified.

Nsp-16 has been predicted to mediate ribose-2'-O-methyltransferase (2'-O-MTase) activity and reverse-genetics experiments have shown that the 2'-O-MTase domain is essential for viral RNA synthesis in HCoV-229E and SARS-CoV. The enzyme may be involved in the production of the cap 1 structures of coronavirus RNAs and it may also cooperate with NendoU and ExoN in other RNA processing pathways. 2'-O-MTase might also methylate specific RNAs to protect them from NendoU-mediated cleavage.

The genomic and protein sequences for nsp-10, -14, -15 and -16 are provided as SEQ ID NO: 2-5 and 6-9, respectively.

SEQ ID NO: 2 (nsp-10 nucleotide sequence – nucleotides 11884-12318 of SEQ ID NO:1)

TCTAAAGGTCACTGAGACAGAGGAAGTGGATGCTGTAGGCATTCTCACTTTGTTCTTTGCAGTA
GATCCTCGGGATACTATATTGTAATATGTGGCAGCAGGTAATCAACCTTTAGGTAACTGTGTTAAA
ATGTTGACAGTACATAATGGTAGTGGTTTGCAATAACATCAAAGCCAAGTCCAACCTCCGGATCAG
GATTCTTATGGAGGAGCTCTGTGTCTTATTGTAGAGCACATATAGCACACACCTGGCGGAGCA
GGAAATTAGATGGACGCTGTCAATTAAAGGTTCTTGTGCAAATACCTACTACGGAGAAAGAT
CCTGTTGGATTCTGTCTACGTAAACAAGGTTGCACGTGTTGTCAGTGGTGGATTGGTTATGGATGT
CAGTGTGATTCACTTAGACAACCTAAACCTTCTGTTCACT

SEQ ID NO: 3 (nsp-14 nucleotide sequence – nucleotides 16938-18500 of SEQ ID NO:1)

GGTACAGGCTTGTAAAATTGCAACAAAGAGTTAGTGGTGTACCCAGCTATGCAGTCACA
ACTAAGGCTTGTGCAACTTATAAAGTTAATGATGAACCTGCTGCACTTGTAAACGTGGAAGCT
GGTCAGAAATAACATATAAACATCTTATTCTTAGGGTTAACATGAGTGTAAATGTTGAA
GGCTGCCACAACATGTTATAACACGTGATGAGGCTATCCGCAACGTAAGAGGTTGGTAGGTTT
GATGTAGAAGCAACACATGCTTGCAGTACTAACATTGTTACAACTGCCTTCCAAGTAGGTTTC
TCTACTGGTGCAGACTTGTAGTTACGCCTGAGGGACTTGTAGATACTCAATAGGAATAATT
GAGCCTGTGAATTCTAAAGCACCTCCAGGTGAACAATTAACTCACTTGAGAGCGTTATTCAAAGT
GCTAAACCTTGGCATGTTGAAGGCCAAGGATTGTGCAAATGTTAGGGATAACCTGTGCAACGTT
TCAGATTGTAGTGTACGTGTCAGTGGTGTACGGCTAGAAACTAACCACTTGCCTATTGTT
AAAATAGGCAAGGACCAAGTTGTTCTTGCAGTACTAACATTAACTCAATGTTACACTCAG
GCTTATGCTTGGAAAGCATTGCTGGTTTGATTGTTATAATCCACTCTTAGTGGATATT
CAACAGTGGGTTATTCTGGTAACCTACAATTAAACCATGATTGCAATTGTTACACTGCAATT
GCACATGTAGCTTCTGCGGATGCTATTATGACGCCTGTCTGCAATTAAATGCATTGCAA
GATGTCAACTGGGATTAACTTACCTCATATAGCAAATGAGGATGAAGTCAATTCTAGCTGTAGA
TATTTCACAACGCATGTATCTTAAATGCATGTGTTGATGCTCTAAAGTAACGTTGTCTATGATA
GGCAACCCTAAAGGTATTAAATGTGTTAGACGTGGAGACTTAAATTAGATTCTATGATAAGAAT
CCAATAGTACCCAAATGTCAAGCAGTTGAGTATGACTATAATCAGCACAAAGATAAGTTGCTGAT

GGTCTTGTATGTTGGAATTGTAATGTGGATTGTTATCCCGACAATTCTTACTTTGTAGGTAC
 GACACACGAAATTGAGTGTAAACCTACCTGGTTGTAATGGTAGCTGTATGTTAACAAAG
 CATGCATTCCACACACCTAAATTGATCGCACTAGCTTCGTAATTGAAAGCTATGCCATTCTT
 TTCTATGACTCATCGCCTTGCAGACCAATTCAATTGGATGGAGTTGCGCAAGACCTGTGTCATTA
 5 GCTACGAAAGATTGATCACAAATGCAACATAGGCGGTGCTGTTGTAAGGACGCACAAATG
 TATGCAGATTGACTTCTTATAATGCAGCTTACTGCTGGTTACTTTGGTTACTAAT
 AATTAAACCATATAATTGAAAGTTTCAGCTCCAG

SEQ ID NO: 4 (nsp-15 nucleotide sequence – nucleotides 18501-19514 of SEQ ID NO:1)

10 TCTATCGACAATATTGCTTATAATATGTATAAGGGTGGTCATTATGATGCTATTGCAGGAGAAATG
 CCCACTATCGTACTGGAGATAAAAGTTTGTATAGATCAAGGCGTAGAAAAGCAGTTTTTT
 AATCAAACAATTCTGCCTACATCTGTAGCGTTGAGCTGTATGCGAAGAGAAATATCGCACACTG
 CCAAACAACCGTATTTGAAAGGTTGGGTGAGATGTGACTAATGGATTGTAATTGGGATTAC
 15 ACGAACCAAACACCATAACCGTAATACTGTTAAGGTATGTCATATACAGACATAGAACCAAAT
 GGCTTAATAGTGTATGATGATAGATGGTATTACCAAGTCTTCTAGCTGCTGATAATGCT
 GTTTAGTTCTACACAGTGTACAAGCGGTATTCTGTATGTAAGAAATACCGTCAAACCTGCTGTT
 CAGAACCGTATTCCGTTAAAGATGGAGCGAACCTGTATGTTATAAGCGTGTAAATGGTGCCTT
 GTTACGCTACCTAACACAATAAACACACAGGTCGAAGTTATGAAACCTTGTAAAGTGT
 20 GTTGAGCGTGATTTCTCGACATGTCAGGGAGAGTTGAGAAAGTATGGTAAAGAATTAGGT
 CTACAGCACATACTGTATGGTGAAGTTGATAAGCCCCAATTAGGTGGTTCCACACTGTTAGGT
 ATGTGCAGACTTTACGTGCGAATAAGTGAACGCAAAGTCTGTTACTAATTCTGATTCTGATGTC
 ATGCAAAATTATTTGTATTGGCAGACAATGGTCTACAAGCAAGTGTACTGTTGTGGATTG
 CTGCTTGATGATTCTAGAACACTGAAACTACTGAAAGAGTATGGTACTAATAAGCTAA
 25 GTTGTAAACAGTGTCAATTGATTACCATAGCATAAATTGACTTGGTTGAAGATGGCATTATT
 AAAACATGTTATCCACAGCTTCAA

SEQ ID NO: 5 (nsp-16 nucleotide sequence – nucleotides 19515-20423 of SEQ ID NO:1)

30 TCAGCATGGACGTGTGGTTATAATATGCCGAACATTATAAGTTCAGAATTGTGTTATGGAACCT
 TGCAACATTCTAATTATGGTGTGGAATAGCGTTGCCAAGTGGTATTATGATGAAATGTGGCAAAG
 TATACACAACCTGTCAATACCTTCGAAAACAACAAATGTGTACCGCATAATATGCGAGTAATG
 CATTTGGAGCTGGAAGTGACAAAGGAGTGGTGCCAGGTAGTACTGTTCTAAACAATGGCTCCCA
 35 GAAGGGACACTCCTGTCGATAATGATATTGTAGACTATGTGTCTGATGCACATGTTCTGTGCTT
 TCAGATTGCAATAAAATATAAGACAGAGCACAAGTTGATCTGTGATATCTGATATGTATACAGAC
 AATGATTCAAAAAGAAAGCATGAAGCGTGATAGCCAATAATGGCAATGATGACGTTTCATATAT
 CTCTCAAGTTCTCGTAATAATTGGCTCTAGGTGGTAGTTGCTGTAAGAGTACAGAGACA
 AGTGGCACGAAGTTTATATGACATTGACAGGATTGTCATGGTGACAATGTTGTACAGCA
 40 GTGAATGCCCTCTTCTCAGAACGATTCTGATTGGTGTAAATTATTGGTGCAAGTGAAAGGTT
 AAGTTAGGGAAAAACGCTGCACGCAAATTATATTTGGAGGAATTGTAATTATTCACAAACC
 TCTGCTTATAGTATATTGACGTTGCTAAGTTGAGATTGAAAGCAACGCCAGTTGTTAAT
 TTGAAAACGACACAGACTTAGTCTTAAATTAAAGTGTGGTAAGTTACTGTTAAG
 GATGTTGGTAACACCTCTTACTAGTGAATCTTGTGTACTATGTAG

45

SEQ ID NO: 6 (nsp-10 amino acid sequence)

50 SKGHETEEVDAVGILSLCSFAVDPADTYCKYVAAGNQPLGNCVKMLTVHNGSGFAITSKPSPTPDQ
 DSYGGASVCLYCRHIAHPGGAGNLDGRQFKGSFVQIPTTEKDPVGFCRLNKVCTVCQCWIGYGC
 QCDSLQRPKPSVQ

SEQ ID NO: 7 (nsp-14 amino acid sequence)

5 GTGLFKICNKEFSGVHPAYAVTTKALAATYKVNDLAAALVNVEAGSEITYKHLISLLGFKMSVNVE
GCHNMFITRDEAIRNVRGVGFVDEATHACGTNIGTNLPFQVGFSTGADFVVTPEGLVDTSIGNNF
EPVNSKAPPGEQFNHLRALFKSAKPWHVRPRIVQMLADNLNCVSDCVVFVTCWCHGLELTTLRYFV
KICKDQVCSCGSRATTFSNSHTQAYACWKHCLGFDFVYNPLLVLDIQQWGYSGNLQFNHDLHCNVGH
AHVASADAIMTRCLAINNAFCQDVNWDLTYPHIANEDEVNSSCRYLQRMYLNACVDALKVNVYDI
GNPKGIKCVRRGDLNFRFYDKNPIVPNVKQFYEYDYNQHKDKFADGLCMFWNCVDCYPDNSLVCRY
DTRNLSVFNLPGCNGGSLYVNKHAFHTPKFDRTSFRNLKAMPFFYDSSPCETIQLDGVAQDLVSL
ATKDCITKCNIGGAVCKKHAQMYADFVTSYNAAVTAGFTFWVTNNFNPYNLWKSFSALQ

10

SEQ ID NO: 8 (nsp-15 amino acid sequence)

15 SIDNIAYNMYKGGHYDAIAGEMPTIVTGDKVFVIDQGVEKAVFFNQTILPTSVAFELYAKRNIRTL
PNNRILKGLGVDTNGFVIWDYTNQTPLYRNTVKVCAYTDIEPNGLIVLYDDRYGDYQSFLAADNA
VLVSTQCYKRYSYVEIPSNLLVQNGIPLKDGANLYVYKRVNGAFVTLPTNLNTQGRSYETFEPRSD
VERDFLDMSEESFVEKYGKELGLQHILYGEVDKPKQLGLHTVIGMCRLRANKLNAKSVTNSDSDV
MQNYFVLADNSYKQVCTVVDLLLDDFLELLRNILKEYGTNKSKVVTVSIDYHSINFMTWFEDGII
KTCYPQLQ

20

SEQ ID NO: 9 (nsp-16 amino acid sequence)

25 SAWTCGYNMPELYKVQNCVMEPCNIPNYGVGIALPSGIMMNVAKYTQLCQYLSKTTMCVPHNMRV
HFGAGSDKGVARGSTVLKQWLPEGTLLVDNDIVDYVSDAHVSVLSDCNKYKTEHKFDLVIISDMYTD
NDSKRKHEGVIANNNGNDDVFIYLSSFLRNNLALGGSFAVKVTETSWHEVLYDIAQDCAWWTMFCTA
VNASSEAFLVGVNYLGASEKVKVSGKTLHANYIFWRNCNYLQTSAYSIFDVAKFDLRLKATPVVN
LKTEQKTDLVFNLIKCGKLLVRDVGNTSFTSDSFVCTM

REDUCED PATHOGENICITY

30 The live, attenuated coronavirus of the present invention comprises a variant replicase gene which causes the virus to have reduced pathogenicity compared to a coronavirus expressing the corresponding wild-type gene.

35 The term "attenuated" as used herein, refers to a virus that exhibits said reduced pathogenicity and may be classified as non-virulent. A live, attenuated virus is a weakened replicating virus still capable of stimulating an immune response and producing immunity but not causing the actual illness.

40 The term "pathogenicity" is used herein according to its normal meaning to refer to the potential of the virus to cause disease in a subject. Typically the pathogenicity of a coronavirus is determined by assaying disease associated symptoms, for example sneezing, snicking and reduction in tracheal ciliary activity.

The term "reduced pathogenicity" is used to describe that the level of pathogenicity of a coronavirus is decreased, lessened or diminished compared to a corresponding, wild-type coronavirus.

- 5 In one embodiment, the coronavirus of the present invention has a reduced pathogenicity compared to the parental M41-CK virus from which it was derived or a control coronavirus. The control coronavirus may be a coronavirus with a known pathogenicity, for example a coronavirus expressing the wild-type replicase protein.
- 10 The pathogenicity of a coronavirus may be assessed utilising methods well-known in the art. Typically, pathogenicity is assessed by assaying clinical symptoms in a subject challenged with the virus, for example a chicken.

As an illustration, the chicken may be challenged at 8-24 days old by nasal or ocular inoculation. Clinical symptoms, associated with IBV infection, may be assessed 3-10 days post-infection. Clinical symptoms commonly assessed to determine the pathogenicity of a coronavirus, for example an IBV, include gasping, coughing, sneezing, snicking, depression, ruffled feathers and loss of tracheal ciliary activity.

- 20 The variant replicase of the present invention, when expressed in a coronavirus, may cause a reduced level of clinical symptoms compared to a coronavirus expressing a wild-type replicase.


For example a coronavirus expressing the variant replicase may cause a number of 25 snicks per bird per minute which is less than 90%, less than 80%, less than 70%, less than 60%, less than 50%, less than 40%, less than 30%, less than 20% or less than 10% of the number of snicks caused by a virus expressing the wild type replicase.

A coronavirus expressing a variant replicase according to the present invention may 30 cause wheezing in less than 70%, less than 60%, less than 50%, less than 40%, less than 30%, less than 20% or less than 10% of the number of birds in a flock infected with the a virus expressing the wild type replicase.

A coronavirus expressing a variant replicase according to the present invention may 35 result in tracheal ciliary activity which is at least 60%, at least 70%, at least 80%, at least 90% or at least 95% of the level of tracheal ciliary activity in uninfected birds.

A coronavirus expressing a variant replicase according to the present invention may cause clinical symptoms, as defined in Table 2, at a lower level than a coronavirus expressing the wild type replicase.

5 Table 2 IBV severity limits based on clinical signs:

Snicking (sneezing)		<p>IBV specific: Mild (N.B. Respiratory signs become apparent from 2-3 dpi if they are going to occur and can continue for up to 7d).</p>
Nasal exudate		
Watery eyes		
Swollen infraorbital sinuses		
Rales (vibration in trachea or bronchi region)		<p>Mild, if exceed 2d increase to moderate</p>
Hunched posture / depressed		
Fluffed up feathers		
Eating and drinking less		<p>IBV specific: Mild, if exceed 24h increase to moderate for a max of 2d. If still drinking in excess then kill by schedule 1 method.</p>
Drinking in excess: evident by fluid filled crop or measured water intake		
Less active but still evade capture		
Weight loss		<p>Mild, if exceed 1d increase to moderate.</p>
Not eating or drinking		
Birds sit alone and does not evade capture		<p>Moderate: birds at end point. Kill by schedule 1 method.</p>
Severe respiratory distress: e.g. excessive gasping		
Snicking and / or rales for 7d in total		
Found dead		<p>Severe: report to project license holder. Full post-mortem to be performed.</p>

The variant replicase of the present invention, when expressed in a coronavirus, may cause the virus to replicate at non-pathogenic levels *in ovo*.

10

While developing vaccines to be administered *in ovo* to chicken embryos, attention must be paid to two points: the effect of maternal antibodies on the vaccines and the effect of the vaccines on the embryo. Maternal antibodies are known to interfere with active immunization. For example, vaccines with mild strains do not induce protective antibody levels when administered to broiler chickens with maternal antibodies as these strains are neutralized by the maternal antibody pool.

15

Thus a viral particle must be sufficiently efficient at replicating and propagating to ensure that it is not neutralized by the maternally-derived antibodies against the virus. Maternally-derived antibodies are a finite pool of effective antibodies, which decrease as the chicken ages, and neutralization of the virus in this manner does not equate to the 5 establishment of long-term immunity for the embryo/chick. In order to develop long-term immunity against the virus, the embryo and hatched chicken must develop an appropriate protective immune response which is distinct to the effect of the maternally-derived antibodies.

10 To be useful for *in ovo* vaccination, the virus must also not replicate and propagate at a level which causes it to be pathogenic to the embryo.

Reduced pathogenicity in terms of the embryo may mean that the coronavirus causes less reduction in hatchability compared to a corresponding, wild-type control coronavirus. 15 Thus the term "without being pathogenic to the embryo" in the context of the present invention may mean "without causing reduced hatchability" when compared to a control coronavirus.

A suitable variant replicase may be identified using methods which are known in the art. 20 For example comparative challenge experiments following *in ovo* vaccination of embryos with or without maternally-derived antibodies may be performed (i.e. wherein the layer has or has not been vaccinated against IBV).

If the variant replicase enables the virus to propagate at a level which is too high, the 25 embryo will not hatch or will not be viable following hatching (i.e. the virus is pathogenic to the embryo). A virus which is pathogenic to the embryo may kill the embryo.

If the variant replicase causes a reduction in viral replication and propagation which is too great, the virus will be neutralised by the maternally-derived antibodies. Subsequent 30 challenge of the chick with IBV will therefore result in the development of clinical symptoms (for example wheezing, snicking, loss of ciliary activity) and the onset of disease in the challenged chick; as it will have failed to develop effective immunity against the virus.

35 VARIANT

As used herein, the term 'variant' is synonymous with 'mutant' and refers to a nucleic acid or amino acid sequence which differs in comparison to the corresponding wild-type sequence.

5 A variant/mutant sequence may arise naturally, or may be created artificially (for example by site-directed mutagenesis). The mutant may have at least 70, 80, 90, 95, 98 or 99% sequence identity with the corresponding portion of the wild type sequence. The mutant may have less than 20, 10, 5, 4, 3, 2 or 1 mutation(s) over the corresponding portion of the wild-type sequence.

10 The term "wild type" is used to mean a gene or protein having a nucleotide or amino acid sequence which is identical with the native gene or protein respectively (i.e. the viral gene or protein).

15 Identity comparisons can be conducted by eye, or more usually, with the aid of readily available sequence comparison programs. These commercially available computer programs can calculate % identity between two or more sequences. A suitable computer program for carrying out such an alignment is the GCG Wisconsin Bestfit package (University of Wisconsin, U.S.A.; Devereux *et al.*, 1984, Nucleic Acids Research 12:387).

20 Examples of other software that can perform sequence comparisons include, but are not limited to, the BLAST package (see Ausubel *et al.*, 1999 *ibid* – Chapter 18), FASTA (Atschul *et al.*, 1990, *J. Mol. Biol.*, 403-410) and the GENEWORKS suite of comparison tools, ClustalX (see Larkin *et al.* (2007) Clustal W and Clustal X version 2.0. *Bioinformatics*, 23:2947-2948). Both BLAST and FASTA are available for offline and 25 online searching (see Ausubel *et al.*, 1999 *ibid*, pages 7-58 to 7-60). However, for some applications, it is preferred to use the GCG Bestfit program. A new tool, called BLAST 2 Sequences is also available for comparing protein and nucleotide sequence (see FEMS *Microbiol. Lett.* 1999 174(2): 247-50; FEMS *Microbiol. Lett.* 1999 177(1): 187-8 and tatiana@ncbi.nlm.nih.gov).

30 The sequence may have one or more deletions, insertions or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent molecule. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues as long as the activity is retained. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity

values include leucine, isoleucine, valine, glycine, alanine, asparagine, glutamine, serine, threonine, phenylalanine, and tyrosine.

Conservative substitutions may be made, for example according to the Table below.

5 Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for each other:

ALIPHATIC	Non-polar	G A P
		I L V
	Polar - uncharged	C S T M
		N Q
Polar - charged		D E
		K R
AROMATIC		H F W Y

10 The coronavirus of the present invention may comprise a variant replicase gene which encodes a protein which comprises a mutation compared to any one of SEQ ID NO: 6, 7, 8 or 9 which, when expressed in a coronavirus, causes the virus to have reduced pathogenicity compared to a coronavirus expressing the corresponding wild-type replicase.

15 The variant replicase gene may encode a protein which comprises at least one or more amino acid mutations in any combination of nsp-10, nsp-14, nsp-15 and nsp-16.

20 The variant replicase gene of the coronavirus of the present invention may encode a protein comprising a mutation as defined in the M41 mod sequences presented in Figure 10.

The variant replicase gene of the coronavirus of the present invention may encode a protein which comprises one or more amino acid mutations selected from the list of:

25 Pro to Leu at position 85 of SEQ ID NO: 6;
 Val to Leu at position 393 of SEQ ID NO: 7;
 Leu to Ile at position 183 of SEQ ID NO: 8;
 Val to Ile at position 209 of SEQ ID NO: 9.

The variant replicase gene of the coronavirus of the present invention may encode a protein which does not comprise a mutation in nsp-2, nsp-3, nsp-6 or nsp-13.

The variant replicase gene of the coronavirus of the present invention may encode a protein which does not comprise a mutation in nsp10 which corresponds to the threonine to isoleucine mutation caused by a mutation at nucleotide position 12,008 in the gene reported by Ammayappan *et al.* (Arch Virol (2009) 154:495-499).

Ammayappan et al (as above) reports the identification of sequence changes responsible for the attenuation of IBV strain Arkansas DPI. The study identified 17 amino acid changes in a variety of IBV proteins following multiple passages, approx. 100, of the virus in embryonated eggs. It was not investigated whether the attenuated virus (Ark DPI 101) is capable of replicating in the presence of maternally-derived antibodies against the virus *in ovo*, without being pathogenic to the embryo. Given that this virus was produced by multiple passage in SPF embryonated eggs, similar methodology for classical IBV vaccines, it is likely that this virus is pathogenic for embryos. The virus may also be sensitive to maternally-derived antibodies if the hens were vaccinated with a similar serotype.

The variant replicase gene of the coronavirus of the present invention may encode a protein which comprises any combination of one or more amino acid mutations provided in the list above.

The variant replicase gene may encode a protein which comprises the amino acid mutation Pro to Leu at position 85 of SEQ ID NO: 6.

The variant replicase gene may encode a protein which comprises the amino acid mutation Val to Leu at position 393 of SEQ ID NO: 7.

The variant replicase gene may encode a protein which comprises the amino acid mutation Leu to Ile at position 183 of SEQ ID NO: 8.

The variant replicase gene may encode a protein which comprises the amino acid mutation Val to Ile at position 209 of SEQ ID NO: 9.

The variant replicase gene may encode a protein which comprises the amino acid mutations Pro to Leu at position 85 of SEQ ID NO: 6, and Val to Leu at position 393 of SEQ ID NO: 7.

5 The variant replicase gene may encode a protein which comprises the amino acid mutations Pro to Leu at position 85 of SEQ ID NO: 6 Leu to Ile at position 183 of SEQ ID NO: 8.

10 The variant replicase gene may encode a protein which comprises the amino acid mutations Pro to Leu at position 85 of SEQ ID NO: 6 and Val to Ile at position 209 of SEQ ID NO: 9.

15 The variant replicase gene may encode a protein which comprises the amino acid mutations Val to Leu at position 393 of SEQ ID NO: 7 and Leu to Ile at position 183 of SEQ ID NO: 8.

The variant replicase gene may encode a protein which comprises the amino acid mutations Val to Leu at position 393 of SEQ ID NO: 7 and Val to Ile at position 209 of SEQ ID NO: 9.

20 The variant replicase gene may encode a protein which comprises the amino acid mutations Leu to Ile at position 183 of SEQ ID NO: 8 and Val to Ile at position 209 of SEQ ID NO: 9.

25 The variant replicase gene may encode a protein which comprises the amino acid mutations Pro to Leu at position 85 of SEQ ID NO: 6, Val to Leu at position 393 of SEQ ID NO: 7 and Leu to Ile at position 183 of SEQ ID NO: 8.

30 The variant replicase gene may encode a protein which comprises the amino acid mutations Pro to Leu at position 85 of SEQ ID NO: 6 Leu to Ile at position 183 of SEQ ID NO: 8 and Val to Ile at position 209 of SEQ ID NO: 9.

35 The variant replicase gene may encode a protein which comprises the amino acid mutations Pro to Leu at position 85 of SEQ ID NO: 6, Val to Leu at position 393 of SEQ ID NO: 7 and Val to Ile at position 209 of SEQ ID NO: 9.

The variant replicase gene may encode a protein which comprises the amino acid mutations Val to Leu at position 393 of SEQ ID NO: 7, Leu to Ile at position 183 of SEQ ID NO: 8 and Val to Ile at position 209 of SEQ ID NO: 9.

- 5 The variant replicase gene may encode a protein which comprises the amino acid mutations Pro to Leu at position 85 of SEQ ID NO: 6, Val to Leu at position 393 of SEQ ID NO: 7, Leu to Ile at position 183 of SEQ ID NO: 8 and Val to Ile at position 209 of SEQ ID NO: 9.
- 10 The variant replicase gene may also be defined at the nucleotide level.

For example the nucleotide sequence of the variant replicase gene of the coronavirus of the present invention may comprise one or more nucleotide substitutions within the regions selected from the list of: 11884-12318, 16938-18500, 18501-19514 and 19515-15 20423 of SEQ ID NO:1.

For example the nucleotide sequence of the variant replicase gene of the coronavirus of the present invention may comprise one or more nucleotide substitutions selected from the list of:

- 20 C to T at nucleotide position 12137;
- G to C at nucleotide position 18114;
- T to A at nucleotide position 19047; and
- G to A at nucleotide position 20139;

compared to the sequence shown as SEQ ID NO: 1.

25 As used herein, the term "substitution" is synonymous with the term mutation and means that the nucleotide at the identified position differs to that of the wild-type nucleotide sequence.

30 The nucleotide sequence may comprise any combination of the nucleotide substitutions selected from the list of:

- C to T at nucleotide position 12137;
- G to C at nucleotide position 18114;
- T to A at nucleotide position 19047; and
- G to A at nucleotide position 20139;

35 compared to the sequence shown as SEQ ID NO: 1.

The nucleotide sequence may comprise the substitution C12137T.

The nucleotide sequence may comprise substitution G18114C.

5 The nucleotide sequence may comprise the substitution T19047A.

The nucleotide sequence may comprise the substitution G20139A.

The nucleotide sequence may comprise the substitutions C12137T and G18114C.

10

The nucleotide sequence may comprise the substitutions C12137T and T19047A.

The nucleotide sequence may comprise the substitutions C12137T and G20139A.

15

The nucleotide sequence may comprise the substitutions G18114C and T19047A.

The nucleotide sequence may comprise the substitutions G18114C and G20139A.

The nucleotide sequence may comprise the substitutions T19047A and G20139A.

20

The nucleotide sequence may comprise the substitutions C12137T, G18114C and T19047A.

25 The nucleotide sequence may comprise the substitutions C12137T, T19047A and G20139A.

The nucleotide sequence may comprise the substitutions C12137T, G18114C and G20139A.

30 The nucleotide sequence may comprise the substitutions G18114C, T19047A and G20139A.

The nucleotide sequence may comprise the substitutions C12137T, G18114C, T19047A and G20139A.

35

The nucleotide sequence may not comprise a substitution which corresponds to the C12008T substitution reported by Ammayappan *et al.* (as above).

The nucleotide sequence may be natural, synthetic or recombinant. It may be double or single stranded, it may be DNA or RNA or combinations thereof. It may, for example, be cDNA, PCR product, genomic sequence or mRNA.

5

The nucleotide sequence may be codon optimised for production in the host/host cell of choice.

It may be isolated, or as part of a plasmid, virus or host cell.

10

PLASMID

A plasmid is an extra-chromosomal DNA molecule separate from the chromosomal DNA which is capable of replicating independently of the chromosomal DNA. They are usually 15 circular and double-stranded.

Plasmids, or vectors (as they are sometimes known), may be used to express a protein in a host cell. For example a bacterial host cell may be transfected with a plasmid capable of encoding a particular protein, in order to express that protein. The term also includes 20 yeast artificial chromosomes and bacterial artificial chromosomes which are capable of accommodating longer portions of DNA.

The plasmid of the present invention comprises a nucleotide sequence capable of encoding a defined region of the replicase protein. It may also comprise one or more 25 additional coronavirus nucleotide sequence(s), or nucleotide sequence(s) capable of encoding one or more other coronavirus proteins such as the S gene and/or gene 3.

The plasmid may also comprise a resistance marker, such as the guanine xanthine phosphoribosyltransferase gene (gpt) from *Escherichia coli*, which confers resistance to 30 mycophenolic acid (MPA) in the presence of xanthine and hypoxanthine and is controlled by the vaccinia virus P7.5 early/late promoter.

RECOMBINANT VACCINIA VIRUS

35 The present invention also relates to a recombinant vaccinia virus (rVV) comprising a variant replicase gene as defined herein.

The recombinant vaccinia virus (rVV) may be made using a vaccinia-virus based reverse genetics system.

In this respect, the present invention also provides a method for making a viral particle by:

- 5 (i) transfected a plasmid as described in the previous section into a host cell;
- (ii) infecting the host cell with a recombining virus comprising the genome of a coronavirus strain with a replicase gene;
- 10 (iii) allowing homologous recombination to occur between the replicase gene sequences in the plasmid and the corresponding sequences in the recombining virus genome to produce a modified replicase gene;
- (iv) selecting for recombining virus comprising the modified replicase gene.

The term 'modified replicase gene' refers to a replicase gene which comprises a variant replicase gene as described in connection with the first aspect of the present invention.

15 Specifically, the term refers to a gene which is derived from a wild-type replicase gene but comprises a nucleotide sequence which causes it to encode a variant replicase protein as defined herein.

20 The recombination may involve all or part of the replicase gene. For example the recombination may involve a nucleotide sequence encoding for any combination of nsp-10, nsp-14, nsp-15 and/or nsp-16. The recombination may involve a nucleotide sequence which encodes for an amino acid mutation or comprises a nucleotide substitution as defined above.

25 The genome of the coronavirus strain may lack the part of the replicase protein corresponding to the part provided by the plasmid, so that a modified protein is formed through insertion of the nucleotide sequence provided by the plasmid.

30 The recombining virus is one suitable to allow homologous recombination between its genome and the plasmid. The vaccinia virus is particularly suitable as homologous recombination is routinely used to insert and delete sequences for the vaccinia virus genome.

The above method optionally includes the step:

35 (v) recovery of recombinant coronavirus comprising the modified replicase gene from the DNA from the recombining virus from step (iv).

Methods for recovering recombinant coronavirus, such as recombinant IBV, are known in the art (See Britton et al (2005) see page 24; and PCT/GB2010/001293).

For example, the DNA from the recombining virus from step (iv) may be inserted into a 5 plasmid and used to transfect cells which express cytoplasmic T7 RNA polymerase. The cells may, for example be pre-infected with a fowlpox virus expressing T7 RNA polymerase. Recombinant coronavirus may then be isolated, for example, from the growth medium.

10 When the plasmid is inserted into the vaccinia virus genome, an unstable intermediate is formed. Recombinants comprising the plasmid may be selected for e.g. using a resistance marker on the plasmid.

15 Positive recombinants may then be verified to contain the modified replicase gene by, for example, PCR and sequencing.

Large stocks of the recombining virus including the modified replicase gene (e.g. recombinant vaccinia virus, (rVV) may be grown up and the DNA extracted in order to carry out step (v)).

20 Suitable reverse genetics systems are known in the art (Casais et al (2001) J. Virol 75:12359-12369; Casais et al (2003) J. Virol. 77:9084-9089; Britton et al (2005) J. Virological Methods 123:203-211; Armesto et al (2008) Methods in Molecular Biology 454:255-273).

25 **CELL**

The coronavirus may be used to infect a cell.

30 Coronavirus particles may be harvested, for example from the supernatant, by methods known in the art, and optionally purified.

The cell may be used to produce the coronavirus particle.

35 Thus the present invention also provides a method for producing a coronavirus which comprises the following steps:

(i) infection of a cell with a coronavirus according to the invention;

- (ii) allowing the virus to replicate in the cell; and
- (iii) harvesting the progeny virus.

5 The present invention also provides a cell capable of producing a coronavirus according to the invention using a reverse genetics system. For example, the cell may comprise a recombining virus genome comprising a nucleotide sequence capable of encoding the replicase gene of the present invention.

10 The cell may be able to produce recombinant recombining virus (e.g. vaccinia virus) containing the replicase gene.

15 Alternatively the cell may be capable of producing recombinant coronavirus by a reverse genetics system. The cell may express or be induced to express T7 polymerase in order to rescue the recombinant viral particle.

15

VACCINE

20 The coronavirus may be used to produce a vaccine. The vaccine may be a live attenuated form of the coronavirus of the present invention and may further comprise a pharmaceutically acceptable carrier. As defined herein, "pharmaceutically acceptable carriers" suitable for use in the invention are well known to those of skill in the art. Such carriers include, without limitation, water, saline, buffered saline, phosphate buffer, alcohol/aqueous solutions, emulsions or suspensions. Other conventionally employed diluents and excipients may be added in accordance with conventional techniques. Such 25 carriers can include ethanol, polyols, and suitable mixtures thereof, vegetable oils, and injectable organic esters. Buffers and pH adjusting agents may also be employed. Buffers include, without limitation, salts prepared from an organic acid or base. Representative buffers include, without limitation, organic acid salts, such as salts of citric acid, e.g., citrates, ascorbic acid, gluconic acid, histidine-HCl, carbonic acid, tartaric acid, 30 succinic acid, acetic acid, or phthalic acid, Tris, trimethanamine hydrochloride, or phosphate buffers. Parenteral carriers can include sodium chloride solution, Ringer's dextrose, dextrose, trehalose, sucrose, and sodium chloride, lactated Ringer's or fixed oils. Intravenous carriers can include fluid and nutrient replenishers, electrolyte replenishers, such as those based on Ringer's dextrose and the like. Preservatives and 35 other additives such as, for example, antimicrobials, antioxidants, chelating agents (e.g., EDTA), inert gases and the like may also be provided in the pharmaceutical carriers. The present invention is not limited by the selection of the carrier. The preparation of these

pharmaceutically acceptable compositions, from the above-described components, having appropriate pH isotonicity, stability and other conventional characteristics is within the skill of the art. See, e.g., texts such as Remington: The Science and Practice of Pharmacy, 20th ed, Lippincott Williams & Wilkins, publ., 2000; and The Handbook of Pharmaceutical Excipients, 4.sup.th edit., eds. R. C. Rowe et al, APhA Publications, 2003.

The vaccine of the invention will be administered in a "therapeutically effective amount", which refers to an amount of an active ingredient, e.g., an agent according to the invention, sufficient to effect beneficial or desired results when administered to a subject or patient. An effective amount can be administered in one or more administrations, applications or dosages. A therapeutically effective amount of a composition according to the invention may be readily determined by one of ordinary skill in the art. In the context of this invention, a "therapeutically effective amount" is one that produces an objectively measured change in one or more parameters associated Infectious Bronchitis condition sufficient to effect beneficial or desired results .An effective amount can be administered in one or more administrations. For purposes of this invention, an effective amount of drug, compound, or pharmaceutical composition is an amount sufficient to reduce the incidence of Infectious Bronchitis. As used herein, the term "therapeutic" encompasses the full spectrum of treatments for a disease, condition or disorder. A "therapeutic" agent of the invention may act in a manner that is prophylactic or preventive, including those that incorporate procedures designed to target animals that can be identified as being at risk (pharmacogenetics); or in a manner that is ameliorative or curative in nature; or may act to slow the rate or extent of the progression of at least one symptom of a disease or disorder being treated.

The present invention also relates to a method for producing such a vaccine which comprises the step of infecting cells, for example Vero cells, with a viral particle comprising a replicase protein as defined in connection with the first aspect of the invention.

VACCINATION METHOD

35 The coronavirus of the present invention may be used to treat and/or prevent a disease.

To "treat" means to administer the vaccine to a subject having an existing disease in order to lessen, reduce or improve at least one symptom associated with the disease and/or to slow down, reduce or block the progression of the disease.

5 To "prevent" means to administer the vaccine to a subject who has not yet contracted the disease and/or who is not showing any symptoms of the disease to prevent or impair the cause of the disease (e.g. infection) or to reduce or prevent development of at least one symptom associated with the disease.

10 The disease may be any disease caused by a coronavirus, such as a respiratory disease and and/or gastroenteritis in humans and hepatitis, gastroenteritis, encephalitis, or a respiratory disease in other animals.

15 The disease may be infectious bronchitis (IB); Porcine epidemic diarrhoea; Transmissible gastroenteritis; Mouse hepatitis virus; Porcine haemagglutinating encephalomyelitis; Severe acute respiratory syndrome (SARS); or Bluecomb disease.

The disease may be infectious bronchitis.

20 The vaccine may be administered to hatched chicks or chickens, for example by eye drop or intranasal administration. Although accurate, these methods can be expensive e.g. for large broiler flocks. Alternatives include spray inoculation of administration to drinking water but it can be difficult to ensure uniform vaccine application using such methods.

25 The vaccine may be provided in a form suitable for its administration, such as an eye-dropper for intra-ocular use.

30 The vaccine may be administered by *in ovo* inoculation, for example by injection of embryonated eggs. *In ovo* vaccination has the advantage that it provides an early stage resistance to the disease. It also facilitates the administration of a uniform dose per subject, unlike spray inoculation and administration via drinking water.

35 The vaccine may be administered to any suitable compartment of the egg, including allantoic fluid, yolk sac, amnion, air cell or embryo. It may be administered below the shell (aircell) membrane and chorioallantoic membrane.

Usually the vaccine is injected into embryonated eggs during late stages of embryonic development, generally during the final quarter of the incubation period, such as 3-4 days prior to hatch. In chickens, the vaccine may be administered between day 15-19 of the 21-day incubation period, for example at day 17 or 18.

5

The process can be automated using a robotic injection process, such as those described in WO 2004/078203.

The vaccine may be administered together with one or more other vaccines, for example, 10 vaccines for other diseases, such as Newcastle disease virus (NDV). The present invention also provides a vaccine composition comprising a vaccine according to the invention together with one or more other vaccine(s). The present invention also provides a kit comprising a vaccine according to the invention together with one or more other vaccine(s) for separate, sequential or simultaneous administration.

15

The vaccine or vaccine composition of the invention may be used to treat a human, animal or avian subject. For example, the subject may be a chick, chicken or mouse (such as a laboratory mouse, e.g. transgenic mouse).

20 Typically, a physician or veterinarian will determine the actual dosage which will be most suitable for an individual subject or group of subjects and it will vary with the age, weight and response of the particular subject(s).

25 The composition may optionally comprise a pharmaceutically acceptable carrier, diluent, excipient or adjuvant. The choice of pharmaceutical carrier, excipient or diluent can be selected with regard to the intended route of administration and standard pharmaceutical practice. The pharmaceutical compositions may comprise as (or in addition to) the carrier, excipient or diluent, any suitable binder(s), lubricant(s), suspending agent(s), coating agent(s), solubilising agent(s), and other carrier agents that may aid or increase 30 the delivery or immunogenicity of the virus.

The invention will now be further described by way of Examples, which are meant to serve to assist one of ordinary skill in the art in carrying out the invention and are not intended in any way to limit the scope of the invention.

35

EXAMPLES

EXAMPLE 1 – Generation of an IBV reverse genetics system based on M41-CK

A M41-CK full-length cDNA was produced by replacement of the Beaudette cDNA in the Vaccinia virus reverse genetics system previously described in PCT/GB2010/001293 (herein incorporated by reference) with synthetic cDNA derived from the M41 consensus sequence.

The IBV cDNA within recombinant Vaccinia virus (rVV) rVV-BeauR-Rep-M41 structure described in Armesto, Cavanagh and Britton (2009). PLoS ONE 4(10): e7384. doi:10.1371/journal.pone.0007384, which consisted of the replicase derived from IBV Beaudette strain and the structural and accessory genes and 3' UTR from IBV M41-CK, was further modified by replacement of the Beaudette 5' UTR-Nsp2-Nsp3 sequence with the corresponding sequence from IBV M41-CK. The resulting IBV cDNA consisted of 5' UTR-Nsp2-Nsp3 from M41, Nsp4-Nsp16 from Beaudette and the structural and accessory genes and 3' UTR from M41. This cDNA was further modified by the deletion of the Beaudette Nsp4-Nsp16 sequence. The resulting cDNA, lacking Nsp4-16, was modified in four further steps in which the deleted Nsps were sequentially replaced with the corresponding sequences from M41-CK, the replacement cDNAs represented M41-CK Nsp4-8, Nsp9-12, Nsp12-14 and finally Nsp15-16. Each replacement cDNA contained approx. 500 nucleotides at the 5' end corresponding to the 3' most M41 sequence previously inserted and approx. 500 nucleotides at the 3' end corresponding to the M41 S gene sequence. This allowed insertion of the M41 cDNA sequence by homologous recombination and sequential addition of contiguous M41 replicase gene sequence. The synthetic cDNAs containing the M41-derived Nsp sequences were added by homologous recombination utilising the inventor's previous described transient dominant selection (TDS) system (see PCT/GB2010/001293). The M41-derived cDNAs containing sequence corresponding to the M41 Nsps-10, -14, -15 and -16 contained the modified amino acids at positions 85, 393, 183 and 209, respectively, as indicated in Figure 10.

A full-length cDNA representing the genome of M41-CK was generated in Vaccinia virus representing the synthetic sequences. Two rIBVs, M41-R-6 and M41-R-12, were rescued and shown to grow in a similar manner as M41-CK (Fig. 1).

EXAMPLE 2 – Determining the pathogenicity of rescued M41 viruses

The viruses rescued in Example 1 were used to infect 8-day-old specific pathogen free (SPF) chicks by ocular and nasal inoculation to test them for pathogenicity, as observed

by clinical signs on a daily basis 3-7 days post-infection and for ciliary activity days 4 and 6 post-infection. Loss of ciliary activity is a well-established method for determining the pathogenicity of IBV. The two M41-R viruses were found to be apathogenic when compared to M41-CK though they did show some clinical signs in comparison to 5 uninfected control chicks (Fig. 2) and some but inconsistent loss in ciliary activity (Fig. 3).

Thus, the M41-R molecular clones of M41-CK were not pathogenic when compared to the parental virus M41-CK.

10 The inventors identified several nucleotide differences in the M41-R compared to the M41-CK sequences. The majority of these were synonymous mutations, as the nucleotide change did not affect the amino acid sequence of the protein associated with the sequence. However, four non-synonymous mutations were identified in the IBV replicase gene specific to Nsp-10, Nsp-14, Nsp-15 and Nsp-16 components of the 15 replicase gene, these mutations resulted in amino acid changes (Table 3).

Table 3. Non-Synonymous mutations identified in the Nsps of M41-R full-length genome

Region of Replicase	Nucleotide position	Nucleotide Mutation	Amino Acid Change
Nsp10	12137	C→T	Pro→Leu
Nsp14	18114	G→C	Val→Leu
Nsp15	19047	T→A	Leu→Ile
Nsp16	20139	G→A	Val→Ile

20

EXAMPLE 3 – Repair of M41-R rIBVs

In order to determine whether the identified mutations were responsible for the loss of 25 pathogenicity associated with M41-R, the Nsp10 mutation was repaired and the mutations in Nsp-14, -15 & -16 were repaired and shown to grow in a similar manner as M41-CK (Fig 9). The inventors thus generated the rIBVs, M41R-nsp10rep and M41R-nsp14, 15, 16rep, using synthetic cDNAs containing the correct nucleotides utilising the inventor's previous described (TDS) system (see PCT/GB2010/001293).

30

The rIBVs were assessed for pathogenicity in chicks as described previously. Both rIBVs showed increased pathogenicity when compared to M41-R but not to the level observed with M41-CK (Figs 4 and 5). M41R-nsp14, 15, 16rep gave more clinical signs and more reduction in ciliary activity than M41R-nsp10rep, overall these results indicated that the 5 changes associated with the four Nsp appear to affect pathogenicity.

To determine the roles of the Nsp in pathogenicity the full-length cDNA corresponding to M41R-nsp10rep was used to repair the mutations in Nsp14, 15 & 16 using a synthetic cDNA containing the correct nucleotides utilising the TDS system.

10

The following rIBVs were produced:-

M41R-nsp10, 15rep – M41-R with the mutations in Nsp-10 and Nsp-15 repaired

M41R-nsp10, 14, 15rep – M41-R with mutations in Nsp-10, -14 and -15 repaired

M41R-nsp10, 14, 16rep – M41-R with mutations in Nsp-10, -14 and -16 repaired

15

M41R-nsp10, 15, 16rep – M41-R with mutations in Nsp-10, -15 and -16 repaired

M41-K - All four mutations, Nsp-10,-14,-15 & -16 repaired in M41-R

The rIBVs were shown to grow in a similar manner as M41-CK (Fig 9) and assessed for pathogenicity as described previously. M41-K (in which all four mutations had been 20 repaired) resulted in clinical signs and 100% loss of ciliary activity (complete ciliostasis) by 4 days post-infection (Fig. 6, 7 & 8). The other rIBVs demonstrated varying levels of pathogenicity, apart from M41R-nsp10, 15, 16rep, which was essentially apathogenic. These results confirmed that repair of all four Nsp restored pathogenicity to M41-R; again supporting the previous evidence that the mutations described in the four Nsp are 25 implicated in attenuating M41-CK.

The inventors also generated rIBV M41R-nsp 10, 14 rep (nsp 10 and 14 are repaired, nsp 15 and 16 contain mutations) and rIBV M41R-nsp 10, 16 rep (nsp 10 and 16 are repaired, nsp 14 and 15 contain mutations) and assessed the pathogenicity of these viruses.

30

rIBV M41R-nsp 10, 14 rep less pathogenic than M41-K but caused around 50% ciliostasis on days 4-6 post-infection. rIBV M41R-nsp 10, 16 rep was almost apathogenic and caused no ciliostasis (see Figure 11a-c).

35 Thus the genome associated with M41-R is a potential backbone genome for a rationally attenuated IBV.

EXAMPLE 4 – Vaccination/Challenge Study with M41-R

Candidate vaccine viruses were tested in studies in which fertilized chicken eggs were vaccinated *in ovo* at 18 days embryonation and in which the hatchability of the inoculated eggs was determined. The clinical health of the chickens was investigated and the 5 chickens were challenged at 21 days of age with a virulent IB M41 challenge virus at $10^{3.65}$ EID₅₀ per dose.

Clinical signs were investigated after challenge protection by the vaccine and a ciliostasis test was performed at 5 days after challenge to investigate the effect of the challenge 10 viruses on movement of the cilia and protection by the vaccine against ciliostasis (inhibition of cilia movement).

In ovo vaccination in commercial broiler eggs

15 The design of the experiment is given in Table 4 and the clinical results are given in Table 5. Hatchability of the eggs inoculated with IB M41-R was good and chickens were healthy. IB M41-R protected against clinical signs after challenge in the broilers (placebo: 19/19 affected, IB M41-R: 3/18 affected and 1 dead). The results of the ciliostasis test are given in Table 6. IB M41-R generated protection against ciliostasis.

20

Table 4 - Design of a hatchability, safety, efficacy study in commercial eggs

Treatment	Treatment Description	EID ₅₀ ¹ per dose	Route of Admin	Day(s) of Admin	Day(s) of Challenge ²	End of Study	Nr. of eggs per treatment
T01	None	NA	NA	NA	NA	NA	30
T02	IB M41-R	10^2	In ovo	18 days embryonation	At 21 days of age, 20 chickens per group	At 26 days of age	30
NTX	Saline	NA	In ovo				30

¹ Dose volume 0.1 ml, NA, not applicable.

² $10^{3.65}$ EID₅₀ per dose.

Table 5 - Hatch percentages and clinical data before and after challenge in commercial chickens, for design see Table 1.

Treatment	Hatch/ total	Vital/ total	Before challenge		After challenge	
			Deaths/ total	Symptoms/ total	Deaths/ total	Symptoms/ total
None	28/30		Euthanized directly after hatch for blood collection			
IB M41-R	28/30	28/28	1/20	0/19	1/19	3/18 ^{1,2}
Saline	29/30	29/29	1/20	0/19	0/19	19/19 ^{1,2,3,4,5,6,7}

5 ¹ Disturbed respiratory system

² Whizzing

³ Change of voice

⁴ Breathing difficult

⁵ Swollen intra-orbital sinuses

10 ⁶ Uneven growth

⁷ Weak

Table 6 - Results of the ciliostasis test after challenge, for design see Table 1.

Treatment	Protected/total	Percentage protection
Saline	0/19	0%
IB M41R	5/18	28%

15 *In ovo vaccination in specific pathogen-free (SPF) eggs*

The design of the study in SPF eggs is given in Table 7 and is similar with the design of the studies with commercial broilers, but the vaccination dose for IB M41-R was higher, (10^5 EID₅₀ per dose).

20 The results (Table 8) show that the hatch percentage for IB M41-R hatch was low, and 19 of 40 hatched and the chicks were weak. Eight chicks died. The remaining 11 chickens were challenged and 11 of the chicks hatched from the eggs which had been inoculated with saline were challenged.

25 In the ciliostasis test after challenge it appeared that all chickens vaccinated *in ovo* with IB M41-R were protected, whereas none of the controls was protected, see Table 9.

Table 7. Design of a hatchability, safety, efficacy study in SPF eggs

Treatment	Treatment Description	EID ₅₀ ¹ per dose	Route of Admin	Day of Admin	Days of Challenge ²	End of Study	Nr. of eggs per treatment
T01	IB M41-R	10 ⁵	In ovo	18 days embryo-nation	At 21 days of age	At 26 days of age	40
T04	Saline	NA	In ovo				40
NTX	NA	NA	NA		NA		10

¹ Dose volume 0.1 ml, NA, not applicable.² Challenge dose 10^{3.3} EID₅₀ in 0.2 ml.

5

Table 8. Hatch percentages and clinical data before and after challenge in SPF chickens, for design see Table 7.

Treatment	Hatch/total	Vital/total	Before challenge		After challenge	
			Deaths/total	Symptoms/total	Deaths/total	Symptoms/total
IB M41-R	19/40	11/40	8/40	weak	0	0
Saline	30/40	30/40	0	-	0	0
NA	9/10	9/10	0	-	-	-

10 Table 9. Results of the ciliostasis test after challenge, for design see Table 7.

Treatment	Protected/total	Percentage protection
Saline	0/11	0%
IB M41R	11/11	100%

In conclusion, IB M41-R was safe in commercial eggs, generated protection against clinical signs and to an extent against ciliostasis.

15

In SPF eggs vaccinated with IB M41 R a relatively low number of chickens hatched. This may be due to the 10⁵ EID₅₀ per egg of IB M41-R used. This was 10-fold higher than the dose used in earlier studies in which there was a higher level of hatchability. The lower hatch percentages may also be caused by a particularly high susceptibility of the batch of 20 SPF eggs for viruses, as in other studies the level of embryo mortality was also higher than had previously been observed.

25 After challenge all surviving chickens after hatch were completely protected against ciliostasis. It is concluded that IB M41-R has great potential as vaccine to be administered *in ovo*.

All publications mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described methods and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology, virology or related fields are intended to be within the scope of the following claims.

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as, an acknowledgement or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.

The claims defining the invention are as follows:

1. A live, attenuated infectious bronchitis virus (IBV) comprising a variant replicase gene encoding polyproteins comprising a mutation in one or more of the non-structural protein(s) (nsp)-10 and nsp-14; wherein the variant replicase gene encodes a protein comprising an amino acid mutation of Pro to Leu at the position corresponding to position 85 of SEQ ID NO: 6, and/or wherein the variant replicase gene encodes a protein comprising an amino acid mutation of Val to Leu at the position corresponding to position 393 of SEQ ID NO: 7.
2. The IBV according to claim 1, further comprising a mutation in one or more of nsp-15 corresponding to SEQ ID NO: 8 and nsp-16 corresponding to SEQ ID NO: 9.
3. The IBV according to claim 2, wherein the variant replicase gene encodes a protein comprising one or more amino acid mutations selected from: an amino acid mutation of Leu to Ile at the position corresponding to position 183 of SEQ ID NO: 8 and an amino acid mutation of Val to Ile at the position corresponding to position 209 of SEQ ID NO: 9.
4. The IBV according to any one of the preceding claims, wherein the variant replicase gene encodes a protein comprising the amino acid mutation Pro to Leu at the position corresponding to position 85 of SEQ ID NO: 6.
5. The IBV according to any one of the preceding claims, wherein the variant replicase gene encodes a protein comprising the amino acid mutations Val to Leu at the position corresponding to position 393 of SEQ ID NO: 7; Leu to Ile at the position corresponding to position 183 of SEQ ID NO: 8; and Val to Ile at the position corresponding to position 209 of SEQ ID NO: 9.
6. The IBV according to any one of the preceding claims, wherein the variant replicase gene encodes a protein comprising the amino acid mutations: Pro to Leu at the position corresponding to position 85 of SEQ ID NO: 6; Val to Leu at the position corresponding to position 393 of SEQ ID NO: 7; Leu to Ile at the position corresponding to position 183 of SEQ ID NO: 8; and Val to Ile at the position corresponding to position 209 of SEQ ID NO: 9.
7. The IBV according to any one of the preceding claims, wherein the variant replicase gene comprises one or more nucleotide substitutions selected from:
C to T at nucleotide position 12137;

G to C at nucleotide position 18114;
T to A at nucleotide position 19047; and
G to A at nucleotide position 20139;
compared to the sequence shown as SEQ ID NO: 1.

8. The IBV according to any one of the preceding claims which is IBV M41.
9. The IBV according to claim 8, which comprises an S protein, at least part of which is from an IBV serotype other than M41.
10. The IBV according to claim 9, wherein the S1 subunit is from an IBV serotype other than M41.
11. The IBV according to claim 9, wherein the S protein is from an IBV serotype other than M41.
12. The IBV according to any one of the preceding claims which has reduced pathogenicity compared to a coronavirus expressing a corresponding wild-type replicase, such that when the virus is administered to an embryonated egg, it is capable of replicating without being pathogenic to the embryo.
13. A variant replicase gene as defined in any one of claims 1 to 7.
14. A protein encoded by a variant replicase gene according to claim 13.
15. A plasmid comprising a variant replicase gene according to claim 13.
16. A method for making the IBV according to any one of claims 1 to 12 which comprises the following steps:
 - (i) transfecting a plasmid according to claim 15 into a host cell;
 - (ii) infecting the host cell with a recombining virus comprising the genome of a coronavirus strain with a replicase gene;
 - (iii) allowing homologous recombination to occur between the replicase gene sequences in the plasmid and the corresponding sequences in the recombining virus genome to produce a modified replicase gene; and
 - (iv) selecting for recombining virus comprising the modified replicase gene.

17. The method according to claim 16, wherein the recombining virus is a vaccinia virus.
18. The method according to claim 16 or 17 which also includes the step:
 - (v) recovering recombinant coronavirus comprising the modified replicase gene from the DNA from the recombining virus from step (iv).
19. A cell capable of producing an IBV according to any one of claims 1 to 12.
20. A vaccine comprising an IBV according to any one of claims 1 to 12 and a pharmaceutically acceptable carrier.
21. A method for treating and/or preventing a disease in a subject which comprises the step of administering a vaccine according to claim 20 to the subject.
22. The use of an IBV according to any one of claims 1 to 12 in the manufacture of a vaccine for treating and/or preventing a disease in a subject.
23. The method or use according to claim 21 or 22 wherein the disease is infectious bronchitis (IB).
24. The method according to claim 21 wherein the method of administration is selected from the group consisting of: eye drop administration, intranasal administration, drinking water administration, post-hatch injection and *in ovo* injection.
25. The method according to claim 24 wherein the vaccination is *in ovo* vaccination.
26. A method for producing a vaccine according to claim 20, which comprises the step of infecting a cell according to claim 19 with an IBV according to any one of claims 1 to 12.

FIGURE 1




FIGURE 2





FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8





FIGURE 9

A



B




FIGURE 10

Nep10

8	20	8	40	8	60	8	80
M41 Nep10: SEGHETEEVIAVGILSLCFIAVTPADTYCKYVRAAEKQPLNCVBNMLTVHNGSGFAITSKPSFTPIQDSYGGASVCLNCRA							
Mod Nep10: SEGHETEEVIAVGILSLCFIAVTPADTYCKYVRAAEKQPLNCVBNMLTVHNGSGFAITSKPSFTPIQDSYGGASVCLNCRA							
8	100	8	120	8	140	8	80
M41 Nep10: SHAFEGGAGNLDGRCQFXGSPFWQIPTTEKDPWGFCLRNWVCTWICWIGCQCDSLRQPKPSVQ							
Mod Nep10: SHAFEGGAGNLDGRCQFXGSPFWQIPTTEKDPWGFCLRNWVCTWICWIGCQCDSLRQPKPSVQ							

Nep14

8	20	8	40	8	60	8	80
M41 Nep14: GTGLEKICNREFSGVNPAYAVITKALARYKVNDEELRALKVNEVEGSEITVNLHLSLIGEKSIVNVEGCHNNEITRDAIP							
Mod Nep14: GTGLEKICNREFSGVNPAYAVITKALARYKVNDEELRALKVNEVEGSEITVNLHLSLIGEKSIVNVEGCHNNEITRDAIP							
8	100	8	120	8	140	8	160
M41 Nep14: NVRGAWVGFIDVETATHACGTNIGTKLPFQNGFSTGAIFVVTPEGLVDTSIGNSFEVRSKAPPGEOFNLHRLFKAAPWY							
Mod Nep14: NVRGAWVGFIDVETATHACGTNIGTKLPFQNGFSTGAIFVVTPEGLVDTSIGNSFEVRSKAPPGEOFNLHRLFKAAPWY							
8	180	8	200	8	220	8	240
M41 Nep14: VRERIVVQLAUNLQVSDCYYFVTCNGLELTTLKTFVKGIDQVSCGGSGATTTFNSHTQAYACSHHCLGFDFVYNPLLV							
Mod Nep14: VRERIVVQLAUNLQVSDCYYFVTCNGLELTTLKTFVKGIDQVSCGGSGATTTFNSHTQAYACSHHCLGFDFVYNPLLV							
8	260	8	280	8	300	8	320
M41 Nep14: DIXXWTSQNLQFNRBLHQNVRGHAYVASARAIIMRCIARNAPCDVWNLITYPHIANEDEVNSSSCRYLGRMILNACVD							
Mod Nep14: DIXXWTSQNLQFNRBLHQNVRGHAYVASARAIIMRCIARNAPCDVWNLITYPHIANEDEVNSSSCRYLGRMILNACVD							
8	340	8	360	8	380	8	400
M41 Nep14: AIKVNWVYDIGNPKGIMCVENGLNFREYIDNPTVPNVKQFEYDYNQKHNIDFAGLCPWNCVBCYPDWSIICRYDTRR							
Mod Nep14: AIKVNWVYDIGNPKGIMCVENGLNFREYIDNPTVPNVKQFEYDYNQKHNIDFAGLCPWNCVBCYPDWSIICRYDTRR							
8	420	8	440	8	460	8	480
M41 Nep14: LSVFNLPGCNGESLTVNKHAFNTPKTFDASFPAKLEMPFFTYESSFCETIQLDGVAQDLVSLAKDCITRCNIGGAVCRK							
Mod Nep14: LSVFNLPGCNGESLTVNKHAFNTPKTFDASFPAKLEMPFFTYESSFCETIQLDGVAQDLVSLAKDCITRCNIGGAVCRK							
8	500	8	520	8	540	8	560
M41 Nep14: HAQMYADFVTSYNAVAVTAGFTFWRVTRNENPYNLNEKFSALQ							
Mod Nep14: HAQMYADFVTSYNAVAVTAGFTFWRVTRNENPYNLNEKFSALQ							

Nsp15


	20	40	60	80	
M41 Nsp15:	SIDWIAAYNNYKGGHYDIAJAGEMPTIVGKXVFVIDQXVEKAVFFNQYIILPTSVAFELYAKRNSIRTLPNRRIILNGLQVNT				
Mod Nsp15:	SIDWIAAYNNYKGGHYDIAJAGEMPTIVGKXVFVIDQXVEKAVFFNQYIILPTSVAFELYAKRNSIRTLPNRRIILNGLQVNT				
	80	100	120	140	160
M41 Nsp15:	NGFVIVIHYTSDTPLYKNTVVECAYTSDIEPGLIVLYDDYGDYOSFLAIDNAVLVSTQCYEBSYVEIPSNLIVQNGIPL				
Mod Nsp15:	NGFVIVIHYTSDTPLYKNTVVECAYTSDIEPGLIVLYDDYGDYOSFLAIDNAVLVSTQCYEBSYVEIPSNLIVQNGIPL				
	160	180	200	220	240
M41 Nsp15:	KIGCANLYVYKRVVGRAPFTLPTNITQGRSYTFFPDSVEEDFLIMSEESFVENTYGEKELCLOHILNGEVNKPQLGGLRTV				
Mod Nsp15:	KIGCANLYVYKRVVGRAPFTLPTNITQGRSYTFFPDSVEEDFLIMSEESFVENTYGEKELCLOHILNGEVNKPQLGGLRTV				
	240	260	280	300	320
M41 Nsp15:	IGKCRILRANKLKAQSVWNSDPMKRFVLAIDNGSGKGVCTVWILLINPFLILRNLKEYGTRNSKVVIVSIDHHSIN				
Mod Nsp15:	IGKCRILRANKLKAQSVWNSDPMKRFVLAIDNGSGKGVCTVWILLINPFLILRNLKEYGTRNSKVVIVSIDHHSIN				
	320	340	360	380	400
M41 Nsp15:	FMKWFEDGILIRTCYPQNG				
Mod Nsp15:	FMKWFEDGILIRTCYPQNG				

Nsp16


	20	40	60	80	
M41 Nsp16:	SAYTCGYKMPFLYKVONCVMEPCRPINPCVGVGIALPSCGIMSVVACYTOOLCQYLSNTTMCVPHNNGKHFQGSIDGVAPGS				
Mod Nsp16:	SAYTCGYKMPFLYKVONCVMEPCRPINPCVGVGIALPSCGIMSVVACYTOOLCQYLSNTTMCVPHNNGKHFQGSIDGVAPGS				
	80	100	120	140	160
M41 Nsp16:	TVLQWNLPEGTLLVNTSIVDVSIDHVSVLSDCIRKTEKPDVLVISIIMYTDNCKNGREGVIANNGNDEVFTYLSSFLR				
Mod Nsp16:	TVLQWNLPEGTLLVNTSIVDVSIDHVSVLSDCIRKTEKPDVLVISIIMYTDNCKNGREGVIANNGNDEVFTYLSSFLR				
	160	180	200	220	240
M41 Nsp16:	KKRLAIGGCFAYVVIETISWREVLVYIILAQDCANWIFCTAVNASSSEAFI				
Mod Nsp16:	KKRLAIGGCFAYVVIETISWREVLVYIILAQDCANWIFCTAVNASSSEAFI				
	240	260	280	300	320
M41 Nsp16:	QTSAYSIIFWVANVTDLKLKATPVVSLKTEQYHLYVFNLIJKCIGKLLVRIWNGNTSFTNISFVCTM				
Mod Nsp16:	QTSAYSIIFWVANVTDLKLKATPVVSLKTEQYHLYVFNLIJKCIGKLLVRIWNGNTSFTNISFVCTM				

FIGURE 11

A)



B)



C)

