

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199957073 B2
(10) Patent No. 771249

(54) Title
Method for purification and manipulation of nucleic acids using paramagnetic particles

(51)⁶ International Patent Classification(s)
C01G 049/02 C01G 049/12

(21) Application No: **199957073** (22) Application Date: **1999.10.27**

(30) Priority Data

(31) Number (32) Date (33) Country
09/183127 1998.10.30 US

(43) Publication Date : **2000.06.01**
(43) Publication Journal Date : **2000.06.01**
(44) Accepted Journal Date : **2004.03.18**

(71) Applicant(s)
Becton Dickinson and Company

(72) Inventor(s)
Matthew P. Collis

(74) Agent/Attorney
F.B. Rice and Co.,139 Rathdowne Street,CARLTON VIC 3053

(56) Related Art
US 5279936
EP 0818461

ABSTRACT

The present invention relates to a composition which is useful for the reversible binding of a nucleic acid molecule. The composition, which may be packaged in a kit, includes a paramagnetic particle in an acidic solution.

10
20
30
40

50

60

70

AUSTRALIA

Patents Act 1990

Becton Dickinson and Company

TRUE COPY

**COMPLETE SPECIFICATION
STANDARD PATENT**

Invention Title:

*Method for purification and manipulation of nucleic acids
using paramagnetic particles*

The following statement is a full description of this invention
including the best method of performing it known to us:-

BACKGROUND OF THE INVENTION

Access to cellular components such as nucleic acids is imperative to a variety of molecular biology methodologies. Such methodologies include nucleic acid sequencing, direct detection of particular nucleic acid sequences by nucleic acid hybridization and nucleic acid sequence amplification techniques.

The preparation and purification of high-purity double-stranded (ds) plasmid DNA, single-stranded (ss) phage DNA, chromosomal DNA, agarose gel-purified DNA fragments and RNA is of critical importance in molecular biology. Ideally, a method for purifying nucleic acids should be simple, rapid and require little, if any, additional sample manipulation. Nucleic acids rendered by such a method should be immediately amenable to transformation, restriction analysis, ligation or sequencing. A method with all of these features would be extremely attractive in the automation of nucleic acid sample preparation, a goal of research and diagnostic laboratories.

Typically, the preparation of plasmid DNA from crude alcohol precipitates is laborious, most often utilizing CsCl gradients, gel filtration, ion exchange chromatography, or RNase, proteinase K and repeated alcohol precipitation steps. These methods also require considerable downstream sample preparation to remove CsCl and other salts, ethidium bromide and alcohol. Similar arguments extend when using any of these methods for purifying DNA fragments. A further problem with these methods is that small, negatively-

charged cellular components can co-purify with the DNA. Thus, the DNA can have an undesirable level of contamination.

Nucleic acids can also be purified using solid phases. Conventional solid phase extraction techniques have utilized surfaces which either (1) fail to attract and hold sufficient quantities of nucleic acid molecules because of surface design to permit easy recovery of the nucleic acid molecules during elution, or (2) excessively adhere nucleic acid molecules to the surface, thereby hindering recovery of the nucleic acid molecules during elution.

Conventional metal surfaces which cause these problems when utilized in solid phase extraction include certain silica surfaces such as glass and Celite. Adequate binding of

10 nucleic acids to these types of surfaces can be achieved only by utilizing high concentrations of chaotropes or alcohols which are generally toxic, caustic, and/or expensive. For example, it is known that DNA will bind to crushed glass powders and to glass fiber filters in the presence of chaotropes. The chaotropic ions typically are washed away with alcohol, and the DNAs are eluted with low-salt solutions or water. Importantly, RNA and protein do not bind. 15 However, a serious drawback in the use of crushed glass powder is that its binding capacity is low. In addition, glass powders often suffer from inconsistent recovery, incompatibility with borate buffers and a tendency to nick large DNAs. Similarly, glass fiber filters provide a nonporous surface with low DNA binding capacity. Other silicas, such as silica gel and glass beads, are not suitable for DNA binding and recovery. Currently, the solid phase of choice 20 for solid phase extraction of DNA is Celite such as found in Prep-A-Gene™ by Bio-Rad Laboratories. As with the crushed glass powders, high concentrations of chaotropes are required for adequate binding of the DNA to the Celite.

However, the hydration of silica substances has, in some instances resulted in elimination of the need for such high concentrations of chaotropes to elute bound DNA from 25 the silica substance as taught in references such as EP 0 512 767, EP 0 585 660, US 5,674,997 and EP 0 832 897.

There are numerous protocols for purifying DNA. For example, U.S. Patent 4,923,978 discloses a process for purifying DNA in which a solution of protein and DNA is passed over a hydroxylated support and the protein is bound and the DNA is eluted. U.S. Patent 4,935,342 discloses purification of DNA by selective binding of DNA to anion
5 exchangers and subsequent elution. U.S. Patent 4,946,952 discloses DNA isolation by precipitation with water-soluble ketones. A DNA purification procedure using chaotropes and dialyzed DNA is disclosed in U.S. Patent 4,900,677.

Diatoms have also been utilized for purification of nucleic acids as evidenced by U.S. Patent No. 5,234,809 to Boom et al. and U.S. Patent No. 5,075,430 to Little et al.

10 Yet a further technique utilized for purification of nucleic acids is binding to specifically adapted paramagnetic particles. Examples of such techniques may be found in references such as European Specification EP 0 446 260 B1 and U.S. Patent 5,512,439 (Hornes et al.) which describe monodisperse, superparamagnetic particles having a particle diameter standard deviation of less than 5%. Each particle carries a plurality of molecules of
15 an oligonucleotide, with each oligonucleotide having a section serving as a probe for a target nucleic acid molecule of interest.

U.S. Patent No. 4,672,040 (Josephson) and U.S. Patent No. 4,695,393 (Whitehead et al.) describe magnetically responsive particles for use in systems to separate certain
20 molecules. The particles have a metal oxide core surrounded by a stable silicone-coating to which organic and/or biological molecules may be coupled.

U.S. Patent No. 3,970,518 (Giaever) describes a method of sorting and separating a select cell population from a mixed cell population. The method utilizes small magnetic particles which are coated with an antibody to the select cell populations.

U.S. Patent No. 4,141,687 (Forrest et al.) describes an automatic apparatus and
25 method to assay fluid samples. The apparatus utilizes a particulate material with a reagent bound thereto. The particulate material is magnetic, and the reagent is a substance which takes part in a reaction in the reaction mixture.

U.S. Patent No. 4,230,685 (Senyei et al.) describes a method for magnetic separation of cells. The method utilizes magnetically-responsive microspheres which are coated with staphylococcal Protein A to which is bound antibody.

U.S. Patent No. 4,774,265 (Ugelstad et al.) describes a process for preparing magnetic
5 polymer particles. The particles are compact or porous polymer particles treated with a solution of iron salts.

U.S. Patent No. 5,232,782 (Charmot) describes magnetizable "core-shell" microspheres which have a core of a magnetizable filler and a shell of crosslinked organopolysiloxane.

10 U.S. Patent No. 5,395,688 (Wang et al.) describes magnetically responsive fluorescent polymer particles which have a polymeric core coated evenly with a layer of polymer containing magnetically responsive metal oxide.

U.S. Patent No. 5,491,068 and U.S. Patent No. 5,695,946 (Benjamin et al.) describe an
15 assay method for detecting the presence of bacteria using magnetic beads with specific antibodies immobilized thereon.

U.S. Patent No. 5,536,644 (Ullman et al.) describes a particle separation method. The
method utilizes magnetic particles with surface functional groups, and optionally, an
additional surface coating.

European Patent Specification EP 0 444 120 B1 (Hornes et al.) describes a method for
20 detection of target RNA or DNA. The method utilizes magnetic particles carrying a single stranded 5'-attached DNA probe capable of binding the target RNA or DNA.

International Publication No. WO 96/18731 (Deggerdal et al.) describes a method for isolating nucleic acid from a sample using a particulate solid support and an anionic detergent.

U.S. patent No. 5,705,628 (Hawkins) describes a method for DNA purification and
25 isolation using magnetic particles with functional group-coated surfaces.

SUMMARY OF THE INVENTION

In order to provide a more effective and efficient technique for the purification and manipulation of nucleic acids, the present invention relates to a composition useful for reversible binding of a nucleic acid molecule. The composition includes a paramagnetic particle in an acidic environment. The invention also includes such composition packaged as a kit, as well as methods utilizing such a composition to reversibly bind a nucleic acid molecule.

- 10 In a first aspect, the present invention provides the composition useful for reversible binding of a nucleic acid molecule, said composition comprising a paramagnetic particle in an acidic solution.

15 In a second aspect, the present invention provides a kit comprising a paramagnetic particle packaged with an acidic solution.

In a third aspect, the present invention provides a method for reversibly binding a nucleic acid molecule to a paramagnetic particles comprising the steps of;

- 20 a) providing a suspension of paramagnetic particles in an acidic solution; and
b) combining said suspension with a nucleic acid molecule.

Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is solely for the purpose of providing a context for the present invention. It is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed before the priority date of each claim of this application.

30 Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

35

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to unique compositions of matter. More specifically, the composition of matter is a paramagnetic particle in an acidic solution, that is, a solution
5 having a pH of less than about 7.0.

The Applicants found that when in an acidic environment, paramagnetic particles will reversibly bind nucleic acid molecules without the necessity of an anionic detergent as taught in International Publication No. WO 96/18731. Although not desiring to be
10 bound by a particular theory, the Applicants believe that an acidic environment increases the electropositive nature of the iron portion of the molecules, and thus increases the binding of the molecules to the electronegative phosphate portion of a nucleic acid molecule.

15 As used herein, the term paramagnetic particles means particles which are capable of having a magnetic moment imparted to them when placed in a magnetic field. Therefore, such paramagnetic particles, when in such a magnetic field, are movable under the action of such a field. Such movement is useful for moving bound nucleic acid molecules for different aspects of a sample processing protocol or other
20 manipulations. Thus, nucleic acid molecules bound to the paramagnetic particles can be moved to different areas for exposure to different reagents and/or conditions with minimal direct contact due to the application of magnetic force.

5
10
15
20

25
30
35
40
45

The Applicants have found that paramagnetic particles useful in the present invention need not be complicated structures. Thus, iron particles are useful in the present invention, and the iron may be an iron oxide of forms such as ferric hydroxide and ferrosoferric oxide, which have low solubility in an aqueous environment. Other iron particles such as iron
5 sulfide and iron chloride may also be suitable for binding and extracting nucleic acids using the conditions described herein.

Similarly, the shape of the paramagnetic particles is not critical to the present invention. Thus, the paramagnetic particles may be of various shapes, including for example, spheres, cubes, oval, capsule-shaped, tablet-shaped, non-descript random shapes, etc., and may be of uniform
10 shape or non-uniform shapes. Whatever the shape of a paramagnetic particle, its diameter at its widest point is generally in the range of from about 0.5 μm to about 20 μm .

The acidic environment in which the paramagnetic particles effectively, reversibly bind nucleic acid molecules can be provided through a variety of means. For example, the paramagnetic particles can be added to an acidic solution, or an acidic solution may be added
15 to the particles. Alternatively a solution or environment in which the paramagnetic particles are located can be acidified by addition of an acidifying agent such as hydrochloric acid, sulfuric acid, acetic acid and citric acid.

Provided that the environment in which the paramagnetic particles are located is of a pH less than about 7.0, the particles will reversibly bind nucleic acid molecules. Furthermore,
20 the Applicants have found that the nucleic acid binding capacity of the paramagnetic particles increases as the pH decreases.

The acidic environment for the paramagnetic particles of the present invention is believed to allow for elimination of the need for detergents as taught in certain references such as International Publication No. WO 96/18731. Without desiring to be held to a particular theory,
25 the Applicant believes that detergents are not necessary for the present invention, because the acidic solution of the present invention promotes the binding of electropositive paramagnetic particles to electronegative nucleic acid molecules in preference to other substances in a sample

such as nucleic acid hybridization and amplification inhibitors. In contrast, the utilization of detergents as taught in references such as International Publication No. WO 96/18731 is solubilize nucleic acid hybridization and amplification inhibitors in order that such inhibitors do not interfere with binding of nucleic acid molecules to paramagnetic particles.

5 As stated above, in an acidic environment, electropositive paramagnetic particles, such as ferric oxide particles, will bind electronegative nucleic acid molecules. Thus, other materials in the environment, such as inhibitors of nucleic acid hybridization and amplification can be separated from the bound nucleic acid molecules. Such separation can be accomplished by means known to those skilled in the art, such as centrifugation, filtering
10 or application of magnetic force.

The bound nucleic acid molecules can then be eluted into an appropriate buffer for further manipulation, such as hybridization or amplification reactions. such elution can be accomplished by heating the environment of the particles with bound nucleic acids and/or raising the pH of such environment. Agents which can be used to aid the elution of nucleic
15 acid from paramagnetic particles include basic solutions such as potassium hydroxide, sodium hydroxide or any compound which will increase the pH of the environment to an extent sufficient that electronegative nucleic acid is displaced from the particles.

The following examples illustrate specific embodiments of the invention described in
20 this document. As would be apparent to skilled artisans, various changes and modifications are possible and are contemplated within the scope of the invention described.

EXAMPLE 1

Nucleic Acid Binding with Iron Oxide

This example was designed to compare binding of nucleic acid with iron oxide to
5 binding of nucleic acid with zirconium at three target input levels.

The following materials were used in this example.




- A Chlamydia trachomatis preparation
- 10 - Phosphate buffer
- Sample buffer for the BDProbeTec™ ET system
- Hydrated Iron Oxide FeO(OH) particles (30 mesh = 300 –1200 microns)
- Hydrated Zirconium Oxide
- Glycine HCl
- 15 - Chlamydia trachomatis Priming and Amplification Wells for the BDProbeTec™ ET system
- Ethyl Alcohol



The procedure followed for the example was as follows.

- 20 Chlamydia trachomatis solutions at 5,000 Elementary Bodies(EB)/ml, 1,000 EB/ml, 500 EB/ml and 0 EB/ml were prepared in phosphate buffer and each solution was transferred to twelve 2.0 ml centrifuge tubes/spike level. The tubes were heated at 105°C for 30 minutes. Hydrated iron oxide particles washed with DiH₂O by gravity were dispensed into six tubes at each of the Chlamydia trachomatis spike levels at 10 mg/tube. Hydrated zirconium particles
- 25 were dispensed into three tubes containing each Chlamydia trachomatis spike level at 10 mg/tube. No particles were added to three tubes of each level. Thirty ul of 3M Glycine-HCl was dispensed into each tube described above, and the tubes were placed on a end over end

15



Hydrated Iron Oxide, centrifuge wash	0	162
Hydrated Iron Oxide, centrifuge wash	500	7421
Hydrated Iron Oxide, centrifuge wash	1,000	6552
Hydrated Iron Oxide, centrifuge wash	5,000	24415
Hydrated Iron Oxide, magnetic wash	0	203
Hydrated Iron Oxide, magnetic wash	500	8956
Hydrated Iron Oxide, magnetic wash	1,000	5243
Hydrated Iron Oxide, magnetic wash	5,000	20294

The conclusions drawn from the results were as follows.

The MOTA values (sum of individual measurement units over time) are an internal

- 5 Becton Dickinson fluorescent unit with an established cutoff level for positivity at 1,000 MOTA. Magnetically separated iron oxide produced the same positivity as the hydrated zirconium particles. The tubes containing no particles produced no carry over results even at

the 5,000 EB/ml level. This experiment demonstrated that hydrated iron oxide binds DNA under acidic conditions.

EXAMPLE 2

5 Elution and Denaturation of Double Stranded DNA from Iron Oxide Particles Using Potassium Hydroxide

10 This example was performed to determine whether DNA could be eluted and denatured using potassium hydroxide (KOH) alone without heat. In addition, the example was performed to determine which neutralization/sample buffer produces optimum MOTA values.

The materials used in this example were as follows.

- 15 - a 70 mM KOH solution
- a 100 mM Bicine, 2X Chlamydia trachomatis sample buffer (20% DMSO, 18% Glycerol, 0.06% Proclin, 0.013mM Tween 20 and 60 mM KPO₄)
- a 200 mM Bicine, 2X Chlamydia trachomatis sample buffer
- a 300 mM Bicine, 2X Chlamydia trachomatis sample buffer
- 20 - a Chlamydia trachomatis sample buffer (10% DMSO, 9% Glycerol, 0.03% Proclin, 0.0065mM Tween 20 and 30 mM KPO₄)
- Hydrated zirconium particles
- Hydrated iron oxide particles
- a Chlamydia trachomatis preparation
- 25 - 95% Ethyl alcohol (ETOH)
- Phosphate buffer

The procedure followed for this example was as follows.

Chlamydia trachomatis solutions were prepared at 1,000 EB/ml and 5,000 EB/ml in phosphate buffer and were dispensed into 32 two ml centrifuge tubes/spike level at 1.0

- 5 ml/tube. Hydrated iron oxide was dispensed into 16 tubes/spike level at 10 mg/tube and 60 ul of 3M glycine HCl was added to each tube. Hydrated zirconium particles were dispensed into 16 tubes/spike level at 10 mg/tube and 30 ul of 3M glycine HCl was added to each tube. The tubes were maintained on an end over end rotator for 30 minutes. The iron oxide tubes were magnetically separated, and then washed with 1.0 ml of ETOH and then 1.0 ml of DiH₂O.
- 10 The zirconium particle tubes were centrifuge washed once with one ml of ETOH and then with one ml of DiH₂O. Twelve tubes of each particle type and spike level had 500 ul of 70 mM KOH added to each tube for 15 minutes. Of the twelve tubes, four tubes of both particle type and spike level had 100 mM 2X Chlamydia trachomatis sample buffer (w/phosphate) added to each tube at 500 ul. Of the twelve tubes, four tubes of both particle types and spike
- 15 levels had 200 mM 2X Chlamydia trachomatis sample buffer (w/phosphate) added to each tube at 500 ul. Of the twelve tubes, four tubes of both particle types and spike levels had 300 mM 2X Chlamydia trachomatis sample buffer (w/phosphate) added to each tube at 500 ul. To the four remaining tubes, 1.0 ml of sample buffer was added. Two tubes of each particle type, spike level and buffer type were placed in a boiling water bath for five minutes. The sample
- 20 fluid was then immediately amplified in the Chlamydia trachomatis BDProbeTec™ ET system.

The results of this example were as follows.

Bicine Concentration	Chlamydia Concentration EB/ml	Iron Oxide No Heat	Iron Oxide Heat	Zirconium No Heat	Zirconium Heat
100 mM	1,000	214	298	1032	11556
200 mM	1,000	3106	123	1320	7250
300 mM	1,000	1318	4871	113	5833
Control	1,000	280	4775	3245	7988
100 mM	5,000	814	3711	24565	38873
200 mM	5,000	5495	19173	11415	32871
300 mM	5,000	13433	15302	18298	18127
Control	5,000	422	17106	4753	20208

5

The conclusions drawn from the results above were as follows.

The results indicate that DNA can be eluted and denatured from hydrated iron oxide and hydrated zirconium oxide particles with KOH alone (w/o heat) and that the neutralization

buffers which produced optimum MOTA values were the 200 and 300 mM Bicine 2X neutralization buffers.

5

EXAMPLE 3

Extraction of Nucleic Acid From Urine Samples Using Hydrated Iron Oxide


10 This example was performed to determine if hydrated iron oxide at low pH can be used to extract Chlamydia trachomatis DNA from spiked urine samples.

The materials used in this example were as follows.

- Urine samples
- Chlamydia trachomatis preparations
- 15 - Sample buffer
- Hydrated zirconium particles
- Hydrated iron oxide particles
- 95% Ethyl Alcohol (ETOH)
- Glycine HCl
- 20 - BDProbeTec™ ET system Chlamydia trachomatis primer wells and amplification wells

The procedure followed for this example was as follows.

Each of ten urine samples were split into three two ml aliquots and the urine samples
25 were spiked with Chlamydia trachomatis preparation at 1,000 EB/ml, 2,500 EB/ml and 5,000 EB/ml. The urine samples were split into 1.0 ml volumes in two ml centrifuge tubes. Then, 10 mg of hydrated iron oxide particles were dispensed into one tube at each spike level for



- each sample. Sixty ul of 3M Glycine HCl was added to each iron oxide tube. The tubes were maintained on an end over end rotator for 30 minutes. In addition, 10 mg of hydrated zirconium particles were dispensed into one tube at each spike level for each sample. Thirty ul of 3 M Glycine HCl was added to each tube containing hydrated zirconium. The tubes
- 5 were maintained on an end over end rotator for 30 minutes. The tubes containing hydrated zirconium particles were centrifuge washed at 10,000 g using one ml of ETOH and then one ml of DiH₂O. The tubes were re-suspended with 1.0 ml of sample buffer. The tubes containing iron oxide particles were magnetically separated, and then washed using one ml of ETOH and then 1.0 ml of DiH₂O and the tubes were re-suspended with 1.0 ml of sample
- 10 buffer. The sample fluid from each tube was boiled for 5 minutes and then amplified using the Chlamydia trachomatis BDProbeTec™ ET system.

The results of this example were as follows.

15	Chlamydia Level	Particle Type	Positives
	1,000 EB/ml	Hydrated Iron Oxide	3/10
	1,000 EB/ml	Hydrated Zirconium	9/10
	2,500 EB/ml	Hydrated Iron Oxide	10/10
	2,500 EB/ml	Hydrated Zirconium	10/10
	5,000 EB/ml	Hydrated Iron Oxide	10/10
	5,000 EB/ml	Hydrated Zirconium	10/10

The conclusions drawn from the above results were as follows.

The detectable positivity rate drops at the same level with either the hydrated zirconium particles or the hydrated iron oxide particles. This indicates similar performance with both particles types. Iron oxide at low pH can extract DNA from urine samples.

EXAMPLE 4

Extraction of Nucleic Acid From Plasma Samples Using Hydrated Iron Oxide

This example was performed to determine if hydrated iron oxide at low pH can be used to extract Chlamydia trachomatis DNA from spiked plasma samples. In addition, this example was performed to compare a plasma DNA extraction method using hydrated zirconium particles to a plasma DNA extraction method using hydrated iron oxide particles.

The materials used for this example were as follows.

- 300 mM Bicine 2X Chlamydia trachomatis sample buffer
- Hydrated iron oxide particles
- Hydrated zirconium oxide particles
- Guanidine isothiocyanate
- Chlamydia trachomatis stock preparation
- Chlamydia trachomatis BDProbeTec™ ET primer wells and amplification wells
- Internal Amplification Control (IAC) BDProbeTec™ ET primer wells and amplification wells
- 150 mM KOH
- Plasma samples
- 95% Ethyl alcohol

The procedures followed for this example was as follows.

Iron Oxide Procedure

5

One ml of DiH₂O was added to eight two ml centrifuge tubes containing 40 mg of hydrated iron oxide particles. Two plasma samples were spiked with Chlamydia trachomatis preparation at 9,000 EB/300 ul, 6,000 EB/300 ul, 3,000 EB/300 ul and 1,500 EB/300 ul.



10

Then, 300 ul of each spiked plasma sample was added to the tubes containing hydrated iron oxide particles. The tubes were heated at 105°C for 30 minutes. Eighty ul of glacial acetic acid was then added to each tube, and the tubes were placed on an end over end rotator for 30 minutes. The tubes were then placed on a magnetic tube rack and the treated sample fluid was removed. The tubes were magnetically separated, and then washed with 1.0 ml of 25 mM acetic acid twice. The DNA was eluted from the hydrated iron oxide particles with 500



15

ul of 150 mM KOH for 15 minutes. The solutions were neutralized with 500 ul of 300 mM 2X sample buffer. The samples were placed in a boiling water bath for five minutes. The samples were then amplified using the BDProbeTec™ ET Chlamydia trachomatis system.



Hydrated Zirconium Procedure

20

The same two plasma samples from the iron oxide procedure above were dispensed at 300ul/tube into four 2.0 ml centrifuge tubes for each plasma sample. Five molar guanidine isothiocyanate (GITC) was added to each tube at 700 ul. The four tubes of each sample were spiked with Chlamydia trachomatis preparation at 9,000 EB/300 ul, 6,000 EB/300 ul, 3,000

25

EB/300 ul and 1,500 EB/300 ul, respectively, and the tubes were maintained at room temperature for 15 minutes. A hydrated zirconium oxide particle slurry was prepared by combining 10 mg of zirconium oxide particles with 30 ul of 3M glycine HCl. Thirty ul of this

slurry was added to each tube and the tubes were placed on an end over end rotator for 30 minutes. The tubes were centrifuged at 10,000 g for 3.0 minutes, the supernatant was removed, and 1.0 ml of 2M GITC was added to each tube. The tubes were centrifuged at 10,000 g for 3.0 minutes, the supernatant was removed and 1.0 ml of 80% ethyl alcohol/20% 5 50 mM Tris buffer was added to each tube. The tubes were washed two additional times by centrifuging at 10,000 g for 3.0 minutes, extracting the supernatant and adding 1.0 ml of DiH₂O. The DNA was eluted in 1.0 ml of 150 mM KOH/300 mM Bicine 2X sample buffer. The samples were placed in a boiling water bath for five minutes. The samples were pop centrifuged at 3,200 g and were amplified using the Chlamydia trachomatis BDProbeTec™ 10 ET system.

Internal Amplification Control Procedure

Sample buffer was spiked with Chlamydia trachomatis preparation at 9,000 EB/300 ul, 15 6,000 EB/300 ul, 3,000 EB/300 ul and 1,000 EB/300 ul. The samples were placed in a boiling water bath for five minutes and were amplified using the Chlamydia trachomatis BDProbeTec™ ET system.

The results of this example were as follows.

Particle Type	Spike Level	Chlamydia MOTA Mean	IAC MOTA Mean
Iron oxide	9,000	14871	14261
Zirconium	9,000	16774	16886
Control	9,000	19808	35077
Iron oxide	6,000	8668	19313

Zirconium	6,000	19115	24339
Control	6,000	22129	34234
Iron oxide	3,000	2436	16128
Zirconium	3,000	20098	33188
Control	3,000	21954	26355
Iron oxide	1,000	492	10021
Zirconium	1,000	15587	25805
Control	1,000	2679	14423

The conclusions drawn from the above results were as follows.

- 5 Using a MOTA positivity value of 1,000, hydrated iron oxide was capable of extracting DNA from plasma samples up to 3,000 EB/300 ul.

EXAMPLE 5

Nucleic Acid Capture Using Ferrosoferric oxide (Fe_3O_4)

10

This example was designed to compare two nucleic acid capture particles: Ferric Hydroxide (hydrated ferric oxide) and Ferrosoferric oxide under neutral and acidic binding conditions.

15


The materials used for this example were as follows.

- Phosphate buffer
- Ferric hydroxide ($\text{FeO}(\text{OH})$) particles

- Ferrosoferric oxide (Fe_3O_4) particles
- Chlamydia trachomatis preparation
- Chlamydia trachomatis BDProbeTec™ ET primer wells and amplification wells
- Internal Amplification Control (IAC) BDProbeTec™ ET primer wells and amplification

5 wells


- Acetic acid
- 150 mM KOH
- 300 mM Bicine, 2X Chlamydia trachomatis sample buffer



10 The procedure followed for this example was as follows.



Ferric hydroxide ($\text{FeO}(\text{OH})$) powder was prepared by mortar and pestle grinding 30-50 mesh material to a fine powder. The powder was allowed to settle in an aqueous solution for four minutes. The supernatant was extracted and the material was allowed to settle for an additional fifteen minutes. The pelleted material was then magnetically separated and was magnetically washed with DiH_2O . The material was then filtered onto 20 μm filter paper and dried overnight at 37°C . Then, 60 μg of the material was transferred to a tube containing 480 μl of DiH_2O . This slurry was dispensed into six 2 ml centrifuge tubes. Also, 240 μg of dried $\text{FeO}(\text{OH})$ material was transferred to a tube containing 480 μl of DiH_2O . The slurry created was transferred to six 2 ml centrifuge tubes.



In addition, 60 μg of Fe_3O_4 (sized at 88-92% passing a 325 sieve) was transferred to a tube containing 480 μl of DiH_2O . The slurry created was dispensed into six 2 ml centrifuge tubes. Another slurry was created by transferring 240 mg of dried Fe_3O_4 material to a tube containing 480 μl of DiH_2O . This slurry was then transferred to six 2 ml centrifuge tubes.

25 Chlamydia trachomatis solutions were prepared at 0 EB/ml, 1,000 EB/ml and 4,000 EB/ml in phosphate buffer. Each concentration of Chlamydia trachomatis solution was dispensed into two tubes of each particle type. The tubes were heated at 105°C for 30 minutes. An 80 μl

aliquot of acetic acid was added to one tube of each particle type, and the tubes were maintained on an end over end rocker for 30 minutes. The tubes were then magnetically separated and washed with 1.0 ml of 25 mM acetic acid/wash twice. A 150 mM KOH aliquot was added to each tube at 500 ul/tube for 15 minutes, and 500 ul of 300 mM Bicine 2X

- 5 sample buffer was added to each tube and the tubes were placed in a boiling water bath for five minutes. The sample fluid was amplified using the Chlamydia trachomatis BDProbeTec™ ET system.

The results from this example were as follows.

10

Chlamydia	Iron Oxide			Chlamydia	IAC
Level	Type	Mass	Acid Volume	MOTA	MOTA
0	FeO(OH)	10	80	187	10076
0	FeO(OH)	40	80	127	7751
0	FeO(OH)	10	0	5	13928
0	FeO(OH)	40	0	111	13814
1,000	FeO(OH)	10	80	3343	11610
1,000	FeO(OH)	40	80	11574	13886
1,000	FeO(OH)	10	0	2209	11883
1,000	FeO(OH)	40	0	2350	13677
4,000	FeO(OH)	10	80	5778	18752
4,000	FeO(OH)	40	80	13997	10866
4,000	FeO(OH)	10	0	2195	17593
4,000	FeO(OH)	40	0	4426	12065

0	Fe ₃ O ₄	10	80	34	13279
0	Fe ₃ O ₄	40	80	37	11336
0	Fe ₃ O ₄	10	0	1	12965
0	Fe ₃ O ₄	40	0	2	5351
1,000	Fe ₃ O ₄	10	80	8507	17116
1,000	Fe ₃ O ₄	40	80	3772	4923
1,000	Fe ₃ O ₄	10	0	48	15623
1,000	Fe ₃ O ₄	40	0	7	925
4,000	Fe ₃ O ₄	10	80	7019	16337
4,000	Fe ₃ O ₄	40	80	21486	17983
4,000	Fe ₃ O ₄	10	0	10438	17613
4,000	Fe ₃ O ₄	40	0	206	10454

The conclusions drawn from the above results were as follows.

- Using a MOTA positivity value of 1,000, Fe₃O₄ can extract nucleic acid to a level of
- 5 1,000 EB/ml, which is comparable to the extraction capability of FeO(OH) with both masses.
- A negative acid volume negatively impacted both the number of positive values at the lower 1,000 EB/ml level and MOTA values.

While the invention has been described with some specificity, modifications apparent to those with ordinary skill in the art may be made without departing from the scope of the invention. Various features of the invention are set forth in the following claims

23

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A composition useful for reversible binding of a nucleic acid molecule, said composition comprising a paramagnetic particle in an acidic solution.
- 5 2. The composition of claim 1 wherein the paramagnetic particle comprises iron.
3. The composition of claim 1 or claim 2, wherein the paramagnetic particle is selected from the group consisting of an iron oxide, iron sulfide and iron chloride.
- 10 4. The composition of claim 4 wherein the iron oxide is ferric hydroxide or ferrosiferic oxide.
- 15 5. A kit comprising a paramagnetic particle packaged with an acidic solution.
6. The kit of claim 5 wherein the paramagnetic particle comprises iron.
7. The kit of claim 5 or claim 6 wherein the paramagnetic particle is selected from the group consisting of an iron oxide, iron sulfide and iron chloride.
- 20 8. A method for reversibly binding a nucleic acid molecule to a paramagnetic particles comprising the steps of;
 - 25 a) providing a suspension of paramagnetic particles in an acidic solution; and
 - b) combining said suspension with a nucleic acid molecule.
9. The method of claim 8 wherein the paramagnetic particles comprises iron.
- 30 10. The method of claim 8 or claim 9 wherein the paramagnetic particles are selected from the group consisting of an iron oxide, iron sulfide and iron chloride.

35

11. A composition useful for reversible binding of a nucleic acid molecule substantially as hereinbefore described with particular reference to the examples and/or the preferred embodiments and excluding, if any, comparative examples.
- 5 12. A kit substantially as hereinbefore described with particular reference to the examples and/or the preferred embodiments and excluding, if any, comparative examples.
- 10 13. A method for reversibly binding a nucleic acid molecule to a paramagnetic particle substantially as hereinbefore described with particular reference to the examples and/or the preferred embodiments and excluding, if any, comparative examples.

Dated this seventeenth day of July 2003

Becton, Dickinson and Company
Patent Attorneys for the Applicant:

F B RICE & CO

15

15