



(72) REICHARD, Raymond, US

(72) BROWN, James Raymond, US

(72) LAWLOR, Elizabeth J., US

(71) SMITHKLINE BEECHAM CORPORATION, US

(71) SMITHKLINE BEECHAM PLC, GB

(51) Int.Cl.⁶ C12N 15/52, A61K 31/70, C12Q 1/68, G01N 33/573,
G01N 33/569, C12N 9/00, A61K 48/00, A61K 38/43, C07K 16/40,
C12Q 1/25, A61K 39/118

(30) 1997/07/23 (08/899,028) US

(54) **NOUVEAUX HISS**

(54) **NOVEL HISS**

(57) La présente invention porte sur des polypeptides hisS et sur l'ADN (ARN) codant les polypeptides hisS ainsi que sur des méthodes pour la production de ces polypeptides par des techniques recombinantes. On divulgue également des méthodes pour utiliser les polypeptides hisS à des fins de dépistage de composés antibactériens.

(57) The invention provides hisS polypeptides and DNA (RNA) encoding hisS polypeptides and methods for producing such polypeptides by recombinant techniques. Also provided are methods for utilizing hisS polypeptides to screen for antibacterial compounds.

ABSTRACT OF THE DISCLOSURE

The invention provides hisS polypeptides and DNA (RNA) encoding hisS polypeptides and methods for producing such polypeptides by recombinant techniques. Also provided are methods for utilizing hisS polypeptides to screen for antibacterial compounds.

NOVEL hisS

FIELD OF THE INVENTION

5 This invention relates to newly identified polynucleotides and polypeptides, and their production and uses, as well as their variants, agonists and antagonists, and their uses. In particular, in these and in other regards, the invention relates to novel polynucleotides and polypeptides of the histidyl tRNA synthetase family, hereinafter referred to as "hisS".

10 BACKGROUND OF THE INVENTION

Chlamydiaceae is a family of obligate intracellular parasites. All members share a common developmental cycle. Chlamydia infect a wide range of vertebrate host, particularly humans.

15 *Chlamydia trachomatis* is one of the two recognized species of Chlamydia. Human infections caused by *Chlamydia trachomatis* are widespread. This species is one of the most common cause of sexually transmitted disease in the world. It is also one of the main causes of infertility in humans.

20 The frequency of *Chlamydia trachomatis* infections has risen dramatically in the past 20 years. This has been attributed to the emergence of multiply antibiotic resistant strains and an increasing population of people with weakened immune systems. It is no longer uncommon to isolate *Chlamydia trachomatis* strains which are resistant to some or all of the standard antibiotics. This has created a demand for both new anti-microbial agents and diagnostic tests for this organism.

25 The t-RNA synthetases have a primary role in protein synthesis according to the following scheme:



in which AA is an amino acid.

Inhibition of this process leads to a reduction in the levels of charged t-RNA and this triggers a cascade of responses known as the stringent response, the result of which is the induction of a state of dormancy in the organism. As such selective inhibitors of bacterial t-RNA synthetase have potential as antibacterial agents. One example of such is mupirocin which is a selective inhibitor of isoleucyl t-RNA synthetase. Other t-RNA synthetases are now being examined as possible anti-bacterial targets, this process being greatly assisted by the isolation of the synthetase.

Clearly, there is a need for factors, such as the novel compounds of the invention, that have a present benefit of being useful to screen compounds for antibiotic activity. Such factors are also useful to determine their role in pathogenesis of infection, dysfunction and disease. There is also a need for identification and characterization of such factors and their antagonists and agonists which can play a role in preventing, ameliorating or correcting infections, dysfunctions or diseases.

The polypeptides of the invention have amino acid sequence homology to a known *Streptococcus equisimilis* histidyl tRNA synthetase protein. (*S.equisimilis* HRS SwissProt P30053; Menguito C.A., Keherly M.J., Tang C.-Y., Papaconstantinou J., Weigel P.H., *Nucleic Acids Res.*, 21:615-620 (1993).)

15

SUMMARY OF THE INVENTION

It is an object of the invention to provide polypeptides that have been identified as novel hisS polypeptides by homology between the amino acid sequence set out in Table 1 [SEQ ID NO: 2] and a known amino acid sequence or sequences of other proteins such as *Streptococcus equisimilis* histidyl tRNA synthetase protein.

It is a further object of the invention to provide polynucleotides that encode hisS polypeptides, particularly polynucleotides that encode the polypeptide herein designated hisS.

In a particularly preferred embodiment of the invention the polynucleotide comprises a region encoding hisS polypeptides comprising the sequence set out in Table 1 [SEQ ID NO:1] which includes a full length gene, or a variant thereof.

25

In another particularly preferred embodiment of the invention there is a novel hisS protein from *Chlamydia trachomatis* comprising the amino acid sequence of Table 1 [SEQ ID NO:2], or a variant thereof.

5 In accordance with another aspect of the invention there is provided an isolated nucleic acid molecule encoding a mature polypeptide expressible by the *Chlamydia trachomatis* D/UW-3/Cx strain.

A further aspect of the invention there are provided isolated nucleic acid molecules encoding hisS, particularly *Chlamydia trachomatis* hisS, including mRNAs, cDNAs, genomic DNAs. Further embodiments of the invention include biologically, diagnostically, 10 prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

In accordance with another aspect of the invention, there is provided the use of a polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic immunization. Among the particularly preferred embodiments of the invention are naturally 15 occurring allelic variants of hisS and polypeptides encoded thereby.

Another aspect of the invention there are provided novel polypeptides of *Chlamydia trachomatis* referred to herein as hisS as well as biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

Among the particularly preferred embodiments of the invention are variants of hisS 20 polypeptide encoded by naturally occurring alleles of the hisS gene.

In a preferred embodiment of the invention there are provided methods for producing the aforementioned hisS polypeptides.

In accordance with yet another aspect of the invention, there are provided inhibitors to such polypeptides, useful as antibacterial agents, including, for example, antibodies.

25 In accordance with certain preferred embodiments of the invention, there are provided products, compositions and methods for assessing hisS expression, treating disease, for example, classic ocular trachoma, inclusion conjunctivitis, genital trachoma, infant pneumonitis, Lymphogranuloma Venerium, incipient trachoma, keratitis, papillary hypertrophy, corneal infiltration, vulvovaginitis, ear infection, mucopurulent rhinitis, salpingitis, cervicitis, cervical 30 follicles, prostatitis, proctitis, urethritis, lymphogranule inguinale, climatic bubo, tropical bubo, and esthiomene., assaying genetic variation, and administering a hisS polypeptide or

polynucleotide to an organism to raise an immunological response against a bacteria, especially a *Chlamydia trachomatis* bacteria.

5 In accordance with certain preferred embodiments of this and other aspects of the invention there are provided polynucleotides that hybridize to hisS polynucleotide sequences, particularly under stringent conditions.

In certain preferred embodiments of the invention there are provided antibodies against hisS polypeptides.

10 In other embodiments of the invention there are provided methods for identifying compounds which bind to or otherwise interact with and inhibit or activate an activity of a polypeptide or polynucleotide of the invention comprising: contacting a polypeptide or polynucleotide of the invention with a compound to be screened under conditions to permit binding to or other interaction between the compound and the polypeptide or polynucleotide to assess the binding to or other interaction with the compound, such binding or interaction being associated with a second component capable of providing a detectable signal in response to the binding or interaction of the polypeptide or polynucleotide with the compound; and determining whether the compound binds to or otherwise interacts with and activates or inhibits an activity of the polypeptide or polynucleotide by detecting the presence or absence of a signal generated from the binding or interaction of the compound with the polypeptide or polynucleotide.

15 In accordance with yet another aspect of the invention, there are provided hisS agonists and antagonists, preferably bacteriostatic or bacteriocidal agonists and antagonists.

20 In a further aspect of the invention there are provided compositions comprising a hisS polynucleotide or a hisS polypeptide for administration to a cell or to a multicellular organism.

25 Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following descriptions and from reading the other parts of the present disclosure.

GLOSSARY

30 The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"Host cell" is a cell which has been transformed or transfected, or is capable of transformation or transfection by an exogenous polynucleotide sequence.

"Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences.

5 In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" and "similarity" can be readily calculated by known methods, including but not limited to those described in (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome*

10 *Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48: 1073 (1988). Preferred methods to determine

15 identity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN, and FASTA (Atschul, S.F. et

20 al., *J. Molec. Biol.* 215: 403-410 (1990). The BLAST X program is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990). As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO: 1 it is intended that the nucleotide sequence of

25 the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: 1. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or

30 a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at

the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. Analogously, by a polypeptide having an amino acid sequence having at least, for example, 95% identity to a reference amino acid sequence of SEQ ID NO:2 is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO: 2. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

"Isolated" means altered "by the hand of man" from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

"Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotide(s)" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions or single-, double- and triple-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded, or triple-stranded regions, or a mixture of single- and double-stranded regions. In addition, "polynucleotide" as used herein refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The strands in such regions may be from the same molecule or from different molecules. The regions may include all of one or more of the molecules, but more typically involve only a region of some of the molecules. One of the molecules of a triple-helical region

often is an oligonucleotide. As used herein, the term "polynucleotide(s)" also includes DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with backbones modified for stability or for other reasons are "polynucleotide(s)" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or modified bases, such as tritylated bases, to name just two examples, are polynucleotides as the term is used herein. It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term "polynucleotide(s)" as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA characteristic of viruses and cells, including, for example, simple and complex cells. "Polynucleotide(s)" also embraces short polynucleotides often referred to as oligonucleotide(s).

"Polypeptide(s)" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds. "Polypeptide(s)" refers to both short chains, commonly referred to as peptides, oligopeptides and oligomers and to longer chains generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene encoded amino acids. "Polypeptide(s)" include those modified either by natural processes, such as processing and other post-translational modifications, but also by chemical modification techniques. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature, and they are well known to those of skill in the art. It will be appreciated that the same type of modification may be present in the same or varying degree at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains, and the amino or carboxyl termini. Modifications include, for example, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid

residues, hydroxylation and ADP-ribosylation, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins, such as arginylation, and ubiquitination. See, for instance, *PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993) and Wold, F., *Posttranslational Protein Modifications: Perspectives and Prospects*, pgs. 1-12 in *POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS*, B. C. Johnson, Ed., Academic Press, New York (1983); Seifter et al., *Meth. Enzymol.* 182:626-646 (1990) and Rattan et al., *Protein Synthesis: Posttranslational Modifications and Aging*, Ann. N.Y. Acad. Sci. 663: 48-62 (1992). Polypeptides may be branched or cyclic, with or without branching. Cyclic, branched and branched circular polypeptides may result from post-translational natural processes and may be made by entirely synthetic methods, as well.

“Variant(s)” as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques, by direct synthesis, and by other recombinant methods known to skilled artisans.

DESCRIPTION OF THE INVENTION

30

The invention relates to novel hisS polypeptides and polynucleotides as described in greater detail below. In particular, the invention relates to polypeptides and polynucleotides of a novel hisS of *Chlamydia trachomatis*, which is related by amino acid sequence homology to Streptococcus equisimilis histidyl tRNA synthetase polypeptide. The invention relates especially to hisS having the nucleotide and amino acid sequences set out in Table 1 [SEQ ID NO: 1] and Table 1 [SEQ ID NO: 2] respectively, and to the hisS nucleotide sequences of the DNA in the strain and amino acid sequences encoded thereby.

TABLE 1

10 hisS Polynucleotide and Polypeptide Sequences

(A) Sequences from *Chlamydia trachomatis* hisS polynucleotide sequence [SEQ ID NO:1].

5' -1 GTGAGAATAG TCCGGTTATA CACAATGAGT AATGCATTGC CAAAAGGCGT

15 51 TTTTGATATT TTTCCCTATG TAACAAGCCC AAAAAATCTT TGGAGAAATT

101 CTTCTCTATG GAAACGTGTT GAGCATGCAG CCCATCGTAT TTGTAATTTA

151 TATGGATTTG ATGAGATCCG AACTCCAGTT TTTGAAAAGA CAGAGACTTT

20 201 TTTACGCGTC GGAGAGTACA GTGATATTGT AAAAAAGGAA GTTTATACCT

251 TCTTAGATAA AAAAAGACGT TCTTTGACTT TGCCTCCAGA AGGGACTGCA

25 301 GCAGTTGTTT GTGCATTGTT GGATCATTCT GCTGATATGC GCAAAGATAA

351 TAAGTTTTAT TATATTTTGC CCATGTTTCG TTACGAGCGG CAACAATCTG

401 GACGTTACCG TCAGCATCAT CAGTTCGGTC TAGAAGCTAT CGGTGTGCGG

30 451 CACCCCTTAC GAGATGCGGA GGTGCTCTCT CTGTTATGGG ATTTTTATGC

501 AGCGGTCGGG CTCAGCATA TGCAAATCCA TGTGAATTTT TTAGGAGGGC

551 AAAAGACTCG GGCTCGTTAT GACGAAGCTT TGCGGGAGTT CTTCCGTAAG
601 GATCTAGACC GGTTATCGCC TCTGAGTCAA GAAAGATATC ATGCGAACTT
5
651 ATTGCGTATA TTAGATTCTA AGGAGCCAGA AGACCAGGAA TTCATTGAAA
701 AAGCTCCCTC AATTTTGGAT TACATAGATG ATCGGGATTT AAGCTATTTT
10
751 GATGCAGTAT TAGCCCAATT AAAGGCTTTA GGGATTTCTT TTGCAATCAA
801 TCCAAGGCTA GTTCGAGGGT TGGATTATTA TACGGATCTT GTATTTGAAG
851 CGGTGACTGT TGTGGGAGAG CACTCCTATG CATTGGGAGG CGGCGGGCGT
15
901 TATGATGAAT TGGTTGCGCA GTCTGGAGGC CCTTCTATGC CAGCTTTTGG
951 TTTTGGAGTG GGATTGGAAA GAGTAATCCA AACGCTATTA GAGCAAGGAA
20
1001 ACTCTTTATC GACCTCTACG CGACGGTTGC GATTAATTCC TATGGATGAG
1051 CAGGCAGATG CATTTTGTTT TTCATGGGCA AATCGTTTAC GTAACCTAGG
1101 CATTGCAACA GAAGTAGATT GGAGCCATAA GAAGCCTAAA TTGTCTCTTA
25
1151 AAGATGCTGC CGATCAACAA GTCAGTTTTG TTTGTCTCTT AGGCGAACAA
1201 GAATTAGCAA CGAAACAATT TATAGTTAAA GATATGTCTT TGCATCAAAG
30
1251 CTTCTCAGGG GCTCAACAAG ATGTAGAACA AAGGTTGGTT TATGAAGTAC
1301 AGAACGCATA A-3'

(B) hisS polypeptide sequence deduced from the polynucleotide sequence in this table
[SEQ ID NO:2].

NH₂-1 MRIVRLYTMS NALPKGVFEDI FPYVTSPKNL WRNSSLWKRV EHAAHRICNL

5 51 YGFDEIRTPV FEKTETFLRV GEYSDIVKKE VYTFLDKKRR SLTLRPEGTA

101 AVVRALLDHS ADMRKDNKFY YILPMFRYER QQSGRYRQHH QFGLEAIGVR

151 HPLRDAEVLS LLWDFYAAVG LQHQIHVNF LGGQKTRARY DEALREFFRK

10 201 DLDRLSPLSQ ERYHANLLRI LDSKEPEDQE FIEKAPSILD YIDDRDLSYF

251 DAVLAQLKAL GISFAINPRL VRGLDYITDL VFEAVTVVGE HSYALGGGGR

15 301 YDELVAQSGG PSMPAFGFGV GLERVIQTLLE EQGNSLSTST RRLRLIPMDE

351 QADAFCSWA NRLRNLIAT EVDWSHKKPK LSLKDAADQQ VSFVCLLGEQ

401 ELATKQFIVK DMSLHQSFSG AQQDVEQRLV YEVQNA*-COOH

20

(C) Polynucleotide sequence embodiments [SEQ ID NO:1].

X-(R₁)_n-1 GTGAGAATAG TCCGGTTATA CACAATGAGT AATGCATTGC CAAAAGGCGT

51 TTTTGATATT TTTCCCTATG TAACAAGCCC AAAAAATCTT TGGAGAAATT

25 101 CTTCTCTATG GAAACGTGTT GAGCATGCAG CCCATCGTAT TTGTAATTTA

151 TATGGATTTG ATGAGATCCG AACTCCAGTT TTTGAAAAGA CAGAGACTTT

30 201 TTTACGCGTC GGAGAGTACA GTGATATTGT AAAAAAGGAA GTTTATACCT

251 TCTTAGATAA AAAAAGACGT TCTTTGACTT TGCCTCCAGA AGGGACTGCA

301 GCAGTTGTTT GTGCATTGTT GGATCATTCT GCTGATATGC GCAAAGATAA

351 TAAGTTTTAT TATATTTTGC CCATGTTTCG TTACGAGCGG CAACAATCTG
 401 GACGTTACCG TCAGCATCAT CAGTTCGGTC TAGAAGCTAT CGGTGTGCGG
 5
 451 CACCCCTTAC GAGATGCGGA GGTGCTCTCT CTGTTATGGG ATTTTTATGC
 501 AGCGGTCGGG CTTCAGCATA TGCAAATCCA TGTGAATTTT TTAGGAGGGC
 10
 551 AAAAGACTCG GGCTCGTTAT GACGAAGCTT TGCGGGAGTT CTTCCGTAAG
 601 GATCTAGACC GGTATCGCC TCTGAGTCAA GAAAGATATC ATGCGAACTT
 651 ATTGCGTATA TTAGATTCTA AGGAGCCAGA AGACCAGGAA TTCATTGAAA
 15
 701 AAGCTCCCTC AATTTTGGAT TACATAGATG ATCGGGATTT AAGCTATTTT
 751 GATGCAGTAT TAGCCCAATT AAAGGCTTTA GGGATTTCTT TTGCAATCAA
 20
 801 TCCAAGGCTA GTTCGAGGGT TGGATTATTA TACGGATCTT GTATTTGAAG
 851 CGGTGACTGT TGTGGGAGAG CACTCCTATG CATTGGGAGG CGGCGGGCGT
 901 TATGATGAAT TGGTTGCGCA GTCTGGAGGC CCTTCTATGC CAGCTTTTGG
 25
 951 TTTTGGAGTG GGATTGGAAA GAGTAATCCA AACGCTATTA GAGCAAGGAA
 1001 ACTCTTTATC GACCTCTACG CGACGGTTGC GATTAATTCC TATGGATGAG
 30
 1051 CAGGCAGATG CATTTTGTTC TTCATGGGCA AATCGTTTAC GTAACCTAGG
 1101 CATTGCAACA GAAGTAGATT GGAGCCATAA GAAGCCTAAA TTGTCTCTTA
 1151 AAGATGCTGC CGATCAACAA GTCAGTTTTG TTTGTCTCTT AGGCGAACAA

1201 GAATTAGCAA CGAAACAATT TATAGTTAAA GATATGTCTT TGCATCAAAG

1251 CTTCTCAGGG GCTCAACAAG ATGTAGAACA AAGGTTGGTT TATGAAGTAC

5

1301 AGAACGCATA A- (R₂)_n-Y

(D) Polypeptide sequence embodiments [SEQ ID NO:2].

X- (R₁)_n-1 MRIVRLYTMS NALPKGVDI FPYVTSPKNL WRNSSLWKRV EHAHRICNL

10

51 YGFDEIRTPV FEKTETFLRV GEYSDIVKKE VYTFLDKKRR SLTLRPEGTA

101 AVVRALLDHS ADMRKDNKFY YILPMFRYER QQSGRYRQHH QFGLEAIGVR

15

151 HPLRDAEVLS LLWDFYA AVG LQHMQIHVNF LGGQKTRARY DEALREFFRK

201 DLDRLSPLSQ ERYHANLLRI LDSKEPEDQE FIEKAPSILD YIDDRDLSYF

251 DAVLAQLKAL GISFAINPRL VRGLDYITDL VFEAVTVVGE HSYALGGGGR

20

301 YDELVAQSGG PSMPAFGFGV GLERVIQTLL EQGNSLSTST RRLRLIPMDE

351 QADAFCSWA NRLRNLIAT EVDWSHKKPK LSLKDAADQQ VSFVCLLGEQ

25

401 ELATKQFIVK DMSLHQSFSG AQQDVEQLV YEVQNA*- (R₂)_n-Y

Polypeptides

The polypeptides of the invention include the polypeptide of Table 1 [SEQ ID NO:2] (in particular the mature polypeptide) as well as polypeptides and fragments, particularly those which have the biological activity of hisS, and also those which have at least 70% identity to the polypeptide of Table 1 [SEQ ID NO:2] or the relevant portion, preferably at least 80% identity to the polypeptide of Table 1 [SEQ ID NO:2], and more preferably at least 90% similarity (more preferably at least 90% identity) to the polypeptide of Table 1 [SEQ ID NO:2] and still more

preferably at least 95% similarity (still more preferably at least 95% identity) to the polypeptide of Table 1 [SEQ ID NO:2] and also include portions of such polypeptides with such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids.

5 The invention also includes polypeptides of the formula set forth in Table 1 (D) wherein, at the amino terminus, X is hydrogen, and at the carboxyl terminus, Y is hydrogen or a metal, R₁ and R₂ is any amino acid residue, and n is an integer between 1 and 1000. Any stretch of amino acid residues denoted by either R group, where R is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

10 A fragment is a variant polypeptide having an amino acid sequence that entirely is the same as part but not all of the amino acid sequence of the aforementioned polypeptides. As with hisS polypeptides fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region, a single larger polypeptide.

15 Preferred fragments include, for example, truncation polypeptides having a portion of the amino acid sequence of Table 1 [SEQ ID NO:2], or of variants thereof, such as a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus. Degradation forms of the polypeptides of the invention in a host cell, particularly a *Chlamydia trachomatis*, are also preferred. Further preferred are fragments
20 characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

25 Also preferred are biologically active fragments which are those fragments that mediate activities of hisS, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those fragments that are antigenic or immunogenic in an animal, especially in a human. Particularly preferred are fragments comprising receptors or domains of enzymes that confer a function essential for viability of
30 *Chlamydia trachomatis* or the ability to initiate, or maintain cause disease in an individual, particularly a human.

Variants that are fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these variants may be employed as intermediates for producing the full-length polypeptides of the invention.

5 **Polynucleotides**

Another aspect of the invention relates to isolated polynucleotides, including the full length gene, that encode the hisS polypeptide having the deduced amino acid sequence of Table 1 [SEQ ID NO:2] and polynucleotides closely related thereto and variants thereof.

Using the information provided herein, such as the polynucleotide sequence set out in
10 Table 1 [SEQ ID NO:1], a polynucleotide of the invention encoding hisS polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and sequencing chromosomal DNA fragments from bacteria using *Chlamydia trachomatis* D/UW-3/Cx cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide sequence of the invention, such as the sequence given in Table 1 [SEQ ID
15 NO:1], typically a library of clones of chromosomal DNA of *Chlamydia trachomatis* D/UW-3/Cx in *E.coli* or some other suitable host is probed with a radiolabeled oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using stringent conditions. By sequencing the individual clones thus identified with sequencing primers designed from the original sequence
20 it is then possible to extend the sequence in both directions to determine the full gene sequence. Conveniently, such sequencing is performed using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are described by Maniatis, T., Fritsch, E.F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see in particular
25 Screening By Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Illustrative of the invention, the polynucleotide set out in Table 1 [SEQ ID NO:1] was discovered in a DNA library derived from *Chlamydia trachomatis* D/UW-3/Cx.

The DNA sequence set out in Table 1 [SEQ ID NO:1] contains an open reading frame encoding a protein having about the number of amino acid residues set forth in Table 1 [SEQ ID
30 NO:2] with a deduced molecular weight that can be calculated using amino acid residue molecular weight values well known in the art. The polynucleotide of SEQ ID NO: 1, between

nucleotide number 1 through number 1308 encodes the polypeptide of SEQ ID NO:2. The stop codon begins at nucleotide number 1309 of SEQ ID NO:1.

The hisS protein of the invention is structurally related to other proteins of the histidyl tRNA synthetase family, as shown by the results of sequencing the DNA encoding hisS of the strain of the invention. The protein exhibits greatest homology to *Streptococcus equisimilis* histidyl tRNA synthetase protein among known proteins. The hisS protein of Table 1 [SEQ ID NO:2] has about 40% identity over its entire length and about 58% similarity over its entire length with the amino acid sequence of *Streptococcus equisimilis* histidyl tRNA synthetase polypeptide.

The invention provides a polynucleotide sequence identical over its entire length to the coding sequence in Table 1 [SEQ ID NO:1]. Also provided by the invention is the coding sequence for the mature polypeptide or a fragment thereof, by itself as well as the coding sequence for the mature polypeptide or a fragment in reading frame with other coding sequence, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence. The polynucleotide may also contain non-coding sequences, including for example, but not limited to non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences, termination signals, ribosome binding sites, sequences that stabilize mRNA, introns, polyadenylation signals, and additional coding sequence which encode additional amino acids. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain embodiments of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), or an HA tag (Wilson *et al.*, *Cell* 37: 767 (1984). Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene expression.

A preferred embodiment of the invention is the polynucleotide of comprising nucleotide 1 to 1308 set forth in SEQ ID NO:1 of Table 1 which encodes the hisS polypeptide.

The invention also includes polynucleotides of the formula set forth in Table 1 (C) wherein, at the 5' end of the molecule, X is hydrogen, and at the 3' end of the molecule, Y is hydrogen or a metal, R₁ and R₂ is any nucleic acid residue, and n is an integer between 1 and 1000. Any stretch of nucleic acid residues denoted by either R group, where R is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

The term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Chlamydia trachomatis* hisS having the amino acid sequence set out in Table 1 [SEQ ID NO:2]. The term also encompasses
5 polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, interrupted by integrated phage or an insertion sequence or editing) together with additional regions, that also may contain coding and/or non-coding sequences.

The invention further relates to variants of the polynucleotides described herein that encode for variants of the polypeptide having the deduced amino acid sequence of Table 1 [SEQ
10 ID NO:2]. Variants that are fragments of the polynucleotides of the invention may be used to synthesize full-length polynucleotides of the invention.

Further particularly preferred embodiments are polynucleotides encoding hisS variants, that have the amino acid sequence of hisS polypeptide of Table 1 [SEQ ID NO:2] in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, deleted or
15 added, in any combination. Especially preferred among these are silent substitutions, additions and deletions, that do not alter the properties and activities of hisS.

Further preferred embodiments of the invention are polynucleotides that are at least 70% identical over their entire length to a polynucleotide encoding hisS polypeptide having the amino acid sequence set out in Table 1 [SEQ ID NO:2], and polynucleotides that are complementary to
20 such polynucleotides. Alternatively, most highly preferred are polynucleotides that comprise a region that is at least 80% identical over its entire length to a polynucleotide encoding hisS polypeptide of the strain and polynucleotides complementary thereto. In this regard, polynucleotides at least 90% identical over their entire length to the same are particularly preferred, and among these particularly preferred polynucleotides, those with at least 95% are
25 especially preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.

Preferred embodiments are polynucleotides that encode polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by the
30 DNA of Table 1 [SEQ ID NO:1].

The invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the invention especially relates to polynucleotides that hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the terms "stringent conditions" and "stringent hybridization conditions" mean hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences. An example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at about 65°C. Hybridization and wash conditions are well known and exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y., (1989), particularly Chapter 11 therein.

The invention also provides a polynucleotide consisting essentially of a polynucleotide sequence obtainable by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in SEQ ID NO:1 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO:1 or a fragment thereof; and isolating said DNA sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers described elsewhere herein.

As discussed additionally herein regarding polynucleotide assays of the invention, for instance, polynucleotides of the invention as discussed above, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding hisS and to isolate cDNA and genomic clones of other genes that have a high sequence similarity to the hisS gene. Such probes generally will comprise at least 15 bases. Preferably, such probes will have at least 30 bases and may have at least 50 bases. Particularly preferred probes will have at least 30 bases and will have 50 bases or less.

For example, the coding region of the hisS gene may be isolated by screening using the DNA sequence provided in SEQ ID NO: 1 to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

The polynucleotides and polypeptides of the invention may be employed, for example, as research reagents and materials for discovery of treatments of and diagnostics for disease, particularly human disease, as further discussed herein relating to polynucleotide assays.

5 Polynucleotides of the invention that are oligonucleotides derived from the sequences of SEQ ID NOS:1 and/or 2 may be used in the processes herein as described, but preferably for PCR, to determine whether or not the polynucleotides identified herein in whole or in part are transcribed in bacteria in infected tissue. It is recognized that such sequences will also have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

10 The invention also provides polynucleotides that may encode a polypeptide that is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in vivo*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

15 A precursor protein, having the mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

20 In sum, a polynucleotide of the invention may encode a mature protein, a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences that are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or more prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

Vectors, host cells, expression

25 The invention also relates to vectors that comprise a polynucleotide or polynucleotides of the invention, host cells that are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques. Cell-free translation

systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a polynucleotide into the host cell can be effected by methods described in many standard laboratory manuals, such as Davis et al., *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection.

Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, enterococci *E. coli*, streptomyces and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal, episomal and virus-derived vectors, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, (*supra*).

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be

incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

Polypeptides of the invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid
5 extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography, and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding protein may be employed to regenerate active conformation when the polypeptide is denatured during
10 isolation and or purification.

Diagnostic Assays

This invention is also related to the use of the hisS polynucleotides of the invention for use as diagnostic reagents. Detection of hisS in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of a disease. Eukaryotes
15 (herein also "individual(s)"), particularly mammals, and especially humans, infected with an organism comprising the hisS gene may be detected at the nucleic acid level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from an infected individual's cells and tissues, such as bone, blood, muscle, cartilage, and skin. Genomic DNA may be used directly for
20 detection or may be amplified enzymatically by using PCR or other amplification technique prior to analysis. RNA or cDNA may also be used in the same ways. Using amplification, characterization of the species and strain of prokaryote present in an individual, may be made by an analysis of the genotype of the prokaryote gene. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the genotype of a reference sequence.
25 Point mutations can be identified by hybridizing amplified DNA to labeled hisS polynucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in the electrophoretic mobility of the DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. See, *e.g.*, Myers et al., *Science*, 230:
30 1242 (1985). Sequence changes at specific locations also may be revealed by nuclease protection

assays, such as RNase and S1 protection or a chemical cleavage method. See, e.g., Cotton et al., *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985).

5 Cells carrying mutations or polymorphisms in the gene of the invention may also be detected at the DNA level by a variety of techniques, to allow for serotyping, for example. For example, RT-PCR can be used to detect mutations. It is particularly preferred to use RT-PCR in conjunction with automated detection systems, such as, for example, GeneScan. RNA or cDNA may also be used for the same purpose, PCR or RT-PCR. As an example, PCR primers complementary to a nucleic acid encoding hisS can be used to identify and analyze mutations. 10 These primers may be used for, among other things, amplifying hisS DNA isolated from a sample derived from an individual. The primers may be used to amplify the gene isolated from an infected individual such that the gene may then be subject to various techniques for elucidation of the DNA sequence. In this way, mutations in the DNA sequence may be detected and used to diagnose infection and to serotype and/or classify the infectious agent.

15 The invention further provides a process for diagnosing, disease, preferably bacterial infections, more preferably infections by *Chlamydia trachomatis*, and most preferably classic ocular trachoma, inclusion conjunctivitis, genital trachoma, infant pneumonitis, Lymphogranuloma Venereum, incipient trachoma, keratitis, papillary hypertrophy, corneal infiltration, vulvovaginitis, ear infection, mucopurulent rhinitis, salpingitis, cervicitis, cervical 20 follicles, prostatitis, proctitis, urethritis, lymphogranule inguinale, climatic bubo, tropical bubo, and esthiomene., comprising determining from a sample derived from an individual a increased level of expression of polynucleotide having the sequence of Table 1 [SEQ ID NO: 1]. Increased or decreased expression of hisS polynucleotide can be measured using any one of the methods well known in the art for the quantitation of polynucleotides, such as, for example, 25 amplification, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods.

In addition, a diagnostic assay in accordance with the invention for detecting over-expression of hisS protein compared to normal control tissue samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a 30 hisS protein, in a sample derived from a host are well-known to those of skill in the art. Such

assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Antibodies

5 The polypeptides of the invention or variants thereof, or cells expressing them can be used as an immunogen to produce antibodies immunospecific for such polypeptides. "Antibodies" as used herein includes monoclonal and polyclonal antibodies, chimeric, single chain, simianized antibodies and humanized antibodies, as well as Fab fragments, including the products of an Fab immunoglobulin expression library.

10 Antibodies generated against the polypeptides of the invention can be obtained by administering the polypeptides or epitope-bearing fragments, analogues or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pg. 77-15 96 in *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms such as other mammals, may be used to express humanized antibodies.

20 Alternatively phage display technology may be utilized to select antibody genes with binding activities towards the polypeptide either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-hisS or from naive libraries (McCafferty, J. *et al.*, (1990), *Nature* 348, 552-554; Marks, J. *et al.*, (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by chain shuffling (Clackson, T. *et al.*, (1991) *Nature* 352, 624-628).

25 If two antigen binding domains are present each domain may be directed against a different epitope - termed 'bispecific' antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptides to purify the polypeptides by affinity chromatography.

30 Thus, among others, antibodies against hisS- polypeptide may be employed to treat infections, particularly bacterial infections and especially classic ocular trachoma, inclusion conjunctivitis, genital trachoma, infant pneumonitis, Lymphogranuloma Venerium, incipient

trachoma, keratitis, papillary hypertrophy, corneal infiltration, vulvovaginitis, ear infection, mucopurulent rhinitis, salpingitis, cervicitis, cervical follicles, prostatitis, proctitis, urethritis, lymphogranule inguinale, climatic bubo, tropical bubo, and esthiomene..

Polypeptide variants include antigenically, epitopically or immunologically equivalent
5 variants that form a particular aspect of this invention. The term "antigenically equivalent
derivative" as used herein encompasses a polypeptide or its equivalent which will be
specifically recognized by certain antibodies which, when raised to the protein or polypeptide
according to the invention, interfere with the immediate physical interaction between pathogen
and mammalian host. The term "immunologically equivalent derivative" as used herein
10 encompasses a peptide or its equivalent which when used in a suitable formulation to raise
antibodies in a vertebrate, the antibodies act to interfere with the immediate physical
interaction between pathogen and mammalian host.

The polypeptide, such as an antigenically or immunologically equivalent derivative or
a fusion protein thereof is used as an antigen to immunize a mouse or other animal such as a
15 rat or chicken. The fusion protein may provide stability to the polypeptide. The antigen may
be associated, for example by conjugation, with an immunogenic carrier protein for example
bovine serum albumin (BSA) or keyhole limpet haemocyanin (KLH). Alternatively a multiple
antigenic peptide comprising multiple copies of the protein or polypeptide, or an antigenically
or immunologically equivalent polypeptide thereof may be sufficiently antigenic to improve
20 immunogenicity so as to obviate the use of a carrier.

Preferably, the antibody or variant thereof is modified to make it less immunogenic in
the individual. For example, if the individual is human the antibody may most preferably be
"humanized"; where the complementarity determining region(s) of the hybridoma-derived
antibody has been transplanted into a human monoclonal antibody, for example as described
25 in Jones, P. et al. (1986), *Nature* 321, 522-525 or Tempest et al., (1991) *Biotechnology* 9, 266-
273.

The use of a polynucleotide of the invention in genetic immunization will preferably
employ a suitable delivery method such as direct injection of plasmid DNA into muscles
(Wolff et al., *Hum Mol Genet* 1992, 1:363, Manthorpe et al., *Hum. Gene Ther.* 1993:4, 419),
30 delivery of DNA complexed with specific protein carriers (Wu et al., *J Biol Chem.* 1989:
264,16985), coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef, *PNAS*

USA, 1986:83,9551), encapsulation of DNA in various forms of liposomes (Kaneda et al., *Science* 1989:243,375), particle bombardment (Tang et al., *Nature* 1992, 356:152, Eisenbraun et al., *DNA Cell Biol* 1993, 12:791) and *in vivo* infection using cloned retroviral vectors (Seeger et al., *PNAS USA* 1984:81,5849).

5 **Antagonists and agonists - assays and molecules**

Polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See, e.g., Coligan et al., *Current Protocols in Immunology 1(2)*:
10 Chapter 5 (1991).

The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of hisS polypeptides or polynucleotides, particularly those compounds that are bacteriostatic and/or bacteriocidal. The method of screening may involve high-throughput techniques. For example, to screen for agonists or
15 antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, comprising hisS polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence or the presence of a candidate molecule that may be a hisS agonist or antagonist. The ability of the candidate molecule to agonize or antagonize the hisS polypeptide is reflected in decreased binding of the labeled ligand or
20 decreased production of product from such substrate. Molecules that bind gratuitously, *i.e.*, without inducing the effects of hisS polypeptide are most likely to be good antagonists. Molecules that bind well and increase the rate of product production from substrate are agonists. Detection of the rate or level of production of product from substrate may be enhanced by using a reporter system. Reporter systems that may be useful in this regard include but are not limited to
25 colorimetric labeled substrate converted into product, a reporter gene that is responsive to changes in hisS polynucleotide or polypeptide activity, and binding assays known in the art.

Another example of an assay for hisS antagonists is a competitive assay that combines hisS and a potential antagonist with hisS-binding molecules, recombinant hisS binding molecules, natural substrates or ligands, or substrate or ligand mimetics, under appropriate conditions for a
30 competitive inhibition assay. The hisS molecule can be labeled, such as by radioactivity or a colorimetric compound, such that the number of hisS molecules bound to a binding molecule or

converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

Potential antagonists include small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide or polypeptide of the invention and thereby inhibit or
5 extinguish its activity. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing hisS-induced activities, thereby preventing the action of hisS by excluding hisS from binding.

Potential antagonists include a small molecule that binds to and occupies the binding site
10 of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, FL (1988), for
15 a description of these molecules). Preferred potential antagonists include compounds related to and variants of hisS.

Each of the DNA sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein, upon expression, can be used as
20 a target for the screening of antibacterial drugs. Additionally, the DNA sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

The invention also provides the use of the polypeptide, polynucleotide or inhibitor of
25 the invention to interfere with the initial physical interaction between a pathogen and mammalian host responsible for sequelae of infection. In particular the molecules of the invention may be used: in the prevention of adhesion of bacteria, in particular gram positive bacteria, to mammalian extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block hisS protein-mediated mammalian cell invasion by, for
30 example, initiating phosphorylation of mammalian tyrosine kinases (Rosenshine *et al.*, *Infect. Immun.* 60:2211 (1992); to block bacterial adhesion between mammalian extracellular matrix proteins and bacterial hisS proteins that mediate tissue damage and; to block the normal

progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

The antagonists and agonists of the invention may be employed, for instance, to inhibit and treat classic ocular trachoma, inclusion conjunctivitis, genital trachoma, infant pneumonitis, Lymphogranuloma Venerium, incipient trachoma, keratitis, papillary hypertrophy, corneal infiltration, vulvovaginitis, ear infection, mucopurulent rhinitis, salpingitis, cervicitis, cervical follicles, prostatitis, proctitis, urethritis, lymphogranule inguinale, climatic bubo, tropical bubo, and esthiomene..

Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal which comprises inoculating the individual with hisS, or a fragment or variant thereof, adequate to produce antibody and/ or T cell immune response to protect said individual from infection, particularly bacterial infection and most particularly *Chlamydia trachomatis* infection. Also provided are methods whereby such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises delivering to such individual a nucleic acid vector to direct expression of hisS, or a fragment or a variant thereof, for expressing hisS, or a fragment or a variant thereof *in vivo* in order to induce an immunological response, such as, to produce antibody and/ or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual from disease, whether that disease is already established within the individual or not. One way of administering the gene is by accelerating it into the desired cells as a coating on particles or otherwise.

Such nucleic acid vector may comprise DNA, RNA, a modified nucleic acid, or a DNA/RNA hybrid.

A further aspect of the invention relates to an immunological composition which, when introduced into an individual capable or having induced within it an immunological response, induces an immunological response in such individual to a hisS or protein coded therefrom, wherein the composition comprises a recombinant hisS or protein coded therefrom comprising DNA which codes for and expresses an antigen of said hisS or protein coded therefrom. The immunological response may be used therapeutically or prophylactically and may take the

form of antibody immunity or cellular immunity such as that arising from CTL or CD4+ T cells.

5 A hisS polypeptide or a fragment thereof may be fused with co-protein which may not by itself produce antibodies, but is capable of stabilizing the first protein and producing a fused protein which will have immunogenic and protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from *Hemophilus influenzae*, Glutathione-S-transferase (GST) or beta-galactosidase, relatively large co-proteins which solubilize the protein and facilitate production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system. The co-protein may be attached to either the amino or carboxy terminus of the first protein.

10 Provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides or polynucleotides of the invention and immunostimulatory DNA sequences, such as those described in Sato, Y. *et al. Science* 273: 352 (1996).

15 Also, provided by this invention are methods using the described polynucleotide or particular fragments thereof which have been shown to encode non-variable regions of bacterial cell surface proteins in DNA constructs used in such genetic immunization experiments in animal models of infection with *Chlamydia trachomatis* will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of monoclonal antibodies of particular value from the requisite organ of the animal successfully resisting or clearing infection for the development of prophylactic agents or therapeutic treatments of bacterial infection, particularly *Chlamydia trachomatis* infection, in mammals, particularly humans.

20 The polypeptide may be used as an antigen for vaccination of a host to produce specific antibodies which protect against invasion of bacteria, for example by blocking adherence of bacteria to damaged tissue. Examples of tissue damage include wounds in skin or connective tissue caused, e.g., by mechanical, chemical or thermal damage or by implantation of indwelling devices, or wounds in the mucous membranes, such as the mouth, mammary glands, urethra or vagina.

30

The invention also includes a vaccine formulation which comprises an immunogenic recombinant protein of the invention together with a suitable carrier. Since the protein may be broken down in the stomach, it is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

While the invention has been described with reference to certain hisS protein, it is to be understood that this covers fragments of the naturally occurring protein and similar proteins with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the recombinant protein.

Compositions, kits and administration

The invention also relates to compositions comprising the polynucleotide or the polypeptides discussed above or their agonists or antagonists. The polypeptides of the invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to a subject. Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide of the invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration. The invention further relates to diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides and other compounds of the invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous, 5 intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.

In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic.

Alternatively the composition may be formulated for topical application for example in the form of ointments, creams, lotions, eye ointments, eye drops, ear drops, 10 mouthwash, impregnated dressings and sutures and aerosols, and may contain appropriate conventional additives, including, for example, preservatives, solvents to assist drug penetration, and emollients in ointments and creams. Such topical formulations may also contain compatible conventional carriers, for example cream or ointment bases, and ethanol or oleyl alcohol for lotions. Such carriers may constitute from about 1% to about 98% by weight 15 of the formulation; more usually they will constitute up to about 80% by weight of the formulation.

For administration to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1 mg/kg. The physician in any event will determine the actual dosage which will be most 20 suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

In-dwelling devices include surgical implants, prosthetic devices and catheters, i.e., 25 devices that are introduced to the body of an individual and remain in position for an extended time. Such devices include, for example, artificial joints, heart valves, pacemakers, vascular grafts, vascular catheters, cerebrospinal fluid shunts, urinary catheters, continuous ambulatory peritoneal dialysis (CAPD) catheters.

The composition of the invention may be administered by injection to achieve a 30 systemic effect against relevant bacteria shortly before insertion of an in-dwelling device. Treatment may be continued after surgery during the in-body time of the device. In addition,

the composition could also be used to broaden perioperative cover for any surgical technique to prevent bacterial wound infections, especially *Chlamydia trachomatis* wound infections.

Many orthopaedic surgeons consider that humans with prosthetic joints should be considered for antibiotic prophylaxis before dental treatment that could produce a bacteremia.

5 Late deep infection is a serious complication sometimes leading to loss of the prosthetic joint and is accompanied by significant morbidity and mortality. It may therefore be possible to extend the use of the active agent as a replacement for prophylactic antibiotics in this situation.

10 In addition to the therapy described above, the compositions of this invention may be used generally as a wound treatment agent to prevent adhesion of bacteria to matrix proteins exposed in wound tissue and for prophylactic use in dental treatment as an alternative to, or in conjunction with, antibiotic prophylaxis.

Alternatively, the composition of the invention may be used to bathe an indwelling device immediately before insertion. The active agent will preferably be present at a concentration of 1µg/ml to 10mg/ml for bathing of wounds or indwelling devices.

15 A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5 microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to
20 suitable individuals.

Each reference disclosed herein is incorporated by reference herein in its entirety. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety.

25 **EXAMPLES**

The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

30 **Example 1 Strain selection, Library Production and Sequencing**

The polynucleotide having the DNA sequence given in SEQ ID NO:1 is obtained, for example from a library of clones of chromosomal DNA of *Chlamydia trachomatis* in *E. coli*. The sequencing data from two or more clones containing overlapping *Chlamydia trachomatis* DNAs is used to construct the contiguous DNA sequence in SEQ ID NO:1. Libraries may be prepared by routine methods, for example:

Methods 1, 2 and 3 below.

Total cellular DNA is isolated from *Chlamydia trachomatis* D/UW-3/Cx according to standard procedures and size-fractionated by either of two methods.

Method 1

Total cellular DNA is mechanically sheared by passage through a needle in order to size-fractionate according to standard procedures. DNA fragments of up to 11kbp in size are rendered blunt by treatment with exonuclease and DNA polymerase, and EcoRI linkers added. Fragments are ligated into the vector Lambda ZapII that has been cut with EcoRI, the library packaged by standard procedures and *E.coli* infected with the packaged library. The library is amplified by standard procedures.

Method 2

Total cellular DNA is partially hydrolyzed with a one or a combination of restriction enzymes appropriate to generate a series of fragments for cloning into library vectors (e.g., RsaI, PstI, AluI, BshI235I), and such fragments are size-fractionated according to standard procedures. EcoRI linkers are ligated to the DNA and the fragments then ligated into the vector Lambda ZapII that have been cut with EcoRI, the library packaged by standard procedures, and *E.coli* infected with the packaged library. The library is amplified by standard procedures.

Method 3

Total cellular DNA is mechanically or enzymatically fragmented to size-fractionate according to standard procedures. DNA fragments of about 1kbp in size, after preparing their ends using standard procedures, are ligated into M13 vector using standard procedures. M13 is introduced into *E.coli* host, such as NM522 (available commercially). Clones with inserts are sequenced using standard procedures.

Example 2 hisS Characterization

The enzyme mediated incorporation of radiolabelled amino acid into tRNA may be measured by the aminoacylation method which measures amino acid-tRNA as trichloroacetic acid-precipitable radioactivity from radiolabelled amino acid in the presence of tRNA and ATP (Hughes J, Mellows G and Soughton S, 1980, FEBS Letters, 122:322-324). Thus inhibitors of
5 histidyl tRNA synthetase can be detected by a reduction in the trichloroacetic acid precipitable radioactivity relative to the control. Alternatively the tRNA synthetase catalysed partial PPi/ATP exchange reaction which measures the formation of radiolabelled ATP from PPi can be used to detect histidyl tRNA synthetase inhibitors (Calender R & Berg P, 1966, Biochemistry, 5, 1681-1690).

10

SEQUENCE LISTING

5 (1) GENERAL INFORMATION:

(i) APPLICANT: Reichard, Raymond W.
Brown, James R.
Lawlor, Elizabeth J.

10 (ii) TITLE OF INVENTION: Novel hisS

(iii) NUMBER OF SEQUENCES: 2

15 (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Dechert, Price & Rhoads
(B) STREET: 4000 Bell Atlantic Tower, 1717 Arch Stre
(C) CITY: Philadelphia
20 (D) STATE: PA
(E) COUNTRY: USA
(F) ZIP: 19103-2793

(v) COMPUTER READABLE FORM:
25 (A) MEDIUM TYPE: Diskette
(B) COMPUTER: IBM Compatible
(C) OPERATING SYSTEM: Windows 95
(D) SOFTWARE: FastSEQ for Windows Version 2.0b

30 (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

35 (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE:

40 (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Falk, Stephen T
(B) REGISTRATION NUMBER: 36,795
(C) REFERENCE/DOCKET NUMBER: GM10048

45 (ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: 215-994-2488
(B) TELEFAX: 215-994-2222
50 (C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1311 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

GTGAGAATAG TCCGGTTATA CACAATGAGT AATGCATTGC CAAAAGGCGT TTTTGATATT      60
TTTCCCTATG TAACAAGCCC AAAAAATCTT TGGAGAAATT CTTCTCTATG GAAACGTGTT      120
GAGCATGCAG CCCATCGTAT TTGTAATTTA TATGGATTTG ATGAGATCCG AACTCCAGTT      180
TTTGAAAAGA CAGAGACTTT TTTACGCGTC GGAGAGTACA GTGATATTGT AAAAAAGGAA      240
GTTTATACCT TCTTAGATAA AAAAAGACGT TCTTTGACTT TCGGTCCAGA AGGGACTGCA      300
GCAGTTGTTT GTGCATTGTT GGATCATTCT GCTGATATGC GCAAAGATAA TAAGTTTTAT      360
TATATTTTGC CCATGTTTCG TTACGAGCGG CAACAATCTG GACGTTACCG TCAGCATCAT      420
CAGTTCGGTC TAGAAGCTAT CGGTGTGCGG CACCCCTTAC GAGATGCGGA GGTGCTCTCT      480
CTGTTATGGG ATTTTTATGC AGCGGTGCGG CTTCAGCATA TGCAAATCCA TGTGAATTTT      540
TTAGGAGGGC AAAAGACTCG GGCTCGTTAT GACGAAGCTT TCGGGGAGTT CTCCGTAAG      600
GATCTAGACC GGTATATCGC TCTGAGTCAA GAAAGATATC ATGCGAACTT ATTGCGTATA      660
TTAGATTCTA AGGAGCCAGA AGACCAGGAA TTCATTGAAA AAGCTCCCTC AATTTTGGAT      720
TACATAGATG ATCGGGATTT AAGCTATTTT GATGCAGTAT TAGCCCAATT AAAGGCTTTA      780
GGGATTTCTT TTGCAATCAA TCCAAGGCTA GTTCGAGGGT TGGATTATTA TACGGATCTT      840
GTATTTGAAG CGGTACTGTG TGTGGGAGAG CACTCCTATG CATTGGGAGG CGGCGGGCGT      900
TATGATGAAT TGGTTGCGCA GTCTGGAGGC CTTTCTATGC CAGCTTTTGG TTTTGGAGTG      960
GGATTGGAAA GAGTAATCCA AACGCTATTA GAGCAAGGAA ACTCTTTATC GACCTCTACG     1020
CGACGGTTGC GATTAATTC TATGGATGAG CAGGCAGATG CATTTTGTTT TTCATGGGCA     1080
AATCGTTTAC GTAACCTAGG CATTGCAACA GAAGTAGATT GGAGCCATAA GAAGCCTAAA     1140
TTGTCTCTTA AAGATGCTGC CGATCAACAA GTCAGTTTTG TTTGTCTCTT AGGCGAACAA     1200
GAATTAGCAA CGAAACAATT TATAGTTAAA GATATGTCTT TGCATCAAAG CTTCTCAGGG     1260
GCTCAACAAG ATGTAGAACA AAGGTTGGTT TATGAAGTAC AGAACGCATA A                1311
    
```

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 436 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Met Arg Ile Val Arg Leu Tyr Thr Met Ser Asn Ala Leu Pro Lys Gly
 1                5                10                15
Val Phe Asp Ile Phe Pro Tyr Val Thr Ser Pro Lys Asn Leu Trp Arg
 20                25                30
Asn Ser Ser Leu Trp Lys Arg Val Glu His Ala Ala His Arg Ile Cys
 35                40                45
    
```

	Asn	Leu	Tyr	Gly	Phe	Asp	Glu	Ile	Arg	Thr	Pro	Val	Phe	Glu	Lys	Thr
	50						55					60				
	Glu	Thr	Phe	Leu	Arg	Val	Gly	Glu	Tyr	Ser	Asp	Ile	Val	Lys	Lys	Glu
	65					70					75					80
5	Val	Tyr	Thr	Phe	Leu	Asp	Lys	Lys	Arg	Arg	Ser	Leu	Thr	Leu	Arg	Pro
					85					90					95	
	Glu	Gly	Thr	Ala	Ala	Val	Val	Arg	Ala	Leu	Leu	Asp	His	Ser	Ala	Asp
				100					105					110		
10	Met	Arg	Lys	Asp	Asn	Lys	Phe	Tyr	Tyr	Ile	Leu	Pro	Met	Phe	Arg	Tyr
			115					120					125			
	Glu	Arg	Gln	Gln	Ser	Gly	Arg	Tyr	Arg	Gln	His	His	Gln	Phe	Gly	Leu
			130				135					140				
15	Glu	Ala	Ile	Gly	Val	Arg	His	Pro	Leu	Arg	Asp	Ala	Glu	Val	Leu	Ser
					150						155					160
	Leu	Leu	Trp	Asp	Phe	Tyr	Ala	Ala	Val	Gly	Leu	Gln	His	Met	Gln	Ile
					165					170					175	
	His	Val	Asn	Phe	Leu	Gly	Gly	Gln	Lys	Thr	Arg	Ala	Arg	Tyr	Asp	Glu
				180					185					190		
20	Ala	Leu	Arg	Glu	Phe	Phe	Arg	Lys	Asp	Leu	Asp	Arg	Leu	Ser	Pro	Leu
			195					200					205			
	Ser	Gln	Glu	Arg	Tyr	His	Ala	Asn	Leu	Leu	Arg	Ile	Leu	Asp	Ser	Lys
			210				215					220				
25	Glu	Pro	Glu	Asp	Gln	Glu	Phe	Ile	Glu	Lys	Ala	Pro	Ser	Ile	Leu	Asp
					230						235					240
	Tyr	Ile	Asp	Asp	Arg	Asp	Leu	Ser	Tyr	Phe	Asp	Ala	Val	Leu	Ala	Gln
					245					250					255	
	Leu	Lys	Ala	Leu	Gly	Ile	Ser	Phe	Ala	Ile	Asn	Pro	Arg	Leu	Val	Arg
				260					265					270		
30	Gly	Leu	Asp	Tyr	Tyr	Thr	Asp	Leu	Val	Phe	Glu	Ala	Val	Thr	Val	Val
			275					280					285			
	Gly	Glu	His	Ser	Tyr	Ala	Leu	Gly	Gly	Gly	Gly	Arg	Tyr	Asp	Glu	Leu
			290				295					300				
35	Val	Ala	Gln	Ser	Gly	Gly	Pro	Ser	Met	Pro	Ala	Phe	Gly	Phe	Gly	Val
					310						315					320
	Gly	Leu	Glu	Arg	Val	Ile	Gln	Thr	Leu	Leu	Glu	Gln	Gly	Asn	Ser	Leu
					325						330				335	
	Ser	Thr	Ser	Thr	Arg	Arg	Leu	Arg	Leu	Ile	Pro	Met	Asp	Glu	Gln	Ala
				340					345					350		
40	Asp	Ala	Phe	Cys	Phe	Ser	Trp	Ala	Asn	Arg	Leu	Arg	Asn	Leu	Gly	Ile
			355					360					365			
	Ala	Thr	Glu	Val	Asp	Trp	Ser	His	Lys	Lys	Pro	Lys	Leu	Ser	Leu	Lys
							375					380				
45	Asp	Ala	Ala	Asp	Gln	Gln	Val	Ser	Phe	Val	Cys	Leu	Leu	Gly	Glu	Gln
					390						395					400
	Glu	Leu	Ala	Thr	Lys	Gln	Phe	Ile	Val	Lys	Asp	Met	Ser	Leu	His	Gln
					405						410				415	
	Ser	Phe	Ser	Gly	Ala	Gln	Gln	Asp	Val	Glu	Gln	Arg	Leu	Val	Tyr	Glu
				420					425					430		
50	Val	Gln	Asn	Ala												
			435													

What is claimed is:

1. An isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of:
 - (a) a polynucleotide having at least a 70% identity to a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:2;
 - (b) a polynucleotide having at least a 70% identity to a polynucleotide encoding the same mature polypeptide expressed by the hisS gene contained in the *Chlamydia trachomatis* of the strain of the invention;
 - (c) a polynucleotide encoding a polypeptide comprising an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID NO:2;
 - (d) a polynucleotide which is complementary to the polynucleotide of (a), (b) or (c);and
 - (e) a polynucleotide comprising at least 15 sequential bases of the polynucleotide of (a), (b) or (c).
2. The polynucleotide of Claim 1 wherein the polynucleotide is DNA.
3. The polynucleotide of Claim 1 wherein the polynucleotide is RNA.
4. The polynucleotide of Claim 2 comprising the nucleic acid sequence set forth in SEQ ID NO:1.
5. The polynucleotide of Claim 2 comprising nucleotide 1 to 1308 set forth in SEQ ID NO:1.
6. The polynucleotide of Claim 2 which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2.
7. A vector comprising the polynucleotide of Claim 1.
8. A host cell comprising the vector of Claim 7.
9. A process for producing a polypeptide comprising: expressing from the host cell of Claim 8 a polypeptide encoded by said DNA.
10. A process for producing a hisS polypeptide or fragment comprising culturing a host of claim 8 under conditions sufficient for the production of said polypeptide or fragment.
11. A polypeptide comprising an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID NO:2.
12. A polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2.

13. An antibody against the polypeptide of claim 11.
14. An antagonist which inhibits the activity or expression of the polypeptide of claim 11.
15. A method for the treatment of an individual in need of hisS polypeptide comprising: administering to the individual a therapeutically effective amount of the polypeptide of claim 11.
16. A method for the treatment of an individual having need to inhibit hisS polypeptide comprising: administering to the individual a therapeutically effective amount of the antagonist of Claim 14.
17. A process for diagnosing a disease related to expression or activity of the polypeptide of claim 11 in an individual comprising:
 - (a) determining a nucleic acid sequence encoding said polypeptide, and/or
 - (b) analyzing for the presence or amount of said polypeptide in a sample derived from the individual.
18. A method for identifying compounds which interact with and inhibit or activate an activity of the polypeptide of claim 11 comprising:

contacting a composition comprising the polypeptide with the compound to be screened under conditions to permit interaction between the compound and the polypeptide to assess the interaction of a compound, such interaction being associated with a second component capable of providing a detectable signal in response to the interaction of the polypeptide with the compound;

and determining whether the compound interacts with and activates or inhibits an activity of the polypeptide by detecting the presence or absence of a signal generated from the interaction of the compound with the polypeptide.
19. A method for inducing an immunological response in a mammal which comprises inoculating the mammal with hisS polypeptide of claim 11, or a fragment or variant thereof, adequate to produce antibody and/or T cell immune response to protect said animal from disease.
20. A method of inducing immunological response in a mammal which comprises delivering a nucleic acid vector to direct expression of hisS polypeptide of claim 11, or fragment or a variant thereof, for expressing said hisS polypeptide, or a fragment or a variant thereof *in*

vivo in order to induce an immunological response to produce antibody and/ or T cell immune response to protect said animal from disease.

21. The use of a therapeutically effective amount of the polypeptide of claim 11 to treat an individual in need of hisS polypeptide.

5 22. The use of a therapeutically effective amount of the antagonist of claim 14 to treat an individual having need to inhibit hisS polypeptide.

23. The use of hisS polypeptide of claim 11, or a fragment or variant thereof, adequate to produce antibody and/or T cell immune response to induce an immunological response in a mammal to protect said mammal from disease.

10

24. The use of a nucleic acid vector to direct expression of hisS polypeptide of claim 11, or fragment or a variant thereof, for expressing said hisS polypeptide, or a fragment or a variant thereof *in vivo* in order to induce an immunological response in a mammal to produce antibody and/or T cell immune response to protect said mammal from disease.