



(51) International Patent Classification:

A23L 1/29 (2006.01) A61K 35/20 (2006.01)
A23L 1/308 (2006.01) A61P 25/00 (2006.01)
A61K 31/702 (2006.01)

(21) International Application Number:

PCT/US2013/059436

(22) International Filing Date:

12 September 2013 (12.09.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

12382354.4 14 September 2012 (14.09.2012) EP

(71) Applicant: ABBOTT LABORATORIES [US/US]; Dept.
377/AP6A-1, 100 Abbott Park Road, Abbott Park, Illinois
60064 (US).

(72) Inventors: HERNANDEZ, Enrique Vazquez; Camino de
Gojar 42, Urb Los Nenufares, E-18151 52 Ogijares (ES).
CABRERA, Ricardo Rueda; C/Brazal, nº 2, E-18008
Granada (ES). BUCK, Rachael; 3710 Peak Ridge Drive,
Gahanna, Ohio 43230 (US). GONZALEZ, Maria
Ramirez; C/Ramon y Cajal 54, Portal 2, 4A, E-18003
Granada (ES).

(74) Agents: ENGLE, Mark R. et al.; Abbott Nutrition a Division
of Abbott Laboratories, 3300 Stelzer Road, Dept
108140 RP3-2, Columbus, Ohio 43219 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,
ZW.


(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: METHODS FOR MODULATING CORTICOSTERONE LEVELS IN PSYCHOLOGICALLY STRESSED INDIVIDUALS

FIG. 2

SERUM CORTICOSTERONE POST RESTRAINT



(57) Abstract: Disclosed are methods of modulating and/or decreasing serum corticosterone levels in an individual affected by stress. Further disclosed are methods of modulating the hypothalamic pituitary adrenal response in an individual. The methods include administration of 2-fucosyl-lactose to an individual.

WO 2014/043330 A1



Published:

— *with international search report (Art. 21(3))*

**METHODS FOR MODULATING CORTICOSTERONE LEVELS IN
PSYCHOLOGICALLY STRESSED INDIVIDUALS**

CROSS-REFERENCE TO RELATED APPLICATION

[0001] This application claims priority to and any other benefit of EP Application No. 12382354.4, filed September 14, 2012, and entitled “METHODS FOR MODULATING CORTICOSTERONE LEVELS IN PSYCHOLOGICALLY STRESSED INDIVIDUALS,” the entire disclosure of which is incorporated by reference herein.

FIELD OF THE DISCLOSURE

[0002] The present disclosure relates to human milk oligosaccharides for reducing exacerbated stress response in individuals, particularly individuals who were affected by acute psychological stress early in life. More particularly, the present disclosure relates to methods of using fucosylated human milk oligosaccharides, and in particular, 2-fucosyl-lactose (2FL), to modulate and/or decrease exacerbated serum corticosterone levels and/or modulate the hypothalamic pituitary adrenal response in an individual affected by stress.

BACKGROUND OF THE DISCLOSURE

[0003] The body's ability to keep a steady homeostatic state is crucial to the health and life of an individual. This involves providing an adequate response to a variety of challenges both physical and mental, such as microbial invasion and emotional distress. Interplay between the neuroendocrine and immune systems is essential in either case. The hypothalamic-pituitary-adrenal (HPA) axis is responsible for initiation of glucocorticoid stress responses in all vertebrate animals.

[0004] Further, there are various mechanisms that modulate HPA activity, particularly by stimulating and inhibiting stress responses, as a hyperactivated HPA response leads to an imbalanced physiological system in the individual, thereby negatively affecting the

immune and gastrointestinal systems, metabolic rates, as well as cognitive function. An imbalanced physiological system can further induce mood disorders and depression.

[0005] A classical view of the neuroendocrine-immune network includes bidirectional interactions in which pro-inflammatory cytokines influence HPA axis-derived hormones (e.g., corticosterone, cortisol, aldosterone) that subsequently affect those cytokines in a feedback mechanism. These cytokines and other hormones are functionally expressed in the brain as well as in other peripheral organs, including immune cells. Because of this intermingled network of molecules that are redundantly expressed, the elucidation of the unique roles of HPA axis-related molecules is one of particular interest.

[0006] Early life stress has been reported to induce long lasting, and in some cases, permanent changes in the central nervous system, inducing hyperactivated HPA responses. This leads to exacerbated stress responses in further stressful situations, as well as induces alterations in behavior, such as increasing anxiety-related responses, aggressiveness, future maternal care abilities, and the like. Early life stress also affects other central nervous system parameters such as the number of neurons in the hippocampus, myelination processes and neuroendocrine responses, as well as other bodily functions, including lipid metabolism and inflammatory responses.

[0007] It would therefore be desirable to provide compositions that provide individual components that will reduce hyperactivated HPA responses in individuals. It would further be beneficial to provide nutritional compositions that modulate and/or decrease exacerbated production of serum corticosterone and/or modulate HPA responses in individuals affected by stress.

SUMMARY OF THE DISCLOSURE

[0008] The present disclosure is directed to methods of reducing stress response in animals, including in individuals, including infants, pediatrics, adults, and older adults, using human milk oligosaccharides, and in particular, 2-fucosyl-lactose. 2-fucosyl-lactose modulates the hypothalamic pituitary adrenal response, thereby decreasing serum corticosterone levels in individuals affected by acute psychological stress.

[0009] In one embodiment, the present disclosure is directed to a method of modulating serum corticosterone levels in an animal. The method comprises administering to the animal a composition comprising 2-fucosyl-lactose.

[0010] In another embodiment, the present disclosure is directed to a method of decreasing serum corticosterone levels in an individual affected by stress. The method comprises administering to the individual a nutritional composition comprising 2-fucosyl-lactose.

[0011] In another embodiment the present disclosure is directed to a method of modulating hypothalamic pituitary adrenal response in an individual. The method comprises administering to the individual a composition comprising 2-fucosyl-lactose.

[0012] It has been unexpectedly discovered that human milk oligosaccharides, and particularly, 2-fucosyl-lactose, can reduce exacerbated stress response in an individual by mediating hyperactive hypothalamic pituitary adrenal (HPA) responses and decreasing serum corticosterone levels in individuals affected by stress, particularly acute psychological stress. As overproduction of corticosterone is closely related to central nervous system function (e.g., cognitive skills, mood disorders, depression), administration of 2-fucosyl-lactose may further promote central nervous system function, especially in stressful situations that may produce mood disorders or depression. These benefits are advantageously achieved without the complications seen with previously used oral synthetic pharmacological approaches.

BRIEF DESCRIPTION OF THE DRAWINGS

[0013] Figure 1 is a graph depicting basal serum corticosterone levels in mice as analyzed in Example 1.

[0014] Figure 2 is a graph depicting post restraint serum corticosterone levels in mice as analyzed in Example 1.

DETAILED DESCRIPTION OF THE DISCLOSURE

[0015] The present disclosure is directed to methods for modulating and/or decreasing serum corticosterone levels and/or modulating the hypothalamic pituitary adrenal response in an individual, thereby maintaining a balanced physiological state. The present

methods include administering 2-fucosyl-lactose to an individual affected by stress, particularly stress early in life, to modulate and/or decrease serum corticosterone levels in response to stress in an individual, and further to prevent/control/reduce and/or treat hyperactivation of the HPA, and to prevent/control/reduce and/or treat malfunctions in the neuroendocrine-immune network. The methods may be useful in maintaining a healthy central nervous system, as well as have implications in the treatment of inflammation and cognitive disorders.

[0016] These and other features of the compositions and methods, as well as some of the many optional variations and additions, are described in detail hereafter.

[0017] The term “animal” as used herein refers to mammals in general, including humans.

[0018] The terms “acute psychological stress” and “acute stress” as used herein, unless otherwise specified, are used interchangeably to refer to a psychological condition (e.g., feeling of strain, pressure, anxiety, being overwhelmed, irritability, nervousness, insecurity, depression, panic, exhaustion) arising in response to a terrifying or traumatic event. A “terrifying event” or “traumatic event” is an experience that causes the individual to experience disturbing or unexpected fear, stress or pain.

[0019] The terms “early stress” or “stress early in life” as used herein, unless otherwise specified, are used interchangeably to refer to the experience of stress early in an individual’s life; that is, during the period ranging from birth to early adolescence. “Early adolescence” refers to the period of from 10 years to 14 years of life.

[0020] The terms “retort packaging” and “retort sterilizing” are used interchangeably herein, and unless otherwise specified, refer to the common practice of filling a container, most typically a metal can or other similar package, with a nutritional liquid and then subjecting the liquid-filled package to the necessary heat sterilization step, to form a sterilized, retort packaged, nutritional liquid product.

[0021] The term “aseptic packaging” as used herein, unless otherwise specified, refers to the manufacture of a packaged product without reliance upon the above-described retort packaging step, wherein the nutritional liquid and package are sterilized separately prior to

filling, and then are combined under sterilized or aseptic processing conditions to form a sterilized, aseptically packaged, nutritional liquid product.

[0022] The terms “fat” and “oil” as used herein, unless otherwise specified, are used interchangeably to refer to lipid materials derived or processed from plants or animals. These terms also include synthetic lipid materials so long as such synthetic materials are suitable for oral administration to humans.

[0023] The term “human milk oligosaccharide” or “HMO”, as used herein, unless otherwise specified, refers generally to a number of complex carbohydrates found in human breast milk that can be in acidic or neutral form, and to precursors thereof. Exemplary non-limiting human milk oligosaccharides include 2-fucosyl-lactose, 3-fucosyl-lactose, 3-sialyl-lactose, 6-sialyl-lactose, and lacto-N-neo-tetraose. Exemplary human milk oligosaccharide precursors include sialic acid and/or fucose.

[0024] The term “shelf stable” as used herein, unless otherwise specified, refers to a nutritional product that remains commercially stable after being packaged and then stored at 18-24°C for at least 3 months, including from about 6 months to about 24 months, and also including from about 12 months to about 18 months.

[0025] The terms “nutritional formulation” or “nutritional composition” as used herein, are used interchangeably and, unless otherwise specified, refer to synthetic formulas including nutritional liquids, nutritional powders, nutritional solids, nutritional semi-solids, nutritional semi-liquids, nutritional supplements, and any other nutritional food product as known in the art. The nutritional powders may be reconstituted to form a nutritional liquid, all of which comprise one or more of fat, protein and carbohydrate and are suitable for oral consumption by a human.

[0026] The term “nutritional liquid” as used herein, unless otherwise specified, refers to nutritional products in ready-to-drink liquid form, concentrated form, and nutritional liquids made by reconstituting the nutritional powders described herein prior to use.

[0027] The term “nutritional powder” as used herein, unless otherwise specified, refers to nutritional products in flowable or scoopable form that can be reconstituted with water or another aqueous liquid prior to consumption and includes both spray dried and drymixed/dryblended powders.

[0028] The term “nutritional semi-solid,” as used herein, unless otherwise specified, refers to nutritional products that are intermediate in properties, such as rigidity, between solids and liquids. Some semi-solids examples include puddings, gelatins, and doughs.

[0029] The term “nutritional semi-liquid,” as used herein, unless otherwise specified, refers to nutritional products that are intermediate in properties, such as flow properties, between liquids and solids. Some semi-liquids examples include thick shakes and liquid gels.

[0030] The terms “susceptible” and “at risk” as used herein, unless otherwise specified, mean having little resistance to a certain condition or disease, including being genetically predisposed, having a family history of, and/or having symptoms of the condition or disease.

[0031] The terms “modulating” or “modulation” or “modulate” as used herein, unless otherwise specified, refer to the targeted movement of a selected characteristic.

[0032] All percentages, parts and ratios as used herein, are by weight of the total composition, unless otherwise specified. All such weights, as they pertain to listed ingredients, are based on the active level and, therefore, do not include solvents or by-products that may be included in commercially available materials, unless otherwise specified.

[0033] Numerical ranges as used herein are intended to include every number and subset of numbers within that range, whether specifically disclosed or not. Further, these numerical ranges should be construed as providing support for a claim directed to any number or subset of numbers in that range. For example, a disclosure of from 1 to 10 should be construed as supporting a range of from 2 to 8, from 3 to 7, from 5 to 6, from 1 to 9, from 3.6 to 4.6, from 3.5 to 9.9, and so forth.

[0034] All references to singular characteristics or limitations of the present disclosure shall include the corresponding plural characteristic or limitation, and vice versa, unless otherwise specified or clearly implied to the contrary by the context in which the reference is made.

[0035] All combinations of method or process steps as used herein can be performed in any order, unless otherwise specified or clearly implied to the contrary by the context in which the referenced combination is made.

[0036] The various embodiments of the compositions for use in the methods of the present disclosure may also be substantially free of any optional or selected ingredient or feature described herein, provided that the nutritional composition still contains all of the required ingredients or features as described herein. In this context, and unless otherwise specified, the term “substantially free” means that the selected nutritional composition contains less than a functional amount of the optional ingredient, typically less than 1%, including less than 0.5%, including less than 0.1%, and also including zero percent, by weight of such optional or selected ingredient.

[0037] The compositions and methods may comprise, consist of, or consist essentially of the essential elements of the compositions and methods as described herein, as well as any additional or optional element described herein or otherwise useful in nutritional product applications.

Product Form

[0038] The compositions used in the methods of the present disclosure include a fucosylated human milk oligosaccharide, particularly 2-fucosyl-lactose (2FL), and may be formulated and administered in any known or otherwise suitable oral product form. Any solid, liquid, semi-solid, semi-liquid, or powder product form, including combinations or variations thereof, are suitable for use herein, provided that such forms allow for safe and effective oral delivery to the individual of the ingredients as also defined herein.

[0039] The compositions used in the methods of the present disclosure are desirably formulated as dietary product forms, which are defined herein as those embodiments comprising the ingredients of the present disclosure in a product form that then contains at least one of fat, protein, and carbohydrate, and preferably also contains vitamins, minerals, or combinations thereof.

[0040] The nutritional compositions may be formulated with sufficient kinds and amounts of nutrients to provide a sole, primary, or supplemental source of nutrition, or to provide a specialized nutritional product for use in individuals afflicted with specific conditions or with a targeted nutritional benefit as described below.

[0041] Some exemplary, non-limiting, examples of specific products that may be suitable for use in accordance with the present disclosure include preterm infant formulas, term

infant formulas, human milk fortifiers, pediatric formulas, adult nutritional formulas, older adult nutritional formulas, medical formulas, geriatric nutritional formulas, diabetic nutritional formulas, and the like.

Nutritional Liquids

[0042] Nutritional liquids include both concentrated and ready-to-feed nutritional liquids. These nutritional liquids are most typically formulated as suspensions or emulsions, although other liquid forms are within the scope of the present disclosure.

[0043] Nutritional emulsions suitable for use may be aqueous emulsions comprising proteins, fats, and carbohydrates. These emulsions are generally flowable or drinkable liquids at from about 1°C to about 25°C and are typically in the form of oil-in-water, water-in-oil, or complex aqueous emulsions, although such emulsions are most typically in the form of oil-in-water emulsions having a continuous aqueous phase and a discontinuous oil phase.

[0044] The nutritional emulsions may be and typically are shelf stable. The nutritional emulsions typically contain up to 95% by weight of water, including from about 50% to 95%, also including from about 60% to about 90%, and also including from about 70% to about 85%, of water by weight of the nutritional emulsions. The nutritional emulsions may have a variety of product densities, but most typically have a density greater than 1.03 g/mL, including greater than 1.04 g/mL, including greater than 1.055 g/mL, including from about 1.06 g/mL to about 1.12 g/mL, and also including from about 1.085 g/mL to about 1.10 g/mL.

[0045] The nutritional emulsions may have a caloric density tailored to the nutritional needs of the ultimate user, although in most instances the emulsions comprise generally at least 19 kcal/fl oz (660 kcal/liter), more typically from about 20 kcal/fl oz (675-680 kcal/liter) to about 25 kcal/fl oz (820 kcal/liter), even more typically from about 20 kcal/fl oz (675-680 kcal/liter) to about 24 kcal/fl oz (800-810 kcal/liter). Generally, the 22-24 kcal/fl oz formulas are more commonly used in preterm or low birth weight infants, and the 20-21 kcal/fl oz (675-680 to 700 kcal/liter) formulas are more often used in term infants. In some embodiments, the emulsion may have a caloric density of from about 50-100 kcal/liter to about 660 kcal/liter, including from about 150 kcal/liter to about 500 kcal/liter. In some specific embodiments, the emulsion may have a caloric density of 25, or 50, or 75, or 100 kcal/liter.

[0046] The nutritional emulsion may have a pH ranging from about 3.5 to about 8, but are most advantageously in a range of from about 4.5 to about 7.5, including from about 5.5 to about 7.3, including from about 6.2 to about 7.2.

[0047] Although the serving size for the nutritional emulsion can vary depending upon a number of variables, a typical serving size is generally at least 1 mL, or even at least 2 mL, or even at least 5 mL, or even at least 10 mL, or even at least 25 mL, including ranges from 1 mL to about 300 mL, including from about 4 mL to about 250 mL, and including from about 10 mL to about 240 mL.

Nutritional Solids

[0048] The nutritional solids may be in any solid form but are typically in the form of flowable or substantially flowable particulate compositions, or at least particulate compositions. Particularly suitable nutritional solid product forms include spray dried, agglomerated and/or dryblended powder compositions. The compositions can easily be scooped and measured with a spoon or similar other device, and can easily be reconstituted by the intended user with a suitable aqueous liquid, typically water, to form a nutritional composition for immediate oral or enteral use. In this context, “immediate” use generally means within about 48 hours, most typically within about 24 hours, preferably right after reconstitution.

[0049] The nutritional powders may be reconstituted with water prior to use to a caloric density tailored to the nutritional needs of the ultimate user, although in most instances the powders are reconstituted with water to form compositions comprising at least 19 kcal/fl oz (660 kcal/liter), more typically from about 20 kcal/fl oz (675-680 kcal/liter) to about 25 kcal/fl oz (820 kcal/liter), even more typically from about 20 kcal/fl oz (675-680 kcal/liter) to about 24 kcal/fl oz (800-810 kcal/liter). Generally, the 22-24 kcal/fl oz formulas are more commonly used in preterm or low birth weight infants, and the 20-21 kcal/fl oz (675-680 to 700 kcal/liter) formulas are more often used in term infants. In some embodiments, the reconstituted powder may have a caloric density of from about 50-100 kcal/liter to about 660 kcal/liter, including from about 150 kcal/liter to about 500 kcal/liter. In some specific embodiments, the emulsion may have a caloric density of 25, or 50, or 75, or 100 kcal/liter.

Methods Of Modulating Hypothalamic Pituitary Adrenal (HPA) Response and Serum Corticosterone Levels

[0050] The methods of the present disclosure use fucosylated human milk oligosaccharides-containing nutritional compositions, and in particular, 2-fucosyl-lactose (2FL)-containing nutritional compositions, to modulate hypothalamic pituitary adrenal (HPA) response in an individual. As noted, HPA response initiates the glucocorticoid stress response in vertebrate animals. During an acute psychological stressful event, stress hormones (e.g., corticosterone) are released from the HPA axis, affecting the release and expression of cytokines in the central nervous system as well as in immune cells. Hyperactivation of HPA leads to an over-release of hormones, which can be damaging to various physiological systems in the body of the individual, negatively affecting, for example, the immune and gastrointestinal systems, metabolic rate, and cognitive function. Particularly, hyperactivation of HPA may induce alterations in behavior, increasing anxiety, periods of depression, and aggressiveness, as well as negatively affect lipid metabolism, inflammatory responses, and gastrointestinal health. The methods of the present disclosure that modulate/prevent/control/reduce and/or treat HPA responses are beneficial for a wide range of individuals, including preterm infants, infants, pediatric individuals, teens, adults, and older adults (adults at least 50 or more years of age).

[0051] The methods further may be useful in preventing/controlling/reducing and/or treating malfunctions in the neuroendocrine-immune network that can result from a hyperactivated HPA response. Thereby, these methods maintain a healthy central nervous system, immunity system and healthy gastrointestinal system. The administration of 2FL may further have implications in the treatment of inflammation and cognitive disorders.

[0052] Additionally, the methods of the present disclosure include modulating and/or decreasing the production of serum corticosterone in an individual, particularly an individual affected by stress. Overproduction of corticosterone, typically a consequence of hyperactivated HPA response, leads to exacerbated stress responses in further stressful situations, as well as induces alterations in behavior, such as increasing anxiety-related responses, aggressiveness, future maternal care abilities, and the like. By modulating and/or reducing the production of serum corticosterone in an individual, including preterm infants,

infants, pediatric individuals, teens, adults, and older adults, these behavioral defects can be prevented/controlled/reduced and/or treated.

[0053] Further, as overproduction of corticosterone is closely related to central nervous system function (e.g., cognitive skills, mood disorders, depression), the methods of the present disclosure may further promote central nervous system function, especially in stressful situations, thereby preventing/controlling/reducing and/or treating occurrences of mood disorders or depression.

[0054] 2-fucosyl-lactose (2FL) may be administered to a subset of individuals in need of modulation or reduction of serum corticosterone levels and/or modulation of the HPA response. Some individuals that are in specific need of modulated or decreased serum corticosterone levels or modulated HPA responses may include infants, pediatrics, teens, or adults who experience acute psychological stress or stressful events (infants, pediatrics, teens, or adults susceptible to or at elevated risk of experiencing acute psychological stress or stressful events), infants, pediatrics, teens, or adults who experienced acute psychological stress early in life, non-breastfed infants, chronically depressed infants, pediatrics, teens, or adults (infants, pediatrics, teens, or adults susceptible to or at elevated risk of chronic depression), infants, pediatrics, teens, or adults affected by post-traumatic stress syndrome (infants, pediatrics, teens, or adults susceptible to or at elevated risk of post-traumatic stress syndrome) and the like. Preterm infants, infants, pediatrics, teens, adults, and older adults may be susceptible to or at elevated risk for experiencing acute psychological stress or stressful events due to family history, age, environment, and/or lifestyle. Based on the foregoing, because some of the method embodiments of the present disclosure are directed to specific subsets or subclasses of identified individuals (that is, the subset or subclass of individuals “in need” of assistance in addressing one or more specific conditions noted herein), not all individuals will fall within the subset or subclass of individuals as described herein for certain diseases or conditions.

[0055] The individual desirably consumes at least one serving of the nutritional composition daily, and in some embodiments, may consume two, three, or even more servings per day. Each serving is desirably administered as a single, undivided dose, although the serving may also be divided into two or more partial or divided servings to be taken at two or more times during the day. The methods of the present disclosure include continuous day after

day administration, as well as periodic or limited administration, although continuous day after day administration is generally desirable. The methods of the present disclosure are preferably applied on a daily basis, wherein the daily administration is maintained continuously for at least 3 days, including at least 5 days, including at least 1 month, including at least 4 weeks, including at least 8 weeks, including at least 2 months, including at least 6 months, desirably for at least 18-24 months, desirably as a long term, continuous, daily, dietary supplement.

2-Fucosyl-lactose (2FL)

[0056] The methods of the present disclosure for modulating and/or decreasing serum corticosterone levels and/or modulating HPA responses utilize compositions that include 2-fucosyl-lactose (2FL). The 2FL used in the composition may be isolated or enriched from milk(s) secreted by mammals including, but not limited to: human, bovine, ovine, porcine, or caprine species. 2FL may also be produced via microbial fermentation, enzymatic processes, chemical synthesis, or combinations thereof.

[0057] 2FL is present in the compositions in an amount (mg of 2FL per mL of composition) of at least 0.001 mg/mL, including at least 0.01 mg/mL, including from 0.001 mg/mL to about 20 mg/mL, including from about 0.01 mg/mL to about 20 mg/mL, including from 0.001 mg/mL to about 10 mg/mL, including from about 0.01 mg/mL to about 10 mg/mL, including from 0.001 mg/mL to about 5 mg/mL, including from about 0.01 mg/mL to about 5 mg/mL, including from 0.001 mg/mL to about 1 mg/mL, including from 0.001 mg/mL to about 0.23 mg/mL, including from about 0.01 mg/mL to about 0.23 mg/mL of 2FL in the composition. Typically, the amount of 2FL present in the composition will depend on the amounts of other components in the compositions, including the amounts any optional other human milk oligosaccharides as described below.

[0058] In one specific embodiment when the composition is a nutritional powder, the concentration of 2FL in the nutritional powder is from about 0.0005% to about 5%, including from about 0.01% to about 1% (by weight of the nutritional powder).

[0059] In another specific embodiment, when the product is a ready-to-feed nutritional liquid, the concentration of 2FL in the ready-to-feed nutritional liquid is from about 0.0001% to about 0.50%, including from about 0.001% to about 0.15%, including from about

0.01% to about 0.10%, and further including from about 0.01% to about 0.03% (by weight of the ready-to-feed nutritional liquid).

[0060] In another specific embodiment when the product is a concentrated nutritional liquid, the concentration of 2FL in the concentrated nutritional liquid is from about 0.0002% to about 0.60%, including from about 0.002% to about 0.30%, including from about 0.02% to about 0.20%, and further including from about 0.02% to about 0.06% (by weight of the concentrated nutritional liquid).

Optional Additional Sialylated or Fucosylated Human Milk Oligosaccharides

[0061] In addition to the 2FL described above, the compositions may optionally include additional sialylated or fucosylated human milk oligosaccharides. The additional human milk oligosaccharide(s) used in the composition may be isolated or enriched from milk(s) secreted by mammals including, but not limited to: human, bovine, ovine, porcine, or caprine species. The human milk oligosaccharides may also be produced via microbial fermentation, enzymatic processes, chemical synthesis, or combinations thereof.

[0062] Suitable sialylated human milk oligosaccharides for optional use in the compositions include at least one sialic acid residue in the oligosaccharide backbone. The sialylated human milk oligosaccharide may include two or more sialic acid residues also. Specific non-limiting examples of sialylated human milk oligosaccharides for use in the present disclosure include sialyl oligosaccharides, sialic acid (i.e., free sialic acid, lipid-bound sialic acid, protein-bound sialic acid), lactosialotetraose, 3'-Sialyl-3-fucosyllactose, Disialomonofucosyllacto-N-neohexaose, Monofucosylmonosialyllacto-N-octaose (sialyl Lea), Sialyllacto-N-fucohexaose II, Disialyllacto-N-fucopentaose II, Monofucosyldisialyllacto-N-tetraose), sialyl fucosyl oligosaccharides, 2'-Sialyllactose, 2-Sialyllactosamine, 3'-Sialyllactose, 3'-Sialyllactosamine, 6'-Sialyllactose, 6'-Sialyllactosamine, Sialyllacto-N-neotetraose c, Monosialyllacto-N-hexaose, Disialyllacto-N-hexaose I, Monosialyllacto-N-neohexaose I, Monosialyllacto-N-neohexaose II, Disialyllacto-N-neohexaose, Disialyllacto-N-tetraose, Disialyllacto-N-hexaose II, Sialyllacto-N-tetraose a, Disialyllacto-N-hexaose I, Sialyllacto-N-tetraose b, sialyl-lacto-N-tetraose a, sialyl-lacto-N-tetraose b, sialyl-lacto-N-tetraose c, sialyl-fucosyl-lacto-N-tetraose I, sialyl-fucosyl-lacto-N-tetraose II, disialyl-lacto-N-tetraose and combinations thereof. Particularly desirable sialylated human milk oligosaccharides include 3'Sialyllactose, 6'Sialyllactose, and combinations thereof.

[0063] Specific non-limiting examples of additional optional fucosylated human milk oligosaccharides for use in the present disclosure include fucosyl oligosaccharides, Lacto-N-fucopentaose I, Lacto-N-fucopentaose II, 3'-Fucosyllactose, Lacto-N-fucopentaose III, Lacto-N-difucohexaose I, Lactodifucotetraose, monofucosyllacto-N-hexaose II, isomeric fucosylated lacto-N-hexaose (1), isomeric fucosylated lacto-N-hexaose (3), isomeric fucosylated lacto-N-hexaose (2), difucosyl-para-lacto-N-neohexaose, difucosyl-para-lacto-N-hexaose, difucosyllacto-N-hexaosemonofucosyllacto-neocataose, monofucosyllacto-N-ocataose, difucosyllacto-N-octaose I, difucosyllacto-N-octaose II, difucosyllacto-N-neocataose II, difucosyllacto-N-neocataose I, lacto-N-fucopentaose V, lacto-N-decaose, trifucosyllacto-N-neooctaose, trifucosyllacto-N-octaose, trifucosyl-iso-lacto-N-octaose, lacto-N-difuco-hexaose II, and combinations thereof.

[0064] Other suitable examples of human milk oligosaccharides that may be included in the compositions for use in the methods of the present disclosure include lacto-N-hexaose, para-lacto-N-hexaose, lacto-N-neohexaose, para-lacto-N-neohexaose, lacto-N-neocataose, para-lacto-N-octanose, iso-lacto-N-octaose, lacto-N-octaose, and combinations thereof.

[0065] The sialylated and fucosylated human milk oligosaccharides (inclusive of 2FL) may be present in the compositions in a total amount of human milk oligosaccharide in the composition (mg of human milk oligosaccharide per mL of composition) of at least 0.001 mg/mL, including at least 0.01 mg/mL, including from 0.001 mg/mL to about 20 mg/mL, including from about 0.01 mg/mL to about 20 mg/mL, including from 0.001 mg/mL to about 10 mg/mL, including from about 0.01 mg/mL to about 10 mg/mL, including from 0.001 mg/mL to about 5 mg/mL, including from about 0.01 mg/mL to about 5 mg/mL, including from 0.001 mg/mL to about 1 mg/mL, including from 0.001 mg/mL to about 0.23 mg/mL, including from about 0.01 mg/mL to about 0.23 mg/mL of total human milk oligosaccharide in the composition. Typically, the amount of specific sialylated human milk oligosaccharide and/or fucosylated human milk oligosaccharide (inclusive of 2FL) present in the composition will depend on the specific human milk oligosaccharide or human milk oligosaccharides present and the amounts of other components in the compositions, including the amounts of any optional human milk oligosaccharides.

Macronutrients

[0066] The compositions including 2FL may be formulated to include at least one of fat, protein, and carbohydrate. In many embodiments, the nutritional compositions will include 2FL with fat, protein, and carbohydrate.

[0067] Although total concentrations or amounts of the fat, protein, and carbohydrates may vary depending upon the product type (i.e., human milk fortifier, preterm infant formula, infant formula, pediatric formula, adult formula, medical formula, etc.), product form (i.e., nutritional solid, powder, ready-to-feed liquid, or concentrated liquid) and targeted dietary needs of the intended user, such concentrations or amounts most typically fall within one of the following embodied ranges, inclusive of any other essential fat, protein, and/or carbohydrate ingredients as described herein.

[0068] For infant and adult formulas, carbohydrate concentrations most typically range from about 5% to about 40%, including from about 7% to about 30%, including from about 10% to about 25%, by weight; fat concentrations most typically range from about 1% to about 30%, including from about 2% to about 15%, and also including from about 3% to about 10%, by weight; and protein concentrations most typically range from about 0.5% to about 30%, including from about 1% to about 15%, and also including from about 2% to about 10%, by weight.

[0069] The amount of fats, proteins, and/or carbohydrates in any of the liquid nutritional compositions described herein may also be characterized in addition to, or in the alternative, as a percentage of total calories in the liquid nutritional composition as set forth in the following table. These macronutrients for liquid nutritional compositions of the present disclosure are most typically formulated within any of the caloric ranges (embodiments A-F) described in the following table (each numerical value is preceded by the term “about”).

Nutrient % Total Cal.	Embodiment A	Embodiment B	Embodiment C
Carbohydrate	0-98	2-96	10-75
Protein	0-98	2-96	5-70
Fat	0-98	2-96	20-85
	Embodiment D	Embodiment E	Embodiment F
Carbohydrate	30-50	25-50	25-50
Protein	15-35	10-30	5-30
Fat	35-55	1-20	2-20

[0070] In one specific example, liquid infant formulas (both ready-to-feed and concentrated liquids) include those embodiments in which the protein component may comprise from about 7.5% to about 25% of the caloric content of the formula; the carbohydrate component may comprise from about 35% to about 50% of the total caloric content of the infant formula; and the fat component may comprise from about 30% to about 60% of the total caloric content of the infant formula. These ranges are provided as examples only, and are not intended to be limiting. Additional suitable ranges are noted in the following table (each numerical value is preceded by the term “about”).

Nutrient % Total Cal.	Embodiment G	Embodiment H	Embodiment I
Carbohydrates:	20-85	30-60	35-55
Fat:	5-70	20-60	25-50
Protein:	2-75	5-50	7-40

[0071] When the nutritional product is a powdered preterm or term infant formula, the protein component is present in an amount of from about 5% to about 35%, including from about 8% to about 12%, and including from about 10% to about 12% by weight of the preterm or term infant formula; the fat component is present in an amount of from about 10% to about 35%, including from about 25% to about 30%, and including from about 26% to about 28% by weight of the preterm or term infant formula; and the carbohydrate component is present in an

amount of from about 30% to about 85%, including from about 45% to about 60%, including from about 50% to about 55% by weight of the preterm or term infant formula.

[0072] For powdered human milk fortifiers the protein component is present in an amount of from about 1% to about 55%, including from about 10% to about 50%, and including from about 10% to about 30% by weight of the human milk fortifier; the fat component is present in an amount of from about 1% to about 30%, including from about 1% to about 25%, and including from about 1% to about 20% by weight of the human milk fortifier; and the carbohydrate component is present in an amount of from about 15% to about 75%, including from about 15% to about 60%, including from about 20% to about 50% by weight of the human milk fortifier.

[0073] The total amount or concentration of fat, carbohydrate, and protein, in the powdered nutritional compositions used herein can vary considerably depending upon the selected composition and dietary or medical needs of the intended user. Additional suitable examples of macronutrient concentrations are set forth below. In this context, the total amount or concentration refers to all fat, carbohydrate, and protein sources in the powdered product. For powdered nutritional compositions, such total amounts or concentrations are most typically and preferably formulated within any of the embodied ranges described in the following table (each numerical value is preceded by the term “about”).

Nutrient % Total Cal.	Embodiment J	Embodiment K	Embodiment L
Carbohydrate	1-85	30-60	35-55
Fat	5-70	20-60	25-50
Protein	2-75	5-50	7-40

Fat

[0074] The nutritional compositions used in the methods of the present disclosure may include a source or sources of fat. Suitable sources of fat for use herein include any fat or fat source that is suitable for use in an oral nutritional product and is compatible with the elements and features of such products. For example, in one specific embodiment, the fat is derived from long chain polyunsaturated fatty acids and/or short chain fatty acids.

[0075] Additional non-limiting examples of suitable fats or sources thereof for use in the nutritional products described herein include coconut oil, fractionated coconut oil, soybean oil, corn oil, olive oil, safflower oil, high oleic safflower oil, oleic acids (EMERSOL 6313 OLEIC ACID, Cognis Oleochemicals, Malaysia), MCT oil (medium chain triglycerides), sunflower oil, high oleic sunflower oil, palm and palm kernel oils, palm olein, canola oil, marine oils, fish oils, fungal oils, algae oils, cottonseed oils, and combinations thereof.

Protein

[0076] The nutritional compositions used in the methods of the present disclosure may optionally further comprise protein. Any protein source that is suitable for use in oral nutritional compositions and is compatible with the elements and features of such products is suitable for use in the nutritional compositions.

[0077] Non-limiting examples of suitable proteins or sources thereof for use in the nutritional products include hydrolyzed, partially hydrolyzed or non-hydrolyzed proteins or protein sources, which may be derived from any known or otherwise suitable source such as milk (e.g., casein, whey), animal (e.g., meat, fish), cereal (e.g., rice, corn), vegetable (e.g., soy, pea) or combinations thereof. Non-limiting examples of such proteins include milk protein isolates, milk protein concentrates as described herein, casein protein isolates, extensively hydrolyzed casein, whey protein, sodium or calcium caseinates, whole cow milk, partially or completely defatted milk, soy protein isolates, soy protein concentrates, intact pea protein concentrates, intact pea protein isolates, hydrolyzed pea protein concentrates, hydrolyzed pea protein isolates, and so forth. In one specific embodiment, the nutritional compositions include a protein source derived from milk proteins of human and/or bovine origin.

Carbohydrate

[0078] The nutritional products as used in the methods of the present disclosure may further optionally comprise any carbohydrates that are suitable for use in an oral nutritional product and are compatible with the elements and features of such products.

[0079] Non-limiting examples of suitable carbohydrates or sources thereof for use in the nutritional products described herein may include maltodextrin, hydrolyzed or modified starch or cornstarch, glucose polymers, corn syrup, corn syrup solids, rice-derived carbohydrates, pea-derived carbohydrates, potato-derived carbohydrates, tapioca, sucrose,

glucose, fructose, lactose, high fructose corn syrup, honey, sugar alcohols (e.g., maltitol, erythritol, sorbitol), artificial sweeteners (e.g., sucralose, acesulfame potassium, stevia) and combinations thereof. A particularly desirable carbohydrate is a low dextrose equivalent (DE) maltodextrin.

Other Optional Ingredients

[0080] The compositions as used in the methods of the present disclosure may further comprise other optional components that may modify the physical, chemical, aesthetic or processing characteristics of the products or serve as pharmaceutical or additional nutritional components when used in the targeted population. Many such optional ingredients are known or otherwise suitable for use in medical food or other nutritional products or pharmaceutical dosage forms and may also be used in the compositions herein, provided that such optional ingredients are safe for oral administration and are compatible with the essential and other ingredients in the selected product form.

[0081] Non-limiting examples of such optional ingredients include preservatives, emulsifying agents, buffers, fructooligosaccharides, galactooligosaccharides, polydextrose, and other prebiotics, probiotics, pharmaceutical actives, anti-inflammatory agents, additional nutrients as described herein, colorants, flavors, thickening agents and stabilizers, emulsifying agents, lubricants, and so forth.

[0082] The nutritional compositions may further comprise a sweetening agent, preferably including at least one sugar alcohol such as maltitol, erythritol, sorbitol, xylitol, mannitol, isomalt, and lactitol, and also preferably including at least one artificial or high potency sweetener such as acesulfame K, aspartame, sucralose, saccharin, stevia, and tagatose. These sweetening agents, especially as a combination of a sugar alcohol and an artificial sweetener, are especially useful in formulating liquid beverage embodiments having a desirable flavor profile. These sweetener combinations are especially effective in masking undesirable flavors sometimes associated with the addition of vegetable proteins to a liquid beverage. Optional sugar alcohol concentrations in the nutritional product may range from at least 0.01%, including from about 0.1% to about 10%, and also including from about 1% to about 6%, by weight of the nutritional product. Optional artificial sweetener concentrations may range from at least 0.01%, including from about 0.05% to about 5%, also including from about 0.1% to about 1.0%, by weight of the nutritional product.

[0083] A flowing agent or anti-caking agent may be included in the nutritional compositions as described herein to retard clumping or caking of the powder over time and to make a powder embodiment flow easily from its container. Any known flowing or anti-caking agents that are known or otherwise suitable for use in a nutritional powder or product form are suitable for use herein, non-limiting examples of which include tricalcium phosphate, silicates, and combinations thereof. The concentration of the flowing agent or anti-caking agent in the nutritional composition varies depending upon the product form, the other selected ingredients, the desired flow properties, and so forth, but most typically range from about 0.1% to about 4%, including from about 0.5% to about 2%, by weight of the nutritional composition.

[0084] A stabilizer may also be included in the nutritional compositions. Any stabilizer that is known or otherwise suitable for use in a nutritional composition is also suitable for use herein, some non-limiting examples of which include gums such as xanthan gum. The stabilizer may represent from about 0.1% to about 5.0%, including from about 0.5% to about 3%, including from about 0.7% to about 1.5%, by weight of the nutritional composition.

[0085] The nutritional compositions may further comprise any of a variety of other vitamins or related nutrients, non-limiting examples of which include vitamin A, vitamin D, vitamin E, vitamin K, thiamine, riboflavin, pyridoxine, vitamin B₁₂, carotenoids (e.g., beta-carotene, zeaxanthin, lutein, lycopene), niacin, folic acid, pantothenic acid, biotin, vitamin C, choline, inositol, salts and derivatives thereof, and combinations thereof.

[0086] The nutritional compositions may further comprise any of a variety of other additional minerals, non-limiting examples of which include calcium, phosphorus, magnesium, iron, zinc, manganese, copper, sodium, potassium, molybdenum, chromium, chloride, and combinations thereof.

Methods of Manufacture

[0087] The nutritional compositions used in the methods of the present disclosure may be prepared by any known or otherwise effective manufacturing technique for preparing the selected product solid or liquid form. Many such techniques are known for any given product form such as nutritional liquids or powders and can easily be applied by one of ordinary skill in the art to the nutritional compositions described herein.

[0088] The nutritional compositions used in the methods of the present disclosure can therefore be prepared by any of a variety of known or otherwise effective formulation or manufacturing methods. In one suitable manufacturing process, for example, at least three separate slurries are prepared, including a protein-in-fat (PIF) slurry, a carbohydrate-mineral (CHO-MIN) slurry, and a protein-in-water (PIW) slurry. The PIF slurry is formed by heating and mixing the oil (e.g., canola oil, corn oil, etc.) and then adding an emulsifier (e.g., lecithin), fat soluble vitamins, and a portion of the total protein (e.g., milk protein concentrate, etc.) with continued heat and agitation. The CHO-MIN slurry is formed by adding with heated agitation to water: minerals (e.g., potassium citrate, dipotassium phosphate, sodium citrate, etc.), trace and ultra trace minerals (TM/UTM premix), thickening or suspending agents (e.g. avicel, gellan, carrageenan). The resulting CHO-MIN slurry is held for 10 minutes with continued heat and agitation before adding additional minerals (e.g., potassium chloride, magnesium carbonate, potassium iodide, etc.), and/or carbohydrates (e.g., 2FL, fructooligosaccharide, sucrose, corn syrup, etc.). The PIW slurry is then formed by mixing with heat and agitation the remaining protein, if any.

[0089] The resulting slurries are then blended together with heated agitation and the pH adjusted to 6.6-7.0, after which the composition is subjected to high-temperature short-time (HTST) processing during which the composition is heat treated, emulsified and homogenized, and then allowed to cool. Water soluble vitamins and ascorbic acid are added, the pH is adjusted to the desired range if necessary, flavors are added, and water is added to achieve the desired total solid level. The composition is then aseptically packaged to form an aseptically packaged nutritional emulsion. This emulsion can then be further diluted, heat-treated, and packaged to form a ready-to-feed or concentrated liquid, or it can be heat-treated and subsequently processed and packaged as a reconstitutable powder, e.g., spray dried, drymixed, agglomerated.

[0090] The nutritional solid, such as a spray dried nutritional powder or drymixed nutritional powder, may be prepared by any collection of known or otherwise effective technique, suitable for making and formulating a nutritional powder.

[0091] For example, when the nutritional powder is a spray dried nutritional powder, the spray drying step may likewise include any spray drying technique that is known for or otherwise suitable for use in the production of nutritional powders. Many different spray

drying methods and techniques are known for use in the nutrition field, all of which are suitable for use in the manufacture of the spray dried nutritional powders herein.

[092] One method of preparing the spray dried nutritional powder comprises forming and homogenizing an aqueous slurry or liquid comprising predigested fat, and optionally protein, carbohydrate, and other sources of fat, and then spray drying the slurry or liquid to produce a spray dried nutritional powder. The method may further comprise the step of spray drying, drymixing, or otherwise adding additional nutritional ingredients, including any one or more of the ingredients described herein, to the spray dried nutritional powder.

[0093] Other suitable methods for making nutritional products are described, for example, in U.S. Pat. No. 6,365,218 (Borschel, et al.), U.S. Pat. No. 6,589,576 (Borschel, et al.), U.S. Pat. No. 6,306,908 (Carlson, et al.), U.S. Patent Application No. 20030118703 A1 (Nguyen, et al.), which descriptions are incorporated herein by reference to the extent that they are consistent herewith.

EXAMPLES

[094] The following examples illustrate specific embodiments and/or features of the nutritional compositions used in the methods of the present disclosure. The examples are given solely for the purpose of illustration and are not to be construed as limitations of the present disclosure, as many variations thereof are possible without departing from the spirit and scope of the disclosure. All exemplified amounts are weight percentages based upon the total weight of the composition, unless otherwise specified.

[095] The exemplified compositions are shelf stable nutritional compositions prepared in accordance with the manufacturing methods described herein, such that each exemplified composition, unless otherwise specified, includes an aseptically processed embodiment and a retort packaged embodiment.

EXAMPLE 1

[0096] In this Example, the effects of 2-fucosyl-lactose on the production of serum corticosterone in early stressed maternally deprived mice were analyzed.

[0097] Initially, mice pups were early stressed using a model of maternal separation. Particularly, one day after delivery, C57BL/6 mice pups were sexed and grouped as sham (non-stressed Control group) or maternally separated (MS – stressed group) animals. From day 2 until day 14 after delivery, the MS pups were removed from the dam and kept isolated for a period of 4 hours per day, typically, from 10:00 AM to 2:00 PM, in a thermostatted cup. During the same time period, pups from the sham group were daily handled for a period of five minutes in order to receive the same grade of habituation to a researcher's hands as the MS pups.

[0098] Each dam had litters of 4 males and 2 females. Females were allowed to stay with the mother, while males were maternally deprived in order to avoid stressing the dam by removing all of the pups. Additionally, pups from the MS group were prematurely weaned, and thus permanently removed from the dam, at postnatal day 17, whereas sham pups were not weaned until postnatal day 21. At postnatal day 22, all pups in the MS group were pooled and four experimental groups were made before starting nutritional intervention. The five total groups are shown in the table below.

Experimental Group	Experimental Conditions
Sham Control Group (Group code E)	Normal mice not MS-stressed receiving AIN-93G diet and water.
MS Control Group (Group code D)	MS-stressed mice receiving AIN-93G diet and water.
MS 2-fucosyllactose Group (Group code C)	MS-stressed mice receiving AIN93G diet supplemented with 2-fucosyllactose in an amount of about 7 mg/mouse/day and water.

[0099] At the nutritional intervention stage, mice were fed experimental diets for a period of eight weeks. After eight weeks, blood samples were taken from facial veins of the mice in order to measure basal stress hormones (i.e., corticosterone, adrenocorticotrophic hormone (ACTH), and melatonin). Stress hormones were measured using luminex technology using a multiplex commercial kit (commercially available from Millipore Corporation, Billerica, Massachusetts.)

[00100] Seven days later, mice were submitted to an acute stress by restraint protocol. Particularly, mice were placed into a multi-drilled 50-ml centrifuge tube for 20 minutes. After this time, mice were extracted from the tube, and again, blood samples were taken in order to

measure serum stress hormones immediately after the acute stressful event (i.e., restraint). Results are shown in Figures 1 & 2.

[00101] As shown in Figures 1 and 2, no effects were found on ACTH or melatonin serum levels in any of the experimental groups post-restraint. The serum corticosterone levels, however, were significantly higher in MS mice that were submitted to early infancy stress as compared to normal sham mice (see Group D). It was further found, however, that early stressed mice fed a diet supplemented with 2-fucosyllactose (Group C) had corticosterone levels in response to restraint that equal that of the normal sham mice and were significantly lower than the levels exhibited by MS mice on the control diet. Accordingly, it is believed that 2-fucosyllactose was able to counteract the exacerbated hypothalamic-pituitary-adrenal (HPA) response in response to acute stressors showed by animals submitted to an early psychological stress during infancy.

EXAMPLES 2-6

[0102] Examples 2-6 illustrate ready-to-feed nutritional emulsions used in the methods of the present disclosure, the ingredients of which are listed in the table below. All ingredient amounts are listed as kilogram per 1000 kilogram batch of product, unless otherwise specified.

Ingredient	Ex. 2	Ex. 3	Ex. 4	Ex. 5	Ex. 6
Water	Q.S.	Q.S.	Q.S.	Q.S.	Q.S.
Condensed Skim Milk	86.64	86.64	86.64	86.64	86.64
Lactose	54.80	54.80	54.80	54.80	54.80
High oleic safflower oil	14.10	14.10	14.10	14.10	14.10
Soybean oil	10.6	10.6	10.6	10.6	10.6
Coconut oil	10.1	10.1	10.1	10.1	10.1
2-fucosyl-lactose (2FL)	0.0948	0.090	0.085	9.479	9.005
Galactooligosaccharides (GOS)	8.63	8.63	8.63	8.63	8.63
Whey protein concentrate	6.40	6.40	6.40	6.40	6.40
Potassium citrate	478.9 g	478.9 g	478.9 g	478.9 g	478.9 g
Calcium carbonate	448.28 g	448.28 g	448.28 g	448.28 g	448.28 g
Soy lecithin	355.74 g	355.74 g	355.74 g	355.74 g	355.74 g
Stabilizer	355.74 g	355.74 g	355.74 g	355.74 g	355.74 g
ARA oil	368.01 g	368.01 g	368.01 g	368.01 g	368.01 g
Nucleotide/chloride premix	293.26 g	293.26 g	293.26 g	293.26 g	293.26 g
Potassium chloride	226.45 g	226.45 g	226.45 g	226.45 g	226.45 g
Ascorbic acid	445.94 g	445.94 g	445.94 g	445.94 g	445.94 g
Vitamin mineral premix	142.88 g	142.88 g	142.88 g	142.88 g	142.88 g
DHA oil	137.8 g	137.8 g	137.8 g	137.8 g	137.8 g
Carrageenan	180.0 g	180.0 g	180.0 g	180.0 g	180.0 g
Magnesium chloride	55.0 g	55.0 g	55.0 g	55.0 g	55.0 g
Ferrous sulfate	58.0 g	58.0 g	58.0 g	58.0 g	58.0 g
Choline chloride	53.9 g	53.9 g	53.9 g	53.9 g	53.9 g

Ingredient	Ex. 2	Ex. 3	Ex. 4	Ex. 5	Ex. 6
Vitamin A, D ₃ , E, K ₁ premix	47.4 g	47.4 g	47.4 g	47.4 g	47.4 g
Citric acid	29.77 g	29.77 g	29.77 g	29.77 g	29.77 g
Mixed carotenoid premix	26.40 g	26.40 g	26.40 g	26.40 g	26.40 g
Sodium chloride	AN	AN	AN	AN	AN
L-carnitine	3.31 g	3.31 g	3.31 g	3.31 g	3.31 g
Tricalcium phosphate	15.65 g	15.65 g	15.65 g	15.65 g	15.65 g
Potassium phosphate monobasic	13.67 g	13.67 g	13.67 g	13.67 g	13.67 g
Riboflavin	2.42 g	2.42 g	2.42 g	2.42 g	2.42 g
Potassium hydroxide	AN	AN	AN	AN	AN

AN = as needed

EXAMPLES 7-11

[0103] Examples 7-11 illustrate ready-to-feed nutritional emulsions for use in the methods of the present disclosure, the ingredients of which are listed in the table below. All ingredient amounts are listed as kilogram per 1000 kilogram batch of product, unless otherwise specified.

Ingredient	Ex. 7	Ex. 8	Ex. 9	Ex. 10	Ex. 11
Water	Q.S.	Q.S.	Q.S.	Q.S.	Q.S.
Condensed Skim Milk	86.64	86.64	86.64	86.64	86.64
Lactose	54.80	54.80	54.80	54.80	54.80
High oleic safflower oil	14.10	14.10	14.10	14.10	14.10
Soybean oil	10.6	10.6	10.6	10.6	10.6
Coconut oil	10.1	10.1	10.1	10.1	10.1
HMO Mixture	0.0948	0.0901	0.0853	9.479	9.0047
6-sialyl-lactose (6SL)	0.0316	0.0300	0.0284	0	0
2-fucosyl-lactose (2FL)	0.0316	0.0300	0.0284	3.159	3.002
Lacto-N-neotetraose (LNnT)	0.0316	0.0300	0.0284	0	0
Galactooligosaccharides (GOS)	8.63	8.63	8.63	8.63	8.63
Whey protein concentrate	6.40	6.40	6.40	6.40	6.40
Potassium citrate	478.9 g	478.9 g	478.9 g	478.9 g	478.9 g
Calcium carbonate	448.28 g	448.28 g	448.28 g	448.28 g	448.28 g
Soy lecithin	355.74 g	355.74 g	355.74 g	355.74 g	355.74 g
Stabilizer	355.74 g	355.74 g	355.74 g	355.74 g	355.74 g
ARA oil	368.01 g	368.01 g	368.01 g	368.01 g	368.01 g
Nucleotide/chloride premix	293.26 g	293.26 g	293.26 g	293.26 g	293.26 g
Potassium chloride	226.45 g	226.45 g	226.45 g	226.45 g	226.45 g
Ascorbic acid	445.94 g	445.94 g	445.94 g	445.94 g	445.94 g
Vitamin mineral premix	142.88 g	142.88 g	142.88 g	142.88 g	142.88 g
DHA oil	137.8 g	137.8 g	137.8 g	137.8 g	137.8 g
Carrageenan	180.0 g	180.0 g	180.0 g	180.0 g	180.0 g
Magnesium chloride	55.0 g	55.0 g	55.0 g	55.0 g	55.0 g
Ferrous sulfate	58.0 g	58.0 g	58.0 g	58.0 g	58.0 g
Choline chloride	53.9 g	53.9 g	53.9 g	53.9 g	53.9 g
Vitamin A, D ₃ , E, K ₁ premix	47.40 g	47.40 g	47.40 g	47.40 g	47.40 g
Citric acid	29.77 g	29.77 g	29.77 g	29.77 g	29.77 g
Mixed carotenoid premix	26.40 g	26.40 g	26.40 g	26.40 g	26.40 g
Sodium chloride	AN	AN	AN	AN	AN
L-carnitine	3.31 g	3.31 g	3.31 g	3.31 g	3.31 g

Ingredient	Ex. 7	Ex. 8	Ex. 9	Ex. 10	Ex. 11
Tricalcium phosphate	15.65 g	15.65 g	15.65 g	15.65 g	15.65 g
Potassium phosphate monobasic	13.67 g	13.67 g	13.67 g	13.67 g	13.67 g
Riboflavin	2.42 g	2.42 g	2.42 g	2.42 g	2.42 g
Potassium hydroxide	AN	AN	AN	AN	AN

AN = as needed

WHAT IS CLAIMED IS

1. A nutritional composition comprising 2-fucosyl-lactose for use in modulating serum corticosterone levels in an animal.
2. A nutritional composition comprising 2-fucosyl-lactose for use in decreasing serum corticosterone levels in an individual affected by stress.
3. A nutritional composition comprising 2-fucosyl-lactose for use in modulating hypothalamic pituitary adrenal response in an individual.
4. The nutritional composition of any one of claims 1-3, wherein the animal or individual is affected by acute physiological stress.
5. The nutritional composition of any one of claims 1-4, wherein the animal or individual has been subjected to stress early in life.
6. The nutritional composition of any one of claims 1-5, wherein the animal or individual is a maternally deprived infant.
7. The nutritional composition of any one of claims 1-6, wherein the animal or individual is an infant.
8. The nutritional composition of any one of claims 1-7, in which the nutritional composition is a liquid and comprises from 0.001 mg/mL to about 20 mg/mL of 2-fucosyl-lactose.
9. The nutritional composition of any one of claims 1-7, in which the nutritional composition is a liquid and comprises from 0.001 mg/mL to about 10 mg/mL of 2-fucosyl-lactose.
10. The nutritional composition of any one of claims 1-7, in which the nutritional composition is a liquid and comprises from 0.001 mg/mL to about 5 mg/mL of 2-fucosyl-lactose.
11. The nutritional composition of any one of claims 1-7, in which the nutritional composition is a powder and comprises from about 0.0005% to about 5% of 2-fucosyl-lactose by weight of the powder.

12. The nutritional composition of any one of claims 1-7, in which the nutritional composition is a powder and comprises from about 0.01% to about 1% of 2-fucosyl-lactose by weight of the powder.

13. The nutritional composition of any one of claims 1-12, in which the nutritional composition further comprises at least one of fat, protein, and carbohydrate.

14. The nutritional composition of any one of claims 1-13, in which the nutritional composition is an infant formula.

1/1

FIG. 1





FIG. 2



INTERNATIONAL SEARCH REPORT

International application No
PCT/US2013/059436

A. CLASSIFICATION OF SUBJECT MATTER
INV. A23L1/29 A23L1/308 A61K31/702 A61K35/20
ADD. A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A23L A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, FSTA, PAJ, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2012/172307 A1 (DAVIS STEVEN R [US] ET AL) 5 July 2012 (2012-07-05) paragraphs [0002], [0013] - [0015], [0032], [0074], [0140] - [0150] -----	1-14
A	US 2012/172327 A1 (BUCK RACHAEL [US] ET AL) 5 July 2012 (2012-07-05) paragraphs [0002], [0003], [0006], [0007], [0012] - [0015], [0033], [0047], [0050], [0075] paragraphs [0143] - [0151] ----- -/-	1-14



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

30 October 2013

Date of mailing of the international search report

11/11/2013

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Couzy, François

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2013/059436

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2011/005681 A1 (CHILDRENS HOSP MEDICAL CENTER [US]; GEN HOSPITAL CORP [US]; INSTITUO N) 13 January 2011 (2011-01-13) page 1, lines 23-26 page 2, lines 9-22 table 1 page 8, lines 6-26 -----	1-14
A	US 2012/172319 A1 (CHOW JOMAY [US] ET AL) 5 July 2012 (2012-07-05) paragraphs [0003], [0005], [0010], [0013], [0016], [0143] - [0149] -----	1-14
A	KUNZ C ET AL: "OLIGOSACCHARIDES IN HUMAN MILK: STRUCTURAL, FUNCTIONAL, AND METABOLIC ASPECTS", ANNUAL REVIEW OF NUTRITION, ANNUAL REVIEWS INC., PALO ALTO, CA, US, vol. 20, 1 January 2000 (2000-01-01), pages 699-722, XP009020860, ISSN: 0199-9885, DOI: 10.1146/ANNUREV.NUTR.20.1.699 abstract pages 706-707 page 710 table 2 -----	1-14
A	Wang, B. et al: "Dietary sialic acid supplementation improves learning and memory in piglets", Am. J. Clin. Nutr., vol. 85, no. 2, 1 January 2007 (2007-01-01), 2007, pages 561-569, XP055050164, Retrieved from the Internet: URL: http://ajcn.nutrition.org/content/85/2/561.full.pdf#page=1&view=FitH [retrieved on 2013-01-17] abstract pages 567-8 page 2; figure 2 -----	1-14

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2013/059436

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2012172307 A1	05-07-2012	CA 2822497 A1 EP 2658549 A1 SG 191394 A1 US 2012172307 A1 WO 2012092159 A1	05-07-2012 06-11-2013 30-08-2013 05-07-2012 05-07-2012
US 2012172327 A1	05-07-2012	CA 2822664 A1 CN 103369974 A EP 2658400 A2 SG 191799 A1 US 2012172327 A1 WO 2012092157 A2	05-07-2012 23-10-2013 06-11-2013 30-08-2013 05-07-2012 05-07-2012
WO 2011005681 A1	13-01-2011	CA 2767043 A1 EP 2451462 A1 JP 2012532195 A US 2012202753 A1 WO 2011005681 A1	13-01-2011 16-05-2012 13-12-2012 09-08-2012 13-01-2011
US 2012172319 A1	05-07-2012	CA 2822219 A1 CN 103338775 A EP 2658547 A1 SG 191392 A1 US 2012172319 A1 WO 2012092156 A1	05-07-2012 02-10-2013 06-11-2013 30-08-2013 05-07-2012 05-07-2012