




(86) Date de dépôt PCT/PCT Filing Date: 1998/08/07
(87) Date publication PCT/PCT Publication Date: 1999/02/18
(45) Date de délivrance/Issue Date: 2010/11/30
(85) Entrée phase nationale/National Entry: 2000/02/08
(86) N° demande PCT/PCT Application No.: EP 1998/005025
(87) N° publication PCT/PCT Publication No.: 1999/007718
(30) Priorités/Priorities: 1997/08/08 (EP197 34 392.9);
1998/05/09 (EP198 20 815.4)

(51) Cl.Int./Int.Cl. *C07H 15/04* (2006.01),
A61K 31/7028 (2006.01), *A61K 31/7042* (2006.01),
A61P 19/02 (2006.01), *A61P 3/10* (2006.01),
A61P 35/04 (2006.01), *A61P 9/10* (2006.01),
C07H 15/12 (2006.01), *C07H 15/18* (2006.01),
C07H 15/26 (2006.01), *C07H 23/00* (2006.01),
C07H 5/10 (2006.01)

(72) Inventeurs/Inventors:
SCHMIDT, WOLFGANG, DE;
HENKE, STEPHAN, DE;
KUNZ, HORST, DE;
KALLUS, CHRISTOPHER, DE;
OPATZ, TILL, DE; ...

(54) Titre : DERIVES SUBSTITUES DU TETRAHYDROPYRANE, LEUR PROCEDE DE PRODUCTION, LEUR
UTILISATION COMME MEDICAMENTS OU AGENTS DIAGNOSTIQUES, ET MEDICAMENTS LES CONTENANT
(54) Title: SUBSTITUTED TETRAHYDROPYRANE DERIVATIVES AND METHOD FOR PRODUCING SAME



(I)

(57) Abrégé/Abstract:

The invention relates to compounds of the formula (I), in which the rests R¹, R², R³, R⁴, R⁵ and X have the meaning given to them in the description, a method for producing the compounds of formula (I) in a solid phase, and their use as medicines.



(72) Inventeurs(suite)/Inventors(continued): WUNBERG, TOBIAS, DE

(73) Propriétaires(suite)/Owners(continued): SANOFI-AVENTIS DEUTSCHLAND GMBH, DE

(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L., S.R.L.

PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION****International Bureau****INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)**

(51) International patent classification ⁶ : C07H	A2	(11) International publication number: WO 99/07718 (43) International publication date: 18 February 1999 (18.02.99)
(21) International application number: PCT/EP98/05025 (22) International filing date: 7 August 1998 (07.08.98) (30) Data relating to the priority: 197 34 392.9 8 August 1997 (08.08.97) DE 198 20 815.4 9 May 1998 (09.05.98) DE (71) Applicant (for all designated States except US): HOECHST MARION ROUSSEL DEUTSCHLAND [DE/DE]; Brüningstrasse 50, D-65929 Frankfurt am Main (DE). (72) Inventors; and (75) Inventors/Applicants (US only): SCHMIDT, Wolfgang [DE/DE]; Johannesallee 14, D-65929 Frankfurtam Main (DE). HENKE, Stephan [DE/DE]; Wingertstrasse 2c, D-65719 Hofheim (DE). KUNZ, Horst [DE/DE]; Gemeindehohl 50, D-55127 Mainz (DE). Kallus, Christopher [DE/DE]; J.J. Becherweg 18 - 20, D-55099 Mainz (DE). OBATZ, Till [DE/DE]; J.J. Becherweg 18 - 20, D-55099 Mainz (DE). WUNBERG, Tobias [DE/DE]; J.J. Becherweg 18 - 20, D-55099 Mainz (DE).		(81) Designated states: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HR, HU, ID, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ZW, ARIPO Patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European Patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI Patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without the International Search Report and to be republished once the report has been received.</i>

As printed

(54) Title: SUBSTITUTED TETRAHYDROPYRANE DERIVATIVES AND METHOD FOR PRODUCING SAME


(54) Bezeichnung: SUBSTITUIERTE TETRAHYDROPYRANDERIVATE, VERFAHREN ZU DEREN HERSTELLUNG, DEREN VERWENDUNG ALS ARZNEIMITTEL ODER DIAGNOSTIKUM SOWIE SIE ENTHALTENDES ARZNEIMITTEL

(57) Abstract

The invention relates to compounds of the formula (I), in which the rests R^1 , R^2 , R^3 , R^4 , R^5 and X have the meaning given to them in the description, a method for producing the compounds of formula (I) in a solid phase, and their use as medicines.

(57) Zusammenfassung

Die Erfindung betrifft Verbindungen der Formel (I), worin die Reste R^1 , R^2 , R^3 , R^4 , R^5 und X die in der Beschreibung genannte Bedeutung haben, ein Verfahren zur Herstellung der Verbindungen der Formel (I) an fester Phase sowie deren Verwendung als Arzneimittel.



WO 99/07718

PCT/EP98/05025

Substituted tetrahydropyrane derivatives and method for producing same

- 5 The invention relates to substituted tetrahydropyran derivatives, processes for their preparation, their use as a pharmaceutical or diagnostic and pharmaceutical comprising them.

10 Peptides and peptide mimetics are a valuable aid for the discovery of new lead structures and the identification of potential active compounds. By fixing side chains in a rigid structure (scaffold), it is hoped, in comparison with the conformationally more flexible peptide chain, for an increase in the affinity of this conformationally fixed ligand for the receptor.

- 15 Very different structural units are already finding use as peptide mimetics.

Owing to their polyvalency and their defined spatial arrangement, carbohydrate units should be particularly highly suitable as structural units for peptide mimetics.

20

Thus, it has recently been shown that a specific monosaccharide mimics, as a conformationally fixed structure, the spatial arrangement of a certain cyclopeptide, somatostatin (K.C. Nicolaou, J.I. Trujillo, K. Chibale, Tetrahedron 1997, 53, 8751-8778).

25

In this connection, starting from a glucose derivative having standard protective groups, a restricted variation of the simply accessible anomeric hydroxyl function and the C-6 hydroxyl function was carried out. The synthesis strategy described there starts from already known sugar units and is restricted by the protective group strategy to a narrow application range of somatostatin. At the same time, the method is not transferable to the targeted variation of the structural unit by solid-phase synthesis.

30

The previous syntheses of carbohydrate derivatives in solution or in the form of substance libraries on a solid phase concentrated, in particular, on the synthesis of oligosaccharides or glycopeptides (L. DeNapoli et al., Tetrahedron Letters 1996, 37, 5007-5010; S.J. Danishefsky et al., Science 1995, 269, 202-204, J.J. Krepinski et al., J. Am. Chem. Soc., 1991, 113, 5095-5097).

35

Compounds synthesized from oligosaccharide or glycopeptide units are, however, of only very restricted use for the discovery of lead structures or as potential active compounds on account of their complexity.

5

Restriction to a monosaccharide as a structural unit, however, combines the positive property of the defined spatial arrangement of the ligands with a low complexity, low molecular weight, low toxicity and further properties which are of importance for potential active compounds.

10

On account of the polyvalency of the monosaccharides, targeted synthesis of selectively functionalized monosaccharides - both in solution and on the solid phase - causes great difficulties.

15 Various protected carbohydrate units are likewise known as a result of the various studies on carbohydrate chemistry (see R.R. Schmidt, Pure & Appl. Chem. 1989, 61, 1257). In the intermediates described there, the hydroxyl groups are temporarily blocked more or less selectively by protective groups which are then deprotected for linkage with other
20 protective groups, as a result of which the synthesis of di- or oligosaccharides takes place.


These intermediates or the polysugars synthesized therefrom are, however, of only restricted use for specific lead structure discovery and as potential active compounds. In some cases, these structures are relatively
25 labile and thus not resistant to degradation or cleavage.

The linkers and activation strategies developed for the preparation of the abovementioned polysaccharides or glycopeptides (D. Kahne et al., J. Am. Chem. Soc. 1994, 116, 6953-6954) are also not generally transferable to
30 the preparation of selectively polysubstituted monosaccharide compounds.

The specific synthesis of selectively functionalized monosaccharide derivatives therefore requires the development of a novel, completely orthogonal protective group strategy, which makes it possible to selectively
35 remove the protective groups of all functional groups, the conditions used for this being stable to the conditions of the synthesis sequence. At the same time, these protective groups must guarantee compatibility with all reaction conditions which are necessary for synthesis in solid-phase synthesis. For synthesis on a solid phase, it is furthermore necessary to

have available a linker system for linking the monosaccharide unit, preferably via the anomeric center, which is compatible with all reaction conditions and can be selectively activated. Such a strategy makes possible the specific different variation of all functionalities of the monosaccharide unit to give stable final products.

The invention thus relates to compounds of the formula I



in which:

R^1, R^2, R^3, R^4, R^5 independently of one another are

1. hydrogen;
2. (C₁-C₁₂)-alkyl;
3. (C₂-C₈)-alkenyl;
4. (C₂-C₈)-alkynyl;
5. (C₁-C₆)-alkylene-(C₃-C₁₀)-cycloalkyl;
6. (C₀-C₆)-alkylene-(C₆-C₁₂)-aryl; preferably phenyl or benzyl;
7. (C₁-C₆)-alkoxy;
8. (C₀-C₆)-alkylene-CO-R⁸;
9. (C₁-C₆)-alkylene-(C₁-C₉)-heteroaryl;
10. carbamoyl;
11. -C(O)NR⁶R⁷;
12. -C(O)OR⁶;
13. a radical defined as in 2.-12., which is mono-, di- or polysubstituted in the alkyl moiety and/or aryl or heteroaryl moiety by a radical from the group consisting of (C₁-C₆)-alkyl, NO₂, CN, halogen, CF₃ or (C₁-C₆)-alkoxy;
14. a radical defined as in 6. and 9., which is substituted in the aryl or heteroaryl moiety by one, two or more halogen atoms;

R^6 and R^7 independently of one another are:

1. hydrogen;
2. (C₁-C₁₂)-alkyl;
3. (C₂-C₈)-alkenyl;
- 5 4. (C₂-C₈)-alkynyl;
5. (C₁-C₆)-alkylene-(C₃-C₁₀)-cycloalkyl;
6. (C₁-C₆)-alkylene-(C₆-C₁₂)-aryl; preferably benzyl;
7. (C₂-C₆)-alkyloxy;
8. (C₀-C₆)-alkylene-CO-R⁸;
- 10 9. (C₁-C₆)-alkylene-(C₁-C₉)-heteroaryl;
10. (C₀-C₆)-alkylene-(C₁-C₆)-alkoxy;
11. (C₃-C₁₀)-cycloalkyl;
12. (C₆-C₁₂)-aryl, preferably phenyl;

- 15 R⁸ is hydrogen, (C₁-C₆)-alkyl, (C₆-C₁₂)-aryl or OR¹²;
 R¹² is hydrogen, (C₁-C₆)-alkyl or (C₆-C₁₂)-aryl;

or

- 20 R² and R³ together or R³ and R⁴ together or R⁴ and R⁵ together are

(C₁-C₃)-alkylene which can be substituted by 1 or 2 (C₁-C₃)-alkyl radicals or optionally substituted (C₆-C₁₂)-aryl radicals;

- 25 X is N or O;

with the proviso that R² is not -C(O)OR⁶ when X is O;

and their physiologically tolerable salts.

30

Preferred compounds of the formula I are those in which the radicals R¹, R², R³, R⁴ and R⁵ do not each have the same meaning, and their physiologically tolerable salts.

- 35 Preferred compounds of the formula I are furthermore those in which only three of the radicals R¹, R², R³, R⁴, R⁵ have the same meaning, and their physiologically tolerable salts.

Particularly preferred compounds of the formula I are those in which only two of the radicals R^1 , R^2 , R^3 , R^4 , R^5 have the same meaning, and their physiologically tolerable salts.

- 5 Very particularly preferred compounds of the formula I are those in which all radicals R^1 , R^2 , R^3 , R^4 , R^5 have a different meaning, and their physiologically tolerable salts.

- 10 Preferred compounds of the formula I are those in which at least one of the radicals R^1 , R^2 , R^3 , R^4 , R^5 is hydrogen, $-C(O)NR^6R^7$, (C_1-C_8) -alkyl, (C_0-C_6) -alkyl- (C_6-C_{12}) -aryl, preferably phenyl or benzyl; the aryl moiety of the (C_1-C_6) -alkyl- (C_6-C_{12}) -aryl radical being unsubstituted or mono-, di- or trisubstituted by (C_1-C_6) -alkyl, cyano, nitro, CF_3 , Cl, Br or (C_1-C_4) -alkoxy, preferably methoxy, and R^6 and R^7 independently of one another are
- 15 hydrogen, (C_1-C_4) -alkyl, benzyl, (C_1-C_3) -alkylene- (C_3-C_7) -cycloalkyl, (C_1-C_3) -alkylene-CO-OR¹², (C_1-C_3) -alkylene- (C_1-C_3) -alkoxy, phenyl, optionally substituted by one or two radicals from the group consisting of CF_3 , Cl, Br, F, nitro, cyano; and R^{12} is as defined above;


- 20 or R^3 and R^4 together or R^4 and R^5 together are $-CH_2-$ which is substituted by 1 or 2 methyl radicals or optionally substituted phenyl radicals, and the other radicals are as defined above,

and their physiologically tolerable salts.

25

Preferred compounds of the formula I are also those in which X is equal to $-O-$ and the other radicals are as defined above, and their physiologically tolerable salts.

- 30 The invention furthermore relates to compounds of the formula II



(II)

in which:

R^1 is a linker group which can be linked via a covalent bond to a carrier functionalized by a heteroatom, for example N, O or Cl;

5

R^2, R^3, R^4, R^5 independently of one another are

a protective group customary in sugar chemistry;

10 Y is O or S, preferably S;

X is O or N, preferably O.

15 Protective groups customary in sugar chemistry are, for example, those such as are described, for example, in T.W. Greene, P.G. Wuts "Protective Groups in Organic Synthesis", 2nd Edition, Wiley/New York, 1991.

20 Suitable protective groups for compounds of the formula II are, for example, silyl protective groups, e.g. TBDPS; alkoxyalkyl groups, e.g. ethoxyethyl; allyl groups; acyl groups such as acetyl or benzoyl; acetals and ketals such as isopropylidene or optionally substituted benzylidene.

25 Preferred compounds of the formula II are those in which the radicals R^2, R^3, R^4 and R^5 are not all the same protective group.

Preferred compounds of the formula II are furthermore those in which only two of the radicals R^2, R^3, R^4, R^5 are an identical protective group.

30 Very particularly preferred compounds of the formula II are those in which the radicals R^2, R^3, R^4, R^5 are each a different protective group.

Preferred compounds of the formula II are also those which have an orthogonal protective group pattern with protective groups from the following different classes:

- 35
- base-labile protective groups, such as the acetate or benzoyl group;
 - acid-labile protective groups, such as acetal- or ketal-like protective groups such as the ethoxyethyl group;
 - fluoride-labile protective groups, such as the tert-butyldimethylsilyl or tert-butyldiphenylsilyl group;

- a protective group removable by transition metal catalysis, such as the allyl group;
- sulfur-containing protective groups, such as in the linker system.

5 The invention also relates to compounds of the formula II, in which:

R^5 is a linker group which can be linked via a covalent bond to a carrier functionalized by a heteroatom, for example N, O or Cl;


10 R^1, R^2, R^3, R^4 independently of one another are
a protective group customary in sugar chemistry;

Y is S or O, preferably S;


15

X is O or N, preferably O.


The invention also relates to compounds of the formulae IIa, IIb and IIc



IIa



IIb



IIc

20

in which:

Y is S or O, preferably S;

X is O or N, preferably O;

25 R^1 is a linker group which can be linked via a covalent bond to a carrier functionalized by a heteroatom, for example N, O or Cl;

R^2 in the case in which X is equal to O,
is a base-labile protective group such as, for example, acetyl or benzoyl;

30 in the case in which X is equal to N,
is a base-labile protective group such as, for example, a phthaloyl protective group, or DDE (1-(4,4-dimethyl-2,6-dioxocyclohexylidene-ethyl) or NDE (2-acetyl-4-nitroindan-1,3-dione);

R^3 is an allyl protective group;

R^4 is an acid-labile protective group, such as acetal- or ketal-like protective groups, for example ethoxyethyl or SEM (2-(trimethylsilyl)ethoxymethyl);

5 R^5 is a suitable silyl protective group, such as, for example, tert-butyldimethylsilyl or tert-butyldiphenylsilyl.

10 In general, suitable silyl protective groups for R^5 are fluoride-labile protective groups, which are more stable, i.e. more difficult to remove, than a trimethylsilyl radical.

15 The invention also relates to compounds of the formula IIa, in which R^4 and R^5 together are an acetal- or ketal-like protective group such as, for example, isopropylidene or benzylidene and the other radicals X, Y, R^1 , R^2 and R^3 are as defined above.

20 The invention additionally relates to compounds of the formula IIb, in which R^3 and R^4 together are an acetal- or ketal-like protective group such as, for example, isopropylidene or benzylidene and the other radicals X, Y, R^1 , R^2 and R^5 are as defined above.

A suitable linker group R^1 or R^5 is, for example, a group of the formula III

25 $(C_1-C_6)\text{-alkylene-[N-C(O)]}_n\text{-[(C}_6\text{-C}_{12}\text{)-arylene]}_p\text{-(C}_0\text{-C}_6\text{)-alkylene-C(O)R}^9$ (III)

in which n and p are 0 or 1, where p and n cannot simultaneously be 1;

30 R^9 is OR^{10} or $NR^{11}R^{11}$, where
 R^{10} is H, $(C_1-C_6)\text{-alkyl}$, $(C_1-C_6)\text{-alkyl-(C}_6\text{-C}_{12}\text{)-aryl}$, and
 R^{11} independently of one another is H, $(C_1-C_6)\text{-alkyl}$, $(C_1-C_6)\text{-alkyl-(C}_6\text{-C}_{12}\text{)-aryl}$.

35 Preferred compounds of the formula I or II are also those in which R^2 and R^3 together, or R^3 and R^4 together or R^4 and R^5 together form a benzylidene radical or isopropylidene radical and the other radicals are as defined above.

Preferred compounds of the formula I and formula II are furthermore those in which the monosaccharide structure is a glucose unit, a galactose unit or a mannose unit.

- 5 The compounds of the formula II and of the formula IIa, IIb or IIc are valuable intermediates for the preparation of compounds of the formula I.

(C₁-C₆)-Aryl is understood, for example, as meaning phenyl, naphthyl or biphenyl.

10

Alkyl, alkenyl, alkynyl, alkylene and radicals derived therefrom such as, for example, alkoxy can be straight-chain or branched, those branched radicals being preferred in which the branching site is not directly situated on the linkage site to the monosaccharide structure.

15

Halogen is preferably fluorine, chlorine or bromine.

- A heteroaryl radical within the meaning of the present invention is the radical of a monocyclic or bicyclic (C₃-C₉)-heteroaromatic which contains one or two N atoms and/or an S or an O atom in the ring system. For the term "heteroaromatic", see Garrat, Vollhardt, Aromatizität [Aromaticity], Stuttgart 1973, pages 131-153. Examples of suitable heteroaryl radicals are the radicals of thiophene, furan, benzo[b]thiophene, benzofuran, pyrrole, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, indole, quinoline, isoquinoline, oxazole, isoxazole, thiazole, isothiazole, isobenzofuran, indolizine, isoindole, indazole, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline and furazan.
- 20
- 25

- As indicated above, aryl, alkyl, heteroaryl and radicals derived therefrom can be monosubstituted or, if chemically possible, alternatively polysubstituted.
- 30

Suitable polymeric solid supports are, for example, crosslinked polystyrenes (e.g. aminomethylpolystyrene (AMPS) or Tentagel.

35

If not stated otherwise, chiral centers can be present in the R or in the S configuration. The invention relates both to the optically pure compounds and to mixtures of stereoisomers such as mixtures of enantiomers and mixtures of diastereomers.

Suitable salts are, in particular, alkali metal and alkaline earth metal salts, salts with physiologically tolerable amines and salts with inorganic or organic acids such as, for example, HCl, HBr, H₂SO₄, maleic acid, fumaric acid.


The abovementioned compounds of the formulae I and II and IIa, IIb or IIc respectively are derivatives of tetrahydropyran which can be synthesized rapidly and in automated form in good yields and high purities on a solid phase with the aid of the combinatorial method described here.

Compounds of the formula I can be prepared, for example, with the aid of intermediates of the formula II which have an orthogonal protective group pattern with one or preferably a number of protective groups from the following different classes:

- base-labile protective groups, such as the acetate or benzoyl group;
- acid-labile protective groups, such as acetal- or ketal-like protective groups such as the ethoxyethyl group;
- 20 - fluoride-labile protective groups, such as the tert-butyldimethylsilyl or tert-butyldiphenylsilyl group;
- a protective group which can be removed by transition metal catalysis, such as the allyl group;
- sulfur-containing protective groups, such as in the linker system.

25

The invention thus relates to a process for the preparation of compounds of the formula I



(I)

30 and their physiologically tolerable salts, in which:

R¹, R², R³, R⁴, R⁵ independently of one another are

1. hydrogen;

2. (C₁-C₁₂)-alkyl;
3. (C₂-C₈)-alkenyl;
4. (C₂-C₈)-alkynyl;
5. (C₁-C₆)-alkylene-(C₃-C₁₀)-cycloalkyl;
- 5 6. (C₀-C₆)-alkylene-(C₆-C₁₂)-aryl; preferably phenyl or benzyl;
7. (C₁-C₆)-alkoxy;
8. (C₀-C₆)-alkylene-CO-R⁸;
9. (C₁-C₆)-alkylene-(C₁-C₉)-heteroaryl;
10. carbamoyl;
- 10 11. -C(O)NR⁶R⁷;
12. -C(O)OR⁶;
13. a radical defined as in 2.-12., which is mono-, di- or polysubstituted in the alkyl moiety and/or aryl or heteroaryl moiety by a radical from the group consisting of (C₁-C₆)-alkyl, NO₂, CN, halogen, CF₃ or (C₁-C₆)-alkoxy;
- 15 14. a radical defined as in 6. and 9., which is substituted in the aryl or heteroaryl moiety by one, two or more halogen atoms;

or

- R² and R³ together or R³ and R⁴ together or R⁴ and R⁵ together are (C₁-C₃)-alkylene which can be substituted by 1 or 2 (C₁-C₃)-alkyl radicals or optionally substituted (C₆-C₁₂)-aryl radicals;

R⁶ and R⁷ independently of one another are:

1. hydrogen;
- 25 2. (C₁-C₁₂)-alkyl;
3. (C₂-C₈)-alkenyl;
4. (C₂-C₈)-alkynyl;
5. (C₁-C₆)-alkylene-(C₃-C₁₀)-cycloalkyl;
6. (C₁-C₆)-alkylene-(C₆-C₁₂)-aryl; preferably benzyl;
- 30 7. (C₂-C₆)-alkoxy;
8. (C₀-C₆)-alkylene-CO-R⁸;
9. (C₁-C₆)-alkylene-(C₁-C₉)-heteroaryl;
10. (C₀-C₆)-alkylene-(C₁-C₆)-alkoxy;
11. (C₃-C₁₀)-cycloalkyl;
- 35 12. (C₆-C₁₂)-aryl, preferably phenyl;

R⁸ is hydrogen, (C₁-C₆)-alkyl, (C₆-C₁₂)-aryl or OR¹²;

12


R^{12} is hydrogen, (C₁-C₆)-alkyl or (C₆-C₁₂)-aryl; and

X is N or O;

5 by

a) introduction of a suitable, preferably sulfur-containing linker on the anomeric center of an unprotected, partially orthogonally protected or completely orthogonally protected monosaccharide derivative of the formula


10



in which R in each case independently of one another is hydrogen or a protective group customary in sugar chemistry; and X is O or N, preferably O;

15

b) reaction of a compound linker-bonded in this way, to give compounds of the formula II



(II)

20

in which R^1 is a linker group which can be linked via a covalent bond to a support functionalized by a heteroatom, for example N, O or Cl, and

R^2 , R^3 , R^4 , R^5 independently of one another are a protective group customary in sugar chemistry, Y is O or S, preferably S and X is O or N, preferably O, by successive or simultaneous introduction of protective groups onto the functional groups -OR or -X-R, the

25

protective groups belonging to identical or different, preferably different, orthogonal protective group classes;

- 5 c) linkage of the monosaccharide derivative of the formula II protected in this way to a polymeric solid support via the linker;
- d) selective deprotection of the functional group to be derivatized on the polymeric solid support;
- 10 e) derivatization of the deprotected functional groups on the polymeric solid support, where the deprotection and subsequent derivatization of the different functional groups can be carried out selectively and also a number of equally protected functional groups can be deprotected and derivatized simultaneously;
- 15 f) removal of the derivatives bonded to the polymeric solid support, for example by activation of the sulfur on the anomeric center by bromine, and subsequent conversion of the compound activated in this way into a compound of the formula I derivatized on the
20 anomeric center.

For the synthesis of the selectively protected monosaccharide derivatives according to formula II or IIa, IIb or IIc respectively on a solid phase, linkage to the anomeric center via a thioglycoside or an O-glycoside, in
25 particular via a thioglycoside, is suitable. In this case, the different monosaccharides such as, for example, glucose, galactose or mannose, differ only slightly in the design of the protective groups and the sequence of their introduction. The differences in the reactivity of the functional groups and the differences associated therewith in the sequence of
30 introduction of the different protective groups is a known problem in carbohydrate chemistry.

The synthesis strategy for the preparation of compounds of the formulae I and II is illustrated in scheme 1 by the example of the glucose derivative
35 and is also transferable to other monosaccharides such as, for example, galactose (see scheme 3) or mannose (see scheme 4) with the abovementioned slight variations.

Compounds of the formula I can also be prepared by carrying out the linkage of selectively protected compounds of the formula II to a polymeric solid support by means of another OH position, for example the 6 OH position, such as shown in scheme 6 by the example of galactose. The linkage of the linker via the 6-OH position is possible, for example, according to the synthesis route shown in scheme 5.

Bonding of a glycosaccharide to the support material (scheme 1)

Reaction of the known 3-O-allyl-protected glucose β -acetate 3 (K. Takeo et al., Carbohydrate Research 133, 1984, 275) with the succinimide linker 2 prepared from 1 leads to compound 4. The β -configured thioglycoside 7 can be prepared analogously from the N-acylated cysteamine 6 by reaction with 3 under BF_3 catalysis. The deacetylation of 4 and 7 affords 8 homogeneously. Silylation on the C-6 hydroxyl group 9 and introduction of the ethoxyethyl protective group affords 10. Hydrolysis of the imide structure in 10 and coupling of the resulting acid 11 to a suitable support such as, for example, aminomethylpolystyrene affords the resin 12 loaded with the protected monosaccharide.

Reaction on the solid phase (see scheme 2)

The removal and reaction of the protected monosaccharide 12 on the solid phase to give compounds of the formula I (13) is shown by way of example in scheme 2. The C-2 hydroxyl function is deacetylated by reaction with hydrazine, the hydroxyl function can then be activated by reaction with potassium tert-butoxide or phosphazene as a base (R. Schwesinger, H. Schlemper, Angew. Chem. 99, 1987, 1212-1214). The activated derivative is trapped by means of electrophiles. An analogous reaction can be carried out in the case of a C-2 amino function. The protective group used here is, for example, the Fmoc group, which can be removed by piperidine.

The removal of the allyl ether on the C-3 hydroxyl group is carried out under zirconocene catalysis (E. Negishi, Tetrahedron Lett. 1986, 27, 2829-2832; E. Negishi, Synthesis, 1988, 1-19). The functionalization is carried out analogously to the manner described above. By this means, the use of strong acids, such as would be necessary in other removal methods familiar to the person skilled in the art, can be avoided and the orthogonality to the other protective groups is guaranteed.

Alternatively to the base-catalyzed functionalization, the allyl ether protective group can be converted into a propyl group by reduction with diimine (see Hüning, H.R. Müller, W. Thier, Angew. Chem. 1965, 77, 368-377). The C-4 hydroxyl function can be removed by transacetalization in an analogous manner to that used in the case of THP acetals (cf. E.J. Corey, H. Niwa, J. Knolle, J. Am. Chem. Soc. 1978, 100, 1942-1943). The functionalization is carried out as described above. The C-6 hydroxyl function is desilylated by reaction with fluoride ions; the reaction with electrophiles is carried out analogously to C-2.

10 The individual steps can be carried out in a different sequence on account of the compatibilities.

After completion of the functionalization of the various groups, the anomeric position is activated by reaction of the polymerically bonded selectively protected monosaccharide with bromine/di-tert-butylpyridine. The 1-bromo derivative is converted into a derivative functionalized on the anomeric center by reaction with alcohol.

20 Synthesis sequence for the preparation of galactose derivatives (see scheme 3)


Starting from galactose pentaacetate 14, the thioglycoside 16 is prepared by boron fluoride-catalyzed reaction with 15. Reaction with sodium methoxide affords 17 with deacetylation. The selective silylation of 17 takes place on the C-6 hydroxyl function. The silyl ether 18 is converted into the isopropylidene-protected derivative 19 using dimethoxypropane. Acetylation by reaction with acetic anhydride affords 20. After removal of the isopropylidene protective group, the dihydroxy compound is converted into the allyl ether derivative protected on C-4 using dibutyltin oxide and allyl bromide. Introduction of the ethoxyethyl protective group affords 21. The ester is hydrolyzed with lithium hydroxide analogously to glucose.

Synthesis sequence for the preparation of mannose derivatives (see scheme 4)

35 Mannose pentaacetate 22 is reacted with thiol 6 with boron trifluoride catalysis to give the thiomannoside 23. Removal of the acetate protective groups by sodium methoxide affords 24. Reaction with dimethoxybenzaldehyde affords the acetal 25. Reaction with dibutyltin





oxide and allyl bromide leads to the 3-O-allyl ester. By acetylation with acetic anhydride, 26 is prepared. The removal of the ketal and subsequent selective silylation on C-6 affords a silyl ether. Introduction of the ethoxyethyl protective group affords 27. The ester in 27 is hydrolyzed with
5 lithium hydroxide analogously to glucose.

Scheme 1




18

Scheme 1 (continuation)




19

Scheme 2



12


- 1) Hydrazine
- 2) Functionalization
- 3) TBAF
- 4) Functionalization
- 5) Acid/alcohol
- 6) Functionalization
- 7) "Zirconocene"
- 8) Functionalization
- 9) Bromine
- 10) Alcohol



13


20

Scheme 3




21

Scheme 4



Bonding to a polymeric solid support via the 6-OH position (scheme 5)

Scheme 5:



5

For anchorage via the primary hydroxyl group, a thioglycoside 28, for example, can be used as a starting material. The removal of the TBDPS group is carried out with tetrabutylammonium fluoride in THF. For the introduction of the linker, for example, etherification according to Mitsunobu is possible. The protection of the 4-hydroxyl group in 30 is carried out with slight warming with SEM-Cl and Hünig's base in dichloromethane.

10


Synthesis sequence for the preparation of galactose derivatives which are bonded to a polymeric solid support via the 1-OH group (scheme 6)

15

After binding of the glycoside to the solid polymeric support, the 1-OH group and the 2-OH group are first to be functionalized in the desired sequence, then the SEM group and the allyl ether are removed successively and the 3- and 4-positions are derivatized. The detachment of the solid polymer, for example using cerium ammonium nitrate (CAN), and the functionalization of the 6-OH function are then carried out.


20

Scheme 6:



- 5 Coupling of a methyl galactosylmercaptobutyrates to an amino-functionalized polymeric support via the 1-position (scheme 7)

Scheme 7



- 10 On account of their polyvalency and their defined spatial arrangement, the compounds of the formulae II, IIa, IIb and IIc are suitable as structural units for biological mimetics, for example peptide mimetics, and are a useful aid for the preparation and/or discovery of new lead structures and the identification of potential active compounds.

24

5

10


15

20

25

30

35



25

5

10


15

20

25

30

35



26

5

10


15

20

25

30

35



27

5

10


15

20

25

30

35



5

10

15

Examples

20 Example 1

N-(2-Thioethyl)succinimide 2

A solution of 24.2 g (0.21 mmol) of cysteamine hydrochloride 1 in 50 ml of H₂O is treated with 19.7 g (0.23 mol) of NaHCO₃ and stirred for 45 min. It is then concentrated in vacuo and the residue is taken up in 100 ml of acetic acid. After addition of 21.3 g (0.21 mol) of succinic anhydride, the suspension is heated under reflux for 3 h. The precipitate resulting on cooling of the solution is filtered off, washed with cold acetic acid and the filtrate is freed from the solvent in vacuo. The product is purified by chromatography on silica gel using petroleum ether/ethyl acetate (1:1). Yield: 13.0 g (38%) of colorless, amorphous solid.

R_f = 0.71 (EtOAc:HOAc = 30:1 v/v), m.p.: 44-45°C.

C₆H₉NO₂S (159.2) Calc.: C 45.26 H 5.70 N 8.80 S 20.14

Found: C 45.16 H 5.76 N 8.71 S 20.20

Example 2

N-[2-S-(2',4',6'-Tri-O-acetyl-3'-O-allyl-D-glucopyranosyl)thioethyl]-succinimide 4

- 5 4 ml (32 mmol) of boron trifluoride etherate in 10 ml of absol. CH₂Cl₂ are added dropwise to a solution of 2.0 g (5.15 mmol) of the acetate 3 and 980 mg (6.18 mmol) of N-(2-thioethyl)succinimide 2 in 55 ml of absol. CH₂Cl₂ cooled to 0°C under argon. The ice cooling is then removed and the reaction is stirred further at room temp. After 16 h, the mixture is
10 treated with 100 ml of CH₂Cl₂ and extracted twice with satd NaHCO₃ solution. The organic phase is dried over MgSO₄ and the solvent is removed in vacuo. The product is purified by chromatography on silica gel using petroleum ether/ethyl acetate (2:1) and the product is then reprecipitated from ethyl acetate/n-pentane. Yield: 1.71 g (75%) of
15 colorless, amorphous solid.

$R_f = 0.27$ (PE:EtOAc = 1:2 v/v), $[\alpha]_D^{25} = -41.8^\circ$ (c 1.0, CHCl₃), m.p.: 108°C

C₂₁H₂₉NO₁₀S (487.5) Calc.: C 51.74 H 6.00 N 2.87 S 6.58
Found: C 51.64 H 6.13 N 2.88 S 6.50

- 200 MHz ¹H-NMR (CDCl₃): [ppm] = 5.80-5.61 (m 1H, CH₂=CH); 5.18-4.97
20 (CH₂=CH, H-2' & H-4'); 4.38 (d, 1H, J_{2,1} = 10.01 Hz, H-1'); 4.19-3.94 (m, 4H, =CH=CH₂, H-6a/b'); 3.79-3.50 (m, 4H, H-3', H-5' & NCH₂ Cya); 2.93-2.79 (m, 1H, SCH₂ Cya); 2.76-2.61 (m, 1H, SCH₂ Cya); 2.65 (s, 4H, COCH₂ Suc); 2.03, 2.02 (2x s, 9H, CH₃ Ac). 100.6 MHz-¹³C-NMR (CDCl₃): [ppm] = 176.6 (COCH₂); 169.1 (COCH₃); 134.2 (CH₂=CH); 116.8
25 (CH₂=CH); 83.3, 81.0, 76.3, 73.0, 71.1, 69.4 (C-1', C-2', C-3', C-4', C-5', =CH-CH₂); 62.3 (C-6'); 38.2 (CH₂CO Suc); 28.0 (NCH₂ Cya); 27.2 (SCH₂ Cya); 20.8, 20.7, 20.6 (CH₃ Ac). The resulting α-anomer can be separated by chromatography. Yield: 0.19 g (8%), colorless oil. $R_f = 0.33$ (PE:EtOAc = 1.2 v/v).

30

Example 3

Monomethyl N-(2-thioethyl)succinamidate 6

- 11.0 g (96.8 mmol) of cysteamine hydrochloride 1 are suspended in 75 ml
35 of absol. acetonitrile under argon. 60 ml (345.0 mmol) of Hünig's base are slowly added dropwise to this with ice-cooling in an argon countercurrent.

- After 5 min, 16.4 ml (130.0 mmol) of trimethylchlorosilane are added in one portion. The mixture is stirred at 0°C for 10 min before a solution of 11.93 ml (96.8 mmol) of monomethyl succinyl chloride 5 in 20 ml of absol. acetonitrile is added dropwise. After 30 min at 0°C and 2 h at room temp., the solution is poured into 200 ml of ice water and the product is extracted twice with 200 ml of ethyl acetate each time. The combined organic phases are washed with 30 ml of 1N HCl, 50 ml of satd NaHCO₃ solution and 50 ml of satd NaCl solution, dried over MgSO₄ and the solvent is removed in vacuo. Yield: 13.1 g (71%), weakly yellowish oil.
- 10 $R_f = 0.54$ (EtOAc), $R_f = 0.58$ (EtOAc:HOAc = 30:1 v/v)
C₇H₁₃NO₃S (191.3)

Calc.:	C 43.96	H 6.85	N 7.32	S 16.76
Found:	C 43.97	H 6.78	N 7.65	S 16.16

- 90 MHz-¹H-NMR (CDCl₃): [ppm] = 3.62 (s, 3H, OCH₃); 3.37 (q, $J_{gem} = 6.26$ Hz, CH₂N Cya); 2.80-2.17 (m, 6H, SCH₂, 2x CH₂CO).

Example 4

Monomethyl N-[2-S-(2',4',6'-tri-O-acetyl-3'-O-allyl-β-D-glucopyranosyl)-thioethyl]succinamdate 7

- 20 6.0 g (15.5 mmol) of 3 are dissolved in 120 ml of absol. CH₂Cl₂. After addition of 3.32 g (18.5 mmol) of the thiol, the solution is cooled to 0°C under argon. A solution of 17.5 ml (139 mmol) of boron trifluoride etherate in 20 ml of absol. CH₂Cl₂ is slowly added dropwise to this mixture. The ice-cooling is then removed and the mixture is stirred at room temp. for 6 h. The reaction mixture is extracted twice with satd NaHCO₃ solution, the organic phase is separated off and dried over MgSO₄, and the solvent is removed in vacuo. After chromatography on silica gel using petroleum ether/ethyl acetate/HOAc (60:30:1), 6.8 g (85%) of a colorless, amorphous solid are obtained.

30 $R_f = 0.44$ (EtOAc), $R_f = 0.51$ (toluene:EtOH = 4:1 v/v), m.p.: 75-77°C, $[\alpha]_D^{25} = -5.3^\circ$ (c 1, CHCl₃). C₂₂H₃₃NO₁₁S (519.57)

Calc.:	C 50.82	H 6.40	N 2.70	S 6.17
Found:	C 50.74	H 6.44	N 2.76	S 6.23

200 MHz-¹H-NMR (CDCl₃): [ppm] = 6.33 (t_b, 1H, J_{gem} = 5.13 Hz, NH); 5.80-5.61 (m, 1H, CH₂=CH); 5.18-4.87 (m, 4H, CH₂=CH, H-2' & H-4'); 4.38 (d, 1H, J_{2,1} = 9.76 Hz, H-1'); 4.11-4.00 (m, 2H, H-6'a/b); 3.62 (s, 3H, CO₂CH₃); 3.59-3.41 (m, 3H, H-3', H-4', H-5'); 3.38-3.24 (m, 2H, CH₂N Cya); 2.89-2.51 (m, 4H, SCH₂ Cya & CH₂CO₂ Suc); 2.43 (t, 2H, J_{gem} = 6.41 Hz, CH₂CON Suc); 2.05, 2.02, 2.01 (3x s, 9H, CH₃ Ac).

Example 5

N-[2-S-(3'-O-Allyl-D-glucopyranosyl)thioethyl]succinimide 8

10

2.6 ml (2.6 mmol) of a 1M NaOMe solution in methanol are added under argon to 7.76 g (14.93 mmol) of thioglycoside 7 (crude product) dissolved in 60 ml of methanol p.a. The reaction solution is stirred at 50°C for 12 h, neutralized (5 min) with acidic ion exchanger Amberlyst® 15, and the ionic exchanger is removed by filtration and washed with methanol. The filtrate is freed from the solvent in vacuo. After chromatography on silica gel using toluene/EtOH (4:1), 4.86 g (86%) of a colorless oil are obtained, which solidifies after some time to give a colorless, amorphous solid.

20 $R_f = 0.27$ (toluene:EtOH = 4:1 v/v), m.p.: 98-99°C, $[\alpha]_D^{25} = -37.2^\circ$ (c 1.0, CHCl₃). CH₁₅H₂₃NO₇S (361.4)

Calc.:	C 49.85	H 6.41	N 3.88	S 8.87
Found:	C 49.89	H 6.62	N 3.86	S 8.83

400MHz ¹H-NMR (CDCl₃): [ppm] = 5.98-5.88 (m, 1H, CH₂=CH); 5.27 (d, 1H, J_{vic,trans} = 17.32 Hz, CH₂=CH); 5.16 (d, 1H, J_{vic,cis} = 10.27 Hz, CH₂=CH); 4.43 (dd, 1H, J_{gem} = 12.62 Hz, J_{vic} = 5.58 Hz, =CH-CH₂); 4.32 (d, 1H, J_{2,1} = 9.69 Hz, H-1'); 4.26 (dd, 1H, J_{gem} = 12.91 Hz, J_{vic} = 5.87 Hz, =CH-CH₂); 3.89-3.86 (m, 1H, H-6'a); 3.79-3.68 (m, 3H, H-2', H-4' & H-6'b); 3.54 (t, 1H, J_{2,3} = J_{4,3} = 9.25 Hz, H-3'); 3.44-3.34 (m, 2H, CH₂N Cya); 3.29 (t, J_{3,4} = J_{5,4} = 8.51 Hz, H-5'); 3.09, 3.03 (2x s_b, 2H, OH); 2.97-2.90 (m, 1H, SCH₂ Cya); 2.83-2.75 (m, 1H, SCH₂ Cya); 2.71 (s, 4H, CH₂CO Suc).

Example 6

N-[2-S-(2'-O-Acetyl-3'-O-allyl-D-glucopyranosyl)thioethyl]succinimide 8a

Variant 1: By deacetylation of the succinimide 4

- 5 1.15 g (2.36 mmol) of thioglycoside are cooled to 0°C under argon in 25 ml of methanol p.a. 13 mg (0.24 mmol) of NaOMe are added to the resulting suspension. After about 2 h, the precipitate dissolves and after a further 45 min the reaction is ended by addition of acidic ion exchanger Amberlyst® 15. The mixture is filtered, the ion exchanger is washed with methanol and
- 10 the combined filtrates are freed from the solvent in vacuo. After chromatography on silica gel using toluene/EtOH (4:1), 895 mg (94%) of a colorless, amorphous solid are obtained.

Variant 2: By deacetylation of the acyclic derivative 7

- 15 A solution of 2.6 g (5.0 mmol) of the glycoside in 50 ml of methanol p.a. is treated with 27.0 mg (0.5 mmol) of NaOMe at -15°C. After 1 h, no conversion can be detected by thin-layer chromatography, so a further 27 mg (0.5 mmol) of NaOMe are added and the temperature is increased to 0°C. As the formation of a further, more polar product (deacetylation in
- 20 the 2-position) can be detected in the thin-layer chromatogram, the reaction is terminated after 6 h by addition of acidic ion exchanger Amberlyst® 15. The solution is filtered, the ion exchanger is washed with methanol and the solvent is removed from the combined filtrates in vacuo. The product is purified by chromatography on silica gel using toluene/EtOH
- 25 (4:1). Yield: 1.2 g (90%), colorless, amorphous solid.

C₁₇H₂₅NO₈S (403.5) Calc.: C 50.61 H 6.25 N 3.47 S 7.95
 Found: C 49.85 H 6.59 N 3.87 S 7.95

- 200 MHz-¹H-NMR (CDCl₃): [ppm] = 5.92-5.73 (m, 1H, CH₂=CH); 5.24-5.08 (m, 2H, CH₂=CH); 4.86 (t, 1H, J_{1.2} = J_{3.2} = 9.52 Hz, H-2'); 4.37 (d, 1H, J_{2.1} = 7.76 Hz, H-1'); 4.25-4.08 (m, 2H, =CH-CH₂); 3.91-3.55 (m, 4H, H-3', H-4', H-6'a/b); 3.49-3.34 (m, 4H, H-5', CH₂N Cya & OH); 2.99-2.71 (m, 2H, SCH₂ Cya); 2.68 (s, 4H, CH₂CO Suc); 2.04 (s, 3H, CH₃ Ac).
- 30

Example 7

N-[2-S-(2'-O-Acetyl-3'-O-allyl-6'-O-tert-butyldiphenylsilyl-D-glucopyranosyl)-thioethyl]succinimide 9

- 5 1.0 g (2.48 mmol) of the succinimide 8a is dissolved in 20 ml of absol. CH₂Cl₂, treated with 583 mg (10.7 mmol) of imidazole, 0.74 ml (3.57 mmol) of tert-butyldiphenylchlorosilane and a spatula-tipful of DMAP and stirred at room temp. After 2 h, the mixture is diluted with 100 ml of CH₂Cl₂ and extracted with 50 ml of 1N HCl and satd NaCl solution. The
10 organic phase is dried over MgSO₄ and the solvent is removed in vacuo. After chromatography on silica gel using petroleum ether/ethyl acetate (1:1), 1.43 g (90%) of a colorless solid are obtained.
R_f = 0.47 (PE:EtOAc = 1:1 v/v), m.p.: 39-40°C

C₃₃H₄₃NO₈SSi (641.9) Calc.: C 61.75 H 6.75 N 2.18 S 5.00
Found: C 61.58 H 7.12 N 2.17 S 4.81

- 15 400MHz-¹H-NMR (CDCl₃): [ppm] = 7.67-7.65 (m, 4H, PhSi); 7.42-7.33 (m, 6H, PhSi); 5.89-5.82 (m, 1H, CH₂=CH); 5.24 (dd, 1H, J_{vic,trans} = 17.22 Hz, J_{gem} = 1.58 Hz, CH₂=CH); 5.14 (d, 1H, J_{vic,cis} = 10.41 Hz, CH₂=CH); 4.91 (t, 1H, J_{1.2} = J_{3.2} = 9.56 Hz, H-2'); 4.42 (d, 1H, J_{2.1} = 9.97 Hz, H-1'); 4.25-
20 4.14 (m, 2H, =CH-CH₂); 3.90 (d, 2H, J = 4.51 Hz, H-6'a/b); 3.76 (t, 1H, J_{3.4} = J_{5.4} = 9.21 Hz, H-4'); 3.70-3.63 (m, 2H, H-3' & H-5'); 3.47-3.41 (m, 2H, CH₂N Cya); 2.89 (s, 1H, OH); 2.87-2.81 (m, 1H, SCH₂ Cya); 2.75-2.62 (m, 1H, SCH₂ Cya); 2.61 (s, 4H, CH₂CO Suc); 2.07 (s, 3H, CH₃ Ac); 1.01 (s, 9H, CH₃ tBuSi).

25

Example 8

N-[2-S-(2'-O-Acetyl-3'-O-allyl-6'-O-tert-butyldiphenylsilyl-4'-O-(1''-(R/S)-ethoxyethyl)-D-glucopyranosyl)thioethyl]succinimide 10

30

- A solution of 1.26 g (1.96 mmol) of the succinimide 9 in 20 ml of absol. CH₂Cl₂ is treated with 0.94 ml (9.80 mmol) of ethyl vinyl ether and 246 mg (0.98 mmol) of pyridinium toluene-4-sulfonate and stirred at room temp. After 3 h, the reaction solution is diluted with 50 ml of CH₂Cl₂ and
35 extracted twice with 30 ml of satd NaHCO₃ solution each time. The organic phase is dried over MgSO₄ and the solvent is removed in vacuo.

Yield: 1.37 g (98%), colorless oil.

$R_f = 0.59$ (PE:EtOAc = 1:1 v/v),

$C_{37}H_{51}NO_9SSi$ (714.0)	Calc.:	C 62.25	H 7.20	N 1.96	S 4.49
	Found:	C 61.61	H 6.86	N 2.00	S 5.06
					(crude product)

- 5 400MHz- ^1H -NMR ($CDCl_3$): [ppm] = 7.70-7.65 (m, 4H, PhSi); 7.40-7.33 (m, 6H, PhSi); 5.85-5.81 (m, 1H, $CH_2=CH$); 5.25-5.08 (m, 2H, $CH_2=CH$); 4.93-4.87 (m, 1H, H-2'); 4.80 (q, 0.5H, $J_{gem} = 5.29$ Hz, $CHCH_3$ EE); 4.65 (q, 0.5H, $J_{gem} = 5.28$ Hz, $CHCH_3$ EE); 4.42 (d, 1H, $J_{2,1} = 9.68$ Hz); 4.23 (dd, 1H, $J_{gem} = 12.62$ Hz, $J_{vic} = 5.57$ Hz, $=CH-CH_2$); 4.16 (dd, 1H, $J_{gem} = 12.62$ Hz, $J_{vic} = 5.58$ Hz, $=CH-CH_2$); 3.90-3.86 (m, 2H, H-6'a/b); 3.75 (t, 1H, $J_{3,4} = J_{5,4} = 9.25$ Hz, H-4'); 3.68-3.58 (m, 3H, H-3' & CH_2N Cya); 3.52-3.41 (m, 3H, H-5' & CH_3CH_2O EE); 2.86-2.81 (m, 1 Hz, SCH_2 Cya); 2.74 (m, 1H, SCH_2 Cya); 2.60 (s, 4H, CH_2CO Suc); 2.06 (s, 3H, CH_3 Ac); 1.28-1.11 (m, 4.5H, $CHCH_3$ & CH_3CH_2O EE); 1.00 (s, 9H, CH_3 tBuSi); 0.89 (t, 15 1.5H, $J = 6.90$ Hz, CH_3CH_2O EE).

Example 9

N^1 -[2-S-(2'-O-Acetyl-3'-O-allyl-6'-O-tert-butyldiphenylsilyl-D-glucopyranosyl)thioethyl]- N^4 -benzylsuccinamide

20

2.8 g (4.36 mmol) of the succinimide 9 are dissolved in 30 ml of THF and cooled to 0°C. After addition of 15 mg of LiOH (4.8 mmol) in 10 ml of H_2O , the mixture is stirred at 0°C for 1.5 h, then acidified to pH = 2.5 with 1N HCl and extracted twice with 50 ml of CH_2Cl_2 each time. The combined organic phases are dried over $MgSO_4$ and freed from the solvent in vacuo.

25

The crude product thus obtained is treated in 30 ml of absol. CH_2Cl_2 with 0.96 ml (8.72 mmol) of benzylamine, 767 mg (6.54 mmol) of N-hydroxysuccinimide and 900 mg (4.36 mmol) of N,N'-dicyclohexylcarbodiimide. After 16 h, the precipitated urea is filtered off and washed with CH_2Cl_2 . The combined filtrates are extracted with 50 ml of 1N HCl and 50 ml of satd $NaHCO_3$ solution. The organic phase is dried over $MgSO_4$ and the solvent is removed in vacuo. The product is purified by chromatography on silica gel using petroleum ether/ethyl acetate mixtures.

30

Yield: 2.13 g (67%), colorless, amorphous solid.

$R_f = 0.53$ (EtOAc), $R_f = 0.33$ (EtOAc:PE:HOAc = 30:30:1 v/v), m.p. 46-47°C,

$C_{40}H_{52}N_2O_8SSi$ (735.0) Calc.: C 65.37 H 7.13 N 3.81 S 4.36
Found: C 63.68 H 6.83 N 3.71 S 4.45

5

200 MHz- ^1H -NMR ($CDCl_3$): [ppm] = 7.77-7.64 (m, 4H, PhSi); 7.50-7.28 (m, 6H, PhSi); 7.25-7.13 (m, 5H, Ph amide); 6.52 (t, 1H, $J_{gem} = 5.37$ Hz, NH); 6.34 (t, 1H, $J_{gem} = 5.37$ Hz, NH); 5.95-5.76 (m, 1H, $CH_2=CH$); 5.28-5.12 (m, 2H, $CH_2=CH$); 4.90 (t, 1H, $J_{1.2} = J_{3.2} = 9.53$ Hz, H-2'); 4.46-4.36 (m, 3H, H-1', CH_2 -Ph amide); 4.28-4.04 (m, 2H, $=CH-CH_2$); 3.91-3.89 (m, 2H, H-6'a/b); 3.69 (t, 1H, $J_{3.4} = J_{5.4} = 9.28$ Hz, H-4'); 3.52-3.26 (m, 5H, H-3', H-5', CH_2N Cya); 2.94 (s, 1H, OH); 2.85-2.58 (m, 2H, SCH_2 Cya); 2.47-2.34 (m, 4H, CH_2CO Suc); 2.08 (s, 3H, CH_3 Ac); 1.03 (s, 9H, CH_3 tBuSi).

15 Preparation of the galactose unit

Example 10

Methyl 4-S-(2',3',4',6'-tetra-O'-acetyl- β -D-galactopyranosyl)mercaptobutyrate 16

20

A solution of 12 g (30 mmol) of 1,2,3,4,6-penta-O-acetylgalactose 14 and 5.5 g of methyl mercaptobutyrate 15 in 150 ml of abs. dichloromethane is prestirred for 1 h with 10 g of thoroughly heated molecular sieve 4Å. The mixture is then cooled to 0°C and 30 ml of boron trifluoride ethyl etherate in 30 ml of abs. dichloromethane are added dropwise to the reaction mixture. It is then allowed to come to room temp. After 24 h, the precipitate is filtered off with suction through Celite® and the organic phase is stirred three times with 300 ml of satd $NaHCO_3$ solution each time. It is then washed with 600 ml of water, dried over $MgSO_4$ and freed from the solvent. The product is purified by chromatography on silica gel (eluent petroleum ether/ethyl acetate 2:1, column 20 × 8 cm). Yield 12.5 g (90%), yellow syrup, $R_f = 0.46$ (petroleum ether/ethyl acetate 1:1).

30

Example 11

Methyl 4-S-(6-O'-tert-butyldiphenylsilyl- β -D-galactopyranosyl)mercaptobutyrate 18

- 5 4.21 g (9.1 mmol) of 16 are dissolved in 40 ml of abs. methanol and 0.098 g (1.82 mmol) of sodium methoxide is added. After 4 h, the mixture is neutralized with acidic ion exchanger Amberlite® IR 120. The resin is filtered off and washed with methanol. After the removal of the solvent by distillation in vacuo and drying in a high vacuum, methyl S- β -D-galactopyranosylmercaptobutyrate 17 is obtained quant. as a colorless solid, R_f = 0.56 (chloroform/methanol 2:1). The crude product is dissolved in 20 ml of DMF and treated with 1.24 g (18.1 mmol) of imidazole and 3.25 ml (12.7 mmol) of tert-butyldiphenylsilyl chloride. The mixture is stirred at room temp. for 5 h and the reaction is then terminated by addition of
- 10 10 ml of water. After 10 min, the mixture is diluted with 60 ml of dichloromethane and washed three times with 40 ml of water each time. The org. phase is dried over $MgSO_4$ and evaporated after filtration in vacuo. The residue is purified by chromatography on silica gel. Yield 4.44 g (92%),
- 15 R_f = 0.23 (petroleum ether/ethyl acetate 1:1), 0.63 (ethyl acetate/acetic acid 30:1)
- 20 400MHz- ^1H -NMR ($CDCl_3$): δ [ppm] = 7.98-7.64, 7.40-7.36 (m, 10H, $SiPh_2$), 4.25 (d, 1H, $J_{1,2}$ = 9.6 Hz, 1-H), 4.10 (s, 1H, 4-H), 3.88 (dd, 1H, J_{gem} = 10.4 Hz, $J_{6a,5}$ = 6.3 Hz, 6- H_a), 3.84 (dd, 1H, J_{gem} = 10.9 Hz, $J_{6b,5}$ = 5.3 Hz, 6- H_b), 3.69 (dd, 1H, $J_{2,1}$ = 9.4 Hz, $J_{2,3}$ = 9.2 Hz, 2-H), 3.59 (s, 3H, $COOCH_3$), 3.56 (d, 1H, $J_{3,4}$ = 3.1 Hz, 3-H), 3.52 (dd, 1H, $J_{5,6a}$ = 5.7 Hz, $J_{5,6b}$ = 5.0 Hz, 5-H), 3.06 (s_{br} , 3H, OH), 2.72 (dt, 1H, J_{gem} = 13.9 Hz, J_{vic} = 7.0 Hz, SCH_a), 2.66 (dt, 1H, J_{gem} = 13.9 Hz, J_{vic} = 7.0 Hz, SCH_b), 2.40-2.36 (m_c , 2H, CH_2COOMe), 1.95-1.89 (m_c , 2H, SCH_2CH_2), 1.02 (s, 9H, $SiC(CH_3)_3$).
- 25 100.6 MHz- ^{13}C -NMR ($CDCl_3$): δ [ppm] = 173.5 ($COOMe$), 135.6; 135.5; 133.1; 133.0; 129.9; 127.8 ($SiPh_2$), 86.0; 78.4; 75.0; 70.5; 69.1; 63.2 (C-1-C-6), 51.5 ($COOCH_3$), 32.7 (SCH_2), 29.2 (CH_2COOMe), 26.8 ($SiC(CH_3)_3$), 25.3 (SCH_2CH_2), 19.1 ($SiC(CH_3)_3$).
- 30

Example 12

Methyl 4-S-(6-O'-tert-butyldiphenylsilyl-3,4-O'-isopropylidene- β -D-galactopyranosyl)mercaptopropionate 19

- 5 4.15 g (7.78 mmol) of 18 are dissolved in 35 ml of acetone dimethyl acetal and the mixture is stirred at room temp. for 4 h after addition of 28 mg of p-TsOH. It is then neutralized with triethylamine. The solution is concentrated in vacuo and the oily residue is freed from impurities by chromatography on silica gel (eluent petroleum ether/ethyl acetate 2:1, column 20 $\times$ 5 cm).
- 10 Yield 4.03 g (90%), colorless oil, R_f = 0.54 1:1 (petroleum ether/ethyl acetate 1:1).
- 400MHz- ^1H -NMR (CDCl_3): δ [ppm] = 7.70-7.66, 7.44-7.33 (m, 10H, Ph-H), 4.31 (dd, 1H, $J_{4,3}$ = 5.3 Hz, $J_{4,5}$ = 1.3 Hz, 4-H), 4.21 (d, 1H, $J_{1,2}$ = 10.3 Hz, 1-H), 4.04 (dd, 1H, $J_{3,2}$ = 7.0 Hz, $J_{3,4}$ = 5.6 Hz, 3-H), 3.90 (dd, 1H, J_{gem} = 10.8 Hz, $J_{6a,5}$ = 1.2 Hz, 6-H_a), 3.88 (dd, 1H, J_{gem} = 10.8 Hz, $J_{6b,5}$ = 4.8 Hz, 6-H_b), 3.87-3.85 (m, 1H, 5-H), 3.60 (s, 3H, COOCH_3), 3.52 (dd, $J_{2,1}$ = 8.8 Hz, $J_{2,3}$ = 8.5 Hz, 2-H), 2.75 (dt, 1H, J_{gem} = 13.2 Hz, J_{vic} = 7.0 Hz, SCH_a), 2.35 (dt, 1H, J_{gem} = 13.2 Hz, J_{vic} = 7.0 Hz, SCH_b), 2.42-2.39 (m, 2H, CH_2COOMe), 1.96-1.89 (m, 2H, SCH_2CH_2), 1.50 (s, 3H, $\text{C}(\text{CH}_3)_2$), 1.34 (s, 3H, $\text{C}(\text{CH}_3)_2$), 1.03 (s, 9H, $\text{C}(\text{CH}_3)_3$). 100.6 MHz- ^{13}C -NMR (CDCl_3): δ [ppm] = 173.3 (COOMe), 135.6; 135.5; 129.7; 127.7; 127.6 (C-Ph), 110.0 ($\text{C}(\text{CH}_3)_3$), 85.6; 79.1; 77.3; 73.3; 72.4; 62.7 (C-1 - C-6), 51.5 (COOCH_3), 32.5 (SCH₂), 29.4 (CH_2COOMe), 28.2 ($\text{C}(\text{CH}_3)_2$), 26.8 (C(CH₃)₃), 26.2 ($\text{C}(\text{CH}_3)_2$), 25.2 (SCH_2CH_2).
- 25

Example 13

Methyl 4-S-(2-O'-acetyl-6-O'-tert-butyldiphenylsilyl-3,4-O'-isopropylidene- β -D-galactopyranosyl)mercaptobutyrate 20

- 30 A solution of 27.6 g (48 mmol) of 19 in 100 ml of acetic anhydride/pyridine (1:1) is stirred for 5 h at room temp. It is then concentrated in a high vacuum and the residue is taken up in 100 ml of dichloromethane. The solution is washed with 50 ml each of 0.5 N HCl, satd NaHCO_3 solution and water. After drying over MgSO_4 , the mixture is freed from the solvent in vacuo. Chromatography on silica gel (eluent petroleum ether/ethyl acetate 4:1, column 30 $\times$ 10 cm) yields the title compound. Yield 24.3 g (82%), colorless oil,
- 35

$R_f = 0.66$ (petroleum ether/ethyl acetate 1:1). 400MHz- ^1H -NMR (CDCl_3):
 $\delta[\text{ppm}] = 7.70\text{-}7.66, 7.43\text{-}7.33$ (m, 10H, Ph-H), 4.98 (dd, $J_{2,1} = 10.0$ Hz,
 $J_{2,3} = 7.3$ Hz, 2-H), 4.34 (d, 1H, $J_{4,3} = 5.3$ Hz, 4-H), 4.31 (d, 1H, $J_{1,2} =$
 10.3 Hz, 1-H), 4.15 (dd, 1H, $J_{3,2} = 7.3$ Hz, $J_{3,4} = 5.2$ Hz, 3-H), 4.10-3.86
 5 (m, 3H, 6- $\text{H}_{a,b}$, 5-H), 3.58 (s, 3H, COOCH_3), 2.72 (dt, 1H, $J_{\text{gem}} = 12.9$ Hz,
 $J_{\text{vic}} = 7.0$ Hz, SCH_a), 2.60 (dt, 1H, $J_{\text{gem}} = 12.6$ Hz, $J_{\text{vic}} = 7.0$ Hz, SCH_b),
 2.41-2.32 (m_c , 2H, CH_2COOMe), 2.08 (s, 3H, COCH_3), 1.93-1.83 (m_c , 2H,
 SCH_2CH_2), 1.53 (s, 3H, $\text{C}(\text{CH}_3)_2$), 1.33 (s, 3H, $\text{C}(\text{CH}_3)_2$), 1.03 (s, 9H,
 $\text{C}(\text{CH}_3)_3$). 100.6 MHz- ^{13}C -NMR (CDCl_3): $\delta[\text{ppm}] = 173.2$ (COOMe), 169.5
 10 (COMe), 135.5; 135.5; 133.4; 133.3; 129.7; 127.6; 127.6 (C-Ph), 110.2
 ($\text{C}(\text{CH}_3)_2$), 82.7; 77.4; 76.7; 73.4; 71.7; 62.6 (C-1 - C-6), 51.4 (COOCH_3),
 32.6 (SCH_2), 29.1 (CH_2COOMe), 27.8 ($\text{C}(\text{CH}_3)_2$), 26.8 ($\text{C}(\text{CH}_3)_3$), 26.3
 ($\text{C}(\text{CH}_3)_2$), 24.9 ($\text{C}(\text{CH}_3)_3$), 20.9 (COCH_3), 19.2 (SCH_2CH_2).

15 Example 14

Methyl 4-S-(3-O'-allyl-2-O'-acetyl-4-O'-[1-(R/S)-ethoxyethyl]-6-O'-tert-
 butyldiphenylsilyl- β -D-galactopyranosyl)mercaptobutyrate 21

20 a) Methyl 4-S-(2-O'-acetyl-6-O'-tert-butyldiphenylsilyl- β -D-galacto-
 pyranosyl)mercaptobutyrate

A solution of 23.97 g (38.86 mmol) of 20 in 370 ml of CHCl_3 is heated to
 reflux with 0.35 g (1.84 mmol) of p-toluenesulfonic acid and 22.70 ml
 (270 mmol) of ethanedithiol. After 5 h, the mixture is cooled to room temp.
 25 and washed with 50 ml each of satd sodium hydrogencarbonate solution,
 0.5 N HCl and water. The organic phase is dried over magnesium sulfate.
 The product is obtained pure by chromatography on silica gel (eluent
 petroleum ether/ethyl acetate 1:1). Yield 16.66 g (74%), colorless oil,
 $R_f = 0.16$ (petroleum ether/ethyl acetate 2:1). 400MHz- ^1H -NMR (CDCl_3):
 30 $\delta[\text{ppm}] = 7.68\text{-}7.63, 7.42\text{-}7.34$ (m, 10H, Ph-H), 5.06 (dd, $J_{2,1} = 9.6$ Hz, $J_{2,3}$
 = 9.6 Hz, 2-H), 4.31 (d, 1H, $J_{1,2} = 9.9$ Hz, 1-H), 4.11 (d, 1H, $J_{4,3} = 5.3$ Hz,
 4-H), 3.92-3.84 (m, 2H, 6- $\text{H}_{a,b}$), 3.61 (d, 1H, $J_{3,4} = 3.4$ Hz, 3-H), 3.58 (s,
 3H, COOCH_3), 3.50 (t, 1H, $J_{5,6} = 5.5$ Hz, 5-H), 2.73 (dt, 1H, $J_{\text{gem}} = 13.0$
 Hz, $J_{\text{vic}} = 7.2$ Hz, SCH_a), 2.59 (dt, 1H, $J_{\text{gem}} = 13.0$ Hz, $J_{\text{vic}} = 7.2$ Hz,
 35 SCH_b), 2.40-2.16 (m_c , 2H, CH_2COOMe), 2.09 (s, 3H, COCH_3), 2.01-1.81
 (m_c , 2H, SCH_2CH_2), 1.03 (s, 9H, $\text{C}(\text{CH}_3)_3$). 100.6 MHz- ^{13}C -NMR (CDCl_3):
 $\delta[\text{ppm}] = 173.4$ (COOMe), 170.8 (COMe), 135.5; 135.4; 132.9; 132.7;
 129.9; 127.8 (C-Ph), 83.2; 78.1; 73.7; 71.2; 69.6; 63.3 (C-1 - C-6), 51.5

(COOCH₃), 32.6 (SCH₂), 28.8 (CH₂COOMe), 26.8 (C(CH₃)₃), 25.1 (C(CH₃)₃), 20.9 (COCH₃), 19.1 (SCH₂CH₂).

5 b) Methyl 4-S-(3-O'-allyl-2-O'-acetyl-6-O'-tert-butyldiphenylsilyl-β-D-galactopyranosyl)mercaptobutyrate

A mixture of 5.7 g (9.8 mmol) of the compound from Ex. 14a) and 3.3 g (9.43 mmol) of dibutyltin oxide in 70 ml of benzene is heated to reflux for 8 h in a water separator. 35 ml of benzene are then removed by distillation and the mixture is treated with 1.76 g (9.43 mmol) of tetrabutylammonium bromide and 1.35 ml (15.6 mmol) of allyl bromide. The solution is stirred at 50°C for 16 h. After addition of 5 ml of methanol, it is largely concentrated in vacuo. The residue is taken up in 50 ml of dichloromethane and washed three times with 10 ml of water each time. After drying over MgSO₄, the solvent is removed in vacuo. The crude product is purified by chromatography (eluent petroleum ether/ethyl acetate 4:1, column 15 × 5 cm) on silica gel. Yield 3.79 g (63%) of yellowish oil, R_f = 0.34 (petroleum ether/ethyl acetate 4:1). 400MHz-¹H-NMR (CDCl₃): δ[ppm] = 7.68-7.65, 7.42-7.34 (m, 10H, Ph-H), 5.90-5.80 (m_c, 1H, =CH), 5.26 (d, 1H, J_{vic} = 17.6 Hz, CH_{trans}=), 5.20 (dd, 1H, J_{2.1} = 10.0 Hz, J_{2.3} = 9.8 Hz, 2-H, R+S), 5.18 (d, 1H, J_{vic} = 11.8 Hz, CH_{cis}=), 4.30 (d, 1H, J_{1.2} = 10.0 Hz, 1-H), 4.16 (d, 1H, J_{4.5} = 2.7 Hz, 4-H), 4.11 (dd, 1H, J_{gem} = 12.9 Hz, J_{vic} = 5.6 Hz, =CH-CH_a), 4.03 (dd, 1H, J_{gem} = 12.9 Hz, J_{vic} = 5.6 Hz, =CH-CH_b), 3.94 (dd, 1H, J_{gem} = 10.3 Hz, J_{6a.5} = 6.5 Hz, 6-H_a), 3.86 (dd, 1H, J_{gem} = 10.3 Hz, J_{6b.5} = 5.6 Hz, 6-H_b), 3.59 (s, 3H, COOCH₃), 3.50 (dd, 1H, J_{5.6a} = J_{5.6b} = 5.9 Hz, 5-H), 3.44 (dd, 1H, J_{3.2} = 9.4 Hz, J_{3.4} = 2.7 Hz, 3-H), 2.75 (dt, 1H, J_{gem} = 13.2 Hz, J_{vic} = 7.0 Hz, SCH_a), 2.61 (dt, 1H, J_{gem} = 13.2 Hz, J_{vic} = 7.0 Hz, SCH_b), 2.37 (m_c, 2H, SCH₂CH₂), 2.07 (s, 3H, COCH₃), 1.88 (m_c, 2H, CH₂COOMe), 1.03 (s, 9H, SiC(CH₃)₃).

30

c) Methyl 4-S-(3-O'-allyl-2-O'-acetyl-4-O'-[1-(R/S)-ethoxyethyl]-6-O'-tert-butyldiphenylsilyl-β-D-galactopyranosyl)mercaptobutyrate

1.95 g (3.15 mmol) of the compound according to Example 14b) are dissolved in 45 ml of dichloromethane and the mixture is stirred at room temp. for 4 h after addition of 45 ml of ethyl vinyl ether and 0.39 g (1.56 mmol) of pyridium p-toluenesulfonate. The mixture is poured into satd NaHCO₃ solution and the aq. phase is extracted with ethyl acetate.

35

The combined org. phases are then dried over MgSO_4 , and freed from the solvent in vacuo. Purification is carried out by chromatography on silica gel (eluent petroleum ether/ethyl acetate 4:1, column 15×2 cm). Yield 1.49 g (69%), colorless oil,

- 5 $R_f = 0.39$ (petroleum ether/ethyl acetate 4:1). 400MHz- ^1H -NMR (CDCl_3): $\delta[\text{ppm}] = 8.03\text{-}8.01$; 7.69-7.36 (m, 10H, Ph-H), 5.80-5.65 (2m_c, 1H, =CH, R+S), 5.56, 5.50 (2dd, 1H, $J_{2.1} = J_{2.3} = 9.8$ Hz, 2-H, R+S), 5.26-5.01 (m, 2H, $\text{CH}_{\text{trans}}=$, $\text{CH}_{\text{cis}}=$, R+S), 4.94 (q, 1H, CHCH_3), 4.52, 4.46 (2d, 1H, $J_{1.2} = 10.3$ Hz, 1-H, R+S), 4.16-3.45 (m, 7H, 4-H, 6-H_{a,b}, 5-H, 3-H, =CH-CH_{a,b}, R+S), 3.54, 3.53 (2s, 3H, COOCH_3 , R+S), 3.69-3.63, 3.32-3.25 (2m_c, 2H, CH_2CH_3 , R+S), 3.04-2.94, 2.89-2.78 (2m_c, 2H, SCH_2 R+S), 2.66, 2.60 (2t, $J_{\text{vic}} = 7.3$ Hz, SCH_2CH_2 , R+S), 1.33, 1.26 (2d, 3H, $J_{\text{vic}} = 5.3$ Hz, CHCH_3 , R+S), 1.17, 0.95 (2t, 3H, $J_{\text{vic}} = 7.0$ Hz, CH_2CH_3 , R+S), 1.07, 1.05 (2s, 9H, $\text{C}(\text{CH}_3)_3$, R+S).

15

The hydrolysis of the ester is carried out as described in the case of glucose.

Preparation of the mannose derivatives

20

Example 15

Monomethyl N-[2-S-(2',3',4',6'-tetra-O-acetyl- α -D-mannopyranosyl)thioethyl]succinamidate 23

- 25 A solution of 11.1 g (34.3 mmol) of monomethyl N-(2-thioethyl)succinamidate 6 and 43 ml (0.34 mol) of boron trifluoride etherate in 70 ml of CH_2Cl_2 is added dropwise with ice-cooling to 10.0 g (25.6 mmol) of the anomer mixture 22 in 300 ml of CH_2Cl_2 . The solution is warmed to room temp. and stirred for 12 h. The mixture is washed twice with 500 ml of satd NaHCO_3 solution in each case and the organic phase is dried over MgSO_4 . After chromatography on silica gel using petroleum ether/ethyl acetate (1:2), 5.5 g (44%) of pure α -product are obtained as a colorless oil and also 2.3 g (19%) of an α,β -mixed fraction as a yellowish oil.

35

$R_f = 0.24$ (PE/EtOAc = 1:2 v/v), $R_f = 0.52$ (EtOAc/HOAc = 30:1 v/v), 200 MHz- ^1H -NMR (CDCl_3): $\delta[\text{ppm}] = 6.28$ (s_b, 1H, NH); 5.30-5.14 (m, 4H, H-1', H-2', H-3' & H-4'); 4.37-4.06 (m, 3H, H-5' & H-6'a/b); 3.65 (s, 3H, CO_2CH_3); 3.57-3.39 (m, 2H, CH_2N Cya); 2.85-2.71 (m, 2H, SCH_2 Cya);

2.63 (t, 2H, $J_{\text{vic}} = 7.08$ Hz, CH_2CO Suc); 2.45 (t, 2H, $J_{\text{vic}} = 7.09$ Hz, CH_2CON Suc); [lacuna].08, 20.7, 20.5 (3x s, 12H, CH_3 Ac).

Example 16

5 N-[2-S-(α -D-Mannopyranosyl)thioethyl]succinimide 24

1.5 ml of a 1 M solution of sodium methoxide in methanol are added under argon and with ice-cooling to 7.9 g (15.2 mmol) of the glycoside 23 in 100 ml of methanol p.a., and the mixture is stirred at room temp. for 2 h
10 and neutralized with Amberlyst® 15. The ion exchanger is filtered off, washed with methanol and the solvent is removed in vacuo. Residues of methanol are removed by codistillation with toluene. 4.8 g (97%) are obtained as a yellowish oil, which still contains slight impurities. The crude product is employed in the next stage without further purification.

15 R_f = (toluene:EtOH = 4.1 v/v).

Example 17

N-[2-S-(4',6'-O-Benzylidene- α -D-mannopyranosyl)thioethyl]succinimide 25

20 5.5 g (10.55 mol) of the thioglycoside 23 are treated at room temp. with 100 mg (1.8 mmol) of NaOMe in 75 ml of methanol. After 2 h, thin-layer chromatographic checking indicates incomplete conversion, for which reason a further 100 mg of NaOMe are added. After a further 1.5 h, the mixture is neutralized with acidic ion exchanger Amberlyst® 15. The ion
25 exchanger is filtered off and the solvent is removed in vacuo.

The crude product is treated in 50 ml of absol. DMF with 3.1 ml (20 mmol) of benzaldehyde dimethyl acetal and 112 mg (1 mmol) of p-toluenesulfonic acid and stirred at 50°C and 50-70 mbar for 2 h. The mixture is neutralized with 10 ml of triethylamine and the solvent is removed in a high vacuum.
30 The residue is taken up in 200 ml of CH_2Cl_2 and extracted twice with 75 ml of NaHCO_3 . The organic phase is separated off and dried over MgSO_4 , and the solvent is removed in vacuo. By chromatography on silica gel using petroleum ether/ethyl acetate mixtures, 3.33 g (77%) of a colorless oil are obtained.

35 R_f = 0.44 (EtOAc/HOAc = 30:1 v/v),

200 MHz- ^1H -NMR (CDCl_3): δ [ppm] = 7.43-7.41 (m, 2H, Ph); 7.33-7.29 (m, 3H, Ph); 5.48 (s, 1H, PhCH); 5.32 (s, 1H, H-1'); 4.19-3.43 (m, 8H, H-2', H-3', H-4', H-5', H-6a/b', NCH_2 Cya); 2.80-2.64 (m, 2H, SCH_2 Cya); 2.58 (s, 4H, COCH_2 Suc).

Example 18

N-[2-S-(2'-O-Acetyl-3'-O-allyl-4',6'-O-benzylidene- α -D-mannopyranosyl)-thioethyl]succinimide 26

5

330 mg (0.81 mmol) of 25 are treated with 227 mg (0.91 mmol) of dibutyltin oxide in 20 ml of methanol. The suspension is heated under reflux for 2.5 h, freed from the solvent in vacuo and the residue is dried in a high vacuum. The tin acetal is treated with 0.12 ml (1.4 mmol) of allyl bromide and 517 mg (1.4 mmol) of TBAI in 20 ml of absol. toluene and the mixture is stirred at 40°C for 6 h. Thin-layer chromatographic checking indicates a low conversion. The reaction is therefore continued at 80°C. After 7 h, the solvent is removed in vacuo and the product is isolated by chromatography on silica gel two times using petroleum ether/ethyl acetate mixtures.

15 300 mg of a rust-colored oil are obtained, which according to the NMR spectrum still contains tin residues.

$R_f = 0.56$ (toluene/EtOH = 4:1 v/v),

200 MHz- ^1H -NMR (CDCl_3): δ [ppm] = 7.48-7.43 (m, 2H, Ph); 7.36-7.31 (m, 3H, Ph); 5.96-5.77 (m, 1H, $\text{CH}_2=\text{CH}$); 5.55 (s, 1H, PhCH); 5.40 (s, 1H, H-1); 5.26 (dd, 1H, $J_{\text{gem,trans}} = 17.09$ Hz, $J_{\text{vic}} = 1.46$ Hz, $\text{CH}_2=\text{CH}$); 5.16 (dd, 1H, $J_{\text{gem,cis}} = 10.26$ Hz, $J_{\text{vic}} = 0.98$ Hz, $\text{CH}_2=\text{CH}$); 4.31-4.01 (m, 5H, H-2, H-6a/b, $=\text{CH}-\text{CH}_2$); 3.88-3.60 (m, 3H, H-4, NCH_2 Cya); 3.29-3.20 (m, 2H, H-3, H-5); 2.85-2.76 (m, 2H, SCH_2 Cya); 2.69 (s, 4H, COCH_2 Suc).

25 300 mg (0.67 mmol) of 25 are stirred at room temp. for 1.5 h with 2 ml of acetic anhydride and a spatula-tipful [lacuna] in 10 ml of pyridine. The solvent is removed in vacuo and the product is purified by chromatography on silica gel using petroleum ether/ethyl acetate (1:1). 191 mg (48% over two stages) of a yellow oil are obtained. $R_f = 0.63$ (toluene/EtOH = 4:1 v/v),

30 200 MHz- ^1H -NMR (CDCl_3): δ [ppm] = 7.47-7.42 (m, 2H, Ph); 7.37-7.32 (m, 3H, Ph); 5.96-5.70 (m, 1H, $\text{CH}_2=\text{CH}$); 5.66 (s, 1H, PhCH); 5.33-5.08 (m, 3H, $\text{CH}_2=\text{CH}$, H-2); 5.25 (s, 1H, H-1); 4.44-4.0 (m, 5H, H-4, H-6a/b, $=\text{CH}-\text{CH}_2$); 3.85-3.45 (m, 4H, H-3, H-5, NCH_2 Cya); 2.87-2.72 (m, 2H, SCH_2 Cya); 2.65 (s, 4H, COCH_2 Suc); 2.12 (s, 3H, CH_3 Ac).

Example 19

N-[2-S-(2'-O-Acetyl-3'-O-allyl-6'-O-tert-butyldiphenylsilyl-4'-O-(1''-(R/S)-ethoxyethyl)- α -D-mannopyranosyl)thioethyl]succinimide 27

- 5 0.12 ml of a 48% strength solution of HBF_4 in water is added to 180 mg (0.37 mmol) of 26 in 10 ml of absol. acetonitrile and the reaction solution is stirred for 2 h at room temp. Thin-layer chromatographic checking shows complete conversion and the mixture is treated with 10 ml of satd NaHCO_3 solution and extracted twice with 50 ml of CH_2Cl_2 . The combined organic
- 10 phases are dried over MgSO_4 and the solvent is removed in vacuo. The residue is taken up in 10 ml of absol. CH_2Cl_2 and treated with 0.19 ml (0.72 mmol) of tert-butyldiphenylchlorosilane and 100 mg (1.5 mmol) of imidazole. After 16 h, the solution is diluted with 50 ml of CH_2Cl_2 and extracted with 0.5 N HCl solution. The organic phase is dried over MgSO_4
- 15 and the solvent is removed in vacuo. By chromatography on silica gel using petroleum ether/ethyl acetate mixtures, 69 mg (30%) of the product are obtained as a colorless oil. $R_f = 0.21$ (toluene/EtOH = 4:1 v/v), 400MHz- ^1H -NMR (CDCl_3): δ [ppm] = 7.69-7.65 (m, 4H, PhSi); 7.41-7.33 (m, 6H, PhSi); 5.89-5.80 (m, 1H, $\text{CH}_2=\text{CH}$); 5.32 (d, 1H, $J_{1,2} = 1.17$ Hz, H-2); 5.29 (s, 1H, H-1); 5.26 (dd, 1H, $J_{\text{vic,trans}} = 16.88$ Hz, $J_{\text{gem}} = 1.62$ Hz, CH_2CH); 5.18 (dd, 1H, $J_{\text{vic,cis}} = 11.74$ Hz, $J_{\text{gem}} = 1.47$ Hz, CH_2CH); 4.11 (dd, 1H, $J_{\text{gem}} = 12.32$ Hz, $J_{\text{vic}} = 5.58$ Hz, $=\text{CH}-\text{CH}_2$); 4.00-3.88 (m, 5H, H-3, H-4 & H-6a/b); 3.78-3.71 (m, 1H, H-5); 3.64-3.58 (m, 2H, $\text{CH}_2\text{N Cya}$); 2.82-2.68 (m, 2H, $\text{SCH}_2\text{ Cya}$); 2.66 (s, 4H, $\text{CH}_2\text{CO Suc}$); 2.08 (s, 3H, CH_3
- 20 Ac); 1.03 (s, (h, 9H, CH_3 tBuSi).
- 25

- 69 mg (0.11 mmol) of the succinimide are reacted with 0.1 ml (1.1 mmol) of ethyl vinyl ether and 27 mg (0.11 mmol) of pyridinium toluene-4-sulfonate in 10 ml of absol. CH_2Cl_2 . After 4 h, the conversion according to the thin-
- 30 layer chromatogram is only about 70%, so a further 0.1 ml of ethyl vinyl ether and 27 mg of pyridinium toluene-4-sulfonate are added and the reaction time is lengthened to 16 h. The mixture is diluted with 30 ml of CH_2Cl_2 and extracted with satd NaHCO_3 solution. The organic phase is dried over MgSO_4 and the solvent is removed in vacuo. By
- 35 chromatography on silica gel using petroleum ether/ethyl acetate (1:1) 42 mg (56%) of a colorless oil are obtained. In addition, some fractions are obtained which contain unreacted starting material. $R_f = 0.44$ (toluene/EtOH = 4:1 v/v), $[\alpha] = +96.1^\circ$ (c 1, CHCl_3).

400MHz-¹H-NMR (CDCl₃): δ [ppm] = 7.70-7.63 (m, 4H, PhSi); 7.38-7.31 (m, 6H, PhSi); 5.88-5.76 (m, 1H, CH₂=CH); 5.32-5.26 (m, 2H, H-1 & H-2); 5.24, 5.18 (dm, 1H, J_{vic,trans} = 16.88 Hz, CH₂=CH); 5.15, 5.12 (dm, 1H, J_{vic,cis} = 10.57 Hz, CH₂=CH); 4.90-4.87 (m, 1H, CHCH₃ EE); 4.10-3.42 (m, 10.5H, H-3, H-4, H-5, H-6a/b, =CH-CH₂, CH₃CH₂O EE & CH₂N Cya); 3.22-3.16 (m, 0.5H, H-5); 2.74-2.70 (m, 2H, SCH₂ Cya); 2.67, 2.65 (2x s, 4H, CH₂CO Suc); 2.10, 2.09 (2x s, 3H, CH₃ Ac); 1.24-1.20 (m, 3H, CHCH₃ EE); 1.15 (t, 1.5H, J_{gem} = 7.05 Hz, CH₃CH₂O EE); 1.04, 1.02 (2x s 9H, CH₃ tBuSi); 0.89 (t, 1.5H, J_{gem} = 7.05 Hz, CH₃CH₂O EE).

10

The hydrolysis of the ester is carried out as described in the case of glucose.

General procedures for synthesis on polymeric supports

15

General procedure for the coupling of the thioglycosides to the amino-functionalized polymeric supports

20

A solution of 14.6 mmol of a thioglycoside is shaken overnight in a solid-phase reactor with 15.4 g (19.8 mmol, 1.28 mmol/g) of aminomethylpolystyrene, 3.7 ml (14.6 mmol) of diisopropylcarbodiimide and 3.86 g (14.6 mmol) of N-hydroxylbenzotriazole. The resin is then filtered off with suction and washed ten times with 50 ml each of DMF and dichloromethane. The loaded polymer is dried in vacuo and the loading with carbohydrate matrix is determined by means of elemental analysis. As a rule, the loading is 50-80% of the maximally possible loading.

25

General procedure for the removal of the glycoside derivatives from the polymeric support

30

a) analytical: A suspension of 80 mg (0.056 mmol) of polymer-bonded derivative in 1.5 ml of abs. dichloromethane is shaken at room temperature in a 5 ml PE syringe (PE frit, plastic cap) with 0.3 ml of a 3.5 percent solution of bromine in abs. dichloromethane and 0.08 ml (0. [lacuna] mmol) of 2,6-di-tert-butylpyridine or a corresponding amount of polymer-bonded 2,6-di-tert-butylpyridine. After 15 min, 0.2 ml of cyclohexene, 0.2 ml of the abs. alcohol to be glycosylated and 25 mg (0.056 mmol) of tetraethylammonium bromide are added. After 2.5 h, the resin is filtered off and washed five times with 1 ml of dichloromethane. The combined

35

filtrates are freed from the solvent in vacuo. The crude product obtained is applied to a silica gel cartridge in a little dichloromethane. It is eluted first with 30 ml of petroleum ether. This fraction is discarded. The product is obtained by eluting with petroleum ether/ethyl acetate (1:1).

5 Characterization by HPLC and MS analysis follows.

b) preparative: A suspension of 400 mg (0.312 mmol) of polymer-bonded galactose derivative in 3 ml of abs. dichloromethane is shaken at room temp. in a 5 ml PE syringe (PE frit, plastic cap) with 1.2 ml of a 3.5 percent solution of bromine in abs. dichloromethane and 0.32 ml (0. [lacuna] mmol) of 2,6-di-tert-butylpyridine. After 15 min, the resin is filtered off and washed five times with 3 ml of dichloromethane each time. 0.5 ml of cyclohexene, 0.5 ml of the abs. alcohol to be glycosylated and 25 mg (0.056 mmol) of tetraethylammonium bromide are added to the filtrate. The organic phase is washed with water, dried over magnesium sulfate and concentrated in vacuo, and the residue is purified by flash chromatography on silica gel.

General procedure for the alkylation of the glycosides on the polymeric support with potassium tert-butoxide

20 A solution of 87 mg (0.78 mmol) of potassium tert-butoxide in abs. DMF is added to a suspension of 100 mg (0.078 mmol) of loaded polymer in 1.5 ml of abs. DMF. The mixture is shaken for 15 min. It is filtered and the filtrate is discarded. 0.78 mmol of the appropriate alkylating agent in 1.5 ml of abs. DMF is then added and the mixture is shaken for 4 h. The resin is separated from the solution and washed five times with 2 ml each of DMF, toluene and dichloromethane.

General procedure for the alkylation of the galactosides on the polymeric support with tert-butyl-P₄ base (Schwesinger base)

30 A solution of 0.22 ml (0.22 mmol) of tert-butyl-P₄ base in abs. DMF is added to a suspension of 80 mg (0.056 mmol) of loaded polymer in 1.5 ml of abs. DMF. The mixture is shaken for 10 min. 0.56 mmol of the appropriate alkylating agent is then added and the mixture is shaken for 2-4 h. The resin is filtered off with suction from the solution and washed five times with 2 ml each of DMF, toluene and dichloromethane.

General procedure for the removal of the acetate protective group on the polymeric support

0.3 ml of a 30% strength sodium methanolate solution is added to a suspension of 0.0050 mmol of loaded polymer in 2 ml of dioxane/methanol. The mixture is shaken at room temperature for 3 h. The
5 resin is filtered off with suction from the solution and first washed five times with methanol/dioxane and then five times with 2 ml each of DMF and dichloromethane.

General procedure for the removal of the tert-butyldiphenylsilyl protective
10 group on the polymeric support

0.56 ml (0.56 mmol, 1M) of tetrabutylammonium fluoride in THF is added to a suspension of 0.056 mmol of loaded polymer in 1.5 ml of THF. The mixture is shaken for 4 h. The resin is filtered off with suction from the
15 solution and washed five times with 2 ml each of DMF and dichloromethane.

General procedure for the removal of the ethoxyethyl ether protective
20 group on the polymeric support

para-Toluenesulfonic acid is added to a suspension of 0.050 mmol of loaded polymer in 1.5 ml of dioxane/methanol and the mixture is stirred at 40°C for 4 h. The resin is filtered off with suction from the solution and washed with 0.5N HCl solution and then washed five times with 2 ml each
25 of DMF and dichloromethane.

The isopropyl protective group can be removed under analogous conditions. The reaction times or temperatures may change in this case.

30 General procedure for the removal of the allyl ether protective group on the polymeric support

A solution of 198 mg of zirconocene dichloride and 0.63 ml of BuLi (1.7 M) in THF is added at -70°C to a suspension of 0.060 mmol of loaded polymer
35 in 1.5 ml of THF. The mixture is stirred at room temperature for 4 h after addition is complete. The resin is filtered off with suction from the solution and washed with 0.5N HCl solution and then washed five times with 2 ml each of DMF and dichloromethane.

Example 20

Methyl 2-O-methyl-6-O-tert-butyldiphenylsilyl-3-O-propyl- α,β -D-glucopyranoside

- 5 166 mg (0.1 mmol) of the polymer are reacted with methyl iodide according to the general working procedure for an alkylation. After filtration on silica gel, 16 mg of a colorless oil are obtained. The product is purified by preparative HPLC and individual fractions are identified by mass spectrometric analysis and NMR spectra.
- 10 $R_f = 0.45, 0.34$ (PE/EtOAc = 4:1 v/v).
 FBA-MS (NBA-pos, LiCl): (m/e) = 495.2 (100%, $[M+Li]^+$, calc.: 295.2); 496.2 (26%, $[M+Li]^+$, ^{13}C , calc.: 496.2).

Example 21

- 15 Methyl 2-O-benzyl-6-O-tert-butyldiphenylsilyl-3-O-propyl- α,β -D-glucopyranoside

- 20 166 mg (0.1 mmol) of the polymer are reacted with benzyl bromide according to the general working procedure for an alkylation and the carbohydrate is removed from the resin. After filtration on silica gel, 15 mg of a colorless oil are obtained. The product is purified by preparative HPLC and individual fractions are identified by mass spectrometric analysis and NMR spectra.
- $R_f = 0.53, 0.47$ (PE/EtOAc = 4:1 v/v).
- 25 $C_{33}H_{44}O_6Si$ (564.8)
 FBA-MS (NBA-pos, LiCl): (m/e) = 571.1 (100%, $[M+Li]^+$, calc.: 571.3); 572.1 (35%, $[M+Li]^+$, ^{13}C , calc.: 572.3).

Example 22

- 30 Methyl 2-O-propyl-6-O-tert-butyldiphenylsilyl-3-O-propyl- α,β -D-glucopyranoside

- 35 166 mg (0.1 mmol) of the polymer are reacted with n-propyl iodide according to the general working procedure for an alkylation and the carbohydrate is removed from the resin. After filtration on silica gel, 18 mg of a colorless oil are obtained.
- $R_f = 0.49, 0.46$ (PE/EtOAc = 4:1 v/v).
 $C_{29}H_{44}O_6Si$ (516.8)

FBA-MS (NBA-pos, LiCl): (m/e) = 523.3 (100%, $[M+Li]^+$, calc.: 523.2); 524.3 (33%, $[M+Li]^+$, calc.: 524.2).

Example 23

- 5 Methyl 2-O-(2'-naphthyl)methyl-6-O-tert-butyldiphenylsilyl-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with (2-naphthyl)methylene bromide according to the general working procedure for an alkylation and the carbohydrate is removed from the resin. After filtration on silica gel, 22 mg of a slightly yellowish oil are obtained. It was possible to detect the desired product by mass spectrometry.

Example 24

- 15 Methyl 2-O-isopropyl-6-O-tert-butyldiphenylsilyl-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with 2-propyl bromide according to the general working procedure for an alkylation and the carbohydrate is removed from the resin. After filtration on silica gel, 17 mg are obtained. The product can be detected by mass spectrometry.

HPLC (gradient 54/80): R_t (min) = 3.61 (15.3%, DTBpy); 14.2 (11.0%, 2-OH); 15.7 (19.2%).

Example 25

Methyl 2-O-(4'-cyanobenzyl)-6-O-tert-butyldiphenylsilyl-3-O-propyl- α,β -D-glucopyranoside

30 166 mg (0.1 mmol) of the polymer are reacted with o-cyanobenzyl bromide according to the general working procedure for an alkylation and the carbohydrate is removed from the resin. After filtration on silica gel, 22 mg of a slightly yellowish oil are obtained. R_f = 0.52, 0.47 (PE/EtOAc = 4:1 v/v).

- 35 $C_{33}H_{43}NO_6Si$ (577.8)

FBA-MS (NBA-pos, LiCl): (m/e) = 241.1 (69%); 596.3 (38%, $[M+Li]^+$, calc.: 596.4); 597.3 (22%, $[M+Li]^+$, ^{13}C , calc.: 597.4); 746.4 (66%, $[M+C_4H_8^{79}BrO+H]^+$, calc.: 746.4); 747.4 (64%, $[M+C_4H_8^{79}BrO+H]^+$, ^{13}C ,

49

calc.: 747.4); 748.4 (100%, $[M+C_4H_8^{81}BrO+H]^+$, calc.: 748.4); 749.4 (57%, $[M+C_4H_8^{81}BrO+H]^+$, ^{13}C , calc.: 749.4).

Example 26

- 5 Methyl 2-O-heptyl-6-O-tert-butyldiphenylsilyl-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with 1-iodoheptane according to the general working procedure for an alkylation and the carbohydrate is removed from the resin. After filtration on silica gel, 25 mg of a colorless oil are obtained.

$R_f = 0.65, 0.49$ (PE/EtOAc = 4:1 v/v).

$C_{33}H_{43}NO_6Si$ (577.79)

FBA-MS (NBA-pos, LiCl): (m/e) = 241.1 (100%); 579.4 (74%, $[M+Li]^+$, calc.: 579.4); 580.4 (31%, $[M+Li]^+$, ^{13}C , calc.: 580.4); 729.4 (72%, $[M+C_4H_8^{79}BrO+H]^+$, calc.: 729.4); 730.4 (38%, $[M+C_4H_8^{79}BrO+H]^+$, ^{13}C , calc.: 730.4); 731.4 (79%, $[M+C_4H_8^{81}BrO+H]^+$, calc.: 731.4); 732.4 (34%, $[M+C_4H_8^{81}BrO+H]^+$, ^{13}C , calc.: 723.4).

20 Example 27

Methyl 2-O-(2'-methoxy-5'-nitrobenzyl)-6-O-tert-butyldiphenylsilyl-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with 2-methoxy-5-nitrobenzyl bromide according to the general working procedure for an alkylation and the carbohydrate is removed from the resin. After filtration on silica gel, 15 mg of a slightly yellowish oil are obtained.

$R_f = 0.48, 0.43$ (PE/EtOAc = 4:1 v/v).

$C_{33}H_{45}NO_9Si$ (627.8)

30

Example 28

Methyl 2-O-methyl-6-O-benzyl-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with methyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is alkylated with benzyl bromide. After removal of the carbohydrate from the resin and filtration on silica gel, 24 mg of a colorless oil are obtained. For confirmed characterization of the

35

product and assignment of the two anomers, the product was purified by preparative HPLC (gradient 90/10).

$C_{18}H_{28}O$ (340.4)

$R_f = 0.63, 0.54$ (PE/EtOAc = 3:1 v/v).

- 5 400MHz- ^1H -NMR ($CDCl_3$): δ [ppm] = 7.32-7.31 (m, 5H, Ph); 4.83 (d, 1H, $J_{2,1} = 3.81$ Hz, H-1); 4.60 (d, 1H, $J_{gem} = 12.32$ Hz, CH_2Ph); 4.55 (d, 1H, $J_{gem} = 12.03$ Hz; CH_2Ph); 4.20 (dd, 0.3H, $J_{1,2} = 3.23$ Hz, $J_{3,2} = 12.03$ Hz, = H-2), 4.12 (dd, 0.7H, $J_{1,2} = 2.94$ Hz, $J_{3,2} = 11.45$ Hz, H-2); 4.00 (dd, 0.3H, $J_{vic} = 11.89$ Hz, $J_{gem} = 6.61$ Hz, OCH_2 Pr); 3.92 (dd, 0.7H, $J_{vic} = 11.44$ Hz, $J_{gem} = 7.34$ Hz, OCH_2 Pr); 3.83-3.82 (m, 1.6H, H-6a/b); 3.81-3.39 (m, 8.4 Hz, H-3, H-4, H-6b); 3.47, 3.41 (2x s, 6H, OCH_3); 3.23-3.21 (m, 1H, H-5); 1.91 (sb, 1H, OH); 1.61-1.54 (m, 2H, OCH_2CH_2 Pr); 0.91 (t, 3H, $J_{gem} = 7.34$ Hz, CH_3 Pr);
- 10

15 Example 29

Benzyl 2-O-methyl-6-O-methyl-3-O-propyl- α,β -D-glucopyranoside

- 166 mg (0.1 mmol) of the polymer are reacted with methyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with methyl iodide. After removal of the carbohydrate from the resin and filtration on silica gel, 20 mg of a colorless oil are obtained.
- 20

$R_f = 0.46, 0.34, 0.1$ (OH) (PE/EtOAc = 3:1 v/v).

$C_{18}H_{28}O_6$ (340.4)

- 25 FBA-MS (NBA-pos, LiCl): (m/e) = 233.1 (Gly^+ , calc.: 233.1); 347.0 ($[M+Li]^+$, calc.: 347.2); 497.2 ($[M+C_4H_8^{79}BrO+Li]^+$, calc.: 497.2); 499.2 ($[M+C_4H_8^{81}BrO+Li]^+$, calc.: 497.2).

Example 30

- 30 Methyl 2-O-methyl-6-O-heptyl-3-O-propyl- α,β -D-glucopyranoside

- 166 mg (0.1 mmol) of the polymer are reacted with methyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with heptyl iodide. After removal of the carbohydrate from the resin and filtration on silica gel, 27 mg of a colorless oil are obtained.
- 35

$R_f = 0.53, 0.42$ (PE/EtOAc = 3:1 v/v).

$C_{18}H_{36}O_6$ (348.5)

FBA-MS (NBA-pos, LiCl): (m/e) = 355.2 (22%, $[M+Li]^+$, calc.: 355.2); 369.2 (88%); 505.2 (100%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 505.2); 507.2 (99%, $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 507.2).

5 Example 31

Isopropyl 2-O-methyl-6-O-heptyl-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with methyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with heptyl iodide. After removal of the carbohydrate from the resin and filtration on silica gel, 21 mg of a colorless oil are obtained.

R_f = 0.69, 0.63 (PE/EtOAc = 3:1 v/v).

$C_{20}H_{40}O_6$ (376.5)

15 FBA-MS (NBA-pos, LiCl): (m/e) = 533.2 (100%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 533.2); 534.2 (28%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 534.2); 535.2 (99%, $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 535.2); 536.2 (26%, $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C , calc.: 536.2).

20 Example 32

Ethyl 2-O-methyl-6-O-(2'-methoxy-5'-nitrobenzyl)-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with methyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with 2-methoxy-5-nitrobenzyl bromide. After removal of the carbohydrate from the resin and filtration on silica gel, 10 mg of a colorless oil are obtained.

R_f = 0.35, 0.27 (PE/EtOAc = 3:1 v/v).

30 $C_{18}H_{36}O_6$ (429.5)

FBA-MS (NBA-pos, LiCl): (m/e) = 384.1 (3%, Gly^+ , calc.: 384.2); 435.1 (22%); 436.1 (22%, $[M+Li]^+$, calc.: 436.2); 586.1 (97%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 586.2); 587.1 (36%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 387.2); 588.1 (100%, $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 588.2); 589.1 (34%, $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C , calc.: 589.2).

Example 33

Methyl 2-O-benzyl-6-O-isopropyl-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with benzyl bromide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with 2-bromopropane. After removal of the carbohydrate from the resin and filtration on silica gel, 9 mg of a slightly yellowish oil are obtained.

$R_f = 0.62$ (PE/EtOAc = 3:1 v/v).

10 $C_{20}H_{32}O_6$ (368.5)

FBA-MS (NBA-pos, LiCl): (m/e) = 301.1 (83%); 373.2 (19%); 525.2 (17%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 525.2); 527.2 (15%, $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 527.2); 573.2 (13%); 575.2 (17%); 667.3 (21%); 669.3 (19%).

15 Example 34

Ethyl 2-O-benzyl-6-O-(4'-cyanobenzyl)-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with benzyl bromide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with 4-cyanobenzyl bromide. After removal of the carbohydrate from the resin and filtration on silica gel, 25 mg of a colorless oil are obtained.

$R_f = 0.66, 0.55$ (PE/EtOAc = 3:1 v/v).

$C_{26}H_{33}NO_6$ (455.6)

25 FBA-MS (NBA-pos, LiCl): (m/e) = 410.2 (5%, Gly⁺, calc.: 410.2); 462.2 (34%, $[M+Li]^+$, calc.: 462.2); 587.2 (13%, ^{79}Br); 589.2 (13%, ^{81}Br); 612.2 (97%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 612.2); 613.2 (43%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 613.2); 614.2 (100%, $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 614.2); 615.2 (32%, $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C , calc.: 615.2); 654.2 (11%, ^{79}Br); 656.2 (13%, ^{81}Br).

Example 35

Methyl 2-O-benzyl-6-O-heptyl-3-O-propyl- α,β -D-glucopyranoside

35 166 mg (0.1 mmol) of the polymer are reacted with benzyl bromide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with heptyl iodide. After removal of the carbohydrate from the resin and filtration on silica gel, 32 mg of a colorless oil are obtained.

$R_f = 0.82, 0.76$ (PE/EtOAc = 3:1 v/v).

$C_{24}H_{40}NO_6$ (424.6)

FBA-MS (NBA-pos, LiCl): (m/e) = 431.3 (14%, $[M+Li]^+$, calc.: 431.2); 581.2 (100%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 581.3); 582.2 (40%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 582.3); 583.2 (100%, $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 583.3); 584.2 (30%, $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C , calc.: 584.3).

Example 36

Isopropyl 2-O-benzyl-6-O-cyclohexylmethyl-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with benzyl bromide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with cyclohexylmethylene bromide. After removal of the carbohydrate from the resin and filtration on silica gel, 15 mg of a colorless oil are obtained.

$R_f = 0.85, 0.61$ (PE/EtOAc = 3:1 v/v).

$C_{26}H_{42}O_6$ (450.6)

FBA-MS (NBA-pos, LiCl): (m/e) = 301.1 (59%); 443.3 (47%); 457.3 (34%, $[M+Li]^+$, calc.: 457.3); 517.2 (22%, ^{79}Br); 519.2 (22%, ^{81}Br); 607.2 (98%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 607.3); 608.2 (42%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 608.3); 609.2 (100%, $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 609.3); 610.2 (34%, $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C , calc.: 610.3); 669.4 (22%).

Example 37

Methyl 2-O-propyl-6-O-(4'-cyanobenzyl)-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with n-propyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with 4-cyanobenzyl bromide. After removal of the carbohydrate from the resin and filtration on silica gel, 17 mg of a colorless oil are obtained.

$R_f = 0.63, 0.53$ (PE/EtOAc = 3:1 v/v).

$C_{25}H_{31}NO_6$ (393.5)

FBA-MS (NBA-pos, LiCl): (m/e) = 382.0 (9%, ^{79}Br); 384.0 (9%, ^{81}Br); 400.2 (28%, $[M+Li]^+$, calc.: 400.2); 414.2 (11%); 442.2 (11%); 477.2 (9%, ^{79}Br); 479.2 (9%, ^{81}Br); 515.2 (9%); 550.2 (100%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 550.2); 551.2 (36%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 551.2); 552.2

(98%, $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 552.2); 553.2 (28%, $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C , calc.: 553.2).

Example 38

5 Isopropyl 2-O-propyl-6-O-benzyl-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with n-propyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with benzyl bromide. After removal of the carbohydrate from the resin and filtration on silica gel, 18 mg of a colorless oil are obtained.

$R_f = 0.73, 0.64$ (PE/EtOAc = 3:1 v/v).

$C_{22}H_{36}O_6$ (396.5)

FBA-MS (NBA-pos, LiCl): (m/e) = 229.1 (94%); 389.2 (66%); 403.2 (43%, $[M+Li]^+$, calc.: 403.3); 445.2 (19%); 505.2 (27%, ^{79}Br); 507.2 (27%, ^{81}Br); 553.2 (82%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 553.2); 554.2 (33%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 554.3); 555.2 (85%, $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 555.3); 556.2 (26%, $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C , calc.: 556.3).

20 Example 39

Benzyl 2-O-propyl-6-O-methyl-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with n-propyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with methyl iodide. After removal of the carbohydrate from the resin and filtration on silica gel, 149 mg of a colorless oil are obtained. The product contains still larger amounts of benzyl alcohol.

$R_f = 0.46, 0.40$ (PE/EtOAc = 3:1 v/v).

30 $C_{20}H_{32}O_6$ (368.5)

FBA-MS (NBA-pos, LiCl): (m/e) = 261.2 (8%, Gly $^+$, calc.: 261.2); 375.2 (25%; $[M+Li]^+$, calc.: 375.2); 525.2 (15%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 525.2); 526.2 (5%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 526.2); 527.2 (15%, $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 527.2); 528.2 (4%, $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C , calc.: 528.2).

Example 40

Methyl 2-O-propyl-6-O-heptyl-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with n-propyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with heptyl iodide. After removal of the carbohydrate from the resin and filtration on silica gel, 24 mg of a colorless oil are obtained.

R_f = 0.58, 0.51 (PE/EtOAc = 3:1 v/v).

10 $C_{20}H_{40}O_6$ (376.5)

FBA-MS (NBA-pos, LiCl): (m/e) = 383.3 (35%; $[M+Li]^+$, calc.: 383.3); 397.3 (100%); 439.3 (28%); 453.4 (48%).

Example 41

15 Ethyl 2-O-pentyl-6-O-heptyl-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with pentyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with heptyl iodide.

20 After removal of the carbohydrate from the resin and filtration on silica gel, 20 mg of a colorless oil are obtained.

R_f = 0.87, 0.80 (PE/EtOAc = 3:1 v/v).

$C_{23}H_{46}O_6$ (418.3)

25 FBA-MS (NBA-pos, LiCl): (m/e) = 425.3 (76%; $[M+Li]^+$, calc.: 452.3); 426.3 (19%, $[M+Li]^+$, ^{13}C , calc.: 426.3); 575.3 (100%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 575.3); 576.3 (38%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 576.3); 577.3 (98%, $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 577.3); 578.3 (28%, $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C , calc.: 578.3).

30 Example 42

Isopropyl 2-O-pentyl-6-O-methyl-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with pentyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with methyl iodide. After removal of the carbohydrate from the resin and filtration on silica gel, 18 mg of a colorless oil are obtained.

R_f = 0.67, 0.64 (PE/EtOAc = 3:1 v/v).

$C_{18}H_{36}O_6$ (348.3)

FBA-MS (NBA-pos, LiCl): (m/e) = 341.0 (100%); 355.1 (62%; $[M+Li]^+$, calc.: 355.2); 505.2 (83%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 505.2); 506.2 (28%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 506.2); 507.2 (80%, $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 507.2); 508.2 (19%, $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C , calc.: 508.2).

5

Example 43

Benzyl 2-O-heptyl-6-O-cyclohexylmethyl-3-O-propyl- α,β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with heptyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with cyclohexylmethylene bromide. After removal of the carbohydrate from the resin and filtration on silica gel, 136 mg of a colorless oil are obtained. The product contains still larger amounts of benzyl alcohol.

15 $R_f = 0.77, 0.67$ (PE/EtOAc = 3:1 v/v).

$C_{30}H_{50}O_6$ (506.7)

FBA-MS (NBA-pos, LiCl): (m/e) = 285.2 (65%); 309.2 (100%); 339.2 (45%); 399.3 (39%, Gly^+ , calc.: 399.3); 451.3 (48%); 513.3 (22%; $[M+Li]^+$, calc.: 513.3); 663.3 (24%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 663.3); 665.3 (34%, $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 665.3).

20

Example 44

Methyl 2-O-heptyl-6-O-benzyl-3-O-propyl- α,β -D-glucopyranoside

25 166 mg (0.1 mmol) of the polymer are reacted with heptyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with benzyl bromide. After removal of the carbohydrate from the resin and filtration on silica gel, 32 mg of a colorless oil are obtained.

30 $R_f = 0.86, 0.77$ (PE/EtOAc = 3:1 v/v).

$C_{24}H_{40}O_6$ (424.6)

FBA-MS (NBA-pos, LiCl): (m/e) = 341.2 (7%; $[6-OH+Li]^+$, calc.: 341.3); 431.3 (13%, $[M+Li]^+$, calc.: 413.3); 459.2 (15%, ^{79}Br); 461.3 (14%, ^{81}Br); 491.2 (16%, $[6-OH+C_4H_8^{79}BrO+Li]^+$, calc.: 491.3); 493.2 (15%, $[6-OH+C_4H_8^{81}BrO+Li]^+$, calc.: 493.3); 581.2 (95%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 581.3); 582.3 (38%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 582.3); 583.2 (100%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 583.2); 584.2 (31%, $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C ,

35

57

calc.: 584.3); 589.2 (68%, ^{79}Br); 590.2 (25%, ^{79}Br , ^{13}C); 591.2 (64%, ^{81}Br); 592.2 (19%, ^{81}Br , ^{13}C).

Example 45

- 5 Isopropyl 2-O-heptyl-6-O-(4'-cyanobenzyl)-3-O-propyl-D- α,β -glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with heptyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with 4-cyanobenzyl bromide. After removal of the carbohydrate from the resin and filtration on silica gel, 21 mg of a colorless oil are obtained.

$R_f = 0.76, 0.74$ (PE/EtOAc = 3:1 v/v).

$\text{C}_{27}\text{H}_{43}\text{NO}_6$ (477.6)

- 15 FBA-MS (NBA-pos, LiCl): (m/e) = 408.2 (10%; Gly $^+$, calc.: 408.2); 470.3 (20%); 484.3 (21%, [M+Li] $^+$, calc.: 484.3); 582.4 (22%); 617.3 (24%, ^{79}Br); 619.3 (24%, ^{81}Br); 634.2 (85%, [M+C₄H₈ $^{79}\text{BrO+Li}$] $^+$, calc.: 634.3); 635.2 (38%, [M+C₄H₈ $^{79}\text{BrO+Li}$] $^+$, ^{13}C , calc.: 635.3); 636.2 (92%, [M+C₄H₈ $^{81}\text{BrO+Li}$] $^+$, calc.: 636.3); 637.2 (31%, [M+C₄H₈ $^{81}\text{BrO+Li}$] $^+$, ^{13}C , calc.: 637.3).

Example 46

Methyl 2-O-heptyl-6-O-isopropyl-3-O-propyl- α,β -D-glucopyranoside

- 25 166 mg (0.1 mmol) of the polymer are reacted with heptyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with n-propyl iodide. After removal of the carbohydrate from the resin and filtration on silica gel, 13 mg of a colorless oil are obtained.

30 $R_f = 0.80, 0.72$ (PE/EtOAc = 3:1 v/v).

$\text{C}_{20}\text{H}_{40}\text{O}_6$ (376.5)

FBA-MS (NBA-pos, LiCl): (m/e) = 383.3 (36%, [M+Li] $^+$, calc.: 383.3); 397.3 (76%); 533.3 (19%, [M+C₄H₈ $^{79}\text{BrO+Li}$] $^+$, calc.: 533.3); 535.3 (16%, [M+C₄H₈ $^{81}\text{BrO+Li}$] $^+$, calc.: 535.3); 611.2 (40%).

Example 47

Methyl 2-O-heptyl-6-O-(4'-bromobenzyl)-3-O-propyl-D-glucopyranoside

166 mg (0.1 mmol) of the polymer are reacted with heptyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with 4-bromobenzyl bromide. After removal of the carbohydrate from the resin and filtration on silica gel, 16 mg of a colorless oil are obtained.

$R_f = 0.84, 0.73$ (PE/EtOAc = 3:1 v/v).

10 $C_{24}H_{39}BrO_6$ (503.5)

FBA-MS (NBA-pos, LiCl): (m/e) = 453.4 (100%); 509.2 (41%, $[M+Li]^+$, ^{79}Br , calc.: 509.3); 511.2 (37%, $[M+Li]^+$, ^{81}Br , calc.: 511.3); 523.2 (84%, ^{79}Br); 525.2 (83%, ^{81}Br); 659.0 (34%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{79}Br , calc.: 659.3); 661.0 (59%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{81}Br , calc.: 661.3); 663.0 (31%,
15 $[M+C_4H_8^{81}BrO+Li]^+$, ^{81}Br , calc.: 663.3).

Example 48

Methyl 2-O-ethyl-6-O-benzyl-3-O-propyl- α,β -D-glucopyranoside

20 166 mg (0.1 mmol) of the polymer are reacted with ethyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with benzyl bromide. After removal of the carbohydrate from the resin and filtration on silica gel, 22 mg of a colorless oil are obtained.

25 $R_f = 0.82, 0.77$ (PE/EtOAc = 3:1 v/v).

$C_{19}H_{30}O_6$ (354.4)

FBA-MS (NBA-pos, LiCl): (m/e) = 271.2 (7%, $[6-OH+Li]^+$, calc.: 271.2); 361.3 (15%, $[M+Li]^+$, calc.: 361.2); 421.2 (11%, $[6-OH+C_4H_8^{79}BrO+Li]^+$, calc.: 421.2); 423.2 (9%, $[6-OH+C_4H_8^{81}BrO+Li]^+$, calc.: 423.2); 449.3
30 (38%); 451.3 (41%); 511.3 (96%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 511.2); 512.3 (36%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 512.2); 513.3 (100%, $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 513.2); 514.3 (26%, $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C , calc.: 514.2).

Example 49

Methyl 2-O-ethyl-6-O-(2'-methoxy-5'-nitrobenzyl)-3-O-propyl- α,β -D-glucopyranoside

- 5 166 mg (0.1 mmol) of the polymer are reacted with ethyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with 2-methoxy-5-nitrobenzyl bromide. After removal of the carbohydrate from the resin and filtration on silica gel, 15 mg of a colorless oil are obtained.
- 10 $R_f = 0.43$ (PE/EtOAc = 3:1 v/v).
 $C_{20}H_{31}NO_9$ (429.5)
 FBA-MS (NBA-pos, LiCl): (m/e) = 436.2 (9%, $[M+Li]^+$, calc.: 436.2); 586.1 (95%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 586.2); 587.1 (35%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 587.2); 588.1 (100%; $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 588.2); 588.1
 15 (28%, $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C , calc.: 589.2).

Example 50

Benzyl 2-O-ethyl-6-O-heptyl-3-O-propyl- α,β -D-glucopyranoside

- 20 166 mg (0.1 mmol) of the polymer are reacted with ethyl iodide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with methyl iodide. After removal of the carbohydrate from the resin and filtration on silica gel, 116 mg of a colorless oil are obtained. The product contains still larger
 25 amounts of benzyl alcohol.
 $R_f = 0.61, 0.54$ (PE/EtOAc = 3:1 v/v).
 $C_{25}H_{42}O_6$ (438.6)
 FBA-MS (NBA-pos, LiCl): (m/e) = 445.3 (29%, $[M+Li]^+$, calc.: 445.3); 595.3 (100%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 595.3); 596.3 (40%,
 30 $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 596.3); 597.3 (98%; $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 597.3); 598.3 (31%, $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C calc.: 595.3).

Example 51

Ethyl 2-O-(2'-cyanobenzyl)-6-O-heptyl-3-O-propyl- α,β -D-glucopyranoside

- 35 166 mg (0.1 mmol) of the polymer are reacted with 2-cyanobenzyl bromide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with

heptyl iodide. After removal of the carbohydrate from the resin and filtration on silica gel, 26 mg of a colorless oil are obtained.

$R_f = 0.78, 0.70$ (PE/EtOAc = 3:1 v/v).

$C_{26}H_{41}NO_6$ (463.6)

- 5 FBA-MS (NBA-pos, LiCl): (m/e) = 418.2 (12%, Gly^+ , calc.: 418.3); 470.3 (100%, $[M+Li]^+$, calc.: 470.3); 471.3 (32%, $[M+Li]^+$, ^{13}C , calc.: 471.3); 568.4 (78%); 603.3 (15%, ^{79}Br); 605.3 (15%, ^{79}Br); 620.2 (57%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 620.3); 621.2 (25%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 621.3); 622.2 (55%; $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 622.3); 623.2 (19%,
10 $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C , calc.: 623.3).

Example 52

Isopropyl 2-O-(2'-cyanobenzyl)-6-O-methyl-3-O-propyl- α,β -D-glucopyranoside

15

166 mg (0.1 mmol) of the polymer are reacted with 2-cyanobenzyl bromide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with methyl iodide. After removal of the carbohydrate from the resin and
20 filtration on silica gel, 21 mg of a colorless oil are obtained.

$R_f = 0.52, 0.47$ (PE/EtOAc = 3:1 v/v).

$C_{21}H_{31}NO_6$ (393.5)

- FBA-MS (NBA-pos, LiCl): (m/e) = 334.1 (8%, Gly^+ , calc.: 334.2); 386.2 (12%); 400.2 (27%, $[M+Li]^+$, calc.: 400.2); 414.2 (12%); 449.2 (10%, ^{79}Br);
25 451.3 (10%; ^{81}Br); 550.2 (98%, $[M+C_4H_8^{79}BrO+Li]^+$, calc.: 550.2); 551.2 (36%, $[M+C_4H_8^{79}BrO+Li]^+$, ^{13}C , calc.: 551.2); 552.2 (100%; $[M+C_4H_8^{81}BrO+Li]^+$, calc.: 552.2); 553.2 (28%, $[M+C_4H_8^{81}BrO+Li]^+$, ^{13}C calc.: 553.2).

30 Example 53

Methyl 2,4-di-O-benzyl-6-O-methyl-3-O-propyl- α,β -D-glucopyranoside

- 166 mg (0.1 mmol) of the polymer are converted into the derivative analogously to the experiments described above. For the removal of the 1-
35 ethoxyethyl protective group, the carbohydrate matrix is suspended in a mixture of 4 ml of dioxane, 0.4 ml of alcohol (methanol, propanol, octanol or benzyl alcohol) and a spatula-tipful of pyridinium toluene-4-sulfonate. The syringe is then closed with a plastic cap and shaken at room temp. for

16 h. Following this, the resin is washed five times with 3 ml of dioxane p.a. each time and twice with 3 ml of DMF.

After the removal of the 1-ethoxyethyl protective group, the 4-position is
5 alkylated with benzyl bromide and the carbohydrate is removed from the resin. 12-16 mg of a colorless oil are obtained. As all crude products contain identical components according to TLC, HPLC and MS, they are combined and purified by chromatography on silica gel using petroleum ether/ethyl acetate (12:1).

10 $C_{25}H_{34}O_6$ (430.6)

FAB-MS (NBA-pos, LiCl): (m/e) = 437.1 (100%, $[M+Li]^+$, calc.: 437.2); 438.1 (28%, $[M+Li]^+$, ^{13}C , calc: 438.2); 513.3 (28%, $[C_{31}H_{38}O_6+Li]^+$, 6-OBn, calc: 506.3).

400MHz- ^1H -NMR ($CDCl_3$): δ [ppm] = 7.50-7.25 (m, 10H, Ph); 4.86 (d, 1H, J_{gem} = 10.86 Hz, CH_2Ph); 4.77 (d, 1H, J_{gem} = 12.32, CH_2Ph); 4.60 (d, 1H, J_{gem} = 12.03 Hz, CH_2Ph); 4.56 (d, 1H, J_{gem} = 10.86 Hz, CH_2Ph); 4.52 (d, 1H, $J_{2,1}$ = 3.81 Hz, H-1); 3.87-3.24 (m, 8H, H-2, H-3, H-4, H-5, H-6a/b, OCH_2Pr); 3.32, 3.30 (2x s, 6H, OCH_3); 1.66-1.61 (m, 2H, OCH_2CH_2Pr); 0.92 (t, J_{gem} = 7.48 Hz, CH_3Pr).

20 6-Benzyl derivative:

400MHz- ^1H -NMR ($CDCl_3$): δ [ppm] = 7.51-7.26 (m, 15H, Ph); 4.86 (d, 1H, J_{gem} = 11.15 Hz, CH_2Ph); 4.80 (d, 1H, J_{gem} = 10.57 Hz, CH_2Ph); 4.67 (d, 1H, J_{gem} = 11.15 Hz, CH_2Ph); 4.61-4.54 (m, 2H, CH_2Ph); 4.50 (d, 1H, J_{gem} = 10.57 Hz, CH_2Ph); 4.25-4.18 (m, 3H, H-1 & OCH_2Pr); 3.80-3.29
25 (m, 6H, H-2, H-3, H-4, H-5, H-6a/b); 3.54 (s, 3H, OCH_3); 1.68-1.60 (m, 2H, OCH_2CH_2Pr); 0.91-0.86 (m, 3H, CH_3Pr).

Example 54

Methyl 2-O-benzyl-4-O-(2'-bromo-1'-ethoxy)ethyl-6-O-methyl-3-O-propyl-
30 α/β -D-glucopyranoside

166 mg (0.1 mmol) of the polymer are converted into the derivative analogously to the experiments described above. By treatment of the resin with various combinations of solvents and 1M citric acid (4 ml of DMF +
35 0.4 ml of 1M citric acid; 4 ml of DMF + 0.4 ml of 1M citric acid + 4 h ultrasound; 4 ml of DMF + 0.04 ml of 1M citric acid; 4 ml of dioxane + 0.04 ml of 1M citric acid; 4 ml of CH_2Cl_2 + 1 ml of acetone + 0.1 ml of 1M citric acid), it is attempted to remove the 1-ethoxyethyl protective group.

The resin is then subjected to a benzylation and the carbohydrate is removed from the polymeric support. 13-17 mg of a colorless oil are obtained. As all crude products contain identical components according to TLC, HPLC and MS, they are combined and purified by chromatography on silica gel using petroleum ether/ethyl acetate (8:1). C₂₂H₃₅BrO₇ (491.4) FAB-MS (NBA-pos, LiCl) of crude product: (m/e) = 497.4 (100%, [M+Li]⁺, ⁷⁹Br, calc.: 497.2); 498.4 (31%, [M+Li]⁺, ⁷⁹Br, ¹³C, calc. 498.2); 499.4 (98%, [M+Li]⁺, ⁸¹Br, calc.: 499.2); 500.4 (22%, [M+Li]⁺, ⁸¹Br, ¹³C, calc.: 500.2); 573.2 (28%, [C₂₈H₃₉⁷⁹BrO₇+Li]⁺ 6-OBn, calc.: 573.3); 575.2 (31%, [C₂₈H₃₉⁸¹BrO₇+Li]⁺ 6-OBn, calc.: 575.3).

1st diastereomer:

400MHz-¹H-NMR (CDCl₃): δ [ppm] = 7.33-7.25 (m, 5H, Ph); 4.87-4.84 (m, 1H, CHCH₂ EEBR); 4.72 (d, 1H, J_{gem} = 12.03 Hz, OCH₂Ph); 4.64 (d, 1H, J_{2,1} = 3.82 Hz, H-1); 4.56 (d, 1H, J_{gem} = 12.03 Hz, OCH₂Ph); 3.90-3.85 (m, 1H, OCH₂); 3.75-3.54 (m, 6H, H-2, H-6a/b, OCH₂); 3.52 (s, 3H, OCH₃); 3.48-3.31 (m, 5H, H-3, H-4, H-5 & CH₂Br EEBR); 3.36 (s, 3H, OCH₃); 1.63-1.57 (m, 2H, OCH₂CH₂ Pr); 1.25-1.18 (m, 3H, OCH₂CH₃ EEBR &); 0.89 (t, J_{gem} = 7.49 Hz, CH₃ Pr).

400MHz-¹H-NMR (CDCl₃): δ [ppm] = 7.34-7.26 (m, 5H, Ph); 4.97-4.95 (m, 1H, CHCH₂ EEBR); 4.72 (d, 1H, J_{gem} = 12.03 Hz, OCH₂Ph); 4.66 (d, 1H, J_{2,1} = 3.52 Hz, H-1); 4.56 (d, 1H, J_{gem} = 12.03 Hz, OCH₂Ph); 3.93-3.87 (m, 1H, OCH₂); 3.70-3.57 (m, 6H, H-2, H-6a/b, OCH₂); 3.54-3.30 (m, 5H, H-3, H-4, H-5 & CH₂Br EEBR); 3.49 (s, 3H, OCH₃); 3.36 (s, 3H, OCH₃); 1.64-1.59 (m, 2H, OCH₂CH₂ Pr); 1.23-1.19 (m, 3H, OCH₂CH₃ EEBR &); 0.89 (t, J_{gem} = 6.75 Hz, CH₃ Pr). HPLC (gradient 90/10):

During chromatography, the derivative benzylation in the 6-position, methyl 2,6-di-O-benzyl-4-O-(2-bromo-1-ethoxy)-ethyl-3-O-propyl-D-glucopyranoside, which has already been observed in the FAB mass spectrum, can be separated off.

C₂₈H₃₉BrO₇ (567.51)

400MHz-¹H-NMR (CDCl₃): δ [ppm] = 7.33-7.25 (m, 10H, Ph); 4.96-4.49 (m, 6H, H-1, OCH₂Ph & CHCH₂Br EEBR); 3.92-3.10 (m, 12H, H-2, H-3, H-4, H-5, H-6a/b, OCH₂ & CH₂Br EEBR); 3.36, 3.33 (s, 3H, OCH₃); 1.64-1.54 (m, 2H, OCH₂CH₂ Pr); 1.25-1.19 (m, 3H, OCH₂CH₃ EEBR &); 0.89 (t, J_{gem} = 7.49 Hz, CH₃ Pr).

Example 55

Methyl 2-O-benzyl-4-O-tert-butyloxycarbonylmethyl-6-O-methyl-3-O-propyl- α/β -D-glucopyranoside

- 5 166 mg (0.1 mmol) of the polymer are reacted with benzyl bromide according to the general working procedure for an alkylation. The TBDPS group is subsequently removed and the product is then alkylated with methyl iodide. After the 1-ethoxyethyl protective group has been removed, the product is reacted with tert-butyl bromoacetate. After removal of the
- 10 carbohydrate from the resin and filtration on silica gel, 23 mg of a colorless oil are obtained. $R_f = 0.64, 0.59$ (PE/EtOAc = 3:1 v/v). $C_{24}H_{38}O_8$ (454.6). FAB-MS (NBA-pos, LiCl): (m/e) = 461.3 (75%, $[M+Li]^+$, calc.: 461.3); 462.3 (23%, $[M+Li]^+$, ^{13}C , calc.: 462.3); 561.3 (100%, $[C_{29}H_{46}O_{10}+Li]^+$, 6-OCH₂CO₂tBu, calc.: 561.3); 562.3 (33%, $[C_{29}H_{46}O_{10}+Li]^+$, ^{13}C , calc.: 562.3).
- 15

Example 56

- Library containing 1,2,6-functionalized glucose derivatives (compound of the formula I where X is equal to O, R^3 is equal to propyl and R^4 is equal to
- 20 hydrogen)

80 mg of the loaded resin were functionalized according to the general working procedures (Table 1)

25 Example 57

- Library containing 1,2,4,6-functionalized glucose derivatives (compound of the formula I where X is equal to O, R^1 is equal to methyl, R^3 is equal to propyl)
- 30 80 mg of the loaded resin are functionalized according to the general working procedures. The alkylation in the 4-position takes place by use of tert-butyl bromoacetate.

Compound	R^2	R^4	R^5	MS analysis (m/e) (FAB, NBA+LiCl)
1	Bn	Bn	Me	437.1
2	Bn	CH ₂ CO ₂ tB	Me	461.3

Table 1

Compound	R ²	R ⁵	R ¹	MS analysis (m/e) (FAB, NBA+LiCl)
1	Me	Bn	Me	347.2
2	Me	Me	Bn	347.2
3	Me	Hep	Me	355.2
4	Me	Hep	iPr	383.2
5	Me	MNBn Et	Et	436.1
6	Bn	iPr	Me	375.1
7	Bn	Cbn	Et	462.2
8	Bn	Hep	Me	431.3
9	Bn	cHex	iPr	457.3
10	Bn	iBu	Me	389.2
11	Pr	Cbn	Me	400.2
12	Pr	Bn	iPr	403.2
13	Pr	Me	Bn	375.2
14	Pr	Hep	Me	383.3
15	Pent	iPr	Me	355.2
16	Pent	Hep	Et	425.3
17	Pent	Me	iPr	355.3
18	Hep	cHex	Bn	513.3
19	Hep	Bn	Me	431.3
20	Hep	Cbn	iPr	484.3
21	Hep	iPr	Me	383.3
22	Hep	BrBn	Me	509.2
23	Et	Bn	Me	361.3
24	Et	MNBn	Me	436.2
25	Et	Hep	Bn	445.3
26	Cbr	iPr	Me	400.3
27	Cbr	Hep	Et	470.3
28	Cbr	Me	iPr	400.2

Abbreviations: MNBn = 2-methoxy-5-nitrobenzyl, Cbn = o-cyanobenzyl,
Pent = pentyl, cHex = cyclohexylmethylene.

Example 58

Methyl S-(6'-O-tert-butyldiphenylsilyl-3',4'-O-isopropylidene- β -D-galactopyranosyl-2'-O-methyl)-4-mercaptobutyrate (32)

- 5 A solution of 3.0 g (5.2 mmol) of 19 in 30 ml of THF is stirred at 0°C for 45 min with 0.6 g (5.3 mmol) of potassium *tert*-butylate under an argon atmosphere and then treated with 0.38 ml (6.0 mmol) of methyl iodide. As only slight conversion is discernible after a few hours by thin-layer chromatographic checking, the same amount of reagents is added again.
- 10 The precipitating solid is dissolved by addition of 15 ml of DMF. After stirring for 12 h, the solution is concentrated in vacuo, and the residue is codistilled with toluene and purified by *flash* chromatography on silica gel (column 18 $\times$ 6 cm, eluent petroleum ether/ethyl acetate 10:1). Yield 1.8 g (59%), colorless oil, $[\alpha]_D^{25} = -15.9^\circ$ ($c = 1$, CHCl_3); $R_F = 0.34$
- 15 (petroleum ether/ethyl acetate 8:1).
 400MHz- ^1H -NMR (CDCl_3): δ [ppm] = 7.69-7.66; 7.41-7.33 (m, 10H, SiPh_2), 4.28 (d, 1H, $J_{1,2} = 9.7$ Hz, 1'-H), 4.27 (dd, 1H, $J_{4,3} = 5.5$ Hz, $J_{4,5} = 2.3$ Hz, 4'-H), 4.09 (dd, 1H, $J_{3,2} = 7.0$ Hz, $J_{3,4} = 5.6$ Hz, 3'-H), 3.89 (m, 2H, 6'-H_{a,b}), 3.79 (ddd, 1H, $J_{5,4} = 2.3$ Hz, $J_{5,6a} \approx J_{5,6b} = 6.2$ Hz, 5'-H), 3.59;
 20 3.55 (2s, 6H, OCH_3), 3.17 (dd, 1H, $J_{2,1} = 10.0$ Hz, $J_{2,3} = 6.8$ Hz, 2'-H), 2.77 (dt, 1H, $J_{\text{vic}} = 6.5$ Hz, $J_{\text{gem}} = 12.9$ Hz, SCH_a), 2.64 (dt, 1H, $J_{\text{vic}} = 6.5$ Hz, $J_{\text{gem}} = 12.9$ Hz, SCH_b), 2.39 (m_c, 2H, CH_2COOMe), 1.91 (m_c, 2H, SCH_2CH_2), 1.50; 1.33 (2s, 6H, $\text{C}(\text{CH}_3)_2$), 1.03 (s, 9H, $\text{SiC}(\text{CH}_3)_3$).
 100.6 MHz- ^{13}C -NMR (CDCl_3): δ [ppm] = 173.3 (CO), 135.6; 135.5 (C_p - SiPh_2), 133.5; 133.4 (C_e - SiPh_2), 129.7; 127.7; 127.6 ($\text{C}_{o,m}$ - SiPh_2), 109.7 ($\text{C}(\text{CH}_3)_2$), 83.6; 81.8; 79.4; 76.8; 73.4; 62.8 (C-1'-C-6'), 59.7 (OCH_3), 51.4 (COOCH_3), 32.7 (SCH_2), 29.6 (CH_2COOMe), 28.0 ($\text{C}(\text{CH}_3)_2$), 26.7 ($\text{SiC}(\text{CH}_3)_3$), 26.2 ($\text{C}(\text{CH}_3)_2$), 25.0 (SCH_2CH_2), 19.2 ($\text{SiC}(\text{CH}_3)_3$).
 $\text{C}_{31}\text{H}_{44}\text{O}_7\text{SSi}$ (588.8) Calc.: C 63.23 H 7.53 S 5.44
 30 Found: C 63.09 H 7.61 S 5.41

Example 59

S-(6'-O-tert-Butyldiphenylsilyl-3',4'-O-isopropylidene- β -D-galactopyranosyl)-4-mercaptobutyric acid, polymer-bonded (33, 34)

35

The thiogalactoside 19 is coupled to 0.390 g (0.6 mmol) of aminomethylpolystyrene to give 33 according to the general procedure.

Loading according to the sulfur content of the elemental analysis: 0.77-0.81 mmol/g.

The thiogalactoside 19 is coupled to 2.0 g (0.56 mmol) of Tentagel to give 34 according to the general procedure.

- 5 Loading (gravimetric): 0.20 mmol/g.

Example 60

S-(6'-O-tert-Butyldiphenylsilyl-3',4'-O-isopropylidene-2'-O- β -D-galactopyranosyl)-4-mercaptobutyric acid, polymer-bonded (35, 36)

10

The 2-O-methylthiogalactoside 32 is coupled to 1.0 g (0.6 mmol) of aminomethylpolystyrene to give 35 according to the general working procedure.

- 15 Loading according to the sulfur content of the elemental analysis:
0.78 mmol/g.

The 2-O-methylthiogalactoside 32 is coupled to 2.0 g (0.56 mmol) of Tentagel to give 36 according to the general working procedure.

Loading (gravimetric): 0.27 mmol/g.

- 20 Example 61

Methyl 6-O-tert-butyldiphenylsilyl-3,4-O-isopropylidene-2-O-methyl- α/β -D-galactopyranoside

- 25 a) by removal of 35: According to the general procedure, 400 mg (0.312 mmol) of 79 are treated with methanol as an alcohol. Purification is carried out by flash chromatography on silica gel (column 18 $\times$ 4 cm, eluent petroleum ether/ethyl acetate 8:1).

- 30 Yield 70 mg (24%) of α -anomer, colorless oil; 134 mg (46%) of β -anomer, colorless oil. By washing the resin again and combining the washings with the mixed fractions, a further 43 mg (15%) are obtained as an anomer mixture.

- 35 b) by removal of 36: According to the general procedure, 80 is glycosylated using methanol as an alcohol. Purification is carried out by flash chromatography on silica gel (column 18 $\times$ 4 cm, eluent petroleum ether/ethyl acetate 8:1).

Yield 12 mg (4%) of α -anomer, colorless oil; 44 mg (15%) of β -anomer, colorless oil.

In addition, 47 mg (17%) of hydrolysis sugar are isolated as an anomer mixture.

- c) by alkylation with sodium hydride/methyl iodide: 370 mg (0.3 mmol) of 33 are suspended in 15 ml of DMF/THF (1:1) and preliminarily shaken at room temp. under an argon atmosphere for 20 min with 0.090 g (3.0 mmol) of sodium hydride (80% in mineral oil). After addition of 0.19 ml (3.0 mmol) of methyl iodide, the mixture is shaken for 12 h, and the resin is filtered off after addition of 5 ml of methanol and washed a number of times with DMF. Removal is carried out according to the general procedure using methanol as an alcohol. Purification is carried out by flash chromatography on silica gel (column 15 × 3 cm, eluent petroleum ether/ethyl acetate 8:1). Yield 92 mg (32%) of α -anomer, clear oil; 43 mg (15%) of β -anomer, clear oil. In addition, 14 mg (5%) of α -anomer and 32 mg (11%) of β -anomer of the nonalkylated methyl glycoside are isolated.
- d) by alkylation with KOtBu/methyl iodide: as under c) at 0°C, 4 h. Yield 55 mg (19%), clear oil, anomer mixture.

Example 62

Methyl 6-O-tert-butyldiphenylsilyl-3,4-O-isopropylidene-2-O-benzyl- β -D-galactopyranoside

- a) by alkylation of 33 with sodium hydride/benzyl bromide: 768 mg (0.6 mmol) of 33 are reacted as above with 180 mg (6.0 mmol) of sodium hydride (80% in mineral oil) and 0.72 ml (6.0 mmol) of benzyl bromide at room temp. under an argon atmosphere for 12 h and removed. Purification is carried out by *flash* chromatography on silica gel (column 15 × 4 cm, eluent petroleum ether/ethyl acetate 15:1). Yield 67 mg (20%) of α -anomer, clear oil; 26 mg (8%) of β -anomer, turbid oil. In addition, 4 mg (2%) are obtained as an anomer mixture.
- b) by alkylation of 34 with sodium hydride/benzyl bromide: 2.5 g (0.5 mmol) of 34 are reacted as above with 900 mg (30.0 mmol) of sodium hydride (80% in mineral oil) and 3.6 ml (30.0 mmol) of benzyl bromide at room temp. under an argon atmosphere for 12 h and removed. Purification is carried out by *flash* chromatography on silica gel (column 18 × 3 cm, eluent petroleum ether/ethyl acetate 10:1). Yield 100 mg (37%) of anomer mixture, yellow oil.
- c) by alkylation with potassium tert-butoxide/benzyl bromide: 640 mg (0.5 mmol) of 33 are swollen in 20 ml of DMF. After addition of 0.56 g (5.0 mmol) of potassium *tert*-butoxide, the mixture is shaken under an

argon atmosphere for 20 min. 0.59 ml (5.0 mmol) of benzyl bromide and 0.22 g (0.6 mmol) of tetrabutylammonium iodide are then added and the mixture is shaken for 16 h. 20 ml of methanol are added, and the resin is filtered off and washed a number of times with DMF and absol. THF. The resin is dried in vacuo. Removal is carried out according to the general procedure with methanol as an alcohol. Purification is carried out by *flash* chromatography on silica gel (column 18 × 3 cm, eluent petroleum ether/ethyl acetate 8:1).

Yield 124 mg (44%) of α -anomer, clear oil; 22 mg (8%) of β -anomer, colorless oil.

d) by alkylation with phosphazene base P₄-t-Bu/benzyl bromide: 384 mg (0.33 mmol) of 33 are swollen in 10 ml of DMF and cooled to 0°C under an argon atmosphere. 1.32 ml (1.32 mmol, 1 M in *n*-hexane) of P₄-*tert*-Bu are added and the mixture is shaken. After 15 min, 0.59 ml (5.0 mmol) of benzyl bromide is added and the mixture is shaken at 0°C for 16 h. The resin is filtered off and washed a number of times with DMF and absol. THF. After drying in vacuo, the compound is removed from the support according to the general procedure using methanol as an alcohol. Purification is carried out by *flash* chromatography on silica gel (column 18 × 3 cm, eluent petroleum ether/ethyl acetate 10:1).

Yield 163 mg (85%) of anomer mixture (α : β about 5:1), colorless oil.

α -Anomer:

$[\alpha]_D^{25} = +44.9$ (c = 1, CHCl₃); R_F = 0.20 (petroleum ether/ethyl acetate 8:1)

β -Anomer:

$[\alpha]_D^{25} = +17.2$ (c = 1, CHCl₃); R_F = 0.30 (petroleum ether/ethyl acetate 8:1).

Example 63

Methyl 2-O-(*o*-cyanobenzyl)-6-O-*tert*-butyldiphenylsilyl-3,4-O-isopropylidene- α / β -D-galactopyranoside

400 mg (0.312 mmol) of 33 are reacted with *o*-cyanobenzyl bromide according to the general working procedure (variant A) and removed with methanol under glycosylating conditions. The crude product is purified by flash chromatography on silica gel (column 20 × 2 cm, eluent petroleum ether/ethyl acetate 12:1).

α -Anomer:

Yield 26 mg (14%), colorless oil, $[\alpha]_D^{25} = +50.5$ ($c = 1.0$, CHCl_3); $R_F = 0.39$ (petroleum ether/ethyl acetate 4:1), HPLC (column C, gradient 1): 20.91 min.

5 β -Anomer:

Yield 12 mg (7%), colorless oil, $[\alpha]_D^{25} = +19.6$ ($c = 0.33$, CHCl_3); $R_F = 0.46$ (petroleum ether/ethyl acetate 4:1), HPLC (column C, gradient 1): 20.91 min.

10 Example 64

Methyl 2-O-(p-bromobenzyl)-6-O-tert-butyldiphenylsilyl-3,4-O-isopropylidene- α/β -D-galactopyranoside

15 400 mg (0.312 mmol) of 33 are reacted with *p*-bromobenzyl bromide according to the general working procedure and the product is removed under glycosylating conditions using methanol. The crude product is purified by *flash* chromatography on silica gel (column 20 $\times$ 2 cm, eluent petroleum ether/ethyl acetate 12:1).

α -Anomer:

20 Yield 12 mg (6%), colorless oil, $[\alpha]_D^{25} = +38.4$ ($c = 1.0$, CHCl_3); $R_F = 0.17$ (petroleum ether/ethyl acetate 12:1), HPLC (column C, gradient 1): 22.13 min.

β -Anomer:

25 Yield 3 mg (2%), colorless oil, $[\alpha]_D^{25} = +39.3$ ($c = 0.1$, CHCl_3); $R_F = 0.23$ (petroleum ether/ethyl acetate 12:1), HPLC (column C, gradient 1): 22.13 min.

Example 65

30 Methyl 2-O-ethyl-6-O-tert-butyldiphenylsilyl-3,4-O-isopropylidene- α/β -D-galactopyranoside

35 400 mg (0.312 mmol) of 33 are reacted with ethyl iodide according to the general working procedure and the product is removed under glycosylating conditions using methanol. The crude product is purified by *flash* chromatography on silica gel (column 15 $\times$ 2.5 cm, eluent petroleum ether/ethyl acetate 12:1).

α -Anomer:

Yield 22 mg (14%), colorless oil, $[\alpha]_D^{25} = +73.2$ ($c = 1$, CHCl_3); $R_F = 0.49$ (petroleum ether/ethyl acetate 4:1), HPLC (column C, gradient 1): 19.62 min.

5 β -Anomer:

Yield 10 mg (7%), colorless oil, $[\alpha]_D^{25} = +1.1$ ($c = 0.33$, CHCl_3); $R_F = 0.49$ (petroleum ether/ethyl acetate 4:1), HPLC (column C, gradient 1): 19.62 min.

10 Example 66

S-(2'-O-Benzyl-3',4'-O-isopropylidene- α/β -D-galactopyranosyl)-4-mercaptobutyric acid, polymer-bonded

15 1 g (0.78 mmol) of 33 is alkylated with benzyl bromide according to the general procedure (variant A) and the silyl protective group is removed. The polymer is dried in vacuo.

Example 67

20 Methyl 2-O-benzyl-6-O-(2'-naphthylmethyl)-3,4-O-isopropyliden- α/β -D-galactopyranoside

400 mg (0.312 mmol) of S-(2'-O-benzyl-3',4'-O-isopropylidene- α/β -D-galactopyranosyl)-4-mercaptobutyric acid - polymer bonded are reacted with 2-bromomethylnaphthalene according to the general working
25 procedures and the product is removed under glycosylating conditions using methanol. The pure product is obtained by *flash* chromatography on silica gel (column 20 $\times$ 2 cm, eluent petroleum ether/ethyl acetate 4:1) as an anomer mixture in the ratio $\alpha:\beta = 10:1$.

Yield 40 mg (38%), yellow oil, $[\alpha]_D^{25} = +79.2$ ($c = 1$, CHCl_3); $R_F = 0.25$
30 (petroleum ether/ethyl acetate 4:1), HPLC (column C, gradient 1): 15.28 min.

Example 68

35 Methyl 2-O-propyl-6-O-tert-butyldiphenylsilyl-3,4-O-isopropylidene- α/β -D-galactopyranoside

400 mg (0.312 mmol) of 33 are reacted with propyl iodide according to the general working procedure and the product is removed under glycosylating

conditions using methanol. The crude product is purified by *flash* chromatography on silica gel (column 17 × 2.5 cm, eluent petroleum ether/ethyl acetate 12:1).

α -Anomer:

- 5 Yield 7 mg (5%), colorless oil, $[\alpha]_D^{25} = +56.9$ (c = 0.2, CHCl₃); R_F = 0.51 (petroleum ether/ethyl acetate 4:1), HPLC (column C, gradient 1): 21.03 min.

β -Anomer:

- Yield 2 mg (2%), colorless oil, $[\alpha]_D^{25} = +2.7$ (c = 0.1, CHCl₃); R_F = 0.57
10 (petroleum ether/ethyl acetate 4:1), HPLC (column C, gradient 1): 19.62 min.

Example 69

- Methyl 2-O-heptyl-6-O-tert-butyldiphenylsilyl-3,4-O-isopropylidene- α/β -D-
15 galactopyranoside

- 400 mg (0.312 mmol) of 33 are reacted with heptyl iodide with addition of 18-crown-6 according to the general working procedure (variant A) and the product is removed under glycosylating conditions using methanol. The
20 crude product is purified by *flash* chromatography on silica gel (column 20 × 2 cm, eluent petroleum ether/ethyl acetate 15:1).

α -Anomer:

- Yield 12 mg (7%), colorless oil, $[\alpha]_D^{25} = +52.6$ (c = 0.6, CHCl₃); R_F = 0.34
25 (petroleum ether/ethyl acetate 10:1), HPLC (column C, gradient 1): 24.27 min.

β -Anomer:

- Yield 8 mg (5%), colorless oil, $[\alpha]_D^{25} = +14.6$ (c = 0.4, CHCl₃); R_F = 0.40
30 (petroleum ether/ethyl acetate 10:1), HPLC (column C (C8), gradient 1): 24.27 min.

Example 70

Ethyl 3,4-O-isopropylidene-1-thio- β -D-galactopyranoside

- A mixture of 19.1 g (85.4 mmol) of ethyl 1-thio- β -D-galactopyranoside,
35 360 ml (2.93 mol) of acetone dimethyl acetal and 0.3 g (16 mmol) of p-toluenesulfonic acid monohydrate is stirred at room temp. After 18 h, 1.2 ml of triethylamine are added and the mixture is concentrated to dryness in vacuo. The residue is suspended in 180 ml of dichloromethane

and treated with 2.4 ml of 50% strength trifluoroacetic acid. After 15 min, 3.6 ml of triethylamine are added and the mixture is concentrated in vacuo. The oil obtained is purified by *flash* chromatography on silica gel (column 30 × 10 cm, eluent petroleum ether/ethyl acetate 2:3).

- 5 Yield 15.7 g (70%), colorless crystals, melting point 89°C, $[\alpha] = +16.5^\circ$ (c = 1, CHCl₃); $R_F = 0.33$ (petroleum ether/ethyl acetate 1:9).

Example 71

10 Ethyl 6-O-tert-butyldiphenylsilyl-3,4-O-isopropylidene-1-thio-β-D-galactopyranoside

- A solution of 9.00 g (34 mmol) of Example compound 70 and 4.62 g (68 mmol) of imidazole in 100 ml of absol. DMF is treated with 10.9 ml (43 mmol) of *tert*-butyldiphenylsilyl chloride and stirred at room temp. for 15 3 h. The reaction is terminated by addition of 50 ml of water. 150 ml of dichloromethane are added and the organic phase is washed three times with 50 ml of water each time. The combined aqueous phases are extracted with 100 ml of dichloromethane, the combined organic phases are dried over magnesium sulfate and the solvent is removed in vacuo. 20 The oil obtained is chromatographed on silica gel (column 20 × 5 cm, eluent petroleum ether/ethyl acetate 4:1). Yield 12.66 g (74%), colorless crystals, $[\alpha]_D^{25} = +0.7^\circ$ (c = 1, CHCl₃); $R_F = 0.28$ (petroleum ether/ethyl acetate 4:1).

25 Example 72

Ethyl 2-O-acetyl-3,4-O-isopropylidene-6-O-tert-butyldiphenylsilyl-1-thio-β-D-galactopyranoside

- 50 ml (520 mmol) of acetyl chloride are added dropwise with ice-cooling to a solution of 12.6 g (25 mmol) of Example compound 71 in 100 ml of absol. pyridine. After 18 h, the solvent is distilled off in vacuo, the residue is taken up in 150 ml of dichloromethane, washed successively with 50 ml each of 0.5 N hydrochloric acid, satd sodium hydrogencarbonate solution and satd sodium chloride solution and the organic phase is dried over 35 magnesium sulfate. After concentrating in vacuo, the residue is separated from impurities by chromatography on silica gel (column 25 × 5 cm, eluent petroleum ether/ethyl acetate 4:1). The oily product solidifies after several days.

Yield 13.6 g (83%), colorless crystals, melting point 76°C, $[\alpha]_D^{25} = +15.6^\circ$ (c = 1, CHCl₃); R_F = 0.62 (petroleum ether/ethyl acetate 4:1).

Example 73

5 Ethyl 2-O-acetyl-6-O-tert-butyldiphenylsilyl-1-thio-β-D-galactopyranoside

a) by deacetalization with acetic acid: A solution of 140 mg (0.26 mmol) of Example compound 72 in 20 ml of 60% strength acetic acid is warmed to 60°C for 2 h with stirring. The mixture is concentrated in
10 vacuo and codistilled with 10 ml of toluene. The residue is taken up in 20 ml of dichloromethane and washed with 10 ml of satd sodium hydrogencarbonate solution. The crude product is purified by chromatography on silica gel (column 15 × 2 cm, eluent petroleum ether/ethyl acetate 2:3).

15 Yield 59 mg (45%), colorless oil; the analytical data agree with those indicated under b).

b) by deacetalization with p-TsOH: 11.36 g (21 mmol) of Example compound 72 are dissolved in 200 ml of chloroform, treated with 0.19 g
20 (1 mmol) of p-toluenesulfonic acid monohydrate and 12.27 ml (146 mmol) of ethanedithiol and the mixture is heated under reflux. After 45 min, it is allowed to cool to room temp. and washed with 50 ml each of satd sodium hydrogencarbonate solution, 0.5 N hydrochloric acid and water. The organic phase is dried over magnesium sulfate and then freed from the
25 solvent. Chromatographic purification (column 30 × 5 cm, eluent petroleum ether/ethyl acetate 2:3) of the crude product yields the title compound after drying in vacuo.

Yield 8.44 g (80%), colorless oil solidifying after days, $[\alpha]_D^{25} = +9.4^\circ$ (c = 1, CHCl₃); R_F = 0.39 (petroleum ether/ethyl acetate 1:1).

30

Example 74

Ethyl 2-O-acetyl-3-O-allyl-6-O-tert-butyldiphenylsilyl-1-thio-β-D-galactopyranoside (28)

35 A solution of 8.33 g (165 mmol) of Example compound 73 and 4.31 g (173 mmol) of dibutyltin oxide in 150 ml of benzene is heated under reflux in a water separator for 16 h. Half of the solvent is then removed by distillation and 2.42 ml (286 mmol) of allyl bromide are added. The mixture is then stirred at 50°C for 5 h. It is then concentrated in vacuo, the residue

is taken up in 150 ml of dichloromethane, and the organic phase is washed three times with 50 ml of water each time and dried over magnesium sulfate. The solvent is concentrated in vacuo and the crude product is purified by chromatography on silica gel (column 25 × 8 cm, eluent petroleum ether/ethyl acetate 4:1).
 5 Yield 7.34 g (82%), yellowish oil, $[\alpha]_D^{25} = +2.4^\circ$ (c = 1, CHCl₃); R_F = 0.35 (petroleum ether/ethyl acetate 4:1).

Example 75

10 Ethyl 2-O-acetyl-3-O-allyl-1-thio- β -D-galactopyranoside (29)

2.5 ml of a 1 M solution of tetrabutylammonium fluoride in THF are added dropwise to a solution of 5.2 g (9.5 mmol) of 28 in 200 ml of THF and the mixture is stirred at room temp. for 1 h. It is then diluted with 500 ml of
 15 dichloromethane, washed with 100 ml of water and the organic phase is dried over magnesium sulfate. After concentrating, the crude product is purified by chromatography on silica gel (column 20 × 5 cm, eluent petroleum ether/ethyl acetate 2:3).
 Yield 2.29 g (79%), yellowish oil, $[\alpha]_D^{25} = -20.6^\circ$ (c = 1, CHCl₃); R_F = 0.35
 20 (petroleum ether/ethyl acetate 4:1).

Example 76

Ethyl 2-O-acetyl-3-O-allyl-6-O-*p*-(4'-methoxycarbonylbutyloxy)phenyl-1-thio- β -D-galactopyranoside (30)

25 A solution of 2.0 ml (13.5 mmol) of diethyl diazodicarboxylate in 5 ml of dichloromethane is added dropwise with stirring to a solution of 2.25 g (9.5 mmol) of 29, 4.7 g (22.5 mmol) of methyl *p*-hydroxyphenoxybutyrate and 6.0 g (22.5 mmol) of triphenylphosphine in 70 ml of dichloromethane.
 30 After 4 h, the mixture is concentrated without further working up and the crude product is purified by chromatography on silica gel (column 20 × 4 cm, eluent petroleum ether/ethyl acetate 3:1).
 Yield 1.92 g (53%), colorless oil, $[\alpha]_D^{25} = +1.5^\circ$ (c = 1, CHCl₃); R_F = 0.52 (petroleum ether/ethyl acetate 1:1).

Example 77

Ethyl 2-O-acetyl-3-O-allyl-6-O-*p*-(4'-methoxycarbonylbutyloxy)phenyl-4-O- β -(trimethylsilyl)ethoxymethyl-1-thio- β -D-galactopyranoside (31)

- 5 2.12 ml (12 mmol) of *N,N*-diisopropylethylamine and 1.59 ml (9 mmol) of β -(trimethylsilyl)ethoxymethyl chloride are added to a solution of 1.51 g (3.02 mmol) of 30 in 20 ml of absol. dichloromethane. The reaction mixture is heated under reflux for 6 h under an argon atmosphere. The reaction is then terminated by addition of 10 ml of methanol and the mixture is
- 10 concentrated in vacuo. Impurities are separated by means of chromatography on silica gel (column 15 $\times$ 2 cm, eluent petroleum ether/ethyl acetate 3:1).

Yield 1.8 g (90%), colorless oil, $[\alpha]_D^{25} = -27.0^\circ$ ($c = 1$, CHCl_3); $R_F = 0.40$ (petroleum ether/ethyl acetate 3:1).

15

Example 78

Preparation of libraries of the type alkyl 3,4-O-isopropylidene-2-O-alkyl-6-O-carbamoylgalactoside

- 20 a) General procedure for the coupling of the methyl galactosylmercaptobutyrate to amino-functionalized polymeric supports

- 1.9 g (30 mmol) of lithium hydroxide are added to a solution of 14.6 mmol
- 25 of the thiogalactoside in 400 ml of THF/methanol/water (2:2:1) and the mixture is stirred at room temp. for 8 h. It is then neutralized (0.5 N hydrochloric acid or phosphate buffer solution), 400 ml of satd sodium chloride solution are added and the mixture is extracted three times with 200 ml of ethyl acetate. The extracts are dried over magnesium sulfate and
- 30 freed from the solvent in vacuo. The residue is taken up in 150 ml of dichloromethane or DMF and the solution obtained is shaken overnight in a solid-phase reactor with 19.8 mmol of the amino-functionalized polymer concerned, 3.7 ml (14.6 mmol) of *N,N'*-diisopropylcarbodiimide and 3.86 g (14.6 mmol) of *N*-hydroxybenzotriazole. The polymer is then filtered off with
- 35 suction and washed ten times with 50 ml each of DMF and dichloromethane. The loaded polymer is dried in vacuo and the loading with carbohydrate matrix is determined by means of elemental analysis.

b) General working procedure for the capping of the polymers

A mixture of 1.56 mmol of the polymer concerned is shaken for 18 h with a solution of 0.45 ml (7.80 mmol) of acetic acid, 2.7 ml (15.60 mmol) of *N,N*-diisopropylethylamine and 3.74 g (7.80 mmol) of PfPyU in 40 ml of DMF. After filtering off the liquid phase with suction, the polymer is washed thoroughly with DMF, dichloromethane and diethyl ether and dried in vacuo.

10 c) General working procedure for the alkylation of capped polymers

0.078 mmol of the polymer concerned is shaken with a solution of 88 mg (0.78 mmol) of potassium *tert*-butoxide in 2 ml of DMF. After 15 min, the polymer is filtered off and the resin is immediately [lacuna] with a solution of 41 mg (0.16 mmol) of 18-crown-6 and 0.78 mmol of the alkyl halide concerned in 2 ml of DMF. The mixture is shaken for 3 h and the polymer is then filtered off with suction. The resin is washed thoroughly with DMF, methanol, dichloromethane and diethyl ether and dried in vacuo.

20 d) General procedure for the removal of the *tert*-butyldiphenylsilyl protective group on the polymeric support.

0.56 ml (0.56 mmol, 1 M) tetrabutylammonium fluoride in THF is added to a suspension of 0.056 mmol of loaded polymer in 1.5 ml of THF. The mixture is shaken for 4 h. The resin is filtered off from the solution and washed five times with 2 ml each of DMF and dichloromethane.

e) General working procedure for carbamoylations

Variant A (using 1,1'-carbonyldiimidazole and amines)

30

A solution of 0.25 g (1.56 mmol) of 1,1'-carbonyldiimidazole in 2 ml of DMF or dioxane is added to 0.078 mmol of polymer with 0.1 ml (0.58 mmol) of *N,N*-diisopropylethylamine, a spatula-tipful of DMAP and a spatula-tipful of KO^tBu. The mixture is shaken for 2 h and filtered. The polymer is then shaken overnight with a solution of 1.56 mmol of the appropriate amine in 2 ml of DMF. It is filtered and washed five times each with DMF and dichloromethane.

Variant B (using isocyanates)

A solution of 1.56 mmol of the appropriate isocyanate in 2 ml of dioxane is added to 0.078 mmol of polymer. A spatula-tipful of DMAP is then added
5 and the mixture is shaken for 3-7 h, depending on the reactivity of the OH group to be reacted. It is filtered and the polymer is washed thoroughly with dioxane, methanol and dichloromethane.


f) General procedure for the removal of the galactose derivatives from
10 the polymeric support

A suspension of 0.056 mmol of polymer-bonded galactose derivative in 1.5 ml of abs. dichloromethane is shaken at room temp. in a 5 ml PE syringe (PE frit, plastic cap) with 0.3 ml of a 3.5% strength (0.36 ml of
15 bromine to 10 ml of dichloromethane or 5-10 eq. of bromine) solution of bromine in abs. dichloromethane and 0.08 ml (0.36 mmol) of 2,6-di-*tert*-butylpyridine or a corresponding amount of polymer-bonded 2,6-di-*tert*-butylpyridine. After 15 min, 0.2 ml of cyclohexene, 0.2 ml of the abs. alcohol to be glycosylated and 25 mg (0.056 mmol) of tetraethylammonium
20 bromide are added. After 4 h, the resin is filtered off and washed five times with 1 ml of dichloromethane. The combined filtrates are freed from the solvent in vacuo. The crude product obtained is applied to a silica gel cartridge in a little dichloromethane. The cartridge is eluted first with 30 ml of petroleum ether. This fraction is discarded. The product is obtained by
25 eluting with suitable petroleum ether/ethyl acetate mixtures. If the mixture is distributed in various alcohols, it is advantageously additionally shaken with bromine solution and 2,6-di-*tert*-butylpyridine. The solution is injected into a vessel from the syringe and washed 5× with abs. dichloromethane. This resulting mixture is distributed in various vessels which contain the
30 ammonium bromide, cyclohexene and the respective alcohol. The vessels are sealed and shaken. Standing for too long in the air is to be avoided in the case of the bromine solutions, as these are moisture-sensitive.

Parallel synthesis


35 6 × 3.1 g (2.42 mmol) of 33 are treated with PfPyU and acetic acid according to the general procedure. The resins thus obtained are alkylated with methyl iodide, ethyl iodide, heptyl iodide, benzyl bromide, *p*-bromobenzyl bromide and *o*-cyanobenzyl bromide according to the general procedure. The silyl protective group is then removed according to the

general procedure. 150 mg each (about 0.119 mmol) of the resins obtained are then reacted either with diethylamine, benzylamine, methoxyethylamine, cyclopropylmethylamine or glycine *tert*-butyl ester according to the general procedure or with *p*-chlorophenyl isocyanate, 5 *o*-trifluoromethylphenyl isocyanate, *o*-nitrophenyl isocyanate, *p*-cyanophenyl isocyanate, *m,p*-dichlorophenyl isocyanate, *m*-fluorophenyl isocyanate, phenyl isocyanate, *n*-propyl isocyanate, *tert*-butyl isocyanate or ethyloxycarbonylmethyl isocyanate according to the general procedure. The polymers are swollen in 2 ml of dichloromethane and shaken for 10 20 min with 0.9 ml in each case of a 3.5% strength bromine solution in dichloromethane and 0.24 ml (1.08 mmol) of 2,6-di-*tert*-butylpyridine. The polymers are filtered off and washed a number of times with dichloromethane. The solutions obtained are divided into three and shaken for 5 h each with 55 mg (0.26 mmol) of tetraethylammonium bromide, 15 0.2 ml of cyclohexene and 0.2 ml of methanol, ethanol or *i*-propanol. The solutions are concentrated in vacuo. The residues are taken up in 250 μ l of dichloromethane each and applied to silica gel cartridges. They are washed with 30 ml each of petroleum ether and the eluate is discarded. They are then eluted with 5 ml each of petroleum ether/ethyl acetate (1:1) 20 and the product obtained is concentrated in vacuo. By activation with CDI (variant A), the following compounds of the formula (A) are obtained:



	R ²	R ⁶	R ⁷	R ¹	Yield	R _T ^{a)}	R _F ^{b)}
1	4-BrBn	CH ₃ O(CH ₂) ₂	H	Me	2 mg (10 %)	18.41	0.60
2	4-BrBn	<i>c</i> -PrCH ₂	H	Me	3 mg (15 %)	17.69	0.58
3	4-BrBn	<i>t</i> -BuOOCCH ₂	H	Me	9 mg (41 %)	18.28, 18.59	0.59
4	2-CNBN	<i>c</i> -PrCH ₂	H	Me	4 mg (23 %)	15.81	0.28
5	2-CNBN	<i>t</i> -BuOOCCH ₂	H	Me	4 mg (20 %)	17.35	0.58
6	Bn	Et	Et	Et	11 mg (64 %)	18.21, 18.48	0.62, 0.68
7	Bn	<i>c</i> -PrCH ₂	H	Et	6mg (35%)	16.94	0.55
8	Bn	Bn	H	Et	10 mg (54 %)	17.69, 18.03	0.60
9	Bn	<i>t</i> -BuOOCCH ₂	H	Et	6 mg (31 %)	17.60, 17.91	0.57
10	4-BrBn	Et	Et	Et	5 mg (25 %)	19.29, 19.72	0.75
11	4-BrBn	CH ₃ O(CH ₂) ₂	H	Et	4 mg (19 %)	19.05	0.36
12	4-BrBn	<i>c</i> -PrCH ₂	H	Et	4 mg (20 %)	18.04	0.70
13	4-BrBn	<i>t</i> -BuOOCCH ₂	H	Et	6 mg (30 %)	18.97, 19.33	0.72
14	2-CNBN	Bn	H	Et	5 mg (26 %)	17.13, 17.34	0.60
15	2-CNBN	CH ₃ O(CH ₂) ₂	H	Et	2 mg (11 %)	13.86	0.21
16	2-CNBN	<i>c</i> -PrCH ₂	H	Et	5 mg (28 %)	16.30, 16.53	0.59
17	2-CNBN	<i>t</i> -BuOOCCH ₂	H	Et	5 mg (25 %)	17.04, 17.23	0.66
18	Bn	Et	Et	<i>i</i> -Pr	9 mg (51 %)	17.79, 17.96	0.74
19	Bn	Bn	H	<i>i</i> -Pr	10 mg (53 %)	18.36, 18.65	0.72
20	Bn	CH ₃ O(CH ₂) ₂	H	<i>i</i> -Pr	3 mg (17 %)	15.84, 16.21	0.43
21	Bn	<i>c</i> -PrCH ₂	H	<i>i</i> -Pr	10 mg (57 %)	17.63, 17.94	0.70
22	Bn	<i>t</i> -BuOOCCH ₂	H	<i>i</i> -Pr	8 mg (40 %)	18.22, 18.51	0.74
23	4-BrBn	Et	Et	<i>i</i> -Pr	3 mg (15 %)	20.48, 20.68	0.48
24	4-BrBn	CH ₃ O(CH ₂) ₂	H	<i>i</i> -Pr	5 mg (24 %)	19.26, 19.70	0.11
25	4-BrBn	<i>c</i> -PrCH ₂	H	<i>i</i> -Pr	3 mg (15 %)	19.10	0.49
26	4-BrBn	<i>t</i> -BuOOCCH ₂	H	<i>i</i> -Pr	12 mg (52 %)	19.26, 19.50	0.50
27	2-CNBN	Bn	H	<i>i</i> -Pr	6 mg (30 %)	17.79, 17.94	0.66
28	2-CNBN	<i>c</i> -PrCH ₂	H	<i>i</i> -Pr	5 mg (27 %)	17.03, 17.18	0.64
29	2-CNBN	<i>t</i> -BuOOCCH ₂	H	<i>i</i> -Pr	5 mg (24 %)	16.23	0.66

By activation with isocyanates (variant B), the following compounds of the formula B are obtained



B

5

	R ⁷	R ²	Yield	R _T ^{a)}	R _F ^{b)}
1	4-ClPh	Me	5 mg	18.75	0.20; 0.16
2	2-CF ₃ Ph	Me	5 mg (30 %)	18.94	0.31; 0.26
3	2-NO ₂ Ph	Me	5 mg (31 %)	18.80	0.30; 0.26
4	4-CN-Ph	Me	3 mg (20 %)	13.93	0.06
5	3,4-Cl ₂ Ph	Me	4 mg (26 %)	19.35	0.20; 0.14
6	4-ClPh	Me	5 mg (31 %)	16.53	0.34
7	2-CF ₃ Ph	Me	5 mg (29 %)	17.63	0.46
8	2-NO ₂ Ph	Me	9 mg (54 %)	16.36	0.46
9	4-CN-Ph	Me	4 mg (25 %)	14.75	0.13
10	3,4-Cl ₂ Ph	Me	5 mg (29 %)	17.84	0.29
11	4-ClPh	Me	9 mg (48 %)	21.35	0.63
12	2-CF ₃ Ph	Me	14 mg (69 %)	21.27	0.61
13	2-NO ₂ Ph	Me	11 mg (57 %)	21.08	0.70
14	4-CN-Ph	Me	5 mg (32 %)	20.04	0.36
15	3,4Cl ₂ Ph	Me	7 mg (35 %)	23.38	0.63
16	4-ClPh	Me	5 mg (23 %)	19.90;20.05	0.30
17	2-CF ₃ Ph	Me	10 mg (43 %)	19.49	0.40
18	2-NO ₂ Ph	Me	9 mg (41 %)	19.78;19.88	0.39
19	4-CN-Ph	Me	3 mg (14 %)	18.53;18.64	0.13
20	3,4Cl ₂ Ph	Me	9 mg (39 %)	20.65;20.95	0.34; 0.26
21	4-ClPh	Me	4 mg	16.12;16.92	0.29; 0.21
22	2-CF ₃ Ph	Me	7 mg (34 %)	20.72	0.33

81

23	2-NO ₂ Ph	Me	6 mg (30 %)	17.83	0.31
24	4-CN-Ph	Me	3 mg	14.46	0.10
25	3,4-Cl ₂ Ph	Me	4 mg (19 %)	19.26;19.69	0.24
26	4-ClPh	Et	4 mg (25 %)	18.98	0.29
27	2-CF ₃ Ph	Et	5 mg (29 %)	15.93;16.09	0.42
28	2-NO ₂ Ph	Et	4 mg (24 %)	14.49	0.40
29	4-CN-Ph	Et	3 mg	12.21	0.15
30	3,4-Cl ₂ Ph	Et	4 mg (23 %)	19.69;19.90	0.33
31	4-ClPh	Et	4 mg (24 %)	17.26	0.54
32	2-CF ₃ Ph	Et	8 mg (44 %)	16.79	0.38
33	2-NO ₂ Ph	Et	7 mg (41 %)	17.24	0.55
34	4-CN-Ph	Et	4 mg (24 %)	15.55	0.27
35	3,4-Cl ₂ Ph	Et	4 mg (22 %)	18.63	0.46
36	4-ClPh	Et	9 mg (46 %)	21.91;22.07	0.75
37	2-CF ₃ Ph	Et	11 mg (53 %)	20.99;21.16	0.71; 0.64
38	2-NO ₂ Ph	Et	9 mg (45 %)	22.14	0.80
39	4-CN-Ph	Et	6 mg (31 %)	20.64;20.79	0.57
40	3,4-Cl ₂ Ph	Et	8 mg (38 %)	22.28	0.77
41	4-ClPh	Et	4 mg (18 %)	18.98	0.46
42	2-CF ₃ Ph	Et	10 mg (42 %)	19.89;20.18	0.55
43	2-NO ₂ Ph	Et	6 mg (27 %)	20.59;20.74	0.55
44	4-CN-Ph	Et	4 mg (18 %)	18.73;19.31	0.25
45	3,4-Cl ₂ Ph	Et	8 mg (34 %)	19.39;19.53	0.45
46	4-ClPh	Et	6 mg (30 %)	16.11	0.35
47	2-CF ₃ Ph	Et	6 mg (29 %)	20.68;21.0 ^{e)}	0.45; 0.36
48	2-NO ₂ Ph	Et	6 mg (29 %)	18.61	0.41
50	3,4-Cl ₂ Ph	Et	3 mg (14 %)	19.68;19.82	0.33
51	4-ClPh	<i>i</i> -Pr	4 mg (24 %)	17.12;18.89	0.35
52	2-CF ₃ Ph	<i>i</i> -Pr	11 mg (61 %)	21.66	0.48
53	2-NO ₂ Ph	<i>i</i> -Pr	8 mg (47 %)	17.16	0.45
55	3,4-Cl ₂ Ph	<i>i</i> -Pr	4 mg (22 %)	19.66	0.35
56	4-ClPh	<i>i</i> -Pr	6 mg (35 %)	17.90;18.07	0.66
57	2-CF ₃ Ph	<i>i</i> -Pr	9 mg (48 %)	17.60;17.84	0.55
58	2-NO ₂ Ph	<i>i</i> -Pr	8 mg (45 %)	17.22;18.06	0.66
59	4-CN-Ph	<i>i</i> -Pr	4 mg (24 %)	16.31	0.38
60	3,4-Cl ₂ Ph	<i>i</i> -Pr	4 mg (22 %)	16.35	0.54
61	4-ClPh	<i>i</i> -Pr	6 mg (30 %)	22.47	0.77

82

62	2-CF ₃ Ph	<i>i</i> -Pr	10 mg (47 %)	22.15;22.39	0.71; 0.64
63	2-NO ₂ Ph	<i>i</i> -Pr	9 mg (44 %)	22.61;22.92	0.79
64	4-CN-Ph	<i>i</i> -Pr	5 mg (25 %)	21.25	0.59
65	3,4-Cl ₂ Ph	<i>i</i> -Pr	7 mg (36 %)	23.47	0.80
66	4-ClPh	<i>i</i> -Pr	7 mg (31 %)	21.04	0.52
67	2-CF ₃ Ph	<i>i</i> -Pr	10 mg (42 %)	23.71 ^{a)}	0.62
68	2-NO ₂ Ph	<i>i</i> -Pr	8 mg (34 %)	21.30	0.57
69	4-CN-Ph	<i>i</i> -Pr	7 mg (31 %)	22.43;22.5 ^{a)}	0.43; 0.33
70	3,4-Cl ₂ Ph	<i>i</i> -Pr	10 mg (41 %)	21.72	0.60; 0.52
72	2-CF ₃ Ph	<i>i</i> -Pr	6 mg (27 %)	23.78 ^{a)}	0.47; 0.41
73	2-NO ₂ Ph	<i>i</i> -Pr	6 mg (28 %)	22.16 ^{a)}	0.54; 0.50
74	4-CN-Ph	<i>i</i> -Pr	3 mg (15 %)	16.78 ^{a)}	0.29
75	3,4-Cl ₂ Ph	<i>i</i> -Pr	7 mg (32 %)	23.13 ^{a)}	0.48; 0.42
76	4-Br-Ph	Me	4 mg (19 %)	15.29;15.73	0.27
77	Pr	Me	5 mg (30 %)	17.27	0.16
79	3-F-Ph	Me	4 mg (21 %)	16.92;17.06	0.25
80	EtOOCCH ₂	Me	4 mg (22 %)	17.28	0.09
81	4-Br-Ph	Et	4 mg (18 %)	18.98	0.45
82	Pr	Et	5 mg (29 %)	16.02;19.41	0.25
83	<i>t</i> -Bu	Et	4 mg (22 %)	17.72	0.46
84	3-F-Ph	Et	4 mg (21 %)	19.35	0.48
85	EtOOCCH ₂	Et	4 mg (21 %)	17.99	0.21
86	4-Br-Ph	<i>i</i> -Pr	3 mg (13 %)	19.44	0.48
87	Pr	<i>i</i> -Pr	5 mg (28 %)	16.84	0.32
88	<i>t</i> -Bu	<i>i</i> -Pr	3 mg (16 %)	13.65	0.48
89	3-F-Ph	<i>i</i> -Pr	3 mg (15 %)	19.90	0.55; 0.48
90	EtOOCCH ₂	<i>i</i> -Pr	2 mg (10 %)	19.16 ^{a)}	0.50

a) Column B, grad. 2; b) Petroleum ether/ethyl acetate (2:1); e) Column B, grad. 3


5 Column B: Beckmann unit (Gold System), Nucleosil 100-5 C18 column measured at 220 and 254 nm

Grad. 2 = CH₃CN:H₂O 10:90 - 90-10, flow 1 ml/min, 0.1% TFA



Grad. 3 = CH₃CN:H₂O 10:90 - 90-10, flow 0.65 ml/min, 0.1% TFA

Example 79

Functionalization of the 1-, 2-, 4- and 6-position



1. a) KOtBu, DMF
b) R²X, 18-crown-6, DMF
2. TBAF, THF
3. Alkylation or carbamoylation (R⁷)
4. PPTS, dioxane/MeOH (10:1)
5. R⁶NCO, DMAP, CH₂Cl₂
6. a) Br₂, DTBP, CH₂Cl₂
b) R¹OH, *c*-hexane, TEAB, CH₂Cl₂



5

The following compounds are obtained:

	R ²	R ⁷	R ⁶	R ¹	Yield (%)
38-1	Pr	<i>o</i> -NO ₂ Ph	<i>o</i> -CF ₃ Ph	Me	20 ^{a)}
38-2	<i>p</i> -BrBn	<i>c</i> -PrCH ₂	<i>o</i> -NO ₂ Ph	Et	15
38-3	Hep	<i>p</i> -CN-Bn	<i>o</i> -CF ₃ Ph	<i>i</i> -Pr	6
39-1	<i>p</i> - <i>t</i> BuBn	CH ₂ COO <i>t</i> Bu	<i>p</i> -ClPh	Me	14

10 Example 80

Methyl 4-S-(2'-O-acetyl-3'-O-benzyl-6'-O-tert-butyldiphenylsilyl-β-D-galactopyranosyl)-4-mercaptopbutyrate

A mixture of 4.0 g (6.9 mmol) of Example compound 14a and 2.1 g

(8.32 mmol) of dibutyltin oxide in 50 ml of benzene is heated under reflux for 2 h in a water separator. 25 ml of benzene are then removed by distillation and the mixture is treated with 3.1 g (8.3 mmol) of tetrabutylammonium bromide and 1.4 ml (11.8 mmol) of benzyl bromide.

- 5 The solution is stirred at 50°C for 16 h and then largely concentrated in vacuo. It is taken up in 50 ml of dichloromethane and washed three times with 10 ml of water each time. After drying over magnesium sulfate, the solvent is removed in vacuo. The crude product is purified by chromatography on silica gel (column 20 × 5 cm, eluent petroleum ether/ethyl acetate 4:1).

10 Yield 2.66 g (58%), yellowish oil, $[\alpha]_D^{25} = +3.0^\circ$ (c = 1, CHCl₃); R_F = 0.25 (petroleum ether/ethyl acetate 4:1).

Example 81

- 15 Methyl S-(2'-O-acetyl-3'-O-benzyl-4'-O-[1''-(R/S)-ethoxyethyl]-6'-O-tert-butylidiphenylsilyl-β-D-galactopyranosyl)-4-mercaptopbutyrate

- A solution of 2.58 g (3.86 mmol) of Example compound 80 in 100 ml of dichloromethane is stirred at room temp. for 4 h after addition of 50 ml of ethyl vinyl ether and 0.49 g (1.93 mmol) of pyridinium *p*-toluenesulfonate. The mixture is poured into satd sodium hydrogencarbonate solution and the aqueous phase is extracted with ethyl acetate. The combined organic phases are then dried over magnesium sulfate and freed from the solvent in vacuo. Purification is carried out by chromatography on silica gel (column 15 × 3 cm, eluent petroleum ether/ethyl acetate 4:1).

25 Yield 1.25 g (45%), yellowish oil, $[\alpha]_D^{25} = -13.3^\circ$ (c = 1, CHCl₃); R_F = 0.22 (petroleum ether/ethyl acetate 4:1).

Example 82

- 30 Methyl S-(3'-O-benzyl-4'-O-[1''-(R/S)-ethoxyethyl]-6'-O-tert-butylidiphenylsilyl-β-D-galactopyranosyl)-4-mercaptopbutyrate, polymer-bound


- According to the general working procedure, 1.45 g (1.60 mmol) of aminomethyl polystyrene are loaded with 1.25 g (1.69 mmol) of Example compound 81.

35 Loading according to sulfur content of the elemental analysis: 0.61 mmol/g.



85

Example 83

Functionalization of the 1, 2, 3, 4 and 6-position



- 1.a) KOtBu, DMF
b) R5X, 18-crown-6, DMF
2. TBAF, THF
3. Alkylation or carbamoylation (R5)
4. PPTS, dioxane/MeOH (10:1)
5. R6NCO, DMAP, dioxane
- 6.a) [Ir(COD)(PmePh₂)₂]PF₆, H₂, dioxane
b) PPTS, dioxane/MeOH (10:1); 50°C
7. R7NCO, DMAP, dioxane
- 8.a) Br₂, DTBP, CH₂Cl₂
b) R1OH, c-hexene, TEAB, CH₂Cl₂



5

The following compounds are obtained:

	R ²	R ⁵	R ⁶	R ⁷	R ¹	Yield (%)
41-1	<i>p</i> -tBuBn	2 × Et (from Et ₂ NH)	<i>p</i> -Cl-Ph	<i>o</i> -NO ₂ -Ph	Me	23
41-2	Hep	<i>o</i> -NO ₂ -Ph	<i>o</i> -CF ₃ -Ph	<i>p</i> -CN-Ph	Me	16*
41-3	<i>p</i> -BrBn	<i>m,p</i> -Cl ₂ -Ph	<i>o</i> -CF ₃ -Ph	<i>p</i> -Cl-Ph	<i>i</i> -Pr	5
42-1	<i>p</i> -BrBn	Hep	<i>o</i> -NO ₂ -Ph	<i>p</i> -Cl-Ph	Me	38
42-2	<i>p</i> -tBuBn	Bu	<i>o</i> -NO ₂ -Ph	<i>p</i> -CN-Ph	<i>i</i> -Pr	4
42-3	Bn	CH ₂ COOtBu	<i>o</i> -NO ₂ -Ph	<i>p</i> -Cl-Ph	Me	6

*) Minor component (according to chromatography)


Example 84

S-(3'-O-Allyl-4'-O-[1''-(R/S)-ethoxyethyl]-6'-O-tert-butyldiphenylsilyl- β -D-galactopyranosyl)-4-mercaptopbutyric acid, polymer-bound (40)


- 5 8.60 g (9.5 mmol) of aminomethylpolystyrene are loaded with 6.70 g (10.0 mmol) of 21 (Ex. 14c) according to the general procedure. Loading according to sulfur content of the elemental analysis: 0.93 mmol/g.

Claims:


1. A compound of the formula IIa, IIb or IIc:



IIa



IIb



IIc

in which:

Y is S or O;

X is O or N;

R¹ is a linker group which can be linked via a covalent bond to a support functionalized by a heteroatom;

R², in the case in which X is equal to O,

is acetyl or benzoyl;

in the case in which X is equal to N,

is a phthaloyl protective group, DDE (1-(4,4-dimethyl-2,6-dioxocyclohexylideneethyl) or NDE (2-acetyl-4-nitroindan-1,3-dione);

R³ is an allyl protective group;

R⁴ is ethoxyethyl or SEM (2-trimethylsilyl)ethoxymethyl);

or R³ and R⁴ together are isopropylidene or benzylidene;

R⁵ is tert-butyldimethylsilyl or tert-butyldiphenylsilyl.

2. A compound of the formula IIa, IIb or IIc as claimed in claim 1 wherein Y is S.

3. A compound of the formula IIb as claimed in claim 1 or 2, in which R^3 and R^4 together are isopropylidene or benzylidene and the other radicals X, Y, R^1 , R^2 and R^5 are as defined in claim 1 or 2.

4. A compound of the formula IIa, IIb or IIc as claimed in any one of claims 1 to 3 in which the linker group R^1 is a group of the formula III




in which n and p are 0 or 1, where p and n cannot simultaneously be 1;

R^9 is OR^{10} or $NR^{11}R^{11}$, where


R^{10} is H, (C_1-C_6) -alkyl, (C_1-C_6) -alkyl- (C_6-C_{12}) -aryl, and

R^{11} independently of one another is H, (C_1-C_6) -alkyl, (C_1-C_6) -alkyl- (C_6-C_{12}) -aryl.


5. A compound of the formula IIa, IIb or IIc



IIa



IIb



IIc

in which:

Y is S or O;

X is O or N;

R^1 is a group of the formula III



in which n and p are 0 or 1, where p and n cannot simultaneously be 1;

R^9 , is $NR^{11}R^{11}$, where one R^{11} is H, (C_1-C_6) -alkyl, (C_1-C_6) -alkyl- (C_6-C_{12}) -aryl, and the other R^{11} is a polymeric support;

R^2 in the case in which X is equal to O,
is acetyl or benzoyl;

in the case in which X is equal to N,
is a phthaloyl protective group, DDE (1-(4,4-dimethyl-2,6-dioxocyclohexylidenethyl) or NDE (2-acetyl-4-nitroindan-1,3-dione);


R^3 is an allyl protective group;

R^4 is ethoxyethyl or SEM (2-trimethylsilyl)ethoxymethyl);

or R^3 and R^4 together are isopropylidene or benzylidene;

R^5 is tert-butyldimethylsilyl or tert-butyldiphenylsilyl.

6. A compound of the formula IIa, IIb or IIc as claimed in claim 5 wherein the polymeric solid support is a cross-linked polystyrene or Tentagel.
7. A compound of the formula IIa, IIb or IIc as claimed in claim 6 wherein the cross-linked polystyrene is aminomethylpolystyrene.



(I)