

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2014/138212 A1

(43) International Publication Date
12 September 2014 (12.09.2014)

WIPO | PCT

(51) International Patent Classification:
A61K 31/437 (2006.01)

(IN). LEHOUX, Isabelle; 190 Hazel Avenue, Millbrae, California 94030 (US).

(21) International Application Number:
PCT/US2014/020700

(74) Agents: MCLEAN, Thomas H. et al.; Choate, Hall & Stewart LLP, Two International Place, Boston, Massachusetts 02110 (US).

(22) International Filing Date:
5 March 2014 (05.03.2014)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(25) Filing Language: English
(26) Publication Language: English
(30) Priority Data:
61/773,779 6 March 2013 (06.03.2013) US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(71) Applicant: ALLOSTERS THERAPEUTICS, INC. [US/US]; 1230 Bordeaux Drive, Sunnyvale, California 94089-1212 (US).

(72) Inventors: LEVY, Daniel E.; 37 Lakewood Circle, San Mateo, California 94402 (US). SCHULMAN, Howard; 560 Pena Court, Palo Alto, California 94306 (US). PARSELLI, Bheema; 6323 Oleander Way, San Diego, California 92130 (US). BRADLEY, Erin; 729 Nevada Avenue, San Mateo, California 94402 (US). NANGUNOORI, Sampath K.; 12-10-335, Dattathreya Nagar Colony, Andhra Pradesh (IN). DABBUGODDU, Brahmaiah; Flat: 305, C-block, Sri Lakshmi Ambica Residency, Netaji Nagar, Kothapet, Hyderabad-500 060, Andhra Pradesh

[Continued on next page]

(54) Title: CaMKII INHIBITORS AND USES THEREOF




Figure 1. CaMKII δ enzyme inhibition curve for compound I-7.

WO 2014/138212 A1

(57) Abstract: The present invention provides compounds useful as inhibitors of Ca²⁺/calmodulin-independent protein kinase (CaMKII), compositions thereof, and methods of using the same. Cardiovascular disease remains the number one cause of death in developed countries. Furthermore, incidence of cardiovascular disease has increased dramatically in developing countries. Although cardiovascular disease usually affects older adults, the antecedents of cardiovascular disease, notably atherosclerosis, begin in early life, making primary prevention efforts necessary from childhood.



Published:

— *with international search report (Art. 21(3))*

CaMKII INHIBITORS AND USES THEREOF

BACKGROUND OF THE INVENTION


[0001] Cardiovascular disease remains the number one cause of death in developed countries. Furthermore, incidence of cardiovascular disease has increased dramatically in developing countries. Although cardiovascular disease usually affects older adults, the antecedents of cardiovascular disease, notably atherosclerosis, begin in early life, making primary prevention efforts necessary from childhood. Therefore, there is increased emphasis on preventing atherosclerosis by modifying risk factors, such as healthy eating, exercise, and avoidance of smoking. It is estimated that 1 in 3 people will die from complications attributable to cardiovascular disease. “Global Atlas on Cardiovascular Disease Prevention and Control”, World Health Organization; January 2012. In order to stem the tide and address the shifting epidemiology of this disease, measures to prevent or reverse cardiovascular disease must be taken.

[0002] Obesity and diabetes mellitus are often linked to cardiovascular disease, as are a history of chronic kidney disease and hypercholesterolemia. In fact, cardiovascular disease is the most life threatening of the diabetic complications and diabetics are two- to four-fold more likely to die of cardiovascular-related causes than nondiabetics.

[0003] Diet and exercise, even when used in conjunction with the current pharmacotherapy, often do not provide sufficient control of cardiovascular symptoms. The continuing and highly prevalent problem of cardiovascular disease highlights the overwhelming need for new drugs to treat this condition and its underlying causes.

SUMMARY OF THE INVENTION

[0004] It has now been found that compounds of this invention, and pharmaceutically acceptable compositions thereof, are effective as inhibitors of Ca^{2+} /calmodulin-dependent protein kinase II (CaMKII). Such compounds have the general formula I:

**I**

or a pharmaceutically acceptable salt thereof, wherein each variable is as defined and described herein.

[0005] Compounds of the present invention, and pharmaceutically acceptable compositions thereof, are useful for treating a variety of diseases, disorders or conditions. For example, provided compounds are useful in treatment of diseases, disorders or conditions associated with the regulation and inhibition of CaMKII. Such diseases, disorders, or conditions include those described herein.

[0006] Compounds provided by this invention are also useful for the study of CaMKII enzymes in biological and pathological phenomena; the study of intracellular signal transduction pathways occurring in cardiac, vascular and other bodily tissues; and the comparative evaluation of new CaMKII inhibitors or other regulators of inflammation *in vitro* or *in vivo*.


BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 presents results of an *in vitro* CaMKII enzyme inhibition assay for compound I-7.

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS

1. General Description of Compounds of the Invention:

[0007] In certain embodiments, the present invention provides inhibitors of CaMKII. In some embodiments, such compounds include those of formula I:

**I**

or a pharmaceutically acceptable salt thereof, wherein:

one of R^{2'} and R^{3'} is -L¹-R^x, and the other is hydrogen;

L¹ is a covalent bond or a straight or branched C₁₋₆ aliphatic group, wherein one or more methylene groups are independently and optionally replaced by -NR^a- or -O-;

R^x is selected from the group consisting of NH₂, guanidino, 4-7 membered optionally substituted saturated heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen or sulfur, and 5-6 membered heteroaromatic ring having 1-2 heteroatoms independently selected from sulfur, nitrogen and oxygen;

each of R³, R⁴, R⁵, R⁶, R⁷, R⁸, R^{4'}, R^{5'}, and R^{6'} is independently selected from the group consisting of hydrogen, halogen, -CN, -CF₃, -OR, -NR₂, -NO₂, -COOR, -CONR₂, and -R;

each R^a is independently hydrogen or C₁₋₃ aliphatic; and

each R is independently hydrogen or optionally substituted C₁₋₆ aliphatic.

2. Compounds and Definitions:

[0008] Compounds of this invention include those described generally above, and are further illustrated by the classes, subclasses, and species disclosed herein. As used herein, the following definitions shall apply unless otherwise indicated. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75th Ed. Additionally, general principles of organic chemistry are described in "Organic Chemistry", Thomas Sorrell, University Science Books, Sausalito: 1999, and "March's Advanced Organic Chemistry", 5th Ed., Ed.: Smith, M.B. and March, J., John Wiley & Sons, New York: 2001, the entire contents of which are hereby incorporated by reference.

[0009] The term "aliphatic" or "aliphatic group", as used herein, means a straight-chain (i.e., unbranched) or branched, substituted or unsubstituted hydrocarbon chain that is completely saturated or that contains one or more units of unsaturation, or a monocyclic hydrocarbon or bicyclic hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic (also referred to herein as "carbocycle," "cycloaliphatic" or "cycloalkyl"), that has a single point of attachment to the rest of the molecule. Unless otherwise specified, aliphatic groups contain 1-6 aliphatic carbon atoms. In some embodiments, aliphatic groups contain 1-5 aliphatic carbon atoms. In other embodiments, aliphatic groups contain 1-4 aliphatic carbon atoms. In still other embodiments, aliphatic groups contain 1-3 aliphatic carbon atoms, and in yet other embodiments, aliphatic groups contain 1-2 aliphatic carbon atoms. In some embodiments, "cycloaliphatic" (or "carbocycle" or "cycloalkyl") refers to a monocyclic C₃-C₆ hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic, that has a single point of attachment to the rest of the molecule. Suitable aliphatic groups include, but are not limited to, linear or branched, substituted or unsubstituted alkyl, alkenyl, alkynyl groups and hybrids thereof such as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or (cycloalkyl)alkenyl.

[0010] The term "lower alkyl" refers to a C₁₋₄ straight or branched alkyl group. Exemplary lower alkyl groups are methyl, ethyl, propyl, isopropyl, butyl, isobutyl, and tert-butyl.

[0011] The term "lower haloalkyl" refers to a C₁₋₄ straight or branched alkyl group that is substituted with one or more halogen atoms.

[0012] The term "heteroatom" means one or more of oxygen, sulfur, nitrogen, phosphorus, or silicon (including, any oxidized form of nitrogen, sulfur, phosphorus, or silicon; the quaternized form of any basic nitrogen or; a substitutable nitrogen of a heterocyclic ring, for example N (as in 3,4-dihydro-2H-pyrrolyl), NH (as in pyrrolidinyl) or NR⁺ (as in N-substituted pyrrolidinyl)).


[0013] The term "unsaturated," as used herein, means that a moiety has one or more units of unsaturation.

[0014] As used herein, the term "bivalent C₁₋₈ (or C₁₋₆) saturated or unsaturated, straight or branched, hydrocarbon chain", refers to bivalent alkylene, alkenylene, and alkynylene chains that are straight or branched as defined herein.

[0015] The term “alkylene” refers to a bivalent alkyl group. An “alkylene chain” is a polymethylene group, i.e., $-(\text{CH}_2)_n-$, wherein n is a positive integer, preferably from 1 to 6, from 1 to 4, from 1 to 3, from 1 to 2, or from 2 to 3. A substituted alkylene chain is a polymethylene group in which one or more methylene hydrogen atoms are replaced with a substituent. Suitable substituents include those described below for a substituted aliphatic group.


[0016] The term “alkenylene” refers to a bivalent alkenyl group. A substituted alkenylene chain is a polymethylene group containing at least one double bond in which one or more hydrogen atoms are replaced with a substituent. Suitable substituents include those described below for a substituted aliphatic group.

[0017] As used herein, the term “cyclopropylenyl” refers to a bivalent cyclopropyl group of the



the following structure:

[0018] As used herein, the term “cyclobutylenyl” refers to a bivalent cyclobutyl group of the



following structure:

[0019] As used herein, the term “oxetanyl” refers to a bivalent oxetanyl group of the



following structure:

[0020] The term “halogen” means F, Cl, Br, or I.

[0021] The term “aryl” used alone or as part of a larger moiety as in “aralkyl,” “aralkoxy,” or “aryloxyalkyl,” refers to monocyclic and bicyclic ring systems having a total of five to 10 ring members, wherein at least one ring in the system is aromatic and wherein each ring in the system contains three to seven ring members. The term “aryl” may be used interchangeably with the term “aryl ring”. In certain embodiments of the present invention, “aryl” refers to an aromatic ring system which includes, but not limited to, phenyl, biphenyl, naphthyl, anthracyl and the like, which may bear one or more substituents. Also included within the scope of the term “aryl,” as it is used herein, is a group in which an aromatic ring is fused to one or more non-aromatic rings, such as indanyl, phthalimidyl, naphthimidyl, phenanthridinyl, or tetrahydronaphthyl, and the like.

[0022] The terms “heteroaryl” and “heteroar–,” used alone or as part of a larger moiety, e.g., “heteroaralkyl,” or “heteroaralkoxy,” refer to groups having 5 to 10 ring atoms, preferably 5, 6, or 9 ring atoms; having 6, 10, or 14 π electrons shared in a cyclic array; and having, in addition to carbon atoms, from one to five heteroatoms. The term “heteroatom” refers to nitrogen, oxygen, or sulfur, and includes any oxidized form of nitrogen or sulfur, and any quaternized form of a basic nitrogen. Heteroaryl groups include, without limitation, thienyl, furanyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolizinyl, purinyl, naphthyridinyl, and pteridinyl. The terms “heteroaryl” and “heteroar–”, as used herein, also include groups in which a heteroaromatic ring is fused to one or more aryl, cycloaliphatic, or heterocyclyl rings, where the radical or point of attachment is on the heteroaromatic ring. Nonlimiting examples include indolyl, isoindolyl, benzothienyl, benzofuranyl, dibenzofuranyl, indazolyl, benzimidazolyl, benzthiazolyl, quinolyl, isoquinolyl, cinnolinyl, phthalazinyl, quinazolinyl, quinoxalinyl, 4H-quinolizinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, phenoxazinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, and pyrido[2,3-b]–1,4–oxazin–3(4H)–one. A heteroaryl group may be mono– or bicyclic. The term “heteroaryl” may be used interchangeably with the terms “heteroaryl ring,” “heteroaryl group,” or “heteroaromatic,” any of which terms include rings that are optionally substituted. The term “heteroaralkyl” refers to an alkyl group substituted by a heteroaryl, wherein the alkyl and heteroaryl portions independently are optionally substituted.

[0023] As used herein, the terms “heterocycle,” “heterocyclyl,” “heterocyclic radical,” and “heterocyclic ring” are used interchangeably and refer to a stable 5– to 7–membered monocyclic or 7–10–membered bicyclic heterocyclic moiety that is either saturated or partially unsaturated, and having, in addition to carbon atoms, one or more, preferably one to four, heteroatoms, as defined above. When used in reference to a ring atom of a heterocycle, the term “nitrogen” includes a substituted nitrogen. As an example, in a saturated or partially unsaturated ring having 0–3 heteroatoms selected from oxygen, sulfur or nitrogen, the nitrogen may be N (as in 3,4–dihydro–2H–pyrrolyl), NH (as in pyrrolidinyl), or ^+NR (as in N –substituted pyrrolidinyl).

[0024] A heterocyclic ring can be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure and any of the ring atoms can be optionally substituted. Examples of such saturated or partially unsaturated heterocyclic radicals include, without

limitation, tetrahydrofuryl, tetrahydrothiophenyl, pyrrolidinyl, piperidinyl, pyrrolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, decahydroquinolinyl, oxazolidinyl, piperazinyl, dioxanyl, dioxolanyl, diazepinyl, oxazepinyl, thiazepinyl, morpholinyl, and quinuclidinyl. The terms “heterocycle,” “heterocycl,” “heterocycl ring,” “heterocyclic group,” “heterocyclic moiety,” and “heterocyclic radical,” are used interchangeably herein, and also include groups in which a heterocycl ring is fused to one or more aryl, heteroaryl, or cycloaliphatic rings, such as indolinyl, 3*H*-indolyl, chromanyl, phenanthridinyl, or tetrahydroquinolinyl, where the radical or point of attachment is on the heterocycl ring. A heterocycl group may be mono- or bicyclic. The term “heterocyclalkyl” refers to an alkyl group substituted by a heterocycl, wherein the alkyl and heterocycl portions independently are optionally substituted.

[0025] As used herein, the term “partially unsaturated” refers to a ring moiety that includes at least one double or triple bond. The term “partially unsaturated” is intended to encompass rings having multiple sites of unsaturation, but is not intended to include aryl or heteroaryl moieties, as herein defined.

[0026] As described herein, compounds of the invention may contain “optionally substituted” moieties. In general, the term “substituted,” whether preceded by the term “optionally” or not, means that one or more hydrogens of the designated moiety are replaced with a suitable substituent. Unless otherwise indicated, an “optionally substituted” group may have a suitable substituent at each substitutable position of the group, and when more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. Combinations of substituents envisioned by this invention are preferably those that result in the formation of stable or chemically feasible compounds. The term “stable,” as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and, in certain embodiments, their recovery, purification, and use for one or more of the purposes disclosed herein.

[0027] Suitable monovalent substituents on a substitutable carbon atom of an “optionally substituted” group are independently halogen; $-(CH_2)_{0-4}R^\circ$; $-(CH_2)_{0-4}OR^\circ$; $-O(CH_2)_{0-4}R^\circ$, $-O-(CH_2)_{0-4}C(O)OR^\circ$; $-(CH_2)_{0-4}CH(OR^\circ)_2$; $-(CH_2)_{0-4}SR^\circ$; $-(CH_2)_{0-4}Ph$, which may be substituted with R° ; $-(CH_2)_{0-4}O(CH_2)_{0-1}Ph$ which may be substituted with R° ; $-CH=CHPh$, which may be substituted with R° ; $-(CH_2)_{0-4}O(CH_2)_{0-1}-pyridyl$ which may be substituted with R° ; $-NO_2$; $-CN$;

$-\text{N}_3;$ $-(\text{CH}_2)_{0-4}\text{N}(\text{R}^\circ)_2;$ $-(\text{CH}_2)_{0-4}\text{N}(\text{R}^\circ)\text{C}(\text{O})\text{R}^\circ;$ $-\text{N}(\text{R}^\circ)\text{C}(\text{S})\text{R}^\circ;$ $-(\text{CH}_2)_{0-4}$
 $\text{N}(\text{R}^\circ)\text{C}(\text{O})\text{NR}^\circ_2;$ $-\text{N}(\text{R}^\circ)\text{C}(\text{S})\text{NR}^\circ_2;$ $-(\text{CH}_2)_{0-4}\text{N}(\text{R}^\circ)\text{C}(\text{O})\text{OR}^\circ;$ $-\text{N}(\text{R}^\circ)\text{N}(\text{R}^\circ)\text{C}(\text{O})\text{R}^\circ;$ $-\text{N}(\text{R}^\circ)\text{N}(\text{R}^\circ)\text{C}(\text{O})\text{NR}^\circ_2;$ $-\text{N}(\text{R}^\circ)\text{N}(\text{R}^\circ)\text{C}(\text{O})\text{OR}^\circ;$ $-(\text{CH}_2)_{0-4}\text{C}(\text{O})\text{R}^\circ;$ $-\text{C}(\text{S})\text{R}^\circ;$ $-(\text{CH}_2)_{0-4}\text{C}(\text{O})\text{OR}^\circ;$ $-(\text{CH}_2)_{0-4}\text{C}(\text{O})\text{SR}^\circ;$ $-(\text{CH}_2)_{0-4}\text{C}(\text{O})\text{OSiR}^\circ_3;$ $-(\text{CH}_2)_{0-4}\text{OC}(\text{O})\text{R}^\circ;$ $-\text{OC}(\text{O})(\text{CH}_2)_{0-4}\text{SR}-;$ $\text{SC}(\text{S})\text{SR}^\circ;$ $-(\text{CH}_2)_{0-4}\text{SC}(\text{O})\text{R}^\circ;$ $-(\text{CH}_2)_{0-4}\text{C}(\text{O})\text{NR}^\circ_2;$ $-\text{C}(\text{S})\text{NR}^\circ_2;$ $-\text{C}(\text{S})\text{SR}^\circ;$ $-\text{SC}(\text{S})\text{SR}^\circ;$ $-(\text{CH}_2)_{0-4}\text{OC}(\text{O})\text{NR}^\circ_2;$ $-\text{C}(\text{O})\text{N}(\text{OR}^\circ)\text{R}^\circ;$ $-\text{C}(\text{O})\text{C}(\text{O})\text{R}^\circ;$ $-\text{C}(\text{O})\text{CH}_2\text{C}(\text{O})\text{R}^\circ;$ $-\text{C}(\text{NOR}^\circ)\text{R}^\circ;$ $-(\text{CH}_2)_{0-4}\text{SSR}^\circ;$ $-(\text{CH}_2)_{0-4}\text{S}(\text{O})_2\text{R}^\circ;$ $-(\text{CH}_2)_{0-4}\text{S}(\text{O})_2\text{OR}^\circ;$ $-(\text{CH}_2)_{0-4}\text{OS}(\text{O})_2\text{R}^\circ;$ $-\text{S}(\text{O})_2\text{NR}^\circ_2;$ $-(\text{CH}_2)_{0-4}\text{S}(\text{O})\text{R}^\circ;$ $-\text{N}(\text{R}^\circ)\text{S}(\text{O})_2\text{NR}^\circ_2;$ $-\text{N}(\text{R}^\circ)\text{S}(\text{O})_2\text{R}^\circ;$ $-\text{N}(\text{OR}^\circ)\text{R}^\circ;$ $-\text{C}(\text{NH})\text{NR}^\circ_2;$ $-\text{P}(\text{O})_2\text{R}^\circ;$ $-\text{P}(\text{O})\text{R}^\circ_2;$ $-\text{OP}(\text{O})\text{R}^\circ_2;$ $-\text{OP}(\text{O})(\text{OR}^\circ)_2;$ $\text{SiR}^\circ_3;$ $-(\text{C}_{1-4}$
 straight or branched alkylene) $\text{O}-\text{N}(\text{R}^\circ)_2;$ or $-(\text{C}_{1-4}$ straight or branched alkylene) $\text{C}(\text{O})\text{O}-\text{N}(\text{R}^\circ)_2,$ wherein each R° may be substituted as defined below and is independently hydrogen, C_{1-6} aliphatic, $-\text{CH}_2\text{Ph},$ $-\text{O}(\text{CH}_2)_{0-1}\text{Ph},$ $-\text{CH}_2$ -(5-6 membered heteroaryl ring), or a 5-6-membered saturated, partially unsaturated, or aryl ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or, notwithstanding the definition above, two independent occurrences of R° , taken together with their intervening atom(s), form a 3-12-membered saturated, partially unsaturated, or aryl mono- or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, which may be substituted as defined below.

[0028] Suitable monovalent substituents on R° (or the ring formed by taking two independent occurrences of R° together with their intervening atoms), are independently halogen, $-(\text{CH}_2)_{0-2}\text{R}^\bullet,$ $-(\text{haloR}^\bullet),$ $-(\text{CH}_2)_{0-2}\text{OH},$ $-(\text{CH}_2)_{0-2}\text{OR}^\bullet,$ $-(\text{CH}_2)_{0-2}\text{CH}(\text{OR}^\bullet)_2;$ $-\text{O}(\text{haloR}^\bullet),$ $-\text{CN},$ $-\text{N}_3,$ $-(\text{CH}_2)_{0-2}\text{C}(\text{O})\text{R}^\bullet,$ $-(\text{CH}_2)_{0-2}\text{C}(\text{O})\text{OH},$ $-(\text{CH}_2)_{0-2}\text{C}(\text{O})\text{OR}^\bullet,$ $-(\text{CH}_2)_{0-2}\text{SR}^\bullet,$ $-(\text{CH}_2)_{0-2}\text{SH},$ $-(\text{CH}_2)_{0-2}\text{NH}_2,$ $-(\text{CH}_2)_{0-2}\text{NHR}^\bullet,$ $-(\text{CH}_2)_{0-2}\text{NR}^\bullet_2,$ $-\text{NO}_2,$ $-\text{SiR}^\bullet_3,$ $-\text{OSiR}^\bullet_3,$ $-\text{C}(\text{O})\text{SR}^\bullet,$ $-(\text{C}_{1-4}$ straight or branched alkylene) $\text{C}(\text{O})\text{OR}^\bullet,$ or $-\text{SSR}^\bullet$ wherein each R^\bullet is unsubstituted or where preceded by “halo” is substituted only with one or more halogens, and is independently selected from C_{1-4} aliphatic, $-\text{CH}_2\text{Ph},$ $-\text{O}(\text{CH}_2)_{0-1}\text{Ph},$ or a 5-6-membered saturated, partially unsaturated, or aryl ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. Suitable divalent substituents on a saturated carbon atom of R° include $=\text{O}$ and $=\text{S}.$

[0029] Suitable divalent substituents on a saturated carbon atom of an “optionally substituted” group include the following: =O, =S, =NNR^{*}₂, =NNHC(O)R^{*}, =NNHC(O)OR^{*}, =NNHS(O)R^{*}₂, =NR^{*}, =NOR^{*}, -O(C(R^{*}₂))₂₋₃O-, or -S(C(R^{*}₂))₂₋₃S-, wherein each independent occurrence of R^{*} is selected from hydrogen, C₁₋₆ aliphatic which may be substituted as defined below, or an unsubstituted 5–6–membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. Suitable divalent substituents that are bound to vicinal substitutable carbons of an “optionally substituted” group include: -O(CR^{*}₂)₂₋₃O-, wherein each independent occurrence of R^{*} is selected from hydrogen, C₁₋₆ aliphatic which may be substituted as defined below, or an unsubstituted 5–6–membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0030] Suitable substituents on the aliphatic group of R^{*} include halogen, -R[•], -(haloR[•]), -OH, -OR[•], -O(haloR[•]), -CN, -C(O)OH, -C(O)OR[•], -NH₂, -NHR[•], -NR[•]₂, or -NO₂, wherein each R[•] is unsubstituted or where preceded by “halo” is substituted only with one or more halogens, and is independently C₁₋₄ aliphatic, -CH₂Ph, -O(CH₂)₀₋₁Ph, or a 5–6–membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0031] Suitable substituents on a substitutable nitrogen of an “optionally substituted” group include -R[†], -NR[†]₂, -C(O)R[†], -C(O)OR[†], -C(O)C(O)R[†], -C(O)CH₂C(O)R[†], -S(O)₂R[†], -S(O)₂NR[†]₂, -C(S)NR[†]₂, -C(NH)NR[†]₂, or -N(R[†])S(O)₂R[†]; wherein each R[†] is independently hydrogen, C₁₋₆ aliphatic which may be substituted as defined below, unsubstituted -OPh, or an unsubstituted 5–6–membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or, notwithstanding the definition above, two independent occurrences of R[†], taken together with their intervening atom(s) form an unsubstituted 3–12–membered saturated, partially unsaturated, or aryl mono– or bicyclic ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0032] Suitable substituents on the aliphatic group of R[†] are independently halogen, -R[•], -(haloR[•]), -OH, -OR[•], -O(haloR[•]), -CN, -C(O)OH, -C(O)OR[•], -NH₂, -NHR[•], -NR[•]₂, or -NO₂, wherein each R[•] is unsubstituted or where preceded by “halo” is substituted only with one or more halogens, and is independently C₁₋₄ aliphatic, -CH₂Ph, -O(CH₂)₀₋₁Ph, or a 5–6–

membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0033] As used herein, the term "pharmaceutically acceptable salt" refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge et al., describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 1977, 66, 1–19, incorporated herein by reference. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy–ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like.


[0034] Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C_{1-4}\text{alkyl})_4$ salts. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate.

[0035] Unless otherwise stated, structures depicted herein are also meant to include all isomeric (e.g., enantiomeric, diastereomeric, and geometric (or conformational)) forms of the

structure; for example, the R and S configurations for each asymmetric center, Z and E double bond isomers, and Z and E conformational isomers. Therefore, single stereochemical isomers as well as enantiomeric, diastereomeric, and geometric (or conformational) mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, all tautomeric forms of the compounds of the invention are within the scope of the invention. Additionally, unless otherwise stated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures including the replacement of hydrogen by deuterium or tritium, or the replacement of a carbon by a ¹³C- or ¹⁴C-enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools, as probes in biological assays, or as therapeutic agents in accordance with the present invention. In certain embodiments, a warhead moiety, R¹, of a provided compound comprises one or more deuterium atoms.

3. Description of Exemplary Embodiments:

[0036] In certain embodiments, the present invention provides inhibitors of CaMKII. In some embodiments, such compounds include those of formula I:



I

or a pharmaceutically acceptable salt thereof, wherein:

one of R² and R³ is -L¹-R^x, and the other is hydrogen;

L¹ is a covalent bond or a straight or branched C₁₋₆ aliphatic group, wherein one or more methylene groups are independently and optionally replaced by -NR^a- or -O-;

R^x is selected from the group consisting of NH₂, guanidino, 4-7 membered optionally substituted saturated heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen or sulfur, and 5-6 membered heteroaromatic ring having 1-2 heteroatoms independently selected from sulfur, nitrogen and oxygen;

each of R³, R⁴, R⁵, R⁶, R⁷, R⁸, R^{4'}, R^{5'}, and R^{6'} is independently selected from the group consisting of hydrogen, halogen, -CN, -CF₃, -OR, -NR₂, -NO₂, -COOR, -CONR₂, and -R; each R^a is independently hydrogen or C₁₋₃ aliphatic; and each R is independently hydrogen or optionally substituted C₁₋₆ aliphatic.

[0037] As defined generally above, one of R² and R³ is -L¹-R^x, and the other is hydrogen. In some embodiments R² is hydrogen and R³ is -L¹-R^x. In some embodiments one of R² is -L¹-R^x and R³ is hydrogen.

[0038] As defined generally above, L¹ is a covalent bond or a straight or branched C₁₋₆ aliphatic group, wherein one or more methylene groups are independently and optionally replaced by -NR^a- or -O-. In some embodiments L¹ is a covalent bond. In some embodiments L¹ is a straight or branched C₁₋₆ aliphatic group wherein one or more methylene groups are independently and optionally replaced by -NR^a- or -O-. In some embodiments L¹ is a C₁₋₃ aliphatic group. In some embodiments L¹ is methylene. In some embodiments L¹ is a C₁₋₄ aliphatic group wherein one methylene group is replaced by -NR^a- or -O-. In some embodiments, L¹ is a C₁₋₄ alkylene group wherein the first methylene is replaced by -NR^a- (i.e. -NR^a-(CH₂)₀₋₃-). In some embodiments L¹ is a C₁₋₄ alkylene group wherein the first methylene is replaced by -O- (i.e. -O-(CH₂)₀₋₃-).


[0039] As defined generally above, R^x is selected from the group consisting of NH₂, guanidino, 4-7 membered optionally substituted saturated heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen or sulfur, and 5-6 membered heteroaromatic ring having 1-2 heteroatoms independently selected from nitrogen and oxygen. In some embodiments R^x is NH₂. In some embodiments R^x is guanidine. In some embodiments R^x is a 4-7 membered optionally substituted saturated heterocyclic group having 1-2 heteroatoms independently selected from nitrogen and oxygen. In some embodiments R^x is piperazino. In some embodiments R^x is a 5-6 membered heteroaromatic ring having 1-2 heteroatoms independently selected from nitrogen and oxygen. In some embodiments, R^x is imidazolo.

[0040] As defined generally above, each of R³, R⁴, R⁵, R⁶, R⁷, R⁸, R^{4'}, R^{5'}, and R^{6'} is independently selected from the group consisting of hydrogen, halogen, -CN, -CF₃, -OR, -NR₂, -NO₂, -COOR, -CONR₂, and -R. In some embodiments both of R⁶ and R⁷ are hydrogen. In some embodiments R⁶ is hydrogen and R⁷ is selected from the group consisting

of halogen, -CN, -CF₃, -OR, -NR₂, -NO₂, -COOR, -CONR₂, and -R. In some embodiments R⁷ is hydrogen and R⁶ is selected from the group consisting of halogen, -CN, -CF₃, -OR, -NR₂, -NO₂, -COOR, -CONR₂, and -R. In some embodiments both R⁶ and R⁷ are independently selected from the group consisting of halogen, -CN, -CF₃, -OR, -NR₂, -NO₂, -COOR, -CONR₂, and -R. In some embodiments R⁶ is halogen. In some embodiments R⁶ is -CN. In some embodiments R⁶ is -OR. In some embodiments R⁶ is -OH. In some embodiments R⁶ is -NR₂. In some embodiments R⁶ is -NO₂. In some embodiments R⁶ is -COOR. In some embodiments R⁶ is -CONR₂. In some embodiments R⁶ is methoxy. In some embodiments R⁶ is -R, wherein R is C₁₋₃ aliphatic optionally substituted by one or more fluorines. In some embodiments R⁶ is methyl. In some embodiments R⁶ is trifluoromethyl. In some embodiments R⁷ is halogen. In some embodiments R⁷ is -CN. In some embodiments R⁷ is -OR. In some embodiments R⁷ is -OH. In some embodiments R⁷ is -NR₂. In some embodiments R⁷ is -NO₂. In some embodiments R⁷ is -COOR. In some embodiments R⁷ is -CONR₂. In some embodiments R⁷ is methoxy. In some embodiments R⁷ is -R, wherein R is C₁₋₃ aliphatic optionally substituted by one or more fluorines. In some embodiments R⁷ is methyl. In some embodiments R⁷ is trifluoromethyl.

[0041] As defined generally above, each R^a is independently hydrogen or C₁₋₃ aliphatic. In some embodiments each R^a is hydrogen. In some embodiments each R^a is C₁₋₃ aliphatic.


[0042] In certain embodiments, the present invention provides a compound of formula **I** wherein R³, R⁴, R⁵, R⁸, R^{4'}, R^{5'}, and R^{6'} are each hydrogen, thereby forming a compound of formula **I-a**:



I-a

or a pharmaceutically acceptable salt thereof, wherein each of R^{2'}, R^{3'}, R⁶, and R⁷ is defined above and described in embodiments herein, both singly and in combination.


[0043] In certain embodiments, the present invention provides a compound of formula **I** wherein R^{2'} is hydrogen and R^{3'} is -L¹-R^x, thereby forming a compound of formula **II**:



II

or a pharmaceutically acceptable salt thereof, wherein each of L¹, R^x, R⁶, and R⁷ is defined above and described in embodiments herein, both singly and in combination.


[0044] In certain embodiments, the present invention provides a compound of formula **I** wherein R² is -L¹-R^x and R³ is hydrogen, thereby forming a compound of formula **III**:



III

or a pharmaceutically acceptable salt thereof, wherein each of L¹, R^x, R⁶, and R⁷ is defined above and described in embodiments herein, both singly and in combination.


[0045] In certain embodiments, the present invention provides a compound of formula **II** wherein L¹ is -NR^a-(CH₂)₀₋₃-, thereby forming a compound of formula **IV**:



IV


or a pharmaceutically acceptable salt thereof, wherein each of R^x, R⁶, R⁷, and R^a is defined above and described in embodiments herein, both singly and in combination.

[0046] In certain embodiments, the present invention provides a compound of formula **III** wherein L¹ is -NR^a-(CH₂)₀₋₃-, thereby forming a compound of formula **V**:



or a pharmaceutically acceptable salt thereof, wherein each of R^x, R⁶, R⁷, and R^a is defined above and described in embodiments herein, both singly and in combination.


[0047] In certain embodiments, the present invention provides a compound of formula **II** wherein L¹ is -O-(CH₂)₀₋₃-, thereby forming a compound of formula **VI**:



VI

or a pharmaceutically acceptable salt thereof, wherein each of R^x, R⁶, and R⁷ is defined above and described in embodiments herein, both singly and in combination.


[0048] In certain embodiments, the present invention provides a compound of formula **III** wherein L¹ is -O-(CH₂)₀₋₃-, thereby forming a compound of formula **VII**:



VII

or a pharmaceutically acceptable salt thereof, wherein each of R^x, R⁶, and R⁷ is defined above and described in embodiments herein, both singly and in combination.


[0049] In certain embodiments, the present invention provides a compound of formula **II** wherein L¹ is a covalent bond, thereby forming a compound of formula **VIII**:



VIII

or a pharmaceutically acceptable salt thereof, wherein each of R^x, R⁶, and R⁷ is defined above and described in embodiments herein, both singly and in combination.

[0050] In certain embodiments, the present invention provides a compound of formula **III** wherein L¹ is a covalent bond, thereby forming a compound of formula **IX**:







IX






or a pharmaceutically acceptable salt thereof, wherein each of R^x, R⁶, and R⁷ is defined above and described in embodiments herein, both singly and in combination.





[0051] Exemplary compounds of formula **I** are set forth in Table 1, below:

Table 1. Exemplary Compounds of Formula I

Compound ID	Compound Structure	Compound ID	Compound Structure
I-1		I-2	

Compound ID	Compound Structure
I-3	
I-4	
I-5	
I-6	
I-7	

Compound ID	Compound Structure
I-8	
I-9	
I-10	
I-11	
I-12	

Compound ID	Compound Structure
I-13	
I-14	
I-15	
I-16	

[0052] In certain embodiments, the present invention provides any compound selected from those depicted in Table 1, above, or a pharmaceutically acceptable salt thereof.

[0053] Compounds or salts thereof provided by the present invention may be utilized in any of a variety of forms. For example, in some embodiments, provided compounds (or salts thereof) are utilized in a solid form; in some such embodiments, provided compounds (or salts thereof) are utilized in an amorphous solid form. In some embodiments, provided compounds are utilized in a crystalline solid form. In some embodiments, provided compounds (or salts thereof) are utilized in a solid form (e.g., a crystalline solid form) that is a solvate or hydrate.

4. Uses, Formulation and Administration and Pharmaceutically Acceptable Compositions

[0054] According to some embodiments, the invention provides a composition comprising a compound of this invention or a pharmaceutically acceptable derivative thereof and a pharmaceutically acceptable carrier, adjuvant, or vehicle.

[0055] In certain embodiments, the invention provides compositions containing an amount of compound effective to measurably inhibit CaMKII, in a biological sample or in a patient. In certain embodiments, the amount of compound in compositions of this invention is such that is effective to measurably inhibit a CaMKII mediated biological process in a biological sample or in a patient. In certain embodiments, provided compositions contain a unit dose amount of a compound described herein, wherein administration of such unit dose amount as part of a therapeutic regimen correlates with a desired pharmacologic and/or therapeutic outcome.

[0056] In certain embodiments, a composition of this invention is formulated for administration to a patient in need of such composition. In some embodiments, a composition of this invention is formulated for oral administration to a patient.

[0057] As used herein, a “dosing regimen” or “therapeutic regimen” refers to a set of unit doses (typically more than one) that are administered individually to a subject, typically separated by periods of time. In some embodiments, a given therapeutic agent has a recommended dosing regimen, which may involve one or more doses. In some embodiments, a dosing regimen comprises a plurality of doses each of which are separated from one another by a time period of the same length; in some embodiments, a dosing regime comprises a plurality of doses and at least two different time periods separating individual doses. In some embodiments,

all doses within a dosing regimen are of the same unit dose amount. In some embodiments, different doses within a dosing regimen are of different amounts. In some embodiments, a dosing regimen comprises a first dose in a first dose amount, followed by one or more additional doses in a second dose amount different from the first dose amount. In some embodiments, a dosing regimen comprises a first dose in a first dose amount, followed by one or more additional doses in a second dose amount same as the first dose amount.

[0058] The term “patient,” as used herein, means an animal, often a mammal, and in many embodiments a human.

[0059] The term “pharmaceutically acceptable carrier, adjuvant, or vehicle”, as used herein, refers to a non-toxic carrier, adjuvant, or vehicle that does not destroy the pharmacological activity of the compound with which it is formulated. Pharmaceutically acceptable carriers, adjuvants or vehicles that may be used in the compositions of this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

[0060] A “pharmaceutically acceptable derivative” means any non-toxic salt, ester, or salt of an ester of a compound of this invention that, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof.

[0061] As used herein, the term "inhibitorily active metabolite or residue thereof" means that a metabolite or residue thereof is also an inhibitor of CaMKII or is retains therapeutic activity in treating the same disease, disorder or condition.

[0062] Compositions of the present invention may be formulated for any appropriate route of administration. For example, in some embodiments, provided compositions may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal,

intrahepatic, intralesional and intracranial injection or infusion techniques. In some embodiments, provided compositions are administered orally, intraperitoneally or intravenously. Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. Such suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents.

[0063] In some embodiments, pharmaceutically acceptable compositions of the invention may be formulated as injectable preparations. Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[0064] In some embodiments, injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[0065] In some embodiments, for example in order to prolong effects of a compound or composition, it may be desirable to slow the absorption of the compound from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the compound then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form. Alternatively or additionally, delayed absorption of a parenterally administered compound form is accomplished by dissolving or suspending the compound in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the compound in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of compound to polymer and the nature of the particular polymer employed, the rate of compound release can be controlled. Examples of other biodegradable polymers include

poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the compound in liposomes or microemulsions that are compatible with body tissues.

[0066] In some embodiments, sterile injectable preparations may be or include a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium.

[0067] For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. Such oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents that are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

[0068] Pharmaceutically acceptable compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

[0069] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating

agents such as agar--agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

[0070] Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[0071] In some embodiments, provided compounds can be in micro-encapsulated form with one or more excipients as noted above. Solid dosage forms such as tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms, the active compound may be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

[0072] Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[0073] Alternatively or additionally, pharmaceutically acceptable compositions of this invention may be administered in the form of suppositories for rectal administration. Such compositions can be prepared by combining a provided compound with a suitable non-irritating excipient that is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

[0074] In some embodiments, pharmaceutically acceptable compositions of this invention may be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

[0075] Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

[0076] For topical applications, provided pharmaceutically acceptable compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively or additionally, provided pharmaceutically acceptable compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more

pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

[0077] Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, ear drops, and eye drops are also contemplated as being within the scope of this invention. Additionally, the present invention contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms can be made by dissolving or dispersing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

[0078] For ophthalmic use, provided pharmaceutically acceptable compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively or additionally, for ophthalmic uses, the pharmaceutically acceptable compositions may be formulated in an ointment such as petrolatum.

[0079] In some embodiments, pharmaceutically acceptable compositions of this invention may be administered by nasal aerosol or inhalation. Such compositions may be prepared according to techniques well-known in the art of pharmaceutical formulation, for example as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

[0080] In some embodiments, pharmaceutically acceptable compositions of this invention are formulated for oral administration. Such formulations may be administered with or without food. In some embodiments, pharmaceutically acceptable compositions of this invention are administered without food. In some embodiments, pharmaceutically acceptable compositions of this invention are administered with food.

[0081] The amount of compounds of the present invention that may be combined with the carrier materials to produce a composition in a single dosage form will vary depending upon the host treated, the particular mode of administration. In some embodiments provided compositions are formulated so that a dosage of between 0.01 - 100 mg/kg body weight/day of the inhibitor can be administered to a patient receiving these compositions.

[0082] It should also be understood that a specific dosage and treatment regimen for any particular patient may depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. In some embodiments, amount of a compound of the present invention included in a composition described herein is determined by activity and/or bioavailability of the particular compound, so that compositions of different compounds may include different absolute amounts of compound.

Uses of Compounds and Pharmaceutically Acceptable Compositions

[0083] Compounds and compositions described herein are useful in the treatment of any of a variety of diseases, disorders, and conditions. In some embodiments, provided compounds and compositions are useful in the treatment of diseases, disorders, or conditions associated with activity of CaMKII.

[0084] Ca^{2+} /calmodulin-dependent protein kinase (CaMKII) is a serine/threonine kinase. Several lines of evidence strongly support the concept of direct inhibition of CaMKII activity as an important therapeutic target for treating a range of diseases including cardiovascular diseases such as atrial fibrillation, ventricular arrhythmia, heart failure, cardiac hypertrophy, atherosclerosis, and in-stent restenosis in coronary artery disease; use in cardioprotection; inflammatory lung diseases such as asthma; neurological diseases and conditions such as pain, stroke, ischemia, hypoxia, opioid tolerance and dependence, and macular degeneration; metabolic disorders such as type II diabetes, insulin resistance, and obesity; cancer and other proliferative disorders, such as osteosarcoma, melanoma, and prostate cancer; bone diseases such as osteoporosis; and inflammatory diseases such as rheumatoid arthritis.

[0085] Ca^{2+} acts as an informational or second messenger to modulate the effects of many hormones and neurotransmitters as well as during depolarization of excitable membranes in

heart, skeletal muscle, and brain. In many instances Ca^{2+} mediates its effects via a Ca^{2+} binding protein termed calmodulin. The Ca^{2+} /calmodulin complex can coordinate multiple cellular functions by activation CaMKII, a major multifunctional protein kinase in diverse tissues, including heart, brain, skeletal muscle, and liver. Strong activation of the kinase by Ca^{2+} can result in a self-phosphorylation or autophosphorylation that makes the kinase persistently active, a state that can also be achieved by oxidation.

[0086] CaMKII refers to four highly conserved isoforms of CaMKII, α , β , γ , and δ . CaMKII α , β , and γ are widely expressed, with the δ isoforms predominating in heart. Upon activation, CaMKII catalyzes a transfer of phosphate from ATP onto specific sites on select substrate proteins, such as the Ryanodine receptor RyR2 and the L-type Ca^{2+} channel in heart. These isoforms have been crystalized as truncated monomeric catalytic domains (Rellos P, Pike ACW, Niesen FH, Salah E, Lee WH, von Delft F, Knapp S. (2010) "Structure of the CaMKIIdelta/Calmodulin Complex Reveals the Molecular Mechanism of CaMKII Kinase Activation" *PLoS Biol* 8:e1000426) as well as in the native multimeric holoenzyme (Chao LH, Stratton MM, Lee I-H, Rosenberg OS, Levitz J, Mandell DJ, Kortemme T, Groves JT, Schulman H, Kuriyan J. (2011) "A Mechanism for Tunable Autoinhibition in the Structure of a Human Ca^{2+} /Calmodulin-Dependent Kinase II Holoenzyme" *Cell* 146:732-745). Comparison of the sequences, three-dimensional structure of the catalytic sites, and limited pharmacology suggest a great deal of homology between the isoforms, and ATP competitive inhibitors of any of the isoforms will likely inhibit all of the others. ATP competitive inhibitors are also known to block both the Ca^{2+} /calmodulin-activated enzyme as well the persistently activated forms generated by autophosphorylation or oxidation by reactive oxygen.

[0087] Pharmacological and genetic inhibition of CaMKII reduces ryanodine receptor-mediated calcium leaks and blocked induction of atrial fibrillation in a mouse model of atrial fibrillation. In atrial cells from atrial fibrillation patients CaMKII activity is increased leading to calcium leaks that promote atrial fibrillation, while inhibition of the kinase reduces the calcium leak. (Dobrev D, et al., "Novel molecular targets for atrial fibrillation therapy" (2012) *Nature Reviews Drug Discovery* 11:275-291; Chelu MG, et al., "Calmodulin kinase II-mediated sarcoplasmic reticulum Ca^{2+} leak promotes atrial fibrillation in mice" (2009) *J Clin Invest* 119:1940-1951; Neef S, et al., "CaMKII-dependent diastolic SR Ca^{2+} leak and elevated diastolic

Ca²⁺ levels in right atrial myocardium of patients with atrial fibrillation" (2010) *Circ Res* 106:1134-1144).

[0088] Pharmacological inhibition of CaMKII is shown to reduce cardiac arrhythmogenesis in vitro and in vivo, including inhibition of torsades that lead to sudden death. (Anderson ME, et al., "KN-93, an inhibitor of multifunctional Ca⁺⁺/Calmodulin-dependent protein kinase, decreases early afterdepolarizations in rabbit heart" (1998) *J Pharmacol Exp Ther* 287:996-1006; Sag CM, et al., "Calcium/calmodulin-dependent protein kinase II contributes to cardiac arrhythmogenesis in heart failure" (2009) *Circ Heart Fail* 2:664-675; Erickson JR, Anderson ME, "CaMKII and Its role in cardiac arrhythmia" (2008) *Journal of Cardiovascular Electrophysiology* 19:1332-1336). CaMKII integrates several proarrhythmic pathways that augment CaMKII activity via Ca²⁺ and reactive oxygen species and which, in turn, acts on the ryanodine receptor, the voltage-dependent calcium channel (Cav1.2), and the Na channel (Nav1.5) to promote arrhythmia (Rokita AG and Anderson ME "New Therapeutic Targets in Cardiology Arrhythmias and Ca²⁺/Calmodulin-Dependent Kinase II (CaMKII)" (2012) *Circulation* 126:2125-2139).

[0089] Studies implicate CaMKII in heart failure and structural heart disease in both mouse models and human heart tissue. Pharmacological and genetic-based inhibition of CaMKII was shown to protect cellular mechanical function and preserves calcium homeostasis after myocardial infarction. CaMKII is increased in cells from failing human heart cells and its pharmacological inhibition improves contractility by an established CaMKII pathway. (Schulman H, Anderson ME, "Ca/Calmodulin-dependent Protein Kinase II in Heart Failure" (2010) *Drug Discovery Today: Disease Mechanisms* 7:e117-e122; Zhang R, et al., "Calmodulin kinase II inhibition protects against structural heart disease" (2005) *Nat Med* 11:409-417; Sossalla S, et al., "Inhibition of Elevated Ca²⁺/Calmodulin-Dependent Protein Kinase II Improves Contractility in Human Failing Myocardium" (2010) *Circulation Research* 107:1150-1161).

[0090] Genetic activation and pharmacological inhibition of CaMKII were used to show that CaMKII mediates cardiac hypertrophy while genetic deletion of δ-CaMKII protected the heart from pathological cardiac hypertrophy and remodeling after pressure overload. (Backs J, et al., "The delta isoform of CaM kinase II is required for pathological cardiac hypertrophy and remodeling after pressure overload" (2009) *Proc Natl Acad Sci USA* 106:2342-2347; Zhang T, et

al., “The cardiac-specific nuclear delta(B) isoform of Ca^{2+} /calmodulin-dependent protein kinase II induces hypertrophy and dilated cardiomyopathy associated with increased protein phosphatase 2A activity” (2002) *J Biol Chem* 277:1261-1267; Anderson ME, et al., “CaMKII in myocardial hypertrophy and heart failure” (2011) *Journal of Molecular and Cellular Cardiology* 51:468-473).

[0091] CaMKII inhibition is found to be effective in several forms of cardioprotection, including from cardiotoxicity caused by cancer therapy (doxorubicin), following heart attack or ischemia-reperfusion, e.g. for the case of acute intervention for heart attack (primary angioplasty), and for patients with mutations leading to sudden death, such as catecholaminergic polymorphic ventricular tachycardia. (Sag CM, et al., “CaMKII-dependent SR Ca leak contributes to doxorubicin-induced impaired Ca handling in isolated cardiac myocytes” (2011) *Journal of Molecular and Cellular Cardiology* 51:749-759; Zhang R, et al., “Calmodulin kinase II inhibition protects against structural heart disease” (2005) *Nat Med* 11:409-417; Liu N, et al., “Calmodulin kinase II inhibition prevents arrhythmias in RyR2(R4496C+/-) mice with catecholaminergic polymorphic ventricular tachycardia” (2011) *Journal of Molecular and Cellular Cardiology* 50:214-222; Joiner, M-L A, et al., “CaMKII determines mitochondrial stress responses in heart” (2012). *Nat Med*, DOI: 10.1038/nature11444, published online October 10, 2012).

[0092] Atherosclerosis pathology includes both the constriction of the vasculature as well as plaque disruption. CaMKII inhibition blocks proliferation of vascular cells as well as mediating ER stressors that lead to apoptosis that underlie plaque disruption. (Timmins JM, et al., “Calcium/calmodulin-dependent protein kinase II links ER stress with Fas and mitochondrial apoptosis pathways” (2009) *The Journal of Clinical Investigation* 119:2925-2941; Li W, et al., “The multifunctional Ca^{2+} /calmodulin-dependent kinase II δ (CaMKII δ) controls neointima formation after carotid ligation and vascular smooth muscle cell proliferation through cell cycle regulation by p21” (2011) *J Biol Chem* 286:7990-7999).

[0093] Studies suggest that CaMKII is an important, but previously unrecognized pro-asthmatic signal, linking the pro-oxidant environment of the asthmatic airways with downstream inflammatory and remodeling events. CaMKII activity in the epithelium may be required for enhancing eosinophilic recruitment to the lung, through a ROS-CaMKII-eotaxin-1 dependent pathway. Inhibition of CaMKII activity may be a novel target in future asthma therapies.

(Sanders PN, et al., “Camkii As A Pro-Asthmatic Signal” (2011) *Am J Respir Care Med* 183:A2795, May 6, 2011 *poster presentation*).

[0094] Smooth muscle proliferation contributes to vascular remodeling and obstructive vasculopathies such as atherosclerosis and restenosis following percutaneous coronary interventions and inhibition of the kinase blocks vascular smooth muscle proliferation and neointimal formation that lead to restenosis. (Li W, et al., “The multifunctional Ca^{2+} /calmodulin-dependent kinase II δ (CaMKII δ) controls neointima formation after carotid ligation and vascular smooth muscle cell proliferation through cell cycle regulation by p21” (2011) *J Biol Chem* 286:7990–7999; House SJ, Singer HA, “CaMKII-delta isoform regulation of neointima formation after vascular injury” (2008) *Arterioscler Thromb Vasc Biol* 28:441-447).

[0095] Pharmacological and genetic suppression of CaMKII has been used to demonstrate a reduction in central and peripheral pain due to injury or inflammation as well as in sensitization to pain. (Zeitz KP, et al., “The contribution of autophosphorylated alpha-calcium-calmodulin kinase II to injury-induced persistent pain” (2004) *Neuroscience* 128:889-898; Luo F, et al., “Reversal of chronic inflammatory pain by acute inhibition of Ca^{2+} /calmodulin-dependent protein kinase II” (2008) *J Pharmacol Exp Ther* 325:267-275; Chen Y, et al., “ Ca^{2+} /Calmodulin-dependent protein kinase II α is required for the initiation and maintenance of opioid-induced hyperalgesia” (2010) *J Neurosci* 30:38-46; Crown ED, et al., “Calcium/calmodulin dependent kinase II contributes to persistent central neuropathic pain following spinal cord injury” (2012) *Pain* 153:710-721).

[0096] Inhibition of CaMKII is neuroprotective, reducing damage due to hypoxia in stroke models. Reduction of atrial fibrillation by Inhibition of CaMKII would also reduce stroke incidence. (Vest RS, et al., “Effective post-insult neuroprotection by a novel CaMKII inhibitor” (2010) *J Biol Chem* 285:20675–20682; Ashpole NM, et al., “Calcium/Calmodulin-dependent Protein Kinase II (CaMKII) Inhibition Induces Neurotoxicity via Dysregulation of Glutamate/Calcium Signaling and Hyperexcitability” (2012) *Journal of Biological Chemistry* 287:8495-8506; Dobrev D, et al., “Novel molecular targets for atrial fibrillation therapy” (2012) *Nature Reviews Drug Discovery* 11:275-291).

[0097] Stimulation of opiate receptors increases CaMKII and leads to tolerance and dependence that are reduced by inhibition of CaMKII. (Liang D, et al., “Increased expression of Ca^{2+} /calmodulin-dependent protein kinase II alpha during chronic morphine exposure” (2004)

Neuroscience 123:769-775; Fan GH, *et al.*, “Inhibition of calcium/calmodulin-dependent protein kinase II in rat hippocampus attenuates morphine tolerance and dependence” (1999) *Mol Pharmacol* 56:39-45).

[0098] Inhibition of CaMKII reduces the VEGF pathway that mediates increased vascularization or angiogenesis of retinal endothelial cells. (Banumathi E, *et al.*, “VEGF-induced retinal angiogenic signalling is critically dependent on Ca^{2+} signalling via Ca^{2+} /calmodulin-dependent protein kinase II” (2011) *Investigative Ophthalmology & Visual Science* 52:3103-3111).

[0099] CaMKII may have several sites of action that support the notion of CaMKII inhibition in type II diabetes. CaMKII modulates insulin signaling that suggest a role in the pathogenesis of insulin resistance. In liver, CaMKII regulates glucose production and its inhibition would be beneficial in diabetes. (Illario M, *et al.*, “Calcium-calmodulin-dependent kinase II (CaMKII) mediates insulin-stimulated proliferation and glucose uptake” (2009) *Cellular Signalling* 21:786-792; Ozcan L, *et al.*, “Calcium Signaling through CaMKII Regulates Hepatic Glucose Production in Fasting and Obesity” (2012) *Cell Metabolism* 15:739-751).

[00100] Studies show that pharmacological inhibition of CaMKII reduces proliferation of osteosarcoma cell lines and indicates changes in signal transduction related to growth. The inhibitor administered to mice with a human osteosarcoma xenograft markedly decreases tumor size. (Yuan K, *et al.*, “ α -CaMKII controls the growth of human osteosarcoma by regulating cell cycle progression” (2007) *Lab Invest* 87:938-950).

[00101] Tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) provides a pathway in melanoma therapy but melanoma is often resistant to TRAIL after metastasis. Inhibition of CaMKII signaling by use of a dominant negative form of the kinase was shown to restore the sensitivity of melanoma to cell death via TRAIL. (Xiao C, *et al.*, “Inhibition of CaMKII-mediated c-FLIP expression sensitizes malignant melanoma cells to TRAIL-induced apoptosis” (2005) *Exp Cell Res* 304:244-255).

[00102] Studies have shown that proliferation and invasion of prostate cancer cell lines is reduced by pharmacological inhibition of CaMKII. Kinase inhibition was used to show that it is important for prostate cancer cell survival and promotes their progression to an androgen-independent state. (Mamaeva OA, *et al.*, “Calcium/calmodulin-dependent kinase II regulates notch-1 signaling in prostate cancer cells” (2009) *J Cell Biochem* 106:25-32; Rokhlin OW, *et al.*,

“Calcium/calmodulin-dependent kinase II plays an important role in prostate cancer cell survival” (2007) *Cancer Biol Ther* 6:732-742).

[00103] Pharmacological inhibition of CaMKII reduces differentiation of osteoclasts and suppresses bone resorption characteristic of osteoporosis. (Ang ESM, *et al.*, “Calcium/calmodulin-dependent kinase activity is required for efficient induction of osteoclast differentiation and bone resorption by receptor activator of nuclear factor kappa B ligand (RANKL)” (2007) *Journal of cellular physiology* 212:787-795).

[00104] Pharmacological and genetic suppression of CaMKII demonstrated its role in the production of proinflammatory cytokines and interferon in macrophages. A small molecule inhibitor of CaMKII was used to show that it required for tumour necrosis factor-related apoptosis inducing ligand (TRAIL)-mediated apoptosis of fibroblast-like synovial cells, suggesting that it is a target for rheumatoid arthritis therapy. (Liu X, *et al.*, “CaMKII promotes TLR-triggered proinflammatory cytokine and type I interferon production by directly binding and activating TAK1 and IRF3 in macrophages” (2008) *Blood* 112:4961-4970; Fujikawa K, *et al.*, “Calcium/calmodulin-dependent protein kinase II (CaMKII) regulates tumour necrosis factor-related apoptosis inducing ligand (TRAIL)-mediated apoptosis of fibroblast-like synovial cells (FLS) by phosphorylation of Akt” (2009) *Clinical and experimental rheumatology* 27:952-957).

[00105] A genetic model of CaMKII inhibition as well as a genetic knockout of the CaMKIIdelta show that the heart is protected from fibrosis and structural remodeling normally induced by myocardial infarction and by pressure overload, respectively (Zhang R, *et al.* “Calmodulin kinase II inhibition protects against structural heart disease” (2005) *Nat Med* 11:409-417; Backs J, *et al.* “The delta isoform of CaM kinase II is required for pathological cardiac hypertrophy and remodeling after pressure overload” (2009) *Proc Natl Acad Sci U S A* 106:2342-2347). These conclusions are consistent with the observation that pharmacological inhibition of CaMKII reduces fibroblast proliferation, secretion of fibroblast growth factor β 1, and decreased expression of several metalloproteinases that underlie cardiac hypertrophy and fibrosis (Zhang W, *et al.* “Inhibition of calcium-calmodulin-dependent kinase II suppresses cardiac fibroblast proliferation and extracellular matrix secretion” (2009) *J Cardiovascular Pharmacology* 55:96-105). Fibrosis is also promoted by activation of CaMKII via reactive oxygen, such as during Angiotensin II stimulation. These data support an indication for CaMKII

inhibitors in reducing fibrosis in a number of possible indications in which fibrosis is prominent, including myocardial fibrosis, cardiac hypertrophy, and myocardial infarction.

[00106] The activity of a compound utilized in this invention as an inhibitor of CaMKII or treatment for a CaMKII-mediated disease, disorder or condition, may be assayed *in vitro* or *in vivo*. An *in vivo* assessment of the efficacy of the compounds of the invention may be made using an animal model of a CaMKII-mediated disease, disorder or condition, e.g., a rodent or primate model. Cell-based assays may be performed using, e.g., a cell line isolated from a tissue that expresses CaMKII. Additionally, biochemical or mechanism-based assays, e.g., transcription assays using a purified protein, Northern blot, RT-PCR, etc., may be performed. *In vitro* assays include assays that determine cell morphology, protein expression, and/or the cytotoxicity, enzyme inhibitory activity, and/or the subsequent functional consequences of treatment of cells with compounds of the invention. Alternate or additional *in vitro* assays may be used to quantitate the ability of the inhibitor to bind to protein or nucleic acid molecules within the cell. Inhibitor binding may be measured by radiolabelling the inhibitor prior to binding, isolating the inhibitor/target molecule complex and determining the amount of radiolabel bound. Alternatively or additionally, inhibitor binding may be determined by running a competition experiment where new inhibitors are incubated with purified proteins or nucleic acids bound to known radioligands. Detailed conditions of exemplary systems for assaying a compound utilized in this invention as an inhibitor of CaMKII are set forth in the Examples below. Such assays are exemplary and not intended to limit the scope of the invention. The skilled practitioner can appreciate that modifications can be made to conventional assays to develop equivalent or other assays that can be employed to comparably assess activity or otherwise characterize compounds and/or compositions as described herein.

[00107] As used herein, the terms “treatment,” “treat,” and “treating” refer to reversing, alleviating, delaying the onset of, reducing incidence or severity, or inhibiting the progress of a disease, disorder or condition, or one or more symptoms thereof, as described herein. In some embodiments, treatment may be administered after one or more symptoms have developed. In other embodiments, treatment may be administered in the absence of symptoms. For example, treatment may be administered to a susceptible individual prior to the onset of symptoms (e.g., in light of a history of symptoms and/or in light of genetic or other susceptibility factors).

Treatment may also be continued after symptoms have resolved, for example to prevent or delay their recurrence.

[00108] Compounds and/or compositions described herein may be administered using any amount and any route of administration effective for treating a disease, disorder, or condition. In some embodiments, compounds and/or compositions are administered in an amount and/or by a route effective for treating a cardiovascular disease, disorder or condition, an inflammatory disease, disorder or condition, a neurological disease, disorder or condition, an ocular disease, disorder or condition, a metabolic disease, disorder or condition, a cancer or other proliferative disease, disorder or condition, a bone disease, disorder or condition, or an addictive disease, disorder, or condition.

[00109] In some embodiments, compounds and/or compositions described herein may be administered using any amount and any route of administration effective for treating or lessening the severity of a disease, disorder or condition associated with CaMKII.

[00110] In some embodiments, compounds and/or compositions may be administered using any amount and any route of administration effective for treating a cardiovascular disease, disorder, or condition. In some embodiments, the cardiovascular disease, disorder or condition is a disease of the heart. In some embodiments, the cardiovascular disease, disorder or condition is a disease of the vasculature. In some embodiments, the cardiovascular disease, disorder or condition is selected from atrial fibrillation, ventricular arrhythmia, heart failure, cardiac hypertrophy, atherosclerosis, fibrosis, or restenosis. In some embodiments the restenosis is in-stent restenosis in coronary artery disease.

[00111] In some embodiments, provided compounds and/or compositions may be administered using any amount and any route of administration effective for achieving cardioprotection from cardiotoxicity. In some embodiments, the cardiotoxicity avoided by administration of the compounds and compositions of the invention is due to drug therapy, heart attack, ischemia-reperfusion injury, or mutations leading to sudden death such as catecholaminergic polymorphic ventricular tachycardia.

[00112] In some embodiments, the compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for treating an inflammatory disease, disorder or condition. In some embodiments, the inflammatory disease, disorder or condition is asthma or rheumatoid arthritis.

[00113] In some embodiments, the compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for treating a neurological disease, disorder or condition. In some embodiments, the neurological disease, disorder or condition is pain or stroke.

[00114] In some embodiments, the compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for treating an addictive disease, disorder or condition. In some embodiments the addictive disease, disorder, or condition is opioid tolerance or dependence.

[00115] In some embodiments, the compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for treating an ocular disease, disorder or condition. In some embodiments, the ocular disease, disorder or condition is macular degeneration.

[00116] In some embodiments, the compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for treating a metabolic disease, disorder or condition. In some embodiments, the metabolic disease, disorder or condition is diabetes. In some embodiments, the diabetes is type II diabetes.

[00117] In some embodiments, the compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for treating a cancer or another proliferative disease, disorder or condition. In some embodiments, the cancer or other proliferative disease, disorder or condition is an osteosarcoma, a melanoma, or a prostate cancer.

[00118] In some embodiments, the compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for treating a bone disease, disorder or condition. In some embodiments, the bone disease, disorder or condition is osteoporosis.

[00119] It will be appreciated by those skilled in the art that the exact amount of a provided compound or composition may vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the infection, the particular agent, its mode of administration, and the like.

[00120] In some embodiments, compounds of the invention are formulated in dosage unit form, for example for ease of administration and uniformity of dosage. The expression "dosage unit form" or "unit dosage" as used herein refers to a physically discrete unit of agent appropriate for the patient to be treated. It will be understood, however, that total daily usage of the compounds and compositions of the present invention may be decided by the attending physician within the scope of sound medical judgment. The specific effective dose level for any particular patient or organism may depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed, and like factors well known in the medical arts.

[00121] According to some embodiments, the invention relates to a method of inhibiting CaMKII in a biological sample comprising the step of contacting said biological sample with a compound of this invention, or a composition comprising said compound.

[00122] The term "biological sample", as used herein, includes, without limitation, cell cultures or extracts thereof; biopsied material obtained from a mammal or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof.

[00123] Inhibition of enzymes in a biological sample is useful for a variety of purposes that are known to one of skill in the art. Examples of such purposes include, but are not limited to biological assays, gene expression studies, and biological target identification.

[00124] Some embodiments of the present invention relate to a method of inhibiting CaMKII in a patient comprising the step of administering to said patient a compound of the present invention, or a composition comprising said compound.

[00125] In some embodiments, the invention relates to a method of inhibiting CaMKII activity in a patient comprising the step of administering to said patient a compound of the present invention, or a composition comprising said compound. In certain embodiments, the present invention provides a method for treating a disease, disorder or condition mediated by CaMKII, in a patient in need thereof, comprising the step of administering to said patient a compound according to the present invention or pharmaceutically acceptable composition thereof. Such diseases, disorders and conditions are described in detail herein.

[00126] In some embodiments compounds and/or compositions of the present invention may be used in a method of treating a cardiovascular disease, disorder, or condition, an inflammatory disease, disorder or condition, a neurological disease, disorder or condition, an ocular disease, disorder or condition, a metabolic disease, disorder or condition, a cancer or other proliferative disease, disorder or condition, or a bone disease, disorder or condition. In certain embodiments the compounds and compositions of the present invention may be used to treat a cardiovascular disease, disorder or condition, an inflammatory disease, disorder or condition, a neurological disease, disorder or condition, an ocular disease, disorder or condition, a metabolic disease, disorder or condition, a cancer or other proliferative disease, disorder or condition, or a bone disease, disorder or condition in a mammal. In certain embodiments the mammal is a human patient.

[00127] In some embodiments the present invention provides a method of treating a cardiovascular disease, disorder or condition, an inflammatory disease, disorder or condition, a neurological disease, disorder or condition, an ocular disease, disorder or condition, a metabolic disease, disorder or condition, a cancer or other proliferative disease, disorder or condition, or a bone disease, disorder or condition, comprising administering a compound or composition of the present invention to a patient in need thereof. In certain embodiments the method of treating a cardiovascular disease, disorder or condition, an inflammatory disease, disorder or condition, a neurological disease, disorder or condition, an ocular disease, disorder or condition, a metabolic disease, disorder or condition, a cancer or other proliferative disease, disorder or condition, or a bone disease, disorder or condition comprises administering compounds and compositions of the present invention to a mammal. In certain embodiments the mammal is a human.

[00128] In certain embodiments, the present invention provides a method of treating a cardiovascular disease, disorder or condition, comprising administering a compound or composition of the present invention to a patient with a cardiovascular disease, disorder or condition. In certain embodiments, the method of treating a cardiovascular disease, disorder or condition comprises administering compounds and compositions of the present invention to a mammal. In certain embodiments, the mammal is a human.

[00129] In certain embodiments, the present invention provides a method of treating a cancer or another proliferative disease, disorder or condition, comprising administering a compound or composition of the present invention to a patient with a cancer or another proliferative disease,

disorder or condition. In certain embodiments, the method of treating a cancer or other proliferative disorder comprises administering compounds and compositions of the present invention to a mammal. In certain embodiments, the mammal is a human.

[00130] As used herein, the terms “treating a cancer” refers to the inhibition of the growth, division, maturation or viability of cancer cells, and/or causing the death of cancer cells, individually or in aggregate with other cancer cells, by cytotoxicity, nutrient depletion, or the induction of apoptosis.

[00131] Examples of tissues containing cancerous cells whose proliferation is inhibited by the compounds and compositions described herein and against which the methods described herein are useful include but are not limited to breast, prostate, brain, blood, bone marrow, bone, liver, pancreas, skin, kidney, colon, ovary, lung, testicle, penis, thyroid, parathyroid, pituitary, thymus, retina, uvea, conjunctiva, spleen, head, neck, trachea, gall bladder, rectum, salivary gland, adrenal gland, throat, esophagus, lymph nodes, sweat glands, sebaceous glands, muscle, heart, and stomach.

[00132] In some embodiments, the cancer treated by compounds or compositions of the invention is a skin cancer, lung cancer, breast cancer, prostate cancer, leukemia, kidney cancer, esophageal cancer, brain cancer, bone cancer or colon cancer. In some embodiments, the cancer treated by the compounds or compositions of the invention is an osteosarcoma, a melanoma or a prostate cancer.

[00133] In certain embodiments, the present invention provides a method of treating a neurological disease, disorder or condition, comprising administering a compound or composition of the present invention to a patient with a neurological disease, disorder or condition. In certain embodiments, the method of treating a neurological disease, disorder or condition comprises administering compounds and compositions of the present invention to a mammal. In certain embodiments, the mammal is a human. In certain embodiments, the neurological disease, disorder or condition is pain or stroke.

[00134] In certain embodiments, the present invention provides a method of treating an inflammatory disease, disorder or condition, comprising administering a compound or composition of the present invention to a patient with an inflammatory disease, disorder or condition. In certain embodiments, the method of treating an inflammatory disease, disorder or condition comprises administering compounds and compositions of the present invention to a

mammal. In certain embodiments, the mammal is a human. In certain embodiments, the neurological disease, disorder or condition is asthma or rheumatoid arthritis.

[00135] In certain embodiments, the present invention provides a method of treating a metabolic disease, disorder or condition, comprising administering a compound or composition of the present invention to a patient with a metabolic disease, disorder or condition. In certain embodiments, the method of treating a metabolic disease, disorder or condition comprises administering compounds and compositions of the present invention to a mammal. In certain embodiments, the mammal is a human. In certain embodiments, the metabolic disease, disorder or condition is diabetes. In some embodiments, the diabetes is type II diabetes.

[00136] In certain embodiments, the present invention provides a method of treating opioid tolerance or dependence, comprising administering a compound or composition of the present invention to an opioid tolerant or dependent patient. In certain embodiments, the method of treating opioid tolerance or dependence comprises administering compounds and compositions of the present invention to a human. In some embodiments the opioid tolerance or dependence is morphine tolerance or dependence.

[00137] In certain embodiments, the present invention provides a method of treating an ocular disease, disorder or condition, comprising administering a compound or composition of the present invention to a patient with an ocular disease, disorder or condition. In certain embodiments, the method of treating an ocular disease, disorder or condition comprises administering compounds and compositions of the present invention to a mammal. In certain embodiments, the mammal is a human. In certain embodiments, the ocular disease, disorder or condition is macular degeneration.

[00138] Depending upon the particular disease, disorder or condition to be treated, additional therapeutic agents, which are normally administered to treat that condition, may be administered in combination with compounds and compositions of this invention. As used herein, additional therapeutic agents that are normally administered to treat a particular disease, or condition, are known as "appropriate for the disease, or condition, being treated".

[00139] In certain embodiments, a provided compound, or composition thereof, is administered in combination with another inhibitor of CaMKII. In some embodiments, a provided compound, or composition thereof, is administered in combination with one or more

other therapeutic agents. Such CaMKII inhibitors include, but are not limited to CaM Kinase II Calmodulin Antagonist peptide, KN-93, and lavendustin C.

[00140] In certain embodiments, a provided compound, or a composition thereof, is administered in combination with another anti-cancer, cytotoxin, or chemotherapeutic agent.

[00141] In certain embodiments, the anti-cancer or chemotherapeutic agents used in combination with compounds or compositions of the invention include, but are not limited to imatinib, nilotinib, gefitinib, sunitinib, carfilzomib, salinosporamide A, retinoic acid, cisplatin, carboplatin, oxaliplatin, mechlorethamine, cyclophosphamide, chlorambucil, ifosfamide, azathioprine, mercaptopurine, doxifluridine, fluorouracil, gemcitabine, methotrexate, tioguanine, vincristine, vinblastine, vinorelbine, vindesine, podophyllotoxin, etoposide, teniposide, tafluposide, paclitaxel, docetaxel, irinotecan, topotecan, amsacrine, actinomycin, doxorubicin, daunorubicin, valrubicin, idarubicin, epirubicin, plicamycin, mitomycin, mitoxantrone, melphalan, busulfan, capecitabine, pemetrexed, epothilones, 13-cis-Retinoic Acid, 2-CdA, 2-Chlorodeoxyadenosine, 5-Azacitidine, 5-Fluorouracil, 5-FU, 6-Mercaptopurine, 6-MP, 6-TG, 6-Thioguanine, Abraxane, Accutane ®, Actinomycin-D, Adriamycin ®, Adrucil ®, Afinitor ®, Agrylin ®, Ala-Cort ®, Aldesleukin, Alemtuzumab, ALIMTA, Alitretinoin, Alkaban-AQ ®, Alkeran ®, All-transretinoic Acid, Alpha Interferon, Altretamine, Amethopterin, Amifostine, Aminoglutethimide, Anagrelide, Anandron ®, Anastrozole, Arabinosylcytosine, Ara-C, Aranesp ®, Aredia ®, Arimidex ®, Aromasin ®, Arranon ®, Arsenic Trioxide, Arzerra™, Asparaginase, ATRA, Avastin ®, Azacitidine, BCG, BCNU, Bendamustine, Bevacizumab, Bexarotene, BEXXAR ®, Bicalutamide, BiCNU, Blenoxane ®, Bleomycin, Bortezomib, Busulfan, Busulfex ®, C225, Calcium Leucovorin, Campath ®, Camptosar ®, Camptothecin-11, Capecitabine, Carac™, Carboplatin, Carmustine, Carmustine Wafer, Casodex ®, CC-5013, CCI-779, CCNU, CDDP, CeeNU, Cerubidine ®, Cetuximab, Chlorambucil, Citrovorum Factor, Cladribine, Cortisone, Cosmegen ®, CPT-11, Cytadren ®, Cytosar-U ®, Cytoxan ®, Dacarbazine, Dacogen, Dactinomycin, Darbepoetin Alfa, Dasatinib, Daunomycin, Daunorubicin Hydrochloride, Daunorubicin Liposomal, DaunoXome ®, Decadron, Decitabine, Delta-Cortef ®, Deltasone ®, Denileukin, Diftitox, DepoCyt™, Dexamethasone, Dexamethasone Acetate, Dexamethasone Sodium Phosphate, Dexasone, Dexrazoxane, DHAD, DIC, Diodex, Docetaxel, Doxil ®, Doxorubicin, Doxorubicin Liposomal, Droxia™, DTIC, DTIC-Dome ®, Duralone ®, Efudex ®, Eligard™, Ellence™, Eloxatin™, Elspar ®, Emcyt ®, Epirubicin, Epoetin Alfa, Erbitux,

Erlotinib, Erwinia L-asparaginase, Estramustine, Ethyol, Etopophos ®, Etoposide, Etoposide Phosphate, Eulexin ®, Everolimus, Evista ®, Exemestane, Fareston ®, Faslodex ®, Femara ®, Filgrastim, Floxuridine, Fludara ®, Fludarabine, Fluoroplex ®, Fluorouracil, Fluorouracil (cream), Fluoxymesterone, Flutamide, Folinic Acid, FUDR ®, Fulvestrant, G-CSF, Gefitinib, Gemcitabine, Gemtuzumab, ozogamicin, ,Gemzar Gleevec ™, Gliadel ® Wafer, GM-CSF, Goserelin, Granulocyte - Colony Stimulating Factor, Granulocyte Macrophage Colony Stimulating Factor, Halotestin ®, Herceptin ®, Hexadrol, Hexalen ®, Hexamethylmelamine, HMM, Hycamtin ®, Hydrea ®, Hydrocort Acetate ®, Hydrocortisone, Hydrocortisone Sodium Phosphate, Hydrocortisone Sodium Succinate, Hydrocortone Phosphate, Hydroxyurea, Ibrutumomab, Ibrutumomab, Tiuxetan, Idamycin ®, Idarubicin Ifex ®, IFN-alpha, Ifosfamide, IL-11, IL-2, Imatinib mesylate, Imidazole Carboxamide, Interferon alfa, Interferon Alfa-2b (PEG Conjugate), Interleukin-2, Interleukin-11, Intron A® (interferon alfa-2b), Iressa ®, Irinotecan, Isotretinoin, Ixabepilone, Ixempra ™, Kidrolase ®, Lanacort ®, Lapatinib, L-asparaginase, LCR, Lenalidomide, Letrozole, Leucovorin, Leukeran, Leukine ™, Leuprolide, Leurocristine, Leustatin ™, Liposomal Ara-C, Liquid Pred ®, Lomustine, L-PAM, L-Sarcolysin, Lupron ®, Lupron Depot ®, Matulane ®, Maxidex, Mechlorethamine, Mechlorethamine Hydrochloride, Medralone ®, Medrol ®, Megace ®, Megestrol, Megestrol Acetate, Melphalan, Mercaptopurine, Mesna, Mesnex ™, Methotrexate, Methotrexate Sodium, Methylprednisolone, Meticorten ®, Mitomycin, Mitomycin-C, Mitoxantrone, M-Prednisol ®, MTC, MTX, Mustargen ®, Mustine, Mutamycin ®, Myleran ®, Mylocel ™, Mylotarg ®, Navelbine ®, Nelarabine, Neosar ®, Neulasta ™, Neumega ®, Neupogen ®, Nexavar ®, Nilandron ®, Nilotinib, Nilutamide, Nipent ®, Nitrogen Mustard, Novaldex ®, Novantrone ®, Nplate, Octreotide, Octreotide acetate, Ofatumumab, Oncospar ®, Oncovin ®, Ontak ®, Onxal ™, Oprelvekin, Orapred ®, Orasone ®, Oxaliplatin, Paclitaxel, Paclitaxel Protein-bound, Pamidronate, Panitumumab, Panretin ®, Paraplatin ®, Pazopanib, Pediapred ®, PEG Interferon, Pegaspargase, Pegfilgrastim, PEG-INTRON ™, PEG-L-asparaginase, PEMETREXED, Pentostatin, Phenylalanine Mustard, Platinol ®, Platinol-AQ ®, Prednisolone, Prednisone, Prelone ®, Procarbazine, PROCRIT ®, Proleukin ®, Prolifeprospan 20 with Carmustine Implant, Purinethol ®, Raloxifene, Revlimid ®, Rheumatrex ®, Rituxan ®, Rituximab, Roferon-A ® (Interferon Alfa-2a), Romiplostim, Rubex ®, Rubidomycin hydrochloride, Sandostatin ®, Sandostatin LAR ®, Sargramostim, Solu-Cortef ®, Solu-Medrol ®, Sorafenib, SPRYCEL ™, STI-571, Streptozocin, SU11248, Sunitinib, Sutent

®, Tamoxifen, Tarceva ®, Targretin ®, Tasigna ®, Taxol ®, Taxotere ®, Temodar ®, Temozolomide, Temsirolimus, Teniposide, TESPA, Thalidomide, Thalomid ®, TheraCys ®, Thioguanine, Thioguanine Tabloid ®, Thiophosphoamide, Thioplex ®, Thiotepa, TICE ®, Toposar ®, Topotecan, Toremifene, Torisel ®, Tositumomab, Trastuzumab, Treanda ®, Tretinoin, Trexall ™, Trisenox ®, TSPA, TYKERB ®, VCR, Vectibix ™, Velban ®, Velcade ®, VePesid ®, Vesanoid ®, Viadur ™, Vidaza ®, Vinblastine, Vinblastine Sulfate, Vincasar Pfs ®, Vincristine, Vinorelbine, Vinorelbine tartrate, VLB, VM-26, Vorinostat, Votrient, VP-16, Vumon ®, Xeloda ®, Zanosar ®, Zevalin ™, Zinecard ®, Zoladex ®, Zoledronic acid, Zolinza, Zometa ®, or combinations of any of the above.

[00142] In certain embodiments, a combination of 2 or more therapeutic agents may be administered together with compounds of the invention. In certain embodiments, a combination of 3 or more therapeutic agents may be administered with compounds of the invention.

[00143] Other examples of agents the inhibitors of this invention may also be combined with include, without limitation: vitamins and nutritional supplements, cancer vaccines, treatments for neutropenia (e.g. G-CSF, filgrastim, lenograstim), treatments for thrombocytopenia (e.g. blood transfusion, erythropoietin), antiemetics (e.g. 5-HT₃ receptor antagonists, dopamine antagonists, NK1 receptor antagonists, histamine receptor antagonists, cannabinoids, benzodiazepines, or anticholinergics), treatments for Alzheimer's Disease such as Aricept® and Excelon®; treatments for Parkinson's Disease such as L-DOPA/carbidopa, entacapone, ropinrole, pramipexole, bromocriptine, pergolide, trihexyphenidyl, and amantadine; agents for treating Multiple Sclerosis (MS) such as beta interferon (e.g., Avonex® and Rebif®), Copaxone®, and mitoxantrone; treatments for asthma such as albuterol and Singulair®; agents for treating schizophrenia such as zyprexa, risperdal, seroquel, and haloperidol; anti-inflammatory agents such as corticosteroids, TNF blockers, IL-1 RA, azathioprine, cyclophosphamide, and sulfasalazine; immunomodulatory and immunosuppressive agents such as cyclosporin, tacrolimus, rapamycin, mycophenolate mofetil, interferons, corticosteroids, cyclophosphamide, azathioprine, and sulfasalazine; neurotrophic factors such as acetylcholinesterase inhibitors, MAO inhibitors, interferons, anti-convulsants, ion channel blockers, riluzole, and anti-Parkinsonian agents; agents for treating cardiovascular disease such as beta-blockers, ACE inhibitors, diuretics, nitrates, calcium channel blockers, and statins, fibrates, cholesterol absorption inhibitors, bile acid sequestrants, and niacin; agents for treating liver disease such as corticosteroids, cholestyramine, interferons, and

anti-viral agents; agents for treating blood disorders such as corticosteroids, anti-leukemic agents, and growth factors; agents for treating immunodeficiency disorders such as gamma globulin; and anti-diabetic agents such as biguanides (metformin, phenformin, buformin), thiazolidinediones (rosiglitazone, pioglitazone, troglitazone), sulfonylureas (tolbutamide, acetohexamide, tolazamide, chlorpropamide, glipizide, glyburide, glimepiride, gliclazide), meglitinides (repaglinide, nateglinide), alpha-glucosidase inhibitors (miglitol, acarbose), incretin mimetics (exenatide, liraglutide, taspoglutide), gastric inhibitory peptide analogs, DPP-4 inhibitors (vildagliptin, sitagliptin, saxagliptin, linagliptin, alogliptin), amylin analogs (pramlintide), and insulin and insulin analogs.

[00144] In certain embodiments, compounds of the present invention, or a pharmaceutically acceptable composition thereof, are administered in combination with antisense agents, a monoclonal or polyclonal antibody or an siRNA therapeutic.

[00145] Those additional agents may be administered separately from an inventive compound-containing composition, as part of a multiple dosage regimen. Alternatively or in addition to those additional agents administered separately, those agents may be part of a single dosage form, mixed together with a compound of this invention in a single composition. If administered as part of a multiple dosage regime, the two active agents may be submitted simultaneously, sequentially or within a period of time from one another, normally within five hours from one another.

[00146] As used herein, the term “combination,” “combined,” and related terms refers to the simultaneous or sequential administration of therapeutic agents in accordance with this invention. For example, a compound of the present invention may be administered with another therapeutic agent simultaneously or sequentially in separate unit dosage forms or together in a single unit dosage form. Accordingly, the present invention provides a single unit dosage form comprising a compound of formula **I**, **I-a**, **II**, **III**, **IV**, **V**, **VI**, **VII**, **VIII**, or **IX**, an additional therapeutic agent, and a pharmaceutically acceptable carrier, adjuvant, or vehicle.

[00147] The amount of both, an inventive compound and additional therapeutic agent (in those compositions which comprise an additional therapeutic agent as described above) that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Preferably, compositions of this

invention should be formulated so that a dosage of between 0.01 - 100 mg/kg body weight/day of an inventive can be administered.

[00148] In those compositions which comprise an additional therapeutic agent, that additional therapeutic agent and the compound of this invention may act synergistically. Therefore, the amount of additional therapeutic agent in such compositions will be less than that required in a monotherapy utilizing only that therapeutic agent. In such compositions a dosage of between 0.01 - 100 μ g/kg body weight/day of the additional therapeutic agent can be administered.


[00149] The amount of additional therapeutic agent present in the compositions of this invention will be no more than the amount that would normally be administered in a composition comprising that therapeutic agent as the only active agent. Preferably the amount of additional therapeutic agent in the presently disclosed compositions will range from about 50% to 100% of the amount normally present in a composition comprising that agent as the only therapeutically active agent.

EXEMPLIFICATION

[00150] As depicted in the Examples below, in certain exemplary embodiments, compounds are prepared according to the following general procedures. It will be appreciated that, although the general methods depict the synthesis of certain compounds of the present invention, the following general methods, and other methods known to one of ordinary skill in the art, can be applied to all compounds and subclasses and species of each of these compounds, as described herein.

[00151] In certain embodiments, compounds of formula I-a are prepared according to the procedure outlined in Scheme 1.

Scheme 1




[00152] Depending on the nature of the functional groups on specific instances of R^2 and R^3' , functional groups on those substituents were modified by protecting groups according to standard procedures known in the art prior to the Suzuki coupling step, and an optional deprotection step was used to remove those protecting groups subsequent to the Suzuki coupling. In some embodiments, primary amine functional groups on substituents R^2' and R^3' may be protected, prior to the coupling step, thereby forming modified groups $R^2\text{-P}$ and $R^3\text{-P}$ respectively, wherein P is a protecting group suitable for primary amines. In some embodiments, protecting group P is a carbamate (e.g. Cbz, BOC, methylcarbamate, ethylcarbamate). In some embodiments, protecting group P is a BOC (t-butoxycarbonyl) group. Methods for the introduction and removal of such primary amine protecting groups are well known in the art, as exemplified by “Greene’s Protective Groups in Organic Synthesis” Wuts PGM, Greene, TW, 4th Ed., 2006, Wiley-Interscience, the entirety of which is incorporated herein by reference. In certain embodiments, compounds and synthetic intermediates thereof of the present invention may be prepared according to methods known to one of ordinary skill in the art. One of ordinary skill in the art will appreciate that compounds of formula I bearing additional substituents R^4' , R^5' , R^6' , R^5 , and R^8 , may be prepared according to similar procedures to those described above for formula I-a.

[00153] The following examples illustrate methods utilized for the preparation of structures relevant to the present invention. Throughout these examples, certain equipment, HPLC columns and solvent systems were utilized in the execution of reactions and the purification of reaction products. Accordingly, microwave reactions were carried out utilizing an AntonPar, Monowave 300 microwave reactor. Preparative HPLC purifications were carried out utilizing a Shimadzu [Prominence LC-20AP], equipped with a Discovery C-18 column (50 x 21.2mm, 5 μ) utilizing the following method: Solvent A = Acetonitrile, Solvent B = Water; Gradient = 95% solvent B to 10% solvent B over 20 min with a flow rate of 10 mL/min. Analytical LCMS data were acquired using a Shimadzu [LCMS-2020] equipped with a SHIMPAK, XR ODS-II column (50 x 2mm) utilizing the following method: Flow Rate = 0.2 mL/min, Solvent A = Acetonitrile, Solvent B = 0.1% TFA in water; Gradient = Initial 95% of solvent B to 10% solvent B over 10 min followed by 10% solvent B for an additional 10 min.

Example 1

[00154] An exemplary procedure for the preparation of compound 9 (I-7) follows.



[00155] Synthesis of 1-(pyridin-2-yl)-1H-benzo[d][1,2,3]triazole (3):

[00156] A suspension of 1H-benzo-[1,2,3] triazole (1, 40 g, 335 mmol) and 2-bromopyridine (2, 105 g, 671 mmol) in toluene (160 mL) was heated at reflux for 18 h after which, the reaction mixture was poured into EtOAc (1 L). The resulting white solid precipitate was dissolved by addition of aqueous KOH (10%, 85 mL). The phases were separated, and the organic layer was washed with aqueous KOH (10%, 2 X 250 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated to dryness. The isolated solids were recrystallized from CH₃OH giving a 62% yield of compound 3. ¹H NMR (500 MHz, CDCl₃) δ ppm: 8.65 (d, 1H), 8.62 (d, 1H), 8.31 (d, 1H), 8.11 (d, 1H), 7.94 (m, 1H), 7.59 (t, 1H), 7.46 (t, 1H), 7.33 (m, 1H); Mass (m/z): 197.2 (M+H).

[00157] Synthesis of 9H-pyrido[2,3-b]indole (4):

[00158] To compound **3** (40 g, 203 mmol) was added polyphosphoric acid (160 g) pre-heated to 160°C. Once gas evolution was complete, H₂O (900 mL) was added and the pH of the solution was adjusted >10 by addition of aqueous NaOH (10M). The mixture was then sonicated at 50°C until the reaction mass was fully suspended. The suspension was poured into H₂O (500 mL) and cooled to room temperature. After 20 min, the resulting solid was collected by filtration, washed with H₂O (2 X 150 mL) and dried under vacuum giving a 35% yield of crude compound **4**. ¹H NMR (500 MHz, DMSO-d₆) δ ppm: 11.72 (br, 1H), 8.44 (d, 1H), 8.37 (d, 1H), 8.11 (d, 1H), 7.47 (dd, 1H), 7.38-7.44 (m, 1H), 7.16-7.20 (m, 1H), 7.15 (d, 1H); Mass (m/z): 169.2 (M+H)

[00159] Synthesis of 9H-pyrido[2,3-b]indole 1-oxide (5):

[00160] Aqueous H₂O₂ (35%, 2.8 g, 83 mmol) was added dropwise to a solution of crude compound **4** (2 g, 11.9 mmol) in CH₃COOH (17 mL). The reaction mixture was refluxed for 4 h. Additional aqueous H₂O₂ (35%, 1 mL) was added dropwise and refluxing was continued for an additional 2 h. The solvent was then removed under vacuum and pH of the oily residue was adjusted to 8 on treatment with saturated aqueous K₂CO₃. The resulting solution was stirred overnight. The resulting solid was collected by filtration, washed with H₂O and dried under giving a 67% yield of compound **5**. ¹H NMR (500 MHz, DMSO-d₆) δ ppm: 12.58 (br, 1H), 8.34(d, 1H), 8.20 (dd, 2H), 7.56 (m, 2H), 7.30 (t, 1H), 7.23 (t, 1H). Mass (m/z): 185.2 (M+H).

[00161] Synthesis of 4-bromo-9H-pyrido[2,3-b]indole (6):

[00162] A suspension of compound **5** (1 g, 5.5 mmol) in anhydrous DMF (10 mL) was cooled to 0 °C and POBr₃ (3.66 g, 12.8 mmol) was added dropwise with stirring. The reaction was stirred for 24 h at room temperature and then poured into H₂O (20 mL). After cooling to 0-5 °C the pH of the solution was adjusted to 8-10 with aqueous KOH (10%). After stirring for an additional 15 min, the resulting precipitate was collected by filtration, washed with H₂O 15 mL and dried under vacuum. The crude material was purified on silica gel (20% EtOAc in Hexane) giving a 50% yield of compound **6**. ¹H NMR (500 MHz, DMSO-d₆) δ ppm: 12.19 (br, 1H), 8.46(d, 1H), 8.27 (d, 1H), 7.57 (m, 2H), 7.46 (d, 1H), 7.32 (m, 1H). Mass (m/z): 247.2 (M+H).

[00163] Synthesis of *tert*-butyl 2-(3-(9H-pyrido[2,3-b]indol-4-yl)phenylamino)ethylcarbamate (8):


[00164] A solution of compound **6** (250 mg, 1.01 mmol), boronate ester **7** (550 mg, 1.52 mmol), aqueous sodium carbonate (2.0M, 1 mL, 2.0 mmol) and dioxane (4 mL) was purged with

nitrogen and $\text{Pd}(\text{PPh}_3)_4$ (58 mg, 0.05 mmol) was added. The resulting mixture was stirred in a microwave reactor at 140 °C for 45 min. After cooling to room temperature, the mixture was poured into water (30 mL) and washed with EtOAc (2 X 50 mL). The combined organic phases were dried over anhydrous Na_2SO_4 , filtered, and concentrated in vacuo. The crude product was purified on silica gel (40% EtOAc in hexane) giving a 40% yield of compound **8**. Mass (m/z): 403.4 (M+H).

[00165] Synthesis of N1-(3-(9H-pyrido[2,3-b]indol-4-yl)phenyl)ethane-1,2-diamine (9, compound I-7):

[00166] Trifluoroacetic acid (114 mg, 1.0 mmol) was added dropwise to a suspension of compound **8** (100 mg, 0.2 mmol) in anhydrous dichloromethane (3 mL). After stirring for 2 h at room temperature, the mixture was concentrated in vacuo giving 110 mg of compound **9**. This material was stirred with desalting resin (MP-carbonate, 150 mg) in MeOH (4 mL) for 2 h. The resin was removed by filtration and the filtrate was concentrated to dryness. Purification of the residue by preparative HPLC gave 22 mg (30% yield) of pure compound **9**. ^1H NMR (500 MHz, DMSO-d_6) δ ppm: 11.9 (br, 1H), 8.41(d, 1H), 7.8 (br, 2H), 7.64 (d, 1H), 7.49 (d, 1H), 7.39 (t, 1H), 7.33 (m, 1H), 7.04 (m, 2H), 6.78-6.85(m, 3H), 6.0 (br, 1H), 3.25(t, 2H), 2.97(t, 2H). Mass (m/z): 303.2 (M+H). Purity: 98.4% by HPLC.

[00167] Synthesis of tert-butyl 2-(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenylamino)-ethylcarbamate (7):



[00168] Synthesis of tert-butyl 2-(3-bromophenylamino) ethylcarbamate (12):

[00169] A mixture of 1,3-dibromobenzene **10** (2.2 g, 9.36 mmol), *tert*-butyl 2-aminoethylcarbamate **11** (1 g, 6.2 mmol), CuI (237 mg, 1.2 mmol), L-proline (215 mg, 1.8 mmol) and potassium carbonate (1.7 g, 12.4 mmol) in DMSO (20 mL) was placed under N_2 and heated to 90°C for 14 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure to dryness. The resulting residue was dissolved in ethyl acetate (100 mL), washed with water (2X25 mL) and washed with brine (2X25 mL). After drying over anhydrous Na_2SO_4 , the organic phase was filtered and concentrated to dryness. The residue was purified on


silica gel (30% EtOAc/hexane) giving compound **12** (1 g, 50% yield) as a brown solid. ¹H NMR (500 MHz, CDCl₃) δ ppm: 7.00 (t, 1H), 6.81(d, 1H), 6.72 (s, 1H), 6.51 (d, 1H), 4.75 (br, 1H), 4.2 (br, 1H), 3.35 (t, 2H), 3.22 (t, 2H), 1.45 (s, 9H). Mass (m/z): 315.1 and 317.1.

[00170] Synthesis of tert-butyl 2-(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenylamino)-ethylcarbamate (7):

[00171] A mixture of *tert*-butyl 2-(3-bromophenylamino)ethylcarbamate **12** (1 g, 3.17 mmol), bis(pinacolato)diboron **13** (1.2 g, 4.76 mmol), palladium(II)acetate (50 mg, 0.2 mmol), BINAP (100 mg, 0.15 mmol) and cesium carbonate (2 g, 6.3 mmol) in dioxane (20 mL) was stirred at 90°C under N₂ for 14 h. After cooling to room temperature, the mixture was filtered through Celite and washed with EtOAc. The filtrate was concentrated to dryness giving crude compound **7** (1.2 g) which was used without purification. Mass (m/z): 363.3 (M+H).

Example 2

[00172] Synthesis of N1-(3-(9H-pyrido[2,3-b]indol-4-yl)phenyl)propane-1,3-diamine (9a, compound I-8):



[00173] Compounds **8a** and **9a** were prepared from compound **6** and compound **7a** according to the protocols described for the preparation of compound **8** and compound **9**.


[00174] Synthesis of *tert*-butyl 2-(3-(9H-pyrido[2,3-b]indol-4-yl)phenylamino)propylcarbamate (8a):

[00175] Compound **8a** was prepared according to the procedure for compound **8** (Example 1) using compound **6** (100 mg, 0.405 mmol), compound **7a** (228 mg, 0.60 mmol), Pd(PPh₃)₄ (23 mg, 0.025 mmol) and proportionate molar equivalents of aqueous sodium carbonate and dioxane. 80 mg (47% yield) of compound **8a** were isolated. Mass (m/z): 417.4 (M+H). Purity: 98.2% by HPLC.

[00176] Synthesis of N1-(3-(9H-pyrido[2,3-b]indol-4-yl)phenyl)propane-1,2-diamine (9a, compound I-8):

[00177] Compound **9a** was prepared according the procedure for compound **9** (Example 1) using compound **8a** (80 mg, 0.19 mmol) and trifluoroacetic acid (109 mg, 0.95 mmol). 12 mg (20% yield) of compound **9a** were isolated. ¹H NMR (500 MHz, DMSO-d₆) δ ppm: 11.91 (br, 1H), 8.41(d, 1H), 7.7 (br, 2H), 7.64 (d, 1H), 7.49 (d, 1H), 7.41 (t, 1H), 7.31 (t, 1H), 7.04-7.06 (m, 2H), 6.83(d, 1H), 6.78 (d, 1H), 5.94(br, 1H), 3.31(t, 2H), 2.90(t, 2H), 1.83-1.85(m, 2H). Mass (m/z): 317.3 (M+H).

[00178] **Synthesis of tert-butyl 3-(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenylamino)-propylcarbamate (7a):**



[00179] Compounds **12a** and **7a** were prepared from 1,3-dibromobenzene **10** and the appropriate mono-Boc protected diamine **11a** according to the protocols described for the preparation of compound **12** and compound **7**.

[00180] **Conversion of compound 10 to compound 12a:**


[00181] Compound **12a** was prepared according to the procedure for compound **12** (Example 1) using compound **11a** (350 mg, 2.96 mmol) and proportionate molar equivalents of compound **10**, CuI, L-proline, potassium carbonate and DMSO. Compound **12a** was isolated in 25% yield. ¹H NMR (500 MHz, CDCl₃) δ ppm: 6.99 (t, 1H), 6.77(d, 1H), 6.73 (s, 1H), 6.50 (d, 1H), 3.22 (d, 2H), 3.15 (t, 2H), 1.75 (m, 2H), 1.45 (s, 9H). Mass (m/z): 329.1, 331.1.

[00182] **Conversion of compound 12a to compound 7a:**

[00183] Compound **7a** was prepared according to the procedure for compound **7** (Example 1) using **12a** (1 g, 3.03 mmol), bis(pinacolato)diboron **13** (1.15 g, 4.55 mmol), palladium(II)acetate (50 mg, 0.2 mmol), BINAP (100 mg, 0.15 mmol) and cesium carbonate (1.98 g, 6.07 mmol) in dioxane (20 mL). crude compound **7a** (1.25 g) was used without purification. Mass (m/z): 377.3 (M+H).

Example 3

[00184] **Synthesis of 3-(9H-pyrido[2,3-b]indol-4-yl)aniline (9b, compound I-6):**



[00185] Compounds **8b** and **9b** were prepared from **6** and compound **7b** according to the protocols described for the preparation of compound **8** and compound **9**.


[00186] Conversion of compound 6 to compound 8b

[00187] Compound **8b** was prepared according to the procedure for compound **8** (Example 1) using compound **6** (100 mg, 0.405 mmol), compound **7b** (194 mg, 0.60 mmol), Pd(PPh₃)₄ (23 mg, 0.025 mmol) and proportionate molar equivalents of aqueous sodium carbonate and dioxane. 67 mg (46% yield) of compound **8b** were isolated. Mass (m/z): 360.4 (M+H). Purity: 97.12% by HPLC.

[00188] Conversion of compound 8b to compound 9b (I-6):

[00189] Compound **9b** was prepared according to the procedure for compound **9** (Example 1) using compound **8b** (67 mg, 0.18 mmol) and trifluoroacetic acid (106 mg, 0.90 mmol). 22 mg (45% yield) of compound **9b** were isolated. ¹H NMR (500 MHz, DMSO-d⁶) δ ppm: 11.89 (br, 1H), 8.41(d, 1H), 7.66 (d, 2H), 7.50 (d, 1H), 7.48 (d, 1H), 7.39 (t, 1H), 7.22 (t, 1H), 7.03 (m, 2H), 6.83(s, 1H), 6.75 (m, 2H), 5.30(br, 2H). Mass (m/z): 260.3 (M+H).

[00190] Synthesis of tert-butyl 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenylcarbamate (7b):



[00191] Synthesis of *tert*-butyl 3-bromophenylcarbamate (12b):

[00192] To a solution of 3-bromoaniline (300 mg, 1.7 mmole) in dichloromethane (5 mL) at 0°C was added triethylamine (3.4 mL, 0.47 mmoles, 2.0 eq.). After stirring for 5-10 min, di-*tert*-butyl borane (1.0 g, 2.0 mmole) was added dropwise. The reaction mixture was stirred for 1 h and then quenched with 10% aqueous NaHSO₃ (10 mL). The organic layer was separated and the aqueous layer was extracted with dichloromethane (3 x 10 mL). The combined organic layers were dried over Na₂SO₄ and concentrated. The residue was purified by column chromatography (silica gel, hexanes/ethyl acetate 9:1) to yield compound 12b (3-bromo-NHBOC-phenyl) (150 mg, 48% yield).


butyl dicarbonate (452 mg, 2 moles, 1.2 eq.) was added slowly and the reaction was allowed to stir at room temperature for 12 h. The reaction was then concentrated to dryness and the residue was purified on silica gel (5% EtOAc/hexane) giving compound **12b** (272 mg, 60% Yield). ¹H NMR (500 MHz, CDCl₃) δ PPM: 7.65 (s, 1H), 7.25 (d, 1H), 7.11-7.13 (m, 2H), 6.51 (br, 1H), 1.46 (s, 9H).

[00193] Conversion of compound **12b to compound **7b**:**

[00194] Compound **7b** was prepared according to the procedure for compound **7** (Example 1) using compound **12b** (200 mg) and proportionate molar equivalents of compound **13**, palladium(II)acetate, BINAP, cesium carbonate and dioxane. 300 mg of crude **7b** were isolated and used without purification.

Example 4

[00195] Synthesis of (3-(9H-pyrido[2,3-b]indol-4-yl)phenyl)methanamine (9c**, compound I-3):**



[00196] Compounds **8c** and **9c** were prepared from compound **6** and compound **7c** according to the protocols described for the preparation of compound **8** and compound **9**.

[00197] Conversion of compound **6 to compound **8c**:**


[00198] Compound **8c** was prepared according to the procedure for compound **8** (Example 1) using compound **6** (100 mg, 0.405 mmol), compound **7c** (199 mg, 0.60 mmol), Pd(PPh₃)₄ (23 mg, 0.025 mmol) and proportionate molar equivalents of aqueous sodium carbonate and dioxane. 75 mg (50% yield) of compound **8c** were isolated. Mass (m/z): 374.4 (M+H). Purity: 95.5% by HPLC.

[00199] Conversion of compound **8c to compound **9c** (I-3):**

[00200] Compound **9c** was prepared according to the procedure for compound **9** (Example 1) using compound **8c** (75 mg, 0.2 mmol), and trifluoroacetic acid (118 mg, 0.90 mmol). 12 mg

(20% yield) of compound **9a** were isolated. ¹H NMR (500 MHz, DMSO-d⁶) δ PPM: 11.89 (br, 1H), 8.48(d, 1H), 7.77 (s, 1H), 7.65-7.68 (m, 3H), 7.53 (dd, 2H), 7.42 (t, 1H), 7.10 (d, 1H), 7.09 (t, 1H), 4.12(s, 2H). Mass (m/z): 274.3 (M+H).

[00201] **Synthesis of tert-butyl 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzylcarbamate (7c):**



[00202] Compounds **12c** and **7c** were prepared from 3-bromobenzylamine hydrochloride and compound **13** according to the protocols described for the preparation of compound **12b** and compound **7b**.

[00203] **Synthesis of tert-butyl 3-bromobenzylcarbamate (12c):**


[00204] Compound **12c** was prepared according to the procedure for compound **12b** (Example 3) using 3-bromobenzylamine hydrochloride (125 mg, 0.56 mmol) and proportionate molar equivalents of dichloromethane, triethylamine and di-*tert*-butyl dicarbonate. 98 mg (60% yield) of compound **12c** were isolated.

[00205] **Conversion of 12c to 7c:**

[00206] Compound **7c** was prepared according to the procedure for compound **7** (Example 1) using compound **12c** (90 mg) and proportionate molar equivalents of compound **13**, palladium(II)acetate, BINAP, cesium carbonate and dioxane. 199 mg of crude **7c** were isolated and used without purification.

Example 5


[00207] **Synthesis of 1-(3-(9H-pyrido[2,3-b]indol-4-yl)phenyl)guanidine (14, compound I-4):**



[00208] To a solution of **9b** (30 mg, 0.115 mmol) in ethanol (2 mL) was added NH₂CN (29 mg, 6 eq.). The mixture was refluxed for 2 days after which, the volatiles were removed under reduced pressure. The resulting crude residue was dissolved in water and the pH was adjusted to ~10 using aqueous 2N NaOH. The aqueous mixture was extracted with EtOAc (5 mL). The organic phase was dried over Na₂SO₄, concentrated and purified by preparative HPLC giving compound **14** (4 mg, 12% yield) as a yellow solid. Mass (m/z): 302.3(M+H). Purity: 90.05% by HPLC.

Example 6


[00209] **Synthesis of 1-(3-(9H-pyrido[2,3-b]indol-4-yl)benzyl)guanidine (14a, compound I-5):**



[00210] Compound **14a** was prepared according to the preparation of compound **14** (Example 5) using compound **9c** (40 mg, 0.146 mmol) and NH₂CN (35 mg, 6 eq.). 4 mg (8.6% yield) of compound **14a** were isolated. Mass (m/z): 316.4 (M+H). Purity: 98.1% by HPLC.

Example 7

[00211] **Synthesis of compound 18 (I-2):**



[00212] Preparation of compound 16:

[00213] A Solution of compound **6** (250 mg, 1.01 mmol), 3-hydroxyphenylboronic acid **15** (167 mg, 1.2 mmol) and aqueous sodium carbonate (2 M, 1 mL, 2.0 mmol) in dioxane (4 mL) was purged with nitrogen. $\text{Pd}(\text{PPh}_3)_4$ (58 mg, 0.05 mmol) was added and the resulting mixture was stirred at 90°C for 5 h. After cooling to room temperature, the mixture was poured into water (30 mL) and extracted with EtOAc (2 X 50 mL). The combined organic phases were dried over Na_2SO_4 , filtered, and concentrated to dryness. The residue was purified on silica gel (40% EtOAc/hexane) giving compound **16** (50% yield). ^1H NMR (500 MHz, DMSO-d_6) δ ppm: 11.89 (br, 1H), 9.68(br, 1H), 8.42 (d, 1H), 7.58 (d, 1H), 7.49 (d, 1H), 7.39 (m, 2H), 7.04 (m, 4H), 6.92 (d, 1H). Mass (m/z): 261.3 ($\text{M}+\text{H}$).

[00214] Synthesis of 3-(9H-pyrido[2,3-b]indol-4-yl)phenyl trifluoromethanesulfonate (17):


[00215] To a solution of compound **16** (200 mg, 0.7 mmole) in dichloromethane (5 mL) at 0°C was added triethylamine (0.26 mL, 1.9 mmole). After stirring for 5-10 min, trifilic anhydride (0.2 mL, 1.5 mmole) was added drop-wise. Stirring was continued at room temperature for 2 h after which, the reaction was concentrated to dryness and the residue was purified on silica gel (20% EtOAc/hexane) giving compound **17** (150 mg, 33% Yield). Mass (m/z): 393.3 ($\text{M}+\text{H}$).

[00216] Synthesis of 4-(3-(1H-imidazol-1-yl)phenyl)-9H-pyrido[2,3-b]indole (18, I-2):

[00217] A mixture of compound **17** (150 mg, 0.38 mmol), imadazole (77 mg, 1.14 mmol), palladium(II)acetate (9 mg, 10 mol%), BINAP (24 mg, 10 mol%) and cesium carbonate (250 mg, 0.76 mmol) in THF (4 mL) was stirred in a microwave reactor at 150°C for 45 min. The mixture was filtered through Celite and the Celite pad was washed with EtOAc. The combined filtrate was concentrated to dryness and the crude product was purified by preparative HPLC to give compound **18** (7 mg, 5.8% yield) as an off white solid. Mass (m/z): 311.2 (M+H). Purity: 99.2% by HPLC.

Example 8

[00218] Synthesis of compound **23** (I-1):



[00219] Synthesis of *tert*-butyl 4-(3-bromophenyl)piperazine-1-carboxylate (**20**):

[00220] A mixture of 1,3-dibromobenzene **10** (1.5 g, 6.35 mmol), *tert*-butyl piperazine-1-carboxylate **19** (1.18g, 6.35 mmol), palladium(II)acetate (14 mg, 0.06 mmol), BINAP (79 mg, 0.12 mmol) and cesium carbonate (4 g, 12.7 mmol) in dioxane (25 mL) was stirred at 90°C under N_2 for 16 h. The mixture was filtered through Celite and the Celite pad was washed with EtOAc. The combined filtrate was concentrated to dryness and the crude product was purified on silica gel (5% EtoAc/hexane) giving compound **20** (500 mg, 23% yield) as an off white solid. ^1H NMR (500 MHz, CDCl_3) δ ppm: 7.10 (t, 1H), 7.08 (s, 1H), 6.98 (d, 1H), 6.81 (d, 1H), 3.55 (t, 4H), 3.13 (t, 4H), 1.45 (s, 9H).

[00221] Synthesis of 3-(4-(*tert*-butoxycarbonyl)piperazin-1-yl) phenylboronic acid (**21**):

[00222] To a solution of compound **20** (500 mg, 1.46 mmol) in anhydrous THF (5 mL) at -78°C was added n-BuLi (2.5M in THF, 0.05 mL, 2.19 mmol) dropwise. After stirring at -78°C for 45 min, triisopropylborate (0.68 mL, 2.93 mmol) was added dropwise. Stirring was continued for 2 h at -78°C and the reaction was allowed to warm to room temperature overnight. The reaction was quenched with aqueous saturated NH₄Cl (5 mL) and extracted with EtOAc (2 X 25 mL). The combined organic phases were dried over Na₂SO₄, filtered and concentrated to dryness. The residue was purified on silica gel (5% MeOH/dichloromethane) giving compound **21** (150 mg, 33% yield). ¹H NMR (500 MHz, CDCl₃) δ ppm: 7.10 (t, 1H), 7.08 (s, 1H), 6.98 (d, 1H), 6.81 (d, 1H), 3.55 (t, 4H), 3.13 (t, 4H), 1.45 (s, 9H). Mass (m/z): 307.3 (M+H).

[00223] **Synthesis of *tert*-butyl 4-(3-(9H-pyrido[2,3-b]indol-4-yl)phenyl)piperazine-1-carboxylate (22):**


[00224] Compound **22** was prepared according to the procedure for compound **16** (Example 7) using compound **6** (100 mg, 0.405 mmol), compound **21** (148 mg, 0.48 mmol) and Pd(PPh₃)₄ (23 mg, 0.025 mmol). 90 mg (52% yield) of compound **22** were isolated. ¹H NMR (500 MHz, DMSO-d₆) δ ppm: 11.94 (br, 1H), 8.45 (d, 1H), 7.50 - 7.58 (m, 3H), 7.42 (t, 1H), 7.25 (s, 1H), 7.19-7.21 (m, 3H), 7.03 (t, 1H), 3.20 (t, 4H), 3.47 (t, 4H), 1.41 (s, 9H). Mass (m/z): 429.3 (M+H).

[00225] **Synthesis of 4-(3-(piperazin-1-yl)phenyl)-9H-pyrido[2,3-b]indole (23, I-1):**

[00226] Compound **23** was prepared according to the procedure for compound **9** (Example 1) using compound **22** (75 mg, 0.2 mmol) and trifluoroacetic acid (118 mg, 0.90 mmol). 12 mg (20% yield) of compound **23** were isolated. ¹H NMR (500 MHz, DMSO-d⁶) δ ppm: 11.97 (br, 1H), 8.87 (b, 1H), 8.45 (d, 1H), 7.54 - 7.58 (m, 3H), 7.42 (d, 1H), 7.25 (s, 1H), 7.19-7.21 (m, 3H), 7.03 (t, 1H), 3.26 (t, 4H), 3.37 (t, 4H). Mass (m/z):

Example 9

[00227] **Synthesis of Compound 28 (I-9):**



[00228] **Synthesis of *tert*-butyl 2-(3-bromophenoxy)ethylcarbamate (25):**

[00229] A mixture of 3-bromophenol **24** (200 mg, 1.15 mmol), *tert*-butyl 2-bromoethylcarbamate (257 mg, 1.15 mmol) and cesium carbonate (753 mg, 2.3 mmol) in DMF (5 mL) was stirred at 90°C under N₂ for 5 h. The mixture was filtered through Celite and the Celite pad was washed with DMF (3 mL). The filtrate was concentrated to dryness and the crude product was purified on silica gel (20% EtOAc/hexane) giving compound **25** in 60% yield. ¹H NMR (500 MHz, CDCl₃) δ PPM: 7.05-7.19 (m, 3H), 6.80 (dd, 1H), 5.0 (br, 1H), 3.98 (t, 2H), 3.52 (t, 2H), 1.46 (s, 9H).

[00230] **Synthesis of *tert*-butyl 2-(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)ethylcarbamate (26):**

[00231] A mixture of *tert*-butyl 2-(3-bromophenoxy)ethylcarbamate **25** (300 mg, 0.94 mmol), bis(Pinacolato)diboron (360 mg, 1.42 mmol), Palladium(II)acetate (12 mg, 0.05 mmol), BINAP (22 mg, 0.03 mmol) and Cesium carbonate (590 mg, 1.8 mmol) in dioxane (20 mL) was stirred at 90°C under N₂ for 16 h. The mixture was filtered through Celite and the Celite pad was washed with EtOAc. The filtrate was concentrated to dryness giving crude compound **26** which was used in the next step without further purification.

[00232] **Synthesis of *tert*-butyl 2-(3-(9H-pyrido[2,3-b]indol-4-yl) phenoxy) ethyl carbamate (27):**

[00233] A mixture of compound **6** (100 mg, 0.40 mmol), compound **26** (220 mg, 0.60 mmol), aqueous sodium carbonate (2 M, 1 mL, 2.0 mmol) in dioxane (4 mL) was purged with nitrogen.


Pd(PPh₃)₄ (23 mg, 0.02 mmol) was added and the resulting mixture was stirred in a microwave reactor at 140°C for 45 min. After cooling to room temperature, the mixture was poured into water (10 mL) and extracted with EtOAc (2 X 25 mL). The combined organic layers were dried over Na₂SO₄, filtered, and concentrated to dryness. The residue was purified on silica gel (40% EtOAc/hexane) giving compound **27** (50% yield). Mass (m/z): 404.4 (M+H).

[00234] Synthesis of 2-(3-(9H-pyrido[2,3-b]indol-4-yl)phenoxy)ethanamine (28, compound I-9):

[00235] To a solution of compound **27** (100 mg, 0.24 mmol) in anhydrous dichloromethane (3 mL) was added trifluoroacetic acid (141 mg, 1.2 mmol) dropwise. After stirring at room temperature for 2 h, the reaction was concentrated to dryness. The residue was stirred with MP-carbonate resin (150 mg) in MeOH (5 mL) for 2 h. The resin was removed by filtration. The filtrate was concentrated to dryness and the residue was purified by preparative HPLC giving compound **28** (14 mg, 18% yield) as a off white solid. ¹H NMR (500 MHz, DMSO-d₆) δ ppm: 11.98 (br, 1H), 8.43(d, 1H), 7.98 (br, 2H), 7.55 (m, 3H), 7.40 (t, 1H), 7.29 (d, 1H), 7.24 (s, 1H), 7.18 (d, 2H), 7.08(d, 3H), 7.02 (t, 1H). Mass (m/z): 304.3 (M+H).

Example 10

[00236] Synthesis of compound 28a (I-10):



[00237] Compound **28a** was prepared from compound **24** according to protocols described for the preparation of compounds **25**, **26**, **27** and **28**.

[00238] Synthesis of *tert*-butyl 3-(3-bromophenoxy)propylcarbamate (25a):

[00239] Compound **25a** was prepared according to the procedure for compound **25** (Example 9) using compound **24** (200 mg, 1.15 mmol), *tert*-butyl 3-bromopropylcarbamate (273 mg, 1.15 mmol), Cs₂CO₃ (753 mg, 2.31 mmol) and DMF (5 mL). 235 mg (62% yield) of compound **25a** were isolated. ¹HNMR (500 MHz, CDCl₃) δ PPM: 7.05-7.19 (m, 3H), 6.80 (dd, 1H), 4.7 (br, 1H), 3.99 (t, 2H), 3.31 (t, 2H), 1.97 (m, 2H), 1.44 (s, 9H).

[00240] **Synthesis of *tert*-butyl 3-(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)propylcarbamate**

[00241] Compound **26a** was prepared according to the procedure for compound **26** (Example 9) using compound **25a** (300 mg, 0.9 mmol), bis(Pinacolato)diboron (346 mg, 1.36 mmol), Palladium(II)acetate (12 mg, 0.05 mmol), BINAP (22 mg, 0.03 mmol) and Cesium carbonate (590 mg, 1.8 mmol) in dioxane (20 mL). Crude compound **26a** (350 mg) was used in the next step without further purification.

[00242] **Synthesis of *tert*-butyl 3-(3-(9H-pyrido[2,3-b]indol-4-yl)phenoxy)propylcarbamate (27a):**


[00243] Compound **27a** was prepared according to the procedure for compound **27** (Example 9) using compound **6** (100 mg, 0.40 mmol), compound **26a** (229 mg, 0.60 mmol), aqueous sodium carbonate (2 M, 1 mL, 2.0 mmol) and Pd(PPh₃)₄ (23 mg, 0.02 mmol) in dioxane (4 mL). Compound **27a** was isolated in 54% yield. Mass (m/z): 418.4 (M+H).

[00244] **Synthesis of 3-(3-(9H-pyrido[2,3-b]indol-4-yl)phenoxy)propan-1-amine (28a, I-10):**

[00245] Compound **28a** was prepared according to the procedure for compound **28** (Example 9) using compound **27a** (100 mg, 0.23 mmol) and trifluoroacetic acid (136 mg, 1.2 mmol) in anhydrous dichloromethane (3 mL). The crude product was purified by preparative HPLC giving compound **28a** (13 mg, 17% yield) as an off white solid. ¹HNMR (500 MHz, DMSO-d₆) δ PPM: 11.97 (br, 1H), 8.44(d, 1H), 7.80 (br, 2H), 7.51-7.54 (m, 2H), 7.39 (t, 1H), 7.25 (d, 1H), 7.19 (s, 1H), 7.08 (d, 2H), 7.07 (dd, 2H), 7.02 (t, 1H); 4.12 (t, 2H), 2.99 (m, 2H), 2.01 (t, 2H). Mass (m/z): 318.3 (M+H). Purity: 97.02% by HPLC.

Example 11

[00246] Scheme for the Preparation of compounds 36a-d (I-11, I-13, I-14, I-12) and General Procedure for the Preparation of Compounds 32a-d:



[00247] General procedure for the synthesis of compounds 32a-d (I-11, I-13, I-14, I-12):

[00248] A mixture of 2,3-dichloro pyridine **29** (1 g, 6.76 mmol), an aniline **30** (7.43 mmol), $\text{Pd}(\text{OAc})_2$ (98 mg, 0.065 mmol), PPh_3 (198 mg, 0.13 mmol) and NaO^tBu (780 mg, 1.2 mmol) in o-Xylene (16 mL) was sparged with nitrogen for 5 min, placed under a nitrogen atmosphere, and heated to 130°C for 3 h in a sealed sample vial. The reaction mixture was cooled to room temperature giving compound **31**. $\text{Pd}(\text{OAc})_2$ (98 mg, 0.065 mmol), PCy_3 (200 mg, 0.1 mmol), DBU (2 g, 2 mmol) and dimethyl acetamide (16 mL) were added to the reaction vessel. The reaction mixture was sparged for 5 min, placed under a nitrogen atmosphere and heated to 160°C for 16 h. The reaction mixture was concentrated to dryness. The residue was dissolved in ethyl

acetate (2 X 200 mL). The mixture was washed with water (3 X 50 mL) and then brine (2 X 50 mL), dried over anhydrous Na_2SO_4 , filtered and concentrated. The residue was purified on silica gel (70% EtOAc/hexane) giving compounds 32.

Example 12

[00249] **Synthesis of 7-methyl-9H-pyrido[2,3-b]indole (32a, I-11):**

[00250] Compound 32a was prepared according to Example 11 using 3-methylaniline (877 mg, 7.43 mmol) 30a and proportionate molar equivalents of compound 29, $\text{Pd}(\text{OAc})_2$, PPh_3 , NaO^tBu and o-xylene. Continuing according to Example 11, crude isolated compound 31a was converted to compound 32a using proportionate molar equivalents of $\text{Pd}(\text{OAc})_2$, PCy_3 , DBU and dimethyl acetamide. 480 mg (40% yield) of compound 32a were isolated. ^1H NMR (500 MHz, DMSO-d_6) δ PPM: 11.61 (br, 1H), 8.41(dd, 2H), 8.02 (d, 1H), 7.38 (s, 1H), 7.21 (d, 1H), 7.15 (d, 1H), 2.45 (s, 3H). Mass (m/z): 183.2 (M+H).

[00251] **Synthesis of 7-methyl-9H-pyrido[2,3-b]indole 1-oxide (33a):**

[00252] Compound 33a was prepared according to the procedure for compound 5 (Example 1) using compound 32a (370 mg, 2.02 mmol) and proportionate molar equivalents of aqueous H_2O_2 (35%) and CH_3COOH . 160 mg (40% yield) of compound 33a were isolated. Mass (m/z): 199.2(M+H).

[00253] **Synthesis of 4-bromo-7-methyl-9H-pyrido[2,3-b]indole (34a):**

[00254] Compound 34a was prepared according to the procedure for compound 6 (Example 1) using compound 33a (155 mg, 0.782 mmol) and proportionate molar equivalents of anhydrous DMF and POBr_3 . 61 mg (30% yield) of compound 34a were isolated. Mass (m/z): 261.1, 263.1.

[00255] **Synthesis of tert-butyl 3-(7-methyl-9H-pyrido[2,3-b]indol-4-yl) phenethyl carbamate (35a):**

[00256] Compound 35a was prepared according to the procedure for compound 8 (Example 1) using compound 34a (61 mg, 0.23 mmol) and proportionate molar equivalents of boronate ester 7, aqueous sodium carbonate (2.0M), dioxane and $\text{Pd}(\text{PPh}_3)_4$. 49 mg (50% yield) of compound 35a were isolated. Mass (m/z): 417.3(M+H). **Synthesis of 2-(3-(7-methyl-9H-pyrido[2,3-b]indol-4-yl)phenyl)ethanamine (36a, I-11):**

[00257] Compound 36a was prepared according to the procedure for compound 9 (Example 1) from compound 35a (49 mg, 0.117 mmol) and treating with proportionate molar equivalents of trifluoroacetic acid in anhydrous dichloromethane. Purification by preparative HPLC gave 14

mg (37% yield) of compound **36a**. ¹H NMR (500 MHz, DMSO-d₆) δ PPM: 11.89 (br, 1H), 8.39(d, 1H), 7.58 (d, 1H), 7.35 (m, 2H), 7.05 (d, 1H), 6.86 (m, 3H), 6.80 (d, 1H), 6.02(br, 1H), 2.98 (t, 2H), 2.40 (s, 3H). ¹H NMR D₂O Exchange (500 MHz, DMSO-d₆) δ PPM: 8.32 (d, 1H), 7.50 (d, 1H), 7.35 (m, 2H), 7.05 (d, 1H), 6.86 (m, 3H), 6.80 (d, 1H), 3.35 (t, 2H), 2.98 (t, 2H), 2.38 (s, 3H). Mass (m/z): 317.3 (M+H).

Example 13

[00258] Synthesis of 6-methyl-9H-pyrido[2,3-b]indole (32b, I-13):

[00259] Compound **32b** was prepared according to Example 11 using 4-methylaniline (877 mg, 7.43 mmol) **30b** and proportionate molar equivalents of compound **29**, Pd(OAc)₂, PPh₃, NaO^tBu and o-xylene. Continuing according to Example 11, crude isolated compound **31b** was converted to compound **32b** using proportionate molar equivalents of Pd(OAc)₂, PCy₃, DBU and dimethyl acetamide. 495 mg (41% yield) of compound **32b** were isolated. ¹H NMR (500 MHz, DMSO-d₆) δ ppm: 11.59 (br, 1H), 8.41(d, 1H), 8.38 (d, 1H), 7.9 (s, 1H), 7.38 (d, 1H), 7.25 (d, 1H), 7.18 (t, 1H), 2.42(s, 3H). Mass (m/z): 183.2 (M+H).

[00260] Synthesis of 6-methyl-9H-pyrido[2,3-b]indole 1-oxide (33b):

[00261] Compound **33b** was prepared according to the procedure for compound **5** (Example 1) using compound **32b** (420 mg, 2.29 mmol) and proportionate molar equivalents of aqueous H₂O₂ (35%) and CH₃COOH. 173 mg (38% yield) of compound **33b** were isolated. Mass (m/z): 199.2 (M+H).

[00262] Synthesis of 4-bromo-6-methyl-9H-pyrido[2,3-b]indole (34b):

[00263] Compound **34b** was prepared according to the procedure for compound **6** (Example 1) using compound **33b** (170 mg, 0.857 mmol) and proportionate molar equivalents of anhydrous DMF and POBr₃. 69 mg (31% yield) of compound **34b** were isolated. Mass (m/z): 261.1, 263.1.

[00264] Synthesis of *tert*-butyl 3-(6-methyl-9H-pyrido[2,3-b]indol-4-yl)phenethylcarbamate (35b):

[00265] Compound **35b** was prepared according to the procedure for compound **8** (Example 1) using compound **34b** (68 mg, 0.256 mmol) and proportionate molar equivalents of boronate ester **7**, aqueous sodium carbonate (2.0M), dioxane and Pd(PPh₃)₄. 54 mg (49% yield) of compound **35b** were isolated. Mass (m/z): 417.3(M+H).

[00266] Synthesis of N1-(3-(6-methyl-9H-pyrido[2,3-b]indol-4-yl)phenyl)ethane-1,2-diamine (36b, I-13):

[00267] Compound **36b** was prepared according to the procedure for compound **9** (Example 1) using compound **35b** (53 mg, 0.126 mmol) and proportionate molar equivalents of trifluoroacetic acid and anhydrous dichloromethane. Purification by preparative HPLC gave 16 mg (40% yield) of compound **36b**. ¹H NMR (500 MHz, DMSO-d₆) δ PPM: 11.95 (br, 1H), 8.62(d, 1H), 7.70(s, 1H), 7.65 (d, 2H), 7.58 (t, 1H), 7.45 (d, 1H), 7.24 (d, 1H), 7.10(dd, 2H), 7.0 (d, 1H), 6.20 (br, 1H), 3.20 (t, 2H), 2.70 (s, 3H). ¹H NMR D₂O Exchange (500 MHz, DMSO-d₆) δ PPM: 8.38 (d, 1H), 7.45 (m, 2H), 7.38 (t, 1H), 7.22 (d, 1H), 7.05 (d, 1H), 6.85 (m, 2H), 6.80 (d 1H), 3.35(t, 2H), 2.98 (t, 2H), 2.22 (s, 3H). Mass (m/z): 317.3 (M+H).

Example 14

[00268] Synthesis of 6 chloro-9H-pyrido[2,3-b]indole (32c):

[00269] Compound **32c** was prepared according to Example 11 using 4-chloroaniline (1.03 g, 7.43 mmol) **30c** and proportionate molar equivalents of compound **29**, Pd(OAc)₂, PPh₃, NaO^tBu and o-xylene. Continuing according to Example 11, crude isolated compound **31c** was converted to compound **32c** using proportionate molar equivalents of Pd(OAc)₂, PCy₃, DBU and dimethyl acetamide. 400 mg (30% yield) of compound **32c** were isolated. ¹H NMR (500 MHz, DMSO-d₆) δ PPM: 11.95 (br, 1H), 8.59(d, 1H), 8.42 (d, 1H), 8.26 (s, 1H), 7.58(d, 1H), 7.49 (d, 1H), 7.21 (t, 1H). Mass (m/z): 203.2, 205.2.

[00270] Synthesis of 6-chloro-9H-pyrido[2,3-b]indole 1-oxide (33c):

[00271] Compound **33c** was prepared according to the procedure for compound **5** (Example 1) using compound **32c** (380 mg, 1.88 mmol) and proportionate molar equivalents of aqueous H₂O₂ (35%) and CH₃COOH. 205 mg (50% yield) of compound **33c** were isolated. Mass (m/z): 219.2, 221.2.

[00272] Synthesis of 4-bromo-6-chloro-9H-pyrido[2,3-b]indole (34c):

[00273] Compound **34c** was prepared according to the procedure for compound **6** (Example 1) using compound **33c** (200 mg, 0.917 mmol) and proportionate molar equivalents of anhydrous DMF and POBr₃. 90 mg (35% yield) of compound **34c** were isolated. Mass (m/z): 283.0, 285.0.

[00274] Synthesis of tert-butyl 3-(6-chloro-9H-pyrido[2,3-b] indol-4-yl) phenethyl carbamate (35c):

[00275] Compound **35c** was prepared according to the procedure for compound **8** (Example 1) using compound **34c** (85 mg, 0.302 mmol) and proportionate molar equivalents of boronate ester **7**, aqueous sodium carbonate (2.0M), dioxane and Pd(PPh₃)₄. 59 mg (45% yield) of compound **35c** were isolated. Mass (m/z): 437.3, 439.3.

[00276] **Synthesis of N1-(3-(6-chloro-9H-pyrido[2,3-b]indol-4-yl)phenyl)ethane-1,2-diamine (36c, I-14):**

[00277] Compound **36c** was prepared according to the procedure for compound **9** (Example 1) using compound **35c** (58 mg, 0.133 mmol) and proportionate molar equivalents of trifluoroacetic acid and anhydrous dichloromethane. Purification by preparative HPLC gave 15 mg (36% yield) of compound **36c**. ¹H NMR (500 MHz, DMSO-d₆) δ PPM: 12.10 (br, 1H), 8.40(d, 1H), 7.80(br, 2H), 7.55 (s, 1H), 7.50 (d, 1H), 7.40 (d, 1H), 7.38 (t, 1H), 7.08(d, 1H), 6.8 (m, 3H), 6.00 (br, 1H), 3.00 (t, 2H). ¹H NMR D₂O Exchange (500 MHz, DMSO-d₆) δ PPM: 8.38 (d, 1H), 7.59 (m, 2H), 7.39 (d, 1H), 7.35 (t, 1H), 7.15 (d, 1H), 6.89 (m, 3H), 3.38 (t, 2H), 2.98 (t, 2H). Mass (m/z): 337.2, 339.2.

Example 15

[00278] **Synthesis of 7-methoxy-9H-pyrido[2,3-b]indole (32d):**

[00279] Compound **32d** was prepared according to Example 11 using 3-methoxyaniline (890 mg, 7.43 mmol) **30d** and proportionate molar equivalents of compound **29**, Pd(OAc)₂, PPh₃, NaO^tBu and o-xylene. Continuing according to Example 11, crude isolated compound **31d** was converted to compound **32d** using proportionate molar equivalents of Pd(OAc)₂, PCy₃, DBU and dimethyl acetamide. 355 mg (30% yield) of compound **32d** were isolated. ¹H NMR (500 MHz, DMSO-d₆) δ PPM: 11.61 (br, 1H), 8.38(d, 1H), 8.34 (d, 1H), 7.18 (d, 1H), 6.90 (s, 1H), 6.81 (t, 1H), 3.82(s, 3H), Mass (m/z): 199.2 (M+H).

[00280] **Synthesis of 7-methoxy-9H-pyrido[2,3-b]indole 1-oxide (33d):**

[00281] Compound **33d** was prepared according to the procedure for compound **5** (Example 1) using compound **32d** (330 mg, 1.67 mmol) and proportionate molar equivalents of aqueous H₂O₂ (35%) and CH₃COOH. 89 mg (25% yield) of compound **33d** were isolated. Mass (m/z): 215.2 (M+H).

[00282] **Synthesis of 4-bromo-7-methoxy-9H-pyrido[2,3-b]indole (34d):**

[00283] Compound **34d** was prepared according to the procedure for compound **6** (Example 1) using compound **33d** (88 mg, 0.411 mmol) and proportionate molar equivalents of anhydrous DMF and POBr₃. 45 mg (40% yield) of compound **34d** were isolated. Mass (m/z): 277.1, 279.1.

[00284] **Synthesis of tert-butyl 3-(6-methoxy-9H-pyrido[2,3-b]indol-4-yl)phenethyl carbamate (35d):**

[00285] Compound **35d** was prepared according to the procedure for compound **8** (Example 1) using compound **34d** (44 mg, 0.158 mmol) and proportionate molar equivalents of boronate ester **7**, aqueous sodium carbonate (2.0M), dioxane and Pd(PPh₃)₄. 27 mg (40% yield) of compound **35d** were isolated. Mass (m/z): 433.3 (M+H).

[00286] **Synthesis N1-(3-(7-methoxy-9H-pyrido[2,3-b]indol-4-yl)phenyl)ethane-1,2-diamine (36d, I-12):**

[00287] Compound **36d** was prepared according to the procedure for compound **9** (Example 1) using compound **35d** (27 mg, 0.062 mmol) and proportionate molar equivalents of trifluoroacetic acid and anhydrous dichloromethane. Purification by preparative HPLC gave 6.3 mg (30% yield) of compound **36d**. Mass (m/z): 333.2 (M+H).

[00288] Additional compounds of formula **I** were prepared in a manner substantially similar to that described above.

Example 16

In Vitro CaMKII δ Activity Assay

[00289] An exemplary procedure for the *in vitro* CaMKII δ inhibition assay, which can be used to determine the inhibitory action of compounds of the invention toward CaMKII, follows. The procedure is taken from Chao LH, et al., (2010) Nat Struct Mol Biol. 17(3): 264–272, which is incorporated herein by reference in its entirety.

[00290] The inhibition of CaMKII activity was evaluated using a coupled assay measuring ADP released following ATP hydrolysis and phosphor-transfer to the peptide substrate AC3 (KKALHRQETVDAL; SEQ ID NO: 1) (1). A full length, C-terminal His/Gln tagged CaMKII δ construct was used (sequence in Table 2 below).

[00291] **Table 2: Amino acid sequence of CaMKII δ construct.**

M	A	S	T	T	T	C	T	R	F	T	D	E	Y	Q	L	F	E	E	L	G	K	G	A	F	S	V	V	R	R	C	M	K	I
P	T	G	Q	E	Y	A	A	K	I	I	N	T	K	K	L	S	A	R	D	H	Q	K	L	E	R	E	A	R	I	C	R	L	L
K	H	P	N	I	V	R	L	H	D	S	I	S	E	E	G	F	H	Y	L	V	F	D	L	V	T	G	G	E	L	F	E	D	I

V A R E Y Y S E A D A S H C I Q Q I L E S V N H C H L N G I V H R D
L K P E N L L A S K S K G A A V K L A D F G L A I E V Q G D Q Q
A W F G F A G T P G Y L S P E V L R K D P Y G K P V D M W A C G V
I L Y I L L V G Y P P F W D E D Q H R L Y Q Q I K A G A Y D F P S P
E W D T V T P E A K D L I N K M L T I N P A K R I T A S E A L K H P
W I C Q R S T V A S M M H R Q E T V D C L K K F N A R R K L K G A
I L T T M L A T R N F S A A K S L L K K P D G V K E S T E S S N T T
I E D E D V K A R K Q E I I K V T E Q L I E A I N N G D F E A Y T K I
C D P G L T A F E P E A L G N L V E G M D F H R F Y F E N A L S K S
N K P I H T I I L N P H V H L V G D D A A C I A Y I R L T Q Y M D G
S G M P K T M Q S E E T R V W H R R D G K W Q N V H F H R S G S P
T V P I K L G S F L D H S F G A R A Q V X G H N H N H (SEQ ID NO: 2)

[00292] Compounds were added in 5 uL volume to wells in UV transparent 96-well plates (1/2 area well size). The final compound concentrations tested ranged from 0.5 nM to 10 uM). Assays were performed in duplicate. CaMKII δ is added to at a final concentration of 16 nM to a mixture containing 100 mM Tris (pH 7.5), 150 mM KCl, 0.27 mM EGTA, 1.3 mM PEP, 0.2 mg/ml AC3, 6.9% (v/v) PK/LDH mixture (Sigma P0294), 0.38 mM NADH and kept on ice. 72 uL of the enzyme mixture was added to the wells containing compounds and the plate was shaken briefly and kept on ice. The assay was initiated by adding 23 uL of a mixture containing 100 mM Tris (pH 7.5), 150 mM KCl, 1.7 mM CaCl2, 48 mM MgCl2, 0.35 mM ATP and 6.7 ug/mL calmodulin. The rate of ADP released was measured as the rate of absorbance decrease at 340 nM at 25°C and plotted against the log of the compound concentration (**Figure 1**). IC₅₀ data were fitted using GraphPad Prism software.

[00293] The results of the *in vitro* CaMKII δ activity assays are set forth in **Table 3**. The compound numbers correspond to the compound numbers in Table 1. Compounds having an activity designated as "A" provided an IC₅₀ \leq 50 nM; compounds having an activity designated as "B" provided an IC₅₀ of 50-250 nM; compounds having an activity designated as "C" provided an IC₅₀ of 250-1000 nM; and compounds having an activity designated as "D" provided an IC₅₀ \geq 1 μ M. "NA" stands for "not assayed." The enzyme inhibition curve for compound I-7 against CaMKII δ is shown in **Figure 1**.

Table 3. Results of *in vitro* CaMKII activity inhibition assays.

Compound ID	CaMKII δ IC ₅₀
I-1	B


Compound ID	CaMKII δ IC ₅₀
I-2	D
I-3	D
I-4	B
I-5	C
I-6	B
I-7	A
I-8	A
I-9	A
I-10	A
I-11	A
I-12	A
I-13	A
I-14	A

[00294] While we have described a number of embodiments of this invention, it is apparent that our basic examples may be altered to provide other embodiments that utilize the compounds and methods of this invention. Therefore, it will be appreciated that the scope of this invention is to be defined by the appended claims rather than by the specific embodiments that have been represented by way of example.

CLAIMS

We claim:

1. A compound of formula **I**:



I

or a pharmaceutically acceptable salt thereof, wherein:

one of R^{2'} and R^{3'} is -L¹-R^x, and the other is hydrogen;

L¹ is a covalent bond or a straight or branched C₁₋₆ aliphatic group, wherein one or more methylene groups are independently and optionally replaced by -NR^a- or -O-;


R^x is selected from the group consisting of NH₂, guanidino, 4-7 membered optionally substituted saturated heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen or sulfur, and 5-6 membered heteroaromatic ring having 1-2 heteroatoms independently selected from sulfur, nitrogen and oxygen;

each of R³, R⁴, R⁵, R⁶, R⁷, R⁸, R^{4'}, R^{5'}, and R^{6'} is independently selected from the group consisting of hydrogen, halogen, -CN, -CF₃, -OR, -NR₂, -NO₂, -COOR, -CONR₂, and -R;

each R^a is independently hydrogen or C₁₋₃ aliphatic; and

each R is independently hydrogen or optionally substituted C₁₋₆ aliphatic.

2. The compound of claim 1 of formula **I-a**:



I-a

or a pharmaceutically acceptable salt thereof, wherein:

one of R^2 and R^3 is $-L^1-R^x$, and the other is hydrogen;

L^1 is a covalent bond or a straight or branched C_{1-6} aliphatic group, wherein one or more methylene groups are independently and optionally replaced by $-NR^a-$ or $-O-$;


R^x is selected from the group consisting of NH_2 , guanidino, 4-7 membered optionally substituted saturated heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen or sulfur, and 5-6 membered heteroaromatic ring having 1-2 heteroatoms independently selected from sulfur, nitrogen and oxygen;

each of R^6 and R^7 is independently selected from the group consisting of hydrogen, halogen, $-CN$, $-CF_3$, $-OR$, $-NR_2$, $-NO_2$, $-COOR$, $-CONR_2$, and $-R$;

each R^a is independently hydrogen or C_{1-3} aliphatic; and

each R is independently hydrogen or optionally substituted C_{1-6} aliphatic.

3. The compound of claim 1 of formula **II**:

**II**

or a pharmaceutically acceptable salt thereof; wherein

L^1 is a covalent bond or a straight or branched C_{1-6} aliphatic group, wherein one or more methylene groups are independently and optionally replaced by $-NR^a-$ or $-O-$;


R^x is selected from the group consisting of NH_2 , guanidino, 4-7 membered optionally substituted saturated heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen or sulfur, and 5-6 membered heteroaromatic ring having 1-2 heteroatoms independently selected from sulfur, nitrogen and oxygen;

each of R^6 and R^7 is independently selected from the group consisting of hydrogen, halogen, $-CN$, $-CF_3$, $-OR$, $-NR_2$, $-NO_2$, $-COOR$, $-CONR_2$, and $-R$;

each R^a is independently hydrogen or C_{1-3} aliphatic; and

each R is independently hydrogen or optionally substituted C₁₋₆ aliphatic.

4. The compound of claim 1 of formula **III**:



III

or a pharmaceutically acceptable salt thereof; wherein

L¹ is a covalent bond or a straight or branched C₁₋₆ aliphatic group, wherein one or more methylene groups are independently and optionally replaced by -NR^a- or -O-;


R^x is selected from the group consisting of NH₂, guanidino, 4-7 membered optionally substituted saturated heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen or sulfur, and 5-6 membered heteroaromatic ring having 1-2 heteroatoms independently selected from sulfur, nitrogen and oxygen;

each of R⁶ and R⁷ is independently selected from the group consisting of hydrogen, halogen, -CN, -CF₃, -OR, -NR₂, -NO₂, -COOR, -CONR₂, and -R;

each R^a is independently hydrogen or C₁₋₃ aliphatic; and

each R is independently hydrogen or optionally substituted C₁₋₆ aliphatic.

5. The compound of claim 3 of formula **IV**:




IV

or a pharmaceutically acceptable salt thereof; wherein


R^x is selected from the group consisting of NH_2 , guanidino, 4-7 membered optionally substituted saturated heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen or sulfur, and 5-6 membered heteroaromatic ring having 1-2 heteroatoms independently selected from sulfur, nitrogen and oxygen; each of R^6 and R^7 is independently selected from the group consisting of hydrogen, halogen, $-CN$, $-CF_3$, $-OR$, $-NR_2$, $-NO_2$, $-COOR$, $-CONR_2$, and $-R$; each R^a is independently hydrogen or C_{1-3} aliphatic; and each R is independently hydrogen or optionally substituted C_{1-6} aliphatic.

6. The compound of claim 4 of formula **V**:



or a pharmaceutically acceptable salt thereof; wherein
 R^x is selected from the group consisting of NH_2 , guanidino, 4-7 membered optionally substituted saturated heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen or sulfur, and 5-6 membered heteroaromatic ring having 1-2 heteroatoms independently selected from sulfur, nitrogen and oxygen; each of R^6 and R^7 is independently selected from the group consisting of hydrogen, halogen, $-CN$, $-CF_3$, $-OR$, $-NR_2$, $-NO_2$, $-COOR$, $-CONR_2$, and $-R$; each R^a is independently hydrogen or C_{1-3} aliphatic; and each R is independently hydrogen or optionally substituted C_{1-6} aliphatic.


7. The compound of claim 3 of formula **VI**:

**VI**

or a pharmaceutically acceptable salt thereof; wherein

R^x is selected from the group consisting of NH₂, guanidino, 4-7 membered optionally substituted saturated heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen or sulfur, and 5-6 membered heteroaromatic ring having 1-2 heteroatoms independently selected from sulfur, nitrogen and oxygen; each of R⁶ and R⁷ is independently selected from the group consisting of hydrogen, halogen, -CN, -CF₃, -OR, -NR₂, -NO₂, -COOR, -CONR₂, and -R; and each R is independently hydrogen or optionally substituted C₁₋₆ aliphatic.


8. The compound of claim 4 of formula **VII**:

**VII**

or a pharmaceutically acceptable salt thereof; wherein

R^x is selected from the group consisting of NH₂, guanidino, 4-7 membered optionally substituted saturated heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen or sulfur, and 5-6 membered heteroaromatic ring having 1-2 heteroatoms independently selected from sulfur, nitrogen and oxygen; each of R⁶ and R⁷ is independently selected from the group consisting of hydrogen, halogen, -CN, -CF₃, -OR, -NR₂, -NO₂, -COOR, -CONR₂, and -R; and each R is independently hydrogen or optionally substituted C₁₋₆ aliphatic.

9. The compound of claim 3 of formula **VIII**:




VIII

or a pharmaceutically acceptable salt thereof; wherein

R^x is selected from the group consisting of NH₂, guanidino, 4-7 membered optionally substituted saturated heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen or sulfur, and 5-6 membered heteroaromatic ring having 1-2 heteroatoms independently selected from sulfur, nitrogen and oxygen;
 each of R⁶ and R⁷ is independently selected from the group consisting of hydrogen, halogen, -CN, -CF₃, -OR, -NR₂, -NO₂, -COOR, -CONR₂, and -R; and
 each R is independently hydrogen of optionally substituted C₁₋₆ aliphatic.

10. The compound of claim 4 of formula **IX**:




IX

or a pharmaceutically acceptable salt thereof; wherein


R^x is selected from the group consisting of NH₂, guanidino, 4-7 membered optionally substituted saturated heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen or sulfur, and 5-6 membered heteroaromatic ring having 1-2 heteroatoms independently selected from sulfur, nitrogen and oxygen;
 each of R⁶ and R⁷ is independently selected from the group consisting of hydrogen, halogen, -CN, -CF₃, -OR, -NR₂, -NO₂, -COOR, -CONR₂, and -R; and
 each R is independently hydrogen of optionally substituted C₁₋₆ aliphatic.

11. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein the compound is selected from the following structures:



12. A pharmaceutical composition comprising a compound according to claim 1 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, adjuvant, or vehicle.

13. A method of synthesizing a compound of the formula **I-a**:



wherein

one of R^{2'} and R^{3'} is -L¹-R^x, and the other is hydrogen;

L¹ is a covalent bond or a straight or branched C₁₋₆ aliphatic group, wherein one or more methylene groups are independently and optionally replaced by -NR^a- or -O-;

R^x is selected from the group consisting of NH₂, guanidino, 4-7 membered optionally substituted saturated heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen or sulfur, and 5-6 membered heteroaromatic ring having 1-2 heteroatoms independently selected from sulfur, nitrogen and oxygen;

each of R⁶ and R⁷ is independently selected from the group consisting of hydrogen,


halogen, -CN, -CF₃, -OR, -NR₂, -NO₂, -COOR, -CONR₂, and -R;

each R^a is independently hydrogen or C₁₋₃ aliphatic; and

each R is independently hydrogen or optionally substituted C₁₋₆ aliphatic;


comprising:

1) contacting a compound of the formula:



R⁶ and R⁷ are defined as above;

with a compound of formula:



R^{2'}-P and R^{3'}-P are as defined for R^{2'} and R^{3'} above with a primary amine protecting group on any primary amine; and

3) removing protecting group P.

14. A method of treating a cardiovascular disease, disorder, or condition, an inflammatory disease, disorder or condition, a neurological disease, disorder or condition, an ocular disease, disorder or condition, a metabolic disease, disorder or condition, a cancer or other proliferative disease, disorder or condition, a bone disease, disorder or condition, or an addictive disease, disorder, or condition, comprising administering to a patient in need thereof the composition according to claim 12.

15. The method of claim 14 wherein the cardiovascular disease, disorder, or condition is selected from atrial fibrillation, ventricular arrhythmia, heart failure, cardiac hypertrophy, atherosclerosis, restenosis; or cardiotoxicity arising from drug therapy, heart attack, ischemia-reperfusion injury, or catecholaminergic polymorphic ventricular tachycardia.

16. The method of claim 14 wherein the inflammatory disease, disorder or condition is asthma or rheumatoid arthritis.

17. The method of claim 14 wherein the neurological disease, disorder or condition is pain or stroke.

18. The method of claim 14 wherein the metabolic disease, disorder, or condition is diabetes.

19. The method of claim 14 wherein the cancer or other proliferative disease, disorder or condition is an osteosarcoma, a melanoma, or a prostate cancer.

20. The method of claim 14 wherein the addictive disease, disorder, or condition is opioid tolerance or dependence.




Figure 1. CaMKII δ enzyme inhibition curve for compound I-7.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2014/020700

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 31/437 (2014.01)

USPC - 514/279

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8) - A61K 31/00, 31/437; C07D 471/00 (2014.01)

USPC - 514/183, 247, 279; 546/26

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

CPC - C07D 401/04, 471/04 (2014.02)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Orbit, STN, Google Patents, Google Scholar, Google, PubChem

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2007/0004684 A1 (SENNHENN et al) 04 January 2007 (04.01.2007) entire document	1-20
A	US 2011/0281862 A1 (GAMBACORTI-PASSERINI et al) 17 November 2011 (17.11.2011) entire document	1-20
A	US 2006/0057639 A1 (PITT et al) 16 March 2006 (16.03.2006) entire document	1-20
A	US 2009/0005356 A1 (BLANEY et al) 01 January 2009 (01.01.2009) entire document	1-20

Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

“A” document defining the general state of the art which is not considered to be of particular relevance

“E” earlier application or patent but published on or after the international filing date

“L” document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

“O” document referring to an oral disclosure, use, exhibition or other means

“P” document published prior to the international filing date but later than the priority date claimed

“T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

“X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

“Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

“&” document member of the same patent family

Date of the actual completion of the international search

13 May 2014

Date of mailing of the international search report

23 MAY 2014

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Blaine R. Copenheaver

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774