

(19) DANMARK

(10) DK/EP 1997481 T3



(12)

Oversættelse af
europæisk patent

Patent- og
Varemærkestyrelsen

(51) Int.Cl.: **A 61 K 41/00 (2006.01)** **A 61 K 39/395 (2006.01)** **A 61 J 3/00 (2006.01)** **A 61 K 9/20 (2006.01)**

(45) Oversættelsen bekendtgjort den: **2017-12-11**

(80) Dato for Den Europæiske Patentmyndigheds
bekendtgørelse om meddelelse af patentet: **2017-09-13**

(86) Europæisk ansøgning nr.: **06799634.8**

(86) Europæisk indleveringsdag: **2006-05-16**

(87) Den europæiske ansøgnings publiceringsdag: **2008-12-03**

(86) International ansøgning nr.: **RU2006000237**

(87) Internationalt publikationsnr.: **WO2007105981**

(30) Prioritet: **2006-03-13 RU 2006107580**

(84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI SK TR**

(73) Patenthaver: **Epshtein, Oleg Iliich, 4 Samotyochny Per., d.3, kv. 72, Moscow 127473, Russiske Føderation**

(72) Opfinder: **Epshtein, Oleg Iliich, 4 Samotyochny Per., d.3, kv. 72, Moscow 127473, Russiske Føderation**

(74) Fuldmægtig i Danmark: **Patrade A/S, Fredens Torv 3A, 8000 Århus C, Danmark**

(54) Benævnelse: **Fast, oral form af et medicinsk præparat og en fremgangsmåde til fremstilling deraf**

(56) Fremdragne publikationer:

EP-A1- 1 295 606

EP-A1- 1 340 493

EP-A1- 1 466 622

EP-A1- 1 550 460

WO-A1-00/32189

RU-C1- 2 182 008

RU-C1- 2 201 254

RU-C2- 2 161 481

SU-A3- 1 836 083

US-B1- 6 248 355

US-B1- 6 326 360

**ABECASSIS J/GAYOT A: "2-1/2-2" In: NEGTIEN G ET AL: "MEDICAMENTS HOMEOPATIQUES", 1986,
MEDICAMENTS HOMEOPATHIQUES. NOTIONS PRATIQUES DE PHARMACIE HOMEOPATIQUE; [GALENICA],
PARIS, TECHNIQUE ET DOCUMENTATION, FR, XP002162252, pages 77-99, * the whole document ***

DK/EP 1997481 T3

DESCRIPTION

[0001] The invention relates to the field of medicine and can be used for technologically simple production of the solid oral form of an antibody-based drug for effective treatment of pathological syndrome without pronounced side effects.

[0002] From prior art, antibody-based drugs are known (sera, immunoglobulins), which are used in therapeutic doses (for example, see Russian Drug Register (in Rus.), Drug Encyclopedia, 7th Ed., 2000, pp. 358-359). However, these drugs are prepared in liquid formulation suitable for injections and are introduced parenterally, since this is the only method of introduction providing bioavailability of these drugs.

[0003] In addition, the field of application of these drugs is limited to etiological treatment of predominantly infectious diseases; their usage could be associated with adverse side effects.

[0004] A method is also known for producing a solid form of a drug, which includes pressing of dry ground components containing active substance and pharmaceutically acceptable additives (RU 2203054 C2, A61K9/20, 2003).

[0005] EP1295606 discloses a pharmaceutical agent for treating a pathological syndrome containing activated forms of ultra-low doses of monoclonal, polyclonal or natural antibodies to an antigen, wherein said activated form is prepared by means of repeated consecutive dilution and external treatment, predominantly based on homeopathic technology.

[0006] Abecassis J/Gayot A, "2-1/2-2", Abecassis J/ Gayot A, Negtien G et al., Medicaments homeopathiques. Notions pratiques de pharmacie homeopathique, Galenica, Paris, Technique de documentation, FR (1986), pages 77 - 99, discloses preparation techniques for homeopathic medicines.

[0007] The invention is aimed at the development of a solid form of an antibody-based drug suitable for oral administration, and a technologically simple method for producing such antibody-based drug in activated form, prepared using homeopathic technique in the form of tablets.

[0008] The subject matter for which protection is sought is as set out in the claims.

[0009] In particular, the present invention provides a method for producing a solid oral form of an antibody-based drug, comprising

1. (a) preparing an aqueous-alcoholic dilution of antibodies in an activated form by combining repeated successive dilutions and external action using homeopathic technique, wherein the antibodies are to an endogenous compound;
2. (b) irrigating a neutral carrier with the aqueous-alcoholic dilution of antibodies in a

- fluidized boiling bed with concurrent drying;
- 3. (c) mixing with pharmaceutically acceptable additives in pharmaceutically acceptable sequence; and
- 4. (d) subsequent tablet formation using direct pressing.

[0010] Solution of the problem is provided by the fact that the solid oral form of the antibody based drug comprises an effective quantity of a carrier, irrigated using a fluidized boiling bed with concurrent drying with the aqueous-alcoholic dilution of antibodies in activated form prepared by combining repeated successive dilution of antibodies with external action and pharmaceutically acceptable additives.

[0011] In this case, aqueous-alcoholic dilution contains the activated form of antibodies to endogenous compound participating in regulation or affecting the mechanisms of formation of pathological syndrome, wherein the activated form is prepared by way of repeated dilution and external action using homeopathic technique.

[0012] It is possible to use mechanical shaking or ultrasonic or electromagnetic treatment of the dilutions as an external action.

[0013] In addition, the drug contains a neutral substance - lactose as a carrier, and a binder and a lubricant as pharmaceutically acceptable additives.

[0014] In this case, as pharmaceutically acceptable additives, the solid oral form additionally contains a neutral carrier - lactose in the amount of 30 to 80 wt. % of the mass of the solid oral form - a tablet, and as a binder, it contains microcrystalline cellulose in the amount of 10.0 to 15.0 wt. % of the mass of the solid oral form - a tablet, and as a lubricant, it contains magnesium stearate in the amount of 0.8 to 1.2 wt. % of the mass of the solid oral form - a tablet.

[0015] The activated form of antibodies constitutes a mixture of various decimal and/or centesimal aqueous-alcoholic homeopathic dilutions.

[0016] The solution of the problem is also achieved by the fact that the method for producing the solid oral form of the antibody-based drug comprises preparation of an aqueous-alcoholic dilution of antibodies in activated form, obtained by combining repeated successive dilutions and external action using homeopathic technique, irrigating of a neutral carrier with this aqueous-alcoholic dilution in a fluidized boiling bed with concurrent drying at the temperature up to 35°C, mixing with pharmaceutically acceptable additives in pharmaceutically acceptable sequence, and subsequent pelletization - tablet formation using direct dry pressing.

[0017] In this case, a neutral substance - lactose is used as a carrier with the particle size between 150 and 250 µm.

[0018] Introduction of antibodies in activated form to the solid drug formulation according to the proposed technique provides a possibility of oral administration of the antibody-based drug while preserving a biological activity.

[0019] Utilization of a mixture of various centesimal alcoholic homeopathic dilutions as activated form of antibodies increases therapeutic efficacy of the proposed drug under condition of individual tolerance (insensibility) of the body with respect to a certain dilution.

[0020] Besides, the qualitative, quantitative, granulometric and structural compositions of the proposed ingredients (components) ensure reliable tablet formation using direct dry pressing while utilizing a limited number of additives.

[0021] In order to prepare a pharmaceutical substance, polyclonal antibodies, monoclonal antibodies and natural antibodies to the substance participating in regulation or affecting the mechanisms of pathological syndrome formation are used.

[0022] Methods for obtaining antibodies are described, for example, in the source: Immunological methods (in Rus.), edited by G. Frimel, Moscow, "Meditina", 1987, pp.9-33.

[0023] Polyclonal antibodies, specifically binding with compounds of different categories, e.g., proteins, polynucleotides, oligosaccharides, glycolipids, etc., as well as interacting with low molecular weight substances (haptens), are produced as a result of active immunization of animals. For this purpose, animals are given a series of injections according to a specifically developed schedule with antigen, which represents either an individually isolated high-molecular substance, or a synthetic conjugate - in case of using haptens. As a result of conducting such procedure, a monospecific antiserum with high antibody content is obtained for further use. If necessary, purification of antibodies present in antiserum is conducted. For this purpose, fractionation with salt precipitation or ion exchange chromatography are used.

[0024] Monoclonal antibodies of various specificity, which interact with low molecular weight haptens as well as with epitopes of high-molecular substances, are obtained using hybridoma technique. The initial stage of the process includes immunization based on principles already developed during preparation of polyclonal antisera. Further stages of work provide for obtaining of hybrid cells producing clones of similar in specificity antibodies. Their individual isolation is done using the same methods as in the case of polyclonal antisera.

[0025] Natural antibodies to exogenous antigens and bioregulators of different nature are isolated from the human blood serum using affinity chromatography. For this purpose, a carrier with covalently bound antigen, which represents either a hapten or a macromolecular compound, is used as immunosorbent. As a result of conducting chromatography, a population of antibodies with narrow distribution in specificity and affinity is obtained.

[0026] Isolated substance or drug antibodies undergo successive repeated dilution in combination with external action, which is predominantly a vertical shaking, until an activated

form is obtained, for example, using homeopathic potentiation technique (see, for example, V. Shvabe, Homeopathic drugs (in Rus.). A description and preparation guide, Moscow, 1967, pp. 12 - 38; or G. Keller, Homeopathy (in Rus.), Moscow, "Meditina", 2000, part 1, pp. 37 - 40). In this case, uniform decrease in concentration is achieved by gradually diluting of 1 volumetric part of initial substance (antibodies) in 9 volumetric parts (for decimal dilution D) or in 99 volumetric parts (for centesimal dilution C), or in 999 volumetric parts (for millesimal dilution) of a neutral solvent - distilled water and/or 70 % ethyl alcohol followed by repeated vertical shaking of each obtained dilution as well as utilization of predominantly separate containers for each subsequent dilution until desired concentration is achieved.

[0027] External action in the process of concentration reduction can also be conducted using ultrasound or electromagnetic action, as well as other mechanical action.

[0028] During the next stage, in order to produce a pharmaceutical substance - an active substance in the form of "saturated" lactose, irrigating of granules of the neutral substance - lactose (milk sugar) with the particle size of 150 - 250 μm , introduced into the fluidized bed, with the activated form of antibodies prepared using a aqueous-alcoholic dilution (preferably, centesimal) with concurrent drying at temperatures up to 35°C is conducted using a fluidized bed apparatus, e.g., "Hüttlin Pilotlab" manufactured by Hüttlin GmbH.

[0029] Calculated amount of prepared pharmaceutical substance - active substance in the form of "saturated" lactose - is loaded into a mixer and mixed with microcrystalline cellulose introduced in the amount of 10.0 - 15.0 wt. % of the total mass of the load. Then, an "unsaturated" lactose is added to the mixture (as required in order to reduce cost and somewhat simplify and speed-up technological process without decline in efficacy of the therapeutic treatment due to reduction in pharmaceutical substance content in the tablet: a aqueous-alcoholic dilution of antibodies in activated form) in the amount of 30 - 80 wt. % of the total mass of the load, and magnesium stearate in the amount of 0.8 - 1.2 wt. % of the total mass of the load, followed by uniform mixing.

[0030] Obtained dry homogeneous mixture is supplied to the pelletizing machine, for example, tablet-press Korsch - XL 400, to form round 150 - 500 mg tablets using direct dry pressing.

[0031] Described below is an example of producing a drug as an activated form of the polyclonal morphine antibodies (antisera), the process of which is conducted in several stages.

[0032] The stage of producing a morphine-ovalbumin conjugate.

[0033] A solution of 50 mg (0.001 mmol) of ovalbumin in 5.0 ml of distilled water is mixed with 2.0 ml of dimethylformamide containing 15.0 mg (0.039 mmol) of morphine 6hemisuccinate, to which, while cooling down, a solution of 15 mg (0.055 mmol) of water-soluble carbodiamide in 3 ml of distilled water is added drop wise. Reaction mixture is incubated for 5 hours at 4°C. Obtained conjugate is isolated using gelchromatography on a column with Sephadex G25 and

liophilicly dried.

[0034] The amount of bound morphine is calculated based on UV-spectra of the initial protein and obtained conjugate using the change in absorption at 280 nm. Based on UF-spectra data, synthesized conjugate contains 12 - 15 moles of hapten per one mole of protein.

[0035] The stage of producing morphine-ovalbumin conjugate monospecific serum.

[0036] Immunization of "Vienna blue" rabbits weighing no more than 2 kg is conducted on the cycling basis with 10-day interval. The maximum number of injection is 4. Conjugate is administered to the front and rear extremities in the area of articular lymph nodes in the amount of 1 mg per immunization. For this purpose, the antigen is preliminary dissolved in 1 ml of complete Freund's adjuvant. The volume of immunization mixture is 2 ml.

[0037] Further immunizations are done using incomplete Freund's adjuvant while maintaining above-specified proportions of antigen and adjuvant. Trial blood sample is taken from the animal's peripheral ear vein 10 days after immunization.

[0038] Rabbit blood serum is obtained by centrifuging at 1000g for 10 minutes at room temperature followed by adding chloroform as a preservative until a final concentration of 13% is reached.

[0039] Obtained antiserum is checked using immune-enzyme analysis for presence of morphine-specific antibodies, which are determined using enzyme-marked antispecies antibody conjugate.

[0040] So obtained antiserum contains specific antibodies in dilution of 1 : 1 000 - 1 : 25 000.

[0041] Next, a g-globulin-rich fraction is isolated from the produced antiserum. For this purpose, protein precipitation with 50% ammonium sulfate is used followed by washing the residue with 30% saline solution, centrifuging and conducting dialysis against phosphate buffer. So prepared fraction, containing given antigen-specific antibodies, is used during the next stage to prepare the drug.

[0042] The stage of producing an activated form of morphine antibodies.

[0043] 0.5 ml of g-globulin fraction of antiserum are placed into a E-61 container, followed by adding 4.5 ml of distilled water and shaking 10 times, which results in 5 ml of the 1st centesimal dilution. Then, 0.05 ml of the 1st centesimal dilution are placed into E-62 container, followed by adding 4.95 ml of distilled water and shaking 10 times, which results in 5.0 ml of the 2nd centesimal dilution. The 3rd through the 29th centesimal dilutions are prepared similar to the 2nd centesimal dilution.

[0044] The final 30th centesimal dilution is obtained by diluting the 29th dilution in 70% ethyl

alcohol.

[0045] Obtained aqueous-alcoholic solution is used for irrigating in a fluidized bed the particles of a neutral carrier - lactose (milk-sugar) when producing a pharmaceutical substance of active compound in the form of "saturated" lactose, which serves as a basis for direct dry press-forming of the solid oral form (round tablets with the mass predominantly 240 and 300 mg) of the proposed drug.

[0046] Provided below are the examples of treating various pathological syndromes using a solid form of the drug based on activated form of antibodies, which are conditionally (based on similarity with the terminology used in homeopathic literature) called potentiated antibodies.

Example 1.

[0047] Patient V., 19 years old. Administers himself with heroin by inhalation at least 3 times a week for 2 months. Based on parental consent, the patient was hospitalized and placed in isolated unit for 24 days. 2 days after admission the patient became irritable and developed sleep disturbance. During conversation, mentioned that he noticed attraction to narcotics. Prescribed: 1 tablet "saturated" with potentiated antibodies - homeopathic dilution C30 of the g-globulin fraction of antiserum to morphine hydrochloride - 6 times a day. 3 weeks later the patient demonstrated steady mood, satisfactory appetite and sleep. During individual session with psychologist, the patient stated that he experienced no attraction. Recommended to continue taking heroin antibodies on a daily basis at 1 tablet a day. 2 months after discharging from the hospital, according to his mother, the patient does not take any narcotics.

Example 2.

[0048] Patient K., 57 years old, suffering from rheumatoid arthritis (RA) for 5 years, class III according to functional classification of the American College of Rheumatology, and was hospitalized due to exacerbation of the disease. Upon admission: complaints about fever, considerable increase in morning stiffness and pain in affected joints, their inflammation. Objectively: temperature 37.5°C, pronounced hyperemia and defiguration of wrist, ankle and proximal interphalangeal joints, pain during palpation.

In blood test: ESR 35 mm/h, rheumatoid factor++. Due to poor tolerance of non-steroid anti-inflammatory drugs, prescribed: 1 tablet "saturated" with potentiated antibodies - a mixture of homeopathic dilutions C50, C200 (1 : 1) of monoclonal antibodies to recombinant human tumor necrosis factor alpha - 3 times a day. 3 days after the beginning of treatment, the patient observed considerable reduction in pain syndrome, body temperature dropped to normothermia. On the 7th day of treatment: morning stiffness observed before hospitalization still remains. The patient was discharged on the 14th day with the clinical-laboratory remission. Recommended prophylactic administration of the drug by taking 1 tablet every other day. 2

months after discharge, the patient's class is changed from III to II according to RA functional classification.

Example 3.

[0049] Patient S., 51 years old, came to see urologist with complains about decrease in libido, erection abnormality, lower satisfaction from sexual intercourse. Mentioned symptoms intensified over the past 2 years. During the last 3 years also observed occasional depression, grizzle, memory impairment and sleep disturbance, decrease in ability to work, palpitation attacks, fluctuations in arterial blood pressure. Objectively: revealed moderate enlargement of prostate gland. Diagnosis: erectile dysfunction with involutional hormonal changes. Prescribed: 1 tablet "saturated" with potentiated antibodies - a mixture of homeopathic dilutions of monoclonal antibodies to a fragment of human endothelium NO synthase D6, C30 and C200 (1 : 1 : 1) - once every 3 days. 2 weeks after the beginning of treatment, the patient demonstrated improvement in erection and increase in libido along with general improvement in health condition: higher overall tonus, better sleep. Recommended taking medicine 1-2 times a week. During follow-up visit 2 months after the beginning of treatment, the patient expressed no previous complaints, observed recovery in libido level, erection and satisfaction from the intercourse.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- RU2203054C2 [0004]
- EP1295606A [0005]

Non-patent literature cited in the description

- Russian Drug RegisterDrug Encyclopedia20000000358-359 [0002]

- **ABECASSIS JGAYOT ANEGTIEN G et al.** Medicaments homeopathiques. Notions pratiques de pharmacie homeopathique Technique de documentation, 1986, 77-99 [0006]
- MeditsinalImmunological methods 198700009-33 [0022]
- **V. SHVABEA** description and preparation guide Homeopathic drugs, 1967, 12-38 [0026]
- **G. KELLER** Meditsina Homeopathy, 2000, 37-40 [0026]

Patentkrav

1. Fremgangsmåde til fremstilling af en fast, oral form af et antistofbaseret lægemiddel, omfattende
 - 5 a) fremstilling af en vandig-alkoholisk fortynding af antistoffer i en aktiveret form ved kombination af gentagne, successive fortyndinger og ekstern virkning ved anvendelse af homøopatisk teknik, hvor antistofferne er til en endogen forbindelse;
 - 10 b) skyllning af en neutral bærer med den vandige-alkoholiske fortynding af antistoffer i et fluidiseret, kogende leje med samtidig tørring;
 - c) blanding med farmaceutisk acceptable tilsetningsstoffer i farmaceutisk acceptabel sekvens; og
 - d) efterfølgende tablet dannelse ved brug af direkte presning.
- 15 2. Fremgangsmåde ifølge krav 1, hvor laktose med en partikelstørrelse fra 150 til 250 μm er anvendt som en bærer.