

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 March 2010 (25.03.2010)

(10) International Publication Number  
WO 2010/033248 A2

(51) International Patent Classification:  
C12N 15/113 (2010.01) A61K 31/7088 (2006.01)

(74) Agent: LOCKHART, Helen, C.; Wolf, Greenfield & Sacks, P.C., Federal Reserve Plaza, 600 Atlantic Avenue, Boston, MA 02210-2206 (US).

(21) International Application Number:  
PCT/US2009/005251

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:  
22 September 2009 (22.09.2009)

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(25) Filing Language: English

Published:

(26) Publication Language: English

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(30) Priority Data:  
61/192,954 22 September 2008 (22.09.2008) US  
61/149,946 4 February 2009 (04.02.2009) US  
61/224,031 8 July 2009 (08.07.2009) US

— with sequence listing part of description (Rule 5.2(a))

(71) Applicant (for all designated States except US): RXI PHARMACEUTICALS CORPORATION [US/US]; 60 Prescott Street, Worcester, MA 01605 (US).

(72) Inventors; and

(71) Applicants : KHVOROVA, Anastasia [US/US]; 4550 Squires Circle, Boulder, CO 80305 (US). SALOMON, William [US/US]; 39 O Street, Apt. #1, Boston, MA 02127 (US). KAMENS, Joanne [US/US]; 21 Wessex Road, Newton, MA 02459 (US). SAMARSKY, Dmitry [US/US]; 15 Wayside Road, Westborough, MA 01581 (US). WOOLF, Tod, M. [US/US]; 14 Babe Ruth Drive, Sudbury, MA 01776-1942 (US). MILLER, Michelle [US/US]; 1302 Village Road East, Norwood, MA 02062 (US). BULOCK, Karen, G. [US/US]; 44 Park Street, Mendon, MA 01756 (US).

(54) Title: NEUTRAL NANOTRANSPORTERS




Figure 11

(57) Abstract: Neutral lipid formulations for nucleic acid delivery are provided according to the invention. The neutral lipid formulations include hydrophobically modified polynucleotides and fat mixtures. Methods of using the neutral lipid formulations are also provided.

## NEUTRAL NANOTRANSPORTERS

## RELATED APPLICATIONS

This application claims the benefit under 35 U.S.C. § 119(e) of U.S. provisional

5 application serial number US 61/192,954, entitled "Chemically Modified Nucleotides and Methods of Using the Same," filed on September 22, 2008, US 61/149,946, entitled "Minimum Length Triggers of RNA Interference," filed on February 4, 2009, and US 61/224,031, entitled "Minimum Length Triggers of RNA Interference," filed on July 8, 2009, the disclosure of each of which is incorporated by reference herein in its entirety.

## 10 FIELD OF INVENTION

This invention pertains to non-charged lipid formulations for nucleic acid delivery.

## BACKGROUND OF INVENTION

Liposome-based formulations are widely used for nucleic acid delivery. Most commercially available lipid or liposome formulations contain at least one positively charged 15 lipid. It has been assumed that the presence of this positively charged lipid is essential for obtaining a high degree of nucleic acid loading and enhancement of liposome fusogenic properties. Several screens have been performed previously to identify optimal positively charged lipid chemistries but all the formulations that have been developed are characterized by the major issues of high levels of toxicity. In fact, *in vivo* limited therapeutic indexes have been 20 reported for liposome formulations containing positively charged lipids at concentrations only slightly higher than concentrations required to achieve silencing. It would be of great benefit to develop non-toxic delivery vehicles for nucleic acids.

## SUMMARY OF INVENTION

Described herein is a novel approach to nucleic acid delivery which enables efficient 25 loading of nucleic acids into neutral fat formulations generating a new class of delivery vehicles. The nucleic acid molecule is modified to increase its hydrophobicity and is mixed with neutral fat formulation, producing efficient encapsulation of nucleic acid molecules in neutral lipid particles. Methods and compositions described herein have widespread applications for *in vivo* delivery of nucleic acids.

30 A composition is provided according to an aspect of the invention. The composition includes a hydrophobic modified polynucleotide; a neutral fatty mixture; and optionally a cargo molecule. The hydrophobic modified polynucleotide and the neutral fatty mixture form a

micelle. In one embodiment the polynucleotide is RNA, such as single stranded or double stranded RNA.

In another embodiment the composition is a pharmaceutical composition and includes a pharmaceutically acceptable carrier. In yet another embodiment the composition is sterile.

5 The neutral fatty mixture is one or more fats which form a non-toxic mixture. In some embodiments the neutral fatty mixture includes a DOPC (dioleoylphosphatidylcholine) and/or a DSPC (distearoylphosphatidylcholine). In other embodiments the neutral fatty mixture further includes a sterol, such as, for instance, cholesterol. In yet other embodiments the neutral fatty mixture includes 20% of a fatty acid derivative of choline. The neutral fatty mixture may 10 also include 20% sterol. In some embodiments the composition includes at least 20% DOPC and at least 20% cholesterol.

15 The hydrophobic portion of the hydrophobic modified polynucleotide may be covalently or non-covalently linked to the polynucleotide. In some embodiments the hydrophobic portion of the hydrophobic modified polynucleotide is a sterol, such as, for instance, a cholesterol, a cholesteryl and/or modified cholesteryl residue. In other embodiments the hydrophobic portion of the hydrophobic modified polynucleotide is one or more of bile acids, cholic acid or taurocholic acid, deoxycholate, oleyl lithocholic acid, oleyl cholenic acid, glycolipids, phospholipids, sphingolipids, isoprenoids, vitamins, saturated fatty acids, unsaturated fatty acids, fatty acid esters, triglycerides, pyrenes, porphyrines, Texaphyrine, adamantane, acridines, biotin, 20 coumarin, fluorescein, rhodamine, Texas-Red, digoxigenin, dimethoxytrityl, t-butyldimethylsilyl, t-butyldiphenylsilyl, cyanine dyes (e.g. Cy3 or Cy5), Hoechst 33258 dye, psoralen, and ibuprofen. In yet other embodiments the hydrophobic portion of the hydrophobic modified polynucleotide is a polycationic molecule, such as, for instance, protamine, arginine rich peptides, and/or spermine.

25 The composition optionally includes a cargo molecule such as a lipid, a peptide, vitamin, and/or a small molecule. In some embodiments the cargo molecule is a commercially available fat emulsions available for a variety of purposes selected from the group consisting of parenteral feeding. In some embodiments the commercially available fat emulsion is an intralipid or a nutralipid. In other embodiments the cargo molecule is a fatty acid mixture containing more 30 than 74% of linoleic acid, a fatty acid mixture containing at least 6% of cardiolipin, or a fatty acid mixture containing at least 74% of linoleic acid and at least 6% of cardiolipin. In another

embodiment the cargo molecule is a fusogenic lipid, such as for example, DOPE, and preferably is at least 10% fusogenic lipid

In some embodiments the polynucleotide includes chemical modifications. For instance it may be at least 40% modified.

5 In other embodiments the polynucleotide is an isolated double stranded nucleic acid molecule having a guide strand, wherein the guide strand is 16-28 nucleotides long and has complementarity to a target gene, wherein the 3' terminal 10 nucleotides of the guide strand include at least two phosphate modifications, and wherein the guide strand has a 5' phosphate modification and includes at least one 2' O-methyl modification or 2'-fluoro modification, and a  
10 10 passenger strand, wherein the passenger strand is 8-28 nucleotides long and has complementarity to the guide strand, wherein the passenger strand is linked to the hydrophobic molecule, wherein the guide strand and the passenger strand form the double stranded nucleic acid molecule.

15 In yet another embodiment the polynucleotide is an isolated double stranded nucleic acid molecule having a guide strand and a passenger strand, wherein the guide strand is from 16-29 nucleotides long and is substantially complementary to a target gene, wherein the passenger strand is from 8-14 nucleotides long and has complementarity to the guide strand, and wherein the guide stand has at least two chemical modifications.

20 The polynucleotide in other embodiments is an isolated double stranded nucleic acid molecule having a guide strand and a passenger strand, wherein the guide strand is from 16-29 nucleotides long and is substantially complementary to a target gene, wherein the passenger strand is from 8-14 nucleotides long and has complementarity to the guide strand, and wherein the guide stand has a single stranded 3' region that is 5 nucleotides or longer.

25 In yet another embodiment the polynucleotide is an isolated double stranded nucleic acid molecule having a guide strand and a passenger strand, wherein the guide strand is from 16-25 nucleotides long and is substantially complementary to a target gene, wherein the passenger strand is from 16-25 nucleotides long and has complementarity to the guide strand, and wherein the double stranded nucleic acid molecule optionally has 3' overhangs.

30 The polynucleotide in other embodiments is an isolated double stranded nucleic acid molecule having a guide strand and a passenger strand, wherein the guide strand is from 24-29 nucleotides long and is substantially complementary to a target gene, wherein the passenger

strand is from 24-29 nucleotides long and has complementarity to the guide strand, and wherein the double stranded nucleic acid molecule optionally has 3' overhangs.

The polynucleotide in some embodiments is an isolated single stranded nucleic acid molecule of 16-29 nucleotides in length and is substantially complementary to a target gene.

5 The polynucleotide may be for instance an antisense ODN, an antagonir, an antiMirs, or a PNA.

In some embodiments the ratio of neutral fatty mixture to polynucleotide is 1:5.

In other embodiments the composition is composed of about 50% cargo molecule.

The composition may have a particle size of 10-140nm.

10 In preferred embodiments the composition is free of cationic lipids.

A composition of a hydrophobic modified polynucleotide; and a neutral fatty mixture is provided in other aspects of the invention. The hydrophobic modified polynucleotide and the neutral fatty mixture form a micelle and wherein between 40-100% of the nucleotides of the polynucleotide are chemically modified nucleotides.

15 An isolated single stranded polynucleotide having at least one phosphorothioate bond and one base modified with 4-pyridyl, 2-pyridyl is provided in other aspects of the invention. In some embodiments all ribose moieties of the polynucleotide are modified. In other embodiments at least 40% of nucleotides of the polynucleotide are chemically modified.

An isolated hydrophobic modified polynucleotide having a sterol-type molecule attached 20 to a polynucleotide, wherein the sterol-type molecule has a polycarbon chain length of 3 - 7 or 9-18 carbons is provided in other aspects of the invention. In some embodiments the sterol-type molecule has a polycarbon chain length of 7.

In another aspect the invention is a method for delivering an polynucleotide to a cell, by contacting a cell with a composition or a polynucleotide of the invention to deliver the 25 polynucleotide to the cell. The method may be performed *in vitro* or *in vivo*.

A method of inducing RNAi in a cell is provided according to other aspects of the invention. The method involves contacting a cell with a composition or a polynucleotide of the invention, wherein the polynucleotide has at least a region of sequence correspondence to a target gene and wherein the polynucleotide induces RNAi of mRNA of the target gene.

30 A method of inducing RNAi in a subject by administering to a subject an effective amount for inducing RNAi of mRNA of a target gene, a composition or a polynucleotide of the invention is provided in other aspects. The polynucleotide has at least a region of sequence

correspondence to the target gene. In some embodiments the subject is a human. In other embodiments the target gene is PPIB, MAP4K4, and/or SOD1.

A method is also provided, in which alteration of lipid content and ratio of the composition of the invention is used to alter pharmacokinetic behavior and tissue distribution of 5 polynucleotides.

In another aspect the invention is a method for delivering a polynucleotide to a target tissue of a subject, by administering to a subject a composition of the invention wherein the composition includes a fatty acid in the neutral fatty mixture or cargo molecule associated with targeting to the target tissue in order to deliver the polynucleotide to the target tissue. In one 10 embodiment the target tissue is cardiomyocytes and the lipid is optionally cardiolipin or linoleic acid. In other embodiments the target tissue is lung, fat or liver.

A method is also provided, in which alteration of lipid content and ratio of the composition of the invention is used to optimize pharmacokinetic behavior and tissue distribution of polynucleotides to cardiomyocytes.

15 A method is also provided, in which alteration of lipid content and ratio of the composition of the invention is used to alter pharmacokinetic behavior and tissue distribution of polynucleotides to fat tissue.

20 A method is also provided, in which alteration of lipid content and ratio of the composition of the invention is used to alter pharmacokinetic behavior and tissue distribution of polynucleotides to lung tissue.

A method is also provided, in which alteration of lipid content and ratio of the composition of the invention is used to alter pharmacokinetic behavior and tissue distribution of polynucleotides to liver.

25 In other aspects the invention is a method of inducing RNAi in a subject by administering to a subject an effective amount for inducing RNAi of mRNA of a target gene, a composition or a polynucleotide of the invention, wherein the polynucleotide has at least a region of sequence correspondence to the target gene, wherein the step of administering is systemic, intravenous, intraperitoneal, intradermal, topical, intranasal, inhalation, oral, intramucosal or intraocular.

30 The invention in other aspects is an isolated hydrophobic modified polynucleotide having a polynucleotide, wherein the polynucleotide is double stranded RNA, attached to a hydrophobic molecule, wherein the hydrophobic molecule is attached to a base, a ribose or a

backbone of a non-terminal nucleotide and wherein the isolated double stranded nucleic acid molecule comprises a guide strand and a passenger strand, wherein the guide strand is from 16-29 nucleotides long and is substantially complementary to a target gene, wherein the passenger strand is from 8-14 nucleotides long and has complementarity to the guide strand. In some 5 embodiments the hydrophobic molecule is attached to the guide strand of the double stranded RNA. In other embodiments the 3' terminal 10 nucleotides of the guide strand include at least two phosphate modifications, and wherein the guide strand has a 5' phosphate modification and includes at least one 2' O-methyl modification or 2'-fluoro modification. In yet other embodiments the hydrophobic molecule is attached to the passenger strand of the double 10 stranded RNA.

In another aspect the invention is an isolated hydrophobic modified polynucleotide having a polynucleotide non-covalently complexed to a hydrophobic molecule, wherein the hydrophobic molecule is a polycationic molecule. The polycationic molecule is protamine, arginine rich peptides, and/or spermine in some embodiments.

15 In yet another aspect the invention is an isolated hydrophobic modified polynucleotide having a polynucleotide, wherein the polynucleotide is double stranded RNA, directly complexed to a hydrophobic molecule without a linker, wherein the hydrophobic molecule is not cholesterol.

Thus, the present invention provides in some aspects and embodiments asymmetric 20 dsRNA molecules. In one embodiment, dsRNA binding domain type proteins are utilized to mask the negative charges of a polynucleotide duplex as a way to facilitate cellular entry for conventional siRNAs and asymmetric dsRNAs. In another embodiment, protamine or other arginine rich peptides are used to mask the negative charges of a polynucleotide duplex as a way 25 to facilitate cellular entry of conventional siRNAs and asymmetric dsRNAs. In another embodiment, spermine (or spermidine type moieties) are used to mask the negative charges of a polynucleotide duplex as a way to facilitate cellular entry of conventional siRNAs and asymmetric dsRNAs.

The present invention in other aspects provides asymmetric dsRNAs molecules, where 30 one or both polynucleotides contain nucleotides where the 5'-position contains hydrophobic modifications selected from the group consisting of (4-pyridyl, 2-pyridyl, indolyl, phenyl (C<sub>6</sub>H<sub>5</sub>OH); tryptophanyl (C<sub>8</sub>H<sub>6</sub>N)CH<sub>2</sub>CH(NH<sub>2</sub>)CO), isobutyl, butyl, aminobenzyl; phenyl; naphthyl, and the like. In a particular embodiment, the asymmetric dsRNA molecule may be in

combination with PS modifications. In another particular embodiment, at least 40% of the asymmetric dsRNA molecule may be chemically-modified.

The present invention also provides single stranded RISC entering and RISC inhibiting modalities, containing hydrophobic modifications. In one embodiment, the single stranded polynucleotides include a combination of phosphorothioates hydrophobicly modified nucleotides and at least 40% of moiety is chemically modified with or without a conjugate. The chemical modification may increase nucleotide hydrophobicity in order to optimize RISC single stranded substrate entry or optimize substrate based inhibition of preloaded RISC complex.

While the chemistry of the compounds stays the same, the chemical modification pattern may favor either RISC entry or substrate inhibition

The present invention also provides a group of sterol type molecules with an extended position 17 attached poly- hydrocarbon tail. In one embodiment, the poly-hydrocarbon tail is 8, 9, 11 or more carbons. In a particular embodiment, the poly-hydrocarbon tail is less than 8 or more than 9 carbons. The chain may be branched. The chain may also be attached through a 3' or 5' or intermolecular linker. In a particular embodiment, at least 40% of the nucleotides are modified. In another particular embodiment, the duplex region contains at least two mismatches.

The present invention also provides methods of utilizing fat emulsions to alter the pharmacokinetic and/or tissue distribution via hydrophobicly modified polynucleotides. In one embodiment, a polynucleotide with increased hydrophobicity is provided in combination with an intralipid type formulation. In a particular embodiment, increased hydrophobicity is achieved by attaching a sterol type modalities, one or more lipids, and/or making one or more chemical modifications. In a particular embodiment, the intralipid type formulation contains more then 74% linoleic acid. In another particular embodiment, a sterol type molecule with a position 17 chain longer than 9 carbons in attached in combination with the use of one or more intralipid type formulations.

The present invention also provides a method of altering the intralipid composition in order to alter the tissue distribution of the polynucleotides of the present invention. In one embodiment, a polynucleotide with a sterol type molecule with a position 17 chain longer than 9 carbons in combination with intralipid formulation containing more then 7% of cardiolipin is provided. In a particular embodiment, the intralipid formulation contains more than 74% of linoleic acid. In another particular embodiment, polynucleotides are provided with sterol type

molecules with a position 17 chain longer than 9 carbons, with at least 40% of nucleotides being chemically modified in combination with intralipid formulation containing more than 74% of linoleic acid and 7% of cardiolipin.

Use of any of the compositions of the invention for inhibiting expression of a target gene 5 or an miRNA is also provided as an aspect of the invention.

A method for manufacturing a medicament of any of the compositions of the invention for inhibiting expression of a target gene in order to treat a disease is also provided.

A composition of an isolated polynucleotide of the invention for inhibiting expression of a target gene or an miRNA is also provided as an aspect of the invention.

10 Each of the limitations of the invention can encompass various embodiments of the invention. It is, therefore, anticipated that each of the limitations of the invention involving any one element or combinations of elements can be included in each aspect of the invention. This invention is not limited in its application to the details of construction and the arrangement of components set forth in the following description or illustrated in the drawings. The invention is 15 capable of other embodiments and of being practiced or of being carried out in various ways. Also, the phraseology and terminology used herein is for the purpose of description and should not be regarded as limiting. The use of "including," "comprising," or "having," "containing", "involving", and variations thereof herein, is meant to encompass the items listed thereafter and equivalents thereof as well as additional items.

20

#### BRIEF DESCRIPTION OF DRAWINGS

FIG. 1 is a schematic depicting proposed structures of asymmetric double stranded RNA molecules (adsRNA). Bold lines represent sequences carrying modification patterns compatible with RISC loading. Striped lines represent polynucleotides carrying modifications compatible 25 with passenger strands. Plain lines represent a single stranded polynucleotide with modification patterns optimized for cell interaction and uptake. FIG 1A depicts adsRNA with extended guide or passenger strands; FIG 1B depicts adsRNA with length variations of a cell penetrating polynucleotide; FIG 1C depicts adsRNA with 3' and 5' conjugates; FIG 1D depicts adsRNAs with mismatches.

30 FIG. 2 is a schematic depicting asymmetric dsRNA molecules with different chemical modification patterns. Several examples of chemical modifications that might be used to

increase hydrophobicity are shown including 4-pyridyl, 2-pyridyl, isobutyl and indolyl based position 5 uridine modifications.

FIG. 3 is a schematic depicting the use of dsRNA binding domains, protamine (or other Arg rich peptides), spermidine or similar chemical structures to block duplex charge to facilitate 5 cellular entry.

FIG. 4 is a schematic depicting positively charged chemicals that might be used for polynucleotide charge blockage.

FIG. 5 is a schematic depicting examples of structural and chemical compositions of single stranded RISC entering polynucleotides. The combination of one or more modifications 10 including 2'd, 2'Ome, 2'F, hydrophobic and phosphorothioate modifications can be used to optimize single strand entry into the RISC.

FIG. 6 is a schematic depicting examples of structural and chemical composition of RISC substrate inhibitors. Combinations of one or more chemical modifications can be used to mediate efficient uptake and efficient binding to preloaded RISC complex.

FIG. 7 is a schematic depicting structures of polynucleotides with sterol type molecules attached, where R represent a polycarbonic tail of 9 carbons or longer. FIG. 7A depicts an 15 adsRNA molecule; FIG. 7B depicts an siRNA molecule of approximately 17-30 bp long; FIG. 7C depicts a RISC entering strand; FIG 7D depicts a substrate analog strand. Chemical modification patterns, as depicted in FIG. 7, can be optimized to promote desired function.

FIG. 8 is a schematic depicting examples of naturally occurring phytosterols with a 20 polycarbon chain that is longer than 8, attached at position 17. More than 250 different types of phytosterols are known.

FIG. 9 is a schematic depicting examples of sterol-like structures, with variations in the size of the polycarbon chains attached at position 17.

FIG. 10 presents schematics and graphs demonstrating that the percentage of liver uptake 25 and plasma clearance of lipid emulsions containing sterol type molecules is directly affected by the size of the polycarbon chain attached at position 17. This figure is adapted from Martins et al, Journal of Lipid Research (1998).

FIG. 11 is a schematic depicting micelle formation. FIG. 11A depicts a polynucleotide 30 with a hydrophobic conjugate; FIG. 11B depicts linoleic acid; FIG. 11C depicts a micelle formed from a mixture of polynucleotides containing hydrophobic conjugates combined with fatty acids.

FIG. 12 is a schematic depicting how alteration in lipid composition can affect pharmacokinetic behavior and tissue distribution of hydrophobically modified and/or hydrophobically conjugated polynucleotides. In particular, use of lipid mixtures enriched in linoleic acid and cardiolipin results in preferential uptake by cardiomyocytes.

5 FIG. 13 presents a graph and schematics of RNAi compounds showing the chemical/structural composition of highly effective sd-rxRNA compounds. Highly effective compounds were found to have the following characteristics: antisense strands of 17-21 nucleotides, sense strands of 10-15 nucleotides, single-stranded regions that contained 2-12 phosphorothioate modifications, preferentially 6-8 phosphorothioate modifications, and sense strands in which the 10 majority of nucleotides were 2'OMe modified, with or without phosphorothioate modification. Any linker chemistry can be used to attach these molecules to hydrophobic moieties such as cholesterol at the 3' end of the sense strand. Version GIIa-b of these RNA compounds demonstrate that elimination of 2'F content has no impact on efficacy.

15 FIG. 14 presents a schematic demonstrating nucleic acids that can be formulated using methods described herein. Nucleic acids compatible with methods associated with the invention can include, for example, traditional siRNA, longer siRNAs (24-29), single stranded oligos, antisense, antogamirs and sd-rxRNA. The nucleic acid molecule is modified such that its hydrophobicity is substantially increased. This can be achieved by modifying bases, sugars or nucleic acid backbone or/and by linking a hydrophobic molecule to the nucleic acid. The 20 hydrophobic molecule can be attached anywhere in the compound and can include, for example, a fatty acid, sterol, vitamin, small molecule or peptide. The hydrophobic molecule can be covalently or non covalently attached.

25 FIG. 15 presents gels that demonstrate a lack of complex formation between hydrophobic oligonucleotides (sd-rxRNA) and neutral fat formulations. Panel A shows a lack of complex formation with a DOPC:DOPE mixture. Panel B demonstrate lack of complex formation with Intralipid. The complex formation is evaluated by complexing reagents and evaluating a shift in oligonucleotide band formation using a non-denaturing polyacrylamide gel. The position of the oligonucleotide is determined by staining.

30 FIG. 16 presents gels demonstrating that the simultaneous presence of DOPC and Cholesterol results in a hydrophobic nucleic acid complexing with neutral lipid formulations. As low as 1:5 weight (lipid to oligonucleotide) ratio was sufficient to produce a significant fraction of encapsulated oligonucleotide. In the context of DOPC:Cholesterol formulation, other neutral

lipids can be added without interfering with particle formation. For example, DOPC:cholesterol: DOPE.

FIG. 17 presents a gel demonstrating that DSPC (saturated fatty acids) may take place of DOPC without any impact on efficiency of complex formation

5 FIG. 18 presents a gel demonstrating DOPC/Intralipid/Cholesterol Formulation involving sd-rxRNA (12386). Complexes were still present when diluted to 500 nM from 1  $\mu$ M with Accell media.

10 FIG. 19 presents a gel demonstrating DOPC/Intralipid/Cholesterol Formulation involving sd-rxRNA (12386). Complexes were still found to be present when diluted to 500 nM from 5  $\mu$ M with Accell media. More RNA appeared to be complexed here than when diluting to 500 nM from 1  $\mu$ M.

15 FIG. 20 presents gels demonstrating that the presence of both DOPC and Cholesterol in significant proportions is required to enable significant formation of Intralipid-containing neutral fat/oligo particles. A slight complex formation is observed with Intralipid:cholesterol but it is minor.

20 FIG. 21 presents gels demonstrating that various additional compounds (lipids, peptides, small molecules) might be encapsulated into the particle as long as formulation comprises at least 20% of DOPC/Cholesterol-type compounds. The demonstrated cargo lipid is intralipid. Variation in the identity, amounts and ratios of cargo lipids will affect cellular uptake and tissue 25 distribution characteristics of these compounds. For example, the length of lipid tails and level of saturability will affect differential uptake to liver, lung, fat and cardiomyocytes. Addition of special hydrophobic molecules like vitamins or different forms of sterols can favor distribution to special tissues which are involved in the metabolism of particular compounds. FIG. 21 also demonstrates that complexes are formed at different oligonucleotide concentrations, with higher concentrations favoring more efficient complex formation.

FIG. 22 presents gels demonstrating that more efficient complex formation is observed at higher oligonucleotide concentrations.

30 FIG. 23 presents a gel demonstrating that only hydrophobically modified oligonucleotides are complexed with neutral fat formulations. When a cholesterol-modified sd-rxRNA compound was mixed with 50:50 DOPC/Cholesterol formulation, the sd-rxRNA quantitatively entered into a complex. When rxRNA (Omethyl modified siRNA), regular siRNA or sd-rxRNA, without a hydrophobic compound is mixed with the same formulation, no

complex was formed, demonstrating that the combination of a hydrophobic modification of the oligonucleotide with neutral fat formulations comprising at least 20% of DOPC:Cholesterol-type compounds are required for efficient encapsulation of oligos into neutral fat formulations.

FIG. 24 presents a fluorescent image and a graph. FIG. 24A demonstrates that neutral 5 lipid based formulations enter cells (HeLa) and effectively silence genes. FIG. 24B demonstrates that the oligonucleotide/lipid ratio and formulation composition affects the level of silencing. Significantly, no toxicity was observed even at 1 uM concentration. This lack of toxicity is a significant improvement over positively charge traditional formulations (i.e., lipofectamine) which exhibit a drastic toxicity at a much lower dose range. This data 10 demonstrates that neutral fat/oligonucleotide formulations are non toxic or have highly reduced toxicity relative to previously described positively charged formulations, and have a wider therapeutic index.

FIG. 25 presents a graph demonstrating various the efficacies of various formulations in vitro with and without serum. Formulation B is DOPC-Cholesterol 50:50; Formulation C is 15 DOPC-DOPE:Cholesterol 33:33:33.

FIG. 26 presents a table describing the peak sizes and zeta potential of RNA:Lipid and DOPC:Chol complexes.

FIG. 27 presents graphs demonstrating size and zeta potential of particles formed upon 20 complexing of neutral fat formulations with hydrophobically modified oligos. While neutral fat by itself forms agglomerates ~ 500-1000 nm in size, addition of increasing concentrations of an oligonucleotide results in the formation of stable small particles (around 60-120nm) which are not charged (Zeta Potential ~ -10). The neutral particles, sized around 50-100 nM, are ideal for systemic administration. The size and charge of the particles is affected by the oligonucleotide/lipid ratio, lipid mixture composition and lipid ratios within formulation.

25 FIG. 28 shows a panel of lipids used for formulation preparations.

FIG. 29 presents schematics demonstrating sterol-type molecules used for formulation 30 preparations. In some instances, some of the formulations comprising longer chain sterol type molecules have a significantly better cellular uptake and tissue distribution properties

FIG. 30 presents schematics demonstrating some hydrophobic molecules which can be linked to an oligonucleotide or included as part of a formulation to improve or alter cellular uptake and tissue distribution.

FIG. 31 presents evidence that the presence of a hydrophobic conjugate is sufficient to get an oligonucleotide formulated in a neutral fat formulation. In particular, sd-rxRNA and cholesterol conjugated siRNA (labeled as Alnylam) both form effective complexes, while the non hydrophobically modified oligo does not. It is worth noting that sd-rxRNA complex 5 incorporation was more efficient with sd-rxRNA rather than regular cholesterol conjugated siRNA. It may be due to better hydrophobicity of the compound.

FIG. 32 presents evidence of efficient incorporation of hydrophobic oligos into neutral fat formulations at high concentrations (160 uM). This concentration enables oligo 10 administration at 20 ng/ kg in vivo. The Figure further demonstrates that use of sonication might be beneficial for improving the particle size and distribution.

FIG. 33 presents data demonstrating that DSPC is more effective in formation of hydrophobic oligonucleotide containing DOPE containing particles than DOPC. The DSPC and DOPC are highly similar molecule with only difference of the single double bone. Apparently 15 these compounds are different in ability to tolerate DOPE as a cargo molecule. We expect that optimization of the fatty acid structure of choline derivative is important for optimizing the formulation for different cargo load and tissue distribution properties.

FIG. 34 presents data showing a highly effective complex formation using DOPC:Retinol and Cholesterol. Surprisingly the presence of Retinol improved the complex 20 formation and is expected to change tissue distribution profiles. This data further support the notion that optimizing of the chemical composition of the formulation is important.

FIG. 35 presents data demonstrating that a variety of sterol type molecules are effectively incorporated in neutral fat formulations. Interestingly use of cholestryloleate actually improved the extent of the complex formation. It is expected that chemical diversity of sterol 25 type compounds is important for cargo loading and tissue distribution and cellular uptake.

FIG. 36 demonstrates efficient incorporation of tocopherol in hydrophobic oligo: neutral fat formulations.

FIG. 37 is a bar graph of data demonstrating efficient cellular uptake and lack of toxicity for variety of neutral fat formulations

FIG. 38 is a bar graph of data demonstrating that different type of hydrophobic 30 oligonucleotides work better with different type of neutral formulations. For example 13766 was more efficacious in a presence of DOPC based formulations compare to DSPC based formulation, while 12884 has better efficacy in DSPC based formulations as compare to DOPC

based formulations. The 12884 has additional hydrophobic molecule attached to the 5' end of the passenger strand in addition to conventional sd-rxRNA structures described previously.

#### DETAILED DESCRIPTION

Neutral fat-based formulations for the efficient delivery of oligonucleotides is disclosed herein. Liposome based formulations are widely used for oligonucleotide delivery. However, most of commercially available lipid or liposome formulations contain at least one positively charged lipid (cationic lipids). The presence of this positively charged lipid is believed to be essential for obtaining a high degree of oligonucleotide loading and for enhancing liposome fusogenic properties. Several methods have been performed and published to identify optimal positively charged lipid chemistries. However, the commercially available liposome formulations containing cationic lipids are characterized by a high level of toxicity. In vivo limited therapeutic indexes have revealed that liposome formulations containing positive charged lipids are associated with toxicity (i.e. elevation in liver enzymes) at concentrations only slightly higher than concentration required to achieve RNA silencing.

The compositions provided herein are referred to as neutral nanotransporters because they enable quantitative oligonucleotide incorporation into non-charged lipids mixtures. The lack of toxic levels of cationic lipids in the neutral nanotransporter compositions of the invention is an important feature.

The neutral nanotransporters compositions of the invention enable efficient loading of oligonucleotide into neutral fat formulation. The composition includes an oligonucleotide that is modified in a manner such that the hydrophobicity of the molecule is increased (for example a hydrophobic molecule is attached (covalently or no-covalently) to a hydrophobic molecule on the oligonucleotide terminus or a non-terminal nucleotide, base, sugar, or backbone), the modified oligonucleotide being mixed with a neutral fat formulation (for example containing at least 25 % of cholesterol and 25% of DOPC or analogs thereof). A cargo molecule, such as another lipid, peptide, vitamin, polymer or small molecule can also be included in the composition. This composition, where part of the formulation is build into the oligonucleotide itself enables efficient encapsulation of oligonucleotide in neutral lipid particles. Any oligonucleotides may be used in the compositions of the invention, For example, the oligonucleotides may be RNA, DNA, single stranded, double stranded etc. As shown in the

examples several sd-rxRNA compounds as well as non-sd-rxRNA compounds have been synthesized and formulated according to the invention.

One of several unexpected observations associated with the invention was that oligonucleotides could effectively be incorporated in a lipid mixture that was free of cationic 5 lipids and that such a composition could effectively deliver the therapeutic oligonucleotide to a cell in a manner that it is functional. Another unexpected observation was the high level of activity observed when the fatty mixture is composed of a phosphatidylcholine base fatty acid and a sterol such as a cholesterol. For instance, one preferred formulation of neutral fatty mixture is composed of at least 20% of DOPC or DSPC and at least 20% of sterol such as 10 cholesterol. Even as low as 1:5 lipid to oligonucleotide ratio was shown to be sufficient to get complete encapsulation of the oligonucleotide in a non charged formulation. The prior art demonstrated only a 1-5% oligonucleotide encapsulation with non-charged formulations, which is not sufficient to get to a desired amount of in vivo efficacy. Compared to the prior art using neutral lipids the level of oligonucleotide delivery to a cell was quite unexpected.

15 As shown in the Examples below stable particles ranging in size from 50 to 140 nm were formed upon complexing of hydrophobic oligonucleotides with preferred formulations. It is interesting to mention that the formulation by itself typically does not form small particles, but rather, forms agglomerates, which are transformed into stable 50-120 nm particles upon addition of the hydrophobic modified oligonucleotide.

20 *Compositions*

The compositions of the invention include a hydrophobic modified polynucleotide, a neutral fatty mixture, and optionally a cargo molecule. A “hydrophobic modified 25 polynucleotide” as used herein is a polynucleotide (described below) that has at least one modification that renders the polynucleotide more hydrophobic than the polynucleotide was prior to modification. The modification may be achieved by attaching (covalently or non-covalently) a hydrophobic molecule to the polynucleotide. In some instances the hydrophobic molecule is or includes a lipophilic group.

The term “lipophilic group” means a group that has a higher affinity for lipids than its affinity for water. Examples of lipophilic groups include, but are not limited to, cholesterol, a 30 cholesteryl or modified cholesteryl residue, adamantine, dihydrotestosterone, long chain alkyl, long chain alkenyl, long chain alkynyl, oleyl-lithocholic, cholenic, oleoyl-cholenic, palmityl, heptadecyl, myrisityl, bile acids, cholic acid or taurocholic acid, deoxycholate, oleyl lithocholic

acid, oleoyl cholenic acid, glycolipids, phospholipids, sphingolipids, isoprenoids, such as steroids, vitamins, such as vitamin E, fatty acids either saturated or unsaturated, fatty acid esters, such as triglycerides, pyrenes, porphyrines, Texaphyrine, adamantane, acridines, biotin, coumarin, fluorescein, rhodamine, Texas-Red, digoxigenin, dimethoxytrityl, t-

5 butyldimethylsilyl, t-butyldiphenylsilyl, cyanine dyes (e.g. Cy3 or Cy5), Hoechst 33258 dye, psoralen, or ibuprofen. The cholesterol moiety may be reduced (e.g. as in cholestan) or may be substituted (e.g. by halogen). A combination of different lipophilic groups in one molecule is also possible.

The hydrophobic molecule may be attached at various positions of the polynucleotide.

10 As described above, the hydrophobic molecule may be linked to the terminal residue of the polynucleotide such as the 3' of 5'-end of the polynucleotide. Alternatively, it may be linked to an internal nucleotide or a nucleotide on a branch of the polynucleotide. The hydrophobic molecule may be attached, for instance to a 2'-position of the nucleotide. The hydrophobic molecule may also be linked to the heterocyclic base, the sugar or the backbone of a nucleotide  
15 of the polynucleotide.

The hydrophobic molecule may be connected to the polynucleotide by a linker moiety. Optionally the linker moiety is a non-nucleotidic linker moiety. Non-nucleotidic linkers are e.g. abasic residues (dSpacer), oligoethyleneglycol, such as triethyleneglycol (spacer 9) or hexaethyleneglycol (spacer 18), or alkane-diol, such as butanediol. The spacer units are

20 preferably linked by phosphodiester or phosphorothioate bonds. The linker units may appear just once in the molecule or may be incorporated several times, e.g. via phosphodiester, phosphorothioate, methylphosphonate, or amide linkages.

Typical conjugation protocols involve the synthesis of polynucleotides bearing an aminolinker at one or more positions of the sequence, however, a linker is not required. The  
25 amino group is then reacted with the molecule being conjugated using appropriate coupling or activating reagents. The conjugation reaction may be performed either with the polynucleotide still bound to a solid support or following cleavage of the polynucleotide in solution phase. Purification of the modified polynucleotide by HPLC typically results in a pure material.

In some embodiments the hydrophobic molecule is a sterol type conjugate, a PhytoSterol  
30 conjugate, cholesterol conjugate, sterol type conjugate with altered side chain length, fatty acid conjugate, any other hydrophobic group conjugate, and/or hydrophobic modifications of the internal nucleoside, which provide sufficient hydrophobicity to be incorporated into micelles.

For purposes of the present invention, the term “sterols”, refers or steroid alcohols are a subgroup of steroids with a hydroxyl group at the 3-position of the A-ring. They are amphipathic lipids synthesized from acetyl-coenzyme A via the HMG-CoA reductase pathway. The overall molecule is quite flat. The hydroxyl group on the A ring is polar. The rest of the aliphatic chain 5 is non-polar. Usually sterols are considered to have an 8 carbon chain at position 17

For purposes of the present invention, the term “sterol type molecules”, refers or steroid alcohols, which are similar in structure to sterols. The main difference is the structure of the ring and number of carbons in a position 17 attached side chain.

For purposes of the present invention, the term “PhytoSterols” (also called plant sterols) 10 are a group of steroid alcohols, phytochemicals naturally occurring in plants. There are more than 200 different known PhytoSterols

For purposes of the present invention, the term “Sterol side chain” refers to a chemical composition of a side chain attached at the position 17 of sterol-type molecule. In a standard definition sterols are limited to a 4 ring structure carrying a 8 carbon chain at position 17. In this 15 invention, the sterol type molecules with side chain longer and shorter than conventional are described. The side chain may branched or contain double back bones.

Thus, sterols useful in the invention, for example, include cholesterols, as well as unique sterols in which position 17 has attached side chain of 2-7 or longer then 9 carbons. In a particular embodiment, the length of the polycarbon tail is varied between 5 and 9 carbons. 20 Figure 9 demonstrates that there is a correlation between plasma clearance, liver uptake and the length of the polycarbon chain. Such conjugates may have significantly better in vivo efficacy, in particular delivery to liver. These types of molecules are expected to work at concentrations 5 to 9 fold lower then oligonucleotides conjugated to conventional cholesterols.

Alternatively the polynucleotide may be bound to a protein, peptide or positively charged 25 chemical that functions as the hydrophobic molecule. Examples are shown in Figure 4. The proteins may be selected from the group consisting of protamine, dsRNA binding domain, and arginine rich peptides. Exemplary positively charged chemicals include spermine, spermidine, cadaverine, and putrescine (Figure 5).

In another embodiment hydrophobic molecule conjugates may demonstrate even higher 30 efficacy when it is combined with optimal chemical modification patterns of the polynucleotide, containing but not limited to hydrophobic modifications.

In another embodiment the sterol type molecule may be a naturally occurring PhytoSterols such as those shown in Figure 8. The polycarbon chain may be longer than 9 and may be linear, branched and/or contain double bonds. Some PhytoSterol containing polynucleotide conjugates may be significantly more potent and active in delivery of 5 polynucleotides to various tissues. Some PhytoSterols may demonstrate tissue preference and thus be used as a way to delivery RNAi specifically to particular tissues.

The hydrophobic modified polynucleotide is mixed with a neutral fatty mixture to form a 10 micelle. The neutral fatty acid mixture is a mixture of fats that has a net neutral or slightly net negative charge at or around physiological pH that can form a micelle with the hydrophobic modified polynucleotide. For purposes of the present invention, the term "micelle" refers to a 15 small nanoparticle formed by a mixture of non charged fatty acids and phospholipids. The neutral fatty mixture may include cationic lipids as long as they are present in an amount that does not cause toxicity. In preferred embodiments the neutral fatty mixture is free of cationic lipids. A mixture that is free of cationic lipids is one that has less than 1% and preferably 0% of 15 the total lipid being cationic lipid. The term "cationic lipid" includes lipids and synthetic lipids having a net positive charge at or around physiological pH. The term "anionic lipid" includes lipids and synthetic lipids having a net negative charge at or around physiological pH.

The neutral fats bind to the oligonucleotides of the invention by a strong but non-covalent attraction (e.g., an electrostatic, van der Waals, pi-stacking, etc. interaction).

20 The neutral fat mixture may include formulations selected from a class of naturally occurring or chemically synthesized or modified saturated and unsaturated fatty acid residues. Fatty acids might exist in a form of triglycerides, diglycerides or individual fatty acids. In another embodiment the use of well-validated mixtures of fatty acids and/or fat emulsions currently used in pharmacology for parenteral nutrition may be utilized.

25 The neutral fatty mixture is preferably a mixture of a choline based fatty acid and a sterol. Choline based fatty acids include for instance, synthetic phosphocholine derivatives such as DDPC, DLPC, DMPC, DPPC, DSPC, DOPC, POPC, and DEPC. DOPC (chemical registry number 4235-95-4) is dioleoylphosphatidylcholine (also known as dielaidoylphosphatidylcholine, dioleoyl-PC, dioleoylphosphocholine, dioleoyl-sn-glycero-3-phosphocholine, dioleylphosphatidylcholine). DSPC (chemical registry number 816-94-4) is 30 distearoylphosphatidylcholine (also known as 1,2-Distearoyl-sn-Glycero-3-phosphocholine).

The sterol in the neutral fatty mixture may be for instance cholesterol. The neutral fatty mixture may be made up completely of a choline based fatty acid and a sterol or it may optionally include a cargo molecule. For instance, the neutral fatty mixture may have at least 20% or 25% fatty acid and 20% or 25% sterol.

5 For purposes of the present invention, the term “Fatty acids” relates to conventional description of fatty acid. They may exist as individual entities or in a form of two-and triglycerides. For purposes of the present invention, the term “fat emulsions” refers to safe fat formulations given intravenously to subjects who are unable to get enough fat in their diet. It is an emulsion of soy bean oil (or other naturally occurring oils) and egg phospholipids. Fat  
10 emulsions are being used for formulation of some insoluble anesthetics. In this disclosure, fat emulsions might be part of commercially available preparations like Intralipid, LipoSyn, Nutrilipid, modified commercial preparations, where they are enriched with particular fatty acids or fully de novo- formulated combinations of fatty acids and phospholipids.

15 In one embodiment, the cells to be contacted with an oligonucleotide composition of the invention are contacted with a mixture comprising the oligonucleotide and a mixture comprising a lipid, *e.g.*, one of the lipids or lipid compositions described supra for between about 12 hours to about 24 hours. In another embodiment, the cells to be contacted with an oligonucleotide composition are contacted with a mixture comprising the oligonucleotide and a mixture comprising a lipid, *e.g.*, one of the lipids or lipid compositions described supra for between  
20 about 1 and about five days. In one embodiment, the cells are contacted with a mixture comprising a lipid and the oligonucleotide for between about three days to as long as about 30 days. In another embodiment, a mixture comprising a lipid is left in contact with the cells for at least about five to about 20 days. In another embodiment, a mixture comprising a lipid is left in contact with the cells for at least about seven to about 15 days.

25 50%-60% of the formulation can optionally be any other lipid or molecule. Such a lipid or molecule is referred to herein as a cargo lipid or cargo molecule. Cargo molecules include but are not limited to intralipid, small molecules, fusogenic peptides or lipids or other small molecules might be added to alter cellular uptake, endosomal release or tissue distribution properties. The ability to tolerate cargo molecules is important for modulation of properties of  
30 these particles, if such properties are desirable. For instance the presence of some tissue specific metabolites might drastically alter tissue distribution profiles. For example use of Intralipid type

formulation enriched in shorter or longer fatty chains with various degrees of saturation affects tissue distribution profiles of these type of formulations (and their loads).

An example of a cargo lipid useful according to the invention is a fusogenic lipid. For instance, the zwitterionic lipid DOPE (chemical registry number 4004-5-1, 1,2-Dioleoyl-sn-5-Glycero-3-phosphoethanolamine) is a preferred cargo lipid.

Intralipid may be comprised of the following composition: 1 000 mL contain: purified soybean oil 90 g, purified egg phospholipids 12 g, glycerol anhydrous 22 g, water for injection q.s. ad 1 000 mL. pH is adjusted with sodium hydroxide to pH approximately 8. Energy content/L: 4.6 MJ (190 kcal). Osmolality (approx.): 300 mOsm/kg water. In another 10 embodiment fat emulsion is Liposyn that contains 5% safflower oil, 5% soybean oil, up to 1.2% egg phosphatides added as an emulsifier and 2.5% glycerin in water for injection. It may also contain sodium hydroxide for pH adjustment. pH 8.0 (6.0 - 9.0). Liposyn has an osmolarity of 276 m Osmol/liter (actual).

Variation in the identity, amounts and ratios of cargo lipids affects the cellular uptake 15 and tissue distribution characteristics of these compounds. For example, the length of lipid tails and level of saturability will affect differential uptake to liver, lung, fat and cardiomyocytes. Addition of special hydrophobic molecules like vitamins or different forms of sterols can favor distribution to special tissues which are involved in the metabolism of particular compounds. Complexes are formed at different oligonucleotide concentrations, with higher concentrations 20 favoring more efficient complex formation (Figs. 21-22).

In another embodiment, the fat emulsion is based on a mixture of lipids. Such lipids may include natural compounds, chemically synthesized compounds, purified fatty acids or any other lipids. In yet another embodiment the composition of fat emulsion is entirely artificial. In a particular embodiment, the fat emulsion is more than 70% linoleic acid. In yet another 25 particular embodiment the fat emulsion is at least 1% of cardiolipin. Linoleic acid (LA) is an unsaturated omega-6 fatty acid. It is a colorless liquid made of a carboxylic acid with an 18-carbon chain and two cis double bonds.

In yet another embodiment of the present invention, the alteration of the composition of the fat emulsion is used as a way to alter tissue distribution of hydrophobicly modified 30 polynucleotides. This methodology provides for the specific delivery of the polynucleotides to particular tissues (Figure 12).

In another embodiment the fat emulsions of the cargo molecule contain more than 70% of Linoleic acid (C18H32O2) and/or cardiolipin are used for specifically delivering RNAi to heart muscle.

Fat emulsions, like intralipid have been used before as a delivery formulation for some 5 non-water soluble drugs (such as Propofol, re-formulated as Diprivan). Unique features of the present invention include (a) the concept of combining modified polynucleotides with the hydrophobic compound(s), so it can be incorporated in the fat micelles and (b) mixing it with the fat emulsions to provide a reversible carrier. After injection into a blood stream, micelles usually bind to serum proteins, including albumin, HDL, LDL and other. This binding is 10 reversible and eventually the fat is absorbed by cells. The polynucleotide, incorporated as a part of the micelle will then be delivered closely to the surface of the cells. After that cellular uptake might be happening though variable mechanisms, including but not limited to sterol type delivery.

For purposes of the present invention, the term, "polynucleotide" includes any molecule 15 that is an organic polymer molecule composed of nucleotide monomers covalently bonded in a chain. DNA (deoxyribonucleic acid) and RNA (ribonucleic acid) are examples of polynucleotides. For purposes of the present invention, the term "polynucleotide" is used synonymously with oligonucleotide and nucleic acid. "Nucleotide" refers to a ribonucleotide or a deoxyribonucleotide or modified form thereof, as well as an analog thereof. Nucleotides 20 include species that comprise purines, e.g., adenine, hypoxanthine, guanine, and their derivatives and analogs, as well as pyrimidines, e.g., cytosine, uracil, thymine, and their derivatives and analogs. Preferably, a "nucleotide" comprises a cytosine, uracil, thymine, adenine, or guanine moiety.

Polynucleotides include any such primers, probes, and/or oligomer fragments. 25 Polynucleotides, include polydeoxyribonucleotides (containing 2-deoxy-D-ribose), to polyribonucleotides (containing D-ribose), and to any other type of polynucleotide which is an N glycoside of a purine or pyrimidine base, or modified purine or pyrimidine base. The term polynucleotide includes any type of nucleic acid and/or oligonucleotide. These terms refer only to the primary structure of the molecule. Thus, these terms include double- and single-stranded 30 DNA, as well as triple-, double-and single-stranded RNA. Exemplary RNA molecules include RNAi, siRNA, miRNA and siRNA inhibitors, single stranded substrates for RISC assembly and

the like. Some of these examples will be discussed in more detail below. Such discussion is exemplary only and is non-limiting.

For purposes of the present invention, the term "deoxynucleotide" refers to a nucleotide or polynucleotide lacking an OH group at the 2' or 3' position of a sugar moiety with appropriate bonding and/or 2',3' terminal dideoxy, instead having a hydrogen bonded to the 2' and/or 3' carbon.

For purposes of the present invention, the terms "deoxyribonucleotide" and "DNA" refer to a nucleotide or polynucleotide comprising at least one ribosyl moiety that has an H at its 2' position of a ribosyl moiety instead of an OH.

For purposes of the present invention, the term "gene" is defined to include both transcribed and non-transcribed elements. Thus, for instance, a gene can include any non-transcribed enhancer and/or promoter (i.e. genomic DNA) that plays a role in determining the level, timing, or tissue specificity of expression of a particular mRNA transcript or non-coding RNA. In addition, the 5' UTR, ORF, 3' UTR, introns, as well as non-coding RNAs such as 15 miRNAs, piRNAs, tRNAs, rRNAs, and more, are included as elements of a gene.

The hydrophobic modified nucleotide may be an siRNA which includes conventional siRNAs, sd-rxRNAs, asymmetric dsRNAs, single stranded RISC entering polynucleotides, and single stranded RISC inhibiting polynucleotides.

Aspects of the invention relate to isolated double stranded nucleic acid molecules comprising a guide (antisense) strand and a passenger (sense) strand. As used herein, the term "double-stranded" refers to one or more nucleic acid molecules in which at least a portion of the nucleomonomers are complementary and hydrogen bond to form a double-stranded region. As used herein, the term "duplex" includes the region of the double-stranded nucleic acid molecule(s) that is (are) hydrogen bonded to a complementary sequence. Double-stranded 25 oligonucleotides of the invention may comprise a nucleotide sequence that is sense to a target gene and a complementary sequence that is antisense to the target gene. The sense and antisense nucleotide sequences correspond to the target gene sequence, *e.g.*, are identical or are sufficiently identical to effect target gene inhibition (*e.g.*, are about at least about 98% identical, 96% identical, 94%, 90% identical, 85% identical, or 80% identical) to the target gene sequence.

For purposes of the present invention, the phrase "guide strand" as used herein, refers to a polynucleotide or region of a polynucleotide that is substantially (i.e., 80% or more) or 30 90% complementary to a target nucleic acid of interest and is capable of efficient loading into the

RISC complex. A guide strand may be single stranded or part of a duplex and may be comprised of a polynucleotide region that is RNA, DNA or chimeric RNA/DNA. For example, an antisense strand may be complementary, in whole or in part, to a molecule of messenger RNA, an RNA sequence that is not mRNA or sequence of DNA that is either coding or non-coding. The guide 5 strand can be modified with a diverse group of small molecules and/or conjugates and one of the embodiments is related to use of chemical modifications, which improve activity of single stranded guide strands..

For purposes of the present invention, the phrase "passenger strand" or "sense strand" refers to a polynucleotide or region that has the same nucleotide sequence, in whole or in part, as 10 a target nucleic acid such as a messenger RNA or a sequence of DNA. When a sequence is provided, by convention, unless otherwise indicated, it is the sense strand (or region), and the presence of the complementary antisense strand (or region) is implicit. It is also implied that the passenger strand is a second non-essential part of the duplex responsible for promotion of RISC entry and the one, which will be lost after initial RISC loading.

15 For example, double-stranded RNA (dsRNA), for instance, may be formulated according to the methods of the invention. In one embodiment, the invention provides a dsRNA molecule such as a conventional siRNA. Conventional ds siRNA can include a duplex structure of between 18 and 25 basepairs (e.g., 21 base pairs). In some embodiments, the dsRNAs include at least one strand that is at least 21 nt long. In other embodiments, the dsRNAs include at least 20 one strand that is at least 15, 16, 17, 18, 19, 20, or more contiguous nucleotides. In conventional siRNA, at least one end of the dsRNA has a single-stranded nucleotide overhang of 1 to 4, generally 1 or 2 nucleotides. Generally, the single-stranded overhang is located at the 3'-terminal end of the antisense strand or, alternatively, at the 3'-terminal end of the sense strand. The dsRNA may also have a blunt end, generally located at the 5'-end of the antisense strand.

25 The dsRNA may be cross-linked in some embodiments. Chemical linking of the two separate dsRNA strands may be achieved by any of a variety of well-known techniques, for example by introducing covalent, ionic or hydrogen bonds; hydrophobic interactions, van der Waals or stacking interactions; by means of metal-ion coordination, or through use of purine analogues. Such chemically linked dsRNAs are suitable for packaging in the association 30 complexes described herein. Generally, the chemical groups that can be used to modify the dsRNA include, without limitation, methylene blue; bifunctional groups, generally bis-(2-chloroethyl)amine; N-acetyl-N'-(p-glyoxylbenzoyl)cystamine; 4-thiouracil; and psoralen. In one

embodiment, the linker is a hexa-ethylene glycol linker. In this case, the dsRNA are produced by solid phase synthesis and the hexa-ethylene glycol linker is incorporated according to standard methods (e.g., Williams, D. J., and K. B. Hall, Biochem. (1996) 35:14665-14670). In a particular embodiment, the 5'-end of the antisense strand and the 3'-end of the sense strand are 5 chemically linked via a hexaethylene glycol linker. In another embodiment, at least one nucleotide of the dsRNA comprises a phosphorothioate or phosphorodithioate groups. The chemical bond at the ends of the dsRNA is generally formed by triple-helix bonds.

dsRNA formulated according to the invention also includes sd-rxRNA. sd-rxRNA refers to a class of RNA molecules described in a PCT application filed on even date entitled 10 "Minimum length triggers of RNA Interference" as well as the provisional applications to which the instant application claims priority, each of which is incorporated by reference. Briefly, sd-rxRNA are asymmetric nucleic acid molecules with a double stranded region of a minimal length such as 8-14 nucleotides, are effective in silencing gene expression. Molecules with such a short double stranded region have not previously been demonstrated to be effective in 15 mediating RNA interference. It had previously been assumed that that there must be a double stranded region of 19 nucleotides or greater. The molecules described herein are optimized through chemical modification, and in some instances through attachment of hydrophobic conjugates. These molecules are highly efficient in silencing of target gene expression and offer significant advantages over previously described RNAi molecules including high activity in the 20 presence of serum, efficient self delivery, compatibility with a wide variety of linkers, and reduced presence or complete absence of chemical modifications that are associated with toxicity.

For purposes of the present invention, the phrase "silencing" is defined as an RNAi-mediated or antisense mediated reduction in gene expression that can be measured by any 25 number of methods including PCR-based methods, Northern blot analysis, Branched DNA, western blot analysis, and other art recognized techniques.

For purposes of the present invention, the term "siRNA" and the phrase "short interfering RNA" refer to unimolecular nucleic acids and to nucleic acids comprised of two separate strands that are capable of performing RNAi and that have a duplex region that is from about 18 to 30 about 30 base pairs in length. Additionally, the term siRNA and the phrase "short interfering RNA" include nucleic acids that also contain moieties other than ribonucleotide moieties,

including, but not limited to, modified nucleotides, modified internucleotide linkages, non-nucleotides, deoxynucleotides and analogs of the aforementioned nucleotides.

For purposes of the present invention, the phrase "RNA interference" and the term "RNAi" are synonymous and refer to the process by which a single, double, or tripartite molecule (e.g. an siRNA, an shRNA, an miRNA, a piRNA) exerts an effect on a biological process by interacting with one or more components of the RNAi pathway including but not limited to Drosha, RISC, Dicer, etc. The process includes, but is not limited to, gene silencing by degrading mRNA, attenuating translation, interactions with tRNA, rRNA, hnRNA, cDNA and genomic DNA, inhibition of as well as methylation of DNA with ancillary proteins. In addition, molecules that modulate RNAi (e.g. siRNA, piRNA, or miRNA inhibitors) are included in the list of molecules that enhance the RNAi pathway (Tomari, Y. et al. *Genes Dev.* 2005, 19(5):517-29).

In certain embodiments, the double-stranded oligonucleotide of the invention is double-stranded over its entire length, *i.e.*, with no overhanging single-stranded sequence at either end of the molecule, *i.e.*, is blunt-ended. In other embodiments, the individual nucleic acid molecules can be of different lengths. In other words, a double-stranded oligonucleotide of the invention is not double-stranded over its entire length. For instance, when two separate nucleic acid molecules are used, one of the molecules, *e.g.*, the first molecule comprising an antisense sequence, can be longer than the second molecule hybridizing thereto (leaving a portion of the molecule single-stranded). Likewise, when a single nucleic acid molecule is used a portion of the molecule at either end can remain single-stranded.

For purposes of the present invention, the term "mismatch" includes a situation in where Watson-Crick base pairing does not take place between a nucleotide of a sense strand and a nucleotide of an antisense strand. An example of a mismatch would be an A across from a G, a C across from an A, a U across from a C, an A across from an A, a G across from a G, a C across from a C, and so on. Mismatches are also meant to include an basic residue across from a nucleotide or modified nucleotide, an acyclic residue across from a nucleotide or modified nucleotide, a gap, or an unpaired loop. In its broadest sense, a mismatch as used herein includes any alteration at a given position which decreases the thermodynamic stability at or in the vicinity of the position where the alteration appears, such that the thermodynamic stability of the duplex at the particular position is less than the thermodynamic stability of a Watson-Crick base pair at that position. Preferred mismatches include a G across from an A, and an A across from a

C. A particularly preferred mismatch comprises an A across from an A, G across from a G, C across from a C, and U across from a U.

In one embodiment, a double-stranded oligonucleotide of the invention contains mismatches and/or loops or bulges, but is double-stranded over at least about 70% of the length 5 of the oligonucleotide. In another embodiment, a double-stranded oligonucleotide of the invention is double-stranded over at least about 80% of the length of the oligonucleotide. In another embodiment, a double-stranded oligonucleotide of the invention is double-stranded over at least about 90%-95% of the length of the oligonucleotide. In another embodiment, a double-stranded oligonucleotide of the invention is double-stranded over at least about 96%-98% of the 10 length of the oligonucleotide. In certain embodiments, the double-stranded oligonucleotide of the invention contains at least or up to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 mismatches.

For purposes of the present invention, the term "complementary" refers to the ability of 15 polynucleotides to form base pairs with one another. Base pairs are typically formed by hydrogen bonds between nucleotide units in antiparallel polynucleotide strands or regions. Complementary polynucleotide strands or regions can base pair in the Watson-Crick manner (e.g., A to T, A to U, C to G, G to U), or in any other manner that allows for the formation of stable duplexes. The guide strand has complementarity to a target gene. Complementarity between the guide strand and the target gene may exist over any portion of the guide strand. 20 Complementarity as used herein may be perfect complementarity or less than perfect complementarity as long as the guide strand is sufficiently complementary to the target that it mediates RNAi. In some embodiments complementarity refers to less than 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, or 1% mismatch between the guide strand and the target. Perfect complementarity refers to 100% complementarity. Perfect complementarity refers to the 25 situation in which each nucleotide unit of one polynucleotide strand or region can hydrogen bond with each nucleotide unit of a second polynucleotide strand or region. Less than perfect complementarity refers to the situation in which some, but not all, nucleotide units of two strands or two regions can hydrogen bond with each other.

Thus the invention has the advantage of being able to tolerate sequence variations that 30 might be expected due to genetic mutation, strain polymorphism, or evolutionary divergence. For example, siRNA sequences with insertions, deletions, and single point mutations relative to the target sequence have also been found to be effective for inhibition. Moreover, not all

positions of a siRNA contribute equally to target recognition. Mismatches in the center of the siRNA are most critical and essentially abolish target RNA cleavage. Mismatches upstream of the center or upstream of the cleavage site referencing the antisense strand are tolerated but significantly reduce target RNA cleavage. Mismatches downstream of the center or cleavage site 5 referencing the antisense strand, preferably located near the 3' end of the antisense strand, e.g. 1, 2, 3, 4, 5 or 6 nucleotides from the 3' end of the antisense strand, are tolerated and reduce target RNA cleavage only slightly.

In contrast to single-stranded polynucleotides, duplex polynucleotides have been difficult to deliver to a cell as they have rigid structures and a large number of negative charges which 10 makes membrane transfer difficult. Unexpectedly, it was found that the polynucleotides of the present invention, although partially double-stranded, are recognized *in vivo* as single-stranded and, as such, are capable of efficiently being delivered across cell membranes. As a result the polynucleotides of the invention are capable in many instances of self delivery. Thus, the polynucleotides of the invention may be formulated in a manner similar to conventional RNAi 15 agents or they may be delivered to the cell or subject alone (or with non-delivery type carriers) and allowed to self deliver. In one embodiment of the present invention, self delivering asymmetric double-stranded RNA molecules are provided in which one portion of the molecule resembles a conventional RNA duplex and a second portion of the molecule is single stranded.

In one aspect of the invention, a longer duplex polynucleotide is provided, including a 20 first polynucleotide that ranges in size from about 16 to about 30 nucleotides; a second polynucleotide that ranges in size from about 26 to about 46 nucleotides, wherein the first polynucleotide (the antisense strand) is complementary to both the second polynucleotide (the sense strand) and a target gene, and wherein both polynucleotides form a duplex and wherein the first polynucleotide contains a single stranded region longer than 6 bases in length and is 25 modified with alternative chemical modification pattern, and/or includes a conjugate moiety that facilitates cellular delivery. In this embodiment, between about 40% to about 90% of the nucleotides of the passenger strand between about 40% to about 90% of the nucleotides of the guide strand, and between about 40% to about 90% of the nucleotides of the single stranded region of the first polynucleotide are chemically modified nucleotides.

30 In an embodiment, the chemically modified nucleotide in the polynucleotide duplex may be any chemically modified nucleotide known in the art, such as those discussed in detail above.

In a particular embodiment, the chemically modified nucleotide is selected from the group consisting of 2' F modified nucleotides, 2'-O-methyl modified and 2'deoxy nucleotides. In another particular embodiment, the chemically modified nucleotides results from "hydrophobic modifications" of the nucleotide base. In another particular embodiment, the chemically

5 modified nucleotides are phosphorothioates. In an additional particular embodiment, chemically modified nucleotides are combination of phosphorothioates, 2'-O-methyl, 2'deoxy, hydrophobic modifications and phosphorothioates. As these groups of modifications refer to modification of the ribose ring, back bone and nucleotide, it is feasible that some modified nucleotides will carry a combination of all three modification types.

10 In another embodiment, the chemical modification is not the same across the various regions of the duplex. In a particular embodiment, the first polynucleotide (the passenger strand), has a large number of diverse chemical modifications in various positions. For this polynucleotide up to 90% of nucleotides might be chemically modified and/or have mismatches introduced.

15 Single stranded modified and non modified RNA, DNA molecules may also be formulated according to the methods of the invention. An example of a single stranded polynucleotide that may be formulated according to the invention is those ODN described in PCT/US2009/004326. Briefly, these ssRNA can form double stranded structures based on internal interactions or on interactions with identical sequences. For instance they may include  
20 two identical single-stranded polynucleotides, each of the single-stranded polynucleotide comprising a 5'-stem sequence having a 5'-end, a 3'-stem sequence having a 3'-end, and a linker sequence linking the 5'-stem sequence and the 3'-stem sequence, wherein: (1) the 5'-stem sequence of a first single-stranded polynucleotide hybridize with the 3'-stem sequence of a second single-stranded polynucleotide to form a first double-stranded stem region; (2) the 5'-  
25 stem sequence of the second single-stranded polynucleotide hybridize with the 3'-stem sequence of the first single-stranded polynucleotide to form a second double-stranded stem region; and, (3) the linker sequences of the first and the second single-stranded polynucleotides form a loop or bulge connecting the first and the second double-stranded stem regions, wherein the 5'-stem sequence and at least a portion of the linker sequence form a guide sequence complementary to a  
30 transcript (such as an mRNA or a non-coding RNA) of a target gene.

Single stranded RNA molecules also include for instance microRNAs (miRNAs).

MicroRNAs are small noncoding RNA molecules that are capable of causing post-

transcriptional silencing of specific genes in cells such as by the inhibition of translation or through degradation of the targeted mRNA. A miRNA can be completely complementary or can have a region of noncomplementarity with a target nucleic acid, consequently resulting in a "bulge" at the region of non-complementarity. The region of noncomplementarity (the bulge) 5 can be flanked by regions of sufficient complementarity, preferably complete complementarity to allow duplex formation. Preferably, the regions of complementarity are at least 8 to 10 nucleotides long (e.g., 8, 9, or 10 nucleotides long). A miRNA can inhibit gene expression by repressing translation, such as when the microRNA is not completely complementary to the target nucleic acid, or by causing target RNA degradation, which is believed to occur only when 10 the miRNA binds its target with perfect complementarity. The invention also can include double-stranded precursors of miRNAs that may or may not form a bulge when bound to their targets.

A miRNA or pre-miRNA can be 16-100 nucleotides in length, and more preferably from 16-80 nucleotides in length. Mature miRNAs can have a length of 16-30 nucleotides, preferably 15 21-25 nucleotides, particularly 21, 22, 23, 24, or 25 nucleotides. MicroRNA precursors can have a length of 70-100 nucleotides and have a hairpin conformation. MicroRNAs can be generated in vivo from pre-miRNAs by enzymes called Dicer and Drosha that specifically process long pre-miRNA into functional miRNA.

Single Stranded DNA molecules include for instance antisense-oligonucleotides. The 20 single-stranded oligonucleotides featured in the invention include antisense nucleic acids. An "antisense" nucleic acid includes a nucleotide sequence that is complementary to a "sense" nucleic acid encoding a gene expression product, e.g., complementary to the coding strand of a double-stranded cDNA molecule or complementary to an RNA sequence, e.g., a pre-mRNA, mRNA, miRNA, or pre-miRNA. Accordingly, an antisense nucleic acid can form hydrogen 25 bonds with a sense nucleic acid target. Given a coding strand sequence (e.g., the sequence of a sense strand of a cDNA molecule), antisense nucleic acids can be designed according to the rules of Watson and Crick base pairing. The antisense nucleic acid molecule can be complementary to a portion of the coding or noncoding region of an RNA, e.g., a pre-mRNA or mRNA. For example, the antisense oligonucleotide can be complementary to the region 30 surrounding the translation start site of a pre-mRNA or mRNA, e.g., the 5' UTR. An antisense oligonucleotide can be, for example, about 10 to 25 nucleotides in length (e.g., 11, 12, 13, 14,

15, 16, 18, 19, 20, 21, 22, 23, or 24 nucleotides in length). An antisense oligonucleotide can also be complementary to a miRNA or pre-miRNA.

An antisense nucleic acid can be constructed using chemical synthesis and/or enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid

5 (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and target nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used. Other appropriate nucleic acid modifications are described herein. Alternatively, the  
10 antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest).

An antisense agent can include ribonucleotides only, deoxyribonucleotides only (e.g., oligodeoxynucleotides), or both deoxyribonucleotides and ribonucleotides. For example, an  
15 antisense agent consisting only of ribonucleotides can hybridize to a complementary RNA, and prevent access of the translation machinery to the target RNA transcript, thereby preventing protein synthesis. An antisense molecule including only deoxyribonucleotides, or deoxyribonucleotides and ribonucleotides, e.g., DNA sequence flanked by RNA sequence at the  
5' and 3' ends of the antisense agent, can hybridize to a complementary RNA, and the RNA  
20 target can be subsequently cleaved by an enzyme, e.g., RNase H. Degradation of the target RNA prevents translation. The flanking RNA sequences can include 2'-O-methylated nucleotides, and phosphorothioate linkages, and the internal DNA sequence can include phosphorothioate internucleotide linkages. The internal DNA sequence is preferably at least five nucleotides in length when targeting by RNaseH activity is desired.

25 Another example of a nucleic acid is a decoy-oligonucleotide, e.g., a decoy RNA. A decoy nucleic acid resembles a natural nucleic acid, but is modified in such a way as to inhibit or interrupt the activity of the natural nucleic acid. For example, a decoy RNA can mimic the natural binding domain for a ligand. The decoy RNA therefore competes with natural binding target for the binding of a specific ligand. The natural binding target can be an endogenous  
30 nucleic acid, e.g., a pre-miRNA, miRNA, premRNA, mRNA or DNA.

Aptamer are also oligonucleotides which may be formulated according to the methods of the invention. An aptamer binds to a non-nucleic acid ligand, such as a small organic molecule

or protein, e.g., a transcription or translation factor, and subsequently modifies (e.g., inhibits) activity. An aptamer can fold into a specific structure that directs the recognition of the targeted binding site on the non-nucleic acid ligand. An aptamer can contain any of the modifications described herein.

5 Antagomirs, which are single stranded, double stranded, partially double stranded and hairpin structured chemically modified oligonucleotides that target a microRNA, are also useful according to the invention. An antagomir may be for instance at least 8 or more contiguous nucleotides substantially complementary to an endogenous miRNA and more particularly agents that include 12 or more contiguous nucleotides substantially complementary to a target sequence 10 of an miRNA or pre-miRNA nucleotide sequence. Preferably, an antagomir featured in the invention includes a nucleotide sequence sufficiently complementary to hybridize to a miRNA target sequence of about 12 to 25 nucleotides, preferably about 15 to 23 nucleotides. An antagomir that is substantially complementary to a nucleotide sequence of an miRNA can be delivered to a cell or a human to inhibit or reduce the activity of an endogenous miRNA, such as 15 when aberrant or undesired miRNA activity, or insufficient activity of a target mRNA that hybridizes to the endogenous miRNA, is linked to a disease or disorder.

For purposes of the present invention, the term asymmetric dsRNA (adsRNA) refers to a duplex, where the length of one strand is substantially higher then the other. As a result, there is an additional single stranded region extending from a duplex. The chemical modification 20 patterns for different regions of the asymmetric dsRNA can be different. For purposes of the present invention, the term "overhang" refers to terminal non-base pairing nucleotide(s) resulting from one strand or region extending beyond the terminus of the complementary strand to which the first strand or region forms a duplex. One or more polynucleotides that are capable of forming a duplex through hydrogen bonding can have overhangs. In a conventional siRNA 25 molecule the overhand length generally doesn't exceed 5 bases in length.

For purposes of the present invention, the term "duplex" refers to a region of double-stranded structure formed by two antiparallel polynucleotide strands as a result of base-pairing between the strands. A duplex may be formed between two separate polynucleotides, or the strands may be contained with a single polynucleotide sequence e.g. a hairpin structure where 30 the "loop" portion of the hairpin allows the two strands to adopt an antiparallel configuration relative to each other.

In certain embodiments, the polynucleotide is unmodified. In other embodiments, at

least one nucleotide is modified. In further embodiments, the modification includes a 2'-H or 2'-modified ribose sugar at the 2nd nucleotide from the 5'-end of the guide sequence. The "2nd nucleotide" is defined as the second nucleotide from the 5'-end of the polynucleotide.

For purposes of the present invention, the term "modification pattern" refers to chemical 5 modification pattern, which is found to be optimal for a particular application. The chemical modification pattern enables generalization of chemical principles for many different sequences. Usually, chemical modification pattern is link to easier functional position or both sequence and functional position. The 2'F modification of every C and U in the guide strand of the duplex, is considered to be an acceptable chemical modification pattern for a guide strand. Another 10 example, is fully Omethyl modified oligo with several phosphorothioates on the ends is an acceptable chemical modification pattern for miRNA inhibitors.

The nucleotides of the invention may be modified at various locations, including the sugar moiety, the phosphodiester linkage, and/or the base.

Sugar moieties include natural, unmodified sugars, *e.g.*, monosaccharide (such as 15 pentose, *e.g.*, ribose, deoxyribose), modified sugars and sugar analogs. In general, possible modifications of nucleomonomers, particularly of a sugar moiety, include, for example, replacement of one or more of the hydroxyl groups with a halogen, a heteroatom, an aliphatic group, or the functionalization of the hydroxyl group as an ether, an amine, a thiol, or the like.

One particularly useful group of modified nucleomonomers are 2'-O-methyl nucleotides. 20 Such 2'-O-methyl nucleotides may be referred to as "methylated," and the corresponding nucleotides may be made from unmethylated nucleotides followed by alkylation or directly from methylated nucleotide reagents. Modified nucleomonomers may be used in combination with unmodified nucleomonomers. For example, an oligonucleotide of the invention may contain both methylated and unmethylated nucleomonomers.

25 Some exemplary modified nucleomonomers include sugar- or backbone-modified ribonucleotides. Modified ribonucleotides may contain a non-naturally occurring base (instead of a naturally occurring base), such as uridines or cytidines modified at the 5'-position, *e.g.*, 5'- (2-amino)propyl uridine and 5'-bromo uridine; adenosines and guanosines modified at the 8-position, *e.g.*, 8-bromo guanosine; deaza nucleotides, *e.g.*, 7-deaza-adenosine; and N-alkylated 30 nucleotides, *e.g.*, N6-methyl adenosine. Also, sugar-modified ribonucleotides may have the 2'-OH group replaced by a H, alkoxy (or OR), R or alkyl, halogen, SH, SR, amino (such as NH<sub>2</sub>, NHR, NR<sub>2</sub>), or CN group, wherein R is lower alkyl, alkenyl, or alkynyl.

Modified ribonucleotides may also have the phosphodiester group connecting to adjacent ribonucleotides replaced by a modified group, *e.g.*, of phosphorothioate group. More generally, the various nucleotide modifications may be combined.

Although the antisense (guide) strand may be substantially identical to at least a portion of the target gene (or genes), at least with respect to the base pairing properties, the sequence need not be perfectly identical to be useful, *e.g.*, to inhibit expression of a target gene's phenotype. Generally, higher homology can be used to compensate for the use of a shorter antisense gene. In some cases, the antisense strand generally will be substantially identical (although in antisense orientation) to the target gene.

The use of 2'-O-methyl modified RNA may also be beneficial in circumstances in which it is desirable to minimize cellular stress responses. RNA having 2'-O-methyl nucleomonomers may not be recognized by cellular machinery that is thought to recognize unmodified RNA. The use of 2'-O-methylated or partially 2'-O-methylated RNA may avoid the interferon response to double-stranded nucleic acids, while maintaining target RNA inhibition. This may be useful, for example, for avoiding the interferon or other cellular stress responses, both in short RNAi (*e.g.*, siRNA) sequences that induce the interferon response, and in longer RNAi sequences that may induce the interferon response.

Overall, modified sugars may include D-ribose, 2'-O-alkyl (including 2'-O-methyl and 2'-O-ethyl), *i.e.*, 2'-alkoxy, 2'-amino, 2'-S-alkyl, 2'-halo (including 2'-fluoro), 2'-methoxyethoxy, 2'-allyloxy (-OCH<sub>2</sub>CH=CH<sub>2</sub>), 2'-propargyl, 2'-propyl, ethynyl, ethenyl, propenyl, and cyano and the like. In one embodiment, the sugar moiety can be a hexose and incorporated into an oligonucleotide as described (Augustyns, K., *et al.*, *Nucl. Acids. Res.* 18:4711 (1992)). Exemplary nucleomonomers can be found, *e.g.*, in U.S. Pat. No. 5,849,902, incorporated by reference herein.

The term "alkyl" includes saturated aliphatic groups, including straight-chain alkyl groups (*e.g.*, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, *etc.*), branched-chain alkyl groups (isopropyl, tert-butyl, isobutyl, *etc.*), cycloalkyl (alicyclic) groups (cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl), alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. In certain embodiments, a straight chain or branched chain alkyl has 6 or fewer carbon atoms in its backbone (*e.g.*, C<sub>1</sub>-C<sub>6</sub> for straight chain, C<sub>3</sub>-C<sub>6</sub> for branched chain), and more preferably 4 or fewer. Likewise, preferred cycloalkyls have

from 3-8 carbon atoms in their ring structure, and more preferably have 5 or 6 carbons in the ring structure. The term C<sub>1</sub>-C<sub>6</sub> includes alkyl groups containing 1 to 6 carbon atoms.

Moreover, unless otherwise specified, the term alkyl includes both “unsubstituted alkyls” and “substituted alkyls,” the latter of which refers to alkyl moieties having independently selected substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonato, phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Cycloalkyls can be further substituted, *e.g.*, with the substituents described above. An “alkylaryl” or an “arylalkyl” moiety is an alkyl substituted with an aryl (*e.g.*, phenylmethyl (benzyl)). The term “alkyl” also includes the side chains of natural and unnatural amino acids. The term “n-alkyl” means a straight chain (*i.e.*, unbranched) unsubstituted alkyl group.

The term “alkenyl” includes unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double bond. For example, the term “alkenyl” includes straight-chain alkenyl groups (*e.g.*, ethylenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl, decenyl, *etc.*), branched-chain alkenyl groups, cycloalkenyl (alicyclic) groups (cyclopropenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl), alkyl or alkenyl substituted cycloalkenyl groups, and cycloalkyl or cycloalkenyl substituted alkenyl groups. In certain embodiments, a straight chain or branched chain alkenyl group has 6 or fewer carbon atoms in its backbone (*e.g.*, C<sub>2</sub>-C<sub>6</sub> for straight chain, C<sub>3</sub>-C<sub>6</sub> for branched chain). Likewise, cycloalkenyl groups may have from 3-8 carbon atoms in their ring structure, and more preferably have 5 or 6 carbons in the ring structure. The term C<sub>2</sub>-C<sub>6</sub> includes alkenyl groups containing 2 to 6 carbon atoms.

Moreover, unless otherwise specified, the term alkenyl includes both “unsubstituted alkenyls” and “substituted alkenyls,” the latter of which refers to alkenyl moieties having independently selected substituents replacing a hydrogen on one or more carbons of the

hydrocarbon backbone. Such substituents can include, for example, alkyl groups, alkynyl groups, halogens, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonato, 5 phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

10 The term “alkynyl” includes unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but which contain at least one triple bond. For example, the term “alkynyl” includes straight-chain alkynyl groups (e.g., ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl, nonynyl, decynyl, *etc.*), branched-chain alkynyl groups, and cycloalkyl or cycloalkenyl substituted alkynyl groups. In certain embodiments, a 15 straight chain or branched chain alkynyl group has 6 or fewer carbon atoms in its backbone (e.g., C<sub>2</sub>-C<sub>6</sub> for straight chain, C<sub>3</sub>-C<sub>6</sub> for branched chain). The term C<sub>2</sub>-C<sub>6</sub> includes alkynyl groups containing 2 to 6 carbon atoms.

Moreover, unless otherwise specified, the term alkynyl includes both “unsubstituted alkynyls” and “substituted alkynyls,” the latter of which refers to alkynyl moieties having 20 independently selected substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, alkyl groups, alkynyl groups, halogens, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonato, 25 phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

30 Unless the number of carbons is otherwise specified, “lower alkyl” as used herein means an alkyl group, as defined above, but having from one to five carbon atoms in its backbone

structure. "Lower alkenyl" and "lower alkynyl" have chain lengths of, for example, 2-5 carbon atoms.

The term "alkoxy" includes substituted and unsubstituted alkyl, alkenyl, and alkynyl groups covalently linked to an oxygen atom. Examples of alkoxy groups include methoxy, ethoxy, isopropoxy, propoxy, butoxy, and pentoxy groups. Examples of substituted alkoxy groups include halogenated alkoxy groups. The alkoxy groups can be substituted with independently selected groups such as alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfmethyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Examples of halogen substituted alkoxy groups include, but are not limited to, fluoromethoxy, difluoromethoxy, trifluoromethoxy, chloromethoxy, dichloromethoxy, trichloromethoxy, etc.

The term "heteroatom" includes atoms of any element other than carbon or hydrogen. Preferred heteroatoms are nitrogen, oxygen, sulfur and phosphorus.

The term "hydroxy" or "hydroxyl" includes groups with an -OH or -O<sup>-</sup> (with an appropriate counterion).

The term "halogen" includes fluorine, bromine, chlorine, iodine, etc. The term "perhalogenated" generally refers to a moiety wherein all hydrogens are replaced by halogen atoms.

The term "substituted" includes independently selected substituents which can be placed on the moiety and which allow the molecule to perform its intended function. Examples of substituents include alkyl, alkenyl, alkynyl, aryl, (CR'R")<sub>0-3</sub>NR'R", (CR'R")<sub>0-3</sub>CN, NO<sub>2</sub>, halogen, (CR'R")<sub>0-3</sub>C(halogen)<sub>3</sub>, (CR'R")<sub>0-3</sub>CH(halogen)<sub>2</sub>, (CR'R")<sub>0-3</sub>CH<sub>2</sub>(halogen), (CR'R")<sub>0-3</sub>CONR'R", (CR'R")<sub>0-3</sub>S(O)<sub>1-2</sub>NR'R", (CR'R")<sub>0-3</sub>CHO, (CR'R")<sub>0-3</sub>O(CR'R")<sub>0-3</sub>H, (CR'R")<sub>0-3</sub>S(O)<sub>0-2</sub>R', (CR'R")<sub>0-3</sub>O(CR'R")<sub>0-3</sub>H, (CR'R")<sub>0-3</sub>COR', (CR'R")<sub>0-3</sub>CO<sub>2</sub>R', or (CR'R")<sub>0-3</sub>OR' groups; wherein each R' and R" are each independently hydrogen, a C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>2</sub>-C<sub>5</sub> alkynyl, or aryl group, or R' and R" taken together are a benzylidene group or a -(CH<sub>2</sub>)<sub>2</sub>O(CH<sub>2</sub>)<sub>2</sub>- group.

The term "amine" or "amino" includes compounds or moieties in which a nitrogen atom is covalently bonded to at least one carbon or heteroatom. The term "alkyl amino" includes groups and compounds wherein the nitrogen is bound to at least one additional alkyl group. The term "dialkyl amino" includes groups wherein the nitrogen atom is bound to at least two additional alkyl groups.

The term "ether" includes compounds or moieties which contain an oxygen bonded to two different carbon atoms or heteroatoms. For example, the term includes "alkoxyalkyl," which refers to an alkyl, alkenyl, or alkynyl group covalently bonded to an oxygen atom which is covalently bonded to another alkyl group.

The term "base" includes the known purine and pyrimidine heterocyclic bases, deazapurines, and analogs (including heterocyclic substituted analogs, *e.g.*, aminoethoxy phenoxazine), derivatives (*e.g.*, 1-alkyl-, 1-alkenyl-, heteroaromatic- and 1-alkynyl derivatives) and tautomers thereof. Examples of purines include adenine, guanine, inosine, diaminopurine, and xanthine and analogs (*e.g.*, 8-oxo-N<sup>6</sup>-methyladenine or 7-diazaxanthine) and derivatives thereof. Pyrimidines include, for example, thymine, uracil, and cytosine, and their analogs (*e.g.*, 5-methylcytosine, 5-methyluracil, 5-(1-propynyl)uracil, 5-(1-propynyl)cytosine and 4,4-ethanocytosine). Other examples of suitable bases include non-purinyl and non-pyrimidinyl bases such as 2-aminopyridine and triazines.

In a preferred embodiment, the nucleomonomers of an oligonucleotide of the invention are RNA nucleotides. In another preferred embodiment, the nucleomonomers of an oligonucleotide of the invention are modified RNA nucleotides. Thus, the oligonucleotides contain modified RNA nucleotides.

The term "nucleoside" includes bases which are covalently attached to a sugar moiety, preferably ribose or deoxyribose. Examples of preferred nucleosides include ribonucleosides and deoxyribonucleosides. Nucleosides also include bases linked to amino acids or amino acid analogs which may comprise free carboxyl groups, free amino groups, or protecting groups. Suitable protecting groups are well known in the art (see P. G. M. Wuts and T. W. Greene, "Protective Groups in Organic Synthesis", 2<sup>nd</sup> Ed., Wiley-Interscience, New York, 1999).

The term "nucleotide" includes nucleosides which further comprise a phosphate group or a phosphate analog.

As used herein, the term "linkage" includes a naturally occurring, unmodified phosphodiester moiety (-O-(PO<sup>2-</sup>)-O-) that covalently couples adjacent nucleomonomers. As

used herein, the term “substitute linkage” includes any analog or derivative of the native phosphodiester group that covalently couples adjacent nucleomonomers. Substitute linkages include phosphodiester analogs, *e.g.*, phosphorothioate, phosphorodithioate, and P-ethyoxyphosphodiester, P-ethoxyphosphodiester, P-alkyloxyphosphotriester,

5 methylphosphonate, and nonphosphorus containing linkages, *e.g.*, acetals and amides. Such substitute linkages are known in the art (*e.g.*, Bjergarde *et al.* 1991. Nucleic Acids Res. 19:5843; Caruthers *et al.* 1991. Nucleosides Nucleotides. 10:47). In certain embodiments, non-hydrolyzable linkages are preferred, such as phosphorothioate linkages.

In certain embodiments, oligonucleotides of the invention comprise 3' and 5' termini 10 (except for circular oligonucleotides). In one embodiment, the 3' and 5' termini of an oligonucleotide can be substantially protected from nucleases *e.g.*, by modifying the 3' or 5' linkages (*e.g.*, U.S. Pat. No. 5,849,902 and WO 98/13526). For example, oligonucleotides can be made resistant by the inclusion of a “blocking group.” The term “blocking group” as used herein refers to substituents (*e.g.*, other than OH groups) that can be attached to oligonucleotides 15 or nucleomonomers, either as protecting groups or coupling groups for synthesis (*e.g.*, FITC, propyl (CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>3</sub>), glycol (-O-CH<sub>2</sub>-CH<sub>2</sub>-O-) phosphate (PO<sub>3</sub><sup>2-</sup>), hydrogen phosphonate, or phosphoramidite). “Blocking groups” also include “end blocking groups” or “exonuclease blocking groups” which protect the 5' and 3' termini of the oligonucleotide, including modified nucleotides and non-nucleotide exonuclease resistant structures.

20 Exemplary end-blocking groups include cap structures (*e.g.*, a 7-methylguanosine cap), inverted nucleomonomers, *e.g.*, with 3'-3' or 5'-5' end inversions (see, *e.g.*, Ortiagao *et al.* 1992. *Antisense Res. Dev.* 2:129), methylphosphonate, phosphoramidite, non-nucleotide groups (*e.g.*, non-nucleotide linkers, amino linkers, conjugates) and the like. The 3' terminal nucleomonomer 25 can comprise a modified sugar moiety. The 3' terminal nucleomonomer comprises a 3'-O that can optionally be substituted by a blocking group that prevents 3'-exonuclease degradation of the oligonucleotide. For example, the 3'-hydroxyl can be esterified to a nucleotide through a 3'→3' internucleotide linkage. For example, the alkyloxy radical can be methoxy, ethoxy, or isopropoxy, and preferably, ethoxy. Optionally, the 3'→3' linked nucleotide at the 3' terminus 30 can be linked by a substitute linkage. To reduce nuclease degradation, the 5' most 3'→5' linkage can be a modified linkage, *e.g.*, a phosphorothioate or a P-alkyloxyphosphotriester linkage. Preferably, the two 5' most 3'→5' linkages are modified linkages. Optionally, the 5' terminal

hydroxy moiety can be esterified with a phosphorus containing moiety, *e.g.*, phosphate, phosphorothioate, or P-ethoxyphosphate.

Another type of conjugates that can be attached to the end (3' or 5' end), the loop region, or any other parts of the miRNA might include a sterol, sterol type molecule, peptide, small molecule, protein, etc. In some embodiments, a miRNA may contain more than one conjugates (same or different chemical nature). In some embodiments, the conjugate is cholesterol.

Another way to increase target gene specificity, or to reduce off-target silencing effect, is to introduce a 2'-modification (such as the 2'-O methyl modification) at a position corresponding to the second 5'-end nucleotide of the guide sequence. This allows the positioning of this 2'-modification in the Dicer-resistant hairpin structure, thus enabling one to design better RNAi constructs with less or no off-target silencing.

In one embodiment, a hairpin polynucleotide of the invention can comprise one nucleic acid portion which is DNA and one nucleic acid portion which is RNA. Antisense (guide) sequences of the invention can be "chimeric oligonucleotides" which comprise an RNA-like and a DNA-like region.

The language "RNase H activating region" includes a region of an oligonucleotide, *e.g.*, a chimeric oligonucleotide, that is capable of recruiting RNase H to cleave the target RNA strand to which the oligonucleotide binds. Typically, the RNase activating region contains a minimal core (of at least about 3-5, typically between about 3-12, more typically, between about 5-12, and more preferably between about 5-10 contiguous nucleomonomers) of DNA or DNA-like nucleomonomers. (See, *e.g.*, U.S. Pat. No. 5,849,902). Preferably, the RNase H activating region comprises about nine contiguous deoxyribose containing nucleomonomers.

The language "non-activating region" includes a region of an antisense sequence, *e.g.*, a chimeric oligonucleotide, that does not recruit or activate RNase H. Preferably, a non-activating region does not comprise phosphorothioate DNA. The oligonucleotides of the invention comprise at least one non-activating region. In one embodiment, the non-activating region can be stabilized against nucleases or can provide specificity for the target by being complementary to the target and forming hydrogen bonds with the target nucleic acid molecule, which is to be bound by the oligonucleotide.

In one embodiment, at least a portion of the contiguous polynucleotides are linked by a substitute linkage, *e.g.*, a phosphorothioate linkage.

In certain embodiments, most or all of the nucleotides beyond the guide sequence (2'-modified or not) are linked by phosphorothioate linkages. Such constructs tend to have improved pharmacokinetics due to their higher affinity for serum proteins. The phosphorothioate linkages in the non-guide sequence portion of the polynucleotide generally do not interfere with guide strand activity, once the latter is loaded into RISC.

5 Antisense (guide) sequences of the present invention may include "morpholino oligonucleotides." Morpholino oligonucleotides are non-ionic and function by an RNase H-independent mechanism. Each of the 4 genetic bases (Adenine, Cytosine, Guanine, and Thymine/Uracil) of the morpholino oligonucleotides is linked to a 6-membered morpholine ring.

10 Morpholino oligonucleotides are made by joining the 4 different subunit types by, e.g., non-ionic phosphorodiamidate inter-subunit linkages. Morpholino oligonucleotides have many advantages including: complete resistance to nucleases (Antisense & Nucl. Acid Drug Dev. 1996. 6:267); predictable targeting (Biochimica Biophysica Acta. 1999. 1489:141); reliable activity in cells (Antisense & Nucl. Acid Drug Dev. 1997. 7:63); excellent sequence specificity

15 (Antisense & Nucl. Acid Drug Dev. 1997. 7:151); minimal non-antisense activity (Biochimica Biophysica Acta. 1999. 1489:141); and simple osmotic or scrape delivery (Antisense & Nucl. Acid Drug Dev. 1997. 7:291). Morpholino oligonucleotides are also preferred because of their non-toxicity at high doses. A discussion of the preparation of morpholino oligonucleotides can be found in Antisense & Nucl. Acid Drug Dev. 1997. 7:187.

20 Some of the preferred chemical modifications described herein are believed to promote single stranded polynucleotide loading into the RISC. Single stranded polynucleotides have been shown to be active in loading into RISC and inducing gene silencing. However, the level of activity for single stranded polynucleotides appears to be 2 to 4 orders of magnitude lower when compared to a duplex polynucleotide.

25 The present invention provides a description of the chemical modification patterns, which may (a) significantly increase stability of the single stranded polynucleotide (b) promote efficient loading of the polynucleotide into the RISC complex and (c) improve uptake of the single stranded nucleotide by the cell. Figure 5 provides some non-limiting examples of the chemical modification patterns which may be beneficial for achieving single stranded

30 polynucleotide efficacy inside the cell. The chemical modification patterns may include combination of ribose, backbone, hydrophobic nucleoside and conjugate type of modifications.

In addition, in some of the embodiments, the 5' end of the single polynucleotide may be chemically phosphorylated.

In yet another embodiment, the present invention provides a description of the chemical modifications patterns, which improve functionality of RISC inhibiting polynucleotides. Single stranded polynucleotides have been shown to inhibit activity of a preloaded RISC complex through the substrate competition mechanism. For these types of molecules, conventionally called antagonists, the activity usually requires high concentration and in vivo delivery is not very effective. The present invention provides a description of the chemical modification patterns, which may (a) significantly increase stability of the single stranded polynucleotide (b) 10 promote efficient recognition of the polynucleotide by the RISC as a substrate and/or (c) improve uptake of the single stranded nucleotide by the cell. Figure 6 provides some non-limiting examples of the chemical modification patterns that may be beneficial for achieving single stranded polynucleotide efficacy inside the cell. The chemical modification patterns may include combination of ribose, backbone, hydrophobic nucleoside and conjugate type of 15 modifications.

The modifications provided by the present invention are applicable to all polynucleotides. This includes single stranded RISC entering polynucleotides, single stranded RISC inhibiting polynucleotides, conventional duplexed polynucleotides of variable length (15-40 bp), asymmetric duplexed polynucleotides, and the like. Polynucleotides may be modified 20 with wide variety of chemical modification patterns, including 5' end, ribose, backbone and hydrophobic nucleoside modifications.

In certain embodiments, the modified RNA polynucleotide of the invention with the above-referenced 5'-end modification exhibits significantly (e.g., at least about 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90% or more) less "off-target" gene 25 silencing when compared to similar constructs without the specified 5'-end modification, thus greatly improving the overall specificity of the RNAi reagent or therapeutics.

For purposes of the present invention, modified bases refer to nucleotide bases such as, for example, adenine, guanine, cytosine, thymine, and uracil, xanthine, inosine, and queuosine that have been modified by the replacement or addition of one or more atoms or groups. Some 30 Examples of types of modifications that can comprise nucleotides that are modified with respect to the base moieties, include but are not limited to, alkylated, halogenated, thiolated, aminated, amidated, or acetylated bases, in various combinations. More specific modified bases include,

for example, 5-propynyluridine, 5-propynylcytidine, 6-methyladenine, 6-methylguanine, N,N,-dimethyladenine, 2-propyladenine, 2-propylguanine, 2-aminoadenine, 1-methylinosine, 3-methyluridine, 5-methylcytidine, 5-methyluridine and other nucleotides having a modification at the 5 position, 5-(2-amino)propyl uridine, 5-halocytidine, 5-halouridine, 4-acetylcytidine, 1-methyladenosine, 2-methyladenosine, 3-methylcytidine, 6-methyluridine, 2-methylguanosine, 7-methylguanosine, 2,2-dimethylguanosine, 5-methylaminoethyluridine, 5-methoxyuridine, deazanucleotides such as 7-deaza-adenosine, 6-azouridine, 6-azocytidine, 6-azothymidine, 5-methyl-2-thiouridine, other thio bases such as 2-thiouridine and 4-thiouridine and 2-thiocytidine, dihydrouridine, pseudouridine, queuosine, archaeosine, naphthyl and substituted naphthyl groups, any O- and N-alkylated purines and pyrimidines such as N6-methyladenosine, 5-methylcarbonylmethyluridine, uridine 5-oxyacetic acid, pyridine-4-one, pyridine-2-one, phenyl and modified phenyl groups such as aminophenol or 2,4,6-trimethoxy benzene, modified cytosines that act as G-clamp nucleotides, 8-substituted adenines and guanines, 5-substituted uracils and thymines, azapyrimidines, carboxyhydroxyalkyl nucleotides, 15 carboxyalkylaminoalkyl nucleotides, and alkylcarbonylalkylated nucleotides. Modified nucleotides also include those nucleotides that are modified with respect to the sugar moiety, as well as nucleotides having sugars or analogs thereof that are not ribosyl. For example, the sugar moieties may be, or be based on, mannoses, arabinoses, glucopyranoses, galactopyranoses, 4-thioribose, and other sugars, heterocycles, or carbocycles. The term nucleotide is also meant to 20 include what are known in the art as universal bases. By way of example, universal bases include but are not limited to 3-nitropyrrole, 5-nitroindole, or nebularine.

In another embodiment, chemical modifications of the first or second polynucleotide include, but not limited to, 5' position modification of Uridine and Cytosine (4-pyridyl, 2-pyridyl, indolyl, phenyl (C<sub>6</sub>H<sub>5</sub>OH); tryptophanyl (C<sub>8</sub>H<sub>6</sub>N)CH<sub>2</sub>CH(NH<sub>2</sub>)CO), isobutyl, butyl, 25 aminobenzyl; phenyl; naphthyl, etc), where the chemical modification might alter base pairing capabilities of a nucleotide. For the guide strand an important feature of this aspect of the invention is the position of the chemical modification relative to the 5' end of the antisense and sequence. For example, chemical phosphorylation of the 5' end of the guide strand is usually beneficial for efficacy. O-methyl modifications in the seed region of the sense strand (position 30 2-7 relative to the 5' end) are not generally well tolerated, whereas 2'F and deoxy are well tolerated. The mid part of the guide strand and the 3' end of the guide strand are more

permissive in a type of chemical modifications applied. Deoxy modifications are not tolerated at the 3' end of the guide strand.

A unique feature of this aspect of the invention involves the use of hydrophobic modification on the bases. In one embodiment, the hydrophobic modifications are preferably positioned near the 5' end of the guide strand, in other embodiments, they localized in the middle of the guides strand, in other embodiment they localized at the 3' end of the guide strand and yet in another embodiment they are distributed thought the whole length of the polynucleotide. The same type of patterns is applicable to the passenger strand of the duplex.

The other part of the molecule is a single stranded region. The single stranded region is expected to range from 7 to 40 nucleotides.

In one embodiment, the single stranded region of the first polynucleotide contains modifications selected from the group consisting of between 40% and 90% hydrophobic base modifications, between 40%-90% phosphorothioates, between 40% -90% modification of the ribose moiety, and any combination of the preceding.

Efficiency of guide strand (first polynucleotide) loading into the RISC complex might be altered for heavily modified polynucleotides, so in one embodiment, the duplex polynucleotide includes a mismatch between nucleotide 9, 11, 12, 13, or 14 on the guide strand (first polynucleotide) and the opposite nucleotide on the sense strand (second polynucleotide) to promote efficient guide strand loading.

## 20 *Synthesis*

Oligonucleotides of the invention can be synthesized by any method known in the art, *e.g.*, using enzymatic synthesis and/or chemical synthesis. The oligonucleotides can be synthesized *in vitro* (*e.g.*, using enzymatic synthesis and chemical synthesis) or *in vivo* (using recombinant DNA technology well known in the art).

25 In a preferred embodiment, chemical synthesis is used for modified polynucleotides. Chemical synthesis of linear oligonucleotides is well known in the art and can be achieved by solution or solid phase techniques. Preferably, synthesis is by solid phase methods. Oligonucleotides can be made by any of several different synthetic procedures including the phosphoramidite, phosphite triester, H-phosphonate, and phosphotriester methods, typically by 30 automated synthesis methods.

Oligonucleotide synthesis protocols are well known in the art and can be found, *e.g.*, in

U.S. Pat. No. 5,830,653; WO 98/13526; Stec *et al.* 1984. *J. Am. Chem. Soc.* 106:6077; Stec *et al.* 1985. *J. Org. Chem.* 50:3908; Stec *et al.* *J. Chromatog.* 1985. 326:263; LaPlanche *et al.* 1986. *Nucl. Acid. Res.* 1986. 14:9081; Fasman G. D., 1989. *Practical Handbook of Biochemistry and Molecular Biology.* 1989. CRC Press, Boca Raton, Fla.; Lamone. 1993. *Biochem. Soc. Trans.* 21:1; U.S. Pat. No. 5,013,830; U.S. Pat. No. 5,214,135; U.S. Pat. No. 5,525,719; Kawasaki *et al.* 1993. *J. Med. Chem.* 36:831; WO 92/03568; U.S. Pat. No. 5,276,019; and U.S. Pat. No. 5,264,423.

The synthesis method selected can depend on the length of the desired oligonucleotide and such choice is within the skill of the ordinary artisan. For example, the phosphoramidite and phosphite triester method can produce oligonucleotides having 175 or more nucleotides, while the H-phosphonate method works well for oligonucleotides of less than 100 nucleotides. If modified bases are incorporated into the oligonucleotide, and particularly if modified phosphodiester linkages are used, then the synthetic procedures are altered as needed according to known procedures. In this regard, Uhlmann *et al.* (1990, *Chemical Reviews* 90:543-584) provide references and outline procedures for making oligonucleotides with modified bases and modified phosphodiester linkages. Other exemplary methods for making oligonucleotides are taught in Sonveaux. 1994. "Protecting Groups in Oligonucleotide Synthesis"; Agrawal. *Methods in Molecular Biology* 26:1. Exemplary synthesis methods are also taught in "Oligonucleotide Synthesis - A Practical Approach" (Gait, M. J. IRL Press at Oxford University Press. 1984). Moreover, linear oligonucleotides of defined sequence, including some sequences with modified nucleotides, are readily available from several commercial sources.

The oligonucleotides may be purified by polyacrylamide gel electrophoresis, or by any of a number of chromatographic methods, including gel chromatography and high pressure liquid chromatography. To confirm a nucleotide sequence, especially unmodified nucleotide sequences, oligonucleotides may be subjected to DNA sequencing by any of the known procedures, including Maxam and Gilbert sequencing, Sanger sequencing, capillary electrophoresis sequencing, the wandering spot sequencing procedure or by using selective chemical degradation of oligonucleotides bound to Hybond paper. Sequences of short oligonucleotides can also be analyzed by laser desorption mass spectroscopy or by fast atom bombardment (McNeal, *et al.*, 1982, *J. Am. Chem. Soc.* 104:976; Viari, *et al.*, 1987, *Biomed. Environ. Mass Spectrom.* 14:83; Grotjahn *et al.*, 1982, *Nuc. Acid Res.* 10:4671). Sequencing methods are also available for RNA oligonucleotides.

The quality of oligonucleotides synthesized can be verified by testing the oligonucleotide by capillary electrophoresis and denaturing strong anion HPLC (SAX-HPLC) using, *e.g.*, the method of Bergot and Egan. 1992. *J. Chrom.* 599:35.

Other exemplary synthesis techniques are well known in the art (see, *e.g.*, Sambrook *et al.*, Molecular Cloning: a Laboratory Manual, Second Edition (1989); DNA Cloning, Volumes I and II (DN Glover Ed. 1985); Oligonucleotide Synthesis (M J Gait Ed, 1984; Nucleic Acid Hybridisation (B D Hames and S J Higgins eds. 1984); A Practical Guide to Molecular Cloning (1984); or the series, Methods in Enzymology (Academic Press, Inc.)).

In certain embodiments, the subject RNAi constructs or at least portions thereof are transcribed from expression vectors encoding the subject constructs. Any art recognized vectors may be used for this purpose. The transcribed RNAi constructs may be isolated and purified, before desired modifications (such as replacing an unmodified sense strand with a modified one, *etc.*) are carried out.

*Delivery/Carrier*

15 Oligonucleotides and oligonucleotide compositions are contacted with (*i.e.*, brought into contact with, also referred to herein as administered or delivered to) and taken up by one or more cells or a cell lysate. The term “cells” includes prokaryotic and eukaryotic cells, preferably vertebrate cells, and, more preferably, mammalian cells. In a preferred embodiment, the oligonucleotide compositions of the invention are contacted with human cells.

20 Oligonucleotide compositions of the invention can be contacted with cells *in vitro*, *e.g.*, in a test tube or culture dish, (and may or may not be introduced into a subject) or *in vivo*, *e.g.*, in a subject such as a mammalian subject. Oligonucleotides are taken up by cells at a slow rate by endocytosis, but endocytosed oligonucleotides are generally sequestered and not available, *e.g.*, for hybridization to a target nucleic acid molecule. In one embodiment, cellular uptake can 25 be facilitated by electroporation or calcium phosphate precipitation. However, these procedures are only useful for *in vitro* or *ex vivo* embodiments, are not convenient and, in some cases, are associated with cell toxicity.

30 In another embodiment, delivery of oligonucleotides into cells can be enhanced by suitable art recognized methods including calcium phosphate, DMSO, glycerol or dextran, electroporation, or by transfection, *e.g.*, using cationic, anionic, or neutral lipid compositions or liposomes using methods known in the art (see *e.g.*, WO 90/14074; WO 91/16024; WO

91/17424; U.S. Pat. No. 4,897,355; Bergan *et al.* 1993. *Nucleic Acids Research*. 21:3567).

Enhanced delivery of oligonucleotides can also be mediated by the use of vectors (See *e.g.*, Shi, Y. 2003. *Trends Genet* 2003 Jan. 19:9; Reichhart J M *et al.* *Genesis*. 2002. 34(1-2):1604, Yu *et al.* 2002. *Proc. Natl. Acad. Sci. USA* 99:6047; Sui *et al.* 2002. *Proc. Natl. Acad. Sci. USA*

5 99:5515) viruses, polyamine or polycation conjugates using compounds such as polylysine, protamine, or Ni, N12-bis (ethyl) spermine (see, *e.g.*, Bartzatt, R. *et al.* 1989. *Biotechnol. Appl. Biochem.* 11:133; Wagner E. *et al.* 1992. *Proc. Natl. Acad. Sci.* 88:4255).

In certain embodiments, the miRNA of the invention may be delivered by using various beta-glucan containing particles, such as those described in US 2005/0281781 A1,

10 WO 2006/007372, and WO 2007/050643 (all incorporated herein by reference). In certain embodiments, the beta-glucan particle is derived from yeast. In certain embodiments, the payload trapping molecule is a polymer, such as those with a molecular weight of at least about 1000 Da, 10,000 Da, 50,000 Da, 100 kDa, 500 kDa, etc. Preferred polymers include (without limitation) cationic polymers, chitosans, or PEI (polyethylenimine), etc.

15 Such beta-glucan based delivery system may be formulated for oral delivery, where the orally delivered beta-glucan / miRNA constructs may be engulfed by macrophages or other related phagocytic cells, which may in turn release the miRNA constructs in selected *in vivo* sites. Alternatively or in addition, the miRNA may change the expression of certain macrophage target genes.

20 The optimal protocol for uptake of oligonucleotides will depend upon a number of factors, the most crucial being the type of cells that are being used. Other factors that are important in uptake include, but are not limited to, the nature and concentration of the oligonucleotide, the confluence of the cells, the type of culture the cells are in (*e.g.*, a suspension culture or plated) and the type of media in which the cells are grown.

25 The delivery of oligonucleotides can also be improved by targeting the oligonucleotides to a cellular receptor. The targeting moieties can be conjugated to the oligonucleotides or attached to a carrier group (*i.e.*, poly(L-lysine) or liposomes) linked to the oligonucleotides. This method is well suited to cells that display specific receptor-mediated endocytosis.

For instance, oligonucleotide conjugates to 6-phosphomannosylated proteins are 30 internalized 20-fold more efficiently by cells expressing mannose 6-phosphate specific receptors than free oligonucleotides. The oligonucleotides may also be coupled to a ligand for a cellular

receptor using a biodegradable linker. In another example, the delivery construct is mannosylated streptavidin which forms a tight complex with biotinylated oligonucleotides. Mannosylated streptavidin was found to increase 20-fold the internalization of biotinylated oligonucleotides. (Vlassov *et al.* 1994. *Biochimica et Biophysica Acta* 1197:95-108).

5 *Administration*

The optimal course of administration or delivery of the oligonucleotides may vary depending upon the desired result and/or on the subject to be treated. As used herein “administration” refers to contacting cells with oligonucleotides and can be performed *in vitro* or *in vivo*. The dosage of oligonucleotides may be adjusted to optimally reduce expression of a 10 protein translated from a target nucleic acid molecule, *e.g.*, as measured by a readout of RNA stability or by a therapeutic response, without undue experimentation.

For example, expression of the protein encoded by the nucleic acid target can be measured to determine whether or not the dosage regimen needs to be adjusted accordingly. In addition, an increase or decrease in RNA or protein levels in a cell or produced by a cell can be 15 measured using any art recognized technique. By determining whether transcription has been decreased, the effectiveness of the oligonucleotide in inducing the cleavage of a target RNA can be determined.

Any of the above-described oligonucleotide compositions can be used alone or in conjunction with a pharmaceutically acceptable carrier. As used herein, “pharmaceutically 20 acceptable carrier” includes appropriate solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, it can be used in the therapeutic compositions. Supplementary active ingredients can also be incorporated into the 25 compositions.

Moreover, the present invention provides for administering the subject oligonucleotides with an osmotic pump providing continuous infusion of such oligonucleotides, for example, as described in Rataiczak *et al.* (1992 *Proc. Natl. Acad. Sci. USA* 89:11823-11827). Such osmotic pumps are commercially available, *e.g.*, from Alzet Inc. (Palo Alto, Calif.). Topical 30 administration and parenteral administration in a cationic lipid carrier are preferred.

With respect to *in vivo* applications, the formulations of the present invention can be

administered to a subject in a variety of forms adapted to the chosen route of administration, e.g., parenterally, orally, or intraperitoneally. Parenteral administration, which is preferred, includes administration by the following routes: intravenous; intramuscular; interstitially; intraarterially; subcutaneous; intra ocular; intrasynovial; trans epithelial, including transdermal; 5 pulmonary via inhalation; ophthalmic; sublingual and buccal; topically, including ophthalmic; dermal; ocular; rectal; and nasal inhalation via insufflation.

Pharmaceutical preparations for parenteral administration include aqueous solutions of the active compounds in water-soluble or water-dispersible form. In addition, suspensions of the active compounds as appropriate oily injection suspensions may be administered. Suitable 10 lipophilic solvents or vehicles include fatty oils, for example, sesame oil, or synthetic fatty acid esters, for example, ethyl oleate or triglycerides. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension include, for example, sodium carboxymethyl cellulose, sorbitol, or dextran, optionally, the suspension may also contain stabilizers. The oligonucleotides of the invention can be formulated in liquid solutions, 15 preferably in physiologically compatible buffers such as Hank's solution or Ringer's solution. In addition, the oligonucleotides may be formulated in solid form and redissolved or suspended immediately prior to use. Lyophilized forms are also included in the invention.

Pharmaceutical preparations for topical administration include transdermal patches, ointments, lotions, creams, gels, drops, sprays, suppositories, liquids and powders. In addition, 20 conventional pharmaceutical carriers, aqueous, powder or oily bases, or thickeners may be used in pharmaceutical preparations for topical administration.

Pharmaceutical preparations for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets or tablets. In addition, thickeners, flavoring agents, diluents, emulsifiers, dispersing aids, or binders may be 25 used in pharmaceutical preparations for oral administration.

For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are known in the art, and include, for example, for transmucosal administration bile salts and fusidic acid derivatives, and detergents. Transmucosal administration may be through nasal sprays or using suppositories. For oral 30 administration, the oligonucleotides are formulated into conventional oral administration forms such as capsules, tablets, and tonics. For topical administration, the oligonucleotides of the

invention are formulated into ointments, salves, gels, or creams as known in the art.

Drug delivery vehicles can be chosen *e.g.*, for *in vitro*, for systemic, or for topical administration. These vehicles can be designed to serve as a slow release reservoir or to deliver their contents directly to the target cell. An advantage of using some direct delivery drug

5 vehicles is that multiple molecules are delivered per uptake. Such vehicles have been shown to increase the circulation half-life of drugs that would otherwise be rapidly cleared from the blood stream. Some examples of such specialized drug delivery vehicles which fall into this category are liposomes, hydrogels, cyclodextrins, biodegradable nanocapsules, and bioadhesive microspheres.

10 The described oligonucleotides may be administered systemically to a subject. Systemic absorption refers to the entry of drugs into the blood stream followed by distribution throughout the entire body. Administration routes which lead to systemic absorption include: intravenous, subcutaneous, intraperitoneal, and intranasal. Each of these administration routes delivers the oligonucleotide to accessible diseased cells. Following subcutaneous administration, the  
15 therapeutic agent drains into local lymph nodes and proceeds through the lymphatic network into the circulation. The rate of entry into the circulation has been shown to be a function of molecular weight or size. The use of a liposome or other drug carrier localizes the oligonucleotide at the lymph node. The oligonucleotide can be modified to diffuse into the cell, or the liposome can directly participate in the delivery of either the unmodified or modified  
20 oligonucleotide into the cell.

The emulsions of the present invention may contain excipients such as emulsifiers, stabilizers, dyes, fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives, and anti-oxidants may also be present in emulsions as needed. These excipients may be present as a solution in either the aqueous phase, oily phase or  
25 itself as a separate phase.

Examples of naturally occurring emulsifiers that may be used in emulsion formulations of the present invention include lanolin, beeswax, phosphatides, lecithin and acacia. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. Examples of finely divided solids that may be used as  
30 emulsifiers include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgite, hectorite, kaolin, montrnorillonite, colloidal aluminum silicate and

colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

Examples of preservatives that may be included in the emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of 5 p-hydroxybenzoic acid, and boric acid. Examples of antioxidants that may be included in the emulsion formulations include free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

10 In one embodiment, the compositions of oligonucleotides are formulated as microemulsions. A microemulsion is a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution. Typically microemulsions are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a 4th component, generally an intermediate chain-length alcohol to form a transparent 15 system.

Surfactants that may be used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol 20 monocaprate (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (S0750), decaglycerol decaoleate (DA0750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules.

25 Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, 30 fatty acid esters, medium chain (C<sub>8</sub>-C<sub>12</sub>) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C<sub>8</sub>-C<sub>10</sub>

glycerides, vegetable oils and silicone oil.

Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both oil/water and water/oil) have been proposed to enhance the oral bioavailability of drugs.

5 Microemulsions offer improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides *et al.*, *Pharmaceutical Research*, 1994, 11:1385; Ho *et al.*, *J. Pharm. Sci.*, 1996, 85:138-143).

10 Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of oligonucleotides from the gastrointestinal tract, as well as improve the local cellular uptake of oligonucleotides within the gastrointestinal tract, vagina, buccal cavity and

15 other areas of administration.

In an embodiment, the present invention employs various penetration enhancers to affect the efficient delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to increasing the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also act to enhance the permeability of lipophilic drugs.

20 Five categories of penetration enhancers that may be used in the present invention include: surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants. Other agents may be utilized to enhance the penetration of the administered oligonucleotides include: glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-15 pyrrol, 25 azones, and terpenes such as limonene, and menthone.

The oligonucleotides, especially in lipid formulations, can also be administered by coating a medical device, for example, a catheter, such as an angioplasty balloon catheter, with a cationic lipid formulation. Coating may be achieved, for example, by dipping the medical device into a lipid formulation or a mixture of a lipid formulation and a suitable solvent, for 30 example, an aqueous-based buffer, an aqueous solvent, ethanol, methylene chloride, chloroform and the like. An amount of the formulation will naturally adhere to the surface of the device

which is subsequently administered to a subject, as appropriate. Alternatively, a lyophilized mixture of a lipid formulation may be specifically bound to the surface of the device. Such binding techniques are described, for example, in K. Ishihara *et al.*, *Journal of Biomedical Materials Research*, Vol. 27, pp. 1309-1314 (1993), the disclosures of which are incorporated 5 herein by reference in their entirety.

The useful dosage to be administered and the particular mode of administration will vary depending upon such factors as the cell type, or for *in vivo* use, the age, weight and the particular animal and region thereof to be treated, the particular oligonucleotide and delivery method used, the therapeutic or diagnostic use contemplated, and the form of the formulation, for example, 10 suspension, emulsion, micelle or liposome, as will be readily apparent to those skilled in the art. Typically, dosage is administered at lower levels and increased until the desired effect is achieved. When lipids are used to deliver the oligonucleotides, the amount of lipid compound that is administered can vary and generally depends upon the amount of oligonucleotide agent being administered. For example, the weight ratio of lipid compound to oligonucleotide agent is 15 preferably from about 1:1 to about 15:1, with a weight ratio of about 5:1 to about 10:1 being more preferred. Generally, the amount of cationic lipid compound which is administered will vary from between about 0.1 milligram (mg) to about 1 gram (g). By way of general guidance, typically between about 0.1 mg and about 10 mg of the particular oligonucleotide agent, and about 1 mg to about 100 mg of the lipid compositions, each per kilogram of subject body 20 weight, is administered, although higher and lower amounts can be used.

The agents of the invention are administered to subjects or contacted with cells in a biologically compatible form suitable for pharmaceutical administration. By “biologically compatible form suitable for administration” is meant that the oligonucleotide is administered in a form in which any toxic effects are outweighed by the therapeutic effects of the 25 oligonucleotide. In one embodiment, oligonucleotides can be administered to subjects. Examples of subjects include mammals, *e.g.*, humans and other primates; cows, pigs, horses, and farming (agricultural) animals; dogs, cats, and other domesticated pets; mice, rats, and transgenic non-human animals.

Administration of an active amount of an oligonucleotide of the present invention is 30 defined as an amount effective, at dosages and for periods of time necessary to achieve the desired result. For example, an active amount of an oligonucleotide may vary according to

factors such as the type of cell, the oligonucleotide used, and for *in vivo* uses the disease state, age, sex, and weight of the individual, and the ability of the oligonucleotide to elicit a desired response in the individual. Establishment of therapeutic levels of oligonucleotides within the cell is dependent upon the rates of uptake and efflux or degradation. Decreasing the degree of 5 degradation prolongs the intracellular half-life of the oligonucleotide. Thus, chemically-modified oligonucleotides, *e.g.*, with modification of the phosphate backbone, may require different dosing.

The exact dosage of an oligonucleotide and number of doses administered will depend upon the data generated experimentally and in clinical trials. Several factors such as the desired 10 effect, the delivery vehicle, disease indication, and the route of administration, will affect the dosage. Dosages can be readily determined by one of ordinary skill in the art and formulated into the subject pharmaceutical compositions. Preferably, the duration of treatment will extend at least through the course of the disease symptoms.

Dosage regimen may be adjusted to provide the optimum therapeutic response. For 15 example, the oligonucleotide may be repeatedly administered, *e.g.*, several doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation. One of ordinary skill in the art will readily be able to determine appropriate doses and schedules of administration of the subject oligonucleotides, whether the oligonucleotides are to be administered to cells or to subjects.

Physical methods of introducing nucleic acids include injection of a solution containing 20 the nucleic acid, bombardment by particles covered by the nucleic acid, soaking the cell or organism in a solution of the nucleic acid, or electroporation of cell membranes in the presence of the nucleic acid. A viral construct packaged into a viral particle would accomplish both efficient introduction of an expression construct into the cell and transcription of nucleic acid 25 encoded by the expression construct. Other methods known in the art for introducing nucleic acids to cells may be used, such as lipid-mediated carrier transport, chemical-mediated transport, such as calcium phosphate, and the like. Thus the nucleic acid may be introduced along with components that perform one or more of the following activities: enhance nucleic acid uptake by the cell, inhibit annealing of single strands, stabilize the single strands, or other-wise increase 30 inhibition of the target gene.

Nucleic acid may be directly introduced into the cell (*i.e.*, intracellularly); or introduced

extracellularly into a cavity, interstitial space, into the circulation of an organism, introduced orally or by inhalation, or may be introduced by bathing a cell or organism in a solution containing the nucleic acid. Vascular or extravascular circulation, the blood or lymph system, and the cerebrospinal fluid are sites where the nucleic acid may be introduced.

5 The cell with the target gene may be derived from or contained in any organism. The organism may a plant, animal, protozoan, bacterium, virus, or fungus. The plant may be a monocot, dicot or gymnosperm; the animal may be a vertebrate or invertebrate. Preferred microbes are those used in agriculture or by industry, and those that are pathogenic for plants or animals.

10 Alternatively, vectors, e.g., transgenes encoding a siRNA of the invention can be engineered into a host cell or transgenic animal using art recognized techniques.

Another use for the nucleic acids of the present invention (or vectors or transgenes encoding same) is a functional analysis to be carried out in eukaryotic cells, or eukaryotic non-human organisms, preferably mammalian cells or organisms and most preferably human cells, 15 e.g. cell lines such as HeLa or 293 or rodents, e.g. rats and mice. By administering a suitable nucleic acid of the invention which is sufficiently complementary to a target mRNA sequence to direct target-specific RNA interference, a specific knockout or knockdown phenotype can be obtained in a target cell, e.g. in cell culture or in a target organism.

Thus, a further subject matter of the invention is a eukaryotic cell or a eukaryotic non-20 human organism exhibiting a target gene-specific knockout or knockdown phenotype comprising a fully or at least partially deficient expression of at least one endogenous target gene wherein said cell or organism is transfected with at least one vector comprising DNA encoding an RNAi agent capable of inhibiting the expression of the target gene. It should be noted that the present invention allows a target-specific knockout or knockdown of several 25 different endogenous genes due to the specificity of the RNAi agent.

Gene-specific knockout or knockdown phenotypes of cells or non-human organisms, particularly of human cells or non-human mammals may be used in analytic to procedures, e.g. in the functional and/or phenotypical analysis of complex physiological processes such as analysis of gene expression profiles and/or proteomes. Preferably the analysis is carried out by 30 high throughput methods using oligonucleotide based chips.

#### *Assays of Oligonucleotide Stability*

In some embodiments, the oligonucleotides of the invention are stabilized, *i.e.*, substantially resistant to endonuclease and exonuclease degradation. An oligonucleotide is defined as being substantially resistant to nucleases when it is at least about 3-fold more resistant to attack by an endogenous cellular nuclease, and is highly nuclease resistant when it is at least 5 about 6-fold more resistant than a corresponding oligonucleotide. This can be demonstrated by showing that the oligonucleotides of the invention are substantially resistant to nucleases using techniques which are known in the art.

One way in which substantial stability can be demonstrated is by showing that the oligonucleotides of the invention function when delivered to a cell, *e.g.*, that they reduce 10 transcription or translation of target nucleic acid molecules, *e.g.*, by measuring protein levels or by measuring cleavage of mRNA. Assays which measure the stability of target RNA can be performed at about 24 hours post-transfection (*e.g.*, using Northern blot techniques, RNase Protection Assays, or QC-PCR assays as known in the art). Alternatively, levels of the target protein can be measured. Preferably, in addition to testing the RNA or protein levels of interest, 15 the RNA or protein levels of a control, non-targeted gene will be measured (*e.g.*, actin, or preferably a control with sequence similarity to the target) as a specificity control. RNA or protein measurements can be made using any art-recognized technique. Preferably, measurements will be made beginning at about 16-24 hours post transfection. (M. Y. Chiang, *et al.* 1991. J Biol Chem. 266:18162-71; T. Fisher, *et al.* 1993. Nucleic Acids Research. 21 3857).

20 The ability of an oligonucleotide composition of the invention to inhibit protein synthesis can be measured using techniques which are known in the art, for example, by detecting an inhibition in gene transcription or protein synthesis. For example, Nuclease S1 mapping can be performed. In another example, Northern blot analysis can be used to measure the presence of RNA encoding a particular protein. For example, total RNA can be prepared over a cesium 25 chloride cushion (see, *e.g.*, Ausebel *et al.*, 1987. Current Protocols in Molecular Biology (Greene & Wiley, New York)). Northern blots can then be made using the RNA and probed (see, *e.g.*, *Id.*). In another example, the level of the specific mRNA produced by the target protein can be measured, *e.g.*, using PCR. In yet another example, Western blots can be used to measure the amount of target protein present. In still another embodiment, a phenotype 30 influenced by the amount of the protein can be detected. Techniques for performing Western blots are well known in the art, see, *e.g.*, Chen *et al.* J. Biol. Chem. 271:28259.

In another example, the promoter sequence of a target gene can be linked to a reporter gene and reporter gene transcription (e.g., as described in more detail below) can be monitored. Alternatively, oligonucleotide compositions that do not target a promoter can be identified by fusing a portion of the target nucleic acid molecule with a reporter gene so that the reporter gene 5 is transcribed. By monitoring a change in the expression of the reporter gene in the presence of the oligonucleotide composition, it is possible to determine the effectiveness of the oligonucleotide composition in inhibiting the expression of the reporter gene. For example, in one embodiment, an effective oligonucleotide composition will reduce the expression of the reporter gene.

10 A "reporter gene" is a nucleic acid that expresses a detectable gene product, which may be RNA or protein. Detection of mRNA expression may be accomplished by Northern blotting and detection of protein may be accomplished by staining with antibodies specific to the protein. Preferred reporter genes produce a readily detectable product. A reporter gene may be operably linked with a regulatory DNA sequence such that detection of the reporter gene product provides 15 a measure of the transcriptional activity of the regulatory sequence. In preferred embodiments, the gene product of the reporter gene is detected by an intrinsic activity associated with that product. For instance, the reporter gene may encode a gene product that, by enzymatic activity, gives rise to a detectable signal based on color, fluorescence, or luminescence. Examples of reporter genes include, but are not limited to, those coding for chloramphenicol acetyl 20 transferase (CAT), luciferase, beta-galactosidase, and alkaline phosphatase.

One skilled in the art would readily recognize numerous reporter genes suitable for use in the present invention. These include, but are not limited to, chloramphenicol acetyltransferase (CAT), luciferase, human growth hormone (hGH), and beta-galactosidase. Examples of such reporter genes can be found in F. A. Ausubel *et al.*, Eds., Current Protocols in Molecular 25 Biology, John Wiley & Sons, New York, (1989). Any gene that encodes a detectable product, e.g., any product having detectable enzymatic activity or against which a specific antibody can be raised, can be used as a reporter gene in the present methods.

One reporter gene system is the firefly luciferase reporter system. (Gould, S. J., and Subramani, S. 1988. *Anal. Biochem.*, 7:404-408 incorporated herein by reference). The 30 luciferase assay is fast and sensitive. In this assay, a lysate of the test cell is prepared and combined with ATP and the substrate luciferin. The encoded enzyme luciferase catalyzes a

rapid, ATP dependent oxidation of the substrate to generate a light-emitting product. The total light output is measured and is proportional to the amount of luciferase present over a wide range of enzyme concentrations.

CAT is another frequently used reporter gene system; a major advantage of this system is 5 that it has been an extensively validated and is widely accepted as a measure of promoter activity. (Gorman C. M., Moffat, L. F., and Howard, B. H. 1982. Mol. Cell. Biol., 2:1044-1051). In this system, test cells are transfected with CAT expression vectors and incubated with the candidate substance within 2-3 days of the initial transfection. Thereafter, cell extracts are prepared. The extracts are incubated with acetyl CoA and radioactive chloramphenicol.

10 Following the incubation, acetylated chloramphenicol is separated from nonacetylated form by thin layer chromatography. In this assay, the degree of acetylation reflects the CAT gene activity with the particular promoter.

Another suitable reporter gene system is based on immunologic detection of hGH. This system is also quick and easy to use. (Selden, R., Burke-Howie, K. Rowe, M. E., Goodman, H. 15 M., and Moore, D. D. (1986), Mol. Cell, Biol., 6:3173-3179 incorporated herein by reference). The hGH system is advantageous in that the expressed hGH polypeptide is assayed in the media, rather than in a cell extract. Thus, this system does not require the destruction of the test cells. It will be appreciated that the principle of this reporter gene system is not limited to hGH but 20 rather adapted for use with any polypeptide for which an antibody of acceptable specificity is available or can be prepared.

In one embodiment, nuclease stability of a double-stranded oligonucleotide of the invention is measured and compared to a control, *e.g.*, an RNAi molecule typically used in the art (*e.g.*, a duplex oligonucleotide of less than 25 nucleotides in length and comprising 2 nucleotide base overhangs) or an unmodified RNA duplex with blunt ends.

25 The target RNA cleavage reaction achieved using the siRNAs of the invention is highly sequence specific. Sequence identity may be determined by sequence comparison and alignment algorithms known in the art. To determine the percent identity of two nucleic acid sequences (or of two amino acid sequences), the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in the first sequence or second sequence for optimal alignment). A 30 preferred, non-limiting example of a local alignment algorithm utilized for the comparison of sequences is the algorithm of Karlin and Altschul (1990) Proc. Natl. Acad. Sci. USA 87:2264-

68, modified as in Karlin and Altschul (1993) Proc. Natl. Acad. Sci. USA 90:5873-77. Such an algorithm is incorporated into the BLAST programs (version 2.0) of Altschul, et al. (1990) J. Mol. Biol. 215:403-10. Additionally, numerous commercial entities, such as Dharmacon, and Invitrogen provide access to algorithms on their website. The Whitehead Institute also offers a  
5 free siRNA Selection Program. Greater than 90% sequence identity, e.g., 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or even 100% sequence identity, between the siRNA and the portion of the target gene is preferred. Alternatively, the siRNA may be defined functionally as a nucleotide sequence (or oligonucleotide sequence) that is capable of hybridizing with a portion of the target gene transcript. Examples of stringency conditions for polynucleotide hybridization  
10 are provided in Sambrook, J., E. F. Fritsch, and T. Maniatis, 1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., chapters 9 and 11, and Current Protocols in Molecular Biology, 1995, F. M. Ausubel et al., eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4, incorporated herein by reference.

*Therapeutic use*

15 By inhibiting the expression of a gene, the oligonucleotide compositions of the present invention can be used to treat any disease involving the expression of a protein. Examples of diseases that can be treated by oligonucleotide compositions, just to illustrate, include: cancer, retinopathies, autoimmune diseases, inflammatory diseases (*i.e.*, ICAM-1 related disorders, Psoriasis, Ulcerative Colitus, Crohn's disease), metabolic, viral diseases (*i.e.*, HIV, Hepatitis C, 20 flu), miRNA disorders, and cardiovascular diseases.

In one embodiment, *in vitro* treatment of cells with oligonucleotides can be used for ex vivo therapy of cells removed from a subject (*e.g.*, for treatment of leukemia or viral infection) or for treatment of cells which did not originate in the subject, but are to be administered to the subject (*e.g.*, to eliminate transplantation antigen expression on cells to be transplanted into a 25 subject). In addition, *in vitro* treatment of cells can be used in non-therapeutic settings, *e.g.*, to evaluate gene function, to study gene regulation and protein synthesis or to evaluate improvements made to oligonucleotides designed to modulate gene expression or protein synthesis. *In vivo* treatment of cells can be useful in certain clinical settings where it is desirable to inhibit the expression of a protein. There are numerous medical conditions for which 30 antisense therapy is reported to be suitable (see, *e.g.*, U.S. Pat. No. 5,830,653) as well as respiratory syncytial virus infection (WO 95/22,553) influenza virus (WO 94/23,028), and

malignancies (WO 94/08,003). Other examples of clinical uses of antisense sequences are reviewed, *e.g.*, in Glaser. 1996. *Genetic Engineering News* 16:1. Exemplary targets for cleavage by oligonucleotides include, *e.g.*, protein kinase Ca, ICAM-1, c-raf kinase, p53, c-myb, and the bcr/abl fusion gene found in chronic myelogenous leukemia.

5 The subject nucleic acids can be used in RNAi-based therapy in any animal having RNAi pathway, such as human, non-human primate, non-human mammal, non-human vertebrates, rodents (mice, rats, hamsters, rabbits, etc.), domestic livestock animals, pets (cats, dogs, etc.), *Xenopus*, fish, insects (*Drosophila*, etc.), and worms (*C. elegans*), etc.

10 The invention provides methods for inhibiting or preventing in a subject, a disease or condition associated with an aberrant or unwanted target gene expression or activity, by administering to the subject a nucleic acid of the invention. If appropriate, subjects are first treated with a priming agent so as to be more responsive to the subsequent RNAi therapy. Subjects at risk for a disease which is caused or contributed to by aberrant or unwanted target gene expression or activity can be identified by, for example, any or a combination of diagnostic 15 or prognostic assays known in the art. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the target gene aberrancy, such that a disease or disorder is prevented or, alternatively, delayed in its progression. Depending on the type of target gene aberrancy, for example, a target gene, target gene agonist or target gene antagonist agent can be used for treating the subject.

20 In another aspect, the invention pertains to methods of modulating target gene expression, protein expression or activity for therapeutic purposes. Accordingly, in an exemplary embodiment, the methods of the invention involve contacting a cell capable of expressing target gene with a nucleic acid of the invention that is specific for the target gene or protein (*e.g.*, is specific for the mRNA encoded by said gene or specifying the amino acid 25 sequence of said protein) such that expression or one or more of the activities of target protein is modulated. These methods can be performed in vitro (*e.g.*, by culturing the cell with the agent), in vivo (*e.g.*, by administering the agent to a subject), or ex vivo. The subjects may be first treated with a priming agent so as to be more responsive to the subsequent RNAi therapy if desired. As such, the present invention provides methods of treating a subject afflicted with a 30 disease or disorder characterized by aberrant or unwanted expression or activity of a target gene polypeptide or nucleic acid molecule. Inhibition of target gene activity is desirable in situations

in which target gene is abnormally unregulated and/or in which decreased target gene activity is likely to have a beneficial effect.

Thus the therapeutic agents of the invention can be administered to subjects to treat (prophylactically or therapeutically) disorders associated with aberrant or unwanted target gene activity. In conjunction with such treatment, pharmacogenomics (i.e., the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, a physician or clinician may consider applying knowledge obtained in relevant pharmacogenomics studies in determining whether to administer a therapeutic agent as well as tailoring the dosage and/or therapeutic regimen of treatment with a therapeutic agent. Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons.

Thus, the present invention provides compositions and methods to provide polynucleotides *in vivo* by (1) synthesizing a polynucleotide, which contains hydrophobic entities (2) mixing this hydrophobicly modified polynucleotide with a fat emulsion to form micelles; and (3) administering the micelle to the patient or animal, for example intravenously, subcutaneously, topically, via catheter locally or orally (Figure 11).

Nucleic acid molecules, or compositions comprising nucleic acid molecules, described herein may be used for delivery to any tissue or target. An example of a tissue or target is skin. The following exemplary discussion relates to use of the compositions of the invention for delivery of polynucleotides to the skin. The discussion is exemplary only and is not intended to limit the type of delivery or target useful according to the invention.

The compositions of the invention may be used for example to promote wound healing (including chronic wounds such as ulcers), and/or for prevention, reduction or inhibition of scarring, and/or promotion of re-epithelialisation of wounds. Such molecules may also be used for treatment or prevention of diseases, disorders or conditions such as treatment of cleft lip and palate (for example in conjunction with surgical repair of such conditions), reduction or inhibition of scarring and accelerated healing of tendons, and promotion of epithelial regeneration at sites of epithelial damage. In some aspects, nucleic acid molecules associated with the invention may also be used in treatment and/or prevention of fibrotic disorders,

including pulmonary fibrosis, liver cirrhosis, scleroderma and glomerulonephritis, lung fibrosis, liver fibrosis, skin fibrosis, muscle fibrosis, radiation fibrosis, kidney fibrosis, proliferative vitreoretinopathy and uterine fibrosis.

A therapeutically effective amount of a nucleic acid molecule described herein may in some embodiments be an amount sufficient to bring about promotion of wound healing and/or inhibition of scarring and/or promotion of epithelial regeneration. The extent of promotion of wound healing and/or inhibition of scarring, or epithelial regeneration will in some instances be determined by, for example, a doctor or clinician. A suitable assessment of the extent of promotion of wound healing and/or the inhibition of scarring, or promotion of epithelial regeneration, may in some instances be determined by the doctor or clinician.

The ability of nucleic acid molecules associated with the invention to promote the healing of wounds may in some instances be measured with reference to properties exhibited by treated wounds. As used herein, a "treated wound" refers to a wound exposed to a therapeutically effective amount of a medicament such as a nucleic acid molecule of the invention, or a wound which has received treatment in accordance with the methods of the invention. In some instances, promotion of the healing of treated wounds may be indicated by an increased rate of epithelialisation as compared to control wounds. A molecule that is effective in accelerating the healing of wounds may in some instances be a molecule that promotes a more rapid re-constitution of a functional epithelial layer over a wounded area than would otherwise be the case. Promotion of healing of treated wounds can also in some embodiments be indicated by decreased width of a wound compared to control wounds at comparable time points.

As used herein, promotion of wound healing can encompass any increase in the rate of healing of a treated wound as compared with the rate of healing occurring in a control-treated or untreated wound. In some instances, promotion of wound healing may be assessed with respect to either comparison of the rate of re-epithelialisation achieved in treated and control wounds, or comparison of the relative width of treated and control wounds at comparable time points. In some aspects, a molecule that promotes wound healing may be a molecule that, upon administration, causes the wound to exhibit an increased rate of re-epithelialisation and/or a reduction of width compared to control wounds at comparable time points. In some embodiments, the promotion of wound healing may give rise to a rate of wound healing that is

5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 100% greater than the rate occurring in control wounds.

Methods and compositions associated with the invention may be used for treatment of wounds of subjects that may otherwise be prone to defective, delayed or otherwise impaired re-epithelialisation, such as dermal wounds in the aged. Other non-limiting examples of conditions or disorders in which wound healing is associated with delayed or otherwise impaired re-epithelialisation include subjects suffering from diabetes, subjects with polypharmacy (for example as a result of old age), post-menopausal women, subjects susceptible to pressure injuries (for example paraplegics), subjects with venous disease, clinically obese subjects, subjects receiving chemotherapy, subjects receiving radiotherapy, subjects receiving steroid treatment, and immuno-compromised subjects. In some instances, defective re-epithelialisation response can contribute to infections at the wound site, and to the formation of chronic wounds such as ulcers.

In some aspects, chronic wounds exhibiting delayed wound healing response may be treated using methods and compositions associated with the invention. As used herein, a wound may be considered chronic if it does not show any healing tendency within approximately eight weeks of formation when subject to appropriate (conventional) therapeutic treatment. Examples of chronic wounds include venous ulcers, diabetic ulcers and decubitus ulcers, however chronic wounds may arise from otherwise normal acute injuries at any time. In some instances, chronic wounds can arise as a result of infection of the wound site, inadequate wound treatment, progressive tissue breakdown caused by venous, arterial, or metabolic vascular disease, pressure, radiation damage, or tumor. In some embodiments, methods associated with the invention may promote the re-epithelialisation of chronic wounds, and may also inhibit scarring associated with wound healing.

Methods associated with the invention are applied to prevention of acute wounds in subjects predisposed to impaired wound healing developing into chronic wounds. In other aspects, methods associated with the invention are applied to promote accelerated wound healing while preventing, reducing or inhibiting scarring for use in general clinical contexts. In some aspects, this can involve the treatment of surgical incisions and application of such methods may result in the prevention, reduction or inhibition of scarring that may otherwise occur on such healing. Such treatment may result in the scars being less noticeable and exhibiting regeneration of a more normal skin structure. In other embodiments, the wound that

is treated is not a wound caused by a surgical incision. The wound may be subject to continued care and continued application of medicaments to encourage re-epithelialisation and closure of the wound.

In some aspects, methods associated with the invention may also be used in the treatment 5 of wounds associated with grafting procedures. This can involve treatment at a graft donor site and/or at a graft recipient site. Grafts can in some embodiments involve skin, artificial skin, or skin substitutes. Methods associated with the invention can also be used for promoting 10 epithelial regeneration. As used herein, promotion of epithelial regeneration encompasses any increase in the rate of epithelial regeneration as compared to the regeneration occurring in a control-treated or untreated epithelium. The rate of epithelial regeneration attained can in some 15 instances be compared with that taking place in control-treated or untreated epithelia using any suitable model of epithelial regeneration known in the art. Promotion of epithelial regeneration may be of use to induce effective re-epithelialisation in contexts in which the re-epithelialisation response is impaired, inhibited, retarded or otherwise defective. Promotion of epithelial 20 regeneration may be also effected to accelerate the rate of defective or normal epithelial regeneration responses in subjects suffering from epithelial damage.

Some instances where re-epithelialisation response may be defective include conditions such as pemphigus, Hailey-Hailey disease (familial benign pemphigus), toxic epidermal 25 necrolysis (TEN)/Lyell's syndrome, epidermolysis bullosa, cutaneous leishmaniasis and actinic keratosis. Defective re-epithelialisation of the lungs may be associated with idiopathic pulmonary fibrosis (IPF) or interstitial lung disease. Defective re-epithelialisation of the eye may be associated with conditions such as partial limbal stem cell deficiency or corneal erosions. Defective re-epithelialisation of the gastrointestinal tract or colon may be associated with 30 conditions such as chronic anal fissures (fissure in ano), ulcerative colitis or Crohn's disease, and other inflammatory bowel disorders.

In some aspects, methods associated with the invention are used to prevent, reduce or otherwise inhibit scarring. This can be applied to any site within the body and any tissue or organ, including the skin, eye, nerves, tendons, ligaments, muscle, and oral cavity (including the lips and palate), as well as internal organs (such as the liver, heart, brain, abdominal cavity, 35 pelvic cavity, thoracic cavity, guts and reproductive tissue). In the skin, treatment may change the morphology and organization of collagen fibers and may result in making the scars less visible and blend in with the surrounding skin. As used herein, prevention, reduction or

inhibition of scarring encompasses any degree of prevention, reduction or inhibition in scarring as compared to the level of scarring occurring in a control-treated or untreated wound.

Prevention, reduction or inhibition of dermal scarring can be assessed and/or measured with reference to microscopic and/or macroscopic characteristic of a treated scar as compared to the appearance of an untreated scar. As used herein, a "treated scar" refers to a scar formed on healing of a treated wound, whereas an "untreated scar" refers to a scar formed on healing of an untreated wound, or a wound treated with placebo or standard care. Suitable controls for comparison may involve matching of scar age, site, size and subject. Macroscopic assessment of scars may involve examination of parameters such as the color, height, surface texture and stiffness of the scar. In some instances, inhibition or reduction of scarring may be demonstrated when the pigmentation or redness of a treated scar more closely resembles that of unscarred skin than does the pigmentation of an untreated scar. In some instances, inhibition or reduction of scarring may be demonstrated when the height of a treated scar more closely resembles that of unscarred skin than does the height of an untreated scar. In some instances, inhibition or reduction of scarring may be demonstrated when the surface texture of a treated scar more closely resembles that of unscarred skin than does the surface texture of an untreated scar. In some instances inhibition or reduction of scarring may be demonstrated when the stiffness of a treated scar more closely resembles that of unscarred skin than does the stiffness of an untreated scar. An overall assessment of scarring can also be made using, for example, a Visual Analogue Scale or a digital assessment scale.

Microscopic assessment of scars may involve examination of parameters such as thickness of extracellular matrix (ECM) fibers, orientation of ECM fibers, ECM composition of the scar, and cellularity of the scar. In some instances, inhibition or reduction of scarring may be demonstrated when the thickness of ECM fibers in a treated scar more closely resembles the thickness of ECM fibers found in unscarred skin than does the thickness of fibers found in an untreated scar. In some instances, inhibition or reduction of scarring may be demonstrated when the orientation of ECM fibers in a treated scar more closely resembles the orientation of ECM fibers found in unscarred skin than does the orientation of such fibers found in an untreated scar. In some instances, inhibition or reduction of scarring may be demonstrated when the composition of ECM fibers in the dermis of a treated scar more closely resembles the composition of such fibers found in unscarred skin than does the composition found in an untreated scar. In some instances, inhibition or reduction of scarring may be demonstrated when

the cellularity of a treated scar more closely resembles the cellularity of unscarred skin than does the cellularity of an untreated scar.

In some aspects, methods associated with the invention are used for cosmetic purposes, particularly when a wound is located at a prominent body site such as the face, neck and hands.

5 Inhibition of scarring at such sites may contribute to improving the cosmetic appearance of the scar. Scarring is also responsible for a number of deleterious effects such as reduction of physical and mechanical function, particularly in the case of contractile scars (such as hypertrophic scars) and/or situations in which scars are formed across joints and alter the mechanical properties of scarred skin. Methods associated with the invention may be used to

10 prevent, reduce or inhibit scarring of wounds covering joints of the body. In other embodiments, methods associated with the invention may be used to promote accelerated wound healing and/or prevent, reduce or inhibit scarring of wounds at increased risk of forming a contractile scar, and/or of wounds located at sites of high skin tension.

In some embodiments, methods associated with the invention can be applied to

15 promoting wound healing and/or preventing, reducing or inhibiting scarring of wounds in which there is an increased risk of pathological scar formation. Pathological scarring, such as hypertrophic scars and keloids, may have more pronounced deleterious effects than normal scarring. For example, wounds of children, such as burns wounds, are also associated with increased hypertrophic scar formation. In some embodiments, methods described herein for

20 promoting accelerated wound healing and/or preventing, reducing or inhibiting scarring are applied to wounds produced by surgical revision of pathological scars.

Aspects of the invention can be applied to wounds caused by burn injuries. Wound healing in response to burn injuries is frequently associated with adverse scarring outcomes, such as the formation of hypertrophic scars. Methods associated with the invention can be

25 applied to treatment of all injuries involving damage to an epithelial layer, such as injuries to the skin in which the epidermis is damaged. Other non-limiting examples of injuries to epithelial tissue include injuries involving the respiratory epithelia, digestive epithelia or epithelia surrounding internal tissues or organs.

In some aspects, methods associated with the invention can also be used

30 prophylactically, for example at sites where no wound exists but where a wound that would otherwise give rise to a scar or chronic wound is to be formed. For example, medicaments in accordance with the invention may be administered to sites that are to undergo wounding as a

result of elective procedures such as surgery, or to sites that are believed to be at elevated risk of wounding. In some embodiments, administration of a medicament may occur around the time of wounding, or immediately prior to the forming of a wound, for example in the period up to six hours before wounding, or the medicaments may be administered at an earlier time before 5 wounding, for example up to 48 hours before a wound is formed. One of ordinary skill in the art would be able to determine, based on a number of factors, the most advantageous time-frame, formulation and route of administration in a given context. Some non-limiting examples of factors that could be assessed for optimization include the formulation and route of administration of the selected medicament, the dosage of the medicament to be administered, the 10 size and nature of the wound to be formed, and the biological status of the subject, including factors such as the subject's age, health, and predisposition to healing complications or adverse scarring.

In some aspects, methods associated with the invention can be applied to promote wound healing and/or inhibit scarring after a wound has been formed. Administration can occur at any 15 time up until the healing process has been completed, even if the wound has already partially healed. The timing of administration to promote accelerated wound healing and/or prevent, reduce or inhibit scarring can depend on several factors including the nature of the wound in question, the degree of damage within the wound, and the size of the wounded area. In some embodiments, if the wound is large, administration of a medicament relatively late in the healing 20 response may still be able to promote wound healing and/or prevent, reduce or inhibit scarring. In some embodiments, administration of a medicament may occur within the first 24-48 hours after a wound is formed. However, in other embodiments, administration of a medicaments of the invention may be administered 3, 4, 5, 6, 7, 8, 9, 10 or more than 10 days after wounding. Methods and medicaments of the invention may be administered on one or more occasions as 25 necessary in order to promote accelerated wound healing and/or prevent, reduce or inhibit scarring. For instance therapeutically effective amounts of the medicaments may be administered to a wound as often as required until the healing process has been sufficiently advanced or completed. For example, in some embodiments, the medicaments of the invention may be administered daily or twice daily to a wound for at least the first three days following the 30 formation of the wound.

In some instances, the methods or medicaments of the invention may be administered both before and after formation of a wound. It will be appreciated that the amount of a

medicament of the invention that should be applied to a wound depends on a number of factors such as the biological activity and bioavailability of the agent present in the medicament, which depends, among other factors, on the nature of the agent and the mode of administration of the medicament. Other factors in determining a suitable therapeutic amount of a medicament may 5 include the half-life of the agent in the subject being treated; the specific condition to be treated, and characteristics of the subject such as the age of the subject. The frequency of administration will also be influenced by the above-mentioned factors including the half-life of the chosen agent. Typically a cream or ointment containing an agent of the invention may be administered to a target tissue such that the concentration of the agent at a wound is maintained at a level 10 suitable for having a therapeutic effect. In some instances, this may require administration daily or several times daily.

Compositions and medicaments associated with the invention may be administered by any suitable route capable of achieving the desired effect of promoting wound healing and/or preventing, reducing or inhibiting scarring. In some embodiments, the medicaments are 15 administered locally at a wound site. For dermal wounds, agents of the invention may be administered by means of intradermal injection. Accordingly, compositions and medicaments associated with the invention may comprise injectable solutions, and in some instances may be injected, for example, around the margins of a site of epithelial damage or a site likely to be damaged. In some embodiments, compositions and medicaments associated with the invention 20 may also be administered in a topical form to promote accelerated wound healing and/or prevention, reduction or inhibition of scarring. In some embodiments, administration of compositions and medicaments associated with the invention comprise part of an initial treatment of a wound or scar, while in other embodiments, administration of compositions and medicaments associated with the invention comprise follow-up care for a wound or scar.

25 *Target genes*

It should be appreciated that based on the RNAi molecules designed and disclosed herein, one of ordinary skill in the art would be able to design such RNAi molecules to target a variety of different genes depending on the context and intended use. For purposes of 30 promoting wound healing or preventing, reducing or inhibiting scarring, one of ordinary skill in the art would appreciate that a variety of suitable target genes could be identified based at least in part on the known or predicted functions of the genes, and/or the known or predicted expression patterns of the genes. Several non-limiting examples of genes that could be targeted

by RNAi molecules for promoting wound healing or preventing, reducing or inhibiting scarring include genes that encode for the following proteins: Transforming growth factor  $\beta$  (TGF $\beta$ 1, TGF $\beta$ 2, TGF $\beta$ 3), Osteopontin, Connective tissue growth factor (CTGF), Platelet-derived growth factor (PDGF), Hypoxia inducible factor-1 $\alpha$  (HIF1 $\alpha$ ), Collagen I and/or III, Prolyl 4-

5 hydroxylase (P4H), Procollagen C-protease (PCP), Matrix metalloproteinase 2, 9 (MMP2, 9), Integrins, Connexin, Histamine H1 receptor, Tissue transglutaminase, Mammalian target of rapamycin (mTOR), HoxB13, VEGF, IL-6, SMAD proteins, Ribosomal protein S6 kinases (RSP6) and Cyclooxygenase-2 (COX-2).

Transforming growth factor  $\beta$  proteins, for which three isoforms exist in mammals

10 (TGF $\beta$ 1, TGF $\beta$ 2, TGF $\beta$ 3), are secreted proteins belonging to a superfamily of growth factors involved in the regulation of many cellular processes including proliferation, migration, apoptosis, adhesion, differentiation, inflammation, immuno-suppression and expression of extracellular proteins. These proteins are produced by a wide range of cell types including epithelial, endothelial, hematopoietic, neuronal, and connective tissue cells. Representative  
15 Genbank accession numbers providing DNA and protein sequence information for human TGF $\beta$ 1, TGF $\beta$ 2 and TGF $\beta$ 3 are BT007245, BC096235, and X14149, respectively.

Osteopontin (OPN), also known as Secreted phosphoprotein 1 (SPP1), Bone Sialoprotein 1 (BSP-1), and early T-lymphocyte activation (ETA-1) is a secreted glycoprotein protein that binds to hydroxyapatite. OPN has been implicated in a variety of biological processes including

20 bone remodeling, immune functions, chemotaxis, cell activation and apoptosis. Osteopontin is produced by a variety of cell types including fibroblasts, preosteoblasts, osteoblasts, osteocytes, odontoblasts, bone marrow cells, hypertrophic chondrocytes, dendritic cells, macrophages, smooth muscle, skeletal muscle myoblasts, endothelial cells, and extraosseous (non-bone) cells in the inner ear, brain, kidney, deciduum, and placenta. Representative Genbank accession  
25 number providing DNA and protein sequence information for human Osteopontin are NM\_000582.2 and X13694.

Connective tissue growth factor (CTGF), also known as Hypertrophic chondrocyte-specific protein 24, is a secreted heparin-binding protein that has been implicated in wound healing and scleroderma. Connective tissue growth factor is active in many cell types including  
30 fibroblasts, myofibroblasts, endothelial and epithelial cells. Representative Genbank accession

number providing DNA and protein sequence information for human CTGF are NM\_001901.2 and M92934.

The Platelet-derived growth factor (PDGF) family of proteins, including several isoforms, are secreted mitogens. PDGF proteins are implicated in wound healing, at least in part, because they are released from platelets following wounding. Representative Genbank accession numbers providing DNA and protein sequence information for human PDGF genes and proteins include X03795 (PDGFA), X02811 (PDGFB), AF091434 (PDGFC), AB033832 (PDGFD).

Hypoxia inducible factor-1 $\alpha$  (HIF1 $\alpha$ ), is a transcription factor involved in cellular response to hypoxia. HIF1 $\alpha$  is implicated in cellular processes such as embryonic vascularization, tumor angiogenesis and pathophysiology of ischemic disease. A representative Genbank accession number providing DNA and protein sequence information for human HIF1 $\alpha$  is U22431.

Collagen proteins are the most abundant mammalian proteins and are found in tissues such as skin, tendon, vascular, ligature, organs, and bone. Collagen I proteins (such as COL1A1 and COL1A2) are detected in scar tissue during wound healing, and are expressed in the skin. Collagen III proteins (including COL3A1) are detected in connective tissue in wounds (granulation tissue), and are also expressed in skin. Representative Genbank accession numbers providing DNA and protein sequence information for human Collagen proteins include: Z74615 (COL1A1), J03464 (COL1A2) and X14420 (COL3A1).

Prolyl 4-hydroxylase (P4H), is involved in production of collagen and in oxygen sensing. A representative Genbank accession number providing DNA and protein sequence information for human P4H is AY198406.

Matrix metalloproteinase 2, 9 (MMP2, 9) belong to the metzincin metalloproteinase superfamily and are zinc-dependent endopeptidases. These proteins are implicated in a variety of cellular processes including tissue repair. Representative Genbank accession numbers providing DNA and protein sequence information for human MMP proteins are M55593 (MMP2) and J05070 (MMP9).

Integrins are a family of proteins involved in interaction and communication between a cell and the extracellular matrix. Vertebrates contain a variety of integrins including  $\alpha_1\beta_1$ ,  $\alpha_2\beta_1$ ,  $\alpha_4\beta_1$ ,  $\alpha_5\beta_1$ ,  $\alpha_6\beta_1$ ,  $\alpha_L\beta_2$ ,  $\alpha_M\beta_2$ ,  $\alpha_{IIb}\beta_3$ ,  $\alpha_v\beta_3$ ,  $\alpha_v\beta_5$ ,  $\alpha_v\beta_6$ ,  $\alpha_6\beta_4$ .

Connexins are a family of vertebrate transmembrane proteins that form gap junctions.

Several examples of Connexins, with the accompanying gene name shown in brackets, include Cx23 (GJE1), Cx25 (GJB7), Cx26 (GJB2), Cx29 (GJE1), Cx30 (GJB6), Cx30.2 (GJC3), Cx30.3 (GJB4), Cx31 (GJB3), Cx31.1 (GJB5), Cx31.9 (GJC1/GJD3), Cx32 (GJB1), Cx33 (GJA6),  
5 Cx36 (GJD2/GJA9), Cx37 (GJA4), Cx39 (GJD4), Cx40 (GJA5), Cx40.1 (GJD4), Cx43 (GJA1), Cx45 (GJC1/GJA7), Cx46 (GJA3), Cx47 (GJC2/GJA12), Cx50 (GJA8), Cx59 (GJA10), and Cx62 (GJA10).

Histamine H1 receptor (HRH1) is a metabotropic G-protein-coupled receptor involved in the phospholipase C and phosphatidylinositol (PIP2) signaling pathways. A representative

10 Genbank accession number providing DNA and protein sequence information for human HRH1 is Z34897.

Tissue transglutaminase, also called Protein-glutamine gamma-glutamyltransferase 2, is involved in protein crosslinking and is implicated in biological processes such as apoptosis, cellular differentiation and matrix stabilization. A representative Genbank accession number 15 providing DNA and protein sequence information for human Tissue transglutaminase is M55153.

Mammalian target of rapamycin (mTOR), also known as Serine/threonine-protein kinase mTOR and FK506 binding protein 12-rapamycin associated protein 1 (FRAP1), is involved in regulating cell growth and survival, cell motility, transcription and translation. A representative 20 Genbank accession number providing DNA and protein sequence information for human mTOR is L34075.

HoxB13 belongs to the family of Homeobox proteins and has been linked to functions such as cutaneous regeneration and fetal skin development. A representative Genbank accession number providing DNA and protein sequence information for human HoxB13 is U57052.

25 Vascular endothelial growth factor (VEGF) proteins are growth factors that bind to tyrosine kinase receptors and are implicated in multiple disorders such as cancer, age-related macular degeneration, rheumatoid arthritis and diabetic retinopathy. Members of this protein family include VEGF-A, VEGF-B, VEGF-C and VEGF-D. Representative Genbank accession numbers providing DNA and protein sequence information for human VEGF proteins are  
30 M32977 (VEGF-A), U43368 (VEGF-B), X94216 (VEGF-C), and D89630 (VEGF-D).

Interleukin-6 (IL-6) is a cytokine involved in stimulating immune response to tissue damage. A representative Genbank accession number providing DNA and protein sequence information for human IL-6 is X04430.

5 SMAD proteins (SMAD1-7, 9) are a family of transcription factors involved in regulation of TGF $\beta$  signaling. Representative Genbank accession numbers providing DNA and protein sequence information for human SMAD proteins are U59912 (SMAD1), U59911 (SMAD2), U68019 (SMAD3), U44378 (SMAD4), U59913 (SMAD5), U59914 (SMAD6), AF015261 (SMAD7), and BC011559 (SMAD9).

10 Ribosomal protein S6 kinases (RSK6) represent a family of serine/threonine kinases involved in activation of the transcription factor CREB. A representative Genbank accession number providing DNA and protein sequence information for human Ribosomal protein S6 kinase alpha-6 is AF184965.

15 Cyclooxygenase-2 (COX-2), also called Prostaglandin G/H synthase 2 (PTGS2), is involved in lipid metabolism and biosynthesis of prostanooids and is implicated in inflammatory disorders such as rheumatoid arthritis. A representative Genbank accession number providing DNA and protein sequence information for human COX-2 is AY462100.

The invention also encompasses diagnostic uses as well as prophylactic, therapeutic and research uses.

20 Formulations include sterile or non-sterile aqueous or non-aqueous solutions, suspensions, and emulsions.

The invention also includes articles, which refers to any one or collection of components. In some embodiments the articles are kits. The articles include pharmaceutical or diagnostic grade compounds of the invention in one or more containers. The article may include instructions or labels promoting or describing the use of the compounds of the invention.

25 As used herein, “promoted” includes all methods of doing business including methods of education, hospital and other clinical instruction, pharmaceutical industry activity including pharmaceutical sales, and any advertising or other promotional activity including written, oral and electronic communication of any form, associated with compositions of the invention in connection with treatment, diagnosis, or prophylaxis of a disease.

30 “Instructions” can define a component of promotion, and typically involve written instructions on or associated with packaging of compositions of the invention. Instructions also can include any oral or electronic instructions provided in any manner.

Thus the agents described herein may, in some embodiments, be assembled into pharmaceutical or diagnostic or research kits to facilitate their use in therapeutic, diagnostic or research applications. A kit may include one or more containers housing the components of the invention and instructions for use. Specifically, such kits may include one or more agents

5 described herein, along with instructions describing the intended therapeutic or diagnostic application and the proper administration of these agents. In certain embodiments agents in a kit may be in a pharmaceutical formulation and dosage suitable for a particular application and for a method of administration of the agents.

The kit may be designed to facilitate use of the methods described herein by physicians

10 and can take many forms. Each of the compositions of the kit, where applicable, may be provided in liquid form (e.g., in solution), or in solid form, (e.g., a dry powder). In certain cases, some of the compositions may be constitutable or otherwise processable (e.g., to an active form), for example, by the addition of a suitable solvent or other species (for example, water or a cell culture medium), which may or may not be provided with the kit. As used herein,

15 "instructions" can define a component of instruction and/or promotion, and typically involve written instructions on or associated with packaging of the invention. Instructions also can include any oral or electronic instructions provided in any manner such that a user will clearly recognize that the instructions are to be associated with the kit, for example, audiovisual (e.g., videotape, DVD, etc.), Internet, and/or web-based communications, etc. The written

20 instructions may be in a form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which instructions can also reflects approval by the agency of manufacture, use or sale for human administration.

The kit may contain any one or more of the components described herein in one or more containers. As an example, in one embodiment, the kit may include instructions for mixing one

25 or more components of the kit and/or isolating and mixing a sample and applying to a subject. The kit may include a container housing agents described herein. The agents may be prepared steriley, packaged in syringe and shipped refrigerated. Alternatively it may be housed in a vial or other container for storage. A second container may have other agents prepared steriley. Alternatively the kit may include the active agents premixed and shipped in a syringe, vial, tube,

30 or other container.

The kit may have a variety of forms, such as a blister pouch, a shrink wrapped pouch, a vacuum sealable pouch, a sealable thermoformed tray, or a similar pouch or tray form, with the accessories loosely packed within the pouch, one or more tubes, containers, a box or a bag. The kit may be sterilized after the accessories are added, thereby allowing the individual accessories  
5 in the container to be otherwise unwrapped. The kits can be sterilized using any appropriate sterilization techniques, such as radiation sterilization, heat sterilization, or other sterilization methods known in the art. The kit may also include other components, depending on the specific application, for example, containers, cell media, salts, buffers, reagents, syringes, needles, a fabric, such as gauze, for applying or removing a disinfecting agent, disposable  
10 gloves, a support for the agents prior to administration etc.

The compositions of the kit may be provided as any suitable form, for example, as liquid solutions or as dried powders. When the composition provided is a dry powder, the powder may be reconstituted by the addition of a suitable solvent, which may also be provided. In  
15 embodiments where liquid forms of the composition are used, the liquid form may be concentrated or ready to use. The solvent will depend on the compound and the mode of use or administration. Suitable solvents for drug compositions are well known and are available in the literature. The solvent will depend on the compound and the mode of use or administration.

The kits, in one set of embodiments, may comprise a carrier means being compartmentalized to receive in close confinement one or more container means such as vials,  
20 tubes, and the like, each of the container means comprising one of the separate elements to be used in the method. For example, one of the containers may comprise a positive control for an assay. Additionally, the kit may include containers for other components, for example, buffers useful in the assay.

The present invention also encompasses a finished packaged and labeled pharmaceutical product. This article of manufacture includes the appropriate unit dosage form in an appropriate vessel or container such as a glass vial or other container that is hermetically sealed. In the case  
25 of dosage forms suitable for parenteral administration the active ingredient is sterile and suitable for administration as a particulate free solution. In other words, the invention encompasses both parenteral solutions and lyophilized powders, each being sterile, and the latter being suitable for reconstitution prior to injection. Alternatively, the unit dosage form may be a solid suitable for  
30 oral, transdermal, topical or mucosal delivery.

In a preferred embodiment, the unit dosage form is suitable for intravenous, intramuscular or subcutaneous delivery. Thus, the invention encompasses solutions, preferably sterile, suitable for each delivery route.

In another preferred embodiment, compositions of the invention are stored in containers 5 with biocompatible detergents, including but not limited to, lecithin, taurocholic acid, and cholesterol; or with other proteins, including but not limited to, gamma globulins and serum albumins. More preferably, compositions of the invention are stored with human serum albumins for human uses, and stored with bovine serum albumins for veterinary uses.

As with any pharmaceutical product, the packaging material and container are designed 10 to protect the stability of the product during storage and shipment. Further, the products of the invention include instructions for use or other informational material that advise the physician, technician or patient on how to appropriately prevent or treat the disease or disorder in question. In other words, the article of manufacture includes instruction means indicating or suggesting a dosing regimen including, but not limited to, actual doses, monitoring procedures (such as 15 methods for monitoring mean absolute lymphocyte counts, tumor cell counts, and tumor size) and other monitoring information.

More specifically, the invention provides an article of manufacture comprising packaging material, such as a box, bottle, tube, vial, container, sprayer, insufflator, intravenous (i.v.) bag, envelope and the like; and at least one unit dosage form of a pharmaceutical agent 20 contained within said packaging material. The invention also provides an article of manufacture comprising packaging material, such as a box, bottle, tube, vial, container, sprayer, insufflator, intravenous (i.v.) bag, envelope and the like; and at least one unit dosage form of each pharmaceutical agent contained within said packaging material. The invention further provides an article of manufacture comprising packaging material, such as a box, bottle, tube, vial, 25 container, sprayer, insufflator, intravenous (i.v.) bag, envelope and the like; and at least one unit dosage form of each pharmaceutical agent contained within said packaging material. The invention further provides an article of manufacture comprising a needle or syringe, preferably packaged in sterile form, for injection of the formulation, and/or a packaged alcohol pad.

In a specific embodiment, an article of manufacture comprises packaging material and a 30 pharmaceutical agent and instructions contained within said packaging material, wherein said instructions indicate a dosing regimen for preventing, treating or managing a subject with a disease.

For the purposes of the invention, ranges may be expressed herein as from "about" one particular value, and/or to "about" another particular value. When such a range is expressed, another embodiment includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by use of the antecedent "about," it will 5 be understood that the particular value forms another embodiment. It will be further understood that the endpoints of each of the ranges are significant both in relation to the other endpoint, and independently of the other endpoint.

Moreover, for the purposes of the present invention, the term "a" or "an" entity refers to one or more of that entity; for example, "a protein" or "a nucleic acid molecule" refers to one or 10 more of those compounds or at least one compound. As such, the terms "a" (or "an"), "one or more" and "at least one" can be used interchangeably herein. It is also to be noted that the terms "comprising", "including", and "having" can be used interchangeably. Furthermore, a compound "selected from the group consisting of" refers to one or more of the compounds in the list that follows, including mixtures (i.e., combinations) of two or more of the compounds.

15 According to the present invention, an isolated, or biologically pure, protein or nucleic acid molecule is a compound that has been removed from its natural milieu. As such, "isolated" and "biologically pure" do not necessarily reflect the extent to which the compound has been purified. An isolated compound of the present invention can be obtained from its natural source, can be produced using molecular biology techniques or can be produced by chemical synthesis.

20 The present invention is further illustrated by the following Examples, which in no way should be construed as further limiting. The entire contents of all of the references (including literature references, issued patents, published patent applications, and co-pending patent applications) cited throughout this application are hereby expressly incorporated by reference.

#### EXAMPLES

##### **Example 1: Neutral nanotransporters for delivery of nucleic acids**

Described herein is the development of neutral fat formulations for nucleic acid delivery. Formulations were developed to allow efficient incorporation of nucleic acids into non-charged mixtures, offering the significant advantage of lower toxicity relative to previously described formulations for nucleic acid delivery.

30 The hydrophobicity of nucleic acids can be increased by conjugating them to sterol-type molecules, as shown in a schematic in Fig. 1C. The sterol-type molecule can be conjugated to either strand of the double stranded nucleic acid molecule. Fig. 2 lists several examples of

chemical modifications that can be used to increase hydrophobicity including 4-pyridyl, 2-pyridyl, isobutyl and indolyl based position 5 uridine modifications. Fig. 2 also shows a schematic depicting double stranded RNA molecules with a variety of chemical modification patterns including hydrophobic modifications that increase the hydrophobicity of the nucleic acid molecule. Proteins or peptides such as protamine (or other Arg rich peptides), spermidine or other similar chemical structures can also be used to block duplex charge and facilitate cellular entry (Fig. 3). Increased hydrophobicity can be achieved through either covalent or non-covalent modifications. Several positively charged chemicals, which might be used for polynucleotide charge blockage are depicted in Fig. 4.

Fig. 5 depicts single stranded polynucleotides, representing a guide strand in a duplex molecule, with a variety of chemical modifications including 2'd, 2'OMe, 2'F, hydrophobic modifications, phosphorothioate modifications, and attachment of conjugates such as "X" in Fig. 5, where X can be a small molecule with high affinity to a PAZ domain, or sterol-type entity. Similarly, Fig. 6 depicts single stranded polynucleotides, representing a passenger strand in a duplex molecule, with proposed structural and chemical compositions of RISC substrate inhibitors. Combinations of chemical modifications can ensure efficient uptake and efficient binding to preloaded RISC complexes.

Fig. 7 depicts structures of polynucleotides, with or without hydrophobic modifications within the polynucleotide, with sterol-type molecules attached. R represents a polycarbonic tail of 9 carbons or longer. Fig. 8 presents examples of naturally occurring phytosterols with a polycarbon chain longer than 8 attached at position 17. More than 250 different types of phytosterols are known. Fig. 9 presents examples of sterol-like structures with variations in the sizes of the polycarbon chains attached at position 17. Optimization of such characteristics can improve uptake properties of the RNAi molecules. Fig. 10 presents data adapted from Martins et al. (J Lipid Research), showing that the percentage of liver uptake and plasma clearance of lipid emulsions containing sterol-type molecules is directly affected by the size of the attached polycarbon chain at position 17. Fig. 11 depicts a micelle formed from a mixture of polynucleotides attached to hydrophobic conjugates and fatty acids. Fig. 12 describes how alteration in lipid composition can affect pharmacokinetic behavior and tissue distribution of hydrophobically modified and/or hydrophobically conjugated polynucleotides. In particular, the use of lipid mixtures that are enriched in linoleic acid and cardiolipin results in preferential uptake by cardiomyocytes.

Nucleic acid molecules that are chemically modified and/or conjugated to sterol-type molecules are capable of self-delivery, as demonstrated in Fig. 13. The double stranded RNA molecules depicted in Fig. 13 are called “sd” RNA molecules, meaning “self-delivering” RNA molecules. Three generations of sd-rxRNA molecules were developed: generation I (GI), 5 generation IIa (GIIa) and generation IIb (GIIb), with variations in the chemical modification patterns incorporated. Highly effective compounds were found to have the following characteristics: antisense strands of 17-21 nucleotides, sense strands of 10-15 nucleotides, single-stranded regions that contained 2-12 phosphorothioate modifications, preferentially 6-8 phosphorothioate modifications, and sense strands in which the majority of nucleotides were 10 2’OMe modified, with or without phosphorothioate modification. As shown in the graph in Fig. 13, these molecules were highly effective in achieving silencing of a target gene. Significantly, any linker chemistry can be used to attach these nucleic acid molecules to hydrophobic moieties such as cholesterol at the 3’ end of the sense strand. Version GIIa-b of these RNA compounds demonstrate that elimination of 2’F content, predicted to significantly decrease toxicity, has no 15 impact on efficacy.

Modifications and conjugations described herein, to increase the hydrophobicity of nucleic acid molecules, can be applied to any type of nucleic acid molecule. Several examples of types of nucleic acids that could be formulated and delivered using methods and compositions described herein are presented in Fig. 14 including conventional siRNAs, longer siRNAs single 20 stranded oligos, antisense, antogamirs and sd-rxRNA. The nucleic acid molecule is modified such that its hydrophobicity is substantially increased. This can be achieved by modifying bases, sugars or nucleic acid backbone or/and by linking a hydrophobic molecule to the nucleic acid. The hydrophobic molecule can be attached anywhere in the compound and can include, for example, a fatty acid, sterol, vitamin, small molecule or peptide. The hydrophobic molecule can 25 be covalently or non covalently attached.

Hydrophobic oligonucleotides (sd-rxRNA) and neutral fat formulations alone were not found to form complexes (Fig. 15). Fig. 15A shows a lack of complex formation with a DOPC:DOPE mixture and Fig. 15B shows a lack of complex formation with Intralipid. The complex formation was evaluated by complexing reagents and evaluating a shift in 30 oligonucleotide band formation using a non-denaturing polyacrylamide gel. The position of the oligonucleotide is determined by staining.

Significantly, the simultaneous presence of DOPC and Cholesterol was found to result in a hydrophobic nucleic acid complexing with neutral lipid formulations (Fig. 16). As low as 1:5 weight (lipid to oligonucleotide) ratio was found to be sufficient to produce a significant fraction of encapsulated oligonucleotide. In the context of DOPC:Cholesterol formulation, other neutral 5 lipids can be added without interfering with particle formation. For example, DOPC:cholesterol: DOPE. Fig. 17 demonstrates that DSPC (saturated fatty acids) can take place of DOPC without any impact on efficiency of complex formation. Figs. 18-19 present results demonstrating DOPC/Intralipid/Cholesterol formulation containing sd-rxRNA (12386). Complexes were still present when diluted to 500 nM from 1  $\mu$ M with Accell media (Fig. 18) 10 and when diluted to 500 nM from 5  $\mu$ M with Accell media (Fig. 19). More RNA appeared to be complexed when diluted to 500 nM from 5  $\mu$ M, than when diluting to 500 nM from 1  $\mu$ M.

Fig. 20 reveals that the presence of both DOPC and Cholesterol in significant proportions was required to enable significant formation of Intralipid-containing neutral fat/oligo particles. A slight complex formation was observed with Intralipid:cholesterol but it 15 was minor compared to the complex formation that occurred in the presence of both DOPC and Cholesterol.

Various additional compounds (i.e., lipids, peptides, small molecules) can be encapsulated into the particle as long as formulation comprises at least 20% of DOPC/Cholesterol-type compounds (Fig. 21). The demonstrated cargo lipid is intralipid. 20 Variation in the identity, amounts and ratios of cargo lipids affects the cellular uptake and tissue distribution characteristics of these compounds. For example, the length of lipid tails and level of saturability will affect differential uptake to liver, lung, fat and cardiomyocytes. Addition of special hydrophobic molecules like vitamins or different forms of sterols can favor distribution to special tissues which are involved in the metabolism of particular compounds. Complexes are 25 formed at different oligonucleotide concentrations, with higher concentrations favoring more efficient complex formation (Figs. 21-22).

Only hydrophobically modified oligonucleotides were found to form complexes with neutral fat formulations (Fig. 23). When a cholesterol-modified sd-rxRNA compound was mixed with 50:50 DOPC/Cholesterol formulation, the sd-rxRNA quantitatively entered into a 30 complex. When rxRNA (Omethyl modified siRNA), regular siRNA or sd-rxRNA, without a hydrophobic compound is mixed with the same formulation, no complex was formed, demonstrating that the combination of a hydrophobic modification of the oligonucleotide with

neutral fat formulations comprising at least 20% of DOPC:Cholesterol-type compounds are required for efficient encapsulation of oligonucleotides into neutral fat formulations.

Neutral lipid based formulations were found to enter cells (i.e., HeLa cells) and effectively silence genes (Fig. 24). Fig. 24B demonstrates that the oligonucleotide/lipid ratio 5 and formulation composition affects the level of silencing. Significantly, no toxicity was observed even at 1 uM concentration. This lack of toxicity is a significant improvement over positively charged traditional formulations (i.e., lipofectamine) which exhibit a drastic toxicity at a much lower dose range. This data demonstrates that neutral fat/oligonucleotide formulations are non toxic or have highly reduced toxicity relative to previously described 10 positively charged formulations, and have a wider therapeutic index.

Fig. 25 demonstrates the efficacies of various formulations in achieving gene silencing *in vitro* with and without serum. Formulation B is DOPC-Cholesterol 50:50 and Formulation C is DOPC-DOPE:Cholesterol 33:33:33.

Figs. 26-27 present data on the size and zeta potential of particles formed upon 15 complexing of neutral fat formulations with hydrophobically modified oligos. While neutral fat by itself forms agglomerates ~ 500-1000 nm in size, addition of increasing concentrations of an oligonucleotide results in the formation of stable small particles (around 60-120nm) which are not charged (Zeta Potential ~ -10). The neutral particles, sized around 50-100 nM, are ideal for systemic administration. The size and charge of the particles is affected by the oligonucleotide/ 20 lipid ratio, lipid mixture composition and lipid ratios within formulation.

A panel of lipids used for formulation preparations is demonstrated in Fig. 28, and 25 schematics of the sterol-type molecules used for formulation preparations is demonstrated in Fig. 29. In some instances, some of the formulations comprising longer chain sterol type molecules have a significantly better cellular uptake and tissue distribution properties. Fig. 30 demonstrates examples of hydrophobic molecules that can be linked to a nucleic acid or included as part of a formulation to improve or alter cellular uptake and tissue distribution.

The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by examples provided, since the examples are intended as a single illustration of one aspect of the invention 30 and other functionally equivalent embodiments are within the scope of the invention. Various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the

appended claims. The advantages and objects of the invention are not necessarily encompassed by each embodiment of the invention.

What is claimed is:

## CLAIMS

1. A composition comprising:  
a hydrophobic modified polynucleotide;  
5 a neutral fatty mixture; and  
optionally a cargo molecule,  
wherein the hydrophobic modified polynucleotide and the neutral fatty mixture forms a  
micelle.
2. The composition of claim 1, wherein the polynucleotide is RNA.
- 10 3. The composition of claim 2, wherein the polynucleotide is double stranded RNA.
4. The composition of any one of claims 1-3, wherein the composition is a  
pharmaceutical composition and includes a pharmaceutically acceptable carrier.
5. The composition of any one of claims 1-4, wherein the composition is sterile.
- 15 6. The composition of any one of claims 1-5, wherein the neutral fatty mixture  
comprises a DOPC (dioleoylphosphatidylcholine).
7. The composition of any one of claims 1-5, wherein the neutral fatty mixture  
comprises a DSPC (distearoylphosphatidylcholine).
8. The composition of any one of claims 6 or 7, wherein the neutral fatty mixture further  
comprises a sterol.
- 20 9. The composition of claim 8, wherein the sterol is cholesterol.
10. The composition of any one of claims 1-5, wherein the neutral fatty mixture  
comprises 20% of a fatty acid derivative of choline.
11. The composition of claim 10, wherein the neutral fatty mixture further comprises  
20% sterol.
- 25 12. The composition of any one of claims 8, 9 or 11, wherein the composition includes  
at least 20% DOPC and at least 20% cholesterol.
13. The composition of any one of claims 1-12, wherein the hydrophobic portion of the  
hydrophobic modified polynucleotide is covalently linked to the polynucleotide.
- 30 14. The composition of any one of claims 1-12, wherein the hydrophobic portion of the  
hydrophobic modified polynucleotide is a sterol.
15. The composition of claim 14, wherein the sterol is a cholesterol, a cholesteryl or  
modified cholesteryl residue.

16. The composition of any one of claims 1-12, wherein the hydrophobic portion of the hydrophobic modified polynucleotide is selected from the group consisting of bile acids, cholic acid or taurocholic acid, deoxycholate, oleyl lithocholic acid, oleyl cholenic acid, glycolipids, phospholipids, sphingolipids, isoprenoids, vitamins, saturated fatty acids, unsaturated fatty acids, 5 fatty acid esters, triglycerides, pyrenes, porphyrines, Texaphyrine, adamantane, acridines, biotin, coumarin, fluorescein, rhodamine, Texas-Red, digoxygenin, dimethoxytrityl, t-butylidimethylsilyl, t-butylidiphenylsilyl, cyanine dyes (e.g. Cy3 or Cy5), Hoechst 33258 dye, psoralen, and ibuprofen.

17. The composition of any one of claims 1-12, wherein the hydrophobic portion of the 10 hydrophobic modified polynucleotide is non-covalently linked to the polynucleotide.

18. The composition of any one of claims 1-13 or 17, wherein the hydrophobic portion of the hydrophobic modified polynucleotide is a polycationic molecule.

19. The composition of claim 18, wherein the polycationic molecule is selected from the group consisting of protamine, arginine rich peptides, and spermine.

15 20. The composition of any one of claims 1-19, wherein the cargo molecule is a lipid, a peptide, vitamin, or a small molecule.

21. The composition of any one of claims 1-19, wherein the cargo molecule is a commercially available fat emulsions available for a variety of purposes selected from the group consisting of parenteral feeding.

20 22. The composition of claim 21, wherein the commercially available fat emulsion is an intralipid or a nutralipid.

23. The composition of any one of claims 1-19, wherein the cargo molecule is a fatty acid mixture containing more than 74% of linoleic acid.

24. The composition of any one of claims 1-19, wherein the cargo molecule is a fatty 25 acid mixture containing at least 6% of cardiolipin.

25. The composition of any one of claims 1-19, wherein the cargo molecule is a fatty acid mixture containing at least 74% of linoleic acid and at least 6% of cardiolipin.

26. The composition of any one of claims 1-25, wherein the polynucleotide is at least 40% modified.

30 27. The composition of any one of claims 1-19, wherein the cargo molecule is a fusogenic lipid and preferably is at least 10% fusogenic lipid.

28. The composition of claims 27, wherein the fusogenic lipid is DOPE.

29. The composition of any one of claims 1-28, wherein the polynucleotide is an isolated double stranded nucleic acid molecule comprising  
a guide strand, wherein the guide strand is 16-28 nucleotides long and has complementarity to a target gene, wherein the 3' terminal 10 nucleotides of the guide strand  
5 include at least two phosphate modifications, and wherein the guide strand has a 5' phosphate modification and includes at least one 2' O-methyl modification or 2'-fluoro modification, and  
a passenger strand, wherein the passenger strand is 8-28 nucleotides long and has complementarity to the guide strand, wherein the passenger strand is linked to the hydrophobic molecule,  
10 wherein the guide strand and the passenger strand form the double stranded nucleic acid molecule.

30. The composition of any one of claims 1-28, wherein the polynucleotide is an isolated double stranded nucleic acid molecule comprising a guide strand and a passenger strand, wherein the guide strand is from 16-29 nucleotides long and is substantially complementary to a  
15 target gene, wherein the passenger strand is from 8-14 nucleotides long and has complementarity to the guide strand, and wherein the guide stand has at least two chemical modifications.

31. The composition of any one of claims 1-28, wherein the polynucleotide is an isolated double stranded nucleic acid molecule comprising a guide strand and a passenger strand,  
20 wherein the guide strand is from 16-29 nucleotides long and is substantially complementary to a target gene, wherein the passenger strand is from 8-14 nucleotides long and has complementarity to the guide strand, and wherein the guide stand has a single stranded 3' region that is 5 nucleotides or longer.

32. The composition of any one of claims 1-28, wherein the polynucleotide is an  
25 isolated double stranded nucleic acid molecule comprising a guide strand and a passenger strand, wherein the guide strand is from 16-25 nucleotides long and is substantially complementary to a target gene, wherein the passenger strand is from 16-25 nucleotides long and has complementarity to the guide strand, and wherein the double stranded nucleic acid molecule optionally has 3' overhangs.

30 33. The composition of any one of claims 1-28, wherein the polynucleotide is an isolated double stranded nucleic acid molecule comprising a guide strand and a passenger strand, wherein the guide strand is from 24-29 nucleotides long and is substantially complementary to a

target gene, wherein the passenger strand is from 24-29 nucleotides long and has complementarity to the guide strand, and wherein the double stranded nucleic acid molecule optionally has 3' overhangs.

34. The composition of any one of claims 1 or 2, wherein the polynucleotide is an 5 isolated single stranded nucleic acid molecule of 16-29 nucleotides in length and is substantially complementary to a target gene.

35. The composition of claim 1, wherein the polynucleotide is selected from the group consisting of an antisense ODN, an antagonir, an antiMirs, and a PNA.

36. The composition of any one of claims 1-35, wherein the ratio of neutral fatty 10 mixture to polynucleotide is 1:5.

37. The composition of any one of claims 1-35, wherein the composition is composed of about 50% cargo molecule.

38. The composition of any one of claims 1-37, wherein the composition has a particle size of 10-140nm.

39. The composition of any one of claims 1-38, wherein the composition is free of 15 cationic lipids.

40. A composition comprising:

a hydrophobic modified polynucleotide; and  
a neutral fatty mixture;

20 wherein the hydrophobic modified polynucleotide and the neutral fatty mixture form a micelle and wherein between 40-100% of the nucleotides of the polynucleotide are chemically modified nucleotides.

41. An isolated single stranded polynucleotide comprising:

at least one phosphorothioate bond and one base modified with 4-pyridyl, 2-pyridyl.

25 42. The isolated single stranded polynucleotide of claim 41, wherein all ribose moieties of the polynucleotide are modified.

43. The isolated single stranded polynucleotide of claim 41, wherein at least 40% of nucleotides of the polynucleotide are chemically modified.

44. An isolated hydrophobic modified polynucleotide comprising:

30 a sterol-type molecule attached to a polynucleotide, wherein the sterol-type molecule has a polycarbon chain length of 3 - 7 or 9-18 carbons.

45. The isolated hydrophobic modified polynucleotide of claim 44, wherein the sterol-type molecule has a polycarbon chain length of 7.

46. A method for delivering an polynucleotide to a cell, comprising:

contacting a cell with a composition of any one of claims 1-37 or an polynucleotide of

5 any one of claims 38-45 to deliver the polynucleotide to the cell.

47. The method of claim 46, wherein the method is performed *in vitro*.

48. The method of claim 46, wherein the method is performed *in vivo*.

49. A method of inducing RNAi in a cell comprising:

contacting a cell with a composition of any one of claims 1-37 or an isolated

10 polynucleotide of any one of claims 38-45, wherein the polynucleotide has at least a region of sequence correspondence to a target gene and wherein the polynucleotide induces RNAi of mRNA of the target gene.

50. A method of inducing RNAi in a subject comprising:

administering to a subject an effective amount for inducing RNAi of mRNA of a target 15 gene, a composition of any one of claims 1-37 or an isolated polynucleotide of any one of claims 38-45, wherein the polynucleotide has at least a region of sequence correspondence to the target gene.

51. The method of claim 50, wherein the subject is a human.

52. The method of claim 50, wherein the target gene is PP1B.

20 53. The method of claim 50, wherein the target gene is MAP4K4.

54. The method of claim 50, wherein the target gene is SOD1.

55. A method, in which alteration of lipid content and ratio of the composition of claim 1 is used to alter pharmacokinetic behavior and tissue distribution of polynucleotides.

56. A method for delivering a polynucleotide to a target tissue of a subject, comprising

25 administering to a subject a composition of claim 1 wherein the composition includes a fatty acid in the neutral fatty mixture or cargo molecule associated with targeting to the target tissue in order to deliver the polynucleotide to the target tissue.

57. The method of claim 56, wherein the target tissue is cardiomyocytes.

58. The method of claim 57, wherein the lipid is cardiolipin or linoleic acid.

30 59. The method of claim 56, wherein the target tissue is lung.

60. The method of claim 56, wherein the target tissue is fat.

61. The method of claim 56, wherein the target tissue is liver.

62. A method, in which alteration of lipid content and ratio of the composition of claim 1 is used to optimize pharmacokinetic behavior and tissue distribution of polynucleotides to cardiomyocytes.

5 63. A method, in which alteration of lipid content and ratio of the composition of claim 1 is used to alter pharmacokinetic behavior and tissue distribution of polynucleotides to fat tissue.

64. A method, in which alteration of lipid content and ratio of the composition of claim 1 is used to alter pharmacokinetic behavior and tissue distribution of polynucleotides to lung tissue.

10 65. A method, in which alteration of lipid content and ratio of the composition of claim 1 is used to alter pharmacokinetic behavior and tissue distribution of polynucleotides to liver.

66. A method of inducing RNAi in a subject comprising:

15 administering to a subject an effective amount for inducing RNAi of mRNA of a target gene, a composition of any one of claims 1-37 or an isolated polynucleotide of any one of claims 38-45, wherein the polynucleotide has at least a region of sequence correspondence to the target gene, wherein the step of administering is systemic, intravenous, intraperitoneal, intradermal, topical, intranasal, inhalation, oral, intramucosal or intraocular.

67. The method of claim 66, wherein the administration is systemic.


68. The method of claim 66, wherein the administration is intravenous.


69. The method of claim 66, wherein the administration is intraperitoneal.


20 70. The method of claim 66, wherein the administration is intradermal.

71. The method of claim 66, wherein the administration is topical.


72. The method of claim 66, wherein the administration is intraocular.

**Figure 1**

**Figure 2**

**Figure 3**

## Figure 4



5/38




Figure 5



**Figure 6**

Figure 7



8/38









**Figure 8**


Figure 9



**Figure 10**





12/38



Intralipid type composition





Preferential delivery to liver

**Figure 12**

Linoleic acid and cardiolipin enriched micelles



Preferential delivery to heart

**Figure 13**

14/38

rxRNA<sup>ori</sup> 25mer with cholesterol based on Map4k4 lead (11546)


- SS: 5' -P.mC.mU.mU.mU.mG.mA.mA.mG.mA.mG.mU.mU.C.U.G.mU.mG.mG.mA.mA.mG.mU.mC.mU.mA-Chol-3'
- AS: 5' -U.A.G.A.C.U.U.C.C.A.C.A.G.A.A.C.U.C.U.U.C.A.A.A.G-3' AND 5'-U.A.G.A.C.U.U.C.C.A.C.A.G.A.A.C.U.C.U\*U\*C\*A\*A\*G-3'

Antagomir-122 (Stoffel paper)

- 5' -mA\*mC\*mA.mA.mA.mC.mA.mC.mC.mA.mU.mU.mG.mU.mC.mA.mC.mA.mC.mU\*mC\*mC\*mA\*-Chol-3'


sd-rxRNA with lipid tail in place of cholesterol

- SS based off 12474 with lipid tail in place of cholesterol  
5'-mC.mU.G.mU.G.G.A.A.G.mU.mC.mU.A.-Fatty Acid-3'
- Duplex with AS 12755)



**Figure 14**

15/38









Figure 16

**Figure 17**

18/38

**Figure 18**

19/38

**Figure 19**

20/38





Figure 20



**Figure 21**






Figure 22

**Figure 23**

24/38

A



1:10 RNA:Lipid,  
50:0:50 DOPC:DOPE:Chol

B





Figure 24



**Figure 26**

26/38


RNA:Lipid	DOPC:Chol	Peak 1 Size (nm)	Peak 2 Size (nm)	Zeta Potential (mV)
1:5	50:50	401 ± 36	95 ± 7	-15 ± 5
1:10	25:75	242 ± 14	64 ± 3	-25 ± 2
1:10	50:50	66 ± 10	- -	-12 ± 3
1:10	75:25	71 ± 10	- -	-9 ± 3
1:20	50:50	121 ± 5	- -	-8 ± 1

27/38

Figure 27

# WITH hydrophobic- RNA


# No hydrophobic- RNA




**Figure 28**

28/38


Material	Supplier	Natural or Synthetic	Category	Ordered	Amount
Intralipid, 20%	Sigma			6/15/2009	100 mL
DOPC (1,2-Dioleoyl-sn-Glycero-3-phosphocholine)	Avanti	Synthetic	Phospholipid	6/18/2009	1 g
DOPC (1,2-Dioleoyl-sn-Glycero-3-phosphoethanolamine)	Avanti	Synthetic	Phospholipid	6/18/2009	1g
Cholesterol	Avanti	Natural	Sterol	6/18/2009	1g
Egg PC (L-alpha-phosphatidylcholine)	Avanti	Natural	Phospholipid	6/29/2009	25 mg
Egg PE (L-alpha-phosphatidylethanolamine)	Avanti	Natural	Phospholipid	6/29/2009	25 mg
Egg PA (L-alpha-phosphatidic acid (sodium salt))	Avanti	Natural	Phospholipid	6/29/2009	25 mg
DLPC (1,2-dilinoleoyl-sn-glycero-3-phosphocholine)	Avanti	Synthetic	Phospholipid	6/29/2009	25 mg
DSPC (1,2-distearoyl-sn-glycero-3-phosphocholine)	Avanti	Synthetic	Phospholipid	6/29/2009	25 mg
DOPA (1,2-dioleoyl-sn-glycero-3-phosphate (sodium salt))	Avanti	Synthetic	Shingolipid	6/29/2009	25 mg
Egg SM (Sphingomyelin)	Avanti	Natural	Shingolipid	6/29/2009	25 mg
Egg Sphingosine	Avanti	Natural	Shingolipid	6/29/2009	5 mg
16:0 DG (1,2-dipalmitoyl-sn-glycerol)	Avanti	Synthetic	Neutral Lipid	6/29/2009	25 mg
18:1 DG (1,2-dioleoyl-sn-glycerol)	Avanti	Synthetic	Neutral Lipid	6/29/2009	25 mg
Retinol (Vitamin A)	Sigma	Synthetic	Vitamin	6/29/2009	25 mg
Tocopherol (Vitamin E)	Sigma	Synthetic	Vitamin	6/29/2009	5 g
Cholecalciferol (Vitamin D)	Sigma	Synthetic	Vitamin	6/29/2009	1 g
Cardiolipin sodium salt from bovine heart	Sigma	Natural	Phospholipid	6/29/2009	10 mg
1a,25-Dihydroxyvitamin D3 (Calcitriol)	Sigma	Synthetic	Vitamin	6/29/2009	0.1 mg
16:0 Ethylene Glycol (1,2-dipalmitoyl ethylene glycol)	Avanti	Synthetic	Neutral Lipid	7/1/2009	10 mg
18:1 Ethylene Glycol (1,2-dioleoyl ethylene glycol)	Avanti	Synthetic	Neutral Lipid	7/1/2009	10 mg
Oleic Acid	Sigma			7/1/2009	1 g
Stearic Acid	Sigma			7/1/2009	1 g
Cholestryloleate	Sigma			7/1/2009	100 mg
Cholestrylstearate	Sigma			7/1/2009	5 g
Glyceryl trioleate	Sigma			7/1/2009	1 g
Glyceryl tristearate	Sigma			7/1/2009	5 g
Soybean oil	Sigma			7/1/2009	1 L

**Figure 29**


30/38




Cardiolipin (bisphosphatidyl glycerol)




carotenoids



Vitamine E



Vitamines D



calcitriol

Figure 30





Figure 32



**Figure 33**





Figure 34



35/38




Figure 35






Figure 36


**Figure 37**

37/38



**Figure 38**

38/38

**sd-rxRNA (13766, Active):Lipid, DOPC or DSPC:DOPE:Cholesterol****sd -rxRNA (12984, Active):Lipid, DOPC or DSPC:DOPE:Cholesterol, Sonicated**