

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 2011223627 B2

(54) Title
Method for producing soluble recombinant interferon protein without denaturing

(51) International Patent Classification(s)
C07K 14/555 (2006.01) **C12N 15/63** (2006.01)
C07K 1/02 (2006.01) **G01N 33/68** (2006.01)
C12N 15/20 (2006.01)

(21) Application No: **2011223627** (22) Date of Filing: **2011.03.02**

(87) WIPO No: **WO11/109556**

(30) Priority Data

(31) Number (32) Date (33) Country
61/310,671 **2010.03.04** **US**

(43) Publication Date: **2011.09.09**
(44) Accepted Journal Date: **2015.06.18**

(71) Applicant(s)
PFENEX INC.

(72) Inventor(s)
Allen, Jeffrey;Feng, Ping-Hua;Patkar, Anant;Haney, Keith L.;Chew, Lawrence;Sengchanthalangsy, Lei Lei Phokham

(74) Agent / Attorney
Griffith Hack, GPO Box 1285, Melbourne, VIC, 3001

(56) Related Art
Russell-Harde, D. et al., Journal of Interferon and Cytokine Research. 1995, vol. 15, pages 31-37.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
9 September 2011 (09.09.2011)(10) International Publication Number
WO 2011/109556 A3

(51) International Patent Classification:

C07K 14/555 (2006.01) *C07K 1/02* (2006.01)
C12N 15/20 (2006.01) *G01N 33/68* (2006.01)
C12N 15/63 (2006.01)

(21) International Application Number:

PCT/US2011/026921

(22) International Filing Date:

2 March 2011 (02.03.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/310,671 4 March 2010 (04.03.2010) US

(71) Applicant (for all designated States except US):
PFENEX INC. [US/US]; 10790 Roselle Street, San Diego, CA 92121 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): ALLEN, Jeffrey [US/US]; 14702 Fairtree Terrace, Poway, CA 92064 (US). FENG, Ping-Hua [CN/US]; 12039 Briarleaf Way, San Diego, CA 92128 (US). PATKAR, Anant [US/US]; 10546 Clasico Court, San Diego, CA 92127 (US). HANEY, Keith, L. [US/US]; 13254 Gabilan Road, San Diego, CA 92128 (US). CHEW, Lawrence [MY/US];

5590 Willowmere Lane, San Diego, CA 92130 (US). SENGCHANTHALANGSY, Lei Lei Phokham [US/US]; 3975 Plateau Place, Carlsbad, CA 92010 (US).

(74) Agents: CHANG, Ardith et al.; Wilson Sonsini Goodrich & Rosati, 650 Page Mill Road, Palo Alto, CA 94304-1050 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CI, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,

[Continued on next page]

(54) Title: METHOD FOR PRODUCING SOLUBLE RECOMBINANT INTERFERON PROTEIN WITHOUT DENATURING

FIGURE 1



(57) Abstract: The present invention relates to the field of recombinant protein production in bacterial hosts. It further relates to extraction of soluble, active recombinant protein from an insoluble fraction without the use of denaturation and without the need for a refolding step. In particular, the present invention relates to a production process for obtaining high levels a soluble recombinant Type 1 interferon protein from a bacterial host.



SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, — *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))* — *with sequence listing part of description (Rule 5.2(a))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

(88) Date of publication of the international search report: 12 January 2012

Published:

- *with international search report (Art. 21(3))*

**METHOD FOR PRODUCING SOLUBLE RECOMBINANT INTERFERON PROTEIN
WITHOUT DENATURING**

CLAIM OF PRIORITY

[0001] This application claims priority under 35 U.S.C. §119(e) to U.S. application Serial No. 5 61/310,671 filed on March 4, 2010. The contents of U.S. application Serial No. 61/310,671 are hereby incorporated by reference in their entirety.

SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII 10 copy, created on February 11, 2011, is named 38194601 (2).txt and is 9,237 bytes in size.

BACKGROUND OF THE INVENTION

[0003] Many heterologous recombinant proteins are produced in a misfolded insoluble form, called 15 inclusion bodies, when expressed in bacterial systems. In general, denaturing reagents must be used to solubilize the recombinant protein in the inclusion bodies. The protein must then be renatured, under conditions that have been optimized for the protein to properly fold. Efforts expended on optimization, as well as the slow refolding process and lowered process yields, add cost and time to the production of a recombinant protein.

[0004] Interferons exhibit antiviral, antiproliferative, immunomodulatory, and other activities. 20 Several distinct types of human interferons, including α , β , and γ , have been distinguished based on, e.g., their anti-viral and anti-proliferative activities. Interferon secretion is induced by signals, including viruses, double-stranded RNAs, other polynucleotides, antigens, and mitogens. Interferon- β is an example of a protein that has been expressed in recombinant form in bacteria, where it is sequestered in inclusion bodies.

[0005] Human interferon- β 1b is a regulatory polypeptide having a molecular weight of about 22 25 kDa and consisting of 165 amino acid residues. It can be produced by many cells in the body, in particular fibroblasts, in response to viral infection or exposure to other biologics. It binds to a multimeric cell surface receptor. Productive receptor binding results in a cascade of intracellular events leading to the expression of interferon- β inducible genes and triggering antiviral, antiproliferative and immunomodulatory activity.

30 [0006] Interferon- β 1b, specifically, Betaseron (h-IFN- β 1b C17S), has been used to treat diseases including multiple sclerosis (MS), hepatitis B and C infections, glioma, and melanoma. Interferon- β has been demonstrated to reduce the number of attacks suffered by patients with relapsing and remitting MS. Substantial amounts of interferon- β 1b are needed for therapeutic use. Recombinant interferon- β 1b has been produced at low levels in

5 mammalian cells, including human fibroblasts and CHO cells. Animal cell expression is typically hindered by technical difficulties including longer process time, easy contamination of cultures, a requirement for maintaining stringent culturing conditions, and the high cost of culture media. As the glycoprotein component has been found to be generally unnecessary for the activity of interferon β , research has turned to the expression of the recombinant protein in the bacterial expression system, *E. coli*. As noted, the inclusion bodies generated in *E. coli* must be solubilized by denaturation, and the interferon- β refolded. Refolding, which is slow, extends process time, adds cost, and lowers yield. To date, a method for quickly and economically producing high levels of soluble recombinant interferon- β in either mammalian or bacterial host cells, without the need for denaturing and refolding steps, has not been described.

10

SUMMARY OF THE INVENTION

15 [0007] The present invention relates to the expression of interferon in *P. fluorescens* and development of a new method to extract active proteins from the fermentation product using mild detergents and without the need for a refolding process.

20 [0008] In particular, in a first aspect, the present invention provides a method for producing a recombinant Type 1 interferon protein, said method comprising expressing the recombinant interferon protein by culturing a *Pseudomonas* or *E. coli* host cell containing an expression construct comprising a nucleic acid comprising a coding sequence that has been optimized for expression in the host cell, lysing the cultured host cell, obtaining an insoluble fraction and a soluble fraction from the lysed host cell, extracting the insoluble fraction by subjecting it to non-denaturing extraction conditions, wherein said non-denaturing extraction conditions comprise a Zwitterionic detergent or octylglucopyranoside, and obtaining an extract pellet and an extract supernatant from the insoluble fraction, wherein the recombinant protein in the extract supernatant is present in soluble form, active form, or a combination thereof, without being further subjected to a renaturing or refolding step.

25 [0008a] A second aspect of the invention provides a recombinant Type 1 interferon protein produced by the method of the first aspect.

30 [0009] In embodiments, the non-denaturing extraction conditions comprise the presence of a mild detergent. In certain embodiments, the mild detergent is a Zwitterionic detergent. In specific embodiments, the Zwitterionic detergent is Zwittergent 3-08, Zwittergent 3-10, Zwittergent 3-12, or Zwittergent 3-14. In embodiments, the non-denaturing extraction conditions comprise about 0.5% to about 2% Zwittergent 3-14. In certain embodiments, the mild detergent is not N-lauroyl-sarcosine (NLS).

5

10

15

20

25

30

35

[0010] In embodiments, the non-denaturing extraction conditions comprise the presence of a mild detergent and further comprise a chaotropic agent and a cosmotropic salt. In certain embodiments, the chaotropic agent is urea or guanidinium hydrochloride, and the cosmotropic salt is NaCl, KCl, or (NH₄)₂SO₄. In specific embodiments, the non-denaturing extraction conditions comprise about 0.5 to about 2% Zwittergent 3-14; about 0 to about 2 M urea; about 0 to about 2 M NaCl; and the pH is about 6.5 to about 8.5. In embodiments, the non-denaturing extraction conditions comprise: 1% Zwittergent 3-14; 2 M urea; 2 M NaCl; and the pH is about 8.2. In other embodiments, the non-denaturing extraction conditions additionally comprise about 1% to about 40% w/v solids. In certain embodiments, the non-denaturing extraction conditions additionally comprise about 5% w/v solids.

[0011] In embodiments, the recombinant Type 1 interferon protein is an interferon- β , an interferon- α , an interferon- κ , an interferon- τ , or an interferon- ω . In specific embodiments, the recombinant Type 1 interferon protein is an interferon- β or an interferon- α . In embodiments, the recombinant Type 1 interferon protein is an interferon- β , and said interferon- β is selected from the group consisting of: a human interferon- β 1b and human interferon- β 1b C17S. In embodiments, wherein the recombinant Type 1 interferon is an interferon- α , the interferon- α is selected from the group consisting of: human interferon- α 2a and human interferon- α 2b.

[0012] In further embodiments the method further comprises measuring the amount of recombinant Type 1 interferon protein in the insoluble fraction and the extract supernatant fractions, wherein the amount of recombinant interferon protein detected in the extract supernatant fraction is about 10% to about 95% of the amount of the recombinant interferon protein detected in the insoluble fraction. In other embodiments, the method further comprises measuring the activity of the recombinant protein, wherein about 40% to about 100% of the recombinant protein present in the extract supernatant is determined to be active. In related embodiments, the recombinant protein is an interferon- β , and the amount of active recombinant protein is determined by Blue Sepharose affinity column chromatography, receptor binding assay, antiviral activity assay, or cytopathic effect assay. In other embodiments, the recombinant protein is an interferon- α , an interferon- κ , or an interferon- ω , and the amount of active recombinant protein is determined by Blue Sepharose affinity column chromatography, receptor binding assay, antiviral activity assay, or cytopathic effect assay.

[0013] Also disclosed are methods for producing a recombinant Type 1 interferon protein, said method comprising expressing the recombinant interferon protein by culturing a *Pseudomonas* or *E. coli* host cell containing an expression construct comprising a coding sequence that has been optimized for expression in the host cell, lysing the host cell,

5 obtaining an insoluble fraction and a soluble fraction from the lysis step, extracting the insoluble fraction by subjecting it to non-denaturing extraction conditions, and obtaining an extract pellet and an extract supernatant from the insoluble fraction, wherein the recombinant protein in the extract supernatant is present in soluble form, active form, or a combination thereof, without being further subjected to a renaturing or refolding step, wherein the recombinant protein is an interferon- β , and further wherein the non-denaturing extraction conditions are optimized using the information in Figure 4B.

10 [0014] In embodiments, the recombinant protein in the extract supernatant is present at a concentration of about 0.3 grams per liter to about 10 grams per liter. In other embodiments, the host cell is cultured in a volume of about 1 to about 20 or more liters. In specific 15 embodiments, the host cell is cultured in a volume of about 1 liter, about 2 liters, about 3 liters, about 4 liters, about 5 liters, about 10 liters, about 15 liters, or about 20 liters.

[0015] In embodiments, the expression construct comprises an inducible promoter. In specific 20 embodiments, the expression construct comprises a lac promoter derivative and expression of the interferon is induced by IPTG.

[0016] In embodiments, the host cell is grown at a temperature of about 25° C to about 33° C, at a 25 pH of about 5.7 to about 6.5, and the IPTG is added to a final concentration of about 0.08 mM to about 0.4 mM, when the OD575 has reached about 80 to about 160. In specific embodiments, the host cell is grown at a temperature of about 32° C, at a pH of about 5.7 to 30 6.25, and the IPTG is added to a final concentration of about 0.2 mM, when the OD575 has reached about 120 to about 160.

[0017] In embodiments, the expression construct comprises a high activity ribosome binding site. In certain embodiments, the host cell is a lon hslUV protease deletion strain. In other 35 embodiments, the Type 1 interferon is expressed in the cytoplasm of the host cell. In related embodiments, the Type 1 interferon is human interferon- β 1b or human interferon- β 1b C17S, and is expressed in the cytoplasm of the host cell.

[0018] Also disclosed are methods for extracting a recombinant Type 1 interferon protein, wherein 40 the recombinant interferon protein is present in an insoluble fraction, said insoluble fraction produced after lysis of a *Pseudomonas* or *E. coli* host cell expressing the recombinant interferon protein, said method comprising subjecting the insoluble fraction to non-denaturing extraction conditions, and obtaining an extract pellet from the insoluble fraction, said extract 45 pellet comprising recombinant interferon protein, wherein the recombinant interferon protein in the extract pellet is in soluble form, active form, or a combination thereof, without being subjected to a renaturing or refolding step.

5 [0019] In embodiments, the recombinant Type 1 interferon protein extracted is an interferon- β , an interferon- α , an interferon- κ , an interferon- τ , or an interferon- ω . In certain embodiments, the recombinant Type 1 interferon protein is an interferon- β or an interferon- α . In embodiments, the recombinant Type 1 interferon protein is an interferon- β , and said interferon- β is selected from the group consisting of: a human interferon- β 1b and human interferon- β 1b C17S. In other embodiments, the interferon- α is selected from the group consisting of: human interferon- α 2a and human interferon- α 2b.

10 [0020] Also disclosed is a method for producing an insoluble fraction comprising a recombinant Type 1 interferon protein, wherein the recombinant interferon protein is expressed in a *Pseudomonas* or *E. coli* host cell from a nucleic acid construct comprising a nucleic acid sequence that is operably linked to a lac derivative promoter, said method comprising growing the host cell at a temperature of about 25° C to about 33° C and at a pH of about 5.7 to about 6.5, to an OD600 of about 80 to about 160, and inducing the host cell at a concentration of about 0.08 mM to about 0.4 mM IPTG, lysing the host cell and centrifuging it to produce the pellet fraction, wherein soluble, active, or soluble and active recombinant interferon protein can be obtained by extracting the pellet fraction under non-denaturing conditions without a subsequent renaturing or refolding step.

15 [0021] In embodiments, the recombinant Type 1 interferon protein comprised by the insoluble fraction is an interferon- β , an interferon- α , an interferon- κ , or an interferon- ω . In specific embodiments, the recombinant Type 1 interferon protein is an interferon- β or an interferon- α . In embodiments, wherein recombinant Type 1 interferon protein is an interferon- β , the interferon- β is selected from the group consisting of: a human interferon- β 1b and human interferon- β 1b C17S. In embodiments, wherein the Type 1 interferon protein is an interferon- α , the interferon- α is selected from the group consisting of: human interferon- α 2a and human interferon- α 2b. In embodiments, in the method for producing an insoluble fraction comprising a recombinant Type 1 interferon protein, the temperature at which the host cell is grown is about 32° C, and the IPTG concentration is about 0.2 mM.

INCORPORATION BY REFERENCE

20 [0022] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

25 [0022a] It is to be understood that, if any prior art publication is referred to herein, such reference does not constitute an admission that the publication forms a part of the common general knowledge in the art, in Australia or any other country.

BRIEF DESCRIPTION OF THE DRAWINGS

[0023] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and in the accompanying drawings.

5

[0024] **Figure 1. Initial CGE evaluation of IFN- β recovered from *P. fluorescens* strain PS530-001.** **A.** Protein analyzed under reducing conditions. **B.** Protein analyzed under non-reducing conditions.

[0025] For both A and B:

[0026] Lane 1. Molecular weight ladder with sizes as indicated.

[0027] Lane 2. Pellet from initial centrifugation after cell lysis (insoluble fraction).

[0028] Lanes 3-9. Supernatant from centrifugation following extraction step. Lanes 3 to 7 represent extraction with PBS buffer, without and with 1% Zwittergent 3-14, as indicated, and Lanes 8 and 9 represent extraction with acetate buffer, without and with 1% Zwittergent 3-14, respectively.

[0029] Lanes 10-13. Pellet from spin following extraction step. Lanes 10 and 11 represent extraction with PBS buffer, without and with 1% Zwittergent 3-14, respectively, and Lanes 12 and 13 represent extraction with acetate buffer, without and with 1% Zwittergent 3-14, respectively.

[0030] **Figure 2. Flowchart of study performed to evaluate extraction of interferon- β using different detergents.**

[0031] **Figure 3. Flowchart of statistically designed study performed to evaluate extraction of interferon- β using different extraction conditions including Zwittergent 3-14.**

[0032] **Figure 4. Results of study performed to evaluate extraction of interferon- β using different extraction conditions including Zwittergent 3-14.**

[0033] A. Statistical summary. B. Ranges of useful extraction conditions.

[0034] **Figure 5. Insoluble IFN- β Production over Post-Induction Time for Replicate Fermentations.**

[0035] The results from three different replicates were plotted.

[0036] **Figure 6. Insoluble IFN- β Production over Post-Induction Time for Alternate pH and OD.**

[0037] The results from three different replicates were plotted.

[0038] **Figure 7. IFN- β 1b Sequence.** A. IFN- β 1b C17S Amino Acid Sequence. (SEQ ID NO: 1) The sequence shows the N-terminal methionine, which is not present in the purified protein. B. IFN- β DNA Sequence with Codons Optimized for *P. fluorescens*. This sequence encodes the amino acid sequence shown in Figure 7A. (SEQ ID NO: 2) C. IFN- β 1b C17S Amino Acid Sequence, without N-terminal methionine. (SEQ ID NO: 3)

[0039] **Figure 8. IFN- α 2a Sequence.** A. IFN- α 2a Amino Acid Sequence. (SEQ ID NO: 4) B. IFN- α 2a DNA Sequence with Codons Optimized for *P. fluorescens*. (SEQ ID NO: 5)

[0040] **Figure 9. IFN- α 2b Sequence.** A. IFN- α 2b Amino Acid Sequence. (SEQ ID NO: 6) B. IFN- α 2b DNA Sequence with Codons Optimized for *P. fluorescens*. (SEQ ID NO: 7)

DETAILED DESCRIPTION OF THE INVENTION

[0041] The present invention relates to a method for producing large amounts of soluble recombinant interferon protein in a *Pseudomonas* expression system. In particular, this

method eliminates the need for the denaturing step and subsequent renaturing/refolding step typically required.

5 [0042] Production of recombinant interferon- β in bacterial expression systems has been hampered by sequestration of the protein in insoluble inclusion bodies. Solubilization of the inclusion bodies requires denaturation, which in turn necessitates the use of a refolding step that is costly, time-consuming, and decreases protein yield. The present invention circumvents the need for a refolding step by providing methods for producing and solubilizing interferon without recourse to denaturation.

10 [0043] Methods for producing a recombinant interferon protein that is soluble, active, or both, in a bacterial expression system, without subjecting the protein to a denaturing step, are provided. In particular, a non-denaturing extraction process that results in soluble interferon protein is described. Interferons expressed in bacterial expression systems are generally localized to an insoluble fraction. In the extraction process of the present invention, this insoluble fraction is subjected to extraction conditions that include non-denaturing 15 concentrations of mild detergents and produce soluble protein.

20 [0044] Also provided by the present invention are methods for producing a recombinant interferon protein wherein growth conditions for the *Pseudomonas* host cell are optimized to maximize yields of the soluble recombinant interferon protein, particularly when the extraction method of the invention is used. Studies of the effect of *E. coli* growth conditions on soluble protein production have been reported. The solubility of a given protein when expressed in *Pseudomonas* can be different from that in *E. coli*. This is illustrated in, e.g., U.S. Pat. App. Pub. No. 2006/0040352, “Expression of Mammalian Proteins in *Pseudomonas Fluorescens*,” which shows side-by-side comparisons of the soluble amounts of several 25 proteins produced using *E. coli* or *P. fluorescens* as the host. Furthermore, there is substantial variation among the solubilities of different proteins even in the same host, as solubility is influenced strongly by protein structure, e.g., amino acid sequence. Previously reported attempts at producing IFN- β in *E. coli* resulted in protein that required refolding. See, e.g., Russell-Harde, 1995, “The Use of Zwittergent 3-14 in the Purification of Recombinant Human Interferon- β Ser17 (Betaseron) et al., J. Interferon and Cytokine Res. 15:31-37, and Ghane, et al., 2008, “Over Expression of Biologically Active Interferon Beta Using Synthetic Gene in *E. coli*,” J. of Sciences, Islamic Republic of Iran 19(3):203-209, both incorporated herein by reference.

30 [0045] The methods further provide optimized growth conditions including growth temperature, OD at time of promoter induction, inducer concentration, and pH. Extraction conditions provided include detergent type and concentration, chaotropic agent, cosmotropic salt, and

pH. Specific values as well as optimal parameter ranges are provided. Also provided are methods for optimizing extraction conditions using the provided ranges.

Bacterial Growth Conditions

[0046] In embodiments of the present invention, the bacterial growth conditions are optimized to increase the amount of soluble interferon protein obtained using the extraction methods as provided herein. Use of the growth conditions of the present invention with other extraction conditions, e.g., other methods described and used in the art, is also contemplated.

[0047] Optimal growth conditions particularly useful in conjunction with the extraction methods of the invention comprise: a temperature of about 25° C to about 33° C; a pH of about 5.7 to about 6.5, and induction with about 0.08 mM to about 0.4 mM IPTG when the culture has reached an OD₅₇₅ of about 80 to about 160.

[0048] The pH of the culture can be maintained using pH buffers and methods known to those of skill in the art. Control of pH during culturing also can be achieved using aqueous ammonia. In embodiments, the pH of the culture is about 5.7 to about 6.5. In certain embodiments, the pH is 5.7, 5.8, 5.9, 6.0, 6.1, 6.2, 6.3, 6.4, or 6.5. In other embodiments, the pH is 5.7 to 5.9, 5.8 to 6.0, 5.9 to 6.1, 6.0 to 6.2, 6.1 to 6.3, or 6.2 to 6.5. In yet other embodiments, the pH is 5.7 to 6.0, 5.8 to 6.1, 5.9 to 6.2, 6.0 to 6.3, 6.1 to 6.4, or 6.2 to 6.5. In certain embodiments, the pH is about 5.7 to about 6.25.

[0049] In embodiments, the growth temperature is maintained at about 25° C to about 33° C. In certain embodiments, the growth temperature is about 25° C, about 26° C, about 27° C, about 28° C, about 29° C, about 30° C, about 31° C, about 32° C, or about 33° C. In other embodiments, the growth temperature is maintained at about 25° C to about 27° C, about 25° C to about 28° C, about 25° C to about 29° C, about 25° C to about 30° C, about 25° C to about 31° C, about 25° C to about 32° C, about 25° C to about 33° C, about 26° C to about 28° C, about 26° C to about 29° C, about 26° C to about 30° C, about 26° C to about 31° C, about 26° C to about 32° C, about 27° C to about 29° C, about 27° C to about 30° C, about 27° C to about 31° C, about 27° C to about 32° C, about 26° C to about 33° C, about 28° C to about 30° C, about 28° C to about 31° C, about 28° C to about 32° C, about 29° C to about 31° C, about 29° C to about 32° C, about 29° C to about 33° C, about 30° C to about 32° C, about 30° C to about 33° C, about 31° C to about 33° C, about 31° C to about 32° C, about 30° C to about 33° C, or about 32° C to about 33° C.

Induction

[0050] As described elsewhere herein, inducible promoters can be used in the expression construct to control expression of the recombinant interferon gene. In the case of the lac promoter derivatives or family members, e.g., the tac promoter, the effector compound is an inducer, such as a gratuitous inducer like IPTG (isopropyl-β-D-1-thiogalactopyranoside, also called

"isopropylthiogalactoside"). In embodiments, a lac promoter derivative is used, and interferon expression is induced by the addition of IPTG to a final concentration of about 0.08 mM to about 0.4 mM, when the cell density has reached a level identified by an OD₅₇₅ of about 80 to about 160.

5 [0051] In embodiments, the OD₅₇₅ at the time of culture induction about 80, about 90, about 100, about 110, about 120, about 130, about 140, about 150, about 160, about 170 or about 180. In other embodiments, the OD₅₇₅ is about 80 to about 100, about 100 to about 120, about 120 to about 140, about 140 to about 160. In other embodiments, the OD₅₇₅ is about 80 to about 120, about 100 to about 140, or about 120 to about 160. In other embodiments, the 10 OD₅₇₅ is about 80 to about 140, or about 100 to 160. The cell density can be measured by other methods and expressed in other units, e.g., in cells per unit volume. For example, an OD₅₇₅ of about 80 to about 160 of a *Pseudomonas fluorescens* culture is equivalent to approximately 8×10^{10} to about 1.6×10^{11} colony forming units per mL or 35 to 70 g/L dry cell weight. In embodiments, the cell density at the time of culture induction is equivalent to the cell density as specified herein by the absorbance at OD₅₇₅, regardless of the method used for determining cell density or the units of measurement. One of skill in the art will know how to make the appropriate conversion for any cell culture.

15 [0052] In embodiments, the final IPTG concentration of the culture is about 0.08 mM, about 0.1 mM, about 0.2 mM, about 0.3 mM, or about 0.4 mM. In other embodiments, the final IPTG concentration of the culture is about 0.08 mM to about 0.1 mM, about .1 mM to about 0.2 mM, about .2 mM to about 0.3 mM, about .3 mM to about 0.4 mM, about .2 mM to about 0.4 mM, or about 0.08 to about 0.2 mM.

20 [0053] In embodiments, the interferon is expressed by induction of a lac promoter or derivative using IPTG, lactose or allolactose, by methods known in the art and described in the literature, e.g., in U.S. Pat. No. 7,759,109, "High density growth of T7 expression strains with auto-induction option," incorporated herein by reference in its entirety.

25 [0054] In embodiments wherein a non-lac type promoter is used, as described herein and in the literature, other inducers or effectors can be used.

30 [0055] After induction is started, cultures are grown for a period of time, typically about 24 hours, during which time the recombinant interferon protein is expressed. Cell cultures can be concentrated by centrifugation, and the culture pellet resuspended in a buffer or solution appropriate for the subsequent lysis procedure.

35 [0056] In embodiments, cells are disrupted using equipment for high pressure mechanical cell disruption (which are available commercially, e.g., Microfluidics Microfluidizer, Constant Cell Disruptor, Niro-Soavi homogenizer or APV-Gaulin homogenizer). Any appropriate method known in the art for lysing cells can be used to release the insoluble fraction. For

example, in embodiments, chemical and/or enzymatic cell lysis reagents, such as cell-wall lytic enzyme and EDTA, can be used. Use of frozen or previously stored cultures is also contemplated in the methods of the invention. Cultures can be OD-normalized prior to lysis.

5 **[0057]** Centrifugation is performed using any appropriate equipment and method. Centrifugation of cell culture or lysate for the purposes of separating a soluble fraction from an insoluble fraction is well-known in the art. For example, lysed cells can be centrifuged at $20,800 \times g$ for 20 minutes (at $4^{\circ} C$), and the supernatants removed using manual or automated liquid handling. The pellet (insoluble) fraction is resuspended in a buffered solution, e.g., phosphate buffered saline (PBS), pH 7.4. Resuspension can be carried out using, e.g., equipment such as impellers connected to an overhead mixer, magnetic stir-bars, rocking shakers, etc.

10 **[0058]** A “soluble fraction,” i.e., the soluble supernatant obtained after centrifugation of a lysate, and an “insoluble fraction,” i.e., the pellet obtained after centrifugation of a lysate, result from lysing and centrifuging the cultures. These two fractions also can be referred to as a “first soluble fraction” and a “first insoluble fraction,” respectively.

15 **[0059]** It is possible to obtain soluble IFN- β using extraction methods according to the invention, from expression cultures prepared by growing cultures under conditions in which the pH and the induction OD are not tightly controlled (see, e.g., Example 2). Optimization of the growth conditions as described herein results in substantially increased production of soluble IFN- β .

Non-Denaturing Extraction Process

[0060] It has been discovered that high levels of soluble interferon protein can be obtained from the insoluble fraction, using non-denaturing extraction methods of the present invention.

25 **[0061]** Non-denaturing extraction conditions identified as particularly useful for producing high levels of soluble recombinant interferon protein comprise: a mild detergent at a non-denaturing concentration, e.g., Zwittergent 3-14 (0.5 to 2% w/v); a chaotropic agent, e.g., urea (0-2M), and a cosmotropic salt, e.g., NaCl (0-2M), at a buffer pH of 6.5 to 8.5 and a solids concentration of 5-20% w/v.

30 **[0062]** After obtaining the soluble fraction and insoluble fraction, as described above, the soluble recombinant interferon protein is extracted from the insoluble fraction by incubating the resuspended insoluble fraction under the non-denaturing extraction conditions described herein. After incubation, the extracted mixture is centrifuged to produce an “extract supernatant” (the soluble supernatant fraction obtained after extraction containing solubilized recombinant protein) and an “extract pellet” (the insoluble pellet fraction

35

obtained after extraction). These fractions can also be referred to as the “second soluble fraction” and the “second insoluble fraction.”

Extraction Conditions

[0063] Many different parameters for the extraction conditions were evaluated for their effect on the amount of soluble protein observed in the extract supernatant, as described in Example 3 herein. It was found that soluble interferon protein was observed when the extraction conditions comprised any of a number of different detergents, at varying concentrations, as well as when other parameters were varied. However, certain parameters had a particularly striking effect on the amount of soluble protein produced.

[0064] Specifically, extraction conditions comprising Zwitterionic detergents (Zwittergents) gave the best soluble protein yields. In particular, use of the Zwitterionic detergents, Zwittergent 3-08, Zwittergent 3-10, Zwittergent 3-12, and especially Zwittergent 3-14, resulted in the highest yields. N-Lauroylsarcosine (NLS) gave a notably high yield, however the soluble protein obtained was found to be inactive based on an affinity assay (Sepharose blue affinity column binding). Therefore, the term “mild detergents” as used herein is intended not to include N-lauroylsarcosine.

[0065] The detergents were tested at non-denaturing concentrations. It was found that a concentration of Zwittergent 3-14 (3-(N,N-dimethylmyristylammonio) propanesulfonate) of at least 0.5% (w/v), and preferably 1%, well above its critical micelle concentration (which is 0.01%) provides the most efficient extraction of soluble interferon protein.

[0066] Therefore, use of non-denaturing concentrations of mild detergents, particularly Zwitterionic detergents, more particularly Zwittergent 3-08, Zwittergent 3-10, Zwittergent 3-12, and Zwittergent 3-14, more particularly Zwittergent 3-14, and not NLS, is contemplated for use in the extraction conditions of the invention.

[0067] In other embodiments of the invention, the non-denaturing extraction conditions comprise a concentration of about 0.5% to about 2% (w/v) Zwittergent 3-14. In embodiments, the w/v concentration of Zwittergent 3-14 is about 0.5% to about 1%, about 1% to about 1.5%, about 1.5% to about 2%, or about 1% to about 2%. In certain embodiments, the w/v concentration of Zwittergent 3-14 is about 0.5%, about 0.6%, about 0.7%, about 0.8%, about 0.9%, about 1.0%, about 1.1%, about 1.2%, about 1.3%, about 1.4%, about 1.5%, about 1.6%, about 1.7%, about 1.8%, about 1.9%, or about 2.0%.

[0068] In other embodiments of the invention, non-denaturing extraction conditions comprise a concentration of about 0.5% to about 2% (w/v) Zwittergent 3-08, Zwittergent 3-10, or Zwittergent 3-12. In embodiments, the w/v concentration of Zwittergent 3-08, Zwittergent 3-10, or Zwittergent 3-12 is about 0.5% to about 1%, about 1% to about 1.5%, about 1.5% to about 2%, or about 1% to about 2%. In certain embodiments, the w/v concentration of

Zwittergent 3-08, Zwittergent 3-10, or Zwittergent 3-12 is about 0.5%, about 0.6%, about 0.7%, about 0.8%, about 0.9%, about 1.0%, about 1.1%, about 1.2%, about 1.3%, about 1.4%, about 1.5%, about 1.6%, about 1.7%, about 1.8%, about 1.9%, or about 2.0%.

[0069] A mild detergent does not cause protein unfolding at low levels, e.g., 2% or less. For 5 example, SDS and NLS are typically considered stronger detergents, as they can denature proteins at these levels. A non-denaturing concentration indicates a concentration of a reagent at which proteins are not denatured. Proteins that are not denatured during processing do not require refolding.

[0070] In embodiments, non-denaturing extraction conditions of the present invention comprise 10 about 0.5 to about 2% Zwittergent 3-14; about 0 to about 2 M urea; about 0 to about 2 M NaCl; and wherein the pH is about 6.5 to about 8.5.

[0071] The following table lists examples of common detergents, including ionic, non-ionic, and 15 zwitter-ionic detergents, and their properties. An important characteristic of a detergent is its critical micelle concentration (CMC), which relates to its protein solubilization capability as well as the ease of subsequent removal of detergents from protein solutions.

Table 1. Examples of Detergents

Detergent	Monomer, MW Da	Micelle, MW Da	CMC % (w/v)	CMC mM
Zwittergent 3-14	364	30,000	0.004-0.015 (0.011)	0.1-0.4 (0.3)
Tween-20	1228		0.007	0.059
Tween-80	1310	76,000	0.0016	0.012
Triton X-100	650	90,000	0.013-0.06 (0.021)	0.2-0.9 (0.3)
Sodium Deoxycholate	432	4,200	0.21	5
Sodium Lauroylsarcosine	293	600	0.4	13.7
NDSB	195	N/A	N/A	N/A
NP-40	617	90000	0.003-0.018,	0.05-0.3
CHAPS	615	6,000	0.37-0.62	6-10
Octyl- β -glucopyranoside	292	8,000	0.73	23

Table 2. Physical Properties of Zwitterionic Detergents

Detergent	Monomer, MW Da	Micelle, MW Da	CMC % (w/v)	CMC mM

Zwittergent 3-08	280	—	9.2	330
Zwittergent 3-10	308	12,500	0.77-1.23	25-40
Zwittergent 3-12	336	18,500	0.067-0.134	2-4
Zwittergent 3-14	364	30,000	0.004-0.015 (0.011)	0.1-0.4 (0.3)
Zwittergent 3-16	392	60,000	0.0004-0.0024	0.01-0.06

[0072] It was further observed that when the non-denaturing extraction conditions comprised the combination of a chaotropic agent, urea, a cosmotropic salt, NaCl, Zwittergent 3-14, and an appropriate buffer range, the extraction yield was increased several-fold compared to the use of Zwittergent 3-14 alone (see Example 3).

Table 3. Selected concentration ranges of extraction components

Component	Permissible Conc. Range	Selected Conc.
Zwittergent 3-		
14	0.5-2% (w/v)	1%
Urea	0-2 M	2 M
NaCl	0-2 M	2 M
Solid Conc.	5-20% (w/v)	5%
Buffer pH	6.5-8.5	8.2

[0073] Chaotropic agents disrupt the 3-dimensional structure of a protein or nucleic acid, causing denaturation. In embodiments, the non-denaturing extraction conditions comprise low, non-denaturing concentrations of chaotropic agents, e.g., urea or guanidinium hydrochloride. In embodiments, the non-denaturing extraction conditions comprise urea at a concentration of about 0.5M to about 2M or higher. We observed that 6M urea denatures IFN- β . Typically, non-denaturing concentrations of urea or guanidinium hydrochloride are below 3M. In embodiments, the non-denaturing extraction conditions comprise urea at a concentration of about 0.5 to about 1M, about 1 to about 1.5M, about 1.5 to about 2M, about 1 to about 2M, about 0.5M, about 0.6M, about 0.7M, about 0.8M, about 0.9M, about 1.0M, about 1.1M, about 1.2M, about 1.3M, about 1.4M, about 1.5M, about 1.6M, about 1.7M, about 1.8M, about 1.9M, or about 2.0M. In other embodiments, the extraction conditions comprise guanidinium hydrochloride at a concentration of 0.5 to 2M. In embodiments, extraction conditions comprise guanidinium hydrochloride at a concentration of 0.5 to 1M, 1 to 1.5M, 1.5 to 2M, 1 to 2M, 0.5M, about 0.6M, about 0.7M, about 0.8M, about 0.9M, about 1.0M,

about 1.1M, about 1.2M, about 1.3M, about 1.4M, about 1.5M, about 1.6M, about 1.7M, about 1.8M, about 1.9M, or about 2.0M.

[0074] Cosmotropic salts contribute to the stability and structure of water-water interactions. Cosmotropes cause water molecules to favorably interact, which also stabilizes intermolecular interactions in macromolecules such as proteins. Any such appropriate agent, as known in the art, can be used in the extraction conditions of the present invention. In embodiments, the non-denaturing extraction conditions comprise a cosmotropic salt selected from NaCl, KCl, and $(\text{NH}_4)_2\text{SO}_4$. In certain embodiments, NaCl is present at a concentration of about 0.15M to about 2M. In embodiments, NaCl is present in the extraction conditions at a concentration of about 0.15 to about 0.5M, about 0.5 to about 10 0.75M, about 0.75M to about 1M, about 1M to about 1.25M, about 1.25M to about 1.5M, about 1.5M to about 1.75M, about 1.75M to about 2M, about 0.15M to about 1M, about 1M to about 1.5M, about 1.5M to about 2M, about 1M to about 2M, about 0.15M, about 0.25M, about 0.5M, about 0.6M, about 0.7M, about 0.8M, about 0.9M, about 1.0M, about 1.1M, about 1.2M, about 1.3M, about 1.4M, about 1.5M, about 1.6M, about 1.7M, about 1.8M, about 1.85M, about 1.9M, or about 2.0M.

[0075] In other embodiments, KCl is present in the non-denaturing extraction conditions at a concentration of about 0.15 to about 0.5M, about 0.5 to about 0.75M, about 0.75M to about 1M, about 1M to about 1.25M, about 1.25M to about 1.5M, about 1.5M to about 1.75M, about 1.75M to about 2M, about 0.15M to about 1M, about 1M to about 1.5M, about 1.5M to about 2M, about 1M to about 2M, about 0.15M, about 0.25M, about 0.5M, about 0.6M, about 0.7M, about 0.8M, about 0.9M, about 1.0M, about 1.1M, about 1.2M, about 1.3M, about 1.4M, about 1.5M, about 1.6M, about 1.7M, about 1.8M, about 1.85M, about 1.9M, or about 2.0M.

[0076] In other embodiments, $(\text{NH}_4)_2\text{SO}_4$ is present in the non-denaturing extraction conditions at a concentration of about 0.15 to about 0.5M, about 0.5 to about 0.75M, about 0.75M to about 1M, about 1M to about 1.25M, about 1.25M to about 1.5M, about 1.5M to about 1.75M, about 1.75M to about 2M, about 0.15M to about 1M, about 1M to about 1.5M, about 1.5M to about 2M, about 1M to about 2M, about 0.15M, about 0.25M, about 0.5M, about 0.6M, about 0.7M, about 0.8M, about 0.9M, about 1.0M, about 1.1M, about 1.2M, about 1.3M, about 1.4M, about 1.5M, about 1.6M, about 1.7M, about 1.8M, about 1.85M, about 1.9M, or about 2.0M.

[0077] The extraction conditions were found to be most effective when the pH was maintained within a range of 6.5 to 8.5. Useful pH buffers are those recommended in standard protein purification texts (e.g., *Protein Purification: Principles and Practice*, by Robert Scopes (Springer, 1993) can be used here, including Tris, Bis-Tris, phosphate, citrate, acetate,

glycine, diethanolamine, 2-amino-2-methyl-1,3-propanediol, triethanolamine, imidazole, histidine, pyridine, etc. Many buffers have been described in the literature and are commercially available. In embodiments, the pH of the non-denaturing extraction conditions is about 6.5, about 6.6, about 6.7, about 6.8, about 6.9, about 7.0, about 7.1, about 5
7.2, about 7.3, about 7.4, about 7.5, about 7.6, about 7.8, about 7.9, about 8.0, about 8.1, about 8.2, about 8.3, about 8.4, or about 8.5. In other embodiments, the pH is about 6.5 to about 6.8, about 6.6 to about 6.9, about 6.7 to about 7.0, about 6.8 to about 7.1, about 6.9 to about 7.2, about 7.0 to about 7.3, about 7.1 to about 7.4, about 7.2 to about 7.5, about 7.3 to about 7.6, about 7.4 to about 7.7, about 7.5 to about 7.8, about 7.6 to about 7.9, about 7.8 to about 10
7.1, about 7.9 to about 8.2, about 8.0 to about 8.3, about 8.1 to about 8.4 or about 8.2 to about 8.5. In other embodiments, the pH is about 6.5 to about 7.0, about 7.0 to about 7.5, or about 7.5 to about 8.0.

15 [0078] The solids concentration in the non-denaturing extraction conditions was also varied. This parameter represents the amount of solid material in the extract incubation mixture. Solids concentration can be determined by weighing the wet pellet (i.e., the insoluble fraction), and comparing this weight with the total weight of the extraction mixture. Specific solids concentrations are achieved by concentrating or diluting the insoluble fraction. High extraction yields were observed across a range of solids concentrations of 5% to 40% (w/v). In embodiments of the invention, the solids in the non-denaturing extraction conditions are present at a w/v concentration of about 5%, about 7.5%, about 10%, about 12.5%, about 20
15%, about 17.5%, about 20%, about 22.5%, about 25%, about 27.5%, about 30%, about 32.5%, about 35%, about 37.5%, or about 40%. In other embodiments of the invention, the solids in the non-denaturing extraction conditions are present at a w/v concentration of about 5% to about 7.5%, about 7.5% to about 10%, about 10% to about 12.5%, about 12.5% to about 15%, about 15% to about 17.5%, about 17.5% to about 20%, about 20% to about 22.5%, about 22.5% to about 25%, about 25% to about 27.5%, about 27.5% to about 30%, about 30% to about 32.5%, about 32.5% to about 35%, about 35% to about 37.5%, about 25
25 37.5% to about 40%, about 5% to about 10%, about 10% to about 15%, about 15% to about 20%, about 20% to about 25%, about 25% to about 30%, about 35% to about 40%, about 5% to about 15%, about 5% to about 25%, about 5% to about 30%, about 5% to about 35%, about 10% to about 20%, about 20% to about 30%, about 30% to about 40%, about 5% to about 20%, or about 20% to about 40%.

30 [0079] In embodiments, the extraction methods of the invention are combined with simultaneous enrichment techniques such as adsorption to further enhance protein yield.

[0080] The solubilized protein can be isolated or purified from other protein and cellular debris by, for example, centrifugation and/or chromatography such as size exclusion, anion or cation exchange, hydrophobic interaction, or affinity chromatography.

Interferons

5 **[0081]** Human interferons have been classified into three major types. Interferon type I: Type I IFNs bind to a specific cell surface receptor complex known as the IFN- α receptor (IFNAR) that consists of IFNAR1 and IFNAR2 chains. Human type I interferons include are IFN- α , IFN- β , IFN- κ , and IFN- ω and IFN- ϵ . Interferon type II: Type II IFNs (human IFN- γ) binds to IFNGR. Interferon type III: type III interferons signal through a receptor complex consisting of IL10R2 (also called CRF2-4) and IFNLR1 (also called CRF2-12).

10 **[0082]** Human Type I interferon appears to bind to two-receptor subunits, IFNAR-1 and -2, which are widely distributed on the cell surface of various cell types. Ligand involvement leads to the induction of the phosphorylation of tyrosine kinases TYK2 and JAK-1, which are coupled to IFNAR-1 and -2 respectively. Once phosphorylated, STAT proteins are released from the receptor and form homodimers as well as heterodimers. Once released, the dimers of STAT associate with interferon Responsive Factor 9 (IRF-9), a DNA binding protein, forming a complex called IFN-stimulated gene factor-3 (ISGF-3), that migrates into the nucleus. Next, the ISGF-3 complex binds to a DNA element existing in the upstream of all IFN inducible genes. Type 1 interferons are described extensively in the literature, e.g., in U.S. Pat. No. 7,625,555, “Recombinant human interferon-like proteins, incorporated herein by reference.”

15 **[0083]** Type 1 IFNs have substantial structural similarity, as evidenced by their sequences and their shared receptor binding capacity. According to Kontsek, P., 1994, “Human type I interferons: structure and function,” Acta Virol. 38(6):345-60, incorporated by reference herein, human type I interferons (13 had been reported at the time) have a 20% overall sequence homology, which determines identical secondary and tertiary folding of polypeptides. Further, Kontsek reports that three-dimensional models suggest that the globular structure of type I IFNs consists of a bundle of 5 α -helices, which might form two polypeptide domains. Disulfide bond Cys 29-Cys 139 stabilizes both domains in a bioactive configuration. Two conservative hydrophilic regions associated with the amino acids (aa) 20-41 and 120-145 are thought to constitute the basic framework of receptor recognition site in type I IFNs, and the different spectra of biological effects among human type I IFNs are speculated to be due to subtle sequential heterogeneity in the segments aa 30-41 and 120-145, and the variable hydrophilic aa regions 23-26, 68-85 and 112-121. A later report by Oritani, et al., 2001, “Type I interferons and limitin: a comparison of structures, receptors, and functions,” Cytokine Growth Factor Rev 12(4):337-48, incorporated by reference

25

30

35

herein, describes family members IFN- α , IFN- β , IFN- π , and IFN- τ . The paper also reports that IFN- α and IFN- β have a globular structure composed of five α -helices, and discusses comparative sequence analyses, a prototypic three-dimensional structure, analysis with monoclonal antibodies, and construction of hybrid molecules and site directed mutagenesis.

5

[0084] Production of any Type 1 interferon protein using the methods of the present invention is contemplated. Type 1 interferon proteins that can be produced using the methods of the invention, include, but not limited to, human IFN- α 2a and 2b, human IFN- β 1b, human IFN- κ (e.g., NM_020124, AAK63835, and described by LaFleur, et al., 2001, "Interferon-kappa, a novel type I interferon expressed in human keratinocytes," J. Biol. Chem. 276 (43), 39765-39771, incorporated herein by reference), and human IFN- ω (e.g., NM_002177, NP_002168, and described in U.S. Pat. No. 7,470,675, "Methods for treating cancer using interferon- ω -expressing polynucleotides," incorporated by reference herein in its entirety). Production of IFN- τ using the methods of the invention is also contemplated. Amino acid and nucleic acid sequences are publicly available, e.g., from GenBank.

10

[0085] Fourteen subtypes of IFN- α proteins have been described: IFNA1, IFNA2, IFNA4, IFNA5, IFNA6, IFNA7, IFNA8, IFNA10, IFNA13, IFNA14, IFNA16, IFNA17, IFNA21. IFN- α is made synthetically as therapeutic agent, as pegylated interferon alfa-2a and pegylated interferon alfa-2b.

15

[0086] IFN- β (IFNB1, or IFN- β 1b) is the main β subtype (see, e.g., GenBank NP002167.1, which provides the mature peptide sequence). Betaseron is an analogue of human IFN- β in which serine was genetically engineered to substitute for cysteine at position 17, is known as IFN- β 1b C17S (described in U.S. Pat. No. 4,588,585, "Human recombinant cysteine depleted interferon- β mutants," incorporated herein by reference). The molecule is a small polypeptide of 165 amino acids with a single disulphide bond, and is produced as a non-glycosylated protein.

20

[0087] IFN- τ is described, and sequences of IFN- τ disclosed, e.g., in U.S. Pat. No. 7,214,367, "Orally-administered interferon-tau compositions and methods," incorporated herein by reference in its entirety.

25

[0088] A number of Type 1 IFNs have been approved by the FDA for use in treating disease in humans. The following table lists examples of Type 1 interferon drugs. In embodiments of the invention, any of these drugs are produced using the methods as claimed or described herein.

Table 4. Examples of Type 1 interferon drugs.

Generic name	Trade name
Interferon α 2a	Roferon A
PEGylated interferon α 2a	Pegasys

PEGylated interferon α 2a	Reiferon Rctard
Interferon α 2b	Intron A/Reliferon
PEGylated interferon α 2b	PegIntron
Human leukocyte Interferon- α (HuIFN- α -Le)	Multiferon
Interferon β 1a, liquid form	Rebif
Interferon β 1a	lyophilized Avonex
Interferon β 1b	Betaseron/Betaferon

[0089] In embodiments, variants and modifications of Type 1 interferon proteins are produced using the methods of the present invention. For example, IFN- β variants are described in U.S. Pat. No. 6,531,122 "Interferon- β variants and conjugates," and U.S. Pat. No. 7,625,555, both incorporated by reference herein. Conjugates include pegylated Type 1 interferons, e.g., the PEGylated agents shown in Table 4, and interferons linked to non-peptide moieties.

[0090] The methods of the invention are expected to be useful for all Type 1 interferons, due to their structural similarities. Certain structurally unrelated proteins, for example, human GCSF, have been found poor candidates for producing using the methods of the present invention. When GCSF was produced and extracted using methods as described herein, less than 5% of the amount of GCSF protein in the insoluble fraction was extracted as soluble protein (data not shown).

[0091] In general, with respect to an amino acid sequence, the term "modification" includes substitutions, insertions, elongations, deletions, and derivatizations alone or in combination. In some embodiments, the peptides may include one or more modifications of a "non-essential" amino acid residue. In this context, a "non-essential" amino acid residue is a residue that can be altered, e.g., deleted or substituted, in the novel amino acid sequence without abolishing or substantially reducing the activity (e.g., the agonist activity) of the peptide (e.g., the analog peptide). In some embodiments, the peptides may include one or more modifications of an "essential" amino acid residue. In this context, an "essential" amino acid residue is a residue that when altered, e.g., deleted or substituted, in the novel amino acid sequence the activity of the reference peptide is substantially reduced or abolished. In such embodiments where an essential amino acid residue is altered, the modified peptide may possess an activity of a Type 1 interferon of interest in the methods provided. The substitutions, insertions and deletions may be at the N-terminal or C-terminal end, or may be at internal portions of the protein. By way of example, the protein may include 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more substitutions, both in a consecutive manner or spaced throughout the peptide molecule. Alone or in combination with the substitutions, the peptide may include 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more insertions, again either in consecutive manner or spaced throughout the peptide molecule. The peptide, alone or in combination with the substitutions and/or insertions, may also include 1, 2, 3, 4, 5, 6, 7, 8, 9,

10, or more deletions, again either in consecutive manner or spaced throughout the peptide molecule. The peptide, alone or in combination with the substitutions, insertions and/or deletions, may also include 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more amino acid additions.

5 [0092] Substitutions include conservative amino acid substitutions. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain, or physicochemical characteristics (e.g., electrostatic, hydrogen bonding, isosteric, hydrophobic features). The amino acids may be naturally occurring or normatural (unnatural). Families of amino acid residues having similar side chains are known in the art. These families include amino acids with basic side chains (e.g. lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, methionine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, tryptophan), β -branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Substitutions may also include non-conservative changes.

10

15

Methods for Selecting Optimal Extraction Conditions

20 [0093] In embodiments of the present invention, the results of the statistical analysis as set forth in Figure 4B are used to further optimize extraction conditions within the ranges of parameter values provided. High level soluble protein production of all Type 1 interferons is expected to be observed when practicing the invention using any values within the ranges set forth. Nonetheless, it is within the capacity of one of skill in the art to utilize the tool represented by Figure 4B to optimize the extraction conditions to fit the need at hand.

Evaluation of product

25 Protein Yield

[0094] Protein yield in the insoluble and soluble fractions as described herein can be determined by methods known to those of skill in the art, for example, by capillary gel electrophoresis (CGE), and Western blot analysis.

30 [0095] Useful measures of protein yield include, e.g., the amount of recombinant protein per culture volume (e.g., grams or milligrams of protein/liter of culture), percent or fraction of recombinant protein measured in the insoluble pellet obtained after lysis (e.g., amount of recombinant protein in extract supernatant/amount of protein in insoluble fraction), percent or fraction of active protein (e.g., amount of active protein/amount protein used in the assay), percent or fraction of total cell protein (tcp), amount of protein/cell, and percent dry biomass. In embodiments, the measure of protein yield as described herein is based on the amount of soluble protein or the amount of active protein, or both, obtained.

35

[0096] In embodiments, the methods of the present invention can be used to obtain an extracted recombinant protein yield of about 0.3 grams per liter to about 10 grams per liter. In certain embodiments, the extracted recombinant protein yield is about 0.3 grams per liter to about 1 gram per liter, about 1 gram per liter to about 2 grams per liter, about 2 grams per liter to about 3 grams per liter, about 3 grams per liter to about 4 grams per liter, about 4 grams per liter to about 5 grams per liter, about 5 grams per liter to about 6 grams per liter, about 6 grams per liter to about 7 grams per liter, about 7 grams per liter to about 8 grams per liter, about 8 grams per liter to about 9 grams per liter, or about 9 grams per liter to about 10 grams per liter. In embodiments, the extracted protein yield is about 1 gram per liter to about 3 grams per liter, about 2 grams per liter to about 4 grams per liter, about 3 grams per liter to about 5 grams per liter, about 4 grams per liter to about 6 grams per liter, about 5 grams per liter to about 7 grams per liter, about 6 grams per liter to about 8 grams per liter, about 7 grams per liter to about 9 grams per liter, or about 8 grams per liter to about 10 grams per liter. In embodiments, the extracted protein yield is about 0.5 grams per liter to about 4 grams per liter, 1 gram per liter to about 5 grams per liter, 2 grams per liter to about 6 grams per liter, about 3 grams per liter to about 7 grams per liter, about 4 grams per liter to about 8 grams per liter, about 5 grams per liter to about 9 grams per liter, or about 6 grams per liter to about 10 grams per liter. In embodiments, the extracted protein yield is about .5 gram per liter to about 5 grams per liter, about 1 grams per liter to about 6 grams per liter, about 2 grams per liter to about 7 grams per liter, about 3 grams per liter to about 8 grams per liter, about 4 grams per liter to about 9 grams per liter, or about 5 grams per liter to about 10 grams per liter.

[0097] In embodiments, the amount of recombinant interferon protein detected in the extracted supernatant fraction is about 10% to about 95% of the amount of the recombinant interferon protein detected in the insoluble fraction. In embodiments, this amount is about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, or about 95%. In embodiments, this amount is about 10% to about 20%, 20% to about 50%, about 25% to about 50%, about 25% to about 50%, about 25% to about 95%, about 30% to about 50%, about 30% to about 40%, about 30% to about 60%, about 30% to about 70%, about 35% to about 50%, about 35% to about 70%, about 35% to about 75%, about 35% to about 95%, about 40% to about 50%, about 40% to about 95%, about 50% to about 75%, about 50% to about 95%, or about 70% to about 95%.

[0098] The protein yield measured in the unextracted insoluble fraction is typically higher than that in the extract supernatant, as material is lost during the extraction procedure. Yields from fermentation cultures are typically higher than those from smaller HTP cultures.

[0099] “Percent total cell protein” is the amount of protein or polypeptide in the host cell as a percentage of aggregate cellular protein. The determination of the percent total cell protein is well known in the art.

[00100] In embodiments, the amount of interferon protein detected in the extracted supernatant fraction produced is about 1% to about 75% of the total cell protein. In certain embodiments, the recombinant protein produced is about 1%, about 2%, about 3%, about 4%, about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 1% to about 5%, about 1% to about 10%, about 1% to about 20%, about 1% to about 30%, about 1% to about 40%, about 1% to about 50%, about 1% to about 60%, about 1% to about 75%, about 2% to about 5%, about 2% to about 10%, about 2% to about 20%, about 2% to about 30%, about 2% to about 40%, about 2% to about 50%, about 2% to about 60%, about 2% to about 75%, about 3% to about 5%, about 3% to about 10%, about 3% to about 20%, about 3% to about 30%, about 3% to about 40%, about 3% to about 50%, about 3% to about 60%, about 3% to about 75%, about 4% to about 10%, about 4% to about 20%, about 4% to about 30%, about 4% to about 40%, about 4% to about 50%, about 4% to about 60%, about 4% to about 75%, about 5% to about 10%, about 5% to about 20%, about 5% to about 30%, about 5% to about 40%, about 5% to about 50%, about 5% to about 60%, about 5% to about 75%, about 10% to about 20%, about 10% to about 30%, about 10% to about 40%, about 10% to about 50%, about 10% to about 60%, about 10% to about 75%, about 20% to about 30%, about 20% to about 40%, about 20% to about 50%, about 20% to about 60%, about 20% to about 75%, about 30% to about 40%, about 30% to about 50%, about 30% to about 60%, about 30% to about 75%, about 40% to about 50%, about 40% to about 60%, about 40% to about 75%, about 50% to about 60%, about 50% to about 75%, about 60% to about 75%, or about 70% to about 75%, of the total cell protein.

[00101] Solubility and Activity

[00102] The “solubility” and “activity” of a protein, though related qualities, are generally determined by different means. Solubility of a protein, particularly a hydrophobic protein, indicates that hydrophobic amino acid residues are improperly located on the outside of the folded protein. Protein activity, which can be evaluated using different methods, e.g., as described below, is another indicator of proper protein conformation. “Soluble, active, or both” as used herein, refers to protein that is determined to be soluble, active, or both soluble and active, by methods known to those of skill in the art.

[00103] Interferon Activity Assays

[00104] Assays for evaluating interferon protein activity are known in the art and include binding activity assays that measure binding of interferon to a standard ligand, e.g., a Blue sepharose column or a specific antibody column.

[00105] Biological activity of interferons can be quantitated using known assays, many of which are available commercially in kits. Most or all interferon species have been shown to exert a similar spectrum of in vitro biological activities in responsive cells, despite the existence of different receptors for type I and type II IFN. Biological activities induced by IFN include antiviral activity, which is mediated via cell receptors and is dependent on the activation of signaling pathways, the expression of specific gene products, and the development of antiviral mechanisms. Sensitivity of cells to IFN-mediated antiviral activity is variable, and depends on a number of factors including cell type, expression of IFN receptors and downstream effector response elements, effectiveness of antiviral mechanisms, and the type of virus used to infect cells.

[00106] Biological activity assays include, e.g., cytopathic effect assays (CPE), antiviral activity assays, assays that measure inhibition of cell proliferation, regulation of functional cellular activities, regulation of cellular differentiation and immunomodulation. Reporter gene assays include the luciferase reporter cell-based assay described herein in the Examples. In a reporter gene assay, the promoter region of an IFN responsive gene is linked with a heterologous reporter gene, for example, firefly luciferase or alkaline phosphatase, and transfected into an IFN-sensitive cell line. Stably transfected cell lines exposed to IFN increase expression of the reporter gene product in direct relation to the dose of IFN, the readout being a measure of this product's enzymic action. Many activity assay tools and kits are available commercially. Biological assays for interferons are described, e.g., by Meager A, "Biological assays for interferons," 1 Mar 2002, J. Immunol. Methods 261(1-2):21-36, incorporated herein by reference.

[00107] In embodiments, activity is represented by the % active protein in the extract supernatant as compared with the total amount assayed. This is based on the amount of protein determined to be active by the assay relative to the total amount of protein used in assay. In other embodiments, activity is represented by the % activity level of the protein compared to a standard, e.g., native protein. This is based on the amount of active protein in supernatant extract sample relative to the amount of active protein in a standard sample (where the same amount of protein from each sample is used in assay).

[00108] In embodiments, about 40% to about 100% of the recombinant interferon protein is determined to be active. In embodiments, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, or about 100% of the recombinant interferon protein is determined to be active. In embodiments, about 40% to about 50%, about 50% to about 60%, about

60% to about 70%, about 70% to about 80%, about 80% to about 90%, about 90% to about 100%, about 50% to about 100%, about 60% to about 100%, about 70% to about 100%, about 80% to about 100%, about 40% to about 90%, about 40% to about 95%, about 50% to about 90%, about 50% to about 95%, about 50% to about 100%, about 60% to about 90%, about 60% to about 95%, about 60% to about 100%, about 70% to about 90%, about 70% to about 95%, about 70% to about 100%, or about 70% to about 100% of the recombinant interferon protein is determined to be active.

5 [00109] In other embodiments, about 75% to about 100% of the recombinant interferon protein is determined to be active. In embodiments, about 75% to about 80%, about 75% to about 85%, about 75% to about 90%, about 75% to about 95%, about 80% to about 85%, about 80% to about 90%, about 80% to about 95%, about 80% to about 100%, about 85% to about 90%, about 85% to about 95%, about 85% to about 100%, about 90% to about 95%, about 90% to about 100%, or about 95% to about 100% of the recombinant interferon protein is determined to be active.

10

15 Expression Systems

10 [00110] Methods for expressing heterologous proteins, including useful regulatory sequences (e.g., promoters, secretion leaders, and ribosome binding sites), in *Pseudomonas* host cells, as well as host cells useful in the methods of the present invention, are described, e.g., in U.S. Pat. App. Pub. No. 2008/0269070 and U.S. Pat. App. Ser. No. 12/610,207, both titled “Method for Rapidly Screening Microbial Hosts to Identify Certain Strains with Improved Yield and/or Quality in the Expression of Heterologous Proteins,” U.S. Pat. App. Pub. No. 2006/0040352, “Expression of Mammalian Proteins in *Pseudomonas Fluorescens*,” and U.S. Pat. App. Pub. No. 2006/0110747, “Process for Improved Protein Expression by Strain Engineering,” all incorporated herein by reference in their entirety. These publications also describe bacterial host strains useful in practicing the methods of the invention, that have been engineered to overexpress folding modulators or wherein protease mutations have been introduced, in order to increase heterologous protein expression. Sequence leaders are described in detail in U.S. Patent App. Pub. No. 2008/0193974, “Bacterial leader sequences for increased expression,” and U.S. Pat. App. Pub. No. 2006/0008877, “Expression systems with Sec-secretion,” both incorporated herein by reference in their entirety, as well as in U.S. Pat. App. Ser. No. 12/610,207.

20

25

30

Promoters

35 [00111] The promoters used in accordance with the present invention may be constitutive promoters or regulated promoters. Common examples of useful regulated promoters include those of the family derived from the lac promoter (i.e. the lacZ promoter), especially the tac and trc promoters described in U.S. Pat. No. 4,551,433 to DeBoer, as well as Ptac16, Ptac17, PtacII,

PlacUV5, and the T7lac promoter. In one embodiment, the promoter is not derived from the host cell organism. In certain embodiments, the promoter is derived from an *E. coli* organism.

5 [00112] Inducible promoter sequences can be used to regulate expression of interferons in accordance with the methods of the invention. In embodiments, inducible promoters useful in the methods of the present invention include those of the family derived from the lac promoter (i.e. the lacZ promoter), especially the tac and trc promoters described in U.S. Pat. No. 4,551,433 to DeBoer, as well as Ptac16, Ptac17, PtacII, PlacUV5, and the T7lac promoter. In one embodiment, the promoter is not derived from the host cell organism. In 10 certain embodiments, the promoter is derived from an *E. coli* organism.

[00113] Common examples of non-lac-type promoters useful in expression systems according to the present invention include, e.g., those listed in Table 5.

Table 5. Examples of non-*lac* Promoters

Promoter	Inducer
P _R	High temperature
P _L	High temperature
Pm	Alkyl- or halo-benzoates
Pu	Alkyl- or halo-toluenes
Psal	Salicylates

15 [00114] See, e.g.: J. Sanchez-Romero & V. De Lorenzo (1999) Manual of Industrial Microbiology and Biotechnology (A. Demain & J. Davies, eds.) pp. 460-74 (ASM Press, Washington, D.C.); H. Schweizer (2001) Current Opinion in Biotechnology, 12:439-445; and R. Slater & R. Williams (2000) Molecular Biology and Biotechnology (J. Walker & R. Rapley, eds.) pp. 125-54 (The Royal Society of Chemistry, Cambridge, UK)). A promoter having the 20 nucleotide sequence of a promoter native to the selected bacterial host cell also may be used to control expression of the transgene encoding the target polypeptide, e.g., a *Pseudomonas* anthranilate or benzoate operon promoter (Pant, Pben). Tandem promoters may also be used in which more than one promoter is covalently attached to another, whether the same or different in sequence, e.g., a Pant-Pben tandem promoter (interpromoter hybrid) or a Plac- 25 Plac tandem promoter, or whether derived from the same or different organisms.

[00115] Regulated promoters utilize promoter regulatory proteins in order to control transcription of the gene of which the promoter is a part. Where a regulated promoter is used herein, a corresponding promoter regulatory protein will also be part of an expression system according to the present invention. Examples of promoter regulatory proteins include: 30 activator proteins, e.g., *E. coli* catabolite activator protein, MalT protein; AraC family

transcriptional activators; repressor proteins, e.g., *E. coli* LacI proteins; and dual-function regulatory proteins, e.g., *E. coli* NagC protein. Many regulated-promoter/promoter-regulatory-protein pairs are known in the art. In one embodiment, the expression construct for the target protein(s) and the heterologous protein of interest are under the control of the same regulatory element.

5 [00116] Promoter regulatory proteins interact with an effector compound, i.e., a compound that reversibly or irreversibly associates with the regulatory protein so as to enable the protein to either release or bind to at least one DNA transcription regulatory region of the gene that is under the control of the promoter, thereby permitting or blocking the action of a transcriptase enzyme in initiating transcription of the gene. Effector compounds are classified as either inducers or co-repressors, and these compounds include native effector compounds and gratuitous inducer compounds. Many regulated-promoter/promoter-regulatory-protein/effector-compound trios are known in the art. Although an effector compound can be used throughout the cell culture or fermentation, in a preferred 10 embodiment in which a regulated promoter is used, after growth of a desired quantity or density of host cell biomass, an appropriate effector compound is added to the culture to directly or indirectly result in expression of the desired gene(s) encoding the protein or polypeptide of interest.

15 [00117] In embodiments wherein a lac family promoter is utilized, a lacI gene can also be present in the system. The lacI gene, which is normally a constitutively expressed gene, encodes the Lac repressor protein LacI protein, which binds to the lac operator of lac family promoters. Thus, where a lac family promoter is utilized, the lacI gene can also be included and expressed in the expression system.

20 [00118] Promoter systems useful in *Pseudomonas* are described in the literature, e.g., in U.S. Pat. App. Pub. No. 2008/0269070, also referenced above.

Other Regulatory Elements

25 [00119] In embodiments, soluble proteins are present in either the cytoplasm or periplasm of the cell during production. Secretion leaders useful for targeting proteins are described elsewhere herein, and in U.S. Pat. App. Pub. No. 2008/0193974, U.S. Pat. App. Pub. No. 2006/0008877, and in U.S. Pat. App. Ser. No. 12/610,207, referenced above.

30 [00120] An expression construct useful in practicing the methods of the present invention can include, in addition to the protein coding sequence, the following regulatory elements operably linked thereto: a promoter, a ribosome binding site (RBS), a transcription terminator, and translational start and stop signals. Useful RBSs can be obtained from any 35 of the species useful as host cells in expression systems according to, e.g., U.S. Pat. App. Pub. No. 2008/0269070 and U.S. Pat. App. Ser. No. 12/610,207. Many specific and a

variety of consensus RBSs are known, e.g., those described in and referenced by D. Frishman et al., Gene 234(2):257-65 (8 Jul. 1999); and B. E. Suzek et al., Bioinformatics 17(12):1123-30 (December 2001). In addition, either native or synthetic RBSs may be used, e.g., those described in: EP 0207459 (synthetic RBSs); O. Ikehata et al., Eur. J. Biochem. 181(3):563-70 (1989) (native RBS sequence of AAGGAAG). Further examples of methods, vectors, and translation and transcription elements, and other elements useful in the present invention are described in, e.g.: U.S. Pat. No. 5,055,294 to Gilroy and U.S. Pat. No. 5,128,130 to Gilroy et al.; U.S. Pat. No. 5,281,532 to Rammller et al.; U.S. Pat. Nos. 4,695,455 and 4,861,595 to Barnes et al.; U.S. Pat. No. 4,755,465 to Gray et al.; and U.S. Pat. No. 5,169,760 to Wilcox.

10 Host Strains

[00121] Bacterial hosts, including *Pseudomonas*, and closely related bacterial organisms are contemplated for use in practicing the methods of the invention. In certain embodiments, the *Pseudomonas* host cell is *Pseudomonas fluorescens*. The host cell can also be an *E. coli* cell.

[00122] *Pseudomonas* and closely related bacteria are generally part of the group defined as "Gram(+) Proteobacteria Subgroup 1" or "Gram-Negative Aerobic Rods and Cocci" (Buchanan and Gibbons (eds.) (1974) Bergey's Manual of Determinative Bacteriology, pp. 217-289). *Pseudomonas* host strains are described in the literature, e.g., in U.S. Pat. App. Pub. No. 2006/0040352, cited above.

[00123] For example, *Pseudomonas* hosts can include cells from the genus *Pseudomonas*, *Pseudomonas enalia* (ATCC 14393), *Pseudomonas nigrifaciensi* (ATCC 19375), and *Pseudomonas putrefaciens* (ATCC 8071), which have been reclassified respectively as *Alteromonas haloplanktis*, *Alteromonas nigrifaciens*, and *Alteromonas putrefaciens*. Similarly, e.g., *Pseudomonas acidovorans* (ATCC 15668) and *Pseudomonas testosteroni* (ATCC 11996) have since been reclassified as *Comamonas acidovorans* and *Comamonas testosteroni*, respectively; and *Pseudomonas nigrifaciens* (ATCC 19375) and *Pseudomonas piscicida* (ATCC 15057) have been reclassified respectively as *Pseudoalteromonas nigrifaciens* and *Pseudoalteromonas piscicida*.

[00124] The host cell can be selected from "Gram-negative Proteobacteria Subgroup 16." "Gram-negative Proteobacteria Subgroup 16" is defined as the group of Proteobacteria of the following *Pseudomonas* species (with the ATCC or other deposit numbers of exemplary strain(s) shown in parenthesis): *Pseudomonas abietaniphila* (ATCC 700689); *Pseudomonas aeruginosa* (ATCC 10145); *Pseudomonas alcaligenes* (ATCC 14909); *Pseudomonas anguilliseptica* (ATCC 33660); *Pseudomonas citronellolis* (ATCC 13674); *Pseudomonas flavaescens* (ATCC 51555); *Pseudomonas mendocina* (ATCC 25411); *Pseudomonas*

nitroreducens (ATCC 33634); *Pseudomonas oleovorans* (ATCC 8062); *Pseudomonas pseudoalcaligenes* (ATCC 17440); *Pseudomonas resinovorans* (ATCC 14235); *Pseudomonas straminea* (ATCC 33636); *Pseudomonas agarici* (ATCC 25941); *Pseudomonas alcaliphila*; *Pseudomonas alginovora*; *Pseudomonas andersonii*; *Pseudomonas asplenii* (ATCC 23835); *Pseudomonas azelaica* (ATCC 27162); *Pseudomonas beyerinckii* (ATCC 19372); *Pseudomonas borealis*; *Pseudomonas boreopolis* (ATCC 33662); *Pseudomonas brassicacearum*; *Pseudomonas butanovora* (ATCC 43655); *Pseudomonas cellulosa* (ATCC 55703); *Pseudomonas aurantiaca* (ATCC 33663); *Pseudomonas chlororaphis* (ATCC 9446, ATCC 13985, ATCC 17418, ATCC 17461); *Pseudomonas fragi* (ATCC 4973); *Pseudomonas lundensis* (ATCC 49968); *Pseudomonas taetrolens* (ATCC 4683); *Pseudomonas cissicola* (ATCC 33616); *Pseudomonas coronafaciens*; *Pseudomonas diterpeniphila*; *Pseudomonas elongata* (ATCC 10144); *Pseudomonas flectens* (ATCC 12775); *Pseudomonas azotoformans*; *Pseudomonas brenneri*; *Pseudomonas cedrella*; *Pseudomonas corrugata* (ATCC 29736); *Pseudomonas extreorientalis*; *Pseudomonas fluorescens* (ATCC 35858); *Pseudomonas gessardii*; *Pseudomonas libanensis*; *Pseudomonas mandelii* (ATCC 700871); *Pseudomonas marginalis* (ATCC 10844); *Pseudomonas migulae*; *Pseudomonas mucidolens* (ATCC 4685); *Pseudomonas orientalis*; *Pseudomonas rhodesiae*; *Pseudomonas synxantha* (ATCC 9890); *Pseudomonas tolaasii* (ATCC 33618); *Pseudomonas veronii* (ATCC 700474); *Pseudomonas frederiksbergensis*; *Pseudomonas geniculata* (ATCC 19374); *Pseudomonas gingeri*; *Pseudomonas graminis*; *Pseudomonas grimontii*; *Pseudomonas halodenitrificans*; *Pseudomonas halophila*; *Pseudomonas hibiscicola* (ATCC 19867); *Pseudomonas huttiensis* (ATCC 14670); *Pseudomonas hydrogenovora*; *Pseudomonas jessenii* (ATCC 700870); *Pseudomonas kilonensis*; *Pseudomonas lanceolata* (ATCC 14669); *Pseudomonas lini*; *Pseudomonas marginata* (ATCC 25417); *Pseudomonas mephitica* (ATCC 33665); *Pseudomonas denitrificans* (ATCC 19244); *Pseudomonas pertucinogena* (ATCC 190); *Pseudomonas pictorum* (ATCC 23328); *Pseudomonas psychrophila*; *Pseudomonas silva* (ATCC 31418); *Pseudomonas monteili* (ATCC 700476); *Pseudomonas mosselii*; *Pseudomonas oryzihabitans* (ATCC 43272); *Pseudomonas plecoglossicida* (ATCC 700383); *Pseudomonas putida* (ATCC 12633); *Pseudomonas reactans*; *Pseudomonas spinosa* (ATCC 14606); *Pseudomonas balearica*; *Pseudomonas luteola* (ATCC 43273); *Pseudomonas stutzeri* (ATCC 17588); *Pseudomonas amygdali* (ATCC 33614); *Pseudomonas avellanae* (ATCC 700331); *Pseudomonas caricapapayae* (ATCC 33615); *Pseudomonas eichorii* (ATCC 10857); *Pseudomonas ficusrectae* (ATCC 35104); *Pseudomonas fuscovaginae*; *Pseudomonas meliae* (ATCC 33050); *Pseudomonas syringae* (ATCC 19310); *Pseudomonas viridiflava* (ATCC 13223); *Pseudomonas*

thermocarboxydovorans (ATCC 35961); *Pseudomonas thermotolerans*; *Pseudomonas thivervalensis*; *Pseudomonas vancouverensis* (ATCC 700688); *Pseudomonas wisconsinensis*; and *Pseudomonas xiamenensis*.

[00125] The host cell can also be selected from "Gram-negative Proteobacteria Subgroup 17." 5 "Gram-negative Proteobacteria Subgroup 17" is defined as the group of Proteobacteria known in the art as the "fluorescent Pseudomonads" including those belonging, e.g., to the following *Pseudomonas* species: *Pseudomonas azotoformans*; *Pseudomonas brenneri*; *Pseudomonas cedrella*; *Pseudomonas corrugata*; *Pseudomonas extremorientalis*; *Pseudomonas fluorescens*; *Pseudomonas gessardii*; *Pseudomonas libanensis*; *Pseudomonas mandelii*; *Pseudomonas marginalis*; *Pseudomonas migulae*; *Pseudomonas mucidolens*; 10 *Pseudomonas orientalis*; *Pseudomonas rhodesiae*; *Pseudomonas synxantha*; *Pseudomonas tolaasii*; and *Pseudomonas veronii*.

Codon Optimization

[00126] Methods for optimizing codons to improve expression in bacterial hosts are known in the art 15 and described in the literature. For example, optimization of codons for expression in a *Pseudomonas* host strain is described, e.g., in U.S. Pat. App. Pub. No. 2007/0292918, "Codon Optimization Method," incorporated herein by reference in its entirety.

[00127] Codon optimization for expression in *E. coli* is described, e.g., by Welch, et al., 2009, PLoS 20 One, "Design Parameters to Control Synthetic Gene Expression in Escherichia coli, 4(9): e7002, Ghane, et al., 2008, "Overexpression of Biologically Active Interferon B Using Synthetic Gene in *E. coli*," Journal of Sciences, Islamic Republic of Iran 19(3): 203-209, and Valente, et al., 2004, "Translational Features of Human Alpha 2b Interferon Production in Escherichia coli," Applied and Environmental Microbiology 70(8): 5033-5036, all incorporated by reference herein.

25 Fermentation Format

[00128] The expression system according to the present invention can be cultured in any 30 fermentation format. For example, batch, fed-batch, semi-continuous, and continuous fermentation modes may be employed herein.

[00129] In embodiments, the fermentation medium may be selected from among rich media, 35 minimal media, and mineral salts media. In other embodiments either a minimal medium or a mineral salts medium is selected. In certain embodiments, a mineral salts medium is selected.

[00130] Mineral salts media consists of mineral salts and a carbon source such as, e.g., glucose, 35 sucrose, or glycerol. Examples of mineral salts media include, e.g., M9 medium, *Pseudomonas* medium (ATCC 179), and Davis and Mingioli medium (see, B D Davis & E S Mingioli (1950) J. Bact. 60:17-28). The mineral salts used to make mineral salts media

5 include those selected from among, e.g., potassium phosphates, ammonium sulfate or chloride, magnesium sulfate or chloride, and trace minerals such as calcium chloride, borate, and sulfates of iron, copper, manganese, and zinc. Typically, no organic nitrogen source, such as peptone, tryptone, amino acids, or a yeast extract, is included in a mineral salts medium. Instead, an inorganic nitrogen source is used and this may be selected from among, e.g., ammonium salts, aqueous ammonia, and gaseous ammonia. A mineral salts medium will typically contain glucose or glycerol as the carbon source. In comparison to mineral salts media, minimal media can also contain mineral salts and a carbon source, but can be supplemented with, e.g., low levels of amino acids, vitamins, peptones, or other 10 ingredients, though these are added at very minimal levels. Media can be prepared using the methods described in the art, e.g., in U.S. Pat. App. Pub. No. 2006/0040352, referenced and incorporated by reference above. Details of cultivation procedures and mineral salts media useful in the methods of the present invention are described by Riesenber, D et al., 1991, "High cell density cultivation of *Escherichia coli* at controlled specific growth rate," J. 15 Biotechnol. 20 (1):17-27.

[00131] Fermentation may be performed at any scale. The expression systems according to the present invention are useful for recombinant protein expression at any scale. Thus, e.g., microliter-scale, centiliter scale, and deciliter scale fermentation volumes may be used, and 1 Liter scale and larger fermentation volumes can be used.

20 [00132] In embodiments, the fermentation volume is at or above about 1 Liter. In embodiments, the fermentation volume is about 1 liter to about 100 liters. In embodiments, the fermentation volume is about 1 liter, about 2 liters, about 3 liters, about 4 liters, about 5 liters, about 6 liters, about 7 liters, about 8 liters, about 9 liters, or about 10 liters. In embodiments, the fermentation volume is about 1 liter to about 5 liters, about 1 liter to about 10 liters, about 1 liter to about 25 liters, about 1 liter to about 50 liters, about 1 liter to about 75 liters, about 25 10 liters to about 25 liters, about 25 liters to about 50 liters, or about 50 liters to about 100 liters. In other embodiments, the fermentation volume is at or above 5 Liters, 10 Liters, 15 Liters, 20 Liters, 25 Liters, 50 Liters, 75 Liters, 100 Liters, 200 Liters, 500 Liters, 1,000 Liters, 2,000 Liters, 5,000 Liters, 10,000 Liters, or 50,000 Liters.

30 [00133] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed 35 in practicing the invention. It is intended that the following claims define the scope of the

invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

EXAMPLES

Example 1: Production of rIFN- β from High Throughput Expression Samples

5 [00134] In the following experiment, IFN- β C17S expression strains were constructed, and the amount of IFN- β in the insoluble fraction obtained for each was quantitated. Based on the resulting data, certain strains were selected for use in optimizing the non-denaturing extraction process of the present invention.

[00135] Construction and Growth of IFN- β Expression Strains

10 [00136] The IFN- β 1b coding sequence was constructed using *P. fluorescens* preferred codons to encode for the human IFN- β amino acid sequence corresponding to the therapeutic Betaseron. Figure 7 shows the amino acid and DNA sequences of the synthetic IFN- β (Betaseron) gene.

[00137] Plasmids were constructed which carry the codon-optimized IFN- β fused to nineteen 15 *P. fluorescens* secretion leaders. The secretion leaders were included to target the protein to the periplasm where it may be recovered in the properly folded and active form. In addition, one plasmid was constructed which carries the codon-optimized IFN- β designed for cytoplasmic expression.

[00138] Expression of IFN- β was driven from the Ptac promoter and translation initiated from either 20 a high (Hi) or medium (Med) activity ribosome binding site (RBS). The resulting 20 plasmids were transformed into 30 *P. fluorescens* host strains (16 protease deletion strains, 13 folding modulator overexpression strains and 1 wild type strain) to produce 600 expression strains (see Tables 6 and 7). Folding modulators, when present, were encoded on a second plasmid and expression driven by a mannitol inducible promoter.

25 [00139] The thirty host strains carrying each of 20 IFN- β expression plasmids (600 expression hosts in total) were grown in triplicate in 96-well plates. Samples harvested 24 hours after induction were used for analysis.

Expression of IFN- β Using Pfenex Expression Technology in 96-Well Format

[00140] Each plasmid (Table 6) was transformed into 30 *P. fluorescens* host strains (Table 7) as 30 follows: Twenty-five microliters of competent cells were thawed and transferred into a 96-well electroporation plate (BTX ECM630 Electroporator), and 1 microliter miniprep plasmid DNA was added to each well. Cells were electroporated at 2.5 KV, 200 Ohms, and 25 μ F. Cells were resuspended in 75 microliters HTP-YE media with trace minerals, transferred to 96-well deep well plate with 500 μ l M9 salts 1% glucose medium (seed culture), and incubated at 30°C, shaking 300 rpm and 50-mm diameter throw for 48 hours.

[00141] Ten microliters of seed culture were transferred into triplicate wells of 96-well deep well plates, each well containing 500 microliters of HTP-YE medium, and incubated as before for 24 hours. Isopropyl- β -D-1-thiogalactopyranoside (IPTG) was added to each well for a final concentration of 0.3 mM to induce the expression of IFN- β . For growth of small cultures in HTP microwells, a specific culture pH is not tightly controlled and the cell density can differ slightly from well to well. Mannitol (Sigma, M1902) was added to each well at a final concentration of 1% to induce the expression of folding modulators in folding modulator over-expressing strains, and the temperature was reduced to 25°C. Twenty four hours after induction, cultures were collected for analysis. For OD normalization, cells were mixed with sterile 1X PBS to obtain a final OD₆₀₀ = 20 in a final volume of 400 microliters using the Biomek liquid handling station (Beckman Coulter). Samples were collected in cluster tube racks.

Sample Preparation and SDS-CGE Analysis

[00142] Soluble fractions (supernatants obtained after centrifugation of lysates) and insoluble fractions (pellets obtained after centrifugation of lysates) were prepared by sonicating the OD-normalized cultures, followed by centrifugation. Frozen, normalized culture broth (400 microliters) was thawed and sonicated for 3.5 minutes. The lysates were centrifuged at 20,800 \times g for 20 minutes (4°C) and the soluble fractions removed using manual or automated liquid handling. The insoluble fractions were frozen and then thawed for re-centrifugation at 20,080 \times g for 20 minutes at 4°C, to remove residual supernatant. The insoluble fractions were then resuspended in 400 μ L of 1X phosphate buffered saline (PBS), pH 7.4. Further dilutions of soluble and insoluble fractions for SDS-CGE analysis were performed in 1X phosphate buffered saline (PBS), pH 7.4. Soluble and insoluble fractions were prepared for analysis by SDS capillary gel electrophoresis (CGE) (Caliper Life Sciences, Protein Express LabChip Kit, Part 760301), in the presence of dithiothreitol (DTT).

[00143] Normalized soluble and insoluble fractions from each well of the 600 strains expressing IFN- β were analyzed by reducing SDS-CGE analysis in one replicate for the soluble fractions and insoluble fractions. No IFN- β signal was detected in the soluble fractions. IFN- β signal varied from no signal to greater than 400 mg/L in the insoluble fractions. Only five of the twenty plasmids tested showed measurable signal of IFN- β in the insoluble fractions of all thirty host strains: p530-001, p530-007, p530-011, p530-018 and p530-020. Valley to valley integration of IFN- β signal using Caliper LabChipGX software was performed in all 150 strains consisting of the five plasmids listed above in the thirty host strains, and data were used to calculate volumetric yields. Volumetric yields of the 150 strains analyzed ranged from 2 to 482 mg/L. Strains carrying p530-020 attained

5 consistently higher yields of IFN- β in the insoluble fraction than other expression strains; however, the protein migrated higher than expected on SDS-CGE, indicating that the secretion leader was not cleaved. High yields were also observed with 2 host strains carrying p530-001. No significant difference in IFN- β in the insoluble fraction was observed among the 30 strains except potentially in one strain, DC441, a lon hslUV protease deletion strain, which showed somewhat higher yields than the other 29 strains.

10 [00144] A subset of 17 top expression strains (Table 8), excluding strains containing plasmid p530-020, was selected for further analyses. The expression strains containing plasmid p530-020 were excluded from further consideration in this experiment due to the potentially unprocessed leader. SDS-CGE analysis was performed on the soluble and insoluble fractions for these strains. Quantification of the SDS-CGE output is shown in Table 8. IFN- β protein concentration ranged from 102 to greater than 482 mg/L. Based upon insoluble yield and processing of either the periplasmic leader sequence or the N-terminal Met from IFN- β , strains were chosen to proceed to fermentation assessment.

15

[00145] Table 6. Plasmids

Plasmid	Expression Vector	Secretion Leader	RBS
p530-001	pDOW5271	None	Hi
p530-002	pDOW5204	Pbp	Med
p530-003	pDOW5206	DsbA	Hi
p530-004	pDOW5207	DsbA	Med
p530-005	pDOW5209	Azu	Hi
p530-006	pDOW5210	Azu	Med
p530-007	pDOW5217	LAO	Hi
p530-008	pDOW5220	Ibp-S31A	Hi
p530-009	pDOW5223	TolB	Hi
p530-010	pDOW5226	Trc	Hi
p530-011	pDOW5232	Ttg2C	Hi
p530-012	pDOW5235	FlgI	Hi
p530-013	pDOW5238	CupC2	Hi
p530-014	pDOW5241	CupB2	Hi
p530-015	pDOW5244	CupA2	Hi
p530-016	pDOW5247	NikA	Hi
p530-017	pDOW5256	PorE	Hi

p530-018	pDOW5259	Pbp-A20V	Hi
p530-019	pDOW5262	DsbC	Hi
p530-020	pDOW5265	Bcc	Hi

[00146] Table 7. IFN- β Expression Strains

Strain Name	Strain Description	Strain Name	Strain Description
DC454	Wild type	DC539	FMO
DC441	PD	DC544	FMO
DC462	FMO	DC547	FMO
DC468	PD	DC548	FMO
DC469	PD	DC552	FMO
DC485	PD	DC565	FMO
DC486	PD	DC566	FMO
DC487	PD	DC567	FMO
DC488	PD	DC568	FMO
DC489	PD	DC575	FMO
DC490	PD	DC584	FMO
DC491	PD	DC598	FMO
DC492	PD	DC599	FMO
DC498	PD	DC667	FMO
DC538	FMO	DC954	PD

PD= protease deletion strain, FMO= folding modulator over-expression strain

5 [00147] Table 8. Calculated Volumetric IFN- β Yields of Top 17 Strains by SDS-CGE

Strain Name	Vol. Yield > 100 ug/ml	Plasmid	Host Strain	Leader
PS530-001	482.3	p530-001	DC441	x
PS530-101	216.5	p530-001	DC485	x
PS530-011	161.1	p530-011	DC441	ttg2C
PS530-071	148.8	p530-011	DC468	ttg2C
PS530-007	141.2	p530-007	DC441	Lao
PS530-031	131.3	p530-011	DC454	ttg2C
PS530-201	122.6	p530-001	DC490	x
PS530-531	121.0	p530-011	DC598	ttg2C
PS530-211	119.8	p530-011	DC490	ttg2C
PS530-151	119.8	p530-011	DC487	ttg2C
PS530-061	119.6	p530-001	DC468	x
PS530-411	114.0	p530-011	DC565	ttg2C
PS530-231	113.3	p530-011	DC491	ttg2C
PS530-391	112.2	p530-011	DC552	ttg2C
PS530-027	104.5	p530-007	DC454	Lao
PS530-291	103.3	p530-011	DC538	ttg2C
PS530-271	102.2	p530-011	DC498	ttg2C

Example 2. Extraction of IFN- β 1b from High Throughput Expression Material

[00148] IFN- β 1b was successfully extracted from insoluble fractions from HTP expression cultures, using extraction conditions comprising Zwittergent 3-14 detergent.

5 [00149] HTP expression plate cultures of *Pseudomonas fluorescens* strains PS530-001 overexpressing cytoplasmic IFN- β 1b and 530-220, overexpressing secreted IFN- β 1b (described in Example 1), were sonicated and centrifuged to obtain an insoluble fraction and a soluble fraction. The pellets were resuspended in extraction buffer 1X PBS, pH 7.4 or sodium acetate at pH 4.5. Each buffer was tested either with or without Zwittergent 3-14 detergent at 1% (w/v). Each of the four combinations of buffer and detergent was incubated for 1-2 hours at room temperature or overnight at 4° C with shaking. After extraction, each sample was centrifuged for 20 minutes at 20,080 x g at 4° C to produce a second insoluble pellet fraction (extract pellet) and a second soluble supernatant fraction (extract supernatant). The first insoluble fraction and first soluble fraction, and the extract pellet fraction and extract supernatant fraction, were analyzed by SDS-CGE. The results are shown in Figures 1A and 1B. As seen in Lane 7, the extraction condition including PBS buffer and Zwittergent 3-14 yielded soluble IFN- β .

10

15

Example 3. Optimization of Conditions for Extraction

[00150] Insoluble fractions from fermentation cultures were extracted under conditions comprising different detergents.

[00151] Frozen cell paste from a 1 L fermentation (grown at 32° C, pH 6.5, and induced using 0.2mM IPTG at an OD₅₇₅ of 100) of strain PS530-001, overexpressing recombinant IFN- β 1b, was resuspended in lysis buffer containing 20 mM sodium phosphate (JT Baker), pH 7.4 to a final solids concentration of 20% (w/v). The well-mixed cell slurry was lysed with two

passes at 38 kpsi through a Constant cell disruptor (Constant Systems, Inc.). The lysate was split in half, and spun by centrifugation at 15,000 x g for 30 minutes at 4 °C (Beckman Coulter, PN# J-20, XPF). The pellets (containing IFN- β and cell debris) were resuspended and each was washed in either Buffer A (20 mM sodium phosphate, pH 7.4) or Buffer B (20 mM sodium acetate, pH 4.0). Samples were spun by centrifugation under the same conditions described for the first spin, supernatants were removed, and the pellets were again resuspended in either Buffer A or B at 20% solid concentration. For each buffer, twenty aliquots of 1 mL each were placed in 1.5 mL conical tubes. Detergent stock solutions were added to the conical tubes at different concentrations. All tubes were 5 incubated at room temperature for 1 hour or overnight (18 hours) at 4 °C with continuous mixing. After extraction, the solutions were centrifuged and the extract supernatant 10 fractions were analyzed for protein concentration by SDS-CGE. Figure 2 provides a flow chart showing how the sample preparation and extraction were carried out.

[00152] Of the detergents tested, Zwittergent 3-14 and N-lauroylsarcosine (NLS), were found to give 15 the best yields regardless of buffer and incubation time (Table 9). However, the product extracted using NLS was not active, as determined by its inability to bind to either a Blue Sepharose affinity column or a cation exchange column (SP HP) (data not shown). The product extracted using Zwittergent 3-14 was determined to be active.

Table 9. Evaluation of Detergents for Extraction

Detergent	Concentration (w/v)	Extracted Product Concentration (ug/mL)			
		Buffer A		Buffer B	
		1hr @RT	18hr @RT	1hr @RT	18hr @RT
Zwittergent 3-14	0.50%	748	557	1011	734
	1.00%	731	392	1060	936
	2.00%	903	398	1548	1146
Lauroylsarcosine	0.20%	1023	643	NA	NA
	0.50%	3104	2125	324	150
	1.50%	2782	2670	2319	2668
NDSB195	10.00%	8	6	11	46
	15.00%	14	13	31	119
NDSB256	5.00%	20	56	15	43
	15.00%	204	233	114	135
Chaps	0.50%	11	36	98	160

	2.00%	75	170	179	250
Octylglucopyranoside	1.00%	83	175	121	169
	5.00%	196	258	164	215
	0.50%	129	237	NA	NA
Tween-20	1.00%	196	274	NA	NA
	0.05%	4	11	NA	6
Tween-80	0.50%	11	37	3	18
	0.01%	4	6	NA	7
	0.10%	5	10	NA	12
Triton-100	0.50%	7	25	3	21
	0.10%	25	68	33	103
	1.00%	40	85	62	176

Evaluation of Zwittergent Analogs

[00153] Using similar methods, Zwittergent analogs were evaluated for their extraction efficiency.

The results are shown in Table 10. The best yields were observed with Zwittergent 3-14.

5 Zwittergent 3-12, Zwittergent 3-10, and Zwittergent 3-08 were also effective.

Table 10. Evaluation of Zwittergent Analogs for Extraction of IFN- β 1b

Detergent	Detergent Conc.	Solid Conc.	Protein (ug/mL)
Zwittergent 3-08	10%	20%	292
Zwittergent 3-10	1.0%	20%	233
Zwittergent 3-12	1.0%	20%	357
Zwittergent 3-16	0.1%	20%	17
Zwittergent 3-14	1.0%	20%	430
Zwittergent 3-14	1.0%	10%	396
Zwittergent 3-14	1.0%	5%	548

Evaluation of the Zwittergent 3-14 Concentration

[00154] To efficiently solubilize proteins, the detergent concentration typically needs to be above its

10 CMC value. The CMC of Zwittergent 3-14 is about 0.01% w/v. Extraction conditions including sodium phosphate buffer at pH 7.4 with increasing concentrations of Zwittergent 3-14 were evaluated. The cell paste used was obtained by growing strain PS530-001 at 32° C, pH 6.5, and induced using 0.2mM IPTG at an OD₅₇₅ of 100. The results in Table 11 show that use of Zwittergent 3-14 at 1% (w/v) concentration resulted in the highest extraction yield.

15

Table 11: Effect of Zwittergent 3-14 Concentration on Extraction of IFN- β 1b

Zwittergent 3-14 Concentration (% w/v)	Extraction Yield microgram/mL	Extraction Yield % IFN- β protein extracted from insoluble pellet (insoluble fraction)
0.01%	10	0%
0.05%	36	1%
0.10%	72	2%
0.50%	341	9%
1.00%	787	21%
2.00%	620	17%

Evaluation of Additional Chemical Reagents

[00155] As shown in Table 11, extraction conditions including Zwittergent 3-14 at 1% (w/v)

5 concentration in sodium phosphate buffer at pH 7.4 yielded 21% of the IFN- β 1b detected in the original insoluble fraction. Further optimization was conducted.

[00156] High concentration (e.g., 6 to 8 M) of some chaotropic reagents like urea and guanidinium hydrochloride commonly have been used as a strong denaturant for solubilization of inclusion bodies. Chaotropes such as urea can increase the detergent critical micelle concentration (CMC) and may potentially improve the extraction efficiency. Low concentrations of urea (up to 2 M) were evaluated in the extraction conditions. Salts, e.g., NaCl, can also affect detergent CMC. Varying Zwittergent 3-14 concentrations were evaluated due to the potential interplay between detergent CMC and the presence of chaotropic reagents and salts. The concentration of insoluble inclusion solids in the extraction conditions was also varied. Lower solids concentration than the 20% (w/v) previously used were evaluated.

10 [00157] In summary, the effect of varying the following parameters on extraction efficiency was tested.

[00158] Sodium Chloride: 150 - 1850 mM

15 [00159] Urea: 0 - 2 M

[00160] Zwittergent 3-14: 0.1 – 1.0% w/v

[00161] Solids: 5 – 20% w/v

[00162] pH: 6.5 – 8.5

20 [00163] The flow chart in Figure 3 describes the preparation and extraction of the first insoluble pellet fraction for this optimization study. Table 12 shows the result of the study. Figures 4A and B summarize the results and significance of the effect of each parameter on the extraction yield. For optimization of extraction of interferon β from the insoluble fraction, a

two-level five-factor half-fractional factorial experimental design was used. JMP software (SAS Institute, Cary, NC) was used for experimental design and analysis. The software estimates the effect of individual factors as well as interactions on experimental output (amount of interferon extracted).

5 TABLE 12. Results of Extraction Study

No.	Solids		pH	NaCl (M)	Urea (M)	Z314 (%)	Interferon- β in extract supernatant	
	(%)	No.					(mg/L)	
1	----+	5	6.5	0.15	0	1	2275	
2	---+-	5	6.5	0.15	2	0.1	896	
3	--+-+	5	6.5	1.85	0	0.1	246	
4	--+++	5	6.5	1.85	2	1	7024	
5	--++-	5	8.5	0.15	0	0.1	638	
6	--+++-	5	8.5	0.15	2	1	5614	
7	--+++-	5	8.5	1.85	0	1	5414	
8	--+++-	5	8.5	1.85	2	0.1	1711	
9	0	12.5	7.5	1	1	0.55	3362	
10	0	12.5	7.5	1	1	0.55	3693	
11	0	12.5	7.5	1	1	0.55	3809	
12	++++	20	6.5	0.15	0	0.1	65	
13	+++++	20	6.5	0.15	2	1	2345	
14	+++-+	20	6.5	1.85	0	1	2149	
15	+++-+	20	6.5	1.85	2	0.1	438	
16	+++-+	20	8.5	0.15	0	1	2350	
17	+++-+	20	8.5	0.15	2	0.1	677	
18	+++-+	20	8.5	1.85	0	0.1	199	
19	+++-+	20	8.5	1.85	2	1	4486	

[00164] Based on the above data, an optimized extraction condition was selected for experiments described hereinafter: 1% (w/v) Zwittergent 3-14, 2 M Urea, 2 M NaCl, Solids 5% w/v, buffer pH 7.5 to 8.5. Using these optimized conditions, the observed extraction yield (in the extract supernatant) was found to be consistently 90% or above (i.e., 90% or more of the amount of recombinant protein measured in the insoluble fraction).

10

Example 4. Production of rIFN- β 1b from Large Scale Fermentation

[00165] Production of recombinant human- β interferon (IFN- β 1b) protein in *Pseudomonas*

fluorescens Pfenex Expression TechnologyTM strain PS530-001 was successfully achieved

in 2 liter fermentors. Multiple fermentation conditions were evaluated resulting in expression of IFN- β 1b up to 9.2 g/L.

[00166] Fermentation cultures were grown in 2 liter fermentors containing a mineral salts medium (as described herein and also by, e.g., Riesenber, D., et al., 1991). Culture conditions were 5 maintained at 32° C and pH 6.5 through the addition of aqueous ammonia. Dissolved oxygen was maintained in excess through increases in agitation and flow of sparged air and oxygen into the fermentor. Glycerol was delivered to the culture throughout the fermentation to maintain excess levels. These conditions were maintained until the target culture optical density (A575) for induction was reached, at which time IPTG was added to 10 initiate IFN- β production. The optical density at induction, the concentration of IPTG, pH and temperature were all varied to determine optimal conditions for expression. After 24 hours, the culture from each fermentor was harvested by centrifugation and the cell pellet frozen at -80 °C.

[00167] Fermentation cultures were induced at 100 OD₅₇₅ using 0.2 mM IPTG, at pH 6.5 and a 15 temperature of 32° C. Replicate fermentations resulted in IFN- β production at 7.5, 8.4 and 7.9 g/L as determined by SDS-CGE of the initial insoluble fraction (Figure 5). When these insoluble fractions were subjected to extraction (under conditions including 1% (w/v) Zwittergent 3-14, 2 M Urea, 2 M NaCl, Solids 5% w/v, and buffer pH 8.2), solubilized IFN- β were observed in the extract supernatant at 2.2, 2.4, and 2.6 g/L. This represents an 20 average extraction yield of 31%.

[00168] Increasing the induction OD to 120 to 160, and decreasing the fermentation pH to 5.7 to 25 6.25, increased IFN- β titers in the initial insoluble fraction to 8.8-9.2 g/L (Figure 6). Extraction of these cell pellets (using the same extraction conditions as for the experiment shown in Figure 5) resulted in 3.1 to 4.0 g/L of IFN- β in the extracted supernatant fraction, an average extraction yield of 39% (Table 13).

Table 13. Extracted Solubilized IFN- β Based on Induction Conditions

Induction OD of 100 and pH 6.5				Induction OD of 120 - 160 and pH 5.7 to 6.25			
	Total Insoluble Titer (g/L)	Extracted Solubilized Titer (g/L)	Extracted Yield (%)		Total Insoluble Titer (g/L)	Extracted Solubilized Titer (g/L)	Extracted Yield (%)
u2	7.5	2.2	29	u2	9.2	4.0	43
u7	8.4	2.4	29	u3	8.8	3.1	35
u8	7.9	2.6	33	u5	8.8	ND	ND
average	7.9	2.4	31	average	8.9	3.5	39
std dev	0.4	0.2	2.3	std dev	0.3	0.6	5.6

5

Example 5. Activity Analysis of IFN- β Extraction Product

[00169] Broth from fermentation of *Pseudomonas fluorescens* strain PS530-001 (1L fermentation at 32° C, pH 6.0, induced at OD₅₇₅ of 100 using 0.2mM IPTG) was centrifuged and the supernatant discarded. The cell paste was resuspended in 20 mM Tris, pH 8.2 (in a ratio of 1:4) and lysed by 10 passing through Microfluidics Microfluidizer M110Y at 15,000 psi. The lysate was centrifuged and the soluble fraction discarded. The insoluble fraction was mixed with extraction buffer (20 mM Tris, 2 M NaCl, 2 M urea, 1% Zwittergent 3-14, pH 8.2) at room temperature for 1 hour and centrifuged to produce an extract supernatant fraction and an extract pellet fraction. The extraction yield of IFN- β (IFN- β in extract supernatant fraction/ IFN- β in the initial insoluble fraction) was 15 close to 100% (>99%) based on SDS-CGE analysis (data not shown).

[00170] The extract supernatant was filtered and loaded on a 5 mL GE Healthcare Blue Sepharose column equilibrated with 20 mM Tris, 2 M NaCl, pH 8.2. The column was washed with the same buffer and the IFN- β eluted with 20 mM Tris, 2 M NaCl, 50% propylene glycol, pH 8.2. The protein in the elution pool was analyzed by SDS-CGE and found to be more than 98% pure IFN- β . 20 Aliquots of the elution pool were exchanged into Buffers C (5 mM glycine pH 3.0) and D (5 mM aspartic acid, 9% trichloroacetic acid, pH 4.0).

[00171] The exchanged samples were analyzed by SDS-CGE as well as with a cell-based assay (PBL Interferon Source, #51100-1). The cell-based assay uses a human cell line (PIL5) sensitized with IFN-type 1 receptor. IFN- β binds to the receptor, which sends a signal via the Jak1/STAT1 signal transduction pathway, activating ISG15-luciferase transcription via the Interferon Sensitive 5 Response Element (ISRE). Cell-based assay kit instructions were followed as per manufacturer's protocol (51100 rev01). The signal was read using conventional plate readers with luminescence detection. Table 14 summarizes the SDS-CGE and cell-based assay results, which indicate that the IFN- β in the samples was fully active.

10 [00172] Table 14. Results of Activity Assays

Sample	SDS-CGE (mg/L)	Cell-based assay (mg/L)
Blue-Sepharose Elution pool in Exchange Buffer A	436	477
Blue-Sepharose Elution pool in Exchange Buffer B	404	404

Example 6. Production of IFN- α 2a and 2b from High Throughput Expression Samples

[00173] IFN- α 2a and IFN- α 2b coding sequences were constructed using *P. fluorescens* preferred codons to encode for the human proteins. Figure 8 shows the amino acid and DNA sequences of the 15 synthetic IFN- α 2a gene, and Figure 9 shows the amino acid and DNA sequences of the synthetic IFN- α 2b gene.

[00174] Plasmids expressing either protein were constructed and transformed into different host strains. Expression strains were tested for their ability to express recombinant protein using HTP analysis, as described with regard to IFN- β herein. A subset of the expression strains are selected 20 for fermentation studies.

[00175] The selected strains were grown and induced according to the present invention. The cells were centrifuged, lysed, and centrifuged again as described herein for IFN- β . The resulting insoluble fraction and first soluble fraction were extracted using extraction conditions described herein. The resulting IFN- α 2a and IFN- α 2b extract supernatants were quantitated using SDS-CGE 25 (data not shown).

Example 7. Extraction of IFN- α 2a and 2b from High Throughput Expression Material

The first insoluble fraction obtained as described in Example 6 is extracted using the extraction conditions of the present invention. IFN- α 2a and 2b in the resulting second soluble fractions are evaluated by CGE and bioactivity assay.

30 **Example 8. Production of IFN- α 2a and 2b from Large Scale Fermentation**

IFN- α 2a and 2b expressing strains selected by HTP analysis are grown in 2 liter fermentors using optimized fermentation conditions of the present invention, e.g., as described in Example 4. The first insoluble fraction is extracted using the methods of the present invention, e.g., as described in Example 4. The IFN- α 2a and 2b protein present in the first insoluble and second soluble fractions are evaluated by CGE and compared.

Example 9. Analysis of IFN- α 2a and 2b Extraction Product

The extraction product obtained in Example 8 is analyzed for IFN- α 2a and 2b bioactivity.

[00176a] In the claims which follow and in the preceding description of the invention, except where the context requires otherwise due to express language or necessary implication, the word “comprise” or variations such as “comprises” or “comprising” is used in an inclusive sense, i.e. to specify the presence of the stated features but not to preclude the presence or addition of further features in various embodiments of the invention.

CLAIMS

1. A method for producing a recombinant Type 1 interferon protein, said method comprising:
expressing the recombinant interferon protein by culturing a *Pseudomonas* or *E. coli* host cell
containing an expression construct, said expression construct comprising a nucleic acid
comprising a coding sequence for the Type 1 interferon protein that has been optimized for
expression in the host cell;
lysing the cultured host cell;
obtaining an insoluble fraction and a soluble fraction from the lysed host cell;
extracting the insoluble fraction by subjecting it to non-denaturing extraction conditions,
wherein said non-denaturing extraction conditions comprise a Zwitterionic detergent or
octylglucopyranoside; and
obtaining an extract pellet and an extract supernatant from the extracted insoluble fraction;
wherein the recombinant protein in the extract supernatant is present in soluble form, active
form, or a combination thereof, without being further subjected to a renaturing or refolding
step.
2. The method of claim 1, wherein the coding sequence for the Type I interferon protein is
operably linked to a lac derivative promoter, wherein the culturing comprises:
growing the host cell at a temperature of about 25° C to about 33° C and at a pH of about 5.7 to
about 6.5, to an OD₆₀₀ of about 80 to about 160; and
inducing the host cell at a concentration of about 0.08 mM to about 0.4 mM IPTG.
3. The method of claim 1 or claim 2, wherein the Zwitterionic detergent is Zwittergent 3-08,
Zwittergent 3-10, Zwittergent 3-12, Zwittergent 3-14, or CHAPS.
4. The method of claim 3, wherein the non-denaturing extraction conditions comprise about 0.5%
to about 2% Zwittergent 3-14.
5. The method of any one of claims 1 to 4, wherein the non-denaturing extraction conditions
further comprise a chaotropic agent and a cosmotropic salt.
6. The method of claim 5, wherein the chaotropic agent is urea or guanidinium hydrochloride, and
wherein the cosmotropic salt is NaCl, KCl, or (NH₄)₂SO₄.

7. The method of any one of claims 4 to 6, wherein the non-denaturing extraction conditions comprise: about 0.5 to about 2% Zwittergent 3-14; about 0 to about 2 M urea; about 0 to about 2 M NaCl; and wherein the pH is about 6.5 to about 8.5.
- 5 8. The method of claim 7, wherein the non-denaturing extraction conditions comprise: about 1% Zwittergent 3-14; about 2 M urea; about 2 M NaCl; and wherein the pH is about 8.2.
9. The method of any one of claims 1 to 8, wherein the non-denaturing extraction conditions additionally comprise about 1% to about 40% w/v solids.
- 10 10. The method of claim 9, wherein the non-denaturing extraction conditions additionally comprise about 5% w/v solids.
11. The method of any one of claims 1 to 10, wherein the recombinant Type 1 interferon protein is an interferon- β , an interferon- α , an interferon- κ , or an interferon- ω .
12. The method of claim 11, wherein the recombinant Type 1 interferon protein is an interferon- β , and wherein said interferon- β is selected from the group consisting of: a human interferon- β 1b and human interferon- β 1b C17S, or wherein the recombinant Type 1 interferon protein is an interferon- α , and wherein the interferon- α is selected from the group consisting of: human interferon- α 2a and human interferon- α 2b.
13. The method of any one of claims 1 to 12, further comprising measuring the amount of recombinant Type 1 interferon protein in a sample of the insoluble fraction, said sample of the insoluble fraction obtained before extracting the insoluble fraction, and measuring the amount of recombinant Type 1 interferon protein in a sample of the extract supernatant, said sample of the extract supernatant obtained after extracting the insoluble fraction, wherein the amount of recombinant interferon protein detected in the sample of the extract supernatant obtained after extracting the insoluble fraction is about 10% to about 95% of the amount of the recombinant interferon protein measured in the insoluble fraction sample obtained before extracting the insoluble fraction.
14. The method of any one of claims 1 to 13, further comprising measuring the activity of the recombinant interferon protein in a sample of the extract supernatant, said sample of the extract supernatant obtained after extracting the insoluble fraction, wherein about 40% to about 100%

of the recombinant interferon protein present in the extract supernatant sample obtained after extracting the insoluble fraction is determined to be active, or wherein about 75% to about 100% of the recombinant interferon protein present in the extract supernatant sample obtained after extracting the insoluble fraction is determined to be active when compared with the amount of active interferon protein in a standard sample containing the same amount of recombinant interferon protein as the extract supernatant sample obtained after extracting the insoluble fraction.

- 5 15. The method of claim 14, wherein the recombinant protein is an interferon- β , and wherein the amount of active recombinant protein is determined by Blue Sepharose affinity column chromatography, receptor binding assay, antiviral activity assay, or cytopathic effect assay.
- 10 16. The method of any one of claims 1 to 15, wherein the recombinant Type 1 interferon protein in the extract supernatant is present at a concentration of about 0.3 grams per liter to about 10 grams per liter.
- 15 17. The method of any one of claims 1 to 16, wherein the host cell is a lon hslUV protease deletion strain.
- 20 18. The method of any one of claims 1 to 17, wherein the Type 1 interferon is expressed in the cytoplasm of the host cell.
- 25 19. The method of claim 12, wherein the Type 1 interferon is human interferon- β 1b or human interferon- β 1b C17S, and wherein the human interferon- β 1b or human interferon- β 1b C17S is expressed in the cytoplasm of the host cell.
- 30 20. The method of any one of claims 1 to 13, further comprising measuring the activity of the recombinant interferon protein in a sample of the extract supernatant, said sample of the extract supernatant obtained after extracting the insoluble fraction, wherein about 80% to about 100%, about 85% to about 100%, or about 90% to about 100%, of the recombinant interferon protein in the extract supernatant is determined to be active.
21. The method of claim 20, wherein the host cell is a lon hslUV protease deletion strain.

22. The method of claim 20 or claim 21, wherein the Type 1 interferon is human interferon- β 1b or human interferon- β 1b C17S, wherein the human interferon- β 1b or human interferon- β 1b C17S is expressed in the cytoplasm of the host cell.
- 5 23. A recombinant Type 1 interferon protein produced by the method of any one of claims 1 to 22.
24. The method of claim 1 or the recombinant Type 1 interferon protein of claim 23, substantially as hereinbefore described with reference to the Examples and/or Figures.


10

WO 2011/109556

PCT/US2011/026921

1/9

FIGURE 1



REPLACEMENT SHEET

AMENDED SHEET (ART. 34)

FIGURE 2





FIGURE 3

FIGURE 4**A**

Term	Estimate	Std Error	t Ratio	Prob> t
Z314	3720.4167	489.4339	7.60	<.0001*
Solids	-92.575	29.36603	-3.15	0.0103*
Urea	615.9375	220.2452	2.80	0.0189*
(Solids-12.5)*(Z314-0.55)	-127.5	65.25785	-1.95	0.0792
NaCl	500.51471	259.112	1.93	0.0822
(NaCl-1)*(Z314-0.55)	1008.3333	575.8046	1.75	0.1105
pH	353.1875	220.2452	1.60	0.1399
(Solids-12.5)*(NaCl-1)	-30.7549	34.54827	-0.89	0.3943


B


FIGURE 5


FIGURE 6

FIGURE 7**A**

MSYNLLGFLQRSSNFQSQKLLWQLNGRLEYCLKDRMNFDIPEEIKQLQQF
QKEDAALTIYEMLQNIFAIFRQDSSSTGWNETIVENLLANVYHQINHLKTV
LEEKLEKEDFTRGKLMSSLHLKRYYGRILHYLKAKEYSHCAWTIVRVEILR
NFYFINRLTGYLRN (SEQ ID NO: 1)

B

Atgtcgtacaacctgttgggcttctgcagcggcctccaacttcaatgcagaagctgtggcaattgaatggcg
cctggaatactgcctgaaggaccgcataactcgacatccctgaagagattaagcaactccagcagttccagaaaga
ggatgcagctgtcgcataatgtcgcagaacatcttcgcgcatttcgcaggacagcagcagcaccgggttgc
gaacgaaaccattgtcgcagaatctgtggccacgttatcaccagattaaccacctaagactgtgtggaaagagaa
gttggagaaagaagatttcacgcgtggcaagttgatgagttcgctgcatttgcatttgcattatggcgtatcctgcatta
cctgaaggccaaagaatacagccactgtgcgtggaccatgttcgcgtggagatctgcgcatttgcattatcatcaat
cggtcaccgggttacccgcact (SEQ ID NO: 2)

C

SYNLLGFLQRSSNFQSQKLLWQLNGRLEYCLKDRMNFDIPEEIKQLQQFQ
KEDAALTIYEMLQNIFAIFRQDSSSTGWNETIVENLLANVYHQINHLKTVL
EEKLEKEDFTRGKLMSSLHLKRYYGRILHYLKAKEYSHCAWTIVRVEILR
NFYFINRLTGYLRN (SEQ ID NO: 3)

FIGURE 8**A**

1	C	D	L	P	Q	T	H	S	L	G	S	R	R	T	L	M	L	L	A	Q
21	M	R	K	I	S	L	F	S	C	L	K	D	R	H	D	F	G	F	P	Q
41	E	E	F	G	N	Q	F	Q	K	A	E	T	I	P	V	L	H	E	M	I
61	Q	Q	I	F	N	L	F	S	T	K	D	S	S	A	A	W	D	E	T	L
81	L	D	K	F	Y	T	E	L	Y	Q	Q	L	N	D	L	E	A	C	V	I
101	Q	G	V	G	V	T	E	T	P	L	M	K	E	D	S	I	L	A	V	R
121	K	Y	F	Q	R	I	T	L	Y	L	K	E	K	K	Y	S	P	C	A	W
141	E	V	V	R	A	E	I	M	R	S	F	S	L	S	T	N	L	Q	E	S
161	L	R	S	K	E	(SEQ ID NO: 4)														

B

1	TGTGACCTGCCTCAGACTCACTCCCTCGTAGCCGCCGGACCCCTGATGCTGTTGGCGCAG
61	ATGCGTAAGATCTCCCTGTTCTCGTGCCTGAAAGACCGCCATGATTCGGCTTCCCGCAG
121	GAAGAATTGGAAACCAGTTTCAGAAGGCTGAAACCATCCCAGTGCTGCACGAGATGATC
181	CAGCAAATTTCAACCTGTTCAGCACCAAGGACAGCTCGGCCGCCTGGGACGAAACGTTG
241	TTGGACAAATTTACACCGAGCTGTACCAACAATGAACGATCTGGAAGCATGCGTTATT
301	CAAGGCCTGGCGTCACCGAAACGCCGCTGATGAAAGAAGATAGCATCCTGGCCGTGCGT
361	AAGTACTTTCAGCGCATCACCTCTACCTGAAAGAGAAGAAGTATTGCCCTGCGCGTGG
421	GAGGTCGTCCCGCGCCGAGATCATGCGGTCTTCAGCCTCTCCACCAATCTGCAGGAAAGT
481	CTCCGCTCGAAAGAA (SEQ ID NO: 5)

FIGURE 9**A**

1	C	D	L	P	Q	T	H	S	L	G	S	R	R	T	L	M	L	L	A	Q
21	M	R	R	I	S	L	F	S	C	L	K	D	R	H	D	F	G	F	P	Q
41	E	E	F	G	N	Q	F	Q	K	A	E	T	I	P	V	L	H	E	M	I
61	Q	Q	I	F	N	L	F	S	T	K	D	S	S	A	A	W	D	E	T	L
81	L	D	K	F	Y	T	E	L	Y	Q	Q	L	N	D	L	E	A	C	V	I
101	Q	G	V	G	V	T	E	T	P	L	M	K	E	D	S	I	L	A	V	R
121	K	Y	F	Q	R	I	T	L	Y	L	K	E	K	K	Y	S	P	C	A	W
141	E	V	V	R	A	E	I	M	R	S	F	S	L	S	T	N	L	Q	E	S
161	L	R	S	K	E	(SEQ ID NO: 6)														

B

1	TGTGACCTGCCTCAGACTCACTCCCTCGGTAGCCGCCGGACCCTGATGCTGTTGGCGCAG
61	ATGCGTCGCATCTCCCTGTTCTCGTCGCTGAAAGACCGCCATGATTCGGCTTCCCGCAG
121	GAAGAATTGGAAACCAGTTTCAGAAGGCTGAAACCATCCCAGTGCTGCACGAGATGATC
181	CAGCAAATTTCACCTGTTCAGCACCAAGGACAGCTCGGCCCTGGGACGAAACGTTG
241	TTGGACAAATTTCACCGAGCTGTACCAACAACGATCTGGAAGCATGCGTTATT
301	CAAGGCGTGGCGTCACCGAAACGCCGCTGATGAAAGAAGATAGCATCCTGCCGTGCGT
361	AAGTACTTTCAGCGCATCACCTCTACCTGAAAGAGAAGAAGTATTGCCCTGCGCGTGG
421	GAGGTCGTCCCGCCGAGATCATGCGGTCTTCAGCCTCTCCACCAATCTGCAGGAAAGT
481	CTCCGCTCGAAAGAA (SEQ ID NO: 7)

2011223627 11 Apr 2013

Sequence_Listing
SEQUENCE LISTING

<110> PFENEX INC.
<120> METHOD FOR PRODUCING SOLUBLE RECOMBINANT INTERFERON PROTEIN
WITHOUT DENATURING
<130> 38194-704.601
<140>
<141>
<150> 61/310,671
<151> 2010-03-04
<160> 7
<170> PatentIn version 3.5
<210> 1
<211> 166
<212> PRT
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: Synthetic
polypeptide
<400> 1
Met Ser Tyr Asn Leu Leu Gly Phe Leu Gln Arg Ser Ser Asn Phe Gln
1 5 10 15

Ser Gln Lys Leu Leu Trp Gln Leu Asn Gly Arg Leu Glu Tyr Cys Leu
20 25 30

Lys Asp Arg Met Asn Phe Asp Ile Pro Glu Glu Ile Lys Gln Leu Gln
35 40 45

Gln Phe Gln Lys Glu Asp Ala Ala Leu Thr Ile Tyr Glu Met Leu Gln
50 55 60

Asn Ile Phe Ala Ile Phe Arg Gln Asp Ser Ser Ser Thr Gly Trp Asn
65 70 75 80

Glu Thr Ile Val Glu Asn Leu Leu Ala Asn Val Tyr His Gln Ile Asn
85 90 95

His Leu Lys Thr Val Leu Glu Glu Lys Leu Glu Lys Glu Asp Phe Thr
100 105 110

Arg Gly Lys Leu Met Ser Ser Leu His Leu Lys Arg Tyr Tyr Gly Arg
115 120 125

Ile Leu His Tyr Leu Lys Ala Lys Glu Tyr Ser His Cys Ala Trp Thr
130 135 140

Ile Val Arg Val Glu Ile Leu Arg Asn Phe Tyr Phe Ile Asn Arg Leu
145 150 155 160

2011223627 11 Apr 2013

Sequence_Listing

Thr Gly Tyr Leu Arg Asn
165

<210> 2
<211> 499
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial sequence: Synthetic
polynucleotide

<400> 2
atgtcgata acctgttggg cttcctgcag cggcctcca actttcaatc gcagaagctg 60
ctgtggcaat tgaatggtcg cctggaatac tgcctgaagg accgcataa cttcgacatc 120
cctgaagaga ttaagcaact ccagcagttc cagaaagagg atgcagctt gacgatctat 180
gaaatgctgc agaacatctt cgcgatctt cgccaggaca gcagcagcac cggttggAAC 240
gaaaccattt tcgagaatct gctggcaac gtctatcacc agattaacca cctcaagact 300
gtgcttggaa agaagtttggaa gaaagaagat ttcacgcgtg gcaagtttgc gagttcgctg 360
catttggaaac gctactatgg tcgtatcctg cattacctga aggccaaaga atacagccac 420
tgtgcgttggaa ccatcggttcg cgtggagatc ctgcgcaact tctacttcat caatcggttc 480
accggttacc tccgcaact 499

<210> 3
<211> 165
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial sequence: Synthetic
polypeptide

<400> 3
Ser Tyr Asn Leu Leu Gly Phe Leu Gln Arg Ser Ser Asn Phe Gln Ser
1 5 10 15

Gln Lys Leu Leu Trp Gln Leu Asn Gly Arg Leu Glu Tyr Cys Leu Lys
20 25 30

Asp Arg Met Asn Phe Asp Ile Pro Glu Glu Ile Lys Gln Leu Gln Gln
35 40 45

Phe Gln Lys Glu Asp Ala Ala Leu Thr Ile Tyr Glu Met Leu Gln Asn
50 55 60

Ile Phe Ala Ile Phe Arg Gln Asp Ser Ser Ser Thr Gly Trp Asn Glu
65 70 75 80

Thr Ile Val Glu Asn Leu Leu Ala Asn Val Tyr His Gln Ile Asn His
85 90 95

2011223627 11 Apr 2013

Sequence_Listing

Leu Lys Thr Val Leu Glu Glu Lys Leu Glu Lys Glu Asp Phe Thr Arg
100 105 110

Gly Lys Leu Met Ser Ser Leu His Leu Lys Arg Tyr Tyr Gly Arg Ile
115 120 125

Leu His Tyr Leu Lys Ala Lys Glu Tyr Ser His Cys Ala Trp Thr Ile
130 135 140

Val Arg Val Glu Ile Leu Arg Asn Phe Tyr Phe Ile Asn Arg Leu Thr
145 150 155 160

Gly Tyr Leu Arg Asn
165

<210> 4

<211> 165

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
polypeptide

<400> 4

Cys Asp Leu Pro Gln Thr His Ser Leu Gly Ser Arg Arg Thr Leu Met
1 5 10 15

Leu Leu Ala Gln Met Arg Lys Ile Ser Leu Phe Ser Cys Leu Lys Asp
20 25 30

Arg His Asp Phe Gly Phe Pro Gln Glu Glu Phe Gly Asn Gln Phe Gln
35 40 45

Lys Ala Glu Thr Ile Pro Val Leu His Glu Met Ile Gln Gln Ile Phe
50 55 60

Asn Leu Phe Ser Thr Lys Asp Ser Ser Ala Ala Trp Asp Glu Thr Leu
65 70 75 80

Leu Asp Lys Phe Tyr Thr Glu Leu Tyr Gln Gln Leu Asn Asp Leu Glu
85 90 95

Ala Cys Val Ile Gln Gly Val Gly Val Thr Glu Thr Pro Leu Met Lys
100 105 110

Glu Asp Ser Ile Leu Ala Val Arg Lys Tyr Phe Gln Arg Ile Thr Leu
115 120 125

Tyr Leu Lys Glu Lys Lys Tyr Ser Pro Cys Ala Trp Glu Val Val Arg
130 135 140

Ala Glu Ile Met Arg Ser Phe Ser Leu Ser Thr Asn Leu Gln Glu Ser
145 150 155 160

2011223627 11 Apr 2013

Sequence_Listing

Leu Arg Ser Lys Glu
165

<210> 5
<211> 495
<212> DNA
<213> Artificial sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 5
tgtgacctgc ctcagactca ctccctcggt agccgcccga ccctgatgct gttggcgcag 60
atgcgttaaga tctccctgtt ctcgtgcctg aaagaccgcg atgatttcgg cttccgcag 120
gaagaattcg ggaaccagggt tcagaaggct gaaaccatcc cagtgcgtca cgagatgatc 180
cagcaaattt tcaacctgtt cagcaccaag gacagctcgg ccgcctggga cgaaacgttg 240
ttggacaaat tttacaccga gctgtaccaa caactgaacg atctggaagc atgcgttatt 300
caaggcgtgg gcgtcaccga aacgcccgtg atgaaagaag atagcatcct ggccgtgcgt 360
aagtactttc agcgcatcac cctctacctg aaagagaaga agtattcgcc ctgcgcgtgg 420
gaggtcgtcc gcgcccagat catgcggtcc ttcagcctct ccaccaatct gcaggaaagt 480
ctccgctcga aagaa 495

<210> 6
<211> 165
<212> PRT
<213> Artificial sequence

<220>
<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 6
Cys Asp Leu Pro Gln Thr His Ser Leu Gly Ser Arg Arg Thr Leu Met
1 5 10 15

Leu Leu Ala Gln Met Arg Arg Ile Ser Leu Phe Ser Cys Leu Lys Asp
20 25 30

Arg His Asp Phe Gly Phe Pro Gln Glu Glu Phe Gly Asn Gln Phe Gln
35 40 45

Lys Ala Glu Thr Ile Pro Val Leu His Glu Met Ile Gln Gln Ile Phe
50 55 60

Asn Leu Phe Ser Thr Lys Asp Ser Ser Ala Ala Trp Asp Glu Thr Leu
65 70 75 80

Leu Asp Lys Phe Tyr Thr Glu Leu Tyr Gln Gln Leu Asn Asp Leu Glu
85 90 95

Sequence_Listing

Ala Cys Val Ile Gln Gly Val Gly Val Thr Glu Thr Pro Leu Met Lys
100 105 110

Glu Asp Ser Ile Leu Ala Val Arg Lys Tyr Phe Gln Arg Ile Thr Leu
115 120 125

Tyr Leu Lys Glu Lys Lys Tyr Ser Pro Cys Ala Trp Glu Val Val Arg
130 135 140

Ala Glu Ile Met Arg Ser Phe Ser Leu Ser Thr Asn Leu Gln Glu Ser
145 150 155 160

Leu Arg Ser Lys Glu
165

<210> 7

<211> 495

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial sequence: Synthetic
polynucleotide

<400> 7

tgtgacctgc ctcagactca ctccctcggt agccgcccga ccctgatgct gttggcgcag	60
atgcgtcgca tctccctgtt ctcgtgcctg aaagaccgcc atgatttcggtt cttcccgccag	120
gaagaattcg ggaaccaggtt tcagaaggct gaaaccatcc cagtgctgca cgagatgatc	180
cagcaaattt tcaacctgtt cagcaccaag gacagctcgg ccgcctggga cgaaacgttgc	240
ttggacaaat tttacaccga gctgtaccaa caactgaacg atctggaagc atgcgttatt	300
caaggcgtgg gcgtcaccga aacgcccgtg atgaaagaag atagcatcct ggccgtgcgt	360
aagtactttc agcgcatcac cctctacctg aaagagaaga agtattcgcc ctgcgcgtgg	420
gaggtcgtcc gcgccgagat catgcgggtcc ttcagcctct ccaccaatct gcaggaaagt	480
ctccgctcga aagaa	495