



US010647731B2

(12) **United States Patent**
Jiang et al.

(10) **Patent No.:** **US 10,647,731 B2**
(45) **Date of Patent:** **May 12, 2020**

(54) **THIAZOLOPYRIMIDINONES AND METHODS OF USE THEREOF**

(71) Applicant: **Genentech, Inc.**, South San Francisco, CA (US)

(72) Inventors: **Yu Jiang**, Beijing (CN); **Guosheng Wu**, Beijing (CN); **Po-Wai Yuen**, Beijing (CN); **Elisia Villemure**, San Francisco, CA (US); **Jacob Schwarz**, San Ramon, CA (US); **Cuong Ly**, Burlingame, CA (US); **Benjamin Sellers**, Larkspur, CA (US); **Matthew Volgraf**, Oakland, CA (US)

(73) Assignee: **Genentech, Inc.**, South San Francisco, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/962,333**

(22) Filed: **Apr. 25, 2018**

(65) **Prior Publication Data**

US 2018/0244694 A1 Aug. 30, 2018

Related U.S. Application Data

(60) Division of application No. 15/094,687, filed on Apr. 8, 2016, now Pat. No. 9,988,400, which is a continuation of application No. PCT/EP2014/071522, filed on Oct. 8, 2014.

(30) **Foreign Application Priority Data**

Oct. 11, 2013 (WO) PCT/CN2013/085031
Sep. 5, 2014 (WO) PCT/CN2014/085959

(51) **Int. Cl.**

C07D 513/14 (2006.01)
C07D 513/04 (2006.01)
C07D 513/10 (2006.01)
C07D 519/00 (2006.01)

(52) **U.S. Cl.**

CPC **C07D 513/14** (2013.01); **C07D 513/04** (2013.01); **C07D 513/10** (2013.01); **C07D 519/00** (2013.01)

(58) **Field of Classification Search**

CPC .. **C07D 513/14**; **C07D 513/04**; **C07D 513/10**;
C07D 519/00
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,958,931 A 9/1999 Adam et al.
5,902,815 A 11/1999 Olney et al.
2007/0155779 A1 5/2007 Verhoest et al.

FOREIGN PATENT DOCUMENTS

WO 2004/074270 A2 9/2004
WO 2014/202493 A1 12/2014
WO 2016/166078 10/2016

OTHER PUBLICATIONS

Agamenzone et al., "Identification of small molecules acting against H1N1 influenza A virus" *Virology* 488:249-258 (Dec. 3, 2015).
Costa et al., "A Novel Family of Negative and Positive Allosteric Modulators of NMDA Receptors" *Journal of Pharmacology and Experimental Therapeutics* 335(3):614-621 (Dec. 1, 2010).
Ferrarini et al., "Study on Affinity Profile toward Native Human and Bovine Adenosine Receptors of a Series of 1,8-Naphthyridine Derivatives" *Journal of Medicinal Chemistry* 47(12):3019-3031 (Jan. 1, 2004).
Guo Chuangxing et al., "Discovery of 2-((1H-benzo[d]imidazol-1-yl)methyl)-4H-pyrido[1,2-a]pyrimidin-4-ones as novel PKM2 activators" *Bioorganic & Medicinal Chemistry Letters* 23(11):3358-3363 (Apr. 1, 2013).
Hackos et al., "Positive Allosteric Modulators of GluN2A-Containing NMDARs with Distinct Modes of Action and Impacts on Circuit Function" *Neuron* 89:983-999 (2016).
Kennis et al., "New substituted 1,2,3,4-tetrahydrobenzofuro[3,2-c]pyridine having highly active and potent central α_2 -antagonistic activity as potential antidepressants" *Bioorganic & Medicinal Chemistry Letters* 10:71-74 (Jan. 1, 2000).
Molnár et al., "Suzuki-Miyaura cross-coupling reactions of halo derivatives of 4H-pyrido[1,2-a]pyrimidin-4-ones" *Organic & Biomolecular Chemistry* 9(19):6559 (Jan. 1, 2011).
Villemure et al., "GluN2A-Selective Pyridopyrimidinone Series of NMDAR Positive Allosteric Modulators with an Improved in Vivo Profile" *ACS Medicinal Chemistry Letters* 8:84-89 (2017).
Volgraf et al., "Discovery of GluN2A-Selective NMDA Receptor Positive Allosteric Modulators (PAMs): Tuning Deactivation Kinetics via Structure-Based Design" *Journal of Medicinal Chemistry* 59:2760-2779 (2016).

Primary Examiner — Zohreh A Fay

(74) *Attorney, Agent, or Firm* — Brian L. Buckwalter

(57) **ABSTRACT**

The present invention relates to certain thiazolopyrimidinone compounds, pharmaceutical compositions comprising such compounds, and methods of treatment using such compounds.

70 Claims, No Drawings

THIAZOLOPYRIMIDINONES AND METHODS OF USE THEREOF

RELATED APPLICATIONS

This application is a divisional of U.S. application Ser. No. 15/094,687, filed Apr. 8, 2016, which is a continuation of International Application No. PCT/EP2014/071522, filed Oct. 8, 2014 claiming priority to International Application No. PCT/CN2013/085031, filed Oct. 11, 2013 and International Application No. PCT/CN2014/085959, filed Sep. 5, 2014, each of which are incorporated herein by reference in its entirety.

FIELD OF INVENTION

The present invention relates to certain thiazolopyrimidinone compounds, pharmaceutical compositions comprising such compounds, and methods of treating neurological and psychiatric conditions, and other diseases and medical conditions, with such compounds and pharmaceutical compositions. The present invention also relates to certain thiazolopyrimidinone compounds for use in modulating NMDA receptor activity.

BACKGROUND OF THE INVENTION

N-Methyl-D-aspartate (NMDA) receptors play an important role in various central nervous system functions, such as synaptic transmission and synaptic plasticity, and underlying functions such as regulation of long-term potentiation, long-term depression, and experience, dependent synaptic refinement. Costa et al., "A Novel Family of Negative and Positive Allosteric Modulators of NMDA Receptors." *J. Pharmacol. Exp. Ther.* 2010, 335, 614-621, at 614. Excitatory nerve transmission in these receptors is regulated by the neurotransmitter, L-glutamate, and the agonist, NMDA. PCT Intl. Publ. No. WO2007/006175, paras. 2-3. NMDA receptors are ligand-gated ion channels comprising seven subunits: GluN1, GluN2A-D, and GluN3A-B. Costa at 615. The NR2A and NR2B subunits have been implicated in glutamate binding to the receptor, while the NR1 subunit may play a role in the binding of the receptor co-agonist, glycine. The three-dimensional structures of the glutamate- and glycine-binding pockets of NMDA receptors have been characterized, allowing for design of more subtype-specific modulators.


Modulation of these receptors effects changes in learning and memory, and modulators of NMDA receptor activity are considered as potential treatments for neurological and psychiatric conditions including pain, neuropathic pain, inflammatory pain, peripheral neuropathy, stroke, epilepsy, neurodegeneration, schizophrenia, drug addiction, mood disorders, post-traumatic stress disorder, seizures, convulsions, age-associated memory impairment, and depression. Costa at 614. Modulation of NMDA receptor activity is linked with a neuroprotective role, with applications in treatments for stroke, traumatic brain injury, ischemia, and neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Creutzfeldt-Jakob disease. Costa at 614-615.

There is a particular need for NMDA receptor modulators that demonstrate subtype selectivity among members of the NMDA receptor family. Selective agents will allow for optimal therapeutic activity with a reduced potential for adverse side effects. Costa at 615.

There remains a need for potent NMDA receptor modulators with desirable pharmaceutical properties. Certain thiazolopyrimidinone derivatives have been found in the context of this invention to have NMDA receptor-modulating activity.

SUMMARY OF THE INVENTION

In one aspect, the invention is directed to a compound of Formula II:



wherein

R^a is C_{1-6} alkyl or C_{2-6} alkenyl, each optionally substituted with one or more R^b substituents; C_{2-6} alkynyl; halo; $-C(O)R^c$; $-NR^dR^e$; $-C(O)NR^dR^e$; $-C(S)NR^dR^e$; $-C(=N-OH)-C_{1-4}$ alkyl; $-OC_{1-4}$ alkyl; $-OC_{1-4}$ haloalkyl; $-SC_{1-4}$ alkyl; $-SO_2C_{1-4}$ alkyl; cyano; C_{3-6} cycloalkyl optionally substituted with one or more R^f substituents; or a phenyl, monocyclic heteroaryl, or heterocycloalkyl ring, each ring optionally substituted with one or more R^g substituents;

wherein each R^b substituent is independently selected from the group consisting of $-OH$, $-C_{1-4}$ alkoxy, $-NR^dR^e$, $-C(O)NR^dR^e$, $-SC_{1-4}$ alkyl, $-SO_2C_{1-4}$ alkyl, cyano, halo, C_{3-6} cycloalkyl, and monocyclic heteroaryl;

R^c is C_{1-4} alkyl, $-C_{1-4}$ haloalkyl, C_{3-6} cycloalkyl, or a monocyclic, carbon-linked heterocycloalkyl;

R^d is H or C_{1-4} alkyl;

R^e is H; C_{1-4} alkyl optionally substituted with $-CN$, $-CF_3$, $-OH$, or a monocyclic heterocycloalkyl; C_{3-6} cycloalkyl; $-OH$; or $-OC_{1-4}$ alkoxy;

or R^d and R^e taken together with the nitrogen to which they are attached form a heterocycloalkyl, optionally substituted with C_{1-4} alkyl or $-OH$;

each R^f substituent is independently selected from the group consisting of: C_{1-4} alkyl optionally substituted with $-OH$, cyano, or C_{1-4} alkoxy; $-OH$; halo; C_{1-4} haloalkyl; $-CONH_2$; and cyano; and

each R^g substituent is independently selected from the group consisting of C_{1-4} alkyl, $-CF_3$, halo, $-NH_2$, $-OCH_3$, cyano, and $-OH$;

R^1 is selected from the group consisting of H, C_{1-6} alkyl, C_{1-4} haloalkyl, C_{3-6} cycloalkyl, halo, $-OC_{1-4}$ alkyl, $-OC_{1-4}$ haloalkyl, cyano, and $-C(O)C_{1-4}$ alkyl; or R^a and R^1 taken together with the carbons to which they are attached form a 5- to 7-membered ring, optionally containing an O or NH, and optionally substituted with one or more R^h substituents;

wherein each R^h substituent is independently $-C(O)NR^jR^k$, cyano, or is C_{1-4} alkyl optionally substituted with $-OH$, $-OCH_3$, cyano, or $-C(O)NR^jR^k$; or two R^h groups attached to the same carbon and taken together with the carbon to which they are attached form a carbonyl or a C_{3-6} cycloalkyl;

3

wherein R^i and R^j are each independently H or C_{1-4} alkyl;

R^2 is $-R^m$, $-OR^m$, or $-NR^mR^n$;

wherein R^m is aryl or heteroaryl, each optionally substituted with one or more R^s substituents;

wherein each R^s substituent is independently selected from the group consisting of C_{1-4} alkyl, C_{2-4} alkenyl (optionally substituted with halo), C_{2-4} alkynyl, C_{1-4} haloalkyl, C_{1-4} alkoxy, C_{1-4} alkyl-OH, C_{1-4} haloalkoxy, halo, cyano, C_{3-6} cycloalkyl (optionally substituted with $-OH$ or halo), monocyclic heteroaryl, $-NH_2$, $-NO_2$, $-NHSO_2C_{1-4}$ alkyl, and $-SO_2C_{1-4}$ alkyl;

R^n is H, C_{1-4} haloalkyl, or C_{1-4} alkyl optionally substituted with $-OH$ or C_{1-4} alkoxy;


or R^m and R^n taken together with the nitrogen to which they are attached form a pyrrolidine or piperidine ring, optionally substituted with C_{1-4} alkyl and optionally fused to phenyl, wherein said phenyl is optionally substituted with halo;

R^3 is H or methyl; and

R^4 is H or fluoro;

or a pharmaceutically acceptable salt thereof.

In one aspect, the invention is directed to a compound of Formula I:



wherein

R^a is C_{1-6} alkyl optionally substituted with one or more R^b substituents; C_{2-6} alkenyl; C_{2-6} alkynyl; halo; $-C(O)R^c$; $-NR^dR^e$; $-C(O)NR^dR^e$; $-C(S)NR^dR^e$; $-C(=N-OH)-C_{1-4}$ alkyl; $-SO_2C_{1-4}$ alkyl; cyano; C_{3-6} cycloalkyl optionally substituted with one or more R^f substituents; or a phenyl, monocyclic heteroaryl, or heterocycloalkyl ring, each ring optionally substituted with one or more R^g substituents;

wherein each R^b substituent is independently selected from the group consisting of $-OH$, $-C_{1-4}$ alkoxy, $-NR^dR^e$, $-C(O)NR^dR^e$, $-SC_{1-4}$ alkyl, $-SO_2C_{1-4}$ alkyl, cyano, halo, and monocyclic heteroaryl;

R^c is C_{1-4} alkyl, $-C_{1-4}$ haloalkyl, C_{3-6} cycloalkyl, or a monocyclic, carbon-linked heterocycloalkyl;

R^d is H or C_{1-4} alkyl;

R^e is H; C_{1-4} alkyl optionally substituted with $-CN$, $-CF_3$, $-OH$, or a monocyclic heterocycloalkyl; C_{3-6} cycloalkyl; $-OH$; or $-OC_{1-4}$ alkoxy;

or R^d and R^e taken together with the nitrogen to which they are attached form a heterocycloalkyl, optionally substituted with C_{1-4} alkyl or $-OH$;

each R^f substituent is independently selected from the group consisting of: C_{1-4} alkyl optionally substituted with $-OH$, cyano, or C_{1-4} alkoxy; C_{1-4} haloalkyl; $-CONH_2$; and cyano; and

each R^g substituent is independently selected from the group consisting of C_{1-4} alkyl, $-CF_3$, halo, $-NH_2$, $-OCH_3$, cyano, and $-OH$;

4

R^1 is selected from the group consisting of H, C_{1-6} alkyl, C_{1-4} haloalkyl, and C_{3-6} cycloalkyl; or R^a and R^1 taken together with the carbons to which they are attached form a 5- to 7-membered ring, optionally containing an O or NH, and optionally substituted with one or more R^h substituents;

wherein each R^h substituent is independently $-C(O)NR^iR^j$, cyano, or is C_{1-4} alkyl optionally substituted with $-OH$, $-OCH_3$, cyano, or $-C(O)NR^iR^j$; or two R^h groups attached to the same carbon and taken together with the carbon to which they are attached form a carbonyl or a C_{3-6} cycloalkyl;

wherein R^i and R^j are each independently H or C_{1-4} alkyl;

R^2 is $-R^m$, $-OR^m$, or $-NR^mR^n$;

wherein R^m is aryl or heteroaryl, each optionally substituted with one or more R^s substituents;

wherein each R^s substituent is independently selected from the group consisting of C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy, C_{1-4} alkyl-OH, C_{1-4} haloalkoxy, halo, cyano, C_{3-6} cycloalkyl, $-NHSO_2C_{1-4}$ alkyl, and $-SO_2C_{1-4}$ alkyl;

R^n is H, C_{1-4} haloalkyl, or C_{1-4} alkyl optionally substituted with $-OH$ or C_{1-4} alkoxy;

or R^m and R^n taken together with the nitrogen to which they are attached form a pyrrolidine or piperidine ring, optionally substituted with C_{1-4} alkyl and optionally fused to phenyl, wherein said phenyl is optionally substituted with halo;

R^3 is H or methyl; and

R^4 is H or fluoro;

or a pharmaceutically acceptable salt thereof.

In a further aspect, the invention relates to pharmaceutical compositions each comprising an effective amount of at least one compound of Formula I or II or a pharmaceutically acceptable salt of a compound of Formula I or II. Pharmaceutical compositions according to the invention may further comprise at least one pharmaceutically acceptable excipient.

In another aspect, the invention is directed to a method of treating a subject suffering from a disease or medical condition mediated by NMDA receptor activity, comprising administering to the subject in need of such treatment an effective amount of at least one compound of Formula I or II or a pharmaceutically acceptable salt of a compound of Formula I or II, or comprising administering to the subject in need of such treatment an effective amount of a pharmaceutical composition comprising an effective amount of at least one compound of Formula I or II or a pharmaceutically acceptable salt of a compound of Formula I or II.

An aspect of the present invention concerns the use of compound of Formula I or II, or a pharmaceutically acceptable salt thereof, for the preparation of a medicament used in the treatment, prevention, inhibition, or elimination of a disease or medical condition mediated by NMDA receptor activity.

An aspect of the present invention concerns the use of a compound of Formula I or II, or a pharmaceutically acceptable salt thereof, for the preparation of a medicament used in the treatment, prevention, inhibition or elimination of a disease or medical condition mediated by NMDA receptor activity.

In another aspect, the compounds of Formula I or II, and pharmaceutically acceptable salts thereof, are useful as NMDA receptor modulators. Thus, the invention is directed to a method for modulating NMDA receptor activity, including when the NMDA receptor is in a subject, comprising exposing the NMDA receptor to an effective amount of at

least one compound of Formula I or II, or a pharmaceutically acceptable salt of a compound of Formula I or II.

In yet another aspect, the present invention is directed to methods of making compounds of Formula I or II, and pharmaceutically acceptable salts thereof.

In certain embodiments of the compounds, pharmaceutical compositions, and methods of the invention, the compound of Formula I or II is a compound selected from those species described or exemplified in the detailed description below, or is a pharmaceutically acceptable salt of such a compound.

Additional embodiments, features, and advantages of the invention will be apparent from the following detailed description and through practice of the invention.

DETAILED DESCRIPTION AND PREFERRED EMBODIMENTS

For the sake of brevity, the disclosures of the publications cited in this specification, including patents and patent applications, are herein incorporated by reference in their entirety.

Most chemical names were generated using IUPAC nomenclature herein. Some chemical names were generated using different nomenclatures or alternative or commercial names known in the art. In the case of conflict between names and structures, the structures prevail.

General Definitions

As used above, and throughout this disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings. If a definition is missing, the conventional definition as known to one skilled in the art controls. If a definition provided herein conflicts or is different from a definition provided in any cited publication, the definition provided herein controls.

As used herein, the terms "including," "containing," and "comprising" are used in their open, non-limiting sense.

As used herein, the singular forms "a," "an," and "the" include plural referents unless the context clearly dictates otherwise.

To provide a more concise description, some of the quantitative expressions given herein are not qualified with the term "about." It is understood that, whether the term "about" is used explicitly or not, every quantity given herein is meant to refer to the actual given value, and it is also meant to refer to the approximation to such given value that would reasonably be inferred based on the ordinary skill in the art, including equivalents and approximations due to the experimental and/or measurement conditions for such given value. Whenever a yield is given as a percentage, such yield refers to a mass of the entity for which the yield is given with respect to the maximum amount of the same entity that could be obtained under the particular stoichiometric conditions. Concentrations that are given as percentages refer to mass ratios, unless indicated differently.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited.

Except as otherwise noted, the methods and techniques of the present embodiments are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification. See, e.g., Loudon, *Organic Chemistry*, 4th edition, New York: Oxford University Press, 2002, pp. 360-361, 1084-1085; Smith and March, *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, 5th edition, Wiley-Interscience, 2001.

Chemical Definitions

As used herein, "alkyl" refers to a saturated, straight- or branched-chain hydrocarbon group having from 1 to 15 carbon atoms. Representative alkyl groups include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, 2-methyl-1-propyl, 2-methyl-2-propyl, 2-methyl-1-butyl, 3-methyl-1-butyl, 2-methyl-3-butyl, 2,2-dimethyl-1-propyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, 2-methyl-2-pentyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 2,2-dimethyl-1-butyl, 3,3-dimethyl-1-butyl, 2-ethyl-1-butyl, butyl, isobutyl, t-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, and the like, and longer alkyl groups, such as heptyl, octyl, and the like. As used herein, "lower alkyl" means an alkyl having from 1 to 6 carbon atoms.

The term "alkenyl" refers to straight chain or branched hydrocarbyl groups having from 2 to 6 carbon atoms and preferably 2 to 4 carbon atoms and having at least 1 and preferably from 1 to 2 sites of double bond unsaturation. This term includes, by way of example, bi-vinyl, allyl, and but-3-en-1-yl. Included within this term are the cis and trans isomers or mixtures of these isomers.

The term "alkynyl" refers to straight or branched mono-valent hydrocarbyl groups having from 2 to 6 carbon atoms and preferably 2 to 3 carbon atoms and having at least 1 and preferably from 1 to 2 sites of triple bond unsaturation. Examples of such alkynyl groups include acetylenyl ($-\text{C}\equiv\text{CH}$), and propargyl ($-\text{CH}_2\text{C}\equiv\text{CH}$).

The term "alkoxy" as used herein includes $-\text{O}(\text{alkyl})$, wherein alkyl is defined above.

"Aryl" means a mono-, bi-, or tricyclic aromatic group, wherein all rings of the group are aromatic and all ring atoms are carbon atoms. For bi- or tricyclic systems, the individual aromatic rings are fused to one another. Examples of aryl groups are 6 and 10 membered aryls. Further examples of aryl groups include, but are not limited to, phenyl, naphthalene, and anthracene.

The term "cyano" as used herein means a substituent having a carbon atom joined to a nitrogen atom by a triple bond.

The term "deuterium" as used herein means a stable isotope of hydrogen having one proton and one neutron.

The term "halo" represents chloro, fluoro, bromo, or iodo. In some embodiments, halo is chloro, fluoro, or bromo. The term "halogen" as used herein refers to fluorine, chlorine, bromine, or iodine.

The term "haloalkyl" represents an alkyl group substituted with one, two, three, or more halogen atoms. Examples of haloalkyl groups include fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, trifluoroethyl, and trifluoropropyl.

The term "hydroxy" means an $-\text{OH}$ group.

The term "oxo" means an $=\text{O}$ group and may be attached to a carbon atom or a sulfur atom.

The term "N-oxide" refers to the oxidized form of a nitrogen atom.

that can occur at any valency-allowed position on the system. In a certain embodiment, one or more substituent means 1, 2, 3, 4, or 5 substituents. In another embodiment, one or more substituent means 1, 2, or 3 substituents.

Any atom that is represented herein with an unsatisfied valence is assumed to have the sufficient number of hydrogen atoms to satisfy the atom's valence.

When any variable (e.g., alkyl or Rⁿ) appears in more than one place in any formula or description provided herein, the definition of that variable on each occurrence is independent of its definition at every other occurrence.

Numerical ranges, as used herein, are intended to include sequential whole numbers. For example, a range expressed as "from 0 to 4" or "0-4" includes 0, 1, 2, 3 and 4.

When a multifunctional moiety is shown, the point of attachment to the remainder of the formula can be at any point on the multifunctional moiety. In some embodiments, the point of attachment is indicated by a line or hyphen. For example, aryloxy- refers to a moiety in which an oxygen atom is the point of attachment to the core molecule while aryl is attached to the oxygen atom.

The nomenclature used herein to name the subject compounds is illustrated in the Examples herein. This nomenclature has generally been derived using the commercially-available LexiChem TK software (OpenEye, Santa Fe, N. Mex.).

Any formula given herein is intended to represent compounds having structures depicted by the structural formula as well as certain variations or forms. For example, compounds of any formula given herein may have asymmetric or chiral centers and therefore exist in different stereoisomeric forms. All stereoisomers, including optical isomers, enantiomers, and diastereomers, of the compounds of the general formula, and mixtures thereof, are considered to fall within the scope of the formula. Furthermore, certain structures may exist as geometric isomers (i.e., cis and trans isomers), as tautomers, or as atropisomers. All such isomeric forms, and mixtures thereof, are contemplated herein as part of the present invention. Thus, any formula given herein is intended to represent a racemate, one or more enantiomeric forms, one or more diastereomeric forms, one or more tautomeric or atropisomeric forms, and mixtures thereof.

The compounds described herein include pharmaceutically acceptable salt forms of compounds of Formula I or II. A "pharmaceutically acceptable salt" refers to a salt form of a free acid or base of a compound of Formula I or II that is non-toxic, is physiologically tolerable, is compatible with the pharmaceutical composition in which it is formulated, and is otherwise suitable for formulation and/or administration to a subject. Reference to a compound herein is understood to include reference to a pharmaceutically acceptable salt of said compound unless otherwise indicated.

Compound salts include acidic salts formed with inorganic and/or organic acids, as well as basic salts formed with inorganic and/or organic bases. In addition, where a given compound contains both a basic moiety, such as, but not limited to, a pyridine or imidazole, and an acidic moiety, such as, but not limited to, a carboxylic acid, one of skill in the art will recognize that the compound may exist as a zwitterion ("inner salt"); such salts are included within the term "salt" as used herein. Salts of the compounds of the invention may be prepared, for example, by reacting a compound with an amount of a suitable acid or base, such as an equivalent amount, in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

Exemplary acid addition salts include acetates, ascorbates, benzoates, benzenesulfonates, bisulfates, borates, butyrates, citrates, camphorates, camphorsulfonates, fumarates, hydrochlorides, hydrobromides, hydroiodides, isonicotinate, lactates, maleates, methanesulfonates, naphthalenesulfonates, nitrates, oxalates, phosphates, propionates, salicylates, succinates, sulfates, tartarates, thiocyanates, toluenesulfonates (also known as tosylates) and the like.

Exemplary basic salts include ammonium salts, alkali metal salts such as sodium, lithium, and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases (for example, organic amines) such as dicyclohexylamines, t-butyl amines, and salts with amino acids such as arginine, lysine and the like.

Additionally, acids and bases which are generally considered suitable for the formation of pharmaceutically useful salts from pharmaceutical compounds are discussed, for example, by P. Stahl et al., Camille G. (eds.) Handbook of Pharmaceutical Salts: Properties, Selection and Use. (2002) Zurich: Wiley-VCH; S. Berge et al., *J. Pharm. Sci.* (1977) 66(1) 1-19. These disclosures are incorporated herein by reference thereto.

Additionally, any compound described herein is intended to refer also to any unsolvated form, or a hydrate or solvate of such a compound, and mixtures thereof, even if such forms are not listed explicitly. "Solvate" means a physical association of a compound of the invention with one or more solvent molecules. This physical association involves varying degrees of ionic and covalent bonding, including hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of a crystalline solid. "Solvate" encompasses both solution-phase and isolatable solvates. Suitable solvates include those formed with pharmaceutically acceptable solvents such as water, ethanol, and the like. In some embodiments, the solvent is water and the solvates are hydrates. A compound of Formula I or II, including any hydrate or solvate forms, may be in the form of a crystalline polymorph, an amorphous solid, or a non-solid form.

The invention also relates to pharmaceutically acceptable prodrugs of the compounds of Formula I or II, and treatment methods employing such pharmaceutically acceptable prodrugs. The term "prodrug" means a precursor of a designated compound that, following administration to a subject, yields the compound in vivo via a chemical or physiological process such as solvolysis or enzymatic cleavage, or under physiological conditions (e.g., a prodrug on being brought to physiological pH is converted to the compound of Formula I or II). A "pharmaceutically acceptable prodrug" is a prodrug that is non-toxic, biologically tolerable, and otherwise suitable for formulation and/or administration to the subject. Illustrative procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985. Prodrugs include, but are not limited to, esters, amides, sulfonates, and phosphonate esters.


The present invention also relates to pharmaceutically active metabolites of compounds of Formula I or II, and uses of such metabolites in the methods of the invention. A "pharmaceutically active metabolite" means a pharmacologically active product of metabolism in the body of a compound of Formula I or II, or salts thereof. Prodrugs and active metabolites of a compound may be determined using routine techniques known or available in the art. See, e.g., Bertolini et al., *J. Med. Chem.* 1997, 40, 2011-2016; Shan et al., *J. Pharm. Sci.* 1997, 86 (7), 765-767; Bagshawe, *Drug*

Dev. Res. 1995, 34, 220-230; Bodor, *Adv. Drug Res.* 1984, 13, 255-331; Bundgaard, *Design of Prodrugs* (Elsevier Press, 1985); and Larsen, *Design and Application of Prodrugs, Drug Design and Development* (Krogsgaard-Larsen et al., eds., Harwood Academic Publishers, 1991).

Any formula given herein is also intended to represent unlabeled forms as well as isotopically labeled forms of the compounds. Isotopically labeled compounds have structures depicted by the formulas given herein except that one or more atoms are replaced by an atom having a selected atomic mass or mass number. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, fluorine, chlorine, and iodine, such as ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , ^{36}Cl , and ^{125}I , respectively. Such isotopically labeled compounds are useful in metabolic studies (for example with ^{14}C), reaction kinetic studies (with, for example ^2H or ^3H), detection or imaging techniques [such as positron emission tomography (PET) or single-photon emission computed tomography (SPECT)] including drug or substrate tissue distribution assays, or in radioactive treatment of patients. In particular, an ^{18}F or ^{11}C labeled compound may be particularly suitable for PET or SPECT studies. Further, substitution with heavier isotopes such as deuterium (i.e., ^2H) may afford certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life or reduced dosage requirements. Isotopically labeled compounds of this invention and prodrugs thereof can generally be prepared by carrying out the procedures disclosed in the schemes or in the examples and preparations described below by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent. Further, substitution with heavier isotopes such as deuterium (i.e., ^2H) may afford certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life or reduced dosage requirements. Isotopically labeled compounds of this invention and prodrugs thereof can generally be prepared by carrying out the procedures disclosed in the schemes or in the examples and preparations described below by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent.

The use of the terms "salt," "solvate," "polymorph," "prodrug," and the like, with respect to the compounds described herein is intended to apply equally to the salt, solvate, polymorph, and prodrug forms of enantiomers, stereoisomers, rotamers, tautomers, atropisomers, and racemates of the inventive compounds.

Also contemplated herein are methods of synthesizing compounds of Formula I or II.
Compounds of the Invention



In some embodiments of (a) Formula I or (b) Formula II, R^a is C_{1-6} alkyl optionally substituted with one or more R^b substituents. In some embodiments, R^a is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pen-

tyl, or isopentyl, each optionally substituted with one or more R^b substituents. In some embodiments, R^a is C_{1-6} alkyl optionally substituted with one or two R^b substituents.

In some embodiments, each R^b is independently —OH, methoxy, ethoxy, —NR^dR^e, —C(O)NR^dR^e, thiomethyl, thioethyl, methanesulfonyl, ethanesulfonyl, cyano, fluoro, chloro, bromo, pyrrolyl, pyrazolyl, imidazolyl, furanyl, thiophenyl, triazolyl, tetrazolyl, oxazolyl, or thiazolyl. In other embodiments, each R^b is independently —OH, —C(O)NHCH₃, —CF₃, methoxy, ethoxy, fluoro, —C(O)NH₂, —C(O)N(CH₃)₂, —N(CH₃)₂, methanesulfonyl, thiomethyl, cyano, pyrazolyl, 6-oxa-1-azaspiro[3.3]heptan-1-yl, azetidiny, 3-hydroxyazetidiny, pyrrolidiny, or hydroxyethylamino.

In other embodiments, R^a is C_{1-6} alkenyl or C_{1-6} alkynyl. In some embodiments, R^a is ethenyl, isopropenyl, or propynyl.

In some embodiments, R^a is halo. In some embodiments, R^a is bromo, chloro, fluoro, or iodo.

In other embodiments, R^a is —C(O)R^c; —NR^dR^e; —C(O)NR^dR^e; —C(S)NR^dR^e; —C(=N—OH)— C_{1-4} alkyl; or —SO₂ C_{1-4} alkyl. In other embodiments, R^a is —C(O)NR^dR^e.

In some embodiments, R^c is methyl, ethyl, propyl, isopropyl, butyl, fluoromethyl, difluoromethyl, trifluoromethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, azetidiny, pyrrolidiny, piperidiny, morpholiny, thiomorpholiny, oxetanyl, tetrahydrofuranyl, or tetrahydropyranyl. In other embodiments, R^c is ethyl, cyclopropyl, methyl, oxetanyl, or trifluoromethyl.

In some embodiments, R^d is H, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, or tert-butyl. In some embodiments, R^e is H, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, cyanomethyl, trifluoroethyl, hydroxyethyl, 2-hydroxy-1-methylethyl, hydroxypropyl, cyclopropyl, hydroxy, methoxy, or oxetanylmethyl. In other embodiments, R^d and R^e taken together with the nitrogen to which they are attached form azetidiny, pyrrolidiny, piperidiny, morpholiny, thiomorpholiny, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, or 6-oxa-1-azaspiro[3.3]heptan-1-yl, each optionally substituted with C_{1-4} alkyl or —OH.

In other embodiments, R^a is cyano.

In other embodiments, R^a is C_{3-6} cycloalkyl optionally substituted with one or more R^f substituents. In some embodiments, R^a is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl, each optionally substituted with one or more R^f substituents. In other embodiments, R^a is cyclopropyl, optionally substituted with one or more R^f substituents.

In some embodiments, each R^f is independently: methyl, ethyl, propyl, or isopropyl, each optionally substituted with —OH, cyano, methoxy, or ethoxy; C_{1-4} fluoroalkyl; —CONH₂; or cyano. In other embodiments, each R^f is independently hydroxymethyl, methyl, cyano, trifluoromethyl, cyanomethyl, methoxymethyl, fluoromethyl, hydroxymethyl, 1-hydroxy-1-methyl-ethyl, or —CONH₂.

In some embodiments, R^a is a phenyl, monocyclic heteroaryl, or heterocycloalkyl ring, each ring optionally substituted with one or more R^g substituents. In other embodiments, R^a is a phenyl, pyrrolyl, furanyl, thiophenyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, triazolyl, pyridyl, pyrimidinyl, pyridazinyl, pyrazinyl, or pyrimidinyl, each optionally substituted with one or more R^g substituents. In some embodiments, R^a is optionally substituted with one or two R^g substituents. In some embodiments, each R^g is independently methyl, ethyl, propyl, isopropyl, —CF₃, fluoro, chloro, —NH₂, —OCH₃, cyano, or —OH. In other

13


embodiments, each R^s is independently fluoro, methyl, $-\text{NH}_2$, $-\text{CF}_3$, chloro, methoxy, or cyano.

In some embodiments, R^a and R^1 taken together with the carbons to which they are attached form a 5- to 7-membered ring, optionally containing an O or NH, and optionally substituted with one or more R^h substituents. In other embodiments, R^a and R^1 taken together with the carbons to which they are attached form cyclopentenyl, cyclohexenyl, dihydrofuranyl, dihydropyranyl, dihydropyrrolyl, or tetrahydropyridine, each optionally substituted with one or more R^h substituents. In some embodiments, each R^h is independently: methyl, ethyl, or propyl, each optionally substituted with hydroxy, cyano, methoxy, or $-\text{C}(\text{O})\text{N}(\text{CH}_3)_2$; $-\text{C}(\text{O})\text{NR}^i\text{R}^j$; or cyano. In other embodiments, each R^h is independently hydroxypropyl, hydroxyethyl, hydroxymethyl, methyl, cyano, methoxymethyl, $-\text{C}(\text{O})\text{NH}_2$, or $-\text{CH}_2\text{C}(\text{O})\text{N}(\text{CH}_3)_2$. Alternatively, two R^h groups attached to the same carbon are taken together with the carbon to which they are attached to form cyclopentyl or a carbonyl.

In some embodiments, R^1 is H, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, trifluoroethyl, cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. In other embodiments, R^1 is H, methyl, isopropyl, trifluoromethyl, or cyclopropyl.

In some embodiments, R^2 is R^m . In other embodiments, R^2 is $-\text{OR}^m$. In other embodiments, R^2 is $-\text{NR}^m\text{R}^n$. In some embodiments, R^m is phenyl, naphthyl, pyridyl, pyrazinyl, pyridazinyl, pyrrolyl, pyrazolyl, triazolyl, imidazolyl, furanyl, oxazolyl, isoxazolyl, thiophenyl, thiazolyl, isothiazolyl, indolyl, indazolyl, quinolinyl, or isoquinolinyl, each optionally substituted with one or more R^s substituents. In other embodiments, R^m is phenyl, naphthyl, pyridyl, indazolyl, or isoquinolinyl, each optionally substituted with one or more R^s substituents. In other embodiments, R^m is pyrazolyl, optionally substituted with one or more R^s substituents. In other embodiments, R^m is phenyl, optionally substituted with one or more R^s substituents. In other embodiments, R^m is optionally substituted with one or two R^s substituents. In some embodiments, each R^s is independently methyl, ethyl, propyl, isopropyl, butyl, fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, trifluoroethyl, methoxy, ethoxy, isopropoxy, hydroxymethyl, hydroxyethyl, trifluoromethoxy, fluoro, chloro, bromo, iodo, cyano, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, $-\text{NHSO}_2\text{C}_{1-2}\text{alkyl}$, or $-\text{SO}_2\text{C}_{1-2}\text{alkyl}$. In other embodiments, each R^s is independently fluoro, chloro, trifluoromethyl, cyano, methyl, methoxy, cyclopropyl, $-\text{NHSO}_2\text{CH}_3$, fluoroethyl, ethyl, propyl, difluoromethyl, hydroxymethyl, fluoromethyl, or methanesulfonyl.

In other embodiments, R^2 is R^m and R^m is



wherein at least one of X^1 , X^2 , and X^3 is N, and the other two are independently N, NR^r , O, S, or $\text{C}-\text{R}^r$;

R^p and R^q are each independently H; $\text{C}_{1-4}\text{haloalkyl}$;

$\text{C}_{1-4}\text{alkyl}$ optionally substituted with $-\text{OH}$; halo; cyano; or $\text{C}_{3-6}\text{cycloalkyl}$; and

R^q is H or fluoro;

14

or R^q and R^r taken together with the carbons to which they are attached form phenyl, optionally substituted with halo.

In some embodiments, X^1 and X^2 are each N and X^3 is $\text{C}-\text{R}^r$. In other embodiments, X^2 is N and X^1 and X^3 are each independently $\text{C}-\text{R}^r$. In other embodiments, X^1 , X^2 , and X^3 are each N.

In some embodiments, R^p and R^r are each independently H, fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, trifluoroethyl, methyl, ethyl, hydroxymethyl, hydroxyethyl, chloro, cyano, cyclopropyl, cyclobutyl, or cyclopentyl. In other embodiments, R^p is trifluoromethyl, chloro, methyl, hydroxyethyl, cyclopropyl, cyano, difluoromethyl, or ethyl. In other embodiments, R^r is ethyl, trifluoromethyl, methyl, chloro, H, hydroxyethyl, cyclopropyl, or cyano.

In some embodiments, R^q is H or fluoro. In other embodiments, R^q and R^r taken together with the carbons to which they are attached form phenyl, optionally substituted with fluoro.


In some embodiments, R^m is H, fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, difluoroethyl, or trifluoroethyl, or is methyl or ethyl optionally substituted with $-\text{OH}$, methoxy, or ethoxy. In other embodiments, R^m is H, methyl, ethyl, fluoroethyl, difluoroethyl, or trifluoroethyl.

In some embodiments, R^m and R^n taken together with the nitrogen to which they are attached form dihydroindole, optionally substituted with methyl or fluoro.

In some embodiments, R^3 is H. In other embodiments, R^3 is methyl.

In some embodiments, R^4 is H. In other embodiments, R^4 is fluoro.

In some embodiments, the compound of Formula (I) is a compound of Formula (I-A):



(I-A)

wherein

R^a is $-\text{C}(\text{O})\text{NR}^d\text{R}^e$;

wherein R^d is H or $\text{C}_{1-4}\text{alkyl}$;

R^e is H; $\text{C}_{1-4}\text{alkyl}$ optionally substituted with $-\text{CN}$, $-\text{CF}_3$, $-\text{OH}$, or a monocyclic heterocycloalkyl; $\text{C}_{3-6}\text{cycloalkyl}$; $-\text{OH}$; or $-\text{OC}_{1-4}\text{alkoxy}$;

or R^d and R^e taken together with the nitrogen to which they are attached form a heterocycloalkyl, optionally substituted with $\text{C}_{1-4}\text{alkyl}$ or $-\text{OH}$;

R^1 is selected from the group consisting of H, $\text{C}_{1-6}\text{alkyl}$, $\text{C}_{1-4}\text{haloalkyl}$, and $\text{C}_{3-6}\text{cycloalkyl}$;

R^2 is $-\text{R}^m$, $-\text{OR}^m$, or $-\text{NR}^m\text{R}^n$;

wherein R^m is aryl or heteroaryl, each optionally substituted with one or more R^s substituents;

wherein each R^s substituent is independently selected from the group consisting of $\text{C}_{1-4}\text{alkyl}$, $\text{C}_{1-4}\text{haloalkyl}$, $\text{C}_{1-4}\text{alkoxy}$, $\text{C}_{1-4}\text{alkyl}-\text{OH}$, $\text{C}_{1-4}\text{haloalkoxy}$, halo, cyano, $\text{C}_{3-6}\text{cycloalkyl}$, $-\text{NHSO}_2\text{C}_{1-4}\text{alkyl}$, and $-\text{SO}_2\text{C}_{1-4}\text{alkyl}$;

R^n is H, $\text{C}_{1-4}\text{haloalkyl}$, or $\text{C}_{1-4}\text{alkyl}$ optionally substituted with $-\text{OH}$ or $\text{C}_{1-4}\text{alkoxy}$;

15

or R^m and Rⁿ taken together with the nitrogen to which they are attached form a pyrrolidine or piperidine ring, optionally substituted with C₁₋₄alkyl and optionally fused to phenyl, wherein said phenyl is optionally substituted with halo;

R³ is H or methyl; and

R⁴ is H or fluoro;


or a pharmaceutically acceptable salt thereof.

In some embodiments of compounds of Formula (I-A), R^d is H, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, or tert-butyl. In some embodiments, R^e is H, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, cyanomethyl, trifluoroethyl, hydroxyethyl, 2-hydroxy-1-methylethyl, hydroxypropyl, cyclopropyl, hydroxy, methoxy, or oxetanyl-methyl. In other embodiments, R^g and R^e taken together with the nitrogen to which they are attached form azetidiny, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, oxetanyl, tetrahydrofuran-yl, tetrahydropyran-yl, or 6-oxa-1-azaspiro[3.3]heptan-1-yl, each optionally substituted with C₁₋₄alkyl or —OH.

In some embodiments, R¹ is H, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, fluoromethyl, fluoromethyl, trifluoromethyl, fluoroethyl, trifluoroethyl, cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. In other embodiments, R¹ is H, methyl, isopropyl, trifluoromethyl, or cyclopropyl.

In some embodiments, R² is R^m. In other embodiments, R² is —OR^m. In other embodiments, R² is —NR^mRⁿ. In some embodiments, R^m is phenyl, naphthyl, pyridyl, pyrazinyl, pyridazinyl, pyrrolyl, pyrazolyl, triazolyl, imidazolyl, furanyl, oxazolyl, isoxazolyl, thiophenyl, thiazolyl, isothiazolyl, indolyl, indazolyl, quinolinyl, or isoquinolinyl, each optionally substituted with one or more R^s substituents. In other embodiments, R^m is phenyl, naphthyl, pyridyl, indazolyl, or isoquinolinyl, each optionally substituted with one or more R^s substituents. In other embodiments, R^m is pyrazolyl, optionally substituted with one or more R^s substituents. In other embodiments, R^m is phenyl, optionally substituted with one or more R^s substituents. In other embodiments, R^m is optionally substituted with one or two R^s substituents. In some embodiments, each R^s is independently methyl, ethyl, propyl, isopropyl, butyl, fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, trifluoroethyl, methoxy, ethoxy, isopropoxy, hydroxymethyl, hydroxyethyl, trifluoromethoxy, fluoro, chloro, bromo, iodo, cyano, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, —NHSO₂C₁₋₂alkyl, or —SO₂C₁₋₂alkyl. In other embodiments, each R^s is independently fluoro, chloro, trifluoromethyl, cyano, methyl, methoxy, cyclopropyl, —NHSO₂CH₃, fluoroethyl, ethyl, propyl, difluoromethyl, hydroxymethyl, fluoromethyl, or methanesulfonyl.

In other embodiments, R² is R^m and R^m is



wherein at least one of X¹, X², and X³ is N, and the other two are independently N, NR^r, O, S, or C—R^r;

R^p and R^r are each independently H; C₁₋₄haloalkyl; C₁₋₄alkyl optionally substituted with —OH; halo; cyano; or C₃₋₆cycloalkyl; and

16

R^q is H or fluoro;

or R^q and R^r taken together with the carbons to which they are attached form phenyl, optionally substituted with halo.

In some embodiments, X¹ and X² are each N and X³ is C—R^r. In other embodiments, X² is N and X¹ and X³ are each independently C—R^r. In other embodiments, X¹, X², and X³ are each N.

In some embodiments, R^p and R^r are each independently H, fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, trifluoroethyl, methyl, ethyl, hydroxymethyl, hydroxyethyl, chloro, cyano, cyclopropyl, cyclobutyl, or cyclopentyl. In other embodiments, R^p is trifluoromethyl, chloro, methyl, hydroxyethyl, cyclopropyl, cyano, difluoromethyl, or ethyl. In other embodiments, R^r is ethyl, trifluoromethyl, methyl, chloro, H, hydroxyethyl, cyclopropyl, or cyano.

In some embodiments, R^q is H or fluoro. In other embodiments, R^q and R^r taken together with the carbons to which they are attached form phenyl, optionally substituted with fluoro.


In some embodiments, Rⁿ is H, fluoromethyl, difluoroethyl, trifluoromethyl, fluoroethyl, difluoroethyl, or trifluoroethyl, or is methyl or ethyl optionally substituted with —OH, methoxy, or ethoxy. In other embodiments, Rⁿ is H, methyl, ethyl, fluoroethyl, difluoroethyl, or trifluoroethyl.

In some embodiments, R^m and Rⁿ taken together with the nitrogen to which they are attached form dihydroindole, optionally substituted with methyl or fluoro.

In some embodiments, R³ is H. In other embodiments, R³ is methyl.

In some embodiments, R⁴ is H. In other embodiments, R⁴ is fluoro.

In some embodiments, the compound of Formula (I) is a compound of Formula (I-B):



(I-B)

wherein

R^a is cyclopropyl, optionally substituted with one or more R^f substituents;

each R^f substituent is independently selected from the group consisting of: C₁₋₄alkyl optionally substituted with —OH, cyano, or C₁₋₄alkoxy; C₁₋₄haloalkyl; —CONH₂; and cyano; and

R¹ is selected from the group consisting of H, C₁₋₆alkyl, C₁₋₄haloalkyl, and C₃₋₆cycloalkyl;

R² is —R^m, —OR^m, or —NR^mRⁿ;

wherein R^m is aryl or heteroaryl, each optionally substituted with one or more R^s substituents;

wherein each R^s substituent is independently selected from the group consisting of C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy, C₁₋₄alkyl-OH, C₁₋₄haloalkoxy, halo, cyano, C₃₋₆cycloalkyl, —NHSO₂C₁₋₄alkyl, and —SO₂C₁₋₄alkyl;

Rⁿ is H, C₁₋₄haloalkyl, or C₁₋₄alkyl optionally substituted with —OH or C₁₋₄alkoxy;

or R^m and Rⁿ taken together with the nitrogen to which they are attached form a pyrrolidine or piperidine

17

ring, optionally substituted with C₁₋₄alkyl and optionally fused to phenyl, wherein said phenyl is optionally substituted with halo;

R³ is H or methyl; and

R⁴ is H or fluoro;


or a pharmaceutically acceptable salt thereof.

In some embodiments of Formula (I-B), each R^f is independently: methyl, ethyl, propyl, or isopropyl, each optionally substituted with —OH, cyano, methoxy, or ethoxy; C₁₋₄fluoroalkyl; —CONH₂; or cyano. In other embodiments, each R^f is independently hydroxymethyl, methyl, cyano, trifluoromethyl, cyanomethyl, methoxymethyl, fluoromethyl, hydroxymethyl, 1-hydroxy-1-methyl-ethyl, or —CONH₂. In some embodiments, R^g is cyclopropyl, optionally substituted with one or two R^f substituents.

In some embodiments, R¹ is H, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, fluoromethyl, fluoromethyl, trifluoromethyl, fluoroethyl, trifluoroethyl, cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. In other embodiments, R¹ is H, methyl, isopropyl, trifluoromethyl, or cyclopropyl.

In some embodiments, R² is R^m. In other embodiments, R² is —OR^m. In other embodiments, R² is —NR^mRⁿ. In some embodiments, R^m is phenyl, naphthyl, pyridyl, pyrazinyl, pyridazinyl, pyrrolyl, pyrazolyl, triazolyl, imidazolyl, furanyl, oxazolyl, isoxazolyl, thiophenyl, thiazolyl, isothiazolyl, indolyl, indazolyl, quinolinyl, or isoquinolinyl, each optionally substituted with one or more R^s substituents. In other embodiments, R^m is phenyl, naphthyl, pyridyl, indazolyl, or isoquinolinyl, each optionally substituted with one or more R^s substituents. In other embodiments, R^m is pyrazolyl, optionally substituted with one or more R^s substituents. In other embodiments, R^m is phenyl, optionally substituted with one or more R^s substituents. In other embodiments, R^m is optionally substituted with one or two R^s substituents. In some embodiments, each R^s is independently methyl, ethyl, propyl, isopropyl, butyl, fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, trifluoroethyl, methoxy, ethoxy, isopropoxy, hydroxymethyl, hydroxyethyl, trifluoromethoxy, fluoro, chloro, bromo, iodo, cyano, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, —NHSO₂C₁₋₂alkyl, or —SO₂C₁₋₂alkyl. In other embodiments, each R^s is independently fluoro, chloro, trifluoromethyl, cyano, methyl, methoxy, cyclopropyl, —NHSO₂CH₃, fluoroethyl, ethyl, propyl, difluoromethyl, hydroxymethyl, fluoromethyl, or methanesulfonyl.

In other embodiments, R² is R^m and R^m is



wherein at least one of X¹, X², and X³ is N, and the other two are independently N, NR^r, O, S, or C—R^r;

R^p and R^r are each independently H; C₁₋₄haloalkyl; C₁₋₄alkyl optionally substituted with —OH; halo; cyano; or C₃₋₆cycloalkyl; and

R^q is H or fluoro;

or R^q and R^r taken together with the carbons to which they are attached form phenyl, optionally substituted with halo.

18

In some embodiments, X¹ and X² are each N and X³ is C—R^r. In other embodiments, X² is N and X¹ and X³ are each independently C—R^r. In other embodiments, X¹, X², and X³ are each N.

In some embodiments, R^p and R^r are each independently H, fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, trifluoroethyl, methyl, ethyl, hydroxymethyl, hydroxyethyl, chloro, cyano, cyclopropyl, cyclobutyl, or cyclopentyl. In other embodiments, R^p is trifluoromethyl, chloro, methyl, hydroxyethyl, cyclopropyl, cyano, difluoromethyl, or ethyl. In other embodiments, R^r is ethyl, trifluoromethyl, methyl, chloro, H, hydroxyethyl, cyclopropyl, or cyano.

In some embodiments, R^q is H or fluoro. In other embodiments, R^q and R^r taken together with the carbons to which they are attached form phenyl, optionally substituted with fluoro.


In some embodiments, Rⁿ is H, fluoromethyl, difluoroethyl, trifluoromethyl, fluoroethyl, difluoroethyl, or trifluoroethyl, or is methyl or ethyl optionally substituted with —OH, methoxy, or ethoxy. In other embodiments, Rⁿ is H, methyl, ethyl, fluoroethyl, difluoroethyl, or trifluoroethyl.

In some embodiments, R^m and Rⁿ taken together with the nitrogen to which they are attached form dihydroindole, optionally substituted with methyl or fluoro.

In some embodiments, R³ is H. In other embodiments, R³ is methyl.

In some embodiments, R⁴ is H. In other embodiments, R⁴ is fluoro.

In some embodiments, the compound of Formula (I) is a compound of Formula (I-C):



(I-C)

R^q is a monocyclic heteroaryl ring, optionally substituted with one or more R^g substituents;

each R^g substituent is independently selected from the group consisting of C₁₋₄alkyl, —CF₃, halo, —NH₂, —OCH₃, cyano, and —OH;

R¹ is selected from the group consisting of H, C₁₋₆alkyl, C₁₋₄haloalkyl, and C₃₋₆cycloalkyl;

R² is —R^m, —OR^m, or —NR^mRⁿ;

wherein R^m is aryl or heteroaryl, each optionally substituted with one or more R^s substituents;

wherein each R^s substituent is independently selected from the group consisting of C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy, C₁₋₄alkyl-OH, C₁₋₄haloalkoxy, halo, cyano, C₃₋₆cycloalkyl, —NHSO₂C₁₋₄alkyl, and —SO₂C₁₋₄alkyl;

Rⁿ is H, C₁₋₄haloalkyl, or C₁₋₄alkyl optionally substituted with —OH or C₁₋₄alkoxy;

or R^m and Rⁿ taken together with the nitrogen to which they are attached form a pyrrolidine or piperidine ring, optionally substituted with C₁₋₄alkyl and optionally fused to phenyl, wherein said phenyl is optionally substituted with halo;

R³ is H or methyl; and

R⁴ is H or fluoro;

or a pharmaceutically acceptable salt thereof.


19

In some embodiments of Formula (I-C), R^q is pyrrolyl, furanyl, thiophenyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, triazolyl, pyridyl, pyrimidinyl, pyridazinyl, pyrazinyl, or pyrimidinyl, each optionally substituted with one or more R^s substituents. In some embodiments, R^q is optionally substituted with one or two R^s substituents. In some embodiments, each R^s is independently methyl, ethyl, propyl, isopropyl, —CF₃, fluoro, chloro, —NH₂, —OCH₃, cyano, or —OH. In other embodiments, each R^s is independently fluoro, methyl, —NH₂, —CF₃, chloro, methoxy, or cyano.

In some embodiments, R¹ is H, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, trifluoroethyl, cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. In other embodiments, R¹ is H, methyl, isopropyl, trifluoromethyl, or cyclopropyl.

In some embodiments, R² is R^m. In other embodiments, R² is —OR^m. In other embodiments, R² is —NR^mRⁿ. In some embodiments, R^m is phenyl, naphthyl, pyridyl, pyrazinyl, pyridazinyl, pyrrolyl, pyrazolyl, triazolyl, imidazolyl, furanyl, oxazolyl, isoxazolyl, thiophenyl, thiazolyl, isothiazolyl, indolyl, indazolyl, quinolinyl, or isoquinolinyl, each optionally substituted with one or more R^s substituents. In other embodiments, R^m is phenyl, naphthyl, pyridyl, indazolyl, or isoquinolinyl, each optionally substituted with one or more R^s substituents. In other embodiments, R^m is pyrazolyl, optionally substituted with one or more R^s substituents. In other embodiments, R^m is phenyl, optionally substituted with one or more R^s substituents. In other embodiments, R^m is optionally substituted with one or two R^s substituents. In some embodiments, each R^s is independently methyl, ethyl, propyl, isopropyl, butyl, fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, trifluoroethyl, methoxy, ethoxy, isopropoxy, hydroxymethyl, hydroxyethyl, trifluoromethoxy, fluoro, chloro, bromo, iodo, cyano, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, —NHSO₂C₁₋₂alkyl, or —SO₂C₁₋₂alkyl. In other embodiments, each R^s is independently fluoro, chloro, trifluoromethyl, cyano, methyl, methoxy, cyclopropyl, —NHSO₂CH₃, fluoroethyl, ethyl, propyl, difluoromethyl, hydroxymethyl, fluoromethyl, or methanesulfonyl.

In other embodiments, R² is R^m and R^m is



wherein at least one of X¹, X², and X³ is N, and the other two are independently N, NR^r, O, S, or C—R^r;

R^p and R^r are each independently H; C₁₋₄haloalkyl; C₁₋₄alkyl optionally substituted with —OH; halo; cyano; or C₃₋₆cycloalkyl; and R^q is H or fluoro;

or R^q and R^r taken together with the carbons to which they are attached form phenyl, optionally substituted with halo.

In some embodiments, X¹ and X² are each N and X³ is C—R^r. In other embodiments, X² is N and X¹ and X³ are each independently C—R^r. In other embodiments, X¹, X², and X³ are each N.

In some embodiments, R^p and R^r are each independently H, fluoromethyl, difluoromethyl, trifluoromethyl, fluoro-

20

ethyl, trifluoroethyl, methyl, ethyl, hydroxymethyl, hydroxyethyl, chloro, cyano, cyclopropyl, cyclobutyl, or cyclopentyl. In other embodiments, R^p is trifluoromethyl, chloro, methyl, hydroxyethyl, cyclopropyl, cyano, difluoromethyl, or ethyl. In other embodiments, R^r is ethyl, trifluoromethyl, methyl, chloro, H, hydroxyethyl, cyclopropyl, or cyano.

In some embodiments, R^q is H or fluoro. In other embodiments, R^q and R^r taken together with the carbons to which they are attached form phenyl, optionally substituted with fluoro.

In some embodiments, Rⁿ is H, fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, difluoroethyl, or trifluoroethyl, or is methyl or ethyl optionally substituted with —OH, methoxy, or ethoxy. In other embodiments, Rⁿ is H, methyl, ethyl, fluoroethyl, difluoroethyl, or trifluoroethyl.


In some embodiments, R^m and Rⁿ taken together with the nitrogen to which they are attached form dihydroindole, optionally substituted with methyl or fluoro.

In some embodiments, R³ is H. In other embodiments, R³ is methyl.

In some embodiments, R⁴ is H. In other embodiments, R⁴ is fluoro.

In some embodiments, compounds described herein are compounds of Formula II or pharmaceutically acceptable salts thereof. Compounds of Formula II include those in which each variable is defined independently as described herein for Formula I, I-A, I-B, or I-C, or combinations of said definitions. Additional embodiments of Formula II include compounds in which R^a is —SCH₃, —CH₂-cyclopropyl, difluorocyclopropyl, hydroxycyclopropyl, —OCH₂CF₃, —CH=CH—CN, or —CH=CH—CONH₂. Additional embodiments of Formula II include compounds in which R¹ is chloro, methoxy, cyano, ethoxy, trifluoroethoxy, or acetyl. Additional embodiments of Formula II include compounds in which R² is fluoro-isopropenyl, ethynyl, hydroxycyclopropyl, fluorocyclopropyl, —NH₂, —NO₂, or thiazolyl.

In other embodiments are compounds of Formula III:



(III)

wherein:

R¹⁰ is C₁₋₄alkyl, C₁₋₄haloalkyl, or cyano, or C₃₋₆cycloalkyl optionally substituted with —C₁₋₄alkyl-OH, R¹¹ is C₁₋₄alkyl; or R¹⁰ and R¹¹ taken together with the carbons to which they are attached form a C₅₋₆cycloalkyl;

R¹² is —H or halo; and

R¹³ is phenyl, optionally substituted with one or more substituents selected from the group consisting of halo, C₁₋₄haloalkyl, and cyano;

and pharmaceutically acceptable salts thereof.

Additional embodiments include pharmaceutical compositions comprising at least one compound of Formula III, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient, and a method of treating a subject suffering from a disease or medical condition mediated by NMDA receptor activity, comprising administering to the subject in need of such treatment an effective amount

of at least one compound of Formula III, or a pharmaceutically acceptable salt thereof.

Embodiments of the invention also include compounds in which each variable is defined independently as described above.

In certain embodiments, the compound of Formula I or II is a compound selected from the group consisting of the compounds in Table 1, and pharmaceutically acceptable salts thereof:

TABLE 1

Ex.	Chemical Name
1.1	N-(cyanomethyl)-7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
1.2	7-(4-Fluorophenoxy)methyl)-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide
1.3	3-[(Azetidin-1-yl)carbonyl]-7-(4-fluorophenoxy)methyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
1.4	N-ethyl-7-(4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide
1.5	7-(4-Fluorophenoxy)methyl)-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide
1.6	7-(3,4-Difluorophenoxy)methyl)-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide
1.7	N-ethyl-7-(4-fluorophenoxy)methyl)-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide
1.8	7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
1.9	7-(4-fluorophenoxy)methyl)-N-hydroxy-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
1.10	7-(4-Fluorophenoxy)methyl)-2-methyl-5-oxo-N-(propan-2-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide
1.11	7-(4-Fluorophenoxy)methyl)-N-(2-hydroxyethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide
1.12	7-(4-Fluorophenoxy)methyl)-N-(1-hydroxypropan-2-yl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide
1.13	7-(4-fluorophenoxy)methyl)-2-methyl-N-(oxetan-3-ylmethyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
1.14	7-(4-fluorophenoxy)methyl)-N-(3-hydroxypropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
1.15	N-cyclopropyl-7-(4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
1.16	7-(4-fluorophenoxy)methyl)-N-methoxy-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
1.17	7-(4-Fluorophenoxy)methyl)-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carbothioamide
2.1	7-(4-fluorophenoxy)methyl)-2-methyl-3-propionyl-5H-thiazolo[3,2-a]pyrimidin-5-one
2.2	7-(4-fluorophenoxy)methyl)-3-(1-hydroxypropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
2.3	7-(4-Fluorophenoxy)methyl)-3-(1-hydroxyethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
2.4	7-(4-Fluorophenoxy)methyl)-3-(2-hydroxypropan-2-yl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
2.5	3-acetyl-7-(4-fluorophenoxy)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
2.6	2-(7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)-N-methylacetamide
2.7	3-Cyclopropanecarbonyl-7-(4-fluorophenoxy)methyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
2.8	7-(4-Fluorophenoxy)methyl)-3-[1-(hydroxyimino)ethyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
2.9	7-(4-fluorophenoxy)methyl)-2-methyl-3-(oxetane-3-carbonyl)-5H-thiazolo[3,2-a]pyrimidin-5-one
2.10	7-(4-fluorophenoxy)methyl)-2-methyl-3-(2,2,2-trifluoro-1-hydroxyethyl)-5H-thiazolo[3,2-a]pyrimidin-5-one
2.11	7-(4-Fluorophenoxy)methyl)-2-methyl-3-(trifluoroacetyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
3.1	2-cyclopropyl-N-ethyl-7-(4-fluorophenoxy)methyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
3.2	7-(4-Fluorophenoxy)methyl)-N-methyl-5-oxo-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide
3.3	2-Cyclopropyl-7-(4-fluorophenoxy)methyl)-N-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide
3.4	N-Ethyl-7-(4-fluorophenoxy)methyl)-5-oxo-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide
3.5	7-(4-fluorophenoxy)methyl)-N-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
3.6	N-ethyl-7-[(5-fluoropyridin-2-yl)oxy]methyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide
4.1	7-(4-Fluorophenoxy)methyl)-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
4.2	7-(4-Fluorophenoxy)methyl)-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (enantiomer 1)

TABLE 1-continued

Ex.	Chemical Name
4.3	7-(4-Fluorophenoxymethyl)-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (enantiomer 2)
4.4	7-((3-fluorophenoxy)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
4.5	7-((4-fluorophenoxy)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
4.6	7-(2,4-Difluorophenoxymethyl)-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
4.7	7-(3,4-Difluorophenoxymethyl)-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
4.8	7-(4-Chlorophenoxymethyl)-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
4.9	7-[(5-Fluoropyridin-2-yl)oxy]methyl-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
4.10	3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-7-((4-(trifluoromethyl)phenoxy)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one
4.11	7-((4-fluorophenoxy)methyl)-2-methyl-3-(oxazol-2-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
4.12	7-((2-fluorophenoxy)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
4.13	4-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methoxy)benzotrile
4.14	7-((4-fluorophenoxy)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-5H-thiazolo[3,2-a]pyrimidin-5-one
4.15	7-((4-fluorophenoxy)methyl)-2-methyl-3-(1H-pyrazol-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
4.16	7-((4-fluorophenoxy)methyl)-2-methyl-3-(4H-1,2,4-triazol-3-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
4.17	3-cyclopropyl-7-[(4-fluorophenoxy)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
4.18	cis-2-[7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carbonitrile
4.18A	trans-2-[7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carbonitrile
4.19	cis-2-[7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carbonitrile (enantiomer 1)
4.20	cis-2-[7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carbonitrile (enantiomer 2)
4.21	trans-2-[7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carbonitrile (enantiomer 1)
4.22	trans-2-[7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carbonitrile (enantiomer 2)
4.23	7-(4-Fluorophenoxymethyl)-3-[cis-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
4.24	trans-7-(4-Fluorophenoxymethyl)-2-methyl-3-[2-(trifluoromethyl)cyclopropyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
4.25	7-(4-Fluorophenoxymethyl)-2-methyl-3-(2-methylcyclopropyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
4.26	trans-2-[2-[7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropyl]acetonitrile
4.27	7-(4-Fluorophenoxymethyl)-3-[2-(methoxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
4.28	3-(2-(fluoromethyl)cyclopropyl)-7-((4-fluorophenoxy)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
4.29	6-fluoro-7-((4-fluorophenoxy)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
4.30	7-(4-Fluorophenoxymethyl)-3-(3-hydroxypropyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
4.31	7-(4-Fluorophenoxymethyl)-3-(methoxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
4.32	7-((4-fluorophenoxy)methyl)-3-(3-hydroxyoxetan-3-yl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
4.33	7-(4-Fluorophenoxymethyl)-3-(4-hydroxybutan-2-yl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
4.34	7-(4-Fluorophenoxymethyl)-3-[2-(2-hydroxypropan-2-yl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
5.1	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-pyrimidin-5-yl-thiazolo[3,2-a]pyrimidin-5-one
5.2	7-(((4-fluorophenyl)(methyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
5.3	7-(((4-fluorophenyl)(2,2,2-trifluoroethyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
5.4	7-(((3,4-difluorophenyl)(ethyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
5.5	7-((ethyl(3-fluorophenyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
5.6	7-(((2,2-difluoroethyl)(4-fluorophenyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one

TABLE 1-continued

Ex.	Chemical Name
5.7	7-((ethyl(pyridine-2-yl)amino)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
5.8	7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(thiazol-2-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
5.9	7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(thiophen-2-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
5.10	3-(ethyl((2-methyl-5-oxo-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)amino)benzotrile
5.11	3-(2-aminopyridin-3-yl)-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
5.12	7-((ethyl(pyridine-2-yl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
5.13	7-((4-fluorophenyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
5.14	3-butyl-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
5.15	2-cyclopropyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-3-pyrimidin-5-yl-thiazolo[3,2-a]pyrimidin-5-one
5.16	2-ethyl-7-((ethyl(4-fluorophenyl)amino)methyl)-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
5.17	7-((ethyl(4-fluorophenyl)amino)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
5.18	7-((ethyl(4-fluorophenyl)amino)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 1)
5.19	7-((ethyl(4-fluorophenyl)amino)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 2)
5.20	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-thiazol-4-yl-thiazolo[3,2-a]pyrimidin-5-one
5.21	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one
5.22	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-phenyl-thiazolo[3,2-a]pyrimidin-5-one
5.23	7-[(N-ethyl-4-fluoro-anilino)methyl]-2,3-dimethyl-thiazolo[3,2-a]pyrimidin-5-one
5.24	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(5-fluoro-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
5.25	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(2-fluoro-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
5.26	3-cyclopropyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
5.27	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-pyrazin-2-yl-thiazolo[3,2-a]pyrimidin-5-one
5.28	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-5-one
5.29	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-isopropenyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
5.30	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-pyridazin-4-yl-thiazolo[3,2-a]pyrimidin-5-one
5.31	3-(5-chloro-3-pyridyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
5.32	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(4-pyridyl)thiazolo[3,2-a]pyrimidin-5-one
5.33	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(1-methylpyrazol-4-yl)thiazolo[3,2-a]pyrimidin-5-one
5.34	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(1H-pyrazol-4-yl)thiazolo[3,2-a]pyrimidin-5-one
5.35	5-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]pyridine-3-carbonitrile
5.36	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-[5-(trifluoromethyl)-3-pyridyl]thiazolo[3,2-a]pyrimidin-5-one
5.37	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-oxazol-2-yl-thiazolo[3,2-a]pyrimidin-5-one
5.38	7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidin-5-one
5.39	7-((ethyl(4-fluorophenyl)amino)methyl)-3-(trans-2-(fluoromethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
5.40	3-ethyl-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
5.41	7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-propyl-5H-thiazolo[3,2-a]pyrimidin-5-one
5.42	7-[[4-fluoro-N-(2-fluoroethyl)anilino]methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 1)
5.43	7-[[4-fluoro-N-(2-fluoroethyl)anilino]methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 2)
5.44	7-((ethyl(4-fluorophenyl)amino)methyl)-3-(furan-3-yl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
5.45	7-((ethyl(4-fluorophenyl)amino)methyl)-3-(furan-2-yl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
5.46	7-((5-fluoro-2-methylindolin-1-yl)methyl)-3-(furan-2-yl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one

TABLE 1-continued

Ex.	Chemical Name
5.47	7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(thiophen-3-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
5.48	7-((5-fluoro-2-methylindolin-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
5.49	7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(4-methylthiazol-2-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
5.50	7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(6-oxo-1,6-dihydropyridin-3-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
5.51	3-(6-aminopyridin-3-yl)-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
5.52	7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(prop-1-ynyl)-5H-thiazolo[3,2-a]pyrimidin-5-one
5.53	7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-vinyl-5H-thiazolo[3,2-a]pyrimidin-5-one
5.54	3-bromo-2-cyclopropyl-7-[(N-ethyl-4-fluoro-anilino)methyl]thiazolo[3,2-a]pyrimidin-5-one
5.55	3-(3,5-difluorophenyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
5.56	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(1-methylpyrazol-3-yl)thiazolo[3,2-a]pyrimidin-5-one
5.57	3-(2-amino-4-pyridyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
5.58	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(5-methoxy-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
6.1	7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-morpholino-5H-thiazolo[3,2-a]pyrimidin-5-one
6.2	3-(dimethylamino)-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
6.3	7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(pyrrolidin-1-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
7.1	7-(((3,4-difluorophenyl)(ethyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile
7.2	7-((ethyl(3-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile
7.3	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carbonitrile
7.4	7-[(N-ethyl-4-fluoro-anilino)methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carbonitrile
7.5	7-((5-fluoro-2-methylindolin-1-yl)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile
7.6	7-(((3-cyanophenyl)(ethyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile
7.7	7-((ethyl(pyridin-2-yl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile
7.8	2-methyl-7-((2-methylindolin-1-yl)methyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile
7.9	7-(((3,5-difluorophenyl)(ethyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile
8.1	6-[(N-ethyl-4-fluoro-anilino)methyl]-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one
8.2	2-[(N-ethyl-4-fluoro-anilino)methyl]-6,7,8,9-tetrahydropyrimido[2,1-b][1,3]benzothiazol-4-one
8.3	6-[(N-ethyl-4-fluoro-anilino)methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one
8.4	6-[(N-ethyl-4-fluoro-anilino)methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one (enantiomer 1)
8.5	6-[(N-ethyl-4-fluoro-anilino)methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one (enantiomer 2)
8.6	6-[(N-ethyl-4-fluoro-anilino)methyl]spiro[2,3-dihydrocyclopenta[3,4]thiazolo[1,4-a]pyrimidine-1,1'-cyclopentane]-8-one
8.7	6-[(N-ethyl-4-fluoro-anilino)methyl]-2,3-dihydrocyclopenta[3,4]thiazolo[1,4-a]pyrimidine-1,8-dione
8.8	6-[(N-ethyl-4-fluoro-anilino)methyl]-1,1-dimethyl-2,3-dihydrocyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one
8.9	2-[(N-ethyl-4-fluoro-anilino)methyl]-8,9-dihydro-6H-pyrano[3,4]thiazolo[1,4-a]pyrimidin-4-one
8.10	2-[(N-ethyl-4-fluoro-anilino)methyl]-7-methyl-8,9-dihydro-6H-pyrido[3,4]thiazolo[1,4-a]pyrimidin-4-one
8.11	6-[(N-ethyl-4-fluoro-anilino)methyl]-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidine-1-carboxamide
8.12	6-[(N-ethyl-4-fluoro-anilino)methyl]-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidine-1-carbonitrile
8.13	2-[6-[(N-ethyl-4-fluoro-anilino)methyl]-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-1-yl]acetonitrile
8.14	6-[(N-ethyl-4-fluoro-anilino)methyl]-1-(2-hydroxyethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one
8.15	2-((ethyl(4-fluorophenyl)amino)methyl)-6-(methoxymethyl)-7,8-dihydrocyclopenta[4,5]thiazolo[3,2-a]pyrimidin-4(6H)-one

TABLE 1-continued

Ex.	Chemical Name
8.16	2-[6-[(N-ethyl-4-fluoro-anilino)methyl]-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-1-yl]acetamide
9.1	3-cyclohexyl-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
9.2	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-isopropyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
9.3	3-cyclopentyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
9.4	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-tetrahydropyran-4-yl-thiazolo[3,2-a]pyrimidin-5-one
10.1	3-cyclobutyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
10.2	3-tert-butyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
10.3	3-acetyl-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
10.4	7-[(N-ethyl-4-fluoro-anilino)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
10.5	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-hydroxy-1-methyl-ethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
10.6	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-hydroxyethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
10.7	3-[(dimethylamino)methyl]-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
10.8	3-(azetidin-1-ylmethyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
10.9	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(pyrrolidin-1-ylmethyl)thiazolo[3,2-a]pyrimidin-5-one
10.10	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-[(3-hydroxyazetidin-1-yl)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
10.11	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(2,2,2-trifluoro-1-hydroxyethyl)thiazolo[3,2-a]pyrimidin-5-one
10.12	7-((ethyl(4-fluorophenyl)amino)methyl)-3-(hydroxymethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
10.13	7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(methoxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
10.14	7-[[Ethyl(4-fluorophenyl)amino]methyl]-2-methyl-3-(1H-pyrazol-1-ylmethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
10.15	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(methylsulfonylmethyl)thiazolo[3,2-a]pyrimidin-5-one
10.16	3-(ethoxymethyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
10.17	2-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]acetonitrile
10.18	3-tert-butyl-7-[(N-ethyl-4-fluoro-anilino)methyl]thiazolo[3,2-a]pyrimidin-5-one
10.19	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-hydroxy-1-methyl-ethyl)thiazolo[3,2-a]pyrimidin-5-one
10.20	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-hydroxyethyl)thiazolo[3,2-a]pyrimidin-5-one
10.21	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(6-oxa-1-azaspiro[3.3]heptan-1-ylmethyl)thiazolo[3,2-a]pyrimidin-5-one
10.22	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-[(2-hydroxyethylamino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
10.23	3-(ethyl-[(3-(hydroxymethyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl]amino)benzonitrile
10.24	3-[ethyl-[[3-(methoxymethyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl)methyl]amino]benzonitrile
10.25	7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-((methylthio)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one
11.1	3-chloro-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
11.2	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-fluoro-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
11.3	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
11.4	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-iodo-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
11.5	3-chloro-7-[(N-ethyl-4-fluoro-anilino)methyl]thiazolo[3,2-a]pyrimidin-5-one
11.6	7-[(N-ethyl-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
12.1	3-(1,3-dihydroxypropyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
12.2	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-fluoro-3-hydroxy-propyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
12.3	3-(1,3-dihydroxypropyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 1)
12.4	3-(1,3-dihydroxypropyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 2)
12.5	3-(1,2-dihydroxyethyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
13.1	7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(3-hydroxypropyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one

TABLE 1-continued

Ex.	Chemical Name
13.2	7-((Ethyl(4-fluorophenyl)amino)methyl)-3-(3-methoxypropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
13.3	7-((Ethyl(4-fluorophenyl)amino)methyl)-3-(4-hydroxybutyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
13.4	7-((Ethyl(4-fluorophenyl)amino)methyl)-3-(2-hydroxyethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
13.5	7-[[Ethyl(4-fluorophenyl)amino]methyl]-3-(2-methoxyethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
13.6	3-(7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)propanamide
13.7	3-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]propanenitrile
13.8	3-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N,N-dimethyl-propanamide
13.9	7-[[Ethyl(4-fluorophenyl)amino]methyl]-3-(3-hydroxy-3-methylbutyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one
14.1	2-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarboxamide
14.2	2-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarboxamide
15.1	7-(5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
15.2	7-((3,5-bis(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
15.3	7-((3-chloro-5-methyl-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
15.4	7-((5-chloro-3-methyl-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
15.5	3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one
15.6	7-((1H-indazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
15.7	7-((5-(2-hydroxyethyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
15.8	7-((3-(2-hydroxyethyl)-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
15.9	7-((3-cyclopropyl-4-fluoro-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
15.10	7-((5-cyclopropyl-4-fluoro-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
15.11	7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
15.12	7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
15.13	7-[[5-cyclopropyl-3-methyl-pyrazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 1)
15.14	7-[[5-cyclopropyl-3-methyl-pyrazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 2)
15.15	7-[[3-cyclopropyl-5-methyl-pyrazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 1)
15.16	7-[[3-cyclopropyl-5-methyl-pyrazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 2)
15.17	7-[[3,5-dicyclopropylpyrazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
15.18	7-((3,5-dimethyl-1H-pyrazol-1-yl)methyl)-6-fluoro-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
15.19	1-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-3-(trifluoromethyl)-1H-pyrazole-5-carbonitrile
15.20	7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-6-fluoro-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
15.21	1-[[3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl)methyl]-5-methyl-pyrazole-3-carbonitrile
15.22	7-[[3-(difluoromethyl)-5-methyl-pyrazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
15.23	7-[[6-fluoroindazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
15.24	7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-isopropyl-thiazolo[3,2-a]pyrimidin-5-one
15.25	7-[[3,5-dimethylpyrazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
15.26	7-[[5-fluoroindazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
15.27	5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
15.28	7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
15.29	7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one

TABLE 1-continued

Ex.	Chemical Name
15.30	7-((3,5-bis(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
15.31	2-methyl-3-(pyrimidin-5-yl)-7-((5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one
15.32	2-methyl-7-((5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
15.33	7-((3,5-bis(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
15.34	7-((1H-indazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
16.1	7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
16.2	7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
16.3	7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-2-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
16.4	7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-2-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
16.5	7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
16.6	2-cyclopropyl-7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
16.7	2-cyclopropyl-7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
17.1	2-methyl-5-oxo-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile
17.2	7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile
17.3	7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile
18.1	3-(1-hydroxyethyl)-2-methyl-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one
18.2	3-acetyl-7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
18.3	3-acetyl-7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
18.4	3-(2-hydroxypropan-2-yl)-2-methyl-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one
18.5	7-((5-fluoro-3-methyl-1H-indazol-1-yl)methyl)-3-(1-hydroxyethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
19.1	7-(1-(5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)ethyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
19.2	7-(1-(5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)ethyl)-3-(2-(hydroxymethyl)-1-methylcyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
20.1	3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile
20.2	3-((6-fluoro-3-(2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile
20.3	2-fluoro-3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile
20.4	3-(trans-2-(hydroxymethyl)cyclopropyl)-7-(isoquinolin-4-ylmethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
20.5	3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-7-(3-(trifluoromethyl)benzyl)-5H-thiazolo[3,2-a]pyrimidin-5-one
20.6	3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-4-methylbenzotrile
20.7	4-fluoro-3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile
20.8	3-fluoro-5-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile
20.9	2-fluoro-5-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile
20.10	3-[[3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl)methyl]-4-methoxy-benzotrile
20.11	3-(2-(hydroxymethyl)cyclopropyl)-2-methyl-7-(4-(trifluoromethyl)benzyl)-5H-thiazolo[3,2-a]pyrimidin-5-one
20.12	4-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)picolinonitrile
20.13	4-cyclopropyl-3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile
21.1	7-(3-cyanobenzyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.2	7-(3-cyano-2-fluorobenzyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.3	7-(3-chloro-2-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide

TABLE 1-continued

Ex.	Chemical Name
21.4	N-ethyl-7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.5	N-ethyl-7-(2-fluoro-3-methylbenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.6	7-(2-chloro-5-cyanobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.7	N-ethyl-2-methyl-5-oxo-7-(3-(trifluoromethyl)benzyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.8	7-(3-cyanobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.9	7-(3-cyano-2-fluorobenzyl)-2-cyclopropyl-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.10	7-(3-cyano-5-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.11	7-(3-cyanobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.12	N-ethyl-7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.13	7-(3-cyano-2-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.14	7-[(3-chloro-2-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.15	N,2-dimethyl-5-oxo-7-[[3-(trifluoromethyl)phenyl]methyl]thiazolo[3,2-a]pyrimidine-3-carboxamide
21.16	7-[(3-chlorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.17	7-[[2-cyclopropyl-5-(trifluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.18	N-ethyl-2-methyl-5-oxo-7-((6-(trifluoromethyl)pyridine-2-yl)methyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.19	N-ethyl-7-(2-ethyl-4,5-difluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.20	7-[[3-(difluoromethyl)-2-fluoro-phenyl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.21	N-ethyl-7-(2-ethyl-4,5-difluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.22	7-((6-cyanopyridin-2-yl)methyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.23	7-[[3-(difluoromethyl)-2-fluoro-phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.24	7-[[2-fluoro-3-(hydroxymethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.25	7-(3-cyclopropyl-2-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.26	7-(3-cyano-2-fluorobenzyl)-N-ethyl-6-fluoro-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.27	7-(3-cyano-2-fluorobenzyl)-6-fluoro-N,N-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.28	7-(3-cyano-2-fluorobenzyl)-N-ethyl-6-fluoro-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.29	6-fluoro-7-(2-fluoro-3-(trifluoromethyl)benzyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.30	7-(5-cyano-2-methylbenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.31	7-(5-cyano-2-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.32	7-(2-chloro-5-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.33	7-(3-cyano-2-fluorobenzyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.34	N-ethyl-7-(5-fluoro-2-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.35	N-ethyl-2-methyl-7-(naphthalen-1-ylmethyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.36	N-ethyl-7-(5-fluoro-2-methylbenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.37	7-(3-cyano-4-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.38	N-ethyl-2-methyl-7-((1-methyl-1H-indazol-4-yl)methyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.39	N-ethyl-2-methyl-7-(3-(methylsulfonamido)benzyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.40	7-(5-cyano-2-(trifluoromethyl)benzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.41	7-(4-chloro-2-methylbenzyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.42	7-(2,5-difluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide

TABLE 1-continued

Ex.	Chemical Name
21.43	7-(3-cyanobenzyl)-2-cyclopropyl-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.44	N-ethyl-7-(2-fluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.45	7-(2,3-difluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.46	N-ethyl-7-(3-fluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.47	7-[(3-chloro-4-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.48	7-[(2,5-dichlorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.49	N,2-dimethyl-5-oxo-7-[[3-(trifluoromethoxy)phenyl]methyl]thiazolo[3,2-a]pyrimidine-3-carboxamide
21.50	7-[(5-cyano-2-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.51	7-[(3-chloro-5-cyano-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.52	7-[(3-cyclopropylphenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.53	7-[(2,5-difluorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.54	7-[(3,4-difluorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.55	7-[(2,3-difluorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.56	7-[(4-chloro-2-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.57	7-[(2,4-dichlorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.58	7-[(3-fluoro-4-methyl-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.59	7-[[4-fluoro-2-(trifluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.60	7-[(2-cyclopropyl-4-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.61	7-(5-cyano-2-(2-fluoroethyl)benzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.62	7-(2-chloro-3-cyanobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.63	N-ethyl-7-(6-ethyl-2,3-difluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.64	7-(5-cyano-2-ethylbenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.65	7-(2-cyclopropyl-5-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.66	7-(5-cyano-2-cyclopropylbenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.67	N-ethyl-7-(5-fluoro-2-propylbenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
21.68	7-[[2-fluoro-3-(fluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
22.1	N-ethyl-7-(1-(2-fluoro-3-(trifluoromethyl)phenyl)ethyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
22.2	N-ethyl-7-(2-fluoro-3-(trifluoromethyl)benzyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
23.1	3-((3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile
23.2	3-((3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzotrile
23.3	6-((3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)picolinotrile
23.4	3-((3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-4-methoxybenzotrile
24.1	2-methyl-3-(pyrimidin-5-yl)-7-(3-(trifluoromethyl)benzyl)-5H-thiazolo[3,2-a]pyrimidin-5-one
24.2	7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
24.3	2-fluoro-3-((2-methyl-5-oxo-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile
24.4	3-((3-cyclopropyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzotrile
24.5	7-(isoquinolin-4-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one
24.6	3-((2-methyl-5-oxo-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile
24.7	7-(5-fluoro-2-methoxybenzyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one

TABLE 1-continued

Ex.	Chemical Name
24.8	2-fluoro-3-((3-(furan-2-yl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile
24.9	3-bromo-2-methyl-7-(3-(methylsulfonyl)benzyl)-5H-thiazolo[3,2-a]pyrimidin-5-one
25.1	7-(3-cyano-2-fluorobenzyl)-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
25.2	N-(cyanomethyl)-7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide
26.1	7-(3-cyanobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile
26.2	7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile
27.1	10-(4-fluorophenoxymethyl)-3-(hydroxymethyl)-7-thia-1,9-diazatricyclo[6.4.0.0 ² [2,6]]dodeca-2(6),8,10-trien-12-one
27.2	10-(4-Fluorophenoxymethyl)-3-(2-hydroxyethyl)-7-thia-1,9-diazatricyclo[6.4.0.0 ² [2,6]]dodeca-2(6),8,10-trien-12-one
27.3	7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-methylsulfonyl-thiazolo[3,2-a]pyrimidin-5-one
27.4	3-(hydroxymethyl)-10-[[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-7-thia-1,9-diazatricyclo[6.4.0.0 ² [2,6]]dodeca-2(6),8,10-trien-12-one
27.5	10-[[5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-7-thia-1,9-diazatricyclo[6.4.0.0 ² [2,6]]dodeca-2(6),8,10-trien-12-one
27.6	10-[[3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-7-thia-1,9-diazatricyclo[6.4.0.0 ² [2,6]]dodeca-2(6),8,10-trien-12-one
27.7	10-[[5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-3,3-dimethyl-7-thia-1,9-diazatricyclo[6.4.0.0 ² [2,6]]dodeca-2(6),8,10-trien-12-one
27.8	10-[[3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-3,3-dimethyl-7-thia-1,9-diazatricyclo[6.4.0.0 ² [2,6]]dodeca-2(6),8,10-trien-12-one
27.9	10-[[3,5-bis(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-3,3-dimethyl-7-thia-1,9-diazatricyclo[6.4.0.0 ² [2,6]]dodeca-2(6),8,10-trien-12-one
27.10	3-((3-acetyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzotrile
27.11	7-(2-fluoro-3-(trifluoromethyl)benzyl)-3-(1-hydroxyethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 1)
27.12	7-(2-fluoro-3-(trifluoromethyl)benzyl)-3-(1-hydroxyethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 2)
27.13	3-((3-acetyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile
27.14	7-(2-fluoro-3-(trifluoromethyl)benzyl)-3-(hydroxymethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one
27.15	7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-3-((methylamino)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one

In certain embodiments, the compound of Formula I or II is a compound selected from the group consisting of the compounds in Table 2, and pharmaceutically acceptable salts thereof:

TABLE 2

Ex.	Chemical Name
1	7-[[3-cyclopropyl-2-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
2	N-ethyl-2-methyl-5-oxo-7-[(2,3,6-trifluorophenyl)methyl]thiazolo[3,2-a]pyrimidine-3-carboxamide
3	2-fluoro-3-[[2-methyl-3-oxazol-2-yl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl)methyl]benzotrile
4	7-[[5-cyano-3-cyclopropyl-2-fluoro-phenyl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
5	N-ethyl-7-[(2-fluoro-3-methoxy-phenyl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
6	7-[[3-cyclopropyl-2-fluoro-phenyl)methyl]-6-fluoro-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
7	2-[7-[(3-chloro-2-fluoro-phenyl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide
8	7-[[3-chloro-2-fluoro-phenyl)methyl]-N-ethyl-6-fluoro-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
	7-[[4,5-difluoro-2-methoxy-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
10	2-fluoro-3-[[2-methyl-3-(2-methylcyclopropyl)-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl)methyl]benzotrile
11	2-[7-[(3-cyclopropyl-2-fluoro-phenyl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide
13	N-ethyl-6-fluoro-7-[[2-fluoro-3-(trifluoromethyl)phenyl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
14	7-[[2-chloro-5-(trifluoromethyl)phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide

TABLE 2-continued

Ex.	Chemical Name
15	7-[(3-cyano-2-fluoro-phenyl)methyl]-6-fluoro-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
16	7-[(2-chloro-4,5-difluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
17	7-[[4,5-difluoro-2-(2-fluoroethyl)phenyl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
18	2-[7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide
19	2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide
20	7-[[5-chloro-3-methyl-pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
21	7-[(3-chloro-5-methyl-pyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
22	7-[(3-chloro-5-cyclopropyl-pyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
23	7-[[5-chloro-3-cyclopropyl-pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
24	3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-7-[[4-(trifluoromethyl)thiazol-2-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one
25	2-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide
26	N-ethyl-2-methyl-7-[[5-methyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
27	N-ethyl-2-methyl-7-[[3-methyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
28	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
29	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
30	7-[[3-(difluoromethyl)-2-fluoro-phenyl]methyl]-6-fluoro-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
31	7-[[3-(difluoromethyl)-2-fluoro-phenyl]methyl]-N-ethyl-6-fluoro-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
32	7-(4-bicyclo[4.2.0]octa-1,3,5-trienylmethyl)-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
33	N-ethyl-7-[[2-fluoro-3-(1-hydroxycyclopropyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
35	N-ethyl-7-[[2-fluoro-3-(1-fluorocyclopropyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
36	N-ethyl-7-[[2-fluoro-3-[1-(fluoromethyl)vinyl]phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
37	7-[(2-ethynyl-4,5-difluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
38	2-fluoro-3-[[2-methyl-5-oxo-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzotrile
39	3-[[3-(2,2-difluorocyclopropyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluoro-benzotrile
40	N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
41	7-[(3-cyano-2-fluoro-phenyl)methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
42	7-[[3-(difluoromethyl)-2-fluoro-phenyl]methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
43	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
44	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
45	N-ethyl-7-[[5-methyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
46	N-ethyl-7-[[3-methyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
47	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-(1H-imidazol-2-yl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
49	7-[(3-cyano-2-fluoro-5-methyl-phenyl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
50	7-[(3-chloro-2-fluoro-phenyl)methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
51	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
52	2-methyl-7-[[5-methyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
53	7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
54	7-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
55	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide

TABLE 2-continued

Ex.	Chemical Name
56	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
57	7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
58	7-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
60	2-chloro-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
61	N-methyl-7-[[5-methyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
62	N-methyl-7-[[3-methyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
63	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
64	7-[[3-chloro-2-fluoro-phenyl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
65	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
66	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
67	2-methyl-7-[[3-methyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
69	2-chloro-7-[[3-cyano-2-fluoro-phenyl]methyl]-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
70	7-[[3-cyano-2-fluoro-phenyl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
71	N-ethyl-2-[[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]acetamide
72	N,2-dimethyl-5-oxo-7-[[4-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidine-3-carboxamide
74	3-[[2-chloro-5-oxo-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluoro-benzonitrile
75	2-[[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide
76	7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
78	2-[[7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide
79	2-[[7-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide
80	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(methylsulfonylmethyl)thiazolo[3,2-a]pyrimidin-5-one
81	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-(1H-imidazol-2-ylmethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
82	N-ethyl-7-[[N-ethyl-4-fluoro-anilino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
83	N-ethyl-7-[[N-ethyl-4-fluoro-anilino]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
84	2-fluoro-3-[[3-(2-methylcyclopropyl)-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzotrile
85	3-[[2-chloro-3-(2-methylcyclopropyl)-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluoro-benzonitrile
87	N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-isopropyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
88	2-fluoro-3-[[5-oxo-2-(trifluoromethyl)-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzotrile
89	6-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one
90	6-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one
91	N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methoxy-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
92	6-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one
93	2-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-N-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
94	2-[[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
95	6-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one
96	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
97	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
98	2-[[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide
99	2-[[7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide

TABLE 2-continued

Ex.	Chemical Name
100	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-(2-hydroxycyclopropyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
102	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
103	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
104	2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
105	2-cyano-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
106	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-N-isopropyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
108	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1-hydroxy-1-methyl-ethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
109	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1-hydroxy-1-methyl-ethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
110	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1-hydroxyethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
111	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1-hydroxyethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
112	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-N-sec-butyl-thiazolo[3,2-a]pyrimidine-3-carboxamide
113	3-[[3-(azetidin-1-yl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluoro-benzonitrile
114	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carbonitrile
115	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carbonitrile
116	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-cyclopropyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
117	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-cyclopropyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
118	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(2-methylcyclopropyl)thiazolo[3,2-a]pyrimidin-5-one
119	3-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
120	3-acetyl-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
121	2-chloro-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-(2-methylcyclopropyl)thiazolo[3,2-a]pyrimidin-5-one
122	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-pyrimidin-5-yl-thiazolo[3,2-a]pyrimidin-5-one
123	7-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
124	7-[[5-(5-chloro-2-pyridyl)-methyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
125	2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 1)
126	2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 2)
127	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methoxy-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
128	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methoxy-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
129	7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methoxy-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
130	7-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methoxy-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
131	2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
132	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(1H-pyrazol-5-yl)thiazolo[3,2-a]pyrimidin-5-one
133	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(1H-pyrazol-5-yl)thiazolo[3,2-a]pyrimidin-5-one
134	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one
135	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-isopropenyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
136	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 1)
137	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 2)
138	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(5-fluoro-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
139	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(5-fluoro-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
140	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-propanoyl-thiazolo[3,2-a]pyrimidin-5-one

TABLE 2-continued

Ex.	Chemical Name
141	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-propanoyl-thiazolo[3,2-a]pyrimidin-5-one
142	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-thiazol-2-yl-thiazolo[3,2-a]pyrimidin-5-one
143	N-ethyl-7-[[[(5-fluoro-2-pyridyl)-methyl-amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
144	2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
145	2-[7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
146	2-fluoro-3-[[5-oxo-2-(trifluoromethyl)-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzoxonitrile
147	N-ethyl-7-[[ethyl-(5-fluoro-2-pyridyl)amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
148	N-ethyl-7-[[ethyl(2-pyridyl)amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
149	3-(5-chloro-3-pyridyl)-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
150	3-(5-chloro-3-pyridyl)-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
151	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-thiazol-4-yl-thiazolo[3,2-a]pyrimidin-5-one
152	7-[[5-chloro-2-pyridyl]-ethyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
153	7-[(5-cyclopropylthiazol-1-yl)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
154	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-[2-methylcyclopropyl]thiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 1)
155	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-[2-methylcyclopropyl]thiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 2)
156	2-ethoxy-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
157	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-fluoro-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
158	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-oxazol-2-yl-thiazolo[3,2-a]pyrimidin-5-one
159	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-oxazol-2-yl-thiazolo[3,2-a]pyrimidin-5-one
160	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
161	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
162	2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
163	7-[[[(5-chloro-2-pyridyl)-methyl-amino]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
164	7-[[3,5-dichloropyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
165	N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(2,2,2-trifluoroethoxy)thiazolo[3,2-a]pyrimidine-3-carboxamide
166	3-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
167	3-acetyl-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
168	3-[(4-chloropyrazol-1-yl)methyl]-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
169	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-isopropenyl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
170	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1H-imidazol-2-yl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
171	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-pyrimidin-5-yl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
172	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-pyrimidin-5-yl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
173	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-pyridyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
174	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-2-(2,2,2-trifluoroethoxy)thiazolo[3,2-a]pyrimidine-3-carboxamide
175	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethoxy-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
176	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethoxy-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
177	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(2-methylpropanoyl)thiazolo[3,2-a]pyrimidin-5-one
178	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(2-methylpropanoyl)thiazolo[3,2-a]pyrimidin-5-one
179	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-N-[(1R)-1-methylpropyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide

TABLE 2-continued

Ex.	Chemical Name
180	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-N-[(1S)-1-methylpropyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
191	7-[[4-chloro-2-pyridyl]-ethyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
182	7-[[5-fluoro-2-pyridyl]-methyl-amino]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
183	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(methoxymethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
184	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-cyclopropyl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
185	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(methylsulfonylmethyl)thiazolo[3,2-a]pyrimidin-5-one
186	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(pyrazol-1-ylmethyl)thiazolo[3,2-a]pyrimidin-5-one
187	2-[7-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
188	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-cyclobutyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
189	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-cyclobutyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
190	2-[7-[[5-chloro-2-pyridyl]-methyl-amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
191	7-[[ethyl-(5-fluoro-2-pyridyl)amino]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
192	3-chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
193	3-chloro-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
194	2-[7-[[4,5-difluoro-2-methoxy-phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
195	2-[7-[[ethyl-(5-fluoro-2-pyridyl)amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
196	7-[[5-chloro-2-pyridyl]-ethyl-amino]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
197	5-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]pyridine-3-carbonitrile
198	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(2,2,2-trifluoroacetyl)thiazolo[3,2-a]pyrimidin-5-one
199	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-methylcyclopropyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
200	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-methylcyclopropyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
201	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-pyrimidin-5-yl-thiazolo[3,2-a]pyrimidin-5-one
202	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one
203	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(2,2,2-trifluoro-1-hydroxy-ethyl)thiazolo[3,2-a]pyrimidin-5-one
204	7-[[5-bromo-2-pyridyl]-ethyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
205	N-ethyl-7-[[ethyl-(5-fluoro-2-pyridyl)amino]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
206	3-(azetid-1-yl)-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
207	2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 1)
208	2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 2)
209	2-[7-[[5-methoxy-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
210	2-[7-[[3-methoxy-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
211	3-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-thiazolo[3,2-a]pyrimidin-5-one
212	3-acetyl-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-thiazolo[3,2-a]pyrimidin-5-one
213	2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
214	2-[7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
215	3-bromo-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
216	3-bromo-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
217	3-chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methoxy-thiazolo[3,2-a]pyrimidin-5-one
218	3-chloro-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methoxy-thiazolo[3,2-a]pyrimidin-5-one

TABLE 2-continued

Ex.	Chemical Name
219	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-methylsulfanyl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
220	2-[7-[(N-ethyl-4-fluoro-anilino)methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-1-(hydroxymethyl)cyclopropanecarbonitrile
221	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(cyclopropylmethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
222	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-hydroxy-1-methyl-ethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
223	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-hydroxy-1-methyl-ethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
224	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(cyclopropanecarbonyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
225	3-bromo-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one
226	3-bromo-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one
227	7-[[3-amino-5-chloro-pyrazol-1-yl]methyl]-3-bromo-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
228	7-[[5-amino-3-chloro-pyrazol-1-yl]methyl]-3-bromo-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
229	2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]acetoneitrile
230	N-ethyl-7-[[5-(5-fluoro-2-pyridyl)-methyl-amino]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
231	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3,3-difluoroazetid-1-yl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
233	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methoxy-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one
234	2-[7-[[5-bromo-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
235	3-chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one
236	3-chloro-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one
237	2-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
238	2-acetyl-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
239	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
240	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(1H-1,2,4-triazol-5-yl)thiazolo[3,2-a]pyrimidin-5-one
241	2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
242	3-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one
243	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2,2,2-trifluoroethoxy)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
244	3-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methoxy-thiazolo[3,2-a]pyrimidin-5-one
245	3-acetyl-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methoxy-thiazolo[3,2-a]pyrimidin-5-one
246	3-bromo-7-[[5-chloro-3-nitro-pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
247	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1H-pyrazol-5-yl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
248	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-thiazol-4-yl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
249	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(5-fluoro-3-pyridyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
250	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-propanoyl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
251	2-[7-[[3,5-dichloropyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
251A	2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]propanenitrile
252	2-[7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]propanenitrile
253	2-fluoro-3-[[3-[2-methylcyclopropyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzonitrile
254	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-fluoroazetid-1-yl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
255	3-(5-chloro-3-pyridyl)-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
256	7-[[3,5-dichloropyrazol-1-yl]methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
257	3-[[3-acetyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluoro-benzonitrile

TABLE 2-continued

Ex.	Chemical Name
258	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(difluoromethyl)-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
259	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(difluoromethyl)-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
260	(Z)-3-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enenitrile
261	(E)-3-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enamide
262	(E)-3-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enenitrile
263	(Z)-3-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enenitrile
264	(E)-3-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enamide
265	N-ethyl-7-[[5-isobutyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
266	2-[2-chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 1)
267	2-[2-chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 2)
268	2-[2-chloro-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
269	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 1)
270	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 2)
271	2-[7-[(4-chloro-1-methyl-pyrazol-3-yl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
272	2-[2-methyl-5-oxo-7-[[3-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 1)
273	2-[2-methyl-5-oxo-7-[[3-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 2)
274	2-[7-[[5-bromo-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 1)
275	2-[7-[[5-bromo-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 2)
276	2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-6-fluoro-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 1)
277	2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-6-fluoro-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 2)
278	2-[2-methyl-7-[[1-methyl-4-(trifluoromethyl)imidazol-2-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
279	(E)-3-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enenitrile
280	7-[(4-fluorophenoxy)methyl]-3-[[2-hydroxyethyl(methyl)amino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
281	7-[(4-fluorophenoxy)methyl]-3-[[2-hydroxyethylamino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
282	2-[7-[(4-fluorophenoxy)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N,N-dimethyl-acetamide
283	7-[(2-cyano-4,5-difluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
284	7-[(2-cyclopropyl-4,5-difluoro-phenyl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
285	3-[2-(azetidin-1-yl)-2-oxo-ethyl]-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
286	7-[(4-fluorophenoxy)methyl]-2-methyl-3-(4H-1,2,4-triazol-3-ylmethyl)thiazolo[3,2-a]pyrimidin-5-one
287	2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-propanamide
288	3-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-propanamide
289	7-[[5-chloro-2-(trifluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
290	7-[[5-ethyl-1,3-benzoxazol-6-yl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
291	7-[(3-chloropyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
292	7-[(5-chloropyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
293	2-[7-[[3-cyano-2-fluoro-phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide
294	N-ethyl-7-[[2-fluoro-3-(1-hydroxypropyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
295	7-[[4,5-difluoro-2-oxazol-2-yl-phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
296	2-fluoro-3-[(2-methyl-5-oxo-3-propanoyl-thiazolo[3,2-a]pyrimidin-7-yl)methyl]benzonitrile

TABLE 2-continued

Ex.	Chemical Name
297	7-[[4,5-difluoro-2-(trifluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
299	7-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
300	7-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
301	7-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
303	3-[[2-chloro-3-cyclopropyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl)methyl]-2-fluorobenzonitrile
304	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(pyrazol-1-yl)methylthiazolo[3,2-a]pyrimidin-5-one
305	N,2-dimethyl-7-[[3-methyl-4-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
306	N,2-dimethyl-7-[[5-methyl-4-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
307	2-fluoro-3-[(8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-6-yl)methyl]benzonitrile
308	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-[hydroxy(thiazol-2-yl)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
309	2-fluoro-3-[(3-methyl-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-6-yl)methyl]benzonitrile
310	2-[7-[(4-fluorophenoxy)methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]-N-methylacetamide
312	2-fluoro-3-[[1-(hydroxymethyl)-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-6-yl]methyl]benzonitrile
313	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-N-(2-hydroxy-1-methyl-ethyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
314	3-[[3-(2,3-dimethylcyclopropyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl)methyl]-2-fluoro-benzonitrile
315	6-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one
316	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(2-oxa-6-azaspiro[3.3]heptan-6-yl)thiazolo[3,2-a]pyrimidin-5-one
317	N-ethyl-6-fluoro-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
318	7-[(4-fluorophenoxy)methyl]-5-oxo-N-(2,2,2-trifluoroethyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
319	N-cyclopentyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
320	7-[[4,5-dichloropyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
321	7-[[3,4-dichloropyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
322	N-ethyl-2-methyl-7-[[methyl(thiazol-2-yl)amino]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
323	7-[[4-chloropyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
324	N-ethyl-2-methyl-7-[[methyl-(1-methylpyrazol-4-yl)amino]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
325	7-[(4-fluorophenoxy)methyl]-2-(trifluoromethyl)-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-5-one
326	7-[[3-ethoxy-2-pyridyl)-methyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
327	7-[[3,5-dimethylisoxazol-4-yl)-methyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
328	3-cyclopropyl-7-[(4-fluorophenoxy)methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
329	7-[(4-fluorophenoxy)methyl]-3-pyrimidin-5-yl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
330	7-[(4-fluorophenoxy)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
331	7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(4-methyl-1,2,4-triazol-3-yl)thiazolo[3,2-a]pyrimidin-5-one
332	2-fluoro-3-[[5-oxo-2-(trifluoromethyl)-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzonitrile
333	N-ethyl-7-[[ethyl(4-pyridyl)amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
334	N-ethyl-7-[[ethyl(3-pyridyl)amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
335	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-methoxy-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
336	2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
337	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-isopropenyl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
338	3-(5-chloro-3-pyridyl)-7-[[5-chloro-2-pyridyl)-methyl-amino]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one

TABLE 2-continued

Ex.	Chemical Name
339	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-pyridyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
340	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-2-(2,2,2-trifluoroethoxy)thiazolo[3,2-a]pyrimidine-3-carboxamide
341	7-[[3-chloro-6-(trifluoromethyl)-2-pyridyl]methyl]-2-methyl-3-(2-methylcyclopropyl)thiazolo[3,2-a]pyrimidin-5-one
342	7-[[5-chloro-2-pyridyl]oxymethyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
343	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one
344	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-pyrrolidin-1-yl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
345	N-ethyl-7-[[5-methoxy-2-pyridyl]-methyl-amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
346	3-(2-chloro-3-pyridyl)-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
347	3-(2-chloro-3-pyridyl)-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
348	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one
349	7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one
350	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-methoxyazetidin-1-yl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
351	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(cyclopropylmethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
352	5-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]pyridine-3-carbonitrile
353	2-fluoro-3-[[3-[2-methylcyclopropyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzotrile
354	7-[[3,5-diisopropylpyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
355	2-[7-[[4-chloro-2-methyl-pyrazol-3-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
356	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-methylazetidin-1-yl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
357	2-[7-[[3,5-dichloropyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 1)
358	2-[7-[[3,5-dichloropyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 2)
359	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(2-hydroxyethyl)cyclopropyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 1)
360	7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(2-hydroxyethyl)cyclopropyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 2)
361	3-chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
362	5-chloro-1-[[3-[2-methylcyclopropyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]pyrazole-3-carbonitrile (cis enantiomer1)
363	5-chloro-1-[[3-[2-methylcyclopropyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]pyrazole-3-carbonitrile (cis enantiomer2)
364	5-chloro-2-[[3-[[2-methylcyclopropyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]pyrazole-3-carbonitrile
365	7-[[5-ethoxy-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
366	N-ethyl-7-[[5-isobutyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
367	2-[2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropyl]acetoneitrile (trans enantiomer 1)
368	2-[2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropyl]acetoneitrile (trans enantiomer 2)

In certain embodiments, the compound of Formula III is a compound selected from the group consisting of the compounds in Table 3, and pharmaceutically acceptable salts thereof:

TABLE 3

Ex. Chemical Name
12 3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-7-[3-(trifluoromethyl)phenoxy]thiazolo[3,2-a]pyrimidin-5-one
34 6-fluoro-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-7-[3-(trifluoromethyl)phenoxy]thiazolo[3,2-a]pyrimidin-5-one
48 N-ethyl-7-[2-fluoro-3-(trifluoromethyl)phenoxy]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
59 7-[2-fluoro-3-(trifluoromethyl)phenoxy]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
68 7-(3-cyano-2-fluoro-phenoxy)-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
73 3-cyclopropyl-7-[2-fluoro-3-(trifluoromethyl)phenoxy]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
77 2-fluoro-3-[2-methyl-5-oxo-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-7-yl]oxy-benzonitrile
86 7-[2-fluoro-3-(trifluoromethyl)phenoxy]-2-methyl-3-(2-methylcyclopropyl)thiazolo[3,2-a]pyrimidin-5-one
101 2-[7-[2-fluoro-3-(trifluoromethyl)phenoxy]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
107 2-fluoro-3-[2-methyl-3-(2-methylcyclopropyl)-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]oxy-benzonitrile
298 N,2-dimethyl-5-oxo-7-[3-(trifluoromethyl)phenoxy]thiazolo[3,2-a]pyrimidine-3-carboxamide
302 N-ethyl-2-methyl-5-oxo-7-[3-(trifluoromethyl)phenoxy]thiazolo[3,2-a]pyrimidine-3-carboxamide
311 2-fluoro-3-[(8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-6-yl)oxy]benzonitrile

disorder or disease, causing regression of the disorder or disease, relieving a condition caused by the disease or disorder, or stopping the symptoms of the disease or disorder.

Pharmaceutical Description

As used herein, the term “subject” encompasses mammals and non-mammals. Examples of mammals include, but are not limited to, any member of the Mammalian class: humans; non-human primates such as chimpanzees, and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, swine; domestic animals such as rabbits, dogs, and cats; and laboratory animals including rodents, such as rats, mice and guinea pigs, and the like. Examples of non-mammals include, but are not limited to, birds, fish and the like. In one embodiment of the present invention, the mammal is a human.

“Patient” encompasses a human or animal subject.

The term “inhibitor” refers to a molecule such as a compound, a drug, an enzyme activator, or a hormone that blocks or otherwise interferes with a particular biologic activity.

The term “modulator” refers to a molecule, such as a compound of the present invention, that increases or decreases, or otherwise affects the activity of a given enzyme or protein.

As used herein, the terms “treat” or “treatment” encompass both “preventative” and “curative” treatment. “Preventative” treatment is meant to indicate a postponement of development of a disease, a symptom of a disease, or medical condition, suppressing symptoms that may appear, or reducing the risk of developing or recurrence of a disease or symptom. “Curative” treatment includes reducing the severity of or suppressing the worsening of an existing disease, symptom, or condition. Thus, treatment includes ameliorating or preventing the worsening of existing disease symptoms, preventing additional symptoms from occurring, ameliorating or preventing the underlying metabolic causes of symptoms, inhibiting the disorder or disease, e.g., arresting the development of the disorder or disease, relieving the

The terms “effective amount” or “therapeutically effective amount” refer to a sufficient amount of the agent to provide the desired biological result. That result can be reduction and/or alleviation of the signs, symptoms, or causes of a disease or medical condition, or any other desired alteration of a biological system. For example, an “effective amount” for therapeutic use is the amount of a compound, or of a composition comprising the compound, that is required to provide a clinically relevant change in a disease state, symptom, or medical condition. An appropriate “effective” amount in any individual case may be determined by one of ordinary skill in the art using routine experimentation. Thus, the expression “effective amount” generally refers to the quantity for which the active substance has a therapeutically desired effect. Effective amounts or doses of the compounds of the embodiments may be ascertained by routine methods, such as modeling, dose escalation, or clinical trials, taking into account routine factors, e.g., the mode or route of administration or drug delivery, the pharmacokinetics of the agent, the severity and course of the infection, the subject’s health status, condition, and weight, and the judgment of the treating physician. An exemplary dose is in the range of about 1 µg to 2 mg of active agent per kilogram of subject’s body weight per day, preferably about 0.05 to 100 mg/kg/day, or about 1 to 35 mg/kg/day, or about 0.1 to 10 mg/kg/day. The total dosage may be given in single or divided dosage units (e.g., BID, TID, QID).

Once improvement of the patient’s disease has occurred, the dose may be adjusted for preventative or maintenance treatment. For example, the dosage or the frequency of administration, or both, may be reduced as a function of the symptoms, to a level at which the desired therapeutic or prophylactic effect is maintained. Of course, if symptoms have been alleviated to an appropriate level, treatment may cease. Patients may, however, require intermittent treatment

on a long-term basis upon any recurrence of symptoms. Patients may also require chronic treatment on a long-term basis.

A pharmaceutical composition according to the invention comprises at least one compound of Formula (I), or a pharmaceutically acceptable salt thereof. The pharmaceutical compositions may further comprise one or more pharmaceutically-acceptable excipients. A pharmaceutically-acceptable excipient is a substance that is non-toxic and otherwise biologically suitable for administration to a subject. Such excipients facilitate administration of the compounds described herein and are compatible with the active ingredient. Examples of pharmaceutically-acceptable excipients include stabilizers, lubricants, anti-caking agents, glidants, surfactants, diluents, anti-oxidants, binders, chelating agents, coating agents, coloring agents, bulking agents, emulsifiers, buffers, pH modifiers, or taste-modifying agents. In preferred embodiments, pharmaceutical compositions according to the embodiments are sterile compositions. Sterile compositions include compositions that are in accord with national and local regulations governing such compositions. Pharmaceutical compositions may be prepared using compounding techniques known or that become available to those skilled in the art.

The pharmaceutical compositions and compounds described herein may be formulated as solutions, emulsions, suspensions, dispersions, or inclusion complexes such as cyclodextrins in suitable pharmaceutical solvents or carriers, or as pills, tablets, lozenges, suppositories, sachets, dragees, granules, powders, powders for reconstitution, or capsules along with solid carriers according to conventional methods known in the art for preparation of various dosage forms. Pharmaceutical compositions of the embodiments may be administered by a suitable route of delivery, such as oral, parenteral, rectal, nasal, topical, or ocular routes, or by inhalation. Preferably, the compositions are formulated for intravenous or oral administration.

A further embodiment of the invention is a method of preparing a pharmaceutical formulation comprising mixing at least one compound of the present invention, and, optionally, one or more pharmaceutically acceptable excipients.

In certain aspects, the invention relates to methods of treating diseases or conditions mediated by activation or deactivation of NMDA receptors, or which are generally mediated by NMDA receptor activity. Such disease or condition is one or more selected from the group consisting of pain, neuropathic pain, inflammatory pain, peripheral neuropathy, stroke, epilepsy, neurodegeneration, schizophrenia, drug addiction, mood disorders, post-traumatic stress disorder, seizures, convulsions, age-associated memory impairment, depression, stroke, traumatic brain injury, ischemia, Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, or Creutzfeldt-Jakob disease. In particular, the disease or condition is schizophrenia.

Still another aspect of this invention is to provide a method for treating, preventing, inhibiting or eliminating a disease or condition in a patient by modulating, activating, or inhibiting NMDA receptor activity in said patient by administering a therapeutically effective amount of at least one compound of this disclosure, wherein said disease or condition is selected from the group consisting of pain, neuropathic pain, inflammatory pain, peripheral neuropathy, stroke, epilepsy, neurodegeneration, schizophrenia, drug addiction, mood disorders, post-traumatic stress disorder, seizures, convulsions, age-associated memory impairment, depression, stroke, traumatic brain injury, ischemia,

Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, or Creutzfeldt-Jakob disease.

Still another aspect of this invention is the use of a compound as described herein as a positive allosteric modulator (PAM) of an NMDA receptor. The invention includes a method of modulating and/or amplifying the activity an NMDA receptor by contacting the receptor at an allosteric binding site with at least one compound as described herein or a pharmaceutical composition comprising such a compound. Further, compounds of the invention are useful as subtype selective for NR2A-containing NMDA receptors. The invention is also directed toward a method of modulating an NR2A-containing NMDA receptor by contacting the receptor with at least one compound of the invention or a pharmaceutical composition comprising such a compound.

The pharmaceutical compositions and compounds described herein may be formulated as solutions, emulsions, suspensions, dispersions, or inclusion complexes such as cyclodextrins in suitable pharmaceutical solvents or carriers, or as pills, tablets, lozenges, suppositories, sachets, dragees, granules, powders, powders for reconstitution, or capsules along with solid carriers according to conventional methods known in the art for preparation of various dosage forms. Pharmaceutical compositions of the embodiments may be administered by a suitable route of delivery, such as oral, parenteral, rectal, nasal, topical, or ocular routes, or by inhalation. Preferably, the compositions are formulated for intravenous or oral administration.

For oral administration, the compounds the embodiments may be provided in a solid form, such as a tablet or capsule, or as a solution, emulsion, or suspension. To prepare the oral compositions, the compounds of the embodiments may be formulated to yield a dosage of, e.g., from about 0.01 to about 50 mg/kg daily, or from about 0.05 to about 20 mg/kg daily, or from about 0.1 to about 10 mg/kg daily. Oral tablets may include the active ingredient(s) mixed with compatible pharmaceutically acceptable excipients such as diluents, disintegrating agents, binding agents, lubricating agents, sweetening agents, flavoring agents, coloring agents and preservative agents. Suitable inert fillers include sodium and calcium carbonate, sodium and calcium phosphate, lactose, starch, sugar, glucose, methyl cellulose, magnesium stearate, mannitol, sorbitol, and the like. Exemplary liquid oral excipients include ethanol, glycerol, water, and the like. Starch, polyvinyl-pyrrolidone (PVP), sodium starch glycolate, microcrystalline cellulose, and alginic acid are exemplary disintegrating agents. Binding agents may include starch and gelatin. The lubricating agent, if present, may be magnesium stearate, stearic acid, or talc. If desired, the tablets may be coated with a material such as glyceryl monostearate or glyceryl distearate to delay absorption in the gastrointestinal tract, or may be coated with an enteric coating.

Capsules for oral administration include hard and soft gelatin capsules. To prepare hard gelatin capsules, active ingredient(s) may be mixed with a solid, semi-solid, or liquid diluent. Soft gelatin capsules may be prepared by mixing the active ingredient with water, an oil such as peanut oil or olive oil, liquid paraffin, a mixture of mono and di-glycerides of short chain fatty acids, polyethylene glycol 400, or propylene glycol.

Liquids for oral administration may be in the form of suspensions, solutions, emulsions, or syrups, or may be lyophilized or presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid compositions may optionally contain: pharmaceutically-ac-

ceptable excipients such as suspending agents (for example, sorbitol, methyl cellulose, sodium alginate, gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminum stearate gel and the like); non-aqueous vehicles, e.g., oil (for example, almond oil or fractionated coconut oil), propylene glycol, ethyl alcohol, or water; preservatives (for example, methyl or propyl p-hydroxybenzoate or sorbic acid); wetting agents such as lecithin; and, if desired, flavoring or coloring agents.

The inventive compositions may be formulated for rectal administration as a suppository. For parenteral use, including intravenous, intramuscular, intraperitoneal, intranasal, or subcutaneous routes, the agents of the embodiments may be provided in sterile aqueous solutions or suspensions, buffered to an appropriate pH and isotonicity or in parenterally acceptable oil. Suitable aqueous vehicles include Ringer's solution and isotonic sodium chloride. Such forms may be presented in unit-dose form such as ampoules or disposable injection devices, in multi-dose forms such as vials from which the appropriate dose may be withdrawn, or in a solid form or pre-concentrate that can be used to prepare an injectable formulation. Illustrative infusion doses range from about 1 to 1000 $\mu\text{g}/\text{kg}/\text{minute}$ of agent admixed with a pharmaceutical carrier over a period ranging from several minutes to several days.

For nasal, inhaled, or oral administration, the inventive pharmaceutical compositions may be administered using, for example, a spray formulation also containing a suitable carrier.

For topical applications, the compounds of the present embodiments are preferably formulated as creams or ointments or a similar vehicle suitable for topical administration. For topical administration, the inventive compounds may be mixed with a pharmaceutical carrier at a concentration of about 0.1% to about 10% of drug to vehicle. Another mode of administering the agents of the embodiments may utilize a patch formulation to effect transdermal delivery.

The actual dosage employed may be varied depending upon the requirements of the patient and the severity of the condition being treated. Determination of the proper dosage regimen for a particular situation is within the skill of the art. For convenience, the total daily dosage may be divided and administered in portions during the day as required.

The amount and frequency of administration of the compounds of the invention and/or the pharmaceutically acceptable salts thereof will be regulated according to the judgment of the attending clinician considering such factors as age, condition and size of the patient as well as severity of the symptoms being treated. A typical recommended daily dosage regimen for oral administration can range from about 1 mg/day to about 500 mg/day, preferably 1 mg/day to 200 mg/day, in two to four divided doses.

Still another embodiment of the invention is a pharmaceutical formulation comprising at least one compound of Formula I or II, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient, and further comprising one or more adjunctive active agent. Methods of treatment as described herein include regimes in which the compound of the invention and at least one adjunctive active agent are administered simultaneously or sequentially.

The expression "adjunctive active agent" generally refers to agents which targets the same or a different disease, symptom, or medical condition as the primary therapeutic agent. Adjunctive active agents may treat, alleviate, relieve, or ameliorate side effects caused by administration of the primary therapeutic agents.

Exemplary, non-limiting, chemical entities and methods useful in preparing compounds of the invention will now be described by reference to the specific examples that follow. Those skilled in the art will appreciate that other synthetic routes may be used to synthesize the compounds according to the invention. Although specific starting materials and reagents are depicted and discussed herein, other starting materials and reagents can be easily substituted to provide a variety of derivatives and/or reaction conditions. In addition, many of the exemplary compounds prepared by the described methods can be further modified in light of this disclosure using conventional chemistry well known to those skilled in the art.

Artisans will recognize that, to obtain the various compounds herein, starting materials may be suitably selected so that the ultimately desired substituents will be carried through the reaction scheme with or without protection as appropriate to yield the desired product. Alternatively, it may be necessary or desirable to employ, in the place of the ultimately desired substituent, a suitable group that may be carried through the reaction scheme and replaced as appropriate with the desired substituent. Each of the reactions depicted in the reaction schemes is preferably run at a temperature from about 0° C. to the reflux temperature of the solvent used.

In the methods of preparing compounds according to the invention, it may be advantageous to separate reaction products from one another and/or from starting materials. The desired products of each step or series of steps may be separated and/or purified to the desired degree of homogeneity by the techniques common in the art. Typically such separations involve multiphase extraction, crystallization from a solvent or solvent mixture, distillation, sublimation, or chromatography. Chromatography can involve any number of methods including, for example: reverse-phase and normal phase; size exclusion; ion exchange; high, medium and low pressure liquid chromatography methods and apparatus; small scale analytical; simulated moving bed (SMB) and preparative thin or thick layer chromatography, as well as techniques of small scale thin layer and flash chromatography.

Diastereomeric mixtures may be separated into their individual diastereomers on the basis of their physical chemical differences by methods well known to those skilled in the art, such as, for example, by chromatography and/or fractional crystallization. Enantiomers may be separated by converting the enantiomeric mixture into a diastereomeric mixture by reaction with an appropriate optically active compound (e.g., chiral auxiliary such as a chiral alcohol or Mosher's acid chloride, or formation of a mixture of diastereomeric salts, for example, with tartaric acid or a chiral amine), separating the diastereomers by, for example, fractional crystallization or chromatography, and converting (e.g., hydrolyzing or de-salting) the individual diastereomers to the corresponding pure enantiomers. Enantiomers may also be separated by use of chiral HPLC column or prepared directly by chiral synthesis. The chiral centers of compounds of the present invention may be designated as "R" or "S" as defined by the IUPAC 1974 Recommendations. Enriched or purified enantiomers can be distinguished by methods used to distinguish other chiral molecules with asymmetric carbon atoms, such as optical rotation and circular dichroism.

General Experimental Conditions

Unless otherwise indicated, ^1H NMR spectra were recorded at ambient temperature using a Varian Unity Inova

65

(400 MHz) spectrometer with a triple resonance 5 mm probe. Chemical shifts are expressed in ppm relative to tetramethylsilane. The following abbreviations have been used: br=broad signal, s=singlet, d=doublet, dd=double doublet, t=triplet, q=quartet, m=multiplet.


Microwave experiments were carried out using a CEM Discover, Smith Synthesiser or a Biotage Initiator 60™, which uses a single-mode resonator and dynamic field tuning, both of which give reproducibility and control. Temperatures from 40-250° C. can be achieved and pressures of up to 30 bars can be reached.

High Pressure Liquid Chromatography-Mass Spectrometry (LCMS) experiments were used to detect associated mass ions. The spectrometers have an electrospray source operating in positive and negative ion mode. Additional detection was achieved using a Sedex 85 evaporative light scattering detector.


The following examples illustrate the preparation of representative compounds of the invention. Unless otherwise specified, all reagents and solvents were of standard commercial grade and were used without further purification. Those having skill in the art will recognize that the starting materials, reagents, and conditions described in the examples may be varied and additional steps employed to produce compounds encompassed by the present inventions.

Method 1

Example 1.1: N-(cyanomethyl)-7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




Step 1: Methyl 3-bromo-2-oxobutanoate



To a solution of methyl 2-oxobutanoate (1.00 g, 8.61 mmol) in chloroform (20 mL) were added hydrogen bromide in acetic acid (40%, 1 mL) and bromine (1.40 g, 8.76 mmol) dropwise with stirring at room temperature. The reaction mixture was stirred for 1 h at 70° C. After cooling down to room temperature, the resulting solution was concentrated in vacuo to afford methyl 3-bromo-2-oxobutanoate as yellow oil (1.60 g, 95%). No LCMS signal.


66

Step 2: Methyl 2-amino-5-methyl-1,3-thiazole-4-carboxylate




To a solution of methyl 3-bromo-2-oxobutanoate (1.60 g, 8.20 mmol) in 1,4-dioxane (30 mL) was added thiourea (625 mg, 8.21 mmol) with stirring. The resulting solution was refluxed for 3 h in an oil bath. After cooling down to room temperature, the solids were collected by filtration and dried in vacuo to afford methyl 2-amino-5-methyl-1,3-thiazole-4-carboxylate as a gray solid (900 mg, 64%). LCMS (ESI): M+H+=173.

Step 3: Methyl 7-(chloromethyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxylate



To a mixture of methyl 2-amino-5-methyl-1,3-thiazole-4-carboxylate (200 mg, 1.16 mmol) and ethyl 4-chloro-3-oxobutanoate (390 mg, 2.32 mmol) was added polyphosphoric acid (5 mL). The reaction mixture was stirred 1 h at 110° C. The reaction was then quenched with water (20 mL). The pH value of the solution was adjusted to pH 8 with sodium hydroxide (aq., 10 mol/L) and extracted with dichloromethane (2x100 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1:1) to afford methyl 7-(chloromethyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxylate as a light yellow solid (120 mg, 38%). LCMS (ESI): M+H+=273.


Step 4: Methyl 7-(4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxylate



67

To a solution of methyl 7-(chloromethyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxylate (200 mg, 0.73 mmol), potassium iodide (60 mg, 0.37 mmol) and potassium carbonate (200 mg, 1.45 mmol) in acetonitrile (15 mL) was added 4-fluorophenol (125 mg, 1.12 mmol). After stirring 2 h at 85° C., the reaction mixture was cooled down to room temperature and concentrated under vacuum. The residue was purified by chromatography with ethyl acetate/petroleum ether (1:1) to afford methyl 7-(4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxylate as a white solid (220 mg, 86%). LCMS (ESI): $M+H^+=349$; 1H NMR (300 MHz, $CDCl_3$) δ 7.02-6.92 (m, 2H), 6.91-6.86 (m, 2H), 6.48 (s, 1H), 4.92 (s, 2H), 3.98 (s, 3H), 2.45 (s, 3H).

Step 5: 7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxylic acid




To a solution of methyl 7-(4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxylate (5.00 g, 14.3 mmol) in tetrahydrofuran (400 mL) and water (200 mL) was added lithium hydroxide (7.00 g, 167 mmol). The resulting solution was stirred at 25° C. for 30 h. After the starting material was consumed (by TLC), the pH of the solution was adjusted to 7 with 2 N hydrogen chloride. Then the solution was concentrated in vacuo until a solid precipitated. The solids were filtered and washed

68

with tetrahydrofuran to afford 7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxylic acid as white solid (1.70 g, 36%). LCMS (ESI): $M+H^+=335$.

Step 6: N-(Cyanomethyl)-7-(4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide







To a solution of 7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxylic acid (100 mg, 0.30 mmol), 2-aminoacetonitrile hydrochloride (56 mg, 0.61 mmol), triethylamine (90 mg, 0.90 mmol), 4-dimethylaminopyridine (4 mg, 0.03 mmol) and 1-hydroxybenzotriazole (80 mg, 0.60 mmol) in N,N-dimethylformamide (8 mL) was added N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (115 mg, 0.6 mmol) with stirring. The resulting solution was stirred overnight and was concentrated under vacuum. The residue was purified on a silica gel column eluting with dichloromethane/methanol (30:1) to afford N-(cyanomethyl)-7-(4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide as white solid (22.7 mg, 20%). LCMS (ESI): $M+H^+=373$; 1H NMR (300 MHz, $DMSO-d_6$) δ 9.22 (m, 1H), 7.18-7.03 (m, 4H), 6.30 (s, 1H), 5.02 (s, 2H), 4.36-4.31 (m, 2H), 2.34 (s, 3H).




The following examples were prepared in a manner similar to Example 1.1:

No.	Structure/Name	LCMS (M + H)	1H NMR
1.2	<p>7-(4-Fluorophenoxy)methyl)-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	348.0	1H NMR (300 MHz, $CDCl_3$) δ 7.01-6.95 (m, 2H), 6.91-6.86 (m, 2H), 6.44 (s, 1H), 6.01 (br, 1H), 4.91 (s, 2H), 3.09-3.04 (m, 3H), 2.42 (s, 3H)

-continued


No.	Structure/Name	LCMS (M + H)	¹ H NMR
1.3	 <p>3-[(Azetidin-1-yl)carbonyl]-7-(4-fluorophenoxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidine-5-one</p>	374.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.03-6.87 (m, 4H), 6.48 (s, 1H), 4.93 (s, 2H), 4.43-4.34 (m, 1H), 4.20-4.02 (m, 2H), 3.93-3.86 (m, 1H), 2.41-2.33 (m, 5H)
1.4	 <p>N-ethyl-7-(4-fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	361.9	¹ H NMR (300 MHz, CDCl ₃) δ 7.01-6.95 (m, 2H), 6.90-6.86 (m, 2H), 6.45 (s, 1H), 5.89 (s, 1H), 4.91 (s, 2H), 3.57-3.50 (m, 2H), 2.42 (s, 3H), 1.31-1.27 (m, 3H)
1.5	 <p>7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	415.9	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.21-9.19 (m, 1H), 7.17-7.04 (m, 4H), 6.29 (s, 1H), 5.01 (s, 2H), 4.05-4.14 (m, 2H), 2.33 (s, 3H)
1.6	 <p>7-(3,4-Difluorophenoxymethyl)-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	365.95	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.38-8.36 (m, 1H), 7.39-7.36 (m, 1H), 7.25-7.24 (m, 1H), 6.91-6.89 (m, 1H), 6.28 (s, 1H), 5.02 (s, 2H), 2.77-2.74 (m, 3H), 2.29 (s, 3H)

-continued




No.	Structure/Name	LCMS (M + H)	¹ H NMR
1.7	 <p>N-ethyl-7-(4-fluorophenoxymethyl)-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	348.0	¹ H NMR (300 MHz, CDCl ₃) δ 9.64 (br, 1H), 8.03 (s, 1H), 6.87-7.03 (m, 4H), 6.58 (s, 1H), 4.96 (s, 2H), 3.42-3.51 (m, 2H), 1.24-1.28 (m, 3H)
1.8	 <p>7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	334.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 7.90 (s, 1H), 7.77 (s, 1H), 7.20-7.10 (m, 2H), 7.10-6.98 (m, 2H), 6.28 (s, 1H), 5.00 (s, 2H), 2.36 (s, 3H).
1.9	 <p>7-((4-fluorophenoxy)methyl)-N-hydroxy-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	350.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.91 (s, 1H), 9.36 (s, 1H), 7.19-7.09 (m, 2H), 7.09-6.98 (m, 2H), 6.29 (s, 1H), 5.00 (s, 2H), 2.35 (d, J = 1.9 Hz, 3H)

45





The following additional compounds were prepared using the methods described above.

No.	Structure/Name	LCMS (M + H)	¹ H NMR
1.10	 <p>7-(4-Fluorophenoxy)methyl)-2-methyl-5-oxo-N-(propan-2-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	376.10	¹ H NMR (300 MHz, CDCl ₃) δ 7.02-6.99 (m, 2H), 6.95-6.91 (m, 2H), 6.48 (s, 1H), 5.74 (s, 1H), 4.94 (s, 2H), 4.36-4.34 (m, 1H), 2.46 (s, 3H), 1.32 (s, 6H)

-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
1.11	 <p data-bbox="397 695 659 779">7-(4-Fluorophenoxymethyl)-N-(2-hydroxyethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	378.0	¹ H NMR (300 MHz, CD ₃ OD) δ 7.04 (m, 4H), 6.45 (s, 1H), 5.01 (s, 2H), 3.80-3.73 (m, 2H), 3.56-3.51 (m, 2H), 2.46 (s, 3H)
1.12	 <p data-bbox="370 1241 683 1325">7-(4-Fluorophenoxymethyl)-N-(1-hydroxypropan-2-yl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	392.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.01-6.96 (m, 2H), 6.91-6.85 (m, 2H), 6.50 (s, 1H), 5.87-5.03 (m, 1H), 4.93 (s, 2H), 4.23-4.25 (m, 1H), 4.08-03 (m, 1H), 3.55-3.50 (m, 1H), 2.45 (s, 3H), 1.30-1.27 (m, 3H)
1.13	 <p data-bbox="370 1808 683 1873">7-((4-fluorophenoxy)methyl)-2-methyl-N-(oxetan-3-ylmethyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	404.0	¹ H NMR (300 MHz, CD ₃ OD) δ 7.09-7.00 (m, 4H), 6.44 (s, 1H), 5.01 (s, 2H), 4.87-4.85 (m, 2H), 4.57-4.53 (m, 2H), 3.72-3.70 (m, 2H), 3.41-3.38 (m, 1H), 2.45 (s, 3H)


-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
1.14	 <p data-bbox="375 701 683 766">7-((4-fluorophenoxy)methyl)-N-(3-hydroxypropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	392.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.02-6.97 (m, 2H), 6.91-6.88 (m, 2H), 6.53 (br, 1H), 6.49 (s, 1H), 4.94 (s, 2H), 3.90-3.85 (m, 2H), 3.70-3.66 (m, 2H), 2.46 (s, 3H), 1.95-1.89 (m, 2H)
1.15	 <p data-bbox="375 1113 678 1199">N-cyclopropyl-7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	374.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.01-6.95 (m, 2H), 6.89-6.85 (m, 2H), 6.45 (s, 1H), 6.03 (s, 1H), 4.91 (s, 2H), 2.93-2.90 (m, 1H), 2.42 (s, 3H), 0.93-0.87 (m, 2H), 0.81-0.74 (m, 2H).
1.16	 <p data-bbox="370 1491 683 1556">7-((4-fluorophenoxy)methyl)-N-methoxy-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	364.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 11.50 (s, 1H), 7.19-7.09 (m, 2H), 7.09-6.99 (m, 2H), 6.30 (s, 1H), 5.01 (s, 2H), 3.73 (s, 3H), 2.36 (s, 3H).
1.17	 <p data-bbox="358 1862 699 1927">7-(4-Fluorophoxymethyl)-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carbothioamide</p>	363.9	¹ H NMR (300 MHz, CD ₃ OD) δ 7.04-6.97 (m, 4H), 6.38 (s, 1H), 5.06 (s, 2H), 3.22 (s, 3H), 2.36 (s, 3H).

77

Method 2


Example 2.1: 7-((4-fluorophenoxy)methyl)-2-methyl-3-propionyl-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of methyl 7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxylate (from Example 1.1, Step 4; 100 mg, 0.29 mmol) in tetrahydrofuran (2 mL) was added ethylmagnesium bromide (0.14 mL, 0.32 mmol). The reaction mixture was stirred for 1 h at room temperature. The reaction was then quenched by addition of water, extracted with dichloromethane (3×30 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified on a silica gel column with ethyl acetate/petroleum ether (1:3) to provide 7-((4-fluorophenoxy)methyl)-2-methyl-3-propionyl-5H-thiazolo[3,2-a]pyrimidin-5-one as a white solid (91.9 mg, 92%). LCMS (ESI): $M+H^+=347.0$.

78


Example 2.2: 7-((4-fluorophenoxy)methyl)-3-(1-hydroxypropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one




To a solution of 7-((4-fluorophenoxy)methyl)-2-methyl-3-propionyl-5H-thiazolo[3,2-a]pyrimidin-5-one (from Example 2.1; 20.0 mg, 0.060 mmol) in methanol (10 mL) was added sodium borohydride (4.70 mg, 0.13 mmol). The reaction mixture was stirred overnight at room temperature. The reaction was then quenched by saturated aqueous ammonium chloride (20 mL), extracted with dichloromethane (3×30 mL), washed with brine, dried over sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with dichloromethane/methanol (50:1) to afford 7-((4-fluorophenoxy)methyl)-3-(1-hydroxypropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one as a white solid (4.50 mg, 21%). LCMS (ESI): $M+H^+=349.0$; 1H NMR (300 MHz, CD_3OD) δ 7.05-7.03 (m, 4H), 6.50 (s, 1H), 5.11-5.01 (m, 1H), 4.89 (s, 2H), 2.51 (s, 3H), 1.96-1.83 (m, 2H), 0.97-0.92 (m, 3H).

The following examples were prepared in a manner similar to Example 2.1 and 2.2:


No.	Structure/Name	LCMS (M + H)	1H NMR
2.3	<p>7-(4-Fluorophenoxy)methyl)-3-(1-hydroxyethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one</p>	335.20	1H NMR (300 MHz, $DMSO-d_6$) δ 7.18-7.12 (m, 2H), 7.08-7.03 (m, 2H), 6.33 (s, 1H), 5.68-5.62 (m, 2H), 5.00 (s, 2H), 2.49 (s, 3H), 1.43-1.41 (m, 3H)
2.4	<p>7-(4-Fluorophenoxy)methyl)-3-(2-hydroxypropan-2-yl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one</p>	349.10	1H NMR (300 MHz, CD_3OD) δ 7.18-7.13 (s, 2H), 7.08-7.05 (s, 2H), 6.66 (s, 1H), 6.41 (s, 1H), 5.02 (s, 2H), 2.54 (s, 3H), 1.65 (s, 6H)

No.	Structure/Name	LCMS (M + H)	¹ H NMR
2.5	 <p>3-acetyl-7-((4-fluorophenoxy)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	333.0	¹ H NMR (300 MHz, CD ₃ OD) δ 7.08-7.01 (m, 4H), 6.46 (s, 1H), 5.06 (s, 2H), 2.50 (s, 3H), 2.40 (s, 3H).

Example 2.6: 2-(7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)-N-methylacetamide




Step 1: 7-((4-fluorophenoxy)methyl)-3-(hydroxymethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one




Into a 25-mL round bottom flask under nitrogen was added a solution of methyl 7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxylate (from Example 1.1, Step 5) (200 mg, 0.57 mmol) in tetrahydrofuran (20 mL) and a solution of diisobutylaluminum hydride in toluene (1.1 mol/L, 1 mL). The resulting solution was stirred overnight at room temperature. The reaction was quenched with water (30 mL), extracted with dichloromethane (3x50 mL), washed with brine, dried over sodium sulfate and concentrated in vacuo. The residue was purified by silica gel chromatography with dichloromethane/methanol (50:1) to afford 7-((4-fluorophenoxy)methyl)-3-(hydroxymethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one as an off-white solid (150 mg, 77%). LCMS (ESI): M+H⁺=321.0; ¹H NMR (300 MHz, CDCl₃) δ 7.02-6.87 (m, 4H), 6.48 (s, 1H), 4.93 (s, 2H), 4.76 (s, 2H), 2.44 (s, 3H).

Step 2: 3-(Chloromethyl)-7-(4-fluorophenoxy)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



To a solution of 7-(4-fluorophenoxy)-3-(hydroxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (200 mg, 0.62 mmol) in dichloromethane (5 mL) was added thionyl chloride (0.5 mL) and N,N-dimethylformamide (10 mg, 0.14 mmol). The resulting solution was stirred overnight at room temperature and then concentrated in vacuo. The residue was purified by chromatography with dichloromethane/ethyl acetate (50/1) to afford 3-(chloromethyl)-7-(4-fluorophenoxy)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a white solid (150 mg, 71%). LCMS (ESI): M+H⁺=339.0; ¹H NMR (400 MHz, CDCl₃) δ 7.01-6.96 (m, 2H), 6.92-6.89 (m, 2H), 6.47 (s, 1H), 5.26 (s, 2H), 4.91 (s, 2H), 2.45 (s, 3H).

Step 3: Methyl 2-[7-(4-fluorophenoxy)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]acetate




To a solution of 3-(chloromethyl)-7-(4-fluorophenoxy)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (200 mg, 0.59 mmol) in methanol (5 mL) was added 1,1'-bis(diphenylphosphino)ferrocenepalladium dichloride (60 mg, 0.08 mmol), potassium carbonate (163 mg, 1.18 mmol). The reaction mixture was stirred for 3 h at 25 °C under carbon monoxide (5 atm) atmosphere. After filtration,

81

the filtrate was concentrated in vacuo. The residue was purified by chromatography with dichloromethane/ethyl acetate (50/1) to afford methyl 2-[7-(4-fluorophenoxy)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]acetate as a white solid (120 mg, 56%). LCMS (ESI): $M+H^+=363.2$; 1H NMR (400 MHz, $CDCl_3$) δ 7.01-6.95 (m, 2H), 6.91-6.86 (m, 2H), 6.37 (s, 1H), 4.89 (s, 2H), 4.19 (s, 2H), 3.75 (s, 3H), 2.33 (s, 3H).

Step 4: 2-[7-(4-Fluorophenoxy)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]-N-methylacetamide






82

Methyl 2-[7-(4-fluorophenoxy)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]acetate (70 mg, 0.19 mmol) and a methylamine in ethanol solution (30%, 5 mL) were added to a 25-mL round-bottom flask. The resulting solution was stirred for 30 min at 40° C. and then concentrated in vacuo. The residue was purified by chromatography with dichloromethane/ethyl acetate (10/1) to afford 2-[7-(4-fluorophenoxy)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]-N-methylacetamide as a white solid (38 mg, 54%). LCMS (ESI): $M+H^+=362.0$; 1H NMR (300 MHz, $CDCl_3$) δ 7.02-6.87 (m, 5H), 6.44 (s, 1H), 4.92 (s, 2H), 4.12 (s, 2H), 2.78-2.76 (m, 3H), 2.51 (s, 3H).

The following examples were prepared using methods analogous to those described above:


No.	Structure/Name	LCMS (M + H)	1H NMR
2.7	<p>3-Cyclopropanecarbonyl-7-(4-fluorophenoxy)methyl-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one</p>	359.0	1H NMR (300 MHz, $CDCl_3$) δ 7.02-6.87 (m, 4H), 6.50 (s, 1H), 4.94 (s, 2H), 2.38 (s, 3H), 2.19-2.10 (m, 1H), 1.38-1.31 (m, 2H), 1.17-1.11 (m, 2H)
2.8	<p>7-(4-Fluorophenoxy)methyl-3-[1-(hydroxyimino)ethyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one</p>	348.0	1H NMR (300 MHz, $CDCl_3$) δ 7.07-6.98 (m, 4H), 6.38 (s, 1H), 4.98 (s, 2H), 2.36 (s, 2H), 2.29 (s, 1H), 2.18 (s, 1H), 2.10 (s, 2H)

-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
2.9	 <p>7-((4-fluorophenoxy)methyl)-2-methyl-3-(oxetane-3-carbonyl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	375.0	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.17-7.13 (m, 4H), 6.28 (s, 1H), 6.04 (s, 1H), 5.69 (s, 1H), 5.16-5.12 (m, 1H), 5.00 (s, 2H), 4.35-4.25 (m, 2H), 2.30 (s, 3H)
2.10	 <p>7-((4-fluorophenoxy)methyl)-2-methyl-3-(2,2,2-trifluoro-1-hydroxyethyl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	389.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.95 (s, 1H), 7.03-6.96 (m, 2H), 6.93-6.87 (m, 2H), 6.66 (s, 1H), 5.35-5.33 (m, 1H), 4.96 (s, 2H), 2.50 (s, 3H)
2.11	 <p>7-(4-Fluorophenoxy)methyl)-2-methyl-3-(trifluoroacetyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one</p>	386.8	¹ H NMR (300 MHz, CDCl ₃) δ 7.03-6.96 (m, 2H), 6.93-6.87 (m, 2H), 6.59 (s, 1H), 4.95 (s, 2H), 2.44 (s, 3H)

Method 3


Example 3.1: 2-cyclopropyl-N-ethyl-7-((4-fluorophenoxy)methyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide



45

Step 1: 7-(chloromethyl)-2-cyclopropyl-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide

50



55


To a solution of 2-amino-5-cyclopropyl-N-ethylthiazole-4-carboxamide (2.53 g, 12.8 mmol) in polyphosphoric acid (16.0 g) was added ethyl 4-chloro-3-oxobutanoate (4.20 g, 25.5 mmol). The resulting solution was stirred for 1 h at 110° C. The reaction was then quenched by water (80 mL) and the pH value of the solution was adjusted to pH 7 with a sodium hydroxide solution (1 mol/L). The reaction mixture was extracted with dichloromethane (3×50 mL), washed with

65

85

brine (30 mL), dried over anhydrous magnesium sulfate and concentrated in vacuo. The residue was purified by chromatography with dichloromethane/ethyl acetate (5/1) to afford 7-(chloromethyl)-2-cyclopropyl-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide as a brown solid (905 mg, 24%). LCMS (ESI): $M+H^+=312.0$.

Step 2: 2-cyclopropyl-N-ethyl-7-((4-fluorophenoxy)methyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




86


A solution of 7-(chloromethyl)-2-cyclopropyl-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide (100 mg, 0.32 mmol) in acetonitrile (10 mL) was treated with potassium iodide (27.0 mg, 0.16 mmol), potassium carbonate (88.0 mg, 0.64 mmol) and 4-fluorophenol (72.0 mg, 0.64 mmol). The reaction mixture was then stirred overnight at 80° C. After cooling down to room temperature, the reaction mixture was concentrated in vacuo. The residue was purified by chromatography with dichloromethane/methanol (50/1) to afford 2-cyclopropyl-N-ethyl-7-(4-fluorophenoxy)methyl-5-oxo-5H-[1,3]Thiazolo[3,2-a]pyrimidine-3-carboxamide as a white solid (36.4 mg, 29%). LCMS (ESI): $M+H^+=388.0$; 1H NMR (300 MHz, $CDCl_3$) δ 7.00-6.95 (m, 2H), 6.90-6.84 (m, 2H), 6.44 (s, 1H), 5.91 (s, 1H), 4.90 (s, 2H), 3.57-3.53 (m, 2H), 2.19-2.09 (m, 1H), 1.32-1.27 (m, 3H), 1.22-1.08 (m, 2H), 0.90-0.83 (m, 2H).

The following examples were prepared in a manner similar to Example 3.1:

No.	Structure/Name	LCMS (M + H)	1H NMR
3.2	<p>7-(4-Fluorophenoxy)methyl-N-methyl-5-oxo-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	401.9	1H NMR (300 MHz, $CDCl_3$) δ 7.02-6.95 (m, 2H), 6.92-6.85 (m, 2H), 6.53 (s, 1H), 5.89 (bs, 1H), 4.92 (s, 2H), 3.10-3.06 (m, 3H)
3.3	<p>2-Cyclopropyl-7-(4-fluorophenoxy)methyl-N-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	374.0	1H NMR (300 MHz, $CDCl_3$) δ 7.02-6.91 (m, 2H), 6.90-6.85 (m, 2H), 6.42 (s, 1H), 6.11 (s, 1H), 4.89 (s, 2H), 3.07 (s, 3H), 2.21-2.12 (m, 1H), 1.29-1.08 (m, 2H), 0.92-0.81 (m, 2H).


-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
3.4	 <p>N-Ethyl-7-(4-fluorophenoxymethyl)-5-oxo-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	415.95	¹ H NMR (300 MHz, CDCl ₃) δ 7.02-6.96 (m, 2H), 6.91-6.85 (m, 2H), 6.53 (s, 1H), 5.87 (br, 1H), 4.92 (s, 2H), 3.60-3.51 (m, 2H), 1.31-1.27 (m, 3H)

3.5	 <p>7-((4-fluorophenoxy)methyl)-N-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	334.0	¹ H NMR (300 MHz, CDCl ₃) δ 8.07 (s, 1H), 7.03-6.96 (m, 2H), 6.94-6.88 (m, 2H), 6.59 (s, 1H), 4.96 (s, 2H), 2.99 (s, 3H)
-----	--	-------	---

45

The following example was prepared using methods analogous to those described above.

No.	Structure/Name	LCMS (M + H)	¹ H NMR
3.6	 <p>N-ethyl-7-[[[(5-fluoropyridin-2-yl)oxy]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	363.15	¹ H NMR (300 MHz, CD ₃ OD) δ 7.99-7.98 (m, 1H), 7.61-7.54 (m, 1H), 7.00-6.96 (m, 1H), 6.32 (s, 1H), 5.27 (s, 2H), 3.45-3.38 (m, 2H), 2.42 (s, 3H), 1.26-1.22 (m, 3H)


89

Method 4


Example 4.1: 7-(4-Fluorophenoxymethyl)-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

Example 4.2: 7-(4-Fluorophenoxymethyl)-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (Enantiomer 1)

Example 4.3: 7-(4-Fluorophenoxymethyl)-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (Enantiomer 2)



Step 1: tert-butyldimethyl((2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)cyclopropyl)methoxy)silane



Diethylzinc (1.0 M in hexane) (200 mL, 200 mmol) was added to freshly distilled dichloromethane (200 mL) under nitrogen. Then a solution of trifluoroacetic acid (15.4 mL, 200 mmol) in dichloromethane (100 mL) was added dropwise at 0° C. Upon stirring for 30 min, a solution of diiodomethane (16.1 mL, 200 mmol) in dichloromethane (100 mL) was added at 0° C. After an additional 30 min of stirring, a solution of (E)-tert-butyldimethyl(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)allyloxy)silane (30.0 g, 100 mmol) in dichloromethane (100 mL) was added at 0° C. The resulting solution was stirred 2 h at room temperature and was then quenched with water. The reaction was extracted with dichloromethane (1000 mL.×2), washed with brine, and the organic layer was then dried over anhydrous sodium sulfate and concentrated to afford tert-butyldimethyl((2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)cyclopropyl)methoxy)silane as a colorless oil (30 g, 96%). ¹H NMR (300 MHz, CDCl₃) δ 3.58-3.53 (m, 1H), 3.44-3.38 (m, 1H), 1.17 (s, 12H), 0.87 (s, 9H), 0.66-0.63 (m, 1H), 0.54-0.49 (m, 1H), 0.06 (s, 6H), -0.35 to -0.25 (m, 2H).

90

Step 2: Potassium


trans-2-(hydroxymethyl)cyclopropyltrifluoroborate



To a solution of tert-butyldimethyl((2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)cyclopropyl)methoxy)silane (30.0 g, 100 mmol) in methanol (300 mL) was added a solution of potassium difluoride (32.0 g, 400 mmol) in water (100 mL) dropwise at 0° C. After stirring 1.5 h at room temperature, the reaction mixture was concentrated under reduce pressure. The resulting solid was suspended in acetone (1 L) and was refluxed 20 min. The heterogeneous mixture was then filtered to remove potassium difluoride and the filtrate was concentrated. The extraction was repeated for the filtered solid. The combined filtrates were concentrated and dissolved in minimal acetone followed by the slow addition of ethyl ether until the solution become cloudy. The mixture was filtered and the solid was collected to provide potassium trans-2-(hydroxymethyl)cyclopropyltrifluoroborate (7.00 g, 40%). ¹H NMR (300 MHz, CDCl₃) δ 4.01-3.97 (m, 1H), 3.44-3.37 (m, 1H), 2.83-2.75 (m, 1H), 0.58-0.48 (m, 1H), 0.01 to -0.03 (m, 1H), -0.21 to -0.25 (m, 1H), -0.94 to -0.97 (m, 1H).


Step 3:

2-(5-methylthiazol-2-yl)isoindoline-1,3-dione



5-Methylthiazol-2-amine (200 g, 1.75 mol) and phthalic acid anhydride (272.4 g, 1.84 mol) were suspended in dioxane (2.5 L) and heated at 110° C. overnight. TLC (DCM/MeOH=20:1) showed the reaction was complete. The mixture was concentrated, and the residue was purified via column chromatography on silica gel (DCM/MeOH=50:1~20:1) to give 2-(5-methylthiazol-2-yl)isoindoline-1,3-dione (240 g, 56%) as an off-white solid. LCMS (ESI): M+H⁺=245.1; ¹H NMR (400 MHz, CDCl₃) δ 8.01-7.99 (m, 2H), 7.84-7.82 (m, 2H), 7.46 (s, 1H), 2.51 (s, 3H).

Step 4: 2-(4-bromo-5-methylthiazol-2-yl)isoindoline-1,3-dione




To a mixture of 2-(5-methylthiazol-2-yl)isoindoline-1,3-dione (250 g, 0.819 mol) in THF (2 L) was added N-bro-

91


mosuccinimide (330 g, 1.85 mol) portionwise at room temperature. Then the mixture was stirred overnight at 30° C. LCMS showed the reaction was complete. The mixture was diluted with water and ethyl acetate. The mixture was filtered and the filter cake was dried to give 2-(4-bromo-5-methylthiazol-2-yl)isoindoline-1,3-dione (210 g, 75%) as a yellow solid. LCMS (ESI): M+H⁺=323.1, 325.1; ¹H NMR (400 MHz, DMSO-d₆) δ 8.03-8.00 (m, 2H), 7.95-7.93 (s, 2H), 2.41 (s, 3H).

Step 5: 4-bromo-5-methylthiazol-2-amine




A mixture of 2-(4-bromo-5-methylthiazol-2-yl)isoindoline-1,3-dione (152 g, 0.471 mol) and hydrazine monohydrate (29.5 g, 0.495 mol) in EtOH (1.5 L) was stirred overnight at 20° C. TLC (100% DCM) showed the reaction was complete. The mixture was then concentrated and the residue was purified via column chromatography on silica (100% DCM) to give 4-bromo-5-methylthiazol-2-amine (67 g, 73%) as a white solid. LCMS (ESI): M+H⁺=193.1; ¹H NMR (400 MHz, CDCl₃) δ 8.52 (br, 1H), 5.11 (br, 1H), 2.23 (s, 1H).

Step 6: 3-bromo-7-(chloromethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



A mixture of 4-bromo-5-methylthiazol-2-amine (60 g, 0.31 mol) and 4-chloro-3-oxo-butanoate (62 g, 0.37 mol) in PPA (500 g) was stirred for 2 h at 110° C. LCMS showed the reaction was complete. The aqueous layer was extracted with DCM (300 mL×3). The combined organic layers were washed with water (200 mL×3) and brine (200 mL), dried over Na₂SO₄ and concentrated. The residue was purified via chromatography on silica gel (DCM/MeOH=100:1~50:1) to give 3-bromo-7-(chloromethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (50 g, 55%) as a brown solid. LCMS (ESI): M+H⁺=293.1, 295.1; ¹H NMR (400 MHz, CDCl₃) δ 6.39 (s, 1H), 4.39 (s, 2H), 2.38 (s, 3H).

Step 7: 3-bromo-7-((4-fluorophenoxy)methyl)-2-methyl-5H-thiazolo[3,2-c]pyrimidin-5-one



92

To a solution of 3-bromo-7-(chloromethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (20 g, 0.068 mmol) in acetonitrile (300 ml) was added 4-fluorophenol (9.20 g, 0.082 mmol), potassium iodide (5.68 g, 0.034 mmol), and potassium carbonate (26.1 g, 0.136 mmol). The mixture was stirred for 3 h at 80° C. and then cooled down room temperature. After filtration and concentration, the residue was purified by chromatography by ethyl acetate/petroleum ether (1/1) to afford 3-bromo-7-((4-fluorophenoxy)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (20 g, 80%) as a yellow solid. LCMS (ESI): M+H⁺=369.1, 371.1; ¹H NMR (300 MHz, CDCl₃) δ 7.01-6.95 (m, 2H), 6.92-6.86 (m, 2H), 6.44 (s, 1H), 4.88 (s, 2H), 2.36 (s, 3H).

Step 8: 7-(4-Fluorophenoxy)methyl-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one and 7-(4-Fluorophenoxy)methyl-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (Enantiomer 1 and Enantiomer 2)





3-Bromo-7-((4-fluorophenoxy)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (500 mg, 1.36 mmol), sodium carbonate (430 mg, 4.07 mmol), 1,1'-bis(diphenylphosphino)-ferrocenepalladiumdichloride (200 mg, 0.27 mmol), potassium organotrifluoroborates (500 mg, 2.80 mmol), 1,4-dioxane (12 mL) and water (3 mL) were placed in a 30-mL sealed tube. The reaction was stirred at 120° C. for 1.5 h under microwave irradiation. The reaction was then extracted with dichloromethane, washed with brine, dried over sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with dichloromethane/ethyl acetate (2/1) to afford 7-(4-fluorophenoxy)methyl-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (Example 4.1) as a light yellow solid (3.00 g, 30%). LCMS (ESI): M+H⁺=361.1; ¹H NMR (300 MHz, CDCl₃) δ 7.04-6.98 (m, 2H), 6.93-6.89 (m, 2H), 6.48 (s, 1H), 4.92 (s, 2H), 4.09-4.05 (m, 1H), 3.16-3.09 (m, 1H), 2.41 (s, 3H), 2.36-2.28 (m, 1H), 1.31-1.28 (m, 1H), 1.07-1.00 (m, 2H).

Example 4.1 was purified by Chiral-Prep-HPLC with the following conditions (Prep-SFC80): Column, Chiralpak IC, 2*25 cm, 5 um; mobile phase, CO₂ and EtOH (0.2% DEA) (hold 65% CO₂ in 13 mins); Detector, UV 220 nm to afford two enantiomers.





Peak (9.93 min): Enantiomer 1 (1.09 g, 10%). LCMS (ESI): M+H⁺=361.0; ¹H NMR (300 MHz, CDCl₃) δ 7.03-6.87 (m, 4H), 6.46 (s, 1H), 4.91 (s, 2H), 4.08-4.03 (m, 1H), 3.16-3.08 (m, 1H), 2.40 (s, 3H), 2.31-2.23 (m, 1H), 1.32-1.26 (m, 1H), 1.07-0.96 (m, 2H).

Peak (11.06 min): Enantiomer 2 (0.96 g, 10%). LCMS (ESI): M+H⁺=361.0; ¹H NMR (300 MHz, CDCl₃) δ 7.02-6.89 (m, 4H), 6.47 (s, 1H), 4.91 (s, 2H), 4.08-4.03 (m, 1H), 3.16-3.09 (m, 1H), 2.40 (s, 3H), 2.30-2.28 (m, 1H), 1.30-1.25 (m, 1H), 1.07-1.01 (m, 2H).





The following examples were prepared in a manner similar to Example 4.1, 4.2, and 4.3:

No.	Structure/Name	LCMS (M + H)	¹ H NMR
4.4	 <p>7-((3-fluorophenoxy)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	361.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.26-7.21 (m, 1H), 6.76-6.64 (m, 3H), 6.45 (s, 1H), 4.93 (s, 2H), 4.08-4.03 (m, 1H), 3.16-3.09 (m, 1H), 2.39 (s, 3H), 2.32-2.26 (m, 1H), 1.31-1.23 (m, 1H), 1.07-0.97 (m, 2H)
4.5	 <p>7-((4-fluorophenoxy)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	369.1	¹ H NMR (300 MHz, CDCl ₃) δ 9.26 (s, 1H), 8.69 (s, 2H), 7.01-6.87 (m, 4H), 6.40 (s, 1H), 4.94 (s, 2H), 2.29 (s, 3H)
4.6	 <p>7-(2,4-Difluorophenoxy)methyl)-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one</p>	379.10	¹ H NMR (300 MHz, CDCl ₃) δ 6.95-6.86 (m, 2H), 6.81-6.74 (m, 1H), 6.50 (s, 1H), 4.96 (s, 2H), 4.08-4.03 (m, 2H), 3.15-3.08 (m, 1H), 2.41 (s, 3H), 2.40-2.27 (m, 1H), 1.30-1.25 (m, 1H), 1.07-0.96 (m, 2H)
4.7	 <p>7-(3,4-Difluorophenoxy)methyl)-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one</p>	379.25	¹ H NMR (300 MHz, CDCl ₃) δ 7.14-7.05 (m, 1H), 6.82-6.75 (m, 1H), 6.68-6.64 (m, 1H), 6.43 (s, 1H), 4.90 (s, 1H), 4.10-4.05 (m, 1H), 3.16-3.09 (m, 1H), 2.41 (s, 3H), 2.37-2.29 (m, 1H), 1.34-1.27 (m, 2H), 1.08-0.98 (m, 2H)



-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
4.8	 <p data-bbox="342 642 675 705">7-(4-Chlorophenoxy)methyl-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one</p>	377.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.26-7.24 (m, 2H), 6.91-6.87 (m, 2H), 6.45 (s, 1H), 4.97 (s, 2H), 4.11-4.03 (m, 1H), 3.16-3.09 (m, 1H), 2.43 (s, 3H), 2.40-2.28 (m, 1H), 1.27-1.19 (m, 1H), 1.07-1.01 (m, 2H)
4.9	 <p data-bbox="342 993 675 1077">7-[(5-Fluoropyridin-2-yl)oxy]methyl-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one</p>	362.0	¹ H NMR (300 MHz, CD ₃ OD) δ 8.02-8.01 (m, 1H), 7.60-7.59 (m, 1H), 7.01-6.97 (m, 1H), 6.27 (s, 1H), 5.25 (s, 2H), 3.64-3.60 (m, 2H), 2.45 (s, 3H), 2.23-2.17 (m, 1H), 1.41-1.32 (m, 1H), 1.07-1.01 (m, 2H)
4.10	 <p data-bbox="342 1476 675 1560">3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-7-((4-(trifluoromethyl)phenoxy)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	411.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.60-7.56 (m, 2H), 7.04-7.01 (m, 2H), 6.45 (s, 1H), 5.04 (s, 2H), 4.07-4.02 (m, 1H), 3.17-3.13 (m, 1H), 2.41 (s, 3H), 2.34-2.27 (m, 1H), 1.30-1.26 (m, 1H), 1.08-1.01 (m, 2H)
4.11	 <p data-bbox="342 1875 675 1929">7-((4-fluorophenoxy)methyl)-2-methyl-3-(oxazol-2-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	358.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.85 (s, 1H), 7.34 (s, 1H), 7.01-6.86 (m, 4H), 6.42 (s, 1H), 4.93 (s, 2H), 2.37 (s, 3H)


-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
4.12	 <p>7-((2-fluorophenoxy)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	361.0	¹ HNMR (300 MHz, CDCl ₃) δ 7.17-6.96 (m, 4H), 6.57 (s, 1H), 5.08 (s, 2H), 4.09-4.08 (m, 1H), 3.20-3.15 (m, 1H), 2.44 (s, 3H), 2.32-2.17 (m, 1H), 1.31-1.28 (m, 1H), 1.09-1.02 (m, 2H)
4.13	 <p>4-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methoxy)benzonitrile</p>	368.0	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.82-7.79 (m, 2H), 7.23-7.20 (m, 2H), 6.22 (s, 1H), 5.07 (s, 2H), 4.59-4.55 (m, 1H), 3.48-3.44 (m, 2H), 2.37 (s, 3H), 2.05-2.01 (m, 1H), 1.33-1.25 (m, 1H), 0.89-0.86 (m, 2H)
4.14	 <p>7-((4-fluorophenoxy)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	347.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 7.18-7.10 (m, 2H), 7.10-7.03 (m, 2H), 7.02 (s, 1H), 6.23 (s, 1H), 4.97 (s, 2H), 4.43 (dd, J = 6.6, 4.6 Hz, 1H), 3.55-3.36 (m, 2H), 2.63-2.54 (m, 1H), 1.41-1.26 (m, 1H), 1.01 (dt, J = 8.5, 5.2 Hz, 1H), 0.87 (dt, J = 8.4, 5.2 Hz, 1H).
4.15	 <p>7-((4-fluorophenoxy)methyl)-2-methyl-3-(1H-pyrazol-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	357.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 12.97 (s, 1H), 7.78 (s, 1H), 7.19-7.09 (m, 2H), 7.09-6.97 (m, 2H), 6.47-6.26 (m, 1H), 6.16 (s, 1H), 4.99 (s, 2H), 2.20 (s, 3H).

-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
4.16	 <p>7-((4-fluorophenoxy)methyl)-2-methyl-3-(4H-1,2,3-triazol-3-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	358.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.64 (s, 1H), 7.18-7.08 (m, 2H), 7.08-7.00 (m, 2H), 6.19 (s, 1H), 5.01 (s, 2H), 2.22 (d, J = 13.2 Hz, 3H).
4.17	 <p>3-cyclopropyl-7-((4-fluorophenoxy)methyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	331.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 7.20-7.09 (m, 2H), 7.09-6.96 (m, 2H), 6.18 (s, 1H), 4.94 (s, 2H), 2.35 (d, J = 1.5 Hz, 3H), 2.16 (tdd, J = 8.5, 5.1, 1.9 Hz, 1H), 0.99-0.87 (m, 2H), 0.75-0.58 (m, 2H).


Example 4.18: cis-2-[7-(4-Fluorophenoxy)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carbonitrile




and

Example 4.18A

trans-2-[7-(4-Fluorophenoxy)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carbonitrile



Step 1: Ethyl 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)cyclopropanecarboxylate



35

40

45

50

55


60

65

To a solution of 2-ethenyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (10.0 g, 4.38 mmol) and palladium acetate (166 mg, 0.44 mmol) in ether (50 mL) was added ethyl 2-diazoacetate (6.60 g, 5.47 mmol) in ether (20 mL) dropwise for 10 min at room temperature. Palladium acetate (166 mg, 0.44 mmol) and ethyl 2-diazoacetate (6.60 g, 5.47 mmol) in ether (20 mL) were again added dropwise for another 10 min. The resulting solution was then stirred for 1 h at room temperature. After filtration through active aluminum oxide, the filtrate was concentrated in vacuo to afford ethyl 2-(tetramethyl-1,3,2-dioxaborolan-2-yl)cyclopropane-1-carboxylate as yellow oil (24.0 g). The crude product was used in the next step without further purification.


101

Step 2: Ethyl 2-[7-(4-fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carboxylate



To a solution of 3-bromo-7-(4-fluorophenoxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 4.1, Step 7) (500 mg, 1.35 mmol), ethyl 2-(4,4,5,5-tetramethyl-1,3-dioxolan-2-yl)cyclopropane-1-carboxylate (2.25 g, 6.37 mmol) and potassium carbonate (697 mg, 5.20 mmol) in 5:1 acetonitrile/water (12 mL) was added 1,1'-bis(diphenylphosphino)ferrocenepalladium dichloride (54.7 mg, 0.06 mmol). The resulting solution was stirred for 1 h at 120° C. in a 20-mL microwave tube. The process was then scaled up to 5 g (10 batches) using the same method. After concentration in vacuo, the residue was purified by chromatography with ethyl acetate/petroleum ether (1/2) to afford ethyl 2-[7-(4-fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carboxylate as a red oil (1.50 g, 28%). LCMS (ESI): $M+H^+=403.0$.


Step 3: 2-[7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carboxylic acid



To a solution of ethyl 2-[7-(4-fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carboxylate (1.50 g, 3.75 mmol) in tetrahydrofuran/water (100/10 mL) was added lithium hydroxide (850 mg, 35.5 mmol) and the solution was stirred overnight at room temperature. The pH value of the solution was adjusted to 4-5 with hydrogen chloride (1 mol/L). The resulting solution was extracted with ethyl acetate (2x200 mL) and dried over anhydrous magnesium sulfate. After filtration, the filtrate was concentrated in vacuo to afford 2-[7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carboxylic acid as a yellow oil (800 mg, 57%). LCMS (ESI): $M+H^+=375.0$.


102

Step 4: 2-(7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)cyclopropanecarboxamide



To a solution of 2-[7-(4-fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carboxylic acid (800 mg, 2.25 mmol), triethylamine (325 mg, 3.25 mmol), in tetrahydrofuran (100 mL) was added propan-2-ylchloroformate (325 mg, 2.75 mmol). The solution was stirred for 20 min at room temperature. Then ammonium hydroxide (10 mL, 2.60 mmol) was added and the solution was stirred for an additional 20 min at room temperature. The resulting solution was extracted with ethyl acetate (2x100 mL), washed with brine, dried over anhydrous magnesium sulfate and concentrated in vacuo. The residue was purified by chromatography with 3% methanol in dichloromethane to afford 2-(7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)cyclopropanecarboxamide as a white solid (500 mg, 62%).

Step 5: 2-(7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)cyclopropanecarbonitrile



To a solution of 2-(7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)cyclopropanecarboxamide (500 mg, 1.20 mmol) in dichloromethane (100 mL) was added 1,8-diazabicyclo[5.4.0]undec-7-ene (100 mL) and ethyl dichlorophosphate (50 mL). After stirring for 30 min at room temperature, the reaction was quenched with water (50 mL) and extracted with dichloromethane (3x20 mL) and the combined organic layer was washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by preparative HPLC with the following conditions: (1#-Pre-HPLC-005(Waters)): Column, SunFire Prep C₁₈ OBD Column, 5 um, 19*150 mm; mobile phase, water with 10 mmol NH₄HCO₃ and CH₃CN (50.0% CH₃CN up to 82.0% in 10 min, down to 50.0% in 2 min); Detector, UV 254/220 nm to

103

afford the cis (Example 4.18; 200 mg, 42%) and trans (Example 4.18A; 60 mg, 12%) isomers.

Example 4.18

LCMS (ESI): $M+H^+=356.2$; 1H NMR (300 MHz, CD_3OD) δ 7.07-7.01 (m, 4H), 6.40 (s, 1H), 4.97 (s, 2H), 3.10-2.92 (m, 1H), 2.42 (s, 3H), 2.05-1.96 (m, 1H), 1.84-1.77 (m, 1H), 1.62-1.55 (m, 1H).

Examples 4.19 and 4.20

The racemic cis isomer (Ex. 4.18) was purified by Chiral-Prep-HPLC with the following conditions (Prep-HPLC-032): Column, Chiralpak IA, 2*25 cm, 5 μ m; mobile phase, Hex (1% TEA)/EtOH (hold 50.0% EtOH in 12 mins), flow, 1.0 mL/min; Detector, UV 254 nm to afford two enantiomers.

Example 4.19 (Cis Enantiomer 1)

Obtained as a white solid (58.6 mg, 16%). Chiral-Prep-HPLC retention time, 6.48 min; LCMS(ESI): $M+H^+=356.2$; 1H NMR (400 MHz, $CDCl_3$) δ 7.03-7.00 (m, 2H), 6.92-6.89 (m, 2H), 6.46 (s, 1H), 4.90 (s, 2H), 3.02-3.01 (m, 1H), 2.42 (s, 3H), 1.88-1.83 (m, 1H), 1.74-1.69 (m, 1H), 1.48-1.42 (m, 1H).

Example 4.20 (Cis Enantiomer 2)

Obtained as a white solid (67.8 mg, 19%). Chiral-Prep-HPLC retention time, 8.80 min; LCMS (ESI): $M+H^+=356.2$; 1H NMR (300 MHz, CD_3OD) δ 7.08-7.01 (m, 4H), 6.40 (s, 1H), 4.96 (s, 2H), 2.98-2.94 (m, 1H), 2.42 (s, 3H), 2.05-1.96 (m, 1H), 1.84-1.77 (m, 1H), 1.62-1.55 (m, 1H).

Examples 4.21 and 4.22

The racemic trans isomer (Ex. 4.18A) was purified by Chiral-Prep-HPLC with the following conditions (Prep-HPLC-032): Column, Chiralpak IA, 2*25 cm, 5 μ m; mobile phase, Hex (1% TEA)/IPA (hold 50.0% IPA in 8 mins), flow, 1.0 mL/min; Detector, UV 254 nm to afford two enantiomers.

Example 4.21 (Trans Enantiomer 1)


Obtained as a white solid (7.1 mg, 1%). Chiral-Prep-HPLC retention time, 2.28 min; LCMS (ESI): $M+H^+=356.2$; 1H NMR (400 MHz, $CDCl_3$) δ 7.05-6.91 (m, 4H), 6.47 (s, 1H), 4.92 (s, 2H), 3.01-2.95 (m, 1H), 2.43 (s, 3H), 2.06-2.00 (m, 1H), 1.86-1.70 (m, 1H), 1.62-1.57 (m, 1H).

Example 4.22 (Trans Enantiomer 2)


Obtained (4.4 mg, 0.5%). Chiral-Prep-HPLC retention time, 3.97 min; LCMS (ESI): $M+H^+=356.2$; 1H NMR (400 MHz, $CDCl_3$) δ 7.05-6.93 (m, 2H), 6.95-6.91 (m, 2H), 6.47 (s, 1H), 4.92 (s, 2H), 3.01-2.95 (m, 1H), 2.43 (s, 3H), 2.06-2.66 (m, 1H), 1.86-1.80 (m, 1H), 1.60-1.52 (m, 1H).

104

Example 4.23: 7-(4-Fluorophenoxymethyl)-3-[cis-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one




Step 1: tert-Butyldimethyl[[3-(tetramethyl-1,3,2-dioxaborolan-2-yl)prop-2-yn-1-yl]oxy]silane



To a solution of tert-butyldimethyl(prop-2-yn-1-yloxy)silane (200 mg, 1.17 mmol) in tetrahydrofuran (6 mL) was added 2.5 M n-butyl lithium (0.57 mL, 1.42 mmol) dropwise at $-78^\circ C$. The resulting solution was stirred for 0.5 h at $-78^\circ C$. Then 4,4,5,5-tetramethyl-2-(propan-2-yloxy)-1,3,2-dioxaborolane (230 mg, 1.24 mmol) was added dropwise at $-78^\circ C$. The resulting solution was allowed to react for an additional 4 h while the temperature was maintained at $-78^\circ C$. The reaction was then quenched by hydrogen chloride in ethyl ether (1 mol/L). After concentration in vacuo, the residue was diluted with ethyl ether (20 mL) and the solids were filtered off. The filtrate was concentrated to afford tert-butyldimethyl[[3-(tetramethyl-1,3,2-dioxaborolan-2-yl)prop-2-yn-1-yl]oxy]silane as a light yellow liquid (260 mg, 75%). 1H NMR (300 MHz, $CDCl_3$) δ 4.35 (s, 2H), 1.27 (s, 12H), 0.90 (s, 9H), 0.12 (s, 6H).

Step 2: tert-butyldimethyl[[[(2Z)-3-(tetramethyl-1,3,2-dioxaborolan-2-yl)prop-2-en-1-yl]oxy]silane




To a suspension of bis(cyclopentadienyl)zirconium chloride hydride (230 mg, 0.88 mmol) in tetrahydrofuran (5 mL) was added tert-butyldimethyl[[3-(tetramethyl-1,3,2-dioxaborolan-2-yl)prop-2-yn-1-yl]oxy]silane (260 mg, 0.88 mmol) in tetrahydrofuran (2 mL) dropwise with stirring at room temperature. The resulting solution was stirred overnight at room temperature and then quenched by water (5 mL). The resulting mixture was stirred for an additional 1 h

106

at room temperature and was extracted with dichloromethane (2x20 mL). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1/50) to afford tert-butyldimethyl[[[(2Z)-3-(tetramethyl-1,3,2-dioxaborolan-2-yl)prop-2-en-1-yl]oxy]silane as a colorless oil (200 mg, 76%). ¹H NMR (400 MHz, CDCl₃) δ 6.58-6.47 (m, 1H), 5.35-5.30 (m, 1H), 4.34-4.33 (m, 2H), 1.27 (s, 12H), 0.90 (s, 9H), 0.12 (s, 6H).

Step 3: tert-Butyldimethyl[[2-(tetramethyl-1,3,2-dioxaborolan-2-yl)cyclopropyl]methoxy]silane



Diethylzinc (1.0 M in hexanes) (4 mL, 4 mmol) was added to freshly distilled dichloromethane (4 mL) under nitrogen. Then trifluoroacetic acid (0.31 mL, 4.00 mmol) in dichloromethane (2 mL) was added dropwise at 0° C. Upon stirring for 20 min, a solution of diethylzinc in hexanes (0.32 mL, 4.00 mmol) in dichloromethane (2 mL) was added at 0° C. After an additional 20 min of stirring, a solution of tert-butyldimethyl[[[(2Z)-3-(tetramethyl-1,3,2-dioxaborolan-2-yl)prop-2-en-1-yl]oxy]silane (600 mg, 2.01 mmol) in dichloromethane (2 mL) was added at 0° C. Then the resulting solution was stirred 1 h at room temperature and was quenched with water. The reaction was extracted with dichloromethane (100 mLx2), washed with brine, dried over anhydrous sodium sulfate and concentrated to afford tert-butyldimethyl((2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)cyclopropyl)methoxy)silane as a colorless oil (600 mg, 96%).


Step 4: Potassium cis-2-(hydroxymethyl)cyclopropyltrifluoroborate



To a solution of tert-butyldimethyl((2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)cyclopropyl)methoxy)silane (600 mg, 2.00 mmol) in methanol (4 mL) was added potassium difluoride (630 mg, 8 mmol) in water (2 mL) drop wise at 0° C. After stirred 1.5 h at room temperature, the reaction mixture was concentrated in vacuo. The resulting solid was suspended in acetone (20 mL) and refluxed 20 min. The heterogeneous mixture was then filtered to remove potassium difluoride and the filtrate was concentrated. The extraction process was repeated for the filtered solid. The combined filtrates were concentrated and dissolved in minimal acetone followed by the slow addition of ethyl ether until the solution become cloudy. The solids were collected by filtration and dried to afford the title compound as a white solid (150 mg, 43%).


106

Step 5: 7-(4-Fluorophenoxymethyl)-3-[cis-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one




3-Bromo-7-(4-fluorophenoxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 4.1, Step 7) (100 mg, 0.27 mmol), 1,1'-bis(diphenylphosphino)ferrocene-palladium dichloride (20.0 mg, 0.03 mmol), sodium carbonate (60.0 mg, 0.57 mmol), potassium cis-2-(hydroxymethyl)cyclopropyltrifluoroborate (100 mg, 0.56 mmol), acetonitrile (3 mL) and water (0.5 mL) were placed in a 10-mL sealed tube. The final reaction mixture was heated in a microwave reactor for 1.5 h at 120° C. The mixture was extracted with dichloromethane (20 mL), washed with brine (10 mL), dried over anhydrous sodium sulfate and concentrated. The residue was purified by chromatography with ethyl acetate/petroleum ether (2:1) to give 7-(4-fluorophenoxymethyl)-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a white solid (23.5 mg, 24%). LCMS (ESI): M+H⁺=361.0; ¹H NMR (300 MHz, CDCl₃) δ 7.02-6.88 (m, 4H), 6.45 (s, 1H), 4.90 (s, 2H), 3.69-3.64 (m, 1H), 3.17-3.11 (m, 1H), 2.55-2.47 (m, 1H), 2.42 (s, 3H), 1.68-1.59 (m, 1H), 1.47-1.40 (m, 1H), 0.83-0.77 (m, 1H).

Example 4.24: trans-7-(4-Fluorophenoxymethyl)-2-methyl-3-[2-(trifluoromethyl)cyclopropyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



Step 1: 2-Diazo-1,1,1-trifluoroethane




To a solution of 2,2,2-trifluoroethan-1-amine hydrochloride (3.24 g, 23.9 mmol) in water (5 mL) and ethyl ether (10 mL) was added dropwise a solution of sodium nitrite (1.84 g, 26.7 mmol) in water (2 mL). The resulting solution was

107


stirred for 3 h at room temperature. The solids were filtered out to afford 2-diazo-1,1,1-trifluoroethane as a light yellow liquid (1.32 g, 51%). No LCMS signal.

Step 2: 4,4,5,5-Tetramethyl-2-[2-(trifluoromethyl)cyclopropyl]-1,3,2-dioxaborolane



To a solution of 2-diazo-1,1,1-trifluoroethane (530 mg, 4.82 mmol) in ether (100 mL) was added palladium acetate (50.0 mg, 0.22 mmol). Then 2-ethenyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (0.66 g, 4.29 mmol) and palladium acetate (50.0 mg, 0.22 mmol) were added with stirring over 20 min. After the resulting solution was stirred for 1 h at room temperature, the solids were filtered off. The resulting solution was concentrated in vacuo to afford 4,4,5,5-tetramethyl-2-[2-(trifluoromethyl)cyclopropyl]-1,3,2-dioxaborolane as dark green oil (580 mg, 51%).


Step 3: trans-7-(4-Fluorophenoxymethyl)-2-methyl-3-[2-(trifluoromethyl)cyclopropyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one




To a solution of 4,4,5,5-tetramethyl-2-[2-(trifluoromethyl)cyclopropyl]-1,3,2-dioxaborolane (384 mg, 1.63 mmol) in acetonitrile (3 mL) and water (1 mL) was added 3-bromo-7-(4-fluorophenoxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (300 mg, 0.81 mmol), 1,1'-bis(diphenylphosphino)ferrocenepalladium dichloride (59 mg, 0.08 mmol) and sodium carbonate (12 mg). The resulting solution was stirred for 1.5 h at 120° C. under nitrogen atmosphere and then diluted with water (5 mL). After extraction with dichloromethane (3x30 mL), the combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The crude product was purified by Prep-HPLC (Conditions: (Prep-HPLC-005): Column, Xbridge Prep C₁₈ OBD Column, 5 um, 19x150 mm; mobile phase, water with 10 mmol ammonium dicarbonate and acetonitrile (35.0% acetonitrile up to 59.0% in 10 min); Detector, UV 254/220 nm) to afford 7-(4-fluorophenoxymethyl)-2-methyl-3-[2-(trifluoromethyl)cyclopropyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a white solid (4.40 mg, 1%). LCMS (ESI): M+H⁺=399.2; ¹H NMR (300 MHz, CDCl₃) δ 7.02-6.96 (m, 2H), 6.92-6.87 (m, 1H), 6.42 (s, 1H), 4.90 (s, 2H), 2.81-2.76 (m, 1H), 2.42 (s, 3H), 1.98-1.87 (m, 1H), 1.55-1.48 (m, 1H), 1.18-1.10 (m, 1H).

108

Example 4.25: 7-(4-Fluorophenoxymethyl)-2-methyl-3-(2-methylcyclopropyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one




Step 1: 4,4,5,5-Tetramethyl-2-(2-methylcyclopropyl)-1,3,2-dioxaborolane



To a solution of 1 M diethylzinc in hexanes (0.36 mL, 0.36 mmol) in dichloromethane (4 mL) was added a solution of trifluoroacetic acid (408 mg, 3.58 mmol) in dichloromethane (4.0 mL), followed by a solution of diiodomethane (957 mg, 3.57 mmol) in dichloromethane (4.0 mL) under nitrogen and the reaction solution was stirred for 40 min at 0° C. Then a solution of 4,4,5,5-tetramethyl-2-[(1E)-prop-1-en-1-yl]-1,3,2-dioxaborolane (300 mg, 1.79 mmol) in dichloromethane (2 mL) was added and the reaction mixture was stirred for an additional 50 min at room temperature. The reaction was quenched by a saturated ammonium chloride solution (10 mL), extracted with petroleum ether (3x20 mL), washed with brine, dried with anhydrous sodium sulfate and concentrated in vacuo to afford 4,4,5,5-tetramethyl-2-(2-methylcyclopropyl)-1,3,2-dioxaborolane as a yellow solid (300 mg). The crude product was used in next step without further purification. LCMS (ESI): M+H⁺=376.1; ¹H NMR (300 MHz, CD₃OD) δ 1.21 (s, 12H), 1.08-1.06 (m, 3H), 0.95-0.91 (m, 1H), 0.69-0.65 (m, 1H), 0.38-0.32 (m, 1H), -0.42 to -0.47 (m, 1H).


Step 2: 7-(4-Fluorophenoxymethyl)-2-methyl-3-(2-methylcyclopropyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one




109

To a solution of 3-bromo-7-(4-fluorophenoxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 4.1, Step 7) (100 mg, 0.27 mmol) in acetonitrile/water (1.5/0.5 mL) was added 4,4,5,5-tetramethyl-2-(2-methylcyclopropyl)-1,3,2-dioxaborolane (100 mg, 0.55 mmol), 1,1'-bis(diphenylphosphino)ferrocenepalladium dichloride (20.3 mg, 0.03 mmol) and sodium carbonate (57.0 mg, 0.54 mmol). The reaction mixture was heated under microwave irradiation for 90 min at 120° C. The reaction was then concentrated under vacuum and the resulting residue was purified by chromatography with dichloromethane/methanol (50/1) to afford 7-(4-fluorophenoxymethyl)-2-methyl-3-(2-methylcyclopropyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as an off-white solid (17.6 mg, 18%). LCMS (ESI): $M+H^+=345.1$; 1H NMR (300 MHz, $CDCl_3$) δ 7.01-6.95 (m, 2H), 6.91-6.86 (m, 2H), 6.38 (s, 1H), 4.88 (s, 2H), 2.36 (s, 3H), 1.94-1.90 (m, 1H), 1.24-1.22 (m, 3H), 1.03-0.95 (m, 1H), 0.93-0.85 (m, 2H).

Example 4.26: trans-2-[2-[7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropyl]acetonitrile



Step 1: [2-[7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropyl]methylmethanesulfonate




To a solution of 7-((3-fluorophenoxy)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (from Example 4.1, Step 8) (40.0 mg, 0.11 mmol) in dichloromethane (51 mL) was added triethylamine

110



(34.0 mg, 0.34 mmol) and methanesulfonyl chloride (38.0 mg). After stirring for 1 h at room temperature, the resulting mixture was concentrated in vacuo. The residue was purified by silica gel chromatography with 2% methanol in dichloromethane to afford [2-[7-(4-fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropyl]methyl methanesulfonate as an off-white solid (40.0 mg, 82%). LCMS (ESI): $M+H^+=439$; 1H NMR (300 MHz, $CDCl_3$) δ 7.03-6.89 (m, 4H), 6.41 (s, 1H), 4.95 (s, 2H), 4.65-4.59 (s, 2H), 2.43 (s, 3H), 2.42-2.21 (m, 1H), 1.56-1.02 (m, 3H).

Step 2: trans-2-[2-[7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropyl]acetonitrile




To a solution of [2-[7-(4-fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropyl]methyl methanesulfonate (50.0 mg, 0.11 mmol) in dimethyl sulfoxide (5 mL) was added sodium cyanide (50.0 mg). The resulting solution was stirred for 1 h at 90° C. After cooling down to room temperature, the reaction mixture was diluted with dichloromethane (10 mL), washed with water (4x5 mL), dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by Prep-HPLC (Conditions: Column, SunFire Prep C_{18} OBD Column, 5 μ m, 19*150 mm; mobile phase, Water with 10 mmol ammonium bicarbonate and acetonitrile (18.0% acetonitrile up to 28.0% in 10 min, up to 95.0% in 2 min, down to 18.0% in 2 min); Detector, UV 254/220 nm) to afford 2-[2-[7-(4-fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropyl]acetonitrile as a white solid (8.90 mg, 21%). LCMS (ESI): $M+H^+=370.0$; 1H NMR (300 MHz, $CDCl_3$) δ 7.04-6.89 (m, 4H), 6.41 (s, 1H), 4.94 (s, 2H), 3.02-2.95 (m, 1H), 2.56-2.45 (m, 1H), 2.42 (s, 3H), 2.06-1.96 (m, 1H), 1.32-1.23 (m, 2H), 1.20-1.09 (m, 1H).


The following examples were prepared in a manner similar to Example 4.26:

No.	Structure/Name	LCMS (M + H)	¹ H NMR
4.27		375.1	¹ H NMR (300 MHz, CDCl ₃) δ 7.01-6.96 (m, 2H), 6.92-6.87 (m, 2H), 6.39 (s, 1H), 4.91 (s, 2H), 3.69-3.64 (m, 1H), 3.38 (s, 3H), 3.32-3.12 (m, 1H), 2.42 (s, 3H), 2.19-2.16 (m, 1H), 1.45-1.34 (m, 1H), 1.13-1.07 (m, 1H), 0.98-0.92 (m, 1H)
4.28		363.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.03-6.89 (m, 4H), 6.38 (s, 1H), 4.95 (s, 2H), 4.73-4.27 (m, 2H), 2.40 (s, 3H), 2.28-2.26 (m, 1H), 1.48-1.40 (m, 1H), 1.26-1.02 (m, 2H)

Example 4.29: 6-fluoro-7-((4-fluorophenoxy)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



Step 1: 3-bromo-7-(chloromethyl)-6-fluoro-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



Into a 30-mL sealed tube purged and maintained with nitrogen was added a solution of 3-bromo-7-(chloromethyl)-


35

2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (from Example 4.1, Step 6) (500 mg, 1.70 mmol) in acetonitrile (20 mL) and 1-chloromethyl-4-fluoro-1,4-diazoniabicyclo [2.2.2]octane bis(tetrafluoroborate) (Selectfluor®; 600 mg, 1.69 mmol). The resulting solution was stirred for 3 h at 75° C. After cooling down to room temperature, the reaction mixture was concentrated in vacuo. The residue was purified by silica gel chromatography with dichloromethane to afford 3-bromo-7-(chloromethyl)-6-fluoro-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as off-white solid (250 mg, 40%). LCMS (ESI): M+H⁺=311.0, 313.0; ¹H NMR (300 MHz, CDCl₃) δ 4.55 (s, 2H), 2.41 (s, 3H).

Step 2: 3-bromo-6-fluoro-7-((4-fluorophenoxy)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one

55

60




To a solution of 3-bromo-7-(chloromethyl)-6-fluoro-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (300 mg, 0.96 mmol) in acetonitrile (20 mL) was added potassium

113


iodide (83.0 mg, 0.48 mmol), potassium carbonate (276 mg, 2.00 mmol) and 4-fluorophenol (224 mg, 2.00 mmol). The reaction mixture was stirred overnight at 80° C. After cooling down to room temperature, the reaction mixture was concentrated in vacuo. The residue was purified on a silica gel column with ethyl acetate/petroleum ether (1/2) to afford 3-bromo-6-fluoro-7-(4-fluorophenoxy)methyl-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as an off-white solid (50.0 mg, 11%). LCMS (ESI): M+H=387.0, 389.0.

Step 3: 6-fluoro-7-((4-fluorophenoxy)methyl)-3-(2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one




Into a 10-mL sealed tube purged and maintained with nitrogen was added a solution of 3-bromo-6-fluoro-7-(4-fluorophenoxy)methyl-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (150 mg, 0.39 mmol) in acetonitrile/water (3/1 mL), 1,1'-bis(diphenylphosphino)ferrocene-palladium-dichloride (29.0 mg, 0.04 mmol), sodium carbonate (82.0 mg, 0.77 mmol) and potassium trans-2-(hydroxymethyl)cyclopropyltrifluoroborate (from Example 4.1, Step 2) (138 mg, 0.78 mmol). The reaction mixture was stirred for 1.5 h at 120° C. After cooling down to room temperature, the reaction mixture was concentrated in vacuo. The residue was purified on a silica gel column with dichloromethane/methanol (100/1) to afford 6-fluoro-7-((4-fluorophenoxy)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one as off-white solid (26.1 mg, 18%). LCMS (ESI): M+H+=379.0; ¹H NMR (300 MHz, CDCl₃) δ 6.98-6.91 (m, 4H), 5.02 (s, 2H), 4.06-4.01 (m, 1H), 3.25-3.18 (m, 1H), 2.39 (s, 3H), 2.33-2.27 (m, 1H), 1.30-1.25 (m, 1H), 1.08-0.99 (m, 2H).

Example 4.30: 7-(4-Fluorophenoxy)methyl-3-(3-hydroxypropyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one




114

Step 1: Ethyl (E)-ethyl 3-(7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)acrylate



To a solution of 3-bromo-7-((3-fluorophenoxy)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (from Example 4.1, Step 7) (205 mg, 0.56 mmol) in acetonitrile (5 mL) was added ethyl prop-2-enoate (110 mg, 1.10 mmol), tri-tolylphosphine (25 mg), tris(dibenzylideneacetone)dipalladium (33.0 mg, 0.04 mmol) and triethylamine (110 mg, 1.09 mmol). The reaction mixture was stirred overnight at 90° C. under a nitrogen atmosphere. After cooling down to room temperature, the reaction mixture was concentrated in vacuo and the residue was purified by chromatography with 3% ethyl acetate in petroleum ether to afford ethyl (E)-ethyl 3-(7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)acrylate as an off-white solid (58.0 mg, 27%). LCMS (ESI): M+H+=389.0.


Step 2: ethyl 3-(7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)propanoate



To a solution of ethyl (E)-ethyl 3-(7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)acrylate (58.0 mg, 0.15 mmol) in methanol (5 mL) was added palladium on carbon (50.0 mg). The reaction mixture was stirred overnight at room temperature under a hydrogen atmosphere (1.5 atm). After the solids were filtered off, the resulting mixture was concentrated in vacuo to afford ethyl 3-(7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)propanoate (52.0 mg, 89%) as a white solid. LCMS (ESI): M+H+=391.0.

115

Step 3: 7-(4-Fluorophenoxymethyl)-3-(3-hydroxypropyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one




116

To a solution of ethyl 3-[7-(4-fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]propanoate (42.0 mg, 0.11 mmol) in tetrahydrofuran (4 mL) and methanol (1 mL) was added lithium borohydride (23.0 mg).
 5 The reaction mixture was stirred overnight at room temperature. The reaction was then quenched with water (5 mL), extracted with dichloromethane (3×10 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by silica gel chromatography with 3% methanol in dichloromethane to afford 7-(4-Fluorophenoxymethyl)-3-(3-hydroxypropyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a white solid (6.3 mg, 1.7%). LCMS (ESI): $M+H^+=349.1$; 1H NMR (300 MHz, $CDCl_3$) δ 7.02-6.91 (m, 4H), 6.46 (s, 1H), 4.92 (s, 2H), 3.70-3.65 (m, 2H), 3.36-3.32 (m, 2H), 2.36 (s, 3H), 1.96-1.89 (m, 2H).

The following compounds were prepared using methods analogous to those described above:


No.	Structure/Name	LCMS (M + H)	1H NMR
4.31	<p>7-(4-Fluorophenoxymethyl)-3-(methoxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one</p>	335.20	1H NMR (300 MHz, CD_3OD) δ 7.08-6.98 (m, 4H), 6.39 (s, 1H), 5.01-4.96 (m, 4H), 3.41 (s, 3H), 2.49 (s, 3H).
4.32	<p>7-((4-fluorophenoxy)methyl)-3-(3-hydroxyoxetan-3-yl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	363.1	1H NMR (400 MHz, $DMSO-d_6$) δ 8.49 (s, 1H), 7.20-7.09 (m, 2H), 7.09-6.93 (m, 2H), 6.25 (s, 1H), 4.98 (s, 2H), 4.92 (d, J = 7.8 Hz, 2H), 4.63-4.45 (m, 2H), 2.26 (s, 3H).
4.33	<p>7-(4-Fluorophenoxymethyl)-3-(4-hydroxybutan-2-yl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one</p>	363.0	1H NMR (400 MHz, $DMSO-d_6$) δ 7.03-6.98 (m, 2H), 6.93-6.90 (m, 2H), 6.46 (s, 1H), 5.06 (s, 1H), 4.94 (s, 2H), 3.70-3.66 (m, 1H), 3.57 (m, 1H), 2.46 (m, 4H), 2.08-1.92 (m, 2H), 1.36-1.33 (m, 3H)

-continued


No.	Structure/Name	LCMS (M + H)	¹ H NMR
4.34	 <p>7-(4-Fluorophenoxymethyl)-3-[2-(2-hydroxypropan-2-yl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one</p>	371.1	¹ H NMR (300 MHz, CDCl ₃) δ 7.01-6.96 (m, 2H), 6.91-6.86 (m, 2H), 6.45 (s, 1H), 4.90 (s, 2H), 2.57-2.50 (m, 1H), 2.39 (s, 3H), 1.28 (s, 3H), 1.25-1.20 (m, 1H), 1.06 (s, 3H), 1.05-0.90 (m, 2H)

Method 5

Example 5.1: 7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-pyrimidin-5-yl-thiazolo[3,2-a]pyrimidin-5-one



Step 1: 3-bromo-7-[(ethyl(4-fluorophenyl)amino)methyl]-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



A mixture of 3-bromo-7-(chloromethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (from Example 4.1, Step 6) (30 g, 0.1 mol), N-ethyl-4-fluoroaniline (18.5 g, 0.13 mol), potassium carbonate (28.2 g, 0.2 mol) and sodium iodide (7.66 g, 0.05 mol) in acetonitrile was heated overnight at 80° C. The mixture was then cooled to room temperature, diluted with a saturated aqueous solution of ammonium chloride (300 mL) and extracted with dichloromethane (200 mL×3). The combined organic layers were washed with water and brine, dried over sodium sulfate and concentrated. The


20

residue was triturated with dichloromethane to give 3-bromo-7-[(ethyl(4-fluorophenyl)amino)methyl]-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (21 g, 54%) as an off-white solid.

25

Step 2: 7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-pyrimidin-5-yl-thiazolo[3,2-a]pyrimidin-5-one

30



35

40

45

To a solution of 3-bromo-7-[(ethyl(4-fluorophenyl)amino)methyl]-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (50 mg, 0.13 mmol) in acetonitrile/H₂O (1.3/0.5 mL) was added (pyrimidin-5-yl) boronic acid (20.3 mg, 0.16 mmol), sodium carbonate (40.5 mg, 0.379 mmol) and [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (9.7 mg, 0.013 mmol). The reaction mixture was heated under microwave irradiation for 20 min at 120° C. After cooling down room temperature, the resulting mixture was concentrated in vacuo. The residue was purified by silica gel chromatography with dichloromethane/methanol (95/5) to afford 7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-pyrimidin-5-yl-thiazolo[3,2-a]pyrimidin-5-one (14.8 mg, 30.0%) as an off-white solid. LCMS (ESI): M+H⁺=396.1; ¹H NMR (400 MHz, DMSO-d₆) δ 9.18 (s, 1H), 8.83 (s, 2H), 6.98 (t, J=8.8 Hz, 2H), 6.62 (dd, J=9.2, 4.4 Hz, 2H), 5.85 (s, 1H), 4.36 (s, 2H), 3.47 (q, J=7.0 Hz, 2H), 2.22 (s, 3H), 1.13 (t, J=7.0 Hz, 3H).





50

55





60

65





The following examples were prepared in a manner similar to Example 5.1:

No.	Structure/Name	LCMS (M + H)	¹ H NMR
5.2	 <p data-bbox="326 638 675 705">7-(((4-fluorophenyl)(methyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	382.1	¹ H NMR (300 MHz, CDCl ₃) δ 9.90 (s, 1H), 9.24 (s, 2H), 6.96-6.88 (m, 2H), 6.63-6.59 (m, 2H), 6.07 (s, 1H), 4.35 (s, 2H), 3.06 (s, 3H), 2.25 (s, 3H).
5.3	 <p data-bbox="326 1037 675 1125">7-(((4-fluorophenyl)(2,2,2-trifluoroethyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	450.1	¹ H NMR (300 MHz, CD ₃ OD) δ 9.13 (s, 1H), 8.76 (s, 2H), 6.94-6.86 (m, 2H), 6.79-6.75 (m, 2H), 5.97 (s, 1H), 5.54 (s, 2H), 4.25-4.16 (m, 2H), 2.25 (s, 3H).
5.4	 <p data-bbox="326 1457 675 1524">7-(((3,4-difluorophenyl)(ethyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	413.8	¹ H NMR (300 MHz, CD ₃ OD) δ 9.17 (s, 1H), 8.81 (s, 2H), 7.07-6.97 (m, 1H), 6.57-6.49 (m, 1H), 6.42-6.38 (m, 1H), 6.01 (s, 1H), 4.40 (s, 2H), 3.52 (m, 2H), 2.29 (s, 3H), 1.22 (m, 3H).
5.5	 <p data-bbox="326 1856 675 1927">7-((ethyl(3-fluorophenyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	396.09	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.19 (s, 1H), 8.84 (s, 2H), 7.15-7.12 (m, 1H), 6.47-6.39 (m, 3H), 5.84 (s, 1H), 4.42 (s, 2H), 3.51-3.49 (m, 2H), 2.23 (s, 3H), 1.16-1.13 (m, 3H).





-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
5.6	 <p data-bbox="326 674 675 758">7-(((2,2-difluoroethyl)(4-fluorophenyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	432.0	¹ H NMR (300 MHz, CD ₃ OD) δ 9.20 (s, 1H), 8.89 (s, 2H), 6.90-6.77 (m, 2H), 6.30-5.93 (m, 2H), 4.63 (s, 2H), 3.93-3.83 (m, 2H), 2.29 (s, 3H).
5.7	 <p data-bbox="326 1083 675 1146">7-((ethyl(pyridine-2-yl)amino)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	371.0	¹ H NMR (300 MHz, CDCl ₃) δ 8.15-8.13 (m, 1H), 7.48-7.42 (m, 1H), 6.60-6.50 (m, 2H), 6.10 (s, 1H), 4.61 (s, 2H), 4.24-4.21 (m, 1H), 4.07-4.01 (m, 1H), 3.62-3.55 (m, 2H), 3.10-3.03 (m, 1H), 2.37 (s, 3H), 2.27-2.25 (m, 1H), 1.26-1.21 (m, 4H), 1.05-0.95 (m, 2H).
5.8	 <p data-bbox="326 1472 675 1535">7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(thiazol-2-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	401.0	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.00-7.91 (m, 2H), 7.00-6.94 (m, 2H), 6.70-6.62 (m, 2H), 5.84 (s, 1H), 4.35 (s, 2H), 3.48-3.45 (m, 2H), 2.20 (s, 3H), 1.14-1.10 (m, 3H).
5.9	 <p data-bbox="326 1860 675 1925">7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(thiophen-2-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	400.1	¹ H NMR (300 MHz, CD ₃ OD) δ 7.61-7.59 (m, 1H), 7.11-7.07 (m, 2H), 6.92-6.85 (m, 2H), 6.67-6.62 (m, 2H), 6.00 (s, 1H), 4.36 (s, 2H), 3.52-3.49 (m, 2H), 2.24 (s, 3H), 1.21-1.18 (m, 3H).





-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
5.10	 <p data-bbox="332 661 672 724">3-((ethyl((2-methyl-5-oxo-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)amino)benzonitrile</p>	402.9	¹ H NMR (300 MHz, CD ₃ OD) δ 9.06 (s, 1H), 8.71 (s, 2H), 7.23-7.18 (m, 1H), 6.87-6.84 (m, 3H), 5.88 (s, 1H), 4.38 (s, 2H), 3.52-3.45 (m, 2H), 2.19 (s, 3H), 1.19-1.12 (m, 3H).
5.11	 <p data-bbox="337 1060 665 1123">3-(2-aminopyridin-3-yl)-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	409.8	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.94-7.90 (m, 1H), 7.25-7.24 (m, 1H), 7.00-6.94 (m, 2H), 6.63-6.52 (m, 3H), 5.90 (br, 2H), 5.74 (s, 1H), 4.31 (s, 2H), 3.46-3.44 (m, 2H), 2.05 (s, 3H), 1.14-1.10 (m, 3H)
5.12	 <p data-bbox="337 1459 665 1522">7-((ethyl(pyridine-2-yl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	379.0	¹ H NMR (300 MHz, CD ₃ OD) δ 9.13 (s, 1H), 8.77 (s, 2H), 7.99-7.97 (m, 1H), 7.51-7.45 (m, 1H), 6.66-6.55 (m, 2H), 5.91 (s, 1H), 4.61 (s, 2H), 3.65-3.57 (m, 2H), 2.26 (s, 3H), 1.21-1.19 (m, 3H).
5.13	 <p data-bbox="328 1864 673 1925">7-((4-fluorophenylamino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	368.0	¹ H NMR (300 MHz, CDCl ₃) δ 9.25 (s, 1H), 8.72 (s, 2H), 6.95-6.86 (m, 2H), 6.69-6.65 (m, 2H), 6.23 (s, 1H), 4.25 (s, 2H), 2.32 (s, 3H).





-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
5.14	 <p data-bbox="337 653 662 716">3-butyl-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	374.1	¹ H NMR (300 MHz, CD ₃ OD) δ 6.91-6.83 (m, 2H), 6.65-6.59 (m, 2H), 6.00 (s, 1H), 4.30 (s, 2H), 3.51-3.44 (m, 2H), 3.14-3.09 (m, 2H), 1.61-1.51 (m, 2H), 1.42-1.30 (m, 2H), 1.17-1.11 (m, 3H), 0.92-0.85 (m, 3H)
5.15	 <p data-bbox="326 1062 675 1125">2-cyclopropyl-7-[(N-ethyl-4-fluoroanilino)methyl]-3-pyrimidin-5-yl-thiazolo[3,2-a]pyrimidine-5-one</p>	422.1	¹ H NMR (300 MHz, CD ₃ OD) δ 9.17 (s, 1H), 8.89 (s, 2H), 6.93-6.87 (m, 2H), 6.68-6.64 (m, 2H), 6.05 (s, 1H), 4.38 (s, 2H), 3.57-3.51 (m, 2H), 1.99-1.91 (m, 1H), 1.23-1.20 (m, 3H), 1.11-1.04 (m, 2H), 0.91-0.79 (m, 2H).
5.16	 <p data-bbox="334 1461 667 1524">2-ethyl-7-((ethyl(4-fluorophenyl)amino)methyl)-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	410.0	¹ H NMR (300 MHz, CD ₃ OD) δ 9.18 (s, 1H), 8.82 (s, 2H), 6.92-6.89 (m, 2H), 6.68-6.64 (m, 2H), 6.04 (s, 1H), 4.39 (s, 2H), 3.37-3.34 (m, 2H), 2.69-2.67 (m, 2H), 1.29-1.14 (m, 6H).
5.17	 <p data-bbox="337 1850 662 1913">7-((ethyl(4-fluorophenyl)amino)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	402.1	¹ H NMR (300 MHz, CDCl ₃) δ 6.94-6.77 (m, 2H), 6.58-6.34 (m, 2H), 6.19 (s, 1H), 4.28 (s, 2H), 4.17-4.03 (m, 2H), 3.50-3.43 (m, 2H), 3.08-3.05 (m, 1H), 2.38 (s, 3H), 2.29-2.24 (m, 1H), 1.29-1.19 (m, 4H), 1.06-0.94 (m, 2H)





-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
5.18	 <p data-bbox="337 659 665 743">7-((ethyl(4-fluorophenyl)amino)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 1)</p>	402.1	¹ H NMR (300 MHz, CDCl ₃) δ 6.94-6.77 (m, 2H), 6.58-6.34 (m, 2H), 6.19 (s, 1H), 4.28 (s, 2H), 4.17-4.03 (m, 2H), 3.50-3.43 (m, 2H), 3.08-3.05 (m, 1H), 2.38 (s, 3H), 2.29-2.24 (m, 1H), 1.29-1.19 (m, 4H), 1.06-0.94 (m, 2H)
5.19	 <p data-bbox="337 1073 665 1157">7-((ethyl(4-fluorophenyl)amino)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 2)</p>	402.1	¹ H NMR (300 MHz, CDCl ₃) δ 6.94-6.77 (m, 2H), 6.58-6.34 (m, 2H), 6.19 (s, 1H), 4.28 (s, 2H), 4.17-4.03 (m, 2H), 3.50-3.43 (m, 2H), 3.08-3.05 (m, 1H), 2.38 (s, 3H), 2.29-2.24 (m, 1H), 1.29-1.19 (m, 4H), 1.06-0.94 (m, 2H)
5.20	 <p data-bbox="337 1488 675 1528">7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-thiazol-4-yl-thiazolo[3,2-a]pyrimidin-5-one</p>	401.2	
5.21	 <p data-bbox="337 1862 675 1906">7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one</p>	395.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.57 (dd, J = 5.0, 1.5 Hz, 1H), 8.54 (d, J = 2.0 Hz, 1H), 7.78 (dt, J = 8.1, 1.9 Hz, 1H), 7.42 (dd, J = 7.8, 5.0 Hz, 1H), 6.97 (t, J = 8.9 Hz, 2H), 6.62 (dd, J = 9.2, 4.3 Hz, 2H), 5.81 (s, 1H), 4.35 (s, 2H), 3.46 (q, J = 7.0 Hz, 2H), 2.16 (s, 3H), 1.13 (t, J = 6.9 Hz, 3H).





-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
5.22	 <p data-bbox="321 667 675 716">7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-phenyl-thiazolo[3,2-a]pyrimidin-5-one</p>	394.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 7.39 (d, J = 5.5 Hz, 3H), 7.32 (d, J = 5.2 Hz, 2H), 6.98 (t, J = 8.6 Hz, 2H), 6.62 (dd, J = 9.1, 4.3 Hz, 2H), 5.78 (s, 1H), 4.34 (s, 2H), 3.47 (q, J = 7.0 Hz, 2H), 2.13 (s, 3H), 1.13 (t, J = 7.0 Hz, 3H).
5.23	 <p data-bbox="342 982 656 1031">7-[(N-ethyl-4-fluoro-anilino)methyl]-2,3-dimethyl-thiazolo[3,2-a]pyrimidin-5-one</p>	332.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 6.98 (t, J = 8.6 Hz, 2H), 6.61 (dd, J = 9.0, 4.3 Hz, 2H), 5.81 (s, 1H), 4.30 (s, 2H), 3.47 (q, J = 7.0 Hz, 2H), 2.59 (s, 3H), 2.28 (s, 3H), 1.13 (t, J = 7.0 Hz, 3H).
5.24	 <p data-bbox="342 1392 656 1461">7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(5-fluoro-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	413.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.62 (d, J = 2.6 Hz, 1H), 8.45 (s, 1H), 7.91-7.79 (m, 1H), 6.98 (t, J = 8.7 Hz, 2H), 6.62 (dd, J = 9.1, 4.3 Hz, 2H), 5.83 (s, 1H), 4.36 (s, 2H), 3.47 (q, J = 7.0 Hz, 2H), 2.19 (s, 3H), 1.13 (t, J = 6.9 Hz, 3H).
5.25	 <p data-bbox="342 1791 656 1860">7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(2-fluoro-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	412	





-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
5.26	 <p data-bbox="354 632 651 695">3-cyclopropyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	358.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 6.98 (t, J = 8.8 Hz, 2H), 6.61 (dd, J = 9.2, 4.3 Hz, 2H), 5.82 (s, 1H), 4.28 (s, 2H), 3.46 (q, J = 7.0 Hz, 2H), 2.34 (s, 3H), 2.14 (p, J = 7.4 Hz, 1H), 1.13 (t, J = 7.0 Hz, 3H), 0.97-0.88 (m, 2H), 0.72-0.60 (m, 2H).
5.27	 <p data-bbox="326 1037 675 1079">7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-pyrazin-2-yl-thiazolo[3,2-a]pyrimidin-5-one</p>	396.1	
5.28	 <p data-bbox="326 1394 675 1457">7-[(N-ethyl-4-fluoro-anilino)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-5-one</p>	374.1	
5.29	 <p data-bbox="354 1814 651 1879">7-[(N-ethyl-4-fluoro-anilino)methyl]-3-isopropenyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	358.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 6.98 (t, J = 8.6 Hz, 2H), 6.61 (dd, J = 9.0, 4.3 Hz, 2H), 5.86 (s, 1H), 5.35 (s, 1H), 5.00 (s, 1H), 4.33 (s, 2H), 3.47 (q, J = 7.0 Hz, 2H), 2.28 (s, 2H), 1.94 (s, 3H), 1.13 (t, J = 7.0 Hz, 3H).

-continued


No.	Structure/Name	LCMS (M + H)	¹ H NMR
5.30	 <p data-bbox="326 663 675 726">7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-pyridazin-4-yl-thiazolo[3,2-a]pyrimidin-5-one</p>	396.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.28 (d, J = 6.7 Hz, 1H), 9.22 (d, J = 1.5 Hz, 1H), 7.72 (d, J = 5.1 Hz, 1H), 7.03-6.91 (m, 2H), 6.69-6.55 (m, 2H), 5.87 (s, 1H), 4.36 (s, 2H), 3.47 (q, J = 7.0 Hz, 2H), 2.22 (s, 3H), 1.13 (t, J = 7.0 Hz, 3H).
5.31	 <p data-bbox="334 1058 664 1125">3-(5-chloro-3-pyridyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	429.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.65 (d, J = 3.0 Hz, 1H), 8.54 (s, 1H), 8.03 (s, 1H), 6.98 (t, J = 8.7 Hz, 2H), 6.62 (dd, J = 9.2, 4.3 Hz, 2H), 5.83 (s, 1H), 4.36 (s, 2H), 3.47 (q, J = 7.0 Hz, 2H), 2.19 (s, 3H), 1.13 (t, J = 7.0 Hz, 3H).
5.32	 <p data-bbox="326 1465 675 1503">7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(4-pyridyl)thiazolo[3,2-a]pyrimidin-5-one</p>	395.1	
5.33	 <p data-bbox="326 1860 675 1929">7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(1-methylpyrazol-4-yl)thiazolo[3,2-a]pyrimidin-5-one</p>	398.2	

-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
5.34	 <p data-bbox="321 653 678 716">7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(1H-pyrazol-4-yl)thiazolo[3,2-a]pyrimidin-5-one</p>	384.1	
5.35	 <p data-bbox="342 1098 659 1161">5-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]pyridine-3-carbonitrile</p>	420.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.04 (s, 1H), 8.87 (d, J = 1.7 Hz, 1H), 8.41 (s, 1H), 6.98 (t, J = 8.7 Hz, 2H), 6.62 (dd, J = 9.3, 4.3 Hz, 2H), 5.84 (s, 1H), 4.36 (s, 2H), 3.47 (q, J = 7.0 Hz, 2H), 2.21 (s, 3H), 1.13 (t, J = 7.0 Hz, 3H).
5.36	 <p data-bbox="321 1518 678 1581">7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-[5-(trifluoromethyl)-3-pyridyl]thiazolo[3,2-a]pyrimidin-5-one</p>	463.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.00 (s, 1H), 8.89 (s, 1H), 8.31 (s, 1H), 6.98 (t, J = 8.7 Hz, 2H), 6.62 (dd, J = 9.0, 4.1 Hz, 2H), 5.83 (s, 1H), 4.36 (s, 2H), 3.47 (q, J = 7.0 Hz, 2H), 2.20 (s, 3H), 1.13 (t, J = 7.0 Hz, 3H).
5.37	 <p data-bbox="321 1896 678 1948">7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-oxazol-2-yl-thiazolo[3,2-a]pyrimidin-5-one</p>	385.1	¹ H NMR (300 MHz, CD ₃ OD) δ 8.09 (s, 1H), 7.38 (s, 1H), 6.94-6.88 (m, 2H), 6.69-6.65 (m, 2H), 6.09 (s, 1H), 4.40 (s, 2H), 3.56-3.51 (m, 2H), 2.36 (s, 3H), 1.24-1.19 (m, 3H)


137

Example 5.38: 7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidin-5-one




To a solution of 3-bromo-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 5.1, Step 1) (200 mg, 0.50 mmol) in N-methyl-pyrrolidone (5 mL) was added copper(I) iodide (12 mg, 0.06 mmol) and methyl 2,2-difluoro-2-(fluorosulfonyl)acetate (146 mg, 0.76 mmol). The reaction solution was stirred for 8 h at 120° C. The resulting mixture was concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/2) to afford 7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-3-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (21.1 mg, 11%) as a white solid. LCMS (ESI): M+H⁺=385.8; ¹H NMR (300 MHz, CDCl₃) δ 6.94-6.88 (m, 2H), 6.59-6.54 (m, 2H), 6.24 (s, 1H), 4.28 (s, 2H), 3.46-3.43 (m, 2H), 2.60-2.51 (m, 3H), 1.24-1.22 (m, 3H).

Example 5.39: 7-((ethyl(4-fluorophenyl)amino)methyl)-3-(trans-2-(fluoromethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one




Step 1: (2-(7-((ethyl(4-fluorocyclohexa-2,4-dienyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)cyclopropyl)methyl methanesulfonate



138


To a solution of 7-((ethyl(4-fluorocyclohexa-2,4-dienyl)amino)methyl)-3-(2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (from Example 5.17) (50 mg, 0.13 mmol) in dichloromethane (5 mL) was added methanesulfonyl chloride (22 mg, 0.19 mmol) and triethylamine (26 mg, 0.26 mmol). The reaction mixture was stirred 30 mins at room temperature. The reaction solution was concentrated in vacuo to afford (2-(7-((ethyl(4-fluorocyclohexa-2,4-dienyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)cyclopropyl)methyl methanesulfonate (60 mg, crude). The crude product was used in next step without further purification.

Step 2: 7-((ethyl(4-fluorophenyl)amino)methyl)-3-(trans-2-(fluoromethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of [2-(7-[[ethyl(4-fluorophenyl)amino]methyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropyl)methyl methanesulfonate (60 mg, 0.13 mmol) in propan-2-ol (0.5 mL) was added cesium fluoride (45.6 mg, 0.30 mmol). The reaction mixture was stirred for 90 min at 80° C. and then concentrated in vacuo. The residue was purified by Prep-HPLC to afford 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-[trans-2-(fluoromethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a yellow solid (3.1 mg, 6.0%). LCMS (ESI): M+H⁺=390.1; ¹H NMR (300 MHz, CDCl₃) δ 6.95-6.89 (m, 2H), 6.72-6.65 (m, 2H), 6.12 (s, 1H), 4.72-4.26 (m, 2H), 3.52-3.45 (m, 2H), 2.41 (s, 3H), 2.28-2.22 (m, 1H), 1.71-1.58 (m, 1H), 1.26-1.15 (m, 4H), 1.08-1.01 (m, 1H).

Example 5.40: 3-ethyl-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 3-ethenyl-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (prepared via a method similar to Example 5.1) (75.0 mg, 0.22 mmol) in methanol (15 mL) was added palladium on carbon (100 mg). The reaction mixture was stirred


139

overnight at room temperature under a hydrogen atmosphere. After filtration, the filtrate was concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/5) to afford 3-ethyl-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]

140



thiazolo[3,2-a]pyrimidin-5-one as a yellow solid (50 mg, 60%). LCMS (ESI): M+H⁺=346.1; ¹H NMR (300 MHz, CDCl₃) δ 6.96-6.90 (m, 2H), 6.73-6.67 (m, 2H), 6.14 (s, 1H), 4.30 (s, 2H), 3.54-3.47 (m, 2H), 3.19-3.14 (m, 2H), 2.32 (s, 3H), 1.38-1.24 (m, 6H).

The following compound was prepared using methods analogous to Example 5.40:





No.	Structure/Name	LCMS (M + H)	¹ H NMR
5.41	 <p>7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-propyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	360.1	¹ H NMR (300 MHz, CDCl ₃) δ 6.92-6.86 (m, 2H), 6.67-6.63 (m, 2H), 6.09 (s, 1H), 4.26 (s, 2H), 3.48-3.44 (m, 2H), 3.10-3.05 (m, 2H), 1.71-1.43 (m, 2H), 1.22-1.17 (m, 3H), 0.95-0.90 (m, 3H)

25





The following compounds were prepared using methods analogous to those described above:

No.	Structure/Name	LCMS (M + H)	¹ H NMR
5.42	 <p>7-[[4-fluoro-N-(2-fluoroethyl)anilino]methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methylthiazolo[3,2-a]pyrimidin-5-one (enantiomer 1)</p>	406.0	¹ H NMR (300 MHz, CDCl ₃) δ 6.95-6.90 (m, 2H), 6.66-6.64 (m, 2H), 6.16 (s, 1H), 4.73-4.71 (m, 1H), 4.62-4.95 (m, 1H), 4.45 (s, 2H), 4.05-4.02 (m, 1H), 4.01-3.72 (m, 2H), 3.14-3.08 (m, 1H), 2.39 (s, 3H), 2.27-2.00 (m, 1H), 1.29-1.24 (m, 1H), 1.05-0.97 (m, 2H).
5.43	 <p>7-[[4-fluoro-N-(2-fluoroethyl)anilino]methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methylthiazolo[3,2-a]pyrimidin-5-one (enantiomer 2)</p>	406.0	¹ H NMR (300 MHz, CDCl ₃) δ 6.94-6.89 (m, 2H), 6.63-6.60 (m, 2H), 6.15 (s, 1H), 4.73-4.70 (m, 1H), 4.61-4.59 (m, 1H), 4.43 (s, 2H), 4.05-4.02 (m, 1H), 3.80-3.71 (m, 2H), 3.12-3.07 (m, 1H), 2.39 (s, 3H), 2.27-2.23 (m, 1H), 1.28-1.25 (m, 1H), 1.05-0.96 (m, 2H).





-continued

No.	Structure/Name	LCMS (M + H) ⁺ ¹ H NMR
5.44	 <p data-bbox="342 653 659 722">7-((ethyl(4-fluorophenyl)amino)methyl)-3-(furan-3-yl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	384.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.49-7.47 (m, 2H), 6.90-6.85 (m, 2H), 6.58-6.54 (m, 2H), 6.45-6.44 (m, 1H), 6.09 (s, 1H), 4.28 (s, 2H), 3.50-3.41 (m, 2H), 2.27 (s, 3H), 1.22-1.18 (m, 3H).
5.45	 <p data-bbox="342 1041 659 1113">7-((ethyl(4-fluorophenyl)amino)methyl)-3-(furan-2-yl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	384.0 ¹ H NMR (300 MHz, CD ₃ OD) δ 7.63 (s, 1H), 6.93-6.87 (m, 2H), 6.67-6.52 (m, 4H), 6.04 (s, 1H), 4.37 (s, 2H), 3.57-3.48 (m, 2H), 2.32 (s, 3H), 1.23-1.20 (m, 3H).
5.46	 <p data-bbox="342 1409 659 1474">7-((5-fluoro-2-methylindolin-1-yl)methyl)-3-(furan-2-yl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	396.0 ¹ H NMR (300 MHz, CD ₃ OD) δ 7.65 (s, 1H), 6.81-6.63 (m, 1H), 6.72-6.54 (m, 3H), 6.26 (s, 1H), 6.24-6.19 (m, 1H), 4.13 (m, 2H), 3.84-3.76 (m, 1H), 3.23-3.15 (m, 1H), 2.71-2.63 (m, 1H), 2.33 (s, 3H), 1.31 (m, 3H).
5.47	 <p data-bbox="342 1793 659 1858">7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(thiophen-3-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	400.0 ¹ H NMR (300 MHz, CD ₃ OD) δ 7.45-7.42 (m, 2H), 7.07-7.06 (m, 1H), 6.93-6.87 (m, 2H), 6.67-6.63 (m, 2H), 6.00 (s, 1H), 4.36 (s, 2H), 3.51 (m, 2H), 2.23 (s, 3H), 1.21 (m, 3H).




-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
5.48	 <p data-bbox="334 604 667 667">7-((5-fluoro-2-methylindolin-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	408.0 ¹ HNMR (300 MHz, CD ₃ OD) δ 9.19 (s, 1H), 8.84 (s, 2H), 6.84-6.81 (m, 1H), 6.71-6.64 (m, 1H), 6.26 (s, 1H), 6.24-6.20 (m, 1H), 4.25-4.05 (m, 2H), 3.84-3.76 (m, 1H), 3.22-3.17 (m, 1H), 2.68-2.62 (m, 1H), 2.31 (s, 3H), 1.35-1.33 (m, 3H).
5.49	 <p data-bbox="334 1014 667 1077">7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(4-methylthiazol-2-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	415.1 ¹ H NMR (300 MHz, CD ₃ OD) δ 7.40 (s, 1H), 6.91-6.84 (m, 2H), 6.65-6.59 (m, 2H), 6.00 (s, 1H), 4.35 (s, 2H), 3.49-3.46 (m, 2H), 2.45 (s, 3H), 2.25 (s, 3H), 1.19-1.15 (m, 3H)
5.50	 <p data-bbox="334 1444 667 1507">7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(6-oxo-1,6-dihydropyridin-3-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	411.2 ¹ H NMR (300 MHz, DMSO-d ₆) δ 7.39-7.34 (m, 2H), 6.70-6.95 (m, 2H), 6.64-6.59 (m, 2H), 6.26 (m, 1H), 5.81 (s, 1H), 4.33 (s, 2H), 3.50-3.43 (m, 2H), 2.21 (s, 3H), 1.13-1.10 (m, 3H)
5.51	 <p data-bbox="334 1875 667 1938">3-(6-aminopyridin-3-yl)-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	409.9 ¹ H NMR (300 MHz, DMSO-d ₆) δ 7.77 (s, 1H), 7.36-7.26 (m, 1H), 6.96-6.90 (m, 2H), 6.59-6.54 (m, 2H), 6.40-6.32 (m, 1H), 6.22 (br, 2H), 5.74 (s, 1H), 4.28 (s, 2H), 3.45-3.38 (m, 2H), 2.12 (s, 3H), 1.10-1.02 (m, 3H)

-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
5.52	 <p data-bbox="337 667 662 730">7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(prop-1-ynyl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	356.0 ¹ H NMR (300 MHz, CDCl ₃) δ 6.93-6.87 (m, 2H), 6.59-6.55 (m, 2H), 6.16 (s, 1H), 4.28 (s, 2H), 3.50-3.43 (m, 2H), 2.45 (s, 3H), 2.23 (s, 3H), 1.27-1.20 (m, 3H)
5.53	 <p data-bbox="337 1014 662 1079">7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-vinyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	344.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.17-7.13 (m, 1H), 6.92-6.87 (m, 2H), 6.59-6.55 (m, 2H), 6.15 (s, 1H), 5.64-5.60 (m, 1H), 5.35-5.29 (m, 1H), 4.28 (s, 2H), 3.50-3.43 (m, 2H), 2.42 (s, 3H), 1.24-1.19 (m, 3H)
5.54	 <p data-bbox="337 1350 662 1409">3-bromo-2-cyclopropyl-7-[(N-ethyl-4-fluoroanilino)methyl]thiazolo[3,2-a]pyrimidin-5-one</p>	424.0 ¹ H NMR (300 MHz, CDCl ₃) δ 6.92-6.86 (m, 2H), 6.58-6.52 (m, 2H), 6.14 (s, 1H), 4.25 (s, 2H), 3.46-3.40 (m, 2H), 2.18-2.10 (m, 1H), 1.25-1.15 (m, 5H), 0.89-0.78 (m, 2H).
5.55	 <p data-bbox="337 1776 662 1839">3-(3,5-difluorophenyl)-7-[(N-ethyl-4-fluoroanilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	450.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 7.29 (tt, J = 9.4, 2.5 Hz, 1H), 7.16 (h, J = 4.8 Hz, 2H), 6.98 (t, J = 8.9 Hz, 2H), 6.62 (dd, J = 9.1, 4.3 Hz, 2H), 5.82 (s, 1H), 4.34 (s, 2H), 3.46 (q, J = 7.0 Hz, 2H), 2.16 (s, 3H), 1.13 (t, J = 7.0 Hz, 3H).


-continued

No.	Structure/Name	LCMS (M + H) ⁺ ¹ H NMR
5.56	 <p data-bbox="326 667 675 730">7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(1-methylpyrazol-3-yl)thiazolo[3,2-a]pyrimidin-5-one</p>	398.34
5.57	 <p data-bbox="334 1245 667 1314">3-(2-amino-4-pyridyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	410.1
5.58	 <p data-bbox="334 1822 667 1892">7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(5-methoxy-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	425.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.31 (d, J = 2.9 Hz, 1H), 8.14 (s, 1H), 7.43 (d, J = 2.3 Hz, 1H), 6.98 (t, J = 8.7 Hz, 2H), 6.62 (dd, J = 9.3, 4.3 Hz, 2H), 5.81 (s, 1H), 4.35 (s, 2H), 3.81 (s, 3H), 3.47 (q, J = 7.0 Hz, 2H), 2.16 (s, 3H), 1.13 (t, J = 7.0 Hz, 3H).

149

Method 6

Example 6.1: 7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-morpholino-5H-thiazolo[3,2-a]pyrimidin-5-one



150

To a solution of 3-bromo-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (250 mg, 0.63 mmol) (from Example 5.1, Step 1) in dimethyl sulfoxide (3 mL) was added potassium phosphate (268 mg, 1.26 mmol), L-proline (22.0 mg, 0.19 mmol), morpholine (165 mg, 1.89 mmol) and cuprous iodide (18.0 mg, 0.09 mmol). The reaction mixture was stirred overnight at 90° C. The resulting mixture was quenched with water (50 mL), extracted with ethyl acetate (30 mL×3), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/2) to afford 7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-3-(morpholin-4-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a white solid (29.5 mg, 12%). LCMS (ESI): M+H⁺=403.0; ¹H NMR (300 MHz, CDCl₃) δ 6.93-6.87 (m, 2H), 6.61-6.57 (m, 2H), 6.15 (s, 1H), 3.91-3.87 (m, 2H), 3.76-3.59 (m, 4H), 3.50-3.43 (m, 2H), 2.64-2.60 (m, 2H), 2.40 (s, 3H), 1.25-1.20 (m, 3H).


The following examples were prepared in a manner similar to Example 6.1:

No. Structure/Name	LCMS (M + H) ¹ H NMR
6.2 3-(dimethylamino)-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one	361.0 ¹ H NMR (300 MHz, CDCl ₃) δ 6.93-6.87 (m, 2H), 6.60-6.55 (m, 2H), 6.12 (s, 1H), 4.28 (s, 2H), 3.50-3.43 (m, 2H), 2.76 (s, 6H), 2.31 (s, 3H), 1.26-1.20 (m, 3H)
6.3 7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(pyrrolidin-1-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one	387.1 ¹ H NMR (300 MHz, CDCl ₃) δ 6.93-6.87 (m, 2H), 6.59-6.54 (m, 2H), 6.07 (s, 1H), 4.29 (s, 2H), 3.49-3.42 (m, 2H), 3.19-3.15 (m, 4H), 2.28 (s, 3H), 1.20-1.16 (m, 4H), 1.26-1.19 (m, 3H)

151

Method 7

Example 7.1: 7-(((3,4-difluorophenyl)(ethyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile







152


To a solution of 3-bromo-7-[[[(3,4-difluorophenyl)(ethyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (prepared via a method similar to Example 5.1, Step 1) (100 mg, 0.24 mmol) in N,N-dimethylformamide (5 mL) was added cuprous cyanide (43.0 mg, 0.48 mmol). The reaction solution was stirred for 1.5 h at 100° C. The reaction solution was quenched by water (50 mL), extracted with dichloromethane, washed with brine, dried over anhydrous magnesium sulfate and concentration in vacuo. The residue was purified by chromatography with dichloromethane/methanol (20/1) to afford 7-[[[(3,4-difluorophenyl)(ethyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carbonitrile (43.1 mg, 47%) as an off-white solid. LCMS (ESI): M+H⁺=361.0; ¹H NMR (400 MHz, CD₃OD) δ 7.08-7.01 (m, 1H), 6.59-6.53 (m, 1H), 6.430-6.40 (m, 1H), 6.16 (s, 1H), 4.41 (s, 2H), 3.56-3.51 (m, 2H), 2.67 (s, 3H), 1.31-1.21 (m, 3H).

The following examples were prepared in a manner similar to Example 7.1:

No.	Structure/Name	LCMS (M + H) ¹ H NMR
7.2	<p>7-((ethyl(3-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile</p>	343.1 ¹ H NMR (300 MHz, CDCl ₃) δ 7.19-7.17 (m, 1H), 6.46-6.32 (m, 3H), 6.25 (s, 1H), 4.34 (s, 2H), 3.53-3.50 (m, 2H), 2.66 (s, 3H), 1.27-1.24 (m, 3H).
7.3	<p>7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carbonitrile</p>	343.0 ¹ H NMR (400 MHz, DMSO-d ₆) δ 6.99 (t, J = 8.7 Hz, 2H), 6.62 (dd, J = 9.1, 4.2 Hz, 2H), 6.01 (s, 1H), 4.36 (s, 2H), 3.48 (q, J = 7.0 Hz, 2H), 2.61 (s, 3H), 1.13 (t, J = 7.0 Hz, 3H).
7.4	<p>7-[(N-ethyl-4-fluoro-anilino)methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carbonitrile</p>	329.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.70 (s, 1H), 7.05-6.87 (m, 2H), 6.71-6.54 (m, 2H), 6.02 (s, 2H), 4.37 (s, 2H), 3.47 (q, J = 7.0 Hz, 2H), 1.13 (t, J = 7.0 Hz, 3H).


The following compounds were prepared using methods analogous to those described above:

No.	Structure/Name	LCMS (M + H) ¹ H NMR
7.5	 <p>7-((5-fluoro-2-methylindolin-1-yl)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile</p>	355.0 ¹ H NMR (300 MHz, CD ₃ OD) δ 6.84-6.81 (m, 1H), 6.70-6.64 (m, 1H), 6.38 (s, 1H), 6.21-6.20 (m, 1H), 4.23-4.04 (m, 2H), 3.80-3.75 (m, 1H), 3.22-3.14 (m, 1H), 2.72-2.67 (m, 1H), 2.66 (s, 3H), 1.34-1.32 (m, 3H).
7.6	 <p>7-((3-cyanophenyl)(ethyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile</p>	350.0 ¹ H NMR (300 MHz, CD ₃ OD) δ 7.35-7.29 (m, 1H), 7.00-6.97 (m, 3H), 6.13 (s, 1H), 4.49 (s, 2H), 3.64-3.56 (m, 2H), 2.66 (s, 3H), 1.30-1.23 (m, 3H).
7.7	 <p>7-((ethyl(pyridin-2-yl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile</p>	326.1 ¹ H NMR (300 MHz, CDCl ₃) δ 8.12-8.11 (m, 1H), 7.50-7.45 (m, 1H), 6.61-6.52 (m, 2H), 6.19 (s, 1H), 4.65 (s, 2H), 3.61-3.54 (m, 2H), 2.65 (s, 3H), 1.26-1.21 (m, 3H).
7.8	 <p>2-methyl-7-((2-methylindolin-1-yl)methyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile</p>	336.8 ¹ H NMR (300 MHz, CD ₃ OD) δ 7.05-7.03 (m, 1H), 6.98-6.93 (m, 1H), 6.65-6.60 (m, 1H), 6.38 (s, 1H), 6.28-6.26 (m, 1H), 4.26-4.09 (m, 2H), 3.83-3.73 (m, 1H), 2.24-2.16 (m, 1H), 2.69-2.66 (m, 1H), 2.64 (s, 3H), 1.34-1.21 (m, 3H).


No.	Structure/Name	LCMS (M + H) ¹ H NMR
7.9	 <p>7-(((3,5-difluorophenyl)(ethyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile</p>	361.1 ¹ H NMR (300 MHz, CDCl ₃) δ 6.19-6.05 (m, 4H), 4.31 (s, 2H), 3.48-3.45 (m, 2H), 2.66 (s, 3H), 1.26-1.23 (m, 3H)

Method 8

Example 8.1: 6-[(N-ethyl-4-fluoro-anilino)methyl]-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one




Step 1: 2-bromocyclopentan-1-one




To a solution of cyclohexanone (5.00 g, 50.9 mmol) in dimethyl sulfoxide (30 mL) was added N-bromosuccinimide (11.1 g, 62.4 mmol). The reaction mixture was stirred 20 min at room temperature and then quenched with water (300 mL). The reaction mixture was extracted with dichloromethane (100 mL \times 2), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo to afford 2-bromocyclopentan-1-one as a light yellow oil (5.6 g, 58%). The crude product was used in next step without further purification. No LCMS signal.

Step 2: 4H,5H,6H-cyclopenta[d][1,3]thiazol-2-amine



To a solution of 2-bromocyclopentan-1-one (5.50 g, 33.7 mmol) in ethanol (50 mL) was added thiourea (3.30 g, 43.4 mmol) and sodium bicarbonate (4.80 g, 57.1 mmol). The reaction mixture was stirred at reflux overnight. After cooling to room temperature, the reaction was quenched with water (200 mL), extracted with dichloromethane (100 mL \times 3), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo to afford 4H,5H,6H-cyclopenta[d][1,3]thiazol-2-amine as a brown solid (2.0 g, 42%). LCMS (ESI): M+H⁺=141.1.


Step 3: 10-(chloromethyl)-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-trien-12-one



A mixture of 4H,5H,6H-cyclopenta[d][1,3]thiazol-2-amine (2.00 g, 14.3 mmol) and ethyl 4-chloro-3-oxobutanoate (3.50 g, 21.3 mmol) in polyphosphoric acid (15 mL) was stirred for 1 h at 110° C. The reaction mixture cooled to room temperature, diluted with water (30 mL) and stirred for 1 h at 80° C. After cooling to room temperature, the reaction was quenched by water (200 mL), and the pH value of the solution was adjusted to pH 8-9 with potassium carbonate, extracted with dichloromethane (100 mL \times 3), washed with brine, dried over sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with 20% ethyl acetate in petroleum ether to afford 10-(chloromethyl)-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-trien-12-one as a brown solid (150 mg, 4.0%). LCMS (ESI): M+H⁺=241.1.

157

Step 4: 6-[(N-ethyl-4-fluoro-anilino)methyl]-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one




To a solution of 10-(chloromethyl)-7-thia-1,9-diazatricyclo[6.4.0.0']_{2,6}]dodeca-2(6),8,10-trien-12-one (150 mg, 0.62 mmol) in acetonitrile (20 mL) was added N-ethyl-4-fluoroaniline (105 mg, 0.75 mmol), potassium carbonate (150 mg, 0.62 mmol), and potassium iodide (52 mg, 0.31 mmol). The reaction mixture was stirred overnight at 60° C. After cooling to room temperature, the reaction was quenched with water (100 mL), extracted with dichloromethane (50 mL×3), washed with brine, dried over sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with 33% ethyl acetate in petroleum ether to afford 6-[(N-ethyl-4-fluoro-anilino)methyl]-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one as a brown solid (47.8 mg, 22%). LCMS (ESI): M+H⁺=343.7; ¹H NMR (300 MHz, DMSO-d₆) δ 7.00-6.94 (m, 2H), 6.61-6.58 (m, 2H), 5.87 (s, 1H), 4.32 (s, 2H), 3.50-3.43 (m, 2H), 3.19-3.13 (m, 2H), 2.85-2.72 (m, 2H), 3.37-2.30 (m, 2H), 1.16-1.09 (m, 3H).

The following examples were prepared in a manner similar to Example 8.1:


No.	Structure/Name	LCMS (M + H) ¹ H NMR
8.2	<p>2-[(N-ethyl-4-fluoro-anilino)methyl]-6,7,8,9-tetrahydropyrimido[2,1-b][1,3]benzothiazol-4-one</p>	358.1 ¹ H NMR (300 MHz, DMSO-d ₆) δ 7.19-6.96 (m, 2H), 6.61-6.57 (m, 2H), 5.80 (s, 1H), 4.29 (s, 2H), 3.46-3.42 (m, 2H), 3.14 (m, 2H), 2.72-2.70 (m, 2H), 1.75 (m, 4H), 1.14-1.09 (m, 3H).

Example 8.3: 6-[(N-ethyl-4-fluoro-anilino)methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one




158

Step 1: ethyl 3-bromo-2-oxocyclopentanecarboxylate



To a solution of ethyl 3-bromo-2-oxocyclopentanecarboxylate (780 mg, 5.00 mmol) in chloroform (15 mL) was added bromine (800 mg, 5.01 mmol) dropwise with stirring. The resulting solution was stirred for 2 h at room temperature. The resulting mixture was concentrated in vacuo to afford ethyl 3-bromo-2-oxocyclopentanecarboxylate as light yellow oil (1.30 g). The crude product was used in the next step without further purification. No LCMS signal.


Step 2: ethyl 2-amino-5,6-dihydro-4H-cyclopenta[d]thiazole-4-carboxylate



To a solution of ethyl 3-bromo-2-oxocyclopentanecarboxylate (1.30 g, 5.53 mmol) in 1,4-dioxane (20 mL) was added thiourea (420 mg, 5.52 mmol). The resulting solution was refluxed for 12 h. After cooled down to room temperature, the reaction mixture was concentrated in vacuo. The residue was purified by chromatography with dichloromethane/methanol (20/1) to afford ethyl-2-amino-5,6-dihydro-4H-cyclopenta[d]thiazole-4-carboxylate as light yellow solid (600 mg, 51%). LCMS (ESI): M+H⁺=213; ¹H NMR (300 MHz, DMSO-d₆) δ 6.90 (br, 2H), 4.12-4.08 (m, 2H), 3.68-3.63 (m, 1H), 2.82-2.40 (m, 4H), 1.29-1.25 (m, 3H).


159

Step 3: Ethyl 6-(chloromethyl)-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidine-1-carboxylate



To a solution of ethyl-2-amino-5,6-dihydro-4H-cyclopenta[d]thiazole-4-carboxylate (1.00 g, 4.70 mmol) and 4-methylbenzenesulfonic acid (170 mg, 1.00 mmol) in toluene (30 mL) was added methyl 4-chloro-3-oxobutanoate (1.40 g, 9.40 mmol). The reaction mixture was stirred for 12 h at 125° C. with a Dean-Stark apparatus to separate water. After cooled down to room temperature, the reaction mixture was concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1:2) to afford ethyl 6-(chloromethyl)-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidine-1-carboxylate as light yellow solid (150 mg, 10%). LCMS (ESI): $M+H^+=313.0$; 1H NMR (300 MHz, $CDCl_3$) δ 6.30 (s, 1H), 4.46-4.40 (m, 1H), 4.33 (s, 2H), 4.22-4.18 (m, 2H), 3.03-2.87 (m, 3H), 2.62-2.54 (m, 1H), 1.29-1.25 (m, 3H).

Step 4: ethyl 2-((ethyl(4-fluorophenyl)amino)methyl)-4-oxo-4,6,7,8-tetrahydrocyclopenta[4,5]thiazolo[3,2-a]pyrimidine-6-carboxylate




To a solution of ethyl 10-(chloromethyl)-12-oxo-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-triene-3-carboxylate (50.0 mg, 0.16 mmol), potassium iodide (14 mg, 0.08 mmol) and potassium carbonate (45 mg, 0.33 mmol) in acetonitrile (5 mL) was added N-ethyl-4-fluoroaniline (33.0 mg, 0.24 mmol). The reaction mixture was stirred overnight at 80° C. After filtration and concentration in vacuo, the residue was purified by chromatography with ethyl acetate/petroleum ether (1/2) to afford ethyl 10-[[ethyl(4-fluorophenyl)amino]methyl]-12-oxo-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-triene-3-carboxylate as a light yellow semi-solid (17.2 mg, 26%). LCMS (ESI): $M+H^+=416.0$; 1H NMR (300 MHz, $CDCl_3$) δ 6.92-6.87 (m, 2H), 6.60-6.55 (m, 2H), 6.14 (s, 1H), 4.50-4.45 (m, 1H), 4.30 (s,

160


2H), 4.22-4.17 (m, 2H), 3.48-3.41 (m, 2H), 3.02-2.85 (m, 3H), 2.60-2.55 (m, 1H), 1.29-1.19 (m, 6H).

Step 5: 10-[[Ethyl(4-fluorophenyl)amino]methyl]-12-oxo-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-triene-3-carboxylic acid



To a solution of ethyl 10-[[ethyl(4-fluorophenyl)amino]methyl]-12-oxo-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-triene-3-carboxylate (40 mg, 0.10 mmol), tetrahydrofuran (2 mL) and water (2 mL) was added lithium hydroxide (12 mg, 0.50 mmol). The resulting solution was stirred overnight at room temperature. The pH value of the solution was adjusted to pH 2 with hydrochloric acid (1 mol/L) and extracted with dichloromethane (2x20 mL). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo to afford 10-[[ethyl(4-fluorophenyl)amino]methyl]-12-oxo-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-triene-3-carboxylic acid (30 mg, 80%) as a light yellow solid. The crude product was used in the next step without further purification.

Step 6: 6-[(N-ethyl-4-fluoro-anilino)methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one




To a solution of 10-[[ethyl(4-fluorophenyl)amino]methyl]-12-oxo-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-triene-3-carboxylic acid (110 mg, 0.28 mmol) and triethylamine (60 mg, 0.57 mmol) in tetrahydrofuran (10 mL) was added chloro(propan-2-yloxy)methanone (70 mg, 0.57 mmol) and the reaction mixture was stirred 0.5 h at room temperature. Then sodium borohydride (22 mg, 0.58 mmol) in water (0.5 mL) was added. The resulting solution was stirred for an additional 1 h at room temperature. The reaction was quenched with saturated aqueous ammonium chloride and extracted with dichloromethane (3x20 mL). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated

161

in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/1) to afford 6-[(N-ethyl-4-fluoro-anilino)methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one (51.2 mg, 48%) as a white solid. LCMS (ESI): $M+H^+=374.0$; 1H NMR (400 MHz, $CDCl_3$) δ 6.94-6.90 (m, 2H), 6.60-6.57 (m, 2H), 6.27 (s, 1H), 4.35 (s, 2H), 3.98-3.91 (m, 2H), 3.75-3.70 (m, 2H), 3.49-3.47 (m, 2H), 3.01-2.95 (m, 1H), 2.88-2.82 (m, 1H), 2.74-2.68 (m, 1H), 2.27-2.22 (m, 1H), 1.26-1.23 (m, 3H).

Examples 8.4 and 8.5: 6-[(N-ethyl-4-fluoro-anilino)methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one (Enantiomers 1 and 2)



162

The product of Example 8.3 was further purified by chiral SFC on a Chiralpak AD (2x15 cm) column eluting with 25% methanol (0.1% NH_4OH)/ CO_2 at 100 bar at a flow rate of 70 mL/min. The peaks isolated were analyzed on Chiralpak AD (50x0.46 cm) column eluting with 25% methanol (0.1% NH_4OH)/ CO_2 , at 120 bar (flow rate 5 mL/min, 220 nm). From this separation two isomers were isolated.

Example 8.4 (Peak 2; Enantiomer 2)

Retention time=1.45 min; LCMS (ESI): $M+H^+=374.0$; 1H NMR (400 MHz, $CDCl_3$) δ 6.94-6.90 (m, 2H), 6.60-6.57 (m, 2H), 6.27 (s, 1H), 4.35 (s, 2H), 3.98-3.91 (m, 2H), 3.75-3.70 (m, 2H), 3.49-3.47 (m, 2H), 3.01-2.95 (m, 1H), 2.88-2.82 (m, 1H), 2.74-2.68 (m, 1H), 2.27-2.22 (m, 1H), 1.26-1.23 (m, 3H).





Example 8.5 (Peak 1, Enantiomer 1)

Retention time=0.59 min; LCMS (ESI): $M+H^+=374.0$ 1H NMR (400 MHz, $CDCl_3$) δ 6.94-6.90 (m, 2H), 6.60-6.57 (m, 2H), 6.27 (s, 1H), 4.35 (s, 2H), 3.98-3.91 (m, 2H), 3.75-3.70 (m, 2H), 3.49-3.47 (m, 2H), 3.01-2.95 (m, 1H), 2.88-2.82 (m, 1H), 2.74-2.68 (m, 1H), 2.27-2.22 (m, 1H), 1.26-1.23 (m, 3H).





The following compounds were prepared using methods analogous to those described above:

No.	Structure/Name	LCMS (M + H) 1H NMR
8.6	<p>6-[(N-ethyl-4-fluoro-anilino)methyl]spiro[2,3-dihydrocyclopenta[3,4]thiazolo[1,4-a]pyrimidine-1,1'-cyclopentane]-8-one</p>	398.0 1H NMR (300 MHz, $CDCl_3$) δ 6.91-6.88 (m, 2H), 6.65-6.60 (m, 2H), 6.14 (s, 1H), 4.30 (s, 2H), 3.50-3.43 (m, 2H), 2.84-2.80 (m, 2H), 2.43-2.34 (m, 4H), 1.98-1.86 (m, 2H), 1.74-1.48 (m, 4H), 1.19-1.23 (m, 3H).
8.7	<p>6-[(N-ethyl-4-fluoro-anilino)methyl]-2,3-dihydrocyclopenta[3,4]thiazolo[1,4-a]pyrimidine-1,8-dione</p>	358.2 1H NMR (300 MHz, $CDCl_3$) δ 7.01-6.74 (m, 4H), 6.17 (s, 1H), 5.54 (s, 2H), 3.47-3.43 (m, 2H), 3.22-3.19 (m, 2H), 3.09-3.07 (m, 2H), 1.20-1.13 (m, 3H)


-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
8.8	 <p data-bbox="337 632 646 716">6-[(N-ethyl-4-fluoro-anilino)methyl]-1,1-dimethyl-2,3-dihydrocyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one</p>	372.0 ¹ H NMR (300 MHz, CD ₃ OD) δ 6.94-6.88 (m, 2H), 6.68-6.63 (m, 2H), 6.09 (s, 1H), 4.36 (s, 2H), 3.56-3.49 (m, 2H), 2.92-2.89 (m, 2H), 2.39-2.37 (m, 2H), 1.49 (s, 6H), 1.22-1.20 (m, 3H).
8.9	 <p data-bbox="337 989 646 1052">2-[(N-ethyl-4-fluoro-anilino)methyl]-8,9-dihydro-6H-pyrano[3,4]thiazolo[1,4-a]pyrimidin-4-one</p>	360.0 ¹ H NMR (300 MHz, CDCl ₃) δ 6.94-6.92 (m, 2H), 6.72 (m, 2H), 6.14 (s, 1H), 4.33-4.27 (m, 2H), 3.50-3.46 (m, 2H), 3.35-3.32 (m, 2H), 2.13-2.19 (m, 2H), 1.28-1.26 (m, 3H).
8.10	 <p data-bbox="337 1325 646 1388">2-[(N-ethyl-4-fluoro-anilino)methyl]-7-methyl-8,9-dihydro-6H-pyrido[3,4]thiazolo[1,4-a]pyrimidin-4-one</p>	373.0 ¹ H NMR (300 MHz, CDCl ₃) δ 6.94-6.86 (m, 2H), 6.59-6.52 (m, 2H), 6.12 (s, 1H), 4.28 (s, 2H), 4.22 (s, 2H), 3.47-3.45 (m, 2H), 2.86 (s, 4H), 2.58 (s, 3H), 1.23-1.21 (m, 3H).
8.11	 <p data-bbox="337 1703 646 1806">6-[(N-ethyl-4-fluoro-anilino)methyl]-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidine-1-carboxamide</p>	387.0 ¹ H NMR (400 MHz, CDCl ₃) δ 7.34 (s, 1H), 6.97-6.93 (m, 2H), 6.72-6.67 (m, 2H), 6.29 (s, 1H), 5.26 (s, 1H), 4.51-4.49 (m, 1H), 4.37 (s, 2H), 3.52-3.49 (m, 2H), 3.26-3.19 (m, 1H), 3.07-3.02 (m, 1H), 2.90-2.83 (m, 1H), 2.68-2.59 (m, 1H), 1.25-1.22 (m, 3H)

-continued


No.	Structure/Name	LCMS (M + H) ¹ H NMR
8.12	 <p data-bbox="326 611 659 695">6-[(N-ethyl-4-fluoro-anilino)methyl]-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidine-1-carbonitrile</p>	369.0 ¹ H NMR (300 MHz, CDCl ₃) δ 6.94-6.87 (m, 2H), 6.60-6.56 (m, 2H), 6.24 (s, 1H), 4.71-4.68 (m, 1H), 4.31 (s, 2H), 3.49-3.46 (m, 2H), 3.21-2.87 (m, 4H), 1.24-1.21 (m, 3H).
8.13	 <p data-bbox="326 1031 659 1115">2-[6-[(N-ethyl-4-fluoro-anilino)methyl]-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-1-yl]acetonitrile</p>	383.3 ¹ H NMR (300 MHz, CD ₃ OD) δ 6.93-6.90 (m, 2H), 6.70-6.65 (m, 2H), 6.14 (s, 1H), 4.40 (s, 2H), 4.02-3.90 (m, 1H), 3.57-3.50 (m, 2H), 3.17-2.85 (m, 5H), 2.48-2.41 (m, 1H), 1.25-1.22 (m, 3H).
8.14	 <p data-bbox="326 1423 659 1507">6-[(N-ethyl-4-fluoro-anilino)methyl]-1-(2-hydroxyethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one</p>	388.2 ¹ H NMR (300 MHz, CDCl ₃) δ 6.96-6.92 (m, 2H), 6.82-6.72 (m, 2H), 6.24 (s, 1H), 4.34 (s, 2H), 3.92-3.90 (m, 1H), 3.66-3.55 (m, 2H), 3.52-3.47 (m, 2H), 2.98-2.90 (m, 1H), 2.84-2.66 (m, 2H), 2.28-2.22 (m, 1H), 1.99-1.89 (m, 2H), 1.28-1.22 (m, 3H).
8.15	 <p data-bbox="326 1856 659 1940">2-[(ethyl(4-fluorophenyl)amino)methyl]-6-(methoxymethyl)-7,8-dihydrocyclopenta[4,5]thiazolo[3,2-a]pyrimidin-4(6H)-one</p>	388.0 ¹ H NMR (300 MHz, CDCl ₃) δ 6.94-6.90 (m, 2H), 6.61-6.58 (m, 2H), 6.18 (s, 1H), 4.33 (s, 2H), 3.91-3.88 (m, 1H), 3.70-3.64 (m, 2H), 3.51-3.48 (m, 2H), 3.32 (s, 3H), 2.99-2.48 (m, 4H), 1.26-1.23 (m, 3H).

-continued


No.	Structure/Name	LCMS (M + H) ¹ H NMR
8.16	 <p>2-[6-[(N-ethyl-4-fluoro-anilino)methyl]-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-1-yl]acetamide</p>	401.2 ¹ H NMR (300 MHz, CD ₃ OD) δ 6.23-6.19 (m, 2H), 6.69-6.65 (m, 2H), 6.12 (s, 1H), 4.38 (s, 2H), 4.08-3.97 (m, 1H), 3.57-3.50 (m, 2H), 2.96-2.73 (m, 4H), 2.48-2.36 (m, 2H), 1.25-1.21 (m, 3H).

Method 9

Example 9.1: 3-cyclohexyl-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



Step 1: 4-(cyclohex-1-en-1-yl)-5-methyl-1,3-thiazol-2-amine




To a solution of 4-bromo-5-methyl-1,3-thiazol-2-amine (from Example 4.1, Step 5) (130 mg, 0.67 mmol) in 1,4-dioxane (2 mL) was added 2-(cyclohex-1-en-1-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (140 mg, 0.67 mmol), potassium phosphate (171 mg, 0.81 mmol) and tetrakis (triphenylphosphine)palladium (78 mg, 0.070 mmol). The reaction mixture was stirred for 2 h at 90° C. and concentrated in vacuo. The residue was purified by silica gel chromatography with 3.3% methanol in dichloromethane to

afford 4-(cyclohex-1-en-1-yl)-5-methyl-1,3-thiazol-2-amine as a yellow solid (140 mg). The crude product was used in next step without further purification. LCMS (ESI): M+H⁺=194.8.

Step 2: 4-cyclohexyl-5-methyl-1,3-thiazol-2-amine

30



35

40


To a solution of 4-(cyclohex-1-en-1-yl)-5-methyl-1,3-thiazol-2-amine (140 mg, 0.72 mmol) in MeOH (20 mL) was added Pd/C (100 mg) and hydrogen chloride (0.1 mL, 12 mol/L). The reaction mixture was stirred overnight at 40° C. under a hydrogen atmosphere (5 atm), then filtered and concentrated in vacuo to afford 4-cyclohexyl-5-methyl-1,3-thiazol-2-amine as a yellow solid (130 mg). The crude product was used in next step without further purification. LCMS (ESI): M+H⁺=197.1.

45

50

Step 3: 7-(chloromethyl)-3-cyclohexyl-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

55



To a solution of 4-cyclohexyl-5-methyl-1,3-thiazol-2-amine (130 mg, 0.66 mmol) in polyphosphoric acid (10 mL)


60

65

169

was added ethyl 4-chloro-3-oxobutanoate (164 mg, 1.00 mmol). The reaction mixture was stirred for 1 h at 100° C. and then quenched by water (100 mL). The pH value was adjusted to pH 8-9 with a sodium hydroxide solution (1 M). The resulting solution was extracted with dichloromethane (50 mL×3), washed with brine, dried over anhydrous sodium sulfate, and concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/1) to afford 7-(chloromethyl)-3-cyclohexyl-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a white solid (49 mg, 25%). LCMS (ESI): M+H⁺=297.1.

Step 4: 3-cyclohexyl-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



170

To a solution of 7-(chloromethyl)-3-cyclohexyl-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (49 mg, 0.17 mmol) in acetonitrile (20 mL) was added potassium carbonate (46 mg, 0.33 mmol), potassium iodide (14 mg, 0.08 mmol) and N-ethyl-4-fluoroaniline (28 mg, 0.20 mmol). The reaction mixture was stirred overnight at 60° C. and then quenched by water (100 mL). The reaction mixture was extracted with dichloromethane (50 mL×3), washed with brine, dried over anhydrous sodium sulfate, and concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/2) to afford 3-cyclohexyl-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a gray solid (8.8 mg, 13%). LCMS (ESI): M+H⁺=399.9; ¹H NMR (300 MHz, CDCl₃) δ 6.94-6.85 (m, 2H), 6.58-6.54 (m, 2H), 6.07 (s, 1H), 4.25 (s, 2H), 3.48-3.41 (m, 2H), 2.41 (s, 3H), 1.87-1.71 (m, 6H), 1.41-1.30 (m, 4H), 1.25-1.20 (m, 3H).


The following example was prepared in a manner similar to Example 9.1:

No.	Structure/Name	LCMS (M + H) ⁺	¹ H NMR
9.2	 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-isopropyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	360.0	¹ H NMR (300 MHz, CD ₃ OD) δ 6.91-6.87 (m, 2H), 6.69-6.19 (m, 2H), 6.05 (s, 1H), 4.33 (s, 2H), 3.55-3.51 (m, 2H), 2.44 (s, 3H), 2.10 (m, 1H), 1.41-1.38 (m, 6H), 1.19-1.21 (m, 3H).

The following compounds were prepared using methods analogous to those described above:

No.	Structure/Name	LCMS (M + H) ⁺	¹ H NMR
9.3	 3-cyclopentyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	386.0	¹ H NMR (300 MHz, CD ₃ OD) δ 6.93-6.87 (m, 2H), 6.67-6.63 (m, 2H), 6.04 (s, 1H), 4.33 (s, 2H), 4.17-4.08 (m, 1H), 3.53-3.47 (m, 2H), 2.41 (s, 3H), 1.96-1.88 (m, 6H), 1.69-1.65 (m, 2H), 1.22-1.19 (m, 3H).


-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
9.4	 <p>7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-tetrahydropyran-4-yl-thiazolo[3,2-a]pyrimidin-5-one</p>	401.8 ¹ H NMR (300 MHz, CDCl ₃) δ 6.92-6.85 (m, 2H), 6.62-6.58 (m, 2H), 6.10 (s, 1H), 4.67 (br, 1H), 4.26 (s, 2H), 4.06-4.01 (m, 2H), 3.55-3.42 (m, 4H), 2.46 (s, 3H), 2.18-2.07 (m, 2H), 1.84-1.80 (m, 2H), 1.30-1.19 (m, 3H).

Method 10

Step 2: 4-Cyclobutyl-5-methyl-1,3-thiazol-2-amine

Example 10.1: 3-cyclobutyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one




25

30

35


40



To a solution of 2-bromo-1-cyclobutylpropan-1-one (3.50 g, 31.3 mmol) in ethanol (30 mL) was added thiourea (1.5 g, 19.71 mmol). The reaction solution was heated to reflux for 1 h and then concentrated in vacuo. The residue was dissolved in dichloromethane (50 mL) and the solids were filtered off. The resulting solution was concentrated in vacuo to afford 4-cyclobutyl-5-methyl-1,3-thiazol-2-amine as a yellow solid (600 mg, 18%). LCMS (ESI): M+H⁺=169.0.

Step 3: 6-(Chloromethyl)-3-cyclobutyl-2-methyl-3aH,4H-thieno[2,3-b]pyridine-4-one


Step 1: 2-Bromo-1-cyclobutylpropan-1-one



45

50

55



To a solution of 1-cyclobutylpropan-1-one (2.50 g, 22.3 mmol) in methanol (50 mL) was added bromine (3.87 g, 24.2 mmol) under nitrogen atmosphere. The reaction solution was stirred overnight at room temperature and was then concentrated in vacuo to afford 2-bromo-1-cyclobutylpropan-1-one as yellow oil (3.5 g). The crude product was used for the next step without further purification.


60

65

To a solution of 4-cyclobutyl-5-methyl-1,3-thiazol-2-amine (335 mg, 1.99 mmol) was added polyphosphoric acid (5 mL) and ethyl 4-chloro-3-oxobutanoate (492 mg, 2.99 mmol). The reaction solution was stirred for 1 h at 110° C. The pH value of the solution was adjusted to pH 9-10 with an aqueous sodium hydroxide solution (2 M). The reaction mixture was extracted with dichloromethane (5×100 mL), washed with brine, dried over sodium sulfate and concentrated in vacuo to afford 6-(chloromethyl)-3-cyclobutyl-2-methyl-3aH,4H-thieno[2,3-b]pyridine-4-one as a brown solid (300 mg). The crude product was used in next step without further purification. LCMS (ESI): M+H⁺=269.0.

173

Step 4: 3-Cyclobutyl-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



To a solution of 7-(chloromethyl)-3-cyclobutyl-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (300 mg, 1.12

174


mmol) in acetonitrile (20 mL) was added potassium iodide (93 mg, 0.56 mmol), potassium carbonate (309 mg, 2.24 mmol), and N-ethyl-4-fluoroaniline (311 mg, 2.23 mmol). The resulting solution was stirred for 5 h at 70° C. After concentrating in vacuo, the crude product was purified by Prep-HPLC with the following conditions (Agilent 1200: Column, X-Brigde C18; mobile phase, 0.05% NH₄HCO₃ in water and CH₃CN (CH₃CN 20% up to 60% in 15 min); Detector, UV254) to afford 3-cyclobutyl-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a yellow solid (36.1 mg, 8%). LCMS (ESI): M+H⁺=371.8; ¹H NMR (300 MHz, CD₃OD) δ 6.93-6.86 (m, 2H), 6.69-6.62 (m, 2H), 6.01 (s, 1H), 4.44-4.35 (m, 1H), 4.31 (s, 2H), 3.53-3.49 (m, 2H), 2.48 (s, 3H), 2.45-2.35 (m, 2H), 1.89-1.80 (m, 1H), 2.06-1.94 (m, 1H), 1.22-1.19 (m, 3H).

The following examples were prepared in a manner similar to Example 10.1:

No.	Structure/Name	LCMS (M+ H) ¹ H NMR
10.2	<p>3-tert-butyl-7-[(N-ethyl-4-fluoroanilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	374.0 ¹ H NMR (300 MHz, CD ₃ OD) δ 6.94-6.91 (m, 2H), 6.68-6.63 (m, 2H), 6.04 (s, 1H), 4.31 (s, 2H), 3.53-3.50 (m, 2H), 2.54 (s, 3H), 1.59 (s, 9H), 1.23-1.20 (m, 3H).
10.3	<p>3-acetyl-7-[(ethyl(4-fluorophenyl)amino)methyl]-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	360.1 ¹ H NMR (300 MHz, CD ₃ OD) δ 6.95-6.89 (m, 2H), 6.70-6.65 (m, 2H), 6.16 (s, 1H), 4.40 (s, 2H), 3.54-3.52 (m, 2H), 2.40 (m, 6H), 1.24-1.21 (m, 3H).
10.4	<p>7-[(N-ethyl-4-fluoroanilino)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	375.2 ¹ H NMR (300 MHz, DMSO-d ₆) δ 8.37-8.36 (m, 1H), 7.14-6.95 (m, 2H), 6.68-6.59 (m, 2H), 5.90 (s, 1H), 4.42 (s, 2H), 3.48-3.35 (m, 2H), 2.73-2.71 (m, 3H), 2.30 (s, 3H), 1.13-1.11 (m, 3H).


175

Example 10.5: 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-hydroxy-1-methyl-ethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one




To a solution of 3-acetyl-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 10.3) (70 mg, 0.19 mmol) in tetrahydrofuran (15 mL) was added methylmagnesium bromide in tetrahydrofuran (1 mol/L, 0.42 mL). The reaction was stirred for 48 h at room temperature and was then quenched by a saturated aqueous ammonium chloride solution (20 mL). The resulting mixture was extracted with dichloromethane (3x30 mL), washed with brine, dried over anhydrous sodium sulfate, and concentrated in vacuo. The residue was purified chromatography with dichloromethane/methanol (50/1) to afford 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(2-hydroxypropan-2-yl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a yellow solid (40 mg, 52%). LCMS (ESI): $M+H^+=376.1$; 1H NMR (300 MHz, CD_3OD) δ 6.95-6.89 (m, 2H), 6.70-6.65 (m, 2H), 6.24 (s, 1H), 4.39 (s, 2H), 3.56-3.50 (m, 2H), 2.58 (s, 3H), 1.73 (s, 6H), 1.24-1.21 (m, 3H).

Example 10.6: 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-hydroxyethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one




Step 1: 7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-carbaldehyde



176


To a solution of 3-bromo-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (from Example 5.1, Step 1) (200 mg, 0.51 mmol) in tetrahydrofuran (10 mL) was added n-butyl lithium (0.3 mL, 2.5 mol/L) at $-78^\circ C.$, then was stirred 30 min at the same temperature. Ethyl formate (75.8 mg, 1.02 mmol) was added to the reaction mixture at $-78^\circ C.$ and allowed to warm to room temperature for 1 hour. The resulting reaction was quenched by water (20 mL), extracted with dichloromethane (30 mLx3), washed with brine, dried over anhydrous sodium sulfate, and concentrated under vacuum. The residue was purified by chromatography with 20% ethyl acetate in petroleum ether to afford 7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-carbaldehyde. LCMS (ESI): $M+H^+=345.1$.

Step 2: 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-hydroxyethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-carbaldehyde (200 mg, 0.58 mmol) (from example 11.4) in tetrahydrofuran (15 mL) was added methylmagnesium bromide (1.3 mL, 0.5 mol/L). The resulting solution was stirred for overnight at room temperature. The reaction was then quenched by ammonium chloride (sat., 20 mL), extracted with dichloromethane (3x30 mL), washed with brine, dried over sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/2) to afford 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(1-hydroxyethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as an off-white solid (67.5 mg, 31%). LCMS (ESI): $M+H^+=362.1$; 1H NMR (300 MHz, CD_3OD) δ 6.95-6.88 (m, 2H), 6.71-6.65 (m, 2H), 6.20 (s, 1H), 5.43-5.36 (m, 1H), 4.39 (s, 2H), 3.56-3.51 (m, 2H), 2.50 (s, 3H), 1.54-1.51 (m, 3H), 1.24-1.21 (m, 3H).

Example 10.7: 3-[(dimethylamino)methyl]-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one




177

To a solution of 7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carbaldehyde (from Example 10.6, Step 1) (100 mg, 0.29 mmol) in methanol (30 mL) was added dimethylamine hydrochloride (118 mg, 1.45 mmol), triethylamine (161 mg, 1.59 mmol) and sodium cyanoborohydride (55 mg, 0.88 mmol). The reaction mixture was stirred overnight at room temperature and then quenched by water (50 mL). The reaction mixture was extracted with dichloromethane (30 mL×3), washed with brine, dried over anhydrous sodium sulfate, and concentrated under vacuum. The residue was purified by chromatography with 25% ethyl acetate in petroleum ether to afford 3-[[dimethylamino]methyl]-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a light yellow solid (11.1 mg, 10%). LCMS [M+H]⁺=374.85; ¹H NMR (300 MHz, CDCl₃) δ 6.92-6.84 (m, 2H), 6.58-6.54 (m, 2H), 6.14 (s, 1H), 4.28 (s, 2H), 4.05 (bs, 2H), 3.48-3.41 (m, 2H), 2.44-2.39 (m, 9H), 1.26-1.19 (m, 3H).

The following examples were prepared in a manner similar to Example 10.7:

178

Example 10.11: 7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(2,2,2-trifluoro-1-hydroxyethyl)thiazolo[3,2-a]pyrimidin-5-one




To a solution of tetrabutylammonium fluoride (113 mg, 0.43 mmol) in tetrahydrofuran (10 mL) was added 4 Å-molecular sieves (200 mg) and then was stirred for 0.5 h at -20°C. A solution of 7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carbal-

No.	Structure/Name	LCMS (M + H)	¹ H NMR
10.8	<p>3-(azetidin-1-ylmethyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	387.1	¹ H NMR (300 MHz, CD ₃ OD) δ 6.93-6.89 (m, 2H), 6.69-6.63 (m, 2H), 6.08 (s, 1H), 4.36 (s, 2H), 4.27 (s, 2H), 3.56-3.52 (m, 2H), 3.33-3.31 (m, 4H), 2.41 (s, 3H), 2.13-2.06 (m, 2H), 1.23-1.21 (m, 3H)
10.9	<p>7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(pyrrolidin-1-ylmethyl)thiazolo[3,2-a]pyrimidin-5-one</p>	401.1	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.01-6.95 (m, 2H), 6.65-6.59 (m, 2H), 5.84 (s, 1H), 4.31 (s, 2H), 4.13 (s, 2H), 3.48-3.43 (m, 2H), 2.50-2.49 (m, 4H), 2.38 (s, 3H), 1.75-1.73 (m, 4H), 1.15-1.10 (m, 3H)
10.10	<p>7-[(N-ethyl-4-fluoro-anilino)methyl]-3-[(3-hydroxyazetidin-1-yl)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	402.9	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.01-6.95 (m, 2H), 6.63-6.59 (m, 2H), 5.83 (s, 1H), 5.22-5.20 (m, 1H), 4.31 (s, 2H), 4.11-4.01 (m, 3H), 3.50-3.37 (m, 4H), 2.83-2.78 (m, 2H), 2.38 (s, 3H), 1.15-1.10 (m, 3H)

179


aldehyde (from Example 10.6, Step 1) (300 mg, 0.87 mmol), trimethyl(trifluoromethyl)silane (617 mg, 4.35 mmol) and 4 Å molecular sieve (100 mg) in tetrahydrofuran (20 mL) was added. The reaction mixture was stirred for an additional 3 h at -30°C . The reaction was quenched with water (50 mL), extracted with dichloromethane (3×50 mL), washed with brine, dried over anhydrous magnesium sulfate and concentrated in vacuo. The residue was purified by silica gel chromatography with dichloromethane/methanol (50/1) to afford 7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-3-(2,2,2-trifluoro-1-hydroxyethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a off-white solid (32.4 mg, 9.0%). LCMS (ESI): $M+H^+=416.0$; $^1\text{H NMR}$ (300 MHz, CD_3OD) δ 6.95-6.89 (m, 2H), 6.69-6.64 (m, 2H), 6.12 (s, 1H), 4.87 (s, 1H), 4.37 (s, 2H), 3.55-3.50 (m, 2H), 2.63 (s, 3H), 1.24-1.20 (m, 3H).

Example 10.12: 7-((ethyl(4-fluorophenyl)amino)methyl)-3-(hydroxymethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-carbaldehyde (from Example 10.6, Step 1) (100 mg, 0.29 mmol) in tetrahydrofuran (10 mL) was added water (2 mL) and sodium borohydride (33 mg, 0.87 mmol). The reaction mixture was stirred overnight at room temperature and then quenched with water (50 mL). The resulting solution was extracted with ethyl acetate ($30 \text{ mL} \times 3$), washed with brine, dried over anhydrous sodium sulfate, and concentrated in vacuo. The residue was purified by chromatography with 50% ethyl acetate in petroleum ether to afford 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(hydroxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a off-white solid (60 mg, 60%). LCMS (ESI): $M+H^+=347.9$; $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 6.93-6.86 (m, 2H), 6.59-6.54 (m, 2H), 6.26 (s, 1H), 4.75-4.73 (m, 2H), 4.49-4.45 (m, 1H), 4.32 (s, 2H), 3.52-3.44 (m, 2H), 2.44 (s, 3H), 1.25-1.19 (m, 3H).


Example 10.13: 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(methoxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



180

To a solution of 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(hydroxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (60 mg, 0.17 mmol) in tetrahydrofuran (10 mL) was added sodium hydride (11 mg, 0.28 mmol) and iodomethane (30 mg, 0.21 mmol). The reaction mixture was stirred overnight at room temperature and then quenched by water (100 mL). The resulting solution was extracted with dichloromethane ($50 \text{ mL} \times 3$), washed with brine, dried over anhydrous sodium sulfate, and concentrated in vacuo. The residue was purified by chromatography with 50% ethyl acetate in petroleum ether to afford 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(methoxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a yellow semi-solid (1.4 mg, 2.0%). LCMS (ESI): $M+H^+=362.1$; $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 6.93-6.87 (m, 2H), 6.61-6.60 (m, 2H), 6.16 (s, 1H), 4.93 (s, 2H), 4.28 (s, 2H), 3.45-3.42 (m, 5H), 2.45 (s, 3H), 1.33-1.21 (m, 3H).


Example 10.14: 7-[[Ethyl(4-fluorophenyl)amino]methyl]-2-methyl-3-(1H-pyrazol-1-ylmethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



To a solution of 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(hydroxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 10.12) (100 mg, 0.29 mmol), 1H-pyrazole (39.2 mg, 0.58 mmol) and triphenylphosphine (135 mg, 0.52 mmol) in tetrahydrofuran (5 mL) was added diisopropyl azodicarboxylate (105 mg, 0.52 mmol). The reaction mixture was stirred at room temperature overnight. After concentrating in vacuo, the crude residue was purified by silica gel chromatography with dichloromethane/methanol (100/1) to afford 7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-3-(1H-pyrazol-1-ylmethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as an off-white solid (10.0 mg, 8.0%). LCMS (ESI): $M+H^+=398.1$; $^1\text{H NMR}$ (300 MHz, CD_3OD) δ 7.76 (s, 1H), 7.45 (s, 1H), 6.93-6.87 (m, 2H), 6.68-6.62 (m, 2H), 6.25 (s, 1H), 6.05 (s, 1H), 5.90 (s, 2H), 4.34 (s, 2H), 3.53-3.47 (m, 2H), 2.58 (s, 3H), 1.22-1.19 (m, 3H).

181


Example 10.15: 7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(methylsulfonylmethyl)thiazolo[3,2-a]pyrimidin-5-one



182

Step 2: 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(methanesulfonylmethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

5




10

15

Step 1: (7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)methyl methanesulfonate

20



25

30

35

To a solution of 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(hydroxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 10.12) (100 mg, 0.28 mmol) in dichloromethane (15 mL) was added triethylamine (58 mg, 0.58 mmol) and methanesulfonyl chloride (50 mg, 0.44 mmol) at 0° C. The reaction mixture was stirred for 30 min at 0° C. and then concentrated under vacuum. The crude product was used directly in the next step.


40

To a solution of (7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)methyl methanesulfonate (120 mg, 0.28 mmol) in ethanol (30 mL) was added sodium methanesulfinate (202 mg, 1.98 mmol). The reaction mixture was stirred at reflux for 2 h at 80° C. and then concentrated in vacuo. The residue was purified by silica gel chromatography with dichloromethane/methanol (30/1) to afford 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(methanesulfonylmethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as an off-white solid (2.8 mg, 2%). LCMS (ESI): $M+H^+=409.9$; 1H NMR (300 MHz, DMSO- d_6) δ 6.94-6.88 (m, 2H), 6.57-6.52 (m, 2H), 5.84 (s, 1H), 5.28 (s, 2H), 4.27 (s, 2H), 3.41-3.36 (m, 2H), 2.93 (s, 3H), 2.39 (s, 3H), 1.08-1.03 (m, 3H) and 3-(ethoxymethyl)-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a off-white solid (22.1 mg, 21%). LCMS (ESI): $M+H^+=409.9$; 1H NMR (300 MHz, DMSO- d_6) δ 6.94-6.88 (m, 2H), 6.57-6.52 (m, 2H), 5.84 (s, 1H), 5.28 (s, 2H), 4.27 (s, 2H), 3.41-3.36 (m, 2H), 2.93 (s, 3H), 2.39 (s, 3H), 1.08-1.03 (m, 3H).

The following examples were prepared in a manner similar to Example 10.15:




No.	Structure/Name	LCMS (M + H) 1H NMR
10.16	<p>3-(ethoxymethyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one</p>	376.1 1H NMR (300 MHz, DMSO- d_6) δ 7.00-6.94 (m, 2H), 6.64-6.60 (m, 2H), 5.86 (s, 1H), 4.87 (s, 2H), 4.32 (s, 2H), 3.49-3.47 (m, 4H), 2.41 (s, 3H), 1.15-1.05 (m, 6H).

-continued





No.	Structure/Name	LCMS (M + H) ¹ H NMR
10.17	 <p>2-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]acetonitrile</p>	357.1 ¹ H NMR (300 MHz, CDCl ₃) δ 6.94-6.87 (m, 2H), 6.59-6.53 (m, 2H), 6.18 (s, 1H), 4.41 (s, 2H), 4.28 (s, 2H), 3.48-3.41 (m, 2H), 2.41 (s, 3H), 1.25-1.19 (m, 3H).

20


The following compounds were prepared using methods analogous to those described above:

No.	Structure/Name	LCMS (M + H) ¹ H NMR
10.18	 <p>3-tert-butyl-7-[(N-ethyl-4-fluoro-anilino)methyl]thiazolo[3,2-a]pyrimidin-5-one</p>	360.0 ¹ H NMR (300 MHz, CDCl ₃) δ 6.92-6.89 (m, 2H), 6.64 (s, 1H), 6.61-6.56 (m, 2H), 6.17 (s, 1H), 4.29 (s, 2H), 3.48-3.45 (m, 2H), 1.55 (s, 9H), 1.24-1.21 (m, 3H).
10.19	 <p>7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-hydroxy-1-methyl-ethyl)thiazolo[3,2-a]pyrimidin-5-one</p>	362.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 7.46-7.36 (m, 1H), 7.03-6.93 (m, 2H), 6.69-6.58 (m, 2H), 6.48 (s, 1H), 6.11 (s, 1H), 4.40 (s, 2H), 3.49 (q, J = 7.0 Hz, 2H), 1.56 (s, 6H), 1.14 (t, J = 7.0 Hz, 3H).
10.20	 <p>7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-hydroxyethyl)thiazolo[3,2-a]pyrimidin-5-one</p>	348.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 7.03-6.92 (m, 2H), 6.67-6.59 (m, 2H), 5.92 (s, 1H), 5.48 (td, J = 6.6, 5.4 Hz, 1H), 7.31-7.25 (m, 1H), 4.34 (s, 2H), 3.47 (q, J = 7.0 Hz, 2H), 1.37 (d, J = 6.3 Hz, 3H), 1.13 (t, J = 7.0 Hz, 3H).

-continued


No.	Structure/Name	LCMS (M + H) ¹ H NMR
10.21	 <p data-bbox="370 709 683 772">7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(6-oxa-1-azaspiro[3.3]heptan-1-ylmethyl)thiazolo[3,2-a]pyrimidin-5-one</p>	429.0 ¹ H NMR (300 MHz, CD ₃ OD) δ 6.96-6.88 (m, 2H), 6.70-6.64 (m, 2H), 6.08 (s, 1H), 5.14-5.10 (m, 2H), 4.66-4.43 (m, 2H), 4.52 (s, 2H), 4.35 (s, 2H), 3.56-3.51 (m, 2H), 3.19-3.16 (m, 2H), 2.46 (s, 3H), 2.34-2.31 (m, 2H), 1.24-1.21 (m, 3H).
10.22	 <p data-bbox="370 1062 683 1125">7-[(N-ethyl-4-fluoro-anilino)methyl]-3-[(2-hydroxyethylamino)methyl]-2-methylthiazolo[3,2-a]pyrimidin-5-one</p>	391.1 ¹ H NMR (300 MHz, CD ₃ OD) δ 6.89-6.83 (m, 2H), 6.64-6.59 (m, 2H), 6.08 (s, 1H), 4.32 (s, 2H), 4.11 (s, 2H), 3.62-3.60 (m, 2H), 3.50-3.45 (m, 2H), 2.69-2.66 (m, 2H), 2.41 (s, 3H), 1.19-1.16 (m, 3H).
10.23	 <p data-bbox="358 1398 699 1461">3-(ethyl((3-(hydroxymethyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)amino)benzonitrile</p>	354.9 ¹ H NMR (300 MHz, DMSO-d ₆) δ 7.34-7.28 (m, 1H), 7.01-6.92 (m, 3H), 5.92 (s, 1H), 4.76 (s, 2H), 4.44 (s, 2H), 3.57-3.50 (m, 2H), 2.41 (s, 3H), 1.16-1.12 (m, 3H).
10.24	 <p data-bbox="370 1766 683 1829">3-[ethyl-[[3-(methoxymethyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl)methyl]amino]benzonitrile</p>	369.0 ¹ H NMR (300 MHz, DMSO-d ₆) δ 7.34-7.28 (m, 1H), 7.02-6.93 (m, 3H), 5.85 (s, 1H), 4.83 (s, 2H), 4.42 (s, 2H), 3.56-3.49 (m, 2H), 3.23 (s, 3H), 2.42 (s, 3H), 1.16-1.11 (m, 3H).

-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
10.25	 <p>7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-((methylthio)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	378.0 ¹ H NMR (300 MHz, DMSO-d ₆) δ 7.01-6.95 (m, 2H), 6.63-6.57 (m, 2H), 5.85 (s, 1H), 4.32 (s, 2H), 4.22 (s, 2H), 3.50-3.43 (m, 2H), 2.37 (s, 3H), 1.98 (s, 3H), 1.14-1.10 (m, 3H).

Method 11



Example 11.1: 3-chloro-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one






To a solution of 3-bromo-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-

20 5-one (from Example 5.1, Step 1) (100 mg, 0.25 mmol) in tetrahydrofuran (20 mL) was added n-butyl lithium (0.12 mL, 2.5 mol/L) dropwise at -80°C . The reaction solution was stirred for 30 min at -80°C . To the reaction was added N-chlorosuccinimide (40 mg, 0.30 mmol) at -80°C . The reaction was slowly warmed to room temperature for 30 min. The reaction was then quenched by the addition of 100 mL of water, extracted with ethyl acetate (30 mL \times 3), washed with brine, dried over anhydrous sodium sulfate, and concentrated in vacuo. The residue was purified by chromatography with 33% ethyl acetate in petroleum ether to afford 3-chloro-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a yellow solid (16.4 mg, 18%). LCMS (ESI): M+H⁺=351.9; ¹H NMR (300 MHz, CDCl₃) δ 6.94-6.88 (m, 2H), 6.61-6.57 (m, 2H), 6.16 (s, 1H), 4.27 (s, 2H), 4.50-4.43 (m, 2H), 2.36 (s, 3H), 1.26-1.20 (m, 3H).

The following examples were prepared in a manner similar to Example 11.1:


No.	Structure/Name	LCMS (M + H) ¹ H NMR
11.2	 <p>7-[(N-ethyl-4-fluoro-anilino)methyl]-3-fluoro-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	336.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.63-7.00 (m, 2H), 6.88-6.70 (m, 2H), 6.46 (s, 1H), 4.20 (s, 2H), 3.30-3.21 (m, 2H), 2.40 (s, 3H), 1.14-1.11 (m, 3H).
11.3	 <p>7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	318.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 7.84 (d, J = 1.8 Hz, 1H), 7.07-6.90 (m, 2H), 6.66-6.50 (m, 2H), 5.93 (s, 1H), 4.35 (s, 2H), 3.47 (q, J = 7.0 Hz, 2H), 2.42 (d, J = 1.4 Hz, 3H), 1.13 (t, J = 7.0 Hz, 3H).

The following compounds were prepared using methods analogous to those described above:

No.	Structure/Name	LCMS (M + H) ¹ H NMR
11.4	 <p>7-[(N-ethyl-4-fluoro-anilino)methyl]-3-iodo-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	443.6 ¹ H NMR (300 MHz, CDCl ₃) δ 6.95-6.90 (m, 2H), 6.71-6.68 (m, 2H), 6.19 (s, 1H), 4.29 (s, 2H), 4.52-4.45 (m, 2H), 2.39 (s, 3H), 1.25-1.21 (m, 3H).
11.5	 <p>3-chloro-7-[(N-ethyl-4-fluoro-anilino)methyl]thiazolo[3,2-a]pyrimidin-5-one</p>	338.1 ¹ H NMR (300 MHz, CDCl ₃) δ 7.57 (s, 1H), 7.55-7.52 (m, 2H), 7.44-7.40 (m, 1H), 4.06-4.00 (m, 3H), 3.22-3.17 (m, 1H), 2.37 (s, 3H), 2.31-2.26 (m, 1H), 1.34-1.25 (m, 1H), 1.06-0.98 (m, 2H).
11.6	 <p>7-[(N-ethyl-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	300.0 ¹ H NMR (400 MHz, DMSO-d ₆) δ 7.84 (s, 1H), 7.14 (t, J = 7.8 Hz, 2H), 6.62 (t, J = 7.5 Hz, 2H), 5.92 (s, 1H), 4.38 (s, 2H), 3.50 (q, J = 7.0 Hz, 2H), 2.42 (s, 3H), 1.15 (t, J = 7.0 Hz, 3H).

Method 12

Example 12.1: 3-(1,3-dihydroxypropyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one




Step 1: 3-[3-(Benzyloxy)-1-hydroxypropyl]-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

45

50

55




To a solution of 3-bromo-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 5.1, Step 1) (600 mg, 1.51 mmol) in tetrahydrofuran (50 mL) was added n-butyl lithium in tetrahydrofuran (2.4 M, 3 mL, 7.2 mmol) at -80° C. After stirring at -80° C. for 0.5 h, 3-(benzyloxy)propanal (500 mg, 3.05 mmol) was added to the reaction. The resulting solution was stirred for 1.5 h at -80° C. The reaction was then quenched by water (50 mL), extracted with dichloromethane

191


(3×50 mL), washed with brine, dried over sodium sulfate and concentrated in vacuo to afford 3-[3-(benzyloxy)-1-hydroxypropyl]-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a yellow solid (500 mg). The crude product was used in next step without purification. LCMS (ESI): $M+H^+=482.0$.

Step 2: 3-(1,3-Dihydroxypropyl)-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one




To a solution of 3-[3-(benzyloxy)-1-hydroxypropyl]-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (500 mg, 1.04 mmol) in dichloromethane (50 mL) was added a solution of boron trichloride (10 mL, 1 mol/L) in dichloromethane (10 mL) at -20°C . The reaction mixture was stirred overnight at room temperature. The reaction was quenched by a saturated aqueous ammonium chloride solution (50 mL), extracted with dichloromethane (3×50 mL), washed with brine, dried over sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/1) to afford 3-(1,3-dihydroxypropyl)-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a off-white solid (136.4 mg, 32%). LCMS (ESI): $M+H^+=392.1$; $^1\text{H NMR}$ (300 MHz, CD_3OD) δ 6.95-6.88 (m, 2H), 6.70-6.65 (m, 2H), 6.22 (s, 1H), 5.32-5.27 (m, 1H), 4.40 (s, 2H), 3.75-3.67 (m, 1H), 3.57-3.49 (m, 3H), 2.51 (s, 3H), 2.22-2.10 (m, 1H), 2.00-1.94 (m, 1H), 1.24-1.22 (m, 3H).

Example 12.2: 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-fluoro-3-hydroxy-propyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one




192

Step 1: 3-(3-(tert-butyl dimethylsilyloxy)-1-fluoropropyl)-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 3-[3-[(tert-butyl dimethylsilyl)oxy]-1-hydroxypropyl]-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (prepared via a similar method as Example 12.1, Step 1) (120 mg, 0.24 mmol) in dichloromethane (20 mL) was added diethylaminosulfur trifluoride (57.4 mg, 0.36 mmol) dropwise at -78°C . The resulting solution was stirred overnight at room temperature. The reaction was then quenched by a saturated aqueous sodium bicarbonate solution (20 mL), extracted with dichloromethane (3×20 mL), washed with brine, dried over sodium sulfate and concentrated in vacuo to afford 3-[3-[(tert-butyl dimethylsilyl)oxy]-1-fluoropropyl]-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a yellow oil (100 mg). LCMS (ESI): $M+H^+=508.1$.

Step 2: 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(1-fluoro-3-hydroxypropyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one




To a solution of 3-[3-[(tert-butyl dimethylsilyl)oxy]-1-fluoropropyl]-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (50 mg, 0.10 mmol) in tetrahydrofuran (15 mL) was added hydrogen chloride (1 M, 2.5 mL) dropwise with stirring. The resulting solution was stirred for 3 h at room temperature. The reaction was then quenched by a saturated aqueous sodium bicarbonate solution (20 mL), extracted with dichloromethane (3×30 mL), washed with brine, dried over sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with dichloromethane/methanol (30/1) to afford 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(1-fluoro-3-hydroxypropyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a light yellow solid (12.2 mg, 30%). LCMS (ESI): $M+H^+=394.1$; $^1\text{H NMR}$ (300 MHz, CD_3OD) δ 6.94-6.76 (m, 3H), 6.68-6.64 (m, 2H), 6.07 (s, 1H), 4.35

193

(s, 2H), 3.79-3.71 (m, 2H), 3.55-3.48 (m, 2H), 2.52 (s, 3H), 2.40-2.52 (m, 2H), 1.23-1.20 (m, 3H).

Examples 12.3 and 12.4: 3-(1,3-dihydroxypropyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (Enantiomers 1 and 2)



The product of Example 12.1 was further purified by chiral SFC on a Chiralpak AD (2x15 cm) column eluting with 25% methanol (0.1% NH₄OH)/CO₂ at 100 bar at a flow rate of 70 mL/min. The peaks isolated were analyzed on Chiralpak AD (50x0.46 cm) column eluting with 25% methanol (0.1% NH₄OH)/CO₂, at 120 bar (flow rate 5 mL/min, 220 nm). From this separation two isomers were isolated.

Example 12.3 (Peak Enantiomer 1)

Retention time=1.54 min; LCMS (ESI): M+H⁺=392.1; ¹H NMR (300 MHz, CD₃OD) δ 6.95-6.88 (m, 2H), 6.70-6.65 (m, 2H), 6.22 (s, 1H), 5.32-5.27 (m, 1H), 4.40 (s, 2H), 3.75-3.67 (m, 1H), 3.57-3.49 (m, 3H), 2.51 (s, 3H), 2.22-2.10 (m, 1H), 2.00-1.94 (m, 1H), 1.24-1.22 (m, 3H).

Example 12.4 (Peak 2, Enantiomer 2)


Retention time=1.63 min; LCMS (ESI): M+H⁺=392.1; ¹H NMR (300 MHz, CD₃OD) δ 6.95-6.88 (m, 2H), 6.70-6.65 (m, 2H), 6.22 (s, 1H), 5.32-5.27 (m, 1H), 4.40 (s, 2H), 3.75-3.67 (m, 1H), 3.57-3.49 (m, 3H), 2.51 (s, 3H), 2.22-2.10 (m, 1H), 2.00-1.94 (m, 1H), 1.24-1.22 (m, 3H).

The following example was prepared in a manner similar to Example 12.1:


194

Method 13

Example 13.1: 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(3-hydroxypropyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one



Step 1: (E)-Ethyl-3-(7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)acrylate




To a solution of 3-bromo-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (from Example 5.1, Step 1) (400 mg, 1.01 mmol), tri(o-tolyl)phosphine (60 mg, 0.20 mmol), triethylamine (200 mg, 1.98 mmol) and tris(dibenzylideneacetone)dipal-

No.	Structure/Name	LCMS (M + H) ¹ H NMR
12.5	<p>3-(1,2-dihydroxyethyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	378.0 ¹ H NMR (300 MHz, CDCl ₃) 6.94-6.88 (m, 2H), 6.60-6.54 (m, 2H), 6.32 (s, 1H), 6.02-5.97 (m, 1H) 4.34 (s, 2H), 4.02-3.96 (m, 1H), 3.74-3.68 (m, 1H), 3.48-3.46 (m, 2H), 2.47 (s, 3H), 1.25-1.20 (m, 3H)

195


ladium(0) (50 mg, 0.05 mmol) in acetonitrile (20 mL) was added ethyl acrylate (200 mg, 2.00 mmol). The reaction mixture was stirred for 3 h at 90° C. and then concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/2) to afford (E)-ethyl-3-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)acrylate as a light yellow solid (280 mg, 67%). LCMS (ESI): M+H⁺=416.0; ¹H NMR (300 MHz, CDCl₃) δ 8.26-8.20 (m, 1H), 6.94-6.90 (m, 2H), 6.62-6.59 (m, 2H), 6.20 (s, 1H), 6.02-5.98 (m, 1H), 4.31-4.27 (m, 4H), 3.50-3.47 (m, 2H), 2.48 (s, 3H), 1.37-1.22 (m, 6H).

Step 2: Ethyl-3-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)propanoate



To a solution of (E)-ethyl-3-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)acrylate in methanol (10 mL) was added 10% palladium on carbon and the reaction solution was stirred 12 h at room temperature under a hydrogen atmosphere (1.5 atm). After filtration the resulting solution was concentrated in vacuo to afford ethyl-3-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)propanoate as a light yellow solid (180 mg, 90%). LCMS (ESI): M+H⁺=418.0; ¹H NMR (300 MHz, CDCl₃) δ 6.94-6.88 (m, 2H), 6.62-6.58 (m, 2H), 6.12 (s, 1H), 4.28 (s, 2H), 4.13-4.10 (m, 2H), 3.51-3.41 (m, 4H), 2.74-2.71 (m, 2H), 2.37 (s, 3H), 1.26-1.20 (m, 6H).


Step 3: 7-((Ethyl(4-fluorophenyl)amino)methyl)-3-(3-hydroxypropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



196


To a solution of ethyl-3-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)propanoate (180 mg, 0.43 mmol) in methanol (10 mL) was added lithiumborohydride (20 mg, 0.91 mmol) at 0° C. The resulting solution was stirred for 5 h at room temperature. The reaction was then quenched by a saturated aqueous ammonium chloride solution (20 mL), extracted with dichloromethane (3×50 mL), washed with brine, dried over anhydrous sodium sulfate, and then concentrated in vacuo. The residue was purified by silica gel chromatography with dichloromethane/methanol (30/1) to afford 7-((ethyl(4-fluorophenyl)amino)methyl)-3-(3-hydroxypropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one as a light yellow solid (100 mg, 62%). LCMS (ESI): M+H⁺=376.0; ¹H NMR (400 MHz, CDCl₃) δ 6.94-6.90 (m, 2H), 6.63-6.59 (m, 2H), 6.15 (s, 1H), 4.31 (s, 2H), 3.68-3.66 (m, 2H), 3.50-3.47 (m, 2H), 3.35-3.32 (m, 2H), 2.36 (s, 3H), 1.95-1.90 (m, 2H), 1.22-1.19 (m, 3H).

Example 13.2: 7-((Ethyl(4-fluorophenyl)amino)methyl)-3-(3-methoxypropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one




To a solution of 7-((ethyl(4-fluorophenyl)amino)methyl)-3-(3-hydroxypropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (50 mg, 0.13 mmol) in tetrahydrofuran (5 mL) was added sodium hydride (7.0 mg, 60%, 0.29 mmol) and stirred for 1 h at room temperature. Then iodomethane (100 mg, 0.70 mmol) was added to the reaction and the resulting solution was stirred 12 h at room temperature. The reaction was quenched by water/ice (10 mL), extracted with dichloromethane (3×30 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/3) to afford 7-((ethyl(4-fluorophenyl)amino)methyl)-3-(3-methoxypropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one as a light yellow semi-solid (7.9 mg, 15%). LCMS (ESI): M+H⁺=390.1; ¹H NMR (300 MHz, CDCl₃) δ 6.91-6.82 (m, 2H), 6.56-6.52 (m, 2H), 6.07 (s, 1H), 4.24 (s, 2H), 3.46-3.33 (m, 4H), 3.30 (s, 3H), 3.21-3.18 (m, 2H), 2.29 (s, 3H), 1.93-1.85 (m, 2H), 1.22-1.19 (m, 3H).


The following example was prepared in a manner similar to Example 13.1 and 13.2:

No.	Structure/Name	LCMS (M + H) ¹ H NMR
13.3	 <p>7-((Ethyl(4-fluorophenyl)amino)methyl)-3-(4-hydroxybutyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	390.1 ¹ H NMR (300 MHz, CDCl ₃) δ 6.91-6.83 (m, 2H), 6.57-6.50 (m, 2H), 6.09 (s, 1H), 4.25 (s, 2H), 3.71-3.68 (m, 2H), 3.45-3.40 (m, 2H), 3.12-3.19 (m, 2H), 2.35 (s, 1H), 2.29 (s, 3H), 1.74-1.60 (m, 4H), 1.23-1.18 (m, 3H).

Example 13.4: 7-((Ethyl(4-fluorophenyl)amino)methyl)-3-(2-hydroxyethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one




Step 1: 3-[(E)-2-ethoxyvinyl]-7-[(N-ethyl-4-fluoroanilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 3-bromo-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 5.1, Step 1) (50 mg, 0.13 mmol) in 1,4-dioxane/water (0.6/0.2 mL) was added 2-[(E)-2-ethoxyethenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (40 mg, 0.20 mmol), potassium phosphate (80 mg, 0.38 mmol) and tetrakis(triphenylphosphine)palladium (20 mg, 0.02

20 mmol). The resulting solution was stirred for 3 h at 90° C. under nitrogen. After cooling down to room temperature, the reaction mixture was concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/5) to afford the title compound as a light yellow solid (16.9 mg, 35%). LCMS (ESI): M+H⁺=388.0; ¹H NMR: (300 MHz, CDCl₃): δ 6.93-6.85 (m, 2H), 6.60-6.51 (m, 2H), 6.52-5.48 (m, 1H), 6.34-6.19 (m, 1H), 6.10 (s, 1H), 4.27 (s, 2H), 3.99-3.92 (m, 2H), 3.53-3.42 (m, 2H), 2.38 (s, 3H), 1.38-1.35 (m, 3H), 2.23-2.19 (m, 3H).


Step 2: 2-(7-((Ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)acetaldehyde



55 To a solution (E)-3-(2-ethoxyvinyl)-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one in acetone was added 3 N hydrogen chloride (15 mL). The resulting solution was refluxed for 3 h in an oil bath. The pH value of the solution was adjusted to pH 8 with a saturated aqueous sodium bicarbonate solution. The mixture was extracted with dichloromethane (3×50 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo to afford 2-(7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)acetaldehyde was obtained as a light yellow oil (180 mg). The crude product was used in the next step without further purification.


199

Step 3: 7-((Ethyl(4-fluorophenyl)amino)methyl)-3-(2-hydroxyethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 2-(7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)acetaldehyde (180 mg, 0.50 mmol) in methanol (10 mL) was added sodium borohydride (40 mg, 1.06 mmol) at 0° C. After stirring overnight at room temperature, the reaction was quenched by saturated aqueous ammonium chloride (20 mL). The resulting solution was extracted with dichloromethane (3x50 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with dichloromethane/methanol (30/1) to afford 7-((ethyl(4-fluorophenyl)amino)methyl)-3-(2-hydroxyethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one as a white solid (90 mg, 50%). LCMS (ESI): M+H⁺=362.0; ¹H NMR (300 MHz, CDCl₃) δ 6.90-6.84 (m, 2H), 6.55-6.50 (m, 2H), 6.14 (s, 1H), 4.26 (s, 2H), 3.89-3.87 (m, 2H), 3.47-3.39 (m, 4H), 2.32 (s, 3H), 1.21-1.18 (m, 3H).


Example 13.5: 7-[[Ethyl(4-fluorophenyl)amino]methyl]-3-(2-methoxyethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



To a solution of 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(2-hydroxyethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (50 mg, 0.14 mmol) in tetrahydrofuran (5 mL) was added sodium hydride (11 mg, 0.46 mmol) at 0° C. After stirred 0.5 h at room temperature, iodomethane (11 mg, 0.08 mmol) was added to the reaction. After stirring 3 h at room temperature, the reaction was quenched with water/ice (20 mL), extracted with dichloromethane (3x30 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/3) to afford 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(2-methoxyethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a gray semi-solid (11.5 mg, 22%). LCMS (ESI): M+H⁺=376.0; ¹H NMR (300 MHz, CDCl₃) δ 6.94-6.85 (m, 2H), 6.60-6.54 (m, 2H), 6.10 (s, 1H), 4.28 (s, 2H), 3.69-3.66 (m, 2H), 3.49-3.39 (m, 4H), 3.31 (s, 3H), 2.33 (s, 3H), 1.23-1.21 (m, 3H).


200

Example 13.6: 3-(7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)propanamide





To a solution of ethyl 3-(7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)propanoate (from Example 13.1, Step 2) (50 mg, 0.12 mmol) in a 10 mL sealed tube was added 1 M ammonia in methanol (3 mL, 3 mmol). The reaction was sealed and stirred overnight at 80° C. and then concentrated in vacuo. The residue was purified by silica gel chromatography with dichloromethane/methanol (20/1) to afford 3-(7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)propanamide as an off-white solid (10 mg, 21%). LCMS (ESI): M+H⁺=389.0; ¹H NMR (300 MHz, CDCl₃) δ 6.97-6.92 (m, 2H), 6.74-6.72 (m, 2H), 6.19 (s, 1H), 4.32 (s, 2H), 3.54-3.43 (m, 4H), 2.60-2.57 (m, 2H), 2.38 (s, 3H), 1.27-1.24 (m, 3H).

Example 13.7: 3-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]propanenitrile




To a solution of 3-(7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)propanamide (50 mg, 0.13 mmol) and triethylamine (30 mg, 0.30 mmol) in dichloromethane (10 mL) was added (trifluoromethane)sulfonyl trifluoromethanesulfonate (40 mg, 0.14 mmol) at 0° C. and then stirred at room temperature for 2 h. The reaction was quenched by saturated aqueous sodium bicarbonate, extracted with dichloromethane (3x30 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/1) to afford 3-(7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)propanenitrile as an off-white solid (14 mg, 29%). LCMS (ESI): M+H⁺=370.9; ¹H NMR (300 MHz, CDCl₃) δ 6.92-6.86 (m, 2H), 6.65-6.60 (m, 2H), 6.13 (s, 1H), 4.27 (s, 2H), 3.49-3.41 (m, 4H), 2.87-2.86 (m, 2H), 2.42 (s, 3H), 1.22-1.19 (m, 3H).

The following compound was prepared using methods analogous to those described above:

No.	Structure/Name	LCMS (M + H) ¹ H NMR
13.8	 <p>3-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N,N-dimethyl-propanamide</p>	417.2 ¹ H NMR (400 MHz, CDCl ₃) δ 6.94-6.89 (m, 2H), 6.60-6.57 (m, 2H), 6.13 (s, 1H), 4.30 (s, 2H), 3.51-3.43 (m, 4H), 3.02 (s, 3H), 2.96 (s, 3H), 2.71-2.69 (m, 2H), 2.37 (s, 3H), 1.24-1.21 (m, 3H).
13.9	 <p>7-[[Ethyl(4-fluorophenyl)amino]methyl]-3-(3-hydroxy-3-methylbutyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one</p>	403.8 ¹ H NMR (400 MHz, CD ₃ OD) δ 6.94-6.89 (m, 2H), 6.68-6.65 (m, 2H), 6.06 (s, 1H), 4.35 (s, 2H), 3.56-3.51 (m, 2H), 3.27-3.23 (m, 2H), 2.38 (s, 3H), 1.77-1.73 (m, 2H), 1.26 (s, 6H), 1.24-1.21 (m, 3H).


Method 14

Example 14.1: 2-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarboxamide



Step 1: ethyl 2-(7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)cyclopropanecarboxylate

40



55


To a microwave tube with a solution of 3-bromo-7-[[ethyl (4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo [3,2-a]pyrimidin-5-one (from Example 5.1, Step 1) (200 mg, 0.51 mmol) in acetonitrile/water (5/1 mL) was added ethyl 2-(4,4,5,5-tetramethyl-1,3-dioxolan-2-yl)cyclopropane-1-carboxylate (from Example 4.8, Step 1) (900 mg, 2.55 mmol), potassium carbonate (279 mg, 2.04 mmol) and [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (19 mg, 0.025 mmol). The resulting solution was stirred for 1 h at 120° C. The resulting mixture was concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/2) to afford ethyl

65

203


2-(7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carboxylate (80 mg, 37%) as a yellow oil. LCMS (ESI): $M+H^+=430.1$.

Step 2: 2-(7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)cyclopropanecarboxylic acid



To a solution of ethyl 2-(7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carboxylate (80 mg, 0.18 mmol) in tetrahydrofuran/water (20/3 mL) was added lithium hydroxide (22 mg, 0.90 mmol). After stirring overnight at room temperature, the pH value of the solution was adjusted to off 4-5 with 1 N HCl. The reaction mixture was extracted with ethyl acetate (200 mL), washed with brine, dried over anhydrous magnesium sulfate, concentrated in vacuo to afford 2-(7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carboxylic acid as a yellow oil (80 mg). The crude product was used in next step without further purification. LCMS (ESI): $M+H^+=402.1$.

Step 3: 2-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarboxamide




To a solution of 2-(7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carboxylic acid (50 mg, 0.12 mmol) in tetrahydrofuran (10 mL) was added triethylamine (25 mg, 0.24 mmol), chloro(propan-2-yloxy)methanone (22.8 mg, 0.19 mmol). The reaction solution was stirred for 10 min at room temperature. To the reaction was added 1 M ammonia in methanol (1 mL, 1 mmol). The reaction solution was stirred for an additional 20 min at room temperature. The resulting mixture was concentrated in vacuo. The residue was purified by chromatography with dichloromethane/methanol (20/1) to afford 2-(7-[[ethyl(4-fluorophenyl)

204

amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carboxamide as a yellow solid (35 mg, 70%). LCMS (ESI): $M+H^+=400.9$; 1H NMR (400 MHz, CD_3OD) δ 6.86-6.83 (m, 2H), 6.67-6.59 (m, 2H), 6.01 (s, 1H), 4.28 (s, 2H), 3.54-3.41 (m, 2H), 2.63-2.61 (m, 1H), 2.38 (s, 3H), 1.87-1.81 (m, 1H), 1.58-1.52 (m, 1H), 1.28-1.15 (m, 4H).


Example 14.2: 2-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarboxamide



To a solution of 2-(7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carboxamide (40 mg, 0.10 mmol) in tetrahydrofuran (10 mL), trifluoroacetic anhydride (105 mg, 0.50 mmol) and triethylamine (60.6 mg, 0.60 mmol) were added. The reaction solution was then stirred overnight at room temperature. The reaction was then quenched with water (20 mL) and extracted with ethyl acetate (100 mL). The combined organic layers were washed with brine, dried over anhydrous magnesium sulfate and concentrated in vacuo. The residue was purified by Prep-HPLC to afford 2-(7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carbonitrile as a off-white solid (5.4 mg, 14%). LCMS (ESI): $M+H^+=383.0$; 1H NMR (400 MHz, CD_3OD) 6.94-6.90 (m, 2H), 6.69-6.56 (m, 2H), 6.11 (s, 1H), 4.34 (s, 2H), 3.56-3.51 (m, 2H), 2.94-2.90 (m, 1H), 1.99-1.94 (m, 1H), 1.80-1.76 (m, 1H), 1.60-1.55 (m, 1H), 1.25-1.21 (m, 3H)


Method 15

Example 15.1: 7-((5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one




205

Step 1: 1,1,1-trifluorohexane-2,4-dione




To a solution of ethyl 2,2,2-trifluoroacetate (4.20 g, 29.6 mmol) in tetrahydrofuran (120 mL) was added (tert-butoxy) potassium (2.70 g, 24.1 mmol), butan-2-one (1.44 g, 20.0 mmol). The resulting solution was stirred for 12 h at room temperature. The reaction was then quenched by water, extracted with ethyl acetate (3×100 mL), washed with brine, dried over anhydrous magnesium sulfate and concentrated under vacuum to afford 1,1,1-trifluorohexane-2,4-dione (600 mg, 18%) as a yellow solid. The crude product was used in next step without further purification.

Step 2: 5-ethyl-3-(trifluoromethyl)-1H-pyrazole



To a solution of 1,1,1-trifluorohexane-2,4-dione (350 mg, 2.08 mmol) in ethanol (20 mL) was added hydrazine monohydrate (135 mg, 2.19 mmol). The resulting solution was stirred for 12 h at 80° C. in an oil bath. The resulting mixture was concentrated under vacuum. The residue was purified by chromatography with dichloromethane/methanol (100:1) to afford 5-ethyl-3-(trifluoromethyl)-1H-pyrazole (140 mg, 41%) as an off-white solid.


Step 3: 3-bromo-7-((5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 3-bromo-7-(chloromethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 4.1, Step 6) (165 mg, 0.56 mmol) in CH₃CN (20 mL) was added potassium iodide (46 mg, 0.28 mmol), potassium carbonate (155 mg, 1.12 mmol) and 5-ethyl-3-(trifluoromethyl)-1H-pyrazole (110 mg, 0.67 mmol). The resulting solution was stirred for 12 h at 80° C. in an oil bath. The resulting mixture was quenched with water (10 mL), extracted with dichloromethane (3×20 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:2.5) to afford 3-bromo-7-[[5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (150 mg, 63%) as a light yellow solid.


206

Step 4: ethyl 2-(7-((5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)cyclopropanecarboxylate



To a solution of 3-bromo-7-[[5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 4.8, Step 1) (160 mg, 0.38 mmol) in CH₃CN (2 mL) was added potassium carbonate (166 mg, 1.20 mmol), [1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloride (30 mg, 0.04 mmol), ethyl 2-(4,4,5,5-tetramethyl-1,3-dioxolan-2-yl)cyclopropane-1-carboxylate (184 mg, 0.76 mmol) and water (0.6 mL). The reaction mixture was irradiated in a microwave for 1 h at 120° C. The resulting mixture was quenched with water (10 mL), extracted with dichloromethane (3×20 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by chromatography with ethyl acetate/petroleum ether (1:2) to afford 2-(7-[[5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carboxylate (50 mg, 29%) as an off-white solid.


Step 5: 2-(7-((5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)cyclopropanecarboxylic acid



To a solution of 2-(7-[[5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carboxylate (50 mg, 0.11 mmol) in tetrahydrofuran (15 mL) was added a solution of lithium hydroxide (8 mg, 0.33 mmol) in water (1 mL). The resulting solution was stirred for 12 h at room temperature. The pH value of the solution was adjusted to pH 5 with aqueous hydrochloric acid. The resulting solution was extracted with 3×20 mL of ethyl acetate, washed with brine, dried over anhydrous magnesium sulfate, and concentrated under vacuum to afford 2-(7-[[5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carboxylic acid (40 mg, crude) as a reddish solid.

207

Step 6: 7-((5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 2-(7-[[5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carboxylic acid (30 mg, 0.07 mmol) in tetrahydrofuran (10 mL) was added triethylamine (14 mg, 0.14 mmol) and chloro(propan-2-yl)oxy



208

methanone (17 mg, 0.14 mmol). The resulting solution was stirred for 2 h at room temperature. Then a solution of sodium borohydride (8 mg, 0.21 mmol) in water (0.5 mL) was added. The resulting solution was stirred for 12 h at room temperature. The reaction was then quenched by the addition of saturated ammonium chloride aqueous solution, extracted with dichloromethane (3×20 mL), washed with brine, dried over anhydrous magnesium sulfate and concentrated under vacuum. The residue was purified by chromatography with ethyl acetate/petroleum ether (1:1.5) to afford 7-[[5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (9.4 mg, 32%) as a light yellow solid. LCMS (ESI): $[M+1]^+413.1$; $^1\text{H NMR}$ (300 MHz, CD_3OD) δ 6.51 (s, 1H), 5.67 (s, 1H), 5.25 (s, 2H), 3.64-3.54 (m, 2H), 2.77-2.69 (m, 2H), 2.42 (s, 3H), 2.15-2.11 (m, 1H), 1.45-1.29 (m, 4H), 1.27-1.12 (m, 2H).



The following examples were prepared in a manner similar to Example 15.1:

No.	Structure/Name	LCMS (M + H) $^1\text{H NMR}$
15.2	<p>7-((3,5-bis(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	453.20 $^1\text{H NMR}$ (300 MHz, CD_3OD) δ 7.30 (s, 1H), 5.84 (s, 1H), 5.48 (s, 2H), 3.61-3.59 (m, 2H), 2.43 (s, 3H), 2.17-2.15 (m, 1H), 1.37-1.30 (m, 1H), 1.05-0.99 (m, 2H)
15.3	<p>7-((3-chloro-5-methyl-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	365.0 $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 6.04 (s, 1H), 5.67 (s, 1H), 5.06 (s, 2H), 4.06-4.01 (m, 1H), 3.14-3.06 (m, 1H), 2.39 (s, 3H), 2.25 (s, 4H), 1.30-1.20 (m, 1H), 1.05-0.97 (m, 2H)
15.4	<p>7-((5-chloro-3-methyl-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	365.0 $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 6.10 (s, 1H), 5.61 (s, 1H), 5.15 (s, 2H), 4.07-4.02 (m, 2H), 3.12-3.05 (m, 1H), 2.38 (s, 3H), 2.27 (s, 3H), 2.23 (s, 1H), 1.30-1.26 (m, 1H), 1.05-0.94 (m, 2H)

-continued


No.	Structure/Name	LCMS (M + H) ¹ H NMR
15.5	 <p>3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	385 ¹ H NMR (300 MHz, CDCl ₃) δ 7.63-7.58 (m, 1H), 6.61-6.60 (m, 1H), 5.86 (s, 1H), 5.22 (s, 2H), 4.05-4.00 (m, 1H), 3.15-3.08 (m, 1H), 2.38 (s, 3H), 2.29-2.24 (m, 1H), 1.28-1.19 (m, 1H), 1.05-0.95 (m, 2H).
15.6	 <p>7-((1H-indazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	367.1 ¹ H NMR (300 MHz, CDCl ₃) δ 8.09 (s, 1H), 7.79-7.76 (m, 1H), 7.42-7.37 (m, 2H), 7.22-7.17 (m, 1H), 5.57 (s, 1H), 5.47 (s, 2H), 4.35 (s, 2H), 3.07-3.04 (m, 1H), 2.37 (s, 3H), 2.23-2.21 (m, 1H), 1.25-1.21 (m, 1H), 1.02-0.92 (m, 2H)

Example 15.7 and 15.8: 7-((5-(2-hydroxyethyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one and 7-((3-(2-hydroxyethyl)-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



Step 1: ((but-3-yn-1-yloxy)methyl)benzene

35




40 A mixture of but-3-yn-1-ol (3.0 g, 42.8 mmol), sodium hydride (3.0 g, 125.0 mmol), (bromomethyl)benzene (7.2 g, 42.1 mmol) in N,N-dimethylformamide (10 mL) was stirred at room temperature overnight. The resulting mixture was extracted with dichloromethane, washed with brine, washed with brine and concentrated under vacuum to afford crude

45 [(but-3-yn-1-yloxy)methyl]benzene (2.6 g, 38%) as yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 7.30-7.27 (m, 5H), 4.81 (s, 2H), 3.64-3.59 (m, 2H), 2.55-2.49 (m, 2H), 2.02-2.00 (m, 1H).

50

Step 2: 6-(benzyloxy)-1,1,1-trifluorohex-3-yn-2-one

55



60


To a solution of [(but-3-yn-1-yloxy)methyl]benzene (1.00 g, 6.24 mmol) in tetrahydrofuran (20 mL) was added butyllithium (2.5 M in hexanes; 3.0 mL, 7.50 mmol) dropwise at

65 -78° C. The resulting solution was stirred for 0.5 h at -78° C. and 2,2,2-trifluoroethyl 2,2,2-trifluoroacetate (1.47 g, 7.50 mmol), trifluoroborane etherate (1.15 g, 7.50 mmol)

211


was added to the reaction mixture and stirred for further 3 h at -78°C . The resulting mixture was washed with brine, extracted with dichloromethane ($3\times 100\text{ mL}$), washed with brine, dried over sodium sulfate and concentrated under vacuum. The residue was purified by column chromatography with ethyl acetate/petroleum ether (1:100) to afford 6-(benzyloxy)-1,1,1-trifluorohex-3-yn-2-one (0.6 g, 38%) as yellow oil. $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 7.40-7.27 (m, 5H), 4.58 (s, 2H), 3.72-3.68 (m, 2H), 2.83-2.79 (m, 2H).

Step 3: 5-(2-(benzyloxy)ethyl)-3-(trifluoromethyl)-1H-pyrazole




A mixture of 6-(benzyloxy)-1,1,1-trifluorohex-3-yn-2-one (0.6 g, 2.34 mmol) in hydrazine monohydrate (0.5 g, 10 mmol) and ethanol (10 mL) was stirred at 85°C for 3 h. The resulting mixture was extracted with dichloromethane ($3\times 100\text{ mL}$), washed with brine, dried over sodium sulfate and concentrated under vacuum to afford 5-[2-(benzyloxy)ethyl]-3-(trifluoromethyl)-1H-pyrazole (0.6 g, 95%) as a yellow oil. $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 7.40-7.29 (m, 5H), 6.32 (s, 1H), 4.57 (s, 2H), 3.76-3.72 (m, 2H), 2.96-2.92 (m, 2H).

Step 4:
2-(3-(trifluoromethyl)-1H-pyrazol-5-yl)ethanol



A mixture of 5-[2-(benzyloxy)ethyl]-3-(trifluoromethyl)-1H-pyrazole (400 mg, 1.48 mmol), palladium on carbon (150 mg), O-(hydroxychloro)oxidanol (0.01 mL) in methanol (15 mL) and tetrahydrofuran (4 mL) was stirred at room temperature for 70 min under hydrogen. The solids were filtered off and the resulting mixture was concentrated under vacuum to afford 2-[3-(trifluoromethyl)-1H-pyrazol-5-yl]ethan-1-ol (250 mg, 95%) as a brown solid. LCMS $[\text{M}+\text{H}]^+ = 181.0$.

Step 5: 3-bromo-7-((5-(2-hydroxyethyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one and 3-bromo-7-((3-(2-hydroxyethyl)-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one




$\text{R}_1, \text{R}_2 = \text{CF}_3, \text{CH}_2\text{CH}_2\text{OH}$

212

A mixture of 2-(3-(trifluoromethyl)-1H-pyrazol-5-yl)ethan-1-ol (480 mg, 2.66 mmol), 3-bromo-7-(chloromethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 4.1, Step 6) (300 mg, 1.02 mmol), potassium carbonate (690 mg, 4.99 mmol), and potassium iodide (0.28 g, 1.70 mmol) in CH_3CN (15 mL) was stirred at 85°C for 2 h. The resulting mixture was extracted with dichloromethane ($3\times 100\text{ mL}$), washed with brine, dried over sodium sulfate and concentrated under vacuum. The residue was purified by silica gel column chromatography with ethyl acetate/petroleum ether (1:2) to afford a mixture of 3-bromo-7-[[5-(2-hydroxyethyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one and 3-bromo-7-((3-(2-hydroxyethyl)-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (470 mg, 40%) as a light yellow solid. LCMS (ESI): $[\text{M}+\text{H}]^+ = 437.0, 439.0$.

Step 6: 7-((5-(2-hydroxyethyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one and 7-((3-(2-hydroxyethyl)-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



A mixture of 3-bromo-7-[[5-(2-hydroxyethyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (200 mg, 0.46 mmol), [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (35 mg, 0.05 mmol), sodium carbonate (100 mg, 0.94 mmol), potassium *trans*-2-(hydroxymethyl)cyclopropyltrifluoroborate (from Example 4.1, Step 2) (160 mg, 0.90 mmol) in CH_3CN (6 mL) and H_2O (2 mL) was irradiated with microwave radiation for 1.5 h at 120°C in a sealed-tube. The resulting mixture was extracted with dichloromethane ($3\times 20\text{ mL}$), washed with brine, dried over sodium sulfate and concentrated under vacuum. The residue was purified by Prep-HPLC (Xselect CSH Prep C18 OBD Column, 5 μm , $19\times 150\text{ mm}$; mobile phase, water with 0.03% $\text{NH}_3\text{--H}_2\text{O}$ and MeCN (23.0% MeCN up to 28.0% in 20 min)) to afford 7-[[5-(2-hydroxyethyl)-3-(trifluoromethyl)-1H-pyrazol-1-

213

yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (19.7 mg, 10.2%) as a white solid and 7-((3-(2-hydroxyethyl)-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (4.3 mg, 2.2%).


Example 15.7

7-[[5-(2-hydroxyethyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one. LCMS (ESI): $[M+H]^+=429.1$; 1H NMR (300 MHz, $CDCl_3$) δ 6.46 (s, 1H), 5.81 (s, 1H), 5.24 (s, 2H), 4.04-3.93 (m, 1H), 3.91-3.89 (m, 2H), 3.13-3.06 (m, 1H), 2.94-2.90 (m, 4H), 2.38 (s, 3H), 2.27-2.21 (m, 1H), 1.27-1.16 (m, 1H), 1.01-0.90 (m, 2H).

Example 15.8


7-((3-(2-hydroxyethyl)-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one. LCMS (ESI): $[M+H]^+=429.0$; 1H NMR (300 MHz, $CDCl_3$) δ 6.91 (s, 1H), 5.56 (s, 1H), 5.26 (s, 2H), 4.06-4.01 (m, 1H), 3.96-3.92 (m, 2H), 3.12-3.06 (m, 1H), 2.94-2.89 (m, 2H), 2.38 (s, 3H), 2.21-2.15 (m, 1H), 1.27-1.16 (m, 1H), 1.03-0.90 (m, 2H).

Example 15.9 and 15.10: 7-((3-cyclopropyl-4-fluoro-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one and 7-((5-cyclopropyl-4-fluoro-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



Step 1:

1-cyclopropyl-4,4,4-trifluorobutane-1,3-dione




214

A mixture of sodium metal (276 mg, 12.0 mmol) and ethanol (20 mL) was stirred for 20 minutes at room temperature. To the reaction mixture was added 1-cyclopropylethanol-1-one (1.42 mg, 0.0200 mmol) and ethyl 2,2,2-trifluoroacetate (840 mg, 5.91 mmol) and the resulting solution was stirred for 2 days at room temperature. The reaction was quenched with water (50 mL), extracted with dichloromethane (3x20 mL) washed with brine, and dried over anhydrous sodium sulfate and concentrated under vacuum to afford 1-cyclopropyl-4,4,4-trifluorobutane-1,3-dione (737 mg, 69%) as colorless oil. The crude product was used in next step without further purification.


Step 2:

3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazole



To a solution of 1-cyclopropyl-4,4,4-trifluorobutane-1,3-dione (400 mg, 2.22 mmol) in ethanol (20 mL) was added hydrazine monohydrate (132 mg, excess) and the resulting solution was stirred for 2 days at 80° C. The resulting mixture was concentrated under vacuum. The residue was purified by chromatography with 2% methanol in dichloromethane to afford 3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazole (258 mg, 66%) as an off-white solid. LCMS (ESI): $[M+H]^+=177.0$; 1H NMR (300 MHz, $CDCl_3$) δ 6.27 (s, 1H), 2.08-1.95 (m, 1H), 2.77-2.64 (m, 2H), 1.11-1.02 (m, 2H).



Step 3: 5-cyclopropyl-4-fluoro-3-(trifluoromethyl)-1H-pyrazole



To a solution of 5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazole (500 mg, 2.84 mmol) in CH_3CN (10 mL) maintained with an inert atmosphere of nitrogen was added Selectfluor® (1.0 g, 2.84 mmol). The resulting solution was stirred for 12 h at 75° C. The resulting mixture was then concentrated under vacuum. The residue was purified by chromatography with 1% methanol in dichloromethane to afford 5-cyclopropyl-4-fluoro-3-(trifluoromethyl)-1H-pyrazole (280 mg, 51%) as a yellow solid. LCMS (ESI): $[M+H]^+=195.0$; 1H NMR (300 MHz, $CDCl_3$) δ 1.81 (m, 1H), 1.03 (m, 2H), 0.85 (m, 2H).


215

Step 4: 3-bromo-7-((3-cyclopropyl-4-fluoro-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one and 3-bromo-7-((5-cyclopropyl-4-fluoro-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 3-bromo-7-(chloromethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 4.1, Step 6) (250 mg, 0.85 mmol) in CH₃CN (20 mL) was added 5-cyclopropyl-4-fluoro-3-(trifluoromethyl)-1H-pyrazole (200 mg, 1.03 mmol), potassium iodide (80 mg, 0.42 mmol), potassium carbonate (250 mg, 1.81 mmol). The resulting solution was stirred for 12 h at 80° C. The resulting mixture was concentrated under vacuum. The residue was purified by chromatography with ethyl acetate/petroleum ether (1:3) to afford 3-bromo-7-[[3-cyclopropyl-4-fluoro-5-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (100 mg, 20%) as a light yellow solid (LCMS (ESI): [M+H]⁺=452.2) and 3-bromo-7-[[5-cyclopropyl-4-fluoro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (200 mg, 50%) as a light yellow solid (LCMS (ESI): [M+H]⁺=452.0).

Step 5: 7-((3-cyclopropyl-4-fluoro-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (Example 15.9)




To a solution of 3-bromo-7-[[3-cyclopropyl-4-fluoro-5-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (90 mg, 0.20 mmol) in CH₃CN (2 mL) under nitrogen, was added sodium carbonate (43 mg, 0.41 mmol), [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (30 mg, 0.04 mmol), potassium trans-2-(hydroxymethyl)cyclopropyltrifluoroborate

216



(from Example 4.1, Step 2) (140 mg, 0.79 mmol) and water (0.6 mL). The reaction mixture was heated under microwave irradiation for 90 min at 120° C. The reaction mixture was then quenched with water (10 mL), extracted with ethyl acetate (3×20 mL), washed with brine, dried over anhydrous magnesium sulfate and concentrated under vacuum. The residue was purified on a silica gel column with ethyl acetate/petroleum ether (1:1), afford 7-[[3-cyclopropyl-4-fluoro-5-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (10.5 mg, 12%) as a light yellow solid. LCMS (ESI): [M+H]⁺=442.9; ¹H NMR (300 MHz, CD₃OD) δ 5.65 (s, 1H), 5.18 (s, 1H), 5.21 (s, 2H), 3.59 (m, 2H), 2.48 (s, 3H), 2.18 (m, 1H), 1.90 (m, 1H), 1.34 (m, 2H), 0.98 (m, 3H), 0.84 (m, 2H).

Step 6: 7-((5-cyclopropyl-4-fluoro-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (Example 15.10)




Into a 10-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed 3-bromo-7-[[5-cyclopropyl-4-fluoro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (100 mg, 0.22 mmol), acetonitrile (2 mL), [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (34 mg, 0.05 mmol), sodium carbonate (48 mg, 0.45 mmol), potassium trans-2-(hydroxymethyl)cyclopropyltrifluoroborate (from Example 4.1, Step 2) (153 mg, 0.86 mmol) and water (0.6 mL). The reaction mixture was irradiated with microwave radiation for 90 min at 120° C. The resulting solution was extracted with ethyl acetate (3×20 mL), washed with brine, dried over anhydrous magnesium sulfate and concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1:1), to afford 7-[[5-cyclopropyl-4-fluoro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (18.5 mg, 19%) as an off-white solid. LCMS (ESI): [M+H]⁺=443.0; ¹H NMR (300 MHz, CD₃OD) δ 5.85 (s, 1H), 5.38 (s, 2H), 3.65 (m, 2H), 2.43 (s, 3H), 2.12 (m, 1H), 1.80 (m, 1H), 1.32 (m, 2H), 0.99 (m, 4H), 0.85 (m, 2H).

The following examples were prepared in a manner similar to Example 15.7-15.10:

No.	Structure/Name	LCMS (M + H) ¹ H NMR
15.11	 <p>7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	425.25 ¹ H NMR (300 MHz, CDCl ₃) δ 6.35 (s, 1H), 5.54 (s, 1H), 5.22 (s, 2H), 4.06-4.02 (m, 2H), 3.12-3.05 (m, 1H), 2.38 (s, 3H), 2.26-2.22 (m, 1H), 2.00-1.91 (m, 1H), 1.32-1.21 (m, 1H), 1.05-0.92 (m, 4H), 0.83-0.74 (m, 2H)
15.12	 <p>7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	425.0 ¹ H NMR (300 MHz, CDCl ₃) δ 6.19 (s, 1H), 5.67 (s, 1H), 5.32 (s, 2H), 4.06-4.01 (m, 1H), 3.14-3.07 (m, 1H), 2.41 (s, 3H), 2.29-2.23 (m, 1H), 1.74-1.66 (m, 1H), 1.28-1.23 (m, 1H), 1.06-0.95 (m, 4H), 0.78-0.69 (m, 2H)

Examples 15.13, 15.14, 15.15, and 15.16: 7-[(5-cyclopropyl-3-methyl-pyrazol-1-yl)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (Enantiomers 1 and 2) and 7-[(3-cyclopropyl-5-methyl-pyrazol-1-yl)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (Enantiomers 1 and 2)



Examples 15.13, 15.14, 15.15 and 15.16 were prepared in a manner analogous to 15.7, where 2-[3-(trifluoromethyl)-1H-pyrazol-5-yl]ethan-1-ol was replaced by 5-cyclopropyl-3-methyl-1H-pyrazole in Step 5. Following the cross-cou-

pling procedure in Step 6, the crude product was purified by Prep-HPLC (Column, XBridge Prep C18 OBD Column, 19*150 mm, 5 μm; mobile phase, CH₃CN and water with 0.5% NH₃H₂O (35% CH₃CN up to 45% in 10 mins); Detector, UV 254/220 nm) to afford a mixture of racemic 7-[(5-cyclopropyl-3-methyl-pyrazol-1-yl)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one and racemic 7-[(3-cyclopropyl-5-methyl-pyrazol-1-yl)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (60 mg, 70%) as a white solid. This material was purified by Chiral-HPLC (Column, CHIRALCEL, OJ-H (2×25 cm, 5 μm); mobile phase, Hex: EtOH=85:15, 25 mins, flow rate, 20 ml/min; Detector, UV 254/220 nm) to afford four isomers:

Example 15.13

7-[(5-cyclopropyl-3-methyl-pyrazol-1-yl)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (peak 1, enantiomer 1). Retention time=8.2 min; Yield=4.2 mg, 5.0%; LCMS (ESI): M+H⁺=271.1; ¹H NMR (300 MHz, CDCl₃) δ 5.75 (s, 1H), 5.68 (s, 1H), 5.31 (s, 2H), 4.06-3.98 (m, 2H), 3.15-3.03 (m, 1H), 2.37 (s, 3H), 2.28-2.20 (m, 4H), 1.63-1.57 (m, 1H), 1.23-1.13 (m, 1H), 1.02-0.83 (m, 4H), 0.70-0.61 (m, 2H).

Example 15.14

7-[(5-cyclopropyl-3-methyl-pyrazol-1-yl)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (peak 2, enantiomer 2). Retention time=10 min; Yield=2.8 mg, 3.3%; LCMS (ESI): M+H⁺=271.1; ¹H NMR (300 MHz, CDCl₃) δ 5.75 (s, 1H), 5.68 (s, 1H), 5.31 (s, 2H), 4.06-3.98 (m, 2H), 3.15-3.03 (m, 1H), 2.37 (s, 3H),

219

2.28-2.20 (m, 4H), 1.63-1.57 (m, 1H), 1.23-1.13 (m, 1H),
1.02-0.83 (m, 4H), 0.70-0.61 (m, 2H).

Example 15.15




7-[(3-cyclopropyl-5-methyl-pyrazol-1-yl)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (peak 3, enantiomer 1). Retention time=14.8 min; Yield=6.9 mg, 8.2%; LCMS (ESI): M+H⁺=271.1; ¹H NMR (300 MHz, CDCl₃) δ 5.75 (s, 1H), 5.57 (s, 1H), 5.08 (s, 2H), 4.04-4.00 (m, 2H), 3.12-3.05 (m, 1H), 2.37 (s, 3H), 2.28-2.19 (m, 4H), 1.94-1.86 (m, 1H), 1.23-1.11 (m, 1H), 1.02-0.86 (m, 4H), 0.71-0.63 (m, 2H).

220





Example 15.16

7-[(3-cyclopropyl-5-methyl-pyrazol-1-yl)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (peak 4, enantiomer 2). Retention time=17.6 min; Yield=14.6 mg, 17.4%; LCMS (ESI): M+H⁺=271.1; ¹H NMR (300 MHz, CDCl₃) δ 5.75 (s, 1H), 5.57 (s, 1H), 5.08 (s, 2H), 4.04-4.00 (m, 2H), 3.12-3.05 (m, 1H), 2.37 (s, 3H), 2.28-2.19 (m, 4H), 1.94-1.86 (m, 1H), 1.23-1.11 (m, 1H), 1.02-0.86 (m, 4H), 0.71-0.63 (m, 2H).




The following examples were prepared in a manner similar to the preceding examples:

No.	Structure/Name	LCMS (M + H) ¹ H NMR
15.17	 <p>7-[(3,5-dicyclopropylpyrazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	397.1 ¹ H NMR (300 MHz, CDCl ₃) δ 5.62 (s, 1H), 5.55 (s, 1H), 5.18 (s, 2H), 4.06-3.98 (m, 2H), 3.12-3.01 (m, 1H), 2.37 (s, 3H), 2.28-2.15 (m, 1H), 1.89-1.80 (m, 1H), 1.60-1.49 (m, 2H), 1.26-1.13 (m, 2H), 1.02-0.80 (m, 5H), 0.71-0.53 (m, 3H)
15.18	 <p>7-((3,5-dimethyl-1H-pyrazol-1-yl)methyl)-6-fluoro-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	363.0 ¹ H NMR (300 MHz, CDCl ₃) δ 5.85 (s, 1H), 5.22-5.19 (m, 2H), 4.09-4.00 (m, 1H), 3.77-3.74 (m, 1H), 3.25-3.18 (m, 1H), 2.38 (s, 3H), 2.31 (s, 3H), 2.29-2.27 (m, 1H), 2.23 (s, 3H), 1.32-1.28 (m, 1H), 1.08-1.01 (m, 2H).
15.19	 <p>1-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-3-(trifluoromethyl)-1H-pyrazole-5-carbonitrile</p>	410.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.12 (s, 2H), 5.93 (s, 1H), 5.39 (s, 2H), 4.06-4.01 (m, 1H), 3.15-3.08 (m, 1H), 2.39 (s, 3H), 2.39-2.23 (m, 1H), 1.33-1.22 (m, 1H), 1.05-1.02 (m, 2H)


-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
15.20	 <p data-bbox="337 680 667 764">7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-6-fluoro-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	443.0 ¹ H NMR (300 MHz, CD ₃ OD) δ 6.29 (s, 1H), 5.53 (s, 2H), 3.71-3.57 (m, 2H), 2.41 (s, 3H), 2.17-2.15 (m, 1H), 2.01-1.94 (m, 1H), 1.46-1.41 (m, 1H), 1.09-1.01 (m, 4H), 0.77-0.72 (m, 2H).
15.21	 <p data-bbox="337 1058 667 1121">1-[[3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]-5-methyl-pyrazole-3-carbonitrile</p>	356.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 6.81 (s, 1H), 5.70 (s, 1H), 5.29 (s, 2H), 4.53 (s, 1H), 3.46 (d, J = 5.9 Hz, 2H), 2.36 (d, J = 1.5 Hz, 3H), 2.32 (s, 3H), 2.00 (dtd, J = 7.1, 5.4, 1.7 Hz, 1H), 1.27 (dp, J = 8.5, 5.8 Hz, 1H), 0.85 (ddt, J = 16.0, 8.6, 5.1 Hz, 2H).
15.22	 <p data-bbox="337 1415 667 1499">7-[[3-(difluoromethyl)-5-methyl-pyrazol-1-yl]methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	381.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 6.91 (t, J = 54.9 Hz, 1H), 6.38 (s, 1H), 5.52 (s, 1H), 5.19 (s, 2H), 4.52 (t, J = 5.7 Hz, 1H), 3.45 (t, J = 5.8 Hz, 2H), 2.36 (d, J = 1.5 Hz, 3H), 2.30 (s, 3H), 2.05-1.91 (m, 1H), 1.27 (dt, J = 8.7, 5.6 Hz, 1H), 0.97-0.66 (m, 3H).
15.23	 <p data-bbox="337 1856 667 1915">7-[[6-fluoroindazol-1-yl]methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one</p>	385.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.18 (s, 1H), 7.85 (dd, J = 8.9, 5.2 Hz, 1H), 7.59 (dd, J = 9.8, 2.2 Hz, 1H), 7.10-7.00 (m, 1H), 5.50 (s, 2H), 5.47 (s, 1H), 4.50 (t, J = 5.5 Hz, 1H), 3.43 (t, J = 5.7 Hz, 2H), 2.35 (d, J = 1.8 Hz, 3H), 2.04-1.89 (m, 1H), 1.25 (dp, J = 9.0, 5.9 Hz, 1H), 0.82 (ddt, J = 15.2, 8.6, 5.0 Hz, 2H).

-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
15.24	 <p>7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-isopropylthiazolo[3,2-a]pyrimidin-5-one</p>	453.2 ¹ H NMR (400 MHz, DMSO-d ₆) δ 6.48 (s, 1H), 5.63 (s, 1H), 5.37 (s, 2H), 4.57 (t, J = 5.6 Hz, 1H), 3.65 (p, J = 6.7 Hz, 1H), 3.55-3.37 (m, 2H), 2.05-1.88 (m, 2H), 1.28 (dq, J = 8.8, 5.6 Hz, 1H), 1.22 (d, J = 6.8 Hz, 6H), 1.01-0.91 (m, 2H), 0.91-0.85 (m, 1H), 0.81 (dt, J = 8.8, 5.2 Hz, 1H), 0.73 (dt, J = 6.7, 4.0 Hz, 2H).
15.25	 <p>7-[(3,5-dimethylpyrazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methylthiazolo[3,2-a]pyrimidin-5-one</p>	345.2 ¹ H NMR (400 MHz, DMSO-d ₆) δ 5.89 (s, 1H), 5.35 (s, 1H), 5.02 (s, 2H), 4.52 (t, J = 5.6 Hz, 1H), 3.45 (t, J = 5.8 Hz, 2H), 2.36 (d, J = 1.6 Hz, 3H), 2.19 (s, 3H), 2.10 (s, 3H), 2.00 (qd, J = 7.4, 6.5, 2.6 Hz, 1H), 1.33-1.20 (m, 1H), 0.95-0.76 (m, 2H).
15.26	 <p>7-[(5-fluorindazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methylthiazolo[3,2-a]pyrimidin-5-one</p>	385.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.14 (s, 1H), 7.74 (dd, J = 9.1, 4.3 Hz, 1H), 7.58 (dd, J = 9.2, 2.4 Hz, 1H), 7.31 (td, J = 9.2, 2.5 Hz, 1H), 5.54 (s, 2H), 5.47 (s, 1H), 4.50 (t, J = 5.5 Hz, 1H), 3.43 (t, J = 5.5 Hz, 2H), 2.35 (s, 3H), 1.97 (ddd, J = 9.6, 6.7, 4.8 Hz, 1H), 1.31-1.15 (m, 1H), 0.82 (ddt, J = 16.2, 8.7, 5.1 Hz, 2H).

Example 15.27: 5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl]-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one




To a solution of 3-bromo-7-[[5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 15.1, Step 3) (80 mg, 0.19


mmol), 1,4-dioxane (2 mL), potassium phosphate (80 mg, 0.38 mmol), tetrakis(triphenylphosphine)palladium (22 mg, 0.02 mmol) and water (0.2 mL) in 1,4-dioxane (2 mL) under nitrogen, was added (pyrimidin-5-yl)boronic acid (47 mg, 0.38 mmol). The resulting solution was stirred for 3 h at 90° C. in an oil bath. The resulting mixture was quenched by water (10 mL), and extracted with dichloromethane (3×30 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by chromatography with ethyl acetate/petroleum ether (1:1) to afford 7-[[5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-3-(pyrimidin-5-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (18.9 mg, 24%) as a yellow solid. LCMS (ESI): [M+H]⁺=462.0; ¹H NMR (300 MHz, CD₃OD) δ 9.18 (s, 1H), 8.82 (s, 2H), 6.52 (s, 1H), 5.70 (s, 1H), 5.31 (s, 2H), 2.80-2.72 (m, 2H), 2.30 (s, 3H), 1.34-1.29 (m, 3H).

225

Example 15.28 and 15.29: 7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one and 7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one



Step 1: 3-bromo-7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one and 3-bromo-7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one




$R_1, R_2 = CF_3, cPr$

To a solution of 3-bromo-7-(chloromethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 4.1, Step 6) (373 mg, 1.27 mmol) in CH_3CN (10 mL) was added 3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazole (from Example 15.9, Step 2) (450 mg, 2.55 mmol), cesium carbonate (223 mg, 0.68 mmol) and potassium iodide (160 mg). The resulting solution was stirred overnight at 80° C. The solids were filtered off and the resulting mixture was concentrated under vacuum. The crude product was purified by Prep-HPLC (Xbridge Shield RP18 OBD Column, 5 μm , 19x150 mm; mobile phase, water with 0.03% $NH_3 \cdot H_2O$ and CH_3CN (10.0% CH_3CN up to 32.0% in 10 min, up to 100.0% in 1 min, hold 100.0% in 1 min, down to 10.0% in 2 min); Detector, uv 254 nm) to afford 3-bromo-7-[[3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (18 mg,

226


3%) as a white solid (LCMS (ESI): $[M+H]^+ = 434.9$; 1H NMR (300 MHz, $CDCl_3$) δ 6.36 (s, 1H), 5.54 (s, 1H), 5.21 (s, 2H), 2.35 (s, 3H), 1.90-2.01 (m, 1H), 0.94-1.00 (m, 2H), 0.71-0.79 (m, 2H)) and 3-bromo-7-[[5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (54 mg, 10%) as a white solid (LCMS (ESI): $[M+H]^+ = 434.9$; 1H NMR (300 MHz, $CDCl_3$) δ 6.18 (s, 1H), 5.66 (s, 1H), 5.30 (s, 2H), 2.36 (s, 3H), 1.72-1.65 (m, 1H), 1.03-0.98 (m, 2H), 0.76-0.71 (m, 2H)).

Step 4: 7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one (Example 15.28)



To a solution of 3-bromo-7-[[3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (12 mg, 0.03 mmol) in CH_3CN (1.5 mL) and water (0.5 mL) was added (pyrimidin-5-yl)boronic acid (7 mg, 0.060 mmol), potassium phosphate (12 mg, 0.06 mmol) and tetrakis(triphenylphosphine)palladium (2.0 mg, 10 μmol %). After stirring 1 h at 100° C. under nitrogen atmosphere, the resulting mixture was concentrated under vacuum. The residue was purified by silica gel chromatography with 1% methanol in dichloromethane. The crude product was purified by Prep-HPLC (SunFire Prep C_{18} OBD Column, 5 μm , 19x150 mm; mobile phase, water with 10 mmol NH_4HCO_3 and CH_3CN (30.0% CH_3CN up to 60.0% in 8 min, up to 95.0% in 2 min, down to 30.0% in 2 min); Detector, UV 254/220 nm) to afford 7-[[3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-3-(pyrimidin-5-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (6.1 mg, 51%) as a white solid. LCMS (ESI): $[M+H]^+ = 433.2$; 1H NMR (300 MHz, $CDCl_3$) δ 9.26 (s, 1H), 8.70 (s, 2H), 6.35 (s, 1H), 5.49 (s, 1H), 5.26 (s, 2H), 2.29 (s, 3H), 1.98-1.88 (m, 1H), 0.99-0.93 (m, 2H), 0.76-0.74 (m, 2H).

Step 5: 7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one (Example 15.29)






227



To a solution of 3-bromo-7-[[5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (33 mg, 0.08 mmol) in CH₃CN (1.5 mL) and water (0.5 mL) was added (pyrimidin-5-yl)boronic acid (21 mg, 0.17 mmol), potassium phosphate (35 mg, 0.16 mmol) and tetrakis(triphenylphosphine)palladium (6 mg, 0.01 mmol). After stirring 1 h at 100° C. under nitrogen atmosphere, the resulting mixture was concentrated under vacuum. The residue was purified by chromatography with 1% methanol in dichloromethane. The crude product

228

NH₄HCO₃ and CH₃CN (30.0% CH₃CN up to 60.0% in 8 min, up to 95.0% in 2 min, down to 30.0% in 2 min); Detector, UV 254/220 nm) to afford 7-[[5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-3-(pyrimidin-5-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (29.2 mg, 89%) as a white solid. LCMS (ESI): [M+H]⁺=433.2; ¹H NMR (300 MHz, CDCl₃) δ 9.26 (s, 1H), 8.70 (s, 2H), 6.18 (s, 1H), 5.64 (s, 1H), 5.35 (s, 2H), 2.29 (s, 3H), 1.77-1.68 (m, 1H), 1.04-0.98 (m, 2H), 0.75-0.70 (m, 2H).



The following examples were prepared in a manner similar to Example 15.27, 15.28, and 15.29:

No.	Structure/Name	LCMS (M + H)	¹ H NMR
15.30	 <p>7-((3,5-bis(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	461.10	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.19 (s, 1H), 8.85 (s, 2H), 7.69 (s, 1H), 5.88 (s, 1H), 5.56 (s, 2H), 2.21 (s, 3H)
15.31	 <p>2-methyl-3-(pyrimidin-5-yl)-7-((5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	393	¹ H NMR (300 MHz, CDCl ₃) δ 9.27 (s, 1H), 8.70 (s, 2H), 7.61-7.60 (m, 1H), 6.61-6.60 (m, 1H), 5.84 (s, 1H), 5.25 (s, 2H), 2.28 (s, 3H)
15.32	 <p>2-methyl-7-((5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	448.15	¹ H NMR (300 MHz, CDCl ₃) δ 9.25 (s, 1H), 8.69 (s, 2H), 6.37 (s, 1H), 5.61 (s, 1H), 5.20 (s, 2H), 2.33 (s, 3H), 2.29 (s, 3H)


No.	Structure/Name	LCMS (M + H)	¹ H NMR
15.33	 <p>7-((3,5-bis(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	382.95	¹ H NMR (300 MHz, CDCl ₃) δ 8.85 (s, 1H), 7.69 (s, 1H), 5.88 (s, 1H), 5.56 (s, 2H), 2.21 (s, 3H)
15.34	 <p>7-((1H-indazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	375	¹ H NMR (300 MHz, CDCl ₃) δ 9.22 (s, 1H), 8.65 (s, 2H), 8.08 (s, 1H), 7.76 (m, 1H), 7.42-7.36 (m, 2H), 7.26-7.16 (m, 1H), 5.30 (s, 2H), 2.27 (s, 3H)

Method 16

Example 16.1 and 16.2: 7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide and 7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




Step 1: ethyl 2-oxobutanoate



Into a 10 L 4-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen was placed

diethyl oxalate (300 g, 2.05 mol, 1.00 equiv) and tetrahydrofuran (4.4 L), followed by ethyl magnesium bromide (740 mL, 1.08 equiv) dropwise with stirring at -10° C. over 2 h. The resulting solution was stirred at -10° C. for 30 min and quenched by the addition of 500 mL 3 M hydrogen chloride. The pH value of the solution was adjusted to pH 4 with hydrogen chloride (3 mol/L) and the resulting solution was extracted with 2×1 L of dichloromethane. The combined organic layers were washed with 1×2 L of sodium chloride, dried over anhydrous sodium sulfate and concentrated under vacuum to afford 200 g (crude) of ethyl 2-oxobutanoate as a yellow oil.


Step 2: ethyl 3-bromo-2-oxobutanoate



Into a 10-L 4-necked round-bottom flask was placed ethyl 2-oxobutanoate (265 g, 2.04 mol, 1.00 equiv), chloroform (5 L), a solution of HBr in AcOH (500 mL), and Br₂ (325 g, 2.03 mol, 1.00 equiv). The resulting solution was stirred at 70° C. for 2 h, cooled to room temperature and concentrated under vacuum to afford 392.9 g (92%) of ethyl 3-bromo-2-oxobutanoate as a brown oil.

231


Step 3: ethyl
2-amino-5-methyl-1,3-thiazole-4-carboxylate



Into a 10-L round-bottom flask was placed ethyl 3-bromo-2-oxobutanoate (392.9 g, 1.88 mol, 1.00 equiv), 1,4-dioxane (3.5 L), and thiourea (143.8 g, 1.89 mol, 1.01 equiv). The resulting solution was stirred overnight at 100° C. and cooled to room temperature. The solids were then collected by filtration, washed with Et₂O and dried under vacuum to afford 310 g (89%) of ethyl 2-amino-5-methyl-1,3-thiazole-4-carboxylate as a light brown solid.


Step 4:

2-amino-N,5-dimethyl-1,3-thiazole-4-carboxamide



Into a 1-L pressure tank reactor was placed ethyl 2-amino-5-methyl-1,3-thiazole-4-carboxylate (140 g, 751.75 mmol, 1.00 equiv) and 30% methylamine in EtOH (500 mL). The resulting solution was stirred overnight at 85° C. and concentrated under vacuum. The residue was purified on a silica gel column eluting with dichloromethane/methanol (20/1) to afford 80 g (62%) of 2-amino-N,5-dimethyl-1,3-thiazole-4-carboxamide as a yellow solid.

Step 5: 7-(chloromethyl)-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide





Into a 3-L 3-necked round-bottom flask was placed 2-amino-N,5-dimethyl-1,3-thiazole-4-carboxamide (80 g, 467.24 mmol, 1.00 equiv), ethyl 4-chloro-3-oxobutanoate (154 g, 935.68 mmol, 2.00 equiv) and polyphosphoric acid (PPA) (800 g). The resulting solution was stirred at 110° C. for 2 h, cooled to 80° C. and quenched by the addition of 100 mL of water. The pH value of the solution was adjusted to pH 8 with sodium hydroxide (10% aq.). The solids were filtered off and the filtrate was extracted with dichloromethane (5 L×5). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated under vacuum. The

232

residue was purified on a silica gel column eluting with dichloromethane/methanol (20/1) to give 80 g (63%) of 7-(chloromethyl)-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide as a tan solid. LCMS (ESI): [M+H]⁺=272; ¹H NMR (400 MHz, DMSO-d₆): δ 8.38 (q, J=4.8 Hz, 1H), 6.39 (s, 1H), 4.60 (s, 2H), 2.74 (d, J=4.8 Hz, 3H), 2.30 (s, 3H).


Step 6: 7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide (Example 16.1) and 7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide (Example 16.2)




To a solution of 7-(chloromethyl)-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (100 mg, 0.37 mmol), potassium iodide (30 mg, 0.19 mmol) and potassium carbonate (100 mg, 0.74 mmol) in CH₃CN (10 mL) was added 5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazole (from Example 15.9, Step 2) (80 mg, 0.45 mmol). The reaction mixture was stirred 3 h at 85° C. in an oil bath. After filtration to remove solids and concentration under vacuum, the residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (2:1) to afford 7-[[5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (50.4 mg, 33%) as a white solid (LCMS (ESI): [M+H]⁺=412.0; ¹H NMR (300 MHz, CDCl₃) δ 6.27 (bs, 1H), 6.19 (s, 1H), 5.64 (s, 1H), 5.32 (s, 2H), 3.03 (m, 3H), 2.40 (s, 3H), 1.71-1.65 (m, 1H), 1.03-0.97 (m, 2H), 0.76-0.66 (m, 2H)) and 7-[[3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (20.7 mg, 14%) as a white solid (LCMS (ESI): [M+H]⁺=411.9; ¹H NMR (300 MHz, CDCl₃) δ 6.36 (s, 1H), 5.94 (br, 1H), 5.56 (s, 1H), 5.24 (s, 2H), 3.03 (m, 3H), 2.42 (s, 3H), 1.98-1.89 (m, 1H), 0.99-0.91 (m, 2H), 0.76-0.66 (m, 2H)).

233

Example 16.3 and 16.4: 7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-2-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide and 7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-2-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




Step 1: methyl
2-amino-5-iodothiazole-4-carboxylate



To a solution of methyl 2-amino-1,3-thiazole-4-carboxylate (20 g, 0.13 mol) in dichloromethane (300 mL) was added N-iodosuccinimide (34 g, 0.15 mol) in portions. The resulting solution was stirred overnight at room temperature. Then the reaction mixture was washed with saturated aqueous Na_2SO_3 (3×150 mL). The organic layer was dried over anhydrous sodium sulfate. After filtration, the filtrate was concentrated to afford methyl 2-amino-5-iodo-1,3-thiazole-4-carboxylate (21 g, 58%) as a reddish solid.


Step 2: methyl 5-iodothiazole-4-carboxylate



234


To a solution of methyl 2-amino-5-iodo-1,3-thiazole-4-carboxylate (21 g, 0.074 mol) in tetrahydrofuran (400 mL) was added t-butyl nitrite (11.5 g, 0.11 mol). The resulting solution was stirred for 1 h at 50° C. in an oil bath. After cooled down to room temperature, the reaction mixture was concentrated under vacuum. The residue was purified by chromatography with ethyl acetate/petroleum ether (1:5) to afford methyl 5-iodo-1,3-thiazole-4-carboxylate (8 g, 40%) as a yellow solid. LCMS (ESI): $[\text{M}+\text{H}]^+=269.9$; $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 8.95 (s, 1H), 3.97 (s, 3H).

Step 3: methyl
5-(trifluoromethyl)thiazole-4-carboxylate



To a solution of methyl 5-iodo-1,3-thiazole-4-carboxylate (8.00 g, 29.7 mmol) and copper iodide (8.70 g, 45.7 mmol) in N,N-dimethylformamide (200 mL) was added methyl 2,2-difluoro-2-(fluorosulfonyl)acetate (8.70 g, 45.3 mmol). The resulting solution was stirred overnight at 80° C. in an oil bath. After filtration remove solids and concentration under vacuum, the residue was purified by chromatography with ethyl acetate/petroleum ether (1:5) to afford methyl 5-(trifluoromethyl)-1,3-thiazole-4-carboxylate (4 g, 64%) as a light yellow solid. LCMS (ESI): $[\text{M}+\text{H}]^+=212.0$; $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 8.91 (s, 1H), 4.00 (s, 3H).


Step 4:
N-ethyl-5-(trifluoromethyl)thiazole-4-carboxamide



A mixture of methyl 5-(trifluoromethyl)-1,3-thiazole-4-carboxylate (1 g, 4.74 mmol) and ethylamine in ethanol (10 mL) was placed in a 30-mL sealed tube. The resulting solution was stirred overnight at 50° C. in an oil bath. After concentration under vacuum, the residue was purified by chromatography with ethyl acetate/petroleum ether (1:5) to afford N-ethyl-5-(trifluoromethyl)-1,3-thiazole-4-carboxamide (400 mg, 38%) as a light yellow oil. LCMS (ESI): $[\text{M}+\text{H}]^+=225.0$; $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 8.79 (s, 1H), 7.42 (br, 1H), 3.55-3.46 (m, 2H), 1.27 (m, 3H).


235

Step 5: 2-bromo-N-ethyl-5-(trifluoromethyl)thiazole-4-carboxamide



To a solution of N-ethyl-5-(trifluoromethyl)-1,3-thiazole-4-carboxamide (200 mg, 0.89 mmol) in tetrahydrofuran (3 mL) was added n-butyllithium (2.5 M in hexanes; 1.1 mL, 2.70 mmol) drop wise with stirring at -78°C . The resulting solution was stirred for 30 min at -78°C . Then carbon tetrabromide (900 mg, 2.70 mmol) in tetrahydrofuran (5 mL) was added drop wise with stirring at -78°C . After stirred 1 h at -78°C , the reaction was quenched with water/ice, extracted with dichloromethane (2x30 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1:10) to afford 2-bromo-N-ethyl-5-(trifluoromethyl)-1,3-thiazole-4-carboxamide (40 mg, 15%) as a light yellow solid. LCMS (ESI): $[M+H]^+=302.8, 304.8$; $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 3.53-3.44 (m, 2H), 1.27 (m, 3H).


Step 6: 2-amino-N-ethyl-5-(trifluoromethyl)thiazole-4-carboxamide



A mixture of 2-bromo-N-ethyl-5-(trifluoromethyl)-1,3-thiazole-4-carboxamide (500 mg, 1.65 mmol), 1,4-dioxane (5 mL) and ammonia (5 mL) was placed in a 30-mL sealed tube. The resulting solution was stirred overnight at 70°C in an oil bath. The resulting mixture was extracted with CH_2Cl_2 (20 mLx2), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1:1) to afford 2-amino-N-ethyl-5-(trifluoromethyl)-1,3-thiazole-4-carboxamide (250 mg, 63%) as a light yellow semi-solid. LCMS (ESI): $[M+H]^+=240.0$; $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 7.15 (br, 1H), 5.24 (br, 2H), 3.48-3.39 (m, 2H), 1.22 (m, 3H).



236

Step 7: 7-(chloromethyl)-N-ethyl-5-oxo-2-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide



To a mixture of 2-amino-N-ethyl-5-(trifluoromethyl)-1,3-thiazole-4-carboxamide (250 mg, 1.05 mmol) and ethyl 4-chloro-3-oxobutanoate (350 mg, 2.13 mmol) was added polyphosphoric acid (5 g, excess). The resulting mixture was stirred for 1.5 h at 110°C in an oil bath. The reaction was then quenched by the addition of water and the pH value of the solution was adjusted to pH 8 with aqueous sodium hydroxide. The resulting mixture was extracted with dichloromethane (100 mLx2), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by silica gel chromatography with dichloromethane/ethyl acetate (10:1) to afford 7-(chloromethyl)-N-ethyl-5-oxo-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (200 mg, 56%) as a white solid. LCMS (ESI): $[M+H]^+=339.9$; $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 6.49 (s, 1H), 5.90 (br, 1H), 4.41 (s, 2H), 3.60-3.51 (m, 2H), 1.22 (m, 3H).

Step 8: 7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-2-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide (Example 16.3) and 7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-2-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide (Example 16.4)



To a solution of 7-(chloromethyl)-N-ethyl-5-oxo-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (80 mg, 0.24 mmol) and potassium carbonate (80 mg,


237

2.00 equiv) in acetonitrile (8 mL) was added 5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazole (from Example 15.9, Step 2) (50 mg, 0.28 mmol). The resulting solution was stirred for 3 h at 80° C. in an oil bath. After filtration to remove solids and concentration under vacuum, the residue was purified by chromatography with ethyl acetate/petroleum ether (1:2) to afford 7-[[5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (26.3 mg, 23%) as a white solid (LCMS (ESI): $[M+H]^+=480.0$; 1H NMR (300 MHz, $CDCl_3$) δ 6.20 (s, 1H), 5.89 (bs, 1H), 5.77 (s, 1H),



238

5.35 (s, 2H), 3.59-3.50 (m, 2H), 1.73-1.61 (m, 1H), 1.22 (m, 3H), 1.07-1.00 (m, 2H), 0.76-0.70 (m, 2H)) and 7-[[3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (16.5 mg, 15%) as a white solid (LCMS (ESI): $[M+H]^+=480.0$; 1H NMR (300 MHz, $CDCl_3$) δ 6.38 (s, 1H), 5.84 (bs, 1H), 5.67 (s, 1H), 5.25 (s, 2H), 3.59-3.50 (m, 2H), 1.98-1.89 (m, 1H), 1.22 (m, 3H), 0.99-0.91 (m, 2H), 0.78-0.73 (m, 2H)).

The following example was prepared in a manner similar to Examples 16.1, 16.2, 16.3, and 16.4:

No.	Structure/Name	LCMS (M + H)	1H NMR
16.5		412.20	1H NMR (300 MHz, $CDCl_3$) δ 9.36 (br, 1H), 7.97 (s, 1H), 6.19 (s, 1H), 5.80 (s, 1H), 5.37 (s, 2H), 3.49-3.40 (m, 2H), 1.75-1.66 (m, 1H), 1.26-1.20 (m, 3H), 1.02-0.95 (m, 2H), 0.74-0.65 (m, 2H)
	7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide		


The following compounds were prepared using methods analogous to those described above:

No.	Structure/Name	LCMS (M + H)	1H NMR
16.6		452.0	1H NMR (300 MHz, $CDCl_3$) δ 6.17 (s, 1H), 5.98 (s, 1H), 5.65 (s, 1H), 5.31 (s, 2H), 3.57-3.48 (m, 2H), 2.18-2.11 (m, 1H), 1.69-1.62 (m, 1H), 1.30-1.22 (m, 3H), 1.19-1.13 (m, 2H), 1.01-0.92 (m, 2H), 0.87-0.81 (m, 2H), 0.72-0.67 (m, 2H)
	2-cyclopropyl-7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide		
16.7		452.0	1H NMR (300 MHz, $CDCl_3$) δ 6.34 (s, 1H), 5.84 (s, 1H), 5.54 (s, 1H), 5.21 (s, 2H), 3.57-3.49 (m, 2H), 2.20-2.13 (m, 1H), 1.96-1.89 (m, 1H), 1.30-1.25 (m, 3H), 1.19-1.12 (m, 2H), 0.98-0.90 (m, 2H), 0.87-0.81 (m, 2H), 0.79-0.71 (m, 2H)
	2-cyclopropyl-7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide		


239

Method 17

Example 17.1: 2-methyl-5-oxo-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile



Step 1: 3-bromo-2-methyl-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 3-bromo-7-(chloromethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (from Example 4.1, Step 6) (300 mg, 0.76 mmol) in CH₃CN (5 mL) was added 3-(trifluoromethyl)-1H-pyrazole (125 mg, 0.92 mmol), potassium carbonate (316 mg, 2.29 mmol) and potassium iodide (63 mg, 0.38 mmol). The resulting solution was heated to reflux overnight. The solids were filtered out and the filtrate was

240

concentrated under vacuum. The residue was purified by silica gel chromatography with dichloromethane/methanol (50/1) to afford 3-bromo-2-methyl-7-[[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (200 mg, 67%) as a white solid.

Step 2: 2-methyl-5-oxo-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile

10


15

20

25



30

35



To a solution of 3-bromo-2-methyl-7-[[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (120 mg, 0.31 mmol) in N,N-dimethylformamide (2 mL) was added copper (I) cyanide (32 mg, 0.36 mmol). The resulting solution was stirred for 1 h at 100° C. The reaction was then quenched by the addition of water. The solids were filtered off and the filtrate was extracted with ethyl acetate (3×10 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/5) to afford 2-methyl-5-oxo-7-[[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carbonitrile (10.2 mg, 10%) as a white solid. LCMS (ESI): [M+H]⁺=340; ¹H NMR (300 MHz, CDCl₃) δ 7.59 (m, 1H), 6.62 (m, 1H), 5.99 (s, 1H), 5.24 (s, 2H), 2.67 (s, 3H).


The following examples were prepared in a manner similar to Example 17.1:

No.	Structure/Name	LCMS (M + H)	¹ H NMR
17.2	 7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile	380.0	¹ H NMR (300 MHz, CD ₃ OD) δ 6.34 (s, 1H), 5.89 (s, 1H), 5.44 (s, 2H), 2.66 (s, 3H), 1.91 (m, 1H), 1.04 (m, 2H), 0.77 (m, 2H)
17.3	 7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile	379.90	¹ H NMR (300 MHz, CD ₃ OD) δ 6.57 (s, 1H), 5.71 (s, 1H), 5.30 (s, 2H), 2.64 (s, 3H), 1.95 (m, 1H), 0.96 (m, 2H), 0.75 (m, 2H)


241

Method 18

Example 18.1: 3-(1-hydroxyethyl)-2-methyl-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one




Step 1: 4-bromopentane-2,3-dione




To a solution of pentane-2,3-dione (1.00 g, 9.99 mmol) in chloroform (30 mL) was added bromine (1.60 g, 10.01 mmol) and hydrogen bromide in acetic acid (33 wt %; 3 drops). The resulting solution was stirred for 3 h at 50° C. The resulting solution was concentrated to afford 4-bromopentane-2,3-dione as a solid (1.79 g). The crude product was used in next step without further purification.

Step 2: 1-(2-amino-5-methylthiazol-4-yl)ethanone



To a solution of 4-bromopentane-2,3-dione (1.79 g, 10.00 mmol) in ethanol (50 mL) was added thiourea (760 mg, 9.98 mmol). The resulting solution was stirred for 1 h at 100° C. and then cooled down room temperature. The mixture was filtered to afford 1-(2-amino-5-methyl-1,3-thiazol-4-yl)ethan-1-one as a off-white solid (1.2 g, 65%). LCMS (ESI): [M+H]⁺=157.0.


Step 3: 3-acetyl-7-(chloromethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



242


To a solution of 1-(2-amino-5-methyl-1,3-thiazol-4-yl)ethan-1-one (300 mg, 1.92 mmol) was added ethyl 4-chloro-3-oxobutanoate (474 mg, 2.88 mmol) and polyphosphoric acid (10 mL). The resulting solution was stirred for 1 h at 110° C. and then cooled down room temperature. The resulting solution was diluted with water (20 mL) and the pH value of the solution was adjusted to 8. The resulting solution was extracted and concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum (1/1) to afford 3-acetyl-7-(chloromethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a light yellow solid (70.0 mg, 12%). LCMS (ESI): [M+H]⁺=257.0; ¹H NMR (300 MHz, CDCl₃) δ 6.46 (s, 1H), 4.42 (s, 2H), 2.52 (s, 3H), 2.39 (s, 3H).

Step 4: 3-acetyl-2-methyl-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 3-acetyl-7-(chloromethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (200 mg, 0.78 mmol) in CH₃CN (25 mL) was added potassium iodide (68 mg, 0.39 mmol), potassium carbonate (220 mg, 1.59 mmol) and 3-(trifluoromethyl)-1H-pyrazole (160 mg, 1.18 mmol). The resulting solution was stirred for 12 h at 80° C. in an oil bath. The resulting mixture was quenched by water (10 mL), extracted with dichloromethane (3×20 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by chromatography with ethyl acetate/petroleum ether (1:2) to afford 3-acetyl-2-methyl-7-[[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (195 mg, 70%) as a yellow solid. LCMS (ESI): [M+H]⁺=357.1; ¹H NMR (300 MHz, CDCl₃) δ 7.61 (m, 1H), 6.62 (m, 1H), 5.91 (s, 1H), 5.25 (s, 2H), 2.45 (s, 3H), 2.38 (s, 3H).

Step 5: 3-(1-hydroxyethyl)-2-methyl-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 3-acetyl-2-methyl-7-[[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (30 mg, 0.08 mmol) in methanol (10 mL) was added sodium borohydride (13 mg, 0.34 mmol). The resulting solution was stirred for 12 h at room temperature in an oil bath. The reaction was then quenched by aqueous ammonium chloride (10 ml), extracted with dichloromethane



243

(3×20 mL), washed with brine, dried over anhydrous magnesium sulfate and concentrated under vacuum. The residue was purified by chromatography with ethyl acetate/petroleum ether (1:1) to afford 3-(1-hydroxyethyl)-2-methyl-7-[[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (11.7 mg, 39%) as a light



244

yellow solid. LCMS (ESI): $[M+H]^+=359.0$; $^1\text{H NMR}$ (300 MHz, CD_3OD) δ 7.94-7.93 (m, 1H), 6.70 (m, 1H), 5.97 (s, 1H), 5.47-5.40 (m, 1H), 5.38 (m, 2H), 2.48 (s, 3H), 1.52 (m, 3H).

The following examples were prepared in a manner similar to Example 18.1:

No.	Structure/Name	LCMS (M + H)	$^1\text{H NMR}$
18.2	 <p>3-acetyl-7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	397.0	$^1\text{H NMR}$ (300 MHz, CD_3OD) δ 6.33 (s, 1H), 5.83 (s, 1H), 5.44 (s, 2H), 2.41 (s, 3H), 2.37 (s, 3H), 1.92 (m, 1H), 1.02 (m, 2H), 0.76 (m, 2H)
18.3	 <p>3-acetyl-7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	397.0	$^1\text{H NMR}$ (300 MHz, CD_3OD) δ 6.58 (s, 1H), 5.69 (s, 1H), 5.32 (s, 2H), 2.41 (s, 3H), 2.37 (s, 3H), 1.98 (m, 1H), 0.98 (m, 2H), 0.78 (m, 2H)


The following compounds were prepared using methods analogous to those described above:

No.	Structure/Name	LCMS (M + H)	$^1\text{H NMR}$
18.4	 <p>3-(2-hydroxypropan-2-yl)-2-methyl-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	373.10	$^1\text{H NMR}$ (300 MHz, CDCl_3) δ 7.60 (m, 1H), 7.15 (m, 1H), 6.61 (m, 1H), 5.98 (s, 1H), 5.26 (s, 2H), 2.52 (s, 3H), 1.72 (s, 6H)
18.5	 <p>7-((5-fluoro-3-methyl-1H-indazol-1-yl)methyl)-3-(1-hydroxyethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	373	$^1\text{H NMR}$ (300 MHz, CDCl_3): δ 7.35-7.25 (m, 2H), 7.19-7.12 (m, 1H), 5.72 (s, 1H), 5.42 (s, 2H), 5.04 (m, 1H), 2.56 (s, 3H), 2.41 (s, 3H), 1.60 (m, 3H)


245

Method 19

Example 19.1: 7-(1-(5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)ethyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



Step 1: 3-(trans-2-((tert-butyldimethylsilyloxy)methyl)cyclopropyl)-7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 7-[[5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 15.11) (120 mg, 0.28 mmol) in dichloromethane (30 mL) was added tert-butyldimethylsilylchloride (177 mg, 1.18 mmol), imidazole (80 mg, cat.) and 4-dimethylaminopyridine (5 mg, cat.). The resulting solution was stirred for overnight at 50° C. After cooling down to room temperature, the reaction mixture was concentrated under vacuum. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:2) to afford 3-(2-[[tert-butyldimethylsilyloxy]methyl]cyclopropyl)-7-[[5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (120 mg, 67%) as yellow oil. LCMS (ESI): [M+H]⁺=539.0. The crude product was used in next step without further purification.

246

Step 2: 3-(trans-2-((tert-butyldimethylsilyloxy)methyl)cyclopropyl)-7-(1-(5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)ethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one

5

10

15

20

25

30

35

40


45

50

55


60

65



To a solution of 3-(2-[[tert-butyldimethylsilyloxy]methyl]cyclopropyl)-7-[[5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (120 mg, 0.22 mmol) in tetrahydrofuran (20 mL) was added n-butyllithium (1.5 mL, 85% in hexanes) and methyl iodide (158 mg, 1.11 mmol). The resulting solution was stirred overnight at room temperature. The reaction was then quenched with water (20 mL), extracted with dichloromethane (3x30 mL), washed with brine, dried over sodium sulfate and concentrated under vacuum. The residue was then purified by chromatography with ethyl acetate/petroleum ether (1:2) to afford 3-(trans-2-[[tert-butyldimethylsilyloxy]methyl]cyclopropyl)-7-[1-(5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)ethyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (50 mg, 35%) as a light yellow solid. LCMS (ESI): [M+H]⁺=553.0.

Step 3: 7-(1-(5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)ethyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 3-(trans-2-[[tert-butyldimethylsilyloxy]methyl]cyclopropyl)-7-[1-(5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)ethyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (50 mg, 0.09 mmol) in ethanol (30 mL) was added 0.5% hydrogen chloride in ethanol (10 mL). The resulting solution was stirred for 2 h at room temperature. After cooling down to room temperature, the reaction mixture was concentrated under vacuum. The residue was purified by chromatography with dichloromethane/methanol (100:1) to afford 7-[1-(5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)ethyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


247

(14.3 mg, 34%) as a white solid. LCMS (ESI): $[M+H]^+$ = 439.0; $^1\text{H NMR}$ (300 MHz, CD_3OD) δ 6.18 (s, 2H), 6.17 (s, 1H), 5.59-5.54 (m, 1H), 4.05-4.01 (m, 1H), 3.14-3.07 (m, 1H), 2.41 (s, 3H), 2.28-2.22 (m, 1H), 1.96-1.93 (m, 3H), 1.69-1.64 (m, 1H), 1.27-1.23 (m, 1H), 1.03-0.72 (m, 4H), 0.73-0.62 (m, 2H).

The following compound was prepared using methods analogous to those described above:


248

with brine (1x30 mL), dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:2) to afford 3-([3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile (102 mg, 17%) as a brown solid. LCMS (ESI): $[M+H]^+$ = 360.0, 362.0; $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 7.65-7.41 (m, 4H), 6.04 (s, 1H), 3.91 (s, 2H), 2.35 (s, 3H).


No.	Structure/Name	LCMS (M + H)	$^1\text{H NMR}$
19.2		439.0	$^1\text{H NMR}$ (300 MHz, CD_3OD) δ 6.18 (s, 1H), 5.73-5.70 (m, 1H), 5.60-5.57 (m, 1H), 4.03-3.96 (m, 1H), 3.42-3.35 (m, 1H), 2.41 (s, 3H), 1.98-1.92 (m, 4H), 1.71-1.63 (m, 1H), 2.52 (s, 3H), 1.45-1.40 (m, 1H), 1.06-0.92 (m, 3H), 0.73-0.65 (m, 3H)
	7-(1-(5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)ethyl)-3-(2-(hydroxymethyl)-1-methylcyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one		

Method 20

Example 20.1: 3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile




Step 1: 3-((3-bromo-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile



To a solution of 3-bromo-7-(chloromethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 4.1, Step 6) (500 mg, 1.70 mmol) in 1,4-dioxane/ H_2O (3/1 mL) was added tetrakis(triphenylphosphine)palladium (198 mg, 0.17 mmol), potassium phosphate (726 mg, 3.42 mmol) and (3-cyanophenyl)boronic acid (302 mg, 2.06 mmol). The resulting solution was stirred overnight at 80° C. After cooling down to room temperature, the resulting mixture was extracted with dichloromethane (3x30 mL), washed


Step 2: 3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile




To a solution of 3-((3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile (80 mg, 0.22 mmol) in $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ (3/1 mL) was added [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (16.7 mg, 0.02 mmol), sodium carbonate (47.2 mg, 0.45 mmol) and potassium trans-2-(hydroxymethyl)cyclopropyltrifluoroborate (from Example 4.1, Step 2) (79.3 mg, 0.45 mmol). The reaction mixture was irradiated in a microwave for 1.5 h at 120° C. The resulting mixture was extracted with dichloromethane (3x20 mL), washed with brine (10 mL), dried over sodium sulfate. After concentration under vacuum, the residue was purified on a silica gel column eluted with dichloromethane/methanol (50:1) to afford 3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile (16.6 mg, 21%) as a white solid. LCMS (ESI): $[M+H]^+$ = 352.0; $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 7.62-7.58 (m, 3H), 7.50-7.44 (m, 1H), 6.02 (s, 1H), 4.04-4.00 (m, 1H), 3.99 (s, 2H), 3.20-3.13 (m, 1H), 2.42 (s, 3H), 2.26-2.25 (m, 1H), 1.28-1.25 (m, 1H), 1.05-1.00 (m, 2H).

249

Example 20.2: 3-((6-fluoro-3-(2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile



Step 1: 3-((3-bromo-6-fluoro-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile




To a solution of 3-bromo-7-(chloromethyl)-6-fluoro-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 4.29, Step 1) (130 mg, 0.42 mmol) in 1,4-dioxane/H₂O (3/1, 4 mL) added tetrakis(triphenylphosphine)palladium (48 mg, 0.04 mmol), potassium phosphate (179 mg, 0.84 mmol), and (3-cyanophenyl)boronic acid (74 mg, 0.50 mmol). The resulting solution was stirred overnight at 80° C., then the resulting mixture was concentrated under vacuum and the residue was purified by chromatography with ethyl acetate/petroleum ether (1:3) to afford 3-((3-

250

bromo-6-fluoro-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile (36 mg, 23%) as a light yellow solid. LCMS (ESI): [M+H]⁺=377.9.

5 Step 2: 3-((6-fluoro-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile





To a solution of 3-((3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzonitrile (100 mg, 0.26 mmol) in CH₃CN/H₂O (3/1 mL) under inert nitrogen atmosphere was added [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (20 mg, 0.03 mmol), sodium carbonate (56.2 mg, 0.53 mmol), and potassium trans-2-(hydroxymethyl)cyclopropyltrifluoroborate (from Example 4.1, Step 2) (94.4 mg, 0.53 mmol). The resulting solution was stirred for 90 min at 120° C. The mixture was concentrated under vacuum, and the residue was purified by chromatography with dichloromethane/methanol (50:1) to afford 2-fluoro-3-((3-[(trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile as a off-white solid (9.9 mg, 10%). LCMS (ESI): [M+H]⁺=370.0; ¹H NMR (300 MHz, CDCl₃) δ 7.57 (s, 1H), 7.55-7.52 (m, 2H), 7.41-7.40 (m, 1H), 4.06-3.96 (m, 3H), 3.22-3.17 (m, 1H), 2.37 (s, 3H), 2.30-2.10 (m, 1H), 1.34-1.25 (m, 1H), 1.06-0.98 (m, 2H).




The following examples were prepared in a manner similar to Example 20.1 and 20.2:

No.	Structure/Name	LCMS (M + H)	¹ H NMR
20.3	<p>2-fluoro-3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile</p>	370.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.64-7.57 (m, 2H), 7.29-7.24 (m, 1H), 6.07 (s, 1H), 4.09-4.04 (m, 1H), 3.99 (s, 2H), 3.18-3.11 (m, 1H), 2.41 (s, 3H), 2.32-2.26 (m, 1H), 1.29-1.24 (m, 1H), 1.06-0.99 (m, 2H)
20.4		378.1	¹ H NMR (300 MHz, CDCl ₃) δ 9.28 (br, 1H), 8.48 (br, 1H), 8.11-8.09 (m, 1H), 8.02-7.99 (m, 1H), 7.86-7.81 (m, 1H), 7.75-7.70 (m, 1H), 5.99 (s, 1H), 4.36 (s, 2H), 4.08-4.00 (m, 2H), 3.09-3.02 (m, 1H), 2.36 (s, 3H), 2.26-2.21 (m, 1H), 1.28-1.20 (m, 1H), 1.03-0.97 (m, 2H)





-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
20.5	 <p>3-(trans-2-(hydroxymethyl)cyclopropyl)-7-(isoquinolin-4-ylmethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	395.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.53-7.41 (m, 4H), 6.02 (s, 1H), 4.14-4.02 (m, 1H), 3.92 (s, 2H), 3.13-3.05 (m, 1H), 2.39 (s, 3H), 2.37-2.26 (m, 1H), 1.30-1.20 (m, 1H), 1.04-0.95 (m, 2H)
	 <p>3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-7-(3-(trifluoromethyl)benzyl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>		


The following compounds were prepared using methods analogous to those described above:

No.	Structure/Name	LCMS (M + H)	¹ H NMR
20.6	 <p>3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-4-methylbenzotrile</p>	366.00	¹ H NMR (300 MHz, CD ₃ OD) δ 7.54 (m, 2H), 7.36 (m, 1H), 5.94 (s, 1H), 3.97 (s, 2H), 3.59 (m, 2H), 2.42 (s, 3H), 2.41 (s, 3H), 2.15 (m, 1H), 1.32 (m, 1H), 1.01 (m, 2H)
20.7	 <p>4-fluoro-3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile</p>	379.95	¹ H NMR (300 MHz, CDCl ₃) δ 7.62-7.57 (m, 2H), 7.19-7.15 (m, 1H), 6.05 (s, 1H), 4.17-4.05 (m, 2H), 3.90 (s, 2H), 3.12-3.07 (m, 1H), 2.38 (s, 3H), 2.28-2.26 (m, 1H), 1.30-1.24 (m, 1H), 1.04-0.98 (m, 2H)
20.8	 <p>3-fluoro-5-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile</p>	370.10	¹ H NMR (300 MHz, CDCl ₃) δ 7.42 (s, 1H), 7.30-7.28 (m, 2H), 6.10 (s, 1H), 4.10-4.05 (m, 1H), 3.91 (s, 2H), 3.18-3.11 (m, 1H), 2.41 (s, 3H), 2.35-2.25 (m, 1H), 1.35-1.25 (m, 1H), 1.07-1.00 (m, 2H)

-continued


No.	Structure/Name	LCMS (M + H)	¹ H NMR
20.9	 <p>2-fluoro-5-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile</p>	370.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.59-7.48 (m, 2H), 7.20-7.14 (m, 1H), 6.06 (s, 1H), 4.14-4.03 (m, 1H), 3.85 (s, 2H), 3.10 (m, 1H), 2.39 (s, 3H), 2.31-2.24 (m, 1H), 1.26-1.23 (m, 1H), 1.06-1.01 (m, 2H)
20.10	 <p>3-[[3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]-4-methoxy-benzotrile</p>	382.0	¹ H NMR (400 MHz, CDCl ₃) δ 7.60-7.58 (m, 1H), 7.50 (s, 1H), 6.94-6.92 (m, 1H), 5.96 (s, 1H), 4.06-4.03 (m, 1H), 3.89-3.85 (m, 5H), 3.13-3.07 (m, 1H), 2.38 (s, 3H), 2.29-2.26 (m, 1H), 1.28-1.25 (m, 2H), 1.05-0.88 (m, 2H)
20.11	 <p>3-(2-(hydroxymethyl)cyclopropyl)-2-methyl-7-(4-(trifluoromethyl)benzyl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	395.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.60-7.59 (m, 2H), 7.40-7.37 (m, 2H), 6.04 (s, 1H), 4.21-4.18 (m, 1H), 4.07-4.01 (m, 1H), 3.91 (s, 2H), 3.11-3.04 (m, 1H), 2.39 (s, 3H), 2.36-2.24 (m, 1H), 1.25-1.19 (m, 1H), 1.04-0.99 (m, 2H)
20.12	 <p>4-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)picolinotrile</p>	352.9	¹ H NMR (300 MHz, CDCl ₃) δ 8.62-8.61 (m, 1H), 7.86 (m, 1H), 7.64-7.62 (m, 1H), 6.21 (s, 1H), 3.99 (s, 2H), 3.63-3.57 (m, 2H), 2.41 (s, 3H), 2.20-2.11 (m, 1H), 1.40-1.28 (m, 1H), 1.05-0.99 (m, 2H)

-continued


No.	Structure/Name	LCMS (M + H)	¹ H NMR
20.13	 <p>4-cyclopropyl-3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile</p>	392.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.47-7.45 (m, 1H), 7.40-7.22 (m, 2H), 5.90 (s, 1H), 4.19 (s, 2H), 4.06-3.98 (m, 1H), 3.14-3.07 (m, 1H), 2.45 (s, 3H), 2.31-2.26 (m, 1H), 1.95-1.86 (m, 1H), 1.30-1.26 (m, 1H), 1.04-0.99 (m, 4H), 0.69-0.68 (m, 2H)

Method 21

Example 21.1: 7-(3-cyanobenzyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




Step 1: 7-(3-cyanobenzyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide








7-(Chloromethyl)-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (from Example 16.1, Step 5) (50 mg, 0.18 mmol), (3-cyanophenyl)boronic acid (56 mg, 0.38 mmol), tetrakis(triphenylphosphine)palladium (20 mg, 0.019 mmol), potassium phosphate (80 mg, 0.38 mmol), 1,4-dioxane (1.5 mL) and water (0.5 mL) were placed in a 10-mL sealed tube. The resulting solution was stirred for 2 h at 80° C. in an oil bath. After cooling down to room temperature, the resulting mixture was extracted with CH₂Cl₂ (20 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with dichloromethane/methanol (20:1) to afford 7-[(3-cyanophenyl)methyl]-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (29.1 mg, 47%) as a white solid. LCMS (ESI): [M+H]⁺ = 339.0; ¹H NMR (300 MHz, CDCl₃) δ 7.61-7.40 (m, 4H), 6.08 (s, 1H), 5.96 (br, 1H), 3.90 (s, 2H), 3.05 (m, 3H), 2.41 (s, 3H).






The following examples were prepared in a manner similar to Example 21.1:

No.	Structure/Name	LCMS (M + H)	¹ H NMR
21.2	 <p>7-(3-cyano-2-fluorobenzyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	356.9	¹ H NMR (300 MHz, DMSO) δ 8.32 (m, 1H), 7.84 (m, 2H), 7.38 (m, 1H), 6.13 (s, 1H), 4.00 (s, 2H), 2.72 (s, 3H), 2.28 (s, 3H)






-continued

No.	Structure/Name	LCMS (M + H)	¹ H NMR
21.3	 <p data-bbox="313 573 667 636">7-(3-chloro-2-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	380.10	¹ H NMR (300 MHz, CDCl ₃) δ 7.34-7.29 (m, 1H), 7.17-7.12 (m, 1H), 7.07-7.02 (m, 1H), 6.04 (s, 1H), 5.88 (br, 1H), 3.94 (s, 2H), 3.56-3.47 (m, 2H), 2.40 (s, 3H), 1.30-1.25 (m, 3H)
21.4	 <p data-bbox="313 873 667 936">N-ethyl-7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	400.0	¹ H NMR (300 MHz, CD ₃ OD) δ 7.63 (m, 2H), 7.32 (m, 1H), 6.14 (s, 1H), 4.05 (s, 2H), 3.92 (s, 3H), 2.38 (s, 3H)
21.5	 <p data-bbox="313 1169 667 1232">N-ethyl-7-(2-fluoro-3-methylbenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	360.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.30-7.26 (m, 1H), 7.12-6.96 (m, 3H), 6.03 (s, 1H), 6.00 (s, 1H), 3.91 (s, 2H), 3.55-3.46 (m, 2H), 2.37 (s, 3H), 2.26 (s, 3H), 1.29-1.24 (m, 3H)
21.6	 <p data-bbox="313 1524 667 1587">7-(2-chloro-5-cyanobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	386.7	¹ H NMR (300 MHz, CDCl ₃) δ 7.59 (s, 1H), 7.50 (s, 2H), 6.05 (s, 1H), 5.82 (m, 1H), 4.27 (s, 2H), 3.55-3.51 (m, 3H), 2.44 (s, 3H), 1.30-1.26 (m, 3H)
21.7	 <p data-bbox="334 1877 651 1940">N-ethyl-2-methyl-5-oxo-7-(3-(trifluoromethyl)benzyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	396.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.43-7.61 (m, 4H), 6.06 (s, 1H), 5.81 (br, 1H), 3.93 (s, 2H), 3.48-3.57 (m, 2H), 2.42 (s, 3H), 1.28 (m, 3H)

-continued


No.	Structure/Name	LCMS (M + H)	¹ H NMR
21.8	 <p>7-(3-cyanobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	353.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.56-7.40 (m, 4H), 6.08 (s, 1H), 5.83 (s, 1H), 3.90 (s, 2H), 3.53 (m, 2H), 2.42 (s, 3H), 1.28 (m, 3H)
21.9	 <p>7-(3-cyano-2-fluorobenzyl)-2-cyclopropyl-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	397.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.56-7.49 (m, 2H), 6.06 (s, 1H), 5.85 (s, 1H), 3.93 (s, 2H), 3.58-3.45 (m, 2H), 2.18-2.09 (m, 1H), 1.18-1.15 (m, 3H), 1.13-1.11 (m, 2H), 0.85-0.79 (m, 2H)
21.10	 <p>7-(3-cyano-5-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	371.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.37 (s, 1H), 7.23-7.22 (m, 2H), 6.10 (s, 1H), 5.82-5.73 (m, 1H), 3.89 (s, 2H), 3.58-3.50 (m, 2H), 2.43 (s, 3H), 1.31-1.26 (m, 3H)
21.11	 <p>7-(3-cyanobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	339.0	¹ H NMR (300 MHz, CD ₃ OD) δ 7.36-7.29 (m, 1H), 7.13-6.94 (m, 3H), 6.15 (s, 1H), 3.93 (s, 2H), 3.41 (m, 2H), 2.40 (s, 3H), 1.24 (m, 3H)
21.12	 <p>N-ethyl-7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	414.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.53 (m, 2H), 7.23 (m, 1H), 6.05 (s, 1H), 5.88 (s, 1H), 3.96 (s, 2H), 3.52 (m, 2H), 2.40 (s, 3H), 1.27 (m, 3H)

-continued


No.	Structure/Name	LCMS (M + H)	¹ H NMR
21.13	 <p>7-(3-cyano-2-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	371.0	¹ H NMR (300 MHz, CDCl ₃) δ 7.61-7.50 (m, 2H), 7.26-7.20 (m, 1H), 6.07 (s, 1H), 5.85 (s, 1H), 3.95 (s, 2H), 3.57-3.48 (m, 2H), 2.42 (s, 3H), 1.28 (m, 3H)
21.14	 <p>7-[(3-chloro-2-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	366.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.30 (q, J = 4.6 Hz, 1H), 7.53-7.45 (m, 1H), 7.39-7.31 (m, 1H), 7.24-7.16 (m, 1H), 6.08 (s, 1H), 2.73 (dd, J = 4.7, 0.9 Hz, 3H), 2.29 (d, J = 0.9 Hz, 3H).
21.15	 <p>N,2-dimethyl-5-oxo-7-[[3-(trifluoromethyl)phenyl]methyl]thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	382.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.29 (d, J = 4.8 Hz, 1H), 7.68 (s, 1H), 7.65-7.52 (m, 3H), 6.19 (s, 1H), 3.98 (s, 2H), 2.73 (d, J = 4.6 Hz, 3H), 2.29 (s, 3H).
21.16	 <p>7-[(3-chlorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	348.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.29 (q, J = 4.6 Hz, 1H), 7.37 (t, J = 1.9 Hz, 1H), 7.34 (d, J = 7.4 Hz, 1H), 7.28 (ddt, J = 11.5, 7.4, 1.6 Hz, 2H), 6.16 (s, 1H), 3.88 (s, 2H), 2.73 (d, J = 4.7 Hz, 3H), 2.29 (s, 3H).
21.17	 <p>7-[[2-cyclopropyl-5-(trifluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	422.1	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.37-8.29 (m, 1H), 7.59 (d, J = 2.0 Hz, 1H), 7.56-7.49 (m, 1H), 7.17 (d, J = 8.1 Hz, 1H), 5.99 (d, J = 0.7 Hz, 1H), 4.18 (s, 2H), 2.73 (d, J = 4.7 Hz, 3H), 2.29 (s, 3H), 2.11-2.03 (m, 1H), 1.02-0.90 (m, 2H), 0.76-0.64 (m, 2H).

263

Example 21.18: N-ethyl-2-methyl-5-oxo-7-((6-(trifluoromethyl)pyridine-2-yl)methyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




Step 1: Bis(pinacolato)diboron



To a solution of 2-bromo-6-(trifluoromethyl)pyridine (500 mg, 2.21 mmol) in 1,4-dioxane (10 mL) under nitrogen, was added potassium acetate (862 mg, 8.78 mmol), [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (330 mg, 0.45 mmol), 4,4,5,5-tetramethyl-2-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (609 mg, 2.40 mmol). The resulting solution was stirred for 12 h at 90° C. The mixture was filtered to remove solids and concentrated under vacuum to afford [6-(trifluoromethyl)pyridine-2-yl]boronic acid (500 mg, crude) as a black solid. The crude product was used in the next step without further purification. LCMS (ESI): [M+H]⁻=191.9.

Step 2: N-ethyl-2-methyl-5-oxo-7-((6-(trifluoromethyl)pyridine-2-yl)methyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




To a solution of 7-(chloromethyl)-N-ethyl-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (prepared in a similar manner to Example 16.1, Step 5) (50 mg, 0.17 mmol) in 1,4-dioxane (2 mL) under nitrogen was added sodium carbonate (36 mg, 0.34 mmol), [6-(trifluoromethyl)pyridine-2-yl]boronic acid (50 mg, 0.26 mmol), [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (13 mg, 0.02 mmol, and water (0.2 mL). The reaction mixture was irradiated with microwave radiation for 20 min at 120° C. The resulting solution was then extracted with ethyl acetate (3×20 mL), washed with brine, dried over anhydrous sodium sulfate, and concentrated


264

under vacuum. The residue was purified by chromatography with dichloromethane/methanol (20:1) to afford N-ethyl-2-methyl-5-oxo-7-[[6-(trifluoromethyl)pyridine-2-yl]methyl]-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (22.8 mg, 33%) as a yellow solid. LCMS (ESI): [M+H]⁺=397.1; ¹H NMR (400 MHz, CDCl₃) δ 7.81 (m, 1H), 7.60 (m, 1H), 7.53 (m, 1H), 6.26 (s, 1H), 5.89 (s, 1H), 4.15 (s, 2H), 3.51 (m, 2H), 2.41 (s, 3H), 1.28 (m, 3H).

Example 21.19: N-ethyl-7-(2-ethyl-4,5-difluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




Step 1: 1-(benzyloxy)-2-bromo-4,5-difluorobenzene



To a mixture of 2-bromo-4,5-difluorophenol (500 mg, 2.39 mmol, 1.00 equiv) and potassium carbonate (800 mg, 5.80 mmol) in CH₃CN (10 mL) was added (bromomethyl)benzene (610 mg, 3.57 mmol). The resulting solution was stirred for 3 h at 80° C. in an oil bath. After filtration to remove solids and concentration, the residue was purified by chromatography with ethyl acetate/petroleum ether (1:30) to afford 1-(benzyloxy)-2-bromo-4,5-difluorobenzene (600 mg, 84%) as colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.46-7.32 (m, 5H), 6.82-6.76 (m, 1H), 5.10 (s, 2H).

Step 2: 1-(benzyloxy)-4,5-difluoro-2-vinylbenzene




1-(Benzyloxy)-2-bromo-4,5-difluorobenzene (560 mg, 1.87 mmol), tetrakis(triphenylphosphine)palladium (200 mg, 0.19 mmol), potassium phosphate (800 mg, 3.78 mmol),

265

1,4-dioxane (10 mL), water (1 mL) and 2-ethenyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (580 mg, 3.77 mmol) were placed in a 30-mL sealed tube. The reaction mixture was stirred for 5 h at 80° C. in an oil bath. After filtration to remove solids and concentration, the residue was purified by chromatography with ethyl acetate/petroleum ether (1:10) to afford 1-(benzyloxy)-2-ethenyl-4,5-difluorobenzene (350 mg, 76%) as light yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 7.69-7.29 (m, 6H), 7.07-6.95 (m, 1H), 6.81-6.71 (m, 1H), 5.69-5.63 (m, 1H), 5.29-5.25 (m, 1H), 5.04 (s, 2H).

Step 3: 2-ethyl-4,5-difluorophenol



To a mixture of 1-(benzyloxy)-2-ethenyl-4,5-difluorobenzene (350 mg, 1.42 mmol) in methanol (10 mL) was added palladium on carbon (20 mg). The resulting reaction was stirred overnight at room temperature under hydrogen atmosphere (1 atm). After filtration to remove catalyst, the filtrate was concentrated under vacuum to afford 2-ethyl-4,5-difluorophenol (200 mg, 89%) as light yellow oil. The crude product was used in the next step without further purification.


Step 4: 2-ethyl-4,5-difluorophenyl trifluoromethanesulfonate



To a mixture of 2-ethyl-4,5-difluorophenol (200 mg, 1.26 mmol) and triethylamine (260 mg, 2.58 mmol) in dichloromethane (10 mL) was added trifluoroacetic anhydride (680 mg, 2.42 mmol) dropwise with stirring at 0° C. The resulting solution was stirred for 30 min at 0° C. The reaction was then quenched by the addition of water, extracted with dichloromethane (10 mL×2), washed with brine, dried over anhydrous sodium sulfate and concentrated in vacuo to afford 2-ethyl-4,5-difluorophenyl trifluoromethanesulfonate (180 mg, 50%) as a light yellow oil. The crude product was used in next step without further purification.


266

Step 5: 2-(2-ethyl-4,5-difluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane



To a mixture of 2-ethyl-4,5-difluorophenyl trifluoromethanesulfonate (180 mg, 0.62 mmol), [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (50 mg, 0.07 mmol), potassium acetate (120 mg, 1.23 mmol) in 1,4-dioxane (5 mL) was added 4,4,5,5-tetramethyl-2-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (240 mg, 0.93 mmol). The resulting mixture was stirred overnight at 90° C. in an oil bath under nitrogen atmosphere. After filtration to remove solids and concentration, the residue was purified by chromatography with ethyl acetate/petroleum ether (1:10) to afford 2-(2-ethyl-4,5-difluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (140 mg, 84%) as a light yellow oil. The crude product was used in next step without further purification.

Step 6: N-ethyl-7-(2-ethyl-4,5-difluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




7-(Chloromethyl)-N-ethyl-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (prepared in a manner similar to Example 16.1, Step 5) (100 mg, 0.35 mmol, 1.00 equiv), 2-(2-ethyl-4,5-difluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (140 mg, 0.52 mmol, 1.50 equiv), [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (30 mg, 0.10 equiv), sodium carbonate (75 mg, 2.00 equiv), 1,4-dioxane (3 mL) and water (0.3 mL) were placed in a 10 mL sealed tube. The reaction mixture was irradiated with microwave radiation for 30 min at 120° C. The resulting solution was diluted with 30 mL of dichloromethane, washed with 2×10 mL of brine, dried over sodium sulfate and concentrated in vacuo. The residue was purified by silica gel chromatography with dichloromethane/methanol (40:1) to afford N-ethyl-7-[(2-ethyl-4,5-difluorophenyl)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (17.9 mg, 13%) as an off-white solid. LCMS (ESI): [M+H]⁺=392.0; ¹H NMR (300 MHz,


267

CDCl_3) δ 7.04-6.91 (m, 2H), 5.88 (s, 1H), 5.82 (br, 1H), 3.84 (s, 2H), 3.58-3.48 (m, 2H), 2.59-2.52 (m, 2H), 2.42 (s, 3H), 1.28 (m, 3H), 1.17 (m, 3H).

Example 21.20: 7-[[3-(difluoromethyl)-2-fluoro-phenyl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide




Step 1: 3-bromo-2-fluorobenzaldehyde



To a solution of (3-bromo-2-fluorophenyl)methanol (2.750 g, 13.413 mmol) in dichloromethane (50 mL) was added manganese dioxide (9.32 g, 107.31 mmol) and the resulting mixture was stirred at 45° C. overnight. Once complete, the reaction was filtered through diatomaceous earth and washed with dichloromethane to obtain 3-bromo-2-fluoro-benzaldehyde as a white solid (2.24 g, 83%). ^1H NMR (400 MHz, DMSO-d_6) δ 10.19 (dt, $J=1.3, 0.7$ Hz, 1H), 8.05 (tdd, $J=6.8, 3.1, 1.5$ Hz, 1H), 7.85 (ddt, $J=7.9, 6.4, 1.5$ Hz, 1H), 7.37 (tdt, $J=7.8, 1.6, 0.8$ Hz, 1H).

Step 2: 1-bromo-3-(difluoromethyl)-2-fluorobenzene




To a solution of 3-bromo-2-fluorobenzaldehyde (1980 mg, 9.5581 mmol) in dichloromethane (30 mL, 468.0 mmol) under inert atmosphere was added diethylaminosulfur trifluoride (2.53 mL, 19.116 mmol) at 0° C., and the mixture was stirred for 1 h at 0° C. The mixture was warmed to room temperature and stirred for 1 h. The reaction was carefully quenched with saturated sodium bicarbonate solution. The reaction mixture was then diluted with ethyl acetate (300 mL) and the organic layer was washed with saturated sodium bicarbonate solution and brine then concentrated to dryness and purified by chromatography (0-50% ethyl acetate in heptane over 20 minutes) to provide 1-bromo-3-(difluoromethyl)-2-fluorobenzene (1.35 g, 61%). ^1H NMR

268


(400 MHz, DMSO-d_6) δ 7.93 (ddq, $J=7.9, 6.8, 1.2$ Hz, 1H), 7.66 (ddq, $J=7.6, 6.4, 1.2$ Hz, 1H), 7.33 (td, $J=7.9, 1.0$ Hz, 1H), 7.24 (t, $J=54.1$ Hz, 1H).

Step 3: 2-[3-(difluoromethyl)-2-fluoro-phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane







A mixture of 1-bromo-3-(difluoromethyl)-2-fluorobenzene (1.18 g, 5.09 mmol) and bis(pinacolato)diboron (2.59 g, 10.2 mmol) 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (280 mg, 0.382 mmol) and potassium acetate (1.50 g, 15.3 mmol) in 1,4-dioxane (15 mL) was heated at 100° C. overnight. The reaction mixture was filtered through diatomaceous earth and concentrated. The crude material was purified by chromatography (0-50% ethyl acetate in heptane) to obtain 2-[3-(difluoromethyl)-2-fluoro-phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane as a light yellow solid (1.11 g, 80%). ^1H NMR (400 MHz, DMSO-d_6) δ 7.86-7.73 (m, 2H), 7.36 (t, $J=7.5$ Hz, 1H), 7.18 (t, $J=54.4$ Hz, 1H), 1.31 (s, 12H).

Step 4: 7-[[3-(difluoromethyl)-2-fluoro-phenyl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide




A mixture of 7-(chloromethyl)-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide (prepared in a similar manner as Example 16.1, Step 5) (75 mg, 0.262 mmol), 2-[3-(difluoromethyl)-2-fluoro-phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (214 mg, 0.787 mmol), 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (14.4 mg, 0.0197 mmol) and potassium carbonate (110 mg, 0.787 mmol) in acetonitrile (3 mL) and water (0.75 mL) was heated at 120° C. in the microwave for 40 minutes. The reaction mixture was filtered through diatomaceous earth and concentrated. The crude material was purified by chromatography (0-100% ethyl acetate in heptane) to obtain 7-[[3-(difluoromethyl)-2-fluoro-phenyl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide (49.6 mg, 47%). LCMS (ESI): $[\text{M}+\text{H}]^+=396.1$; ^1H NMR (400 MHz, DMSO-d_6) δ 8.37 (t, $J=5.7$ Hz, 1H), 7.55 (q, $J=8.0$ Hz, 2H), 7.38-7.02 (m, 2H), 6.08 (s, 1H), 3.97 (s, 2H), 3.27-3.16 (m, 2H), 2.29 (s, 3H), 1.09 (t, $J=7.2$ Hz, 3H).

The following examples were prepared in a manner similar to Example 21.18, 21.19, and 21.20:

No.	Structure/Name	LCMS (M + H) ¹ H NMR
21.21	 <p data-bbox="285 646 623 730">N-ethyl-7-(2-ethyl-4,5-difluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	374.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.24-7.14 (m, 1H), 6.95-6.84 (m, 2H), 5.87 (s, 1H), 5.84 (s, 1H), 3.91 (s, 2H), 3.56-3.47 (m, 2H), 2.60-2.53 (m, 2H), 2.41 (s, 3H), 1.29-1.23 (m, 3H), 1.19-1.14 (m, 3H)
21.22	 <p data-bbox="285 1010 623 1094">7-((6-cyanopyridin-2-yl)methyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	354.05 ¹ H NMR (300 MHz, CDCl ₃) δ 7.61-7.50 (m, 2H), 7.26-7.20 (m, 1H), 6.07 (s, 1H), 5.85 (s, 1H), 3.95 (s, 2H), 3.57-3.48 (m, 2H), 2.42 (s, 3H), 1.28 (m, 2H)
21.23	 <p data-bbox="285 1373 623 1457">7-[[3-(difluoromethyl)-2-fluorophenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	382.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.31 (t, J = 5.3 Hz, 1H), 7.38-7.28 (m, 1H), 7.22 (t, J = 52 Hz, 2H), 6.08 (s, 1H), 4.06 (s, 2H), 2.73 (d, J = 4.6 Hz, 3H), 2.29 (s, 3H).
21.24	 <p data-bbox="285 1724 623 1835">7-[[2-fluoro-3-(hydroxymethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	362.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.31 (q, J = 4.7 Hz, 1H), 7.37 (td, J = 7.3, 1.8 Hz, 1H), 7.24 (td, J = 7.4, 1.8 Hz, 1H), 7.14 (t, J = 7.6 Hz, 1H), 6.00 (s, 1H), 5.20 (brs, 1H), 4.53 (s, 2H), 3.90 (s, 2H), 2.73 (d, J = 4.8 Hz, 3H), 2.29 (d, J = 1.0 Hz, 3H).


271

Example 21.25: 7-(3-cyclopropyl-2-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




To a solution of 7-[(3-chloro-2-fluorophenyl)methyl]-N-ethyl-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (prepared in a similar manner as Example 21.14) (70 mg, 0.18 mmol) in dioxane (2 mL) and water (0.5 mL) was added cyclopropylboronic acid (35 mg, 0.41 mmol), palladium acetate (28 mg, 0.12 mmol) and tricyclohexylphosphine (21 mg). After stirring overnight at 90° C. under nitrogen atmosphere, the resulting mixture was concentrated under vacuum. The residue was purified by chromatography with 2% methanol in dichloromethane. The crude product was purified by Prep-HPLC (SunFire Prep C₁₈ OBD Column, 5 μm, 19×150 mm; mobile phase A, water with 10 mmol NH₄HCO₃ and mobile phase B, CH₃CN; 50.0% CH₃CN up to 82.0% in 10 min, down to 50.0% in 2 min; Detector, UV 254/220 nm) to afford 7-[(3-cyclopropyl-2-fluorophenyl)methyl]-N-ethyl-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (16.8 mg, 24%) as a white solid. LCMS (ESI): [M+H]⁺=386.10; ¹H NMR (300 MHz, CDCl₃) δ 7.26-6.96 (m, 2H), 6.96-6.78 (m, 1H), 6.05 (s, 1H), 5.85 (br, 1H), 3.95 (s, 2H), 3.58-3.42 (m, 2H), 2.42 (s, 3H), 2.10-2.02 (m, 1H), 1.28 (m, 3H), 1.00-0.94 (m, 2H), 0.73-0.68 (m, 2H).

Example 21.26: 7-(3-cyano-2-fluorobenzyl)-N-ethyl-6-fluoro-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




Step 1: 7-(3-cyano-2-fluorobenzyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide



272


To a solution of 7-(chloromethyl)-N-ethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (prepared in a similar manner as Example 16.1, Step 5) (100 mg, 0.37 mmol) in 1,4-dioxane/water (2 mL/0.5 mL) was added 3-cyano-2-fluorophenylboronic acid (121 mg, 0.73 mmol), [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (28 mg, 0.04 mmol), and sodium carbonate (78 mg, 0.74 mmol). The reaction mixture was irradiated with microwave radiation for 45 min at 100° C. The resulting mixture was concentrated under vacuum and purified by chromatography with dichloromethane/methanol (30:1) to afford 7-[(3-cyano-2-fluorophenyl)methyl]-N-ethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (69 mg, 53%) as a yellow solid. LCMS (ESI): [M+H]⁺=357.0.

Step 2: 7-(3-cyano-2-fluorobenzyl)-N-ethyl-6-fluoro-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




To a solution of 7-[(3-cyano-2-fluorophenyl)methyl]-N-ethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (30 mg, 0.08 mmol) in CH₃CN (3 mL) was added Selectfluor® (30 mg, 0.24 mmol). The resulting solution was stirred for 1.5 h at 75° C. The resulting mixture was cooled to room temperature and concentrated under vacuum, and the residue was purified by chromatography with dichloromethane/methanol (30:1) to afford 7-[(3-cyano-2-fluorophenyl)methyl]-N-ethyl-6-fluoro-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (8 mg, 25%) as a white solid. LCMS (ESI): [M+H]⁺=375.0; ¹H NMR (300 MHz, CDCl₃) δ 7.73-7.65 (m, 2H), 7.58 (s, 1H), 7.36-7.31 (m, 1H), 4.20 (s, 2H), 3.33-3.31 (m, 2H), 1.29-1.10 (m, 3H).

Example 21.27: 7-(3-cyano-2-fluorobenzyl)-6-fluoro-N,N-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide



273


Step 1: 7-(chloromethyl)-6-fluoro-N,N-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide



To a solution of 7-(chloromethyl)-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (from Example 16.1, Step 5) (500 mg, 1.84 mmol) in CH₃CN (10 mL) was added Selectfluor® (980 mg, 2.76 mmol) and the resulting solution was stirred for 4 h at 75° C. The resulting mixture was concentrated under vacuum. The residue was purified by chromatography with dichloromethane/ethyl acetate (5:1) to afford 7-(chloromethyl)-6-fluoro-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (110 mg, 21%) as a light yellow solid. LCMS (ESI): [M+H]⁺=290.0.



274

Step 2: 7-(3-cyano-2-fluorobenzyl)-6-fluoro-N,N-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide







To a solution of 7-(chloromethyl)-6-fluoro-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (100 mg, 0.35 mmol) in 1,4-dioxane (2 mL) under nitrogen was added (3-cyano-2-fluorophenyl)boronic acid (86 mg, 0.52 mmol), sodium carbonate (74 mg, 0.70 mmol), [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (26 mg, 0.04 mmol), and water (0.2 mL). The reaction mixture was irradiated with microwave radiation for 20 min at 120° C. The resulting solution was extracted with ethyl acetate (3×20 mL), washed with brine, dried over anhydrous sodium sulfate, and concentrated under vacuum. The residue was purified by chromatography with dichloromethane/ethyl acetate (2:3), to afford 7-[(3-cyano-2-fluorophenyl)methyl]-6-fluoro-N,N-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (18 mg, 14%) as a off-white solid. LCMS (ESI): [M+H]⁺=374.9; ¹H NMR (300 MHz, CDCl₃) δ 7.56 (m, 2H), 7.21 (m, 1H), 5.97 (s, 1H), 4.09 (m, 2H), 3.05 (m, 3H), 2.41 (s, 3H).





The following examples were prepared in a manner similar to Example 21.26 and 21.27:

No.	Structure/Name	LCMS (M + H) ¹ H NMR
21.28	 7-(3-cyano-2-fluorobenzyl)-N-ethyl-6-fluoro-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide	388.9 ¹ H NMR (300 MHz, CDCl ₃) δ 7.56-7.50 (m, 2H), 7.23-7.19 (m, 1H), 5.86 (s, 1H), 4.09 (s, 2H), 3.58-3.51 (m, 2H), 2.40 (s, 3H), 1.32-1.28 (m, 3H)
21.29	 6-fluoro-7-(2-fluoro-3-(trifluoromethyl)benzyl)-N,N-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide	418.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.51 (m, 2H), 7.21 (m, 1H), 5.91 (s, 1H), 4.10 (s, 2H), 3.05 (m, 3H), 2.41 (s, 3H)





The following compounds were prepared using methods analogous to those described above:

No.	Structure/Name	LCMS (M + H) ¹ H NMR
21.30		367.0 ¹ H NMR (300 MHz, CD ₃ OD) δ 7.54 (m, 2H), 7.39 (m, 1H), 6.03 (s, 1H), 4.09 (s, 2H), 3.39 (m, 2H), 2.39 (s, 6H), 1.22 (m, 3H)
	7-(5-cyano-2-methylbenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide	
21.31		371.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.66-7.58 (m, 2H), 7.25-7.20 (m, 1H), 6.12 (s, 1H), 5.90-5.80 (m, 1H), 3.96 (s, 2H), 3.60-3.50 (m, 2H), 2.45 (s, 3H), 1.33-1.29 (m, 3H)
	7-(5-cyano-2-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide	
21.32		379.95 ¹ H NMR (300 MHz, CDCl ₃) δ 7.36-7.32 (m, 1H), 7.03-6.91 (m, 2H), 5.99 (s, 1H), 5.87 (br, 1H), 3.99 (s, 2H), 3.57-3.48 (m, 2H), 2.41 (s, 3H), 1.28 (m, 3H)
	7-(2-chloro-5-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide	
21.33		357.0 ¹ H NMR (300 MHz, CDCl ₃) δ 9.66 (br, 1H), 8.02 (s, 1H), 7.58-7.55 (m, 2H), 7.27 (s, 1H), 6.19 (s, 1H), 4.00 (s, 2H), 3.50-3.41 (m, 2H), 1.27-1.17 (m, 3H)
	7-(3-cyano-2-fluorobenzyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide	






-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
21.34	 <p data-bbox="331 684 672 768">N-ethyl-7-(5-fluoro-2-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	421.20 ¹ H NMR (300 MHz, CDCl ₃) δ 7.70-7.66 (m, 1H), 7.09-7.04 (m, 2H), 5.95-5.92 (m, 2H), 4.08 (s, 2H), 3.56-3.47 (m, 2H), 2.41 (s, 3H), 1.30-1.18 (m, 3H)
21.35	 <p data-bbox="331 1100 672 1184">N-ethyl-2-methyl-7-(naphthalen-1-ylmethyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	378.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.87-7.79 (m, 3H), 7.52-7.40 (m, 4H), 5.95-5.84 (m, 1H), 5.81 (s, 1H), 4.36 (s, 2H), 3.66-3.47 (m, 2H), 2.37 (s, 3H), 1.25-1.16 (t, 3H)
21.36	 <p data-bbox="331 1495 672 1579">N-ethyl-7-(5-fluoro-2-methylbenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	360.0 ¹ H NMR (300 MHz, CD ₃ OD) δ 7.19 (m, 1H), 6.94 (m, 2H), 5.95 (s, 1H), 3.95 (s, 2H), 3.40 (m, 2H), 2.40 (s, 3H), 2.26 (s, 3H), 1.23 (m, 3H)
21.37	 <p data-bbox="331 1852 672 1936">7-(3-cyano-4-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	371.0 ¹ H NMR (400 MHz, CDCl ₃) δ 7.54-7.47 (m, 2H), 7.19-7.14 (m, 1H), 6.09 (s, 1H), 5.84 (br, 1H), 3.86 (s, 2H), 3.57-3.50 (m, 2H), 2.42 (s, 3H), 1.27 (m, 3H)






-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
21.38	 <p data-bbox="326 642 672 716">N-ethyl-2-methyl-7-((1-methyl-1H-indazol-4-yl)methyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	382.20 ¹ H NMR (400 MHz, CDCl ₃) δ 7.98 (s, 1H), 7.37-7.30 (m, 2H), 7.02-7.01 (m, 1H), 6.01 (s, 1H), 5.83 (s, 1H), 4.25 (s, 2H), 4.12 (s, 3H), 3.51-3.50 (m, 2H), 2.45-2.35 (m, 3H), 1.27-1.15 (m, 3H)
21.39	 <p data-bbox="347 1068 656 1142">N-ethyl-2-methyl-7-(3-(methylsulfonamido)benzyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	421.0 ¹ H NMR (300 MHz, CD ₃ OD) δ 7.29 (m, 2H), 7.09 (m, 2H), 6.13 (s, 1H), 3.91 (s, 2H), 3.40 (m, 2H), 2.94 (s, 3H), 2.39 (s, 3H), 1.23 (m, 3H)
21.40	 <p data-bbox="347 1493 656 1568">7-(5-cyano-2-(trifluoromethyl)benzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	421.20 ¹ H NMR (300 MHz, CDCl ₃) δ 7.81-7.78 (m, 1H), 7.69-7.67 (m, 1H), 5.60 (s, 1H), 5.82 (s, 1H), 4.11 (s, 2H), 3.58-3.52 (m, 2H), 2.44 (s, 3H), 1.31-1.26 (m, 3H)
21.41	 <p data-bbox="331 1856 672 1932">7-(4-chloro-2-methylbenzyl)-N,N-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	362.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.13 (m, 3H), 5.99 (s, 1H), 5.87 (s, 1H), 4.10 (s, 2H), 3.03 (s, 3H), 2.30 (s, 3H), 2.24 (s, 3H)






-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
21.42	 <p data-bbox="337 583 662 659">7-(2,5-difluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	364.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.05-6.89 (m, 3H), 6.10 (s, 1H), 6.04 (s, 1H), 3.88 (s, 2H), 3.55-3.46 (m, 2H), 2.36 (s, 3H), 1.29-1.27 (m, 3H)
21.43	 <p data-bbox="337 919 662 995">7-(3-cyanobenzyl)-2-cyclopropyl-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	379.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.54-7.53 (m, 2H), 7.49-7.38 (m, 2H), 6.06 (s, 1H), 5.89 (s, 1H), 3.88 (s, 2H), 3.56-3.49 (m, 2H), 2.19-2.09 (m, 1H), 1.31-1.21 (m, 3H), 1.18-1.11 (m, 2H), 0.88-0.78 (m, 2H)
21.44	 <p data-bbox="337 1255 662 1310">N-ethyl-7-(2-fluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	346.0 ¹ H NMR (400 MHz, CDCl ₃) δ 7.48-7.46 (m, 1H), 7.30-7.21 (m, 1H), 7.14-7.03 (m, 2H), 6.06 (s, 1H), 6.01 (s, 1H), 3.93 (s, 2H), 3.56-3.46 (m, 2H), 2.38 (s, 3H), 1.30-1.25 (m, 3H)
21.45	 <p data-bbox="337 1549 662 1625">7-(2,3-difluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	364.0 ¹ H NMR (400 MHz, CDCl ₃) δ 7.11-6.99 (m, 3H), 6.04 (s, 1H), 5.90 (s, 1H), 3.95 (s, 2H), 3.55-3.48 (m, 2H), 2.40 (s, 3H), 1.29-1.16 (m, 3H)
21.46	 <p data-bbox="337 1885 662 1940">N-ethyl-7-(3-fluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	346.0 ¹ H NMR (300 MHz, CD ₃ OD) δ 7.36-7.29 (m, 1H), 7.13-6.94 (m, 3H), 6.15 (s, 1H), 3.93 (s, 2H), 3.41 (m, 2H), 2.40 (s, 3H), 1.24 (m, 3H)





-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
21.47	 <p data-bbox="334 558 667 625">7-[(3-chloro-4-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	366.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.29 (t, J = 4.8 Hz, 1H), 7.52 (dd, J = 7.2, 2.0 Hz, 1H), 7.39-7.28 (m, 2H), 6.17 (s, 1H), 2.73 (d, J = 4.7 Hz, 3H), 2.29 (s, 3H).
21.48	 <p data-bbox="334 905 667 972">7-[(2,5-dichlorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	382.0 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.32 (q, J = 4.7 Hz, 1H), 7.52-7.48 (m, 2H), 7.39 (dd, J = 8.6, 2.6 Hz, 1H), 5.97 (s, 1H), 4.02 (s, 2H), 2.73 (d, J = 4.8 Hz, 3H), 2.29 (s, 3H).
21.49	 <p data-bbox="334 1209 667 1276">N,2-dimethyl-5-oxo-7-[(3-(trifluoromethoxy)phenyl)methyl]thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	398.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.35-8.25 (m, 1H), 7.45 (t, J = 7.9 Hz, 1H), 7.33 (dd, J = 9.9, 2.4 Hz, 2H), 7.27-7.19 (m, 1H), 6.16 (d, J = 1.2 Hz, 1H), 3.93 (s, 2H), 2.73 (d, J = 4.6 Hz, 3H), 2.29 (s, 3H).
21.50	 <p data-bbox="334 1535 667 1602">7-[(5-cyano-2-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	357.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.30 (d, J = 4.9 Hz, 1H), 7.92 (dd, J = 6.9, 2.2 Hz, 1H), 7.85 (ddd, J = 8.5, 4.8, 2.2 Hz, 1H), 7.44 (dd, J = 9.7, 8.6 Hz, 1H), 6.11 (s, 1H), 3.97 (s, 2H), 2.73 (d, J = 4.7 Hz, 3H), 2.29 (s, 3H).
21.51	 <p data-bbox="334 1860 667 1936">7-[(3-chloro-5-cyano-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	373.0 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.31-8.25 (m, 1H), 7.91 (t, J = 1.8 Hz, 1H), 7.76 (dt, J = 7.0, 1.6 Hz, 2H), 6.22 (s, 1H), 3.94 (s, 2H), 2.74 (d, J = 4.7 Hz, 3H), 2.29 (s, 3H).




-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
21.52	 <p data-bbox="334 583 669 646">7-[(3-cyclopropylphenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	354.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.30 (q, J = 4.7 Hz, 1H), 7.20-7.14 (m, 1H), 7.03 (dd, J = 6.8, 1.4 Hz, 2H), 6.91 (dt, J = 7.9, 1.5 Hz, 1H), 6.08 (s, 1H), 3.80 (s, 2H), 2.73 (d, J = 4.7 Hz, 3H), 2.28 (s, 3H), 1.88 (tt, J = 8.4, 5.1 Hz, 1H), 0.97-0.88 (m, 2H), 0.68-0.61 (m, 2H).
21.53	 <p data-bbox="334 932 669 999">7-[(2,5-difluorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	350.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.31 (q, J = 4.7 Hz, 1H), 7.24 (tt, J = 8.8, 4.1 Hz, 2H), 7.15 (tt, J = 8.8, 3.6 Hz, 1H), 6.06 (s, 1H), 3.91 (s, 2H), 2.73 (d, J = 4.7 Hz, 3H), 2.29 (s, 3H).
21.54	 <p data-bbox="334 1247 669 1310">7-[(3,4-difluorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	350.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.30 (q, J = 4.7 Hz, 1H), 7.43-7.29 (m, 2H), 7.15 (dd, J = 4.6, 2.2 Hz, 1H), 6.15 (s, 1H), 3.86 (s, 2H), 2.78-2.68 (m, 3H), 2.34 (s, 3H).
21.55	 <p data-bbox="334 1558 669 1621">7-[(2,3-difluorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	350.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.31 (t, J = 5.3 Hz, 1H), 7.38-7.28 (m, 1H), 7.22-7.13 (m, 2H), 6.08 (s, 1H), 4.06 (s, 2H), 2.73 (d, J = 4.6 Hz, 3H), 2.29 (s, 3H).
21.56	 <p data-bbox="334 1869 669 1932">7-[(4-chloro-2-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	366.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.32 (q, J = 4.7 Hz, 1H), 7.46-7.41 (m, 1H), 7.41-7.36 (m, 1H), 7.26 (dd, J = 8.3, 2.1 Hz, 1H), 6.06 (s, 1H), 3.91 (s, 2H), 2.73 (d, J = 4.7 Hz, 3H), 2.29 (s, 3H).





-continued

No.	Structure/Name	LCMS (M + H) ⁺ ¹ H NMR
21.57	 <p data-bbox="326 604 675 688">7-[(2,4-dichlorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	382.0 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.29 (d, J = 5.4 Hz, 1H), 7.61-7.55 (m, 2H) 7.30 (dd, J = 8.3, 2.1 Hz, 1H), 6.19 (s, 1H), 3.88 (s, 2H), 2.73 (d, J = 4.7 Hz, 3H), 2.29 (s, 3H).
21.58	 <p data-bbox="326 961 675 1045">7-[(3-fluoro-4-methyl-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	346.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.29 (d, J = 5.2 Hz, 1H), 7.21 (t, J = 8.0 Hz, 1H), 7.09-6.99 (m, 2H), 6.12 (s, 1H), 3.83 (s, 2H), 2.73 (dt, J = 4.8, 0.8 Hz, 3H), 2.29 (s, 3H), 2.23 (s, 3H).
21.59	 <p data-bbox="326 1360 675 1480">7-[[4-fluoro-2-(trifluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	400.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.32-8.26 (m, 1H), 7.73 (dd, J = 7.1, 2.2 Hz, 1H), 7.67 (ddd, J = 7.8, 4.9, 2.3 Hz, 1H), 7.45 (dd, J = 11.0, 8.4 Hz, 1H), 6.19 (s, 1H), 3.96 (s, 2H), 2.73 (d, J = 4.6 Hz, 3H), 2.29 (s, 3H).
21.60	 <p data-bbox="326 1822 675 1904">7-[(2-cyclopropyl-4-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	372.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.36 (q, J = 4.7 Hz, 1H), 7.24 (dd, J = 8.5, 6.2 Hz, 1H), 6.96 (td, J = 8.5, 2.8 Hz, 1H), 6.78 (dd, J = 10.7, 2.7 Hz, 1H), 5.92 (d, J = 0.7 Hz, 1H), 4.04 (s, 2H), 2.73 (d, J = 4.7 Hz, 3H), 2.29 (s, 3H), 1.98 (ddd, J = 13.7, 8.6, 5.3 Hz, 1H), 0.94-0.85 (m, 2H), 0.69-0.60 (m, 2H).

-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
21.61		425.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.65 (m, 3H), 5.93 (s, 1H), 5.85 (s, 1H), 4.72 (m, 1H), 4.57 (m, 1H), 3.96 (s, 2H), 3.52 (m, 2H), 3.13 (m, 1H), 3.05 (m, 1H), 2.42 (s, 3H), 1.25 (m, 3H)
21.62	7-(2-chloro-3-cyanobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide	386.9 ¹ H NMR (300 MHz, DMSO-d ₆) δ 8.42-8.39 (m, 1H), 7.93-7.90 (m, 1H), 7.78-7.76 (m, 1H), 7.57-7.52 (m, 1H), 6.02 (s, 1H), 4.11 (s, 2H), 3.31-3.18 (m, 2H), 2.30 (s, 3H), 1.12-1.07 (m, 3H)
21.63		392.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.04 (m, 2H), 5.94 (s, 1H), 5.82 (s, 1H), 4.01 (s, 2H), 3.51 (m, 2H), 2.58 (m, 2H), 2.41 (s, 3H), 1.26 (m, 3H), 1.15 (m, 3H)
21.64		381.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.53-7.51 (m, 1H), 7.44 (s, 1H), 7.33-7.26 (m, 1H), 5.89-5.87 (m, 2H), 3.94 (s, 2H), 3.56-3.49 (m, 2H), 2.71-2.65 (m, 2H), 2.41 (s, 3H), 1.29-1.23 (m, 3H), 1.21-1.17 (m, 3H)


-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
21.65		386.10 ¹ H NMR (300 MHz, CDCl ₃) δ 7.27-7.00 (m, 1H), 6.92-6.68 (m, 2H), 5.95 (s, 1H), 5.86 (br, 1H), 4.10 (s, 2H), 3.58-3.49 (m, 2H), 2.43 (s, 3H), 1.84-1.75 (m, 1H), 1.31-1.19 (m, 3H), 0.92-0.88 (m, 2H), 0.67-0.55 (m, 2H)
7-(2-cyclopropyl-5-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide		
21.66		393.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.47-7.44 (m, 1H), 7.32-7.19 (m, 2H), 5.94 (s, 1H), 5.83-5.81 (m, 1H), 4.16 (s, 2H), 3.56-3.52 (m, 2H), 2.41 (s, 3H), 1.92-1.82 (m, 1H), 1.32-1.23 (m, 3H), 1.02-0.96 (m, 2H), 0.70-0.65 (m, 2H)
7-(5-cyano-2-cyclopropylbenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide		
21.67		388.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.16-7.11 (m, 1H), 6.93-6.83 (m, 2H), 5.86 (s, 1H), 5.86 (s, 1H), 3.90 (s, 2H), 3.53-3.51 (m, 2H), 2.54-2.49 (m, 2H), 2.42 (s, 3H), 1.59-1.51 (m, 2H), 1.29-1.24 (m, 3H), 0.96-0.91 (m, 3H)
N-ethyl-7-(5-fluoro-2-propylbenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide		
21.68		364.1 ¹ H NMR (400 MHz, DMSO-d ₆) δ 8.31 (q, J = 4.6 Hz, 1H), 7.43 (tq, J = 7.4, 2.2 Hz, 2H), 7.26-7.18 (m, 1H), 6.04 (s, 1H), 5.48 (dd, J = 47.6, 1.2 Hz, 2H), 3.95 (s, 2H), 2.73 (d, J = 4.7 Hz, 3H), 2.29 (s, 3H).
7-[[2-fluoro-3-(fluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide		


293

Method 22

Example 22.1: N-ethyl-7-(1-(2-fluoro-3-(trifluoromethyl)phenyl)ethyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide



Step 1: N-ethyl-7-(1-(2-fluoro-3-(trifluoromethyl)phenyl)ethyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




To a -78°C . solution of N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (from Example 21.12) (200 mg, 0.48 mmol) in tetrahydrofuran (20 mL) was added n-butyllithium (2.5 M in hexanes; 1 mL). After 30 min at -78°C ., iodomethane (235 mg, 1.66 mmol) was added. The resulting solution was stirred for 3 h at room temperature. The reaction was then quenched by water (20 mL), extracted with dichloromethane (3x30 mL), washed with brine, dried over sodium sulfate and concentrated in vacuo. The residue was purified by chromatography with dichloromethane/methanol (50:1) to afford N-ethyl-7-[1-(2-fluoro-3-(trifluoromethyl)phenyl)ethyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (52.7 mg, 25%) as a white solid. LCMS (ESI): $[M+H]^+=428.0$; $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 7.60-7.48 (m, 2H), 7.20-7.19 (m, 1H), 6.13 (s, 1H), 5.92 (m, 1H), 4.42-4.34 (m, 2H), 3.56-3.48 (m, 2H), 2.40 (s, 3H), 1.64-1.62 (m, 3H), 1.30-1.25 (m, 3H).

The following compound was prepared using methods analogous to those described above:


294

Method 23

Example 23.1: 3-((3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile



Step 1: 3-((3-bromo-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile




To a solution of 3-bromo-7-(chloromethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 4.1, Step 6) (500 mg, 1.70 mmol) in 1,4-dioxane/ H_2O (3/1 mL) was added (3-cyanophenyl)boronic acid (300 mg, 2.04 mmol), tetrakis(triphenylphosphine)palladium (197 mg, 0.17 mmol) and potassium phosphate (730 mg, 3.44 mmol). The resulting solution was stirred overnight at 80°C . After cooling down to room temperature, the resulting mixture was washed with brine (30 mL), extracted with dichloromethane (3x20 mL), dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by silica gel chromatography with ethyl acetate/petroleum ether (1:2) to afford 3-((3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile as a light brown solid (522 mg, 85%). LCMS $[M+H]^+=360.0$, 362.0.

No.	Structure/Name	LCMS (M + H) $^1\text{H NMR}$
22.2	<p>N-ethyl-7-(2-fluoro-3-(trifluoromethyl)benzyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	428.0 $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 7.55-7.48 (m, 2H), 7.26-7.18 (m, 1H), 6.05 (s, 1H), 3.97 (s, 2H), 3.77-3.68 (m, 0.5H), 3.54-3.45 (m, 0.5H), 3.26-3.18 (m, 1H), 3.11 (s, 1H), 2.86 (s, 2H), 2.34 (s, 3H), 1.30-1.25 (m, 2H), 1.17-1.12 (m, 1H).


295

Step 2: ethyl 2-(7-(3-cyanobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)cyclopropanecarboxylate



To a solution of 3-([3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl)methyl]benzonitrile (200 mg, 0.56 mmol) in CH₃CN/H₂O (3/1 mL) was added ethyl 2-(tetramethyl-1,3,2-dioxaborolan-2-yl)cyclopropane-1-carboxylate (from Example 4.18, Step 1) (267 mg, 1.11 mmol), [1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloride (42 mg, 0.06 mmol) and potassium carbonate (154 mg, 1.11 mmol). The reaction mixture was heated under microwave irradiation for 1.5 h at 120° C. The resulting mixture was washed with brine (20 mL), extracted with dichloromethane (3×20 mL), dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by chromatography with dichloromethane/methanol (50:1) to afford ethyl 7-[(3-cyanophenyl)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-carboxylate (68 mg 35%) as a brown solid. LCMS (ESI): [M+H]⁺=394.0.


Step 3: 2-(7-(3-cyanobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)cyclopropanecarboxylic acid



To a solution of 7-[(3-cyanophenyl)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carboxylate (68 mg, 0.17 mmol) in THF/H₂O (2/1 mL) was added lithium hydroxide (73 mg, 1.7 mmol). The resulting solution was stirred overnight at room temperature. The pH of the solution was adjusted to pH 7 with hydrochloric acid solution (aq.) and the resulting mixture was extracted with ethyl acetate (3×20 mL), washed with brine, dried over sodium sulfate and concentrated under vacuum to afford 2-[7-[(3-cyanophenyl)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carboxylic acid (80 mg, crude) as a brown solid. The crude product was used in next step without further purification. LCMS (ESI): [M+H]⁺=366.0.


296

Step 4: 2-(7-(3-cyanobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)cyclopropanecarboxamide




To a solution of 2-[7-[(3-cyanophenyl)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carboxylic acid (80 mg, 0.22 mmol) in tetrahydrofuran (5 mL) was added propan-2-yl chloroformate (40.4 mg, 0.33 mmol), triethylamine (44 mg, 0.43 mmol) and ammonia (2 mL, 25 weight % in water). The resulting solution was stirred for 1 h at room temperature and concentrated under vacuum, and the residue was purified by chromatography with dichloromethane/methanol (30:1) to afford 2-[7-[(3-cyanophenyl)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carboxamide (21 mg, 26%) as a brown solid. LCMS (ESI): [M+H]⁺=364.9.

Step 5: 3-((3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile




To a solution of 2-[7-[(3-cyanophenyl)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carboxamide (17 mg, 0.05 mmol) in methylene chloride (10 mL) was added 1,8-diazabicyclo[5.4.0]undec-7-ene (0.25 mL) and ethoxyphosphonyl dichloride (0.25 mL). The resulting solution was stirred for 1 h at room temperature. The reaction was then quenched by water (30 mL), extracted with dichloromethane (3×20 mL), washed with brine, and concentrated under vacuum. The residue was purified on a silica gel column eluted with dichloromethane/methanol (50:1) to afford 3-[(3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl)methyl]benzonitrile (12.6 mg, 78%) as a off-white solid. LCMS (ESI): [M+H]⁺=347.0; ¹H NMR (300 MHz, CDCl₃) δ 7.69-7.60 (m, 3H), 7.53-7.47 (m, 1H), 6.16 (s, 1H), 3.94 (s, 2H), 2.92-2.88 (m, 1H), 2.39 (s, 3H), 1.96-1.92 (m, 1H), 1.81-1.74 (m, 1H), 1.59-1.52 (m, 1H).

Example 23.2: 3-((3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzonitrile



297


Step 1: 3-((3-bromo-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzonitrile



A mixture of 3-bromo-7-(chloromethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from Example 4.1, Step 6) (500 mg, 1.70 mmol), potassium phosphate (733 mg, 3.45 mmol), tetrakis(triphenylphosphine)palladium (198 mg, 0.17 mmol), (3-cyano-2-fluorophenyl)boronic acid (339 mg, 2.06 mmol), 1,4-dioxane (6 mL) and water (1 mL) was stirred overnight at 80° C. in a 30-mL sealed tube. The resulting mixture was diluted with brine and extracted with 3x30 mL of dichloromethane, and the combined organic layers were washed with brine, dried over anhydrous sodium sulfate, and concentrated under vacuum. The residue was purified by silica gel chromatography with dichloromethane/methanol (80:1) to afford 3-((3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzonitrile (70 mg, 11%) as a yellow solid. LCMS (ESI): [M+H]⁺=378.

298

Step 2: 3-((3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzonitrile



To a solution of 3-((3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzonitrile (100 mg, 0.26 mmol) in 1,4-dioxane/H₂O (2 mL/0.5 mL) added tetrakis(triphenylphosphine)palladium (31 mg, 0.03 mmol), potassium phosphate (112 mg, 0.53 mmol), and potassium trans-2-cyanocyclopropyltrifluoroborate (prepared in a manner similar to Example 4.1, Step 2) (92 mg, 0.53 mmol). The resulting solution was stirred for 3 h at 80° C. After filtration to remove solids, the filtrate was concentrated under vacuum and purified by chromatography with dichloromethane/methanol (100:1) to afford 3-((3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzonitrile (16 mg, 17%) as a white solid. LCMS (ESI): [M+H]⁺=364.9; ¹H NMR (300 MHz, CDCl₃) δ 7.60-7.54 (m, 2H), 7.28-7.21 (m, 1H), 6.03 (s, 1H), 3.94 (s, 2H), 3.00-2.94 (m, 1H), 2.43 (s, 3H), 1.90-1.80 (m, 1H), 1.79-1.70 (m, 1H), 1.45-1.35 (m, 1H).

The following example was prepared in a manner similar to Example 23.1 and 23.2:

No.	Structure/Name	LCMS (M + H) ¹ H NMR
23.3	<p>6-((3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)picolinonitrile</p>	348.05 ¹ H NMR (300 MHz, CDCl ₃) δ 7.95-7.92 (m, 1H), 7.77-7.69 (m, 2H), 6.21 (s, 1H), 4.86 (s, 2H), 2.86-2.98 (m, 1H), 2.39 (s, 3H), 1.28-2.02 (m, 3H)

45


The following compound was prepared using methods analogous to those described above:

No.	Structure/Name	LCMS (M + H) ¹ H NMR
23.4	<p>3-((3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-4-methoxybenzonitrile</p>	753.15 ¹ H NMR (300 MHz, CDCl ₃) δ 7.62 (m, 1H), 7.50 (s, 1H), 6.96 (m, 1H), 6.01 (s, 1H), 3.91 (s, 3H), 3.88 (s, 2H), 3.01 (m, 1H), 2.41 (s, 3H), 1.81 (m, 1H), 1.74 (m, 1H), 1.41 (m, 1H)

299

Method 24

Example 24.1: 2-methyl-3-(pyrimidin-5-yl)-7-(3-(trifluoromethyl)benzyl)-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 3-bromo-2-methyl-7-[[3-(trifluoromethyl)phenyl]methyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-





300


one (prepared in a manner similar to Example 21.1, Step 1) (60 mg, 0.15 mmol) in 1,4-dioxane (2 mL) was added (pyrimidin-5-yl)boronic acid (37 mg, 0.30 mmol), potassium phosphate (64 mg, 0.30 mmol), tetrakis(triphenylphosphine)palladium (17 mg, 0.01 mmol) and water (0.2 mL). The resulting solution was stirred for 3 h at 90° C. in an oil bath. The resulting solution was quenched with water (10 mL), extracted with dichloromethane (3×20 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was then purified by chromatography with ethyl acetate/petroleum ether (1:2.5) to afford 2-methyl-3-(pyrimidin-5-yl)-7-[[3-(trifluoromethyl)phenyl]methyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (27.7 mg, 46%) as a off-white solid. LCMS (ESI): [M+H+41]⁺ 444.1; ¹H NMR (300 MHz, CD₃OD) δ 9.18 (s, 1H), 8.82 (s, 2H), 7.65-7.55 (m, 4H), 6.11 (s, 1H), 4.03 (s, 2H), 2.28 (s, 3H).

The following examples were prepared in a manner similar to Example 24.1:

No.	Structure/Name	LCMS (M + H) ¹ H NMR
24.2	<p>7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	421.0 ¹ H NMR (300 MHz, CDCl ₃) δ 9.25 (s, 1H), 8.70 (s, 2H), 7.56-7.50 (m, 2H), 7.22-7.19 (m, 1H), 5.99 (s, 1H), 3.99 (s, 2H), 2.27 (s, 3H)
24.3	<p>2-fluoro-3-((2-methyl-5-oxo-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile</p>	378.0 ¹ H NMR (300 MHz, CDCl ₃) δ 9.25 (s, 1H), 8.69 (s, 2H), 7.60-7.53 (m, 2H), 7.23-7.21 (m, 1H), 6.00 (s, 1H), 3.96 (s, 2H), 2.26 (s, 3H)
24.4	<p>3-((3-cyclopropyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzonitrile</p>	340.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.60-7.51 (m, 2H), 7.24-7.18 (m, 1H), 5.96 (s, 1H), 3.90 (s, 2H), 2.36 (s, 3H), 2.26-2.25 (m, 1H), 1.10-1.03 (m, 2H), 0.69-0.65 (m, 2H)


The following compounds were prepared using methods analogous to those described above:

No.	Structure/Name	LCMS (M + H) ¹ H NMR
24.5	 <p data-bbox="277 709 649 785">7-(isoquinolin-4-ylmethyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	386.0 ¹ H NMR (300 MHz, CDCl ₃) δ 9.22 (s, 2H), 8.66 (s, 2H), 8.49 (s, 1H), 8.03 (m, 1H), 7.95 (m, 1H), 7.76-7.62 (m, 2H), 5.87 (s, 1H), 4.32 (s, 2H), 2.26 (s, 3H)
24.6	 <p data-bbox="277 1077 649 1157">3-((2-methyl-5-oxo-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile</p>	401.15 ¹ H NMR (300 MHz, CDCl ₃) δ 9.26 (s, 1H), 8.76 (s, 2H), 7.68-7.41 (m, 3H), 7.29-7.26 (m, 1H), 6.02 (s, 1H), 3.92 (s, 2H), 2.27 (s, 3H)
24.7	 <p data-bbox="277 1503 649 1577">7-(5-fluoro-2-methoxybenzyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	382.95 ¹ H NMR (300 MHz, CD ₃ OD) δ 9.21 (s, 1H), 8.81 (s, 2H), 7.07-6.95 (m, 3H), 5.92 (s, 1H), 3.91 (s, 2H), 3.79 (s, 3H), 2.17 (s, 3H)
24.8	 <p data-bbox="277 1860 649 1934">2-fluoro-3-((3-(furan-2-yl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile</p>	366.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.58-7.52 (m, 3H), 7.24-7.19 (m, 1H), 6.52-6.50 (m, 2H), 6.01 (s, 1H), 3.95 (s, 2H), 2.30 (s, 3H)


No.	Structure/Name	LCMS (M + H) ¹ H NMR
24.9	 <p>3-bromo-2-methyl-7-(3-(methylsulfonyl)benzyl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	415 ¹ H NMR (300 MHz, CDCl ₃) δ 7.87-7.84 (m, 3H), 7.56-7.53 (m, 1H), 6.04 (s, 1H), 3.95 (s, 2H), 3.06 (s, 3H), 2.35 (s, 3H)

Method 25

Example 25.1: 7-(3-cyano-2-fluorobenzyl)-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




Step 1: ethyl 7-(3-cyano-2-fluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxylate




To a solution of ethyl 7-(chloromethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxylate (500 mg, 1.74 mmol) in dioxane (2 mL) and water (0.5 mL) was added (3-cyano-2-fluorophenyl)boronic acid (375 mg, 2.27 mmol), sodium carbonate (370 mg, 3.49 mmol) and [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (77 mg, 0.11 mmol). After stirring 20 minutes at 120° C. under nitrogen atmosphere, the resulting mixture was concentrated under vacuum. The residue was purified by chromatography with 10% ethyl acetate in dichloromethane to afford ethyl 7-[(3-cyano-2-fluorophenyl)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxylate (23.1 mg, 18%) as a white solid. LCMS (ESI): [M+H]⁺=372.0; ¹H NMR (300 MHz, CDCl₃) δ 7.72-7.66 (m, 2H), 7.36-7.31 (m, 1H), 6.19 (s, 1H), 4.44-4.37 (m, 2H), 4.06 (s, 2H), 2.43 (s, 3H), 1.36 (m, 3H).

Step 2: 7-(3-cyano-2-fluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxylic acid



To a solution of ethyl 7-[(3-cyano-2-fluorophenyl)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxylate (prepared in a manner similar to Example 21.1) (100 mg, 0.27 mmol) in tetrahydrofuran (3 mL) was added lithium hydroxide (64 mg, 2.67 mmol) in water (3 mL). The resulting solution stirred for 3 days at room temperature. The reaction mixture was diluted with dichloromethane (20 mL) and water (5 mL). The pH value of the water layer was adjusted to pH 6 with hydrogen chloride. The resulting solution was extracted with dichloromethane (3×10 mL), washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum to afford 7-[(3-cyano-2-fluorophenyl)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxylic acid (22 mg, 24%) as a brown solid. The crude product was used in next step without further purification. LCMS (ESI): [M+H]⁺=344.0.

Step 3: 7-(3-cyano-2-fluorobenzyl)-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




To a solution of 7-[(3-cyano-2-fluorophenyl)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxylic acid (60 mg, 0.17 mmol) in N,N-dimethylformamide (9 mL) was added 2,2,2-trifluoroethan-1-amine (36 mg, 0.36

305


mmol), N-hydroxybenzotriazole (36 mg, 0.27 mmol), N,N-diisopropylethylamine (66 mg, 0.51 mmol) and 1,1'-thiocarbonyldiimidazole (48 mg, 0.27 mmol). After stirred for 5 h at room temperature, the mixture was diluted with water (5 mL). The resulting solution was extracted with dichloromethane (20 mL×3), washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by chromatography with 1% methanol in dichloromethane. The crude product was purified by Prep-HPLC (SunFire Prep C₁₈ OBD Column, 5 μm, 19×150 mm; mobile phase, water with 10 mmol NH₄HCO₃ and CH₃CN (50.0% CH₃CN up to 82.0% in 10 min, down to 50.0% in 2 min); Detector, UV 254/220 nm) to afford 7-[(3-cyano-2-fluorophenyl)methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (20.8 mg, 28%) as a white solid. LCMS (ESI): [M+H]⁺ 424.95; ¹H NMR (300 MHz, CD₃OD) δ 7.72-7.66 (m, 2H), 7.36-7.31 (m, 1H), 6.19 (s, 1H), 4.15-4.06 (m, 4H), 2.40 (s, 3H).

The following example was prepared in a manner similar to Example 25.1:

No.	Structure/Name	LCMS (M + H) ⁺ ¹ H NMR
25.2	 <p>N-(cyanomethyl)-7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide</p>	424.95 (300 MHz, DMSO-d ₆) δ 9.15 (m, 1H), 7.75-7.68 (m, 2H), 7.42-7.37 (m, 1H), 6.16 (s, 1H), 4.32 (m, 2H), 4.03 (s, 2H), 2.31 (s, 3H)


Method 26

Example 26.1: 7-(3-cyanobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile




added copper cyanide (49 mg, 0.55 mmol). The resulting solution was stirred for 1 h at 100° C. After cooling to room temperature, the reaction mixture was concentrated under vacuum. The residue was purified by chromatography with dichloromethane/methanol (50:1) to afford 7-[(3-cyanophenyl)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carbonitrile (8.4 mg, 10%) as a off-white solid. LCMS (ESI): [M+H]⁺=306.9; ¹H NMR (300 MHz, CDCl₃) δ 7.60-7.56 (m, 2H), 7.51-7.41 (m, 2H), 6.17 (s, 1H), 3.91 (s, 2H), 2.65 (s, 3H).

The following example was prepared in a manner similar to Example 26.1:

No.	Structure/Name	LCMS (M + H) ⁺ ¹ H NMR
26.2	 <p>7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile</p>	368.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.57-7.48 (m, 2H), 7.23-7.21 (m, 1H), 6.16 (s, 1H), 3.98 (s, 2H), 2.65 (s, 3H)





306

Step 1: 7-(3-cyanobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile








To a solution of 3-[(3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl)methyl]benzonitrile (from Example 21.1, Step 1) (100 mg, 0.28 mmol) in N,N-dimethylformamide (10 mL) under inert atmosphere was






The following compounds were prepared using methods analogous to those described herein.

No.	Structure/Name	LCMS (M + H) ¹ H NMR
27.1	 <p>10-(4-fluorophenoxymethyl)-3-(hydroxymethyl)-7-thia-1,9-diazatricyclo[6.4.0.0']dodeca-2(6),8,10-trien-12-one</p>	347 ¹ H NMR (300 MHz, CDCl ₃) δ 7.01-6.87 (m, 4H), 6.52 (s, 1H), 4.95 (s, 2H), 3.96-3.91 (m, 2H), 3.77-3.66 (m, 1H), 2.98-2.64 (m, 3H), 2.28-2.11 (m, 1H)
27.2	 <p>10-(4-Fluorophenoxymethyl)-3-(2-hydroxyethyl)-7-thia-1,9-diazatricyclo[6.4.0.0']dodeca-2(6),8,10-trien-12-one</p>	361.10 ¹ H NMR (300 MHz, CDCl ₃) δ 7.02-6.88 (m, 4H), 6.48 (s, 1H), 4.94 (s, 2H), 3.95-3.91 (m, 1H), 3.68-3.62 (m, 2H), 2.99-2.91 (m, 1H), 2.85-2.69 (m, 2H), 2.30-2.24 (m, 1H), 2.00-1.89 (m, 2H)
27.3	 <p>7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-methylsulfonyl-thiazolo[3,2-a]pyrimidin-5-one</p>	396.1
27.4	 <p>3-(hydroxymethyl)-10-[[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-7-thia-1,9-diazatricyclo[6.4.0.0']dodeca-2(6),8,10-trien-12-one</p>	388.15 ¹ H NMR (400 MHz, CD ₃ OD) δ 7.94 (s, 1H), 6.70 (s, 1H), 5.93 (s, 1H), 5.36 (s, 2H), 3.88-3.76 (m, 3H), 3.02-2.40 (m, 4H)


-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
27.5	 <p>10-([5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl)-7-thia-1,9-diazatricyclo[6.4.0.0']_{2,6}dodeca-2(6),8,10-trien-12-one</p>	381.00 ¹ H NMR (300 MHz, CD ₃ OD) δ 6.32 (s, 1H), 5.72 (s, 1H), 5.41 (m, 2H), 3.54-3.28 (m, 2H), 2.93-2.87 (m, 2H), 2.54-2.45 (m, 2H), 1.95-1.86 (m, 1H), 1.05-0.99 (m, 2H), 0.77-0.72 (m, 2H)
27.6	 <p>10-([3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl]methyl)-7-thia-1,9-diazatricyclo[6.4.0.0']_{2,6}dodeca-2(6),8,10-trien-12-one</p>	381.00 ¹ H NMR (300 MHz, CD ₃ OD) δ 6.58 (s, 1H), 5.58 (s, 1H), 5.21 (m, 2H), 3.54-3.28 (m, 2H), 2.93-2.87 (m, 2H), 2.59-2.41 (m, 2H), 2.01-1.93 (m, 1H), 0.98-0.96 (m, 2H), 0.79-0.77 (m, 2H)
27.7	 <p>10-([5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl)-3,3-dimethyl-7-thia-1,9-diazatricyclo[6.4.0.0']_{2,6}dodeca-2(6),8,10-trien-12-one</p>	409.1 ¹ H NMR (300 MHz, CDCl ₃) δ 6.17 (s, 1H), 5.65 (s, 1H), 5.33 (s, 2H), 2.85 (m, 2H), 2.38 (m, 2H), 1.75 (m, 1H), 1.50 (s, 6H), 1.01 (m, 2H), 0.75 (m, 2H)
27.8	 <p>10-([3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl]methyl)-3,3-dimethyl-7-thia-1,9-diazatricyclo[6.4.0.0']_{2,6}dodeca-2(6),8,10-trien-12-one</p>	409.0 ¹ H NMR (300 MHz, CDCl ₃) δ 6.34 (s, 1H), 5.51 (s, 1H), 5.23 (s, 2H), 2.85 (m, 2H), 2.36 (m, 2H), 1.94 (m, 1H), 1.49 (s, 6H), 0.98 (m, 2H), 0.77 (m, 2H)
27.9	 <p>10-([3,5-bis(trifluoromethyl)-1H-pyrazol-1-yl]methyl)-3,3-dimethyl-7-thia-1,9-diazatricyclo[6.4.0.0']_{2,6}dodeca-2(6),8,10-trien-12-one</p>	437.0 ¹ H NMR (300 MHz, CD ₃ OD) δ 7.30 (s, 1H), 5.88 (s, 1H), 5.50 (s, 2H), 2.90 (m, 2H), 2.39 (m, 2H), 1.50 (s, 6H)


-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
27.10	 <p data-bbox="347 562 654 638">3-((3-acetyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzonitrile</p>	341.9 ¹ H NMR (400 MHz, CDCl ₃) δ 7.58-7.54 (m, 2H), 7.26-7.22 (m, 1H), 6.10 (s, 1H), 3.97 (s, 2H), 2.46 (s, 3H), 2.36 (s, 3H)
27.11	 <p data-bbox="347 877 654 982">7-(2-fluoro-3-(trifluoromethyl)benzyl)-3-(1-hydroxyethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 1)</p>	387.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.60-7.48 (m, 2H), 7.28-7.19 (m, 1H), 6.17 (s, 1H), 5.83 (m, 1H), 5.08-5.03 (m, 1H), 4.00 (s, 2H), 2.40 (s, 3H), 1.56-1.49 (m, 3H)
27.12	 <p data-bbox="347 1222 654 1318">7-(2-fluoro-3-(trifluoromethyl)benzyl)-3-(1-hydroxyethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (enantiomer 2)</p>	387.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.60-7.48 (m, 2H), 7.22-7.19 (m, 1H), 6.16 (s, 1H), 5.84 (m, 1H), 5.08-5.02 (m, 1H), 4.00 (s, 2H), 2.42 (s, 3H), 1.56-1.49 (m, 3H)
27.13	 <p data-bbox="380 1558 621 1633">3-((3-acetyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile</p>	323.95 ¹ H NMR (300 MHz, CDCl ₃) δ 7.59-7.51 (m, 3H), 7.46-7.41 (m, 1H), 6.10 (s, 1H), 3.93 (s, 2H), 2.47 (s, 3H), 2.37 (s, 3H)
27.14	 <p data-bbox="347 1852 654 1934">7-(2-fluoro-3-(trifluoromethyl)benzyl)-3-(hydroxymethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	273.10 ¹ H NMR (300 MHz, CDCl ₃) δ 7.57-7.48 (m, 2H), 7.26-7.19 (m, 1H), 6.12 (s, 1H), 4.74 (s, 2H), 4.42 (br, 1H), 3.99 (s, 2H), 2.42 (s, 3H)


-continued

No.	Structure/Name	LCMS (M + H) ¹ H NMR
27.15	 <p>7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-3-((methylamino)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one</p>	386.0 ¹ H NMR (300 MHz, CDCl ₃) δ 7.61-7.47 (m, 2H), 7.26-7.19 (m, 1H), 6.04 (s, 1H), 4.06 (s, 2H), 3.96 (s, 2H), 2.46 (s, 3H), 2.43 (s, 3H)

Example 12: 3-[2-(Hydroxymethyl)cyclopropyl]-2-methyl-7-[3-(trifluoromethyl)phenoxy]thiazolo[3,2-a]pyrimidin-5-one




Step 1: 7-Hydroxy-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 5-methyl-1,3-thiazol-2-amine (10 g, 87.6 mmol) in xylene (300 mL) was added 1,3-dimethyl propanedioate (23 g, 174 mmol), and stirred overnight at 150° C. The resulting mixture was concentrated in vacuo to afford 7-hydroxy-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (14.6 g, 91%) as a light brown solid. LCMS (ESI): M+H⁺=183.0;


Step 2: 7-Chloro-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 7-hydroxy-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (30 g, 165 mmol) in phosphorus oxy-


chloride (250 mL, 2.68 mol) was stirred for 5 h at 110° C. The resulting solution was concentrated in vacuo. The residue was purified by chromatography on silica gel eluting with dichloromethane/ethyl acetate (30/1) to afford 7-chloro-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (2.34 g, 7%) as a yellow solid. LCMS (ESI): M+H⁺=201.0.

Step 3: 2-Methyl-7-(3-(trifluoromethyl)phenoxy)-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 7-chloro-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (500 mg, 2.50 mmol) in N,N-dimethylformamide (50 mL) was added 3-(trifluoromethyl)phenol (808 mg, 5 mmol) and potassium carbonate (1.03 g, 7.5 mmol). The resulting solution was stirred overnight at 100° C. and then concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/methanol (50/1) to afford 2-methyl-7-(3-(trifluoromethyl)phenoxy)-5H-thiazolo[3,2-a]pyrimidin-5-one (500 mg, 61%) as a yellow solid. LCMS (ESI): M+H⁺=327.0.

Step 4: 3-Bromo-2-methyl-7-[3-(trifluoromethyl)phenoxy]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one




To a solution of 2-methyl-7-[3-(trifluoromethyl)phenoxy]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (500 mg, 1.53 mmol) in tetrahydrofuran (30 mL) was dropwise added n-butyllithium (1.5 mL, 2.5 mol/L) at -78° C. under an inert atmosphere of nitrogen, and the reaction was stirred for 30 mins at -78° C. 1-Bromopyrrolidine-2,5-dione (300 mg, 1.69 mmol) was added at -78° C. and the resulting solution

315


was raised slowly to room temperature. The reaction was quenched by 30 mL of water, then extracted and concentrated in vacuo. The residue was purified by chromatography on silica gel eluting with dichloromethane/petroleum ether (10/1) to afford 3-bromo-2-methyl-7-[3-(trifluoromethyl)phenoxy]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (100 mg, 13%) as an off-white solid. LCMS (ESI): $M+H^+=405.0$.

Step 5: 3-[2-(Hydroxymethyl)cyclopropyl]-2-methyl-7-[3-(trifluoromethyl)phenoxy]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one




To a solution of 3-bromo-2-methyl-7-[3-(trifluoromethyl)phenoxy]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (30 mg, 0.07 mmol) in acetonitrile/water (2 mL/0.5 mL) was added bis(diphenylphosphino)ferrocene]palladium(II) dichloride (6 mg, 0.01 mmol), sodium carbonate (16 mg, 0.15 mmol), and potassium trans-2-(hydroxymethyl)cyclopropyltrifluoroborate (26 mg, 0.15 mmol). The reaction mixture was irradiated with microwave radiation for 90 min at 120° C. and then concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/methanol (50/1) to afford 3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-7-[3-(trifluoromethyl)phenoxy]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (1.2 mg, 4%) as an off-white solid. LCMS (ESI): $M+H^+=396.9$; 1H NMR (300 MHz, $CDCl_3$) δ 7.57-7.52 (m, 2H), 7.40 (s, 1H), 7.35-7.31 (m, 1H), 5.58 (s, 1H), 4.08-4.05 (m, 1H), 3.12-3.07 (m, 1H), 2.38 (s, 3H), 2.29-2.26 (m, 1H), 1.28-1.26 (m, 1H), 1.06-0.99 (m, 2H).

Example 24: 3-[2-(Hydroxymethyl)cyclopropyl]-2-methyl-7-[4-(trifluoromethyl)thiazol-2-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one




Step 1: Methyl 2-[3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl]acetate



316


To a solution of 3-bromo-7-(chloromethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (19.4 g, 66.1 mmol) in methanol (200 mL) was added [1,1''-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (5.0 g, 6.83 mmol) and triethylamine (20 g, 197 mmol), and the reaction was placed under an atmosphere of CO (g) at 5 atm. The resulting solution was stirred for 12 h at room temperature and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/3) to afford methyl 2-[3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl]acetate (12.4 g, 59%) as a yellow solid. LCMS (ESI): $M+H^+=317.0$.

Step 2: 2-[3-Bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl]acetamide



To a solution of methyl 2-[3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl]acetate (3 g, 9.46 mmol) in methanol (20 mL) was added NH_3 in methanol (40 mL, 5 mol/L). The reaction tube was sealed and the resulting solution was stirred for 6 h at 60° C. and then concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/methanol (5/1) to afford 2-[3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl]acetamide (689 mg, 24%) as a brown solid. LCMS (ESI): $M+H^+=302.0$.


Step 3: 2-[3-Bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl]ethanethioamide



To a solution of 2-[3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl]acetamide (689 mg, 2.28 mmol) in tetrahydrofuran (20 mL) was added 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide (2.8 g, 6.92 mmol). The resulting solution was stirred for 1 h at 65° C. and then concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/ethyl acetate (3/1) to afford 2-[3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl]ethanethioamide (334 mg, 46%) as a yellow solid. LCMS (ESI): $M+H^+=318.0$.


317

Step 4: 3-Bromo-2-methyl-7-[[4-(trifluoromethyl)-1,3-thiazol-2-yl]methyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



To a solution of 2-[3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl]ethanethioamide (200 mg, 0.63 mmol) in ethanol (10 mL) was added 3-bromo-1,1,1-trifluoropropan-2-one (180 mg, 0.94 mmol). The resulting solution was stirred for 1 h at 100° C., and the resulting mixture was concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/ethyl acetate (40/1) to afford 3-bromo-2-methyl-7-[[4-(trifluoromethyl)-1,3-thiazol-2-yl]methyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (164 mg, 64%) as a yellow solid. LCMS (ESI): $M+H^+=410.0$.

Step 5: 3-[2-(Hydroxymethyl)cyclopropyl]-2-methyl-7-[[4-(trifluoromethyl)-1,3-thiazol-2-yl]methyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



To a solution of 3-bromo-2-methyl-7-[[4-(trifluoromethyl)-1,3-thiazol-2-yl]methyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (164 mg, 0.40 mmol) in acetonitrile/water (2 mL/0.5 mL) was added potassium trans-2-(hydroxymethyl)cyclopropyltrifluoroborate (143 mg, 0.80 mmol), [1,1"-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (30 mg, 0.04 mmol), and sodium carbonate (85 mg, 0.80 mmol). The reaction mixture was irradiated with microwave radiation for 1.5 h at 120° C., and the resulting mixture was concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/methanol (50/1) to afford 3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-7-[[4-(trifluoromethyl)-1,3-thiazol-2-yl]methyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (7.7 mg, 5%) as an off-white solid. LCMS (ESI): $M+H^+=402.0$; 1H NMR (300 MHz, $CDCl_3$) δ 7.71 (s, 1H), 6.22 (s, 1H), 4.30 (s, 2H), 4.08-4.04 (m, 1H), 3.16-3.09 (m, 1H), 2.39 (s, 3H), 2.30-2.22 (m, 1H), 1.28-1.07 (m, 1H), 1.05-1.03 (m, 2H).


318

Example 28: 7-[[5-Chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide


5

10

15




Step 1: Ethyl 3-bromo-2-oxobutanoate



Into each of two 3000-mL 3-necked round-bottom flasks purged and maintained with an inert atmosphere of nitrogen was placed dichloromethane (1500 mL) and ethyl 2-oxobutanoate (286 g, 2.20 mol, 1.00 equiv) followed by the addition of dibromane (352 g, 2.20 mol, 1.00 equiv) dropwise with stirring at 0-5° C. The resulting solutions were stirred at 25° C. overnight. The combined reactions were quenched by the addition of 3000 mL of saturated aqueous sodium bicarbonate. The resulting solution was extracted with 2x500 mL of dichloromethane and the combined organic layers were dried over anhydrous Na_2SO_4 and concentrated under vacuum to afford 780 g (85%) of ethyl 3-bromo-2-oxobutanoate as yellow oil.

45

Step 2: Ethyl 2-amino-5-methylthiazole-4-carboxylate hydrobromide




60

65

Into each of three 3000-mL round-bottom flasks was placed ethanol (2100 mL), thiourea (101.5 g, 1.33 mol, 1.03 equiv), and ethyl 3-bromo-2-oxobutanoate (270 g, 1.29 mol, 1.00 equiv). The resulting solutions were stirred at 85° C. in an oil bath overnight. The combined reaction mixtures were cooled to room temperature and filtered to afford 840 g (crude) of ethyl 2-amino-5-methylthiazole-4-carboxylate hydrobromide as a light yellow solid.


319

Step 3: 2-Amino-N-ethyl-5-methyl-1,3-thiazole-4-carboxamide




Into a 50000-mL pressure tank reactor was placed a solution of ethanamine in ethanol (3500 mL) and ethyl 2-amino-5-methylthiazole-4-carboxylate hydrobromide (400 g, 2.15 mol, 1.00 equiv). The resulting solution was stirred at 120° C. for 64 h. This reaction was repeated once. The combined resulting mixtures were concentrated under vacuum. The residue was applied onto a silica gel column eluting with dichloromethane/methanol (20:1) to afford 300 g (38%) of 2-amino-N-ethyl-5-methyl-1,3-thiazole-4-carboxamide as a yellow solid.

Step 4: 7-(Chloromethyl)-N-ethyl-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide



Into each of two 3000-mL 3-necked round-bottom flasks was placed PPA (1500 g), ethyl 4-chloro-3-oxobutanoate (603 g, 3.66 mol, 4.50 equiv), and 2-amino-N-ethyl-5-methyl-1,3-thiazole-4-carboxamide (150 g, 809.73 mmol, 1.00 equiv). The resulting solutions were stirred at 110° C. for 2 h. The combined reaction mixtures were cooled to 80° C. and quenched carefully by the addition of 450 mL of water. The pH of the solution was adjusted to 8 with saturated aqueous sodium carbonate. The solids were collected by filtration and washed with DCM to afford 260 g (56%) of 7-(chloromethyl)-N-ethyl-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide as a yellow solid.


Step 5: 7-[[5-Chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide




320

Into each of two 3000-mL round-bottom flasks purged and maintained with an inert atmosphere of nitrogen was placed CH₃CN (1500 mL), potassium carbonate (126 g, 911.66 mmol, 2.00 equiv), KI (38 g, 0.50 equiv), 5-chloro-3-(trifluoromethyl)-1H-pyrazole (160 g, 938.31 mmol, 2.00 equiv), and 7-(chloromethyl)-N-ethyl-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (130 g, 454.94 mmol, 1.00 equiv). The resulting solutions were stirred at 80° C. for 4 h. The combined reaction mixtures were cooled to room temperature, concentrated under vacuum, and dissolved in 8 L of ethyl acetate. The solids were filtered out. The resulting filtrate was concentrated under vacuum. The crude product was re-crystallized from EtOAc. The residue was applied onto a silica gel column eluted with petroleum ether/EtOAc/DCM (1:1:1). The crude product was purified by HPLC (Column: SO230330-2, C18, 330 g, 20-45 um, 100 Å; 254 nm, 220 nm; CH₃CN: 0.05% TFA/H₂O=40%-65%, 20 min) to give 50 g (13%) of 7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide as a white solid. LCMS (ESI): M+H⁺=420; ¹H NMR (300 MHz, CDCl₃) δ 6.58 (s, 1H), 5.93 (s, 1H), 5.72 (s, 1H), 3.47-3.56 (m, 2H), 2.42 (s, 3H), 1.25-1.30 (t, J=7.2 Hz, 3H).

Example 39: 3-[[3-(2,2-Difluorocyclopropyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluoro-benzonitrile




Step 1: 3-((3-Bromo-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzonitrile



To a solution of 3-bromo-7-(chloromethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (500 mg, 1.70 mmol) in 1,4-dioxane/water (10 mL/1 mL) in a sealed tube was added (3-cyano-2-fluorophenyl)boronic acid (420 mg, 2.55 mmol), bis(diphenylphosphino)ferrocene/palladium(II) dichloride (125 mg, 0.17 mmol), and sodium carbonate (370 mg, 3.49 mmol). The resulting solution was stirred for 12 h at 80° C., and then concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/ethyl acetate (80/1) to afford 3-((3-bromo-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzonitrile (150 mg, 23%) as a light yellow solid. LCMS (ESI): [M+H]⁺=378.0.


321

Step 2: 3-([3-Ethenyl-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl]methyl)-2-fluorobenzonitrile



To a solution of 2-ethenyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (244 mg, 1.58 mmol) in 1,4-dioxane/water (1.5 mL/0.5 mL) was added 3-([3-bromo-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl]methyl)-2-fluorobenzonitrile (300 mg, 0.79 mmol), sodium carbonate (168 mg, 1.59 mmol), and 1,1'-bis(diphenylphosphino)ferrocenepalladiumdichloride (50 mg, 0.07 mmol). The resulting solution was stirred for 2 h at 90° C. The reaction was quenched by water, then extracted with dichloromethane and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/methanol (40/1) to afford of 2-fluoro-3-([2-methyl-5-oxo-3-[2-(trifluoromethyl)cyclopropyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl]methyl)benzonitrile (178 mg, 69%) as a white solid. LCMS (ESI): M+H⁺=326.0.


Step 3: 3-[[3-(2,2-Difluorocyclopropyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluorobenzonitrile




To a solution of 3-([3-ethenyl-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl]methyl)-2-fluorobenzonitrile (80 mg, 0.25 mmol) in tetrahydrofuran (8 mL) was added sodium iodide (72 mg, 0.48 mmol) and trimethyl(trifluoromethyl)silane (176 mg, 1.29 mmol). The resulting solution was stirred for 2 h at 65° C. The reaction was quenched by water, then extracted with dichloromethane and concentrated in vacuo. The residue was purified by preparative HPLC (Column, XBridge Prep C₁₈ OBD Column, 5 μm, 19×150 mm; mobile phase, water with 10 mmol monosodium hydrogen carbonate and acetonitrile (24.0% acetonitrile up to 46.0% in 10 min); Detector, UV 254/220 nm) to afford 3-[[3-(2,2-difluorocyclopropyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluorobenzonitrile as a white solid (15.4 mg, 17%). LCMS (ESI): M+H⁺=376.0; ¹H NMR (300 MHz, CDCl₃) δ 7.61-7.53 (m, 2H), 7.26-7.20 (m, 1H), 5.99 (s, 1H), 3.93 (s, 2H), 2.80-2.74 (m, 1H), 2.40 (s, 3H), 1.98-1.84 (m, 1H), 1.60-1.46 (m, 1H).

322

Example 47: 7-[[2-Fluoro-3-(trifluoromethyl)phenyl]methyl]-3-(1H-imidazol-2-yl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one hydrate




Step 1: Methyl 7-(chloromethyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxylate



To a solution of methyl 2-amino-5-methylthiazole-4-carboxylate (5 g, 26.8 mmol) in PPA (30 mL) was added ethyl 4-chloro-3-oxobutanoate (8.82 g, 53.6 mmol), and stirred for 1 h at 110° C. The mixture was quenched by water, and the pH value of the solution was adjusted to 7 with sodium hydroxide (1 mol/L). The resulting solution was extracted with dichloromethane and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/ethyl acetate (10/1) to afford methyl 7-(chloromethyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxylate (3.2 g, 45%) as a yellow solid. LCMS (ESI): M+H⁺=273.0.

Step 2: Methyl 7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxylate




To a solution of methyl 7-(chloromethyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxylate (900 mg, 3.14 mmol) in 1,4-dioxane/water (10 mL/0.5 mL) was added 2-fluoro-3-(trifluoromethyl)phenylboronic acid (982 mg, 4.72 mmol), bis(diphenylphosphino)ferrocene]palladium(II) dichloride (150 mg, 0.21 mmol), and potassium carbonate (869 mg, 6.29 mmol). The resulting solution was stirred for 14 h at 90° C. and then concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/1) to afford methyl 7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-

323


thiazolo[3,2-a]pyrimidine-3-carboxylate (800 mg, 62%) as a yellow solid. LCMS (ESI): $M+H^+=401.0$.

Step 3: 7-(2-Fluoro-3-(trifluoromethyl)benzyl)-3-(hydroxymethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one




To a solution of methyl 7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxylate (300 mg, 0.72 mmol) in dichloromethane (10 ml) was added DIBAL-H (2 mL, 1 mol/L in toluene) at -78°C . The reaction solution was stirred for 2 h at room temperature, and then quenched by water. The resulting solution was extracted with dichloromethane and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/ethyl acetate (2/1) to afford 7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbaldehyde (136 mg, 50%) as a white solid. LCMS (ESI): $M+H^+=373.0$.

Step 4: 7-(2-Fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbaldehyde



To a solution of 7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-(hydroxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (50 mg, 0.13 mmol) in dichloromethane (5 mL) was added 1,1-bis(acetyloxy)-3-oxo-3H-1H-[5],2-benziodaoxol-1-yl acetate (85 mg, 0.20 mmol). The resulting solution was stirred overnight at room temperature. After the reaction was quenched with water, then extracted with dichloromethane and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/methanol (40/1) to afford 7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carbaldehyde (38 mg, 76%) as an off-white solid. LCMS (ESI): $M+H^+=326.0$.


Step 5: 7-[[2-Fluoro-3-(trifluoromethyl)phenyl]methyl]-3-(1H-imidazol-2-yl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one hydrate




324

To a solution of 7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carbaldehyde (38 mg, 0.10 mmol) in ethanol (2 mL) was added ammonia (0.2 mL, 5.71 mmol) and oxalaldehyde (200 mg, 3.45 mmol). The resulting solution was stirred overnight at room temperature. After the reaction was quenched with water, then extracted with dichloromethane and concentrated in vacuo. The residue was purified by preparative HPLC (Column, XBridge Prep C_{18} OBD Column, 5 μm , 19×150 mm; mobile phase, water with 10 mmol monosodium hydrogen carbonate and acetonitrile (24.0% acetonitrile up to 46.0% in 10 min); Detector, UV 254/220 nm) to afford 7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-(1H-imidazol-2-yl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one monohydrate as an off-white solid (4.5 mg, 10%). LCMS $[M+H]^+$ 371.0; ^1H NMR (300 MHz, CDCl_3) δ 10.82 (br, 1H), 7.56-7.48 (m, 2H), 7.25-7.17 (m, 3H), 6.03 (s, 1H), 5.02 (br, 2H), 3.98 (s, 2H), 2.56 (s, 3H).

Example 91: N-Ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methoxy-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide




Step 1: 2-Amino-N-ethyl-5-methoxy-1,3-thiazole-4-carboxamide



To a solution of 2-amino-5-chloro-N-ethyl-1,3-thiazole-4-carboxamide (1.00 g, 4.86 mmol) in methanol (5 mL) was added sodium methoxide (1.05 g, 19.4 mmol). The resulting solution was stirred for 2 h at 50°C . and then quenched with water. The resulting solution was extracted with dichloromethane and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/methanol (20/1) to afford of 2-amino-N-ethyl-5-methoxy-1,3-thiazole-4-carboxamide (300 mg, 31%) as an off-white solid. LCMS (ESI): $M+H^+=202.0$.


325

Step 2: 7-(Chloromethyl)-N-ethyl-2-methoxy-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide



To a solution of 2-amino-N-ethyl-5-methoxy-1,3-thiazole-4-carboxamide (2.50 g, 12.4 mmol) in polyphosphoric acid (30 mL) was added ethyl 4-chloro-3-oxobutanoate (4.09 g, 24.8 mmol). The resulting solution was stirred for 3 h at 60° C. The reaction mixture was diluted with methanol (50 mL) and the pH value of the solution was adjusted to 7 with triethylamine. The mixture was added water (200 mL), and the resulting solution was extracted with dichloromethane and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/methanol (50/1) to afford 7-(chloromethyl)-N-ethyl-2-methoxy-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (190 mg, 5%) as an off-white solid. LCMS (ESI): $M+H^+=302.0$.


Step 3: N-Ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methoxy-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide




To a solution of 7-(chloromethyl)-N-ethyl-2-methoxy-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (280 mg, 0.93 mmol) in 1,4-dioxane/water (1.5 mL/0.5 mL) was added [2-fluoro-3-(trifluoromethyl)phenyl]boronic acid (400 mg, 1.92 mmol), tricyclohexylphosphane (260 mg, 0.93 mmol), diacetoxypalladium (100 mg, 0.45 mmol), and potassium phosphate (400 mg, 1.88 mmol). The resulting solution was stirred overnight at 90° C. and then concentrated in vacuo. The residue was purified by preparative HPLC with the following conditions (1#-Pre-HPLC-005 (Waters)): Column, SunFire Prep C₁₈ OBD Column, 5 μ m, 19*150 mm; mobile phase, water with 10 mmol monosodium hydrogen carbonate and acetonitrile (50.0% acetonitrile up to 82.0% in 10 min, down to 50.0% in 2 min); Detector, UV 254/220 nm to afford N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methoxy-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (31.6 mg, 8%) as an off-white solid. LCMS (ESI): $M+H^+=430.1$; ¹H NMR (400 MHz, CD₃OD) δ 7.68-7.61 (m, 2H), 7.36-7.32 (m, 1H), 6.19 (s, 1H), 4.08 (s, 5H), 3.41-3.32 (m, 2H), 1.25-1.21 (m, 3H).

326

Example 105: 2-Cyano-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide




Step 1: 2-Amino-N-ethylthiazole-4-carboxamide



A solution of ethyl 2-aminothiazole-4-carboxylate (11.5 g, 72.7 mmol) in ethanamine/ethanol (100 mL, 30%) was stirred for 2 h at 110° C. in a sealed tube, and the resulting solution was concentrated in vacuo to afford 2-amino-N-ethylthiazole-4-carboxamide (12.3 g, 99%). LCMS (ESI): $M+H^+=172.0$.


Step 2:
2-Amino-5-chloro-N-ethylthiazole-4-carboxamide



To a solution of 2-amino-N-ethylthiazole-4-carboxamide (12.3 g, 71.8 mmol) in N,N-dimethylformamide (100 mL) was added N-chlorosuccinimide (10.5 g, 79.0 mmol). The resulting solution was stirred overnight at 50° C. and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/ethyl acetate (8/1) to afford 2-amino-5-chloro-N-ethylthiazole-4-carboxamide (7.6 g, 51%) as a brown solid. LCMS (ESI): $M+H^+=206.0$.


327

Step 3: 2-Chloro-7-(chloromethyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide



To a solution of 2-amino-5-chloro-N-ethylthiazole-4-carboxamide (12.4 g, 60.3 mmol) in polyphosphoric acid (30 mL) was added ethyl-4-chloro-3-oxobutanoate (20 g, 121 mmol). The reaction mixture was stirred for 1 h at 110° C. The reaction was quenched by water/ice, and the pH of the solution was adjusted to 7 with sodium hydroxide (1 mol/L). The resulting solution was extracted with dichloromethane and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/methanol (50/1) to afford 2-chloro-7-(chloromethyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide (4.7 g, 25%) as a brown solid. LCMS (ESI): $M+H^+$ =305.0, 307.0.


Step 4: 2-Chloro-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide



To a solution of 2-chloro-7-(chloromethyl)-N-ethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (100 mg, 0.33 mmol) in 1,4-dioxane/water (1.5 mL/0.5 mL) was added 2-fluoro-3-(trifluoromethyl)phenylboronic acid (100 mg, 0.48 mmol), potassium carbonate (90 mg, 0.65 mmol), and [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (50 mg, 0.07 mmol). The resulting solution was stirred overnight at 90° C. and then concentrated in vacuo. The residue was purified by preparative HPLC (Column, XBridge Prep C₁₈ OBD Column, 5 μm, 19×150 mm; mobile phase, water with 10 mmol monosodium hydrogen carbonate and acetonitrile (24.0% acetonitrile up to 46.0% in 10 min); Detector, UV 254/220 nm) to afford 2-chloro-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (13.4 mg, 9%) as a white solid. LCMS (ESI): $M+H^+$ =434.0; ¹H NMR (300 MHz, CDCl₃) δ 7.61-7.44 (m, 2H), 7.31-7.19 (m, 1H), 6.10 (s, 1H), 5.84 (br, 1H), 3.96 (s, 2H), 3.59-3.50 (m, 2H), 1.32-1.20 (m, 3H).


328

Step 5: Methyl 3-(ethylcarbamoyl)-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-2-carboxylate



To a solution of 2-chloro-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (500 mg, 1.15 mmol) in methanol (10 mL) was added triethylamine (233 mg, 2.30 mmol) and [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (85 mg, 0.12 mmol). The resulting solution was stirred for 6 h at 50° C. under carbon monoxide atmosphere about 10 atm and then concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/methanol (20/1) to afford methyl 3-(ethylcarbamoyl)-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-2-carboxylate (350 mg, 66%) as a brown solid. LCMS (ESI): $M+H^+$ =458.0.


Step 6: 3-(Ethylcarbamoyl)-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-2-carboxylic acid



To a solution of methyl 3-(ethylcarbamoyl)-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-2-carboxylate (170 mg, 0.37 mmol) in tetrahydrofuran/water (6 mL/2 mL) was added lithium hydroxide (110 mg, 4.59 mmol). The reaction mixture was stirred for 1 h at room temperature. The pH of the solution was adjusted to 6 with aqueous HCl solution (1 mol/L). The resulting solution was extracted with dichloromethane and concentrated in vacuo to afford 3-(ethylcarbamoyl)-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-2-carboxylic acid (110 mg, 67%) as a brown solid. LCMS (ESI): $M-H^-$ =442.0.


329

Step 7: 3-N-Ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-2,3-dicarboxamide



To a solution of 3-(ethylcarbamoyl)-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-2-carboxylic acid (300 mg, 0.68 mmol) in dichloromethane (5 mL) was added triethylamine (200 mg, 1.98 mmol) and chloro(propan-2-yloxy)methane (166 mg, 1.35 mmol) at 0° C. The mixture was stirred for 20 min at room temperature, and then ammonium hydroxide (0.5 mL, 30%) was added. The resulting solution was stirred for 2 h at room temperature and then concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/methanol (50/1) to afford 3-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-2,3-dicarboxamide (152 mg, 51%) as a brown solid. LCMS (ESI): $M+H^+=443.0$.

Step 8: 2-Cyano-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide



To a solution of 3-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-2,3-dicarboxamide (152 mg) in dichloromethane (5 mL) was added triethylamine (0.5 mL) and trifluoroacetic anhydride (0.2 mL, 0.93 mmol). The resulting solution was stirred for 30 min at room temperature and concentrated in vacuo. The residue was purified by preparative HPLC (Column, SunFire Prep C₁₈ OBD Column, 5 μm, 19*150 mm; mobile phase, Water with 10 mmol monosodium hydrogen carbonate and acetonitrile (50.0% acetonitrile up to 82.0% in 10 min, down to 50.0% in 2 min); Detector, UV 254/220 nm) to afford 2-cyano-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (7.4 mg, 5%) as an off-white solid. LCMS (ESI): $M+H^+=424.9$; ¹H NMR (300 MHz, CDCl₃) δ 7.61-7.54 (m, 1H), 7.50-7.46 (m, 1H), 7.26-7.21 (m, 1H), 6.55 (br, 1H), 6.18 (s, 1H), 3.99 (s, 2H), 3.61-3.52 (m, 2H), 1.33-1.26 (m, 3H).


330

Example 153: 7-[(5-Cyclopropyltriazol-1-yl)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one


5

10

15




Step 1: 7-(Azidomethyl)-3-bromo-2-methyl-thiazolo[3,2-a]pyrimidin-5-one



Sodium azide (134 mg, 2.04 mmol) in water (0.85 mL) was added to a solution of 3-bromo-7-(chloromethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (200 mg, 0.68 mmol) in acetonitrile (3.4 mL, 0.2 M). The mixture was stirred at room temperature for 20 h and at 50° C. for another 20 h. Water (10 mL) was added and the product was recovered by filtration. The solid was washed with cold water and heptane and was dried under vacuum to afford 7-(azidomethyl)-3-bromo-2-methyl-thiazolo[3,2-a]pyrimidin-5-one as a beige solid (167 mg, 82%). LCMS (ESI): $M+H^+=300.1, 302.1$; ¹H NMR (400 MHz, DMSO-d₆) δ 6.21 (d, J=0.8 Hz, 1H), 4.35 (s, 2H), 2.33 (s, 3H).

Step 2: 3-Bromo-7-[(5-cyclopropyltriazol-1-yl)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one




A degassed solution of chloro(pentamethylcyclopentadienyl)bis(triphenylphosphine)ruthenium(II) (12 mg, 0.01 mmol) in 1,4-dioxane (1.5 mL) with N₂ was added to a solution of 7-(azidomethyl)-3-bromo-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (89 mg, 0.29 mmol) and ethynylcyclopropane (59 mg, 0.89 mmol) in 1,4-dioxane (1.5 mL) under N₂. The mixture was capped and stirred at 60° C. for 20 h. The crude mixture was adsorbed on diatomaceous earth and

331


purified by flash chromatography (0-5% MeOH/DCM gradient) to yield 3-bromo-7-[(5-cyclopropyltriazol-1-yl)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one as a beige solid (90 mg, 83%). LCMS (ESI): $M+H^+=366.1$, 368.1 ; 1H NMR (400 MHz, Chloroform- d) δ 7.35 (s, 1H), 5.78 (s, 1H), 5.45 (s, 2H), 2.36 (s, 3H), 1.71-1.65 (m, 1H), 1.07-0.97 (m, 2H), 0.76-0.67 (m, 2H).

Step 3: 7-[(5-Cyclopropyltriazol-1-yl)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one



$Pd[dppf]Cl_2$ (18 mg, 0.02 mmol) was added to a solution of 3-bromo-7-[(5-cyclopropyltriazol-1-yl)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one (90 mg, 0.24 mmol), potassium trifluoro-[2-(hydroxymethyl)cyclopropyl]borane (65 mg, 0.37 mmol), and K_2CO_3 (68 mg, 0.49 mmol) in 1,4-dioxane/water (1.5 mL, 10:1) under N_2 in a microwave vessel. The vial was capped and heated at $120^\circ C$. for 45 min in the microwave. The crude reaction was filtered through a pad of diatomaceous earth and the filtrate was concentrated to dryness. The crude product was purified by flash chromatography (0-10% MeOH/DCM gradient) to give 7-[(5-cyclopropyltriazol-1-yl)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one as a white solid (16 mg, 18%). LCMS (ESI): $M+H^+=358.2$; 1H NMR (400 MHz, DMSO- d_6) δ 7.44 (s, 1H), 5.73 (s, 1H), 5.51 (s, 2H), 4.54 (t, $J=5.5$ Hz, 1H), 3.46 (t, $J=5.7$ Hz, 2H), 2.37 (s, 3H), 2.06-1.95 (m, 1H), 1.93-1.82 (m, 1H), 1.34-1.21 (m, 1H), 0.99-0.91 (m, 2H), 0.90-0.78 (m, 2H), 0.71-0.60 (m, 2H).

Example 206: 3-(Azetidin-1-yl)-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one




A mixture of 3-bromo-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one (50 mg, 0.10 mmol), azetidine hydrochloride (12 mg, 0.12 mmol), and K_2CO_3 (36 mg, 0.26 mmol) in


332

acetonitrile (0.5 mL) was stirred at $80^\circ C$. for 3 h and at room temperature for 20 h. The mixture was filtered to remove the salts and adsorbed on diatomaceous earth. The crude product was purified by flash chromatography (10% EtOAc/heptane) followed by a second purification by preparative HPLC to provide 3-(azetidin-1-yl)-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one is a yellow lyophilized solid (11 mg, 24%). LCMS (ESI): $M+H^+=458.0$; 1H NMR (400 MHz, DMSO- d_6) δ 7.14 (s, 1H), 5.86 (s, 1H), 5.37 (s, 2H), 4.20 (td, $J=7.6$, 7.1, 1.8 Hz, 4H), 2.24 (p, $J=7.7$ Hz, 2H).

Examples 207 and 208: 2-(7-((5-Chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-(trifluoromethyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)cyclopropanecarbonitrile (Enantiomers)




Step 1: N,N-Dimethyl-3-(trifluoromethyl)-1H-pyrazole-1-sulfonamide



Into a 20-L 4-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen was placed 3-(trifluoromethyl)-1H-pyrazole (1000 g, 7.35 mol, 1.00 equiv), CH_3CN (10 L), and 1,4-diazabicyclo[2.2.2]octane (990 g, 8.83 mol, 1.20 equiv) followed by the addition of N,N-dimethylsulfamoyl chloride (1156 g, 8.05 mol, 1.10 equiv) dropwise with stirring at $0^\circ C$. The resulting solution was stirred at room temperature for 3 h, concentrated under vacuum, diluted with 10 L of H_2O , and extracted with 3×5 L of ethyl acetate. The combined organic layers were washed with 2×5 L of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column eluted with dichloromethane to afford 1700 g (95%) of N,N-dimethyl-3-(trifluoromethyl)-1H-pyrazole-1-sulfonamide as colorless oil.


333

Step 2: 5-Chloro-N,N-dimethyl-3-(trifluoromethyl)-1H-pyrazole-1-sulfonamide



Into a 20-L 4-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen was placed N,N-dimethyl-3-(trifluoromethyl)-1H-pyrazole-1-sulfonamide (1200 g, 4.93 mol, 1.00 equiv) and tetrahydrofuran (10 L) followed by the addition of n-BuLi (2.5 M in hexane) (2.37 L, 1.20 equiv) dropwise with stirring at -78°C . The mixture was stirred at -70 to -80°C . for 1 h. To this was added a solution of C_2Cl_6 (1605 g, 1.40 equiv) in tetrahydrofuran (2.5 L) dropwise with stirring at -70°C . The resulting solution was stirred at -70°C . for 3 h, quenched by the addition of 1000 mL of saturated aqueous NH_4Cl , and extracted with 2×1 L of ethyl acetate. The combined organic layers were washed with 2×3 L of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column eluted with ethyl acetate/petroleum ether (1:30) to afford 1120 g (82%) of 5-chloro-N,N-dimethyl-3-(trifluoromethyl)-1H-pyrazole-1-sulfonamide as light yellow oil.


Step 3: 5-Chloro-3-(trifluoromethyl)-1H-pyrazole



Into a 10-L 4-necked round-bottom flask was placed 5-chloro-N,N-dimethyl-3-(trifluoromethyl)-1H-pyrazole-1-sulfonamide (2200 g, 7.92 mol, 1.00 equiv) and dichloromethane (2000 mL) followed by the addition of trifluoroacetic acid (1500 mL, 3.00 equiv) dropwise with stirring at 0°C . The resulting solution was stirred at room temperature for 5 h, concentrated under vacuum, and diluted with 6 L of H_2O . The pH of the solution was adjusted to 8-9 with sodium bicarbonate. The resulting solution was extracted with 3×4 L of ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under vacuum. The crude product was purified by distillation under reduced pressure (15 mm Hg) and the fraction was collected at 52 - 65°C . The crude product was recrystallized from DCM/n-hexane (1:50) to afford 520 g (38%) of 5-chloro-3-(trifluoromethyl)-1H-pyrazole as a white solid. LCMS (ESI): $\text{M}+\text{H}^+=171$.

334

Step 4: 2-(Tetramethyl-1,3,2-dioxaborolan-2-yl)cyclopropane-1-carbonitrile




Into a 30-mL sealed tube purged and maintained with an inert atmosphere of nitrogen a solution of cyclopropanecarbonitrile (1.0 g, 14.9 mmol) in tetrahydrofuran (12.2 mL) was added $[\text{Ir}(\text{COD})\text{OMe}]_2$ (320 mg, 0.25 mmol), bis(pinacolato) diboron (1.59 g, 12.5 mmol) and 2,9-dimethylphenanthroline (50.5 mg, 0.49 mmol). The reaction mixture was stirred at 90°C . for 18 h and concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1:4) to afford 2-(tetramethyl-1,3,2-dioxaborolan-2-yl)cyclopropane-1-carbonitrile as a light yellow oil (1 g, crude).

Step 5: Potassium 2-(cyano)cyclopropyltrifluoroborate



To a solution of 2-(tetramethyl-1,3,2-dioxaborolan-2-yl)cyclopropane-1-carbonitrile (180 g, crude) in methanol (4.5 L) was added difluorane potassium (9.98 g, 129 mmol) in H_2O (2 L). The resulting reaction mixture stirred at room temperature for 12 h and concentrated in vacuo. The residue was washed with propan-2-one (6×1.5 L). The filtrate was concentrated in vacuo, dissolved with water (5 L), and washed with DCM (3×3 L) and EtOAc (3×3 L). The water layer was freeze-dry to afford 2-(trifluoro- λ -boranyl)cyclopropane-1-carbonitrile potassium as a white solid (151.9 g, about 30% in two steps).


Step 6: 5-Bromo-1,3-thiazol-2-amine



To a mixture of sodium bicarbonate (5.8 kg, 69.04 mol, 3.00 equiv) in water (30 L) and dichloromethane (20 L) was added 5-bromo-1,3-thiazol-2-amine hydrobromide (6 kg, 23.08 mol, 1.00 equiv) in batches. The resulting mixture was stirred at room temperature for 4 h and extracted with dichloromethane. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under vacuum to afford 5-bromo-1,3-thiazol-2-amine as a gray solid (2.9 kg, 70%).


335

Step 7: 5-Bromo-2-chloro-1,3-thiazole



To a solution of 5-bromo-1,3-thiazol-2-amine (1 kg, 5.59 mol, 1.00 equiv) in CH₃CN (7 L) was added CuCl (0.83 kg, 8.4 mol, 1.5 equiv) followed by the addition of t-BuONO (1.15 kg, 11.2 mol, 2.00 equiv) dropwise with stirring. The reaction mixture was stirred at 70° C. overnight, cooled to room temperature, quenched with water, and extracted with dichloromethane. The combined organic layers were dried over anhydrous sodium sulfate and concentrated. The residue was purified by chromatography eluted with diethyl ether/petroleum ether (1/20) to afford 5-bromo-2-chloro-1,3-thiazole as a yellow solid (0.4 kg, 36%).


Step 8: 4-Bromo-2-chloro-5-iodo-1,3-thiazole



To a solution of 5-bromo-2-chloro-1,3-thiazole (500 g, 2.52 mol, 1.00 equiv) in tetrahydrofuran (8 L) was added LDA (1517 mL, 2 mol/L, 3 mmol, 1.20 equiv) dropwise with stirring at -70° C. under nitrogen atmosphere. The resulting solution was stirred at -70° C. for 2 h. To this reaction mixture was added a solution of I₂ (967 g, 3.81 mol, 1.50 equiv) in tetrahydrofuran (3 L) dropwise with stirring at -70° C. The reaction mixture was stirred at room temperature overnight, quenched with water, and extracted with dichloromethane. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under vacuum to give a residue, which was purified by chromatography eluted with ethyl acetate/petroleum ether (1:50) to afford 4-bromo-2-chloro-5-iodo-1,3-thiazole (500 g, 61%) as a gray solid.

Step 9:

4-Bromo-2-chloro-5-(trifluoromethyl)-1,3-thiazole




To a mixture of 4-bromo-2-chloro-5-iodo-1,3-thiazole (500 g, 1.54 mol, 1.00 equiv) in N,N-dimethylformamide (5 L) was added CuI (440 g, 2.31 mol, 1.50 equiv) and methyl 2,2-difluoro-2-(fluorosulfonyl)acetate (593 g, 3.09 mol, 2.00 equiv). The reaction mixture was stirred at 80° C. overnight under nitrogen atmosphere, cooled to room temperature, and filtered. The filtrate was diluted with water and extracted

336

with ether. The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum to give a residue, which was purified by chromatography eluting with diethyl ether/petroleum ether (1/20) to afford 4-bromo-2-chloro-5-(trifluoromethyl)-1,3-thiazole as a light yellow oil (300 g, crude), which was used in next step without further purification.


Step 10:

4-Bromo-5-(trifluoromethyl)-1,3-thiazol-2-amine




To a solution of 4-bromo-2-chloro-5-(trifluoromethyl)-1,3-thiazole (300 g, 1.13 mol, 1.00 equiv) in 1,4-dioxane (2 L) was added NH₃/H₂O (28%, 2 L). The resulting solution was stirred at 50° C. overnight. The resulting solution was extracted with ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under vacuum to give a residue, which was purified by chromatography eluted with ethyl acetate/petroleum ether (1:8) to afford 4-bromo-5-(trifluoromethyl)-1,3-thiazol-2-amine as a light yellow solid (160 g, 42% in 2 steps).

Step 11: 3-Bromo-7-(chloromethyl)-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



To a mixture of 4-bromo-5-(trifluoromethyl)-1,3-thiazol-2-amine (320 g, 1.30 mol, 1.00 equiv) in PPA (3200 g) was added ethyl 4-chloro-3-oxobutanoate (1068 g, 6.49 mol, 5.00 equiv). The resulting mixture was stirred at 130° C. for 2 h, quenched by the addition of water, and extracted with ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under vacuum to give a residue, which was purified by chromatography eluted with ethyl acetate/petroleum ether (1/10) to afford 3-bromo-7-(chloromethyl)-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a light yellow solid (306 g, 68%). LCMS (ESI): M+H⁺=348.9.


Step 12: 3-Bromo-7-((5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidin-5-one



337

To a mixture of 3-bromo-7-(chloromethyl)-2-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidin-5-one (1 g, 2.88 mmol) in acetonitrile (10 mL) was added sodium carbonate (610 mg, 5.75 mmol) and 5-chloro-3-(trifluoromethyl)-1H-pyrazole (590 mg, 3.45 mmol). The resulting mixture was stirred overnight at 80° C. After 30 iterations on the same scale, the mixtures were combined, then filtered and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1:20) to afford 3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidin-5-one (13.5 g, 40%) as a light yellow solid and 3-bromo-7-((3-chloro-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidin-5-one (6.0 g, 15%) a light yellow solid. LCMS (ESI): M+H⁺=480.9.

Step 13: 2-(7-[[5-Chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carbonitrile (Cis Enantiomers)




To a mixture of 3-bromo-7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (500 mg, 1.03 mmol) in 1,4-dioxane/water (10 mL/1 mL) was added [bis(diphenylphosphino)ferrocene]palladium(II) dichloride (171 mg, 0.23 mmol), sodium carbonate (500 mg, 4.68 mmol), and 2-(trifluoro-lambda4-boranyl)cyclopropane-1-carbonitrile potassium (500 mg, 2.87 mmol). The resulting mixture was stirred overnight at 85° C. The reaction was repeated 20 times on the same scale and combined. The resulting mixture was extracted with dichloromethane and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/2) to afford the racemic product (4.5 g, 46%). Then the racemic product was separated with supercritical fluid chromatography (Column: Phenomenex Lux 5u Cellulose-4,250*50 mm; Mobile Phase: CO₂:MeOH=50:50; Flow rate: 160 mL/min; detector: 220 nm) to afford two enantiomers. Enantiomer 1 (Peak 1, 1.88 g, 20%, white solid): Retention Time: 4.43 min; LCMS (ESI): M+H⁺=468.0; ¹H NMR (400 MHz, CDCl₃) δ 6.60 (s, 1H), 5.81 (s, 1H), 5.27 (s, 2H), 3.11-3.00 (m, 1H), 1.91-1.80 (m, 2H), 1.65-1.62 (m, 1H). Peak 2 (Enantiomer 2, 1.89 g, 20%, white solid): Retention Time: 5.59 min; LCMS (ESI): M+H⁺=468.0; ¹H NMR (400 MHz,


338

CDCl₃) δ 6.63 (s, 1H), 5.84 (s, 1H), 5.29 (s, 2H), 3.15-3.07 (m, 1H), 1.95-1.86 (m, 2H), 1.68-1.59 (m, 1H).

Example 220: 2-[7-[(N-Ethyl-4-fluoro-anilino)methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-1-(hydroxymethyl)cyclopropanecarbonitrile




Step 1: 7-(Chloromethyl)-3-(hydroxymethyl)thiazolo[3,2-a]pyrimidin-5-one



To methyl 7-(chloromethyl)-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxylate (0.2 g, 0.77 mmol) in 4 mL methanol at rt was added sodium borohydride (0.11 g, 2.9 mmol) over 1 min resulting in exotherm. After 5 min, additional NaBH₄ (50 mg) was added, resulting in gas evolution. After 1 h, the mixture was partitioned between CH₂Cl₂ and 1 N HCl (aq). The phases were separated and the aqueous phase was extracted with CH₂Cl₂. The combined organic phases were dried with Na₂SO₄ and concentrated onto silica gel for purification using CombiFlash® (12 g column, 0 to 80% EtOAc in CH₂Cl₂, 15 min) to afford 36 mg (20%) of 7-(chloromethyl)-3-(hydroxymethyl)thiazolo[3,2-a]pyrimidin-5-one as a tan solid.

Step 2: 7-(Chloromethyl)-5-oxo-thiazolo[3,2-a]pyrimidine-3-carbaldehyde




To 7-(chloromethyl)-3-(hydroxymethyl)thiazolo[3,2-a]pyrimidin-5-one (36 mg, 0.16 mmol) in 5 mL CH₂Cl₂ was added alumina (0.34 g, 3.3 mmol), then pyridinium chlorochromate (0.17 g, 0.78 mmol). The mixture was stirred overnight, then filtered through diatomaceous earth and concentrated onto silica gel for purification using CombiFlash® (4 g column, 0 to 80% EtOAc in CH₂Cl₂, 15 min)

339


to afford 28 mg (78%) of 7-(chloromethyl)-5-oxo-thiazolo[3,2-a]pyrimidine-3-carbaldehyde as a colorless solid.

Step 3: (E)-3-[7-(Chloromethyl)-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-2-cyano-prop-2-enoate




To a solution of 7-(chloromethyl)-5-oxo-thiazolo[3,2-a]pyrimidine-3-carbaldehyde (80 mg, 0.35 mmol) and ethyl cyanoacetate (40 mg, 0.35 mmol) in 5 mL CH₂Cl₂ was added 1 drop of piperidine resulting in a bright yellow color. After 2 h, the mixture was concentrated onto silica gel for purification using CombiFlash® (4 g column, 0 to 40% EtOAc in CH₂Cl₂, 15 min) to afford 66 mg (58%) of ethyl (E)-3-[7-(chloromethyl)-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-2-cyano-prop-2-enoate as a yellow solid.

Step 4: Ethyl 2-[7-(chloromethyl)-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-1-cyano-cyclopropanecarboxylate



To a solution of trimethylsulfoxonium iodide (69 mg, 0.31 mmol) in 1 mL DMSO at rt was added sodium hydride (60% dispersion in paraffin liquid, 12 mg, 0.31 mmol). The mixture was stirred 5 min at which time ethyl (E)-3-[7-(chloromethyl)-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-2-cyano-prop-2-enoate (66 mg, 0.20 mmol) in 1 mL DMSO was added quickly dropwise resulting in a dark orange color. The mixture was stirred 1 h, then partitioned between EtOAc and water. The phases were separated and the aqueous phase extracted with EtOAc. The combined organic phases were washed with brine, dried with Na₂SO₄, and concentrated onto silica for purification using CombiFlash® (12 g column, 0 to 80% EtOAc in CH₂Cl₂, 15 min) to afford 14 mg (20%) of ethyl 2-[7-(chloromethyl)-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-1-cyano-cyclopropanecarboxylate as a single diastereomer.


Step 5: 2-[7-(Chloromethyl)-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-1-(hydroxymethyl)cyclopropanecarbonitrile



340


To a solution of ethyl 2-[7-(chloromethyl)-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-1-cyano-cyclopropanecarboxylate (14 mg, 0.04 mmol) in 2 mL MeOH at 0° C. was added sodium borohydride (22 mg, 0.58 mmol) in one portion. The mixture was warmed to room temperature and stirred overnight. The mixture was concentrated, then partitioned between CH₂Cl₂ and 1N HCl (aq). The phases were separated, and the aqueous phase extracted with CH₂Cl₂. The combined organic phases were dried with Na₂SO₄ and concentrated. The above process was repeated (3 mL MeOH and 40 mg NaBH₄ added at room temperature) to effect complete conversion. Workup as before and concentration onto silica gel for purification using CombiFlash® (4 g column, 0 to 100% EtOAc in CH₂Cl₂, 15 min) afforded 11 mg (90%) of 2-[7-(chloromethyl)-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-1-(hydroxymethyl)cyclopropanecarbonitrile as a colorless solid.

Step 6: 2-[7-[(N-Ethyl-4-fluoro-anilino)methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-1-(hydroxymethyl)cyclopropanecarbonitrile




A mixture of N-ethyl-4-fluoro-aniline (8 mg, 0.06 mmol) and 2-[7-(chloromethyl)-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-1-(hydroxymethyl)cyclopropanecarbonitrile (11 mg, 0.04 mmol) in 2 mL acetonitrile was stirred for 3 d. The mixture was concentrated onto silica for purification using CombiFlash® (4 g column, 0 to 100% EtOAc in CH₂Cl₂, 15 min) to afford 7 mg (47%) of 2-[7-[(N-ethyl-4-fluoro-anilino)methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-1-(hydroxymethyl)cyclopropanecarbonitrile as a colorless solid. ¹H NMR (400 MHz, Chloroform-d) δ 6.93 (m, 2H), 6.7 (s, 1H), 6.61 (m, 2H), 6.25 (s, 1H), 4.33 (s, 2H), 4.16 (m, 2H), 3.48 (m, 2H), 3.32 (d, J=11.2 Hz, 1H), 3.25 (ddd, J=8.5, 7.1, 1.5 Hz, 1H), 1.76 (m, 1H), 1.55 (dd, J=8.3, 6.3 Hz, 1H), 1.24 (t, J=7.1 Hz, 3H). MS m/z 399.13 (M+H).

Example 222: 7-[[5-Chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-hydroxy-1-methyl-ethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one




341

Step 1: Methyl 2-methyl-3-oxopentanoate




To a solution of methyl 3-oxopentanoate (10 g, 76.8 mmol) in tetrahydrofuran (20 mL) was added potassium carbonate (21 g, 152 mmol) and iodomethane (12 g, 84.5 mmol). The resulting mixture was stirred for 5 h at 70° C., and then cooled to room temperature. The mixture was concentrated in vacuo to afford methyl 2-methyl-3-oxopentanoate as yellow oil (12 g, crude).

Step 2: Methyl 4-bromo-2-methyl-3-oxopentanoate



To a solution of methyl 2-methyl-3-oxopentanoate (9.00 g, 62.4 mmol) in chloroform (25 mL, 310 mmol) was added Br₂ (12 g, 75.1 mmol). The resulting solution was stirred for 12 h at 25° C., and then concentrated in vacuo to afford methyl 4-bromo-2-methyl-3-oxopentanoate (13 g, 93%) as yellow oil.


Step 3: Ethyl 2-(2-amino-5-methyl-1,3-thiazol-4-yl)propanoate



To a solution of methyl 4-bromo-2-methyl-3-oxopentanoate (10 g, 44.8 mmol) in ethanol (100 mL, 1.72 mol) was added thiourea (4.8 g, 63.1 mmol). The resulting solution was stirred for 12 h at 110° C. The reaction mixture was cooled to room temperature and then concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/3) to afford ethyl 2-(2-amino-5-methyl-1,3-thiazol-4-yl)propanoate (8 g, 83%) as a yellow oil. LCMS (ESI): M+H⁺=215.1.


342

Step 4: Ethyl 2-[7-(chloromethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]propanoate



To a solution of ethyl 2-(2-amino-5-methyl-1,3-thiazol-4-yl)propanoate (10 g, 46.7 mmol) in polyphosphoric acid (54 g, 469 mmol) was added ethyl 4-chloro-3-oxobutanoate (6.6 g, 40.1 mmol). The resulting solution was stirred for 1 h at 110° C. and then cooled to room temperature. The pH of the solution was adjusted to 7 with sodium carbonate (5%). The resulting solution was extracted with ethyl acetate and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/3) to afford ethyl 2-[7-(chloromethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]propanoate as a brown solid (13 g, 88%). LCMS (ESI): M+H⁺=315.0.


Step 5: Ethyl 2-(7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)propanoate



To a solution of ethyl 2-[7-(chloromethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]propanoate (2 g, 6.35 mmol) in acetonitrile (30 mL) was added 5-chloro-3-(trifluoromethyl)-1H-pyrazole (1.1 g, 6.45 mmol), potassium iodide (1.06 g, 6.39 mmol), and potassium carbonate (2.2 g, 15.9 mmol). The resulting mixture was stirred for 2 h at 90° C., cooled to room temperature and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/4) to afford ethyl 2-(7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)propanoate (2.9 g, crude) as a brown solid. LCMS (ESI): M+H⁺=449.0.


343

Step 6: 7-[[5-Chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-3-(1-hydroxypropan-2-yl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one




To a solution of methyl 2-(7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)propanoate (60 mg, 0.14 mmol) in dichloromethane (6 mL) was added DIBAL-H (40 mg, 0.28 mmol) at 0° C. The resulting solution was stirred for 12 h at 25° C., and the reaction was then quenched by water. The resulting solution was extracted with ethyl acetate and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/ethyl acetate (2/1) to afford 7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-3-(1-hydroxypropan-2-yl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (28.7 mg, 51%) as a light yellow oil. LCMS (ESI): M+H⁺=407.1; ¹H NMR (400 MHz, CDCl₃) δ 6.60 (s, 1H), 5.68 (s, 1H), 5.27 (s, 2H), 3.94-3.90 (m, 2H), 2.43 (s, 3H), 1.74 (m, 1H), 1.38-1.37 (m, 3H).

Example 240: 7-[[2-Fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(1H-1,2,4-triazol-5-yl)thiazolo[3,2-a]pyrimidin-5-one



Step 1. Methyl 7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxylate




Pd(dppf)Cl₂ (108 mg, 0.15 mmol) was added to a solution of methyl 7-(chloromethyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxylate (400 mg, 1.47 mmol), [2-fluoro-3-(trifluoromethyl)phenyl]boronic acid (457 mg, 2.20

344


mmol), and K₂CO₃ (405 mg, 2.93 mmol) in 1,4-dioxane/water (9 mL, 10:1) under N₂. The mixture was stirred at 90° C. for 20 h. The reaction mixture was filtered through diatomaceous earth and washed with EtOAc. The crude product was adsorbed on diatomaceous earth and purified by flash chromatography (0-60% EtOAc/heptane gradient) to afford methyl 7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxylate as an orange solid (409 mg, 70%). LCMS (ESI): M+H⁺=401.0; ¹H NMR (400 MHz, DMSO-d₆) δ 7.70 (q, J=8.1 Hz, 2H), 7.38 (t, J=7.8 Hz, 1H), 6.18 (s, 1H), 4.03 (s, 2H), 3.84 (s, 3H), 2.38 (s, 3H).

Step 2. 7-[[2-Fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxylic acid



Methyl 7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxylate (365 mg, 0.91 mmol) was dissolved in THF (9 mL) and LiOH 2 M (4.5 mL, 9.12 mmol) was added. The mixture was stirred at 60° C. for 6 h. The mixture was extracted with DCM (3×20 mL). The aqueous layer was acidified with 1 N HCl and extracted with DCM (4×20 mL). The organics were dried with MgSO₄, filtered and concentrated to give 7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxylic acid as a crude beige solid (186 mg, 53%). LCMS (ESI): M+H⁺=387.1.

Step 3. 7-[[2-Fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide




Oxalyl chloride (42 μL, 0.47 mmol) was added to a solution of 7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxylic acid (90 mg, 0.23 mmol) and DMF (2 μL, 0.02 mmol) in DCM (1.5 mL) at 0° C. The mixture was stirred at room temperature for 30 min and then the reaction mixture was concentrated to dryness. The solvent was switched for THF (0.6 mL) and a solution of ammonia in 1,4-dioxane (0.5 mol/L, 4 mL, 1.86 mmol) was added at 0° C. The mixture was stirred at room temperature for 30 min. The reaction was partitioned in water/DCM and extracted with DCM (3×10 mL). The organics were washed with brine, dried with MgSO₄, filtered and concentrated to obtain 7-[[2-fluoro-3-

345


(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo [3,2-a]pyrimidine-3-carboxamide as a crude pale yellow solid (73 mg, 81%). LCMS (ESI): $M+H^+=386.2$.

Step 4: 7-[[2-Fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(1H-1,2,4-triazol-5-yl)thiazolo [3,2-a]pyrimidin-5-one




7-[[2-Fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide (73 mg, 0.19 mmol) in *N,N*-dimethylformamide dimethyl acetal (1.9 mL, 14.2 mmol) was stirred at 100° C. for 4 h. The mixture was concentrated, treated with acetic acid (1.9 mL, 32.7 mmol) and hydrazine (120 μ L, 3.78 mmol), and stirred at 100° C. for 2 h. The reaction mixture was concentrated and purified by preparative HPLC and lyophilization to provide 7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(1H-1,2,4-triazol-5-yl)thiazolo[3,2-a]pyrimidin-5-one as a beige solid (19 mg, 25%). LCMS (ESI): $M+H^+=410.1$; 1H NMR (~2:1 triazole tautomer ratio, * denotes minor tautomer peaks, 400 MHz, DMSO- d_6) δ 14.19 (s, 1H), 8.52 (s, 1H), 7.79-7.65 (m, 2H), 7.39 (t, $J=7.8$ Hz, 1H), 6.13* (s, 0.3H), 6.04 (s, 0.7H), 4.02 (s, 2H), 2.33* (s, 1H), 2.18 (s, 2H).

Example 244: 3-Acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methoxy-thiazolo[3,2-a]pyrimidin-5-one



Step 1: 5-Methoxythiazol-2-amine




To a solution of 5-bromo-1,3-thiazol-2-amine hydrobromide (26 g, 100 mmol) in methanol (100 mL) was added dropwise sodium methoxide (12 g, 222 mmol) in 40 mL methanol at 0° C. The resulting solution was stirred for 1 h at room temperature and the reaction was diluted with ethyl

346


acetate. The solids were filtered out and the filtrate was concentrated in vacuo to afford 5-methoxy-1,3-thiazol-2-amine (6.5 g, crude) as a tan solid. LCMS (ESI): $M+H^+=131.0$.

Step 2:
2-(5-Methoxythiazol-2-yl)isoindoline-1,3-dione




To a solution of 5-methoxy-1,3-thiazol-2-amine (6.50 g, crude) in acetonitrile (100 mL) was added ethyl 1,3-dioxo-2,3-dihydro-1H-isoindole-2-carboxylate (10.9 g, 49.9 mmol). The resulting solution was stirred overnight at 50° C. and then concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with dichloromethane/petroleum ether (3/1) to afford 2-(5-methoxy-1,3-thiazol-2-yl)-2,3-dihydro-1H-isoindole-1,3-dione (5 g, 38%) as a light brown solid. LCMS (ESI): $M+H^+=261.0$.

Step 3: 2-(4-Chloro-5-methoxythiazol-2-yl)isoindoline-1,3-dione



A solution of 2-(5-methoxy-1,3-thiazol-2-yl)-2,3-dihydro-1H-isoindole-1,3-dione (4.00 g, 15.4 mmol) in acetonitrile (100 mL) was treated with *N*-chlorosuccinimide (2.16 g, 16.2 mmol) and then stirred for 2 h at 90° C. The resulting mixture was concentrated in vacuo to afford 2-(4-chloro-5-methoxy-1,3-thiazol-2-yl)-2,3-dihydro-1H-isoindole-1,3-dione (4.6 g, crude) as an orange solid. LCMS (ESI): $M+H^+=295.0$.


Step 4: 4-Chloro-5-methoxythiazol-2-amine



To a solution of 2-(4-chloro-5-methoxy-1,3-thiazol-2-yl)-2,3-dihydro-1H-isoindole-1,3-dione (4.6 g, 15.6 mmol) in methanol (50 mL) was added $NH_2NH_2 \cdot H_2O$ (15 mL, 308 mmol). The resulting solution was stirred for 4 h at room temperature. The solids were filtered out and the filtrate was concentrated in vacuo to afford 4-chloro-5-methoxy-1,3-thiazol-2-amine (2 g, 78%) as a light brown solid. LCMS (ESI): $M+H^+=165.0$.


347

Step 5: 3-Chloro-7-(chloromethyl)-2-methoxy-5H-thiazolo[3,2-a]pyrimidin-5-one




To a solution of 4-chloro-5-methoxy-1,3-thiazol-2-amine (2.00 g, 12.1 mmol) in PPA (30 g, 260 mmol) was added ethyl 4-chloro-3-oxobutanoate (6.00 g, 36.4 mmol). The resulting solution was stirred for 2 h at 60° C. and then quenched with water/ice. The resulting solution was extracted with dichloromethane and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/3) to afford 3-chloro-7-(chloromethyl)-2-methoxy-5H-thiazolo[3,2-a]pyrimidin-5-one (1.5 g, 47%) as a light brown solid. LCMS (ESI): M+H⁺=265.0.

Step 6: 3-Chloro-7-((5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methoxy-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 3-chloro-7-(chloromethyl)-2-methoxy-5H-thiazolo[3,2-a]pyrimidin-5-one (100 mg, 0.38 mmol) in acetonitrile (2 mL) was added 5-chloro-3-(trifluoromethyl)-1H-pyrazole (78 mg, 0.46 mmol) and potassium carbonate (104 mg, 0.75 mmol). The resulting solution was stirred for 2 h at 80° C. and then concentrated in vacuo. The residue was purified by preparative TLC with ethyl acetate/petroleum ether (1/2) to afford 3-chloro-7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methoxy-5H-thiazolo[3,2-a]pyrimidin-5-one (24.4 mg, 16%) as a white solid. LCMS (ESI): M+H⁺=399.9; ¹H NMR (300 MHz, CDCl₃) δ 6.58 (s, 1H), 5.73 (s, 1H), 5.23 (s, 2H), 4.02 (s, 3H).


Step 7: 7-((5-Chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(1-ethoxyvinyl)-2-methoxy-5H-thiazolo[3,2-a]pyrimidin-5-one



348


To a solution of 3-chloro-7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-methoxy-5H-thiazolo[3,2-a]pyrimidin-5-one (300 mg, 0.752 mmol) in 1,4-dioxane (8 mL) was added tributyl(1-ethoxyethyl)stannane (543 mg, 1.504 mmol), DIEA (194 mg, 1.50 mmol), and bis(diphenylphosphino)ferrocene]palladium(II) dichloride (106 mg, 0.151 mmol). The resulting solution was stirred overnight at 80° C. and then concentrated in vacuo. The residue was purified by preparative TLC with ethyl acetate/petroleum ether (1:2) to afford 7-((5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(1-ethoxyvinyl)-2-methoxy-5H-thiazolo[3,2-a]pyrimidin-5-one (180 mg, 55%) as light yellow oil. LCMS (ESI): M+H⁺=435.0.

Step 8: 3-Acetyl-7-((5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methoxy-5H-thiazolo[3,2-a]pyrimidin-5-one




To a solution of 7-((5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(1-ethoxyvinyl)-2-methoxy-5H-thiazolo[3,2-a]pyrimidin-5-one (180 mg, 0.414 mmol) in dichloromethane (10 mL) was added a solution of hydrogen chloride in 1,4-dioxane (0.5 mL, saturated). The resulting solution was stirred for 30 min at room temperature and then concentrated in vacuo. The residue was purified by preparative TLC with ethyl acetate/petroleum ether (1/1) to afford 3-acetyl-7-((5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methoxy-5H-thiazolo[3,2-a]pyrimidin-5-one (64.7 mg, 38%) as a white solid. LCMS (ESI): M+H⁺=407.0; ¹H NMR (300 MHz, CDCl₃) δ 6.58 (s, 1H), 5.81 (s, 1H), 5.27 (s, 2H), 4.05 (s, 3H), 2.50 (s, 3H).

Example 250: 7-[[5-Chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-propanoyl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one




349

Step 1: Methyl 2-[[tert-butoxy]carbonyl]amino]-1,3-thiazole-4-carboxylate




To a solution of 2-amino-1,3-thiazole-4-carboxylate (20 g, 126 mmol) and 4-dimethylaminopyridine (1.54 g, 12.6 mmol) in dichloromethane/tetrahydrofuran (200 mL/200 mL) was added di-tert-butyl dicarbonate (33 g, 151 mmol). The resulting solution was stirred for 12 h at room temperature and then concentrated in vacuo. The residue was purified by chromatography with ethyl acetate/petroleum ether (1/2) to afford methyl 2-[[tert-butoxy]carbonyl]amino]-1,3-thiazole-4-carboxylate (28 g, 86%) as an off-white solid. LCMS (ESI): $M+H^+=259.0$.

Step 2: 2-[[tert-Butoxy]carbonyl]amino]-1,3-thiazole-4-carboxylic acid



To a solution of methyl 2-[[tert-butoxy]carbonyl]amino]-1,3-thiazole-4-carboxylate (28 g, 108 mmol) in tetrahydrofuran (300 mL) was added a solution of lithium hydroxide (10.4 g, 433 mmol) in water (150 mL). The resulting mixture was stirred for 12 h at room temperature. The pH of the solution was adjusted to 4 with hydrochloric acid (2 mol/L). The solids were collected by filtration to afford 2-[[tert-butoxy]carbonyl]amino]-1,3-thiazole-4-carboxylic acid (24 g, 76%) as an off-white solid. LCMS (ESI): $M+H^+=245.0$.

Step 3: tert-Butyl N-[4-[methoxy(methyl)carbamoyl]-1,3-thiazol-2-yl]carbamate




To a mixture of 2-[[tert-butoxy]carbonyl]amino]-1,3-thiazole-4-carboxylic acid (20.0 g, 81.9 mmol) in dichloromethane (400 mL) was added methoxy(methyl)amine hydrochloride (16.0 g, 164 mmol), HATU (37.4 g, 98.3 mmol), and triethylamine (16.6 g, 164 mmol). The resulting mixture was stirred for 4 h at room temperature. The reaction

350


was then quenched by water, then extracted with ethyl acetate and concentrated in vacuo to afford tert-butyl N-[4-[methoxy(methyl)carbamoyl]-1,3-thiazol-2-yl]carbamate (20 g, 85%) as a light red solid. LCMS (ESI): $M+H^+=288.0$.

Step 4: tert-Butyl N-(4-propanoyl-1,3-thiazol-2-yl)carbamate




To a solution of tert-butyl N-[4-[methoxy(methyl)carbamoyl]-1,3-thiazol-2-yl]carbamate (5.00 g, 17.4 mmol) in tetrahydrofuran (100 mL, 1.23 mol) was added bromo(ethyl)magnesium (4.59 mL, 34.8 mmol) at -70°C . The resulting solution was stirred for 12 h at room temperature. The reaction was diluted with saturated aqueous NH_4Cl (30 mL), extracted with ethyl acetate, and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/3) to afford tert-butyl N-(4-propanoyl-1,3-thiazol-2-yl)carbamate (1.3 g, 29%) as a light yellow solid. LCMS (ESI): $M+H^+=257.0$.

Step 5: 1-(2-Amino-1,3-thiazol-4-yl)propan-1-one



To a solution of tert-butyl N-(4-propanoyl-1,3-thiazol-2-yl)carbamate (1.3 g, 5.07 mmol) in dichloromethane (15 mL) was added trifluoroacetic acid (24 mL). The resulting solution was stirred for 2 h at 0°C and then concentrated in vacuo. The residue was dissolved in dichloromethane and then washed with sodium bicarbonate (1 mol/L). The organic layers were concentrated in vacuo to afford 1-(2-amino-1,3-thiazol-4-yl)propan-1-one (750 mg, 90%) as a light yellow solid. LCMS (ESI): $M+H^+=157.0$.


Step 6: 1-(2-Amino-5-iodo-1,3-thiazol-4-yl)propan-1-one



351


To a solution of 1-(2-amino-1,3-thiazol-4-yl)propan-1-one (850 mg, 5.44 mmol) in dichloromethane (20 mL) was added N-iodo-succinimide (1.35 g, 5.99 mmol). The resulting mixture was stirred for 12 h at room temperature and then concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/3) to afford 1-(2-amino-5-iodo-1,3-thiazol-4-yl)propan-1-one (1 g, 65%) as a brown solid. LCMS (ESI): $M+H^+=283.0$.

Step 7: 2-(5-Iodo-4-propanoyl-1,3-thiazol-2-yl)-2,3-dihydro-1H-isoindole-1,3-dione



To a solution of 1-(2-amino-5-iodo-1,3-thiazol-4-yl)propan-1-one (500 mg, 1.77 mmol) and triethylamine (89.7 mg, 0.89 mmol) in dichloromethane (30 mL) was added ethyl 1,3-dioxo-2,3-dihydro-1H-isoindole-2-carboxylate (777 mg, 3.54 mmol). The resulting mixture was stirred for 12 h at 40° C., and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/2) to afford 2-(5-iodo-4-propanoyl-1,3-thiazol-2-yl)-2,3-dihydro-1H-isoindole-1,3-dione (450 mg, 62%) as a light red solid. LCMS (ESI): $M+H^+=413.0$.

Step 8: 2-(5-Iodo-4-propanoyl-1,3-thiazol-2-yl)-2,3-dihydro-1H-isoindole-1,3-dione



To a solution of 2-(5-iodo-4-propanoyl-1,3-thiazol-2-yl)-2,3-dihydro-1H-isoindole-1,3-dione (450 mg, 1.09 mmol) in N,N-dimethylformamide (10 mL) was added ethyl 2,2-difluoro-2-(fluorosulfonyl)acetate (450 mg, 2.18 mmol) and copper(I) iodide (416 mg, 2.18 mmol). The resulting mixture was stirred for 2 h at 80° C. and then concentrated in vacuo. The residue was dissolved in dichloromethane (20 mL), and the solids were filtered out. The resulting solution was concentrated in vacuo to afford 2-[4-propanoyl-5-(trifluoromethyl)-1,3-thiazol-2-yl]-2,3-dihydro-1H-isoindole-1,3-dione (230 mg, 59%) as a yellow solid. LCMS (ESI): $M+H^+=355.0$.


352

Step 9: 1-[2-Amino-5-(trifluoromethyl)-1,3-thiazol-4-yl]propan-1-one




To a solution of 2-[4-propanoyl-5-(trifluoromethyl)-1,3-thiazol-2-yl]-2,3-dihydro-1H-isoindole-1,3-dione (230 mg, 0.65 mmol) in acetonitrile (10 mL) was added hydrazine monohydrate (0.31 mL, 6.38 mmol). The resulting solution was stirred for 30 min at room temperature. After concentration, the residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/1) to afford 1-[2-amino-5-(trifluoromethyl)-1,3-thiazol-4-yl]propan-1-one (60 mg, 41%) as a light yellow oil. LCMS (ESI): $M+H^+=225.0$.

Step 10: 7-(Chloromethyl)-3-propanoyl-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



To a mixture of 1-[2-amino-5-(trifluoromethyl)-1,3-thiazol-4-yl]propan-1-one (60 mg, 0.27 mmol) in polyphosphoric acid (1 g, 8.69 mmol) was added ethyl 4-chloro-3-oxobutanoate (220 mg, 1.34 mmol). The resulting mixture was stirred for 12 h at 130° C. The reaction was then quenched by water (25 mL), extracted with ethyl acetate and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/2) to afford 7-(chloromethyl)-3-propanoyl-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (15 mg, 17%) as a yellow solid. LCMS (ESI): $M+H^+=325.0$.


Step 11: 7-[[5-Chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-3-propanoyl-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one




353

To a mixture of 7-(chloromethyl)-3-propanoyl-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (15 mg, 0.05 mmol) in acetonitrile (3 mL, 57.1 mmol) was added potassium carbonate (13 mg, 0.09 mmol) and 5-chloro-3-(trifluoromethyl)-1H-pyrazole (10 mg, 0.06 mmol). The resulting mixture was stirred for 2 h at 80° C. After filtration and concentration, the residue was purified by chromatography with ethyl acetate/petroleum ether (1/3) to afford 7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-3-propanoyl-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (4.5 mg, 21%) as a light brown solid. LCMS (ESI): $M+H^+=459.0$; 1H NMR (400 MHz, $CDCl_3$) δ 6.60 (s, 1H), 5.83 (s, 1H), 5.31 (s, 2H), 2.97-2.83 (m, 2H), 1.29-1.25 (m, 3H).

Example 251: 2-[7-[(3,5-Dichloropyrazol-1-yl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile




Step 1: N,N-Dimethyl-1H-pyrazole-1-sulfonamide



A solution of 1H-pyrazole (30 g, 440 mmol) in tetrahydrofuran (500 mL) was treated with sodium hydride (26 g, 648 mmol, 60%) at 0° C., and then stirred for 1 h at 0° C. N,N-Dimethylsulfamoyl chloride (95 g, 661 mmol) was added dropwise at 0° C. The resulting solution was stirred for additional 2 h at room temperature and then quenched by water. The resulting solution was extracted with dichloromethane and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/6) to afford N,N-dimethyl-1H-pyrazole-1-sulfonamide (58 g, 75%) as colorless oil. LCMS (ESI): $M+H^+=175.0$.

Step 2:


5-Chloro-N,N-dimethyl-1H-pyrazole-1-sulfonamide



354


To a solution of N,N-dimethyl-1H-pyrazole-1-sulfonamide (37.2 g, 212 mmol) in tetrahydrofuran (600 mL) was added dropwise n-BuLi (127 mL, 2.5 mmol/L) at -78° C. The resulting solution was stirred for 1 h at -78° C. Hexachloroethane (75.4 g, 318 mmol) in tetrahydrofuran (400 mL) was added dropwise at -78° C. The reaction mixture was allowed to warm to room temperature and was stirred overnight at room temperature. The reaction was then quenched by water, then extracted with dichloromethane and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/6) to afford 5-chloro-N,N-dimethyl-1H-pyrazole-1-sulfonamide as a red oil (39.5 g, 89%). LCMS (ESI): $M+H^+=209.0$.

Step 3: 5-Chloro-1H-pyrazole




To a solution of 5-chloro-N,N-dimethyl-1H-pyrazole-1-sulfonamide (30 g, 143 mmol) in dichloromethane (500 mL) was added trifluoroacetic acid (45.7 g, 401 mmol). The reaction mixture was stirred for 2 h at room temperature and quenched by water. The pH of the solution was adjusted to 8 with saturated sodium bicarbonate. The resulting solution was extracted with dichloromethane and concentrated in vacuo to afford 5-chloro-1H-pyrazole as a reddish solid (14 g, 95%). LCMS (ESI): $M+H^+=103.0$.

Step 4: 5-Chloro-1-nitro-1H-pyrazole



To a solution of 5-chloro-1H-pyrazole (14 g, 136 mmol) in acetic acid/acetic anhydride (36 mL/92 mL) was added fuming nitric acid (36 mL). The resulting solution was stirred overnight at room temperature, and then diluted with water (500 mL). The solids were collected by filtration to afford 5-chloro-1-nitro-1H-pyrazole as a yellow solid (7 g, 35%).


Step 5: 5-Chloro-3-nitro-1H-pyrazole



355


A solution of 5-chloro-1-nitro-1H-pyrazole (3 g, 20.3 mmol) in anisole (53.6 mL) was stirred overnight at 130° C. The resulting solution was diluted with of H₂O:petroleum ether (1:1), then extracted with sodium hydroxide (10%) and the aqueous layers combined. The pH of the solution was adjusted to 2 with hydrochloric acid (3 mol/L). The resulting solution was extracted with ethyl acetate and concentrated in vacuo to afford 5-chloro-3-nitro-1H-pyrazole as a yellow solid (2.7 g, 90%).

Step 6: 3-Bromo-7-[(5-chloro-3-nitro-1H-pyrazol-1-yl)methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



To a solution of 3-bromo-7-(chloromethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (400 mg, 1.36 mmol) in acetonitrile (50 mL) was added 5-chloro-3-nitro-1H-pyrazole (211 mg, 1.43 mmol), KI (113 mg, 0.68 mmol), and potassium carbonate (565 mg, 4.09 mmol). The resulting solution was stirred for 2 h at 80° C., cooled, extracted with dichloromethane, and then concentrated in vacuo to afford 3-bromo-7-[(5-chloro-3-nitro-1H-pyrazol-1-yl)methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a gray solid (20 mg, 4%). LCMS (ESI): M+H⁺=405.0.


Step 7: 7-[(3-Amino-5-chloro-1H-pyrazol-1-yl)methyl]-3-bromo-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 3-bromo-7-[(5-chloro-3-nitro-1H-pyrazol-1-yl)methyl]-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (30 mg, 0.066 mmol) in ethanol/water (5 mL/1 mL) was added iron powder (29 mg, 0.52 mmol) and ammonium chloride (35 mg, 0.660 mmol). The reaction mixture was stirred for 2 h at 80° C., cooled and extracted with dichloromethane. The combined organic phase was concentrated in vacuo to afford 7-[(3-amino-5-chloro-1H-pyrazol-1-yl)methyl]-3-bromo-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one as a gray solid (10 mg, 4%). LCMS (ESI): M+H⁺=375.0.


356

Step 8: 3-Bromo-7-[(3,5-dichloro-1H-pyrazol-1-yl)methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



To a solution of 7-[(5-amino-3-chloro-1H-pyrazol-1-yl)methyl]-3-bromo-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (1 g, 2.67 mmol) in acetonitrile (10 mL) was added tert-butyl nitrite (495 mg, 4.80 mmol) and copper(I) chloride (715 mg, 5.32 mmol). The resulting mixture was stirred for 1 h at 25° C. and then concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/1) to afford 3-bromo-7-[(3,5-dichloro-1H-pyrazol-1-yl)methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (200 mg, 19%) as a yellow solid. LCMS (ESI): M+H⁺=395.0.


Step 9: 2-[7-[(3,5-Dichloro-1H-pyrazol-1-yl)methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carbonitrile




To a solution of 3-bromo-7-[(3,5-dichloro-1H-pyrazol-1-yl)methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (150 mg, 0.38 mmol) in acetonitrile/water (4 mL/0.4 mL) was added 2-(trifluoro-lambda4-boranyl)cyclopropane-1-carbonitrile potassium (263 mg, 1.52 mmol), sodium carbonate (80 mg, 0.75 mmol), and [bis(diphenylphosphino)ferrocene]palladium(II) dichloride (27 mg, 0.037 mmol). The resulting solution was stirred for 14 h at 90° C. and then concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/1) to afford the racemic product (45 mg, 31%). The racemic product was separated by chiral HPLC with the following conditions (Column: Chiralpak ic 0.46*25 cm, 5 um; Mobile Phase: 100% MeOH-HPLC; Flow rate: 1 mL/min; detector: 254 nm) to afford the title compound as a white solid (13.3 mg, 9%). Retention Time: 9.18 min; LCMS (ESI): M+H⁺=381.0; ¹H NMR (300 MHz, CDCl₃) δ 6.25 (s, 1H), 5.71 (s, 1H), 5.14 (s, 2H), 2.99-2.94 (m, 1H), 2.39 (s, 3H), 1.84-1.79 (m, 1H), 1.69-1.66 (m, 1H), 1.45-1.35 (m 1H).

357

Example 258: 7-[[5-Chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(difluoromethyl)-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide




Step 1: 2-Chloro-7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-N-ethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide



To a solution of 2-chloro-7-(chloromethyl)-N-ethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (2 g, 6.53 mmol) in acetonitrile (10 mL) was added 5-chloro-3-(trifluoromethyl)-1H-pyrazole (872 mg, 5.11 mmol), potassium iodide (542 mg, 3.26 mmol), and potassium carbonate (1.8 g, 13 mmol). The resulting mixture was stirred for 1 h at 80° C. and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/1) to afford of 2-chloro-7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-N-ethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (1.1 g, 38%) as a yellow solid. LCMS (ESI): M+H⁺=441.0.

Step 2: 7-[[5-Chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-ethenyl-N-ethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide




To a solution of 2-chloro-7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-N-ethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (500 mg, 1.14

358


mmol) in 1,4-dioxane/water (15 mL/1 mL) was added 2-ethenyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (349 mg, 2.27 mmol), sodium carbonate, (238 mg, 2.25 mmol) and [bis(diphenylphosphino)ferrocene]palladium(II) dichloride (83 mg, 0.11 mmol). The resulting solution was stirred for 14 h at 90° C. and then concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/1) to afford 7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-ethenyl-N-ethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide as a yellow solid (200 mg, 41%). LCMS (ESI): M+H⁺=432.0.

Step 3: 7-[[5-Chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-N-ethyl-2-formyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide



To a solution of 7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-ethenyl-N-ethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (175 mg, 0.41 mmol) in 1,4-dioxane/water (5 mL/3 mL) was added osmium tetroxide (1.03 mg, 0.004 mmol), N-methylmorpholine-N-oxide (94.83 mg, 0.811 mmol), and sodium periodate (173 mg, 0.81 mmol). The resulting solution was stirred for 14 h at 25° C. and concentrated. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/2) to afford 7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-N-ethyl-2-formyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (50 mg 28%) as a light yellow solid. LCMS (ESI): M+H⁺=456.0.

Step 4: 7-((5-Chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-(difluoromethyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide




To a solution of 7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-N-ethyl-2-formyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide (100 mg, 0.23 mmol) in dichloromethane (20 mL) was added bis(2-methoxyethyl)amino)sulfur trifluoride (510 mg, 2.31 mmol) at 0° C. The resulting solution was stirred for 14 h at 25° C., and then quenched by water. The resulting solution was


359

extracted with ethyl acetate and concentrated in vacuo to afford 7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-2-(difluoromethyl)-N-ethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide as a white solid (21.2 mg, 20%). LCMS (ESI): $M+H^+=456.0$; 1H NMR (400 MHz, $CDCl_3$) δ 7.10-6.83 (m, 1H), 6.59 (s, 1H), 6.37 (s, 1H), 5.80 (s, 1H), 5.30 (s, 2H), 3.54-3.47 (m, 2H), 1.28-1.25 (m, 3H).

Examples 266 and 267: 2-[2-Chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (Cis Enantiomers)




Step 1: tert-Butyl N-(4-bromo-5-chloro-1,3-thiazol-2-yl)carbamate



To a solution of tert-butyl N-(5-bromo-1,3-thiazol-2-yl) carbamate (5 g, 17.9 mmol) in tetrahydrofuran (100 ml) was added dropwise LDA (29.4 ml, 2 mol/L) at $-78^\circ C$, and the resulting mixture was stirred for 1 h at $-78^\circ C$. Then the mixture was added a solution of hexachloroethane (14 g, 59.1 mmol) in tetrahydrofuran (50 ml) at $-78^\circ C$. The reaction was stirred for additional 15 h at room temperature. The reaction was quenched by water, then extracted with dichloromethane and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/9) to afford tert-butyl N-(4-bromo-5-chloro-1,3-thiazol-2-yl)carbamate (4.07 g, 72%) as brown oil. LCMS (ESI): $M+H^+=313.0$.

Step 2: 4-Bromo-5-chloro-1,3-thiazol-2-amine




To a solution of tert-butyl N-(4-bromo-5-chloro-1,3-thiazol-2-yl)carbamate (4.07 g, 13.0 mmol) in dichloromethane (20 mL) was added trifluoroacetic acid (29.7 g, 260 mmol), and the reaction was stirred for 15 h at room temperature. The pH of the solution was adjusted to 7 with saturated

360


sodium bicarbonate, then extracted with dichloromethane and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/5) to afford 4-bromo-5-chloro-1,3-thiazol-2-amine (1.02 g, 37%) as a brown solid. LCMS (ESI): $M+H^+=213.0$.

Step 3: 3-Bromo-2-chloro-7-(chloromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one



To a solution of 4-bromo-5-chloro-1,3-thiazol-2-amine (700 mg, 3.28 mmol) in polyphosphoric acid (2.81 g, 24.4 mmol) was added ethyl 4-chloro-3-oxobutanoate (1.08 g, 6.56 mmol). The reaction mixture was stirred for 1 h at $110^\circ C$ and cooled to room temperature. The reaction was then quenched by water and the pH of the solution was adjusted to 7 with sodium hydroxide (1 mol/L). The resulting solution was extracted with dichloromethane and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/3) to afford 3-bromo-2-chloro-7-(chloromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (900 mg, 87%) as a brown solid. LCMS (ESI): $M+H^+=313.0$.


Step 4: 3-Bromo-2-chloro-7-((5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one



To a solution of 3-bromo-2-chloro-7-(chloromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (500 mg, 1.59 mmol) in acetonitrile (10 mL) was added 5-chloro-3-(trifluoromethyl)-1H-pyrazole (327 mg, 1.92 mmol), potassium iodide (133 mg, 0.80 mmol), and potassium carbonate (442 mg, 3.20 mmol). The resulting solution was stirred for 2 h at $80^\circ C$ and cooled. The solid was filtered off and the filtrate was concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/9) to afford 3-bromo-2-chloro-7-(chloromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as a brown solid (300 mg, 60%). LCMS (ESI): $M+H^+=448.0$.

361

Step 5: 2-(2-Chloro-7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)cyclopropane-1-carbonitrile (Enantiomer 1)



To a solution of 3-bromo-2-chloro-7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (100 mg, 0.22 mmol) in 1,4-dioxane/water (3 mL/0.3 mL) was added potassium 2-(cyano) cyclopropyltrifluoroborate (77.2 mg, 0.45 mmol), [bis (diphenylphosphino)ferrocene]palladium(II) dichloride

362

(16.3 mg, 0.022 mmol), and potassium phosphate (94.8 mg, 0.45 mmol). The resulting solution was stirred for 14 h at 85° C. and concentrated in vacuo. The residue was purified by flash chromatography on silica gel eluting with ethyl acetate/petroleum ether (1/1) to afford the racemic product (40 mg, 41%). Then the racemic product was separated with chiral HPLC with the following conditions (Column: Phenomenex Lux 5u Cellulose-4, AXIA Packed 250*21.2 mm, 5 um; Mobile Phase: 100% MeOH; Flow rate: 20 mL/min; detector: 254 nm/220 nm) to afford two enantiomers. Enantiomer 1 (Peak 1, white solid, 7.5 mg, 8%): Retention Time: 1.54 min; LCMS (ESI): M+H⁺=434.0; ¹H NMR (400 MHz, CDCl₃) δ 6.59 (s, 1H), 5.74 (s, 1H), 5.24 (s, 2H), 2.98-2.93 (m, 1H), 1.88-1.82 (m, 2H), 1.66-1.61 (m, 1H). Enantiomer 2 (Peak 2, 6.1 mg, 6%): Retention Time: 2.06 min; LCMS (ESI): M+H⁺=434.0; ¹H NMR (400 MHz, CDCl₃) δ 6.59 (s, 1H), 5.75 (s, 1H), 5.24 (s, 2H), 2.99-2.89 (m, 1H), 1.88-1.82 (m, 2H), 1.66-1.61 (m, 1H).

The following examples were prepared using methods analogous to those described in the appropriate Reference Method or Example (Ref. Method or Ex.) column. Satisfactory analytical data was obtained for each compound.

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
1		7-[(3-cyclopropyl-2-fluorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
2		N-ethyl-2-methyl-5-oxo-7-[(2,3,6-trifluorophenyl)methyl]thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
3		2-fluoro-3-[(2-methyl-3-oxazol-2-yl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl)methyl]benzonitrile	Method 24
4		7-[(5-cyano-3-cyclopropyl-2-fluorophenyl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
5		N-ethyl-7-[(2-fluoro-3-methoxyphenyl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
6		7-[(3-cyclopropyl-2-fluorophenyl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
7		2-[7-[(3-chloro-2-fluorophenyl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide	Method 2
8		7-[(3-chloro-2-fluorophenyl)methyl]-N-ethyl-6-fluoro-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
9		7-[(4,5-difluoro-2-methoxyphenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
10		2-fluoro-3-[[[2-methyl-3-(2-methylcyclopropyl)-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzonitrile	Method 20
11		2-[7-[(3-cyclopropyl-2-fluorophenyl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide	Method 2






-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
13		N-ethyl-6-fluoro-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
14		7-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
15		7-[[3-(cyano-2-fluoro-phenyl)methyl]-6-fluoro-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
16		7-[(2-chloro-4,5-difluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
17		7-[[4,5-difluoro-2-(2-fluoroethyl)phenyl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
18		2-[7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide	Method 2
19		2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide	Method 2
20		7-[(5-chloro-3-methyl-pyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
21		7-[(3-chloro-5-methyl-pyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
22		7-[(3-chloro-5-cyclopropyl-pyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
23		7-[(5-chloro-3-cyclopropyl-pyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
25		2-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide	Method 2
26		N-ethyl-2-methyl-7-[[5-methyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
27		N-ethyl-2-methyl-7-[[3-methyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
29		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
30		7-[[3-(difluoromethyl)-2-fluoro-phenyl]methyl]-6-fluoro-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
31		7-[[3-(difluoromethyl)-2-fluorophenyl]methyl]-N-ethyl-6-fluoro-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
32		7-(4-bicyclo[4.2.0]octa-1,3,5-trienylmethyl)-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
33		N-ethyl-7-[[2-fluoro-3-(1-hydroxycyclopropyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
34		6-fluoro-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-7-[3-(trifluoromethyl)phenoxy]thiazolo[3,2-a]pyrimidin-5-one	Ex. 12
35		N-ethyl-7-[[2-fluoro-3-(1-fluorocyclopropyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
36		N-ethyl-7-[[2-fluoro-3-[1-(fluoromethyl)viny]phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
37		7-[(2-ethynyl-4,5-difluorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
38		2-fluoro-3-[[2-methyl-5-oxo-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzonitrile	Method 20
40		N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
41		7-[(3-cyano-2-fluoro-phenyl)methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
42		7-[[3-(difluoromethyl)-2-fluoro-phenyl]methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
43		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
44		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28







-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
45		N-ethyl-7-[[5-methyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
46		N-ethyl-7-[[3-methyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[1,2-a]pyrimidine-3-carboxamide	Ex. 28
48		N-ethyl-7-[2-fluoro-3-(trifluoromethyl)phenoxy]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 12
49		7-[(3-cyano-2-fluoro-5-methylphenyl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
50		7-[(3-chloro-2-fluoro-phenyl)methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
51		7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 25






-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
52		2-methyl-7-[[5-methyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 25
53		7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
54		7-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
55		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
56		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
57		7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 25
58		7-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 25
59		7-[2-fluoro-3-(trifluoromethyl)phenoxy]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 12
60		2-chloro-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 258
61		N-methyl-7-[[5-methyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
62		N-methyl-7-[[3-methyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28







-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
63		7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
64		7-[(3-chloro-2-fluoro-phenyl)methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
65		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 25
66		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 25
67		2-methyl-7-[[3-methyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 25

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
68		7-(3-cyano-2-fluoro-phenoxy)-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 12
69		2-chloro-7-[(3-cyano-2-fluoro-phenyl)methyl]-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 258
70		7-[(3-cyano-2-fluoro-phenyl)methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
71		N-ethyl-2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]acetamide	Method 2
72		N,N-dimethyl-5-oxo-7-[[4-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
73		3-cyclopropyl-7-[2-fluoro-3-(trifluoromethyl)phenoxy]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Ex. 12
74		3-[[2-chloro-5-oxo-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluoro-benzonitrile	Ex. 267

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
75		2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide	Method 2
76		7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
77		2-fluoro-3-[2-methyl-5-oxo-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-7-yl]oxy-benzonitrile	Ex. 12
78		2-[7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide	Method 2
79		2-[7-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide	Method 2
80		7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(methylsulfonylmethyl)thiazolo[3,2-a]pyrimidin-5-one	Method 10

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
81		7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-(1H-imidazol-2-ylmethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 10
82		N-ethyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
83		N-ethyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
84		2-fluoro-3-[[3-(2-methylcyclopropyl)-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzonitrile	Ex. 207
85		3-[[2-chloro-3-(2-methylcyclopropyl)-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluoro-benzonitrile	Ex. 267
86		7-[2-fluoro-3-(trifluoromethyl)phenoxy]-2-methyl-3-(2-methylcyclopropyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 12
87		N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-isopropyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
88		2-fluoro-3-[[5-oxo-2-(trifluoromethyl)-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzonitrile	Ex. 207
89		6-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one	Method 27
90		6-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one	Method 27
92		6-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one	Method 27
93		2-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-N-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
94		2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Ex. 207

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
95		6-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one	Method 27
96		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
97		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
98		2-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide	Method 2
99		2-[[7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide	Method 2
100		7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-(2-(2-hydroxycyclopropyl)-2-methylthiazolo[3,2-a]pyrimidin-5-one	Method 20







-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
101		2-[7-[2-fluoro-3-(trifluoromethyl)phenoxy]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Ex. 12
102		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
103		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
104		2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Method 15
106		7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-N-isopropyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
107		2-fluoro-3-[2-methyl-3-(2-methylcyclopropyl)-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]oxy-benzonitrile	Ex. 12







-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
108		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1-hydroxy-1-methyl-ethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 18
109		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1-hydroxy-1-methyl-ethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 18
110		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1-hydroxyethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 18
111		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1-hydroxyethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 18
112		7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-N-sec-butyl-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
113		3-[[3-(azetidin-1-yl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluoro-benzonitrile	Method 6







-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
114		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carbonitrile	Method 17
115		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carbonitrile	Method 17
116		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-cyclopropyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
117		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-cyclopropyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
118		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(2-methylcyclopropyl)thiazolo[3,2-a]pyrimidin-5-one	Method 15
119		3-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 18

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
120		3-acetyl-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 18
121		2-chloro-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-(2-methylcyclopropyl)thiazolo[3,2-a]pyrimidin-5-one	Method 33
122		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-pyrimidin-5-yl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
123		7-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
124		7-[[5-chloro-2-pyridyl]-methylamino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
125		2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 1)	Method 15

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
126		2-[[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 2)	Method 15
127		7-[[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methoxy-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 91
128		7-[[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methoxy-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 91
129		7-[[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methoxy-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 91
130		7-[[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methoxy-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 91
131		2-[[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Method 20

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
132		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(1H-pyrazol-5-yl)thiazolo[3,2-a]pyrimidin-5-one	Method 15
133		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(1H-pyrazol-5-yl)thiazolo[3,2-a]pyrimidin-5-one	Method 15
134		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one	Method 15
135		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-isopropenyl-2-methylthiazolo[3,2-a]pyrimidin-5-one	Method 15
136		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methylthiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 1)	Method 15
137		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methylthiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 2)	Method 15







-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
138		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(5-fluoro-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
139		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(5-fluoro-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
140		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-propanoyl-thiazolo[3,2-a]pyrimidin-5-one	Method 18
141		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-propanoyl-thiazolo[3,2-a]pyrimidin-5-one	Method 18
142		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-thiazol-2-yl-thiazolo[3,2-a]pyrimidin-5-one	Method 15

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
143		N-ethyl-7-[[[(5-fluoro-2-pyridyl)-methyl-amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
144		2-[7-[[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Ex. 207
145		2-[7-[[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Ex. 207
146		2-fluoro-3-[[[5-oxo-2-(trifluoromethyl)-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzonitrile	Ex. 207
147		N-ethyl-7-[[[ethyl-(5-fluoro-2-pyridyl)amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
148		N-ethyl-7-[[[ethyl(2-pyridyl)amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28







-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
149		3-(5-chloro-3-pyridyl)-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
150		3-(5-chloro-3-pyridyl)-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
151		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-thiazol-4-yl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
152		7-[[5-chloro-2-pyridyl]-ethyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
154		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-[2-methylcyclopropyl]thiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 1)	Method 15
155		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-[2-methylcyclopropyl]thiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 2)	Method 15

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
156		2-ethoxy-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 91
157		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-fluoro-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
158		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-oxazol-2-yl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
159		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-oxazol-2-yl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
160		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Method 15
161		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Method 15

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
162		2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Ex. 207
163		7-[[[(5-chloro-2-pyridyl)-methyl-amino]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methylthiazolo[3,2-a]pyrimidin-5-one	Method 14
164		7-[[3,5-dichloropyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
165		N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(2,2,2-trifluoroethoxy)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 91
166		3-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 207
167		3-acetyl-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 207

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
168		3-[[4-chloropyrazol-1-yl)methyl]-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 10
169		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl)methyl]-3-isopropenyl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 207
170		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl)methyl]-3-(1H-imidazol-2-yl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Ex. 258
171		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl)methyl]-3-pyrimidin-5-yl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 207
172		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl)methyl]-3-pyrimidin-5-yl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 207

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
173		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-pyridyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 207
174		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-2-(2,2,2-trifluoroethoxy)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 91
175		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethoxy-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 91
176		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethoxy-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 91
177		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(2-methylpropanoyl)thiazolo[3,2-a]pyrimidin-5-one	Method 18

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
178		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(2-methylpropanoyl)thiazolo[3,2-a]pyrimidin-5-one	Method 18
179		7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-N-[(1R)-1-methylpropyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
180		7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-N-[(1S)-1-methylpropyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 21
191		7-[[[(4-chloro-2-pyridyl)-ethylamino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
182		7-[[[(5-fluoro-2-pyridyl)-methylamino]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
183		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(methoxymethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 10

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
184		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-cyclopropyl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 91
185		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(methylsulfonylmethyl)thiazolo[3,2-a]pyrimidin-5-one	Method 10
186		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(pyrazol-1-ylmethyl)thiazolo[3,2-a]pyrimidin-5-one	Method 10
187		2-[7-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Method 20
188		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-cyclobutyl-2-methylthiazolo[3,2-a]pyrimidin-5-one	Ex. 28
189		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-cyclobutyl-2-methylthiazolo[3,2-a]pyrimidin-5-one	Ex. 28

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
190		2-[7-[(5-chloro-2-pyridyl)-methyl-amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Method 14
191		7-[[ethyl-(5-fluoro-2-pyridyl)amino]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 14
192		3-chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
193		3-chloro-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
194		2-[7-[(4,5-difluoro-2-methoxyphenyl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Method 23
195		2-[7-[[ethyl-(5-fluoro-2-pyridyl)amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Method 14

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
196		7-[[[(5-chloro-2-pyridyl)-ethyl-amino)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 14
197		5-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]pyridine-3-carbonitrile	Method 15
198		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(2,2,2-trifluoroacetyl)thiazolo[3,2-a]pyrimidin-5-one	Method 10
199		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-methylcyclopropyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 207
200		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-methylcyclopropyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Method 15
201		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-pyrimidin-5-yl-thiazolo[3,2-a]pyrimidin-5-one	Method 15

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
202		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one	Method 23
203		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(2,2,2-trifluoro-1-hydroxy-ethyl)thiazolo[3,2-a]pyrimidin-5-one	Method 14
204		7-[[5-(2-bromo-2-pyridyl)-ethyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 14
205		N-ethyl-7-[[ethyl-(5-fluoro-2-pyridyl)amino]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Method 15
209		2-[7-[[5-methoxy-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Method 15
210		2-[7-[[3-methoxy-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Method 15







-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
211		3-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-thiazolo[3,2-a]pyrimidin-5-one	Method 18
212		3-acetyl-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-thiazolo[3,2-a]pyrimidin-5-one	Method 18
213		2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Method 15
214		2-[7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Method 15
215		3-bromo-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
216		3-bromo-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15







-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
217		3-chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methoxy-thiazolo[3,2-a]pyrimidin-5-one	Ex. 244
218		3-chloro-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methoxy-thiazolo[3,2-a]pyrimidin-5-one	Ex. 244
219		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-methylsulfanyl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 206
221		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(cyclopropylmethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
223		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-hydroxy-1-methyl-ethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Ex. 222
224		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(cyclopropanecarbonyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 10







-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
225		3-bromo-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one	Method 15
226		3-bromo-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one	Method 15
227		7-[(3-amino-5-chloro-pyrazol-1-yl)methyl]-3-bromo-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
228		7-[(5-amino-3-chloro-pyrazol-1-yl)methyl]-3-bromo-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
229		2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]acetonitrile	Method 10
230		N-ethyl-7-[[[(5-fluoro-2-pyridyl)-methyl-amino]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28







-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
231		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3,3-difluoroazetidin-1-yl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 206
233		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methoxy-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 244
234		2-[7-[[5-bromo-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Method 15
235		3-chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one	Method 15
236		3-chloro-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one	Method 15
237		2-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-carboxamide	Ex. 258

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
238		2-acetyl-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 258
239		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Method 15
241		2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Method 15
242		3-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one	Method 18
243		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2,2,2-trifluoroethoxy)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 206
245		3-acetyl-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methoxy-thiazolo[3,2-a]pyrimidin-5-one	Ex. 244







-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
246		3-bromo-7-[(5-chloro-3-nitro-pyrazol-1-yl)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one	Method 15
247		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1H-pyrazol-5-yl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 207
248		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-thiazol-4-yl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 207
249		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(5-fluoro-3-pyridyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 207
251A		2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]propanenitrile	Ex. 222
252		2-[7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]propanenitrile	Ex. 222

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
253		2-fluoro-3-[[3-[2-methylcyclopropyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzonitrile	Ex. 207
254		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-fluoroazetidin-1-yl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 206
255		3-(5-chloro-3-pyridyl)-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one	Ex. 207
256		7-[(3,5-dichloropyrazol-1-yl)methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28
257		3-[[3-acetyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluorobenzonitrile	Ex. 207
259		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(difluoromethyl)-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 258

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
260		(Z)-3-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enitrile	Method 5
261		(E)-3-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enamide	Method 5
262		(E)-3-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enitrile	Method 5
263		(Z)-3-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enitrile	Method 5
264		(E)-3-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enamide	Method 5
265		N-ethyl-7-[[5-isobutyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide	Ex. 28

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
268		2-[2-chloro-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Ex. 267
269		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 1)	Ex. 207
270		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 2)	Ex. 207
271		2-[7-((4-chloro-1-methyl-pyrazol-3-yl)methyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Method 20
272		2-[2-methyl-5-oxo-7-[[3-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 1)	Method 15
273		2-[2-methyl-5-oxo-7-[[3-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 2)	Method 15

-continued

Ex.	Structure/Name	Chemical Name	Ref. Method or Ex.
274		2-[7-[[5-bromo-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 1)	Method 15
275		2-[7-[[5-bromo-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 2)	Method 15
276		2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-6-fluoro-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 1)	Ex. 207
277		2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-6-fluoro-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 2)	Ex. 207
278		2-[2-methyl-7-[[1-methyl-4-(trifluoromethyl)imidazol-2-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile	Ex. 24
279		(E)-3-[7-[(N-ethyl-4-fluoroanilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enenitrile	Method 5

The following compounds were prepared using methods analogous to those described herein. Satisfactory analytical data was obtained for each compound.

Ex.	Structure	Name
280		7-[(4-fluorophenoxy)methyl]-3-[[2-hydroxyethyl(methyl)amino]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
281		7-[(4-fluorophenoxy)methyl]-3-[(2-hydroxyethylamino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
282		2-[7-[(4-fluorophenoxy)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N,N-dimethyl-acetamide
283		7-[(2-cyano-4,5-difluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
284		7-[(2-cyclopropyl-4,5-difluoro-phenyl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide

-continued

Ex.	Structure	Name
285		3-[2-(azetidin-1-yl)-2-oxo-ethyl]-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
286		7-[(4-fluorophenoxy)methyl]-2-methyl-3-(4H-1,2,4-triazol-3-ylmethyl)thiazolo[3,2-a]pyrimidin-5-one
287		2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-propanamide
288		3-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-propanamide
289		7-[[5-chloro-2-(trifluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
290		7-[(5-ethyl-1,3-benzoxazol-6-yl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide

-continued

Ex.	Structure	Name
291		7-[(3-chloropyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
292		7-[(5-chloropyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
293		2-[7-[(3-cyano-2-fluoro-phenyl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide
294		N-ethyl-7-[[2-fluoro-3-(1-hydroxypropyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
295		7-[[4,5-difluoro-2-oxazol-2-yl-phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
296		2-fluoro-3-[[2-methyl-5-oxo-3-propanoyl-thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzonitrile
297		7-[[4,5-difluoro-2-(trifluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide

-continued

Ex.	Structure	Name
298		N,2-dimethyl-5-oxo-7-[3-(trifluoromethyl)phenoxy]thiazolo[3,2-a]pyrimidine-3-carboxamide
299		7-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
300		7-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
301		7-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
302		N-ethyl-2-methyl-5-oxo-7-[3-(trifluoromethyl)phenoxy]thiazolo[3,2-a]pyrimidine-3-carboxamide
303		3-[(2-chloro-3-cyclopropyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl)methyl]-2-fluoro-benzonitrile
304		7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(pyrazol-1-ylmethyl)thiazolo[3,2-a]pyrimidin-5-one







-continued

Ex.	Structure	Name
305		N,2-dimethyl-7-[[3-methyl-4-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
306		N,2-dimethyl-7-[[5-methyl-4-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
307		2-fluoro-3-[(8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-6-yl)methyl]benzonitrile
308		7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-[hydroxy(thiazol-2-yl)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
309		2-fluoro-3-[(3-methyl-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-6-yl)methyl]benzonitrile
310		2-[7-[(4-fluorophenoxy)methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide
311		2-fluoro-3-[(8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-6-yl)oxy]benzonitrile

-continued

Ex.	Structure	Name
312		2-fluoro-3-[[1-(hydroxymethyl)-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-6-yl]methyl]benzonitrile
313		7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-N-(2-hydroxy-1-methyl-ethyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
314		3-[[3-(2,3-dimethylcyclopropyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluorobenzonitrile
315		6-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one
316		7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(2-oxa-6-azaspiro[3.3]heptan-6-yl)thiazolo[3,2-a]pyrimidin-5-one
317		N-ethyl-6-fluoro-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide

-continued

Ex.	Structure	Name
318		7-[(4-fluorophenoxy)methyl]-5-oxo-N-(2,2,2-trifluoroethyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
319		N-cyclopentyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
320		7-[(4,5-dichloropyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
321		7-[(3,4-dichloropyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
322		N-ethyl-2-methyl-7-[[methyl(thiazol-2-yl)amino]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
323		7-[(4-chloropyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide







-continued

Ex.	Structure	Name
324		N-ethyl-2-methyl-7-[[methyl-(1-methylpyrazol-4-yl)amino]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
325		7-[(4-fluorophenoxy)methyl]-2-(trifluoromethyl)-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-5-one
326		7-[(3-ethoxy-2-pyridyl)-methyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
327		7-[(3,5-dimethylisoxazol-4-yl)-methyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
328		3-cyclopropyl-7-[(4-fluorophenoxy)methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one

-continued

Ex.	Structure	Name
329		7-[(4-fluorophenoxy)methyl]-3-pyrimidin-5-yl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
330		7-[(4-fluorophenoxy)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
331		7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(4-methyl-1,2,4-triazol-3-yl)thiazolo[3,2-a]pyrimidin-5-one
332		2-fluoro-3-[[5-oxo-2-(trifluoromethyl)-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzonitrile
333		N-ethyl-7-[[ethyl(4-pyridyl)amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
334		N-ethyl-7-[[ethyl(3-pyridyl)amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide

-continued

Ex.	Structure	Name
335		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-methoxy-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
336		2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
337		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-isopropenyl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
338		3-(5-chloro-3-pyridyl)-7-[[5-chloro-2-pyridyl)methyl-amino]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
339		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-pyridyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
340		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-2-(2,2,2-trifluoroethoxy)thiazolo[3,2-a]pyrimidine-3-carboxamide

-continued

Ex.	Structure	Name
341		7-[[3-chloro-6-(trifluoromethyl)-2-pyridyl]methyl]-2-methyl-3-(2-methylcyclopropyl)thiazolo[3,2-a]pyrimidin-5-one
342		7-[(5-chloro-2-pyridyl)oxymethyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
343		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one
344		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-pyrrolidin-1-yl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
345		N-ethyl-7-[[5-(methoxy-2-pyridyl)-methylamino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
346		3-(2-chloro-3-pyridyl)-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one

-continued

Ex.	Structure	Name
347		3-(2-chloro-3-pyridyl)-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one
348		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one
349		7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one
350		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-methoxyazetidin-1-yl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
351		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(cyclopropylmethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one

-continued

Ex.	Structure	Name
352		5-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]pyridine-3-carbonitrile
353		2-fluoro-3-[[3-[2-methylcyclopropyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzonitrile
354		7-[(3,5-diisopropylpyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
355		2-[7-[(4-chloro-2-methyl-pyrazol-3-yl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile
356		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-methylazetidin-1-yl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
357		2-[7-[(3,5-dichloropyrazol-1-yl)methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 1)

-continued

Ex.	Structure	Name
358		2-[7-[(3,5-dichloropyrazol-1-yl)methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile (cis enantiomer 2)
359		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(2-hydroxyethyl)cyclopropyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 1)
360		7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(2-hydroxyethyl)cyclopropyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one (trans enantiomer 2)
361		3-chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one
362		5-chloro-1-[[3-[2-methylcyclopropyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]pyrazole-3-carbonitrile (cis enantiomer 1)
363		5-chloro-1-[[3-[2-methylcyclopropyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]pyrazole-3-carbonitrile (cis enantiomer 2)

-continued

Ex.	Structure	Name
364		5-chloro-2-[[3-[2-methylcyclopropyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]pyrazole-3-carbonitrile
365		7-[[5-ethoxy-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide
366		N-ethyl-7-[[5-isobutyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide
367		2-[2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropyl]acetonitrile (trans enantiomer 1)
368		2-[2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropyl]acetonitrile (trans enantiomer 2)

It is understood that the person skilled in the art will be able to prepare the compounds of the present invention using methods known in the art along with the general method of synthesis described herein.

Assay 1: Cell-Based Assay

HEK cells stably transfected with tetracycline inducible hNR1 and hNR2A were seeded into clear bottom 384 well poly-D-lysine coated plates (2.5×10⁴ cells per well) in Minimum Essential Media (MEM; without L-) including 7.5 μg mL⁻¹ doxycycline and 500 μM (+)-ketamine. The cells were incubated at 37° C. in 5% CO₂ for 24 h. For measurement of changes in cytosolic calcium, the seeding media was removed and the cells incubated at 37° C. for 60 min with 1× Becton Dickinson Calcium Assay Kit reagent in Hanks Balanced Salt Solution (HBSS; w/o magnesium, including 1.8 mM calcium, 0.65 mg ml⁻¹ probenecid and 10 μM (+)-ketamine, pH 7.15) then allowed to equilibrate to rt for 30 min. Concentration-effect curves to Positive Allosteric Modulators (PAMs) were constructed by adding different concentrations (with 30 μM glycine and 300 nM L-glutamate (EC₃₀)) to different wells in HBSS. Compounds were added after a 10 second baseline read and maximum level of relative fluorescence units (RFU) was measured over a 5 min period. Responses were scaled relative to 100 μM L-glutamate maximal response (100%) and 0 μM L-glutamate (0%). EC₅₀ values are provided for compounds reaching maximal response plateaus, and the max % (EC₅₀ (- -)) only if no plateau was reached.

A four-parameter Hill equation was fitted to individual concentration-effect curves:

$$Y = S_0 + \frac{S_{inf} - S_0}{1 + \left(\frac{10^{\log AC_{50}}}{10^c}\right)^n}$$

in which Y, S₀, S_{inf}, AC₅₀, n and c were effect, lower-asymptote, upper-asymptote, mid-point location, slope parameter, and concentration respectively.

Data for compounds tested in this assay are shown below in Table 2.

TABLE 2

No.	EC ₅₀ (uM)	Max %
1.1	9.6	63%
1.2	—	92%
1.3	—	45%
1.4	11.2	59.5%
1.5	23	51.4%
1.6	31.6	64%
1.7	27	44.8%
1.8	26.3	47.4%
1.9	—	99.5%
1.10	—	44%
1.11	—	73%
1.12	—	45.5%
1.13	—	41.6%
1.14	—	42.7%
1.15	—	55.1%
1.16	—	56.5%
1.17	4	68.6%
2.1	5.2	51.3%
2.2	—	96.5%
2.3	12.6	49.8%
2.4	3.0	44.6%
2.5	32.9	115%
2.6	19.6	59.9%
2.7	26	61.3%

TABLE 2-continued

No.	EC ₅₀ (uM)	Max %
2.8	—	137%
2.9	—	56.4%
2.10	—	119%
2.11	—	87.4%
3.1	41	56.2%
3.2	10.1	42.7%
3.3	—	61.8%
3.4	—	44.6%
3.5	—	65.6%
3.6	—	68%
4.1	5.1	141%
4.2	1.3	137%
4.3	20.5	71.6%
4.4	2.5	138%
4.5	5.7	42.2%
4.6	7.7	128%
4.7	5.8	134%
4.8	4.4	99.3%
4.9	4.4	123%
4.10	4.3	108%
4.11	4.8	40.8%
4.12	30.6	127%
4.13	31.1	72.8%
4.14	7.8	93%
4.15	13.9	50%
4.16	—	46.2%
4.17	—	40.3%
4.18	5.0	102%
4.19	9.0	46.9%
4.20	2.4	93.9%
4.21	16.3	47.4

TABLE 2-continued

No.	EC ₅₀ (uM)	Max %
4.22	—	84.2
4.23	26.5	62.9%
4.24	2.1	80%
4.25	2.8	109%
4.26	—	98.8
4.27	—	103%
4.28	3.4	89%
4.29	17.5	134%
4.30	15.5	75.4%
4.31	—	43.3%
4.32	—	50.2%
4.33	13.8	121%
4.34	—	49.2%
5.1	0.560	145.1
5.2	5.7	148%
5.3	2.2	177%
5.4	0.956	172%
5.5	2.4	168%
5.6	0.766	164%
5.7	0.867	160%
5.8	0.669	156%
5.9	1.3	143%
5.10	18	133%
5.11	2.9	118%
5.12	—	106%
5.13	—	87.9%
5.14	7.5	111%
5.15	7	144%
5.16	2.5	157%
5.17	0.108	176%
5.18	0.091	164%

US 10,647,731 B2

481

TABLE 2-continued

No.	EC ₅₀ (uM)	Max %
5.19	1.7	141%
5.20	0.815	157%
5.21	0.584	141%
5.22	1.6	142%
5.23	2.6	135%
5.24	0.731	142%
5.25	0.7	141
5.26	0.563	149%
5.27	1.1	139%
5.28	.0952	176%
5.29	0.445	147%
5.30	4.3	132%
5.31	2.7	132%
5.32	4.3	123%
5.33	4.2	120
5.34	2.8	118%
5.35	2.7	136%
5.36	—	61.3%
5.37	0.515	152%
5.38	1.1	147%
5.39	0.952	167%
5.40	1.9	149%
5.41	2	125%
5.42	0.0329	162%
5.43	0.821	145%
5.44	1.1	158%
5.45	0.214	139%
5.46	3.8	155%
5.47	0.976	151%
5.48	6.2	136%
5.49	12.9	127%
5.50	—	96.8%
5.51	40.7	80.3%
5.52	1.6	58%
5.53	1.3	153%
5.54	51.5	129%
5.55	4.7	111%
5.56	—	110%
5.57	6.4	120%
5.58	5.1	120%
6.1	10	123%
6.2	5	121%
6.3	10.1	91.9%
7.1	0.853	184%
7.2	0.612	134%
7.3	0.376	156%
7.4	5.7	166%
7.5	2.1	162%
7.6	3.7	151%
7.7	21	148%
7.8	2.5	126%
7.9	1.3	121%
8.1	5.2	146%
8.2	7.2	122%
8.3	1.3	150%
8.4	0.569	164%
8.5	16.9	114%
8.6	—	60.5%
8.7	—	81.4%
8.8	8.7	108%
8.9	11.1	124%
8.10	12.5	127%
8.11	2.7	154%
8.12	0.385	161%
8.13	2.2	162%
8.14	1.1	154%
8.15	2.3	138%
8.16	13	137%
9.1	—	48.1%
9.2	2.9	117%
9.3	7.0	108%
9.4	9.2	114%
10.1	4.4	130%
10.2	5.3	110%
10.3	0.535	177%
10.4	0.593	186%
10.5	0.495	158%
10.6	0.548	176%

482

TABLE 2-continued

No.	EC ₅₀ (uM)	Max %
10.7	23.4	127%
10.8	—	63.5%
10.9	—	72.1%
10.10	18.8	132%
10.11	1.2	167%
10.12	2.1	149%
10.13	0.345	165%
10.14	1.5	172%
10.15	1.3	156%
10.16	2.7	147%
10.17	1.2	174%
10.18	16.9	88.2%
10.19	0.837	165%
10.20	2.2	178%
10.21	—	117%
10.22	7.5	156%
10.23	22.9	169%
10.24	8.9	160%
10.25	0.345	165%
11.1	2	159%
11.2	—	107%
11.3	3.2	132%
11.4	—	103%
11.5	4.6	167%
11.6	5.0	133%
12.1	0.723	166%
12.2	0.27	192%
12.3	1.6	183%
12.4	0.591	176%
12.5	0.974	153%
13.1	0.434	161%
13.2	0.913	130%
13.3	0.783	137%
13.4	0.777	185%
13.5	1	152%
13.6	1.4	125%
13.7	0.547	137%
13.8	—	61.2%
13.9	5.0	166%
14.1	2.6	163%
14.2	0.122	205%
15.1	2.2	144%
15.2	1	167%
15.3	0.489	144%
15.4	0.782	137%
15.5	1.7	131%
15.6	9.2	138%
15.7	0.826	158%
15.8	30.8	122%
15.9	18.6	117%
15.10	3.1	133%
15.11	0.151	159%
15.12	2.5	143%
15.13	—	67.6%
15.14	3.1	159%
15.15	—	61.9%
15.16	17.7	115%
15.17	3.4	137%
15.18	10.5	143%
15.19	0.512	126%
15.20	0.314	153%
15.21	1.6	161%
15.22	0.704	149%
15.23	1.9	143%
15.24	1.4	138%
15.25	16.5	137%
15.26	3.8	123%
15.27	12.3	137%
15.28	—	80.8%
15.29	7	135%
15.30	22	116%
15.31	—	57.8%
15.32	—	94.8%
15.33	18.8	51.8%
15.34	—	48.2%
16.1	2.3	140%
16.2	11.7	110%
16.3	1.4	160%

483

TABLE 2-continued

No.	EC ₅₀ (uM)	Max %
16.4	9.2	130%
16.5	10.6	149%
16.6	5.5	140%
16.7	36.2	67.5%
17.1	—	107%
17.2	2	119%
17.3	—	104%
18.1	—	94.8%
18.2	2.2	144%
18.3	25.4	129%
18.4	—	127%
18.5	—	40.7%
19.1	12.2	116%
19.2	—	64.2%
20.1	3.1	137%
20.2	7.4	145%
20.3	1.8	148%
20.4	8.5	147%
20.5	24	124%
20.6	7.2	137%
20.7	10.8	137%
20.8	19	134%
20.9	13.9	133%
20.10	9.1	127%
20.11	31.2	78.7%
20.12	—	74%
20.13	5.8	146%
21.1	—	96.6%
21.2	21.6	133%
21.3	7.6	131%
21.4	7.4	129%
21.5	18.9	124%
21.6	45.3	114%
21.7	—	100%
21.8	—	99.3%
21.9	—	90.1%
21.10	—	70.3%
21.11	—	47%
21.12	6.7	112%
21.13	12.3	131%
21.14	11	112%
21.15	—	112%
21.16	14.3	107%
21.17	23.7	107%
21.18	—	58.5
21.19	7.4	145%
21.20	5.2	114%
21.21	33.6	134%
21.22	—	60.1%
21.23	6.5	107%
21.24	34.5	102%
21.25	3.9	105%
21.26	—	76.5%
21.27	16	122%
21.28	11.4	126%
21.29	4.7	98.2%
21.30	—	100%
21.31	—	88.8%
21.32	—	84.3%
21.33	—	82.5%
21.34	—	79.9%
21.35	—	77.8%
21.36	—	71.8%
21.37	—	68.4%
21.38	—	66.5%
21.39	—	62.6%
21.40	—	61.6%
21.41	—	59.7%
21.42	—	54.5%
21.43	—	53.5%
21.44	—	51.9%
21.45	—	49.1%
21.46	—	43.2%
21.47	—	103%
21.48	8.2	100%
21.49	—	82.3%
21.50	—	76.1
21.51	—	75.0

484

TABLE 2-continued

No.	EC ₅₀ (uM)	Max %
21.52	—	72.9%
21.53	—	49%
21.54	—	46%
21.55	—	43.8
21.56	—	41%
21.57	—	64.1%
21.58	—	46.1
21.59	—	42.5%
21.60	—	78%
21.61	33.1	131%
21.62	—	45.6%
21.63	—	124%
21.64	—	120%
21.65	—	96.3%
21.66	—	92.4%
21.67	—	75%
21.68	4.2	111%
22.1	—	96.9%
22.2	—	41.4%
23.1	18.1	124%
23.2	2.6	134%
23.3	14.8	133%
23.4	6.5	124%
24.1	—	78.5%
24.2	—	116%
24.3	—	87.7%
24.4	15.3	103%
24.5	—	59.3%
24.6	—	50.5
24.7	—	47.4%
24.8	8.9	94.8%
24.9	—	48.5%
25.1	18.7	86.2%
25.2	15.2	140%
26.1	—	80.6%
26.2	9	71%
27.1	—	45.6%
27.2	—	83.2%
27.3	—	93.3%
27.4	—	58.4%
27.5	30.4	53.2%
27.6	—	91.6%
27.7	26.5	68.5%
27.8	30.4	53.2%
27.9	—	47.8%
27.10	29.1	131%
27.11	25.3	93.8%
27.12	40.8	41.4%
27.13	—	48.8%
27.14	44.5	93%
27.15	—	65.3%

Additional data for compounds tested in this assay are shown in Table 3.

Ex.	EC ₅₀ (uM)	Max %
1	0.932	84.0
2	33.7	50.6
3	10.9	55.9
4	3.88	47.5
5	15.1	105
6	0.753	66.8
7	9.73	77.9
8	2.55	86.5
9	8.04	86.4
10	0.471	138
11	1.14	78.9
12	2.05	113
13	0.858	89.9
14	1.56	56.2
15	9.02	91.5
16	15.5	88.0
17	7.04	176
18	2.25	192

US 10,647,731 B2

485

-continued

486

-continued

Ex.	EC ₅₀ (uM)	Max %		Ex.	EC ₅₀ (uM)	Max %
19	5.45	99.5		96	1.21	119
20	10.2	136	5	97	5.11	154
21	7.88	112		98	1.57	130
22	4.13	172		99	3.69	148
23	3.48	90.0		100	11.6	94.1
24	1.33	129		101	3.71	107
25	0.292	204		102	0.0547	129
26	4.82	130	10	103	0.251	151
27	10.6	126		104	0.0341	137
28	0.388	125		105	3.9	96.1
29	1.76	165		106	3.4	69.7
30	2.57	106		107	0.994	121
31	4.17	111		108	0.247	111
32	5.95	110	15	109	0.734	121
33	6.43	75.8		110	1.88	126
34	24.5	78.4		111	6.0	143
35	3.99	70.7		112	0.825	55.7
36	1.72	80.5		113	2.37	69.3
37	9.02	69.4		114	1.34	116
38	0.533	129		115	2.75	128
39	6.1	101	20	116	0.351	101
40	1.32	61.4		117	1.03	107
41	4.78	101		118	0.169	128
42	1.57	105		119	0.325	131
43	0.362	131		120	0.995	151
44	0.685	114		121	17.6	38.0
45	1.25	130	25	122	9.14	109
46	7.04	123		123	2.59	110
47	5.14	89.3		124	5.9	108
48	7.66	92.3		125	0.0278	154
49	17.5	85.2		126	0.249	125
50	6.4	95.2		127	4.87	155
51	3.16	62.5	30	128	0.682	108
52	8.76	129		129	0.774	149
53	0.635	162		130	6.86	102
54	2.2	96.2		131	0.681	113
55	1.16	136		132	2.04	119
56	2.27	139		133	10.1	135
57	0.627	174	35	134	0.501	76.6
58	6.53	126		135	0.397	115
59	8.22	96.7		136	0.0391	142
60	3.22	106		137	0.42	129
61	8.06	127		138	0.775	104
62	24.6	108		139	6.36	124
63	3.71	94.5		140	0.147	129
64	12.1	114	40	141	0.328	131
65	0.551	128		142	2.09	126
66	1.69	149		143	12.0	137
67	36.6	119		144	0.0411	151
68	30.4	105		145	0.132	175
69	11.2	114		146	0.156	140
70	2.92	68.7	45	147	1.98	179
71	1.37	61.6		148	20.8	159
72	62.6	41.0		149	0.729	107
73	40.1	53.9		150	17.3	114
74	0.507	148		151	0.538	115
75	28.9	71.9		152	2.85	140
76	0.436	145	50	153	73.5	100
77	0.271	129		154	0.504	126
78	2.96	174		155	0.0616	136
79	14.3	56.5		156	1.47	74.2
80	12.2	43.1		157	13.0	99.8
81	52.0	53.8		158	1.42	130
82	0.0787	192	55	159	7.59	149
83	0.154	185		160	0.753	133
84	11.2	109		161	1.49	141
85	2.39	128		162	0.199	130
86	10.5	114		163	0.359	143
87	42.0	92.4		164	0.562	131
88	3.7	118	60	165	1.37	46.3
89	9.04	79.2		166	2.69	126
90	1.22	45.7		167	5.26	132
91	1.03	89.4		168	34.1	73.3
92	47.1	94.5		169	7.46	108
93	1.83	120		170	0.814	135
94	2.08	125	65	171	1.21	67.5
95	9.2	135		172	27.6	83.3

US 10,647,731 B2

487

-continued

488

-continued

Ex.	EC ₅₀ (uM)	Max %		Ex.	EC ₅₀ (uM)	Max %
173	24.7	78.6		251	0.42	133
174	1.67	90.5	5	251A	2.62	106
175	0.864	90.6		252	5.76	140
176	8.92	131		253	9.95	111
177	0.123	133		254	1.37	104
178	0.354	163		255	7.86	66.2
179	0.698	54.1		256	0.611	145
180	2.94	59.1	10	257	20.5	93.5
191	7.34	120		258	0.227	133
182	0.175	169		259	0.749	159
183	5.29	129		260	2.45	138
184	9.99	118		261	1.54	126
185	17.8	97.9		262	1.11	112
186	3.85	94.1	15	263	0.501	117
187	11.1	104		264	5.09	82
188	3.14	129		265	4.84	101
189	7.45	121		266	0.0249	158
190	0.779	136		267	0.157	141
191	0.172	187		268	0.182	166
192	0.198	96.3	20	269	0.0831	144
193	0.243	80.9		270	0.518	120
194	1.28	117		271	3.57	136
195	0.0742	177		272	0.252	143
196	0.0642	171		273	2.08	124
197	0.327	89.8		274	0.012	
198	0.201	108	25	275	0.0737	
199	1.39	119		276	0.0364	156
200	3.13	124		277	0.155	140
201	1.4	88.4		278	10.6	129
202	23.9	100		279	1.82	141
203	5.85	135		280	—	28.3
204	5.12	137		281	—	30.9
205	1.12	163	30	282	—	41.2
206	7.19	121		283	—	56.2
207	0.0074	150		284	—	149
208	0.37	120		285	—	38.0
209	0.0666	157		286	—	59.0
210	8.92	161		287	—	123
211	0.31	123	35	288	—	121
212	5.84	133		289	—	129
213	0.066	152		290	—	50.5
214	0.18	154		291	—	86.6
215	0.0911	95.6		292	—	46.3
216	0.0594	64.9		293	—	96.3
217	4.74	125	40	294	—	82.8
218	83.0	112		295	—	49.4
219	10.5	118		296	—	111
220	47.0	95.6		297	—	93.1
221	3.76	96.5		298	—	82.3
222	2.92	115		299	—	53.1
223	10.9	128	45	300	—	72.1
224	0.366	115		301	—	35.8
225	8.69	99.6		302	—	85.4
226	1.42	48.4		303	—	116
227	13.0	59.3		304	—	34.8
228	32.2	66.5		305	—	32.9
229	6.46	107		306	—	32.2
230	6.86	130	50	307	—	41.7
231	61.3	116		308	—	35.3
233	6.14	98.6		309	—	47.5
234	0.0342	163		310	—	32.5
235	21.2	127		311	—	28.9
236	44.5	103		312	—	67.0
237	0.889	98.7	55	313	—	88.2
238	2.32	123		314	—	98.4
239	14.2	131		315	—	54.4
240	37.1	120		316	—	49.0
241	0.313	134		317	—	51.1
242	11.4	134		318	—	27.0
243	0.864	122	60	319	—	28.2
244	0.392	134		320	—	26.9
245	0.409	139		321	—	26.6
246	0.0999	148		322	—	36.1
247	7.7	108		323	—	53.7
248	1.45	124		324	—	62.2
249	6.33	99.2	65	325	—	30.9
250	13.8	111		326	—	36.2

489


-continued

Ex.	EC ₅₀ (uM)	Max %
327	—	35.5
328	—	39.3
329	—	29.3
330	—	38.3
331	—	47.5
332	—	75.4
333	—	42.3
334	—	48.0
335	—	53.8
336	—	81.5
337	—	103
338	—	65.5
339	—	71.5
340	—	131
341	—	95.2
342	—	66.1
343	—	103
344	—	38.4
345	—	32.3
346	—	51.1
347	—	41.1
348	—	82.1
349	—	35.4
350	—	87.9
351	3.76	96.5
352	—	83.7
353	—	34.9
354	—	NT
355	—	46.9

While the present invention has been described in conjunction with the specific embodiments set forth above, many alternatives, modifications and other variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present invention.

The invention claimed is:

1. A method of treating a subject suffering from or diagnosed with depression or anxiety mediated by NR2A activity, comprising administering to the subject in need of such treatment an effective amount a compound of Formula (II):



wherein

R^a is C₁₋₆alkyl or C₂₋₆alkenyl, each optionally substituted with one or more R^b substituents; C₂₋₆alkynyl; halo; —C(O)R^c; —NR^dR^e; —C(O)NR^dR^e; —C(S)NR^dR^e; —C(=N—OH)—C₁₋₄alkyl; —OC₁₋₄alkyl; —OC₁₋₄haloalkyl; —SC₁₋₄alkyl; SO₂C₁₋₄alkyl; cyano; C₃₋₆cycloalkyl optionally substituted with one or more R^f substituents; or a phenyl, monocyclic heteroaryl, or heterocycloalkyl ring, each ring optionally substituted with one or more R^g substituents;

wherein each R^b substituent is independently selected from the group consisting of —OH, —C₁₋₄alkoxy,

490

—NR^dR^e, —C(O)NR^dR^e, —SC₁₋₄alkyl, —SO₂C₁₋₄alkyl, cyano, halo, C₃₋₆cycloalkyl, and monocyclic heteroaryl;

R^c is C₁₋₄alkyl, —C₁₋₄haloalkyl, C₃₋₆cycloalkyl, or a monocyclic, carbon-linked heterocycloalkyl;

R^d is H or C₁₋₄alkyl;

R^e is H; C₁₋₄alkyl optionally substituted with —CN, —CF₃, —OH, or a monocyclic heterocycloalkyl; C₃₋₆cycloalkyl; —OH; or —OC₁₋₄alkoxy;

or R^d and R^e taken together with the nitrogen to which they are attached form a heterocycloalkyl, optionally substituted with C₁₋₄alkyl or OH;

each R^f substituent is independently selected from the group consisting of: C₁₋₄alkyl optionally substituted with —OH, cyano, or C₁₋₄alkoxy; —OH; halo; C₁₋₄haloalkyl; —CONH₂; and cyano; and

each R^g substituent is independently selected from the group consisting of C₁₋₄alkyl, CF₃, halo, —NH₂, —OCH₃, cyano, and —OH;

R¹ is selected from the group consisting of H, C₁₋₆alkyl, C₁₋₄haloalkyl, C₃₋₆cycloalkyl, halo, —OC₁₋₄alkyl, —OC₁₋₄haloalkyl, cyano, and —C(O)C₁₋₄alkyl; or R^a and R¹ taken together with the carbons to which they are attached form a 5- to 7-membered ring, optionally containing an O or NH, and optionally substituted with one or more R^h substituents;

wherein each R^h substituent is independently —C(O)NRⁱR^j, cyano, or is C₁₋₄alkyl optionally substituted with —OH, OCH₃, cyano, or —C(O)NRⁱR^j; or two R^h groups attached to the same carbon and taken together with the carbon to which they are attached form a carbonyl or a C₃₋₆cycloalkyl;

wherein Rⁱ and R^j are each independently H or C₁₋₄alkyl;

R² is —R^m, —OR^m, or —NR^mRⁿ;

wherein R^m is phenyl, naphthyl, pyridyl, pyrazinyl, pyridazinyl, pyrrolyl, pyrazolyl, triazolyl, imidazolyl, furanyl, oxazolyl, isoxazolyl, thiophenyl, thiazolyl, isothiazolyl, indolyl, indazolyl, quinolyl, or isoquinolyl, each optionally substituted with one or more R^s substituents;

wherein each R^s substituent is independently selected from the group consisting of C₁₋₄alkyl, C₂₋₄alkenyl (optionally substituted with halo), C₂₋₄alkynyl, C₁₋₄haloalkyl, C₁₋₄alkoxy, C₁₋₄alkyl-OH, C₁₋₄haloalkoxy, halo, cyano, C₃₋₆cycloalkyl (optionally substituted with —OH or halo), monocyclic heteroaryl, —NH₂, —NO₂, NHSO₂C₁₋₄alkyl, and —SO₂C₁₋₄alkyl;

Rⁿ is H, C₁₋₄haloalkyl, or C₁₋₄alkyl optionally substituted with —OH or C₁₋₄alkoxy;

or R^m and Rⁿ taken together with the nitrogen to which they are attached form a pyrrolidine or piperidine ring, optionally substituted with C₁₋₄alkyl and optionally fused to phenyl, wherein said phenyl is optionally substituted with halo;


R³ is H or methyl; and

R⁴ is H or fluoro;

or a pharmaceutically acceptable salt thereof.

2. The method of claim 1, wherein the compound of Formula II is a compound of Formula (I):

491



(I)

wherein

R^a is C_{1-6} alkyl optionally substituted with one or more R^b substituents; C_{2-6} alkenyl; C_{2-6} alkynyl; halo; $-C(O)R^c$; $-NR^dR^e$; $-C(O)NR^dR^e$; $-C(S)NR^dR^e$;

$-C(=N-OH)-C_{1-4}$ alkyl; SO_2C_{1-4} alkyl; cyano; C_{3-6} cycloalkyl optionally substituted with one or more R^f substituents; or a phenyl, monocyclic heteroaryl, or heterocycloalkyl ring, each ring optionally substituted with one or more R^g substituents;

wherein each R^b substituent is independently selected from the group consisting of $-OH$, $-C_{1-4}$ alkoxy, $-NR^dR^e$, $-C(O)NR^dR^e$, $-SC_{1-4}$ alkyl, $-SO_2C_{1-4}$ alkyl, cyano, halo, and monocyclic heteroaryl;

R^c is C_{1-4} alkyl, $-C_{1-4}$ haloalkyl, C_{3-6} cycloalkyl, or a monocyclic, carbon-linked heterocycloalkyl;

R^d is H or C_{1-4} alkyl;

R^e is H; C_{1-4} alkyl optionally substituted with $-CN$, $-CF_3$, $-OH$, or a monocyclic heterocycloalkyl; C_{3-6} cycloalkyl; $-OH$; or $-OC_{1-4}$ alkoxy;

or R^d and R^e taken together with the nitrogen to which they are attached form a heterocycloalkyl, optionally substituted with C_{1-4} alkyl or OH;

each R^f substituent is independently selected from the group consisting of: C_{1-4} alkyl optionally substituted with $-OH$, cyano, or C_{1-4} alkoxy; C_{1-4} haloalkyl; $-CONH_2$; and cyano; and

each R^g substituent is independently selected from the group consisting of C_{1-4} alkyl, CF_3 , halo, $-NH_2$, $-OCH_3$, cyano, and $-OH$;

R^1 is selected from the group consisting of H, C_{1-6} alkyl, C_{1-4} haloalkyl, and C_{3-6} cycloalkyl; or R^a and R^1 taken together with the carbons to which they are attached form a 5- to 7-membered ring, optionally containing an O or NH, and optionally substituted with one or more R^h substituents;

wherein each R^h substituent is independently $-C(O)NR^iR^j$, cyano, or is C_{1-4} alkyl optionally substituted with $-OH$, OCH_3 , cyano, or $-C(O)NR^iR^j$; or two R^h groups attached to the same carbon and taken together with the carbon to which they are attached form a carbonyl or a C_{3-6} cycloalkyl;

wherein R^i and R^j are each independently H or C_{1-4} alkyl;

R^2 is $-R^m$, $-OR^m$, or $-NR^mR^n$;

wherein R^m is phenyl, naphthyl, pyridyl, pyrazinyl, pyridazinyl, pyrrolyl, pyrazolyl, triazolyl, imidazolyl, furanyl, oxazolyl, isoxazolyl, thiophenyl, thiazolyl, isothiazolyl, indolyl, indazolyl, quinolinyl, or isoquinolinyl, each optionally substituted with one or more R^s substituents;

wherein each R^s substituent is independently selected from the group consisting of C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy, C_{1-4} alkyl-OH, C_{1-4} haloalkoxy, halo, cyano, C_{3-6} cycloalkyl, $NHSO_2C_{1-4}$ alkyl, and $-SO_2C_{1-4}$ alkyl;

492

R^n is H, C_{1-4} haloalkyl, or C_{1-4} alkyl optionally substituted with $-OH$ or C_{1-4} alkoxy;

or R^m and R^n taken together with the nitrogen to which they are attached form a pyrrolidine or piperidine ring, optionally substituted with C_{1-4} alkyl and optionally fused to phenyl, wherein said phenyl is optionally substituted with halo;

R^3 is H or methyl; and

R^4 is H or fluoro;

10 or a pharmaceutically acceptable salt thereof.

3. The method of claim 1, wherein R^a is C_{1-6} alkyl optionally substituted with one or more R^b substituents.

4. The method of claim 1, wherein R^a is C_{1-6} alkyl optionally substituted with one or two R^b substituents.

5. The method of claim 1, wherein R^a is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, or isopentyl, each optionally substituted with one or more R^b substituents.

6. The method of claim 1, wherein each R^b is independently $-OH$, methoxy, ethoxy, $-NR^dR^e$, $-C(O)NR^dR^e$, thiomethyl, thioethyl, methanesulfonyl, ethanesulfonyl, cyano, fluoro, chloro, bromo, pyrrolyl, pyrazolyl, imidazolyl, furanyl, thiophenyl, triazolyl, tetrazolyl, oxazolyl, or thiazolyl.

7. The method of claim 1, wherein each R^b is independently $-OH$, $-C(O)NHCH_3$, $-CF_3$, methoxy, ethoxy, fluoro, $-C(O)NH_2$, $-C(O)N(CH_3)_2$, $-N(CH_3)_2$, methanesulfonyl, thiomethyl, cyano, pyrazolyl, 6-oxa-1-azaspiro [3.3]heptan-1-yl, azetidiny, 3-hydroxyazetidiny, pyrrolidinyl, or hydroxyethylamino.

8. The method of claim 1, wherein R^a is C_{1-6} alkenyl or C_{1-6} alkynyl.

9. The method of claim 1, wherein R^a is ethenyl, isopropenyl, or propynyl.

10. The method of claim 1, wherein R^a is halo.

11. The method of claim 1, wherein R^a is bromo, chloro, fluoro, or iodo.

12. The method of claim 1, wherein R^a is $-C(O)R^c$; $-NR^dR^e$; $-C(O)NR^dR^e$; $C(S)NR^dR^e$; $-C(=N-OH)-C_{1-4}$ alkyl; or SO_2C_{1-4} alkyl.

13. The method of claim 1, wherein R^e is methyl, ethyl, propyl, isopropyl, butyl, fluoromethyl, difluoromethyl, trifluoromethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, azetidiny, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, oxetanyl, tetrahydrofuranyl, or tetrahydropyranlyl.

14. The method of claim 1 wherein R^c is ethyl, cyclopropyl, methyl, oxetanyl, or trifluoromethyl.

15. The method of claim 1, wherein R^d is H, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, or tert-butyl.

16. The method of claim 1, wherein R^e is H, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, cyanomethyl, trifluoroethyl, hydroxyethyl, 2-hydroxy-1-methylethyl, hydroxypropyl, cyclopropyl, hydroxy, methoxy, or oxetanylmethyl.

17. The method of claim 1, wherein R^d and R^e taken together with the nitrogen to which they are attached form azetidiny, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranlyl, or 6-oxa-1-azaspiro[3.3]heptan-1-yl, each optionally substituted with C_{1-4} alkyl or $-OH$.

18. The method of claim 1, wherein R^a is cyano.

19. The method of claim 1, wherein R^a is C_{3-6} cycloalkyl optionally substituted with one or more R^f substituents.

20. The method of claim 1, wherein R^a is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl, each optionally substituted with one or more R^f substituents.

493

21. The method of claim 1, wherein R^a is cyclopropyl, optionally substituted with one or more R^f substituents.

22. The method of claim 1, wherein each R^f is independently: methyl, ethyl, propyl, or isopropyl, each optionally substituted with —OH, cyano, methoxy, or ethoxy; C_{1-4} fluoroalkyl; —CONH₂; or cyano.

23. The method of claim 1, wherein each R^f is independently hydroxymethyl, methyl, cyano, trifluoromethyl, cyanomethyl, methoxymethyl, fluoromethyl, hydroxymethyl, 1-hydroxy-1-methyl-ethyl, or —CONH₂.

24. The method of claim 1, wherein R^a is a phenyl, monocyclic heteroaryl, or heterocycloalkyl ring, each ring optionally substituted with one or more R^g substituents.

25. The method of claim 1, wherein R^a is a phenyl, pyrrolyl, furanyl, thiophenyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, triazolyl, pyridyl, pyrimidinyl, pyridazinyl, pyrazinyl, or pyrimidinyl, each optionally substituted with one or more R^g substituents.

26. The method of claim 1, wherein R^a is optionally substituted with one or two R^g substituents.

27. The method of claim 1, wherein each R^g is independently methyl, ethyl, propyl, isopropyl, —CF₃, fluoro, chloro, —NH₂, —OCH₃, cyano, or —OH.

28. The method of claim 1, wherein each R^g is independently fluoro, methyl, —NH₂, —CF₃, chloro, methoxy, or cyano.

29. The method of claim 1, wherein R^a and R^1 taken together with the carbons to which they are attached form a 5- to 7-membered ring, optionally containing an O or NH, and optionally substituted with one or more R^h substituents.

30. The method of claim 1, wherein R^a and R^1 taken together with the carbons to which they are attached form cyclopentenyl, cyclohexenyl, dihydrofuranyl, dihydropyranyl, dihydropyrrolyl, or tetrahydropyridine, each optionally substituted with one or more R^h substituents.

31. The method of claim 1, wherein each R^h is independently: methyl, ethyl, or propyl, each optionally substituted with hydroxy, cyano, methoxy, or —C(O)N(CH₃)₂; C(O)NR^{*h*}R^{*h*}; or cyano.

32. The method of claim 1, wherein each R^h is independently hydroxypropyl, hydroxyethyl, hydroxymethyl, methyl, cyano, methoxymethyl, —C(O)NH₂, or CH₂C(O)N(CH₃)₂.

33. The method of claim 1, wherein two R^h groups attached to the same carbon are taken together with the carbon to which they are attached to form cyclopentyl or a carbonyl.

34. The method of claim 1, wherein R^1 is H, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, fluoromethyl, fluoromethyl, trifluoromethyl, fluoroethyl, trifluoroethyl, cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.

35. The method of claim 1, wherein R^1 is H, methyl, isopropyl, trifluoromethyl, or cyclopropyl.

36. The method of claim 1, wherein R^2 is R^m .

37. The method of claim 1, wherein R^2 is —OR^{*m*}.

38. The method of claim 1, wherein R^2 is —NR^{*m*}R^{*m*}.

39. The method of claim 1, wherein R^m is phenyl, naphthyl, pyridyl, indazolyl, or isoquinolinyl, each optionally substituted with one or more R^s substituents.

40. The method of claim 1, wherein R^m is pyrazolyl, optionally substituted with one or more R^s substituents.

41. The method of claim 1, wherein R^m is phenyl, optionally substituted with one or more R^s substituents.


42. The method of claim 1, wherein R^m is optionally substituted with one or two R^s substituents.

494

43. The method of claim 1, wherein each R^s is independently methyl, ethyl, propyl, isopropyl, butyl, fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, trifluoroethyl, methoxy, ethoxy, isopropoxy, hydroxymethyl, hydroxyethyl, trifluoromethoxy, fluoro, chloro, bromo, iodo, cyano, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, NHSO₂C₁₋₂alkyl, or —SO₂C₁₋₂alkyl.

44. The method of claim 1, wherein each R^s is independently fluoro, chloro, trifluoromethyl, cyano, methyl, methoxy, cyclopropyl, —NHSO₂CH₃, fluoroethyl, ethyl, propyl, difluoromethyl, hydroxymethyl, fluoromethyl, or methanesulfonyl.

45. The method of claim 1, wherein R^2 is R^m and R^m is



wherein at least one of X¹, X², and X³ is N, and the other two are independently N, NR, O, S, or C—R^{*r*};

R^{*p*} and R^{*r*} are each independently H; C₁₋₄haloalkyl; C₁₋₄alkyl optionally substituted with OH; halo; cyano; or C₃₋₆cycloalkyl; and

R^{*q*} is H or fluoro;

or R^{*q*} and R^{*r*} taken together with the carbons to which they are attached form phenyl, optionally substituted with halo.

46. The method of claim 45, wherein X¹ and X² are each N and X³ is C—R^{*r*}.

47. The method of claim 45, wherein X² is N and X¹ and X³ are each independently C—R^{*r*}.

48. The method of claim 45, wherein X¹, X², and X³ are each N.

49. The method of claim 45, wherein R^{*p*} and R^{*r*} are each independently H, fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, trifluoroethyl, methyl, ethyl, hydroxymethyl, hydroxyethyl, chloro, cyano, cyclopropyl, cyclobutyl, or cyclopentyl.

50. The method of claim 45, wherein R^{*p*} is trifluoromethyl, chloro, methyl, hydroxyethyl, cyclopropyl, cyano, difluoromethyl, or ethyl.

51. The method of claim 45, wherein R^{*r*} is ethyl, trifluoromethyl, methyl, chloro, H, hydroxyethyl, cyclopropyl, or cyano.

52. The method of claim 45, wherein R^{*q*} is H or fluoro.

53. The method of claim 45, wherein R^{*q*} and R^{*r*} taken together with the carbons to which they are attached form phenyl, optionally substituted with fluoro.

54. The method of claim 1, wherein R^{*r*} is H, fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, difluoroethyl, or trifluoroethyl, or is methyl or ethyl optionally substituted with —OH, methoxy, or ethoxy.

55. The method of claim 1, wherein R^{*r*} is H, methyl, ethyl, fluoroethyl, difluoroethyl, or trifluoroethyl.

56. The method of claim 1, wherein R^{*m*} and R^{*n*} taken together with the nitrogen to which they are attached form dihydroindole, optionally substituted with methyl or fluoro.

57. The method of claim 1 wherein R³ is H.

58. The method of claim 1, wherein R³ is methyl.

59. The method of claim 1, wherein R⁴ is H.

60. The method of claim 1, wherein R⁴ is fluoro.

61. A method of treating a subject suffering from or diagnosed with depression or anxiety mediated by NR2A activity, comprising administering to the subject in need of such treatment an effective amount of a compound selected from:

N-(cyanomethyl)-7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(4-Fluorophenoxymethyl)-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide;

3-[(Azetidin-1-yl)carbonyl]-7-(4-fluorophenoxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

N-ethyl-7-(4-fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(3,4-Difluorophenoxymethyl)-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-7-(4-fluorophenoxymethyl)-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-((4-fluorophenoxy)methyl)-N-hydroxy-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-N-(propan-2-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(4-Fluorophenoxymethyl)-N-(2-hydroxyethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(4-Fluorophenoxymethyl)-N-(1-hydroxypropan-2-yl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-((4-fluorophenoxy)methyl)-2-methyl-N-(oxetan-3-ylmethyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-((4-fluorophenoxy)methyl)-N-(3-hydroxypropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-cyclopropyl-7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-((4-fluorophenoxy)methyl)-N-methoxy-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(4-Fluorophenoxymethyl)-N,2-dimethyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carbothioamide;

7-((4-fluorophenoxy)methyl)-2-methyl-3-propionyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((4-fluorophenoxy)methyl)-3-(1-hydroxypropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-(4-Fluorophenoxymethyl)-3-(1-hydroxyethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

7-(4-Fluorophenoxymethyl)-3-(2-hydroxypropan-2-yl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

3-acetyl-7-((4-fluorophenoxy)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

2-(7-((4-fluorophenoxy)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-3-yl)-N-methylacetamide;

3-Cyclopropanecarbonyl-7-(4-fluorophenoxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

7-(4-Fluorophenoxymethyl)-3-[1-(hydroxyimino)ethyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

7-((4-fluorophenoxy)methyl)-2-methyl-3-(oxetane-3-carbonyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((4-fluorophenoxy)methyl)-2-methyl-3-(2,2,2-trifluoro-1-hydroxyethyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-(4-Fluorophenoxymethyl)-2-methyl-3-(trifluoroacetyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

2-cyclopropyl-N-ethyl-7-((4-fluorophenoxy)methyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(4-Fluorophenoxymethyl)-N-methyl-5-oxo-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide;

2-Cyclopropyl-7-(4-fluorophenoxymethyl)-N-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-Ethyl-7-(4-fluorophenoxymethyl)-5-oxo-2-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-((4-fluorophenoxy)methyl)-N-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-7-[[5-fluoropyridin-2-yl]oxy]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(4-Fluorophenoxymethyl)-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

7-((3-fluorophenoxy)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((4-fluorophenoxy)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-(2,4-Difluorophenoxymethyl)-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

7-(3,4-Difluorophenoxymethyl)-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

7-(4-Chlorophenoxymethyl)-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-Fluoropyridin-2-yl]oxy]methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-7-((4-trifluoromethyl)phenoxy)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((4-fluorophenoxy)methyl)-2-methyl-3-(oxazol-2-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((2-fluorophenoxy)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

4-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methoxy)benzotrile;

7-((4-fluorophenoxy)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((4-fluorophenoxy)methyl)-2-methyl-3-(1H-pyrazol-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((4-fluorophenoxy)methyl)-2-methyl-3-(4H-1,2,4-triazol-3-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

3-cyclopropyl-7-[(4-fluorophenoxy)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

cis-2-[7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carbonitrile;

trans-2-[7-(4-Fluorophenoxymethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropane-1-carbonitrile;

7-(4-Fluorophenoxymethyl)-3-[cis-2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

trans-7-(4-Fluorophenoxyethyl)-2-methyl-3-[2-(trifluoromethyl)cyclopropyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

7-(4-Fluorophenoxyethyl)-2-methyl-3-(2-methylcyclopropyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

trans-2-[2-[7-(4-Fluorophenoxyethyl)-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropyl]acetonitrile;

7-(4-Fluorophenoxyethyl)-3-[2-(methoxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

3-(2-(fluoromethyl)cyclopropyl)-7-((4-fluorophenoxy)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

6-fluoro-7-((4-fluorophenoxy)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-(4-Fluorophenoxyethyl)-3-(3-hydroxypropyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

7-(4-Fluorophenoxyethyl)-3-(methoxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

7-((4-fluorophenoxy)methyl)-3-(3-hydroxyoxetan-3-yl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-(4-Fluorophenoxyethyl)-3-(4-hydroxybutan-2-yl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

7-(4-Fluorophenoxyethyl)-3-[2-(2-hydroxypropan-2-yl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-pyrimidin-5-yl-thiazolo[3,2-a]pyrimidin-5-one;

7-(((4-fluorophenyl)(methyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-(((4-fluorophenyl)(2,2,2-trifluoroethyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-(((3,4-difluorophenyl)(ethyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((ethyl(3-fluorophenyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-(((2,2-difluoroethyl)(4-fluorophenyl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((ethyl(pyridine-2-yl)amino)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(thiazol-2-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(thiophen-2-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

3-(ethyl((2-methyl-5-oxo-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)amino)benzotrile;

3-(2-aminopyridin-3-yl)-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((ethyl(pyridine-2-yl)amino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((4-fluorophenylamino)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

3-butyl-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

2-cyclopropyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-3-pyrimidin-5-yl-thiazolo[3,2-a]pyrimidin-5-one;

2-ethyl-7-((ethyl(4-fluorophenyl)amino)methyl)-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((ethyl(4-fluorophenyl)amino)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-thiazol-4-yl-thiazolo[3,2-a]pyrimidin-5-one;

7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-phenylthiazolo[3,2-a]pyrimidin-5-one;

7-[(N-ethyl-4-fluoro-anilino)methyl]-2,3-dimethyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(5-fluoro-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(2-fluoro-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

3-cyclopropyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-pyrazin-2-yl-thiazolo[3,2-a]pyrimidin-5-one;

7-[(N-ethyl-4-fluoro-anilino)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-5-one;

7-[(N-ethyl-4-fluoro-anilino)methyl]-3-isopropenyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-pyridazin-4-yl-thiazolo[3,2-a]pyrimidin-5-one;

3-(5-chloro-3-pyridyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(4-pyridyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(1-methylpyrazol-4-yl)thiazolo[3,2-a]pyrimidin-5-one;

7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(1H-pyrazol-4-yl)thiazolo[3,2-a]pyrimidin-5-one;

5-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]pyridine-3-carbonitrile;

7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-[5-(trifluoromethyl)-3-pyridyl]thiazolo[3,2-a]pyrimidin-5-one;

7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-oxazol-2-yl-thiazolo[3,2-a]pyrimidin-5-one;

7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((ethyl(4-fluorophenyl)amino)methyl)-3-(trans-2-(fluoromethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

3-ethyl-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-propyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-[[4-fluoro-N-(2-fluoroethyl)anilino]methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-((ethyl(4-fluorophenyl)amino)methyl)-3-(furan-3-yl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((ethyl(4-fluorophenyl)amino)methyl)-3-(furan-2-yl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((5-fluoro-2-methylindolin-1-yl)methyl)-3-(furan-2-yl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(thiophen-3-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((5-fluoro-2-methylindolin-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(4-methylthiazol-2-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(6-oxo-1,6-dihydropyridin-3-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

3-(6-aminopyridin-3-yl)-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
 7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(prop-1-ynyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;
 7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-vinyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
 3-bromo-2-cyclopropyl-7-[(N-ethyl-4-fluoro-anilino)methyl]thiazolo[3,2-a]pyrimidin-5-one;
 3-(3,5-difluorophenyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(1-methylpyrazolo-3-yl)thiazolo[3,2-a]pyrimidin-5-one;
 3-(2-amino-4-pyridyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(5-methoxy-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
 7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-morpholino-5H-thiazolo[3,2-a]pyrimidin-5-one;
 3-(dimethylamino)-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
 7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-(pyrrolidin-1-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;
 7-(((3,4-difluorophenyl)(ethyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile;
 7-((ethyl(3-fluorophenyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carbonitrile;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carbonitrile;
 7-((5-fluoro-2-methylindolin-1-yl)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile;
 7-(((3-cyanophenyl)(ethyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile;
 7-((ethyl(pyridin-2-yl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile;
 2-methyl-7-((2-methylindolin-1-yl)methyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile;
 7-(((3,5-difluorophenyl)(ethyl)amino)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile;
 6-[(N-ethyl-4-fluoro-anilino)methyl]-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one;
 2-[(N-ethyl-4-fluoro-anilino)methyl]-6,7,8,9-tetrahydropyrimido[2,1-b][1,3]benzothiazol-4-one;
 6-[(N-ethyl-4-fluoro-anilino)methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one;
 6-[(N-ethyl-4-fluoro-anilino)methyl]spiro[2,3-dihydrocyclopenta[3,4]thiazolo[1,4-a]pyrimidine-1,1'-cyclopentane]-8-one;
 6-[(N-ethyl-4-fluoro-anilino)methyl]-2,3-dihydrocyclopenta[3,4]thiazolo[1,4-a]pyrimidine-1,8-dione;
 6-[(N-ethyl-4-fluoro-anilino)methyl]-1,1-dimethyl-2,3-dihydrocyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one;
 2-[(N-ethyl-4-fluoro-anilino)methyl]-8,9-dihydro-6H-pyrano[3,4]thiazolo[1,4-a]pyrimidin-4-one;
 2-[(N-ethyl-4-fluoro-anilino)methyl]-7-methyl-8,9-dihydro-6H-pyrido[3,4]thiazolo[1,4-a]pyrimidin-4-one;
 6-[(N-ethyl-4-fluoro-anilino)methyl]-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidine-1-carboxamide;
 6-[(N-ethyl-4-fluoro-anilino)methyl]-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidine-1-carbonitrile;

2-[6-[(N-ethyl-4-fluoro-anilino)methyl]-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-1-yl]acetoneitrile;
 6-[(N-ethyl-4-fluoro-anilino)methyl]-1-(2-hydroxyethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one;
 2-((ethyl(4-fluorophenyl)amino)methyl)-6-(methoxymethyl)-7,8-dihydrocyclopenta[4,5]thiazolo[3,2-a]pyrimidin-4(6H)-one;
 2-[6-[(N-ethyl-4-fluoro-anilino)methyl]-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-1-yl]acetamide;
 3-cyclohexyl-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-isopropyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
 3-cyclopentyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-tetrahydropyran-4-yl-thiazolo[3,2-a]pyrimidin-5-one;
 3-cyclobutyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
 3-tert-butyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
 3-acetyl-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-hydroxy-1-methyl-ethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-hydroxyethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
 3-[[dimethylamino)methyl]-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 3-(azetidin-1-ylmethyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(pyrrolidin-1-ylmethyl)thiazolo[3,2-a]pyrimidin-5-one;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-[(3-hydroxyazetidin-1-yl)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(2,2,2-trifluoro-1-hydroxy-ethyl)thiazolo[3,2-a]pyrimidin-5-one;
 7-((ethyl(4-fluorophenyl)amino)methyl)-3-(hydroxymethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
 7-[[ethyl(4-fluorophenyl)amino]methyl]-3-(methoxymethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[[Ethyl(4-fluorophenyl)amino]methyl]-2-methyl-3-(1H-pyrazol-1-ylmethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(methylsulfonylmethyl)thiazolo[3,2-a]pyrimidin-5-one;
 3-(ethoxymethyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
 2-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]acetoneitrile;
 3-tert-butyl-7-[(N-ethyl-4-fluoro-anilino)methyl]thiazolo[3,2-a]pyrimidin-5-one;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-hydroxy-1-methyl-ethyl)thiazolo[3,2-a]pyrimidin-5-one;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-hydroxyethyl)thiazolo[3,2-a]pyrimidin-5-one;

501

- 7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-(6-oxa-1-azaspiro[3.3]heptan-1-ylmethyl)thiazolo[3,2-a]pyrimidin-5-one;
- 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-[(2-hydroxyethylamino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 3-(ethyl((3-(hydroxymethyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)amino)benzoni-trile;
- 3-[ethyl-[[3-(methoxymethyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]amino]benzoni-trile;
- 7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-3-((methylthio)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 3-chloro-7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
- 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-fluoro-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-iodo-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 3-chloro-7-[(N-ethyl-4-fluoro-anilino)methyl]thiazolo[3,2-a]pyrimidin-5-one;
- 7-[(N-ethylamino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 3-(1,3-dihydroxypropyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(1-fluoro-3-hydroxy-propyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 3-(1,2-dihydroxyethyl)-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 7-[(N-ethyl-4-fluoro-anilino)methyl]-3-(3-hydroxypropyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 7-((Ethyl(4-fluorophenyl)amino)methyl)-3-(3-methoxypropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 7-((Ethyl(4-fluorophenyl)amino)methyl)-3-(4-hydroxybutyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 7-((Ethyl(4-fluorophenyl)amino)methyl)-3-(2-hydroxyethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 7-[[Ethyl(4-fluorophenyl)amino]methyl]-3-(2-methoxyethyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
- 3-(7-[[ethyl(4-fluorophenyl)amino]methyl]-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidin-3-yl)propanamide;
- 3-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]propanenitrile;
- 3-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N,N-dimethyl-propanamide;
- 7-[[Ethyl(4-fluorophenyl)amino]methyl]-3-(3-hydroxy-3-methylbutyl)-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
- 2-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarboxamide;
- 7-((5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 7-((3,5-bis(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 7-((3-chloro-5-methyl-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

502

- 7-((5-chloro-3-methyl-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 7-((1H-indazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 7-((5-(2-hydroxyethyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 7-((3-(2-hydroxyethyl)-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 7-((3-cyclopropyl-4-fluoro-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 7-((5-cyclopropyl-4-fluoro-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 7-[[5-cyclopropyl-3-methyl-pyrazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 7-[[3-cyclopropyl-5-methyl-pyrazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 7-[[3,5-dicyclopropylpyrazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 7-((3,5-dimethyl-1H-pyrazol-1-yl)methyl)-6-fluoro-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 1-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-3-(trifluoromethyl)-1H-pyrazole-5-carbonitrile;
- 7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-6-fluoro-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
- 1-[[3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]-5-methyl-pyrazole-3-carbonitrile;
- 7-[[3-(difluoromethyl)-5-methyl-pyrazol-1-yl]methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 7-[[6-fluoroindazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-isopropyl-thiazolo[3,2-a]pyrimidin-5-one;
- 7-[[3,5-dimethylpyrazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 7-[[5-fluoroindazol-1-yl)methyl]-3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
- 5-ethyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

503

7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((3,5-bis(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

2-methyl-3-(pyrimidin-5-yl)-7-((5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

2-methyl-7-((5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((3,5-bis(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((1H-indazol-1-yl)methyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-2-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-2-(trifluoromethyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

2-cyclopropyl-7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

2-cyclopropyl-7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

2-methyl-5-oxo-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile;

7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile;

7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile;

3-(1-hydroxyethyl)-2-methyl-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

3-acetyl-7-((5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

3-acetyl-7-((3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

3-(2-hydroxypropan-2-yl)-2-methyl-7-((3-(trifluoromethyl)-1H-pyrazol-1-yl)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-((5-fluoro-3-methyl-1H-indazol-1-yl)methyl)-3-(1-hydroxyethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

7-(1-(5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)ethyl)-3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

504

7-(1-(5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl)ethyl)-3-(2-(hydroxymethyl)-1-methylcyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzoxonitrile;

3-((6-fluoro-3-(2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzoxonitrile;

2-fluoro-3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzoxonitrile;

3-(trans-2-(hydroxymethyl)cyclopropyl)-7-(isoquinolin-4-ylmethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;

3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-7-(3-(trifluoromethyl)benzyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-4-methylbenzoxonitrile;

4-fluoro-3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzoxonitrile;

3-fluoro-5-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzoxonitrile;

2-fluoro-5-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzoxonitrile;

3-[[3-[trans-2-(hydroxymethyl)cyclopropyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]-4-methoxy-benzoxonitrile;

3-(2-(hydroxymethyl)cyclopropyl)-2-methyl-7-(4-(trifluoromethyl)benzyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;

4-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)picolinonitrile;

4-cyclopropyl-3-((3-(trans-2-(hydroxymethyl)cyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzoxonitrile;

7-(3-cyanobenzyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(3-cyano-2-fluorobenzyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(3-chloro-2-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-7-(2-fluoro-3-methylbenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(2-chloro-5-cyanobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-2-methyl-5-oxo-7-(3-(trifluoromethyl)benzyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(3-cyanobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(3-cyano-2-fluorobenzyl)-2-cyclopropyl-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(3-cyano-5-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(3-cyano-2-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[3-(chloro-2-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

505

N,2-dimethyl-5-oxo-7-[[3-(trifluoromethyl)phenyl]methyl]thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(3-chlorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[[2-cyclopropyl-5-(trifluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-2-methyl-5-oxo-7-((6-(trifluoromethyl)pyridine-2-yl)methyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-7-(2-ethyl-4,5-difluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[[3-(difluoromethyl)-2-fluoro-phenyl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-((6-cyanopyridin-2-yl)methyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[[3-(difluoromethyl)-2-fluoro-phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[[2-fluoro-3-(hydroxymethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(3-cyclopropyl-2-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(3-cyano-2-fluorobenzyl)-N-ethyl-6-fluoro-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(3-cyano-2-fluorobenzyl)-6-fluoro-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(3-cyano-2-fluorobenzyl)-N-ethyl-6-fluoro-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 6-fluoro-7-(2-fluoro-3-(trifluoromethyl)benzyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(5-cyano-2-methylbenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(5-cyano-2-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(2-chloro-5-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(3-cyano-2-fluorobenzyl)-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-7-(5-fluoro-2-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-2-methyl-7-(naphthalen-1-ylmethyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-7-(5-fluoro-2-methylbenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(3-cyano-4-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-2-methyl-7-(1-methyl-1H-indazol-4-yl)methyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-2-methyl-7-(3-(methylsulfonamido)benzyl)-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(5-cyano-2-(trifluoromethyl)benzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(4-chloro-2-methylbenzyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(2,5-difluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(3-cyanobenzyl)-2-cyclopropyl-N-ethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-7-(2-fluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(2,3-difluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;

506

N-ethyl-7-(3-fluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(3-chloro-4-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[[2,5-dichlorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N,2-dimethyl-5-oxo-7-[[3-(trifluoromethoxy)phenyl]methyl]thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(5-cyano-2-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(3-chloro-5-cyano-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(3-cyclopropylphenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(2,5-difluorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(3,4-difluorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(2,3-difluorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(4-chloro-2-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(2,4-dichlorophenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(3-fluoro-4-methyl-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[[4-fluoro-2-(trifluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(2-cyclopropyl-4-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(5-cyano-2-(2-fluoroethyl)benzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(2-chloro-3-cyanobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-7-(6-ethyl-2,3-difluorobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(5-cyano-2-ethylbenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(2-cyclopropyl-5-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(5-cyano-2-cyclopropylbenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-7-(5-fluoro-2-propylbenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[[2-fluoro-3-(fluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-7-(1-(2-fluoro-3-(trifluoromethyl)phenyl)ethyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-7-(2-fluoro-3-(trifluoromethyl)benzyl)-N,2-dimethyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 3-((3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile;
 3-((3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzotrile;
 6-((3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)picolinonitrile;
 3-((3-(2-cyanocyclopropyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-4-methoxybenzotrile;
 2-methyl-3-(pyrimidin-5-yl)-7-(3-(trifluoromethyl)benzyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;
 7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;
 2-fluoro-3-((2-methyl-5-oxo-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzotrile;

507

3-((3-cyclopropyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzonitrile;
 7-(isoquinolin-4-ylmethyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;
 3-((2-methyl-5-oxo-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile;
 7-(5-fluoro-2-methoxybenzyl)-2-methyl-3-(pyrimidin-5-yl)-5H-thiazolo[3,2-a]pyrimidin-5-one;
 2-fluoro-3-((3-(furan-2-yl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile;
 3-bromo-2-methyl-7-(3-(methylsulfonyl)benzyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;
 7-(3-cyano-2-fluorobenzyl)-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-(cyanomethyl)-7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-(3-cyanobenzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile;
 7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carbonitrile;
 10-(4-fluorophenoxymethyl)-3-(hydroxymethyl)-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-trien-12-one;
 10-(4-Fluorophenoxymethyl)-3-(2-hydroxyethyl)-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-trien-12-one;
 7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-3-methylsulfonyl-thiazolo[3,2-a]pyrimidin-5-one;
 3-(hydroxymethyl)-10-[[3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-trien-12-one;
 10-[[5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-trien-12-one;
 10-[[3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-trien-12-one;
 10-[[5-cyclopropyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-3,3-dimethyl-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-trien-12-one;
 10-[[3-cyclopropyl-5-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-3,3-dimethyl-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-trien-12-one;
 10-[[3,5-bis(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-3,3-dimethyl-7-thia-1,9-diazatricyclo[6.4.0.0^{2,6}]dodeca-2(6),8,10-trien-12-one;
 3-((3-acetyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)-2-fluorobenzonitrile;
 7-(2-fluoro-3-(trifluoromethyl)benzyl)-3-(1-hydroxyethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
 3-((3-acetyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)methyl)benzonitrile;
 7-(2-fluoro-3-(trifluoromethyl)benzyl)-3-(hydroxymethyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one;
 7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-3-((methylamino)methyl)-5H-thiazolo[3,2-a]pyrimidin-5-one;
 7-[(3-cyclopropyl-2-fluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-2-methyl-5-oxo-7-[(2,3,6-trifluorophenyl)methyl]thiazolo[3,2-a]pyrimidine-3-carboxamide;
 2-fluoro-3-[(2-methyl-3-oxazol-2-yl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl)methyl]benzonitrile;

508

7-[(5-cyano-3-cyclopropyl-2-fluoro-phenyl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-7-[(2-fluoro-3-methoxy-phenyl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(3-cyclopropyl-2-fluoro-phenyl)methyl]-6-fluoro-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 2-[7-[(3-chloro-2-fluoro-phenyl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide;
 7-[(3-chloro-2-fluoro-phenyl)methyl]-N-ethyl-6-fluoro-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(4,5-difluoro-2-methoxy-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 2-fluoro-3-[[2-methyl-3-(2-methylcyclopropyl)-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzonitrile;
 2-[7-[(3-cyclopropyl-2-fluoro-phenyl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide;
 N-ethyl-6-fluoro-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(3-cyano-2-fluoro-phenyl)methyl]-6-fluoro-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(2-chloro-4,5-difluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[[4,5-difluoro-2-(2-fluoroethyl)phenyl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 2-[7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide;
 2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide;
 7-[(5-chloro-3-methyl-pyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(3-chloro-5-methyl-pyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(3-chloro-5-cyclopropyl-pyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(5-chloro-3-cyclopropyl-pyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-7-[[4-(trifluoromethyl)thiazol-2-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one;
 2-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide;
 N-ethyl-2-methyl-7-[[5-methyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-2-methyl-7-[[3-methyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[3-(difluoromethyl)-2-fluoro-phenyl]methyl]-6-fluoro-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[3-(difluoromethyl)-2-fluoro-phenyl]methyl]-N-ethyl-6-fluoro-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-(4-bicyclo[4.2.0]octa-1,3,5-trienylmethyl)-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-7-[[2-fluoro-3-(1-hydroxycyclopropyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-7-[[2-fluoro-3-(1-fluorocyclopropyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-7-[[2-fluoro-3-[1-(fluoromethyl)vinyl]phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[(2-ethynyl-4,5-difluoro-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

2-fluoro-3-[[2-methyl-5-oxo-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzonitrile;

3-[[3-(2,2-difluorocyclopropyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluoro-benzonitrile;

N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[(3-cyano-2-fluoro-phenyl)methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[3-(difluoromethyl)-2-fluoro-phenyl]methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-7-[[5-methyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-7-[[3-methyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-(1H-imidazol-2-yl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[(3-cyano-2-fluoro-5-methyl-phenyl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[(3-chloro-2-fluoro-phenyl)methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

2-methyl-7-[[5-methyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

2-chloro-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-methyl-7-[[5-methyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-methyl-7-[[3-methyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[(3-chloro-2-fluoro-phenyl)methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

2-methyl-7-[[3-methyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

2-chloro-7-[(3-cyano-2-fluoro-phenyl)methyl]-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[(3-cyano-2-fluoro-phenyl)methyl]-N-methyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]acetamide;

N,2-dimethyl-5-oxo-7-[[4-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidine-3-carboxamide;

3-[[2-chloro-5-oxo-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluoro-benzonitrile;

2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide;

7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

2-[7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide;

2-[7-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide;

511

7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(methylsulfonylmethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-(1H-imidazol-2-ylmethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one; 5

N-ethyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-7-[(N-ethyl-4-fluoro-anilino)methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide; 10

2-fluoro-3-[[3-(2-methylcyclopropyl)-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzotriazole;

3-[[2-chloro-3-(2-methylcyclopropyl)-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluoro-benzonitrile; 15

N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-isopropyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

6-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one; 20

6-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one; 25

N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methoxy-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

6-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one; 30

2-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-N-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

6-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one; 35

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide; 40

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide; 45

2-[7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide;

7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-(2-hydroxycyclopropyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one; 50

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one; 55

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile; 60

2-cyano-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-N-isopropyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide; 65

512

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1-hydroxy-1-methyl-ethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1-hydroxy-1-methyl-ethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1-hydroxyethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1-hydroxyethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-N-sec-butyl-thiazolo[3,2-a]pyrimidine-3-carboxamide;

3-[[3-(azetidin-1-yl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluoro-benzonitrile;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carbonitrile;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carbonitrile;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-cyclopropyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one; 25

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-cyclopropyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(2-methylcyclopropyl)thiazolo[3,2-a]pyrimidin-5-one;

3-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

3-acetyl-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

2-chloro-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-(2-methylcyclopropyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-pyrimidin-5-yl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-chloro-2-pyridyl]-methyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

cis-2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methoxy-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methoxy-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-cyclopropyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methoxy-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methoxy-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(1H-pyrazol-5-yl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(1H-pyrazol-5-yl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-isopropenyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

trans-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(5-fluoro-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(5-fluoro-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-propanoyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-propanoyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-thiazol-2-yl-thiazolo[3,2-a]pyrimidin-5-one;

N-ethyl-7-[[5-fluoro-2-pyridyl]-methyl-amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

2-[7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

N-ethyl-7-[[ethyl-(5-fluoro-2-pyridyl)amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-7-[[ethyl(2-pyridyl)amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

3-(5-chloro-3-pyridyl)-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

3-(5-chloro-3-pyridyl)-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-thiazol-4-yl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-2-pyridyl]-ethyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-cyclopropyltriazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

trans-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-[2-methylcyclopropyl]thiazolo[3,2-a]pyrimidin-5-one;

2-ethoxy-N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-fluoro-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-oxazol-2-yl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-oxazol-2-yl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-2-pyridyl]-methyl-amino]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[3,5-dichloropyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(2,2,2-trifluoroethoxy)thiazolo[3,2-a]pyrimidine-3-carboxamide;

3-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

3-acetyl-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

3-[[4-chloropyrazol-1-yl]methyl]-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-isopropenyl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1H-imidazol-2-yl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-pyrimidin-5-yl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-pyrimidin-5-yl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-pyridyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-2-(2,2,2-trifluoroethoxy)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethoxy-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethoxy-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(2-methylpropanoyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(2-methylpropanoyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-N-[(1R)-1-methylpropyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-N-[(1S)-1-methylpropyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[4-chloro-2-pyridyl]-ethyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-fluoro-2-pyridyl]-methyl-amino]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(methoxymethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

515

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-cyclopropyl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(methylsulfonylmethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(pyrazol-1-ylmethyl)thiazolo[3,2-a]pyrimidin-5-one;

2-[7-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-cyclobutyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-cyclobutyl-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

2-[7-[[5-chloro-2-pyridyl]-methyl-amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

7-[[ethyl-(5-fluoro-2-pyridyl)amino]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

3-chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

3-chloro-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

2-[7-(4,5-difluoro-2-methoxy-phenyl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

2-[7-[[ethyl-(5-fluoro-2-pyridyl)amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

7-[[5-chloro-2-pyridyl]-ethyl-amino]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

5-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]pyridine-3-carbonitrile;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(2,2,2-trifluoroacetyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-methylcyclopropyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-methylcyclopropyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-pyrimidin-5-yl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-3-(2,2,2-trifluoro-1-hydroxy-ethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-bromo-2-pyridyl]-ethyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-7-[[ethyl-(5-fluoro-2-pyridyl)amino]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

3-(azetidin-1-yl)-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

cis-2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

516

2-[7-[[5-methoxy-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

2-[7-[[3-methoxy-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

3-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-thiazolo[3,2-a]pyrimidin-5-one;

3-acetyl-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-thiazolo[3,2-a]pyrimidin-5-one;

2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

2-[7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

3-bromo-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

3-bromo-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

3-chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methoxy-thiazolo[3,2-a]pyrimidin-5-one;

3-chloro-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methoxy-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-methylsulfonyl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

2-[7-(N-ethyl-4-fluoro-anilino)methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-1-(hydroxymethyl)cyclopropanecarbonitrile;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-hydroxy-1-methyl-ethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-hydroxy-1-methyl-ethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(cyclopropanecarbonyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

3-bromo-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one;

3-bromo-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-amino-5-chloro-pyrazol-1-yl]methyl]-3-bromo-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-amino-3-chloro-pyrazol-1-yl]methyl]-3-bromo-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]acetoneitrile;

N-ethyl-7-[[5-fluoro-2-pyridyl]-methyl-amino]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3,3-difluoroazetidin-1-yl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methoxy-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one;

2-[7-[[5-bromo-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

3-chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one;

3-chloro-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one;

2-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

2-acetyl-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(1H-1,2,4-triazol-5-yl)thiazolo[3,2-a]pyrimidin-5-one;

2-[[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

3-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2,2,2-trifluoroethoxy)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

3-acetyl-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methoxy-thiazolo[3,2-a]pyrimidin-5-one;

3-acetyl-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methoxy-thiazolo[3,2-a]pyrimidin-5-one;

3-bromo-7-[(5-chloro-3-nitro-pyrazol-1-yl)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(1H-pyrazol-5-yl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-thiazol-4-yl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(5-fluoro-3-pyridyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-propanoyl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

2-[[7-[(3,5-dichloropyrazol-1-yl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile];

2-[[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]propanenitrile];

2-[[7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]propanenitrile];

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-fluoroazetidin-1-yl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

3-(5-chloro-3-pyridyl)-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[(3,5-dichloropyrazol-1-yl)methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

3-[[3-acetyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluoro-benzonitrile;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(difluoromethyl)-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(difluoromethyl)-N-ethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

(Z)-3-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enenitrile;

(E)-3-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enamide;

(E)-3-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enenitrile;

(Z)-3-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enenitrile;

(E)-3-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enamide;

N-ethyl-7-[[5-isobutyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

cis-2-[2-chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

2-[2-chloro-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

trans-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

2-[7-[(4-chloro-1-methyl-pyrazol-3-yl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

cis-2-[2-methyl-5-oxo-7-[[3-(trifluoromethyl)pyrazol-1-yl]methyl]thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

cis-2-[7-[[5-bromo-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

cis-2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-6-fluoro-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

2-[2-methyl-7-[[1-methyl-4-(trifluoromethyl)imidazol-2-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

(E)-3-[7-[(N-ethyl-4-fluoro-anilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]prop-2-enenitrile;

7-[[4-fluorophenoxy)methyl]-3-[[2-hydroxyethyl(methyl)amino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[4-fluorophenoxy)methyl]-3-[[2-hydroxyethyl(methyl)amino)methyl]-3-[[2-hydroxyethyl(methyl)amino)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

2-[[7-[[4-fluorophenoxy)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N,N-dimethylacetamide];

7-[[2-cyano-4,5-difluoro-phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[2-cyclopropyl-4,5-difluoro-phenyl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

3-[2-(azetidin-1-yl)-2-oxo-ethyl]-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[4-fluorophenoxy)methyl]-2-methyl-3-(4H-1,2,4-triazol-3-yl)methyl]thiazolo[3,2-a]pyrimidin-5-one;

2-[[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-propanamide];

3-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-propanamide];

7-[[5-chloro-2-(trifluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[5-ethyl-1,3-benzoxazol-6-yl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[(3-chloropyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(5-chloropyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 2-[7-[(3-cyano-2-fluoro-phenyl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide;
 N-ethyl-7-[[2-fluoro-3-(1-hydroxypropyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(4,5-difluoro-2-oxazol-2-yl-phenyl)methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 2-fluoro-3-[(2-methyl-5-oxo-3-propanoyl-thiazolo[3,2-a]pyrimidin-7-yl)methyl]benzotrile;
 7-[[4,5-difluoro-2-(trifluoromethyl)phenyl]methyl]-N,2-dimethyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-N-ethyl-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-N-(2,2,2-trifluoroethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;
 3-[(2-chloro-3-cyclopropyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl)methyl]-2-fluoro-benzotrile;
 7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(pyrazol-1-ylmethyl)thiazolo[3,2-a]pyrimidin-5-one;
 N,2-dimethyl-7-[[3-methyl-4-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N,2-dimethyl-7-[[5-methyl-4-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 2-fluoro-3-[(8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-6-yl)methyl]benzotrile;
 7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-3-[hydroxy(thiazol-2-yl)methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
 2-fluoro-3-[(3-methyl-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-6-yl)methyl]benzotrile;
 2-[7-[(4-fluorophenoxy)methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]-N-methyl-acetamide;
 2-fluoro-3-[[1-(hydroxymethyl)-8-oxo-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-6-yl]methyl]benzotrile;
 7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-N-(2-hydroxy-1-methyl-ethyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 3-[[3-(2,3-dimethylcyclopropyl)-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-7-yl]methyl]-2-fluoro-benzotrile;
 6-[[3-cyclopropyl-5-(trifluoromethyl)pyrazol-1-yl]methyl]-1-(hydroxymethyl)-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one;
 7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(2-oxa-6-azaspiro[3.3]heptan-6-yl)thiazolo[3,2-a]pyrimidin-5-one;
 N-ethyl-6-fluoro-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[(4-fluorophenoxy)methyl]-5-oxo-N-(2,2,2-trifluoroethyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-cyclopentyl-7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(4,5-dichloropyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(3,4-dichloropyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-2-methyl-7-[[methyl(thiazol-2-yl)amino]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(4-chloropyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-2-methyl-7-[[methyl-(1-methylpyrazol-4-yl)amino]methyl]-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[(4-fluorophenoxy)methyl]-2-(trifluoromethyl)-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-5-one;
 7-[[3-ethoxy-2-pyridyl]-methyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[[3,5-dimethylisoxazol-4-yl)-methyl-amino]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 3-cyclopropyl-7-[(4-fluorophenoxy)methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;
 7-[(4-fluorophenoxy)methyl]-3-pyrimidin-5-yl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;
 7-[(4-fluorophenoxy)methyl]-3-[2-(hydroxymethyl)cyclopropyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;
 7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-3-(4-methyl-1,2,4-triazol-3-yl)thiazolo[3,2-a]pyrimidin-5-one;
 2-fluoro-3-[[5-oxo-2-(trifluoromethyl)-3-[2-(trifluoromethyl)cyclopropyl]thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzotrile;
 N-ethyl-7-[[ethyl(4-pyridyl)amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 N-ethyl-7-[[ethyl(3-pyridyl)amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-methoxy-3-pyridyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
 2-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;
 7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-isopropenyl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;
 3-(5-chloro-3-pyridyl)-7-[[5-chloro-2-pyridyl)-methyl-amino]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;
 7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-pyridyl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;
 7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-5-oxo-2-(2,2,2-trifluoroethoxy)thiazolo[3,2-a]pyrimidine-3-carboxamide;
 7-[[3-chloro-6-(trifluoromethyl)-2-pyridyl]methyl]-2-methyl-3-(2-methylcyclopropyl)thiazolo[3,2-a]pyrimidin-5-one;
 7-[(5-chloro-2-pyridyl)oxymethyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-ethyl-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-pyrrolidin-1-yl-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

N-ethyl-7-[[5-methoxy-2-pyridyl]-methyl-amino]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

3-(2-chloro-3-pyridyl)-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

3-(2-chloro-3-pyridyl)-7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[3-chloro-5-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-pyridyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(3-methoxyazetidin-1-yl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(cyclopropylmethyl)-2-methyl-thiazolo[3,2-a]pyrimidin-5-one;

5-[7-[[2-fluoro-3-(trifluoromethyl)phenyl]methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]pyridine-3-carbonitrile;

2-fluoro-3-[[3-[2-methylcyclopropyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]benzoxonitrile;

7-[(3,5-diisopropylpyrazol-1-yl)methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

2-[7-[(4-chloro-2-methyl-pyrazol-3-yl)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-(2-methylazetidin-1-yl)-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

cis-2-[7-[(3,5-dichloropyrazol-1-yl)methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile;

trans-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-3-[2-(2-hydroxyethyl)cyclopropyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

3-chloro-7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-5-one;

cis-5-chloro-1-[[3-[2-methylcyclopropyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]pyrazole-3-carbonitrile;

5-chloro-2-[[3-[(2-methylcyclopropyl)-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-7-yl]methyl]pyrazole-3-carbonitrile;

7-[[5-ethoxy-3-(trifluoromethyl)pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide;

N-ethyl-7-[[5-isobutyl-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidine-3-carboxamide; and

trans-2-[2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropyl]acetoneitrile;

or a pharmaceutically acceptable salt thereof.

62. A method of treating a subject suffering from or diagnosed with depression or anxiety mediated by NR2A activity, comprising administering to the subject in need of such treatment an effective amount of 7-(4-fluorophenoxymethyl)-3-[2-(hydroxymethyl)cyclopropyl]-2-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one, or a pharmaceutically acceptable salt thereof.

63. A method of treating a subject suffering from or diagnosed with depression or anxiety mediated by NR2A activity, comprising administering to the subject in need of such treatment an effective amount of 7-((ethyl(4-fluorophenyl)amino)methyl)-3-(2-(hydroxymethyl)cyclopropyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one, or a pharmaceutically acceptable salt thereof.

64. A method of treating a subject suffering from or diagnosed with depression or anxiety mediated by NR2A activity, comprising administering to the subject in need of such treatment an effective amount of 6-[(N-ethyl-4-fluoroanilino)methyl]-2,3-dihydro-1H-cyclopenta[3,4]thiazolo[1,4-a]pyrimidin-8-one, or a pharmaceutically acceptable salt thereof.

65. A method of treating a subject suffering from or diagnosed with depression or anxiety mediated by NR2A activity, comprising administering to the subject in need of such treatment an effective amount of 3-acetyl-7-((ethyl(4-fluorophenyl)amino)methyl)-2-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one, or a pharmaceutically acceptable salt thereof.

66. A method of treating a subject suffering from or diagnosed with depression or anxiety mediated by NR2A activity, comprising administering to the subject in need of such treatment an effective amount of N-ethyl-7-(2-fluoro-3-(trifluoromethyl)benzyl)-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide, or a pharmaceutically acceptable salt thereof.

67. A method of treating a subject suffering from or diagnosed with depression or anxiety mediated by NR2A activity, comprising administering to the subject in need of such treatment an effective amount of 7-[[5-chloro-3-(trifluoromethyl)-1H-pyrazol-1-yl]methyl]-N-ethyl-2-methyl-5-oxo-5H-[1,3]thiazolo[3,2-a]pyrimidine-3-carboxamide, or a pharmaceutically acceptable salt thereof.

68. A method of treating a subject suffering from or diagnosed with depression or anxiety mediated by NR2A activity, comprising administering to the subject in need of such treatment an effective amount of N-ethyl-7-[(N-ethyl-4-fluoroanilino)methyl]-2-methyl-5-oxo-thiazolo[3,2-a]pyrimidine-3-carboxamide, or a pharmaceutically acceptable salt thereof.

69. A method of treating a subject suffering from or diagnosed with depression or anxiety mediated by NR2A activity, comprising administering to the subject in need of such treatment an effective amount of 2-[7-[[5-chloro-3-(trifluoromethyl)pyrazol-1-yl]methyl]-5-oxo-2-(trifluoromethyl)thiazolo[3,2-a]pyrimidin-3-yl]cyclopropanecarbonitrile, or a pharmaceutically acceptable salt thereof.

70. A method of treating a subject suffering from or diagnosed with depression or anxiety mediated by NR2A activity, comprising administering to the subject in need of such treatment an effective amount of 7-(3-cyano-2-fluorobenzyl)-N-ethyl-2-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidine-3-carboxamide, or a pharmaceutically acceptable salt thereof.