



(86) **Date de dépôt PCT/PCT Filing Date:** 2015/02/07
(87) **Date publication PCT/PCT Publication Date:** 2015/08/13
(85) **Entrée phase nationale/National Entry:** 2016/08/02
(86) **N° demande PCT/PCT Application No.:** US 2015/014939
(87) **N° publication PCT/PCT Publication No.:** 2015/120350
(30) **Priorités/Priorities:** 2014/02/07 (US61/937,272);
2014/06/10 (US62/010,004); 2014/08/14 (US62/037,497)

(51) **Cl.Int./Int.Cl.** **A61K 31/519** (2006.01),
A61K 31/357 (2006.01), **A61K 31/382** (2006.01),
A61K 31/404 (2006.01), **A61K 31/436** (2006.01),
A61K 31/4418 (2006.01), **A61K 31/713** (2006.01),
A61K 45/00 (2006.01), **A61P 19/04** (2006.01),
C12Q 1/00 (2006.01), **C12Q 1/68** (2006.01),
G01N 33/48 (2006.01)

(71) **Demandeur/Applicant:**
EFFECTOR THERAPEUTICS, INC., US

(72) **Inventeurs/Inventors:**
APPLEMAN, JAMES, US;
THOMPSON, PEGGY A., US

(74) **Agent:** GOWLING WLG (CANADA) LLP

(54) **Titre : COMPOSITIONS ET METHODES DE TRAITEMENT DE MALADIES FIBREUSES**

(54) **Title: COMPOSITIONS AND METHODS FOR TREATING FIBROTIC DISEASE**




Fig. 7

(57) **Abrégé/Abstract:**

The present disclosure relates to methods for treating or preventing a fibrotic disorder or disease, wherein differentially expressed genes in subjects with said fibrotic disorder or disease are target for therapy using gene or protein modulators.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

13 August 2015 (13.08.2015)



(10) International Publication Number

WO 2015/120350 A3

(51) International Patent Classification:

C12Q 1/68 (2006.01)

(21) International Application Number:

PCT/US2015/014939

(22) International Filing Date:

7 February 2015 (07.02.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/937,272 7 February 2014 (07.02.2014) US

62/010,004 10 June 2014 (10.06.2014) US

62/037,497 14 August 2014 (14.08.2014) US

(71) Applicant: EFFECTOR THERAPEUTICS, INC.

[US/US]; 11180 Roselle Street, San Diego, California 92121 (US).

(72) Inventors: APPLEMAN, James; 4818 Longford Street,

San Diego, California 92117 (US). THOMPSON, Peggy, A.; 1016 Bush Street, San Diego, California 92103 (US).

(74) Agents: PEPE, Jeffrey, C. et al.; Seed Intellectual Prop-

erty Law Group PLLC, Suite 5400, 701 Fifth Avenue, Seattle, Washington 98104-7064 (US).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,

HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

— with international search report (Art. 21(3))

— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

— with sequence listing part of description (Rule 5.2(a))

[Continued on next page]

(54) Title: METHODS FOR TREATING FIBROTIC DISEASE




Fig. 7

(57) Abstract: The present disclosure relates to methods for treating or preventing a fibrotic disorder or disease, wherein differentially expressed genes in subjects with said fibrotic disorder or disease are target for therapy using gene or protein modulators.

WO 2015/120350 A3



(88) Date of publication of the international search report:
1 October 2015

COMPOSITIONS AND METHODS FOR TREATING FIBROTIC DISEASE

STATEMENT REGARDING SEQUENCE LISTING

The Sequence Listing associated with this application is provided in text format in lieu of a paper copy, and is hereby incorporated by reference into the specification.

- 5 The name of the text file containing the Sequence Listing is
300078_403WO_SEQUENCE_LISTING.txt. The text file is 2.3 KB, was created on
February 7, 2015, and is being submitted electronically via EFS-Web.

BACKGROUND

- Progressive scarring (fibrosis) is a pathological feature of many chronic
10 inflammatory diseases, and is an important cause of morbidity and mortality worldwide. Fibrosis is characterized by the accumulation of excess extracellular matrix components (*e.g.*, collagen, fibronectin) that forms fibrous connective tissue in and around an inflamed or damaged tissue. Fibrosis may cause overgrowth, hardening, and/or scarring that disrupts the architecture of the underlying organ or tissue. While controlled tissue
15 remodeling and scarring is part of the normal wound healing process promoted by transdifferentiation of fibroblasts into myofibroblasts, excessive and persistent scarring due to severe or repetitive injury or dysregulated wound healing (*e.g.*, persistence of myofibroblasts) can eventually result in permanent scarring, organ dysfunction and failure, and even death.

- 20 Fibrotic changes can occur in vascular disorders (*e.g.*, peripheral vascular disease, cardiac disease, cerebral disease) and in all main tissue and organ systems (*e.g.*, lung, liver, kidney, heart, skin). Fibrotic disorders include a wide range of clinical presentations, including multisystemic disorders, such as systemic sclerosis, multifocal fibrosclerosis, and organ-specific disorders, such as pulmonary, liver, and kidney
25 fibrosis (Rosenbloom *et al.*, *Ann. Intern. Med.* 152:159, 2010; Wynn, *Nat. Rev. Immunol.* 4:583, 2004). While the etiology and causative mechanisms of individual fibrotic disorders may vary (*e.g.*, ischemic event, exposure to a chemical, radiation, or infectious agent) and are poorly understood, they all share the common feature of

abnormal and excessive deposition of extracellular matrix in affected tissues (Wynn and Ramalingam, *Nat. Med.* 18:1028, 2012).

There are no effective therapies on the market today in the U.S. for treating or preventing fibrotic disorders. Current treatments generally target the inflammatory cascade that contribute to the progression of fibrosis and may temporarily improve symptoms, but are not effective in the long run (Wynn, 2004). Furthermore, the lack of biomarkers for assessing fibrosis progression or regression and therapeutic response has impeded rapid clinical screening of potential therapeutics (Schuppan and Pinzani, *J. Hepatol.* 56:S66, 2012; Castro and Jimenez, *Biomark Med.* 4:133, 2010).

There is clearly a need in the art for new, effective methods of treating or preventing fibrotic disorders and for identifying biomarkers for use in developing therapeutic agents and assessing therapeutic response. The present disclosure meets such needs, and further provides other related advantages.

BRIEF SUMMARY

In one aspect, the present disclosure provides a method for preventing, treating or ameliorating a fibrotic disease, comprising administering to a subject having a fibrotic disorder a therapeutically effective amount of a modulator specific for any one of the genes or encoded products listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5, Table 6, Table 7, an eIF2 component or regulator, an eIF4F complex or regulator, an eIF4F complex component or regulator, an eIF5A or regulator, or any combination thereof.

In other aspects, the present disclosure provides a method for reducing the risk of developing a fibrotic disease, comprising administering to a subject at risk of developing a fibrotic disorder a therapeutically effective amount of a modulator specific for any one of the genes or encoded products listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5, Table 6, Table 7, an eIF2 component or regulator, an eIF4F complex or regulator, an eIF4F complex component or regulator, an eIF5A or regulator, or any combination thereof.

In further aspects, the present disclosure provides a method for reducing myofibroblasts, comprising administering to a subject at risk of developing or having a

fibrotic disorder a therapeutically effective amount of a modulator specific for any one of the genes or encoded products listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5, Table 6, Table 7, an eIF2 component or regulator, an eIF4F complex or regulator, an eIF4F complex component or regulator, an eIF5A or regulator, or any combination thereof.

In still further aspects, the present disclosure provides a method for inhibiting or reversing transdifferentiation of a fibroblast to a myofibroblast, comprising administering to a subject at risk of developing or having a fibrotic disorder a therapeutically effective amount of a modulator specific for any one of the genes or encoded products listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5, Table 6, Table 7, an eIF2 component or regulator, an eIF4F complex or regulator, an eIF4F complex component or regulator, an eIF5A or regulator, or any combination thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the induction of procollagen secreted from fibroblasts by TGF β . Procollagen type 1 levels (Procollagen Type 1C-Peptide, "PIPC") were measured after 24 hours of treating fibroblasts with various concentrations of mTOR inhibitor PP242 and 10ng/mL TGF β . The difference in absorbance at 450 and 540 nm (y-axis) is proportional to the procollagen concentration.

Figure 2 shows a Western blot of protein phosphorylation levels during fibroblast transformation. Western blot analysis of fibroblast transformation as monitored by measuring α -smooth muscle actin (α -SMA) levels after 24 hrs of treating cells with various concentrations of mTOR inhibitor PP242 and 10ng/mL TGF β .

Figure 3 shows the translational and transcriptional profile of fibroblasts treated with TGF β . Comparison of changes in mRNA levels (RNA) and translational rate (RPF) in fibroblasts treated with TGF β . Data points in black have $p \leq 0.05$ for changes in translational efficiency.

Figures 4A to 4C show schematics from Ingenuity Pathway Analysis (IPA) depicting the genes differentially regulated at the mRNA level in the hepatic fibrosis/hepatic stellate cell activation pathway. (A) Early signaling events in hepatic stellate cells. (B) Signaling events in activated hepatic stellate cells. Gene list used and

gene signature identified in analysis is based on p-value from differential concentrations of protein-coding mRNAs from control and TGF β treated fibroblasts. (C) Provides the list of differentially regulated genes from (A) and (B) and shows the magnitude of change (log ratio) and p-value for each gene in their respective pathway (results from 5 biological replicates).

Figures 5A to 5C show schematics from IPA depicting the translational rate differential regulation of genes in the hepatic fibrosis/hepatic stellate cell activation pathway. (A) Early signaling events in hepatic stellate cells. (B) Signaling events in activated hepatic stellate cells. Gene list used and gene signature identified in analysis is based on p-value from differential translation rates from control and TGF β treated fibroblasts. (C) Provides the list of differentially regulated genes from (A) and (B) and shows the magnitude of change (log ratio) and p-value for each gene in their respective pathway. (results from 5 biological replicates).

Figures 6A and 6B show (A) a schematic from IPA depicting the genes regulated in the eIF2 signaling pathway. The gene list used in this analysis is based upon a p-value from differential translation efficiencies from control versus TGF β -treated fibroblasts. (B) Provides the list of differentially regulated genes from (A) and shows the magnitude of change (log ratio) and p-value for each gene in their respective pathway (results from 5 biological replicates).

Figure 7 shows the normalization of translational efficiencies of fibrotic disorder-associated gene signature in the eIF2 signaling pathway. The bar graph shows the translational efficiencies of fibrotic disorder-associated gene signature in fibroblasts treated with TGF β and fibroblasts treated with TGF β and PP242. The normal fibroblast translational efficiency is set at zero, and each of the 12 genes shown had an altered translational efficiency in TGF β -treated fibroblasts (p-value ≤ 0.05), which is a representative result from a single experimental replicate.

Figure 8 shows a bar graph of the translational efficiencies of all 141 fibrotic disorder-associated genes (order as presented in Table 5) showing a differential translational profile in (A) TGF β -treated fibroblasts as compared to untreated (normal) fibroblasts (value set at zero), and (B) how fibroblasts treated with TGF β and mTOR inhibitor PP242 normalizes (*i.e.*, differential closer to zero) many of the 141 fibrotic

disorder-associated genes. The p-value for change in translational efficiency upon TGF β treatment was ≤ 0.05 for this gene signature, which is a representative result from a single experimental replicate.

Figure 9 shows the effect of silvestrol on the induction of procollagen secretion from fibroblasts treated with TGF β . Procollagen type 1 levels (Procollagen Type 1C-Peptide, "PIPC") were measured after 24 hrs. of treating fibroblasts with various concentrations of eIF4A inhibitor silvestrol and 10ng/mL TGF β .

Figure 10 shows the effect of silvestrol on smooth muscle actin (α -SMA) levels in fibroblasts treated with TGF β . The ratio of α -SMA to β -actin (control) was measured after 24 hrs. of treating fibroblasts with various concentrations of eIF4A inhibitor silvestrol and 10ng/mL TGF β .

Figure 11 shows the western blot analysis of α -SMA levels upon fibroblast transformation in the presence of an unrelated siRNA control (siCont) or an siRNA specific for eIF4A1 (siEIF4A1). TGF β -induces transdifferentiation of fibroblasts into myofibroblasts as shown by the large increase in α -SMA levels (see column 2). The levels of negative control, β -actin, are unaffected by the presence of TGF β and/or siRNAs. Knockdown of eIF4A expression with siRNA results in a reduced level of α -SMA production by fibroblasts in the presence of TGF β (compare columns 2 and 4), which essentially prevents fibroblast transdifferentiation into myofibroblasts. The level of eIF4A1 mRNA knockdown was about 80%, as measured by qPCR (data not shown). As shown in the western blot, a corresponding decrease in eIF4A1 protein levels was also observed (see columns 3 and 4).

Figures 12A to 12D show immunofluorescent staining of α -SMA in fibroblasts and TGF β transformed myofibroblasts. TGF β -induced transdifferentiation of fibroblasts into myofibroblasts stimulates an increase in production of α -SMA fibrils (compare A and B). The knockdown of eIF4A prevents TGF β induced transformation of fibroblasts to myofibroblasts and inhibits α -SMA production, which essentially eliminates fibril formation (compare A and C). The presence of either siRNA in the absence of TGF β has no effect on α -SMA production (see B and D).

Figure 13 shows the correlation of translational efficiency measurements between representative experiments in the form of a scatter plot. The alignment of the

dots in a straight line indicate that the genes identified as translationally regulated are highly correlated between experiments, even when the change in translational efficiency for certain genes is not statistically significant in every experiment.

Figure 14 shows a heat map of the differential translational profile for genes modulated (renormalized) by PP242, silvestrol, pirfenidone, CPX, or TSA ($\Delta\log_2$ fold change (fibrotic cells vs. treated fibrotic cells) cut off of ≥ 1). Genes that show the most renormalization appear "white" in the heat map, while genes that are not renormalized appear "black" in the heat map and the genes that are "gray" had an intermediate level of renormalization.

10 DETAILED DESCRIPTION

The instant disclosure provides compositions and methods for identifying agents and validating targets for preventing, ameliorating or treating a fibrotic disorder or disease. For example, translational profiles may be used to (a) identify a candidate therapeutic against an eIF2 pathway protein or regulator (*e.g.*, eIF2 kinase, such as eIF2AK1), an eIF4F complex or component (*e.g.*, eIF4A, eIF4E), an eIF5A protein or hypusination-related protein (*e.g.*, deoxyhypusine synthase, DHPS; deoxyhypusine hydroxylase DOHH), or any combination thereof, for normalizing a translational profile associated with a fibrotic disease, (b) validate a target in or regulator of the eIF2 pathway, an eIF4F complex or component (*e.g.*, eIF4A), or any combination thereof for normalizing a translational profile associated with a fibrotic disease, or (c) identify a subject having or at risk of developing a fibrotic disease as a candidate subject for treating or preventing the fibrotic disease with a therapeutic agent against an eIF2 pathway protein or regulator, an eIF4F complex or component (*e.g.*, eIF4A), or any combination thereof.

By way of background, an injury is an event that damages tissue and initiates the wound healing process. After injury, both mechanical (*i.e.*, extracellular stress caused by disruption of the extracellular matrix, ECM) and chemical signals (*e.g.*, inflammatory mediators like TGF β) activate fibroblastic cells to increase production of extra cellular matrix (ECM) components, which begins the process of fibroblast differentiation into myofibroblasts (Tomasek *et al.*, *Nat. Rev. Mol. Cell Biol.* 3:349,

2002; Werner *et al.*, *Physiol.Rev.* 83:835, 2003). Depending on the type of tissue being remodeled, the fibroblasts that differentiate may come from different sources, including locally present fibroblasts, pericytes, smooth muscle cells, fibrocytes from bone marrow, and from epithelial-mesenchymal transition (EMT) (Hinz *et al.*, *Am. J. Pathol.*

5 170:1807, 2007). In early stages of differentiation, fibroblasts also increase production of focal adhesion proteins and form stress fibers (made up primarily of actin), which is considered a proto-myofibroblast phenotype (Tomasek *et al.*, 2002). Further differentiation to the myofibroblast phenotype occurs when TGF β accumulates, specialized ECM components are present (such as the ED-A variant of fibronectin), and
 10 extracellular stress from ECM and cell remodeling (Tomasek *et al.*, 2002). A hallmark of myofibroblasts is the production of α -smooth muscle actin (α -SMA) (Hinz, *J. Invest. Dermatol.* 127:526, 2007). Wound healing is complete when the newly formed, crosslinked ECM takes over the mechanical load, which is a signal to myofibroblasts to undergo apoptosis (Tomasek *et al.*, 2002; Carlson *et al.*, *J. Surg. Res.* 110:304, 2003).

15 If the injury is severe or repetitive, or if the wound-healing process is dysregulated, fibrosis becomes pathogenic, resulting in permanent scarring or hardening of the tissue, organ malfunction or failure, and ultimately death. For example, idiopathic pulmonary fibrosis (IPF) is a poorly understood progressive and fatal lung disease that has no effective treatment other than lung transplantation (Mason *et al.*,
 20 *Ann. Thorac. Surg.* 84:1121-8, 2007). Median survival of five years after diagnosis is less than 20%. Most forms of interstitial lung diseases and other forms of pulmonary fibrosis are characterized by fibrotic lesions, progressive distortion of alveolar architecture occurs and replacement with fibrotic or scar tissues with excess ECM deposition (American Thoracic Society, *Am. J. Respir. Crit. Care Med.* 161:646, 2000;
 25 Noble *et al.*, *Clin. Chest Med.* 25:749, 2004; Selman *et al.*, *Ann. Intern. Med.* 134:136, 2001). This results in progressive dyspnea and loss of lung function. A hallmark morphological lesion is spatial and temporal heterogeneity incorporating areas of normal lung being directly adjacent to areas of fully established fibrosis, microscopic honeycombing, and areas of evolving fibrosis containing collagen-producing
 30 fibroblasts/myofibroblasts, often referred to as "fibrotic foci." Myofibroblasts are

present in abundance within fibrotic lesions and, therefore, contribute to the excessive scarring found in such lesions of fibrotic disease (Gabbiani, *J. Pathol.* 200:500, 2003).

Prior to setting forth this disclosure in more detail, it may be helpful to an understanding thereof to provide definitions of certain terms to be used herein.

5 Additional definitions are set forth throughout this disclosure.

In the present description, any concentration range, percentage range, ratio range, or integer range is to be understood to include the value of any integer within the recited range and, when appropriate, fractions thereof (such as one tenth and one hundredth of an integer), unless otherwise indicated. Also, any number range recited
10 herein relating to any physical feature, such as polymer subunits, size or thickness, are to be understood to include any integer within the recited range, unless otherwise indicated. As used herein, the term "about" means $\pm 20\%$ of the indicated range, value, or structure, unless otherwise indicated. The term "consisting essentially of" limits the scope of a claim to the specified materials or steps, or to those that do not materially
15 affect the basic and novel characteristics of the claimed invention. It should be understood that the terms "a" and "an" as used herein refer to "one or more" of the enumerated components. The use of the alternative (*e.g.*, "or") should be understood to mean either one, both, or any combination thereof of the alternatives. As used herein, the terms "include," "have" and "comprise" are used synonymously, which terms and
20 variants thereof are intended to be construed as non-limiting.

As used herein, the term "translational profile" refers to the amount of protein that is translated (*i.e.*, translational level) for each gene in a given set of genes in a biological sample, collectively representing a set of individual translational rate values, translational efficiency values, or both translational rate and translational efficiency
25 values for each of one or more genes in a given set of genes. In some embodiments, a translational profile comprises translational levels for a plurality of genes in a biological sample (*e.g.*, cells), *e.g.*, for at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000, 10,000 genes or more, or for at least about 1%, 2%, 3%, 4%,
30 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 50% or more of all genes in the sample. In some embodiments, a translational profile comprises a genome-wide measurement of

translational rate, translational efficiency or both in a biological sample. In certain embodiments, a translational profile refers to a quantitative measure of the amount of mRNA associated with one or more ribosomes for each gene (*i.e.*, translational rate, efficiency or both) in a given set of genes in a biological sample, wherein the amount of ribosome-associated mRNA correlates to the amount of protein that is translated (*i.e.*, translational level).

As used herein, "translation rate" or "rate of translation" or "translational rate" refers to the total count of ribosome engagement, association or occupancy of mRNA for a particular gene as compared to the total count of ribosome engagement, association or occupancy of mRNA for at least one other gene or set of genes, wherein the count of total ribosomal occupancy correlates to the level of protein synthesis. Examination of translation rate across individual genes may be quantitative or qualitative, which will reveal differences in translation. In certain embodiments, translational rate provides a measure of protein synthesis for one or more genes, a plurality of genes, or across an entire genome. In particular embodiments, a translation rate is the amount of mRNA fragments protected by ribosomes for a particular gene relative to the amount of mRNA fragments protected by ribosomes for one or more other genes or groups of genes. For example, the mRNA fragments protected by ribosomes may correspond to a portion of the 5'-untranslated region, a portion of the coding region, a portion of a splice variant coding region, or combinations thereof. In further embodiments, the translation rate is a measure of one, a plurality or all mRNA variants of a particular gene. Translation rates can be established for one or more selected genes or groups of genes within a single composition (*e.g.*, biological sample), between different compositions, or between a composition that has been split into at least two portions and each portion exposed to different conditions.

As used herein, "mRNA level" refers to the amount, abundance, or concentration of mRNA or portions thereof for a particular gene in a composition (*e.g.*, biological sample). In certain embodiments, mRNA level refers to a count of one mRNA, a plurality of mRNA or all mRNA forms or fragments for a particular gene, including pre-mRNA, mature mRNA, or splice variants thereof. In particular embodiments, an mRNA level for one or more genes or groups of genes corresponds to

counts of unique mRNA sequences or portions thereof for a particular gene that map to a 5'-untranslated region, a coding region, a splice variant coding region, or any combination thereof.

As used herein, "translation efficiency" or "translational efficiency" refers to the ratio of the translation rate for a particular gene to the mRNA level for a particular gene in a given set of genes. For example, gene X may produce an equal abundance of mRNA (*i.e.*, same or similar mRNA level) in normal and diseased tissue, but the amount of protein X produced may be greater in diseased tissue as compared to normal tissue. In this situation, the message for gene X is more efficiently translated in diseased tissue than in normal tissue (*i.e.*, an increased translation rate without an increase in mRNA level). In another example, gene Y may produce half the mRNA level in normal tissue as compared to diseased tissue, and the amount of protein Y produced in normal tissue is half the amount of protein Y produced in diseased tissue. In this second situation, the message for gene Y is translated equally efficiently in normal and diseased tissue (*i.e.*, a change in translation rate in diseased tissue that is proportional to the increase in mRNA level and, therefore, the translational efficiency is unchanged). In other words, the expression of gene X is altered at the translational level, while gene Y is altered at the transcriptional level. In certain situations, both the amount of mRNA and protein may change such that mRNA abundance (transcription), translation rate, translation efficiency, or a combination thereof is altered relative to a particular reference or standard.

In certain embodiments, translational efficiency may be standardized by measuring a ratio of ribosome-associated mRNA read density (*i.e.*, translation level) to mRNA abundance read density (*i.e.*, transcription level) for a particular gene (*see, e.g.*, Example 3). As used herein, "read density" is a measure of mRNA abundance and protein synthesis (*e.g.*, ribosome profiling reads) for a particular gene, wherein at least 5, 10, 15, 20, 25, 50, 100, 150, 175, 200, 225, 250, 300 reads or more per unique mRNA or portion thereof is performed in relevant samples to obtain single-gene quantification for one or more treatment conditions. In certain embodiments, translational efficiency is scaled to standardize or normalize the translational efficiency of a median gene to 1.0 after excluding regulated genes (*e.g.*, $\log_2$ fold-change ± 1.5

after normalizing for the all-gene median), which corrects for differences in the absolute number of sequencing reads obtained for different libraries. In further embodiments, changes in protein synthesis, mRNA abundance and translational efficiency are similarly computed as the ratio of read densities between different
 5 samples and normalized to give a median gene a ratio of 1.0, normalized to the mean, normalized to the mean or median of log values, or the like.

As used herein, "gene signature" refers to a plurality of genes that exhibit a generally coherent, systematic, coordinated, unified, collective, congruent, or signature expression pattern or translation efficiency. In certain embodiments, a gene signature is
 10 (a) a plurality of genes that together comprise at least a detectable or identifiable portion of a biological pathway (*e.g.*, 2, 3, 4, 5, or more genes; a fibrotic disease gene signature comprising 11 or 12 genes from the eIF2 translation pathway as illustrated in Figures 6 and 7, or a plurality of genes regulated by the eIF4F complex or component thereof, such as eIF4A), (b) a complete set of genes associated with a biological pathway, or (c)
 15 a cluster or grouping of independent genes having a recognized pattern of expression (*e.g.*, response to a known drug or active compound; related to a disease state such as a fibrotic disorder). One or more genes from a particular gene signature may be part of a different gene signature (*e.g.*, a cell migration pathway may share a gene with a cell adhesion pathway) – that is, gene signatures may intersect or overlap but each signature
 20 can still be independently defined by its unique translation profile.

The term "modulate" or "modulator," as used with reference to altering an activity of a target gene or signaling pathway, refers to increasing (*e.g.*, activating, facilitating, enhancing, agonizing, sensitizing, potentiating, or up regulating) or decreasing (*e.g.*, preventing, blocking, inactivating, delaying activation, desensitizing,
 25 antagonizing, attenuating, or down regulating) the activity of the target gene or signaling pathway. In certain embodiments, a modulator alters a translational profile at the translational level (*i.e.*, increases or decreases translation rate, translation efficiency or both, as described herein), at the transcriptional level, or both.

As used herein, a modulator or agent that "specifically binds" or is "specific for"
 30 a target refers to an association or union of a modulator or agent (*e.g.*, siRNA, chemical compound) to a target molecule (*e.g.*, a nucleic acid molecule encoding a target, a target

product encoded by a nucleic acid molecule, or a target activity), which may be a covalent or non-covalent association, while not significantly associating or uniting with any other molecules or components in a cell, tissue, biological sample, or subject. A modulator or agent specific for a target (*e.g.*, translation machinery component, such as
5 eIF2, eIF4A, eIF4E, eIF5A; translation machinery regulator, such as eIF2AK1, eIF2AK2, eIF2AK3, eIF2AK4, DHPS, DOHH) includes analogs and derivatives thereof. In certain embodiments, a modulator specific for a translation machinery component (*e.g.*, eIF4A, eIF5A) or translation machinery regulator (*e.g.*, eIF2AK1, DOHH) is a siRNA molecule.

10 As used herein, the term "derivative" refers to a modification of a compound by chemical or biological means, with or without an enzyme, which modified compound is structurally similar to a parent compound and (actually or theoretically) derivable from that parent compound. Generally, a "derivative" differs from an "analog" in that a parent compound may be the starting material to generate a "derivative," whereas the
15 parent compound may not necessarily be used as the starting material to generate an "analog." An analog or derivative may have different chemical, biological or physical properties from the parent compound, such as being more hydrophilic or having altered reactivity. Derivatization (*i.e.*, modification) may involve substitution of one or more moieties of a molecule (*e.g.*, a change in functional group). For example, a hydrogen
20 may be substituted with a halogen, such as fluorine or chlorine, or a hydroxyl group (-OH) may be replaced with a carboxylic acid moiety (-COOH). Other exemplary derivatizations include glycosylation, alkylation, acylation, acetylation, ubiquitination, esterification, and amidation.

The term "derivative" also refers to all solvates, for example, hydrates or
25 adducts (*e.g.*, adducts with alcohols), active metabolites, and salts of a parent compound. The type of salt depends on the nature of the moieties within the compound. For example, acidic groups, such as carboxylic acid groups, can form alkali metal salts or alkaline earth metal salts (*e.g.*, sodium salts, potassium salts, magnesium salts, calcium salts, and also salts with physiologically tolerable quaternary ammonium
30 ions and acid addition salts with ammonia and physiologically tolerable organic amines such as, for example, triethylamine, ethanolamine or tris-(2-hydroxyethyl)amine).

Basic groups can form acid addition salts with, for example, inorganic acids such as hydrochloric acid, sulfuric acid or phosphoric acid, or with organic carboxylic acids or sulfonic acids such as acetic acid, citric acid, lactic acid, benzoic acid, maleic acid, fumaric acid, tartaric acid, methanesulfonic acid or p-toluenesulfonic acid. Compounds
5 that simultaneously contain a basic group and an acidic group, for example, a carboxyl group in addition to basic nitrogen atoms, can be present as zwitterions. Salts can be obtained by customary methods known to those skilled in the art, for example, by combining a compound with an inorganic or organic acid or base in a solvent or diluent, or from other salts by cation exchange or anion exchange.

10 In some embodiments, an agent that modulates translation in a fibrotic disease is identified as suitable for use when one or more genes of one or more biological pathways, gene signatures or combinations thereof are differentially translated by at least 1.5-fold (*e.g.*, at least 1.5-fold, at least 2-fold, at least 2.5-fold, at least 3-fold, at least 3.5-fold, at least 4-fold, at least 4.5-fold, at least 5-fold, at least 6-fold, at least 7-
15 fold, at least 8-fold, at least 9-fold, at least 10-fold or more) in a first translational profile (*e.g.*, treated fibrotic disease sample or normal sample) as compared to a second translational profile (*e.g.*, untreated fibrotic disease sample). In some embodiments, an agent that modulates translation in a fibrotic disease is identified as suitable for use when the translational rate, translational efficiency or both for one or more genes of one
20 or more biological pathways, gene signatures or combinations thereof are increased or decreased by at least 1.5-fold (*e.g.*, at least 1.5-fold, at least 2-fold, at least 2.5-fold, at least 3-fold, at least 3.5-fold, at least 4-fold, at least 4.5-fold, at least 5-fold, at least 6-fold, at least 7-fold, at least 8-fold, at least 9-fold, at least 10-fold or more) in a first translational profile as compared to a second translational profile.

25 A "biological sample" includes blood and blood fractions or products (*e.g.*, serum, plasma, platelets, red blood cells, or the like); sputum or saliva; kidney, lung, liver, heart, brain, nervous tissue, thyroid, eye, skeletal muscle, cartilage, or bone tissue; cultured cells, *e.g.*, primary cultures, explants, and transformed cells, stem cells, stool, urine, *etc.* Such biological samples (*e.g.*, disease samples or normal samples) also
30 include sections of tissues, such as a biopsy or autopsy sample, frozen sections taken for histologic purposes, or cells or other biological material used to model disease or to be

representative of a pathogenic state (*e.g.*, TGF β treated fibroblasts as a model system for fibrosis). In certain embodiments, a biological sample is obtained from a "subject," *e.g.*, a eukaryotic organism, most preferably a mammal such as a primate, *e.g.*, chimpanzee or human; cow; dog; cat; rodent, *e.g.*, guinea pig, rat, or mouse; rabbit;
 5 bird; reptile; or fish.

As used herein, the term "normalize" or "normalizing" or "normalization" refers to adjusting the translational rate, translational efficiency, or both of one or more genes in a biological sample from a subject (*e.g.*, a disease sample from one or more subjects, tissues or organs) to a level that is more similar, closer to, or comparable to the
 10 translational rate, translational efficiency, or both of those same one or more genes in a control sample (*e.g.*, a non-diseased or normal sample from the same or different subject, tissue or organ). In certain embodiments, normalization refers to modulation of one or more translational regulators or translational system components to adjust or shift the translational rate, efficiency or both of one or more genes in a biological
 15 sample (*e.g.*, diseased, abnormal or other biologically altered condition) to a translational efficiency that is more similar, closer to or comparable to the translational efficiency of those one or more genes in a non-diseased or normal control sample. In some embodiments, normalization is evaluated by determining a translational rate, translational efficiency or both of one or more genes in a biological sample (*e.g.*,
 20 disease sample) from a subject before and after an agent (*e.g.*, therapeutic or known active agent) is administered to the subject and comparing the translational rate, translational efficiency or both before and after administration to the translational rate, translational efficiency or both from a control sample in the absence or presence of the agent. Exemplary methods of evaluating normalization of a translational profile
 25 associated with a disease or disorder includes observing a shift in a gene signature or evaluating a translational profile shift due to a therapeutic intervention in a fibrotic or fibrotic-associated condition, disease or disorder.

As used herein, the phrase "differentially translated" refers to a change or difference (*e.g.*, increase, decrease or a combination thereof) in translation rate,
 30 translation efficiency, or both of one gene, a plurality of genes, a set of genes of interest, one or more gene clusters, or one or more gene signatures under a particular

condition as compared to the translation rate, translation efficiency, or both of the same gene, plurality of genes, set of genes of interest, gene clusters, or gene signatures under a different condition, which is observed as a difference in expression pattern. For example, a translational profile of a diseased cell may reveal that one or more genes
5 have higher translation rates, higher translation efficiencies, or both (*e.g.*, higher ribosome engagement of mRNA or higher protein abundance) than observed in a control or normal cell. Another exemplary translational profile of a diseased cell may reveal that one or more genes have lower translation rates, lower translation
10 efficiencies, or both (*e.g.*, lower ribosome engagement of mRNA or lower protein abundance) than observed in a control or normal cell. In still another example, a translational profile of a diseased cell may reveal that one or more genes have higher translation rates, one or more genes have higher translation efficiencies, one or more genes have lower translation rates, one or more genes have lower translation
15 efficiencies, or any combination thereof than observed in a control or normal cell. In some embodiments, one or more gene signatures, gene clusters or sets of genes of interest are differentially translated in a first translational profile as compared to one or more other translational profiles. In further embodiments, one or more genes, gene signatures, gene clusters or sets of genes of interest in a first translational profile show at least a 1.5-fold translation differential or at least a 1.0 log₂ change (*i.e.*, increase or
20 decrease) as compared to the same one or more genes in at least one other different (*e.g.*, second, third, etc.) translational profile.

In some embodiments, two or more translational profiles are generated and compared to each other to determine the differences (*i.e.*, increases and/or decreases in translational rate, translational efficiency, or both) for each gene in a given set of genes
25 between the two or more translational profiles. The comparison between the two or more translational profiles is referred to as the "differential translational profile." In certain embodiments, a differential translational profile comprises one or more genes, gene clusters, or gene signatures (*e.g.*, a fibrotic disease-associated pathway), or combinations thereof.

30 In certain embodiments, differential translation between genes or translational profiles may involve or result in a biological (*e.g.*, phenotypic, physiological, clinical,

therapeutic, prophylactic) benefit. For example, when identifying a therapeutic, validating a target, or treating a subject having a fibrotic disorder or disease, a "biological benefit" means that the effect on translation rate, translation efficiency or both, or the effect on the translation rate, translation efficiency or both of one or more genes of a translational profile allows for intervention or management of the fibrotic disorder or disease of a subject (*e.g.*, a human or non-human mammal, such as a primate, horse, dog, mouse, rat). In general, one or more differential translations or differential translation profiles indicate that a "biological benefit" will be in the form, for example, of an improved clinical outcome; lessening or alleviation of symptoms associated with fibrotic disease; decreased occurrence of symptoms; improved quality of life; longer disease-free status; diminishment of extent of fibrotic disease; stabilization of a fibrotic disease; delay of fibrotic disease progression; remission; survival; or prolonging survival. In certain embodiments, a biological benefit comprises normalization of a differential translation profile, or comprises a shift in translational profile to one closer to or comparable to a translational profile induced by a known active compound or therapeutic, or comprises inducing, stimulating or promoting a desired phenotype or outcome (*e.g.*, reversal of transformation, induction of a quiescent state, apoptosis, necrosis, cytotoxicity), or reducing, inhibiting or preventing an undesired phenotype or outcome (*e.g.*, activation, transformation, proliferation, migration).

In some embodiments, a biological benefit comprises reducing the amount of myofibroblast cells associated with a fibrotic disorder or disease by reducing, blocking or reversing transdifferentiation of fibroblast cells into myofibroblast cells, or by promoting apoptosis, senescence or quiescence of myofibroblast cells. In further embodiments, a biological benefit comprises inhibiting or blocking the production, secretion or both of organ damaging (*i.e.*, toxic in a fibrotic disorder) proteins produced by myofibroblasts (*e.g.*, ECM related proteins) in a subject having a fibrotic disorder or disease. As used herein, the phrase "organ damaging proteins" refers to one or more proteins associated with myofibroblasts in the context of a fibrotic disorder under these circumstances can be toxic, whereas such proteins under normal conditions would not be organ damaging.

In some embodiments, less than about 20% of the genes in the genome are differentially translated by at least 1.5-fold in a first translational profile as compared to a second translational profile. In some embodiments, less than about 5% of the genes in the genome are differentially translated by at least 2-fold or at least 3-fold in a first
 5 translational profile as compared to a second translational profile. In some embodiments, less than about 1% of the genes in the genome are differentially translated by at least 4-fold or at least 5-fold in a first translational profile as compared to a second translational profile.

As described herein, differentially translated genes between first and second
 10 translational profiles under a first condition may exhibit translational profiles "closer to" each other (*i.e.*, identified through a series of pair-wise comparisons to confirm a similarity of pattern) under one or more different conditions (*e.g.*, differentially translated genes between a normal sample and a fibrotic disease sample may have a more similar translational profile when the normal sample is compared to a fibrotic
 15 disease sample contacted with a candidate agent; differentially translated genes between a fibrotic disease sample and a fibrotic disease sample treated with a known active agent may have a more similar translational profile when the disease sample treated with a known active agent is compared to the disease sample contacted with a candidate agent). In certain embodiments, a test translational profile is "closer to" a reference
 20 translational profile when at least 99%, 95%, 90%, 80%, 70%, 60%, 50%, 25%, or 10% of a selected portion of differentially translated genes, a majority of differentially translated genes, or all differentially translated genes show a translational profile within 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, or 25%, respectively, of their corresponding genes in the reference translational profile. In further embodiments, a
 25 selected portion of differentially translated genes, a majority of differentially translated genes, or all differentially translated genes from an experimental translational profile have a translational profile "closer to" the translational profile of the same genes in a reference translational profile when the amount of protein translated in the experimental and reference translational profiles are within about $3.0 \log_2$, $2.5 \log_2$, $2.0 \log_2$, $1.5 \log_2$,
 30 $1.1 \log_2$, $0.5 \log_2$, $0.2 \log_2$ or closer. In still further embodiments, a selected portion of differentially translated genes, a majority of differentially translated genes, or all

differentially translated genes from an experimental translational profile have a translational profile "closer to" the translational profile of the same genes in a reference translational profile when the amount of protein translated in the experimental and reference translational profiles differs by no more than about 50%, 45%, 40%, 35%,
 5 30%, 25%, 20%, 15%, 10%, 5%, 1% or less.

In some embodiments, an experimental differential profile as compared to a reference differential translational profile of interest has at least a 1.0 $\log_2$ change in translational rate, translational efficiency, or both for at least 0.05%, at least 0.1%, at least 0.25%, at least 0.5%, at least 1%, at least 2%, at least 3%, at least 4%, at least 5%,
 10 at least 6%, at least 7%, at least 8%, at least 9%, at least 10%, at least 15%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90% or more of a set of selected differentially translated genes or for the entire set of selected differentially translated genes. In some embodiments, an experimental differential profile as compared to a reference differential translational
 15 profile of interest has at least a 2 $\log_2$ change in translational rate, translational efficiency, or both for at least 0.05%, at least 0.1%, at least 0.25%, at least 0.5%, at least 1%, at least 5%, at least 10%, at least 15%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90% or more of a set of selected differentially translated genes or for the entire set of differentially
 20 translated genes. In some embodiments, an experimental differential profile as compared to a reference differential translational profile of interest has at least a 3 $\log_2$ change in translational rate, translational efficiency, or both for at least 0.05%, at least 0.1%, at least 0.25%, at least 0.5%, at least 1%, at least 5%, at least 10%, at least 15%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least
 25 80%, or at least 90% or more of a set of selected differentially translated genes or for the entire set of selected differentially translated genes. In some embodiments, an experimental differential profile as compared to a reference differential translational profile of interest has at least a 4 $\log_2$ change in translational levels for at least 0.05%, at least 0.1%, at least 0.25%, at least 0.5%, at least 1%, at least 5%, at least 10%, at least
 30 15%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%,

at least 80%, or at least 90% or more of a set of selected differentially translated genes or for the entire set of selected differentially translated genes.

As described herein, a differential translational profile between a first sample and a control may be "comparable" to a differential translational profile between a second sample and the control (*e.g.*, the differential profile between a fibrotic disease sample and the fibrotic disease sample treated with a known active compound may be comparable to the differential profile between the fibrotic disease sample and the fibrotic disease sample contacted with a candidate agent; the differential profile between a fibrotic disease sample and a non-diseased (normal) sample may be comparable to the differential profile between the fibrotic disease sample and the fibrotic disease sample contacted with a candidate agent). In certain embodiments, a test differential translational profile is "comparable to" a reference differential translational profile when at least 99%, 95%, 90%, 80%, 70%, 60%, 50%, 25%, or 10% of a selected portion of differentially translated genes, a majority of differentially translated genes, or all differentially translated genes show a translational profile within 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, or 25%, respectively, of their corresponding genes in the reference translational profile. In further embodiments, a differential translational profile comprising a selected portion of the differentially translated genes or all the differentially translated genes has a differential translational profile "comparable to" the differential translational profile of the same genes in a reference differential translational profile when the amount of protein translated in the experimental and reference differential translational profiles are within about 3.0 log₂, 2.5 log₂, 2.0 log₂, 1.5 log₂, 1.0 log₂, 0.5 log₂, 0.2 log₂ or closer. In still further embodiments, a differential translational profile comprising a selected portion of the differentially translated genes or all the differentially translated genes has a differential translational profile "comparable to" the differential translational profile of the same genes in a reference differential translational profile when the amount of protein translated in the experimental and reference differential translational profiles differs by no more than about 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, 5%, 1% or less.

As used herein, "fibroblast" refers to a mesenchymal-derived connective tissue cell that secretes extracellular matrix components, such as collagen, and other

macromolecules. Fibroblasts have a spindle-shaped morphology and can play a role in organismal development and wound healing. For example, fibroblasts may be identified by global DNase I hypersensitive site mapping which establishes a lineage association (Stamatoyannopoulos *et al.*, *Nat. Genet.* 43:264, 2011), or by the production
5 of certain biomarkers (*e.g.*, Discoidin Domain Receptor 2, DDR2 and vimentin). Fibroblasts are different from endothelial cells and hematopoietic cells.

The term "fibrotic disorder" or "fibrotic disease" refers to a medical condition featuring progressive and/or irreversible fibrosis, wherein excessive deposition of extracellular matrix occurs in and around inflamed or damaged tissue. In certain
10 embodiments, a fibrotic disorder or disease is associated with the persistent presence of myofibroblasts in and around fibrotic foci or lesions. Excessive and persistent fibrosis can progressively remodel and destroy normal tissue, which may lead to dysfunction and failure of affected organs, and ultimately death. A fibrotic disorder may affect any tissue in the body and is generally initiated by an injury and the transdifferentiation of
15 fibroblasts into myofibroblasts. As used herein, "transdifferentiation" refers to the direct conversion of one cell type into another. It is to be understood that fibrosis alone triggered by normal wound healing processes that has not progressed to a pathogenic state is not considered a fibrotic disorder or disease of this disclosure. A "fibrotic lesion" or "fibrotic plaque" refers to a focal area of fibrosis.

20 As used herein, "injury" refers to an event that damages tissue and initiates fibrosis. An injury may be caused by an external factor, such as mechanical insult (*e.g.*, cut, surgery), exposure to radiation, chemicals (*e.g.*, chemotherapy, toxins, irritants, smoke), or infectious agent (*e.g.*, bacteria, virus, or parasite). An injury may be caused by, for example, chronic autoimmune inflammation, allergic response, HLA
25 mismatching (*e.g.*, transplant recipients), or ischemia (*i.e.*, an "ischemic event" or "ischemia" refers to an injury that restricts in blood supply to a tissue, resulting in damage to or dysfunction of tissue, which may be caused by problems with blood vessels, atherosclerosis, thrombosis or embolism, and may affect a variety of tissues and organs; an ischemic event may include, for example, a myocardial infarction,
30 stroke, organ or tissue transplant, or renal artery stenosis). In certain embodiments, an injury leading to a fibrotic disorder may be of unknown etiology (*i.e.*, idiopathic).

Non-limiting examples of fibrotic disorders or fibrotic diseases include pulmonary fibrosis, idiopathic pulmonary fibrosis, cystic fibrosis, liver fibrosis (*e.g.*, cirrhosis), cardiac fibrosis, endomyocardial fibrosis, vascular fibrosis (*e.g.*, atherosclerosis, stenosis, restenosis), atrial fibrosis, mediastinal fibrosis, myelofibrosis, retroperitoneal fibrosis, progressive massive fibrosis (*e.g.*, lungs), chronic kidney disease, nephrogenic systemic fibrosis, Crohn's disease, hypertrophic scarring, keloid, scleroderma, systemic sclerosis (*e.g.*, skin, lungs), athrofibrosis (*e.g.*, knee, shoulder, other joints), Peyronie's disease, Dupuytren's contracture, adhesive capsulitis, organ transplant associated fibrosis, ischemia associated fibrosis, or the like.

Reference to "pulmonary fibrotic disorder" means diseases or disorders characterized by fibrotic hypertrophy or fibrosis of lung tissue. Exemplary pulmonary fibrotic disorders include pulmonary fibrosis, idiopathic pulmonary fibrosis, interstitial lung disease, interstitial pulmonary fibrosis, chronic interstitial pneumonitis, Hamman-Rich Syndrome, usual interstitial pneumonitis (UIP), fibrosing alveolitis, pulmonary sarcoidosis, progressive massive fibrosis (*e.g.*, lungs), systemic sclerosis (*e.g.*, lungs), lung transplant associated fibrosis, or the like.

"Treatment," "treating" or "ameliorating" refers to medical management of a disease, disorder, or condition of a subject (*i.e.*, patient), which may be therapeutic, prophylactic/preventative, or a combination treatment thereof. A treatment may improve or decrease the severity at least one symptom of fibrotic disease, delay worsening or progression of a disease, delay or prevent onset of additional associated diseases, or improve remodeling of fibrotic lesions into functional (partially or fully) tissue. "Reducing the risk of developing a fibrotic disorder" refers to preventing or delaying onset of a fibrotic disorder or reoccurrence of one or more symptoms of the fibrotic disorder.

A "therapeutically effective amount (or dose)" or "effective amount (or dose)" of a compound refers to that amount sufficient to result in amelioration of one or more symptoms of the disease being treated in a statistically significant manner. When referring to an individual active ingredient administered alone, a therapeutically effective dose refers to that ingredient alone. When referring to a combination, a

therapeutically effective dose refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered serially or simultaneously.

The term "pharmaceutically acceptable" refers to molecular entities and compositions that do not produce allergic or other serious adverse reactions when
5 administered to a subject using routes well-known in the art.

A "subject in need" refers to a subject at risk of, or suffering from, a disease, disorder or condition (*e.g.*, fibrosis) that is amenable to treatment or amelioration with a compound or a composition thereof provided herein. In certain embodiments, a subject in need is a human.

10 The "percent identity" between two or more nucleic acid sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % identity=number of identical positions/total number of positions x 100), taking into account the number of gaps, and the length of each gap that needs to be introduced to optimize alignment of two or more sequences. The comparison of sequences and determination of percent
15 identity between two or more sequences can be accomplished using a mathematical algorithm, such as BLAST and Gapped BLAST programs at their default parameters (*e.g.*, Altschul *et al.*, *J. Mol. Biol.* 215:403, 1990; *see also* BLASTN at www.ncbi.nlm.nih.gov/BLAST).

A "conservative substitution" is recognized in the art as a substitution of one
20 amino acid for another amino acid that has similar properties. Exemplary conservative substitutions are well known in the art (*see, e.g.*, WO 97/09433, p. 10; Lehninger, *Biochemistry*, 2nd Edition; Worth Publishers, Inc. NY:NY (1975), pp.71-77; Lewin, *Genes IV*, Oxford University Press, NY and Cell Press, Cambridge, MA (1990), p. 8).

Fibrotic Disorder or Disease

25 In one aspect, the present disclosure provides a method for preventing, treating or ameliorating a fibrotic disorder, comprising administering to a subject having a fibrotic disorder a therapeutically effective amount of a modulator specific for any one or more of the genes (including any alleles, homologs, or orthologs) or any encoded products (including any active fragments or splice variants thereof) listed in Figure 7,
30 Table 1, Table 3A, Table 3B, Table 5, Table 6, Table 7, an eIF2 component or regulator, an eIF4F complex, an eIF4F complex component (such as eIF4A or eIF4E)

or regulator, or an eIF5A or regulator (such as DHPS or DOHH). In certain embodiments, the present disclosure provides a method for reducing the risk of developing a fibrotic disorder, comprising: administering to a subject at risk of developing a fibrotic disorder a therapeutically effective amount of a modulator specific
 5 for any one or more of the genes (including any alleles, homologs, or orthologs) or any encoded products (including any active fragments or splice variants thereof) listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5, Table 6, Table 7, an eIF2 component or regulator (such as eIF2AK1, eIF2AK2, eIFAK3 or eIFAK4), an eIF4F complex, an eIF4F complex component (such as eIF4A or eIF4E) or regulator, an eIF5A or regulator
 10 (such as DHPS or DOHH).

In other aspects, the present disclosure provides a method for reducing myofibroblasts, comprising administering to a subject at risk of developing or having a fibrotic disorder a therapeutically effective amount of a modulator specific for any one of the genes (including any alleles, homologs, or orthologs) or any encoded products
 15 (including any active fragments or splice variants thereof) listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5, Table 6, Table 7, an eIF2 component or regulator, an eIF4F complex or regulator, an eIF4F complex component or regulator, an eIF5A or regulator, or any combination thereof.

In still other aspects, the present disclosure provides a method for inhibiting or
 20 reversing fibroblast transdifferentiation into a myofibroblast, comprising administering to a subject at risk of developing or having a fibrotic disorder a therapeutically effective amount of a modulator specific for any one of the genes (including any alleles, homologs, or orthologs) or any encoded products (including any active fragments or splice variants thereof) listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5, Table
 25 6, Table 7, an eIF2 component or regulator, an eIF4F complex or regulator, an eIF4F complex component or regulator, an eIF5A or regulator, or any combination thereof.

In another aspect, the instant disclosure provides a method for identifying a candidate therapeutic for normalizing a translational profile associated with a fibrotic disease, comprising (a) determining three independent translational profiles, each for a
 30 plurality of genes, wherein (i) a first translational profile is from a fibrotic disease sample, (ii) a second translational profile is from (1) a control non-diseased sample or

(2) a control non-diseased sample contacted with a candidate agent, and (iii) a third translational profile is from the fibrotic disease sample contacted with a candidate agent; (b) determining a first differential translational profile comprising one or more genes differentially translated in the first translational profile as compared to the second translational profile, and determining a second differential translational profile comprising one or more genes differentially translated in the first translational profile as compared to the third translational profile, wherein the one or more differentially translated genes are selected from EIF2AK1, EIF2AK2, EIF2AK3, EIF2AK4, EIF5A, mTOR, DOHH, DHPS, HDAC6, SIRT2, RSK, AHCY, or the genes listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5, Table 6 or Table 7; and (c) identifying the agent as a candidate therapeutic for normalizing a translational profile associated with the fibrotic disorder when the first differential translational profile is comparable to the second differential translational profile.

In still another aspect, the instant disclosure provides a method for validating a target for normalizing a translational profile associated with a fibrotic disease, the method comprising (a) determining three independent translational profiles, each for a plurality of genes, wherein (i) a first translational profile is from a fibrotic disease sample, (ii) a second translational profile is from (1) a control non-diseased sample or (2) a control non-diseased sample contacted with an agent that modulates a target (*e.g.*, an eIF2 component; an eIF2 regulator such as EIF2AK1, EIF2AK2, EIF2AK3 or EIF2AK4; an eIF4F complex component such as eIF4A or eIF4E; eIF5A or a regulator of eIF5A such as DHPS or DOHH), and (iii) a third translational profile is from the fibrotic disease sample contacted with the agent that modulates the target; (b) determining a first differential translational profile comprising one or more genes differentially translated in the first translational profile as compared to the second translational profile, and determining a second differential translational profile comprising one or more genes differentially translated in the first translational profile as compared to the third translational profile, wherein the one or more differentially translated genes are selected from, for example, EIF2AK1, EIF2AK2, EIF2AK3, EIF2AK4, EIF5A, mTOR, DOHH, DHPS, HDAC6, SIRT2, RSK, AHCY, or the genes listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5 or Table 7; and (c) validating

the target as a target for normalizing a translational profile associated with the fibrotic disease when the first differential translational profile is comparable to the second differential translational profile.

In certain aspects, a target comprises a translation machinery element, a
 5 regulator of a translation machinery component, or combinations thereof. In certain embodiments, a target comprises an eIF2 component (*e.g.*, eIF2 α , eIF2 β , eIF2 γ), an eIF4F complex, an eIF4F complex component (*e.g.*, eIF4E, eIF4A, eIF4G), or any combination thereof. In particular embodiments, a target comprises eIF2 α , eIF2 β , eIF2 γ , eIF4A, eIF4E, eIF5A, rpS6, or any combination thereof. In further
 10 embodiments, a target comprises an eIF2 α kinase (*e.g.*, EIF2AK1 or heme-regulated inhibitor kinase (HRI), EIF2AK2 or double-stranded RNA-dependent kinase (protein kinase R, PKR), EIF2AK3 or PKR-like endoplasmic reticulum kinase (PERK), EIF2AK4 or general control nonderepressible 2 (GCN2)), mTOR, deoxyhypusine hydroxylase (DOHH), deoxyhypusine synthase (DHPS), histone deacetylase 6
 15 (HDAC6), NAD-dependent deacetylase sirtuin-2 (SIRT2), p90 Ribosomal S6 kinase (RSK), adenosylhomocysteinase (AHCY), or any combination thereof.

In certain embodiments, a therapeutically effective amount of a modulator specifically targeting eIF2AK1 activity, eIF2AK2 activity, eIF2AK3 activity, eIF2AK4 activity, eIF4A activity, eIF5A activity, or DOHH activity is administered to a subject
 20 at risk of developing or having a fibrotic disorder in order to reduce the amount of myofibroblasts contributing to the fibrotic disorder. The reduction may be due to promoting apoptosis of the existing myofibroblasts, reducing or inhibiting transdifferentiation of fibroblasts into myofibroblasts, promoting transdifferentiation of myofibroblasts into another cell type, inducing senescence or quiescence of
 25 myofibroblasts, or any combination thereof. In other embodiments, a therapeutically effective amount of a modulator specifically targeting eIF2AK1 activity, eIF2AK2 activity, eIF2AK3 activity, eIF2AK4 activity, eIF4A activity, eIF5A activity, or DOHH activity is administered to a subject having a fibrotic disorder in order to inhibit or block the production, secretion or both of organ damaging proteins from myofibroblasts
 30 (*e.g.*, ECM related proteins). In further embodiments, a modulator specifically targets a nucleic acid encoding the eIF2AK1 activity, eIF2AK2 activity, eIF2AK3 activity,

eIF2AK4 activity, eIF4A activity, eIF5A activity, or DOHH activity. For example, a specific modulator may be an siRNA or a derivative thereof (*e.g.*, nuclease resistant modifications, such as phosphorothioate, locked nucleic acids (LNA), 2'-O-methyl modifications, morpholino linkages, or the like).

5 In particular embodiments, a modulator specifically targeting DOHH activity does not modulate or minimally modulates other protein hydroxylases, such as lysyl-hydroxylase, prolyl-hydroxylase and aspartyl/asparaginyl hydroxylase. By way of background, during the formation and maintenance of fibrocellular scar tissue, certain proteins like collagen and the chaperone LTBP are hydroxylated (*e.g.*, prolyl-4-
10 hydroxylase modifies proline to 4-hydroxyproline (Hyp) on collagen and the presence of Hyp is required for collagen structural stability). A surprising result of this disclosure is that specific inhibition of DOHH alone (without affecting other hydroxylases) will inhibit or reverse transdifferentiation of fibroblasts into myofibroblasts, inhibit the production and/or secretion of organ damaging proteins
15 produced by myofibroblasts – therefore, specific inhibition of DOHH can be used to treat or prevent fibrotic disorders.

 In further particular embodiments, a modulator specifically targeting eIF2AK1 activity will inhibit or reverse transdifferentiation of fibroblasts into myofibroblasts and, therefore, treat or prevent fibrotic disorders. In certain embodiments, a modulator
20 specifically targeting eIF2AK1 activity is used to treat or ameliorate fibrotic disorders by inhibiting or blocking the production, secretion or both of organ damaging proteins from myofibroblasts.

 In further particular embodiments, a modulator specifically targeting eIF2AK2 activity will inhibit or reverse transdifferentiation of fibroblasts into myofibroblasts and, therefore, treat or prevent fibrotic disorders. In certain embodiments, a modulator
25 specifically targeting eIF2AK2 activity is used to treat or ameliorate fibrotic disorders by inhibiting or blocking the production, secretion or both of organ damaging proteins from myofibroblasts.

 In further particular embodiments, a modulator specifically targeting eIF2AK3
30 activity will inhibit or reverse transdifferentiation of fibroblasts into myofibroblasts and, therefore, treat or prevent fibrotic disorders. In certain embodiments, a modulator

specifically targeting eIF2AK3 activity is used to treat or ameliorate fibrotic disorders by inhibiting or blocking the production, secretion or both of organ damaging proteins from myofibroblasts.

In further particular embodiments, a modulator specifically targeting eIF2AK4
5 activity will inhibit or reverse transdifferentiation of fibroblasts into myofibroblasts and, therefore, treat or prevent fibrotic disorders. In certain embodiments, a modulator specifically targeting eIF2AK4 activity is used to treat or ameliorate fibrotic disorders by inhibiting or blocking the production, secretion or both of organ damaging proteins from myofibroblasts.

10 In still further particular embodiments, a modulator specifically targeting eIF4A activity will inhibit or reverse transdifferentiation of fibroblasts into myofibroblasts and, therefore, treat or prevent fibrotic disorders. In certain embodiments, a modulator specifically targeting eIF4A activity is used to treat or ameliorate fibrotic disorders by inhibiting or blocking the production, secretion or both of organ damaging proteins
15 from myofibroblasts.

In yet further particular embodiments, a modulator specifically targeting eIF5A activity will inhibit or reverse transdifferentiation of fibroblasts into myofibroblasts and, therefore, treat or prevent fibrotic disorders. In certain embodiments, a modulator specifically targeting eIF5A activity is used to treat or ameliorate fibrotic disorders by
20 inhibiting or blocking the production, secretion or both of organ damaging proteins from myofibroblasts.

In certain embodiments, a target comprises CREB5, DIAPH3, LGALS1, NACA, RPL12, RPL13A, RPL17, RPL21, RPL22L1, RPL23, RPL26, RPL27A, RPL28, RPL3, RPL30, RPL34, RPL36, RPL37, RPL37A, RPL4, RPL7A, RPS9,
25 RPLP1, RPLP2, RPS10, RPS16, RPS19, RPS27, RPS5, RPS8, RPS9, SLC25A6, SOX6, STS, TKT or any combination thereof. In further embodiments, a target comprises ABCA6, ANKH, CARD16, CEP192, DDX60, DNASE1L1, DYNC2H1, EDN1, HBEGF, HOMER1, INHBA, KDM6B, LENG9, MATN3, MYO19, NRG1, PABPC4, PLD1, PLEKHA5, RASD1, SGIP1, SLC2A12, SNRPA, TEN1, TOP2A,
30 TRERF1 or any combination thereof. In still further embodiments, a target comprises C9orf85, EIF3E, GAPDH, HNRNPA1, MRPL45 or any combination thereof. In yet

further embodiments, a target comprises CES1, LAMP5, PAQR5, PLEKHG1, ROBO2, TOMM7 or any combination thereof. In more embodiments, a target comprises AOX1, ARPC1A, AURKA, C12orf57, GPSM2, KITLG, MAP3K5, MURC, NOV, RPL14, SLC15A3, SOX5, ZNF608 or any combination thereof.

5 In certain embodiments, a target comprises ANKDD1A, ATP5G2, CHCHD10, DNAJC22, FGF5, FMO2, GNB2L1, GLTSCR2, HIGD2A, IFIH1, MTUS1, RPS18, RPL18A, RPL31, RPL35A, RPL5, RPS18, RPS29, SPATA6 or any combination thereof. In further embodiments, a target comprises RPS6KA5, BIVM, ACTA1, KRT7, AMIGO3, CCDC102B, RPL10, TAF1D, ADAMTS5, LAMB3, CLCF1,
10 EPB41L1, GAS2L3, IRAK3, LPAR3, PCBP2, PDE7B, TMTC1, FRMD4A, GDF10, OBSCN, PLEKHA6, SHC3 or any combination thereof. In still further embodiments, a target comprises THBS3, RPS28, EEF1A1, EEF2, EIF4B, FMO2, RAB3D, CIT, PDE4B, PPARG, SLC40A1, ASPM, CA5B, GLCCI1, GLTSCR2, P2RX7, STAMBPL1 or any combination thereof.

15 In some aspects, the instant disclosure provides a method of identifying a subject as a candidate for preventing, treating or ameliorating a fibrotic disease with a therapeutic agent, the method comprising (a) determining a first translational profile for a plurality of genes in a sample from a subject having or suspected of having a fibrotic disease; (b) determining a second translational profile for a plurality of genes in a
20 control sample, wherein the control sample is from a subject known to respond to the therapeutic agent and wherein the sample has not been contacted with the therapeutic agent; and (c) identifying the subject as a candidate for treating fibrotic disease with the therapeutic agent when the translational profile for one or more genes selected from, for example, EIF2AK1, EIF2AK2, EIF2AK3, EIF2AK4, EIF5A, mTOR, DOHH, DHPS,
25 HDAC6, SIRT2, RSK, AHCY, or Figure 7, Table 1, Table 3A, Table 3B, Table 5 or Table 7 of the first translational profile are comparable to the translational profile of the corresponding genes in the second translational profile. In a related aspect, the instant disclosure provides a method for preventing, treating or ameliorating a fibrotic disease, comprising administering a therapeutic agent to a subject identified according to the
30 method of identifying a subject as a candidate for preventing, treating or ameliorating a fibrotic disease, thereby treating the subject. In certain embodiments, a method of

identifying a subject as a candidate for preventing, treating or ameliorating a fibrotic disease with a therapeutic agent comprises using any one or more of the combinations of modulators described herein.

In any of the aforementioned embodiments, the fibrotic disease or disorder may
5 be due to injury or may be idiopathic. In some embodiments, the injury is an ischemic event or due to exposure to radiation, a chemical, or an infectious agent. In any of these embodiments, a specific modulator for an eIF2 pathway protein or regulator (such as EIF2AK1, EIF2AK2, EIF2AK3 or EIF2AK4), a specific modulator of an eIF4F
10 complex or component thereof (such as eIF4A or eIF4E), a specific modulator of an eIF5A or a regulator of eIF5A (such as DHPS or DOHH), or any combination thereof is administered before or after a fibrotic lesion has developed in the subject. In further embodiments, a modulator specific for a target of interest is formulated with a pharmaceutically acceptable diluent, carrier or excipient.

In certain embodiments, the present disclosure provides methods for treating
15 fibrotic disorders or for inhibiting transdifferentiation of fibroblasts into myofibroblasts by administering a modulator of a first target combined with one or more modulators of one or more different targets (*e.g.*, two, three, four, five, or six targets). In some embodiments, the first target is a first translation machinery element and the second target is a second a translation machinery element; or the first target is a translation
20 machinery element and the second target is a regulator of a translation machinery element; or the first target is a regulator of a first translation machinery element and the second target is a regulator of a second translation machinery element; or the first target is a first regulator of a first translation machinery element and the second target is a second regulator of the first translation machinery element, or any combination thereof.
25 Any of the combination embodiments described herein may include one or more modulators that are specific for the named target.

Exemplary methods for treating fibrotic disorders or for inhibiting
transdifferentiation of fibroblasts into myofibroblasts or for reducing the presence of myofibroblasts or for inhibiting or blocking the production and/or secretion of organ
30 damaging proteins from myofibroblasts, or any combination thereof, comprise administering combinations of two or more modulators (for the same or different

targets) of this disclosure. In any of these embodiments, a modulator may be specific for its target.

Representative combinations of modulators for use in the methods described herein include (1) a modulator of eIF4A and a modulator of eIF2A, (2) a modulator of eIF4A and a modulator of eIF2AK1, (3) a modulator of eIF4A and a modulator of eIF5A, (4) a modulator of eIF4A and a modulator of DHPS, (5) a modulator of eIF4A and a modulator of DOHH, (6) a modulator of eIF4A and a modulator of eEF1A1, (7) a modulator of eIF4A and a modulator of eEF2, (8) a modulator of eIF4A and a modulator of eEF2K, (9) a modulator of eIF4A and a modulator of eIF4B, (10) a modulator of eIF4A and a modulator of eIF4G, or (11) any combination thereof.

In further embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of eIF4E and a modulator of eIF2A, (2) a modulator of eIF4E and a modulator of eIF2AK1, (3) a modulator of eIF4E and a modulator of eIF5A, (4) a modulator of eIF4E and a modulator of DHPS, (5) a modulator of eIF4E and a modulator of DOHH, (6) a modulator of eIF4E and a modulator of eEF1A1, (7) a modulator of eIF4E and a modulator of eEF2, (8) a modulator of eIF4E and a modulator of eEF2K, (9) a modulator of eIF4E and a modulator of eIF4B, (10) a modulator of eIF4E and a modulator of eIF4G, or (11) any combination thereof.

In other embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of eIF4G and a modulator of eIF2A, (2) a modulator of eIF4G and a modulator of eIF2AK1, (3) a modulator of eIF4G and a modulator of eIF5A, (4) a modulator of eIF4G and a modulator of DHPS, (5) a modulator of eIF4G and a modulator of DOHH, (6) a modulator of eIF4G and a modulator of eEF1A1, (7) a modulator of eIF4G and a modulator of eEF2, (8) a modulator of eIF4G and a modulator of eEF2K, (9) a modulator of eIF4G and a modulator of eIF4B, or (10) any combination thereof.

In still further embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of eIF4B and a modulator of eIF2A, (2) a modulator of eIF4B and a modulator of eIF2AK1, (3) a modulator of eIF4B and a modulator of eIF5A, (4) a modulator of eIF4B and a modulator of DHPS, (5) a

modulator of eIF4B and a modulator of DOHH, (6) a modulator of eIF4B and a modulator of eEF1A1, (7) a modulator of eIF4B and a modulator of eEF2, (8) a modulator of eIF4B and a modulator of eEF2K, or (9) any combination thereof.

In further embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of eIF2A and a modulator of eIF2AK, (2) a modulator of eIF2A and a modulator of eIF5A, (3) a modulator of eIF2A and a modulator of DHPS, (4) a modulator of eIF2A and a modulator of DOHH, (5) a modulator of eIF2A and a modulator of eEF1A1, (6) a modulator of eIF2A and a modulator of eEF2, (7) a modulator of eIF2A and a modulator of eEF2K, or (8) any combination thereof.

In yet further embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of eIF2AK and a modulator of eIF5A, (2) a modulator of eIF2AK and a modulator of DHPS, (3) a modulator of eIF2AK and a modulator of DOHH, (4) a modulator of eIF2AK and a modulator of eEF1A1, (5) a modulator of eIF2AK and a modulator of eEF2, (6) a modulator of eIF2AK and a modulator of eEF2K, or (7) any combination thereof.

In certain embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of DHPS and a modulator of eIF5A, (2) a modulator of DHPS and a modulator of DOHH, (3) a modulator of DHPS and a modulator of eEF1A1, (4) a modulator of DHPS and a modulator of eEF2, (5) a modulator of DHPS and a modulator of eEF2K, or (6) any combination thereof.

In more embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of DOHH and a modulator of eIF5A, (2) a modulator of DOHH and a modulator of eEF1A1, (3) a modulator of DOHH and a modulator of eEF2, (4) a modulator of DOHH and a modulator of eEF2K, or (5) any combination thereof.

In still more embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of eIF5A and a modulator of eEF1A1, (2) a modulator of eIF5A and a modulator of eEF2, (3) a modulator of eIF5A and a modulator of eEF2K, or (4) any combination thereof.

More exemplary methods for treating fibrotic disorders or for inhibiting transdifferentiation of fibroblasts into myofibroblasts or for reducing the presence of myofibroblasts or for inhibiting or blocking the production and/or secretion of organ damaging proteins from myofibroblasts, or any combination thereof, comprise

5 administering combinations of modulators targeting at least two different regulators of translation machinery elements, such as (1) a modulator of eIF2AK1 and a modulator of a PI3K, (2) a modulator of eIF2AK1 and a modulator of an AKT, (3) a modulator of eIF2AK1 and a modulator of mTOR, (4) a modulator of eIF2AK1 and a modulator of a S6K70, (5) a modulator of eIF2AK1 and a modulator of an MNK, (6) a modulator of

10 eIF2AK1 and a modulator of a MEK1/2, (7) a modulator of eIF2AK1 and a modulator of an ERK, (8) a modulator of eIF2AK1 and a modulator of a RSK90, (9) a modulator of eIF2AK1 and a modulator of eEF2K, or (10) any combination thereof.

In further embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of eIF2AK2 and a modulator of a PI3K, (2) a

15 modulator of eIF2AK2 and a modulator of an AKT, (3) a modulator of eIF2AK2 and a modulator of mTOR, (4) a modulator of eIF2AK2 and a modulator of a S6K70, (5) a modulator of eIF2AK2 and a modulator of an MNK, (6) a modulator of eIF2AK2 and a modulator of a MEK1/2, (7) a modulator of eIF2AK2 and a modulator of an ERK, (8) a modulator of eIF2AK2 and a modulator of a RSK90, (9) a modulator of eIF2AK2 and a

20 modulator of eEF2K, or (10) any combination thereof.

In still further embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of eIF2AK3 and a modulator of a PI3K, (2) a modulator of eIF2AK3 and a modulator of an AKT, (3) a modulator of eIF2AK3 and a modulator of mTOR, (4) a modulator of eIF2AK3 and a modulator of a S6K70, (5) a

25 modulator of eIF2AK3 and a modulator of an MNK, (6) a modulator of eIF2AK3 and a modulator of a MEK1/2, (7) a modulator of eIF2AK3 and a modulator of an ERK, (8) a modulator of eIF2AK3 and a modulator of a RSK90, (9) a modulator of eIF2AK3 and a modulator of eEF2K, or (10) any combination thereof.

In yet further embodiments, combinations of modulators for use in the methods

30 described herein comprise (1) a modulator of eIF2AK4 and a modulator of a PI3K, (2) a modulator of eIF2AK4 and a modulator of an AKT, (3) a modulator of eIF2AK4 and a

modulator of mTOR, (4) a modulator of eIF2AK4 and a modulator of a S6K70, (5) a
 modulator of eIF2AK4 and a modulator of an MNK, (6) a modulator of eIF2AK4 and a
 modulator of a MEK1/2, (7) a modulator of eIF2AK4 and a modulator of an ERK, (8) a
 modulator of eIF2AK4 and a modulator of a RSK90, (9) a modulator of eIF2AK4 and a
 5 modulator of eEF2K, or (10) any combination thereof.

In more embodiments, combinations of modulators for use in the methods
 described herein comprise (1) a modulator of GADD34 and a modulator of a PI3K, (2)
 a modulator of GADD34 and a modulator of an AKT, (3) a modulator of GADD34 and
 a modulator of mTOR, (4) a modulator of GADD34 and a modulator of a S6K70, (5) a
 10 modulator of GADD34 and a modulator of an MNK, (6) a modulator of GADD34 and a
 modulator of a MEK1/2, (7) a modulator of GADD34 and a modulator of an ERK, (8) a
 modulator of GADD34 and a modulator of a RSK90, (9) a modulator of GADD34 and
 a modulator of eEF2K, or (10) any combination thereof.

In other embodiments, combinations of modulators for use in the methods
 15 described herein comprise (1) a modulator of GSK3 β and a modulator of a PI3K, (2) a
 modulator of GSK3 β and a modulator of an AKT, (3) a modulator of GSK3 β and a
 modulator of mTOR, (4) a modulator of GSK3 β and a modulator of a S6K70, (5) a
 modulator of GSK3 β and a modulator of an MNK, (6) a modulator of GSK3 β and a
 modulator of a MEK1/2, (7) a modulator of GSK3 β and a modulator of an ERK, (8) a
 20 modulator of GSK3 β and a modulator of a RSK90, (9) a modulator of GSK3 β and a
 modulator of eEF2K, or (10) any combination thereof.

In certain embodiments, combinations of modulators for use in the methods
 described herein comprise (1) a modulator of a PI3K and a modulator of a MNK, (2) a
 modulator of a PI3K and a modulator of DHPS, (3) a modulator of a PI3K and a
 25 modulator of DOHH, (4) a modulator of a PI3K and a modulator of a MEK1/2, (5) a
 modulator of a PI3K and a modulator of an ERK, (6) a modulator of a PI3K and a
 modulator of a RSK90, (7) a modulator of a PI3K and a modulator of eEF2K, or (8) any
 combination thereof.

In other embodiments, combinations of modulators for use in the methods
 30 described herein comprise (1) a modulator of an AKT and a modulator of a MNK, (2) a
 modulator of an AKT and a modulator of DHPS, (3) a modulator of an AKT and a

modulator of DOHH, (4) a modulator of an AKT and a modulator of a MEK1/2, (5) a modulator of an AKT and a modulator of an ERK, (6) a modulator of an AKT and a modulator of a RSK90, (7) a modulator of an AKT and a modulator of eEF2K, or (8) any combination thereof.

5 In still other embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of mTOR and a modulator of a MNK, (2) a modulator of mTOR and a modulator of DHPS, (3) a modulator of mTOR and a modulator of DOHH, (4) a modulator of mTOR and a modulator of a MEK1/2, (5) a modulator of mTOR and a modulator of an ERK, (6) a modulator of mTOR and a
10 modulator of a RSK90, (7) a modulator of mTOR and a modulator of eEF2K, or (8) any combination thereof.

In yet other embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of S6K70 and a modulator of a MNK, (2) a modulator of S6K70 and a modulator of DHPS, (3) a modulator of S6K70 and a
15 modulator of DOHH, (4) a modulator of S6K70 and a modulator of a MEK1/2, (5) a modulator of S6K70 and a modulator of an ERK, (6) a modulator of S6K70 and a modulator of a RSK90, (7) a modulator of S6K70 and a modulator of eEF2K, or (8) any combination thereof.

In further embodiments, combinations of modulators for use in the methods
20 described herein comprise (1) a modulator of a MNK and a modulator of DHPS, (2) a modulator of a MNK and a modulator of DOHH, (3) a modulator of a MNK and a modulator of a MEK1/2, (4) a modulator of a MNK and a modulator of an ERK, (5) a modulator of a MNK and a modulator of a RSK90, (6) a modulator of a MNK and a modulator of eEF2K, or (7) any combination thereof.

25 In other embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of DHPS and a modulator of a MEK1/2, (2) a modulator of DHPS and a modulator of an ERK, (3) a modulator of DHPS and a modulator of a RSK90, (4) a modulator of DHPS and a modulator of eEF2K, or (5) any combination thereof.

30 In more embodiments, combinations of modulators for use in the methods described herein comprise administering (1) a modulator of DOHH and a modulator of

a MEK1/2, (2) a modulator of DOHH and a modulator of an ERK, (3) a modulator of DOHH and a modulator of a RSK90, (4) a modulator of DOHH and a modulator of eEF2K, or (5) any combination thereof.

In even more embodiments, combinations of modulators for use in the methods described herein comprise administering (1) a modulator of eEF2K and a modulator of a MEK1/2, (2) a modulator of eEF2K and a modulator of an ERK, (3) a modulator of eEF2K and a modulator of a RSK90, or (4) any combination thereof.

Further exemplary methods for treating fibrotic disorders or for inhibiting transdifferentiation of fibroblasts into myofibroblasts or for reducing the presence of myofibroblasts or for inhibiting or blocking the production and/or secretion of organ damaging proteins from myofibroblasts, or any combination thereof, comprise administering combinations of modulators targeting at least two different translation machinery elements, such as (1) a modulator of an eIF4A and a modulator of a eEF1, (2) a modulator of an eIF4A and a modulator of an eEF2, (3) a modulator of an eIF4A and a modulator of eIF5A, or (4) any combination thereof.

In certain embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of an eIF4E and a modulator of a eEF1, (2) a modulator of an eIF4E and a modulator of an eEF2, (3) a modulator of an eIF4E and a modulator of eIF5A, or (4) any combination thereof.

In further embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of an eIF5A and a modulator of a eEF1, (2) a modulator of an eIF5A and a modulator of an eEF2, or (3) any combination thereof.

Additional exemplary methods for treating fibrotic disorders or for inhibiting transdifferentiation of fibroblasts into myofibroblasts or for reducing the presence of myofibroblasts or for inhibiting or blocking the production and/or secretion of organ damaging proteins from myofibroblasts, or any combination thereof, comprise administering combinations of modulators targeting at least one translation machinery element and at least one regulator of translation machinery elements, such as (1) a modulator of eIF2AK1 and a modulator of an eIF4A, (2) a modulator of eIF2AK1 and a modulator of an eIF4E, (3) a modulator of eIF2AK1 and a modulator of eEF1, (4) a

modulator of eIF2AK1 and a modulator of a eEF2, (5) a modulator of eIF2AK1 and a modulator of an eIF5A, or (6) any combination thereof.

In certain embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of eIF2AK2 and a modulator of an eIF4A, 5 (2) a modulator of eIF2AK2 and a modulator of an eIF4E, (3) a modulator of eIF2AK2 and a modulator of eEF1, (4) a modulator of eIF2AK2 and a modulator of a eEF2, (5) a modulator of eIF2AK2 and a modulator of an eIF5A, or (6) any combination thereof.

In further embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of eIF2AK3 and a modulator of an eIF4A, 10 (2) a modulator of eIF2AK3 and a modulator of an eIF4E, (3) a modulator of eIF2AK3 and a modulator of eEF1, (4) a modulator of eIF2AK3 and a modulator of a eEF2, (5) a modulator of eIF2AK3 and a modulator of an eIF5A, or (6) any combination thereof.

In more embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of eIF2AK4 and a modulator of an eIF4A, 15 (2) a modulator of eIF2AK4 and a modulator of an eIF4E, (3) a modulator of eIF2AK4 and a modulator of eEF1, (4) a modulator of eIF2AK4 and a modulator of a eEF2, (5) a modulator of eIF2AK4 and a modulator of an eIF5A, or (6) any combination thereof.

In other embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of GADD34 and a modulator of an eIF4A, 20 (2) a modulator of GADD34 and a modulator of an eIF4E, (3) a modulator of GADD34 and a modulator of eEF1, (4) a modulator of GADD34 and a modulator of a eEF2, (5) a modulator of GADD34 and a modulator of an eIF5A, or (6) any combination thereof.

In still more embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of GSK3 β and a modulator of an eIF4A, (2) 25 a modulator of GSK3 β and a modulator of an eIF4E, (3) a modulator of GSK3 β and a modulator of eEF1, (4) a modulator of GSK3 β and a modulator of a eEF2, (5) a modulator of GSK3 β and a modulator of an eIF5A, or (6) any combination thereof.

In certain embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of a PI3K and a modulator of an eIF4A, (2) a 30 modulator of a PI3K and a modulator of an eIF4E, (3) a modulator of a PI3K and a

modulator of eEF1, (4) a modulator of a PI3K and a modulator of a eEF2, (5) a modulator of a PI3K and a modulator of an eIF5A, or (6) any combination thereof.

In further embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of an AKT and a modulator of an eIF4A, (2) a modulator of an AKT and a modulator of an eIF4E, (3) a modulator of an AKT and a modulator of eEF1, (4) a modulator of an AKT and a modulator of a eEF2, (5) a modulator of an AKT and a modulator of an eIF5A, or (6) any combination thereof.

In more embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of mTOR and a modulator of an eIF4A, (2) a modulator of mTOR and a modulator of an eIF4E, (3) a modulator of mTOR and a modulator of eEF1, (4) a modulator of mTOR and a modulator of a eEF2, (5) a modulator of mTOR and a modulator of an eIF5A, or (6) any combination thereof.

In more embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of S6K70 and a modulator of an eIF4A, (2) a modulator of S6K70 and a modulator of an eIF4E, (3) a modulator of S6K70 and a modulator of eEF1, (4) a modulator of S6K70 and a modulator of a eEF2, (5) a modulator of S6K70 and a modulator of an eIF5A, or (6) any combination thereof.

In certain embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of a MNK and a modulator of an eIF4A, (2) a modulator of a MNK and a modulator of an eIF4E, (3) a modulator of a MNK and a modulator of eEF1, (4) a modulator of a MNK and a modulator of a eEF2, (5) a modulator of a MNK and a modulator of an eIF5A, or (6) any combination thereof.

In certain embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of a MEK1/2 and a modulator of an eIF4A, (2) a modulator of a MEK1/2 and a modulator of an eIF4E, (3) a modulator of a MEK1/2 and a modulator of eEF1, (4) a modulator of a MEK1/2 and a modulator of a eEF2, (5) a modulator of a MEK1/2 and a modulator of an eIF5A, or (6) any combination thereof.

In further embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of an ERK and a modulator of an eIF4A, (2) a modulator of an ERK and a modulator of an eIF4E, (3) a modulator of an ERK and a

modulator of eEF1, (4) a modulator of an ERK and a modulator of a eEF2, (5) a modulator of an ERK and a modulator of an eIF5A, or (6) any combination thereof.

In more embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of RSK90 and a modulator of an eIF4A, (2) a modulator of RSK90 and a modulator of an eIF4E, (3) a modulator of RSK90 and a modulator of eEF1, (4) a modulator of RSK90 and a modulator of a eEF2, (5) a modulator of RSK90 and a modulator of an eIF5A, or (6) any combination thereof.

In even more embodiments, combinations of modulators for use in the methods described herein comprise (1) a modulator of eEF2K and a modulator of an eIF4A, (2) a modulator of eEF2K and a modulator of an eIF4E, (3) a modulator of eEF2K and a modulator of an eIF5A, or (4) any combination thereof.

In any of the combination therapies described herein, a combination of modulators can be administered serially, simultaneously, or concurrently. When administering serially, a first modulator or pharmaceutical composition thereof is formulated in a separate composition from a second (or third, etc.) modulator or pharmaceutical composition thereof. When administering simultaneously or concurrently, a first and second (or third, etc.) modulator may be formulated in separate compositions or formulated in a single composition. In any of these embodiments, the single or combination modulator therapies can be administered as a single dose unit or administered as a single dose unit a plurality of times (daily, weekly, biweekly, monthly, biannually, annually, etc., or any combination thereof).

In certain embodiments, a combination therapy described herein is used in a method for treating a fibrotic disorder or disease. In further embodiments, a combination therapy described herein is used in a method for inhibiting transdifferentiation of fibroblasts into myofibroblasts. In still further embodiments, a combination therapy described herein is used in a method for reducing the presence of myofibroblasts. In yet further embodiments, a combination therapy described herein is used in a method for inhibiting or blocking the production and/or secretion of organ damaging proteins from myofibroblasts.

In certain embodiments, a combination of modulators described herein is used in a method of identifying a subject as a candidate for preventing, treating or ameliorating a fibrotic disease with a therapeutic agent.

In still further embodiments, a modulator of a target of interest or any of the
5 aforementioned combinations, which may be specific for their target, are administered in combination with one or more adjunctive therapeutic agents, such as angiotensin converting enzyme inhibitor, nintedanib (BIBF-1120), STX-100, QAX576, CNTO-888, SD-208, SB-525334, GC1008, BMS-986202, AM152, lebrikizumab, tralokinumab, SAR156597, PRM-151, simtuzumab (AB0024, GS-6624), GSK2126458, FG-3019,
10 captopril, genistein, EUK-207, silvestrol or derivatives thereof, pateamine A or derivatives thereof, hippuristanol, or pirfenidone, for the treatment of, for example, a fibrotic disorder (such as idiopathic pulmonary fibrosis).

In any of the aforementioned embodiments, the fibrotic disease or disorder is a pulmonary fibrosis, idiopathic pulmonary fibrosis, cystic fibrosis, liver fibrosis, cardiac
15 fibrosis, endomyocardial fibrosis, atrial fibrosis, mediastinal fibrosis, myelofibrosis, retroperitoneal fibrosis, chronic kidney disease, nephrogenic systemic fibrosis, Chron's disease, hypertrophic scarring, keloid, scleroderma, organ transplant-associated fibrosis, ischemia-associated fibrosis, or a combination thereof. In any of the aforementioned embodiments, the subject is a human.

20 Subjects in need of administration of therapeutic agents as described herein include subjects at high risk for developing a fibrotic disorder as well as subjects presenting with an existing fibrotic disorder. A subject may be at high risk for developing a fibrotic disorder if the subject has experienced an injury, for example: exposure to radiation, environmental or occupational pollutant, chemical, irritant,
25 certain medications, infectious agent; having a certain genetic mutation; chronic autoimmune response; or an ischemic event. Subjects suffering from or suspected of having a fibrotic disorder can be identified using methods as described herein. A subject may be any organism capable of developing a fibrotic disorder, such as humans, pets, livestock, show animals, zoo specimens, or other animals. For example, a subject
30 may be a human, a non-human primate, dog, cat, rabbit, horse, or the like.

The therapeutic agents or pharmaceutical compositions that treat or reduce the risk of developing a fibrotic disorder provided herein are administered to a subject who has or is at risk of developing a fibrotic disorder at a therapeutically effective amount or dose. Such a dose may be determined or adjusted depending on various factors

5 including the specific therapeutic agents or pharmaceutical compositions, the routes of administration, the subject's condition, that is, stage of the disease, severity of symptoms caused by the disease, general health status, as well as age, gender, and weight, and other factors apparent to a person skilled in the medical art. Similarly, the dose of the therapeutic for treating a disease or disorder may be determined according

10 to parameters understood by a person skilled in the medical art. When referring to a combination, a therapeutically effective dose refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered serially or simultaneously (in the same formulation or concurrently in separate formulations). Optimal doses may generally be determined using experimental models and/or clinical

15 trials. Design and execution of pre-clinical and clinical studies for a therapeutic agent (including when administered for prophylactic benefit) described herein are well within the skill of a person skilled in the relevant art.

Generally, the therapeutic agent is administered at a therapeutically effective amount or dose. A therapeutically effective amount or dose will vary according to

20 several factors, including the chosen route of administration, formulation of the composition, patient response, severity of the condition, the subject's weight, and the judgment of the prescribing physician. The dosage can be increased or decreased over time, as required by an individual patient. In certain instances, a patient initially is given a low dose, which is then increased to an efficacious dosage tolerable to the

25 patient. Determination of an effective amount is well within the capability of those skilled in the art.

The route of administration of a therapeutic agent can be oral, intraperitoneal, transdermal, subcutaneous, by intravenous or intramuscular injection, by inhalation, topical, intralesional, infusion; liposome-mediated delivery; topical, intrathecal,

30 gingival pocket, rectal, intrabronchial, nasal, transmucosal, intestinal, ocular or otic delivery, or any other methods known in the art.

In some embodiments, a therapeutic agent is formulated as a pharmaceutical composition. In some embodiments, a pharmaceutical composition incorporates particulate forms, protective coatings, protease inhibitors, or permeation enhancers for various routes of administration, including parenteral, pulmonary, nasal and oral. The pharmaceutical compositions can be administered in a variety of unit dosage forms depending upon the method/mode of administration. Suitable unit dosage forms, including powders, tablets, pills, capsules, lozenges, suppositories, patches, nasal sprays, injectables, implantable sustained-release formulations, *etc.*

In some embodiments, a pharmaceutical composition comprises an acceptable diluent, carrier or excipient. A pharmaceutically acceptable carrier includes any solvent, dispersion media, or coating that are physiologically compatible and that preferably do not interfere with or otherwise inhibit the activity of the therapeutic agent. Preferably, a carrier is suitable for intravenous, intramuscular, oral, intraperitoneal, transdermal, topical, or subcutaneous administration. Pharmaceutically acceptable carriers can contain one or more physiologically acceptable compound(s) that act, for example, to stabilize the composition or to increase or decrease the absorption of the active agent(s). Physiologically acceptable compounds can include, for example, carbohydrates, such as glucose, sucrose, or dextrans, antioxidants, such as ascorbic acid or glutathione, chelating agents, low molecular weight proteins, compositions that reduce the clearance or hydrolysis of the active agents, or excipients or other stabilizers and/or buffers. Other pharmaceutically acceptable carriers and their formulations are well-known and generally described in, for example, *Remington: The Science and Practice of Pharmacy*, 21st Edition, Philadelphia, PA. Lippincott Williams & Wilkins, 2005. Various pharmaceutically acceptable excipients are well-known in the art and can be found in, for example, *Handbook of Pharmaceutical Excipients* (5th ed., Ed. Rowe *et al.*, Pharmaceutical Press, Washington, D.C.).

EXAMPLES

EXAMPLE 1

EFFECT OF MTOR INHIBITION ON FIBROTIC DISEASE DEVELOPMENT

TGF β -mediated transformation of fibroblasts into fibrotic myofibroblasts is
 5 well-established as an essential step in fibroplasia, a key component of many fibrotic
 disorders (Blobe *et al.*, *N. Engl. J. Med.* 342:1350, 2000; Border and Noble, *N. Engl. J.*
Med. 331:1286, 1994). This TGF β -mediated transformation of fibroblasts was used as
 a model for examining fibrotic disorders.

Briefly, normal human lung fibroblasts (Lonza #CC-2512; cell passage numbers
 10 2 through 5 were used for all experiments) were seeded (Day 0) and cultured in
 Dulbecco's modified Eagle medium (DMEM) supplemented with 10% fetal bovine
 serum (FBS), penicillin, streptomycin and glutamax (Invitrogen) at 37°C in a
 humidified incubator with 5% CO₂ overnight. On Day 1, cells were harvested, washed
 with phosphate buffered saline (PBS), and then incubated for 48 hours in fresh
 15 serum-free DMEM supplemented with penicillin, streptomycin, and glutamax. On Day
 3, cells were harvested, resuspended in fresh serum-free DMEM containing PP242 (10,
 2.5, 0.625, 0.156, 0.039, 0.01 or 0.002 μ M) and 10 ng/ml TGF β , and cultured for 24
 hours. Controls included untreated cells, cells treated with TGF β only, and cells treated
 with PP242 only.

20 After this 24 hour incubation, procollagen type 1 levels were measured by
 collecting culture media, centrifuging to pellet cellular debris, and stored at -80°C.
 Procollagen Type 1 C-Peptide (PIPC) was quantified using the (PIP) EIA kit (Clontech,
 Cat. No. MK101) according to manufacturer's instructions. The TGF β -treated
 fibroblasts of this example were examined by ribosomal profiling (about 6 x 10⁶
 25 cells/10 cm plate) and western blot analysis (about 1 x 10⁶ cells/well of a 6-well plate).

Results

Transformation of fibroblasts into fibrotic myofibroblasts by treatment with
 TGF β for 24 hours was accompanied by an approximately 7-fold increase in
 procollagen production, while treatment with PP242 was able to block this increase

(EC₅₀ of about 0.2 μ M) (Figure 1). Expression of TGF β induced myofibroblast differentiation marker, α -smooth muscle actin (α -SMA), was also analyzed by Western blot analysis (*see* Example 3; Figure 2). After 24 hours of TGF β stimulation, increased α -SMA protein levels were detectable, while the level of β -actin did not change. As
5 with procollagen, co-incubation of cells with TGF- β and PP242 caused a reduction of the α -SMA protein level in a dose dependent manner.

Conclusion

Co-administration of TGF- β with mTOR inhibitor PP242 reverses or prevents the changes observed in a fibrotic disorder-related pathway as evidenced by an
10 inhibition of increased production of fibrotic disorder biomarker proteins, type 1 procollagen and α -SMA (which are both hallmarks of TGF β -mediated fibroblast transformation into myofibroblasts).

EXAMPLE 2

PHOSPHORYLATION OF PROTEIN TRANSLATION COMPONENTS:

15 EFFECT OF MTOR INHIBITION ON FIBROTIC DISEASE DEVELOPMENT

TGF β -dependent activation of the PI3K/Akt/mTOR and ERK pathways were also examined by western blot analysis (about 1×10^6 cells/well from a 6-well plate). Briefly, cells were washed with PBS and lysed in 1x cell lysis buffer (Cell Signaling Technology, Inc., Danvers, MA) for 15 minutes at 4°C. Lysates were briefly sonicated,
20 clarified by centrifugation for 15 minutes at 14,000 rpm, and then supernatants were collected. Protein concentration in the soluble fraction was determined by BCA protein assay (Thermo Scientific, Rockford, IL). Samples of protein (20 μ g) were resolved on 4-20% Bis-Tris gradient gel (Invitrogen, Carlsbad, CA) and transferred to nitrocellulose membrane. The resulting blots were blocked for 1 hour at room temperature with
25 Odyssey blocking solution (LI-COR) and then incubated with primary antibodies at 4°C overnight. The following day, each blot was washed three times for 10 minutes in TBST, and then incubated with goat anti-rabbit fluorescent conjugated secondary antibody (IRDye 800 CW at 1:20,000; LI-COR) for 1 hour at room temperature. The blots were washed and scanned, and then specific proteins were detected by using the

LI-COR Odyssey infrared imager. The following antibodies were used at 1:1000 dilution: anti- α -actin (Sigma #A2547), anti-phospho-4EBP(Ser65), anti-phospho-rpS6(Ser235/236)(#4858), anti-phospho-ERK1/2(Thr202/Tyr204)(#4370), anti-phospho-p70S6K(Thr421/Ser424)(#9204), anti-phospho-pAKT(Ser473), anti-phospho-MNK(Thr197/202), and anti- β -actin (#4970). Unless otherwise indicated, the antibodies were from Cell Signaling Technology, Inc. (Danvers, MA).

Results

Figure 2 shows that phosphorylation of AKT, 4EBP, S6K, and S6 of the mTOR pathway was strongly stimulated in fibroblasts treated with TGF β , whereas only a moderate increase in ERK phosphorylation was observed. Co-incubation of cells with mTOR inhibitor PP242 (0.625 μ M) was sufficient to abolish TGF β -dependent increases in phosphorylation of AKT, 4EBP, S6K, and S6, as well as causing a decrease in α -SMA to pretreatment levels (Figure 2).

Conclusion

Co-administration of TGF β with mTOR inhibitor PP242 reverses or prevents the changes observed in a fibrotic disorder-related pathway (*i.e.*, normalizes the translational efficiencies of the genes) and inhibits increased production of fibrotic disorder biomarker proteins, type 1 procollagen and α -smooth muscle actin (which are both hallmarks of TGF β -mediated fibroblast transformation to myofibroblasts).

20

EXAMPLE 3

TRANSLATIONAL PROFILING: EFFECT OF mTOR INHIBITION ON FIBROTIC DISEASE DEVELOPMENT

Ribosomal profiling allows for measurement of changes in transcription and translation on a genome-wide basis accompanying TGF β -dependent transformation of fibroblasts to myofibroblasts. Ribosomal profiles of the TGF β -treated fibroblasts from Example 1 (about 6×10^6 cells/10 cm plate) were prepared and analyzed for changes in translational efficiencies with respect to potential disease-associated cellular changes accompanying this TGF β -induced transformation.

Briefly, cells were washed with cold PBS supplemented with cycloheximide and lysed with 1x mammalian cell lysis buffer for 10 minutes on ice. Lysates were clarified by centrifugation for 10 minutes at 14,000 rpm and supernatants were collected. Cell lysates were processed to generate ribosomal protected fragments and total mRNA
5 according to the instructions included with the ARTseq Ribosome Profiling Kit (Illumina, San Diego, CA). Sequencing of total RNA (RNA) and of ribosome-protected fragments of RNA (RPF) was carried out using RNA-Seq methodology according to the manufacturer's instructions (Illumina). To analyze the ribosomal profiles, RNA-Seq reads were processed with tools from the FASTX-Toolkit
10 (fastq_quality_trimmer, fastx_clipper and fastx_trimmer). Unprocessed and processed reads were evaluated for a variety of quality measures using FastQC, and processed reads were mapped to the human genome using TopHat (*see, e.g.,* Trapnell *et al.*, Bioinformatics 25:1105, 2009). Gene-by-gene assessment of the number of fragments strictly and uniquely mapping to the coding region of each gene was conducted using
15 HTSeq-count, a component of the HTSeq package. Differential analyses of the transforming effect of TGF β on fibroblasts and the effect of PP242 treatment on this transformation were carried out with the software packages DESeq for transcription (RNA counts) and translational rate (RPF counts) and BABEL for translational efficiency based upon ribosomal occupancy as a function of RNA level (RNA and RPF
20 counts). Genes with low counts in either RPF or RNA were excluded from differential analyses. Pathway and network analyses of differential data were conducted using Ingenuity Pathway Analysis (IPA).

Results

Genes identified in TGF β treated fibroblasts by Differential Expression Analysis
25 (differential mRNA amount (transcription), differential translational rate (RPF counts) and differential translational efficiency (ratio of translational rate to mRNA amount)) are listed in Tables 1-3 (Table 3A shows genes with an altered translational efficiency identified after one profiling experiment and Table 3B has a refined gene list based on five replicate profiling experiments). These three gene signatures were analyzed for
30 pathway and network connections using Ingenuity Pathway Analysis (IPA).

Table 1. Gene Signature with Altered Translation Rate

ENSEMBL ID	HGNC ID	log₂FC RPF(M) / RPF(Ctrl)	RPF p-value
ENSG00000049540	ELN	5.36941753	1.08E-07
ENSG00000163661	PTX3	-5.3307069	1.41E-07
ENSG00000107796	ACTA2	5.26064763	1.67E-07
ENSG00000146674	IGFBP3	4.82169249	1.06E-06
ENSG00000123610	TNFAIP6	4.75270158	1.41E-06
ENSG00000106366	SERPINE1	4.33934001	6.65E-06
ENSG00000171793	CTPS	4.09973744	2.08E-05
ENSG00000138735	PDE5A	-4.0235025	2.74E-05
ENSG00000140416	TPM1	3.97312305	2.76E-05
ENSG00000148848	ADAM12	3.93485546	3.60E-05
ENSG00000076706	MCAM	4.03592992	3.84E-05
ENSG00000139211	AMIGO2	3.93283643	4.20E-05
ENSG00000130707	ASS1	3.94677625	5.20E-05
ENSG00000103489	XYLT1	3.66194114	0.0001064
ENSG00000149591	TAGLN	3.60417531	0.00010993
ENSG00000135074	ADAM19	3.65751476	0.00011063
ENSG00000177283	FZD8	3.66199763	0.00012173
ENSG00000197442	MAP3K5	-3.6700003	0.00015186
ENSG00000162591	MEGF6	3.52392664	0.00018903
ENSG00000170558	CDH2	3.37710278	0.00027648
ENSG00000211455	STK38L	3.35942032	0.00033916
ENSG00000164761	TNFRSF11B	-3.3013119	0.0003522
ENSG00000127863	TNFRSF19	-3.3472307	0.00036218
ENSG00000006016	CRLF1	3.35876314	0.0003716
ENSG00000187498	COL4A1	3.22671553	0.00043313
ENSG00000128965	CHAC1	3.32922215	0.00043675
ENSG00000172986	GXYLT2	3.34594516	0.0004446
ENSG00000113140	SPARC	3.18391733	0.00050358
ENSG00000189184	PCDH18	-3.2023222	0.00052431
ENSG00000099860	GADD45B	3.15900734	0.0007182
ENSG00000197321	SVIL	-3.1352953	0.00074287
ENSG00000176170	SPHK1	3.17785195	0.00083387
ENSG00000079308	TNS1	3.0455728	0.00083771
ENSG00000115884	SDC1	3.08392695	0.00087241
ENSG00000137809	ITGA11	3.01124359	0.00103089
ENSG00000135269	TES	3.05095571	0.00108443

ENSEMBL ID	HGNC ID	log ₂ FC RPF(M) / RPF(Ctrl)	RPF p-value
ENSG00000106211	HSPB1	2.94411799	0.00117519
ENSG00000122786	CALD1	2.92932598	0.00122331
ENSG00000137124	ALDH1B1	2.99287373	0.0012512
ENSG00000107957	SH3PXD2A	2.9304952	0.00130719
ENSG00000206190	ATP10A	3.00335812	0.00131835
ENSG00000163453	IGFBP7	2.8797874	0.00145005
ENSG00000137331	IER3	2.83980944	0.00184245
ENSG00000120708	TGFBI	2.80133447	0.00187418
ENSG00000115902	SLC1A4	2.85901341	0.00197395
ENSG00000103257	SLC7A5	2.80503703	0.00215617
ENSG00000070371	CLTCL1	2.90823707	0.00218709
ENSG00000127241	MASP1	-2.7577773	0.00241631
ENSG00000164292	RHOBTB3	-2.7504911	0.00246705
ENSG00000171617	ENC1	2.74021006	0.00248563
ENSG00000158186	MRAS	2.92224323	0.00271268
ENSG00000182752	PAPPA	-2.7035058	0.00294361
ENSG00000134853	PDGFRA	-2.6580228	0.00312982
ENSG00000165029	ABCA1	-2.6467049	0.00326207
ENSG00000118523	CTGF	2.62517157	0.00334596
ENSG00000198853	RUSC2	2.65438677	0.00352266
ENSG00000188641	DPYD	-2.6916522	0.00352337
ENSG00000147872	PLIN2	-2.6415827	0.00357964
ENSG00000119408	NEK6	-2.6438108	0.00399688
ENSG00000070669	ASNS	2.58331253	0.00400905
ENSG00000065911	MTHFD2	2.5641977	0.00439044
ENSG00000131435	PDLIM4	2.55921868	0.00458365
ENSG00000197594	ENPP1	2.56938165	0.00466656
ENSG00000165124	SVEP1	-2.5106635	0.00496929
ENSG00000104368	PLAT	-2.5054577	0.00517753
ENSG00000162804	SNED1	-2.5154081	0.00558283
ENSG00000087008	ACOX3	2.49632218	0.00613453
ENSG00000156265	C21orf7	2.56814445	0.00614118
ENSG00000139329	LUM	-2.4449067	0.00616621
ENSG00000183010	PYCR1	2.46985869	0.00638868
ENSG00000214517	PPME1	2.44759383	0.0065779
ENSG00000111186	WNT5B	2.46335531	0.00661341
ENSG00000122694	GLIPR2	2.49429147	0.00678059

ENSEMBL ID	HGNC ID	log ₂ FC RPF(M) / RPF(Ctrl)	RPF p-value
ENSG00000100234	TIMP3	2.40382243	0.00681395
ENSG00000136542	GALNT5	-2.4288148	0.00695359
ENSG00000198121	LPAR1	-2.4357876	0.00713053
ENSG00000154553	PDLIM3	2.45374968	0.00729239
ENSG00000162520	SYNC	2.53882525	0.00736301
ENSG00000135919	SERPINE2	2.37546587	0.00738884
ENSG00000115963	RND3	-2.3750235	0.00749913
ENSG00000112096	SOD2	-2.4089889	0.00750824
ENSG00000168994	PXDC1	2.41631019	0.00757789
ENSG00000130635	COL5A1	2.36463839	0.00765854
ENSG00000125257	ABCC4	-2.3915739	0.00767406
ENSG00000175899	A2M	-2.3564983	0.00789551
ENSG00000147224	PRPS1	2.42227839	0.00792948
ENSG00000134352	IL6ST	-2.3544877	0.00798272
ENSG00000144655	CSRNPI	2.42466893	0.00830203
ENSG00000121068	TBX2	-2.3897652	0.00831686
ENSG00000183963	SMTN	2.35351506	0.008544
ENSG00000161638	ITGA5	2.33254716	0.00856086
ENSG00000073712	FERMT2	2.3323394	0.00864574
ENSG00000128590	DNAJB9	2.35073771	0.00877118
ENSG00000133816	MICAL2	2.31968643	0.00902167
ENSG00000113083	LOX	2.31217587	0.00910434
ENSG00000105329	TGFB1	2.31202161	0.00921304
ENSG00000134871	COL4A2	2.29661847	0.00939314
ENSG00000106772	PRUNE2	2.32751681	0.00944999
ENSG00000152952	PLOD2	2.28476269	0.00993745
ENSG00000110328	GALNTL4	2.45799148	0.00994213
ENSG00000147852	VLDLR	2.32508202	0.01043418
ENSG00000011422	PLAUR	2.29496533	0.01050673
ENSG00000186340	THBS2	2.26219171	0.01052617
ENSG00000164465	DCBLD1	2.29448412	0.01058626
ENSG00000100889	PCK2	2.28919489	0.01072773
ENSG00000171223	JUNB	2.24626722	0.01131056
ENSG00000134285	FKBP11	2.28988985	0.01148763
ENSG00000163110	PDLIM5	2.23835271	0.01163489
ENSG00000181104	F2R	-2.2106721	0.01226847
ENSG00000106799	TGFBR1	2.21730253	0.01229295

ENSEMBL ID	HGNC ID	log ₂ FC RPF(M) / RPF(Ctrl)	RPF p-value
ENSG00000136205	TNS3	-2.2199359	0.01255329
ENSG00000135048	TMEM2	2.22084958	0.01287039
ENSG00000099250	NRP1	-2.1921383	0.01318577
ENSG00000196923	PDLIM7	2.19174283	0.01327444
ENSG00000175183	CSRP2	2.23683689	0.01398546
ENSG00000128591	FLNC	2.16561759	0.01399886
ENSG00000165072	MAMDC2	2.19235097	0.01402561
ENSG00000135069	PSAT1	2.1684862	0.01430236
ENSG00000166888	STAT6	-2.2016097	0.0143223
ENSG00000134668	SPOCD1	2.30705749	0.01433835
ENSG00000142552	RCN3	2.14884922	0.01537399
ENSG00000165801	ARHGEF40	2.140188	0.0157106
ENSG00000132688	NES	2.12201961	0.01691507
ENSG00000141753	IGFBP4	-2.1015663	0.01722583
ENSG00000158966	CACHD1	2.19269046	0.01756212
ENSG00000114850	SSR3	2.08950932	0.01804143
ENSG00000112902	SEMA5A	-2.1061326	0.01873853
ENSG00000067057	PFKP	2.070317	0.01875024
ENSG00000072682	P4HA2	2.06604184	0.01900324
ENSG00000113361	CDH6	2.07218406	0.01905514
ENSG00000129038	LOXL1	2.07073937	0.01937479
ENSG00000154122	ANKH	2.09170046	0.01954498
ENSG00000142871	CYR61	2.0468831	0.01960644
ENSG00000136603	SKIL	2.07686006	0.01962497
ENSG00000135905	DOCK10	2.13008136	0.01976733
ENSG00000149256	ODZ4	-2.0726023	0.01995218
ENSG00000065054	SLC9A3R2	2.06603755	0.02075587
ENSG00000107249	GLIS3	2.20934781	0.02122407
ENSG00000174099	MSRB3	2.05355244	0.02238316
ENSG00000198467	TPM2	1.9923924	0.02282632
ENSG00000135931	ARMC9	-2.039883	0.02397523
ENSG00000100139	MICALL1	2.04996518	0.02417362
ENSG00000187840	EIF4EBP1	2.02361386	0.02463796
ENSG00000127418	FGFRL1	2.04284709	0.02477047
ENSG00000100600	LGMN	1.98781976	0.02480798
ENSG00000129116	PALLD	1.9624052	0.02496631
ENSG00000109861	CTSC	-1.973874	0.02538713

ENSEMBL ID	HGNC ID	log ₂ FC RPF(M) / RPF(Ctrl)	RPF p-value
ENSG00000152377	SPOCK1	1.95829856	0.02546107
ENSG00000164574	GALNT10	1.9899263	0.02554389
ENSG00000006327	TNFRSF12A	1.95211091	0.02613436
ENSG00000117143	UAP1	1.93366364	0.02778912
ENSG00000156642	NPTN	1.93440065	0.02800529
ENSG00000163697	APBB2	1.95832684	0.02848863
ENSG00000162616	DNAJB4	1.9411517	0.02867804
ENSG00000132329	RAMP1	1.96899533	0.02869506
ENSG00000104635	SLC39A14	1.93105847	0.02891801
ENSG00000052802	MSMO1	1.97230039	0.02897764
ENSG00000117519	CNN3	1.9058952	0.02919559
ENSG00000152990	GPR125	-2.0967778	0.02924709
ENSG00000196352	CD55	1.9238114	0.02934175
ENSG00000165323	FAT3	2.00120682	0.02982447
ENSG00000106105	GARS	1.89960819	0.02989086
ENSG00000182492	BGN	1.89388618	0.03008093
ENSG00000092841	MYL6	1.89193012	0.03024906
ENSG00000072110	ACTN1	1.8896353	0.03037721
ENSG00000131236	CAP1	1.88826534	0.03083029
ENSG00000135424	ITGA7	1.90531417	0.03096942
ENSG00000115380	EFEMP1	-1.872144	0.03198116
ENSG00000136052	SLC41A2	2.03094837	0.03254325
ENSG00000104332	SFRP1	-1.8639924	0.03405278
ENSG00000117152	RGS4	1.9636591	0.03413922
ENSG00000065413	ANKRD44	1.93808539	0.03475794
ENSG00000115129	TP53I3	1.88563054	0.03483187
ENSG00000164949	GEM	1.8466341	0.03627579
ENSG00000007933	FMO3	-1.8668129	0.03664315
ENSG00000070495	JMJD6	1.9411517	0.03666109
ENSG00000131981	LGALS3	-1.8278832	0.03684358
ENSG00000173846	PLK3	1.87764875	0.03706531
ENSG00000100596	SPTLC2	1.85354905	0.03718866
ENSG00000131711	MAP1B	1.81244936	0.0376664
ENSG00000117114	LPHN2	-1.9045313	0.03767691
ENSG00000144724	PTPRG	-1.8259538	0.03907041
ENSG00000049323	LTBP1	1.79396749	0.03916814
ENSG00000155304	HSPA13	1.80267254	0.03950001

ENSEMBL ID	HGNC ID	log ₂ FC RPF(M) / RPF(Ctrl)	RPF p-value
ENSG00000135047	CTSL1	-1.7963856	0.0401029
ENSG00000159363	ATP13A2	1.85868954	0.0405056
ENSG00000101871	MID1	-1.9027541	0.04064161
ENSG00000140682	TGFB1I1	1.79894207	0.04076587
ENSG00000181019	NQO1	-1.791716	0.04097796
ENSG00000165996	PTPLA	1.90508244	0.04142984
ENSG00000182054	IDH2	1.84983653	0.04158138
ENSG00000116016	EPAS1	-1.8287135	0.04167834
ENSG00000142089	IFITM3	-1.7819417	0.04248534
ENSG00000198018	ENTPD7	1.91258254	0.04261311
ENSG00000213949	ITGA1	1.77199225	0.04266232
ENSG00000166147	FBN1	1.75788331	0.04302846
ENSG00000163297	ANTXR2	-1.7756523	0.043628
ENSG00000104518	GSDMD	-1.8925912	0.04396197
ENSG00000188643	S100A16	1.76850759	0.04440116
ENSG00000087303	NID2	-1.7533753	0.0447325
ENSG00000138061	CYP1B1	-1.7576119	0.04494945
ENSG00000132432	SEC61G	1.77185056	0.04531989
ENSG00000169756	LIMS1	1.77771073	0.046255
ENSG00000060982	BCAT1	1.75102124	0.04686101
ENSG00000197622	CDC42SE1	1.85639408	0.0476266
ENSG00000100219	XBP1	1.74004838	0.04778771
ENSG00000136010	ALDH1L2	1.72220246	0.04905653
ENSG00000198856	OSTC	1.730833	0.04911071
ENSG00000151729	SLC25A4	1.81741633	0.04914682
ENSG00000152818	UTRN	-1.7488228	0.04936291
ENSG00000168268	NT5DC2	1.77572809	0.04980796

Table 2. Gene Signature with Altered Transcription

ENSEMBL ID	HGNC ID	log ₂ FC RNA(M) / RNA(Ctrl)	RNA p-value
ENSG00000107796	ACTA2	5.69	8.88E-07
ENSG00000049540	ELN	5.35	3.02E-06
ENSG00000106366	SERPINE1	4.45	5.47E-05
ENSG00000146674	IGFBP3	4.50	5.47E-05
ENSG00000123610	TNFAIP6	4.12	0.00017735

ENSEMBL ID	HGNC ID	log2FC RNA(M) / RNA(Ctrl)	RNA p-value
ENSG00000076706	MCAM	4.12	0.00026551
ENSG00000135074	ADAM19	4.00	0.00032434
ENSG00000140416	TPM1	3.72	0.00051342
ENSG00000130707	ASS1	3.81	0.00052605
ENSG00000107957	SH3PXD2A	3.70	0.00063215
ENSG00000187498	COL4A1	3.57	0.00080623
ENSG00000139211	AMIGO2	3.61	0.00085494
ENSG00000148848	ADAM12	3.58	0.00091659
ENSG00000176170	SPHK1	3.54	0.00104044
ENSG00000164761	TNFRSF11B	-3.41	0.00130277
ENSG00000149591	TAGLN	3.40	0.00131241
ENSG00000006016	CRLF1	3.48	0.00153949
ENSG00000177283	FZD8	3.46	0.00160105
ENSG00000079308	TNS1	3.33	0.0021064
ENSG00000099860	GADD45B	3.38	0.00218565
ENSG00000158186	MRAS	3.37	0.00231937
ENSG00000211455	STK38L	3.30	0.00244545
ENSG00000172986	GXYLT2	3.13	0.00364601
ENSG00000189184	PCDH18	-3.08	0.00368398
ENSG00000131435	PDLIM4	3.05	0.00385193
ENSG00000106211	HSPB1	3.01	0.00438968
ENSG00000170558	CDH2	2.98	0.00526545
ENSG00000171793	CTPS	3.00	0.00558329
ENSG00000139329	LUM	-2.90	0.00567798
ENSG00000113140	SPARC	2.88	0.00571107
ENSG00000162591	MEGF6	2.95	0.00596424
ENSG00000103489	XYLT1	2.97	0.00602805
ENSG00000115884	SDC1	2.88	0.0072085
ENSG00000163453	IGFBP7	2.72	0.00865684
ENSG00000134853	PDGFRA	-2.70	0.00918479
ENSG00000175899	A2M	-2.68	0.00920233
ENSG00000206190	ATP10A	2.74	0.00933257
ENSG00000163661	PTX3	-2.59	0.01218007
ENSG00000137809	ITGA11	2.65	0.01264373
ENSG00000137124	ALDH1B1	2.70	0.01272813
ENSG00000122786	CALD1	2.55	0.01365603
ENSG00000127241	MASP1	-2.57	0.01389162

ENSEMBL ID	HGNC ID	log2FC RNA(M) / RNA(Ctrl)	RNA p-value
ENSG00000110328	GALNTL4	2.66	0.01464927
ENSG00000196923	PDLIM7	2.53	0.01542871
ENSG00000025708	TYMP	2.59	0.01700225
ENSG00000152952	PLOD2	2.45	0.01776819
ENSG00000165072	MAMDC2	2.51	0.01829777
ENSG00000197594	ENPP1	2.53	0.01861632
ENSG00000135919	SERPINE2	2.40	0.01922323
ENSG00000135269	TES	2.47	0.01973346
ENSG00000104368	PLAT	-2.42	0.02041801
ENSG00000129038	LOXL1	2.40	0.02049761
ENSG00000128965	CHAC1	2.54	0.02099663
ENSG00000198832	RP3- 412A9.11.1	2.37	0.02162873
ENSG00000115902	SLC1A4	2.50	0.02246419
ENSG00000122694	GLIPR2	2.38	0.02572646
ENSG00000011422	PLAUR	2.33	0.02681845
ENSG00000107249	GLIS3	2.37	0.02732877
ENSG00000183010	PYCR1	2.28	0.02757996
ENSG00000111186	WNT5B	2.29	0.02786211
ENSG00000162520	SYNC	2.35	0.02977719
ENSG00000163697	APBB2	2.32	0.03055306
ENSG00000100234	TIMP3	2.22	0.03067964
ENSG00000106772	PRUNE2	2.31	0.03076204
ENSG00000168994	PXDC1	2.23	0.03111411
ENSG00000168268	NT5DC2	2.24	0.03137418
ENSG00000065054	SLC9A3R2	2.23	0.0315624
ENSG00000147224	PRPS1	2.29	0.03202045
ENSG00000164465	DCBLD1	2.22	0.03220594
ENSG00000130635	COL5A1	2.18	0.03264192
ENSG00000171617	ENC1	2.21	0.03379284
ENSG00000103257	SLC7A5	2.26	0.03503209
ENSG00000175183	CSRP2	2.26	0.0350484
ENSG00000154553	PDLIM3	2.21	0.03515586
ENSG00000133816	MICAL2	2.15	0.03618208
ENSG00000129116	PALLD	2.14	0.03660141
ENSG00000131711	MAP1B	2.20	0.03708651
ENSG00000134871	COL4A2	2.11	0.03797089
ENSG00000006327	TNFRSF12A	2.12	0.03816414

ENSEMBL ID	HGNC ID	log2FC RNA(M) / RNA(Ctrl)	RNA p-value
ENSG00000108821	COL1A1	2.10	0.03826416
ENSG00000187840	EIF4EBP1	2.22	0.03834287
ENSG00000165124	SVEP1	-2.14	0.03854141
ENSG00000105329	TGFB1	2.17	0.03974378
ENSG00000106799	TGFBR1	2.10	0.03990703
ENSG00000136010	ALDH1L2	2.18	0.04014228
ENSG00000147872	PLIN2	-2.10	0.04036349
ENSG00000171223	JUNB	2.10	0.04105211
ENSG00000121068	TBX2	-2.12	0.04242393
ENSG00000164574	GALNT10	2.11	0.04286773
ENSG00000165029	ABCA1	-2.06	0.044136
ENSG00000116016	EPAS1	-2.11	0.04452792
ENSG00000060982	BCAT1	2.09	0.04457308
ENSG00000131981	LGALS3	-2.06	0.04466942
ENSG00000100139	MICALL1	2.13	0.04562388
ENSG00000118523	CTGF	2.02	0.04636739
ENSG00000065911	MTHFD2	2.03	0.0469468
ENSG00000186340	THBS2	2.02	0.04716845
ENSG00000117410	ATP6V0B	2.03	0.04779432
ENSG00000120708	TGFBI	2.00	0.04817508
ENSG00000173457	PPP1R14B	2.02	0.0494489
ENSG00000174099	MSRB3	2.10	0.04957009
ENSG00000181104	F2R	-2.00	0.04989444
ENSG00000161638	ITGA5	1.99	0.05000656
ENSG00000099250	NRP1	-2.01	0.05023084
ENSG00000173540	GMPPB	2.15	0.0505729
ENSG00000144655	CSRN1P1	2.06	0.05123146
ENSG00000092964	DPYSL2	-2.02	0.05155695
ENSG00000135048	TMEM2	2.01	0.05188932
ENSG00000198853	RUSC2	2.03	0.05303444
ENSG00000132329	RAMP1	2.02	0.05315947
ENSG00000104635	SLC39A14	1.98	0.0536051
ENSG00000164932	CTHRC1	1.97	0.05405677
ENSG00000112769	LAMA4	-1.95	0.05420429
ENSG00000134285	FKBP11	2.07	0.05432854
ENSG00000145050	MANF	1.98	0.05494105
ENSG00000134668	SPOCD1	2.06	0.05509074

ENSEMBL ID	HGNC ID	log2FC RNA(M) / RNA(Ctrl)	RNA p-value
ENSG00000142552	RCN3	1.97	0.05510327
ENSG00000087008	ACOX3	2.00	0.05524263
ENSG00000018510	AGPS	2.06	0.05540727
ENSG00000163110	PDLIM5	2.00	0.05572685
ENSG00000165801	ARHGEF40	1.95	0.05592368
ENSG00000070669	ASNS	1.96	0.05660313
ENSG00000007933	FMO3	-2.01	0.05694724
ENSG00000156265	C21orf7	2.01	0.05708367
ENSG00000165996	PTPLA	2.00	0.05830933
ENSG00000090520	DNAJB11	2.00	0.05899164
ENSG00000152377	SPOCK1	1.95	0.05984535
ENSG00000125753	VASP	1.95	0.06037993
ENSG00000155304	HSPA13	1.92	0.06156215
ENSG00000128590	DNAJB9	1.97	0.06179333
ENSG00000151327	FAM177A1	1.98	0.06266866
ENSG00000138735	PDE5A	-1.89	0.06344404
ENSG00000109861	CTSC	-1.90	0.06413627
ENSG00000163513	TGFBR2	-1.91	0.06421189
ENSG00000151729	SLC25A4	2.01	0.06432535
ENSG00000100600	LGMN	1.92	0.06502239
ENSG00000113361	CDH6	1.91	0.06517739
ENSG00000100596	SPTLC2	1.94	0.06546002
ENSG00000144724	PTPRG	-1.92	0.06605951
ENSG00000183963	SMTN	1.88	0.06648929
ENSG00000147852	VLDLR	1.99	0.067094
ENSG00000131018	SYNE1	1.86	0.06722868
ENSG00000198833	UBE2J1	1.92	0.06787646
ENSG00000169756	LIMS1	1.99	0.06804636
ENSG00000185000	DGAT1	1.94	0.06842024
ENSG00000112096	SOD2	-1.90	0.06877769
ENSG00000052802	MSMO1	1.93	0.06924478
ENSG00000100889	PCK2	1.88	0.06938457
ENSG00000136052	SLC41A2	1.96	0.07000892
ENSG00000127334	DYRK2	1.91	0.07030249
ENSG00000182054	IDH2	1.87	0.07097019
ENSG00000122729	ACO1	-1.86	0.07120481
ENSG00000087303	NID2	-1.87	0.07131478

ENSEMBL ID	HGNC ID	log2FC RNA(M) / RNA(Ctrl)	RNA p-value
ENSG00000162616	DNAJB4	1.92	0.07154524
ENSG00000092841	MYL6	1.81	0.07161024
ENSG00000156642	NPTN	1.84	0.07238219
ENSG00000214517	PPME1	1.93	0.07294207
ENSG00000101825	MXRA5	1.92	0.07301266
ENSG00000198467	TPM2	1.80	0.07356735
ENSG00000144810	COL8A1	1.80	0.07628075
ENSG00000181019	NQO1	-1.80	0.07693505
ENSG00000134030	CTIF	1.88	0.07713563
ENSG00000115380	EFEMP1	-1.78	0.07802011
ENSG00000196072	BLOC1S2	1.86	0.07837829
ENSG00000118508	RAB32	1.78	0.07878147
ENSG00000159363	ATP13A2	1.83	0.07900456
ENSG00000144746	ARL6IP5	-1.87	0.08073223
ENSG00000137331	IER3	1.80	0.08080819
ENSG00000142871	CYR61	1.77	0.08166504
ENSG00000154122	ANKH	1.81	0.0822588
ENSG00000073712	FERMT2	1.76	0.08290623
ENSG00000136802	LRRC8A	1.77	0.08363724
ENSG00000108106	UBE2S	1.84	0.08470435
ENSG00000114850	SSR3	1.75	0.08485595
ENSG00000239672	NME1	1.82	0.08581195
ENSG00000140682	TGFB1I1	1.79	0.08583217
ENSG00000196576	PLXNB2	-1.73	0.0874579
ENSG00000115129	TP53I3	1.75	0.08790906
ENSG00000070371	CLTCL1	1.80	0.08850262
ENSG00000126524	SBDS	1.77	0.08865039
ENSG00000198018	ENTPD7	1.81	0.08984177
ENSG00000130513	GDF15	-1.72	0.09015532
ENSG00000166888	STAT6	-1.79	0.09068891
ENSG00000127418	FGFRL1	1.79	0.09101936
ENSG00000167996	FTH1	-1.69	0.09230109
ENSG00000101955	SRPX	-1.79	0.09230885
ENSG00000115963	RND3	-1.70	0.09246072
ENSG00000072110	ACTN1	1.70	0.09255263
ENSG00000162704	ARPC5	1.69	0.09583256
ENSG00000127863	TNFRSF19	-1.73	0.0958679

ENSEMBL ID	HGNC ID	log2FC RNA(M) / RNA(Ctrl)	RNA p-value
ENSG00000161091	MFSD12	1.74	0.09725927
ENSG00000198542	ITGBL1	1.68	0.09754718
ENSG00000196352	CD55	1.75	0.09941386
ENSG00000116260	QSOX1	-1.65	0.09954164

Table 3A. Gene Signature (1 replicate) with Altered Translational Efficiency

ENSEMBL ID	HGNC ID	TE	BABEL p-value
ENSG00000163661	PTX3	-2.7413506	4.80E-10
ENSG00000182752	PAPPA	-2.031799	1.15E-07
ENSG00000173230	GOLGB1	-1.4747045	3.65E-07
ENSG00000008441	NFIX	-1.6325086	3.25E-06
ENSG00000108055	SMC3	-1.5202935	6.61E-06
ENSG00000112902	SEMA5A	-1.5159033	1.90E-05
ENSG00000168724	DNAJC21	-1.6152699	2.36E-05
ENSG00000138735	PDE5A	-2.1383717	3.96E-05
ENSG00000102908	NFAT5	-1.2539722	8.13E-05
ENSG00000100644	HIF1A	1.29634616	8.38E-05
ENSG00000166833	NAV2	-1.4353025	0.000100353
ENSG00000136542	GALNT5	-1.2478982	0.000102241
ENSG00000138385	SSB	-1.276029	0.000112036
ENSG00000047410	TPR	-1.1002148	0.000115647
ENSG00000100815	TRIP11	-1.253114	0.000115691
ENSG00000135905	DOCK10	1.41487073	0.000116804
ENSG00000187446	AC012652.1.1	1.4975647	0.000169735
ENSG00000108106	UBE2S	-1.9808659	0.000238753
ENSG00000108107	RPL28	1.23778107	0.00027262
ENSG00000197442	MAP3K5	-2.166021	0.000273661
ENSG00000198121	LPAR1	-1.8734643	0.000278857
ENSG00000101871	MID1	-1.6869885	0.000281936
ENSG00000138081	FBXO11	-1.2032936	0.000367153
ENSG00000111011	RSRC2	-1.2385789	0.000370672
ENSG00000127863	TNFRSF19	-1.6203293	0.000384143
ENSG00000118058	MLL	-1.1124833	0.000407516
ENSG00000197321	SVIL	-1.4581886	0.000458188
ENSG00000119414	PPP6C	1.16868462	0.000575527

ENSEMBL ID	HGNC ID	TE	BABEL p-value
ENSG00000102241	HTATSF1	-1.1116943	0.000665853
ENSG00000119408	NEK6	-1.5858303	0.000679474
ENSG00000144674	GOLGA4	-1.1337996	0.000700442
ENSG00000008952	SEC62	-1.0264219	0.000755397
ENSG00000005339	CREBBP	-1.1959303	0.000899463
ENSG00000101972	STAG2	-1.1875122	0.000970595
ENSG00000173812	EIF1	0.74705692	0.001004834
ENSG00000168172	HOOK3	1.12108562	0.001093288
ENSG00000151067	CACNA1C	-1.4996202	0.001110948
ENSG00000136819	C9orf78	-1.1803341	0.001114894
ENSG00000136244	IL6	1.40808063	0.0011622
ENSG00000137831	UACA	-1.0308231	0.001200576
ENSG00000126777	KTN1	-1.1586794	0.00141984
ENSG00000085224	ATRX	-0.9250079	0.00147298
ENSG00000181722	ZBTB20	-1.3561731	0.001599887
ENSG00000151914	DST	-1.2345227	0.001733711
ENSG00000171793	CTPS	1.09773458	0.001903283
ENSG00000164292	RHOBTB3	-1.7514562	0.002336358
ENSG00000018510	AGPS	-1.1321327	0.002431067
ENSG00000148218	ALAD	-1.1916613	0.002478664
ENSG00000170027	YWHAG	0.85191988	0.002717768
ENSG00000101040	ZMYND8	-1.3571925	0.002752192
ENSG00000093167	LRRFIP2	-1.2205154	0.002887031
ENSG00000176105	YES1	-1.2351384	0.002892828
ENSG00000026025	VIM	1.08993981	0.003294757
ENSG00000152818	UTRN	-1.213851	0.003490212
ENSG00000167658	EEF2	1.75181457	0.003801433
ENSG00000153922	CHD1	-0.9896109	0.004445999
ENSG00000147677	EIF3H	1.33689856	0.004466087
ENSG00000135316	SYNCRIP	-0.8986489	0.00454662
ENSG00000105373	GLTSCR2	1.93251408	0.004658713
ENSG00000151461	UPF2	-1.0525428	0.004708017
ENSG00000143374	TARS2	-1.2193317	0.00478043
ENSG00000114439	BBX	-1.0539237	0.004795963
ENSG00000133318	RTN3	0.84613845	0.005442031
ENSG00000049618	ARID1B	-1.0178473	0.005566074
ENSG00000197170	PSMD12	-0.852153	0.005683822

ENSEMBL ID	HGNC ID	TE	BABEL p-value
ENSG00000158615	PPP1R15B	1.23383146	0.005756279
ENSG00000153774	CFDP1	-1.1005412	0.005942768
ENSG00000188641	DPYD	-1.2580397	0.006287312
ENSG00000136770	DNAJC1	1.03702109	0.00652172
ENSG00000121940	CLCC1	-1.0098191	0.006536917
ENSG00000100154	TTC28	-1.0182753	0.006608779
ENSG00000113161	HMGCR	0.94424256	0.006719302
ENSG00000185068	GTF2H5	-1.0829484	0.006949215
ENSG00000005955	GGNBP2	-0.8936361	0.00729981
ENSG00000080603	SRCAP	-0.9154367	0.00733119
ENSG00000001497	LAS1L	-1.0314198	0.007743017
ENSG00000138061	CYP1B1	-0.8401242	0.007885734
ENSG00000164715	LMTK2	1.29018489	0.007969877
ENSG00000143545	RAB13	-1.0411451	0.007980361
ENSG00000026508	CD44	0.68665057	0.008092007
ENSG00000117054	ACADM	-1.0717708	0.008211763
ENSG00000092201	SUPT16H	-0.8692281	0.008387717
ENSG00000110367	DDX6	-0.879027	0.009106782
ENSG00000143924	EML4	-0.9889802	0.0092973
ENSG00000164294	GPX8	0.89184246	0.009363418
ENSG00000241685	ARPC1A	1.50567956	0.009392258
ENSG00000164548	TRA2A	0.93385802	0.00952787
ENSG00000138688	KIAA1109	-1.0676173	0.009640977
ENSG00000100528	CNIH	0.8304318	0.009829269
ENSG00000127603	MACF1	-0.943354	0.009856068
ENSG00000115652	UXS1	-1.0666256	0.0099403
ENSG00000118523	CTGF	0.60443779	0.010005062
ENSG00000182240	BACE2	-1.222017	0.01056114
ENSG00000119004	CYP20A1	-0.9860759	0.011089198
ENSG00000136146	MED4	-0.992898	0.011091961
ENSG00000124786	SLC35B3	-1.1042075	0.01133021
ENSG00000077097	TOP2B	-0.8579402	0.011387162
ENSG00000185418	TARSL2	-1.1697798	0.011458991
ENSG00000125257	ABCC4	-1.2299933	0.011565549
ENSG00000104419	NDRG1	0.84121291	0.012271036
ENSG00000143995	MEIS1	-0.9850066	0.012417371
ENSG00000112972	HMGCS1	1.04256421	0.012443231

ENSEMBL ID	HGNC ID	TE	BABEL p-value
ENSG00000164830	OXR1	-0.9587293	0.013955428
ENSG00000067900	ROCK1	-0.8067914	0.014145159
ENSG00000122406	RPL5	1.49266971	0.01448355
ENSG00000137331	IER3	1.04042945	0.014705463
ENSG00000136521	NDUFB5	1.18012704	0.014891357
ENSG00000020577	SAMD4A	-1.0959325	0.015243102
ENSG00000180398	MCFD2	-0.7756358	0.015377781
ENSG00000169100	SLC25A6	1.34004011	0.01579095
ENSG00000135047	CTSL1	-0.8906545	0.016073112
ENSG00000140497	SCAMP2	0.80895096	0.016513709
ENSG00000144029	MRPS5	-0.9350724	0.016712818
ENSG00000132561	MATN2	0.8592783	0.016779427
ENSG00000117523	PRRC2C	-0.5599625	0.016819351
ENSG00000118689	FOXO3	-1.0512663	0.017059692
ENSG00000080345	RIF1	-0.7907628	0.017236668
ENSG00000105894	PTN	-0.8244564	0.017342288
ENSG00000213949	ITGA1	0.7423431	0.017353023
ENSG00000147548	WHSC1L1	-0.9352328	0.018230486
ENSG00000163946	FAM208A	-1.0185676	0.018771848
ENSG00000106261	ZKSCAN1	-1.1775938	0.018843712
ENSG00000105193	RPS16	1.10211287	0.019637409
ENSG00000139726	DENR	-0.8974717	0.019791894
ENSG00000136205	TNS3	-0.8280863	0.020242812
ENSG00000145555	MYO10	-0.8048733	0.021421007
ENSG00000152022	LIX1L	-0.8124639	0.0218285
ENSG00000142669	SH3BGRL3	0.55526839	0.021867346
ENSG00000141367	CLTC	0.77612663	0.022218177
ENSG00000162804	SNED1	-0.8185651	0.022559589
ENSG00000104408	EIF3E	1.32647476	0.022886033
ENSG00000110344	UBE4A	0.87653113	0.023224777
ENSG00000213064	SFT2D2	-0.755902	0.02323942
ENSG00000225921	NOL7	-0.8815021	0.023433926
ENSG00000138593	SECISBP2L	-0.7761533	0.023443449
ENSG00000065150	IPO5	0.99464329	0.02369603
ENSG00000115806	GORASP2	0.67835655	0.023794157
ENSG00000100129	EIF3L	1.62687625	0.024524439
ENSG00000196914	ARHGEF12	-1.1675105	0.024891831

ENSEMBL ID	HGNC ID	TE	BABEL p-value
ENSG00000204217	BMPR2	0.69246757	0.025438073
ENSG00000109390	NDUFC1	-0.8952606	0.025491324
ENSG00000196367	TRRAP	-0.7594148	0.025604694
ENSG00000144036	EXOC6B	-1.0757792	0.026021627
ENSG00000169946	ZFPM2	-1.0479271	0.02631718
ENSG00000137509	PRCP	0.87652803	0.026354393
ENSG00000116747	TROVE2	-0.726508	0.026552043
ENSG00000060339	CCAR1	-0.6997234	0.026685169
ENSG00000146247	PHIP	-0.8332369	0.026820782
ENSG00000084652	TXLNA	0.73915572	0.027068209
ENSG00000130741	EIF2S3	1.05466806	0.027133332
ENSG00000150961	SEC24D	0.62765591	0.027298647
ENSG00000187240	DYNC2H1	-1.06679	0.027761828
ENSG00000146433	TMEM181	-1.0414552	0.027796354
ENSG00000179295	PTPN11	-0.9140165	0.0288127
ENSG00000113282	CLINT1	-0.711755	0.028973307
ENSG00000171988	JMJD1C	-0.7790863	0.028992676
ENSG00000090054	SPTLC1	0.83148984	0.029632806
ENSG00000197579	TOPORS	-0.8458666	0.029699333
ENSG00000142864	SERBP1	-0.7646724	0.029899373
ENSG00000034713	GABARAPL2	0.68823059	0.029970584
ENSG00000090520	DNAJB11	-0.7810914	0.0300775
ENSG00000166855	CLPX	-0.8488647	0.030097263
ENSG00000135046	ANXA1	0.86394566	0.030138772
ENSG00000196935	SRGAP1	-0.8040381	0.030202802
ENSG00000106538	RARRES2	-0.9688629	0.030236136
ENSG00000099783	HNRNPM	-0.5956752	0.030351889
ENSG00000141458	NPC1	0.74822754	0.030844322
ENSG00000177606	JUN	0.71000106	0.030935017
ENSG00000097021	ACOT7	0.80866661	0.031064001
ENSG00000197111	PCBP2	0.68049208	0.031080336
ENSG00000100316	RPL3	1.67662485	0.03114172
ENSG00000197056	ZMYM1	-0.9388698	0.031399769
ENSG00000198042	MAK16	-0.8940405	0.031501821
ENSG00000148396	SEC16A	0.58297597	0.031545262
ENSG00000260916	CCPG1	-0.7102265	0.031823744
ENSG00000188191	PRKAR1B	-1.027972	0.032129214

ENSEMBL ID	HGNC ID	TE	BABEL p-value
ENSG00000198146	ZNF770	-0.8146313	0.032249599
ENSG00000107949	BCCIP	-0.8019017	0.032357337
ENSG00000140443	IGF1R	-0.8736668	0.03237335
ENSG00000169738	DCXR	-1.0509859	0.032502003
ENSG00000107317	PTGDS	-0.5958188	0.032502353
ENSG00000101079	NDRG3	-0.9225847	0.032934071
ENSG00000138095	LRPPRC	0.84253686	0.03303189
ENSG00000136153	LMO7	-0.9201728	0.033102447
ENSG00000146109	ABT1	-1.1549145	0.033713746
ENSG00000136156	ITM2B	0.61379782	0.033862692
ENSG00000142599	RERE	-0.8238852	0.033895821
ENSG00000154767	XPC	-0.7641246	0.034070785
ENSG00000107819	SFXN3	-0.8139634	0.034640721
ENSG00000107951	MTPAP	-0.8791984	0.034849044
ENSG00000107862	GBF1	0.63999556	0.03505938
ENSG00000113194	FAF2	0.71045312	0.035949786
ENSG00000136485	DCAF7	-0.6757873	0.036070299
ENSG00000189241	TSPYL1	0.7062045	0.036103289
ENSG00000138293	NCOA4	0.88765663	0.036332045
ENSG00000070371	CLTCL1	1.10443252	0.036773077
ENSG00000172954	LCLAT1	-0.8524558	0.036863783
ENSG00000196683	TOMM7	1.3075298	0.037353358
ENSG00000113048	MRPS27	1.0170333	0.037417425
ENSG00000113163	COL4A3BP	-0.7578152	0.037499686
ENSG00000170242	USP47	-0.8028916	0.037635249
ENSG00000127914	AKAP9	-0.6744918	0.038026974
ENSG00000141279	NPEPPS	-0.8812721	0.038230406
ENSG00000134313	KIDINS220	-0.7488098	0.038375301
ENSG00000023287	RB1CC1	-0.8487863	0.03971745
ENSG00000118181	RPS25	1.58310188	0.040307595
ENSG00000029363	BCLAF1	-0.6873768	0.041239905
ENSG00000152332	UHMK1	-0.9690309	0.041660214
ENSG00000170871	KIAA0232	-0.9350598	0.041663582
ENSG00000132792	CTNNBL1	-0.7352665	0.041674551
ENSG00000013441	CLK1	-0.9194947	0.042371585
ENSG00000169972	PUSL1	-1.0350875	0.043097174
ENSG00000128591	FLNC	0.6848822	0.043354638

ENSEMBL ID	HGNC ID	TE	BABEL p-value
ENSG00000135931	ARMC9	-0.944761	0.043425047
ENSG00000128965	CHAC1	0.79389693	0.043641321
ENSG00000131171	SH3BGRL	0.59583837	0.043953482
ENSG00000138246	DNAJC13	-0.7333811	0.044369031
ENSG00000150593	PDCD4	-0.6243741	0.044464047
ENSG00000129292	PHF20L1	-0.7749239	0.044871229
ENSG00000104067	TJP1	-0.642144	0.044883401
ENSG00000171858	RPS21	1.35698428	0.044918379
ENSG00000132688	NES	0.54614566	0.045111901
ENSG00000144713	RPL32	1.55450933	0.045432473
ENSG00000103222	ABCC1	-0.7216554	0.045567887
ENSG00000078699	CBFA2T2	-0.9232918	0.045900617
ENSG00000179387	ELMOD2	0.77525851	0.046344506
ENSG00000131389	SLC6A6	-0.6368225	0.04645475
ENSG00000122126	OCRL	0.69883954	0.046768488
ENSG00000115364	MRPL19	-0.711517	0.047054746
ENSG00000115355	CCDC88A	-0.6782108	0.047073125
ENSG00000048649	RSF1	-0.6764684	0.047138641
ENSG00000156256	USP16	-0.8521086	0.047196289
ENSG00000188739	RBM34	-0.8515963	0.047979658
ENSG00000173726	TOMM20	0.94716704	0.048911642
ENSG00000135720	DYNC1LI2	-0.713868	0.049159881
ENSG00000126945	HNRNPH2	0.68135224	0.049532841
ENSG00000105887	MTPN	0.61407664	0.049580426
ENSG00000141753	IGFBP4	-0.6583794	0.049591031

Table 3B. Gene Signature (5 Biological Replicates) with Altered Translational Efficiency

ENSEMBL ID	HGNC	Log2 TE	BABEL p-value
ENSG00000250479	CHCHD10	2.51943	7.09E-12
ENSG00000118473	SGIP1	-2.76752	1.51E-11
ENSG00000150636	CCDC102B	-2.97708	1.62E-11
ENSG00000122966	CIT	-3.08447	2.07E-11
ENSG00000108107	RPL28	1.869423	1.81E-10
ENSG00000143632	ACTA1	-2.01787	8.62E-10

ENSEMBL ID	HGNC	Log2 TE	BABEL p-value
ENSG00000170681	MURC	3.678236	6.92E-09
ENSG00000105664	COMP	1.317102	7.17E-09
ENSG00000173641	HSPB7	1.292289	1.04E-08
ENSG00000054654	SYNE2	-1.79281	3.79E-08
ENSG00000132510	KDM6B	1.602992	5.15E-08
ENSG00000132170	PPARG	-3.52005	6.11E-08
ENSG00000119508	NR4A3	-1.93064	7.13E-08
ENSG00000143850	PLEKHA6	-2.06925	9.33E-08
ENSG00000175505	CLCF1	1.594058	1.45E-07
ENSG00000132031	MATN3	1.340341	1.85E-07
ENSG00000169100	SLC25A6	1.909395	2.02E-07
ENSG00000196616	ADH1B	-2.79469	2.8E-07
ENSG00000197111	PCBP2	1.29269	4.67E-07
ENSG00000171408	PDE7B	-2.78241	5.78E-07
ENSG00000137331	IER3	1.702028	6.76E-07
ENSG00000049130	KITLG	-2.02679	9.38E-07
ENSG00000139734	DIAPH3	1.658951	1.01E-06
ENSG00000167658	EEF2	2.322838	1.61E-06
ENSG00000077312	SNRPA	1.461712	1.69E-06
ENSG00000132122	SPATA6	-1.92545	3.83E-06
ENSG00000124496	TRERF1	-1.89102	4.18E-06
ENSG00000162599	NFIA	-1.2934	5.92E-06
ENSG00000222009	BTBD19	1.524069	7.57E-06
ENSG00000135480	KRT7	2.000334	8.38E-06
ENSG00000105193	RPS16	1.989185	8.68E-06
ENSG00000138675	FGF5	-1.16325	1.03E-05
ENSG00000108551	RASD1	1.836079	1.15E-05
ENSG00000066279	ASPM	-2.175	1.26E-05
ENSG00000174279	EVX2	2.887275	1.64E-05
ENSG00000145029	NICN1	1.402987	1.84E-05
ENSG00000139354	GAS2L3	-2.34161	1.96E-05
ENSG00000106415	GLCCI1	-1.84191	0.000026
ENSG00000197442	MAP3K5	-2.0153	2.66E-05
ENSG00000094963	FMO2	-2.14622	0.000031

ENSEMBL ID	HGNC	Log2 TE	BABEL p-value
ENSG00000146411	SLC2A12	-1.9708	4.09E-05
ENSG00000115267	IFIH1	-1.41561	4.77E-05
ENSG00000124749	COL21A1	-1.44522	5.07E-05
ENSG00000144218	AFF3	1.146175	5.18E-05
ENSG00000154262	ABCA6	-2.16651	5.24E-05
ENSG00000233927	RPS28	1.937376	0.000055
ENSG00000136244	IL6	1.369	7.79E-05
ENSG00000138735	PDE5A	-1.59351	8.35E-05
ENSG00000187605	TET3	1.540855	8.43E-05
ENSG00000125869	LAMP5	1.298433	8.54E-05
ENSG00000146066	HIGD2A	1.195557	9.35E-05
ENSG00000189367	KIAA0408	-1.28312	0.000103
ENSG00000157168	NRG1	1.162042	0.000106
ENSG00000155962	CLIC2	-2.39184	0.000106
ENSG00000188846	RPL14	1.745447	0.000125
ENSG00000137033	IL33	-2.48811	0.000128
ENSG00000129226	CD68	1.047995	0.00014
ENSG00000171793	CTPS	1.080465	0.000147
ENSG00000152154	TMEM178	2.01577	0.000152
ENSG00000117724	CENPF	-1.23716	0.000153
ENSG00000112276	BVES	1.172691	0.00016
ENSG00000131747	TOP2A	-1.31413	0.000181
ENSG00000134419	RPS15A	2.194763	0.000184
ENSG00000128016	ZFP36	1.187628	0.000188
ENSG00000135390	ATP5G2	1.380759	0.000188
ENSG00000049192	ADAMTS6	2.862797	0.000209
ENSG00000105373	GLTSCR2	2.066729	0.000211
ENSG00000115268	RPS15	1.300314	0.000224
ENSG00000124406	ATP8A1	-2.12205	0.000228
ENSG00000168769	TET2	1.37689	0.000237
ENSG00000166441	RPL27A	2.090323	0.000238
ENSG00000168916	ZNF608	-1.58959	0.00024
ENSG00000147677	EIF3H	1.328509	0.000247
ENSG00000198121	LPAR1	-1.44836	0.000253

ENSEMBL ID	HGNC	Log2 TE	BABEL p-value
ENSG00000198932	GPRASP1	-1.23921	0.000259
ENSG00000104408	EIF3E	1.469386	0.000287
ENSG00000198848	CES1	1.084318	0.000306
ENSG00000133321	RARRES3	-1.99537	0.000312
ENSG00000122406	RPL5	1.592522	0.000392
ENSG00000070756	PABPC1	1.207052	0.000397
ENSG00000135486	HNRNPA1	1.08564	0.000404
ENSG00000137628	DDX60	-1.21094	0.000417
ENSG00000164292	RHOBTB3	-1.52866	0.000434
ENSG00000133687	TMTC1	-1.90695	0.000441
ENSG00000204628	GNB2L1	1.732459	0.000477
ENSG00000171858	RPS21	1.846388	0.00048
ENSG00000182899	RPL35A	1.626674	0.000484
ENSG00000144713	RPL32	1.915935	0.000484
ENSG00000198963	RORB	-1.76565	0.000505
ENSG00000109846	CRYAB	1.107197	0.000535
ENSG00000177600	RPLP2	1.69427	0.000538
ENSG00000156508	EEF1A1	1.850575	0.000545
ENSG00000152818	UTRN	-1.08497	0.00056
ENSG00000244242	IFITM10	-1.29903	0.000572
ENSG00000154654	NCAM2	-1.13535	0.000618
ENSG00000129422	MTUS1	-1.14926	0.000642
ENSG00000155621	C9orf85	1.111901	0.000643
ENSG00000097021	ACOT7	1.010887	0.000688
ENSG00000050426	LETMD1	1.270162	0.000706
ENSG00000229117	RPL41	1.473651	0.000712
ENSG00000100316	RPL3	1.780059	0.000743
ENSG00000164342	TLR3	-1.37101	0.000771
ENSG00000169239	CA5B	-1.79072	0.000794
ENSG00000170537	TMC7	1.744989	0.000811
ENSG00000185033	SEMA4B	-1.41345	0.000833
ENSG00000147403	RPL10	1.690799	0.000923
ENSG00000187240	DYNC2H1	-1.22778	0.000952
ENSG00000125375	ATP5S	1.075005	0.000968

ENSEMBL ID	HGNC	Log2 TE	BABEL p-value
ENSG00000184661	CDCA2	-1.57267	0.001062
ENSG00000171517	LPAR3	-2.38727	0.001079
ENSG00000026025	VIM	1.001738	0.001082
ENSG00000109452	INPP4B	-1.56908	0.001104
ENSG00000197756	RPL37A	2.028577	0.001124
ENSG00000110700	RPS13	1.538546	0.00115
ENSG00000088367	EPB41L1	1.018214	0.001179
ENSG00000152527	PLEKHH2	-1.38949	0.00121
ENSG00000064042	LIMCH1	-1.22217	0.001224
ENSG00000100575	TIMM9	1.113218	0.001229
ENSG00000121957	GPSM2	-1.40775	0.001239
ENSG00000118898	PPL	-1.53977	0.00126
ENSG00000197958	RPL12	1.878187	0.001271
ENSG00000087586	AURKA	-1.73649	0.001312
ENSG00000118181	RPS25	1.697268	0.001415
ENSG00000089351	GRAMD1A	1.068576	0.001647
ENSG00000129675	ARHGEF6	-1.10045	0.001654
ENSG00000171425	ZNF581	1.450193	0.001733
ENSG00000175315	CST6	1.241221	0.001753
ENSG00000138182	KIF20B	-1.28121	0.001776
ENSG00000139209	SLC38A4	-1.14245	0.00181
ENSG00000163219	ARHGAP25	1.165653	0.00184
ENSG00000125967	NECAB3	1.402793	0.001844
ENSG00000148303	RPL7A	1.650224	0.001928
ENSG00000125744	RTN2	-1.14254	0.001929
ENSG00000189180	ZNF33A	-1.05822	0.002062
ENSG00000170889	RPS9	1.6139	0.002078
ENSG00000063177	RPL18	1.611619	0.002079
ENSG00000128394	APOBEC3F	-1.12736	0.002113
ENSG00000100784	RPS6KA5	-1.43095	0.002244
ENSG00000174748	RPL15	1.253839	0.002262
ENSG00000197959	DNM3	-1.16065	0.002439
ENSG00000182405	PGBD4	1.055482	0.002447
ENSG00000109475	RPL34	1.607573	0.002479

ENSEMBL ID	HGNC	Log2 TE	BABEL p-value
ENSG00000176020	AMIGO3	-1.31388	0.002484
ENSG00000138336	TET1	1.853326	0.00249
ENSG00000185437	SH3BGR	1.121445	0.002612
ENSG00000163931	TKT	1.142358	0.002636
ENSG00000146757	ZNF92	-1.23798	0.002664
ENSG00000198182	ZNF607	-1.10762	0.002724
ENSG00000171863	RPS7	1.549128	0.002765
ENSG00000241685	ARPC1A	1.287518	0.00277
ENSG00000088756	ARHGAP28	-1.02406	0.002782
ENSG00000156482	RPL30	1.622306	0.002854
ENSG00000153707	PTPRD	-1.18636	0.002863
ENSG00000128849	CGNL1	-1.02286	0.002892
ENSG00000231500	RPS18	2.003068	0.002929
ENSG00000145592	RPL37	1.808138	0.003002
ENSG00000105640	RPL18A	1.669484	0.003009
ENSG00000127863	TNFRSF19	-1.09317	0.003027
ENSG00000139372	TDG	1.024894	0.003073
ENSG00000104205	SGK3	-1.01117	0.003075
ENSG00000170962	PDGFD	-1.02886	0.003142
ENSG00000162244	RPL29	1.588943	0.003207
ENSG00000172053	QARS	1.361835	0.003231
ENSG00000151773	CCDC122	-1.00405	0.003236
ENSG00000131469	RPL27	1.518506	0.003252
ENSG00000174444	RPL4	1.688063	0.003284
ENSG00000120156	TEK	-1.65221	0.003299
ENSG00000100129	EIF3L	1.429034	0.003336
ENSG00000138326	RPS24	1.443788	0.003411
ENSG00000113272	THG1L	1.155349	0.003452
ENSG00000169330	KIAA1024	2.216185	0.003475
ENSG00000175390	EIF3F	1.290244	0.003503
ENSG00000169231	THBS3	1.187649	0.00351
ENSG00000183765	CHEK2	-1.45694	0.003579
ENSG00000134532	SOX5	-1.66694	0.003634
ENSG00000130255	RPL36	1.410987	0.003671

ENSEMBL ID	HGNC	Log2 TE	BABEL p-value
ENSG00000184702	5-Sep	1.209162	0.003794
ENSG00000104529	EEF1D	1.61919	0.003976
ENSG00000164587	RPS14	1.683901	0.003995
ENSG00000161016	RPL8	1.352789	0.00415
ENSG00000109674	NEIL3	-2.63009	0.004174
ENSG00000138449	SLC40A1	1.114207	0.004183
ENSG00000142541	RPL13A	1.796824	0.004223
ENSG00000184588	PDE4B	-2.02072	0.004281
ENSG00000155816	FMN2	-1.24637	0.004306
ENSG00000163584	RPL22L1	1.012263	0.004479
ENSG00000089289	IGBP1	1.26553	0.004627
ENSG00000138356	AOX1	-1.14295	0.004762
ENSG00000108932	SLC16A6	-1.3497	0.004807
ENSG00000204520	MICA	1.143447	0.004864
ENSG00000142937	RPS8	1.610846	0.004896
ENSG00000113070	HBEGF	1.088046	0.004959
ENSG00000145741	BTF3	1.130214	0.005066
ENSG00000248905	FMN1	-1.90638	0.00514
ENSG00000083845	RPS5	1.532714	0.005184
ENSG00000198918	RPL39	1.535891	0.005218
ENSG00000105372	RPS19	1.549677	0.005369
ENSG00000163682	RPL9	1.613142	0.005382
ENSG00000142676	RPL11	1.396155	0.005383
ENSG00000165171	WBSCR27	1.116069	0.005577
ENSG00000111640	GAPDH	1.262613	0.005634
ENSG00000117543	DPH5	1.114344	0.005946
ENSG00000137154	RPS6	1.598032	0.005993
ENSG00000143947	RPS27A	1.275373	0.006092
ENSG00000170275	CRTAP	1.021832	0.006428
ENSG00000075651	PLD1	-1.03241	0.006527
ENSG00000111678	C12orf57	1.172975	0.006548
ENSG00000172809	RPL38	1.353068	0.006956
ENSG00000130741	EIF2S3	1.067385	0.00704
ENSG00000125691	RPL23	1.603975	0.007277

ENSEMBL ID	HGNC	Log2 TE	BABEL p-value
ENSG00000196683	TOMM7	1.126267	0.007329
ENSG00000178401	DNAJC22	-1.18938	0.00741
ENSG00000136040	PLXNC1	-1.0996	0.007562
ENSG00000090376	IRAK3	-1.01671	0.007597
ENSG00000198755	RPL10A	1.509807	0.007598
ENSG00000114391	RPL24	1.304469	0.007667
ENSG00000071082	RPL31	1.599814	0.007781
ENSG00000213741	RPS29	1.570822	0.007815
ENSG00000186184	POLR1D	1.229911	0.007848
ENSG00000161970	RPL26	1.654815	0.007913
ENSG00000198467	TPM2	1.077188	0.008168
ENSG00000115392	FANCL	-1.23269	0.008213
ENSG00000137818	RPLP1	1.460829	0.008299
ENSG00000187987	ZSCAN23	-1.04823	0.008574
ENSG00000197728	RPS26	1.090113	0.008582
ENSG00000198034	RPS4X	1.533681	0.008759
ENSG00000198242	RPL23A	1.394537	0.009726
ENSG00000204397	CARD16	-1.07621	0.009841
ENSG00000149273	RPS3	1.517901	0.010391
ENSG00000117461	PIK3R3	-1.0571	0.010612
ENSG00000184254	ALDH1A3	-1.32678	0.010897
ENSG00000176438	C14orf49	-1.04426	0.010906
ENSG00000115594	IL1R1	-1.00334	0.011677
ENSG00000137819	PAQR5	-1.44514	0.013098
ENSG00000053747	LAMA3	-1.23334	0.014709
ENSG00000257949	TEN1	1.361387	0.014841
ENSG00000175084	DES	1.20409	0.015111
ENSG00000101639	CEP192	-1.19163	0.015211
ENSG00000196531	NACA	1.082663	0.01541
ENSG00000140988	RPS2	1.606507	0.015474
ENSG00000156467	UQCRB	1.047919	0.015711
ENSG00000047648	ARHGAP6	-1.08196	0.015848
ENSG00000164116	GUCY1A3	-1.93203	0.015978
ENSG00000180667	YOD1	1.146396	0.01633

ENSEMBL ID	HGNC	Log2 TE	BABEL p-value
ENSG00000186468	RPS23	1.302477	0.016725
ENSG00000136942	RPL35	1.279749	0.017369
ENSG00000250722	SEPP1	-1.03545	0.01764
ENSG00000139132	FGD4	-1.14179	0.017942
ENSG00000114942	EEF1B2	1.434033	0.018005
ENSG00000215472	RPL17	1.29905	0.018135
ENSG00000138134	STAMBPL1	-1.13882	0.018424
ENSG00000167526	RPL13	1.465152	0.018815
ENSG00000196323	ZBTB44	-1.22176	0.019332
ENSG00000221983	UBA52	1.125432	0.020091
ENSG00000177954	RPS27	1.428562	0.020692
ENSG00000148082	SHC3	-1.54133	0.021037
ENSG00000145425	RPS3A	1.660815	0.021105
ENSG00000163661	PTX3	-1.57581	0.021121
ENSG00000089009	RPL6	1.106565	0.021763
ENSG00000142534	RPS11	1.314223	0.02235
ENSG00000254772	EEF1G	1.550899	0.023278
ENSG00000112306	RPS12	1.642226	0.023317
ENSG00000232859	C17orf108	-1.32937	0.026416
ENSG00000108298	RPL19	1.075199	0.027309
ENSG00000111057	KRT18	1.811152	0.027619
ENSG00000106789	CORO2A	1.211664	0.027779
ENSG00000008988	RPS20	1.435033	0.030941
ENSG00000104321	TRPA1	-1.25586	0.031417
ENSG00000168028	RPSA	1.358592	0.031477
ENSG000000081377	CDC14B	-1.33116	0.03257
ENSG00000146966	DENND2A	-1.15633	0.033148
ENSG00000124614	RPS10	1.210967	0.035745
ENSG00000138386	NAB1	-1.01045	0.036666
ENSG00000128602	SMO	-1.25412	0.037773
ENSG00000141837	CACNA1A	-1.05541	0.044943
ENSG00000152413	HOMER1	1.003421	0.04711
ENSG00000135749	PCNXL2	-1.26064	0.048103

Some characteristics of these gene signatures, including the identity of the pathway with the highest statistical association for each signature, are listed in Table 4 (while these are the most significant, it is notable that significant association of these gene lists with other pathways were observed).

5 Table 4. Properties of Fibrotic Disorder Gene Signatures from IPA Analysis
(Results from Single Representative Experimental Replicate)

	RNA (Transcriptome)	RPF (Translational Rate)	Translational Efficiency
p-value threshold for differential	0.1	0.05	0.05
No. of genes meeting threshold	194	211	238
% of total gene set	4.20%	4.50%	5.10%
Most significant pathway / category from IPA analysis	Hepatic Fibrosis / Hepatic Stellate Cell Activation	Hepatic Fibrosis / Hepatic Stellate Cell Activation	Canonical eIF2 Signaling Pathway

In particular, genes showing changes in RNA levels and translational rates were most strongly associated with Hepatic Fibrosis/Hepatic Stellate Cell Activation (*see* Figures 4 and 5). This action of TGF β in fibroblasts recapitulates much of the behavior observed in liver fibrosis. In contrast, a 12 gene signature showing a change in translational efficiency was most strongly associated with regulation of eIF2 signaling (Figure 7). All 12 genes showed a significant increase in translational efficiency (TE) (Table 5; Figures 7 and 8), which extends far beyond these genes. For example, 118 of the 141 genes in the pathway evaluable in this study moved in concert, showing an increase in translational efficiency (*see* Table 5; Figure 8A). The translational efficiencies of the 141 pathway-associated genes in fibroblasts before treatment with TGF β were low (mean value -1.70 log₂ relative to population mean); the impact of TGF β induced transformation was to increase the translational efficiency of many genes in this signature (mean value of signature upon TGF β treatment was -1.05). Nonetheless, this was still two-fold lower than the overall population and indicates this pathway is a bottleneck in cellular transformation.

Table 5. Effect of TGF β and PP242 Treatment on Translational Efficiency of Genes in Canonical eIF2 Signaling Pathway

Symbol	Entrez Gene Name	p-value ^s	Log ₂ FC TE	
			TGF- β	TGF- β + PP242
EIF2C2	eukaryotic translation initiation factor 2C	0.849	0.12	0.49
AKT1	v-akt murine thymoma viral oncogene homolog 1	0.604	-0.15	0.23
AKT2	v-akt murine thymoma viral oncogene homolog 2	0.634	-0.16	1.05
AKT3	v-akt murine thymoma viral oncogene homolog 3	0.734	0.35	0.09
ATM	ataxia telangiectasia mutated	0.657	-0.16	-0.62
EIF1	eukaryotic translation initiation factor 1	0.001	0.75	0.40
EIF5	eukaryotic translation initiation factor 5	0.205	0.56	-0.10
EIF2A	eukaryotic translation initiation factor 2A, 65kDa	0.137*	0.74	-0.51
EIF2AK1	eukaryotic translation initiation factor 2-alpha kinase 1	0.505	-0.25	-0.66
EIF2AK3	eukaryotic translation initiation factor 2-alpha kinase 3	0.210	0.67	-0.33
EIF2AK4	eukaryotic translation initiation factor 2 alpha kinase 4	0.484	-0.28	-0.22
EIF2B1	eukaryotic translation initiation factor 2B, subunit 1 alpha, 26kDa	0.185	0.57	-0.11
EIF2B2	eukaryotic translation initiation factor 2B, subunit 2 beta, 39kDa	0.825*	0.16	0.77
EIF2B3	eukaryotic translation initiation factor 2B, subunit 3 gamma, 58kDa	0.281	0.54	0.51
EIF2B4	eukaryotic translation initiation factor 2B, subunit 4 delta, 67kDa	0.366	-0.35	0.34
EIF2B5	eukaryotic translation initiation factor 2B, subunit 5 epsilon, 82kDa	0.432	0.46	0.22
EIF2S1	eukaryotic translation initiation factor 2, subunit 1 alpha, 35kDa	0.622	-0.18	-0.33
EIF2S2	eukaryotic translation initiation factor 2, subunit 2 beta, 38kDa	0.290	-0.43	-1.41
EIF2S3	eukaryotic translation initiation factor 2, subunit 3 gamma, 52kDa	0.027*	1.05	-0.26
EIF3A	eukaryotic translation initiation factor 3, subunit A	0.758	0.04	-1.26
EIF3B	eukaryotic translation initiation factor 3, subunit B	0.671	0.30	0.18
EIF3D	eukaryotic translation initiation factor 3, subunit D	0.239*	0.71	-0.08
EIF3E	eukaryotic translation initiation factor 3, subunit E	0.023*	1.33	-0.67
EIF3F	eukaryotic translation initiation factor 3, subunit F	0.202*	0.92	0.46
EIF3G	eukaryotic translation initiation factor 3, subunit G	0.513*	0.55	0.48
EIF3H	eukaryotic translation initiation factor 3, subunit H	0.004*	1.34	-0.20

Symbol	Entrez Gene Name	p-value [§]	Log ₂ FC TE	
			TGF-β	TGF-β + PP242
EIF3I	eukaryotic translation initiation factor 3, subunit I	0.654	0.29	0.42
EIF3K	eukaryotic translation initiation factor 3, subunit K	0.267*	0.73	1.09
EIF3L	eukaryotic translation initiation factor 3, subunit L	0.025*	1.63	0.27
EIF3M	eukaryotic translation initiation factor 3, subunit M	0.121*	0.90	-0.42
EIF4A1	eukaryotic translation initiation factor 4A1	0.659	0.35	0.58
EIF4A2	eukaryotic translation initiation factor 4A2	0.507	0.41	-0.63
EIF4A3	eukaryotic translation initiation factor 4A3	0.765	0.19	0.35
EIF4E	eukaryotic translation initiation factor 4E	0.219	-0.51	-1.20
EIF4G1	eukaryotic translation initiation factor 4 gamma, 1	0.725	0.10	-0.01
EIF4G2	eukaryotic translation initiation factor 4 gamma, 2	0.080	0.87	-0.09
EIF4G3	eukaryotic translation initiation factor 4 gamma, 3	0.126	-0.54	-0.93
FAU	Finkel-Biskis-Reilly murine sarcoma virus (FBR-MuSV) ubiquitously expressed	0.718*	0.35	0.47
GRB2	growth factor receptor-bound protein 2	0.874	0.05	0.19
GSK3B	glycogen synthase kinase 3 beta	0.741	-0.13	-0.01
HRAS	Harvey rat sarcoma viral oncogene homolog	0.992	0.13	0.43
MAP2K1	mitogen-activated protein kinase kinase 1	0.630	-0.18	-0.59
MAP2K2	mitogen-activated protein kinase kinase 2	0.835	0.02	0.54
MAPK1	mitogen-activated protein kinase 1	0.799	0.12	0.25
MAPK3	mitogen-activated protein kinase 3	0.483	0.29	0.90
MRAS	muscle RAS oncogene homolog	0.271	-0.45	-0.30
NRAS	neuroblastoma RAS viral (v-ras) oncogene homolog	0.708	0.19	-0.89
PABPC1	poly(A) binding protein, cytoplasmic 1	0.824*	0.16	-1.48
PAIP1	poly(A) binding protein interacting protein 1	0.115	-0.69	-1.28
PIK3C3	phosphatidylinositol 3-kinase, catalytic subunit type 3	0.770	-0.09	-0.50
PIK3C2A	phosphatidylinositol-4-phosphate 3-kinase, catalytic subunit type 2 alpha	0.116	-0.64	-0.77
PIK3CA	phosphatidylinositol-4,5-bisphosphate 3-kinase, catalytic subunit alpha	0.135	-0.66	-0.94
PIK3R4	phosphoinositide-3-kinase, regulatory subunit 4	0.070	-0.69	-0.57
PPP1CA	protein phosphatase 1, catalytic subunit, alpha isozyme	0.907	0.04	0.92
PPP1CB	protein phosphatase 1, catalytic subunit, beta isozyme	0.685	0.13	-0.80
PPP1CC	protein phosphatase 1, catalytic subunit, gamma isozyme	0.583	0.27	0.04
PPP1R15A	protein phosphatase 1, regulatory subunit 15A	0.748	-0.09	-0.51
RAF1	v-raf-1 murine leukemia viral oncogene homolog 1	0.926	0.16	0.30

Symbol	Entrez Gene Name	p-value [§]	Log ₂ FC TE	
			TGF-β	TGF-β + PP242
RPL3	ribosomal protein L3	0.031*	1.68	0.09
RPL4	ribosomal protein L4	0.150*	1.23	-0.42
RPL5	ribosomal protein L5	0.014*	1.49	-0.37
RPL6	ribosomal protein L6	0.093	1.13	-0.36
RPL7	ribosomal protein L7	0.583	0.59	-1.33
RPL8	ribosomal protein L8	0.333*	0.69	0.90
RPL9	ribosomal protein L9	0.117*	1.41	0.69
RPL10	ribosomal protein L10	0.335*	0.74	0.09
RPL11	ribosomal protein L11	0.323*	0.78	-0.47
RPL12	ribosomal protein L12	0.123*	1.28	0.02
RPL13	ribosomal protein L13	0.482*	0.83	1.06
RPL14	ribosomal protein L14	0.114*	1.08	-0.35
RPL15	ribosomal protein L15	0.092*	0.95	-0.14
RPL17	ribosomal protein L17	0.267*	0.97	-1.28
RPL18	ribosomal protein L18	0.376*	0.81	0.58
RPL19	ribosomal protein L19	0.283*	0.71	0.31
RPL21	ribosomal protein L21	0.503*	0.83	-0.74
RPL22	ribosomal protein L22	0.849	0.22	-0.96
RPL23	ribosomal protein L23	0.363*	0.79	-0.78
RPL24	ribosomal protein L24	0.138*	1.09	0.28
RPL26	ribosomal protein L26	0.152*	1.42	-0.08
RPL27	ribosomal protein L27	0.099*	1.17	0.38
RPL28	ribosomal protein L28	0.000*	1.24	0.60
RPL29	ribosomal protein L29	0.139*	1.16	0.45
RPL30	ribosomal protein L30	0.125*	1.20	0.22
RPL31	ribosomal protein L31	0.322*	0.80	-0.71
RPL32	ribosomal protein L32	0.045	1.55	0.71
RPL34	ribosomal protein L34	0.098*	1.21	-0.44
RPL35	ribosomal protein L35	0.560*	0.59	0.07
RPL36	ribosomal protein L36	0.341*	0.76	0.46
RPL37	ribosomal protein L37	0.144*	1.31	0.20
RPL38	ribosomal protein L38	0.143*	1.00	0.50
RPL41	ribosomal protein L41	0.203*	0.85	0.77
RPL10A	ribosomal protein L10a	0.393*	0.92	0.39
RPL13A	ribosomal protein L13a	0.233*	1.26	0.42
RPL18A	ribosomal protein L18a	0.254*	1.06	1.14

Symbol	Entrez Gene Name	p-value [§]	Log ₂ FC TE	
			TGF-β	TGF-β + PP242
RPL22L1	ribosomal protein L22-like 1	0.076*	0.84	-0.42
RPL23A	ribosomal protein L23a	0.226*	1.03	0.23
RPL27A	ribosomal protein L27a	0.158*	1.17	0.31
RPL35A	ribosomal protein L35a	0.079*	1.22	0.56
RPL36A L	ribosomal protein L36a-like	0.792	-0.08	-0.99
RPL37A	ribosomal protein L37a	0.175*	1.33	0.71
RPL7A	ribosomal protein L7a	0.080*	1.37	0.84
RPLP0	ribosomal protein, large, P0	0.641	0.73	0.76
RPLP1	ribosomal protein, large, P1	0.546*	0.50	1.00
RPLP2	ribosomal protein, large, P2	0.243*	0.93	0.58
RPS2	ribosomal protein S2	0.441*	0.88	-0.07
RPS3	ribosomal protein S3	0.198*	1.22	0.42
RPS5	ribosomal protein S5	0.510*	0.75	0.78
RPS6	ribosomal protein S6	0.079*	1.59	0.48
RPS7	ribosomal protein S7	0.152*	1.13	0.13
RPS8	ribosomal protein S8	0.180*	1.20	-0.08
RPS9	ribosomal protein S9	0.288*	0.97	0.87
RPS10	ribosomal protein S10	0.519*	0.74	0.75
RPS11	ribosomal protein S11	0.339*	0.97	0.53
RPS12	ribosomal protein S12	0.500*	0.94	0.12
RPS13	ribosomal protein S13	0.098*	1.15	0.36
RPS14	ribosomal protein S14	0.309*	1.01	0.44
RPS15	ribosomal protein S15	0.096*	0.74	0.06
RPS16	ribosomal protein S16	0.020*	1.10	3.46
RPS18	ribosomal protein S18	0.349*	1.18	0.48
RPS19	ribosomal protein S19	0.321*	0.99	0.61
RPS20	ribosomal protein S20	0.194*	1.39	0.29
RPS21	ribosomal protein S21	0.045*	1.36	0.73
RPS23	ribosomal protein S23	0.415*	0.74	-0.02
RPS24	ribosomal protein S24	0.343*	0.66	-0.64
RPS25	ribosomal protein S25	0.040*	1.58	0.05
RPS26	ribosomal protein S26	0.242*	0.71	0.60
RPS27	ribosomal protein S27	0.205*	1.08	-0.09
RPS28	ribosomal protein S28	0.336*	0.74	0.15
RPS29	ribosomal protein S29	0.329*	0.89	0.64
RPS15A	ribosomal protein S15a	0.074*	1.53	-0.04

Symbol	Entrez Gene Name	p-value [§]	Log ₂ FC TE	
			TGF- β	TGF- β + PP242
RPS27A	ribosomal protein S27a	0.346*	0.71	-0.06
RPS27L	ribosomal protein S27-like	0.687	-0.15	-0.89
RPS3A	ribosomal protein S3A	0.197*	1.48	0.59
RPS4X	ribosomal protein S4, X-linked	0.252*	1.14	0.18
RPS4Y1	ribosomal protein S4, Y-linked 1	0.280	0.78	0.07
RPSA	ribosomal protein SA	0.407*	0.99	1.30
RRAS	related RAS viral (r-ras) oncogene homolog	0.965	0.00	0.70
RRAS2	related RAS viral (r-ras) oncogene homolog 2	0.914	0.02	-0.41
SHC1	SHC (Src homology 2 domain containing) transforming protein 1	0.777	-0.04	0.78
SOS2	son of sevenless homolog 2 (Drosophila)	0.419	0.49	-0.63
UBA52	ubiquitin A-52 residue ribosomal protein fusion product 1	0.249*	0.87	0.71

[§]p-value is for the difference between TGF β -treated and control cells. Gene names in **BOLD** have p values ≤ 0.05 .

* p-values marked with an asterisk have values ≤ 0.05 based upon data from five biological control and five TGF β -treated replicates.

5 The linkage of fibroblast transformation by TGF β to the eIF2 pathway can be identified by changes in translational rate. For example, the mean log₂FC (log₂ fold change) in TE for this set of 141 genes was 0.65 as compared to 0.02 for the overall gene population. The mean change in translational rate was of the same order (0.76 for the 141 genes as compared to 0.24 for the overall gene population). But in this system, 10 the relatively large number of changes in translational rate driven primarily by changes in mRNA levels obscured this relationship. In contrast, there is little evidence of linkage to the eIF2 pathway in the transcriptome data (*e.g.*, mean change was 0.11 for the 141 genes as compared to 0.22 for the overall gene population).

15 When fibroblasts were treated with TGF β and the mTOR inhibitor PP242, 11 out of 12 of the subset of genes found to be associated with the eIF2 signaling pathway moved toward the untransformed, normal state (Figure 7). The mean increase in translational efficiency in these 12 genes caused by TGF β treatment of fibroblasts was 1.3 log₂, but the presence of PP242 decreased this value to 0.4 log₂. Similar results were seen for all genes in the pathway, wherein 104 of 141 genes moved toward normal

(Figure 8). The mean increase in translational efficiency of these 12 genes in fibroblasts treated with TGF β was 0.65 log₂, while the presence of PP242 decreased this value to 0.09 log₂. Clearly, the presence of PP242 maintains the translational efficiencies of the genes in the eIF2 pathway at their normal levels in fibroblasts.

5 Moreover, PP242 inhibition of mTOR directly regulates the translational efficiencies of a number of genes in the eIF2 pathway in other disease cell systems (*e.g.*, prostate (PC3) and colon (SW620) cancer cells). Normalization of this pathway by mTOR inhibitor PP242 is due to substantially inhibiting TGF β induced transformation of fibroblasts to fibrotic myofibroblasts.

10 Conclusion

TGF β -dependent transformation of fibroblasts to myofibroblasts is known to be driven in large part by transcriptional activation. Here, changes in translational rate and RNA levels on a genome-wide level were shown to be highly correlated (*see* Figure 3). In contrast, changes in translational efficiency were relatively independent of

15 transcriptional and translational rate changes. Thus, in this case, measurements of translational efficiency provide a unique window into the cellular biology of fibrotic disorders. Correspondingly, the outcome of pathway analysis based on gene identification via changes in translational efficiency upon TGF β treatment is quite distinct from analyses based upon transcription or translational rate.

20 Ribosomal profiling showed that the effect of PP242 on both procollagen and α -SMA were a consequence of preventing fibroblast transformation and transcriptional regulation (instead of a decrease in translation efficiency of mRNA to protein). Specifically, the translational efficiencies of the procollagen and α -SMA were essentially independent of TGF β and PP242 treatment. Co-administration of TGF β

25 with mTOR inhibitor PP242 reverses or prevents the changes observed in the eIF2 pathway (*i.e.*, normalizes the translational efficiencies of the genes) and inhibits increased production of fibrotic disorder biomarker proteins, type 1 procollagen and smooth muscle α -actin (which are both hallmarks of TGF- β -mediated fibroblast transformation into myofibroblasts). Although these two biomarkers are only affected

30 at the transcriptional level and not the translational level, they nonetheless provide a

means to monitor the pathogenic state of the cell that is mediated by other fibrosis-related genes that are affected at the translational level.

Comparison of translational efficiencies between the normal, healthy state (fibroblasts) and the pathogenic state (fibrotic myofibroblasts induced by TGF β treatment) identified a novel pathway previously not associated with fibrosis, which is a new insight into a key role of translational efficiency in the pathogenesis of fibrotic disease. Further, an mTOR inhibitor (such as PP242) that modulates this fibrotic disorder-associated pathway and prevents TGF β -mediated fibroblast to myofibroblast transformation confirms the association of this pathway with fibrotic disease and, thus, shows that components and regulators of this pathway are new targets. The methods of the instant disclosure show that new gene signatures having altered translational profiles (*e.g.*, altered translational efficiency) may be identified using such methods. Furthermore, these data show that an agent or therapeutic that normalizes a translational profile may also be identified. Finally, these data show that targets not previously validated for a particular disorder (in this case, fibrosis), can be identified and validated using the methods of this disclosure.

EXAMPLE 4

EFFECT OF EIF4A INHIBITION ON FIBROTIC DISEASE DEVELOPMENT

Silvestrol, a cyclopenta[b]benzofuran compound, is a natural product known to inhibit eIF4A, which is the DEAD-box RNA helicase of the eIF4F complex. The TGF β -mediated transdifferentiation of fibroblasts as described in Example 1 was used as a model to examine whether modulation of eIF4A might have a role in fibrotic disorders.

Briefly, normal human lung fibroblasts (Lonza; cell passage numbers 2 through 5 were used for all experiments) were seeded (Day 0) and cultured in Dulbecco's modified Eagle medium (DMEM) supplemented with 10% fetal bovine serum (FBS), penicillin, streptomycin and glutamax (Invitrogen) at 37°C in a humidified incubator with 5% CO₂ conditions overnight. On Day 1, cells were harvested, washed with phosphate buffered saline (PBS), and then incubated for 48 hours in fresh serum-free

DMEM supplemented with penicillin, streptomycin, and glutamax. On Day 3, cells were harvested, resuspended in fresh serum-free DMEM containing silvestrol (1, 5, 10, 75, 100, 500, or 1000 nM) and 10 ng/ml TGF β , and cultured for 24 hours. Controls included untreated cells and cells treated with only TGF β .

5 After this 24 hour incubation, procollagen type 1 levels were measured by collecting culture media, centrifuging to pellet cellular debris, and stored at -80°C. Procollagen Type 1 C-Peptide (PIPC) was quantified using the (PIP) EIA kit (Clontech) according to manufacturer's instructions. The TGF β -treated fibroblasts of this example are examined by ribosomal profiling (about 6 x 10⁶ cells/10 cm plate) and western blot
10 analysis (about 1 x 10⁶ cells/well of a 6-well plate).

Results

Transdifferentiation of fibroblasts into fibrotic myofibroblasts by treatment with TGF β for 24 hours was accompanied by an approximately 75% increase in procollagen production, while treatment with silvestrol was able to block this increase (EC₅₀ of
15 about 12 nM) (Figure 9). Inhibition of procollagen by silvestrol appears saturable, with the extent of inhibition at higher concentrations reaching about 90% of untreated TGF β -transformed myofibroblasts (equivalent to about 80% inhibition of untreated, untransformed control fibroblasts). Expression of TGF β -induced myofibroblast differentiation marker, smooth muscle actin (α -SMA), was also analyzed by western
20 blot analysis. After 24 hours of TGF- β stimulation, increased α -SMA protein levels were detectable, while the level of β -actin did not change. As with procollagen, co-incubation of cells with TGF β and silvestrol caused a reduction of the α -SMA protein level in a dose dependent manner (EC₅₀ of about 11 nM) such that the ratio of α -SMA to β -actin at higher silvestrol concentrations was the same as in untransformed,
25 untreated fibroblasts (Figure 10).

Conclusion

Co-administration of TGF β with eIF4A inhibitor silvestrol reverses or prevents the changes observed in a fibrotic disorder-related pathway as evidenced by an inhibition of increased production of fibrotic disorder biomarker proteins, type 1
30 procollagen and α -smooth muscle actin (which are both hallmarks of TGF β -mediated fibroblast transdifferentiation into myofibroblasts). While silvestrol is known to have

anti-tumor activity (*see, e.g., Cencic et al., PLoS One 4:e5223, 2009*), the usefulness of an eIF4A inhibitor like silvestrol in the treatment of fibrosis was an unexpected result.

EXAMPLE 5

EFFECT OF siRNA KNOCKDOWN OF EIF4A ON TRANSDIFFERENTIATION OF FIBROBLASTS INTO MYOFIBROBLASTS

Normal human lung fibroblasts were seeded (Day 0) and cultured in Dulbecco's modified Eagle medium (DMEM) supplemented with 10% fetal bovine serum (FBS), penicillin, streptomycin and glutamax (Invitrogen) at 37°C in a humidified incubator with 5% CO₂ overnight. On Day 1, cells were washed with phosphate buffered saline (PBS) and then incubated for 24 hours in fresh serum-free DMEM supplemented with penicillin, streptomycin, and glutamax. On Day 2, cells were transfected with siRNA against eIF4A1 (sieIF4A1; sense sequence: GCGAGCCAUUCUACCUUGUtt (SEQ ID NO.: 5); antisense sequence: ACAAGGUAGAAUGGCUCGCtg (SEQ ID NO.: 6)) or unrelated control siRNA (siCont) in serum-free DMEM and cultured for 24 hours. On Day 3, serum-free DMEM in the presence or absence of 10 ng/ml TGFβ was added and cultured for an additional 24 hours.

The levels of α-SMA and procollagen type 1, markers of fibroblast transdifferentiation into myofibroblasts and a surrogate for fibrotic disease progression, were measured after a 24 hour incubation of fibroblasts in the presence or absence of TGFβ sieIF4A1 or siCont. The knockdown efficiency was determined by quantitating the eIF4A mRNA level by qPCR (data not shown) and assessing the protein level by western analysis (see Figure 11).

For the imaging of α-smooth muscle actin and F-actin, normal human lung fibroblasts transfected with siControl or sieIF4A ± TGFβ as described above were seeded on coverslips for immunofluorescence microscopy (IF). The following day, cells were washed with PBS, fixed in 4% paraformaldehyde, permeabilized in PBS containing 0.3% Triton X-100®, and blocked in 10% goat serum. To stain for α-smooth muscle actin (α-SMA) and F-actin, coverslips were then incubated with the anti-α-SMA antibody followed by fluorescent conjugated secondary antibody and

fluorescent conjugated phalloidin. Finally, coverslips were washed and mounted with mounting media containing DAPI to visualize the nuclei. Immunofluorescence images were captured using the EVOS FL Cell Imaging System (Life Technologies) (Figure 12).

5 Results

Transdifferentiation of fibroblasts into myofibroblasts by treatment with TGF β was verified by an increase in production of both α -smooth muscle actin (α -SMA) and procollagen, which was not affected in samples transfected with control siRNA (siCont) alone. In contrast, cells transfected with eIF4A siRNA resulted in approximately 80%
10 knockdown of eIF4A (as determined by either qPCR or western analysis). The specific knockdown of eIF4A with siRNA substantially inhibited TGF β -induced increases in α -SMA and procollagen levels (Figure 11).

Examination of cell morphology by immunofluorescent staining of α -SMA shows an increase in α -SMA fibril formation associated with TGF β -induced
15 transdifferentiation of fibroblasts into myofibroblasts, which appear as long white fibrils with nuclei appearing as white spots in the middle of the fibril (see Figure 12A). The specific knockdown of eIF4A with siRNA inhibits the TGF β -induced transdifferentiation of fibroblasts into myofibroblasts, which results in decreased production and staining of α -SMA with primarily only the nuclei visible (see Figure
20 12C). The presence of either siRNA in the absence of TGF β treatment has no effect on the fibroblast phenotype and no effect on α -SMA production (see Figures 12B and 12D).

Conclusion

Specific knockdown of the translation initiation target, eIF4A, reverses or
25 prevents the changes observed in a fibrotic disorder-related pathway as evidenced by an inhibition of increased production of fibrotic disorder biomarker proteins, α -smooth muscle actin and type 1 procollagen (which are both hallmarks of TGF β -mediated fibroblast transdifferentiation into myofibroblasts).

EXAMPLE 6

EFFECT OF VARIOUS COMPOUNDS ON FIBROTIC DISEASE DEVELOPMENT

The TGF β -mediated transformation of fibroblasts as described in Examples 1 and 4 was used as a model to examine whether compounds known to treat fibrosis as well as compounds with no previously known association with the disease might have a role in treating fibrotic disorders. Compounds shown to be effective at treating fibrosis *in vitro* and *in vivo* and tested here include: pirfenidone (5-methyl-1-phenylpyridin-2-one, approved for the treatment of idiopathic pulmonary fibrosis), trichostatin A (TSA, 7-[4-(dimethylamino)phenyl]-N-hydroxy-4,6-dimethyl-7-oxohepta-2,4-dienamide) and rapamycin (an mTOR inhibitor). While pirfenidone has been shown to have antifibrotic and anti-inflammatory properties in various *in vitro* and *in vivo* models (Schaefer *et al.*, *Eur. Respir. Rev.* 20:85, 2011), it is unknown what biological molecule is targeted by this drug. TSA inhibits the class I and II histone deacetylase enzymes and has been shown to prevent the accumulation of extracellular matrix in mouse fibrosis models (Huber *et al.*, *Arthritis Rheum.* 56:2755, 2007; Van Beneden *et al.*, *Tox. Appl. Pharma.* 271:276, 2013).

Inhibitors and activators of specific targets with no previously known association with fibrosis were also examined in the TGF β -mediated fibroblast transformation assay. For example, agents that modulate translation factors were tested, including silvestrol (an eIF4A inhibitor), 1-(benzo[c][1,2,5]oxadiazol-5-yl)-3-(4-chlorophenyl)urea (BOCPU, an activator of eIF2 α K), siRNA specific for knocking down eIF4A and eIF4E. In addition, N1-guanyl-1,7-diaminoheptane (GC7), ciclopirox oloamine (CPX), and siRNA specific for knocking down DOHH were tested, which are inhibitors of enzymes that post-translationally modify eIF5A. Also tested was 3-deaza-adenosine (DZA), which indirectly inhibits the enzyme responsible for 5'-cap methylation of mRNA as well as a number of other methylating enzymes. Finally, also evaluated was 2-[(3,5-Difluoro-4-hydroxyphenyl)amino]-7,8-dihydro-5,7-dimethyl-8-(3-methylbutyl)-6(5H)-pteridone (BI-D1870), an inhibitor of the p90 ribosomal s6K enzyme within the RAS/MEK pathway.

Table 6. Effect of Various Compounds on Fibrosis Markers

Target	Translation Factor	Agent	% Renormalization*	
			α -SMA	Collagen
–	–	pirfenidone	+++	+++
Tyrosine kinase inhibitor	–	Nintedanib	+++	+++
eIF2 α K	eIF2 α	BOCPU	+++	ND [†]
eIF4A	eIF4A	Silvestrol	+++	+++
eIF4A	eIF4A	siRNA ^{††}	+++	++
mTOR	eIF4E	PP242	+++	+++
eIF4E	eIF4E	siRNA [§]	+	++
mTOR	rpS6	Rapamycin	+	++
Deoxyhypusine hydroxylase (DOHH)	eIF5A	CPX	+++	+++
		siRNA [#]	++	++
Histone deacetylase (HDAC)	eIF5A	TSA	++	+++
Deoxyhypusine synthase (DHPS)	eIF5A	GC7	+	+
mTOR	rpS6	Rapamycin	+	++
p90 Ribosomal S6 kinase (RSK)	rpS6	BI-D1870	++	+++
Adenosylhomocysteinase (AHCY)	Cap methylation	DZA	+	+

* + = 0-30%; ++ = 30-60%; +++ = >60%

[†] Not determined

^{††} 80% knockdown of eIF4A target

[§] 70% knockdown of eIF4E target

[#] 50% knockdown of DOHH target

5

10

15

As shown in the previous examples, transformation of fibroblasts into fibrotic myofibroblasts by treatment with TGF β for 24 hours was accompanied by an increase in procollagen and smooth muscle actin (α -SMA) production, as measured by ELISA and western blot analysis, respectively. Both PP242 (Example 1) and silvestrol (Example 4) were able to block this increase in procollagen and α -SMA – that is, renormalize production levels of these fibrosis biomarkers (*see, also*, Table 6). Interestingly, mTOR inhibitors PP242 and rapamycin do not promote renormalization to the same level, which may be due to the different mechanisms of action (rapamycin is an allosteric inhibitor of mTOR, while PP242 is an ATP-competitive inhibitor that

directly targets the mTOR catalytic site). The compounds most effective at promoting renormalization of the fibrosis markers in fibroblasts treated with TGFβ (PP242, silvestrol, pirfenidone, CPX, and TSA) were further analyzed by ribosomal profiling as described in Example 3.

5

EXAMPLE 7

TRANSLATIONAL PROFILING: REVERSAL (RENORMALIZATION)
OF FIBROTIC DISEASE DEVELOPMENT

Ribosomal profiles of TGFβ-treated fibroblasts in the presence or absence of PP242, silvestrol, pirfenidone, CPX, or TSA individually were prepared in duplicate
10 and analyzed for changes in translational efficiencies as described in Example 3.

Genes showing TGFβ-induced modulation of their translational profiles, including both up and down modulated profiles, as compared to untreated cells are provided in Table 7 (see column labeled "Control vs. TGFβ"). The gene set listed in Table 7 represents those genes having a log₂ fold change in translational efficiency of ≥
15 1 and ≤ -1 (translational efficiency p-value ≤ 0.01).

Table 7. Differential Effect of Various Compounds on Translational Renormalization of TGFβ Modulation

ENSEMBL ID	HGNC ID	Control vs. TGFβ [†]	Compound [‡]					Exp. (NS)		
			PP242	Pirfenidone	Silvestrol	CPX	TSA	1	2	3
ENSG00000154262	ABCA6	A	-	-	++	-	-	*		*
ENSG00000097021	ACOT7	D	-	-	-	-	-			*
ENSG00000143632	ACTA1	A	-	+		+	-			*
ENSG00000158859	ADAMTS4	D	-	-		-	-	*		*
ENSG00000154736	ADAMTS5	C	-	-	+	+	-			
ENSG00000049192	ADAMTS6	F	-	-	-	-	-			
ENSG00000176020	AMIGO3	C	-	+	-	-	+		*	*
ENSG00000101935	AMMECR1	D	-	-	-	-	-	*		*
ENSG00000166839	ANKDD1A	C	+	+	-	-	-			
ENSG00000154122	ANKH	D	-	-	+	-	-	*		*

ENSEMBL ID	HGNC ID	Control vs. TGFβ [†]	Compound [‡]					Exp. (NS)		
			PP242	Pirfenidone	Silvestrol	CPX	TSA	1	2	3
ENSG00000138356	AOX1	B	-	-	-	-	++			
ENSG00000129675	ARHGEF6	C	-	-	-	-	-			*
ENSG00000241685	ARPC1A	D	-	-	-	-	+	*		
ENSG00000066279	ASPM	A	+	+	+	-	+			*
ENSG00000130707	ASS1	D	-	-	-	-	-	*		
ENSG00000099624	ATP5D	D	-	-	-	-	-			
ENSG00000135390	ATP5G2	E	+	+	-	-	-			
ENSG00000087586	AURKA	B	-	-	-	-	++			
ENSG00000134897	BIVM	B	+	-	-	+	-	*	*	
ENSG00000145741	BTF3	D	-	-	-	-	-			
ENSG00000111678	C12orf57	D	-	-	-	-	+			*
ENSG00000176438	C14orf49	C	-	-	-	-	-			
ENSG00000104979	C19orf53	D	-	-	-	-	-			*
ENSG00000183172	C22orf32	D	-	-	-	-	-		*	
ENSG00000155621	C9orf85	D	+	-	-	-	-		*	
ENSG00000169239	CA5B	B	+	++	+	-	+			*
ENSG00000162545	CAMK2N1	C	-	-	-	-	-	*		
ENSG00000204397	CARD16	B	-	-	+	-	-	*		*
ENSG00000150636	CCDC102B	A	-	++	-	-	+			
ENSG00000101639	CEP192	C	-	-	+	-	-	*		*
ENSG00000198848	CES1	E	-	-		+	-			
ENSG00000250479	CHCHD10	F	+++	+++		-	-			*
ENSG00000122966	CIT	A	-	+	+++	-	+++			
ENSG00000175505	CLCF1	E	-	-	+	-	+			
ENSG00000105664	COMP	E	-	-	-	-	-			*
ENSG00000106789	CORO2A	E	-	-		-	-	*		*
ENSG00000127184	COX7C	D	-	-	-	-	-		*	
ENSG00000146592	CREB5	D	-	+	-	-	-		*	
ENSG00000109846	CRYAB	D	-	-	-	-	-			
ENSG00000171793	CTPS	D	-	-	-	-	-			*
ENSG00000137628	DDX60	C	-	-	+	-	-	*		*
ENSG00000146966	DENND2A	C	-	-	-	-	-	*		*
ENSG00000175084	DES	E	-	-	-	-	-	*		
ENSG00000139734	DIAPH3	E	-	++	-	-	-			
ENSG00000178401	DNAJC22	B	+	+	-	-	-		*	*

ENSEMBL ID	HGNC ID	Control vs. TGFβ [†]	Compound [‡]					Exp. (NS)		
			PP242	Pirfenidone	Silvestrol	CPX	TSA	1	2	3
ENSG00000013563	DNASE1L1	D	-	-	+	-	-		*	*
ENSG000000117543	DPH5	D	-	-	-	-	-	*	*	
ENSG000000187240	DYNC2H1	B	-	-	++	-	-	*		*
ENSG000000078401	EDN1	E	-	-	++	-	-	*		
ENSG000000156508	EEF1A1	E	+	+	-	-	+			
ENSG000000114942	EEF1B2	E	-	-	-	-	-			
ENSG000000104529	EEF1D	E	-	-	-	-	-			
ENSG000000254772	EEF1G	E	-	-	-	-	-	*		*
ENSG000000167658	EEF2	F	+	+	-	-	+			
ENSG000000130741	EIF2S3	D	-	-	-	-	-	*		
ENSG000000104408	EIF3E	E	+	-	-	-	-			
ENSG000000175390	EIF3F	D	-	-	-	-	-			
ENSG000000147677	EIF3H	E	-	-	-	-	-			
ENSG000000100129	EIF3L	D	-	-	-	-	-	*		
ENSG000000149100	EIF3M	D	-	-	-	-	-	*		
ENSG000000063046	EIF4B	E	+	+	-	-	+	*		
ENSG000000088367	EPB41L1	D	-	-	+	-	+			
ENSG000000149806	FAU	D	-	-	-	-	-	*		*
ENSG000000138675	FGF5	B	++	+	-	-	-			
ENSG000000094963	FMO2	A	+++	+++	-	-	+			*
ENSG000000151474	FRMD4A	C	-	-	-	+	+	*		
ENSG000000111640	GAPDH	E	+	-	-	-	-			
ENSG000000139354	GAS2L3	A	-	-	+	-	+	*		*
ENSG000000107623	GDF10	C	-	-	-	+	+	*		
ENSG000000130513	GDF15	D	-	-	-	-	-			*
ENSG000000106415	GLCCI1	B	+	+	+	-	+			
ENSG000000105373	GLTSCR2	E	+	+	-	+	+			
ENSG000000204628	GNB2L1	E	+	+	-	-	-			
ENSG000000121957	GPSM2	B	-	-	-	-	++			
ENSG000000113070	HBEGF	D	-	-	+	-	-	*		*
ENSG000000164588	HCN1	D	-	-	-					*
ENSG000000146066	HIGD2A	D	+	+	-	-	-			*
ENSG000000064393	HIPK2	C	-	-	-	-	-	*		
ENSG000000135486	HNRNPA1	D	+	-	-	-	-			
ENSG000000152413	HOMER1	D	-	-	+	-	-	*		*

ENSEMBL ID	HGNC ID	Control vs. TGFβ [†]	Compound [‡]					Exp. (NS)		
			PP242	Pirfenidone	Silvestrol	CPX	TSA	1	2	3
ENSG00000170801	HTRA3	C	-	-	-	-	-	*		
ENSG00000137331	IER3	E	-	-	-	-	-			
ENSG00000115267	IFIH1	B	+	+	-	-	-			*
ENSG00000089289	IGBP1	E	-	-	-	-	-	*		
ENSG00000163453	IGFBP7	E	-	-	-	-	-			
ENSG00000136244	IL6	E	-	-	-	-	-			
ENSG00000178035	IMPDH2	D	-	-	-	-	-	*		*
ENSG00000122641	INHBA	D	-	-	+	-	-			*
ENSG00000090376	IRAK3	C	-	-	+	-	+		*	*
ENSG00000149596	JPH2	F	-					*		*
ENSG00000089094	KDM2B	C	-	-	-	-	-		*	*
ENSG00000132510	KDM6B	E	-	-	+	-	-			
ENSG00000169330	KIAA1024	F				-	-	*	*	
ENSG00000049130	KITLG	A	-	-	-	-	+			*
ENSG00000135480	KRT7	E	-	++	-	++	-			
ENSG00000196878	LAMB3	D	-	-	+	+	-	*		
ENSG00000125869	LAMP5	E	-	-	-	+	-			
ENSG00000182909	LENG9	D	-	-	+	-	-			*
ENSG00000050426	LETMD1	D	-	-	-	-	-			
ENSG00000100097	LGALS1	D	-	+	-	-	-			
ENSG00000121897	LIAS	D	-	-	-	-	-		*	
ENSG00000128342	LIF	D	-	-	-	-	-	*		*
ENSG00000198121	LPAR1	C	-	-	-	-	-			
ENSG00000171517	LPAR3	A	-	-	+	-	++		*	*
ENSG00000188906	LRRK2	C	-	-	-	-	-	*		*
ENSG00000197442	MAP3K5	B	-	-	-	-	+			*
ENSG00000132031	MATN3	D	-	-	+	-	-			
ENSG00000112559	MDFI	A	-	-	-		-			
ENSG00000204520	MICA	E	-	-	-	-	-			
ENSG00000174100	MRPL45	D	+	-	-	-	-		*	*
ENSG00000129422	MTUS1	B	+	++	-	-	-			
ENSG00000170681	MURC	F	-	-	-	-	+			*
ENSG00000141140	MYO19	D	-	-	+	-	-	*		*
ENSG00000196531	NACA	E	-	+	-	-	-			
ENSG00000109065	NAT9	D	-	-	-	-	-		*	

ENSEMBL ID	HGNC ID	Control vs. TGFβ [†]	Compound [‡]					Exp. (NS)		
			PP242	Pirfenidone	Silvestrol	CPX	TSA	1	2	3
ENSG00000125967	NECAB3	E	-	-	-	-	-		*	*
ENSG00000136999	NOV	C	-	-	-	-	+		*	
ENSG00000183971	NPW	D	-	-	-		-			*
ENSG00000157168	NRG1	D	-	-	+	-	-			
ENSG00000154358	OBSCN	B	-	-	-	+	+++	*		
ENSG00000155463	OXA1L	D	-	-	-	-	-		*	*
ENSG00000089041	P2RX7	B	+	++	-	++	++	*	*	
ENSG00000070756	PABPC1	D	-	-	-	-	-			*
ENSG00000090621	PABPC4	D	-	-	+	-	-	*		*
ENSG00000137819	PAQR5	B	-	-	-	+	-	*		
ENSG00000197111	PCBP2	E	-	-	+	-	+			
ENSG00000184588	PDE4B	B	-	+	+++	-	+	*		*
ENSG00000138735	PDE5A	B	-	-	-	-	-	*		*
ENSG00000171408	PDE7B	A	-	-	+	-	++			
ENSG00000075651	PLD1	B	-	-	++	-	-	*		*
ENSG00000052126	PLEKHA5	C	-	-	++	-	-	*		*
ENSG00000143850	PLEKHA6	B	-	-	-	+	+			
ENSG00000120278	PLEKHG1	B	-	-	-	+	-	*		
ENSG00000152527	PLEKHH2	B	-	-	-	-	-	*		*
ENSG00000198523	PLN	F	-	-		-	-			*
ENSG00000102007	PLP2	D	-	-	-	-	-			
ENSG00000186184	POLR1D	E	-	-	-	-	-	*	*	
ENSG00000132170	PPARG	A	-	+++	+++	-	+			
ENSG00000118898	PPL	B	-	-	-	-	-	*		
ENSG00000185920	PTCH1	D	-	-	-	-	-	*		*
ENSG00000172053	QARS	D	-	-	-	-	-			
ENSG00000105514	RAB3D	B	+	+	-	-	++	*	*	
ENSG00000133321	RARRES3	B	-	-	-	-	-			
ENSG00000108551	RASD1	E	-	-	+++	-	-			*
ENSG00000068028	RASSF1	D	-	-	-	-	-			
ENSG00000164292	RHOBTB3	B	-	-	-	-	-	*		*
ENSG00000185008	ROBO2	C	-	-	-	+	-	*		
ENSG00000166503	RP11-382A20.3.1	C	-	-	-	-	-	*		*
ENSG00000147403	RPL10	E	-	+	-	-	+			
ENSG00000198755	RPL10A	E	-	-	-	-	-			

ENSEMBL ID	HGNC ID	Control vs. TGFβ [†]	Compound [‡]					Exp. (NS)		
			PP242	Pirfenidone	Silvestrol	CPX	TSA	1	2	3
ENSG00000142676	RPL11	E	-	-	-	-	-			
ENSG00000197958	RPL12	E	-	+	-	-	-			
ENSG00000167526	RPL13	E	-	-	-	-	-	*		*
ENSG00000142541	RPL13A	E	-	+	-	-	-			
ENSG00000188846	RPL14	E	-	-	-	-	+			
ENSG00000174748	RPL15	D	-	-	-	-	-			
ENSG00000215472	RPL17	E	-	+	-	-	-			
ENSG00000063177	RPL18	E	+	+	-	-	-			
ENSG00000105640	RPL18A	E	+	+	-	-	-			
ENSG00000108298	RPL19	D	-	-	-	-	-	*		*
ENSG00000122026	RPL21	E	-	+	-	-	-	*		*
ENSG00000163584	RPL22L1	D	-	+	-	-	-			
ENSG00000125691	RPL23	E	-	+	-	-	-			
ENSG00000198242	RPL23A	E	-	-	-	-	-	*		*
ENSG00000114391	RPL24	E	-	-	-	-	-			
ENSG00000161970	RPL26	E	-	+	-	-	-			
ENSG00000131469	RPL27	E	-	-	-	-	-	*		
ENSG00000166441	RPL27A	F	-	+	-	-	-			
ENSG00000108107	RPL28	E	-	+	-	-	-			
ENSG00000162244	RPL29	E	-	-	-	-	-			
ENSG00000100316	RPL3	E	-	+	-	-	-			
ENSG00000156482	RPL30	E	-	+	-	-	-			
ENSG00000071082	RPL31	E	+	+	-	-	-			
ENSG00000144713	RPL32	E	-	-	-	-	-			
ENSG00000109475	RPL34	E	-	+	-	-	-			
ENSG00000136942	RPL35	E	-	-	-	-	-			*
ENSG00000182899	RPL35A	E	+	+	-	-	-			
ENSG00000130255	RPL36	E	-	+	-	-	-			
ENSG00000165502	RPL36AL	D	-		-	-	-			
ENSG00000145592	RPL37	E	-	+	-	-	-			
ENSG00000197756	RPL37A	E	-	+	-	-	-			
ENSG00000172809	RPL38	E	-	-	-	-	-	*		
ENSG00000198918	RPL39	E	-	-	-	-	-			
ENSG00000174444	RPL4	E	-	+	-	-	-			
ENSG00000229117	RPL41	E	-	-	-	-	-			*

ENSEMBL ID	HGNC ID	Control vs. TGFβ [†]	Compound [‡]					Exp. (NS)		
			PP242	Pirfenidone	Silvestrol	CPX	TSA	1	2	3
ENSG00000122406	RPL5	E	+	+	-	-	-			
ENSG00000089009	RPL6	D	-	-	-	-	-	*		*
ENSG00000148303	RPL7A	E	-	+	-	-	-			
ENSG00000161016	RPL8	E	-	-	-	-	-			
ENSG00000163682	RPL9	E	-	+	-	-	-			
ENSG00000089157	RPLP0	D	-	-	-	-	-	*		*
ENSG00000137818	RPLP1	E	-	+	-	-	-	*		*
ENSG00000177600	RPLP2	E	-	+	-	-	-			
ENSG00000124614	RPS10	D	-	+	-	-	-	*		*
ENSG00000142534	RPS11	D	-	-	-	-	-	*		*
ENSG00000112306	RPS12	E	-	-	-	-	-	*		*
ENSG00000110700	RPS13	E	-	-	-	-	-			
ENSG00000164587	RPS14	E	-	-	-	-	-			
ENSG00000115268	RPS15	D	+	+	-	-	-			*
ENSG00000134419	RPS15A	F	+	+	-	-	-			
ENSG00000105193	RPS16	F	-	+	-	-	-			
ENSG00000231500	RPS18	E	+	+	-	-	-			
ENSG00000105372	RPS19	E	-	+	-	-	-			
ENSG00000140988	RPS2	E	-	-	-	-	-	*		*
ENSG00000008988	RPS20	E	-	-	-	-	-	*		
ENSG00000171858	RPS21	E	-	-	-	-	-			
ENSG00000186468	RPS23	E	-	-	-	-	-			
ENSG00000138326	RPS24	E	-	-	-	-	-			
ENSG00000118181	RPS25	E	-	-	-	-	-			
ENSG00000197728	RPS26	D	-	-	-	-	-			*
ENSG00000177954	RPS27	E	-	+	-	-	-	*		
ENSG00000143947	RPS27A	E	-	-	-	-	-			
ENSG00000233927	RPS28	E	+	+	-	+	-			
ENSG00000213741	RPS29	E	+	+	-	-	-			
ENSG00000149273	RPS3	E	-	-	-	-	-	*		
ENSG00000145425	RPS3A	E	-	-	-	-	-	*		*
ENSG00000198034	RPS4X	E	-	-	-	-	-			
ENSG00000129824	RPS4Y1	D	-	-	-	-	-			
ENSG00000083845	RPS5	E	-	+	-	-	-			
ENSG00000137154	RPS6	E	-	-	-	-	-			

ENSEMBL ID	HGNC ID	Control vs. TGFβ [†]	Compound [‡]					Exp. (NS)		
			PP242	Pirfenidone	Silvestrol	CPX	TSA	1	2	3
ENSG00000100784	RPS6KA5	B	+	-	+	-	-			*
ENSG00000171863	RPS7	E	-	-	-	-	-			
ENSG00000142937	RPS8	E	-	+	-	-	-			*
ENSG00000170889	RPS9	E	-	+	-	-	-			
ENSG00000168028	RPSA	D	-	-	-	-	-	*		*
ENSG00000125744	RTN2	C	-	-	-	-	-			*
ENSG00000075213	SEMA3A	C	-	-	-	-	-	*		*
ENSG00000118473	SGIP1	B	-	-	+	-	-			*
ENSG00000185437	SH3BGR	D	-	-		-	-			*
ENSG00000148082	SHC3	B	-	-	-	+++	+	*		
ENSG00000110446	SLC15A3	B	-	-	-	-	++	*		
ENSG00000169100	SLC25A6	E	-	+	-	-	-			
ENSG00000146411	SLC2A12	B	-	-	++	-	-	*		*
ENSG00000138449	SLC40A1	E	-	++	-	++	+			
ENSG00000155465	SLC7A7	C	-	-	-	-	-	*		
ENSG00000128602	SMO	C	-	-	-	-	-	*		*
ENSG00000077312	SNRPA	E	-	-	+	-	-			*
ENSG00000134532	SOX5	B	-	-	-	-	+	*		
ENSG00000110693	SOX6	E	-	+	-	-	-		*	
ENSG00000132122	SPATA6	B	+	+	-	-	-			*
ENSG00000138134	STAMBPL1	C	-	+	+	+	+	*		*
ENSG00000101846	STS	C	-	+	-	-	-		*	
ENSG00000054654	SYNE2	B	-	-	-	-	-			
ENSG00000166012	TAF1D	E	-	+	-	-	+		*	
ENSG00000139372	TDG	D	-	-	-	-	-			
ENSG00000257949	TEN1	E	-	-	+	-	-			
ENSG00000168769	TET2	E	-	-	-	-	-	*		*
ENSG00000187605	TET3	E	-	-	-	-	-			*
ENSG00000120708	TGFBI	D	-	-	-	-	-	*		
ENSG00000169231	THBS3	D	+	+	+	-	-		*	*
ENSG00000113272	THG1L	D	-	-	-	-	-	*	*	
ENSG00000100575	TIMM9	D	-	-	-	-	-		*	
ENSG00000163931	TKT	D	-	+	-	-	-			
ENSG00000133687	TMTC1	B	-	-	+++	-	+			*
ENSG00000196683	TOMM7	D	-	-	-	+	-			*

ENSEMBL ID	HGNC ID	Control vs. TGFβ [†]	Compound [‡]					Exp. (NS)		
			PP242	Pirfenidone	Silvestrol	CPX	TSA	1	2	3
ENSG00000131747	TOP2A	B	-	-	+++	-	-	*		*
ENSG00000141933	TPGS1	D	-	-	-	-	-	*		
ENSG00000198467	TPM2	D	-	-	-	-	-	*		
ENSG00000133112	TPT1	D	-	-	-	-	-	*		*
ENSG00000124496	TRERF1	B	-	-	+	-	-			*
ENSG00000221983	UBA52	D	-	-	-	-	-	*		*
ENSG00000156467	UQCRB	E	-	-	-	-	-			
ENSG00000152818	UTRN	C	-	-	-	-	-	*		*
ENSG00000026025	VIM	D	-	-	-	-	-			
ENSG00000171425	ZNF581	E	-	-	-	-	-			*
ENSG00000168916	ZNF608	C	-	-	-	-	+			

- 5 [†] The letters A-F each represent a log₂ fold change in translational profile caused by TGFβ, with the following values: A = < -2.5; B = < -1.5 to -2.5; C = < -1.0 to -1.5; D = < 1.0 to 1.5; E = < 1.5 to 2.5; F = > 2.5. Letters A-C each represent a decrease in translation, while D-F each represent an increase in translation.
- 10 [‡] The effect of each compound in renormalizing the translational profile of each gene is shown as log₂ fold change as follows: – = < 0.9; + = 0.9 to 1.5; ++ = 1.5 to 2.0; +++ = > 2.0. Cells that are blank indicate no data was obtained.
- 10 ^{||} The change in translational profile for each gene was not statistically significant (NS) in each of the three TGFβ modulation experiments (Exp), but each trended in the same direction and a scatter plot (*see* Figure 13) shows that the translational regulation was highly correlated between experiments. Cells that are blank indicate the value obtained for that gene was statistically significant for that experiment.

Each of the compounds tested (PP242, silvestrol, pirfenidone, CPX, and TSA) showed a pronounced effect on renormalizing the levels of the fibrosis biomarkers procollagen and α-SMA, which had both increased in the presence of TGFβ (*see* Table 6). While these two biomarkers were not observed to be directly regulated at the translational level, ribosome profiling showed that expression of other genes was modulated at the translational level (*see* Table 7). Indeed, an analysis of the differential translational profile of healthy cells as compared to cells having a fibrotic disease state (*i.e.*, TGFβ-modulated) as compared to the diseased cells treated with PP242, silvestrol,

pirfenidone, CPX or TSA showed that each of these compounds is an agent that modulates translation in a fibrotic disease.

Moreover, a heat map of the differential translational profile for these compounds ($\Delta\log_2$ fold change (fibrotic cells vs. treated fibrotic cells) cut off of ≥ 1) reveals that each agent (PP242, silvestrol, pirfenidone, CPX, and TSA) has a unique renormalization profile (*see* Figure 14). Genes that show the most renormalization appear "white" in the heat map, while genes that are not renormalized appear "black" in the heat map and the genes that are "gray" had an intermediate level of renormalization. For example, pirfenidone renormalizes a number of genes that are also renormalized to the same degree by PP242, but many of the modulated genes are unique to pirfenidone. In contrast, each of silvestrol, CPX and TSA have very few genes that overlap with PP242 or pirfenidone, but each has its own strongly unique fingerprints.

Overall, each compound produces a unique gene signature associated with translational efficiency. Pirfenidone uniquely modulated the translational efficiency of 35 genes (CREB5, DIAPH3, LGALS1, NACA, RPL12, RPL13A, RPL17, RPL21, RPL22L1, RPL23, RPL26, RPL27A, RPL28, RPL3, RPL30, RPL34, RPL36, RPL37, RPL37A, RPL4, RPL7A, RPS9, RPLP1, RPLP2, RPS10, RPS16, RPS19, RPS27, RPS5, RPS8, RPS9, SLC25A6, SOX6, STS, TKT) (*i.e.*, not modulated, minimally modulated or not statistically significantly modulated by any of the other compounds tested), silvestrol uniquely modulated 26 genes (ABCA6, ANKH, CARD16, CEP192, DDX60, DNASE1L1, DYNC2H1, EDN1, HBEGF, HOMER1, INHBA, KDM6B, LENG9, MATN3, MYO19, NRG1, PABPC4, PLD1, PLEKHA5, RASD1, SGIP1, SLC2A12, SNRPA, TEN1, TOP2A, TRERF1), PP242 uniquely modulated five genes (C9orf85, EIF3E, GAPDH, HNRNPA1, MRPL45), CPX uniquely modulated six genes (CES1, LAMP5, PAQR5, PLEKHG1, ROBO2, TOMM7) and TSA uniquely modulated 13 genes (AOX1, ARPC1A, AURKA, C12orf57, GPSM2, KITLG, MAP3K5, MURC, NOV, RPL14, SLC15A3, SOX5, ZNF608).

In addition, there are unique gene signatures that overlap between two compounds. For example, there are a total of 19 genes (ANKDD1A, ATP5G2, CHCHD10, DNAJC22, FGF5, FMO2, GNB2L1, GLTSCR2, HIGD2A, IFIH, MTUS1, RPS18, RPL18A, RPL31, RPL35A, RPL5, RPS18, RPS29, SPATA6) modulated by

both pirfenidone and PP242; one gene (RPS6KA5) modulated by both silvestrol and PP242; one gene (BIVM) modulated by both PP242 and CPX; two genes (ACTA1, KRT7) modulated by both pirfenidone and CPX; four genes (AMIGO3, CCDC102B, RPL10, TAF1D) modulated by both pirfenidone and TSA; two genes (ADAMTS5, LAMB3) modulated by both silvestrol and CPX; eight genes (CLCF1, EPB41L1, GAS2L3, IRAK3, LPAR3, PCBP2, PDE7B, TMTC1) modulated by silvestrol and TSA; five genes (FRMD4A, GDF10, OBSCN, PLEKHA6, SHC3) modulated by CPX and TSA.

Finally, there are unique gene signatures that overlap between groups of three or four compounds. For example, one gene (THBS3) was modulated by pirfenidone, silvestrol and PP242; one gene (RPS28) was modulated by pirfenidone, CPX and PP242; five genes (EEF1A1, EEF2, EIF4B, FMO2, RAB3D) were modulated by pirfenidone, TSA and PP242; three genes (CIT, PDE4B, PPARG) were modulated by pirfenidone, TSA and silvestrol; one gene (SLC40A1) was modulated by pirfenidone, TSA and CPX; three genes (ASPM, CA5B, GLCCI1) were modulated by pirfenidone, PP242, silvestrol and TSA; two genes (GLTSCR2, P2RX7) were modulated by pirfenidone, PP242, CPX and TSA; and one gene (STAMBPL1) was modulated by pirfenidone, silvestrol, CPX and TSA.

As is evident, the modulated genes (*i.e.*, targets for drug development) include those that encode proteins of the translation machinery, regulators of the translation machinery, proteins that modify disease, and downstream proteins.

The various embodiments described above can be combined to provide further embodiments. All of the U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification and/or listed in the Application Data Sheet, including but not limited to U.S. Patent Application No. 61/937,272, United States Patent Application No. 62/010,004, and United States Patent Application No. 62/037,497, are incorporated herein by reference in their entirety. Aspects of the embodiments can be modified, if necessary, to employ concepts of the various patents, applications and publications to provide further embodiments.

These and other changes can be made to the embodiments in light of the above-detailed description. In general, in the following claims, the terms used should not be construed to limit the claims to the specific embodiments disclosed in the specification and the claims, but should be construed to include all possible embodiments along with
5 the full scope of equivalents to which such claims are entitled. Accordingly, the claims are not limited by the disclosure.

CLAIMS

What is claimed is:

1. A method for preventing, treating or ameliorating a fibrotic disease, comprising administering to a subject having a fibrotic disorder a therapeutically effective amount of a modulator specific for any one of the genes or encoded products listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5, Table 6, Table 7, an eIF2 component or regulator, an eIF4F complex or regulator, an eIF4F complex component or regulator, an eIF5A or regulator, or any combination thereof.

2. A method for reducing the risk of developing a fibrotic disease, comprising administering to a subject at risk of developing a fibrotic disorder a therapeutically effective amount of a modulator specific for any one of the genes or encoded products listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5, Table 6, Table 7, an eIF2 component or regulator, an eIF4F complex or regulator, an eIF4F complex component or regulator, an eIF5A or regulator, or any combination thereof.

3. A method for reducing myofibroblasts, comprising administering to a subject at risk of developing or having a fibrotic disorder a therapeutically effective amount of a modulator specific for any one of the genes or encoded products listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5, Table 6, Table 7, an eIF2 component or regulator, an eIF4F complex or regulator, an eIF4F complex component or regulator, an eIF5A or regulator, or any combination thereof.

4. A method for inhibiting or reversing transdifferentiation of a fibroblast to a myofibroblast, comprising administering to a subject at risk of developing or having a fibrotic disorder a therapeutically effective amount of a modulator specific for any one of the genes or encoded products listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5, Table 6, Table 7, an eIF2 component or regulator, an eIF4F complex or regulator, an eIF4F complex component or regulator, an eIF5A or regulator, or any combination thereof.

5. The method according to any one of claims 1-4, wherein the gene or encoded product comprises a translation machinery element, a regulator of a translation machinery element, or combinations thereof.

6. The method according to claim 5, wherein the translation machinery element comprises an eIF2 component, an eIF4F complex, an eIF4F complex component, eIF5A, rpS6, or any combination thereof.

7. The method according to claim 6, wherein the translation machinery element is eIF2 α , eIF2 β , eIF2 γ , eIF4A, eIF4E, eIF5A, rpS6, or any combination thereof.

8. The method according to claim 5, wherein the regulator of a translation machinery element comprises an EIF2AK1, EIF2AK2, EIF2AK3, EIF2AK4, mTOR, deoxyhypusine hydroxylase (DOHH), deoxyhypusine synthase (DHPS), histone deacetylase 6 (HDAC6), NAD-dependent deacetylase sirtuin-2 (SIRT2), p90 Ribosomal S6 kinase (RSK), adenosylhomocysteinase (AHCY), or any combination thereof.

9. The method according to any one of claims 1-4, wherein the gene or encoded product comprises (a) CREB5, DIAPH3, LGALS1, NACA, RPL12, RPL13A, RPL17, RPL21, RPL22L1, RPL23, RPL26, RPL27A, RPL28, RPL3, RPL30, RPL34, RPL36, RPL37, RPL37A, RPL4, RPL7A, RPS9, RPLP1, RPLP2, RPS10, RPS16, RPS19, RPS27, RPS5, RPS8, RPS9, SLC25A6, SOX6, STS, TKT or any combination thereof; (b) ABCA6, ANKH, CARD16, CEP192, DDX60, DNASE1L1, DYNC2H1, EDN1, HBEGF, HOMER1, INHBA, KDM6B, LENG9, MATN3, MYO19, NRG1, PABPC4, PLD1, PLEKHA5, RASD1, SGIP1, SLC2A12, SNRPA, TEN1, TOP2A, TRERF1 or any combination thereof; (c) CES1, LAMP5, PAQR5, PLEKHG1, ROBO2, TOMM7 or any combination thereof; or (d) AOX1, ARPC1A, AURKA, C12orf57, GPSM2, KITLG, MAP3K5, MURC, NOV, RPL14, SLC15A3, SOX5, ZNF608 or any combination thereof.

10. The method according to any one of claims 1-4, wherein the gene or encoded product comprises (a) ANKDD1A, ATP5G2, CHCHD10, DNAJC22, FGF5, FMO2, GNB2L1, GLTSCR2, HIGD2A, IFIH, MTUS1, RPS18, RPL18A, RPL31, RPL35A, RPL5, RPS18, RPS29, SPATA6 or any combination thereof; or (b) RPS6KA5, BIVM, ACTA1, KRT7, AMIGO3, CCDC102B, RPL10, TAF1D, ADAMTS5, LAMB3, CLCF1, EPB41L1, GAS2L3, IRAK3, LPAR3, PCBP2, PDE7B, TMTC1, FRMD4A, GDF10, OBSCN, PLEKHA6, SHC3 or any combination thereof.

11. The method according to any one of claims 1-4, wherein the gene or encoded product comprises THBS3, RPS28, EEF1A1, EEF2, EIF4B, FMO2, RAB3D, CIT, PDE4B, PPARG, SLC40A1, ASPM, CA5B, GLCCI1, GLTSCR2, P2RX7, STAMBPL1 or any combination thereof.

12. The method according to any one of claims 1 to 11, wherein the fibrotic disease is due to injury or is idiopathic.

13. The method according to claim 12, wherein the injury is an ischemic event or due to exposure to radiation, a chemical, or an infectious agent.

14. The method according to any one of claims 1 to 11, wherein the modulator is administered after a fibrotic lesion has developed in the subject.

15. The method according to any of the preceding claims, wherein the modulator is formulated with a pharmaceutically acceptable excipient.

16. The method according to any of the preceding claims, wherein the modulator is administered in combination with one or more adjunctive therapeutic agents.

17. The method according to claim 16, wherein the one or more adjunctive therapeutic agents is selected from angiotensin converting enzyme inhibitor, nintedanib,

STX-100, QAX576, CNTO-888, SD-208, SB-525334, GC1008, BMS-986202, AM152, lebrikizumab, tralokinumab, SAR156597, PRM-151, simtuzumab (AB0024, GS-6624), GSK2126458, FG-3019, captopril, genistein, silvestrol or derivatives thereof, pirfenidone, pateamine A or derivatives thereof, hippuristanol, or EUK-207.

18. The method according to any of the preceding claims, wherein the fibrotic disease is selected from pulmonary fibrosis, idiopathic pulmonary fibrosis, cystic fibrosis, liver fibrosis, cardiac fibrosis, endomyocardial fibrosis, atrial fibrosis, mediastinal fibrosis, myelofibrosis, retroperitoneal fibrosis, chronic kidney disease, nephrogenic systemic fibrosis, Chron's disease, hypertrophic scarring, keloid, scleroderma, organ transplant-associated fibrosis, or ischemia-associated fibrosis.

19. The method according to any of the preceding claims, wherein the subject is a human.

20. A method for identifying a candidate therapeutic for normalizing a translational profile associated with a fibrotic disease, comprising:

- (a) determining three independent translational profiles, each for a plurality of genes, wherein (i) a first translational profile is from a fibrotic disease sample, (ii) a second translational profile is from (1) a control non-diseased sample or (2) a control non-diseased sample contacted with a candidate agent, and (iii) a third translational profile is from the fibrotic disease sample contacted with a candidate agent;
- (b) determining a first differential translational profile comprising one or more genes differentially translated in the first translational profile as compared to the second translational profile, and determining a second differential translational profile comprising one or more genes differentially translated in the first translational profile as compared to the third translational profile, wherein the one or more differentially translated genes are selected from EIF2AK1, EIF2AK2, EIF2AK3, EIF2AK4, EIF5A, mTOR, DOHH, DHPS, HDAC6, SIRT2, RSK, AHCY, or the genes listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5 or Table 7; and

(c) identifying the agent as a candidate therapeutic for normalizing a translational profile associated with the fibrotic disease when the first differential translational profile is comparable to the second differential translational profile.

21. The method according to claim 20, wherein the gene or encoded product comprises a translation machinery element, a regulator of a translation machinery element, or combinations thereof.

22. The method according to claim 21, wherein the translation machinery element comprises an eIF2 component, an eIF4F complex, an eIF4F complex component, eIF5A, rpS6, or any combination thereof.

23. The method according to claim 22, wherein the translation machinery element is eIF2 α , eIF2 β , eIF2 γ , eIF4A, eIF4E, eIF5A, rpS6, or any combination thereof.

24. The method according to claim 21, wherein the regulator of a translation machinery element comprises an EIF2AK1, EIF2AK2, EIF2AK3, EIF2AK4, mTOR, deoxyhypusine hydroxylase (DOHH), deoxyhypusine synthase (DHPS), histone deacetylase 6 (HDAC6), NAD-dependent deacetylase sirtuin-2 (SIRT2), p90 Ribosomal S6 kinase (RSK), adenosylhomocysteinase (AHCY), or any combination thereof.

25. The method according to claim 20, wherein the gene or encoded product comprises (a) CREB5, DIAPH3, LGALS1, NACA, RPL12, RPL13A, RPL17, RPL21, RPL22L1, RPL23, RPL26, RPL27A, RPL28, RPL3, RPL30, RPL34, RPL36, RPL37, RPL37A, RPL4, RPL7A, RPS9, RPLP1, RPLP2, RPS10, RPS16, RPS19, RPS27, RPS5, RPS8, RPS9, SLC25A6, SOX6, STS, TKT or any combination thereof; (b) ABCA6, ANKH, CARD16, CEP192, DDX60, DNASE1L1, DYNC2H1, EDN1, HBEGF, HOMER1, INHBA, KDM6B, LENG9, MATN3, MYO19, NRG1, PABPC4, PLD1, PLEKHA5, RASD1, SGIP1, SLC2A12, SNRPA, TEN1, TOP2A, TRERF1 or

any combination thereof; (c) CES1, LAMP5, PAQR5, PLEKHG1, ROBO2, TOMM7 or any combination thereof; or (d) AOX1, ARPC1A, AURKA, C12orf57, GPSM2, KITLG, MAP3K5, MURC, NOV, RPL14, SLC15A3, SOX5, ZNF608 or any combination thereof.

26. The method according to claim 20, wherein the gene or encoded product comprises (a) ANKDD1A, ATP5G2, CHCHD10, DNAJC22, FGF5, FMO2, GNB2L1, GLTSCR2, HIGD2A, IFIH, MTUS1, RPS18, RPL18A, RPL31, RPL35A, RPL5, RPS18, RPS29, SPATA6 or any combination thereof; or (b) RPS6KA5, BIVM, ACTA1, KRT7, AMIGO3, CCDC102B, RPL10, TAF1D, ADAMTS5, LAMB3, CLCF1, EPB41L1, GAS2L3, IRAK3, LPAR3, PCBP2, PDE7B, TMTC1, FRMD4A, GDF10, OBSCN, PLEKHA6, SHC3 or any combination thereof.

27. The method according to claim 20, wherein the gene or encoded product comprises THBS3, RPS28, EEF1A1, EEF2, EIF4B, FMO2, RAB3D, CIT, PDE4B, PPARG, SLC40A1, ASPM, CA5B, GLCCI1, GLTSCR2, P2RX7, STAMBPL1 or any combination thereof.

28. The method according to any one of claims 20 to 27, wherein the fibrotic disease is selected from pulmonary fibrosis, idiopathic pulmonary fibrosis, cystic fibrosis, liver fibrosis, cardiac fibrosis, endomyocardial fibrosis, atrial fibrosis, mediastinal fibrosis, myelofibrosis, retroperitoneal fibrosis, chronic kidney disease, nephrogenic systemic fibrosis, Chron's disease, hypertrophic scarring, keloid, scleroderma, organ transplant-associated fibrosis, or ischemia-associated fibrosis.

29. A method for validating a target for normalizing a translational profile associated with a fibrotic disease, the method comprising:

(a) determining three independent translational profiles, each for a plurality of genes, wherein (i) a first translational profile is from a fibrotic disease sample, (ii) a second translational profile is from (1) a control non-diseased sample or (2) a control non-diseased sample contacted with an agent that modulates a target, and

(iii) a third translational profile is from the fibrotic disease sample contacted with the agent that modulates the target;

(b) determining a first differential translational profile comprising one or more genes differentially translated in the first translational profile as compared to the second translational profile, and determining a second differential translational profile comprising one or more genes differentially translated in the first translational profile as compared to the third translational profile, wherein the one or more differentially translated genes are selected from EIF2AK1, EIF2AK2, EIF2AK3, EIF2AK4, EIF5A, mTOR, DOHH, DHPS, HDAC6, SIRT2, RSK, AHCY, or the genes listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5, or Table 7; and

(c) validating the target as a target for normalizing a translational profile associated with the fibrotic disease when the first differential translational profile is comparable to the second differential translational profile.

30. The method according to claim 29, wherein the gene encodes a translation machinery element, a regulator of a translation machinery element, or combinations thereof.

31. The method according to claim 30, wherein the translation machinery element comprises an eIF2 component, an eIF4F complex, an eIF4F complex component, eIF5A, rpS6, or any combination thereof.

32. The method according to claim 31, wherein the translation machinery element is eIF2 α , eIF2 β , eIF2 γ , eIF4A, eIF4E, eIF5A, rpS6, or any combination thereof.

33. The method according to claim 30, wherein the regulator of a translation machinery element comprises an EIF2AK1, EIF2AK2, EIF2AK3, EIF2AK4, mTOR, deoxyhypusine hydroxylase (DOHH), deoxyhypusine synthase (DHPS), histone deacetylase 6 (HDAC6), NAD-dependent deacetylase sirtuin-2 (SIRT2), p90 Ribosomal S6 kinase (RSK), adenosylhomocysteinase (AHCY), or any combination thereof.

34. The method according to claim 29, wherein the gene is (a) CREB5, DIAPH3, LGALS1, NACA, RPL12, RPL13A, RPL17, RPL21, RPL22L1, RPL23, RPL26, RPL27A, RPL28, RPL3, RPL30, RPL34, RPL36, RPL37, RPL37A, RPL4, RPL7A, RPS9, RPLP1, RPLP2, RPS10, RPS16, RPS19, RPS27, RPS5, RPS8, RPS9, SLC25A6, SOX6, STS, TKT or any combination thereof; (b) ABCA6, ANKH, CARD16, CEP192, DDX60, DNASE1L1, DYNC2H1, EDN1, HBEGF, HOMER1, INHBA, KDM6B, LENG9, MATN3, MYO19, NRG1, PABPC4, PLD1, PLEKHA5, RASD1, SGIP1, SLC2A12, SNRPA, TEN1, TOP2A, TRERF1 or any combination thereof; (c) CES1, LAMP5, PAQR5, PLEKHG1, ROBO2, TOMM7 or any combination thereof; or (d) AOX1, ARPC1A, AURKA, C12orf57, GPSM2, KITLG, MAP3K5, MURC, NOV, RPL14, SLC15A3, SOX5, ZNF608 or any combination thereof.

35. The method according to claim 29, wherein the gene is (a) ANKDD1A, ATP5G2, CHCHD10, DNAJC22, FGF5, FMO2, GNB2L1, GLTSCR2, HIGD2A, IFIH1, MTUS1, RPS18, RPL18A, RPL31, RPL35A, RPL5, RPS18, RPS29, SPATA6 or any combination thereof; or (b) RPS6KA5, BIVM, ACTA1, KRT7, AMIGO3, CCDC102B, RPL10, TAF1D, ADAMTS5, LAMB3, CLCF1, EPB41L1, GAS2L3, IRAK3, LPAR3, PCBP2, PDE7B, TMTC1, FRMD4A, GDF10, OBSCN, PLEKHA6, SHC3 or any combination thereof.

36. The method according to claim 29, wherein the gene is THBS3, RPS28, EEF1A1, EEF2, EIF4B, FMO2, RAB3D, CIT, PDE4B, PPARG, SLC40A1, ASPM, CA5B, GLCCI1, GLTSCR2, P2RX7, STAMBPL1 or any combination thereof.

37. The method according to any one of claims 21 to 36, wherein the fibrotic disease is selected from pulmonary fibrosis, idiopathic pulmonary fibrosis, cystic fibrosis, liver fibrosis, cardiac fibrosis, endomyocardial fibrosis, atrial fibrosis, mediastinal fibrosis, myelofibrosis, retroperitoneal fibrosis, chronic kidney disease, nephrogenic systemic fibrosis, Chron's disease, hypertrophic scarring, keloid, scleroderma, organ transplant-associated fibrosis, or ischemia-associated fibrosis.

38. A method of identifying a subject as a candidate for treating a fibrotic disease with a therapeutic agent, the method comprising:

(a) determining a first translational profile for a plurality of genes in a sample from a subject having or suspected of having a fibrotic disease;

(b) determining a second translational profile for a plurality of genes in a control sample, wherein the control sample is from a subject known to respond to the therapeutic agent and wherein the sample has not been contacted with the therapeutic agent; and

(c) identifying the subject as a candidate for treating fibrotic disease with the therapeutic agent when the translational profile for one or more genes selected from EIF2AK1, EIF2AK2, EIF2AK3, EIF2AK4, EIF5A, mTOR, DOHH, DHPS, HDAC6, SIRT2, RSK, AHCY, or the genes listed in Figure 7, Table 1, Table 3A, Table 3B, Table 5 or Table 7 of the first translational profile are comparable to the translational profile of the corresponding genes in the second translational profile.

39. A method for treating a fibrotic disease, comprising administering a therapeutic agent to a subject identified according to the method of claim 14, thereby treating the subject.

40. The method according to claim 38 or 39, wherein the gene encodes a translation machinery element, a regulator of a translation machinery element, or combinations thereof.

41. The method according to claim 40, wherein the translation machinery element comprises an eIF2 component, an eIF4F complex, an eIF4F complex component, eIF5A, rpS6, or any combination thereof.

42. The method according to claim 41, wherein the translation machinery element is eIF2 α , eIF2 β , eIF2 γ , eIF4A, eIF4E, eIF5A, rpS6, or any combination thereof.


43. The method according to claim 40, wherein the regulator of a translation machinery element comprises an EIF2AK1, EIF2AK2, EIF2AK3, EIF2AK4, mTOR, deoxyhypusine hydroxylase (DOHH), deoxyhypusine synthase (DHPS), histone deacetylase 6 (HDAC6), NAD-dependent deacetylase sirtuin-2 (SIRT2), p90 Ribosomal S6 kinase (RSK), adenosylhomocysteinase (AHCY), or any combination thereof.

44. The method according to claim 38 or 39, wherein the gene is (a) CREB5, DIAPH3, LGALS1, NACA, RPL12, RPL13A, RPL17, RPL21, RPL22L1, RPL23, RPL26, RPL27A, RPL28, RPL3, RPL30, RPL34, RPL36, RPL37, RPL37A, RPL4, RPL7A, RPS9, RPLP1, RPLP2, RPS10, RPS16, RPS19, RPS27, RPS5, RPS8, RPS9, SLC25A6, SOX6, STS, TKT or any combination thereof; (b) ABCA6, ANKH, CARD16, CEP192, DDX60, DNASE1L1, DYNC2H1, EDN1, HBEGF, HOMER1, INHBA, KDM6B, LENG9, MATN3, MYO19, NRG1, PABPC4, PLD1, PLEKHA5, RASD1, SGIP1, SLC2A12, SNRPA, TEN1, TOP2A, TRERF1 or any combination thereof; (c) CES1, LAMP5, PAQR5, PLEKHG1, ROBO2, TOMM7 or any combination thereof; or (d) AOX1, ARPC1A, AURKA, C12orf57, GPSM2, KITLG, MAP3K5, MURC, NOV, RPL14, SLC15A3, SOX5, ZNF608 or any combination thereof.

45. The method according to claim 38 or 39, wherein the gene is (a) ANKDD1A, ATP5G2, CHCHD10, DNAJC22, FGF5, FMO2, GNB2L1, GLTSCR2, HIGD2A, IFIH, MTUS1, RPS18, RPL18A, RPL31, RPL35A, RPL5, RPS18, RPS29, SPATA6 or any combination thereof; or (b) RPS6KA5, BIVM, ACTA1, KRT7, AMIGO3, CCDC102B, RPL10, TAF1D, ADAMTS5, LAMB3, CLCF1, EPB41L1, GAS2L3, IRAK3, LPAR3, PCBP2, PDE7B, TMTC1, FRMD4A, GDF10, OBSCN, PLEKHA6, SHC3 or any combination thereof.

46. The method according to claim 38 or 39, wherein the gene is THBS3, RPS28, EEF1A1, EEF2, EIF4B, FMO2, RAB3D, CIT, PDE4B, PPARG, SLC40A1, ASPM, CA5B, GLCCI1, GLTSCR2, P2RX7, STAMBPL1 or any combination thereof.

47. The method according to any one of claims 38 to 46, wherein the fibrotic disease is selected from pulmonary fibrosis, idiopathic pulmonary fibrosis, cystic fibrosis, liver fibrosis, cardiac fibrosis, endomyocardial fibrosis, atrial fibrosis, mediastinal fibrosis, myelofibrosis, retroperitoneal fibrosis, chronic kidney disease, nephrogenic systemic fibrosis, Chron's disease, hypertrophic scarring, keloid, scleroderma, organ transplant-associated fibrosis, or ischemia-associated fibrosis.

1 / 28**PIPC after 24 hr. treatment*****Fig. 1***






Fig. 2


3 / 28***Fig. 3***

4 / 28

Early signaling events in hepatic stellate cells

**Fig. 4A**

5 / 28

**Fig. 4A (Cont. 1)**

6 / 28

CXCL9 is one of the only chemokines
to exert an antifibrotic effect in liver
fibrosis.






Fig. 4A (Cont. 2)

7 / 28


Signaling events in activated HSC (Myofibroblasts)

**Fig. 4B**

8 / 28

**Fig. 4B (Cont. 1)**

9 / 28


**Fig. 4B (Cont. 2)**

Target	Log Ratio	p-value	Target	Log Ratio	p-value
A2M	-2.075	9.99E-15	IGFBP3	5.015	1.45E-81
ACTA2	6.372	1.20E-119	IGFBP4	-1.687	1.08E-09
CCL2	-4.057	1.99E-19	IL1R1	-1.642	6.72E-11
COL11A1	2.713	7.58E-08	IL6R	-3.584	1.36E-12
COL16A1	1.720	5.95E-09	MYH11	5.684	1.68E-15
COL1A1	2.043	3.42E-15	MYL6	1.884	3.00E-11
COL4A1	3.940	9.03E-53	MYL9	2.533	3.22E-21
COL4A2	2.283	8.14E-22	NGFR	7.698	9.57E-77
COL4A5	-2.099	2.65E-11	PDGFA	3.938	2.26E-48
COL4A6	-1.813	3.76E-07	PDGFD	-1.776	5.14E-08
COL5A1	2.380	5.70E-23	PDGFRA	-2.279	8.12E-21
COL7A1	2.913	6.35E-20	SERPINE1	5.579	5.79E-36
COL8A1	2.167	1.22E-18	SMAD3	-3.521	5.50E-37
COL8A2	4.032	3.35E-50	SMAD7	2.034	2.33E-16
CTGF	2.866	7.18E-11	TGFB1	2.028	2.67E-14
EDN1	6.424	4.38E-14	TGFBR1	2.337	1.12E-11
FGF1	3.101	4.50E-14	TNFRSF11B	-2.172	8.44E-15
HGF	-3.538	4.77E-33	TNFRSF1B	-5.279	1.80E-23
ICAM1	-3.106	3.97E-13	VCAM1	-4.070	4.41E-27
IGF1	5.316	7.89E-14			

Fig. 4C

11 / 28

Early signaling events in hepatic stellate cells

**Fig. 5A**

12 / 28




Fig. 5A (Cont. 1)

13 / 28

CXCL9 is one of the only chemokines
to exert an antifibrotic effect in liver
fibrosis.






Fig. 5A (Cont. 2)

14 / 28


Signaling events in activated HSC (Myofibroblasts)

**Fig. 5B**

15 / 28

**Fig. 5B (Cont. 1)**

16 / 28

**Fig. 5B (Cont. 2)**

17 / 28

Target	Log Ratio	p-value	Target	Log Ratio	p-value
A2M	-2.116	4.00E-19	IGFBP3	5.216	1.76E-189
ACTA2	5.893	4.36E-112	IGFBP4	-1.832	1.23E-42
CCL2	-3.854	4.18E-55	IL1R1	-2.645	1.03E-24
COL10A1	3.007	2.25E-15	IL6R	-3.772	1.94E-34
COL11A1	3.482	9.18E-67	MMP1	-1.889	2.47E-27
COL16A1	1.938	1.19E-20	MYH9	1.547	1.64E-14
COL1A1	1.949	7.64E-15	MYH10	1.657	2.51E-09
COL21A1	-2.991	2.12E-08	MYH11	5.878	5.64E-29
COL4A1	3.869	3.22E-27	MYL6	2.088	1.31E-31
COL4A2	2.700	3.90E-23	MYL9	3.119	1.52E-99
COL4A5	-2.181	6.65E-28	NGFR	7.071	4.55E-151
COL4A6	-2.073	2.51E-29	PDGFA	4.505	1.93E-31
COL5A1	2.585	6.00E-21	PDGFD	-2.805	1.14E-14
COL7A1	3.575	1.25E-26	PDGFRA	-2.432	2.65E-42
COL8A1	2.002	4.63E-38	SERPINE1	5.366	9.74E-49
COL8A2	4.162	1.08E-44	SMAD3	-3.835	3.47E-84
COL9A3	3.158	1.42E-10	SMAD7	2.327	2.73E-26
CTGF	3.338	5.65E-48	TGFB1	2.660	5.39E-42
EDN1	7.236	8.49E-28	TGFBR1	2.257	2.86E-21
EDNRA	-1.548	3.47E-17	TGFBR2	-1.579	3.68E-34
EDNRB	-2.445	9.96E-11	TNFRSF11B	-2.063	3.91E-12
FGF1	3.778	9.42E-17	TNFRSF1B	-5.364	4.95E-35
FLT1	-1.628	4.94E-08	TNFSF10	-3.807	6.07E-14
HGF	-4.352	2.91E-181	TNFSF18	-2.908	1.92E-13
ICAM1	-3.363	6.64E-28	VCAM1	-4.582	6.15E-56
IGF1	5.659	3.06E-23	VEGFA	1.598	9.18E-24

Fig. 5C




Fig. 6A




Fig. 6A (Cont.)

20 / 28

Target	Log Ratio	p-value	Target	Log Ratio	p-value	Target	Log Ratio	p-value
EIF1	0.784	5.21E-04	RPL28	1.869	1.81E-10	RPLP2	1.694	5.38E-04
EIF2S3	1.067	7.04E-03	RPL29	1.589	3.21E-03	RPS5	1.533	5.18E-03
EIF3E	1.469	2.87E-04	RPL30	1.622	2.85E-03	RPS6	1.598	5.99E-03
EIF3F	1.290	3.50E-03	RPL31	1.600	7.78E-03	RPS7	1.549	2.77E-03
EIF3H	1.329	2.47E-04	RPL32	1.916	4.84E-04	RPS8	1.611	4.90E-03
EIF3L	1.429	3.34E-03	RPL34	1.608	2.48E-03	RPS9	1.614	2.08E-03
PABPC1	1.207	3.97E-04	RPL36	1.411	3.67E-03	RPS13	1.539	1.15E-03
RPL3	1.780	7.43E-04	RPL37	1.808	3.00E-03	RPS14	1.684	3.99E-03
RPL4	1.688	3.28E-03	RPL38	1.353	6.96E-03	RPS15	1.300	2.24E-04
RPL5	1.593	3.92E-04	RPL39	1.536	5.22E-03	RPS16	1.989	8.68E-06
RPL8	1.353	4.15E-03	RPL41	1.474	7.12E-04	RPS18	2.003	2.93E-03
RPL9	1.613	5.38E-03	RPL10A	1.510	7.60E-03	RPS19	1.550	5.37E-03
RPL10	1.691	9.23E-04	RPL13A	1.797	4.22E-03	RPS21	1.846	4.80E-04
RPL11	1.396	5.38E-03	RPL18A	1.669	3.01E-03	RPS24	1.444	3.41E-03
RPL12	1.878	1.27E-03	RPL22L1	1.012	4.48E-03	RPS25	1.697	1.41E-03
RPL14	1.745	1.25E-04	RPL23A	1.395	9.73E-03	RPS26	1.090	8.58E-03
RPL15	1.254	2.26E-03	RPL27A	2.090	2.38E-04	RPS28	1.937	5.50E-05
RPL18	1.612	2.08E-03	RPL35A	1.627	4.84E-04	RPS29	1.571	7.82E-03
RPL23	1.604	7.28E-03	RPL36AL	0.794	8.34E-03	RPS15A	2.195	1.84E-04
RPL24	1.304	7.67E-03	RPL37A	2.029	1.12E-03	RPS27A	1.275	6.09E-03
RPL26	1.655	7.91E-03	RPL7A	1.650	1.93E-03	RPS4X	1.534	8.76E-03
RPL27	1.519	3.25E-03	RPLP1	1.461	8.30E-03	RPLP2	1.694	5.38E-04

Fig. 6B




Fig. 7






Fig. 8

23 / 28**Fig. 9**

24 / 28***Fig. 10***







Fig. 11

26 / 28

**Fig. 12**

27 / 28**Fig. 13**

28 / 28

***Fig. 14***