

(19) DANMARK

(10) DK/EP 2224954 T3



(12)

Oversættelse af  
europæisk patent

Patent- og  
Varemærkestyrelsen

---

(51) Int.Cl.: **A 61 K 39/395 (2006.01)** **A 61 P 35/00 (2006.01)** **C 07 K 16/28 (2006.01)**

(45) Oversættelsen bekendtgjort den: **2014-03-03**

(80) Dato for Den Europæiske Patentmyndigheds  
bekendtgørelse om meddelelse af patentet: **2014-01-08**

(86) Europæisk ansøgning nr.: **08848354.0**

(86) Europæisk indleveringsdag: **2008-11-07**

(87) Den europæiske ansøgnings publiceringsdag: **2010-09-08**

(86) International ansøgning nr.: **US2008082745**

(87) Internationalt publikationsnr.: **WO2009061996**

(30) Prioritet: **2007-11-07 US 2253 P** **2007-11-07 US 2253**  
**2008-09-10 US 191551 P** **2008-09-10 US 191551**

(84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC**  
**MT NL NO PL PT RO SE SI SK TR**

(73) Patenthaver: **Celldex Therapeutics, Inc., 119 Fourth Avenue, Needham, MA 02494-2725, USA**

(72) Opfinder: **KELER, Tibor, 30 Park Road, Ottsville, PA 18942, USA**  
**HE, Lizhen, 1675 Ridgeview Drive, Allentown, PA 18104, USA**  
**RAMAKRISHNA, Venky, c/o Celldex Therapeutics, Inc., 19 Fourth Avenue, Needham, MA 02494-2725, USA**  
**VITALE, Laura A., 4194 Milords Lane, Doylestown, PA 18901, USA**

(74) Fuldmægtig i Danmark: **HJERRILD & LEVIN A/S, Vedbæk Strandvej 341, 2950 Vedbæk, Danmark**

(54) Benævnelse: **Antistoffer, der binder human dendritisk og epitelcelle 205 (DEC-205)**

(56) Fremdragne publikationer:  
**WO-A-2004/035619**  
**WO-A-2004/074432**  
**WO-A-2005/018610**  
**US-A1- 2004 258 688**  
**US-A1- 2005 186 612**  
**BADIEE ET AL: "Enhanced delivery of immunoliposomes to human dendritic cells by targeting the multilectin receptor DEC-205" VACCINE, BUTTERWORTH SCIENTIFIC, GUILDFORD, GB, vol. 25, no. 25, 30 May 2007 (2007-05-30), pages 4757-4766, XP022098619 ISSN: 0264-410X**  
**GUO M ET AL: "A monoclonal antibody to the DEC-205 endocytosis receptor on human dendritic cells" HUMAN IMMUNOLOGY, NEW YORK, NY, US, vol. 61, no. 8, 1 August 2000 (2000-08-01), pages 729-738, XP002319045 ISSN: 0198-8859**

**DK/EP 2224954 T3**

**Description****Related Applications**

5 [0001] This application claims priority to U.S. Provisional Application No.: 61/002253, filed on November 7, 2007 and U.S. Provisional Application No.: 61/191551, filed on September 10, 2008.

**Background of the Invention**

10 [0002] Dendritic cells (DCs) are specialized cells of the immune system. DCs have the unique capacity for initiating primary and secondary T and B lymphocyte responses by presenting antigens in the form of peptides bound to cell-surface major histocompatibility complex (MHC) molecules. The antigen-presentation function of dendritic cells has been correlated with the high-level expression of human dendritic and epithelial cell 205 receptor (DEC-205) (Jiang et al. (1995) *Nature* 375(11)151).

15 [0003] DEC-205 is an endocytic receptor found primarily on dendritic cells, but is also found on B cells, brain capillaries, bone marrow stroma, epithelia of intestinal villi and pulmonary airways, as well as the cortical epithelium of the thymus and the dendritic cells in the T cell areas of peripheral lymphoid organs. DEC-205 is expressed at high levels on DCs in the T cell areas of lymphoid organs (Kraal et al. (1986) *J. Exp. Med.* 163:981; Witmer-Pack et al. (1995) *Cell. Immunol.* 163:157). DEC-205 has ten membrane-external, contiguous C-type lectin domains (*Id.*; Mahnke et al. (2000) *J. Cell Biol.* 151:673) which mediate the efficient processing and presentation of antigens on MHC class II products *in vivo* (Hawiger et al. (2001) *J. Exp. Med.* 194:769). It has been shown that small amounts of injected antigen, targeted to DCs by the DEC-205 adsorptive pathway, are able to induce solid peripheral CD8<sup>+</sup> T cell tolerance (Bonifaz et al. (2002) *J. Exp. Med.* 196(12):1627).

20 [0004] Despite recent advances in the characterization of dendritic cells, very little is known regarding dendritic cell-specific receptors, such as DEC-205, and few reagents are available which are specific to dendritic cells. Reagents, in particular antibodies, which react specifically or preferentially with dendritic cells, such as through DEC-205, have great potential as targeting agents to induce potent immune responses to tumor or infectious disease antigens. These cell-specific targeting agents could also be engineered to deliver toxins to eliminate potent antigen presenting cells (e.g., dendritic cells) in bone marrow and organ transplantations or other autoimmune disorders. Accordingly, such dendritic 25 cell-specific binding agents possess great therapeutic and diagnostic value.

**Summary of the Invention**

30 [0005] The present invention provides an isolated monoclonal antibody which binds to human Dendritic and Epithelial Cell 205 receptor (DEC-205) and comprises: a heavy chain variable region CDR1 comprising SEQ ID NO: 29; a heavy chain variable region CDR2 comprising SEQ ID NO:30; a heavy chain variable region CDR3 comprising SEQ ID NO: 31; a light chain variable region CDR1 comprising SEQ ID NO: 35; a light chain variable region CDR2 comprising SEQ ID NO: 36; and a light chain variable region CDR3 comprising SEQ ID NO: 37. The present invention also provides vaccine conjugates, bispecific molecules, and therapeutic compositions containing such antibodies. Accordingly, the 35 antibodies and compositions of the invention can be used in a variety of dendritic cell-targeted therapies, for example, to enhance antigen presentation and/or induce T cell responses, such as cytotoxic T cell (CTL) responses, against a variety of target cells or pathogens, or to treat antigen presenting cell (APC)-mediated diseases.

40 [0006] In one embodiment, the antibodies of the present invention are human monoclonal antibodies that bind to human DEC-205 with an affinity constant of at least 10<sup>8</sup> M<sup>-1</sup> as measured by surface plasmon resonance and optionally 45 exhibit at least one of the following properties: (a) internalization after binding to human dendritic cells expressing DEC-205; (b) generation or enhancement of human T-cell responses to an antigen (which may be linked to the antibody), suitably mediated by either MHC Class I and/or Class II pathways; (c) generation or enhancement of human CTL or NKT responses to an antigen; (d) localization to antigen processing compartments in dendritic cells; and (e) induction of peripheral CD8<sup>+</sup> T cell tolerance; or (f) binding to an epitope located on the extracellular domain of human DEC-205, 50 for example, on one or a combination of the cysteine rich domain, the FnII domain, or one or more of the ten C-type lectin-like domains. Furthermore, the antibodies may cross-react with DEC-205 on non-human primate dendritic cells or those of other species.

55 [0007] In a further embodiment, isolated antibodies of the invention bind to human DEC-205 and include a heavy chain variable region and a light chain variable region including the amino acid sequences of SEQ ID NOs: 28 and 34, respectively.

[0008] Isolated antibodies which include heavy and light chain variable regions having at least 80%, or at least 85%, or at least 90%, or at least 95%, or at least 96%, or at least 97%, or at least 98%, or at least 99%, or more sequence identity to any of the above sequences are also included in the present invention. Ranges intermediate to the above-

recited values, e.g., heavy and light chain variable regions having at least 80-85%, 85-90%, 90-95% or 95-100% sequence identity to any of the above sequences are also intended to be encompassed by the present invention.

[0009] The isolated antibody may include a heavy chain variable region comprising an amino acid sequence of SEQ ID NO: 28, or sequences where at least one amino acid residue in the framework region of the heavy chain variable region is substituted with the corresponding germline residue. The antibody may further include a light chain variable region comprising an amino acid sequence of SEQ ID NO: 34, or sequences where at least one amino acid residue in the framework region of the light chain variable region is substituted with the corresponding germline residue. The substituted amino acid residue can include: a residue that non-covalently binds antigen directly; a residue adjacent to a CDR; a CDR-interacting residue; a residue participating in the VL-VH interface, a canonical residue, a vernier zone residue, or an interchain packing residue.

[0010] The antibodies of the invention can either be full-length, for example, any of the following isotypes: IgG1, IgG2, IgG3, IgG4, IgM, IgA1, IgA2, IgAsec, IgD, and IgE. Alternatively, the antibodies can be fragments such as an antigen-binding portion or a single chain antibody (e.g., a Fab, F(ab')<sub>2</sub>, Fv, a single chain Fv fragment, an isolated complementarity determining region (CDR) or a combination of two or more isolated CDRs).

[0011] The invention also provides a molecular conjugate comprising an antibody of the invention linked to an antigen (including fragments, epitopes and antigenic determinants), such as component of a pathogen, a tumor antigen or an autoantigen. For example, the antigen may include a tumor antigen, such as  $\beta$ hCG, gp100 or Pmel17, CEA, gp100, TRP-2, NY-BR-1, NY-CO-58, MN (gp250), idiotype, Tyrosinase, Telomerase, SSX2, MUC-1, MAGE-A3, and high molecular weight-melanoma associated antigen (HMW-MAA) MART1, melan-A, NY-ESO-1, MAGE-1, MAGE-3, WT1, Her2, mesothelin or high molecular weight-melanoma associated antigen (HMW-MAA).

[0012] The term "tumor antigen" as used herein preferably means any antigen or antigenic determinant which is present on (or associated with) a tumor cell and not typically on normal cells, or an antigen or antigenic determinant which is present on or associated with tumor cells in greater amounts than on normal (non-tumor) cells, or an antigen or antigenic determinant which is present on tumor cells in a different form than that found on normal (non-tumor) cells. The term thus includes tumor-specific antigens including tumor-specific membrane antigens, tumor-associated antigens, including tumor-associated membrane antigens, embryonic antigens on tumors, growth factor receptors, growth factor ligands, and any other type of antigen that is associated with cancer. A tumor antigen may be, for example, an epithelial cancer antigen, (e.g., breast, gastrointestinal, lung), a prostate specific cancer antigen (PSA) or prostate specific membrane antigen (PSMA), a bladder cancer antigen, a lung (e.g., small cell lung) cancer antigen, a colon cancer antigen, an ovarian cancer antigen, a brain cancer antigen, a gastric cancer antigen, a renal cell carcinoma antigen, a pancreatic cancer antigen, a liver cancer antigen, an esophageal cancer antigen, a head and neck cancer antigen, or a colorectal cancer antigen.

[0013] The term "fragment" refers to an amino acid sequence that is a portion of a full-length protein or polypeptide, for example between about 8 and about 1500 amino acids in length, suitably between about 8 and about 745 amino acids in length, suitably about 8 to about 300, for example about 8 to about 200 amino acids, or about 10 to about 50 or 100 amino acids in length.

[0014] In another embodiment, the molecular complex further includes a therapeutic agent, such as a cytotoxic agent, an immunosuppressive agent, or a chemotherapeutic agent.

[0015] The invention also provides a bispecific molecule comprising an antibody of the invention linked to a second functional moiety having a different binding specificity than said antibody.

[0016] Compositions including an antibody, a molecular conjugate or a bispecific molecule described herein, and a pharmaceutically effective carrier, are also provided. The compositions may further include a therapeutic agent (e.g., an immunosuppressive agent or an antibody different from an antibody of the invention).

[0017] Nucleic acid molecules encoding the antibodies of the invention are also encompassed by the invention, as well as expression vectors comprising such nucleic acids and host cells comprising such expression vectors. Moreover, the invention provides a transgenic mouse comprising human immunoglobulin heavy and light chain transgenes, wherein the mouse expresses an antibody of the invention, as well as hybridomas prepared from such a mouse, wherein the hybridoma produces the antibody of the invention.

[0018] Antibodies of the present invention are useful in methods for targeting an antigen to a cell, e.g., a cell capable of antigen presentation (such as peripheral blood mononuclear cells (PBMC), monocytes (such as THP-1), B lymphoblastoid cells (such as C1R.A2, 1518 B-LCL) and monocyte-derived DCs in a subject by administering a molecule which binds a receptor on the cell (e.g., the previously described DEC-205 antibodies) linked to an antigen. The targeted cell (which may be a B-cell) may stimulate MHC Class I restricted T-cells.

[0019] The antibodies and other compositions of the present invention can also be used to induce or enhance an immune response (e.g., a T cell-mediated immune response) against an antigen in a subject. Accordingly, in one embodiment, the present invention provides a method for inducing or enhancing a CTL response against an antigen by forming a conjugate of the antigen and a antibody which binds to a receptor on an antigen presenting cell, e.g., human DEC-205. The conjugate is then contacted, either *in vivo* or *ex vivo*, with cells expressing human DEC-205 such that

the antigen is internalized, processed and presented to T cells in a manner which induces or enhances a CTL response (e.g., a response mediated by CD8<sup>+</sup> cytotoxic T cells) against the antigen. In another embodiment, this serves also to induce a helper T cell response (e.g., a response mediated by CD4<sup>+</sup> helper T cells) against the antigen. Thus, the immune response may be induced through both MHC class I and MHC class II pathways. The cells expressing DEC-205 5 can also be contacted with an adjuvant, a cytokine which stimulates proliferation of dendritic cells, and/or an immunostimulatory agent to further enhance the immune response.

[0020] In another embodiment, methods of detecting the presence of DEC-205, or a cell expressing DEC-205, in a sample are provided by: (a) contacting the sample with the antibody of the invention under conditions that allow for formation of a complex between the antibody and DEC-205; and (b) detecting the formation of a complex between the antibody and DEC-205 in the sample.

[0021] Also within the scope of the invention are kits comprising the compositions (e.g., antibodies, molecular conjugates, multispecific and bispecific molecules) of the invention and, optionally, instructions for use. The kit can further contain a least one additional reagent, such as a cytokine or complement, or one or more additional human antibodies of the invention (e.g., a human antibody having a complementary activity which binds to an epitope on dendritic cells 15 distinct from the first human antibody).

[0022] Other features and advantages of the instant invention will be apparent from the following detailed description and claims.

#### Brief Description of the Drawings

[0023]

Figures 1A-1I include graphs showing the binding of human anti-DEC-205 antibodies (3D6-2F4, 3D6-4C8, 3G9-2D2, 5A8-1F1, 2D3-1F5-2A9, 3C7-3A3, 5D12-5G1, 1G6-1G6 and 3A4-1C10) to CHO-S cells expressing human DEC-205 by fluorescence analysis using a LSR<sup>TM</sup> instrument (BD Biosciences, NJ, USA).

Figures 2A-2I include graphs showing the binding of human anti-DEC-205 antibodies (3D6-2F4, 3D6-4C8, 3G9-2D2, 5A8-1F1, 2D3-1F5-2A9, 3C7-3A3, 5D12-5G1, 1G6-1G6 and 3A4-1C10) to DEC-205 on human dendritic cells by flow cytometry.

Figure 3 is a graph showing the binding of human anti-DEC-205 antibodies (3D6-2F4, 3D6-4C8, 3G9-2D2, 5A8-1F1, 2D3-1F5-2A9, 3C7-3A3, 5D12-5G1, 1G6-1G6 and 3A4-1C10) to DEC-205 using ELISA.

Figures 4A-4C show internalization into the dendritic cells of FITC-labelled HuMab (FITC-3G9-2D2) compared to the control (FITC-human IgG1) using confocal microscopy.

Figure 5 is an alignment of human VH and VK Germline Sequences with VH and VK sequences of anti-DEC-205 antibodies (3D6-2F4, 3D6-4C8, 3G9-2D2, 5A8-1F1, 2D3-1F5-2A9, 3C7-3A3, 5C3-2-3F6, 1E6-3D10). Figure discloses SEQ ID NOS 92, 34, 46, 58, 93, 82, 22, 94, 10, 95, 4, 16, 103-105, 76, 88, 96, 106 and 70, respectively, in order of appearance.

Figure 6 shows alignments of VH CDR1, CDR2 and CDR3 sequences of human anti-DEC-205 antibodies (3D6-2F4, 3D6-4C8, 3G9-2D2, 5A8-1F1, 3C7-3A3, 2D3-1F5-2A9, 1E6-3D10, 5C3-2-3F6, 5D12-5G1).

Figure 7 shows alignments of human anti-DEC-205 HuMab VK CDR1, CDR2 and CDR3 sequences of human anti-DEC-205 antibodies (3D6-2F4, 3D6-4C8, 3G9-2D2, 5A8-1F1, 3C7-3A3, 5C3-2-3F6).

Figure 8 shows a schematic representation of an example of an anti-DEC-205/antigen fusion APC targeted vaccine construct.

Figures 9A and B include graphs showing antigen-specific activity using 3G9-βhCG APC-targeted vaccine conjugate in peripheral blood mononuclear cells (PBMC), monocytes (THP-1), B lymphoblastoid cells (C1R.A2, 1518 B-LCL) and monocyte-derived DCs.

#### Detailed Description of the Invention

[0024] The present invention provides antibodies (e.g., human antibodies) which bind to human DEC-205. In certain embodiments, the antibodies exhibit a variety of functional properties, e.g., binding to human DEC-205 with an affinity

## EP 2 224 954 B1

constant of at least  $10^8$  M<sup>-1</sup> as measured by surface plasmon resonance, internalization after binding to human dendritic cells expressing DEC-205, generating or enhancing human T-cell responses, for example CD4+ or CD8<sup>+</sup> (CTL) or NKT cell responses, to an antigen which may be linked to the antibody, e.g., CTL responses mediated by both MHC Class I and Class II pathways; localization to antigen processing compartments in dendritic cells; induction of peripheral CD8<sup>+</sup> T cell tolerance; or cross-reaction with DEC-205 on non-human primate dendritic cells or those of other species. In other embodiments, the antibodies include heavy and light chain variable regions which utilize particular human germline genes and include particular structural features such as, particular CDR sequences. The invention further provides methods of making such antibodies, molecular conjugates and bispecific molecules including such antibodies, as well as compositions containing the antibodies. The invention also provides methods of targeting antigens to antigen presenting cells (e.g., peripheral blood mononuclear cells (PBMC), monocytes (such as THP-1), B lymphoblastoid cells (such as C1R.A2, 1518 B-LCL) and monocyte-derived DCs either *in vitro* or *in vivo*, for example, by using the anti-DEC-205 antibodies of the present invention. Methods of the present invention also include methods of inducing and enhancing an immune response (e.g., a T cell-mediated immune response) against an antigen in a subject. Such methods include the presentation of the antigen *via* a receptor on an antigen presenting cell (e.g., DEC-205) as a component of an MHC-I and/or MHC-II conjugate (e.g., the T cell response is mediated by both CD4+ and CD8+ T cells or by cytotoxic T cells or helper T cells). In one embodiment the targeted cell (which may be a B-cell) stimulates MHC Class I restricted T-cells.

**[0025]** In order that the present invention may be more readily understood, certain terms are first defined. Additional definitions are set forth throughout the detailed description.

**[0026]** The term "human Dendritic and Epithelial Cell 205 receptor" (DEC-205) includes any variants or isoforms of DEC-205 which are naturally expressed by cells (e.g., human DEC-205 deposited with GENBANK® having accession no. AAC17636, and mouse DEC-205 deposited with GENBANK® having accession no. AAL81722). Accordingly, human antibodies of the invention may cross-react with DEC-205 from species other than human. Alternatively, the antibodies may be specific for human DEC-205 and may not exhibit any cross-reactivity with other species. DEC-205 or any variants and isoforms thereof, may either be isolated from cells or tissues which naturally express them (e.g. human, mouse and cynomolgous monkey cells) or be recombinantly produced using well-known techniques in the art and/or those described herein.

Genbank® (Accession No. AAC17636A) reports the amino acid sequence of human DEC-205 as follows (SEQ ID NO: 1):

30 1 mrtgwatprr pagllmlfw ffdlaepsga aandptivh gntgkckpv ygwivaddcd  
 61 etedklwkvv sqhrlfhlhs qkclgllditk svnelrmfsc dssamlwwkc ehhslygaar  
 121 yrlalkdghg taisnasdvw kkggseeslc dqpyheiyytr dgnsygrpce fpflidgtwh  
 181 hdcildedhs gpwcattlny eydrkwgicl kpengcednw ekneqfgscy qfntqtalsw  
 241 keavyvscqnq gadllsinsa aeltylkeke giakifwigl nqlysargwe wsdhkplnfl

40

45

50

55

301 nwdpdrpsap tiggssarm daesglwqsf sceaqlpvc rkplnntvel tdvwtystr  
 361 cdagwlpnng fcylvnnesn swdkahakck afssdlisih sladvevvvt klhnedikee  
 5 421 vwiglnini ptlfqwsdgt evltywden epnvpynktp ncvsylgelg qwkvqsceek  
 481 lkyvckrkge kldassdkm cppdegwkrh getcykiyed evpfgtncln titsrfeqey  
 541 lndlmkkydk slrkyfwtgl rdvdscgeyn watvgrrra vtfsnwnfle paspggcavam  
 10 601 stgksvgkwe vkdcrsfkal sickkmsgpl gpeeaspkpd dpcpegwqsf paslscykvf  
 661 haerivrkrn weeaerfcqa lgahlssfsh vdeikeflhf ltdqfsgqhw lwiglnkrsp  
 721 dlqgswqwsd rtpvstiimp nefqqdydir dcaavkvfhr pwrrgwhfyd drefiylrpf  
 15 781 acdtklewvc qipkgrtpkt pdwypndrag ihgppliieg seywfavadlh lnyeeavlyc  
 841 asnhsflati tsfvglkaik nkianisgdg qkwwirise piddhftysr ypwhrfpvtf  
 901 geeclymsak twlidlgkpt ddstklpfic ekynvsslek yspdsaakvq cseqwipfqn  
 20 961 kcflkikpvs ltfqsasdtc hsyggtlpsv lsqieqdfit sllpdmeatl wiglrwtaye  
 1021 kinkwtdnre ltysnfhpll vsgrlripen ffeeesryhc alilnlqksp ftgtwnftsc  
 1081 serhfvsllcq kysevksrq lqnasetvky lnnlykiipk tltwhsakre clksnmqlvs  
 25 1141 itdpypyqqafl svqallhnss lwiglfsqdd elnfgwsdgk rlhfsrwaet ngqledcvvl  
 1201 dtdgfwktd cndnqpgaic yysgneteke vkpvdsvkcp spvlntpwip fqncycnff  
 1261 tkrnrhmattq devhtkcqkl npkshilsir dekennfle qlyfnymas wvmlgityrn  
 1321 nslmwfdktp lsythwragr ptiknekfla glstdgfwdi qtfkvieeav yfhqhsilac  
 1381 kiemvdykee hntlpqfmp yedgiyviq kkvtwyealn mcsqsgghla svhnqngqlf  
 1441 ledivkrdgf plwvglssh gsesfewsd gstdyipwk gqtspgncvl ldpkgtwkhe  
 1501 kcnsvkdgai cykptkskkl srlyssrcp aakengsrwi qykgchcyksd qalhsfseak  
 1561 klcskhdsat tivsikdeder nkfvslmre nnnitmrwl gisqhsvdqs wswldgsevt  
 40 1621 fvkwenksks gvgrcsmlia snetworkvec ehgfgrvvck vplgpdytai aiivatlsil  
 1681 vlmgqliwfl fqrhrlhlag fssvryaqgv nedeimlpsf hd

[0027] The major domains of human DEC-205 can be represented as follows:

45

**N-CR-FNI1-CTLD1-CTLD2-CTLD3-CTLD4-CTLD5-CTLD6-CTLD7-CTLD8-CTLD9-CTLD10-TMC**

Where N is the N-terminus, CR represents the "Cys Rich" domain, FNII represents the "Fibronectin Type II" domain, CTLD1 to CTLD10 represent the ten "C-Type Lectin-Like" domains and TMC represents the transmembrane and cytoplasmic domains.

[0028] The term "dendritic cell" as used herein, includes immature and mature dendritic cells and related myeloid progenitor cells that are capable of differentiating into dendritic cells, or related antigen presenting cells (e.g., monocytes and macrophages) in that they express antigens in common with dendritic cells. As used herein, the term "related" includes a cell that is derived from a common progenitor cell or cell lineage. In one embodiment, binding of an antibody of the invention to dendritic cells mediates an effect on dendritic cell growth and/or function by targeting molecules or cells with defined functions (e.g., tumor cells, effector cells, microbial pathogens) to dendritic cells. In a further embodiment, binding of an antibody of the invention to a dendritic cell results in internalization of the antibody by the dendritic cell.

[0029] "MHC molecules" include two types of molecules, MHC class I and MHC class II. MHC class I molecules present

antigen to specific CD8<sup>+</sup> T cells and MHC class II molecules present antigen to specific CD4<sup>+</sup> T cells. Antigens delivered exogenously to APCs are processed primarily for association with MHC class II. In contrast, antigens delivered endogenously to APCs are processed primarily for association with MHC class I. However, under specific conditions, DCs have the unique capacity to allow exogenous antigens access to internal compartments for binding to MHC class I molecules, in addition to MHC class II molecules. This process is called "cross-priming" or "cross-presentation."

**[0030]** As used herein, the term "immunostimulatory agent" refers to compounds capable of stimulating APCs, such as DCs and macrophages. For example, suitable immunostimulatory agents for use in the present invention are capable of stimulating APCs so that the maturation process of the APCs is accelerated, the proliferation of APCs is increased, and/or the recruitment or release of co-stimulatory molecules (e.g., CD80, CD86, ICAM-1, MHC molecules and CCR7) and pro-inflammatory cytokines (e.g., IL-1 $\beta$ , IL-6, IL-12, IL-15, and IFN- $\gamma$ ) is upregulated. Suitable immunostimulatory agents are also capable of increasing T cell proliferation. Such immunostimulatory agents include, but are not be limited to, CD40 ligand; cytokines, such as IFN- $\alpha$ , IFN- $\beta$ , IFN- $\gamma$  and IL-2; colony-stimulating factors, such as G-CSF (granulocyte colony-stimulating factor) and GM-CSF (granulocyte-macrophage colony-stimulating factor); an anti-CTLA-4 antibody; LPS (endotoxin); ssRNA; dsRNA; Bacille Calmette-Guerin (BCG); Levamisole hydrochloride; and intravenous immune globulins. In one embodiment an immunostimulatory agent may be a Toll-like Receptor (TLR) agonist. For example the immunostimulatory agent may be a TLR3 agonist such as double-stranded inosine:cytosine polynucleotide (Poly I:C, for example available as Ampligen<sup>TM</sup> from Hemispherx Biopharma, PA, US) or Poly A:U; a TLR4 agonist such as monophosphoryl lipid A (MPL) or RC-529 (for example as available from GSK, UK); a TLR5 agonist such as flagellin; a TLR7 or TLR8 agonist such as an imidazoquinoline TLR7 or TLR 8 agonist, for example imiquimod (eg Aldara<sup>TM</sup>) or resiquimod and related imidazoquinoline agents (for example as available from 3M Corporation); or a TLR 9 agonist such as a deoxynucleotide with unmethylated CpG motifs (so-called "CpGs", for example as available from Coley Pharmaceutical). Such immunostimulatory agents may be administered simultaneously, separately or sequentially with the antibodies and constructs of the present invention and may also be physically linked to the antibodies and constructs.

**[0031]** As used herein, the term "linked" refers to the association of two or more molecules. The linkage can be covalent or non-covalent. The linkage also can be genetic (i.e., recombinantly fused). Such linkages can be achieved using a wide variety of art recognized techniques, such as chemical conjugation and recombinant protein production.

**[0032]** As used herein, the term antigen "cross-presentation" refers to presentation of exogenous protein antigens to T cells via MHC class I and class II molecules on APCs.

**[0033]** As used herein, the term "T cell-mediated response" refers to any response mediated by T cells, including effector T cells (e.g., CD8<sup>+</sup> cells) and helper T cells (e.g., CD4<sup>+</sup> cells). T cell mediated responses include, for example, T cell cytotoxicity and proliferation.

**[0034]** As used herein, the term "cytotoxic T lymphocyte (CTL) response" refers to an immune response induced by cytotoxic T cells. CTL responses are mediated primarily by CD8<sup>+</sup> T cells.

**[0035]** The term "antibody" as referred to herein includes whole antibodies and any antigen binding fragment (i.e., "antigen-binding portion") or single chain thereof. An "antibody" refers, in one preferred embodiment, to a glycoprotein comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds, or an antigen binding portion thereof. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as V<sub>H</sub>) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as V<sub>L</sub>) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The V<sub>H</sub> and V<sub>L</sub> regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each V<sub>H</sub> and V<sub>L</sub> is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The constant regions of the antibodies may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (e.g., effector cells) and the first component (C1q) of the classical complement system.

**[0036]** The term "antigen-binding portion" of an antibody (or simply "antibody portion"), as used herein, refers to one or more fragments of an antibody that retain the ability to specifically bind to an antigen (e.g., human DEC-205). It has been shown that the antigen-binding function of an antibody can be performed by fragments of a full-length antibody. Examples of binding fragments encompassed within the term "antigen-binding portion" of an antibody include (i) a Fab fragment, a monovalent fragment consisting of the V<sub>L</sub>, V<sub>H</sub>, CL and CH1 domains; (ii) a F(ab')<sub>2</sub> fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the V<sub>H</sub> and CH1 domains; (iv) a Fv fragment consisting of the V<sub>L</sub> and V<sub>H</sub> domains of a single arm of an antibody, (v) a dAb fragment (Ward et al., (1989) *Nature* 341:544-546), which consists of a V<sub>H</sub> domain; and (vi) an isolated complementarity determining region (CDR) or (vii) a combination of two or more isolated CDRs which may optionally be joined by a synthetic linker. Furthermore, although the two domains of the Fv fragment, V<sub>L</sub> and V<sub>H</sub>, are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the V<sub>L</sub> and V<sub>H</sub> regions pair to form monovalent molecules (known as single chain Fv (scFv); see

e.g., Bird et al. (1988) *Science* 242:423-426; and Huston et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883). Such single chain antibodies are also intended to be encompassed within the term "antigen-binding portion" of an antibody. These antibody fragments are obtained using conventional techniques known to those with skill in the art, and the fragments are screened for utility in the same manner as are intact antibodies. Antigen-binding portions can be produced by recombinant DNA techniques, or by enzymatic or chemical cleavage of intact immunoglobulins.

**[0037]** A "bispecific" or "bifunctional antibody" is an artificial hybrid antibody having two different heavy/light chain pairs and two different binding sites. Bispecific antibodies can be produced by a variety of methods including fusion of hybridomas or linking of Fab' fragments. See, e.g., Songsivilai & Lachmann, *Clin. Exp. Immunol.* 79:315-321 (1990); Kostelny et al., *J. Immunol.* 148, 1547-1553 (1992).

**[0038]** The term "monoclonal antibody," as used herein, refers to an antibody which displays a single binding specificity and affinity for a particular epitope. Accordingly, the term "human monoclonal antibody" refers to an antibody which displays a single binding specificity and which has variable and optional constant regions derived from human germline immunoglobulin sequences. In one embodiment, human monoclonal antibodies are produced by a hybridoma which includes a B cell obtained from a transgenic non-human animal, e.g., a transgenic mouse, having a genome comprising a human heavy chain transgene and a light chain transgene fused to an immortalized cell.

**[0039]** The term "recombinant human antibody," as used herein, includes all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as (a) antibodies isolated from an animal (e.g., a mouse) that is transgenic or transchromosomal for human immunoglobulin genes or a hybridoma prepared therefrom, (b) antibodies isolated from a host cell transformed to express the antibody, e.g., from a transfectoma, (c) antibodies isolated from a recombinant, combinatorial human antibody library, and (d) antibodies prepared, expressed, created or isolated by any other means that involve splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies comprise variable and constant regions that utilize particular human germline immunoglobulin sequences are encoded by the germline genes, but include subsequent rearrangements and mutations which occur, for example, during antibody maturation. As known in the art (see, e.g., Lonberg (2005) *Nature Biotech.* 23(9): 1117-1125), the variable region contains the antigen binding domain, which is encoded by various genes that rearrange to form an antibody specific for a foreign antigen. In addition to rearrangement, the variable region can be further modified by multiple single amino acid changes (referred to as somatic mutation or hypermutation) to increase the affinity of the antibody to the foreign antigen. The constant region will change in further response to an antigen (i.e., isotype switch). Therefore, the rearranged and somatically mutated nucleic acid molecules that encode the light chain and heavy chain immunoglobulin polypeptides in response to an antigen may not have sequence identity with the original nucleic acid molecules, but instead will be substantially identical or similar (i.e., have at least 80% identity).

**[0040]** The term "human antibody" includes antibodies having variable and constant regions (if present) of human germline immunoglobulin sequences. Human antibodies of the invention can include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*) (see, Lonberg, N. et al. (1994) *Nature* 368(6474): 856-859); Lonberg, N. (1994) *Handbook of Experimental Pharmacology* 113:49-101; Lonberg, N. and Huszar, D. (1995) *Intern. Rev. Immunol.* Vol. 13: 65-93, and Harding, F. and Lonberg, N. (1995) *Ann. N.Y. Acad. Sci* 764:536-546). However, the term "human antibody" does not include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences (i.e., humanized antibodies).

**[0041]** As used herein, a "heterologous antibody" is defined in relation to the transgenic non-human organism producing such an antibody. This term refers to an antibody having an amino acid sequence or an encoding nucleic acid sequence corresponding to that found in an organism not consisting of the transgenic non-human animal, and generally from a species other than that of the transgenic non-human animal.

**[0042]** An "isolated antibody," as used herein, is intended to refer to an antibody which is substantially free of other antibodies having different antigenic specificities (e.g., an isolated antibody that specifically binds to human DEC-205 is substantially free of antibodies that specifically bind antigens other than human DEC-205). An isolated antibody that specifically binds to an epitope of may, however, have cross-reactivity to other DEC-205 proteins from different species. However, the antibody preferably always binds to human DEC-205. In addition, an isolated antibody is typically substantially free of other cellular material and/or chemicals. In one embodiment of the invention, a combination of "isolated" antibodies having different DEC-205 specificities is combined in a well defined composition.

**[0043]** The term "epitope" or "antigenic determinant" refers to a site on an antigen to which an immunoglobulin or antibody specifically binds. Epitopes can be formed both from contiguous amino acids or noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids are typically retained on exposure to denaturing solvents, whereas epitopes formed by tertiary folding are typically lost on treatment with denaturing solvents. An epitope typically includes at least 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acids in a unique spatial conformation. Methods of determining spatial conformation of epitopes include techniques in the art and those described herein, for example, x-ray crystallography and 2-dimensional nuclear magnetic resonance (see, e.g., *Epitope Mapping Protocols in Methods in Molecular Biology*, Vol. 66, G. E. Morris, Ed. (1996)). In the present case an epitope is preferably

located in the extracellular domain of human DEC-205, for example in one or a combination of the cysteine rich domain, the FnII domain or one or more of the ten C-type lectin-like domains of human DEC-205.

[0044] As used herein, the terms "specific binding," "selective binding," "selectively binds," and "specifically binds," refer to antibody binding to an epitope on a predetermined antigen. Typically, the antibody binds with an equilibrium dissociation constant ( $K_D$ ) of approximately less than  $10^{-7}$  M, such as approximately less than  $10^{-8}$  M,  $10^{-9}$  M or  $10^{-10}$  M or even lower when determined by surface plasmon resonance (SPR) technology in a BIACORE 2000 instrument using recombinant human DEC-205 as the analyte and the antibody as the ligand and binds to the predetermined antigen with an affinity that is at least two-fold greater than its affinity for binding to a non-specific antigen (e.g., BSA, casein) other than the predetermined antigen or a closely-related antigen. The phrases "an antibody recognizing an antigen" and "an antibody specific for an antigen" are used interchangeably herein with the term "an antibody which binds specifically to an antigen."

[0045] Also encompassed by the present invention are antibodies that bind the same epitope and/or antibodies that compete for binding to human DEC-205 with the antibodies described herein. Antibodies that recognize the same epitope or compete for binding can be identified using routine techniques. Such techniques include, for example, an immunoassay, which shows the ability of one antibody to block the binding of another antibody to a target antigen, *i.e.*, a competitive binding assay. Competitive binding is determined in an assay in which the immunoglobulin under test inhibits specific binding of a reference antibody to a common antigen, such as DEC-205. Numerous types of competitive binding assays are known, for example: solid phase direct or indirect radioimmunoassay (RIA), solid phase direct or indirect enzyme immunoassay (EIA), sandwich competition assay (see Stahli et al., *Methods in Enzymology* 9:242 (1983)); solid phase direct biotin-avidin EIA (see Kirkland et al., *J. Immunol.* 137:3614 (1986)); solid phase direct labeled assay, solid phase direct labeled sandwich assay (see Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press (1988)); solid phase direct label RIA using 1-125 label (see Morel et al., *Mol. Immunol.* 25(1):7 (1988)); solid phase direct biotin-avidin EIA (Cheung et al., *Virology* 176:546 (1990)); and direct labeled RIA. (Moldenhauer et al., *Scand. J. Immunol.* 32:77 (1990)). Typically, such an assay involves the use of purified antigen bound to a solid surface or cells bearing either of these, an unlabeled test immunoglobulin and a labeled reference immunoglobulin. Competitive inhibition is measured by determining the amount of label bound to the solid surface or cells in the presence of the test immunoglobulin. Usually the test immunoglobulin is present in excess. Usually, when a competing antibody is present in excess, it will inhibit specific binding of a reference antibody to a common antigen by at least 50-55%, 55-60%, 60-65%, 65-70% 70-75% or more.

[0046] Other techniques include, for example, epitope mapping methods, such as, x-ray analyses of crystals of antigen; antibody complexes which provides atomic resolution of the epitope. Other methods monitor the binding of the antibody to antigen fragments or mutated variations of the antigen where loss of binding due to a modification of an amino acid residue within the antigen sequence is often considered an indication of an epitope component. In addition, computational combinatorial methods for epitope mapping can also be used. These methods rely on the ability of the antibody of interest to affinity isolate specific short peptides from combinatorial phage display peptide libraries. The peptides are then regarded as leads for the definition of the epitope corresponding to the antibody used to screen the peptide library. For epitope mapping, computational algorithms have also been developed which have been shown to map conformational discontinuous epitopes.

[0047] The term " $K_D$ ," as used herein, is intended to refer to the dissociation equilibrium constant of a particular antibody-antigen interaction. Typically, the human antibodies of the invention bind to DEC-205 with a dissociation equilibrium constant ( $K_D$ ) of approximately  $10^{-8}$  M or less, such as less than  $10^{-9}$  M or  $10^{-10}$  M or even lower when determined by surface plasmon resonance (SPR) technology in a BIACORE 2000 instrument using recombinant human DEC-205 as the analyte and the antibody as the ligand.

[0048] The term "kd" as used herein, is intended to refer to the off rate constant for the dissociation of an antibody from the antibody/antigen complex.

[0049] The term "ka" as used herein, is intended to refer to the on rate constant for the association of an antibody with the antigen.

[0050] The term "EC50," as used herein, refers to the concentration of an antibody or an antigen-binding portion thereof, which induces a response, either in an *in vitro* or an *in vivo* assay, which is 50% of the maximal response, *i.e.*, halfway between the maximal response and the baseline.

[0051] As used herein, "isotype" refers to the antibody class (e.g., IgM or IgG1) that is encoded by heavy chain constant region genes. In one embodiment, a human monoclonal antibody of the invention is of the IgG1 isotype. In another embodiment, a human monoclonal antibody of the invention is of the IgG2 isotype.

[0052] The term "binds to immobilized DEC-205," refers to the ability of a human antibody of the invention to bind to DEC-205, for example, expressed on the surface of a cell or which is attached to a solid support.

[0053] The term "cross-reacts," as used herein, refers to the ability of an antibody of the invention to bind to DEC-205 from a different species. For example, an antibody of the present invention which binds human DEC-205 may also bind cynomolgous DEC-205. As used herein, cross-reactivity is measured by detecting a specific reactivity with purified

antigen in binding assays (e.g., SPR, ELISA) or binding to, or otherwise functionally interacting with, cells physiologically expressing DEC-205. Methods for determining cross-reactivity include standard binding assays as described herein, for example, by Biacore™ surface plasmon resonance (SPR) analysis using a Biacore™ 2000 SPR instrument (Biacore AB, Uppsala, Sweden), or binding to DEC-205 expressing cells from the species concerned (e.g., dendritic cells) by, for example, flow cytometric techniques.

5 [0054] As used herein, "isotype switching" refers to the phenomenon by which the class, or isotype, of an antibody changes from one Ig class to one of the other Ig classes.

10 [0055] As used herein, "nonswitched isotype" refers to the isotypic class of heavy chain that is produced when no isotype switching has taken place; the CH gene encoding the nonswitched isotype is typically the first CH gene immediately downstream from the functionally rearranged VDJ gene. Isotype switching has been classified as classical or non-classical isotype switching. Classical isotype switching occurs by recombination events which involve at least one switch sequence region in the transgene. Non-classical isotype switching may occur by, for example, homologous recombination between human  $\sigma_{\mu}$  and human  $\Sigma_{\mu}$ . ( $\delta$ -associated deletion). Alternative non-classical switching mechanisms, such as intertransgene and/or interchromosomal recombination, among others, may occur and effectuate isotype switching.

15 [0056] As used herein, the term "switch sequence" refers to those DNA sequences responsible for switch recombination. A "switch donor" sequence, typically a  $\mu$  switch region, will be 5' (i.e., upstream) of the construct region to be deleted during the switch recombination. The "switch acceptor" region will be between the construct region to be deleted and the replacement constant region (e.g.,  $\gamma$ ,  $\epsilon$ , etc.). As there is no specific site where recombination always occurs, the final gene sequence will typically not be predictable from the construct.

20 [0057] As used herein, "glycosylation pattern" is defined as the pattern of carbohydrate units that are covalently attached to a protein, more specifically to an immunoglobulin protein. A glycosylation pattern of a heterologous antibody can be characterized as being substantially similar to glycosylation patterns which occur naturally on antibodies produced by the species of the nonhuman transgenic animal, when one of ordinary skill in the art would recognize the glycosylation pattern of the heterologous antibody as being more similar to said pattern of glycosylation in the species of the nonhuman 25 transgenic animal than to the species from which the CH genes of the transgene were derived.

[0058] The term "naturally-occurring" as used herein as applied to an object refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be isolated from a source in nature and which has not been intentionally modified by man in the laboratory is naturally-occurring.

30 [0059] The term "rearranged" as used herein refers to a configuration of a heavy chain or light chain immunoglobulin locus wherein a V segment is positioned immediately adjacent to a D-J or J segment in a conformation encoding essentially a complete  $V_H$  or  $V_L$  domain, respectively. A rearranged immunoglobulin gene locus can be identified by comparison to germline DNA; a rearranged locus will have at least one recombined heptamer/nonamer homology element.

35 [0060] The term "unrearranged" or "germline configuration" as used herein in reference to a V segment refers to the configuration wherein the V segment is not recombined so as to be immediately adjacent to a D or J segment.

[0061] The term "nucleic acid molecule," as used herein, is intended to include DNA molecules and RNA molecules. A nucleic acid molecule may be single-stranded or double-stranded, but preferably is double-stranded DNA.

40 [0062] The term "isolated nucleic acid molecule," as used herein in reference to nucleic acids encoding antibodies or antibody portions (e.g.,  $V_H$ ,  $V_L$ , CDR3) that bind to DEC-205, is intended to refer to a nucleic acid molecule in which the nucleotide sequences encoding the antibody or antibody portion are free of other nucleotide sequences encoding antibodies or antibody portions that bind antigens other than DEC-205, which other sequences may naturally flank the nucleic acid in human genomic DNA. For example, SEQ ID NOs: 2, 3 (with signal peptide) / 4 (without signal peptide), and SEQ ID NOs: 8, 9 (with signal peptide) / 10 (without signal peptide) correspond, respectively, to the nucleotide and 45 amino acid sequences comprising the heavy chain ( $V_H$ ) and light chain ( $V_L$ ) variable regions of the human anti-DEC-205 antibody 3D6-2F4 of the invention. In particular, SEQ ID NO: 2 and 3/4 correspond to the nucleotide and amino acid sequence, respectively, of  $V_H$  of the 3D6-2F4 antibody, SEQ ID NO: 8 and 9/10 correspond to the nucleotide and amino acid sequence, respectively, of  $V_L$  of the 3D6-2F4 antibody.

50 [0063] "Conservative sequence modifications" of the sequences set forth in SEQ ID NOs: 2-91 are nucleotide and amino acid sequence modifications which do not abrogate the binding of the antibody encoded by the nucleotide sequence or containing the amino acid sequence, to the antigen. Such conservative sequence modifications include conservative nucleotide and amino acid substitutions, as well as, nucleotide and amino acid additions and deletions. For example, modifications can be introduced into SEQ ID NOs: 2-91 by standard techniques known in the art, such as site-directed mutagenesis and PCR-mediated mutagenesis. Conservative amino acid substitutions include ones in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine, tryptophan), nonpolar side chains (e.g., alanine, valine,

leucine, isoleucine, proline, phenylalanine, methionine), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in a human anti-DEC-205 antibody is preferably replaced with another amino acid residue from the same side chain family. Methods of identifying nucleotide and amino acid conservative substitutions which do not eliminate antigen binding are well-known in the art (see, e.g., Brummell et al., *Biochem.* 32:1180-1187 (1993); Kobayashi et al. *Protein Eng.* 12(10):879-884 (1999); and Burks et al. *Proc. Natl. Acad. Sci. USA* 94:412-417 (1997))

5 [0064] Alternatively, in another embodiment, mutations can be introduced randomly along all or part of an anti-DEC-205 antibody coding sequence, such as by saturation mutagenesis, and the resulting modified anti-DEC-205 antibodies can be screened for binding activity.

10 [0065] For nucleic acids, the term "substantial homology" indicates that two nucleic acids, or designated sequences thereof, when optimally aligned and compared, are identical, with appropriate nucleotide insertions or deletions, in at least about 80% of the nucleotides, usually at least about 90% to 95%, and more preferably at least about 98% to 99.5% of the nucleotides. Alternatively, substantial homology exists when the segments will hybridize under selective hybridization conditions, to the complement of the strand.

15 [0066] The percent identity between two sequences is a function of the number of identical positions shared by the sequences (i.e., % homology = # of identical positions/total # of positions x 100), taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences. The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm, as described in the non-limiting examples below.

20 [0067] The percent identity between two nucleotide sequences can be determined using the GAP program in the GCG software package (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. The percent identity between two nucleotide or amino acid sequences can also be determined using the algorithm of E. Meyers and W. Miller (CABIOS, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. In addition, the percent identity between two amino acid sequences can be determined using the Needleman and Wunsch (J. Mol. Biol. (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6.

25 [0068] The nucleic acid and protein sequences of the present invention can further be used as a "query sequence" to perform a search against public databases to, for example, identify related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, et al. (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

30 [0069] The nucleic acids may be present in whole cells, in a cell lysate, or in a partially purified or substantially pure form. A nucleic acid is "isolated" or "rendered substantially pure" when purified away from other cellular components or other contaminants, e.g., other cellular nucleic acids or proteins, by standard techniques, including alkaline/SDS treatment, CsCl banding, column chromatography, agarose gel electrophoresis and others well known in the art. See, F. Ausubel, et al., ed. *Current Protocols in Molecular Biology*, Greene Publishing and Wiley Interscience, New York (1987).

35 [0070] The nucleic acid compositions of the present invention, while often in a native sequence (except for modified restriction sites and the like), from either cDNA, genomic or mixtures thereof may be mutated, in accordance with standard techniques to provide gene sequences. For coding sequences, these mutations, may affect amino acid sequence as desired. In particular, DNA sequences substantially homologous to or derived from native V, D, J, constant, switches and other such sequences described herein are contemplated (where "derived" indicates that a sequence is identical or modified from another sequence).

40 [0071] A nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For instance, a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence. With respect to transcription regulatory sequences, operably linked means that the DNA sequences being linked are contiguous and, where necessary to join two protein coding regions, contiguous and in reading frame. For switch sequences, operably linked indicates that the sequences are capable of effecting switch recombination.

45 [0072] The term "vector," as used herein, is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid," which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host

cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein 5 as "recombinant expression vectors" (or simply, "expression vectors"). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" may be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

10 [0073] The term "recombinant host cell" (or simply "host cell"), as used herein, is intended to refer to a cell into which a recombinant expression vector has been introduced. It should be understood that such terms are intended to refer not only to the particular subject cell but to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term "host cell" as used herein.

15 [0074] The term "antigen presenting cell" or "APC" is a cell that displays foreign antigen complexed with MHC on its surface. T-cells recognize this complex using T-cell receptor (TCR). Examples of APCs include, but are not limited to, dendritic cells (DCs), peripheral blood mononuclear cells (PBMC), monocytes (such as THP-1), B lymphoblastoid cells (such as C1R.A2, 1518 B-LCL) and monocyte-derived dendritic cells (DCs). Some APCs internalize antigens either by phagocytosis or by receptor-mediated endocytosis. Examples of APC receptors include, but are not limited to C-type 20 lectins, such as, the human Dendritic and Epithelial Cell 205 receptor (DEC-205), and the human macrophage mannose receptor.

[0075] The term "antigen presentation" refers to the process by which APCs capture antigens and enables their recognition by T-cells, e.g., as a component of an MHC-I and/or MHC-II conjugate.

25 [0076] The terms "inducing an immune response" and "enhancing an immune response" are used interchangeably and refer to the stimulation of an immune response (i.e., either passive or adaptive) to a particular antigen.

[0077] The terms "treat," "treating," and "treatment," as used herein, refer to therapeutic or preventative measures described herein. The methods of "treatment" employ administration to a subject, in need of such treatment, a human antibody of the present invention, for example, a subject in need of an enhanced immune response against a particular antigen or a subject who ultimately may acquire such a disorder, in order to prevent, cure, delay, reduce the severity of, 30 or ameliorate one or more symptoms of the disorder or recurring disorder, or in order to prolong the survival of a subject beyond that expected in the absence of such treatment.

[0078] The term "effective dose" or "effective dosage" is defined as an amount sufficient to achieve or at least partially achieve the desired effect. The term "therapeutically effective dose" is defined as an amount sufficient to cure or at least partially arrest the disease and its complications in a patient already suffering from the disease. Amounts effective for 35 this use will depend upon the severity of the disorder being treated and the general state of the patient's own immune system.

[0079] The term "patient" includes human and other mammalian subjects that receive either prophylactic or therapeutic treatment.

40 [0080] As used herein, the term "subject" includes any human or non-human animal. For example, the methods and compositions of the present invention can be used to treat a subject with an immune disorder. The term "non-human animal" includes all vertebrates, e.g., mammals and non-mammals, such as non-human primates, sheep, dog, cow, chickens, amphibians, reptiles, etc.

[0081] Various aspects of the invention are described in further detail in the following subsections.

45 I. Production of Antibodies to DEC-205

[0082] The present invention encompasses antibodies, e.g., fully human antibodies, that bind DEC-205, e.g., human DEC-205. Exemplary monoclonal antibodies that bind DEC-205 include 3D6-2F4, 3D6-4C8, 3G9-2D2, 5A8-1F1, 2D3-1F5-2A9, 3C7-3A3, 5D12-5G1, 1G6-1G6, 5C3-2-3F6, 1E6-3D10 and 3A4-1C10.

50 [0083] Monoclonal antibodies of the invention can be produced using a variety of known techniques, such as the standard somatic cell hybridization technique described by Kohler and Milstein, *Nature* 256: 495 (1975). Although somatic cell hybridization procedures are preferred, in principle, other techniques for producing monoclonal antibodies also can be employed, e.g., viral or oncogenic transformation of B lymphocytes, phage display technique using libraries of human antibody genes.

55 [0084] Accordingly, in one embodiment, a hybridoma method is used for producing an antibody that binds human DEC-205. In this method, a mouse or other appropriate host animal can be immunized with a suitable antigen in order to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the antigen used for immunization. Alternatively, lymphocytes may be immunized *in vitro*. Lymphocytes can then be fused with myeloma

cells using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)). Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against the antigen. After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, *Monoclonal Antibodies: Principles and Practice*, pp. 59-103 (Academic Press, 1986)). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may be grown *in vivo* as ascites tumors in an animal. The monoclonal antibodies secreted by the subclones can be separated from the culture medium, ascites fluid, or serum by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

5 [0085] In another embodiment, antibodies and antibody portions that bind human DEC-205 can be isolated from antibody phage libraries generated using the techniques described in, for example, McCafferty et al., *Nature*, 348:552-554 (1990). Clackson et al., *Nature*, 352:624-628 (1991), Marks et al., *J. Mol. Biol.*, 222:581-597 (1991) and Hoet et al (2005) *Nature Biotechnology* 23, 344-348 ; U.S. Patent Nos. 5,223,409; 5,403,484; and 5,571,698 to Ladner et al.; U.S. Patent 10 Nos. 5,427,908 and 5,580,717 to Dower et al.; U.S. Patent Nos. 5,969,108 and 6,172,197 to McCafferty et al.; and U.S. Patent Nos. 5,885,793; 6,521,404; 6,544,731; 6,555,313; 6,582,915 and 6,593,081 to Griffiths et al. Additionally, production of high affinity (nM range) human antibodies by chain shuffling (Marks et al., *Bio/Technology*, 10:779-783 (1992)), as well as combinatorial infection and *in vivo* recombination as a strategy for constructing very large phage libraries (Waterhouse et al., *Nuc. Acids. Res.*, 21:2265-2266 (1993)) may also be used.

20 [0086] In a particular embodiment, the antibody that binds human DEC-205 is produced using the phage display technique described by Hoet et al., *supra*. This technique involves the generation of a human Fab library having a unique combination of immunoglobulin sequences isolated from human donors and having synthetic diversity in the heavy-chain CDRs is generated. The library is then screened for Fabs that bind to human DEC-205.

25 [0087] The preferred animal system for generating hybridomas which produce antibodies of the invention is the murine system. Hybridoma production in the mouse is well known in the art, including immunization protocols and techniques for isolating and fusing immunized splenocytes.

30 [0088] In one embodiment, antibodies directed against DEC-205 are generated using transgenic or transchromosomal mice currying parts of the human immune system rather than the mouse system. In one embodiment, the invention employs transgenic mice, referred to herein as "HuMAb mice" which contain a human immunoglobulin gene miniloci 35 that encodes unarranged human heavy ( $\mu$  and  $\gamma$ ) and  $\kappa$  light chain immunoglobulin sequences, together with targeted mutations that inactivate the endogenous  $\mu$  and  $\kappa$  chain loci (Lonberg, N. et al. (1994) *Nature* 368(6474): 856-859). Accordingly, the mice exhibit reduced expression of mouse IgM or  $\kappa$ , and in response to immunization, the introduced human heavy and light chain transgenes undergo class switching and somatic mutation to generate high affinity human IgG $\kappa$  monoclonal antibodies (Lonberg, N. et al. (1994), *supra*; reviewed in Lonberg, N. (1994) *Handbook of Experimental Pharmacology* 113:49-101; Lonberg, N. and Huszar, D. (1995) *Intern. Rev. Immunol.* Vol. 13: 65-93, and Harding, F. and Lonberg, N. (1995) *Ann. N.Y. Acad. Sci* 764:536-546). The preparation of HuMAb mice is described in detail in Section II below and in Taylor, L. et al. (1992) *Nucleic Acids Research* 20:6287-6295; Chen, J. et al. (1993) *International Immunology* 5: 647-656; Tuailon et al. (1993) *Proc. Natl. Acad. Sci USA* 90:3720-3724; Choi et al. (1993) *Nature Genetics* 4:117-123; Chen, J. et al. (1993) *EMBO J.* 12: 821-830; Tuailon et al. (1994) *J. Immunol.* 152:2912-2920; Lonberg et al., (1994) *Nature* 368(6474): 856-859; Lonberg, N. (1994) *Handbook of Experimental Pharmacology* 113:49-101; Taylor, L. et al. (1994) *International Immunology* 6: 579-591; Lonberg, N. and Huszar, D. (1995) *Intern. Rev. Immunol.* Vol. 13: 65-93; Harding, F. and Lonberg, N. (1995) *Ann. N.Y. Acad. Sci* 764:536-546; Fishwild, D. et al. (1996) *Nature Biotechnology* 14: 845-851. See further, U.S. Patent Nos. 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,789,650; 5,877,397; 5,661,016; 5,814,318; 5,874,299; and 5,770,429; all to Lonberg and Kay, and GenPharm International; U.S. Patent No. 5,545,807 to Surani et al.; International Publication Nos. WO 98/24884, published on June 11, 1998; WO 94/25585, published November 10, 1994; WO 93/1227, published June 24, 1993; WO 92/22645, published December 23, 1992; WO 92/03918, published March 19, 1992.

#### 50 *Immunizations*

55 [0089] To generate fully human antibodies to DEC-205, transgenic or transchromosomal mice containing human immunoglobulin genes (e.g., HCo12, HCo7 or KM mice) can be immunized with a purified or enriched preparation of the DEC-205 antigen and/or cells expressing DEC-205, as described, for example, by Lonberg et al. (1994) *Nature* 368 (6474): 856-859; Fishwild et al. (1996) *Nature Biotechnology* 14: 845-851 and WO 98/24884. As described herein, HuMAb mice are immunized either with recombinant DEC-205 proteins or cell lines expressing DEC-205 as immunogens. Alternatively, mice can be immunized with DNA encoding human DEC-205. Preferably, the mice will be 6-16 weeks of age upon the first infusion. For example, a purified or enriched preparation (5-50  $\mu$ g) of the recombinant DEC-205 antigen can be used to immunize the HuMAb mice intraperitoneally. In the event that immunizations using a purified or enriched

## EP 2 224 954 B1

preparation of the DEC-205 antigen do not result in antibodies, mice can also be immunized with cells expressing DEC-205, e.g., a cell line, to promote immune responses. Exemplary cell lines include DEC-205-overexpressing stable CHO and Raji cell lines.

[0090] Cumulative experience with various antigens has shown that the HuMAb transgenic mice respond best when initially immunized intraperitoneally (IP) or subcutaneously (SC) with antigen in complete Freund's adjuvant, followed by every other week IP/SC immunizations (up to a total of 10) with antigen in incomplete Freund's adjuvant. The immune response can be monitored over the course of the immunization protocol with plasma samples being obtained by retro-orbital bleeds. The plasma can be screened by ELISA (as described below), and mice with sufficient titers of anti-DEC-205 human immunoglobulin can be used for fusions. Mice can be boosted intravenously with antigen 3 days before sacrifice and removal of the spleen.

*Generation of Hybridomas Producing Monoclonal Antibodies to DEC-205*

[0091] To generate hybridomas producing monoclonal antibodies to DEC-205, splenocytes and lymph node cells from immunized mice can be isolated and fused to an appropriate immortalized cell line, such as a mouse myeloma cell line. The resulting hybridomas can then be screened for the production of antigen-specific antibodies. For example, single cell suspensions of splenic lymphocytes from immunized mice can be fused to SP2/0-Ag8.653 nonsecreting mouse myeloma cells (ATCC, CRL 1580) with 50% PEG (w/v). Cells can be plated at approximately  $1 \times 10^5$  in flat bottom microtiter plate, followed by a two week incubation in selective medium containing besides usual reagents 10% fetal Clone Serum, 5-10% origin hybridoma cloning factor (IGEN) and 1X HAT (Sigma). After approximately two weeks, cells can be cultured in medium in which the HAT is replaced with HT. Individual wells can then be screened by ELISA for human anti-DEC-205 monoclonal IgM and IgG antibodies, or for binding to the surface of cells expressing DEC-205, e.g., a CHO cell line expressing DEC-205, by FLISA (fluorescence-linked immunosorbent assay). Once extensive hybridoma growth occurs, medium can be observed usually after 10-14 days. The antibody secreting hybridomas can be replated, screened again, and if still positive for IgG, anti-DEC-205 monoclonal antibodies can be subcloned at least twice by limiting dilution. The stable subclones can then be cultured *in vitro* to generate antibody in tissue culture medium for characterization.

*Generation of Transfectomas Producing Monoclonal Antibodies to DEC-205*

[0092] Antibodies of the invention also can be produced in a host cell transfectoma using, for example, a combination of recombinant DNA techniques and gene transfection methods as is well known in the art (Morrison, S. (1985) *Science* 229:1202).

[0093] For example, in one embodiment, the gene(s) of interest, e.g., human antibody genes, can be ligated into an expression vector such as a eukaryotic expression plasmid such as used by GS gene expression system disclosed in WO 87/04462, WO 89/01036 and EP 338 841 or other expression systems well known in the art. The purified plasmid with the cloned antibody genes can be introduced in eukaryotic host cells such as CHO-cells or NSO-cells or alternatively other eukaryotic cells like a plant derived cells, fungi or yeast cells. The method used to introduce these genes could be methods described in the art such as electroporation, lipofectine, lipofectamine or other. After introducing these antibody genes in the host cells, cells expressing the antibody can be identified and selected. These cells represent the transfectomas which can then be amplified for their expression level and upscaled to produce antibodies. Recombinant antibodies can be isolated and purified from these culture supernatants and/or cells.

[0094] Alternatively these cloned antibody genes can be expressed in other expression systems such as *E. coli* or in complete organisms or can be synthetically expressed.

*Use of Partial Antibody Sequences to Express Intact Antibodies*

[0095] Antibodies interact with target antigens predominantly through amino acid residues that are located in the six heavy and light chain complementarity determining regions (CDRs). For this reason, the amino acid sequences within CDRs are more diverse between individual antibodies than sequences outside of CDRs. Because CDR sequences are responsible for most antibody-antigen interactions, it is possible to express recombinant antibodies that mimic the properties of specific naturally occurring antibodies by constructing expression vectors that include CDR sequences from the specific naturally occurring antibody grafted onto framework sequences from a different antibody with different properties (see, e.g., Riechmann, L. et al., 1998, *Nature* 332:323-327; Jones, P. et al., 1986, *Nature* 321:522-525; and Queen, C. et al., 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86:10029-10033). Such framework sequences can be obtained from public DNA databases that include germline antibody gene sequences. These germline sequences will differ from mature antibody gene sequences because they will not include completely assembled variable genes, which are formed by V(D)J joining during B cell maturation. Germline gene sequences will also differ from the sequences of a high affinity

secondary repertoire antibody at individual evenly across the variable region. For example, somatic mutations are relatively infrequent in the amino-terminal portion of framework region. For example, somatic mutations are relatively infrequent in the amino terminal portion of framework region 1 and in the carboxy-terminal portion of framework region 4. Furthermore, many somatic mutations do not significantly alter the binding properties of the antibody. For this reason, 5 it is not necessary to obtain the entire DNA sequence of a particular antibody in order to recreate an intact recombinant antibody having binding properties similar to those of the original antibody (see PCT/US99/05535 filed on March 12, 1999). Partial heavy and light chain sequence spanning the CDR regions is typically sufficient for this purpose. The partial sequence is used to determine which germline variable and joining gene segments contributed to the recombined antibody variable genes. The germline sequence is then used to fill in missing portions of the variable regions. Heavy 10 and light chain leader sequences are cleaved during protein maturation and do not contribute to the properties of the final antibody. To add missing sequences, cloned cDNA sequences can be combined with synthetic oligonucleotides by ligation or PCR amplification. Alternatively, the entire variable region can be synthesized as a set of short, overlapping, oligonucleotides and combined by PCR amplification to create an entirely synthetic variable region clone. This process has certain advantages such as elimination or inclusion or particular restriction sites, or optimization of particular codons. 15 [0096] The nucleotide sequences of heavy and light chain transcripts from a hybridoma are used to design an overlapping set of synthetic oligonucleotides to create synthetic V sequences with identical amino acid coding capacities as the natural sequences. The synthetic heavy and kappa chain sequences can differ from the natural sequences in three ways: strings of repeated nucleotide bases are interrupted to facilitate oligonucleotide synthesis and PCR amplification; optimal translation initiation sites are incorporated according to Kozak's rules (Kozak, 1991, J. Biol. Chem. 266: 20 19867-19870); and, HindIII sites are engineered upstream of the translation initiation sites. [0097] For both the heavy and light chain variable regions, the optimized coding, and corresponding non-coding, strand 25 sequences are broken down into 30 - 50 nucleotide approximately the midpoint of the corresponding non-coding oligonucleotide. Thus, for each chain, the oligonucleotides can be assembled into overlapping double stranded sets that span segments of 150 - 400 nucleotides. The pools are then used as templates to produce PCR amplification products of 150 - 400 nucleotides. Typically, a single variable region oligonucleotide set will be broken down into two pools which are 30 separately amplified to generate two overlapping PCR products. These overlapping products are then combined by PCR amplification to form the complete variable region. It may also be desirable to include an overlapping fragment of the heavy or light chain constant region (including the BbsI site of the kappa light chain, or the AgeI site if the gamma heavy chain) in the PCR amplification to generate fragments that can easily be cloned into the expression vector constructs. 35 [0098] The reconstructed heavy and light chain variable regions are then combined with cloned promoter, leader sequence, translation initiation, leader sequence, constant region, 3' untranslated, polyadenylation, and transcription termination, sequences to form expression vector constructs. The heavy and light chain expression constructs can be combined into a single vector, co-transfected, serially transfected, or separately transfected into host cells which are then fused to form a host cell expressing both chains. [0099] Plasmids for use in construction of expression vectors were constructed so that PCR amplified V heavy and V 40 kappa light chain cDNA sequences could be used to reconstruct complete heavy and light chain minigenes. These plasmids can be used to express completely human IgG<sub>1</sub>κ or IgG<sub>4</sub>κ antibodies. Fully human and chimeric antibodies of the present invention also include IgG2, IgG3, IgE, IgA, IgM, and IgD antibodies. Similar plasmids can be constructed for expression of other heavy chain isotypes, or for expression of antibodies comprising lambda light chains. [0100] Thus, in another aspect of the invention, structural features of anti-DEC-205 antibodies of the invention are 45 used to create structurally related anti-DEC-205 antibodies that retain at least one functional property of the antibodies of the invention, such as, for example, binding to human DEC-205 with an affinity constant of at least 10<sup>8</sup> M<sup>-1</sup> as measured by surface plasmon resonance; internalizing after binding to human dendritic cells expressing DEC-205; localizing to antigen processing compartments in human dendritic cells; activating human dendritic cells expressing DEC-205; cross-reacting with DEC-205 on non-human primate dendritic cells or those of other species; and generating or enhancing human T cell, such as CTL, responses to an antigen, preferably CTL responses mediated by both MHC Class I and Class II pathways. [0101] In one embodiment, one or more CDR regions of antibodies of the invention can be combined recombinantly 50 with known framework regions and CDRs to create additional, recombinantly-engineered, anti-DEC-205 antibodies of the invention. The heavy and light chain variable framework regions can be derived from the same or different antibody sequences. The antibody sequences can be the sequences of naturally occurring antibodies or can be consensus sequences of several antibodies. See Kettleborough et al., Protein Engineering 4:773 (1991); Kolbinger et al., Protein Engineering 6:971 (1993) and Carter et al., WO 92/22653. [0102] Accordingly, in another embodiment, the invention provides a method for preparing an anti-DEC-205 antibody 55 including:

preparing an antibody including (1) heavy chain framework regions and heavy chain CDRs, where the heavy chain CDRs comprise the amino acid sequences shown in SEQ ID NOs: 29, 30 and 31; and (2) light chain framework

regions and light chain CDRs, where the light chain CDRs comprise the amino acid sequences shown in SEQ ID NOs: 35, 36 and 37,

5 where the antibody retains the ability to bind to DEC-205. The ability of the antibody to bind DEC-205 can be determined using standard binding assays, such as those set forth in the Examples (e.g., an ELISA or a FLISA).

[0103] 10 It is well known in the art that antibody heavy and light chain CDR3 domains play a particularly important role in the binding specificity/affinity of an antibody for an antigen (see, Hall et al., J. Immunol., 149:1605-1612 (1992); Polymenis et al., J. Immunol., 152:5318-5329 (1994); Jahn et al., Immunobiol., 193:400-419 (1995); Klimka et al., Brit. J. Cancer, 83:252-260 (2000); Beboer et al., J. Mol. Biol., 296:833-849 (2000); Rader et al., Proc. Natl. Acad. Sci. USA, 95:8910-8915 (1998); Barbas et al., J. Am. Chem. Soc., 116:2161-2162 (1994); Ditzel et al., J. Immunol., 157:739-749 (1996)). Accordingly, the recombinant antibodies of the invention prepared as set forth above preferably comprise the heavy and/or light chain CDRs of antibody 3G9-2D2.

15 *Generation of Antibodies Having Modified Sequences*

[0104] 20 In another embodiment, the variable region sequences, or portions thereof, of the anti-DEC-205 antibodies of the invention are modified to create structurally related anti-DEC-205 antibodies that retain binding (i.e., to the same epitope as the unmodified antibody) and, thus, are functionally equivalent. Methods for identifying residues that can be altered without removing antigen binding are well-known in the art (see, e.g., Marks et al. (Biotechnology (1992) 10(7): 779-83 (monoclonal antibodies diversification by shuffling light chain variable regions, then heavy chain variable regions with fixed CDR3 sequence changes), Jespers et al. (1994) Biotechnology 12(9):899-903 (selection of human antibodies from phage display repertoires to a single epitope of an antigen), Sharon et al. (1986) PNAS USA 83(8):2628-31 (site-directed mutagenesis of an invariant amino acid residue at the variable-diversity segments junction of an antibody); Casson et al. (1995) J. Immunol. 155(12):5647-54 (evolution of loss and change of specificity resulting from random mutagenesis of an antibody heavy chain variable region).

[0105] 25 The CDR1, 2, and/or 3 regions of engineered antibodies can comprise the exact amino acid sequence(s) as those of antibodies 3D6-2F4, 3D6-4C8, 3G9-2D2, 5A8-1F1, 2D3-1F5-2A9, 3C7-3A3, 5D12-5G1, 1G6-1G6 and 3A4-1C10 disclosed herein. Antibodies comprising derivatives from the exact CDR sequences of 3D6-2F4, 3D6-4C8, 3G9-2D2, 5A8-1F1, 2D3-1F5-2A9, 3C7-3A3, 5D12-5G1, 1G6-1G6 and 3A4-1C10 may still retain the ability of to bind DEC-205 effectively. Such sequence modifications may include one or more amino acid additions, deletions, or substitutions, e.g., conservative sequence modifications as described above. Sequence modifications may also be based on the consensus sequences described above for the particular CDR1, CDR2, and CDR3 sequences of antibodies 3D6-2F4, 3D6-4C8, 3G9-2D2, 5A8-1F1, 2D3-1F5-2A9, 3C7-3A3, 5D12-5G1, 1G6-1G6 and 3A4-1C10.

[0106] 30 Engineered antibodies may be composed of one or more CDRs that are, for example, 90%, 95%, 98% or 99.5% identical to one or more CDRs of antibodies 3D6-2F4, 3D6-4C8, 3G9-2D2, 5A8-1F1, 2D3-1F5-2A9, 3C7-3A3, 5D12-5G1, 1G6-1G6 and 3A4-1C10.

[0107] 35 One or more residues of a CDR may be altered to modify binding to achieve a more favored on-rate of binding, a more favored off-rate of binding, or both, such that an idealized binding constant is achieved. Using this strategy, an antibody having ultra high binding affinity of, for example,  $10^{10}$  M<sup>-1</sup> or more, can be achieved. Affinity maturation techniques, well known in the art and those described herein, can be used to alter the CDR region(s) followed by screening of the resultant binding molecules for the desired change in binding. Accordingly, as CDR(s) are altered, changes in binding affinity as well as immunogenicity can be monitored and scored such that an antibody optimized for the best combined binding and low immunogenicity are achieved.

[0108] 40 In addition to or instead of modifications within the CDRs, modifications can also be made within one or more of the framework regions, FR1, FR2, FR3 and FR4, of the heavy and/or the light chain variable regions of a antibody, so long as these modifications do not eliminate the binding affinity of the antibody. For example, one or more non-germline amino acid residues in the framework regions of the heavy and/or the light chain variable region of a antibody of the invention, is substituted with a germline amino acid residue, i.e., the corresponding amino acid residue in the human germline sequence for the heavy or the light chain variable region, which the antibody has significant sequence identity with. For example, a antibody chain can be aligned to a germline antibody chain which it shares significant sequence identity with, and the amino acid residues which do not match between antibody framework sequence and the germline chain framework can be substituted with corresponding residues from the germline sequence. When an amino acid differs between a antibody variable framework region and an equivalent human germline sequence variable framework region, the antibody framework amino acid should usually be substituted by the equivalent human germline sequence amino acid if it is reasonably expected that the amino acid falls within one of the following categories:

- (1) an amino acid residue which noncovalently binds antigen directly,
- (2) an amino acid residue which is adjacent to a CDR region,

(3) an amino acid residue which otherwise interacts with a CDR region (e.g., is within about 3-6 Å of a CDR region as determined by computer modeling), or  
(4) an amino acid residue which participates in the VL-VH interface.

5 [0109] Residues which "noncovalently bind antigen directly" include amino acids in positions in framework regions which have a good probability of directly interacting with amino acids on the antigen according to established chemical forces, for example, by hydrogen bonding, Van der Waals forces, hydrophobic interactions, and the like. Accordingly, in one embodiment, an amino acid residue in the framework region of an antibody of the invention is substituted with the corresponding germline amino acid residue which noncovalently binds antigen directly.

10 [0110] Residues which are "adjacent to a CDR region" include amino acid residues in positions immediately adjacent to one or more of the CDRs in the primary sequence of the antibody, for example, in positions immediately adjacent to a CDR as defined by Kabat, or a CDR as defined by Chothia (see e.g., Chothia and Lesk *J. Mol. Biol.* 196:901 (1987)). Accordingly, in one embodiment, an amino acid residue within the framework region of an antibody of the invention is substituted with a corresponding germline amino acid residue which is adjacent to a CDR region.

15 [0111] Residues that "otherwise interact with a CDR region" include those that are determined by secondary structural analysis to be in a spatial orientation sufficient to affect a CDR region. Such amino acids will generally have a side chain atom within about 3 angstrom units (Å) of some atom in the CDRs and must contain an atom that could interact with the CDR atoms according to established chemical forces, such as those listed above. Accordingly, in one embodiment, an amino acid residue within the framework region of an antibody of the invention is substituted with the corresponding germline amino acid residue which otherwise interacts with a CDR region.

20 [0112] The amino acids at several positions in the framework are known to be important for determining CDR confirmation (e.g., capable of interacting with the CDRs) in many antibodies (Chothia and Lesk, *supra*, Chothia *et al.*, *supra* and Tramontano *et al.*, *J. Mol. Biol.* 215:175 (1990)). These authors identified conserved framework residues important for CDR conformation by analysis of the structures of several known antibodies. The antibodies analyzed fell into a limited number of structural or "canonical" classes based on the conformation of the CDRs. Conserved framework residues within members of a canonical class are referred to as "canonical" residues. Canonical residues include residues 2, 25, 29, 30, 33, 48, 64, 71, 90, 94 and 95 of the light chain and residues 24, 26, 29, 34, 54, 55, 71 and 94 of the heavy chain. Additional residues (e.g., CDR structure-determining residues) can be identified according to the methodology of Martin and Thornton (1996) *J. Mol. Biol.* 263:800. Notably, the amino acids at positions 2, 48, 64 and 71 of the light chain and 26-30, 71 and 94 of the heavy chain (numbering according to Kabat) are known to be capable of interacting with the CDRs in many antibodies. The amino acids at positions 35 in the light chain and 93 and 103 in the heavy chain are also likely to interact with the CDRs. Additional residues which may effect conformation of the CDRs can be identified according to the methodology of Foote and Winter (1992) *J. Mol. Biol.* 224:487. Such residues are termed "vernier" residues and are those residues in the framework region closely underlying (i.e., forming a "platform" under) the CDRs.

25 [0113] Residues which "participate in the VL-VH interface" or "packing residues" include those residues at the interface between VL and VH as defined, for example, by Novotny and Haber, *Proc. Natl. Acad. Sci. USA*, 82:4592-66 (1985) or Chothia *et al.*, *supra*.

30 [0114] Occasionally, there is some ambiguity about whether a particular amino acid falls within one or more of the above-mentioned categories. In such instances, alternative variant antibodies are produced, one of which has that particular substitution, the other of which does not. Alternative variant antibodies so produced can be tested in any of the assays described herein for the desired activity, and the preferred antibody selected.

35 [0115] Additional candidates for substitution within the framework region are amino acids that are unusual or "rare" for an antibody at that position. These amino acids can be substituted with amino acids from the equivalent position of the human germline sequence or from the equivalent positions of more typical antibodies. For example, substitution may be desirable when the amino acid in a framework region of the antibody is rare for that position and the corresponding amino acid in the germline sequence is common for that position in immunoglobulin sequences; or when the amino acid in the antibody is rare for that position and the corresponding amino acid in the germline sequence is also rare, relative to other sequences. It is contemplated that by replacing an unusual amino acid with an amino acid from the germline sequence that happens to be typical for antibodies, the antibody may be made less immunogenic.

40 [0116] The term "rare", as used herein, indicates an amino acid occurring at that position in less than about 20%, preferably less than about 10%, more preferably less than about 5%, even more preferably less than about 3%, even more preferably less than about 2% and even more preferably less than about 1% of sequences in a representative sample of sequences, and the term "common", as used herein, indicates an amino acid occurring in more than about 25% but usually more than about 50% of sequences in a representative sample. For example, all light and heavy chain variable region sequences are respectively grouped into "subgroups" of sequences that are especially homologous to each other and have the same amino acids at certain critical positions (Kabat *et al.*, *supra*). When deciding whether an amino acid in an antibody sequence is "rare" or "common" among sequences, it will often be preferable to consider only those sequences in the same subgroup as the antibody sequence.

[0117] In general, the framework regions of antibodies are usually substantially identical, and more usually, identical to the framework regions of the human germline sequences from which they were derived. Of course, many of the amino acids in the framework region make little or no direct contribution to the specificity or affinity of an antibody. Thus, many individual conservative substitutions of framework residues can be tolerated without appreciable change of the specificity or affinity of the resulting immunoglobulin. Thus, in one embodiment the variable framework region of the antibody shares at least 85% sequence identity to a human germline variable framework region sequence or consensus of such sequences. In another embodiment, the variable framework region of the antibody shares at least 90%, 95%, 96%, 97%, 98% or 99% sequence identity to a human germline variable framework region sequence or consensus of such sequences.

[0118] In addition to simply binding DEC-205, an antibody may be selected for its retention of other functional properties of antibodies of the invention, such as, for example:

- (a) binding to human DEC-205 with an affinity constant of at least  $10^8 \text{ M}^{-1}$  as measured by surface plasmon resonance;
- (b) internalizing after binding to human dendritic cells expressing DEC-205;
- (c) localizing to antigen processing compartments in the dendritic cells;
- (d) activating human dendritic cells expressing DEC-205;
- (e) cross-reacting with DEC-205 on non-human primate dendritic cells or those of other species;
- (f) generating or enhancing human T-cell responses, preferably T-cell responses mediated by both MHC Class I and Class II pathways;
- (g) generating or enhancing human CD4+, CD8+ or NKT cell responses; and
- (h) induces peripheral CD8<sup>+</sup> T cell tolerance.

*Characterization of Monoclonal Antibodies to DEC-205*

[0119] Monoclonal antibodies of the invention can be characterized for binding to DEC-205 using a variety of known techniques. Generally, the antibodies are initially characterized by ELISA. Briefly, microtiter plates can be coated with purified DEC-205 in PBS, and then blocked with irrelevant proteins such as bovine serum albumin (BSA) diluted in PBS. Dilutions of plasma from DEC-205-immunized mice are added to each well and incubated for 1-2 hours at 37°C. The plates are washed with PBS/Tween 20 and then incubated with a goat-anti-human IgG Fc-specific polyclonal reagent conjugated to alkaline phosphatase for 1 hour at 37°C. After washing, the plates are developed with ABTS substrate, and analyzed at OD of 405. Preferably, mice which develop the highest titers will be used for fusions.

[0120] An ELISA assay as described above can be used to screen for antibodies and, thus, hybridomas that produce antibodies that show positive reactivity with the DEC-205 immunogen. Hybridomas that bind, preferably with high affinity, to DEC-205 can then be subcloned and further characterized. One clone from each hybridoma, which retains the reactivity of the parent cells (by ELISA), can then be chosen for making a cell bank, and for antibody purification.

[0121] To purify anti-DEC-205 antibodies, selected hybridomas can be grown in roller bottles, two-liter spinner-flasks or other culture systems. Supernatants can be filtered and concentrated before affinity chromatography with protein A-sepharose (Pharmacia, Piscataway, NJ) to purify the protein. After buffer exchange to PBS, the concentration can be determined by OD<sub>280</sub> using 1.43 extinction coefficient or preferably by nephelometric analysis. IgG can be checked by gel electrophoresis and by antigen specific method.

[0122] To determine if the selected anti-DEC-205 monoclonal antibodies bind to unique epitopes, each antibody can be biotinylated using commercially available reagents (Pierce, Rockford, IL). Biotinylated MAb binding can be detected with a streptavidin labeled probe. To determine the isotype of purified antibodies, isotype ELISAs can be performed using art recognized techniques. For example, wells of microtiter plates can be coated with 10 µg/ml of anti- Ig overnight at 4°C. After blocking with 5% BSA, the plates are reacted with 10 µg/ml of monoclonal antibodies or purified isotype controls, at ambient temperature for two hours. The wells can then be reacted with either IgG1 or other isotype specific conjugated probes. Plates are developed and analyzed as described above.

[0123] To test the binding of monoclonal antibodies to live cells expressing DEC-205, flow cytometry can be used. Briefly, cell lines and/or human PBMCs expressing membrane-bound DEC-205 (grown under standard growth conditions) are mixed with various concentrations of monoclonal antibodies in PBS containing 0.1% BSA and 0.01% NaN3 at 4°C for 1 hour. After washing, the cells are reacted with Fluorescein-labeled anti- IgG antibody under the same conditions as the primary antibody staining. The samples can be analyzed by FACScan instrument using light and side scatter properties to gate on single cells and binding of the labeled antibodies is determined. An alternative assay using fluorescence microscopy may be used (in addition to or instead of) the flow cytometry assay. Cells can be stained exactly as described above and examined by fluorescence microscopy. This method allows visualization of individual cells, but may have diminished sensitivity depending on the density of the antigen.

[0124] Anti-DEC-205 IgGs can be further tested for reactivity with the DEC-205 antigen by Western blotting. Briefly, cell extracts from cells expressing DEC-205 can be prepared and subjected to sodium dodecyl sulfate polyacrylamide

gelectrophoresis. After electrophoresis, the separated antigens will be transferred to nitrocellulose membranes, blocked with 20% mouse serum, and probed with the monoclonal antibodies to be tested. IgG binding can be detected using anti- IgG alkaline phosphatase and developed with BCIP/NBT substrate tablets (Sigma Chem. Co., St. Louis, MO).

5 [0125] Methods for analyzing binding affinity, cross-reactivity, and binding kinetics of various anti-DEC-205 antibodies include standard assays known in the art, for example, Biacore™ surface plasmon resonance (SPR) analysis using a Biacore™ 2000 SPR instrument (Biacore AB, Uppsala, Sweden), as described in Example 2 herein.

## II. Molecular Conjugates/Immunotoxins

10 [0126] The present invention provides a variety of therapeutic molecular conjugates (e.g., vaccine conjugates) which include an antigen, such as a tumor or viral antigen, linked to an antibody that binds to a receptor on an APC, for example, an antibody which binds to DEC-205. This allows for targeting of the antigen to APCs, such as cells expressing DEC-205 (e.g., dendritic cells and B cells) to enhance processing, presentation and, ultimately, an immune response against the antigen(s), e.g., a CTL response. A schematic representation of such a conjugate is shown in Figure 8. In the example 15 shown, an antigen is genetically fused to the CH3 domain of each of the heavy chains of a substantially complete anti-DEC-205 antibody. However, it will be appreciated that the antigen may alternatively be joined to other parts of such an antibody or fragment thereof, and that other forms of conjugation, such as chemical conjugation, may also be employed, as discussed further below.

20 [0127] Suitable antigens for use in the present invention include, for example, infectious disease antigens and tumor antigens, against which protective or therapeutic immune responses are desired, e.g., antigens expressed by a tumor cell or a pathogenic organism or infectious disease antigens. For example, suitable antigens include tumor-associated antigens for the prevention or treatment of cancers. Examples of tumor-associated antigens include, but are not limited to, sequences comprising all or part of the sequences of  $\beta$ hCG, gp100 or Pmel17, HER2/neu, WT1, mesothelin, CEA, gp100, MART1, TRP-2, melan-A, NY-ESO-1, NY-BR-1, NY-CO-58, MN (gp250), idiotype, MAGE-1, MAGE-3, MAGE-25 A3, Tyrosinase, Telomerase, SSX2 and MUC-1 antigens, and germ cell derived tumor antigens. Tumor associated antigens also include the blood group antigens, for example, Le<sup>a</sup>, Le<sup>b</sup>, LeX, LeY, H-2, B-1, B-2 antigens. Alternatively, more than one antigen can be included within the antigen-antibody constructs of the invention. For example, a MAGE antigen can be combined with other antigens such as melanin A, tyrosinase, and gp100 along with adjuvants such as GM-CSF or IL-12, and linked to an anti-APC antibody.

30 [0128] Other suitable antigens include viral antigens for the prevention or treatment of viral diseases. Examples of viral antigens include, but are not limited to, HIV-1 gag, HIV-1 env, HIV-1 nef, HBV (surface or core antigens), HPV, FAS, HSV-1, HSV-2, p17, ORF2 and ORF3 antigens. Examples of bacterial antigens include, but are not limited to, *Toxoplasma gondii* or *Treponema pallidum*. The antibody-bacterial antigen conjugates of the invention can be in the treatment or prevention of various bacterial diseases such as Anthrax, Botulism, Tetanus, Chlamydia, Cholera, Diphtheria, 35 Lyme Disease, Syphilis and Tuberculosis.

40 [0129] Sequences of the foregoing antigens are well known in the art. For example, an example of a MAGE-3 cDNA sequence is provided in US 6,235,525 (Ludwig Institute for Cancer Research); examples of NY-ESO-1 nucleic acid and protein sequences are provided in US 5,804,381 and US 6,069,233 (Ludwig Institute for Cancer Research); examples of Melan-A nucleic acid and protein sequences are provided in US 5,620,886 and US 5,854,203 (Ludwig Institute for Cancer Research); examples of NY-BR-1 nucleic acid and protein sequences are provided in US 6,774,226 and US 6,911,529 (Ludwig Institute for Cancer Research) and examples of NY-CO-58 nucleic acid and protein sequences are provided in WO 02090986 (Ludwig Institute for Cancer Research); an example of an amino acid sequence for the HER-2/neu protein is available at GENBANK® Accession No. AAA58637; and a nucleotide sequence (mRNA) for human 45 carcinoembryonic antigen-like 1 (CEA-1) is available at GENBANK® Accession No. NM\_020219.

50 [0130] An HPV antigen that may be used in the pharmaceutical compositions and the methods of the invention may include, for example an HPV-16 antigen, an HPV-18 antigen, an HPV-31 antigen, an HPV-33 antigen and/or an HPV-35 antigen; and is suitably an HPV-16 antigen and/or HPV-18 antigen. A genome of HPV-16 is described in Virology, 145:181- 185 (1985) and DNA sequences encoding HPV-18 are described in US Patent No. 5,840,306. HPV-16 antigens (e.g., seroreactive regions of the E1 and/or E2 proteins of HPV-16) are described in US Patent No. 6,531,127, and HPV-18 antigens (e.g., seroreactive regions of the L1 and/or L2 proteins of HPV-18) are described in US Patent No. 5,840,306. Similarly, a complete genome for HBV is available at GENBANK® Accession No. NC\_003977. The genome of HCV is described in European Patent Application No. 318 216. PCT/US90/01348 discloses sequence information of clones of the HCV genome, amino acid sequences of HCV viral proteins and methods of making and using such compositions for HCV vaccines comprising HCV proteins and peptides derived there from.

55 [0131] Antigenic peptides of proteins (i.e., those containing T cell epitopes) can be identified in a variety of manners well known in the art. For example, T cell epitopes can be predicted by analyzing the sequence of the protein using web-based predictive algorithms (BIMAS & SYFPEITHI) to generate potential MHC class I and II- binding peptides that match an internal database of 10,000 well characterized MHC binding peptides previously defined by CTLs. High scoring

peptides can be ranked and selected as "interesting" on the basis of high affinity to a given MHC molecule.

[0132] Another method for identifying antigenic peptides containing T cell epitopes is by dividing the protein into non-overlapping peptides of desired length or overlapping peptides of desired lengths which can be produced recombinantly, synthetically, or in certain limited situations, by chemical cleavage of the protein and tested for immunogenic properties, 5 e.g., eliciting a T cell response (i.e., proliferation or lymphokine secretion).

[0133] In order to determine precise T cell epitopes of the protein by, for example, fine mapping techniques, a peptide having T cell stimulating activity and thus comprising at least one T cell epitope, as determined by T cell biology techniques, can be modified by addition or deletion of amino acid residues at either the amino or carboxy terminus of the peptide and tested to determine a change in T cell reactivity to the modified peptide. If two or more peptides which share an 10 area of overlap in the native protein sequence are found to have human T cell stimulating activity, as determined by T cell biology techniques, additional peptides can be produced comprising all or a portion of such peptides and these additional peptides can be tested by a similar procedure. Following this technique, peptides are selected and produced recombinantly or synthetically. Peptides are selected based on various factors, including the strength of the T cell 15 response to the peptide (e.g., stimulation index). The physical and chemical properties of these selected peptides (e.g., solubility, stability) can then be examined to determine whether the peptides are suitable for use in therapeutic compositions or whether the peptides require modification.

[0134] In addition, the vaccine conjugate can include one or more immunostimulatory agents that also enhance the immune response against the antigen. Antibody-antigen vaccine conjugates of the invention can be made genetically 20 or chemically. In either case, the antibody portion of the conjugate may consist of the whole antibody or a portion of the antibody, such as the Fab fragment or single-chain Fv. In addition, more than one antigen and/or immunostimulatory agent can be included in the conjugate.

[0135] Chemically constructed antibody-antigen conjugates can be made using a variety of well known and readily 25 available cross-linking reagents. These cross-linking reagents can be homofunctional or heterofunctional compounds, such as N-succinimidyl-3-(2-pyridylidithio)propionate (SPDP), N-succinimidyl-S-acetyl-thioacetate (SATA), sulfosuccinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate (sulfo-SMCC), 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB), that 30 form covalent linkages with different reactive amino acid or carbohydrate side chains on the anti-dendritic antibody and selected antigen. Other coupling and cross-linking agents also can be used to generate covalent linkages, such as protein A, carbodiimide, and o-phenylenedimaleimide (oPDM); (see e.g., Karpovsky et al. (1984) J. Exp. Med. 160:1686; Liu, MA et al. (1985) Proc. Natl. Acad. Sci. USA 82:8648). Other methods include those described by Paulus (Behring 35 Ins. Mitt. (1985) No. 78, 118-132); Brennan et al. (Science (1985) 229:81-83), and Glennie et al. (J. Immunol. (1987) 139: 2367-2375). Preferred conjugating agents are SATA and sulfo-SMCC, both available from Pierce Chemical Co. (Rockford, IL). Immunostimulatory agents can also be chemically linked to the molecular conjugates of the present 40 invention using the same linking methods described above.

[0136] In another embodiment, the antibodies of the present invention are linked to a therapeutic moiety, such as a 45 cytotoxin, a drug or a radioisotope. When conjugated to a cytotoxin, these antibody conjugates are referred to as "immunotoxins." A cytotoxin or cytotoxic agent includes any agent that is detrimental to (e.g., kills) cells. Examples include taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicine, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 50 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine). An antibody of the present invention can be conjugated to a radioisotope, e.g., radioactive iodine, to generate cytotoxic radiopharmaceuticals for treating a dendritic-related disorder, such as an autoimmune or inflammatory disease, or graft versus host disease.

[0137] The antibody conjugates of the invention can be used to modify a given biological response, and the drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be 55 a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, an enzymatically active toxin, or active fragment thereof, such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor or interferon- $\gamma$ ; or, biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors.

[0138] Techniques for conjugating such therapeutic moiety to antibodies are well known, see, e.g., Arnon et al., "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in Monoclonal Antibodies And Cancer Therapy, Reisfeld et al. (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom et al., "Antibodies For Drug Delivery", in Controlled Drug Delivery (2nd Ed.), Robinson et al. (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody

Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in Monoclonal Antibodies '84: Biological And Clinical Applications, Pinchera et al. (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in Monoclonal Antibodies For Cancer Detection And Therapy, Baldwin et al. (eds.), pp. 303-16 (Academic Press 1985), and Thorpe et al., "The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates", Immunol. Rev., 62:119-58 (1982).

[0139] In another embodiment, the antibodies of the present invention can be used to directly target whole cells, e.g., a tumor cell, an effector cell or a microbial pathogen, to dendritic cells. For example, anti-DEC-205 antibodies can be directly expressed on the surface of a cell, for example, by transfection or transduction of a cell with a vector containing nucleic acid sequences encoding an antibody of the invention. This can be done, for example, by transfecting the target cell with a nucleic acid encoding a fusion protein containing a transmembrane domain and a anti-dendritic cell antibody, or antigen binding fragment thereof. Methods for generating such nucleic acids, fusion proteins, and cells expressing such fusion proteins are described, for example, in U.S. Patent Application Serial No: 09/203,958. Alternatively, anti-dendritic cell antibodies, or antigen binding fragments thereof, can be bound to a cell or a pathogen by the use of chemical linkers, lipid tags, or other related methods (deKruif, J. et al. (2000) Nat. Med. 6:223-227; Nizard, P. et al. (1998) FEBS Lett. 433:83-88). Cells with surface-anchored anti-DEC-205 antibodies may be used to induce specific immune responses against the cell, e.g., a tumor cell or microbial pathogen.

### III. Pharmaceutical Compositions

[0140] In another embodiment, the present invention provides a composition, e.g., a pharmaceutical composition, containing one or a combination of monoclonal antibodies of the present invention, formulated together with a pharmaceutically acceptable carrier. Compositions containing bispecific molecules or molecular conjugates which comprise an antibody of the present invention are also provided. In one embodiment, the compositions include a combination of multiple (e.g., two or more) isolated antibodies of the invention. Preferably, each of the antibodies of the composition binds to a distinct, pre-selected epitope of DEC-205.

[0141] Pharmaceutical compositions of the invention also can be administered in combination therapy, i.e., combined with other agents. For example, the combination therapy can include a composition of the present invention with at least one or more additional therapeutic agents, such as anti-inflammatory agents, DMARDs (disease-modifying anti-rheumatic drugs), immunosuppressive agents, and chemotherapeutics. The pharmaceutical compositions of the invention can also be administered in conjunction with radiation therapy. Co-administration with other antibodies, such as CD4 specific antibodies, or IL-2 specific antibodies, are also encompassed by the invention. Such combinations with CD4 specific antibodies or IL-2 specific antibodies are considered particularly useful for treating autoimmune diseases and transplant rejections. Combinations with antibodies to CTLA4, CD40 etc particularly useful in cancer and infectious disease treatments.

[0142] In another embodiment, a vaccine conjugate that is rapidly internalized by APCs can be combined with a monoclonal antibody that enhances antigen presenting cell activities of dendritic cells, e.g., release of immunostimulatory cytokines.

[0143] As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. Preferably, the carrier is suitable for intravenous, intramuscular, subcutaneous, parenteral, spinal or epidermal administration (e.g., by injection or infusion). Depending on the route of administration, the active compound, i.e., antibody, bispecific and multispecific molecule, may be coated in a material to protect the compound from the action of acids and other natural conditions that may inactivate the compound.

[0144] Examples of adjuvants which may be used with the antibodies and constructs of the present invention include: Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, Mich.); Merck Adjuvant 65 (Merck and Company, Inc., Rahway, N.J.); AS-2 (SmithKline Beecham, Philadelphia, Pa.); aluminum salts such as aluminum hydroxide gel (alum) or aluminum phosphate; salts of calcium, iron or zinc; an insoluble suspension of acylated tyrosine; acylated sugars; cationically or anionically derivatised polysaccharides; polyphosphazenes; biodegradable microspheres; cytokines, such as GM-CSF, interleukin-2, -7, -12, and other like factors; 3D-MPL; CpG oligonucleotide; and monophosphoryl lipid A, for example 3-de-O-acylated monophosphoryl lipid A.

[0145] MPL adjuvants are available from Corixa Corporation (Seattle, Wash; see, for example, U.S. Pat. Nos. 4,436,727; 4,877,611; 4,866,034 and 4,912,094). CpG-containing oligonucleotides (in which the CpG dinucleotide is unmethylated) are well known and are described, for example, in WO 96/02555, WO 99/33488 and U.S. Pat. Nos. 6,008,200 and 5,856,462. Immunostimulatory DNA sequences are also described, for example, by Sato et al., Science 273:352, 1996.

[0146] Further alternative adjuvants include, for example, saponins, such as Quil A, or derivatives thereof, including QS21 and QS7 (Aquila Biopharmaceuticals Inc., Framingham, Mass.); Escin; Digitonin; or Gypsophila or Chenopodium quinoa saponins; Montanide ISA 720 (Seppic, France); SAF (Chiron, California, United States); ISCOMS (CSL), MF-59

(Chiron); the SBAS series of adjuvants (e.g., SBAS-2 or SBAS-4, available from SmithKline Beecham, Rixensart, Belgium); Detox (Enhantyn™) (Corixa, Hamilton, Mont.); RC-529 (Corixa, Hamilton, Mont.) and other aminoalkyl glucosamine 4-phosphates (AGPs); polyoxyethylene ether adjuvants such as those described in WO 99/52549A1; synthetic imidazoquinolines such as imiquimod [S-26308, R-837], (Harrison, et al., Vaccine 19: 1820-1826, 2001; and resiquimod [S-28463, R-848] (Vasilakos, et al., Cellular immunology 204: 64-74, 2000; Schiff bases of carbonyls and amines that are constitutively expressed on antigen presenting cell and T-cell surfaces, such as tucaresol (Rhodes, J. et al., Nature 377: 71-75, 1995); cytokine, chemokine and co-stimulatory molecules as either protein or peptide, including for example pro-inflammatory cytokines such as Interferon, GM-CSF, IL-1 alpha, IL-1 beta, TGF-alpha and TGF-beta, Th1 inducers such as interferon gamma, IL-2, IL-12, IL-15, IL-18 and IL-21, Th2 inducers such as IL-4, IL-5, IL-6, IL-10 and IL-13 and other chemokine and co-stimulatory genes such as MCP-1, MIP-1 alpha, MIP-1 beta, RANTES, TCA-3, CD80, CD86 and CD40L; immunostimulatory agents targeting ligands such as CTLA-4 and L-selectin, apoptosis stimulating proteins and peptides such as Fas; synthetic lipid based adjuvants, such as vaxfectin, (Reyes et al., Vaccine 19: 3778-3786, 2001) squalene, alpha-tocopherol, polysorbate 80, DOPC and cholesterol; endotoxin, [LPS], (Beutler, B., Current Opinion in Microbiology 3: 23-30, 2000); ligands that trigger Toll receptors to produce Th1-inducing cytokines, such as synthetic Mycobacterial lipoproteins, Mycobacterial protein p19, peptidoglycan, teichoic acid and lipid A; and CT (cholera toxin, subunits A and B) and LT (heat labile enterotoxin from *E. coli*, subunits A and B), heat shock protein family (HSPs), and LLO (listeriolysin O; WO 01/72329). These and various further Toll-like Receptor (TLR) agonists are described for example in Kanzler et al, Nature Medicine, May 2007, Vol 13, No 5.

**[0147]** A "pharmaceutically acceptable salt" refers to a salt that retains the desired biological activity of the parent compound and does not impart any undesired toxicological effects (see e.g., Berge, S.M., et al. (1977) *J. Pharm. Sci.* 66:1-19). Examples of such salts include acid addition salts and base addition salts. Acid addition salts include those derived from nontoxic inorganic acids, such as hydrochloric, nitric, phosphoric, sulfuric, hydrobromic, hydroiodic, phosphorous and the like, as well as from nontoxic organic acids such as aliphatic mono- and dicarboxylic acids, phenyl-substituted alkanoic acids, hydroxy alkanoic acids, aromatic acids, aliphatic and aromatic sulfonic acids and the like. Base addition salts include those derived from alkaline earth metals, such as sodium, potassium, magnesium, calcium and the like, as well as from nontoxic organic amines, such as N,N'-dibenzylethylenediamine, N-methylglucamine, chloroprocaine, choline, diethanolamine, ethylenediamine, procaine and the like.

**[0148]** A composition of the present invention can be administered by a variety of methods known in the art. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. The active compounds can be prepared with carriers that will protect the compound against rapid release, such as a controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation of such formulations are patented or generally known to those skilled in the art. See, e.g., *Sustained and Controlled Release Drug Delivery Systems*, J.R. Robinson, ed., Marcel Dekker, Inc., New York, 1978.

**[0149]** To administer a compound of the invention by certain routes of administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation. For example, the compound may be administered to a subject in an appropriate carrier, for example, liposomes, or a diluent. Pharmaceutically acceptable diluents include saline and aqueous buffer solutions. Liposomes include water-in-oil-in-water CGF emulsions as well as conventional liposomes (Strejan et al. (1984) *J. Neuroimmunol.* 7:27).

**[0150]** Pharmaceutically acceptable carriers include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the pharmaceutical compositions of the invention is contemplated. Supplementary active compounds can also be incorporated into the compositions.

**[0151]** Therapeutic compositions typically must be sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, microemulsion, liposome, or other ordered structure suitable to high drug concentration. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.

**[0152]** Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by sterilization microfiltration. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of

sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying (lyophilization) that yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

**[0153]** Dosage regimens are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. For example, the antibodies of the invention may be administered once or twice weekly by subcutaneous injection or once or twice monthly by subcutaneous injection.

**[0154]** It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

**[0155]** Examples of pharmaceutically-acceptable antioxidants include: (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

**[0156]** For the therapeutic compositions, formulations of the present invention include those suitable for oral, nasal, topical (including buccal and sublingual), rectal, vaginal and/or parenteral administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods known in the art of pharmacy. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will vary depending upon the subject being treated, and the particular mode of administration. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will generally be that amount of the composition which produces a therapeutic effect. Generally, out of one hundred per cent, this amount will range from about 0.001 per cent to about ninety percent of active ingredient, preferably from about 0.005 per cent to about 70 per cent, most preferably from about 0.01 per cent to about 30 per cent.

**[0157]** Formulations of the present invention which are suitable for vaginal administration also include pessaries, tampons, creams, gels, pastes, foams or spray formulations containing such carriers as are known in the art to be appropriate. Dosage forms for the topical or transdermal administration of compositions of this invention include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. The active compound may be mixed under sterile conditions with a pharmaceutically acceptable carrier, and with any preservatives, buffers, or propellants which may be required.

**[0158]** The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, epidural and intrasternal injection and infusion.

**[0159]** Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

**[0160]** These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of presence of microorganisms may be ensured both by sterilization procedures, supra, and by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

**[0161]** When the compounds of the present invention are administered as pharmaceuticals, to humans and animals, they can be given alone or as a pharmaceutical composition containing, for example, 0.001 to 90% (more preferably, 0.005 to 70%, such as 0.01 to 30%) of active ingredient in combination with a pharmaceutically acceptable carrier.

**[0162]** Regardless of the route of administration selected, the compounds of the present invention, which may be used in a suitable hydrated form, and/or the pharmaceutical compositions of the present invention, are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art.

**[0163]** Actual dosage levels of the active ingredients in the pharmaceutical compositions of the present invention may

be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient. The selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, or the ester, salt or amide thereof, the route of administration, the time of administration, 5 the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts. A physician or veterinarian having ordinary skill in the art can readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the compounds of the 10 invention employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved. In general, a suitable daily dose of a composition of the invention will be that amount of the compound which is the lowest dose effective to produce a therapeutic effect. Such an effective dose will generally depend upon the factors described above. It is preferred that administration be intravenous, intramuscular, intraperitoneal, or subcutaneous, preferably administered proximal to the 15 site of the target. If desired, the effective daily dose of a therapeutic composition may be administered as two, three, four, five, six or more sub-doses administered separately at appropriate intervals throughout the day, optionally, in unit dosage forms. While it is possible for a compound of the present invention to be administered alone, it is preferable to administer the compound as a pharmaceutical formulation (composition).

**[0164]** Therapeutic compositions can be administered with medical devices known in the art. For example, in a preferred 20 embodiment, a therapeutic composition of the invention can be administered with a needleless hypodermic injection device, such as the devices disclosed in U.S. Patent Nos. 5,399,163, 5,383,851, 5,312,335, 5,064,413, 4,941,880, 4,790,824, or 4,596,556. Examples of well-known implants and modules useful in the present invention include: U.S. Patent No. 4,487,603, which discloses an implantable micro-infusion pump for dispensing medication at a controlled rate; U.S. Patent No. 4,486,194, which discloses a therapeutic device for administering medicants through the skin; 25 U.S. Patent No. 4,447,233, which discloses a medication infusion pump for delivering medication at a precise infusion rate; U.S. Patent No. 4,447,224, which discloses a variable flow implantable infusion apparatus for continuous drug delivery; U.S. Patent No. 4,439,196, which discloses an osmotic drug delivery system having multi-chamber compartments; and U.S. Patent No. 4,475,196, which discloses an osmotic drug delivery system. Many other such implants, delivery systems, and modules are known to those skilled in the art.

**[0165]** In certain embodiments, the antibodies of the invention can be formulated to ensure proper distribution *in vivo*. 30 For example, the blood-brain barrier (BBB) excludes many highly hydrophilic compounds. To ensure that the therapeutic compounds of the invention cross the BBB (if desired), they can be formulated, for example, in liposomes. For methods of manufacturing liposomes, see, e.g., U.S. Patents 4,522,811; 5,374,548; and 5,399,331. The liposomes may comprise one or more moieties which are selectively transported into specific cells or organs, thus enhance targeted drug delivery 35 (see, e.g., V.V. Ranade (1989) *J. Clin. Pharmacol.* 29:685). Exemplary targeting moieties include folate or biotin (see, e.g., U.S. Patent 5,416,016 to Low et al.); mannosides (Umezawa et al., (1988) *Biochem. Biophys. Res. Commun.* 153: 1038); antibodies (P.G. Bloeman et al. (1995) *FEBS Lett.* 357:140; M. Owais et al. (1995) *Antimicrob. Agents Chemother.* 39:180); surfactant protein A receptor (Briscoe et al. (1995) *Am. J. Physiol.* 1233:134), different species of which may 40 comprise the formulations of the inventions, as well as components of the invented molecules; p120 (Schreier et al. (1994) *J. Biol. Chem.* 269:9090); see also K. Keinanen; M.L. Laukkonen (1994) *FEBS Lett.* 346:123; J.J. Killion; I.J. Fidler (1994) *Immunomethods* 4:273. In one embodiment of the invention, the therapeutic compounds of the invention 45 are formulated in liposomes; in a more preferred embodiment, the liposomes include a targeting moiety. In a most preferred embodiment, the therapeutic compounds in the liposomes are delivered by bolus injection to a site proximal to the tumor or infection. The composition must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi.

**[0166]** The ability of a compound to inhibit cancer can be evaluated in an animal model system predictive of efficacy 50 in human tumors. Alternatively, this property of a composition can be evaluated by examining the ability of the compound to inhibit, such inhibition *in vitro* by assays known to the skilled practitioner. A therapeutically effective amount of a therapeutic compound can decrease tumor size, or otherwise ameliorate symptoms in a subject. One of ordinary skill in the art would be able to determine such amounts based on such factors as the subject's size, the severity of the subject's symptoms, and the particular composition or route of administration selected.

**[0167]** The ability of the antibodies to enhance antigen presentation or induce cytotoxic T cell (CTL) responses against 55 a variety of target cells or pathogens can also be evaluated according to methods well known in the art.

**[0168]** The composition must be sterile and fluid to the extent that the composition is deliverable by syringe. In addition to water, the carrier can be an isotonic buffered saline solution, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. Proper fluidity can be maintained, for example, by use of coating such as lecithin, by maintenance of required particle size in the case of dispersion and by use of

surfactants. In many cases, it is preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol or sorbitol, and sodium chloride in the composition. Long-term absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate or gelatin.

5 [0169] When the active compound is suitably protected, as described above, the compound may be orally administered, for example, with an inert diluent or an assimilable edible carrier.

#### IV. Uses and Methods of the Invention

10 [0170] In one embodiment, the antibodies, bispecific molecules, and molecular conjugates of the present invention can be used to treat and/or prevent (e.g., immunize against) a variety of diseases and conditions.

15 [0171] One of the primary disease indications that can be treated using antibodies of the invention is cancer. This includes, but is not limited to, colon cancer, melanoma, lymphoma, prostate carcinoma, pancreatic carcinoma, bladder carcinoma, fibrosarcoma, rhabdomyosarcoma, mastocytoma, mammary adenocarcinoma, leukemia, or rheumatoid fibroblastsoma. Another primary disease indication is infectious diseases including, but not limited to, HIV, Hepatitis (e.g.,

16 A, B, & C), Influenza, Herpes, Giardia, Malaria, Leishmania, Staphylococcus Aureus, Pseudomonas aeruginosa. Another primary disease indication includes autoimmune diseases.

20 [0172] For use in therapy, vaccine conjugates of the invention can be administered to a subject directly (i.e., *in vivo*), either alone or with an immunostimulatory agent. In one aspect, the immunostimulatory agent is linked to the conjugate. Alternatively, the conjugates can be administered to a subject indirectly by first contacting the conjugates (e.g., by 25 culturing or incubating) with APCs, such as dendritic cells, and then administering the cells to the subject (i.e., *ex vivo*). The contacting and delivering of the conjugates to APCs, such that they are processed and presented by the APCs prior to administration, is also referred to as antigen or cell "loading." Techniques for loading antigens to APCs are well known in the art and include, for example, Gunzer and Grabbe, Crit Rev Immunol 21 (1-3):133-45 (2001) and Steinman, Exp Hematol 24(8): 859-62 (1996).

25 [0173] In all cases, the vaccine conjugates and the immunostimulatory agents are administered in an effective amount to exert their desired therapeutic effect. The term "effective amount" refers to that amount necessary or sufficient to realize a desired biologic effect. For example, an effective amount could be that amount necessary to eliminate a tumor, cancer, or bacterial, viral or fungal infection. The effective amount for any particular application can vary depending on such factors as the disease or condition being treated, the particular conjugate being administered, the size of the subject, 30 or the severity of the disease or condition. One of ordinary skill in the art can empirically determine the effective amount of a particular multispecific molecule without necessitating undue experimentation.

35 [0174] Preferred routes of administration for the vaccine conjugates include, for example, injection (e.g., subcutaneous, intravenous, parenteral, intraperitoneal, intrathecal). The injection can be in a bolus or a continuous infusion. Other routes of administration include oral administration.

40 [0175] Vaccine conjugates of the invention also can be coadministered with adjuvants and other therapeutic agents. It will be appreciated that the term "coadministered" as used herein includes any or all of simultaneous, separate, or sequential administration of the antibodies and conjugates of the present invention with adjuvants and other agents, including administration as part of a dosing regimen. The conjugates are typically formulated in a pharmaceutically acceptable carrier alone or in combination with such agents. Examples of such carriers include solutions, solvents, dispersion media, delay agents, emulsions and the like. The use of such media for pharmaceutically active substances is well known in the art. Any other conventional carrier suitable for use with the molecules falls within the scope of the instant invention.

45 [0176] Suitable agents for coadministration with the vaccine conjugates include other antibodies, cytotoxins and/or drugs. In one embodiment, the agent is an anti-CTLA-4 antibody which is known to aid or induce immune responses. In another embodiment, the agent is a chemotherapeutic agent. The vaccine conjugates also can be administered in combination with radiation.

50 [0177] Chemotherapeutic agents suitable for coadministration with the antibodies and conjugates of the present invention in the treatment of tumors include, for example: taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicine, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Further agents include, for example, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclothosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine) and temozolamide.

55 [0178] Agents that delete or inhibit immunosuppressive activities, for example, by immune cells (for example regulatory

T-cells, NKT cells, macrophages, myeloid-derived suppressor cells, immature or suppressive dendritic cells) or suppressive factors produced by the tumor or host cells in the local microenvironment of the tumor (for example, TGFbeta, indoleamine 2,3 dioxygenase - IDO), may also be administered with the antibodies and conjugates of the present invention. Such agents include antibodies and small molecule drugs such as IDO inhibitors such as 1 methyl tryptophan or derivatives.

**[0179]** In another embodiment, the antibodies of the present invention can be used to treat a subject with an autoimmune, immune system, or inflammatory disorder, e.g., a disorder characterized by aberrant or unwanted immune activity associated with immunomodulation by dendritic cells. Autoimmune, immune system, and inflammatory disorders that may benefit from treatment with the anti-dendritic cells of the invention include rheumatoid arthritis, multiple sclerosis, immune-mediated or Type 1 diabetes mellitus, myasthenia gravis, pernicious anemia, Addison's disease, Sjogren's syndrome, psoriasis, lupus erythematosus, inflammatory bowel diseases such as Crohn's disease and ulcerative colitis, scleroderma/Raynaud's syndrome, Reiter's syndrome, and autoimmune thyroid diseases such as Hashimoto's thyroiditis and Graves's disease. For example, a subject suffering from an autoimmune disorder may benefit from inhibition of dendritic cell mediated presentation of an autoantigen..

**[0180]** The antibodies of the present invention may also be used for preventing and treating all forms of allergy and allergic disorder, including without limitation: ophthalmic allergic disorders, including allergic conjunctivitis, vernal conjunctivitis, vernal keratoconjunctivitis, and giant papillary conjunctivitis; nasal allergic disorders, including allergic rhinitis and sinusitis; otic allergic disorders, including eustachian tube itching; allergic disorders of the upper and lower airways, including intrinsic and extrinsic asthma; allergic disorders of the skin, including dermatitis, eczema and urticaria; and allergic disorders of the gastrointestinal tract.

**[0181]** Suitable agents for coadministration with the antibodies of the present invention for treatment of such immune disorders include for example, immunosuppressive agents such as rapamycin, cyclosporin and FK506; anti-TNF $\alpha$  agents such as etanercept, adalimumab and infliximab; and steroids. Examples of specific natural and synthetic steroids include, for example: aldosterone, beclomethasone, betamethasone, budesonide, clobrednol, cortisone, cortivazol, deoxycorticosterone, desonide, desoximetasone, dexamethasone, difluorocortolone, fluclorolone, flumethasone, flunisolide, fluocinolone, fluocinonide, fluocortin butyl, fluorocortisone, fluorocortolone, fluorometholone, flurandrenolone, fluticasone, halcinonide, hydrocortisone, icomethasone, meprednisone, methylprednisolone, paramethasone, prednisolone, prednisone, tixocortol and triamcinolone.

**[0182]** Other examples of diseases that can be treated using the anti-DEC-205 antibodies of the invention include transplant rejection and graft versus host disease.

#### Transplant Rejection

**[0183]** Over recent years there has been a considerable improvement in the efficiency of surgical techniques for transplanting tissues and organs such as skin, kidney, liver, heart, lung, pancreas and bone marrow. Perhaps the principal outstanding problem is the lack of satisfactory agents for inducing immune-tolerance in the recipient to the transplanted allograft or organ. When allogeneic cells or organs are transplanted into a host (i.e., the donor and donee are different individual from the same species), the host immune system is likely to mount an immune response to foreign antigens in the transplant (host-versus-graft disease) leading to destruction of the transplanted tissue. CD8+ cells, CD4+ cells and monocytes are all involved in the rejection of transplant tissues. The antibodies of the present invention are useful to inhibit dendritic cell mediated alloantigen-induced immune responses in the donee thereby preventing such cells from participating in the destruction of the transplanted tissue or organ.

#### Graft Versus Host Disease

**[0184]** A related use for the antibodies of the present invention is in modulating the immune response involved in "graft versus host" disease (GVHD). GVHD is a potentially fatal disease that occurs when immunologically competent cells are transferred to an allogeneic recipient. In this situation, the donor's immunocompetent cells may attack tissues in the recipient. Tissues of the skin, gut epithelia and liver are frequent targets and may be destroyed during the course of GVHD. The disease presents an especially severe problem when immune tissue is being transplanted, such as in bone marrow transplantation; but less severe GVHD has also been reported in other cases as well, including heart and liver transplants. The therapeutic agents of the present invention are used to inhibit the activity of host antigen presenting cells, e.g., dendritic cells.

**[0185]** The present invention is further illustrated by the following examples which should not be construed as further limiting.

**EXAMPLES****Example 1**5 **Generation of DEC-205-Specific Human Monoclonal Antibodies (HuMabs)**

[0186] Human anti-DEC-205 monoclonal antibodies were generated by immunizing the HC2/KCo7 strain of HuMab® transgenic mice ("HuMab" is a Trade Mark of Medarex, Inc., Princeton, New Jersey) with a soluble human DEC-205 antigen. HC2/KCo7 HuMab mice were generated as described in U.S. Pat. Nos. 5,770,429 and 5,545,806.

10 [0187] Antigen and Immunization: The antigen was a soluble fusion protein comprising a DEC-205 extracellular domain (comprising all ten lectin-binding domains) fused with an antibody Fc domain. A nucleic acid and amino acid sequence of human DEC-205 is provided in PCT Patent Publication No WO 96023882 (Steinman). The antigen was mixed with Complete Freund's (Sigma) adjuvant for the first immunization. Thereafter, the antigen was mixed with Incomplete Freund's (Sigma). Additional mice were immunized with the soluble DEC-205 protein in RIBI MPL plus TDM adjuvant 15 system (Sigma). 5-25 micrograms soluble recombinant DEC-205 antigen in PBS or  $5 \times 10^6$  CHO cells transfected for surface expression of human DEC-205 in PBS were mixed 1:1 with the adjuvant. Mice were injected with 100 microliters of the prepared antigen into the peritoneal cavity every 14 days. Animals that developed anti-DEC-205 titers were given an iv injection of 10 micrograms soluble recombinant DEC-205 antigen three to four days prior to fusion. Mouse spleens were harvested, and the isolated splenocytes used for hybridoma preparation.

20 [0188] Hybridoma Preparation: The P3x63Ag8.653 murine myeloma cell line (ATCC CRL 1580) was used for the fusions. RPMI 1640 (Invitrogen) containing 10% FBS, and was used to culture the myeloma cells. Additional media supplements were added to the Hybridoma growth media, which included: 3% Ongen-Hybridoma Cloning Factor (Igen), 10% FBS (Sigma), L-glutamine (Gibco) 0.1 % gentamycin (Gibco), 2-mercaptoethanol (Gibco), HAT (Sigma;  $1.0 \times 10^{-4}$  M hypoxanthine,  $4.0 \times 10^{-7}$  M aminopterin,  $1.6 \times 10^{-5}$  M thymidine), or HT (Sigma;  $1.0 \times 10^{-4}$  M hypoxanthine,  $1.6 \times 10^{-5}$  M thymidine) media.

25 [0189] Spleen cells were mixed with the 653 myeloma cells in a 6:1 ratio and pelleted by centrifugation. Polyethylene glycol was added dropwise with careful mixing to facilitate fusion. Hybridomas were allowed to grow out for one to two weeks until visible colonies become established. Supernatant was harvested and used for initial screening for human IgG via ELISA using a human kappa chain specific capture and a human Fc specific detection. IgG positive supernatants 30 were then assayed for DEC-205 specificity via flow cytometry or using a DEC-205 ELISA.

[0190] Hybridomas producing specific HuMab IgG were subcloned and expanded. The HuMabs produced were then purified by protein A column chromatography according to standard conditions which led to the isolation of a number of antibodies of particular interest.

35 **Example 2****Determination of Affinity and Rate Constants of HuMabs by Surface Plasmon Resonance (SPR)**

[0191] Binding affinity and binding kinetics of various human anti-DEC-205 antibodies from Example 1 were examined 40 by Biacore™ surface plasmon resonance (SPR) analysis using a Biacore™ 2000 SPR instrument (Biacore AB, Uppsala, Sweden) according to the manufacturer's guidelines.

45 [0192] Purified recombinant human DEC-205 fusion (or control) protein was covalently linked to a Biacore™ CM5 sensor chip (carboxymethylated dextran covalently attached to a gold surface; Biacore Product No. BR-1000-14) using standard amine coupling chemistry with an Amine Coupling Kit provided by Biacore according to the manufacturer's guidelines (Biacore Product No. BR-1000-50, comprising coupling reagents N-hydroxysuccinimide (NHS) and 1-Ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (EDC)). Low levels of ligand were immobilised to limit any effects of mass transport of analyte on kinetic parameters, such that the  $R_{MAX}$  observed was in the order of 200 RU.

50 [0193] Binding was measured by flowing the antibodies over the sensor chip in HBS-NP buffer (HBS-N buffer, Biacore Product No. BR-1003-69: 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) 0.24%, sodium chloride 0.88%, qs water, filtered/de-gassed and pre-equilibrated to room temperature with a 1:2000 dilution of Surfactant P20) at concentrations ranging from 1.25 to 200 nM and at a flow rate of 35  $\mu$ l/minute. The antigen-antibody association and dissociation kinetics were followed for approximately 300 to 600 seconds in each case.

55 [0194] Corresponding controls were conducted in each case using an unrelated protein for "background" subtraction. A single injection of 18 mM NaOH for 17 seconds at 35  $\mu$ l/min was used as the regeneration conditions throughout the study.

[0195] Biacore's kinetics wizard was used in each case to derive kinetic parameters from the concentration series of analyte diluted in HBS-NP running buffer. The association and dissociation curves were fitted to a 1:1 Langmuir binding model using Biacore™ kinetics wizard software (Biacore AB) according to the manufacturer's guidelines. The affinity

and kinetic parameters (with background subtracted) as determined are shown in Table 1 below. For each antibody, the figures shown are the mean of two separate series of experiments, using separately prepared sensor chips in each case, (where  $k_a$  = rate constant of association,  $k_d$  = rate constant of dissociation,  $K_D$  = dissociation equilibrium constant (measure of affinity),  $K_A$  = association equilibrium constant,  $R_{max}$  = maximum SPR response signal).

5

10

15

20

**Table 1**

mAb #	mAb ID	$k_a$ (1/Ms)	$k_d$ (1/s)	$K_A$ (1/M)	$K_D$ (M)	$R_{max}$ (RU)
#1	3A4-1C10	$1.5 \times 10^6$	$9.6 \times 10^{-5}$	$1.6 \times 10^{10}$	$6.6 \times 10^{-11}$	278
#2	5A8-1F1	$3.6 \times 10^5$	$2.0 \times 10^{-4}$	$2.1 \times 10^9$	$1.5 \times 10^{-9}$	172
#3	3C7-3A3	$1.7 \times 10^5$	$7.6 \times 10^{-4}$	$5.2 \times 10^8$	$5.6 \times 10^{-9}$	133
#4	2D3-1F5	$3.3 \times 10^5$	$2.2 \times 10^{-5}$	$1.5 \times 10^{10}$	$6.8 \times 10^{-11}$	275
#5	3D6-2F4	$1.8 \times 10^6$	$1.2 \times 10^{-4}$	$1.5 \times 10^{10}$	$8.0 \times 10^{-11}$	294
#6	5D12-5G1	$5.4 \times 10^5$	$3.2 \times 10^{-4}$	$2.0 \times 10^9$	$7.0 \times 10^{-10}$	272
#7	1G6-1G6	$1.4 \times 10^6$	$3.0 \times 10^{-4}$	$4.7 \times 10^9$	$2.3 \times 10^{-10}$	249
#8	3G9-2D2	$9.0 \times 10^5$	$1.9 \times 10^{-4}$	$4.7 \times 10^9$	$2.4 \times 10^{-10}$	268

**Example 3****Binding of HuMabs to cells expressing human DEC-205**

25

**[0196]** The ability of anti-DEC-205 HuMabs to bind to DEC-205 on CHO-S cells expressing human DEC-205 on their surface was investigated by flow cytometry as follows.

30

**[0197]** Antibodies were tested for binding to CHO-S cells expressing human DEC-205 on their surface. Protein A purified HuMabs 3D6-2F4, 3D6-4C8, 3G9-2D2, 5A8-1F1, 2D3-1F5-2A9, 3C7-3A3, 5D12-5G1, 1G6-1G6 and 3A4-1C10 were incubated with the CHO-S cells expressing human DEC-205, as well as CHO-S control cells at 4°C. All antibodies were used at saturating concentrations. After 1 hour, the cells were washed with PBS containing 0.1% BSA and 0.05%  $\text{NaN}_3$  (PBA) and the bound antibodies were detected by incubating the cells with a PE labeled goat anti-human IgG Fc-specific probe, at 4°C. The excess probe was washed from the cells with PBA and the cell associated fluorescence was determined by analysis using a LSR™ instrument (BD Biosciences, NJ, USA) according to the manufacturer's directions. Results are shown in Figure 1.

35

**[0198]** As shown in Figure 1, the HuMabs demonstrated high level binding to CHO-S cells expressing human DEC-205. These data demonstrate that these antibodies bind efficiently and specifically to human DEC-205 expressed on live CHO-S cells compared to the control cells.

40

**Example 4****Binding of HuMabs to human dendritic cells**

45

**[0199]** Human peripheral blood mononuclear cells (PBMCs) were obtained by density gradient centrifugation of Leukopak platelet apheresis preparations. Monocytes were isolated by adherence to tissue culture flasks for two hours, and then differentiated into dendritic cells by incubation with 2 ng/ml GM-CSF and 10 ng/ml IL-4 in macrophage serum free media (Gibco) for 5 to 7 days.

50

**[0200]** The ability of anti-DEC-205 HuMabs to bind to DEC-205 on human dendritic cells prepared as above was investigated by flow cytometry as follows.

55

**[0201]** Protein A purified HuMabs 3D6-2F2, 3D6-4C8, 3G9-2D2, 5A8-1F1, 2D3-1F5-2A9, 3C7-3A3, 5D12-5G1, 1G6-1G6 and 3A4-1C10, and an isotype control (human IgG) were incubated with the human dendritic cells at 4°C. All antibodies were used at saturating concentrations. After 1 hour, the cells were washed with PBS containing 0.1% BSA and 0.05%  $\text{NaN}_3$  (PBA) and the bound antibodies were detected by incubating the cells with a PE labeled goat anti-human IgG Fc-specific probe, at 4°C. The excess probe was washed from the cells with PBA and the cell associated fluorescence was determined by analysis using a LSR™ instrument (BD Biosciences, NJ, USA) according to the manufacturer's directions. Results are shown in Figure 2, which shows that the HuMabs demonstrated high level binding to human dendritic cells compared to the isotype control.

**Example 5****ELISA Assay to Determine HuMAb Binding Characteristics on DEC-205**

5 [0202] Microtiter plates were coated with soluble DEC-205/Fc fusion protein in PBS, and then blocked with 5% bovine serum albumin in PBS. Protein A purified HuMabs and an isotype control were added at saturating concentrations and incubated at 37°C. The plates were washed with PBS/Tween and then incubated with a goat-anti-human IgG Fc-specific polyclonal reagent conjugated to alkaline phosphatase at 37°C. After washing, the plates were developed with pNPP substrate (1 mg/ml), and analyzed at OD 405-650 using a microtiter plate reader. Results are shown in Figure 3, which shows that the HuMabs demonstrated high level binding compared to the isotype control.

**Example 6****Antibody internalization assay**

15 [0203] Human monocyte-derived dendritic cells 5 x 10<sup>5</sup> per aliquot were incubated with human IgG (1 mg/ml) to block non-specific binding. Cells were then incubated for 30 minutes on ice with 100 µg/ml of FITC-conjugated anti-Dec-205 HuMab 3G9-2D2 in blocking buffer for binding, and subsequently transferred to 37°C for 0, 10, 30, 60 and 120 minutes for internalization. FITC-conjugated human IgG1 at same concentration was used as control. Cells were then washed 20 and fixed with 1% paraformaldehyde. Fixed cells were washed, resuspended in water, and cytospon onto microscope slides. Images were taken with a Zeiss LSM 510 Meta confocal microscope. Results are shown in Figure 4, which shows that the FITC-labelled HuMabs demonstrated efficient internalization into the dendritic cells compared to the control.

**Example 7****Antibody Sequencing**

25 [0204] As described above in Example 1, HuMabs from hybridomas producing specific HuMab IgG were purified by protein A column chromatography which led to the isolation of eight antibodies ("HuMabs") of particular interest. The V<sub>H</sub> and V<sub>L</sub> coding regions of HuMabs 3D6-2F4, 3D6-4C8, 3G9-2D2, 5A8-1F1, 2D3-1F5-2A9 (V<sub>H</sub> region), 3C7-3A3, 1E6-3D10 (V<sub>H</sub> region) and 5C3-2-3F6 were identified using RNA from the corresponding hybridomas. RNA was reverse transcribed to cDNA, the V coding regions were amplified by PCR and the PCR product was sequenced. The following are the nucleic and amino acid sequences of the V<sub>H</sub> and V<sub>L</sub> regions of the HuMabs (in the case of the amino acid sequences, the Complementarity Determining Regions (CDRs) are underlined).

35 3D6-2F4 V<sub>H</sub> nucleic acid sequence (VH3, locus 3-33; JH4) (SEQ ID NO:2):

40 atggagttgggctgagctgggtttcctcggtcttttaagagggtgtccagtgtcaggtgcagctgtggaggtctggggag  
gcgtggccagccctgggagggtccctgagactctcctgtcagcgctgtggattcatcttcagttatctatggcatgcactgggtcccg  
ccaggctccaggcaaggggctggagtgggtggcagttatatgtatggaaagtaataactatgcagactccgtgaagg  
gccgattcaccatctccagagacaattccaagaacacgcgttatctgcaaatacgtggccagagccgaggacacggctgtg  
45 tattactgtgcgagagctcctcactttgactactggggccagggaaacctggtcaccgtctcctcagctagc

3D6-2F4 V<sub>H</sub> amino acid sequence (SEQ ID NO:3) including signal peptide:

50 MEGGLSWVFLVALLRGVQCQVQLVESGGVVQPGRSRLSCAASGFIFSIYGM  
HWVRQAPGKGLEWVAVIWYDGSNKYYADSVKGRFTISRDNSKNTLYLQMNSL  
RAEDTAVYYCARAPHFDYWQGTLTVSS

55 3D6-2F4 V<sub>H</sub> "mature" amino acid sequence (SEQ ID NO: 4) excluding signal peptide:  
QVQLVESGGVVQPGRSRLSCAASGFIFSIYGMHWVRQAPGKGLEWVAWTWY  
DGSNKYYADSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCARAPHFDYW GQGTLTVSS

## EP 2 224 954 B1

3D6-2F4 V<sub>H</sub> CDR1 (SEQ ID NO: 5): IYGMH

3D6-2F4 V<sub>H</sub> CDR2 (SEQ ID NO: 6): VIWYDGSNKYYADSVKG

5 3D6-2F4 V<sub>H</sub> CDR3 (SEQ ID NO: 7): APHFDY

3D6-2F4 V<sub>L</sub> nucleic acid sequence (VK1, locus L15; JK2) (SEQ ID NO: 8):

10 atggatggagctgtatcatctgttccctgtggccacagcaaccgggtgtccactccgacatccagatgaccaggacttccatcc  
cactgtctgcacatgttggagacagagtaccatcacttgtcggggcgggtcaggattagcagctggtagcctggatcaga  
gaaaccagagaaagccctaagtccctgatctatgcacatccagggtttgcaaagtgggtccatcaagggtcagcggcagtg  
15 gatctggacagatttcacttcaccatcagcagcctgcagcctgaagatttgcaacttattactgccaacagtataatagttacc  
cgtacactttggccaggggaccaagctggagatcaaacgtacg

3D6-2F4 V<sub>L</sub> amino acid sequence (SEQ ID NO: 9) including signal peptide:

20 MDMRVLAQLLGLLLCFPGARCDIQMTQSPSSLSASVGDRVITCRASQGISSW  
LAWYQQKPEKAPKSLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYY  
25 CQQYNSYPYTFGQGTKLEIK

3D6-2F4 V<sub>L</sub> "mature" amino acid sequence (SEQ ID NO: 10) excluding signal peptide:

30 DIQMTQSPSSLSASVGDRVITCRASQGISSWLAWYQQKPEKAPKSLIYAASSLQ  
SGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYNSYPYTFGQGTKLEIK

3D6-2F4 V<sub>L</sub> CDR1 (SEQ ID NO: 11): RASQGISSWLA

35 3D6-2F4 V<sub>L</sub> CDR2 (SEQ ID NO: 12): AASSLQS

3D6-2F4 V<sub>L</sub> CDR3 (SEQ ID NO: 13): QQYNSYPY

40 3D6-4C8 V<sub>H</sub> nucleic acid sequence (VH3, locus 3-33; JH4) (SEQ ID NO: 14):

45 atggagttggctgagctgggtttccctgttgcttttaagagggtgtccagtgtcaggcgtggagttctggggag  
gctgggtccagccctggagggtccctgagactctctgtcagcgtctggattcatcttcagttatctatggcatgcactgggtccg  
ccaggctccaggcaaggggctggagtgggtggcagttatatggatgaaagtaataactatgcagactccgtgaagg  
50 gccgattcaccatctccagagacaattccaagaacacgcgttatctgaaatgaacagcctgagagccgaggacacggctgtg  
tattactgtgcgagagctcctactttgactactggggccagggaaacctggtcaccgtctccatgcctccaccaaggcccc  
tcggcttccccctggcac

3D6-4C8 V<sub>H</sub> amino acid sequence (SEQ ID NO: 15) including signal peptide:

55

5 MEFGLSWVFLVALLRGVQCQVQLVESGGGVVQPGRSRLSCAASGFIFSIYGM  
HWVRQAPGKGLEWVAVIWYDGSNKYYADSVKGRFTISRDNSKNTLYLQMNSL  
RAEDTAVYYCARAPHFDYWQGTLVTVSS

3D6-4C8 V<sub>H</sub> "mature" amino acid sequence (SEQ ID NO: 16) excluding signal peptide:

10 QVQLVESGGGVVQPGRSRLSCAASGFIFSIYGMHWVRQAPGKGLEWVAVIWY  
DGSNKYYADSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCARAPHFDYW  
GQGTLVTVSS

15 3D6-4C8 V<sub>H</sub>CDR1 (SEQ ID NO: 17): IYGMH

3D6-4C8 V<sub>H</sub>CDR2 (SEQ ID NO: 18): VIWYDGSNKYYADSVKG

20 3D6-4C8 V<sub>H</sub>CDR3 (SEQ ID NO: 19): APHFDY

25 3D6-4C8 V<sub>L</sub> nucleic acid sequence (VK1, locus L4; JK4) (SEQ ID NO: 20):

atggacatgagggtccccgctcagctcctgggcttctgctgctggctccagggtgccagatgtgccatccagttgacccag  
tctccatcctccctgtctgcacatctgttaggagacagagtcaccatcacttgccgggcaagtcagggcattagcagtgccttagcct  
ggatcagcagaaccaggaaagctctaagctctgatctatgatgcctccagttggaaagtgggtccatcaagggtca  
gccccagtgatctggacagatttcacttcaccatcagcagcctgcagcctgaagatttgcaacttattactgtcaacagtt  
30 aatagttaccctctactttcggcggaggaccaggtggagatcaaa

3D6-4C8 V<sub>L</sub> amino acid sequence (SEQ ID NO: 21) including signal peptide:

35 MDMRVPAQLLGLLLWLPGARCAIQLTQSPSSLASAVGDRVITCRASQGISSAL  
AWYQQKPGKAPKLLIYDASSLESGVPSRFSGSGTDFLTISLQPEDFATYYC  
QQFNSYPLTFGGGTKVEIK

40 3D6-4C8 V<sub>L</sub> "mature" amino acid sequence (SEQ ID NO: 22) excluding signal peptide:

45 AIQLTQSPSSLASAVGDRVITCRASQGISSALAWYQQKPGKAPKLLIYDASSLES  
GVPSRFSGSGTDFLTISLQPEDFATYYCQQFNSYPLTFGGGTKVEIK

3D6-4C8 V<sub>L</sub>CDR1 (SEQ ID NO: 23): RASQGISSALA

50 3D6-4C8 V<sub>L</sub>CDR2 (SEQ ID NO: 24): DASSLES

3D6-4C8 V<sub>L</sub>CDR3 (SEQ ID NO: 25): QQFNSYPLT

3G9-2D2, V<sub>H</sub> nucleic acid sequence (VH3, locus 3-33; D undetermined; JH4) (SEQ ID NO: 26):

55

5 atggagttgggctgagctgggtttcctcgcttttaagagggtgtccagtgtcaggtcagctggtagtctggggag  
 gcgtggccagccctgggagggtccctgagactctcctgtcagcgtctggattcacccatgtactatggcatgtactgggtccg  
 ccaggctccaggcaaggggctggagtggcagttatatgtatggagaataaaatactatgcagactccgtgaagg  
 10 gccgattcaccatctccagagacaattcaagaacacacgctgtatctgcaaatacgaacagcctgagagccgaggacacggctgt  
 tattactgtcgagagatctctgggatgtactttgacttggccaggaaacctggtcaccgtctcagctagc

10 3G9-2D2,  $V_H$  amino acid sequence (SEQ ID NO: 27) including signal peptide:

MEFGLSWVFLVALLRGVQCQVQLVESGGVVQPGRLRLSCAASGFTFSNYGM  
 15 YWVRQAPGKGLEWVAVIWYDGSNKYYADSVKGRFTISRDNSKNTLYLQMNSL  
RAEDTAVYYCARDLWGWYFDYWGQGTLVTVSSASTKGPSVFPLA

3G9-2D2,  $V_H$  "mature" amino acid sequence (SEQ ID NO: 28) excluding signal peptide:

20 QVQLVESGGVVQPGRLRLSCAASGFTFSNYGMYWVRQAPGKGLEWVAVIW  
YDGSNKYYADSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCARDLWGWY  
 25 FDYWGQGTLVTVSSASTKGPSVFPLA

3G9-2D2,  $V_H$ CDR1 (SEQ ID NO: 29): NYGM

3G9-2D2,  $V_H$  CDR2 (SEQ ID NO: 30): VIWYDGSNKYYADSVKG

30 3G9-2D2,  $V_H$ CDR3 (SEQ ID NO: 31): DLWGWYFDY

3G9-2D2,  $V_L$  nucleic acid sequence (VK3, locus L6; JK4) (SEQ ID NO: 32):

35 atgggatggagctgtatcatctgttcctcggtggccacagcaaccgggtccactccgaaattgtgttgcacacagtctccagcca  
 ccctgtcttgcctccaggggaaagagccaccctctctgcaggcccagtcagatgttagcagacttagcctggtaccaac  
 40 agaaacctggccaggctccaggctctatgtatgcataccaaacagggccactggcatccagccagggttcagtgccagt  
 gggctggacagacttcacttcaccatcagcagcctagagcctgaagatttgcaattactgtcagcagcgtcgcaact  
 gcccgtcacttcggcggaggaccaggtggagatcaaacgtacg

45 3G9-2D2,  $V_L$  amino acid sequence (SEQ ID NO: 33) including signal peptide:

MEAPAQLLFLLLLWLPDTTGEIVLTQSPATLSLSPGERATLSCRASQSVSSYLA  
 50 YQQKPGQAPRLLIYDASNRATGIPARFSGSGSTDFLTISLLEPEDFAVYYCQQ  
RRNWPLTFGGGTKVEIK

3G9-2D2,  $V_L$  "mature" amino acid sequence (SEQ ID NO: 34) excluding signal peptide:

55 EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYYQQKPGQAPRLLIYDASNRA  
TGIPARFSGSGSTDFLTISLLEPEDFAVYYCQQRNWPLTFGGGTKVEIK

3G9-2D2, V<sub>L</sub> CDR1 (SEQ ID NO: 35): RASQSVSSYLA

3G9-2D2, V<sub>L</sub> CDR2 (SEQ ID NO: 36): DASNRAT

5 3G9-2D2, V<sub>L</sub> CDR3 (SEQ ID NO: 37): QQRRNWPLT

5A8-1F1, V<sub>H</sub> nucleic acid sequence (VH3, locus 3-33; JH2) (SEQ ID NO: 38):

10 atggagtttgggctgacctgggtttccctcggtgccttttaagaggtgtccagtgtcaggtgcagctgggtggagtcggggagg  
cgtggtccagcctgggaggccctgagactctccgtgcagcgtctggattcaccttcagtagcatggcatgcactgggtccg  
ccaggctccaggcaaggggctggagtggatggcaattatatggatgatggaggttaataactatgcagactccgtgaagg  
15 gccgattcaccatctccagagacaattccaagaacacgcgttatctgcaaataatgaacagcctgagagccgaggacacggctgtg  
tattactgtgcgagagacttctactggtacttcgtatctgggcccgtggcaccctggactgtctccctcagcctccaccaagg  
20 cccatcggtctccctggcaagg

5A8-1F1, V<sub>H</sub> amino acid sequence (SEQ ID NO: 39) including signal peptide:

25 MEFGLTWVFLVALLRGVQCQVQLVESGGVVQPGRLRLSCAASGFTFSTYGM  
HWVRQAPGKGLEWVAIIWYDGGNKYYADSVKGRFTISRDNSKNTLYLQMNSL.  
RAEDTAVYYCARDFYWYFDLWGRGTLVTVSSASTKGPSVFPLA

30 5A8-1F1, V<sub>H</sub> "mature" amino acid sequence (SEQ ID NO: 40) excluding signal peptide:

35 QVQLVESGGVVQPGRLRLSCAASGFTFSTYGMHWVRQAPGKGLEWVAIIW  
YDGGNKYYADSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCARDFYWYF  
DLWGRGTLVTVSSASTKGPSVFPLA

5A8-1F1, V<sub>H</sub> CDR1 (SEQ ID NO: 41): TYGMH

40 5A8-1F1, V<sub>H</sub> CDR2 (SEQ ID NO: 42): IIWYDGGNKYYADSVKG

5A8-1F1, V<sub>H</sub> CDR3 (SEQ ID NO: 43): DFYWYFDL

5A8-1F1, V<sub>L</sub> nucleic acid sequence (VK3, locus L6; JK1) (SEQ ID NO: 44):

45 atggaagccccagctcagttctttccctgtactctggctccagataccaccggagaaattgtgttgcacacagtctccagc  
caccctgtttgtctccaggggaaagagaccaccctctctgcaggccagtcagagtgttagcagctacttagcctggatcca  
50 acagaaacctggccaggctccaggctccatctatgtatgcattccaaacaggccactggcatcccagccagggttcagtgca  
gtgggtctggacagacttcacttcaccatcagcagccatggctgaagatttgcagtttattactgtcagcagcgtggac  
gttcggccaaggggaccaagggtggaaatcaaacga

55

5A8-1F1, V<sub>L</sub> amino acid sequence (SEQ ID NO: 45) including signal peptide:

5 MEAPAQLLFLLLLWLPDTTGEIVLTQSPATLSLSPGERATLSCRASQSVSSYLA  
 YQQKPGQAPRLLIYD**ASNRAT**GIPARFSGSGSGTDFTLTISSL**EPEDFAVYYCQQ**  
**RRTFGQGTKVEIK**

5A8-1F1, V<sub>L</sub> "mature" amino acid sequence (SEQ ID NO: 46) excluding signal peptide:

10 EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDASNRAT  
**TGIPARFSGSGSGTDFTLTISSL**EPEDFAVYYCQQRRTFGQGTKVEIK****

15 5A8-1F1, V<sub>L</sub> CDR1 (SEQ ID NO: 47): **RASQSVSSYLA**

5A8-1F1, V<sub>L</sub> CDR2 (SEQ ID NO: 48): **DASNRAT**

5A8-1F1, V<sub>L</sub> CDR3 (SEQ ID NO: 49): **QQRRT**

20 3C7-3A3, V<sub>H</sub> nucleic acid sequence (VH3, locus 3-33; JH2) (SEQ ID NO: 50):

25 atggagttgggctgagctgggtttcctcggtgcatttaagagggtccagtgtcaggcagctggtagtctggggag  
 gcgtggtccagccctgggagggtccctgagacttcctgtcagcgtctggattcacccctcagtagctataacatgcactgggtcc  
 gccaggctccaggcaaggggctggagtggtggcatttatatgtatgtatggaaagtaataactatggagactccgtgaag  
 ggccgattcaccatctccagagacaattccaaaacacgcgttatctgcaaatacgtatggaaacagcctgagagccgaggacacggctgt  
 30 gtattactgtcgagagaagagctgggatcgggtggacttcgatctctggggccgtggaccctggactgtctccctcagc  
 ctccaccaaggggccatcggtctccccctggcac

3C7-3A3, V<sub>H</sub> amino acid sequence (SEQ ID NO: 51) including signal peptide:

35 MEFGLSWVFLVALLRGVQCQVQLVESGGVVQPGRLRLSCAASGFTFSSYNM  
**HWVRQAPGKGLEWVAFI**WYDGSNKYYGDSVKGRFTISRDNSKNTLYLQMNSL  
 RAEDTAVYYCAREEL**GIGWYFDL**WGRGTLVTVSSASTKGPSVFPLA

40 3C7-3A3, V<sub>H</sub> "mature" amino acid sequence (SEQ ID NO: 52) excluding signal peptide:

45 QVQLVESGGVVQPGRLRLSCAASGFTFSSYNMHWVRQAPGKGLEWVAFI  
**WYDGSNKYYGDSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCAREELGIGW**  
**YFDL**WGRGTLVTVSSASTKGPSVFPLA

50 3C7-3A3, V<sub>H</sub> CDR1 (SEQ ID NO: 53): **SYNMH**

3C7-3A3, V<sub>H</sub> CDR2 (SEQ ID NO: 54): **F**WYDGSNKYYGDSVKG

3C7-3A3, V<sub>H</sub> CDR3 (SEQ ID NO: 55): **EELGIGWYFDL**

55 3C7-3A3, V<sub>L</sub> nucleic acid sequence (VK3, locus L6; JK1) (SEQ ID NO: 56):

5 atggaagccccagctcagttcttcttcctgactctggctccagataccaccggagaaattgtgttgacacagtctccagc  
 caccctgtcttgcagggaaagagaccaccctctctgcagggccagtcagagttagcagctacttagcctggatcca  
 10 acagaaacctggccaggctccaggctccatctatcatccaaacaggccactggcatcccagccagggtcagtgcc  
 gtgggtctgggacagacttcacttcaccatcagcagcctagagcctgaagatttgcagtttattactgtcagcagcgtaggac  
 gttggccaagggaccaagggtggaaatcaaacgaactgtggctgaccatctgtctcatctccgcatctgtatgagcagtt  
 gaaatctggaactgcctctgttgtgcctgc

3C7-3A3,  $V_L$  amino acid sequence (SEQ ID NO: 57) including signal peptide:

15 MEAPAQLLFLLLLWLPDTTGEIVLTQSPATLSLSPGERATLSCRASQSVSSYLA  
 YQQKPGQAPRLLIYDASNRATGIPARFSGSGSGTDFTLTISLEPEDFAVYYCQQ  
 20 RRTFGQGTKVEIK

3C7-3A3,  $V_L$  "mature" amino acid sequence (SEQ ID NO: 58) excluding signal peptide:

25 EIVLTQSPATLSLSPGERATLSCRASQSVSSYLA  
 YQQKPGQAPRLLIYDASNRATGIPARFSGSGSGTDFTLTISLEPEDFAVYYCQQRRTFGQGTKVEIK

3C7-3A3,  $V_L$  CDR1 (SEQ ID NO: 59): RASQSVSSYLA

30 3C7-3A3,  $V_L$  CDR2 (SEQ ID NO: 60): DASN RAT

3C7-3A3,  $V_L$  CDR3 (SEQ ID NO: 61): QQR RT

2D3-1F5-2A9,  $V_H$  nucleic acid sequence (VH3, locus Orph-C16; JH3) (SEQ ID NO: 62):

35 atggagtttgtgctgagctgggttccttgtctatattaaaagggtgtccagtgtaggttcagctggcagtcggggagg  
 ctggtagatccctgggggtccctgagacttcctgtgcaggctctggattcacccatcactatgtatgcactgggtcgcc  
 40 aggctccaggaaaagggtctggatgggtatcaactattgtactgggtggcacaccctatgcagactccgtgaaggggccgc  
 ttcaccatctccagagacaatgccaagaactcctgtatctcaaatgaacagcctgagagccgaggacatggctgttattact  
 gtgcatttaagtgtttgtctggggcaaggacatggtcaccgtcttcagccctccaccaagggccatggcttccc  
 45 cctggcac

2D3-1F5-2A9,  $V_H$  amino acid sequence (SEQ ID NO: 63) including signal peptide:

50 MEFVLSWVLLVAILKGVQCEVQLVQSGGLVHPGGSLRLSCAGSGFTFSNYAM  
 HWVRQAPGKGLEWVSTIGGGTPYADSVKGRFTISRDNAKNSLYLQMNSLRA  
 EDMAVYYCALSAFDVWGQGTMVTVSSASTKGPSVFPLA

55 2D3-1F5-2A9,  $V_H$  "mature" amino acid sequence (SEQ ID NO: 64) excluding signal peptide:

EVQLVQSGGLVHPGSLRLSCAGSGFTFSNYAMHWVRQAPGKGLEWVSTIGT  
GGGTPYADSVKGRFTISRDNAKNSLYLQMNSLRAEDMAVYYCALSAFDVWGQ  
5 GTMVTVSSASTKGPSVFPLA

2D3-1F5-2A9, V<sub>H</sub>CDR1 (SEQ ID NO: 65): NYAMH

2D3-1F5-2A9, V<sub>H</sub>CDR2 (SEQ ID NO: 66): TIGTGGGTPYADSVKG

10 2D3-1F5-2A9, V<sub>H</sub>CDR3 (SEQ ID NO: 67): SAFDV

1E6-3D10 V<sub>H</sub> nucleic acid sequence (VH3, locus Orph-HC16; JH4) (SEQ ID NO: 68):

15 Atggagtttgtctgagctgggtttcccttgtctgtatattaaaagggtccagtgtgagggtcagctgggtcagtcggggagg  
cttggtacatcctgggggtccctgagactctcctgtcaggctctggattcacccatcagtagctatgctatgcactgggtcgcc  
20 aggctccaggaaaagggtctggagtggtatcagctattggtaactgggtttacacatactatgttagactccgtgaagggccgatt  
caccatctccagagacaatgccaagaagtccctgtatctcaaataatgaacagcctgagagccgaggacatggctgttattactgt  
gcaagagagccgtttacgatatttgactggtaattcccccatactttgactactgggccaggaaacctggtcaccgtctccca  
25 gcctccaccaaggggccatcggtctccccctggcac

1E6-3D10 V<sub>H</sub> amino acid sequence (SEQ ID NO: 69) including signal peptide:

30 MEFVLSWVFLVAILKGVQCEVQLVQSGGLVHPGSLRLSCAGSGFTSSYAM  
HWVRQAPGKGLEWVSAIGTGGYTYVDSVKGRFTISRDNAKKSLYLQMNSLR  
AEDMAVYYCAREPFYDILTGYSPYFDYWGQTLTVSS

35 1E6-3D10 V<sub>H</sub> "mature" amino acid sequence (SEQ ID NO: 70) excluding signal peptide:

40 EVQLVQSGGLVHPGSLRLSCAGSGFTSSYAMWVRQAPGKGLEWVSAIGT  
GGYTYVDSVKGRFTISRDNAKKSLYLQMNSLRAEDMAVYYCAREPFYDILTG  
YSPYFDYWGQTLTVSS

45 1E6-3D10 V<sub>H</sub>CDR1: (SEQ ID NO: 71):SYAMH

1E6-3D10 V<sub>H</sub>CDR2 (SEQ ID NO: 72): AIGTGGYTYVDSVKG

45 1E6-3D10 V<sub>H</sub>CDR3 (SEQ ID NO: 73): EPFYDILTGYSPYFDY

50 5C3-2-3F6 V<sub>H</sub> nucleic acid sequence (VH3, locus 3-33; JH2) (SEQ ID NO: 74):

Atggagttgggctgagctgggtttcctcgltgcttttaagaggtgtccagtgtaggtgcagctggggag  
 5 gcgtggtccagcctgggaggccctgagactctcctgtcagcgtctggattcacccctcagtagctataacatgcactgggtcc  
 gccaggctccaggcaaggggctggagtggtggcagttatatggatggaagtaataactatggagactccgtgaag  
 10 ggccgattcaccatctccagagacaattccaagaacacgctgtatctgcaaatacgaacagcctgagagccgaggacacggctgt  
 gtattactgtcgagagaagagctggggatcgggtgtacttcgatctggggccgtggcaccctggtaactgtctcctcagc  
 ctccaccaagggccatcggtttccccctggcac

5C3-2-3F6 V<sub>H</sub> amino acid sequence (SEQ ID NO: 75) including signal peptide:

15 MEGFLSWVFLVALLRGVQCQVQLVESGGGVQPGRSRLSCAASGFTSSYNM  
HWVRQAPGKGLEWVAVIWYDGSNKYYGDSVKGRFTISRDNSKNTLYLQMNSL  
RAEDTAVYYCAREELGIGWYFDLWGRGTLTVSS

20 5C3-2-3F6 V<sub>H</sub> "mature" amino acid sequence (SEQ ID NO: 76) excluding signal peptide:

25 QVQLVESGGGVQPGRSRLSCAASGFTSSYNMHWVRQAPGKGLEWVAVIW  
YDGSNKYYGDSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCAREELGIGW  
YFDLWGRGTLTVSS

30 5C3-2-3F6 V<sub>H</sub>CDR1 (SEQ ID NO: 77): SYNMH

35 5C3-2-3F6 V<sub>H</sub>CDR2 (SEQ ID NO: 78): VIWYDGSNKYYGDSVKG

50 5C3-2-3F6 V<sub>H</sub>CDR3 (SEQ ID NO: 79): EELGIGWYFDL

5C3-2-3F6 VK V<sub>L</sub> nucleic acid sequence (VK1, locus L18; JK5) (SEQ ID NO: 80):

40 Atggacatgagggccccgctcagctccctggggcttcgtgtctggctccagggtgcacatccagggtggaccca  
 gtctccatcctccctgtctgcacatctgttaggagacagactcaccatcacttgcggggcaagtcaggcattagcagtgccttagcc  
 tggatcagcagaaccaggaaagctctaagctccatgtatgcctccagggtggaaagtgggtccatcaagggttc  
 45 agcggcagtggatctggacagatttcacttcaccatcagcgcctgcagcctgaagatttgcaacttattactgtcaacagtt  
 taatagttaccctcacttcggccaaggacacgactggagattaaacgaactgtggctgcaccatctgtttcatctcccgccat  
 ctgatgagcagttgaaatctggaaactgcctgtgtgcctgcaaggc

50 5C3-2-3F6 VK V<sub>L</sub> amino acid sequence (SEQ ID NO: 81) including signal peptide:

MDMRVPAQLLGLLLWLPGARCAIQLTQSPSSLASAVGDRVTITCRASQGISSAL  
AWYQQKPGKAPKLIYDASSLESGVPSRFSGSGTDFLTISLQPEDFATYYC  
QQFNSYPHFGQGTRLEIK

5C3-2-3F6 VK V<sub>L</sub> "mature" amino acid sequence (SEQ ID NO: 82) excluding signal peptide:

AIQLTQSPSSLSASVGDRVТИCRASQGISSALAWYQQKPGKAPKLLIYDASSLES  
GVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQFNSYPHFGQGTRLEIK

5 5C3-2-3F6 V<sub>L</sub>CDR1 (SEQ ID NO: 83): RASQGISSALA

5C3-2-3F6 V<sub>L</sub>CDR2 (SEQ ID NO: 84): DASSLES

10 5C3-2-3F6 V<sub>L</sub>CDR3 (SEQ ID NO: 85): QQFNSYPH

5D12-5G1 VH nucleic acid sequence (VH3, locus 3-33; JH2) (SEQ ID NO: 86):

15 Atggagttgggctgagctgggtttcctcggtgccttttaagaggtgtccagtgccaggcagctgggtggagctggggag  
gcgtggccaggcctgggagggtccctgagactcctgtgcagcgtctggattcaccttcagtagctatggcatgcactgggtcc  
gccaggcaggcaaggggctggagtggtggcagttatatggatgatggaagtaataactatgcagactccgtgaag  
20 gcccatttcaccatctccagagacaattccaagaacacgcgtatctgcattgcacagcctgagagccgaggacacggctgt  
gtattactgtgcgagaggccccctcggtacttcgatctctggggccgtggcaccctggtaactgtctcctcagccctccaccaa  
25 gggccatcggtcttccctggcac

5D12-5G1 VH amino acid sequence (SEQ ID NO: 87) including signal peptide:

30 MEFGLSWVFLVALLRGVQCQLVESGGVVQPGRSRLSCAASGFTSSYGM  
HWVRQAPGKGLEWVAVIWYDGSNKYYADSVKGRFTISRDNSKNTLYLQMNSL  
RAEDTAVYYCARGPPRYFDLWGRGTLTVSS

35 5D12-5G1 VH "mature" amino acid sequence (SEQ ID NO: 88) excluding signal peptide:

40 QVQLVESGGVVQPGRSRLSCAASGFTSSYGMHWVRQAPGKGLEWVAVIW  
YDGSNKYYADSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCARGPPRYFD  
LWGRGTLTVSS

5D12-5G1 VH CDR1 (SEQ ID NO: 89): SYGMH

45 5D12-5G1 VH CDR2 (SEQ ID NO: 90): VIWYDGSNKYYADSVKG

5D12-5G1 VH CDR3 (SEQ ID NO: 91): GPPRYFDL

50 [0205] For reference, the amino acid sequences of the proposed corresponding germline sequences (assigned without prejudice) are as follows:

Germline L6 (SEQ ID NO: 92):

55 EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDASNRA  
TGIPARFSGSGSGTDFTLTISSLEPEDFAVYYCQQRSNWP

Germline L4 (SEQ ID NO: 93):

AIQLTQSPSSLSASVGDRVITCRASQGISSALAWYQQKPGKAPKLLIYDASSLES  
 5  
 GVPSRFSGSGSGTDFLTISLQPEDFATYYCQQFNSYP

Germline L15 (SEQ ID NO: 94):

DIQMTQSPSSLSASVGDRVITCRARQGISSWLAWYQQKPEKAPKSLIYAASSLQ  
 10  
 SGVPSRFSGSGSGTDFLTISLQPEDFATYYCQQYNSYP

Germline V<sub>H</sub>3-33 (SEQ ID NO: 95):

QVQLVESGGVVQPGRSRLSCAASGFTFSSYGMHWVRQAPGKGLEWVAVIW  
 20  
 YDGSNKYYADSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCAR

Germline Orph-C16 (SEQ ID NO: 96):

EVQLVQSGGGLVHPGGSLRLSCAGSGFTFSSYAMHWVRQAPGKGLEWV  
 25  
 SAI GTGGGTYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDMAVYYCAR

30 [0206] Sequence alignments of the V<sub>L</sub> and V<sub>H</sub> sequences against the proposed corresponding germline sequences are shown in Figure 5, for illustration purposes only.

#### Example 8

##### 3G9- $\beta$ hCG APC-targeted vaccine conjugate

35 [0207] A DEC-205 targeted vaccine conjugate was generated by linking the  $\beta$ hCG antigen to HuFab 3G9-2D2 (also determined to be cross-reactive with cynomolgous DEC-205) from Example 7 above. Linkage was accomplished by covalently attaching the antigen to the heavy chain of the antibody by genetic fusion.

40 [0208] A plasmid containing neomycin and dihydrofolate reductase genes was generated containing the  $\beta$ hCG coding sequence fused to antibody 3G9-2D2 heavy chain at the CH3 domain and the 3G9-2D2 light chain. The resulting plasmid construct was transfected into CHO cells using a standardized protocol (Qiagen Inc, Valencia, CA). Transfected cells were selected in media containing the antibiotic G418. After selection, the cells were cloned by limiting dilution, and stable clonal lines were used to generate cell banks for further studies. To confirm expression of the 3G9- $\beta$ hCG constructs, Western Blot analysis of proteins run on SDS-PAGE under reducing and non-reducing conditions was performed. This fusion protein was observed to be of the expected molecular weight and to be properly assembled (*i.e.*, to contain both the heavy chain fusion and the light chain). Specifically, the vaccine conjugate and the antibody alone were analyzed by SDS-PAGE using denaturing conditions and detected by Western blot analysis. The blot was then probed separately using goat anti-human IgG, and with a mAb (US Biologicals) specific to the  $\beta$ hCG C-terminal peptide. The results confirmed that the transformed CHO cells specifically expressed the 3G9- $\beta$ hCG vaccine conjugate as evidenced by the appropriate size and composition of the fusion product.

#### Example 9

##### Antigen-specific activity using 3G9- $\beta$ hCG APC-targeted vaccine conjugate

55 [0209] Cells capable of antigen presentation were human in origin and varied from peripheral blood mononuclear cells (PBMC), monocytes (THP-1), B lymphoblastoid cells (C1R.A2, 1518 B-LCL) and monocyte-derived DCs. All cells were positive for cell surface expression of DEC-205 as assessed by flow cytometry.

[0210] The vector pk: 3G9-hCG $\beta$  was transfected into CHO cells. Stable clones were selected with G418 and subsequently subcloned. The fusion protein produced by the cells (3G9- $\beta$ hCG vaccine conjugate; Example 8) was collected in the supernatant and purified over Protein A column.

[0211] T cells were obtained from leukopacks of normal healthy donors. Antigen-specific T cells were generated in vitro by 2-3 weekly stimulations with autologous DCs targeted with 3G9-hCG $\beta$  and enriched for CD8+ and CD4+ T cells before testing for antigen-specific activity with a variety of APCs (as described above) by GrB or IFNy ELISpot assays (MabTech). Cytokines IL-7 and IL-2 were added to maintain effector propagation and activity every 3-4 days. Antigen-specific T cells were expanded on Miltenyi-MACS T cell expansion kit for 10-12 days in the presence of low dose of IL-2. CD40L (Alexis Biochemicals) was used to induce maturation of DCs. As shown in Figure 9A, CD8+ T cell responses were achieved in DCs and monocytes (THP-1), as well as B lymphoblastoid cells (Figure 9B). Accordingly, antigen targeting via the DEC-205 receptor to B cells resulted in the stimulation of MHC-class I restricted T cells.

## SEQUENCE LISTING

15 [0212]

<110> Celldex Therapeutics Inc., et al.

<120> ANTIBODIES THAT BIND HUMAN DENDRITIC AND EPITHELIAL CELL 205 (DEC-205)

20 <130> CDJ-346PC

<140> Not Yet Assigned

<141> Concurrently Herewith

25 <150> 61/002,253  
<151> 2007-11-07

30 <150> 61/191,551  
<151> 2008-09-10

<160> 106

35 <170> PatentIn version 3.5

<210> 1  
<211> 1722  
<212> PRT  
<213> Homo sapiens

40 <400> 1

45

50

55

## EP 2 224 954 B1

Met Arg Thr Gly Trp Ala Thr Pro Arg Arg Pro Ala Gly Leu Leu Met  
1 5 . 10 15

5 Leu Leu Phe Trp Phe Phe Asp Leu Ala Glu Pro Ser Gly Arg Ala Ala  
20 25 30

10 Asn Asp Pro Phe Thr Ile Val His Gly Asn Thr Gly Lys Cys Ile Lys  
35 40 45

15 Pro Val Tyr Gly Trp Ile Val Ala Asp Asp Cys Asp Glu Thr Glu Asp  
50 55 60

20 Lys Leu Trp Lys Trp Val Ser Gln His Arg Leu Phe His Leu His Ser  
65 70 75 80

25 Gln Lys Cys Leu Gly Leu Asp Ile Thr Lys Ser Val Asn Glu Leu Arg  
85 90 95

30 Met Phe Ser Cys Asp Ser Ser Ala Met Leu Trp Trp Lys Cys Glu His  
100 105 110

His Ser Leu Tyr Gly Ala Ala Arg Tyr Arg Leu Ala Leu Lys Asp Gly  
115 120 125

35 His Gly Thr Ala Ile Ser Asn Ala Ser Asp Val Trp Lys Lys Gly Gly

40

45

50

55

## EP 2 224 954 B1

	130	135	140
5	Ser Glu Glu Ser Leu Cys Asp Gln Pro Tyr His Glu Ile Tyr Thr Arg		
	145	150	155
	160		
	Asp Gly Asn Ser Tyr Gly Arg Pro Cys Glu Phe Pro Phe Leu Ile Asp		
	165	170	175
10	Gly Thr Trp His His Asp Cys Ile Leu Asp Glu Asp His Ser Gly Pro		
	180	185	190
15	Trp Cys Ala Thr Thr Leu Asn Tyr Glu Tyr Asp Arg Lys Trp Gly Ile		
	195	200	205
20	Cys Leu Lys Pro Glu Asn Gly Cys Glu Asp Asn Trp Glu Lys Asn Glu		
	210	215	220
	Gln Phe Gly Ser Cys Tyr Gln Phe Asn Thr Gln Thr Ala Leu Ser Trp		
	225	230	235
	240		
25	Lys Glu Ala Tyr Val Ser Cys Gln Asn Gln Gly Ala Asp Leu Leu Ser		
	245	250	255
30	Ile Asn Ser Ala Ala Glu Leu Thr Tyr Leu Lys Glu Lys Gly Ile		
	260	265	270
	Ala Lys Ile Phe Trp Ile Gly Leu Asn Gln Leu Tyr Ser Ala Arg Gly		
35	275	280	285
	Trp Glu Trp Ser Asp His Lys Pro Leu Asn Phe Leu Asn Trp Asp Pro		
	290	295	300
40	Asp Arg Pro Ser Ala Pro Thr Ile Gly Gly Ser Ser Cys Ala Arg Met		
	305	310	315
	320		
45	Asp Ala Glu Ser Gly Leu Trp Gln Ser Phe Ser Cys Glu Ala Gln Leu		
	325	330	335
	Pro Tyr Val Cys Arg Lys Pro Leu Asn Asn Thr Val Glu Leu Thr Asp		
50	340	345	350
	Val Trp Thr Tyr Ser Asp Thr Arg Cys Asp Ala Gly Trp Leu Pro Asn		
	355	360	365
55	Asn Gly Phe Cys Tyr Leu Leu Val Asn Glu Ser Asn Ser Trp Asp Lys		
	370	375	380

## EP 2 224 954 B1

Ala His Ala Lys Cys Lys Ala Phe Ser Ser Asp Leu Ile Ser Ile His  
 385 390 395 400

5 Ser Leu Ala Asp Val Glu Val Val Val Thr Lys Leu His Asn Glu Asp  
 405 410 415

10 Ile Lys Glu Glu Val Trp Ile Gly Leu Lys Asn Ile Asn Ile Pro Thr  
 420 425 430

15 Leu Phe Gln Trp Ser Asp Gly Thr Glu Val Thr Leu Thr Tyr Trp Asp  
 435 440 445

20 Glu Asn Glu Pro Asn Val Pro Tyr Asn Lys Thr Pro Asn Cys Val Ser  
 450 455 460

25 Tyr Leu Gly Glu Leu Gly Gln Trp Lys Val Gln Ser Cys Glu Glu Lys  
 465 470 475 480

30 Leu Lys Tyr Val Cys Lys Arg Lys Gly Glu Lys Leu Asn Asp Ala Ser  
 485 490 495

Ser Asp Lys Met Cys Pro Pro Asp Glu Gly Trp Lys Arg His Gly Glu  
 500 505 510

35 Thr Cys Tyr Lys Ile Tyr Glu Asp Glu Val Pro Phe Gly Thr Asn Cys  
 515 520 525

Asn Leu Thr Ile Thr Ser Arg Phe Glu Gln Glu Tyr Leu Asn Asp Leu  
 530 535 540

40 Met Lys Lys Tyr Asp Lys Ser Leu Arg Lys Tyr Phe Trp Thr Gly Leu  
 545 550 555 560

Arg Asp Val Asp Ser Cys Gly Glu Tyr Asn Trp Ala Thr Val Gly Gly  
 565 570 575

45 Arg Arg Arg Ala Val Thr Phe Ser Asn Trp Asn Phe Leu Glu Pro Ala  
 580 585 590

50 Ser Pro Gly Gly Cys Val Ala Met Ser Thr Gly Lys Ser Val Gly Lys  
 595 600 605

55 Trp Glu Val Lys Asp Cys Arg Ser Phe Lys Ala Leu Ser Ile Cys Lys  
 610 615 620

## EP 2 224 954 B1

Lys Met Ser Gly Pro Leu Gly Pro Glu Glu Ala Ser Pro Lys Pro Asp  
 625 630 635 640  
  
 5 Asp Pro Cys Pro Glu Gly Trp Gln Ser Phe Pro Ala Ser Leu Ser Cys  
 645 650 655  
  
 10 Tyr Lys Val Phe His Ala Glu Arg Ile Val Arg Lys Arg Asn Trp Glu  
 660 665 670  
  
 15 Glu Ala Glu Arg Phe Cys Gln Ala Leu Gly Ala His Leu Ser Ser Phe  
 675 680 685  
  
 20 Ser His Val Asp Glu Ile Lys Glu Phe Leu His Phe Leu Thr Asp Gln  
 690 695 700  
  
 25 Phe Ser Gly Gln His Trp Leu Trp Ile Gly Leu Asn Lys Arg Ser Pro  
 705 710 715 720  
  
 30 Asp Leu Gln Gly Ser Trp Gln Trp Ser Asp Arg Thr Pro Val Ser Thr  
 725 730 735  
  
 Ile Ile Met Pro Asn Glu Phe Gln Gln Asp Tyr Asp Ile Arg Asp Cys  
 740 745 750  
  
 35 Ala Ala Val Lys Val Phe His Arg Pro Trp Arg Arg Gly Trp His Phe  
 755 760 765  
  
 Tyr Asp Asp Arg Glu Phe Ile Tyr Leu Arg Pro Phe Ala Cys Asp Thr  
 770 775 780  
  
 40 Lys Leu Glu Trp Val Cys Gln Ile Pro Lys Gly Arg Thr Pro Lys Thr  
 785 790 795 800  
  
 45 Pro Asp Trp Tyr Asn Pro Asp Arg Ala Gly Ile His Gly Pro Pro Leu  
 805 810 815  
  
 50 Ile Ile Glu Gly Ser Glu Tyr Trp Phe Val Ala Asp Leu His Leu Asn  
 820 825 830  
  
 Tyr Glu Glu Ala Val Leu Tyr Cys Ala Ser Asn His Ser Phe Leu Ala  
 835 840 845  
  
 55 Thr Ile Thr Ser Phe Val Gly Leu Lys Ala Ile Lys Asn Lys Ile Ala  
 850 855 860

## EP 2 224 954 B1

Asn	Ile	Ser	Gly	Asp	Gly	Gln	Lys	Trp	Trp	Ile	Arg	Ile	Ser	Glu	Trp	
865					870					875				880		
5 Pro Ile Asp Asp His Phe Thr Tyr Ser Arg Tyr Pro Trp His Arg Phe																
					885					890				895		
10 Pro Val Thr Phe Gly Glu Glu Cys Leu Tyr Met Ser Ala Lys Thr Trp																
					900					905				910		
15 Leu Ile Asp Leu Gly Lys Pro Thr Asp Cys Ser Thr Lys Leu Pro Phe																
					915					920				925		
20 Ile Cys Glu Lys Tyr Asn Val Ser Ser Leu Glu Lys Tyr Ser Pro Asp																
					930					935				940		
25 Ser Ala Ala Lys Val Gln Cys Ser Glu Gln Trp Ile Pro Phe Gln Asn																
					945					950				955		960
30 Lys Cys Phe Leu Lys Ile Lys Pro Val Ser Leu Thr Phe Ser Gln Ala																
					965					970				975		
35 Ser Asp Thr Cys His Ser Tyr Gly Gly Thr Leu Pro Ser Val Leu Ser																
					980					985				990		
40 Gln Ile Glu Gln Asp Phe Ile Thr Ser Leu Leu Pro Asp Met Glu Ala																
					995					1000				1005		
45 Thr Leu Trp Ile Gly Leu Arg Trp Thr Ala Tyr Glu Lys Ile Asn																
					1010					1015				1020		
50 Lys Trp Thr Asp Asn Arg Glu Leu Thr Tyr Ser Asn Phe His Pro																
					1025					1030				1035		
55 Leu Leu Val Ser Gly Arg Leu Arg Ile Pro Glu Asn Phe Phe Glu																
					1040					1045				1050		
60 Glu Glu Ser Arg Tyr His Cys Ala Leu Ile Leu Asn Leu Gln Lys																
					1055					1060				1065		
65 Ser Pro Phe Thr Gly Thr Trp Asn Phe Thr Ser Cys Ser Glu Arg																
					1070					1075				1080		
70 His Phe Val Ser Leu Cys Gln Lys Tyr Ser Glu Val Lys Ser Arg																
					1085					1090				1095		
75 Gln Thr Leu Gln Asn Ala Ser Glu Thr Val Lys Tyr Leu Asn Asn																

## EP 2 224 954 B1

	1100	1105	1110
5	Leu Tyr Lys Ile Ile Pro Lys Thr Leu Thr Trp His Ser Ala Lys		
	1115	1120	1125
10	Arg Glu Cys Leu Lys Ser Asn Met Gln Leu Val Ser Ile Thr Asp		
	1130	1135	1140
15	Pro Tyr Gln Gln Ala Phe Leu Ser Val Gln Ala Leu Leu His Asn		
	1145	1150	1155
20	Ser Ser Leu Trp Ile Gly Leu Phe Ser Gln Asp Asp Glu Leu Asn		
	1160	1165	1170
25	Phe Gly Trp Ser Asp Gly Lys Arg Leu His Phe Ser Arg Trp Ala		
	1175	1180	1185
30	Glu Thr Asn Gly Gln Leu Glu Asp Cys Val Val Leu Asp Thr Asp		
	1190	1195	1200
35	Gly Phe Trp Lys Thr Val Asp Cys Asn Asp Asn Gln Pro Gly Ala		
	1205	1210	1215
40	Ile Cys Tyr Tyr Ser Gly Asn Glu Thr Glu Lys Glu Val Lys Pro		
	1220	1225	1230
45	Val Asp Ser Val Lys Cys Pro Ser Pro Val Leu Asn Thr Pro Trp		
	1235	1240	1245
50	Ile Pro Phe Gln Asn Cys Cys Tyr Asn Phe Ile Ile Thr Lys Asn		
	1250	1255	1260
55	Arg His Met Ala Thr Thr Gln Asp Glu Val His Thr Lys Cys Gln		
	1265	1270	1275
60	Lys Leu Asn Pro Lys Ser His Ile Leu Ser Ile Arg Asp Glu Lys		
	1280	1285	1290
65	Glu Asn Asn Phe Val Leu Glu Gln Leu Leu Tyr Phe Asn Tyr Met		
	1295	1300	1305
70	Ala Ser Trp Val Met Leu Gly Ile Thr Tyr Arg Asn Asn Ser Leu		
	1310	1315	1320
75	Met Trp Phe Asp Lys Thr Pro Leu Ser Tyr Thr His Trp Arg Ala		
	1325	1330	1335

## EP 2 224 954 B1

Gly Arg Pro Thr Ile Lys Asn Glu Lys Phe Leu Ala Gly Leu Ser  
 1340 1345 1350

5 Thr Asp Gly Phe Trp Asp Ile Gln Thr Phe Lys Val Ile Glu Glu  
 1355 1360 1365

10 Ala Val Tyr Phe His Gln His Ser Ile Leu Ala Cys Lys Ile Glu  
 1370 1375 1380

15 Met Val Asp Tyr Lys Glu Glu His Asn Thr Thr Leu Pro Gln Phe  
 1385 1390 1395

20 Met Pro Tyr Glu Asp Gly Ile Tyr Ser Val Ile Gln Lys Lys Val  
 1400 1405 1410

25 Thr Trp Tyr Glu Ala Leu Asn Met Cys Ser Gln Ser Gly Gly His  
 1415 1420 1425

30 Leu Ala Ser Val His Asn Gln Asn Gly Gln Leu Phe Leu Glu Asp  
 1430 1435 1440

35 Ile Val Lys Arg Asp Gly Phe Pro Leu Trp Val Gly Leu Ser Ser  
 1445 1450 1455

40 His Asp Gly Ser Glu Ser Ser Phe Glu Trp Ser Asp Gly Ser Thr  
 1460 1465 1470

45 Phe Asp Tyr Ile Pro Trp Lys Gly Gln Thr Ser Pro Gly Asn Cys  
 1475 1480 1485

50 Val Leu Leu Asp Pro Lys Gly Thr Trp Lys His Glu Lys Cys Asn  
 1490 1495 1500

55 Ser Val Lys Asp Gly Ala Ile Cys Tyr Lys Pro Thr Lys Ser Lys  
 1505 1510 1515

Lys Leu Ser Arg Leu Thr Tyr Ser Ser Arg Cys Pro Ala Ala Lys  
 1520 1525 1530

60 Glu Asn Gly Ser Arg Trp Ile Gln Tyr Lys Gly His Cys Tyr Lys  
 1535 1540 1545

65 Ser Asp Gln Ala Leu His Ser Phe Ser Glu Ala Lys Lys Leu Cys  
 1550 1555 1560

## EP 2 224 954 B1

Ser Lys His Asp His Ser Ala Thr Ile Val Ser Ile Lys Asp Glu  
 1565 1570 1575

5 Asp Glu Asn Lys Phe Val Ser Arg Leu Met Arg Glu Asn Asn Asn  
 1580 1585 1590

10 Ile Thr Met Arg Val Trp Leu Gly Leu Ser Gln His Ser Val Asp  
 1595 1600 1605

15 Gln Ser Trp Ser Trp Leu Asp Gly Ser Glu Val Thr Phe Val Lys  
 1610 1615 1620

20 Trp Glu Asn Lys Ser Lys Ser Gly Val Gly Arg Cys Ser Met Leu  
 1625 1630 1635

25 Ile Ala Ser Asn Glu Thr Trp Lys Lys Val Glu Cys Glu His Gly  
 1640 1645 1650

30 Phe Gly Arg Val Val Cys Lys Val Pro Leu Gly Pro Asp Tyr Thr  
 1655 1660 1665

Ala Ile Ala Ile Ile Val Ala Thr Leu Ser Ile Leu Val Leu Met  
 1670 1675 1680

Gly Gly Leu Ile Trp Phe Leu Phe Gln Arg His Arg Leu His Leu  
 1685 1690 1695

35 Ala Gly Phe Ser Ser Val Arg Tyr Ala Gln Gly Val Asn Glu Asp  
 1700 1705 1710

40 Glu Ile Met Leu Pro Ser Phe His Asp  
 1715 1720

<210> 2  
 <211> 408  
 45 <212> DNA  
 <213> Homo sapiens

<400> 2

50 atggagtttg ggctgagctg ggttttcctc gttgctcttt taagaggtgt ccagtgtcag 60  
 gtgcagctgg tggagctctgg gggaggcggt gtccagcctg ggaggtccct gagactctcc 120  
 tgtgcagcgt ctggattcat cttcagttatc tatggcatgc actgggtccg ccaggctcca 180  
 55 ggcagaaggggc tggagtggtt ggcagttata tggatgtatg gaagtaataa atactatgca 240  
 gactccgtga agggccgatt caccatctcc agagacaatt ccaagaacac gctgtatctg 300

## EP 2 224 954 B1

caaatgaaca gcctgagagc cgaggacacg gctgtgtatt actgtgcag agtcctcac 360  
 tttgactact ggggccaggg aaccctggtc accgtctcct cagctagc 408  
 5  
 <210> 3  
 <211> 134  
 <212> PRT  
 <213> Homo sapiens  
 10  
 <400> 3

Met	Glu	Phe	Gly	Leu	Ser	Trp	Val	Phe	Leu	Val	Ala	Leu	Leu	Arg	Gly
1															

15  
 Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln  
 20 25 30

Pro	Gly	Arg	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ile	Phe
35															

25  
 Ser Ile Tyr Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu  
 50 55 60

Glu	Trp	Val	Ala	Val	Ile	Trp	Tyr	Asp	Gly	Ser	Asn	Lys	Tyr	Tyr	Ala
65															

30  
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn  
 85 90 95

Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val
100															

35  
 Tyr Tyr Cys Ala Arg Ala Pro His Phe Asp Tyr Trp Gly Gln Gly Thr  
 115 120 125

Leu	Val	Thr	Val	Ser	Ser										
130															

40  
 <210> 4  
 <211> 115  
 <212> PRT  
 <213> Homo sapiens  
 45  
 <400> 4

## EP 2 224 954 B1

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg  
 1 5 10 15

5 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ile Phe Ser Ile Tyr  
 20 25 30

10 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

15 Ala Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val  
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
 65 70 75 80

20 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95

25 Ala Arg Ala Pro His Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr  
 100 105 110

30 Val Ser Ser  
 115

35 <210> 5  
 <211> 5  
 <212> PRT  
 <213> Homo sapiens

40 <400> 5

Ile Tyr Gly Met His  
 1 5

45 <210> 6  
 <211> 17  
 <212> PRT  
 <213> Homo sapiens

50 <400> 6

Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys  
 1 5 10 15

55 Gly

<210> 7  
 <211> 6  
 <212> PRT  
 <213> Homo sapiens

&lt;400&gt; 7

	Ala	Pro	His	Phe	Asp	Tyr
1					5	

5

&lt;210&gt; 8

&lt;211&gt; 384

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

10

&lt;400&gt; 8

	atgggatgga	gctgtatcat	cctgttcctc	gtggccacag	caaccggtgt	ccactccgac	60
15	atccagatga	cccagtctcc	atccctactg	tctgcatctg	ttggagacag	agtcaccatc	120
	acttgtcggg	cgagtcaggg	tattagcagc	tggtagcct	ggtatcagca	gaaaccagag	180
20	aaagcccta	agtccctgat	ctatgctgca	tccagttgc	aaagtgggt	cccatcaagg	240
	ttcagcggca	gtggatctgg	gacagatttc	actctcacca	tcagcagcct	gcagcctgaa	300
	gattttgcaa	cttattactg	ccaacagtat	aatagttacc	cgtacactt	tggccaggg	360
25	accaagctgg	agatcaaacg	tacg				384

&lt;210&gt; 9

&lt;211&gt; 129

&lt;212&gt; PRT

30

&lt;213&gt; Homo sapiens

&lt;400&gt; 9

35

40

45

50

55

## EP 2 224 954 B1

Met	Asp	Met	Arg	Val	Leu	Ala	Gln	Leu	Leu	Gly	Leu	Leu	Leu	Cys		
1				5				10					15			
5	Phe	Pro	Gly	Ala	Arg	Cys	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser
					20				25				30			
10	Leu	Ser	Ala	Ser	Val	Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser
					35				40				45			
15	Gln	Gly	Ile	Ser	Ser	Trp	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Glu	Lys
					50				55				60			
20	Ala	Pro	Lys	Ser	Leu	Ile	Tyr	Ala	Ala	Ser	Ser	Leu	Gln	Ser	Gly	Val
					65		70			75			80			
25	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr
					85				90				95			
30	Ile	Ser	Ser	Leu	Gln	Pro	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln
					100				105				110			
35	Tyr	Asn	Ser	Tyr	Pro	Tyr	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile
					115				120				125			
40	Lys															
45																
50																
55																

<210> 10  
 <211> 107  
 <212> PRT  
 <213> Homo sapiens

<400> 10

40

## EP 2 224 954 B1

	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly	
1															15		
	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Gly	Ile	Ser	Ser	Trp	
5															30		
	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Glu	Lys	Ala	Pro	Lys	Ser	Leu	Ile	
10															45		
	Tyr	Ala	Ala	Ser	Ser	Leu	Gln	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	
														50	60		
15		Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
															80		
20		Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Tyr	Asn	Ser	Tyr	Pro	Tyr
															95		
	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys						
25												100		105			
	<210> 11																
	<211> 11																
	<212> PRT																
30	<213> Homo sapiens																
	<400> 11																
		Arg	Ala	Ser	Gln	Gly	Ile	Ser	Ser	Trp	Leu	Ala					
35												1		5	10		
	<210> 12																
	<211> 7																
	<212> PRT																
40	<213> Homo sapiens																
	<400> 12																
		Ala	Ala	Ser	Ser	Leu	Gln	Ser									
45									1		5						
	<210> 13																
	<211> 9																
	<212> PRT																
50	<213> Homo sapiens																
	<400> 13																
		Gln	Gln	Tyr	Asn	Ser	Tyr	Pro	Tyr	Thr							
55											1		5				
	<210> 14																
	<211> 439																

## EP 2 224 954 B1

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 14

5

atggagtttgcg	ggctgagctgg	gttttcctc	gttgctcttta	taagagggtgt	ccagtgtcag	60
gtgcagctgg	tggagtctgg	gggaggcgtg	gtccagcctg	ggaggtccct	gagactctcc	120
10 tgcgcgt	ctggattcat	cttcagtatc	tatggcatgc	actgggtccg	ccaggctcca	180
ggcaagggc	tggagtgggt	ggcagttata	tggtatgatg	gaagtaataa	atactatgca	240
15 gactccgtga	agggccgatt	caccatctcc	agagacaatt	ccaagaacac	gctgtatctg	300
caaataaca	gcctgagagc	cgaggacacg	gctgtgtatt	actgtgcgag	agctcctcac	360
20 tttgactact	ggggccaggg	aaccctggtc	accgtctcct	cagcctccac	caagggccca	420
.	tcggtcttcc	ccctggcac				439

&lt;210&gt; 15

&lt;211&gt; 134

&lt;212&gt; PRT

25

&lt;213&gt; Homo sapiens

&lt;400&gt; 15

30

Met	Glu	Phe	Gly	Leu	Ser	Trp	Val	Phe	Leu	Val	Ala	Leu	Leu	Arg	Gly
1				5					10				15		

35

Val	Gln	Cys	Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Val	Val	Gln
				20				25					30		

40

Pro	Gly	Arg	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ile	Phe
				35				40				45			

45

Ser	Ile	Tyr	Gly	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu
				50				55			60				

50

Glu	Trp	Val	Ala	Val	Ile	Trp	Tyr	Asp	Gly	Ser	Asn	Lys	Tyr	Tyr	Ala
				65				70			75		80		

55

Asp	Ser	Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn
					85				90			95			

55

Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val
					100				105			110			

Tyr	Tyr	Cys	Ala	Arg	Ala	Pro	His	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr
						115			120			125			

## EP 2 224 954 B1

Leu Val Thr Val Ser Ser  
130

<210> 16

5 <211> 115

<212> PRT

<213> Homo sapiens

<400> 16

10

Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg  
1 5 10 15

15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ile Phe Ser Ile Tyr  
20 25 30

20

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
35 40 45

25

Ala Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val  
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
65 70 75 80

30

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
85 90 95

35

Ala Arg Ala Pro His Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr  
100 105 110

40

Val Ser Ser  
115

<210> 17

<211> 5

<212> PRT

<213> Homo sapiens

45

<400> 17

50

Ile Tyr Gly Met His  
1 5

<210> 18

<211> 17

<212> PRT

<213> Homo sapiens

55

<400> 18

## EP 2 224 954 B1

Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys

5

1

5

10

15

Gly

10

&lt;210&gt; 19

&lt;211&gt; 6

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

15

&lt;400&gt; 19

Ala	Pro	His	Phe	Asp	Tyr
1				5	

20

&lt;210&gt; 20

&lt;211&gt; 387

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

25

&lt;400&gt; 20

atggacatga gggccccgc tcagctcctg gggcttctgc tgctctggct cccaggtgcc 60

30

agatgtgcca tccagttgac ccagtctcca tcctccctgt ctgcacatctgt aggagacaga 120

gtcaccatca cttgccgggc aagtcaaggc attagcagtg ctttagcctg gtatcagcag 180

aaaccaggga aagctcctaa gctcctgatc tatgatgcct ccagtttggaa aagtggggtc 240

35

ccatcaaggt tcagcggcag tggatctggg acagatttca ctctcaccat cagcagcctg 300

cagcctgaag attttgcaac ttattactgt caacagttta atagttaccc tctcactttc 360

ggcggaggga ccaaggtggaa gatcaaa 387

40

&lt;210&gt; 21

&lt;211&gt; 129

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

45

&lt;400&gt; 21

50

55

## EP 2 224 954 B1

Met	Asp	Met	Arg	Val	Pro	Ala	Gln	Leu	Leu	Gly	Leu	Leu	Leu	Trp		
1					5				10					15		
5	Leu	Pro	Gly	Ala	Arg	Cys	Ala	Ile	Gln	Leu	Thr	Gln	Ser	Pro	Ser	Ser
						20			25					30		
10	Leu	Ser	Ala	Ser	Val	Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser
						35			40				45			
15	Gln	Gly	Ile	Ser	Ser	Ala	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys
						50			55				60			
20	Ala	Pro	Lys	Leu	Leu	Ile	Tyr	Asp	Ala	Ser	Ser	Leu	Glu	Ser	Gly	Val
						65			70				75			80
25	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr
						85					90				95	
30	Ile	Ser	Ser	Leu	Gln	Pro	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln
					100				105					110		
35	Phe	Asn	Ser	Tyr	Pro	Leu	Thr	Phe	Gly	Gly	Thr	Lys	Val	Glu	Ile	
						115			120				125			
40	Lys															
	<210> 22															
	<211> 107															
	<212> PRT															
	<213> Homo sapiens															
45	<400> 22															
	50															
	55															

## EP 2 224 954 B1

	Ala	Ile	Gln	Leu	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
1					5						10				15	
5																
	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Gly	Ile	Ser	Ser	Ala
					20						25				30	
10																
	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile
					35					40				45		
15																
	Tyr	Asp	Ala	Ser	Ser	Leu	Glu	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
					50					55			60			
20																
	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
					65			70			75				80	
25																
	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Phe	Asn	Ser	Tyr	Pro	Leu
					85					90				95		
30																
	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Val	Glu	Ile	Lys					
					100					105						
35																
	<210> 23															
	<211> 11															
	<212> PRT															
40	<213> Homo sapiens															
	<400> 23															
45																
	Arg	Ala	Ser	Gln	Gly	Ile	Ser	Ser	Ala	Leu	Ala					
					1		5				10					
50																
	<210> 24															
	<211> 7															
	<212> PRT															
55	<213> Homo sapiens															
	<400> 24															
60																
	Asp	Ala	Ser	Ser	Leu	Glu	Ser									
					1		5									
65																
	<210> 25															
	<211> 9															
	<212> PRT															
70	<213> Homo sapiens															
	<400> 25															
75																
	Gln	Gln	Phe	Asn	Ser	Tyr	Pro	Leu	Thr							
					1		5									
80																
	<210> 26															
	<211> 417															

## EP 2 224 954 B1

<212> DNA  
 <213> Homo sapiens

<400> 26

5

atggagtttggctgagctggttttcctcgttgctctttaagaggtgtccagtgtcag 60  
 gtgcagctggtggagtcgtggggaggcggtgtccagcctggaggtccctgagactctcc 120  
 10 tgcgcagcgtctggattcaccttcagtaattatggcatgtactgggtccgccaggctcca 180  
 ggcaaggggctggagtggttgcagttataatgtatgtgaagtaataatactatgca 240  
 gactccgtgaaggccgattcaccatctccagacaattccaagaacacgctgtatctg 300  
 15 caaatgaacagcctgagagccaggacacggtgttattactgtgcgagagatctctgg 360  
 ggatggtaactttgactatttggccaggaccctggtaacccttcctcagctagc 417

20

<210> 27

<211> 149

<212> PRT

<213> Homo sapiens

25

<400> 27

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Leu Leu Arg Gly  
 1 5 10 15

30

Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln  
 20 25 30

35

Pro Gly Arg Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe  
 35 40 45

40

45

50

55

## EP 2 224 954 B1

Ser Asn Tyr Gly Met Tyr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu  
 50 55 60

5 Glu Trp Val Ala Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala  
 65 70 75 80

10 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn  
 85 90 95

15 Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val  
 100 105 110

20 Tyr Tyr Cys Ala Arg Asp Leu Trp Gly Trp Tyr Phe Asp Tyr Trp Gly  
 115 120 125

25 Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser  
 130 135 140.

30 Val Phe Pro Leu Ala  
 145

<210> 28  
 <211> 130  
 <212> PRT  
 35 <213> Homo sapiens

<400> 28

35 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg  
 1 5 10 15

40 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr  
 20 25 30

45 Gly Met Tyr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

50 Ala Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val  
 50 55 60

55 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
 65 70 75 80

55 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95

Ala Arg Asp Leu Trp Gly Trp Tyr Phe Asp Tyr Trp Gly Gln Gly Thr  
 100 105 110

## EP 2 224 954 B1

Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
 115 120 125

5 Leu Ala  
 130

<210> 29  
 <211> 5  
 10 <212> PRT  
 <213> Homo sapiens

<400> 29

15 Asn Tyr Gly Met Tyr  
 1 5

<210> 30  
 <211> 17  
 20 <212> PRT  
 <213> Homo sapiens

<400> 30

25 Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys  
 1 5 10 15

Gly

30 <210> 31  
 <211> 9  
 <212> PRT  
 <213> Homo sapiens

35 <400> 31

Asp Leu Trp Gly Trp Tyr Phe Asp Tyr  
 1 5

40 <210> 32  
 <211> 384  
 <212> DNA  
 <213> Homo sapiens

45 <400> 32

atggatgga	gctgtatcat	cctgttcctc	gtggccacag	caaccgtgt	ccactccgaa	60
attgtgttga	cacagtctcc	agccaccctg	tctttgtctc	cagggaaag	agccaccctc	120
50 tcctgcaggg	ccagtcagag	tgttagcagc	tacttagcct	ggtaccaaca	gaaacctggc	180
caggctccca	ggctcctcat	ctatgtgca	tccaacaggg	ccactggcat	cccagccagg	240
55 ttcagtggca	gtgggtctgg	gacagacttc	actctcacca	tcagcagcct	agagcctgaa	300
gattttgcag	tttattactg	tcagcagcgt	cgcaactggc	cgctcacttt	cggcggaggg	360

accaagggtgg agatcaaacg tacg

384

&lt;210&gt; 33

5 &lt;211&gt; 127

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 33

10

Met	Glu	Ala	Pro	Ala	Gln	Leu	Leu	Phe	Leu	Leu	Leu	Leu	Trp	Leu	Pro
1															15

15

Asp	Thr	Thr	Gly	Glu	Ile	Val	Leu	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser
															30
									20					25	

20

Leu	Ser	Pro	Gly	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser
															45
									35					40	

25

Val	Ser	Ser	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro
															50
									55					60	

Arg	Leu	Leu	Ile	Tyr	Asp	Ala	Ser	Asn	Arg	Ala	Thr	Gly	Ile	Pro	Ala
															65
									70					75	

30

Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser
															85
														90	

35

Ser	Leu	Glu	Pro	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	Arg	Arg
															100
														105	

110

40

Asn	Trp	Pro	Leu	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Val	Glu	Ile	Lys	
															115
														120	

125

45

&lt;210&gt; 34

&lt;211&gt; 107

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 34

50

55

## EP 2 224 954 B1

	Glu	Ile	Val	Leu	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser	Leu	Ser	Pro	Gly
1					5						10					15
5	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	Val	Ser	Ser	Tyr
					20					25					30	
10	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Arg	Leu	Leu	Ile
					35					40					45	
15	Tyr	Asp	Ala	Ser	Asn	Arg	Ala	Thr	Gly	Ile	Pro	Ala	Arg	Phe	Ser	Gly
					50					55					60	
20	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Glu	Pro
					65					70					80	
25	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	Arg	Arg	Asn	Trp	Pro	Leu
					85					90					95	
30	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Val	Glu	Ile	Lys					
					100					105						
35	<210> 35															
	<211> 11															
	<212> PRT															
40	<213> Homo sapiens															
	<400> 35															
45	Arg	Ala	Ser	Gln	Ser	Val	Ser	Ser	Tyr	Leu	Ala					
	1				5						10					
50	<210> 36															
	<211> 7															
	<212> PRT															
55	<213> Homo sapiens															
	<400> 36															
60	Asp	Ala	Ser	Asn	Arg	Ala	Thr									
	1				5											
65	<210> 37															
	<211> 9															
	<212> PRT															
70	<213> Homo sapiens															
	<400> 37															
75	Gln	Gln	Arg	Arg	Asn	Trp	Pro	Leu	Thr							
	1				5											
80	<210> 38															
	<211> 447															

<212> DNA  
 <213> Homo sapiens

5	<400> 38	
	atggagttt ggctgacactg ggtttcctc gttgctctt taagaggtgt ccagtgtcag	60
	gtgcagctgg tggagtctgg gggaggcgtg gtccagcctg ggaggtccct gagactctcc	120
10	tgtgcagcgt ctggattcac cttcagtacc tatggcatgc actgggtccg ccaggctcca	180
	ggcaaggggc tggagtgggt ggcaattata tggtatgatg gaggtataaa atactatgca	240
	gactccgtga agggccgatt caccatctcc agagacaatt ccaagaacac gctgtatctg	300
15	caaatgaaca gcctgagagc cgaggacacg gctgtgtatt actgtgcgag agacttctac	360
	tggtacttcg atctctgggg ccgtggcacc ctggtcactg tctcctcagc ctccaccaag	420
20	ggcccatcgg tcttccccct ggcaagg	447

<210> 39  
 <211> 148  
 <212> PRT  
 25 <213> Homo sapiens

<400> 39

30

35

40

45

50

55

## EP 2 224 954 B1

Met	Glu	Phe	Gly	Leu	Thr	Trp	Val	Phe	Leu	Val	Ala	Leu	Leu	Arg	Gly		
1														15			
5	Val	Gln	Cys	Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Val	Val	Gln	
														25	30		
10	Pro	Gly	Arg	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	
														35	40	45	
15	Ser	Thr	Tyr	Gly	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	
														55	60		
20	Glu	Trp	Val	Ala	Ile	Ile	Trp	Tyr	Asp	Gly	Gly	Asn	Lys	Tyr	Tyr	Ala	
														65	70	75	80
25	Asp	Ser	Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	
														85	90	95	
30	Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	
														100	105	110	
35	Tyr	Tyr	Cys	Ala	Arg	Asp	Phe	Tyr	Trp	Tyr	Phe	Asp	Leu	Trp	Gly	Arg	
														115	120	125	
40	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	
														130	135	140	
45	Phe	Pro	Leu	Ala													
														145			
50	<210>	40															
	<211>	129															
	<212>	PRT															
	<213>	Homo sapiens															
	<400>	40															
	Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Val	Val	Gln	Pro	Gly	Arg		
	1					5								10		15	

55

## EP 2 224 954 B1

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Thr Tyr  
 20 25 30

5 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

10 Ala Ile Ile Trp Tyr Asp Gly Gly Asn Lys Tyr Tyr Ala Asp Ser Val  
 50 55 60

15 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
 65 70 75 80

20 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95

25 Ala Arg Asp Phe Tyr Trp Tyr Phe Asp Leu Trp Gly Arg Gly Thr Leu  
 100 105 110

30 Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu  
 115 120 125

Ala

35 <210> 41  
 <211> 5  
 <212> PRT  
 <213> Homo sapiens

40 <400> 41

Thr Tyr Gly Met His  
 1 5

45 <210> 42  
 <211> 17  
 <212> PRT  
 <213> Homo sapiens

50 <400> 42

Ile Ile Trp Tyr Asp Gly Gly Asn Lys Tyr Tyr Ala Asp Ser Val Lys  
 1 5 10 15

55 Gly  
 <210> 43  
 <211> 8  
 <212> PRT  
 <213> Homo sapiens

<400> 43

## EP 2 224 954 B1

Asp	Phe	Tyr	Trp	Tyr	Phe	Asp	Leu
1					5		

5 &lt;210&gt; 44

&lt;211&gt; 372

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

10 &lt;400&gt; 44

atggaagccc	cagctcagct	tctcttcctc	ctgctactct	ggctcccaga	taccaccgga	60
gaaaattgtgt	tgacacagtc	tccagccacc	ctgtcttgc	ctccagggga	aagagccacc	120
15 ctctcctgca	gggccagtca	gagtgttagc	agctacttag	cctggtagca	acagaaacct	180
ggccaggctc	ccaggctcct	catctatgtat	gcattccaaca	ggccactgg	catcccagcc	240
20 aggttcagtgc	gcagtgggtc	tgggacagac	ttcactctca	ccatcagcag	cctagagcct	300
gaagattttgc	cagtttattat	ctgtcagcag	cgtggacgt	tcggccaagg	gaccaagggtg	360
gaaatcaaacc	ga					372

25 &lt;210&gt; 45

&lt;211&gt; 123

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

30 &lt;400&gt; 45

Met	Glu	Ala	Pro	Ala	Gln	Leu	Leu	Phe	Leu	Leu	Leu	Leu	Trp	Leu	Pro
1					5				10				15		

35

Asp	Thr	Thr	Gly	Glu	Ile	Val	Leu	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser
					20			25				30			

40

Leu	Ser	Pro	Gly	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser
					35			40		45					

45

Val	Ser	Ser	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro
					50			55		60					

50

Arg	Leu	Leu	Ile	Tyr	Asp	Ala	Ser	Asn	Arg	Ala	Thr	Gly	Ile	Pro	Ala
					65			70		75		80			

55

Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser
								85		90		95			

55

Ser	Leu	Glu	Pro	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	Arg	Arg
					100				105			110			

## EP 2 224 954 B1

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys  
 115 120

<210> 46

5 <211> 103

<212> PRT

<213> Homo sapiens

<400> 46

10

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly  
 1 5 10 15

15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr  
 20 25 30

20

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile  
 35 40 45

25

Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly  
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro  
 65 70 75 80

30

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Arg Thr Phe Gly Gln  
 85 90 95

35

Gly Thr Lys Val Glu Ile Lys  
 100

<210> 47

<211> 11

<212> PRT

40 <213> Homo sapiens

<400> 47

45

Arg Ala Ser Gln Ser Val Ser Ser Tyr Leu Ala  
 1 5 10

<210> 48

<211> 7

<212> PRT

50 <213> Homo sapiens

<400> 48

55

Asp Ala Ser Asn Arg Ala Thr  
 1 5

<210> 49

<211> 5

## EP 2 224 954 B1

<212> PRT  
 <213> Homo sapiens

<400> 49

5

Gln	Gln	Arg	Arg	Thr
1				5

<210> 50

10

<211> 454

<212> DNA

<213> Homo sapiens

<400> 50

15

atggagtttggctgagctgggtttcctcgttgctcttttaagagggtgtccagtgtcag 60

gtgcagctggtggagtctggggaggcggtgtccagcctggaggtccctgagactctcc 120

20

tgtgcagcgtctggattcaccttcagtagctataacatgcactgggtccgcaggctcca 180

ggcaaggggcgtggagtgggtggcatttatatgtatgatgaaagtaataatactatgga 240

gactccgtgaagggccgattcaccatctccagacaaattccaaaaacacgctgtatctg 300

25

caaatgaacagcctgagagccgacacggtgtgttactgtgcgagagaagagctg 360

gggatcgggtgtacttcgtatctggggcgtggcaccctgtcactgtctctcagcc 420

30

tccaccaaggccccatcggtttccccctggcac 454

<210> 51

<211> 151

<212> PRT

35

<213> Homo sapiens

<400> 51

40

45

50

55

## EP 2 224 954 B1

Met	Glu	Phe	Gly	Leu	Ser	Trp	Val	Phe	Leu	Val	Ala	Leu	Leu	Arg	Gly		
1														15			
5	Val	Gln	Cys	Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Val	Val	Gln	
														25	30		
10	Pro	Gly	Arg	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	
														35	40	45	
15	Ser	Ser	Tyr	Asn	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	
														50	55	60	
20	Glu	Trp	Val	Ala	Phe	Ile	Trp	Tyr	Asp	Gly	Ser	Asn	Lys	Tyr	Tyr	Gly	
														65	70	75	80
25	Asp	Ser	Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	
														85	90	95	
30	Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	
35	100														105		110
40	Tyr	Tyr	Cys	Ala	Arg	Glu	Glu	Leu	Gly	Ile	Gly	Trp	Tyr	Phe	Asp	Leu	
														115	120	125	
45	Trp	Gly	Arg	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	
														130	135	140	
50	Pro	Ser	Val	Phe	Pro	Leu	Ala										
														145	150		
55	<210> 52																
	<211> 132																
	<212> PRT																
	<213> Homo sapiens																
	<400> 52																

## EP 2 224 954 B1

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg  
 1 5 10 15

5 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr  
 20 25 30

10 Asn Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

15 Ala Phe Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Gly Asp Ser Val  
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
 65 70 75 80

20 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95

25 Ala Arg Glu Glu Leu Gly Ile Gly Trp Tyr Phe Asp Leu Trp Gly Arg  
 100 105 110

30 Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val  
 115 120 125

Phe Pro Leu Ala  
 130

35 <210> 53  
 <211> 5  
 <212> PRT  
 <213> Homo sapiens

40 <400> 53

Ser Tyr Asn Met His  
 1 5

45 <210> 54  
 <211> 17  
 <212> PRT  
 <213> Homo sapiens

50 <400> 54

Phe Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Gly Asp Ser Val Lys  
 1 5 10 15

55 Gly

<210> 55  
 <211> 11

## EP 2 224 954 B1

<212> PRT  
 <213> Homo sapiens

<400> 55

5

	Glu	Glu	Leu	Gly	Ile	Gly	Trp	Tyr	Phe	Asp	Leu
	1				5						10

<210> 56

10

<211> 454  
 <212> DNA  
 <213> Homo sapiens

<400> 56

15

atggaagccc cagctcagct tctttcctc ctgctactct ggctccaga taccaccgga 60

gaaattgtgt tgacacagtc tccagccacc ctgtcttgc ctccagggga aagagccacc 120

20

ctctcctgca gggccagtc gagtgtagc agctacttag cctggtagca acagaaacct 180

ggccaggctc ccaggctcct catctatgtat gcatccaaca gggccactgg catcccagcc 240

aggttcagtg gcagtgggtc tgggacagac ttcactctca ccatcagcag cctagagcct 300

25

gaagattttg cagtttatta ctgtcagcag ctagggacgt tcggccaagg gaccaagggtg 360

gaaatcaaac gaactgtggc tgcaccatct gtcttcatct tcccgccatc tcatgagcag 420

30

ttgaaatctg gaactgcctc tgggtgtgc ctgc 454

<210> 57

<211> 123

<212> PRT

35

<213> Homo sapiens

<400> 57

40

Met	Glu	Ala	Pro	Ala	Gln	Leu	Leu	Phe	Leu	Leu	Leu	Leu	Trp	Leu	Pro
1					5				10				15		

45

50

55

## EP 2 224 954 B1

	Asp Thr Thr Gly Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser			
20		25	30	
5	Leu Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser			
	35	40	45	
10	Val Ser Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro			
	50	55	60	
	Arg Leu Leu Ile Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala			
65	70	75	80	
15	Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser			
	85	90	95	
20	Ser Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Arg			
	100	105	110	
25	Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys			
	115	120		
	<210> 58			
	<211> 103			
	<212> PRT			
30	<213> Homo sapiens			
	<400> 58			
35	Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly			
	1	5	10	15
40	Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr			
	20	25	30	
	Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile			
	35	40	45	
45	Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly			
	50	55	60	
50	Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro			
	65	70	75	80
55	Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Arg Thr Phe Gly Gln			
	85	90	95	
	Gly Thr Lys Val Glu Ile Lys			

## EP 2 224 954 B1

<210> 59  
 <211> 11  
 <212> PRT  
 <213> Homo sapiens

5

&lt;400&gt; 59

Arg Ala Ser Gln Ser Val Ser Ser Tyr Leu Ala  
 1 5 10

10

<210> 60  
 <211> 7  
 <212> PRT  
 <213> Homo sapiens

15

&lt;400&gt; 60

Asp Ala Ser Asn Arg Ala Thr  
 1 5

20

<210> 61  
 <211> 5  
 <212> PRT  
 <213> Homo sapiens

25

&lt;400&gt; 61

Gln Gln Arg Arg Thr  
 1 5

30

<210> 62  
 <211> 433  
 <212> DNA  
 <213> Homo sapiens

35

&lt;400&gt; 62

atggagttt gctgagctg gtttcctt gttgctatataaaaagggtgt ccagtgtgag 60

40

gttcagctgg tgcagtctgg gggaggctt g tacatcctg ggggtccct gagactctcc 120

tgtgcaggct ctggattcac cttcagtaac tatgctatgc actgggtcg ccaggctcca 180

ggaaaaggta tggagtgggt atcaactatt ggtactggtg gtggcacacc ctatgcagac 240

45

tccgtgaagg gccgcttcac catctccaga gacaatgccca agaactcctt gtatcttcaa 300

atgaacagcc tgagagccga ggacatggct gtgtattact gtgcattaa tgctttgtat 360

50

gtctggggcc aaggacaat ggtcaccgtc tcttcagcct ccaccaaggg cccatcggtc 420

ttccccctgg cac 433

<210> 63  
 <211> 144  
 <212> PRT  
 <213> Homo sapiens

&lt;400&gt; 63

5	Met	Glu	Phe	Val	Leu	Ser	Trp	Val	Leu	Leu	Val	Ala	Ile	Leu	Lys	Gly
	1				5					10					15	
10	Val	Gln	Cys	Glu	Val	Gln	Leu	Val	Gln	Ser	Gly	Gly	Gly	Leu	Val	His
					20				25					30		
15	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Gly	Ser	Gly	Phe	Thr	Phe
					35			40					45			
20	Ser	Asn	Tyr	Ala	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu
					50			55				60				
25	Glu	Trp	Val	Ser	Thr	Ile	Gly	Thr	Gly	Gly	Thr	Pro	Tyr	Ala	Asp	
					65		70		75				80			
30	Ser	Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Ser
					85			90				95				
35	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Met	Ala	Val	Tyr
					100			105				110				
40	Tyr	Cys	Ala	Leu	Ser	Ala	Phe	Asp	Val	Trp	Gly	Gln	Gly	Thr	Met	Val
					115			120				125				
45	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala
					130			135				140				
50	<210> 64															
	<211> 125															
	<212> PRT															
	<213> Homo sapiens															
55	<400> 64															

## EP 2 224 954 B1

Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly  
 1 5 10 15

5 Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Asn Tyr  
 20 25 30

10 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

15 Ser Thr Ile Gly Thr Gly Gly Thr Pro Tyr Ala Asp Ser Val Lys  
 50 55 60

Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr Leu  
 65 70 75 80

20 Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys Ala  
 85 90 95

25 Leu Ser Ala Phe Asp Val Trp Gly Gln Gly Thr Met Val Thr Val Ser  
 100 105 110

30 Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala  
 115 120 125

35 <210> 65  
 <211> 5  
 <212> PRT  
 <213> Homo sapiens

40 <400> 65

Asn Tyr Ala Met His  
 1 5

45 <210> 66  
 <211> 16  
 <212> PRT  
 <213> Homo sapiens

50 <400> 66

Thr Ile Gly Thr Gly Gly Thr Pro Tyr Ala Asp Ser Val Lys Gly  
 1 5 10 15

55 <210> 67  
 <211> 5  
 <212> PRT  
 <213> Homo sapiens

<400> 67

## EP 2 224 954 B1

Ser	Ala	Phe	Asp	Val
1				5

&lt;210&gt; 68

&lt;211&gt; 466

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 68

10

atggagtttg tgctgagctg ggtttcctt gttgctatat taaaagggtgt ccagtgtgag 60

gttcagctgg tgcagtcgg gggaggctt gtagatcctg gggggccct gagactctcc 120

15

tgtgcaggct ctggattcac cttagttagc tatgctatgc actgggtcg ccaggctcca 180

ggaaaaggc tggagtgggt atcagctatt ggtactggtg gttacacata ctatgttagac 240

tccgtgaagg gccgattcac catctccaga gacaatgcc agaagtccct gtatctcaa 300

20

atgaacagcc tgagagccga ggacatggct gtgttattact gtgcaagaga gccgtttac 360

gatattttga ctggttattc cccatacttt gactactggg gccaggaaac cctggtcacc 420

25

gtctcctcag cctccaccaa gggccatcg gtctccccc tggcac 466

&lt;210&gt; 69

&lt;211&gt; 143

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 69

35

40

45

50

55

## EP 2 224 954 B1

Met	Glu	Phe	Val	Leu	Ser	Trp	Val	Phe	Leu	Val	Ala	Ile	Leu	Lys	Gly	
1				5					10					15		
5	Val	Gln	Cys	Glu	Val	Gln	Leu	Val	Gln	Ser	Gly	Gly	Gly	Leu	Val	His
					20				25					30		
10	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Gly	Ser	Gly	Phe	Thr	Phe
					35				40				45			
15	Ser	Ser	Tyr	Ala	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu
					50				55				60			
20	Glu	Trp	Val	Ser	Ala	Ile	Gly	Thr	Gly	Tyr	Thr	Tyr	Tyr	Val	Asp	
					65			70		75				80		
25	Ser	Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Lys	Ser
					85				90				95			
30	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Met	Ala	Val	Tyr
					100				105				110			
35	Tyr	Cys	Ala	Arg	Glu	Pro	Phe	Tyr	Asp	Ile	Leu	Thr	Gly	Tyr	Ser	Pro
					115				120				125			
40	Tyr	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	
					130			135				140				
45	<210>	70	<211>	124	<212>	PRT	<213>	Homo sapiens								
50	<400>	70														
55	Glu	Val	Gln	Leu	Val	Gln	Ser	Gly	Gly	Gly	Leu	Val	His	Pro	Gly	Gly
					1		5			10			15			
60	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Gly	Ser	Gly	Phe	Thr	Phe	Ser	Ser	Tyr
					20			25				30				

## EP 2 224 954 B1

Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

5 Ser Ala Ile Gly' Thr Gly Gly Tyr Thr Tyr Tyr Val Asp Ser Val Lys  
 50 55 60

10 Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Lys Ser Leu Tyr Leu  
 65 70 75 80

15 Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys Ala  
 85 90 95

20 Arg Glu Pro Phe Tyr Asp Ile Leu Thr Gly Tyr Ser Pro Tyr Phe Asp  
 100 105 110

25 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser  
 115 120

<210> 71  
 <211> 5  
 <212> PRT  
 <213> Homo sapiens

30 <400> 71

35 Ser Tyr Ala Met His  
 1 5

<210> 72  
 <211> 16  
 <212> PRT  
 <213> Homo sapiens

40 <400> 72

45 Ala Ile Gly Thr Gly Gly Tyr Thr Tyr Tyr Val Asp Ser Val Lys Gly  
 1 5 10 15

<210> 73  
 <211> 16  
 <212> PRT  
 <213> Homo sapiens

50 <400> 73

55 Glu Pro Phe Tyr Asp Ile Leu Thr Gly Tyr Ser Pro Tyr Phe Asp Tyr  
 1 5 10 15

<210> 74  
 <211> 454  
 <212> DNA  
 <213> Homo sapiens

## EP 2 224 954 B1

&lt;400&gt; 74

	atggagtttggctgagctggtttcctcgttgctctttaagagggtgtccagtgtcag	60
5	gtgcagctggtggagtcgtggggaggcgtgtccagcctggaggtccctgagactctcc	120
	tgtgcagcgtctggattcaccttcagtagctataacatgcactgggtccgccaggctcca	180
10	ggcaaggggctggagtggtggcagttatatgtatgtgaagtaataatactatggagactccgtgaagggccgattcaccatctccagacaattccaagaacacgctgtatctg	240
	caaataacaacgcctgagagcaggacacggtgttattactgtgcagagaagagctg	300
15	gggatcgggtgtacttcgatctctggggccgtggcaccctgtcactgtctcctcagcc	360
	tccaccaaggccccatcggtcttccccctggcac	420
		454

&lt;210&gt; 75

20 &lt;211&gt; 139

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 75

25

Met	Glu	Phe	Gly	Leu	Ser	Trp	Val	Phe	Leu	Val	Ala	Leu	Leu	Arg	Gly
1														15	

30

Val	Gln	Cys	Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Val	Val	Gln
														30	
20								25							

35

Pro	Gly	Arg	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe
														45	
35								40							

40

Ser	Ser	Tyr	Asn	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu
															60
50								55							

Glu	Trp	Val	Ala	Val	Ile	Trp	Tyr	Asp	Gly	Ser	Asn	Lys	Tyr	Tyr	Gly
														80	
65					70				75						

45

Asp	Ser	Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn
															95
85								90							

50

Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val
															110
100								105							

55

Tyr	Tyr	Cys	Ala	Arg	Glu	Glu	Leu	Gly	Ile	Gly	Trp	Tyr	Phe	Asp	Leu
															125
115								120							

Trp	Gly	Arg	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser					
130								135							

## EP 2 224 954 B1

<210> 76  
 <211> 120  
 <212> PRT  
 <213> Homo sapiens

5

&lt;400&gt; 76

10 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg  
 1 5 10 15

10

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr  
 20 25 30

15

Asn Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

20

Ala Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Gly Asp Ser Val  
 50 55 60

25

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
 65 70 75 80

30

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95

35

Gly Thr Leu Val Thr Val Ser Ser  
 115 120

&lt;210&gt; 77

&lt;211&gt; 5

40

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 77

45

Ser Tyr Asn Met His  
 1 5

50

<210> 78  
 <211> 17  
 <212> PRT  
 <213> Homo sapiens

55

<400> 78  
 Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Gly Asp Ser Val Lys  
 1 5 10 15

Gly

## EP 2 224 954 B1

<210> 79  
 <211> 11  
 <212> PRT  
 <213> Homo sapiens

5

&lt;400&gt; 79

Glu	Glu	Leu	Gly	Ile	Gly	Trp	Tyr	Phe	Asp	Leu
1					5					10

10

<210> 80  
 <211> 475  
 <212> DNA  
 <213> Homo sapiens

15

&lt;400&gt; 80

atggacatga	gggtccccgc	tca	gctcctg	ggc	ttctgc	tgc	tctggct	ccc	agggtgcc	60
20	agatgtgcca	tcc	agttgac	cc	agtctcca	tc	ctccctgt	ct	gcac	120
	gtcaccatca	ctt	gccgggc	aag	tca	at	tcagg	ctt	atcagcag	180
25	aaaccaggga	aag	ctcctaa	g	ctc	tgat	gcct	cc	atgggg	240
	ccatcaaggt	tc	agcggcag	tgg	at	ctgg	ac	at	tc	300
	cagcctgaag	at	tttgcaac	tt	at	ttactgt	ca	ac	ttaccc	360
30	caagggacac	gac	tggagat	taa	ac	gtgg	ctgcac	cat	ttcccg	420
	ccatctgatg	agc	atgtgaa	at	ctgg	act	gc	ctgtt	tgtgc	475

35

<210> 81  
 <211> 128  
 <212> PRT  
 <213> Homo sapiens

40

&lt;400&gt; 81

45

50

55

## EP 2 224 954 B1

Met	Asp	Met	Arg	Val	Pro	Ala	Gln	Leu	Leu	Gly	Leu	Leu	Leu	Trp	
1														15	
5															
Leu	Pro	Gly	Ala	Arg	Cys	Ala	Ile	Gln	Leu	Thr	Gln	Ser	Pro	Ser	Ser
															30
10															
Leu	Ser	Ala	Ser	Val	Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser
															45
15															
Gln	Gly	Ile	Ser	Ser	Ala	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys
															60
20															
Ala	Pro	Lys	Leu	Leu	Ile	Tyr	Asp	Ala	Ser	Ser	Leu	Glu	Ser	Gly	Val
															80
25															
Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr
															95
30															
Ile	Ser	Ser	Leu	Gln	Pro	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln
															110
35															
Phe	Asn	Ser	Tyr	Pro	His	Phe	Gly	Gln	Gly	Thr	Arg	Leu	Glu	Ile	Lys
															125
40															
<210> 82															
<211> 106															
<212> PRT															
<213> Homo sapiens															
45															
<400> 82															
50															
55															

## EP 2 224 954 B1

	Ala	Ile	Gln	Leu	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
1															15	
	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Gly	Ile	Ser	Ser	Ala
5															30	
	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile
10															45	
	Tyr	Asp	Ala	Ser	Ser	Leu	Glu	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
15														50		
	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
20															80	
	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Phe	Asn	Ser	Tyr	Pro	His
25															95	
	Phe	Gly	Gln	Gly	Thr	Arg	Leu	Glu	Ile	Lys						
30											100				105	
	<210> 83															
	<211> 11															
	<212> PRT															
35	<213> Homo sapiens															
	<400> 83															
	Arg	Ala	Ser	Gln	Gly	Ile	Ser	Ser	Ala	Leu	Ala					
40												1		5	10	
	<210> 84															
	<211> 7															
	<212> PRT															
45	<213> Homo sapiens															
	<400> 84															
	Asp	Ala	Ser	Ser	Leu	Glu	Ser									
50								1		5						
	<210> 85															
	<211> 8															
	<212> PRT															
55	<213> Homo sapiens															
	<400> 85															
	Gln	Gln	Phe	Asn	Ser	Tyr	Pro	His								
60									1		5					
	<210> 86															
	<211> 445															

## EP 2 224 954 B1

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 86

5

atggagtttgcg	ggctgagctgg	gttttcctc	gttgctcttta	aaagggtgt	ccagtgtcag	60	
gtgcagctgg	tggagtcgtgg	gggaggcgtg	gtccagcctg	ggaggctccct	gagactctcc	120	
10	tgtgcagcgt	ctggattcac	cttcagtagc	tatggcatgc	actgggtccg	ccaggctcca	180
ggcaaggggc	tggagtgggt	ggcagttata	tggtatgtg	gaagtaataa	atactatgca	240	
15	gactccgtga	agggccgatt	caccatctcc	agagacaatt	ccaagaacac	gctgtatctg	300
caaataaca	gcctgagagc	cgaggacacg	gctgtgtatt	actgtgcgag	aggccccct	360	
20	cgtacttcg	atctctgggg	ccgtggcacc	ctggtcactg	tctcctcagc	ctccaccaag	420
ggcccatcgg	tcttccccct	ggcac					445

&lt;210&gt; 87

&lt;211&gt; 136

&lt;212&gt; PRT

25

&lt;213&gt; Homo sapiens

&lt;400&gt; 87

30

Met	Glu	Phe	Gly	Leu	Ser	Trp	Val	Phe	Leu	Val	Ala	Leu	Leu	Arg	Gly
1				5					10				15		

35

Val	Gln	Cys	Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Val	Val	Gln
				20				25				30		

40

Pro	Gly	Arg	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe
				35				40				45			

45

Ser	Ser	Tyr	Gly	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu
				50				55			60				

50

Glu	Trp	Val	Ala	Val	Ile	Trp	Tyr	Asp	Gly	Ser	Asn	Lys	Tyr	Tyr	Ala
				65				70			75		80		

55

## EP 2 224 954 B1

	Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn
	85 90 95
5	Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
	100 105 110
10	Tyr Tyr Cys Ala Arg Gly Pro Pro Arg Tyr Phe Asp Leu Trp Gly Arg
	115 120 125
	Gly Thr Leu Val Thr Val Ser Ser
	130 135
15	<210> 88
	<211> 117
	<212> PRT
	<213> Homo sapiens
20	<400> 88
25	Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
	1 5 10 15
	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
	20 25 30
30	Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
	35 40 45
35	Ala Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
	50 55 60
40	Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
	65 70 75 80
	Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
	85 90 95
45	Ala Arg Gly Pro Pro Arg Tyr Phe Asp Leu Trp Gly Arg Gly Thr Leu
	100 105 110
50	Val Thr Val Ser Ser
	115
	<210> 89
	<211> 5
55	<212> PRT
	<213> Homo sapiens
	<400> 89

## EP 2 224 954 B1

Ser Tyr Gly Met His

	1	5		
5				
	<210> 90			
	<211> 17			
	<212> PRT			
	<213> Homo sapiens			
10				
	<400> 90			
15				
	Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys			
	1	5	10	15
	Gly			
20	<210> 91			
	<211> 8			
	<212> PRT			
	<213> Homo sapiens			
25	<400> 91			
	Gly Pro Pro Arg Tyr Phe Asp Leu			
	1	5		
30	<210> 92			
	<211> 95			
	<212> PRT			
	<213> Homo sapiens			
35	<400> 92			
	Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly			
	1	5	10	15
40				
	Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr			
	20	25	30	
45	Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile			
	35	40	45	
50	Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly			
	50	55	60	
	Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro			
	65	70	75	80
55				
	Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro			
	85	90	95	

## EP 2 224 954 B1

<210> 93  
<211> 95  
<212> PRT  
<213> Homo sapiens

5

&lt;400&gt; 93

10 Ala Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly  
1 5 10 15

15 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Ala  
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile  
35 40 45

20 Tyr Asp Ala Ser Ser Leu Glu Ser Gly Val Pro Ser Arg Phe Ser Gly  
50 55 60

25 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro  
65 70 75 80

30 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Phe Asn Ser Tyr Pro  
85 90 95

35

<210> 94  
<211> 95  
<212> PRT  
<213> Homo sapiens

&lt;400&gt; 94

40

45

50

55

## EP 2 224 954 B1

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly	
1					5					10				15		
5	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Arg	Gln	Gly	Ile	Ser	Ser	Trp
					20					25				30		
10	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Glu	Lys	Ala	Pro	Lys	Ser	Leu	Ile
					35				40				45			
15	Tyr	Ala	Ala	Ser	Ser	Leu	Gln	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
					50				55			60				
20	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
					65			70			75			80		
25	<210>	95	<211>	98	<212>	PRT	<213> Homo sapiens									
30	<400> 95															
35	Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Val	Val	Gln	Pro	Gly	Arg	
					1			5			10			15		
40	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Ser	Tyr
					20				25				30			
45	Gly	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val
					35				40			45				
50	Ala	Val	Ile	Trp	Tyr	Asp	Gly	Ser	Asn	Lys	Tyr	Tyr	Ala	Asp	Ser	Val
					50				55			60				
55	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
					65			70			75			80		
60	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
						85				90			95			
65	Ala Arg															
70	<210> 96															
	<211> 97															

<212> PRT  
<213> Homo sapiens

<400> 96

5

Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly  
1 5 10 15

10

Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Ser Tyr  
20 25 30

15

Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
35 40 45

20

Ser Ala Ile Gly Thr Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys  
50 55 60

25

Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr Leu  
65 70 75 80

30

Arg

35

<210> 97  
<211> 5  
<212> PRT  
<213> Artificial sequence

<220>  
<221> source  
<223> /note="Description of artificial sequence: Synthetic consensus sequence"

40

<220>  
<221> VARIANT  
<222> (1)..(1)  
<223> /replace="Asn" or "Thr" or "Ser"

45

<220>  
<221> misc\_feature  
<222> (1)..(1)  
<223> /note="Residue given in the sequence has no preference with respect to those in the annotation for said position"

50

<220>  
<221> VARIANT  
<222> (3)..(3)  
<223> /replace="Asn" or "Ala"

<220>  
<221> misc\_feature

## EP 2 224 954 B1

<222> (3)..(3)  
 <223> /note="Residue given in the sequence has no preference with respect to those in the annotation for said position"

5 <220>  
 <221> VARIANT  
 <222> (5)..(5)  
 <223> /replace="Tyr"

10 <220>  
 <221> misc\_feature  
 <222> (5)..(5)  
 <223> /note="Residue given in the sequence has no preference with respect to those in the annotation for said position"

15 <400> 97

Ile	Tyr	Gly	Met	His
1			5	

20 <210> 98  
 <211> 17  
 <212> PRT  
 <213> Artificial sequence

25 <220>  
 <221> source  
 <223> /note="Description of artificial sequence: Synthetic consensus sequence"

30 <220>  
 <221> VARIANT  
 <222> (1)..(1)  
 <223> /replace="Ile" or "Phe" or "Thr" or "Ala"

35 <220>  
 <221> misc\_feature  
 <222> (1)..(1)  
 <223> /note="Residue given in the sequence has no preference with respect to those in the annotation for said position"

40 <220>  
 <221> VARIANT  
 <222> (3)..(3)  
 <223> /replace="Gly"

45 <220>  
 <221> VARIANT  
 <222> (4)..(4)  
 <223> /replace="Thr"

50 <220>  
 <221> VARIANT  
 <222> (5)..(5)  
 <223> /replace="Gly"

55 <220>  
 <221> misc\_feature  
 <222> (3)..(5)

## EP 2 224 954 B1

<223> /note="Residues given in the sequence have no preference with respect to those in the annotations for said positions"

5 <220>

<221> VARIANT

<222> (7)..(7)

<223> /replace="Gly" or "Tyr"

10 <220>

<221> VARIANT

<222> (8)..(8)

<223> /replace="Thr"

15 <220>

<221> VARIANT

<222> (9)..(9)

<223> /replace="Pro"

20 <220>

<221> misc\_feature

<222> (7)..(9)

<223> /note="Residues given in the sequence have no preference with respect to those in the annotations for said positions"

25 <220>

<221> VARIANT

<222> (11)..(11)

<223> /replace="Ala" or "Val"

30 <220>

<221> VARIANT

<222> (12)..(12)

<223> /replace="Gly" or " "

35 <220>

<221> misc\_feature

<222> (11)..(12)

<223> /note="Residues given in the sequence have no preference with respect to those in the annotations for said positions"

40 <400> 98

Val	Ile	Trp	Tyr	Asp	Gly	Ser	Asn	Lys	Tyr	Tyr	Ala	Asp	Ser	Val	Lys
1					5				10					15	

45

Gly

50 <210> 99

<211> 6

<212> PRT

<213> Artificial sequence

55 <220>

<221> source

<223> /note="Description of artificial sequence: Synthetic consensus sequence"

<220>

<221> VARIANT  
 <222> (1)..(1)  
 <223> /replace="Gly" or "Tyr" or "Ser" or "Pro" or "  
  
 5 <220>  
 <221> VARIANT  
 <222> (2)..(2)  
 <223> /replace="Trp" or "Ser" or "Arg"  
  
 10 <220>  
 <221> VARIANT  
 <222> (3)..(3)  
 <223> /replace="Ala" or "His"  
  
 15 <220>  
 <221> misc\_feature  
 <222> (1)..(3)  
 <223> /note="Residues given in the sequence have no preference with respect to those in the annotations for said positions"  
  
 20 <220>  
 <221> VARIANT  
 <222> (6)..(6)  
 <223> /replace="Leu" or "Val"  
  
 25 <220>  
 <221> misc\_feature  
 <222> (6)..(6)  
 <223> /note="Residue given in the sequence has no preference with respect to those in the annotation for said position"  
  
 30 <400> 99

Ala	Pro	Tyr	Phe	Asp	Tyr
1				5	

35 <210> 100  
 <211> 11  
 <212> PRT  
 40 <213> Artificial sequence

<220>  
 <221> source  
 <223> /note="Description of artificial sequence: Synthetic consensus sequence"

45 <220>  
 <221> VARIANT  
 <222> (5)..(5)  
 <223> /replace="Gly"

50 <220>  
 <221> VARIANT  
 <222> (6)..(6)  
 <223> /replace="Val"

55 <220>  
 <221> misc\_feature  
 <222> (5)..(6)

## EP 2 224 954 B1

<223> /note="Residues given in the sequence have no preference with respect to those in the annotations for said positions"

5 <220>

<221> VARIANT

<222> (9)..(9)

<223> /replace="Trp" or "Ala"

10 <220>

<221> misc\_feature

<222> (9)..(9)

<223> /note="Residue given in the sequence has no preference with respect to those in the annotation for said position"

15 <400> 100

Arg	Ala	Ser	Gln	Ser	Ile	Ser	Ser	Tyr	Leu	Ala
1				5					10	

20 <210> 101

<211> 7

<212> PRT

<213> Artificial sequence

25 <220>

<221> source

<223> /note="Description of artificial sequence: Synthetic consensus sequence"

30 <220>

<221> VARIANT

<222> (1)..(1)

<223> /replace="Ala"

35 <220>

<221> misc\_feature

<222> (1)..(1)

<223> /note="Residue given in the sequence has no preference with respect to those in the annotation for said position"

40 <220>

<221> VARIANT

<222> (4)..(4)

<223> /replace="Ser"

45 <220>

<221> VARIANT

<222> (5)..(5)

<223> /replace="Leu"

50 <220>

<221> VARIANT

<222> (6)..(6)

<223> /replace="Gln" or "Glu"

55 <220>

<221> VARIANT

<222> (7)..(7)

<223> /replace="Ser"

5                   <220>  
 <221> misc\_feature  
 <222> (4)..(7)  
 <223> /note="Residues given in the sequence have no preference with respect to those in the annotations for said  
 positions"  
 10                <400> 101  
 10                   Asp Ala Ser Asn Arg Ala Thr  
 1                    1                   5  
 15                <210> 102  
 <211> 9  
 <212> PRT  
 <213> Artificial sequence  
 20                <220>  
 <221> source  
 <223> /note="Description of artificial sequence: Synthetic consensus sequence"  
 20                <220>  
 <221> VARIANT  
 <222> (3)..(3)  
 <223> /replace="Tyr" or "Phe"  
 25                <220>  
 <221> VARIANT  
 <222> (4)..(4)  
 <223> /replace="Asn"  
 30                <220>  
 <221> VARIANT  
 <222> (5)..(5)  
 <223> /replace="Ser" or "Asn"  
 35                <220>  
 <221> VARIANT  
 <222> (6)..(6)  
 <223> /replace="Trp" or " "  
 40                <220>  
 <221> VARIANT  
 <222> (7)..(7)  
 <223> /replace=" "  
 45                <220>  
 <221> VARIANT  
 <222> (8)..(8)  
 <223> /replace="Leu" or "His" or " "  
 50                <220>  
 <221> VARIANT  
 <222> (9)..(9)  
 <223> /replace=" "  
 55                <220>  
 <221> misc\_feature  
 <222> (3)..(9)

## EP 2 224 954 B1

<223> /note="Residues given in the sequence have no preference with respect to those in the annotations for said positions"

5 <400> 102

1 Gln Gln Arg Arg Thr Tyr Pro Tyr Thr  
5

10 <210> 103

<211> 118

<212> PRT

<213> Homo sapiens

15 <400> 103

1 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg  
5 10 15

20

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr  
20 25 30

25

Gly Met Tyr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
35 40 45

30

Ala Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val  
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
65 70 75 80

35

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
85 90 95

40

Ala Arg Asp Leu Trp Gly Trp Tyr Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

45

Leu Val Thr Val Ser Ser  
115

<210> 104

<211> 117

<212> PRT

<213> Homo sapiens

50

<400> 104

55

## EP 2 224 954 B1

1	10	15
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg		
5	5	10
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Thr Tyr		
20	25	30
5		
Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val		
10	35	40
45		
10		
Ala Ile Ile Trp Tyr Asp Gly Gly Asn Lys Tyr Tyr Ala Asp Ser Val		
15	50	55
60		
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr		
20	65	70
75		
20		
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys		
25	85	90
95		
25		
Ala Arg Asp Phe Tyr Trp Tyr Phe Asp Leu Trp Gly Arg Gly Thr Leu		
30	100	105
110		
30		
Val Thr Val Ser Ser		
35	115	
<210> 105		
<211> 120		
<212> PRT		
<213> Homo sapiens		
40	<400> 105	
40		
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg		
45	1	5
10		
15		

50

55

## EP 2 224 954 B1

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr  
 20 25 30

5 Asn Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

10 Ala Phe Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Gly Asp Ser Val  
 50 55 60

15 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
 65 70 75 80

20 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95

25 Ala Arg Glu Glu Leu Gly Ile Gly Trp Tyr Phe Asp Leu Trp Gly Arg  
 100 105 110

30 Gly Thr Leu Val Thr Val Ser Ser  
 115 120

<210> 106  
 <211> 113  
 <212> PRT  
 30 <213> Homo sapiens

<400> 106

35 Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly  
 1 5 10 15

40 Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Asn Tyr  
 20 25 30

Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

45 Ser Thr Ile Gly Thr Gly Gly Thr Pro Tyr Ala Asp Ser Val Lys  
 50 55 60

50 Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr Leu  
 65 70 75 80

55 Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys Ala  
 85 90 95

Leu Ser Ala Phe Asp Val Trp Gly Gln Gly Thr Met Val Thr Val Ser  
 100 105 110

**DK/EP 2224954 T3**

**EP 2 224 954 B1**

Ser

**Patentkrav**

**1.** Et isoleret, monoklonalt antistof, som binder til human dendritisk og epitelcelle 205 receptor (DEC-205) og omfatter:

en tung kæde variabel region CDR1 omfattende SEQ ID NO: 29;

en tung kæde variabel region CDR2 omfattende SEQ ID NO: 30;

en tung kæde variabel region CDR3 omfattende SEQ ID NO: 31;

en let kæde variabel region CDR1 omfattende SEQ ID NO: 35;

en let kæde variabel region CDR2 omfattende SEQ ID NO: 36, og

en let kæde variabel region CDR3 omfattende SEQ ID NO: 37.

**2.** Antistoffet ifølge krav 1 omfattende en tung kæde variabel region omfattende SEQ ID NO: 28 og en let kæde variabel region omfattende SEQ ID NO: 34, eller en tung kæde variabel region og en let kæde variabel region, der kodes af henholdsvis SEQ ID NO: 26 og SEQ ID NO:32.

**3.** Antistoffet ifølge krav 1, som er et human monoklonalt antistof, der binder human DEC-205 med en affinitetskonstant på mindst  $10^8 \text{ M}^{-1}$  som bestemt ved overfladeplasmon-resonans, og eventuelt udviser mindst en af de følgende egenskaber:

(a) internaliseres efter binding til human dendritiske celler, der udtrykker DEC-205;

(b) frembringer, eller forøger, human CD4<sup>+</sup> T-celle svar mod et antigen, hvori human T-celle svaret eventuelt er medieret af enten MHC klasse I eller MHC klasse II stiveje, eller af dem begge;

(c) genererer, eller forbedrer, et human CTL eller NKT svar på et antigen;

- (d) lokaliseres til antigen-processerende rum i dendritiske celler;
- (e) inducerer perifer CD8<sup>+</sup> T-celle-tolerance, eller
- (f) binder til en epitop lokaliseret i det ekstracellulære domæne af human DEC-205.

**4.** Antistoffet ifølge krav 1 eller 2, hvor antistoffet er et humant antistof.

**5.** Antistoffet ifølge krav 1 eller 2, hvor antistoffet er et kimært antistof, som er et IgG2, IgG3, IgE, IgA, IgM eller IgD-antistof, eller er et antistof valgt fra gruppen bestående af et IgG1, et IgG2, et IgG3, et IgG4, et IgM, et IgA1, et IgA2, et IgAsec, et IgD og et IgE-antistof.

**6.** En ekspressionsvektor omfattende en nukleotidsekvens, der koder for antistoffet ifølge et hvilket som helst af kravene 1 til 5.

**7.** En celle, der er transformert med en ekspressions-vektor ifølge krav 6.

**8.** Et molekylært konjugat omfattende antistoffet ifølge et hvilket som helst af kravene 1 til 7 bundet til et antigen, et allergen eller et autoantigen.

**9.** Det molekylære konjugat ifølge krav 8, hvor antigenet er valgt fra gruppen bestående af en komponent af et patogen, såsom HIV, HPV, HBV eller HCV, et tumor-antigen, såsom βhCG, gp100 eller Pmel17, HER2/neu, WT1, mesothelin, CEA, gp100, MART1, TRP-2, NY-BR-1, NY-CO-58, MN (gp250), idiotype, Tyrosinase, Telomerase, SSX2, MUC-1, MART1, melan-A, NY-ESO-1, MAGE-1, MAGE-3, MAGE-A3, og høj molekylvægt

melanom-associeret antigen (HMW-MAA).

**10.** Det molekulære konjugat ifølge krav 8 eller 9 yderligere omfattende et terapeutisk middel valgt fra gruppen bestående af et cytotoxisk middel, et immunundertrykkende middel, og et kemoterapeutisk middel.

**11.** Et bispecifikt molekyle omfattende antistoffet ifølge et hvilket som helst af kravene 1 til 5 bundet til et molekyle, der har en bindingsspecificitet, som er forskellig fra antistoffet.

**12.** En sammensætning omfattende antistoffet ifølge et hvilket som helst af kravene 1 til 5 og en farmaceutisk effektiv bærer, og eventuelt yderligere omfattende et terapeutisk middel.


**13.** Anvendelse af sammensætningen ifølge krav 12, eller af konjugatet ifølge et hvilket som helst af kravene 8 til 10, ved fremstillingen af et lægemiddel til anvendelse i behandling af cancer eller en autoimmun sygdom.


**14.** Sammensætningen ifølge krav 12, eller konjugatet ifølge et hvilket som helst af kravene 8 til 10, til anvendelse ved behandling af en lidelse i et individ, til immunisering af et individ, eller til induktion eller forøgelse af et immunsvær til et antigen i et individ.


**15.** En fremgangsmåde til påvisning af tilstedeværelsen eller fraværet af DEC-205 i en biologisk prøve, omfattende:

(a) at kontakte en biologisk prøve med antistoffet ifølge et hvilket som helst af kravene 1 til 5, hvor antistoffet er mærket med et påviseligt stof, og

(b) påvisning af antistoffet bundet til DEC-205 for derved at detektere tilstedeværelsen eller fraværet af DEC-205 i den biologiske prøve.


**Fig. 1**

*Fig. 2*



**Fig. 3**

## Dec205 mAb Binding and Internalization in Dendritic Cells



Monocyte-derived human DCs were pulsed with FITC-3G9-2D2 or FITC-human IgG1 for 30 minutes on ice. Cells were then incubated at 37°C for the indicated periods to allow for internalization. Images were captured by confocal microscopy, and green staining revealed the presence of FITC-labeled molecules.

**Fig. 4**

## Human $V_H$ and $V_K$ Alignments and Germline Sequences

### $V_K$ Alignments

		CDR1	V	CDR2	J
<u>L6 germ.</u> 3G9-2D2		EIVLTO SPATLS SGERATLS C S Q V S S Y L A V Y Q Q K P G Q A P R L I V A S N P A T I P A R F S G S G G I D E F T L I S S L E P D F A V Y Y C Q P O R S N P P			I F G G G T R V E I K J K 4
5A8-1F1		EIVLTO SPATLS SGERATLS C S Q V S S Y L A V Y Q Q K P G Q A P R L I V A S N P A T I P A R F S G S G G I D E F T L I S S L E P D F A V Y Y C Q P O R S N P P			I F G G G T R V E I K J K 4
3C7-3A3		EIVLTO SPATLS SGERATLS C S Q V S S Y L A V Y Q Q K P G Q A P R L I V A S N P A T I P A R F S G S G G I D E F T L I S S L E P D F A V Y Y C Q P O R S N P P			I F G G G T R V E I K J K 4
 <u>L4 germ.</u> 5C3-2-3F6					
5D6-4C8		A I Q L T O S P S S I L A S V G D R V T I T C A S Q G I S S A L A V Y Q Q K P G K A P K L I V I P A S S L E S G V P S R F S G S G G S G T D F T L I S S L Q P E D F A T Y Y C Q O N S P P			I F G G G T R V E I K J K 5
		A I Q L T O S P S S I L A S V G D R V T I T C A S Q G I S S A L A V Y Q Q K P G K A P K L I V I P A S S L E S G V P S R F S G S G G S G T D F T L I S S L Q P E D F A T Y Y C Q O N S P P			I F G G G T R V E I K J K 5
 <u>L15 germ.</u> 3D6-2E4					
		D I Q M T O S P S S I L A S V G D R V T I T C A S Q G I S S A L A V Y Q Q K P G K A P K L I V I P A S S L E S G V P S R F S G S G G S G T D F T L I S S L Q P E D F A T Y Y C Q O N S P P			I F G G G T R V E I K J K 2
		D I Q M T O S P S S I L A S V G D R V T I T C A S Q G I S S A L A V Y Q Q K P G K A P K L I V I P A S S L E S G V P S R F S G S G G S G T D F T L I S S L Q P E D F A T Y Y C Q O N S P P			I F G G G T R V E I K J K 2

### $V_H$ Alignments

		CDR1	V	CDR2	D	CDR3	J
<u>3-33 germ.</u> 3D6-2F4		Q V Q L V E S G G G V V Q P G R S L R U S C A A S G F T S Y G M H V R Q A P G K G L E V N A V I M D G S N K Y A D S V K G R E F T I S R D N S K I T L Y Q M N S L R A E D T A V Y C A R			A P H	F D I M Q G Q T I L V T V S J K 4	
3D6-4C8		Q V Q L V E S G G G V V Q P G R S L R U S C A A S G F T S Y G M H V R Q A P G K G L E V N A V I M D G S N K Y A D S V K G R E F T I S R D N S K I T L Y Q M N S L R A E D T A V Y C A R			A P H	F D I M Q G Q T I L V T V S J K 4	
3G9-2D2		Q V Q L V E S G G G V V Q P G R S L R U S C A A S G F T S Y G M H V R Q A P G K G L E V N A V I M D G S N K Y A D S V K G R E F T I S R D N S K I T L Y Q M N S L R A E D T A V Y C A R			D I R G M	Y F D I M Q G Q T I L V T V S J K 4	
5A8-1F1		Q V Q L V E S G G G V V Q P G R S L R U S C A A S G F T S Y G M H V R Q A P G K G L E V N A V I M D G S N K Y A D S V K G R E F T I S R D N S K I T L Y Q M N S L R A E D T A V Y C A R			D E	Y F D I M Q G Q T I L V T V S J K 2	
3C7-3A3		Q V Q L V E S G G G V V Q P G R S L R U S C A A S G F T S Y G M H V R Q A P G K G L E V N A V I M D G S N K Y A D S V K G R E F T I S R D N S K I T L Y Q M N S L R A E D T A V Y C A R			E E L G G	W F D I M Q G Q T I L V T V S J K 2	
5C3-2-3F6		Q V Q L V E S G G G V V Q P G R S L R U S C A A S G F T S Y G M H V R Q A P G K G L E V N A V I M D G S N K Y A D S V K G R E F T I S R D N S K I T L Y Q M N S L R A E D T A V Y C A R			E E L G G	W F D I M Q G Q T I L V T V S J K 2	
5D12-5G1		Q V Q L V E S G G G V V Q P G R S L R U S C A A S G F T S Y G M H V R Q A P G K G L E V N A V I M D G S N K Y A D S V K G R E F T I S R D N S K I T L Y Q M N S L R A E D T A V Y C A R			G P P R	Y F D I M Q G Q T I L V T V S J K 2	
 <u>Orp-HC16</u> 2D3-1F5-2A9		E V Q L V Q S G G G G L V H P P G S S I R L S C A G G G F T S S Y A M H V R Q A P G K G L E V N A V I M D G S N K Y A D S V K G R E F T I S R D N A K S I L Y Q M N S L R A E D N A V Y C A R					
1E6-3D10		E V Q L V Q S G G G G L V H P P G S S I R L S C A G G G F T S S Y A M H V R Q A P G K G L E V N A V I M D G S N K Y A D S V K G R E F T I S R D N A K S I L Y Q M N S L R A E D N A V Y C A R			S	E P E D I L G Y S P	
		E V Q L V Q S G G G G L V H P P G S S I R L S C A G G G F T S S Y A M H V R Q A P G K G L E V N A V I M D G S N K Y A D S V K G R E F T I S R D N A K S I L Y Q M N S L R A E D N A V Y C A R				D3-9	

**Fig. 5**

## Human $V_H$ CDR Consensus Sequences

3D6-2F4 VH CDR1 (SEQ ID NO: 5):	IYGMH
3D6-4C8 VH CDR1 (SEQ ID NO: 17):	IYGMH
3G9-2D2 VH CDR1 (SEQ ID NO: 29):	NYGMY
5A8-1F1 VH CDR1 (SEQ ID NO: 41):	TYGMH
3C7-3A3, VH CDR1 (SEQ ID NO: 53):	SYNMH
2D3-1F5-2A9, VH CDR1 (SEQ ID NO: 65):	NYAMH
1E6-3D10 VH CDR1: (SEQ ID NO: 71):	SYAMH
5C3-2-3F6 VH CDR1 (SEQ ID NO: 77):	SYNMH
5D12-5G1 VH CDR1 (SEQ ID NO: 89):	SYGMH
VH CDR1 CONSENSUS (SEQ ID NO: 97): (I, N, T, S) Y (G, N, A) M (H, Y)	
3D6-2F4 VH CDR2 (SEQ ID NO: 6):	VIWYDGSNKKYADSVKG
3D6-4C8 VH CDR2 (SEQ ID NO: 18):	VIWYDGSNKKYADSVKG
3G9-2D2, VH CDR2 (SEQ ID NO: 30):	VIWYDGSNKKYADSVKG
5A8-1F1, VH CDR2 (SEQ ID NO: 42):	VIWYDGGNKKYADSVKG
3C7-3A3, VH CDR2 (SEQ ID NO: 54):	FIWYDGSNKKYGDSVKG
2D3-1F5-2A9, VH CDR2 (SEQ ID NO: 66):	TIGIGGGIPYAA-DSVKG
1E6-3D10 VH CDR2 (SEQ ID NO: 72):	AIGTGGGTYYY-DSVKG
5C3-2-3F6 VH CDR2 (SEQ ID NO: 78):	VIWYDGSNKKYGDSVKG
5D12-5G1 VH CDR2 (SEQ ID NO: 90):	VIWYDGSNKKYADSVKG
VH CDR2 CONSENSUS (SEQ ID NO: 98): (V, I, F, T, A) I (W, G) (Y, T) (D, G)	
G (S, G, Y) (N, T) (K, P) Y (Y, A, V) (A, G, -) D S V K G	
3D6-2F4 VH CDR3 (SEQ ID NO: 7):	APHEDY
3D6-4C8 VH CDR3 (SEQ ID NO: 19):	APHEDY
3G9-2D2, VH CDR3 (SEQ ID NO: 31):	DLMGWYFDY
5A8-1F1, VH CDR3 (SEQ ID NO: 43):	DEYWYIFDL
3C7-3A3, VH CDR3 (SEQ ID NO: 55):	EELGIGWYFDL
2D3-1F5-2A9, VH CDR3 (SEQ ID NO: 67):	SAFDV
1E6-3D10 VH CDR3 (SEQ ID NO: 73):	EPFYDILIGSPYFDY
5C3-2-3F6 VH CDR3 (SEQ ID NO: 79):	EELGIGWYFDL
5D12-5G1 VH CDR3 (SEQ ID NO: 91):	GPPRYIFDL
VH CDR3 (CORE) CONSENSUS (SEQ ID NO: 99): (A, G, Y, S, P, -) (P, W, S, R) (Y, A, H) F D (Y, L, V)	

(Where “-“ denotes option of no amino acid residue present at that position)

## Fig. 6


## Human VL CDR Consensus Sequences

3D6-2F4 VL CDR1 (SEQ ID NO: 11) :	RASQGISSWLA
3D6-4C8 VL CDR1 (SEQ ID NO: 23) :	RASQGISSALA
3G9-2D2, VL CDR1 (SEQ ID NO: 35) :	RASQSVSSYIA
5A8-1F1, VL CDR1 (SEQ ID NO: 47) :	RASQSVSSYIA
3C7-3A3, VL CDR1 (SEQ ID NO: 59) :	RASQSVSSYIA
5C3-2-3F6 VL CDR1 (SEQ ID NO: 83) :	RASQGISSALA
 VL CDR1 CONSENSUS (SEQ ID NO: 100) : R A S Q (S, G) (I, V) S S (Y, W, A) L A	
3D6-2F4 VL CDR2 (SEQ ID NO: 12) :	AASSLQS
3D6-4C8 VL CDR2 (SEQ ID NO: 24) :	DASSLES
3G9-2D2, VL CDR2 (SEQ ID NO: 36) :	DASN RAT
5A8-1F1, VL CDR2 (SEQ ID NO: 48) :	DASN RAT
3C7-3A3, VL CDR2 (SEQ ID NO: 60) :	DASN RAT
5C3-2-3F6 VL CDR2 (SEQ ID NO: 84) :	DASSLES
 VL CDR2 CONSENSUS (SEQ ID NO: 101) : (D, A) A S (N, S) (R, L) (A, Q, E) (T, S)	
3D6-2F4 VL CDR3 (SEQ ID NO: 13) :	QQNSYPYT
3D6-4C8 VL CDR3 (SEQ ID NO: 25) :	QQFNSYPLT
3G9-2D2, VL CDR3 (SEQ ID NO: 37) :	QQRRNWPLT
5A8-1F1, VL CDR3 (SEQ ID NO: 49) :	QQRRT----
3C7-3A3, VL CDR3 (SEQ ID NO: 61) :	QQRRT----
5C3-2-3F6 VL CDR3 (SEQ ID NO: 85) :	QQFNSYPH-
 VL CDR3 CONSENSUS (SEQ ID NO: 102) : Q Q (R, Y, F) (R, N) (T, S, N) (Y, W, -) (P, -) (Y, L, H, -) (T, -)	

(Where "—" denotes option of no amino acid residue present at that position)

# Fig. 7

Example of anti-DEC-205/antigen fusion  
APC targeted vaccine construct (schematic representation)



*Fig. 8*




Fig. 9A




Fig. 9B