

(51) International Patent Classification:

B01J 23/46 (2006.01)

(21) International Application Number:

PCT/US2012/040608

(22) International Filing Date:

1 June 2012 (01.06.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/492,834

3 June 2011 (03.06.2011)

US

(71) Applicant (for all designated States except US): **EMORY UNIVERSITY** [US/US]; Office of Technology Transfer, 1599 Clifton Road NE, 4th Floor, Atlanta, Georgia 30322 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DAVIES, Huw M. L.** [US/US]; 1802 Berkshire Eve Drive, Duluth, Georgia 30097 (US). **HANSEN, Jørn H** [US/US]; Office of Technology Transfer, 1599 Clifton Road NE, 4th Floor, Atlanta, Georgia 30322 (US). **QIN, Changming** [CN/US]; 1131 Highland Lake Circle, Decatur, Georgia 30033 (US).

(74) Agents: **MASON, James C.** et al.; Emory University, Office of Technology Transfer, 1599 Clifton Road NE, 4th Floor, Atlanta, Georgia 30322 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: DIRHODIUM CATALYST COMPOSITIONS AND SYNTHETIC PROCESSES RELATED THERETO





FIGURE 1



(57) Abstract: This disclosure relates to compositions comprising dirhodium catalysts and uses related thereto, e.g., in enantioselective transformations of donor/acceptor carbenoids. In certain embodiments, the dirhodium catalyst comprises a cyclopropyl ring substituted with a carboxylic acid ligand. In certain embodiments, the disclosure relates to compositions comprising a compound of the following formula, or salts thereof wherein, R1, R2, and R3 are defined herein.

**Declarations under Rule 4.17:**

— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

Published:

— *without international search report and to be republished upon receipt of that report (Rule 48.2(g))*

DIRHODIUM CATALYST COMPOSITIONS AND SYNTHETIC PROCESSES RELATED THERETO

STATEMENT REGARDING FEDERALLY FUNDED RESEARCH

This invention was made with government support under Grants CHE 0750273 awarded by the National Science Foundation. The government has certain rights in the invention.

CROSS REFERENCE TO RELATED APPLICATIONS


This application claims priority to U.S. provisional application number 61/492,834 filed June 3, 2011, hereby incorporated by reference in its entirety.

BACKGROUND

Dirhodium catalysts have played a prominent role in the metal-catalyzed reactions of diazo compounds. Effort has been made to prepare various chiral dirhodium tetracarboxylate and dirhodium tetracarboxamidate catalysts. A series of proline-based dirhodium catalysts, such as $\text{Rh}_2(\text{S-DOSP})_4$ (See Figure 1, compound 1), have been prepared and examined for their capacity to induce asymmetric induction in the reactions of donor/acceptor carbenoids. See Davies et al., JACS, 1996, 118:6897-6907. The asymmetric induction in these catalyzed reactions is sensitive to solvent and substrate structure. Asymmetric induction with $\text{Rh}_2(\text{S-DOSP})_4$ as catalyst requires the use of a methyl ester as the acceptor group in the carbenoid and nonpolar solvents such as hydrocarbons. The phthalimidocarboxylate $\text{Rh}_2(\text{S-PTAD})_4$ (See Figure 1, compound 2) and the bridged dicarboxylate catalyst, $\text{Rh}_2(\text{S-biTISP})_2$ (See Figure 1, compound 3) may be prepared by relatively tedious synthetic routes. Thus, there is a need for improved catalysts.


SUMMARY

This disclosure relates to compositions comprising dirhodium catalysts and uses related thereto, e.g., in enantioselective transformations of donor/acceptor carbenoids. In certain embodiments, the dirhodium catalyst comprises a cyclopropyl ring substituted with a carboxylic acid ligand. In certain embodiments, the disclosure relates to compositions comprising a compound of the following formula,



or derivatives or salts thereof wherein, R^1 , R^2 , and R^3 are defined herein.

In certain embodiments, the disclosure relates to compositions comprising $Rh_2(S$ -
 5 biTISP) $_2$, $Rh_2(R$ -biTISP) $_2$, or a compound of formula



In certain embodiments, the disclosure relates to methods of preparing dirhodium
 10 catalysts or intermediates disclosed herein by mixing starting materials with catalysts
 under conditions such that the catalysts or intermediates are formed.

In certain embodiments, the disclosure contemplates enantioselective reactions of
 donor/acceptor carbenoids, such as cyclopropanations, formal [4 + 3] cycloadditions, C-H
 functionalizations, and ylide transformations comprising mixing a compound comprising a
 15 carbenoid precursor, e.g., a diazo compound, and catalysts disclosed herein and reactive
 compounds under conditions such that a synthetic compound is formed.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 illustrates certain chiral dirhodium catalysts.

20 Figure 2 illustrates three distinct ligand orientations used to rationalize
 enantioselectivity in dirhodium carboxylate-catalyzed reactions.

Figure 3 illustrates $Rh_2(R$ -BTPCP) $_4$ (axial ligands omitted for clarity) indicated by
 data from an X-ray crystal structure study.

Figure 4 illustrates the calculated structure catalytically active sites of $Rh_2(R$ -
 25 BTPCP) $_4$ (X-ray) and $Rh_2(S$ -PTTL) $_4$.

Figure 5 illustrates data from a DFT model of s-cis and s-trans carbene conformers.

Figure 6 illustrates ONIOM partitioning of $Rh_2(R$ -BTPCP) $_4$ -carbene complex.

Figure 7 illustrates the calculated lowest-energy conformation of s-trans carbene, top view (left) and side view (right). Hydrogens atoms were omitted for clarity.

Figure 8 illustrates a predictive stereochemical model for $Rh_2(R-BTPCP)_4$ -catalyzed transformation.

5 Figure 9 illustrates additional embodiments of the disclosure.

DETAILED DESCRIPTION

Terms

Unless defined otherwise, all technical and scientific terms used herein have the
10 same meaning as is commonly understood by one of skill in the art. The following definitions are provided to help interpret the disclosure and claims of this application. In the event a definition in this section is not consistent with definitions elsewhere, the definition set forth in this section will control.

As used herein, the term "derivative" refers to a structurally similar compound that
15 retains sufficient functional attributes of the identified analogue. The derivative may be structurally similar because it is lacking one or more atoms, substituted, a salt, in different hydration/oxidation states, or because one or more atoms within the molecule are switched, such as, but not limited to, replacing an oxygen atom with a sulfur atom or replacing an amino group with a hydroxy group. Derivatives may be prepared by any
20 variety of synthetic methods or appropriate adaptations presented in synthetic or organic chemistry text books, such as those provided in March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure, Wiley, 6th Edition (2007) Michael B. Smith or Domino Reactions in Organic Synthesis, Wiley (2006) Lutz F. Tietze, hereby incorporated by reference.

25 The term "substituted" refers to a molecule wherein at least one hydrogen atom is replaced with a substituent. When substituted, one or more of the groups are "substituents." The molecule may be multiply substituted. In the case of an oxo substituent ("=O"), two hydrogen atoms are replaced. Example substituents within this context may include halogen, hydroxy, alkyl, alkoxy, nitro, cyano, oxo, carbocyclyl, carbocycloalkyl,
30 heterocarbocyclyl, heterocarbocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, -NRaRb, -NRaC(=O)Rb, -NRaC(=O)NRaNRb, -NRaC(=O)ORb, -NRaSO₂Rb, -C(=O)Ra, -C(=O)ORa, -C(=O)NRaRb, -OC(=O)NRaRb, -ORa, -SRa, -SORa, -S(=O)₂Ra, -OS(=O)₂Ra and -S(=O)₂ORa. Ra and Rb in this context may be the same or different and independently hydrogen, halogen hydroxy, alkyl, alkoxy, alkyl, amino, alkylamino,

dialkylamino, carbocyclyl, carbocycloalkyl, heterocarbocyclyl, heterocarbocycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl.

When used in reference to compound(s) disclosed herein, "salts" refer to derivatives of the disclosed compound(s) where the parent compound is modified making acid or base salts thereof. Examples of salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines, alkylamines, or dialkylamines; alkali or organic salts of acidic residues such as carboxylic acids; and the like.

As used herein, "alkyl" means a noncyclic straight chain or branched, unsaturated or saturated hydrocarbon such as those containing from 1 to 10 carbon atoms. Representative saturated straight chain alkyls include methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, n-septyl, n-octyl, n-nonyl, and the like; while saturated branched alkyls include isopropyl, sec-butyl, isobutyl, tert-butyl, isopentyl, and the like. Unsaturated alkyls contain at least one double or triple bond between adjacent carbon atoms (referred to as an "alkenyl" or "alkynyl", respectively). Representative straight chain and branched alkenyls include ethylenyl, propylenyl, 1-butenyl, 2-butenyl, isobutylenyl, 1-pentenyl, 2-pentenyl, 3-methyl-1-butenyl, 2-methyl-2-butenyl, 2,3-dimethyl-2-butenyl, and the like; while representative straight chain and branched alkynyls include acetylenyl, propynyl, 1-butyne, 2-butyne, 1-pentyne, 2-pentyne, 3-methyl-1-butyne, and the like.

Non-aromatic mono or polycyclic alkyls are referred to herein as "carbocycles" or "carbocyclyl" groups. Representative saturated carbocycles include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and the like; while unsaturated carbocycles include cyclopentenyl and cyclohexenyl, and the like.

"Heterocarbocycles" or heterocarbocyclyl" groups are carbocycles which contain from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and sulfur which may be saturated or unsaturated (but not aromatic), monocyclic or polycyclic, and wherein the nitrogen and sulfur heteroatoms may be optionally oxidized, and the nitrogen heteroatom may be optionally quaternized. Heterocarbocycles include morpholinyl, pyrrolidinonyl, pyrrolidinyl, piperidinyl, hydantoinyl, valerolactamyl, oxiranyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydropyridinyl, tetrahydroprimidinyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, and the like.

"Aryl" means an aromatic carbocyclic monocyclic or polycyclic ring such as phenyl or naphthyl. Polycyclic ring systems may, but are not required to, contain one or more non-aromatic rings, as long as one of the rings is aromatic.

As used herein, "heteroaryl" or "heteroaromatic" refers an aromatic heterocarbocycle having 1 to 4 heteroatoms selected from nitrogen, oxygen and sulfur, and containing at least 1 carbon atom, including both mono- and polycyclic ring systems. Polycyclic ring systems may, but are not required to, contain one or more non-aromatic rings, as long as one of the rings is aromatic. Representative heteroaryls are furyl, benzofuranyl, thiophenyl, benzothiophenyl, pyrrolyl, indolyl, isoindolyl, azaindolyl, pyridyl, quinoliny, isoquinoliny, oxazolyl, isooxazolyl, benzoxazolyl, pyrazolyl, imidazolyl, benzimidazolyl, thiazolyl, benzothiazolyl, isothiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, cinnoliny, phthalazinyl, and quinazolinyl. It is contemplated that the use of the term "heteroaryl" includes N-alkylated derivatives such as a 1-methylimidazol-5-yl substituent.

As used herein, "heterocycle" or "heterocyclyl" refers to mono- and polycyclic ring systems having 1 to 4 heteroatoms selected from nitrogen, oxygen and sulfur, and containing at least 1 carbon atom. The mono- and polycyclic ring systems may be aromatic, non-aromatic or mixtures of aromatic and non-aromatic rings. Heterocycle includes heterocarbocycles, heteroaryls, and the like.

"Alkylthio" refers to an alkyl group as defined above attached through a sulfur bridge. An example of an alkylthio is methylthio, (i.e., -S-CH₃).

"Alkoxy" refers to an alkyl group as defined above attached through an oxygen bridge. Examples of alkoxy include, but are not limited to, methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, s-butoxy, t-butoxy, n-pentoxy, and s-pentoxy. Preferred alkoxy groups are methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, s-butoxy, t-butoxy.

"Alkylamino" refers an alkyl group as defined above attached through an amino bridge. An example of an alkylamino is methylamino, (i.e., -NH-CH₃).

"Alkanoyl" refers to an alkyl as defined above attached through a carbonyl bridge (i.e., -(C=O)alkyl).

"Alkylsulfonyl" refers to an alkyl as defined above attached through a sulfonyl bridge (i.e., -S(=O)₂alkyl) such as mesyl and the like, and "Arylsulfonyl" refers to an aryl attached through a sulfonyl bridge (i.e., -S(=O)₂aryl).


"Alkylsulfamoyl" refers to an alkyl as defined above attached through a sulfamoyl bridge (i.e., -S(=O)₂NHalkyl), and an "Arylsulfamoyl" refers to an alkyl attached through a sulfamoyl bridge (i.e., -S(=O)₂NHaryl).

"Alkylsulfinyl" refers to an alkyl as defined above with the indicated number of carbon atoms attached through a sulfinyl bridge (i.e. -S(=O)alkyl).

The terms "halogen" and "halo" refer to fluorine, chlorine, bromine, and iodine.

Dirhodium catalysts

In certain embodiments, the disclosure relates to a dirhodium catalyst comprising a
 5 cyclopropyl ring substituted with a carboxylic acid ligand. In certain embodiments, the disclosure relates to compositions comprising a compound of the following formula,



10 or salts thereof wherein,

R^1 is hydrogen, alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R^1 is optionally substituted with one or more, the same or different, R^4 ;

15 R^2 is hydrogen, alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R^2 is optionally substituted with one or more, the same or different, R^4 ;


R^3 is hydrogen, alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy,
 20 carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R^3 is optionally substituted with one or more, the same or different, R^4 ;

R^4 is alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl,
 25 alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R^4 is optionally substituted with one or more, the same or different, R^5 ;

R^5 is alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R^5 is optionally
 30 substituted with one or more, the same or different, R^6 ; and

R^6 is halogen, nitro, cyano, hydroxy, trifluoromethoxy, trifluoromethyl, amino, formyl, carboxy, carbamoyl, mercapto, sulfamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxymethyl, methylamino, ethylamino, dimethylamino, diethylamino, N-methyl-N-ethylamino, acetylamino, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, N-methyl-N-ethylcarbamoyl, methylthio, ethylthio, methylsulfinyl, ethylsulfinyl, mesyl, ethylsulfonyl, methoxycarbonyl, ethoxycarbonyl, N-methylsulfamoyl, N-ethylsulfamoyl, N,N-dimethylsulfamoyl, N,N-diethylsulfamoyl, N-methyl-N-ethylsulfamoyl, carbocyclyl, aryl, or heterocyclyl.


In certain embodiments, the catalyst is in a composition with greater than 55%, 75%, 85%, 95%, 98%, or 99% enantiomeric excess of the following stereoisomers:



- In certain embodiments, R^1 is carbocyclyl, aryl, or heterocyclyl.
- In certain embodiments, R^1 is an aryl or an aromatic heterocyclyl.
- In certain embodiments, R^1 is a phenyl optionally substituted with one or more R^4 .
- In certain embodiments, R^1 is a phenyl optionally substituted with one or more halogen, alkyl, or alkoxy.
- In certain embodiments, R^1 is hydrogen.
- In certain embodiments, R^2 is carbocyclyl, aryl, or heterocyclyl.
- In certain embodiments, R^2 is an aryl or an aromatic heterocyclyl.
- In certain embodiments, R^2 is a phenyl optionally substituted with one or more R^4 .
- In certain embodiments, R^2 is a phenyl optionally substituted with one or more halogen, alkyl, or alkoxy.
- In certain embodiments, R^3 is carbocyclyl, aryl, or heterocyclyl.
- In certain embodiments, R^3 is an aryl or an aromatic heterocyclyl.
- In certain embodiments, R^3 is a phenyl optionally substituted with one or more R^4 .
- In certain embodiments, R^3 is a phenyl optionally substituted with one or more halogen, alkyl, or alkoxy.
- In certain embodiments, R^3 is a phenyl substituted with a halogen.

In certain embodiments, R^3 is a phenyl substituted with a halogen in the ortho or para position.

In certain embodiments, the disclosure relates to compositions comprising a compound,




5

wherein X is an electron withdrawing group.

In certain embodiments, the catalyst is in a composition with greater than 55%, 75%, 85%, 95%, 98%, or 99% enantiomeric excess of the following stereoisomers:


10



In certain embodiments, X is a halogen.


In certain embodiments, the disclosure relates to compositions comprising a compound

15



In certain embodiments, the catalyst is in a composition with greater than 55%, 75%, 85%, 95%, 98%, or 99% enantiomeric excess of the following stereoisomers:

20



Synthetic Processes

Many enantioselective reactions of donor/acceptor carbenoids may be utilized with dirhodium catalysts disclosed herein. In certain embodiments, the dirhodium can be used to make intermediates for pharmaceutical agents as exemplified herein. See Davies & Manning, *Nature*, 2008, 451, 417–424; Davies & Denton, *Chem. Soc. Rev.*, 2009, 38, 3061–3071; Davies, et al., *J. Am. Chem. Soc.*, 2006, 128, 2485–2490; and Davies & Walji, *Angew. Chem., Int. Ed.*, 2005, 44, 1733–1735, all hereby incorporated by reference.

In certain embodiments, the disclosure contemplates reactions of donor/acceptor carbenoids, such as cyclopropanations, formal [4 + 3] cycloadditions, C–H functionalizations, and ylide transformations comprising mixing a compound comprising a carbenoid precursor, e.g., a diazo compound, and catalysts disclosed herein and reactive compounds under conditions such that a synthetic compound is formed.

In certain embodiments, the disclosure relates to methods of making a synthetic compound comprising mixing a) a diazo compound, b) a compound with a carbon hydrogen bond, and c) a dirhodium catalyst disclosed herein, under conditions such that a synthetic compound is formed comprising a carbon to carbon bond between the diazo compound and the compound with a carbon hydrogen bond. The synthetic compound may be the result of an inter or an intra-molecular reaction.

In certain embodiments, the disclosure relates to methods of making a synthetic compound comprising mixing a) a diazo compound, b) a compound with a nitrogen hydrogen bond, and c) a dirhodium catalyst disclosed herein, under conditions such that a synthetic compound is formed comprising a carbon to nitrogen bond between the diazo compound and the compound with a nitrogen hydrogen bond. The synthetic compound may be the result of an inter or an intra-molecular reaction.

In certain embodiments, the disclosure relates to cyclopropanation reactions using catalysts disclosed herein. See Davies et al., *J. Am. Chem. Soc.* 1996, 118, 6897– 6907, hereby incorporated by reference. In certain embodiments, the disclosure relates to

- methods of making a synthetic compound comprising mixing a) a diazo compound such as vinyl diazomethane and vinyl diazoacetates optionally substituted with one or more substituents, b) a double bond compound such as an alkene or diene optionally substituted with one or more substituent, and c) a compound comprising a dirhodium catalyst
- 5 disclosed herein under conditions such that a synthetic compound is formed comprising a cyclopropyl ring and carbon to carbon bond between the diazo compound and the double bond compound.

In certain embodiments, the diazo compound has the following formula,



the double bond compound has the following formula,



and the synthetic compound has the following formula,



- 20 wherein, R^1 is alkyl, carbocyclyl, aryl, or heterocyclyl, wherein R^1 is optionally substituted with one or more, the same or different, R^4 ;

- R^2 is alkyl, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R^2 is optionally substituted with one or more,
- 25 the same or different, R^4 ;

R^3 is hydrogen, alkyl, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R^3 is optionally substituted with one or more, the same or different, R^4 ;

R⁴ is alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁴ is optionally substituted with one or more, the same or different, R⁵;

5 R⁵ is alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁵ is optionally substituted with one or more, the same or different, R⁶; and

R⁶ is halogen, nitro, cyano, hydroxy, trifluoromethoxy, trifluoromethyl, amino, formyl, carboxy, carbamoyl, mercapto, sulfamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxymethyl, methylamino, ethylamino, dimethylamino, diethylamino, N-methyl-N-ethylamino, acetylamino, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, N-methyl-N-ethylcarbamoyl, methylthio, ethylthio, methylsulfinyl, ethylsulfinyl, mesyl, ethylsulfonyl, methoxycarbonyl, ethoxycarbonyl, N-methylsulfamoyl, N-ethylsulfamoyl, N,N-dimethylsulfamoyl, N,N-diethylsulfamoyl, N-methyl-N-ethylsulfamoyl, carbocyclyl, aryl, or heterocyclyl.

In certain embodiments, R¹ is alkyl optionally substituted with one or more, the same or different, R⁴.


20 In certain embodiments, R² is an electron donating group selected from cyano, alkenyl, alkynyl, formyl, carbamoyl, carboxy, alkylsulfinyl, alkylsulfonyl, or arylsulfonyl, aryl, or an aromatic heterocyclyl optionally substituted with one or more, the same or different, R⁴.

In certain embodiments, R² is aryl or an aromatic heterocyclyl optionally substituted with one or more, the same or different, R⁴.


25 In certain embodiments, R³ is an electron donating group selected from cyano, alkenyl, alkynyl, formyl, carbamoyl, carboxy, alkylsulfinyl, alkylsulfonyl, or arylsulfonyl, aryl, or an aromatic heterocyclyl optionally substituted with one or more, the same or different, R⁴.

30 In certain embodiments, R³ is aryl or an aromatic heterocyclyl optionally substituted with one or more, the same or different, R⁴.

In certain embodiments, the synthetic product is in a composition with greater than 55%, 75%, 85%, 95%, 98%, or 99% enantiomeric excess of the following stereoisomers:



In certain embodiments, the diazo compound has the following formula,




5

the double bond compound has the following formula,



10


and the synthetic compound has the following formula,



or salts thereof wherein


- 15 R^1 is phenyl optionally substituted with one or more substituent and
 R^2 is phenyl optionally substituted with one or more substituent.

In certain embodiments, the synthetic product is in a composition with greater than 55%, 75%, 85%, 95%, 98%, or 99% enantiomeric excess of the following stereoisomers:



20

In certain embodiments, the diazo compound has the following formula,




5 the double bond compound has the following formula,




and the synthetic compound has the following formula,

10




wherein Ar^1 and Ar^2 are each, the same or different, aryl optionally substituted with one or more substituents.

In certain embodiments, the synthetic product is in a composition with greater than
 15 55%, 75%, 85%, 95%, 98%, or 99% enantiomeric excess of the following stereoisomers:




In certain embodiments, the disclosure contemplates the use of dirhodium catalysts
 20 disclosed herein in [4 + 3] cycloaddition reactions, e.g., mixing a diene and a
 vinyldiazoacetate and a dirhodium catalyst disclosed herein under conditions such that a
 cyclic compound is formed. See Davies et al, J. Am. Chem. Soc. 1998, 120, 3326– 3331
 and Reddy & Davies, J. Am. Chem. Soc., 2007, 129 (34), 10312–10313.

In certain embodiments, the diazo compound is a vinyl diazoacetate of the following formula,




5

the diene compound has the following formula,



10

and the synthetic compound has the following formula,



wherein, R¹ is alkyl, carbocyclyl, aryl, or heterocyclyl, wherein R¹ is optionally substituted with one or more, the same or different, R⁹;

R² is hydrogen, alkyl, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R² is optionally substituted with one or more, the same or different, R⁹;

R³ is hydrogen, alkyl, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R³ is optionally substituted with one or more, the same or different, R⁹;

R⁴ is hydrogen, alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl,

alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁴ is optionally substituted with one or more, the same or different, R⁹;

R⁵ is hydrogen, alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁵ is optionally substituted with one or more, the same or different, R⁹;

R⁶ is hydrogen, alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁶ is optionally substituted with one or more, the same or different, R⁹;


R⁷ is hydrogen, alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁷ is optionally substituted with one or more, the same or different, R⁹;

R⁸ is hydrogen, alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁸ is optionally substituted with one or more, the same or different, R⁹;

R⁹ is alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁹ is optionally substituted with one or more, the same or different, R¹⁰; and

R¹⁰ is halogen, nitro, cyano, hydroxy, trifluoromethoxy, trifluoromethyl, amino, formyl, carboxy, carbamoyl, mercapto, sulfamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxymethyl, methylamino, ethylamino, dimethylamino, diethylamino, N-methyl-N-ethylamino, acetylamino, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, N-methyl-N-ethylcarbamoyl, methylthio, ethylthio, methylsulfinyl, ethylsulfinyl, mesyl, ethylsulfonyl, methoxycarbonyl, ethoxycarbonyl, N-methylsulfamoyl, N-ethylsulfamoyl, N,N-dimethylsulfamoyl, N,N-diethylsulfamoyl, N-methyl-N-ethylsulfamoyl, carbocyclyl, aryl, or heterocyclyl.

In certain embodiments, the synthetic product is in a composition with greater than 55%, 75%, 85%, 95%, 98%, or 99% enantiomeric excess of the following stereoisomers:



In certain embodiments, R^1 is alkyl optionally substituted with one or more, the same or different, R^9 .

- 5 In certain embodiments, R^2 is an electron donating group selected from cyano, alkenyl, alkynyl, formyl, carbamoyl, carboxy, alkylsulfinyl, alkylsulfonyl, or arylsulfonyl, aryl, or an aromatic heterocyclyl optionally substituted with one or more, the same or different, R^9 .

- 10 In certain embodiments, R^2 is aryl or an aromatic heterocyclyl optionally substituted with one or more, the same or different, R^9 .

In certain embodiments, R^3 is an electron donating group selected from cyano, alkenyl, alkynyl, formyl, carbamoyl, carboxy, alkylsulfinyl, alkylsulfonyl, or arylsulfonyl, aryl, or an aromatic heterocyclyl optionally substituted with one or more, the same or different, R^9 .

- 15 In certain embodiments, R^3 is aryl or an aromatic heterocyclyl optionally substituted with one or more, the same or different, R^9 .

In certain embodiments, R^4 is hydrogen.


In certain embodiments, R^5 is hydrogen.

In certain embodiments, R^6 is hydrogen.

- 20 In certain embodiments, R^7 is hydrogen.


In certain embodiments, R^8 is hydrogen.

In certain embodiments, the diazo compound is a vinyl diazoacetate of the following formula,




25

the diene compound has the following formula,




and the synthetic compound has the following formula,

5




Wherein R is phenyl optionally substituted with one or more substituent.

In certain embodiments, the synthetic product is in a composition with greater than
 10 55%, 75%, 85%, 95%, 98%, or 99% enantiomeric excess of the following stereoisomers:




In certain embodiments, the diazo compound has the following formula,

15




the double bond compound has the following formula,



20

and the synthetic compound has the following formula,



wherein, R¹ is alkyl, carbocyclyl, aryl, or heterocyclyl, wherein R¹ is optionally substituted with one or more, the same or different, R⁷;

5 R² is alkyl, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R² is optionally substituted with one or more, the same or different, R⁷;

10 R³ is hydrogen, alkyl, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R³ is optionally substituted with one or more, the same or different, R⁷;

15 R⁴ is hydrogen, alkyl, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁴ is optionally substituted with one or more, the same or different, R⁷;

20 R⁵ is hydrogen, alkyl, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁵ is optionally substituted with one or more, the same or different, R⁷; or R⁴ and R⁵ form a ring selected from a carbocyclyl, aryl, or heterocyclyl wherein the ring is optionally substituted with one or more, the same or different, R⁷

25 R⁶ is hydrogen, alkyl, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁶ is optionally substituted with one or more, the same or different, R⁷;

30 R⁷ is alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁷ is optionally substituted with one or more, the same or different, R⁸;

R⁸ is alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁸ is optionally substituted with one or more, the same or different, R⁹; and

- 5 R⁹ is halogen, nitro, cyano, hydroxy, trifluoromethoxy, trifluoromethyl, amino, formyl, carboxy, carbamoyl, mercapto, sulfamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxymethyl, methylamino, ethylamino, dimethylamino, diethylamino, N-methyl-N-ethylamino, acetylamino, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, N-methyl-N-ethylcarbamoyl, methylthio, 10 ethylthio, methylsulfinyl, ethylsulfinyl, mesyl, ethylsulfonyl, methoxycarbonyl, ethoxycarbonyl, N-methylsulfamoyl, N-ethylsulfamoyl, N,N-dimethylsulfamoyl, N,N-diethylsulfamoyl, N-methyl-N-ethylsulfamoyl, carbocyclyl, aryl, or heterocyclyl.

In certain embodiments, R¹ is alkyl optionally substituted with one or more, the same or different, R⁷.


- 15 In certain embodiments, R² is an electron donating group selected from cyano, alkenyl, alkynyl, formyl, carbamoyl, carboxy, alkylsulfinyl, alkylsulfonyl, or arylsulfonyl, aryl, or an aromatic heterocyclyl optionally substituted with one or more, the same or different, R⁷.

- In certain embodiments, R² is aryl or an aromatic heterocyclyl optionally substituted with one or more, the same or different, R⁷.
- 20

In certain embodiments, R³ is an electron donating group selected from cyano, alkenyl, alkynyl, formyl, carbamoyl, carboxy, alkylsulfinyl, alkylsulfonyl, or arylsulfonyl, aryl, or an aromatic heterocyclyl optionally substituted with one or more, the same or different, R⁷.


- 25 In certain embodiments, R³ is hydrogen or alkyl optionally substituted with one or more, the same or different, R⁷.

In certain embodiments, the disclosure relates to diazo compound of the following formula:




- 30 wherein EDG is an electron withdrawing group and EWG is an electron donating group. Such a diazo compound may react with a compound with a C-H bond or a N-H bond as further exemplified below. See Davies & Morton, Chem. Soc. Rev., 2011, 40, 1857–1869, hereby incorporated by reference.

In certain embodiments, the method comprises mixing the diazo compound with a compound of the following formula:



under conditions such that a synthetic compound of the following formula is formed,



5

wherein X is carbon or nitrogen, wherein R³ is absent if X is nitrogen,

R¹ is alkyl, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R¹ is optionally substituted with one or more, the same or different, R⁴;

10

R² is alkyl, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R² is optionally substituted with one or more, the same or different, R⁴;

15

or R¹ and R² form a carbocyclyl, aryl, or heterocyclyl ring optionally substituted with one or more, the same or different, R⁴; or when X is carbon R¹, R², and R³ form a multicyclic carbocyclyl;

20

R³ is hydrogen, alkyl, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R³ is optionally substituted with one or more, the same or different, R⁴;

25

R⁴ is alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁴ is optionally substituted with one or more, the same or different, R⁵;

30

R⁵ is alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁵ is optionally substituted with one or more, the same or different, R⁶; and
R⁶ is halogen, nitro, cyano, hydroxy, trifluoromethoxy, trifluoromethyl, amino, formyl, carboxy, carbamoyl, mercapto, sulfamoyl, methyl, ethyl, methoxy, ethoxy, acetyl,


acetoxy, methylamino, ethylamino, dimethylamino, diethylamino, N-methyl-N-ethylamino, acetylamino, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, N-methyl-N-ethylcarbamoyl, methylthio, ethylthio, methylsulfinyl, ethylsulfinyl, mesyl, ethylsulfonyl, methoxycarbonyl, ethoxycarbonyl, N-methylsulfamoyl, N-ethylsulfamoyl, N,N-dimethylsulfamoyl, N,N-diethylsulfamoyl, N-methyl-N-ethylsulfamoyl, carbocyclyl, aryl, or heterocyclyl.

In certain embodiments, the disclosure contemplates the use of dirhodium catalysts disclosed herein for the intra-molecular insertion reactions for the formation of carbon to carbon bonds. In certain embodiment, disclosure contemplates a method of making an ring structure comprising mixing a dirhodium catalyst disclosed herein and a compound comprising a diazo group and the same compound comprising a C-H bond configure about the molecule to form a five or six membered ring.

In certain embodiments, the disclosure contemplates the use of dirhodium catalysts disclosed herein in C-H aminations. In certain embodiments, the disclosure contemplates the use of dirhodium catalysts disclosed herein for the insertion reactions for the formation of carbon to nitrogen bonds, e.g., inter and intra-molecular C-H aminations. See Reddy & Davies, Org. Lett., 2006, 8, 5013– 5016, hereby incorporated by reference.


In certain embodiments, the disclosure relates to a method of making an synthetic compound comprising mixing a compound with an aromatic compound, $\text{PhI}(\text{OAc})_2$, NsNH_2 (4-nitrobenzenesulfonamide), and a dirhodium catalysts disclosed herein under conditions such that a synthetic compound with a C-N bond is formed.

In certain embodiments, the aromatic compound is a has the following formula



25

and the synthetic compound is



30

wherein, Y is 4-nitrobenzenesulfonamide or 2-nitrobenzenesulfonamide;

R^1 is aryl or aromatic heterocyclyl wherein R^1 is optionally substituted with one or more, the same or different, R^3 ;

R^2 is hydrogen alkyl, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R^2 is optionally substituted with one or more, the same or different, R^3 ;


- 5 or R^1 and R^2 together with the attached atoms form a carbocyclyl, aryl, or heterocyclyl optionally substituted with one or more, the same or different, R^3 ;

R^3 is hydrogen, alkyl, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R^3 is optionally substituted with one or more, the same or different, R^4 ;


R^4 is hydrogen, alkyl, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R^4 is optionally substituted with one or more, the same or different, R^5 ;

15 R^5 is halogen, nitro, cyano, hydroxy, trifluoromethoxy, trifluoromethyl, amino, formyl, carboxy, carbamoyl, mercapto, sulfamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxymethyl, methylamino, ethylamino, dimethylamino, diethylamino, N-methyl-N-ethylamino, acetylamino, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, N-methyl-N-ethylcarbamoyl, methylthio, ethylthio, methylsulfinyl, ethylsulfinyl, mesyl, ethylsulfonyl, methoxycarbonyl, ethoxycarbonyl, N-methylsulfamoyl, N-ethylsulfamoyl, N,N-dimethylsulfamoyl, N,N-diethylsulfamoyl, N-methyl-N-ethylsulfamoyl, carbocyclyl, aryl, or heterocyclyl.

In certain embodiments, the synthetic product is in a composition with greater than 55%, 75%, 85%, 95%, 98%, or 99% enantiomeric excess of the following stereoisomers:



In certain embodiments, the disclosure relates to a method of making Rasagiline comprising the step of mixing 2,3-dihydro-1H-indene, $\text{PhI}(\text{OAc})_2$, YNH_2 , MgO , and a dirhodium catalyst disclosed herein, wherein Y is 4-nitrobenzenesulfonamide or 2-nitrobenzenesulfonamide, under conditions such that a compound of the following formula is formed,



Reaction of this product with propargyl bromide in potassium carbonate provides the N substituted propargyl product. Removal of the nitrobenzenesulfonamide with DBU and HSCH₂CH₂OH provides Rasagiline. See Reddy & Davies, Org. Lett., 2006, 8, 5013–

5 5016, hereby incorporated by reference.


EXPERIMENTAL

The following is intended to provide examples on methods of making and using
10 embodiments of the disclosure. It is not intended to limit the scope.

*D*₂-Symmetric Dirhodium Catalyst Derived from a 1,2-Triarylcyclopropanecarboxylate Ligand


Donor/acceptor carbenoids are also capable of generating highly substituted
15 cyclopropanes with enantioselectivity. Cyclopropane carboxylic acids are useful ligands for dirhodium catalysts. The cyclopropane ring limits the possible conformations of the ligands and in this way addresses certain limitations observed with the Rh₂(S-DOSP)₄ catalyst. See Figure 1.

To evaluate the utility of cyclopropanecarboxylate catalysts, the 1,2-diphenylcyclo
20 propane carboxylate complex 8, Rh₂(R-DPCP)₄, was prepared.



The selectivity of Rh₂(R-DPCP)₄ was evaluated in the standard cyclopropanation reaction
25 between styrene 6 and styryldiazoacetate 9 using dichloromethane as solvent. The


styrylcyclopropane 10 was formed in 81% yield and with high diastereoselectivity (>20: 1 dr). The asymmetric induction exhibited by $\text{Rh}_2(\text{R-DPCP})_4$ was 11% ee.



5

Experiments to identify a more effective cyclopropanecarboxylate ligand were instituted. Even though the cyclopropanecarboxylate in 8 has two stereogenic centers, the two phenyl rings are on the opposite side of the ring to the carboxylate. A ligand with an additional phenyl ring syn- to the carboxylate was made. The


- 10 triarylcyclopropanecarboxylate, was prepared from $\text{Rh}_2(\text{R-DOSP})_4$ catalyzed cyclopropanation reactions between 1,1-diphenylethylene (11) and para-bromophenyl diazoacetate 12.



15

Cyclopropanation proceeds with enantioselectivity. The cyclopropane ester was converted to the corresponding carboxylic acid by $t\text{BuOK}$ in dimethyl sulfoxide without erosion of enantioselectivity and was enriched to 99% ee after recrystallization. Standard ligand exchange conditions generated $\text{Rh}_2(\text{R-BTPCP})_4$ (4) in 63% yield as a green solid.

- 20 With $\text{Rh}_2(\text{R-BTPCP})_4$ in hand, its efficiency was examined in the standard reaction between the styryldiazoacetate 9 and styrene 6.

Table 1. Initial Evaluation of $\text{Rh}_2(\text{R-BTPCP})_4$ - and $\text{Rh}_2(\text{R-DOSP})_4$ -Catalyzed

5 Cyclopropanationa

entry	solvent	temp. (°C)	time (h)	$\text{Rh}_2(\text{R-BTPCP})_4$		$\text{Rh}_2(\text{R-DOSP})_4$	
				yield (%)	ee (%)	yield (%)	ee (%)
1	pentane	23	1.5	82	84 ^b	85	92
2	CH_2Cl_2	23	1.5	86	91	80	81
3	CH_2Cl_2	0	2.0	72	92 ^c	81	84
4	CH_2Cl_2	-78	12	71	91	78	87
5 ^d	CH_2Cl_2	23	24	77	92	69	81
6 ^e	CH_2Cl_2	23	60	46	92	38	81

a- Standard conditions: 9 (0.4 mmol) was added to a solution of Rh(II) catalyst and styrene (2.0 mmol) under argon over 1 h. Reported yields were obtained after chromatographic purification. dr was determined by ^1H NMR prior to chromatography, and ee was determined by chiral HPLC. Absolute configuration of 10 was assigned according to previous studies.

b- 89% ee was obtained when $\text{Rh}_2(\text{S-BTPCP})_4$ was used.

c- $\text{Rh}_2(\text{S-BTPCP})_4$ was used.

d- 0.1 mol % catalyst.

e- 0.01 mol % catalyst.

The influence of the syn-phenyl group in the ligand was dramatic as the reaction in dichloromethane at 23 °C generated the cyclopropane 10 in 91% ee, which is better than

the same reaction catalyzed by $\text{Rh}_2(\text{R-DOSP})_4$ (81% ee), (Table 1, entry 2). Further experiments comparing $\text{Rh}_2(\text{R-BTPCP})_4$ and $\text{Rh}_2(\text{R-DOSP})_4$ are summarized in Table 1. Temperature did not influence the enantioselectivity of the $\text{Rh}_2(\text{R-BTPCP})_4$ -catalyzed reaction between 9 and styrene 6 as 10 was produced in 91% ee over a reaction

- 5 temperature range from 23 to -78°C (Table 1, entries 2, 4). Furthermore, decreasing the catalyst loading from 1 to 0.01 mol % did not alter the enantioselectivity (Table 1, entries 5, 6), but the reaction did not go to completion at the 0.01 mol % catalyst loading.

High asymmetric induction with $\text{Rh}_2(\text{S-DOSP})_4$ is obtained when the electron-withdrawing group is a methyl ester. Changing the methyl ester to a *tert*-butyl ester caused a drop in the enantioselectivity. Therefore, a study was undertaken to determine
10 how $\text{Rh}_2(\text{R-BTPCP})_4$ -catalyzed reactions responded to the size of the ester. See Table 2.




Table 2. Effect of Ester Size on Asymmetric Cyclopropanation

entry	starting material	R	catalyst	product	yield (%)	ee (%)
1	9	Me	$\text{Rh}_2(\text{R-BTPCP})_4$	10	86	91
2	13a	Et	$\text{Rh}_2(\text{R-BTPCP})_4$	14a	88	93
3	13b	<i>i</i> -Pr	$\text{Rh}_2(\text{R-BTPCP})_4$	14b	89	96
4	13c	<i>t</i> -Bu	$\text{Rh}_2(\text{R-BTPCP})_4$	14c	87	95
5	9	Me	$\text{Rh}_2(\text{R-DOSP})_4$	10	80	81
6	13a	Et	$\text{Rh}_2(\text{R-DOSP})_4$	14a	84	75
7	13b	<i>i</i> -Pr	$\text{Rh}_2(\text{R-DOSP})_4$	14b	82	55
8	13c	<i>t</i> -Bu	$\text{Rh}_2(\text{R-DOSP})_4$	14c	80	16

15

a- Standard conditions: Diazo compound (0.4 mmol) was added to a solution of Rh(II) catalyst and styrene (2.0 mmol) in dichloromethane under argon over 1 h.

Reported yields were obtained after chromatographic purification. *dr* was determined by ^1H NMR prior to chromatography, and *ee* was determined by chiral HPLC.

5 $\text{Rh}_2(\text{R-BTPCP})_4$ exhibited good tolerance to ester size, and the enantioselectivity actually improved on increasing the ester size from methyl to *tert*-butyl (Table 2, entries 1–4, from 91% *ee* to 95% *ee*). In contrast, hindered esters were confirmed to have a detrimental effect on asymmetric induction with $\text{Rh}_2(\text{R-DOSP})_4$, leading to a steady drop in enantioselectivity from 81% *ee* to 16% *ee* (Table 2, entries, 5–8).

10 To probe the generality of this catalyst in enantioselective cyclopropanation with donor/acceptor carbenoid intermediates, various combinations of aryl- or vinyl diazoacetates with terminal alkenes were evaluated. The results for representative vinyl diazoacetates and terminal alkenes are summarized in Table 3.




Table 3. Rh₂(*R*-BTPCP)₄-Catalyzed Asymmetric Cyclopropanation with Styryldiazoacetates

entry	R ¹	diazo compound	R ²	product	yield (%)
1	<i>p</i> -CF ₃ C ₆ H ₄	9	Ph	16a	87
2	<i>p</i> -NO ₂ C ₆ H ₄	9	Ph	16b	92
3	<i>p</i> -ClC ₆ H ₄	9	Ph	16c	86
4	3,4-diClC ₆ H ₃	9	Ph	16d	88
5	<i>p</i> -MeOC ₆ H ₄	9	Ph	16e	94
6	<i>o</i> -CH ₃ C ₆ H ₄	9	Ph	16f	88
7	<i>o</i> -MeO ₂ CC ₆ H ₄	9	Ph	16g	59
8	H ₃ C(CH ₂) ₃	9	Ph	16h	60
9	Ph	15a	<i>p</i> -ClC ₆ H ₄	16i	82
10	Ph	15c	3,4-diClC ₆ H ₃	16j	79
11	Ph	15d	3,4-diMeOC ₆ H ₃	16k	70
<i>ee</i> (%)					
	97				
	95				
	95				
	93				
	84(93) ^b				
	93				
	89				
	90 ^c				
	90				
	86				
	−93 ^d				

5

a- Standard conditions: Diazo compound (0.4 mmol) was added to a solution of Rh(II) catalyst and styrene (2.0 mmol) in dichloromethane under argon over 1 h. Reported yields were obtained after chromatographic purification. *dr* was determined by ¹H NMR prior to chromatography, and *ee* was determined by chiral HPLC.

10

b- Reaction was conducted at −40 °C.

c- *dr* >10:1.

d- Rh₂(*S*-BTPCP)₄ was used as catalyst.

The reactions proceeded smoothly in dichloromethane, affording the corresponding cyclopropanes 16a-k with high diastereo- and enantioselectivity with yields ranging from 59% to 94%. Carbenoid precursors with electron-donating substituents on the donor group (Table 3, entry 11) resulted in higher asymmetric induction than those having electron-withdrawing substituents on the donor group (Table 3, entries 9, 10). Electron-deficient alkenes gave better enantioselectivity than electron-rich alkenes. An optimum system was identified as para-trifluoromethylstyrene and its reaction with styryldiazoacetate 9 which produced the cyclopropane 16a in 97% ee (Table 3, entry 1). In contrast, the reaction with para-methoxystyrene produced the corresponding cyclopropane 16e in 83% ee at room temperature, but this could be improved to 93% ee when the reaction was conducted at -40 °C (Table 3, entry 5). Inclusion of an ortho substituent on the styrene had little influence on the enantioselectivity, as reaction of ortho-methylstyrene and methyl 2-vinylbenzoate with the styryldiazoacetate 9 provided the cyclopropanes 16f and 16g in 93% ee and 89% ee, respectively (Table 3, entries 6, 7). Unactivated alkenes are also suitable substrates as 1-hexene (Table 3, entry 8) gave the cyclopropane 16h in 60% yield and 90% ee.

The cyclopropanation reactions of representative aryldiazoacetates are summarized in Table 4.




Table 4. Rh₂(*R*-BTPCP)₄-Catalyzed Asymmetric Cyclopropanation with Aryldiazoacetates

entry	starting material	R	product	yield (%)	ee (%)
1	7	Ph	18a	82	83
2	12	<i>p</i> -BrC ₆ H ₄	18b	80	85
3	17a	<i>p</i> -MeOC ₆ H ₄	18c	74	91
4	17b	<i>p</i> -CF ₃ C ₆ H ₄	18d	86	89

a- Standard conditions: Diazo compound (0.4 mmol) was added to a solution of Rh(II) catalyst and styrene (2.0 mmol) in dichloromethane under argon over 1 h. Reported yields were obtained after chromatographic purification. dr was determined by ^1H NMR prior to chromatography, and ee was determined by chiral HPLC.

b- Reaction was conducted at $-40\text{ }^\circ\text{C}$.

c- dr >10:1.

d- $\text{Rh}_2(\text{S-BTPCP})_4$ was used as catalyst.

The reactions generated the cyclopropanes 18a–d in high yield and stereoselectivity (>20:1 dr, 83–91% ee), with p-methoxyphenyl derivative 17a giving the highest enantioselectivity (91% ee). These results are similar to the reported results for the $\text{Rh}_2(\text{S-DOSP})_4$ -catalyzed cyclopropanations of aryldiazoacetates in hexane as solvent.

Encouraged by the cyclopropanation results, it was decided to investigate the scope of the catalyst further by looking at more elaborate reactions of vinyldiazoacetates. When vinyldiazoacetates react with dienes, a formal [4 + 3] cycloaddition occurs by a tandem cyclopropanation/Cope rearrangement. See Table 5.




Table 5. $\text{Rh}_2(\text{R-BTPCP})_4$ Catalyzed Tandem Cyclopropanation/Cope Rearrangement

entry	starting diene	R	product	yield (%)	ee (%)
1	19a	Ph	20a	56	87
2	19b	<i>p</i> -CF ₃ C ₆ H ₄	20b	71	91
3	19c	<i>p</i> -MeOC ₆ H ₄	20c	60	89

a- Standard conditions: Diazo compound (0.4 mmol) was added to a solution of Rh(II) catalyst and styrene (2.0 mmol) in dichloromethane at $-40\text{ }^\circ\text{C}$ under argon


over 1 h. Reported yields were obtained after chromatographic purification. dr was determined by ^1H NMR prior to chromatography, and ee was determined by chiral HPLC. Absolute configuration of the products was assigned according to previous studies.

5

The $\text{Rh}_2(\text{R-BTPCP})_4$ -catalyzed reactions of styryldiazoacetate **9** with dienes **19** gave rise to the cycloheptadienes **20** in 56–71% yield and the enantioselectivity was 87–91% ee. Compounds **20** were formed as single diastereomers.


Vinyldiazoacetates are useful in a range of C–H functionalization reactions. An example is the formal C–H functionalization of dihydronaphthalene **21** to form **22**.

10



Catalysts with cyclopropane rings containing three aromatic rings as gave asymmetric induction, see Table 6 and Figure 9.

15



20 Table 6. Dirhodium Catalysts with Different Substituents

entry	catalyst	yield (%)	ee (%)
1	Rh ₂ (<i>R</i> -DPCP) ₄	81	11
2	Rh ₂ (<i>R</i> -TPCP) ₄	84	79
3	Rh ₂ (<i>R</i> -BTPCP) ₄	86	91
4	Rh ₂ (1 <i>R</i> ,2 <i>S</i> -CMCP) ₄	81	22
5	Rh ₂ (<i>R</i> -NPCP) ₄	77	94
6	Rh ₂ (<i>R</i> -BPPCP) ₄	76	84
7	Rh ₂ (<i>S</i> -BNPCP) ₄	74	-87

Although it is not intended that certain embodiments, of the disclosure be limited by any particular mechanism, it is believed that this reaction proceeds by a sequence involving a combined C–H functionalization/Cope rearrangement followed by a reverse Cope rearrangement. When this reaction was catalyzed by Rh₂(*R*-BTPCP)₄, the product 22 was formed in 92% yield as a single diastereomer and in 98% ee. This result compares favorably to the Rh₂(*S*-DOSP)₄-catalyzed formation of 22 with hexane as solvent.

Even though dirhodium tetracarboxylates contain two potentially reactive rhodium sites and four identical ligands of C₁ symmetry, they are capable of very high asymmetric induction. Although it is not intended that embodiments of this disclosure be limited by any particular mechanism, it is believed that the dirhodium tetracarboxylates are in conformations of higher symmetry than the ligands themselves. For Rh₂(*S*-DOSP)₄-catalyzed reactions, the catalyst is proposed to preferentially exist in a D₂ symmetric “up-down-up-down” conformation, providing identical C₂ symmetric binding sites at both rhodium faces of the catalyst (Figure 2). Hashimoto rationalized the selectivity of his phthalimido amino acid derived catalysts by proposing a C₂ symmetric “up-up-down-down” arrangement for the catalyst. More recently, Fox reported that the tert-butyl catalyst Rh₂(*S*-PTTL)₄, the most broadly used of Hashimoto’s catalysts, exists in an “all-up” distorted C₄ conformation. According to this model, the bulky tert-butyl groups are necessary to limit reactivity to only one of the Rh faces of the catalyst, whereas a distorted C₄ symmetric chiral crown-like ligand arrangement guides the facial selectivity at the open Rh-face. Recently, other groups have reported X-ray structures of related catalysts to Rh₂(*S*-PTTL)₄, and all these catalysts adopt an “all-up” C₄ conformation. Because Rh₂(*R*-BTPCP)₄ is structurally different from the other above-mentioned chiral dirhodium tetracarboxylate catalysts, a study to determine what made it such an effective chiral catalyst was undertaken.

A single crystal X-ray diffraction analysis of $\text{Rh}_2(\text{R-BTPCP})_4$ provided valuable insights into the catalyst geometry. In the X-ray structure, the large chiral ligands of $\text{Rh}_2(\text{R-BTPCP})_4$ are organized in an overall D2 symmetric arrangement, forming identical rectangular orthogonal (approximately $8.5 \times 10.5 \text{ \AA}$) binding cavities of C2 symmetry at the two catalytically active axial termini of the rhodium dimer (Figure 3).

$\text{Rh}_2(\text{R-BTPCP})_4$ and $\text{Rh}_2(\text{S-PTTL})_4$ have different overall symmetry, D2 and distorted C4, respectively, but they possess a related rectangular binding cavity at the reactive axial positions (Figure 4). The rectangular binding cavity for $\text{Rh}_2(\text{R-BTPCP})_4$ ($8.5 \times 10.5 \text{ \AA}$) is significantly smaller than the cavity for $\text{Rh}_2(\text{S-PTTL})_4$ ($12.8 \times 14.1 \text{ \AA}$). Furthermore, both Rh faces of $\text{Rh}_2(\text{R-BTPCP})_4$ have identical binding cavities of C2 symmetry. However, for $\text{Rh}_2(\text{S-PTTL})_4$ only one Rh face has the rectangular binding cavity, whereas the other face is considered to be blocked by the tert-butyl groups. According to the $\text{Rh}_2(\text{S-PTTL})_4$ model proposed by Fox, a carbene would align with the wide dimension of the catalyst leaving the Si face open for the attack because of the wider gap between two of the ligands. To test if the same model could be utilized to explain the stereochemical outcome of $\text{Rh}_2(\text{R-BTPCP})_4$ -catalyzed transformations, a computational analysis of the catalyst-carbene complex was conducted.

Traditional computational approaches to analyze dirhodium carbene complexes have tended to rely on the X-ray structure of the catalyst in combination with DFT optimized geometries of the carbenoid with a simplified, achiral, catalyst model. Although such analysis may provide preliminary insights into the catalyst selectivity, more precise studies accounting for interactions between the carbene and ligands, as well as ligand mobility in these systems would give a much better understanding of the role of the chiral catalyst. In fact, our preliminary DFT calculations on the simplified model of the dirhodium(tetracarboxylate)-carbene complex (Figure 5) demonstrates the importance of such studies. A comparison of the DFT optimized geometries, given in Figure 5, with the reported size of the catalyst binding cavity (Figure 4) reveals that even the shortest s-trans carbenoid conformation would not fit into the tight environment (the dimensions shown on Figures 4 and 5 were measured from atomic centers and do not account for atomic radii) of the $\text{Rh}_2(\text{R-BTPCP})_4$ catalyst.

Performing DFT calculations on the complete rhodium carbene complex (approximately 200 atoms) would be impractical unless very small basis sets were used, and these may not give a realistic representation of the complex. Therefore, we employ an alternative ONIOM approach to study the carbenoid reaction. A two-layer ONIOM

(QM:MM) method used in these studies divides the whole system into two subsystems (denoted "model" and "real" layers). The ONIOM partitioning of the catalyst-carbene system that was used is shown in Figure 6. In these studies the B3LYP and UFF methods were used for the calculations of the "model" (in red) and "real" (in blue) layers,
5 respectively. Such partition provided an description for the central rhodium carboxylate-carbene complex and accounted for the steric influence of the surrounding ligands.

Our computational results support the importance of the steric environment within the catalyst. Thus even the best matching combination of the shortest s-trans carbene aligned with the widest dimension of the orthogonal binding cavity caused steric
10 repulsions significant enough to rotate two para-bromophenyl groups in either conrotary or disrotary directions. A similar result was obtained when the s-trans carbene was aligned with the shortest catalyst dimension; this time the phenyl groups rotated in a similar way. Sixteen distinct minima were located on the potential energy surface corresponding to different conformations of the s-trans carbene with $\text{Rh}_2(\text{R-BTPCP})_4$. The lowest energy
15 conformation, separated from the closest one by $\Delta E = 1.8$ kcal/mol is depicted in Figure 7. In this conformer two ligands rotated in a conrotary fashion to minimize steric interactions with the carbene, whereas the other two ligands remained in the upward position to reduce steric repulsions with the neighboring ligands. This arrangement resulted in a C2
symmetric environment at the carbene site cavity containing two phenyl rings and two
20 para-bromophenyl groups. One of the rings is blocking the donor group (aryl, styryl) while the other one is positioned next to the acceptor group (ester). The same ligand conformation having the s-cis carbene geometry was found to be 2.5 kcal/mol higher.

Although it is not intended that embodiments of the disclosure be limited by any mechanism, a stereochemical model that explains the selectivity observed in $\text{Rh}_2(\text{R-BTPCP})_4$ catalyzed transformations is proposed in Figure 8 basis on the computational
25 data above. The substrate approaches donor/acceptor-substituted rhodium-carbenoids over the donor group. The ester group aligns perpendicular to the carbene plane, and blocks attack on its side. When the substrate approaches over the donor group, the Re-face is blocked by the aryl ring of the ligand leaving the Si-face open for the attack. This model
30 predicts correctly the observed absolute configuration of the products.

**Synthesis of Dirhodium tetrakis((1S,2R)-1,2-diphenylcyclopropanecarboxylate (8).
(1S,2R)-methyl 1,2-diphenylcyclopropanecarboxylate**

To a flame-dried round bottom flask kept under a dry atmosphere of argon, was added $\text{Rh}_2(\text{RDOSP})_4$ (0.01 equiv., 379 mg), styrene (2.32 equiv., 13.7 mL), and dry
5 degassed pentane (100 mL) at room temperature. A solution of methyl 2-diazo-2-phenylacetate 7 (1.0 equiv., 3.52 g) in dry, degassed pentane (100 mL) was then added to the former solution drop-wise over 3 hours under argon atmosphere. The mixture was then allowed to stir for overnight, and concentrated in vacuo. The crude residue was analyzed by ^1H NMR and purified by flash column chromatography eluting with a 50:1 mixture of
10 hexanes/ethyl acetate to provide the product as a white solid (4.08 g, 81% yield, 91% ee). Recrystallization with a 100:1 mixture of hexane/ethyl acetate afforded the enantioenriched product (2.82 g, 69% yield, >99% ee). ^1H NMR (400 MHz, CDCl_3): δ 7.17–7.15 (m, 3H), 7.09–7.05 (m, 5H), 6.81–6.79 (m, 2H), 3.69 (s, 3H), 3.15 (dd, $J = 9.4$, 7.3 Hz, 1H), 2.18 (dd, $J = 9.4$, 5.0 Hz, 1H), 1.92 (dd, $J = 7.3$, 5.0 Hz, 1H); HPLC (S,S-
15 Whelk, 1.5% isopropanol in hexane, 0.7 mL/min, 1 mg/mL, 30 min, UV 254 nm) retention times of 12.9 min (major) and 15.4 min (minor), 91% ee.

(1S,2R)-1,2-diphenylcyclopropanecarboxylic acid

20 To a round-bottom flask at room temperature was added (1S,2R)-methyl 1,2-diphenylcyclopropanecarboxylate (11.5 mmol, 2.91 g, 1.0 equiv.) in dry DMSO (24 mL). $t\text{-BuOK}$ (25.4 mmol, 2.85 g, 2.2 equiv.) was added in several portions over 30 minutes under argon. The reaction was monitored by TLC technique until the starting material was consumed completely. The reaction mixture was cooled with ice bath and acidified by
25 careful addition of saturated ammonium chloride (10 mL), followed by a slow addition of 1 N HCl aqueous (35 mL) with vigorous stirring until the pH value reached 3–4. Light yellow solid precipitate was collected by filtration, washed with water (3x5 mL), dissolved in ethyl acetate (100 mL), washed with brine (3x5 mL), dried over MgSO_4 , and concentrated in vacuo. The crude material was purified using flash column
30 chromatography eluting with a 10:1 mixture of hexanes/ethyl acetate to provide the desired product as a white solid (1.64 g, 60% yield, >99% ee). ^1H NMR (400 MHz, CDCl_3) δ 7.15–7.14 (m, 3H), 7.08–7.03 (m, 5H), 7.06–6.79 (m, 2H), 3.18 (dd, $J = 9.4$, 7.4 Hz, 1H), 2.21 (dd, $J = 9.4$, 4.9 Hz, 1H), 1.97 (dd, $J = 7.4$, 4.9 Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 179.8, 136.1, 134.2, 132.1, 128.3, 127.99, 127.97, 127.5, 126.7, 37.4,

34.1, 20.9; HPLC (to improve the separation, the product was converted to the corresponding methyl ester prepared using dimethylsulfate/KOH in tetrahydrofuran under reflux), (S,S)-Whelk, 1.5% isopropanol in hexane, 0.7 mL/min, 1 mg/mL, 30 min, UV 254 nm) retention times of 12.5 min (major) and 15.5 min (minor), >99% ee.

5

Dirhodium tetrakis((1S,2R)-1,2-diphenylcyclopropanecarboxylate (8)

A solution of sodium rhodium carbonate $[\text{Na}_4\text{Rh}_2(\text{CO}_3)_4] \cdot 2.5\text{H}_2\text{O}$ (501 mg, 1.0 equiv.) and (1S,2R)-1,2-diphenylcyclopropanecarboxylic acid (1.64 g, 8.0 equiv.) in 35 mL distilled water was refluxed for 2 days under argon, and then the solution was
 10 extracted with dichloromethane (3x50 mL), organic extracts were combined, and then washed with saturated sodium bicarbonate (3x10 mL), brine (3x10 mL), dried over anhydrous MgSO_4 , and concentrated in vacuo. The crude material was purified using flash column chromatography eluting with a 100:1 mixture of toluene and acetonitrile mixtures to provide the desired catalyst 8 as a green solid (650 mg, 65%). mp: 222-224 °C; R_f =
 15 0.20 (toluene/acetonitrile = 20/1); $[\alpha]_{20}^{\text{D}} 265.1^\circ$ (c = 0.02, CHCl_3); ^1H NMR (400 MHz, CDCl_3): δ 7.10-7.05 (m, 24H), 6.94-6.92 (m, 8H), 6.74-6.72 (m, 8H), 2.96 (dd, J = 9.2, 7.2 Hz, 1H), 2.04 (dd, J = 9.2, 4.8 Hz, 1H), 1.77 (dd, J = 7.2, 4.8 Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 191.6, 136.8, 135.4, 131.7, 127.9, 127.6, 127.5, 126.7, 126.1, 60.3, 39.5, 32.9, 19.7; IR (neat): 3028, 1682, 1577,
 20 1497, 1398; HRMS (ESI) calcd for $\text{C}_{64}\text{H}_{52}\text{O}_8\text{ClRh}_2$ ($\text{M}+\text{Cl}$) $^+$ 1189.14663 found 1189.14675.

Synthesis of Dirhodium Tetrakis(1-(4-bromophenyl)-2,2-diphenylcyclopropane carboxylate (4)

25 (R)-methyl 1-(4-bromophenyl)-2,2-diphenylcyclopropanecarboxylate

To a flame-dried round bottom flask kept under a dry atmosphere of argon, was added $\text{Rh}_2(\text{RDOSP})_4$ (0.01 equiv., 382 mg), 1,1-diphenylethylene 11 (2.32 equiv., 8.16 mL), and dry degassed pentane (100 mL). A solution of freshly prepared methyl 2-(4-bromophenyl)-2-diazoacetate 12 (1.0 equiv., 5.1 g) in dry, degassed pentane (150 mL) was
 30 added to the former solution drop-wise over 3 hours at room temperature. The mixture was allowed to stir overnight, and then concentrated in vacuo. The crude material was purified using flash column chromatography eluting with a 50:1 mixture of hexanes/ethyl acetate to provide the desired product as a white foam (7.2 g, 88% yield, 98% ee). mp: 82-85 °C; R_f = 0.55 (hexanes/ethyl acetate = 7/1); $[\alpha]_{20}^{\text{D}} -289.2^\circ$ (c = 1.17, CHCl_3); ^1H NMR (400

MHz, CDCl₃) δ 7.48-7.47 (m, 2H), 7.35-7.31 (m, 2H), 7.27-7.18 (m, 5H), 7.02-6.97 (m, 5H), 3.34 (s, 3H), 2.68 (d, J = 5.5 Hz, 1H), 2.39 (d, J = 5.9 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 170.9, 141.7, 139.1, 134.8, 133.5, 130.6, 129.8, 128.5, 128.3, 127.7, 126.9, 126.3, 121.1, 52.2, 44.6, 42.4, 22.6; IR (neat): 3057, 3024, 2947, 1724, 1490, 1217;
 5 HRMS (APCI) calcd for C₂₃H₂₀BrO₂.(M+H)⁺ 407.06412 found 407.06432; HPLC (S,S-Whelk, 5% isopropanol in hexane, 0.8 mL/min, 1 mg/mL, 60 min, UV 254 nm) retention times of 10.6 min (major) and 50.9 min (minor), 98% ee. (S)-methyl 1-(4-bromophenyl)-2,2-diphenylcyclopropanecarboxylate was prepared by the same procedure with Rh₂(S-DOSP)₄ as the catalyst, [α]_D²⁰: 219.6° (c = 1.32, CHCl₃).

10

(R)-1-(4-bromophenyl)-2,2-diphenylcyclopropanecarboxylic acid

To a round-bottom flask at room temperature was added (R)-methyl 1-(4-bromophenyl)-2,2-diphenylcyclopropanecarboxylate (17.9 mmol, 7.3 g, 1.0 equiv) in dry DMSO (35 mL). t-BuOK (39.6 mmol, 4.4 g, 2.2 equiv) was added in several portions over
 15 30 minutes under argon. The reaction was monitored by TLC technique until the starting material was consumed completely. The reaction mixture was cooled with ice bath and acidified by saturated ammonium chloride aqueous (15 mL), followed by a slow addition of 1 N HCl (50 mL) with vigorous stirring until the pH value reached 3-4. Sticky solid precipitate was collected by filtration, washed with water (3x5 mL), dissolved in ethyl
 20 acetate (150 mL), washed with brine (3x10 mL), dried over anhydrous MgSO₄, and concentrated in vacuo. The crude material was purified using flash column chromatography eluting with a 4:1 mixture of hexanes/ethyl acetate to provide the desired product as a white solid (4.9 g, 69% yield). Recrystallization in pentane/ethyl acetate (50/1) provided the enantioenriched product as a white solid (4.46 g, 64% yield, 99% ee).
 25 (S)-1-(4-bromophenyl)-2,2-diphenylcyclopropanecarboxylic acid was prepared by the same procedure,
 [α]_D²⁰ 260.4° (c = 1.37, CHCl₃). mp: 180-182 °C; R_f = 0.21 (hexanes/ethyl acetate = 5/1);
 [α]_D²⁰ -224.8° (c = 3.77, CHCl₃); ¹H NMR (400 MHz; CDCl₃) δ 7.44-7.42 (m, 2H),
 30 7.29-7.22 (m, 5H), 7.13-7.11 (m, 2H), 6.99-6.93 (m, 5H), 2.56 (d, J = 5.6 Hz, 1H), 2.41 (d, J = 5.6 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 176.7, 141.2, 138.9, 134.3, 133.5, 130.6, 129.7, 128.5, 128.4, 127.8, 127.1, 126.5, 121.3, 45.8, 41.9, 23.1; IR (neat): 3024, 1688, 1489, 1449; HRMS (APCI) calcd for C₂₂H₁₆BrO₂ (M-H)⁺ 391.03391 found 391.03446; HPLC (to improve the separation, the product was converted to the corresponding methyl

ester prepared using dimethylsulfate/KOH in tetrahydrofuran under reflux), (S,S-Whelk, 5% isopropanol in hexane, 0.8 mL/min, 1 mg/mL, 60 min, UV 254 nm) retention times of 10.7 min (major) and 49.8 min (minor), 99% ee.

5 Dirhodium tetrakis((R)-1-(4-bromophenyl)-2,2-diphenylcyclopropanecarboxylate) (4)

A solution of sodium rhodium carbonate $[\text{Na}_4\text{Rh}_2(\text{CO}_3)_4] \cdot 2.5\text{H}_2\text{O}$ (557 mg, 1.0 equiv.) and (R)-1-(4-bromophenyl)-2,2-diphenylcyclopropanecarboxylic acid (3.0 g, 8.0 equiv.) in 60 mL distilled water was refluxed for 3 days under argon, and then the solution
 10 was extracted with dichloromethane (3x100mL), organic extracts were combined, washed with saturated sodium bicarbonate (3x10 mL), 10% sodium hydroxide (3x10 mL), brine (3x10 mL), dried over anhydrous MgSO_4 , and concentrated in vacuo. The crude material was purified using flash column chromatography eluting with a 50:1 mixture of toluene and acetonitrile to provide the desired catalyst 4 as a green solid (1.07 g, 63% yield). mp:
 15 235-237 °C; $R_f = 0.25$ (toluene/acetonitrile = 50/1); $[\alpha]_D^{20} -41.7^\circ$ ($c = 0.02$, CHCl_3); ^1H NMR (400 MHz, CDCl_3): δ 7.23–7.15 (m, 7H), 6.90–6.81 (m, 7H), 2.30 (d, 1H, $J = 5.0$ Hz), 2.39 (d, 1H, $J = 5.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ 188.9, 142.4, 140.8, 135.7, 132.8, 130.6, 129.9, 129.2, 128.3, 127.9, 126.6, 126.4, 120.9, 46.9, 42.7, 23.6; IR (neat): 1577, 1490, 1447, 1397, 1379, 1277, 1156, 1076, 1010, 991, 906, 823, 777, 733; HRMS
 20 (ESI) calcd for $\text{C}_{88}\text{H}_{64}\text{Br}_4\text{O}_8$ (M^+) 1769.9444 found 1769.94392. $\text{Rh}_2(\text{S-BTPCP})_4$ was prepared with the same procedure, $[\alpha]_D^{20} 140.9^\circ$ ($c = 0.02$, CHCl_3).

Preparation of $[\text{Rh}_2(\text{R-NPCP})_4]$

(R)-methyl 1-(naphthalen-2-yl)-2,2-diphenylcyclopropanecarboxylate

To a flame-dried round bottom flask kept under a dry atmosphere of argon, was
 25 added $\text{Rh}_2(\text{R-DOSP})_4$ (0.01 equiv., 474 mg), 1,1-diphenylethyene (2.32 equiv., 10.5 g, 58 mmol), and dry degassed pentane (100 mL). A solution of freshly prepared methyl 2-diazo-2-(naphthalen-2-yl)acetate (1.0 equiv., 5.65 g, 25 mmol) in dry, degassed pentane (250 mL) was added to the former solution drop-wise over 3 hours at -40 °C. The mixture was allowed to stir overnight to room temperature, and then concentrated *in vacuo*. The
 30 crude material was purified using flash column chromatography eluting with a 40:1 mixture of hexanes/ethyl acetate to provide the desired product as a white solid (7.0 g, 74% yield, >99% ee). mp: 138-139 °C; $[\alpha]_D^{20} -347.9^\circ$ ($c = 1.0$, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.78 (s, 1H), 7.74-7.71 (m, 1H), 7.50-7.36 (m, 4H), 7.24-7.22 (m, 1H),

7.06-7.04 (m, 2H), 6.96-6.86 (m, 3H), 3.38 (s, 3H), 2.79 (d, $J = 5.6$ Hz, 1H), 2.57 (d, $J = 5.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 171.4, 141.9, 139.4, 133.6, 132.8, 132.3, 130.4, 130.2, 129.9, 128.7, 128.3, 127.7, 127.6, 127.5, 126.9, 126.7, 126.1, 125.8, 125.6, 141.9, 139.4, 133.6, 132.8, 132.2, 130.4, 130.2, 129.9, 128.7, 128.3, 127.7, 127.6, 127.5, 126.9, 126.7, 126.1, 125.8, 125.6, 52.2, 44.5, 43.2, 23.1; IR (neat): 3057, 3024, 2947, 1724, 1433, 1215; HRMS (APCI) calcd for $\text{C}_{27}\text{H}_{23}\text{O}_2$ ($\text{M}+\text{H}$) $^+$ 379.16926 found 379.16952; HPLC (ODH, 0.5% isopropanol in hexane, 1.0 mL/min, 1 mg/mL, 30 min, UV 254 nm) retention times of 8.6 min (major) and 9.6 min (minor), >99% ee.

10 **(*R*)-1-(naphthalen-2-yl)-2,2-diphenylcyclopropanecarboxylic acid**

To a round-bottom flask at room temperature was added (*R*)-methyl 1-(naphthalen-2-yl)-2,2-diphenylcyclopropanecarboxylate (18.5 mmol, 7.0 g, 1.0 equiv) in dry DMSO (50 mL). $t\text{-BuOK}$ (41.0 mmol, 4.6 g, 2.2 equiv) was added in several portions over 30 minutes under argon. The reaction was monitored by TLC technique until the starting material was consumed completely. The reaction mixture was cooled with ice bath and acidified by saturated ammonium chloride aqueous (15 mL), followed by a slow addition of 1 N HCl (40 mL) with vigorous stirring until the pH value reached 3-4. Sticky solid precipitate was collected by filtration, washed with water (3x5 mL), dissolved in ethyl acetate (150 mL), washed with brine (3x10 mL), dried over anhydrous MgSO_4 , and concentrated *in vacuo*. The crude material was purified using flash column chromatography eluting with a 4:1 mixture of hexanes/ethyl acetate to provide the desired product as a white solid (4.1 g, 61% yield. mp: 195-196 °C; $[\alpha]_{\text{D}}^{20}$ -361.7° ($c = 1.0$, CHCl_3); ^1H NMR (400 MHz; CDCl_3) δ 7.80-7.76 (m, 3H), 7.65-7.59 (m, 3H), 7.53-7.46 (m, 3H), 7.42-7.34 (m, 3H), 7.11-7.09 (m, 2H), 7.0-6.9 (m, 3H), 2.73 (d, $J = 5.2$ Hz, 1H), 2.65 (d, $J = 5.2$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 177.4, 141.5, 139.1, 133.1, 132.7, 132.3, 130.3, 130.1, 129.8, 128.6, 128.4, 127.7, 127.6, 127.5, 126.9, 126.7, 126.2, 125.8, 125.7, 42.7, 23.4; IR (neat): 3058, 1687, 1448, 1300; HRMS (APCI) calcd for $\text{C}_{26}\text{H}_{21}\text{O}_2$ ($\text{M}+\text{H}$) $^+$ 365.15361 found 365.15349; HPLC (to improve the separation, the product was converted to the corresponding methyl ester prepared using dimethylsulfate/KOH in tetrahydrofuran under reflux), HPLC (ODH, 0.5% isopropanol in hexane, 1.0 mL/min, 1 mg/mL, 30 min, UV 254 nm) retention times of 8.6 min (major) and 9.6 min (minor), >99% ee.

Dirhodium Tetrakis[(R)-1-(naphthalen-2-yl)-2,2-diphenylcyclopropanecarboxylate]

A solution of sodium rhodium carbonate $[\text{Na}_4\text{Rh}_2(\text{CO}_3)_4] \cdot 2.5\text{H}_2\text{O}$ (599.6 mg, 1.0 equiv.) and (R)-1-(naphthalen-2-yl)-2,2-diphenylcyclopropanecarboxylic acid (3.0 g, 8.0 equiv.) in 60 mL distilled water was refluxed for 2 days under argon, and then the solution
5 was extracted with dichloromethane (3x100mL), organic extracts were combined, washed with saturated sodium bicarbonate (3x10 mL), 10% sodium hydroxide (3x10 mL), brine (3x10 mL), dried over anhydrous MgSO_4 , and concentrated *in vacuo*. The crude material was purified using flash column chromatography eluting with a 50:1 mixture of toluene and acetonitrile to provide the desired catalyst as a green solid (989 mg, 58% yield). mp:
10 249-251 °C; $[\alpha]_D^{20}$ 91.2° (c = 0.02, CHCl_3); ^1H NMR (400 MHz, CDCl_3): δ 7.70-7.67 (m, 8 H), 7.54 (s, 4H), 7.43-7.29 (m, 32 H), 7.24-7.21 (m, 4H), 7.10 (d, J = 8.4 Hz, 4H), 6.92-6.85 (m, 12 H), 6.75-6.65 (m, 12H), 2.48 (d, J = 4.8 Hz, 4H), 2.02 (d, J = 4.8 Hz, 4H). ^{13}C NMR (100 MHz, CDCl_3) δ 188.9, 142.6, 140.9, 134.5, 132.6, 132.1, 130.9, 129.8, 129.0, 128.2, 128.1, 127.7, 127.6, 127.4, 127.3, 126.2, 126.1, 125.7, 125.4, 125.3, 46.6, 43.1,
15 23.5; IR (neat): 3056, 3023, 1579, 1389, 1348, 908, 744, 755, 704; HRMS (ESI) calcd for $\text{C}_{104}\text{H}_{76}\text{O}_8\text{Cl}(\text{M}+\text{Cl})^+$ 1693.33443 found 1693.33556.

General procedure for catalyzed cyclopropanation

To a 25 mL flame-dried round-bottom flask, kept under a dry atmosphere of argon,
20 was added alkene (2.0 mmol, 5.0 equiv.), dry and degassed CH_2Cl_2 (1.0 mL) and then $\text{Rh}_2(\text{R-BTPCP})_4$ (7.0 mg, 0.01 equiv.). The diazo compound (0.4 mmol, 1.0 equiv.), dissolved in dry and degassed CH_2Cl_2 (2 mL), was then added to the former solution dropwise over 1 hour at room temperature. The mixture was allowed to stir for at least 30 min after the addition; when the diazo compound was fully consumed by TLC analysis, the
25 reaction mixture was concentrated in vacuo. The crude residue was analyzed by ^1H NMR and purified by flash column chromatography (hexanes/ethyl acetate) to afford the cyclopropane.

(1R,2R)-methyl 2-phenyl-1-((E)-styryl)cyclopropanecarboxylate (10)

30 This compound was prepared according to the general procedure for $\text{Rh}_2(\text{R-BTPCP})_4$ -catalyzed cyclopropanation. Using (E)-methyl 2-diazo-4-phenylbut-3-enoate 9 (0.4 mmol, 80.8 mg, 1.0 equiv.) and styrene 6 (2.0 mmol, 208mg, 5.0 equiv.). The crude residue was analyzed by ^1H NMR and purified by flash column chromatography

(hexanes/ethyl acetate = 50/1) to afford 10 as a white solid (95.3 mg, 86% yield). ¹H NMR (400 MHz, CDCl₃): δ 7.25–7.21 (m, 4H), 7.19–7.13 (m, 6H), 6.35 (d, J = 16.2 Hz, 1H), 6.14 (d, J = 16.2 Hz, 1H), 3.77 (s, 3H), 3.04–2.99 (m, 1H), 2.03 (dd, J = 9.2, 5.0 Hz, 1H), 1.88 (dd, J = 7.3, 5.0 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 174.1, 137.0, 135.5, 133.0, 129.1, 128.3, 127.9, 127.3, 126.7, 126.2, 124.0, 52.4, 34.9, 33.2, 18.6; HPLC (OJH, 1.5% isopropanol in hexane, 1.0 mL/min, 1 mg/mL, 30 min, UV 254 nm) retention times of 15.1 min (major) and 19.9 min (minor), 91% ee.

General procedure for Rh₂(R-BTPCP)₄-catalyzed cyclopropanation/Cope rearrangement

To a 25 mL flame-dried round-bottom flask, kept under a dry atmosphere of argon, was added diene (2 mmol, 5.0 equiv.), dry and degassed CH₂Cl₂ (1.0 mL) and Rh₂(R-BTPCP)₄ (7.0 mg, 0.01 equiv.). The given diazo compound (0.4 mmol, 1.0 equiv.), dissolved in dry and degassed CH₂Cl₂ (2 mL), was then added to the former solution dropwise over 1 hour at -40 °C. The mixture was allowed to stir to room temperature overnight, and was then concentrated in vacuo. The crude residue was analyzed by ¹H NMR and purified by flash column chromatography (hexanes/ethyl acetate) to afford the desired 1,4-cycloheptadiene.

(3R,4S)-methyl 3,4-diphenylcyclohepta-1,5-dienecarboxylate (20a)

This compound was prepared according to the general procedure for Rh₂(R-BTPCP)₄-catalyzed cyclopropanation/Cope rearrangement. Using (E)-methyl 2-diazo-4-phenylbut-3-enoate 9 (0.4 mmol, 80.8 mg, 1.0 equiv.) and (E)-buta-1,3-dien-1-ylbenzene 19a (2.0 mmol, 260 mg, 5.0 equiv.). The crude residue was analyzed by ¹H NMR and purified by flash column chromatography (hexanes/ethyl acetate = 50/1) to afford 20a as a colorless oil (68.3 mg, 56% yield). ¹H NMR (400 MHz, CDCl₃): δ 7.23–7.12 (m, 7H), 6.90–6.88 (m, 2H), 6.82–6.80 (m, 2H), 6.01–5.95 (m, 1H), 5.78 (ddd, J = 11.4, 5.6, 2.7 Hz, 1H), 4.42 (s, 1H), 3.89 (s, 1H), 3.78 (s, 3H), 3.55–3.39 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 167.7, 144.5, 140.5, 139.8, 133.1, 132.3, 129.8, 129.1, 127.8, 127.5, 126.7, 126.6, 126.3, 52.0, 50.1, 49.6, 25.7; HPLC (S,S-Whelk, 1.5% isopropanol in hexane, 0.7 mL/min, 1 mg/mL, 30 min, UV 254 nm) retention times of 14.1 min (minor) and 15.8 min (major), 87% ee.

(3R,4S)-methyl 3-phenyl-4-(4-(trifluoromethyl)phenyl)cyclohepta-1,5-dienecarboxylate (20b)

This compound was prepared according to the general procedure for Rh₂(R-BTPCP)₄-catalyzed cyclopropanation/Cope rearrangement. Using (E)-methyl 2-diazo-4-phenylbut-3-enoate 9 (0.4 mmol, 80.8 mg, 1.0 equiv.) and (E)-1-(buta-1,3-dien-1-yl)-4-(trifluoromethyl) benzene 19b (2.0 mmol, 396 mg, 5.0 equiv.). The crude residue was analyzed by ¹H NMR and purified by flash column chromatography (hexanes/ethyl acetate = 50/1) to afford 20b as a colorless oil (105.2 mg, 71% yield). R_f = 0.53 (hexanes/ethyl acetate = 8/1); [α]_D²⁰ -60.7° (c = 3.57, CHCl₃); ¹H NMR (400 MHz, CDCl₃): δ 7.40 (d, J = 8.2 Hz, 2H), 7.26-7.19 (m, 3H), 7.08 (dd, J = 6.7, 2.7 Hz, 1H), 6.89-6.86 (m, 4H), 6.03-5.97 (m, 1H), 5.72 (ddd, J = 11.4, 5.6, 2.9, 1H), 4.44 (s, 1H), 3.91-3.89 (m, 1H), 3.77 (s, 3H), 3.56-3.49 (m, 1H), 3.44-3.37 (m, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 167.5, 143.9, 143.6, 140.0, 132.7, 132.0, 130.1, 129.4, 129.1, 128.9, 128.8, 128.6, 128.5, 128.0, 126.9, 125.6, 124.4, 124.3, 122.9, 52.0, 50.0, 49.2, 25.7; IR (neat): 3026, 2952, 1712, 1325; HRMS (APCI) calcd for C₂₂H₂₁F₃O₂ (M+H)⁺ 373.14153 found 373.14045; HPLC (OJH, 1.0% isopropanol in hexane, 1.0 mL/min, 1 mg/mL, 30 min, UV 254 nm) retention times of 6.2 min (minor) and 7.7 min (major), 91% ee.

(3R,4S)-methyl 4-(4-methoxyphenyl)-3-phenylcyclohepta-1,5-dienecarboxylate (20c)

This compound was prepared according to the general procedure for Rh₂(R-BTPCP)₄-catalyzed cyclopropanation/Cope rearrangement. Using (E)-methyl 2-diazo-4-phenylbut-3-enoate 9 (0.4 mmol, 80.8 mg, 1.0 equiv.) and (E)-1-(buta-1,3-dien-1-yl)-4-methoxybenzene 19c (2.0 mmol, 320 mg, 5.0 equiv.). The crude residue was analyzed by ¹H NMR and purified by flash column chromatography (hexanes/ethyl acetate = 50/1) to afford 20c as a colorless oil (79.6 mg, 60% yield). R_f = 0.51 (hexanes/ethyl acetate = 8/1); [α]_D²⁰ -2.1° (c = 3.01, MeOH); ¹H NMR (400 MHz, CDCl₃): δ 7.27-7.21 (m, 3H), 7.11 (dd, J = 6.7, 2.8 Hz, 1H), 6.90 (dd, J = 6.7, 2.8 Hz, 1H), 6.71 (s, 4H), 5.95-5.94 (m, 1H), 5.74 (ddd, J = 11.4, 5.6, 2.7 Hz, 1H), 4.39 (s, 1H), 3.79 (s, 3H), 3.77 (s, 3H), 3.52-3.46 (m, 1H), 3.42-3.37 (m, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 167.7, 158.4, 144.7, 140.8, 133.4, 132.4, 131.8, 130.8, 129.0, 127.8, 126.7, 125.9, 112.9, 55.1, 52.0, 49.7, 49.4, 25.7; IR (neat): 3010, 2950, 2359, 1708, 1509; HRMS (APCI) calcd for C₂₂H₂₄O₃ (M+H)⁺ 335.16471 found 335.16378; HPLC (S,S-Whelk, 1.5% isopropanol in hexane, 0.7 mL/min, 1 mg/mL, 30 min, UV 254 nm) retention times of 23.9 min (minor) and 26.0 min (major), 89% ee.

Procedure for Rh₂(R-BTPCP)₄-catalyzed combined C-H Cope/retro Cope rearrangement


(R,E)-methyl 2-((S)-4-methyl-1,2-dihydronaphthalen-2-yl)-4-phenylbut-3-enoate (22)

5 To a 25 mL flame-dried round-bottom flask, kept under a dry atmosphere of argon, was added 4-methyl-1,2-dihydronaphthalene (2 mmol, 5.0 equiv.), dry and degassed CH₂Cl₂ (1.0 mL) and Rh₂(R-BTPCP)₄ (7.0 mg, 0.01 equiv.). The given diazo compound (0.4 mmol, 1.0 equiv.), dissolved in dry and degassed CH₂Cl₂ (2 mL), was then added to the former solution drop-wise over 1 hour at room temperature. The mixture was allowed
10 to stir for overnight, and was then concentrated in vacuo. The crude residue was analyzed by ¹H NMR and purified by flash column chromatography (hexanes/ethyl acetate = 50/1) to afford the desired product 22 as a colorless oil (117.3 mg, 92% yield). The NMR spectral data are consistent with previously published results. R_f = 0.60 (hexanes/ethyl acetate = 8/1); ¹H NMR (400 MHz, CDCl₃): δ 7.36-7.07 (m, 9H), 6.42 (d, J = 15.9 Hz, 1H), 6.14 (dd, J = 15.9, 9.8 Hz, 1H), 5.73 (d, J = 3.1 Hz, 1H), 3.70 (s, 3H), 3.11 (t, J = 9.5 Hz, 1H), 2.89-2.83 (m, 2H), 2.74-2.70 (m, 1H), 2.07 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 173.9, 136.9, 135.5, 134.6, 133.9, 128.8, 128.0, 127.9, 127.4, 126.8, 126.6, 126.5, 126.3, 123.3, 53.4, 52.1, 36.3, 31.9, 19.6; HPLC (OD, 2.0% isopropanol in hexane, 0.8 mL/min, 1 mg/mL, 30 min, UV 254 nm) retention times of 8.9 min (minor) and 10.5 min (major),
20 98% ee.

CLAIMS

What is Claimed:

1. A dirhodium catalyst comprising a cyclopropyl ring substituted with a carboxylic acid ligand.
2. A composition comprising a compound of the following formula,



or salts thereof wherein,

R^1 is hydrogen, alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R^1 is optionally substituted with one or more, the same or different, R^4 ;

R^2 is hydrogen, alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R^2 is optionally substituted with one or more, the same or different, R^4 ;

R^3 is hydrogen, alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R^3 is optionally substituted with one or more, the same or different, R^4 ;

R^4 is alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl, alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R^4 is optionally substituted with one or more, the same or different, R^5 ;

R^5 is alkyl, halogen, cyano, hydroxy, amino, mercapto, formyl, carboxy, carbamoyl, alkoxy, alkanoyl, alkylthio, alkylamino, (alkyl)₂amino, alkylsulfinyl,

alkylsulfonyl, arylsulfonyl, carbocyclyl, aryl, or heterocyclyl, wherein R⁵ is optionally substituted with one or more, the same or different, R⁶; and

R⁶ is halogen, nitro, cyano, hydroxy, trifluoromethoxy, trifluoromethyl, amino, formyl, carboxy, carbamoyl, mercapto, sulfamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxymethyl, methylamino, ethylamino, dimethylamino, diethylamino, N-methyl-N-ethylamino, acetylamino, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, N-methyl-N-ethylcarbamoyl, methylthio, ethylthio, methylsulfinyl, ethylsulfinyl, mesyl, ethylsulfonyl, methoxycarbonyl, ethoxycarbonyl, N-methylsulfamoyl, N-ethylsulfamoyl, N,N-dimethylsulfamoyl, N,N-diethylsulfamoyl, N-methyl-N-ethylsulfamoyl, carbocyclyl, aryl, or heterocyclyl.

3. The composition of Claims 2-2, wherein R¹ is carbocyclyl, aryl, or heterocyclyl.
4. The composition of Claims 2-3, wherein R¹ is an aryl or an aromatic heterocyclyl.
5. The composition of Claims 2-4, wherein R¹ is a phenyl optionally substituted with one or more R⁴.
6. The composition of Claims 2-5, wherein R¹ is a phenyl optionally substituted with one or more halogen, alkyl, or alkoxy.
7. The composition of Claims 2-6, wherein R² is carbocyclyl, aryl, or heterocyclyl.
8. The composition of Claims 2-7, wherein R² is an aryl or an aromatic heterocyclyl.
9. The composition of Claims 2-8, wherein R² is a phenyl optionally substituted with one or more R⁴.
10. The composition of Claims 2-9, wherein R² is a phenyl optionally substituted with one or more halogen, alkyl, or alkoxy.
11. The composition of Claims 2-10, wherein R³ is carbocyclyl, aryl, or heterocyclyl.
12. The composition of Claims 2-11, wherein R³ is an aryl or an aromatic heterocyclyl.


13. The composition of Claims 2-12, wherein R^3 is a phenyl optionally substituted with one or more R^4 .

14. The composition of Claims 2-13, wherein R^3 is a phenyl optionally substituted with one or more halogen, alkyl, or alkoxy.

15. The composition of Claims 2-14, wherein R^3 is a phenyl substituted with a halogen.

16. The composition of Claims 2-15, wherein R^3 is a phenyl substituted with a halogen in the ortho or para position.


17. A composition comprising a compound,



wherein X is an electron withdrawing group.

18. The composition of Claim 17, wherein X is a halogen.

19. A composition comprising a compound



20. A method of making a synthetic compound comprising mixing

- a diazo compound,
- a compound with a carbon hydrogen bond, and
- a compound comprising a compound as in Claims 1-19,

under conditions such that a synthetic compound is formed comprising a carbon to carbon bond between the diazo compound and the compound with a carbon hydrogen bond.

21. A method of making a synthetic compound comprising mixing

- a) a diazo compound,
- b) a compound with a nitrogen hydrogen bond, and
- c) a compound comprising a compound as in Claims 1-19,

under conditions such that a synthetic compound is formed comprising a carbon to nitrogen bond between the diazo compound and the compound with a nitrogen hydrogen bond.




FIGURE 1

D_2 C_2 C_4

Davies'
 $\text{Rh}_2(\text{S-DOSP})_4$
model Hashimoto's
 $\text{Rh}_2(\text{S-PTPA})_4$
model Fox's
 $\text{Rh}_2(\text{S-PTTL})_4$
model

FIGURE 2

Figure 1 consists of two parts. Part (a) is a 3D ball-and-stick model of the [Rh2(μ₃-O)₂(μ₂-O)₂] core, showing two rhodium atoms (Rh) bridged by two μ₃-O atoms and two μ₂-O atoms. Part (b) is a schematic representation of the same core, showing the coordination of various ligands: two phenyl (Ph) groups, two 4-bromophenyl (Br-Ph) groups, and two 4-phenylphenyl (Ph-Ph) groups, along with two aryl (Ar) groups.

FIGURE 3







FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8




FIGURE 9