

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 January 2009 (08.01.2009)

PCT

(10) International Publication Number
WO 2009/006086 A2

- (51) International Patent Classification:
C07D 239/46 (2006.01) C07D 239/36 (2006.01)
- (21) International Application Number:
PCT/US2008/067963
- (22) International Filing Date: 24 June 2008 (24.06.2008)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/947,256 29 June 2007 (29.06.2007) US
- (71) Applicant (for all designated States except US): SANOFI-AVENTIS [FR/FR]; 174 avenue de France, F-75013 Paris (FR).

Elizabeth [US/US]; 55 Corporate Drive, Bridgewater, New Jersey 08807 (US). **POWERS, Matthew R.** [US/US]; 55 Corporate Drive, Bridgewater, New Jersey 088007 (US).


- (74) Agents: LIN, Jiang et al.; Sanofi-Aventis U.S. Inc., Route 202-206, P. O. Box 6800, Bridgewater, NJ 08807 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (72) Inventors; and
- (75) Inventors/Applicants (for US only): HUANG, Bao-Guo [US/US]; 55 Corporate Drive, Bridgewater, New Jersey 08807 (US). HANNA, Reda G. [US/US]; 55 Corporate Drive, Bridgewater, New Jersey 08807 (US). SECORD,

- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

[Continued on next page]

(54) Title: A NEW PROCESS FOR PREPARING 2-(3-{6-[2-(2,4-DICHLOROPHENYL)-ETHYLAMINO]-2-METHOXPYRIMIDIN-4-YL}-PHENYL)-2-METHYLPROPIONIC ACID



(57) Abstract: This invention is directed to a process for preparing 2-(3- {6-[2-(2,4-dichlorophenyl)-ethylamino]-2-methoxy-pyrimidin-4-yl}-phenyl)-2-methyl-propionic acid.

Area Percent Report

Signal 1: DAD1 D, Sig=220,4 Ref=500,12

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	7.089	MM	0.1323	6.41000	8.07685e-1	0.0761
2	9.849	MM	0.1574	7.28500	7.71627e-1	0.0865
3	11.360	MM	0.2757	8402.71777	508.05106	99.7740
4	14.849	MM	0.2001	5.33806	4.44598e-1	0.0634

Totals : 8421.75084 510.07497

Results obtained with enhanced integrator!

*** End of Report ***

WO 2009/006086 A2



European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

Declaration under Rule 4.17:

— *of inventorship (Rule 4.17(iv))*

**A NEW PROCESS FOR PREPARING 2-(3-{6-[2-(2,4-DICHLOROPHENYL)-
ETHYLAMINO]-2-METHOXYPYRIMIDIN-4-YL}-PHENYL)-
2-METHYLPROPIONIC ACID**

5


Field of the Invention

This invention is directed to a process for preparing 2-(3-{6-[2-(2,4-dichlorophenyl)-ethylamino]-2-methoxypyrimidin-4-yl}-phenyl)-2-methylpropionic acid.

10

Background of the Invention

Patent application WO 2006044732 (hereinafter the '732 application), hereby incorporated by reference, discloses pyrimidines of Formula (I),



15

(I)

wherein R^1 , L^1 , L^2 , Cy^1 and Cy^2 are as defined therein, their preparation, pharmaceutical compositions containing these compounds, and their pharmaceutical use in the treatment of disease states capable of being modulated by the inhibition of the prostaglandin D2 receptor, such as asthma [Arimura A, Yasui K, Kishino J, Asanuma F, Hasegawa H, Kakudo S, Ohtani M, Arita H, Prevention of allergic inflammation by a novel prostaglandin receptor antagonist, S-5751, *J Pharmacol Exp Ther.* **298**(2), 411-9, 2001], allergic rhinitis [Jones, T. R., Savoie, C., Robichaud, A., Sturino, C., Scheigetz, J., Lachance, N., Roy, B., Boyd, M., Abraham, W., Studies with a DP receptor antagonist in sheep and guinea pig models of allergic rhinitis, *Am. J. Resp. Crit. Care Med.* **167**, A218, 2003; and Arimura A, Yasui K, Kishino J, Asanuma F, Hasegawa H, Kakudo S, Ohtani M, Arita H, Prevention of allergic inflammation by a novel prostaglandin receptor antagonist, S-5751. *J. Pharmacol. Exp. Ther.* **298**(2), 411-9, 2001], and allergic conjunctivitis and allergic dermatitis [Arimura A, Yasui K, Kishino J, Asanuma F, Hasegawa H, Kakudo S, Ohtani M, Arita H, Prevention of allergic inflammation by a novel prostaglandin receptor antagonist, S-5751. *J. Pharmacol. Exp. Ther.* **298**(2), 411-9, 2001; and


30

Torisu K, Kobayashi K, Iwahashi M, Nakai Y, Onoda T, Nagase T, Sugimoto I, Okada Y, Matsumoto R, Nanbu F, Ohuchida S, Nakai H, Toda M, Discovery of a new class of potent, selective, and orally active prostaglandin D₂ receptor antagonists, *Bioorg. & Med. Chem.* 12, 5361-5378, 2004].

5


The '732 application discloses a process for preparing pyrimidines of Formula (I) wherein L² is a bond, by reacting a corresponding compound of Formula (V), wherein X² is a halogen, preferably chlorine, or a triflate group, with a boronic acid of Formula (VI), or a boronic acid pinacol ester of formula (XVII) in the presence of cesium carbonate and a complex metal catalyst such as tetrakis(triphenylphosphine)palladium (0) or *pddf*.

10




15

The '732 application further discloses that the compound of Formula (V) wherein X² is chlorine may be prepared by coupling a corresponding dichloropyrimidine of Formula (X) with a corresponding amine of Formula (IV), in the presence of a suitable base, such as sodium bicarbonate.



20


The '732 application more particularly discloses 2-(3-{6-[2-(2,4-dichlorophenyl)-ethylamino]-2-methoxypyrimidin-4-yl}-phenyl)-2-methylpropionic acid of formula (A),



and its preparation as shown in Scheme I.


5

Scheme I




10 The '732 application also discloses an alternative process for preparing pyrimidines of Formula (I) by reacting a corresponding compound of Formula (III), wherein X^1 is a halogen, particularly chlorine, or a triflate group, with a corresponding amine of Formula (IV), in the a suitable base, such as sodium bicarbonate, in an inert solvent, such as 1-methyl-2-pyrrolidinone, and at a temperature at about 160°C.

15



More particularly, Example 43(b) of the '732 application exemplifies the synthesis of [2-methoxy-6-(3-methoxy-phenyl)-pyrimidin-4-yl]-[2-(4-nitro-phenyl)-ethyl]-amine by such alternative process, as shown in Scheme II below:

Scheme II



10

Patent application WO 2007047378 (hereinafter '378 application) also specifically discloses dihydrogen phosphate salt of 2-(3-{6-[2-(2,4-dichlorophenyl)-ethylamino]-2-methoxypyrimidin-4-yl}-phenyl)-2-methylpropionic acid, and its preparation. However, the dihydrogen salt prepared in THF according to the procedures described therein contains about 1400-1600 ppm residual THF, which exceeds the limit of International Conference on Harmonization (ICH) guideline of 700 ppm. Furthermore, recrystallization from acetone is not practical for large scale preparation due to the low solubility.

15

A process for the preparation of a pharmaceutical active ingredient used in a medicament on an industrial scale has to fulfill various requirements. The process and the obtained product have to be in line with the regulatory requirements and have to be reproducible and validated.

5 In particular, regulatory authorities stipulate a precise degree of purity of the obtained drug substance and an acceptable level (e.g. single digit ppm) for heavy metals such as Pd or Ni. On the other hand, a process performed on an industrial scale for preparing a marketed product should of course be as simple, cost and labor effective as possible. If possible, it should thus avoid the use of expensive starting materials, physiologically unacceptable toxic
10 materials, difficult technical operations, long reaction time, or multiple procedural steps.

2-(3-{6-[2-(2,4-Dichloro-phenyl)-ethylamino]-2-methoxy-pyrimidin-4-yl}-phenyl)-2-methyl-propionic acid prepared by following the procedures described in the '732 application requires chromatographic purification, which is not desirable. In addition, it is also not desirable to
15 remove the residual Palladium catalyst in the last step of the process.

Furthermore, the procedures described in Example 43(b) of the '732 application utilize a high amount (5 mol%) of unstable catalyst, palladium tetrakis(triphenyl-phosphine). In addition, the intermediate and product formed therein also need to be purified by chromatography.

20 Some steps in the process also require long reaction time or use of microwave equipment, which lead to a very low time-space yield and a technical difficulty for large scale operation.


Thus, there is a need for a simpler and improved process for manufacturing the compound of formula A. The present invention satisfies this need by providing such a process.

25

Summary of the Invention

The present invention is directed to a process for preparing 2-(3-{6-[2-(2,4-dichlorophenyl)-ethylamino]-2-methoxypyrimidin-4-yl}-phenyl)-2-methylpropionic acid of formula (A),

30



(A)

comprising:

(1) coupling 4,6-dichloro-2-methoxypyrimidine with 3-(1-carboxy-1-methyl-ethyl)-phenyl boronic acid in the presence of a suitable Palladium catalyst to provide 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methylpropionic acid,

(2) coupling 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methyl-propionic acid with 2,4-dichlorophenethylamine to provide 2-(3-{6-[2-(2,4-dichlorophenyl)-ethylamino]-2-methoxy-pyrimidin-4-yl}-phenyl)-2-methylpropionic acid.

Another aspect of the invention is the compound 2-[3-(6-chloro-2-hydroxypyrimidin-4-yl)-phenyl]-2-methylpropionic acid, or a salt thereof.

The present invention is more fully discussed with the aid of the following figures and detailed description below.

Brief Description of the Drawings

FIGURE 1 is an HPLC spectrum of 2-(3-{6-[2-(2,4-dichloro-phenyl)-ethylamino]-2-methoxy-pyrimidin-4-yl}-phenyl)-2-methyl-propionic acid prepared by the process of the invention.

Detailed Description of the Invention

The present invention will be better appreciated by reference to the following detailed description.

Definitions and Abbreviations

As used above, and throughout the description of the invention, the following abbreviations, unless otherwise indicated, shall be understood to have the following meanings:

	DME	1,2-dimethoxyethane
5	g	gram
	h	hour
	HCl	Hydrochloric acid
	K ₂ CO ₃	Potassium carbonate
	Na ₂ CO ₃	Sodium carbonate
10	Cs ₂ CO ₃	Cesium carbonate
	K ₃ PO ₄	Potassium Phosphate
	Na ₃ PO ₄	Sodium Phosphate
	NaOH	Sodium hydroxide
	KOH	Potassium hydroxide
15	mg	milligram
	min	minute
	mL	milliliter
	n-BuOAc	n-butyl acetate
	<i>pddf</i>	1,1'-Bis(diphenylphosphino)ferrocene-palladium(II)dichloride
20		dichloromethane complex
	Pd(OAc) ₂	Palladium (II) acetate
	Pd(PPh ₃) ₄	tetrakis(triphenylphosphine)palladium (0)
	PPh ₃	triphenylphosphine
	ppm	parts per million
25	TMT	trithiocyanuric acid
	THF	tetrahydrofuran
	~	approximately

As used above, and throughout the description of the invention, the following terms, unless otherwise indicated, shall be understood to have the following meanings.

“Suitable Palladium catalyst” includes, for example, Pd, PdCl₂, Pd(OAc)₂, or *pddf*.

“Suitable base” includes inorganic base, for example, Na_2CO_3 , K_2CO_3 , Cs_2CO_3 , K_3PO_4 , KOH or NaOH, more particularly K_2CO_3 , and organic base, for example, triethylamine or diisopropyl ethylamine.

5 “Suitable solvent system” include organic solvent, for example, alcohol, ethyl acetate, THF, DME, toluene, etc., or a mixture of water soluble organic solvent with water, for example, a mixture of DME and water, THF and water, or alcohol and water, more particularly a mixture of DME and water.

10 Particular Embodiments of the Invention

In a particular embodiment of the invention, the first step coupling reaction is carried out in the presence of Pd, PdCl_2 , $\text{Pd}(\text{PPh}_3)_4$, *pdf*, or $\text{Pd}(\text{OAc})_2$; more particularly $\text{Pd}(\text{OAc})_2$.

15 In another particular embodiment of the invention, the first step coupling reaction is carried out in the presence of $\text{Pd}(\text{OAc})_2$ and PPh_3 .

In another particular embodiment of the invention, the first step coupling reaction is carried out in the presence of about 0.01 mol% to about 0.5 mol% of $\text{Pd}(\text{OAc})_2$ and about 0.02 mol%
20 to about 1.0 mol% PPh_3 , more particularly, about 0.1 mol% of $\text{Pd}(\text{OAc})_2$ and about 0.2 mol% PPh_3 .

In another particular embodiment of the invention, the first step coupling reaction is carried out by adding 3-(1-carboxy-1-methyl-ethyl)-phenyl boronic acid to 4,6-dichloro-2-
25 methoxypyrimidine slowly over a period of time, preferably about 2-6 hrs, thus, significantly reducing the over reaction byproduct and thermal decomposition of boronic acid.

In another particular embodiment of the invention, the first step coupling reaction is carried out in the presence of a suitable base.

30

In another particular embodiment of the invention, the first step coupling reaction is carried out in the presence of a base, wherein the amount of the base is in the range of about 2-4 equivalents, more particularly, about 3.0 equivalents.

In another particular embodiment of the invention, the first step coupling reaction is carried out using about 1.0-2.0 equivalents of 4,6-dichloro-2-methoxypyrimidine, more particularly about 1.2 equivalents.

5

In another particular embodiment of the invention, the first step coupling reaction is carried out in the presence of a suitable solvent system.

In another particular embodiment of the invention, the first step coupling product, 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methylpropionic acid, is purified by:
10 removing excess 4,6-dichloro-2-methoxypyrimidine by a phase separation, adjusting pH of the phase containing 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methylpropionic acid to about 6.5 to about 7.5, more particularly about 7.2, and
15 extracting the phase containing 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methylpropionic acid with a water-immiscible organic solvent, for example, n-butyl acetate, ethyl acetate or toluene.

In another particular embodiment of the invention, wherein the Palladium catalyst is removed
20 from 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methyl-propionic acid prior to the second step, for example, by treating 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methyl-propionic acid with TMT and charcoal, in an organic solvent, such as n-butyl acetate at about 50-70°C, particularly, at about 70°C.

25 In another particular embodiment of the invention, the first step product, 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methyl-propionic acid, is used directly in the second step as a solution.

In another particular embodiment of the invention, the second step is carried out in water.

30

In another particular embodiment of the invention, the second step product, 2-(3-{6-[2-(2,4-dichlorophenyl)-ethylamino]-2-methoxypyrimidin-4-yl}-phenyl)-2-methylpropionic acid, is

purified by recrystallization from an organic solvent and water, for example, THF/water or DME/water, more particularly, DME/water.

In another particular embodiment of the invention, the compound of formula (A) is converted
5 into dihydrogen phosphate salt thereof in methanol.

The present invention offers an improved process for preparing 2-(3-{6-[2-(2,4-dichloro-
phenyl)-ethylamino]-2-methoxypyrimidin-4-yl}-phenyl)-2-methylpropionic acid in high
yields and high purities yet requiring no expensive and time consuming column
10 chromatographic purification as used in the '732 application. The present process uses a less
amount of and more stable catalyst. A simpler phase separation is used to separate the step-1
product from excess reagents, and the product purification is achieved by a pH adjustment.
The residual Pd removal is also reduced to the desired level (5 ppm) in step-1, thus avoiding
handling the problem in the final stage since the final product is less soluble in common
15 organic solvents, making it harder to operate. Furthermore, the step-1 product is concatenated
into step-2 reaction without isolation. Overall, the present invention provides a much more
efficient process for preparing the desired product in a high purity (>99% pure by HPLC) and
high time-space yield. The salt formation carried out in methanol instead of THF as disclosed
in the '378 application also eliminates the residual solvent issue.


20

Examples

The present invention may be better understood by reference to the following non-limiting
Examples, which are exemplary of the invention. They should in no way be construed,
25 however, as limiting the broad scope of the invention.

Example 1:

2-(3-{6-[2-(2,4-Dichlorophenyl)-ethylamino]-2-methoxypyrimidin-4-yl}-phenyl)-2-
methylpropionic acid



Step 1: To an 1-L reactor are added 4,6-dichloro-2-methoxy-pyrimidine (50.1 g), K₂CO₃ (96.4
 5 g.), Pd(OAc)₂ (48 mg, 0.1 mol%) and PPh₃ (120 mg, 0.2 mol%) in DME/water (100 mL/200 mL). The mixture is heated to 50°C, while 3-(1-carboxy-1-methyl-ethyl)-phenyl boronic acid (51.6 g, 94wt% purity) in DME (50 mL)/water (100 mL) is added over 2 h via a metering pump. The reaction is held at 50°C for an additional hour. The mixture is cooled to 25°C and toluene (250 mL) is added. The mixture is stirred for 15 min and the layers are separated. To
 10 the aqueous layer is added n-BuOAc (300 mL) and pH is adjusted to 7.2 with 4 M aqueous HCl (~ 190 mL). The aqueous layer is extracted with n-BuOAc (2 x 300 mL). The combined organic layers are treated with TMT (3.2 g) and charcoal (6.4 g) at 70°C for 3 h. The mixture is allowed to cool to 25°C and filtered through celite. The filtrate is extracted twice with an aqueous K₂CO₃ solution (31.8 g of K₂CO₃ in 320 mL of water). The combined aqueous
 15 layers are used directly in the next step.

Step 2: To the above aqueous layer is added 2,4-dichlorophenethylamine (43.5 g). The resulting mixture is heated to 95°C for 6 h. The mixture is cooled to 50°C and n-BuOAc (150 mL) is added. The pH is adjusted to ~ 4.94 with an aqueous solution of 4 M HCl (~ 204 mL).
 20 The mixture is then cooled to 25°C over 30 min and further cooled to 4°C over 30 min. The mixture is filtered and the wet cake is washed with water (2 x 200 mL) and dried to afford 2-(3-(6-[2-(2,4-dichloro-phenyl)-ethylamino]-2-methoxy-pyrimidin-4-yl)-phenyl)-2-methyl-

propionic acid (76.2 g, 97.5% purity by HPLC), which is further purified by recrystallization from DME/water to provide over 99% purity by HPLC (see FIGURE 1).

Example 2:

5 Dihydrogen phosphate salt of 2-(3-{6-[2-(2,4-dichlorophenyl)-ethylamino]-2-methoxy-pyrimidin-4-yl}-phenyl)-2-methylpropionic acid

A 1-L jacketed reactor is charged with 2-(3-{6-[2-(2,4-dichloro-phenyl)-ethylamino]-2-methoxy-pyrimidin-4-yl}-phenyl)-2-methylpropionic acid (50.0 g, 0.109 mol) and methanol (500 mL). The mixture is heated to 37.5±2 °C. A 1:10 phosphoric acid:MeOH solution (110
10 mL) is then added to the mixture over 3 hrs. The mixture is then cooled to 10°C and the temperature is maintained for 0.5 hrs. The solid is collected by filtration and the filter cake is washed with MeOH (120 mL) and dried to give 2-(3-{6-[2-(2,4-dichloro-phenyl)-ethylamino]-2-methoxy-pyrimidin-4-yl}-phenyl)-2-methylpropionic acid phosphoric acid salt (57.9 g, 95.6% yield). Gas Chromatography shows that the product contains 201 ppm residual
15 methanol, well below the limit of ICH guideline of 3000 ppm.

The purity of the compound is analyzed by HPLC using the following conditions:

Instrument: Agilent 1100 series HPLC

Column: Phenomenex Synergi 4µ Hydro-RP, 150 x 4.6mm

20 Conditions: Mobile phase: A: 0.1%TFA in acetonitrile; B: 0.1% TFA in water

Flow rate: 1.5 mL/min

Detector: 220 nm

Injection: 10 µL

Temperature: 25°C

25 Run time: 18 min

Gradient:

Time (min)	Mobile Phase A	Mobile Phase B
0	30	70
2.0	30	70
15.0	70	30
18.0	90	10


Sample preparation:

30 Dissolve testing sample in water/0.1% TFA in acetonitrile (20/80).

We claim:

1. A process for preparing 2-(3-{6-[2-(2,4-dichloro-phenyl)-ethylamino]-2-methoxy-pyrimidin-4-yl}-phenyl)-2-methyl-propionic acid of formula (A),

5



(A)

comprising:

- (1) a coupling 4,6-dichloro-2-methoxy-pyrimidine with 3-(1-carboxy-1-methyl-ethyl)-phenyl boronic acid in the presence of a suitable Palladium catalyst to provide 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methyl-propionic acid,
- 10 (2) coupling 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methyl-propionic acid with 2,4-dichlorophenethylamine to provide 2-(3-{6-[2-(2,4-dichloro-phenyl)-ethylamino]-2-methoxy-pyrimidin-4-yl}-phenyl)-2-methyl-propionic acid.

15

2. The process according to claim 1, wherein the suitable Palladium catalyst is Pd, PdCl₂, Pd(PPh₃)₄, *pddf*, or Pd(OAc)₂.

3. The process according to claim 1, wherein the suitable Palladium catalyst is Pd(OAc)₂.

20

4. The process according to claim 3, wherein the coupling of 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methyl-propionic acid with 2,4-dichlorophenethylamine is carried out in the presence of PPh₃.

25

5. The process according to claim 4, wherein the amount of Pd(OAc)₂ is about 0.01 mol% to about 0.5 mol%, and the amount of PPh₃ is about 0.02 mol% to about 1.0 mol%.

6. The process according to claim 4, wherein the amount of Pd(OAc)₂ is about 0.01 mol%, and the amount of PPh₃ is about 0.02 mol%.

7. The process according to claim 1, wherein the first step coupling reaction is carried out by adding 3-(1-carboxy-1-methyl-ethyl)-phenyl boronic acid to 4,6-dichloro-2-methoxypyrimidine over a period of time of about 2-6 hours.

8. The process according to claim 1, wherein the first step coupling reaction is carried out in the presence of a suitable base.

9. The process according to claim 1, wherein the first step coupling reaction is carried out in the presence of K₂CO₃.

10. The process according to claim 8, wherein the amount of the base is about 2 to 4 equivalents.

11. The process according to claim 8, wherein the amount of the base is about 3.0 equivalents.

12. The process according to claim 1, wherein the first step coupling reaction is carried out using about 1.0 to about 2.0 equivalents of 4,6-dichloro-2-methoxypyrimidine.

13. The process according to claim 1, wherein the first step coupling reaction is carried out using about 1.2 equivalents of 4,6-dichloro-2-methoxypyrimidine.

14. The process according to claim 1, wherein the first step coupling reaction is carried out in the presence of a suitable solvent system.

15. The process according to claim 1, wherein the first step coupling reaction is carried out in a mixture of 1,2-dimethoxyethane and water.

16. The process according to claim 1, wherein before coupling 2-[3-(6-chloro-2-hydroxypyrimidin-4-yl)-phenyl]-2-methyl-propionic acid with 2,4-dichlorophenethylamine, 2-[3-(6-

chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methylpropionic acid is purified by steps comprising:

removing excess 4,6-dichloro-2-methoxypyrimidine by a phase separation,
adjusting pH of the phase containing 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-
5 phenyl]-2-methylpropionic acid to about 6.5 to about 7.5, more particularly about 7.2,
and
extracting the phase containing 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-
methylpropionic acid with a water-immiscible organic solvent, for example, n-butyl
acetate, ethyl acetate or toluene.

10

17. The process according to claim 1, wherein the Palladium catalyst is removed from 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methyl-propionic acid before coupling 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methyl-propionic acid with 2,4-dichlorophenethylamine.

15

18. The process according to claim 17, wherein the Palladium catalyst is removed by treating 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methyl-propionic acid with TMT and charcoal.

20

19. The process according to claim 17, wherein the Palladium catalyst is removed by treating 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methyl-propionic acid with TMT and charcoal in an organic solvent at about 50-70°C.

25

20. The process according to claim 1, wherein 2-[3-(6-chloro-2-hydroxy-pyrimidin-4-yl)-phenyl]-2-methyl-propionic acid is used directly in the second step as a solution.

21. The process according to claim 1, wherein the second step is carried out in water.

30

22. The process according to claim 1, further comprising purifying 2-(3-(6-[2-(2,4-dichlorophenyl)-ethylamino]-2-methoxypyrimidin-4-yl)-phenyl)-2-methylpropionic acid by recrystallization from an organic solvent and water.

23. The process according to claim 1, further comprising purifying 2-(3-{6-[2-(2,4-dichlorophenyl)-ethylamino]-2-methoxypyrimidin-4-yl}-phenyl)-2-methylpropionic acid by recrystallization from 1,2-dimethoxyethane and water.
- 5 24. A process for preparing dihydrogen phosphate salt of 2-(3-{6-[2-(2,4-dichlorophenyl)-ethylamino]-2-methoxypyrimidin-4-yl}-phenyl)-2-methylpropionic acid, comprising reacting 2-(3-{6-[2-(2,4-dichlorophenyl)-ethylamino]-2-methoxypyrimidin-4-yl}-phenyl)-2-methylpropionic acid with phosphoric acid in methanol.
- 10 25. A compound, which is 2-[3-(6-chloro-2-hydroxypyrimidin-4-yl)-phenyl]-2-methylpropionic acid, or a salt thereof.
26. The compound according to claim 25, which is 2-[3-(6-chloro-2-hydroxypyrimidin-4-yl)-phenyl]-2-methylpropionic acid.

