



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N		A2	(11) International Publication Number: WO 00/50562
			(43) International Publication Date: 31 August 2000 (31.08.00)
(21) International Application Number: PCT/US00/04413			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 22 February 2000 (22.02.00)			
(30) Priority Data: 09/255,376 22 February 1999 (22.02.99) US 09/387,699 13 August 1999 (13.08.99) US			
(71) Applicant: SYNAPTIC PHARMACEUTICAL CORPORATION [US/US]; 215 College Road, Paramus, NJ 07652-1431 (US).			
(72) Inventors: BONINI, James, A.; 80 Manito Avenue, Oakland, NJ 07436 (US). BOROWSKY, Beth, E.; 218 Park Street, Montclair, NJ 07042 (US). ADHAM, Nika; 301 Mastin Place, Ridgewood, NJ 07450 (US). BOYLE, Noel; 369 Edgewater Road, Apt. 2, Cliffside Park, NJ 07010 (US). THOMPSON, Thelma, O.; 77 Brook Avenue, Apt. G20, Passaic Park, NJ 07055 (US).			(Published) <i>Without international search report and to be republished upon receipt of that report.</i>
(74) Agent: WHITE, John, P.; Cooper & Dunham LLP, 1185 Avenue of the Americas, New York, NY 10036 (US).			

(54) Title: DNA ENCODING SNORF25 RECEPTOR

(57) Abstract

This invention provides isolated nucleic acids encoding mammalian SNORF25 receptors, purified mammalian SNORF25 receptors, vectors comprising nucleic acid encoding mammalian SNORF25 receptors, cells comprising such vectors, antibodies directed to mammalian SNORF25 receptors, nucleic acid probes useful for detecting nucleic acid encoding mammalian SNORF25 receptors, antisense oligonucleotides complementary to unique sequences of nucleic acid encoding mammalian SNORF25 receptors, transgenic, nonhuman animals which express DNA encoding normal or mutant mammalian SNORF25 receptors, methods of isolating mammalian SNORF25 receptors, methods of treating an abnormality that is linked to the activity of the mammalian SNORF25 receptors, as well as methods of determining binding of compounds to mammalian SNORF25 receptors, methods of identifying agonists and antagonists of SNORF25 receptors, and agonists and antagonists so identified.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

DNA ENCODING SNORF25 RECEPTOR5 **BACKGROUND OF THE INVENTION**

This application claims priority of U.S. Serial No. 09/387,699, filed August 13, 1999, which is a continuation-in-part of U.S. Serial No. 09/255,376, filed February 22, 1999, the contents of which are hereby incorporated by 10 reference into the subject application.

Throughout this application various publications are referred to by partial citations within parentheses. Full citations for these publications may be found at the end of the 15 specification immediately preceding the claims. The disclosures of these publications, in their entireties, are hereby incorporated by reference into this application in order to more fully describe the state of the art to which the invention pertains.

20

Neuroregulators comprise a diverse group of natural products that subserve or modulate communication in the nervous system. They include, but are not limited to, neuropeptides, amino acids, biogenic amines, lipids, and lipid metabolites, 25 and other metabolic byproducts. Many of these neuroregulator substances interact with specific cell surface receptors, which transduce signals from the outside to the inside of the cell. G-protein coupled receptors (GPCRs) represent a major class of cell surface receptors with which many 30 neurotransmitters interact to mediate their effects. GPCRs are characterized by seven membrane-spanning domains and are coupled to their effectors via G-proteins linking receptor activation with intracellular biochemical sequelae such as stimulation of adenylyl cyclase.

35

Vitamin A₁ (all-trans-retinol) is oxidized to vitamin A₁ aldehyde (all-trans-retinal) by an alcohol dehydrogenase.

All-*trans*-retinal is critical for the synthesis of rhodopsin in retinal cells, where it plays a key role in the visual system. All-*trans*-retinal can also be converted to all-*trans*-retinoic acid (ATRA) by aldehyde dehydrogenase and 5 oxidase in other cell types (Bowman, W.C. and Rand, M.J., 1980).

Historically, ATRA and the other active metabolites of vitamin A, 9-*cis*-retinoic acid (9CRA), were thought to only 10 mediate their cellular effects through the action of nuclear retinoic acid receptors (RAR α , β , γ) and retinoid X receptors (RXR α , β , γ) (Mangelsdorf, D.J., et al, 1994). These receptors are members of a superfamily of ligand-dependent transcription factors, which include the vitamin D receptor 15 (VDR), thyroid hormone receptor (TR), and peroxisome proliferator activator receptors (PPAR). They form heterodimers and homodimers that bind to DNA response elements in the absence of ligand. In response to ligand binding the dimer changes conformation which leads to 20 transactivation and regulation of transcription of a set(s) of cell type-specific genes (Mangelsdorf, D.J., et al, 1994; Hofman, C. and Eichele, G., 1994; and Gudas, L.J. et al, 1994).

25 Since retinoic acid produces a wide variety of biological effects, it is not surprising that it is proposed to play an important role in various physiological and pathophysiological processes. Retinoids control critical physiological events including cell growth, differentiation, 30 reproduction, metabolism, and hematopoiesis in a wide variety of tissues. At a cellular level, retinoids are capable of inhibiting cell proliferation, inducing differentiation, and inducing apoptosis (Breitman, T. et al, 1980; Sporn, M. and Roberts, A., 1984, and Martin, S., et al, 1990). These 35 diverse effects of retinoid treatment prompted a series of investigations evaluating retinoids for cancer chemotherapy as well as cancer chemoprevention. Clinically, retinoids are

used for the treatment of a wide variety of malignant diseases including: acute promyelocytic leukemia (APL), cutaneous T-cell malignancies, dermatological malignancies, squamous cell carcinomas of skin and of the cervix and 5 neuroblastomas (Redfern, C.P. et al, 1995 for review). Retinoids have also been examined for their ability to suppress carcinogenesis and prevent development of invasive cancer. 13-cis retinoic acid reverses oral leukoplakia, the most common premalignant lesion of the aerodigestive tract, 10 and is also used in the chemoprevention of bladder cancer (Sabichi, A.L. et al, 1998, for review). Also, 13-cis retinoic acid treatment as adjuvant therapy after surgery and radiation in head and neck cancer caused a significant delay in the occurrence of second primary cancers (Gottardis, M.M. 15 et al, 1996, for review).

Interestingly, retinoids also have an effect on pancreatic function. It has been demonstrated that retinoic acid (or retinol) is required for insulin secretion from isolated 20 islets (Chertow, B.S., et al, 1987) and from RINm5F rat insulinoma cells (Chertow, B.S., et al, 1989). Retinoic acid may also have an effect on cell-to-cell adhesion and aggregation (Chertow, B.S., et al, 1983). In addition, a single intragastric administration of 9CRA (but not ATRA) 25 induced a wave of DNA synthesis in the pancreatic acinar cells and in the proximal tubular epithelial cells of the kidneys (Ohmura, T., et al, 1997). Therefore, retinoic acid could play a role in the normal pancreatic function and possibly in the development of diabetes. There is also some 30 evidence that retinoids could be useful in the treatment of pancreatic malignancies (El-Metwally, T.H. et al, 1999; Rosenwicz, S. et al, 1997; and Rosenwicz, S. et al, 1995).

Retinoids have been shown to affect epidermal cell growth and 35 differentiation as well as sebaceous gland activity and exhibit immunomodulatory and anti-inflammatory properties. Therefore, retinoids have been increasingly used for treatment of a variety of skin disorders including: psoriasis

and other hyperkeratotic and parakeratotic skin disorders, keratotic genodermatoses, severe acne and acne-related dermatoses, and also for therapy and/or chemoprevention of skin cancer and other neoplasia (Orfanos, C.E., et al, 1997 5 for review).

Retinoids are also involved in lung development. Fetal lung branching leading to development of the alveolar tree is accelerated by retinoic acid. Currently, prematurely 10 delivered infants who have immature lungs are treated with vitamin A, but other applications may exist that require further investigation (Chytil, F., 1996).

Lastly, there is some evidence that suggests that retinoids 15 may play a role in schizophrenia (Goodman, A.B. 1998) and Alzheimer's disease (Connor, M.J. and Sidell, N., 1997).

The extensive list of retinoid-mediated effects indicate that 20 retinoic acid receptors (non-nuclear) are attractive as targets for therapeutic intervention for several disorders and would be useful in developing drugs with higher specificity and fewer side effects for a wide variety of diseases.

SUMMARY OF THE INVENTION

This invention provides an isolated nucleic acid encoding a mammalian SNORF25 receptor.

5

This invention further provides a purified mammalian SNORF25 receptor protein.

10 This invention also provides a vector comprising a nucleic acid in accordance with this invention.

This invention still further provides a cell comprising a vector in accordance with this invention.

15 This invention additionally provides a membrane preparation isolated from a cell in accordance with this invention.

Furthermore, this invention provides a nucleic acid probe comprising at least 15 nucleotides, which probe specifically 20 hybridizes with a nucleic acid encoding a mammalian SNORF25 receptor, wherein the probe has a sequence complementary to a unique sequence present within one of the two strands of the nucleic acid encoding the mammalian SNORF25 receptor contained in plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 25 203495).

This invention further provides a nucleic acid probe comprising at least 15 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF25 receptor, wherein the probe has a sequence complementary to a unique sequence present within (a) the nucleic acid sequence shown in Figures 1A-1B (SEQ ID NO: 1) or (b) the reverse complement thereof.

35 This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to RNA encoding a mammalian SNORF25 receptor, so as to prevent translation of such RNA.

This invention further provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to genomic DNA encoding a mammalian SNORF25 receptor, so as to prevent transcription of such genomic DNA.

5

This invention also provides an antibody capable of binding to a mammalian SNORF25 receptor encoded by a nucleic acid in accordance with this invention.

10 Moreover, this invention provides an agent capable of competitively inhibiting the binding of an antibody in accordance with this invention to a mammalian SNORF25 receptor.

15 This invention still further provides a pharmaceutical composition comprising (a) an amount of an oligonucleotide in accordance with this invention capable of passing through a cell membrane and effective to reduce expression of a mammalian SNORF25 receptor and (b) a pharmaceutically acceptable carrier capable of passing through the cell membrane.

25 This invention also provides a pharmaceutical composition which comprises an amount of an antibody in accordance with this invention effective to block binding of a ligand to a human SNORF25 receptor and a pharmaceutically acceptable carrier.

30 This invention further provides a transgenic, nonhuman mammal expressing DNA encoding a mammalian SNORF25 receptor in accordance with this invention.

35 This invention still further provides a transgenic, nonhuman mammal comprising a homologous recombination knockout of a native mammalian SNORF25 receptor.

This invention further provides a transgenic, nonhuman mammal whose genome comprises antisense DNA complementary to DNA

encoding a mammalian SNORF25 receptor in accordance with this invention so placed within such genome as to be transcribed into antisense mRNA which is complementary to and hybridizes with mRNA encoding the mammalian SNORF25 receptor so as to 5 reduce translation of of such mRNA and expression of such receptor.

This invention provides a process for identifying a chemical compound which specifically binds to a mammalian SNORF25 10 receptor which comprises contacting cells containing DNA encoding, and expressing on their cell surface, the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, with the compound under conditions suitable for binding, and detecting specific 15 binding of the chemical compound to the mammalian SNORF25 receptor.

This invention further provides a process for identifying a chemical compound which specifically binds to a mammalian 20 SNORF25 receptor which comprises contacting a membrane preparation from cells containing DNA encoding, and expressing on their cell surface, the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, with the compound under 25 conditions suitable for binding, and detecting specific binding of the chemical compound to the mammalian SNORF25 receptor.

This invention still further provides a process involving 30 competitive binding for identifying a chemical compound which specifically binds to a mammalian SNORF25 receptor which comprises separately contacting cells expressing on their cell surface the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, 35 with both the chemical compound and a second chemical compound known to bind to the receptor, and with only the second chemical compound, under conditions suitable for binding of such compounds to the receptor, and detecting

specific binding of the chemical compound to the mammalian SNORF25 receptor, a decrease in the binding of the second chemical compound to the mammalian SNORF25 receptor in the presence of the chemical compound being tested indicating 5 that such chemical compound binds to the mammalian SNORF25 receptor.

This invention further provides a process involving competitive binding for identifying a chemical compound which 10 specifically binds to a mammalian SNORF25 receptor which comprises separately contacting a membrane preparation from cells expressing on their cell surface the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, with both the chemical compound 15 and a second chemical compound known to bind to the receptor, and with only the second chemical compound, under conditions suitable for binding of such compounds to the receptor, and detecting specific binding of the chemical compound to the mammalian SNORF25 receptor, a decrease in the binding of the 20 second chemical compound to the mammalian SNORF25 receptor in the presence of the chemical compound indicating that the chemical compound binds to the mammalian SNORF25 receptor.

This invention further provides a compound identified by one 25 of the processes of this invention.

This invention provides a method of screening a plurality of chemical compounds not known to bind to a mammalian SNORF25 receptor to identify a compound which specifically binds to 30 the mammalian SNORF25 receptor, which comprises (a) contacting cells transfected with, and expressing, DNA encoding the mammalian SNORF25 receptor with a compound known to bind specifically to the mammalian SNORF25 receptor; (b) contacting the cells of step (a) with the plurality of compounds not 35 known to bind specifically to the mammalian SNORF25 receptor, under conditions permitting binding of compounds known to bind to the mammalian SNORF25 receptor; (c) determining whether the binding of the compound known to bind to the

mammalian SNORF25 receptor is reduced in the presence of the plurality of compounds, relative to the binding of the compound in the absence of the plurality of compounds; and if so (d) separately determining the binding to the mammalian SNORF25 receptor of each compound included in the plurality of compounds, so as to thereby identify any compound included therein which specifically binds to the mammalian SNORF25 receptor.

This invention further provides a method of screening a plurality of chemical compounds not known to bind to a mammalian SNORF25 receptor to identify a compound which specifically binds to the mammalian SNORF25 receptor, which comprises (a) contacting a membrane preparation from cells transfected with, and expressing, DNA encoding the mammalian SNORF25 receptor with the plurality of compounds not known to bind specifically to the mammalian SNORF25 receptor under conditions permitting binding of compounds known to bind to the mammalian SNORF25 receptor; (b) determining whether the binding of a compound known to bind to the mammalian SNORF25 receptor is reduced in the presence of any compound within the plurality of compounds, relative to the binding of the compound in the absence of the plurality of compounds; and if so (c) separately determining the binding to the mammalian SNORF25 receptor of each compound included in the plurality of compounds, so as to thereby identify any compound included therein which specifically binds to the mammalian SNORF25 receptor.

This invention also provides a method of detecting expression of a mammalian SNORF25 receptor by detecting the presence of mRNA coding for the mammalian SNORF25 receptor which comprises obtaining total mRNA from the cell and contacting the mRNA so obtained with a nucleic acid probe according to this invention under hybridizing conditions, detecting the presence of mRNA hybridized to the probe, and thereby detecting the expression of the mammalian SNORF25 receptor by the cell.

This invention further provides a method of detecting the presence of a mammalian SNORF25 receptor on the surface of a cell which comprises contacting the cell with an antibody according to this invention under conditions permitting 5 binding of the antibody to the receptor, detecting the presence of the antibody bound to the cell, and thereby detecting the presence of the mammalian SNORF25 receptor on the surface of the cell.

10 This invention still further provides a method of determining the physiological effects of varying levels of activity of mammalian SNORF25 receptors which comprises producing a transgenic, nonhuman mammal in accordance with this invention whose levels of mammalian SNORF25 receptor activity are 15 varied by use of an inducible promoter which regulates mammalian SNORF25 receptor expression.

This invention additionally provides a method of determining the physiological effects of varying levels of activity of 20 mammalian SNORF25 receptors which comprises producing a panel of transgenic, nonhuman mammals in accordance with this invention each expressing a different amount of mammalian SNORF25 receptor.

25 Moreover, this invention provides a method for identifying an antagonist capable of alleviating an abnormality wherein the abnormality is alleviated by decreasing the activity of a mammalian SNORF25 receptor comprising administering a compound to a transgenic, nonhuman mammal according to this 30 invention, and determining whether the compound alleviates any physiological and/or behavioral abnormality displayed by the transgenic, nonhuman mammal as a result of overactivity of a mammalian SNORF25 receptor, the alleviation of such an abnormality identifying the compound as an antagonist.

35

This invention also provides an antagonist identified by the preceding method.

This invention further provides a composition, e.g. a pharmaceutical composition, comprising an antagonist according to this invention and a carrier, e.g. a pharmaceutically acceptable carrier.

5

This invention additionally provides a method of treating an abnormality in a subject wherein the abnormality is alleviated by decreasing the activity of a mammalian SNORF25 receptor which comprises administering to the subject an effective amount of the pharmaceutical composition according to this invention so as to thereby treat the abnormality.

In addition, this invention provides a method for identifying an agonist capable of alleviating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian SNORF25 receptor comprising administering a compound to a transgenic, nonhuman mammal according to this invention, and determining whether the compound alleviates any physiological and/or behavioral abnormality displayed by the transgenic, nonhuman mammal, the alleviation of such an abnormality identifying the compound as an agonist.

This invention further provides an agonist identified by the preceding method.

This invention still further provides a composition, e.g. a pharmaceutical composition, comprising an agonist according to this invention and a carrier, e.g. pharmaceutically acceptable carrier.

Moreover, this invention provides a method of treating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian SNORF25 receptor which comprises administering to the subject an effective amount of the pharmaceutical composition according to this invention so as to thereby treat the abnormality.

Yet further, this invention provides a method for diagnosing a predisposition to a disorder associated with the activity of a specific mammalian allele which comprises: (a) obtaining DNA of subjects suffering from the disorder; 5 (b) performing a restriction digest of the DNA with a panel of restriction enzymes; (c) electrophoretically separating the resulting DNA fragments on a sizing gel; (d) contacting the resulting gel with a nucleic acid probe capable of specifically hybridizing with a unique sequence included 10 within the sequence of a nucleic acid molecule encoding a mammalian SNORF25 receptor and labeled with a detectable marker; (e) detecting labeled bands which have hybridized to the DNA encoding a mammalian SNORF25 receptor to create a unique band pattern specific to the DNA of subjects suffering 15 from the disorder; (f) repeating steps (a)-(e) with DNA obtained for diagnosis from subjects not yet suffering from the disorder; and (g) comparing the unique band pattern specific to the DNA of subjects suffering from the disorder from step (e) with the band pattern from step (f) for 20 subjects not yet suffering from the disorder so as to determine whether the patterns are the same or different and thereby diagnose predisposition to the disorder if the patterns are the same.

25 This invention also provides a method of preparing a purified mammalian SNORF25 receptor according to the invention which comprises: (a) culturing cells which express the mammalian SNORF25 receptor; (b) recovering the mammalian SNORF25 receptor from the cells; and (c) purifying the mammalian 30 SNORF25 receptor so recovered.

This invention further provides a method of preparing the purified mammalian SNORF25 receptor according to the invention which comprises: (a) inserting a nucleic acid 35 encoding the mammalian SNORF25 receptor into a suitable expression vector; (b) introducing the resulting vector into a suitable host cell; (c) placing the resulting host cell in suitable conditions permitting the production of the

mammalian SNORF25 receptor; (d) recovering the mammalian SNORF25 receptor so produced; and optionally (e) isolating and/or purifying the mammalian SNORF25 receptor so recovered.

5 Furthermore, this invention provides a process for determining whether a chemical compound is a mammalian SNORF25 receptor agonist which comprises contacting cells transfected with and expressing DNA encoding the mammalian SNORF25 receptor with the compound under conditions
10 permitting the activation of the mammalian SNORF25 receptor, and detecting any increase in mammalian SNORF25 receptor activity, so as to thereby determine whether the compound is a mammalian SNORF25 receptor agonist.

15 This invention also provides a process for determining whether a chemical compound is a mammalian SNORF25 receptor antagonist which comprises contacting cells transfected with and expressing DNA encoding the mammalian SNORF25 receptor with the compound in the presence of a known mammalian
20 SNORF25 receptor agonist, under conditions permitting the activation of the mammalian SNORF25 receptor, and detecting any decrease in mammalian SNORF25 receptor activity, so as to thereby determine whether the compound is a mammalian SNORF25 receptor antagonist.

25 This invention still further provides a composition, for example a pharmaceutical composition, which comprises an amount of a mammalian SNORF25 receptor agonist determined by a process according to this invention effective to increase
30 activity of a mammalian SNORF25 receptor and a carrier, for example, a pharmaceutically acceptable carrier.

Also, this invention provides a composition, for example a pharmaceutical composition, which comprises an amount of a
35 mammalian SNORF25 receptor antagonist determined by a process according to this invention effective to reduce activity of a mammalian SNORF25 receptor and a carrier, for example, a pharmaceutically acceptable carrier.

This invention moreover provides a process for determining whether a chemical compound specifically binds to and activates a mammalian SNORF25 receptor, which comprises contacting cells producing a second messenger response and 5 expressing on their cell surface the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, with the chemical compound under conditions suitable for activation of the mammalian SNORF25 receptor, and measuring the second messenger response in the 10 presence and in the absence of the chemical compound, a change, e.g. an increase, in the second messenger response in the presence of the chemical compound indicating that the compound activates the mammalian SNORF25 receptor.

15 This invention still further provides a process for determining whether a chemical compound specifically binds to and inhibits activation of a mammalian SNORF25 receptor, which comprises separately contacting cells producing a second messenger response and expressing on their cell 20 surface the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, with both the chemical compound and a second chemical compound known to activate the mammalian SNORF25 receptor, and with only the second chemical compound, under conditions suitable 25 for activation of the mammalian SNORF25 receptor, and measuring the second messenger response in the presence of only the second chemical compound and in the presence of both the second chemical compound and the chemical compound, a smaller change, e.g. increase, in the second messenger 30 response in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound indicating that the chemical compound inhibits activation of the mammalian SNORF25 receptor.

35

Further, this invention provides a compound determined by a process according to the invention and a composition, for example, a pharmaceutical composition, which comprises an

amount of a mammalian SNORF25 receptor agonist determined to be such by a process according to the invention, effective to increase activity of the mammalian SNORF25 receptor and a carrier, for example, a pharmaceutically acceptable 5 carrier.

This invention also provides a composition, for example, a pharmaceutical composition, which comprises an amount of a mammalian SNORF25 receptor antagonist determined to be such 10 by a process according to the invention, effective to reduce activity of the mammalian SNORF25 receptor and a carrier, for example, a pharmaceutically acceptable carrier.

This invention yet further provides a method of screening a 15 plurality of chemical compounds not known to activate a mammalian SNORF25 receptor to identify a compound which activates the mammalian SNORF25 receptor which comprises: (a) contacting cells transfected with and expressing the mammalian SNORF25 receptor with the plurality of compounds 20 not known to activate the mammalian SNORF25 receptor, under conditions permitting activation of the mammalian SNORF25 receptor; (b) determining whether the activity of the mammalian SNORF25 receptor is increased in the presence of one or more the compounds; and if so (c) separately 25 determining whether the activation of the mammalian SNORF25 receptor is increased by any compound included in the plurality of compounds, so as to thereby identify each compound which activates the mammalian SNORF25 receptor.

30 This invention provides a method of screening a plurality of chemical compounds not known to inhibit the activation of a mammalian SNORF25 receptor to identify a compound which inhibits the activation of the mammalian SNORF25 receptor, which comprises: (a) contacting cells transfected with and 35 expressing the mammalian SNORF25 receptor with the plurality of compounds in the presence of a known mammalian SNORF25 receptor agonist, under conditions permitting activation of the mammalian SNORF25 receptor; (b) determining whether the

extent or amount of activation of the mammalian SNORF25 receptor is reduced in the presence of one or more of the compounds, relative to the extent or amount of activation of the mammalian SNORF25 receptor in the absence of such one or 5 more compounds; and if so (c) separately determining whether each such compound inhibits activation of the mammalian SNORF25 receptor for each compound included in the plurality of compounds, so as to thereby identify any compound included in such plurality of compounds which inhibits the activation 10 of the mammalian SNORF25 receptor.

This invention also provides a composition, for example a pharmaceutical composition, comprising a compound identified by a method according to this invention in an amount 15 effective to increase mammalian SNORF25 receptor activity and a carrier, for example, a pharmaceutically acceptable carrier.

This invention still further provides a composition, for 20 example, a pharmaceutical composition, comprising a compound identified by a method according to this invention in an amount effective to decrease mammalian SNORF25 receptor activity and a carrier, for example, a pharmaceutically acceptable carrier.

25 Furthermore, this invention provides a method of treating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian SNORF25 receptor which comprises administering to the subject a 30 compound which is a mammalian SNORF25 receptor agonist in an amount effective to treat the abnormality.

This invention additionally provides a method of treating an abnormality in a subject wherein the abnormality is 35 alleviated by decreasing the activity of a mammalian SNORF25 receptor which comprises administering to the subject a compound which is a mammalian SNORF25 receptor antagonist in an amount effective to treat the abnormality.

This invention also provides a process for making a composition of matter which specifically binds to a mammalian SNORF25 receptor which comprises identifying a chemical 5 compound using a process in accordance with this invention and then synthesizing the chemical compound or a novel structural and functional analog or homolog thereof.

This invention further provides a process for preparing a 10 composition, for example, a pharmaceutical composition which comprises admixing a carrier, for example, a pharmaceutically acceptable carrier, and a pharmaceutically effective amount of a chemical compound identified by a process in accordance with this invention or a novel structural and functional 15 analog or homolog thereof.

BRIEF DESCRIPTION OF THE FIGURESFigures 1A-1B

Nucleotide sequence including sequence encoding a human
5 SNORF25 receptor (SEQ ID NO: 1). Putative open reading
frames including the shortest open reading frame are
indicated by underlining one start (ATG) codon (at positions
61-63) and the stop codon (at positions 1066-1068). In
addition, partial 5' and 3' untranslated sequences are shown.

10

Figures 2A-2B

Deduced amino acid sequence (SEQ ID NO: 2) of the human
SNORF25 receptor encoded by the longest open reading frame
indicated in the nucleotide sequence shown in Figures 1A-1B
15 (SEQ ID NO: 1). The seven putative transmembrane (TM) regions
are underlined.

Figures 3A-3B

Nucleotide sequence including sequence encoding a rat SNORF25
20 receptor (SEQ ID NO: 3). Putative open reading frames
including the shortest open reading frame are indicated by
underlining one start (ATG) codon (at positions 49-51) and
the stop codon (at positions 1054-1056). In addition,
partial 5' and 3' untranslated sequences are shown.

25

Figures 4A-4B

Deduced amino acid sequence (SEQ ID NO: 4) of the rat SNORF25
receptor encoded by the longest open reading frame indicated
in the nucleotide sequence shown in Figures 3A-3B (SEQ ID NO:
30 3). The seven putative transmembrane (TM) regions are
underlined.

Figure 5

Comparison of basal cAMP levels of SNORF25-and
35 mock-transfected CHO cells. SNORF25 or empty vector (mock)
DNA was transfected into CHO cells as described in Materials
and Methods. The transfectants were plated into 96-well
plates, and assayed for cAMP release as described. The

results of a representative experiment are shown.

Figure 6

5 Modulation of cAMP release by ATRA, vitamin A₁ and forskolin in SNORF25-expressing mock-transfected CHO cells. The transfectants were plated into 96-well plates, challenged with 10 μ M concentrations of drugs and assayed for cAMP release as described. The results of a representative 10 experiment involving known cyclase stimulatory receptors are shown. Results are means \pm S.E.M of triplicate determinations with the exception of vitamin A₁ which is a single point. Results are normalized to % basal cAMP release.

15 Figure 7

Specificity of ATRA cAMP response in Cos-7 cells. The transfectants were plated into 96-well plates, challenged with 10 μ M concentrations of ATRA and assayed for cAMP release as described. The results of a representative 20 experiment are shown. Results are means \pm S.E.M of triplicate determinations.

Figure 8

ATRA Dose-response curve in transiently-transfected Cos-7 25 cells. A representative example of dose-response effect of ATRA to increase cAMP release in SNORF25- (■) and mock- (□) transfected cells.

Figures 9A-9C

30 Stimulation of CFTR by ATRA in oocytes expressing SNORF25. Voltage clamp recording from oocyte previously injected with SNORF25 receptor mRNA and CFTR (Figure 9A), and from control (CFTR alone) oocyte (Figure 9B). Application of epinephrine (1 μ M) evokes a similar current in other oocytes expressing 35 the B2 adrenergic receptor (B2AR) and CFTR (Figure 9C). Holding potential was -70 mV for all recordings.

Figure 10

Mean current amplitudes stimulated by ATRA (10 μ M) in control (CFTR alone) oocytes (n = 16) and oocytes injected with mRNA encoding SNORF25 and CFTR (n = 17).

DETAILED DESCRIPTION OF THE INVENTION

This invention provides a recombinant nucleic acid comprising a nucleic acid encoding a mammalian SNORF25 receptor, wherein

5 the mammalian receptor-encoding nucleic acid hybridizes under high stringency conditions to (a) a nucleic acid encoding a human SNORF25 receptor and having a sequence identical to the sequence of the human SNORF25 receptor-encoding nucleic acid contained in plasmid pEXJT3T7-hSNORF25 (ATCC Accession No.

10 203495) or (b) a nucleic acid encoding a rat SNORF25 receptor and having a sequence identical to the sequence of the rat SNORF25 receptor-encoding nucleic acid contained in plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494).

15 This invention further provides a recombinant nucleic acid comprising a nucleic acid encoding a human SNORF25 receptor, wherein the human SNORF25 receptor comprises an amino acid sequence identical to the sequence of the human SNORF25 receptor encoded by the shortest open reading frame indicated

20 in Figures 1A-1B (SEQ ID NO: 1).

This invention also provides a recombinant nucleic acid comprising a nucleic acid encoding a rat SNORF25 receptor, wherein the rat SNORF25 receptor comprises an amino acid sequence identical to the sequence of the rat SNORF25 receptor encoded by the shortest open reading frame indicated

25 in Figures 3A-3B (SEQ ID NO: 3).

Plasmid pEXJT3T7-hSNORF25 and plasmid pcDNA3.1-rSNORF25 were

30 both deposited on November 24, 1998, with the American Type Culture Collection (ATCC), 10801 University Blvd., Manassas, Virginia 20110-2209, U.S.A. under the provisions of the Budapest Treaty for the International Recognition of the Deposit of Microorganisms for the Purposes of Patent

35 Procedure and were accorded ATCC Accession Nos. 203495 and 203494, respectively.

Hybridization methods are well known to those of skill in the

art. For purposes of this invention, hybridization under high stringency conditions means hybridization performed at 40°C in a hybridization buffer containing 50% formamide, 5X SSC, 7mM Tris, 1X Denhardt's, 25 µg/ml salmon sperm DNA; wash at 5 50°C in 0.1X SSC, 0.1%SDS.

Throughout this application, the following standard abbreviations are used to indicate specific nucleotide bases:

A = adenine
10 G = guanine
C = cytosine
T = thymine
M = adenine or cytosine
R = adenine or guanine
15 W = adenine or thymine
S = cytosine or guanine
Y = cytosine or thymine
K = guanine or thymine
V = adenine, cytosine, or guanine (not thymine)
20 H = adenine, cytosine, or thymine (not cytosine)
B = cytosine, guanine, or thymine (not adenine)
N = adenine, cytosine, guanine, or thymine (or other
modified base such as inosine)
I = inosine
25

Furthermore, the term "agonist" is used throughout this application to indicate any peptide or non-peptidyl compound which increases the activity of any of the polypeptides of the subject invention. The term "antagonist" is used 30 throughout this application to indicate any peptide or non-peptidyl compound which decreases the activity of any of the polypeptides of the subject invention.

Furthermore, as used herein, the phrase "pharmaceutically acceptable carrier" means any of the standard pharmaceutically acceptable carriers. Examples include, but are not limited to, phosphate buffered saline, physiological saline, water, and emulsions, such as oil/water emulsions.

It is possible that the mammalian SNORF25 receptor gene contains introns and furthermore, the possibility exists that additional introns could exist in coding or non-coding regions. In addition, spliced form(s) of mRNA may encode 5 additional amino acids either upstream of the currently defined starting methionine or within the coding region. Further, the existence and use of alternative exons is possible, whereby the mRNA may encode different amino acids within the region comprising the exon. In addition, single 10 amino acid substitutions may arise via the mechanism of RNA editing such that the amino acid sequence of the expressed protein is different than that encoded by the original gene. (Burns, et al., 1996; Chu, et al., 1996). Such variants may exhibit pharmacologic properties differing from the 15 polypeptide encoded by the original gene.

This invention provides splice variants of the mammalian SNORF25 receptors disclosed herein. This invention further provides alternate translation initiation sites and 20 alternately spliced or edited variants of nucleic acids encoding the SNORF25 receptors in accordance with this invention.

This invention also contemplates recombinant nucleic acids 25 which comprise nucleic acids encoding naturally occurring allelic variants of the SNORF25 receptors disclosed herein.

The nucleic acids of the subject invention also include nucleic acid analogs of the human SNORF25 receptor genes, 30 wherein the human SNORF25 receptor gene comprises the nucleic acid sequence shown in Figures 1A-1B or contained in plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495). Nucleic acid analogs of the human SNORF25 receptor genes differ from the human SNORF25 receptor genes described herein in terms of the 35 identity or location of one or more nucleic acid bases (deletion analogs containing less than all of the nucleic acid bases shown in Figures 1A-1B or contained in plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495), substitution

analogs wherein one or more nucleic acid bases shown in Figures 1A-1B or contained in plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495), are replaced by other nucleic acid bases, and addition analogs, wherein one or more nucleic acid bases are added to a terminal or medial portion of the nucleic acid sequence) and which encode proteins which share some or all of the properties of the proteins encoded by the nucleic acid sequences shown in Figure 1A-1B or contained in plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495). In one embodiment of the present invention, the nucleic acid analog encodes a protein which has an amino acid sequence identical to that shown in Figures 2A-2B or encoded by the nucleic acid sequence contained in plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495). In another embodiment, the nucleic acid analog encodes a protein having an amino acid sequence which differs from the amino acid sequences shown in Figures 2A-2B or encoded by the nucleic acid contained in plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495). In a further embodiment, the protein encoded by the nucleic acid analog has a function which is the same as the function of the receptor proteins having the amino acid sequence shown in Figures 2A-2B. In another embodiment, the function of the protein encoded by the nucleic acid analog differs from the function of the receptor protein having the amino acid sequence shown in Figures 2A-2B. In another embodiment, the variation in the nucleic acid sequence occurs within the transmembrane (TM) region of the protein. In a further embodiment, the variation in the nucleic acid sequence occurs outside of the TM region.

30

The nucleic acids of the subject invention also include nucleic acid analogs of the rat SNORF25 receptor genes, wherein the rat SNORF25 receptor gene comprises the nucleic acid sequence shown in Figures 3A-3B or contained in plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494). Nucleic acid analogs of the rat SNORF25 receptor genes differ from the rat SNORF25 receptor genes described herein in terms of the identity or location of one or more nucleic acid bases

(deletion analogs containing less than all of the nucleic acid bases shown in Figures 3A-3B or contained in plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494) substitution analogs wherein one or more nucleic acid bases shown in Figures 3A-3B or contained in plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494), are replaced by other nucleic acid bases, and addition analogs, wherein one or more nucleic acid bases are added to a terminal or medial portion of the nucleic acid sequence) and which encode proteins which share some or all of the properties of the proteins encoded by the nucleic acid sequences shown in Figure 3A-3B or contained in plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494). In one embodiment of the present invention, the nucleic acid analog encodes a protein which has an amino acid sequence identical to that shown in Figures 4A-4B or encoded by the nucleic acid sequence contained in plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494). In another embodiment, the nucleic acid analog encodes a protein having an amino acid sequence which differs from the amino acid sequences shown in Figures 4A-4B or encoded by the nucleic acid contained in plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494). In a further embodiment, the protein encoded by the nucleic acid analog has a function which is the same as the function of the receptor proteins having the amino acid sequence shown in Figures 4A-4B. In another embodiment, the function of the protein encoded by the nucleic acid analog differs from the function of the receptor protein having the amino acid sequence shown in Figures 4A-4B. In another embodiment, the variation in the nucleic acid sequence occurs within the transmembrane (TM) region of the protein. In a further embodiment, the variation in the nucleic acid sequence occurs outside of the TM region.

This invention provides the above-described isolated nucleic acid, wherein the nucleic acid is DNA. In an embodiment, the DNA is cDNA. In another embodiment, the DNA is genomic DNA. In still another embodiment, the nucleic acid is RNA. Methods for production and manipulation of nucleic acid

molecules are well known in the art.

This invention further provides nucleic acid which is degenerate with respect to the DNA encoding any of the 5 polypeptides described herein. In an embodiment, the nucleic acid comprises a nucleotide sequence which is degenerate with respect to the nucleotide sequence shown in Figures 1A-1B (SEQ ID NO: 1) or the nucleotide sequence contained in the plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495), that 10 is, a nucleotide sequence which is translated into the same amino acid sequence.

This invention further provides nucleic acid which is degenerate with respect to the DNA encoding any of the 15 polypeptides described herein. In an embodiment, the nucleic acid comprises a nucleotide sequence which is degenerate with respect to the nucleotide sequence shown in Figures 3A-3B (SEQ ID NO: 3) or the nucleotide sequence contained in the plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494), that 20 is, a nucleotide sequence which is translated into the same amino acid sequence.

This invention also encompasses DNAs and cDNAs which encode amino acid sequences which differ from those of the 25 polypeptides according to this invention, but which should not produce phenotypic changes. Alternately, this invention also encompasses DNAs, cDNAs, and RNAs which hybridize with the DNA, cDNA, and RNA according to the subject invention. Hybridization methods are well known to those of skill in the 30 art.

The nucleic acids according to the subject invention also include nucleic acid molecules coding for polypeptide analogs, fragments or derivatives of antigenic polypeptides 35 which differ from naturally-occurring forms in terms of the identity or location of one or more amino acid residues (deletion analogs containing less than all of the residues specified for the protein, substitution analogs wherein one

or more residues specified are replaced by other residues and addition analogs wherein one or more amino acid residues is added to a terminal or medial portion of the polypeptides) and which share some or all properties of naturally-occurring 5 forms. These molecules include: the incorporation of codons "preferred" for expression by selected non-mammalian hosts; the provision of sites for cleavage by restriction endonuclease enzymes; and the provision of additional initial, terminal or intermediate DNA sequences that 10 facilitate construction of readily expressed vectors. The creation of polypeptide analogs is well known to those of skill in the art (Spurney, R. F. et al. (1997); Fong, T.M. et al. (1995); Underwood, D.J. et al. (1994); Graziano, M.P. et al. (1996); Guan X.M. et al. (1995)).

15

The modified polypeptides according to this invention may be transfected into cells either transiently or stably using methods well-known in the art, examples of which are disclosed herein. This invention also provides binding 20 assays using the modified polypeptides, in which the polypeptide is expressed either transiently or in stable cell lines. This invention further provides a compound identified using a modified polypeptide in a binding assay such as the binding assays described herein.

25

The nucleic acids described and claimed herein are useful for the information which they provide concerning the amino acid sequence of the polypeptide and as products for the large scale synthesis of the polypeptides by a variety of 30 recombinant techniques. The nucleic acid molecule is useful for generating new cloning and expression vectors, transformed and transfected prokaryotic and eukaryotic host cells, and new and useful methods for cultured growth of such host cells capable of expression of the polypeptide and 35 related products.

This invention also provides an isolated nucleic acid encoding species homologs of the SNORF25 receptors encoded

by the nucleic acid sequence shown in Figures 1A-1B (SEQ ID NO: 1) or encoded by the plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495). In one embodiment, the nucleic acid encodes a mammalian SNORF25 receptor homolog which has 5 substantially the same amino acid sequence as does the SNORF25 receptor encoded by the plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495). In another embodiment, the nucleic acid encodes a mammalian SNORF25 receptor homolog which has above 75% amino acid identity to the SNORF25 10 receptor encoded by the plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495); preferably above 85% amino acid identity to the SNORF25 receptor encoded by the plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495); most preferably above 95% amino acid identity to the SNORF25 15 receptor encoded by the plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495). In another embodiment, the mammalian SNORF25 receptor homolog has above 70% nucleic acid identity to the SNORF25 receptor gene contained in plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495); preferably 20 above 80% nucleic acid identity to the SNORF25 receptor gene contained in the plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495); more preferably above 90% nucleic acid identity to the SNORF25 receptor gene contained in the plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495). Examples of 25 methods for isolating and purifying species homologs are described elsewhere (e.g., U.S. Patent No. 5,602,024, WO94/14957, WO97/26853, WO98/15570).

This invention also provides an isolated nucleic acid 30 encoding species homologs of the SNORF25 receptors encoded by the nucleic acid sequence shown in Figures 3A-3B (SEQ ID NO: 3) or encoded by the plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494). In one embodiment, the nucleic acid encodes a mammalian SNORF25 receptor homolog which has 35 substantially the same amino acid sequence as does the SNORF25 receptor encoded by the plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494). In another embodiment, the nucleic acid encodes a mammalian SNORF25 receptor homolog

which has above 75% amino acid identity to the SNORF25 receptor encoded by the plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494); preferably above 85% amino acid identity to the SNORF25 receptor encoded by the plasmid 5 pcDNA3.1-rSNORF25 (ATCC Accession No. 203494); most preferably above 95% amino acid identity to the SNORF25 receptor encoded by the plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494). In another embodiment, the mammalian SNORF25 receptor homolog has above 70% nucleic acid identity 10 to the SNORF25 receptor gene contained in plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494); preferably above 80% nucleic acid identity to the SNORF25 receptor gene contained in the plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494); more preferably above 90% nucleic acid identity 15 to the SNORF25 receptor gene contained in the plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494).

This invention provides an isolated nucleic acid encoding a modified mammalian SNORF25 receptor, which differs from a 20 mammalian SNORF25 receptor by having an amino acid(s) deletion, replacement, or addition in the third intracellular domain.

This invention provides an isolated nucleic acid encoding a 25 mammalian SNORF25 receptor. In one embodiment, the nucleic acid is DNA. In another embodiment, the DNA is cDNA. In another embodiment, the DNA is genomic DNA. In another embodiment, the nucleic acid is RNA. In another embodiment, the mammalian SNORF25 receptor is a human SNORF25 receptor. 30 In another embodiment, the human SNORF25 receptor has an amino acid sequence identical to that encoded by the plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495). In another embodiment, the human SNORF25 receptor has an amino acid sequence identical to the amino acid sequence shown in 35 Figures 2A-2B (SEQ ID NO: 2).

In an embodiment, the mammalian SNORF25 receptor is a rat SNORF25 receptor. In another embodiment, the rat SNORF25

receptor has an amino acid sequence identical to that encoded by the plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494). In another embodiment, the rat SNORF25 receptor has an amino acid sequence identical to the amino acid sequence shown in 5 Figures 4A-4B (SEQ ID NO: 4).

This invention provides a purified mammalian SNORF25 receptor protein. In one embodiment, the SNORF25 receptor protein is a human SNORF25 receptor protein. In a further embodiment, 10 the SNORF25 receptor protein is a rat SNORF25 receptor protein.

This invention provides a vector comprising a nucleic acid in accordance with this invention. This invention further 15 provides a vector adapted for expression in a cell which comprises the regulatory elements necessary for expression of the nucleic acid in the cell operatively linked to the nucleic acid encoding the receptor so as to permit expression thereof, wherein the cell is a bacterial, amphibian, yeast, 20 insect or mammalian cell. In one embodiment, the vector is a baculovirus. In another embodiment, the vector is a plasmid.

This invention provides a plasmid designated 25 PEXJT3T7-hSNORF25 (ATCC Accession No. 203495). This invention also provides a plasmid designated pcDNA3.1-rSNORF25 (ATCC Accession No. 203494).

This invention further provides any vector or plasmid which 30 comprises modified untranslated sequences, which are beneficial for expression in desired host cells or for use in binding or functional assays. For example, a vector or plasmid with untranslated sequences of varying lengths may express differing amounts of the polypeptide depending upon 35 the host cell used. In an embodiment, the vector or plasmid comprises the coding sequence of the polypeptide and the regulatory elements necessary for expression in the host cell.

This invention provides a cell comprising a vector in accordance with this invention. In one embodiment, the cell is a non-mammalian cell. In one embodiment, the non-mammalian cell is a *Xenopus* oocyte cell or a *Xenopus* melanophore cell.

5 In another embodiment, the cell is a mammalian cell. In another embodiment, the cell is a COS-7 cell, a 293 human embryonic kidney cell, a NIH-3T3 cell, a LM(tk-) cell, a mouse Y1 cell, or a CHO cell. In another embodiment, the cell is an insect cell. In another embodiment, the insect cell is an Sf9 cell, an Sf21 cell or a *Trichoplusia ni* 5B-4 cell.

10

This invention provides a membrane preparation isolated from a cell in accordance with this invention.

15

Furthermore, this invention provides a nucleic acid probe comprising at least 15 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF25 receptor, wherein the probe has a sequence complementary to a unique sequence present within one of the two strands of the nucleic acid encoding the mammalian SNORF25 receptor contained in plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495) or plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494).

25

This invention further provides a nucleic acid probe comprising at least 15 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF25 receptor, wherein the probe has a sequence complementary to a unique sequence present within (a) the nucleic acid sequence shown in Figures 1A-1B (SEQ ID NO: 1) or (b) the reverse complement thereof. This invention also provides a nucleic acid probe comprising at least 15 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF25 receptor, wherein the probe has a sequence complementary to a unique sequence present within (a) the nucleic acid sequence shown in Figures 3A-3B (SEQ ID NO: 3) or (b) the reverse complement thereof. In one

embodiment, the nucleic acid is DNA. In another embodiment, the nucleic acid is RNA.

As used herein, the phrase "specifically hybridizing" means 5 the ability of a nucleic acid molecule to recognize a nucleic acid sequence complementary to its own and to form double-helical segments through hydrogen bonding between complementary base pairs.

- 10 The nucleic acids according to this invention may be used as probes to obtain homologous nucleic acids from other species and to detect the existence of nucleic acids having complementary sequences in samples.
- 15 The nucleic acids may also be used to express the receptors they encode in transfected cells.

The use of a constitutively active receptor encoded by SNORF25 either occurring naturally without further 20 modification or after appropriate point mutations, deletions or the like, allows screening for antagonists and *in vivo* use of such antagonists to attribute a role to receptor SNORF25 without prior knowledge of the endogenous ligand.

- 25 Use of the nucleic acids further enables elucidation of possible receptor diversity and of the existence of multiple subtypes within a family of receptors of which SNORF25 is a member.
- 30 Finally, it is contemplated that this receptor will serve as a valuable tool for designing drugs for treating various pathophysiological conditions such as chronic and acute inflammation, arthritis, autoimmune diseases, transplant rejection, graft vs. host disease, bacterial, fungal, 35 protozoan and viral infections, septicemia, AIDS, pain, psychotic and neurological disorders, including anxiety, depression, schizophrenia, dementia, mental retardation, memory loss, epilepsy, neurological disorders, neuromotor

disorders, respiratory disorders, asthma, eating/body weight disorders including obesity, bulimia, diabetes, anorexia, nausea, hypertension, hypotension, vascular and cardiovascular disorders, ischemia, stroke, cancers, ulcers, 5 urinary retention, sexual/reproductive disorders, circadian rhythm disorders, renal disorders, bone diseases including osteoporosis, benign prostatic hypertrophy, gastrointestinal disorders, nasal congestion, dermatological disorders such as psoriasis, allergies, Parkinson's disease, Alzheimer's 10 disease, acute heart failure, angina disorders, delirium, dyskinesias such as Huntington's disease or Gilles de la Tourette's syndrome, among others and diagnostic assays for such conditions. This receptor may also serve as a valuable tool for designing drugs for chemoprevention.

15

Methods of transfecting cells e.g. mammalian cells, with such nucleic acid to obtain cells in which the receptor is expressed on the surface of the cell are well known in the art. (See, for example, U.S. Patent Nos. 5,053,337; 20 5,155,218; 5,360,735; 5,472,866; 5,476,782; 5,516,653; 5,545,549; 5,556,753; 5,595,880; 5,602,024; 5,639,652; 5,652,113; 5,661,024; 5,766,879; 5,786,155; and 5,786,157, the disclosures of which are hereby incorporated by reference in their entireties into this application.)

25

Such transfected cells may also be used to test compounds and screen compound libraries to obtain compounds which bind to the SNORF25 receptor, as well as compounds which activate or inhibit activation of functional responses in such cells, and 30 therefore are likely to do so in vivo. (See, for example, U.S. Patent Nos. 5,053,337; 5,155,218; 5,360,735; 5,472,866; 5,476,782; 5,516,653; 5,545,549; 5,556,753; 5,595,880; 5,602,024; 5,639,652; 5,652,113; 5,661,024; 5,766,879; 5,786,155; and 5,786,157, the disclosures of which are hereby 35 incorporated by reference in their entireties into this application.)

This invention further provides an antibody capable of

binding to a mammalian receptor encoded by a nucleic acid encoding a mammalian receptor. In one embodiment, the mammalian receptor is a human receptor. In a further embodiment, the mammalian receptor is a rat receptor. This 5 invention also provides an agent capable of competitively inhibiting the binding of an antibody to a mammalian receptor. In one embodiment, the antibody is a monoclonal antibody or antisera.

10 Methods of preparing and employing antisense oligonucleotides, antibodies, nucleic acid probes and transgenic animals directed to the SNORF25 receptor are well known in the art. (See, for example, U.S. Patent Nos. 5,053,337; 5,155,218; 5,360,735; 5,472,866; 5,476,782; 15 5,516,653; 5,545,549; 5,556,753; 5,595,880; 5,602,024; 5,639,652; 5,652,113; 5,661,024; 5,766,879; 5,786,155; and 5,786,157, the disclosures of which are hereby incorporated by reference in their entireties into this application.)

20 This invention also provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to RNA encoding a mammalian SNORF25 receptor, so as to prevent translation of such RNA. This invention further provides an antisense oligonucleotide having a sequence capable of 25 specifically hybridizing to genomic DNA encoding a mammalian SNORF25 receptor, so as to prevent transcription of such genomic DNA. In one embodiment, the oligonucleotide comprises chemically modified nucleotides or nucleotide analogues.

30 This invention provides an antibody capable of binding to a mammalian SNORF25 receptor encoded by a nucleic acid in accordance with this invention. In one embodiment, the mammalian SNORF25 receptor is a human SNORF25 receptor. In 35 a further embodiment, the mammalian SNORF25 receptor is a rat SNORF25 receptor.

Moreover, this invention provides an agent capable of

competitively inhibiting the binding of an antibody in accordance with this invention to a mammalian SNORF25 receptor. In one embodiment, the antibody is a monoclonal antibody or antisera.

5

This invention still further provides a pharmaceutical composition comprising (a) an amount of an oligonucleotide in accordance with this invention capable of passing through a cell membrane and effective to reduce expression of a 10 mammalian SNORF25 receptor and (b) a pharmaceutically acceptable carrier capable of passing through the cell membrane.

In one embodiment, the oligonucleotide is coupled to a 15 substance which inactivates mRNA. In another embodiment, the substance which inactivates mRNA is a ribozyme. In another embodiment, the pharmaceutically acceptable carrier comprises a structure which binds to a mammalian SNORF25 receptor on a cell capable of being taken up by the cells after binding 20 to the structure. In another embodiment, the pharmaceutically acceptable carrier is capable of binding to a mammalian SNORF25 receptor which is specific for a selected cell type.

25 This invention also provides a pharmaceutical composition which comprises an amount of an antibody in accordance with this invention effective to block binding of a ligand to a human SNORF25 receptor and a pharmaceutically acceptable carrier.

30

This invention further provides a transgenic, nonhuman mammal expressing DNA encoding a mammalian SNORF25 receptor in accordance with this invention. This invention provides a transgenic, nonhuman mammal comprising a homologous 35 recombination knockout of a native mammalian SNORF25 receptor. This invention further provides a transgenic, nonhuman mammal whose genome comprises antisense DNA complementary to DNA encoding a mammalian SNORF25 receptor

in accordance with this invention so placed within such genome as to be transcribed into antisense mRNA which is complementary to and hybridizes with mRNA encoding the mammalian SNORF25 receptor so as to reduce translation of 5 such mRNA and expression of such receptor. In one embodiment, the DNA encoding the mammalian SNORF25 receptor additionally comprises an inducible promoter. In another embodiment, the DNA encoding the mammalian SNORF25 receptor additionally comprises tissue specific regulatory elements. 10 In another embodiment, the transgenic, nonhuman mammal is a mouse.

This invention provides a process for identifying a chemical compound which specifically binds to a mammalian SNORF25 15 receptor which comprises contacting cells containing DNA encoding, and expressing on their cell surface, the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, with the compound under conditions suitable for binding, and detecting specific 20 binding of the chemical compound to the mammalian SNORF25 receptor. This invention further provides a process for identifying a chemical compound which specifically binds to a mammalian SNORF25 receptor which comprises contacting a membrane preparation from cells containing DNA encoding, and 25 expressing on their cell surface, the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, with the compound under conditions suitable for binding, and detecting specific binding of the chemical compound to the mammalian SNORF25 30 receptor.

In one embodiment, the mammalian SNORF25 receptor is a human SNORF25 receptor. In another embodiment, the mammalian SNORF25 receptor has substantially the same amino acid 35 sequence as the human SNORF25 receptor encoded by plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495). In another embodiment, the mammalian SNORF25 receptor has substantially the same amino acid sequence as that shown in Figures 2A-2B

(SEQ ID NO: 2). In another embodiment, the mammalian SNORF25 receptor has the amino acid sequence shown in Figures 2A-2B (SEQ ID NO: 2).

5 In another embodiment, the mammalian SNORF25 receptor is a rat SNORF25 receptor. In another embodiment, the mammalian SNORF25 receptor has substantially the same amino acid sequence as the rat SNORF25 receptor encoded by plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494). In another 10 embodiment, the mammalian SNORF25 receptor has substantially the same amino acid sequence as that shown in Figures 4A-4B (SEQ ID NO: 4). In another embodiment, the mammalian SNORF25 receptor has the amino acid sequence shown in Figures 4A-4B (SEQ ID NO: 4).

15

In one embodiment, the compound is not previously known to bind to a mammalian SNORF25 receptor. In one embodiment, the cell is an insect cell. In one embodiment, the cell is a mammalian cell. In another embodiment, the cell is 20 nonneuronal in origin. In another embodiment, the nonneuronal cell is a COS-7 cell, 293 human embryonic kidney cell, a CHO cell, a NIH-3T3 cell, a mouse Y1 cell, or a LM(tk-) cell. In another embodiment, the compound is a compound not previously known to bind to a mammalian SNORF25 25 receptor. This invention provides a compound identified by the preceding process of this invention.

This invention still further provides a process involving competitive binding for identifying a chemical compound which 30 specifically binds to a mammalian SNORF25 receptor which comprises separately contacting cells expressing on their cell surface the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, with both the chemical compound and a second chemical 35 compound known to bind to the receptor, and with only the second chemical compound, under conditions suitable for binding of such compounds to the receptor, and detecting specific binding of the chemical compound to the mammalian

SNORF25 receptor, a decrease in the binding of the second chemical compound to the mammalian SNORF25 receptor in the presence of the chemical compound being tested indicating that such chemical compound binds to the mammalian SNORF25 receptor.

This invention provides a process involving competitive binding for identifying a chemical compound which specifically binds to a mammalian SNORF25 receptor which 10 comprises separately contacting a membrane preparation from cells expressing on their cell surface the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, with both the chemical compound and a second chemical compound known to bind to the receptor, 15 and with only the second chemical compound, under conditions suitable for binding of such compounds to the receptor, and detecting specific binding of the chemical compound to the mammalian SNORF25 receptor, a decrease in the binding of the second chemical compound to the mammalian SNORF25 receptor 20 in the presence of the chemical compound being tested indicating that such chemical compound binds to the mammalian SNORF25 receptor.

In one embodiment, the mammalian SNORF25 receptor is a human 25 SNORF25 receptor. In another embodiment, the mammalian SNORF25 receptor is a rat SNORF25 receptor. In a further embodiment, the cell is an insect cell. In another embodiment, the cell is a mammalian cell. In another embodiment, the cell is nonneuronal in origin. In another 30 embodiment, the nonneuronal cell is a COS-7 cell, 293 human embryonic kidney cell, a CHO cell, a NIH-3T3 cell, a mouse Y1 cell, or a LM(tk-) cell. In another embodiment, the compound is not previously known to bind to a mammalian SNORF25 receptor. This invention provides a compound 35 identified by the preceding process of this invention.

This invention provides a method of screening a plurality of chemical compounds not known to bind to a mammalian SNORF25

receptor to identify a compound which specifically binds to the mammalian SNORF25 receptor, which comprises (a) contacting cells transfected with, and expressing, DNA encoding the mammalian SNORF25 receptor with a compound known 5 to bind specifically to the mammalian SNORF25 receptor; (b) contacting the cells of step (a) with the plurality of compounds not known to bind specifically to the mammalian SNORF25 receptor, under conditions permitting binding of compounds known to bind to the mammalian SNORF25 receptor; 10 (c) determining whether the binding of the compound known to bind to the mammalian SNORF25 receptor is reduced in the presence of the plurality of compounds, relative to the binding of the compound in the absence of the plurality of compounds; and if so (d) separately determining the binding 15 to the mammalian SNORF25 receptor of each compound included in the plurality of compounds, so as to thereby identify any compound included therein which specifically binds to the mammalian SNORF25 receptor.

20 This invention provides a method of screening a plurality of chemical compounds not known to bind to a mammalian SNORF25 receptor to identify a compound which specifically binds to the mammalian SNORF25 receptor, which comprises (a) contacting a membrane preparation from cells transfected 25 with, and expressing, DNA encoding the mammalian SNORF25 receptor with the plurality of compounds not known to bind specifically to the mammalian SNORF25 receptor under conditions permitting binding of compounds known to bind to the mammalian SNORF25 receptor; (b) determining whether the 30 binding of a compound known to bind to the mammalian SNORF25 receptor is reduced in the presence of the plurality of compounds, relative to the binding of the compound in the absence of the plurality of compounds; and if so (c) separately determining the binding to the mammalian SNORF25 35 receptor of each compound included in the plurality of compounds, so as to thereby identify any compound included therein which specifically binds to the mammalian SNORF25 receptor.

In one embodiment, the mammalian SNORF25 receptor is a human SNORF25 receptor. In a further embodiment, the mammalian SNORF25 receptor is a rat SNORF25 receptor. In another embodiment, the cell is a mammalian cell. In another 5 embodiment, the mammalian cell is non-neuronal in origin. In a further embodiment, the non-neuronal cell is a COS-7 cell, a 293 human embryonic kidney cell, a LM(tk-) cell, a CHO cell, a mouse Y1 cell, or an NIH-3T3 cell.

10 This invention provides a method of detecting expression of a mammalian SNORF25 receptor by detecting the presence of mRNA coding for the mammalian SNORF25 receptor which comprises obtaining total mRNA from the cell and contacting the mRNA so obtained with a nucleic acid probe according to 15 this invention under hybridizing conditions, detecting the presence of mRNA hybridized to the probe, and thereby detecting the expression of the mammalian SNORF25 receptor by the cell.

20 This invention provides a method of detecting the presence of a mammalian SNORF25 receptor on the surface of a cell which comprises contacting the cell with an antibody according to this invention under conditions permitting binding of the antibody to the receptor, detecting the 25 presence of the antibody bound to the cell, and thereby detecting the presence of the mammalian SNORF25 receptor on the surface of the cell.

This invention provides a method of determining the 30 physiological effects of varying levels of activity of mammalian SNORF25 receptors which comprises producing a transgenic, nonhuman mammal in accordance with this invention whose levels of mammalian SNORF25 receptor activity are varied by use of an inducible promoter which regulates 35 mammalian SNORF25 receptor expression.

This invention provides a method of determining the physiological effects of varying levels of activity of

mammalian SNORF25 receptors which comprises producing a panel of transgenic, nonhuman mammals in accordance with this invention each expressing a different amount of mammalian SNORF25 receptor.

5

This invention provides a method for identifying an antagonist capable of alleviating an abnormality wherein the abnormality is alleviated by decreasing the activity of a mammalian SNORF25 receptor comprising administering a compound to a transgenic, nonhuman mammal according to this invention, and determining whether the compound alleviates any physiological and/or behavioral abnormality displayed by the transgenic, nonhuman mammal as a result of overactivity of a mammalian SNORF25 receptor, the alleviation of such abnormality identifying the compound as an antagonist. In one embodiment, the mammalian SNORF25 receptor is a human SNORF25 receptor. In a further embodiment, the mammalian SNORF25 receptor is a rat SNORF25 receptor. The invention provides an antagonist identified by the preceding method according to this invention. This invention provides a composition, e.g. a pharmaceutical composition, comprising an antagonist according to this invention and a carrier, e.g. a pharmaceutically acceptable carrier. This invention provides a method of treating an abnormality in a subject wherein the abnormality is alleviated by decreasing the activity of a mammalian SNORF25 receptor which comprises administering to the subject an effective amount of the pharmaceutical composition according to this invention so as to thereby treat the abnormality.

30

This invention provides a method for identifying an agonist capable of alleviating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian SNORF25 receptor comprising administering a compound to a transgenic, nonhuman mammal according to this invention, and determining whether the compound alleviates any physiological and/or behavioral abnormality displayed by the transgenic, nonhuman mammal, the alleviation of such an

abnormality identifying the compound as an agonist. In one embodiment, the mammalian SNORF25 receptor is a human SNORF25 receptor. In a further embodiment, the mammalian SNORF25 receptor is a rat SNORF25 receptor. This invention provides 5 an agonist identified by the preceding method according to this invention. This invention provides a composition, e.g. a pharmaceutical composition, comprising an agonist identified by the method according to this invention and a carrier, e.g. a pharmaceutically acceptable carrier.

10

This invention provides a method of treating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian SNORF25 receptor which comprises administering to the subject an effective 15 amount of the pharmaceutical composition according to this invention so as to thereby treat the abnormality.

This invention provides a method for diagnosing a predisposition to a disorder associated with the activity of 20 a specific mammalian allele which comprises: (a) obtaining DNA of subjects suffering from the disorder; (b) performing a restriction digest of the DNA with a panel of restriction enzymes; (c) electrophoretically separating the resulting DNA fragments on a sizing gel; (d) contacting the resulting gel 25 with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a mammalian SNORF25 receptor and labeled with a detectable marker; (e) detecting labeled bands which have hybridized to the DNA encoding a mammalian 30 SNORF25 receptor to create a unique band pattern specific to the DNA of subjects suffering from the disorder; (f) repeating steps (a)-(e) with DNA obtained for diagnosis from subjects not yet suffering from the disorder; and (g) comparing the unique band pattern specific to the DNA of 35 subjects suffering from the disorder from step (e) with the band pattern from step (f) for subjects not yet suffering from the disorder so as to determine whether the patterns are the same or different and thereby diagnose predisposition to

the disorder if the patterns are the same.

In one embodiment, the disorder is a disorder associated with the activity of a specific mammalian allele is diagnosed.

5

This invention provides a method of preparing a purified mammalian SNORF25 receptor according to this invention which comprises: (a) culturing cells which express the mammalian SNORF25 receptor; (b) recovering the mammalian SNORF25 receptor from the cells; and (c) purifying the mammalian SNORF25 receptor so recovered.

This invention provides a method of preparing the purified mammalian SNORF25 receptor according to this invention which comprises: (a) inserting a nucleic acid encoding the mammalian SNORF25 receptor into a suitable expression vector; (b) introducing the resulting vector into a suitable host cell; (c) placing the resulting host cell in suitable conditions permitting the production of the mammalian SNORF25 receptor; (d) recovering the mammalian SNORF25 receptor so produced; and optionally (e) isolating and/or purifying the mammalian SNORF25 receptor so recovered.

This invention provides a process for determining whether a chemical compound is a mammalian SNORF25 receptor agonist which comprises contacting cells transfected with and expressing DNA encoding the mammalian SNORF25 receptor with the compound under conditions permitting the activation of the mammalian SNORF25 receptor, and detecting any increase in mammalian SNORF25 receptor activity, so as to thereby determine whether the compound is a mammalian SNORF25 receptor agonist.

This invention provides a process for determining whether a chemical compound is a mammalian SNORF25 receptor antagonist which comprises contacting cells transfected with and expressing DNA encoding the mammalian SNORF25 receptor with the compound in the presence of a known mammalian SNORF25

receptor agonist, under conditions permitting the activation of the mammalian SNORF25 receptor, and detecting any decrease in mammalian SNORF25 receptor activity, so as to thereby determine whether the compound is a mammalian SNORF25 receptor antagonist.

In one embodiment, the mammalian SNORF25 receptor is a human SNORF25 receptor. In another embodiment, the mammalian SNORF25 receptor is a rat SNORF25 receptor.

10

This invention provides a composition, for example a pharmaceutical composition, which comprises an amount of a mammalian SNORF25 receptor agonist determined by a process according to this invention effective to increase activity of a mammalian SNORF25 receptor and a carrier, for example, a pharmaceutically acceptable carrier. In one embodiment, the mammalian SNORF25 receptor agonist is not previously known.

20 This invention provides a composition, for example a pharmaceutical composition, which comprises an amount of a mammalian SNORF25 receptor antagonist determined by a process according to this invention effective to reduce activity of a mammalian SNORF25 receptor and a carrier, for example, a pharmaceutically acceptable carrier. In one embodiment, the mammalian SNORF25 receptor antagonist is not previously known.

30 This invention provides a process for determining whether a chemical compound specifically binds to and activates a mammalian SNORF25 receptor, which comprises contacting cells producing a second messenger response and expressing on their cell surface the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, 35 with the chemical compound under conditions suitable for activation of the mammalian SNORF25 receptor, and measuring the second messenger response in the presence and in the absence of the chemical compound, a change, e.g. an increase,

in the second messenger response in the presence of the chemical compound indicating that the compound activates the mammalian SNORF25 receptor.

- 5 In one embodiment, the second messenger response comprises chloride channel activation and the change in second messenger is an increase in the level of chloride current.
- 10 In another embodiment, the second messenger response comprises change in intracellular calcium levels and the change in second messenger is an increase in the measure of intracellular calcium. In another embodiment, the second messenger response comprises release of inositol phosphate and the change in second messenger is an increase in the level of inositol phosphate. In another embodiment, the
- 15 second messenger response comprises release of arachidonic acid and the change in second messenger is an increase in the level of arachidonic acid. In yet another embodiment, the second messenger response comprises GTP γ S ligand binding and the change in second messenger is an increase in GTP γ S ligand
- 20 binding. In another embodiment, the second messenger response comprises activation of MAP kinase and the change in second messenger response is an increase in MAP kinase activation. In a further embodiment, the second messenger response comprises cAMP accumulation and the change in second
- 25 messenger response is an increase in cAMP accumulation.

This invention provides a process for determining whether a chemical compound specifically binds to and inhibits activation of a mammalian SNORF25 receptor, which comprises

- 30 separately contacting cells producing a second messenger response and expressing on their cell surface the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, with both the chemical compound and a second chemical compound known to activate the
- 35 mammalian SNORF25 receptor, and with only the second chemical compound, under conditions suitable for activation of the mammalian SNORF25 receptor, and measuring the second messenger response in the presence of only the second

chemical compound and in the presence of both the second chemical compound and the chemical compound, a smaller change, e.g. increase, in the second messenger response in the presence of both the chemical compound and the second 5 chemical compound than in the presence of only the second chemical compound indicating that the chemical compound inhibits activation of the mammalian SNORF25 receptor.

In one embodiment, the second messenger response comprises 10 chloride channel activation and the change in second messenger response is a smaller increase in the level of chloride current in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound. In another 15 embodiment, the second messenger response comprises change in intracellular calcium levels and the change in second messenger response is a smaller increase in the measure of intracellular calcium in the presence of both the chemical compound and the second chemical compound than in the 20 presence of only the second chemical compound. In another embodiment, the second messenger response comprises release of inositol phosphate and the change in second messenger response is a smaller increase in the level of inositol phosphate in the presence of both the chemical compound and 25 the second chemical compound than in the presence of only the second chemical compound.

In one embodiment, the second messenger response comprises activation of MAP kinase and the change in second messenger 30 response is a smaller increase in the level of MAP kinase activation in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound. In another embodiment, the second messenger response comprises change in cAMP levels and the 35 change in second messenger response is a smaller change in the level of cAMP in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound. In another

embodiment, the second messenger response comprises release of arachidonic acid and the change in second messenger response is an increase in the level of arachidonic acid levels in the presence of both the chemical compound and the 5 second chemical compound than in the presence of only the second chemical compound. In a further embodiment, the second messenger response comprises GTP γ S ligand binding and the change in second messenger is a smaller increase in GTP γ S ligand binding in the presence of both the chemical compound 10 and the second chemical compound than in the presence of only the second chemical compound.

In one embodiment, the mammalian SNORF25 receptor is a human SNORF25 receptor. In a further embodiment, the mammalian 15 SNORF25 receptor is a rat SNORF25 receptor. In another embodiment, the cell is an insect cell. In another embodiment, the cell is a mammalian cell. In another embodiment, the mammalian cell is nonneuronal in origin. In another embodiment, the nonneuronal cell is a COS-7 cell, CHO 20 cell, 293 human embryonic kidney cell, NIH-3T3 cell or LM(tk-) cell. In another embodiment, the compound is not previously known to bind to a mammalian SNORF25 receptor.

This invention provides a compound determined by a process 25 according to this invention and a composition, for example, a pharmaceutical composition, which comprises an amount of a mammalian SNORF25 receptor agonist determined to be such by a process according to this invention effective to increase activity of the mammalian SNORF25 receptor and a 30 carrier, for example, a pharmaceutically acceptable carrier. In one embodiment, the mammalian SNORF25 receptor agonist is not previously known.

This invention provides a composition, for example, a 35 pharmaceutical composition, which comprises an amount of a mammalian SNORF25 receptor antagonist determined to be such by a process according to this invention, effective to reduce activity of the mammalian SNORF25 receptor and a carrier, for

example, a pharmaceutically acceptable carrier. In one embodiment, the mammalian SNORF25 receptor antagonist is not previously known.

- 5 This invention provides a method of screening a plurality of chemical compounds not known to activate a mammalian SNORF25 receptor to identify a compound which activates the mammalian SNORF25 receptor which comprises: (a) contacting cells transfected with and expressing the mammalian SNORF25
- 10 receptor with the plurality of compounds not known to activate the mammalian SNORF25 receptor, under conditions permitting activation of the mammalian SNORF25 receptor; (b) determining whether the activity of the mammalian SNORF25 receptor is increased in the presence of one or more of the
- 15 compounds; and if so (c) separately determining whether the activation of the mammalian SNORF25 receptor is increased by any compound included in the plurality of compounds, so as to thereby identify each compound which activates the mammalian SNORF25 receptor. In one embodiment, the mammalian
- 20 SNORF25 receptor is a human SNORF25 receptor. In a further embodiment, the mammalian SNORF25 receptor is a rat SNORF25 receptor.

This invention provides a method of screening a plurality of chemical compounds not known to inhibit the activation of a mammalian SNORF25 receptor to identify a compound which inhibits the activation of the mammalian SNORF25 receptor, which comprises: (a) contacting cells transfected with and expressing the mammalian SNORF25 receptor with the plurality of compounds in the presence of a known mammalian SNORF25 receptor agonist, under conditions permitting activation of the mammalian SNORF25 receptor; (b) determining whether the extent or amount of activation of the mammalian SNORF25 receptor is reduced in the presence of one or more of the

30 compounds, relative to the extent or amount of activation of the mammalian SNORF25 receptor in the absence of such one or more compounds; and if so (c) separately determining whether each such compound inhibits activation of the mammalian

35

SNORF25 receptor for each compound included in the plurality of compounds, so as to thereby identify any compound included in such plurality of compounds which inhibits the activation of the mammalian SNORF25 receptor.

5

In one embodiment, the mammalian SNORF25 receptor is a human SNORF25 receptor. In a further embodiment, the mammalian SNORF25 receptor is a rat SNORF25 receptor. In another embodiment, wherein the cell is a mammalian cell. In another 10 embodiment, the mammalian cell is non-neuronal in origin. In another embodiment, the non-neuronal cell is a COS-7 cell, a 293 human embryonic kidney cell, a LM(tk-) cell or an NIH-3T3 cell.

15 This invention provides a composition, for example a pharmaceutical composition, comprising a compound identified by a method according to this invention in an amount effective to increase mammalian SNORF25 receptor activity and a carrier, for example, a pharmaceutically acceptable 20 carrier.

This invention provides a composition, for example, a pharmaceutical composition, comprising a compound identified by a method according to this invention in an amount 25 effective to decrease mammalian SNORF25 receptor activity and a carrier, for example, a pharmaceutically acceptable carrier.

This invention provides a method of treating an abnormality 30 in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian SNORF25 receptor which comprises administering to the subject a compound which is a mammalian SNORF25 receptor agonist in an amount effective to treat the abnormality. In one embodiment, the abnormality 35 is a regulation of a steroid hormone disorder, an epinephrine release disorder, a gastrointestinal disorder, a cardiovascular disorder, an electrolyte balance disorder, hypertension, diabetes, a respiratory disorder, asthma, a

reproductive function disorder, an immune disorder, an endocrine disorder, a musculoskeletal disorder, a neuroendocrine disorder, a cognitive disorder, a memory disorder, somatosensory and neurotransmission disorders, a 5 motor coordination disorder, a sensory integration disorder, a motor integration disorder, a dopaminergic function disorder, an appetite disorder, such as anorexia or obesity, a sensory transmission disorder, an olfaction disorder, an autonomic nervous system disorder, pain, psychotic behavior, 10 affective disorder, migraine, cancer, proliferative diseases, wound healing, tissue regeneration, blood coagulation-related disorders, developmental disorders, or ischemia-reperfusion injury-related diseases.

15 This invention provides a method of treating an abnormality in a subject wherein the abnormality is alleviated by decreasing the activity of a mammalian SNORF25 receptor which comprises administering to the subject a compound which is a mammalian SNORF25 receptor antagonist in an amount 20 effective to treat the abnormality. In one embodiment, the abnormality is a regulation of a steroid hormone disorder, an epinephrine release disorder, a gastrointestinal disorder, a cardiovascular disorder, an electrolyte balance disorder, hypertension, diabetes, a respiratory disorder, asthma, a 25 reproductive function disorder, an immune disorder, an endocrine disorder, a musculoskeletal disorder, a neuroendocrine disorder, a cognitive disorder, a memory disorder, somatosensory and neurotransmission disorders, a motor coordination disorder, a sensory integration disorder, 30 a motor integration disorder, a dopaminergic function disorder, an appetite disorder, such as anorexia or obesity, a somatosensory neurotransmission disorder, an olfaction disorder, an autonomic nervous system disorder, pain, psychotic behavior, affective disorder, migraine, cancer, 35 proliferative diseases, wound healing, tissue regeneration, blood coagulation-related disorders, developmental disorders, or ischemia-reperfusion injury-related diseases.

This invention provides a process for making a composition of matter which specifically binds to a mammalian SNORF25 receptor which comprises identifying a chemical compound using a process in accordance with this invention and then 5 synthesizing the chemical compound or a novel structural and functional analog or homolog thereof. In one embodiment, the mammalian SNORF25 receptor is a human SNORF25 receptor. In another embodiment, the mammalian SNORF25 receptor is a rat SNORF25 receptor.

10

This invention provides a process for preparing a composition, for example, a pharmaceutical composition which comprises admixing a carrier, for example, a pharmaceutically acceptable carrier, and a pharmaceutically effective amount 15 of a chemical compound identified by a process in accordance with this invention or a novel structural and functional analog or homolog thereof. In one embodiment, the mammalian SNORF25 receptor is a human SNORF25 receptor. In another embodiment, the mammalian SNORF25 receptor is a rat SNORF25 20 receptor.

Thus, once the gene for a targeted receptor subtype is cloned, it is placed into a recipient cell which then expresses the targeted receptor subtype on its surface. This 25 cell, which expresses a single population of the targeted human receptor subtype, is then propagated resulting in the establishment of a cell line. This cell line, which constitutes a drug discovery system, is used in two different types of assays: binding assays and functional assays. In 30 binding assays, the affinity of a compound for both the receptor subtype that is the target of a particular drug discovery program and other receptor subtypes that could be associated with side effects are measured. These measurements enable one to predict the potency of a compound, 35 as well as the degree of selectivity that the compound has for the targeted receptor subtype over other receptor subtypes. The data obtained from binding assays also enable chemists to design compounds toward or away from one or more

of the relevant subtypes, as appropriate, for optimal therapeutic efficacy. In functional assays, the nature of the response of the receptor subtype to the compound is determined. Data from the functional assays show whether the 5 compound is acting to inhibit or enhance the activity of the receptor subtype, thus enabling pharmacologists to evaluate compounds rapidly at their ultimate human receptor subtypes targets permitting chemists to rationally design drugs that will be more effective and have fewer or substantially less 10 severe side effects than existing drugs.

Approaches to designing and synthesizing receptor subtype-selective compounds are well known and include traditional medicinal chemistry and the newer technology of 15 combinatorial chemistry, both of which are supported by computer-assisted molecular modeling. With such approaches, chemists and pharmacologists use their knowledge of the structures of the targeted receptor subtype and compounds determined to bind and/or activate or inhibit activation of 20 the receptor subtype to design and synthesize structures that will have activity at these receptor subtypes.

Combinatorial chemistry involves automated synthesis of a variety of novel compounds by assembling them using different 25 combinations of chemical building blocks. The use of combinatorial chemistry greatly accelerates the process of generating compounds. The resulting arrays of compounds are called libraries and are used to screen for compounds ("lead compounds") that demonstrate a sufficient level of activity 30 at receptors of interest. Using combinatorial chemistry it is possible to synthesize "focused" libraries of compounds anticipated to be highly biased toward the receptor target of interest.

35 Once lead compounds are identified, whether through the use of combinatorial chemistry or traditional medicinal chemistry or otherwise, a variety of homologs and analogs are prepared to facilitate an understanding of the relationship between

chemical structure and biological or functional activity. These studies define structure activity relationships which are then used to design drugs with improved potency, selectivity and pharmacokinetic properties. Combinatorial 5 chemistry is also used to rapidly generate a variety of structures for lead optimization. Traditional medicinal chemistry, which involves the synthesis of compounds one at a time, is also used for further refinement and to generate compounds not accessible by automated techniques. Once such 10 drugs are defined the production is scaled up using standard chemical manufacturing methodologies utilized throughout the pharmaceutical and chemistry industry.

This invention will be better understood from the 15 Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the invention as described more fully in the claims which follow thereafter.

EXPERIMENTAL DETAILSMaterials and Methods5 Mixed Oligonucleotide Primed Amplification of cDNA (MOPAC)

Mixed Oligonucleotide Primed Amplification of cDNA (MOPAC) was performed on several DNA templates including: rat genomic DNA, cDNA reverse-transcribed from mRNA isolated from the GH1 10 cell line, and the Rin14b cell line. The MOPAC reaction was performed using Taq DNA polymerase (Boehringer-Mannheim, Indianapolis, IN) and the following degenerate oligonucleotides: JAB55, designed based on the third transmembrane domain of the galanin, somatostatin, and opiate 15 receptor families; and TL1020, designed based on the 7th transmembrane domain of the galanin receptor family.

The conditions for the MOPAC PCR reaction were as follows: 3 minute hold at 94° C; 10 cycles of 1 minute at 94° C, 1 20 minute 45 seconds at 44° C, 2 minutes at 72° C; 30 cycles of 94° C for 1 minute, 49° C for 1 minute 45 seconds, 2 minutes at 72° C; 4 minute hold at 72° C; 4° C hold until ready for agarose gel electrophoresis.

25 The products were run on a 1% agarose TAE gel and bands of the expected size (~500-600 bp) were cut from the gel, purified using the QIAQUICK gel extraction kit (QIAGEN, Chatsworth, CA), and subcloned into the TA cloning vector (Invitrogen, San Diego, CA). White (insert-containing) 30 colonies were picked and subjected to PCR using pCR2.1 vector primers JAB1 and JAB2 using the following protocol: 94° C hold for 3 minutes; 35 cycles of 94° C for 1 minute, 68° C for 1 minute 15 seconds; 2 minute hold at 68° C, 4° C hold until the products were ready for purification. PCR products 35 were purified by isopropanol precipitation (10 μl PCR product, 18 μl low TE, 10.5 μl 2M NaClO₄, and 21.5 μl isopropanol) and sequenced using the ABI Big Dye cycle

sequencing protocol and ABI 377 sequencers (ABI, Foster City, CA). One of these PCR products, later named SNORF25, was determined to be a novel G protein-coupled receptor-like sequence based on database searches and its homology to other 5 known G protein-coupled receptors (~29% identity to the known receptors dopamine D1, beta-adrenergic 2b and 5-HT1f; 34% identity to the 5-HT4l receptor).

5' and 3' RACE

10

To determine the full-length coding sequence of SNORF25, the Clontech Marathon cDNA Amplification kit (Clontech, Palo Alto, CA) for 5'/3' Rapid Amplification of cDNA ends (RACE) was utilized. Total RNA from Rin14b cells was PolyA⁺ selected using a FastTrack mRNA Isolation Kit (Invitrogen). For 5'RACE, double-stranded cDNA was synthesized from 1 μ g polyA⁺ RNA using primer JAB73, a reverse primer from the putative fifth transmembrane domain of the PCR fragment described above (SNORF25). Adaptor ligation and nested PCR 15 were performed according to the Marathon cDNA Amplification protocol using Advantage KlenTaq Polymerase (Clontech, Palo Alto, CA). The initial PCR was performed on a 50-fold dilution of the ligated cDNA using the supplier's Adaptor Primer 1 and JAB71, a reverse primer from the 5'end of the 20 putative fifth transmembrane domain of the PCR fragment described above. One μ l of this initial PCR reaction was re-amplified using the Adaptor Primer 2 and JAB69, a reverse primer just downstream of the fourth transmembrane domain. The 25 conditions for PCR were 1 minute at 94° C; 5 cycles of 94° C for 15 seconds and 72° C for 1 minute 30 seconds; 5 cycles of 94° C for 15 seconds and 70° C for 1 minute 30 seconds; 22 cycles of 94° C for 15 seconds and 68° C for 1 minute 30 seconds; 68° C hold for 5 minutes, and 4° C hold until the 30 products were ready for analysis. A 600 bp fragment from the nested PCR was isolated from a 1% agarose TAE gel using the QIAQUICK kit and sequenced using ABI 377 sequencers and 35 BigDye termination cycle sequencing as described above. Sequences were analyzed using the Wisconsin Package (GCG,

Genetics Computer Group, Madison, WI).

For 3' RACE, double stranded cDNA was synthesized from 1 μ g polyA⁺ RNA using the cDNA synthesis primer CDS supplied with 5 the Marathon cDNA Amplification Kit (Clontech). PCR conditions for the 3' RACE reactions were similar to the 5' RACE reactions, except that JAB74 and JAB72, forward primers from the sequence located between the fifth and sixth transmembrane domains of the novel PCR fragment from MOPAC 10 described above, were used in place of JAB 71 and JAB73, respectively. A 1.4 kb fragment from the nested PCR was isolated from a 1% agarose TAE gel using the QIAQUICK gel purification kit (QIAGEN) and sequenced as above.

15 After determining the full-length coding sequence of this receptor sequence, the entire coding region was amplified from Rin14b cell line cDNA and rat genomic DNA using the Expand Long PCR system (Boehringer-Mannheim). The primers for this reaction were specific to the 5' and 3' untranslated 20 regions of SNORF25 with *Bam*HI and *Hind*III restriction sites incorporated into the 5' ends of the 5' (JAB86) and 3' (JAB84) primers, respectively. The products from this reaction were then digested with *Bam*HI and *Hind*III, subcloned into the *Bam*HI/*Hind*III site of the expression vector pcDNA3.1 25 (-), and sequenced in both directions using vector- and gene-specific primers. Double-stranded sequence from the Rin14b-cloned SNORF25 product agreed with the sequence of the same gene amplified from rat genomic DNA. This receptor/expression vector construct of rat SNORF25 in 30 pcDNA3.1(-) was named pcDNA3.1-rSNORF25.

Homology cloning of the human homolog of SNORF25

To clone the human homolog of SNORF25, two oligonucleotide 35 probes were designed based on the second (BB426) and fifth (BB427) transmembrane domains (TMs) of the rat SNORF25 sequence, and used to probe a human genomic cosmid library

(Clontech). Both primers were end-labeled with $\alpha^{32}\text{P}$ -dATP and terminal transferase (Promega, Madison, WI). Hybridization was performed under medium stringency conditions: 40° C in a solution containing 37.5% formamide 5x SSC (1X SSC is 0.15M sodium chloride, 0.015M sodium citrate), 1x Denhardt's solution (0.02% polyvinylpyrrolindone, 0.02% Ficoll, 0.02% bovine serum albumin), 7 mM Tris, and 25 $\mu\text{g}/\text{ml}$ sonicated salmon sperm DNA. The filters were washed three times for 20 minutes at room temperature in a buffer containing 2x SSC/0.1% sodium dodecyl sulfate; two times for 20 minutes in a buffer containing 0.1x SSC/0.1% sodium dodecyl sulfate, and exposed at -70°C to Kodak BioMax MS film in the presence of an intensifying screen.

15 Cosmid clones hybridizing with the probes were picked, streaked on plates, and screened a second time with the same probes to verify and isolate the individual positive colonies under the same conditions. Cosmid DNA from positive colonies was digested with *Bam*HI and *Hind*III, run on an agarose gel,

20 transferred to nitrocellulose, and probed with ^{32}P -labelled BB426. A fragment of approximately 1.9kb from clone #45a (COS4 library) that hybridized to the probe was subcloned into the *Bam*HI/*Hind*III site of pEXJT3T7, an Okayama and Berg expression vector modified from pcEXV (Miller and Germain, 1986) to contain *Bst*XI and other additional restriction sites as well as T3 and T7 promoters (Stratagene), and sequenced on both strands as described above. The construct of the human SNORF25 receptor in this vector is named pEXJT3T7-hSNORF25. Human SNORF25 was analyzed using the GCG 25 software and was determined to contain the full-length sequence of human SNORF25, having 80% amino acid identity and 83% nucleotide identity to the rat receptor.

30

Oligonucleotide primers

35 The following is a list of primers and their associated sequences which were used in the cloning of these receptors:
JAB55: 5'-TBD SYVYIGAYM GITAY VTKG-3' (SEQ ID NO: 5)
TL1020: 5'-GAIRSIARIGMRTAIAYIAKIGGRTT-3' (SEQ ID NO: 6)

JAB1: 5'-TTATGCTTCGGCTCGTATGTTGTG-3' (SEQ ID NO: 7)
JAB2: 5'-ATGTGCTGCAAGGCGATTTAAGTTGGG-3' (SEQ ID NO: 8)
JAB69: 5'-TGGTCTGCTGGAATATGGAG-3' (SEQ ID NO: 9)
JAB71: 5'-CTTGGGTGAAACACAGCAAAGAAGG-3' (SEQ ID NO: 10)
5 JAB72: 5'-ATGGAACATGCAGGAGCCATGGTTGG-3' (SEQ ID NO: 11)
JAB73: 5'-AAGACAAAGAGGAGCACAGCTGGG-3' (SEQ ID NO: 12)
JAB74: 5'-GCTCAAGATTGCCTCTGTGCACAG-3' (SEQ ID NO: 13)
JAB84: 5'-ATCTATAAGCTTAGGCACTTGGAAACATCCATTCC-3' (SEQ ID NO:
14)
10 JAB86: 5'-ATCTATGGATCCTGTGAGAATCTGAGCTAAGACCC-3' (SEQ ID NO:
15)
BB426: 5'-TTCACCTTAAATCTGGCCGTGGCTGATAACCTTGAT-
TGGCGTGGCTATTCTGGGCTAG-3' (SEQ ID NO: 16)
BB427: 5'-GCTGTGTTCACCCAAGGTTGTGCTGACCCTCTC-
15 CTGTGCTGGCTTCTCCCAGCTGTGC-3' (SEQ ID NO: 17)

Isolation of other species homologs of SNORF25 receptor cDNA
A nucleic acid sequence encoding a SNORF25 receptor cDNA from
other species may be isolated using standard molecular
20 biology techniques and approaches such as those described
below:

Approach #1: A genomic library (e.g., cosmid, phage, P1, BAC,
YAC) generated from the species of interest may be screened
25 with a ³²P-labeled oligonucleotide probe corresponding to a
fragment of the human or rat SNORF25 receptors whose sequence
is shown in Figures 1A-1B and 3A-3B to isolate a genomic
clone. The full-length sequence may be obtained by
sequencing this genomic clone. If one or more introns are
30 present in the gene, the full-length intronless gene may be
obtained from cDNA using standard molecular biology
techniques. For example, a forward PCR primer designed in
the 5'UT and a reverse PCR primer designed in the 3'UT may
be used to amplify a full-length, intronless receptor from
35 cDNA. Standard molecular biology techniques could be used
to subclone this gene into a mammalian expression vector.

Approach #2: Standard molecular biology techniques may be

used to screen commercial cDNA phage libraries of the species of interest by hybridization under reduced stringency with a ^{32}P -labeled oligonucleotide probe corresponding to a fragment of the sequences shown in Figures 1A-1B or 3A-3B.

5 One may isolate a full-length SNORF25 receptor by obtaining a plaque purified clone from the lambda libraries and then subjecting the clone to direct DNA sequencing. Alternatively, standard molecular biology techniques could be used to screen cDNA plasmid libraries by PCR amplification

10 of library pools using primers designed against a partial species homolog sequence. A full-length clone may be isolated by Southern hybridization of colony lifts of positive pools with a ^{32}P -oligonucleotide probe.

15 Approach #3: 3' and 5' RACE may be utilized to generate PCR products from cDNA derived from the species of interest expressing SNORF25 which contain the additional sequence of SNORF25. These RACE PCR products may then be sequenced to determine the additional sequence. This new sequence is then

20 used to design a forward PCR primer in the 5'UT and a reverse primer in the 3'UT. These primers are then used to amplify a full-length SNORF25 clone from cDNA.

Examples of other species include, but are not limited to,
25 mouse, dog, monkey, hamster and guinea pig.

Host cells

A broad variety of host cells can be used to study heterologously expressed proteins. These cells include but
30 are not limited to mammalian cell lines such as; Cos-7, CHO, LM(tk^-), HEK293, etc.; insect cell lines such as; Sf9, Sf21, etc.; amphibian cells such as *Xenopus* oocytes; assorted yeast strains; assorted bacterial cell strains; and others. Culture conditions for each of these cell types is specific
35 and is known to those familiar with the art. The cells used to express SNORF25 receptor were Cos-7 and Chinese hamster ovary (CHO) cells.

COS-7 cells are grown on 150 mm plates in DMEM with supplements (Dulbecco's Modified Eagle Medium with 10% bovine calf serum, 4 mM glutamine, 100 units/ml penicillin/100 µg/ml streptomycin) at 37°C, 5% CO₂. Stock plates of COS-7 cells 5 are trypsinized and split 1:6 every 3-4 days.

CHO cells are grown on 150 mm plates in HAM's F-12 medium with supplements (10% bovine calf serum, 4 mM L-glutamine and 100 units/ml penicillin/ 100 µg/ml streptomycin) at 37°C, 5% 10 CO₂. Stock plates of CHO cells are trypsinized and split 1:8 every 3-4 days.

Transient expression

DNA encoding proteins to be studied can be transiently 15 expressed in a variety of mammalian, insect, amphibian, yeast, bacterial and other cell lines by several methods, such as, calcium phosphate-mediated, DEAE-dextran mediated, liposomal-mediated, viral-mediated, electroporation-mediated and microinjection delivery. Each of these methods may 20 require optimization of assorted experimental parameters depending on the DNA, cell line, and the type of assay to be subsequently employed. The electroporation method was used to transiently transfect various cell lines with SNORF25 cDNA.

25 A typical protocol for the electroporation method as applied to Cos-7 cells is described as follows. Cells to be used for transfection are split 24 hours prior to the transfection to provide flasks which are subconfluent at the time of 30 transfection. The cells are harvested by trypsinization resuspended in their growth media and counted. 5 x 10⁶ cells are suspended in 300 µl of DMEM and placed into an electroporation cuvette. 8 µg of receptor DNA plus 8 µg of any additional DNA needed (e.g. G protein expression vector, 35 reporter construct, antibiotic resistance marker, mock vector, etc.) is added to the cell suspension, the cuvette is placed into a BioRad Gene Pulser and subjected to an electrical pulse (Gene Pulser settings: 0.25 kV voltage, 950

μ F capacitance). Following the pulse, 800 μ l of complete DMEM is added to each cuvette and the suspension transferred to a sterile tube. Complete medium is added to each tube to bring the final cell concentration to 1 x 10⁵ cells/100 μ l.

5 The cells are then plated as needed depending upon the type of assay to be performed.

Stable expression

Heterologous DNA can be stably incorporated into host cells, 10 causing the cell to perpetually express a foreign protein. Methods for the delivery of the DNA into the cell are similar to those described above for transient expression but require the co-transfection of an ancillary gene to confer drug resistance on the targeted host cell. The ensuing drug 15 resistance can be exploited to select and maintain cells that have taken up the DNA. An assortment of resistance genes are available including but not restricted to neomycin, kanamycin, and hygromycin. For the purposes of studies concerning the receptor of this invention, stable expression 20 of a heterologous receptor protein is typically carried out in, mammalian cells including but not necessarily restricted to, CHO, HEK293, LM(tk-), etc.

In addition native cell lines that naturally carry and 25 express the nucleic acid sequences for the receptor may be used without the need to engineer the receptor complement.

Membrane preparations

Cell membranes expressing the receptor protein according to 30 this invention are useful for certain types of assays including but not restricted to ligand binding assays, GTP- γ -S binding assays, and others. The specifics of preparing such cell membranes may in some cases be determined by the nature of the ensuing assay but typically involve 35 harvesting whole cells and disrupting the cell pellet by sonication in ice cold buffer (e.g. 20 mM Tris-HCl, 5 mM EDTA, pH 7.4). The resulting crude cell lysate is cleared of cell debris by low speed centrifugation at 200xg for 5 min

at 4°C. The cleared supernatant is then centrifuged at 40,000xg for 20 min at 4°C, and the resulting membrane pellet is washed by suspending in ice cold buffer and repeating the high speed centrifugation step. The final washed membrane 5 pellet is resuspended in assay buffer. Protein concentrations are determined by the method of Bradford (1976) using bovine serum albumin as a standard. The membranes may be used immediately or frozen for later use.

10 Generation of baculovirus

The coding region of DNA encoding the human receptor disclosed herein may be subcloned into pBlueBacIII into existing restriction sites or sites engineered into sequences 5' and 3' to the coding region of the polypeptides. To 15 generate baculovirus, 0.5 µg of viral DNA (BaculoGold) and 3 µg of DNA construct encoding a polypeptide may be co-transfected into 2×10^6 *Spodoptera frugiperda* insect Sf9 cells by the calcium phosphate co-precipitation method, as outlined by Pharmingen (in "Baculovirus Expression Vector 20 System: Procedures and Methods Manual"). The cells then are incubated for 5 days at 27°C.

The supernatant of the co-transfection plate may be collected by centrifugation and the recombinant virus plaque purified. 25 The procedure to infect cells with virus, to prepare stocks of virus and to titer the virus stocks are as described in Pharmingen's manual.

Labeled ligand binding assays

30 Cells expressing the receptor according to this invention may be used to screen for ligands for said receptors, for example, by labeled ligand binding assays. Once a ligand is identified the same assays may be used to identify agonists or antagonists of the receptor that may be employed for a 35 variety of therapeutic purposes.

In an embodiment, labeled ligands are placed in contact with either membrane preparations or intact cells expressing the

receptor in multi-well microtiter plates, together with unlabeled compounds, and binding buffer. Binding reaction mixtures are incubated for times and temperatures determined to be optimal in separate equilibrium binding assays. The 5 reaction is stopped by filtration through GF/B filters, using a cell harvester, or by directly measuring the bound ligand. If the ligand was labeled with a radioactive isotope such as ^3H , ^{14}C , ^{125}I , ^{35}S , ^{32}P , ^{33}P , etc., the bound ligand may be detected by using liquid scintillation counting, 10 scintillation proximity, or any other method of detection for radioactive isotopes. If the ligand was labeled with a fluorescent compound, the bound labeled ligand may be measured by methods such as, but not restricted to, fluorescence intensity, time resolved fluorescence, 15 fluorescence polarization, fluorescence transfer, or fluorescence correlation spectroscopy. In this manner agonist or antagonist compounds that bind to the receptor may be identified as they inhibit the binding of the labeled ligand to the membrane protein or intact cells expressing the 20 receptor. Non-specific binding is defined as the amount of labeled ligand remaining after incubation of membrane protein in the presence of a high concentration (e.g., 100-1000 X K_D) of unlabeled ligand. In equilibrium saturation binding assays membrane preparations or intact cells transfected with 25 the receptor are incubated in the presence of increasing concentrations of the labeled compound to determine the binding affinity of the labeled ligand. The binding affinities of unlabeled compounds may be determined in equilibrium competition binding assays, using a fixed 30 concentration of labeled compound in the presence of varying concentrations of the displacing ligands.

Functional assays

Cells expressing the SNORF25 receptor DNA may be used to 35 screen for ligands to SNORF25 receptor using functional assays. Once a ligand is identified the same assays may be used to identify agonists or antagonists of the SNORF25 receptor that may be employed for a variety of therapeutic

purposes. It is well known to those in the art that the over-expression of a GPCR can result in the constitutive activation of intracellular signaling pathways. In the same manner, over-expression of the SNORF25 receptor in any cell 5 line as described above, can result in the activation of the functional responses described below, and any of the assays herein described can be used to screen for both agonist and antagonist ligands of the SNORF25 receptor.

10 A wide spectrum of assays can be employed to screen for the presence of SNORF25 receptor ligands. These assays range from traditional measurements of total inositol phosphate accumulation, cAMP levels, intracellular calcium mobilization, and potassium currents, for example; to systems 15 measuring these same second messengers but which have been modified or adapted to be of higher throughput, more generic and more sensitive; to cell based assays reporting more general cellular events resulting from receptor activation such as metabolic changes, differentiation, cell 20 division/proliferation. Description of several such assays follow.

Cyclic AMP (cAMP) assay

The receptor-mediated stimulation or inhibition of cyclic AMP 25 (cAMP) formation may be assayed in cells expressing the receptors. Cells are plated in 96-well plates or other vessels and preincubated in a buffer such as HEPES buffered saline (NaCl (150 mM), CaCl₂ (1 mM), KCl (5 mM), glucose (10 mM)) supplemented with a phosphodiesterase inhibitor such as 30 5mM theophylline, with or without protease inhibitor cocktail (For example, a typical inhibitor cocktail contains 2 µg/ml aprotinin, 0.5 mg/ml leupeptin, and 10 µg/ml phosphoramidon.) for 20 min at 37°C, in 5% CO₂. Test compounds are added with or without 10 mM forskolin and incubated for an additional 35 10 min at 37°C. The medium is then aspirated and the reaction stopped by the addition of 100 mM HCl or other methods. The plates are stored at 4°C for 15 min, and the cAMP content in the stopping solution is measured by

radioimmunoassay. Radioactivity may be quantified using a gamma counter equipped with data reduction software. Specific modifications may be performed to optimize the assay for the receptor or to alter the detection method of cAMP.

5

Arachidonic acid release assay

Cells expressing the receptor are seeded into 96 well plates or other vessels and grown for 3 days in medium with supplements. ^3H -arachidonic acid (specific activity = 0.75 10 $\mu\text{Ci}/\text{ml}$) is delivered as a 100 μL aliquot to each well and samples are incubated at 37°C, 5% CO₂ for 18 hours. The labeled cells are washed three times with medium. The wells are then filled with medium and the assay is initiated with the addition of test compounds or buffer in a total volume 15 of 250 μL . Cells are incubated for 30 min at 37°C, 5% CO₂. Supernatants are transferred to a microtiter plate and evaporated to dryness at 75°C in a vacuum oven. Samples are then dissolved and resuspended in 25 μL distilled water. Scintillant (300 μL) is added to each well and samples are 20 counted for ^3H in a Trilux plate reader. Data are analyzed using nonlinear regression and statistical techniques available in the GraphPAD Prism package (San Diego, CA).

Inositol phosphate assay

25 SNORF25 receptor-mediated activation of the inositol phosphate (IP) second messenger pathways can be assessed by radiometric measurement of IP products.

30 In a 96 well microplate format assay, cells are plated at a density of 70,000 cells per well and allowed to incubate for 24 hours. The cells are then labeled with 0.5 μCi [^3H]-myo-inositol overnight at 37°C, 5% CO₂. Immediately before the assay, the medium is removed and replaced with 90 μL of PBS containing 10 mM LiCl. The plates are then 35 incubated for 15 min at 37°C, 5% CO₂. Following the incubation, the cells are challenged with agonist (10 $\mu\text{l}/\text{well}$; 10x concentration) for 30 min at 37°C, 5% CO₂. The challenge is terminated by the addition of 100 μL of 50% v/v

trichloroacetic acid, followed by incubation at 4°C for greater than 30 minutes. Total IPs are isolated from the lysate by ion exchange chromatography. Briefly, the lysed contents of the wells are transferred to a Multiscreen HV 5 filter plate (Millipore) containing Dowex AG1-X8 (200-400 mesh, formate form). The filter plates are prepared adding 100 µL of Dowex AG1-X8 suspension (50% v/v, water: resin) to each well. The filter plates are placed on a vacuum manifold to wash or elute the resin bed. Each well is first washed 10 2 times with 200 µL of 5 mM myo-inositol. Total [³H]inositol phosphates are eluted with 75 µL of 1.2M ammonium formate/0.1M formic acid solution into 96-well plates. 200 µL of scintillation cocktail is added to each well, and the radioactivity is determined by liquid scintillation counting.

15

Intracellular calcium mobilization assays

The intracellular free calcium concentration may be measured by microspectrofluorimetry using the fluorescent indicator dye Fura-2/AM (Bush et al, 1991). Cells expressing the 20 receptor are seeded onto a 35 mm culture dish containing a glass coverslip insert and allowed to adhere overnight. Cells are then washed with HBS and loaded with 100 µL of Fura-2/AM (10 µM) for 20 to 40 min. After washing with HBS to remove the Fura-2/AM solution, cells are equilibrated in 25 HBS for 10 to 20 min. Cells are then visualized under the 40X objective of a Leitz Fluovert FS microscope and fluorescence emission is determined at 510 nM with excitation wavelengths alternating between 340 nM and 380 nM. Raw fluorescence data are converted to calcium concentrations 30 using standard calcium concentration curves and software analysis techniques.

In another method, the measurement of intracellular calcium can also be performed on a 96-well (or higher) format and 35 with alternative calcium-sensitive indicators, preferred examples of these are: aequorin, Fluo-3, Fluo-4, Fluo-5, Calcium Green-1, Oregon Green, and 488 BAPTA. After activation of the receptors with agonist ligands the emission

elicited by the change of intracellular calcium concentration can be measured by a luminometer, or a fluorescence imager; a preferred example of this is the fluorescence imager plate reader (FLIPR).

5

Cells expressing the receptor of interest are plated into clear, flat-bottom, black-wall 96-well plates (Costar) at a density of 80,000-150,000 cells per well and allowed to incubate for 48 hr at 5% CO₂, 37°C. The growth medium is 10 aspirated and 100 µl of loading medium containing fluo-3 dye is added to each well. The loading medium contains: Hank's BSS (without phenol red) (Gibco), 20 mM HEPES (Sigma), 0.1 or 1% BSA (Sigma), dye/pluronic acid mixture (e.g. 1 mM Flou-3, AM (Molecular Probes) and 10% pluronic acid (Molecular 15 Probes) mixed immediately before use), and 2.5 mM probenecid (Sigma) (prepared fresh). The cells are allowed to incubate for about 1 hour at 5% CO₂, 37°C.

During the dye loading incubation the compound plate is 20 prepared. The compounds are diluted in wash buffer (Hank's BSS (without phenol red), 20 mM HEPES, 2.5 mM probenecid) to a 4X final concentration and aliquoted into a clear v-bottom plate (Nunc). Following the incubation the cells are washed 25 to remove the excess dye. A Denley plate washer is used to gently wash the cells 4 times and leave a 100 µl final volume of wash buffer in each well. The cell plate is placed in the center tray and the compound plate is placed in the right tray of the FLIPR. The FLIPR software is setup for the experiment, the experiment is run and the data are collected. 30 The data are then analyzed using an excel spreadsheet program.

Antagonist ligands are identified by the inhibition of the signal elicited by agonist ligands.

35

GTpS functional assay

Membranes from cells expressing the receptor are suspended in assay buffer (e.g., 50 mM Tris, 100 mM NaCl, 5 mM MgCl₂,

10 μ M GDP, pH 7.4) with or without protease inhibitors (e.g., 0.1% bacitracin). Membranes are incubated on ice for 20 minutes, transferred to a 96-well Millipore microtiter GF/C filter plate and mixed with GTP γ ³⁵S (e.g., 250,000 5 cpm/sample, specific activity ~1000 Ci/mmol) plus or minus unlabeled GTP γ S (final concentration = 100 μ M). Final membrane protein concentration \approx 90 μ g/ml. Samples are incubated in the presence or absence of test compounds for 30 min. at room temperature, then filtered on a Millipore 10 vacuum manifold and washed three times with cold (4⁰C) assay buffer. Samples collected in the filter plate are treated with scintillant and counted for ³⁵S in a Trilux (Wallac) liquid scintillation counter. It is expected that optimal 15 results are obtained when the receptor membrane preparation is derived from an appropriately engineered heterologous expression system, i.e., an expression system resulting in high levels of expression of the receptor and/or expressing G-proteins having high turnover rates (for the exchange of GDP for GTP). GTP γ S assays are well-known to those skilled 20 in the art, and it is contemplated that variations on the method described above, such as are described by Tian et al. (1994) or Lazareno and Birdsall (1993), may be used.

Microphysiometric assay

25 Because cellular metabolism is intricately involved in a broad range of cellular events (including receptor activation of multiple messenger pathways), the use of microphysiometric measurements of cell metabolism can in principle provide a generic assay of cellular activity arising from the 30 activation of any receptor regardless of the specifics of the receptor's signaling pathway.

General guidelines for transient receptor expression, cell preparation and microphysiometric recording are described 35 elsewhere (Salon, J.A. and Owicki, J.A., 1996). Typically cells expressing receptors are harvested and seeded at 3 x 10⁵ cells per microphysiometer capsule in complete media 24 hours prior to an experiment. The media is replaced with

serum free media 16 hours prior to recording to minimize non-specific metabolic stimulation by assorted and ill-defined serum factors. On the day of the experiment the cell capsules are transferred to the microphysiometer and 5 allowed to equilibrate in recording media (low buffer RPMI 1640, no bicarbonate, no serum (Molecular Devices Corporation, Sunnyvale, CA) containing 0.1% fatty acid free BSA), during which a baseline measurement of basal metabolic activity is established.

10

A standard recording protocol specifies a 100 μ l/min flow rate, with a 2 min total pump cycle which includes a 30 sec flow interruption during which the acidification rate measurement is taken. Ligand challenges involve a 1 min 20 sec exposure to the sample just prior to the first post challenge rate measurement being taken, followed by two additional pump cycles for a total of 5 min 20 sec sample exposure. Typically, drugs in a primary screen are presented to the cells at 10 μ M final concentration. Follow up 15 experiments to examine dose-dependency of active compounds are then done by sequentially challenging the cells with a drug concentration range that exceeds the amount needed to generate responses ranging from threshold to maximal levels. Ligand samples are then washed out and the acidification 20 rates reported are expressed as a percentage increase of the peak response over the baseline rate observed just prior to challenge.

MAP kinase assay

30 MAP kinase (mitogen activated kinase) may be monitored to evaluate receptor activation. MAP kinase is activated by multiple pathways in the cell. A primary mode of activation involves the ras/raf/MEK/MAP kinase pathway. Growth factor (tyrosine kinase) receptors feed into this pathway via 35 SHC/Grb-2/SOS/ras. Gi coupled receptors are also known to activate ras and subsequently produce an activation of MAP kinase. Receptors that activate phospholipase C (such as Gq/G11-coupled) produce diacylglycerol (DAG) as a consequence

of phosphatidyl inositol hydrolysis. DAG activates protein kinase C which in turn phosphorylates MAP kinase.

MAP kinase activation can be detected by several approaches.

5 One approach is based on an evaluation of the phosphorylation state, either unphosphorylated (inactive) or phosphorylated (active). The phosphorylated protein has a slower mobility in SDS-PAGE and can therefore be compared with the unstimulated protein using Western blotting. Alternatively,

10 antibodies specific for the phosphorylated protein are available (New England Biolabs) which can be used to detect an increase in the phosphorylated kinase. In either method, cells are stimulated with the test compound and then extracted with Laemmli buffer. The soluble fraction is

15 applied to an SDS-PAGE gel and proteins are transferred electrophoretically to nitrocellulose or Immobilon. Immunoreactive bands are detected by standard Western blotting technique. Visible or chemiluminescent signals are recorded on film and may be quantified by densitometry.

20 Another approach is based on evaluation of the MAP kinase activity via a phosphorylation assay. Cells are stimulated with the test compound and a soluble extract is prepared. The extract is incubated at 30°C for 10 min with

25 gamma-³²P-ATP, an ATP regenerating system, and a specific substrate for MAP kinase such as phosphorylated heat and acid stable protein regulated by insulin, or PHAS-I. The reaction is terminated by the addition of H₃PO₄ and samples are transferred to ice. An aliquot is spotted onto Whatman P81

30 chromatography paper, which retains the phosphorylated protein. The chromatography paper is washed and counted for ³²P in a liquid scintillation counter. Alternatively, the cell extract is incubated with gamma-³²P-ATP, an ATP regenerating system, and biotinylated myelin basic protein

35 bound by streptavidin to a filter support. The myelin basic protein is a substrate for activated MAP kinase. The phosphorylation reaction is carried out for 10 min at 30°C. The extract can then be aspirated through the filter, which

retains the phosphorylated myelin basic protein. The filter is washed and counted for ^{32}P by liquid scintillation counting.

5 Cell proliferation assay

Receptor activation of the receptor may lead to a mitogenic or proliferative response which can be monitored via ^{3}H -thymidine uptake. When cultured cells are incubated with ^{3}H -thymidine, the thymidine translocates into the nuclei 10 where it is phosphorylated to thymidine triphosphate. The nucleotide triphosphate is then incorporated into the cellular DNA at a rate that is proportional to the rate of cell growth. Typically, cells are grown in culture for 1-3 days. Cells are forced into quiescence by the removal of 15 serum for 24 hrs. A mitogenic agent is then added to the media. Twenty-four hours later, the cells are incubated with ^{3}H -thymidine at specific activities ranging from 1 to 10 $\mu\text{Ci}/\text{ml}$ for 2-6 hrs. Harvesting procedures may involve trypsinization and trapping of cells by filtration over GF/C 20 filters with or without a prior incubation in TCA to extract soluble thymidine. The filters are processed with scintillant and counted for ^{3}H by liquid scintillation counting. Alternatively, adherent cells are fixed in MeOH 25 or TCA, washed in water, and solubilized in 0.05% deoxycholate/0.1 N NaOH. The soluble extract is transferred to scintillation vials and counted for ^{3}H by liquid scintillation counting.

Alternatively, cell proliferation can be assayed by measuring 30 the expression of an endogenous or heterologous gene product, expressed by the cell line used to transfect the receptor, which can be detected by methods such as, but not limited to, fluorescence intensity, enzymatic activity, immunoreactivity, DNA hybridization, polymerase chain reaction, etc.

35

Promiscuous second messenger assays

It is not possible to predict, a priori and based solely upon the GPCR sequence, which of the cell's many different

signaling pathways any given receptor will naturally use. It is possible, however, to coax receptors of different functional classes to signal through a pre-selected pathway through the use of promiscuous G_{α} subunits. For example, by

5 providing a cell based receptor assay system with an endogenously supplied promiscuous G_{α} subunit such as $G_{\alpha 15}$ or $G_{\alpha 16}$ or a chimeric G_{α} subunit such as $G_{\alpha QZ}$, a GPCR, which might normally prefer to couple through a specific signaling pathway (e.g., G_s , G_i , G_q , G_0 , etc.), can be made to couple

10 through the pathway defined by the promiscuous G_{α} subunit and upon agonist activation produce the second messenger associated with that subunit's pathway. In the case of $G_{\alpha 15}$, $G_{\alpha 16}$ and/or $G_{\alpha QZ}$ this would involve activation of the G_q pathway and production of the second messenger IP_3 . Through

15 the use of similar strategies and tools, it is possible to bias receptor signaling through pathways producing other second messengers such as Ca^{++} , cAMP, and K^+ currents, for example (Milligan, 1999).

20 It follows that the promiscuous interaction of the exogenously supplied G_{α} subunit with the receptor alleviates the need to carry out a different assay for each possible signaling pathway and increases the chances of detecting a functional signal upon receptor activation.

25

Methods for recording currents in *Xenopus* oocytes

Oocytes are harvested from *Xenopus laevis* and injected with mRNA transcripts as previously described (Quick and Lester, 1994; Smith et al., 1997). The test receptor of this

30 invention and G_{α} subunit RNA transcripts are synthesized using the T7 polymerase ("Message Machine," Ambion) from linearized plasmids or PCR products containing the complete coding region of the genes. Oocytes are injected with 10 ng synthetic receptor RNA and incubated for 3-8 days at 17

35 degrees. Three to eight hours prior to recording, oocytes are injected with 500 pg promiscuous G_{α} subunits mRNA in order to observe coupling to Ca^{++} activated Cl^- currents. Dual electrode voltage clamp (Axon Instruments Inc.) is

performed using 3 M KCl-filled glass microelectrodes having resistances of 1-2 M Ω m. Unless otherwise specified, oocytes are voltage clamped at a holding potential of -80 mV. During recordings, oocytes are bathed in continuously flowing (1-3 ml/min) medium containing 96 mM NaCl, 2 mM KCl, 1.8 mM CaCl₂, 1 mM MgCl₂, and 5 mM HEPES, pH 7.5 (ND96). Drugs are applied either by local perfusion from a 10 μ l glass capillary tube fixed at a distance of 0.5 mm from the oocyte, or by switching from a series of gravity fed perfusion lines.

10

Other oocytes may be injected with a mixture of receptor mRNAs and synthetic mRNA encoding the genes for G-protein-activated inward rectifier channels (GIRK1 and GIRK4, U.S. Patent Nos. 5,734,021 and 5,728,535 or GIRK1 and GIRK2) or any other appropriate combinations (see, e.g., Inanobe et al., 1999). Genes encoding G-protein inwardly rectifying K⁺ (GIRK) channels 1, 2 and 4 (GIRK1, GIRK2, and GIRK4) may be obtained by PCR using the published sequences (Kubo et al., 1993; Dascal et al., 1993; Krapivinsky et al., 1995 and 1995b) to derive appropriate 5' and 3' primers. Human heart or brain cDNA may be used as template together with appropriate primers.

Heterologous expression of GPCRs in *Xenopus* oocytes has been widely used to determine the identity of signaling pathways activated by agonist stimulation (Gundersen et al., 1983; Takahashi et al., 1987). Activation of the phospholipase C (PLC) pathway is assayed by applying a test compound in ND96 solution to oocytes previously injected with mRNA for the SNORF25 receptor and observing inward currents at a holding potential of approximately -80 mV. The appearance of currents that reverse at -25 mV and display other properties of the Ca⁺⁺-activated Cl⁻ channel is indicative of receptor-activation of PLC and release of IP₃ and intracellular Ca⁺⁺. Such activity is exhibited by GPCRs that couple to G_q or G₁₁.

Involvement of the G_{i/o} class of G-proteins in GPCR-stimulated

Ca^{++} -activated Cl^- currents is evaluated using PTX, a toxin which inactivates $\text{G}_{i/o}$ G-proteins. Oocytes are injected with 25 ng PTX/oocyte and modulation of Ca^{++} -activated Cl^- currents by SNORF25 receptor is evaluated 2-5 h subsequently.

5

Elevation of intracellular cAMP can be monitored in oocytes by expression of the cystic fibrosis transmembrane conductance regulator (CFTR) whose Cl^- -selective pore opens in response to phosphorylation by protein kinase A (Riordan, 10 1993). In order to prepare RNA transcripts for expression in oocytes, a template was created by PCR using 5' and 3' primers derived from the published sequence of the CFTR gene (Riordan, 1993). The 5' primer included the sequence coding for T7 polymerase so that transcripts could be generated 15 directly from the PCR products without cloning. Oocytes were injected with 10 ng of CFTR mRNA in addition to 10-15 ng mRNA for SNORF25. Electrophysiological recordings were made in ND96 solution after a 2-3 day incubation at 18°C. Currents are recorded under dual electrode voltage clamp (Axon 20 Instruments Inc.) with 3 M KCl-filled glass microelectrodes having resistances of 1-2 Mohm. Unless otherwise specified, oocytes are voltage clamped at a holding potential of -80 mV. During recordings, oocytes are bathed in continuously flowing (1-3 ml/min) medium containing 96 mM NaCl, 2 mM KCl, 1.8 mM 25 CaCl_2 , 1 mM MgCl_2 , and 5 mM HEPES, pH 7.5 (ND96). Drugs are applied either by local perfusion from a 10 μl glass capillary tube fixed at a distance of 0.5 mm from the oocyte, or by switching from a series of gravity fed perfusion lines.

30 Activation of G-protein G_i and G_o can be monitored by measuring the activity of inwardly rectifying K^+ (potassium) channels (GIRKs). Activity may be monitored in oocytes that have been co-injected with mRNAs encoding the mammalian receptor plus GIRK subunits. GIRK gene products co-assemble 35 to form a G-protein activated potassium channel known to be activated (i.e., stimulated) by a number of GPCRs that couple to G_i or G_o (Kubo et al., 1993; Dascal et al., 1993). Oocytes expressing the mammalian receptor plus the GIRK

subunits are tested for test compound responsivity by measuring K⁺ currents in elevated K⁺ solution containing 49 mM K⁺.

5 **Localization of mRNA coding for human and rat SNORF25.**

Methods: Quantitative RT-PCR using a fluorogenic probe with real time detection.

10 Quantitative RT-PCR using fluorogenic probes and a panel of mRNA extracted from human and rat tissue was used to characterize the localization of SNORF25 rat and human RNA.

15 This assay utilizes two oligonucleotides for conventional PCR amplification and a third specific oligonucleotide probe that is labeled with a reporter at the 5' end and a quencher at the 3' end of the oligonucleotide. In the instant invention, FAM (6-carboxyfluorescein) and JOE (6

20 carboxy-4,5-dichloro-2,7-dimethoxyfluorescein) were the two reporters that were utilized and TAMRA (6-carboxy-4,7,2,7'-tetramethylrhodamine) was the quencher. As amplification progresses, the labelled oligonucleotide probe hybridizes to the gene sequence between the two

25 oligonucleotides used for amplification. The nuclease activity of Taq, or rTth thermostable DNA polymerases is utilized to cleave the labelled probe. This separates the quencher from the reporter and generates a fluorescent signal that is directly proportional to the amount of amplicon

30 generated. This labelled probe confers a high degree of specificity. Non-specific amplification is not detected as the labelled probe does not hybridize. All experiments were conducted in a PE7700 Sequence Detection System (Perkin Elmer, Foster City, CA).

35

Quantitative RT-PCR

For the detection of RNA encoding SNORF25, quantitative

RT-PCR was performed on mRNA extracted from tissue. Reverse transcription and PCR reactions were carried out in 50 μ l volumes using rTth thermostable DNA polymerase (Perkin Elmer). Primers with the following sequences were used:

5

SNORF 25 human:

Forward primer:

SNORF25H-765F

10 5'-CCTCTACCTAGTGCTGGAACGG-3' (SEQ ID NO: 18)

Reverse primer

SNORF25H-868R

5'-GCTGCAGTCGCACCTCCT-3' (SEQ ID NO: 19)

15

Fluorogenic oligonucleotide probe:

SNORF25H-814T

5' (6-FAM) -TCCCTGCTAACCCACTCATCTATGCCTATT- (TAMRA) 3' (SEQ ID NO: 20)

20

SNORF25 rat

forward primer

SNORF25R-231F

25 5'-GTGTAGCCTCGGATGGCA-3' (SEQ ID NO: 21)

reverse primer

SNORF25R-329R

5'-GGCTGCTTAATGGCCAGGTAC-3' (SEQ ID NO: 22)

30

Fluorogenic oligonucleotide probe:

SNORF25R-278T

5' (6-FAM) -TCCTCACGGTCATGCTGATTGCCTTT- (TAMRA) 3' (SEQ ID NO: 23)

35

Using these primer pairs, amplicon length is 104 bp for human SNORF25 and 99 bp for rat SNORF25. Each RT-PCR reaction contained 50 ng mRNA. Oligonucleotide concentrations were:

500 nM of forward and reverse primers, and 200 nM of fluorogenic probe. Concentrations of reagents in each reaction were: 300 μ M each of dGTP; dATP; dCTP; 600 μ M UTP; 3.0mM Mn(OAc)₂; 50 mM Bicine; 115 mM potassium acetate, 8% 5 glycerol, 5 units rTth thermostable DNA polymerase, and 0.5 units of uracil N-glycosylase. Buffer for RT-PCR reactions also contained a fluor used as a passive reference (ROX: Perkin Elmer proprietary passive reference I). All reagents for RT-PCR (except mRNA and oligonucleotide primers) were 10 obtained from Perkin Elmer (Foster City, CA). Reactions were carried using the following thermal cycler profile: 50°C 2 min., 60°C 30 min., 95°C 5 min., followed by 40 cycles of: 94°C, 20 sec., 62°C 1 min.

15 Positive controls for PCR reactions consisted of amplification of the target sequence from a plasmid construct. Standard curves for quantitation were constructed using the human SNORF25 gene in a plasmid vector or RNA extracted from pancreas as a template for amplification.

20 Negative controls consisted of mRNA blanks, as well as primer and mRNA blanks. To confirm that the mRNA was not contaminated with genomic DNA, PCR reactions were carried out without reverse transcription using Taq DNA polymerase.

25 Integrity of RNA was assessed by amplification of mRNA coding for cyclophilin or glyceraldehyde 3-phosphate dehydrogenase (GAPDH). Following reverse transcription and PCR amplification, data was analyzed using Perkin Elmer sequence detection software. The fluorescent signal from each well was normalized using an internal passive reference, and data 30 was fitted to a standard curve to obtain relative quantities of SNORF25 mRNA expression.

RESULTS AND DISCUSSIONCloning of the full-length sequence of SNORF25

5 Genomic DNA and cDNA prepared from several tissues (including GH1 cells and Rin14b cells) was subjected to MOPAC PCR with two degenerate primers designed based on the third transmembrane domain of the members of the galanin, somatostatin, and opioid receptor families and the seventh
10 transmembrane domain of members of the galanin receptor family. Three products from this reaction were found to be the same clone in either orientation (forward or reverse), which was a novel sequence not found in the Genbank, SwissProtPlus, GSS, EST, or STS databases. It contained
15 significant homology to other known G protein-coupled receptors (~29% identity to the known receptors dopamine D1, beta-adrenergic 2b and 5-HT_{1F}; 34% identity to 5-HT receptor). This receptor sequence was later named SNORF25,
20 and was used to design primers for 5' and 3' Rapid Amplification of cDNA Ends (RACE), as described in the Methods section above. The 5' RACE reaction yielded sequence information through the first transmembrane domain and a putative in-frame initiating methionine-coding sequence surrounded by a kozak consensus sequence (ACCATGG).
25

The 3' RACE reaction yielded a 600 bp band by agarose gel electrophoresis. This band was subcloned into the TA cloning kit, and isolated colonies were sequenced. The sequence of these products revealed the presence of an in-frame stop
30 codon downstream from the region coding for the seventh transmembrane domain. The entire size of the coding sequence of SNORF25 was determined to be 1005 bp, coding for a protein of 335 amino acids. Two primers, JAB86 and JAB84, were used to amplify the entire coding sequence from Rin14b cell line
35 cDNA and rat genomic DNA using the Expand Long PCR system. The primers for this reaction were specific to the 5' and 3' untranslated regions of SNORF25 with *Bam*HI and *Hind*III restriction sites incorporated into the 5' ends of the 5' and

3' primers, respectively. When the products of these reactions were subcloned into pcDNA3.1(-) and sequenced, the sequence of the Rin14b clone and the genomic clone were found to be identical, and the vector construct containing rat 5 SNORF25 was named pcDNA3.1-rSNORF25.

Hydophobicity (Kyte-Doolittle) analysis of the amino acid sequence of the full-length clone indicates the presence of seven hydrophobic regions, which is consistent with the seven 10 transmembrane domains of a G protein-coupled receptor. The seven expected transmembrane domains are indicated in Figure 4. A comparison of nucleotide and peptide sequences of rat SNORF25 with sequences contained in the Genbank, EMBL, and SwissProtPlus databases reveals that the amino acid sequence 15 of this receptor is most related to histamine, adenosine, serotonin, beta adrenergic, and dopamine receptor families, displaying between 25-30% overall amino acid identity with these receptors. The N- and C-termini are relatively short, much like the adenosine receptor family. However, 20 transmembrane domain analysis indicates that this receptor shares a significant degree of identity to other GPCRs in its transmembrane domains. A comparison of all of the transmembrane domains of SNORF25 simultaneously with a comprehensive list of GPCR transmembrane domains would 25 suggest that the transmembrane domains of SNORF25 have the highest degree of identity with the beta adrenergic receptors 1 and 2 of 31% and 32%, respectively, as well as 5-HT₇ and 5-hT_{5B} receptors of 32% and 36.6%, respectively. When transmembrane domains are analyzed individually by a FASTA 30 search, SNORF25 exhibits considerable similarity to the transmembrane domains of a variety of known G protein-coupled receptors.

In order to clone the human homolog of SNORF25, a human 35 genomic cosmid library was screened at medium stringency with labelled oligonucleotide probes designed based on the second and fifth transmembrane domains of rat SNORF25. Out of roughly 225,000 colonies screened, two colonies hybridized

to the probes. After isolation and analysis of each colony, these two clones were determined to be identical cosmid clones containing the human homolog of SNORF25. Southern blot analysis of several restriction digests of this cosmid and subsequent sequencing of positive bands indicated that a *Bam*HI/*Hind*III digest of this cosmid yielded a 1.9 kb fragment containing the full-length coding sequence of this human clone. The construct of the human receptor subcloned into the *Bam*HI/*Hind*III site of the pEXJT3T7 vector is named 10 pEXJT3T7-hSNORF25. Human SNORF25 exhibits an 80% DNA identity and 83% amino acid identity to rat SNORF25. Like the rat receptor, the protein-coding region of human SNORF25 is 1005 nucleotides (Figures 1A-1B), coding for a protein of 335 amino acids (Figures 2A-2B). The DNA and amino acid 15 sequences of rat SNORF25 are shown in Figures 3A-3B and 4A-4B, respectively.

A search of the GenEMBL, SwissProtPlus, EST, STS and GSS databases confirmed that human SNORF25 is also a unique novel 20 sequence. Other than its identity with rat SNORF25, it shares 28-30% overall identity with adenosine 2a, 5-HT_{4L}, 5-HT_{4S}, 5-H_{T6}, and 5-HT₇, dopamine D₁ and D₅, and somatostatin 5 receptors. It also shares 25-26% identity with adenosine A1, histamine H1 and 2, beta adrenergic 1, and somatostatin 25 2 and 3 receptors. A comparison of all of the transmembrane domains of human SNORF25 simultaneously with a comprehensive list of GPCR transmembrane domains would suggest that the transmembrane domains of human SNORF25 have the highest degree of identity with the beta 1 and 2 adrenergic receptors 30 (29% and 32%, respectively) and 5-HT₄. Individual transmembrane domains of human SNORF25 share significant identity with transmembrane domains from several other G protein-coupled receptors. .

35 Both rat and human SNORF25 have several potential protein kinase C (PKC) phosphorylation motifs throughout their amino acid sequences. For both receptors, threonine 73, serine 79, and serine 309 are potential PKC phosphorylation sites. The

human receptor has an additional putative PKC phosphorylation site at serine 214, which is a proline in rat SNORF25. Both receptors share a potential casein kinase II (CKII) phosphorylation site at serine 329. The human SNORF25 also 5 contains two more potential CKII phosphorylation sites, threonine 217 and serine 331, that are not present in the rat receptor. Conversely, rat SNORF25 contains a potential tyrosine phosphorylation site at tyrosine 323, which is not present in the human receptor.

10

cAMP response of SNORF25-transfected cells

The expression vector (pcDNA) containing the SNORF25 cDNA was transfected by electroporation method into CHO cells. After plating, the transfectants were challenged with a ligand 15 library that included, among other things, several of the traditional neurotransmitters such as histamine, adenosine, serotonin, norepinephrine, and dopamine, based on homology of SNORF25 to the receptors of these ligands (see above), and tested for their ability to stimulate cAMP or IP release 20 above mock-transfected cells. Interestingly, the basal cAMP levels of SNORF25-transfected cells were significantly higher (>10-fold) than mock-transfected cells (Figure 5). This observation suggested that SNORF25 receptor may functionally be coupled to a cAMP stimulatory pathway. Among the ligands 25 tested, only all-trans retinoic acid (ATRA) produced a significant increase in cAMP but not IP release in SNORF25-transfected cells, without affecting these parameters in mock-transfected CHO cells. The response produced at 10 μ M concentration of ATRA (2- to 5-fold above basal) was 30 comparable to that produced by forskolin, a potent direct stimulator of adenylyl cyclase (Figure 6) (n=3).

Responses to forskolin in both mock- and SNORF25-transfected Cos-7 cells were almost identical (Figure 6), suggesting that 35 the enhanced maximal response to ATRA observed in SNORF25-expressing cells, as compared to mock DNA-transfected cells, was not due to a change in cell density or in the intrinsic properties of the cells. All-trans retinol

(vitamin A₁), a close analogue of ATRA failed to produce an increase in cAMP at 10 μ M (Figure 6).

Subsequent experiments demonstrated that the ATRA-induced 5 increase in cAMP formation was independent of host cell as it was observed also in Cos-7 cells (n=3) (Figure 7). All-trans-retinoic acid produced no response in Cos-7 cells transfected with other known cyclase-stimulatory receptors including dopamine D₁, D₅, serotonin 5-HT₄ and 5-HT₆ 10 receptors, indicating that the response observed to ATRA is specific to SNORF25-transfected cells (Figure 7).

The cAMP response to ATRA in Cos-7 cells was concentration-dependent with EC₅₀ values ranging from 15 approximately 0.2 to 1 μ M and E_{max} of approximately 200-300% (Figure 8).

Activation of calcium-activated Cl⁻ currents in SNORF25 expressing Xenopus oocytes

20 Elevation of intracellular cAMP can be monitored in oocytes by expression of the cystic fibrosis transmembrane conductance regulator (CFTR) whose Cl⁻-selective pore opens in response to phosphorylation by protein kinase A (Riordan, 1993). The activity of SNORF25 was therefore tested in 25 oocytes co-injected with mRNA encoding SNORF25 and mRNA encoding CFTR. In 17 out of 39 of these oocytes an inward Cl⁻ current (105 \pm 20 nA) was measured in response to the application of 10 μ M all-trans-retinoic acid (See Figures 9A-9C and 10).

30 This response was specific to the expression of SNORF25 since no such current was observed in other oocytes injected with only mRNA encoding the CFTR channel. Similar currents were observed in oocytes injected with the β 2-adrenergic receptor 35 (B2AR) (See Figure 9C), although the currents generated by SNORF25-expressing oocytes were generally 2-3 fold slower and smaller. All-trans-retinoic acid did not stimulate Cl⁻ currents in oocytes lacking CFTR, indicating that the

Gq-mediated phospholipase C pathway was not activated. Responses also were not evoked in oocytes expressing chimeric G-proteins which are able to couple Gi and Go coupled GPCRs to the phospholipase C pathway. Taken together, these 5 observations support the hypothesis that SNORF25 encodes a GPCR which binds all-trans-retinoic acid and stimulates the production of cAMP, presumably via activation of Gs.

In other systems, all-trans-retinoic acid stimulates one of 10 several nuclear receptors (see background). This results in the enhancement of transcription of one or more genes. SNORF25 expression in oocytes could result in the expression of a nuclear receptor for all-trans-retinoic acid, not normally present in uninjected oocytes, that when stimulated 15 produces an elevation of cAMP. If this were the case, then retinoic acid would not necessarily bind the SNORF25 receptor, but would act on a previously known or novel nuclear receptor for retinoic acid. This indirect mechanism of action of retinoic acid may explain why the ligand failed to 20 elicit a CFTR response in 3 out of 6 batches of oocytes (17 of 39 oocytes), and why the kinetics of CFTR activation were 2-3 times slower than those observed under conditions where responses were evoked by activation of well-characterized GPCRs such as the B2 adrenergic receptor (Figure 9C). 25 Nevertheless, the delay for activation of CFTR by retinoic acid was on the order of 10 seconds, and the activation of nuclear receptors is typically in the range of several minutes to hours. Thus, while we cannot rule out an indirect mechanism of action of retinoic acid, the relatively rapid 30 onset of the response in SNORF25-expressing oocytes suggests that such a mechanism is unlikely.

Detection of mRNA coding for human SNORF25:

35 mRNA was isolated from multiple tissues (listed in Table 1) and assayed as described.

Quantitative RT-PCR using a fluorogenic probe demonstrated expression of mRNA encoding human SNORF25 in most tissues assayed (Table 1). Highest levels of human SNORF25 mRNA are found in the pancreas, stomach, small intestine and fetal 5 liver, with lower levels detected elsewhere. Most nervous system structures showed little expression of SNORF25 mRNA as compared to peripheral organs.

The highest levels of SNORF25 expression are found in the 10 pancreas. The pancreas secretes a variety of broadly active substances (including insulin), indicating that SNORF25 may play a role in regulating multiple metabolic functions, potentially via endocrine mechanisms. SNORF25 expression in the pancreas is not surprising as SNORF25 is also expressed 15 in a rat insulinoma cell line. This finding as well as the detection of SNORF25 mRNA in liver indicate a possible role in the regulation of glucose levels and possibly diabetes.

Other organs with high levels of SNORF25 mRNA are stomach and 20 small intestine. The distribution to these structures is consistent with functions relating to gastrointestinal motility or absorption. It is not known at this time if SNORF25 mRNA is localized to smooth muscle or to mucosal/submucosal layers.

25 Although detected in very low levels, the presence of SNORF25 mRNA in multiple regions of the CNS including the thalamus and hippocampal formation (where levels are highest in the CNS) and other functionally diverse areas, indicate a diffuse 30 regulatory function or regional functionality for this receptor.

Human SNORF25 mRNA appears to be developmentally regulated. In fetal liver, levels of mRNA approach those measured in 35 adult pancreas (83%). However in adult tissue, this drops to less than 1% of the amount found in the pancreas. The profound change of SNORF25 mRNA during development implies a role in the maturation of the liver, or a role in the

regulation of glucose demands/levels during development. The time course of this increase has not been examined and would be important in understanding the function of this receptor.

- 5 In summary, the distribution of SNORF25 receptor mRNA implies broad regulatory functions that involves multiple organ systems, endocrine mechanisms, as well as the central nervous system.

10 **Detection of mRNA coding for rat SNORF25**

Unlike the restricted distribution of human SNORF25 mRNA, the distribution of SNORF25 mRNA in the rat is widespread. One striking difference in the distribution between rat and human 15 is the high levels of SNORF25 mRNA detected in the rat central nervous system. In the human, the highest concentrations of SNORF25 mRNA are found in the pancreas, with very low levels found in CNS structures. In the rat the highest levels of SNORF25 mRNA are found in the hippocampal 20 formation, closely followed by levels detected in the cerebral cortex, cerebellum, hypothalamus, choroid plexus and medulla. SNORF25 mRNA is also detected in both dorsal root and trigeminal ganglia. Although SNORF25 mRNA is detected in rat pancreas and other peripheral organs, it is present 25 there in much lower levels than in the CNS.

Rat SNORF25 was detected in most tissues assayed. In addition to the pancreas it is expressed in appreciable amounts in lung, colon, duodenum, ovary, kidney and the 30 adrenal glands. It was detected in other tissues in decreasing amounts as shown in Table 2.

In summary, the broad distribution of rat SNORF25 receptor mRNA implies broad regulatory functions that involve multiple 35 organ systems, endocrine mechanisms as well as the central nervous system. The difference in the distribution pattern seen between human and rat suggests a broader, and potentially different role for this receptor in the rat as

compared to human.

Table 1

Distribution of mRNA coding for human SNORF25 receptors using qRT-PCR

mRNA encoding SNORF25h is expressed as % of highest
5 expressing tissue.

Region	qRT-PCR % of max	Potential applications
heart	0.31	cardiovascular indications
kidney	0.62	hypertension, electrolyte balance
10 liver	0.18	diabetes
lung	0.32	respiratory disorders, asthma
pancreas	100	diabetes, endocrine disorders
15 pituitary	0.03	endocrine/neuroendocrine regulation
placenta	0.42	gestational abnormalities
small intestine	4.63	gastrointestinal disorders
15 spleen	1.50	immune disorders
stomach	12.60	gastrointestinal disorders
striated muscle	0.32	musculoskeletal disorders
amygdala	0.18	depression, phobias, anxiety, mood disorders
20 caudate-putamen	0.17	modulation of dopaminergic function
cerebellum	0.06	motor coordination
cerebral cortex	0.01	sensory and motor integration, cognition
hippocampus	0.27	cognition/memory
25 spinal cord	0.00	analgesia, sensory modulation and transmission

5	substantia nigra	0.05	modulation of dopaminergic function. modulation of motor coordination.
	thalamus	0.60	sensory integration
	fetal brain	0.14	developmental disorders
	fetal lung	0.04	developmental disorders
	fetal kidney	0.90	developmental disorders
	fetal liver	82.63	developmental disorders

Table 2

Distribution of mRNA coding for rat SNORF25 receptors using qRT-PCR

mRNA encoding SNORF25r is expressed as % of highest
5 expressing tissue.

Tissue	qRT-PCR % of max	Potential applications
adipose tissue	9.08	metabolic disorders
adrenal cortex	8.78	regulation of steroid hormones
adrenal medulla	16.34	regulation of epinephrine release
colon	24.15	gastrointestinal disorders
duodenum	18.89	gastrointestinal disorders
heart	11.98	cardiovascular indications
kidney	15.86	electrolyte balance, hypertension
liver	trace	diabetes
lung	32.57	respiratory disorders, asthma
ovary	17.74	reproductive function
pancreas	30.45	diabetes, endocrine disorders
spleen	n o t detected	immune disorders
stomach	3.44	gastrointestinal disorders
striated muscle	1.04	musculoskeletal disorders

5	testes	5.10	reproductive function
	urinary bladder	7.87	urinary incontinence
	vas deferens	7.16	reproductive function
	celiac plexus	17.82	modulation of autonomic innervation
	cerebellum	84.14	motor coordination
	cerebral cortex	83.54	Sensory and motor integration, cognition
	choroid plexus	66.59	regulation of cerebrospinal fluid
	dorsal root ganglia	38.14	sensory transmission
	hippocampus	100	cognition/memory
	hypothalamus	67.19	appetite/obesity, neuroendocrine regulation
	medulla	52.66	analgesia, motor coordination
	olfactory bulb	6.66	olfaction
	pineal gland	41.16	regulation of melatonin release
	spinal cord	31.72	analgesia, sensory modulation and transmission
	trigeminal ganglia	42.98	sensory transmission

REFERENCES

Bowman, W.C. and Rand, M.J., eds., "The eye and drugs affecting ocular function", In: Textbook of Pharmacology.
5 Second Edition, London: Blackwell Scientific Publications.
p.29.22-29.27 (1980).

Bradford, M.M., "A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding", *Anal. Biochem.* 72: 248-254
10 (1976).

Breitman, T., et al., "Induction of differentiation of human promyelocytic leukemia cell line (HL-60) by retinoic acid",
15 *Proc. Natl. Acad. Sci.* 77: 2936-2940 (1980).

Burns, C.C., et al., "Identification and deletion of sequences required for feline leukemia virus RNA packaging and construction of a high-titer feline leukemia virus
20 packaging cell line", *Virology* 222(1): 14-20 (1996).

Bush, et al., "Nerve growth factor potentiates bradykinin-induced calcium influx and release in PC12 cells",
J. Neurochem. 57: 562-574 (1991).
25

Chertow, B.S., et al., "Cellular mechanisms of insulin release: effects of retinoids on rat islet cell-to-cell adhesion, reaggregation, and insulin release", *Diabetes* 32:
568-574 (1983).

30 Chertow, B.S., et al., "Effects of vitamin A deficiency and repletion on rat insulin secretion in vivo and in vitro from isolated islets", *J. Clin. Invest.* 79: 163-169 (1987).

35 Chertow, B.S., et al., "Cytoplasmic retinoid-binding proteins

and retinoid effects on insulin release in RINm5F -cells", *Diabetes* 38: 1544-1548 (1989).

5 Chu, Y.Y., et al., "Characterization of the rat A2a adenosine receptor gene", *DNA Cell Biol.* 15(4): 329-337 (1996).

Chytil, F., "Retinoids in lung development", *FASEB J.* 10(9): 986-992 (1996).

10 Connor, M.J. and Sidell, N., "Retinoic acid synthesis in normal and Alzheimer diseased brain and human neural cells", *Mol. Chem. Neurophathol.* 30(3): 239-252 (1997).

15 Dascal, N., et al., "A trial G protein-activated K⁺ channel: expression cloning and molecular properties", *Proc. Natl. Acad. Sci. USA* 90:10235-10239 (1993).

20 El-Matwally, T.H. and T.E. Adrian (1999) Optimization of treatment conditions for studying the anticancer effects of retinoids using pancreatic adenocarcinoma as a model. *Biochem. Biophys. Res. Commun.* 257(2): 596-603.

25 Fong, T.M., et al., "Mutational analysis of neurokinin receptor function" *Can. J. Physio. Pharmacol.* 73(7): 860-865 (1995).

Goodman, A.B., "Three independent lines of evidence suggest retinoids as causal to schizophrenia", *PNAS* 95(13): 7240-7244 (1998).

30 Gottardis, M.M., et al., "The efficacy of 9-cis retinoic acid in experimental models of cancer", *Breast Cancer Res. and Treat.* 38: 85-96 (1996).

35 Graziano, M.P. et al., "The amino terminal domain of the glucagon-like peptide-1 receptor is a critical determinant

of subtype specificity" *Receptors Channels* 4(1): 9-17 (1996).

5 Guan, X.M., et al., "Determination of Structural Domains for G Protein Coupling and Ligand Binding in (3 - Adrenergic Receptor" *Mol. Pharmacol.* 48(3): 492-498 (1995).

10 Gudas, L.J., et al. "Cellular biology and biochemistry of the retinoids" In: Sporn, M.B., Roberts, A.B., Goodman, D.S. eds. The retinoids: Biology, chemistry, and medicine. 2nd ed. New York: Raven Press. p. 443-520 (1994).

15 Gundersen, C.B., et al., "Serotonin receptors induced by exogenous messenger RNA in *Xenopus* oocytes" *Proc. R. Soc. Lond. B. Biol. Sci.* 219(1214): 103-109 (1983).

20 Hofman, C. and Eichele, G., "Retinoids in development" In: Sporn, M.B., Roberts, A.B., Goodman, D.S. eds. The retinoids: Biology, chemistry, and medicine. 2nd ed. New York: Raven Press. p. 387-441 (1994).

25 Inanobe, A., et al., "Characterization of G-protein-gated K⁺ channels composed of Kir3.2 subunits in dopaminergic neurons of the substantia nigra" *J. of Neurosci.* 19(3): 1006-1017 (1999).

30 Krapivinsky, G., et al., "The G-protein-gated atrial K⁺ channel IKACH is a heteromultimer of two inwardly rectifying K(+)-channel proteins" *Nature* 374:135-141 (1995).

35 Krapivinsky, G., et al., "The cardiac inward rectifier K⁺ channel subunit, CIR, does not comprise the ATP-sensitive K⁺ channel, IKATP", *J. Biol. Chem.* 270:28777-28779 (1995b).

Kubo, Y., et al., "Primary structure and functional expression of a rat G-protein-coupled muscarinic potassium

channel" *Nature* 364:802-806 (1993).

Lazareno, S. and Birdsall, N.J.M., "Pharmacological characterization of acetylcholine stimulated [³⁵S]-GTPγS binding mediated by human muscarinic m₁-m₄ receptors: antagonist studies", *Br. J. Pharmacol.* 109: 1120-1127 (1993).

Mangelsdorf, D.J., et al. "The retinoid receptors", In: Sporn, M.B., Roberts, A.B., Goodman, D.S. eds. The retinoids: Biology, chemistry, and medicine. 2nd ed. New York: Raven Press. p. 319-349 (1994). Martin, S., et al., "HL-60 cells induced to differentiate towards neutrophils subsequently die via apoptosis" *Clin. Exp. Immunol.* 79: 448-453 (1990).

15 Miller, J., and Germain, R.N. "Efficient cell surface expression of class II MHC molecules in the absence of associated invariant chain", *J. Exp. Med.* 164(5): 1478-1489 (1986).

20 Milligan, G., et al., "Use of chimeric G(proteins in drug discovery" *TIPS* (In press).

25 Ohmura, T., et al., "Induction of cellular DNA synthesis in the pancreas and kidneys of rats by peroxisome proliferators, 9-cis retinoic acid, and 3,3',5-triiodo-L-thyronine", *Cancer Res.* 57: 795-798 (1997).

30 Orfanos, C.E., et al., "Current use and future potential role of retinoids in dermatology", *Drugs* 53(3): 358-388 (1997).

Quick, M.W. and Lester, H.A., "Methods for expression of excitability proteins in *Xenopus* oocytes", *Meth. Neurosci.* 19: 261-279 (1994).

35 Redfern, C.P., et al., "Gene expression and neuroblastoma cell differentiation in response to retinoic acid:

differential effects of 9-cis and all-trans retinoic acid"
Eur. J. Cancer 31A(4): 486-494 (1995).

Riordan, J.R., "The cystic fibrosis transmembrane conductance
5 regulator", *Ann. Rev. Physiol.* 55: 609-630 (1993).

Rosenwicz, S., Wollbergs, K., Von Lampe, B., Matthes, H.,
Kaiser, A., and E.O. Riecken (1997) Retinoids inhibit
adhesion to laminin in human pancreatic carcinoma cells via
10 the alpha 6 beta 1-integrin receptor. *Gastroenterology*
112(2): 532-542.

Rosenwicz, S., Stier, U., Brembeck, F., Kaiser, A.,
Papadimitriou, C.A., Berdel, W.E., Wiedenmann, B., and E.O.
15 Riecken (1995) Retinoids: effects on growth,
differentiation, and nuclear receptor expression in human
pancreatic carcinoma cell lines. *Gastroenterology* 109(5):
1646-1660.

20 Sabichi, A.L., et al., "Retinoids in the chemoprevention of
bladder cancer", *Curr. Opin. Oncol.* 10(5): 479-484 (1998).

Salon, J.A. and Owicki, J.A., "Real-time measurements of
receptor activity: Application of microphysiometric
25 techniques to receptor biology" *Meth. Neurosci.* 25: 201-224
(1996).

Smith, K.E., et al., "Expression cloning of a rat
hypothalamic galanin receptor coupled to phosphoinositide
30 turnover", *J. Biol. Chem.* 272: 24612-24616 (1997).

Sporn, M., and Roberts, A., "Role of retinoids in
differentiation and carcinogenesis" *J. Natl. Cancer Inst.* 73:
1381-1387 (1984).

Spurney, R.F., et al., "The C-terminus of the thromboxane receptor contributes to coupling and desensitization in a mouse mesangial cell line", *J. Pharmacol. Exp. Ther.* 283(1): 207-215 (1997).

5

Takahashi, T., et al., ARat brain serotonin receptors in *Xenopus* oocytes are coupled by intracellular calcium to endogenous channels@ *Proc. Natl. Acad. Sci. USA* 84(14): 5063-5067 (1987)..

10

Tian, W., et al., "Determinants of alpha-Adrenergic Receptor Activation of G protein: Evidence for a Precoupled Receptor/G protein State" *Molecular Pharm.* 45: 524-553 (1994) .

15

Underwood, D.J. et al., "Structural model of antagonist and agonist binding to the angiotensin II, AT1 subtype, G protein coupled receptor", *Chem. Biol.* 1(4): 211-221 (1994) .

20

25

30

35

What is claimed is:

1. An isolated nucleic acid encoding a mammalian SNORF25 receptor.

5

2. The nucleic acid of claim 1, wherein the nucleic acid is DNA.

10

3. The DNA of claim 2, wherein the DNA is cDNA.

10

4. The DNA of claim 2, wherein the DNA is genomic DNA.

5. The nucleic acid of claim 1, wherein the nucleic acid is RNA.

15

6. The nucleic acid of claim 1, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor.

20

7. The nucleic acid of claim 6, wherein the human SNORF25 receptor has an amino acid sequence identical to that encoded by the plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495).

25

8. The nucleic acid of claim 6, wherein the human SNORF25 receptor has an amino acid sequence identical to the amino acid sequence shown in Figures 2A-2B (SEQ ID NO: 2).

30

9. The nucleic acid of claim 1, wherein the mammalian SNORF25 receptor is a rat SNORF25 receptor.

35

10. The nucleic acid of claim 9, wherein the human SNORF25 receptor has an amino acid sequence identical to that encoded by the plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494).

11. The nucleic acid of claim 9, wherein the human SNORF25 receptor has an amino acid sequence identical to the amino acid sequence shown in Figures 4A-4B (SEQ ID NO: 4).

5

12. A purified mammalian SNORF25 receptor protein.

10 13. The purified mammalian SNORF25 receptor protein of claim 9, wherein the SNORF25 receptor protein is a human SNORF25 receptor protein.

14. The purified mammalian SNORF25 receptor protein of claim 9, wherein the SNORF25 receptor protein is a rat SNORF25 receptor protein.

15

15. A vector comprising the nucleic acid of claim 1.

16. A vector comprising the nucleic acid of claim 6.

20 17. A vector of claim 15 or 16 adapted for expression in a cell which comprises the regulatory elements necessary for expression of the nucleic acid in the cell operatively linked to the nucleic acid encoding the receptor so as to permit expression thereof, wherein the cell is a bacterial, amphibian, yeast, insect or mammalian cell.

25 18. The vector of claim 17, wherein the vector is a baculovirus.

30

19. The vector of claim 15, wherein the vector is a plasmid.

35 20. The plasmid of claim 19 designated pEXJT3T7-hSNORF25 (ATCC Accession No. 203495).

21. The plasmid of claim 20 designated pcDNA3.1-rSNORF25 (ATCC Accession No. 203494).

22. A cell comprising the vector of claim 18.
23. A cell of claim 22, wherein the cell is a non-mammalian cell.
5
24. A cell of claim 23, wherein the non-mammalian cell is a *Xenopus* oocyte cell or a *Xenopus* melanophore cell.
25. A cell of claim 22, wherein the cell is a mammalian cell.
10
26. A mammalian cell of claim 25, wherein the cell is a COS-7 cell, a 293 human embryonic kidney cell, a NIH-3T3 cell, a LM(tk-) cell, a mouse Y1 cell, or a CHO cell.
15
27. A cell of claim 22, wherein the cell is an insect cell.
20
28. An insect cell of claim 27, wherein the insect cell is an Sf9 cell, an Sf21 cell or a *Trichoplusia ni* 5B-4 cell.
29. A membrane preparation isolated from the cell of any one of claims 22, 23, 25, 26, 27 or 28.
25
30. A nucleic acid probe comprising at least 15 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF25 receptor, wherein the probe has a sequence complementary to a unique sequence present within one of the two strands of the nucleic acid encoding the mammalian SNORF25 receptor contained in plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495).
30
31. A nucleic acid probe comprising at least 15
35

5 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF25 receptor, wherein the probe has a sequence complementary to a unique sequence present within one of the two strands of the nucleic acid encoding the mammalian SNORF25 receptor contained in plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494).

10 32. A nucleic acid probe comprising at least 15 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF25 receptor, wherein the probe has a sequence complementary to a unique sequence present within (a) the nucleic acid sequence shown in Figures 1A-1B (SEQ ID NO: 1) or (b) 15 the reverse complement thereof.

20 33. A nucleic acid probe comprising at least 15 nucleotides, which probe specifically hybridizes with a nucleic acid encoding a mammalian SNORF25 receptor, wherein the probe has a sequence complementary to a unique sequence present within (a) the nucleic acid sequence shown in Figures 3A-3B (SEQ ID NO: 3) or (b) 25 the reverse complement thereof.

34. The nucleic acid probe of claim 32 or 33, wherein the nucleic acid is DNA.

35. The nucleic acid probe of claim 32 or 33, wherein the nucleic acid is RNA.

30 36. An antisense oligonucleotide having a sequence capable of specifically hybridizing to the RNA of claim 5, so as to prevent translation of the RNA.

35 37. An antisense oligonucleotide having a sequence capable of specifically hybridizing to the genomic DNA of claim 4, so as to prevent transcription of the genomic DNA.

38. An antisense oligonucleotide of claim 36 or 37, wherein the oligonucleotide comprises chemically modified nucleotides or nucleotide analogues.
- 5 39. An antibody capable of binding to a mammalian SNORF25 receptor encoded by the nucleic acid of claim 1.
- 10 40. An antibody of claim 39, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor or a rat SNORF25 receptor.
- 15 41. An agent capable of competitively inhibiting the binding of the antibody of claim 39 to a mammalian SNORF25 receptor.
42. An antibody of claim 39, wherein the antibody is a monoclonal antibody or antisera.
- 20 43. A pharmaceutical composition comprising (a) an amount of the oligonucleotide of claim 36 capable of passing through a cell membrane and effective to reduce expression of a mammalian SNORF25 receptor and (b) a pharmaceutically acceptable carrier capable of passing through the cell membrane.
- 25 44. A pharmaceutical composition of claim 43, wherein the oligonucleotide is coupled to a substance which inactivates mRNA.
- 30 45. A pharmaceutical composition of claim 44, wherein the substance which inactivates mRNA is a ribozyme.
46. A pharmaceutical composition of claim 44, wherein the pharmaceutically acceptable carrier comprises a structure which binds to a mammalian SNORF25 receptor on a cell capable of being taken up by the cells after

binding to the structure.

47. A pharmaceutical composition of claim 46, wherein the pharmaceutically acceptable carrier is capable of binding to a mammalian SNORF25 receptor which is specific for a selected cell type.
- 5
48. A pharmaceutical composition which comprises an amount of the antibody of claim 39 effective to block binding of a ligand to a human SNORF25 receptor and a pharmaceutically acceptable carrier.
- 10
49. A transgenic, nonhuman mammal expressing DNA encoding a mammalian SNORF25 receptor of claim 1.
- 15
50. A transgenic, nonhuman mammal comprising a homologous recombination knockout of the native mammalian SNORF25 receptor.
- 20
51. A transgenic, nonhuman mammal whose genome comprises antisense DNA complementary to the DNA encoding a mammalian SNORF25 receptor of claim 1 so placed within the genome as to be transcribed into antisense mRNA which is complementary to mRNA encoding the mammalian SNORF25 receptor and which hybridizes with mRNA encoding the mammalian SNORF25 receptor so as to thereby reduce translation of the mRNA and expression of the receptor.
- 25
- 30
52. The transgenic, nonhuman mammal of claim 49 or 50, wherein the DNA encoding the mammalian SNORF25 receptor additionally comprises an inducible promoter.
- 35
53. The transgenic, nonhuman mammal of claim 49 or 50, wherein the DNA encoding the mammalian SNORF25 receptor additionally comprises tissue specific regulatory elements.

54. A transgenic, nonhuman mammal of claim 49, 50, or 51, wherein the transgenic, nonhuman mammal is a mouse.

55. A process for identifying a chemical compound which specifically binds to a mammalian SNORF25 receptor which comprises contacting cells containing DNA encoding and expressing on their cell surface the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, with the compound under conditions suitable for binding, and detecting specific binding of the chemical compound to the mammalian SNORF25 receptor.

56. A process for identifying a chemical compound which specifically binds to a mammalian SNORF25 receptor which comprises contacting a membrane preparation from cells containing DNA encoding and expressing on their cell surface the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, with the compound under conditions suitable for binding, and detecting specific binding of the chemical compound to the mammalian SNORF25 receptor.

57. The process of claim 55 or 56, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor.

58. The process of claim 55 or 56, wherein the mammalian SNORF25 receptor has substantially the same amino acid sequence as the human SNORF25 receptor encoded by plasmid pEXJT3T7-hSNORF25 (ATCC Accession No. 203495).

59. The process of claim 55 or 56, wherein the mammalian SNORF25 receptor has substantially the same amino acid sequence as that shown in Figures 2A-2B (SEQ ID NO: 2).

60. The process of claim 55 or 56, wherein the mammalian SNORF25 receptor has the amino acid sequence shown in Figures 2A-2B (SEQ ID NO: 2).
- 5 61. The process of claim 55 or 56, wherein the mammalian SNORF25 receptor is a rat SNORF25 receptor.
- 10 62. The process of claim 55 or 56, wherein the mammalian SNORF25 receptor has substantially the same amino acid sequence as the human SNORF25 receptor encoded by plasmid pcDNA3.1-rSNORF25 (ATCC Accession No. 203494).
- 15 63. The process of claim 55 or 56, wherein the mammalian SNORF25 receptor has substantially the same amino acid sequence as that shown in Figures 4A-4B (SEQ ID NO: 4).
- 20 64. The process of claim 55 or 56, wherein the mammalian SNORF25 receptor has the amino acid sequence shown in Figures 4A-4B (SEQ ID NO: 4).
- 25 65. The process of claim 55 or 56, wherein the compound is not previously known to bind to a mammalian SNORF25 receptor.
66. A compound identified by the process of claim 65.
- 30 67. A process of claim 55 or 56, wherein the cell is an insect cell.
68. The process of claim 55 or 56, wherein the cell is a mammalian cell.
- 35 69. The process of claim 68, wherein the cell is nonneuronal in origin.
70. The process of claim 69, wherein the nonneuronal cell is a COS-7 cell, 293 human embryonic kidney cell, a

CHO cell, a NIH-3T3 cell, a mouse Y1 cell, or a LM(tk-) cell.

71. A process of claim 68, wherein the compound is a
5 compound not previously known to bind to a mammalian
SNORF25 receptor.

72. A compound identified by the process of claim 71.

10 73. A process involving competitive binding for
identifying a chemical compound which specifically
binds to a mammalian SNORF25 receptor which comprises
separately contacting cells expressing on their cell
surface the mammalian SNORF25 receptor, wherein such
15 cells do not normally express the mammalian SNORF25
receptor, with both the chemical compound and a second
chemical compound known to bind to the receptor, and
with only the second chemical compound, under
conditions suitable for binding of such compounds to
20 the receptor, and detecting specific binding of the
chemical compound to the mammalian SNORF25 receptor,
a decrease in the binding of the second chemical
compound to the mammalian SNORF25 receptor in the
presence of the chemical compound being tested
25 indicating that such chemical compound binds to the
mammalian SNORF25 receptor.

74. A process involving competitive binding for
identifying a chemical compound which specifically
30 binds to a mammalian SNORF25 receptor which comprises
separately contacting a membrane preparation from
cells expressing on their cell surface the mammalian
SNORF25 receptor, wherein such cells do not normally
35 express the mammalian SNORF25 receptor, with both the
chemical compound and a second chemical compound known
to bind to the receptor, and with only the second

chemical compound, under conditions suitable for binding of such compounds to the receptor, and detecting specific binding of the chemical compound to the mammalian SNORF25 receptor, a decrease in the 5 binding of the second chemical compound to the mammalian SNORF25 receptor in the presence of the chemical compound being tested indicating that such chemical compound binds to the mammalian SNORF25 receptor.

10

75. A process of claim 73 or 74, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor.

15

76. A process of claim 73 or 74, wherein the mammalian SNORF25 receptor is a rat SNORF25 receptor.

77. The process of claim 73 or 74, wherein the cell is an insect cell.

20

78. The process of claim 73 or 74, wherein the cell is a mammalian cell.

79. The process of claim 77, wherein the cell is nonneuronal in origin.

25

80. The process of claim 79, wherein the nonneuronal cell is a COS-7 cell, 293 human embryonic kidney cell, a CHO cell, a NIH-3T3 cell, a mouse Y1 cell, or a LM(tk-) cell.

30

81. The process of claim 80, wherein the compound is not previously known to bind to a mammalian SNORF25 receptor.

35

82. A compound identified by the process of claim 81.

83. A method of screening a plurality of chemical compounds not known to bind to a mammalian SNORF25

receptor to identify a compound which specifically binds to the mammalian SNORF25 receptor, which comprises

5 (a) contacting cells transfected with and expressing DNA encoding the mammalian SNORF25 receptor with a compound known to bind specifically to the mammalian SNORF25 receptor;

10 (b) contacting the cells of step (a) with the plurality of compounds not known to bind specifically to the mammalian SNORF25 receptor, under conditions permitting binding of compounds known to bind to the mammalian SNORF25 receptor;

15 (c) determining whether the binding of the compound known to bind to the mammalian SNORF25 receptor is reduced in the presence of the plurality of compounds, relative to the binding of the compound in the absence of the plurality of compounds; and if so

20 (d) separately determining the binding to the mammalian SNORF25 receptor of each compound included in the plurality of compounds, so as to thereby identify any compound included therein which specifically binds to the mammalian SNORF25 receptor.

25 84. A method of screening a plurality of chemical compounds not known to bind to a mammalian SNORF25 receptor to identify a compound which specifically binds to the mammalian SNORF25 receptor, which comprises

30 35 (a) contacting a membrane preparation from cells

transfected with, and expressing, DNA encoding the mammalian SNORF25 receptor with the plurality of compounds not known to bind specifically to the mammalian SNORF25 receptor under conditions permitting binding of compounds known to bind to the mammalian SNORF25 receptor;

10 (b) determining whether the binding of a compound known to bind to the mammalian SNORF25 receptor is reduced in the presence of the plurality of compounds, relative to the binding of the compound in the absence of the plurality of compounds; and if so

15 (c) separately determining the binding to the mammalian SNORF25 receptor of each compound included in the plurality of compounds, so as to thereby identify any compound included therein which specifically binds to the mammalian SNORF25 receptor.

20

85. A method of claim 83 or 84, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor.

25 86. A method of claim 83 or 84, wherein the mammalian
SNORF25 receptor is a rat SNORF25 receptor.

87. A method of claim 83 or 84, wherein the cell is a mammalian cell

30
88. A method of claim 87, wherein the mammalian cell is
non-neuronal in origin.

89. The method of claim 88, wherein the non-neuronal cell
35 is a COS-7 cell, a 293 human embryonic kidney cell, a
LM(tk-) cell, a CHO cell, a mouse Y1 cell, or an
NIH-3T3 cell.

90. A method of detecting expression of a mammalian SNORF25 receptor by detecting the presence of mRNA coding for the mammalian SNORF25 receptor which comprises obtaining total mRNA from the cell and 5 contacting the mRNA so obtained with the nucleic acid probe of claim 30, 31, 32 or 33 under hybridizing conditions, detecting the presence of mRNA hybridized to the probe, and thereby detecting the expression of the mammalian SNORF25 receptor by the cell.

10 91. A method of detecting the presence of a mammalian SNORF25 receptor on the surface of a cell which comprises contacting the cell with the antibody of 15 claim 39 under conditions permitting binding of the antibody to the receptor, detecting the presence of the antibody bound to the cell, and thereby detecting the presence of the mammalian SNORF25 receptor on the surface of the cell.

20 92. A method of determining the physiological effects of varying levels of activity of mammalian SNORF25 receptors which comprises producing a transgenic, nonhuman mammal of claim 49 whose levels of mammalian SNORF25 receptor activity are varied by use of an 25 inducible promoter which regulates mammalian SNORF25 receptor expression.

30 93. A method of determining the physiological effects of varying levels of activity of mammalian SNORF25 receptors which comprises producing a panel of transgenic, nonhuman mammals of claim 49 each expressing a different amount of mammalian SNORF25 receptor.

35 94. A method for identifying an antagonist capable of alleviating an abnormality wherein the abnormality is

alleviated by decreasing the activity of a mammalian SNORF25 receptor comprising administering a compound to the transgenic, nonhuman mammal of claim 49, 50, or 51, and determining whether the compound alleviates any physiological and/or behavioral abnormality displayed by the transgenic, nonhuman mammal as a result of overactivity of a mammalian SNORF25 receptor, the alleviation of such an abnormality identifying the compound as an antagonist.

10 95. The method of claim 94, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor or a rat SNORF25 receptor.

15 96. An antagonist identified by the method of claim 94.

97. A composition comprising an antagonist of claim 96 and a carrier.

20 98. A method of treating an abnormality in a subject wherein the abnormality is alleviated by decreasing the activity of a mammalian SNORF25 receptor which comprises administering to the subject an effective amount of the pharmaceutical composition of claim 97 so as to thereby treat the abnormality.

25 99. A method for identifying an agonist capable of alleviating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian SNORF25 receptor comprising administering a compound to the transgenic, nonhuman mammal of claim 49, 50, or 51, and determining whether the compound alleviates the physiological and/or behavioral abnormalities displayed by the transgenic, nonhuman mammal, the alleviation of the abnormality identifying the compound as an agonist.

30 35 100. The method of claim 99, wherein the mammalian SNORF25

receptor is a human SNORF25 receptor or a rat SNORF25 receptor.

101. An agonist identified by the method of claim 99.

5

102. A composition comprising an agonist identified by the method of claim 101 and a carrier.

103. A method of treating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian SNORF25 receptor which comprises administering to the subject an effective amount of the composition of claim 102, thereby treating the abnormality.

15

104. A method for diagnosing a predisposition to a disorder associated with the activity of a specific mammalian allele which comprises:

20 (a) obtaining DNA of subjects suffering from the disorder;

25 (b) performing a restriction digest of the DNA with a panel of restriction enzymes;

(c) electrophoretically separating the resulting DNA fragments on a sizing gel;

30 (d) contacting the resulting gel with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a mammalian SNORF25 receptor and labeled with a detectable marker;

35

(e) detecting labeled bands which have hybridized to

the DNA encoding a mammalian SNORF25 receptor of claim 1 to create a unique band pattern specific to the DNA of subjects suffering from the disorder;

5

(f) repeating steps (a)-(e) with DNA obtained for diagnosis from subjects not yet suffering from the disorder; and

10

(g) comparing the unique band pattern specific to the DNA of subjects suffering from the disorder from step (e) with the band pattern from step (f) for subjects not yet suffering from the disorder so as to determine whether the patterns are the same or different and thereby diagnose predisposition to the disorder if the patterns are the same.

105. The method of claim 104, wherein a disorder associated with the activity of a specific mammalian allele is diagnosed.

106. A method of preparing the purified mammalian SNORF25 receptor of claim 12 which comprises:

25 (a) culturing cells which express the mammalian SNORF25 receptor;

(b) recovering the mammalian SNORF25 receptor from the cells; and

30 (c) purifying the mammalian SNORF25 receptor so recovered.

107. A method of preparing the purified mammalian SNORF25 receptor of claim 12 which comprises:

(a) inserting a nucleic acid encoding the mammalian SNORF25 receptor into a suitable expression

vector;

5 (b) introducing the resulting vector into a suitable host cell;

10 (c) placing the resulting host cell in suitable conditions permitting the production of the mammalian SNORF25 receptor;

15 (d) recovering the mammalian SNORF25 receptor so produced; and optionally

(e) isolating and/or purifying the mammalian SNORF25 receptor so recovered.

15 108. A process for determining whether a chemical compound is a mammalian SNORF25 receptor agonist which comprises contacting cells transfected with and expressing DNA encoding the mammalian SNORF25 receptor with the compound under conditions permitting the activation of the mammalian SNORF25 receptor, and detecting any increase in mammalian SNORF25 receptor activity, so as to thereby determine whether the compound is a mammalian SNORF25 receptor agonist.

25 109. A process for determining whether a chemical compound is a mammalian SNORF25 receptor antagonist which comprises contacting cells transfected with and expressing DNA encoding the mammalian SNORF25 receptor with the compound in the presence of a known mammalian SNORF25 receptor agonist, under conditions permitting the activation of the mammalian SNORF25 receptor, and detecting any decrease in mammalian SNORF25 receptor activity, so as to thereby determine whether the compound is a mammalian SNORF25 receptor antagonist.

110. A process of claim 108 or 109, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor or a rat SNORF25 receptor.
- 5 111. A composition which comprises an amount of a mammalian SNORF25 receptor agonist determined by the process of claim 108 effective to increase activity of a mammalian SNORF25 receptor and a carrier.
- 10 112. A composition of claim 111, wherein the mammalian SNORF25 receptor agonist is not previously known.
113. A composition which comprises an amount of a mammalian SNORF25 receptor antagonist determined by the process of claim 109 effective to reduce activity of a mammalian SNORF25 receptor and a carrier.
- 15 114. A composition of claim 113, wherein the mammalian SNORF25 receptor antagonist is not previously known.
- 20 115. A process for determining whether a chemical compound specifically binds to and activates a mammalian SNORF25 receptor, which comprises contacting cells producing a second messenger response and expressing on their cell surface the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, with the chemical compound under conditions suitable for activation of the mammalian SNORF25 receptor, and measuring the second messenger response in the presence and in the absence of the chemical compound, a change in the second messenger response in the presence of the chemical compound indicating that the compound activates the mammalian SNORF25 receptor.
- 25 30 35 116. The process of claim 115, wherein the second messenger response comprises chloride channel activation and the change in second messenger is an increase in the level

of chloride current.

117. The process of claim 115, wherein, the second messenger response comprises change in intracellular calcium levels and the change in second messenger is an increase in the measure of intracellular calcium.
5
118. The process of claim 115, wherein the second messenger response comprises release of inositol phosphate and the change in second messenger is an increase in the level of inositol phosphate.
10
119. A process for determining whether a chemical compound specifically binds to and inhibits activation of a mammalian SNORF25 receptor, which comprises separately contacting cells producing a second messenger response and expressing on their cell surface the mammalian SNORF25 receptor, wherein such cells do not normally express the mammalian SNORF25 receptor, with both the chemical compound and a second chemical compound known to activate the mammalian SNORF25 receptor, and with only the second chemical compound, under conditions suitable for activation of the mammalian SNORF25 receptor, and measuring the second messenger response in the presence of only the second chemical compound and in the presence of both the second chemical compound and the chemical compound, a smaller change in the second messenger response in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound indicating that the chemical compound inhibits activation of the mammalian SNORF25 receptor.
15
20
25
30
- 35 120. The process of claim 119, wherein the second messenger response comprises chloride channel activation and the

change in second messenger response is a smaller increase in the level of chloride current in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound.

- 5 121. The process of claim 119, wherein the second messenger response comprises change in intracellular calcium levels and the change in second messenger response is a smaller increase in the measure of intracellular calcium in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound.
- 10 122. The process of claim 119, wherein the second messenger response comprises release of inositol phosphate and the change in second messenger response is a smaller increase in the level of inositol phosphate in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound.
- 15 123. A process of any of claims 115, 116, 117, 118, 119, 120, 121, or 122, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor or a rat SNORF25 receptor.
- 20 124. The process of any of claims 115, 116, 117, 118, 119, 120, 121, or 122, wherein the cell is an insect cell.
- 25 125. The process of any of claims 115, 116, 117, 118, 119, 120, 121, or 122, wherein the cell is a mammalian cell.
- 30 126. The process of claim 125, wherein the mammalian cell is nonneuronal in origin.
- 35 127. The process of claim 126, wherein the nonneuronal cell

is a COS-7 cell, CHO cell, 293 human embryonic kidney cell, NIH-3T3 cell or LM(tk-) cell.

128. The process of claim 115, 116, 117, 118, 119, 120,
5 121, or 122, wherein the compound is not previously known to bind to a mammalian SNORF25 receptor.

129. A compound determined by the process of claim 128.

10 130. A composition which comprises an amount of a mammalian SNORF25 receptor agonist determined to be such by the process of claim 115, 116, 117, or 118, effective to increase activity of a mammalian SNORF25 receptor and a carrier.

15

131. A composition of claim 130, wherein the mammalian SNORF25 receptor agonist is not previously known.

20

132. A composition which comprises an amount of a mammalian SNORF25 receptor antagonist determined to be such by the process of claim 119, 120, 121, or 122, effective to reduce activity of a mammalian SNORF25 receptor and a carrier.

25

133. A composition of claim 132, wherein the mammalian SNORF25 receptor antagonist is not previously known.

30

134. A method of screening a plurality of chemical compounds not known to activate a mammalian SNORF25 receptor to identify a compound which activates the mammalian SNORF25 receptor which comprises:

35

(a) contacting cells transfected with and expressing the mammalian SNORF25 receptor with the plurality of compounds not known to activate the mammalian SNORF25 receptor, under conditions permitting

activation of the mammalian SNORF25 receptor;

5 (b) determining whether the activity of the mammalian SNORF25 receptor is increased in the presence of one or more of the compounds; and if so

10 (c) separately determining whether the activation of the mammalian SNORF25 receptor is increased by any compound included in the plurality of compounds, so as to thereby identify each compound which activates the mammalian SNORF25 receptor.

135. A method of claim 134, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor or a rat SNORF25 receptor.

20 136. A method of screening a plurality of chemical compounds not known to inhibit the activation of a mammalian SNORF25 receptor to identify a compound which inhibits the activation of the mammalian SNORF25 receptor, which comprises:

25 (a) contacting cells transfected with and expressing the mammalian SNORF25 receptor with the plurality of compounds in the presence of a known mammalian SNORF25 receptor agonist, under conditions permitting activation of the mammalian SNORF25 receptor;

30 (b) determining whether the extent or amount of activation of the mammalian SNORF25 receptor is reduced in the presence of one or more of the compounds, relative to the extent or amount of activation of the mammalian SNORF25 receptor in the absence of such one or more compounds; and if so

5 (c) separately determining whether each such compound inhibits activation of the mammalian SNORF25 receptor for each compound included in the plurality of compounds, so as to thereby identify any compound included in such plurality of compounds which inhibits the activation of the mammalian SNORF25 receptor.

10 137. A method of claim 136, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor.

138. A method of any of claims 134, 135, 136, 137, wherein the cell is a mammalian cell.

15 139. A method of claim 138, wherein the mammalian cell is non-neuronal in origin.

20 140. The method of claim 139, wherein the non-neuronal cell is a COS-7 cell, a 293 human embryonic kidney cell, a LM(tk-) cell or an NIH-3T3 cell.

25 141. A composition comprising a compound identified by the method of claim 134 or 135 in an amount effective to increase mammalian SNORF25 receptor activity and a carrier.

30 142. A composition comprising a compound identified by the method of claim 136 or 137 effective to decrease mammalian SNORF25 receptor activity and a carrier.

35 143. A method of treating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian SNORF25 receptor which comprises administering to the subject a compound which is a mammalian SNORF25 receptor agonist in an amount effective to treat the abnormality.

144. A method of treating an abnormality in a subject wherein the abnormality is alleviated by decreasing the activity of a mammalian SNORF25 receptor which comprises administering to the subject a compound which is a mammalian SNORF25 receptor antagonist in an amount effective to treat the abnormality.

5

145. A process for making a composition of matter which specifically binds to a mammalian SNORF25 receptor which comprises identifying a chemical compound using the process of any of claims 55, 56, 73, 74, 83 or 84 and then synthesizing the chemical compound or a novel structural and functional analog or homolog thereof.

10

146. The process of claims 145, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor or a rat SNORF25 receptor.

15

147. A process for making a composition of matter which specifically binds to a mammalian SNORF25 receptor which comprises identifying a chemical compound using the process of any of claims 108, 115, or 134 and then synthesizing the chemical compound or a novel structural and functional analog or homolog thereof.

20

25

148. The process of claim 147, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor or a rat SNORF25 receptor.

30

149. A process for making a composition of matter which specifically binds to a mammalian SNORF25 receptor which comprises identifying a chemical compound using the process of any of claims 109, 119 or 156 and then synthesizing the chemical compound or a novel structural and functional analog or homolog thereof.

35

150. The process of claim 149, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor or a rat

SNORF25 receptor.

151. A process for preparing a composition which comprises admixing a carrier and a pharmaceutically effective amount of a chemical compound identified by the process of any of claims 55, 56, 73, 74, 83 or 84 or a novel structural and functional analog or homolog thereof.
5
- 10 152. The process of claim 151, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor or a rat SNORF25 receptor.
- 15 153. A process for preparing a composition which comprises admixing a carrier and a pharmaceutically effective amount of a chemical compound identified by the process of any of claims 108, 115, or 134 or a novel structural and functional analog or homolog thereof.
- 20 154. The process of claim 153, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor or a rat SNORF25 receptor.
155. A process for preparing a composition which comprises admixing a carrier and a pharmaceutically effective amount of a chemical compound identified by the process of any of claims 109, 119 or 136 or a novel structural and functional analog or homolog thereof.
25
- 30 156. The process of claim 155, wherein the mammalian SNORF25 receptor is a human SNORF25 receptor or a rat SNORF25 receptor.

1/14

FIGURE 1A

1 TGAGAATTTCAGCTGGAGAGATTAGCATGCCCTGGTAAGTGAAGTCCTGCCACTTCGAGAC 60
61 ATGGAATCATCTTCTCATTGGAGTGATCCTGGCTGCTGGCTGGCTGCTGCTCATTCATTGCT 120
121 ACTAACACACTAGTGGCTGCTGCTGATCCACAAAGAACATGATGGTGTCACT 180
181 CTCTGCTTCACCTTGAAATCTGGCTGACACCTTGATTGGTGTGGCCATCTCTGGC 240
241 CTACTCACAGACCAGCTCTCCAGCCCTTCGGCCCCACACAGAACACCCCTGTGCAGGCCTG 300
301 CGGATGGCATTGTCACTTCCCTGCAGCTGCCTCAGGTCACTGCTGATCACC 360
361 TTTGACAGGTACCTTGCATCAAGCAGCCCTCCGGCTACTTGAAGATCATGAGTGGGTTTC 420
421 GTGGCCGGGGCCATTGCATTGGCTGGGGCTGTGGTTAGTGTCTTACCTCATGGCTTCCTCCCCA 480
481 CTCGGAATCCCCATGTTCCAGCAGACTGCCTACAAAGGGCAGTCAGCTTGTGCTGTA 540
541 TTTCACCCCTCACTTCGTTGCTGACCCCTCTGGCTTCTCCAGCCATGCTCCTC 600

FIGURE 1B

601 TTTGTCTTCTACTGGCACATGCTCAAAGATGCCTCCATGCCAGCAGATTCGA 660
661 AAGATGGAACATGCAGGAGCCATGGCTGGAGGTTATCGATCCCCACGGACTCCAGCGAC 720
721 TTCAAAAGCTCTCCGTACTGTGTCTCATGGGAGCTTGTCTATCCTGGACCCCC 780
781 TTCCTTATCACTGGCATTGGCAGGTGCCCTGCCAGGAGTGTCAACCCACTCATCTATGCC 840
841 GAAACGGTACCTGTGGCTGCTCGGGCAACTCCCTGCTCAACCCACTCATCTATGCC 900
901 TATTGGCAGAAGGAGGTGGCACTGCAGCTCTACCAACATGGCCCTAGGAGTGAAGAAAGGTG 960
961 CTCACCTCATTCCTCTTCTCTGGCCAGGAATTGTGGCCAGAGGCCAGGGAA 1020
1021 AGTTCCCTGTCAACATCGTCACTATCTCCAGCTCAAGTTGATGGCTAAGACGGTAAGGGC 1080
1081 AGAGAAGTTCAAAGTGCCTTCTCCACTCTGGAGCCCCAACTAG 1129

3/14

FIGURE 2A

1	M	E	S	S	F	S	F	G	V	I	L	A	V	L	A	S	L	I	I	A	20
21	T	N	T	L	V	A	V	L	L	I	H	K	N	D	G	V	S	S	40		
41	L	C	F	T	L	N	L	A	V	A	D	T	L	I	G	V	A	I	S	G	60
61	L	L	T	D	Q	L	S	S	P	S	R	P	T	Q	K	T	L	C	S	L	80
81	R	M	A	F	V	T	S	S	A	A	A	S	V	L	T	V	M	L	I	T	100
101	E	D	R	Y	L	A	I	K	Q	P	F	R	Y	L	K	I	M	S	G	F	120
121	V	A	G	A	C	I	A	G	L	W	L	V	S	Y	L	I	G	F	L	P	140
141	L	G	I	P	M	F	Q	Q	T	A	Y	K	G	Q	C	S	F	F	A	V	160
161	F	H	P	H	F	V	L	T	L	S	C	V	G	F	F	P	A	M	L	L	180
181	F	V	F	F	Y	C	D	M	L	K	I	A	S	M	H	S	Q	Q	I	R	200
201	K	M	E	H	A	G	A	M	A	G	G	Y	R	S	P	R	T	P	S	D	220

4/14

FIGURE 2B

221	F	K	A	L	R	T	V	S	V	L	I	G	S	F	A	L	S	W	T	P	240
241	<u>F</u>	<u>L</u>	<u>I</u>	<u>T</u>	<u>G</u>	<u>I</u>	<u>V</u>	<u>Q</u>	<u>V</u>	<u>A</u>	<u>C</u>	<u>Q</u>	<u>E</u>	<u>C</u>	<u>H</u>	<u>L</u>	<u>Y</u>	<u>L</u>	<u>V</u>	<u>L</u>	260
261	E	R	<u>Y</u>	<u>L</u>	<u>W</u>	<u>L</u>	<u>L</u>	<u>G</u>	<u>V</u>	<u>G</u>	<u>N</u>	<u>S</u>	<u>L</u>	<u>L</u>	<u>N</u>	<u>P</u>	<u>L</u>	<u>I</u>	<u>Y</u>	<u>A</u>	280
281	<u>Y</u>	<u>W</u>	<u>Q</u>	<u>K</u>	<u>E</u>	<u>V</u>	<u>R</u>	<u>L</u>	<u>Q</u>	<u>L</u>	<u>Y</u>	<u>H</u>	<u>M</u>	<u>A</u>	<u>L</u>	<u>G</u>	<u>V</u>	<u>K</u>	<u>K</u>	<u>V</u>	300
301	L	T	S	F	L	L	F	L	S	A	R	N	C	G	P	E	R	P	R	E	320
321	S	S	C	H	I	V	T	I	S	S	E	F	D	G						335	

5/14

FIGURE 3A

1	TCAAGACCCAGCATTGCCCTTATAAGTGGGAGTCCCTGCTAACCTCGAACCATGGAGTCATCT	60
61	TTCTCATTTGGAGTGAATCCTTGCTGCTTATCAATTCCCTGACCATTATGCTTAATGGCGCTG	120
121	GTGGTTGGCTATGCTGCTTATCAAGAATGATGGTGGCTTGGCCTTGGCTTACACC	180
181	TTAAATCTGGCGTGGCTGATACCTTGATTGGCTGGCTATTCTGGCTAGTTACAGAC	240
241	CAGCTCTCCAGCTCTGCTCAGCACACAGAACACAGAACAGAC	300
301	GTCACTTCTCTGCAGCCGCCCTCTGTCCTCACGGTCATGCTGATTGCCCTTGACAGGTAC	360
361	CTGGCCATTAAAGCAGCCCCCTCCGTTACTTCCAGATCATGAATGGGCTTGTAGCCGGAGGA	420
421	TGCATTGCAGGGCTGGTGTGATATCTTACCTTACGGCTTCCACTTGGAGTCCTCC	480
481	ATATTCCAGCAGACCACCTACCATGGGCCCTGCACCTTCTTGTGTTACCCAAGG	540
541	TTTGTGCTGACCCCTCCTGTGGCTGGCTTCTTGTGCTTGTCTTGTCTTC	600
601	TACTGTGACATGCTCAAGATGGCTCAGGCCACATCCGGAAAGATGGAAACAT	660

6/14

FIGURE 3B

661	GCAGGAGCCATGGTTGGAGCCTTGGCCGGCCCCACGGCCCTGTCAATGACTTCAGGCTGTC	720
721	CGGACTGTATCTGTCCATTGGAGCTTCACCCCTGGTCTCCGTTCTCATCACT	780
781	AGCATTGTGCAGGTGGCCTGCCACAAATGCTGCCACTACCAAGTGCTGGAAAATACCTC	840
841	TGGCTCCTTGGCAACTCCCTGCTCAACCCACTCATCTATGCCTATTGGCAGAGG	900
901	GAGGTTGGCAGCAGCTCTGCCACATGCCCTGGGGTGAAGAAAGTTCTTACTTCAATC	960
961	TTCCCTCCTCTCGGCCAGGAATCGTGGTCCACAGAGGACCCGAGAAAGCTCCTATCAC	1020
1021	ATCGTCACTATCAGCCAGCGAGCTCGATGGCTAGGATGGTAAGGAATGGATGTTCCA	1080
1081	AG	1082

FIGURE 4A

1 M E S S F S F G V I L A V L T I L I I A 20

21 V N A L V V V A M L L S I Y K N D G V G 40

41 L C F T L N L A V A D T L I G V A I S G 60

61 L V T D Q L S S S A Q H T Q K T L C S L 80

81 R M A F V T S S A A A S V L T V M L I A 100

101 E D R Y L A I K Q P L R Y F Q I M N G L 120

121 V A G G C I A G L W L I S Y L I G F L P 140

141 L G V S I F Q Q T T Y H G P C T F F A V 160

161 F H P R F V L T L S C A G F F P A V L L 180

181 F V F F Y C D M L K I A S V H S Q H I R 200

201 K M E H A G A M V G A C R P P R P V N D 220

7/14


8/14

FIGURE 4B

221	F	K	A	V	R	T	V	S	V	L	I	G	S	E	T	L	S	W	S	E	240	
241	F	I	I	T	T	S	I	V	O	V	A	C	H	K	C	C	L	Y	Q	V	L	260
261	E	K	Y	L	W	L	L	G	V	G	N	S	L	L	N	P	L	I	Y	A	280	
281	Y	W	Q	R	E	V	R	Q	Q	L	C	H	M	A	L	G	V	K	K	F	300	
301	F	T	S	I	F	L	L	S	A	R	N	R	G	P	Q	R	T	R	E	320		
321	S	S	Y	H	I	V	T	I	S	Q	P	E	L	D	G					335		

9/14


FIGURE 5



10/14

FIGURE 6


**Specificity of ATRA on cAMP
release in
SNORF25-transfected CHO cells**



11/14

FIGURE 7


Effect of ATRA on cAMP release in Cos-7 cells transfected with various cyclase stimulatory receptors



12/14

FIGURE 8

ATRA dose-response curve for cAMP release in SNORF25 and mock-transfected Cos-7 cells



13/14

SNORF25 + CFTR

Figure 9A



CFTR alone

Figure 9B



B2AR + CFTR


Figure 9C



14/14

FIGURE 10

**Mean Peak Response to ATRA in 17
Responding Oocytes**



SEQUENCE LISTING

<110> Bonini, James A.
Borowsky, Beth E.
Adham, Nika
Boyle, Noel

<120> DNA Encoding SNORF25 Receptor

<130> 56095-A

<140>

<141>

<150> 09/255,376

<151> 1999-02-22

<160> 23

<170> PatentIn Ver. 2.0 - beta

<210> 1

<211> 1129

<212> DNA

<213> Homo sapiens

<400> 1

tgagaatttc agctggagag atagcatgcc ctggtaagtgc aagtccctgcc acttcgagac 60
atggaatcat ctttctcatt tggagtgtac cttgtgtcc tggcctccct catcattgct 120
actaacacac tagtggctgt ggctgtgtc ctgttgatcc acaagaatga tggtgtca 180
ctctgcttca ctttgaatct ggctgtggct gacaccttga ttggtgtggc catctctggc 240
ctactcacag accagctctc cagcccttct cggcccacac agaagaccct gtgcagcctg 300
cggatggcat ttgtcacttc ctccgcagct gcctctgtcc tcacggatcat gctgatcacc 360
tttgacaggt accttgccat caagcagccc ttccgctact tgaagatcat gagtgggttc 420
gtggccgggg cctgcattgc cgggctgtgg tttagtgcattt acctcattgg cttccctccca 480
ctcggaaatcc ccatgttcca gcagactgcc tacaaaggcc agtgcagctt ctttgcgtta 540
tttcacccttc acttcgtgtc gaccctctcc tgcgttgct tcttcccagc catgctcctc 600
tttgcatttct tctactgcga catgctcaag attgcctcca tgcacagcca gcagattcga 660
aagatggAAC atgcaggAGC catggctggA ggttatcgat ccccacggAC tcccagcgAC 720
ttcaaAGCTC tccgtactgt gtctgttctc attggagct ttgtctatc ctggacccccc 780
ttccattatca ctggcattgt gcagggtggcc tgccaggagt gtcacctcta cctagtgcgt 840
gaacggtaCC tggctgtgc cggcgtggc aactccctgc tcaacccact catctatgcc 900
tattggcaga aggagggtgcg actgcagctc taccacatgg ccctaggagt gaagaagggtg 960
ctcacctcat tcctccttct tctctcgcc aggaattgtg gcccagagag gcccaggaa 1020
agttcctgtc acatcgtaac tatctccagc tcagagttt atggctaaga cggtaaggc 1080
agagaagttt caaaagtgcct ttctccccc actctggagc cccaaactag 1129

<210> 2

<211> 335

<212> PRT

<213> Homo sapiens

<400> 2

Met Glu Ser Ser Phe Ser Phe Gly Val Ile Leu Ala Val Leu Ala Ser
1 5 10 15

Leu Ile Ile Ala Thr Asn Thr Leu Val Ala Val Ala Val Leu Leu Leu
20 25 30

Ile His Lys Asn Asp Gly Val Ser Leu Cys Phe Thr Leu Asn Leu Ala
35 40 45

Val Ala Asp Thr Leu Ile Gly Val Ala Ile Ser Gly Leu Leu Thr Asp
50 55 60

Gln Leu Ser Ser Pro Ser Arg Pro Thr Gln Lys Thr Leu Cys Ser Leu
65 70 75 80

Arg Met Ala Phe Val Thr Ser Ser Ala Ala Ala Ser Val Leu Thr Val
85 90 95

Met Leu Ile Thr Phe Asp Arg Tyr Leu Ala Ile Lys Gln Pro Phe Arg
100 105 110

Tyr Leu Lys Ile Met Ser Gly Phe Val Ala Gly Ala Cys Ile Ala Gly
115 120 125

Leu Trp Leu Val Ser Tyr Leu Ile Gly Phe Leu Pro Leu Gly Ile Pro
130 135 140

Met Phe Gln Gln Thr Ala Tyr Lys Gly Gln Cys Ser Phe Phe Ala Val
145 150 155 160

Phe His Pro His Phe Val Leu Thr Leu Ser Cys Val Gly Phe Phe Pro
165 170 175

Ala Met Leu Leu Phe Val Phe Phe Tyr Cys Asp Met Leu Lys Ile Ala
180 185 190

Ser Met His Ser Gln Gln Ile Arg Lys Met Glu His Ala Gly Ala Met
195 200 205

Ala Gly Gly Tyr Arg Ser Pro Arg Thr Pro Ser Asp Phe Lys Ala Leu
210 215 220

Arg Thr Val Ser Val Leu Ile Gly Ser Phe Ala Leu Ser Trp Thr Pro
225 230 235 240

Phe Leu Ile Thr Gly Ile Val Gln Val Ala Cys Gln Glu Cys His Leu
245 250 255

Tyr Leu Val Leu Glu Arg Tyr Leu Trp Leu Leu Gly Val Gly Asn Ser
 260 265 270

Leu Leu Asn Pro Leu Ile Tyr Ala Tyr Trp Gln Lys Glu Val Arg Leu
 275 280 285

Gln Leu Tyr His Met Ala Leu Gly Val Lys Lys Val Leu Thr Ser Phe
 290 295 300

Leu Leu Phe Leu Ser Ala Arg Asn Cys Gly Pro Glu Arg Pro Arg Glu
 305 310 315 320

Ser Ser Cys His Ile Val Thr Ile Ser Ser Ser Glu Phe Asp Gly
 325 330 335

<210> 3

<211> 1082

<212> DNA

<213> Rattus norvegicus

<400> 3

tcaagaccca gcatgccctt ataagtggga gtcctgctac ctcgaaccat ggagtcatct 60
 ttctcatttg gagtgtaccc ttgctgtcctg accatccta tcattgtgtt taatgcgtg 120
 gtggttgtgg ctatgtgtct atcaatctac aagaatgtat gtgttggcct ttgcttcacc 180
 ttaaatctgg ccgtggctga taccttgatt ggcgtggcta tttctggctt agttacagac 240
 cagctctcca gctctgctca gcacacacag aagaccttgtt gtagccttcg gatggcattc 300
 gtcacttctt ctgcagccgc ctctgtcctc acggcatgc tgattgcctt tgacaggtac 360
 ctggccatta agcagccccct ccgttacttc cagatcatga atggcttgtt agccggagga 420
 tgcattgcag ggctgtgggtt gatatcttac cttatcggtt tcctcccact tggagtctcc 480
 atattccagc agaccaccta ccatggggccc tgcaccttctt ttgctgtgtt tcacccaagg 540
 tttgtgtga ccctctcctt tgctggcttc ttcccagctt tgctcctctt tgtcttctt 600
 tactgtgaca tgctcaagat tgccctgtt cacagccagc acatccggaa gatggaacat 660
 gcaggagcca tgggtggagc ttgccccccc ccacggctt tcaatgactt caaggctgtc 720
 cggactgtat ctgtccttat tggagcttc accctgtcctt ggtctccgtt tctcatact 780
 agcattgtgc agtgtggctt ccacaaaatgc tgcctctacc aagtgtgtt aaaaatacctc 840
 tggctccttggc gagttggcaa ctccctgttcc aacccactca tctatgccta ttggcagagg 900
 gaggttcggc agcagctgtt ccacatggcc ctgggggttga agaagttctt tacttcaatc 960
 ttccctccttc tctcggccag gaatcgtgtt ccacagagga cccgagaaag ctcctatcac 1020
 atcgtcaacta tcagccagcc ggagctcgat ggcttaggatg gtaaggaatg gatgtttcca 1080
 ag 1082

<210> 4

<211> 335

<212> PRT

<213> Rattus norvegicus

<400> 4

Met Glu Ser Ser Phe Ser Phe Gly Val Ile Leu Ala Val Leu Thr Ile

1	5	10	15
Leu Ile Ile Ala Val Asn Ala Leu Val Val Val Ala Met	Leu Leu Ser		
20	25		30
Ile Tyr Lys Asn Asp Gly Val Gly Leu Cys Phe Thr	Leu Asn Leu Ala		
35	40		45
Val Ala Asp Thr Leu Ile Gly Val Ala Ile Ser Gly	Leu Val Thr Asp		
50	55		60
Gln Leu Ser Ser Ser Ala Gln His Thr Gln Lys Thr	Leu Cys Ser Leu		
65	70		75
Arg Met Ala Phe Val Thr Ser Ser Ala Ala Ala Ser	Val Leu Thr Val		
85	90		95
Met Leu Ile Ala Phe Asp Arg Tyr Leu Ala Ile Lys	Gln Pro Leu Arg		
100	105		110
Tyr Phe Gln Ile Met Asn Gly Leu Val Ala Gly Gly	Cys Ile Ala Gly		
115	120		125
Leu Trp Leu Ile Ser Tyr Leu Ile Gly Phe Leu Pro	Leu Gly Val Ser		
130	135		140
Ile Phe Gln Gln Thr Thr Tyr His Gly Pro Cys Thr	Phe Phe Ala Val		
145	150		160
Phe His Pro Arg Phe Val Leu Thr Leu Ser Cys Ala	Gly Phe Phe Pro		
165	170		175
Ala Val Leu Leu Phe Val Phe Phe Tyr Cys Asp Met	Leu Lys Ile Ala		
180	185		190
Ser Val His Ser Gln His Ile Arg Lys Met Glu His	Ala Gly Ala Met		
195	200		205
Val Gly Ala Cys Arg Pro Pro Arg Pro Val Asn Asp	Phe Lys Ala Val		
210	215		220
Arg Thr Val Ser Val Leu Ile Gly Ser Phe Thr	Leu Ser Trp Ser Pro		
225	230		240
Phe Leu Ile Thr Ser Ile Val Gln Val Ala Cys His	Lys Cys Cys Leu		
245	250		255
Tyr Gln Val Leu Glu Lys Tyr Leu Trp Leu Leu Gly	Val Gly Asn Ser		
260	265		270

Leu Leu Asn Pro Leu Ile Tyr Ala Tyr Trp Gln Arg Glu Val Arg Gln
275 280 285

Gln Leu Cys His Met Ala Leu Gly Val Lys Lys Phe Phe Thr Ser Ile
290 295 300

Phe Leu Leu Leu Ser Ala Arg Asn Arg Gly Pro Gln Arg Thr Arg Glu
305 310 315 320

Ser Ser Tyr His Ile Val Thr Ile Ser Gln Pro Glu Leu Asp Gly
325 330 335

<210> 5

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> n = inosine

<220>

<223> Description of Artificial Sequence: primer/probe

<400> 5

tbdsyvynga ymgntayvtk g

21

<210> 6

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> n = inosine

<220>

<223> Description of Artificial Sequence: primer/probe

<400> 6

ganrsnarng mrtanaynak nggrtt

26

<210> 7

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer/probe

<400> 7

ttatgcttcc ggctcgatg ttgtg 25

<210> 8
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 8
atgtgctgca aggcgattta agttggg 27

<210> 9
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 9
tggtctgctg gaatatggag 20

<210> 10
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 10
cttgggtgaa acacagcaaa gaagg 25

<210> 11
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 11
atggaacatg caggagccat ggttgg 26

<210> 12
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 12
aagacaaaga ggagcacagc tggg 24

<210> 13
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 13
gctcaagatt gcctctgtgc acag 24

<210> 14
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 14
atctataaggc tttaggcactt ggaaacatcc attcc 35

<210> 15
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 15
atctatggat cctgtgagaa tctgagctca agaccc 36

<210> 16
<211> 58
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 16
ttcaccttaa atctggccgt ggctgatacc ttgattggcg tggctatttc tggcttag 58

<210> 17
<211> 61
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 17
gctgtgtttc acccaagggtt tgtgctgacc ctctcctgtg ctggcttctt cccagctgtg 60
C 61

<210> 18
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 18
cctctaccta gtgctggaac gg 22

<210> 19
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 19
gctgcagtcg cacccctt 18

<210> 20
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 20
tccctgctca acccactcat ctatgcctat t 31

<210> 21
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 21
gtgttagcctt cggatggca 19

<210> 22
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 22
ggcttgcttaa tggccaggta c 21

<210> 23
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer/probe

<400> 23
tcctcacggt catgctgatt gccttt 26