



(86) Date de dépôt PCT/PCT Filing Date: 2012/07/26  
(87) Date publication PCT/PCT Publication Date: 2013/01/31  
(85) Entrée phase nationale/National Entry: 2014/01/23  
(86) N° demande PCT/PCT Application No.: CN 2012/001001  
(87) N° publication PCT/PCT Publication No.: 2013/013505  
(30) Priorité/Priority: 2011/07/26 (CN201110222016.3)

(51) Cl.Int./Int.Cl. *C07C 235/66* (2006.01),  
*A61K 31/337* (2006.01), *A61K 31/397* (2006.01),  
*A61K 31/40* (2006.01), *A61K 31/403* (2006.01),  
*A61K 31/65* (2006.01), *A61P 3/10* (2006.01),  
*A61P 31/00* (2006.01), *A61P 35/00* (2006.01),  
*C07D 205/04* (2006.01), *C07D 205/12* (2006.01),  
*C07D 209/52* (2006.01), *C07D 209/54* (2006.01),  
*C07D 221/20* (2006.01)


(71) Demandeur/Applicant:  
KBP BIOSCIENCES CO., LTD., CN

(72) Inventeurs/Inventors:  
ZHANG, HUI, CN;  
DONG, YANYAN, CN

(74) Agent: SIM & MCBURNEY

(54) Titre : COMPOSE TETRACYCLINE SUBSTITUE PAR 9-AMINOMETHYLE

(54) Title: 9-AMINOMETHYL SUBSTITUTED TETRACYCLINE COMPOUNDS




(57) Abrégé/Abstract:

The present invention relates to a 9-aminomethyl substituted tetracycline compound represented by general formula (I), pharmaceutically acceptable salts, solvates or isomers thereof, wherein  $R^{2a}$ ,  $R^{2b}$ ,  $R^3$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^5$ ,  $R^{6a}$ ,  $R^{6b}$ ,  $R^7$ ,  $R^8$ ,  $R^{9a}$ ,  $R^{9b}$ ,  $R^{10}$ ,  $R^{11}$ ,  $R^{12}$ ,  $R^{13a}$  and  $R^{13b}$  are as defined in the specification; the present invention also relates to a method for preparing the compounds; a pharmaceutical composition comprising the compounds; and use of the compounds in the preparation of medicaments for treatment and/or prevention of diseases responsive to tetracyclines.

## ABSTRACT

The present invention relates to 9-aminomethyl substituted tetracycline compounds represented by formula (I), or pharmaceutically acceptable salt, prodrug, solvate or isomer thereof, as well as a method for preparing these compounds and a pharmaceutical composition comprising the same. The present invention relates also to a use of these compounds in the preparation of a medicament for the treatment and/or prophylaxis of tetracycline drug-sensitive disease.



wherein,  $R^{2a}$ ,  $R^{2b}$ ,  $R^3$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^5$ ,  $R^{6a}$ ,  $R^{6b}$ ,  $R^7$ ,  $R^8$ ,  $R^{9a}$ ,  $R^{9b}$ ,  $R^{10}$ ,  $R^{11}$ ,  $R^{12}$ ,  $R^{13a}$  and  $R^{13b}$  are each independently as defined in the description.

FPCH12160074P

## 9-Aminomethyl Substituted Tetracycline Compounds

5 The present application claims priority to the Chinese application of No.201110222016.3, filed on July 26, 2011. The contents of this prior application are incorporated herein as a whole by reference. The contents of all the documents cited herein constitute a part of the present application.

10

### [Technical Field]

The present invention relates to 9-aminomethyl substituted tetracycline compounds, or pharmaceutically acceptable salt, prodrug, solvate or isomer thereof, as well as a method for preparing these  
15 compounds and a pharmaceutical composition comprising the same. The present invention relates also to a use of these compounds in the preparation of a medicament for the treatment and/or prophylaxis of tetracycline drug-sensitive disease.


### 20 [Background Art]

Tetracycline antibiotics are a kind of broad-spectrum antibiotics for oral use, which are generated by Actinomycete-Streptomyces fermentation. They have good pharmacological action against rickettsiales, many Gram-positive and Gram-negative bacteria, lymphogranuloma  
25 venereum pathogens, inclusion conjunctivitis pathogens and psittacosis pathogens.

The first tetracycline antibiotic is aureomycin extracted from Streptomyces aureofaciens in 1948, and then oxytetracycline, tetracycline and demeclocycline, which are all nature products have been developed  
30 successively. However, these drugs have high drug resistance and many side effects. Studies on the chemical structures of these compounds have been conducted later, and as a result, demethyl-tetracycline and


dimethylamino-tetracycline have been synthesized. But the wide use of tetracyclines brings about more and more serious drug resistance of bacterium, which renders the use reduction of tetracycline antibiotics.

In the early 1990s, a new, third-generation tetracycline named as glycyclines were developed, and the representative drug was tigecycline(GAR-936), which had broad antimicrobial spectrum. Tigecycline has similar antibacterial activity with the prior tetracyclines, and further, due to efflux mechanism and ribosomal protection mechanisms, it has antibacterial activity even to the pathogens resistant against tetracyclines. However, tigecycline can not be used orally, but only by intravenous drip for twice a day. Therefore, the use of tigecycline is inconvenient and painful for patients. There is no oral-usable tigecycline commercially available now.




Tigecycline

The PCT application of WO2004/064728 has disclosed the compounds having structures as below, and studied the antibacterial activity thereof, but found they have poor antibacterial activity.



PTK-0796 is an oral-usable tetracycline antibiotic developed by Paratek Company, and it is being at phase III clinical trials for treating complex skin and skin soft-tissue infection. This drug may be


administered via injection or oral use once a day, and has broad antimicrobial spectrum. It has strong antibacterial activity to Gram-positive and Gram-negative bacteria, anaerobic bacteria, and atypical bacteria which are sensitive or resistance to many drugs. However, there are few tetracycline antibiotics similar with PTK-0796, and as such there are many limitations in the current clinic drug use.



Therefore, there is an insistent demand to develop new tetracycline antibiotics which have broad antimicrobial spectrum and strong antibacterial activity, and may be administered for one time or orally, and may be easily synthesized and suitable for industrial production.

#### [Summary of the invention]

The present invention relates to a compound represented by the following general formula I:



or pharmaceutically acceptable salt, prodrug, solvate or isomer thereof,

wherein,  $R^{2a}$ ,  $R^{2b}$ ,  $R^3$ ,  $R^{10}$ ,  $R^{11}$  and  $R^{12}$  are each independently hydrogen;

$R^5$ ,  $R^{6a}$ ,  $R^{6b}$  and  $R^8$  are each independently hydrogen, mercapto, halogen, hydroxy, amino, carboxyl,  $C_{1-6}$ alkyl, halo $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{1-6}$ alkoxy, halo $C_{1-6}$ alkoxy,  $C_{1-6}$ alkylthio,  $C_{1-6}$ alkylamino, di( $C_{1-6}$ alkyl)amino, hydroxy $C_{1-6}$ alkyl, carboxyl $C_{1-6}$ alkyl, amino $C_{1-6}$ alkyl,  $C_{1-6}$ alkylamino $C_{1-6}$ alkyl,  $C_{1-6}$ alkylcarbonyl,  $C_{1-6}$ alkylcarbonyloxy,  $C_{1-6}$ alkoxycarbonyl,  $C_{1-6}$ alkylsulfinyl,  $C_{1-6}$ alkylsulfonyl, sulfonic, sulfonyl $C_{1-6}$ alkyl, sulfoamino, sulfoamino $C_{1-6}$ alkyl,  $C_{1-6}$ alkylsulfoamino, aminosulfonyl,  $C_{1-6}$ alkylaminosulfonyl, di( $C_{1-6}$ alkyl)aminosulfonyl, aminosulfonyl $C_{1-6}$ alkyl,  $C_{1-6}$ alkylformylamino,  $C_{1-6}$ alkylcarbamoyl, di( $C_{1-6}$ alkyl)carbamoyl, carbamoyl, carbamoyl $C_{1-6}$ alkyl, 3- to 8-membered cycloalkyl, 6- to 14-membered aryl, 6- to 14-membered aryl- $C_{1-6}$ alkyl, 6- to 14-membered aryl-formyl, 6- to 14-membered aryl-formyloxy, 3- to 8-membered heterocyclyl, 3- to 8-membered heterocyclyl- $C_{1-6}$ alkyl, 6- to 14-membered heteroaryl, or 6- to 14-membered heteroaryl- $C_{1-6}$ alkyl;

$R^7$  is  $NR^{7a}R^{7b}$ ;

$R^{4a}$ ,  $R^{4b}$ ,  $R^{7a}$  and  $R^{7b}$  are each independently hydrogen,  $C_{1-6}$ alkyl, halo $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{1-6}$ alkoxy, halo $C_{1-6}$ alkoxy,  $C_{1-6}$ alkylthio, hydroxy $C_{1-6}$ alkyl, carboxyl $C_{1-6}$ alkyl, amino $C_{1-6}$ alkyl,  $C_{1-6}$ alkylamino $C_{1-6}$ alkyl,  $C_{1-6}$ alkylcarbonyl,  $C_{1-6}$ alkoxycarbonyl,  $C_{1-6}$ alkylsulfinyl,  $C_{1-6}$ alkylsulfonyl, sulfonic, sulfonyl $C_{1-6}$ alkyl, sulfoamino $C_{1-6}$ alkyl, aminosulfonyl,  $C_{1-6}$ alkylaminosulfonyl, di( $C_{1-6}$ alkyl)aminosulfonyl, aminosulfonyl $C_{1-6}$ alkyl,  $C_{1-6}$ alkylcarbamoyl, di( $C_{1-6}$ alkyl)carbamoyl, carbamoyl, carbamoyl $C_{1-6}$ alkyl, 3- to 8-membered cycloalkyl, 6- to 14-membered aryl, 6- to 14-membered aryl- $C_{1-6}$ alkyl, 3- to 14-membered heterocyclyl or 3- to 14-membered heterocyclyl- $C_{1-6}$ alkyl;

$R^{9a}$  and  $R^{9b}$  are each independently:

(1) hydrogen, provided that  $R^{9a}$  and  $R^{9b}$  can not be hydrogen at the same time,

(2) cyclopropyl, cyclobutyl, 6- to 12-membered spirocyclic group, 6-membered endocyclic group, 8- to 12-membered endocyclic group or 6- to 10-membered saturated fused ring group, the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same

or different and selected from  $Q^1$ , and carbon(s) in said group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, CO, S, SO, SO<sub>2</sub>, N and NR<sup>c</sup>, R<sup>c</sup> represents hydrogen or C<sub>1-6</sub>alkyl;

5 (3) cyclobutylC<sub>1-6</sub>alkyl, cyclopentylC<sub>1-6</sub>alkyl, 6- to 12-membered spirocyclic group C<sub>1-6</sub>alkyl, 6- to 9-membered endocyclic group C<sub>1-6</sub>alkyl or 6- to 10-membered saturated fused ring group C<sub>1-6</sub>alkyl, the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same or different and selected from  $Q^1$ , and carbon(s) in said  
10 cyclobutyl, 6- to 12-membered spirocyclic group and 6- to 9-membered endocyclic group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, CO, S, SO, SO<sub>2</sub>, N and NR<sup>c</sup>, carbon(s) in said 6- to 10-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s) which may be the same  
15 or different and selected from CO, S, SO, SO<sub>2</sub>, N and NR<sup>c</sup>, R<sup>c</sup> represents hydrogen or C<sub>1-6</sub>alkyl;


Alternatively, R<sup>9a</sup> and R<sup>9b</sup> together with the nitrogen atom to which they are attached form azetidiny, 6- to 9-membered spirocyclic group or 6- to 9-membered saturated fused ring group, the above groups are  
20 unsubstituted or substituted by 1 to 3  $Q^2$  substituent(s) which may be the same or different, and carbon(s) in said azetidiny, 6- to 9-membered spirocyclic group and 6- to 9-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, CO, S, SO, SO<sub>2</sub>, N and NR<sup>c</sup>, R<sup>c</sup> represents hydrogen  
25 or C<sub>1-6</sub>alkyl;

$Q^1$  and  $Q^2$  are independently selected from halogen, hydroxy, amino, carboxyl, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy, haloC<sub>1-6</sub>alkyl, haloC<sub>1-6</sub>alkoxy, hydroxyC<sub>1-6</sub>alkyl, aminoC<sub>1-6</sub>alkyl, carboxylC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylamino, di(C<sub>1-6</sub>alkyl)amino, aminosulfonyl, aminosulfonylC<sub>1-6</sub>alkyl, carbamoyl,  
30 carbamoylC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, C<sub>1-6</sub>alkoxycarbonyl, phenyl, 3- to 8-membered cycloalkyl or 3- to 8-membered heterocyclyl;

$R^{13a}$  and  $R^{13b}$  are each independently hydrogen,  $C_{1-6}$ alkyl or 3- to 8-membered cycloalkyl.

Preferably, Formula (I) has a structure represented by Formula (II) as below:


5



$R^{9a}$  and  $R^{9b}$  are each independently as defined in above Formula (I).

Preferably, Formula (I) has a structure represented by Formula (III) as

10 below:



$R^{9a}$  and  $R^{9b}$  are each independently as defined in above Formula (I).

15 More preferably, in the above formulas (I) ~ (III)

$R^{9a}$  and  $R^{9b}$  are each independently:

(1) hydrogen, provided that  $R^{9a}$  and  $R^{9b}$  can not be hydrogen at the same time,

(2) cyclopropyl, cyclobutyl, 6- to 12-membered spirocyclic group, 6-membered endocyclic group or 6- to 10-membered saturated fused ring group, the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same or different and selected from  $Q^1$ , and carbon(s) in said cyclopropyl, cyclobutyl, 6- to 12-membered spirocyclic group, 6-membered endocyclic group and 6- to 10-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s)

which may be the same or different and selected from O, CO, S, N and  $\text{NR}^c$ ,  $\text{R}^c$  represents hydrogen or  $\text{C}_{1-4}$ alkyl,

(3) cyclobutyl $\text{C}_{1-4}$ alkyl, cyclopentyl $\text{C}_{1-4}$ alkyl, 6- to 12-membered spirocyclic group  $\text{C}_{1-4}$ alkyl, 6- to 9-membered endocyclic group  $\text{C}_{1-4}$ alkyl  
 5 or 6- to 10-membered saturated fused ring group- $\text{C}_{1-4}$ alkyl, the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same or different and selected from  $\text{Q}^1$ , and carbon(s) in said cyclobutyl, 6- to 12-membered spirocyclic group and 6- to 9-membered endocyclic group may be replaced by 1 to 3 atom(s) or group(s) which  
 10 may be the same or different and selected from O, CO, S, N and  $\text{NR}^c$ , carbon(s) in said 6- to 10-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from CO, S, N and  $\text{NR}^c$ ,  $\text{R}^c$  represents hydrogen or  $\text{C}_{1-4}$ alkyl;

Alternatively,  $\text{R}^{9a}$  and  $\text{R}^{9b}$  together with the nitrogen atom to which  
 15 they are attached form azetidiny, 6- to 9-membered spirocyclic group or 6- to 9-membered saturated fused ring group, the above groups are unsubstituted or substituted by 1 to 3  $\text{Q}^2$  substituent(s) which may be the same or different, and carbon(s) in said azetidiny, 6- to 9-membered spirocyclic group and 6- to 9-membered saturated fused ring group may be  
 20 replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, CO, S, N and  $\text{NR}^c$ ,  $\text{R}^c$  represents hydrogen or  $\text{C}_{1-4}$ alkyl; and

$\text{Q}^1$  and  $\text{Q}^2$  are independently selected from halogen, hydroxy, amino, carboxyl,  $\text{C}_{1-4}$ alkyl,  $\text{C}_{1-4}$ alkoxy, halo $\text{C}_{1-4}$ alkyl, halo $\text{C}_{1-4}$ alkoxy, phenyl, 3-  
 25 to 8-membered cycloalkyl or 3- to 8-membered heterocyclyl.

More preferably, in the above formulas (I) ~ (III) ,

$\text{R}^{9a}$  and  $\text{R}^{9b}$  are each independently

(1) hydrogen, provided that  $\text{R}^{9a}$  and  $\text{R}^{9b}$  can not be hydrogen at the same time,

30 (2) cyclopropyl, cyclobutyl, 6- to 10-membered spirocyclic group, 6-membered endocyclic group or 6- to 10-membered saturated fused ring group, the above groups are unsubstituted or substituted by 1 to 3

substituent(s) which may be the same or different and selected from  $Q^1$ , and carbon(s) in said cyclopropyl, cyclobutyl, 6- to 10-membered spirocyclic group, 6-membered endocyclic group and 6- to 10-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, CO, N and  $NR^c$ ,  $R^c$  represents hydrogen or  $C_{1-4}$ alkyl,

(3) cyclobutyl $C_{1-4}$ alkyl, cyclopentyl $C_{1-4}$ alkyl, 6- to 10-membered spirocyclic group  $C_{1-4}$ alkyl, 6- to 9-membered endocyclic group  $C_{1-4}$ alkyl or 6- to 10-membered saturated fused ring group- $C_{1-4}$ alkyl, the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same or different and selected from  $Q^1$ , and carbon(s) in said cyclobutyl, 6- to 10-membered spirocyclic group and 6- to 9-membered endocyclic group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, CO, N and  $NR^c$ , and carbon(s) in said 6- to 10-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from CO, N and  $NR^c$ ,  $R^c$  represents hydrogen or  $C_{1-4}$ alkyl;

Alternatively,  $R^{9a}$  and  $R^{9b}$  together with the nitrogen atom to which they are attached form azetidiny, 6- to 9-membered spirocyclic group or 6- to 9-membered saturated fused ring group, the above groups are unsubstituted or substituted by 1 to 3  $Q^2$  substituent(s) which may be the same or different, and carbon(s) in said azetidiny, 6- to 9-membered spirocyclic group and 6- to 9-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, CO, N and  $NR^c$ ,  $R^c$  represents hydrogen or  $C_{1-4}$ alkyl; and

$Q^1$  and  $Q^2$  are independently selected from halogen, hydroxy, amino, carboxyl,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy or halo $C_{1-4}$ alkyl.

Further preferably, in the above formulas (I) ~ (III) ,

$R^{9a}$  and  $R^{9b}$  are each independently:

(1) hydrogen, provided that  $R^{9a}$  and  $R^{9b}$  can not be hydrogen at the same time,

(2) cyclopropyl, cyclobutyl, 6- to 10-membered spirocyclic group, 6-membered endocyclic group or 6- to 10-membered saturated fused ring group, the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same or different and selected from  $Q^1$ ,  
 5 and carbon(s) in said cyclopropyl, cyclobutyl, 6- to 10-membered spirocyclic group, 6-membered endocyclic group and 6- to 10-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, N and  $NR^c$ ,  $R^c$  represents hydrogen or  $C_{1-4}$ alkyl;

10 (3) cyclobutyl $C_{1-3}$ alkyl, cyclopentyl $C_{1-3}$ alkyl, 6- to 10-membered spirocyclic group  $C_{1-3}$ alkyl, 6- to 9-membered endocyclic group  $C_{1-3}$ alkyl or 6- to 10-membered saturated fused ring group  $C_{1-3}$ alkyl, the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same or different and selected from  $Q^1$ , and carbon(s) in said  
 15 cyclobutyl, 6- to 10-membered spirocyclic group and 6- to 9-membered endocyclic group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, N and  $NR^c$ , carbon(s) in said 6- to 10-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected  
 20 from N and  $NR^c$ ,  $R^c$  represents hydrogen or  $C_{1-4}$ alkyl;

Alternatively,  $R^{9a}$  and  $R^{9b}$  together with the nitrogen atom to which they are attached form azetidiny, 6- to 9-membered spirocyclic group or 6- to 9-membered saturated fused ring group, the above groups are unsubstituted or substituted by 1 to 3  $Q^2$  substituent(s) which may be the  
 25 same or different, and carbon(s) in said azetidiny, 6- to 9-membered spirocyclic group and 6- to 9-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, N and  $NR^c$ ,  $R^c$  represents hydrogen or  $C_{1-4}$ alkyl; and

$Q^1$  and  $Q^2$  are independently selected from halogen,  $C_{1-4}$ alkyl,  
 30 halo $C_{1-4}$ alkyl or amino.

Further more preferably, in the above formulas (I) ~ (III) ,

$R^{9a}$  and  $R^{9b}$  are each independently:

(1) hydrogen, provided that  $R^{9a}$  and  $R^{9b}$  can not be hydrogen at the same time,

(2) cyclopropyl, cyclobutyl, 6- to 8-membered spirocyclic group or 6-membered endocyclic group, the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same or different and selected from  $Q^1$ , and carbon(s) in said cyclopropyl, cyclobutyl, 6- to 8-membered spirocyclic group and 6-membered endocyclic group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from N and  $NR^c$ ,  $R^c$  represents hydrogen, methyl or ethyl;

(3) cyclobutyl $C_{1-3}$ alkyl, cyclopentyl $C_{1-3}$ alkyl, 6- to 8-membered spirocyclic group  $C_{1-3}$ alkyl, 6- to 8-membered endocyclic group  $C_{1-3}$ alkyl, the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same or different and selected from  $Q^1$ , and carbon(s) in said cyclobutyl, 6- to 8-membered spirocyclic group and 6- to 8-membered endocyclic group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, N and  $NR^c$ ,  $R^c$  represents hydrogen, methyl or ethyl;


Alternatively,  $R^{9a}$  and  $R^{9b}$  together with the nitrogen atom to which they are attached form azetidiny, 6- to 8-membered spirocyclic group or 6- to 9-membered saturated fused ring group, the above groups are unsubstituted or substituted by 1 to 3  $Q^2$  substituent(s) which may be the same or different, and carbon(s) in said azetidiny, 6- to 8-membered spirocyclic group and 6- to 9-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, N and  $NR^c$ ,  $R^c$  represents hydrogen, methyl or ethyl; and


$Q^1$  and  $Q^2$  are independently selected from halogen,  $C_{1-4}$ alkyl or amino.

Particularly preferably, in the above formulas (I) ~ (III) ,




$R^{9a}$  and  $R^{9b}$  are each independently:

(1) hydrogen, provided that  $R^{9a}$  and  $R^{9b}$  can not be hydrogen at the same time,

(2) , the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same or different and selected from Q<sup>1</sup>;

(3) , the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same or different and selected from Q<sup>1</sup>;

Alternatively, R<sup>9a</sup> and R<sup>9b</sup> together with the nitrogen atom to which

they are attached form , , or ,

the above groups are unsubstituted or substituted by 1 to 3 Q<sup>2</sup> substituent(s) which may be the same or different;

Q<sup>1</sup> and Q<sup>2</sup> are independently selected from fluorine, chlorine, methyl or amino.

[Detailed description of the invention]

The term "C<sub>1-6</sub> alkyl" as used herein means linear or branched alkyl having 1 to 6 carbon atoms and includes "C<sub>1-4</sub> alkyl", "C<sub>1-3</sub> alkyl", "C<sub>1-2</sub> alkyl" and the like. Examples include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, n-butyl, 2-methylpropyl, 1-methylpropyl, 1,1-dimethylethyl, n-pentyl, 3-methylbutyl, 2-methylbutyl, 1-methylbutyl, 1-ethylpropyl, n-hexyl, 4-methylpentyl, 3-methylpentyl, 2-methylpentyl, 1-methylpentyl, 3,3-dimethylbutyl, 2,2-dimethylbutyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,3-dimethylbutyl, 2-ethylbutyl, 1,2-dimethylpropyl and the like.

The term "C<sub>2-6</sub> alkenyl" as used herein means linear or branched or cyclic alkenyl having 2 to 6 carbon atoms and containing double bonds. It

includes “C<sub>2-5</sub>alkenyl”, “C<sub>2-4</sub>alkenyl”, “C<sub>2-3</sub>alkenyl” and “C<sub>3-6</sub>cycloalkenyl” etc., Examples include, but are not limited to, ethenyl 1-propenyl, 2-propenyl, 1-methylethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 2-methyl-1-propenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-methyl-1-butenyl, 2-methyl-1-butenyl, 3-methyl-1-butenyl, 1-methyl-2-butenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1-methyl-3-butenyl, 2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,1-dimethyl-2-propenyl, 1,2-dimethyl-1-propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-1-propenyl, 1-ethyl-2-propenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-1-pentenyl, 2-methyl-1-pentenyl, 3-methyl-1-pentenyl, 4-methyl-1-pentenyl, 1-methyl-2-pentenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl, 4-methyl-2-pentenyl, 1-methyl-3-pentenyl, 2-methyl-3-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1-methyl-4-pentenyl, 2-methyl-4-pentenyl, 3-methyl-4-pentenyl, 4-methyl-4-pentenyl, 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl, 1,2-dimethyl-1-butenyl, 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl, 1,3-dimethyl-1-butenyl, 1,3-dimethyl-2-butenyl, 1,3-dimethyl-3-butenyl, 2,2-dimethyl-3-butenyl, 2,3-dimethyl-1-butenyl, 2,3-dimethyl-2-butenyl, 2,3-dimethyl-3-butenyl, 3,3-dimethyl-1-butenyl, 3,3-dimethyl-2-butenyl, 1-ethyl-1-butenyl, 1-ethyl-2-butenyl, 1-ethyl-3-butenyl, 2-ethyl-1-butenyl, 2-ethyl-2-butenyl, 2-ethyl-3-butenyl, 1,1,2-trimethyl-2-propenyl, 1-ethyl-1-methyl-2-propenyl, 1-ethyl-2-methyl-1-propenyl, 1-ethyl-2-methyl-2-propenyl, 1,3-butadienyl, 1,3-pentadienyl, 1,4-pentadienyl, 1,4-hexadienyl, cyclopropenyl, cyclobutenyl, cyclopentenyl, 1,3-cyclopentadienyl, cyclohexenyl, 1,4-cyclohexadienyl and the like.

The term “C<sub>2-6</sub>alkynyl” as used herein means linear or branched alkynyl having 2 to 6 carbon atoms and containing triple bonds. It includes “C<sub>2-5</sub>alkynyl”, “C<sub>2-4</sub>alkynyl” and “C<sub>2-3</sub>alkynyl” etc. . Examples include, but are not limited to, ethynyl, 2-propynyl, 2-butyne, 3-butyne, 1-methyl-2-propynyl, 2-pentyne, 3-pentyne, 4-pentyne,

1-methyl-2-butynyl, 1-methyl-3-butynyl, 2-methyl-3-butynyl,  
 1,1-dimethyl-2-propynyl, 1-ethyl-2-propynyl, 2-hexynyl, 3-hexynyl,  
 4-hexynyl, 5-hexynyl, 1-methyl-2-pentynyl, 1-methyl-3-pentynyl,  
 1-methyl-4-pentynyl, 2-methyl-3-pentynyl, 2-methyl-4-pentynyl,  
 5 3-methyl-4-pentynyl, 4-methyl-2-pentynyl, 1,1-dimethyl-2-butynyl,  
 1,1-dimethyl-3-butynyl, 1,2-dimethyl-3-butynyl, 2,2-dimethyl-3-butynyl,  
 1-ethyl-2-butynyl, 1-ethyl-3-butynyl, 2-ethyl-3-butynyl,  
 1-ethyl-1-methyl-2-propynyl and the like.

The terms “C<sub>1-6</sub>alkoxy”, “C<sub>1-6</sub>alkylthio”, “C<sub>1-6</sub>alkylamino”,  
 10 “di(C<sub>1-6</sub>alkyl)amino”, “C<sub>1-6</sub>alkylcarbonyloxy”, “C<sub>1-6</sub>alkoxycarbonyl”,  
 “C<sub>1-6</sub>alkylcarbonyl”, “C<sub>1-6</sub>alkylsulfonyl”, “C<sub>1-6</sub>alkylsulfinyl”,  
 “C<sub>1-6</sub>alkylsulfoamino”, “C<sub>1-6</sub>alkylformylamino”, “C<sub>1-6</sub>alkylcarbamoyl”,  
 “di(C<sub>1-6</sub>alkyl)carbamoyl”, “C<sub>1-6</sub>alkylaminosulfonyl”,  
 “di(C<sub>1-6</sub>alkyl)aminosulfonyl” as used herein respectively refer to  
 15 “C<sub>1-6</sub>alkyl-O-”, “C<sub>1-6</sub>alkyl-S-”, “C<sub>1-6</sub>alkyl-NH-”, “(C<sub>1-6</sub>alkyl)<sub>2</sub>N-”,  
 “C<sub>1-6</sub>alkyl-C(O)-O-”, “C<sub>1-6</sub>alkyl-O-C(O)-”, “C<sub>1-6</sub>alkyl-C(O)-”,  
 “C<sub>1-6</sub>alkyl-SO<sub>2</sub>-”, “C<sub>1-6</sub>alkyl-SO-”, “C<sub>1-6</sub>alkyl-SO<sub>2</sub>-NH-”,  
 “C<sub>1-6</sub>alkyl-C(O)-NH-”, “C<sub>1-6</sub>alkyl-NH-C(O)-”, “(C<sub>1-6</sub>alkyl)<sub>2</sub>N-C(O)-”,  
 “C<sub>1-6</sub>alkyl-NH-SO<sub>2</sub>-”, “(C<sub>1-6</sub>alkyl)<sub>2</sub>N-SO<sub>2</sub>-”, wherein “C<sub>1-6</sub>alkyl” is defined  
 20 as above.

The terms “C<sub>1-4</sub>alkoxy”, “C<sub>1-4</sub>alkylthio”, “C<sub>1-4</sub>alkylamino”,  
 “di(C<sub>1-4</sub>alkyl)amino”, “C<sub>1-4</sub>alkylcarbonyloxy”, “C<sub>1-4</sub>alkoxycarbonyl”,  
 “C<sub>1-4</sub>alkylcarbonyl”, “C<sub>1-4</sub>alkylsulfonyl”, “C<sub>1-4</sub>alkylsulfinyl”,  
 “C<sub>1-4</sub>alkylsulfoamino”, “C<sub>1-4</sub>alkylformylamino”, “C<sub>1-4</sub>alkylcarbamoyl”,  
 25 “di(C<sub>1-4</sub>alkyl)carbamoyl”, “C<sub>1-4</sub>alkylaminosulfonyl”,  
 “di(C<sub>1-4</sub>alkyl)aminosulfonyl” as used herein respectively refer to  
 “C<sub>1-4</sub>alkyl-O-”, “C<sub>1-4</sub>alkyl-S-”, “C<sub>1-4</sub>alkyl-NH-”, “(C<sub>1-4</sub>alkyl)<sub>2</sub>N-”,  
 “C<sub>1-4</sub>alkyl-C(O)-O-”, “C<sub>1-4</sub>alkyl-O-C(O)-”, “C<sub>1-4</sub>alkyl-C(O)-”,  
 “C<sub>1-4</sub>alkyl-SO<sub>2</sub>-”, “C<sub>1-4</sub>alkyl-SO-”, “C<sub>1-4</sub>alkyl-SO<sub>2</sub>-NH-”,  
 30 “C<sub>1-4</sub>alkyl-C(O)-NH-”, “C<sub>1-4</sub>alkyl-NH-C(O)-”, “(C<sub>1-4</sub>alkyl)<sub>2</sub>N-C(O)-”,  
 “C<sub>1-4</sub>alkyl-NH-SO<sub>2</sub>-”, “(C<sub>1-4</sub>alkyl)<sub>2</sub>N-SO<sub>2</sub>-”, wherein “C<sub>1-4</sub>alkyl” is defined  
 as above.

The term “hydroxyC<sub>1-6</sub>alkyl”, “carboxylC<sub>1-6</sub>alkyl”, “aminoC<sub>1-6</sub>alkyl”, “C<sub>1-6</sub>alkylaminoC<sub>1-6</sub>alkyl”, “sulfonylC<sub>1-6</sub>alkyl”, “sulfoaminoC<sub>1-6</sub>alkyl”, “aminosulfonylC<sub>1-6</sub>alkyl”, “carbamoylC<sub>1-6</sub>alkyl” as used herein respectively refer to C<sub>1-6</sub>alkyl substituted by hydroxy, carboxyl, amino, C<sub>1-6</sub>alkylamino, sulfonyl, sulfoamino, aminosulfonyl, carbamoyl, wherein “C<sub>1-6</sub>alkyl” is defined as above.

The terms “hydroxyC<sub>1-4</sub>alkyl”, “carboxylC<sub>1-4</sub>alkyl”, “aminoC<sub>1-4</sub>alkyl”, “C<sub>1-4</sub>alkylaminoC<sub>1-4</sub>alkyl”, “sulfonylC<sub>1-4</sub>alkyl”, “sulfoaminoC<sub>1-4</sub>alkyl”, “aminosulfonylC<sub>1-4</sub>alkyl”, “carbamoylC<sub>1-4</sub>alkyl” as used herein respectively refer to C<sub>1-4</sub>alkyl substituted by hydroxy, carboxyl, amino, C<sub>1-4</sub>alkylamino, sulfonyl, sulfoamino, aminosulfonyl, carbamoyl, wherein “C<sub>1-4</sub>alkyl” is defined as above.

The term “halogen” as used herein refers to fluorine, chlorine, bromine or iodine.



The term “haloC<sub>1-6</sub>alkyl” as used herein refers to “C<sub>1-6</sub>alkyl” substituted by one or more “halogen” atom(s), the term “haloC<sub>1-4</sub>alkyl” as used herein refers to “C<sub>1-4</sub>alkyl” substituted by one or more “halogen” atom(s), wherein “halogen” and “C<sub>1-6</sub>alkyl”, “C<sub>1-4</sub>alkyl” are defined as above.








The term “haloC<sub>1-6</sub>alkoxy” as used herein refers to “C<sub>1-6</sub>alkoxy” substituted by one or more “halogen” atom(s), the term “haloC<sub>1-4</sub>alkoxy” as used herein refers to “C<sub>1-4</sub>alkoxy” substituted by one or more “halogen” atom(s), wherein “halogen”, “C<sub>1-6</sub>alkoxy”, “C<sub>1-4</sub>alkoxy” are defined as above.

The term “3- to 8-membered cycloalkyl” as used herein refers to a cyclic alkyl group, all the ring atoms of which are carbon atoms, and one hydrogen atom is removed from the ring, including for example 3- to 7-membered cycloalkyl group, 3- to 6-membered cycloalkyl, 4- to 6-membered cycloalkyl, 5- to 6-membered cycloalkyl; Examples include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, etc..


The term “6- to 12-membered spirocyclic group” as used herein refers to a structure containing 6 - 12 carbon atoms and/or hetero atoms and at



least two rings share one atom, wherein the hetero atoms include nitrogen, oxygen and sulfur and the like. The group includes for example, 6- to 10-membered spirocyclic group, 6- to 9-membered spirocyclic group, 6- to 8-membered spirocyclic group, 7- to 8-membered spirocyclic group and


5 the like. Examples include, but are not limited to, , ,



The terms “6-membered endocyclic group” and “8- to 12- membered endocyclic group” as used herein refer to a structure containing 6 or 8 - 12 carbon atoms and/or hetero atoms and any two rings share two non-adjacent atoms, wherein the hetero atoms include nitrogen, oxygen

and sulfur and the like. Examples include, but are not limited to, ,





The term “6- to 10-membered saturated fused ring group” as used herein refers to a saturated fused ring group containing 6-10 carbon atoms or/and a hetero atom, and formed by two or more cyclic structures sharing two adjacent atoms, wherein the hetero atoms include nitrogen, oxygen and sulfur and the like. The group includes 6- to 9-membered saturated fused ring group, 6- to 8-membered saturated fused ring group and the like. Examples include, but are not limited to e.g. bicyclo[3.1.0]hexyl, 3-azabicyclo[3.1.0]hexyl, 2-azabicyclo[3.1.0]hexyl, 3-azabicyclo[3.2.0]heptyl, 2-octahydrocyclopentane[C]pyrrolyl, bicyclo[3.2.0]heptyl, 3-azabicyclo[3.2.0]heptyl, octahydrocyclopentadienyl, octahydrocyclopenta[c]pyrrolyl, octahydropyrrolo[3,4-*c*]pyrrolyl, bicyclo[4.2.0]octyl, 3-azabicyclo[4.2.0]octyl, bicyclo[4.1.0]heptyl, octahydro-1*H*-indenyl, octahydro-1*H*-isoindolyl, decahydronaphthyl, decahydroisoquinolyl and the like.

The term “6- to 14-membered aryl” as used herein refers to a cyclic aromatic group containing 6-14 carbon atoms, including 6- to 8-membered aryl, 8- to 14-membered aryl and the like. Examples include, but are not limited to, phenyl, naphthyl, phenanthryl, 2,3-dihydro-1*H*-indenyl, 1*H*-indenyl, 1,2,3,4-tetrahydronaphthyl, 1,4-dihydronaphthyl and the like.










The term “3- to 8-membered heterocyclyl” as used herein refers to monocyclic heterocyclic group having 3-8 ring atoms, and wherein at least one atom is hetero atom. The group includes 5- to 8-membered heterocyclyl, 5- to 6- membered heterocyclyl and the like. Examples include, but are not limited to, furyl, thienyl, pyrrolyl, thiazolyl, thiadiazolyl, oxazolyl, oxadiazolyl, imidazolyl, pyrazolyl, pyridyl, pyrimidinyl, 1,4-dioxinyl, 2*H*-1,2-oxazinyl, 4*H*-1,2-oxazinyl, 6*H*-1,2-oxazinyl, 4*H*-1,3-oxazinyl, 6*H*-1,3-oxazinyl, 4*H*-1,4-oxazinyl, pyridazinyl, pyrazinyl, 1,2,3-triazinyl, 1,2,4-triazinyl, 1,3,5-triazinyl,











1,2,4,5-tetrazinyl, oxepinyl, thiepinyl, azepinyl, 1,3-diazepinyl, azacyclooctatetraenyl, 2,5-dihydrothienyl, 4,5-dihydropyrazolyl, 3,4-dihydro-2*H*-pyranyl, 5,6-dihydro-4*H*-1,3-oxazinyl, aziridinyl, azetidiny, thiacyclobutanyl, tetrahydrofuryl, tetrahydropyrrolyl, imidazolidinyl, pyrazolidinyl, tetrahydrofuranyl, 1,4-dioxanyl, 1,3-dioxanyl, 1,3-dithianyl, morpholinyl, piperazinyl and the like.











The term “6- to 14-membered heteroaryl” as used herein refers to a fused ring structure having 6-14 ring atoms, and wherein at least one atom is hetero atom, formed by two or more cyclic structures sharing two adjacent atoms. The group includes 8- to 12-membered heteroaryl, “9- to 10-membered heteroaryl and the like, for example, benzene ring fused with 3- to 8-membered heterocyclyl group, 3- to 8-membered heterocyclyl group fused with 3- to 8-membered heterocyclyl group. Examples include, but are not limited to, benzofuryl, benzisofuryl, benzothienyl, indolyl, benzoxazolyl, benzimidazolyl, indazolyl, benzotriazolyl, quinolyl, isoquinolyl, acridinyl, phenanthridinyl, benzopyridazinyl, phthalazinyl, quinazolinyl, quinoxalinyl, phenazinyl, pteridinyl, purinyl, naphthyridinyl, 1,3-dihydrobenzofuryl, benzo[*d*][1.3]dioxolyl, isoindolinyl, chromanyl, 1,2,3,4-tetrahydropyrrolo[3,4-*c*]pyrrolyl, 5,6-dihydroimidazo[1.2-*a*]pyrazin-7(8*H*)-yl, 5,6-dihydro-1,7-naphthyridin-7(8*H*)-yl, 5*H*-pyrrolo[3.4-*b*]pyridin-6(7*H*)-yl, 7,8-dihydropyridino[4.3-*d*]pyrimidin-6(5*H*)-yl, 2,3,6,7-tetrahydro-1*H*-pyrazolo[4.3-*c*]pyridin-5(4*H*)-yl, 6,7-dihydrothiazolo[5.4-*c*]pyridin-5(4*H*)-yl, 3-methyl-6,7-dihydro-3*H*-pyrazolo[4.5-*c*]pyridin-5(4*H*)-yl, 2-methylhexahydrocyclopenta[*c*]pyrrol-5-yl and the like.

Particularly preferred compounds of the present invention are shown in Table 1 below.


Table 1

| No. | Structure                                                                            |
|-----|--------------------------------------------------------------------------------------|
| 1   |    |
| 2   |    |
| 3   |  |
| 4   |  |
| 5   |  |
| 6   |  |
| 7   |  |
| 8   |  |
| 9   |  |

| No. | Structure                                                                            |
|-----|--------------------------------------------------------------------------------------|
| 10  |    |
| 11  |    |
| 12  |   |
| 13  |  |
| 14  |  |
| 15  |  |
| 16  |  |
| 17  |  |
| 18  |  |
| 19  |  |

| No. | Structure                                                                            |
|-----|--------------------------------------------------------------------------------------|
| 20  |    |
| 21  |    |
| 22  |   |
| 23  |  |
| 24  |  |
| 25  |  |
| 26  |  |
| 27  |  |
| 28  |  |
| 29  |  |

The present invention also relates to the preparation of the above compound of general formula (I), said method comprising the following steps:



5

The reaction steps:

(1) Preparation of Intermediate 1

Raw material 1 (commercially available) was added to the acidic catalyst, after the complete dissolvation, added N-iodosuccinimide (NIS). The reaction solution was poured into ice-water in which sodium thiosulfate was dissolved and stirred. The resulting mixture was extracted with an organic solvent, and the organic phase was rotary evaporated to remove solvents, and dried to give Intermediate 1.

15 (2) Preparation of Intermediate 2.

The intermediate 1, anhydrous sodium carbonate, a palladium catalyst

and a metal complex ligand were placed in an anhydrous organic solvent and maintained a positive pressure of carbon monoxide atmosphere, further added triethylsilane or tri-n-butyl tin to generate hydrogen, or directly blown a mixed gas of carbon monoxide and hydrogen. After the  
 5 completion of the reaction, the reaction was purified via a reversed-phase column to give Intermediate 2.

### (3) Preparation of the compound of formula (I)

The intermediate 2 was dissolved in an organic solvent and the raw  
 10 material 2 or a salt thereof was added, and further added an alkali. After the mixture was stirred at room temperature, the reducing agent was added and stirred, separated to give the compound of formula (I).

Said acidic catalyst is selected from the group consisting of, for  
 15 example, but not limited to, methanesulfonic acid, p-toluenesulfonic acid, benzenesulfonic acid, etc..

Said palladium catalyst is selected from the group consisting of, for example, but not limited to, palladium acetate, palladium acetylacetonate ( II ), bis (triphenylphosphino) palladium dichloride, tetrakis  
 20 (triphenylphosphino) palladium.

Said metal complex ligand is selected from the group consisting of, for example, but not limited to, 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene, bis(diphenylphosphino)methane, etc. .

Said organic solvent is selected from the group consisting of, for  
 25 example, but not limited to, acetonitrile, dimethylformamide, dimethyl sulfoxide, acetone, 1,3-dimethyl-2-imidazolinone, 1,3-dimethyl-3,4,5,6-tetrahydro-2-pyrimidone, toluene, ethyl acetate, chloroform, diethyl ether, N-methylpyrrolidone, etc..

Said bases include organic bases and inorganic bases, and inorganic  
 30 bases are selected from the group consisting of, for example, but are not limited to, potassium hydroxide, sodium hydroxide, zinc hydroxide, calcium hydroxide, potassium carbonate, potassium bicarbonate, sodium

carbonate, sodium bicarbonate and the like; said organic bases are selected from the group consisting of, for example, but not limited to, an amine compound such as methylamine, trimethylamine, diethylamine, triethylamine, diisopropylamine, ethylenediamine, triethanolamine, 5 diisopropylethylamine, tributylamine, N,N-dimethylaniline, dicyclohexylamine, dibenzylamine, N-benzyl-  $\beta$  -phenylethyl amine, 1-diphenyl hydroxymethyl amine, N,N' -dibenzylethylene diamine, diethanolamine, dimethylethanolamine, 2-(diethylamino) ethanol, 2-aminoethanol, tromethamine; alkali metal salts of alcohols such as 10 sodium methoxide, potassium ethoxide, potassium tert-butoxide, etc.; alkyl lithium compound such as ethyl lithium, n-butyl lithium, sec-butyl lithium, tert-butyl lithium, etc.; lithium amide compound such as lithium diisopropylamide, lithium hexamethyldisilyl amide.

Said reducing agent is selected from the group consisting of, for 15 example, but are not limited to, lithium aluminum hydride, sodium cyanoborohydride, sodium triacetoxyboronhydride.

$R^{2a}$ ,  $R^{2b}$ ,  $R^3$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^5$ ,  $R^{6a}$ ,  $R^{6b}$ ,  $R^7$ ,  $R^8$ ,  $R^{9a}$ ,  $R^{9b}$ ,  $R^{10}$ ,  $R^{11}$  and  $R^{12}$  in the above reaction equation are defined as above. When necessary, the 20 needed functional groups can be protected, and the protecting group can be subsequently removed by a conventional method.

The term "pharmaceutically acceptable salt of the compound of formula (I) " refers to a salt prepared from a suitable inorganic or organic cation(base) when the compound of formula (I) includes an acidic group (eg. -COOH, -OH,  $SO_3H$ ), including a salt formed with alkali metal such 25 as sodium, potassium and lithium, a salt formed with alkaline earth such as calcium and magnesium, a salt formed with other metals such as aluminum, iron, zinc and copper, a salt formed with inorganic bases such as ammonium, a salt formed with organic bases such as tertiary-octyl amine, dibenzylamine, morpholine, glucamine, phenyl glycine alkyl ester, 30 ethylenediamine, N-methylglucamine, guanidine, diethylamine, triethylamine, dicyclohexylamine, N,N' -dibenzyl-ethylenediamine, chlorprocaine, procaine, diethanol amine, N-benzyl-phenylethylamine,

piperazine, tetramethylamine, tri(hydroxymethyl)aminomethane and the like. Alternatively, the term "pharmaceutically acceptable salt of the compound of formula (I)" refers to a salt prepared from a suitable inorganic or organic anion (acid) when the compound of formula (I) includes a basic functional group (eg.  $\text{-NH}_2$ ), including a salt formed with inorganic acids such as nitric acid, perchloric acid, sulfuric acid, phosphoric acid, hydrofluoric acid, hydrochloric acid, hydrobromic acid, hydroiodic acid, a salt formed with sulfonic acid such as methanesulfonic acid, trifluoromethanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid and the like; a salt formed with organic acids such as acetic acid, malic acid, fumaric acid, succinic acid, citric acid, tartaric acid, oxalic acid, maleic acid and the like; a salt formed with amino acids such as glycine, trimethyl glycine, arginine, ornithine, glutamic acid, aspartic acid and the like.

15 The term "solvate" of the present compound of formula (I) refers to the substance formed by associating with a solvent. The solvent may be an organic solvent (e.g., ethanol, methanol, propanol, acetonitrile, etc.), water, etc.. For example, the compound of the present invention can form an ethanolate with ethanol, and a hydrate with water.

20 When one or more asymmetric carbon atoms exist in the compound of formula (I) of the present invention, there are diastereomers. When the compound contains an alkenyl group or a cyclic structure, there are cis/trans isomers. When the compound contains a ketone or an oxime, there are tautomers. All these enantiomorphs, diastereomers, racemic isomers, cis-trans isomers, tautomers, geometric isomers, epimeride and mixture thereof are included within the scope of the present invention.

The term "prodrug" of the present compound of formula (I) refers to the compounds that can be converted in vivo to the active form of the compound of the present invention (see R. B. Silverman, 1992, "The Organic Chemistry of Drug Design and Drug Action", Academic Press, Chp. 8). The prodrug can be used to change the biodistribution or pharmacokinetics. For example, a hydroxyl group or a carboxyl group is

esterified to form an ester, and when the ester is administered to a patient, the ester is enzymatically or non-enzymatically hydrolyzed, and the ester group is removed via reduction or hydrolysis.

The compound of formula (I) of the present invention, the  
5 pharmaceutically acceptable salt, prodrug, solvate, or isomer thereof can be made into pharmaceutical preparations with one or more pharmaceutically acceptable carrier(s). Said pharmaceutical preparations refer to conventional preparations in the clinical use, and can be orally or parenterally applied to patients in need of such treatment. For oral  
10 administration, they can be made into conventional solid preparations such as tablets, capsules, pills, granules, etc., as well as oral liquid preparations, such as oral solutions, oral suspensions, syrups, etc.. For parenteral administration, they can be made into injections, including injection solution, a sterile powder for injection, concentrated solution for injection  
15 and suspension for injection. For rectal administration, they can be made into suppositories and the like. For transpulmonary administration, they can be made into inhalations or aerosols and the like. For topical or percutaneous administration, they can be made into ointments, pastes, creams, lotions, gels, powders, solutions or transdermal stickers and the  
20 like. These preparations can be prepared by a conventional method, adding pharmaceutically acceptable carriers such as excipients, binders, moisturizers, disintegrating agents, thickeners and the like.

The compounds of formula (I) or pharmaceutically acceptable salt, prodrug, solvate or isomer thereof can be used for the treatment and/or  
25 prophylaxis of tetracycline drug-sensitive disease including infections (e.g. infections of rickettsiales pathogens, lymphogranuloma venereum, inclusion conjunctivitis, psittacosis pathogens and other tetracycline compound resistant infections), cancers, diabetes and any other diseases which have been found to be treatable and/or preventable by tetracycline  
30 compounds. The mentioned tetracycline compounds refer to the compounds having tetracycline ring structure. Examples include aureomycin, terramycin, demeclocycline, methacycline, sancycline,

rolitetracycline, guamecycline, minocycline, doxycycline, chelocardin and the like.

The administration amount and frequency of the compound of the present invention can be adjusted according to the judgment of the clinician or pharmacist, for example according to the patient's age, weight, the severity of the symptoms. Generally, the daily dose of the compound of the present invention when administrated in a single dose or divided doses may be for example 0.2 to 3mg/kg body weight, preferably 0.5-2.5mg/kg body weight. In one embodiment, for example, a patient is daily administered with approximately 10mg-200mg, preferably 30mg-180mg of the compound of the present invention in a single dose or multiple doses. In another embodiment, a patient is firstly administered with for example, 50mg-150mg, preferably 80mg-120mg of the compound of the present invention, then with 20mg-100mg/time, preferably 50mg -80mg/time in divided doses (e.g., one to four times a day).

The compound of the present invention is a broad spectrum antibacterial and has strong antibacterial activity against both Gram-positive and-negative bacteria, including aerobic and anaerobic bacteria, and further has good pharmacokinetics and high oral bioavailability. In addition, the administration of the compound of the present invention is convenience and able to meet clinical needs. Further, the synthesis of the compound of the present invention is simple, which is advantageous for industrial production.

## Examples

Hereinafter, the present invention will be further illustrated in details by the following specific examples. However, these examples are illustrative only and should not be construed as limiting the invention in any way, and any functionally equivalent embodiments are within the scope of this invention.

In the examples, the abbreviations have the following meanings:

DAST: diethylaminosulfur trifluoride;  
DCC: N,N'-dicyclohexylcarbodiimide;  
DCM: dichloromethane;  
DMF: N,N-dimethylformamide;  
5 DMA: dimethyl adipate;  
Et<sub>3</sub>N: triethylamine;  
EtOAc: ethyl acetate;  
TEA: triethanolamine;  
TFA: trifluoroacetic acid;  
10 THF: tetrahydrofuran;  
NMP: N-methylpyrrolidone;  
NIS: N-iodosuccinimide;  
NaBH<sub>3</sub>CN: sodium cyanoborohydride;  
NaBH<sub>4</sub>: sodium borohydride;  
15 InCl<sub>3</sub>: indium trichloride;  
NH<sub>4</sub>Cl: ammonium chloride;  
MeOH: methanol;  
CCl<sub>3</sub>COCl: trichloroacetyl chloride;  
Me<sub>2</sub>S: dimethyl sulfide ;  
20 NH<sub>3</sub> • H<sub>2</sub>O: ammonia;  
STAB: sodium triacetoxymborohydride;  
Boc-: *tert*-butoxycarbonyl;  
Cbz-: benzyloxycarbonyl;  
Ph-: phenyl;  
25 Ms-: mesyl.

Minocycline hydrochloride used in the following examples was purchased from Suzhou Juli Chemical Co. Ltd and Hubei Prosperity Galaxy Chemical Co. Ltd.


30

### Example 1

Preparation of (4S,4aS,5aR,12aS)-9-formyl-4,7-bis(dimethylamino)-


3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound A)

1) : (4S,4aS,5aR,12aS)-9-iodo-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound B)



To methanesulfonic acid (200mL) was added minocycline hydrochloride (28g) slowly portionwise. After complete dissolution of the compound, NIS(38g, 168.9mmol) was added portionwise at room temperature within 3 hours. Then the reaction mixture was poured into 20mL of ice-water in which 17.9g of sodium thiosulfate had been dissolved, and followed by vigorous stirring for 30 minutes. The obtained mixture was washed with ethyl acetate, and the aqueous phase was poured into a mixture of sodium bicarbonate (260g) and n-butanol (300mL), stirred, kept stand and separated. The aqueous phase was extracted with n-butanol again. The organic phases were combined and washed with water and a saturated aqueous solution of sodium chloride once respectively, and then rotary evaporated to remove solvents, dried in vacuo to give 22.8g of Compound B as a yellow solid.

## 2) Compound A



The above obtained Compound B (14.6g), anhydrous sodium

carbonate (10.6g, 100mmol), palladium acetate (0.11g, 0.5mmol) and 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene(0.29g, 0.5mmol) were placed in anhydrous NMP(100mL), and maintained a positive pressure of carbon monoxide atmosphere. The reaction mixture was heated to 70°C, and triethylsilane (4.44mL, 27.5mmol) was added dropwise within 90 minutes using a syringe. After the completion of addition, the mixture was purified by medium-pressure reverse-phase preparative column (water/acetonitrile) to give 4.8g of compound A as a yellow solid.


LC-MS (M+H): 486 (Found)

10

## Example 2

Preparation of (4S,4aS,5aR,12aS)-9-((3,3-dimethylazetidin-1-yl)methyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 1)


15 1) Diethyl 2,2-dimethylmalonate (Compound 1-2)



Sodium (36g, 1.56mol) was dissolved in 800mL of anhydrous ethanol and was cooled with an ice-water bath, diethyl malonate (Compound 1-1) (100g, 0.62mol) was slowly added dropwise. After the completion of addition, methyl iodide(97.3mL, 1.56mol) was further added dropwise. The ice-water bath was removed after the completion of addition, the mixture was stirred at room temperature overnight. 1000mL of water was added to the reaction mixture, and then extracted three times with 1000mL of ethyl acetate, the organic phases were combined, washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, the desiccant was removed by filtration, and concentrated to give the crude Compound 1-2 as a pale yellow oil (102.1g, 87.5% yield) and used directly in the next reaction.

30


## 2) 2,2-dimethylpropane-1,3-diol (Compound 1-3)



The above obtained Compound 1-2 (101.8g, 0.54mol) was dissolved  
 5 in 750mL of dry tetrahydrofuran, cooled with an ice-water bath, lithium  
 aluminium hydride (30.8g, 0.81mol) was added portionwise to the above  
 solution. After the addition was completed, the ice-water bath was  
 removed, and the mixture was stirred at room temperature  
 overnight. Added 100mL of ethyl acetate and stirred for 2 hours to  
 10 quench the reaction, then adjusted the pH to acidic using hydrochloric  
 acid, and the solvent was removed by rotary evaporation. The resulting  
 oil was purified by a short silica gel column (dichloromethane/methanol  
 = 10: 1), and then concentrated to give the crude Compound 1-3 as a pale  
 yellow oil, 36g, yield 64%.

15


## 3) 2,2-dimethylpropane-1,3-diyl dimethanesulfonate (Compound 1-4)



The above obtained Compound 1-3 (56.3g, 0.54mol) and  
 20 triethylamine (302mL, 2.17mol) were dissolved in 600mL of methylene  
 chloride, cooled with an ice-water bath, mesyl chloride (168.7mL, 2.17  
 mol) was added dropwise to the above mixture, and stirred at room  
 temperature overnight. Added 1000mL of water, stirred and separated. The  
 aqueous phase was extracted with 300mL of dichloromethane twice, and  
 25 the organic phases were combined, washed successively with water and a  
 saturated aqueous solution of sodium chloride, dried over anhydrous  
 sodium sulfate, the desiccant was removed by filtration, the solvent was  
 removed by rotary evaporation, and the residue was recrystallized with  
 acetic acid acetate/cyclohexane (volume ratio 10: 1) to give Compound 1-4

(76.6g, 54.5% yield) as a brown solid.


4) 1,3-diiodo-2,2-dimethylpropane (Compound 1-5)



Compound 1-4 (50.4g, 0.19mol) and potassium iodide (193g, 1.16mol) were dissolved in 250mL of DMF, heated to 110 °C and stirred overnight. Added 2000mL of water, extracted three times with 1000mL of ethyl acetate. The organic phases were combined, washed successively  
 10 with water and brine, dried over anhydrous sodium sulfate, the desiccant was removed by filtration, and concentrated to give the crude Compound 1-5 as a black oil (62.6g, crude).

5) 3,3-dimethyl-1-tosylazetidine (Compound 1-6)


15



The above obtained Compound 1-5 (64.8g, 0.2mol), anhydrous potassium carbonate (82.2g, 0.6mol) and p-toluenesulfonamide (34.2g, 0.2mol) were dissolved in 200mL of DMF, and reacted at 110°C for 3  
 20 hours. Added 2000mL of water and extracted three times with 1000mL of petroleum ether. The organic phases were combined, washed successively with water and brine, dried over anhydrous sodium sulfate and filtered to remove the desiccant, and concentrated to give Compound 1-6 (21.9g, 46% yield) as a white solid.


25

6) 3,3-dimethylazetidine hydrochloride (Compound 1-7)



Compound 1-6 (7.2g, 30mmol) was dissolved in 100mL of n-amyl alcohol, the reaction solution was controlled at 110°C, and was added portionwise 6.9g of sodium metal. After the sodium disappeared, continued maintaining the temperature for 1 hour, and then cooled. Thereto was poured 100mL of water, separated the water phase, the organic phase was washed with 450mL of 2N hydrochloric acid three times. The solvent was removed by rotary evaporation and the resulting residue was dissolved in 100mL of a 2N aqueous solution of NaOH, then extracted three times with 300mL of dichloromethane, the organic phases were combined and washed twice with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate. Added 100mL of 2N hydrochloric acid and concentrated to remove solvents to give Compound 1-7 (0.9g, 24.7% yield) as a white solid.

## 7) Compound 1




Compound A (0.5g, 1.0mmol) was dissolved in DMF (10mL), added Compound 1-7 (0.3g, 2.5mmol), triethylamine (500mg, 5mmol) and anhydrous indium chloride (10mg), and the mixture was stirred at room temperature for 30 minutes, then added sodium cyanoborohydride (0.3g, 5.8mmol), stirred for 0.5h at room temperature and then separated by HPLC to give Compound 1 (35mg).

<sup>1</sup>H NMR (D<sub>2</sub>O, 400 MHz) δ : 7.42 (s, 1H), 4.29 (s, 2H), 3.78 - 3.89

(m, 4H), 3.74 (s, 1H), 3.03 (m, 1H), 2.82 (s, 7H), 2.62 (m, 1H), 2.48 - 2.56 (m, 6H), 2.26 (m, 1H), 2.13 (m, 1H), 1.53 - 1.68 (m, 1H), 1.11 - 1.28 (m, 6H)

### 5 Example 3

Preparation of (4S,4aS,5aR,12aS)-9-((3,3-difluorocyclobutylamino) methyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 2)



Compound A (0.8g, 1.65mmol), 3,3-difluorocyclobutylamine (0.5g, 3.30mmol), Et<sub>3</sub>N (0.3g, 3.30mmol) and InCl<sub>3</sub>(73mg, 0.33mmol) were dissolved in 10mL of DMF. The mixture was stirred at room temperature for 2h and then added NaBH<sub>3</sub>CN (209mg, 3.30mmol). The mixture was stirred at room temperature overnight, concentrated and isolated, then purified by preparative chromatography to obtain the objective Compound 2 as a yellow solid (0.508g).

15


LC-MS (M+H): 577.2 (Found)

20 <sup>1</sup>H-NMR (CD<sub>3</sub>OD, 400 MHz) δ : 7.65 (s, 1H), 4.24 (s, 2H), 4.09(s, 1H), 3.84(m, 1H), 3.40(m, 1H), 2.83-3.22 (m, 12H), 2.62-2.82(m, 6H), 2.15-2.40(m, 2H), 1.67(m, 1H)

### Example 4


25 Preparation of (4S,4aS,5aR,12aS)-9-((spiro[2.3]hexan-5-ylamino) methyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 3)

1) *tert*-Butyl 3-methylenecyclobutylcarbamate (Compound 3-2)



Triphenylphosphonium bromide (11g, 30.8mmol) was suspended in 250mL of tetrahydrofuran and cooled in an ice-salt bath to below -5°C, added potassium tert-butoxide (4g, 35.6mmol) in four portions. After the addition, warmed to room temperature and stirred for 1 hour, then cooled to -5°C. To the suspension was added dropwise a solution of tert-butyl 3-oxocyclobutylcarbamate (Compound 3-1) (5g, 27mmol) dissolved in 40mL of tetrahydrofuran. After the addition, warmed to room temperature and stirred for 2 hours, then suction filtered to remove insolubles, rotary evaporated to remove solvents, the resulting residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate = 10: 1), to give Compound 3-2 (3.5g, yield 70.7%).


2) *tert*-Butyl spiro[2.3]hexan-5-ylcarbamate (Compound 3-3)



Diethyl zinc (40mL, 1N hexane solution) was added to 100mL of anhydrous dichloromethane under nitrogen. After the addition, cooled to -78°C with dry ice-acetone, and diiodomethane (15g, 56mmol) was slowly added dropwise. After the addition stirring was continued for 30 minutes, and then changed into the ice-water bath. A solution of Compound 3-2 (1.7g, 9.3mmol) dissolved in 100mL of dichloromethane was further added dropwise and stirred overnight, added 200mL of water to quench the reaction, liquid separated, and the aqueous phase was extracted with 200mL of dichloromethane three times. The organic phases were combined, rotary evaporated to remove solvents, and purified by silica gel column chromatography (petroleum ether/ethyl acetate = 10 : 1) to give


Compound 3-3 (1.03g, 56.1% yield).

### 3) Spiro[2.3]hexan-5-amine trifluoroacetate (Compound 3-4)



The above obtained Compound 3-3 (1g, 5.1mmol) was dissolved in 5mL of trifluoroacetic acid, stirred for 10 minutes, rotary evaporated to remove trifluoroacetic acid to give 1.1g of Compound 3-4 and used directly in the next step reactions.

### 4) Compound 3



Compound A (0.5g, 1.03mmol) was dissolved in DMF (10mL), added Compound 3-4 (0.7g, 3.3mmol), triethylamine (500mg, 5mmol) and anhydrous indium trichloride (10mg), and the mixture was stirred at room temperature for 30 minutes, added sodium cyanoborohydride (0.3g, 4.8mmol), then stirred at room temperature for 0.5h, separated by HPLC to give Compound 3 (41mg).

$^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ , 400 MHz)  $\delta$  : 7.82 (s, 1H), 4.08 (s, 2H), 3.92(s, 1H), 3.84(s, 1H), 3.10 (s, 7H), 2.73 - 2.98 (m, 8H), 2.42 (m, 1H), 2.38 (m, 2H), 2.13 (m, 3H), 1.45 - 1.60 (m, 1H), 0.30 (m, 4H)


### Example 5

Preparation of (4S,4aS,5aR,12aS)-9-((spiro[3.3]heptan-2-

ylmethylamino)methyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,1-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide  
(Compound 4)


1) 3-methylenecyclobutylmethylamine (Compound 4-2)

5




AlLiH<sub>4</sub> (2.45g, 64.5mmol) was dissolved in 50mL of anhydrous tetrahydrofuran, added a solution of 3-methylenecyclobutylcarbonitrile (compound 4-1) (5g, 53.7mmol) dissolved in 20mL of tetrahydrofuran, heated to reflux and stirred for 2 hours. After cooling, water (5mL, 0.7mmol) was slowly added dropwise under cooling in an ice-water bath, after the completion of addition, stirring was continued for 2 hours, and filtrated through Celite to remove insolubles. The resulting filtrate was rotary evaporated to remove solvents to give Compound 4-2 (3.5g, 67.1% yield).

2) Benzyl 3-methylenecyclobutylmethylcarbamate (Compound 4-3)




Compound 4-2 (3g, 30.9mmol) was dissolved in 50mL of tetrahydrofuran, added Na<sub>2</sub>CO<sub>3</sub> (6.3g, 60mmol) and benzyl chloroformate (5.25g, 30.8mmol). The mixture was stirred for 12 hours at room temperature, added 100mL of water, extracted three times with 300mL of ethyl acetate, combined the organic phases, rotary evaporated to remove solvents and purified by column chromatography (petroleum ether/ethyl acetate = 10: 1) to give Compound 4-3 ( 4g, 56% yield).

3) Benzyl (5,5-dichloro-6-oxospiro[3.3]heptan-2-yl)methylcarbamate (Compound 4-4)



5 Compound 4-3 (3g, 12.98mmol) and zinc-copper alloy (7.78g, 64.9mmol) were placed in 100mL of diethyl ether, added dropwise a solution of  $\text{CCl}_3\text{COCl}$  (6.97g, 38.33mmol) dissolved in 15mL of DMA with stirring slowly. After the completion of addition, the mixture was stirred at room temperature for 18 hours. To the reaction mixture was  
 10 poured 100mL of saturated aqueous solution of sodium bicarbonate, and suction filtrated to remove insolubles, liquid separated, the aqueous phase was extracted with diethyl ether. The organic phases were combined, dried over anhydrous sodium sulfate, suction filtrated to remove the desiccant, and rotary evaporated to remove solvents, the resulting crude Compound  
 15 4-4 was used directly in the next reaction.


4) Benzyl (6-oxospiro[3.3]heptan-2-yl)methylcarbamate (Compound 4-5)



20


The above obtained Compound 4-4, zinc powder (2.18g, 33.3mmol) and  $\text{NH}_4\text{Cl}$  (1.38g, 25.8mmol) were added to 50mL of methanol and refluxed for 4 hours. Suction filtered to remove insolubles, and rotary evaporated to remove solvents. The mixture was purified by column  
 25 chromatography (petroleum ether/ethyl acetate = 5: 1) to give Compound 4-5 (2.1g, yield 59.2% (two steps)).

5) Spiro[3.3]heptan-2-ylmethanamine (Compound 4-6)



5 Compound 4-5 (2g, 7.32mmol) and hydrazine monohydrate (0.828g, 16.5mmol) and NaOH (0.585g, 14.63mmol) were added to triethylene glycol (20mL) and heated under reflux for 1 hour. Then remove the condenser, heated to 200°C in an oil bath and maintained for 3 hours, and after cooling purified by column chromatography to give Compound 4-6  
10 (0.67g, 73% yield).

## 6) Synthesis of Compound 4




Compound A (0.230g, 0.474mmol), Compound 4-6 (0.65g, 5.2mmol), triethylamine (500mg, 5mmol) and anhydrous indium trichloride (10mg) were dissolved in DMF and stirred for 1 hour, added sodium cyanoborohydride(0.090g, 1.43mmol) and stirred for 4 hours, then separated by a reverse-phase preparative chromatography to give Compound 4 (13mg).

<sup>1</sup>H-NMR (D<sub>2</sub>O, 400 MHz) δ : 7.69 (s, 1H), 4.12 - 4.27 (m, 2H), 3.77 - 3.88 (m, 1H), 2.73 - 3.07 (m, 17H), 2.41 - 2.45 (m, 1H), 2.19 (m, 1H), 2.05 - 2.14 (m, 1H), 1.87 - 1.96 (m, 2H), 1.74 - 1.82 (m, 2H), 1.58 - 1.73 (m, 7H)

## Example 6


Preparation of (4S,4aS,5aR,12aS)-9-((spiro[2.5]octan-6-ylamino)methyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 7)

1) *tert*-Butyl 4-methylenecyclohexylcarbamate (Compound 7-2)



Triphenylmethylphosphonium bromide (53.7g, 0.15mol) was dissolved in 500mL of tetrahydrofuran, potassium *tert*-butoxide (16.8g, 0.15mol) was added at -20°C, reacted for 0.5h after the temperature was raised to 0°C. Then, *tert*-butyl 4-oxo-cyclohexylcarbamate (Compound 7-1) (21.3g, 0.1mol) was dissolved in 100mL of THF and added dropwise into the flask under nitrogen. After 3 hours of reaction at room temperature, a small amount of water was added to dissolve the solids. The mixture was rotary evaporated to remove THF, extracted with anhydrous diethyl ether, and dried, concentrated, the concentrate was dissolved in *n*-hexane, and filtered by silica gel, then concentrated to give Compound 7-2 as a colorless liquid (19.5g, 92.3% yield).


2) *tert*-Butyl spiro[2.5]octan-6-ylcarbamate (Compound 7-3)



Diethyl zinc (40mL, 1 M solution in hexane) was slowly added to 300mL of anhydrous DCM at -78°C under nitrogen, further dimethyl


iodide (15g, 56mmol) was added slowly. After 30 minutes, warmed to room temperature and maintained for 30 minutes, cooled with an ice-water bath. Then a solution of Compound 7-2 (2g, 9.5mmol) dissolved in 10mL of methylene chloride was added and reacted overnight. To the reaction mixture was poured 100mL of saturated aqueous solution of ammonium chloride, and liquid separated, the organic phase was extracted with ethyl acetate (100mL) twice, the organic phases were combined and dried over anhydrous sodium sulfate. Suction filtered to remove desiccant, and rotary evaporated to remove solvents. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate = 10: 1) to give Compound 7-3 as a white solid (0.8g, 37.4% yield) .

### 3) Spiro[2.5]octan-6-amine trifluoroacetate (Compound 7-4)



Compound 7-3 (1.5g, 6.7mmol) was dissolved in 15mL of DCM and TFA (2.5mL) was added and stirred for 1 hour to give Compound 7-4 (1.6g, 99.8% yield).

### 4) Compound 7



Compound A (1.0g, 2.1mmol), Compound 7-4 (300.0mg, 1.25mmol), 1mL of TEA and 10mg of anhydrous indium trichloride were dissolved in 10mL of DMF and reacted at room temperature for 0.5h, 400mg of


sodium triacetoxyborohydride were added slowly and the reaction was continued for 1.5 hours. The reaction solution was diluted with 1L of water, and separated using reverse-phase preparative column to obtain 600mg of crude product. The crude product was further purified by semi-preparative  
 5 high-pressure chromatography to give Compound 7 (83mg) .

$^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ , 400 MHz)  $\delta$  : 7.48 (s, 1H), 4.28 (br. s., 2H), 3.75 (s, 1H), 2.98 - 3.19(m, 2H), 2.69 - 2.95(m, 7H), 2.45 - 2.65 (m, 8H), 2.28 (m, 1H), 2.13(m, 1H), 2.01 (m, 2H), 1.40 - 1.76 (m, 6H), 0.93 (m, 2H), 0.25  
 10 (m, 2H), 0.16 (m, 2H)

### Example 7

Preparation of (4S,4aS,5aR,12aS)-9-((spiro[2.5]octan-6-ylmethylamino)methyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-  
 15 1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide  
 (Compound 8)

1) Ethyl 4-methylenecyclohexylcarboxylate(Compound 8-2)




20 Triphenylmethylphosphonium bromide (53.7g, 0.15mol) was dissolved in 500mL of THF, and potassium tert-butoxide (16.8g, 0.15mol) was added at  $-20^{\circ}\text{C}$ . Reacted for 0.5h after the temperature was raised to  $0^{\circ}\text{C}$ . Subsequently, ethyl 4-oxo-cyclohexylcarboxylate (Compound 8-1) (17g, 0.1mol) was dissolved in 100mL of THF and added dropwise to the  
 25 flask under nitrogen, reacted at room temperature for 3 hours, then a small amount of water was added to dissolve the solid, and rotary evaporated to remove THF, extracted with anhydrous diethyl ether, dried, and concentrated, the concentrate was dissolved in n-hexane, and filtered by

silica gel, then concentrated to give Compound 8-2 as a colorless liquid (16.1g, 95.7% yield).

## 2) 4-methylenecyclohexanecarboxamide (Compound 8-3)

5




Compound 8-2 (16.1g, 0.096mol) and sodium hydroxide (8g, 0.2mol) were dissolved in a mixed solvent of 100mL of methanol and water (volume ratio 1: 1), reacted at 50°C for 0.5h, and rotary evaporated to  
 10 remove methanol. Then added 200mL of water, and adjusted the pH to acidic, extracted with dichloromethane, washed with saturated brine, and then dried over anhydrous sodium sulfate, the desiccant was removed by filtration, the solvent was removed by rotary evaporation, and concentrated to give a white solid.

15 The above obtained white solid and N-hydroxysuccinimide (13.2g, 0.115mol) were dissolved in 200mL of DCM, DCC (23.65g, 0.114mol) was added slowly at -10°C. After the addition, the reaction mixture was stirred for 1 hour at room temperature, suction filtered, and rotary evaporated to remove solvents. The resulting residue was dispersed in  
 20 200mL of ammonia cooled with an ice-water bath, and then heated to 80°C, maintained at that temperature for 2 hours, and then pressurized to remove most of the ammonia gas, and extracted with ethyl acetate, dried and concentrated, then crystallized from ethyl acetate/petroleum ether to give Compound 8-3 (9.5g, 71.1% yield).

25


## 3) Benzyl 4-methylenecyclohexylmethylcarbamate (Compound 8-4)



Compound 8-3 (9.5g, 0.068mol) was dissolved in 300mL of THF, and lithium aluminium hydride(2.62g, 0.069mol) was added slowly at  $-10^{\circ}\text{C}$ . After the addition, reacted at room temperature for 2 hours, and added 5 mL of water after completion of the reaction of raw materials, suction filtered through Celite and the filtrate was concentrated to give an oil. The obtained oil and triethylamine (10.7mL, 0.075mol) were dissolved in 100mL of DCM, and added dropwise benzyl chloroformate (11.6g, 0.068mol) slowly at  $-10^{\circ}\text{C}$ , then further reacted for 2 hours at room temperature after the addition was completed. The reaction mixture was washed with 100mL of water and 100mL of saturated brine once respectively, dried and concentrated and purified by column chromatography (petroleum ether/ethyl acetate = 10: 1) to give Compound 8-4 as a white solid (6.3g, yield 35.7%).

15


#### 4) Benzyl spiro[2.5]octan-6-ylmethylcarbamate (Compound 8-5)



Diethyl zinc (25mL, 1M hexane solution) was slowly added to 100mL of anhydrous DCM at  $-40^{\circ}\text{C}$  under nitrogen, and further slowly added TFA (2.85g, 0.025mol) and reacted for 0.5h, then diiodomethane (6.67g, 0.025mol) was slowly added thereto. The reaction was continued for 0.5h, then Compound 8-4 (3g, 0.012mol) was dissolved in 20mL of anhydrous DCM and added dropwise to the reaction mixture under nitrogen, reacted

overnight at  $-5^{\circ}\text{C}$ , added saturated ammonium chloride solution to quench the reaction, followed by liquid separated, the organic phase was concentrated and purified by column chromatography (petroleum ether/ethyl acetate = 10: 1) to give Compound 8-5 as a white solid  
 5 (2.85g, yield 86.9%) .


### 5) Spiro[2.5]octan-6-ylmethanamine hydrochloride (Compound 8-6)



10 Compound 8-5 (2.85g, 10.4mmol) was dissolved in 30mL of methanol, added 0.2g of palladium on carbon and 1mL of concentrated hydrochloric acid, and subjected to catalytic hydrogenation reaction overnight and filtered to remove insolubles, then concentrated to give Compound 8-6 ( 1.8g, yield 98.5%).

15

### 6) Compound 8




20 Compound A (1g, 2.06mmol), Compound 8-6 (1g, 5.69mmol), 2mL of TEA and 30mg of anhydrous indium trichloride were dissolved in 10mL of DMF, reacted at room temperature for 0.5h, then slowly added 1g of sodium cyanoborohydride, and further reacted for 0.5h. The reaction mixture was diluted with 1 L of water, separated by reverse-phase preparative chromatography to give Compound 8 (320mg).

<sup>1</sup>H-NMR (CD<sub>3</sub>OD, 400 MHz) δ : 7.52 (s, 1H), 4.20 - 4.32 (m, 2H), 3.62 (br. s., 1H), 3.41-3.35(m, 1H), 2.98 (m, 4H), 2.81 (m, 5H), 2.54 - 2.75 (m, 7H), 2.06 - 2.28 (m, 2H), 1.59 - 1.86 (m, 6H), 1.17 - 1.25 (m, 2H),  
5 0.89 - 1.03 (m, 2H), 0.17 - 0.36 (m, 4H)

### Example 8

Preparation of (4*S*,4*aS*,5*aR*,12*aS*)-9-(2-azaspiro[3.3]heptan-2-ylmethyl)-4,7-bis(dimethylamino)-3,10,12,12*a*-tetrahydroxy-1,11-dioxo-1, 10 4,4*a*,5,5*a*,6,11,12*a*-octahydrotetracene-2-carboxamide (Compound 9)





Compound A (0.230g, 0.474mmol) was dissolved in DMF (2mmol), and added 2-azaspiro[3.3]heptane hydrochloride (0.126g, 0.948mmol) and triethylamine (180mg, 1.8mmol). After the mixture was stirred at room temperature for 30 minutes, added sodium cyanoborohydride (0.2g, 1.7mmol), then stirred for 4 hours at room temperature, separated by HPLC to give Compound 9 (60mg).  
15


<sup>1</sup>H-NMR (CD<sub>3</sub>OD, 400 MHz) δ : 7.57 (s, 1H), 4.35 (br. s., 2H), 4.16 (br. s., 4H), 3.69 (s, 1H), 3.34 - 3.50 (m, 1H), 2.82 - 2.97 (m, 7H), 2.77 (d, 1H), 2.56 - 2.70 (m, 6H), 2.29 (t, 4H), 2.20 (d, 2H), 1.85 (m, 2H), 1.57 - 1.71 (m, 1H)  
20

### Example 9

Preparation of (4*S*,4*aS*,5*aR*,12*aS*)-9-(((3,3-difluorocyclobutyl)methylamino)methyl)-4,7-bis(dimethylamino)-3,10,12,12*a*-tetrahydroxy-1, 11-dioxo-1,4,4*a*,5,5*a*,6,11,12*a*-octahydrotetracene-2-carboxamide (Compound 10)  
25




### 1) 3-oxocyclobutanecarbonitrile (Compound 10-2)



5

Compound 10-1 (10g, 0.108mol) was dissolved in a mixed solvent of 50mL of MeOH and 50mL of DCM. To the reaction system was blown  $O_3$  gas, cooled to  $-78^\circ\text{C}$ , the solution became blue. The end of the reaction was monitored with TLC (petroleum ether/ethyl acetate = 2: 1). To the reaction system was blown  $O_2$  gas for 0.5h, and then  $N_2$  gas for 0.5h to remove the excess  $O_3$  gas. 15mL of  $Me_2S$  was added to quench the reaction, and stirred at room temperature overnight, rotary dried to give the crude product, then purified by silica gel column chromatography (petroleum ether/ethyl acetate = 15: 1-7: 1) to give Compound 10-2 (7.5g, yield 73.0%) as a white solid.


### 2) 3,3-difluorocyclobutanecarbonitrile (Compound 10-3)



Compound 10-2 (8.0g, 84.21mmol) was dissolved in DCM (80mL) at 0 °C, added DAST (27g, 0.168mol). The reaction mixture was stirred overnight at room temperature, and the end of the reaction was monitored by TLC (petroleum ether/ethyl acetate = 3: 1). Ice water was added to the reaction solution, extracted with DCM, washed with saturated brine, and then dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The organic layer was concentrated under reduced pressure and purified by column chromatography (petroleum ether/ethyl acetate = 150: 1-30:1) to give the crude Compound 10-3 (8.1g) as a brown oil.


10

3 ) 3,3-difluorocyclobutylmethylamine hydrochloride (Compound 10-4)



Compound 10-3 (7.0g, 59.78mmol), NH<sub>3</sub> • H<sub>2</sub>O (7mL) and Raney nickel (7.0g, 100%/W) were dissolved in ethanol (70mL) and the mixture was stirred for 3 hours in H<sub>2</sub> (50 psi) atmosphere at room temperature. Then the mixture was filtered, added 10mL of 4M hydrochloric acid methanol solution, and concentrated to give Compound 10-4 (5.5g, 58.4%) as a white solid.

#### 4) Compound 10



Compound A (1.0g, 2.06mmol), Compound 10-4 (0.65g, 4.12mmol), Et<sub>3</sub>N (0.416g, 4.12mmol) and InCl<sub>3</sub> (91mg, 0.41mmol) were dissolved in DMF (10mL) and the mixture was stirred at room temperature for 2 hours. NaBH<sub>3</sub>CN (260mg, 4.14mmol) was added, and the mixture was stirred at room temperature overnight, purified by preparative chromatography to give Compound 10 (0.5g) as a yellow solid.


LC-MS (M+H): 591.3 (Found)

<sup>1</sup>H-NMR (D<sub>2</sub>O, 400MHz)  $\delta$ : 7.95 (s, 1H), 4.32 (s, 2H), 4.11 (s, 1H), 3.37 - 3.28 (m, 3H), 3.20 - 2.92 (m, 13H), 2.79 (m, 2H), 2.65 - 2.54 (m, 1H), 2.52 - 2.40 (m, 4H), 2.31 - 2.15 (m, 1H), 1.68 (m, 1H)

### Example 10


Preparation of (4S,4aS,5aR,12aS)-9-(4-azaspiro[2.4]heptan-4-ylmethyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 11)

1) 4-azaspiro[2.4]heptane trifluoroacetate (Compound 11-2)



*tert*-Butyl 4-azaspiro[2.4]heptane-4-carboxylate (Compound 11-1) (0.75g, 3.80mmol) and TFA (3mL) were added in DCM (10mL) and the mixture was stirred at room temperature for 1 hour and concentrated to give Compound 11-2 (800mg, 99.6%) as a yellow oil.

2) Compound 11




Compound A (0.8g, 1.65mmol), Compound 11-2 (0.7g, 3.30mmol), Et<sub>3</sub>N (0.33g, 3.30mmol) and InCl<sub>3</sub> (73mg, 0.33mmol) were dissolved in DMF (8mL), and stirred at room temperature for 2 hours. NaBH<sub>3</sub>CN (0.21g, 3.30mmol) was added, and the mixture was stirred at room temperature overnight, purified by preparative chromatography to give Compound 11 (0.21g) as a yellow solid.

LC-MS (M+H): 567.3 (Found)

<sup>1</sup>H-NMR (CD<sub>3</sub>OD, 400 MHz) δ : 7.71 (d, 1H), 4.30 - 4.23(m, 2H), 4.08(s, 1H), 3.49 - 3.36 (m, 3H), 3.25 - 2.85 (m, 9H), 2.76 (m, 5H), 2.48 - 1.90 (m, 6H), 1.65 (m, 2H), 1.39 (m, 1H), 1.02 (m, 2H)

### Example 11


Preparation of (4S,4aS,5aR,12aS)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-9-(((3-methyloxetan-3-yl)methylamino)methyl)-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 14)



4.34 (m, 2H), 4.09(s, 1H), 3.44(m, 3H), 2.89-3.12(m, 8H), 2.74(s, 6H), 2.15 - 2.40 (m, 2H), 1.67 (m, 1H), 1.46 (s, 3H)

## Example 12

5 Preparation of  
(4S,4aS,5aR,12aS)-9-((6,6-difluoro-2-azaspiro[3.3]heptan-2-yl)methyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 16)




10 The entitled compound was prepared according to the same method of Example 8 but replacing 2-azaspiro[3.3]heptane hydrochloride with 6,6-difluoro-2-azaspiro[3.3]heptane hydrochloride.

15  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ , 400 MHz)  $\delta$  : 7.48 - 7.57 (m, 1H), 4.40 - 4.49 (m, 2H), 4.24 - 4.36 (m, 4H), 4.02 - 4.13 (m, 1H), 3.37 - 3.50 (m, 1H), 2.85 - 3.05 (m, 10H), 2.56 - 2.70 (m, 8H), 2.14 - 2.33 (m, 2H), 1.57 - 1.75 (m, 1H)

## 20 Example 13

Preparation of (4S,4aS,5aR,12aS)-9-(2-oxa-6-azaspiro[3.3]heptan-6-ylmethyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 17)




25 The entitled compound was prepared according to the same method of

Example 8 but replacing 2-azaspiro[3.3]heptane hydrochloride with 2-oxa-6-azaspiro[3.3]heptane hydrochloride.

$^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ , 400 MHz)  $\delta$  : 7.85 (s, 1H), 4.27 - 4.37 (m, 2H), 3.80 - 4.00 (m, 4H), 3.51(s, 2H), 3.41(s, 2H), 2.65 - 3.20(m, 15H), 2.40 (m, 1H), 2.09(m, 1H), 1.50(m, 1H)

#### Example 14

Preparation of (4S,4aS,5aR,12aS)-9-((3,3-difluoroazetidin-1-yl)methyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 18)



Compound A (0.8g, 1.65mmol), 3,3-difluoroazetidine hydrochloride (0.43g, 3.30mmol),  $\text{Et}_3\text{N}$  (0.33g, 3.30mmol) and  $\text{InCl}_3$  (73mg, 0.33mmol) were dissolved in DMF (10mL). The mixture was stirred at room temperature for 2 hours, added  $\text{NaBH}_3\text{CN}$  (208mg, 3.30mmol), and stirred at room temperature overnight, purified by preparative chromatography to give Compound 18 (0.377g) as a yellow solid.

$^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ , 400MHz)  $\delta$  : 7.63 (br. s., 1H), 4.78 (m, 4H), 4.58 (br. s., 2H), 4.08 (br. s., 1H), 3.40(m, 1H), 2.85 - 3.08 (m, 8H), 2.72 (s, 6H), 2.13-2.45 (m, 2H), 1.68 (m, 1H)

#### Example 15

Preparation of (4S,4aS,5aR,12aS)-4,7-bis(dimethylamino)-9-((3-fluoroazetidin-1-yl)methyl)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 19)




## 5




15

2) 1-benzhydryl-3-fluoroazetidine (Compound 19-3)




DAST (26.46g, 0.164mol) was added to a solution of Compound 19-2 (13.1g, 0.0547mol) in dry DCM (200mL) at 0°C. The reaction mixture was stirred at room temperature overnight. The end of the reaction was monitored with TLC (petroleum ether/ethyl acetate = 5: 1). The reaction mixture was poured into ice water, extracted with DCM, washed with saturated brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, and the organic layer was concentrated under reduced pressure, the resulting crude product was purified by column chromatography (petroleum ether/ethyl acetate = 150: 1) to give Compound 19-3 (4.0g, 30% yield) as a white solid.

### 3) 3-fluoroazetidine hydrochloride (Compound 19-4)



Pd(OH)<sub>2</sub> (8g, 0.057mol) was added to a solution of Compound 19-3 (6g, 0.025mol) in methanol (70mL) under argon, evacuated to remove gases, and then purged with hydrogen gas for several times. The mixture was stirred overnight under H<sub>2</sub>(50 psi) at 30°C, the end of the reaction was monitored with TLC (petroleum ether/ethyl acetate = 10: 1). After filtration, to the filtrate was added dropwise 30mL of 1 mol/L hydrochloric acid-methanol solution, and concentrated to give Compound 19-4 (1.96g, 70.3% yield) as a white solid.

### 4) Compound 19




Compound A (0.8g, 1.65mmol), Compound 19-4 (0.366g, 3.30mmol), Et<sub>3</sub>N (0.333g, 3.30mmol) and InCl<sub>3</sub> (73mg, 0.33mmol) were dissolved in DMF (15mL) and stirred at room temperature for 2 hours, added NaBH<sub>3</sub>CN (208mg, 3.30mmol), further stirred at room temperature overnight, purified by preparative chromatography to give Compound 19 (0.242g) a yellow solid.

<sup>1</sup>H-NMR (CD<sub>3</sub>OD, 400MHz) δ : 7.62 (br. s., 1H), 5.40 (m, 1H), 4.30 - 4.70(m, 6H), 4.08(s, 1H), 3.33(m, 1H), 2.99(m, 8H), 2.73(m, 6H), 2.15-2.40 (m, 2H), 1.61 (m, 1H)

### Example 16

Preparation of (4S,4aS,5aR,12aS)-9-(6-azaspiro[2.5]octan-6-ylmethyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 20)




The entitled compound was prepared according to the same method of Example 10 Step 2) but replacing 4-azaspiro[2.4]heptane trifluoroacetate with 6-azaspiro[2.5]octane trifluoroacetate.

<sup>1</sup>H-NMR (D<sub>2</sub>O, 400 MHz) δ : 7.49 - 7.69 (m, 1H), 4.24 (br. s., 2H),

3.72 (s, 1H), 3.36 (m, 2H), 2.93 - 3.13 (m, 3H), 2.55 - 2.93 (m, 13H), 2.29 (m, 1H), 2.11 (m, 1H), 1.97 (m, 2H), 1.48 - 1.67 (m, 1H), 0.94 - 1.14 (m, 2H), 0.13 - 0.44 (m, 4H)

### 5 Example 17

Preparation of (4S,4aS,5aR,12aS)-9-(5-azaspiro[2.4]heptan-5-ylmethyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 21)




The entitled compound was prepared according to the same method of Example 10 Step 2) but replacing 4-azaspiro[2.4]heptane trifluoroacetate with 5-azaspiro[2.4]heptane trifluoroacetate.

15  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ , 400 MHz)  $\delta$  : 7.44 - 7.58 (m, 1H), 4.64 (m, 1H), 4.20 - 4.45 (m, 2H), 3.49 (s, 2H), 3.09 - 3.27 (m, 4H), 2.48 - 2.85 (m, 11H), 1.94 - 2.11 (m, 4H), 1.48 - 1.64 (m, 1H), 0.64 - 0.81 (m, 4H)

### Example 18

20 Preparation of (4S,4aS,5aR,12aS)-9-(6-azaspiro[3.4]octan-6-ylmethyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 22)




25 The entitled compound was prepared according to the same method of

Example 10 Step 2) but replacing 4-azaspiro[2.4]heptane trifluoroacetate with 6-azaspiro[3.4]octane trifluoroacetate.


$^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ , 400 MHz)  $\delta$  : 7.54 (s, 1H), 4.38 (s, 2H), 3.71 (br. s., 1H), 3.36 - 3.54 (m, 4H), 2.89 - 3.02 (m, 1H), 2.84 (s, 6H), 2.70 - 2.78 (m, 1H), 2.64 (s, 6H), 2.11 - 2.33 (m, 5H), 1.59 - 1.76 (m, 1H), 1.49 (m, 1H), 1.29 (s, 4H)

### Example 19

Preparation of (4S,4aS,5aR,12aS)-9-(3-azabicyclo[3.1.0]hexan-3-ylmethyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 23)




1) 3-benzyl-3-azabicyclo[3.1.0]hexane-2,4-dione (Compound 23-2)



3-oxabicyclo[3.1.0]hexane-2,4-dione (23-1) (20g, 0.179mol) was dissolved in acetic acid (100mL), and benzylamine (28.8g, 0.269mol) was added dropwise under cooling in a water bath, after the completion of addition, the temperature was raised to reflux and stirred overnight. The reaction mixture was cooled to room temperature, poured into 1 L of water


to precipitate large amount of white solid, filtered, and the filter cake was recrystallized from isopropanol to give Compound 23-2 (27g).

2 ) 3-benzyl-3-azabicyclo[3.1.0]hexane hydrochloride (Compound 23-3)



Sodium bis(2-methoxyethoxy)aluminum dihydride (trade name Red-A1) (70% toluene solution) (100mL, 0.36mol) was dissolved in 100mL of THF under nitrogen, and a solution of Compound 23-2 (16g, 0.08mol) in THF (100mL) was added dropwise under cooling with an ice bath. The mixture was stirred for 0.5h after the completion of addition, and warmed to room temperature and stirred overnight. Water was added carefully to quench the reaction, extracted with ethyl acetate, and rotary evaporated to dryness. Ethanol was added to the concentrate to dissolve it. Hydrogen chloride - ethanol solution was added to adjust the pH to be strongly acidic, and rotary evaporated to dryness to give a white solid, washed with ethyl acetate to give a crude product of Compound 23-3(18.5g) and used directly in the next step reactions.

3) 3-azabicyclo[3.1.0]hexane hydrochloride (Compound 23-4)




Compound 23-3 (18g, 0.086mol) was dissolved in 270mL of ethanol, added 10% Pd/C(2.8g), the atmosphere was replaced with hydrogen, followed by stirring at room temperature overnight. The reaction mixture

was filtered and the filtrate was rotary evaporated to dryness to give a white solid, washed with ethyl acetate to give Compound 23-3 (9.4g).

#### 4) Compound 23

5



Compound A (4.85g, 10mmol), and 3-azabicyclo[3.1.0]hexane hydrochloride(1.43g, 12mmol) were dissolved in 80mL of dichloromethane. NaBH(OAc)<sub>3</sub> (6.36g, 30mmol) was added portionwise at  
 10 room temperature, reacted for 30 minutes after the addition. 30mL of methanol was added to quench the reaction, and the reaction mixture was evaporated to dryness, added 100mL of acetone and adjusted the pH to 3~4 with trifluoroacetic acid, filtered and added excess hydrogen  
 15 chloride-ethanol solution (7 mol/L) to the filtrate until a large amount of solid precipitated, then filtered and dried to give 5g of solid. The solid was separated by a reverse-phase column to give 0.8g of a crude product, and further purified by a reverse-phase column to give Compound 23 (0.5g).

<sup>1</sup>H-NMR (CD<sub>3</sub>OD, 400 MHz) δ : 7.42(s, 1H), 3.93-4.05 (Abq, 2H) ,  
 20 3.23-3.35 (m, 3H) , 3.16 (s, 1H) , 2.92-3.06 (m, 2H) , 2.73-2.84 (m, 1H) ,  
 2.59 (s, 6H) , 2.55 (s, 6H) , 2.53-2.61 (m, 1H) , 1.96-2.07 (m, 2H) ,  
 1.52-1.63 (m, 3H) , 0.81-0.86 (m, 1H) , 0.54-0.61 (m, 1H)

#### 5) Hydrochloride of Compound 23


25 Compound 23 (0.3g) was added into a single-neck flask, and added 4.5mL of acetone, further added dropwise excess hydrogen chloride-ethanol solution (7mol/L) under cooling in an ice bath until a large amount of white solid precipitated. Stirring was further continued for

15 min, filtered, rinsed with 2mL of acetone, dried to give 0.28g of hydrochloride of Compound 23.

<sup>1</sup>H-NMR (CD<sub>3</sub>OD, 400 MHz) δ : 7.52 (s, 1H), 4.30 - 4.45 (m, 2H),  
 5 3.62 (br. s., 1H), 3.56 (m, 4H), 3.36 (m, 1H), 2.77 - 3.02 (m, 6H), 2.69 (m,  
 1H), 2.58 (s, 6H), 2.09 - 2.26 (m, 2H), 1.83 (m, 1H), 1.52 - 1.70 (m, 1H),  
 0.90 (m, 2H), 0.82 (m, 1H), 0.52 - 0.63 (m, 1H)

### Example 20

10 Preparation of (4S,4aS,5aR,12aS)-9-(3-azabicyclo[3.1.1]heptan-3-ylmethyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 24)



15 In a 25mL single-neck flask, Compound A (0.5g), 3-azabicyclo[3.1.1]heptane trifluoroacetate (150mg), 1mL TEA and 10mg of anhydrous indium trichloride were dissolved in 10mL of DMF and reacted at room temperature for 0.5h. 300mg of sodium cyanoborohydride was added slowly, and further reacted for 0.5h. Then the reaction mixture  
 20 was diluted with 1L of water, separated by a reverse-phase column to give 300mg of a crude product, further purified by high-pressure semi-preparative chromatographed to give Compound 24 (74mg).


<sup>1</sup>H-NMR (CD<sub>3</sub>OD, 400 MHz) δ : 7.63 (s, 1H), 4.41 - 4.57 (m, 2H),  
 25 4.08 (s, 1H), 3.53 - 3.91 (m, 4H), 3.38 - 3.49 (m, 1H), 2.92 - 3.06 (m, 8H),  
 2.59 - 2.70 (m, 6H), 2.53 (m, 2H), 2.08 - 2.32 (m, 2H), 1.55 - 1.75 (m, 1H),  
 1.23 - 1.40 (m, 4H)

## Example 21

## Preparation

of

(4S,4aS,5aR,12aS)-9-((6-amino-3-azabicyclo[3.1.0]hexan-3-yl)methyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide hydrochloride (the hydrochloride of Compound 26)



In a 25mL single-neck flask, Compound A (1g) and tert-butyl 3-azabicyclo[3.1.0]hexan-6-ylcarbamate (1g) were dissolved in 5mL of DMF, and reacted at room temperature for 0.5h, 1.4g of sodium triacetoxy borohydride was added slowly, and further reacted for 0.5h. Then the reaction solution was mixed with 10g of C18 fillers, packed into a column, separated by a quick preparative chromatography of ISCO (acetonitrile: water = 1-10: 100), collected the fraction which was confirmed by thin layer chromatography (TLC) to contain Compound 26. 10mL of concentrated hydrochloric acid was added and stirred at room temperature for 0.5h. After enrichment, the mixture was concentrated and freeze-dried to give hydrochloride of Compound 26 (58mg, pale yellow powder).


<sup>1</sup>H NMR(D<sub>2</sub>O, 400MHz) δ : 7.93 (s, 1H), 4.50 (m, 3H), 3.97 (s, 1H), 3.67(br, 2H), 3.18 (s, 6H), 2.86-3.09(m, 10H), 2.49(t, 1H), 2.32(s, 2H), 2.17(d, 2H), 1.58(m, 1H)

## Example 22

## Preparation

of

(4S,4aS,5aR,12aS)-9-((6-dimethyl-3-azabicyclo[3.1.0]hexan-3-yl)methyl)-4,7-bis(dimethylamino)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 27)




In a 25mL single-neck flask, Compound A (1 g) and 6,6-dimethyl-3-azabicyclo[3.1.0]hexane (1g) were dissolved in 5mL of DMF, and reacted at room temperature for 0.5h. 1.4g of sodium triacetoxyborohydride was added slowly and further reacted for 0.5h. Then the reaction solution was mixed with 10g of C18 fillers, packed into a column, separated by a quick preparative chromatography of ISCO (acetonitrile: water = 1-10:100), collected the fraction which was confirmed by thin layer chromatography (TLC) to contain Compound 27. After enrichment, the mixture was concentrated and freeze-dried to give Compound 27 (60mg, a pale yellow powder).

LC-MS(M+1): 581 (Found)

<sup>1</sup>H NMR(D<sub>2</sub>O, 400MHz) δ : 7.34 (s, 1H), 4.17 (s, 2H), 3.68 (s, 1H), 3.53(br, 2H), 2.96 (m, 1H), 2.79(s, 6H), 2.55-2.73(m, 4H), 2.43(s, 6H), 2.07-2.13(m, 2H), 1.66(s, 2H), 1.50-1.60(m, 1H), 0.99(s, 3H), 0.93(s, 3H)

### Example 23

Preparation of (4S,4aS,5aR,12aS)-4,7-bis(dimethylamino)-9-((hexahydrocyclopenta[*c*]pyrrol-2(1*H*)-yl)methyl)-3,10,12,12a-tetrahydroxy-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydrotetracene-2-carboxamide (Compound 28)




In a 25mL single-neck flask, Compound A (1g) and octahydrocyclopenta[*c*]pyrrole (1g) were dissolved in 5mL of DMF, and reacted at room temperature for 0.5h. 1.4g of sodium triacetoxyborohydride was added slowly, and further reacted for 0.5h.  
 5 Then the reaction solution was mixed with 10g of C18 fillers, packed into a column, separated by a quick preparative chromatography of ISCO (acetonitrile: water = 1-10:100), collected the fraction which was confirmed by thin layer chromatography (TLC) to contain Compound 28. After enrichment, the mixture was concentrated and freeze-dried to give  
 10 Compound 28 (50mg, a pale yellow powder).

LC-MS(M+1):581 (Found)

<sup>1</sup>H NMR(D<sub>2</sub>O, 400MHz) δ : 7.42 (s, 1H), 4.18 (s, 2H), 3.66 (s, 1H), 3.49(br, 2H), 2.93-2.98 (m, 1H), 2.79(s, 6H), 2.55-2.73(m, 5H), 2.43(s,  
 15 6H), 2.08(t, 2H), 1.38-1.56(m, 8H)

#### Example 24

Preparation of (4*S*,4*aS*,5*aR*,12*aS*)-9-((1*H*-isoindol-2(3*H*,3*aH*,4*H*,5*H*, 6*H*,7*H*,7*aH*) -yl)methyl)-4,7-bis(dimethylamino)-3,10,12,12*a*-tetrahydroxy  
 20 -1,11-dioxo-1,4,4*a*,5,5*a*,6,11,12*a*-octahydrotetracene-2-carboxamide  
 (Compound 29)



25 In a 25mL single-neck flask, Compound A (1g) and octahydro-1*H*-isoindole (1g) were dissolved in 5mL of DMF, and reacted at room temperature for 0.5h. 1.4g of sodium triacetoxyborohydride was added slowly, and further reacted for 0.5h. Then the reaction solution was mixed with 10g of C18 fillers, packed into a column, separated by a quick

preparative chromatography of ISCO (acetonitrile: water = 1-10:100), collected the fraction which was confirmed by thin layer chromatography (TLC) to contain Compound 29. After enrichment, the mixture was concentrated and freeze-dried to give Compound 29 (160mg, a pale yellow powder).

LC-MS(M+1):595 (Found)

$^1\text{H}$  NMR(MeOD,400MHz)  $\delta$  : 7.58 (s, 1H), 4.44 (s, 2H), 3.69 (s, 1H), 3.48 (s, 2H), 3.37-3.42 (m, 2H), 2.87-2.99 (m, 1H), 2.83(s, 6H), 2.65-2.71 (d, 1H), 2.62 (s, 6H), 2.48 (s, 2H), 2.14-2.26 (m, 2H), 1.14-1.71 (m, 10H)

The beneficial effects of the compound of the present invention will be further elaborated by the following in vitro and in vivo antibacterial experiments and pharmacokinetic experimental determination, but this should not be understood that the compound of the present invention has only the following beneficial effects.

#### Test Example 1


The antibacterial spectrum and in vitro antibacterial activity of the compound of the present invention

Test strains:

| Strains             | Category                                                | Source                                                             |
|---------------------|---------------------------------------------------------|--------------------------------------------------------------------|
| G+                  | Methicillin-resistant Staphylococcus aureus (MRSA)      | Shanghai Renji Hosptital                                           |
|                     | Methicillin-resistant Staphylococcus epidermidis (MRSE) | Shanghai Changzheng Hosptital                                      |
|                     | Staphylococcus aureus                                   | Southwest Hospital affiliated to Third Military Medical University |
|                     | Enterococcus faecalis                                   | Shanghai Changzheng Hosptital                                      |
|                     | Enterococcus faecium                                    | Shanghai Renji Hosptital                                           |
| G-                  | Klebsiella pneumoniae                                   | Southwest Hospital affiliated to Third Military Medical University |
|                     | Escherichia coli                                        | Southwest Hospital affiliated to Third Military Medical University |
|                     | Acinetobacter baumannii                                 | Jilin Province People Hospital                                     |
|                     | Klebsiella oxytoca                                      | Jilin Province People Hospital                                     |
|                     | Acinetobacter calcoaceticus                             | Jilin Province People Hospital                                     |
|                     | Hemophilus influenzae                                   | Southwest Hospital affiliated to Third Military Medical University |
| fastidious bacteria | Streptococcus agalactiae                                | Beijing Chaoyang Hospital                                          |
|                     | Streptococcus pyogenes                                  | Shanghai Renji Hosptital                                           |
|                     | Streptococcus constellatus                              | Sun Yat-Sen Memorial Hosptital                                     |


Test substance:

control drugs: (1) tigecycline, (2) PTK-0796, see above structural  
disclosed in the "Background Art" section;  
(3)



(Hereinafter referred to as Compound c) ,

(4)



(Hereinafter referred to as Compound d) ,

5

Compound c and Compound d were prepared according to the preparation method disclosed in WO2004/064728.

The chemical names, structural formulas and preparation methods of the present compounds, see Examples.

10 Experimental method: the agar dilution method, with reference to Clinical And Laboratory Standards Institute. Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically; Approved Standard-Seventh Edition. CLSI Document M7-A7.Vol 26, no. 2, Wayne, PA: Clinical and Laboratory Standards Institute, 2006.

15

Experimental results and conclusions:

Table 2

| Test substances | antibacterial activity MIC ( $\mu$ g/mL) |       |                  |                       |                         |                        |
|-----------------|------------------------------------------|-------|------------------|-----------------------|-------------------------|------------------------|
|                 | MR SA                                    | MR SE | Escherichia coli | Klebsiella pneumoniae | Acinetobacter baumannii | Streptococcus pyogenes |
| PTK-0796        | 0.5                                      | 0.5   | 1                | 4                     | 2                       | 0.25                   |
| Compound 2      | 0.5                                      | 0.25  | 1                | 4                     | 0.5                     | 0.125                  |
| Compound 3      | 0.5                                      | 0.5   | 1                | 2                     | 4                       | 0.06                   |
| Compound 10     | 1                                        | 0.5   | 1                | 2                     | 4                       | 0.125                  |

20

Table 3

| Test substances | antibacterial activity MIC ( μ g/mL) |       |                  |                       |                         |                    |
|-----------------|--------------------------------------|-------|------------------|-----------------------|-------------------------|--------------------|
|                 | MR SA                                | MR SE | Escherichia coli | Klebsiella pneumoniae | Acinetobacter baumannii | Klebsiella oxytoca |
| Compound d      | 1                                    | 2     | 2                | 8                     | 4                       | 8                  |
| Compound 9      | 0.5                                  | 0.5   | 0.5              | 1                     | 4                       | 1                  |
| Compound 16     | 0.5                                  | 0.5   | 1                | 1                     | 1                       | 1                  |
| Compound 20     | 0.5                                  | 1     | 0.5              | 2                     | 1                       | 1                  |
| Compound 1      | 0.5                                  | 1     | 1                | 2                     | 1                       | 1                  |
| Compound 19     | 0.5                                  | 0.5   | 1                | 1                     | 0.25                    | 0.25               |

Table 4

5

| Test substance s | antibacterial activity MIC ( μ g/mL) |                         |                             |                  |                          |                       |                       |
|------------------|--------------------------------------|-------------------------|-----------------------------|------------------|--------------------------|-----------------------|-----------------------|
|                  | MR SA                                | Acinetobacter baumannii | Acinetobacter calcoaceticus | Escherichia coli | Streptococcus agalactiae | Klebsiella pneumoniae | Enterococcus faecalis |
| PTK-0796         | 2                                    | 4                       | 0.5                         | 4                | 0.5                      | 8                     | 0.25                  |
| Compound c       | 1                                    | 4                       | 0.5                         | 4                | 0.25                     | 16                    | 0.5                   |
| Compound 23      | 0.25                                 | 1                       | 0.125                       | 0.25             | 0.06                     | 1                     | 0.06                  |
| Compound 24      | 0.5                                  | 1                       | 0.25                        | 1                | 0.125                    | 2                     | 0.25                  |
| Compound 27      | 0.5                                  | 0.5                     | 0.125                       | 2                | 0.125                    | 4                     | 0.125                 |
| Compound 28      | 0.5                                  | 1                       | 0.125                       | 1                | 0.125                    | 2                     | 0.06                  |
| Compound 29      | 1                                    | 2                       | 0.25                        | 1                | 0.25                     | 4                     | 0.125                 |

Table 5

| Test substances | antibacterial activity MIC <sub>90</sub> ( μ g/mL) |                            |                       |                         |                       |
|-----------------|----------------------------------------------------|----------------------------|-----------------------|-------------------------|-----------------------|
|                 | MRSA                                               | Streptococcus constellatus | Staphylococcus aureus | Acinetobacter baumannii | Hemophilus influenzae |
| tigecycline     | 1                                                  | 0.25                       | 0.5                   | 4                       | 0.5                   |
| PTK-0796        | 2                                                  | 0.5                        | 0.5                   | 8                       | 1                     |
| Compound 23     | 0.5                                                | 0.125                      | 0.25                  | 2                       | 0.5                   |

10       Wherein MIC represents the minimum inhibitory concentration, MIC<sub>90</sub> represents the 90% inhibitory concentration.

As shown in Tables 2-5, the compounds of the present invention have

good activity to both Gram-positive bacteria and Gram-negative bacteria, and the antibacterial activity thereof is higher than that of the control drugs.

5        Test Example 2    In vivo pharmacokinetics experiment of the compounds of the present invention in rats

Test animals: male SD rats, weighing 230-250g, purchased from Shanghai SLAC Laboratory Animals Co., Ltd., Certificate of Conformity:  
10 SCXK (SH) 2007-0005 22045; Six rats were tested for each compound.

Test substances: Compound 2 of the present invention, prepared in accordance with the method of Example 3; dissolved in physiological saline.

Internal standard substance: The acetonitrile solution of 50ng/ml  
15 KBP-5747 was used as an internal standard solution in the measurement of Compound 2.

Experimental Methods:

Administration of Compound 2 of the present invention: intravenous  
20 bolus injection (iv), dose: 5mg/kg, administrating volume: 5mL/kg; gavage (po), dose: 5mg/kg, administrating volume: 5mL/kg. Prior to administration help but water fasting for 12 hours, 4 hours after administration to food.

Blood sampling time point

25        iv blood sampling time point: denoted by 0 min before administration, 5min, 15min, 30min, 1h, 2h, 4h, 8h, 24h, 30h, 48h after administration

po blood sampling time point: denoted by 0 min before administration, 5min, 15min, 30min, 1h, 2h, 4h, 8h, 24h, 30h, 48h after administration

Each time point to take around 150  $\mu$ l whole blood, centrifugated at  
30 2000  $\times$  g at 4°C for 5 minutes in a low-temperature high-speed centrifuge to separate plasma, the resulting plasma frozen was at -70°C in a refrigerator.

Plasma samples analysis: 30  $\mu$ l of plasma was taken and added 100  $\mu$

1 of internal standard solution, subjected to vortex at 14,000 rev/min for 5 minutes, then to centrifugation at 12,000 rpm/min for 5 minutes. The supernatant was taken and analyzed using LC-MS/MS.

Formula

$$5 \quad \text{Absolute Bioavailability } F\% = \frac{[AUC]_{\text{INF (po)}} \times \text{Dose}_{(\text{iv})}}{[AUC]_{\text{INF (iv)}} \times \text{Dose}_{(\text{po})}}$$

Experimental results and conclusions

10 Table 6 The evaluation results of pharmacokinetics of the present compound (iv)

| Test substance | dose (mg/kg) | AUC <sub>inf</sub> (ng/mL/h) | V <sub>ss</sub> (L/kg) | T <sub>1/2</sub> (h) | CL (L/kg/h) |
|----------------|--------------|------------------------------|------------------------|----------------------|-------------|
| Compound 2     | 5            | 2293                         | 5.12                   | 4.07                 | 2.23        |

15 Table 7 The evaluation results of pharmacokinetics of the present compound (po) in rats

|            | dosage (mg/kg) | AUC <sub>inf</sub> (ng/mL/h) | C <sub>max</sub> (ng/mL) | T <sub>max</sub> (h) | T <sub>1/2</sub> (h) | F%   |
|------------|----------------|------------------------------|--------------------------|----------------------|----------------------|------|
| Compound 2 | 5              | 777                          | 251                      | 1.00                 | 2.32                 | 33.9 |

AUC<sub>inf</sub> represents the area under the plasma concentration versus time curve from zero to infinity;

CL represents the plasma clearance;

20 V<sub>ss</sub> represents the apparent volume of distribution;

T<sub>1/2</sub> represents the half-life;

T<sub>max</sub> represents the time to maximum plasma concentration;

C<sub>max</sub> represents the maximum plasma concentration;

F% represents the absolute bioavailability.

25

Tables 6 and 7 show that the compounds of the present invention have

good pharmacokinetics property and high bioavailability, and therefore have great clinical value.

Test Example 3 In vivo pharmacokinetic experiment of the present  
5 compound in rats

Test animals: male SD rats, weighing 180-280g, purchased from Beijing Vital River Laboratory Animal Technology Co., Ltd.; animal Certificate of Conformity: SCXK (Beijing) 2006-0009. Six rats were  
10 tested for each compound.

Test substances:

the compounds of the present invention, prepared in accordance with the methods of above Examples, dissolved in physiological saline.

tigecycline, dissolved in physiological saline.

15 Internal standard substances:

The methanol solution of 50ng/ml alogliptin was used as an internal standard solution in the measurement of Compound 9.

The 0.1% formic acid -methanol solution of 50ng/ml alogliptin was used as an internal standard solution in the measurement of Compound 20.

20 The methanol solution of 20ng/ml alogliptin was used as an internal standard solution in the measurement of Compound 23.

The methanol solution of 200ng/ml alogliptin was used as an internal standard solution in the measurement of tigecycline.

25 Experimental Methods

Administration of the compounds of the present invention: intravenous bolus injection (iv) , dose: 5mg/kg, administrating volume: 5mL/kg; gavage (po), dose: 5mg/kg, administrating volume: 5mL/kg.

Administration of tigecycline: gavage (po), dose: 10mg/kg,  
30 administrating volume: 5mL/kg.

Prior to administration help but water fasting for 16 hours, 4 hours after administration to food.

## Blood sampling

Compound 9: iv blood sampling time point: denoted by 0 min before administration, 5min, 15min, 30min, 1h, 2h, 4h, 8h, 24h, 30h after administration; po blood sampling time point: denoted by 0 min before administration, 5min, 15min, 30min, 1h, 2h, 4h, 8h, 24h, 30h after administration;

Compound 20: iv blood sampling time point: 5min, 15min, 30min, 1h, 2h, 4h, 8h, 24h, 30h after administration; po blood sampling time point: 10min, 30min, 1h, 2h, 4h, 8h, 24h, 30h after administration;

Compound 23: iv blood sampling time point: denoted by 0 min before administration, 5min, 15min, 30min, 1h, 2h, 4h, 8h, 24h, 30h, 48h after administration; po blood sampling time point: denoted by 0 min before administration, 5min, 15min, 30min, 1h, 2h, 4h, 8h, 24h, 30h, 48h after administration;

Tigecycline: iv blood sampling time point: 5min, 15min, 30min, 1h, 2h, 4h, 6h, 8h after administration; po blood sampling time point: 10min, 30min, 1h, 2h, 4h, 6h, 8h, 24h after administration.

Each time point is taken around 100  $\mu$ l whole blood and placed in a heparinized tube, centrifugated at 8000rpm/min at 4°C for 6 minutes in a low-temperature high-speed centrifuge to separate plasma, the resulting plasma was frozen at -80°C refrigerator.

Plasma samples analysis: 20  $\mu$ l of plasma was taken and added 200  $\mu$ l of internal standard solution, subjected to vortex for 3 minutes, then to centrifugation at 12,000rpm/min for 5 minutes. 100  $\mu$ l of supernatant was taken and analyzed using LC-MS/MS.

## Formula

$$\text{Absolute Bioavailability } F\% = \frac{[\text{AUC}]_{(\text{po})} \times \text{Dose}_{(\text{iv})}}{[\text{AUC}]_{(\text{iv})} \times \text{Dose}_{(\text{po})}}$$

Experimental results: see Tables 8 and 9.

Table 8 The evaluation results of pharmacokinetics of the present compounds (iv) in rats

|             | dosage<br>(mg/kg) | AUC<br>(h × ng/mL) | V <sub>ss</sub><br>(L/kg) | T <sub>1/2</sub><br>(h) | CL<br>(L/kg/h) |
|-------------|-------------------|--------------------|---------------------------|-------------------------|----------------|
| Compound 9  | 5                 | 6071 <sup>①</sup>  | 4.65                      | 5.85                    | 0.80           |
| Compound 20 | 5                 | 11592 <sup>①</sup> | 2.22                      | 4.98                    | 0.44           |
| Compound 23 | 5                 | 9110 <sup>②</sup>  | 5.00                      | 9.94                    | 0.56           |
| tigecycline | 5                 | 5121 <sup>②</sup>  | 4.47                      | 3.8                     | 0.99           |

Table 9 The evaluation results of pharmacokinetics of the present compound (po) in rats

5

|             | dosage<br>(mg/kg) | AUC<br>(h × ng/mL) | C <sub>max</sub><br>(ng/mL) | T <sub>max</sub><br>(h) | F%   |
|-------------|-------------------|--------------------|-----------------------------|-------------------------|------|
| Compound 9  | 5                 | 216 <sup>①</sup>   | 35.6                        | 1.00                    | 3.56 |
| Compound 20 | 5                 | 605 <sup>①</sup>   | 116                         | 2.00                    | 5.22 |
| Compound 23 | 5                 | 2003 <sup>②</sup>  | 199.4                       | 1.00                    | 22.0 |
| tigecycline | 10                | 296 <sup>②</sup>   | 31.8                        | 0.5                     | 1.8  |

□ AUC<sub>last</sub> represents the area under the plasma concentration-time curve to the last measurable sampling time;

② AUC<sub>inf</sub> represents the area under the plasma concentration versus time curve from zero to infinity;

10 CL represents the plasma clearance;

V<sub>ss</sub> represents the apparent volume of distribution;

T<sub>1/2</sub> represents the half-life;

T<sub>max</sub> represents the time to maximum plasma concentration;

C<sub>max</sub> represents the maximum plasma concentration;

15 F% represents the absolute bioavailability.

In the third paragraph, page 2 of CENTER FOR DRUG EVALUATION AND RESEARCH *APPLICATION NUMBER* : 21-821 MEDICAL REVIEW(S) , it was reported that tigecycline can not be

absorbed in an oral use, and therefore it can be administrated only by intravenous injection. In addition, in the text of the paper on page 9, item of "PHARMACOKINETICS/TOXICOKINETICS" , the pharmacokinetics of tigecycline admistrated orally to a monkey was investigated and revealed tigecycline can not be absorbed in vivo when orally used. The pharmacokinetics of tigecycline administrated by intravenous injection was also investigated and the results were disclosed in page 10 of the paper as below:

| Animals | number | sex  | time<br>(h) | dosage<br>(mg/kg) | AUC<br>(ng×hr/mL) | T <sub>1/2</sub> |
|---------|--------|------|-------------|-------------------|-------------------|------------------|
| SD rats | 4      | male | 0-24        | 5(iv)             | 3550              | 1.0              |

10

As can be seen from Table 8, the intravenous administration of the compounds of the present invention provided better pharmacokinetic indicators than that actually measured in tigecycline and the above literature values, which indicated the pharmacokinetics properties of Compounds 9, 20 and 23 of the present invention are better than that of tigecycline.

As can be seen from Table 9, when the dose of tigecycline was 2 times than the dose of the present compounds, F% of the present compounds was larger than that of tigecycline. Because “F%” is the most important indicator to measure oral medication pharmacokinetic properties, it can be seen, when administered orally, the compounds of the present invention have good pharmacokinetic properties and high bioavailability, and therefore they are suitable for oral formulations.

#### 25 Test Example 4

Bactericidal effects of the present compounds in vivo in mouse thigh model with Staphylococcus aureus infection

##### 1. Test animals and strains

SPF grade CD-1 (ICR) female mice housed in SPF environment,

weighing  $25 \pm 2$ g, purchased from Beijing Vital River Laboratory Animal Technology Co., Ltd., licensed cell No: SCXK (Beijing) 2006-0009.

Staphylococcus aureus: purchased from Shanghai Renji Hospital.

5

## 2. Test substances

(1) control drugs: tigecycline, PTK-0796

(2) The compounds of the present invention

10

## 3. Test Method

### 3.1 Neutropenia mouse model

Four days before bacterial infection, intraperitoneal (ip) injection of cyclophosphamide 150mg/kg;

1 day before bacterial infection, intraperitoneal (ip) injection of  
15 cyclophosphamide 100mg/kg;

Administration volume: 10ml/kg.

### 3.2 Neutropenic mouse thigh infection model

Picked fresh Staphylococcus aureus single colonies which were plate cultured for 18h, and inoculated into broth and cultured at 35°C for 18h.  
20 The bacterial suspension was diluted with broth, and 0.5 McFarland turbidity standard indicator bacteria suspension was prepared with a McFarland turbiditor, 10-fold diluted with broth and injected into the thigh muscle of mice in an amount of 0.1mL for each thigh. The concentration of the bacteria suspension was about  $10^6$  - $10^7$  CFU/ml. The inoculum  
25 concentration was determined by an agar-plate count method.

### 3.3 Test operation

Blank Group: the animals were sacrificed by dislocation at 2h (recorded as 0h) after the inoculation of bacteria.

Solvent group: The first administration time was set at 2h (denoted as  
30 0h) after the animals were infected with bacteria, the solvent was subcutaneously administrated, the animals were sacrificed by dislocation at 24h after the administration.

Treatment group: The first administration time was set at 2h (denoted as 0h). The animals were subcutaneously administrated at a dose and in a frequency as shown in Table 10, and were sacrificed by dislocation at 24h after treatment.

5 The thigh muscle tissue was separated from the sacrificed animals, and 5mL of sodium chloride injection was added, homogenated. The homogenate obtained were diluted series, taken 0.05mL and inoculated to agar plates, two plates were inoculated in parallel for each gradient bacterial suspension, and incubated at 35°C for 18h. Observe the growth  
10 of the colonies on each plate, and select the plates with the number of colonies between 30-300 for colony count. The number of colonies was recorded as n1, n2. N represents the diluted fold. The number of colonies (units CFU) for each thigh was calculated according to the formula of " $(n_1+n_2)/2/0.05 \times 5 \times N$ ". A logarithm to the base 10 for the number of  
15 colonies of each thigh was calculated to obtain  $\log_{10}$ CFU value for each thigh.

#### 4. Results and Discussions

The  $\log_{10}$ CFU values and standard deviations for each thigh at 24h after drug administration in different drug administrated groups are shown  
20 in Table 10 as below:


Table 10

| Test substances | dosage mg/kg/day | Dose times | MIC values (ug/ml) | blank group mean (0h) | mean $\pm$ standard deviations (24h) | change values |
|-----------------|------------------|------------|--------------------|-----------------------|--------------------------------------|---------------|
| Compound 23     | 2                | 2          | 0.25               | 6.39 $\pm$ 0.07       | 7.67 $\pm$ 0.17                      | 1.28          |
|                 | 6                | 2          |                    |                       | 6.27 $\pm$ 0.41                      | -0.12         |
|                 | 18               | 2          |                    |                       | 4.35 $\pm$ 0.45                      | -2.04         |
| tigecycline     | 2                | 2          | 0.25               |                       | 8.02 $\pm$ 0.24                      | 1.63          |
|                 | 6                | 2          |                    |                       | 7.02 $\pm$ 0.43                      | 0.63          |
|                 | 18               | 2          |                    |                       | 4.20 $\pm$ 0.25                      | -2.19         |
| PTK-0796        | 2                | 2          | 0.5                |                       | 8.16 $\pm$ 0.13                      | 1.77          |
|                 | 6                | 2          |                    |                       | 7.59 $\pm$ 0.21                      | 1.20          |
|                 | 18               | 2          |                    |                       | 5.34 $\pm$ 0.79                      | -1.05         |
| solvent         | 0                | 2          | /                  |                       | 8.93 $\pm$ 0.14                      | 2.54          |

Data of Table 10 shows that when the dosage was 6mg/kg/day, Compound 23 had bactericidal effect against *Staphylococcus aureus* in vivo in mice, it's bacteria reducing value of each thigh is 0.12, whereas the  
5 control drugs tigecycline and PTK-0796 did not show bactericidal effect, their bacteria growing values of each thigh were 0.63 and 1.20 respectively. The control drugs tigecycline and PTK-0796 did not show bactericidal effect until the dosage reached 18mg/kg/day. From the above results, it can be concluded that the subcutaneous administration of  
10 compound 23 provided a better bactericidal effect than the control drugs tigecycline and PTK-0796 in vivo.

## CLAIMS

1. A compound of the general formula I:



or pharmaceutically acceptable salt, prodrug, solvate or isomer thereof,

wherein

$R^{2a}$ ,  $R^{2b}$ ,  $R^3$ ,  $R^{10}$ ,  $R^{11}$  and  $R^{12}$  are each independently hydrogen;

$R^5$ ,  $R^{6a}$ ,  $R^{6b}$  and  $R^8$  are each independently hydrogen, mercapto, halogen, hydroxy, amino, carboxyl,  $C_{1-6}$ alkyl, halo $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{1-6}$ alkoxy, halo $C_{1-6}$ alkoxy,  $C_{1-6}$ alkylthio,  $C_{1-6}$ alkylamino, di( $C_{1-6}$ alkyl)amino, hydroxy $C_{1-6}$ alkyl, carboxyl $C_{1-6}$ alkyl, amino $C_{1-6}$ alkyl,  $C_{1-6}$ alkylamino $C_{1-6}$ alkyl,  $C_{1-6}$ alkylcarbonyl,  $C_{1-6}$ alkylcarbonyloxy,  $C_{1-6}$ alkoxycarbonyl,  $C_{1-6}$ alkylsulfinyl,  $C_{1-6}$ alkylsulfonyl, sulfonic, sulfonyl $C_{1-6}$ alkyl, sulfoamino, sulfoamino $C_{1-6}$ alkyl,  $C_{1-6}$ alkylsulfoamino, aminosulfonyl,  $C_{1-6}$ alkylaminosulfonyl, di( $C_{1-6}$ alkyl)aminosulfonyl, aminosulfonyl $C_{1-6}$ alkyl,  $C_{1-6}$ alkylformylamino,  $C_{1-6}$ alkylcarbamoyl, di( $C_{1-6}$ alkyl)carbamoyl, carbamoyl, carbamoyl $C_{1-6}$ alkyl, 3- to 8-membered cycloalkyl, 6- to 14-membered aryl, 6- to 14-membered aryl- $C_{1-6}$ alkyl, 6- to 14-membered aryl-formyl, 6- to 14-membered aryl-formyloxy, 3- to 8-membered heterocyclyl, 3- to 8-membered heterocyclyl- $C_{1-6}$ alkyl, 6- to 14-membered heteroaryl or 6- to 14-membered heteroaryl- $C_{1-6}$ alkyl;

$R^7$  is  $NR^{7a}R^{7b}$ ;

$R^{4a}$ ,  $R^{4b}$ ,  $R^{7a}$  and  $R^{7b}$  are each independently hydrogen,  $C_{1-6}$ alkyl, halo $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{1-6}$ alkoxy, halo $C_{1-6}$ alkoxy,  $C_{1-6}$ alkylthio, hydroxy $C_{1-6}$ alkyl, carboxyl $C_{1-6}$ alkyl, amino $C_{1-6}$ alkyl,

C<sub>1-6</sub>alkylaminoC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkoxycarbonyl,  
 C<sub>1-6</sub>alkylsulfinyl, C<sub>1-6</sub>alkylsulfonyl, sulfonic, sulfonylC<sub>1-6</sub>alkyl,  
 sulfoaminoC<sub>1-6</sub>alkyl, aminosulfonyl, C<sub>1-6</sub>alkylaminosulfonyl,  
 di(C<sub>1-6</sub>alkyl)aminosulfonyl, aminosulfonylC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbamoyl,  
 5 di(C<sub>1-6</sub>alkyl)carbamoyl, carbamoyl, carbamoylC<sub>1-6</sub>alkyl, 3- to 8-membered  
 cycloalkyl, 6- to 14-membered aryl, 6- to 14-membered aryl, 6- to  
 14-membered aryl-C<sub>1-6</sub>alkyl, 3- to 14-membered heterocyclyl or 3- to  
 14-membered heterocyclyl-C<sub>1-6</sub>alkyl;

R<sup>9a</sup> and R<sup>9b</sup> are each independently:

10 (1) hydrogen, provided that R<sup>9a</sup> and R<sup>9b</sup> can not be Hydrogen at the  
 same time,

(2) cyclopropyl, cyclobutyl, 6- to 12-membered spirocyclic group,  
 6-membered endocyclic group, 8- to 12-membered endocyclic group or 6-  
 to 10-membered saturated fused ring group, the above groups are  
 15 unsubstituted or substituted by 1 to 3 substituent(s) which may be the same  
 or different and selected from Q<sup>1</sup>, and carbon(s) in said cyclopropyl,  
 cyclobutyl, 6- to 12-membered spirocyclic group, 6-membered endocyclic  
 group, 8- to 12-membered endocyclic group and 6- to 10-membered  
 saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s)  
 20 which may be the same or different and selected from O, CO, S, SO, SO<sub>2</sub>,  
 N and NR<sup>c</sup>, R<sup>c</sup> represents hydrogen or C<sub>1-6</sub>alkyl;

(3) cyclobutylC<sub>1-6</sub>alkyl, cyclopentylC<sub>1-6</sub>alkyl, 6- to 12-membered  
 spirocyclic group C<sub>1-6</sub>alkyl, 6- to 9-membered endocyclic group C<sub>1-6</sub>alkyl  
 or 6- to 10-membered saturated fused ring group C<sub>1-6</sub>alkyl, the above  
 25 groups are unsubstituted or substituted by 1 to 3 substituent(s) which may  
 be the same or different and selected from Q<sup>1</sup>, and carbon(s) in said  
 cyclobutyl, 6- to 12-membered spirocyclic group and 6- to 9-membered  
 endocyclic group may be replaced by 1 to 3 atom(s) or group(s) which  
 may be the same or different and selected from O, CO, S, SO, SO<sub>2</sub>, N and  
 30 NR<sup>c</sup>, and carbon(s) in said 6- to 10-membered saturated fused ring group  
 may be replaced by 1 to 3 atom(s) or group(s) which may be the same  
 or different and selected from CO, S, SO, SO<sub>2</sub>, N and NR<sup>c</sup>, R<sup>c</sup> represents


hydrogen or C<sub>1-6</sub>alkyl;

Alternatively, R<sup>9a</sup> and R<sup>9b</sup> together with the nitrogen atom to which they are attached form azetidiny, 6- to 9-membered spirocyclic group or 6- to 9-membered saturated fused ring group, the above groups are  
 5 unsubstituted or substituted by 1 to 3 Q<sup>2</sup> substituent(s) which may be the same or different, and carbon(s) in said azetidiny, 6- to 9-membered spirocyclic group and 6- to 9-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, CO, S, SO, SO<sub>2</sub>, N and NR<sup>c</sup>, R<sup>c</sup> represents hydrogen  
 10 or C<sub>1-6</sub>alkyl;

Q<sup>1</sup> and Q<sup>2</sup> are independently selected from halogen, hydroxy, amino, carboxyl, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy, haloC<sub>1-6</sub>alkyl, haloC<sub>1-6</sub>alkoxy, hydroxyC<sub>1-6</sub>alkyl, aminoC<sub>1-6</sub>alkyl, carboxylC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylamino, di(C<sub>1-6</sub>alkyl)amino, aminosulfonyl, aminosulfonylC<sub>1-6</sub>alkyl, carbamoyl,  
 15 carbamoylC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, C<sub>1-6</sub>alkoxycarbonyl, phenyl, 3- to 8-membered cycloalkyl or 3- to 8-membered heterocyclyl;

R<sup>13a</sup> and R<sup>13b</sup> are each independently hydrogen, C<sub>1-6</sub>alkyl or 3- to 8-membered cycloalkyl.


20 2. The compound of claim 1, or pharmaceutically acceptable salt, prodrug, solvate or isomer thereof, wherein formula (I) has a structure represented by formula (II) as below:



25

wherein R<sup>9a</sup> and R<sup>9b</sup> are each independently as defined in claim 1.

3. The compound of claim 1, or pharmaceutically acceptable salt, prodrug, solvate or isomer thereof, wherein formula (I) has a structure represented by formula (III) as below:



wherein  $R^{9a}$  and  $R^{9b}$  are each independently as defined in claim 1.

5        4. The compound of claim 3, or pharmaceutically acceptable salt, prodrug, solvate or isomer thereof,

wherein

$R^{9a}$  and  $R^{9b}$  are each independently:

10        (1) hydrogen, provided that  $R^{9a}$  and  $R^{9b}$  can not be hydrogen at the same time,

      (2) cyclopropyl, cyclobutyl, 6- to 10-membered spirocyclic group, 6-membered endocyclic group or 6- to 10-membered saturated fused ring group, the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same or different and selected from  $Q^1$ , and carbon(s) in said cyclopropyl, cyclobutyl, 6- to 10-membered spirocyclic group, 6-membered endocyclic group and 6- to 10-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, CO, N and  $NR^c$ ,  $R^c$  represents hydrogen or  $C_{1-4}$ alkyl,

20        (3) cyclobutyl $C_{1-4}$ alkyl, cyclopentyl $C_{1-4}$ alkyl, 6- to 10-membered spirocyclic group  $C_{1-4}$ alkyl, 6- to 9-membered endocyclic group  $C_{1-4}$ alkyl or 6- to 10-membered saturated fused ring group  $C_{1-4}$ alkyl, the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same or different and selected from  $Q^1$ , and carbon(s) in said cyclobutyl, 6- to 10-membered spirocyclic group and 6- to 9-membered endocyclic group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, CO, N and  $NR^c$ , carbon(s) in said 6- to 10-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different

and selected from CO, N and  $\text{NR}^c$ ,  $\text{R}^c$  represents hydrogen or  $\text{C}_{1-4}$ alkyl;

Alternatively,  $\text{R}^{9a}$  and  $\text{R}^{9b}$  together with the nitrogen atom to which they are attached form azetidiny, 6- to 9-membered spirocyclic group or 6- to 9-membered saturated fused ring group, the above groups are  
 5 unsubstituted or substituted by 1 to 3  $\text{Q}^2$  substituent(s) which may be the same or different, and carbon(s) in said azetidiny, 6- to 9-membered spirocyclic group and 6- to 9-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, CO, N and  $\text{NR}^c$ ,  $\text{R}^c$  represents hydrogen or  $\text{C}_{1-4}$ alkyl;

10  $\text{Q}^1$  and  $\text{Q}^2$  are independently selected from halogen, hydroxy, amino, carboxyl,  $\text{C}_{1-4}$ alkyl,  $\text{C}_{1-4}$ alkoxy or halo $\text{C}_{1-4}$ alkyl.

5. The compound of claim 4, or pharmaceutically acceptable salt, prodrug, solvate or isomer thereof,

wherein

15  $\text{R}^{9a}$  and  $\text{R}^{9b}$  are each independently:

(1) hydrogen, provided that  $\text{R}^{9a}$  and  $\text{R}^{9b}$  can not be hydrogen at the same time,

(2) cyclopropyl, cyclobutyl, 6- to 8-membered spirocyclic group or 6-membered endocyclic group, the above groups are unsubstituted or  
 20 substituted by 1 to 3 substituent(s) which may be the same or different and selected from  $\text{Q}^1$ , and carbon(s) in said cyclopropyl, cyclobutyl, 6- to 8-membered spirocyclic group and 6-membered endocyclic group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, N and  $\text{NR}^c$ ,  $\text{R}^c$  represents hydrogen, methyl or ethyl;

25 (3) cyclobutyl $\text{C}_{1-3}$ alkyl, cyclopentyl $\text{C}_{1-3}$ alkyl, 6- to 8-membered spirocyclic group  $\text{C}_{1-3}$ alkyl, 6- to 8-membered endocyclic group  $\text{C}_{1-3}$ alkyl, the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same or different and selected from  $\text{Q}^1$ , and carbon(s) in  
 30 said cyclobutyl, 6- to 8-membered spirocyclic group and 6- to 8-membered endocyclic group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, N and  $\text{NR}^c$ ,  $\text{R}^c$  represents hydrogen, methyl or ethyl;


Alternatively,  $R^{9a}$  and  $R^{9b}$  together with the nitrogen atom to which they are attached form azetidiny, 6- to 8-membered spirocyclic group or 6- to 9-membered saturated fused ring group, the above groups are unsubstituted or substituted by 1 to 3  $Q^2$  substituent(s) which may be the same or different, and carbon(s) in said azetidiny, 6- to 8-membered spirocyclic group and 6- to 9-membered saturated fused ring group may be replaced by 1 to 3 atom(s) or group(s) which may be the same or different and selected from O, N and  $NR^c$ ,  $R^c$  represents hydrogen, methyl or ethyl; and

10  $Q^1$  and  $Q^2$  are independently selected from halogen,  $C_{1-4}$ alkyl or amino.


6. The compound of claim 5, or pharmaceutically acceptable salt, prodrug, solvate or isomer thereof,


wherein  $R^{9a}$  and  $R^{9b}$  are each independently:

15 (1) hydrogen, provided that  $R^{9a}$  and  $R^{9b}$  can not be hydrogen at the same time,






the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same or different and selected from  $Q^1$ ;



or , the above groups are unsubstituted or substituted by 1 to 3 substituent(s) which may be the same or different and selected from  $Q^1$ ;


Alternatively,  $R^{9a}$  and  $R^{9b}$  together with the nitrogen atom to which




, the above groups are unsubstituted or substituted by 1 to 3  $Q^2$  substituent(s) which may be the same or different;

$Q^1$  and  $Q^2$  are independently selected from fluorine, chlorine, methyl or amino.

- 5        7. The compound of claim 1, or pharmaceutically acceptable salt, prodrug, solvate or isomer thereof, wherein the compound is selected from:






1, or pharmaceutically acceptable salt, prodrug, solvate or isomer thereof, and a pharmaceutically acceptable carrier and/or diluent.

9. A use of the compound of claim 1, or pharmaceutically acceptable salt, prodrug, solvate or isomer thereof, in the preparation of a medicament  
5 for the treatment and/or prophylaxis of tetracycline drug-sensitive disease, such as infections, cancers, diabetes and any other diseases which have been found to be treatable and/or preventable by tetracycline compounds.

10. A process for preparing the compound of general formula (I), pharmaceutically acceptable salt, prodrug, solvate, or isomer thereof,  
10 comprising reacting a raw material compound 1 with N-iodosuccinimide to give an intermediate compound 1, converting the intermediate compound 1 into an intermediate compound 2 with a catalyst, and then reacting the intermediate compound 2 with a raw material compound 2 to give the compound of general formula (I), the pharmaceutically acceptable salt,  
15 prodrug, solvate or isomer thereof,



wherein  $R^{2a}$ ,  $R^{2b}$ ,  $R^3$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^5$ ,  $R^{6a}$ ,  $R^{6b}$ ,  $R^7$ ,  $R^8$ ,  $R^{9a}$ ,  $R^{9b}$ ,  $R^{10}$ ,  $R^{11}$  and  $R^{12}$  are each independently as defined in claim 1.

