



(86) Date de dépôt PCT/PCT Filing Date: 1994/06/23  
 (87) Date publication PCT/PCT Publication Date: 1995/01/05  
 (45) Date de délivrance/Issue Date: 2004/10/26  
 (85) Entrée phase nationale/National Entry: 1995/02/21  
 (86) N° demande PCT/PCT Application No.: IT 1994/000095  
 (87) N° publication PCT/PCT Publication No.: 1995/000852  
 (30) Priorité/Priority: 1993/06/23 (RM93A000409) IT

(51) Cl.Int.<sup>6</sup>/Int.Cl.<sup>6</sup> G01N 33/68, C07K 14/52, G01N 33/74,  
G01N 33/566  
 (72) Inventeurs/Inventors:  
SAVINO, ROCCO, IT;  
LAHM, ARMIN, IT;  
CILIBERTO, GENNARO, IT  
 (73) Propriétaire/Owner:  
ISTITUTO DI RICERCHE DI BIOLOGIA MOLECOLARE  
P. ANGELETTI S.P.A., IT  
 (74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : METHODE DE SELECTION DE SUPERAGONISTES, ANTAGONISTES ET DE SUPERAGONISTES  
D'HORMONES DONT LE COMPLEXE DE RECEPTEURS INCLUENT LE GP130  
 (54) Title: A METHODOLOGY FOR SELECTING SUPERAGONISTS, ANTAGONISTS AND SUPERANTAGONISTS OF  
HORMONES WHOSE RECEPTOR COMPLEX INCLUDES GP 130

(57) **Abrégé/Abstract:**

It has been found that the ligands of the group of cytokins similar to Interleukin 6 (IL-6), that is Oncostatin M (OSM), Leukemia Inhibitory Factor (LIF), Ciliary Neurotrophic Factor (CNTF) and Interleukin 11 (IL-11), induce formation of a receptor complex of which the membrane molecule gp 130 is a part. It is thus possible to formulate the hypothesis that two different surfaces of the protein (in this class of hormones), known as site 1 and site 2 bind to two different molecules: site 1 to the specific receptor and site 2 to gp 130. The invention allows the identification of these sites and the isolation of variants having, with respect to the wild hormone, a greater affinity for the specific receptor (superagonists and superantagonists) or affinity for gp 130 reduced or abolished (antagonists and superantagonists). The figure shows the construct pHenΔ hIL-6 and the production of phasmidic particles that can be used to select agonists of Interleukin 6.





PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)


(51) International Patent Classification <sup>5</sup> : G01N 33/68, 33/74, C12N 15/12, 15/18, C07K 7/26	A3	(11) International Publication Number: <b>WO 95/00852</b> (43) International Publication Date: 5 January 1995 (05.01.95)
(21) International Application Number: PCT/IT94/00095 (22) International Filing Date: 23 June 1994 (23.06.94) (30) Priority Data: RM93A000409 23 June 1993 (23.06.93) IT (71) Applicant (for all designated States except US): ISTITUTO DI RICERCHE DI BIOLOGIA MOLECOLARE P. AN- GELETTI S.P.A. [IT/IT]; Via Pontina Km 30.600, I-00040 Pomezia (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): SAVINO, Rocco [IT/IT]; Via della Tecnica, 76, I-00040 Pomezia (IT). LAHM, Armin [IT/IT]; Piazza S. Maria Ausiliatrice, 18, I-00153 Roma (IT). CILIBERTO, Gennaro [IT/IT]; Viale Gorgia di Leontini, 330/19, I-00124 Casalpalocco (IT). (74) Agents: DI CERBO, Mario et al.; Società Italiana Brevetti S.p.A., Piazza di Pietra, 39, I-00186 Roma (IT).	(81) Designated States: AU, CA, CN, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments. (88) Date of publication of the international search report: 30 March 1995 (30.03.95)	

(54) Title: A METHODOLOGY FOR SELECTING SUPERAGONISTS, ANTAGONISTS AND SUPERANTAGONISTS OF HORMONES WHOSE RECEPTOR COMPLEX INCLUDES GP 130

## (57) Abstract

It has been found that the ligands of the group of cytokins similar to Interleukin 6 (IL-6), that is Oncostatin M (OSM), Leukemia Inhibitory Factor (LIF), Ciliary Neurotrophic Factor (CNTF) and Interleukin 11 (IL-11), induce formation of a receptor complex of which the membrane molecule gp 130 is a part. It is thus possible to formulate the hypothesis that two different surfaces of the protein (in this class of hormones), known as site 1 and site 2, bind to two different molecules: site 1 to the specific receptor and site 2 to gp 130. The invention allows the identification of these sites and the isolation of variants having, with respect to the wild hormone, a greater affinity for the specific receptor (superagonists and superantagonists) or affinity for gp 130 reduced or abolished (antagonists and superantagonists). The figure shows the construct pHenΔ hIL-6 and the production of phasmidic particles that can be used to select agonists of Interleukin 6.

## pHenΔ hIL6 AND PRODUCTION OF PHAGEMID PARTICLES



- 1 -

A METHODOLOGY FOR SELECTING SUPERAGONISTS, ANTAGONISTS AND SUPERANTAGONISTS OF HORMONES WHOSE RECEPTOR COMPLEX INCLUDES GP 130

DESCRIPTION

The present invention relates to a methodology for selecting superagonists, antagonists and superantagonists of hormones whose receptor complex includes gp 130.

As is known, WO 92/21029 to Genentec Inc. teaches a method for determination of agonists or antagonists of growth hormones and ligands with a similar structural conformation. The potential agonists and antagonists are put into contact with a receptor for the hormone and this causes formation of a ternary complex consisting of a molecule of the potential agonist or antagonist and two molecules of such receptor for the hormone to be agonized or antagonized. Dimerization of receptors induced by a ligand molecule allows to conclude that the ligand has two different interaction sites (site 1 and site 2), on which it is possible to operate using mutagenesis to generate agonists or antagonists.

It has now been surprisingly found that the ligands in the group of cytokines similar to Interleukin 6 (IL-6), that is Oncostatin M (OSM), Leukemia Inhibitory Factor (LIF), Ciliary Neurotrophic Factor (CNTF), and Interleukin 11 (IL-11), induce the formation of a receptor complex of which the membrane molecule gp 130 is a part. In this receptor complex the specific receptor for each of these cytokines and the membrane molecule gp 130 are always present as common elements. It is thus possible to formulate the hypothesis that site 1 and site 2 bind to two different molecules in this class of hormones: site 1 to the specific receptor and site 2 to gp 130.

Identification of the two sites is made possible, as will be seen more clearly from the following, by

**2142973**

- 2 -

construction of a three-dimensional model of the receptor complex based on the functional similarity between sequences of the human growth hormone receptor (hGH) and sequences of the receptors for the hormones in question. Isolation of variants that, with respect to the wild type hormone, have a greater affinity for the specific receptor (superagonists or superantagonists) is obtained by construction of filamentous phage libraries, for example M13, carrying the hormone, both in the wild type and mutant version.

The difference between the three-dimensional model, for example of IL-6, adopted here and the one adopted in W092/21029 leads to hypothesize different residues in helix A and C as constituents of site 2.

Modelling of the human interleukin 6 molecule is performed as follows. Knowing, from data available in scientific literature, that human interleukin 6 belongs to a class of cytokines that have four helices forming the core of their three-dimensional structure, the amino acidic sequence of human interleukin 6 was analyzed to identify the four regions in which there was the highest probability of a helix formation. In a following stage, these four helix regions of the interleukin 6 molecule were modelled in a computerized interactive graphic unit. To start, it was assumed that the orientation of the four helices might be similar to that seen in hormones such as the growth hormone or the macrophage granulocyte colony stimulation factor. To optimize packaging of the hydrophobic amino acids in the space between the four helices, adjustments to the relative positions of the helices were made. Subsequently, the loops connecting the four helices were modelled.

This three-dimensional model of interleukin 6 has enabled the identification of the two sites of interaction between human interleukin 6 and its two receptors: the low affinity receptor gp 80 (site 1) and

- 3 -

the high affinity receptor for gp 130 signal transduction (site 2). The following procedure was used to identify the two sites. From comparison of sequences it is known that all the members of one family of hematopoietic receptors are related to each other by the fact that they share a domain, known as the cytokine recognition domain. This similarity of sequences also indicates a high probability of structural similarity in corresponding parts of the various receptors, including the two interleukin 6 receptors, gp 80 and gp 130. The observation that the cytokines that bind to these receptors all have (or can be predicted to have) a similar structure, that is a four helix matrix, strongly supports the hypothesis that the interaction between these cytokines and their receptors, by means of the cytokin recognition domain, must be very similar in biologically active complexes.

Considering that the three-dimensional structure of one of these compounds (the complex made by growth hormone and the extra-cellular domain of the dimeric receptor for the growth hormone) has been determined by means of X-ray crystallography, our model of human interleukin 6 allows us to identify the potential sites of interaction between interleukin 6 and its two receptors gp 80 (site 1) and gp 130 (site 2). This by comparison with the complex involving the growth hormone and assuming that the functionally important amino acids are located in similar positions on the surface of the two hormones.

The need to provide a methodology the production of agonists, antagonists and superantagonists for hormones of the immune system whose receptor complex includes gp 130, will be explained with reference to the case of interleukin-6.

As is known, interleukin 6 is a polypeptide of 184 amino acids which, as described, belongs to the class of helical cytokines. Interleukin 6 is a multi-

**2142973**

- 4 -

functional cytokine produced by various types of cell. It acts as a differentiation and growth factor on cells of various types, such as for example cells in the immune system, hepatocytes, kidney cells, hemopoietic staminal cells, keratinocytes and neurones.

Production of superagonists of interleukin 6 would allow the use of lower therapeutic doses than those required with wild interleukin 6 in the treatment of numerous serious diseases. In fact, interleukin 6 has important and promising applications in the treatment of breast cancer, leukemia, and infectious diseases or diseases connected with disorders of the cells producing bone marrow.

On the other hand the production of antagonists or superantagonists of human interleukin 6 would allow inhibition of interleukin 6 in numerous diseases characterized by its excessive production, such as chronic autoimmune diseases, myeloma/plasmacytoma, post-menopausal osteoporosis and cancer cachexy.

The methodology for the selection of superagonists, antagonists or superantagonists of a hormone using the membrane molecule gp 130 to activate the mechanisms regulating cell physiology, according to the present invention, comprises the following operations:

- comparing the amino acid sequences of the growth hormone with the sequences of said hormone;

- comparing the amino acid sequences of the growth hormone receptor with those of the two receptors of the hormone in question, that is with the hormone-specific receptor and gp 130;

- on the basis of the above comparisons, formulating a three-dimensional model of the receptor complex based on the functional similarity between sequences of the growth hormone receptor and the two hormone receptors in question; and

- identifying the residues of the wild type

- 5 -

hormone in question that are a part of the site of interaction with the specific receptor and of the site of interaction with gp 130, respectively, introducing a mutation to said residues and evaluating biological activity of resulting mutants.

The hormone in question can be chosen from the group comprising Interleukin 6 (IL-6), Oncostatin M (OSM), Leukemia Inhibitory Factor (LIF), Ciliary Neurotrophic Factor (CNTF) and Interleukin 11 (IL-11).

For selection of superagonists of interleukin 6, the methodology according to the present invention further comprises the following additional operations:

- production of a series of phage libraries containing mutations of the following wild type residues of interleukin 6, present in the form of fusion product with filamentous phage proteins Glu 42, Glu 51, Ser 52, Ser 53, Lys 54, Glu 55, Asn 63, Lys 66, Met 67, Ala 68, Glu 69, Lys 70, Asp 71, Phe 170, Gln 175, Ser 176, Ser 177, Leu 181, Gln 183;
- generation of a phage library, each phage having a mutant interleukin 6 sequence;
- selection, from the mixed population of phages expressing interleukin 6 mutants, of that or those with an affinity for the specific receptor greater than that of wild type interleukin; and
- identification of the best amino acid sequence or sequences binding the receptor by sequencing of the DNA extracted from the selected phage particles.

In this case, a series of phage libraries can be produced containing mutations of said wild residues of interleukin 6 present as a product of fusion with the protein pIII of M13.

The methodology for selecting antagonists of interleukin 6 according to the present invention comprises - along with the operations indicated above - the following operations:

- mutation of the residues identified in claim 1, to form part of the site of interaction with gp 130 (Arg 30, Tyr 31, Gly 35, Ser 37, Ala 38, Ser 118, Lys

**2142973**

- 6 -

120, Val 121, Gln 124, Phe 125, Gln 127, Lys 128 and Lys 129), using conventional molecular biology techniques;

- evaluation of biological activity and affinity for the specific interleukin 6 receptor of the mutants produced as above, in order to identify variants of interleukin 6 whose affinity to the specific receptor is intact and that show reduction or loss of the biological activity; and

- evaluation of the above variants of interleukin 6 as antagonists for the biological activity of wild interleukin 6.

In case of selection of superantagonists of interleukin 6 by combination of the variations of amino acid sequences responsible for antagonist activity, indicated above, with amino-acid variations responsible for an increased affinity of the specific receptor for interleukin 6.

In the methodology for selecting antagonists or superantagonists of interleukin 6, the mutagenesis of the residues identified as above can be performed using a molecular biology technique chosen from the group comprising Polymerase Chain Reaction, Primer Extension, Oligonucleotide Directed Mutagenesis, and combinations thereof.

The present invention is not limited to the methodology for selection of agonists, antagonists or superantagonists of interleukin 6. On the contrary, it extends to the agonists, antagonists and superantagonists of hormones using the membrane molecule gp 130 to activate mechanisms regulating cell physiology, and that can be obtained using the selection methodology described above.

Up to this point a general description of the subject of the present invention has been given. With the aid of the following examples a detailed description of specific embodiments of the invention

- 7 -

will now be given, with the purpose of giving a better understanding of the objects, characteristics, advantages and methods of application thereof.

Figure 1 shows the construct pHen $\Delta$ hIL-6 and the production of phasmidic particles (see example 1, "Vector Construction"), used for selection of agonists of hIL-6.

Figure 2 shows the antagonistic activity of mutant IL-6 Tyr31Asp/Gly35Phe/Ser118Arg/Val121Asp with the increase of its concentration.

#### DEPOSITS

E.Coli K12 bacteria - transformed using the plasmid pHen $\Delta$ hIL-6 containing, from the site for recognition of the restriction enzyme Sall to that for the restriction enzyme NotI, a nucleotidic sequence coding for the amino acid sequence of wild type human interleukin-6 - have been deposited on 10/6/1993 with The National Collection of Industrial and Marine Bacteria Ltd. (NCIMB), Aberdeen, Scotland, UK, with access number NCIMB 40563.

#### Example 1

Application of the methodology according to the present invention for the selection of agonists of interleukin-6

#### 1) VECTOR CONSTRUCTION

The strategy consists in construction of a hybrid gene containing all the region coding for hIL-6 followed by the last 157 amino acids of protein pIII of the phage M13 and preceded by the sequence Pel B, which vectors the synthesized protein to the periplasmic space.

Expression of the hybrid gene is driven by the promotor lacZ. The construction is made in the context of vector pHen $\Delta$ e, and takes the name of pHen $\Delta$ hIL-6 (see the only figure enclosed). This plasmid also contains a phage replication origin. If a bacterial cell containing this plasmid is infected by a

2142973

- 8 -

bacteriophage known as a "helper", such as M13K07, single filament copies will be produced by the plasmid, and will be coated with the phage proteins, just like a true phage genome. These phage particles containing the plasmid are known as phasmids. Along with the normal pIII molecules, they also contain hIL6-pIII fusion molecules. A hIL-6 molecule and the gene coding its amino acid sequence are thus contained within the same unit. In the case of mutant molecules, it will be possible to determine the amino acid sequence of the molecules exposed on the surfaces of the phages obtained from selection processes, simply by sequencing the phasmid DNA.

A further characteristic of the construction pHen hIL-6 is the presence of a translation codon stop between the IL-6 gene and the pIII gene. Production of the hybrid protein is carried out in bacterial strains capable of suppressing this Stop codon. Vice versa the use of non-suppressor strains allows production of hIL-6 alone, directly in the periplasmic space.

The following experiments demonstrate the ability to use shrIL-6R to purify, from among the vast range of mutant interleukin-6 exposed on the phage using pHen hIL-6, those having the greatest affinity with said receptor, by means of amplification selection cycles.

## 2) SYSTEM CHARACTERIZATION EXPERIMENTS

### a) ELISA Test

The wells in ELISA plates were coated with hIL-6 phasmids or with M13K07 and made to react with shrIL-6R (soluble Human Recombinant IL-6R). After repeated washing the presence of the receptor was indicated using a specific monoclonal antibody conjugated with alkaline phosphatase. The signal obtained is greater in the case of hIL-6 phasmids and increases as the amount of receptor used increases.

### b) Enrichment of the hIL-6 phasmid with phasmid-K07 mixtures using shrIL-6R

2142973

- 9 -

hIL-6 phasmid particles were mixed with particles of M13K07 at a ratio of 1:100. This mixture was incubated with polystyrene balls coated with shrIL-6R. After repeated washing at a neutral pH the balls underwent stringent washing at pH 4.2, followed by elution at pH 3.6. The proportion of hIL-6 phasmids:M13K07 in the resulting eluant was then determined. The proportion is found to be 1:10, with a consequent enrichment by ten times of the hIL-6 phasmids with respect to M13K07.

c) Selection from mixtures of mutant interleukin-6 of those having the highest affinity for the receptor gp 80

Phasmid particles were produced having on their surface mutant molecules of hIL-6 with a higher (176 Arg) or lower (179 Ala) affinity with the receptor, compared with the natural version. These particles were mixed in a ratio of 1:1. The mixture was incubated with the receptor in a solid phase (as described in point b). In the eluate at pH 3.6, the ratio between the two types of particle was found to be 15:1 in favor of 176 Arg.

d) Determination of the relative affinity of M13K07, hIL-6 phasmid, 176 Arg phasmid and 179 Ala phasmid particles for the receptor gp 80

Equal amounts of particles of the various types listed above were incubated separately with the balls coated with receptor. After the usual washing sequence the number of particles recovered in the eluant at 3.6 was determined for each type of phage. Considering the value of particles recovered for the phasmid hIL-6 as 1, a value of 3 is obtained for 176 Arg, 0.2 for 179 Ala and 0.18 for M13K07. These values reflect the relative affinities of the molecules of IL-6 non exposed on the phage. It is, in fact, known that 176 Arg binds the receptor with an affinity three times greater than that of the natural molecule, whereas in

2142973

- 10 -

the case of 179 Ala the affinity for the receptor is reduced almost to zero (in fact a phasmid with this mutant, as regards binding to the receptor, behaves like M13K07).

This group of experiments has shown that it is possible, using the method according to the present invention, to select from mixtures of mutants exposed on phage those having the greatest affinity for the receptor.

#### Example 2

#### Generation and selection of antagonists of interleukin 6 using the methodology according to the present invention

The plasmid Phen $\Delta$ hIL-6 was used as a template for all the mutagenesis reactions. This plasmid is a derivative of the plasmid PHEN1 and contains the area coding for human IL-6 (SEQ ID NO:1) upstream the area coding for the carboxy-terminal portion (from codon 250 to the COOH end) of the protein pIII of the bacteriophage M13; the two coding areas are in reading frame and are separated by an Amber UAG Stop codon, which can be suppressed in bacterial strains that have the gene of the suppressor SupE integrated in their genome. The area coding for human IL-6 is also in reading frame, below the peptide PelB, a signal sequence for secretion, and the whole gene is under the control of the promotor LacZ. Finally, a single site for the restriction enzyme SacI has been introduced in the nucleotide sequence coding for the amino acids 20-21-22 of hIL-6 without changing their identity and, equally, a single site for the restriction enzyme BfrI has been introduced in the area coding for the amino acids 38-39-40 of hIL-6, likewise without changing them.

A PCR strategy (Polymerase Chain Reaction) was used to generate mutations within the codons selected in the area coding for human interleukin 6. The

- 11 -

primer downstream is H9, a 32 nucleotide primer, corresponding to positions 426-457 (antisense filament) of hIL-6 cDNA (taking the first nucleotide of the first codon of the mature polypeptide to be 1). The primer hybridization site is downstream the recognition site for the enzyme XbaI, naturally present in the cDNA. The mutagenetic primer upstream is IL-6 31D/35 PCR, a 70 nucleotides primer, whose sequence is SEQ ID NO:2.

The primer IL-6 31D/35 PCR extends from position 55 to position 124 (sense filament) of the hIL-6 cDNA and introduces degenerations into the codons coding for the amino acid 31 (wild type tyrosine) and 35 (wild type glycine). A DNA fragment of 403 pairs of bases is amplified using PCR according to standard PCR amplification protocols. Amplification is performed in 35 cycles. Each cycle consists of incubation for 2 minutes at 94°C for denaturation of the template, 2 minutes at 50°C for hybridization of the oligonucleotide and 3 minutes at 72°C for extension of the chain. The amplified fragment is digested with SacI and XbaI and purified using 2% agarose gel. The fragment generated by PCR and digested by the two enzymes is ligated into the vector pHen hIL-6 digested with the same two enzymes, purified on 0.8% agarose gel, to replace the wild type sequence.

The following Table 1 shows the biological activity, in human hepatoma cells, and binding to the receptor gp 80 for wild type interleukin-6 and mutations thereof, the mutations being indicated with an asterix.

**B**

TABLE I

Receptor binding properties and biological activity  
of wild type interleukin 6 and its mutants in helix A

	27	30	35	40	Biological activity	Receptor binding
a)	Lys* Gln Ile Arg Arg Tyr* Ile Leu Asp Gly* Ile Ser Ala Leu Arg Lys Glu	Ile Arg Arg Tyr* Ile Leu Asp Gly* Ile Ser Ala Leu Arg Lys Glu	Gln Ile Leu Asp Asp Ile Ile Ser Ala Leu Arg Lys Glu	Arg Arg Tyr* Ile Leu Asp Gly* Ile Ser Ala Leu Arg Lys Glu	100%	100%
b)	Ala Gln Ile Arg Arg Tyr Ile Leu Asp Gln Thr Ser Ala Leu Arg Lys Glu	Ile Arg Arg Tyr Ile Leu Asp Gln Thr Ser Ala Leu Arg Lys Glu	Gln Ile Leu Asp Asp Ile Ile Ser Ala Leu Arg Lys Glu	Arg Arg Tyr Ile Leu Asp Gln Thr Ser Ala Leu Arg Lys Glu	80 ■ 19%	100 ■ 4%
c)	Lys Gln Ile Arg Arg Asp Ile Leu Asp Tyr Ile Ser Ala Leu Arg Lys Glu	Ile Arg Arg Asp Ile Leu Asp Tyr Ile Ser Ala Leu Arg Lys Glu	Gln Ile Leu Asp Asp Ile Ile Ser Ala Leu Arg Lys Glu	Arg Arg Tyr Ile Leu Asp Tyr Ile Ser Ala Leu Arg Lys Glu	4.5 ■ 1%	83%
d)	Lys Gln Ile Arg Arg Asp Ile Leu Asp Phe Ile Ser Ala Leu Arg Lys Glu	Ile Arg Arg Asp Ile Leu Asp Phe Ile Ser Ala Leu Arg Lys Glu	Gln Ile Leu Asp Asp Ile Ile Ser Ala Leu Arg Lys Glu	Arg Arg Tyr Ile Leu Asp Phe Ile Ser Ala Leu Arg Lys Glu	2 ■ 0.5%	80 ■ 2%
e)	Lys Gln Ile Arg Arg Asp Ile Leu Asp Leu Ile Ser Ala Leu Arg Lys Ala	Ile Arg Arg Asp Ile Leu Asp Leu Ile Ser Ala Leu Arg Lys Ala	Gln Ile Leu Asp Asp Ile Ile Ser Ala Leu Arg Lys Ala	Arg Arg Tyr Ile Leu Asp Leu Ile Ser Ala Leu Arg Lys Ala	6 ■ 4%	177 ■ 7%
f)	Lys Gln Ile Arg Arg Asp Ile Leu Asp His Ile Ser Ala Leu Arg Lys Glu	Ile Arg Arg Asp Ile Leu Asp His Ile Ser Ala Leu Arg Lys Glu	Gln Ile Leu Asp Asp Ile Ile Ser Ala Leu Arg Lys Glu	Arg Arg Tyr Ile Leu Asp His Ile Ser Ala Leu Arg Lys Glu	38 ■ 8%	85 ■ 15%
g)	Lys Gln Ile Arg Arg Asp Ile Leu Asp Cys Ile Ser Ala Leu Arg Lys Glu	Ile Arg Arg Asp Ile Leu Asp Cys Ile Ser Ala Leu Arg Lys Glu	Gln Ile Leu Asp Asp Ile Ile Ser Ala Leu Arg Lys Glu	Arg Arg Tyr Ile Leu Asp Cys Ile Ser Ala Leu Arg Lys Glu	30 ■ 1%	82 ■ 18%

- a) is SEQ ID NO: 8
- b) is SEQ ID NO: 9
- c) is SEQ ID NO: 10
- d) is SEQ ID NO: 11
- e) is SEQ ID NO: 12
- f) is SEQ ID NO: 13
- g) is SEQ ID NO: 14

2142973

- 13 -

As can be seen from the table, the mutants (Tyr31Asp, Gly35Tyr), (Tyr31Asp, Gly35Phe) and (Tyr31Asp, Gly35Leu, Glu42Ala) show lower biological activity when compared with wild type interleukin 6. In all three cases the residual activity is 2-5% that of wild type interleukin-6. The three mutants maintain their full ability to bind to the interleukin 6 receptor gp 80. In conclusion, the three mutants have a decreasing signal transduction activity in hepatoma cells. In other words, they are antagonists of wild interleukin 6. In particular, the mutant (Tyr31Asp, Gly35Phe) is a particularly effective antagonist, as it is capable of reducing the activity of wild type interleukin 6 when used at 50 fold molar excess in tests on human hepatoma cells.

### Example 3

#### Generation and selection of further antagonists of interleukin 6 using the methodology according to the present invention

The plasmid pHEN $\Delta$ hIL-6, described in the preceding example, was used as a template for all the mutagen reactions. A PCR strategy (Polymerase Chain Reaction) was used to generate mutations within the codons selected for the area coding for human interleukin 6. The primer above is HP/1, a primer with 29 nucleotides, corresponding to positions 1-19 (sense filament) of hIL-6 cDNA (taking the first nucleotide of the first codon of the mature polypeptide to be 1). The primer hybridization site is upstream to the site recognizing the enzyme SacI, artificially introduced in the cDNA without changing the sequence coded by the latter, as described in example 2. The mutagenetic primer below is IL-6 118RCLF/121VD, a primer with 72 nucleotides, whose sequence is SEQ ID NO 3.

The primer IL-6 118RCLF/121VD extends from position 334 to position 405 (antisense filament) of the cDNA of hIL-6 and introduces degenerations into

the codons coding for the amino acid 118 (wild type serine) and 121 (wild type valine). A DNA fragment of 415 pairs of bases is amplified using PCR according to standard PCR amplification protocols. Amplification is performed in 35 cycles. Each cycle consists of incubation for 2 minutes at 94°C for denaturation of the template, 2 minutes at 50°C for hybridization of the oligonucleotide and 3 minutes at 72°C for extension of the chain. The amplified fragment is digested with SacI and XbaI and purified using 2% agarose gel. The fragment generated by PCR and digested by the two enzymes is ligated into the vector pHen $\Delta$ hIL-6 digested with the same two enzymes, purified on 0.8% agarose gel, to replace the wild type sequence.

The following Table 2 shows the biological activity, in human hepatoma cells, and binding to the receptor for wild interleukin-6 and versions thereof with mutations in the residues indicated.

TABLE 2

Receptor binding properties and biological activity of wild-type interleukin 6 and helix C mutations thereof

118	121	Biological activity	Receptor binding
Ser	Val	100%	100%
Arg	Val	66%	81%
Leu	Asp	36■4%	92%
Arg	Asp	3.5■0.5%	66■5%
Ser	Asp	58■23%	78■2%

It can be seen that the mutant IL-6 Ser118Arg/Val121Asp has characteristics very similar to those of the mutant IL-6 Tyr31Asp/Gly35Phe, described in Table 1, that is to say it binds normally to the

- 15 -

type I receptor of interleukin 6, but cytokin biological activity is reduced approximately thirty-fold.

#### Example 4

#### Generation and selection of more powerful antagonists of interleukin 6 using the methodology according to the present invention

In this case, the plasmid pHEN  $\Delta$  hIL-6 Tyr31Asp/Gly35Phe, obtained as described in Example 2, was used as a template for all the mutagenic reactions.

A PCR strategy (Polymerase Chain Reaction) was used to generate mutations within the codons selected for the area coding for human interleukin 6. The primer above is HP/1, described in the previous example. The mutagenetic primer below is IL-6 118RCLF/121VD, also described in the previous example. A DNA fragment of 415 pairs of bases is amplified using PCR according to standard PCR amplification protocols. Amplification is performed in 35 cycles. Each cycle consists of incubation for 2 minutes at 94°C for denaturation of the template, 2 minutes at 50°C for hybridization of the oligonucleotide and 3 minutes at 72°C for extension of the chain. The amplified fragment is digested with SacI and XbaI and purified using 2% agarose gel. The fragment generated by PCR and digested by the two enzymes is ligated into the vector pHen hIL-6 digested with the same two enzymes, purified on 0.8% agarose gel, to replace the wild type sequence.

The following Table 3 shows the biological activity, in human hepatoma cells, and binding to the receptor for wild interleukin-6 and versions thereof with mutations in the residues indicated.

TABLE 3

Receptor binding properties and biological activity of wild interleukin 6 and helix A and C mutants thereof

- 16 -

Helix A		Helix C		Biological activity	Receptor binding
31	35	118	121		
Tyr	Gly	Ser	Val	100%	100%
Asp	Phe	Leu	Val	1.4%	N.D.
Asp	Phe	Arg	Val	5.4■1.1%	66■2%
Asp	Phe	Leu	Asp	0%	63■4%
Asp	Phe	Arg	Asp	0%	97■15%
Asp	Phe	Phe	Asp	0%	76■26%

N.D. = Not determined

Three of the variants containing mutations both on helix A and on helix C show no sign of biological activity in human hepatoma cells, whereas they maintain their ability to bind the receptor gp80. Among these three proteins, the mutant IL-6 Tyr31Asp/Gly35Phe/Ser118Arg/Val121Asp was chosen for competition experiments with wild type interleukin 6 biological activity in human hepatoma cells. The cells were stimulated with wild interleukin 6 at 4 nanograms per milliliter (ng/ml) of culture medium, in the presence of increasing concentrations of mutant. As illustrated in figure 2 (in which biological activity of wild type interleukin 6 is expressed in arbitrary units), increasing concentrations of the mutant manage to fully antagonize the effects of wild interleukin 6 on human hepatoma cells.

Table 4 shows the levels of inhibition (in percentage) of the biological activity of wild interleukin 6 according to the concentration of antagonist (expressed in nanograms per milliliter, molar excess compared with wild type interleukin 6 and nanomoles per liter, respectively).

- 17 -

TABLE 4

Inhibition of the biological activity of wild interleukin 6 according to the concentration of antagonist

IL-6 Tyr31Asp/Gly35Phe/Ser118Arg/Val121Asp g/ml	molar excess	concentration (nM)	Inhibition of wild IL-6 (%)
16	4 X	0.7 nM	23%
60	15 X	2.7 nM	28%
164	41 X	7.5 nM	50%
500	125 X	22.7 nM	75%
1000	250 X	45.5 nM	82%
2000	500 X	90.0 nM	90%
4000	1000 X	181.8 nM	96%

Example 5

The interleukin 6 antagonist, generated and selected using the methodology described in the present invention, inhibit interleukin 6-dependent growth of human myeloma cells.

In the previous example it was shown that one of the mutants, namely IL-6 Tyr31Asp/Gly35Phe/Ser118Arg/Val121Asp was able to inhibit interleukin 6 biological activity (stimulation of transcription by an interleukin 6 inducible promoter) on human hepatoma cells. In the introductory part of this patent application it is stated that development of interleukin 6 antagonists or superantagonists would have practical application because they could be used to inhibit interleukin 6 activity in diseases characterized by interleukin 6 overproduction, like various forms of multiple myeloma/plasmacytoma. We provide here a further example by showing that the three mutants

IL-6 Tyr31Asp/Gly35Phe/Ser118Arg/Val121Asp (DFRD),

IL-6 Tyr31Asp/Gly35Phe/Ser118Phe/Val121Asp (DFFD),

**2142973**

- 18 -

IL-6 Tyr31Asp/Gly35Phe/Ser118Leu/Val121Asp (DFLD) fully inhibit the interleukin 6-dependent growth of a human myeloma cell line, called XG-1, derived from freshly isolated myeloma cells from a patient with terminal disease. The XG-1 myeloma cell line growth is strictly dependent on exogenously added interleukin 6, similarly to what has been shown for fresh myeloma cells, therefore this cell line can be considered an excellent in vitro model of the multiple myeloma disease (Jourdan, M., Zhang, X-G., Portier, M., Boiron, J.-M., Bataille, R. and Klein, B. (1991) *J. Immunol.* 147, 4402-4407). To test mutants antagonism on wild type interleukin 6, XG-1 myeloma cells were cultured in 96-well microtiter plates at 6000 cell/microwell with wild type interleukin 6 at 0.1 nanograms per milliliter (ng/ml) of culture medium, in the presence of increasing concentrations of each one of the three mutants. After 7 days of culture, cell numbers were evaluated by colorimetric determination of hexosaminidase levels (Landegren, U. (1984) *J. Immunol. Methods* 67, 379-388). The following Table 5 shows the inhibition of wild type interleukin 6 activity as a function of the three antagonist concentrations (expressed in nanograms of mutant per milliliter of culture medium).

- 19 -

TABLE 5

Inhibition of wild type interleukin 6 activity (stimulation of growth of XG-1 human myeloma cells) as a function of the three antagonists concentration.

Antagonist	Inhibition of wild type interleukin 6 produced by concentration		
	DFRD	DFFD	DFLD
3.3 ng/ml	6%	0%	0%
10 ng/ml	12%	7%	3%
30 ng/ml	25%	18%	10%
90 ng/ml	42%	35%	31%
270 ng/ml	64%	59%	53%
810 ng/ml	84%	84%	71%
2430 ng/ml	90%	90%	76%
7290 ng/ml	93%	95%	81%

As can be seen from the table, all mutants are particularly effective antagonists of interleukin 6 biological activity on human myeloma cells.

#### Example 6

Application of the methodology according to the present invention for the selection of novel superagonists of interleukin 6

A phage library (containing mutations of residues Gln 175, Ser 177, Leu 181 and Leu 183 of wild type interleukin 6, present in the form of fusion product with filamentous phage protein) was constructed using the Primer Extension molecular biology technique. The mutagenetic oligonucleotide is IL-6 QSLQ (AS), a 62 nucleotides oligo, whose sequence is SEQ ID NO:4. Primer IL-6 QSLQ (AS) extends from position 507 to the stop codon of the interleukin 6 cDNA (antisense strand), it introduces degenerations into codons coding for the amino acids 175 (wild type Gln), 177 (wild type Ser), 181 (wild type Leu) and 183 (wild type Gln) and it also introduces a NotI restriction site downstream

**2142973**

- 20 -

of the interleukin 6 stop codon. Oligonucleotide IL-6 QSLQ pr. Bam, whose sequence is SEQ ID NO:5, was used as primer for the Primer Extension reaction. Oligonucleotide IL-6 QSLQ pr. Bam extends from position 503 to position 522 (sense strand) of the interleukin 6 cDNA and it contains a BamHI recognition site within the 5' nine nucleotides. The two oligonucleotides are complementary to each other on a region corresponding to position 507 through position 522 of the interleukin 6 cDNA. The two oligonucleotides were annealed in vitro, and the annealed oligonucleotides were used as substrate for a Primer Extension reaction, performed using the Klenow enzyme. The double-stranded DNA fragment thus obtained was then digested with BamHI (compatible with BglII) and with NotI and ligated into the plasmid phen $\Delta$ hIL-6, digested with BglII (compatible with BamHI) and with NotI, in order to replace the wild type sequence with the mutated ones. The ligation product was inserted in bacteria, yielding roughly one million independent transformants ("trasformant" is the definition given to a bacterium which has incorporated a recombinant plasmid). The transformed bacteria were infected by with the M13K07 helper bacteriophage to generate the phage library (a library of phasmids) as described in the example 1.

The library underwent selection by incubation with polystyrene balls coated with shrIL-6R, as described in the example 1. The phage population eluted at pH 3.6 was then amplified in bacteria. After five cycles of selection-amplification, randomly selected phages were sequenced over the mutagenized region, the corresponding mutant interleukin 6 proteins were produced in the periplasmic space of the appropriate bacterial strain (as described in the example 1) and tested for both receptor binding and biological activity on human hepatoma cells. Table 6 below shows that, by using using the methodology to the present invention, it is

- 21 -

possible to select superagonists of interleukin 6, mutant molecules which have increased both receptor binding and biological activity on human hepatoma cells.

TABLE 6

Receptor binding properties and biological activities of wild type intyerleukin 6 and of its mutants in the helix D

Position	175	177	181	183	Receptor binding (%)	Biological activity (%)
wild type	Gln	Ser	Leu	Gln	100%	100%
phage 5-4	Gln	Ser	Leu	Tyr	240%	130%
phage 5-8	Gln	Ser	Leu	Ala	240%	120%
phage 5-2	Ile	Ser	Leu	Ala	260%	150%
phage 4-8	Gln	Ser	Ile	Asn	100%	N. D.
phage 5-3	Ile	Ser	Val	His	80%	N. D.

N. D.: not determined

The mutations selected by the methodology according to the present invention can be used as starting point for the development of more potent interleukin 6 superagonists. This is shown in this example, in which the mutation identified in the phage 5-2 are combined with the mutation Ser176Arg, which by prior art is known to increase both receptor binding and biological activity (see International Publication WO 94/11402 of a PCT application of the present Applicant, filed 2.11.93 with Italian priority of 6.11.92). The three mutations were grouped on the same cDNA by mean of Oligonucleotide Directed Mutagenesis. Oligonucleotide 175I/176R/183A (S), whose sequence is SEQ ID NO:6 is 73 nucleotides long which extends from position 499 to the stop codon (sense strand) of the interleukin 6 cDNA present in phen hIL-6 and which has a recognition site for the enzyme NotI downstream of

2142973

- 22 -

the stop codon. Oligonucleotide 175I/176R/183A (AS), whose sequence is SEQ ID NO. . is 73 nucleotides long which extends from position 499 to the stop codon (antisense strand) of the interleukin 6 cDNA present in phen $\Delta$ hIL-6 and which has a recognition site for the enzyme NotI upstream of the stop codon. The two oligonucleotide are complementary to each other and they both encode the amino acid Isoleucine in position 175, the amino amino acid Serine in position 176 and the amino acid Alanine in position 183. The two oligonucleotides were annealed in vitro, the double-stranded DNA fragment thus obtained was digested with the restriction enzymes BglII and Not I and ligated into the vector phen $\Delta$ hIL-6 digested with the same two enzymes, in order to replace the wild type sequence with the mutated one. The corresponding interleukin 6 variant carrying the three desired mutations was produced in the periplasmic space of the appropriate bacterial strain (as described in the example 1) and tested for both receptor binding and biological activity on human hepatoma cells. The following Table 7 shows the receptor binding properties and the biological activity on human hepatoma cells of both the new triple mutant and of the parental double mutant.

TABLE 7

Receptor binding properties and biological activities of wild type intyerleukin 6 and of both double and triple mutants in helix D

Position	175	176	183	Receptor binding (%)	Biological activity (%)
wild type	Gln	Ser	Gln	100%	100%
phage 5-2	Ile	Ser	Ala	260%	150%
triple mutant	Ile	Arg	Ala	450%	260%

As can be seen from the table, the mutant carrying

- 23 -

the substitutions Gln175Ile/Ser176Arg/Gln183Ala is a much more effective interleukin 6 superagonist than the parental mutant carrying only the two substitutions Gln175Ile/Gln183Ala.

#### Example 7

#### Application of the methodology according to the present invention for the selection of superantagonists of interleukin 6

The three mutations Gln175Ile/Ser176Arg/Gln183Ala, identified in the example 6, which strongly increases the receptor binding capacity, were combined with the four mutations Tyr31Asp/Gly35Phe/Ser118Arg/Val121Asp, described in examples 4 and 5, which show the strongest antagonistic behaviour, by mean of a PCR strategy. The corresponding mutant protein, called SAnt 1 and containing all the seven mutations, was tested both for receptor binding and for antagonistic behaviour on human hepatoma and myeloma cells. the following table 8 shows the receptor binding properties of both SAnt 1 and DFRD (the mutant from which SAnt 1 was derived), together with the amounts (in nanograms of mutant per milliliter of colture medium) of mutant necessary to inhibit 50% of interleukin 6 biological activity (hepatoma cells were stimulated with 4 nanograms of wild type interleukin 6 per milliliter of colture medium, while myeloma cells were stimulated with 0.1 nanograms of interleukin 6 per milliliter of colture medium, due to the higher sensitivity of the latter cells to wild type interleukin 6).

2142973

- 24 -

TABLE 8

Inhibition of wild type interleukin 6 activity on both human hepatoma and myeloma cells as a function of the antagonists receptor binding capacity.

---

Antagonist	Receptor binding (% of wt)	50% inhibition of interleukin 6 activity on		
		hepatoma cells Hep3B	hepatoma cells HepG2	myeloma cells XG-1
DFRD	97%	164 ng/ml	132 ng/ml	190 ng/ml
SAnt1	406%	19 ng/ml	32 ng/ml	22 ng/ml

---

As can be seen from the table, the introduction of the three mutations described in the example 6 (Gln175Ile/Ser176Arg/Gln183Ala) has at once increased the receptor binding capacity of the parental mutant DFRD and strongly decreased the amount of antagonist needed to inhibit 50% of interleukin 6 biological activity on all cell lines tested, therefore generating a very effective interleukin 6 superantagonist.

## SEQUENCE LISTING

## GENERAL INFORMATION

- (i) APPLICANT: ISTITUTO DI RICERCHE DI BIOLOGIA MOLECOLARE P. ANGELETTI S.p.A.
- (ii) TITLE OF INVENTION: METHOD FOR SELECTING SUPERAGONISTS, ANTAGONISTS AND SUPERANTAGONISTS FOR HORMONES HAVING GP 130 AS PART OF THEIR RECEPTOR COMPLEX
- (iii) NUMBER OF SEQUENCES: 14
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Società Italiana Brevetti
  - (B) STREET: Piazza di Pietra, 39
  - (C) CITY: Rome
  - (D) COUNTRY: Italy
  - (E) POSTAL CODE: I-00186
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk 3.5" 1.44 MBYTES
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS Rev. 5.0
  - (D) SOFTWARE: Microsoft Wordstar 4.0
- (viii) ATTORNEY INFORMATION
  - (A) NAME: DI CERBO, Mario (Dr.)
  - (C) REFERENCE: RM/X88217/PC-DC
- (ix) TELECOMMUNICATION INFORMATION
  - (A) TELEPHONE: 06/6785941
  - (B) TELEFAX: 06/6794692
  - (C) TELEX: 612287 ROPAT
- (1) INFORMATION FOR SEQ ID NO: 1:
  - (i) SEQUENCE CHARACTERISTICS
    - (A) LENGTH: 555 base pairs
    - (B) TYPE: nucleic acid
    - (C) STRANDEDNESS: double
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: cDNA
  - (iii) HYPOTHETICAL: no
  - (iv) ANTISENSE: no
  - (v) FRAGMENT TYPE: internal

## (vii) IMMEDIATE SOURCE:

(A) SYNTHESIS: production in bacteria

## (ix) FEATURE:

(A) NAME: IL-6 cDNA

(C) IDENTIFICATION METHOD: polyacrylamide gel

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CCA GTA CCC CCA GGA GAA GAT TCC AAA GAT GTA GCC GCC CCA CAC AGA 48  
 Pro Val Pro Pro Gly Glu Asp Ser Lys Asp Val Ala Ala Pro His Arg  
 1                   5                   10                   15  
 CAG CCA CTC ACG AGC TCA GAA CGA ATT GAC AAA CAA ATT CGG TAC ATC 96  
 Gln Pro Leu Thr Ser Ser Glu Arg Ile Asp Lys Gln Ile Arg Tyr Ile  
                   20                   25                   30  
 CTC GAC GGC ATC TCA GCC TTA AGA AAG GAG ACA TGT AAC AAG AGT AAC 144  
 Leu Asp Gly Ile Ser Ala Leu Arg Lys Glu Thr Cys Asn Lys Ser Asn  
                   35                   40                   45  
 ATG TGT GAA AGC AGC AAA GAG GCA CTG GCA GAA AAC AAC CTG AAC CTT 192  
 Met Cys Glu Ser Ser Lys Glu Ala Leu Ala Glu Asn Asn Leu Asn Leu  
                   50                   55                   60  
 CCA AAG ATG GCT GAA AAA GAT GGA TGC TTC CAA TCT GGA TTC AAT GAG 240  
 Pro Lys Met Ala Glu Lys Asp Gly Cys Phe Gln Ser Gly Phe Asn Glu  
 65                   70                   75                   80  
 GAG ACT TGC CTG GTG AAA ATC ATC ACT GGT CTT TTG GAG TTT GAG GTA 288  
 Glu Thr Cys Leu Val Lys Ile Ile Thr Gly Leu Leu Glu Phe Glu Val  
                   85                   90                   95  
 TAC CTA GAG TAC CTC CAG AAC AGA TTT GAG AGT AGT GAG GAA CAA GCC 336  
 Tyr Leu Glu Tyr Leu Gln Asn Arg Phe Glu Ser Ser Glu Glu Gln Ala  
                   100                   105                   110  
 AGA GCT GTC CAG ATG AGT ACA AAA GTC CTG ATC CAG TTC CTG CAG AAA 384  
 Arg Ala Val Gln Met Ser Thr Lys Val Leu Ile Gln Phe Leu Gln Lys  
                   115                   120                   125  
 AAG GCA AAG AAT CTA GAT GCA ATA ACC ACC CCT GAC CCA ACC ACA AAT 432  
 Lys Ala Lys Asn Leu Asp Ala Ile Thr Thr Pro Asp Pro Thr Thr Asn  
                   130                   135                   140  
 GCC AGC CTG CTG ACG AAG CTG CAG GCA CAG AAC CAG TGG CTG CAG GAC 480  
 Ala Ser Leu Leu Thr Lys Leu Gln Ala Gln Asn Gln Trp Leu Gln Asp  
 145                   150                   155                   160

ATG ACA ACT CAT CTC ATT CTG AGA TCT TTT AAG GAG TTC CTG CAG TCC 528  
 Met Thr Thr His Leu Ile Leu Arg Ser Phe Lys Glu Phe Leu Gln Ser

165

170

175

AGC CTG AGG GCT CTT CGG CAA ATG TAG

555

Ser Leu Arg Ala Leu Arg Gln Met

180

## (2) INFORMATION FOR SEQ ID NO: 2:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 70 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

(iii) HYPOTHETICAL: no

(iv) ANTISENSE: no

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE:

(A) SYNTHESIS: oligonucleotide synthesizer

(ix) FEATURE:

(A) NAME: IL-6 31D35 PCR

(C) IDENTIFICATION METHOD: polyacrylamide gel

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CGCACGAGCT CAGAACGAAT TGACAAACAA ATTCGGKACA TCCTCGACYD TATCTCAGCC 60

TTAAGAAAGG

70

## (3) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 72 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

(iii) HYPOTHETICAL: no

(iv) ANTISENSE: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE:

(A) SYNTHESIS: oligonucleotidic synthesizer

(ix) FEATURE:

(A) NAME: IL-6 118RCLF/121VD

(C) IDENTIFICATION METHOD: polyacrylamide gel

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CGCATCTAGA TTCTTTGCCT TTTTCTGCAG GAACTGGATC AGGWCTTTTG TGMRCATCTG 60  
CACAGCTCTG GC 72

(4) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 62 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

(iii) HYPOTHETICAL: no

(iv) ANTISENSE: yes

(v) FRAGMENT TYPE: carboxi-terminal

(vii) IMMEDIATE SOURCE:

(A) SYNTHESIS: oligonucleotide synthesizer

(ix) FEATURE:

(A) NAME : IL-6 QSLQ (AS)

(C) IDENTIFICATION METHOD: polyacrylamide gel

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CCGGGCGGCC GCCCTACATM NNCCGMNAG CCCTCAGMNN GGAMNNCAGG AACTCCTTAA 60  
AG 62

(5) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

(iii) HYPOTHETICAL: no

(iv) ANTISENSE: no

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE:

(A) SYNTHESIS: oligonucleotide synthesizer

(ix) FEATURE:

(A) NAME : IL-6 QSLQ pr. Bam

(C) IDENTIFICATION METHOD: polyacrylamide gel

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

CGCGGATCCT TTAAGGAGTT CCTG 24

(6) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 73 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

(iii) HYPOTHETICAL: no

(iv) ANTISENSE: no

(v) FRAGMENT TYPE: carboxi-terminal

(vii) IMMEDIATE SOURCE:

(A) SYNTHESIS: oligonucleotide synthesizer

(ix) FEATURE:

(A) NAME : 175I/167R/183A (S)

(C) IDENTIFICATION METHOD: polyacrylamide gel

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

GCCTGAGATC TTTTAAGGAG TTCCTGATCC GTAGCCTGAG GGCTCTTCGG GCTATGTAGG 60

GCGGCCGCAT GGC 73

(7) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 73 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single stranded

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

(iii) HYPOTHETICAL: no

(iv) ANTISENSE: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) SYNTHESIS: oligonucleotide synthesizer

(ix) FEATURE:

(A) NAME : 175I/167R/183A (AS)

(C) IDENTIFICATION METHOD: polyacrylamide gel



(D) OTHER INFORMATION: sequence of a mutant form of interleukin 6, from position 27 to position 42

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Ala Gln Ile Arg Tyr Ile Leu Asp Gln Thr Ser Ala Leu Arg Lys Glu  
1                   5                   10                   15

(10) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: no

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE:

(A) SYNTHESIS: production in bacteria as recombinant protein

(ix) FEATURE:

(A) NAME : Tyr31Asp, Gly35Phe

(C) IDENTIFICATION METHOD: polyacrylamide gel

(D) OTHER INFORMATION: sequence of a mutant form of interleukin 6, from position 27 to position 42

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Lys Gln Ile Arg Asp Ile Leu Asp Tyr Ile Ser Ala Leu Arg Lys Glu  
1                   5                   10                   15

(11) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: no

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE:

(A) SYNTHESIS: production in bacteria as recombinant protein

## (ix) FEATURE:

- (A) NAME : Tyr31Asp, Gly35Phe
- (C) IDENTIFICATION METHOD: polyacrylamide gel
- (D) OTHER INFORMATION: sequence of a mutant form of interleukin 6, from position 27 to position 42

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

Lys Gln Ile Arg Asp Ile Leu Asp Phe Ile Ser Ala Leu Arg Lys Glu  
 1                    5                    10                    15

## (12) INFORMATION FOR SEQ ID NO: 12:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: no

(v) FRAGMENT TYPE: internal

## (vii) IMMEDIATE SOURCE:

(A) SYNTHESIS: production in bacteria as recombinant protein

## (ix) FEATURE:

- (A) NAME : Tyr31Asp, Gly35Leu, Glu42Ala
- (C) IDENTIFICATION METHOD: polyacrylamide gel
- (D) OTHER INFORMATION: sequence of a mutant form of interleukin 6, from position 27 to position 42

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

Lys Gln Ile Arg Asp Ile Leu Asp Leu Ile Ser Ala Leu Arg Lys Ala  
 1                    5                    10                    15

## (13) INFORMATION FOR SEQ ID NO: 13:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: no

(v) FRAGMENT TYPE: internal

## (vii) IMMEDIATE SOURCE:

(A) SYNTHESIS: production in bacteria as recombinant protein

(ix) FEATURE:

(A) NAME : Tyr31Asp, Gly35His

(C) IDENTIFICATION METHOD: polyacrylamide gel

(D) OTHER INFORMATION: sequence of a mutant form of interleukin 6, from position 27 to position 42

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

Lys Gln Ile Arg Asp Ile Leu Asp His Ile Ser Ala Leu Arg Lys Glu  
 1                    5                    10                    15

(14) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: no

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE:

(A) SYNTHESIS: production in bacteria as recombinant protein

(ix) FEATURE:

(A) NAME : Tyr31Asp, Gly35Cys

(C) IDENTIFICATION METHOD: polyacrylamide gel

(D) OTHER INFORMATION: sequence of a mutant form of interleukin 6, from position 27 to position 42

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

Lys Gln Ile Arg Asp Ile Leu Asp Cys Ile Ser Ala Leu Arg Lys Glu  
 1                    5                    10                    15

- 30 -

WE CLAIM:

1. A methodology for selecting superagonists, antagonists and superantagonists of interleukin 6 using a membrane molecule gp 130 to activate the mechanisms regulating cell physiology, comprising the following operations:

- comparing amino acid sequences of a growth hormone with amino acid sequences of interleukin 6;
- comparing amino acid sequences of a growth hormone receptor with amino acid sequences of two receptors of interleukin 6, that is with an interleukin 6-specific receptor and gp 130;
- on the basis of the above comparisons, formulating a three-dimensional model of an interleukin 6 receptor complex based on the functional similarity between sequences on the growth hormone receptor and the two receptors of interleukin 6;
- identifying the residues of wild type interleukin 6 that form a part of a site of interaction with the interleukin 6-specific receptor and a site of interaction with gp 130, respectively;
- introducing a mutation to residues of interleukin 6 that form a part of the sites of interaction; and
- evaluating biological activity of resulting mutants.

2. The methodology for selecting superagonists of interleukin 6 according to claim 1, further comprising the following additional operations:

- production of a series of phage libraries containing mutations of the following wild type residues of interleukin 6 present as a fusion product with filamentous phage proteins, Glu 42, Glu 51, Ser 52, Ser 53, Lys 54, Glu 55, Asn 63, Lys 66, Met 67, Ala 68, Glu 69, Lys 70, Asp 71, Phe 170, Gln 175, Ser 176, Ser 177, Leu 181, Gln 183;
- generation of a phage library, each phage having a mutant interleukin 6 sequence;
- selection, from a mixed population of phages expressing interleukin 6 mutants, of that or those with an affinity for the interleukin 6-specific receptor greater than that of wild type interleukin 6; and

- 31 -

- identification of the best amino acid sequence or sequences binding the interleukin 6-specific receptor by sequencing of DNA extracted from the selected phage particles.

3. The methodology for selecting superagonists of interleukin 6 according to claim 2, in which a series of phage libraries are produced, containing mutants of said wild type residues of interleukin 6 present as a product of fusion with protein pIII of M13.

4. The methodology for selecting antagonists of interleukin 6 according to claim 1, further comprising the following additional operations:

- mutation of residues of interleukin 6 identified to form part of the site of interaction with gp 130 selected from: Arg 30, Tyr 31, Gly 35, Ser 37, Ala 38, Ser 118, Lys 120, Val 121, Gln 124, Phe 125, Gln 127, Lys 128 and Lys 129, using conventional molecular biology techniques;

- evaluation of biological activity and affinity with the interleukin 6-specific receptor of resulting mutants, in order to identify variants of interleukin 6 whose affinity to the interleukin 6-specific receptor is intact and that show reduction or loss of the biological activity; and

- evaluation of the variants of interleukin 6 as antagonists for biological activity of wild type interleukin 6.

5. The methodology for selecting superantagonists of interleukin 6 according to claim 1 by combination of the variations of amino acid sequences responsible for antagonist activity with amino-acid variations responsible for an increased affinity for the interleukin 6-specific receptor.

6. A methodology for selecting antagonists or superantagonists of interleukin 6 according to claim 5, in which the mutagenesis of the residues is performed using a molecular biology technique chosen from Polymerase Chain Reaction, Primer Extension, Oligonucleotide Directed Mutagenesis, or combinations thereof.

# pHenΔ hIL6 AND PRODUCTION OF PHAGEMID PARTICLES




FIG.1




FIG.2