US 20040086884A1

(19) United States

(12) Patent Application Publication (10) Pub. No.: US 2004/0086884 A1

Beach et al. (43) Pub. Date: May 6, 2004

(54) METHODS AND COMPOSITIONS FOR RNA INTERFERENCE

(75) Inventors: **David H. Beach**, Boston, MA (US); **Emily Bernstein**, Huntington, NY (US); **Amy Caudy**, St. Louis, NY (US); **Scott Hammond**, Pittsboro, NC (US); **Gregory J. Hannon**, Huntington, NY (US); **Patrick J. Paddison**, Cold Spring Harbor, NY (US); **Douglas Conklin**, Centerport, NY (US)

Correspondence Address:
ROPES & GRAY LLP
ONE INTERNATIONAL PLACE
BOSTON, MA 02110-2624 (US)

(73) Assignees: **Genetica, Inc.**, Cambridge, MA (US); **Cold Spring Harbor Laboratory**, Cold Spring Harbor, NY (US)

(21) Appl. No.: **10/350,798**

(22) Filed: **Jan. 24, 2003**

Related U.S. Application Data

(63) Continuation-in-part of application No. 10/055,797, filed on Jan. 22, 2002, which is a continuation-in-part of application No. PCT/US01/08435, filed on Mar. 16, 2001.
Continuation-in-part of application No. 09/866,557, filed on May 24, 2001.
Continuation-in-part of application No. 09/858,862, filed on May 16, 2001.

(60) Provisional application No. 60/243,097, filed on Oct. 24, 2000. Provisional application No. 60/189,739, filed on Mar. 16, 2000.

Publication Classification

(51) **Int. Cl.**[7] **C12Q 1/68**; C12N 15/85

(52) **U.S. Cl.** **435/6**; 435/455

(57) ABSTRACT

The present invention provides methods for attenuating gene expression in a cell using gene-targeted double stranded RNA (dsRNA). The dsRNA contains a nucleotide sequence that hybridizes under physiologic conditions of the cell to the nucleotide sequence of at least a portion of the gene to be inhibited (the "target" gene).




Fig. 1A




Fig. 1B




Fig. 1C




Fig. 2A




Fig. 2B




Fig. 2C




Fig. 2D




Fig. 3



Fig. 4A




Fig. 4B




Fig. 5A




Fig. 5B




Fig. 5C




Fig. 6A




Fig. 6C




Fig. 6B




Fig. 6D




Fig. 6E




Fig. 6F




Fig. 7A Fig. 7B




Fig. 7C



R3
R3
dsRBD
zap
helicase

Fig. 8A



D.m. Dicer
helicase
ZAP
RIII a
RIII b dsrm
1
606
1099 1246
1742
2239
C.e. Dicer
helicase
ZAP
RIII a
RIII b dsrm
1
498
759 915
1325
1806
H.s. Dicer
helicase
ZAP
RIII a
RIII b dsrm
1
606
1099 1044
1315
1914
A.t. CAF
helicase
ZAP
RIII a
RIII b dsrm dsrm
1
767
1180 1318
1375
1904
S.p. Dicer
helicase
ZAP
RIII a
RIII b
1
467
930
1233
RDE-1/Qde-2 family
ZAP
PIWI

Fig. 8B



CELL FREE EXTRACT
200K SPIN
RIBOSOME PELLET
HIGH SALT EXTRACTION
RIBOSOME ASSOCIATED PROTEINS
LOW SALT PRECIPITATION
PELLET
RNAi ACTIVITY
SUPERNATANT
RESOLUBILIZE HIGH SALT
CHROMATOGRAPHY
SUPEROSE 6
MONO S
MONO Q
HYDROXYAPATITE

Fig. 9



size std, kD
669
440
232 158
51
216
129
91
Fraction 24
Luciferase transcript
Cyclin E (control) transcript
dArgo-2 immunoblot.

Fig. 10



Fraction 5
24
Luciferase transcript
dArgo-2 immunoblot
216
129
91

Fig. 11



Fraction 5
24
Luciferase transcript
dArgo-2 immunoblot
216
129
91

Fig. 12




Fig. 13



At-argonaute-1
Hs-eIF2C
Nc-qde-2
Ce-rde-1
Dm-piwi
Dm-sting
Dm-argonaute-1
Dm-argonaute-2
PAZ
Piwi

Fig. 14




Fig. 15




Fig. 16



S10
S100
S10
S100
luc
cyclin E
S10
S100
mRNA degradation
dsRNA processing

Fig. 17




Fig. 18



Resource Phenyl
Q-sepharose
HAP
Superose
S-sepharose
Purification of the 22-mer generating enzyme

Fig. 19




Fig. 20A




Fig. 20C




Fig. 20B




Fig. 21




Fig. 22A




Fig. 22B




Fig. 23

MGKKDKNKKGGQDSAAAPQPQQQQKQQQQRQQQPQQLQQPQQLQQPQQLQQPQQQQQQ
QPHQQQQQSSRQQPSTSSGGSRASGFQQGGQQQKSQDAEGWTAQKKQGKQQVQGWTKQ
GQQGGHQQGRQGQDGGYQQRPPGQQQGGHQQGRQGQEGGYQQRPPGQQQGGHQQGRQG
QEGGYQQRPSGQQQGGHQQGRQGQEGGYQQRPPGQQQGGHQQGRQGQEGGYQQRPSGQ
QQGGHQQGRQGQEGGYQQRPSGQQQGGHQQGRQGQEGGYQQRPSGQQQGGHQQGRQGQ
EGGYQQRPPGQQPNQTQSQGQYQSRGPPQQQQAAPLPLPPQPAGSIKRGTIGKPGQVG
INYLDLDLSKMPSVAYHYDVKIMPERPKKFYRQAFEQFRVDQLGGAVLAYDGKASCYS
VDKLPLNSQNPEVTVTDRNGRTLRYTIEIK**ETGDSTIDLK**SLTTYMNDR**IFDKPMR**AM
QCVEVVLASPCHNKAIRVGR**SFFK**MSDPNNRHELDDGYEALVGLYQAFMLGDRPFLNV
DISHKSFPISMPMIEYLERFSLKAK**INNTTNLDYSR**RFLEPFLRGINVVYTPPQSFQS
APRVYRVNGLSR**APASSETFEHDGK**KVTIASYFHSRNYPLKFPQLHCLNVGSSIKSIL
LPIELCSIEEGQALNRKDGATQVANMIKYAATSTNVRKRKIMNLLQYFQHNLDPTISR
FGIRIANDFIVVSTRVLSPPQVEYHSKRFTMVKNGSWRMDGMK**FLEPKPK**AHKCAVLY
CDPRSGRKMNYTQLNDFGNLIISQGKAVNISLDSDVTYRPFTDDERSLDTIFADLKRS
QHDLAIVIIPQFRISYDTIKQKAELQHGILTQCIKQFTVERKCNNQTIGNILLKINSK
LNGINHKIKDDPRLPMMKNTMYIGADVTHPSPDQREIPSVVGVAASHDPYGASYNMQY
RLQRGALEEIEDMFSITLEHLRVYKEYRNAYPDHIIYYRDGVSDGQFPKIKNEELRCI
KQACDKVGCKPKICCVIVVKRHHTRFFPSGDVTTSNKFNNVDPGTVVDRTIVHPNEMQ
FFMVSHQAIQGTAKPTRYNVIENTGNLDIDLLQQLTYNLCHMFPRCNRSVSYPAPAYL
AHLVAARGR**VYLTGTNR**FLDLKKEYAKRTIVPEFMKKNPMYFV

Fig. 24



PREDICTED INTRON
REPEATS
PAZ
Piwi
S2 genomic
S2 cDNA
Library clone #7
Argo-2/plZ
No template
Embryo
Adult
S2

Fig. 25



Fig. 26



STRATEGY FOR STABLE EXPRESSION OF dsRNA IN CULTURED MAMMALIAN CELLS
CRE RECOMBINASE
IN VIVO EXPRESSION OF RNA HAIRPIN
PRODUCTION OF dsDNA HOMOLOGOUS TO TARGET mRNA

Fig. 27




Fig. 28




Fig. 29A




Fig. 29B




Fig. 29C




Fig. 29D



STABLE SUPPRESSION OF TRANSGENE EXPRESSION
IN MAMMALIAN CELLS
RFP
GFP
RFP
GFP
wt P19
PUTATIVE P19 CLONES STABLY
EXPRESSING 500mer GFP HAIRPIN
GFP
GFP
CO-TRANSFECTION WITH pRFP And pGFP, 42 HRS POST-TRANSFECTION

Fig. 30



RATIO TARGETED Luc: CONTROL Luc
1.2
1
0.8
0.6
0.4
0.2
0
FF ds500
Ren ds500
GFP ds500

Fig. 31




Fig. 32A




Fig. 32B




Fig. 32C



DUAL LUCIFERASE ASSAY 21 HRS POST-TRANSFECTION
(.4ug dsRNA)
FF-LUC/REN-LUC
VALUES NORMALIZED
TO dsGFP CONTROL
1
0.9
0.8
0.7
0.6
0.5
0.4
0.3
0.2
0.1
0
5x
7x
dsLUC
dsGFP
Hela
CHO
P19

Fig. 33




Fig. 34A





Fig. 34B



P19 GFP HAIRPIN CLONE NUMBER #10
48 HRS POST-TRANSFECTION
FLUORESCENT MICROSCOPY SUPERIMPOSED WITH BRIGHT FIELD

Fig. 34C




Fig. 34D



dsFF
dsGFP
RATIO FF:REN
1.4
1.2
1
0.8
0.6
0.4
0.2
0
0 µg
.01 µg
.1 µg
1 µg
µg dsRNA ADDED TO EXTRACT

Fig. 35



SILENCING IS SPECIFIC AND REQUIRES dsRNA
RATIO FF/REN
1.6
1.4
1.2
1
0.8
0.6
0.4
0.2
0
dsFF
ssFF
asFF
dsGFP
ssGFP
asGFP
RATIO REN/FF
1.8
1.6
1.4
1.2
1
0.8
0.6
0.4
0.2
0
dsGFP
ssGFP
asGFP
dsREN
ssREN
asREN

Fig. 36



P19 CELLS SOAKED WITH IN dsRNA FOR
12 HRS IN 2mL GROWTH MEDIUM (ALPHA MEM, 10% FBS)
dsLUC
dsGFP
FF-LUC/REN-LUC
VALUES NORMALIZED
TO 0 MICROG CONTROL
1.2
1
0.8
0.6
0.4
0.2
0
1
5
20
40
MICROGRAMS OF dsRNA

Fig. 37



IN VITRO SYNTHESIS OF siRNAs by T7 RNA POLYMERASE
a.
T7
b.
T7
TRANSCRIBE/ANNEAL
TREAT CELLS
ASSAY PHENOTYPE
DNA SYNTHESIS/RNA TRANSCRIPTION
~$16/siRNA VERSUS~$400/siRNA FOR CHEMICAL SYNTHESIS
BRINGS LARGE-SCALE PROJECTS WITHIN REASONABLE BUDGET RANGE

Fig. 38

siRNA

```
  UCGAAGUACUCAGCGUAAGUG
AAAGCUUCAUGAGUCGCAUUC
```

cshFf

```
                                  U
CAUCGACUGAAAUCCCUGGUAAUCCGUUG U
GUAGCUGACUUUAGGGACCAUUAGGCAAC A
                                  A
```

cshFf-L7

```
                                  ----------      U
CAUCGACUGAAAUCCCUGGUAAUCCGUUU            GGGGC \
GUAGCUGAUUUUAGGGACUAUUAGGUAAA            UCCCG C
                                  UAGGGUAUCG      U
```

cshFf-L7m

```
                  GCC             ----------      U
CAUCGACUGAAAUCCC    GUAAUCCGUUU           GGGGC \
GUAGCUGAUUUUAGGG    UAUUAGGUAAA           UCCCG C
                  AC-             UAGGGUAUCG      U
```

Fig. 39A




Fig. 39B




Fig. 39C



293T
HeLa
Cos-1
NIH 3T3
1.2
1
0.8
0.6
0.4
0.2
0
GFP
siRNA
cshFf
cshFf-L7
cshFf-L7m

Fig. 40

siRNA

UCGAAGUACUCAGCGUAAGUG
AAAGCUUCAUGAGUCGCAUUC

T7siRNA

GGUCGAAGUACUCAGCGUAAGAA
AAAGCUUCATGAGUCGCAUUCGG

T7siFf-2

GGUUGUGGAUCUGGAUACCGG
UUCCAACACCUAGACCUAUGG

T7siFf-3

GGUGCCAACCCUAUUCUCCUU
GACCACGGUUGGGAUAAGAGG

T7siFf-8

GGCUAUGAAGAGAGUACGCCCU
UUCCGAUACUUCUCUCAUGCGG

Fig. 41A




Fig. 41B

T7shFf29

```
GGU|                              U
   CGAAGUACUCAGCGUAAGUGAUGUCCAC U
   GUUUUGUGGGUUGUGUUUGUUGUGGGUG A
  G^                               A
```

T7shFf27

```
GGU|                            U
   CGAAGUACUCAGCGUAAGUGAUGUCC U
   GUUUUGUGGGUUGUGUUUGUUGUGGG A
  G^                             A
```

T7shFf25

```
GGU|                          U
   CGAAGUACUCAGCGUAAGUGAUGU U
   GUUUUGUGGGUUGUGUUUGUUGUG A
  G^                           A
```

T7shFf22

```
GGU|                        U
   CGAAGUACUCAGCGUAAGUGA U
   GUUUUGUGGGUUGUGUUUGUU A
  G^                       A
```

T7shFf29-5'T

```
GGCUCGAGU|                                U
         CGAAGUACUCAGCGUAAGUGAUGUCCAC U
         GUUUUGUGGGUUGUGUUUGUUGUGGGUG A
G--------^                                A
```

T7shFf29-3'T

```
-----G|                                 U
      GUCGAAGUACUCAGCGUAAGUGAUGUCCAC U
      CGGUUUUGUGGGUUGUGUUUGUUGUGGGUG A
 GAGCU^                                 A
```

Fig. 41C




Fig. 41D




Fig. 42A




Fig. 42B




Fig. 42C




Fig. 43




Fig. 44A




Fig. 44B



SIMULTANEOUS INTRODUCTION OF MULTIPLE HAIRPINS DOES NOT PRODUCE SYNERGY
SERIES 1
FF/RENILLA RATIC
50
45
40
35
30
25
20
15
10
5
0
CONTROL
CONTROL
siRNA
HP #1
HP #2
MIX

Fig. 45



ENCODED SHORT HAIRPINS FUNCTION *in vivo*
29 bp
PLASMID
~70 bp
LUCIFERASE ACTIVITY (%RENILLA)
3
2.5
2
1.5
1
0.5
0
siRNA
Luc SIMPLE
U6 VECTOR
siOligo1-2
siOligo1-6
siOligo1-10
siOligo1-19
siOligo2-23
siOligo2-36

Fig. 46



STABLE SUPPRESSION BY SHORT dsRNAs - STABLE EXPRESSION STRATEGIES
T7 GIVES SITE-SPECIFIC INITIATION. 3' END FORMATION ACHIEVED WITH RIBOZYME (E.G. HEPATITIS DELTA VIRUS RIBOZYME).
POLIII GIVES SITE-SPECIFIC INITIATION.
EXAMPLE PROMOTERS-U6 snRNA, H1 RNA, SRP RNAs (7SL)
3' END FORMATION
ACHIEVED WITH NATIVE TERMINATOR (e.g. TTTTT). LEAVES THE LAST TT, SO THAT COULD BE USED TO PAIR TO TRANSCRIPT.
COULD ALSO USE VA1, tRNA ETC. BUT WOULD HAVE TO COUPLE WITH RIBOZYME SINCE THOSE PROMOTERS NEED ALSO INTERNAL ELEMENTS.
POLIII GIVES SITE-SPECIFIC INITIATION
EXAMPLE PROMOTERS WOULD BE U1 snRNA, PROMOTERS, CMV ETC...
3' END FORMATION ACHIEVED WITH RIBOZYME
(e.g. HEPATITIS DELTA VIRUS RIBOZYME).

Fig. 47



STABLE SUPPRESSION BY SHORT dsRNAs - CLONING STRATEGY
PCR-1
+
PCR-2
AUTOMATIC SUBCLONING INTO VECTOR OF CHOICE

Fig. 48




Fig. 49



EARLY PASSAGE PKR -/- MEFs: DUAL LUCIFERASE ASSAY WITH LONG dsRNA
(~500nt)
RATION REN-FF LUCIFERASE
1.4
1.2
1
0.8
0.6
0.4
0.2
0
12HRS
42HRS
82HRS
HOURS POST-TRANSFECTION
dsGFP
dsREN

Fig. 50



MOUSE TYROSINASE PROMOTER
Enhancer
~12 kb
-1404 to -1007
-1003 to -506
-505 to -85
+1

Fig. 51



pGL3-CONTROL:
SV40
polyA SITE
FIREFLY LUCIFERASE
pLuc-NS5B:
SV40
polyA SITE
FIREFLY LUCIFERASE
HCV NS5B
LUCIFERASE siRNA:
5'-CUUACGCUGAGUACUUCGATT-3'
3'-TTGAAUGCGACUCAUGAAGCU-5'
UNRELATED siRNA:
5'-AGCUUCAUAAGGCGCAUGCTT-3'
3'-TTUCGAAGUAUUCCGCGUACG-5'
NS5B siRNA:
5'-CUGUGAGAUCUACGGAGCCUGTT-3'
3'-TTGACACUCUAGAUGCCUCGGAC-5'
pShh1-Ff1:
5' ---|
GAA
GGAUUCCAAUUCAGCGGGAGCCACCUGAU G
CCUAAGGUUGAGUCGCUCUCGGUGGGCUA C
3'-UUA^
GUU

Fig. 52




Fig. 53A




Fig. 53B




Fig. 54A




Fig. 54B




Fig. 55A




Fig. 55B




Fig. 55C




Fig. 56A




Fig. 56B

METHODS AND COMPOSITIONS FOR RNA INTERFERENCE

RELATED APPLICATIONS

[0001] This application is a continuation-in-part of U.S. application Ser. No. 10/055,797, filed Jan. 22, 2002, which is a continuation-in-part of International Application PCT/US01/08435, filed Mar. 16, 2001, which claims the benefit of priority from U.S. Provisional Application Nos. 60/189,739, filed Mar. 16, 2000, and 60/243,097, filed Oct. 24, 2000. This application is also a continuation-in-part of U.S. application Ser. No. 09/866,557 filed May 24, 2001 and a continuation-in-part of U.S. application Ser. No. 09/858,862 filed May 16, 2001. The specifications of such applications are incorporated by reference herein. International Application PCT/US01/08435 was published under PCT Article 21(2) in English.

GOVERNMENT SUPPORT

[0002] Work described herein was supported by National Institutes of Health Grant R01-GM62534. The United States Government may have certain rights in the invention.

BACKGROUND OF THE INVENTION

[0003] "RNA interference", "post-transcriptional gene silencing", "quelling"—these different names describe similar effects that result from the overexpression or misexpression of transgenes, or from the deliberate introduction of double-stranded RNA into cells (reviewed in Fire A (1999) Trends Genet 15: 358-363; Sharp P A (1999) *Genes Dev* 13: 139-141; Hunter C (1999) *Curr Biol* 9: R440-R442; Baulcombe D C (1999) *Curr Biol* 9: R599-R601; Vaucheret et al. (1998) *Plant J* 16: 651-659). The injection of double-stranded RNA into the nematode *Caenorhabditis elegans*, for example, acts systemically to cause the post-transcriptional depletion of the homologous endogenous RNA (Fire et al. (1998) *Nature* 391: 806-811; and Montgomery et al. (1998) *PNAS* 95: 15502-15507). RNA interference, commonly referred to as RNAi, offers a way of specifically and potently inactivating a cloned gene, and is proving a powerful tool for investigating gene function. Although the phenomenon is interesting in its own right; the mechanism has been rather mysterious, but recent research—for example that recently reported by Smardon et al. (2000) *Curr Biol* 10: 169-178—is beginning to shed light on the nature and evolution of the biological processes that underlie RNAi.

[0004] RNAi was discovered when researchers attempting to use the antisense RNA approach to inactivate a *C. elegans* gene found that injection of sense-strand RNA was actually as effective as the antisense RNA at inhibiting gene function (Guo et al. (1995) *Cell* 81: 611-620). Further investigation revealed that the active agent was modest amounts of double-stranded RNA that contaminate in vitro RNA preparations. Researchers quickly determined the 'rules' and effects of RNAi which have become the paradigm for thinking about the mechanism which mediates this affect. Exon sequences are required, whereas introns and promoter sequences, while ineffective, do not appear to compromise RNAi (though there may be gene-specific exceptions to this rule). RNAi acts systemically—injection into one tissue inhibits gene function in cells throughout the animal. The results of a variety of experiments, in *C. elegans* and other organisms, indicate that RNAi acts to destabilize cellular RNA after RNA processing.

[0005] The potency of RNAi inspired Timmons and Fire (1998 *Nature* 395: 854) to do a simple experiment that produced an astonishing result. They fed to nematodes bacteria that had been engineered to express double-stranded RNA corresponding to the *C. elegans* unc-22 gene. Amazingly, these nematodes developed a phenotype similar to that of unc-22 mutants that was dependent on their food source. The ability to conditionally expose large numbers of nematodes to gene-specific double-stranded RNA formed the basis for a very powerful screen to select for RNAi-defective *C. elegans* mutants and then to identify the corresponding genes.

[0006] Double-stranded RNAs (dsRNAs) can provoke gene silencing in numerous in vivo contexts including Drosophila, *Caenorhabditis elegans*, planaria, hydra, trypanosomes, fungi and plants. However, the ability to recapitulate this phenomenon in higher eukaryotes, particularly mammalian cells, has not been accomplished in the art. Nor has the prior art demonstrated that this phenomena can be observed in cultured eukaryotic cells. Additionally, the 'rules' established by the prior art have taught that RNAi requires exon sequences, and thus constructs consisting of intronic or promoter sequences were not believed to be effective reagents in mediating RNAi. The present invention aims to address each of these deficiencies in the prior art and provides evidence both that RNAi can be observed in cultured eukaryotic cells and that RNAi constructs consisting of non-exon sequences can effectively repress gene expression.

SUMMARY OF THE INVENTION

[0007] One aspect of the present invention provides a method for attenuating expression of a target gene in cultured cells, comprising introducing double stranded RNA (dsRNA) into the cells in an amount sufficient to attenuate expression of the target gene, wherein the dsRNA comprises a nucleotide sequence that hybridizes under stringent conditions to a nucleotide sequence of the target gene.

[0008] Another aspect of the present invention provides a method for attenuating expression of a target gene in a mammalian cell, comprising

[0009] (i) activating one or both of a Dicer activity or an Argonaut activity in the cell, and

[0010] (ii) introducing into the cell a double stranded RNA (dsRNA) in an amount sufficient to attenuate expression of the target gene, wherein the dsRNA comprises a nucleotide sequence that hybridizes under stringent conditions to a nucleotide sequence of the target gene.

[0011] In certain embodiments, the cell is suspended in culture; while in other embodiments the cell is in a whole animal, such as a non-human mammal.

[0012] In certain preferred embodiments, the cell is engineered with (i) a recombinant gene encoding a Dicer activity, (ii) a recombinant gene encoding an Argonaut activity, or (iii) both. For instance, the recombinant gene may encode, for a example, a protein which includes an amino acid

sequence at least 50 percent identical to SEQ ID No. 2 or 4; or be defined by a coding sequence which hybridizes under wash conditions of 2×SSC at 22° C. to SEQ ID No. 1 or 3. In certain embodiments, the recombinant gene may encode, for a example, a protein which includes an amino acid sequence at least 50 percent identical to the Argonaut sequence shown in **FIG. 24**. In certain embodiments, the recombinant gene may encode a protein which includes an amino acid sequence at least 60 percent, 70 percent, 80 percent, 85 percent, 90 percent, or 95 percent identical to SEQ ID No. 2 or 4. In certain embodiments, the recombinant gene may be defined by a coding sequence which hybridizes under stringent conditions, including a wash step selected from 0.2×-2.0×SSC at from 50° C.-65° C., to SEQ ID No. 1 or 3.

[**0013**] In certain embodiments, rather than use a heterologous expression construct(s), an endogenous Dicer gene or Argonaut gene can be activated, e.g., by gene activation technology, expression of activated transcription factors or other signal transduction protein(s), which induces expression of the gene, or by treatment with an endogenous factor which upregulates the level of expression of the protein or inhibits the degradation of the protein.

[**0014**] In certain preferred embodiments, the target gene is an endogenous gene of the cell. In other embodiments, the target gene is a heterologous gene relative to the genome of the cell, such as a pathogen gene, e.g., a viral gene.

[**0015**] In certain embodiments, the cell is treated with an agent that inhibits protein kinase RNA-activated (PKR) apoptosis, such as by treatment with agents which inhibit expression of PKR, cause its destruction, and/or inhibit the kinase activity of PKR.

[**0016**] In certain preferred embodiments, the cell is a primate cell, such as a human cell.

[**0017**] In certain preferred embodiments, the length of the dsRNA is at least 20, 21 or 22 nucleotides in length, e.g., corresponding in size to RNA products produced by Dicer-dependent cleavage. In certain embodiments, the dsRNA construct is at least 25, 50, 100, 200, 300 or 400 bases. In certain embodiments, the dsRNA construct is 400-800 bases in length.

[**0018**] In certain preferred embodiments, expression of the target gene is attenuated by at least 5 fold, and more preferably at least 10, 20 or even 50 fold, e.g., relative to the untreated cell or a cell treated with a dsRNA construct which does not correspond to the target gene.

[**0019**] Yet another aspect of the present invention provides a method for attenuating expression of a target gene in cultured cells, comprising introducing an expression vector having a "coding sequence" which, when transcribed, produces double stranded RNA (dsRNA) in the cell in an amount sufficient to attenuate expression of the target gene, wherein the dsRNA comprises a nucleotide sequence that hybridizes under stringent conditions to a nucleotide sequence of the target gene. In certain embodiments, the vector includes a single coding sequence for the dsRNA which is operably linked to (two) transcriptional regulatory sequences which cause transcription in both directions to form complementary transcripts of the coding sequence. In other embodiments, the vector includes two coding sequences which, respectively, give rise to the two complementary sequences which form the dsRNA when annealed. In still other embodiments, the vector includes a coding sequence which forms a hairpin. In certain embodiments, the vectors are episomal, e.g., and transfection is transient. In other embodiments, the vectors are chromosomally integrated, e.g., to produce a stably transfected cell line. Preferred vectors for forming such stable cell lines are described in U.S. Pat. No. 6,025,192 and PCT publication WO/9812339, which are incorporated by reference herein.

[**0020**] Another aspect of the present invention provides a method for attenuating expression of a target gene in cultured cells, comprising introducing an expression vector having a "non-coding sequence" which, when transcribed, produces double stranded RNA (dsRNA) in the cell in an amount sufficient to attenuate expression of the target gene. The non-coding sequence may include intronic or promoter sequence of the target gene of interest, and the dsRNA comprises a nucleotide sequence that hybridizes under stringent conditions to a nucleotide sequence of the promoter or intron of the target gene. In certain embodiments, the vector includes a single sequence for the dsRNA which is operably linked to (two) transcriptional regulatory sequences which cause transcription in both directions to form complementary transcripts of the sequence. In other embodiments, the vector includes two sequences which, respectively, give rise to the two complementary sequences which form the dsRNA when annealed. In still other embodiments, the vector includes a coding sequence which forms a hairpin. In certain embodiments, the vectors are episomal, e.g., and transfection is transient. In other embodiments, the vectors are chromosomally integrated, e.g., to produce a stably transfected cell line. Preferred vectors for forming such stable cell lines are described in U.S. Pat. No. 6,025,192 and PCT publication WO/9812339, which are incorporated by reference herein.

[**0021**] Another aspect the present invention provides a double stranded (ds) RNA for inhibiting expression of a mammalian gene. The dsRNA comprises a first nucleotide sequence that hybridizes under stringent conditions, including a wash step of 0.2×SSC at 65° C., to a nucleotide sequence of at least one mammalian gene and a second nucleotide sequence which is complementary to the first nucleotide sequence.

[**0022**] In one embodiment, the first nucleotide sequence of said double-stranded RNA is at least 20, 21, 22, 25, 50, 100, 200, 300, 400, 500, 800 nucleotides in length.

[**0023**] In another embodiment, the first nucleotide sequence of said double-stranded RNA is identical to at least one mammalian gene. In another embodiment, the first nucleotide sequence of said double-stranded RNA is identical to one mammalian gene. In yet another embodiment, the first nucleotide sequence of said double-stranded RNA hybridizes under stringent conditions to at least one human gene. In still another embodiment, the first nucleotide sequence of said double-stranded RNA is identical to at least one human gene. In still another embodiment, the first nucleotide sequence of said double-stranded RNA is identical to one human gene.

[**0024**] The double-stranded RNA may be an siRNA or a hairpin, and may be expressed transiently or stably. In one embodiment, the double-stranded RNA is a hairpin comprising a first nucleotide sequence that hybridizes under

stringent conditions to a nucleotide sequence of at least one mammalian gene, and a second nucleotide sequence which is a complementary inverted repeat of said first nucleotide sequence and hybridizes to said first nucleotide sequence to form a hairpin structure.

[0025] The first nucleotide sequence of said double-stranded RNA can hybridize to either coding or non-coding sequence of at least one mammalian gene. In one embodiment, the first nucleotide sequence of said double-stranded RNA hybridizes to a coding sequence of at least one mammalian gene. In another embodiment, the first nucleotide sequence of said double-stranded RNA hybridizes to a coding sequence of at least one human gene. In another embodiment, the first nucleotide sequence of said double-stranded RNA is identical to a coding sequence of at least one mammalian gene. In still another embodiment, the first nucleotide sequence of said double-stranded RNA is identical to a coding sequence of at least one human gene.

[0026] In another embodiment, the first nucleotide sequence of said double-stranded RNA hybridizes to a non-coding sequence of at least one mammalian gene. In another enbodiment, the first nucleotide sequence of said double-stranded RNA hybridizes to a non-coding sequence of at least one human gene. In another embodiment, the first nucleotide sequence of said double-stranded RNA is identical to a non-coding sequence of at least one mammalian gene. In still another embodiment, the first nucleotide sequence of said double-stranded RNA is identical to a non-coding sequence of at least one human gene. In any of the foregoing embodiments, the non-coding sequence may be a non-transcribed sequence.

[0027] Still another aspect of the present invention provides an assay for identifying nucleic acid sequences, either coding or non-coding sequences, responsible for conferring a particular phenotype in a cell, comprising

[0028] (i) constructing a variegated library of nucleic acid sequences from a cell in an orientation relative to a promoter to produce double stranded DNA;

[0029] (ii) introducing the variegated dsRNA library into a culture of target cells;

[0030] (iii) identifying members of the library which confer a particular phenotype on the cell, and identifying the sequence from a cell which correspond, such as being identical or homologous, to the library member.

[0031] Yet another aspect of the present invention provides a method of conducting a drug discovery business comprising:

[0032] (i) identifying, by the subject assay, a target gene which provides a phenotypically desirable response when inhibited by RNAi;

[0033] (ii) identifying agents by their ability to inhibit expression of the target gene or the activity of an expression product of the target gene;

[0034] (iii) conducting therapeutic profiling of agents identified in step (b), or further analogs thereof, for efficacy and toxicity in animals; and

[0035] (iv) formulating a pharmaceutical preparation including one or more agents identified in step (iii) as having an acceptable therapeutic profile.

[0036] The method may include an additional step of establishing a distribution system for distributing the pharmaceutical preparation for sale, and may optionally include establishing a sales group for marketing the pharmaceutical preparation.

[0037] Another aspect of the present invention provides a method of conducting a target discovery business comprising:

[0038] (i) identifying, by the subject assay, a target gene which provides a phenotypically desirable response when inhibited by RNAi;

[0039] (ii) (optionally) conducting therapeutic profiling of the target gene for efficacy and toxicity in animals; and

[0040] (iii) licensing, to a third party, the rights for further drug development of inhibitors of the target gene.

[0041] Another aspect of the invention provides a method for inhibiting RNAi by inhibiting the expression or activity of an RNAi enzyme. Thus, the subject method may include inhibiting the activity of Dicer and/or the 22-mer RNA.

[0042] Still another aspect relates to a method for altering the specificity of an RNAi by modifying the sequence of the RNA component of the RNAi enzyme.

[0043] In another aspect, gene expression in an undifferentiated stem cell, or the differentiated progeny thereof, is altered by introducing dsRNA of the present invention. In one embodiment, the stem cells are embryonic stem cells. Preferably, the embryonic stem cells are derived from mammals, more preferably from non-human primates, and most preferably from humans.

[0044] The embryonic stem cells may be isolated by methods known to one of skill in the art from the inner cell mass (ICM) of blastocyst stage embryos. In one embodiment the embryonic stem cells are obtained from previously established cell lines. In a second embodiment, the embryonic stem cells are derived de novo by standard methods.

[0045] In another aspect, the embryonic stem cells are the result of nuclear transfer. The donor nuclei are obtained from any adult, fetal, or embryonic tissue by methods well known in the art. In one embodiment, the donor nuclei is transferred to a recipient oocyte which had previously been modified. In one embodiment, the oocyte is modified using one or more dsRNAs. Exemplary modifications of the recipient oocyte include any changes in gene or protein expression that prevent an embryo derived from said modified oocyte from successfully implanting in the uterine wall. Since implantation in the uterine wall is essential for fertilized mammalian embryos to progress from beyond the blastocyst stage, embryos made from such modified oocytes could not give rise to viable organisms. Non-limiting examples of such modifications include those that decrease or eliminate expression of cell surface receptors (i.e., integrins) required for the recognition between the blastocyst and the uterine wall, modifications that decrease or eliminate expression of proteases (i.e., collagenase, stromelysin, and plasminogen activator) required to digest matrix in the uterine lining and thus allow proper implantation, and modifications that decrease or eliminate expression of proteases (i.e., strypsin)

necessary for the blastocyst to hatch from the zona pellucida. Such hatching is required for implantation.

[0046] In another embodiment, embryonic stem cells, embryonic stem cells obtained from fertilization of modified oocytes, or the differentiated progeny thereof, can be modified or further modified with one or more dsRNAs. In a preferred embodiment, the modification decreases or eliminates MHC expression. Cells modified in this way will be tolerated by the recipient, thus avoiding complications arising from graft rejection. Such modified cells are suitable for transplantation into a related or unrelated patient to treat a condition characterized by cell damage or cell loss.

[0047] In another aspect of the invention, the undifferentiated stem cell is an adult stem cell. Exemplary adult stem cells include, but are not limited to, hematopoietic stem cells, mesenchymal stem cells, cardiac stem cells, pancreatic stem cells, and neural stem cells. Exemplary adult stem cells include any stem cell capable of forming differentiated ectodermal, mesodermal, or endodermal derivatives. Non-limiting examples of differentiated cell types which arise from adult stem cells include: blood, skeletal muscle, myocardium, endocardium, pericardium, bone, cartilage, tendon, ligament, connective tissue, adipose tissue, liver, pancreas, skin, neural tissue, lung, small intestine, large intestine, gall bladder, rectum, anus, bladder, female or male reproductive tract, genitals, and the linings of the body cavity.

[0048] In one embodiment, an undifferentiated adult stem cell, or the differentiated progeny thereof, is altered with one or more dsRNAs to decrease or eliminate MHC expression. Cells modified in this way will be tolerated by the recipient, thus avoiding complications arising from graft rejection. Such modified cells are suitable for transplantation into a related or unrelated patient to treat a condition characterized by cell damage or cell loss.

[0049] In another aspect of the invention, an embryonic stem cell, an undifferentiated adult stem cell, or the differentiated progeny of either an embryonic or adult stem cell is altered with one or more dsRNA to decrease or eliminate expression of genes required for HIV infection. In a preferred embodiment, the stem cell is one capable of giving rise to hematopoietic cells. Modified cells with hematopoietic potential can be transplanted into a patient as a preventative therapy or treatment for HIV or AIDS.

[0050] Another aspect of the invention relates to purified or semi-purified preparations of the RNAi enzyme or components thereof. In certain embodiments, the preparations are used for identifying compounds, especially small organic molecules, which inhibit or potentiate the RNAi activity. Small molecule inhibitors, for example, can be used to inhibit dsRNA responses in cells which are purposefully being transfected with a virus which produces double stranded RNA.

[0051] The dsRNA construct may comprise one or more strands of polymerized ribonucleotide. It may include modifications to either the phosphate-sugar backbone or the nucleoside. The double-stranded structure may be formed by a single self-complementary RNA strand or two complementary RNA strands. RNA duplex formation may be initiated either inside or outside the cell. The dsRNA construct may be introduced in an amount which allows delivery of at least one copy per cell. Higher doses of double-stranded material may yield more effective inhibition. Inhibition is sequence-specific in that nucleotide sequences corresponding to the duplex region of the RNA are targeted for genetic inhibition. In certain embodiments, dsRNA constructs containing a nucleotide sequences identical to a portion of the target gene are preferred for inhibition. RNA sequences with insertions, deletions, and single point mutations relative to the target sequence (i.e., RNA sequences similar to the target sequence) have also been found to be effective for inhibition. Thus, sequence identity may be optimized by alignment algorithms known in the art and calculating the percent difference between the nucleotide sequences. Alternatively, the duplex region of the RNA may be defined functionally as a nucleotide sequence that is capable of hybridizing with a portion of the target gene transcript. In another embodiment, dsRNA constructs containing nucleotide sequences identical to a non-coding portion of the target gene are preferred for inhibition. Exemplary non-coding regions include introns and the promoter region. Sequences with insertions, deletions, and single point mutations relative to the target non-coding sequence may also be used.

[0052] Yet another aspect of the invention pertains to transgenic non-human mammals which include a transgene encoding a dsRNA construct, wherein the dsRNA is identical or similar to either the coding or non-coding sequence of the target gene, preferably which is stably integrated into the genome of cells in which it occurs. The animals can be derived by oocyte microinjection, for example, in which case all of the nucleated cells of the animal will include the transgene, or can be derived using embryonic stem (ES) cells which have been transfected with the transgene, in which case the animal is a chimera and only a portion of its nucleated cells will include the transgene. In certain instances, the sequence-independent dsRNA response, e.g., the PKR response, is also inhibited in those cells including the transgene.

[0053] In still other embodiments, dsRNA itself can be introduced into an ES cell in order to effect gene silencing, and that phenotype will be carried for at least several rounds of division, e.g., into the progeny of that cell.

BRIEF DESCRIPTION OF THE DRAWINGS

[0054] **FIG. 1**: RNAi in S2 cells. (a) Drosophila S2 cells were transfected with a plasmid that directs lacZ expression from the copia promoter in combination with dsRNAs corresponding to either human CD8 or lacZ, or with no dsRNA, as indicated. (b) S2 cells were co-transfected with a plasmid that directs expression of a GFP-US9 fusion protein and dsRNAs of either lacZ or cyclin E, as indicated. Upper panels show FACS profiles of the bulk population. Lower panels show FACS profiles from GFP-positive cells. (c) Total RNA was extracted from cells transfected with lacZ, cyclin E, fizzy or cyclin A dsRNAs, as indicated. Northern blots were hybridized with sequences not present in the transfected dsRNAs.

[0055] **FIG. 2**: RNAi in vitro. (a) Transcripts corresponding to either the first 600 nucleotides of Drosophila cyclin E (E600) or the first 800 nucleotides of lacZ (Z800) were incubated in lysates derived from cells that had been transfected with either lacZ or cyclin E (cycE) dsRNAs, as indicated. Time points were 0, 10, 20, 30, 40 and 60 min for cyclin E and 0, 10, 20, 30 and 60 min for lacZ. (b)

Transcripts were incubated in an extract of S2 cells that had been transfected with cyclin E dsRNA (cross-hatched box, below). Transcripts corresponded to the first 800 nucleotides of lacZ or the first 600, 300, 220 or 100 nucleotides of cyclin E, as indicated. Eout is a transcript derived from the portion of the cyclin E cDNA not contained within the transfected dsRNA. E-ds is identical to the dsRNA that had been transfected into S2 cells. Time points were 0 and 30 min. (c) Synthetic transcripts complementary to the complete cyclin E cDNA (Eas) or the final 600 nucleotides (Eas600) or 300 nucleotides (Eas300) were incubated in extract for 0 or 30 min.

[0056] **FIG. 3**: Substrate requirements of the RISC. Extracts were prepared from cells transfected with cyclin E dsRNA. Aliquots were incubated for 30 min at 30° C. before the addition of either the cyclin E (E600) or lacZ (Z800) substrate. Individual 20 μl aliquots, as indicated, were pre-incubated with 1 mM $CaCl_2$ and 5 mM EGTA, 1 mM $CaCl_2$, 5 mM EGTA and 60 U of micrococcal nuclease, 1 mM $CaCl_2$ and 60 U of micrococcal nuclease or 10 U of DNase I (Promega) and 5 mM EGTA. After the 30 min pre-incubation, EGTA was added to those samples that lacked it. Yeast tRNA (1 μg) was added to all samples. Time points were at 0 and 30 min.

[0057] **FIG. 4**: The RISC contains a potential guide RNA. (a) Northern blots of RNA from either a crude lysate or the S100 fraction (containing the soluble nuclease activity, see Methods) were hybridized to a riboprobe derived from the sense strand of the cyclin E mRNA. (b) Soluble cyclin-E-specific nuclease activity was fractionated as described in Methods. Fractions from the anion-exchange resin were incubated with the lacZ, control substrate (upper panel) or the cyclin E substrate (centre panel). Lower panel, RNA from each fraction was analysed by northern blotting with a uniformly labelled transcript derived from sense strand of the cyclin E cDNA. DNA oligonucleotides were used as size markers.

[0058] **FIG. 5**: Generation of 22mers and degradation of mRNA are carried out by distinct enzymatic complexes. (a) Extracts prepared either from 0-12 hour Drosophila embryos or Drosophila S2 cells (see Methods) were incubated for 0, 15, 30, or 60 minutes (left to right) with a uniformly-labeled double-stranded RNA corresponding to the first 500 nucleotides of the Drosophila cyclin E coding region. M indicates a marker prepared by in vitro transcription of a synthetic template. The template was designed to yield a 22 nucleotide transcript. The doublet most probably results from improper initiation at the +1 position. (b) Whole-cell extracts were prepared from S2 cells that had been transfected with a dsRNA corresponding to the first 500 nt. of the luciferase coding region. S10 extracts were spun at 30,000×g for 20 minutes which represents our standard RISC extract. S100 extracts were prepared by further centrifugation of S10 extracts for 60 minutes at 100,000×g. Assays for mRNA degradation were carried out as described previously for 0, 30 or 60 minutes (left to right in each set) with either a single-stranded luciferase mRNA or a single-stranded cyclin E mRNA, as indicated. (c) S10 or S100 extracts were incubated with cyclin E dsRNAs for 0, 60 or 120 minutes (L to R).

[0059] **FIG. 6**: Production of 22mers by recombinant CG4792/Dicer. (a) Drosophila S2 cells were transfected with plasmids that direct the expression of T7-epitope tagged versions of Drosha, CG4792/Dicer-1 and Homeless. Tagged proteins were purified from cell lysates by immunoprecipitation and were incubated with cyclin E dsRNA. For comparison, reactions were also performed in Drosophila embryo and S2 cell extracts. As a negative control, immunoprecipitates were prepared from cells transfected with a β-galactosidase expression vector. Pairs of lanes show reactions performed for 0 or 60 minutes. The synthetic marker (M) is as described in the legend to **FIG. 1**. (b) Diagrammatic representations of the domain structures of CG4792/Dicer-1, Drosha and Homeless are shown. (c) Immunoprecipitates were prepared from detergent lysates of S2 cells using an antiserum raised against the C-terminal 8 amino acids of Drosophila Dicer-1 (CG4792). As controls, similar preparations were made with a pre-immune serum and with an immune serum that had been pre-incubated with an excess of antigenic peptide. Cleavage reactions in which each of these precipitates was incubated with an ~500 nt. fragment of Drosophila cyclin E are shown. For comparsion, an incubation of the substrate in Drosophila embryo extract was electrophoresed in parallel. (d) Dicer immunoprecipitates were incubated with dsRNA substrates in the presence or absence of ATP. For comparison, the same substrate was incubated with S2 extracts that either contained added ATP or that were depleted of ATP using glucose and hexokinase (see methods). (e) Drosophila S2 cells were transfected with uniformly, ^{32}P-labelled dsRNA corresponding to the first 500 nt. of GFP. RISC complex was affinity purified using a histidine-tagged version of Drosophila Ago-2, a recently identified component of the RISC complex (Hammond et al., in prep). RISC was isolated either under conditions in which it remains ribosome associated (ls, low salt) or under conditions that extract it from the ribosome in a soluble form (hs, high salt). For comparison, the spectrum of labelled RNAs in the total lysate is shown. (f) Guide RNAs produced by incubation of dsRNA with a Dicer immunoprecipitate are compared to guide RNAs present in an affinity-purified RISC complex. These precisely comigrate on a gel that has single-nucleotide resolution. The lane labelled control is an affinity selection for RISC from a cell that had been transfected with labeled dsRNA but not with the epitope-tagged Drosophila Ago-2.

[0060] **FIG. 7**: Dicer participates in RNAi. (a) Drosophila S2 cells were transfected with dsRNAs corresponding to the two Drosophila Dicers (CG4792 and CG6493) or with a control dsRNA corresponding to murine caspase 9. Cytoplasmic extracts of these cells were tested for Dicer activity. Transfection with Dicer dsRNA reduced activity in lysates by 7.4-fold. (b) The Dicer-1 antiserum (CG4792) was used to prepare immunoprecipitates from S2 cells that had been treated as described above. Dicer dsRNA reduced the activity of Dicer-1 in this assay by 6.2-fold. (c) Cells that had been transfected two days previously with either mouse caspase 9 dsRNA or with Dicer dsRNA were cotransfected with a GFP expression plasmid and either control, luciferase dsRNA or GFP dsRNA. Three independent experiments were quantified by FACS. A comparison of the relative percentage of GFP-positive cells is shown for control (GFP plasmid plus luciferase dsRNA) or silenced (GFP plamsid plus GFP dsRNA) populations in cells that had previously been transfected with either control (caspase 9) or Dicer dsRNAs.

[0061] **FIG. 8**: Dicer is an evolutionarily conserved ribonuclease. (a) A model for production of 22mers by Dicer. Based upon the proposed mechanism of action of Ribonuclease III, we propose that Dicer acts on its substrate as a dimer. The positioning of the two ribonuclease domains (RIIIa and RIIIb) within the enzyme would thus determine the size of the cleavage product. An equally plausible alternative model could be derived in which the RIIIa and RIIIb domains of each Dicer enzyme would cleave in concert at a single position. In this model, the size of the cleavage product would be determined by interaction between two neighboring Dicer enzymes. (b) Comparison of the domain structures of potential Dicer homologs in various organisms (Drosophila—CG4792, CG6493, *C. elegans*—K12H4.8, Arabidopsis—CARPEL FACTORY, T25K16.4, AC012328_1, human Helicase-MOI and *S. pombe*—YC9A_SCHPO). The ZAP domains were identified both by analysis of individual sequences with Pfam and by Psi-blast searches. The ZAP domain in the putative *S. pombe* Dicer is not detected by PFAM but is identified by Psi-Blast and is thus shown in a different color. For comparison, a domain structure of the RDE1/QDE2/ARGONAUTE family is shown. It should be noted that the ZAP domains are more similar within each of the Dicer and ARGONAUTE families than they are between the two groups. (c) An alignment of the ZAP domains in selected Dicer and Argonaute family members is shown. The alignment was produced using ClustalW.

[0062] **FIG. 9**: Purification strategy for RISC. (second step in RNAi model).

[0063] **FIG. 10**: Fractionation of RISC activity over sizing column. Activity fractionates as 500 KDa complex. Also, antibody to Drosophila argonaute 2 cofractionates with activity.

[0064] FIGS. **11-13**: Fractionation of RISC over monoS, monoQ, Hydroxyapatite columns. Drosophila argonaute 2 protein also cofactionates.

[0065] **FIG. 14**: Alignment of Drosophila argonaute 2 with other family members.

[0066] **FIG. 15**: Confirmation of Drosophila argonaute 2. S2 cells were transfected with labeled dsRNA and His tagged argonaute. Argonaute was isolated on nickel agarose and RNA component was identified on 15% acrylamide gel.

[0067] **FIG. 16**: S2 cell and embryo extracts were assayed for 22mer generating activity.

[0068] **FIG. 17**: RISC can be separated from 22mer generating activity (dicer). Spinning extracts (S100) can clear RISC activity from supernatant (left panel) however, S100 spins still contain dicer activity (right panel).

[0069] **FIG. 18**: Dicer is specific for dsRNA and prefers longer substrates.

[0070] **FIG. 19**: Dicer was fractionated over several columns.

[0071] **FIG. 20**: Identification of dicer as enzyme which can process dsRNA into 22mers. Various RNaseIII family members were expressed with n terminal tags, immunoprecipitated, and assayed for 22mer generating activity (left panel). In right panel, antibodies to dicer could also precipitate 22mer generating activity.

[0072] **FIG. 21**: Dicer requires ATP.

[0073] **FIG. 22**: Dicer produces RNAs that are the same size as RNAs present in RISC.

[0074] **FIG. 23**: Human dicer homolog when expressed and immunoprecipitated has 22mer generating activity.

[0075] **FIG. 24**: Sequence of Drosophila argonaute 2. Peptides identified by microsequencing are shown in underline.

[0076] **FIG. 25**: Molecular charaterization of Drosophila argonaute 2. The presence of an intron in coding sequence was determined by northern blotting using intron probe. This results in a different 5' reading frame then the published genome seqeunce. Number of polyglutamine repeats was determined by genomic PCR.

[0077] **FIG. 26**: Dicer activity can be created in human cells by expression of human dicer gene. Host cell was 293. Crude extracts had dicer activity, while activity was absent from untransfected cells. Activity is not dissimilar to that seen in drosophila embryo extracts.

[0078] **FIG. 27**: A ~500 nt. fragment of the gene that is to be silenced (X) is inserted into the modified vector as a stable direct repeat using standard cloning procedures. Treatment with commercially available cre recombinase reverses sequences within the loxP sites (L) to create an inverted repeat. This can be stably maintained and amplified in an sbc mutant bacterial strain (DL759). Transcription in vivo from the promoter of choice (P) yields a hairpin RNA that causes silencing. A zeocin resistance marker is included to insure maintenance of the direct and inverted repeat structures; however this is non-essential in vivo and could be removed by pre-mRNA splicing if desired. (Smith et al. (2000) *Nature* 407: 319-20).

[0079] **FIG. 28**: RNAi in P19 embryonal carcinoma cells. Ten-centimeter plates of P19 cells were transfected by using 5 μg of GFP plasmid and 40 μg of the indicated dsRNA (or no RNA). Cells were photographed by fluorescent (tope panel) and phase-contrast microscopy (bottom panel) at 72 h after transfection; silencing was also clearly evident at 48 h posttransfection.

[0080] **FIG. 29**: RNAi of firefly and Renilla luciferase in P19 cells. (A and B) P19 cells were transfected with plasmids that direct the expression of firefly and Renilla luciferase and dsRNA 500 mers (25 or 250 ng, as indicated in A and B, respectively), that were either homologous to the firefly luciferase mRNA (dsFF) or nonhomologous (dsGFP). Luciferase activities were assayed at various times after transfection, as indicated. Ratios of firefly to Renilla activity are normalized to dsGFP controls. (C and D) P19 cells in 12-well culture dishes (2 ml of media) were transfected with 0.25 μg of a 9:1 mix of pGL3-Control and pRL-SV40 as well as 2 μg of the indicated RNA. Extracts were prepared 9 h after transfection. (C) Ratio of firefly to Renilla luciferase is shown. (D) Ratio of Renilla to firefly luciferase is shown. Values are normalized to dsGFP. The average of three independent experiments is shown; error bars indicate standard deviation.

[0081] **FIG. 30**: The panels at the right show expression of either RFP or GFP following transient transfection into wild type P19 cells. The panels at the left demonstrate the specific suppression of GFP expression in P19 clones which stably

express a 500 nt double stranded GFP hairpin. P19 clones which stably express the double stranded GFP hairpin were transiently transfected with RFP or GFP, and expression of RFP or GFP was assessed by visual inspection.

[0082] **FIG. 31**: Specific silencing of luciferase expression by dsRNA in murine embryonic stem cells. Mouse embryonic stem cells in 12-well culture dishes (1 ml of media) were transfected with 1.5 μg of dsRNA along with 0.25 μg of a 10:1 mixture of the reporter plasmids pGL3-Control and pRL-SV40. Extracts were prepared and assayed 20 h after transfection. The ratio of firefly to Renilla luciferase expression is shown for FF ds500; the ratio of Renilla to firefly is shown for Ren ds500. Both are normalized to ratios from the dsGFP transfection. The average of three independent experiments is shown; error bars indicate standard deviation.

[0083] **FIG. 32**: RNAi in C2C12 murine myoblast cells. (A) Mouse C2C12-cells in 12-well culture dishes (1 ml of media) were transfected with 1 μg of the indicated dsRNA along with 0.250 μg of the reporter plasmids pGL3-Control and pRL-SV40. Extracts were prepared and assayed 24 h after transfection. The ratio of firefly to Renilla luciferase expression is shown; values are normalized to ratios from the no dsRNA control. The average of three independent experiments is shown; error bars indicate standard deviation. (B) C2C12 cells cotransfected with 1 μg of either plasmid alone or a plasmid containing a hyperactive mutant of vaccinia virus K3L (Kawagishi-Kobayashi et al. 2000, Virology 276: 424-434). The absolute counts of Renilla and firefly luciferase activity are shown. (C) The ratios of firefly/Renilla activity from B, normalized to no dsRNA controls.

[0084] **FIG. 33**: Hela, Chinese hamster ovary, and P19 (pluripotent, mouse embryonic carcinoma) cell lines transfected with plasmids expressing *Photinus pyralis* (firefly) and *Renilla reniformis* (sea pansy) luciferases and with dsRNA 500mers (400 ng), homologous to either firefly luciferase mRNA (dsLUC) or non-homologous (dsGFP). Dual luciferase assays were carried out using an Analytical Scientific Instruments model 3010 Luminometer. In this assay Renilla luciferase serves as an internal control for dsRNA-specific suppression of firefly luciferase activity. These data demonstrate that 500mer dsRNA can specifically suppress cognate gene expression in vivo.

[0085] **FIG. 34**: Expression of a hairpin RNA produces P19 EC cell lines that stably silence GFP. (A) A cartoon of the FLIP cassette used to construct the GFP hairpin. GFP represents the first 500 coding base pairs of EGFP. Zeo, zeocin resistance gene; L, Lox; P, the cytomegalovirus promoter in the expression plasmid pcDNA3. Homologous GFP fragments are first cloned as direct repeats into the FLIP cassette. To create inverted repeats for hairpin production, the second repeat is flipped by using Cre recombinase. When transcribed, the inverted repeat forms a GFP dsRNA with a hairpin loop. (B) P19 cell lines stably expressing the GFP hairpin plasmid, GFPhp.1 (clone 10) and GFPhp.2 (clone 12), along with wt P19 were transfected with 0.25 μg each of GFP and RFP reporter genes. Fluorescence micrographs were taken by using filters appropriate for GFP and RFP. Magnification is 200×. (C) P19 GFPhp.1 cells were transfected with pEGFP and 0, 0.5, or 1 μg of Dicer or firefly dsRNA. Fluorescence micrographs were taken at 48 h posttransfection and are superimposed with bright field images to reveal non-GFP expressing cells. Magnification is 100×. (D) In vitro and in vivo processing of dsRNA in P19 cells. In vitro Dicer assays were performed on S2 cells and three independently prepared P19 extracts by using ^{32}P-labeled dsRNA (30° C. for 30 min). A Northern blot of RNA extracted from control and GFPhp.1 P19 cells shows the production of ≈22mer RNA species in hairpin-expressing cells but not in control cells. Blots were probed with a ^{32}P-labeled "sense" GFP transcript.

[0086] **FIG. 35**: dsRNA induces silencing at the posttranscriptional level. P19 cell extracts were used for in vitro translation of firefly and Renilla luciferase mRNA (100 ng each). Translation reactions were programmed with various amounts of dsRNA 500mers, either homologous to firefly luciferase mRNA (dsLUC) or nonhomologous (dsGFP). Luciferase assays were carried out after a 1 h incubation at 30° C. Ratios of firefly to Renilla activity are normalized to no dsRNA controls. Standard deviations from the mean are shown.

[0087] **FIG. 36**: S10 fractions from P19 cell lysates were used for in vitro translations of mRNA coding for *Photinus pyralis* (firefly) and *Renilla reniformis* (sea pansy) luciferases. Translation reactions were programmed with dsRNA, ssRNA, or asRNA 500mers, either complementary to firefly luciferase mRNA (dsFF, ssFF, or asFF), complementary to Renilla luciferase (dsREN, ssREN, or asREN) or non-complementary (dsGFP). Reactions were carried out at 30° C. for 1 hour, after a 30 min preincubation with dsRNA, ssRNA, or asRNA. Dual luciferase assays were carried out using an Analytical Scientific Instruments model 3010 Luminometer. On the left, Renilla luciferase serves as an internal control for dsRNA-specific suppression of firefly luciferase activity. On the right, firefly luciferase serves as an internal control for dsRNA-specific suppression of Renilla luciferase activity. These data demonstrate that 500mer double-stranded RNA (dsRNA) but not single-stranded (ssRNA) or anti-sense RNA (asRNA) suppresses cognate gene expression in vitro in a manner consistent with post-transcriptional gene silencing.

[0088] **FIG. 37**: P19 cells were grown in 6-well tissue culture plates to approximately 60% confluence. Various amounts of dsRNA, either homologous to firefly luciferase mRNA (dsLUC) or non-homologous (dsGFP), were added to each well and incubated for 12 hrs under normal tissue culture conditions. Cells were then transfected with plasmids expressing *Photinus pyralis* (firefly) and *Renilla reniformis* (sea pansy) luciferases and with dsRNA 500mers (500 ng). Dual luciferase assays were carried out 12 hrs post-transfection using an Analytical Scientific Instruments model 3010 Luminometer. In this assay Renilla luciferase serves as an internal control for dsRNA-specific suppression of firefly luciferase activity. These data show that 500mer dsRNA can specifically suppress cognate gene expression in vivo without transfection under normal tissue culture conditions.

[0089] **FIG. 38**: Previous methods for generating siRNAs required costly chemical synthesis. The invention provides an in vitro method for synthesizing siRNAs using standard RNA transcription reactions.

[0090] **FIG. 39**: Short hairpins suppress gene expression in Drosophila S2 cells. (A) Sequences and predicted secondary structure of representative chemically synthesized

RNAs. Sequences correspond to positions 112-134 (siRNA) and 463-491 (shRNAs) of Firefly luciferase carried on pGL3-Control. An siRNA targeted to position 463-485 of the luciferase sequence was virtually identical to the 112-134 siRNA in suppressing expression, but is not shown. (B) Exogenously supplied short hairpins suppress expression of the targeted Firefly luciferase gene in vivo. Six-well plates of S2 cells were transfected with 250 ng/well of plasmids that direct the expression of firefly and Renilla luciferase and 500 ng/well of the indicated RNA. Luciferase activities were assayed 48 h after transfection. Ratios of firefly to Renilla luciferase activity were normalized to a control transfected with an siRNA directed at the green fluorescent protein (GFP). The average of three independent experiments is shown; error bars indicate standard deviation. (C) Short hairpins are processed by the Drosophila Dicer enzyme. T7 transcribed hairpins shFfL22, shFfL29, and shFfS29 were incubated with (+) and without (−) 0-2-h Drosophila embryo extracts. Those incubated with extract produced ~22-nt siRNAs, consistent with the ability of these hairpins to induce RNA interference. A long dsRNA input (cyclin E 500-mer) was used as a control. Cleavage reactions were performed as described in Bernstein et al., 2001, Nature, 409:363-366.

[0091] **FIG. 40**: Short hairpins function in mammalian cells. HEK 293T, HeLa, COS-1, and NIH 3T3 cells were transfected with plasmids and RNAs as in **FIG. 1** and subjected to dual luciferase assays 48 h posttransfection. The ratios of firefly to Renilla luciferase activity are normalized to a control transfected with an siRNA directed at the green fluorescent protein (GFP). The average of three independent experiments is shown; error bars indicate standard deviation.

[0092] **FIG. 41**: siRNAs and short hairpins transcribed in vitro suppress gene expression in mammalian cells. (A) Sequences and predicted secondary structure of representative in vitro transcribed siRNAs. Sequences correspond to positions 112-134 (siRNA) and 463-491 (shRNAs) of firefly luciferase carried on pGL3-Control. (B) In vitro transcribed siRNAs suppress expression of the targeted firefly luciferase gene in vivo. HEK 293T cells were transfected with plasmids as in **FIG. 2**. The presence of non-base-paired guanosine residues at the 5' end of siRNAs significantly alters the predicted end structure and abolishes siRNA activity. (C) Sequences and predicted secondary structure of representative in vitro transcribed shRNAs. Sequences correspond to positions 112-141 of firefly luciferase carried on pGL3-Control. (D) Short hairpins transcribed in vitro suppress expression of the targeted firefly luciferase gene in vivo. HEK 293T cells were transfected with plasmids as in **FIG. 2**.

[0093] **FIG. 42**: Transcription of functional shRNAs in vivo. (A) Schematic of the pShh1 vector. Sequences encoding shRNAs with between 19 and 29 bases of homology to the targeted gene are synthesized as 60-75-bp double-stranded DNA oligonucleotides and ligated into an EcoRV site immediately downstream of the U6 promoter. (B) Sequence and predicted secondary structure of the Ff1 hairpin. (C) An shRNA expressed from the pShh1 vector suppresses luciferase expression in mammalian cells. HEK 293T, HeLa, COS-1, and NIH 3T3 cells were transfected with reporter plasmids as in **FIG. 1**, and pShh1 vector, firefly siRNA, or pshh1 firefly shRNA constructs as indicated. The ratios of firefly to Renilla luciferase activity were determined 48 h after transfection and represent the average of three independent experiments; error bars indicate standard deviation.

[0094] **FIG. 43**: Dicer is required for shRNA-mediated gene silencing. HEK 293T cells were transfected with luciferase reporter plasmids as well as pShh1-Ff1 and an siRNA targeting human Dicer either alone or in combination, as indicated. The Dicer siRNA sequence (TCA ACC AGC CAC TGC TGG A) corresponds to coordinates 3137-3155 of the human Dicer sequence. The ratios of firefly to Renilla luciferase activity were determined 26 h after transfection and represent the average of three independent experiments; error bars indicate standard deviation.

[0095] **FIG. 44**: Stable shRNA-mediated gene silencing of an endogenous gene. (A) Sequence and predicted secondary structure of the p53 hairpin. The 5' shRNA stem contains a 27-nt sequence derived from mouse p53 (nucleotides 166-192), whereas the 3' stem harbors the complimentary antisense sequence. (B) Senescence bypass in primary mouse embryo fibroblasts (MEFs) expressing an shRNA targeted at p53. Wild-type MEFs, passage 5, were transfected with pBabe-RasV12 with control plasmid or with p53hp (5 μg each with FuGENE; Roche). Two days after transfection, cells were trypsinized, counted, and plated at a density of 1×10^5/10-cm plate in media containing 2.0 μg/mL of puromycin. Control cells cease proliferation and show a senescent morphology (left panel). Cells expressing the p53 hairpin continue to grow (right panel). Photos were taken 14 d posttransfection.

[0096] **FIG. 45**: A mixture of two short hairpins, both corresponding to firefly luciferase, does not result in a synergistic suppression of gene expression. Suppression of firefly luciferase gene expression resulting from transfection of a mixture of two different short hairpins (HP #1 and HP #2) was examined. The mixture of HP #1 and HP #2 did not have a more robust effect on the suppression of firefly luciferase gene expression than expression of HP #1 alone.

[0097] **FIG. 46**: Encoded short hairpins specifically suppress gene expression in vivo. DNA oligonucleotides encoding 29 nucleotide hairpins corresponding to firefly luciferase were inserted into a vector containing the U6 promoter. Three independent constructs were examined for their ability to specifically suppress firefly luciferase gene expression in 293T cells. siOligo1-2, siOligo1-6, and siOligo1-19 (construct in the correct orientation) each suppressed gene expression as effectively as siRNA. In contrast, siOligo1-10 (construct in the incorrect orientation) did not suppress gene expression. An independent construct targeted to a different portion of the firefly luciferase gene did not effectively suppress gene expression in either orientation (siOligo2-23, siOligo2-36).

[0098] FIGS. **47-49**: Strategies for stable expression of short dsRNAs.

[0099] **FIG. 50**: Dual luciferase assays were performed as described in detail in FIGS. **28-35**, however the cells used in these experiments were PKR−/− murine embryonic fibroblasts (MEFs). Briefly, RNAi using long dsRNAs typically envokes a non-specific response in MEFs (due to PKR activity). To evaluate the effect of long dsRNA constructs to specifically inhibit gene expression in MEFs, RNAi was examined in PKR−/− MEFs. Such cells do not respond to

dsRNA with a non-specific response. The data summarized in this figure demonstrates that in the absence of the non-specific PKR response, long dsRNA constructs specifically suppress gene expression in MEFs.

[0100] **FIG. 51**: Is a schematic representation of the mouse tyrosinase promoter. Primers were used to amplify three separate regions in the proximal promoter, or to amplify sequence corresponding to an enhancer located approximately 12 kb upstream.

[0101] **FIG. 52**: Reporter expression plasmids and siRNA sequences used in Figures X and Y. PGL-3-Control and Pluc-NS5B are the expression plasmids used for transfection into mouse liver. The nucleotide sequences of the siRNAs used in the study are shown underneath.

[0102] **FIG. 53**: RNA interference in adult mice using siRNAs. (a) Representative images of light emitted from mice co-transfected with the luciferase plasmid pGL3-control and either no siRNA, luciferase siRNA or unrelated siRNA. A pseudocolour image representing intensity of emitted light (red, most intense; blue, least intense) superimposed on a greyscale reference image (for orientation) shows that RNAi functions in adult mice. Annealed 21-nucleotide siRNAs (40 μg; Dharmacon) were co-injected into the livers of mice with 2 μg pGL3-control DNA (Promega) and 800 units of RNasin (Promega) in 1.8 ml PBS buffer in 5-7 s. After 72 h, mice were anaesthetized and given 3 mg luciferin intraperitoneally 15 min before imaging. (b) siRNA results (six mice per group) from a representative experiment. Mice receiving luciferase siRNA emitted significantly less light than reporter-alone controls (one-way ANOVA with post hoc Fisher's test). Results for reporter alone and unrelated siRNA were statistically similar. Animals were treated according to the US National Institutes of Health's guidelines for animal care and the guidelines of Stanford University.

[0103] **FIG. 54**: RNA interference in adult mice using shRNAs. (a) Representative images of light emitted from mice co-transfected with the luciferase plasmid control, pShh1-Ff1, and pShh1-Ff1rev. pShh1-Ff1, but not pShh1-Ff1rev, reduced luciferase expression in mice relative to the reporter-alone control. pShh1-Ff1 or pShh1-rev (10 μg) were co-injected with 2 μg pGL3-control in 1.8 ml PBS buffer. (b) Average of three independent shRNA experiments (n=5). Average values for the reporter-alone group are designated as 100% in each of the three experiments. Animals were treated according to the US National Institutes of Health's guidelines for animal care and the guidelines of Stanford University.

[0104] **FIG. 55**: Heritable repression of Neil1 expression by RNAi in several tissues. (a) Expression of Neil1 mRNA in the livers of three mice containing the Neil1 shRNA transgene (shRNA-positive) or three siblings lacking the transgene (shRNA-negative) was assayed by RT-PCR (top row is Neil1). An RT-PCR of β-actin was done to ensure that equal quantities of mRNAs were tested for each mouse (second row). Expression of the neomycin resistance gene (neo), carried on the shRNA vector, was tested similarly (third row). Finally, the mice were genotyped using genomic DNA that was PCR-amplified with vector-specific primers (bottom row). (b) Similar studies were performed in the heart. (c) Similar studies were performed in the spleen. Animal procedures have been approved by the SUNY, Stony Brook Institutional Animal Care and Use Committee (IACUC).

[0105] **FIG. 56**: Reduction in Neil1 protein correlates with the presence of siRNAs. (a) Expression of Neil1 protein was examined in protein extracts from the livers of mice carrying the shRNA transgene (shRNA-positive) or siblings lacking the transgene (shRNA-negative) by western blotting with Neil1-specific antiserum. A western blot for PCNA was used to standardize loading. (b) The presence of siRNAs in RNA derived from the livers of transgenic mice as assayed by northern blotting using a 300 nt probe, part of which was complementary to the shRNA sequence. We note siRNAs only in mice transgenic for the shRNA expression cassette.

DETAILED DESCRIPTION OF CERTAIN PREFERRED EMBODIMENTS

[0106] I. Overview

[0107] The present invention provides methods for attenuating gene expression in a cell using gene-targeted double stranded RNA (dsRNA). The dsRNA contains a nucleotide sequence that hybridizes under physiologic conditions of the cell to the nucleotide sequence of at least a portion of the gene to be inhibited (the "target" gene). The nucleotide sequence can hybridize to either coding or non-coding sequence of the target gene.

[0108] A significant aspect to certain embodiments of the present invention relates to the demonstration in the present application that RNAi can in fact be accomplished both in cultured mammalian cells and in whole organisms. This had not been previously described in the art.

[0109] Another salient feature of the present invention concerns the ability to carry out RNAi in higher eukaryotes, particularly in non-oocytic cells of mammals, e.g., cells from adult mammals as an example.

[0110] Furthermore, in contrast to the teachings of the prior art, we demonstrate that RNAi in mammalian systems can be mediated with dsRNA identical or similar to non-coding sequence of a target gene. It was previously believed that although dsRNA identical or similar to non-coding sequences (i.e., promoter, enhancer, or intronic sequences) did not inhibit RNAi, such dsRNAs were not thought to mediate RNAi.

[0111] As described in further detail below, the present invention(s) are based on the discovery that the RNAi phenomenum is mediated by a set of enzyme activities, including an essential RNA component, that are evolutionarily conserved in eukaryotes ranging from plants to mammals.

[0112] One enzyme contains an essential RNA component. After partial purification, a multi-component nuclease (herein "RISC nuclease") co-fractionates with a discrete, 22-nucleotide RNA species which may confer specificity to the nuclease through homology to the substrate mRNAs. The short RNA molecules are generated by a processing reaction from the longer input dsRNA. Without wishing to be bound by any particular theory, these 22mer guide RNAs may serve as guide sequences that instruct the RISC nuclease to destroy specific mRNAs corresponding to the dsRNA sequences.

[0113] As illustrated, double stranded forms of the 22-mer guide RNA can be sufficient in length to induce sequence-dependent dsRNA inhibition of gene expression. In the illustrated example, dsRNA contructs are administered to cells having a recombinant luciferase reporter gene. In the control cell, e.g., no exogeneously added RNA, the level of expression of the luciferase reporter is normalized to be the value of "1". As illustrated, both long (500-mer) and short (22-mer) dsRNA constructs complementary to the luciferase gene could inhibit expression of that gene product relative to the control cell. On the other hand, similarly sized dsRNA complementary to the coding sequence for another protein, green fluorescence protein (GFP), did not significantly effect the expression of luciferase—indicating that the inhibitory phenomena was in each case sequence-dependent. Likewise, single stranded 22-mers of luciferase did not inhibit expression of that gene—indicating that the inhibitory phenomena is double stranded-dependent.

[0114] The appended examples also identify an enzyme, Dicer, that can produce the putative guide RNAs. Dicer is a member of the RNAse III family of nucleases that specifically cleave dsRNA and is evolutionarily conserved in worms, flies, plants, fungi and, as described herein, mammals. The enzyme has a distinctive structure which includes a helicase domain and dual RNAse III motifs. Dicer also contains a region of homology to the RDE1/QDE2/ARGONAUTE family, which have been genetically linked to RNAi in lower eukaryotes. Indeed, activation of, or overexpression of Dicer may be sufficient in many cases to permit RNA interference in otherwise non-receptive cells, such as cultured eukaryotic cells, or mammalian (non-oocytic) cells in culture or in whole organisms.

[0115] In certain embodiments, the cells can be treated with an agent(s) that inhibits the general double-stranded RNA response(s) by the host cells, such as may give rise to sequence-independent apoptosis. For instance, the cells can be treated with agents that inhibit the dsRNA-dependent protein kinase known as PKR (protein kinase RNA-activated). Double stranded RNAs in mammalian cells typically activate protein kinase PKR and lead to apoptosis. The mechanism of action of PKR includes phosphorylation and inactivation of eIF2α (Fire (1999) *Trends Genet* 15: 358). It has also been reported that induction of NF-κB by PKR is involved in apoptosis commitment and this process is mediated through activation of the IKK complex. This sequence-independent response may reflect a form of primitive immune response, since the presence of dsRNA is a common feature of many viral lifecycles.

[0116] As described herein, Applicants have demonstrated that the PKR response can be overcome in favor of the sequence-specific RNAi response. However, in certain instances, it may be desirable to treat the cells with agents which inhibit expression of PKR, cause its destruction, and/or inhibit the kinase activity of PKR, and such methods are specifically contemplated for use in the present invention. Likewise, overexpression of agents which ectopically activate eIF2α can be used. Other agents which can be used to suppress the PKR response include inhibitors of IκB phosphorylation of IκB, inhibitors of IκB ubiquitination, inhibitors of IκB degradation, inhibitors of NF-κB nuclear translocation, and inhibitors of NF-κB interaction with κB response elements.

[0117] Other inhibitors of sequence-independent dsRNA response in cells include the gene product of the vaccinia virus E3L. The E3L gene product contains two distinct domains. A conserved carboxy-terminal domain has been shown to bind double-stranded RNA (dsRNA) and inhibit the antiviral dsRNA response by cells. Expression of at least that portion of the E3L gene in the host cell, or the use of polypeptide or peptidomimetics thereof, can be used to suppress the general dsRNA response. Caspase inhibitors sensitize cells to killing by double-stranded RNA. Accordingly, ectopic expression or activation of caspases in the host cell can be used to suppress the general dsRNA response.

[0118] In other embodiments, the subject method is carried out in cells which have little or no general response to double stranded RNA, e.g., have no PKR-dependent dsRNA response, at least under the culture conditions. As illustrated in FIGS. **28-32**, CHO and P19 cells can be used without having to inhibit PKR or other general dsRNA responses.

[0119] Thus, the present invention provides a process and compositions for inhibiting expression of a target gene in a cell, especially a mammalian cell. In certain embodiments, the process comprises introduction of RNA (the "dsRNA construct") with partial or fully double-stranded character into the cell or into the extracellular environment. Inhibition is specific in that a nucleotide sequence from a portion of the target gene is chosen to produce the dsRNA construct. The dsRNA may be identical or similar to coding or non-coding sequence of the target gene. In preferred embodiments, the method utilizes a cell in which Dicer and/or Argonaute activities are recombinantly expressed or otherwise ectopically activated. This process can be (1) effective in attenuating gene expression, (2) specific to the targeted gene, and (3) general in allowing inhibition of many different types of target gene.

[0120] II. Definitions

[0121] For convenience, certain terms employed in the specification, examples, and appended claims are collected here.

[0122] As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a genomic integrated vector, or "integrated vector", which can become integrated into the chromsomal DNA of the host cell. Another type of vector is an episomal vector, i.e., a nucleic acid capable of extra-chromosomal replication. Vectors capable of directing the expression of genes to which they are operatively linked are referred to herein as "expression vectors". In the present specification, "plasmid" and "vector" are used interchangeably unless otherwise clear from the context.

[0123] As used herein, the term "nucleic acid" refers to polynucleotides such as deoxyribonucleic acid (DNA), and, where appropriate, ribonucleic acid (RNA). The term should also be understood to include, as applicable to the embodiment being described, single-stranded (such as sense or antisense) and double-stranded polynucleotides.

[0124] As used herein, the term "gene" or "recombinant gene" refers to a nucleic acid comprising an open reading frame encoding a polypeptide of the present invention, including both exon and (optionally) intron sequences. The nucleic acid may also optionally include non-coding

sequences such as promoter or enhancer sequences. A "recombinant gene" refers to nucleic acid encoding such regulatory polypeptides, that may optionally include intron sequences that are derived from chromosomal DNA. The term "intron" refers to a DNA sequence present in a given gene that is not translated into protein and is generally found between exons.

[0125] A "protein coding sequence" or a sequence that "encodes" a particular polypeptide or peptide, is a nucleic acid sequence that is transcribed (in the case of DNA) and is translated (in the case of mRNA) into a polypeptide in vitro or in vivo when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A coding sequence can include, but is not limited to, cDNA from procaryotic or eukaryotic mRNA, genomic DNA sequences from procaryotic or eukaryotic DNA, and even synthetic DNA sequences. A transcription termination sequence will usually be located 3' to the coding sequence.

[0126] Likewise, "encodes", unless evident from its context, will be meant to include DNA sequences that encode a polypeptide, as the term is typically used, as well as DNA sequences that are transcribed into inhibitory antisense molecules.

[0127] The term "loss-of-function", as it refers to genes inhibited by the subject RNAi method, refers to a diminishment in the level of expression of a gene(s) in the presence of one or more dsRNA construct(s) when compared to the level in the absense of such dsRNA construct(s).

[0128] The term "expression" with respect to a gene sequence refers to transcription of the gene and, as appropriate, translation of the resulting mRNA transcript to a protein. Thus, as will be clear from the context, expression of a protein coding sequence results from transcription and translation of the coding sequence.

[0129] "Cells,""host cells" or "recombinant host cells" are terms used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

[0130] The term "cultured cells" refers to cells suspended in culture, e.g., dispersed in culture or in the form tissue. It does not, however, include oocytes or whole embryos (including blastocysts and the like) which may be provided in culture. In certain embodiments, the cultured cells are adults cells, e.g., non-embryonic.

[0131] By "recombinant virus" is meant a virus that has been genetically altered, e.g., by the addition or insertion of a heterologous nucleic acid construct into the particle.

[0132] As used herein, the terms "transduction" and "transfection" are art recognized and mean the introduction of a nucleic acid, e.g., an expression vector, into a recipient cell by nucleic acid-mediated gene transfer. "Transformation", as used herein, refers to a process in which a cell's genotype is changed as a result of the cellular uptake of exogenous DNA or RNA, and, for example, the transformed cell expresses a dsRNA contruct.

[0133] "Transient transfection" refers to cases where exogenous DNA does not integrate into the genome of a transfected cell, e.g., where episomal DNA is transcribed into mRNA and translated into protein.

[0134] A cell has been "stably transfected" with a nucleic acid construct when the nucleic acid construct is capable of being inherited by daughter cells.

[0135] As used herein, a "reporter gene construct" is a nucleic acid that includes a "reporter gene" operatively linked to at least one transcriptional regulatory sequence. Transcription of the reporter gene is controlled by these sequences to which they are linked. The activity of at least one or more of these control sequences can be directly or indirectly regulated by the target receptor protein. Exemplary transcriptional control sequences are promoter sequences. A reporter gene is meant to include a promoter-reporter gene construct that is heterologously expressed in a cell.

[0136] As used herein, "transformed cells" refers to cells that have spontaneously converted to a state of unrestrained growth, i.e., they have acquired the ability to grow through an indefinite number of divisions in culture. Transformed cells may be characterized by such terms as neoplastic, anaplastic and/or hyperplastic, with respect to their loss of growth control. For purposes of this invention, the terms "transformed phenotype of malignant mammalian cells" and "transformed phenotype" are intended to encompass, but not be limited to, any of the following phenotypic traits associated with cellular transformation of mammalian cells: immortalization, morphological or growth transformation, and tumorigenicity, as detected by prolonged growth in cell culture, growth in semi-solid media, or tumorigenic growth in immuno-incompetent or syngeneic animals.

[0137] As used herein, "proliferating" and "proliferation" refer to cells undergoing mitosis.

[0138] As used herein, "immortalized cells" refers to cells that have been altered via chemical, genetic, and/or recombinant means such that the cells have the ability to grow through an indefinite number of divisions in culture.

[0139] The "growth state" of a cell refers to the rate of proliferation of the cell and the state of differentiation of the cell.

[0140] "MHC antigen", as used herein, refers to a protein product of one or more MHC genes; the term includes fragments or analogs of products of MHC genes which can evoke an immune response in a recipient organism. Examples of MHC antigens include the products (and fragments or analogs thereof) of the human MHC genes, i.e., the HLA genes.

[0141] The term "histocompatibility" refers to the similarity of tissue between different individuals. The level of histocompatibility describes how well matched the patient and donor are. The major histocompatibility determinants are the human leukocyte antigens (HLA). HLA typing is performed between the potential marrow donor and the potential transplant recipient to determine how close a HLA match the two are. The closer the match the less the donated marrow and the patient's body will react against each other.

[0142] The term "human leukocyte antigens" or "HLA", refers to proteins (antigens) found on the surface of white blood cells and other tissues that are used to match donor and patient. For instances, a patient and potential donor may have their white blood cells tested for such HLA antigens as, HLA-A, B and DR. Each individual has two sets of these antigens, one set inherited from each parent. For this reason, it is much more likely for a brother or sister to match the patient than an unrelated individual, and much more likely for persons of the same racial and ethnic backgrounds to match each other.

[0143] III. Exemplary Embodiments of Isolation Method

[0144] One aspect of the invention provides a method for potentiating RNAi by induction or ectopic activation of an RNAi enzyme in a cell (in vivo or in vitro) or cell-free mixtures. In preferred embodiments, the RNAi activity is activated or added to a mammalian cell, e.g., a human cell, which cell may be provided in vitro or as part of a whole organism. In other embodiments, the subject method is carried out using eukaryotic cells generally (except for oocytes) in culture. For instance, the Dicer enzyme may be activated by virtue of being recombinantly expressed or it may be activated by use of an agent which (i) induces expression of the endogenous gene, (ii) stabilizes the protein from degradation, and/or (iii) allosterically modifies the enzyme to increase its activity (by altering its Kcat, Km or both).

[0145] A. Dicer and Argonaut Activities

[0146] In certain embodiments, at least one of the activated RNAi enzymes is Dicer, or a homolog thereof. In certain preferred embodiments, the present method provides for ectopic activation of Dicer. As used herein, the term "Dicer" refers to a protein which (a) mediates an RNAi response and (b) has an amino acid sequence at least 50 percent identical, and more preferably at least 75, 85, 90 or 95 percent identical to SEQ ID No. 2 or 4, and/or which can be encoded by a nucleic acid which hybridizes under wash conditions of 2×SSC at 22° C., and more preferably 0.2× SSC at 65° C., to a nucleotide represented by SEQ ID No. 1 or 3. Accordingly, the method may comprise introducing a dsRNA contruct into a cell in which Dicer has been recombinantly expressed or otherwise ectopically activated.

[0147] In certain embodiment, at least one of the activated RNAi enzymes is Argonaut, or a homolog thereof. In certain preferred embodiments, the present method provides for ectopic activation of Argonaut. As used herein, the term "Argonaut" refers to a protein which (a) mediates an RNAi response and (b) has an amino acid sequence at least 50 percent identical, and more preferably at least 75, 85, 90 or 95 percent identical to the amino acid sequence shown in **FIG. 24**. Accordingly, the method may comprise introducing a dsRNA contruct into a cell in which Argonaut has been recombinantly expressed or otherwise ectopically activated.

[0148] This invention also provides expression vectors containing a nucleic acid encoding a Dicer or Argonaut polypeptide, operably linked to at least one transcriptional regulatory sequence. Operably linked is intended to mean that the nucleotide sequence is linked to a regulatory sequence in a manner which allows expression of the nucleotide sequence. Regulatory sequences are art-recognized and are selected to direct expression of the subject Dicer or Argonaut proteins. Accordingly, the term transcriptional regulatory sequence includes promoters, enhancers and other expression control elements. Such regulatory sequences are described in Goeddel; Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, Calif. (1990). For instance, any of a wide variety of expression control sequences, sequences that control the expression of a DNA sequence when operatively linked to it, may be used in these vectors to express DNA sequences encoding Dicer or Argonaut polypeptides of this invention. Such useful expression control sequences, include, for example, a viral LTR, such as the LTR of the Moloney murine leukemia virus, the early and late promoters of SV40, adenovirus or cytomegalovirus immediate early promoter, the lac system, the trp system, the TAC or TRC system, T7 promoter whose expression is directed by T7 RNA polymerase, the major operator and promoter regions of phage λ, the control regions for fd coat protein, the promoter for 3-phosphoglycerate kinase or other glycolytic enzymes, the promoters of acid phosphatase, e.g., Pho5, the promoters of the yeast α-mating factors, the polyhedron promoter of the baculovirus system and other sequences known to control the expression of genes of prokaryotic or eukaryotic cells or their viruses, and various combinations thereof. It should be understood that the design of the expression vector may depend on such factors as the choice of the host cell to be transformed and/or the type of protein desired to be expressed.

[0149] Moreover, the vector's copy number, the ability to control that copy number and the expression of any other proteins encoded by the vector, such as antibiotic markers, should also be considered.

[0150] The recombinant Dicer or Argonaut genes can be produced by ligating a nucleic acid encoding a Dicer or Argonaut polypeptide into a vector suitable for expression in either prokaryotic cells, eukaryotic cells, or both. Expression vectors for production of recombinant forms of the subject Dicer or Argonaut polypeptides include plasmids and other vectors. For instance, suitable vectors for the expression of a Dicer or Argonaut polypeptide include plasmids of the types: pBR322-derived plasmids, pEMBL-derived plasmids, pEX-derived plasmids, pBTac-derived plasmids and pUC-derived plasmids for expression in prokaryotic cells, such as *E. coli.*

[0151] A number of vectors exist for the expression of recombinant proteins in yeast. For instance, YEP24, YIP5, YEP51, YEP52, pYES2, and YRP17 are cloning and expression vehicles useful in the introduction of genetic constructs into *S. cerevisiae* (see, for example, Broach et al. (1983) in Experimental Manipulation of Gene Expression, ed. M. Inouye Academic Press, p. 83, incorporated by reference herein). These vectors can replicate in *E. coli* due the presence of the pBR322 ori, and in *S. cerevisiae* due to the replication determinant of the yeast 2 micron plasmid. In addition, drug resistance markers such as ampicillin can be used. In an illustrative embodiment, a Dicer or Argonaut polypeptide is produced recombinantly utilizing an expression vector generated by sub-cloning the coding sequence of a Dicer or Argonaut gene.

[0152] The preferred mammalian expression vectors contain both prokaryotic sequences, to facilitate the propagation of the vector in bacteria, and one or more eukaryotic

transcription units that are expressed in eukaryotic cells. The pcDNAI/amp, pcDNAI/neo, pRc/CMV, pSV2gpt, pSV2neo, pSV2-dhfr, pTk2, pRSVneo, pMSG, pSVT7, pko-neo and pHyg derived vectors are examples of mammalian expression vectors suitable for transfection of eukaryotic cells. Some of these vectors are modified with sequences from bacterial plasmids, such as pBR322, to facilitate replication and drug resistance selection in both prokaryotic and eukaryotic cells. Alternatively, derivatives of viruses such as the bovine papillomavirus (BPV-1), or Epstein-Barr virus (pHEBo, pREP-derived and p205) can be used for transient expression of proteins in eukaryotic cells. The various methods employed in the preparation of the plasmids and transformation of host organisms are well known in the art. For other suitable expression systems for both prokaryotic and eukaryotic cells, as well as general recombinant procedures, see Molecular Cloning A Laboratory Manual, 2nd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press: 1989) Chapters 16 and 17.

[0153] In yet another embodiment, the subject invention provides a "gene activation" construct which, by homologous recombination with a genomic DNA, alters the transcriptional regulatory sequences of an endogenous Dicer or Argonaut gene. For instance, the gene activation construct can replace the endogenous promoter of a Dicer or Argonaut gene with a heterologous promoter, e.g., one which causes constitutive expression of the Dicer or Argonaut gene or which causes inducible expression of the gene under conditions different from the normal expression pattern of Dicer or Argonaut. A variety of different formats for the gene activation constructs are available. See, for example, the Transkaryotic Therapies, Inc PCT publications WO93/09222, WO95/31560, WO96/29411, WO95/31560 and WO94/12650.

[0154] In preferred embodiments, the nucleotide sequence used as the gene activation construct can be comprised of (1) DNA from some portion of the endogenous Dicer or Argonaut gene (exon sequence, intron sequence, promoter sequences, etc.) which direct recombination and (2) heterologous transcriptional regulatory sequence(s) which is to be operably linked to the coding sequence for the genomic Dicer or Argonaut gene upon recombination of the gene activation construct. For use in generating cultures of Dicer or Argonaut producing cells, the construct may further include a reporter gene to detect the presence of the knockout construct in the cell.

[0155] The gene activation construct is inserted into a cell, and integrates with the genomic DNA of the cell in such a position so as to provide the heterologous regulatory sequences in operative association with the native Dicer or Argonaut gene. Such insertion occurs by homologous recombination, i.e., recombination regions of the activation construct that are homologous to the endogenous Dicer or Argonaut gene sequence hybridize to the genomic DNA and recombine with the genomic sequences so that the construct is incorporated into the corresponding position of the genomic DNA.

[0156] The terms "recombination region" or "targeting sequence" refer to a segment (i.e., a portion) of a gene activation construct having a sequence that is substantially identical to or substantially complementary to a genomic gene sequence, e.g., including 5' flanking sequences of the genomic gene, and can facilitate homologous recombination between the genomic sequence and the targeting transgene construct.

[0157] As used herein, the term "replacement region" refers to a portion of a activation construct which becomes integrated into an endogenous chromosomal location following homologous recombination between a recombination region and a genomic sequence.

[0158] The heterologous regulatory sequences, e.g., which are provided in the replacement region, can include one or more of a variety of elements, including: promoters (such as constitutive or inducible promoters), enhancers, negative regulatory elements, locus control regions, transcription factor binding sites, or combinations thereof.

[0159] Promoters/enhancers which may be used to control the expression of the targeted gene in vivo include, but are not limited to, the cytomegalovirus (CMV) promoter/enhancer (Karasuyama et al. (1989) *J. Exp. Med* 169: 13), the human β-actin promoter (Gunning et al. (1987) *PNAS* 84: 4831-4835), the glucocorticoid-inducible promoter present in the mouse mammary tumor virus long terminal repeat (MMTV LTR) (Klessig et al. (1984) *Mol. Cell Biol.* 4: 1354-1362), the long terminal repeat sequences of Moloney murine leukemia virus (MuLV LTR) (Weiss et al. (1985) *RNA Tumor Viruses*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.), the SV40 early or late region promoter (Bemoist et al. (1981) *Nature* 290: 304-310; Templeton et al. (1984) *Mol. Cell Biol.* 4: 817; and Sprague et al. (1983) *J. Virol.* 45: 773), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (RSV) (Yamamoto et al. (1980) *Cell* 22: 787-797), the herpes simplex virus (HSV) thymidine kinase promoter/enhancer (Wagner et al. (1981) *PNAS* 82: 3567-71), and the herpes simplex virus LAT promoter (Wolfe et al. (1992) *Nature Genetics* 1: 379-384).

[0160] In still other embodiments, the replacement region merely deletes a negative transcriptional control element of the native gene, e.g., to activate expression, or ablates a positive control element, e.g., to inhibit expression of the targeted gene.

[0161] B. Cell/Organism

[0162] The cell with the target gene may be derived from or contained in any organism (e.g., plant, animal, protozoan, virus, bacterium, or fungus). The dsRNA construct may be synthesized either in vivo or in vitro. Endogenous RNA polymerase of the cell may mediate transcription in vivo, or cloned RNA polymerase can be used for transcription in vivo or in vitro. For generating double stranded transcripts from a transgene in vivo, a regulatory region may be used to transcribe the RNA strand (or strands). Furthermore, dsRNA can be generated by transcribing an RNA strand which forms a hairpin, thus producing a dsRNA.

[0163] Genetic manipulation becomes possible in organisms that are not classical genetic models. Breeding and screening programs may be accelerated by the ability to rapidly assay the consequences of a specific, targeted gene disruption. Gene disruptions may be used to discover the function of the target gene, to produce disease models in which the target gene are involved in causing or preventing a pathological condition, and to produce organisms with improved economic properties.

[0164] The cell with the target gene may be derived from or contained in any organism. The organism may be a plant, animal, protozoan, bacterium, virus, or fungus. The plant may be a monocot, dicot or gymnosperm; the animal may be a vertebrate or invertebrate. Preferred microbes are those used in agriculture or by industry, and those that are pathogenic for plants or animals. Fungi include organisms in both the mold and yeast morphologies.

[0165] Plants include arabidopsis; field crops (e.g., alfalfa, barley, bean, corn, cotton, flax, pea, rape, rice, rye, safflower, sorghum, soybean, sunflower, tobacco, and wheat); vegetable crops (e.g., asparagus, beet, broccoli, cabbage, carrot, cauliflower, celery, cucumber, eggplant, lettuce, onion, pepper, potato, pumpkin, radish, spinach, squash, taro, tomato, and zucchini); fruit and nut crops (e.g., almond, apple, apricot, banana, blackberry, blueberry, cacao, cherry, coconut, cranberry, date, fajoa, filbert, grape, grapefruit, guava, kiwi, lemon, lime, mango, melon, nectarine, orange, papaya, passion fruit, peach, peanut, pear, pineapple, pistachio, plum, raspberry, strawberry, tangerine, walnut, and watermelon); and ornamentals (e.g., alder, ash, aspen, azalea, birch, boxwood, camellia, carnation, chrysanthemum, elm, fir, ivy, jasmine, juniper, oak, palm, poplar, pine, redwood, rhododendron, rose, and rubber).

[0166] Examples of vertebrate animals include fish, mammal, cattle, goat, pig, sheep, rodent, hamster, mouse, rat, primate, and human.

[0167] Invertebrate animals include nematodes, other worms, drosophila, and other insects. Representative genera of nematodes include those that infect animals (e.g., Ancylostoma, Ascaridia, Ascaris, Bunostomum, Caenorhabditis, Capillaria, Chabertia, Cooperia, Dictyocaulus, Haernonchus, Heterakis, Nematodirus, Oesophagostomum, Ostertagia, Oxyuris, Parascaris, Strongylus, Toxascaris, Trichuris, Trichostrongylus, Tflichonema, Toxocara, Uncinaria) and those that infect plants (e.g., Bursaphalenchus, Criconerriella, Diiylenchus, Ditylenchus, Globodera, Helicotylenchus, Heterodera, Longidorus, Melodoigyne, Nacobbus, Paratylenchus, Pratylenchus, Radopholus, Rotelynchus, Tylenchus, and Xiphinerna). Representative orders of insects include Coleoptera, Diptera, Lepidoptera, and Homoptera.

[0168] The cell having the target gene may be from the germ line or somatic, totipotent or pluripotent, dividing or non-dividing, parenchyma or epithelium, immortalized or transformed, or the like. The cell may be a stem cell or a differentiated cell. Cell types that are differentiated include adipocytes, fibroblasts, myocytes, cardiomyocytes, endothelium, neurons, glia, blood cells, megakaryocytes, lymphocytes, macrophages, neutrophils, eosinophils, basophils, mast cells, leukocytes, granulocytes, keratinocytes, chondrocytes, osteoblasts, osteoclasts, hepatocytes, and cells of the endocrine or exocrine glands.

[0169] C. Targeted Genes

[0170] The target gene may be a gene derived from the cell, an endogenous gene, a transgene, or a gene of a pathogen which is present in the cell after infection thereof. Depending on the particular target gene and the dose of double stranded RNA material delivered, the procedure may provide partial or complete loss of function for the target gene. Lower doses of injected material and longer times after administration of dsRNA may result in inhibition in a smaller fraction of cells. Quantitation of gene expression in a cell may show similar amounts of inhibition at the level of accumulation of target mRNA or translation of target protein.

[0171] "Inhibition of gene expression" refers to the absence (or observable decrease) in the level of protein and/or mRNA product from a target gene. "Specificity" refers to the ability to inhibit the target gene without manifest effects on other genes of the cell. The consequences of inhibition can be confirmed by examination of the outward properties of the cell or organism (as presented below in the examples) or by biochemical techniques such as RNA solution hybridization, nuclease protection, Northern hybridization, reverse transcription, gene expression monitoring with a microarray, antibody binding, enzyme linked immunosorbent assay (ELISA), Western blotting, radioimmunoassay (RIA), other immunoassays, and fluorescence activated cell analysis (FACS). For RNA-mediated inhibition in a cell line or whole organism, gene expression is conveniently assayed by use of a reporter or drug resistance gene whose protein product is easily assayed. Such reporter genes include acetohydroxyacid synthase (AHAS), alkaline phosphatase (AP), beta galactosidase (LacZ), beta glucoronidase (GUS), chloramphenicol acetyltransferase (CAT), green fluorescent protein (GFP), horseradish peroxidase (HRP), luciferase (Luc), nopaline synthase (NOS), octopine synthase (OCS), and derivatives thereof. Multiple selectable markers are available that confer resistance to ampicillin, bleomycin, chloramphenicol, gentamycin, hygromycin, kanamycin, lincomycin, methotrexate, phosphinothricin, puromycin, and tetracyclin.

[0172] Depending on the assay, quantitation of the amount of gene expression allows one to determine a degree of inhibition which is greater than 10%, 33%, 50%, 90%, 95% or 99% as compared to a cell not treated according to the present invention. Lower doses of injected material and longer times after administration of dsRNA may result in inhibition in a smaller fraction of cells (e.g., at least 10%, 20%, 50%, 75%, 90%, or 95% of targeted cells). Quantitation of gene expression in a cell may show similar amounts of inhibition at the level of accumulation of target mRNA or translation of target protein. As an example, the efficiency of inhibition may be determined by assessing the amount of gene product in the cell: mRNA may be detected with a hybridization probe having a nucleotide sequence outside the region used for the inhibitory double-stranded RNA, or translated polypeptide may be detected with an antibody raised against the polypeptide sequence of that region.

[0173] As disclosed herein, the present invention is not limited to any type of target gene or nucleotide sequence. But the following classes of possible target genes are listed for illustrative purposes: developmental genes (e.g., adhesion molecules, cyclin kinase inhibitors, Writ family members, Pax family members, Winged helix family members, Hox family members, cytokines/lymphokines and their receptors, growth/differentiation factors and their receptors, neurotransmitters and their receptors); oncogenes (e.g., ABLI, BCLI, BCL2, BCL6, CBFA2, CBL, CSFIR, ERBA, ERBB, EBRB2, ETSI, ETS1, ETV6, FGR, FOS, FYN, HCR, HRAS, JUN, KRAS, LCK, LYN, MDM2, MLL, MYB, MYC, MYCLI, MYCN, NRAS, PIM 1, PML, RET, SRC, TALI, TCL3, and YES); tumor suppressor genes (e.g.,

APC, BRCA 1, BRCA2, MADH4, MCC, NF 1, NF2, RB 1, TP53, and WTI); and enzymes (e.g., ACC synthases and oxidases, ACP desaturases and hydroxylases, ADP-glucose pyrophorylases, ATPases, alcohol dehydrogenases, amylases, amyloglucosidases, catalases, cellulases, chalcone synthases, chitinases, cyclooxygenases, decarboxylases, dextrinases, DNA and RNA polymerases, galactosidases, glucanases, glucose oxidases, granule-bound starch synthases, GTPases, helicases, hemicellulases, integrases, inulinases, invertases, isomerases, kinases, lactases, lipases, lipoxygenases, lysozymes, nopaline synthases, octopine synthases, pectinesterases, peroxidases, phosphatases, phospholipases, phosphorylases, phytases, plant growth regulator synthases, polygalacturonases, proteinases and peptidases, pullanases, recombinases, reverse transcriptases, RUBISCOs, topoisomerases, and xylanases).

[0174] D. dsRNA Constructs

[0175] The dsRNA construct may comprise one or more strands of polymerized ribonucleotide. It may include modifications to either the phosphate-sugar backbone or the nucleoside. For example, the phosphodiester linkages of natural RNA may be modified to include at least one of a nitrogen or sulfur heteroatom. Modifications in RNA structure may be tailored to allow specific genetic inhibition while avoiding a general panic response in some organisms which is generated by dsRNA. Likewise, bases may be modified to block the activity of adenosine deaminase. The dsRNA construct may be produced enzymatically or by partial/total organic synthesis, any modified ribonucleotide can be introduced by in vitro enzymatic or organic synthesis.

[0176] The dsRNA construct may be directly introduced into the cell (i.e., intracellularly); or introduced extracellularly into a cavity, interstitial space, into the circulation of an organism, introduced orally, or may be introduced by bathing an organism in a solution containing RNA. Methods for oral introduction include direct mixing of RNA with food of the organism, as well as engineered approaches in which a species that is used as food is engineered to express an RNA, then fed to the organism to be affected. Physical methods of introducing nucleic acids include injection of an RNA solution directly into the cell or extracellular injection into the organism.

[0177] The double-stranded structure may be formed by a single self-complementary RNA strand or two complementary RNA strands. RNA duplex formation may be initiated either inside or outside the cell. The RNA may be introduced in an amount which allows delivery of at least one copy per cell. Higher doses (e.g., at least 5, 10, 100, 500 or 1000 copies per cell) of double-stranded material may yield more effective inhibition; lower doses may also be useful for specific applications. Inhibition is sequence-specific in that nucleotide sequences corresponding to the duplex region of the RNA are targeted for genetic inhibition.

[0178] dsRNA constructs containing a nucleotide sequences identical to a portion, of either coding or non-coding sequence, of the target gene are preferred for inhibition. RNA sequences with insertions, deletions, and single point mutations relative to the target sequence (ds RNA similar to the target gene) have also been found to be effective for inhibition. Thus, sequence identity may be optimized by sequence comparison and alignment algorithms known in the art (see Gribskov and Devereux, Sequence Analysis Primer, Stockton Press, 1991, and references cited therein) and calculating the percent difference between the nucleotide sequences by, for example, the Smith-Waterman algorithm as implemented in the BESTFIT software program using default parameters (e.g., University of Wisconsin Genetic Computing Group). Greater than 90% sequence identity, or even 100% sequence identity, between the inhibitory RNA and the portion of the target gene is preferred. Alternatively, the duplex region of the RNA may be defined functionally as a nucleotide sequence that is capable of hybridizing with a portion of the target gene transcript (e.g., 400 mM NaCl, 40 mM PIPES pH 6.4, 1 mM EDTA, 50° C. or 70° C. hybridization for 12-16 hours; followed by washing). In certain preferred embodiments, the length of the dsRNA is at least 20, 21 or 22 nucleotides in length, e.g., corresponding in size to RNA products produced by Dicer-dependent cleavage. In certain embodiments, the dsRNA construct is at least 25, 50, 100, 200, 300 or 400 bases. In certain embodiments, the dsRNA construct is 400-800 bases in length.

[0179] 100% sequence identity between the RNA and the target gene is not required to practice the present invention. Thus the invention has the advantage of being able to tolerate sequence variations that might be expected due to genetic mutation, strain polymorphism, or evolutionary divergence.

[0180] The dsRNA construct may be synthesized either in vivo or in vitro. Endogenous RNA polymerase of the cell may mediate transcription in vivo, or cloned RNA polymerase can be used for transcription in vivo or in vitro. For transcription from a transgene in vivo or an expression construct, a regulatory region (e.g., promoter, enhancer, silencer, splice donor and acceptor, polyadenylation) may be used to transcribe the dsRNA strand (or strands). Inhibition may be targeted by specific transcription in an organ, tissue, or cell type; stimulation of an environmental condition (e.g., infection, stress, temperature, chemical inducers); and/or engineering transcription at a developmental stage or age. The RNA strands may or may not be polyadenylated; the RNA strands may or may not be capable of being translated into a polypeptide by a cell's translational apparatus. The dsRNA construct may be chemically or enzymatically synthesized by manual or automated reactions. The dsRNA construct may be synthesized by a cellular RNA polymerase or a bacteriophage RNA polymerase (e.g., T3, T7, SP6). The use and production of an expression construct are known in the art (see also WO 97/32016; U.S. Pat. Nos. 5,593,874, 5,698,425, 5,712,135, 5,789,214, and 5,804,693; and the references cited therein). If synthesized chemically or by in vitro enzymatic synthesis, the RNA may be purified prior to introduction into the cell. For example, RNA can be purified from a mixture by extraction with a solvent or resin, precipitation, electrophoresis, chromatography or a combination thereof. Alternatively, the dsRNA construct may be used with no or a minimum of purification to avoid losses due to sample processing. The dsRNA construct may be dried for storage or dissolved in an aqueous solution. The solution may contain buffers or salts to promote annealing, and/or stabilization of the duplex strands.

[0181] Physical methods of introducing nucleic acids include injection of a solution containing the dsRNA construct, bombardment by particles covered by the dsRNA construct, soaking the cell or organism in a solution of the

RNA, or electroporation of cell membranes in the presence of the dsRNA construct. A viral construct packaged into a viral particle would accomplish both efficient introduction of an expression construct into the cell and transcription of dsRNA construct encoded by the expression construct. Other methods known in the art for introducing nucleic acids to cells may be used, such as lipid-mediated carrier transport, chemical mediated transport, such as calcium phosphate, and the like. Thus the dsRNA construct may be introduced along with components that perform one or more of the following activities: enhance RNA uptake by the cell, promote annealing of the duplex strands, stabilize the annealed strands, or other-wise increase inhibition of the target gene.

[0182] E. Illustrative Uses

[0183] One utility of the present invention is as a method of identifying gene function in an organism, especially higher eukaryotes, comprising the use of double-stranded RNA to inhibit the activity of a target gene of previously unknown function. Instead of the time consuming and laborious isolation of mutants by traditional genetic screening, functional genomics would envision determining the function of uncharacterized genes by employing the invention to reduce the amount and/or alter the timing of target gene activity. The invention could be used in determining potential targets for pharmaceuticals, understanding normal and pathological events associated with development, determining signaling pathways responsible for postnatal development/aging, and the like. The increasing speed of acquiring nucleotide sequence information from genomic and expressed gene sources, including total sequences for mammalian genomes, can be coupled with the invention to determine gene function in a cell or in a whole organism. The preference of different organisms to use particular codons, searching sequence databases for related gene products, correlating the linkage map of genetic traits with the physical map from which the nucleotide sequences are derived, and artificial intelligence methods may be used to define putative open reading frames from the nucleotide sequences acquired in such sequencing projects.

[0184] A simple assay would be to inhibit gene expression according to the partial sequence available from an expressed sequence tag (EST). Functional alterations in growth, development, metabolism, disease resistance, or other biological processes would be indicative of the normal role of the EST's gene product.

[0185] The ease with which the dsRNA construct can be introduced into an intact cell/organism containing the target gene allows the present invention to be used in high throughput screening (HTS). For example, duplex RNA can be produced by an amplification reaction using primers flanking the inserts of any gene library derived from the target cell or organism. Inserts may be derived from genomic DNA or mRNA (e.g., cDNA and cRNA). Individual clones from the library can be replicated and then isolated in separate reactions, but preferably the library is maintained in individual reaction vessels (e.g., a 96 well microtiter plate) to minimize the number of steps required to practice the invention and to allow automation of the process.

[0186] In an exemplary embodiment, the subject invention provides an arrayed library of RNAi constructs. The array may be in the form of solutions, such as multi-well plates, or may be "printed" on solid substrates upon which cells can be grown. To illustrate, solutions containing duplex RNAs that are capable of inhibiting the different expressed genes can be placed into individual wells positioned on a microtiter plate as an ordered array, and intact cells/organisms in each well can be assayed for any changes or modifications in behavior or development due to inhibition of target gene activity.

[0187] In one embodiment, the subject method uses an arrayed library of RNAi constructs to screen for combinations of RNAi that are lethal to host cells. Synthetic lethality is a bedrock principle of experimental genetics. A synthetic lethality describes the properties of two mutations which, individually, are tolerated by the organism but which, in combination, are lethal. The subject arrays can be used to identify loss-of-function mutations that are lethal in combination with alterations in other genes, such as activated oncogenes or loss-of-function mutations to tumor suppressors. To achieve this, one can create "phenotype arrays" using cultured cells. Expression of each of a set of genes, such as the host cell's genome, can be individually systematically disrupted using RNA interference. Combination with alterations in oncogene and tumor suppressor pathways can be used to identify synthetic lethal interactions that may identify novel therapeutic targets.

[0188] In certain embodiments, the RNAi constructs can be fed directly to, or injected into, the cell/organism containing the target gene. Alternatively, the duplex RNA can be produced by in vivo or in vitro transcription from an expression construct used to produce the library. The construct can be replicated as individual clones of the library and transcribed to produce the RNA; each clone can then be fed to, injected into, or delivered by another method known in the art to the cell/organism containing the target gene. The function of the target gene can be assayed from the effects it has on the cell/organism when gene activity is inhibited. This screening could be amenable to small subjects that can be processed in large number, for example, tissue culture cells derived from mammals, especially primates, and most preferably humans.

[0189] If a characteristic of an organism is determined to be genetically linked to a polymorphism through RFLP or QTL analysis, the present invention can be used to gain insight regarding whether that genetic polymorphism might be directly responsible for the characteristic. For example, a fragment defining the genetic polymorphism or sequences in the vicinity of such a genetic polymorphism can be amplified to produce an RNA, the duplex RNA can be introduced to the organism or cell, and whether an alteration in the characteristic is correlated with inhibition can be determined. Of course, there may be trivial explanations for negative results with this type of assay, for example: inhibition of the target gene causes lethality, inhibition of the target gene may not result in any observable alteration, the fragment contains nucleotide sequences that are not capable of inhibiting the target gene, or the target gene's activity is redundant.

[0190] The present invention may be useful in allowing the inhibition of essential genes. Such genes may be required for cell or organism viability at only particular stages of development or only in specific cellular compartments or tissues. The functional equivalent of conditional mutations

may be produced by inhibiting activity of the target gene when or where it is not required for viability. The invention allows addition of RNA at specific times of development and locations in the organism without introducing permanent mutations into the target genome.

[0191] The present invention may be useful in allowing the inhibition of genes that have been difficult to inhibit using other methods due to gene redundancy. Since the present methods may be used to deliver more than one dsRNA to a cell or organism, dsRNA identical or similar to more than one gene, wherein the genes have a redundant function during normal development, may be delivered.

[0192] If alternative splicing produced a family of transcripts that were distinguished by usage of characteristic exons, the present invention can target inhibition through the appropriate exons to specifically inhibit or to distinguish among the functions of family members. For example, a protein factor that contained an alternatively spliced transmembrane domain may be expressed in both membrane bound and secreted forms. Instead of isolating a nonsense mutation that terminates translation before the transmembrane domain, the functional consequences of having only secreted hormone can be determined according to the invention by targeting the exon containing the transmembrane domain and thereby inhibiting expression of membrane-bound hormone. That is, the subject method can be used for selected ablation of splicing variants.

[0193] The present invention may be used alone or as a component of a kit having at least one of the reagents necessary to carry out the in vitro or in vivo introduction of RNA to test samples or subjects. Preferred components are the dsRNA and a vehicle that promotes introduction of the dsRNA. Such a kit may also include instructions to allow a user of the kit to practice the invention.

[0194] Alternatively, an organism may be engineered to produce dsRNA which produces commercially or medically beneficial results, for example, resistance to a pathogen or its pathogenic effects, improved growth, or novel developmental patterns.

IV. Exemplification

[0195] The invention, now being generally described, will be more readily understood by reference to the following examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present invention and are not intended to limit the invention.

EXAMPLE 1

An RNA-Directed Nuclease Mediates RNAi Gene Silencing

[0196] In a diverse group of organisms that includes *Caenorhabditis elegans*, Drosophila, planaria, hydra, trypanosomes, fungi and plants, the introduction of double-stranded RNAs inhibits gene expression in a sequence-specific manner (Sharp (1999) *Genes and Development* 13: 139-141; Sanchez-Alvarado and Newmark (1999) *PNAS* 96: 5049-5054; Lohman et al. (1999) *Developmental Biology* 214: 211-214; Cogoni and Macino (1999) *Nature* 399: 166-169; Waterhouse et al. (1998) *PNAS* 95: 13959-13964; Montgomery and Fire (1998) *Trends Genet.* 14: 225-228; Ngo et al. (1998) *PNAS* 95: 14687-14692). These responses, called RNA interference or post-transcriptional gene silencing, may provide anti-viral defence, modulate transposition or regulate gene expression (Sharp (1999) *Genes and Development* 13: 139-141; Montgomery and Fire (1998) *Trends Genet.* 14: 225-228; Tabara et al. (1999) *Cell* 99: 123-132; Ketting et al. (1999) *Cell* 99: 133-141; Ratcliff et al. (1997) *Science* 276: 1558-1560). We have taken a biochemical approach towards elucidating the mechanisms underlying this genetic phenomenon. Here we show that 'loss-of-function' phenotypes can be created in cultured Drosophila cells by transfection with specific double-stranded RNAs. This coincides with a marked reduction in the level of cognate cellular messenger RNAs. Extracts of transfected cells contain a nuclease activity that specifically degrades exogenous transcripts homologous to transfected double-stranded RNA. This enzyme contains an essential RNA component. After partial purification, the sequence-specific nuclease co-fractionates with a discrete, ~25-nucleotide RNA species which may confer specificity to the enzyme through homology to the substrate mRNAs.

[0197] Although double-stranded RNAs (dsRNAs) can provoke gene silencing in numerous biological contexts including Drosophila (Kennerdell et al. (1998) *Cell* 95: 1017-1026; Misquitta and Paterson (1999) *PNAS* 96: 1451-1456), the mechanisms underlying this phenomenon have remained mostly unknown. We therefore wanted to establish a biochemically tractable model in which such mechanisms could be investigated.

[0198] Transient transfection of cultured, Drosophila S2 cells with a lacZ expression vector resulted in β-galactosidase activity that was easily detectable by an in situ assay (**FIG. 1***a*). This activity was greatly reduced by co-transfection with a dsRNA corresponding to the first 300 nucleotides of the lacZ sequence, whereas co-transfection with a control dsRNA (CD8) (**FIG. 1***a*) or with single-stranded RNAs of either sense or antisense orientation (data not shown) had little or no effect. This indicated that dsRNAs could interfere, in a sequence-specific fashion, with gene expression in cultured cells.

[0199] To determine whether RNA interference (RNAi) could be used to target endogenous genes, we transfected S2 cells with a dsRNA corresponding to the first 540 nucleotides of Drosophila cyclin E, a gene that is essential for progression into S phase of the cell cycle. During log-phase growth, untreated S2 cells reside primarily in G2/M (**FIG. 1***b*). Transfection with lacZ dsRNA had no effect on cell-cycle distribution, but transfection with the cyclin E dsRNA caused a G1-phase cell-cycle arrest (**FIG. 1***b*). The ability of cyclin E dsRNA to provoke this response was length-dependent. Double-stranded RNAs of 540 and 400 nucleotides were quite effective, whereas dsRNAs of 200 and 300 nucleotides were less potent. Double-stranded cyclin E RNAs of 50 or 100 nucleotides were inert in our assay, and transfection with a single-stranded, antisense cyclin E RNA had virtually no effect.

[0200] One hallmark of RNAi is a reduction in the level of mRNAs that are homologous to the dsRNA. Cells transfected with the cyclin E dsRNA (bulk population) showed diminished endogenous cyclin E mRNA as compared with control cells (**FIG. 1***c*). Similarly, transfection of cells with dsRNAs homologous to fizzy, a component of the anaphase-

promoting complex (APC) or cyclin A, a cyclin that acts in S, G2 and M, also caused reduction of their cognate mRNAs (**FIG. 1***c*). The modest reduction in fizzy mRNA levels in cells transfected with cyclin A dsRNA probably resulted from arrest at a point in the division cycle at which fizzy transcription is low (Wolf and Jackson (1998) *Current Biology* 8: R637-R639; Kramer et al. (1998) *Current Biology* 8: 1207-1210). These results indicate that RNAi may be a generally applicable method for probing gene function in cultured Drosophila cells.

[**0201**] The decrease in mRNA levels observed upon transfection of specific dsRNAs into Drosophila cells could be explained by effects at transcriptional or post-transcriptional levels. Data from other systems have indicated that some elements of the dsRNA response may affect mRNA directly (reviewed in Sharp (1999) *Genes and Development* 13: 139-141; Montgomery and Fire (1998) *Trends Genet.* 14: 225-228). We therefore sought to develop a cell-free assay that reflected, at least in part, RNAi.

[**0202**] S2 cells were transfected with dsRNAs corresponding to either cyclin E or lacZ. Cellular extracts were incubated with synthetic mRNAs of lacZ or cyclin E. Extracts prepared from cells transfected with the 540-nucleotide cyclin E dsRNA efficiently degraded the cyclin E transcript; however, the lacZ transcript was stable in these lysates (**FIG. 2***a*). Conversely, lysates from cells transfected with the lacZ dsRNA degraded the lacZ transcript but left the cyclin E mRNA intact. These results indicate that RNAi ablates target mRNAs through the generation of a sequence-specific nuclease activity. We have termed this enzyme RISC (RNA-induced silencing complex). Although we occasionally observed possible intermediates in the degradation process (see **FIG. 2**), the absence of stable cleavage end-products indicates an exonuclease (perhaps coupled to an endonuclease). However, it is possible that the RNAi nuclease makes an initial endonucleolytic cut and that non-specific exonucleases in the extract complete the degradation process (Shuttleworth and Colman (1988) *EMBO J.* 7: 427-434). In addition, our ability to create an extract that targets lacZ in vitro indicates that the presence of an endogenous gene is not required for the RNAi response.

[**0203**] To examine the substrate requirements for the dsRNA-induced, sequence-specific nuclease activity, we incubated a variety of cyclin-E-derived transcripts with an extract derived from cells that had been transfected with the 540-nucleotide cyclin E dsRNA (**FIG. 2***b, c*). Just as a length requirement was observed for the transfected dsRNA, the RNAi nuclease activity showed a dependence on the size of the RNA substrate. Both a 600-nucleotide transcript that extends slightly beyond the targeted region (**FIG. 2***b*) and an ~1-kilobase (kb) transcript that contains the entire coding sequence (data not shown) were completely destroyed by the extract. Surprisingly, shorter substrates were not degraded as efficiently. Reduced activity was observed against either a 300- or a 220-nucleotide transcript, and a ~100-nucleotide transcript was resistant to nuclease in our assay. This was not due solely to position effects because ~100-nucleotide transcripts derived from other portions of the transfected dsRNA behaved similarly (data not shown). As expected, the nuclease activity (or activities) present in the extract could also recognize the antisense strand of the cyclin E mRNA. Again, substrates that contained a substantial portion of the targeted region were degraded efficiently whereas those that contained a shorter stretch of homologous sequence (~130 nucleotides) were recognized inefficiently (**FIG. 2***c*, as600). For both the sense and antisense strands, transcripts that had no homology with the transfected dsRNA (**FIG. 2***b*, Eout; **FIG. 2***c*, as300) were not degraded. Although we cannot exclude the possibility that nuclease specificity could have migrated beyond the targeted region, the resistance of transcripts that do not contain homology to the dsRNA is consistent with data from *C. elegans*. Double-stranded RNAs homologous to an upstream cistron have little or no effect on a linked downstream cistron, despite the fact that unprocessed, polycistronic mRNAs can be readily detected (Tabara et al. (1998) *Science* 282: 430-432; Bosher et al. (1999) *Genetics* 153: 1245-1256). Furthermore, the nuclease was inactive against a dsRNA identical to that used to provoke the RNAi response in vivo (**FIG. 2***b*). In the in vitro system, neither a 5' cap nor a poly(A) tail was required, as such transcripts were degraded as efficiently as uncapped and non-polyadenylated RNAs.

[**0204**] Gene silencing provoked by dsRNA is sequence specific. A plausible mechanism for determining specificity would be incorporation of nucleic-acid guide sequences into the complexes that accomplish silencing (Hamilton and Baulcombe (1999) *Science* 286: 950-952). In accord with this idea, pre-treatment of extracts with a Ca^{2+}-dependent nuclease (micrococcal nuclease) abolished the ability of these extracts to degrade cognate mRNAs (**FIG. 3**). Activity could not be rescued by addition of non-specific RNAs such as yeast transfer RNA. Although micrococcal nuclease can degrade both DNA and RNA, treatment of the extract with DNAse I had no effect (**FIG. 3**). Sequence-specific nuclease activity, however, did require protein (data not shown). Together, our results support the possibility that the RNAi nuclease is a ribonucleoprotein, requiring both RNA and protein components. Biochemical fractionation (see below) is consistent with these components being associated in extract rather than being assembled on the target mRNA after its addition.

[**0205**] In plants, the phenomenon of co-suppression has been associated with the existence of small (~25-nucleotide) RNAs that correspond to the gene that is being silenced (Hamilton and Baulcombe (1999) *Science* 286: 950-952). To address the possibility that a similar RNA might exist in Drosophila and guide the sequence-specific nuclease in the choice of substrate, we partially purified our activity through several fractionation steps. Crude extracts contained both sequence-specific nuclease activity and abundant, heterogeneous RNAs homologous to the transfected dsRNA (**FIGS. 2 and 4***a*). The RNAi nuclease fractionated with ribosomes in a high-speed centrifugation step. Activity could be extracted by treatment with high salt, and ribosomes could be removed by an additional centrifugation step. Chromatography of soluble nuclease over an anion-exchange column resulted in a discrete peak of activity (**FIG. 4***b*, cyclin E). This retained specificity as it was inactive against a heterologous mRNA (**FIG. 4***b*, lacZ). Active fractions also contained an RNA species of 25 nucleotides that is homologous to the cyclin E target (**FIG. 4***b*, northern). The band observed on northern blots may represent a family of discrete RNAs because it could be detected with probes specific for both the sense and antisense cyclin E sequences and with probes derived from distinct segments of the dsRNA (data not shown). At present, we cannot determine

whether the 25-nucleotide RNA is present in the nuclease complex in a double-stranded or single-stranded form.

[0206] RNA interference allows an adaptive defence against both exogenous and endogenous dsRNAs, providing something akin to a dsRNA immune response. Our data, and that of others (Hamilton and Baulcombe (1999) *Science* 286: 950-952), is consistent with a model in which dsRNAs present in a cell are converted, either through processing or replication, into small specificity determinants of discrete size in a manner analogous to antigen processing. Our results suggest that the post-transcriptional component of dsRNA-dependent gene silencing is accomplished by a sequence-specific nuclease that incorporates these small RNAs as guides that target specific messages based upon sequence recognition. The identical size of putative specificity determinants in plants (Hamilton and Baulcombe (1999) *Science* 286: 950-952) and animals predicts a conservation of both the mechanisms and the components of dsRNA-induced, post-transcriptional gene silencing in diverse organisms. In plants, dsRNAs provoke not only post-transcriptional gene silencing but also chromatin remodelling and transcriptional repression (Jones et al. (1998) *EMBO J.* 17: 6385-6393; Jones et al. (1999) *Plant Cell* 11: 2291-2301). It is now critical to determine whether conservation of gene-silencing mechanisms also exists at the transcriptional level and whether chromatin remodelling can be directed in a sequence-specific fashion by these same dsRNA-derived guide sequences.

[0207] Methods:

[0208] Cell culture and RNA methods S2 cells (Schneider (1972) *J. Embryol Exp Morpho* 27: 353-365) were cultured at 27° C. in 90% Schneider's insect media (Sigma), 10% heat inactivated fetal bovine serum (FBS). Cells were transfected with dsRNA and plasmid DNA by calcium phosphate co-precipitation (DiNocera and Dawid (1983) *PNAS* 80: 7095-7098). Identical results were observed when cells were transfected using lipid reagents (for example, Superfect, Qiagen). For FACS analysis, cells were additionally transfected with a vector that directs expression of a green fluorescent protein (GFP)-US9 fusion protein (Kalejta et al. (1999) *Exp Cell Res.* 248: 322-328). These cells were fixed in 90% ice-cold ethanol and stained with propidium iodide at 25 μg/ml. FACS was performed on an Elite flow cytometer (Coulter). For northern blotting, equal loading was ensured by over-probing blots with a control complementary DNA (RP49). For the production of dsRNA, transcription templates were generated by polymerase chain reaction such that they contained T7 promoter sequences on each end of the template. RNA was prepared using the RiboMax kit (Promega). Confirmation that RNAs were double stranded came from their complete sensitivity to RNAse III (a gift from A. Nicholson). Target mRNA transcripts were synthesized using the Riboprobe kit (Promega) and were gel purified before use.

[0209] Extract preparation Log-phase S2 cells were plated on 15-cm tissue culture dishes and transfected with 30 μg dsRNA and 30 μg carrier plasmid DNA. Seventy-two hours after transfection, cells were harvested in PBS containing 5 mM EGTA, washed twice in PBS and once in hypotonic buffer (10 mM HEPES pH 7.3, 6 mM β-mercaptoethanol). Cells were suspended in 0.7 packed-cell volumes of hypotonic buffer containing Complete protease inhibitors (Boehringer) and 0.5 units/ml of RNasin (Promega). Cells were disrupted in a dounce homogenizer with a type B pestle, and lysates were centrifuged at 30,000 g for 20 min. Supernatants were used in an in vitro assay containing 20 mM HEPES pH 7.3, 110 mM KOAc, 1 mM $Mg(OAc)_2$, 3 mM EGTA, 2 mM $CaCl_2$, 1 mM DTT. Typically, 5 μl extract was used in a 10 μl assay that contained also 10,000 c.p.m. synthetic mRNA substrate.

[0210] Extract fractionation Extracts were centrifuged at 200,000 g for 3 h and the resulting pellet (containing ribosomes) was extracted in hypotonic buffer containing also 1 mM $MgCl_2$ and 300 mM KOAc. The extracted material was spun at 100,000 g for 1 h and the resulting supernatant was fractionated on Source 15Q column (Pharmacia) using a KCl gradient in buffer A (20 mM HEPES pH 7.0, 1 mM dithiothreitol, 1 mM $MgCl_2$). Fractions were assayed for nuclease activity as described above. For northern blotting, fractions were proteinase K/SDS treated, phenol extracted, and resolved on 15% acrylamide 8M urea gels. RNA was electroblotted onto Hybond N+ and probed with strand-specific riboprobes derived from cyclin E mRNA. Hybridization was carried out in 500 mM $NaPO_4$ pH 7.0, 15% formamide, 7% SDS, 1% BSA. Blots were washed in 1×SSC at 37-45° C.

EXAMPLE 2

Role for a Bidentate Ribonuclease in the Initiation Step of RNA Interference

[0211] Genetic approaches in worms, fungi and plants have identified a group of proteins that are essential for double-stranded RNA-induced gene silencing. Among these are ARGONAUTE family members (e.g. RDE1, QDE2) (Tabara et al. (1999) *Cell* 99: 123-132; Catalanotto et al. (2000) *Nature* 404: 245; Fagard et al. (2000) *PNAS* 97: 11650-11654), recQ-family helicases (MUT-7, QDE3) (Ketting et al. (1999) *Cell* 99: 133-141; Cogoni and Macino. (1999) *Science* 286: 2342-2344), and RNA-dependent RNA polymerases (e.g. EGO-1, QDE1, SGS2/SDE1) (Cogoni and Macino (1999) *Nature* 399: 166-169; Smardon et al. (2000) *Current Biology* 10: 169-178; Mourrain et al. (2000) *Cell* 101: 533-542; Dalmay et al. (2000) *Cell* 101: 543-553). While potential roles have been proposed, none of these genes has been assigned a definitive function in the silencing process. Biochemical studies have suggested that PTGS is accomplished by a multicomponent nuclease that targets mRNAs for degradation (Hammond et al. (2000) *Nature* 404: 293-296; Zamore et al. (2000) *Cell* 101 25-33; Tuschl et al. (1999) *Genes and Development* 13: 3191-3197). We have shown that the specificity of this complex may derive from the incorporation of a small guide sequence that is homologous to the mRNA substrate (Hammond et al. (2000) *Nature* 404: 293-296). Originally identified in plants that were actively silencing transgenes (Hamilton and Baulcombe (1999) *Science* 286: 950-952), these ~22 nt. RNAs have been produced during RNAi in vitro using an extract prepared from Drosophila embryos (Zamore et al. (2000) *Cell* 101 25-33). Putative guide RNAs can also be produced in extracts from Drosophila S2 cells (**FIG. 5*a***). With the goal of understanding the mechanism of post-transcriptional gene silencing, we have undertaken both biochemical fractionation and candidate gene approaches to identify the enzymes that execute each step of RNAi.

[0212] Our previous studies resulted in the partial purification of a nuclease, RISC, that is an effector of RNA interference. See Example 1. This enzyme was isolated from Drosophila S2 cells in which RNAi had been initiated in vivo by transfection with dsRNA. We first sought to determine whether the RISC enzyme and the enzyme that initiates RNAi via processing of dsRNA into 22mers are distinct activities. RISC activity could be largely cleared from extracts by high-speed centrifugation (100,000×g for 60 min.) while the activity that produces 22mers remained in the supernatant (**FIG. 5***b,c*). This simple fractionation indicated that RISC and the 22mer-generating activity are separable and thus distinct enzymes. However, it seems likely that they might interact at some point during the silencing process.

[0213] RNAse III family members are among the few nucleases that show specificity for double-stranded RNA (Nicholson (1999) *FEMS Microbiol Rev* 23: 371-390). Analysis of the Drosophila and *C. elegans* genomes reveals several types of RNAse III enzymes. First is the canonical RNAse III which contains a single RNAse III signature motif and a double-stranded RNA binding domain (dsRBD; e.g. RNC_CAEEL). Second is a class represented by Drosha (Filippov et al. (2000) *Gene* 245: 213-221), a Drosophila enzyme that contains two RNAse III motifs and a dsRBD (CeDrosha in *C. elegans*). A third class contains two RNAse III signatures and an amino terminal helicase domain (e.g. Drosophila CG4792, CG6493, *C. elegans* K12H4.8), and these had previously been proposed by Bass as candidate RNAi nucleases (Bass (2000) *Cell* 101: 235-238). Representatives of all three classes were tested for the ability to produce discrete, ~22 nt. RNAs from dsRNA substrates.

[0214] Partial digestion of a 500 nt. cyclin E dsRNA with purified, bacterial RNAse III produced a smear of products while nearly complete digestion produced a heterogeneous group of 11-17 nucleotide RNAs (not shown). In order to test the dual-RNAse III enzymes, we prepared T7 epitope-tagged versions of Drosha and CG4792. These were expressed in transfected S2 cells and isolated by immunoprecipitation using antibody-agarose conjugates. Treatment of the dsRNA with the CG4792 immunoprecipitate yielded ~22 nt. fragments similar to those produced in either S2 or embryo extracts (**FIG. 6***a*). Neither activity in extract nor activity in immunoprecipitates depended on the sequence of the RNA substrate since dsRNAs derived from several genes were processed equivalently (see Supplement 1). Negative results were obtained with Drosha and with immunoprecipitates of a DExH box helicase (Homeless (Gillespie et al. (1995) *Genes and Development* 9: 2495-2508); see **FIG. 6***a,b*). Western blotting confirmed that each of the tagged proteins was expressed and immunoprecipitated similarly (see Supplement 2). Thus, we conclude that CG4792 may carry out the initiation step of RNA interference by producing ~22 nt. guide sequences from dsRNAs. Because of its ability to digest dsRNA into uniformly sized, small RNAs, we have named this enzyme Dicer (Dcr). Dicer mRNA is expressed in embryos, in S2 cells, and in adult flies, consistent with the presence of functional RNAi machinery in all of these contexts (see Supplement 3).

[0215] The possibility that Dicer might be the nuclease responsible for the production of guide RNAs from dsRNAs prompted us to raise an antiserum directed against the carboxy-terminus of the Dicer protein (Dicer-1, CG4792). This antiserum could immunoprecipitate a nuclease activity from either Drosophila embryo extracts or from S2 cell lysates that produced ~22 nt. RNAs from dsRNA substrates (**FIG. 6C**). The putative guide RNAs that are produced by the Dicer-1 enzyme precisely comigrate with 22mers that are produced in extract and with 22mers that are associated with the RISC enzyme (**FIG. 6D,F**). It had previously been shown that the enzyme that produced guide RNAs in Drosophila embryo extracts was ATP-dependent (Zamore et al. (2000) *Cell* 101 25-33). Depletion of this cofactor resulted in an ~6-fold lower rate of dsRNA cleavage and in the production of RNAs with a slightly lower mobility. Of interest was the fact that both Dicer-I immunoprecipitates and extracts from S2 cells require ATP for the production of 22mers (**FIG. 6D**). We do not observe the accumulation of lower mobility products in these cases, although we do routinely observe these in ATP-depleted embryo extracts. The requirement of this nuclease for ATP is a quite unusual property. We hypothesize that this requirement could indicate that the enzyme may act processively on the dsRNA, with the helicase domain harnessing the energy of ATP hydrolysis both for unwinding guide RNAs and for translocation along the substrate.

[0216] Efficient induction of RNA interference in *C. elegans* and in Drosophila has several requirements. For example, the initiating RNA must be double-stranded, and it must be several hundred nucleotides in length. To determine whether these requirements are dictated by Dicer, we characterized the ability of extracts and of immunoprecipitated enzyme to digest various RNA substrates. Dicer was inactive against single stranded RNAs regardless of length (see Supplement 4). The enyzme could digest both 200 and 500 nucleotide dsRNAs but was significantly less active with shorter substrates (see Supplement 4). Double-stranded RNAs as short as 35 nucleotides could be cut by the enzyme, albeit very inefficiently (data not shown). In contrast, *E. coli* RNAse III could digest to completion dsRNAs of 35 or 22 nucleotides (not shown). This suggests that the substrate preferences of the Dicer enzyme may contribute to but not wholly determine the size dependence of RNAi.

[0217] To determine whether the Dicer enzyme indeed played a role in RNAi in vivo, we sought to deplete Dicer activity from S2 cells and test the effect on dsRNA-induced gene silencing. Transfection of S2 cells with a mixture of dsRNAs homologous to the two Drosophila Dicer genes (CG4792 and CG6493) resulted in an ~6-7 fold reduction of Dicer activity either in whole cell lysates or in Dicer-1 immunoprecipitates (**FIG. 7A,B**). Transfection with a control dsRNA (murine caspase 9) had no effect. Qualitatively similar results were seen if Dicer was examined by Northern blotting (not shown). Depletion of Dicer in this manner substantially compromised the ability of cells to silence subsequently an exogenous, GFP transgene by RNAi (**FIG. 7C**). These results indicate that Dicer is involved in RNAi in vivo. The lack of complete inhibition of silencing could result from an incomplete suppression of Dicer (which is itself required for RNAi) or could indicate that in vivo, guide RNAs can be produced by more than one mechanism (e.g. through the action of RNA-dependent RNA polymerases).

[0218] Our results indicate that the process of RNA interference can be divided into at least two distinct steps. According to this model, initiation of PTGS would occur upon processing of a double-stranded RNA by Dicer into

~22 nucleotide guide sequences, although we cannot formally exclude the possibility that another, Dicer-associated nuclease may participate in this process. These guide RNAs would be incorporated into a distinct nuclease complex (RISC) that targets single-stranded mRNAs for degradation. An implication of this model is that guide sequences are themselves derived directly from the dsRNA that triggers the response. In accord with this model, we have demonstrated that ^{32}P-labeled, exogenous dsRNAs that have been introduced into S2 cells by transfection are incorporated into the RISC enzyme as 22 mers (**FIG. 7E**). However, we cannot exclude the possibility that RNA-dependent RNA polymerases might amplify 22mers once they have been generated or provide an alternative method for producing guide RNAs.

[**0219**] The structure of the Dicer enzyme provokes speculation on the mechanism by which the enzyme might produce discretely sized fragments irrespective of the sequence of the dsRNA (see Supplement 1, **FIG. 8***a*). It has been established that bacterial RNAse III acts on its substrate as a dimer (Nicholson (1999) *FEMS Microbiol Rev* 23: 371-390; Robertson et al. (1968) *J Biol Chem* 243: 82-91; Dunn (1976) *J Biol Chem* 251: 3807-3814). Similarly, a dimer of Dicer enzymes may be required for cleavage of dsRNAs into ~22 nt. pieces. According to one model, the cleavage interval would be determined by the physical arrangement of the two RNAse III domains within Dicer enzyme (**FIG. 8***a*). A plausible alternative model would dictate that cleavage was directed at a single position by the two RIII domains in a single Dicer protein. The 22 nucleotide interval could be dictated by interaction of neighboring Dicer enzymes or by translocation along the mRNA substrate. The presence of an integral helicase domain suggests that the products of Dicer cleavage might be single-stranded 22 mers that are incorporated into the RISC enzyme as such.

[**0220**] A notable feature of the Dicer family is its evolutionary conservation. Homologs are found in *C. elegans* (K12H4.8), Arabidopsis (e.g., CARPEL FACTORY (Jacobson et al. (1999) *Development* 126: 5231-5243), T25K16.4, AC012328_1), mammals (Helicase-MOI (Matsuda et al. (2000) *Biochim Biophys Acta* 1490: 163-169) and *S. pombe* (YC9A_SCHPO) (**FIG. 8***b*, see Supplements 6,7 for sequence comparisons). In fact, the human Dicer family member is capable of generating ~22 nt. RNAs from dsRNA substrates (Supplement 5) suggesting that these structurally similar proteins may all share similar biochemical functions. It has been demonstrated that exogenous dsRNAs can affect gene function in early mouse embryos (Wianny et al. (2000) *Nature Cell Biology* 2: 70-75), and our results suggest that this regulation may be accomplished by an evolutionarily conserved RNAi machinery.

[**0221**] In addition to RNAseIII and helicase motifs, searches of the PFAM database indicate that each Dicer family member also contains a ZAP domain (**FIG. 8***c*) (Sonnhammer et al. (1997) *Proteins* 28: 405-420). This sequence was defined based solely upon its conservation in the Zwille/ARGONAUTE/Piwi family that has been implicated in RNAi by mutations in *C. elegans* (Rde-1) and Neurospora (Qde-2) (Tabara et al. (1999) *Cell* 99: 123-132; Catalanotto et al (2000) *Nature* 404: 245). Although the function of this domain is unknown, it is intriguing that this region of homology is restricted to two gene families that participate in dsRNA-dependent silencing. Both the ARGONAUTE and Dicer families have also been implicated in common biological processes, namely the determination of stem-cell fates. A hypomorphic allele of carpel factory, a member of the Dicer family in Arabidopsis, is characterized by increased proliferation in floral meristems (Jacobsen et al. (1999) *Development* 126 5231-5243). This phenotype and a number of other characteristic features are also shared by Arabidopsis ARGONAUTE (ago1-1) mutants (Bohmert et al. (1998) *EMBO J* 17: 170-180; C. Kidner and R. Martienssen, pers. comm.). These genetic analyses begin to provide evidence that RNAi may be more than a defensive response to unusual RNAs but may also play important roles in the regulation of endogenous genes.

[**0222**] With the identification of Dicer as a catalyst of the initiation step of RNAi, we have begun to unravel the biochemical basis of this unusual mechanism of gene regulation. It will be of critical importance to determine whether the conserved family members from other organisms, particularly mammals, also play a role in dsRNA-mediated gene regulation.

[**0223**] Methods:

[**0224**] Plasmid constructs. A full-length cDNA encoding Drosha was obtained by PCR from an EST sequenced by the Berkeley Drosophila genome project. The Homeless clone was a gift from Gillespie and Berg (Univ. Washington). The T7 epitope-tag was added to the amino terminus of each by PCR, and the tagged cDNAs were cloned into pRIP, a retroviral vector designed specifically for expression in insect cells (E. Bernstein, unpublished). In this vector, expression is driven by the *Orgyia pseudotsugata* IE2 promoter (Invitrogen). Since no cDNA was available for CG4792/Dicer, a genomic clone was amplified from a bacmid (BACR23F10; obtained from the BACPAC Resource Center in the Dept. of Human Genetics at the Roswell Park Cancer Institute). Again, during amplification, a T7 epitope tag was added at the amino terminus of the coding sequence. The human Dicer gene was isolated from a cDNA library prepared from HaCaT cells (GJH, unpublished). A T7-tagged version of the complete coding sequence was cloned into pCDNA3 (Invitrogen) for expression in human cells (LinX-A).

[**0225**] Cell culture and extract preparation. S2 and embryo culture. S2 cells were cultured at 27° C. in 5% CO_2 in Schneider's insect media supplemented with 10% heat inactivated fetal bovine serum (Gemini) and 1% antibiotic-antimycotic solution (Gibco BRL). Cells were harvested for extract preparation at 10×10^6 cells/ml. The cells were washed 1× in PBS and were resuspended in a hypotonic buffer (10 mM Hepes pH 7.0, 2 mM $MgCl_2$, 6 mM PME) and dounced. Cell lysates were spun 20,000×g for 20 minutes. Extracts were stored at −80° C. Drosophila embryos were reared in fly cages by standard methodologies and were collected every 12 hours. The embryos were dechorionated in 50% chlorox bleach and washed thoroughly with distilled water. Lysis buffer (10 mM Hepes, 10 mM KCl, 1.5 mM $MgCl_2$, 0.5 mM EGTA, 10 mM β-glycerophosphate, 1 mM DTT, 0.2 mM PMSF) was added to the embryos, and extracts were prepared by homogenization in a tissue grinder. Lysates were spun for two hours at 200,000×g and were frozen at −80° C. LinX-A cells, a highly-transfectable derivative of human 293 cells, (Lin Xie and GJH, unpublished) were maintained in DMEM/10% FCS.

[0226] Transfections and immunoprecipitations. S2 cells were transfected using a calcium phosphate procedure essentially as previously described (Hammond et al. (2000) *Nature* 404: 293-296). Transfection rates were ~90% as monitored in controls using an in situ β-galactosidase assay. LinX-A cells were also transfected by calcium phosphate co-precipitation. For immunoprecipitations, cells (~5×10^6 per IP) were transfected with various clones and lysed three days later in IP buffer (125 mM KOAc, 1 mM MgOAc, 1 mM $CaCl_2$, 5 mM EGTA, 20 mM Hepes pH 7.0, 1 mM DTT, 1% NP-40 plus Complete protease inhibitors (Roche)). Lysates were spun for 10 minutes at 14,000×g and supernatants were added to T7 antibody-agarose beads (Novagen). Antibody binding proceeded for 4 hours at 4° C. Beads were centrifuged and washed in lysis buffer three times, and once in reaction buffer. The Dicer antiserum was raised in rabbits using a KLH-conjugated peptide corresponding to the C-terminal 8 amino acids of Drosophila Dicer-1 (CG4792).

[0227] Cleavage reactions. RNA preparation. Templates to be transcribed into dsRNA were generated by PCR with forward and reverse primers, each containing a T7 promoter sequence. RNAs were produced using Riboprobe (Promega) kits and were uniformly labeling during the transcription reaction with ^{32}P-UTP. Single-stranded RNAs were purified from 1% agarose gels. dsRNA cleavage. Five microliters of embryo or S2 extracts were incubated for one hour at 30° C. with dsRNA in a reaction containing 20 mM Hepes pH 7.0, 2 mM MgOAc, 2 mM DTT, 1 mM ATP and 5% Superasin (Ambion). Immunoprecipitates were treated similarly except that a minimal volume of reaction buffer (including ATP and Superasin) and dsRNA were added to beads that had been washed in reaction buffer (see above). For ATP depletion, Drosophila embryo extracts were incubated for 20 minutes at 30° C. with 2 mM glucose and 0.375 U of hexokinase (Roche) prior to the addition of dsRNA.

[0228] Northern and Western analysis. Total RNA was prepared from Drosophila embryos (0-12 hour), from adult flies, and from S2 cells using Trizol (Lifetech). Messenger RNA was isolated by affinity selection using magnetic oligo-dT beads (Dynal). RNAs were electrophoresed on denaturing formaldehyde/agarose gels, blotted and probed with randomly primed DNAs corresponding to Dicer. For Western analysis, T7-tagged proteins were immunoprecipitated from whole cell lysates in IP buffer using anti-T7-antibody-agarose conjugates. Proteins were released from the beads by boiling in Laemmli buffer and were separated by electrophoresis on 8% SDS PAGE. Following transfer to nitrocellulose, proteins were visualized using an HRP-conjugated anti-T7 antibody (Novagen) and chemiluminescent detection (Supersignal, Pierce).

[0229] RNAi of Dicer. Drosophila S2 cells were transfected either with a dsRNA corresponding to mouse caspase 9 or with a mixture of two dsRNAs corresponding to Drosophila Dicer-1 and Dicer-2 (CG4792 and CG6493). Two days after the initial transfection, cells were again transfected with a mixture containing a GFP expression plasmid and either luciferase dsRNA or GFP dsRNA as previously described (Hammond et al. (2000) *Nature* 404: 293-296). Cells were assayed for Dicer activity or fluorescence three days after the second transfection. Quantification of fluorescent cells was done on a Coulter EPICS cell sorter after fixation. Control transfections indicated that Dicer activity was not affected by the introduction of caspase 9 dsRNA.

EXAMPLE 3

A Simplified Method for the Creation of Hairpin Constructs for RNA Interference

[0230] In numerous model organisms, double stranded RNAs have been shown to cause effective and specific suppression of gene function (Bosher and Labouesse (2000) *Nature* Cell Biology 2: E31-E36). This response, termed RNA interference or post-transcriptional gene silencing, has evolved into a highly effective reverse genetic tool in *C. elegans*, Drosophila, plants and numerous other systems. In these cases, double-stranded RNAs can be introduced by injection, transfection or feeding; however, in all cases, the response is both transient and systemic. Recently, stable interference with gene expression has been achieved by expression of RNAs that form snap-back or hairpin structures (Fortier and Belote (2000) *Genesis* 26: 240-244; Kennerdell and Carthew (2000) *Nature Biotechnology* 18: 896-898; Lam and Thummel (2000) *Current Biology* 10: 957-963; Shi et al. (2000) *RNA* 6: 1069-1076; Smith et al. (2000) *Nature* 407: 319-320; Tavemarakis et al. (2000) *Nature Genetics* 24: 180-183). This has the potential not only to allow stable silencing of gene expression but also inducible silencing as has been observed in trypanosomes and adult Drosophila (Fortier and Belote (2000) *Genesis* 26: 240-244; Lam and Thummel (2000) *Current Biology* 10: 957-963; Shi et al. (2000) *RNA* 6: 1069-1076). The utility of this approach is somewhat hampered by the difficulties that arise in the construction of bacterial plasmids containing the long inverted repeats that are necessary to provoke silencing. In a recent report, it was stated that more than 1,000 putative clones were screened to identify the desired construct (Tavemarakis et al. (2000) *Nature Genetics* 24: 180-183).

[0231] The presence of hairpin structures often induces plasmid rearrangement, in part due to the *E. coli* sbc proteins that recognize and cleave cruciform DNA structures (Connelly et al. (1996) *Genes Cell* 1: 285-291). We have developed a method for the construction of hairpins that does not require cloning of inverted repeats, per se. Instead, the fragment of the gene that is to be silenced is cloned as a direct repeat, and the inversion is accomplished by treatment with a site-specific recombinase, either in vitro (or potentially in vivo) (see **FIG. 27**). Following recombination, the inverted repeat structure is stable in a bacterial strain that lacks an intact SBC system (DL759). We have successfully used this strategy to construct numerous hairpin expression constructs that have been successfully used to provoke gene silencing in Drosophila cells.

[0232] In the following examples, we use this method to express long dsRNAs in a variety of mammalian cell types. We show that such long dsRNAs mediate RNAi in a variety of cell types. Additionally, since the vector described in **FIG. 27** contains a selectable marker, dsRNAs produced in this manner can be stably expressed in cells. Accordingly, this method allows not only the examination of transient effects of RNA suppression in a cell, but also the effects of stable and prolonged RNA suppression.

[0233] Methods:

[0234] Plasmids expressing hairpin RNAs were constructed by cloning the first 500 basepairs of the GFP coding region into the FLIP cassette of pRIP-FLIP as a direct repeat. The FLIP cassette contains two directional cloning sites, the second of which is flanked by LoxP sites. The Zeocin gene, present between the cloning sites, maintains selection and stability. To create an inverted repeat for hairpin production, the direct repeat clones were exposed to Cre recombinase (Stratagene) in vitro and, afterwards, transformed into DL759 *E. Coli*. These bacteria permit the replication of DNA containing cruciform structures, which tend to form inverted repeats.

EXAMPLE 4

Long dsRNAs Suppress Gene Expression in Mammalian Cells

[0235] Previous experiments have demonstrated that dsRNA, produced using a variety of methods including via the construction of hairpins, can suppress gene expression in Drosophila cells. We now demonstrate that dsRNA can also suppress gene expression in mammalian cells in culture. Additionally, the power of RNAi as a genetic tool would be greatly enhanced by the ability to engineer stable silencing of gene expression. We therefore undertook an effort to identify mammalian cells in which long dsRNAs could be used as RNAi triggers in the hope that these same cell lines would provide a platform upon which to develop stable silencing strategies. We demonstrate that RNA suppression can be mediated by stably expressing a long hairpin in a mammalian cell line. The ability to engineer stable silencing of gene expression in cultured mammalian cells, in addition to the ability to transiently silence gene expression, has many important applications.

[0236] A. RNAi in Pluripotent Murine P19 Cells.

[0237] We first sought to determine whether long dsRNA triggers could induce sequence-specific silencing in cultured murine cells, both to develop this approach as a tool for probing gene function and to allow mechanistic studies of dsRNA-induced silencing to be propagated to mammalian systems. We, therefore, attempted to extend previous studies in mouse embryos (Wianny et al. 2000, Nat. Cell Biol. 2: 70-75; Svoboda et al. 2000, Development 127: 4147-4156) by searching for RNAi-like mechanisms in pluripotent, embryonic cell types. We surveyed a number of cell lines of embryonic origin for the degree to which generalized suppression of gene expression occurred upon introduction of dsRNA. As an assay, we tested the effects of dsRNA on the expression of GFP as measured in situ by counting fluorescent cells. As expected, in both human embryonic kidney cells (293) and mouse embryo fibroblasts, GFP expression was virtually eliminated irrespective of the sequence of the cotransfected dsRNA. In some pluripotent teratocarcinoma and teratoma cell lines (e.g., N-Teral, F9), the PKR response was attenuated but still evident; however, in contrast, transfection of nonhomologous dsRNAs had no effect on the expression of reporter genes (e.g., GFP or luciferase) either in mouse embryonic stem cells or in p19 embryonal carcinoma cells (**FIG. 28**).

[0238] Transfection of P19 embryonal carcinoma cells with GFP in the presence of cognate dsRNA corresponding to the first ≈500 nts of the GFP coding sequence had a strikingly different effect. GFP expression was eliminated in the vast majority of cotransfected cells (**FIG. 28**), suggesting that these cultured murine cells might respond to dsRNA in a manner similar to that which we had previously demonstrated in cultured, Drosophila S2 cells (Hammond et al. 2000, Nature 404: 293-296).

[0239] To quantify the extent to which dsRNA could induce sequence-specific gene silencing, we used a dual luciferase reporter assay similar to that which had first been used to demonstrate RNAi in Drosophila embryo extracts (Tuschl et al. 1999, Genes Dev. 13: 3191-3197). P19 EC cells were transfected with a mixture of two plasmids that individually direct the expression of firefly luciferase and Renilla luciferase. These were cotransfected with no dsRNA, with dsRNA that corresponds to the first ≈500 nts of the firefly luciferase, or with dsRNA corresponding to the first ≈500 nts of GFP as a control. Cotransfection with GFP dsRNA gave luciferase activities that were similar to the no-dsRNA control, both in the firefly/Renilla activity ratio and in the absolute values of both activities. In contrast, in cells that received the firefly luciferase dsRNA, the ratio of firefly to Renilla luciferase activity was reduced by up to 30-fold (250 ng, **FIG. 29B**). For comparison, we carried out an identical set of experiments in Drosophila S2 cells. Although qualitatively similar results were obtained, the silencing response was more potent. At equivalent levels of dsRNA, S2 cells suppressed firefly luciferase activity to virtually background levels.

[0240] The complementary experiment, in which dsRNA was homologous to Renilla luciferase, was also performed. Again, in this case, suppression of the expression of the Renilla enzyme was ≈10-fold (**FIG. 29D**). Thus, the dsRNA response in P19 cells was flexible, and the silencing machinery was able to adapt to dsRNAs directed against any of the reporters that were tested.

[0241] We took two approaches to test whether this response was specific for dsRNA. Pretreatment of the trigger with purified RNase III, a dsRNA-specific ribonuclease, before transfection greatly reduced its ability to provoke silencing. Furthermore, transfection of cells with single-stranded antisense RNAs directed against either firefly or Renilla luciferase had little or no effect on expression of the reporters (**FIGS. 29C and 29D**). Considered together, these results provided a strong indication that double-stranded RNAs provoke a potent and specific silencing response in P19 embryonal carcinoma cells. Efficient silencing could be provoked with relatively low concentrations of dsRNA (25 ng/ml culture media; see **FIG. 29A**). The response was concentration-dependent, with maximal suppression of ≈20-fold being achieved at a dose of 1.5 μg/ml culture media. Silencing was established rapidly and was evident by 9 h posttransfection (the earliest time point examined). Furthermore, the response persisted without significant changes in the degree of suppression for up to 72 h following a single dose of dsRNA.

[0242] **FIG. 30** further shows wildtype P19 cells which have been co-transfected with either RFP or GFP (right panel). Note the robust expression of RFP or GFR respectively approximately 42 hours post-transfection. We isolated P19 clones which stably express a 500 nt. GFP hairpin. Such clones were then transfected with either RFP or GFP, and

expression of RFP or GFP was assessed by visual inspection of the cells. The left panel demonstrates that a 500 nt GFP hairpin specifically suppresses expression of GFP in P19 cells.

[0243] B. RNAi in Embryonic Stem Cells.

[0244] To assess whether the presence of a sequence-specific response to dsRNA was a peculiarity of P19 cells or whether it also extended to normal murine embryonic cells, we performed similar silencing assays in mouse embryonic stem cells. Cotransfection of embryonic stem cells with noncognate dsRNAs (e.g., GFP), again, had no dramatic effect on either the absolute values or the ratios of Renilla and firefly luciferase activity (**FIG. 31**). However, transfection with either firefly or Renilla luciferase dsRNA dramatically and specifically reduced the activity of the targeted enzyme (**FIG. 31**).

[0245] This result suggests that RNAi can operate in multiple murine cell types of embryonic origin, including normal embryonic stem cells. The ability to provoke silencing in a cell type that is normally used for the generation of genetic, mosaic animals suggests the possibility of eventually testing the biological effects of silencing both in culture and in reconstituted animal models. Our ability to successfully manipulate ES cell via RNAi allows the use of RNAi in the generation of transgenic and knock-out mice.

[0246] C. RNAI in Murine Somatic Cells.

[0247] RNAi effector pathways are likely to be present in mammalian somatic cells, based on the ability of siRNAs to induce transient silencing (Elbashir et al. 2001, Nature 411: 494-498). Furthermore, we have shown that RNAi initiator and effector pathways clearly exist in embryonic cells that can enforce silencing in response to long dsRNA triggers. We therefore sought to test whether the RNAi machinery might exist intact in some somatic cell lines.

[0248] Transfection of HeLa cells with luciferase reporters in combination with long dsRNA triggers caused a nearly complete suppression of activity, irrespective of the RNA sequence. In a murine myoblast cell line, C2C12, we noted a mixture of two responses. dsRNAs homologous to firefly luciferase provoked a sequence-specific effect, producing a degree of suppression that was slightly more potent than was observed upon transfection with cognate ≈21-nt siRNA (Elbashir et al. 2001, Nature 411: 494-498) (see **FIG. 32A**). However, with long dsRNA triggers, the specific effect was superimposed upon a generalized suppression of reporter gene expression that was presumably because of PKR activation (**FIG. 32B**).

[0249] Numerous mammalian viruses have evolved the ability to block PKR as an aid to efficient infection. For example, adenoviruses express VA RNAs, which mimic dsRNA with respect to binding but not to activation of PKR (Clarke et al. 1995, RNA 1: 7-20). Vaccinia virus uses two strategies to evade PKR. The first is expression of E3L, which binds and masks dsRNAs (Kawagishi-Kobayashi et al. 2000, Virology 276: 424-434). The second is expression of K3L, which binds and inhibits PKR via its ability to mimic the natural substrate of this enzyme, eIF2α (Kawagishi-Kobayashi et al. 2000, supra).

[0250] Transfection of C2C12 cells with a vector that directs K3L expression attenuates the generalized repression of reporter genes in response to dsRNA. However, this protein had no effect on the magnitude of specific inhibition by RNAi (**FIG. 32C**).

[0251] **FIG. 33** further shows the results of a transient co-transfection assay performed in Hela cells, CHO cells, and P19 cells. The cell lines were each transfected with plasmids expressing *Photinus pyralis* (firefly) and *Renila reniformis* (sea pansy) luciferases. The cells lines were additionally transfected with 400 ng of 500nt dsRNAs corresponding to either firefly luciferase (dsLUC) or dsGFP. The results demonstrate that dsRNA can specifically mediate suppression in a multiple mammalian cells types in culture.

[0252] These results raise the possibility that, at least in some cell lines and/or cell types, blocking nonspecific responses to dsRNA will enable the use of long dsRNAs for the study of gene function. This might be accomplished through the use of viral inhibitors, as described here, or through the use of cells isolated from animals that are genetically modified to lack undesirable responses.

[0253] D. Stable Suppression of Gene Expression Using RNAi.

[0254] To date, dsRNAs have been used to induce sequence-specific gene silencing in either cultured mammalian cells or in embryos only in a transient fashion. However, the most powerful applications of genetic manipulation are realized only with the creation of stable mutants. The ability to induce silencing by using long dsRNAs offers the opportunity to translate into mammalian cells work from model systems such as Drosophila, plants, and *C. elegans* wherein stable silencing has been achieved by enforced expression of hairpin RNAs (Kennerdell et al. 2000, Nat. Biotechnol. 18: 896-898; Smith et al. 2000, Nature 407: 319-320; Tavemarakis et al. 2000, Nat. Genet. 24: 180-183).

[0255] P19 EC cells were transfected with a control vector or with an expression vector that directs expression of a ≈500-nt GFP hairpin RNA from an RNA polymerase II promoter (cytomegalovirus). Colonies arising from cells that had stably integrated either construct were selected and expanded into clonal cell lines. Each cell line was assayed for persistent RNAi by transient cotransfection with a mixture of two reporter genes, dsRED to mark transfected cells and GFP to test for stable silencing.

[0256] Transfection of clonal P19 EC cells that had stably integrated the control vector produced equal numbers of red and green cells, as would be expected in the absence of any specific silencing response (**FIG. 34B**), whereas cells that express the GFP hairpin RNA gave a very different result. These cells expressed the dsRED protein with an efficiency comparable to that observed in cells containing the control vector. However, the cells failed to express the cotransfected GFP reporter (**FIG. 34B**). These data provide a strong indication that continuous expression of a hairpin dsRNA can provoke stable, sequence-specific silencing of a target gene.

[0257] In Drosophila S2 cells and *C. elegans*, RNAi is initiated by the Dicer enzyme, which processes dsRNA into ≈22-nt siRNAs (Bernstein et al. 2001, Nature 409: 363-366; Grishok et al. 2001, Cell 106: 23-34; Hutvagner et al. 2001, Science 293: 834-838; Ketting et al. 2001, Genes Dev. 15: 2654-2659; Knight et al. 2001, Science 293: 2269-2271). In both, S2 cells and *C. elegans* experiments by using dsRNA

to target Dicer suppress the RNAi response. Whether Dicer plays a central role in hairpin-induced gene silencing in P19 cells was tested by transfecting P19 cells stably transfected with GFP hairpin constructs with mouse Dicer dsRNA. Treatment with Dicer dsRNA, but not control dsRNA, resulted in derepression of GFP (**FIG. 34C**).

[**0258**] E. dsRNA Induces Posttranscriptional Silencing.

[**0259**] A key feature of RNAi is that it exerts its effect at the posttranscriptional level by destruction of targeted mRNAs (Hammond et al. 2001, Nat. Rev. Genet. 2: 110-119). To test whether dsRNAs induced silencing in mouse cells via posttranscriptional mechanisms, we used an assay identical to that, used initially to characterize RNAi responses in Drosophila embryo extracts (Tuschl et al. 1999, Genes Dev. 13: 3191-3197). We prepared lysates from P19 EC cells that were competent for in vitro translation of capped mRNAs corresponding to Renilla and firefly luciferase. Addition of nonspecific dsRNAs to these extracts had no substantial effect on either the absolute amount of luciferase expression or on the ratio of firefly to Renilla luciferase (**FIG. 35**). In contrast, addition of dsRNA homologous to the firefly luciferase induced a dramatic and dose-dependent suppression of activity. Addition of RNA corresponding to only the antisense strand of the dsRNA had little effect, comparable to a nonspecific dsRNA control, and pretreatment of the dsRNA silencing trigger with RNase III greatly reduced its potential to induce silencing in vitro. A second hallmark of RNAi is the production of small, ≈22-nt siRNAs, which determine the specificity of silencing. We found that such RNA species were generated from dsRNA in P19 cell extracts (**FIG. 34**D, in vitro), indicative of the presence of a mouse Dicer activity. These species were also produced in cells that stably express GFP hairpin RNAs (**FIG. 34**D, in vivo). Considered together, the posttranscriptional nature of dsRNA-induced silencing, the association of silencing with the production of ≈22-nt siRNAs, and the dependence of this response on Dicer, a key player in the RNAi pathway, strongly suggests that dsRNA suppresses gene expression in murine cells via a conventional RNAi mechanism.

[**0260**] F RNAi-Mediated Gene Silencing is Specific and Requires dsRNAs.

[**0261**] We carried out experiments to verify that the suppressive effects observed in the in vitro system were specific to double stranded RNA. Briefly, experiments were performed in accordance with the methods outlined above. Either dsRNA (ds), single-stranded RNA (ss), or antisense-RNA (as) corresponding to firefly (FF) or Renilla (Ren) luciferase was added to the translation reaction. Following reactions performed at 30° C. for 1 hour, dual luciferase assays were performed using an Analytical Scientific Instruments model 3010 Luminometer.

[**0262**] **FIG. 36** summarizes the results of these experiments which demonstrate that the suppression of gene expression observed in this in vitro assay is specific for dsRNA. These results further support the conclusion that dsRNA suppresses gene expression in this mammalian in vitro system in a manner consistent with post-transcriptional silencing.

[**0263**] G. Mammalian Cells Soaked with dsRNAs Results in Gene Silencing.

[**0264**] Studies of post-transcriptional silencing in invertebrates have demonstrated that transfection or injection of the dsRNA is not necessary to achieve the suppressive affects. For example, dsRNA suppression in *C. elegans* can be observed by either soaking the worms in dsRNA, or by feeding the worms bacteria expressing the dsRNA of interest. We addressed whether dsRNA suppression in mammalian cells could be observed without transfection of the dsRNA. Such a result would present additional potential for easily using dsRNA suppression in mammalian cells, and would also allow the use of dsRNA to suppress gene expression in cell types which have been difficult to transfect (i.e., cell types with a low transfection efficiency, or cell types which have proven difficult to transfect at all).

[**0265**] P19 cells were grown in 6-well tissue culture plates to approximately 60% confluency in growth media (αMEM/10% FBS). Varying concentrations of firefly dsRNA were added to the cultures, and cells were cultured for 12 hours in growth media+dsRNA. Cells were then transfected with plasmids expressing firefly or sea pansy luciferase, as described in detail above. Dual luciferase assays were carried out 12 hours post-transfection using an Analytical Scientific Instruments model 3010 Luminometer.

[**0266**] **FIG. 37** summarizes these results which demonstrate that dsRNA can suppress gene expression in mammalian cells without transfection. Culturing cells in the presence of dsRNA resulted in a dose dependent suppression of firefly luciferase gene expression.

[**0267**] Methods:

[**0268**] Cell Culture. P19 mouse embryonic carcinoma cells (American Type Culture Collection, CRL-1825) were cultured in α-MEM (GIBCO/BRL) supplemented with 10% heat-inactivated FBS and 1% antibiotic/antimycotic solution (GIBCO/BRL). Mouse embryo stem cells (J1, provided by S. Kim, Cold Spring Harbor Laboratory) were cultured in DMEM containing ESgro (Chemicon) according to the manufacturer's instructions. C2C12 murine myoblast cells (gift of N. Tonks, Cold Spring Harbor Laboratory) were cultured in DMEM (GIBCO/BRL) supplemented with 10% heat-inactivated FBS and 1% antibiotic/antimycotic solution (GIBCO/BRL).

[**0269**] RNA Preparation. For the production of dsRNA, transcription templates were generated by PCR; they contained T7 promoter sequences on each end of the template (see Hammond et al. 2000, Nature 404: 293-296). dsRNAs were prepared by using the RiboMax kit (Ambion, Austin, Tex.). Firefly and Renilla luciferase mRNA transcripts were synthesized by using the Riboprobe kit (Promega) and were gel purified before use.

[**0270**] Transfection and Gene Silencing Assays. Cells were transfected with indicated amounts of dsRNA and plasmid DNA by using FuGENE6 (Roche Biochemicals) according to the manufacturer's instructions. Cells were transfected at 50-70% confluence in 12-well plates containing either 1 or 2 ml of medium per well. Dual luciferase assays (Promega) were carried out by cotransfecting cells with plasmids contain firefly luciferase under the control of SV40 promoter (pGL3-Control, Promega) and Renilla luciferase under the control of the SV40 early enhancer/

promoter region (pSV40, Promega). These plasmids were cotransfected by using a 1:1 or 10:1 ratio of pGL3-control (250 ng/well) to pRL-SV40. Both ratios yielded similar results. For some experiments, cells were transfected with vectors that direct expression of enhanced green fluorescent protein (EGFP)-US9 fusion protein (Kalejta et al. 1999, Exp. Cell Res. 248: 322-328) or red fluorescent protein (RFP) (pDsRed N1, CLONTECH). RNAi in S2 cells was performed as described (Hammond et al. 2000, Nature 404: 293-296).

[0271] Plasmids expressing hairpin RNAs (RNAs with a self-complimentary stem loop) were constructed by cloning the first 500 bp of the EGFP coding region (CLONTECH) into the FLIP cassette of pRIP-FLIP as a direct repeat. The FLIP cassette contains two directional cloning sites, the second of which sports flanking LoxP sites (see **FIG. 35A**). The Zeocin gene (Stratagene), present between the cloning sites, maintains selection and, thus, stability of the FLIP cassette. The FLIP cassette containing EGFP direct repeats was subcloned into pcDNA3 (Invitrogen). To create an inverted repeat for hairpin production, EGFP direct repeat clones were exposed to Cre recombinase (Stratagene) in vitro and, afterward, transformed into DL759 *Escherichia coli* (Connelly et al. 1996, Genes Cells 1: 285-291). These bacteria permit the replication of DNA containing cruciform structures, which tend to form from inverted repeats. DL759 transformants were screened for plasmids containing inverted repeats (≈50%).

[0272] Silencing of Dicer was accomplished by using a dsRNA comprising exon 25 of the mouse Dicer gene and corresponding to nucleotides 5284-5552 of the human Dicer cDNA.

[0273] In Vitro Translation and in Vitro Dicer Assays. Logarithmically growing cells were harvested in PBS containing 5 mM EGTA washed twice in PBS and once in hypotonic buffer (10 mM Hepes, pH 7.3/6 mM β-mercaptoethanol). Cells were suspended in 0.7 packed-cell volumes of hypotonic buffer containing Complete protease inhibitors (Roche Molecular Biochemicals) and 0.5 units/ml of RNasin (Promega). Cells were disrupted in a Dounce homogenizer with a type B pestle, and lysates were centrifuged at 30,000×g for 20 min. Supernatants were used in an in vitro translation assay containing capped m7G(5')pppG firefly and Renilla luciferase mRNA or in in vitro Dicer assays containing ^{32}P-labeled dsRNA. For in vitro translation assays, 5 μl of extract were mixed with 100 ng of firefly and Renilla mRNA along with 1 μg of dsRNA (or buffer)/10 mM DTT/0.5 mM spermidine/200 mM Hepes, 3.3 mM MgOAc/800 mM KOAc/1 mM ATP/1 mM GTP/4 units of Rnasin/215 μg of creatine phosphate/1 μg of creatine phosphate kinase/1 mM amino acids (Promega). Reactions were carried out for 1 h at 30° C. and quenched by adding 1× passive lysis buffer (Promega). Extracts were then assayed for luciferase activity. In vitro assays for Dicer activity were performed as described (Bernstein et al. 2001, Nature 409: 363-366).

[0274] Construction of Stable Silencing Lines. Ten-centimeter plates of P19 cells were transfected with 5 μg of GFP hairpin expression plasmid and selected for stable integrants by using G-418 (300 ng/ml) for 14 days. Clones were selected and screened for silencing of GFP.

EXAMPLE 5

Compositions and Methods for Synthesizing siRNAs

[0275] Previous results have indicated that short synthetic RNAs (siRNAs) can efficiently induce RNA suppression. Since short RNAs do not activate the non-specific PKR response, they offer a means for efficiently silencing gene expression in a range of cell types. However, the current state of the art with respect to siRNAs has several limitations. Firstly, siRNAs are currently chemically synthesized at great cost (approx. $400/siRNA). Such high costs make siRNAs impractical for either small laboratories or for use in large scale screening efforts. Accordingly, there is a need in the art for methods for generating siRNAs at reduced cost.

[0276] We provide compositions and methods for synthesizing siRNAs by T7 polymerase. This approach allows for the efficient sythesis of siRNAs at a cost consistent with standard RNA transcription reactions (approx. $16/siRNA). This greatly reduced cost makes the use of siRNA a reasonable approach for small laboratories, and also will facilitate their use in large-scale screening projects.

[0277] **FIG. 38** shows the method for producing siRNAs using T7 polymerase. Briefly, T7 polymerase is used to transcribe both a sense and antisense transcript. The transcripts are then annealed to provide an siRNA. One of skill in the art will recognize that any one of the available RNA polymerases can be readily substituted for T7 to practice the invention (i.e., T3, Sp6, etc.).

[0278] This approach is amenable to the generation of a single siRNA species, as well as to the generation of a library of siRNAs. Such a library of siRNAs can be used in any number of high-throughput screens including cell based phenotypic screens and gene array based screens.

EXAMPLE 6

Generation of Short Hairpin dsRNA and Suppression of Gene Expression Using Such Short Hairpins

[0279] Since the realization that small, endogenously encoded hairpin RNAs could regulate gene expression via elements of the RNAi machinery, we have sought to exploit this biological mechanism for the regulation of desired target genes. Here we show that short hairpin RNAs (shRNAs) can induce sequence-specific gene silencing in mammalian cells. As is normally done with siRNAs, silencing can be provoked by transfecting exogenously synthesized hairpins into cells. However, silencing can also be triggered by endogenous expression of shRNAs. This observation opens the door to the production of continuous cells lines in which RNAi is used to stably suppress gene expression in mammalian cells. Furthermore, similar approaches should prove efficacious in the creation of transgenic animals and potentially in therapeutic strategies in which long-term suppression of gene function is essential to produce a desired effect.

[0280] Several groups (Grishok et al. 2001, Cell 106: 23-34; Ketting et al. 2001, Genes & Dev. 15: 2654-2659; Knight et al. 2001, Science 293: 2269-2271; Hutvagner et al. 2001, Science 293: 834-838) have shown that endogenous

triggers of gene silencing, specifically small temporal RNAs (stRNAs) let-7 and lin-4, function at least in part through RNAi pathways. Specifically, these small RNAs are encoded by hairpin precursors that are processed by Dicer into mature, ~21-nt forms. Moreover, genetic studies in *C. elegans* have shown a requirement for Argonaute-family proteins in stRNA function. Specifically, alg-1 and alg-2, members of the EIF2c subfamily, are implicated both in stRNA processing and in their downstream effector functions (Grishok et al., 2001, supra). We have recently shown that a component of RISC, the effector nuclease of RNAi, is a member of the Argonaute family, prompting a model in which stRNAs may function through RISC-like complexes, which regulate mRNA translation rather than mRNA stability (Hammond et al. 2001, Science 293: 1146-1150).

[0281] A. Short Hairpin RNAs Triggeedr Gene Silencing in Drosophila Cells.

[0282] We wished to test the possibility that we might retarget these small, endogenously encoded hairpin RNAs to regulate genes of choice with the ultimate goal of subverting this regulatory system for manipulating gene expression stably in mammalian cell lines and in transgenic animals. Whether triggered by long dsRNAs or by siRNAs, RNAi is generally more potent in the suppression of gene expression in Drosophila S2 cells than in mammalian cells. We therefore chose this model system in which to test the efficacy of short hairpin RNAs (shRNAs) as inducers of gene silencing.

[0283] Neither stRNAs nor the broader group of miRNAs that has recently been discovered form perfect hairpin structures. Indeed, each of these RNAs is predicted to contain several bulged nucleotides within their rather short (~30-nt) stem structures. Because the position and character of these bulged nucleotides have been conserved throughout evolution and among at least a subset of miRNAs, we sought to design retargeted miRNA mimics to conserve these predicted structural features. Only the let-7 and lin-4 miRNAs have known miRNA targets (Wightman et al. 1993, Cell 75: 855-862; Slack et al. 2000, Mol. Cell 5: 659-669). In both cases, pairing to binding sites within the regulated transcripts is imperfect, and in the case of lin-4, the presence of a bulged nucleotide is critical to suppression (Ha et al. 1996, Genes & Dev. 10: 3041-3050). We therefore also designed shRNAs that paired imperfectly with their target substrates. A subset of these shRNAs is depicted in **FIG. 39A**.

[0284] To permit rapid testing of large numbers of shRNA variants and quantitative comparison of the efficacy of suppression, we chose to use a dual-luciferase reporter system, as previously described for assays of RNAi in both Drosophila extracts (Tuschl et al. 1999, Genes & Dev. 13: 3191-3197) and mammalian cells (Caplen et al. 2001, Proc. Natl. Acad. Sci. 98: 9742-9747; Elbashir et al., 2001, Nature 411: 494-498). Cotransfection of firefly and Renilla luciferase reporter plasmids with either long dsRNAs or with siRNAs homologous to the firefly luciferase gene yielded an ~95% suppression of firefly luciferase without effect on Renilla luciferase (**FIG. 39B**; data not shown). Firefly luciferase could also be specifically silenced by cotransfection with homologous shRNAs. The most potent inhibitors were those composed of simple hairpin structures with complete homology to the substrate. Introduction of G-U basepairs either within the stem or within the substrate recognition sequence had little or no effect (**FIGS. 39A and 39B**; data not shown).

[0285] These results show that short hairpin RNAs can induce gene silencing in Drosophila S2 cells with potency similar to that of siRNAs (**FIG. 39B**). However, in our initial observation of RNA interference in Drosophila S2 cells, we noted a profound dependence of the efficiency of silencing on the length of the dsRNA trigger (Hammond et al. 2000, Nature 404: 293-296). Indeed, dsRNAs of fewer than ~200 nt triggered silencing very inefficiently. Silencing is initiated by an RNase III family nuclease, Dicer, that processes long dsRNAs into ~22-nt siRNAs. In accord with their varying potency as initiators of silencing, long dsRNAs are processed much more readily than short RNAs by the Dicer enzyme (Bernstein et al. 2001, Nature 409: 363-366). We therefore tested whether shRNAs were substrates for the Dicer enzyme.

[0286] We had noted previously that let-7 (Ketting et al. 2001, Genes & Dev. 15: 2654-2659) and other miRNAs (E. Bernstein, unpublished data) are processed by Dicer with an unexpectedly high efficiency as compared with short, non-hairpin dsRNAs. Similarly, Dicer efficiently processed shRNAs that targeted firefly luciferase, irrespective of whether they were designed to mimic a natural Dicer substrate (let-7) or whether they were simple hairpin structures (**FIG. 39C**). These data suggest that recombinant shRNAs can be processed by Dicer into siRNAs and are consistent with the idea that these short hairpins trigger gene silencing via an RNAi pathway.

[0287] B. Short Hairpin RNAs Activated Gene Silencing in Mammalian Cells.

[0288] Mammalian cells contain several endogenous systems that were predicted to hamper the application of RNAi. Chief among these is a dsRNA-activated protein kinase, PKR, which effects a general suppression of translation via phosphorylation of EIF-2α (Williams 1997, Biochem. Soc. Trans. 25: 509-513; Gil et al. 2000, Apoptosis 5: 107-114). Activation of these, and other dsRNA-responsive pathways, generally requires duplexes exceeding 30 bp in length, possibly to permit dimerization of the enzyme on its allosteric activator (e.g., Clarke et al. 1995, RNA 1: 7-20). Small RNAs that mimic Dicer products, siRNAs, presumably escape this limit and trigger specific silencing, in part because of their size. However, short duplex RNAs that lack signature features of siRNAs can efficiently induce silencing in Drosophila S2 cells but not in mammalian cells (A. A. Caudy, unpublished data). Endogenously encoded miRNAs may also escape PKR surveillance because of their size but perhaps also because of the discontinuity of their duplex structure. Given that shRNAs of <30 bp were effective inducers of RNAi in Drosophila S2 cells, we tested whether these RNAs could also induce sequence-specific silencing in mammalian cells.

[0289] Human embryonic kidney (HEK293T) cells were cotransfected with chemically synthesized shRNAs and with a mixture of firefly and Renilla luciferase reporter plasmids. As had been observed in S2 cells, shRNAs were effective inducers of gene silencing. Once again, hairpins designed to mimic let-7 were consistently less effective than were simple hairpin RNAs, and the introduction of mismatches between the antisense strand of the shRNA and the mRNA target abolished silencing (**FIG. 40A**; data not shown). Overall, shRNAs were somewhat less potent silencing triggers than were siRNAs. Whereas siRNAs homologous to firefly

luciferase routinely yielded ~90%-95% suppression of gene expression, suppression levels achieved with shRNAs ranged from 80%-90% on average. As we also observe with siRNAs, the most important determinant of the potency of the silencing trigger is its sequence. We find that roughly 50% of both siRNAs and shRNAs are competent for suppressing gene expression. However, neither analysis of the predicted structures of the target mRNA nor analysis of alternative structures in siRNA duplexes or shRNA hairpins has proved of predictive value for choosing effective inhibitors of gene expression.

[0290] We have adopted as a standard, shRNA duplexes containing 29 bp. However, the size of the helix can be reduced to ~25 nt without significant loss of potency. Duplexes as short as 22 bp can still provoke detectable silencing, but do so less efficiently than do longer duplexes. In no case did we observe a reduction in the internal control reporter (Renilla luciferase) that would be consistent with an induction of nonspecific dsRNA responses.

[0291] The ability of shRNAs to induce gene silencing was not confined to 293T cells. Similar results were also obtained in a variety of other mammalian cell lines, including human cancer cells (HeLa), transformed monkey epithelial cells (COS-1), murine fibroblasts (NIH 3T3), and diploid human fibroblasts (IMR90; **FIG. 40**; data not shown).

[0292] C. Synthesis of Effective Inhibitors of Gene Expression Using T7 RNA Polymerse.

[0293] The use of siRNAs to provoke gene silencing is developing into a standard methodology for investigating gene function in mammalian cells. To date, siRNAs have been produced exclusively by chemical synthesis (e.g., Caplen et al. 2001, Proc. Natl. Acad. Sci. 98: 9742-9747; Elbashir et al. 2001, Nature 411: 494-498). However, the costs associated with this approach are significant, limiting its potential utility as a tool for investigating in parallel the functions of large numbers of genes. Short hairpin RNAs are presumably processed into active siRNAs in vivo by Dicer. Thus, these may be more tolerant of terminal structures, both with respect to nucleotide overhangs and with respect to phosphate termini. We therefore tested whether shRNAs could be prepared by in vitro transcription with T7 RNA polymerase.

[0294] Transcription templates that were predicted to generate siRNAs and shRNAs similar to those prepared by chemical RNA synthesis were prepared by DNA synthesis (**FIG. 41A,C**). These were tested for efficacy both in S2 cells (data not shown) and in human 293 cells (**FIG. 41B,D**). Overall, the performance of the T7-synthesized hairpin or siRNAs closely matched the performance of either produced by chemical synthesis, both with respect to the magnitude of inhibition and with respect to the relative efficiency of differing sequences. Because T7 polymerase prefers to initiate at twin guanosine residues, however, it was critical to consider initiation context when designing in vitro transcribed siRNAs (**FIG. 41B**). In contrast, shRNAs, which are processed by Dicer (see **FIG. 39C**), tolerate the addition of these bases at the 5' end of the transcript.

[0295] Studies in Drosophila embryo extracts indicate that siRNAs possess 5' phosphorylated termini, consistent with their production by an RNase III family nuclease. In vitro, this terminus is critical to the induction of RNAi by synthetic RNA oligonucleotides (Elbashir et al. 2001, EMBO J. 20: 6877-6888; Nykanen et al. 2001, Cell 107: 309-321). Chemically synthesized siRNAs are nonphosphorylated, and enzymatic addition of a 5' phosphate group in vitro prior to transfection does not increase the potency of the silencing effect (A. A. Caudy, unpublished data). This suggests either that the requirement for phosphorylated termini is less stringent in mammalian cells or that a kinase efficiently phosphorylates siRNAs in vivo. RNAs synthesized with T7 RNA polymerase, however, possess 5' triphosphate termini. We therefore explored the possibility of synthesizing siRNAs with T7 polymerase followed by treatment in vitro with pyrophosphatase to modify the termini to resemble those of siRNAs. Surprisingly, monophosphorylated siRNAs (data not shown) were as potent in inducing gene silencing as transcription products bearing triphosphate termini (**FIG. 41B**). This may suggest either that the requirement for monophosphorylated termini is less stringent in mammalian cells or that siRNAs are modified in vivo to achieve an appropriate terminal structure.

[0296] Considered together, our data suggest that both shRNAs and siRNA duplexes can be prepared by synthesis with T7 RNA polymerase in vitro. This significantly reduces the cost of RNAi in mammalian cells and paves the way for application of RNAi on a whole-genome scale.

[0297] D. Transcription of Small Hairpin RNAs In Vivo by RNA Polymerase III.

[0298] Although siRNAs are an undeniably effective tool for probing gene function in mammalian cells, their suppressive effects are by definition of limited duration. Delivery of siRNAs can be accomplished by any of a number of transient transfection methodologies, and both the timing of peak suppression and the recovery of protein levels as silencing decays can vary with both the cell-type and the target gene. Therefore, one limitation on siRNAs is the development of continuous cell lines in which the expression of a desired target is stably silenced.

[0299] Hairpin RNAs, consisting of long duplex structures, have been proved as effective triggers of stable gene silencing in plants, in *C. elegans*, and in Drosophila (Kennerdell et al. 2000, Nat. Biotechnol. 18: 896-898; Smith et al. 2000, Nature 407: 319-320; Tavernarakis et al. 2000, Nat. Genet. 24: 180-183). We have recently shown stable suppression of gene expression in cultured mammalian cells by continuous expression of a long hairpin RNA (Paddison et al. 2002, Proc. Natl. Acad. Sci. 99: 1443-1448). However, the scope of this approach was limited by the necessity of expressing such hairpins only in cells that lack a detectable PKR response. In principle, shRNAs could bypass such limitations and provide a tool for evoking stable suppression by RNA in mammalian somatic cells.

[0300] To test this possibility, we initially cloned sequences encoding a firefly luciferase shRNA into a CMV-based expression plasmid. This was predicted to generate a capped, polyadenylated RNA polymerase II transcript in which the hairpin was extended on both the 5' and 3' ends by vector sequences and poly(A). This construct was completely inert in silencing assays in 293T cells.

[0301] During our studies on chemically and T7-synthesized shRNAs, we noted that the presence of significant

single-stranded extensions (either 5' or 3' of the duplex) reduced the efficacy of shRNAs. We therefore explored the use of alternative promoter strategies in an effort to produce more defined hairpin RNAs. In particular, RNA polymerase III promoters have well-defined initiation and termination sites and naturally produce a variety of small, stable RNA species. Although many Pol III promoters contain essential elements within the transcribed region, limiting their utility for our purposes; class III promoters use exclusively non-transcribed promoter sequences. Of these, the U6 snRNA promoter and the H1 RNA promoter have been well studied (Lobo et al. 1990, Nucleic Acids Res. 18: 2891-2899; Hannon et al. 1991, J. Biol. Chem. 266: 22796-22799; Chong et al. 2001, J. Biol. Chem. 276: 20727-20734).

[0302] By placing a convenient cloning site immediately behind the U6 snRNA promoter, we have constructed pShh-1, an expression vector in which short hairpins are harnessed for gene silencing. Into this vector either of two shRNA sequences derived from firefly luciferase were cloned from synthetic oligonucleotides. These were cotransfected with firefly and Renilla luciferase expression plasmids into 293T cells. One of the two encoded shRNAs provoked effective silencing of firefly luciferase without altering the expression of the internal control (**FIG. 42C**). The second encoded shRNA also produced detectable, albeit weak, repression. In both cases, silencing was dependent on insertion of the shRNA in the correct orientation with respect to the promoter (**FIG. 42C**; data not shown). Although the shRNA itself is bilaterally symmetric, insertion in the incorrect orientation would affect Pol III termination and is predicted to produce a hairpin with both 5' and 3' single-stranded extensions. Similar results were also obtained in a number of other mammalian cell lines including HeLa, COS-1, NIH 3T3, and IMR90 (**FIG. 42**; data not shown). pShh1-Ff1 was, however, incapable of effecting suppression of the luciferase reporter in Drosophila cells, in which the human U6 promoter is inactive.

[0303] E. Dicer is Required for shRNA-Mediated Gene Silencing.

[0304] As a definitive test of whether the plasmid-encoded shRNAs brought about gene silencing via the mammalian RNAi pathway, we assessed the dependence of suppression on an essential component of the RNAi pathway. We transfected pShh1-Ff1 along with an siRNA homologous to human Dicer. **FIG. 43** shows that treatment of cells with Dicer siRNAs is able to completely depress the silencing induced by pShh1-Ff1. Addition of an unrelated siRNA had no effect on the magnitude of suppression by pShh1-Ff1. Importantly, Dicer siRNAs had no effect on siRNA-induced silencing of firefly luciferase. These results are consistent with shRNAs operating via an RNAi pathway similar to those provoked by stRNAs and long dsRNAs. Furthermore, it suggests that siRNA-mediated silencing is less sensitive to depletion of the Dicer enzyme.

[0305] F. Stable shRNA-Mediated Gene Silencing of an Endogenous Gene.

[0306] The ultimate utility of encoded short hairpins will be in the creation of stable mutants that permit the study of the resulting phenotypes. We therefore tested whether we could create a cellular phenotype through stable suppression. Expression of activated alleles of the ras oncogene in primary mouse embryo fibroblasts (MEFs) induces a stable growth arrest that resembles, as a terminal phenotype, replicative senescence (Serrano et al. 1997, Cell 88: 593-602). Cells cease dividing and assume a typical large, flattened morphology. Senescence can be countered by mutations that inactivate the p53 tumor suppressor pathway (Serrano et al. 1997, supra). As a test of the ability of vector-encoded shRNAs to stably suppress an endogenous cellular gene, we generated a hairpin that was targeted to the mouse p53 gene. As shown in **FIG. 44**, MEFs transfected with pBabe-RasV12 fail to proliferate and show a senescent morphology when cotransfected with an empty control vector. As noted previously by Serrano et al., the terminally arrested state is achieved in 100% of drug-selected cells in culture by 8 d posttransfection. However, upon cotransfection of an activated ras expression construct with the pShh-p53, cells emerged from drug selection that not only fail to adopt a senescent morphology but also maintain the ability to proliferate for a minimum of several weeks in culture (**FIG. 44**). These data strongly suggest that shRNA expression constructs can be used for the creation of continuous mammalian cell lines in which selected target genes are stably suppressed.

[0307] G. Simultaneous Introduction of Multiple Hairpin RNAs Does Not Produce Synergy.

[0308] In an attempt to further understand the mechanisms by which short hairpins suppress gene expression, we examined the effects of transfecting cells with a mixture of two different short hairpins corresponding to firefly luciferase. **FIG. 45** summarizes the results of experiments which suggest that there is no synergistic affects on suppression of firefly luciferase gene expression obtained when cells are exposed to a mixture of such short hairpins.

[0309] Methods:

[0310] Cell culture. HEK 293T, HeLa, COS-1, MEF, and IMR90 cells were cultured in DMEM (GIBCO BRL) supplemented with 10% heat-inactivated fetal bovine serum (FBS) and 1% antibiotic/antimycotic solution (GIBCO BRL). NIH 3T3 cells were cultured in DMEM supplemented with 10% heat-inactivated calf serum and 1% antibiotic/antimycotic solution.

[0311] RNA preparation. Both shRNAs and siRNAs were produced in vitro using chemically synthesized DNA oligonucleotide templates (Sigma) and the T7 Megashortscript kit (Ambion). Transcription templates were designed such that they contained T7 promoter sequences at the 5' end. shRNA transcripts subjected to in vitro Dicer processing were synthesized using a Riboprobe kit (Promega). Chemically synthesized RNAs were obtained from Dharmacon, Inc.

[0312] Transfection and gene silencing assays. Cells were transfected with indicated amounts of siRNA, shRNA, and plasmid DNA using standard calcium phosphate procedures at 50%-70% confluence in 6-well plates. Dual luciferase assays (Promega) were carried out by cotransfecting cells with plasmids containing firefly luciferase under the control of the SV40 promoter (pGL3-Control, Promega) and Renilla luciferase under the control of the SV40 early enhancer/promoter region (pSV40, Promega). Plasmids were cotransfected using a 1:1 ratio of pGL3-Control (250 ng/well) to pRL-SV40. RNAi in S2 cells was performed as previously described (Hammond et al., 2000, Nature 404: 293-296). For stable silencing, primary MEFs (a gift from S. Lowe, Cold

Spring Harbor Laboratory, NY) were cotransfected using Fugene 6 with pBabe-Ha-rasV12 and pShh-p53 (no resistance marker), according to the manufacturer's recommendations. Selection was for the presence of the activated Ha-rasV12 plasmid, which carries a puromycin-resistance marker. The pShh-p53 plasmid was present in excess, as is standard in a cotransfection experiment. We have now generated a version of the U6 promoter vector (pSHAG-1) that is compatible with the GATEWAY system (Invitrogen), and this can be used to transport the shRNA expression cassette into a variety of recipient vectors that carry cis-linked selectable markers. Furthermore, we have validated delivery of shRNAs using retroviral vectors. Updated plasmid information can be obtained at http://www.cshl.org/public/science/hannon.html.

[0313] Plasmids expressing hairpin RNAs. The U6 promoter region from −265 to +1 was amplified by PCR, adding 5' KpnI and 3' EcoRV sites for cloning into pBSSK+. A linker/terminator oligonucleotide set bearing the U6 terminator sequence and linker ends of 5' EcoRV and 3' NotI was cloned into the promoter construct, resulting in a U6 cassette with an EcoRV site for insertion of new sequences. This vector has been named pShh1. Blunt-ended, double-stranded DNA oligonucleotides encoding shRNAs with between 19 and 29 bases of homology to the targeted gene were ligated into the EcoRV site to produce expression constructs. The oligonucleotide sequence used to construct Ff1 was: TCCAATTCAGCGGGAGCCACCTGATGAAGCTTGATCGGGTGGCTCTCGCTGAG TTGGAATCCATTTTTTTT. This sequence is preceded by the sequence GGAT, which is supplied by the vector, and contains a tract of more than five Ts as a Pol III terminator.

[0314] In vitro Dicer assays. In vitro assays for Dicer activity were performed as described (Bernstein et al., 2001, Nature 409: 363-366).

EXAMPLE 7

Encoded Short Hairpins Function in Vivo

[0315] An object of the present invention is to improve methods for generating siRNAs and short hairpins for use in specifically suppressing gene expression. Example 6 demonstrates that siRNAs and short hairpins are highly effective in specifically suppressing gene expression. Accordingly, it would be advantageous to combine the efficient suppression of gene expression attainable using short hairpins and siRNAs with a method to encode such RNA on a plasmid and express it either transiently or stably.

[0316] **FIG. 46** demonstrates that short hairpins encoded on a plasmid are effective in suppressing gene expression. DNA oligonucleotides encoding 29 nucleotide hairpins corresponding to firefly luciferase were inserted into a vector containing the U6 promoter. Three independent constructs were examined for their ability to specifically suppress firefly luciferase gene expression in 293T cells. siOligo1-2, siOligo1-6, and siOligo1-19 (construct in the correct orientation) each suppressed gene expression as effectively as siRNA. In contrast, siOligo1-10 (construct in the incorrect orientation) did not suppress gene expression. Additionally, an independent construct targeted to a different portion of the firefly luciferase gene did not effectively suppress gene expression in either orientation (siOligo2-23, siOligo2-36).

[0317] The results summarized in **FIG. 46** demonstrate that transient expression of siRNAs and short hairpins encoded on a plasmid can efficiently suppress gene expression. One of skill can choose from amongst a range of vectors to either transiently or stably express an siRNA or short hairpin. Non-limiting examples of vectors and strategies to stably express short dsRNAs are presented in FIGS. **47-49**.

EXAMPLE 8

dsRNA Suppression in the Absence of a PKR Response

[0318] One potential impediment to the use of RNAi to suppress gene expression in some cell types, is the non-specific PKR response that can be triggered by long dsRNAs. Numerous mammalian viruses have evolvd the ability to block PKR inorder to aid in the infection of potential host cells. For example, adenoviruses express RNAs which mimic dsRNA but do not activate the PKR response. Vaccinia virus uses two strategies to evade PKR: the expression of E3L which binds and masks dsRNA; the expression of K3L to mimic the natural PKR substrate eIF2α.

[0319] Our understanding of the mechanisms by which viruses avoid the PKR response allows us to design approaches to circumvent the PKR response in cell types in which in might be advantageous to suppression gene expression with long dsRNAs. Possible approaches include treating cells with an agent that inhibits protein kinase RNA-activated (PKR) apoptosis, such as by treatment with agents which inhibit expression of PKR, cause its destruction, and/or inhibit the kinase activity of PKR. Accordingly, RNAi suppression of gene expression in such cell types could involve first inhibiting the PKR response, and then delivering a dsRNA identical or similar to a target gene.

[0320] A. In a murine myoblast cell line, C2C12, we noted that the cells responded to long dsRNAs with a mixture of specific and non-specific (presumably PKR) responses. In order to attenuate the non-specific PKR response while maintaining the robust and specific suppression due to the long dsRNA, C2C12 cells were transfected with a vector that directs K3L expression. This additional step successfully attenuated the PKR response, however expression of K3L protein had no effect on the magnitude of specific inhibition.

[0321] B. However, since the efficacy of such a two step approach had not been previously demonstrated, it was formerly possible that dsRNA suppression would not be possible in cells with a PKR response. **FIG. 50** summarizes results which demonstrate that such a two step approach is possible, and that robust and specific dsRNA mediated suppression is possible in cells which had formerly possessed a robust PKR response.

[0322] Briefly, dual luciferase assay were carried out as described in detail above. The experiments were carried out using PKR−/− MEFs harvested from E13.5 PKR−/− mosue embryos. MEFs typically have a robust PKR response, and thus treatment with long dsRNAs typically results in non-specific suppression of gene expression and apoptosis. However, in PKR−/− cells examined 12, 42, and 82 hours after transfection, expression of dsRenilla luciferase RNA spe-

cifically suppresses expression *Renilla reniformis* (sea pansy) luciferase. This suppression is stable over time.

[0323] These results demonstrate that the non-specific PKR response can be blocked without affecting specific suppression of gene expression mediated by dsRNA. This allows the use of long dsRNAs to suppress gene expression in a diverse range of cell types, including those that would be previously intractable due to the confounding influences of the non-specific PKR response to long dsRNA.

EXAMPLE 9

Suppression of Gene Expression Using dsRNA which Corresponds to Non-Coding Sequence

[0324] Current models for the mechanisms which drive RNAi have suggested that the dsRNA construct must contain coding sequence corresponding to the gene of interest. Although evidence has demonstrated that such coding sequence need not be a perfect match to the endogenous coding sequence (i.e., it may be similar), it has been widely held that the dsRNA construct must correspond to coding sequence. We present evidence that contradicts the teachings of the prior art, and demonstrate that dsRNA corresponding to non-coding regions of a gene can suppress gene function in vivo. These results are significant not only because they demonstrate that dsRNA identical or similar to non-coding sequences (i.e., promoter sequences, enhancer sequences, or intronic sequences) can mediate suppression, but also because we demonstrate the in vivo suppression of gene expression using dsRNA technology in a mouse model.

[0325] We generated doubled stranded RNA corresponding to four segments of the mouse tyrosinase gene promoter. Three of these segments correspond to the proximal promoter and one corresponds to an enhancer (**FIG. 51**). The tyrosinase gene encodes the rate limiting enzyme involved in the melanin biosynthetic pathway (Bilodeau et al. (2001) *Pigment Cell Research* 14: 328-336). Accordingly, suppression of the tyrosinase gene is expected to inhibit pigmentation.

[0326] Double stranded RNA corresponding to each of the above promoter segments was injected into the pronuclei of fertilized eggs. Pups were born after 19 days. In total 42/136 (31%) of the embryos were carried to term. This number is within the expected range for transgenesis (30-40%). Two pups out of 42 (5%) appear totally unpigmented at birth, consistent with suppression of tyrosinase function.

[0327] Methods:

[0328] dsRNA from non-coding promoter region of tyrosinase gene. Four segments of the mouse tyrosinase gene promoter were amplified by PCR using primers which incorporated T7 RNA polymerase promoters into the PCR products (shown in bold—**FIG. 51**). Sequences of the mouse tyrosinase gene 5' flanking regions were obtained from GenBank (accession number D00439 and X51743). The sequence of the tyrosinase enhancer, located approximately 12 kb upstream of the transcriptional start site, was also obtained from GenBank (accession number X76647).

[0329] The sequences of the primers used were as follows: note the sequence of the T7 RNA polymerase promoter is shown in bold

(a) Tyrosinase enhancer (~12 kb upstream):
5'**TAATACGACTCACTATAGGG**CAAGGTCATAGTTCCTGCCAGCTG 3'
5'**TAATACGACTCACTATAGGG**CAGATATTTTCTTACCACCCACCC 3'

(b) -1404 to -1007:
5'**TAATACGACTCACTATAGGG**TTAAGTTTAACAGGAGAAGCTGGA 3'
5'**TAATACGACTCACTATAGGG**AAATCATTGCTTTCCTGATAATGC 3'

(c) -1003 to -506:
5'**TAATACGACTCACTATAGGG**TAGATTTCCGCAGCCCCAGTGTTC 3'
5'**TAATACGACTCACTATAGGG**GTTGCCTCTCATTTTTCCTTGATT 3'

(d) -505 to -85:
5'**TAATACGACTCACTATAGGG**TATTTTAGACTGATTACTTTTATAA 3'
5'**TAATACGACTCACTATAGGG**TCACATGTTTTGGCTAAGACCTAT 3'

[0330] PCR products were gel purified from 1% TAE agarose gels using QiaExII Gel Extraction Kit (Qiagen). Double stranded RNA was produced from these templates using T7-Megashortscript Kit (Ambion). Enzymes and unincorporated nucleotides were removed using Qiaquick MinElute PCR Purification Kit. RNA was phenol/chloroform extracted twice, and ethanol precipitated. Pellets were resuspended in injection buffer ((10 mM Tris (pH 7.5), 0.15 nM EDTA (pH 8.0)) at a concentration of 20 ng/ul and run on a 1% TAE agarose gel to confirm integrity.

[0331] Generation of mice: An equal mixture of double stranded RNA from each of the above primer sets was injected into the pronuclei of fertilized eggs from C57BL6J mice. A total of 136 injections was performed, and 34 embryos were implanted into each of 4 pseudopregnant CD-1 females. Pups were born after 19 days. In total, 42/136 (31%) of the embryos were carried to term. 2/42 pups (5%) appear totally unpigmented at birth.

[0332] It is not clear whether the RNAi mediated by dsRNA identical or similar to non-coding sequence works via the same mechanism as PTGS observed in the presence of dsRNA identical or similar to coding sequence. However, whether these results ultimately reveal similar or differing mechanisms does not diminish the tremendous utility of the compositions and methods of the present invention to suppress expression of one or more genes in vitro or in vivo.

[0333] The present invention demonstrates that dsRNA ranging in length from 20-500 nt can readily suppress expression of target genes both in vitro and in vivo. Furthermore, the present invention demonstrates that the dsRNAs can be generated using a variety of methods including the formation of hairpins, and that these dsRNAs can be expressed either stably or transiently. Finally, the present invention demonstrates that dsRNA identical or similar to non-coding sequences can suppress target gene expression.

EXAMPLE 10

RNA Interference in Adult Mice

[0334] RNA interference is an evolutionarily conserved surveillance mechanism that responds to double-stranded RNA by sequence-specific silencing of homologous genes. Here we show that transgene expression can be suppressed in adult mice by synthetic small interfering RNAs and by small-hairpin RNAs transcribed in vivo from DNA templates. We also show the therapeutic potential of this technique by demonstrating effective targeting of a sequence from hepatitis C virus by RNA interference in vivo.

[0335] Small interfering RNAs (siRNAs) mimic intermediates in the RNA-interference (RNAi) pathway and can silence genes in somatic cells without activating non-specific suppression by double-stranded RNA-dependent protein kinase (Elbashir et al. 2001, Nature 411: 494-498). To investigate whether siRNAs also inhibit gene expression in vivo, we used a modification of hydrodynamic transfection methods (Zhang et al. 1999, Hum. Gene Therapy 10: 1735-1737; Liu et al. 1999, Gene Therapy 6: 1258-1266; Chang et al. 2001, J. Virol. 75: 3469-3473) to deliver naked siRNAs to the livers of adult mice. Either an siRNA derived from firefly luciferase or an unrelated siRNA was co-injected with a luciferase-expression plasmid (for construct description and sequences, see **FIG. 52**). We monitored luciferase expression in living animals using quantitative whole-body imaging (Contag, et al. 1997, Photochem. Photobiol. 66: 523-531) (see **FIGS. 53***a*, **54***a*), and found that it was dependent on reporter-plasmid dose.

[0336] In each experiment, serum measurements of a co-injected human χ-1 antitrypsin (hAAT) plasmid (Yant et al. 2000, Nature Genet. 25: 35-41) served to normalize transfection efficiency and to monitor non-specific translational inhibition. Average serum concentrations of HAAT after 74 h were similar in all groups.

[0337] Our results indicate that there was specific, siRNA-mediated inhibition of luciferase expression in adult mice ($P<0.0115$) and that unrelated siRNAs had no effect ($P<0.864$; **FIGS. 53***a*, **53***b*). In 11 independent experiments, luciferase siRNAs reduced luciferase expression (as judged by emitted light) by an average of 81% (±2.2%). These findings indicate that RNAi can downregulate gene expression in adult mice.

[0338] As RNAi degrades respiratory syncitial virus RNAs in culture (Bitko et al. 2001, BMC Microbiol. 1: 34), we investigated whether RNAi could be directed against a human pathogenic RNA expressed in a mouse, namely that of hepatitis C virus (HCV). Infection by HCV (an RNA virus that infects 1 in 40 people worldwide) is the most common reason for liver transplantation in the United States and Europe. We fused the NS5B region (non-structural protein 5B, viral-polymerase-encoding region) of this virus with luciferase RNA and monitored RNAi by co-transfection in vivo. An siRNA targeting the NS5B region reduced luciferase expression from the chimaeric HCV NS5B protein-luciferase fusion by 75% (±6.8%; 6 animals per group). This result suggests that it may be feasible to use RNAi as a therapy against other important human pathogens.

[0339] Although our results show that siRNAs are functional in mice, delivery remains a major obstacle. Unlike siRNAs, functional small-hairpin RNAs (shRNAs) can be expressed in vivo from DNA templates using RNA polymerase III promoters (Paddison et al. 2002, Genes Dev. 16: 948-958; Tuschl 2002, Nature Biotechnol. 20: 446-448); they are as effective as siRNAs in inducing gene suppression. Expression of a cognate shRNA (pShh1-Ff1) inhibited luciferase expression by up to 98% (±0.6%), with an average suppression of 92.8% (±3.39%) in three independent experiments (see **FIGS. 54***a*, **54***b*). An empty shRNA-expression vector had no effect; reversing the orientation of the shRNA (pShh1-Ff1rev) insert prevents gene silencing because it alters the termination by RNA polymerase III and generates an improperly structured shRNA. These findings indicate that plasmid-encoded shRNAs can induce a potent and specific RNAi response in adult mice.

[0340] RNAi may find application in functional genomics or in identifying targets for designer drugs. It is a more promising system than gene-knockout mice because groups of genes can be simultaneously rendered ineffective without the need for time-consuming crosses. Gene therapy currently depends on the ectopic expression of exogenous proteins; however, RNAi may eventually complement this gain-of-function approach by silencing disease-related genes with DNA constructs that direct the expression of shRNAs. Our method of RNAi delivery could also be tailored to take advantage of developing viral and non-viral gene-transfer vectors in a clinical context.

EXAMPLE 11

Germline Transmission of RNAi in Mice

[0341] MicroRNA molecules (miRNAs) are small, non-coding RNA molecules that have been found in a diverse array of eukaryotes, including mammals. miRNA precursors share a characteristic secondary structure, forming short 'hairpin' RNAs. Genetic and biochemical studies have indicated that miRNAs are processed to their mature forms by Dicer, an RNAse III family nuclease, and function through RNA-mediated interference (RNAi) and related pathways to regulate the expression of target genes (Hannon 2002, Nature 418: 244-251; Pasquinelli et al. 2002, Annu. Rev. Cell. Dev. Biol. 18: 495-513). Recently, we and others have remodeled miRNAs to permit experimental manipulation of gene expression in mammalian cells and have dubbed these synthetic silencing triggers 'short hairpin RNAs' (shRNAs) (Paddison et al. 2002, Cancer Cell 2: 17-23). Silencing by shRNAs requires the RNAi machinery and correlates with the production of small interfering RNAs (siRNAs), which are a signature of RNAi.

[0342] Expression of shRNAs can elicit either transient or stable silencing, depending upon whether the expression cassette is integrated into the genome of the recipient cultured cell (Paddison et al. 2002, Cancer Cell 2: 17-23). shRNA expression vectors also induce gene silencing in adult mice following transient delivery (Lewis et al. 2002, Nat. Genet. 32: 107-108; McCaffrey et al. 2002, Nature 418: 38-39). However, for shRNAs to be a viable genetic tool in mice, stable manipulation of gene expression is essential. Hemann and colleagues have demonstrated long-term suppression of gene expression in vivo following retroviral delivery of shRNA-expression cassettes to hematopoietic stem cells (Hemann et al. 2003, Nat. Genet. in the press). Here we sought to test whether shRNA-expression cassettes that were passed through the mouse germline could enforce heritable gene silencing.

[0343] We began by taking standard transgenesis approaches (Gordon et al. 1993, Methods Enzymol. 225: 747-771) using shRNAs directed against a variety of targets with expected phenotypes, including the genes encoding tyrosinase (albino), myosin VIIa (shaker), Bmp-5 (crinkled ears), Hox a-10 (limb defects), homogentisate 1,2,-dioxygenase (urine turns black upon exposure to air), Hairless (hair loss) and melanocortin 1 receptor (yellow). Three constructs per gene were linearized and injected into pronuclei to produce transgenic founder animals. Although we

noted the presence of the transgene in some animals, virtually none showed a distinct or reproducible phenotype that was expected for a hypomorphic allele of the targeted gene.

[0344] Therefore, we decided to take another approach: verifying the presence of the shRNA and its activity toward a target gene in cultured embryonic stem (ES) cells and then asking whether those cells retained suppression in a chimeric animal in vivo. We also planned to test whether such cells could pass a functional RNAi-inducing construct through the mouse germline. For these studies, we chose to examine a novel gene, Neil1, which is proposed to have a role in DNA repair. Oxidative damage accounts for 10,000 DNA lesions per cell per day in humans and is thought to contribute to carcinogenesis, aging and tissue damage following ischemia (Ames et al. 1993, Proc. Natl. Acad. Sci. USA 90: 7915-7922; Jackson et al. 2001, Mutat. Res. 477: 7-21). Oxidative DNA damage includes abasic sites, strand breaks and at least 20 oxidized bases, many of which are cytotoxic or pro-mutagenic (Dizdaroglu et al. 2002, Free Radic. Biol. Med. 32: 1102-1115). DNA N-glycosylases initiate the base excision repair pathway by recognizing specific bases in DNA and cleaving the sugar base bond to release the damaged base (David et al. 1998, Chem. Rev. 98: 1221-1262).

[0345] The Neil genes are a newly discovered family of mammalian DNA N-glycosylases related to the Fpg/Nei family of proteins from *Escherichia coli* (Hazra et al. 2002, Proc. Natl. Acad. Sci. USA 99: 3523-3528; Bandaru et al. 2002, DNA Repair 1: 517-529). Neil1 recognizes and removes a wide spectrum of oxidized pyrimidines and ring-opened purines from DNA, including thymine glycol (Tg), 2,6-diamino-4-hydroxy-5-formamidopyrimidine (FapyG) and 4,6-diamino-5-formidopyrimidine (FapyA). Tg, FapyG and FapyA are among the most prevalent oxidized bases produced by ionizing radiation (Dizdaroglu et al. 2002, Free Radic. Biol. Med. 32: 1102-1115) and can block replicative DNA polymerases, which can, in turn, cause cell death (Asagoshi et al. 2002, J. Biol. Chem. 277: 14589-14597; Clark et al. 1989, Biochemistry 28: 775-779).

[0346] The Nth1 and Ogg1 glycosylases each remove subsets of oxidized DNA bases that overlap with substrates of Neil1 (Nishimura 2002, Free Radic. Biol. Med. 32: 813-821; Asagoshi et al. 2000, Biochemistry 39: 11389-11398; Dizdaroglu et al. 1999, Biochemistry 38: 243-246). However, mice with null mutations in either Nth1 (Ocampo et al. 2002, Mol. Cell. Biol. 22: 6111-6121; Takao et al. 2002, EMBO J. 21: 3486-3493) or Ogg1 (Klungland et al. 1999, Proc. Natl. Acad. Sci. USA 96: 13300-13305; Minowa et al. 2000, Proc. Natl. Acad. Sci. USA 97: 4156-4161) are viable, raising the possibility that Neil1 activity tempers the loss of Nth1 or Ogg1. Recently, a residual Tg-DNA glycosylase activity in $Nth1^{-/-}$ mice has been identified as Neil1 (Takao et al. 2002, J. Biol. Chem. 277: 42205-42213).

[0347] We constructed a single shRNA expression vector targeting a sequence near the 5' end of the Neil1 coding region. This vector was introduced into mouse embryonic stem cells by electroporation, and individual stable integrants were tested for expression of the Neil1 protein (see the weblink: http://www.cshl.edu/public/SCIENCE/hannon.html for detailed procedures). The majority of cell lines showed an ~80% reduction in Neil1 protein, which correlated with a similar change in levels of Neil1 mRNA. These cells showed an approximately two-fold increase in their sensitivity to ionizing radiation, consistent with a role for Neil1 in DNA repair. Two independent ES cell lines were injected into BL/6 blastocysts, and several high-percentage chimeras were obtained. These chimeras were out-crossed, and germline transmission of the shRNA-expression construct was noted in numerous F_1 progeny (13/27 for one line and 12/26 for the other).

[0348] To determine whether the silencing of Neil1 that had been observed in ES cells was transmitted faithfully, we examined Neil1 mRNA and protein levels. Both were reduced by approximately the same extent that had been observed in the engineered ES cells (**FIGS. 55, 56**). Consistent with this having occurred through the RNAi pathway, we detected the presence of siRNAs corresponding to the shRNA sequence in F_1 animals that carry the shRNA expression vector but not in those that lack the vector (**FIG. 56*b***).

[0349] The aforementioned data demonstrate that shRNAs can be used to create germline transgenic mice in which RNAi has silenced a target gene. These observations open the door to using of RNAi as a complement to standard knock-out methodologies and provide a means to rapidly assess the consequences of suppressing a gene of interest in a living animal. Coupled with activator-dependent U6 promoters, the use of shRNAs will ultimately provide methods for tissue-specific, inducible and reversible suppression of gene expression in mice.

[0350] V. Equivalents

[0351] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

[0352] All of the above-cited references and publications are hereby incorporated by reference.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 45

<210> SEQ ID NO 1
<211> LENGTH: 5775
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(5775)

-continued

<400> SEQUENCE: 1

```
atg aaa agc cct gct ttg caa ccc ctc agc atg gca ggc ctg cag ctc       48
Met Lys Ser Pro Ala Leu Gln Pro Leu Ser Met Ala Gly Leu Gln Leu
1               5                  10                  15

atg acc cct gct tcc tca cca atg ggt cct ttc ttt gga ctg cca tgg       96
Met Thr Pro Ala Ser Ser Pro Met Gly Pro Phe Phe Gly Leu Pro Trp
            20                  25                  30

caa caa gaa gca att cat gat aac att tat acg cca aga aaa tat cag      144
Gln Gln Glu Ala Ile His Asp Asn Ile Tyr Thr Pro Arg Lys Tyr Gln
        35                  40                  45

gtt gaa ctg ctt gaa gca gct ctg gat cat aat acc atc gtc tgt tta      192
Val Glu Leu Leu Glu Ala Ala Leu Asp His Asn Thr Ile Val Cys Leu
    50                  55                  60

aac act ggc tca ggg aag aca ttt att gct agt act act cta cta aag      240
Asn Thr Gly Ser Gly Lys Thr Phe Ile Ala Ser Thr Thr Leu Leu Lys
65                  70                  75                  80

agc tgt ctc tat cta gat cta ggg gag act tca gct aga aat gga aaa      288
Ser Cys Leu Tyr Leu Asp Leu Gly Glu Thr Ser Ala Arg Asn Gly Lys
                85                  90                  95

agg acg gtg ttc ttg gtc aac tct gca aac cag gtt gct caa caa gtg      336
Arg Thr Val Phe Leu Val Asn Ser Ala Asn Gln Val Ala Gln Gln Val
            100                 105                 110

tca gct gtc aga act cat tca gat ctc aag gtt ggg gaa tac tca aac      384
Ser Ala Val Arg Thr His Ser Asp Leu Lys Val Gly Glu Tyr Ser Asn
        115                 120                 125

cta gaa gta aat gca tct tgg aca aaa gag aga tgg aac caa gag ttt      432
Leu Glu Val Asn Ala Ser Trp Thr Lys Glu Arg Trp Asn Gln Glu Phe
    130                 135                 140

act aag cac cag gtt ctc att atg act tgc tat gtc gcc ttg aat gtt      480
Thr Lys His Gln Val Leu Ile Met Thr Cys Tyr Val Ala Leu Asn Val
145                 150                 155                 160

ttg aaa aat ggt tac tta tca ctg tca gac att aac ctt ttg gtg ttt      528
Leu Lys Asn Gly Tyr Leu Ser Leu Ser Asp Ile Asn Leu Leu Val Phe
                165                 170                 175

gat gag tgt cat ctt gca atc cta gac cac ccc tat cga gaa ttt atg      576
Asp Glu Cys His Leu Ala Ile Leu Asp His Pro Tyr Arg Glu Phe Met
            180                 185                 190

aag ctc tgt gaa att tgt cca tca tgt cct cgc att ttg gga cta act      624
Lys Leu Cys Glu Ile Cys Pro Ser Cys Pro Arg Ile Leu Gly Leu Thr
        195                 200                 205

gct tcc att tta aat ggg aaa tgg gat cca gag gat ttg gaa gaa aag      672
Ala Ser Ile Leu Asn Gly Lys Trp Asp Pro Glu Asp Leu Glu Glu Lys
    210                 215                 220

ttt cag aaa cta gag aaa att ctt aag agt aat gct gaa act gca act      720
Phe Gln Lys Leu Glu Lys Ile Leu Lys Ser Asn Ala Glu Thr Ala Thr
225                 230                 235                 240

gac ctg gtg gtc tta gac agg tat act tct cag cca tgt gag att gtg      768
Asp Leu Val Val Leu Asp Arg Tyr Thr Ser Gln Pro Cys Glu Ile Val
                245                 250                 255

gtg gat tgt gga cca ttt act gac aga agt ggg ctt tat gaa aga ctg      816
Val Asp Cys Gly Pro Phe Thr Asp Arg Ser Gly Leu Tyr Glu Arg Leu
            260                 265                 270

ctg atg gaa tta gaa gaa gca ctt aat ttt atc aat gat tgt aat ata      864
Leu Met Glu Leu Glu Glu Ala Leu Asn Phe Ile Asn Asp Cys Asn Ile
        275                 280                 285

tct gta cat tca aaa gaa aga gat tct act tta att tcg aaa cag ata      912
```

-continued

```
Ser Val His Ser Lys Glu Arg Asp Ser Thr Leu Ile Ser Lys Gln Ile
    290                 295                 300

cta tca gac tgt cgt gcc gta ttg gta gtt ctg gga ccc tgg tgt gca      960
Leu Ser Asp Cys Arg Ala Val Leu Val Val Leu Gly Pro Trp Cys Ala
305                 310                 315                 320

gat aaa gta gct gga atg atg gta aga gaa cta cag aaa tac atc aaa     1008
Asp Lys Val Ala Gly Met Met Val Arg Glu Leu Gln Lys Tyr Ile Lys
                325                 330                 335

cat gag caa gag gag ctg cac agg aaa ttt tta ttg ttt aca gac act     1056
His Glu Gln Glu Glu Leu His Arg Lys Phe Leu Leu Phe Thr Asp Thr
            340                 345                 350

ttc cta agg aaa ata cat gca cta tgt gaa gag cac ttc tca cct gcc     1104
Phe Leu Arg Lys Ile His Ala Leu Cys Glu Glu His Phe Ser Pro Ala
        355                 360                 365

tca ctt gac ctg aaa ttt gta act cct aaa gta atc aaa ctg ctc gaa     1152
Ser Leu Asp Leu Lys Phe Val Thr Pro Lys Val Ile Lys Leu Leu Glu
    370                 375                 380

atc tta cgc aaa tat aaa cca tat gag cga cac agt ttt gaa agc gtt     1200
Ile Leu Arg Lys Tyr Lys Pro Tyr Glu Arg His Ser Phe Glu Ser Val
385                 390                 395                 400

gag tgg tat aat aat aga aat cag gat aat tat gtg tca tgg agt gat     1248
Glu Trp Tyr Asn Asn Arg Asn Gln Asp Asn Tyr Val Ser Trp Ser Asp
                405                 410                 415

tct gag gat gat gat gag gat gaa gaa att gaa gaa aaa gag aag cca     1296
Ser Glu Asp Asp Asp Glu Asp Glu Glu Ile Glu Glu Lys Glu Lys Pro
            420                 425                 430

gag aca aat ttt cct tct cct ttt acc aac att ttg tgc gga att att     1344
Glu Thr Asn Phe Pro Ser Pro Phe Thr Asn Ile Leu Cys Gly Ile Ile
        435                 440                 445

ttt gtg gaa aga aga tac aca gca gtt gtc tta aac aga ttg ata aag     1392
Phe Val Glu Arg Arg Tyr Thr Ala Val Val Leu Asn Arg Leu Ile Lys
    450                 455                 460

gaa gct ggc aaa caa gat cca gag ctg gct tat atc agt agc aat ttc     1440
Glu Ala Gly Lys Gln Asp Pro Glu Leu Ala Tyr Ile Ser Ser Asn Phe
465                 470                 475                 480

ata act gga cat ggc att ggg aag aat cag cct cgc aac aac acg atg     1488
Ile Thr Gly His Gly Ile Gly Lys Asn Gln Pro Arg Asn Asn Thr Met
                485                 490                 495

gaa gca gaa ttc aga aaa cag gaa gag gta ctt agg aaa ttt cga gca     1536
Glu Ala Glu Phe Arg Lys Gln Glu Glu Val Leu Arg Lys Phe Arg Ala
            500                 505                 510

cat gag acc aac ctg ctt att gca aca agt att gta gaa gag ggt gtt     1584
His Glu Thr Asn Leu Leu Ile Ala Thr Ser Ile Val Glu Glu Gly Val
        515                 520                 525

gat ata cca aaa tgc aac ttg gtg gtt cgt ttt gat ttg ccc aca gaa     1632
Asp Ile Pro Lys Cys Asn Leu Val Val Arg Phe Asp Leu Pro Thr Glu
    530                 535                 540

tat cga tcc tat gtt caa tct aaa gga aga gca agg gca ccc atc tct     1680
Tyr Arg Ser Tyr Val Gln Ser Lys Gly Arg Ala Arg Ala Pro Ile Ser
545                 550                 555                 560

aat tat ata atg tta gcg gat aca gac aaa ata aaa agt ttt gaa gaa     1728
Asn Tyr Ile Met Leu Ala Asp Thr Asp Lys Ile Lys Ser Phe Glu Glu
                565                 570                 575

gac ctt aaa acc tac aaa gct att gaa aag atc ttg aga aac aag tgt     1776
Asp Leu Lys Thr Tyr Lys Ala Ile Glu Lys Ile Leu Arg Asn Lys Cys
            580                 585                 590

tcc aag tcg gtt gat act ggt gag act gac att gat cct gtc atg gat     1824
```

-continued

```
Ser Lys Ser Val Asp Thr Gly Glu Thr Asp Ile Asp Pro Val Met Asp
        595                 600                 605

gat gat cac gtt ttc cca cca tat gtg ttg agg cct gac gat ggt ggt     1872
Asp Asp His Val Phe Pro Pro Tyr Val Leu Arg Pro Asp Asp Gly Gly
    610                 615                 620

cca cga gtc aca atc aac acg gcc att gga cac atc aat aga tac tgt     1920
Pro Arg Val Thr Ile Asn Thr Ala Ile Gly His Ile Asn Arg Tyr Cys
625                 630                 635                 640

gct aga tta cca agt gat ccg ttt act cat cta gct cct aaa tgc aga     1968
Ala Arg Leu Pro Ser Asp Pro Phe Thr His Leu Ala Pro Lys Cys Arg
                645                 650                 655

acc cga gag ttg cct gat ggt aca ttt tat tca act ctt tat ctg cca     2016
Thr Arg Glu Leu Pro Asp Gly Thr Phe Tyr Ser Thr Leu Tyr Leu Pro
            660                 665                 670

att aac tca cct ctt cga gcc tcc att gtt ggt cca cca atg agc tgt     2064
Ile Asn Ser Pro Leu Arg Ala Ser Ile Val Gly Pro Pro Met Ser Cys
        675                 680                 685

gta cga ttg gct gaa aga gtt gtc gct ctc att tgc tgt gag aaa ctg     2112
Val Arg Leu Ala Glu Arg Val Val Ala Leu Ile Cys Cys Glu Lys Leu
    690                 695                 700

cac aaa att ggc gaa ctg gat gac cat ttg atg cca gtt ggg aaa gag     2160
His Lys Ile Gly Glu Leu Asp Asp His Leu Met Pro Val Gly Lys Glu
705                 710                 715                 720

act gtt aaa tat gaa gag gag ctt gat ttg cat gat gaa gaa gag acc     2208
Thr Val Lys Tyr Glu Glu Glu Leu Asp Leu His Asp Glu Glu Glu Thr
                725                 730                 735

agt gtt cca gga aga cca ggt tcc acg aaa cga agg cag tgc tac cca     2256
Ser Val Pro Gly Arg Pro Gly Ser Thr Lys Arg Arg Gln Cys Tyr Pro
            740                 745                 750

aaa gca att cca gag tgt ttg agg gat agt tat ccc aga cct gat cag     2304
Lys Ala Ile Pro Glu Cys Leu Arg Asp Ser Tyr Pro Arg Pro Asp Gln
        755                 760                 765

ccc tgt tac ctg tat gtg ata gga atg gtt tta act aca cct tta cct     2352
Pro Cys Tyr Leu Tyr Val Ile Gly Met Val Leu Thr Thr Pro Leu Pro
    770                 775                 780

gat gaa ctc aac ttt aga agg cgg aag ctc tat cct cct gaa gat acc     2400
Asp Glu Leu Asn Phe Arg Arg Arg Lys Leu Tyr Pro Pro Glu Asp Thr
785                 790                 795                 800

aca aga tgc ttt gga ata ctg acg gcc aaa ccc ata cct cag att cca     2448
Thr Arg Cys Phe Gly Ile Leu Thr Ala Lys Pro Ile Pro Gln Ile Pro
                805                 810                 815

cac ttt cct gtg tac aca cgc tct gga gag gtt acc ata tcc att gag     2496
His Phe Pro Val Tyr Thr Arg Ser Gly Glu Val Thr Ile Ser Ile Glu
            820                 825                 830

ttg aag aag tct ggt ttc atg ttg tct cta caa atg ctt gag ttg att     2544
Leu Lys Lys Ser Gly Phe Met Leu Ser Leu Gln Met Leu Glu Leu Ile
        835                 840                 845

aca aga ctt cac cag tat ata ttc tca cat att ctt cgg ctt gaa aaa     2592
Thr Arg Leu His Gln Tyr Ile Phe Ser His Ile Leu Arg Leu Glu Lys
    850                 855                 860

cct gca cta gaa ttt aaa cct aca gac gct gat tca gca tac tgt gtt     2640
Pro Ala Leu Glu Phe Lys Pro Thr Asp Ala Asp Ser Ala Tyr Cys Val
865                 870                 875                 880

cta cct ctt aat gtt gtt aat gac tcc agc act ttg gat att gac ttt     2688
Leu Pro Leu Asn Val Val Asn Asp Ser Ser Thr Leu Asp Ile Asp Phe
                885                 890                 895

aaa ttc atg gaa gat att gag aag tct gaa gct cgc ata ggc att ccc     2736
```

-continued

```
Lys Phe Met Glu Asp Ile Glu Lys Ser Glu Ala Arg Ile Gly Ile Pro
           900                 905                 910

agt aca aag tat aca aaa gaa aca ccc ttt gtt ttt aaa tta gaa gat     2784
Ser Thr Lys Tyr Thr Lys Glu Thr Pro Phe Val Phe Lys Leu Glu Asp
        915                 920                 925

tac caa gat gcc gtt atc att cca aga tat cgc aat ttt gat cag cct     2832
Tyr Gln Asp Ala Val Ile Ile Pro Arg Tyr Arg Asn Phe Asp Gln Pro
    930                 935                 940

cat cga ttt tat gta gct gat gtg tac act gat ctt acc cca ctc agt     2880
His Arg Phe Tyr Val Ala Asp Val Tyr Thr Asp Leu Thr Pro Leu Ser
945                 950                 955                 960

aaa ttt cct tcc cct gag tat gaa act ttt gca gaa tat tat aaa aca     2928
Lys Phe Pro Ser Pro Glu Tyr Glu Thr Phe Ala Glu Tyr Tyr Lys Thr
                965                 970                 975

aag tac aac ctt gac cta acc aat ctc aac cag cca ctg ctg gat gtg     2976
Lys Tyr Asn Leu Asp Leu Thr Asn Leu Asn Gln Pro Leu Leu Asp Val
            980                 985                 990

gac cac aca tct tca aga ctt aat  ctt ttg aca cct cga  cat ttg aat   3024
Asp His Thr Ser Ser Arg Leu Asn  Leu Leu Thr Pro Arg  His Leu Asn
        995                 1000                1005

cag aag  ggg aaa gcg ctt cct  tta agc agt gct gag  aag agg aaa      3069
Gln Lys  Gly Lys Ala Leu Pro  Leu Ser Ser Ala Glu  Lys Arg Lys
    1010                1015                1020

gcc aaa  tgg gaa agt ctg cag  aat aaa cag ata ctg  gtt cca gaa      3114
Ala Lys  Trp Glu Ser Leu Gln  Asn Lys Gln Ile Leu  Val Pro Glu
    1025                1030                1035

ctc tgt  gct ata cat cca att  cca gca tca ctg tgg  aga aaa gct      3159
Leu Cys  Ala Ile His Pro Ile  Pro Ala Ser Leu Trp  Arg Lys Ala
    1040                1045                1050

gtt tgt  ctc ccc agc ata ctt  tat cgc ctt cac tgc  ctt ttg act      3204
Val Cys  Leu Pro Ser Ile Leu  Tyr Arg Leu His Cys  Leu Leu Thr
    1055                1060                1065

gca gag  gag cta aga gcc cag  act gcc agc gat gct  ggc gtg gga      3249
Ala Glu  Glu Leu Arg Ala Gln  Thr Ala Ser Asp Ala  Gly Val Gly
    1070                1075                1080

gtc aga  tca ctt cct gcg gat  ttt aga tac cct aac  tta gac ttc      3294
Val Arg  Ser Leu Pro Ala Asp  Phe Arg Tyr Pro Asn  Leu Asp Phe
    1085                1090                1095

ggg tgg  aaa aaa tct att gac  agc aaa tct ttc atc  tca att tct      3339
Gly Trp  Lys Lys Ser Ile Asp  Ser Lys Ser Phe Ile  Ser Ile Ser
    1100                1105                1110

aac tcc  tct tca gct gaa aat  gat aat tac tgt aag  cac agc aca      3384
Asn Ser  Ser Ser Ala Glu Asn  Asp Asn Tyr Cys Lys  His Ser Thr
    1115                1120                1125

att gtc  cct gaa aat gct gca  cat caa ggt gct aat  aga acc tcc      3429
Ile Val  Pro Glu Asn Ala Ala  His Gln Gly Ala Asn  Arg Thr Ser
    1130                1135                1140

tct cta  gaa aat cat gac caa  atg tct gtg aac tgc  aga acg ttg      3474
Ser Leu  Glu Asn His Asp Gln  Met Ser Val Asn Cys  Arg Thr Leu
    1145                1150                1155

ctc agc  gag tcc cct ggt aag  ctc cac gtt gaa gtt  tca gca gat      3519
Leu Ser  Glu Ser Pro Gly Lys  Leu His Val Glu Val  Ser Ala Asp
    1160                1165                1170

ctt aca  gca att aat ggt ctt  tct tac aat caa aat  ctc gcc aat      3564
Leu Thr  Ala Ile Asn Gly Leu  Ser Tyr Asn Gln Asn  Leu Ala Asn
    1175                1180                1185

ggc agt  tat gat tta gct aac  aga gac ttt tgc caa  gga aat cag      3609
```

-continued

```
Gly Ser  Tyr Asp Leu Ala Asn  Arg Asp Phe Cys Gln  Gly Asn Gln
    1190                1195                1200

cta aat  tac tac aag cag gaa  ata ccc gtg caa cca  act acc tca      3654
Leu Asn  Tyr Tyr Lys Gln Glu  Ile Pro Val Gln Pro  Thr Thr Ser
    1205                1210                1215

tat tcc  att cag aat tta tac  agt tac gag aac cag  ccc cag ccc      3699
Tyr Ser  Ile Gln Asn Leu Tyr  Ser Tyr Glu Asn Gln  Pro Gln Pro
    1220                1225                1230

agc gat  gaa tgt act ctc ctg  agt aat aaa tac ctt  gat gga aat      3744
Ser Asp  Glu Cys Thr Leu Leu  Ser Asn Lys Tyr Leu  Asp Gly Asn
    1235                1240                1245

gct aac  aaa tct acc tca gat  gga agt cct gtg atg  gcc gta atg      3789
Ala Asn  Lys Ser Thr Ser Asp  Gly Ser Pro Val Met  Ala Val Met
    1250                1255                1260

cct ggt  acg aca gac act att  caa gtg ctc aag ggc  agg atg gat      3834
Pro Gly  Thr Thr Asp Thr Ile  Gln Val Leu Lys Gly  Arg Met Asp
    1265                1270                1275

tct gag  cag agc cct tct att  ggg tac tcc tca agg  act ctt ggc      3879
Ser Glu  Gln Ser Pro Ser Ile  Gly Tyr Ser Ser Arg  Thr Leu Gly
    1280                1285                1290

ccc aat  cct gga ctt att ctt  cag gct ttg act ctg  tca aac gct      3924
Pro Asn  Pro Gly Leu Ile Leu  Gln Ala Leu Thr Leu  Ser Asn Ala
    1295                1300                1305

agt gat  gga ttt aac ctg gag  cgg ctt gaa atg ctt  ggc gac tcc      3969
Ser Asp  Gly Phe Asn Leu Glu  Arg Leu Glu Met Leu  Gly Asp Ser
    1310                1315                1320

ttt tta  aag cat gcc atc acc  aca tat cta ttt tgc  act tac cct      4014
Phe Leu  Lys His Ala Ile Thr  Thr Tyr Leu Phe Cys  Thr Tyr Pro
    1325                1330                1335

gat gcg  cat gag ggc cgc ctt  tca tat atg aga agc  aaa aag gtc      4059
Asp Ala  His Glu Gly Arg Leu  Ser Tyr Met Arg Ser  Lys Lys Val
    1340                1345                1350

agc aac  tgt aat ctg tat cgc  ctt gga aaa aag aag  gga cta ccc      4104
Ser Asn  Cys Asn Leu Tyr Arg  Leu Gly Lys Lys Lys  Gly Leu Pro
    1355                1360                1365

agc cgc  atg gtg gtg tca ata  ttt gat ccc cct gtg  aat tgg ctt      4149
Ser Arg  Met Val Val Ser Ile  Phe Asp Pro Pro Val  Asn Trp Leu
    1370                1375                1380

cct cct  ggt tat gta gta aat  caa gac aaa agc aac  aca gat aaa      4194
Pro Pro  Gly Tyr Val Val Asn  Gln Asp Lys Ser Asn  Thr Asp Lys
    1385                1390                1395

tgg gaa  aaa gat gaa atg aca  aaa gac tgc atg ctg  gcg aat ggc      4239
Trp Glu  Lys Asp Glu Met Thr  Lys Asp Cys Met Leu  Ala Asn Gly
    1400                1405                1410

aaa ctg  gat gag gat tac gag  gag gag gat gag gag  gag gag agc      4284
Lys Leu  Asp Glu Asp Tyr Glu  Glu Glu Asp Glu Glu  Glu Glu Ser
    1415                1420                1425

ctg atg  tgg agg gct ccg aag  gaa gag gct gac tat  gaa gat gat      4329
Leu Met  Trp Arg Ala Pro Lys  Glu Glu Ala Asp Tyr  Glu Asp Asp
    1430                1435                1440

ttc ctg  gag tat gat cag gaa  cat atc aga ttt ata  gat aat atg      4374
Phe Leu  Glu Tyr Asp Gln Glu  His Ile Arg Phe Ile  Asp Asn Met
    1445                1450                1455

tta atg  ggg tca gga gct ttt  gta aag aaa atc tct  ctt tct cct      4419
Leu Met  Gly Ser Gly Ala Phe  Val Lys Lys Ile Ser  Leu Ser Pro
    1460                1465                1470

ttt tca  acc act gat tct gca  tat gaa tgg aaa atg  ccc aaa aaa      4464
```

-continued

```
Phe Ser  Thr Thr Asp Ser Ala  Tyr Glu Trp Lys Met  Pro Lys Lys
    1475                1480                1485

tcc tcc  tta ggt agt atg cca  ttt tca tca gat ttt  gag gat ttt      4509
Ser Ser  Leu Gly Ser Met Pro  Phe Ser Ser Asp Phe  Glu Asp Phe
    1490                1495                1500

gac tac  agc tct tgg gat gca  atg tgc tat ctg gat  cct agc aaa      4554
Asp Tyr  Ser Ser Trp Asp Ala  Met Cys Tyr Leu Asp  Pro Ser Lys
    1505                1510                1515

gct gtt  gaa gaa gat gac ttt  gtg gtg ggg ttc tgg  aat cca tca      4599
Ala Val  Glu Glu Asp Asp Phe  Val Val Gly Phe Trp  Asn Pro Ser
    1520                1525                1530

gaa gaa  aac tgt ggt gtt gac  acg gga aag cag tcc  att tct tac      4644
Glu Glu  Asn Cys Gly Val Asp  Thr Gly Lys Gln Ser  Ile Ser Tyr
    1535                1540                1545

gac ttg  cac act gag cag tgt  att gct gac aaa agc  ata gcg gac      4689
Asp Leu  His Thr Glu Gln Cys  Ile Ala Asp Lys Ser  Ile Ala Asp
    1550                1555                1560

tgt gtg  gaa gcc ctg ctg ggc  tgc tat tta acc agc  tgt ggg gag      4734
Cys Val  Glu Ala Leu Leu Gly  Cys Tyr Leu Thr Ser  Cys Gly Glu
    1565                1570                1575

agg gct  gct cag ctt ttc ctc  tgt tca ctg ggg ctg  aag gtg ctc      4779
Arg Ala  Ala Gln Leu Phe Leu  Cys Ser Leu Gly Leu  Lys Val Leu
    1580                1585                1590

ccg gta  att aaa agg act gat  cgg gaa aag gcc ctg  tgc cct act      4824
Pro Val  Ile Lys Arg Thr Asp  Arg Glu Lys Ala Leu  Cys Pro Thr
    1595                1600                1605

cgg gag  aat ttc aac agc caa  caa aag aac ctt tca  gtg agc tgt      4869
Arg Glu  Asn Phe Asn Ser Gln  Gln Lys Asn Leu Ser  Val Ser Cys
    1610                1615                1620

gct gct  gct tct gtg gcc agt  tca cgc tct tct gta  ttg aaa gac      4914
Ala Ala  Ala Ser Val Ala Ser  Ser Arg Ser Ser Val  Leu Lys Asp
    1625                1630                1635

tcg gaa  tat ggt tgt ttg aag  att cca cca aga tgt  atg ttt gat      4959
Ser Glu  Tyr Gly Cys Leu Lys  Ile Pro Pro Arg Cys  Met Phe Asp
    1640                1645                1650

cat cca  gat gca gat aaa aca  ctg aat cac ctt ata  tcg ggg ttt      5004
His Pro  Asp Ala Asp Lys Thr  Leu Asn His Leu Ile  Ser Gly Phe
    1655                1660                1665

gaa aat  ttt gaa aag aaa atc  aac tac aga ttc aag  aat aag gct      5049
Glu Asn  Phe Glu Lys Lys Ile  Asn Tyr Arg Phe Lys  Asn Lys Ala
    1670                1675                1680

tac ctt  ctc cag gct ttt aca  cat gcc tcc tac cac  tac aat act      5094
Tyr Leu  Leu Gln Ala Phe Thr  His Ala Ser Tyr His  Tyr Asn Thr
    1685                1690                1695

atc act  gat tgt tac cag cgc  tta gaa ttc ctg gga  gat gcg att      5139
Ile Thr  Asp Cys Tyr Gln Arg  Leu Glu Phe Leu Gly  Asp Ala Ile
    1700                1705                1710

ttg gac  tac ctc ata acc aag  cac ctt tat gaa gac  ccg cgg cag      5184
Leu Asp  Tyr Leu Ile Thr Lys  His Leu Tyr Glu Asp  Pro Arg Gln
    1715                1720                1725

cac tcc  ccg ggg gtc ctg aca  gac ctg cgg tct gcc  ctg gtc aac      5229
His Ser  Pro Gly Val Leu Thr  Asp Leu Arg Ser Ala  Leu Val Asn
    1730                1735                1740

aac acc  atc ttt gca tcg ctg  gct gta aag tac gac  tac cac aag      5274
Asn Thr  Ile Phe Ala Ser Leu  Ala Val Lys Tyr Asp  Tyr His Lys
    1745                1750                1755

tac ttc  aaa gct gtc tct cct  gag ctc ttc cat gtc  att gat gac      5319
```

-continued

```
Tyr Phe  Lys Ala Val Ser Pro  Glu Leu Phe His Val  Ile Asp Asp
    1760                1765                1770

ttt gtg  cag ttt cag ctt gag  aag aat gaa atg caa  gga atg gat      5364
Phe Val  Gln Phe Gln Leu Glu  Lys Asn Glu Met Gln  Gly Met Asp
    1775                1780                1785

tct gag  ctt agg aga tct gag  gag gat gaa gag aaa  gaa gag gat      5409
Ser Glu  Leu Arg Arg Ser Glu  Glu Asp Glu Glu Lys  Glu Glu Asp
    1790                1795                1800

att gaa  gtt cca aag gcc atg  ggg gat att ttt gag  tcg ctt gct      5454
Ile Glu  Val Pro Lys Ala Met  Gly Asp Ile Phe Glu  Ser Leu Ala
    1805                1810                1815

ggt gcc  att tac atg gat agt  ggg atg tca ctg gag  aca gtc tgg      5499
Gly Ala  Ile Tyr Met Asp Ser  Gly Met Ser Leu Glu  Thr Val Trp
    1820                1825                1830

cag gtg  tac tat ccc atg atg  cgg cca cta ata gaa  aag ttt tct      5544
Gln Val  Tyr Tyr Pro Met Met  Arg Pro Leu Ile Glu  Lys Phe Ser
    1835                1840                1845

gca aat  gta ccc cgt tcc cct  gtg cga gaa ttg ctt  gaa atg gaa      5589
Ala Asn  Val Pro Arg Ser Pro  Val Arg Glu Leu Leu  Glu Met Glu
    1850                1855                1860

cca gaa  act gcc aaa ttt agc  ccg gct gag aga act  tac gac ggg      5634
Pro Glu  Thr Ala Lys Phe Ser  Pro Ala Glu Arg Thr  Tyr Asp Gly
    1865                1870                1875

aag gtc  aga gtc act gtg gaa  gta gta gga aag ggg  aaa ttt aaa      5679
Lys Val  Arg Val Thr Val Glu  Val Val Gly Lys Gly  Lys Phe Lys
    1880                1885                1890

ggt gtt  ggt cga agt tac agg  att gcc aaa tct gca  gca gca aga      5724
Gly Val  Gly Arg Ser Tyr Arg  Ile Ala Lys Ser Ala  Ala Ala Arg
    1895                1900                1905

aga gcc  ctc cga agc ctc aaa  gct aat caa cct cag  gtt ccc aat      5769
Arg Ala  Leu Arg Ser Leu Lys  Ala Asn Gln Pro Gln  Val Pro Asn
    1910                1915                1920

agc tga                                                          5775
Ser
```

<210> SEQ ID NO 2
<211> LENGTH: 1924
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```
Met Lys Ser Pro Ala Leu Gln Pro Leu Ser Met Ala Gly Leu Gln Leu
1               5                  10                  15

Met Thr Pro Ala Ser Ser Pro Met Gly Pro Phe Phe Gly Leu Pro Trp
            20                  25                  30

Gln Gln Glu Ala Ile His Asp Asn Ile Tyr Thr Pro Arg Lys Tyr Gln
        35                  40                  45

Val Glu Leu Leu Glu Ala Ala Leu Asp His Asn Thr Ile Val Cys Leu
    50                  55                  60

Asn Thr Gly Ser Gly Lys Thr Phe Ile Ala Ser Thr Thr Leu Leu Lys
65                  70                  75                  80

Ser Cys Leu Tyr Leu Asp Leu Gly Glu Thr Ser Ala Arg Asn Gly Lys
                85                  90                  95

Arg Thr Val Phe Leu Val Asn Ser Ala Asn Gln Val Ala Gln Gln Val
            100                 105                 110

Ser Ala Val Arg Thr His Ser Asp Leu Lys Val Gly Glu Tyr Ser Asn
```

-continued

```
        115                 120                 125

Leu Glu Val Asn Ala Ser Trp Thr Lys Glu Arg Trp Asn Gln Glu Phe
    130                 135                 140

Thr Lys His Gln Val Leu Ile Met Thr Cys Tyr Val Ala Leu Asn Val
145                 150                 155                 160

Leu Lys Asn Gly Tyr Leu Ser Leu Ser Asp Ile Asn Leu Leu Val Phe
                165                 170                 175

Asp Glu Cys His Leu Ala Ile Leu Asp His Pro Tyr Arg Glu Phe Met
            180                 185                 190

Lys Leu Cys Glu Ile Cys Pro Ser Cys Pro Arg Ile Leu Gly Leu Thr
        195                 200                 205

Ala Ser Ile Leu Asn Gly Lys Trp Asp Pro Glu Asp Leu Glu Glu Lys
    210                 215                 220

Phe Gln Lys Leu Glu Lys Ile Leu Lys Ser Asn Ala Glu Thr Ala Thr
225                 230                 235                 240

Asp Leu Val Val Leu Asp Arg Tyr Thr Ser Gln Pro Cys Glu Ile Val
                245                 250                 255

Val Asp Cys Gly Pro Phe Thr Asp Arg Ser Gly Leu Tyr Glu Arg Leu
            260                 265                 270

Leu Met Glu Leu Glu Glu Ala Leu Asn Phe Ile Asn Asp Cys Asn Ile
        275                 280                 285

Ser Val His Ser Lys Glu Arg Asp Ser Thr Leu Ile Ser Lys Gln Ile
    290                 295                 300

Leu Ser Asp Cys Arg Ala Val Leu Val Val Leu Gly Pro Trp Cys Ala
305                 310                 315                 320

Asp Lys Val Ala Gly Met Met Val Arg Glu Leu Gln Lys Tyr Ile Lys
                325                 330                 335

His Glu Gln Glu Glu Leu His Arg Lys Phe Leu Leu Phe Thr Asp Thr
            340                 345                 350

Phe Leu Arg Lys Ile His Ala Leu Cys Glu Glu His Phe Ser Pro Ala
        355                 360                 365

Ser Leu Asp Leu Lys Phe Val Thr Pro Lys Val Ile Lys Leu Leu Glu
    370                 375                 380

Ile Leu Arg Lys Tyr Lys Pro Tyr Glu Arg His Ser Phe Glu Ser Val
385                 390                 395                 400

Glu Trp Tyr Asn Asn Arg Asn Gln Asp Asn Tyr Val Ser Trp Ser Asp
                405                 410                 415

Ser Glu Asp Asp Asp Glu Asp Glu Glu Ile Glu Glu Lys Glu Lys Pro
            420                 425                 430

Glu Thr Asn Phe Pro Ser Pro Phe Thr Asn Ile Leu Cys Gly Ile Ile
        435                 440                 445

Phe Val Glu Arg Arg Tyr Thr Ala Val Val Leu Asn Arg Leu Ile Lys
    450                 455                 460

Glu Ala Gly Lys Gln Asp Pro Glu Leu Ala Tyr Ile Ser Ser Asn Phe
465                 470                 475                 480

Ile Thr Gly His Gly Ile Gly Lys Asn Gln Pro Arg Asn Asn Thr Met
                485                 490                 495

Glu Ala Glu Phe Arg Lys Gln Glu Glu Val Leu Arg Lys Phe Arg Ala
            500                 505                 510

His Glu Thr Asn Leu Leu Ile Ala Thr Ser Ile Val Glu Glu Gly Val
        515                 520                 525
```

-continued

```
Asp Ile Pro Lys Cys Asn Leu Val Val Arg Phe Asp Leu Pro Thr Glu
    530                 535                 540

Tyr Arg Ser Tyr Val Gln Ser Lys Gly Arg Ala Arg Ala Pro Ile Ser
545                 550                 555                 560

Asn Tyr Ile Met Leu Ala Asp Thr Asp Lys Ile Lys Ser Phe Glu Glu
                565                 570                 575

Asp Leu Lys Thr Tyr Lys Ala Ile Glu Lys Ile Leu Arg Asn Lys Cys
            580                 585                 590

Ser Lys Ser Val Asp Thr Gly Glu Thr Asp Ile Asp Pro Val Met Asp
        595                 600                 605

Asp Asp His Val Phe Pro Pro Tyr Val Leu Arg Pro Asp Asp Gly Gly
    610                 615                 620

Pro Arg Val Thr Ile Asn Thr Ala Ile Gly His Ile Asn Arg Tyr Cys
625                 630                 635                 640

Ala Arg Leu Pro Ser Asp Pro Phe Thr His Leu Ala Pro Lys Cys Arg
                645                 650                 655

Thr Arg Glu Leu Pro Asp Gly Thr Phe Tyr Ser Thr Leu Tyr Leu Pro
            660                 665                 670

Ile Asn Ser Pro Leu Arg Ala Ser Ile Val Gly Pro Pro Met Ser Cys
        675                 680                 685

Val Arg Leu Ala Glu Arg Val Val Ala Leu Ile Cys Cys Glu Lys Leu
    690                 695                 700

His Lys Ile Gly Glu Leu Asp Asp His Leu Met Pro Val Gly Lys Glu
705                 710                 715                 720

Thr Val Lys Tyr Glu Glu Glu Leu Asp Leu His Asp Glu Glu Glu Thr
                725                 730                 735

Ser Val Pro Gly Arg Pro Gly Ser Thr Lys Arg Arg Gln Cys Tyr Pro
            740                 745                 750

Lys Ala Ile Pro Glu Cys Leu Arg Asp Ser Tyr Pro Arg Pro Asp Gln
        755                 760                 765

Pro Cys Tyr Leu Tyr Val Ile Gly Met Val Leu Thr Thr Pro Leu Pro
    770                 775                 780

Asp Glu Leu Asn Phe Arg Arg Arg Lys Leu Tyr Pro Pro Glu Asp Thr
785                 790                 795                 800

Thr Arg Cys Phe Gly Ile Leu Thr Ala Lys Pro Ile Pro Gln Ile Pro
                805                 810                 815

His Phe Pro Val Tyr Thr Arg Ser Gly Glu Val Thr Ile Ser Ile Glu
            820                 825                 830

Leu Lys Lys Ser Gly Phe Met Leu Ser Leu Gln Met Leu Glu Leu Ile
        835                 840                 845

Thr Arg Leu His Gln Tyr Ile Phe Ser His Ile Leu Arg Leu Glu Lys
    850                 855                 860

Pro Ala Leu Glu Phe Lys Pro Thr Asp Ala Asp Ser Ala Tyr Cys Val
865                 870                 875                 880

Leu Pro Leu Asn Val Val Asn Asp Ser Ser Thr Leu Asp Ile Asp Phe
                885                 890                 895

Lys Phe Met Glu Asp Ile Glu Lys Ser Glu Ala Arg Ile Gly Ile Pro
            900                 905                 910

Ser Thr Lys Tyr Thr Lys Glu Thr Pro Phe Val Phe Lys Leu Glu Asp
        915                 920                 925
```

-continued

```
Tyr Gln Asp Ala Val Ile Ile Pro Arg Tyr Arg Asn Phe Asp Gln Pro
    930                 935                 940

His Arg Phe Tyr Val Ala Asp Val Tyr Thr Asp Leu Thr Pro Leu Ser
945                 950                 955                 960

Lys Phe Pro Ser Pro Glu Tyr Glu Thr Phe Ala Glu Tyr Tyr Lys Thr
                965                 970                 975

Lys Tyr Asn Leu Asp Leu Thr Asn Leu Asn Gln Pro Leu Leu Asp Val
            980                 985                 990

Asp His Thr Ser Ser Arg Leu Asn  Leu Leu Thr Pro Arg  His Leu Asn
        995                 1000                1005

Gln Lys  Gly Lys Ala Leu Pro  Leu Ser Ser Ala Glu  Lys Arg Lys
    1010                1015                1020

Ala Lys  Trp Glu Ser Leu Gln  Asn Lys Gln Ile Leu  Val Pro Glu
    1025                1030                1035

Leu Cys  Ala Ile His Pro Ile  Pro Ala Ser Leu Trp  Arg Lys Ala
    1040                1045                1050

Val Cys  Leu Pro Ser Ile Leu  Tyr Arg Leu His Cys  Leu Leu Thr
    1055                1060                1065

Ala Glu  Glu Leu Arg Ala Gln  Thr Ala Ser Asp Ala  Gly Val Gly
    1070                1075                1080

Val Arg  Ser Leu Pro Ala Asp  Phe Arg Tyr Pro Asn  Leu Asp Phe
    1085                1090                1095

Gly Trp  Lys Lys Ser Ile Asp  Ser Lys Ser Phe Ile  Ser Ile Ser
    1100                1105                1110

Asn Ser  Ser Ser Ala Glu Asn  Asp Asn Tyr Cys Lys  His Ser Thr
    1115                1120                1125

Ile Val  Pro Glu Asn Ala Ala  His Gln Gly Ala Asn  Arg Thr Ser
    1130                1135                1140

Ser Leu  Glu Asn His Asp Gln  Met Ser Val Asn Cys  Arg Thr Leu
    1145                1150                1155

Leu Ser  Glu Ser Pro Gly Lys  Leu His Val Glu Val  Ser Ala Asp
    1160                1165                1170

Leu Thr  Ala Ile Asn Gly Leu  Ser Tyr Asn Gln Asn  Leu Ala Asn
    1175                1180                1185

Gly Ser  Tyr Asp Leu Ala Asn  Arg Asp Phe Cys Gln  Gly Asn Gln
    1190                1195                1200

Leu Asn  Tyr Tyr Lys Gln Glu  Ile Pro Val Gln Pro  Thr Thr Ser
    1205                1210                1215

Tyr Ser  Ile Gln Asn Leu Tyr  Ser Tyr Glu Asn Gln  Pro Gln Pro
    1220                1225                1230

Ser Asp  Glu Cys Thr Leu Leu  Ser Asn Lys Tyr Leu  Asp Gly Asn
    1235                1240                1245

Ala Asn  Lys Ser Thr Ser Asp  Gly Ser Pro Val Met  Ala Val Met
    1250                1255                1260

Pro Gly  Thr Thr Asp Thr Ile  Gln Val Leu Lys Gly  Arg Met Asp
    1265                1270                1275

Ser Glu  Gln Ser Pro Ser Ile  Gly Tyr Ser Ser Arg  Thr Leu Gly
    1280                1285                1290

Pro Asn  Pro Gly Leu Ile Leu  Gln Ala Leu Thr Leu  Ser Asn Ala
    1295                1300                1305

Ser Asp  Gly Phe Asn Leu Glu  Arg Leu Glu Met Leu  Gly Asp Ser
```

-continued

```
    1310                1315                1320

Phe Leu  Lys His Ala Ile Thr  Thr Tyr Leu Phe Cys  Thr Tyr Pro
    1325                1330                1335

Asp Ala  His Glu Gly Arg Leu  Ser Tyr Met Arg Ser  Lys Lys Val
    1340                1345                1350

Ser Asn  Cys Asn Leu Tyr Arg  Leu Gly Lys Lys Lys  Gly Leu Pro
    1355                1360                1365

Ser Arg  Met Val Val Ser Ile  Phe Asp Pro Pro Val  Asn Trp Leu
    1370                1375                1380

Pro Pro  Gly Tyr Val Val Asn  Gln Asp Lys Ser Asn  Thr Asp Lys
    1385                1390                1395

Trp Glu  Lys Asp Glu Met Thr  Lys Asp Cys Met Leu  Ala Asn Gly
    1400                1405                1410

Lys Leu  Asp Glu Asp Tyr Glu  Glu Glu Asp Glu Glu  Glu Glu Ser
    1415                1420                1425

Leu Met  Trp Arg Ala Pro Lys  Glu Glu Ala Asp Tyr  Glu Asp Asp
    1430                1435                1440

Phe Leu  Glu Tyr Asp Gln Glu  His Ile Arg Phe Ile  Asp Asn Met
    1445                1450                1455

Leu Met  Gly Ser Gly Ala Phe  Val Lys Lys Ile Ser  Leu Ser Pro
    1460                1465                1470

Phe Ser  Thr Thr Asp Ser Ala  Tyr Glu Trp Lys Met  Pro Lys Lys
    1475                1480                1485

Ser Ser  Leu Gly Ser Met Pro  Phe Ser Ser Asp Phe  Glu Asp Phe
    1490                1495                1500

Asp Tyr  Ser Ser Trp Asp Ala  Met Cys Tyr Leu Asp  Pro Ser Lys
    1505                1510                1515

Ala Val  Glu Glu Asp Asp Phe  Val Val Gly Phe Trp  Asn Pro Ser
    1520                1525                1530

Glu Glu  Asn Cys Gly Val Asp  Thr Gly Lys Gln Ser  Ile Ser Tyr
    1535                1540                1545

Asp Leu  His Thr Glu Gln Cys  Ile Ala Asp Lys Ser  Ile Ala Asp
    1550                1555                1560

Cys Val  Glu Ala Leu Leu Gly  Cys Tyr Leu Thr Ser  Cys Gly Glu
    1565                1570                1575

Arg Ala  Ala Gln Leu Phe Leu  Cys Ser Leu Gly Leu  Lys Val Leu
    1580                1585                1590

Pro Val  Ile Lys Arg Thr Asp  Arg Glu Lys Ala Leu  Cys Pro Thr
    1595                1600                1605

Arg Glu  Asn Phe Asn Ser Gln  Gln Lys Asn Leu Ser  Val Ser Cys
    1610                1615                1620

Ala Ala  Ala Ser Val Ala Ser  Ser Arg Ser Ser Val  Leu Lys Asp
    1625                1630                1635

Ser Glu  Tyr Gly Cys Leu Lys  Ile Pro Pro Arg Cys  Met Phe Asp
    1640                1645                1650

His Pro  Asp Ala Asp Lys Thr  Leu Asn His Leu Ile  Ser Gly Phe
    1655                1660                1665

Glu Asn  Phe Glu Lys Lys Ile  Asn Tyr Arg Phe Lys  Asn Lys Ala
    1670                1675                1680

Tyr Leu  Leu Gln Ala Phe Thr  His Ala Ser Tyr His  Tyr Asn Thr
    1685                1690                1695
```

-continued

```
Ile Thr  Asp Cys Tyr Gln Arg  Leu Glu Phe Leu Gly  Asp Ala Ile
    1700                1705                1710

Leu Asp  Tyr Leu Ile Thr Lys  His Leu Tyr Glu Asp  Pro Arg Gln
    1715                1720                1725

His Ser  Pro Gly Val Leu Thr  Asp Leu Arg Ser Ala  Leu Val Asn
    1730                1735                1740

Asn Thr  Ile Phe Ala Ser Leu  Ala Val Lys Tyr Asp  Tyr His Lys
    1745                1750                1755

Tyr Phe  Lys Ala Val Ser Pro  Glu Leu Phe His Val  Ile Asp Asp
    1760                1765                1770

Phe Val  Gln Phe Gln Leu Glu  Lys Asn Glu Met Gln  Gly Met Asp
    1775                1780                1785

Ser Glu  Leu Arg Arg Ser Glu  Glu Asp Glu Glu Lys  Glu Glu Asp
    1790                1795                1800

Ile Glu  Val Pro Lys Ala Met  Gly Asp Ile Phe Glu  Ser Leu Ala
    1805                1810                1815

Gly Ala  Ile Tyr Met Asp Ser  Gly Met Ser Leu Glu  Thr Val Trp
    1820                1825                1830

Gln Val  Tyr Tyr Pro Met Met  Arg Pro Leu Ile Glu  Lys Phe Ser
    1835                1840                1845

Ala Asn  Val Pro Arg Ser Pro  Val Arg Glu Leu Leu  Glu Met Glu
    1850                1855                1860

Pro Glu  Thr Ala Lys Phe Ser  Pro Ala Glu Arg Thr  Tyr Asp Gly
    1865                1870                1875

Lys Val  Arg Val Thr Val Glu  Val Val Gly Lys Gly  Lys Phe Lys
    1880                1885                1890

Gly Val  Gly Arg Ser Tyr Arg  Ile Ala Lys Ser Ala  Ala Ala Arg
    1895                1900                1905

Arg Ala  Leu Arg Ser Leu Lys  Ala Asn Gln Pro Gln  Val Pro Asn
    1910                1915                1920

Ser
```

<210> SEQ ID NO 3
<211> LENGTH: 6750
<212> TYPE: DNA
<213> ORGANISM: Drosophila melanogaster
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(6750)

<400> SEQUENCE: 3

```
atg gcg ttc cac tgg tgc gac aac aat ctg cac acc acc gtg ttc acg       48
Met Ala Phe His Trp Cys Asp Asn Asn Leu His Thr Thr Val Phe Thr
1               5                   10                  15

ccg cgc gac ttt cag gtg gag cta ctg gcc acc gcc tac gag cgg aac       96
Pro Arg Asp Phe Gln Val Glu Leu Leu Ala Thr Ala Tyr Glu Arg Asn
            20                  25                  30

acg att att tgc ctg ggc cat cga agt tcc aag gag ttt ata gcc ctc      144
Thr Ile Ile Cys Leu Gly His Arg Ser Ser Lys Glu Phe Ile Ala Leu
        35                  40                  45

aag ctg ctc cag gag ctg tcg cgt cga gca cgc cga cat ggt cgt gtc      192
Lys Leu Leu Gln Glu Leu Ser Arg Arg Ala Arg Arg His Gly Arg Val
    50                  55                  60

agt gtc tat ctc agt tgc gag gtt ggc acc agc acg gaa cca tgc tcc      240
```

-continued

```
Ser Val Tyr Leu Ser Cys Glu Val Gly Thr Ser Thr Glu Pro Cys Ser
65                  70                  75                  80

atc tac acg atg ctc acc cac ttg act gac ctg cgg gtg tgg cag gag      288
Ile Tyr Thr Met Leu Thr His Leu Thr Asp Leu Arg Val Trp Gln Glu
                85                  90                  95

cag ccg gat atg caa att ccc ttt gat cat tgc tgg acg gac tat cac      336
Gln Pro Asp Met Gln Ile Pro Phe Asp His Cys Trp Thr Asp Tyr His
            100                 105                 110

gtt tcc atc cta cgg cca gag gga ttt ctt tat ctg ctc gaa act cgc      384
Val Ser Ile Leu Arg Pro Glu Gly Phe Leu Tyr Leu Leu Glu Thr Arg
        115                 120                 125

gag ctg ctg ctg agc agc gtc gaa ctg atc gtg ctg gaa gat tgt cat      432
Glu Leu Leu Leu Ser Ser Val Glu Leu Ile Val Leu Glu Asp Cys His
    130                 135                 140

gac agc gcc gtt tat cag agg ata agg cct ctg ttc gag aat cac att      480
Asp Ser Ala Val Tyr Gln Arg Ile Arg Pro Leu Phe Glu Asn His Ile
145                 150                 155                 160

atg cca gcg cca ccg gcg gac agg cca cgg att ctc gga ctc gct gga      528
Met Pro Ala Pro Pro Ala Asp Arg Pro Arg Ile Leu Gly Leu Ala Gly
                165                 170                 175

ccg ctg cac agc gcc gga tgt gag ctg cag caa ctg agc gcc atg ctg      576
Pro Leu His Ser Ala Gly Cys Glu Leu Gln Gln Leu Ser Ala Met Leu
            180                 185                 190

gcc acc ctg gag cag agt gtg ctt tgc cag atc gag acg gcc agt gat      624
Ala Thr Leu Glu Gln Ser Val Leu Cys Gln Ile Glu Thr Ala Ser Asp
        195                 200                 205

att gtc acc gtg ttg cgt tac tgt tcc cga ccg cac gaa tac atc gta      672
Ile Val Thr Val Leu Arg Tyr Cys Ser Arg Pro His Glu Tyr Ile Val
    210                 215                 220

cag tgc gcc ccc ttc gag atg gac gaa ctg tcc ctg gtg ctt gcc gat      720
Gln Cys Ala Pro Phe Glu Met Asp Glu Leu Ser Leu Val Leu Ala Asp
225                 230                 235                 240

gtg ctc aac aca cac aag tcc ttt tta ttg gac cac cgc tac gat ccc      768
Val Leu Asn Thr His Lys Ser Phe Leu Leu Asp His Arg Tyr Asp Pro
                245                 250                 255

tac gaa atc tac ggc aca gac cag ttt atg gac gaa ctg aaa gac ata      816
Tyr Glu Ile Tyr Gly Thr Asp Gln Phe Met Asp Glu Leu Lys Asp Ile
            260                 265                 270

ccc gat ccc aag gtg gac ccc ctg aac gtc atc aac tca cta ctg gtc      864
Pro Asp Pro Lys Val Asp Pro Leu Asn Val Ile Asn Ser Leu Leu Val
        275                 280                 285

gtg ctg cac gag atg ggt cct tgg tgc acg cag cgg gct gca cat cac      912
Val Leu His Glu Met Gly Pro Trp Cys Thr Gln Arg Ala Ala His His
    290                 295                 300

ttt tac caa tgc aat gag aag tta aag gtg aag acg ccg cac gaa cgt      960
Phe Tyr Gln Cys Asn Glu Lys Leu Lys Val Lys Thr Pro His Glu Arg
305                 310                 315                 320

cac tac ttg ctg tac tgc cta gtg agc acg gcc ctt atc caa ctg tac     1008
His Tyr Leu Leu Tyr Cys Leu Val Ser Thr Ala Leu Ile Gln Leu Tyr
                325                 330                 335

tcc ctc tgc gaa cac gca ttc cat cga cat tta gga agt ggc agc gat     1056
Ser Leu Cys Glu His Ala Phe His Arg His Leu Gly Ser Gly Ser Asp
            340                 345                 350

tca cgc cag acc atc gaa cgc tat tcc agc ccc aag gtg cga cgt ctg     1104
Ser Arg Gln Thr Ile Glu Arg Tyr Ser Ser Pro Lys Val Arg Arg Leu
        355                 360                 365

ttg cag aca ctg agg tgc ttc aag ccg gaa gag gtg cac acc caa gcg     1152
```

-continued

```
Leu Gln Thr Leu Arg Cys Phe Lys Pro Glu Glu Val His Thr Gln Ala
    370                 375                 380

gac gga ctg cgc aga atg cgg cat cag gtg gat cag gcg gac ttc aat     1200
Asp Gly Leu Arg Arg Met Arg His Gln Val Asp Gln Ala Asp Phe Asn
385                 390                 395                 400

cgg tta tct cat acg ctg gaa agc aag tgc cga atg gtg gat caa atg     1248
Arg Leu Ser His Thr Leu Glu Ser Lys Cys Arg Met Val Asp Gln Met
                405                 410                 415

gac caa ccg ccg acg gag aca cga gcc ctg gtg gcc act ctt gag cag     1296
Asp Gln Pro Pro Thr Glu Thr Arg Ala Leu Val Ala Thr Leu Glu Gln
            420                 425                 430

att ctg cac acg aca gag gac agg cag acg aac aga agc gcc gct cgg     1344
Ile Leu His Thr Thr Glu Asp Arg Gln Thr Asn Arg Ser Ala Ala Arg
        435                 440                 445

gtg act cct act cct act ccc gct cat gcg aag ccg aaa cct agc tct     1392
Val Thr Pro Thr Pro Thr Pro Ala His Ala Lys Pro Lys Pro Ser Ser
    450                 455                 460

ggt gcc aac act gca caa cca cga act cgt aga cgt gtg tac acc agg     1440
Gly Ala Asn Thr Ala Gln Pro Arg Thr Arg Arg Arg Val Tyr Thr Arg
465                 470                 475                 480

cgc cac cac cgg gat cac aat gat ggc agc gac acg ctc tgc gca ctg     1488
Arg His His Arg Asp His Asn Asp Gly Ser Asp Thr Leu Cys Ala Leu
                485                 490                 495

att tac tgc aac cag aac cac acg gct cgc gtg ctc ttt gag ctt cta     1536
Ile Tyr Cys Asn Gln Asn His Thr Ala Arg Val Leu Phe Glu Leu Leu
            500                 505                 510

gcg gag att agc aga cgt gat ccc gat ctc aag ttc cta cgc tgc cag     1584
Ala Glu Ile Ser Arg Arg Asp Pro Asp Leu Lys Phe Leu Arg Cys Gln
        515                 520                 525

tac acc acg gac cgg gtg gca gat ccc acc acg gag ccc aaa gag gct     1632
Tyr Thr Thr Asp Arg Val Ala Asp Pro Thr Thr Glu Pro Lys Glu Ala
    530                 535                 540

gag ttg gag cac cgg cgg cag gaa gag gtg cta aag cgc ttc cgc atg     1680
Glu Leu Glu His Arg Arg Gln Glu Glu Val Leu Lys Arg Phe Arg Met
545                 550                 555                 560

cat gac tgc aat gtc ctg atc ggt act tcg gtg ctg gaa gag ggc atc     1728
His Asp Cys Asn Val Leu Ile Gly Thr Ser Val Leu Glu Glu Gly Ile
                565                 570                 575

gat gtg ccc aag tgc aat ttg gtt gtg cgc tgg gat ccg cca acc aca     1776
Asp Val Pro Lys Cys Asn Leu Val Val Arg Trp Asp Pro Pro Thr Thr
            580                 585                 590

tat cgc agt tac gtt cag tgc aaa ggt cga gcc cgt gct gct cca gcc     1824
Tyr Arg Ser Tyr Val Gln Cys Lys Gly Arg Ala Arg Ala Ala Pro Ala
        595                 600                 605

tat cat gtc att ctc gtc gct ccg agt tat aaa agc cca act gtg ggg     1872
Tyr His Val Ile Leu Val Ala Pro Ser Tyr Lys Ser Pro Thr Val Gly
    610                 615                 620

tca gtg cag ctg acc gat cgg agt cat cgg tat att tgc gcg act ggt     1920
Ser Val Gln Leu Thr Asp Arg Ser His Arg Tyr Ile Cys Ala Thr Gly
625                 630                 635                 640

gat act aca gag gcg gac agc gac tct gat gat tca gcg atg cca aac     1968
Asp Thr Thr Glu Ala Asp Ser Asp Ser Asp Asp Ser Ala Met Pro Asn
                645                 650                 655

tcg tcc ggc tcg gat ccc tat act ttt ggc acg gca cgc gga acc gtg     2016
Ser Ser Gly Ser Asp Pro Tyr Thr Phe Gly Thr Ala Arg Gly Thr Val
            660                 665                 670

aag atc ctc aac ccc gaa gtg ttc agt aaa caa cca ccg aca gcg tgc     2064
```

-continued

```
Lys Ile Leu Asn Pro Glu Val Phe Ser Lys Gln Pro Pro Thr Ala Cys
        675                 680                 685

gac att aag ctg cag gag atc cag gac gaa ttg cca gcc gca gcg cag     2112
Asp Ile Lys Leu Gln Glu Ile Gln Asp Glu Leu Pro Ala Ala Ala Gln
    690                 695                 700

ctg gat acg agc aac tcc agc gac gaa gcc gtc agc atg agt aac acg     2160
Leu Asp Thr Ser Asn Ser Ser Asp Glu Ala Val Ser Met Ser Asn Thr
705                 710                 715                 720

tct cca agc gag agc agt aca gaa caa aaa tcc aga cgc ttc cag tgc     2208
Ser Pro Ser Glu Ser Ser Thr Glu Gln Lys Ser Arg Arg Phe Gln Cys
                725                 730                 735

gag ctg agc tct tta acg gag cca gaa gac aca agt gat act aca gcc     2256
Glu Leu Ser Ser Leu Thr Glu Pro Glu Asp Thr Ser Asp Thr Thr Ala
            740                 745                 750

gaa atc gat act gct cat agt tta gcc agc acc acg aag gac ttg gtg     2304
Glu Ile Asp Thr Ala His Ser Leu Ala Ser Thr Thr Lys Asp Leu Val
        755                 760                 765

cat caa atg gca cag tat cgc gaa atc gag cag atg ctg cta tcc aag     2352
His Gln Met Ala Gln Tyr Arg Glu Ile Glu Gln Met Leu Leu Ser Lys
    770                 775                 780

tgc gcc aac aca gag ccg ccg gag cag gag cag agt gag gcg gaa cgt     2400
Cys Ala Asn Thr Glu Pro Pro Glu Gln Glu Gln Ser Glu Ala Glu Arg
785                 790                 795                 800

ttt agt gcc tgc ctg gcc gca tac cga ccc aag ccg cac ctg cta aca     2448
Phe Ser Ala Cys Leu Ala Ala Tyr Arg Pro Lys Pro His Leu Leu Thr
                805                 810                 815

ggc gcc tcc gtg gat ctg ggt tct gct ata gct ttg gtc aac aag tac     2496
Gly Ala Ser Val Asp Leu Gly Ser Ala Ile Ala Leu Val Asn Lys Tyr
            820                 825                 830

tgc gcc cga ctg cca agc gac acg ttc acc aag ttg acg gcg ttg tgg     2544
Cys Ala Arg Leu Pro Ser Asp Thr Phe Thr Lys Leu Thr Ala Leu Trp
        835                 840                 845

cgc tgc acc cga aac gaa agg gct gga gtg acc ctg ttt cag tac aca     2592
Arg Cys Thr Arg Asn Glu Arg Ala Gly Val Thr Leu Phe Gln Tyr Thr
    850                 855                 860

ctc cgt ctg ccc atc aac tcg cca ttg aag cat gac att gtg ggt ctt     2640
Leu Arg Leu Pro Ile Asn Ser Pro Leu Lys His Asp Ile Val Gly Leu
865                 870                 875                 880

ccg atg cca act caa aca ttg gcc cgc cga ctg gct gcc ttg cag gct     2688
Pro Met Pro Thr Gln Thr Leu Ala Arg Arg Leu Ala Ala Leu Gln Ala
                885                 890                 895

tgc gtg gaa ctg cac agg atc ggt gag tta gac gat cag ttg cag cct     2736
Cys Val Glu Leu His Arg Ile Gly Glu Leu Asp Asp Gln Leu Gln Pro
            900                 905                 910

atc ggc aag gag gga ttt cgt gcc ctg gag ccg gac tgg gag tgc ttt     2784
Ile Gly Lys Glu Gly Phe Arg Ala Leu Glu Pro Asp Trp Glu Cys Phe
        915                 920                 925

gaa ctg gag cca gag gac gaa cag att gtg cag cta agc gat gaa cca     2832
Glu Leu Glu Pro Glu Asp Glu Gln Ile Val Gln Leu Ser Asp Glu Pro
    930                 935                 940

cgt ccg gga aca acg aag cgt cgt cag tac tat tac aaa cgc att gca     2880
Arg Pro Gly Thr Thr Lys Arg Arg Gln Tyr Tyr Tyr Lys Arg Ile Ala
945                 950                 955                 960

tcc gaa ttt tgc gat tgc cgt ccc gtt gcc gga gcg cca tgc tat ttg     2928
Ser Glu Phe Cys Asp Cys Arg Pro Val Ala Gly Ala Pro Cys Tyr Leu
                965                 970                 975

tac ttt atc caa ctg acg ctc caa tgt ccg att ccc gaa gag caa aac     2976
```

-continued

```
Tyr Phe Ile Gln Leu Thr Leu Gln Cys Pro Ile Pro Glu Glu Gln Asn
           980                 985                 990

acg cgg gga cgc aag att tat ccg  ccc gaa gat gcg cag  cag gga ttc   3024
Thr Arg Gly Arg Lys Ile Tyr Pro  Pro Glu Asp Ala Gln  Gln Gly Phe
        995                1000                1005

ggc att  cta acc acc aaa cgg  ata ccc aag ctg agt  gct ttc tcg      3069
Gly Ile  Leu Thr Thr Lys Arg  Ile Pro Lys Leu Ser  Ala Phe Ser
    1010                1015                1020

ata ttc  acg cgt tcc ggt gag  gtg aag gtt tcc ctg  gag tta gct      3114
Ile Phe  Thr Arg Ser Gly Glu  Val Lys Val Ser Leu  Glu Leu Ala
    1025                1030                1035

aag gaa  cgc gtg att cta act  agc gaa caa ata gtc  tgc atc aac      3159
Lys Glu  Arg Val Ile Leu Thr  Ser Glu Gln Ile Val  Cys Ile Asn
    1040                1045                1050

gga ttt  tta aac tac acg ttc  acc aat gta ctg cgt  ttg caa aag      3204
Gly Phe  Leu Asn Tyr Thr Phe  Thr Asn Val Leu Arg  Leu Gln Lys
    1055                1060                1065

ttt ctg  atg ctc ttc gat ccg  gac tcc acg gaa aat  tgt gta ttc      3249
Phe Leu  Met Leu Phe Asp Pro  Asp Ser Thr Glu Asn  Cys Val Phe
    1070                1075                1080

att gtg  ccc acc gtg aag gca  cca gct ggc ggc aag  cac atc gac      3294
Ile Val  Pro Thr Val Lys Ala  Pro Ala Gly Gly Lys  His Ile Asp
    1085                1090                1095

tgg cag  ttt ctg gag ctg atc  caa gcg aat gga aat  aca atg cca      3339
Trp Gln  Phe Leu Glu Leu Ile  Gln Ala Asn Gly Asn  Thr Met Pro
    1100                1105                1110

cgg gca  gtg ccc gat gag gag  cgc cag gcg cag ccg  ttt gat ccg      3384
Arg Ala  Val Pro Asp Glu Glu  Arg Gln Ala Gln Pro  Phe Asp Pro
    1115                1120                1125

caa cgc  ttc cag gat gcc gtc  gtt atg ccg tgg tat  cgc aac cag      3429
Gln Arg  Phe Gln Asp Ala Val  Val Met Pro Trp Tyr  Arg Asn Gln
    1130                1135                1140

gat caa  ccg cag tat ttc tat  gtg gcg gag ata tgt  cca cat cta      3474
Asp Gln  Pro Gln Tyr Phe Tyr  Val Ala Glu Ile Cys  Pro His Leu
    1145                1150                1155

tcc cca  ctc agc tgc ttc cct  ggt gac aac tac cgc  acg ttc aag      3519
Ser Pro  Leu Ser Cys Phe Pro  Gly Asp Asn Tyr Arg  Thr Phe Lys
    1160                1165                1170

cac tac  tac ctc gtc aag tat  ggt ctg acc ata cag  aat acc tcg      3564
His Tyr  Tyr Leu Val Lys Tyr  Gly Leu Thr Ile Gln  Asn Thr Ser
    1175                1180                1185

cag ccg  cta ttg gac gtg gat  cac acc agt gcg cgg  tta aac ttc      3609
Gln Pro  Leu Leu Asp Val Asp  His Thr Ser Ala Arg  Leu Asn Phe
    1190                1195                1200

ctc acg  cca cga tac gtt aat  cgc aag ggc gtt gct  ctg ccc act      3654
Leu Thr  Pro Arg Tyr Val Asn  Arg Lys Gly Val Ala  Leu Pro Thr
    1205                1210                1215

agt tcg  gag gag aca aag cgg  gca aag cgc gag aat  ctc gaa cag      3699
Ser Ser  Glu Glu Thr Lys Arg  Ala Lys Arg Glu Asn  Leu Glu Gln
    1220                1225                1230

aag cag  atc ctt gtg cca gag  ctc tgc act gtg cat  cca ttc ccc      3744
Lys Gln  Ile Leu Val Pro Glu  Leu Cys Thr Val His  Pro Phe Pro
    1235                1240                1245

gcc tcc  ttg tgg cga act gcc  gtg tgc ctg ccc tgc  atc ctg tac      3789
Ala Ser  Leu Trp Arg Thr Ala  Val Cys Leu Pro Cys  Ile Leu Tyr
    1250                1255                1260

cgc ata  aat ggt ctt cta ttg  gcc gac gat att cgg  aaa cag gtt      3834
```

-continued

```
Arg Ile  Asn Gly Leu Leu Leu  Ala Asp Asp Ile Arg  Lys Gln Val
    1265                1270                1275

tct gcg  gat ctg ggg ctg gga  agg caa cag atc gaa  gat gag gat      3879
Ser Ala  Asp Leu Gly Leu Gly  Arg Gln Gln Ile Glu  Asp Glu Asp
    1280                1285                1290

ttc gag  tgg ccc atg ctg gac  ttt ggg tgg agt cta  tcg gag gtg      3924
Phe Glu  Trp Pro Met Leu Asp  Phe Gly Trp Ser Leu  Ser Glu Val
    1295                1300                1305

ctc aag  aaa tcg cgg gag tcc  aaa caa aag gag tcc  ctt aag gat      3969
Leu Lys  Lys Ser Arg Glu Ser  Lys Gln Lys Glu Ser  Leu Lys Asp
    1310                1315                1320

gat act  att aat ggc aaa gac  tta gct gat gtt gaa  aag aaa ccg      4014
Asp Thr  Ile Asn Gly Lys Asp  Leu Ala Asp Val Glu  Lys Lys Pro
    1325                1330                1335

act agc  gag gag acc caa cta  gat aag gat tca aaa  gac gat aag      4059
Thr Ser  Glu Glu Thr Gln Leu  Asp Lys Asp Ser Lys  Asp Asp Lys
    1340                1345                1350

gtt gag  aaa agt gct att gaa  cta atc att gag gga  gag gag aag      4104
Val Glu  Lys Ser Ala Ile Glu  Leu Ile Ile Glu Gly  Glu Glu Lys
    1355                1360                1365

ctg caa  gag gct gat gac ttc  att gag ata ggc act  tgg tca aac      4149
Leu Gln  Glu Ala Asp Asp Phe  Ile Glu Ile Gly Thr  Trp Ser Asn
    1370                1375                1380

gat atg  gcc gac gat ata gct  agt ttt aac caa gaa  gac gac gac      4194
Asp Met  Ala Asp Asp Ile Ala  Ser Phe Asn Gln Glu  Asp Asp Asp
    1385                1390                1395

gag gat  gac gcc ttc cat ctc  cca gtt tta ccg gca  aac gtt aag      4239
Glu Asp  Asp Ala Phe His Leu  Pro Val Leu Pro Ala  Asn Val Lys
    1400                1405                1410

ttc tgt  gat cag caa acg cgc  tac ggt tcg ccc aca  ttt tgg gat      4284
Phe Cys  Asp Gln Gln Thr Arg  Tyr Gly Ser Pro Thr  Phe Trp Asp
    1415                1420                1425

gtg agc  aat ggc gaa agc ggc  ttc aag ggt cca aag  agc agt cag      4329
Val Ser  Asn Gly Glu Ser Gly  Phe Lys Gly Pro Lys  Ser Ser Gln
    1430                1435                1440

aat aag  cag ggt ggc aag ggc  aaa gca aag ggt ccg  gca aag ccc      4374
Asn Lys  Gln Gly Gly Lys Gly  Lys Ala Lys Gly Pro  Ala Lys Pro
    1445                1450                1455

aca ttt  aac tat tat gac tcg  gac aat tcg ctg ggt  tcc agc tac      4419
Thr Phe  Asn Tyr Tyr Asp Ser  Asp Asn Ser Leu Gly  Ser Ser Tyr
    1460                1465                1470

gat gac  gac gat aac gca ggt  ccg ctc aat tac atg  cat cac aac      4464
Asp Asp  Asp Asp Asn Ala Gly  Pro Leu Asn Tyr Met  His His Asn
    1475                1480                1485

tac agt  tcg gat gac gac gat  gtg gca gat gat atc  gat gcg gga      4509
Tyr Ser  Ser Asp Asp Asp Asp  Val Ala Asp Asp Ile  Asp Ala Gly
    1490                1495                1500

cgc att  gcg ttc acc tcc aag  aat gaa gcg gag act  att gaa acc      4554
Arg Ile  Ala Phe Thr Ser Lys  Asn Glu Ala Glu Thr  Ile Glu Thr
    1505                1510                1515

gca cag  gaa gtg gaa aag cgc  cag aag cag ctg tcc  atc atc cag      4599
Ala Gln  Glu Val Glu Lys Arg  Gln Lys Gln Leu Ser  Ile Ile Gln
    1520                1525                1530

gcg acc  aat gct aac gag cgg  cag tat cag cag aca  aag aac ctg      4644
Ala Thr  Asn Ala Asn Glu Arg  Gln Tyr Gln Gln Thr  Lys Asn Leu
    1535                1540                1545

ctc att  gga ttc aat ttt aag  cat gag gac cag aag  gaa cct gcc      4689
```

-continued

```
Leu Ile  Gly Phe Asn Phe Lys  His Glu Asp Gln Lys  Glu Pro Ala
    1550                1555                1560

act ata  aga tat gaa gaa tcc  ata gct aag ctc aaa  acg gaa ata      4734
Thr Ile  Arg Tyr Glu Glu Ser  Ile Ala Lys Leu Lys  Thr Glu Ile
    1565                1570                1575

gaa tcc  ggc ggc atg ttg gtg  ccg cac gac cag cag  ttg gtt cta      4779
Glu Ser  Gly Gly Met Leu Val  Pro His Asp Gln Gln  Leu Val Leu
    1580                1585                1590

aaa aga  agt gat gcc gct gag  gct cag gtt gca aag  gta tcg atg      4824
Lys Arg  Ser Asp Ala Ala Glu  Ala Gln Val Ala Lys  Val Ser Met
    1595                1600                1605

atg gag  cta ttg aag cag ctg  ctg ccg tat gta aat  gaa gat gtg      4869
Met Glu  Leu Leu Lys Gln Leu  Leu Pro Tyr Val Asn  Glu Asp Val
    1610                1615                1620

ctg gcc  aaa aag ctg ggt gat  agg cgc gag ctt ctg  ctg tcg gat      4914
Leu Ala  Lys Lys Leu Gly Asp  Arg Arg Glu Leu Leu  Leu Ser Asp
    1625                1630                1635

ttg gta  gag cta aat gca gat  tgg gta gcg cga cat  gag cag gag      4959
Leu Val  Glu Leu Asn Ala Asp  Trp Val Ala Arg His  Glu Gln Glu
    1640                1645                1650

acc tac  aat gta atg gga tgc  gga gat agt ttt gac  aac tat aac      5004
Thr Tyr  Asn Val Met Gly Cys  Gly Asp Ser Phe Asp  Asn Tyr Asn
    1655                1660                1665

gat cat  cat cgg ctg aac ttg  gat gaa aag caa ctg  aaa ctg caa      5049
Asp His  His Arg Leu Asn Leu  Asp Glu Lys Gln Leu  Lys Leu Gln
    1670                1675                1680

tac gaa  cga att gaa att gag  cca cct act tcc acg  aag gcc ata      5094
Tyr Glu  Arg Ile Glu Ile Glu  Pro Pro Thr Ser Thr  Lys Ala Ile
    1685                1690                1695

acc tca  gcc ata tta cca gct  ggc ttc agt ttc gat  cga caa ccg      5139
Thr Ser  Ala Ile Leu Pro Ala  Gly Phe Ser Phe Asp  Arg Gln Pro
    1700                1705                1710

gat cta  gtg ggc cat cca gga  ccc agt ccc agc atc  att ttg caa      5184
Asp Leu  Val Gly His Pro Gly  Pro Ser Pro Ser Ile  Ile Leu Gln
    1715                1720                1725

gcc ctc  aca atg tcc aat gct  aac gat ggc atc aat  ctg gag cga      5229
Ala Leu  Thr Met Ser Asn Ala  Asn Asp Gly Ile Asn  Leu Glu Arg
    1730                1735                1740

ctg gag  aca att gga gat tcc  ttt cta aag tat gcc  att acc acc      5274
Leu Glu  Thr Ile Gly Asp Ser  Phe Leu Lys Tyr Ala  Ile Thr Thr
    1745                1750                1755

tac ttg  tac atc acc tac gag  aat gtg cac gag gga  aaa cta agt      5319
Tyr Leu  Tyr Ile Thr Tyr Glu  Asn Val His Glu Gly  Lys Leu Ser
    1760                1765                1770

cac ctg  cgc tcc aag cag gtt  gcc aat ctc aat ctc  tat cgt ctg      5364
His Leu  Arg Ser Lys Gln Val  Ala Asn Leu Asn Leu  Tyr Arg Leu
    1775                1780                1785

ggc aga  cgt aag aga ctg ggt  gaa tat atg ata gcc  act aaa ttc      5409
Gly Arg  Arg Lys Arg Leu Gly  Glu Tyr Met Ile Ala  Thr Lys Phe
    1790                1795                1800

gag ccg  cac gac aat tgg ctg  cca ccc tgc tac tac  gtg cca aag      5454
Glu Pro  His Asp Asn Trp Leu  Pro Pro Cys Tyr Tyr  Val Pro Lys
    1805                1810                1815

gag cta  gag aag gcg ctc atc  gag gcg aag atc ccc  act cac cat      5499
Glu Leu  Glu Lys Ala Leu Ile  Glu Ala Lys Ile Pro  Thr His His
    1820                1825                1830

tgg aag  ctg gcc gat ctg cta  gac att aag aac cta  agc agt gtg      5544
```

-continued

```
Trp Lys  Leu Ala Asp Leu Leu  Asp Ile Lys Asn Leu  Ser Ser Val
    1835                1840                1845

caa atc  tgc gag atg gtt cgc  gaa aaa gcc gat gcc  ctg ggc ttg      5589
Gln Ile  Cys Glu Met Val Arg  Glu Lys Ala Asp Ala  Leu Gly Leu
    1850                1855                1860

gag cag  aat ggg ggt gcc caa  aat gga caa ctt gac  gac tcc aat      5634
Glu Gln  Asn Gly Gly Ala Gln  Asn Gly Gln Leu Asp  Asp Ser Asn
    1865                1870                1875

gat agc  tgc aat gat ttt agc  tgt ttt att ccc tac  aac ctt gtt      5679
Asp Ser  Cys Asn Asp Phe Ser  Cys Phe Ile Pro Tyr  Asn Leu Val
    1880                1885                1890

tcg caa  cac agc att ccg gat  aag tct att gcc gat  tgc gtc gaa      5724
Ser Gln  His Ser Ile Pro Asp  Lys Ser Ile Ala Asp  Cys Val Glu
    1895                1900                1905

gcc ctc  att gga gcc tat ctc  att gag tgc gga ccc  cga ggg gct      5769
Ala Leu  Ile Gly Ala Tyr Leu  Ile Glu Cys Gly Pro  Arg Gly Ala
    1910                1915                1920

tta ctc  ttt atg gcc tgg ctg  ggc gtg aga gtg ctc  cct atc aca      5814
Leu Leu  Phe Met Ala Trp Leu  Gly Val Arg Val Leu  Pro Ile Thr
    1925                1930                1935

agg cag  ttg gac ggg ggt aac  cag gag caa cga ata  ccc ggt agc      5859
Arg Gln  Leu Asp Gly Gly Asn  Gln Glu Gln Arg Ile  Pro Gly Ser
    1940                1945                1950

aca aaa  ccg aat gcc gaa aat  gtg gtc acc gtt tac  ggt gca tgg      5904
Thr Lys  Pro Asn Ala Glu Asn  Val Val Thr Val Tyr  Gly Ala Trp
    1955                1960                1965

ccc acg  ccg cgt agt cca ctg  ctg cac ttt gct cca  aat gct acg      5949
Pro Thr  Pro Arg Ser Pro Leu  Leu His Phe Ala Pro  Asn Ala Thr
    1970                1975                1980

gag gag  ctg gac cag tta cta  agc ggc ttt gag gag  ttt gag gag      5994
Glu Glu  Leu Asp Gln Leu Leu  Ser Gly Phe Glu Glu  Phe Glu Glu
    1985                1990                1995

agt ttg  gga tac aag ttc cgg  gat cgg tcg tac ctg  ttg caa gcc      6039
Ser Leu  Gly Tyr Lys Phe Arg  Asp Arg Ser Tyr Leu  Leu Gln Ala
    2000                2005                2010

atg aca  cat gcc agt tac acg  ccc aat cga ttg acg  gat tgc tat      6084
Met Thr  His Ala Ser Tyr Thr  Pro Asn Arg Leu Thr  Asp Cys Tyr
    2015                2020                2025

cag cgt  ctg gag ttc ctg ggc  gat gct gtt cta gat  tac ctc att      6129
Gln Arg  Leu Glu Phe Leu Gly  Asp Ala Val Leu Asp  Tyr Leu Ile
    2030                2035                2040

acg cgg  cat tta tac gaa gat  ccc cgc cag cat tct  cca ggc gca      6174
Thr Arg  His Leu Tyr Glu Asp  Pro Arg Gln His Ser  Pro Gly Ala
    2045                2050                2055

tta acg  gat ttg cgg tca gca  ctg gtg aat aat aca  ata ttc gcc      6219
Leu Thr  Asp Leu Arg Ser Ala  Leu Val Asn Asn Thr  Ile Phe Ala
    2060                2065                2070

tcc ctg  gct gtt cgc cat ggc  ttc cac aag ttc ttc  cgg cac ctc      6264
Ser Leu  Ala Val Arg His Gly  Phe His Lys Phe Phe  Arg His Leu
    2075                2080                2085

tcg ccg  ggc ctt aac gat gtg  att gac cgt ttt gtg  cgg atc cag      6309
Ser Pro  Gly Leu Asn Asp Val  Ile Asp Arg Phe Val  Arg Ile Gln
    2090                2095                2100

cag gag  aat gga cac tgc atc  agt gag gag tac tac  tta ttg tcc      6354
Gln Glu  Asn Gly His Cys Ile  Ser Glu Glu Tyr Tyr  Leu Leu Ser
    2105                2110                2115

gag gag  gag tgc gat gac gcc  gag gac gtt gag gtg  ccc aag gca      6399
```

-continued

```
Glu Glu  Glu Cys Asp Asp Ala  Glu Asp Val Glu Val  Pro Lys Ala
    2120                2125                2130

ttg ggc  gac gtt ttc gag tcg  atc gca ggt gcc att  ttt ctc gac      6444
Leu Gly  Asp Val Phe Glu Ser  Ile Ala Gly Ala Ile  Phe Leu Asp
    2135                2140                2145

tca aac  atg tcg ctg gac gtg  gtt tgg cac gta tat  agc aac atg      6489
Ser Asn  Met Ser Leu Asp Val  Val Trp His Val Tyr  Ser Asn Met
    2150                2155                2160

atg agc  ccg gag atc gag cag  ttc agc aac tca gtg  cca aaa tcg      6534
Met Ser  Pro Glu Ile Glu Gln  Phe Ser Asn Ser Val  Pro Lys Ser
    2165                2170                2175

ccc att  cgg gag ctc ctc gag  ctg gag ccg gaa acc  gcc aag ttc      6579
Pro Ile  Arg Glu Leu Leu Glu  Leu Glu Pro Glu Thr  Ala Lys Phe
    2180                2185                2190

ggc aag  ccc gag aag ctg gcg  gat ggg cga cgg gtg  cgc gtt acc      6624
Gly Lys  Pro Glu Lys Leu Ala  Asp Gly Arg Arg Val  Arg Val Thr
    2195                2200                2205

gtg gat  gtc ttc tgc aaa gga  acc ttc cgt ggc atc  gga cgc aac      6669
Val Asp  Val Phe Cys Lys Gly  Thr Phe Arg Gly Ile  Gly Arg Asn
    2210                2215                2220

tat cgc  att gcc aag tgc acg  gcg gcc aaa tgc gca  ttg cgc caa      6714
Tyr Arg  Ile Ala Lys Cys Thr  Ala Ala Lys Cys Ala  Leu Arg Gln
    2225                2230                2235

ctc aaa  aag cag ggc ttg ata  gcc aaa aaa gac taa                  6750
Leu Lys  Lys Gln Gly Leu Ile  Ala Lys Lys Asp
    2240                2245
```

<210> SEQ ID NO 4
<211> LENGTH: 2249
<212> TYPE: PRT
<213> ORGANISM: Drosophila melanogaster

<400> SEQUENCE: 4

```
Met Ala Phe His Trp Cys Asp Asn Asn Leu His Thr Thr Val Phe Thr
1               5                   10                  15

Pro Arg Asp Phe Gln Val Glu Leu Leu Ala Thr Ala Tyr Glu Arg Asn
            20                  25                  30

Thr Ile Ile Cys Leu Gly His Arg Ser Ser Lys Glu Phe Ile Ala Leu
        35                  40                  45

Lys Leu Leu Gln Glu Leu Ser Arg Arg Ala Arg Arg His Gly Arg Val
    50                  55                  60

Ser Val Tyr Leu Ser Cys Glu Val Gly Thr Ser Thr Glu Pro Cys Ser
65                  70                  75                  80

Ile Tyr Thr Met Leu Thr His Leu Thr Asp Leu Arg Val Trp Gln Glu
                85                  90                  95

Gln Pro Asp Met Gln Ile Pro Phe Asp His Cys Trp Thr Asp Tyr His
            100                 105                 110

Val Ser Ile Leu Arg Pro Glu Gly Phe Leu Tyr Leu Leu Glu Thr Arg
        115                 120                 125

Glu Leu Leu Leu Ser Ser Val Glu Leu Ile Val Leu Glu Asp Cys His
    130                 135                 140

Asp Ser Ala Val Tyr Gln Arg Ile Arg Pro Leu Phe Glu Asn His Ile
145                 150                 155                 160

Met Pro Ala Pro Pro Ala Asp Arg Pro Arg Ile Leu Gly Leu Ala Gly
                165                 170                 175
```

-continued

```
Pro Leu His Ser Ala Gly Cys Glu Leu Gln Gln Leu Ser Ala Met Leu
            180                 185                 190

Ala Thr Leu Glu Gln Ser Val Leu Cys Gln Ile Glu Thr Ala Ser Asp
        195                 200                 205

Ile Val Thr Val Leu Arg Tyr Cys Ser Arg Pro His Glu Tyr Ile Val
    210                 215                 220

Gln Cys Ala Pro Phe Glu Met Asp Glu Leu Ser Leu Val Leu Ala Asp
225                 230                 235                 240

Val Leu Asn Thr His Lys Ser Phe Leu Leu Asp His Arg Tyr Asp Pro
                245                 250                 255

Tyr Glu Ile Tyr Gly Thr Asp Gln Phe Met Asp Glu Leu Lys Asp Ile
            260                 265                 270

Pro Asp Pro Lys Val Asp Pro Leu Asn Val Ile Asn Ser Leu Leu Val
        275                 280                 285

Val Leu His Glu Met Gly Pro Trp Cys Thr Gln Arg Ala Ala His His
    290                 295                 300

Phe Tyr Gln Cys Asn Glu Lys Leu Lys Val Lys Thr Pro His Glu Arg
305                 310                 315                 320

His Tyr Leu Leu Tyr Cys Leu Val Ser Thr Ala Leu Ile Gln Leu Tyr
                325                 330                 335

Ser Leu Cys Glu His Ala Phe His Arg His Leu Gly Ser Gly Ser Asp
            340                 345                 350

Ser Arg Gln Thr Ile Glu Arg Tyr Ser Ser Pro Lys Val Arg Arg Leu
        355                 360                 365

Leu Gln Thr Leu Arg Cys Phe Lys Pro Glu Glu Val His Thr Gln Ala
    370                 375                 380

Asp Gly Leu Arg Arg Met Arg His Gln Val Asp Gln Ala Asp Phe Asn
385                 390                 395                 400

Arg Leu Ser His Thr Leu Glu Ser Lys Cys Arg Met Val Asp Gln Met
                405                 410                 415

Asp Gln Pro Pro Thr Glu Thr Arg Ala Leu Val Ala Thr Leu Glu Gln
            420                 425                 430

Ile Leu His Thr Thr Glu Asp Arg Gln Thr Asn Arg Ser Ala Ala Arg
        435                 440                 445

Val Thr Pro Thr Pro Thr Pro Ala His Ala Lys Pro Lys Pro Ser Ser
    450                 455                 460

Gly Ala Asn Thr Ala Gln Pro Arg Thr Arg Arg Arg Val Tyr Thr Arg
465                 470                 475                 480

Arg His His Arg Asp His Asn Asp Gly Ser Asp Thr Leu Cys Ala Leu
                485                 490                 495

Ile Tyr Cys Asn Gln Asn His Thr Ala Arg Val Leu Phe Glu Leu Leu
            500                 505                 510

Ala Glu Ile Ser Arg Arg Asp Pro Asp Leu Lys Phe Leu Arg Cys Gln
        515                 520                 525

Tyr Thr Thr Asp Arg Val Ala Asp Pro Thr Thr Glu Pro Lys Glu Ala
    530                 535                 540

Glu Leu Glu His Arg Arg Gln Glu Glu Val Leu Lys Arg Phe Arg Met
545                 550                 555                 560

His Asp Cys Asn Val Leu Ile Gly Thr Ser Val Leu Glu Glu Gly Ile
                565                 570                 575

Asp Val Pro Lys Cys Asn Leu Val Val Arg Trp Asp Pro Pro Thr Thr
```

-continued

```
           580                 585                 590

Tyr Arg Ser Tyr Val Gln Cys Lys Gly Arg Ala Arg Ala Ala Pro Ala
        595                 600                 605

Tyr His Val Ile Leu Val Ala Pro Ser Tyr Lys Ser Pro Thr Val Gly
    610                 615                 620

Ser Val Gln Leu Thr Asp Arg Ser His Arg Tyr Ile Cys Ala Thr Gly
625                 630                 635                 640

Asp Thr Thr Glu Ala Asp Ser Asp Ser Asp Asp Ser Ala Met Pro Asn
                645                 650                 655

Ser Ser Gly Ser Asp Pro Tyr Thr Phe Gly Thr Ala Arg Gly Thr Val
            660                 665                 670

Lys Ile Leu Asn Pro Glu Val Phe Ser Lys Gln Pro Pro Thr Ala Cys
        675                 680                 685

Asp Ile Lys Leu Gln Glu Ile Gln Asp Glu Leu Pro Ala Ala Ala Gln
    690                 695                 700

Leu Asp Thr Ser Asn Ser Ser Asp Glu Ala Val Ser Met Ser Asn Thr
705                 710                 715                 720

Ser Pro Ser Glu Ser Ser Thr Glu Gln Lys Ser Arg Arg Phe Gln Cys
                725                 730                 735

Glu Leu Ser Ser Leu Thr Glu Pro Glu Asp Thr Ser Asp Thr Thr Ala
            740                 745                 750

Glu Ile Asp Thr Ala His Ser Leu Ala Ser Thr Thr Lys Asp Leu Val
        755                 760                 765

His Gln Met Ala Gln Tyr Arg Glu Ile Glu Gln Met Leu Leu Ser Lys
    770                 775                 780

Cys Ala Asn Thr Glu Pro Pro Glu Gln Glu Gln Ser Glu Ala Glu Arg
785                 790                 795                 800

Phe Ser Ala Cys Leu Ala Ala Tyr Arg Pro Lys Pro His Leu Leu Thr
                805                 810                 815

Gly Ala Ser Val Asp Leu Gly Ser Ala Ile Ala Leu Val Asn Lys Tyr
            820                 825                 830

Cys Ala Arg Leu Pro Ser Asp Thr Phe Thr Lys Leu Thr Ala Leu Trp
        835                 840                 845

Arg Cys Thr Arg Asn Glu Arg Ala Gly Val Thr Leu Phe Gln Tyr Thr
    850                 855                 860

Leu Arg Leu Pro Ile Asn Ser Pro Leu Lys His Asp Ile Val Gly Leu
865                 870                 875                 880

Pro Met Pro Thr Gln Thr Leu Ala Arg Arg Leu Ala Ala Leu Gln Ala
                885                 890                 895

Cys Val Glu Leu His Arg Ile Gly Glu Leu Asp Asp Gln Leu Gln Pro
            900                 905                 910

Ile Gly Lys Glu Gly Phe Arg Ala Leu Glu Pro Asp Trp Glu Cys Phe
        915                 920                 925

Glu Leu Glu Pro Glu Asp Glu Gln Ile Val Gln Leu Ser Asp Glu Pro
    930                 935                 940

Arg Pro Gly Thr Thr Lys Arg Arg Gln Tyr Tyr Tyr Lys Arg Ile Ala
945                 950                 955                 960

Ser Glu Phe Cys Asp Cys Arg Pro Val Ala Gly Ala Pro Cys Tyr Leu
                965                 970                 975

Tyr Phe Ile Gln Leu Thr Leu Gln Cys Pro Ile Pro Glu Glu Gln Asn
            980                 985                 990
```

-continued

```
Thr Arg Gly Arg Lys Ile Tyr Pro  Pro Glu Asp Ala Gln  Gln Gly Phe
        995                 1000                1005

Gly Ile  Leu Thr Thr Lys Arg  Ile Pro Lys Leu Ser  Ala Phe Ser
    1010                1015                1020

Ile Phe  Thr Arg Ser Gly Glu  Val Lys Val Ser Leu  Glu Leu Ala
    1025                1030                1035

Lys Glu  Arg Val Ile Leu Thr  Ser Glu Gln Ile Val  Cys Ile Asn
    1040                1045                1050

Gly Phe  Leu Asn Tyr Thr Phe  Thr Asn Val Leu Arg  Leu Gln Lys
    1055                1060                1065

Phe Leu  Met Leu Phe Asp Pro  Asp Ser Thr Glu Asn  Cys Val Phe
    1070                1075                1080

Ile Val  Pro Thr Val Lys Ala  Pro Ala Gly Gly Lys  His Ile Asp
    1085                1090                1095

Trp Gln  Phe Leu Glu Leu Ile  Gln Ala Asn Gly Asn  Thr Met Pro
    1100                1105                1110

Arg Ala  Val Pro Asp Glu Glu  Arg Gln Ala Gln Pro  Phe Asp Pro
    1115                1120                1125

Gln Arg  Phe Gln Asp Ala Val  Val Met Pro Trp Tyr  Arg Asn Gln
    1130                1135                1140

Asp Gln  Pro Gln Tyr Phe Tyr  Val Ala Glu Ile Cys  Pro His Leu
    1145                1150                1155

Ser Pro  Leu Ser Cys Phe Pro  Gly Asp Asn Tyr Arg  Thr Phe Lys
    1160                1165                1170

His Tyr  Tyr Leu Val Lys Tyr  Gly Leu Thr Ile Gln  Asn Thr Ser
    1175                1180                1185

Gln Pro  Leu Leu Asp Val Asp  His Thr Ser Ala Arg  Leu Asn Phe
    1190                1195                1200

Leu Thr  Pro Arg Tyr Val Asn  Arg Lys Gly Val Ala  Leu Pro Thr
    1205                1210                1215

Ser Ser  Glu Glu Thr Lys Arg  Ala Lys Arg Glu Asn  Leu Glu Gln
    1220                1225                1230

Lys Gln  Ile Leu Val Pro Glu  Leu Cys Thr Val His  Pro Phe Pro
    1235                1240                1245

Ala Ser  Leu Trp Arg Thr Ala  Val Cys Leu Pro Cys  Ile Leu Tyr
    1250                1255                1260

Arg Ile  Asn Gly Leu Leu Leu  Ala Asp Asp Ile Arg  Lys Gln Val
    1265                1270                1275

Ser Ala  Asp Leu Gly Leu Gly  Arg Gln Gln Ile Glu  Asp Glu Asp
    1280                1285                1290

Phe Glu  Trp Pro Met Leu Asp  Phe Gly Trp Ser Leu  Ser Glu Val
    1295                1300                1305

Leu Lys  Lys Ser Arg Glu Ser  Lys Gln Lys Glu Ser  Leu Lys Asp
    1310                1315                1320

Asp Thr  Ile Asn Gly Lys Asp  Leu Ala Asp Val Glu  Lys Lys Pro
    1325                1330                1335

Thr Ser  Glu Glu Thr Gln Leu  Asp Lys Asp Ser Lys  Asp Asp Lys
    1340                1345                1350

Val Glu  Lys Ser Ala Ile Glu  Leu Ile Ile Glu Gly  Glu Glu Lys
    1355                1360                1365
```

-continued

```
Leu Gln  Glu Ala Asp Asp Phe  Ile Glu Ile Gly Thr  Trp Ser Asn
    1370                1375                1380

Asp Met  Ala Asp Asp Ile Ala  Ser Phe Asn Gln Glu  Asp Asp Asp
    1385                1390                1395

Glu Asp  Asp Ala Phe His Leu  Pro Val Leu Pro Ala  Asn Val Lys
    1400                1405                1410

Phe Cys  Asp Gln Gln Thr Arg  Tyr Gly Ser Pro Thr  Phe Trp Asp
    1415                1420                1425

Val Ser  Asn Gly Glu Ser Gly  Phe Lys Gly Pro Lys  Ser Ser Gln
    1430                1435                1440

Asn Lys  Gln Gly Gly Lys Gly  Lys Ala Lys Gly Pro  Ala Lys Pro
    1445                1450                1455

Thr Phe  Asn Tyr Tyr Asp Ser  Asp Asn Ser Leu Gly  Ser Ser Tyr
    1460                1465                1470

Asp Asp  Asp Asp Asn Ala Gly  Pro Leu Asn Tyr Met  His His Asn
    1475                1480                1485

Tyr Ser  Ser Asp Asp Asp Asp  Val Ala Asp Asp Ile  Asp Ala Gly
    1490                1495                1500

Arg Ile  Ala Phe Thr Ser Lys  Asn Glu Ala Glu Thr  Ile Glu Thr
    1505                1510                1515

Ala Gln  Glu Val Glu Lys Arg  Gln Lys Gln Leu Ser  Ile Ile Gln
    1520                1525                1530

Ala Thr  Asn Ala Asn Glu Arg  Gln Tyr Gln Gln Thr  Lys Asn Leu
    1535                1540                1545

Leu Ile  Gly Phe Asn Phe Lys  His Glu Asp Gln Lys  Glu Pro Ala
    1550                1555                1560

Thr Ile  Arg Tyr Glu Glu Ser  Ile Ala Lys Leu Lys  Thr Glu Ile
    1565                1570                1575

Glu Ser  Gly Gly Met Leu Val  Pro His Asp Gln Gln  Leu Val Leu
    1580                1585                1590

Lys Arg  Ser Asp Ala Ala Glu  Ala Gln Val Ala Lys  Val Ser Met
    1595                1600                1605

Met Glu  Leu Leu Lys Gln Leu  Leu Pro Tyr Val Asn  Glu Asp Val
    1610                1615                1620

Leu Ala  Lys Lys Leu Gly Asp  Arg Arg Glu Leu Leu  Leu Ser Asp
    1625                1630                1635

Leu Val  Glu Leu Asn Ala Asp  Trp Val Ala Arg His  Glu Gln Glu
    1640                1645                1650

Thr Tyr  Asn Val Met Gly Cys  Gly Asp Ser Phe Asp  Asn Tyr Asn
    1655                1660                1665

Asp His  His Arg Leu Asn Leu  Asp Glu Lys Gln Leu  Lys Leu Gln
    1670                1675                1680

Tyr Glu  Arg Ile Glu Ile Glu  Pro Pro Thr Ser Thr  Lys Ala Ile
    1685                1690                1695

Thr Ser  Ala Ile Leu Pro Ala  Gly Phe Ser Phe Asp  Arg Gln Pro
    1700                1705                1710

Asp Leu  Val Gly His Pro Gly  Pro Ser Pro Ser Ile  Ile Leu Gln
    1715                1720                1725

Ala Leu  Thr Met Ser Asn Ala  Asn Asp Gly Ile Asn  Leu Glu Arg
    1730                1735                1740

Leu Glu  Thr Ile Gly Asp Ser  Phe Leu Lys Tyr Ala  Ile Thr Thr
```

-continued

```
    1745                1750                1755

Tyr Leu  Tyr Ile Thr Tyr Glu  Asn Val His Glu Gly  Lys Leu Ser
    1760                1765                1770

His Leu  Arg Ser Lys Gln Val  Ala Asn Leu Asn Leu  Tyr Arg Leu
    1775                1780                1785

Gly Arg  Arg Lys Arg Leu Gly  Glu Tyr Met Ile Ala  Thr Lys Phe
    1790                1795                1800

Glu Pro  His Asp Asn Trp Leu  Pro Pro Cys Tyr Tyr  Val Pro Lys
    1805                1810                1815

Glu Leu  Glu Lys Ala Leu Ile  Glu Ala Lys Ile Pro  Thr His His
    1820                1825                1830

Trp Lys  Leu Ala Asp Leu Leu  Asp Ile Lys Asn Leu  Ser Ser Val
    1835                1840                1845

Gln Ile  Cys Glu Met Val Arg  Glu Lys Ala Asp Ala  Leu Gly Leu
    1850                1855                1860

Glu Gln  Asn Gly Gly Ala Gln  Asn Gly Gln Leu Asp  Asp Ser Asn
    1865                1870                1875

Asp Ser  Cys Asn Asp Phe Ser  Cys Phe Ile Pro Tyr  Asn Leu Val
    1880                1885                1890

Ser Gln  His Ser Ile Pro Asp  Lys Ser Ile Ala Asp  Cys Val Glu
    1895                1900                1905

Ala Leu  Ile Gly Ala Tyr Leu  Ile Glu Cys Gly Pro  Arg Gly Ala
    1910                1915                1920

Leu Leu  Phe Met Ala Trp Leu  Gly Val Arg Val Leu  Pro Ile Thr
    1925                1930                1935

Arg Gln  Leu Asp Gly Gly Asn  Gln Glu Gln Arg Ile  Pro Gly Ser
    1940                1945                1950

Thr Lys  Pro Asn Ala Glu Asn  Val Val Thr Val Tyr  Gly Ala Trp
    1955                1960                1965

Pro Thr  Pro Arg Ser Pro Leu  Leu His Phe Ala Pro  Asn Ala Thr
    1970                1975                1980

Glu Glu  Leu Asp Gln Leu Leu  Ser Gly Phe Glu Glu  Phe Glu Glu
    1985                1990                1995

Ser Leu  Gly Tyr Lys Phe Arg  Asp Arg Ser Tyr Leu  Leu Gln Ala
    2000                2005                2010

Met Thr  His Ala Ser Tyr Thr  Pro Asn Arg Leu Thr  Asp Cys Tyr
    2015                2020                2025

Gln Arg  Leu Glu Phe Leu Gly  Asp Ala Val Leu Asp  Tyr Leu Ile
    2030                2035                2040

Thr Arg  His Leu Tyr Glu Asp  Pro Arg Gln His Ser  Pro Gly Ala
    2045                2050                2055

Leu Thr  Asp Leu Arg Ser Ala  Leu Val Asn Asn Thr  Ile Phe Ala
    2060                2065                2070

Ser Leu  Ala Val Arg His Gly  Phe His Lys Phe Phe  Arg His Leu
    2075                2080                2085

Ser Pro  Gly Leu Asn Asp Val  Ile Asp Arg Phe Val  Arg Ile Gln
    2090                2095                2100

Gln Glu  Asn Gly His Cys Ile  Ser Glu Glu Tyr Tyr  Leu Leu Ser
    2105                2110                2115

Glu Glu  Glu Cys Asp Asp Ala  Glu Asp Val Glu Val  Pro Lys Ala
    2120                2125                2130
```

-continued

```
Leu Gly  Asp Val Phe Glu Ser  Ile Ala Gly Ala Ile  Phe Leu Asp
    2135                2140                2145

Ser Asn  Met Ser Leu Asp Val  Val Trp His Val Tyr  Ser Asn Met
    2150                2155                2160

Met Ser  Pro Glu Ile Glu Gln  Phe Ser Asn Ser Val  Pro Lys Ser
    2165                2170                2175

Pro Ile  Arg Glu Leu Leu Glu  Leu Glu Pro Glu Thr  Ala Lys Phe
    2180                2185                2190

Gly Lys  Pro Glu Lys Leu Ala  Asp Gly Arg Arg Val  Arg Val Thr
    2195                2200                2205

Val Asp  Val Phe Cys Lys Gly  Thr Phe Arg Gly Ile  Gly Arg Asn
    2210                2215                2220

Tyr Arg  Ile Ala Lys Cys Thr  Ala Ala Lys Cys Ala  Leu Arg Gln
    2225                2230                2235

Leu Lys  Lys Gln Gly Leu Ile  Ala Lys Lys Asp
    2240                2245


<210> SEQ ID NO 5
<211> LENGTH: 1145
<212> TYPE: PRT
<213> ORGANISM: Drosophila melanogaster

<400> SEQUENCE: 5

Met Gly Lys Lys Asp Lys Asn Lys Lys Gly Gly Gln Asp Ser Ala Ala
1               5                   10                  15

Ala Pro Gln Pro Gln Gln Gln Gln Lys Gln Gln Gln Gln Arg Gln Gln
            20                  25                  30

Gln Pro Gln Gln Leu Gln Gln Pro Gln Gln Leu Gln Gln Pro Gln Gln
        35                  40                  45

Leu Gln Gln Pro Gln Gln Gln Gln Gln Gln Gln Pro His Gln Gln Gln
    50                  55                  60

Gln Gln Ser Ser Arg Gln Gln Pro Ser Thr Ser Ser Gly Gly Ser Arg
65                  70                  75                  80

Ala Ser Gly Phe Gln Gln Gly Gly Gln Gln Gln Lys Ser Gln Asp Ala
                85                  90                  95

Glu Gly Trp Thr Ala Gln Lys Lys Gln Gly Lys Gln Gln Val Gln Gly
            100                 105                 110

Trp Thr Lys Gln Gly Gln Gln Gly Gly His Gln Gln Gly Arg Gln Gly
        115                 120                 125

Gln Asp Gly Gly Tyr Gln Gln Arg Pro Pro Gly Gln Gln Gln Gly Gly
    130                 135                 140

His Gln Gln Gly Arg Gln Gly Gln Glu Gly Gly Tyr Gln Gln Arg Pro
145                 150                 155                 160

Pro Gly Gln Gln Gln Gly Gly His Gln Gln Gly Arg Gln Gly Gln Glu
                165                 170                 175

Gly Gly Tyr Gln Gln Arg Pro Ser Gly Gln Gln Gln Gly Gly His Gln
            180                 185                 190

Gln Gly Arg Gln Gly Gln Glu Gly Gly Tyr Gln Gln Arg Pro Pro Gly
        195                 200                 205

Gln Gln Gln Gly Gly His Gln Gln Gly Arg Gln Gly Gln Glu Gly Gly
    210                 215                 220

Tyr Gln Gln Arg Pro Ser Gly Gln Gln Gln Gly Gly His Gln Gln Gly
```

-continued

```
225                 230                 235                 240

Arg Gln Gly Gln Glu Gly Gly Tyr Gln Gln Arg Pro Ser Gly Gln Gln
                245                 250                 255

Gln Gly Gly His Gln Gln Gly Arg Gln Gly Gln Glu Gly Gly Tyr Gln
            260                 265                 270

Gln Arg Pro Ser Gly Gln Gln Gln Gly Gly His Gln Gln Gly Arg Gln
        275                 280                 285

Gly Gln Glu Gly Gly Tyr Gln Gln Arg Pro Pro Gly Gln Gln Pro Asn
    290                 295                 300

Gln Thr Gln Ser Gln Gly Gln Tyr Gln Ser Arg Gly Pro Pro Gln Gln
305                 310                 315                 320

Gln Gln Ala Ala Pro Leu Pro Leu Pro Pro Gln Pro Ala Gly Ser Ile
                325                 330                 335

Lys Arg Gly Thr Ile Gly Lys Pro Gly Gln Val Gly Ile Asn Tyr Leu
            340                 345                 350

Asp Leu Asp Leu Ser Lys Met Pro Ser Val Ala Tyr His Tyr Asp Val
        355                 360                 365

Lys Ile Met Pro Glu Arg Pro Lys Lys Phe Tyr Arg Gln Ala Phe Glu
    370                 375                 380

Gln Phe Arg Val Asp Gln Leu Gly Gly Ala Val Leu Ala Tyr Asp Gly
385                 390                 395                 400

Lys Ala Ser Cys Tyr Ser Val Asp Lys Leu Pro Leu Asn Ser Gln Asn
                405                 410                 415

Pro Glu Val Thr Val Thr Asp Arg Asn Gly Arg Thr Leu Arg Tyr Thr
            420                 425                 430

Ile Glu Ile Lys Glu Thr Gly Asp Ser Thr Ile Asp Leu Lys Ser Leu
        435                 440                 445

Thr Thr Tyr Met Asn Asp Arg Ile Phe Asp Lys Pro Met Arg Ala Met
    450                 455                 460

Gln Cys Val Glu Val Val Leu Ala Ser Pro Cys His Asn Lys Ala Ile
465                 470                 475                 480

Arg Val Gly Arg Ser Phe Phe Lys Met Ser Asp Pro Asn Asn Arg His
                485                 490                 495

Glu Leu Asp Asp Gly Tyr Glu Ala Leu Val Gly Leu Tyr Gln Ala Phe
            500                 505                 510

Met Leu Gly Asp Arg Pro Phe Leu Asn Val Asp Ile Ser His Lys Ser
        515                 520                 525

Phe Pro Ile Ser Met Pro Met Ile Glu Tyr Leu Glu Arg Phe Ser Leu
    530                 535                 540

Lys Ala Lys Ile Asn Asn Thr Thr Asn Leu Asp Tyr Ser Arg Arg Phe
545                 550                 555                 560

Leu Glu Pro Phe Leu Arg Gly Ile Asn Val Val Tyr Thr Pro Pro Gln
                565                 570                 575

Ser Phe Gln Ser Ala Pro Arg Val Tyr Arg Val Asn Gly Leu Ser Arg
            580                 585                 590

Ala Pro Ala Ser Ser Glu Thr Phe Glu His Asp Gly Lys Lys Val Thr
        595                 600                 605

Ile Ala Ser Tyr Phe His Ser Arg Asn Tyr Pro Leu Lys Phe Pro Gln
    610                 615                 620

Leu His Cys Leu Asn Val Gly Ser Ser Ile Lys Ser Ile Leu Leu Pro
625                 630                 635                 640
```

-continued

```
Ile Glu Leu Cys Ser Ile Glu Glu Gly Gln Ala Leu Asn Arg Lys Asp
                645                 650                 655

Gly Ala Thr Gln Val Ala Asn Met Ile Lys Tyr Ala Ala Thr Ser Thr
            660                 665                 670

Asn Val Arg Lys Arg Lys Ile Met Asn Leu Leu Gln Tyr Phe Gln His
        675                 680                 685

Asn Leu Asp Pro Thr Ile Ser Arg Phe Gly Ile Arg Ile Ala Asn Asp
    690                 695                 700

Phe Ile Val Val Ser Thr Arg Val Leu Ser Pro Pro Gln Val Glu Tyr
705                 710                 715                 720

His Ser Lys Arg Phe Thr Met Val Lys Asn Gly Ser Trp Arg Met Asp
                725                 730                 735

Gly Met Lys Phe Leu Glu Pro Lys Pro Lys Ala His Lys Cys Ala Val
            740                 745                 750

Leu Tyr Cys Asp Pro Arg Ser Gly Arg Lys Met Asn Tyr Thr Gln Leu
        755                 760                 765

Asn Asp Phe Gly Asn Leu Ile Ile Ser Gln Gly Lys Ala Val Asn Ile
    770                 775                 780

Ser Leu Asp Ser Asp Val Thr Tyr Arg Pro Phe Thr Asp Asp Glu Arg
785                 790                 795                 800

Ser Leu Asp Thr Ile Phe Ala Asp Leu Lys Arg Ser Gln His Asp Leu
                805                 810                 815

Ala Ile Val Ile Ile Pro Gln Phe Arg Ile Ser Tyr Asp Thr Ile Lys
            820                 825                 830

Gln Lys Ala Glu Leu Gln His Gly Ile Leu Thr Gln Cys Ile Lys Gln
        835                 840                 845

Phe Thr Val Glu Arg Lys Cys Asn Asn Gln Thr Ile Gly Asn Ile Leu
    850                 855                 860

Leu Lys Ile Asn Ser Lys Leu Asn Gly Ile Asn His Lys Ile Lys Asp
865                 870                 875                 880

Asp Pro Arg Leu Pro Met Met Lys Asn Thr Met Tyr Ile Gly Ala Asp
                885                 890                 895

Val Thr His Pro Ser Pro Asp Gln Arg Glu Ile Pro Ser Val Val Gly
            900                 905                 910

Val Ala Ala Ser His Asp Pro Tyr Gly Ala Ser Tyr Asn Met Gln Tyr
        915                 920                 925

Arg Leu Gln Arg Gly Ala Leu Glu Glu Ile Glu Asp Met Phe Ser Ile
    930                 935                 940

Thr Leu Glu His Leu Arg Val Tyr Lys Glu Tyr Arg Asn Ala Tyr Pro
945                 950                 955                 960

Asp His Ile Ile Tyr Tyr Arg Asp Gly Val Ser Asp Gly Gln Phe Pro
                965                 970                 975

Lys Ile Lys Asn Glu Glu Leu Arg Cys Ile Lys Gln Ala Cys Asp Lys
            980                 985                 990

Val Gly Cys Lys Pro Lys Ile Cys  Cys Val Ile Val Val  Lys Arg His
        995                 1000                1005

His Thr  Arg Phe Phe Pro Ser  Gly Asp Val Thr Thr  Ser Asn Lys
    1010                1015                1020

Phe Asn  Asn Val Asp Pro Gly  Thr Val Val Asp Arg  Thr Ile Val
    1025                1030                1035
```

-continued

```
His Pro  Asn Glu Met Gln Phe  Phe Met Val Ser Gly  Gln Ala Ile
    1040                1045                1050

Gln Gly  Thr Ala Lys Pro Thr  Arg Tyr Asn Val Ile  Glu Asn Thr
    1055                1060                1065

Gly Asn  Leu Asp Ile Asp Leu  Leu Gln Gln Leu Thr  Tyr Asn Leu
    1070                1075                1080

Cys His  Met Phe Pro Arg Cys  Asn Arg Ser Val Ser  Tyr Pro Ala
    1085                1090                1095

Pro Ala  Tyr Leu Ala His Leu  Val Ala Ala Arg Gly  Arg Val Tyr
    1100                1105                1110

Leu Thr  Gly Thr Asn Arg Phe  Leu Asp Leu Lys Lys  Glu Tyr Ala
    1115                1120                1125

Lys Arg  Thr Ile Val Pro Glu  Phe Met Lys Lys Asn  Pro Met Tyr
    1130                1135                1140

Phe Val
    1145
```

<210> SEQ ID NO 6
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: photinus pyralis

<400> SEQUENCE: 6

```
ucgaaguacu cagcguaagu ga                                              22
```

<210> SEQ ID NO 7
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: photinus pyralis

<400> SEQUENCE: 7

```
uaaagcuuca ugagucgcau uc                                              22
```

<210> SEQ ID NO 8
<211> LENGTH: 83
<212> TYPE: RNA
<213> ORGANISM: photinus pyralis

<400> SEQUENCE: 8

```
caucgacuga aaucccuccc ugguaauccg uuugggcucu gcccugcuau gggauaaaug     60

guuaucaggg auuuuagucg auc                                             83
```

<210> SEQ ID NO 9
<211> LENGTH: 62
<212> TYPE: RNA
<213> ORGANISM: photinus pyralis

<400> SEQUENCE: 9

```
caucgacuga aaucccuggu aauccguugu uaacaacgga uuaccaggga uuucagucga     60

ug                                                                    62
```

<210> SEQ ID NO 10
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

-continued

<400> SEQUENCE: 10

taatacgact cactataggg caaggtcata gttcctgcca gctg 44

<210> SEQ ID NO 11
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 11

taatacgact cactataggg cagatatttt cttaccaccc accc 44

<210> SEQ ID NO 12
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 12

taatacgact cactataggg ttaagtttaa caggagaagc tgga 44

<210> SEQ ID NO 13
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 13

taatacgact cactataggg aaatcattgc tttcctgata atgc 44

<210> SEQ ID NO 14
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 14

taatacgact cactataggg tagatttccg cagccccagt gttc 44

<210> SEQ ID NO 15
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 15

taatacgact cactataggg gttgcctctc atttttcctt gatt 44

<210> SEQ ID NO 16
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 16

taatacgact cactataggg tattttagac tgattacttt tataa 45

-continued

<210> SEQ ID NO 17
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 17

taatacgact cactataggg tcacatgttt tggctaagac ctat 44

<210> SEQ ID NO 18
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Dicer siRNA

<400> SEQUENCE: 18

tcaaccagcc actgctgga 19

<210> SEQ ID NO 19
<211> LENGTH: 71
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide Primer

<400> SEQUENCE: 19

tccaattcag cgggagccac ctgatgaagc ttgatcgggt ggctctcgct gagttggaat 60

ccatttttt t 71

<210> SEQ ID NO 20
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: photinus pyralis

<400> SEQUENCE: 20

ucgaaguacu cagcguaagu g 21

<210> SEQ ID NO 21
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 21

aaagcuucau gagucgcauu c 21

<210> SEQ ID NO 22
<211> LENGTH: 81
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 22

caucgacuga aaucccuggu aauccguuug gggcucugcc cugcuauggg auaaauggau 60

uaucagggau uuuagucgau g 81

<210> SEQ ID NO 23
<211> LENGTH: 82
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 23

-continued

caucgacuga aaucccgccg uaauccguuu ggggcucugc ccugcuaugg gauaaaugga 60

uuaucaggga uuuuagucga ug 82

<210> SEQ ID NO 24
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 24

ggucgaagua cucagcguaa gaa 23

<210> SEQ ID NO 25
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 25

aaagcuucat gagucgcauu cgg 23

<210> SEQ ID NO 26
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 26

gguuguggau cuggauaccg g 21

<210> SEQ ID NO 27
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 27

uuccaacacc uagaccuaug g 21

<210> SEQ ID NO 28
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 28

ggugccaacc cuauucuccu u 21

<210> SEQ ID NO 29
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 29

gaccacgguu gggauaagag g 21

<210> SEQ ID NO 30
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 30

ggcuaugaag agaguacgcc cu 22

<210> SEQ ID NO 31

-continued

<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 31

uuccgauacu ucucucaugc gg 22

<210> SEQ ID NO 32
<211> LENGTH: 64
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 32

ggucgaagua cucagcguaa gugaugucca cuuaaguggg uguuguuugu guuggguguu 60

uugg 64

<210> SEQ ID NO 33
<211> LENGTH: 60
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 33

ggucgaagua cucagcguaa gugauguccu uaaggguguu guuuguguug gguguuuugg 60

<210> SEQ ID NO 34
<211> LENGTH: 56
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 34

gggcgaagua cucagcguaa gugauguuua aguguuguuu guguugggug uuuugg 56

<210> SEQ ID NO 35
<211> LENGTH: 50
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 35

ggucgaagua cucagcguaa gugauuaauu guuuguguug gguguuuugg 50

<210> SEQ ID NO 36
<211> LENGTH: 69
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 36

ggcucgaguc gaaguacuca gcguaaguga uguccauuaa guggguguug uuuguguugg 60

guguuuugg 69

<210> SEQ ID NO 37
<211> LENGTH: 70
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 37

ggucgaagua cucagcguaa gugaugucca cuuaaguggg uguuguuugu guuggguguu 60

uuggcucgag 70

<210> SEQ ID NO 38

-continued

<211> LENGTH: 69
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 38

ggauuccaau ucagcgggag ccaccugaug aagcuugauc ggguggcucu cgcugaguug 60

gaauccauu 69

<210> SEQ ID NO 39
<211> LENGTH: 64
<212> TYPE: RNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 39

ggucuaagug gagcccuucg aguguuagaa gcuugugaca cucggagggc uucacuuggg 60

ccuu 64

<210> SEQ ID NO 40
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 40

cuuacgcuga guacuucgat t 21

<210> SEQ ID NO 41
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 41

ucgaaguacu cagcguaagt t 21

<210> SEQ ID NO 42
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 42

agcuucauaa ggcgcaugct t 21

<210> SEQ ID NO 43
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 43

gcaugcgccu uaugaagcut t 21

<210> SEQ ID NO 44
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 44

cugugagauc uacggagccu gtt 23

<210> SEQ ID NO 45
<211> LENGTH: 23
<212> TYPE: DNA

-continued

<213> ORGANISM: Photinus Pyralis

<400> SEQUENCE: 45

caggcuccgu agaucucaca gtt 23

We claim:

1. A method for attenuating expression of a target gene in host cells, comprising introducing double stranded RNA (dsRNA) into the host cells in an amount sufficient to attenuate expression of the target gene, wherein the dsRNA comprises a nucleotide sequence that hybridizes under stringent conditions to an untranslated sequence of the target gene.

2. A method for attenuating expression of a target gene in host cells, comprising introducing double stranded RNA (dsRNA) into the host cells in an amount sufficient to attenuate expression of the target gene, wherein the dsRNA comprises a nucleotide sequence that hybridizes under stringent conditions to an intronic sequence of the target gene.

3. A composition for attenuating expression of a target gene, comprising a double stranded RNA (dsRNA) having a nucleotide sequence that hybridizes under stringent conditions to an untranslated sequence of the target gene.

4. A composition for attenuating expression of a target gene, comprising an expression vector which, when expressed, produces a double stranded RNA (dsRNA) having a nucleotide sequence that hybridizes under stringent conditions to an untranslated sequence of the target gene.

5. A composition for attenuating expression of a target gene, comprising a double stranded RNA (dsRNA) having a nucleotide sequence that hybridizes under stringent conditions to an intronic sequence of the target gene.

6. A composition for attenuating expression of a target gene, comprising an expression vector which, when expressed, produces a double stranded RNA (dsRNA) having a nucleotide sequence that hybridizes under stringent conditions to an intronic sequence of the target gene.

7. The method of claim 1 or the composition of claim 3 or **4**, wherein the untranslated sequence of the target gene to which the dsRNA hybridizes is selected from the group consisting of a promoter sequence and an enhancer sequence.

8. The method of any of claims **1** and **2**, wherein the host cells are suspended in culture.

9. The method of any of claims **1** and **2**, wherein the host cells are in a whole animal.

10. The method of any of claims **1** and **2**, wherein the target gene is a genomically integrated gene of the host cells.

11. The method of any of claims **1** and **2**, wherein the target gene is a heterologous gene

12. The method of claim 11, wherein the target gene is a viral gene.

13. The method of any of claims **1** and **2**, wherein the host cells is a primate cells.

14. The method of claim 13, wherein the host cells are human cells.

15. The method of any of claims **1** and **2**, or the composition of claim 3 or **5**, wherein the dsRNA is short interfering double stranded RNA which does not produce a significant PKR-dependent response in the host cells at concentrations effective for attenuating expression of the target gene.

16. The method or the composition of claim 15, wherein the dsRNA is 15-45 basepairs in length.

17. The method or the composition of claim 15, wherein the dsRNA is 19-30 basepairs in length.

18. The method of any of claims **1** and **2**, wherein the dsRNA is produced in the cell by an expression vector having a coding sequence that is transcribed to produce one or more transcriptional products that are processed to an siRNA species by said host cells.

19. The method of any of claims **1** and **2**, or the composition of claim 3 or **5**, wherein the dsRNA is a hairpin RNA that is processed to an siRNA species by said host cells.

20. The method of any of claims **1** and **2**, wherein expression of the target gene is attenuated by at least 10 fold.

21. A pharmaceutical package comprising:

a pharmaceutically preparation of a double stranded RNA (DSRNA) that is, or gives rise to, a short interfering double stranded RNA (siRNA) in a mammalian cell and attenuates expression of a target gene, which dsRNA does not produce a significant PKR-dependent response in the mammalian cell at concentrations effective for attenuating expression of the target gene; and

label or instructions (written and/or pictorial) for administering the preparation to a patient.

22. The pharmaceutical package of claim 21, wherein the siRNA hybridizes to an untranscribed sequence or an non-coding sequence of the target gene.

23. A method of attenuating expression of a target gene in the cells of a mammal, comprising administering the mammal a composition of any of claims **3-6**.

24. The method of claim 23, wherein attenuating expression of the target gene reduces unwanted growth or differentiation of the cells.

25. A method for generating short interfering double-stranded RNA (siRNA), comprising:

(i) providing an in vitro transcription system including:

a double-stranded nucleic acid having complementary sense and antisense target sequences, which target sequence is flanked by promoters for an RNA polymerase,

the RNA polymerase,

wherein the sense and antisense target sequences are transcribed and can anneal to form an siRNA; and

(ii) isolating said siRNA from said in vitro transcription system.

26. The method of claim 25, wherein the RNA polymerase is a bacteriophage RNA polymerase.

27. The method of claim 26, wherein the RNA polymerase is selected from the group consisting of T3 polymerase, T7 polymerase and SP6 polymerase.

28. The method of claim 25, wherein the in vitro transcription system includes a variegated library of target sequences to produce a variegated library of siRNA species.

29. A method for performing a business of distributing short interfering double-stranded RNA (siRNA), comprising:

(i) accepting orders for siRNA species having a sequence designated by user;

(ii) providing an in vitro transcription system including:

a double-stranded nucleic acid having a target sequence of complementary sense and antisense sequences for said user designated siRNA species, which target sequence is flanked by promoters for an RNA polymerase, and

the RNA polymerase,

wherein the sense and antisense target sequences are transcribed and can anneal to form an siRNA;

(iii) isolating said siRNA from said in vitro transcription system; and

(iv) packaging and shipping said siRNA to said user.

30. A method for altering the MHC phenotype of a donor stem cell or the progeny thereof, comprising introducing a double stranded RNA (dsRNA) into a stem cell in an amount sufficient to attenuate expression of an MHC gene otherwise expressed by said stem cell or the progeny thereof.

31. The method of claim 30, wherein the dsRNA reduces expression of one or more human leukocyte antigens (HLA) otherwise expressed by said stem cell or the progeny thereof.

32. A culture of donor stem cells or the progeny thereof, having an altered MHC phenotype resulting from stable attenuation of expression of one or more MHC genes by introduction of double stranded RNA (dsRNA) into the donor stem cells.

33. A method of transplanting a patient, comprising:

(i) generating an ex vivo cell or tissue culture of donor stem cells, or the progeny thereof, having an altered MHC phenotype resulting from stable attenuation of expression of one or more MHC genes by introduction of double stranded RNA (dsRNA) into the donor stem cells; and

(ii) transplanting said patient with said cell or tissue culture.

34. Use of a stem cell in the manufacture of a cellular medicament for transplantation to a patient, which cellular medicament comprises stem cells or the progeny thereof which have an altered MHC phenotype resulting from stable attenuation of expression of one or more MHC genes by introduction of double stranded RNA (dsRNA) into the stem cells.

35. Use of the compositions of any of claims **3-6** in the manufacture of a medicament for attenuating expression of one or more genes in vivo.

36. A method for reducing the susceptibility of host cells to infection by pathogen, comprising introducing a double stranded RNA (dsRNA) into said host cells in an amount sufficient to attenuate expression of one or more genes necessary for expression by said pathogen.

37. The method of claim 36, wherein the pathogen is a virus, and the dsRNA attenuates expression of a cellular surface protein necessary for infection of the host cell by the virus.

38. A non-human transgenic mammal having germline and/or somatic cells comprising a transgene encoding a double stranded RNA transcriptional produce that is processed to an short interfering double-stranded RNA (siRNA) species, wherein transcription of said transgene attenuates expression of an endogenous target gene in at least one cell-type of the animal.

39. The transgenic animal of claim 38, which is chimeric for said transgene.

40. The transgenic animal of claim 38, wherein said transgene is chromosomally incorporated.

41. The transgenic animal of claim 38, wherein the transgene transcribes separate complementary transcripts that anneal to form said siRNA, which siRNA does not produce a significant PKR-dependent response at concentrations effective for attenuating expression of the target gene.

42. The transgenic animal of claim 38, wherein the transgene transcribes a hairpin RNA that is processed to said siRNA.

43. A hairpin RNA comprising a first nucleotide sequence that hybridizes under stringent conditions to a nucleotide sequence of a target gene, and a second nucleotide sequence which is a complementary inverted repeat of said first nucleotide sequence and hybridizes to said first nucleotide sequence to form a hairpin structure, which hairpin RNA attenuates expression of said target gene and does not produce a significant PKR-dependent response at concentrations effective for attenuating expression of the target gene.

44. The hairpin RNA of claim 43, wherein the hairpin RNA is chemically synthesized.

45. The hairpin RNA of claim 43, wherein the hairpin RNA is enzymatically synthesized in vitro.

46. The hairpin RNA of claim 45, wherein the hairpin RNA is synthesized by T7 RNA polymerase in vitro.

47. The hairpin RNA of claim 43, wherein the hairpin RNA is enzymatically synthesized in vivo.

48. The hairpin RNA of claim 47, wherein the hairpin RNA is synthesized by RNA polymerase III in vivo.

49. The hairpin RNA of claim 43, wherein the hairpin RNA is produced by a vector.

50. The hairpin RNA of claim 43, wherein the hairpin RNA has a size of about 20-50 nucleotides.

51. The hairpin RNA of claim 43, wherein the hairpin RNA has a size of about 50-100 nucleotides.

52. The hairpin RNA of claim 43, wherein the hairpin RNA has a size of about 100-500 nucleotides.

53. The hairpin RNA of claim 43, which includes a restriction enzyme site in the loop of the hairpin.

* * * * *