



(51) International Patent Classification:

A01N 55/08 (2006.01) A61K 31/47 (2006.01)
A61K 31/69 (2006.01)

(21) International Application Number:

PCT/US2010/030272

(22) International Filing Date:

7 April 2010 (07.04.2010)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/167,456 7 April 2009 (07.04.2009) US

(71) Applicant (for all designated States except US): **INFINITI PHARMACEUTICALS, INC.** [US/US]; 780 Memorial Drive, Cambridge, MA 02139 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CASTRO, Alfredo, C.** [US/US]; 43 Wildwood Street, Winchester, MA 01890 (US). **GROGAN, Michael, J.** [US/US]; 27 Dunster Road, Winchester, MA 01890 (US). **SNYDER, Daniel, A.** [US/US]; 94 Jaques Street, Somerville, MA 02145 (US).

(74) Agent: **ROBIDOUX, Andrea, L., C.**; Choate, Hall & Stewart LLP, Two International Place, Boston, MA 02110 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: INHIBITORS OF FATTY ACID AMIDE HYDROLASE



(57) Abstract: Provided herein are compounds of formula (I) or pharmaceutically acceptable salts, solvates or prodrugs thereof, or mixtures thereof, wherein Z^1 , Z^2 , X^1 , X^2 , X^3 , R^1 , R^2 , R^3 , m and n are defined herein. Also provided are pharmaceutically acceptable compositions that include a compound of formula I and a pharmaceutically acceptable excipient. Also provided are methods for treating FAAH-mediated disorders comprising administering to a subject in need thereof a therapeutically effective amount of a compound or composition of the present invention.

INHIBITORS OF FATTY ACID AMIDE HYDROLASE

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority to United States Provisional Patent Application serial number 61/167,456, filed April 7, 2009, the entirety of which is hereby incorporated herein by reference.

BACKGROUND


Fatty acid amide hydrolase (FAAH), also referred to as oleamide hydrolase and anandamide amidohydrolase, is an integral membrane protein that degrades fatty acid primary amides and ethanolamides, including oleamide and anandamide. FAAH degrades neuromodulating fatty acid amides at their sites of action and is intimately involved in their regulation.

FAAH has been demonstrated to be involved in a number of biological processes and its inhibition has been shown to be effective in treating a variety of conditions. For example, inhibiting FAAH has been shown to be useful in treating chronic pain, acute pain, neuropathic pain, anxiety, depression, feeding behaviors, movement disorders, glaucoma, neuroprotection and cardiovascular disease.

SUMMARY

Compounds described herein, and pharmaceutically acceptable compositions thereof, are effective inhibitors of fatty acid amide hydrolase (FAAH).

In one aspect, provided herein are compounds of formula I:



I

or pharmaceutically acceptable salts, solvates or prodrugs thereof, or mixtures thereof, wherein:

----- is selected from a single bond and a double bond;

when ----- is a single bond, X^1 is selected from CR^4R^5 and NR^6 , and X^2 is selected from CR^7R^8 and NR^9 ;

5 when ----- is a double bond, X^1 is selected from CR^4 and N, and X^2 is selected from CR^7 and N;

X^3 is selected from $CR^{10}R^{11}$ and NR^{12} ;

provided that at least one of X^1 , X^2 and X^3 is selected from N, NR^6 , NR^9 , and N^{12} ;

m is 0 or 1;

Z^1 is selected from $-OR^{13}$ and C_{1-6} alkyl;

10 Z^2 is selected from $-OR^{14}$ and C_{1-6} alkyl;

or alternatively, Z^1 and Z^2 , together with the B to which they are bound, form an 5- to 8-membered ring having at least one O atom directly attached to B, wherein the ring is comprised of carbon atoms and optionally one or more additional heteroatoms independently selected from the group consisting of N, S, and O;

15 n is 0, 1, 2 or 3;

R^1 and R^2 each independently is selected from H, halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} perhaloalkyl, -CN and $-OR^{15}$;

or alternatively, R^1 and R^2 , taken together with the carbon to which they are bound, form a carbonyl group;

20 R^3 , at each occurrence, independently is selected from halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} perhaloalkyl, -OH, C_{1-6} alkoxy and -CN;

R^4 , R^5 , R^7 , R^8 , R^{10} , and R^{11} each independently is selected from H, halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} perhaloalkyl, -CN, $-OR^{16}$, $NR^{17}R^{18}$, $-C(O)R^{19}$, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

25 R^6 , R^9 , and R^{12} each independently is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, $-C(O)R^{20}$, $-C(O)OR^{21}$, $S(O)_2R^{22}$, C_{3-10} carbocyclyl, C_{6-10} aryl, 3-10 membered heterocyclyl, 5-10 membered heteroaryl, and $-(CR^{27}R^{28})_p-R^{23}$;

R^{13} and R^{14} , at each occurrence, each independently is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl;

R^{15} and R^{16} , at each occurrence, each independently is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

R^{17} and R^{18} , at each occurrence, each independently is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, $-C(O)R^{24}$, $-C(O)OR^{25}$, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

R^{19} , R^{20} and R^{21} , at each occurrence, each independently is selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

R^{22} is selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, C_{6-10} aryl, 3-10 membered heterocyclyl, 5-10 membered heteroaryl, and $-(CR^{29}R^{30})_q-R^{26}$;

R^{23} , at each occurrence, independently is selected from C_{3-10} carbocyclyl, C_{6-10} aryl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;


R^{24} and R^{25} , at each occurrence, each independently is selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

R^{26} , at each occurrence, independently is selected from C_{3-10} carbocyclyl, C_{6-10} aryl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

R^{27} , R^{28} , R^{29} and R^{30} , at each occurrence, each independently is selected from H and C_{1-6} alkyl; and

p and q, at each occurrence, each independently is selected from 1, 2, 3, 4, 5 and 6.


In certain embodiments, the compound of the formula **Ia**:



Ia


or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

In certain embodiments, the compound is of the formula **Ib**:

**Ib**



or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

In certain embodiments, the compound is of the formula **V**:

**V**


or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

In certain embodiments, the compound is of the formulae **Va** or **Vb**:

**Va****Vb**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.



In certain embodiments, the compound is of the formula **IX**:

**IX**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.


In certain embodiments, the compound is of the formulae **IXa** or **IXb**:

5

**IXa****IXb**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.



In certain embodiments, the compound is of the formula **XI**:

**XI**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.


In certain embodiments, the compound is of the formulae **XIa** or **XIb**:

10

**XIa****XIb**



or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

In certain embodiments, the compound is of the formula **VII**:

**VII**


or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

In certain embodiments, the compound is of the formulae **VIIa** or **VIIb**:

**VIIa****VIIb**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.



In certain embodiments, the compound is of the formula **XV**:

**XV**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

In certain embodiments, the compound is of the formulae **XVa** or **XVb**:

5

**XVa****XVb**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

In certain embodiments, ----- is a double bond. In certain embodiments, X¹ is CR⁴. In
 10 certain embodiments, R⁴ is selected from H, C₁₋₆ alkyl and C₇₋₁₀ aralkyl. However, in certain
 embodiments, X¹ is N. In certain embodiments, X² is CR⁷. In certain embodiments, R⁷ is
 selected from H, C₁₋₆ alkyl and C₇₋₁₀ aralkyl. However, in certain embodiments, X² is N.

In certain embodiments, ----- is a single bond. In certain embodiments, X¹ is CR⁴R⁵. In
 certain embodiments, R⁴ is H and R⁵ is H. In certain embodiments, X² is CR⁷R⁸. In certain
 15 embodiments, R⁷ is H and R⁸ is H. However, in certain embodiments, X² is NR⁹. In certain
 embodiments, R⁹ is selected from H, C₁₋₆ alkyl and -(CR²⁷R²⁸)_p-R²³. In certain embodiments,
 R²³ is phenyl.


In certain embodiments, m is 1 and X³ is NR¹². In certain embodiments, R¹² is selected
 from H, C₁₋₆ alkyl and -(CR²⁷R²⁸)_p-R²³. In certain embodiments, R²³ is phenyl. However, in
 20 certain embodiments, there is no X³, *i.e.*, in certain embodiments, m is 0.

In certain embodiments, R^1 and R^2 , taken together with the carbon to which they are bound, form a carbonyl group (*i.e.*, $=O$).

In certain embodiments, there is no R^3 (*i.e.*, wherein n is 0).

In certain embodiments, Z^1 is $-OR^{13}$ and Z^2 is $-OR^{14}$. In certain embodiments, R^{13} is H
5 and R^{14} is H (*i.e.*, wherein Z^1 and Z^2 are both $-OH$).

In certain embodiments, the compound is selected from:



or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

In another aspect, provided herein are pharmaceutical compositions comprising a compound of the present invention or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, and a pharmaceutically acceptable excipient.

5 In yet another aspect, provided herein are methods of treating an FAAH-mediated disorder comprising administering to a subject in need thereof a therapeutically effective amount of a compound of the present invention or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, or a pharmaceutical composition thereof.

10 In certain embodiments, the FAAH-mediated disorder is selected from a painful disorder, an inflammatory disorder, an immune disorder, depression, anxiety, an anxiety-related disorder, a sleep disorder, a feeding behavior, a movement disorder, glaucoma, neuroprotection and cardiovascular disease.


In certain embodiments, the FAAH-mediated disorder is a painful disorder. In certain embodiments, the painful disorder is selected from neuropathic pain, central pain, deafferentiation pain, chronic pain, stimulus of nociceptive receptors, acute pain, non-inflammatory pain, inflammatory pain, pain associated with cancer, preoperative pain, arthritic pain, lumbosacral pain, musculo-skeletal pain, headache, migraine, muscle ache, lower back and neck pain, and toothache. In certain embodiments, the painful disorder is neuropathic pain. In certain embodiments, the painful disorder is arthritic pain. In certain embodiments, the arthritic pain is osteoarthritic pain. In certain embodiments, the arthritic pain is rheumatoid arthritic pain.

20 In certain embodiments, the inflammatory pain is associated with an inflammatory disorder.

In certain embodiments, the FAAH-mediated disorder is an inflammatory disorder. In certain embodiments, the inflammatory disorder is irritable bowel disease.

DETAILED DESCRIPTION

25 Provided are inhibitors of FAAH that contain at least one Lewis acidic boron head group, such as a boronic acid, boronic ester, borinic acid or borinic ester head group. Such compounds include compounds of formula I or a pharmaceutically acceptable salt, solvate, or prodrug thereof, or a mixture thereof:



wherein:

----- is selected from a single bond and a double bond;

when ----- is a single bond, X^1 is selected from CR^4R^5 and NR^6 , and X^2 is selected from CR^7R^8 and NR^9 ;

when ----- is a double bond, X^1 is selected from CR^4 and N, and X^2 is selected from CR^7 and N;

X^3 is selected from $CR^{10}R^{11}$ and NR^{12} ;

provided that at least one of X^1 , X^2 and X^3 is selected from N, NR^6 , NR^9 or NR^{12} ;

m is 0 or 1;

Z^1 is selected from $-OR^{13}$ and C_{1-6} alkyl;

Z^2 is selected from $-OR^{14}$ and C_{1-6} alkyl;

or alternatively, Z^1 and Z^2 , together with the B to which they are bound, form an 5- to 8-membered ring having at least one O atom directly attached to B, wherein the ring is comprised of carbon atoms and optionally one or more additional heteroatoms independently selected from the group consisting of N, S, and O;

n is 0, 1, 2 or 3;

R^1 and R^2 each independently is selected from H, halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} perhaloalkyl, -CN and $-OR^{15}$;

or alternatively, R^1 and R^2 , taken together with the carbon to which they are bound, form a carbonyl group;

R^3 , at each occurrence, independently is selected from halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} perhaloalkyl, -OH, C_{1-6} alkoxy and -CN;

R^4 , R^5 , R^7 , R^8 , R^{10} , and R^{11} each independently is selected from H, halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} perhaloalkyl, -CN, $-OR^{16}$, $NR^{17}R^{18}$, $-C(O)R^{19}$, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

R^6 , R^9 , and R^{12} each independently is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, $-C(O)R^{20}$, $-C(O)OR^{21}$, $S(O)_2R^{22}$, C_{3-10} carbocyclyl, C_{6-10} aryl, 3-10 membered heterocyclyl, 5-10 membered heteroaryl, and $-(CR^{27}R^{28})_p-R^{23}$;

5 R^{13} and R^{14} , at each occurrence, each independently is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl;

R^{15} and R^{16} , at each occurrence, each independently is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

10 R^{17} and R^{18} , at each occurrence, each independently is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, $-C(O)R^{24}$, $-C(O)OR^{25}$, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

R^{19} , R^{20} and R^{21} , at each occurrence, each independently is selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

15 R^{22} is selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, C_{6-10} aryl, 3-10 membered heterocyclyl, 5-10 membered heteroaryl, and $-(CR^{29}R^{30})_q-R^{26}$;

R^{23} , at each occurrence, independently is selected from C_{3-10} carbocyclyl, C_{6-10} aryl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;



20 R^{24} and R^{25} , at each occurrence, each independently is selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

R^{26} , at each occurrence, independently is selected from C_{3-10} carbocyclyl, C_{6-10} aryl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

25 R^{27} , R^{28} , R^{29} and R^{30} , at each occurrence, each independently is selected from H and C_{1-6} alkyl; and



p, and q, at each occurrence, each independently is selected from 1, 2, 3, 4, 5 and 6.

Embodiments of compounds of formula **I** include compounds of the formulae **Ia** or **Ib**:


**Ia****Ib**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof,
 wherein -----, Z^1 , Z^2 , X^1 , X^2 , X^3 , R^1 , R^2 , R^3 , m and n are as defined above and herein. In certain
 5 preferred embodiments, Z^1 and Z^2 are both OH.

In some embodiments of compounds of formulae **I**, **Ia**, or **Ib**, R^1 and R^2 , taken together
 with the carbon to which they are bound, form a carbonyl group, *i.e.*, compounds of the formula
II, **IIa**, or **IIb**:


**II****IIa**

10

**IIb**


or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, wherein ----- , Z^1 , Z^2 , X^1 , X^2 , X^3 , R^3 , m and n are as defined above and herein. In certain preferred embodiments, Z^1 and Z^2 are both OH.



In some embodiments of compounds of formula **I**, m is 1 and X^3 is NR^{12} , *i.e.*, compounds of the formula **III**:

**III**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, wherein ----- , Z^1 , Z^2 , X^1 , X^2 , R^1 , R^2 , R^3 , R^{12} and n are as defined above and herein. In certain preferred embodiments, Z^1 and Z^2 are both OH.


In some embodiments of compounds of formula **III**, ----- is a double bond, X^1 is CR^4 , and X^2 is CR^7 , *i.e.*, compounds of formulae **IV**, **IVa**, or **IVb**:



**IV**

**IVa****IVb**

5 or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, wherein Z^1 , Z^2 , R^1 , R^2 , R^3 , R^4 , R^7 , R^{12} and n are as defined above and herein. In certain preferred embodiments, Z^1 and Z^2 are both OH.

In certain embodiments of these compounds, R^1 and R^2 , taken together with the carbon to which they are bound, form a carbonyl group, *i.e.*, compounds of formulae **V**, **Va**, or **Vb**:


**V**



**Va****Vb**

5 or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, wherein Z^1 , Z^2 , R^3 , R^4 , R^7 , R^{12} and n are as defined above and herein. In certain preferred embodiments, Z^1 and Z^2 are both OH.

In some embodiments of compounds of formulae **IV**, **IVa**, **IVb**, **V**, **Va**, or **Vb**, R^4 and R^7 each independently is selected from H and C_{1-6} alkyl. In some embodiments, R^4 is H and R^7 is H. In some embodiments of compounds of formulae **IV**, **IVa**, **IVb**, **V**, **Va**, or **Vb**, n is 0.


10 In certain embodiments of compounds of formula **III**, ----- is a single bond, X^1 is CR^4R^5 , and X^2 is CR^7R^8 , *i.e.*, compounds of formulae **VI**, **VIa**, or **VIb**:


**VI**

**VIa****VIb**


5 or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, wherein Z^1 , Z^2 , R^1 , R^2 , R^3 , R^4 , R^5 , R^7 , R^8 , R^{12} and n are as defined above and herein. In certain preferred embodiments, Z^1 and Z^2 are both OH.

In certain embodiments of these compounds, R^1 and R^2 , taken together with the carbon to which they are bound, form a carbonyl group, *i.e.*, compounds of formulae **VII**, **VIIa**, or **VIIb**:

**VII**



VIIa






VIIb

5 or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, wherein Z^1 , Z^2 , R^3 , R^4 , R^5 , R^7 , R^8 , R^{12} and n are as defined above and herein. In certain preferred embodiments, Z^1 and Z^2 are both OH.

In some embodiments of compounds of formulae **VI**, **VIa**, **VIb**, **VII**, **VIIa**, or **VIIb**, R^4 , R^5 , R^7 and R^8 each independently is selected from H and C_{1-6} alkyl. In some embodiments, R^4 is H, R^5 is H, R^7 is H, and R^8 is H. In some embodiments of compounds of formulae **VI**, **VIa**, **VIb**, **VII**, **VIIa**, or **VIIb**, n is 0. In some embodiments of compounds of formulae **III**, **IV**, **IVa**, **IVb**, **V**, **Va**, **Vb**, **VI**, **VIa**, **VIb**, **VII**, **VIIa**, or **VIIb**, R^{12} is not H. In certain embodiments, R^{12} is not $-CH_3$. In certain embodiments, R^{12} is C_{1-6} alkyl or $-(CR^{27}R^{28})_p-R^{23}$. In some embodiments, R^{12} is C_{1-6} alkyl (*e.g.*, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, hexyl, and the like). In other embodiments, R^{12} is $-(CR^{27}R^{28})_p-R^{23}$, wherein R^{23} is C_{6-10} aryl or 5-10 membered heteroaryl. In some embodiments, R^{23} is phenyl. In some embodiments, p is 1, and in others p is 2. In some embodiments, each R^{27} is H and each R^{28} is H. In other embodiments, at least one R^{27} is C_{1-6} alkyl (*e.g.*, methyl) and at least one R^{28} is C_{1-6} alkyl (*e.g.*, methyl). In still other embodiments, at least one R^{27} is C_{1-6} alkyl (*e.g.*, methyl) and each R^{28} is H. In some




embodiments, R^{12} is benzyl, 1-phenethyl, 2-phenethyl, 1-phenylpropyl, 2-phenylpropyl, or 1-phenylpropan-2-yl.

In certain embodiments of compounds of formula **III**, ----- is a double bond, X^1 is N, and X^2 is CR^7 , *i.e.*, compounds of formulae **VIII**, **VIIIa**, or **VIIIb**:

**VIII****VIIIa****VIIIb**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, wherein Z^1 , Z^2 , R^1 , R^2 , R^3 , R^7 , R^{12} and n are as defined above and herein. In certain preferred embodiments, Z^1 and Z^2 are both OH.

In certain embodiments of these compounds, R^1 and R^2 , taken together with the carbon to which they are bound, form a carbonyl group, *i.e.*, compounds of formulae **IX**, **IXa**, or **IXb**:



**IX****IXa****IXb**


or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, wherein Z^1 , Z^2 , R^3 , R^7 , R^{12} and n are as defined above and herein. In certain preferred embodiments, Z^1 and Z^2 are both OH.

- 10 In some embodiments of compounds of formulae **VIII**, **VIIIa**, **VIIIb**, **IX**, **IXa**, or **IXb**, R^7 is H, C_{1-6} alkyl, or C_{7-12} aralkyl. In some embodiments, R^7 is H. In other embodiments, R^7 is C_{1-6} alkyl (*e.g.*, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, and the like). In other embodiments, R^7 is C_{7-12} aralkyl (*e.g.*, benzyl, 1-phenethyl, 2-phenethyl, 1-phenylpropyl, 2-phenylpropyl, 1-phenylpropan-2-yl, and the like). In some embodiments of compounds of
- 15 formulae **VIII**, **VIIIa**, **VIIIb**, **IX**, **IXa**, or **IXb**, n is 0.

In some embodiments of compounds of formulae **VIII**, **VIIIa**, **VIIIb**, **IX**, **IXa**, or **IXb**, R^{12} is not H. In certain embodiments, R^{12} is not $-CH_3$. In certain embodiments, R^{12} is H, C_{1-6} alkyl or $-(CR^{27}R^{28})_p-R^{23}$. In some embodiments, R^{12} is C_{1-6} alkyl (e.g., methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, hexyl, and the like). In other embodiments, R^{12} is $-(CR^{27}R^{28})_p-R^{23}$, wherein R^{23} is C_{6-10} aryl or 5-10 membered heteroaryl. In some embodiments, R^{23} is phenyl. In some embodiments, p is 1, and in others p is 2. In some embodiments, each R^{27} is H and each R^{28} is H. In other embodiments, at least one R^{27} is C_{1-6} alkyl (e.g., methyl) and at least one R^{28} is C_{1-6} alkyl (e.g., methyl). In still other embodiments, at least one R^{27} is C_{1-6} alkyl (e.g., methyl) and each R^{28} is H. In some embodiments, R^{12} is benzyl, 1-phenethyl, 2-phenethyl, 1-phenylpropyl, 2-phenylpropyl, or 1-phenylpropan-2-yl. In other embodiments, R^{12} is H and R^7 is C_{1-6} alkyl (e.g., methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, and the like) or C_{7-12} aralkyl (e.g., benzyl, 1-phenethyl, 2-phenethyl, 1-phenylpropyl, 2-phenylpropyl, 1-phenylpropan-2-yl, and the like).



In certain embodiments of compounds of formula **III**, ----- is a double bond, X^1 is CR^4 , and X^2 is N, i.e., compounds of formulae **X**, **Xa**, or **Xb**:




**Xb**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, wherein Z^1 , Z^2 , R^1 , R^2 , R^3 , R^4 , R^{12} and n are as defined above and herein. In certain preferred embodiments, Z^1 and Z^2 are both OH.

In certain embodiments of these compounds, R^1 and R^2 , taken together with the carbon to which they are bound, form a carbonyl group, *i.e.*, compounds of formulae **XI**, **XIa**, or **XIb**:

**XI****XIa**





**XIb**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, wherein Z^1 , Z^2 , R^3 , R^4 , R^{12} and n are as defined above and herein. In certain preferred
 5 embodiments, Z^1 and Z^2 are both OH.

In some embodiments of compounds of formulae **X**, **Xa**, **Xb**, **XI**, **XIa**, or **XIb**, R^4 is H or C_{1-6} alkyl. In some embodiments, R^4 is H. In some embodiments of compounds of formulae **X**, **Xa**, **Xb**, **XI**, **XIa**, or **XIb**, n is 0.

In some embodiments of compounds of formulae **X**, **Xa**, **Xb**, **XI**, **XIa**, or **XIb**, R^{12} is not
 10 H. In some embodiments of compounds of formulae **X**, **Xa**, **Xb**, **XI**, **XIa**, or **XIb**, R^{12} is not $-CH_3$. In certain embodiments, R^{12} is C_{1-6} alkyl or $-(CR^{27}R^{28})_p-R^{23}$. In some embodiments, R^{12} is C_{1-6} alkyl (*e.g.*, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, hexyl, and the like). In other embodiments, R^{12} is $-(CR^{27}R^{28})_p-R^{23}$, wherein R^{23} is C_{6-10} aryl or 5-10
 15 membered heteroaryl. In some embodiments, R^{23} is phenyl. In some embodiments, p is 1, and in others p is 2. In some embodiments, each R^{27} is H and each R^{28} is H. In other embodiments, at least one R^{27} is C_{1-6} alkyl (*e.g.*, methyl) and at least one R^{28} is C_{1-6} alkyl (*e.g.*, methyl). In still other embodiments, at least one R^{27} is C_{1-6} alkyl (*e.g.*, methyl) and each R^{28} is H. In some
 20 embodiments, R^{12} is benzyl, 1-phenethyl, 2-phenethyl, 1-phenylpropyl, 2-phenylpropyl or 1-phenylpropan-2-yl.

In certain embodiments, a compound of formula **I** or a subset thereof (*e.g.*, a compound of formulae **Ia**, **Ib**, **II**, **IIa**, **IIb**, **III**, **IV**, **IVa**, **IVb**, **V**, **Va**, **Vb**, **VI**, **VIa**, **VIb**, **VII**, **VIIa**, **VIIb**, **VIII**, **VIIa**, **VIIb**, **IX**, **IXa**, or **IXb**) is not any one of the following compounds:




5

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof,



In some embodiments of compounds of formula I, m is 0, *i.e.*, compounds of the formula

XII:

**XII**


or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, wherein Z^1 , Z^2 , R^1 , R^2 , R^3 , X^1 , X^2 and n are as defined above and herein. In certain preferred embodiments, Z^1 and Z^2 are both OH.



Embodiments of compounds of formula **XII** include compounds of formulae **XIIa** or **XIIb**:

**XIIa****XIIb**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, wherein Z^1 , Z^2 , R^1 , R^2 , R^3 , X^1 , X^2 and n are as defined above and herein. In certain preferred embodiments, Z^1 and Z^2 are both OH.


In some embodiments of compounds of formulae **XII**, **XIIa**, or **XIIb**, R^1 and R^2 , taken together with the carbon to which they are bound, form a carbonyl group, *i.e.*, compounds of formulae **XIII**, **XIIIa**, or **XIIIb**:

**XIII**



**XIIIa****XIIIb**

5 or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, wherein ----- , Z^1 , Z^2 , R^3 , X^1 , X^2 and n are as defined above and herein. In certain preferred embodiments, Z^1 and Z^2 are both OH.

In some embodiments of compounds of formula **XII**, ----- is a single bond, X^1 is CR^4R^5 , and X^2 is NR^9 , *i.e.*, compounds of formulae **XIV**, **XIVa**, or **XIVb**:


**XIV**



10

**XIVa****XIVb**

5 or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, wherein Z^1 , Z^2 , R^1 , R^2 , R^3 , R^4 , R^5 , R^9 and n are as defined above and herein. In certain preferred embodiments, Z^1 and Z^2 are both OH.

In certain embodiments of these compounds, R^1 and R^2 , taken together with the carbon to which they are bound, form a carbonyl group, *i.e.*, compounds of formulae **XV**, **XVa**, or **XVb**:

**XV**

**XVa****XVb**


5 or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, wherein Z^1 , Z^2 , R^3 , R^4 , R^5 , R^9 and n are as defined above and herein. In certain preferred embodiments, Z^1 and Z^2 are both OH.

In some embodiments of compounds of formulae **XIV**, **XIVa**, **XIVb**, **XV**, **XVa**, or **XVb**, R^4 and R^5 each independently is selected from H and C_{1-6} alkyl. In some embodiments, R^4 is H and R^5 is H. In some embodiments of compounds of formulae **XIV**, **XIVa**, **XIVb**, **XV**, **XVa**, or **XVb**, n is 0.

In some embodiments of compounds of formulae **XIV**, **XIVa**, **XIVb**, **XV**, **XVa**, or **XVb**, R^9 is C_{1-6} alkyl or $-(CR^{27}R^{28})_p-R^{23}$. In some embodiments, R^9 is C_{1-6} alkyl (*e.g.*, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, hexyl, and the like). However, in certain
 15 embodiments, R^9 is not H. In certain embodiments, R^9 is not $-CH^3$. In other embodiments, R^9 is $-(CR^{27}R^{28})_p-R^{23}$, wherein R^{23} is C_{6-10} aryl or 5-10 membered heteroaryl. In some embodiments, R^{23} is phenyl. In some embodiments, p is 1, and in others p is 2. In some embodiments, each R^{27} is H and each R^{28} is H. In other embodiments, at least one R^{27} is C_{1-6} alkyl (*e.g.*, methyl) and at least one R^{28} is C_{1-6} alkyl (*e.g.*, methyl). In still other embodiments, at least one R^{27} is C_{1-6}

alkyl (*e.g.*, methyl) and each R^{28} is H. In some embodiments, R^9 is benzyl, 1-phenethyl, 2-phenethyl, 1-phenylpropyl, 2-phenylpropyl or 1-phenylpropan-2-yl.

In certain embodiments, a compound of formula **I** or a subset thereof (*e.g.*, a compound of formulae **XII**, **XIIa**, **XIIb**, **XIII**, **XIIIa**, **XIIIb**, **XIX**, **XIXa**, **XIXb**, **XV**, **XVa**, or **XVb**,) is not any one of the following compounds:




or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

In some embodiments of compounds of formulae **I**, **Ia**, **Ib**, **II**, **IIa**, **IIb**, **III**, **IV**, **IVa**, **IVb**, **V**, **Va**, **Vb**, **VI**, **VIa**, **VIb**, **VII**, **VIIa**, **VIIb**, **VIII**, **VIIIa**, **VIIIb**, **IX**, **IXa**, **IXb**, **X**, **Xa**, **Xb**, **XI**, **XIa**, **XIb**, **XII**, **XIIa**, **XIIb**, **XIII**, **XIIIa**, **XIIIb**, **XIV**, **XIVa**, **XIVb**, **XV**, **XVa**, or **XVb**, Z^1 is OR^{13} and Z^2 is OR^{14} . In some embodiments, R^{13} is H and R^{14} is H (*i.e.*, Z^1 is OH and Z^2 is OH).

In other embodiments, Z^1 and Z^2 taken together with the boron atom to which they are bound, form a 5- to 8-membered ring having at least one O, S, N or NR^A atom directly attached to the boron atom, wherein R^A is selected from hydrogen, $-SO_2R^B$, $-SOR^B$, $-C(O)R^B$, $-CO_2R^B$, $-C(O)N(R^B)_2$, C_{1-8} alkyl, C_{2-6} alkenyl, C_{2-8} alkynyl, C_{3-10} carbocyclyl, 3-10 membered heterocyclyl, C_{6-10} aryl, and 5-10 membered heteroaryl; and each instance of R^B is, independently, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, 3-10 membered heterocyclyl, C_{6-10} aryl, or 5-10 membered heteroaryl. In some embodiments, the 5- to 8-membered ring is with one or more groups selected from halogen, oxo ($=O$), $-SO_2R^C$, $-SOR^C$, $-C(O)R^C$, $-C(O)OR^C$, $-C(O)N(R^C)_2$, $-C(O)NHR^C$, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, 3-10 membered heterocyclyl, C_{6-10} aryl, or 5-10 membered heteroaryl groups, or two groups present on the ring are joined to form a 5- to 8-membered monocyclic or bicyclic ring optionally containing one or more heteroatoms selected from O, S, N or NR^A ; wherein each instance of R^C is independently, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, 3-10 membered heterocyclyl, C_{6-10} aryl, or 5-10 membered heteroaryl.

For example, in certain embodiments, Z^1 and Z^2 , taken together with the boron atom to which they are bound, form an optionally substituted 5-membered ring having at least one O, S or NR^A atom directly attached to the boron atom. Exemplary 5-membered rings include, but are not limited to:


5




wherein R^A is as defined herein.

In other embodiments, Z^1 and Z^2 , taken together with the boron atom to which they are bound, form a 6-membered ring having at least one O, S or NR^A atom directly attached to the boron atom. Exemplary 6-membered rings include, but are not limited to:

10



In yet other embodiments, Z^1 and Z^2 form an 8-membered ring having at least one O, S or NR^A atom directly attached to the boron atom. Exemplary 8-membered ring structures include, but are not limited to:



5 wherein R^A is as defined herein.

Definitions

Definitions of specific functional groups and chemical terms are described in more detail below. The chemical elements are identified in accordance with the Periodic Table of the
10 Elements, CAS version, Handbook of Chemistry and Physics, 75th Ed., inside cover, and specific functional groups are generally defined as described therein. Additionally, general principles of organic chemistry, as well as specific functional moieties and reactivity, are described in *Organic Chemistry*, Thomas Sorrell, University Science Books, Sausalito, 1999; Smith and March *March's Advanced Organic Chemistry*, 5th Edition, John Wiley & Sons, Inc., New York,
15 2001; Larock, *Comprehensive Organic Transformations*, VCH Publishers, Inc., New York, 1989; and Carruthers, *Some Modern Methods of Organic Synthesis*, 3rd Edition, Cambridge University Press, Cambridge, 1987.

Certain compounds of the present invention can comprise one or more asymmetric centers, and thus can exist in various isomeric forms, *e.g.*, enantiomers and/or diastereomers.
20 The compounds provided herein can be in the form of an individual enantiomer, diastereomer or geometric isomer, or can be in the form of a mixture of stereoisomers, including racemic mixtures and mixtures enriched in one or more stereoisomer. In certain embodiments, the compounds of the invention are enantiopure compounds. In certain other embodiments, mixtures of stereoisomers are provided.

25 Furthermore, certain compounds, as described herein can have one or more double bonds that can exist as either the *cis* or *trans*, or the *E* or *Z* isomer, unless otherwise indicated. The invention additionally encompasses the compounds as individual isomers substantially free of

other isomers, and alternatively, as mixtures of various isomers, *e.g.*, racemic mixtures of E/Z isomers or mixtures enriched in one E/Z isomer.

Where a particular enantiomer is preferred, it can be provided substantially free of the corresponding enantiomer, *i.e.*, optically enriched. “Optically-enriched,” as used herein, means that the compound is made up of a greater proportion of one enantiomer compared to the other. In certain embodiments the compound is made up of at least about 90% by weight of a preferred enantiomer. In other embodiments the compound is made up of at least about 95%, 98%, or 99% by weight of a preferred enantiomer. Preferred enantiomers can be isolated from mixtures by methods known to those skilled in the art, including chiral high pressure liquid chromatography (HPLC) and the formation and crystallization of chiral salts; or preferred enantiomers can be prepared by asymmetric syntheses. See, for example, Jacques, *et al.*, *Enantiomers, Racemates and Resolutions* (Wiley Interscience, New York, 1981); Wilen, S.H., *et al.*, *Tetrahedron* 33:2725 (1977); Eliel, E.L. *Stereochemistry of Carbon Compounds* (McGraw-Hill, NY, 1962); and Wilen, S.H. *Tables of Resolving Agents and Optical Resolutions* p. 268 (E.L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, IN 1972).

When a range of values is listed, it is intended to encompass each value and sub-range within the range. For example “C₁₋₆ alkyl” is intended to encompass, C₁, C₂, C₃, C₄, C₅, C₆, C₁₋₆, C₁₋₅, C₁₋₄, C₁₋₃, C₁₋₂, C₂₋₆, C₂₋₅, C₂₋₄, C₂₋₃, C₃₋₆, C₃₋₅, C₃₋₄, C₄₋₆, C₄₋₅, and C₅₋₆ alkyl.

As used herein a “direct bond” or “covalent bond” refers to a single bond.

As used herein, the term “boronic acid” refers to any chemical compound comprising a –B(OH)₂ moiety. Arylboronic acid compounds readily form oligomeric anhydrides by dehydration of the boronic acid moiety (see, for example, Snyder *et al.*, *J. Am. Chem. Soc.* (1958) 80: 3611). Thus, unless otherwise apparent from context, the term “boronic acid” is expressly intended to encompass free boronic acids, oligomeric anhydrides, including, but not limited to, dimers, trimers, and tetramers, and mixtures thereof.

The terms “boronic ester”, “borinic acid” and “borinic ester” are art understood terms referring to a –B(OR)₂ moiety, a –B(R)OH moiety and a –B(R)OR moiety, respectively, wherein R is a group other than hydrogen (*e.g.*, an C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, carbocycyl, heterocycyl, aryl, or heteroaryl group; or two R groups are joined to form a 5- to 8-membered ring optionally containing 1 to 4 heteroatoms).

As used herein, alone or as part of another group, “halo” and “halogen” refer to fluorine (fluoro, -F), chlorine (chloro, -Cl), bromine (bromo, -Br), or iodine (iodo, -I).

As used herein, alone or as part of another group, “alkyl” refers to a monoradical of a straight-chain or branched saturated hydrocarbon group having from 1 to 8 carbon atoms (“C₁₋₈ alkyl”). In some embodiments, an alkyl group can have from 1 to 6 carbon atoms (“C₁₋₆ alkyl”). In some embodiments, an alkyl group can have from 1 to 4 carbon atoms (“C₁₋₄ alkyl”). Examples of C₁₋₄ alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl and *tert*-butyl. Examples of C₁₋₆ alkyl groups include the aforementioned C₁₋₄ alkyl groups as well as pentyl, isopentyl, neopentyl, hexyl and the like. Additional examples of alkyl groups include heptyl, octyl and the like. Unless otherwise specified, each instance of an alkyl group is independently unsubstituted or substituted with 1–5 groups as described below.

As used herein, alone or as part of another group, “alkenyl” refers to a monoradical of a straight-chain or branched hydrocarbon group having from 2 to 8 carbon atoms and one or more carbon-carbon double bonds (“C₂₋₈ alkenyl”). In some embodiments, an alkenyl group can have from 2 to 6 carbon atoms (“C₂₋₆ alkenyl”). In some embodiments, an alkenyl group can have from 2 to 4 carbon atoms (“C₂₋₄ alkenyl”). The one or more carbon-carbon double bonds can be internal (such as in 2-butenyl) or terminal (such as in 1-butenyl). Examples of C₂₋₄ alkenyl groups include ethenyl, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl, butadienyl and the like. Examples of C₂₋₆ alkenyl groups include the aforementioned C₂₋₄ alkenyl groups as well as pentenyl, pentadienyl, hexenyl and the like. Additional examples of alkenyl include heptenyl, octenyl, octatrienyl and the like. Unless otherwise specified, each instance of an alkenyl group is independently unsubstituted or substituted with 1–5 groups as described below.

As used herein, alone or as part of another group, “alkynyl” refers to a monoradical of a straight-chain or branched hydrocarbon group having from 2 to 8 carbon atoms and one or more carbon-carbon triple bonds (“C₂₋₈ alkynyl”). In some embodiments, an alkynyl group can have from 2 to 6 carbon atoms (“C₂₋₆ alkynyl”). In some embodiments, an alkynyl group can have from 2 to 4 carbon atoms (“C₂₋₄ alkynyl”). The one or more carbon-carbon triple bonds can be internal (such as in 2-butyne) or terminal (such as in 1-butyne). Examples of C₂₋₄ alkynyl groups include ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne and the like. Examples of C₂₋₆ alkynyl groups include the aforementioned C₂₋₄ alkynyl groups as well as pentynyl, hexynyl and the like. Additional examples of alkynyl include heptynyl, octynyl and the like.

Unless otherwise specified, each instance of an alkynyl group is independently unsubstituted or substituted with 1–5 groups as described below.

As used herein, alone or as part of another group, “alkylene” refers to a diradical of a straight-chain or branched saturated alkyl group having from 1 to 6 carbon atoms (“C₁₋₆ alkylene”). In some embodiments, an alkylene group can have from 1 to 4 carbon atoms (“C₁₋₄ alkylene”). In some embodiments, an alkylene group can have from 1 to 2 carbon atoms (“C₁₋₂ alkylene”). Examples of C₁₋₂ alkylene groups include methylene and ethylene. Examples of C₁₋₄ alkylene groups include the aforementioned C₁₋₂ alkylene groups as well as trimethylene (1,3-propanediyl), propylene (1,2-propanediyl), tetramethylene (1,4-butanediyl), butylene (1,2-butanediyl), 1,3-butanediyl, 2-methyl-1,3-propanediyl and the like. Examples of C₁₋₆ alkylene groups include the aforementioned C₁₋₄ alkylene groups as well as pentamethylene (1,5-pentanediyl), pentylene (1,2-pentanediyl), hexamethylene (1,6-hexanediyl), hexylene (1,2-hexanediyl), 2,3-dimethyl-1,4-butanediyl and the like. In some embodiments, an alkylene group is an α,ω -diradical. Examples of α,ω -diradical alkylene groups include methylene, ethylene, trimethylene, tetramethylene, pentamethylene and hexamethylene. Unless otherwise specified, each instance of an alkylene group is independently unsubstituted or substituted with 1–5 groups as described below.

As used herein, alone or as part of another group, “alkenylene” refers to a diradical of a straight-chain or branched alkenyl group having from 2 to 6 carbon atoms and one or more carbon-carbon double bonds (“C₂₋₆ alkenylene”). In some embodiments, an alkenylene group can have from 2 to 4 carbon atoms (“C₂₋₄ alkenylene”). In some embodiments, an alkenylene group can have 2 carbon atoms, i.e., ethenediyl. The one or more carbon-carbon double bonds can be internal (such as in 1,4-but-2-enediyl) or terminal (such as in 1,4-but-1-enediyl). Examples of C₂₋₄ alkenylene groups include ethenediyl, 1,2-propenediyl, 1,3-propenediyl, 1,4-but-1-enediyl, 1,4-but-2-enediyl and the like. Examples of C₂₋₆ alkenylene groups include the aforementioned C₂₋₄ alkenylene groups as well as 1,5-pent-1-enediyl, 1,4-pent-2-enediyl, 1,6-hex-2-enediyl, 2,5-hex-3-enediyl, 2-methyl-1,4-pent-2-enediyl and the like. In some embodiments, an alkenylene group is an α,ω -diradical. Examples of α,ω -diradical alkenylene groups include ethenediyl, 1,3-propenediyl, 1,4-but-2-enediyl, 1,5-pent-1-enediyl, 1,6-hex-3-enediyl and the like. Unless otherwise specified, each instance of an alkenylene group is independently unsubstituted or substituted with 1–5 groups as described below.

As used herein, alone or as part of another group, “alkynylene” refers to a diradical of a straight-chain or branched alkynyl group having from 2 to 6 carbon atoms and one or more carbon-carbon triple bonds (“C₂₋₆ alkynylene”). In some embodiments, an alkynylene group can have from 2 to 4 carbon atoms (“C₂₋₄ alkynylene”). In some embodiments, an alkynylene group can have 2 carbon atoms, i.e., ethynediyl. The one or more carbon-carbon triple bonds can be internal (such as in 1,4-but-2-ynediyl) or terminal (such as in 1,4-but-1-ynediyl). Examples of C₂₋₄ alkynylene groups include ethynediyl, propynediyl, 1,4-but-1-ynediyl, 1,4-but-2-ynediyl and the like. Examples of C₂₋₆ alkynylene groups include the aforementioned C₂₋₄ alkynylene groups as well as 1,5-pent-1-ynediyl, 1,4-pent-2-ynediyl, 1,6-hex-2-ynediyl, 2,5-hex-3-ynediyl, 3-methyl-1,5-hex-1-ynediyl and the like. In some embodiments, an alkynylene group is an α,ω -diradical. Examples of α,ω -diradical alkynylene groups include ethynediyl, propynediyl, 1,4-but-2-ynediyl, 1,5-pent-1-ynediyl, 1,6-hex-3-ynediyl and the like. Unless otherwise specified, each instance of an alkynylene group is independently unsubstituted or substituted with 1-5 groups as described below.

As used herein, alone or as part of another group, “perhaloalkyl” refers to an alkyl group having from 1 to 6 carbon atoms, wherein all of the hydrogen atoms are each independently replaced with fluoro or chloro. In some embodiments, all of the hydrogen atoms are each replaced with fluoro. In some embodiments, all of the hydrogen atoms are each replaced with chloro. Examples of perhaloalkyl groups include -CF₃, -CF₂CF₃, -CF₂CF₂CF₃, -CCl₃, -CFC₂Cl, -CF₂Cl and the like.

As used herein, alone or as part of another group, “alkoxy” or “alkyloxy” refers to a -O-alkyl group having from 1 to 8 carbon atoms (“C₁₋₈ alkoxy”). In some embodiments, an alkoxy group can have from 1 to 6 carbon atoms (“C₁₋₆ alkoxy”). In some embodiments, an alkoxy group can have from 1 to 4 carbon atoms (“C₁₋₄ alkoxy”). Examples of C₁₋₄ alkoxy groups include methoxy, ethoxy, propoxy, isopropoxy, butoxy, *tert*-butoxy and the like. Examples of C₁₋₆ alkoxy groups include the aforementioned C₁₋₄ alkoxy groups as well as pentyloxy, isopentyloxy, neopentyloxy, hexyloxy and the like. Additional examples of alkoxy groups include heptyloxy, octyloxy and the like. Unless otherwise specified, each instance of an alkoxy group is independently unsubstituted or substituted with 1-5 groups as described below.

As used herein, alone or as part of another group, “perhaloalkoxy” refers to an alkoxy group having from 1 to 3 carbon atoms, wherein all of the hydrogen atoms are each

independently replaced with fluoro or chloro. In some embodiments, all of the hydrogen atoms are each replaced with fluoro. In some embodiments, all of the hydrogen atoms are each replaced with chloro. Examples of perhaloalkoxy groups include $-\text{OCF}_3$, $-\text{OCF}_2\text{CF}_3$, $-\text{OCF}_2\text{CF}_2\text{CF}_3$, $-\text{OCCl}_3$, $-\text{OCFCl}_2$, $-\text{OCF}_2\text{Cl}$ and the like.

5 As used herein, alone or as part of another group, "alkylthio" refers to an $-\text{S}-$ alkyl group having from 1 to 8 carbon atoms. In some embodiments, an alkylthio group can have from 1 to 6 carbon atoms. In some embodiments, an alkylthio group can have from 1 to 4 carbon atoms. Examples of C_{1-4} alkylthio groups include methylthio, ethylthio, propylthio, isopropylthio, butylthio, isobutylthio and the like. Examples of C_{1-6} alkylthio groups include the
10 aforementioned C_{1-4} alkylthio groups as well as pentylthio, isopentylthio, hexylthio and the like. Additional examples of alkylthio groups include heptylthio, octylthio and the like. Unless otherwise specified, each instance of an alkylthio group is independently unsubstituted or substituted with 1–5 groups as described below.

As used herein, alone or as part of another group, "carbocyclyl" or "carbocycle" refers to
15 a radical of a non-aromatic cyclic hydrocarbon group having from 3 to 10 ring carbon atoms (" C_{3-10} carbocyclyl"). In some embodiments, a carbocyclyl group can have from 3 to 8 ring carbon atoms (" C_{3-8} carbocyclyl"). In some embodiments, a carbocyclyl group can have from 3 to 6 ring carbon atoms (" C_{3-6} carbocyclyl"). Examples of C_{3-6} carbocyclyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cyclohexadienyl
20 and the like. Examples of C_{3-8} carbocyclyl groups include the aforementioned C_{3-6} carbocyclyl groups as well as cycloheptyl, cycloheptadienyl, cycloheptatrienyl, cyclooctyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl and the like. Examples of C_{3-10} carbocyclyl groups include the aforementioned C_{3-8} carbocyclyl groups as well as octahydro-1*H*-indenyl, decahydronaphthalenyl, spiro[4.5]decanyl and the like. As the foregoing examples illustrate, in
25 some embodiments a carbocyclyl group can be monocyclic ("monocyclic carbocyclyl") or bicyclic ("bicyclic carbocyclyl", e.g., containing a fused, bridged or spiro ring system), and can be saturated or can contain one or more carbon-carbon double or triple bonds. "Carbocyclyl" also refers to a phenyl group (as defined below) fused to a monocyclic carbocyclyl group. Examples of such carbocyclyl groups include 1,2,3,4-tetrahydronaphthalene (e.g., 1,2,3,4-tetra-
30 hydronaphthalen-1-yl, 1,2,3,4-tetrahydronaphthalen-5-yl, and the like), 2,3-dihydro-1*H*-indene (e.g., 2,3-dihydro-1*H*-inden-1-yl, 2,3-dihydro-1*H*-inden-4-yl, and the like), indene (e.g., 1*H*-

inden-1-yl, 1H-inden-7-yl, and the like), 5,6,7,8-tetrahydroquinoline (e.g., 5,6,7,8-tetrahydroquinolin-5-yl, 5,6,7,8-tetrahydroquinolin-2-yl, and the like), 4,5,6,7-tetrahydro-1H-indole (e.g., 4,5,6,7-tetrahydro-1H-indol-4-yl, 4,5,6,7-tetrahydro-1H-indol-3-yl, and the like), 4,5,6,7-tetrahydrobenzofuran (e.g., 4,5,6,7-tetrahydrobenzofuran-7-yl, 4,5,6,7-tetrahydrobenzofuran-2-yl, and the like) and the like. Unless otherwise specified, each instance of a carbocyclyl or carbocycle group is independently unsubstituted or substituted with 1-5 groups as described below.

In some embodiments, “carbocyclyl” or “carbocycle” can refer to a monocyclic, saturated carbocyclyl group (“cycloalkyl”) having from 3 to 10 ring carbon atoms (“C₃₋₁₀ cycloalkyl”). In some embodiments, a cycloalkyl group can have from 3 to 6 ring carbon atoms (“C₃₋₆ cycloalkyl”). In some embodiments, a cycloalkyl group can have from 5 to 6 ring carbon atoms (“C₅₋₆ cycloalkyl”). Examples of C₅₋₆ cycloalkyl groups include cyclopentyl and cyclohexyl. Examples of C₃₋₆ cycloalkyl groups include the aforementioned C₅₋₆ cycloalkyl groups as well as cyclopropyl and cyclobutyl. Examples of C₃₋₈ cycloalkyl groups include the aforementioned C₃₋₆ cycloalkyl groups as well as cycloheptyl and cyclooctyl. Unless otherwise specified, each instance of a cycloalkyl group is independently unsubstituted or substituted with 1-5 groups as described below.

As used herein, alone or as part of another group, “heterocyclyl” or “heterocycle” refers to a radical of a 3– to 10–membered non–aromatic ring system having ring carbon atoms and 1 to 4 ring heteroatoms, each heteroatom independently selected from nitrogen, oxygen and sulfur. In some embodiments, a heterocyclyl group can have from 3 to 7 ring atoms selected from carbon atoms and 1 to 3 heteroatoms, each heteroatom independently selected from nitrogen, oxygen and sulfur. In some embodiments, a heterocyclyl group can have from 5 to 7 ring atoms selected from carbon atoms and 1 or 2 heteroatoms, each heteroatom independently selected from nitrogen, oxygen and sulfur. In some embodiments, a heterocyclyl group can have from 5 to 6 ring atoms selected from carbon atoms and 1 to 3 heteroatoms, each heteroatom independently selected from nitrogen, oxygen and sulfur. Heterocyclyl groups can be saturated or can contain one or more carbon–carbon double bonds, carbon-nitrogen double bonds, or carbon-carbon triple bonds. In heterocyclyl groups that contain one or more nitrogen atoms, the point of attachment can be a carbon or nitrogen atom, as valency permits.

Examples of heterocyclyl groups with 1–2 ring heteroatoms include oxiranyl, aziridinyl, oxetanyl, azetidiny, pyrrolidinyl, dihydropyrrolyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydrothiophenyl, dihydrothiophenyl, pyrazolidinyl, imidazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, piperidinyl, tetrahydropyridinyl, dihydropyridinyl, piperazinyl, tetrahydropyranyl, dioxanyl, morpholinyl, azepanyl, diazepanyl, diazepinyl, oxepanyl, dioxepanyl, oxazepanyl, oxazepinyl and the like. Examples of heterocyclyl groups with 1–3 heteroatoms include the aforementioned heterocyclyl groups as well as triazolidinyl, oxadiazolidinyl, triazinanyl and the like. Heterocyclyl groups can be monocyclic (“monocyclic heterocyclyl”) as in the aforementioned examples, bicyclic (“bicyclic heterocyclyl”), or tricyclic (“tricyclic heterocyclyl”). Bicyclic heterocyclyl groups can include one or more heteroatoms in one or both rings. Examples of such bicyclic heterocyclyl groups include tetrahydroindolyl, decahydroquinolyl, decahydroisoquinolyl, octahydrochromenyl, octahydroisochromenyl, decahydronaphthyridinyl, decahydro–1,8–naphthyridinyl, octahydropyrrolo[3,2–b]pyrrole and the like.

“Heterocyclyl” or “heterocycle” also refers to a radical of a 5– to 10–membered fused ring system having ring carbon atoms and 1 to 4 ring heteroatoms, each heteroatom independently selected from nitrogen, oxygen and sulfur, wherein one ring is aromatic and the other is non–aromatic. In some embodiments, at least one heteroatom is present in either the aromatic or non–aromatic ring, while in other embodiments, at least one heteroatom is present in both rings. In heterocyclyl groups that contain one or more nitrogen atoms, the point of attachment can be a carbon or nitrogen atom, as valency permits. Examples of such heterocyclyl groups include indolyl (e.g., indolin–1–yl, indolin–4–yl, and the like), isoindolyl (e.g., isoindolin–1–yl, isoindolin–4–yl, and the like), 4,5,6,7–tetrahydro–1H–indolyl (e.g., tetrahydro–1H–indol–2–yl, 4,5,6,7–tetrahydro–1H–indol–4–yl, and the like), dihydrobenzofuranyl (e.g., dihydrobenzofuran–3–yl, dihydrobenzofuran–5–yl, and the like), 4,5,6,7–tetrahydrobenzofuranyl (e.g., 4,5,6,7–tetrahydrobenzofuran–2–yl, 4,5,6,7–tetrahydrobenzofuran–5–yl, and the like), dihydrobenzothienyl (e.g., dihydrobenzothien–2–yl, dihydrobenzothien–4–yl, and the like), 4,5,6,7–tetrahydrobenzothiophenyl (e.g., 4,5,6,7–tetrahydrobenzothiophen–2–yl, 4,5,6,7–tetrahydrobenzothiophen–7–yl, and the like), 1,2,3,4–tetrahydroquinolyl (e.g., 1,2,3,4–tetrahydroquinolin–1–yl, 1,2,3,4–tetrahydroquinolin–7–yl, and the like), chromanyl (e.g., chroman–2–yl, chroman–5–yl, and the like), chromenyl (chromen–4–yl, chromen–8–yl, and the

like), thiochromanyl (e.g., thiochroman-3-yl, isochroman-7-yl, and the like), 1H-benzo[e][1,4]diazepinyl (e.g., 1H-benzo[e][1,4]diazepin-2-yl, 1H-benzo[e][1,4]diazepin-6-yl, and the like), 2,3-dihydro-1H-pyrrolo[2,3-b]pyridinyl (e.g., 2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl, 2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-4-yl, and the like), 4,5,6,7-tetrahydro-1H-pyrrolo[2,3-b]pyridinyl (e.g., 4,5,6,7-tetrahydro-1H-pyrrolo[2,3-b]pyridin-2-yl, 4,5,6,7-tetrahydro-1H-pyrrolo[2,3-b]pyridin-4-yl, and the like), 1,4,5,7-tetrahydropyrano[3,4-b]pyrrolyl (e.g., 1,4,5,7-tetrahydropyrano[3,4-b]pyrrol-2-yl, 1,4,5,7-tetrahydropyrano[3,4-b]pyrrol-4-yl, and the like), 2,3-dihydrofuro[2,3-b]pyridinyl (e.g., 2,3-dihydrofuro[2,3-b]pyridin-3-yl, 2,3-dihydrofuro[2,3-b]pyridin-5-yl, and the like), 4,5,6,7-tetrahydrofuro[3,2-c]pyridinyl (e.g., 4,5,6,7-tetrahydrofuro[3,2-c]pyridin-2-yl, 4,5,6,7-tetrahydrofuro[3,2-c]pyridin-5-yl, and the like), 4,5,6,7-tetrahydrothieno[3,2-b]pyridinyl (e.g., 4,5,6,7-tetrahydrothieno[3,2-b]pyridin-2-yl, 4,5,6,7-tetrahydrothieno[3,2-b]pyridin-7-yl, and the like), 5,6-dihydro-4H-furo[3,2-b]pyrrolyl (e.g., 5,6-dihydro-4H-furo[3,2-b]pyrrol-6-yl, 5,6-dihydro-4H-furo[3,2-b]pyrrol-2-yl, and the like), 6,7-dihydro-5H-furo[3,2-b]pyranyl (e.g., 6,7-dihydro-5H-furo[3,2-b]pyran-2-yl, 6,7-dihydro-5H-furo[3,2-b]pyran-6-yl, and the like), 5,7-dihydro-4H-thieno[2,3-c]pyranyl (e.g., 5,7-dihydro-4H-thieno[2,3-c]pyran-2-yl, 5,7-dihydro-4H-thieno[2,3-c]pyran-4-yl, and the like), 1,2,3,4-tetrahydro-1,6-naphthyridinyl (e.g., 1,2,3,4-tetrahydro-1,6-naphthyridin-3-yl, 1,2,3,4-tetrahydro-1,6-naphthyridin-8-yl, and the like), and the like.

Unless otherwise specified, each instance of a heterocyclcyl group is independently unsubstituted or substituted with 1-5 groups as described below.

As used herein, alone or as part of another group, "aryl" refers to a radical of an aromatic monocyclic or bicyclic ring system having 6 or 10 ring carbon atoms. Examples of such aryl groups include phenyl, 1-naphthyl and 2-naphthyl. Unless otherwise specified, each instance of an aryl group is independently unsubstituted or substituted with 1-5 groups as described below.

The term "aralkyl" refers to an alkyl group substituted by an aryl group, wherein the alkyl and aryl portions independently are as described below.

As used herein, alone or as part of another group, "heteroaryl" refers to a radical of a 5- to 10-membered aromatic ring system having ring carbon atoms and 1 to 4 ring heteroatoms, each heteroatom independently selected from nitrogen, oxygen and sulfur. Examples of such heteroaryl groups include pyrrolyl, furanyl (furyl), thiophenyl (thienyl), pyrazolyl, imidazolyl,

oxazolyl, isoxazolyl, thiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridinyl (pyridyl), pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, indolyl, benzofuranyl, benzothiophenyl (benzothienyl), indazolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, quinolinyl, isoquinolinyl, cinnolinyl, quinazolinyl, quinoxalinyl, phthalazinyl, naphthyridinyl and the like. As the foregoing examples illustrate, in some embodiments a heteroaryl group can be monocyclic ("monocyclic heteroaryl"), and in some embodiments a heteroaryl group can be bicyclic ("bicyclic heteroaryl"). For bicyclic heteroaryl groups wherein one ring does not contain a heteroatom (e.g., indolyl, quinolinyl, and the like) the point of attachment can be on either ring, i.e., either the ring bearing a heteroatom (e.g., 2-indolyl) or the ring that does not contain a heteroatom (e.g., 5-indolyl). Unless otherwise specified, each instance of an heteroaryl group is independently unsubstituted or substituted with 1–5 groups as described below.

The term "heteroaralkyl" refers to an alkyl group substituted by a heteroaryl group, wherein the alkyl and heteroaryl portions independently are as described below.

As used herein, the term "partially unsaturated" refers to a ring moiety that includes at least one double or triple bond. The term "partially unsaturated" is intended to encompass rings having multiple sites of unsaturation, but is not intended to include aryl or heteroaryl moieties, as herein defined.

As described herein, alkyl, alkenyl, alkynyl, alkylene, alkenylene, alkynylene, alkoxy, alkylthio, carbocycle, heterocycle, aryl, aralkyl, heteroaryl, and heteroaralkyl groups as described above and herein are substituted or unsubstituted (*i.e.*, optionally substituted). In general, the term "substituted", whether preceded by the term "optionally" or not, means that one or more hydrogens of the designated moiety are replaced with a suitable substituent. Unless otherwise indicated, a substituted group can have a suitable substituent at each substitutable position of the group, and when more than one position in any given structure is substituted with more than one substituent, then the substituent can be either the same or different at these positions.

Combinations of substituents envisioned by this invention are preferably those that result in the formation of stable compounds. The term "stable", as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and, in certain embodiments, their recovery, purification, and/or use for one or more of the purposes disclosed herein.

Suitable monovalent substituents on a carbon atom are independently halogen; $-(CH_2)_{0-4}R^\circ$; $-(CH_2)_{0-4}OR^\circ$; $-O-(CH_2)_{0-4}C(O)OR^\circ$; $-(CH_2)_{0-4}CH(OR^\circ)_2$; $-(CH_2)_{0-4}SR^\circ$; $-(CH_2)_{0-4}Ph$, which can be substituted with one or more R° ; $-(CH_2)_{0-4}O(CH_2)_{0-1}Ph$ which can be substituted with R° ; $-CH=CHPh$, which can be substituted with one or more R° ; $-NO_2$; $-CN$; $-N_3$; $-(CH_2)_{0-4}N(R^\circ)_2$; $-(CH_2)_{0-4}N(R^\circ)C(O)R^\circ$; $-N(R^\circ)C(S)R^\circ$; $-(CH_2)_{0-4}N(R^\circ)C(O)NR^\circ_2$; $-N(R^\circ)C(S)NR^\circ_2$; $-(CH_2)_{0-4}N(R^\circ)C(O)OR^\circ$; $-N(R^\circ)N(R^\circ)C(O)R^\circ$; $-N(R^\circ)N(R^\circ)C(O)NR^\circ_2$; $-N(R^\circ)N(R^\circ)C(O)OR^\circ$; $-(CH_2)_{0-4}C(O)R^\circ$; $-C(S)R^\circ$; $-(CH_2)_{0-4}C(O)OR^\circ$; $-(CH_2)_{0-4}C(O)SR^\circ$; $-(CH_2)_{0-4}C(O)OSiR^\circ_3$; $-(CH_2)_{0-4}OC(O)R^\circ$; $-OC(O)(CH_2)_{0-4}SR$, $SC(S)SR^\circ$; $-(CH_2)_{0-4}SC(O)R^\circ$; $-(CH_2)_{0-4}C(O)NR^\circ_2$; $-C(S)NR^\circ_2$; $-C(S)SR^\circ$; $-SC(S)SR^\circ$; $-(CH_2)_{0-4}OC(O)NR^\circ_2$; $-C(O)N(OR^\circ)R^\circ$; $-C(O)C(O)R^\circ$; $-C(O)CH_2C(O)R^\circ$; $-C(NOR^\circ)R^\circ$; $-(CH_2)_{0-4}SSR^\circ$; $-(CH_2)_{0-4}S(O)_2R^\circ$; $-(CH_2)_{0-4}S(O)_2OR^\circ$; $-(CH_2)_{0-4}OS(O)_2R^\circ$; $-S(O)_2NR^\circ_2$; $-(CH_2)_{0-4}S(O)R^\circ$; $-N(R^\circ)S(O)_2NR^\circ_2$; $-N(R^\circ)S(O)_2R^\circ$; $-N(OR^\circ)R^\circ$; $-C(NH)NR^\circ_2$; $-P(O)_2R^\circ$; $-P(O)R^\circ_2$; $-OP(O)R^\circ_2$; $-OP(O)(OR^\circ)_2$; SiR°_3 ; $-(C_{1-4} \text{ alkylene})O-N(R^\circ)_2$; or $-(C_{1-4} \text{ alkylene})C(O)O-N(R^\circ)_2$, wherein each R° can be substituted as defined below and is independently hydrogen, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{1-8} heteroalkyl, C_{2-8} heteroalkenyl, C_{2-8} heteroalkynyl, $-CH_2Ph$, $-O(CH_2)_{0-1}Ph$, or a 5- or 6-membered saturated, partially unsaturated, or aromatic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or, notwithstanding the definition above, two independent occurrences of R° , taken together with the atom(s) to which they are bound, form a 3- to 12-membered saturated, partially unsaturated, or aromatic mono- or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, which can be substituted as defined below.

Suitable monovalent substituents on $R^\bullet$ (or the ring formed by two independent occurrences of $R^\bullet$ together with the atoms to which they are bound), are independently halogen, $-(CH_2)_{0-2}R^\bullet$, $-(CH_2)_{0-2}OH$, $-(CH_2)_{0-2}OR^\bullet$, $-(CH_2)_{0-2}CH(OR^\bullet)_2$, $-CN$, $-N_3$, $-(CH_2)_{0-2}C(O)R^\bullet$, $-(CH_2)_{0-2}C(O)OH$, $-(CH_2)_{0-2}C(O)OR^\bullet$, $-(CH_2)_{0-2}SR^\bullet$, $-(CH_2)_{0-2}SH$, $-(CH_2)_{0-2}NH_2$, $-(CH_2)_{0-2}NHR^\bullet$, $-(CH_2)_{0-2}NR^\bullet_2$, $-NO_2$, $-SiR^\bullet_3$, $-OSiR^\bullet_3$, $-C(O)SR^\bullet$, $-(C_{1-4} \text{ alkylene})C(O)OR^\bullet$, or $-SSR^\bullet$ wherein each $R^\bullet$ is unsubstituted or substituted with one or more halogens, and is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, $-CH_2Ph$, $-O(CH_2)_{0-1}Ph$, or a 5- or 6-membered saturated, partially unsaturated, or aromatic ring having 0-4 heteroatoms

independently selected from nitrogen, oxygen, or sulfur. Suitable divalent substituents on a saturated carbon atom of R° include $=O$ and $=S$.

Suitable divalent substituents on a saturated carbon atom include the following: $=O$, $=S$, $=NNR^*_2$, $=NNHC(O)R^*$, $=NNHC(O)OR^*$, $=NNHS(O)_2R^*$, $=NR^*$, $=NOR^*$, $-O(C(R^*_2))_{2-3}O-$, or $-S(C(R^*_2))_{2-3}S-$, wherein each independent occurrence of R^* is selected from hydrogen; C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, each of which can be substituted as defined below; or an unsubstituted 5- or 6-membered saturated, partially unsaturated, or aromatic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. Suitable divalent substituents that are bound to vicinal substitutable carbons include: $-O(CR^*_2)_{2-3}O-$, wherein each independent occurrence of R^* is selected from hydrogen; $-$, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, each of which can be substituted as defined below; or an unsubstituted 5- or 6-membered saturated, partially unsaturated, or aromatic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

Suitable substituents on a C_{1-6} alkyl, C_{2-6} alkenyl, or C_{2-6} alkynyl $R^\bullet$ group include halogen, $-R^\bullet$, $-(haloR^\bullet)$, $-OH$, $-OR^\bullet$, $-CN$, $-C(O)OH$, $-C(O)OR^\bullet$, $-NH_2$, $-NHR^\bullet$, $-NR^\bullet_2$, or $-NO_2$, wherein each $R^\bullet$ is unsubstituted or substituted with one or more halogens, and is independently C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, $-CH_2Ph$, $-O(CH_2)_{0-1}Ph$, or a 5- or 6-membered saturated, partially unsaturated, or aromatic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

Suitable substituents on a substitutable nitrogen include $-R^\dagger$, $-NR^\dagger_2$, $-C(O)R^\dagger$, $-C(O)OR^\dagger$, $-C(O)C(O)R^\dagger$, $-C(O)CH_2C(O)R^\dagger$, $-S(O)_2R^\dagger$, $-S(O)_2NR^\dagger_2$, $-C(S)NR^\dagger_2$, $-C(NH)NR^\dagger_2$, or $-N(R^\dagger)S(O)_2R^\dagger$; wherein each $R^\dagger$ is independently hydrogen; C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, each of which can be substituted as defined below; unsubstituted $-OPh$; or an unsubstituted 5- or 6-membered saturated, partially unsaturated, or aromatic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or, notwithstanding the definition above, two independent occurrences of $R^\dagger$, taken together with the atom(s) to which they are bound form an unsubstituted 3- to 12-membered saturated, partially unsaturated, or aromatic mono- or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

Suitable substituents on a C_{1-6} alkyl, C_{2-6} alkenyl, or C_{2-6} alkynyl $R^\dagger$ group are independently halogen, $-R^\bullet$, $-OH$, $-OR^\bullet$, $-CN$, $-C(O)OH$, $-C(O)OR^\bullet$, $-NH_2$, $-NHR^\bullet$, $-NR^\bullet_2$,

or $-\text{NO}_2$, wherein each $\text{R}^\bullet$ is unsubstituted or substituted with one or more halogens, and is independently C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, $-\text{CH}_2\text{Ph}$, $-\text{O}(\text{CH}_2)_{0-1}\text{Ph}$, or a 5- or 6-membered saturated, partially unsaturated, or aromatic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

5 In certain embodiments, each C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl, alone or as part of another group, is independently and with 1-5 substituents selected from halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} perhaloalkyl, $-\text{OH}$, C_{1-6} alkoxy, $-\text{NH}_2$, $-\text{NH}(\text{C}_{1-6} \text{ alkyl})$, $-\text{N}(\text{C}_{1-6} \text{ alkyl})_2$, and $-\text{CN}$.

10 As used herein, “solvate” refers to a compound of the present invention or a pharmaceutically acceptable salt or prodrug thereof, that further includes a stoichiometric or non-stoichiometric amount of solvent bound by non-covalent intermolecular forces. Where the solvent is water, the solvate is a hydrate.

 As used herein, the term “pharmaceutically acceptable salt” refers to those salts which
15 are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge et al., describe pharmaceutically acceptable salts in detail in J. Pharmaceutical Sciences, 1977, 66, 1-19, incorporated herein by reference.

20 Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid
25 or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide,
30 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate,

palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C_{1-4}alkyl)_4$ salts. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate.

As used herein, and unless otherwise specified, the term “prodrug” means a biologically active derivative of a compound that can hydrolyze, oxidize, or otherwise react under physiological conditions (in vitro or in vivo) to provide the pharmacologically active compound. Additionally, prodrugs can be converted to the compounds of the invention by chemical or biochemical methods in an ex vivo environment. For example, prodrugs can be slowly converted to the compounds of the invention when placed in a transdermal patch reservoir with a suitable enzyme or chemical reagent. Prodrugs are often useful because, in some situations, they may be easier to administer than the parent drug. They may, for instance, be bioavailable by oral administration whereas the parent drug is not. The prodrug may also have improved solubility in pharmaceutical compositions over the parent drug. Examples of prodrugs include, but are not limited to, compounds that comprise biohydrolyzable moieties such as biohydrolyzable amides, biohydrolyzable esters, biohydrolyzable carbamates, biohydrolyzable carbonates, biohydrolyzable ureides, and biohydrolyzable phosphate analogues. Other examples of prodrugs include compounds that comprise --NO, --NO₂, --ONO, or --ONO₂ moieties. The compounds of the invention readily undergo dehydration to form oligomeric anhydrides by dehydration of the boronic acid moiety to form dimers, trimers, and tetramers, and mixtures thereof. These oligomeric species hydrolyze under physiological conditions to reform the boronic acid. As such, the oligomeric anhydrides are contemplated as a “prodrug” of the compounds described herein, and can be used in the treatment of disorder and/or conditions wherein the inhibition of FAAH provides a therapeutic effect.

Exemplary prodrugs of the compounds described herein include, but are not limited to, compounds wherein Z¹ and Z² taken together form a 5- to 8-membered ring having at least one heteroatom atom selected from nitrogen, oxygen and sulfur directly attached to boron, wherein

the ring is comprised of carbon atoms and optionally one or more additional heteroatoms independently selected from nitrogen, oxygen and sulfur.

Other examples of prodrugs of the compounds described herein are trifluoroborate prodrugs which hydrolyze to the boronic acid (*i.e.*, $-\text{BF}_3$ hydrolyzing to $-\text{B}(\text{OH})_2$) at acidic pH.

5 Salt forms of the boronic acid (*e.g.*, Na^+ , Li^+ , Mg^{2+} , Ca^{2+} , and the like) are also considered prodrugs. Amino acids can be used to form prodrugs, such as, for example, serine and cysteine protected boronic acids. 1,2 and 1,3 hydroxy sugars can be used to form prodrugs, such as, for example, glycerol, erythritol, threitol, ribitol, arabinitol, xylitol, allitol, altritol, galactitol, sorbitol, mannitol, and iditol protected boronic acids. Other sugars which are useful in the
10 formation of prodrugs include, but are not limited to, maltitol, lactitol, and isomalt; other monosaccharides which include hexoses (*e.g.*, allose, altrose, glucose, mannose, gulose, idose, galactose, talose) and pentoses (*e.g.*, ribose, arabinaose, xylose, lyxose); pentaerythritols and structural derivatives thereof, such as methylated, ethylated, acetate, ethoxylate, and propoxylate derivatives; and phenolic polyols such as 1,2,4 benzenetriol, 5-methyl benzene1,2,3-triol, 2,3,4-
15 trihydroxybenzaldehyde, and 3,4,5-trihydroxybenzamide. Prodrugs also include NMIDA-derivatives.

Pharmaceutically Acceptable Compositions and Formulations

In certain embodiments, the present invention provides a pharmaceutically acceptable
20 composition comprising a compound of any of formulae **I**, **Ia**, **Ib**, **II**, **IIa**, **IIb**, **III**, **IV**, **IVa**, **IVb**, **V**, **Va**, **Vb**, **VI**, **VIa**, **VIb**, **VII**, **VIIa**, **VIIb**, **VIII**, **VIIIa**, **VIIIb**, **IX**, **IXa**, **IXb**, **X**, **Xa**, **Xb**, **XI**, **XIa**, **XIb**, **XII**, **XIIa**, **XIIb**, **XIII**, **XIIIa**, **XIIIb**, **XIV**, **XIVa**, **XIVb**, **XV**, **XVa**, or **XVb**, or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, and a pharmaceutically acceptable excipient.

25 Pharmaceutically acceptable excipients include any and all solvents, diluents or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. General considerations in the formulation and/or manufacture of pharmaceutically acceptable compositions agents can be found, for example, in *Remington's*
30 *Pharmaceutical Sciences*, Sixteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa.,

1980), and *Remington: The Science and Practice of Pharmacy*, 21st Edition (Lippincott Williams & Wilkins, 2005).

Pharmaceutically acceptable compositions described herein can be prepared by any method known in the art of pharmacology. In general, such preparatory methods include the steps of bringing the active ingredient into association with an excipient and/or one or more other accessory ingredients, and then, if necessary and/or desirable, shaping and/or packaging the product into a desired single- or multi-dose unit.

Pharmaceutically acceptable compositions can be prepared, packaged, and/or sold in bulk, as a single unit dose, and/or as a plurality of single unit doses. As used herein, a “unit dose” is discrete amount of the pharmaceutically acceptable composition comprising a predetermined amount of the active ingredient. The amount of the active ingredient is generally equal to the dosage of the active ingredient which would be administered to a subject and/or a convenient fraction of such a dosage such as, for example, one-half or one-third of such a dosage.

Relative amounts of the active ingredient, the pharmaceutically acceptable excipient, and/or any additional ingredients in a pharmaceutically acceptable composition of the invention will vary, depending upon the identity, size, and/or condition of the subject treated and further depending upon the route by which the composition is to be administered. By way of example, the composition can comprise between about 0.1% and about 100% (w/w) active ingredient, or between about 2% and about 90% (w/w) active ingredient, or between about 5% and about 80% (w/w) active ingredient.

Pharmaceutically acceptable excipients used in the manufacture of provided pharmaceutically acceptable compositions include inert diluents, dispersing and/or granulating agents, surface active agents and/or emulsifiers, disintegrating agents, binding agents, preservatives, buffering agents, lubricating agents, and/or oils. Excipients such as cocoa butter and suppository waxes, coloring agents, coating agents, sweetening, flavoring, and perfuming agents can also be present in the composition.

Exemplary diluents include calcium carbonate, sodium carbonate, calcium phosphate, dicalcium phosphate, calcium sulfate, calcium hydrogen phosphate, sodium phosphate lactose, sucrose, cellulose, microcrystalline cellulose, kaolin, mannitol, sorbitol, inositol, sodium chloride, dry starch, cornstarch, powdered sugar, *etc.*, and combinations thereof.

Exemplary granulating and/or dispersing agents include potato starch, corn starch, tapioca starch, sodium starch glycolate, clays, alginic acid, guar gum, citrus pulp, agar, bentonite, cellulose and wood products, natural sponge, cation-exchange resins, calcium carbonate, silicates, sodium carbonate, cross-linked poly(vinyl-pyrrolidone) (crospovidone), sodium carboxymethyl starch (sodium starch glycolate), carboxymethyl cellulose, cross-linked sodium carboxymethyl cellulose (croscarmellose), methylcellulose, pregelatinized starch (starch 1500), microcrystalline starch, water insoluble starch, calcium carboxymethyl cellulose, magnesium aluminum silicate (Veegum), sodium lauryl sulfate, quaternary ammonium compounds, *etc.*, and combinations thereof.

Exemplary surface active agents and/or emulsifiers include natural emulsifiers (*e.g.* acacia, agar, alginic acid, sodium alginate, tragacanth, chondrux, cholesterol, xanthan, pectin, gelatin, egg yolk, casein, wool fat, cholesterol, wax, and lecithin), colloidal clays (*e.g.* bentonite [aluminum silicate] and Veegum [magnesium aluminum silicate]), long chain amino acid derivatives, high molecular weight alcohols (*e.g.* stearyl alcohol, cetyl alcohol, oleyl alcohol, triacetin monostearate, ethylene glycol distearate, glyceryl monostearate, and propylene glycol monostearate, polyvinyl alcohol), carbomers (*e.g.* carboxy polymethylene, polyacrylic acid, acrylic acid polymer, and carboxyvinyl polymer), carrageenan, cellulosic derivatives (*e.g.* carboxymethylcellulose sodium, powdered cellulose, hydroxymethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, methylcellulose), sorbitan fatty acid esters (*e.g.* polyoxyethylene sorbitan monolaurate [Tween 20], polyoxyethylene sorbitan [Tween 60], polyoxyethylene sorbitan monooleate [Tween 80], sorbitan monopalmitate [Span 40], sorbitan monostearate [Span 60], sorbitan tristearate [Span 65], glyceryl monooleate, sorbitan monooleate [Span 80]), polyoxyethylene esters (*e.g.* polyoxyethylene monostearate [Myrj 45], polyoxyethylene hydrogenated castor oil, polyethoxylated castor oil, polyoxymethylene stearate, and Solutol), sucrose fatty acid esters, polyethylene glycol fatty acid esters (*e.g.* Cremophor), polyoxyethylene ethers, (*e.g.* polyoxyethylene lauryl ether [Brij 30]), poly(vinyl-pyrrolidone), diethylene glycol monolaurate, triethanolamine oleate, sodium oleate, potassium oleate, ethyl oleate, oleic acid, ethyl laurate, sodium lauryl sulfate, Pluronic F 68, Poloxamer 188, cetrimonium bromide, cetylpyridinium chloride, benzalkonium chloride, docusate sodium, *etc.* and/or combinations thereof.

Exemplary binding agents include starch (*e.g.* cornstarch and starch paste), gelatin, sugars (*e.g.* sucrose, glucose, dextrose, dextrin, molasses, lactose, lactitol, mannitol, *etc.*), natural and synthetic gums (*e.g.* acacia, sodium alginate, extract of Irish moss, panwar gum, ghatti gum, mucilage of isapol husks, carboxymethylcellulose, methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, microcrystalline cellulose, cellulose acetate, poly(vinyl-pyrrolidone), magnesium aluminum silicate (Veegum), and larch arabogalactan), alginates, polyethylene oxide, polyethylene glycol, inorganic calcium salts, silicic acid, polymethacrylates, waxes, water, alcohol, *etc.*, and/or combinations thereof.

Exemplary preservatives include antioxidants, chelating agents, antimicrobial preservatives, antifungal preservatives, alcohol preservatives, acidic preservatives, and other preservatives.

Exemplary antioxidants include alpha tocopherol, ascorbic acid, ascorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, monothioglycerol, potassium metabisulfite, propionic acid, propyl gallate, sodium ascorbate, sodium bisulfite, sodium metabisulfite, and sodium sulfite.

Exemplary chelating agents include ethylenediaminetetraacetic acid (EDTA) and salts and hydrates thereof (*e.g.*, sodium edetate, disodium edetate, trisodium edetate, calcium disodium edetate, dipotassium edetate, and the like), citric acid and salts and hydrates thereof (*e.g.*, citric acid monohydrate), fumaric acid and salts and hydrates thereof, malic acid and salts and hydrates thereof, phosphoric acid and salts and hydrates thereof, and tartaric acid and salts and hydrates thereof. Exemplary antimicrobial preservatives include benzalkonium chloride, benzethonium chloride, benzyl alcohol, bronopol, cetrimide, cetylpyridinium chloride, chlorhexidine, chlorobutanol, chlorocresol, chloroxylenol, cresol, ethyl alcohol, glycerin, hexetidine, imidurea, phenol, phenoxyethanol, phenylethyl alcohol, phenylmercuric nitrate, propylene glycol, and thimerosal.

Exemplary antifungal preservatives include butyl paraben, methyl paraben, ethyl paraben, propyl paraben, benzoic acid, hydroxybenzoic acid, potassium benzoate, potassium sorbate, sodium benzoate, sodium propionate, and sorbic acid.

Exemplary alcohol preservatives include ethanol, polyethylene glycol, phenol, phenolic compounds, bisphenol, chlorobutanol, hydroxybenzoate, and phenylethyl alcohol.

Exemplary acidic preservatives include vitamin A, vitamin C, vitamin E, beta-carotene, citric acid, acetic acid, dehydroacetic acid, ascorbic acid, sorbic acid, and phytic acid.

Other preservatives include tocopherol, tocopherol acetate, deteroxime mesylate, cetrimide, butylated hydroxyanisol (BHA), butylated hydroxytoluene (BHT), ethylenediamine, sodium lauryl sulfate (SLS), sodium lauryl ether sulfate (SLES), sodium bisulfite, sodium metabisulfite, potassium sulfite, potassium metabisulfite, Glydant Plus, Phenonip, methylparaben, Germall 115, Germaben II, Neolone, Kathon, and Euxyl. In certain embodiments, the preservative is an anti-oxidant. In other embodiments, the preservative is a chelating agent.

Exemplary buffering agents include citrate buffer solutions, acetate buffer solutions, phosphate buffer solutions, ammonium chloride, calcium carbonate, calcium chloride, calcium citrate, calcium gluconate, calcium gluceptate, calcium gluconate, D-gluconic acid, calcium glycerophosphate, calcium lactate, propanoic acid, calcium levulinate, pentanoic acid, dibasic calcium phosphate, phosphoric acid, tribasic calcium phosphate, calcium hydroxide phosphate, potassium acetate, potassium chloride, potassium gluconate, potassium mixtures, dibasic potassium phosphate, monobasic potassium phosphate, potassium phosphate mixtures, sodium acetate, sodium bicarbonate, sodium chloride, sodium citrate, sodium lactate, dibasic sodium phosphate, monobasic sodium phosphate, sodium phosphate mixtures, tromethamine, magnesium hydroxide, aluminum hydroxide, alginic acid, pyrogen-free water, isotonic saline, Ringer's solution, ethyl alcohol, *etc.*, and combinations thereof.

Exemplary lubricating agents include magnesium stearate, calcium stearate, stearic acid, silica, talc, malt, glyceryl behenate, hydrogenated vegetable oils, polyethylene glycol, sodium benzoate, sodium acetate, sodium chloride, leucine, magnesium lauryl sulfate, sodium lauryl sulfate, *etc.*, and combinations thereof.

Exemplary oils include almond, apricot kernel, avocado, babassu, bergamot, black current seed, borage, cade, camomile, canola, caraway, carnauba, castor, cinnamon, cocoa butter, coconut, cod liver, coffee, corn, cotton seed, emu, eucalyptus, evening primrose, fish, flaxseed, geraniol, gourd, grape seed, hazel nut, hyssop, isopropyl myristate, jojoba, kukui nut, lavandin, lavender, lemon, litsea cubeba, macademia nut, mallow, mango seed, meadowfoam seed, mint, nutmeg, olive, orange, orange roughy, palm, palm kernel, peach kernel, peanut, poppy seed, pumpkin seed, rapeseed, rice bran, rosemary, safflower, sandalwood, sasquana, savoury, sea

buckthorn, sesame, shea butter, silicone, soybean, sunflower, tea tree, thistle, tsubaki, vetiver, walnut, and wheat germ oils. Exemplary oils include, but are not limited to, butyl stearate, caprylic triglyceride, capric triglyceride, cyclomethicone, diethyl sebacate, dimethicone 360, isopropyl myristate, mineral oil, octyldodecanol, oleyl alcohol, silicone oil, and combinations thereof.

Liquid dosage forms for oral and parenteral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredients, the liquid dosage forms can comprise inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (e.g., cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents. In certain embodiments for parenteral administration, the conjugates of the invention are mixed with solubilizing agents such as Cremophor, alcohols, oils, modified oils, glycols, polysorbates, cyclodextrins, polymers, and combinations thereof.

Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions can be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation can be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

In order to prolong the effect of a drug, it is often desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This can be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, can depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Compositions for rectal or vaginal administration are typically suppositories which can be prepared by mixing the conjugates of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active ingredient.

Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active ingredient is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form can comprise buffering agents.

Solid compositions of a similar type can be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They can optionally comprise opacifying agents and can be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples

of embedding compositions which can be used include polymeric substances and waxes. Solid compositions of a similar type can be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

5 The active ingredients can be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active ingredient can be admixed with at least one inert diluent such as sucrose, lactose or starch. Such
10 dosage forms can comprise, as is normal practice, additional substances other than inert diluents, *e.g.*, tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms can comprise buffering agents. They can optionally comprise opacifying agents and can be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract,
15 optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

Dosage forms for topical and/or transdermal administration of a compound of this invention can include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants and/or patches. Generally, the active ingredient is admixed under sterile conditions
20 with a pharmaceutically acceptable carrier and/or any needed preservatives and/or buffers as may be required. Additionally, the present invention contemplates the use of transdermal patches, which often have the added advantage of providing controlled delivery of an active ingredient to the body. Such dosage forms can be prepared, for example, by dissolving and/or dispensing the active ingredient in the proper medium. Alternatively or additionally, the rate can be controlled
25 by either providing a rate controlling membrane and/or by dispersing the active ingredient in a polymer matrix and/or gel.

Suitable devices for use in delivering intradermal pharmaceutically acceptable compositions described herein include short needle devices such as those described in U.S. Patents 4,886,499; 5,190,521; 5,328,483; 5,527,288; 4,270,537; 5,015,235; 5,141,496; and
30 5,417,662. Intradermal compositions can be administered by devices which limit the effective penetration length of a needle into the skin, such as those described in PCT publication WO

99/34850 and functional equivalents thereof. Jet injection devices which deliver liquid vaccines to the dermis via a liquid jet injector and/or via a needle which pierces the stratum corneum and produces a jet which reaches the dermis are suitable. Jet injection devices are described, for example, in U.S. Patents 5,480,381; 5,599,302; 5,334,144; 5,993,412; 5,649,912; 5,569,189; 5,704,911; 5,383,851; 5,893,397; 5,466,220; 5,339,163; 5,312,335; 5,503,627; 5,064,413; 5,520,639; 4,596,556; 4,790,824; 4,941,880; 4,940,460; and PCT publications WO 97/37705 and WO 97/13537. Ballistic powder/particle delivery devices which use compressed gas to accelerate vaccine in powder form through the outer layers of the skin to the dermis are suitable. Alternatively or additionally, conventional syringes can be used in the classical mantoux method of intradermal administration.

Formulations suitable for topical administration include, but are not limited to, liquid and/or semi liquid preparations such as liniments, lotions, oil in water and/or water in oil emulsions such as creams, ointments and/or pastes, and/or solutions and/or suspensions. Topically-administrable formulations can, for example, comprise from about 1% to about 10% (w/w) active ingredient, although the concentration of the active ingredient can be as high as the solubility limit of the active ingredient in the solvent. Formulations for topical administration can further comprise one or more of the additional ingredients described herein.

A pharmaceutically acceptable composition of the invention can be prepared, packaged, and/or sold in a formulation suitable for pulmonary administration via the buccal cavity. Such a formulation can comprise dry particles which comprise the active ingredient and which have a diameter in the range from about 0.5 to about 7 nanometers or from about 1 to about 6 nanometers. Such compositions are conveniently in the form of dry powders for administration using a device comprising a dry powder reservoir to which a stream of propellant can be directed to disperse the powder and/or using a self propelling solvent/powder dispensing container such as a device comprising the active ingredient dissolved and/or suspended in a low-boiling propellant in a sealed container. Such powders comprise particles wherein at least 98% of the particles by weight have a diameter greater than 0.5 nanometers and at least 95% of the particles by number have a diameter less than 7 nanometers. Alternatively, at least 95% of the particles by weight have a diameter greater than 1 nanometer and at least 90% of the particles by number have a diameter less than 6 nanometers. Dry powder compositions can include a solid fine powder diluent such as sugar and are conveniently provided in a unit dose form.

Low boiling propellants generally include liquid propellants having a boiling point of below 65 °F at atmospheric pressure. Generally the propellant can constitute 50 to 99.9% (w/w) of the composition, and the active ingredient can constitute 0.1 to 20% (w/w) of the composition. The propellant can further comprise additional ingredients such as a liquid non-ionic and/or solid anionic surfactant and/or a solid diluent (which can have a particle size of the same order as particles comprising the active ingredient).

Pharmaceutically acceptable compositions of the invention formulated for pulmonary delivery can provide the active ingredient in the form of droplets of a solution and/or suspension. Such formulations can be prepared, packaged, and/or sold as aqueous and/or dilute alcoholic solutions and/or suspensions, optionally sterile, comprising the active ingredient, and can conveniently be administered using any nebulization and/or atomization device. Such formulations can further comprise one or more additional ingredients including, but not limited to, a flavoring agent such as saccharin sodium, a volatile oil, a buffering agent, a surface active agent, and/or a preservative such as methylhydroxybenzoate. The droplets provided by this route of administration can have an average diameter in the range from about 0.1 to about 200 nanometers.

The formulations described herein as being useful for pulmonary delivery are useful for intranasal delivery of a pharmaceutically acceptable composition of the invention. Another formulation suitable for intranasal administration is a coarse powder comprising the active ingredient and having an average particle from about 0.2 to 500 micrometers. Such a formulation is administered by rapid inhalation through the nasal passage from a container of the powder held close to the nares.

Formulations suitable for nasal administration can, for example, comprise from about as little as 0.1% (w/w) and as much as 100% (w/w) of the active ingredient, and can comprise one or more of the additional ingredients described herein. A pharmaceutically acceptable composition of the invention can be prepared, packaged, and/or sold in a formulation suitable for buccal administration. Such formulations can, for example, be in the form of tablets and/or lozenges made using conventional methods, and can comprise, for example, 0.1 to 20% (w/w) active ingredient, the balance comprising an orally dissolvable and/or degradable composition and, optionally, one or more of the additional ingredients described herein. Alternately, formulations suitable for buccal administration can comprise a powder and/or an aerosolized

and/or atomized solution and/or suspension comprising the active ingredient. Such powdered, aerosolized, and/or aerosolized formulations, when dispersed, can have an average particle and/or droplet size in the range from about 0.1 to about 200 nanometers, and can further comprise one or more of the additional ingredients described herein.

5 A pharmaceutically acceptable composition of the invention can be prepared, packaged, and/or sold in a formulation suitable for ophthalmic administration. Such formulations can, for example, be in the form of eye drops including, for example, a 0.1/1.0% (w/w) solution and/or suspension of the active ingredient in an aqueous or oily liquid carrier. Such drops can further comprise buffering agents, salts, and/or one or more other of the additional ingredients described
10 herein. Other ophthalmically-administrable formulations which are useful include those which comprise the active ingredient in microcrystalline form and/or in a liposomal preparation. Ear drops and/or eye drops are contemplated as being within the scope of this invention.

Although the descriptions of pharmaceutically acceptable compositions provided herein are principally directed to pharmaceutical compositions which are suitable for administration to
15 humans, it will be understood by the skilled artisan that such compositions are generally suitable for administration to animals of all sorts. Modification of pharmaceutically acceptable compositions suitable for administration to humans in order to render the compositions suitable for administration to various animals is well understood, and the ordinarily skilled veterinary pharmacologist can design and/or perform such modification with ordinary experimentation.

20 Also provided are kits comprising one or more compounds of the invention (or pharmaceutically acceptable salts or prodrugs thereof), and/or one or more pharmaceutically acceptable compositions as described herein. Kits are typically provided in a suitable container (*e.g.*, for example, a foil, plastic, or cardboard package). In certain embodiments, a kit can include one or more pharmaceutical excipients, pharmaceutical additives, therapeutically active
25 agents, and the like, as described herein. In certain embodiments, a kit can include means for proper administration, such as, for example, graduated cups, syringes, needles, cleaning aids, and the like. In certain embodiments, a kit can include instructions for proper administration and/or preparation for proper administration.

30 ***Methods of Treatment***

The present invention also provides methods for treating an FAAH-mediated disorder by administering a therapeutically effective amount of a compound of any of formulae **I**, **Ia**, **Ib**, **II**, **IIa**, **IIb**, **III**, **IV**, **IVa**, **IVb**, **V**, **Va**, **Vb**, **VI**, **VIa**, **VIb**, **VII**, **VIIa**, **VIIb**, **VIII**, **VIIIa**, **VIIIb**, **IX**, **IXa**, **IXb**, **X**, **Xa**, **Xb**, **XI**, **XIa**, **XIb**, **XII**, **XIIa**, **XIIb**, **XIII**, **XIIIa**, **XIIIb**, **XIV**, **XIVa**, **XIVb**, **XV**, **XVa**, or **XVb**, or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, to a subject in need thereof.

Additionally, the present invention provides methods for inhibiting FAAH in a subject by administering a therapeutically effective amount of a compound of any of formulae **I**, **Ia**, **Ib**, **II**, **IIa**, **IIb**, **III**, **IV**, **IVa**, **IVb**, **V**, **Va**, **Vb**, **VI**, **VIa**, **VIb**, **VII**, **VIIa**, **VIIb**, **VIII**, **VIIIa**, **VIIIb**, **IX**, **IXa**, **IXb**, **X**, **Xa**, **Xb**, **XI**, **XIa**, **XIb**, **XII**, **XIIa**, **XIIb**, **XIII**, **XIIIa**, **XIIIb**, **XIV**, **XIVa**, **XIVb**, **XV**, **XVa**, or **XVb**, or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, to a subject in need thereof.

As used herein, and unless otherwise specified, the terms “treat,” “treating” and “treatment” contemplate an action that occurs while a subject is suffering from the specified disease, condition, or disorder, which reduces the severity of the disease or disorder, or retards or slows the progression of the disease or disorder. Treatment can be *via* prophylactic or therapeutic therapy.

As used herein, and unless otherwise specified, a “therapeutically effective amount” of a compound is an amount sufficient to provide a therapeutic benefit in the treatment or management of a disease, disorder, or condition, or to delay or minimize one or more symptoms associated with the disease, disorder, or condition. A therapeutically effective amount of a compound means an amount of therapeutic agent, alone or in combination with other therapies, which provides a therapeutic benefit in the treatment or management of the disease or condition. The term “therapeutically effective amount” can encompass an amount that improves overall therapy, reduces or avoids symptoms or causes of disease, disorder, or condition, or enhances the therapeutic efficacy of another therapeutic agent. The therapeutically effective amount in a subject will vary depending on the compound, the disease and its severity, and the age, weight, etc., of the subject to be treated. In some embodiments, a “therapeutically effective amount” can encompass a “prophylactically effective amount.”

As used herein, and unless otherwise specified, a “prophylactically effective amount” of a compound is an amount sufficient to prevent a disease, disorder, or condition, or one or more

symptoms associated with the disease, disorder, or condition, or prevent its recurrence. A prophylactically effective amount of a compound means an amount of therapeutic agent, alone or in combination with other agents, which provides a prophylactic benefit in the prevention of the disease. The term “prophylactically effective amount” can encompass an amount that improves overall prophylaxis or enhances the prophylactic efficacy of another prophylactic agent.

As used herein, and unless otherwise specified, the terms “manage,” “managing” and “management” encompass preventing the recurrence of the specified disease, disorder, or condition in a subject who has already suffered from the disease or disorder, and/or lengthening the time that a subject who has suffered from the disease, disorder, or condition remains in remission. The terms encompass modulating the threshold, development and/or duration of the disease, disorder or condition, or changing the way that a subject responds to the disease, disorder, or condition.

The term “subject” is defined herein to include animals such as mammals, including, but not limited to, primates (*e.g.*, humans (*e.g.*, male, female, infant, child, adolescent, adult, elderly, *etc.*)), cows, sheep, goats, horses, dogs, cats, birds, rabbits, rats, mice and the like. In preferred embodiments, the subject is a human.

In other embodiments, the present invention provides a method for inhibiting FAAH in a biological sample comprising the step of contacting said sample with a compound of any of formulae **I**, **Ia**, **Ib**, **II**, **IIa**, **IIb**, **III**, **IV**, **IVa**, **IVb**, **V**, **Va**, **Vb**, **VI**, **VIa**, **VIb**, **VII**, **VIIa**, **VIIb**, **VIII**, **VIIIa**, **VIIIb**, **IX**, **IXa**, **IXb**, **X**, **Xa**, **Xb**, **XI**, **XIa**, **XIb**, **XII**, **XIIa**, **XIIb**, **XIII**, **XIIIa**, **XIIIb**, **XIV**, **XIVa**, **XIVb**, **XV**, **XVa**, or **XVb**, or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

The phrases “FAAH-mediated conditions,” “FAAH-mediated diseases,” and “FAAH-mediated disorders,” as used interchangeably, include, but are not limited to, painful conditions, painful diseases or painful disorders, inflammatory disorders, immune disorders, depression, anxiety, anxiety-related disorders, sleep disorders, feeding behaviors, movement disorders, glaucoma, neuroprotection and cardiovascular disease. The terms “disease,” “disorder,” and “condition” are used interchangeably.

In certain embodiments, the FAAH-mediated disorder is a painful disorder. As used herein, a “painful disorder” includes, but is not limited to, neuropathic pain (*e.g.*, peripheral neuropathic pain), central pain, deafferentiation pain, chronic pain (*e.g.*, chronic nociceptive

pain, and other forms of chronic pain such as post-operative pain), stimulus of nociceptive receptors, acute pain (*e.g.*, phantom and transient acute pain), non-inflammatory pain, inflammatory pain, pain associated with cancer, wound pain, burn pain, post-operative pain, pain associated with medical procedures, arthritic pain (*e.g.*, pain associated with rheumatoid arthritis, osteoarthritis), lumbosacral pain, musculo-skeletal pain, headache, migraine, muscle ache, lower back and neck pain, toothache and the like.

In certain embodiments, the painful disorder is neuropathic pain. The term “neuropathic pain” refers to pain resulting from injury to a nerve. Neuropathic pain is distinguished from nociceptive pain, which is the pain caused by acute tissue injury involving small cutaneous nerves or small nerves in muscle or connective tissue. Neuropathic pain typically is long-lasting or chronic and often develops days or months following an initial acute tissue injury. Neuropathic pain can involve persistent, spontaneous pain as well as allodynia, which is a painful response to a stimulus that normally is not painful. Neuropathic pain also can be characterized by hyperalgesia, in which there is an accentuated response to a painful stimulus that usually is trivial, such as a pin prick. Neuropathic pain conditions can develop following neuronal injury and the resulting pain may persist for months or years, even after the original injury has healed. Neuronal injury may occur in the peripheral nerves, dorsal roots, spinal cord or certain regions in the brain. Neuropathic pain conditions include: diabetic neuropathy; sciatica; non-specific lower back pain; multiple sclerosis pain; fibromyalgia; HIV-related neuropathy; neuralgia, such as post-herpetic neuralgia and trigeminal neuralgia; and pain resulting from physical trauma, amputation, cancer, chemotherapy-induced pain, chemotherapy, surgery, invasive medical procedures, toxins burns, infection, or chronic inflammatory conditions. Neuropathic pain can result from a peripheral nerve disorder such as neuroma; nerve compression; nerve crush, nerve stretch or incomplete nerve transection; mononeuropathy or polyneuropathy. Neuropathic pain can also result from a disorder such as dorsal root ganglion compression; inflammation of the spinal cord; contusion, tumor or hemisection of the spinal cord; tumors of the brainstem, thalamus or cortex; or trauma to the brainstem, thalamus or cortex.

The symptoms of neuropathic pain are heterogeneous and are often described as spontaneous shooting and lancinating pain, or ongoing, burning pain. In addition, there is pain associated with normally non-painful sensations such as “pins and needles” (paraesthesias and

dysesthesias), increased sensitivity to touch (hyperesthesia), painful sensation following innocuous stimulation (dynamic, static or thermal allodynia), increased sensitivity to noxious stimuli (thermal, cold, mechanical hyperalgesia), continuing pain sensation after removal of the stimulation (hyperpathia) or an absence of or deficit in selective sensory pathways (hypoalgesia).

5 In certain embodiments, the painful disorder is non-inflammatory pain and/or inflammatory pain. The types of non-inflammatory pain include, without limitation, peripheral neuropathic pain (*e.g.*, pain caused by a lesion or dysfunction in the peripheral nervous system), central pain (*e.g.*, pain caused by a lesion or dysfunction of the central nervous system), deafferentation pain (*e.g.*, pain due to loss of sensory input to the central nervous system),
10 chronic nociceptive pain (*e.g.*, certain types of cancer pain), noxious stimulus of nociceptive receptors (*e.g.*, pain felt in response to tissue damage or impending tissue damage), phantom pain (*e.g.*, pain felt in a part of the body that no longer exists, such as a limb that has been amputated), pain felt by psychiatric patients (*e.g.*, pain where no physical cause may exist), and wandering pain (*e.g.*, wherein the pain repeatedly changes location in the body). In certain embodiments,
15 non-inflammatory pain and/or inflammatory pain are associated with disorders such as inflammatory diseases (*e.g.*, autoimmune disease).

In certain embodiments, the FAAH-mediated disorder is an inflammatory disorder. The term "inflammatory disorders" refers to those diseases or conditions that are characterized by signs of pain (dolor, from the generation of noxious substances and the stimulation of nerves),
20 heat (calor, from vasodilatation), redness (rubor, from vasodilatation and increased blood flow), swelling (tumor, from excessive inflow or restricted outflow of fluid), and/or loss of function (functio laesa, which may be partial or complete, temporary or permanent). Inflammatory disorders include, without limitation, those affecting the blood vessels (*e.g.*, polyarteritis, temporal arteritis); joints (*e.g.*, arthritis: crystalline, osteo-, psoriatic, reactive, rheumatoid,
25 Reiter's syndrome); gastrointestinal tract (*e.g.*, Crohn's disease, ulcerative colitis); skin (*e.g.*, dermatitis); or multiple organs and tissues (*e.g.*, systemic lupus erythematosus). Inflammatory disorders include, but are not limited to, inflammation associated with vascular diseases, migraine headaches, tension headaches, arteritis, thyroiditis, aplastic anemia, Hodgkin's disease, scleroderma, rheumatic fever, type I diabetes, myasthenia gravis, sarcoidosis, nephrotic
30 syndrome, Behcet's syndrome, polymyositis, gingivitis, hypersensitivity, conjunctivitis, multiple sclerosis, and ischemia (*e.g.*, myocardial ischemia), and the like. The compounds and

compositions can be useful for treating neuroinflammation associated with brain disorders (*e.g.*, Parkinson's disease and Alzheimer's disease) and chronic inflammation associated with cranial radiation injury. The compounds can be useful for treating acute inflammatory conditions (*e.g.*, conditions resulting from infection) and chronic inflammatory conditions (*e.g.*, conditions
5 resulting from asthma, arthritis and inflammatory bowel disease). The compounds can also be useful in treating inflammation associated with trauma and non-inflammatory myalgia. Inflammation takes on many forms and includes, but is not limited to, acute, adhesive, atrophic, catarrhal, chronic, cirrhotic, diffuse, disseminated, exudative, fibrinous, fibrosing, focal, granulomatous, hyperplastic, hypertrophic, interstitial, metastatic, necrotic, obliterative,
10 parenchymatous, plastic, productive, proliferous, pseudomembranous, purulent, sclerosing, seroplastic, serous, simple, specific, subacute, suppurative, toxic, traumatic, and/or ulcerative inflammation.

In certain embodiments, the FAAH-mediated disorder is an immune disorder. Immune disorders, such as auto-immune disorders, include, but are not limited to, arthritis (including
15 rheumatoid arthritis, spondyloarthropathies, gouty arthritis, degenerative joint diseases such as osteoarthritis, systemic lupus erythematosus, Sjogren's syndrome, ankylosing spondylitis, undifferentiated spondylitis, Behcet's disease, haemolytic autoimmune anaemias, multiple sclerosis, amyotrophic lateral sclerosis, amylosis, acute painful shoulder, psoriatic, and juvenile arthritis), asthma, atherosclerosis, osteoporosis, bronchitis, tendonitis, bursitis, skin inflammation
20 disorders (*e.g.*, psoriasis, eczema, burns, dermatitis), enuresis, eosinophilic disease, gastrointestinal disorders (*e.g.*, inflammatory bowel disease (IBD), peptic ulcers, regional enteritis, diverticulitis, gastrointestinal bleeding, Crohn's disease, gastritis, diarrhoea, irritable bowel syndrome and ulcerative colitis), and disorders ameliorated by a gastropromotory agent (*e.g.*, ileus, postoperative ileus and ileus during sepsis; gastroesophageal reflux disease (GORD, or its synonym GERD); eosinophilic esophagitis, gastroparesis such as diabetic gastroparesis;
25 food intolerances and food allergies and other functional bowel disorders, such as non-ulcerative dyspepsia (NUD) and non-cardiac chest pain (NCCP, including costo-chondritis)).

In certain embodiments, the immune disorder is a gastrointestinal disorder. In some
embodiments, the immune disorder is inflammatory bowel disease (*e.g.*, Crohn's disease and/or
30 ulcerative colitis), peptic ulcers, regional enteritis, diverticulitis, gastrointestinal bleeding,

Crohn's disease, gastritis, diarrhea, irritable bowel syndrome and ulcerative colitis. In other embodiments, the immune disorder is inflammatory bowel disease (IBD).

In certain embodiments, the FAAH-mediated disorder is a skin disorder. In some embodiments, the skin disorder is pruritus (itch), psoriasis, eczema, burns or dermatitis. In certain embodiments, the skin disorder is psoriasis. In certain embodiments, the skin disorder is pruritis.

In certain embodiments, the FAAH-mediated disorder is anxiety. "Anxiety," as used herein, includes, but is not limited to anxiety and anxiety disorders or conditions, such as, for example, clinical anxiety, panic disorder, agoraphobia, generalized anxiety disorder, specific phobia, social phobia, obsessive-compulsive disorder, acute stress disorder, and post-traumatic stress disorder; and adjustment disorders with anxious features, anxiety disorders associated with depression, anxiety disorders due to general medical conditions, and substance-induced anxiety disorders. This treatment can also be to induce or promote sleep in a patient (*e.g.*, for example, a patient with anxiety).

In certain embodiments, the FAAH-mediated disorder is a sleep disorder. "Sleep disorders" include, but are not limited to, insomnia, sleep apnea, restless legs syndrome (RLS), delayed sleep phase syndrome (DSPS), periodic limb movement disorder (PLMD), hypopnea syndrome, rapid eye movement behavior disorder (RBD), shift work sleep disorder (SWSD), and sleep problems (*e.g.*, parasomnias) such as nightmares, night terrors, sleep talking, head banging, snoring, and clenched jaw and/or grinding of teeth (bruxism).

In certain embodiments, the FAAH-mediated disorder is depression. "Depression," as used herein, includes, but is not limited to, depressive disorders or conditions, such as, for example, major depressive disorders (unipolar depression), dysthymic disorders (chronic, mild depression) and bipolar disorders (manic-depression). The depression can be clinical or subclinical depression.

In certain embodiments, the FAAH-mediated disorder is feeding behavior. "Feeding behavior," as used herein, includes but is not limited to, eating disorders (*e.g.*, anorexias and cachexias of various natures, over-eating leading to obesity), weight loss associated with cancer, weight loss associated with other general medical conditions, weight loss associated with failure to thrive, and other wasting conditions. The compounds disclosed herein can also be used to reduce body fat and for treating or preventing obesity in a mammal. The compounds disclosed

herein can also be used for preventing or treating the diseases associated with these health conditions.

In certain embodiments, the FAAH-mediated disorder is a movement disorder. In other embodiments, the FAAH-mediated disorder is glaucoma. In yet other embodiments, the
5 FAAH-mediated disorder is neuroprotection. In still yet other embodiments, the FAAH-mediated disorder is cardiovascular disease.

Administration

Provided compounds can be administered using any amount and any route of
10 administration effective for treatment. The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the infection, the particular composition, its mode of administration, its mode of activity, and the like.

Compounds provided herein are typically formulated in dosage unit form for ease of
15 administration and uniformity of dosage. It will be understood, however, that the total daily usage of the compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular subject or organism will depend upon a variety of factors including the disease, disorder, or condition being treated and the severity of the disorder; the activity of the specific
20 active ingredient employed; the specific composition employed; the age, body weight, general health, sex and diet of the subject; the time of administration, route of administration, and rate of excretion of the specific active ingredient employed; the duration of the treatment; drugs used in combination or coincidental with the specific active ingredient employed; and like factors well known in the medical arts.

The compounds and compositions provided herein can be administered by any route,
25 including oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, subcutaneous, intraventricular, transdermal, interdermal, rectal, intravaginal, intraperitoneal, topical (as by powders, ointments, creams, and/or drops), mucosal, nasal, bucal, enteral, sublingual; by intratracheal instillation, bronchial instillation, and/or inhalation; and/or as an oral
30 spray, nasal spray, and/or aerosol. Specifically contemplated routes are systemic intravenous injection, regional administration via blood and/or lymph supply, and/or direct administration to

an affected site. In general the most appropriate route of administration will depend upon a variety of factors including the nature of the agent (*e.g.*, its stability in the environment of the gastrointestinal tract), the condition of the subject (*e.g.*, whether the subject is able to tolerate oral administration), *etc.*

5 The exact amount of a compound required to achieve a therapeutically effective amount will vary from subject to subject, depending, for example, on species, age, and general condition of a subject, severity of the side effects or disorder, identity of the particular compound(s), mode of administration, and the like. The desired dosage can be delivered three times a day, two times a day, once a day, every other day, every third day, every week, every two weeks, every three
10 weeks, or every four weeks. In certain embodiments, the desired dosage can be delivered using multiple administrations (*e.g.*, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, or more administrations).

 In certain embodiments, a therapeutically effective amount of a compound for administration one or more times a day to a 70 kg adult human can comprise about 0.0001 mg to
15 about 1000 mg of an inventive compound per unit dosage form. For example, a therapeutically effective amount of a compound of the present invention can comprise about 0.01 mg, about 0.5 mg, about 1 mg, about 2 mg, about 3 mg, about 5 mg, about 10 mg, about 25 mg, about 50 mg, about 70 mg, about 100 mg, about 200 mg, about 300 mg, about 400 mg, about 500 mg, about 600 mg, about 700 mg, about 800 mg, about 900 mg, or about 1000 mg depending on the
20 compound, the disease and its severity, and the age, weight, *etc.*, of the subject to be treated. It will be appreciated that dose ranges as described herein provide guidance for the administration of provided pharmaceutically acceptable compositions to an adult. The amount to be administered to, for example, a child or an adolescent can be determined by a medical practitioner or person skilled in the art and can be lower or the same as that administered to an
25 adult.

 It will be also appreciated that a compound or composition, as described herein, can be administered in combination with one or more additional therapeutically active agents. The compound or composition can be administered concurrently with, prior to, or subsequent to, one or more additional therapeutically active agents. In general, each agent will be administered at a
30 dose and/or on a time schedule determined for that agent. It will further be appreciated that the additional therapeutically active agent utilized in this combination can be administered together

in a single composition or administered separately in different compositions. The particular combination to employ in a regimen will take into account compatibility of the inventive compound with the additional therapeutically active agent and/or the desired therapeutic effect to be achieved. In general, it is expected that additional therapeutically active agents utilized in combination be utilized at levels that do not exceed the levels at which they are utilized individually. In some embodiments, the levels utilized in combination will be lower than those utilized individually.

The compounds or compositions can be administered in combination with agents that improve their bioavailability, reduce and/or modify their metabolism, inhibit their excretion, and/or modify their distribution within the body. It will also be appreciated that therapy employed can achieve a desired effect for the same disorder (for example, a compound can be administered in combination with an anti-inflammatory, anti-anxiety and/or anti-depressive agent, *etc.*), and/or it can achieve different effects (*e.g.*, control of adverse side-effects).

Exemplary active agents include, but are not limited to, anti-cancer agents, antibiotics, anti-viral agents, anesthetics, anti-coagulants, inhibitors of an enzyme, steroidal agents, steroidal or non-steroidal anti-inflammatory agents, antihistamine, immunosuppressant agents, anti-neoplastic agents, antigens, vaccines, antibodies, decongestants, sedatives, opioids, pain-relieving agents, analgesics, anti-pyretics, hormones, prostaglandins, progestational agents, anti-glaucoma agents, ophthalmic agents, anti-cholinergics, anti-depressants, anti-psychotics, hypnotics, tranquilizers, anti-convulsants, muscle relaxants, anti-spasmodics, muscle contractants, channel blockers, miotic agents, anti-secretory agents, anti-thrombotic agents, anticoagulants, anti-cholinergics, β -adrenergic blocking agents, diuretics, cardiovascular active agents, vasoactive agents, vasodilating agents, anti-hypertensive agents, angiogenic agents, modulators of cell-extracellular matrix interactions (*e.g.* cell growth inhibitors and anti-adhesion molecules), or inhibitors/intercalators of DNA, RNA, protein-protein interactions, protein-receptor interactions, *etc.* Active agents include small organic molecules such as drug compounds (*e.g.*, compounds approved by the Food and Drugs Administration as provided in the Code of Federal Regulations (CFR)), peptides, proteins, carbohydrates, monosaccharides, oligosaccharides, polysaccharides, nucleoproteins, mucoproteins, lipoproteins, synthetic polypeptides or proteins, small molecules linked to proteins, glycoproteins, steroids, nucleic

acids, DNAs, RNAs, nucleotides, nucleosides, oligonucleotides, antisense oligonucleotides, lipids, hormones, vitamins and cells.

In certain embodiments, the additional therapeutically active agent is a pain-relieving agent. In other embodiments, the additional therapeutically active agent is an anti-inflammatory agent.

Methods of Determining Biological Activity

Methods of determining the activity of the compounds provided herein for various therapeutic uses are known in the art. These include, but are not limited to, high throughput screening to identify compounds that bind to and/or modulate the activity of isolated FAAH, as well as *in vitro* and *in vivo* models of therapies.

Assays useful for screening the compounds provided herein can detect the binding of the inhibitor to FAAH or the release of a reaction product (*e.g.*, fatty acid amide or ethanolamine) produced by the hydrolysis of a substrate such as oleoylethanolamide or ananadamide. The substrate can be labeled to facilitate detection of the released reaction products. U.S. Pat. No. 5,559,410 discloses high throughput screening methods for proteins, and U.S. Pat. Nos. 5,576,220 and 5,541,061 disclose high throughput methods of screening for ligand/antibody binding.

Methods for screening FAAH inhibitors for an antinociceptive effect are known in the art. For example, compounds can be tested in the mouse hot-plate test and the mouse formalin test, and the nociceptive reactions to thermal or chemical tissue damage measured (for example, see U.S. Pat. No. 6,326,156 for a description of methods of screening for antinociceptive activity; see also Cravatt *et al.* *Proc. Natl. Acad. Sci. U.S.A.* (2001) 98:9371–9376).

Two pharmacologically validated animal models of anxiety are the elevated zero maze test, and the isolation-induced ultrasonic emission test. The zero maze consists of an elevated annular platform with two open and two closed quadrants and is based on the conflict between an animal's instinct to explore its environment and its fear of open spaces (see, for example, Bickerdike, M. J. *et al.*, *Eur. J. Pharmacol.*, (1994) 271, 403–411; Shepherd, J. K. *et al.*, *Psychopharmacology*, (1994) 116, 56–64). Clinically used anxiolytic drugs, such as the benzodiazepines, increase the proportion of time spent in, and the number of entries made into, the open compartments.

A second test for an anti-anxiety compound is the ultrasonic vocalization emission model, which measures the number of stress-induced vocalizations emitted by rat pups removed from their nest (see, for example, Insel, T. R. et al., *Pharmacol. Biochem. Behav.*, 24, 1263–1267 (1986); Miczek, K. A. et al., *Psychopharmacology*, 121, 38–56 (1995); Winslow, J. T. et al.,
5 *Biol. Psychiatry*, 15, 745–757 (1991).

The effect of the compounds provided herein in the treatment of depression can be tested in the model of chronic mild stress induced anhedonia in rats. This model is based on the observation that chronic mild stress causes a gradual decrease in sensitivity to rewards, for example consumption of sucrose, and that this decrease is dose-dependently reversed by chronic
10 treatment with antidepressants. See, e.g., Willner, Paul, *Psychopharmacology*, 1997, 134, 319–329.

Another test for antidepressant activity is the forced swimming test (*Nature* 266, 730–732, 1977). In this test, animals are administered an agent 30 or 60 minutes before being placed in container of water, and the time during which they remain immobile is recorded. A decrease
15 in the immobility time of the mice is indicative of antidepressant activity.

A similar test for antidepressant activity is the mouse caudal suspension test (*Psychopharmacology*, 85, 367–370, 1985). In this test, animals are administered an agent 30 or 60 minutes before being suspended by the tail, and their immobility time is recorded. A decrease in the immobility time of the mice is indicative of antidepressant activity.

20 Animal models are available for assessing anticonvulsant activity of test compounds (see, e.g., U.S. Pat. Nos. 6,309,406 and 6,326,156).

Inhibition of FAAH has been reported to induce sleep in test animals (see, e.g., U.S. Pat. No. 6,096,784). Methods for studying sleep inducing compounds are known in the art (see, e.g., U.S. Pat. Nos. 6,096,784 and 6,271,015). Compounds can be administered to a test animal (e.g.,
25 rat or mouse) or a human and the subsequent time (e.g., onset, duration) spent sleeping (e.g., eyes closed, motor quiescence) can be monitored. See also WO 98/24396.

Methods for screening FAAH inhibitors which induce catalepsy are also well known in the art (see, e.g., Quistand et al. in *Toxicology and Applied Pharmacology* 173: 48–55 (2001); Cravatt et al. *Proc. Natl. Acad. Sci. U.S.A.* 98:9371–9376 (2001)).

30 Methods of assessing appetitive behavior are known in the art (see, e.g., U.S. Pat. No. 6,344,474). One method of assessing the effect on appetite behavior is to administer a FAAH

inhibitor to a rat and assess its effect on the intake of a sucrose solution (see, e.g., W. C. Lynch et al., *Physiol. Behav.*, 1993, 54, 877–880).

Methods of Synthesis

5 The reaction of an organometallic species with an organic borate, such as trimethyl borate, can be used to synthesize boronate esters. Suitable organometallic species include, but are not limited to, alkyl lithium and Grignard reagents. Other methods for the synthesis of boronates are employed when the boronate contains sensitive functionality that can not tolerate alkyl lithium reagents or Grignard reagents. These methods include palladium coupling
10 reactions of aryl or akenyl halides and diboronates or dialkoxo boranes and hydroboration of alkenes or alkynes. Using these methods a diverse collection of boronates can be synthesized. Boronates can be readily transformed in to boronic acids by hydrolyzing the boronate under aqueous acidic conditions using a suitable acid. Suitable acids include, but are not limited to HCl, H₂SO₄, and HBr. Another method of hydrolyzing boronates is an oxidative hydrolysis
15 employing an oxidizing agent, such as NaIO₄. The boronic acid compounds of the present invention readily form boronic esters when exposed to alcohols. The resulting boronic esters can also be used in the methods provided herein. Cyclic boronates are formed when certain diols (e.g., 1,2– and 1,3–diols) are used. Boronic acid compounds provided herein readily form oligomeric anhydrides by dehydration of the boronic acid moiety to form dimers, trimers, and
20 tetramers, and mixtures thereof. These species in the presence of water and under physiological conditions convert back to the boronic acid by hydrolysis.


EXEMPLIFICATION

The invention now being generally described, it will be more readily understood by reference to the following examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present invention, and are not intended to limit the invention.


General Synthetic Methods

The following is a description of general synthetic routes that can be used to prepare compounds of the present invention. Additionally, one of skill in the art will recognize that protecting groups may be necessary for the preparation of certain compounds and will be aware of those conditions compatible with a selected protecting group. Examples of such protecting groups include, for example, those set forth in *Protective Groups in Organic Synthesis*, Greene, T. W.; Wuts, P. G. M., John Wiley & Sons, New York, N.Y., (3rd Edition, 1999). Accordingly, the exemplary methods and the examples described herein are illustrative of the present invention and are not to be construed as limiting the scope thereof.

Method 1:




General conditions for the preparation of boronic acid pinacol esters: A dry flask under argon atmosphere is charged with aryl bromide (1.0 equiv), 1,1"-Bis(diphenylphosphino)-ferrocenedichloropalladium(II) (0.05 equiv), potassium acetate (1.0 equiv), cesium carbonate (3.0 equiv), and bis(pinacolato)diboron (2.0 equiv). The mixture is suspended with dimethylsulfoxide (0.1 M with respect to aryl bromide) and heated at 80 °C for 2-8 hours. Upon completion as judged by thin layer chromatography analysis, the reaction is split between water and ethyl acetate, and the organic layer is washed with brine, dried over sodium sulfate, and concentrated *in vacuo*. The concentrated reaction mixture is purified by flash silica gel chromatography (ethyl acetate / hexanes) to provide boronic acid pinacol ester.

Method 2:

General conditions for the conversion of boronate esters to boronic acids: The boronate ester (1.0 eq), sodium periodate (5.0 eq) and ammonium acetate (5.0 eq) are dissolved in acetone/water 2:1 (0.05 M boronate ester) and stirred for 12 h at 23 °C until TLC or LCMS indicates conversion to the boronic acid is complete. One option for isolation is to precipitate the product by dilution of the mixture with 1N aqueous HCl and collection of the solid boronic acid by filtration. Alternately, the mixture is partitioned between water and ethyl acetate, and the organic layer is washed with brine, dried over sodium sulfate, and concentrated *in vacuo*. The residue is purified either by recrystallization and trituration (heptane, acetonitrile, or other solvents) or by flash silica gel chromatography (e.g., using 0.5% to 10% methanol/dichloromethane) to afford pure boronic acid.


Examples

Exemplary compounds are set forth in the Examples provided below. Compounds were assayed as inhibitors of human FAAH using the method described in detail in Example 16. Activity designated as “A” refers to compounds having a K_i of less than or equal to 0.01 μM , “B” refers to compounds having a K_i of between 0.01 μM and 0.1 μM , “C” refers to compounds having a K_i of between 0.1 μM and 1 μM , and “D” refers to compounds having a K_i of greater than 1 μM .

Example 1:


6-Bromo-1-hydroxyisoquinoline (140 mg, 0.63 mmol, 1.0 equiv), tetrabutylammonium bromide (30 mg, 0.09 mmol, 0.15 equiv), and iodomethane (50 μ L, 0.75 mmol, 1.2 equiv) were combined in toluene (10 mL) and treated with 50% aq. NaOH (2 mL), and the resulting was mixture stirred rapidly at ambient temperature. After an additional 14h at room temperature, the mixture was diluted with diethyl ether and washed with water, saturated NaHCO₃, and brine. The organic phase was dried over Na₂SO₄ and concentrated to give a white solid of 6-bromo-2-methylisoquinolin-1-one (104 mg, 0.44 mmol, 70%). This crude material was taken forward without further purification and converted, via Methods 1 and 2, to compound **1** (45 mg, 38 %) and isolated as a beige solid. [M-H]⁻ = 202.1 m/z. Activity: B

Example 2:




2

6-Bromo-1-hydroxyisoquinoline (150 mg, 0.67 mmol, 1.0 equiv), tetrabutylammonium bromide (25 mg, 0.07 mmol, 0.1 equiv), and benzyl bromide (95 μ L, 0.8 mmol, 1.2 equiv) were combined in toluene (7 mL), and treated with 50% aq. NaOH (2 mL) after which point the resulting mixture was stirred rapidly at 90 °C. After 80 min, the mixture was diluted with MTBE and washed with water and brine. The organic phase was dried over Na₂SO₄. Addition of silica gel, concentration, and purification of the residue by flash silica gel chromatography (gradient of 2-12% ethyl acetate/hexanes) provided 2-benzyl-6-bromo-isoquinolin-1-one (221 mg, 0.67 mmol, quantitative) which was then converted, via Methods 1 and 2, to compound **2** (91 mg, 49 %) and isolated as a light-brown solid. [M-H]⁻ = 278.1 m/z. Activity: B


Example 3:

6-Bromo-1-hydroxyisoquinoline (300 mg, 1.34 mmol, 1.0 equiv), tetrabutylammonium bromide (50 mg, 0.13 mmol, 0.10 equiv), and phenethyl bromide (220 μ L, 1.6 mmol, 1.2 eq) were combined in toluene (14 mL), treated with 50% aq. NaOH (3 mL) after which the resulting mixture stirred rapidly at ambient temperature. After stirring for an additional 14 hours, the mixture was diluted with MTBE and washed with water, saturated NaHCO₃, and brine. The organic phase was dried over Na₂SO₄. Addition of silica gel, concentration, and purification of the residue using flash silica gel chromatography (gradient of 3-15% ethyl acetate/hexanes) provided 6-bromo-2-phenethyl-isoquinolin-1-one (225 mg, 0.69 mmol, 51%) as a white solid. This isoquinolinone (225 mg, 0.69 mmol) was then converted, via Methods 1 and 2, to compound **3** (95 mg, 47%) and isolated as a white solid. [M-H]⁻ = 292.1 m/z. Activity: A


Example 4:

6-Bromo-1-hydroxyisoquinoline (150 mg, 0.67 mmol, 1.0 equiv), tetrabutylammonium bromide (25 mg, 0.07 mmol, 0.1 equiv), and 1-bromoethylbenzene (160 μ L, 0.8 mmol, 1.2 equiv) were combined in toluene (7 mL) and treated with 50% aq. NaOH (2 mL) and the resulting mixture was stirred rapidly while heating to 70 °C. After additional stirring for 14 hours, the mixture was diluted with diethyl ether and washed with water and brine. The organic phase was dried over Na₂SO₄. Addition of silica gel, concentration, and purification of the residue using flash silica gel chromatography (gradient 2-12% ethyl acetate/hexanes) gave 6-bromo-2-(1-phenylethyl)-isoquinolin-1-one (227 mg, 0.67 mmol, quant.) as a yellowish oil. This

isoquinolinone (220 mg, 0.67 mmol) was converted, via Methods 1 and 2, to compound **4** (109 mg, 56%) and isolated as a white solid. $[M-H]^- = 292.1$ m/z. Activity: B


Example 5:

5
6-Bromo-1-hydroxyisoquinoline (150 mg, 0.67 mmol, 1.0 equiv), tetrabutylammonium bromide (25 mg, 0.07 mmol, 0.1 equiv), and 1-bromo-2-phenylpropane (160 μ L, 0.8 mmol, 1.2 equiv) were combined in toluene (10 mL) and treated with 50% aq. NaOH (2 mL), and the resulting mixture was stirred rapidly at ambient temperature for 48 hours, then heated at 60 °C
10 for 24h. The mixture was diluted with diethyl ether and washed with water and brine. The organic phase was dried over Na_2SO_4 . Concentration and purification of the residue using silica gel flash chromatography (2-12% EtOAc/hex.) gave white crystals of 6-bromo-2-(2-phenylpropyl)-isoquinolin-1-one (169 mg, 0.49 mmol, 74%). This isoquinolinone (169 mg, 0.49 mmol) was converted, via Methods 1 and 2, to compound **5** (65 mg, 43%) which was isolated as
15 white solid. $[M-H]^- = 306.1$ m/z. Activity: B


Example 6:

6-Bromo-1-hydroxyisoquinoline (150 mg, 0.67 mmol, 1.0 equiv), tetrabutylammonium bromide (25 mg, 0.07 mmol, 0.1 equiv), and 1-bromo-2-phenylpropane (160 μ L, 0.8 mmol, 1.2 equiv) were combined in toluene (10 mL) and treated with 50% aq. NaOH (2 mL), and the resulting mixture was stirred rapidly and heated at 70 °C. After heating for an additional 14 hours, the mixture was diluted with diethyl ether and washed with water and brine. The organic phase was dried over Na_2SO_4 and concentrated. Purification of the residue using flash silica gel
20

chromatography (gradient of 2-15% ethyl acetate/hexanes) gave 6-bromo-2-(1-phenyl-2-propyl)-isoquinolin-1-one (74 mg, 0.22 mmol, 32%) as a colorless oil. This isoquinolinone (74 mg, 0.22 mmol) was converted, via Methods 1 and 2, to a 1:1 mixture of compound **6** and 6-hydroxy-2-(1-phenyl-2-propyl)-isoquinolin-1-one that was isolated as 44 mg of a white solid (33% of **6**). [M-H]⁻ = 306.1 m/z. Activity: B


Example 7:

6-Bromo-1-hydroxyisoquinoline (200 mg, 0.89 mmol, 1.0 equiv), tetrabutylammonium bromide (30 mg, 0.09 mmol, 0.1 equiv), and hexyl bromide (150 μ L, 1.1 mmol, 1.2 equiv) were combined in toluene (10 mL) and treated with 50% aq. NaOH (2 mL), and the resulting mixture was stirred rapidly at ambient temperature. After an additional 14 hours of stirring at room temperature, the mixture was diluted with MTBE and washed with water, sat. NaHCO₃, and brine. The organic phase was dried over Na₂SO₄. Addition of silica gel, concentration, and purification of the residue using flash silica gel chromatography (gradient of 3-12% ethyl acetate/hexanes) gave 6-bromo-2-(2-phenylpropyl)-isoquinolin-1-one (87 mg, 0.28 mmol, 32%) as white crystals. This isoquinolinone (87 mg, 0.28 mmol) was converted, via Methods 1 and 2, to compound **7** (27 mg, 35%) which was isolated as a white solid. [M-H]⁻ = 372.2 m/z. Activity: A


Example 8:

7-Bromo-2-methyl-1H-quinoxaline-4-one (200 mg, 0.84 mmol, 1.0 equiv), tetrabutylammonium bromide (30 mg, 0.09 mmol, 0.1 equiv), and iodomethane (350 μ L, 5.5

mmol, 7.0 equiv) were combined in toluene (10 mL) and treated with 50% aq. NaOH (2 mL), and the resulting mixture was stirred rapidly at ambient temperature for 24 hours, after which it was heated to 35 °C for an additional 16 hours. At this point, the mixture was diluted with diethyl ether and washed with water, saturated NaHCO₃, and brine. The organic phase was dried over Na₂SO₄ and concentrated to give a white solid (193 mg, 0.76 mmol, 91%). This crude bromide (193 mg, 0.76 mmol) was converted, via Methods 1 and 2, to compound **8** (21 mg, 13%) which was isolated as a light yellow solid. [M-H]⁻ = 217.1 m/z. Activity: D


Example 9:

7-Bromo-2-methyl-1H-quinazolin-4-one (200 mg, 0.84 mmol, 1.0 equiv), tetrabutylammonium bromide (30 mg, 0.09 mmol, 0.1 equiv), and phenethyl bromide (150 uL, 1.0 mmol, 1.2 equiv) were combined in toluene (10 mL) and treated with 50% aq. NaOH (2 mL), and the resulting mixture was refluxed for 14 hours after which point the reaction mixture was diluted with ethyl acetate and washed with water and brine. The organic phase was dried over Na₂SO₄. Addition of silica gel, concentration, and purification of the residue using flash silica gel chromatography (gradient of 3->20% ethyl acetate/hexanes) gave a white solid (226 mg, 0.66 mmol, 79%). This bromide (226 mg, 0.66 mmol) was converted, via Methods 1 and 2, to compound **9** (69 mg, 34%) which was isolated as a white solid. [M-H]⁻ = 307.1 m/z. Activity: B

Example 10:


7-Bromo-1H-quinoxaline-4-one (200 mg, 0.89 mmol, 1.0 equiv), tetrabutylammonium bromide (30 mg, 0.09 mmol, 0.1 equiv), and phenethyl bromide (150 μ L, 1.0 mmol, 1.2 equiv) were combined in toluene (10 mL) and treated with 50% aq. NaOH (2 mL), and the resulting mixture was refluxed for 2 hours, after which the reaction mixture was diluted with ethyl acetate and washed with water. The organic phase was dried over Na₂SO₄. Addition of silica gel, concentration, and purification of the crude residue using flash silica gel chromatography (gradient of 3->25% ethyl acetates/hexanes) gave a white solid (197 mg, 0.60 mmol, 67%). This bromide (197 mg, 0.60 mmol) was converted, via Methods 1 and 2, to compound **10** (52 mg, 30%) which was isolated as a white solid. [M-H]⁻ = 293.1 m/z. Activity: A

Example 11:




11

7-Bromo-1-hydroxyisoquinoline (300 mg, 1.34 mmol, 1.0 equiv), tetrabutylammonium bromide (50 mg, 0.13 mmol, 0.10 equiv), and phenethyl bromide (220 μ L, 1.6 mmol, 1.2 equiv) were combined in toluene (14 mL) and treated with 50% aq. NaOH (3 mL), and the resulting mixture stirred rapidly while heating to 80 °C. After 1.5 hours, the mixture was diluted with MTBE and washed with water, saturated NaHCO₃, and brine. The organic phase was dried over Na₂SO₄. Addition of silica gel, concentration, and purification of the crude residue using flash silica gel chromatography (gradient of 2-30% ethyl acetates/hexanes) provided 7-bromo-2-phenethyl-isoquinolin-1-one (340 mg, 1.04 mmol, 77%) as a white solid. This isoquinolinone (340 mg, 1.04 mmol) was converted, via Methods 1 and 2, to compound **11** (141 mg, 46%) which was isolated as a white solid. Activity: B


Example 12:**12**

6-Bromo-1-hydroxy-3,4-dihydroisoquinoline (300 mg, 1.33 mmol, 1.0 equiv), tetrabutylammonium bromide (50 mg, 0.13 mmol, 0.10 equiv), and phenethyl bromide (220 μ L, 1.6 mmol, 1.2 equiv) were combined in toluene (14 mL) and treated with 50% aq. NaOH (3 mL), and the resulting mixture stirred rapidly while heating to 70 °C. After heating for 14 hours, the mixture was diluted with MTBE and washed with water, saturated NaHCO₃, and brine. The organic phase was dried over Na₂SO₄. Addition of silica gel, concentration, and purification of the residue using flash silica gel chromatography (gradient of 3->18% ethyl acetate/hexanes) provided 6-bromo-2-phenethyl-3,4-dihydroisoquinolin-1-one (201 mg, 0.61 mmol, 46%) as a white solid. This isoquinolinone (201 mg, 0.61 mmol) was converted, via Methods 1 and 2, to compound **12** (90 mg, 50%) which was isolated as a white solid. Activity: A

Example 13:**13**


A solution of 5-bromophthalide (1.0g, 4.7 mmol, 1.0 equiv) and NBS (850 mg, 4.7 mmol, 1.0 equiv) in carbon tetrachloride (15 mL) was treated with AIBN (50 mg, cat.) and heated to reflux overnight. The hot reaction mixture was filtered and the filtrate concentrated to a yellowish solid (1.38g, quant.). Crude ¹H NMR indicated this material contained ca. 80% of the desired 3,5-dibromophthalide, and it was carried on without further purification. An aliquot of this crude material (250 mg, 0.69 mmol, 1.0 equiv) in 95% ethanol (6 mL) was treated with phenelzine sulfate (175 mg, 0.75 mmol, 1.1 equiv) and NaHCO₃ (250 mg, 2.8 mmol, 4.0 equiv). After stirring overnight at ambient temperature, the reaction mixture was diluted with diethyl

ether, washed with dilute NaOH, dilute HCl, and saturated NaHCO₃, then dried over Na₂SO₄ and concentrated. Purification of the residue using flash silica gel chromatography (gradient of 3-12% ethyl acetate/hexanes) provided 6-bromo-2-phenethyl-1-phthalazinone (147 mg, 0.45 mmol, 65%) as a white solid. This crude phthalazinone (147 mg, 0.45 mmol) was converted, via Methods 1 and 2, to compound **13** (81 mg, 62%) which was isolated as a white solid. [M-H]⁻ = 293.1 m/z. Activity: A

Example 14:**14**

2-Amino-5-bromobenzoic acid (2.0 g, 9.3 mmol, 1.8 equiv) in ether (50 mL) was treated with hydrocinnamoyl chloride (760 μ L, 0.51 mmol, 1.0 equiv) and stirred overnight. The solution was filtered and washed sparingly with ether. The filtrate, upon concentration, gave a white amorphous powder which was recrystallized from ethanol/water to give fine yellowish needles of 5-bromo-2-(3-phenylpropanamido)benzoic acid (1.58g, 4.5 mmol, 89%). This material (1.58g, 4.5 mmol, 1.0 equiv) was added to acetic anhydride (8 mL) and heated to 100 °C for 30 min. to dissolve all the material, then 160 °C until ca. 20% of the volume had distilled. The reaction mixture was then concentrated *in vacuo* to provide 6-bromo-2-phenethylbenzo[3,1]oxazin-4-one as a yellowish waxy solid.

A solution of DMF (5 mL) and concentrated ammonium hydroxide (5 mL) was heated in to 80 °C in an oil bath and crude benzoxazinone (500 mg, 1.5 mmol, 1.0 equiv) was added in one portion. When dissolution was complete, a 6M KOH solution (300 μ L) was added and the mixture heated for 1 hour. The mixture was cooled and diluted with water (5 mL). The resulting solid was collected by filtration and washed with water to yield 6-bromo-2-phenethylquinazolinone (265 mg, 0.81 mmol, 53%) as white crystals. This crude quinazolinone (265 mg, 0.81 mmol) was converted, via Methods 1 and 2, to compound **14** (40 mg, 1.4 mmol, 17%) which was isolated as a beige solid. [M-H]⁻ = 293.1 m/z. Activity: C

Example 15:

5-Bromophthalide (500 mg, 2.3 mmol) was placed in a small tube, dissolved in neat phenethylamine (1 mL), sealed, and heated at 150 °C overnight. The resulting black solution was diluted with diethyl ether, washed with dilute HCl and saturated sodium bicarbonate, dried on Na₂SO₄, and concentrated to give crude *N*-phenethyl-2-hydroxymethyl-4-bromobenzamide as a yellow solid (721 mg, 92%). This crude material (250 mg, 0.75 mmol) was taken up in toluene (8 mL) and treated with tetrabutylammonium bromide (25 mg, 0.08 mmol, 0.1 eq) and *p*-toluenesulfonyl chloride (160 mg, 0.82 mmol, 1.1 eq). A sodium hydroxide solution (50%, 2 mL) was added and the biphasic mixture stirred rapidly at ambient temperature. After the reaction was allowed to stir overnight, LC/MS showed complete consumption of the starting material. The mixture was diluted with diethyl ether, washed with dilute HCl and saturated NaHCO₃, dried over Na₂SO₄, and concentrated to give crude 2-phenethyl-5-bromo-2,3-dihydroisoindole-1-one as a yellowish solid (149 mg, 63%). This crude material (149 mg, 0.47 mmol) was converted using Method 3 followed by Method 5, to form boronic acid **15** (21 mg, 16%) as a white solid. Activity: A.

Example 16: Inhibition of Rat and Human FAAH

The following assays were used to determine the inhibition of FAAH: (1) a fluorescence-based assay (Manjunath *et al.*, *Analytical Biochemistry* (2005) 343:143–151); and (2) a microsome-based fluorescent assay (Wang *et al.*, *Biomolecular Screening* (2006) 1–9).

Rat FAAH Preparation: Five rat livers were homogenized in five fold volume with ice cold Tris (20 mM pH 8.0) and 0.32 M Sucrose solution via an Ultra Turrax T25 homogenizer. All subsequent preparation steps were carried out at 4 °C. The homogenate was centrifuged at 6000 g, for 20 minutes and the pellet, containing nuclear debris and mitochondria was discarded. The supernatant was centrifuged at 40,000 g for 30 minutes. The supernatant was discarded and the pellet solubilized via a dounce homogenizer in resuspension buffer (20 mM Hepes pH 7.8, 10% v/v glycerol, 1 mM EDTA, 1% triton X-100) overnight at 4 °C to resolubilize membrane

bound FAAH. The solution was centrifuged at 40,000 g for 30 minutes and the pellet discarded. The supernatant containing rat FAAH was aliquoted and flash frozen with liquid nitrogen and stored for long term usage at -80°C .

Human FAAH Preparation: COS-7 cells were split the day before, 1:5 into 150 mm x 25 mm cell culture dishes (Corning Inc., Cat. No. 430599). Transient transfection took place at 30–40% confluency according to FuGENE 6 Transfection Reagent (Roche, Cat. No. 11814 443 001).

Transfection Procedure: The FuGENE transfection 6 reagent (45uL) was added to 1410 μL of media (DMEM, serum free without pen/strep) in a 15 mL conical tube and incubated at room temp for 5 minutes, followed by the addition of FAAH plasmid DNA (15 μg) (OriGene Cat. No. TC119221, Genbank Accession No. NM_001441.1, 0.67 $\mu\text{g}/\mu\text{L}$) and a further incubation of 15 minutes at room temperature. The resulting solution was added into one dish of 30–40% confluent COS-7 cells in a drop-wise manner. The COS-7 cell dish was subsequently incubated for 48 hours. The cells were then harvested.

Harvest procedure: Media was aspirated from the dishes and the cells rinsed with 10mL PBS. The PBS was removed and 3 mL of PBS added to the dish. The dish was scraped to resuspend the cells, and the subsequent cell suspension collected into a 15 mL conical tube. The cells were pelleted by centrifugation at 1200 rpm for 5 minutes in a bench top centrifuge. PBS was removed and the cell pellet snap frozen in liquid nitrogen and stored at -80°C .

COS-7 cells – FAAH purification:

(1) Fractionation: Frozen cell pellets from transient transfections were thawed on ice and resuspended in 12.5mM Hepes pH 8.0, 100mM NaCl, 1mM EDTA (10 mL/0.2 g cell pellet). The pellets were dounce homogenized and then sonicated to produce cell extract. The cell extract was subsequently centrifuged at 1000 g to remove cellular debris. The pellet was discarded and the supernatant centrifuged at 13,000 g for 20 minutes. The pellet contained membrane bound FAAH. The supernatant was discarded and the pellet resolubilized.

(2) Re-solubilization: The fraction of interest, (13,000g, membrane fraction) was re-suspended in 2.3 mL re-suspension buffer (20mM Hepes pH 7.8, 10%v/v Glycerol, 1 mM EDTA, 1% Triton X-100) and the sample incubated on ice for 1 hour and then centrifuged to remove any particulate matter. The supernatant containing

solubilized human FAAH was aliquoted and snap frozen in liquid nitrogen and stored at –80 °C until use.

(3) Characterization: Protein Concentration determined by Bradford assay.

SDS gel and Western blot to confirm presence of FAAH

FAAH activity assay

K_m determination – 96–well assay

Linear dependence – 96–well assay

Standard compound K_i determination – 384–well assay

Rat FAAH Biochemical Inhibition Assay; Materials and methods: Rat FAAH

biochemical assays were carried out in a 96 well flat bottom black non–treated polystyrene plates (Corning Costar Catalogue # 3915). FAAH reaction buffer: 50 mM Hepes (pH 7.5), 1 mM EDTA, 0.2% Triton X–100. FAAH substrate– AMC Arachidonoyl Amide (Cayman Chemicals Company, Catalog # 10005098). The reaction was read in an Envision microtiter plate reader [Excitation filter 355 nm (40 nm bandpass); Emmision filter 460 nm (25 nm bandpass)]. The raw fluorescence was plotted on the y axis and the inhibitor concentration on the x axis to give a dose response inhibition curve. The data was fitted to a single site competitive inhibition equation, fixing the K_m for the rat and human enzyme to 12 μM and 9 μM respectively.

Rat FAAH Biochemical Inhibition Assay; Experimental Protocol: The principle of this assay was the hydrolysis of AMC–Arichodonoyl, a fluorescent analogue of Anandamide, which results in the formation of Arachidonic acid and AMC. The formation of AMC results in an increase in fluorescence (see, for example, Manjunath *et al.*, *Analytical Biochemistry* (2005) 343:143–151; and Wang *et al.*, *Biomolecular Screening* (2006) 1–9). The inhibition of product formation and hence fluorescence as a function of inhibitor concentration enables the determination of K_i for the compounds.

A 0.49 mg/ml Rat liver FAAH solution was made up in FAAH reaction buffer, and 78 ul pipetted into a 96 well plate. To this was added 2 uL of a 3 fold serially diluted inhibitor from a DMSO stock solution. The FAAH solution and inhibitor were incubated for 30 minutes at room temperature. The FAAH reaction was initiated by the addition of 80 μL of 40 μM AMC Arachidonoyl Amide in FAAH reaction buffer, yielding a final reaction FAAH rat liver preparation concentration of 0.25 mg/mL and AMC–Arachidonoyl substrate concentration of 20 μM, reaction volume 160 μL. The reaction was allowed to proceed for 4 hours at room

temperature. The reaction was stopped by the addition of 80 μ L 12 μ M α -ketoheterocycle (Cayman Chemicals, catalogue # 10435). The microtiter plate was read in the envision plate reader.

Human FAAH assay; Experimental Protocol: A 0.1 mg/mL Human FAAH solution was made up in FAAH reaction buffer, and 24 μ L pipetted into a 384 well plate. To this was added 1 μ L of a 3 fold serially diluted inhibitor from a DMSO stock solution. The FAAH solution and inhibitor were incubated for 30 minutes at room temperature. The FAAH reaction was initiated by the addition of 25 μ L of 40 μ M AMC Arachidonoyl Amide in FAAH reaction buffer, yielding a final reaction human FAAH preparation concentration of 0.05 mg/ml and AMC–Arachidonoyl substrate concentration of 20 μ M, reaction volume 50 μ L. The reaction was allowed to proceed for 4 hours at room temperature. The reaction was stopped by the addition of 25 μ L 12 μ M α -ketoheterocycle (Cayman Chemicals, catalogue # 10435). The microtiter plate was read in the envision plate reader.

The raw fluorescence was plotted on the y axis and the inhibitor concentration on the x axis to give a dose response inhibition curve. The data was fitted to a single site competitive inhibition equation, fixing the K_m for the rat and human enzyme to 12 μ M and 9 μ M respectively.


The contents of all references, pending patent applications and published patent applications, cited throughout this application are hereby incorporated by reference in their entirety as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

CLAIMS

We claim:

1. A compound of formula I:



I

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof,

wherein:

----- is selected from a single bond and a double bond;

when ----- is a single bond, X¹ is selected from CR⁴R⁵ and NR⁶, and X² is selected from CR⁷R⁸ and NR⁹;

when ----- is a double bond, X¹ is selected from CR⁴ and N, and X² is selected from CR⁷ and N;

X³ is selected from CR¹⁰R¹¹ and NR¹²;

provided that at least one of X¹, X² and X³ is selected from N, NR⁶, NR⁹, and N¹²;

m is 0 or 1;

Z¹ is selected from -OR¹³ and C₁₋₆ alkyl;

Z² is selected from -OR¹⁴ and C₁₋₆ alkyl;

or alternatively, Z¹ and Z², together with the B to which they are bound, form an 5- to 8-membered ring having at least one O atom directly attached to B, wherein the ring is comprised of carbon atoms and optionally one or more additional heteroatoms independently selected from the group consisting of N, S, and O;

n is 0, 1, 2 or 3;

R¹ and R² each independently is selected from H, halogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ perhaloalkyl, -CN and -OR¹⁵;

or alternatively, R¹ and R², taken together with the carbon to which they are bound, form a carbonyl group;

R^3 , at each occurrence, independently is selected from halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} perhaloalkyl, -OH, C_{1-6} alkoxy and -CN;

R^4 , R^5 , R^7 , R^8 , R^{10} , and R^{11} each independently is selected from H, halogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} perhaloalkyl, -CN, -OR¹⁶, NR¹⁷R¹⁸, -C(O)R¹⁹, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

R^6 , R^9 , and R^{12} each independently is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, -C(O)R²⁰, -C(O)OR²¹, S(O)₂R²², C_{3-10} carbocyclyl, C_{6-10} aryl, 3-10 membered heterocyclyl, 5-10 membered heteroaryl, and -(CR²⁷R²⁸)_p-R²³;

R^{13} and R^{14} , at each occurrence, each independently is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl;

R^{15} and R^{16} , at each occurrence, each independently is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

R^{17} and R^{18} , at each occurrence, each independently is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, -C(O)R²⁴, -C(O)OR²⁵, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

R^{19} , R^{20} and R^{21} , at each occurrence, each independently is selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

R^{22} is selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, C_{6-10} aryl, 3-10 membered heterocyclyl, 5-10 membered heteroaryl, and -(CR²⁹R³⁰)_q-R²⁶;

R^{23} , at each occurrence, independently is selected from C_{3-10} carbocyclyl, C_{6-10} aryl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

R^{24} and R^{25} , at each occurrence, each independently is selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, C_{6-10} aryl, C_{7-12} aralkyl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

R^{26} , at each occurrence, independently is selected from C_{3-10} carbocyclyl, C_{6-10} aryl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl;

R^{27} , R^{28} , R^{29} and R^{30} , at each occurrence, each independently is selected from H and C_{1-6} alkyl; and

p, and q, at each occurrence, each independently is selected from 1, 2, 3, 4, 5 and 6..

2. The compound of claim 1, wherein m is 1 and X^3 is NR^{12} .
3. The compound of claim 2, wherein ----- is a double bond.
- 5 4. The compound of claim 3, wherein X^1 is CR^4 .
5. The compound of claim 4, wherein X^2 is CR^7 .
- 10 6. The compound of claim 5, wherein R^7 is selected from H, C_{1-6} alkyl and C_{7-10} aralkyl.
7. The compound of claim 4, wherein X^2 is N.
8. The compound of any one of claims 4 to 7, wherein R^4 is selected from H, C_{1-6} alkyl and
15 C_{7-10} aralkyl.
9. The compound of claim 3, wherein X^1 is N.
10. The compound of claim 9, wherein X^2 is CR^7 .
- 20 11. The compound of claim 10, wherein R^7 is selected from H, C_{1-6} alkyl and C_{7-10} aralkyl.
12. The compound of claim 2, wherein ----- is a single bond.
- 25 13. The compound of claim 12, wherein X^1 is CR^4R^5 .
14. The compound of claim 12 or 13, wherein X^2 is CR^7R^8 .
15. The compound of claim 14, wherein R^7 is H and R^8 is H.
- 30 16. The compound of any one of claims 13 to 15, wherein R^4 is H and R^5 is H.

17. The compound of any one of claims 2 to 16, wherein R^{12} is selected from H, C_{1-6} alkyl and $-(CR^{27}R^{28})_p-R^{23}$.

5 18. The compound of claim 1, wherein m is 0.

19. The compound of claim 18, wherein ----- is a single bond.

20. The compound of claim 19, wherein X^2 is NR^9 .

10

21. The compound of claim 20, wherein R^9 is selected from H, C_{1-6} alkyl and $-(CR^{27}R^{28})_p-R^{23}$.

22. The compound of any one of claims 19 to 21, wherein X^1 is CR^4R^5 .

15


23. The compound of claim 22, wherein R^4 is H and R^5 is H.

24. The compound of any one of claims 1 to 23, wherein R^1 and R^2 , taken together with the carbon to which they are bound, form a carbonyl group.

20

25. The compound of any one of claims 1 to 24, wherein n is 0.

26. The compound of any one of claims 1 to 25, wherein the compound is of the formula **Ia**:




25

Ia


or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

27. The compound of any one of claims 1 to 25, wherein the compound is of the formula **Ib**:

**Ib**



- 5 or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

28. The compound of claim 1, wherein the compound is of the formula **V**:

**V**


- 10 or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

29. The compound of claim 28, wherein the compound is of the formulae **Va** or **Vb**:

**Va****Vb**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.


30. The compound of claim 1, wherein the compound is of the formula **IX**:




IX

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

31. The compound of claim 30, wherein the compound is of the formulae **IXa** or **IXb**:




IXa



IXb

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

32. The compound of claim 1, wherein the compound is of the formula **XI**:




XI

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.


33. The compound of claim 32, wherein the compound is of the formulae **XIa** or **XIb**:

5





or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

- 10 34. The compound of claim 1, wherein the compound is of the formula **VII**:



or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

- 15 35. The compound of claim 34, wherein the compound is of the formulae **VIIa** or **VIIb**:


**VIIa****VIIb**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

36. The compound of any one of claims 1 to 35, wherein R^{12} is selected from H, C_{1-6} alkyl and $-(CH_2)_p-R^{23}$.



37. The compound of claim 36, wherein R^{23} is phenyl.

38. The compound of claim 1, wherein the compound is of the formula **XV**:

**XV**


or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.


39. The compound of claim 38, wherein the compound is of the formulae **XVa** or **XVb**:

**XVa****XVb**

or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

- 5 40. The compound of claim 38 or 39, wherein R^9 is selected from H, C_{1-6} alkyl and $-(CH_2)_p-R^{23}$.
41. The compound of claim 40, wherein R^{23} is phenyl.
- 10 42. The compound of any one of claims 1 to 41, wherein Z^1 is $-OR^{13}$ and Z^2 is $-OR^{14}$.
43. The compound of claim 42, wherein R^{13} is H and R^{14} is H.
44. The compound of claim 1, wherein the compound is selected from:





or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof.

5

45. A pharmaceutical composition comprising a compound of any one of claims 1 to 44 or a pharmaceutically acceptable salt, solvate or prodrug thereof, or mixture thereof, and a pharmaceutically acceptable excipient.

10 46. A method of treating an FAAH-mediated disorder comprising administering to a subject in need thereof a therapeutically effective amount of a compound or composition of any one of claims 1 to 45.

15 47. The method of claim 46, wherein the FAAH-mediated disorder is selected from a painful disorder, an inflammatory disorder, an immune disorder, depression, anxiety, an anxiety-related disorder, a sleep disorder, a feeding behavior, a movement disorder, glaucoma, neuroprotection and cardiovascular disease.

48. The method of claim 47, wherein the FAAH-mediated disorder is a painful disorder.

20

49. The method of 48, wherein the painful disorder is selected from neuropathic pain, central pain, deafferentiation pain, chronic pain, stimulus of nociceptive receptors, acute pain, non-

inflammatory pain, inflammatory pain, pain associated with cancer, preoperative pain, arthritic pain, lumbosacral pain, musculo-skeletal pain, headache, migraine, muscle ache, lower back and neck pain, and toothache.

5 50. The method of claim 49, wherein the painful disorder is neuropathic pain.

51. The method of claim 49, wherein the painful disorder is arthritic pain.

52. The method of claim 49, wherein the arthritic pain is osteoarthritic pain.

10

53. The method of claim 49, wherein the arthritic pain is rheumatoid arthritic pain.

54. The method of claim 49, wherein the inflammatory pain is associated with an inflammatory disorder.

15

55. The method of claim 46, wherein the FAAH-mediated disorder is an inflammatory disorder.

56. The method of claim 51, wherein the inflammatory disorder is irritable bowel disease.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 10/30272

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A01N 55/08; A61K 31/69, 31/47 (2010.01)

USPC - 514/64, 307

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

USPC - 514/64, 307

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
USPC - 514/187-188, 253.05 (see search terms below)Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Electronic Database Searched: PubWest (PGPB,USPT,USOC,EPAB,JPAB), Google. search terms: Quinazolinone, quinoline, quinazoline, boronic acid, boron**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2005/0182045 A1 (Nagase et al.) 18 August 2005 (18.08.2005) especially para [0013]-[0014]	1-15, 18-23, 28-35, 38-41 and 44
Y	US 2006/0234981 A1 (Baker et al.) 19 October 2006 (19.10.2006) especially para [0092], compound 6	1-15, 18-23, 28-35, 38-41 and 44
A	US 5,616,582 A (Baker) 01 April 1997 (01.04.1997) entire document	1-15, 18-23, 28-35, 38-41 and 44
A	US 5,089,499 A (Baker et al.) 18 February 1992 (18.02.1992) entire document	1-15, 18-23, 28-35, 38-41 and 44

☐ Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

22 May 2010 (22.05.2010)

Date of mailing of the international search report

02 JUN 2010

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 10/30272

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☒ Claims Nos.: 16-17, 24-27, 36-37, 42-43 and 45-56
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.