

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 April 2003 (10.04.2003)

PCT

(10) International Publication Number
WO 03/028543 A2

(51) International Patent Classification⁷: **A61B**

(21) International Application Number: PCT/US02/31642

(22) International Filing Date: 3 October 2002 (03.10.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/326,708 3 October 2001 (03.10.2001) US

(71) Applicants (*for all designated States except US*): **PFIZER PRODUCTS INC.** [US/US]; Eastern Point Road, Groton, CT 06340-5196 (US). **OXFORD GLYCOSCIENCES (UK) LTD.** [GB/GB]; The Forum, 86 Milton Park, Abingdon, OX14 4RY (GB).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **DURHAM, L., Kathryn** [US/US]; 94 Glenwood Avenue, New London, CT 06320 (US). **FRIEDMAN, David, L.** [US/US]; 368 Bartlett Drive, Madison, CT 06433 (US). **HERATH, Herath, Mudiyansele, Athula, Chandrasiri** [GB/GB]; Oxford Glycosciences (UK) LTD., The Forum, 86 Milton Park, Abingdon, Oxfordshire OX14 4RY (GB). **KIM-MEL, Lida, H.** [US/US]; 55 Bokum Road, Chester, CT 06412 (US). **PAREKH, Rajesh, Bhikhu** [GB/GB]; Oxford Glycosciences (UK) LTD., The Forum, 86 Milton Park, Abingdon, Oxfordshire OX14 4RY (GB). **POTTER, David, M.** [US/US]; 33 Chriswood Trace, Ledyard, CT 06339 (US). **ROHLFF, Christian** [DE/GB]; Oxford Glycosciences (UK) LTD., The Forum, 86 Milton Park, Abingdon, Oxfordshire OX14 4RY (GB). **SILBER, B., Michael** [US/US]; 104 Randi Drive, Madison, CT 06433 (US). **SNYDER, Peter, Jeffrey** [US/US]; 98 Arundel Avenue, West Hartford, CT 06107 (US). **SOARES, Holly,**

Daria [US/US]; 108 Seneca Drive, Noank, CT 06340 (US). **STIGER, Thomas, R.** [US/US]; 93 Castle Hill Road, Pawcatuck, CT 06379 (US). **SUNDERLAND, P., Trey** [US/US]; 4718 Cumberland Avenue, Chevy Chase, MD 20815 (US). **TOWNSEND, Robert, Reid** [US/GB]; Oxford Glycosciences (UK) LTD., The Forum, 86 Milton Park, Abingdon, Oxfordshire OX14 4RY (GB). **WHITE, W., Frost** [US/US]; 65 Homestead Road, Ledyard, CT 06339 (US). **WILLIAMS, Stephen, A.** [GB/US]; 114 Colony Road, Groton, CT 06340 (US).

(74) Agents: **ARNOLD, Beth, E.** et al.; Patent Group, Foley Hoag LLP, 155 Seaport Boulevard, Boston, MA 02210 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NUCLEIC ACID MOLECULES, POLYPEPTIDES AND USES THEREFOR, INCLUDING DIAGNOSIS AND TREATMENT OF ALZHEIMER'S DISEASE

(57) Abstract: The present invention provides methods and compositions for screening, diagnosis and prognosis of Alzheimer's disease, for monitoring the effectiveness of Alzheimer's disease treatment, and for drug development. Alzheimer's Disease-Associated features (AFs), detectable by two-dimensional electrophoresis of cerebrospinal fluid, serum or plasma are described. The invention further provides Alzheimer's Disease-Associated Protein Isoforms (APIs) detectable in cerebrospinal fluid, serum or plasma, preparations comprising isolated APIs antibodies immunospecific for APIs, pharmaceutical compositions, diagnostic and therapeutic methods, and kits comprising or based on the same.



WO 03/028543 A2

NUCLEIC ACID MOLECULES, POLYPEPTIDES AND USES THEREFOR, INCLUDING DIAGNOSIS AND TREATMENT OF ALZHEIMER'S DISEASE

1. INTRODUCTION

5 The present invention relates to the identification of proteins and protein isoforms that are associated with predisposition to Alzheimer's Disease and its onset and development, and of genes and nucleic acid molecules, encoding the same, and to their use for *e.g.*, clinical screening, diagnosis, treatment, as well as for drug screening and drug development.

2. BACKGROUND OF THE INVENTION

10 Alzheimer's Disease (AD) is an increasingly prevalent form of neurodegeneration that accounts for approximately 50 % - 60 % of the overall cases of dementia among people over 65 years of age. It currently affects an estimated 15 million people worldwide and
15 owing to the relative increase of elderly people in the population its prevalence is likely to increase over the next 2 to 3 decades. AD is a progressive disorder with a mean duration of around 8.5 years between onset of clinical symptoms and death. Death of pyramidal neurons and loss of neuronal synapses in brains regions associated with higher mental functions results in the typical symptomology, characterized by gross and progressive
20 impairment of cognitive function (Francis et al., 1999, J. Neurol. Neurosurg. Psychiatry 66:137-47). Currently, a diagnosis of AD requires a careful medical history and physical examination; a detailed neurological and psychiatric examination; laboratory blood studies to exclude underlying metabolic and medical illnesses that masquerade as AD; a mental status assessment and formal cognitive tests; and a computed tomographic scan or magnetic
25 resonance image of the brain (Growdon, JH., 1995, Advances in the diagnosis of AD. In: Iqbal, K., Mortimer, JA., Winblad, B., Wisniewski, HM eds Research Advances in Alzheimer's Disease and Related Disorders. New York, NY: John Wiley & Sons Inc. 1995:139-153). Due to the time consuming nature of these tests, their expense, and their inconvenience to patients, it would be highly desirable to measure a substance or substances
30 in body samples, such as samples of cerebrospinal fluid (CSF), blood or urine, that would lead to a positive diagnosis of AD or that would help to exclude AD from the differential diagnosis. Since the CSF bathes the brain, changes in its protein composition may most accurately reveal alterations in brain protein expression patterns that are causatively or diagnostically linked to the disease.

Current candidate biomarkers for AD include: (1) mutations in presenilin 1 (PS1), presenilin 2 (PS2) and amyloid precursor protein (APP) genes; (2) the detection of alleles of apolipoprotein E (ApoE); and (3) altered concentrations of amyloid β -peptides (A β), tau protein, and neuronal thread protein (NTP) in the CSF. See, *e.g.*, Neurobiology of Aging 19:109-116 (1998) for a review. Mutations in PS1, PS2 and APP genes are indicative of early-onset familial AD. However, early-onset familial AD is relatively rare; only 120 families worldwide are currently known to carry deterministic mutations (Neurobiology of Aging 19:109-116 (1998)). The detection of the ϵ 4 allele of ApoE has been shown to correlate with late-onset and sporadic forms of AD. However, ϵ 4 alone cannot be used as a biomarker for AD since ϵ 4 has been detected in many individuals not suffering from AD and the absence of ϵ 4 does not exclude AD (Neurobiology of Aging 19:109-116 (1998)).

A decrease in the A β peptide A β 42 and an increase in tau protein in the CSF of AD have been shown to correlate with the presence of AD (Neurobiology of Aging 19:109-116 (1998)). However, the specificity and sensitivity of A β 42 and tau protein as biomarkers of AD are modest. For example, it has been difficult to determine a cut-off level of CSF tau protein that is diagnostically informative. Also, elevated levels of NTP in the CSF of postmortem subjects have been shown to correlate with the presence of AD (Neurobiology of Aging 19:109-116 (1998)). Therefore, a need exists to identify sensitive and specific biomarkers for the diagnosis of AD in living subjects.

3. SUMMARY OF THE INVENTION

The present invention provides methods and compositions for screening, diagnosis and treatment of AD and for screening and development of drugs for treatment of AD.

Therefore, one aspect of the invention provides methods for identification of AD that comprise analyzing a sample of CSF by two-dimensional electrophoresis to detect the presence or level of at least one Alzheimer's Disease-Associated Feature (AF), *e.g.*, one or more of the AFs disclosed herein, or any combination thereof. These methods are also suitable for clinical screening, prognosis, monitoring the results of therapy, for identifying patients most likely to respond to a particular therapeutic treatment, drug screening and development, and identification of new targets for drug treatment.

A further aspect of the invention provides methods for diagnosis of AD that comprise detecting in a sample of CSF the presence or level of at least one Alzheimer's

Disease-Associated Protein Isoform (API), *e.g.*, one or more of the APIs disclosed herein or any combination thereof.

Another aspect of the invention provides antibodies, *e.g.*, monoclonal, polyclonal, humanised, chimeric or bispecific antibodies capable of immunospecific binding to an API, *e.g.*, an API disclosed herein.

A further aspect of the invention provides a preparation comprising an isolated API, *i.e.*, an API substantially free from proteins or protein isoforms having a significantly different isoelectric point or a significantly different apparent molecular weight from the API.

Another aspect of the invention provides kits that may be used in the above recited methods and that may comprise single or multiple preparations, or antibodies, together with other reagents, labels, substrates, if needed, and directions for use. The kits may be used for diagnosis of disease, or may be assays for the identification of new diagnostic and/or therapeutic agents.

An additional aspect of the invention provides methods of treating AD, comprising administering to a subject a therapeutically effective amount of an agent that modulates (*e.g.*, upregulates or downregulates) the expression or activity (*e.g.* enzymatic or binding activity), or both, of an API in subjects having AD.

A further aspect of the invention provides methods of screening for agents that modulate a characteristic of, *e.g.*, the expression or the enzymatic or binding activity, of an API, an API analog, or an API-related polypeptide.

Other objects and advantages will become apparent from a review of the ensuing detailed description taken in conjunction with the following illustrative drawings.

4. BRIEF DESCRIPTION OF THE FIGURE

Figure 1 is an image obtained from 2-dimensional electrophoresis of normal CSF, which has been annotated to identify twelve landmark features, designated CSF1 to CSF12, and which are illustrative of an embodiment of an aspect of the present invention.

Figure 2 is a flow chart depicting the characterization of a Feature and relationship of a Feature and Protein Isoform(s). A feature may be further characterized as or by a Protein Isoform having a particular peptide sequence associated with its pI and MW. As depicted herein, a feature may comprise one or more protein isoform(s), which have indistinguishable pIs and MWs using the Preferred Technology, but which comprise

distinct peptide sequences. The peptide sequence(s) of the protein isoform can be utilized to search database(s) for previously identified proteins comprising such peptide sequence(s), for which previously identified protein it can be ascertained whether a commercially available antibody exists, which may recognize the previously identified protein and/or a member of its protein family.

5. DETAILED DESCRIPTION OF THE INVENTION

The present invention described in detail below provides methods, compositions and kits useful, *e.g.*, for screening, diagnosis and treatment of Alzheimer's disease (AD) in a mammalian subject, and for drug screening and drug development. The invention also encompasses the administration of therapeutic compositions to a mammalian subject to treat or prevent AD a human adult, *i.e.* a human subject at least 21 (more particularly at least 35, at least 50, at least 60, at least 70, or at least 80) years old. For clarity of disclosure, and not by way of limitation, the invention will be described with respect to the analysis of CSF samples. However, as one skilled in the art will appreciate, based on the present description the assays and techniques described below can be applied to other types of samples, including a body fluid (*e.g.* blood, serum, plasma or saliva), a tissue sample from a subject at risk of having or developing AD (*e.g.* a biopsy such as a brain biopsy) or homogenate thereof. The methods and compositions of the present invention are useful, such as for example, screening, diagnosis and treatment of a living subject, but may also be used for post-mortem diagnosis of a subject, for example, to identify if family members of the subject may be at risk of developing the same disease.

The following definitions are provided to assist in the review of the instant disclosure.

5.1. DEFINITIONS

"Feature" refers to a spot identified in a 2D gel, and the term "Alzheimer's disease-Associated Feature" (AF) refers to a feature that is differentially present in a first sample or sample set from a subject having AD compared with a second sample or sample set from a subject free from AD. A feature or spot identified in a 2D gel is characterized by its isoelectric point (pI) and apparent molecular weight (MW) as determined by 2D gel electrophoresis, particularly utilizing the Preferred Technology described herein. As used herein, a feature is "differentially present" in a first sample or sample set with respect to a second sample or sample set when a method for detecting the said feature (*e.g.*, 2D

electrophoresis) gives a different signal when applied to the first and second samples or sample sets. An AF, (or a Protein Isoform, i.e. API, as defined *infra*) is "increased" in the first sample or sample set with respect to the second sample or sample set if the method of detection indicates that the AF, or API is more abundant in the first sample or sample set than in the second sample or sample set, or if the AF, or API is detectable in the first sample or sample set and substantially undetectable in the second sample or sample set. Conversely, an AF, or API is "decreased" in the first sample or sample set with respect to the second sample or sample set if the method of detection indicates that the AF, or API is less abundant in the first sample or sample set than in the second sample or sample set or if the AF, or API is undetectable in the first sample or sample set and detectable in the second sample or sample set.

Particularly, the relative abundance of a feature in the two samples or sample sets is determined in reference to its normalized signal, in two steps. First, the signal obtained upon detecting the feature in a first sample or sample set is normalized by reference to a suitable background parameter, *e.g.*, (a) to the total protein in the sample being analyzed (*e.g.*, total protein loaded onto a gel); (b) to an Expression Reference Feature (ERF) i.e., a feature whose abundance is substantially invariant, within the limits of variability of the Preferred Technology, in the population of subjects being examined, *e.g.* the ERFs disclosed in Table III, or (c) more preferably to the total signal detected as the sum of each of all proteins in the sample.

Secondly, the normalized signal for the feature in the first sample or sample set is compared with the normalized signal for the same feature in the second sample or sample set in order to identify features that are "differentially present" in the first sample or sample set with respect to the second sample or sample set.

"Fold change" includes "fold increase" and "fold decrease" and refers to the relative increase or decrease in abundance of a MSF or the relative increase or decrease in expression or activity of a polypeptide (*e.g.* an API, as defined *infra*.) in a first sample or sample set compared to a second sample or sample set. An AF or polypeptide fold change may be measured by any technique known to those of skill in the art, albeit the observed increase or decrease will vary depending upon the technique used. Preferably, fold change is determined herein as described in the Examples *infra*.

"Alzheimer's Disease-Associated Protein Isoform" (API) refers to a polypeptide that is differentially present in a first sample or sample set from a subject having AD

compared with a second sample or sample set from a subject free from AD. As used herein, an API is "differentially present" in a first sample or sample set with respect to a second sample or sample set when a method for detecting the said feature, (*e.g.*, 2D electrophoresis or immunoassay) gives a different signal when applied to the first and second samples or sample sets (as described above in relation to AFs). An API is characterised by one or more peptide sequences of which it is comprised, and further by a pI and MW, preferably determined by 2D electrophoresis, particularly utilising the Preferred Technology as described herein. Typically, APIs are identified or characterised by the amino acid sequencing of AFs (Figure 2).

An API is characterized as, or by, a particular peptide sequence associated with its pI and MW. As depicted herein, an AF may comprise one or more API(s), which have indistinguishable pIs and MWs using the Preferred Technology, but which have distinct peptide sequences. The peptide sequence(s) of the API can be utilized to search database(s) for previously identified proteins comprising such peptide sequence(s). In some instances, it can be ascertained whether a commercially available antibody exists which may recognize the previously identified protein and/or a variant thereof. Preferably the API corresponds to the previously identified protein, or be a variant of the previously identified protein.

"Variant" as used herein refers to a polypeptide which is a member of a family of polypeptides that are encoded by a single gene or from a gene sequence within a family of related genes and which differ in their pI or MW, or both. Such variants can differ in their amino acid composition (*e.g.* as a result of alternative mRNA or premRNA processing, *e.g.* alternative splicing or limited proteolysis) and in addition, or in the alternative, may arise from differential post-translational modification (*e.g.*, glycosylation, acylation, phosphorylation).

"Modulate" in reference to expression or activity of an AF, API or API-related polypeptide refers to any change, *e.g.*, upregulation or downregulation, increase or decrease, of the expression or activity of the AF, API or API-related polypeptide. Those skilled in the art, based on the present disclosure, will understand that such modulation can be determined by assays known to those of skill in the art.

"API analog" refers to a polypeptide that possesses similar or identical function(s) as as API but need not necessarily comprise an amino acid sequence that is similar or identical to the amino acid sequence of the API, or possess a structure that is similar or identical to that of the API. As used herein, an amino acid sequence of a polypeptide is

"similar" to that of an API if it satisfies at least one of the following criteria: (a) the polypeptide has an amino acid sequence that is at least 30% (more preferably, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or at least 99%)

5 identical to the amino acid sequence of the API; (b) the polypeptide is encoded by a nucleotide sequence that hybridizes under stringent conditions to a nucleotide sequence encoding at least 5 amino acid residues (more preferably, at least 10 amino acid residues, at least 15 amino acid residues, at least 20 amino acid residues, at least 25 amino acid residues, at least 40 amino acid residues, at least 50 amino acid residues, at least 60 amino residues, at least 70 amino acid residues, at least 80 amino acid residues, at least 90 amino
10 acid residues, at least 100 amino acid residues, at least 125 amino acid residues, or at least 150 amino acid residues) of the API; or (c) the polypeptide is encoded by a nucleotide sequence that is at least 30% (more preferably, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%,
15 at least 85%, at least 90%, at least 95% or at least 99%) identical to the nucleotide sequence encoding the MSPI. As used herein, a polypeptide with "similar structure" to that of a API refers to a polypeptide that has a similar secondary, tertiary or quarternary structure as that of the API. The structure of a polypeptide can determined by methods known to those skilled in the art, including but not limited to, X-ray crystallography, nuclear magnetic
20 resonance, and crystallographic electron microscopy.

"API fusion protein" refers to a polypeptide that comprises (i) an amino acid sequence of an API, API fragment, API-related polypeptide or a fragment of an API-related polypeptide and (ii) an amino acid sequence of a heterologous polypeptide (i.e., a non-API, non-API fragment or non-API-related polypeptide).

25 "API homolog" refers to a polypeptide that comprises an amino acid sequence similar to that of a API but does not necessarily possess a similar or identical function as the API.

"API ortholog" refers to a non-human polypeptide that (i) comprises an amino acid sequence similar to that of an API and (ii) possesses a similar or identical function to that of
30 the API.

"API-related polypeptide" refers to an API homolog, an API analog, a variant of an API, an API ortholog, or any combination thereof.

“Chimeric Antibody” refers to a molecule in which different portions are derived from different animal species, such as those having a human immunoglobulin constant region and a variable region derived from a murine mAb. (See, *e.g.*, Cabilly et al., U.S. Patent No. 4,816,567; and Boss et al., U.S. Patent No. 4,816,397). For example, a portion of the antibody may be fused with the constant domain of immunoglobulins (IgA, IgE, IgG, IgM), or portions thereof (CH1, CH2, CH3, or any combination thereof and portions thereof) resulting in chimeric antibodies.

“Humanised Antibody” refers to a molecule from non-human species having one or more complementarily determining regions (CDRs) from the non-human species and a framework region from a human immunoglobulin molecule.

“Derivative” refers to a polypeptide that comprises an amino acid sequence of a second polypeptide that has been altered by the introduction of at least one amino acid residue substitution, deletion or addition. The derivative polypeptide possesses a similar or identical function as the second polypeptide.

“Fragment” refers to a peptide or polypeptide comprising an amino acid sequence of at least 5 amino acid residues (preferably, at least 10 amino acid residues, at least 15 amino acid residues, at least 20 amino acid residues, at least 25 amino acid residues, at least 40 amino acid residues, at least 50 amino acid residues, at least 60 amino acid residues, at least 70 amino acid residues, at least 80 amino acid residues, at least 90 amino acid residues, at least 100 amino acid residues, at least 125 amino acid residues, at least 150 amino acid residues, at least 175 amino acid residues, at least 200 amino acid residues, or at least 250 amino acid residues) of the amino acid sequence of a second polypeptide. Preferably the fragment of a MSPI possesses the functional activity of the MSPI.

The “percent identity” of two amino acid sequences or of two nucleic acid sequences can be or is generally determined by aligning the sequences for optimal comparison purposes (*e.g.*, gaps can be introduced in either sequences for best alignment with the other sequence) and comparing the amino acid residues or nucleotides at corresponding positions. The “best alignment” is an alignment of two sequences that results in the highest percent identity. The percent identity is determined by the number of identical amino acid residues or nucleotides in the sequences being compared (*i.e.*, % identity = # of identical positions/total # of positions x 100).

The determination of percent identity between two sequences can be accomplished using a mathematical algorithm known to those of skill in the art. An example of a

mathematical algorithm for comparing two sequences is the algorithm of Karlin and Altschul (1990) Proc. Natl. Acad. Sci. USA 87:2264-2268, modified as in Karlin and Altschul (1993) Proc. Natl. Acad. Sci. USA 90:5873-5877. The NBLAST and XBLAST programs of Altschul, et al. (1990) J. Mol. Biol. 215:403-410 have incorporated such an algorithm. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to a nucleic acid molecule of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to a protein molecule of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (1997) Nucleic Acids Res. 25:3389-3402. Alternatively, PSI-Blast can be used to perform an iterated search which detects distant relationships between molecules (*Id.*). When utilizing BLAST, Gapped BLAST, and PSI-BLAST programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used.

Another example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, CABIOS (1989). The ALIGN program (version 2.0) which is part of the GCG sequence alignment software package has incorporated such an algorithm. Other algorithms for sequence analysis known in the art include ADVANCE and ADAM as described in Torellis and Robotti (1994) Comput. Appl. Biosci., 10 :3-5; and FASTA described in Pearson and Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-8. Within FASTA, ktup is a control option that sets the sensitivity and speed of the search.

“Diagnosis” refers to diagnosis, prognosis, monitoring, characterizing, selecting patients, including participants in clinical trials, and identifying patients at risk for or having a particular disorder or those most likely to respond to a particular therapeutic treatment, or for assessing or monitoring a patient’s response to a particular therapeutic treatment.

“Treatment” refers to therapy, prevention and prophylaxis and particularly refers to the administration of medicine or the performance of medical procedures with respect to a patient, for either prophylaxis (prevention) or to cure the infirmity or malady in the instance where the patient is afflicted.

“Agent” refers to all materials that may be used to prepare pharmaceutical and diagnostic compositions, or that may be compounds, agonists, antagonists, nucleic acids,

polypeptides, fragments, isoforms, variants, or other materials that may be used independently for such purposes, all in accordance with the present invention.

"Highly stringent conditions" refers to hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65 °C, and washing in 0.1xSSC/0.1% SDS at 68 °C (Ausubel F.M. et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & Sons, Inc., New York, at p. 2.10.3)

For some applications, less stringent conditions for duplex formation are required. As used herein "moderately stringent conditions" refers to washing in 0.2xSSC/0.1% SDS at 42 °C (Ausubel et al., 1989, supra).

In the context of the present invention, the term "sample" can refer to a sample obtained from any source, such as a serum sample, a CSF sample or a tissue sample, e.g. brain tissue. In addition, as used in this specification and the appended claims, the singular forms "a", "an", and "the" include plural references unless the context clearly dictates otherwise. Thus for example, references to "Alzheimer disease-associated protein" includes one or more of such proteins, reference to "the method" includes one or more methods, and/or steps of the type described herein and/or which will become apparent to those persons skilled in the art upon reading this disclosure, references to a "sample" includes sets of more than one sample and so forth.

"Cerebrospinal fluid", (CSF), refers to the fluid that surrounds the bulk of the central nervous system, as described in Physiological Basis of Medical Practice (J.B. West, ed., Williams and Wilkins, Baltimore, MD 1985). CSF includes ventricular CSF and lumbar CSF.

"Two-dimensional electrophoresis" (2D-electrophoresis) means a technique comprising isoelectric focusing, followed by denaturing electrophoresis; this generates a two-dimensional gel (2D-gel) containing a plurality of separated proteins.

5.2 The "Preferred Technology"

Preferably, the step of denaturing electrophoresis uses polyacrylamide electrophoresis in the presence of sodium dodecyl sulfate (SDS-PAGE). Especially preferred are the highly accurate and automatable methods and apparatus ("the Preferred Technology") described in WO 98/23950 and in U.S. Patent Nos 6,064,654, and 6,278,794, each of which is incorporated herein by reference in its entirety with particular reference to

the protocol at pages 23-35 of WO 98/23950. Briefly, the Preferred Technology provides efficient, computer-assisted methods and apparatus for identifying, selecting and characterising biomolecules (e.g. proteins, including glycoproteins) in a biological sample. A two-dimensional array is generated by separating biomolecules on a two-dimensional gel according to their electrophoretic mobility and isoelectric point. A computer-generated digital profile of the array is generated, representing the identity, apparent molecular weight, isoelectric point, and relative abundance of a plurality of biomolecules detected in the two-dimensional array, thereby permitting computer-mediated comparison of profiles from multiple biological samples, as well as computer aided excision of separated proteins of interest.

A preferred scanner for detecting fluorescently labeled proteins is described in WO 96/36882 and in the Ph.D. thesis of David A. Basiji, entitled "Development of a High-throughput Fluorescence Scanner Employing Internal Reflection Optics and Phase-sensitive Detection (Total Internal Reflection, Electrophoresis)", University of Washington (1997), Volume 58/12-B of Dissertation Abstracts International, page 6686, the contents of each of which are incorporated herein by reference. These documents describe an image scanner designed specifically for automated, integrated operation at high speeds. The scanner can image gels that have been stained with fluorescent dyes or silver stains, as well as storage phosphor screens. The Basiji thesis provides a phase-sensitive detection system for discriminating modulated fluorescence from baseline noise due to laser scatter or homogeneous fluorescence, but the scanner can also be operated in a non-phase-sensitive mode. This phase-sensitive detection capability would increase the sensitivity of the instrument by an order of magnitude or more compared to conventional fluorescence imaging systems. The increased sensitivity would reduce the sample-preparation load on the upstream instruments while the enhanced image quality simplifies image analysis downstream in the process.

A more highly preferred scanner is a modified version of the above described scanner. In the preferred scanner, the gel is transported through the scanner on a precision lead-screw drive system. This is preferable to laying the glass plate on the belt-driven system that is described in the Basiji thesis, as it provides a reproducible means of accurately transporting the gel past the imaging optics.

In the preferred scanner, the gel is secured against three alignment stops that rigidly hold the glass plate in a known position. By doing this in conjunction with the above

precision transport system, the absolute position of the gel can be predicted and recorded. This ensures that co-ordinates of each feature on the gel can be determined more accurately and communicated, if desired, to a cutting robot for excision of the feature. In the preferred scanner, the carrier that holds the gel has four integral fluorescent markers for use to correct the image geometry. These markers are a quality control feature that confirms that the scanning has been performed correctly.

In comparison to the scanner described in the Basiji thesis, the optical components of the preferred scanner have been inverted. In the preferred scanner, the laser, mirror, waveguide and other optical components are above the glass plate being scanned. The scanner described in the Basiji thesis has these components underneath. In the preferred scanner, the glass plate is mounted onto the scanner gel side down, so that the optical path remains through the glass plate. By doing this, any particles of gel that may break away from the glass plate will fall onto the base of the instrument rather than into the optics. This does not affect the functionality of the system, but increases its reliability.

Still more preferred is a modified version of the preferred scanner, in which the signal output is digitised to the full 16-bit data without any peak saturation or without square root encoding of the signal. A compensation algorithm has also been applied to correct for any variation in detection sensitivity along the path of the scanning beam. This variation is due to anomalies in the optics and differences in collection efficiency across the waveguide. A calibration is performed using a perspex plate with an even fluorescence throughout. The data received from a scan of this plate are used to determine the multiplication factors needed to increase the signal from each pixel level to a target level. These factors are then used in subsequent scans of gels to remove any internal optical variations.

5.3 Alzheimer's Disease-Associated Features (AFs)

In one aspect of the invention, two-dimensional electrophoresis is used to analyze CSF from a subject, preferably a living subject, in order to detect or quantify the expression of one or more Alzheimer's Disease-Associated Features (AFs) for screening, treatment or diagnosis of AD.

By way of example and not of limitation, using the Preferred Technology, a number of samples from subjects having AD and samples from subjects free from AD are separated by two-dimensional electrophoresis, and the fluorescent digital images of the resulting gels are matched to a chosen representative primary master gel image. This process allows any

gel feature, characterized by its pI and MW, to be identified and examined on any gel of the study. In particular, the amount of protein present in a given feature can be measured in each gel; this feature abundance can be averaged amongst gels from similar samples (e.g. gels from samples from subjects having AD). Finally, statistical analyses can be conducted on the thus created sample sets, in order to compare 2 or more sample sets to each other.

In accordance with an aspect of the present invention, the AFs disclosed herein have been identified by comparing CSF samples from subjects having AD against CSF samples from subjects free from AD. Subjects free from AD include subjects with no known disease or condition (normal subjects) and subjects with diseases (including neurological and neurodegenerative diseases) other than AD.

Two groups of AFs have been identified through the methods and apparatus of the Preferred Technology. The first group consists of AFs that are decreased in the CSF of subjects having AD as compared with the CSF of subjects free from AD. These AFs can be described by apparent molecular weight (MW) and isoelectric point (pI) as provided in Table I.

Table I. AFs Decreased in CSF of Subjects Having AD

(a) AFs Identified From Mastergroup Analysis

Table I (a)

AF#	Fold Decrease [#]	pI	MW (Da)	p value*
AF-200	2.217	6.32	18300	0.000 ⁽¹⁾
AF-201	2.132	8.32	63264	0.000 ⁽¹⁾
AF-202	2.132	8.32	63264	0.000 ⁽¹⁾
AF-203	1.991	5.53	74605	0.000 ⁽¹⁾
AF-204	1.955	7.97	60878	0.000 ⁽¹⁾
AF-205	1.876	6.82	61514	0.000 ⁽¹⁾
AF-206	1.807	5.55	28801	0.000 ⁽¹⁾
AF-207	1.790	6.65	33533	0.000 ⁽¹⁾
AF-208	1.747	5.75	29376	0.000 ⁽¹⁾
AF-209	1.741	8.4	59188	0.000 ⁽¹⁾
AF-210	1.674	5.96	29414	0.000 ⁽¹⁾
AF-211	1.673	8.07	42384	0.000 ⁽¹⁾
AF-212	1.670	6.37	62904	0.000 ⁽¹⁾

Table I (a)

AF#	Fold Decrease[#]	pI	MW (Da)	p value*
AF-213	1.669	6.53	62318	0.000 ⁽¹⁾
AF-214	1.658	4.28	18482	0.000 ⁽¹⁾
AF-215	1.617	8.01	44616	0.004 ⁽¹⁾
AF-216	1.603	6.73	56999	0.000 ⁽¹⁾
AF-217	1.572	6.4	26203	0.000 ⁽¹⁾
AF-218	1.558	6.74	33401	0.000 ⁽¹⁾
AF-219	1.555	7.93	63355	0.000 ⁽¹⁾
AF-220	1.555	6.76	54345	0.000 ⁽¹⁾
AF-221	1.542	5.51	139627	0.000 ⁽¹⁾
AF-222	1.532	5.62	132758	0.000 ⁽¹⁾
AF-223	1.513	5.67	131516	0.000 ⁽¹⁾
AF-224	1.513	6.86	49289	0.000 ⁽¹⁾
AF-225	1.495	10.85	58257	0.000 ⁽¹⁾
AF-226	1.448	7.23	19847	0.000 ⁽¹⁾
AF-227	1.444	4.38	21160	0.000 ⁽¹⁾
AF-228	1.444	4.38	21160	0.000 ⁽¹⁾
AF-229	1.431	5.29	29663	0.000 ^(1,4)
AF-230	1.392	5.55	135815	0.005 ^(1,4)
AF-231	1.349	7.72	175333	0.000 ⁽¹⁾
AF-232	1.341	4.41	42104	0.000 ⁽¹⁾
AF-233	1.326	5.19	46876	0.000 ⁽¹⁾
AF-234	1.320	5.04	18662	0.000 ^(1,4)
AF-235	1.301	5.38	62756	0.000 ⁽¹⁾
AF-236	1.285	7.35	64620	0.000 ⁽¹⁾
AF-237	1.283	7.86	59059	0.000 ⁽¹⁾
AF-238	1.276	8.1	34846	0.000 ^(1,4)
AF-239	1.269	6.05	184171	0.005 ⁽¹⁾
AF-240	1.264	5.01	46760	0.000 ⁽¹⁾
AF-241	1.262	5.46	16541	0.000 ⁽¹⁾
AF-242	1.258	4.48	34420	0.000 ⁽¹⁾

Table I (a)

AF#	Fold Decrease[#]	pI	MW (Da)	p value*
AF-243	1.256	5.25	119795	0.000 ⁽¹⁾
AF-244	1.251	7.12	54519	0.006 ⁽¹⁾
AF-245	1.244	4.87	49219	0.000 ^(1,4)
AF-246	1.233	4.24	73972	0.018 ⁽¹⁾
AF-247	1.232	6.46	92692	0.017 ⁽¹⁾
AF-248	1.222	5.52	39355	0.002 ⁽¹⁾
AF-249	1.210	4.25	20787	0.008 ⁽¹⁾
AF-250	1.210	7.88	14371	0.535 ⁽³⁾
AF-251	1.206	4.95	54997	0.001 ⁽¹⁾
AF-252	1.205	4.53	12315	0.004 ⁽¹⁾
AF-253	1.200	5.36	22610	0.000 ⁽¹⁾
AF-254	1.199	7.67	31952	0.045 ⁽¹⁾
AF-255	1.196	5.92	185771	0.048 ⁽¹⁾
AF-256	1.187	5.04	185128	0.012 ⁽¹⁾
AF-257	1.179	8.19	16738	0.01 ⁽¹⁾
AF-258	1.178	5.47	15411	0.0001 ^(1,2)
AF-259	1.167	4.81	30575	0.024 ⁽¹⁾
AF-260	1.166	4.93	154156	0.069 ⁽¹⁾
AF-261	1.152	6.71	16524	0.022 ⁽¹⁾
AF-262	1.151	5.81	114225	0.000 ⁽¹⁾
AF-263	1.150	4.43	16818	0.001 ⁽¹⁾
AF-264	1.146	5.12	184512	0.045 ⁽¹⁾
AF-265	1.137	6.13	87723	0.033 ⁽¹⁾
AF-266	1.131	6.47	35322	0.018 ⁽¹⁾
AF-267	1.130	5.25	179949	0.074 ^(1,3)
AF-268	1.130	9.25	17981	0.032 ⁽¹⁾
AF-269	1.125	6.58	170682	0.011(1)
AF-270	1.119	5.76	45729	0.123 ⁽³⁾
AF-271	1.113	4.99	63787	0.044 ⁽¹⁾
AF-272	1.103	6.38	37769	0.085 ⁽¹⁾

Table I (a)

AF#	Fold Decrease [#]	pI	MW (Da)	p value*
AF-273	1.100	4.62	82833	0.101 ⁽³⁾
AF-274	1.100	9.38	35170	0.232 ⁽³⁾
AF-275	1.099	5.76	189295	0.049 ⁽¹⁾
AF-276	1.099	6.49	43668	0.132 ⁽³⁾
AF-277	1.099	6.05	29055	0.256 ⁽³⁾
AF-278	1.097	5.67	21864	0.003 ⁽¹⁾
AF-279	1.092	5.39	95475	0.098 ⁽¹⁾
AF-280	1.089	5.3	102319	0.142 ⁽³⁾
AF-281	1.087	6.84	151156	0.026 ⁽¹⁾
AF-282	1.080	6.03	40754	0.146 ⁽³⁾
AF-283	1.077	6.22	70359	0.066 ⁽¹⁾
AF-284	1.076	5.84	187223	0.121 ⁽³⁾
AF-285	1.075	5.33	183136	0.255 ⁽³⁾
AF-286	1.072	6.36	20758	0.148 ⁽³⁾
AF-287	1.069	6.79	14728	0.0001 ^(1,2)
AF-288	1.067	5.22	37843	0.679 ⁽³⁾
AF-289	1.065	6.18	66668	0.066 ⁽¹⁾
AF-290	1.065	5.78	19705	0.150 ⁽³⁾
AF-291	1.063	4.88	135547	0.269 ⁽³⁾
AF-292	1.062	5.18	184053	0.338 ⁽³⁾
AF-293	1.061	5.79	21435	0.658 ⁽³⁾
AF-294	1.059	6.38	21221	0.343 ⁽³⁾
AF-295	1.052	6.57	114704	0.389 ⁽³⁾
AF-296	1.050	6.2	35622	0.119 ⁽³⁾
AF-297	1.046	5.91	16055	0.349 ⁽³⁾
AF-298	1.037	4.98	91522	0.445 ⁽³⁾
AF-299	1.032	4.61	29548	0.464 ⁽³⁾
AF-300	1.027	6.19	174157	0.538 ⁽³⁾
AF-301	1.025	6.46	25118	0.705 ⁽³⁾
AF-302	1.025	5	18011	0.642 ⁽³⁾

Table I (a)

AF#	Fold Decrease [#]	pI	MW (Da)	p value*
AF-303	1.019	6.13	175113	0.0001 ^(1,2)
AF-304	1.014	5.97	60157	0.902 ⁽³⁾
AF-305	1.013	4.63	21103	0.873 ⁽³⁾
AF-306	1.010	6.58	27953	0.836 ⁽³⁾
AF-307	1.006	6.79	30719	0.969 ⁽³⁾
AF-308	1.005	5.57	179672	0.924 ⁽³⁾
AF-309	1.000	10.69	18098	0.998 ⁽³⁾

* The statistical technique used to calculate a given p value is indicated by a footnote for each p value. The statistical techniques used for the group analysis were comparisons between patients with AD and control patients with no AD. Analysis included (1) a linear model, controlling for age and gender; (2) Stepwise multivariate linear discriminant analysis; (3) nearest neighbor and (4) longitudinal analysis.

(b) AFs Identified By Mastergroup Analysis Which Can Be Used In Combination With The Proteins Of Table I (a) and Table II (a)

Table I (b)

AF#	Fold Decrease [#]	pI	MW (Da)	p value*
AF-1	1.41	4.79	150081	0.001695 ⁽¹⁾
AF-2	1.34	4.28	21349	0.000133 ⁽¹⁾
AF-3	1.47	8.10	34846	0.000083 ⁽¹⁾
AF-4	1.56	4.38	21160	0.001192 ⁽¹⁾
AF-5	1.51	7.34	36554	0.000010 ⁽¹⁾
AF-6	1.46	4.91	29812	0.000003 ⁽¹⁾
AF-7	1.24	4.25	20787	0.003558 ⁽¹⁾
AF-8	1.44	4.93	187927	0.002221 ⁽¹⁾
AF-9	1.34	5.21	136768	0.000799 ⁽¹⁾
AF-10	1.30	5.19	17694	0.000400 ⁽¹⁾
AF-13	1.37	6.01	184530	0.000100 ⁽¹⁾
AF-14	1.52	4.72	63166	0.009387 ⁽¹⁾
AF-15	1.38	4.47	38970	0.000437 ⁽¹⁾

Table I (b)

AF#	Fold Decrease [#]	pI	MW (Da)	p value*
AF-16	1.22	5.19	46876	0.000697 ⁽¹⁾
AF-17	1.38	5.82	50294	0.000126 ⁽¹⁾
AF-18	1.30	4.87	49219	0.020661 ⁽¹⁾
AF-19	1.24	4.82	12454	0.00146 ⁽¹⁾
AF-20	1.30	4.43	16818	0.000322 ⁽¹⁾
AF-21	1.30	5.40	141094	0.000560 ⁽¹⁾
AF-22	1.36	4.93	133773	0.011000 ⁽¹⁾
AF-23	1.21	4.50	32473	0.000209 ⁽⁴⁾
AF-24	1.20	5.31	46663	0.000871 ⁽¹⁾
AF-25	1.19	5.68	36700	0.00251 ⁽¹⁾
AF-26	1.31	8.11	32305	0.002204 ⁽¹⁾
AF-27	1.26	5.33	141371	0.010447 ⁽¹⁾
AF-28	1.06	5.13	158568	0.000100 ⁽¹⁾
AF-29	1.27	9.22	47059	0.000028 ⁽¹⁾
AF-30	1.13	5.67	48057	0.000100 ⁽¹⁾
AF-31	1.15	6.07	91258	0.012712 ⁽⁴⁾
AF-32	1.15	6.17	48958	0.008321 ⁽⁴⁾
AF-33	1.06	4.41	42104	0.000126 ⁽¹⁾
AF-34	1.14	4.54	145408	0.017268 ⁽⁴⁾
AF-35	1.22	5.21	18623	0.001094 ⁽¹⁾
AF-36	1.17	5.78	14416	0.010744 ⁽¹⁾
AF-37	1.29	6.91	33523	0.000087 ⁽¹⁾
AF-38	1.18	6.47	29535	0.002759 ⁽¹⁾
AF-39	1.30	7.50	35510	0.002858 ⁽¹⁾
AF-40	1.22	7.29	38617	0.001187 ⁽¹⁾
AF-41	1.11	5.85	17345	0.016690 ⁽⁴⁾
AF-42	1.10	5.04	18662	0.002252 ⁽¹⁾
AF-43	1.13	9.83	14065	0.003303 ⁽¹⁾
AF-44	1.10	6.63	102328	0.020753 ⁽⁴⁾

Table I (b)

AF#	Fold Decrease [#]	pI	MW (Da)	p value*
AF-45	1.09	6.04	46998	0.031910 ⁽⁴⁾
AF-46	1.09	4.71	19802	0.008437 ⁽⁴⁾
AF-47	1.09	5.99	49664	0.002187 ⁽¹⁾
AF-48	1.32	5.32	122332	0.006582 ⁽¹⁾
AF-49	1.07	6.94	27576	0.010068 ⁽⁴⁾
AF-50	1.07	6.82	71337	0.035409 ⁽⁴⁾
AF-51	1.04	5.70	34388	0.006156 ⁽⁴⁾
AF-76	1.11	5.59	45537	0.001973 ⁽¹⁾
AF-149	1.15	4.82	190721	0.003541 ⁽¹⁾
AF-150	1.24	6.87	157592	0.000100 ⁽¹⁾
AF-152	1.09	5.04	81703	0.002800 ⁽¹⁾
AF-154	1.06	5.03	67307	0.000100 ⁽¹⁾
AF-155	1.38	9.21	64021	0.000070 ⁽¹⁾
AF-156	9.75	4.36	58083	0.001568 ⁽¹⁾
AF-159	1.18	5.08	52008	0.000100 ⁽¹⁾
AF-160	1.06	5.76	45729	0.004123 ⁽¹⁾
AF-162	1.23	5.47	38663	0.001073 ⁽¹⁾
AF-163	1.39	4.45	34879	0.001228 ⁽¹⁾
AF-164	1.51	5.00	33485	0.01179 ⁽¹⁾
AF-169	1.00	8.00	34362	0.000004 ⁽¹⁾
AF-170	1.38	5.41	31886	0.005600 ⁽¹⁾
AF-172	1.53	6.71	28747	0.01051 ⁽¹⁾
AF-173	10.91	7.67	27476	0.003738 ⁽¹⁾
AF-174	1.03	4.67	27811	0.002423 ⁽¹⁾
AF-175	1.03	5.33	24936	0.044270 ⁽¹⁾
AF-176	1.15	4.86	22248	0.012144 ⁽¹⁾
AF-177	1.14	4.63	21103	0.006564 ⁽¹⁾
AF-178	1.08	6.03	22247	0.05097 ⁽¹⁾
AF-181	1.21	5.72	16336	0.004745 ⁽¹⁾

Table I (b)

AF#	Fold Decrease [#]	pI	MW (Da)	p value*
AF-183	1.44	10.36	11160	0.000270 ⁽¹⁾
AF-184	1.08	5.31	48769	0.004689 ⁽¹⁾
AF-186	2.76	4.71	29693	0.002446 ⁽¹⁾
AF-187	2.09	4.93	154156	0.000750 ⁽¹⁾
AF-188	1.35	5.52	39355	0.007175 ⁽¹⁾
AF-189	1.29	6.79	30719	0.000377 ⁽¹⁾
AF-190	1.26	5.29	29663	0.000178 ⁽¹⁾
AF-191	1.12	5.31	46663	0.000654 ⁽¹⁾

* The statistical technique used to calculate a given p value is indicated by a footnote for each p value. The statistical techniques used for these group analyzes were (1) a linear model, controlling for age and gender; (2) classification trees; (3) a logistic regression model and (4) longitudinal analysis.

5 # Fold changes reported here are those calculated before adjustment for age and gender.

(c) AFs Identified From Pooled Gel Analysis Which Can Be Used In Combination With The Proteins From Table I (a) and Table II (a)

Table I (c)

AF#	Fold Decrease [#]	pI	MW (Da)
AF-78	2.80	5.59	158937
AF-79	2.62	5.52	142378
AF-80	2.78	5.56	142378
AF-81	2.60	5.43	78299
AF-82	7.25	6.69	74838
AF-83	4.07	6.81	71920
AF-84	4.44	6.94	73402
AF-85	16.77	7.10	73878
AF-86	4.86	5.24	67676
AF-87	4.17	5.95	64179

Table I (c)

AF#	Fold Decrease[#]	pI	MW (Da)
AF-88	6.48	5.36	66979
AF-89	2.64	5.39	65155
AF-90	4.81	7.61	62945
AF-91	3.10	8.16	56352
AF-92	2.63	6.18	50860
AF-93	6.33	6.28	49268
AF-94	11.17	4.38	45882
AF-95	9.28	5.60	46036
AF-96	11.07	4.43	44966
AF-98	5.51	6.72	44664
AF-99	10.1	5.92	44365
AF-100	15.39	6.08	44068
AF-101	9.82	4.47	44216
AF-102	3.14	6.02	44216
AF-103	3.57	5.93	42722
AF-104	22.78	5.09	42184
AF-105	4.11	5.19	42184
AF-107	13.3	7.26	33226
AF-108	19.0	7.54	33136
AF-110	9.21	5.39	28237
AF-111	14.85	5.68	27835
AF-112	2.60	6.00	20681
AF-114	8.98	6.80	18741
AF-115	2.85	6.04	17422
AF-116	6.85	6.68	14031
AF-117	2.90	4.65	13983
AF-118	15.19	6.94	11739
AF-119	16.27	7.23	11699

These features were identified as having a differential presence that is significant on the basis of having at least a 2-fold difference in mean intensity (*i.e.* a fold change threshold of at least 2) between Alzheimer's CSF and normal CSF.

- 5 The second group consists of AFs that are increased in the CSF of subjects having AD as compared with the CSF of subjects free from AD. These AFs can be described by apparent molecular weight (MW) and isoelectric point (pI) as provided in Table II.

Table II. AFs Increased in CSF of Subjects Having AD

- 10 (a) *AFs Identified From Mastergroup Analysis*

Table II(a)

AF#	Fold Increase [#]	pI	MW (Da)	p value*
AF-310	1.888	4.65	30092	0.001 ⁽¹⁾
AF-311	1.752	5.84	69278	0.000 ⁽¹⁾
AF-312	1.337	5.51	20016	0.014 ⁽¹⁾
AF-313	1.211	6.2	76215	0.006 ⁽¹⁾
AF-314	1.208	5.65	80079	0.003 ⁽¹⁾
AF-315	1.161	5.75	27709	0.015 ⁽¹⁾
AF-316	1.158	4.66	14580	0.000 ⁽¹⁾
AF-317	1.153	4.82	12454	0.003 ⁽¹⁾
AF-318	1.142	5.88	68454	0.002 ⁽¹⁾
AF-319	1.135	7.53	39490	0.027 ⁽¹⁾
AF-320	1.131	6.52	59328	0.26 ⁽³⁾
AF-321	1.131	4.94	12487	0.014 ⁽¹⁾
AF-322	1.125	5.78	81219	0.046 ⁽¹⁾
AF-323	1.109	5.79	148765	0.124 ⁽³⁾
AF-324	1.103	4.69	59610	0.108 ⁽³⁾
AF-325	1.094	10.5	28059	0.0013 ^(1,2)
AF-326	1.088	5.47	38663	0.284 ⁽³⁾
AF-327	1.084	5.59	44069	0.121 ⁽³⁾
AF-328	1.072	5.39	90721	0.083 ⁽¹⁾
AF-329	1.069	5.28	42179	0.201 ⁽³⁾

Table II(a)

AF#	Fold Increase [#]	pI	MW (Da)	p value*
AF-330	1.064	5.88	28511	0.336 ⁽³⁾
AF-331	1.064	4.79	18423	0.226 ⁽³⁾
AF-332	1.061	6.65	39644	0.682 ⁽³⁾
AF-333	1.049	5.29	61283	0.427 ⁽³⁾
AF-334	1.044	6.67	47683	0.264 ⁽³⁾
AF-335	1.032	4.82	23513	0.706 ⁽³⁾
AF-336	1.025	5.94	35737	0.643 ⁽³⁾
AF-337	1.021	6.82	51089	0.683 ⁽¹⁾
AF-338	1.006	5.7	43923	0.934 ⁽³⁾
AF-339	1.005	6.41	32196	0.917 ⁽³⁾

* The statistical technique used to calculate a given p value is indicated by a footnote for each p value. The statistical techniques used for the group analysis were comparisons between patients with AD and control patients with no AD. Analysis included (1) a linear model, controlling for age and gender; (2) Stepwise multivariate linear discriminant analysis; (3) nearest neighbor and (4) longitudinal analysis.

5

(b) AFs Identified By Mastergroup Analysis Which Can Be Used In Combination With The Proteins Of Table I (a) and Table II (a)

Table II (b)

AF#	Fold Increase [#]	pI	MW (Da)	p value*
AF-52	2.81	6.30	32573	0.000009 ⁽⁴⁾
AF-53	1.80	5.84	45302	0.016106 ⁽⁴⁾
AF-54	1.76	5.12	17520	0.003235 ⁽¹⁾
AF-55	1.29	8.10	12361	0.000482 ⁽¹⁾
AF-56	1.49	8.56	52128	0.005771 ⁽¹⁾
AF-57	1.46	6.30	68549	0.000274 ⁽¹⁾
AF-58	1.40	5.01	14507	0.01182 ⁽¹⁾
AF-59	1.37	6.74	33401	0.001351 ⁽¹⁾
AF-60	1.38	5.39	33873	0.009818 ⁽¹⁾

Table II (b)

AF#	Fold Increase#	pI	MW (Da)	p value*
AF-61	1.34	6.76	54345	Bag Tree 4 Analysis ⁽²⁾
AF-62	1.31	6.60	31004	0.000027 ⁽¹⁾
AF-63	1.24	5.97	14897	0.10696 ⁽¹⁾
AF-64	1.20	6.67	68119	0.000731 ⁽¹⁾
AF-65	1.22	7.19	58620	0.005833 ⁽⁴⁾
AF-66	1.09	10.05	30092	0.000100 ⁽¹⁾
AF-67	1.21	5.02	13735	0.006391 ⁽⁴⁾
AF-68	1.21	9.06	35351	0.003575 ⁽¹⁾
AF-69	1.19	5.01	46760	0.005125 ⁽⁴⁾
AF-70	1.19	8.91	38789	0.021552 ⁽⁴⁾
AF-71	1.20	6.44	68579	0.003848 ⁽¹⁾
AF-72	1.15	5.00	43788	0.014917 ⁽⁴⁾
AF-73	1.19	5.21	31615	0.000008 ⁽¹⁾
AF-74	1.14	6.19	51934	0.054917 ⁽¹⁾
AF-75	1.12	5.03	33671	0.002399 ⁽¹⁾
AF-77	1.09	6.41	32196	0.035148 ⁽⁴⁾
AF-151	1.13	5.28	137531	Bag Tree 1 Analysis ⁽²⁾
AF-153	1.23	9.85	69630	0.004906 ⁽¹⁾
AF-157	1.70	4.99	55449	0.006272 ⁽¹⁾
AF-161	1.00	5.18	44404	0.000600 ⁽¹⁾
AF-165	1.88	7.17	34230	0.000035 ⁽¹⁾
AF-166	1.20	8.54	33657	0.000009 ⁽¹⁾
AF-167	1.31	5.69	33621	0.005400 ⁽¹⁾
AF-168	1.00	7.66	33920	0.000013 ⁽¹⁾
AF-171	1.10	4.98	29658	0.004242 ⁽¹⁾
AF-179	1.64	5.26	20115	Stepwise Analysis ⁽¹⁾
AF-180	1.62	6.17	16255	0.005047 ⁽¹⁾

Table II (b)

AF#	Fold Increase#	pI	MW (Da)	p value*
AF-182	1.37	4.89	13651	0.005380 ⁽¹⁾
AF-185	6.00	5.32	40323	0.005520 ⁽¹⁾
AF-192	1.04	5.38	62756	0.000213 ⁽¹⁾

* The statistical technique used to calculate a given p value is indicated by a footnote for each p value. The statistical techniques used for these group analyzes were (1) a linear model, controlling for age and gender; (2) classification trees; (3) a logistic regression model and (4) longitudinal analysis.

- 5 # Fold changes reported here are those calculated before adjustment for age and gender.

(c) AFs Identified From Pooled Gel Analysis Which Can Be Used In Combination With The Proteins From Table I (a) and Table II (a)

Table II (c)

AF#	Fold Increase#	pI	MW (Da)
AF-121	11.7	5.42	105108
AF-122	2.20	5.27	71060
AF-123	6.35	7.31	64933
AF-124	6.86	7.47	64736
AF-125	2.34	4.77	61297
AF-126	48.84	4.11	60374
AF-127	6.79	4.98	59649
AF-128	2.36	6.60	57865
AF-129	2.90	5.29	54625
AF-130	2.94	5.08	51880
AF-131	5.19	6.54	50944
AF-132	3.41	4.72	47414
AF-133	3.08	5.12	44068
AF-134	2.34	5.00	43516
AF-137	4.41	4.98	36855
AF-139	4.11	5.00	34295

Table II (c)

AF#	Fold Increase#	pI	MW (Da)
AF-140	110.32	6.80	32080
AF-141	3.58	7.50	28440
AF-142	2.66	6.75	27279
AF-143	5.68	7.44	26066
AF-144	2.71	6.56	20744
AF-145	4.43	4.76	18069
AF-146	2.18	4.94	12790
AF-147	2.29	4.81	12790
AF-148	5.04	5.32	11382

These features were identified as having a differential presence that is significant on the basis of having at least a 2-fold difference in mean intensity (*i.e.* a fold change threshold of at least 2) between Alzheimer's CSF and normal CSF.

5 For any given AF, the signal obtained upon analyzing CSF from subjects having AD relative to the signal obtained upon analyzing CSF from subjects free from AD will depend upon the particular analytical protocol and detection technique that is used. Accordingly, those skilled in the art will understand that any laboratory, based on the present description, can establish a suitable reference range for any AF in subjects free from AD according to

10 the analytical protocol and detection technique in use. In particular, at least one positive control CSF sample from a subject known to have AD or at least one negative control CSF sample from a subject known to be free from AD (and more preferably both positive and negative control samples) are included in each batch of test samples analyzed. In one

15 embodiment, the level of expression of a feature is determined relative to a background value, which is defined as the level of signal obtained from a proximal region of the image that (a) is equivalent in area to the particular feature in question; and (b) contains no substantial discernable protein feature.

In a preferred embodiment, the signal associated with an AF in the CSF of a subject (*e.g.*, a subject suspected of having or known to have AD) is normalized with reference to

20 one or more Expression Reference Features (ERFs) detected in the same 2D gel. As will be apparent to one of ordinary skill in the art, such ERFs may readily be determined by

comparing different samples using techniques and protocols such as the Preferred Technology. Suitable ERFs include (but are not limited to) that described in the following table.

5 Table III. Expression Reference Features

ERF#	pI	MW (Da)
ERF-1	5.94	18860
ERF-2	6.04	47450

As those of skill in the art will readily appreciate, the measured MW and pI of a given feature or protein isoform will vary to some extent depending on the precise protocol used for each step of the 2D electrophoresis and for landmark matching. As used herein, the terms "MW" and "pI" are defined, respectively, to mean the apparent molecular weight in Daltons and the apparent isoelectric point of a feature or protein isoform as measured in careful accordance with the Reference Protocol identified in Section 6 below. When the Reference Protocol is followed and when samples are run in duplicate or a higher number of replicates, variation in the measured mean pI of an AF or API is typically less than 3% and variation in the measured mean MW of an AF or API is typically less than 5%. Where the skilled artisan wishes to diverge from the Reference Protocol, calibration experiments should be performed to compare the MW and pI for each AF or protein isoform as detected (a) by the Reference Protocol and (b) by the divergent.

The AFs of the invention can be used, for example, for detection, treatment, diagnosis, or the drug development or pharmaceutical products. In one embodiment of the invention, CSF from a subject (*e.g.*, a subject suspected of having AD) is analyzed by 2D electrophoresis for quantitative detection of one or more of the AFs as defined in Table I (a), optionally in combination with one or more of the Afs as defined in Tables I (b) and (c). A decreased abundance of said one or more AFs in any suitable combination in the CSF from the subject relative to CSF from a subject or subjects free from AD (*e.g.*, a control sample or a previously determined reference range) indicates the presence of AD.

In a particular embodiment hereof, CSF from a subject (*e.g.*, a subject suspected of having AD) is analyzed by 2D electrophoresis for quantitative detection of one or more of the AFs are set forth herein in Table I (a). A decreased abundance of one or more in any suitable combination of such AFs in the CSF from the subject relative to CSF from a

subject or subjects free from AD (*e.g.*, a control sample or a previously determined reference range) indicates the presence of AD.

In another embodiment of the invention, CSF from a subject is analyzed by 2D electrophoresis for quantitative detection of one or more of the Afs as defined in Table II (a), optionally in combination with one or more Afs as defined in Tables II (b) and (c). An increased abundance of said one or more AFs in any suitable combination in the CSF from the subject relative to CSF from a subject or subjects free from AD (*e.g.*, a control sample or a previously determined reference range) indicates the presence of AD. In a particular embodiment hereof, CSF from a subject is analyzed by 2D electrophoresis for quantitative detection of one or more of the AFs set forth herein in Table II (a). An increased abundance of said one or more AFs in any suitable combination in the CSF from the subject relative to CSF from a subject or subjects free from AD (*e.g.*, a control sample or a previously determined reference range) indicates the presence of AD.

In yet another embodiment, CSF from a subject is analyzed by 2D electrophoresis for quantitative detection of (a) one or more AFs or any suitable combination of them, whose decreased abundance indicates the presence of AD, *i.e.*, preferably AFs as set forth in Tables I (a), optionally in combination with the AFs set out in Tables I (b), or I (c); and (b) one or more AFs or any combination of them, whose increased abundance indicates the presence of AD *i.e.*, preferably AFs set forth in Tables II (a), optionally in combination with the AFs set out in Tables II (b), or II (c).

In yet another embodiment of the invention, CSF from a subject is analyzed by 2D electrophoresis for quantitative detection of one or more of the AFs set forth in Table I (a) or II (a), optionally in combination with the AFs set forth in Tables I (b), I (c), II (b), or II (c) wherein the ratio of the one or more AFs relative to an Expression Reference Feature (ERF) indicates that AD is present. In a specific embodiment, a decrease in one or more AF/ERF ratios in a sample being tested relative to the AF/ERF ratios in a control sample or a reference range indicates the presence of AD; the AFs set forth in Table I (a), optionally combined with those set forth in Tables I (b), or I (c) are suitable are suitable AFs for this purpose. In another specific embodiment, one may measure one or more AFs in a test sample, and compare them to an ERF, as a method for detecting the presence of AD. Thus, an increase in one or more AF/ERF ratios in a test sample relative to the AF/ERF ratios in a control sample or a reference range indicates the presence of AD, the AFs set forth in Table

II (a), optionally combined with the AFs set forth in Tables II (b) or II (c) are suitable AFs for this purpose.

In a further embodiment of the invention, CSF from a subject is analyzed by 2D electrophoresis for quantitative detection of (a) one or more AFs, or any suitable combination of them, whose decreased AF/ERF ratio(s) in a test sample relative to the AF/ERF ratio(s) in a control sample indicates the presence of AD, *i.e.*, preferably the AFs set forth in Table I (a), optionally combined with the AFs set forth in Tables I (b) or I (c); (b) one or more AFs, or any combination of them, whose increased AF/ERF ratio(s) in a test sample relative to the AF/ERF ratio(s) in a control sample indicates the presence of AD, *i.e.*, preferably the AFs as set forth in Table II (a), optionally combined with the AFs set forth in Tables II (b), or II (c).

In a preferred embodiment, CSF from a subject is analyzed for quantitative detection of a plurality of AFs.

5.4 Alzheimer's Disease-Associated Protein Isoforms (APIs)

In another aspect of the invention, CSF from a subject, is analyzed for quantitative detection of one or more Alzheimer's Disease-Associated Protein Isoforms (APIs), *e.g.* for screening, treatment or diagnosis of AD or for development of pharmaceutical products. As is well known in the art, a given protein may be expressed as one or more variants forms (isoforms) that differ in amino acid composition (*e.g.* as a result of alternative mRNA or premRNA processing, *e.g.* alternative splicing or limited proteolysis) or as a result of differential post-translational modification (*e.g.*, glycosylation, phosphorylation, acylation), or both, so that proteins of identical amino acid sequence can differ in their pI, MW, or both. "Alzheimer's Disease -Associated Protein Isoform" refers to a protein isoform that is differentially present in CSF from a subject having AD compared with CSF from a subject free from AD.

Two groups of APIs are described herein by the amino acid sequencing of AFs as depicted in Figure 2 and described above. APIs were isolated, subjected to proteolysis, and analyzed by mass spectrometry using in this instance the methods and apparatus of the Preferred Technology, it being understood that the preferred technology is set forth as representative but not restrictive of the invention. One skilled in the art can identify sequence information from proteins analyzed by mass spectrometry and/or tandem mass spectrometry using various spectral interpretation methods and database searching tools.

Examples of some of these methods and tools can be found at the Swiss Institute of Bioinformatics web site at <http://www.expasy.ch/>, and the European Molecular Biology Laboratory web site at www.mann.embl-heidelberg.de/Services/PeptideSearch/.

Identification of APIs was performed using the SEQUEST search program (Eng et al., 1994, J. Am. Soc. Mass Spectrom. 5:976-989) and the method described in PCT Application No. PCT/GB01/04034.

The first group comprises of APIs that are decreased in the CSF of subjects having AD as compared with the CSF of subjects free from AD. The amino acid sequences of peptides produced from these APIs by proteolysis using trypsin and identified by tandem mass spectrometry and database searching using the SEQUEST program are listed in Table IV, in addition to their corresponding pIs and MWs. For one API, the partial sequence information derived from tandem mass spectrometry was not found to be described in any known public database. This API is listed as 'NOVEL' in Table IV, and the partial amino acid sequence information derived from manually interpreting the MS/MS spectrum of tryptic peptides of this API as described in the Example *infra*, is given in Table IX.

Table IV. APIs Decreased in CSF of Subjects Having AD

(a) APIs Decreased in CSF of Subjects Having AD

AF#	API#	pI	MW	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
AF-201	API-375	8.32	63264	YGPFGPEMTNPLR VGYGPTFK LVGQLMDGLK	SEQ ID NO: 448 SEQ ID NO: 411 SEQ ID NO: 274
AF-204	API-300	7.97	60878	GQLVFMNR LNMGLTDLQGLR VGDTLNLNLR	SEQ ID NO: 166 SEQ ID NO: 254 SEQ ID NO: 408
AF-205	API-301	6.82	61514	LAQWQSFQLEGGLK QFSFPLSSEPFQGSYK NEDSLVFVQTDK	SEQ ID NO: 220 SEQ ID NO: 319 SEQ ID NO: 289
AF-206	API-302	5.55	28801	TYLHTYESEL VWNYFQR	SEQ ID NO: 392 SEQ ID NO: 433
AF-207	API-303	6.65	33533	AQGFTEDTLVFLPQTDK TMLLQPAGSLGYSYR	SEQ ID NO: 30 SEQ ID NO: 379
AF-208	API-304	5.75	29376	WVEELMK	SEQ ID NO: 441

AF#	API#	pI	MW	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
				TYLHTYESEL VWNYFQR	SEQ ID NO: 392 SEQ ID NO: 433
AF-209	API-305	8.4	59188	VGDTLNLNLR LNMGLTDLQGLR	SEQ ID NO: 408 SEQ ID NO: 254
AF-209	API-306	8.4	59188	SGFSFGFK LPGLFELGLSSQSDR EYESYSDFER LPLEYSYGEYR	SEQ ID NO: 345 SEQ ID NO: 257 SEQ ID NO: 114 SEQ ID NO: 259
AF-210	API-307	5.96	29414	WVEELMK TYLHTYESEL VWNYFQR	SEQ ID NO: 441 SEQ ID NO: 392 SEQ ID NO: 433
AF-211	API-308	8.07	42384	EVQGFESATFLGYFK QTQVSVLPEGGETPLFK HVVPNEVVQR	SEQ ID NO: 112 SEQ ID NO: 332 SEQ ID NO: 187
AF-212	API-309	6.37	62904	NEDSLVFVQTDK LAQWQSFQLEGGLK	SEQ ID NO: 289 SEQ ID NO: 220
AF-213	API-310	6.53	62318	QFSFPLSSEPFQGSYK NEDSLVFVQTDK	SEQ ID NO: 319 SEQ ID NO: 289
AF-214	API-376	4.28	18482	NSEPQDEGELFQGVDP	SEQ ID NO: 302
AF-214	API-377	4.28	18482	TQSSLVPALTDFVR	SEQ ID NO: 383
AF-215	API-311	8.01	44616	VGDTLNLNLR GQLVFMNR LNMGLTDLQGLR	SEQ ID NO: 408 SEQ ID NO: 166 SEQ ID NO: 254
AF-216	API-312	6.73	56999	QTQVSVLPEGGETPLFK EPGLQLWR HVVPNEVVQR	SEQ ID NO: 332 SEQ ID NO: 105 SEQ ID NO: 187
AF-218	API-313	6.74	33401	LSYEGETVK TSLEDFYLDEER	SEQ ID NO: 267 SEQ ID NO: 385
AF-219	API-378	7.93	63355	PGSTGTWNPSSER HPDEAAFFDTASTGK GLLDEVNQDFTNR GGSTSYGTGSETESPR ADSGEGDFLAEGGGVR	SEQ ID NO: 309 SEQ ID NO: 182 SEQ ID NO: 157 SEQ ID NO: 150 SEQ ID NO: 8
AF-220	API-314	6.76	54345	LPLEDGSGEVVLSR	SEQ ID NO: 258

AF#	API#	pI	MW	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
				TLYTPGSTVLRYR TVMVNLENPEGLPVK QELSEAEQATR	SEQ ID NO: 378 SEQ ID NO: 390 SEQ ID NO: 316
AF-224	API-379	6.86	49289	PDDLFPSTVVPLK EHSSLAFWK ATVVYQGER AGLLENGAVR	SEQ ID NO: 308 SEQ ID NO: 89 SEQ ID NO: 37 SEQ ID NO: 18
AF-225	API-315	10.85	58257	VGDTLNLNLR TTNLQGLNLLFSSR LNMGLTDLQGLR GQLVFMNR	SEQ ID NO: 408 SEQ ID NO: 387 SEQ ID NO: 254 SEQ ID NO: 166
AF-226	API-316	7.23	19847	AQGFTEDTLVFLPQTDK TMLLQPAGSLGSSYSYR	SEQ ID NO: 30 SEQ ID NO: 379
AF-227	API-380	4.38	21160	SGELEQEEER EEEEEEMAVVPQGLFR	SEQ ID NO: 344 SEQ ID NO: 81
AF-229	API-317	5.29	29663	VAQLPLSLK LVEAFGGATK NNYMYAR ELDVLQGR	SEQ ID NO: 397 SEQ ID NO: 270 SEQ ID NO: 300 SEQ ID NO: 95
AF-229	API-369	5.29	29663	AQGFTEDTIVFLPQTDK TMLLQPAGSLGSSYSYR	SEQ ID NO: 29 SEQ ID NO: 379
AF-230	API-318	5.55	135815	LPPNVVEESAR ALGYLNTGYQR	SEQ ID NO: 260 SEQ ID NO: 22
AF-231	API-319	7.72	175333	STGGLSVPGPMGPSGPR GESGPSGPAGPTGAR SLSKQLENLR NSVAYMDQETGNLK GETGPAGPAGPVGPVGA R	SEQ ID NO: 358 SEQ ID NO: 145 SEQ ID NO: 352 SEQ ID NO: 303 SEQ ID NO: 146
AF-232	API-320	4.41	42104	GLSAEPGWQAK EEEEEEMAVVPQGLFR	SEQ ID NO: 159 SEQ ID NO: 81
AF-233	API-321	5.19	46876	EGPVLLGR	SEQ ID NO: 88
AF-234	API-322	5.04	18662	FSGTWYAMAK LLVHNGYCDGR	SEQ ID NO: 128 SEQ ID NO: 249

AF#	API#	pI	MW	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
AF-235	API-323	5.38	62756	QTQVSVLPEGGETPLFK TGAQELLR YLETDPANR	SEQ ID NO: 332 SEQ ID NO: 370 SEQ ID NO: 455
AF-236	API-381	7.35	64620	PGSTGTWNPSSER HPDEAAFFDTASTGK GLLDEVNQDFTNR GGSTSYGTGSETESPR DSHSLTTNLMELLR ALTDMPQMR ADSGEGDFLAEGGGVR	SEQ ID NO: 309 SEQ ID NO: 182 SEQ ID NO: 157 SEQ ID NO: 150 SEQ ID NO: 70 SEQ ID NO: 24 SEQ ID NO: 8
AF-237	API-382	7.86	59059	YVLPNFEVK LNMGLTDLQGLR EMSGSPASGLPVK	SEQ ID NO: 462 SEQ ID NO: 254 SEQ ID NO: 101
AF-237	API-383	7.86	59059	TLYTPGSTVLRY LPLEDGSGEVVLSR	SEQ ID NO: 378 SEQ ID NO: 258
AF-237	API-384	7.86	59059	TNSFEGPVLDR SLMLHYEFLQR SGFSFGFK SFEGPVLDR SDLEVAHYK QALEEFQK NVESYTPQTQGK LPLEYSYGEYR LPGLFELGLSSQSDR EYESYSDFER	SEQ ID NO: 381 SEQ ID NO: 351 SEQ ID NO: 345 SEQ ID NO: 343 SEQ ID NO: 341 SEQ ID NO: 312 SEQ ID NO: 304 SEQ ID NO: 259 SEQ ID NO: 257 SEQ ID NO: 114
AF-238	API-324	8.1	34846	VLVWVNNLGEK LVNLYDSMPLR YLELFQR	SEQ ID NO: 419 SEQ ID NO: 280 SEQ ID NO: 454
AF-239	API-325	6.05	184171	VGFYESDVMGR ALGYLNTGYQR LPPNVVEESAR	SEQ ID NO: 409 SEQ ID NO: 22 SEQ ID NO: 260
AF-240	API-326	5.01	46760	ALVQQMEQLR TQVNTQAEQLR	SEQ ID NO: 25 SEQ ID NO: 384
AF-241	API-327	5.46	16541	GEVQAMLGQSTEELR	SEQ ID NO: 147

AF#	API#	pI	MW	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
				FWDYLR SELEEQLTPVAEETR KVEQAVETEPEPELR	SEQ ID NO: 134 SEQ ID NO: 342 SEQ ID NO: 215
AF-242	API-328	4.48	34420	EDEEEEEGENYQK GEAGAPGEEDLQGPTK	SEQ ID NO: 79 SEQ ID NO: 142
AF-243	API-329	5.25	119795	TYFEGER AFLFQDTPR	SEQ ID NO: 391 SEQ ID NO: 12
AF-244	API-330	7.12	54519	QELSEAEQATR TLYTPGSTVLYR	SEQ ID NO: 316 SEQ ID NO: 378
AF-245	API-331	4.87	49219	LEEELGDEAR LVLGMDVAASEFYR LGAEVYHTLK	SEQ ID NO: 225 SEQ ID NO: 276 SEQ ID NO: 233
AF-248	API-385	5.52	39355	WEAEPVYVQR LDVHWTR	SEQ ID NO: 438 SEQ ID NO: 223
AF-249	API-332	4.25	20787	EEEEEMA VVPQGLFR	SEQ ID NO: 81
AF-250	API-333	7.88	14371	VSFELFADK FEDENFLLK	SEQ ID NO: 427 SEQ ID NO: 118
AF-252	API-334	4.53	12315	ASEEEPEYGEELK GYPGVQAPEDLEWER	SEQ ID NO: 32 SEQ ID NO: 173
AF-253	API-335	5.36	22610	DYVSQFEGSALGK VQPYLDDFQK WQEEMELYR DEPPQSPWDR THLAPYSDEL R	SEQ ID NO: 76 SEQ ID NO: 425 SEQ ID NO: 440 SEQ ID NO: 52 SEQ ID NO: 372
AF-254	API-336	7.67	31952	AQGFTEDTLVFLPQTDK TMLLQPAGSLGSSYSYR	SEQ ID NO: 30 SEQ ID NO: 379
AF-255	API-337	5.92	185771	ALGYLNTGYQR LPPNVVEESAR	SEQ ID NO: 22 SEQ ID NO: 260
AF-256	API-338	5.04	185128	DGNPFYFTDHR VLAVNEVGR VEEVKPLEGR DGNPFYFTDHR	SEQ ID NO: 56 SEQ ID NO: 415 SEQ ID NO: 399 SEQ ID NO: 56
AF-257	API-339	8.19	16738	AQGFTEDTLVFLPQTDK TMLLQPAGSLGSSYSYR	SEQ ID NO: 30 SEQ ID NO: 379

AF#	API#	pI	MW	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
				SVVAPATDGGLNLTSTF LR	SEQ ID NO: 360
AF-258	API-340	5.47	15411	QAVETEPEPELR VEQAVETEPEPELR QQTEWQSGQR GEVQAMLGQSTEELR SELEEQLTPVAEETR KVEQAVETEPEPELR	SEQ ID NO: 313 SEQ ID NO: 403 SEQ ID NO: 327 SEQ ID NO: 147 SEQ ID NO: 342 SEQ ID NO: 215
AF-259	API-341	4.81	30575	AQGFTEDTLVFLPQTDK TMLLQPAGSLGSYSYR	SEQ ID NO: 30 SEQ ID NO: 379
AF-260	API-342	4.93	154156	LLMLFTDGGEER GYYYELPSLGALR	SEQ ID NO: 248 SEQ ID NO: 176
AF-261	API-386	6.71	16524	TYSYLNK LVVEWQLQDDK	SEQ ID NO: 393 SEQ ID NO: 281
AF-262	API-387	5.81	114225	YGPFGPEMTNPLR TYLHTYESEL LEDLHLLVER DLEHLTSLDFFR	SEQ ID NO: 448 SEQ ID NO: 392 SEQ ID NO: 224 SEQ ID NO: 58
AF-263	API-343	4.43	16818	GKLEEDSEVLMMMLK TQSSLVPALTDFVR	SEQ ID NO: 153 SEQ ID NO: 383
AF-264	API-344	5.12	184512	VLAVNEVGR SMEQNGPGLEYR VEEVKPLEGR DGNPFYFTDHR	SEQ ID NO: 415 SEQ ID NO: 353 SEQ ID NO: 399 SEQ ID NO: 56
AF-265	API-388	6.13	87723	YLETDPANR TGAQELLR QTQVSVLPEGGETPLFK PALPAGTEDTAK HVVVPNEVVVQR EPGLQLWR DPDQTDGLGLSYLSSH AGALNSNDAFVLK	SEQ ID NO: 455 SEQ ID NO: 370 SEQ ID NO: 332 SEQ ID NO: 307 SEQ ID NO: 187 SEQ ID NO: 105 SEQ ID NO: 63 SEQ ID NO: 15
AF-267	API-345	5.25	179949	VLAVNEVGR VEEVKPLEGR	SEQ ID NO: 415 SEQ ID NO: 399

AF#	API#	pI	MW	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
				DGNPFYFTDHR	SEQ ID NO: 56
AF-268	API-346	9.25	17981	AQGFTEDTLVFLPQTDK	SEQ ID NO: 30
				TMLLQPAGSLGSYSYR	SEQ ID NO: 379
AF-269	API-389	6.58	170682	RPYFPVAVGK	SEQ ID NO: 336
				NGFYPATR	SEQ ID NO: 294
				ELMENYNLALR	SEQ ID NO: 98
AF-270	API-390	5.76	45729	VPQVSTPTLVEVSR	SEQ ID NO: 422
				VPEVSTPTLVEVSR	SEQ ID NO: 421
				KVPQVSTPTLVEVSR	SEQ ID NO: 217
				HPDYSVLLLLR	SEQ ID NO: 184
				FQNALLVR	SEQ ID NO: 126
				AVMDDFAAFVEK	SEQ ID NO: 40
AF-271	API-347	4.99	63787	TVGSDTFYSFK	SEQ ID NO: 388
				YNSQNQSNNQFVLYR	SEQ ID NO: 458
AF-272	API-391	6.38	37769	LNHGLLYDEEK	SEQ ID NO: 252
				ELMENYNLALR	SEQ ID NO: 98
AF-274	API-348	9.38	35170	LVGGPMDASVEEEGVR	SEQ ID NO: 273
				ALDFAVGEYNK	SEQ ID NO: 20
AF-275	API-349	5.76	189295	LPPNVVEESAR	SEQ ID NO: 260
				ALGYLNTGYQR	SEQ ID NO: 22
AF-277	API-350	6.05	29055	VTELWQEVQR	SEQ ID NO: 431
				AVVVHAGEDDLGR	SEQ ID NO: 41
AF-278	API-351	5.67	21864	DYVSQFEGSALGK	SEQ ID NO: 76
				THLAPYSDELRL	SEQ ID NO: 372
AF-281	API-352	6.84	151156	LLVQAQPEWLK	SEQ ID NO: 250
				QGSTHWQTAR	SEQ ID NO: 321
				GPPGPPGGVVVR	SEQ ID NO: 165
AF-283	API-392	6.22	70359	YLETDPANR	SEQ ID NO: 455
				HVVPNEVVQR	SEQ ID NO: 187
AF-284	API-393	5.84	187223	YGAATFTR	SEQ ID NO: 447
				VGFYESDVMGR	SEQ ID NO: 409
				NEDSLVFVQTDK	SEQ ID NO: 289
				LPPNVVEESAR	SEQ ID NO: 260
				HYDGSYSTFGER	SEQ ID NO: 188

AF#	API#	pI	MW	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
				ALGYLNTGYQR	SEQ ID NO: 22
AF-285	API-394	5.33	183136	QYTDSTFR	SEQ ID NO: 334
				QSEDSTFYLGFR	SEQ ID NO: 328
				GAYPLSLEPLGVR	SEQ ID NO: 136
AF-285	API-395	5.33	183136	VLAVNEVGR	SEQ ID NO: 415
				TAVTANLDR	SEQ ID NO: 365
				SMEQNGPGLEYR	SEQ ID NO: 353
				PLTEESSTLGEGSK	SEQ ID NO: 310
				GDLYFANVEEK	SEQ ID NO: 139
AF-289	API-353	6.18	66668	ELSHLPSLYDYSAYR	SEQ ID NO: 100
				VLFYVDSEK	SEQ ID NO: 417
				DGFVQDEGTMFPVGK	SEQ ID NO: 55
AF-291	API-354	4.88	135547	GNLAGLTLR	SEQ ID NO: 161
				EGLDLQVLEDSGR	SEQ ID NO: 86
AF-295	API-396	6.57	114704	YGPFGPEMTNPLR	SEQ ID NO: 448
				VWNYFQR	SEQ ID NO: 433
AF-296	API-397	6.20	35622	LEVLYTPTAMLRPD	SEQ ID NO: 230
				GNPVPQQYLWEK	SEQ ID NO: 162
AF-299	API-355	4.61	29548	CPNPPVQENFDVNK	SEQ ID NO: 47
				NPNLPPETVDSLK	SEQ ID NO: 301
AF-300	API-398	6.19	174157	SSNLLLLLEHLK	SEQ ID NO: 357
				RPYFPVAVGK	SEQ ID NO: 336
				NGFYFATR	SEQ ID NO: 294
				NDFTWFK	SEQ ID NO: 288
				ELMENYNLALR	SEQ ID NO: 98
				EFDHNSNLR	SEQ ID NO: 84
AF-303	API-399	6.13	175113	SSNLLLLLEHLK	SEQ ID NO: 357
				ELMENYNLALR	SEQ ID NO: 98
AF-306	API-400	6.58	27953	YSSLAEAASK	SEQ ID NO: 459
				VLDALQALK	SEQ ID NO: 416
				LYPLANGNNQSPVDLK	SEQ ID NO: 283
AF-307	API-356	6.79	30719	VEYGFQVK	SEQ ID NO: 405
				LTQVLHFTK	SEQ ID NO: 268
				GLQDEDGYR	SEQ ID NO: 158

AF#	API#	pI	MW	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
				FACYYP	SEQ ID NO: 115
AF-308	API-357	5.57	179672	ELMENYNLALR NGFYFATR RPYFPVAVGK	SEQ ID NO: 98 SEQ ID NO: 294 SEQ ID NO: 336
AF-309	API-401	10.69	18098	LVGGPMDASVEEEGVR	SEQ ID NO: 273

(b) APIs which can be used in combination with the APIs in Table IV (a) and Table V (a)

AF#	API#	pI	MW (Da)	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
AF-1	API-47	4.79	150081	QPEYAVVQR GKPPPSFSWTR EDYICYAR	SEQ ID NO: 325 SEQ ID NO: 154 SEQ ID NO: 80
AF-1	API-242	4.79	150081	FVVTDGGITR IIMLFTDGGEER	SEQ ID NO: 133 SEQ ID NO: 195
AF-2	API-1	4.28	21349	EEEEEMAVVPQGLFR SGELEQEEER	SEQ ID NO: 81 SEQ ID NO: 344
AF-3	API-48	8.1	34846	YLELFQR VIVVWNNIGEK LVNIYDSMPLR	SEQ ID NO: 454 SEQ ID NO: 414 SEQ ID NO: 279
AF-5	API-49	7.34	36554	DCSGVSLHLTR	SEQ ID NO: 51
AF-6	API-2	4.91	29812	TEAYLEAIR	SEQ ID NO: 367
AF-8	API-194	4.93	187927	DGNPFYFTDHR	SEQ ID NO: 56
AF-9	API-3	5.21	136768	AETYESGVYQCTAR	SEQ ID NO: 11
AF-10	API-50	5.19	17694	SELEEQLTPVAEETR KVEQAVETEPEPELR GEVQAMLGQSTEELR FWDYLR	SEQ ID NO: 342 SEQ ID NO: 215 SEQ ID NO: 147 SEQ ID NO: 134
AF-10	API-51	5.19	17694	VNSDGGLVALR	SEQ ID NO: 420
AF-13	API-4	6.01	184530	LPPNVVEESAR VGFYESDVMGR HYDGSYSTFGER	SEQ ID NO: 260 SEQ ID NO: 409 SEQ ID NO: 188
AF-14	API-52	4.72	63166	EIGELYLPK	SEQ ID NO: 90

AF#	API#	pI	MW (Da)	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
				ADLSGITGAR	SEQ ID NO: 6
AF-14	API-243	4.72	63166	FEDGVLDPDYPR	SEQ ID NO: 119
AF-15	API-53	4.47	38970	ELDESLQVAER	SEQ ID NO: 94
AF-15	API-244	4.47	38970	TEVQLEHLSR	SEQ ID NO: 369
AF-16	API-54	5.19	46876	EGPVLILGR	SEQ ID NO: 87
AF-17	API-5	5.82	50294	VAMHLVCPSR YEAAPDPR TALASGGVLDASGDYR EPGEFALLR	SEQ ID NO: 394 SEQ ID NO: 442 SEQ ID NO: 363 SEQ ID NO: 104
AF-18	API-55	4.87	49219	LGAEVYHTLK IVIGMDVAASEFYR	SEQ ID NO: 233 SEQ ID NO: 206
AF-18	API-245	4.87	49219	VEQATQAIPMER	SEQ ID NO: 402
AF-21	API-6	5.4	141094	IDGDTIIFSNVQER GKPPPSFSWTR AETYEGVYQCTAR LSPYVNYSFR	SEQ ID NO: 190 SEQ ID NO: 154 SEQ ID NO: 11 SEQ ID NO: 266
AF-22	API-56	4.93	133773	RTMRDQDTGK LICSELNGR EGLDLQVLEDSGR	SEQ ID NO: 337 SEQ ID NO: 239 SEQ ID NO: 86
AF-22	API-57	4.93	133773	ILDDLSPR NGIDIYSLTVDSR SPEQQETVLDGNLIIR YIFHNFMER	SEQ ID NO: 196 SEQ ID NO: 295 SEQ ID NO: 356 SEQ ID NO: 452
AF-23	API-7	4.5	32473	IPTTFENGR	SEQ ID NO: 199
AF-23	API-8	4.5	32473	HLEEPGETQNAFLNER GEAGAPGEEDIQGPTK EDEEEEEGENYQK	SEQ ID NO: 179 SEQ ID NO: 141 SEQ ID NO: 79
AF-24	API-9	5.31	46663	NNLVIFHR IVQFSPSGK EGPVLILGR	SEQ ID NO: 299 SEQ ID NO: 207 SEQ ID NO: 87
AF-25	API-10	5.68	36700	ASSIIDELFQDR	SEQ ID NO: 34
AF-26	API-14	8.11	32305	APEAQVSVQPNFQQDK TMLLPAGSLGSYSYR	SEQ ID NO: 28 SEQ ID NO: 379
AF-27	API-15	5.33	141371	WLQGSQELPR	SEQ ID NO: 439

AF#	API#	pI	MW (Da)	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
AF-27	API-58	5.33	141371	NALGAIHHTISVR IDGDTIIFSNVQER GKPPPSFSWTR AETYEGVYQCTAR LSPYVNYSFR	SEQ ID NO: 287 SEQ ID NO: 190 SEQ ID NO: 154 SEQ ID NO: 11 SEQ ID NO: 266
AF-28	API-16	5.13	158568	IALVITDGR	SEQ ID NO: 189
AF-28	API-59	5.13	158568	GAYPLSIEPIGVR QSEDSTFYLGFR ALYLQYTDFTFR	SEQ ID NO: 135 SEQ ID NO: 328 SEQ ID NO: 26
AF-29	API-196	9.22	47059	ALDFAVGEYNK LVGGPMDASVEEEGVR	SEQ ID NO: 20 SEQ ID NO: 273
AF-30	API-17	5.67	48057	TSLEDFYLDEER	SEQ ID NO: 385
				LAAAVSNFGYDLFR	SEQ ID NO: 219
AF-31	API-60	6.07	91258	HVVPNEVVVQR AGALNSNDAFVLK YIETDPANR	SEQ ID NO: 187 SEQ ID NO: 15 SEQ ID NO: 451
AF-32	API-18	6.17	48958	VAMHLVCPSR TALASGGVLDASGDYR EPGEFALLR	SEQ ID NO: 394 SEQ ID NO: 363 SEQ ID NO: 104
AF-34	API-61	4.54	145408	FFEECDPNK TGLEAISNKK	SEQ ID NO: 120 SEQ ID NO: 371
AF-35	API-62	5.21	18623	SELEEQLTPVAEETR QQTEWQSGQR	SEQ ID NO: 342 SEQ ID NO: 327
AF-37	API-19	6.91	33523	EVGVYEALK FVEGLPINDFSR ENFSCLTR DVIATDKEDVAFK	SEQ ID NO: 111 SEQ ID NO: 132 SEQ ID NO: 102 SEQ ID NO: 74
AF-38	API-63	6.47	29535	YTNWIKK EKPGVYTNVCR LVHGGPCDK LSELIQPLPLER	SEQ ID NO: 461 SEQ ID NO: 93 SEQ ID NO: 275 SEQ ID NO: 264
AF-39	API-64	7.5	35510	CSVFYGAPSK	SEQ ID NO: 48
AF-39	API-65	7.5	35510	YLELFQR LVNIYDSMPLR	SEQ ID NO: 454 SEQ ID NO: 279

AF#	API#	pI	MW (Da)	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
AF-40	API-20	7.29	38617	HIYLLPSGR NFGLYNER IGADFLAR VGGVQSLGGTGALR ITWSNPQAQGAR	SEQ ID NO: 177 SEQ ID NO: 291 SEQ ID NO: 194 SEQ ID NO: 410 SEQ ID NO: 205
AF-41	API-22	5.85	17345	LEGEACGVYTPR	SEQ ID NO: 227
AF-42	API-66	5.04	18662	LIVHNGYCDGR	SEQ ID NO: 241
AF-43	API-67	9.83	14065	LEEQAQQIR LGPLVEQGR	SEQ ID NO: 226 SEQ ID NO: 236
AF-43	API-68	9.83	14065	LVGGPMDASVEEEGVR	SEQ ID NO: 273
AF-44	API-69	6.63	102328	EELPAQDIK	SEQ ID NO: 82
AF-44	API-70	6.63	102328	AASGTQNNVLR GCPTEEGCGER	SEQ ID NO: 3 SEQ ID NO: 137
AF-45	API-23	6.04	46998	TSLEDFYLDEER LAAAVSNFGYDLYR ELLDTVTAPQK ALYYDLISSPDIHGTYK	SEQ ID NO: 385 SEQ ID NO: 219 SEQ ID NO: 96 SEQ ID NO: 27
AF-46	API-24	4.71	19802	THPHFVIPYR	SEQ ID NO: 374
AF-46	API-197	4.71	19802	YENEVALR QSLEASLAETEGR	SEQ ID NO: 445 SEQ ID NO: 329
AF-46	API-198	4.71	19802	YEELQQTAGR	SEQ ID NO: 444
AF-47	API-25	5.99	49664	VAMHLVCPSR YEAAVPDPR TALASGGVLDASGDYR EPGEFALLR	SEQ ID NO: 394 SEQ ID NO: 442 SEQ ID NO: 363 SEQ ID NO: 104
AF-48	API-71	5.32	122332	AFLFQESPR YLELESSGHR	SEQ ID NO: 13 SEQ ID NO: 453
AF-49	API-26	6.94	27576	DSCQGDSGGPLVCGDHL R EKPGVYTNVCR GLVSWGNI PCGSK	SEQ ID NO: 68 SEQ ID NO: 93 SEQ ID NO: 160
AF-49	API-27	6.94	27576	TMLLQPAGSLGYSYR	SEQ ID NO: 379
AF-50	API-72	6.82	71337	NVPLPVIAELPPK	SEQ ID NO: 305
AF-50	API-73	6.82	71337	EQPPSLTR	SEQ ID NO: 107

AF#	API#	pI	MW (Da)	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
				CFEPQLLR	SEQ ID NO: 43
AF-50	API-199	6.82	71337	GGEGTGYFVDFSVR DSPVLIDFFEDTER YWNDCEPPDSR	SEQ ID NO: 149 SEQ ID NO: 71 SEQ ID NO: 465
AF-50	API-200	6.82	71337	CISIIYSER VYLFDPEGK	SEQ ID NO: 45 SEQ ID NO: 435
AF-51	API-28	5.7	34388	ASSIIDELFQDR	SEQ ID NO: 34
AF-51	API-30	5.7	34388	LIAPVAEEEEATVPNNK	SEQ ID NO: 238
				LKDDEVAQLK SADTLWDIQK	SEQ ID NO: 242 SEQ ID NO: 339
AF-76	API-86	5.59	45537	NNLVIFHR EGPVLILGR	SEQ ID NO: 299 SEQ ID NO: 87
AF-79	API-201	5.52	142378	LPPNVVEESAR	SEQ ID NO: 260
AF-81	API-88	5.43	78299	LVESGGGLVQPGGSLR	SEQ ID NO: 271
AF-81	API-202	5.43	78299	VSSQNIQDFPSVLR GEASVCVEDWESGDR	SEQ ID NO: 428 SEQ ID NO: 143
AF-82	API-89	6.69	74838	HSTVLENLPDK LLEACTFHSAK	SEQ ID NO: 185 SEQ ID NO: 244
AF-83	API-90	6.81	71920	HSTVLENLPDK WCAIGHEETQK EPVDNAENCHLAR FDQFFGEGCAPGSQR WCTISNQEANK CGLVPVLAENYK GSNFQWNQLQGK SPDFQLFSSSHGK IECVSAENTEDCIAK QMDFELLCCQNGAR DQYELLCR	SEQ ID NO: 185 SEQ ID NO: 436 SEQ ID NO: 106 SEQ ID NO: 116 SEQ ID NO: 437 SEQ ID NO: 44 SEQ ID NO: 168 SEQ ID NO: 355 SEQ ID NO: 193 SEQ ID NO: 324 SEQ ID NO: 67
AF-84	API-91	6.94	73402	HSTVLENLPDK FDQFFGEGCAPGSQR EPVDNAENCHLAR WCTISNQEANK VTCVAEELLK	SEQ ID NO: 185 SEQ ID NO: 116 SEQ ID NO: 106 SEQ ID NO: 437 SEQ ID NO: 430

AF#	API#	pI	MW (Da)	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
				QMDFELLQNGAR DQYELLCR DNPQTHYYAVAVVK	SEQ ID NO: 324 SEQ ID NO: 67 SEQ ID NO: 62
AF-85	API-92	7.1	73878	IPIEDGSGEVVLSR	SEQ ID NO: 198
AF-85	API-93	7.1	73878	HSTVLENLPDK CGLVPVLAENYK EPVDNAENCHLAR SPDFQLFSSSHGK FDQFFGEGCAPGSQR DQYELLCR	SEQ ID NO: 185 SEQ ID NO: 44 SEQ ID NO: 106 SEQ ID NO: 355 SEQ ID NO: 116 SEQ ID NO: 67
AF-87	API-95	5.95	64179	FQPLVDEPK DVFLGTFLYEYSR LGEYGFQNALIVR IYEATLEDCCAK ECCHGDLLECADDR	SEQ ID NO: 127 SEQ ID NO: 73 SEQ ID NO: 234 SEQ ID NO: 209 SEQ ID NO: 77
AF-89	API-97	5.39	65155	AGDFLEANYMNLQR GYTQQLAFR	SEQ ID NO: 16 SEQ ID NO: 175
AF-90	API-98	7.61	62945	LPLEYSYGEYR	SEQ ID NO: 259
AF-91	API-99	8.16	56352	GDYPLEAVR GIFPVLCK DPVQEAWAEDVDLR LFEELVR	SEQ ID NO: 140 SEQ ID NO: 152 SEQ ID NO: 65 SEQ ID NO: 231
AF-100	API-101	6.08	44068	HPDYSVSLLLR DVFLGTFLYEYSR CCTESLVNR YICENQDTISTK LGEYGFQNALIVR LSCAEDYLSLVLR	SEQ ID NO: 183 SEQ ID NO: 73 SEQ ID NO: 42 SEQ ID NO: 450 SEQ ID NO: 234 SEQ ID NO: 262
AF-103	API-102	5.93	42722	ITCTEEGWSPTPK EIMENYNIALR INHGILYDEEK	SEQ ID NO: 202 SEQ ID NO: 92 SEQ ID NO: 197
AF-104	API-103	5.09	42184	YVMLPVADQEK	SEQ ID NO: 463
AF-105	API-104	5.19	42184	GSPAINVAVHVFR	SEQ ID NO: 169
AF-107	API-107	7.26	33226	DNLAIQTR	SEQ ID NO: 61

AF#	API#	pI	MW (Da)	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
				ITVVDALHEIPVK	SEQ ID NO: 204
AF-107	API-210	7.26	33226	KLIVENVDVLTQMR	SEQ ID NO: 211
AF-108	API-108	7.54	33136	AGAAAGGPGVSGVCVC K GTCEQGPSIVTPPK GYCAPGMECVK	SEQ ID NO: 14 SEQ ID NO: 170 SEQ ID NO: 172
AF-117	API-113	4.65	13983	SELEEQLTPVAEETR KVEQAVETEPEPELR	SEQ ID NO: 342 SEQ ID NO: 215
AF-119	API-114	7.23	11699	LLVVYPWTQR VVAGVANALAHK GTFATLSELHCDK	SEQ ID NO: 251 SEQ ID NO: 432 SEQ ID NO: 171
AF-149	API-214	4.82	190721	VEEVKPLEGR	SEQ ID NO: 399
AF-150	API-144	6.87	157592	VEVLAGDLR GPPGPPGGVVVR	SEQ ID NO: 404 SEQ ID NO: 165
AF-152	API-146	5.04	81703	FTDSENVQER FTFEYSR	SEQ ID NO: 130 SEQ ID NO: 131
AF-152	API-147	5.04	81703	VIALINDQR	SEQ ID NO: 413
AF-152	API-148	5.04	81703	ELLESYIDGR TATSEYQTFNPR	SEQ ID NO: 97 SEQ ID NO: 364
AF-154	API-150	5.03	67307	LCQDLGPGAFR QEDDLANINQWVK	SEQ ID NO: 222 SEQ ID NO: 314
AF-154	API-151	5.03	67307	AIEDYINEFSVR DVVLTTTFVDDIK	SEQ ID NO: 19 SEQ ID NO: 75
AF-154	API-152	5.03	67307	WLQGSQELPR	SEQ ID NO: 439
AF-155	API-215	9.21	64021	ALDFAVGEYNK LVGGPMDASVEEEGVR	SEQ ID NO: 20 SEQ ID NO: 273
AF-156	API-153	4.36	58083	DQDGEILLPR	SEQ ID NO: 66
AF-159	API-158	5.08	52008	TSLEDFYLDEER TALASGGVLDASGDYR EPGEFALLR QVFGATK YYTVFDR	SEQ ID NO: 385 SEQ ID NO: 363 SEQ ID NO: 104 SEQ ID NO: 333 SEQ ID NO: 467
AF-163	API-165	4.45	34879	NPNLPPETVDSLK NILTSNNIDVK	SEQ ID NO: 301 SEQ ID NO: 296

AF#	API#	pI	MW (Da)	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
				CPNPPVQENFDVNK IPTTFENGR	SEQ ID NO: 47 SEQ ID NO: 199
AF-163	API-166	4.45	34879	GEAGAPGEEDIQGPTK	SEQ ID NO: 141
AF-164	API-167	5	33485	EILSVDCSTNNPSQAK FMETVAEK ELDESLQVAER	SEQ ID NO: 91 SEQ ID NO: 124 SEQ ID NO: 94
AF-169	API-173	8	34362	EELVYELNPLDHR GSFEFPVGDAVSK LGQYASPTAK	SEQ ID NO: 83 SEQ ID NO: 167 SEQ ID NO: 237
AF-170	API-174	5.41	31886	ELDESLQVAER	SEQ ID NO: 94
AF-170	API-175	5.41	31886	AADDTWEPFASGK GSPAINVAVHVFR	SEQ ID NO: 1 SEQ ID NO: 169
AF-170	API-176	5.41	31886	AATVGSLAGQPLQER SELEEQLTPVAEETR LEEQAQQIR LGADMEDVCGR SWFEPLVEDMQR	SEQ ID NO: 4 SEQ ID NO: 342 SEQ ID NO: 226 SEQ ID NO: 232 SEQ ID NO: 361
AF-172	API-179	6.71	28747	HLDSVLQQLQTEVYR GPCWCVDR	SEQ ID NO: 178 SEQ ID NO: 164
AF-172	API-180	6.71	28747	DSCQGDSGGPLVCGDHL R EKPGVYTNVCR GLVSWGNI PCGSK KPNLQVFLGK	SEQ ID NO: 68 SEQ ID NO: 93 SEQ ID NO: 160 SEQ ID NO: 213
AF-173	API-181	7.67	27476	NEQVEIR SNLDEDI AEENIVSR	SEQ ID NO: 290 SEQ ID NO: 354
AF-174	API-182	4.67	27811	SVTEQGAELSNEER	SEQ ID NO: 359
AF-175	API-183	5.33	24936	AQGFTEDTIVFLPQTDK TMLLQPAGSLG SYSYR APEAQVSVQPNFQQDK	SEQ ID NO: 29 SEQ ID NO: 379 SEQ ID NO: 28
AF-176	API-184	4.86	22248	AQGFTEDTIVFLPQTDK TMLLQPAGSLG SYSYR	SEQ ID NO: 29 SEQ ID NO: 379
AF-178	API-185	6.03	22247	LPFVINDGK	SEQ ID NO: 256
AF-178	API-217	6.03	22247	APEAQVSVQPNFQQDK	SEQ ID NO: 28

AF#	API#	pI	MW (Da)	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
				AQGFTEDTIVFLPQTDK TMLLQPAGSLGSSYR	SEQ ID NO: 29 SEQ ID NO: 379
AF-178	API-219	6.03	22247	TQGFTEDAIVFLPQTDK	SEQ ID NO: 382
AF-181	API-187	5.72	16336	GDGPVQGIINFEQK HVGDLGNVTADK	SEQ ID NO: 138 SEQ ID NO: 186
AF-183	API-189	10.36	11160	ALDFAVGEYNK LVGGPMDASVEEEGVR	SEQ ID NO: 20 SEQ ID NO: 273
AF-184	API-190	5.31	48769	TSLEDFYLDEER ELLDTVTAPQK	SEQ ID NO: 385 SEQ ID NO: 96
AF-186	API-238	4.71	29693	IPTTFENGR	SEQ ID NO: 199
AF-187	API-239	4.93	154156	QPEYAVVQR	SEQ ID NO: 325
AF-190	API-240	5.29	29663	NNYMYAR ELDVLQGR	SEQ ID NO: 300 SEQ ID NO: 95

The second group comprises APIs that are increased in the CSF of subjects having AD as compared with the CSF of subjects free from AD. The amino acid sequences of peptides produced from these APIs by proteolysis using trypsin and identified by tandem mass spectrometry and database searching using the SEQUEST program are listed in Table V, in addition to their corresponding pIs and MWs.

Table V. APIs Increased in CSF of Subjects Having AD

(a) APIs Increased in CSF of Subjects Having AD

AF#	API#	pI	MW (Da)	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
AF-310	API-358	4.65	30092	THLPEVFLSK YTFELSR	SEQ ID NO: 373 SEQ ID NO: 460
AF-311	API-402	5.84	69278	LSLELEQLELQR	SEQ ID NO: 265
AF-311	API-403	5.84	69278	NFPSPVDAAFR GGYTLVSGYPK	SEQ ID NO: 293 SEQ ID NO: 151
AF-312	API-359	5.51	20016	LSEDYGVLK TDEGLAYR QLTVNDLPVGR	SEQ ID NO: 263 SEQ ID NO: 366 SEQ ID NO: 323

AF#	API#	pI	MW (Da)	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
AF-313	API-360	6.2	76215	YVTSAPMPEPQAPGR VSVFVPPR DGFFGNPR	SEQ ID NO: 464 SEQ ID NO: 429 SEQ ID NO: 54
AF-317	API-404	4.82	12454	VQVTSQEYSAR LVESYQLR	SEQ ID NO: 426 SEQ ID NO: 272
AF-318	API-405	5.88	68454	QSSSYSFFK MALEVYK LSLELEQLELQR AAPMDLTLTETR	SEQ ID NO: 330 SEQ ID NO: 284 SEQ ID NO: 265 SEQ ID NO: 2
AF-319	API-406	7.53	39490	LVMGLPTFGR LLGQQVPYATK GNQWVGYYDDQESVK FSNTDYAVGYMLR FPLTNALK	SEQ ID NO: 278 SEQ ID NO: 246 SEQ ID NO: 163 SEQ ID NO: 129 SEQ ID NO: 125
AF-323	API-407	5.79	148765	VGFIYESDVMGR LPPNVVEESAR	SEQ ID NO: 409 SEQ ID NO: 260
AF-323	API-408	5.79	148765	WLQGSQELPR	SEQ ID NO: 439
AF-323	API-409	5.79	148765	TLLSDDWK MNNGDVDLTSDR LVESYQLR LLALAPTFEMNPMK FLPLLPLPER ATSVALTWSR	SEQ ID NO: 376 SEQ ID NO: 285 SEQ ID NO: 272 SEQ ID NO: 243 SEQ ID NO: 123 SEQ ID NO: 36
AF-324	API-361	4.69	59610	NLAVSQVVHK AVLDVFEETGEASAATAVK LYGSEAFATDFQDSAAAK	SEQ ID NO: 297 SEQ ID NO: 39 SEQ ID NO: 282
AF-325	API-362	10.5	28059	GNQWVGYYDDQESVK FSNTDYAVGYMLR LVMGLPTFGR	SEQ ID NO: 163 SEQ ID NO: 129 SEQ ID NO: 278
AF-326	API-363	5.47	38663	DVFLGMFLYEYAR FQNALLVR	SEQ ID NO: 72 SEQ ID NO: 126
AF-327	API-410	5.59	44069	VPQVSTPTLVEVSR VPEVSTPTLVEVSR QTALVELVK	SEQ ID NO: 422 SEQ ID NO: 421 SEQ ID NO: 331

AF#	API#	pI	MW- (Da)	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
				PLVEEPQNLLK KVPEVSTPTLVEVSR HPDYSVVLLLR FQNALLVR DVFLGMFLYEYAR AVMDDFAAFVEK	SEQ ID NO: 311 SEQ ID NO: 216 SEQ ID NO: 184 SEQ ID NO: 126 SEQ ID NO: 72 SEQ ID NO: 40
AF-328	API-411	5.39	90721	NFMLDSNGELLLR MNNGDVDLTSDR LLALAPTFEMNPMK FLPLLPLPER ASPPVYK	SEQ ID NO: 292 SEQ ID NO: 285 SEQ ID NO: 243 SEQ ID NO: 123 SEQ ID NO: 33
AF-330	API-364	5.88	28511	AQGFTEDTLVFLPQTDK TMLLQPAGSLGSYSYR	SEQ ID NO: 30 SEQ ID NO: 379
AF-331	API-412	4.79	18423	YEVSSPYFK VNSDGGLVALR	SEQ ID NO: 446 SEQ ID NO: 420
AF-331	API-413	4.79	18423	EVAGLWLK	SEQ ID NO: 109
AF-332	API-365	6.65	39644	NLDYVATSLHEAVTK LTWSNPPAQGAR VGGVQSLGGTGALR HLYLLPSGR	SEQ ID NO: 298 SEQ ID NO: 269 SEQ ID NO: 410 SEQ ID NO: 181
AF-333	API-366	5.29	61283	SGNTFHPEVR WLQGSQELPR QEPSQGTTTFAVTSLLR	SEQ ID NO: 347 SEQ ID NO: 439 SEQ ID NO: 317
AF-334	API-414	6.67	47683	VWVYPPEK GGYTLVSGYPK	SEQ ID NO: 434 SEQ ID NO: 151
AF-334	API-415	6.67	47683	YEAAVPDPR VPTFDER TALASGGVLDASGDYR	SEQ ID NO: 442 SEQ ID NO: 424 SEQ ID NO: 363
AF-335	API-367	4.82	23513	VQPYLDDFQK DYVSQFEGSALGK THLAPYSDELRL	SEQ ID NO: 425 SEQ ID NO: 76 SEQ ID NO: 372
AF-335	API-370	4.82	23513	AQGFTEDTIVFLPQTDK TMLLQPAGSLGSYSYR	SEQ ID NO: 29 SEQ ID NO: 379
AF-336	API-416	5.94	35737	KTLLSNLEEAK	SEQ ID NO: 214

AF#	API#	pI	MW (Da)	Amino Acid Sequence of Tryptic Digest Peptides	SEQ ID NO:
				ASSLLDELQDR	SEQ ID NO: 35
AF-336	API-417	5.94	35737	TMLLQPAGSLGSYSYR APEAQVSVQPNFQQDK	SEQ ID NO: 379 SEQ ID NO: 28
AF-337	API-418	6.82	51089	EHSSLAFWK ATVVYQGER AGLLENGAVR	SEQ ID NO: 89 SEQ ID NO: 37 SEQ ID NO: 18
AF-338	API-368	5.7	43923	AVMDDFAAFVEK VPQVSTPTLVEVSR DVFLGMFLYEYAR FQNALLVR	SEQ ID NO: 40 SEQ ID NO: 422 SEQ ID NO: 72 SEQ ID NO: 126

(b) APIs which can be used in combination with the APIs in Table IV (a) and Table V (a)

AF#	API#	pI	MW (Da)	Amino Acid Sequences of Tryptic Digest Peptides	SEQ ID NO:
AF-52	API-74	6.3	32573	FACYYPYR GLQDEDGYR	SEQ ID NO: 115 SEQ ID NO: 158
AF-53	API-33	5.84	45302	YICENQDSISSK AVMDDFAAFVEK	SEQ ID NO: 449 SEQ ID NO: 40
AF-54	API-221	5.12	17520	SELEEQLTPVAEETR	SEQ ID NO: 342
AF-55	API-34	8.1	12361	ALDFAVGEYNK LVGGPMDASVEEEGVR	SEQ ID NO: 20 SEQ ID NO: 273
AF-56	API-75	8.56	52128	DNDGWLTSDPR LESDVSAQMEYCR IRPFFPQQ NYCGLPGYEWLGNDK	SEQ ID NO: 60 SEQ ID NO: 229 SEQ ID NO: 200 SEQ ID NO: 306
AF-56	API-246	8.56	52128	TGAQELLR AGALNSNDAFVLK	SEQ ID NO: 370 SEQ ID NO: 15
AF-57	API-35	6.3	68549	MTLDDFR	SEQ ID NO: 286
AF-57	API-76	6.3	68549	VFLDCCNYITELR	SEQ ID NO: 406

AF#	API#	pI	MW (Da)	Amino Acid Sequences of Tryptic Digest Peptides	SEQ ID NO:
AF-57	API-222	6.3	68549	QSLEASLAETEGR	SEQ ID NO: 329
AF-58	API-77	5.01	14507	KVEQAVETEPEPELR	SEQ ID NO: 215
AF-59	API-36	6.74	33401	TSLEDFYLDDEER	SEQ ID NO: 385
AF-60	API-37	5.39	33873	SELEEQLTPVAEETR	SEQ ID NO: 342
				KVEQAVETEPEPELR	SEQ ID NO: 215
				GEVQAMLGQSTEELR	SEQ ID NO: 147
AF-61	API-78	6.76	54345	IPIEDGSGEVVLSR	SEQ ID NO: 198
				TIYTPGSTVLVLR	SEQ ID NO: 375
				QELSEAEQATR	SEQ ID NO: 316
AF-62	API-38	6.6	31004	FACYYP	SEQ ID NO: 115
				ITQVLHFTK	SEQ ID NO: 203
				GLQDEDGYR	SEQ ID NO: 158
AF-63	API-79	5.97	14897	DSCVGSLLVVK	SEQ ID NO: 69
				EVVADSVWVDVK	SEQ ID NO: 113
				QPVPGQQMTLK	SEQ ID NO: 326
				IWDVVEK	SEQ ID NO: 208
AF-64	API-80	6.67	68119	IPIEDGSGEVVLSR	SEQ ID NO: 198
				SNLDEDIIAEENIVSR	SEQ ID NO: 354
				DFDFVPPVVR	SEQ ID NO: 53
AF-65	API-81	7.19	58620	FLCTGGVSPYADPNTCR	SEQ ID NO: 122
				CLVNLIEK	SEQ ID NO: 46
AF-65	API-223	7.19	58620	VGDTLNLNLR	SEQ ID NO: 408
AF-66	API-82	10.05	30092	FISLGEACK	SEQ ID NO: 121
				VFLDCCNYITELR	SEQ ID NO: 406
AF-66	API-83	10.05	30092	ALDFAVGEYNK	SEQ ID NO: 20
				LVGGPMDASVEEEGVR	SEQ ID NO: 273
AF-67	API-39	5.02	13735	AADDTWEPFASGK	SEQ ID NO: 1
AF-68	API-84	9.06	35351	VPTANVSVVDLTCR	SEQ ID NO: 423
				LISWYDNEFGYSNR	SEQ ID NO: 240
AF-68	API-85	9.06	35351	ISYQSSSTEER	SEQ ID NO: 201
AF-69	API-40	5.01	46760	ALVQQMEQLR	SEQ ID NO: 25
				TQVNTQAEQLR	SEQ ID NO: 384
AF-69	API-247	5.01	46760	QGSFQGGFR	SEQ ID NO: 320
				AEMADQAAAWLTR	SEQ ID NO: 9

AF#	API#	pI	MW (Da)	Amino Acid Sequences of Tryptic Digest Peptides	SEQ ID NO:
				VLSLAQEQVGGSPEK	SEQ ID NO: 418
AF-70	API-41	8.91	38789	QHFTTLIK GNQWVGYYDDQESVK FSNTDYAVGYMLR EGDGSCFPDALDR LVMGIPTFGR	SEQ ID NO: 322 SEQ ID NO: 163 SEQ ID NO: 129 SEQ ID NO: 85 SEQ ID NO: 277
AF-70	API-224	8.91	38789	YLYEIAR DAIPEDLPPLTADFAEDK	SEQ ID NO: 457 SEQ ID NO: 49
AF-71	API-42	6.44	68579	GYTQQLAFR SNLDEDIIEENIVSR VFLDCCNYITELR	SEQ ID NO: 175 SEQ ID NO: 354 SEQ ID NO: 406
AF-72	API-43	5	43788	LEPYADQLR ALVQQMEQLR TQVNTQAEQLR IDQTVEELR	SEQ ID NO: 228 SEQ ID NO: 25 SEQ ID NO: 384 SEQ ID NO: 191
AF-73	API-44	5.21	31615	AADDTWEPFASGK	SEQ ID NO: 1
AF-74	API-45	6.19	51934	YYCFQGNQFLR GECQAEGVLFFQGDR	SEQ ID NO: 466 SEQ ID NO: 144
AF-74	API-248	6.19	51934	TVMVNIENPEGIPVK TIYTPGSTVLYR	SEQ ID NO: 389 SEQ ID NO: 375
AF-75	API-46	5.03	33671	EILSVCSTNNPSQAK ELDESLQVAER	SEQ ID NO: 91 SEQ ID NO: 94
AF-75	API-225	5.03	33671	AATVGSLAQPLQER LGPLVEQGR	SEQ ID NO: 4 SEQ ID NO: 236
AF-121	API-116	5.42	105108	RLDGSVDFK YLQEIYNSNNQK VELEDWNGR VAQLEAQCQEPCK TSTADYAMFK YEASILTHDSSIR DNCCILDER	SEQ ID NO: 335 SEQ ID NO: 456 SEQ ID NO: 400 SEQ ID NO: 396 SEQ ID NO: 386 SEQ ID NO: 443 SEQ ID NO: 59
AF-123	API-118	7.31	64933	ADSGEGDFLAEGGGVR GLIDEVNQDFTNR	SEQ ID NO: 8 SEQ ID NO: 156
AF-124	API-119	7.47	64736	ESSSHHPGIAEFPSR	SEQ ID NO: 108

AF#	API#	pI	MW (Da)	Amino Acid Sequences of Tryptic Digest Peptides	SEQ ID NO:
				GLIDEVQNQDFTNR	SEQ ID NO: 156
AF-125	API-120	4.77	61297	SGNENGFEFYLR	SEQ ID NO: 346
AF-126	API-121	4.11	60374	DCQPGLCCAFQR	SEQ ID NO: 50
				DQDGEILLPR	SEQ ID NO: 66
AF-126	API-122	4.11	60374	DQDGEILLPR	SEQ ID NO: 66
AF-127	API-123	4.98	59649	ALQDQLVLVAAK	SEQ ID NO: 23
				SLDFTELDVAAEK	SEQ ID NO: 350
AF-128	API-124	6.6	57865	VGDTLNLNLR	SEQ ID NO: 408
				LNMGITDLQGLR	SEQ ID NO: 253
AF-129	API-125	5.29	54625	LCDNLSTK	SEQ ID NO: 221
				YTFELSR	SEQ ID NO: 460
				VCSQYAAAYGEK	SEQ ID NO: 398
				LPEATPTELAK	SEQ ID NO: 255
				FEDCCQEK	SEQ ID NO: 117
				HLSLLTTLSNR	SEQ ID NO: 180
				THLPEVFLSK	SEQ ID NO: 373
				ELPEHTVK	SEQ ID NO: 99
				KLCMAALK	SEQ ID NO: 210
AF-129	API-126	5.29	54625	DPTFIPAPIQAK	SEQ ID NO: 64
				SLDFTELDVAAEK	SEQ ID NO: 350
AF-130	API-127	5.08	51880	TSLED FYLDEER	SEQ ID NO: 385
				LAAAVSNFGYDLYR	SEQ ID NO: 219
				LQSLFDSPDFSK	SEQ ID NO: 261
AF-130	API-128	5.08	51880	VAMHLVCPSR	SEQ ID NO: 394
				TALASGGVLDASGDYR	SEQ ID NO: 363
				EPGEFALLR	SEQ ID NO: 104
AF-132	API-130	4.72	47414	AEMADQASAWLTR	SEQ ID NO: 10
				ADGSYAAWLSR	SEQ ID NO: 5
				QGSFQGGFR	SEQ ID NO: 320
				VLSLAQEQVGGSPK	SEQ ID NO: 418
				TEQWSTLPPETK	SEQ ID NO: 368
				DHAVDLIQK	SEQ ID NO: 57
AF-133	API-131	5.12	44068	LEPYADQLR	SEQ ID NO: 228
				TQVNTQAEQLR	SEQ ID NO: 384

AF#	API#	pI	MW (Da)	Amino Acid Sequences of Tryptic Digest Peptides	SEQ ID NO:
AF-134	API-132	5	43516	LEPYADQLR	SEQ ID NO: 228
AF-137	API-134	4.98	36855	KYNELLK	SEQ ID NO: 218
				ELDESLQVAER	SEQ ID NO: 94
AF-137	API-135	4.98	36855	VFSLQWGEVK	SEQ ID NO: 407
				AQLGDLPWQVAIK	SEQ ID NO: 31
AF-137	API-232	4.98	36855	LGPIEAIQK	SEQ ID NO: 235
AF-137	API-233	4.98	36855	LEEQAQQIR	SEQ ID NO: 226
				LGPLVEQGR	SEQ ID NO: 236
AF-137	API-234	4.98	36855	KMEENEK	SEQ ID NO: 212
AF-139	API-136	5	34295	TLLSNLEEAK	SEQ ID NO: 377
				EILSVDCSTNNPSQAK	SEQ ID NO: 91
				EDALNETRESETKLK	SEQ ID NO: 78
				IDSLLENDR	SEQ ID NO: 192
				ELDESLQVAER	SEQ ID NO: 94
AF-139	API-137	5	34295	AATVGSLAGQPLQER	SEQ ID NO: 4
				SELEEQLTPVAEETR	SEQ ID NO: 342
AF-140	API-138	6.8	32080	FACYYPYR	SEQ ID NO: 115
				GLQDEDGYR	SEQ ID NO: 158
AF-141	API-139	7.5	28440	TNFDNDIALVR	SEQ ID NO: 380
				LLEVPEGR	SEQ ID NO: 245
AF-142	API-140	6.75	27279	VELLHNPAFCSLATTK	SEQ ID NO: 401
				SNLDEDIIEENIVSR	SEQ ID NO: 354
AF-142	API-141	6.75	27279	LSELIQPLPLER	SEQ ID NO: 264
AF-143	API-142	7.44	26066	SGTASVVCLLNNFYPR	SEQ ID NO: 348
				LLIYWASTR	SEQ ID NO: 247
AF-144	API-143	6.56	20744	SDGSCAWYR	SEQ ID NO: 340
				EVDSGNDIYGNPIK	SEQ ID NO: 110
AF-151	API-145	5.28	137531	IDGDTIIFSNVQER	SEQ ID NO: 190
				GKPPPSFSWTR	SEQ ID NO: 154
				AETYEGVYQCTAR	SEQ ID NO: 11
AF-153	API-149	9.85	69630	VGDTLNLNLR	SEQ ID NO: 408
				LNMGITDLQGLR	SEQ ID NO: 253
AF-157	API-155	4.99	55449	YEAAVPDPR	SEQ ID NO: 442
				TALASGGVLDASGDYR	SEQ ID NO: 363

AF#	API#	pI	MW (Da)	Amino Acid Sequences of Tryptic Digest Peptides	SEQ ID NO:
				EPGEFALLR	SEQ ID NO: 104
AF-161	API-161	5.18	44404	RVEPYGENFNK	SEQ ID NO: 338
				LEPYADQLR	SEQ ID NO: 228
				ALVQQMEQLR	SEQ ID NO: 25
				SLAPYAQDTQEK	SEQ ID NO: 349
				TQVNTQAEQLR	SEQ ID NO: 384
				IDQTVHEELR	SEQ ID NO: 191
AF-161	API-162	5.18	44404	TSLEDGYLDEER	SEQ ID NO: 385
AF-161	API-163	5.18	44404	VAPEEHPVLLTEAPLNPK	SEQ ID NO: 395
				QEYDESGPSIVHR	SEQ ID NO: 318
				GYSFTTTAER	SEQ ID NO: 174
				AGFAGDDAPR	SEQ ID NO: 17
				SYELPDGQVITIGNER	SEQ ID NO: 362
				AVFPSIVGR	SEQ ID NO: 38
AF-165	API-168	7.17	34230	EPFLSCCQFAESLR	SEQ ID NO: 103
				EELVYELNPLDHR	SEQ ID NO: 83
AF-166	API-169	8.54	33657	EELVYELNPLDHR	SEQ ID NO: 83
				GLCVATPVQLR	SEQ ID NO: 155
AF-167	API-170	5.69	33621	TLLSNLEEAK	SEQ ID NO: 377
				ASSIIDELFQDR	SEQ ID NO: 34
AF-167	API-171	5.69	33621	SELEEQLTPVAEETR	SEQ ID NO: 342
				GEVQAMLGQSTEELRLEEQ	SEQ ID NO: 148
				AQQIR	
AF-168	API-237	7.66	33920	ALEESNYELEGK	SEQ ID NO: 21
AF-168	API-172	7.66	33920	EPFLSCCQFAESLR	SEQ ID NO: 103
				EELVYELNPLDHR	SEQ ID NO: 83
				GLCVATPVQLR	SEQ ID NO: 155
				GSFEFPVGDAVSK	SEQ ID NO: 167
AF-171	API-177	4.98	29658	AQGFTEDTIVFLPQTDK	SEQ ID NO: 29
				TMLLQPAGSLGSYSYR	SEQ ID NO: 379
AF-171	API-178	4.98	29658	AADDTWEPFASGK	SEQ ID NO: 1
				GSPAINVAVHVFR	SEQ ID NO: 169
AF-179	API-186	5.26	20115	FSGTWYAMAK	SEQ ID NO: 128
				QEELCLAR	SEQ ID NO: 315

AF#	API#	pI	MW (Da)	Amino Acid Sequences of Tryptic Digest Peptides	SEQ ID NO:
				LIVHNGYCDGR	SEQ ID NO: 241
AF-180	API-220	6.17	16255	GLQDEDGYR CSVFYGAPSK	SEQ ID NO: 158 SEQ ID NO: 48
AF-182	API-188	4.89	13651	AADDTWEPFASGK	SEQ ID NO: 1
AF-185	API-191	5.32	40323	VGYYSGWGR	SEQ ID NO: 412
AF-185	API-192	5.32	40323	ADQVCINLR SGNENGEFYLR	SEQ ID NO: 7 SEQ ID NO: 346

Those skilled in the art will understand, based upon the present description, that a given API can be described according to the data provided for that API in Table IV or V. The API is a protein comprising a peptide sequence described for that API (preferably comprising a plurality of, more preferably all of, the peptide sequences described for that API) and has a pI of about the value stated for that API (preferably within about 10%, more preferably within about 5% still more preferably within about 1% of the stated value) and has a MW of about the value stated for that API (preferably within about 10%, more preferably within about 5%, still more preferably within about 1% of the stated value).

In one embodiment, CSF from a subject is analyzed for quantitative detection of one or more of the APIs as defined in Table IV(a), optionally combined with one or more of the APIs as defined in Table IV (b): wherein a decreased abundance of the API or APIs (or any suitable combination of them) in the CSF from the subject relative to CSF from a subject or subjects free from AD (*e.g.*, a control sample or a previously determined reference range) indicates the presence of AD.

In another embodiment of the invention, CSF from a subject is analyzed for quantitative detection of one or more of the APIs as defined in Table V(a), optionally combined with the APIs as defined in Table V(b), wherein an increased abundance of the API or APIs (or any suitable combination of them) in CSF from the subject relative to CSF from a subject or subjects free from AD (*e.g.*, a control sample or a previously determined reference range) indicates the presence of AD.

In a further embodiment, CSF from a subject is analyzed for quantitative detection of (a) one or more APIs, or any suitable combination of them, whose decreased abundance indicates the presence of AD, *i.e.*, preferably the APIs set forth in Table IV (a), optionally combined with the APIs set forth in Table IV (b); and (b) one or more APIs, or any suitable

combination of them, whose increased abundance indicates the presence of AD, *i.e.*, preferably the APIs set forth in Table V (a), optionally combined with the APIs as set forth in Table V (b).

In yet a further embodiment, CSF from a subject is analyzed for quantitative detection of one or more APIs and one or more previously known biomarkers of AD (*e.g.*, tau, NTP, A β). In accordance with this embodiment, the abundance of each API and known biomarker relative to a control or reference range indicates whether a subject has AD.

Preferably, the abundance of an API is normalized to an Expression Reference Protein Isoform (ERPI). ERPIs can be identified by partial amino acid sequence characterization of ERFs, which are described above, and which may be accomplished using *e.g.* the methods and apparatus of the Preferred Technology. The partial amino acid sequences of an ERPI are presented in Table VI.

Table VI Expression Reference Protein Isoforms

ERPI#	ERF#	Amino Acid Sequences of Tryptic Digest Peptides	SEQ ID NO:
ERPI-1	ERF-2	ELLDTVTAPQK	SEQ ID NO: 96
		LAAAVSNFGYDLYR	SEQ ID NO: 219
		TSLEDFYLDEER	SEQ ID NO: 385
		ALYYDLISSPDIHGTYK	SEQ ID NO: 27

As shown above, the APIs described herein include previously unknown proteins, as well as isoforms of known proteins where the isoforms were not previously known to be associated with AD. For each API, the present invention additionally provides: (a) a preparation comprising the isolated API; (b) a preparation comprising one or more fragments of an API; and (c) antibodies that bind to said API, to said fragments, or both to said API and to said fragments. As used herein, an API is "isolated" when it is present in a preparation that is substantially free of other proteins, *i.e.*, a preparation in which less than 10% (particularly less than 5%, more particularly less than 1%) of the total protein present is contaminating protein(s). Another protein is a protein or protein isoform having a significantly different pI or MW from those of the isolated API, as determined by 2D electrophoresis. As used herein, a "significantly different" pI or MW is one that permits the

other protein to be resolved from the API on 2D electrophoresis, performed according to the Reference Protocol.

In one embodiment, an isolated protein is provided, that comprises a peptide with the amino acid sequence identified in Table IV or V for an API, said protein having a pI and MW within 10% (particularly within 5%, more particularly within 1%) of the values identified in Table IV or V for that API.

The APIs of the invention can be qualitatively or quantitatively detected by any method known to those skilled in the art, including but not limited to the Preferred Technology described herein, kinase assays, enzyme assays, binding assays and other functional assays, immunoassays, and western blotting. In one embodiment, the APIs are separated on a 2-D gel by virtue of their MWs and pIs and are visualized by staining the gel. In one embodiment, the APIs are stained with a fluorescent dye and imaged with a fluorescence scanner. Sypro Red (Molecular Probes, Inc., Eugene, Oregon) is a suitable dye for this purpose. A preferred fluorescent dye is Pyridinium, 4-[2-[4-(dipentylamino)-2-trifluoromethylphenyl] ethenyl]-1-(sulfobutyl)-, inner salt. See U.S. 6,335,446.

Alternatively, APIs can be detected in an immunoassay. In one embodiment, an immunoassay is performed by contacting a sample with an anti-API capture reagent, e.g. an antibody under conditions such that immunospecific binding can occur if the API is present, and detecting or measuring the amount of any immunospecific binding by the capture reagent, e.g. an antibody. Anti-API antibodies can be produced by the methods and techniques described herein; examples of such antibodies known in the art are set forth in Table VII. These antibodies shown in Table VII are already known to bind to the protein of which the API is itself a family member. Particularly, the anti-API antibody preferentially binds to the API rather than to other isoforms of the same protein. In a particular embodiment, the anti-API antibody binds to the API with at least 2-fold greater affinity, more particularly at least 5-fold greater affinity, still more particularly at least 10-fold greater affinity, than to said other isoforms of the same protein. When the antibodies shown in Table VII do not display the required preferential selectivity for the target API, one skilled in the art can generate additional antibodies by using the API itself for the generation of such antibodies

APIs can be transferred from a gel to a suitable membrane (e.g. a PVDF membrane) and subsequently probed in suitable assays that include, without limitation, competitive and non-competitive assay systems using techniques such as western blots and "sandwich"

immunoassays using anti-API antibodies as described herein, *e.g.*, the antibodies identified in Table VII, or others raised against the APIs of interest as those skilled in the art will appreciate based on the present description. The immunoblots can be used to identify those anti-API antibodies displaying the selectivity required to immuno-specifically differentiate an API from other isoforms encoded by the same gene.

Table VII. Known Antibodies That Recognize APIs or API-Related Polypeptides

Table VII

API #	Antibody	Manufacturer	Cat. No.
API-1	Chromogranin A	BIODESIGN INTERNATIONAL	M54219M
API-3	ANTI-Human CD56 ANTIGEN (NEURAL CELL ADHESION MOLECULE)	RDI RESEARCH DIAGNOSTICS, INC	RDI-CBL159
API-4	Gel	DAKO - 1998 CATALOGUE	A0033
API-6	ANTI-Human CD56 ANTIGEN (NEURAL CELL ADHESION MOLECULE)	RDI RESEARCH DIAGNOSTICS, INC	RDI-CBL159
API-7	Apolipoprotein D, Clone: 36C6, Mab anti-Human, paraffin, IH/WB	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	MED- CLA457
API-10	Goat anti-Clusterin (human)	RDI RESEARCH DIAGNOSTICS, INC	RDI- CLUSTRCabG
API-15	Monoclonal mouse anti- human IgA1	RDI RESEARCH DIAGNOSTICS, INC	RDI-TRK1A2- 2B5
API-16	Monoclonal anti Human Collagen Type VI	BIODESIGN INTERNATIONAL	M22090M
API-22	RABBIT anti-human INSULIN GROWTH FACTOR BINDING PROTEIN 2	RDI RESEARCH DIAGNOSTICS, INC	RDI-IGFBP2abr

Table VII

API #	Antibody	Manufacturer	Cat. No.
API-28	Goat anti-Clusterin (human)	RDI RESEARCH DIAGNOSTICS, INC	RDI- CLUSTRCabG
API-30	Lactic Dehydrogenase (LDH) (H-subunit), Clone: HH-17, Mab anti- Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	BYA- 6019-1
API-33	Albumin, Human, Chicken anti-	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-026-02
API-34	Cystatin C, Rabbit anti- Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	AXL- 574
API-37	Apolipoprotein E, LDL, VLDL, Clone: 3D12, Mab anti-Human, frozen/paraffin	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YM- 5029
API-38	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-39	Transthyretin, Prealbumin, 55kD, Rabbit anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	MED- CLA 193
API-40	Apolipoprotein A1 (HDL), Plasminogen absorbed, Sheep anti- Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	ACL- 20076A
API-42	C3 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-001-02
API-43	Apolipoprotein A1 (HDL), Plasminogen absorbed, Sheep anti- Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	ACL- 20076A
API-44	Transthyretin, Prealbumin, 55kD, Rabbit anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	MED- CLA 193

Table VII

API #	Antibody	Manufacturer	Cat. No.
API-45	Hemopexin, Beta-1, Rabbit anti-Human, precipitating	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YN- RHHPX
API-46	Goat anti-Clusterin (human)	RDI RESEARCH DIAGNOSTICS, INC	RDI- CLUSTRCabG
API-47	ANTI-Human CD56 ANTIGEN (NEURAL CELL ADHESION MOLECULE)	RDI RESEARCH DIAGNOSTICS, INC	RDI-CBL159
API-50	Apolipoprotein E, LDL, VLDL, Clone: 3D12, Mab anti-Human, frozen/paraffin	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YM- 5029
API-52	Alpha-1- Antichymotrypsin, Rabbit anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	AXL- 145/2
API-53	Goat anti-Clusterin (human)	RDI RESEARCH DIAGNOSTICS, INC	RDI- CLUSTRCabG
API-55	Monoclonal anti-Neuron Specific Enolase	BIODESIGN INTERNATIONAL	M37403M
API-58	ANTI-Human CD56 ANTIGEN (NEURAL CELL ADHESION MOLECULE)	RDI RESEARCH DIAGNOSTICS, INC	RDI-CBL159
API-60	Gelsolin, plasma + cytoplasmic, Sheep anti-	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YBG- 4628- 6210
API-62	Apolipoprotein E, LDL, VLDL, Clone: 3D12, Mab anti-Human, frozen/paraffin	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YM- 5029
API-64	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02

Table VII

API #	Antibody	Manufacturer	Cat. No.
API-66	Retinol Binding Protein, Rabbit anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	AXL- 163/2
API-67	Apolipoprotein E, LDL, VLDL, Clone: 3D12, Mab anti-Human, frozen/paraffin	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YM- 5029
API-69	Complement Factor B, C3 proactivator, Rabbit anti- Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	AXL- 466/2
API-72	Gel	DAKO - 1998 CATALOGUE	A0475
API-74	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-75	Fibrinogen, Fibrin I, B- beta chain (B β 1-42), Clone: 18C6, Mab anti- Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	NYB- 18C6
API-76	C3 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-001-02
API-77	Apolipoprotein E, LDL, VLDL, Clone: 3D12, Mab anti-Human, frozen/paraffin	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YM- 5029
API-78	C3 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-001-02
API-79	C3 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-001-02
API-80	C3 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-001-02
API-81	Complement Factor B, C3 proactivator, Rabbit anti- Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	AXL- 466/2
API-82	C3 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-001-02

Table VII

API #	Antibody	Manufacturer	Cat. No.
API-84	Glyceraldehyde-3- Phosphate Dehydrogenase	BIODESIGN INTERNATIONAL	H86504M
API-90	Monoclonal mouse anti- lactoferrin	RDI RESEARCH DIAGNOSTICS, INC	RDI-TRK4L2- LF2B8
API-92	C3 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-001-02
API-93	Monoclonal mouse anti- lactoferrin	RDI RESEARCH DIAGNOSTICS, INC	RDI-TRK4L2- LF2B8
API-95	Albumin, Human, Chicken anti-	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-026-02
API-97	C3 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-001-02
API-98	C8 Complement, Goat anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	BMD- G35
API-101	Albumin, Human, Chicken anti-	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-026-02
API-102	Factor H (Complement), Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-066-02
API-103	Goat anti-Haptoglobin	BIODESIGN INTERNATIONAL	L15320G
API-104	Transferrin, Prealbumin, 55kD, Rabbit anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	MED- CLA 193
API-113	Apolipoprotein E, LDL, VLDL, Clone: 3D12, Mab anti-Human, frozen/paraffin	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YM- 5029
API-118	Monoclonal anti-human Fibrinogen	BIODESIGN INTERNATIONAL	N77190M
API-119	Monoclonal anti-human Fibrinogen	BIODESIGN INTERNATIONAL	N77190M
API-123	AT1 (306)	SANTA CRUZ BIOTECHNOLOGY, INC - RESEARCH ANTIBODIES 98/99	sc-579

Table VII

API #	Antibody	Manufacturer	Cat. No.
API-124	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-126	AT1 (306)	SANTA CRUZ BIOTECHNOLOGY, INC - RESEARCH ANTIBODIES 98/99	sc-579
API-130	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-131	Apolipoprotein A1 (HDL), Plasminogen absorbed, Sheep anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	ACL- 20076A
API-132	Apolipoprotein A1 (HDL), Plasminogen absorbed, Sheep anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	ACL- 20076A
API-134	Goat anti-Clusterin (human)	RDI RESEARCH DIAGNOSTICS, INC	RDI- CLUSTRCabG
API-136	Goat anti-Clusterin (human)	RDI RESEARCH DIAGNOSTICS, INC	RDI- CLUSTRCabG
API-137	Apolipoprotein E, LDL, VLDL, Clone: 3D12, Mab anti-Human, frozen/paraffin	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YM- 5029
API-138	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-140	C3 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-001-02
API-142	Kappa Chain, Mab anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	BMD- 021D

Table VII

API #	Antibody	Manufacturer	Cat. No.
API-143	Tissue Inhibitor of Matrix Metalloproteinase 2 (TIMP2) (NO X w/TIMP1), Clone: 3A4, Mab anti-Human, paraffin, IH	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	MED- CLA498
API-145	ANTI-Human CD56 ANTIGEN (NEURAL CELL ADHESION MOLECULE)	RDI RESEARCH DIAGNOSTICS, INC	RDI-CBL159
API-149	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-150	Sheep anti-Alpha 2 Antiplasmin	BIODESIGN INTERNATIONAL	K90038C
API-161	Apolipoprotein A1 (HDL), Plasminogen absorbed, Sheep anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	ACL- 20076A
API-165	Apolipoprotein D, Clone: 36C6, Mab anti-Human, paraffin, IH/WB	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	MED- CLA457
API-167	Goat anti-Clusterin (human)	RDI RESEARCH DIAGNOSTICS, INC	RDI- CLUSTRCabG
API-168	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-169	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-170	Goat anti-Clusterin (human)	RDI RESEARCH DIAGNOSTICS, INC	RDI- CLUSTRCabG
API-171	Apolipoprotein E, LDL, VLDL, Clone: 3D12, Mab anti-Human, frozen/paraffin	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YM- 5029

Table VII

API #	Antibody	Manufacturer	Cat. No.
API-172	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-173	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-174	Goat anti-Clusterin (human)	RDI RESEARCH DIAGNOSTICS, INC	RDI- CLUSTRCabG
API-175	Transthyretin, Prealbumin, 55kD, Rabbit anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	MED- CLA 193
API-176	Apolipoprotein E, LDL, VLDL, Clone: 3D12, Mab anti-Human, frozen/paraffin	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YM- 5029
API-178	Transthyretin, Prealbumin, 55kD, Rabbit anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	MED- CLA 193
API-179	IGFBP6 (M-20)	SANTA CRUZ BIOTECHNOLOGY, INC - RESEARCH ANTIBODIES 98/99	sc-6008
API-181	C3 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-001-02
API-182	Rabbit anti-14-3-3B (Broadly Reactive)	RDI RESEARCH DIAGNOSTICS, INC	RDI-1433BNabr
API-186	Retinol Binding Protein, Rabbit anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	AXL- 163/2
API-187	Anti-Superoxide Dismutase (Cu/Zn-SOD) IgG fraction (POLYCLONAL)	RDI RESEARCH DIAGNOSTICS, INC	RDI-SODabg
API-188	Transthyretin, Prealbumin, 55kD, Rabbit anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	MED- CLA 193
API-189	Cystatin C, Rabbit anti- Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	AXL- 574

Table VII

API #	Antibody	Manufacturer	Cat. No.
API-191	Goat anti-Haptoglobin	BIODESIGN INTERNATIONAL	L15320G
API-194	ANTI-Human CD56 ANTIGEN (NEURAL CELL ADHESION MOLECULE)	RDI RESEARCH DIAGNOSTICS, INC	RDI-CBL159
API-196	Cystatin C, Rabbit anti- Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	AXL- 574
API-201	Gel	DAKO - 1998 CATALOGUE	A0033
API-215	Cystatin C, Rabbit anti- Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	AXL- 574
API-220	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-221	Apolipoprotein E, LDL, VLDL, Clone: 3D12, Mab anti-Human, frozen/paraffin	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YM- 5029
API-223	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-225	Apolipoprotein E, LDL, VLDL, Clone: 3D12, Mab anti-Human, frozen/paraffin	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YM- 5029
API-233	Apolipoprotein E, LDL, VLDL, Clone: 3D12, Mab anti-Human, frozen/paraffin	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YM- 5029
API-238	Apolipoprotein D, Clone: 36C6, Mab anti-Human, paraffin, IH/WB	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	MED- CLA457
API-239	ANTI-Human CD56 ANTIGEN (NEURAL CELL ADHESION MOLECULE)	RDI RESEARCH DIAGNOSTICS, INC	RDI-CBL159

Table VII

API #	Antibody	Manufacturer	Cat. No.
API-300	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-301	rabbit anti-human alpha-2-macroglobulin	Cambio Ltd.	CA-0427
API-303	Anti-prostaglandin D synthase/Prostaglandin H2-D isomerase	Oxford Biomedical Research	PD-01
API-305	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-308	Gelsolin, plasma + cytoplasmic, Sheep anti-	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YBG- 4628-6210
API-309	rabbit anti-human alpha-2-macroglobulin	Cambio Ltd.	CA-0427
API-310	rabbit anti-human alpha-2-macroglobulin	Cambio Ltd.	CA-0427
API-311	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-312	Gelsolin, plasma + cytoplasmic, Sheep anti-	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YBG- 4628-6210
API-313	Mouse anti-Human pigment epithelium-derived factor (PEDF) (monoclonal) Clone 10F12.2	Chemicon International	MAB1059
API-314	C3 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-001-02
API-315	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-316	Anti-prostaglandin D synthase/Prostaglandin H2-D isomerase	Oxford Biomedical Research	PD-01
API-318	rabbit anti-human alpha-2-macroglobulin	Cambio Ltd.	CA-0427

Table VII

API #	Antibody	Manufacturer	Cat. No.
API-320	Chromogranin A	BIODESIGN INTERNATIONAL	M54219M
API-322	Retinol Binding Protein, Rabbit anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	AXL- 163/2
API-323	Gelsolin, plasma + cytoplasmic, Sheep anti-	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YBG- 4628-6210
API-325	rabbit anti-human alpha-2- macroglobulin	Cambio Ltd.	CA-0427
API-326	Apolipoprotein A1 (HDL), Sheep anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	ACL- 20075AP
API-327	Apolipoprotein E, LDL, VLDL, Clone: 3D12, Mab anti-Human, frozen/paraffin	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YM- 5029
API-330	C3 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-001-02
API-332	Chromogranin A	BIODESIGN INTERNATIONAL	M54219M
API-335	Apolipoprotein A1 (HDL), Sheep anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	ACL- 20075AP
API-336	Anti-prostaglandin D synthase/Prostaglandin H2-D isomerase	Oxford Biomedical Research	PD-01
API-337	rabbit anti-human alpha-2- macroglobulin	Cambio Ltd.	CA-0427
API-339	Anti-prostaglandin D synthase/Prostaglandin H2-D isomerase	Oxford Biomedical Research	PD-01
API-340	Apolipoprotein E, LDL, VLDL, Clone: 3D12, Mab anti-Human, frozen/paraffin	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YM- 5029
API-341	Anti-prostaglandin D synthase/Prostaglandin H2-D isomerase	Oxford Biomedical Research	PD-01

Table VII

API #	Antibody	Manufacturer	Cat. No.
API-346	Anti-prostaglandin D synthase/Prostaglandin H2-D isomerase	Oxford Biomedical Research	PD-01
API-348	Cystatin C, Rabbit anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	AXL- 574
API-349	rabbit anti-human alpha-2-macroglobulin	Cambio Ltd.	CA-0427
API-351	Apolipoprotein A1 (HDL), Sheep anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	ACL- 20075AP
API-353	C7 Complement, Goat anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	BMD- G34
API-355	Apolipoprotein D, Clone: 36C6, Mab anti-Human, paraffin, IH/WB	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	MED- CLA457
API-356	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-361	Alpha-1-Antichymotrypsin, Rabbit anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	AXL- 145/2
API-363	Albumin, Human, Chicken anti-	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-026-02
API-364	Anti-prostaglandin D synthase/Prostaglandin H2-D isomerase	Oxford Biomedical Research	PD-01
API-366	Monoclonal mouse anti-human IgA1	RDI RESEARCH DIAGNOSTICS, INC	RDI-TRK1A2-2B5
API-367	Apolipoprotein A1 (HDL), Sheep anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	ACL- 20075AP
API-368	Albumin, Human, Chicken anti-	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-026-02
API-369	Anti-prostaglandin D synthase/Prostaglandin H2-D isomerase	Oxford Biomedical Research	PD-01

Table VII

API #	Antibody	Manufacturer	Cat. No.
API-370	Anti-prostaglandin D synthase/Prostaglandin H2-D isomerase	Oxford Biomedical Research	PD-01
API-378	Monoclonal anti-human Fibrinogen	BIODESIGN INTERNATIONAL	N77190M
API-380	Chromogranin A	BIODESIGN INTERNATIONAL	M54219M
API-381	Monoclonal anti-human Fibrinogen	BIODESIGN INTERNATIONAL	N77190M
API-382	C4 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-032-02
API-383	C3 Complement, Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-001-02
API-388	Gelsolin, plasma + cytoplasmic, Sheep anti-	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YBG- 4628-6210
API-390	Albumin, Human, Chicken anti-	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-026-02
API-391	Factor H (Complement), Chicken anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-066-02
API-392	Gelsolin, plasma + cytoplasmic, Sheep anti-	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YBG- 4628-6210
API-393	rabbit anti-human alpha-2-macroglobulin	Cambio Ltd.	CA-0427
API-394	Rabbit Polyclonal Anti-Human Ceruloplasmin	DAKO CORPORATION	A0031
API-400	Goat anti-human Carbonic Anhydrase I (polyclonal)	Abcam Ltd.	ab6618
API-401	Cystatin C, Rabbit anti-Human	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	AXL- 574
API-403	Hemopexin, Beta-1, Rabbit anti-Human, precipitating	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YN- RHHPX
API-407	rabbit anti-human alpha-2-macroglobulin	Cambio Ltd.	CA-0427

Table VII

API #	Antibody	Manufacturer	Cat. No.
API-408	Monoclonal mouse anti-human IgA1	RDI RESEARCH DIAGNOSTICS, INC	RDI-TRK1A2-2B5
API-409	Anti-Human Contactin, Mouse monoclonal	BD Biosciences	610579
API-410	Albumin, Human, Chicken anti-	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	IMS- 01-026-02
API-411	Anti-Human Contactin, Mouse monoclonal	BD Biosciences	610579
API-414	Hemopexin, Beta-1, Rabbit anti-Human, precipitating	ACCURATE CHEMICAL & SCIENTIFIC CORPORATION	YN- RHHPX
API-415	goat anti-Pyruvate Kinase (polyclonal)	SANTA CRUZ BIOTECHNOLOGY, INC - RESEARCH ANTIBODIES 98/99	sc-7140
API-416	Goat anti-Clusterin (human)	RDI RESEARCH DIAGNOSTICS, INC	RDI-CLUSTRCabG
API-417	Anti-prostaglandin D synthase/Prostaglandin H2-D isomerase	Oxford Biomedical Research	PD-01

*Further information about these antibodies can be obtained from their commercial sources at: ACCURATE CHEMICAL & SCIENTIFIC CORPORATION

<http://www.accuratechemical.com/>; BIODESIGN INTERNATIONAL -

<http://www.biodesign.com/>; RDI RESEARCH DIAGNOSTICS, INC -

5 <http://www.researchd.com/>; SANTA CRUZ BIOTECHNOLOGY, INC -

<http://www.scbt.com/>.

In one embodiment, binding of antibody in tissue sections can be used to detect API localization or the level of one or more APIs. In a specific embodiment, antibody to an API can be used to assay a tissue sample (*e.g.*, a brain biopsy) from a subject for the level of the API where a substantially changed level of API is indicative of AD. As used herein, a "substantially changed level" means a level that is increased or decreased compared with the level in a subject free from AD or a reference level. If desired, the comparison can be performed with a matched sample from the same subject, taken from a portion of the body

not affected by AD.

Any suitable immunoassay can be used to detect an API, including, without limitation, competitive and non-competitive assay systems using techniques such as western blots, radioimmunoassays, ELISAs (enzyme linked immunosorbent assays),
5 “sandwich” immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays and protein A immunoassays.

For example, an API can be detected in a fluid sample (*e.g.*, CSF, blood, urine, or
10 tissue homogenate) by means of a two-step sandwich assay. In the first step, a capture reagent (*e.g.*, an anti-API antibody) is used to capture the API. Examples of such antibodies known in the art are set forth in Table VII. The capture reagent can optionally be immobilized on a solid phase. In the second step, a directly or indirectly labeled detection reagent is used to detect the captured API. In one embodiment, the detection
15 reagent is a lectin. A lectin can be used for this purpose that preferentially binds to the API rather than to other isoforms that have the same core protein as the API or to other proteins that share the antigenic determinant recognized by the antibody. In a preferred embodiment, the chosen lectin binds to the API with at least 2-fold greater affinity, more particularly at least 5-fold greater affinity, still more preferably at least 10-fold greater
20 affinity, than to said other isoforms that have the same core protein as the API or to said other proteins that share the antigenic determinant recognized by the antibody. Based on the present description, a lectin that is suitable for detecting a given API can readily be identified by those skilled in the art using methods well known in the art, for instance upon testing one or more lectins enumerated in Table I on pages 158-159 of Sumar et al., *Lectins as Indicators of Disease-Associated Glycoforms*, In: Gabius H-J & Gabius S (eds.), 1993,
25 *Lectins and Glycobiology*, at pp. 158-174. Lectins with the desired oligosaccharide specificity can be identified, for example, by their ability to detect the API in a 2D gel, in a replica of a 2D gel following transfer to a suitable solid substrate such as a nitrocellulose membrane, or in a two-step assay following capture by an antibody. In an alternative
30 embodiment, the detection reagent is an antibody, *e.g.*, an antibody that immunospecifically detects other post-translational modifications, such as an antibody that immunospecifically binds to phosphorylated amino acids. Examples of such antibodies include those that bind to phosphotyrosine (BD Transduction Laboratories, catalog nos.: P11230-050/P11230-150;

P11120; P38820; P39020), those that bind to phosphoserine (Zymed Laboratories Inc., South San Francisco, CA, catalog no. 61-8100) and those that bind to phosphothreonine (Zymed Laboratories Inc., South San Francisco, CA, catalog nos. 71-8200, 13-9200).

If desired, a gene encoding an API, a related gene (e.g. a gene having sequence
5 homology), or related nucleic acid sequences or subsequences, including complementary sequences, can also be used in hybridization assays. A nucleotide encoding an API, or subsequences thereof comprising at least 8 nucleotides, preferably at least 12 nucleotides, and most preferably at least 15 nucleotides can be used as a hybridization probe.

Hybridization assays can be used for detection, treatment, diagnosis, or monitoring of
10 conditions, disorders, or disease states, associated with aberrant expression of genes encoding APIs, or for differential diagnosis of subjects with signs or symptoms suggestive of AD. In particular, such a hybridization assay can be carried out by a method comprising contacting a subject's sample containing nucleic acid with a nucleic acid probe capable of hybridizing to a DNA or RNA that encodes an API, under conditions such that
15 hybridization can occur, and detecting or measuring any resulting hybridization.

Nucleotides can be used for therapy of subjects having AD, as described below.

The invention also provides diagnostic kits, comprising an anti-API capture reagent. In addition, such a kit may optionally comprise one or more of the following: (1) instructions for using the anti-API capture reagent for diagnosis, therapeutic monitoring or
20 any suitable combination of these applications; (2) a labeled binding partner to the capture reagent; (3) a solid phase (such as a reagent strip) upon which the anti-API capture reagent is immobilized; and (4) a label or insert indicating regulatory approval for diagnostic, prognostic or therapeutic use or any suitable combination thereof. If no labeled binding partner to the antibody is provided, the anti-API capture reagent itself can be labeled with a
25 detectable marker, *e.g.*, a chemiluminescent, enzymatic, fluorescent, or radioactive moiety.

The invention also provides a kit comprising a nucleic acid probe capable of hybridizing to RNA encoding an API. In a specific embodiment, a kit comprises in one or more containers a pair of primers (*e.g.*, each in the size range of 6-30 nucleotides, more preferably 10-30 nucleotides and still more preferably 10-20 nucleotides) that under
30 appropriate reaction conditions can prime amplification of at least a portion of a nucleic acid encoding an API, such as by polymerase chain reaction (see, *e.g.*, Innis et al., 1990, PCR Protocols, Academic Press, Inc., San Diego, CA), ligase chain reaction (see EP 320,308) use of Q β replicase, cyclic probe reaction, or other methods known in the art.

Kits are also provided which allow for the detection of a plurality of APIs or a plurality of nucleic acids each encoding an API. A kit can optionally further comprise a predetermined amount of an isolated API protein or a nucleic acid encoding an API, *e.g.*, for use as a standard or control.

5

5.5 Statistical Techniques for Identifying AFs, APIs and API Clusters

Uni-variate differential analysis tools, such as fold changes, wilcoxon rank sum test and t-test, are useful in identifying individual AFs or APIs that are diagnostically associated with AD or in identifying individual APIs that regulate the disease process. However, those skilled in the art will appreciate that the disease process is associated with a suitable combination of AFs or APIs (and to be regulated by a suitable combination of APIs), rather than individual AFs and APIs in isolation. The strategies for discovering such suitable combinations of AFs and APIs differ from those for discovering individual AFs and APIs. In such cases, each individual AF and API can be regarded as one variable and the disease can be regarded as a joint, multi-variate effect caused by interaction of these variables.

10

15

The following steps can be used to identify markers from data produced by the Preferred Technology.

The first step is to identify a collection of AFs or APIs that individually show significant association with AD. The association between the identified individual AFs or individual APIs and A need not be as highly significant when a collection of AFs and APIs is used as a diagnostic as is desirable when an individual AF or API is used as a diagnostic. Any of the tests discussed above (fold changes, Wilcoxon rank sum test, etc.) can be used at this stage. Once a suitable collection of AFs or APIs has been identified, a sophisticated multi-variate analysis capable of identifying clusters can then be used to estimate the significant multivariate associations with AD.

20

25

Linear Discriminant Analysis (LDA) is one such procedure, which can be used to detect significant association between a cluster of variables (*i.e.*, AFs or APIs) and AD. In performing LDA, a set of weights is associated with each variable (*i.e.*, AF or API) so that the linear combination of weights and the measured values of the variables can identify the disease state by discriminating between subjects having AD and subjects free from AD. Enhancements to the LDA allow stepwise inclusion (or removal) of variables to optimize the discriminant power of the model. The result of the LDA is therefore a cluster of AFs, or APIs, which can be used for diagnosis, treatment or development of pharmaceutical

30

products. Other enhanced variations of LDA, such as Flexible Discriminant Analysis permit the use of non-linear combinations of variables to discriminate a disease state from a state in which there is no disease. The results of the discriminant analysis can be verified by post-hoc tests and also by repeating the analysis using alternative techniques such as classification trees.

A further category of AFs or APIs can be identified by qualitative measures by comparing the percentage feature presence of an AF or API of one group of samples (*e.g.*, samples from diseased subjects) with the percentage feature presence of an AF or API in another group of samples (*e.g.*, samples from control subjects). The “percentage feature presence” of an AF or API is the percentage of samples in a group of samples in which the AF or API is detectable by the detection method of choice. For example, if an AF is detectable in 95 percent of samples from diseased subjects, the percentage feature presence of that AF in that sample group is 95 percent. If only 5 percent of samples from non-diseased subjects have detectable levels of the same AF, detection of that AF in the sample of a subject would suggest that it is likely that the subject has AD.

Similarly, the use of “nearest neighbor analysis” comprises the visual inspection of a particular group or cluster of AFs or APIs and the consequent association of particular such features or isoforms on the basis of their physical closeness and similarity in appearance, to other features or isoforms that are found to demonstrate a significant presence in samples taken from diseased subjects.

5.6. Use in Clinical Studies

The diagnostic methods and compositions of the present invention can assist in monitoring a clinical study, *e.g.* to evaluate therapies for AD. In one embodiment, chemical compounds are tested for their ability to restore AF or API levels in a subject having AD to levels found in subjects free from AD or, in a treated subject (*e.g.* after treatment with a cholinesterase inhibitor), to preserve AF or API levels at or near levels seen in subjects free from AD. The levels of one or more AFs or APIs can be assayed.

In another embodiment, the methods and compositions of the present invention are used to screen individuals for entry into a clinical study to identify individuals having AD; individuals already having AD can then be excluded from the study or can be placed in a separate cohort for treatment or analysis. If desired, the candidates can concurrently be screened to identify individuals with Lewy Body disease and/or senile dementia or a known

measure of AD; procedures for these screens are well known in the art (Harding and Halliday, 1998, Neuropathol. Appl. Neurobiol. 24:195-201).

5.7. Purification of APIs

5 In particular aspects, the invention provides isolated mammalian APIs, preferably human APIs, API fragments, API -related polypeptides or API -fusion proteins which comprise an antigenic determinant (*i.e.*, can be recognized by an antibody) or which are otherwise functionally active, as well as nucleic acid sequences encoding the foregoing. “Functionally active” as used herein refers to material displaying one or more functional
10 activities associated with a full-length (wild-type) API, *e.g.*, binding to an API substrate or API binding partner, antigenicity (binding to an anti-API antibody), immunogenicity, enzymatic activity and the like.

In specific embodiments, the invention provides fragments of an API comprising at least 5 amino acids, at least 10 amino acids, at least 50 amino acids, or at least 75 amino
15 acids. Fragments lacking some or all of the regions of an API are also provided, as are proteins (*e.g.*, fusion proteins) comprising such fragments. Nucleic acids encoding the foregoing are provided.

Once a recombinant nucleic acid which encodes the API, a portion of the API, or a precursor of the API is identified, the gene product can be analyzed. This can be achieved
20 by assays based on the physical or functional properties of the given product, including, for example, radioactive labeling of the product followed by analysis by gel electrophoresis, immunoassay, etc.

The APIs identified herein can be isolated and purified by standard methods including chromatography (*e.g.*, ion exchange, affinity, and sizing column
25 chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins.

Alternatively, once a recombinant nucleic acid that encodes the API is identified, the entire amino acid sequence of the API can be deduced from the nucleotide sequence of the gene-coding region contained in the recombinant nucleic acid. As a result, the protein
30 can be synthesized by standard chemical methods known in the art (*e.g.*, see Hunkapiller et al., 1984, Nature 310:105-111).

In another alternative embodiment, native APIs can be purified from natural sources, by standard methods such as those described above (*e.g.*, immunoaffinity

purification).

In a preferred embodiment, APIs are isolated by the Preferred Technology described *supra*. For preparative-scale runs, a narrow-range "zoom gel" having a pH range of 2 pH units or less is preferred for the isoelectric step, according to the method described in
5 Westermeier, 1993, Electrophoresis in Practice (VCH, Weinheim, Germany), pp. 197-209 (which is incorporated herein by reference in its entirety); this modification permits a larger quantity of a target protein to be loaded onto the gel, and thereby increases the quantity of isolated API that can be recovered from the gel. When used in this way for preparative-
scale runs, the Preferred Technology typically provides up to 100 ng, and can provide up to
10 1000 ng, of an isolated API in a single run. Those of skill in the art will appreciate that a zoom gel can be used in any separation strategy which employs gel isoelectric focusing.

The invention thus provides an isolated API, an isolated API fragment, an isolated API -related polypeptide or an isolated API -fusion protein; any of the foregoing can be produced by recombinant DNA techniques or by chemical synthetic methods.

5.8 Isolation of DNA Encoding an API

Particular embodiments for the cloning of a gene encoding an API are presented below by way of example and not of limitation.

The nucleotide sequences of the present invention, including DNA and RNA, and
20 comprising a sequence encoding an API, API fragment, API -related polypeptide or API -fusion protein, may be synthesized using methods known in the art, such as using conventional chemical approaches or polymerase chain reaction (PCR) amplification. The nucleotide sequences of the present invention also permit the identification and cloning of the gene encoding an API homolog or API ortholog including, for example, by screening
25 cDNA libraries, genomic libraries or expression libraries.

For example, to clone a gene encoding an API by PCR techniques, anchored degenerate oligonucleotides (or a set of most likely oligonucleotides) can be designed for all API peptide fragments identified as part of the same protein. PCR reactions under a variety of conditions can be performed with relevant cDNA and genomic DNAs (*e.g.*, from
30 brain tissue or from cells of the immune system) from one or more species. Also vectorette reactions can be performed on any available cDNA and genomic DNA using the oligonucleotides (which preferably are nested) as above. Vectorette PCR is a method that enables the amplification of specific DNA fragments in situations where the sequence of

only one primer is known. Thus, it extends the application of PCR to stretches of DNA where the sequence information is only available at one end. (Arnold C, 1991, PCR Methods Appl. 1(1): 39-42; Dyer KD, Biotechniques, 1995, 19(4):550-2). Vectorette PCR may be performed with probes that are, for example, anchored degenerate oligonucleotides (or most likely oligonucleotides) coding for API peptide fragments, using as a template a genomic library or cDNA library pools.

Anchored degenerate oligonucleotides (and most likely oligonucleotides) can be designed for all API peptide fragments. These oligonucleotides may be labelled and hybridized to filters containing cDNA and genomic DNA libraries. Oligonucleotides to different peptides from the same protein will often identify the same members of the library. The cDNA and genomic DNA libraries may be obtained from any suitable or desired mammalian species, for example from humans.

Nucleotide sequences comprising a nucleotide sequence encoding an API or API fragment of the present invention are useful, for example, for their ability to hybridize selectively with complementary stretches of genes encoding other proteins. Depending on the application, a variety of hybridization conditions may be employed to obtain nucleotide sequences at least about 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 99% identical, or 100% identical, to the sequence of a nucleotide encoding an API.

For a high degree of selectivity, relatively stringent conditions are used to form the duplexes, such as low salt or high temperature conditions.

Hybridization conditions can also be rendered more stringent by the addition of increasing amounts of formamide, to destabilize the hybrid duplex. Thus, particular hybridization conditions can be readily manipulated, and will generally be chosen depending on the desired results. In general, convenient hybridization temperatures in the presence of 50% formamide are: 42°C for a probe which is 95 to 100% identical to the fragment of a gene encoding an API, 37°C for 90 to 95% identity and 32°C for 70 to 90% identity.

In the preparation of genomic libraries, DNA fragments are generated, some of which will encode parts or the whole of an API. Any suitable method for preparing DNA fragments may be used in the present invention. For example, the DNA may be cleaved at specific sites using various restriction enzymes. Alternatively, one may use DNase in the presence of manganese to fragment the DNA, or the DNA can be physically sheared, as for

example, by sonication. The DNA fragments can then be separated according to size by standard techniques, including but not limited to agarose and polyacrylamide gel electrophoresis, column chromatography and sucrose gradient centrifugation. The DNA fragments can then be inserted into suitable vectors, including but not limited to plasmids, cosmids, bacteriophages lambda or T4, and yeast artificial chromosome (YAC). (See, e.g., Sambrook et al., 1989, *Molecular Cloning, A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; Glover, D.M. (ed.), 1985, *DNA Cloning: A Practical Approach*, MRL Press, Ltd., Oxford, U.K. Vol. I, II; Ausubel F.M. et al., eds., 1989, *Current Protocols in Molecular Biology*, Vol. I, Green Publishing Associates, Inc., and John Wiley & sons, Inc., New York). The genomic library may be screened by nucleic acid hybridization to labeled probe (Benton and Davis, 1977, *Science* 196:180; Grunstein and Hogness, 1975, *Proc. Natl. Acad. Sci. U.S.A.* 72:3961).

Based on the present description, the genomic libraries may be screened with labeled degenerate oligonucleotide probes corresponding to the amino acid sequence of any peptide of the API using optimal approaches well known in the art. Any probe used is at least 10 nucleotides, at least 15 nucleotides, at least 20 nucleotides, at least 25 nucleotides, at least 30 nucleotides, at least 40 nucleotides, at least 50 nucleotides, at least 60 nucleotides, at least 70 nucleotides, at least 80 nucleotides, or at least 100 nucleotides. Preferably a probe is 10 nucleotides or longer, and more preferably 15 nucleotides or longer.

In Tables IV and V above, some APIs disclosed herein correspond to isoforms of previously identified proteins encoded by genes whose sequences are publicly known. To screen such a gene, any probe may be used that is complementary to the gene or its complement; preferably the probe is 10 nucleotides or longer, more preferably 15 nucleotides or longer. The SWISS-PROT and trEMBL databases (held by the Swiss Institute of Bioinformatics (SIB) and the European Bioinformatics Institute (EBI) which are available at <http://www.expasy.ch/>) and the GenBank database (held by the National Institute of Health (NIH) which is available at <http://www.ncbi.nlm.nih.gov/>) provide protein sequences comprising the amino acid sequences listed for the APIs in Tables IV and V under the following accession numbers and each sequence is incorporated herein by reference:

*Table VIII. Nucleotide sequences encoding APIs, API Related Proteins, or ERPIs***Table VIII**

AF#	API#	Accession Numbers of Identified Sequences
AF-1	API-47	O15179
AF-1	API-242	AAF03259
AF-2	API-1	P10645
AF-3	API-48	Q9UBQ6
AF-5	API-49	P19021
AF-6	API-2	P47868
AF-8	API-194	AAB60937
AF-9	API-3	O15179
AF-10	API-50	P02649
AF-10	API-51	P55290
AF-13	API-4	P01023
AF-14	API-52	P01011
AF-14	API-243	P04004
AF-15	API-53	P10909
AF-15	API-244	AAC50896
AF-16	API-54	P19021
AF-17	API-5	O43505
AF-18	API-55	P09104
AF-18	API-245	P51693
AF-21	API-6	O15179
AF-22	API-56	O94985
AF-22	API-57	AAD05198
AF-23	API-7	P05090
AF-23	API-8	P05060
AF-24	API-9	P19021
AF-25	API-10	P10909
AF-26	API-14	P41222

Table VIII

AF#	API#	Accession Numbers of Identified Sequences
AF-27	API-15	P20758
AF-27	API-58	O15179
AF-28	API-16	Q04857
AF-28	API-59	P00450
AF-29	API-196	P01034
AF-30	API-17	P36955
AF-31	API-60	P06396
AF-32	API-18	O43505
AF-34	API-61	Q14800
AF-35	API-62	P02649
AF-37	API-19	P40925
AF-38	API-63	Q92876
AF-39	API-64	P01028
AF-39	API-65	Q9UBQ6
AF-40	API-20	P17174
AF-41	API-22	P18065
AF-42	API-66	P02753
AF-43	API-67	P02649
AF-43	API-68	P01034
AF-44	API-69	P00751
AF-44	API-70	P10643
AF-45	API-23	P36955
AF-46	API-24	P05067
AF-46	API-197	P13645
AF-46	API-198	P13647
AF-47	API-25	O43505
AF-48	API-71	Q99435
AF-49	API-26	Q92876
AF-49	API-27	P41222
AF-50	API-72	P01871

Table VIII

AF#	API#	Accession Numbers of Identified Sequences
AF-50	API-73	P00748
AF-50	API-199	P04196
AF-50	API-200	AAC34741
AF-51	API-28	P10909
AF-51	API-30	P07195
AF-52	API-74	P01028
AF-53	API-33	P02768
AF-54	API-221	P02649
AF-55	API-34	P01034
AF-56	API-75	P02675
AF-56	API-246	P06396
AF-57	API-35	P35527
AF-57	API-76	P01024
AF-57	API-222	P13645
AF-58	API-77	P02649
AF-59	API-36	P36955
AF-60	API-37	P02649
AF-61	API-78	P01024
AF-62	API-38	P01028
AF-63	API-79	P01024
AF-64	API-80	P01024
AF-65	API-81	P00751
AF-65	API-223	P01028
AF-66	API-82	P01024
AF-66	API-83	P01034
AF-67	API-39	P02766
AF-68	API-84	P04406
AF-68	API-85	P04279
AF-69	API-40	P06727
AF-69	API-247	CAB89302

Table VIII

AF#	API#	Accession Numbers of Identified Sequences
AF-70	API-41	P36222
AF-70	API-224	229552 (gb)
AF-71	API-42	P01024
AF-72	API-43	P06727
AF-73	API-44	P02766
AF-74	API-45	P02790
AF-74	API-248	P01024
AF-75	API-46	P10909
AF-75	API-225	P02649
AF-76	API-86	P19021
AF-79	API-201	P01023
AF-81	API-88	AAA52900
AF-81	API-202	AAC48775
AF-82	API-89	P09571
AF-83	API-90	P09571
AF-84	API-91	P09571
AF-85	API-92	P01024
AF-85	API-93	P09571
AF-87	API-95	P08835
AF-89	API-97	P01024
AF-90	API-98	P07358
AF-91	API-99	S64635
AF-100	API-101	P08835
AF-103	API-102	Q03591
AF-104	API-103	P06866
AF-105	API-104	P02766
AF-107	API-107	Q16270
AF-107	API-210	BAA25513
AF-108	API-108	O88812
AF-117	API-113	P02649

Table VIII

AF#	API#	Accession Numbers of Identified Sequences
AF-119	API-114	P02023
AF-121	API-116	P04469
AF-123	API-118	P02671
AF-124	API-119	P02671
AF-125	API-120	Q12805
AF-126	API-121	O43532
AF-126	API-122	AAF02676
AF-127	API-123	P01019
AF-128	API-124	P01028
AF-129	API-125	P02774
AF-129	API-126	P01019
AF-130	API-127	P36955
AF-130	API-128	O43505
AF-132	API-130	P01028
AF-133	API-131	P06727
AF-134	API-132	P06727
AF-137	API-134	P10909
AF-137	API-135	P05156
AF-137	API-232	Q9Y6R4
AF-137	API-233	P02649
AF-137	API-234	P33176
AF-139	API-136	P10909
AF-139	API-137	P02649
AF-140	API-138	P01028
AF-141	API-139	P09871
AF-142	API-140	P01024
AF-142	API-141	Q92876
AF-143	API-142	751423A
AF-144	API-143	P16035
AF-149	API-214	AAB60937

Table VIII

AF#	API#	Accession Numbers of Identified Sequences
AF-150	API-144	Q02246
AF-151	API-145	O15179
AF-152	API-146	P43652
AF-152	API-147	P51693
AF-152	API-148	P00734
AF-153	API-149	P01028
AF-154	API-150	P08697
AF-154	API-151	P02748
AF-154	API-152	P01877
AF-155	API-215	P01034
AF-156	API-153	AF177396
AF-157	API-155	O43505
AF-159	API-158	P36955
AF-159	API-159	O43505
AF-159	API-160	P07339
AF-161	API-161	P06727
AF-161	API-162	P36955
AF-161	API-163	P02570
AF-163	API-165	P05090
AF-163	API-166	P05060
AF-164	API-167	P10909
AF-165	API-168	P01028
AF-166	API-169	P01028
AF-167	API-170	P10909
AF-167	API-171	AAD02505
AF-168	API-237	P13645
AF-168	API-172	P01028
AF-169	API-173	P01028
AF-170	API-174	P10909
AF-170	API-175	P02766

Table VIII

AF#	API#	Accession Numbers of Identified Sequences
AF-170	API-176	AAD02505
AF-171	API-177	P41222
AF-171	API-178	P02766
AF-172	API-179	P24592
AF-172	API-180	AAD51475
AF-173	API-181	P01024
AF-174	API-182	P29361
AF-175	API-183	P41222
AF-176	API-184	P41222
AF-178	API-185	P47971
AF-178	API-217	P41222
AF-178	API-219	Q29562
AF-179	API-186	P02753
AF-180	API-220	P01028
AF-181	API-187	P00441
AF-182	API-188	P02766
AF-183	API-189	P01034
AF-184	API-190	P36955
AF-185	API-191	P00737
AF-185	API-192	Q12805
AF-186	API-238	P05090
AF-187	API-239	O15179
AF-190	API-240	NP_055108 (gb)
AF-192	API-241	P19021
AF-201	API-375	Q15117
AF-204	API-300	P01028
AF-205	API-301	P01023
AF-206	API-302	1160616
AF-207	API-303	P41222
AF-208	API-304	1160616

Table VIII

AF#	API#	Accession Numbers of Identified Sequences
AF-209	API-305	P01028
AF-209	API-306	P07358
AF-210	API-307	1160616
AF-211	API-308	P06396
AF-212	API-309	P01023
AF-213	API-310	P01023
AF-214	API-376	4507889
AF-214	API-377	Q9NYQ9
AF-215	API-311	P01028
AF-216	API-312	P06396
AF-218	API-313	P36955
AF-219	API-378	P02671
AF-220	API-314	P01024
AF-224	API-379	P02749
AF-225	API-315	P01028
AF-226	API-316	P41222
AF-227	API-380	P10645
AF-229	API-317	10092631
AF-229	API-369	P41222
AF-230	API-318	P01023
AF-231	API-319	4502945
AF-232	API-320	P10645
AF-233	API-321	P19021
AF-234	API-322	P02753
AF-235	API-323	P06396
AF-236	API-381	P02671
AF-237	API-382	P01028
AF-237	API-383	P01024
AF-237	API-384	P07358
AF-238	API-324	Q9UBQ6

Table VIII

AF#	API#	Accession Numbers of Identified Sequences
AF-239	API-325	P01023
AF-240	API-326	P06727
AF-241	API-327	P02649
AF-242	API-328	P05060
AF-243	API-329	Q99435
AF-244	API-330	P01024
AF-245	API-331	1732416
AF-248	API-385	P25311
AF-249	API-332	P10645
AF-250	API-333	P05092
AF-252	API-334	P05060
AF-253	API-335	P02647
AF-254	API-336	P41222
AF-255	API-337	P01023
AF-256	API-338	2190958
AF-257	API-339	P41222
AF-258	API-340	P02649
AF-259	API-341	P41222
AF-260	API-342	179762
AF-261	API-386	Q15668
AF-262	API-387	Q15117
AF-263	API-343	7341255
AF-264	API-344	2190958
AF-265	API-388	P06396
AF-267	API-345	2190958
AF-268	API-346	P41222
AF-269	API-389	P08603
AF-270	API-390	P02768
AF-271	API-347	P01043
AF-272	API-391	Q03591

Table VIII

AF#	API#	Accession Numbers of Identified Sequences
AF-274	API-348	P01034
AF-275	API-349	P01023
AF-277	API-350	P08294
AF-278	API-351	P02647
AF-281	API-352	Q02246
AF-283	API-392	P06396
AF-284	API-393	P01023
AF-285	API-394	P00450
AF-285	API-395	O00533
AF-289	API-353	P10643
AF-291	API-354	7662374
AF-295	API-396	Q15117
AF-296	API-397	15636798
AF-299	API-355	P05090
AF-300	API-398	P08603
AF-303	API-399	P08603
AF-306	API-400	P00915
AF-307	API-356	P01028
AF-308	API-357	P08603
AF-309	API-401	P01034
AF-310	API-358	P02774
AF-311	API-402	P04003
AF-311	API-403	P02790
AF-312	API-359	P32119
AF-313	API-360	P04220
AF-317	API-404	P12960
AF-318	API-405	P04003
AF-319	API-406	14919433
AF-323	API-407	P01023
AF-323	API-408	P01876

Table VIII

AF#	API#	Accession Numbers of Identified Sequences
AF-323	API-409	Q12860
AF-324	API-361	P01011
AF-325	API-362	P36222
AF-326	API-363	P02768
AF-327	API-410	P02768
AF-328	API-411	Q12860
AF-330	API-364	P41222
AF-331	API-412	P55290
AF-331	API-413	P17900
AF-332	API-365	P17174
AF-333	API-366	P01876
AF-334	API-414	P02790
AF-334	API-415	2745741
AF-335	API-367	P02647
AF-335	API-370	P41222
AF-336	API-416	P10909
AF-336	API-417	P41222
AF-337	API-418	P02749
AF-338	API-368	P02768

When no nucleotide sequence is known that encodes a protein comprising an amino acid sequence of a given API, degenerate probes can be used for screening. In Table IX, a degenerate set of probes is provided for API-111 and API-112. The partial amino acid sequences listed in Table IX were derived from manual interpretation of the tandem mass spectra of tryptic digest peptides of the API. In the method of tandem mass spectroscopy used for sequencing peptides in the present invention, the following pairs of amino acids could not be distinguished from each other: leucine and isoleucine; and, under certain circumstances, phenylalanine and oxidized methionine. As used herein, an amino acid sequence "as determined by mass spectrometry" refers to the set of amino acid sequences

containing at the indicated positions, one or other member of the designated pairs of amino acids. For example, the amino acid sequence P[L/I]A indicates the amino acid sequences PLA and PIA. As will be obvious to one of skill in the art, a sequence containing n designated pairs indicates 2^n amino acid sequences. In Table IX, each possible amino acid sequence is listed for each sequence determined by mass spectroscopy, and preferred and fully degenerate sets of probes for each possible amino acid sequence are provided.

Table IX. Amino Acid Sequences and Probes for APIs

AF#	API#	Amino Acid Sequences of Tryptic Digest Peptides as Determined by Mass Spectrometry				Preferred Probes	Degenerate Probes
		Mass of singly protonated peptide (Da)*	Partial sequence	N-terminal Mass (Da)*	C-terminal Mass (Da)*		
AF-114	API-111	1097.57	HQV	0	733.50	CACCAGG TG	CAYCARG TN
AF-114	API-112	1547.74	PGLGM (SEQ ID NO 468)	0	1076.63	CCCGGCC TGGGCAT G (SEQ ID NO: 469)	CCNGGNY TNGGNAT G (SEQ ID NO: 470)
AF-114	API-112	1547.74	PGLGF (SEQ ID NO: 471)	0	1076.63	CCCGGCC TGGGCTT C (SEQ ID NO: 472)	CCNGGNY TNGGNTT Y (SEQ ID NO: 473)
AF-114	API-112	1547.74	PGIGM (SEQ ID NO: 474)	0	1076.63	CCCGGCA TCGGCAT G (SEQ ID NO: 475)	CCNGGNA THGGNAT G (SEQ ID NO: 476)
AF-114	API-112	1547.74	PGIGF (SEQ ID NO: 477)	0	1076.63	CCCGGCA TCGGCTT C (SEQ ID NO: 478)	CCNGGNA THGGNTT Y (SEQ ID NO: 479)
AF-114	API-112	1547.74	GPLGM (SEQ ID NO: 480)	0	1076.63	GGCCCCC TGGGCAT G (SEQ ID	GGNCCNY TNGGNAT G (SEQ ID

AF#	API#	Amino Acid Sequences of Tryptic Digest Peptides as Determined by Mass Spectrometry				Preferred Probes	Degenerate Probes
		Mass of singly protonated peptide (Da)*	Partial sequence	N-terminal Mass (Da)*	C-terminal Mass (Da)*		
						NO: 481)	NO: 482)
AF-114	API-112	1547.74	GPLGF (SEQ ID NO: 483)	0	1076.63	GGCCCCC TGGGCTT C (SEQ ID NO: 484)	GGNCCNY TNGGNTT Y (SEQ ID NO: 485)
AF-114	API-112	1547.74	GPIGM (SEQ ID NO: 486)	0	1076.63	GGCCCCA TCGGCAT G (SEQ ID NO: 487)	GGNCCNA THGGNAT G (SEQ ID NO: 488)
AF-114	API-112	1547.74	GPIGF (SEQ ID NO: 489)	0	1076.63	GGCCCCA TCGGCTT C (SEQ ID NO: 490)	GGNCCNA THGGNTT Y (SEQ ID NO: 491)

*The masses determined by mass spectrometry have an error of mass measurement of 100 parts-per-million (ppm) or less. For a given measured mass, M, having an error of mass measurement of z ppm, the error of mass measurement can be calculated as $(M \times z \div 1000000)$.

5

As used herein, the “mass of the singly protonated peptide” is the mass of the singly protonated tryptic digest peptide measured by mass spectrometry (having an error of measurement of approximately 100 parts-per-million or less) and corresponds to the total mass of the constituent amino acid residues of the peptide with the addition of a water molecule (H₂O) and a single proton (H⁺). As used herein, an “amino acid residue” refers to

10 an amino acid residue of the general structure: -NH-CHR-CO- and which have the

following symbols, elemental compositions and monoisotopic masses:

Amino acid residue elemental compositions and monoisotopic masses

Amino Acid	Symbol	Elemental Composition	Monoisotopic mass (Da)
Alanine	A	C ₃ H ₅ NO	71.037114
Arginine	R	C ₆ H ₁₂ N ₄ O	156.10111
Asparagine	N	C ₄ H ₆ N ₂ O ₂	114.042927
Aspartic Acid	D	C ₄ H ₅ NO ₃	115.026943
Carboxyamido	C	C ₅ H ₈ N ₂ O ₂ S	160.03065
Cysteine ¹			
Glutamic Acid	E	C ₅ H ₇ NO ₃	129.042593
Glutamine	Q	C ₃ H ₈ N ₂ O ₂	128.058577
Glycine	G	C ₂ H ₃ NO	57.021464
Histadine	H	C ₆ H ₇ N ₃ O	137.058912
Isoleucine	I	C ₆ H ₁₁ NO	113.084064
Leucine	L	C ₆ H ₁₁ NO	113.084064
Lysine	K	C ₆ H ₁₂ N ₂ O	128.094963
Methionine	M	C ₅ H ₉ NOS	131.040485
Oxidised Methionine	M*	C ₅ H ₉ NO ₂ S	147.035340
Phenylalanine	F	C ₉ H ₉ NO	147.068414
Proline	P	C ₅ H ₇ NO	97.052764
Serine	S	C ₃ H ₅ NO ₂	87.032028
Threonine	T	C ₄ H ₇ NO ₂	101.047678
Tryptophan	W	C ₁₁ H ₁₀ N ₂ O	186.079313
Tyrosine	Y	C ₉ H ₉ NO ₂	163.063328
Valine	V	C ₅ H ₉ NO	99.068414

¹All Cysteines are modified to the carboxyamino derivative during our production process.

As used herein “tryptic digest peptides” are peptides produced through treatment of the protein with the enzyme trypsin. Trypsin cleaves specifically at the carboxyl side of

lysine (Lys) and arginine (Arg) residues, so that the tryptic digest peptides generated should have a Lys or Arg as the C-terminal amino acid, unless the peptide fragment was obtained from the C-terminal of the protein. Similarly, the amino acid directly preceding the N-terminal amino acid of the tryptic digest peptides should also be a Lys or Arg, unless the peptide was obtained from the N-terminal of the protein. The mass of a tryptic digest peptide corresponds to the total mass of the constituent amino acid residues of the peptide with the addition of a water molecule (H₂O). As used herein, the "partial sequence" is an amino acid sequence within the tryptic digest peptide determined from the interpretation of the tandem mass spectrum of the peptide. As used herein, the "N-terminal mass" is the mass measured by mass spectrometry (having an error of measurement of approximately 100 parts-per-million or less) of the portion of the tryptic digest peptide extending from the start of the partial sequence to the N-terminus of the peptide. This is a neutral mass corresponding to the total mass of the constituent amino acid residues extending from the partial sequence to the N-terminus of the peptide. As used herein, the "C-terminal mass" is the mass measured by mass spectrometry (having an error of measurement of approximately 100 parts-per-million or less) of the portion of the tryptic digest peptide extending from the end of the partial sequence to the C-terminus of the peptide. This mass corresponds to the total mass of the constituent amino acid residues extending from the end of the partial sequence to the C-terminus of the peptide with the addition of a water molecular (H₂O), and a single proton (H⁺).

In Table IX, *supra*, the preferred and degenerate sets of probes are described using GCG Nucleotide Ambiguity Codes as employed in GCG SeqWebTM sequence analysis software (SeqWebTM version 1.1, part of Wisconsin Package Version 10, Genetics Computer Group, Inc.). These Nucleotide Ambiguity Codes have the following meaning:

GCG Code	Meaning
A	A
C	C
G	G
T	T
U	T
M	A or C
R	A or G

	W	A or T
	S	C or G
	Y	C or T
	K	G or T
5	V	A or C or G
	H	A or C or T
	D	A or G or T
	B	C or G or T
	X	G or A or T or C
10	N	G or A or T or
		C

GCG uses the letter codes for amino acid codes and nucleotide ambiguity proposed by IUPAC-IUB. These codes are compatible with the codes used by the EMBL, GenBank, and PIR databases. See IUPAC, Commission on Nomenclature of Organic Chemistry. A Guide to IUPAC Nomenclature of Organic Compounds (Recommendations 1993), Blackwell Scientific publications, 1993.

When a library is screened, clones with insert DNA encoding the API of interest or a fragment thereof will hybridize to one or more members of the corresponding set of degenerate oligonucleotide probes (or their complement). Hybridization of such oligonucleotide probes to genomic libraries is carried out using methods known in the art. For example, hybridization with one of the above-mentioned degenerate sets of oligonucleotide probes, or their complement (or with any member of such a set, or its complement) can be performed under highly stringent or moderately stringent conditions as defined above, or can be carried out in 2X SSC, 1.0% SDS at 50°C and washed using the washing conditions described *supra* for highly stringent or moderately stringent hybridization.

In yet another aspect of the invention, clones containing nucleotide sequences encoding the entire API, a fragment of an API, an API-related polypeptide, or a fragment of an API-related polypeptide or any of the foregoing may also be obtained by screening expression libraries. For example, DNA from the relevant source is isolated and random fragments are prepared and ligated into an expression vector (e.g., a bacteriophage,

plasmid, phagemid or cosmid) such that the inserted sequence in the vector is capable of being expressed by the host cell into which the vector is then introduced. Various screening assays can then be used to select for the expressed API or API-related polypeptides. In one embodiment, the various anti-API antibodies of the invention can be used to identify the
5 desired clones using methods known in the art. See, for example, Harlow and Lane, 1988, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, Appendix IV. Colonies or plaques from the library are brought into contact with the antibodies to identify those clones that bind antibody.

In an embodiment, colonies or plaques containing DNA that encodes an API, API
10 fragment, API -related polypeptide or API -fusion protein can be detected using DYNA Beads according to Olsvick et al., 29th ICAAC, Houston, Tex. 1989, incorporated herein by reference. Anti-API antibodies are crosslinked to tosylated DYNA Beads M280, and these antibody-containing beads are then contacted with colonies or plaques expressing recombinant polypeptides. Colonies or plaques expressing an API, API fragment, API -
15 related polypeptide or API -fusion protein are identified as any of those that bind the beads.

Alternatively, the anti-API antibodies can be nonspecifically immobilized to a suitable support, such as silica or Celite® resin. This material is then used to adsorb to bacterial colonies expressing the API, API fragment, API -related polypeptide or API -
fusion protein ptide as described herein.

20 In another aspect, PCR amplification may be used to isolate from genomic DNA a substantially pure DNA (*i.e.*, a DNA substantially free of contaminating nucleic acids) encoding the entire API or a part thereof. Preferably such a DNA is at least 95% pure, more preferably at least 99% pure. Oligonucleotide sequences, degenerate or otherwise, that correspond to peptide sequences of APIs disclosed herein can be used as primers.

25 PCR can be carried out, *e.g.*, by use of a Perkin-Elmer Cetus thermal cycler and Taq polymerase (Gene Amp® or AmpliTaq DNA polymerase). One can choose to synthesize several different degenerate primers, for use in the PCR reactions. It is also possible to vary the stringency of hybridization conditions used in priming the PCR reactions, to allow for greater or lesser degrees of nucleotide sequence similarity between the degenerate primers
30 and the corresponding sequences in the DNA. After successful amplification of a segment of the sequence encoding an API, that segment may be molecularly cloned and sequenced, and utilized as a probe to isolate a complete genomic clone. This, in turn, will permit the determination of the gene's complete nucleotide sequence, the analysis of its expression,

and the production of its protein product for functional analysis, as described *infra*.

The gene encoding an API can also be identified by mRNA selection by nucleic acid hybridization followed by *in vitro* translation. In this procedure, fragments are used to isolate complementary mRNAs by hybridization. Such DNA fragments may represent available, purified DNA encoding an API of another species (*e.g.*, mouse, human). Immunoprecipitation analysis or functional assays (*e.g.*, aggregation ability *in vitro*; binding to receptor) of the *in vitro* translation products of the isolated products of the isolated mRNAs identifies the mRNA and, therefore, the complementary DNA fragments that contain the desired sequences. In addition, specific mRNAs may be selected by adsorption of polysomes isolated from cells to immobilized antibodies that specifically recognize an API. A radiolabelled cDNA encoding an API can be synthesized using the selected mRNA (from the adsorbed polysomes) as a template. The radiolabelled mRNA or cDNA may then be used as a probe to identify the DNA fragments encoding an API from among other genomic DNA fragments.

Alternatives to isolating genomic DNA encoding an API include, but are not limited to, chemically synthesizing the gene sequence itself from a known sequence or making cDNA to the mRNA which encodes the API. For example, RNA for cDNA cloning of the gene encoding an API can be isolated from cells which express the API. Those skilled in the art will understand from the present description that other methods may be used and are within the scope of the invention.

Any suitable eukaryotic cell can serve as the nucleic acid source for the molecular cloning of the gene encoding an API. The nucleic acid sequences encoding the API can be isolated from vertebrate, mammalian, primate, human, porcine, bovine, feline, avian, equine, canine or murine sources. The DNA may be obtained by standard procedures known in the art from cloned DNA (*e.g.*, a DNA "library"), by chemical synthesis, by cDNA cloning, or by the cloning of genomic DNA, or fragments thereof, purified from the desired cell. (See, *e.g.*, Sambrook et al., 1989, *Molecular Cloning, A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; Glover, D.M. (ed.), 1985, *DNA Cloning: A Practical Approach*, MRL Press, Ltd., Oxford, U.K. Vol. I, II.). Clones derived from genomic DNA may contain regulatory and intron DNA regions in addition to coding regions; clones derived from cDNA will contain only exon sequences.

The identified and isolated gene or cDNA can then be inserted into any suitable

cloning vector. A large number of vector-host systems known in the art may be used. As those skilled in the art will appreciate, the vector system chosen should be compatible with the host cell used. Such vectors include, but are not limited to, bacteriophages such as lambda derivatives, plasmids such as PBR322 or pUC plasmid derivatives or the Bluescript
5 vector (Stratagene) or modified viruses such as adenoviruses, adeno-associated viruses or retroviruses. The insertion into a cloning vector can be accomplished, for example, by ligating the DNA fragment into a cloning vector which has complementary cohesive termini. However, if the complementary restriction sites used to fragment the DNA are not present in the cloning vector, the ends of the DNA molecules may be enzymatically
10 modified. Alternatively, any site desired may be produced by ligating nucleotide sequences (linkers) onto the DNA termini; these ligated linkers may comprise specific chemically synthesized oligonucleotides encoding restriction endonuclease recognition sequences. In an alternative method, the cleaved vector and the gene encoding an API may be modified by homopolymeric tailing. Recombinant molecules can be introduced into host cells via
15 transformation, transfection, infection, electroporation, etc., so that many copies of the gene sequence are generated.

In specific embodiments, transformation of host cells with recombinant DNA molecules that incorporate the isolated gene encoding the API, cDNA, or synthesized DNA sequence enables generation of multiple copies of the gene. Thus, the gene may be
20 obtained in large quantities by growing transformants, isolating the recombinant DNA molecules from the transformants and, when necessary, retrieving the inserted gene from the isolated recombinant DNA.

The nucleotide sequences of the present invention include nucleotide sequences encoding amino acid sequences with substantially the same amino acid sequences as native
25 APIs, nucleotide sequences encoding amino acid sequences with functionally equivalent amino acids, nucleotide sequences encoding APIs, API fragments, API -related polypeptides or fragments of API-related polypeptides.

In a specific embodiment, an isolated nucleic acid molecule encoding an API-related polypeptide can be created by introducing one or more nucleotide substitutions,
30 additions or deletions into the nucleotide sequence of an API, such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein. Standard techniques known to those of skill in the art can be used to introduce mutations, including, for example, site-directed mutagenesis and PCR-mediated mutagenesis. Preferably,

conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a side chain with a similar charge. Families of amino acid residues having side chains with similar charges have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Alternatively, mutations can be introduced randomly along all or part of the coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for biological activity to identify mutants that retain activity. Following mutagenesis, the encoded protein can be expressed and the activity of the protein can be determined.

5.9 Expression of DNA Encoding APIs

The nucleotide sequence coding for an API, API fragment, API -related polypeptide or API -fusion protein, can be inserted into an appropriate expression vector, *i.e.*, a vector which contains the necessary elements for the transcription and translation of the inserted protein-coding sequence. The necessary transcriptional and translational signals can also be supplied by the native gene encoding the API, or its flanking regions, or the native gene encoding the API-related polypeptide or its flanking regions. A variety of host-vector systems may be utilized in the present invention to express the protein-coding sequence. These include but are not limited to mammalian cell systems infected with virus (*e.g.*, vaccinia virus, adenovirus, etc.); insect cell systems infected with virus (*e.g.*, baculovirus); microorganisms such as yeast containing yeast vectors; or bacteria transformed with bacteriophage, DNA, plasmid DNA, or cosmid DNA. The expression elements of vectors vary in their strengths and specificities. Depending on the host-vector system utilized, any one of a number of suitable transcription and translation elements may be used. In specific embodiments, a nucleotide sequence encoding a human gene (or a nucleotide sequence encoding a functionally active portion of a human API) is expressed. In yet another embodiment, a fragment of an API comprising a domain of the API is expressed.

Any of the methods previously described for the insertion of DNA fragments into a

vector may be used to construct expression vectors containing a chimeric gene consisting of appropriate transcriptional and translational control signals and the protein coding sequences. These methods may include *in vitro* recombinant DNA and synthetic techniques and *in vivo* recombinants (genetic recombination). Expression of nucleic acid sequence encoding an API, API fragment, API -related polypeptide or API -fusion protein may be regulated by a second nucleic acid sequence so that the API, API fragment, API -related polypeptide or API -fusion protein is expressed in a host transformed with the recombinant DNA molecule. For example, expression of an API may be controlled by any promoter or enhancer element known in the art. Promoters which may be used to control the expression of the gene encoding an API, API fragment, API -related polypeptide or API -fusion protein include, but are not limited to, the SV40 early promoter region (Bernoist and Chambon, 1981, Nature 290:304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, et al., 1980, Cell 22:787-797), the herpes thymidine kinase promoter (Wagner et al., 1981, Proc. Natl. Acad. Sci. U.S.A. 78:1441-1445), the regulatory sequences of the metallothionein gene (Brinster et al., 1982, Nature 296:39-42), the tetracycline (Tet) promoter (Gossen et al., 1995, Proc. Nat. Acad. Sci. USA 89:5547-5551); prokaryotic expression vectors such as the β -lactamase promoter (Villa-Kamaroff, et al., 1978, Proc. Natl. Acad. Sci. U.S.A. 75:3727-3731), or the tac promoter (DeBoer, et al., 1983, Proc. Natl. Acad. Sci. U.S.A. 80:21-25; see also "Useful proteins from recombinant bacteria" in Scientific American, 1980, 242:74-94); plant expression vectors comprising the nopaline synthetase promoter region (Herrera-Estrella et al., Nature 303:209-213) or the cauliflower mosaic virus 35S RNA promoter (Gardner, et al., 1981, Nucl. Acids Res. 9:2871), and the promoter of the photosynthetic enzyme ribulose biphosphate carboxylase (Herrera-Estrella et al., 1984, Nature 310:115-120); promoter elements from yeast or other fungi such as the Gal 4 promoter, the ADC (alcohol dehydrogenase) promoter, PGK (phosphoglycerol kinase) promoter, alkaline phosphatase promoter, and the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: elastase I gene control region which is active in pancreatic acinar cells (Swift et al., 1984, Cell 38:639-646; Ornitz et al., 1986, Cold Spring Harbor Symp. Quant. Biol. 50:399-409; MacDonald, 1987, Hepatology 7:425-515); insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, Nature 315:115-122), immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., 1984, Cell 38:647-658; Adames et al., 1985, Nature 318:533-538; Alexander et al.,

1987, Mol. Cell. Biol. 7:1436-1444), mouse mammary tumor virus control region which is active in testicular, breast, lymphoid and mast cells (Leder et al., 1986, Cell 45:485-495), un gene control region which is active in liver (Pinkert et al., 1987, Genes and Devel. 1:268-276), alpha-fetoprotein gene control region which is active in liver (Krumlauf et al., 1985, Mol. Cell. Biol. 5:1639-1648; Hammer et al., 1987, Science 235:53-58; alpha 1-antitrypsin gene control region which is active in the liver (Kelsey et al., 1987, Genes and Devel. 1:161-171), beta-globin gene control region which is active in myeloid cells (Mogam et al., 1985, Nature 315:338-340; Kollias et al., 1986, Cell 46:89-94; myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead et al., 1987, Cell 48:703-712); myosin light chain-2 gene control region which is active in skeletal muscle (Sani, 1985, Nature 314:283-286); neuronal-specific enolase (NSE) which is active in neuronal cells (Morelli et al., 1999, Gen. Virol. 80:571-83); brain-derived neurotrophic factor (BDNF) gene control region which is active in neuronal cells (Tabuchi et al., 1998, Biochem. Biophysic. Res. Com. 253:818-823); glial fibrillary acidic protein (GFAP) promoter which is active in astrocytes (Gomes et al., 1999, Braz J Med Biol Res 32(5):619-631; Morelli et al., 1999, Gen. Virol. 80:571-83) and gonadotropic releasing hormone gene control region which is active in the hypothalamus (Mason et al., 1986, Science 234:1372-1378).

In a specific embodiment, a vector is used that comprises a promoter operably linked to an API-encoding nucleic acid, one or more origins of replication, and, optionally, one or more selectable markers (*e.g.*, an antibiotic resistance gene).

In a specific embodiment, an expression construct is made by subcloning an API or an API-related polypeptide coding sequence into the EcoRI restriction site of each of the three pGEX vectors (Glutathione S-Transferase expression vectors; Smith and Johnson, 1988, Gene 7:31-40). This allows for the expression of the API product or API-related polypeptide from the subclone in the correct reading frame.

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the API coding sequence or API-related polypeptide coding sequence may be ligated to an adenovirus transcription/translation control complex, *e.g.*, the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (*e.g.*, region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the

antibody molecule in infected hosts (*e.g.*, see Logan & Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:355-359). Specific initiation signals may also be required for efficient translation of inserted antibody coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading
5 frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bittner et al., 1987, Methods in Enzymol. 153:51-544).

10 Expression vectors containing inserts of a gene encoding an API or an API-related polypeptide can be identified, for example, by three general approaches: (a) nucleic acid hybridization, (b) presence or absence of "marker" gene functions, and (c) expression of inserted sequences. In the first approach, the presence of a gene encoding an API inserted in an expression vector can be detected by nucleic acid hybridization using probes
15 comprising sequences that are homologous to an inserted gene encoding an API. In the second approach, the recombinant vector/host system can be identified and selected based upon the presence or absence of certain "marker" gene functions (*e.g.*, thymidine kinase activity, resistance to antibiotics, transformation phenotype, occlusion body formation in baculovirus, etc.) caused by the insertion of a gene encoding an API in the vector. For
20 example, if the gene encoding the API is inserted within the marker gene sequence of the vector, recombinants containing the gene encoding the API insert can be identified by the absence of the marker gene function. In the third approach, recombinant expression vectors can be identified by assaying the gene product (*i.e.*, API) expressed by the recombinant. Such assays can be based, for example, on the physical or functional properties of the API
25 in *in vitro* assay systems, *e.g.*, binding with anti-API antibody.

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Expression from certain promoters can be elevated in the presence of certain inducers; thus, expression of the genetically engineered API or API-related polypeptide
30 may be controlled. Furthermore, different host cells have characteristic and specific mechanisms for the translational and post-translational processing and modification (*e.g.*, glycosylation, phosphorylation of proteins). Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed.

For example, expression in a bacterial system will produce an unglycosylated product and expression in yeast will produce a glycosylated product. Eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERO, BHK, Hela, COS, MDCK, HEK 293, 3T3, WI38, and in particular, neuronal cell lines such as, for example, SK-N-AS, SK-N-FI, SK-N-DZ human neuroblastomas (Sugimoto et al., 1984, J. Natl. Cancer Inst. 73: 51-57), SK-N-SH human neuroblastoma (Biochim. Biophys. Acta, 1982, 704: 450-460), Daoy human cerebellar medulloblastoma (He et al., 1992, Cancer Res. 52: 1144-1148) DBTRG-05MG glioblastoma cells (Kruse et al., 1992, *In vitro* Cell. Dev. Biol. 28A: 609-614), IMR-32 human neuroblastoma (Cancer Res., 1970, 30: 2110-2118), 1321N1 human astrocytoma (Proc. Natl Acad. Sci. USA, 1977, 74: 4816), MOG-G-CCM human astrocytoma (Br. J. Cancer, 1984, 49: 269), U87MG human glioblastoma-astrocytoma (Acta Pathol. Microbiol. Scand., 1968, 74: 465-486), A172 human glioblastoma (Olopade et al., 1992, Cancer Res. 52: 2523-2529), C6 rat glioma cells (Benda et al., 1968, Science 161: 370-371), Neuro-2a mouse neuroblastoma (Proc. Natl. Acad. Sci. USA, 1970, 65: 129-136), NB41A3 mouse neuroblastoma (Proc. Natl. Acad. Sci. USA, 1962, 48: 1184-1190), SCP sheep choroid plexus (Bolin et al., 1994, J. Virol. Methods 48: 211-221), G355-5, PG-4 Cat normal astrocyte (Haapala et al., 1985, J. Virol. 53: 827-833), Mpf ferret brain (Trowbridge et al., 1982, *In vitro* 18: 952-960), and normal cell lines such as, for example, CTX TNA2 rat normal cortex brain (Radany et al., 1992, Proc. Natl. Acad. Sci. USA 89: 6467-6471) such as, for example, CRL7030 and Hs578Bst. Furthermore, different vector/host expression systems may effect processing reactions to different extents.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the differentially expressed or pathway gene protein may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (*e.g.*, promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched medium, and then are switched to a selective medium. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be

cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the differentially expressed or pathway gene protein. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the differentially expressed or pathway gene protein.

5 A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy, et al., 1980, Cell 22:817) genes can be employed in tk-, hgp^rt- or ap^rt- cells, respectively. Also, antimetabolite resistance
10 can be used as the basis of selection for dhfr, which confers resistance to methotrexate (Wigler, et al., 1980, Natl. Acad. Sci. USA 77:3567; O'Hare, et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., 1981, J. Mol. Biol. 150:1); and hygro,
15 which confers resistance to hygromycin (Santerre, et al., 1984, Gene 30:147) genes.

In other embodiments, the API, API fragment or API-related polypeptide may be expressed as a fusion, or chimeric protein product (comprising the protein, fragment, analog, or derivative joined via a peptide bond to a heterologous protein sequence). For example, the polypeptides of the present invention may be fused with the constant domain
20 of immunoglobulins (IgA, IgE, IgG, IgM), or portions thereof (CH1, CH2, CH3, or any combination thereof and portions thereof) resulting in chimeric polypeptides. Such fusion proteins may facilitate purification, increase half-life *in vivo*, and enhance the delivery of an antigen across an epithelial barrier to the immune system. An increase in the half-life *in vivo* and facilitated purification has been shown for chimeric proteins consisting of the first
25 two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. See, e.g., EP 394,827; Traunecker et al., Nature, 331:84-86 (1988). Enhanced delivery of an antigen across the epithelial barrier to the immune system has been demonstrated for antigens (e.g., insulin) conjugated to an FcRn binding partner such as IgG or Fc fragments (see, e.g., PCT
30 publications WO 96/22024 and WO 99/04813).

Nucleic acids encoding an API, a fragment of an API, an API-related polypeptide, or a fragment of an API-related polypeptide can be fused to an epitope tag (e.g., the hemagglutinin ("HA") tag or flag tag) to aid in detection and purification of the expressed

polypeptide. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht et al., 1991, Proc. Natl. Acad. Sci. USA 88:8972-897).

An API fusion protein can be made by ligating the appropriate nucleic acid sequences encoding the desired amino acid sequences to each other by methods known in the art, in the proper coding frame, and expressing the chimeric product by methods commonly known in the art. Alternatively, an API fusion protein may be made by protein synthetic techniques, *e.g.*, by use of a peptide synthesizer.

Both cDNA and genomic sequences can be cloned and expressed.

5.10 Domain Structure of APIs

Domains of some of the APIs provided by the present invention are known in the art and have been described in the scientific literature. Moreover, domains of an API can be identified using techniques known to those of skill in the art. For example, one or more domains of an API can be identified by using one or more of the following programs: ProDom, TMPred, and SAPS. ProDom compares the amino acid sequence of a polypeptide to a database of compiled domains (see, *e.g.*, <http://www.toulouse.inra.fr/prodom.html>; Corpet F., Gouzy J. & Kahn D., 1999, Nucleic Acids Res., 27:263-267). TMPred predicts membrane-spanning regions of a polypeptide and their orientation. This program uses an algorithm that is based on the statistical analysis of TMbase, a database of naturally occurring transmembrane proteins (see, *e.g.*, http://www.ch.embnet.org/software/TMPRED_form.html; Hofmann & Stoffel. (1993) "TMbase - A database of membrane spanning proteins segments." Biol. Chem. Hoppe-Seyler 347,166). The SAPS program analyzes polypeptides for statistically significant features like charge-clusters, repeats, hydrophobic regions, compositional domains (see, *e.g.*, Brendel et al., 1992, Proc. Natl. Acad. Sci. USA 89: 2002-2006). Thus, based on the present description, those skilled in the art can identify domains of an API having enzymatic or binding activity, and further can identify nucleotide sequences encoding such domains. These nucleotide sequences can then be used for recombinant expression of an API fragment that retains the enzymatic or binding activity of the API.

Based on the present description, those skilled in the art can identify domains of an API having enzymatic or binding activity, and further can identify nucleotide sequences encoding such domains. These nucleotide sequences can then be used for recombinant

expression of API fragments that retain the enzymatic or binding activity of the API.

In one embodiment, an API has an amino acid sequence sufficiently similar to an identified domain of a known polypeptide. As used herein, the term "functionally similar" refers to a first amino acid or nucleotide sequence which contains a sufficient number of
5 identical or equivalent (*e.g.*, with a similar side chain) amino acid residues or nucleotides to a second amino acid or nucleotide sequence such that the first and second amino acid or nucleotide sequences have or encode a common structural domain or common functional activity or both.

An API domain can be assessed for its function using techniques well known to
10 those of skill in the art. For example, a domain can be assessed for its kinase activity or for its ability to bind to DNA using techniques known to the skilled artisan. Kinase activity can be assessed, for example, by measuring the ability of a polypeptide to phosphorylate a substrate. DNA binding activity can be assessed, for example, by measuring the ability of a polypeptide to bind to a DNA binding element in a electromobility shift assay. In a
15 preferred embodiment, the function of a domain of an API is determined using an assay described in one or more of the references identified in Table IX, *infra*.

5.11 Production of Antibodies to APIs

According to the invention API, API fragment, API -related polypeptide or API -
20 fusion protein may be used as an immunogen to generate antibodies which immunospecifically bind such an immunogen. Such immunogens can be isolated by any convenient means, including the methods described above. Antibodies of the invention include, but are not limited to polyclonal, monoclonal, bispecific, humanized or chimeric antibodies, single chain antibodies, Fab fragments and F(ab') fragments, fragments
25 produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, *i.e.*, molecules that contain an antigen-binding site that specifically binds an antigen. The immunoglobulin molecules of the invention can be of any class (*e.g.*, IgG, IgE, IgM, IgD and IgA) or subclass of immunoglobulin molecule.
30

In one embodiment, antibodies that recognize gene products of genes encoding APIs may be prepared. For example, antibodies that recognize these APIs and/or their isoforms include antibodies recognizing API-1, API-3, API-4, API-6, API-7, API-10, API-15, API-

16, API-22, API-28, API-30, API-33, API-34, API-37, API-38, API-39, API-40, API-42, API-43, API-44, API-45, API-46, API-47, API-50, API-52, API-53, API-55, API-58, API-60, API-62, API-64, API-66, API-67, API-69, API-72, API-74, API-75, API-76, API-77, API-78, API-79, API-80, API-81, API-82, API-84, API-90, API-92, API-93, API-95, API-97, API-98, API-101, API-102, API-103, API-104, API-113, API-118, API-119, API-123, API-124, API-126, API-130, API-131, API-132, API-134, API-136, API-137, API-138, API-140, API-142, API-143, API-145, API-149, API-150, API-161, API-165, API-167, API-168, API-169, API-170, API-171, API-172, API-173, API-174, API-175, API-176, API-178, API-179, API-181, API-182, API-186, API-188, API-189, API-191, API-194, API-196, API-201, API-215, API-220, API-221, API-223, API-225, API-233, API-234, API-237, API-238, API-239, API-240, API-241, API-242, API-243, API-244, API-245, API-246, API-247, API-248, API-300, API-301, API-302, API-303, API-304, API-305, API-306, API-307, API-308, API-309, API-310, API-311, API-312, API-313, API-314, API-315, API-316, API-317, API-318, API-319, API-320, API-321, API-322, API-323, API-324, API-325, API-326, API-327, API-328, API-329, API-330, API-331, API-332, API-333, API-334, API-335, API-336, API-337, API-338, API-339, API-340, API-341, API-342, API-343, API-344, API-345, API-346, API-347, API-348, API-349, API-350, API-351, API-352, API-353, API-354, API-355, API-356, API-357, API-358, API-359, API-360, API-361, API-362, API-363, API-364, API-365, API-366, API-367 or API-368.

Certain antibodies are already known and can be purchased from commercial sources as shown in Table VII above. In another embodiment, methods known to those skilled in the art are used to produce antibodies that recognize an API, an API analog, an API-related polypeptide, or a derivative or fragment of any of the foregoing.

In one embodiment of the invention, antibodies to a specific domain of an API are produced. In a specific embodiment, hydrophilic fragments of an API are used as immunogens for antibody production.

In the production of antibodies, screening for the desired antibody can be accomplished by techniques known in the art, *e.g.* ELISA (enzyme-linked immunosorbent assay). For example, to select antibodies which recognize a specific domain of an API, one may assay generated hybridomas for a product which binds to an API fragment containing such domain. For selection of an antibody that specifically binds a first API homolog but which does not specifically bind to (or binds less avidly to) a second API homolog, one can select on the basis of positive binding to the first API homolog and a lack of binding to (or

reduced binding to) the second API homolog. Similarly, for selection of an antibody that specifically binds an API but which does not specifically bind to (or binds less avidly to) a different isoform of the same protein (such as a different glycoform having the same core peptide as the API), one can select on the basis of positive binding to the API and a lack of binding to (or reduced binding to) the different isoform (*e.g.*, a different glycoform). Thus, the present invention provides an antibody (particularly a monoclonal antibody) that binds with greater affinity (particularly at least 2-fold, more particularly at least 5-fold still more particularly at least 10-fold greater affinity) to an API than to a different isoform or isoforms (*e.g.*, glycoforms) of the API.

Polyclonal antibodies, which may be used in the methods of the invention, are heterogeneous populations of antibody molecules derived from the sera of immunized animals. Unfractionated immune serum can also be used. Various procedures known in the art may be used for the production of polyclonal antibodies to an API, API fragment, API-related polypeptide or API-fusion protein. In a particular embodiment, rabbit polyclonal antibodies to an epitope of an API, API fragment, API-related polypeptide or API-fusion protein can be obtained. For example, for the production of polyclonal or monoclonal antibodies, various host animals can be immunized by injection with the native or a synthetic (*e.g.*, recombinant) version of an API, API fragment, API-related polypeptide or API-fusion protein, including but not limited to rabbits, mice, rats, etc. Isolated APIs suitable for such immunization may be obtained by the use of discovery techniques, such as the preferred technology described herein. If the API is purified by gel electrophoresis, the API can be used for immunization with or without prior extraction from the polyacrylamide gel. Various adjuvants may be used to enhance the immunological response, depending on the host species, including, but not limited to, complete or incomplete Freund's adjuvant, a mineral gel such as aluminum hydroxide, surface active substance such as lysolecithin, pluronic polyol, a polyanion, a peptide, an oil emulsion, keyhole limpet hemocyanin, dinitrophenol, and an adjuvant such as BCG (*bacille Calmette-Guerin*) or *corynebacterium parvum*. Additional adjuvants are also well known in the art.

For preparation of monoclonal antibodies (mAbs) directed toward an API, API fragment, API-related polypeptide or API-fusion protein, any technique which provides for the production of antibody molecules by continuous cell lines in culture may be used. For example, the hybridoma technique originally developed by Kohler and Milstein (1975, *Nature* 256:495-497), as well as the trioma technique, the human B-cell hybridoma

technique (Kozbor et al., 1983, Immunology Today 4:72), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al., 1985, in Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof.

- 5 The hybridoma producing the mAbs of the invention may be cultivated *in vitro* or *in vivo*. In an additional embodiment of the invention, monoclonal antibodies can be produced in germ-free animals utilizing known technology (PCT/US90/02545, incorporated herein by reference).

The monoclonal antibodies include but are not limited to human monoclonal
10 antibodies and chimeric monoclonal antibodies (*e.g.*, human-mouse chimeras). Humanized antibodies are antibody molecules from non-human species having one or more complementarily determining regions (CDRs) from the non-human species and a framework region from a human immunoglobulin molecule. (See, *e.g.*, Queen, U.S. Patent No. 5,585,089, which is incorporated herein by reference in its entirety.)

15 Chimeric and humanized monoclonal antibodies can be produced by recombinant DNA techniques known in the art, for example using methods described in PCT Publication No. WO 87/02671; European Patent Application 184,187; European Patent Application 171,496; European Patent Application 173,494; PCT Publication No. WO 86/01533; U.S. Patent No. 4,816,567; European Patent Application 125,023; Better et al., 1988, Science
20 240:1041-1043; Liu et al., 1987, Proc. Natl. Acad. Sci. USA., 84:3439-3443; Liu et al., 1987, J. Immunol. 139:3521-3526; Sun et al., 1987, Proc. Natl. Acad. Sci. USA., 84:214-218; Nishimura et al., 1987, Canc. Res. 47:999-1005; Wood et al., 1985, Nature 314:446-449; and Shaw et al., 1988, J. Natl. Cancer Inst. 80:1553-1559; Morrison, 1985, Science 229:1202-1207; Oi et al., 1986, Bio/Techniques 4:214; U.S. Patent 5,225,539; Jones et al.,
25 1986, Nature 321:552-525; Verhoeyan et al. (1988) Science 239:1534; and Beidler et al., 1988, J. Immunol. 141:4053-4060.

Completely human antibodies (antibodies derived solely from human antigenic material) are particularly desirable for therapeutic treatment of human subjects. Such antibodies can be produced using transgenic mice which are incapable of expressing
30 endogenous immunoglobulin heavy and light chains genes, but which can express human heavy and light chain genes. The transgenic mice are immunized in the normal fashion with selected antigens, *e.g.*, all or a portion of an API of the invention. Monoclonal antibodies directed against the antigen can be obtained using conventional hybridoma

technology. The human immunoglobulin transgenes harbored by the transgenic mice rearrange during B cell differentiation, and subsequently undergo class switching and somatic mutation. Thus, using such a technique, it is possible to produce therapeutically useful IgG, IgA, IgM and IgE antibodies. For an overview of this technology for producing human antibodies, see Lonberg and Huszar (1995, *Int. Rev. Immunol.* 13:65-93). For a detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, see, *e.g.*, U.S. Patent 5,625,126; U.S. Patent 5,633,425; U.S. Patent 5,569,825; U.S. Patent 5,661,016; and U.S. Patent 5,545,806. In addition, companies such as Abgenix, Inc. (Freemont, CA) and Genpharm (San Jose, CA) can be engaged to provide human antibodies directed against a selected antigen using technology similar to that described above.

Completely human antibodies which recognize a selected epitope can be generated using a technique referred to as "guided selection." In this approach a selected non-human monoclonal antibody, *e.g.*, a mouse antibody, is used to guide the selection of a completely human antibody recognizing the same epitope. (Jespers et al. (1994) *Biotechnology* 12:899-903).

The antibodies of the present invention can also be generated using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of phage particles which carry the polynucleotide sequences encoding them. In a particular, such phage can be utilized to display antigen-binding domains expressed from a repertoire or combinatorial antibody library (*e.g.*, human or murine). Phage expressing an antigen binding domain that binds the antigen of interest can be selected or identified with antigen, *e.g.*, using labeled antigen or antigen bound or captured to a solid surface or bead. Phage used in these methods are typically filamentous phage including fd and M13 binding domains expressed from phage with Fab, Fv or disulfide stabilized Fv antibody domains recombinantly fused to either the phage gene III or gene VIII protein. Phage display methods that can be used to make the antibodies of the present invention include those disclosed in Brinkman et al., *J. Immunol. Methods* 182:41-50 (1995); Ames et al., *J. Immunol. Methods* 184:177-186 (1995); Kettleborough et al., *Eur. J. Immunol.* 24:952-958 (1994); Persic et al., *Gene* 187 9-18 (1997); Burton et al., *Advances in Immunology* 57:191-280 (1994); PCT Application No. PCT/GB91/01134; PCT Publications WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and U.S. Patent Nos. 5,698,426; 5,223,409;

5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780,225; 5,658,727; 5,733,743 and 5,969,108; each of which is incorporated herein by reference in its entirety.

As described in the above references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies, including human antibodies, or any other desired antigen binding fragment, and expressed in any desired host, including mammalian cells, insect cells, plant cells, yeast, and bacteria, *e.g.*, as described in detail below. For example, techniques to recombinantly produce Fab, Fab' and F(ab')₂ fragments can also be employed using methods known in the art such as those disclosed in PCT publication WO 92/22324; Mullinax et al., *BioTechniques* 12(6):864-869 (1992); and Sawai et al., *AJRI* 34:26-34 (1995); and Better et al., *Science* 240:1041-1043 (1988) (said references incorporated by reference in their entireties).

Examples of suitable techniques which can be used to produce single-chain Fvs and antibodies against APIs of the present invention include those described in U.S. Patents 4,946,778 and 5,258,498; Huston et al., *Methods in Enzymology* 203:46-88 (1991); Shu et al., *PNAS* 90:7995-7999 (1993); and Skerra et al., *Science* 240:1038-1040 (1988).

The invention further provides for the use of bispecific antibodies, which can be made by methods known in the art. Traditional production of full-length bispecific antibodies is based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities (Milstein et al., 1983, *Nature* 305:537-539). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. Purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker et al., 1991, *EMBO J.* 10:3655-3659.

According to a different and more preferred approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH₂, and CH₃ regions. It is preferred to have the first heavy-chain constant region (CH₁) containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the

immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance.

In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in WO 94/04690 published March 3, 1994. For further details for generating bispecific antibodies see, for example, Suresh et al., *Methods in Enzymology*, 1986, 121:210.

The invention provides functionally active fragments, derivatives or analogs of the anti-API immunoglobulin molecules. Functionally active means that the fragment, derivative or analog is able to elicit anti-anti-idiotypic antibodies (*i.e.*, tertiary antibodies) that recognize the same antigen that is recognized by the antibody from which the fragment, derivative or analog is derived. Specifically, in a preferred embodiment the antigenicity of the idiotype of the immunoglobulin molecule may be enhanced by deletion of framework and CDR sequences that are C-terminal to the CDR sequence that specifically recognizes the antigen. To determine which CDR sequences bind the antigen, synthetic peptides containing the CDR sequences can be used in binding assays with the antigen by any suitable binding assay known in the art.

The present invention provides antibody fragments such as, but not limited to, F(ab')₂ fragments and Fab fragments. Antibody fragments which recognize specific epitopes may be generated by known techniques. F(ab')₂ fragments consist of the variable region, the light chain constant region and the CH1 domain of the heavy chain and are generated by pepsin digestion of the antibody molecule. Fab fragments are generated by reducing the disulfide bridges of the F(ab')₂ fragments. The invention also provides heavy chain and light chain dimers of the antibodies of the invention, or any minimal fragment

thereof such as Fvs or single chain antibodies (SCAs) (*e.g.*, as described in U.S. Patent 4,946,778; Bird, 1988, Science 242:423-42; Huston et al., 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; and Ward et al., 1989, Nature 334:544-54), or any other molecule with the same specificity as the antibody of the invention. Single chain antibodies are formed by
5 linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide. Techniques for the assembly of functional Fv fragments in *E. coli* may be used (Skerra et al., 1988, Science 242:1038-1041).

In other embodiments, the invention provides fusion proteins of the immunoglobulins of the invention (or functionally active fragments thereof), for example in
10 which the immunoglobulin is fused via a covalent bond (*e.g.*, a peptide bond), at either the N-terminus or the C-terminus to an amino acid sequence of another protein (or portion thereof, preferably at least 10, 20 or 50 amino acid portion of the protein) that is not the immunoglobulin. Preferably the immunoglobulin, or fragment thereof, is covalently linked to the other protein at the N-terminus of the constant domain. As stated above, such fusion
15 proteins may facilitate purification, increase half-life *in vivo*, and enhance the delivery of an antigen across an epithelial barrier to the immune system.

The immunoglobulins of the invention include analogs and derivatives that are either modified, *i.e.*, by the covalent attachment of any type of molecule as long as such covalent attachment that does not impair immunospecific binding. For example, but not by
20 way of limitation, the derivatives and analogs of the immunoglobulins include those that have been further modified, *e.g.*, by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to specific chemical
25 cleavage, acetylation, formylation, etc. Additionally, the analog or derivative may contain one or more non-classical or unnatural amino acids.

The foregoing antibodies can be used in methods known in the art relating to the localization and activity of the APIs of the invention, *e.g.*, for imaging these proteins, measuring levels thereof in appropriate physiological samples, in diagnostic methods, etc.

5.12 Expression Of Antibodies

The antibodies of the invention can be produced by any suitable method known in the art for the synthesis of antibodies, in particular, by chemical synthesis or by

recombinant expression, and are preferably produced by recombinant expression techniques.

Recombinant expression of antibodies, or fragments, derivatives or analogs thereof, requires construction of a nucleic acid that encodes the antibody. If the nucleotide
5 sequence of the antibody is known, a nucleic acid encoding the antibody may be assembled from chemically synthesized oligonucleotides (*e.g.*, as described in Kutmeier et al., 1994, BioTechniques 17:242), which, briefly, involves the synthesis of overlapping oligonucleotides containing portions of the sequence encoding antibody, annealing and ligation of those oligonucleotides, and then amplification of the ligated oligonucleotides by
10 PCR.

Alternatively, the nucleic acid encoding the antibody may be obtained by cloning the antibody. If a clone containing the nucleic acid encoding the particular antibody is not available, but the sequence of the antibody molecule is known, a nucleic acid encoding the antibody may be obtained from a suitable source (*e.g.*, an antibody cDNA library, or cDNA
15 library generated from any tissue or cells expressing the antibody) by PCR amplification using synthetic primers hybridizable to the 3' and 5' ends of the sequence or by cloning using an oligonucleotide probe specific for the particular gene sequence.

If an antibody molecule that specifically recognizes a particular antigen is not available (or a source for a cDNA library for cloning a nucleic acid encoding such an
20 antibody), antibodies specific for a particular antigen may be generated by any method known in the art, for example, by immunizing an animal, such as a rabbit, to generate polyclonal antibodies or, more preferably, by generating monoclonal antibodies.

Alternatively, a clone encoding at least the Fab portion of the antibody may be obtained by screening Fab expression libraries (*e.g.*, as described in Huse et al., 1989, Science
25 246:1275-1281) for clones of Fab fragments that bind the specific antigen or by screening antibody libraries (See, *e.g.*, Clackson et al., 1991, Nature 352:624; Hane et al., 1997 Proc. Natl. Acad. Sci. USA 94:4937).

Once a nucleic acid encoding at least the variable domain of the antibody molecule is obtained, it may be introduced into a vector containing the nucleotide sequence encoding
30 the constant region of the antibody molecule (see, *e.g.*, WO 86/05807; WO 89/01036; and U.S. 5,122,464). Vectors containing the complete light or heavy chain for co-expression with the nucleic acid to allow the expression of a complete antibody molecule are also available. Then, the nucleic acid encoding the antibody can be used to introduce the

nucleotide substitution(s) or deletion(s) necessary to substitute (or delete) the one or more variable region cysteine residues participating in an intrachain disulfide bond with an amino acid residue that does not contain a sulfhydryl group. Such modifications can be carried out by any method known in the art for the introduction of specific mutations or deletions in a nucleotide sequence, for example, but not limited to, chemical mutagenesis, *in vitro* site directed mutagenesis (Hutchinson et al., 1978, J. Biol. Chem. 253:6551), PCR based methods, etc.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., 1984, Proc. Natl. Acad. Sci. 81:851-855; Neuberger *et al.*, 1984, Nature 312:604-608; Takeda *et al.*, 1985, Nature 314:452-454) by splicing genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. As described *supra*, a chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human antibody constant region, *e.g.*, humanized antibodies.

Once a nucleic acid encoding an antibody molecule of the invention has been obtained, the vector for the production of the antibody molecule may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing the protein of the invention by expressing nucleic acid containing the antibody molecule sequences are described herein. Methods which are well known to those skilled in the art, can be used to construct expression vectors containing an antibody molecule coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. See, for example, the techniques described in Sambrook et al. (1990, Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY) and Ausubel et al. (eds., 1998, Current Protocols in Molecular Biology, John Wiley & Sons, NY).

The expression vector is transferred to a host cell by conventional techniques and the transfected cells are then cultured by conventional techniques to produce an antibody of the invention.

The host cells used to express a recombinant antibody of the invention may be either bacterial cells such as *Escherichia coli*, or, preferably, eukaryotic cells, especially for the expression of whole recombinant antibody molecule. In particular, mammalian cells such

as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus is an effective expression system for antibodies (Foecking *et al.*, 1986, *Gene* 45:101; Cockett *et al.*, 1990, *Bio/Technology* 8:2).

5 A variety of host-expression vector systems may be utilized to express an antibody molecule of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express the antibody molecule of the invention in situ. These include but are not
10 limited to microorganisms such as bacteria (*e.g.*, *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing antibody coding sequences; yeast (*e.g.*, *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing antibody coding sequences; insect cell systems infected with recombinant virus expression vectors (*e.g.*, baculovirus)
15 containing the antibody coding sequences; plant cell systems infected with recombinant virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (*e.g.*, Ti plasmid) containing antibody coding sequences; or mammalian cell systems (*e.g.*, COS, CHO, BHK, HEK 293, 3T3 cells) harboring recombinant expression constructs containing promoters
20 derived from the genome of mammalian cells (*e.g.*, metallothionein promoter) or from mammalian viruses (*e.g.*, the adenovirus late promoter; the vaccinia virus 7.5K promoter).

 In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the antibody molecule being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of
25 pharmaceutical compositions comprising an antibody molecule, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther *et al.*, 1983, *EMBO J.* 2:1791), in which the antibody coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a
30 fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, *Nucleic Acids Res.* 13:3101-3109; Van Heeke & Schuster, 1989, *J. Biol. Chem.* 24:5503-5509); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily

be purified from lysed cells by adsorption and binding to a matrix glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

5 In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The antibody coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). In mammalian host cells, a number of viral-based
10 expression systems (e.g., an adenovirus expression system) may be utilized.

As discussed above, a host cell strain may be chosen based on the present description which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the
15 function of the protein.

For long-term, high-yield production of recombinant antibodies, stable expression is preferred. For example, cells lines that stably express an antibody of interest can be produced by transfecting the cells with an expression vector comprising the nucleotide sequence of the antibody and the nucleotide sequence of a selectable (e.g., neomycin or
20 hygromycin), and selecting for expression of the selectable marker. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that interact directly or indirectly with the antibody molecule.

The expression levels of the antibody molecule can be increased by vector amplification (for a review, see Bebbington and Hentschel, The use of vectors based on
25 gene amplification for the expression of cloned genes in mammalian cells in DNA cloning, Vol.3. (Academic Press, New York, 1987)). When a marker in the vector system expressing antibody is amplifiable, increase in the level of inhibitor present in culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the antibody gene, production of the antibody will also increase (Crouse
30 et al., 1983, Mol. Cell. Biol. 3:257).

The host cell may be co-transfected with two expression vectors of the invention, the first vector encoding a heavy chain derived polypeptide and the second vector encoding a light chain derived polypeptide. The two vectors may contain identical selectable markers

which enable equal expression of heavy and light chain polypeptides. Alternatively, a single vector may be used which encodes both heavy and light chain polypeptides. In such situations, the light chain should be placed before the heavy chain to avoid an excess of toxic free heavy chain (Proudfoot, 1986, Nature 322:52; Kohler, 1980, Proc. Natl. Acad. Sci. USA 77:2197). The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.

Once the antibody molecule of the invention has been recombinantly expressed, it may be purified by any method known in the art for purification of an antibody molecule, for example, by chromatography (*e.g.*, ion exchange chromatography, affinity chromatography such as with protein A or specific antigen, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins.

Alternatively, any fusion protein may be readily purified by utilizing an antibody specific for the fusion protein being expressed. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht *et al.*, 1991, Proc. Natl. Acad. Sci. USA 88:8972-897). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the open reading frame of the gene is translationally fused to an amino-terminal tag consisting of six histidine residues. The tag serves as a matrix binding domain for the fusion protein. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni²⁺ nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

5.13 Conjugated Antibodies

In a preferred embodiment, anti-API antibodies or fragments thereof are conjugated to a diagnostic or a therapeutic molecule. The antibodies can be used, for example, for diagnosis or to determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, radioactive nuclides, positron emitting metals (for use in positron emission tomography), and nonradioactive paramagnetic metal ions. See generally U.S. Patent No. 4,741,900 for metal ions which can be conjugated to antibodies for use as diagnostics according to the present invention. Suitable enzymes include

horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; suitable prosthetic groups include streptavidin, avidin and biotin; suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride and phycoerythrin; suitable luminescent materials include luminol; suitable bioluminescent materials include luciferase, luciferin, and aequorin; and suitable radioactive nuclides include ^{125}I , ^{131}I , ^{111}In and ^{99}Tc .

Anti-API antibodies or fragments thereof can be conjugated to a therapeutic agent or pharmaceutical product to modify a given biological response. The therapeutic agent or drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor, α -interferon, β -interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator, a thrombotic agent or an anti-angiogenic agent, *e.g.*, angiostatin or endostatin; or, a biological response modifier such as a lymphokine, interleukin-1 (IL-1), interleukin-2 (IL-2), interleukin-6 (IL-6), granulocyte macrophage colony stimulating factor (GM-CSF), granulocyte colony stimulating factor (G-CSF), nerve growth factor (NGF) or other growth factor.

Techniques for conjugating such therapeutic moiety to antibodies are well known, see, *e.g.*, Arnon et al., "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld et al. (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom et al., "Antibodies For Drug Delivery", in *Controlled Drug Delivery* (2nd Ed.), Robinson et al. (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera et al. (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin et al. (eds.), pp. 303-16 (Academic Press 1985), and Thorpe et al., "The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates", *Immunol. Rev.*, 62:119-58 (1982). These references are incorporated herein in their entirety.

Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described in U.S. 4,676,980.

An antibody with or without a therapeutic moiety conjugated to it can be used as a therapeutic that is administered alone or in combination with cytotoxic factor(s) and/or cytokine(s).

5 5.14 Diagnosis of AD

In accordance with the present invention, suitable test samples, *e.g.*, of cerebrospinal fluid (CSF), serum, plasma or urine obtained from a subject suspected of having or known to have AD can be used for diagnosis. In one embodiment, a decreased abundance of one or more AFs or APIs (or any combination of them) in a test sample relative to a control
10 sample (from a subject or subjects free from AD) or a previously determined reference range indicates the presence of AD; AFs and APIs suitable for this purpose are identified in Tables I and IV, respectively, as described in detail above. In another embodiment of the invention, an increased abundance of one or more AFs or APIs (or any combination of them) in a test sample compared to a control sample or a previously determined reference
15 range indicates the presence of AD; AFs and APIs suitable for this purpose are identified in Tables II and V, respectively, as described in detail above. In another embodiment, the relative abundance of one or more AFs or APIs (or any combination of them) in a test sample compared to a control sample or a previously determined reference range indicates a subtype of AD (*e.g.*, familial or sporadic AD). In yet another embodiment, the relative
20 abundance of one or more AFs or APIs (or any combination of them) in a test sample relative to a control sample or a previously determined reference range indicates the degree or severity of AD. In any of the aforesaid methods, detection of one or more APIs described herein may optionally be combined with detection of one or more additional biomarkers for AD including, but not limited to apolipoprotein E (ApoE), amyloid β -
25 peptides (A β), tau and neural thread protein (NTP). Any suitable method in the art can be employed to measure the level of AFs and APIs, including but not limited to the Preferred Technology described herein, kinase assays, immunoassays to detect and/or visualize the APIs (*e.g.*, Western blot, immunoprecipitation followed by sodium dodecyl sulfate polyacrylamide gel electrophoresis, immunocytochemistry, etc.). In cases where an API
30 has a known function, an assay for that function may be used to measure API expression. In a further embodiment, a decreased abundance of mRNA encoding one or more APIs identified in Table IV (or any combination of them) in a test sample relative to a control sample or a previously determined reference range indicates the presence of AD. In yet a

further embodiment, an increased abundance of mRNA encoding one or more APIs identified in Table V (or any combination of them) in a test sample relative to a control sample or previously determined reference range indicates the presence of AD. Any suitable hybridization assay can be used to detect API expression by detecting and/or visualizing mRNA encoding the API (*e.g.*, Northern assays, dot blots, in situ hybridization, etc.).

In another embodiment of the invention, labeled antibodies, derivatives and analogs thereof, which specifically bind to an API can be used for diagnostic purposes, *e.g.*, to detect, diagnose, or monitor AD. Preferably, AD is detected in an animal, more preferably in a mammal and most preferably in a human.

5.15 Screening Assays

The invention provides methods for identifying active agents (*e.g.*, chemical compounds, proteins, or peptides) that bind to an API or have a stimulatory or inhibitory effect on the expression or activity of an API. The term “active agent” includes APIs, API fragments, API-related polypeptides, anti-API antibodies, fragments of anti-API antibodies and agents which modulate the expression of APIs for example, but without limitation, agonists or antagonists of APIs. The invention also provides methods of identifying candidate agents that bind to an API-related polypeptide or an API fusion protein or have a stimulatory or inhibitory effect on the expression or activity of an API-related polypeptide or an API fusion protein. Examples of active agents or candidate agents include, but are not limited to, nucleic acids (*e.g.*, DNA and RNA), carbohydrates, lipids, proteins, peptides, peptidomimetics, small molecules and other drugs. Candidate agents can be obtained using any of the numerous suitable approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the “one-bead one-compound” library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam, 1997, Anticancer Drug Des. 12:145; U.S. Patent No. 5,738,996; and U.S. Patent No. 5,807,683, each of which is incorporated herein in its entirety by reference).

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt et al., 1993, Proc. Natl. Acad. Sci. USA 90:6909; Erb et al., 1994,

Proc. Natl. Acad. Sci. USA 91:11422; Zuckermann et al., 1994, J. Med. Chem. 37:2678; Cho et al., 1993, Science 261:1303; Carrell et al., 1994, Angew. Chem. Int. Ed. Engl. 33:2059; Carrell et al., 1994, Angew. Chem. Int. Ed. Engl. 33:2061; and Gallop et al., 1994, J. Med. Chem. 37:1233, each of which is incorporated herein in its entirety by reference.

5 Libraries of compounds may be presented, *e.g.*, presented in solution (*e.g.*, Houghten, 1992, Bio/Techniques 13:412-421), or on beads (Lam, 1991, Nature 354:82-84), chips (Fodor, 1993, Nature 364:555-556), bacteria (U.S. Patent No. 5,223,409), spores (Patent Nos. 5,571,698; 5,403,484; and 5,223,409), plasmids (Cull et al., 1992, Proc. Natl. Acad. Sci. USA 89:1865-1869) or phage (Scott and Smith, 1990, Science 249:386-390; 10 Devlin, 1990, Science 249:404-406; Cwirla et al., 1990, Proc. Natl. Acad. Sci. USA 87:6378-6382; and Felici, 1991, J. Mol. Biol. 222:301-310).

In one embodiment, candidate agents that interact with APIs, API fragments, API - related polypeptides or API -fusion proteins are identified in a cell-based assay system. In accordance with this embodiment, cells expressing an API, API fragment, API-related 15 polypeptide, or API-fusion protein are contacted with a candidate agent or a control agent and the ability of the candidate agent to interact with the API is determined. If desired, this assay may be used to screen a plurality (*e.g.* a library) of candidate agents. The cell, for example, can be of prokaryotic origin (*e.g.*, *E. coli*) or eukaryotic origin (*e.g.*, yeast or mammalian). Further, the cells can express the an API, API fragment, API-related 20 polypeptide, or API-fusion protein endogenously or be genetically engineered to express the API, API fragment, API-related polypeptide, or API-fusion protein. In some embodiments, the API, fragment of the API, API-related polypeptide, a fragment of the API-related polypeptide, or an API fusion protein or the candidate agent is labeled, for example with a radioactive label (such as ³²P, ³⁵S or ¹²⁵I) or a fluorescent label (such as 25 fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, o-phthaldehyde or fluorescamine) to enable detection of an interaction between an API and a candidate agent. The ability of the candidate agent to interact directly or indirectly with an API, API fragment, API-related polypeptide, or API-fusion protein can be determined by methods known to those of skill in the art. For example, the interaction between a 30 candidate agent and an an API, API fragment, API-related polypeptide, or API-fusion protein can be determined by flow cytometry, a scintillation assay, immunoprecipitation or western blot analysis.

In another embodiment, agents that interact with (*i.e.*, bind to) an API, API

fragment, API-related polypeptide, or API-fusion protein are identified in a cell-free assay system. In accordance with this embodiment, a native or recombinant API or fragment thereof, or a native or recombinant API-related polypeptide or fragment thereof, or an API-fusion protein or fragment thereof, is contacted with a candidate agent or a control
5 compound and the ability of the candidate compound to interact with the API, API fragment, API-related polypeptide, or API-fusion protein is determined. If desired, this assay may be used to screen a plurality (*e.g.* a library) of candidate compounds. Preferably, the API, API fragment, API-related polypeptide, or API-fusion protein is first immobilized, by, for example, contacting the API, API fragment, API-related polypeptide, or API-fusion
10 protein with an immobilized antibody which specifically recognizes and binds it, or by contacting a purified preparation of the API, API fragment, API-related polypeptide, or API-fusion protein with a surface designed to bind proteins. The API, API fragment, API-related polypeptide, or API-fusion protein may be partially or completely purified (*e.g.*, partially or completely free of other polypeptides) or part of a cell lysate. Further, the API,
15 API fragment, API-related polypeptide, a fragment of an API-related polypeptide may be a fusion protein comprising the API or a biologically active portion thereof, or API-related polypeptide and a domain such as glutathione-S-transferase. Alternatively, the API, API fragment, API-related polypeptide, fragment of an API-related polypeptide or API fusion protein can be biotinylated using techniques well known to those of skill in the art (*e.g.*,
20 biotinylation kit, Pierce Chemicals; Rockford, IL). The ability of the candidate compound to interact with an API, API fragment, API-related polypeptide, or API-fusion protein can be determined by methods known to those of skill in the art.

In another embodiment, a cell-based assay system is used to identify agents that bind to or modulate the activity of a protein, such as an enzyme, or a biologically active
25 portion thereof, which is responsible for the production or degradation of an API or is responsible for the post- translational modification of an API. In a primary screen, a plurality (*e.g.*, a library) of compounds are contacted with cells that naturally or recombinantly express: (i) an API, API fragment, API-related polypeptide, or API-fusion protein; and (ii) a protein that is responsible for processing of the API, API fragment, API-
30 related polypeptide, or API-fusion protein in order to identify compounds that modulate the production, degradation, or post-translational modification of the API, API fragment, API-related polypeptide, or API-fusion protein. If desired, compounds identified in the primary screen can then be assayed in a secondary screen against cells naturally or recombinantly

expressing the specific API of interest. The ability of the candidate compound to modulate the production, degradation or post-translational modification of an API, API fragment, API-related polypeptide, or API-fusion protein can be determined by methods known to those of skill in the art, including without limitation, flow cytometry, a scintillation assay, immunoprecipitation and western blot analysis.

In another embodiment, agents that competitively interact with an API, API fragment, API-related polypeptide, or API-fusion protein are identified in a competitive binding assay. In accordance with this embodiment, cells expressing an API, API fragment, API-related polypeptide, or API-fusion protein are contacted with a candidate compound and a compound known to interact with the API, API fragment, API-related polypeptide, or API-fusion protein; the ability of the candidate compound to competitively interact with the API, API fragment, API-related polypeptide, or API-fusion protein is then determined. Alternatively, agents that competitively interact with an API, API fragment, API-related polypeptide, or API-fusion protein are identified in a cell-free assay system by contacting an API, API fragment, API-related polypeptide, or API-fusion protein with a candidate compound and a compound known to interact with the API, API fragment, API-related polypeptide, or API-fusion protein. As stated above, the ability of the candidate compound to interact with an API, API fragment, API-related polypeptide, or API-fusion protein can be determined by methods known to those of skill in the art. These assays, whether cell-based or cell-free, can be used to screen a plurality (*e.g.*, a library) of candidate compounds.

In another embodiment, agents that modulate the expression of an API, API fragment, API-related polypeptide, or API-fusion protein are identified by contacting cells (*e.g.*, cells of prokaryotic origin or eukaryotic origin) expressing the API, API fragment, API-related polypeptide, or API-fusion protein with a candidate agent or a control agent (*e.g.*, phosphate buffered saline (PBS)) and determining the expression of the API, API fragment, API-related polypeptide, or API-fusion protein, or the expression of mRNA encoding the API, API fragment, API-related polypeptide, or API-fusion protein. The level of expression of a selected API, API fragment, API-related polypeptide, or API-fusion protein or mRNA encoding the API, API fragment, API-related polypeptide, or API-fusion protein in the presence of the candidate compound is compared to the level of expression of the API, API fragment, API-related polypeptide, or API-fusion protein or mRNA encoding the API, API fragment, API-related polypeptide, or API-fusion protein in the absence of the

candidate agent (*e.g.*, in the presence of a control agent). The candidate agent can then be identified as a modulator of the expression of the API, API fragment, API-related polypeptide, or API-fusion protein or mRNA encoding the API, API fragment, API-related polypeptide, or API-fusion protein based on this comparison. For example, when
5 expression of the API or mRNA is significantly greater in the presence of the candidate agent than in its absence, the candidate agent is identified as a stimulator of expression of the API or its mRNA. Alternatively, when expression of the API or mRNA is significantly less in the presence of the candidate agent than in its absence, the candidate agent is identified as an inhibitor of the expression of the API or its mRNA. The level of expression
10 of an API or the mRNA that encodes it can be determined by methods known to those of skill in the art based on the present description. For example, mRNA expression can be assessed by Northern blot analysis or RT-PCR, and protein levels can be assessed by western blot analysis.

In another embodiment, agents that modulate the activity of an API, API fragment, API-related polypeptide, or API-fusion protein are identified by contacting a preparation
15 containing the API, API fragment, API-related polypeptide, or API-fusion protein, or cells (*e.g.*, prokaryotic or eukaryotic cells) expressing the API, API fragment, API-related polypeptide, or API-fusion protein with a candidate agent or a control agent and determining the ability of the candidate agent to modulate the activity of the API, API
20 fragment, API-related polypeptide, or API-fusion protein. The activity of an API, API fragment, API-related polypeptide, or API-fusion protein can be assessed by detecting induction of a downstream effector of the API, API fragment, API-related polypeptide, or API-fusion protein (*e.g.*, intracellular Ca^{2+} , diacylglycerol, IP3, etc.), detecting catalytic or enzymatic activity of the target on a suitable substrate, detecting the induction of a reporter
25 gene (*e.g.*, a regulatory element that is responsive to an API, API fragment, API-related polypeptide, or API-fusion protein and is operably linked to a nucleic acid encoding a detectable marker, *e.g.*, luciferase), or detecting a cellular response, for example, cellular differentiation, or cell proliferation as the case may be, based on the present description, techniques known to those of skill in the art can be used for measuring these activities (see,
30 *e.g.*, U.S. 5,401,639). The candidate agent can then be identified as a modulator of the activity of an API, API fragment, API-related polypeptide, or API-fusion protein by comparing the effects of the candidate agent to the control agent. Suitable control agents include phosphate buffered saline (PBS) and normal saline (NS).

In another embodiment, agents that modulate the expression, activity or both the expression and activity of an API, API fragment, API-related polypeptide, or API-fusion protein are identified in an animal model. Examples of suitable animals include, but are not limited to, mice, rats, rabbits, monkeys, guinea pigs, dogs and cats. Preferably, the animal used represent a model of AD (*e.g.*, animals that express human familial AD (FAD) β -amyloid precursor (APP), animals that overexpress human wild-type APP, animals that overexpress β -amyloid 1-42 (β A), animals that express FAD presenillin-1 (PS-1). See, *e.g.*, Higgins, LS, 1999, *Molecular Medicine Today* 5:274-276. In accordance with this embodiment, the candidate agent or a control agent is administered (*e.g.*, orally, rectally or parenterally such as intraperitoneally or intravenously) to a suitable animal and the effect on the expression, activity or both expression and activity of the API, API fragment, API-related polypeptide, or API-fusion protein is determined. Changes in the expression of an API, API fragment, API-related polypeptide, or API-fusion protein can be assessed by any suitable method described above, based on the present description.

In yet another embodiment, an API, API fragment, API-related polypeptide, or API-fusion protein is used as a "bait protein" in a two-hybrid assay or three hybrid assay to identify other proteins that bind to or interact with an API, API fragment, API-related polypeptide, or API-fusion protein (see, *e.g.*, U.S. Patent No. 5,283,317; Zervos et al. (1993) *Cell* 72:223-232; Madura et al. (1993) *J. Biol. Chem.* 268:12046-12054; Bartel et al. (1993) *Bio/Techniques* 14:920-924; Iwabuchi et al. (1993) *Oncogene* 8:1693-1696; and PCT Publication No. WO 94/10300). As those skilled in the art will appreciate, such binding proteins are also likely to be involved in the propagation of signals by the APIs of the invention as, for example, upstream or downstream elements of a signaling pathway involving the APIs of the invention.

As those skilled in the art will appreciate, Table X enumerates scientific publications describing suitable assays for detecting or quantifying enzymatic or binding activity of an API, API fragment, API-related polypeptide, or API-fusion protein. Each such reference is hereby incorporated in its entirety. In a preferred embodiment, an assay referenced in Table X is used in the screens and assays described herein, for example, to screen for or to identify an agent that modulates the activity of (or that modulates both the expression and activity of) an API, API fragment, API-related polypeptide, or API-fusion protein.

Table X.

Table X

API	References
API-39, API-44, API-178, API-188	Structural Biology 7, 312-321, 2000 J. Am. Chem. Soc. 122, 2178-2192, 2000
API-123 API-126	Neuroendocrinology 1992 Mar 55:3 308-16
API-182	Biochem J 1997 Mar 1 322 (Pt 2): 455-60; Biochem Soc Trans 1997 Nov 25:4 S591; Biochim Biophys Acta 1986 Oct 10 888:3 325-31 http://www.promega.com
API-322 API-186	"Assay of human transthyretin-bound holo-retinol-binding protein with reversed-phase high-performance liquid chromatography." J Chromatogr 1991 Jul 5 567:2 369-80 "Liquid-chromatographic assay for free and transthyretin-bound retinol-binding protein in serum from normal humans." Clin Chem 1989 Apr 35:4 582-6

Table X

API	References
API-300 API-305 API-315 API-311 API-356 API-38 API-74 API-105 API-124 API-130 API-138 API-169 API-172	"Development of a rapid kinetic assay for the function of the classical pathway of the complement system and for C2 and C4." J Clin Lab Immunol 1986 Dec 21:4 201-7
API-361 API-52	"High-performance liquid chromatographic assay of chymotrypsin and chymotrypsin-like enzyme activity." J Chromatogr 1987 411:498-501 "Assay for chymotrypsin and trypsin. II. On a few considerations for the assay of the commercial anti-inflammatri enzyme preparations." Eisei Shikenjo Hokoku 1972 90:89-92 "Effect of the reactant mixing sequence on the chymotrypsin inhibition assay." Analyst 1990 Aug 115:8 1143-4
API-330 API-314 API-76 API-78 API-79 API-80 API-82	"Time-resolved immunofluorometric assay of complement C3: application to cerebrospinal fluid." Clin Chem 1993 Feb 39:2 309-12 "A fluorimetric assay for native C3. The hemolytically active form of the third component of human complement."

Table X

API	References
API -140	J Immunol Methods 1987, 102(1):7-14

This invention further provides novel agents identified by the above-described screening assays and uses thereof for treatments as described herein.

5

5.16 Therapeutic Uses Of APIs

The invention provides for treatment or prevention of various diseases and disorders by administration of a therapeutic agent. Such agents include but are not limited to: APIs, API analogs, API-related polypeptides and derivatives (including fragments) thereof; antibodies to the foregoing; nucleic acids encoding API, API fragment, API-related polypeptide, or API-fusion protein; antisense nucleic acids to a gene encoding an API, API fragment, API-related polypeptide, or API-fusion protein; and modulator (*e.g.*, agonists and antagonists) of a gene encoding an API or API-related polypeptide. An important feature of the present invention is the identification of genes encoding APIs involved in AD. AD can be treated (*e.g.* to ameliorate symptoms or to retard onset or progression) or prevented by administration of a therapeutic compound that promotes function or expression of one or more APIs that are decreased in the CSF of subjects having AD, or by administration of a therapeutic compound that reduces function or expression of one or more APIs that are increased in the CSF of subjects having AD.

In one embodiment, one or more antibodies each specifically binding to an API are administered alone or in combination with one or more additional therapeutic compounds or treatments. Examples of such therapeutic compounds or treatments include, but are not limited to, tacrine, donepezil, α -tocopherol, selegeline, NSAIDs, estrogen replacement therapy, physostigmine, rivastigmine, hepastigmine, metrifonate, ENA-713, ginkgo biloba extract, physostigmine, amridin, talsaclidine, zifrosilone, eptastigmine, methanesulfonyl chloride, nefiracetam, ALCAR, talsachidine, xanomeline, galanthamine, and propentofylline.

Preferably, a biological product such as an antibody is allogeneic to the subject to which it is administered. In a preferred embodiment, a human API or a human API-related polypeptide, a nucleotide sequence encoding a human API or a human API-related

30

polypeptide, or an antibody to a human API or a human API-related polypeptide, is administered to a human subject for therapy (*e.g.* to ameliorate symptoms or to retard onset or progression) or prophylaxis.

5 *5.16.1 Treatment And Prevention Of AD*

AD can be treated or prevented by administration to a subject suspected of having or known to have AD or to be at risk of developing AD following administration of an agent that modulates (*i.e.*, increases or decreases) the level or activity (*i.e.*, function) of one or more APIs or the level of one or more AFs that are differentially present in the CSF of subjects having AD compared with CSF of subjects free from AD. In one embodiment, AD is treated by administering to a subject suspected of having or known to have AD or to be at risk of developing AD an agent that upregulates (*i.e.*, increases) the level or activity (*i.e.*, function) of one or more APIs or the level of one or more AFs that are decreased in the CSF of subjects having AD. In another embodiment, an agent is administered that downregulates the level or activity (*i.e.*, function) of one or more APIs - or the level of one or more AFs - that are increased in the CSF of subjects having AD. Examples of such a compound include but are not limited to: APIs, API fragments and API-related polypeptides; nucleic acids encoding an API, an API fragment and an API-related polypeptide (*e.g.*, for use in gene therapy); and, for those APIs or API-related polypeptides with enzymatic activity, compounds or molecules known to modulate that enzymatic activity. Other compounds that can be used, *e.g.*, API agonists, can be identified using *in vitro* assays, as defined or described above or earlier.

AD is also treated or prevented by administration to a subject suspected of having or known to have AD or to be at risk of developing AD of a compound that downregulates the level or activity of one or more APIs - or the level of one or more AFs - that are increased in the CSF of subjects having AD. In another embodiment, a compound is administered that upregulates the level or activity of one or more APIs - or the level of one or more AFs - that are decreased in the CSF of subjects having AD. Examples of such a compound include, but are not limited to, API antisense oligonucleotides, ribozymes, antibodies directed against APIs, and compounds that inhibit the enzymatic activity of an API. Other useful compounds *e.g.*, API antagonists and small molecule API antagonists, can be identified using *in vitro* assays.

In a preferred embodiment, therapy or prophylaxis is tailored to the needs of an

individual subject. Thus, in specific embodiments, compounds that promote the level or function of one or more APIs, or the level of one or more AFs, are therapeutically or prophylactically administered to a subject suspected of having or known to have AD, in whom the levels or functions of said one or more APIs, or levels of said one or more AFs, are absent or are decreased relative to a control or normal reference range. In further embodiments, compounds that promote the level or function of one or more APIs, or the level of one or more AFs, are therapeutically or prophylactically administered to a subject suspected of having or known to have AD in whom the levels or functions of said one or more APIs, or levels of said one or more AFs, are increased relative to a control or to a reference range. In further embodiments, compounds that decrease the level or function of one or more APIs, or the level of one or more AFs, are therapeutically or prophylactically administered to a subject suspected of having or known to have AD in whom the levels or functions of said one or more APIs, or levels of said one or more AFs, are increased relative to a control or to a reference range. In further embodiments, compounds that decrease the level or function of one or more APIs, or the level of one or more AFs, are therapeutically or prophylactically administered to a subject suspected of having or known to have AD in whom the levels or functions of said one or more APIs, or levels of said one or more AFs, are decreased relative to a control or to a reference range. The change in API function or level, or AF level, due to the administration of such compounds can be readily detected, *e.g.*, by obtaining a sample (*e.g.*, a sample of CSF, blood or urine or a tissue sample such as biopsy tissue) and assaying *in vitro* the levels of said AFs or the levels or activities of said APIs, or the levels of mRNAs encoding said APIs. or any combination of the foregoing. Such assays can be performed before and after the administration of the compound as described herein.

The compounds of the invention include but are not limited to any compound, *e.g.*, a small organic molecule, protein, peptide, antibody, nucleic acid, etc. that restores the AD API or AF profile towards normal with the proviso that such compound is not an acetylcholinesterase (AChE) inhibitor (*e.g.*, tacrine, donepezil, rivastigmine, heparastigmine, Metrigonate, physostigmine, Amridin, Talsaclidine, KA-672, Huperzine, P-11012, P-11149, Zifrosilone, Eptastigmine, Methanesulfonyl chloride, and S-9977), an acetylcholine receptor agonist (*e.g.*, Nefiracetam, LU-25109, and NS2330), a muscarinic receptor agonist (*e.g.*, SB-20206, Talsachidine, AF-1025B, and SR-46559A), a nicotonic cholinergic receptor agonist (*e.g.*, ABT-418), an acetylcholine modulator (*e.g.*, FKS-508 and

Galantamine) or propentofylline.

5.16.2 Gene Therapy

In another embodiment, nucleic acids comprising a sequence encoding an API, an API fragment, API-related polypeptide or fragment of an API-related polypeptide, are administered to promote API function by way of gene therapy. Gene therapy refers to the administration of an expressed or expressible nucleic acid to a subject. In this embodiment, the nucleic acid produces its encoded polypeptide and the polypeptide mediates a therapeutic effect by promoting API function.

Any suitable methods for gene therapy available in the art can be used according to the present invention.

For general reviews of the methods of gene therapy, see Goldspiel et al., 1993, Clinical Pharmacy 12:488-505; Wu and Wu, 1991, Biotherapy 3:87-95; Tolstoshev, 1993, Ann. Rev. Pharmacol. Toxicol. 32:573-596; Mulligan, 1993, Science 260:926-932; and Morgan and Anderson, 1993, Ann. Rev. Biochem. 62:191-217; May, 1993, TIBTECH 11(5):155-215. Methods commonly known in the art of recombinant DNA technology which can be used in the present invention are described in Ausubel et al. (eds.), 1993, Current Protocols in Molecular Biology, John Wiley & Sons, NY; and Kriegler, 1990, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY.

In a particular aspect, the compound comprises a nucleic acid encoding an API or fragment or chimeric protein thereof, said nucleic acid being part of an expression vector that expresses an API or fragment or chimeric protein thereof in a suitable host. In particular, such a nucleic acid has a promoter operably linked to the API coding region, said promoter being inducible or constitutive (and, optionally, tissue-specific). In another particular embodiment, a nucleic acid molecule is used in which the API coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the API nucleic acid (Koller and Smithies, 1989, Proc. Natl. Acad. Sci. USA 86:8932-8935; Zijlstra et al., 1989, Nature 342:435-438).

Delivery of the nucleic acid into a subject may be direct, in which case the subject is directly exposed to the nucleic acid or nucleic acid-carrying vector; this approach is known as *in vivo* gene therapy. Alternatively, delivery of the nucleic acid into the subject may be indirect, in which case cells are first transformed with the nucleic acid *in vitro* and then

transplanted into the subject, known as "*ex vivo* gene therapy".

In another embodiment, the nucleic acid is directly administered *in vivo*, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, *e.g.*, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, *e.g.*, by infection using a defective or attenuated retroviral or other viral vector (see U.S. Patent No. 4,980,286); by direct injection of naked DNA; by use of microparticle bombardment (*e.g.*, a gene gun; Biolistic, Dupont); by coating with lipids, cell-surface receptors or transfecting agents; by encapsulation in liposomes, microparticles or microcapsules; by administering it in linkage to a peptide which is known to enter the nucleus; or by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see, *e.g.*, Wu and Wu, 1987, J. Biol. Chem. 262:4429-4432), which can be used to target cell types specifically expressing the receptors. In another embodiment, a nucleic acid-ligand complex can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted *in vivo* for cell specific uptake and expression, by targeting a specific receptor (see, *e.g.*, PCT Publications WO 92/06180 dated April 16, 1992 (Wu et al.); WO 92/22635 dated December 23, 1992 (Wilson et al.); WO92/20316 dated November 26, 1992 (Findeis et al.); WO93/14188 dated July 22, 1993 (Clarke et al.), WO 93/20221 dated October 14, 1993 (Young)). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (Koller and Smithies, 1989, Proc. Natl. Acad. Sci. USA 86:8932-8935; Zijlstra et al., 1989, Nature 342:435-438).

In a further embodiment, a viral vector that contains a nucleic acid encoding an API is used. For example, a retroviral vector can be used (see Miller et al., 1993, Meth. Enzymol. 217:581-599). These retroviral vectors have been modified to delete retroviral sequences that are not necessary for packaging of the viral genome and integration into host cell DNA. The nucleic acid encoding the API to be used in gene therapy is cloned into the vector, which facilitates delivery of the gene into a subject. More detail about retroviral vectors can be found in Boesen et al., 1994, Biotherapy 6:291-302, which describes the use of a retroviral vector to deliver the *mdr1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes et al., 1994, J. Clin. Invest. 93:644-651;

Kiem et al., 1994, Blood 83:1467-1473; Salmons and Gunzberg, 1993, Human Gene Therapy 4:129-141; and Grossman and Wilson, 1993, Curr. Opin. in Genetics and Devel. 3:110-114.

Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson, 1993, Current Opinion in Genetics and Development 3:499-503 present a review of adenovirus-based gene therapy. Bout et al., 1994, Human Gene Therapy 5:3-10 demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld et al., 1991, Science 252:431-434; Rosenfeld et al., 1992, Cell 68:143-155; Mastrangeli et al., 1993, J. Clin. Invest. 91:225-234; PCT Publication WO94/12649; and Wang, et al., 1995, Gene Therapy 2:775-783.

Adeno-associated virus (AAV) has also been proposed for use in gene therapy (Walsh et al., 1993, Proc. Soc. Exp. Biol. Med. 204:289-300; U.S. Patent No. 5,436,146).

Another suitable approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a subject.

In this embodiment, the nucleic acid is introduced into a cell prior to administration *in vivo* of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign genes into cells (see, *e.g.*, Loeffler and Behr, 1993, Meth. Enzymol. 217:599-618; Cohen et al., 1993, Meth. Enzymol. 217:618-644; Cline, 1985, Pharmac. Ther. 29:69-92) and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not

disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

The resulting recombinant cells can be delivered to a subject by various methods known in the art. In a preferred embodiment, epithelial cells are injected, *e.g.*, subcutaneously. In another embodiment, recombinant skin cells may be applied as a skin graft onto the subject. Recombinant blood cells (*e.g.*, hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, the condition of the subject, etc., and can be determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to neuronal cells, glial cells (*e.g.*, oligodendrocytes or astrocytes), epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, *e.g.*, as obtained from bone marrow, umbilical cord blood, peripheral blood or fetal liver.

In a preferred embodiment, the cell used for gene therapy is autologous to the subject that is treated.

In an embodiment in which recombinant cells are used in gene therapy, a nucleic acid encoding an API is introduced into the cells such that it is expressible by the cells or their progeny, and the recombinant cells are then administered *in vivo* for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem or progenitor cells which can be isolated and maintained *in vitro* can be used in accordance with this embodiment of the present invention (see *e.g.* PCT Publication WO 94/08598, dated April 28, 1994; Stemple and Anderson, 1992, Cell 71:973-985; Rheinwald, 1980, Meth. Cell Bio. 21A:229; and Pittelkow and Scott, 1986, Mayo Clinic Proc. 61:771).

In another embodiment, the nucleic acid to be introduced for purposes of gene therapy may comprise an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription.

Direct injection of a DNA coding for an API may also be performed according to,

for example, the techniques described in United States Patent No. 5,589,466. These techniques involve the injection of "naked DNA", *i.e.*, isolated DNA molecules in the absence of liposomes, cells, or any other material besides a suitable carrier. The injection of DNA encoding a protein and operably linked to a suitable promoter results in the production of the protein in cells near the site of injection and the elicitation of an immune response in the subject to the protein encoded by the injected DNA. In a preferred embodiment, naked DNA comprising (a) DNA encoding an API and (b) a promoter are injected into a subject to elicit an immune response to the API.

5.16.3 Inhibition Of APIs To Treat AD

In one embodiment of the invention, AD is treated or prevented by administration of a compound that antagonizes (inhibits) the level(s) and/or function(s) of one or more APIs which are elevated in the CSF of subjects having AD as compared with CSF of subjects free from AD. Compounds useful for this purpose include but are not limited to anti-API antibodies (and fragments and derivatives containing the binding region thereof), API antisense or ribozyme nucleic acids, and nucleic acids encoding dysfunctional APIs that are used to "knockout" endogenous API function by homologous recombination (see, *e.g.*, Capecchi, 1989, Science 244:1288-1292). Other compounds that inhibit API function can be identified by use of known *in vitro* assays, *e.g.*, assays for the ability of a test compound to inhibit binding of an API to another protein or a binding partner, or to inhibit a known API function. Preferably such inhibition is assayed *in vitro* or in cell culture, but genetic assays may also be employed. The Preferred Technology can also be used to detect levels of the API before and after the administration of the compound. Preferably, suitable *in vitro* or *in vivo* assays are utilized to determine the effect of a specific compound and whether its administration is indicated for treatment of the affected tissue, as described in more detail below.

In a particular embodiment, a compound that inhibits an API function is administered therapeutically or prophylactically to a subject in whom an increased CSF level or functional activity of the API (*e.g.*, greater than the normal level or desired level) is detected as compared with CSF of subjects free from AD or a predetermined reference range. Methods standard in the art can be employed to measure the increase in an API level or function, as outlined above. Preferred API inhibitor compositions include small molecules, *i.e.*, molecules of 1000 daltons or less. Such small molecules can be identified

by the screening methods described herein.

5.16.4 *Antisense Regulation of APIs*

In a further embodiment, API expression is inhibited by use of API antisense
5 nucleic acids. The present invention provides the therapeutic or prophylactic use of nucleic acids comprising at least six nucleotides that are antisense to a gene or cDNA encoding an API or a portion thereof. As used herein, an API "antisense" nucleic acid refers to a nucleic acid capable of hybridizing by virtue of some sequence complementarity to a portion of an RNA (preferably mRNA) encoding an API. The antisense nucleic acid may be
10 complementary to a coding and/or noncoding region of an mRNA encoding an API. Such antisense nucleic acids have utility as compounds that inhibit API expression, and can be used in the treatment or prevention of AD.

The antisense nucleic acids of the invention are double-stranded or single-stranded oligonucleotides, RNA or DNA or a modification or derivative thereof, and can be directly
15 administered to a cell or produced intracellularly by transcription of exogenous, introduced sequences.

The invention further provides pharmaceutical compositions comprising a therapeutically effective amount of an API antisense nucleic acid, and a pharmaceutically-acceptable carrier, vehicle or diluent.

20 In another embodiment, the invention provides methods for inhibiting the expression of an API nucleic acid sequence in a prokaryotic or eukaryotic cell comprising providing the cell with an effective amount of a composition comprising an API antisense nucleic acid of the invention.

API antisense nucleic acids and their uses are described in detail below.

25

5.16.5 *API Antisense Nucleic Acids*

The API antisense nucleic acids are of at least six nucleotides and are preferably oligonucleotides ranging from 6 to about 50 oligonucleotides. In specific aspects, the oligonucleotide is at least 10 nucleotides, at least 15 nucleotides, at least 100 nucleotides, or
30 at least 200 nucleotides. The oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof and can be single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone. The oligonucleotide may include other appended groups such as peptides;

agents that facilitate transport across the cell membrane (see, *e.g.*, Letsinger et al., 1989, Proc. Natl. Acad. Sci. USA 86:6553-6556; Lemaitre et al., 1987, Proc. Natl. Acad. Sci. 84:648-652; PCT Publication No. WO 88/09810, published December 15, 1988) or blood-brain barrier (see, *e.g.*, PCT Publication No. WO 89/10134, published April 25, 1988);
5 hybridization-triggered cleavage agents (see, *e.g.*, Krol et al., 1988, BioTechniques 6:958-976) or intercalating agents (see, *e.g.*, Zon, 1988, Pharm. Res. 5:539-549).

In a particular aspect of the invention, an API antisense oligonucleotide is provided, preferably of single-stranded DNA. The oligonucleotide may be modified at any position on its structure with substituents generally known in the art.

10 The API antisense oligonucleotide may comprise any suitable of the following modified base moieties, *e.g.* 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-
15 methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-
20 methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, 2,6-diaminopurine, and other base analogs.

In another embodiment, the oligonucleotide comprises at least one modified sugar moiety, *e.g.*, one of the following sugar moieties: arabinose, 2-fluoroarabinose, xylulose,
25 and hexose.

In yet another embodiment, the oligonucleotide comprises at least one of the following modified phosphate backbones: a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, a formacetal, or an analog of formacetal.

30 In yet another embodiment, the oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gautier et al., 1987, Nucl. Acids Res. 15:6625-6641).

The oligonucleotide may be conjugated to another molecule, *e.g.*, a peptide, hybridization triggered cross-linking agent, transport agent, or hybridization-triggered cleavage agent.

Oligonucleotides of the invention may be synthesized by standard methods known in the art, *e.g.*, by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein et al. (1988, Nucl. Acids Res. 16:3209), and methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. USA 85:7448-7451).

In another embodiment, the API antisense nucleic acid of the invention is produced intracellularly by transcription from an exogenous sequence. For example, a vector can be introduced *in vivo* such that it is taken up by a cell, within which cell the vector or a portion thereof is transcribed, producing an antisense nucleic acid (RNA) of the invention. Such a vector would contain a sequence encoding the API antisense nucleic acid. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in mammalian cells. Expression of the sequence encoding the API antisense RNA can be by any promoter known in the art to act in mammalian, preferably human, cells. Such promoters can be inducible or constitutive. Examples of such promoters are outlined above.

The antisense nucleic acids of the invention comprise a sequence complementary to at least a portion of an RNA transcript of a gene encoding an API, preferably a human gene encoding an API. However, absolute complementarity, although preferred, is not required. A sequence "complementary to at least a portion of an RNA," as referred to herein, means a sequence having sufficient complementarity to be able to hybridize under stringent conditions (*e.g.*, highly stringent conditions comprising hybridization in 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65 °C and washing in 0.1xSSC/0.1% SDS at 68 °C, or moderately stringent conditions comprising washing in 0.2xSSC/0.1% SDS at 42 °C) with the RNA, forming a stable duplex; in the case of double-stranded API antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the longer the

hybridizing nucleic acid, the more base mismatches with an RNA encoding an API it may contain and still form a stable duplex (or triplex, as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

5

5.16.6 Therapeutic Use Of API Antisense Nucleic Acids

The API antisense nucleic acids can be used to treat or prevent AD when the target API is overexpressed in the CSF of subjects suspected of having or suffering from AD. In a preferred embodiment, a single-stranded DNA antisense API oligonucleotide is used.

10 Cell types which express or overexpress RNA encoding an API can be identified by various methods known in the art. Such cell types include but are not limited to leukocytes (*e.g.*, neutrophils, macrophages, monocytes) and resident cells (*e.g.*, astrocytes, glial cells, neuronal cells, and ependymal cells). Such methods include, but are not limited to, hybridization with an API-specific nucleic acid (*e.g.*, by Northern hybridization, dot blot
15 hybridization, in situ hybridization), observing the ability of RNA from the cell type to be translated *in vitro* into an API, immunoassay, etc. In a preferred aspect, primary tissue from a subject can be assayed for API expression prior to treatment, *e.g.*, by immunocytochemistry or in situ hybridization.

Pharmaceutical compositions of the invention, comprising an effective amount of an
20 API antisense nucleic acid in a pharmaceutically acceptable carrier, vehicle or diluent can be administered to a subject having AD.

The amount of API antisense nucleic acid which will be effective in the treatment of AD can be determined by standard clinical techniques.

In a specific embodiment, pharmaceutical compositions comprising one or more
25 API antisense nucleic acids are administered via liposomes, microparticles, or microcapsules. In various embodiments of the invention, such compositions may be used to achieve sustained release of the API antisense nucleic acids.

5.16.7 Inhibitory Ribozyme And Triple Helix Approaches

30 In another embodiment, symptoms of AD may be ameliorated by decreasing the level of an API or API activity by using gene sequences encoding the API in conjunction with well-known gene "knock-out," ribozyme or triple helix methods to decrease gene expression of an API. In this approach ribozyme or triple helix molecules are used to

modulate the activity, expression or synthesis of the gene encoding the API, and thus to ameliorate the symptoms of AD. Such molecules may be designed to reduce or inhibit expression of a mutant or non-mutant target gene. Techniques for the production and use of such molecules are well known to those of skill in the art.

5 Ribozyme molecules designed to catalytically cleave gene mRNA transcripts encoding an API can be used to prevent translation of target gene mRNA and, therefore, expression of the gene product. (See, *e.g.*, PCT International Publication WO90/11364, published October 4, 1990; Sarver et al., 1990, Science 247:1222-1225).

10 Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. (For a review, see Rossi, 1994, Current Biology 4, 469-471). The mechanism of ribozyme action involves sequence specific hybridization of the ribozyme molecule to complementary target RNA, followed by an endonucleolytic cleavage event. The composition of ribozyme molecules must include one or more sequences complementary to the target gene mRNA, and must include the well known catalytic
15 sequence responsible for mRNA cleavage. For this sequence, see, *e.g.*, U.S. Patent No. 5,093,246, which is incorporated herein by reference in its entirety.

 While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy mRNAs encoding an API, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form
20 complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Myers, 1995, Molecular Biology and Biotechnology: A Comprehensive Desk Reference, VCH Publishers, New York, (see especially Figure 4, page 833) and in Haseloff and
25 Gerlach, 1988, Nature, 334, 585-591, each of which is incorporated herein by reference in its entirety.

 Preferably the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the mRNA encoding the API, *i.e.*, to increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

30 The ribozymes of the present invention also include RNA endoribonucleases (hereinafter "Cech-type ribozymes") such as the one that occurs naturally in Tetrahymena thermophila (known as the IVS, or L-19 IVS RNA) and that has been extensively described by Thomas Cech and collaborators (Zaug, et al., 1984, Science, 224, 574-578; Zaug and

Cech, 1986, Science, 231, 470-475; Zaug, et al., 1986, Nature, 324, 429-433; published International patent application No. WO 88/04300 by University Patents Inc.; Been and Cech, 1986, Cell, 47, 207-216). The Cech-type ribozymes have an eight base pair active site which hybridizes to a target RNA sequence whereafter cleavage of the target RNA takes place. The invention encompasses those Cech-type ribozymes which target eight base-pair active site sequences that are present in the gene encoding the API.

As in the antisense approach, the ribozymes can be composed of modified oligonucleotides (*e.g.*, for improved stability, targeting, etc.) and should be delivered to cells that express the API *in vivo*. A preferred method of delivery involves using a DNA construct "encoding" the ribozyme under the control of a strong constitutive pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous mRNA encoding the API and inhibit translation. Because ribozymes, unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficacy.

Endogenous API expression can also be reduced by inactivating or "knocking out" the gene encoding the API, or the promoter of such a gene, using targeted homologous recombination (*e.g.*, see Smithies, et al., 1985, Nature 317:230-234; Thomas and Capecchi, 1987, Cell 51:503-512; Thompson et al., 1989, Cell 5:313-321; and Zijlstra et al., 1989, Nature 342:435-438, each of which is incorporated by reference herein in its entirety). For example, a mutant gene encoding a non-functional API (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous gene (either the coding regions or regulatory regions of the gene encoding the API) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express the target gene *in vivo*. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the target gene. Such approaches are particularly suited in the agricultural field where modifications to ES (embryonic stem) cells can be used to generate animal offspring with an inactive target gene (*e.g.*, see Thomas and Capecchi, 1987 and Thompson, 1989, *supra*). However, this approach can be adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors.

Alternatively, the endogenous expression of a gene encoding an API can be reduced by targeting deoxyribonucleotide sequences complementary to the regulatory region of the gene (*i.e.*, the gene promoter and/or enhancers) to form triple helical structures that prevent

transcription of the gene encoding the API in target cells in the body. (See generally, Helene, 1991, *Anticancer Drug Des.*, 6(6), 569-584; Helene, et al., 1992, *Ann. N.Y. Acad. Sci.*, 660, 27-36; and Maher, 1992, *Bioassays* 14(12), 807-815).

5 Nucleic acid molecules to be used in triplex helix formation for the inhibition of transcription in the present invention should be single stranded and composed of deoxynucleotides. The base composition of these oligonucleotides must be designed to promote triple helix formation via Hoogsteen base pairing rules, which generally require sizeable stretches of either purines or pyrimidines to be present on one strand of a duplex. Nucleotide sequences may be pyrimidine-based, which will result in TAT and CGC+
10 triplets across the three associated strands of the resulting triple helix. The pyrimidine-rich molecules provide base complementarity to a purine-rich region of a single strand of the duplex in a parallel orientation to that strand. In addition, nucleic acid molecules may be chosen that are purine-rich, for example, contain a stretch of G residues. These molecules will form a triple helix with a DNA duplex that is rich in GC pairs, in which the majority of
15 the purine residues are located on a single strand of the targeted duplex, resulting in GGC triplets across the three strands in the triplex.

Alternatively, the potential sequences that can be targeted for triple helix formation may be increased by creating a so called "switchback" nucleic acid molecule. Switchback molecules are synthesized in an alternating 5'-3', 3'-5' manner, such that they base pair with
20 first one strand of a duplex and then the other, eliminating the necessity for a sizeable stretch of either purines or pyrimidines to be present on one strand of a duplex.

In one embodiment, wherein the antisense, ribozyme, or triple helix molecules described herein are utilized to inhibit mutant gene expression, it is possible that the technique may so efficiently reduce or inhibit the transcription (triple helix) or translation
25 (antisense, ribozyme) of mRNA produced by normal gene alleles of an API that the situation may arise wherein the concentration of API present may be lower than is necessary for a normal phenotype. In such cases, to ensure that substantially normal levels of activity of a gene encoding an API are maintained, gene therapy may be used to introduce into cells nucleic acid molecules that encode and express the API that exhibit
30 normal gene activity and that do not contain sequences susceptible to whatever antisense, ribozyme, or triple helix treatments are being utilized. Alternatively, in instances whereby the gene encodes an extracellular protein, normal API can be co-administered in order to maintain the requisite level of API activity.

Antisense RNA and DNA, ribozyme, and triple helix molecules of the invention may be prepared by any method known in the art for the synthesis of DNA and RNA molecules, as discussed above. These include techniques for chemically synthesizing oligodeoxyri-bonucleotides and oligoribonucleotides well known in the art such as for example solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding the antisense RNA molecule. Such DNA sequences may be incorporated into a wide variety of vectors that incorporate suitable RNA polymerase promoters such as the T7 or SP6 polymerase promoters. Alternatively, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly, depending on the promoter used, can be introduced stably into cell lines.

5.17 Assays For Therapeutic Or Prophylactic Compounds

The present invention also provides assays for use in discovery of pharmaceutical products in order to identify or verify the efficacy of compounds for treatment or prevention of AD. Agents can be assayed for their ability to restore AF or API levels in a subject having AD towards levels found in subjects free from AD or to produce similar changes in experimental animal models of AD. Compounds able to restore AF or API levels in a subject having AD towards levels found in subjects free from AD or to produce similar changes in experimental animal models of Alzheimer's disease can be used as lead compounds for further drug discovery, or used therapeutically. AF and API expression can be assayed by the Preferred Technology, immunoassays, gel electrophoresis followed by visualization, detection of API activity, or any other method taught herein or known to those skilled in the art. Such assays can be used to screen candidate drugs, in clinical monitoring or in drug development, where abundance of an AF or API can serve as a surrogate marker for clinical disease.

In various embodiments, *in vitro* assays can be carried out with cells representative of cell types involved in a subject's disorder, to determine if a compound has a desired effect upon such cell types.

Compounds for use in therapy can be tested in suitable animal model systems prior to testing in humans, including but not limited to rats, mice, chicken, cows, monkeys, rabbits, etc. For *in vivo* testing, prior to administration to humans, any animal model system known in the art may be used. Examples of animal models of AD include, but are

not limited to, animals that express human familial AD (FAD) β -amyloid precursor (APP), animals that overexpress human wild-type APP, animals that overexpress β -amyloid 1-42 (β A), animals that express FAD presenillin-1 (PS-1) (see, *e.g.*, Higgins, LS, 1999, Molecular Medicine Today 5:274-276). Further, animal models for Downs syndrome (*e.g.*, TgSOD1, TgPFFKL, TgS100 β , TgAPP, TgEts2, TgHMG14, TgMNB, Ts65Dn, and Ts1Cje (see, *e.g.*, Kola et al., 1999, Molecular Medicine Today 5:276-277) can be utilized to test compounds that modulate AF or API levels since the neuropathology exhibited by individuals with Downs syndrome is similar to that of AD. It is also apparent to the skilled artisan that, based upon the present disclosure, transgenic animals can be produced with "knock-out" mutations of the gene or genes encoding one or more APIs. A "knock-out" mutation of a gene is a mutation that causes the mutated gene to not be expressed, or expressed in an aberrant form or at a low level, such that the activity associated with the gene product is nearly or entirely absent. Preferably, the transgenic animal is a mammal, more preferably, the transgenic animal is a mouse.

In one embodiment, test compounds that modulate the expression of an API are identified in non-human animals (*e.g.*, mice, rats, monkeys, rabbits, and guinea pigs), preferably non-human animal models for AD or Downs syndrome, expressing the API. In accordance with this embodiment, a test compound or a control compound is administered to the animals, and the effect of the test compound on expression of one or more APIs is determined. A test compound that alters the expression of an API (or a plurality of APIs) can be identified by comparing the level of the selected API or APIs (or mRNA(s) encoding the same) in an animal or group of animals treated with a test compound with the level of the API(s) or mRNA(s) in an animal or group of animals treated with a control compound. Techniques known to those of skill in the art can be used to determine the mRNA and protein levels, for example, in situ hybridization. The animals may or may not be sacrificed to assay the effects of a test compound.

In another embodiment, test compounds that modulate the activity of an API or a biologically active portion thereof are identified in non-human animals (*e.g.*, mice, rats, monkeys, rabbits, and guinea pigs), preferably non-human animal models for AD or Downs syndrome, expressing the API. In accordance with this embodiment, a test compound or a control compound is administered to the animals, and the effect of a test compound on the activity of an API is determined. A test compound that alters the activity of an API (or a plurality of APIs) can be identified by assaying animals treated with a control compound

and animals treated with the test compound. The activity of the API can be assessed by detecting induction of a downstream effector of the API (*e.g.*, intracellular Ca²⁺, diacylglycerol, IP₃, etc.), detecting catalytic or enzymatic activity of the API or binding partner thereof, detecting the induction of a reporter gene (*e.g.*, a regulatory element that is responsive to an API of the invention operably linked to a nucleic acid encoding a detectable marker, such as luciferase or green fluorescent protein), or detecting a cellular response (*e.g.*, cellular differentiation or cell proliferation). Techniques known to those of skill in the art can be utilized to detect changes in the activity of an API (see, *e.g.*, U.S. Patent No. 5,401,639, which is incorporated herein in its entirety by reference).

In yet another embodiment, test compounds that modulate the level or expression of an API (or plurality of APIs) are identified in human subjects having AD or Downs syndrome, preferably those having mild to severe AD and most preferably those having mild AD. In accordance with this embodiment, a test compound or a control compound is administered to the human subject, and the effect of a test compound on API expression is determined by analyzing the expression of the API or the mRNA encoding the same in a biological sample (*e.g.*, CSF, serum, plasma, or urine). A test compound that alters the expression of an API can be identified by comparing the level of the API or mRNA encoding the same in a subject or group of subjects treated with a control compound to that in a subject or group of subjects treated with a test compound. Alternatively, alterations in the expression of an API can be identified by comparing the level of the API or mRNA encoding the same in a subject or group of subjects before and after the administration of a test compound. Any suitable techniques known to those of skill in the art can be used to obtain the biological sample and analyze the mRNA or protein expression. For example, the Preferred Technology described herein can be used to assess changes in the level of an API.

In another embodiment, test compounds that modulate the activity of an API (or plurality of APIs) are identified in human subjects having AD or Downs syndrome, preferably those having mild to severe AD and most preferably those with mild AD. In this embodiment, a test compound or a control compound is administered to the human subject, and the effect of a test compound on the activity of an API is determined. A test compound that alters the activity of an API can be identified by comparing biological samples from subjects treated with a control compound to samples from subjects treated with the test compound. Alternatively, alterations in the activity of an API can be identified by

comparing the activity of an API in a subject or group of subjects before and after the administration of a test compound. The activity of the API can be assessed by detecting in a biological sample (*e.g.*, CSF, serum, plasma, or urine) induction of a downstream effector of the API (*e.g.*, intracellular Ca²⁺, diacylglycerol, IP₃, etc.), catalytic or enzymatic activity of the API or a binding partner thereof, or a cellular response, for example, cellular differentiation, or cell proliferation. Techniques known to those of skill in the art can be used to detect changes in the induction of a second messenger of an API or changes in a cellular response. For example, RT-PCR can be used to detect changes in the induction of a cellular second messenger.

In a particular embodiment, an agent that changes the level or expression of an API towards levels detected in control subjects (*e.g.*, humans free from AD) is selected for further testing or therapeutic use. In another preferred embodiment, a test compound that changes the activity of an API towards the activity found in control subjects (*e.g.*, humans free from AD) is selected for further testing or therapeutic use.

In another embodiment, test compounds that reduce the severity of one or more symptoms associated with AD are identified in human subjects having AD or Downs syndrome, preferably subjects having mild to severe AD and most preferably subjects with mild AD. In accordance with this embodiment, a test compound or a control compound is administered to the subjects, and the effect of a test compound on one or more symptoms of AD is determined. A test compound that reduces one or more symptoms can be identified by comparing the subjects treated with a control compound to the subjects treated with the test compound. Techniques known to physicians familiar with AD can be used to determine whether a test compound reduces one or more symptoms associated with AD. For example, a test compound that enhances memory or reduces confusion in a subject having AD will be beneficial for treating subjects having AD.

In a preferred embodiment, an agent that reduces the severity of one or more symptoms associated with AD in a human having AD is selected for further testing or therapeutic use.

5.18 Therapeutic and Prophylactic Compositions and Their Use

The invention provides methods of treatment comprising administering to a subject an effective amount of an agent of the invention. In a preferred aspect, the compound is substantially purified (*e.g.*, substantially free from substances that limit its effect or produce

undesired side-effects). The subject is preferably an animal, including but not limited to animals such as cows, pigs, horses, chickens, cats, dogs, etc., and is preferably a mammal, and most preferably human. In a specific embodiment, a non-human mammal is the subject.

5 Formulations and methods of administration that can be employed when the compound comprises a nucleic acid are described above; additional appropriate formulations and routes of administration are described below.

 Various delivery systems are known and can be used to administer a compound of the invention, *e.g.*, encapsulation in liposomes, microparticles, microcapsules, recombinant
10 cells capable of expressing the compound, receptor-mediated endocytosis (see, *e.g.*, Wu and Wu, 1987, J. Biol. Chem. 262:4429-4432), construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of introduction can be enteral or parenteral and include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The compounds may be administered
15 by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (*e.g.*, oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, it may be desirable to introduce the pharmaceutical compositions of the invention into the central nervous system by any suitable route,
20 including intraventricular and intrathecal injection; intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir. Pulmonary administration can also be employed, *e.g.*, by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

 In a specific embodiment, it may be desirable to administer the pharmaceutical
25 compositions of the invention locally to the area in need of treatment; this may be achieved, for example, and not by way of limitation, by local infusion during surgery, topical application, *e.g.*, by injection, by means of a catheter, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. In one embodiment, administration can be by direct
30 injection into CSF or at the site (or former site) of neurodegeneration or to CNS tissue.

 In another embodiment, the compound can be delivered in a vesicle, in particular a liposome (see Langer, 1990, Science 249:1527-1533; Treat et al., in Liposomes in the Therapy of Infectious Disease and Cancer, Lopez-Berestein and Fidler (eds.), Liss, New

York, pp. 353-365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*)

In yet another embodiment, the compound can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, *supra*; Sefton, 1987, CRC Crit. Ref. Biomed. Eng. 14:201; Buchwald et al., 1980, Surgery 88:507; Saudek et al., 1989, N. Engl. J. Med. 321:574). In another embodiment, polymeric materials can be used (see Medical Applications of Controlled Release, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); Controlled Drug Bioavailability, Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, J., 1983, Macromol. Sci. Rev. Macromol. Chem. 23:61; see also Levy et al., 1985, Science 228:190; During et al., 1989, Ann. Neurol. 25:351; Howard et al., 1989, J. Neurosurg. 71:105). In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target, *i.e.*, the brain, thus requiring only a fraction of the systemic dose (see, *e.g.*, Goodson, in Medical Applications of Controlled Release, *supra*, vol. 2, pp. 115-138 (1984)).

Other suitable controlled release systems are discussed in the review by Langer (1990, Science 249:1527-1533).

In another embodiment where the compound of the invention is a nucleic acid encoding a protein, the nucleic acid can be administered *in vivo* to promote expression of its encoded protein, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, *e.g.*, by use of a retroviral vector (see U.S. Patent No. 4,980,286), or by direct injection, or by use of microparticle bombardment (*e.g.*, a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox-like peptide which is known to enter the nucleus (see *e.g.*, Joliot et al., 1991, Proc. Natl. Acad. Sci. USA 88:1864-1868), etc. Alternatively, a nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination.

The present invention also provides pharmaceutical compositions. Such compositions comprise a therapeutically effective amount of an agent, and a pharmaceutically acceptable carrier. In a particular embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is

administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" Ed. E.W. Martin, ISBN: 0-912734-04-3, Mack Publishing Co. Such compositions will contain a therapeutically effective amount of the compound, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the subject. The formulation should suit the mode of administration.

In a preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lidocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

The compounds of the invention can be formulated as neutral or salt forms.

Pharmaceutically acceptable salts include those formed with free amino groups such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with free carboxyl groups such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

The amount of the compound of the invention which will be effective in the treatment of AD can be determined by standard clinical techniques based on the present description. In addition, *in vitro* assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each subject's circumstances. However, suitable dosage ranges for intravenous administration are generally about 20-500 micrograms of active compound per kilogram body weight. Suitable dosage ranges for intranasal administration are generally about 0.01 pg/kg body weight to 1 mg/kg body weight. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

Suppositories generally contain active ingredient in the range of 0.5% to 10% by weight; oral formulations preferably contain 10% to 95% active ingredient.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects (a) approval by the agency of manufacture, use or sale for human administration, (b) directions for use, or both.

6. EXAMPLE: IDENTIFICATION OF PROTEINS DIFFERENTIALLY EXPRESSED IN THE CSF IN AD

Using the following exemplary and non-limiting procedure, proteins in CSF samples from (a) 148 subjects having AD, (b) 60 family members of these AD subjects, and (c) 32 unrelated controls were separated by isoelectric focusing followed by SDS-PAGE and analyzed. From some subjects, serial samples were taken over time. Parts 6.1.1 to 6.1.9 (inclusive) of the procedure set forth below are hereby designated as the "Reference Protocol".

6.1 MATERIALS AND METHODS

6.1.1 Sample Preparation

A protein assay (Pierce BCA Cat # 23225) was performed on each CSF sample as received. Prior to protein separation, each sample was processed for selective depletion of certain proteins, in order to enhance and simplify protein separation and facilitate analysis by removing proteins that may interfere with or limit analysis of proteins of interest. See International Patent Application No. PCT/GB99/01742, filed June 1, 1999, which is incorporated by reference in its entirety, with particular reference to pages 3 and 6.

Removal of albumin, haptoglobin, transferrin and immunoglobulin G (IgG) from CSF ("CSF depletion") was achieved by an affinity chromatography purification step in which the sample was passed through a series of 'Hi-Trap' columns containing immobilized antibodies for selective removal of albumin, haptoglobin and transferrin, and protein G for selective removal of immunoglobulin G. Two affinity columns in a tandem assembly were prepared by coupling antibodies to protein G-sepharose contained in Hi-Trap columns (Protein G-Sepharose Hi-Trap columns (1 ml) Pharmacia Cat. No. 17-0404-01). This was done by circulating the following solutions sequentially through the columns: (1) Dulbecco's Phosphate Buffered Saline (Gibco BRL Cat. No. 14190-094); (2) concentrated antibody solution; (3) 200 mM sodium carbonate buffer, pH 8.35; (4) cross-linking solution (200 mM sodium carbonate buffer, pH 8.35, 20 mM dimethylpimelimidate); and (5) 500 mM ethanolamine, 500 mM NaCl. A third (un-derivatised) protein G Hi-Trap column was then attached to the lower end of the tandem column assembly.

The chromatographic procedure was automated using an Akta Fast Protein Liquid Chromatography (FPLC) System such that a series of up to seven runs could be performed sequentially. The samples were passed through the series of 3 Hi-Trap columns in which the affinity chromatography media selectively bind the above proteins thereby removing them from the sample. Fractions (typically 3 ml per tube) were collected of unbound material ("Flowthrough fractions") that eluted through the column during column loading and washing stages and of bound proteins ("Bound/Eluted fractions") that were eluted by step elution with Immunopure Gentle Ag/Ab Elution Buffer (Pierce Cat. No. 21013). The eluate containing unbound material was collected in fractions which were pooled, desalted/concentrated by centrifugal ultrafiltration and stored to await further analysis by 2D PAGE.

A volume of depleted CSF containing approximately 300 µg of total protein was aliquoted and an equal volume of 10% (w/v) SDS (Fluka 71729), 2.3% (w/v) dithiothreitol (BDH 443852A) was added. The sample was heated at 95°C for 5 mins, and then allowed to cool to 20°C. 125µl of the following buffer was then added to the sample:

- 5 8M urea (BDH 452043w)
- 4% CHAPS (Sigma C3023)
- 65mM dithiotheitol (DTT)
- 2% (v/v) Resolytes 3.5-10 (BDH 44338 2x)

 This mixture was vortexed, and centrifuged at 13000 rpm for 5 mins at 15°C, and the
10 supernatant was analyzed by isoelectric focusing.

6.1.2 *Isoelectric Focusing*

Isoelectric focusing (IEF), was performed using the Immobiline® DryStrip Kit (Pharmacia BioTech), following the procedure described in the manufacturer's instructions,
15 see Instructions for Immobiline® DryStrip Kit, Pharmacia, # 18-1038-63, Edition AB (incorporated herein by reference in its entirety). Immobilized pH Gradient (IPG) strips (18cm, pH 3-10 non-linear strips; Pharmacia Cat. # 17-1235-01) were rehydrated overnight at 20°C in a solution of 8M urea, 2% (w/v) CHAPS, 10mM DTT, 2% (v/v) Resolytes 3.5-10, as described in the Immobiline DryStrip Users Manual. For IEF, 50µl of supernatant
20 (prepared as above) was loaded onto a strip, with the cup-loading units being placed at the basic end of the strip. The loaded gels were then covered with mineral oil (Pharmacia 17-3335-01) and a voltage was immediately applied to the strips according to the following profile, using a Pharmacia EPS3500XL power supply (Cat 19-3500-01):

- 25 Initial voltage = 300V for 2 hrs
- Linear Ramp from 300V to 3500V over 3hrs
- Hold at 3500V for 19hrs

 For all stages of the process, the current limit was set to 10mA for 12 gels, and the
30 wattage limit to 5W. The temperature was held at 20°C throughout the run.

6.1.3 Gel Equilibration and SDS-PAGE

After the final 19hr step, the strips were immediately removed and immersed for 10 mins at 20°C in a first solution of the following composition: 6M urea; 2% (w/v) DTT; 2% (w/v) SDS; 30% (v/v) glycerol (Fluka 49767); 0.05M Tris/HCl, pH 6.8 (Sigma Cat T-1503).

- 5 The strips were removed from the first solution and immersed for 10 mins at 20°C in a second solution of the following composition: 6M urea; 2% (w/v) iodoacetamide (Sigma I-6125); 2% (w/v) SDS; 30% (v/v) glycerol; 0.05M Tris/HCl, pH 6.8. After removal from the second solution, the strips were loaded onto supported gels for SDS-PAGE according to Hochstrasser et al., 1988, *Analytical Biochemistry* 173: 412-423 (incorporated herein by
10 reference in its entirety), with modifications as specified below.

6.1.4 Preparation of supported gels

- The gels were cast between two glass plates of the following dimensions: 23cm wide x 24cm long (back plate); 23cm wide x 24cm long with a 2cm deep notch in the central
15 19cm (front plate). To promote covalent attachment of SDS-PAGE gels, the back plate was treated with a 0.4% solution of γ -methacryl-oxypropyltrimethoxysilane in ethanol (BindSilaneTM; Pharmacia Cat. # 17-1330-01). The front plate was treated with a 2% solution of dimethyldichlorosilane dissolved in octamethyl cyclo-octasilane (RepelSilaneTM Pharmacia Cat. # 17-1332-01) to reduce adhesion of the gel. Excess reagent was removed
20 by washing with water, and the plates were allowed to dry. At this stage, both as identification for the gel, and as a marker to identify the coated face of the plate, an adhesive bar-code was attached to the back plate in a position such that it would not come into contact with the gel matrix.

- The dried plates were assembled into a casting box with a capacity of 13 gel
25 sandwiches. The top and bottom plates of each sandwich were spaced by means of 1mm thick spacers, 2.5 cm wide. The sandwiches were interleaved with acetate sheets to facilitate separation of the sandwiches after gel polymerization. Casting was then carried out according to Hochstrasser et al., *op. cit.*

- A 9-16% linear polyacrylamide gradient was cast, extending up to a point 2cm
30 below the level of the notch in the front plate, using the Angelique gradient casting system (Large Scale Biology). Stock solutions were as follows. Acrylamide (40% in water) was from Serva (Cat. # 10677). The cross-linking agent was PDA (BioRad 161-0202), at a concentration of 2.6% (w/w) of the total starting monomer content. The gel buffer was

0.375M Tris/HCl, pH 8.8. The polymerization catalyst was 0.05% (v/v) TEMED (BioRad 161-0801), and the initiator was 0.1% (w/v) APS (BioRad 161-0700). No SDS was included in the gel and no stacking gel was used. The cast gels were allowed to polymerize at 20°C overnight, and then stored at 4°C in sealed polyethylene bags with 6ml of gel buffer, and were used within 4 weeks.

6.1.5 SDS-PAGE

A solution of 0.5% (w/v) agarose (Fluka Cat 05075) was prepared in running buffer (0.025M Tris, 0.198M glycine (Fluka 50050), 1% (w/v) SDS, supplemented by a trace of bromophenol blue). The agarose suspension was heated to 70°C with stirring, until the agarose had dissolved. The top of the supported 2nd D gel was filled with the agarose solution, and the equilibrated strip was placed into the agarose, and tapped gently with a palette knife until the gel was intimately in contact with the 2nd D gel. The gels were placed in the 2nd D running tank, as described by Amess et al., 1995, Electrophoresis 16: 1255-1267 (incorporated herein by reference in its entirety). The tank was filled with running buffer (as above) until the level of the buffer was just higher than the top of the region of the 2nd D gels which contained polyacrylamide, so as to achieve efficient cooling of the active gel area. Running buffer was added to the top buffer compartments formed by the gels, and then voltage was applied immediately to the gels using a Consort E-833 power supply. For 1 hour, the gels were run at 20mA/gel. The wattage limit was set to 150W for a tank containing 6 gels, and the voltage limit was set to 600V. After 1 hour, the gels were then run at 40mA/gel, with the same voltage and wattage limits as before, until the bromophenol blue line was 0.5cm from the bottom of the gel. The temperature of the buffer was held at 16°C throughout the run. Gels were not run in duplicate.

6.1.6 Staining

Upon completion of the electrophoresis run, the gels were immediately removed from the tank for fixation. The top plate of the gel cassette was carefully removed, leaving the gel bonded to the bottom plate. The bottom plate with its attached gel was then placed into a staining apparatus, which can accommodate 12 gels. The gels were completely immersed in fixative solution of 40% (v/v) ethanol (BDH 28719), 10% (v/v) acetic acid (BDH 100016X), 50% (v/v) water (MilliQ-Millipore), which was continuously circulated over the gels. After an overnight incubation, the fixative was drained from the tank, and the

gels were primed by immersion in 7.5% (v/v) acetic acid, 0.05% (w/v) SDS, 92.5% (v/v) water for 30 mins. The priming solution was then drained, and the gels were stained by complete immersion for 4 hours in a staining solution of Pyridinium, 4-[2-[4-(dipentylamino)-2-trifluoromethylphenyl] ethenyl]-1-(sulfobutyl)-, inner salt, prepared by diluting a stock solution of this dye (2mg/ml in DMSO) in 7.5% (v/v) aqueous acetic acid to give a final concentration of 1.2 mg/l; the staining solution was vacuum filtered through a 0.4µm filter (Duropore) before use.

6.1.7 *Imaging of the gel*

A computer-readable output was produced by imaging the fluorescently stained gels with the Apollo 2 scanner (Oxford Glycosciences, Oxford, UK) described in section 5.1, *supra*. This scanner has a gel carrier with four integral fluorescent markers (Designated M1, M2, M3, M4) that are used to correct the image geometry and are a quality control feature to confirm that the scanning has been performed correctly.

For scanning, the gels were removed from the stain, rinsed with water and allowed to air dry briefly, and imaged on the Apollo 2. After imaging, the gels were sealed in polyethylene bags containing a small volume of staining solution, and then stored at 4°C.

6.1.8 *Digital Analysis of the Data*

The data were processed as described in U.S. Application Serial No. 08/980,574, (published as WO 98/23950) at Sections 5.4 and 5.5 (incorporated herein by reference), as set forth more particularly below.

The output from the scanner was first processed using the MELANIE® II 2D PAGE analysis program (Release 2.2, 1997, BioRad Laboratories, Hercules, California, Cat. # 170-7566) to autodetect the registration points, M1, M2, M3 and M4; to autocrop the images (*i.e.*, to eliminate signals originating from areas of the scanned image lying outside the boundaries of the gel, *e.g.* the reference frame); to filter out artifacts due to dust; to detect and quantify features; and to create image files in GIF format. Features were detected using the following parameters:

Smooths = 2

Laplacian threshold 50

Partials threshold 1

Saturation = 100

Peakedness = 0

Minimum Perimeter = 10

6.1.9 Assignment of pI and MW Values

- 5 Landmark identification was used to determine the pI and MW of features detected in the images. Twelve landmark features, designated CSF1 to CSF12, were identified in a standard CSF image obtained from a pooled sample. These landmark features are identified in Figure 1 and were assigned the pI and/or MW values identified in Table XI.

10 Table X. Landmark Features Used In This Study

Name	pI	MW (Da)	Name	pI	MW (Da)
CSF1	5.96	185230	CSF7	4.78	41340
CSF2	5.39	141700	CSF8	9.2	40000
CSF3	6.29	100730	CSF9	5.5	31900
CSF4	5.06	71270	CSF10	6.94	27440
CSF5	7.68	68370	CSF11	5.9	23990
CSF6	5.67	48090	CSF12	6.43	10960

- As many of these landmarks as possible were identified in each gel image of the dataset. Each feature in the study gels was then assigned a pI value by linear interpolation or extrapolation (using the MELANIE®-II software) to the two nearest landmarks, and was
 15 assigned a MW value by linear interpolation or extrapolation (using the MELANIE®-II software) to the two nearest landmarks.

6.1.10 Matching With Primary Master Image

- 20 Images were edited to remove gross artifacts such as dust, to reject images which had gross abnormalities such as smearing of protein features, or were of too low a loading or

overall image intensity to allow identification of more than the most intense features, or were of too poor a resolution to allow accurate detection of features. Images were then compared by pairing with one common image from the whole sample set. This common image, the "primary master image", was selected on the basis of protein load (maximum
5 load consistent with maximum feature detection), a well resolved myoglobin region, (myoglobin was used as an internal standard), and general image quality. Additionally, the primary master image was chosen to be an image which appeared to be generally representative of all those to be included in the analysis. (This process by which a primary master gel was judged to be representative of the study gels was rechecked by the method
10 described below and in the event that the primary master gel was seen to be unrepresentative, it was rejected and the process repeated until a representative primary master gel was found.)

Each of the remaining study gel images was individually matched to the primary master image such that common protein features were paired between the primary master
15 image and each individual study gel image as described below.

6.1.11 Cross-matching Between Samples

To facilitate statistical analysis of large numbers of samples for purposes of identifying features that are differentially expressed, the geometry of each study gel was
20 adjusted for maximum alignment between its pattern of protein features, and that of the primary master, as follows. Each of the study gel images was individually transformed into the geometry of the primary master image using a multi-resolution warping procedure. This procedure corrects the image geometry for the distortions brought about by small changes in the physical parameters of the electrophoresis separation process from one sample to
25 another. The observed changes are such that the distortions found are not simple geometric distortions, but rather a smooth flow, with variations at both local and global scale.

The fundamental principle in multi-resolution modeling is that smooth signals may be modeled as an evolution through 'scale space', in which details at successively finer scales are added to a low resolution approximation to obtain the high resolution signal. This
30 type of model is applied to the flow field of vectors (defined at each pixel position on the reference image) and allows flows of arbitrary smoothness to be modeled with relatively few degrees of freedom. Each image is first reduced to a stack, or pyramid, of images derived from the initial image, but smoothed and reduced in resolution by a factor of 2 in

each direction at every level (Gaussian pyramid) and a corresponding difference image is also computed at each level, representing the difference between the smoothed image and its progenitor (Laplacian pyramid). Thus the Laplacian images represent the details in the image at different scales.

5 To estimate the distortion between any 2 given images, a calculation was performed at level 7 in the pyramid (*i.e.* after 7 successive reductions in resolution). The Laplacian images were segmented into a grid of 16x16 pixels, with 50% overlap between adjacent grid positions in both directions, and the cross correlation between corresponding grid squares on the reference and the test images was computed. The distortion displacement was then
10 given by the location of the maximum in the correlation matrix. After all displacements had been calculated at a particular level, they were interpolated to the next level in the pyramid, applied to the test image, and then further corrections to the displacements were calculated at the next scale.

The warping process brought about good alignment between the common features in
15 the primary master image, and the images for the other samples. The MELANIE® II 2D PAGE analysis program was used to calculate and record approximately 500-700 matched feature pairs between the primary master and each of the other images. The accuracy of this program was significantly enhanced by the alignment of the images in the manner described above. To improve accuracy still further, all pairings were finally examined by
20 eye in the MelView interactive editing program and residual recognizably incorrect pairings were removed. Where the number of such recognizably incorrect pairings exceeded the overall reproducibility of the Preferred Technology (as measured by repeat analysis of the same biological sample) the gel selected to be the primary master gel was judged to be insufficiently representative of the study gels to serve as a primary master gel. In that case,
25 the gel chosen as the primary master gel was rejected, and different gel was selected as the primary master gel, and the process was repeated.

All the images were then added together to create a composite master image, and the positions and shapes of all the gel features of all the component images were super-imposed onto this composite master as described below.

30 Once all the initial pairs had been computed, corrected and saved, a second pass was performed whereby the original (unwarped) images were transformed a second time to the geometry of the primary master, this time using a flow field computed by smooth interpolation of the multiple tie-points defined by the centroids of the paired gel features. A

composite master image was thus generated by initialising the primary master image with its feature descriptors. As each image was transformed into the primary master geometry, it was digitally summed pixel by pixel into the composite master image, and the features that had not been paired by the procedure outlined above were likewise added to the composite master image description, with their centroids adjusted to the master geometry using the flow field correction.

The final stage of processing was applied to the composite master image and its feature descriptors, which now represent all the features from all the images in the study transformed to a common geometry. The features were grouped together into linked sets or "clusters", according to the degree of overlap between them. Each cluster was then given a unique identifying index, the molecular cluster index (MCI).

An MCI identifies a set of matched features on different images. Thus an MCI represents a protein or proteins eluting at equivalent positions in the 2D separation in different samples.

6.1.12. Construction of Profiles

After matching all component gels in the study to the final composite master image, the intensity of each feature was measured and stored. The end result of this analysis was the generation of a digital profile which contained, for each identified feature: 1) a unique identification code relative to corresponding feature within the composite master image (MCI), 2) the x, y coordinates of the features within the gel, 3) the isoelectric point (pI) of the AFs, 4) the apparent molecular weight (MW) of the AFs, 5) the signal value, 6) the standard deviation for each of the preceding measurements, and 7) a method of linking the MCI of each feature to the master gel to which this feature was matched. By virtue of a Laboratory Information Management System (LIMS), this MCI profile was traceable to the actual stored gel from which it was generated, so that proteins identified by computer analysis of gel profile databases could be retrieved. The LIMS also permitted the profile to be traced back to an original sample or patient.

6.1.13. Differential Analysis of the Profiles

For the pooled gel data within each sample set (Alzheimer's CSF and normal CSF), the profiles were analyzed to identify and select those features differentially present in the profiles. These selected features were then assembled into an Alzheimer's pooled gel feature

set. Matching features of each feature set were then compared to identify those features showing at least a 2-fold difference in mean intensity between Alzheimer's CSF and normal CSF. Differentially present features were identified as Alzheimer's Disease Associated Features (AFs).

5

6.1.14. Statistical Analysis of the Profiles

The MCI data was represented in statistical models in two forms: 1) percent of total protein volume for a given gel (PCTVOL) and 2) absolute volume, scaled by the total volume loaded on the gel (VOL). A value of 0 was entered for PCTVOL and VOL if an MCI did not appear on a particular gel. For most analyzes, in order for an MCI to be considered in the statistical model, it had to have non-zero values for PCTVOL and VOL in at least 75% of gels in at least one of the diagnosis groups in the analysis (described below).

The complementary statistical strategies specified below were used to identify AFs from the MCIs within the mastergroup.

(I) Group Analysis

The purpose of these analyses was to characterize differences among gels from individuals with different clinical diagnoses. The diagnosis groups were 1) autopsy-confirmed (AD) vs. normal controls (NCO) at their first sample, 2) Dementia Alzheimer's type (DAT) with an initial sample within 3 years of disease onset vs. NCO, and 3) last sample of first-degree relatives of individuals diagnosed with dementia of Alzheimer's type without a clinical diagnosis of dementia (NCF) vs. NCO.

The following statistical techniques were used in the group analyses:

(1) Linear model

A linear model controlling for age and gender that compared a DAT group vs the NCO group with regard to the rank of the volume.

(2) Classification trees

Classification trees were used with the MCI volumes as predictors, and clinical diagnosis as the response. The algorithm looks for 'split points' in the predictors that partition the data into homogeneous sets according to the response variable. After evaluating all possible splits for a given node of the tree, the split is chosen that maximizes the change in deviance according to a multinomial likelihood model. Tree models were fit to both the original data and data from bootstrap samples of the original data (sampling with replacement). The

statistical test involved whether a given MCI proved to be an important 'split point' to determine diagnosis, either in the original data tree or a bootstrap sample tree.

(3) Logistic regression model

5 A logistic regression model was used to model the probability of being AD. The volumes of the various MCI's were used as the explanatory variables. A stepwise procedure was used to select 5 MCI's.

Criteria for inclusion based on the group analyses:

Information from all of the above described analyses were used to select MCI's that:

- 10
1. Were among the 5-6 MCI's with the smallest p-value for a given analysis
 2. Appeared in the smallest 100 p-values for 2 or more analysis
 3. Appeared as an important split-point in a classification tree
 4. Had desired distributional properties

15 (4) Nearest Neighbor Analysis

In addition to the techniques described above, the gels were examined with nearest neighbor analysis. In this analysis, the gels were visually examined and MCIs located proximate to other MCIs that demonstrated a volume or other characteristics that are indicative of disease, and that themselves demonstrated a similar or like appearance and intensity, were noted and categorized as of like significance. These MCIs were thus included in the group for further examination and analysis in relation to their role in the disease.

20

(II) Longitudinal Analysis

25 The purpose of the longitudinal analyses was to identify AF's associated with changes in disease state as measured by the MMSEM, a combination of the MMSE, CDR, and GDS assessment measures. DAT subjects with two or more samples were used in these analyses.

There were two models employed in the longitudinal analyses. In the first, the goal was to identify AF's for which changes in volume were significantly correlated with changes in the MMSEM. For each AF, MMSEM was regressed on the rank of the volume after controlling for age and subject. AF's with p-values less than 0.05, in the top 100 of

30

any of the group analyses, and consistent with the group analyses in terms of up or down regulation were included.

The goal of the second model was to identify AF's for which volume in a subject's first sample was a significant predictor of disease progression rate during the period following the time of the first sample. First, a simple linear regression model was used to estimate a progression rate based on the MMSEM for each subject. Only subjects with an initial MMSEM greater than or equal to 12 and with greater than four months between the first and last samples were used. In addition, only samples within the first three years of the first sample were used. Regression modeling and split-sample validation were then used to identify significant AF's. More specifically, subjects were first randomly divided into two groups. For each group, stepwise weighted least-squares (WLS) regression using the rank of volume from each subject's first sample was used to select the five best AF's for predicting progression rate. If an AF was in the top five in one group and yielded a slope estimate with the same sign when included in the other group, it was included. In addition, the top five AF's from a stepwise WLS on both groups combined were included.

6.1.15 Recovery and analysis of selected proteins

Proteins in AFs were robotically excised and processed to generate tryptic digest peptides. Tryptic peptides were analyzed by mass spectrometry using a PerSeptive Biosystems Voyager-DETM STR Matrix-Assisted Laser Desorption Ionization Time-of-Flight (MALDI-TOF) mass spectrometer, and selected tryptic peptides were analyzed by tandem mass spectrometry (MS/MS) using a Micromass Quadrupole Time-of-Flight (Q-TOF) mass spectrometer (Micromass, Altrincham, U.K.) equipped with a nanoflowTM electrospray Z-spray source. For partial amino acid sequencing and identification of APIs uninterpreted tandem mass spectra of tryptic peptides were searched using the SEQUEST search program (Eng et al., 1994, J. Am. Soc. Mass Spectrom. 5:976-989), version v.C.1. Criteria for database identification included: the cleavage specificity of trypsin; the detection of a suite of a, b and y ions in peptides returned from the database, and a mass increment for all Cys residues to account for carbamidomethylation. The database searched was database constructed of protein entries in the non-redundant database held by the National Centre for Biotechnology Information (NCBI) which is accessible at <http://www.ncbi.nlm.nih.gov/>. Following identification of proteins through spectral-spectral correlation using the SEQUEST program, masses detected in MALDI-TOF mass spectra

were assigned to tryptic digest peptides within the proteins identified. In cases where no proteins could be identified through searching with uninterpreted MS/MS spectra of tryptic digest peptides using the SEQUEST program, tandem mass spectra of the peptides were interpreted manually, using methods known in the art. (In the case of interpretation of low-energy fragmentation mass spectra of peptide ions see Gaskell et al., 1992, Rapid Commun. Mass Spectrom. 6:658-662). The method described in PCT Application No. PCT/GB01/04034, which is incorporated herein by reference in its entirety, was also used to interpret mass spectra.

6.2 RESULTS

These initial experiments identified 111 features that were decreased and 30 features that were increased in AD CSF as compared with normal CSF. Details of these AFs are provided in Tables I (a) and II (a). Each AF was differentially present in AD CSF as compared with normal CSF. For some preferred AFs (AF-200, AF-201, AF-203, AF-204, AF-205, AF-206, AF-207, AF-208, AF-209, AF-210, AF-211, AF-212, AF-213, AF-214, AF-215, AF-216, AF-217, AF-218, AF-219, AF-220, AF-221, AF-222, AF-223, AF-224, AF-225, AF-226, AF-227, AF-227, AF-229, AF-230, AF-231, AF-232, AF-233, AF-234, AF-235, AF-236, AF-237, AF-238, AF-239, AF-240, AF-241, AF-242, AF-243, AF-244, AF-245, AF-248, AF-249, AF-251, AF-252, AF-253, AF-258, AF-262, AF-263, AF-278, AF-287, AF-303, AF-310, AF-311, AF-313, AF-314, AF-316, AF-317, AF-318, AF-325) the difference was highly significant ($p < 0.01$), and for certain highly preferred AFs (AF-200, AF-201, AF-203, AF-204, AF-205, AF-206, AF-207, AF-208, AF-209, AF-210, AF-211, AF-212, AF-213, AF-214, AF-216, AF-217, AF-218, AF-219, AF-220, AF-221, AF-222, AF-223, AF-224, AF-225, AF-226, AF-227, AF-227, AF-229, AF-231, AF-232, AF-233, AF-234, AF-235, AF-236, AF-237, AF-238, AF-240, AF-241, AF-242, AF-243, AF-245, AF-253, AF-258, AF-262, AF-287, AF-303, AF-311, AF-316), the difference was still more significant ($p < 0.001$).

Partial amino acid sequences were determined for the differentially present APIs in these AFs. Details of these APIs are provided in Tables IV and V. Computer searches of public databases identified at least one API for which neither the partial amino acid sequence, nor any oligonucleotide encoding such a peptide sequence, was described in any public database examined.

7. EXAMPLE: DIAGNOSIS AND TREATMENT OF AD

The following example illustrate the use of an API of the invention for screening, treatment or diagnosis of AD. The following example also illustrates the use of modulators (e.g., agonist or antagonists) of an API of the invention to treat or prevent AD.

Pigment epithelium-derived factor (PEDF) is a neurotrophic protein synthesized and secreted by retinal pigment epithelial cells in early embryogenesis and has been shown to be present in the extracellular matrix between the RPE cells and the neural retina. It induces neuronal differentiation and promotes survival of neurons of the central nervous system from degeneration caused by serum withdrawal or glutamate cytotoxicity. PEDF has been shown to protect immature but not mature cerebellar cells from apoptotic death, acting as a survival factor for such cells, as well as protecting them against glutamate and hydrogen peroxide toxicity. PEDF binds to glycosaminoglycans and to an 80 kDa receptor present on the surface of retinoblastoma and cerebellar granule cells. PEDF binding to the 80 kDa receptor, as well as PEDF activity, may be blocked by antibodies recognizing PEDF, and by a 44 amino acid fragment (amino acids 78-121) of PEDF.

The expression of an isoform of PEDF with a molecular weight of 33,401 kDa and pI of 6.74 has been shown herein to be significantly increased in the cerebrospinal fluid (CSF) of subjects having AD as compared with the CSF of subjects free from AD (AF-218/API-313, see Tables I(a) and IV(a)). Thus, quantitative detection of PEDF in CSF can be used to diagnose AD, determine the progression of AD or monitor the effectiveness of a therapy for AD.

In one embodiment of the invention, compounds that modulate (*i.e.*, upregulate or downregulate) the expression, activity or both the expression and activity of PEDF are administered to a subject in need of treatment or for prophylaxis of AD. Antibodies that modulate the expression, activity or both the expression and activity of PEDF are suitable for this purpose. In addition, nucleic acids coding for all or a portion of PEDF, or nucleic acids complementary to all or a portion of PEDF, may be administered. PEDF, or fragments of the PEDF polypeptide may also be administered.

The invention also provides screening assays to identify additional compounds that modulate the expression of PEDF or activity of PEDF. Compounds that modulate the expression of PEDF *in vitro* can be identified by comparing the expression of PEDF in cells treated with a test compound to the expression of PEDF in cells treated with a control

compound (*e.g.*, saline). Methods for detecting expression of PEDF are known in the art and include measuring the level of PEDF RNA (*e.g.*, by northern blot analysis or RT-PCR) and measuring PEDF protein (*e.g.*, by immunoassay or western blot analysis). Compounds that modulate the activity of PEDF can be identified by comparing the ability of a test
5 compound to agonize or antagonize a function of PEDF, such as its neurotrophic activity or its binding to the 80 kDa receptor, to the ability of a control compound (*e.g.*, saline) to inhibit the same function of PEDF. Compounds capable of modulating PEDF binding to its receptor or PEDF activity are identified as compounds suitable for further development as a compound useful for the treatment of AD.

10 Binding between PEDF and its receptor can be determined by, for example, contacting PEDF with cells known to express the PEDF receptor and assaying the extent of binding between PEDF and the cell surface receptor, or by contacting PEDF with its receptor in a cell-free assay, *i.e.*, an assay where the PEDF and PEDF receptor are isolated, and, preferably, recombinantly produced, and assaying the extent of binding between PEDF
15 and its receptor. Through the use of such assays, candidate compounds may be tested for their ability to agonize or antagonize the binding of PEDF to its receptor.

Compounds identified *in vitro* that affect the expression or activity of PEDF can be tested *in vivo* in animal models of AD or Downs syndrome, or in subjects having AD, to determine their therapeutic efficacy.

20 The present invention is not to be limited in terms of the particular embodiments described in this application, which are intended as single illustrations of individual aspects of the invention. Functionally equivalent methods and apparatus within the scope of the invention, in addition to those enumerated herein, will be apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications and
25 variations are intended to fall within the scope of the appended claims. The contents of each reference, patent and patent application cited in this application is hereby incorporated by reference in its entirety.

WE CLAIM:

1. A method for screening or diagnosis of Alzheimer's disease in a subject, for determining the stage or severity of Alzheimer's disease in a subject, for identifying a subject at risk of developing Alzheimer's disease, or for monitoring the effect of therapy administered to a subject having Alzheimer's disease, said method comprising:
 - (a) analysing a test sample of body fluid from the subject by two-dimensional electrophoresis to generate a two-dimensional array of features, said array comprising one or more of the following Alzheimer's disease-Associated Features (AFs): AF-200, AF-201, AF-202, AF-203, AF-204, AF-205, AF-206, AF-207, AF-208, AF-209, AF-210, AF-211, AF-212, AF-213, AF-214, AF-215, AF-216, AF-217, AF-218, AF-219, AF-220, AF-221, AF-222, AF-223, AF-224, AF-225, AF-226, AF-227, AF-228, AF-229, AF-230, AF-231, AF-232, AF-233, AF-234, AF-235, AF-236, AF-237, AF-238, AF-239, AF-240, AF-241, AF-242, AF-243, AF-244, AF-245, AF-246, AF-247, AF-248, AF-249, AF-250, AF-251, AF-252, AF-253, AF-254, AF-255, AF-256, AF-257, AF-258, AF-259, AF-260, AF-261, AF-262, AF-263, AF-264, AF-265, AF-266, AF-267, AF-268, AF-269, AF-270, AF-271, AF-272, AF-273, AF-274, AF-275, AF-276, AF-277, AF-278, AF-279, AF-280, AF-281, AF-282, AF-283, AF-284, AF-285, AF-286, AF-287, AF-288, AF-289, AF-290, AF-291, AF-292, AF-293, AF-294, AF-295, AF-296, AF-297, AF-298, AF-299, AF-300, AF-301, AF-302, AF-303, AF-304, AF-305, AF-306, AF-307, AF-308, AF-309, AF-310, AF-311, AF-312, AF-313, AF-314, AF-315, AF-316, AF-317, AF-318, AF-319, AF-320, AF-321, AF-322, AF-323, AF-324, AF-325, AF-326, AF-327, AF-328, AF-329, AF-330, AF-331, AF-332, AF-333, AF-334, AF-335, AF-336, AF-337, AF-338, AF-339, ; optionally in combination with one or more of the following AFs: , AF-1, AF-2, AF-3, AF-4, AF-5, AF-6, AF-7, AF-8, AF-9, AF-10, AF-13, AF-14, AF-15, AF-16, AF-17, AF-18, AF-19, AF-20, AF-21, AF-22, AF-23, AF-24, AF-25, AF-26, AF-27, AF-28, AF-29, AF-30, AF-31, AF-32, AF-33, AF-34, AF-35, AF-36, AF-37, AF-38, AF-39, AF-40, AF-41, AF-42, AF-43, AF-44, AF-45, AF-46, AF-47, AF-48, AF-49, AF-50, AF-51, AF-76, AF-149, AF-150, AF-152, AF-154, AF-155, AF-156, AF-159, AF-160, AF-162, AF-163, AF-164, AF-169, AF-170, AF-172, AF-173, AF-174, AF-175, AF-176, AF-177, AF-178, AF-181, AF-183, AF-184, AF-186, AF-187, AF-188, AF-189, AF-190, AF-191, AF-78, AF-79, AF-80, AF-81, AF-82, AF-83, AF-84, AF-85, AF-86, AF-87, AF-88, AF-89, AF-90, AF-91, AF-92, AF-93, AF-94, AF-95, AF-96, AF-98, AF-99, AF-100, AF-101, AF-102, AF-103, AF-104, AF-105, AF-107, AF-108, AF-110, AF-111, AF-112, AF-114, AF-115, AF-116, AF-

117, AF-118, AF-119, AF-52, AF-53, AF-54, AF-55, AF-56, AF-57, AF-58, AF-59, AF-60, AF-61, AF-62, AF-63, AF-64, AF-65, AF-66, AF-67, AF-68, AF-69, AF-70, AF-71, AF-72, AF-73, AF-74, AF-75, AF-77, AF-151, AF-153, AF-157, AF-161, AF-165, AF-166, AF-167, AF-168, AF-171, AF-179, AF-180, AF-182, AF-185, AF-192, AF-121, AF-122, AF-123, AF-124, AF-125, AF-126, AF-127, AF-128, AF-129, AF-130, AF-131, AF-132, AF-133, AF-134, AF-137, AF-139, AF-140, AF-141, AF-142, AF-143, AF-144, AF-145, AF-146, AF-147, AF-148; and

(b) comparing the abundance of the one or more DAFs in the test sample with the abundance of the one or more DAFs in a body fluid sample from one or more subjects free from Alzheimer's disease, or with a previously determined reference range for that feature in subjects free from Alzheimer's disease, or with the abundance at least one Expression Reference Feature (ERF) in the test sample.

2. A method for screening or diagnosis of Alzheimer's disease in a subject, for determining the stage or severity of Alzheimer's disease in a subject, for identifying a subject at risk of developing Alzheimer's disease, or for monitoring the effect of therapy administered to a subject having Alzheimer's disease, said method comprising quantitatively detecting, in a test sample of body fluid from the subject, one or more of the following Alzheimer's disease-Associated Protein Isoforms (APIs): API-375, API-300, API-301, API-302, API-303, API-304, API-305, API-306, API-307, API-308, API-309, API-310, API-376, API-377, API-311, API-312, API-313, API-378, API-314, API-379, API-315, API-316, API-380, API-317, API-369, API-318, API-319, API-320, API-321, API-322, API-323, API-381, API-382, API-383, API-384, API-324, API-325, API-326, API-327, API-328, API-329, API-330, API-331, API-385, API-332, API-333, API-334, API-335, API-336, API-337, API-338, API-339, API-340, API-341, API-342, API-386, API-387, API-343, API-344, API-388, API-345, API-346, API-389, API-390, API-347, API-391, API-348, API-349, API-350, API-351, API-352, API-392, API-393, API-394, API-395, API-353, API-354, API-396, API-397, API-355, API-398, API-399, API-400, API-356, API-357, API-401, API-358, API-402, API-403, API-359, API-360, API-404, API-405, API-406, API-407, API-408, API-409, API-361, API-362, API-363, API-410, API-411, API-364, API-412, API-413, API-365, API-366, API-414, API-415, API-367, API-370, API-416, API-417, API-418, API-368, XXXXXXXXXX; optionally combined with one or more of the following APIs: API-47, API-242, API-1, API-48, API-49, API-2, API-194,

API-3, API-50, API-51, API-4, API-52, API-243, API-53, API-244, API-54, API-5, API-55, API-245, API-6, API-56, API-57, API-7, API-8, API-9, API-10, API-14, API-15, API-58, API-16, API-59, API-196, API-17, API-60, API-18, API-61, API-62, API-19, API-63, API-64, API-65, API-20, API-22, API-66, API-67, API-68, API-69, API-70, API-23, API-24, API-197, API-198, API-25, API-71, API-26, API-27, API-72, API-73, API-199, API-200, API-28, API-30, API-86, API-201, API-88, API-202, API-89, API-90, API-91, API-92, API-93, API-95, API-97, API-98, API-99, API-101, API-102, API-103, API-104, API-107, API-210, API-108, API-111, API-112, API-113, API-114, API-214, API-144, API-146, API-147, API-148, API-150, API-151, API-152, API-215, API-153, API-158, API-159, API-160, API-165, API-166, API-167, API-173, API-174, API-175, API-176, API-179, API-180, API-181, API-182, API-183, API-184, API-185, API-217, API-219, API-187, API-189, API-190, API-238, API-239, API-240, API-74, API-33, API-221, API-34, API-75, API-246, API-35, API-76, API-222, API-77, API-36, API-37, API-78, API-38, API-79, API-80, API-81, API-223, API-82, API-83, API-39, API-84, API-85, API-40, API-247, API-41, API-224, API-42, API-43, API-44, API-45, API-248, API-46, API-225, API-116, API-118, API-119, API-120, API-121, API-122, API-123, API-124, API-125, API-126, API-127, API-128, API-130, API-131, API-132, API-134, API-135, API-232, API-233, API-234, API-136, API-137, API-138, API-139, API-140, API-141, API-142, API-143, API-145, API-149, API-155, API-161, API-162, API-163, API-168, API-169, API-170, API-171, API-237, API-172, API-177, API-178, API-186, API-220, API-188, API-191, API-192

3. The method according to claim 1 or 2 where the body fluid is Cerebrospinal Fluid.
4. The method according to claim 2 or 3 where the abundance of the one or more APIs in the test sample is compared with the abundance of the one or more APIs in a sample from one or more subjects free from Alzheimer's disease, or with a previously determined reference range for that feature in subjects free from Alzheimer's disease, or with the abundance at least one Expression Reference Feature (ERF) in the test sample.
5. The method according to any one of claims 2 to 4, wherein the step of quantitatively detecting comprises testing at least one aliquot of the first sample, said step of testing comprising:

- (a) contacting the aliquot with an antibody that is immunospecific for a API;
- (b) quantitatively measuring the binding of the antibody and the API; and
- (c) comparing the results of step (b) with a predetermined reference range.

5 6. The method according to claim 5, wherein the step of quantitatively detecting comprises testing a plurality of aliquots with a plurality of antibodies cognate for a plurality of preselected APIs.

7. A pharmaceutical composition comprising an Alzheimer's disease-Associated
10 Protein Isoform (API) as defined in claim 2, or a nucleic acid encoding an API, and a pharmaceutically acceptable carrier.

8. The pharmaceutical composition according to claim 7, wherein the Alzheimer's disease-Associated Protein Isoform (API) is in recombinant form.

15

9. An antibody capable of immunospecific binding to a Alzheimer's disease-Associated Protein Isoform (API) as defined in claim 2.

10. The method according to claim 5 or 6 or an antibody according to claim 9, wherein
20 the antibody is a monoclonal antibody.

11. The method according to claim 5 or 6 or an antibody according to claim 9 or 10, wherein the antibody is a chimeric antibody.

25 12. The method according to claim 5 or 6 or an antibody according to claim 9 or 10, wherein the antibody is a bispecific antibody.

13. The method according to claim 5 or 6 or an antibody according to claim 9 or 10, wherein the antibody is a humanised antibody.

30

14. The method according to claim 5 or 6 or an antibody according to any one of claims 9 to 13, wherein the antibody binds to the API with greater affinity than to another isoform of the API.

15. A kit comprising one or more antibodies as claimed in any one of claims 9 to 14 and/or one or more APIs as defined in claim 2, other reagents and instructions for use.
- 5 16. The kit of claim 15 for use in the screening or diagnosis of Alzheimer's disease in a subject, for determining the stage or severity of Alzheimer's disease in a subject, for identifying a subject at risk of developing Alzheimer's disease, or for monitoring the effect of therapy administered to a subject having Alzheimer's disease.
- 10 17. The kit according to claim 15 or 16 comprising a plurality of antibodies as claimed in any one of claims 9 to 14 and/or a plurality of APIs as defined in claim 2.
18. A pharmaceutical composition comprising a therapeutically effective amount of an antibody, or a fragment or derivative of an antibody according to any one of claims 9 to 14
15 and a pharmaceutically acceptable carrier.
19. A method of treating or preventing Alzheimer's disease comprising administering to a subject in need of such treatment a therapeutically effective amount of an antibody as claimed in any one of claims 9 to 14.
20
20. A method of treating or preventing Alzheimer's disease comprising administering to a subject in need of such treatment or prevention a therapeutically effective amount of one or more of the Alzheimer's disease-Associated Protein Isoforms (APIs) as defined in claim 2 and/or a nucleic acid encoding said APIs.
25
21. A method of treating or preventing Alzheimer's disease comprising administering to a subject in need of such treatment or prevention a therapeutically effective amount of a nucleic acid that inhibits the function of one or more of the Alzheimer's disease-Associated Protein Isoforms (APIs) as defined in claim 2.
30
22. The method according to claim 21, wherein the nucleic acid is a API antisense nucleic acid or ribozyme.

23. A method of screening for agents that interact with one or more a Alzheimer's disease-Associated Protein Isoforms (APIs) as defined in claim 2, fragments of APIs (API fragment), polypeptides related to APIs (API-related polypeptide), or API-fusion proteins said method comprising:

- 5 (a) contacting an API, a fragment of an API, an API -related polypeptide, or an API -fusion protein with a candidate agent; and
- (b) determining whether or not the candidate agent interacts with the API, the API fragment, the API -related polypeptide, or the API -fusion protein.

10 24. The method according to claim 23, wherein the determination of interaction between the candidate agent and the API, API fragment, API -related polypeptide or API -fusion protein comprises quantitatively detecting binding of the candidate agent and the API, API fragment, API -related polypeptide or API -fusion protein.

15 25. A method of screening for or identifying agents that modulate the expression or activity of one or more Alzheimer's disease-Associated Protein Isoforms (APIs) as defined in claim 2, fragments of API (API fragment), polypeptides related to APIs (API -related polypeptide) or API -fusion proteins comprising:

- (a) contacting a first population of cells expressing the API, API fragment, API-
20 related polypeptide, or API-fusion protein with a candidate agent;
- (b) contacting a second population of cells expressing said API, API fragment, API-related polypeptide, or API-fusion protein with a control agent; and
- (c) comparing the level of said API, API fragment, API-related polypeptide, or
25 API- fusion protein or mRNA encoding said API, API fragment, API-
 related polypeptide, or API-fusion protein in the first and second populations of cells, or comparing the level of induction of a downstream effector in the first and second populations of cells.

26. A method of screening for or identifying agents that modulate the expression or
30 activity of one or more Alzheimer's disease-Associated Protein Isoforms (APIs) as defined in claim 2, fragments of APIs (API fragment), polypeptides related to APIs (API-related polypeptide) or API-fusion proteins said method comprising:

- (a) administering a candidate agent to a first mammal or group of mammals;

- (b) administering a control agent to a second mammal or group of mammals;
and
(c) comparing the level of expression of the API, API fragment, API-related polypeptide or API-fusion protein, or mRNA encoding said API, API fragment, API-related polypeptide or API-fusion protein in the first and second groups, or comparing the level of induction of a downstream effector in the first and second groups.

27. The method as claimed in claim 26, wherein the mammals are animal models for Alzheimer's disease.
28. The method according to any one of claims 25 to 27, wherein administration of a candidate agent results in an increase in the level of said API, API fragment, API-related polypeptide or API-fusion protein, or mRNA encoding said API, API fragment, API-related polypeptide, or API-fusion protein, or said downstream effector in the first population of cells or mammals compared to the second population of cells or mammals.
29. The method according to any one of claims 25 to 27, wherein administration of a candidate agent results in a decrease in the level of said API, API fragment, API-related polypeptide, or API-fusion protein, or mRNA encoding said API, API fragment, API-related polypeptide, or API-fusion protein, or said downstream effector in the first population of cells or mammals compared to the second population of cells or mammals.
30. The method as claimed in claim 25 or 27, wherein the levels of said API, API fragment, API-related polypeptide, or API-fusion protein, or mRNA encoding said API, API fragment, API-related polypeptide, or API-fusion protein, or of said downstream effector in the first and second groups are further compared to the level of said API, API fragment, API-related polypeptide or API fusion protein, or mRNA encoding said API, API fragment, API-related polypeptide or API fusion protein in normal control mammals.
31. The method according to claim 30, wherein said mammals are human subjects having Alzheimer's disease.

32. A method of screening for or identifying agents that modulate the activity of one or more of the Alzheimer's disease-Associated Proteins Isoforms (APIs) as defined in claim 2, fragments of APIs (API fragment), polypeptides related to APIs (API-related polypeptide) or API-fusion proteins said method comprising:

- 5 (a) in a first aliquot, contacting a candidate agent with the API, API fragment, API-related polypeptide or API fusion protein, and
- (b) determining and comparing the activity of the API, API fragment, API-related polypeptide or API fusion protein in the first aliquot after addition of the candidate agent with the activity of the API, API fragment, API-related
- 10 polypeptide or API fusion protein in a control aliquot, or with a previously determined reference range.

33. The method according to any one of claims 20 or 23 to 32, wherein the API, API fragment, API-related polypeptide, or API fusion protein is a recombinant protein.

15

34. The method according to any one of claims 23, 24 or 32, wherein the API, API fragment, API-related polypeptide or API fusion protein is immobilized on a solid phase.

35. A method for screening or diagnosis of Alzheimer's disease in a subject or for monitoring the effect of an anti-Alzheimer's disease drug or therapy administered to a subject, comprising:

- 20 (a) contacting at least one oligonucleotide probe comprising 10 or more consecutive nucleotides complementary to a nucleotide sequence encoding an API as defined in claim 2 with RNA obtained from a biological sample
- 25 from the subject or with cDNA copied from the RNA wherein said contacting occurs under conditions that permit hybridization of the probe to the nucleotide sequence if present;
- (b) detecting hybridization, if any, between the probe and the nucleotide sequence; and
- 30 (c) comparing the hybridization, if any, detected in step (b) with the hybridization detected in a control sample, or with a previously determined reference range.

36. The method as claimed in claim 35, wherein step (a) includes the step of hybridizing the nucleotide sequence to a DNA array, wherein one or more members of the array are the probes complementary to a plurality of nucleotide sequences encoding distinct APIs.

5 37. A method of modulating the activity of one or more of the Alzheimer's disease-Associated Protein Isoforms as defined in claim 2 comprising administering to a subject an agent identified by any one of claims 23 to 34.

38. A method of treating or preventing Alzheimer's disease comprising administering to
10 a subject in need of such treatment or prevention a therapeutically effective dose of an agent that modulates the activity of one or more of the Alzheimer's disease-Associated Protein Isoforms as defined in claim 2; whereby the symptoms of Alzheimer's disease are ameliorated.

15 39. A method for identifying targets for therapeutic modulation of Alzheimer's disease wherein the activity of one or more of the Alzheimer's disease-Associated Protein Isoforms as defined in claim 2 is utilized as a measure to determine whether a candidate target is effective for modulation of Alzheimer's disease.

20

FIGURE 1




FIGURE 2

