



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C12N 9/12, 15/54, 1/21, 5/10, 15/00, 15/63, C07K 16/40, A61K 38/45, C12Q 1/68, 1/48, C07K 14/00</p>	A1	<p>(11) International Publication Number: WO 99/09146</p> <p>(43) International Publication Date: 25 February 1999 (25.02.99)</p>
<p>(21) International Application Number: PCT/US98/17248</p> <p>(22) International Filing Date: 20 August 1998 (20.08.98)</p> <p>(30) Priority Data: 60/056,112 20 August 1997 (20.08.97) US</p> <p>(71) Applicant: SMITHKLINE BEECHAM CORPORATION [US/US]; One Franklin Plaza, Philadelphia, PA 19103 (US).</p> <p>(72) Inventors: ANDERSON, Karen, M.; 319 Joy Lane, West Chester, PA 19380 (US). JACKSON, Jeffrey, R.; 134 Spruce Lane, Collegeville, PA 19426 (US). HANSBURY, Michael, J.; 885 Haddon Avenue, Collingswood, NJ 08108 (US). NERURKAR, Sandhya, S.; Apartment #J415, 261 Avon Road, Devon, PA 19333 (US). ROSHAK, Amy, K.; 3004 Green Ridge Drive, East Norriton, PA 19403 (US). BOUZYK, Mark; 2 Ford Cottages, Little Hadham, Herts SG11 2BN (GB).</p> <p>(74) Agents: ANDERSEN, Robert, L. et al.; Ratner & Prestia, 301 One Westlakes, Berwyn, P.O. Box 980, Valley Forge, PA 19482-0980 (US).</p>	<p>(81) Designated States: CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report.</i></p>	
<p>(54) Title: HUMAN SERUM INDUCIBLE KINASE (SNK)</p>		
<p>(57) Abstract</p> <p>The Serum Inducible Kinase (Snk) polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing Serum Inducible Kinase (Snk) polypeptides and polynucleotides in therapy, and diagnostic assays for such.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

HUMAN SERUM INDUCIBLE KINASE (SNK)

This application claims the benefit of U.S. Provisional Application No. 60/056,112, filed August 20, 1998, whose contents are herein incorporated by reference in their entirety.

Field of the Invention

This invention relates to newly identified polypeptides and polynucleotides encoding such polypeptides, to their use in therapy and in identifying compounds which may be agonists, antagonists and/or inhibitors which are potentially useful in therapy, and to production of such polypeptides and polynucleotides.

Background of the Invention

The drug discovery process is currently undergoing a fundamental revolution as it embraces 'functional genomics', that is, high throughput genome- or gene-based biology. This approach is rapidly superseding earlier approaches based on 'positional cloning'. A phenotype, that is a biological function or genetic disease, would be identified and this would then be tracked back to the responsible gene, based on its genetic map position.

Functional genomics relies heavily on the various tools of bioinformatics to identify gene sequences of potential interest from the many molecular biology databases now available. There is a continuing need to identify and characterise further genes and their related polypeptides/proteins, as targets for drug discovery.

Protein phosphorylation plays a critical role in promoting cell cycle progression. Most prominent among the regulators of the cell cycle is a family of cyclins, cyclin dependent kinases (CDKs), CDK regulatory kinases, and phosphatases (See Lees, E., *Curr. Opin. Cell Biol.* 1995, 7:773-780; Piwinica-Worms, H., *J. Lab. Clin. Med.* 1996, 128:350-354). A new family of cell cycle regulators, the polo-like kinases, has been identified and shown to be essential for progression through the cell cycle (Lane, H. A., *Trends in Cell Biol.* 1997, 7:63-68). This subfamily of serine/threonine kinases contains the following related but distinct members: (1) Plk (polo-like kinase; human) and its homologs Polo (*Drosophila*), *cdc5* (*S. cerevisiae*), Plx (*Xenopus*), and Plo (*S. pombe*); (2) Prk (polo-related kinase; human) and its murine homolog Fnk; and (3) Snk (serum-inducible kinase; murine). Known functions of these genes include regulation of spindle assembly (human plk1,

- 2 -

Drosophila polo, *S. pombe plo1*) and late nuclear division (*S. cerevisiae cdc5*). PLK1 expression correlates with the mitotic index (Holtrich U., Proc. Natl. Acad. Sci. 1994, 91:1736-1740) and mutations of the *Drosophila polo* or *S. cerevisiae cdc5* gene cause mitotic arrest. In addition, antibodies directed against human PLK1 cause impaired mitosis.

5 Progression from the G2 phase to the M phase of the cell cycle requires the activity of cdc25 phosphatase. PLX1 (*Xenopus*) phosphorylates and, thereby, activates cdc25-c, an isoform of cdc25 (Dunphy W.G., Science 1996. 273:1377-1380). The murine Snk is an early growth response gene which reportedly phosphorylates heterologous (although unidentified) substrates (Simmons D.L., Mol. Cell. Biol. 1992, 12:4164-4169). Identification of the

10 consensus sequence of the polo-like family in the amino-terminal putative catalytic domain of Snk (published murine sequence and present invention) and the consensus polo box sequence in the carboxy terminus place this protein in the polo-like family and suggest that this enzyme is potentially a critical regulator of cell cycle progression.

Summary of the Invention

15 The present invention relates to Serum Inducible Kinase (Snk), in particular Serum Inducible Kinase (Snk) polypeptides and Serum Inducible Kinase (Snk) polynucleotides, recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including the treatment

20 of proliferative diseases such as leukemia, solid tumor cancers and metastases; chronic inflammatory proliferative diseases such as psoriasis and rheumatoid arthritis; proliferative cardiovascular diseases such as restenosis; proliferative ocular disorders such as diabetic retinopathy; and benign hyperproliferative diseases such as hemangiomas, hereinafter referred to as "the Diseases", amongst others. In a further aspect, the invention relates to methods for identifying agonists and antagonists/inhibitors using the materials provided

25 by the invention, and treating conditions associated with Serum Inducible Kinase (Snk) imbalance with the identified compounds. In a still further aspect, the invention relates to diagnostic assays for detecting diseases associated with inappropriate Serum Inducible Kinase (Snk) activity or levels.

- 3 -

Description of the Invention

In a first aspect, the present invention relates to Serum Inducible Kinase (Snk) polypeptides. Such peptides include isolated polypeptides comprising an amino acid sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, to that of SEQ ID NO:2 over the entire length of SEQ ID NO:2. Such polypeptides include those comprising the amino acid of SEQ ID NO:2.

Further peptides of the present invention include isolated polypeptides in which the amino acid sequence has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, to the amino acid sequence of SEQ ID NO:2 over the entire length of SEQ ID NO:2. Such polypeptides include the polypeptide of SEQ ID NO:2.

Further peptides of the present invention include isolated polypeptides encoded by a polynucleotide comprising the sequence contained in SEQ ID NO:1.

Polypeptides of the present invention are believed to be members of the Polo-like Kinase family of polypeptides. They are therefore of interest because the Polo-like Kinase family has an established, proven history as therapeutic targets. Clearly there is a need for identification and characterization of further members of the Polo-like Kinase family. These properties are hereinafter referred to as "Serum Inducible Kinase (Snk) activity" or "Serum Inducible Kinase (Snk) polypeptide activity" or "biological activity of Serum Inducible Kinase (Snk)". Also included amongst these activities are antigenic and immunogenic activities of said Serum Inducible Kinase (Snk) polypeptides, in particular the antigenic and immunogenic activities of the polypeptide of SEQ ID NO:2. Preferably, a polypeptide of the present invention exhibits at least one biological activity of Serum Inducible Kinase (Snk).

The polypeptides of the present invention may be in the form of the "mature" protein or may be a part of a larger protein such as a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or

- 4 -

leader sequences, pro-sequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

The present invention also includes include variants of the aforementioned polypeptides, that is polypeptides that vary from the referents by conservative amino acid
5 substitutions, whereby a residue is substituted by another with like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acids are substituted, deleted, or added in any combination.

10 Polypeptides of the present invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

15 In a further aspect, the present invention relates to Serum Inducible Kinase (Snk) polynucleotides. Such polynucleotides include isolated polynucleotides comprising a nucleotide sequence encoding a polypeptide which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, to the amino acid sequence of SEQ ID NO:2, over the entire length of SEQ
20 ID NO:2. In this regard, polypeptides which have at least 97% identity are highly preferred, whilst those with at least 98-99% identity are more highly preferred, and those with at least 99% identity are most highly preferred. Such polynucleotides include a polynucleotide comprising the nucleotide sequence contained in SEQ ID NO:1 encoding the polypeptide of SEQ ID NO:2.

25 Further polynucleotides of the present invention include isolated polynucleotides comprising a nucleotide sequence that has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, to a nucleotide sequence encoding a polypeptide of SEQ ID NO:2, over the entire coding region. In this regard, polynucleotides which have at least 97% identity are highly preferred,

- 5 -

whilst those with at least 98-99 % identity are more highly preferred, and those with at least 99 % identity are most highly preferred.

Further polynucleotides of the present invention include isolated polynucleotides comprising a nucleotide sequence which has at least 70 % identity, preferably at least 80 %
5 identity, more preferably at least 90 % identity, yet more preferably at least 95 % identity, to SEQ ID NO:1 over the entire length of SEQ ID NO:1. In this regard, polynucleotides which have at least 97 % identity are highly preferred, whilst those with at least 98-99 % identity are more highly preferred, and those with at least 99 % identity are most highly preferred. Such polynucleotides include a polynucleotide comprising the polynucleotide of
10 SEQ ID NO:1 as well as the polynucleotide of SEQ ID NO:1.

The invention also provides polynucleotides which are complementary to all the above described polynucleotides.

The nucleotide sequence of SEQ ID NO:1 shows homology with Murine Serum Inducible Kinase (Simmons et al., Mol. Cell. Biol. 12(9):4164-4169, 1992). The
15 nucleotide sequence of SEQ ID NO:1 is a cDNA sequence and comprises a polypeptide encoding sequence (nucleotide 124 to 2181 encoding a polypeptide of 685 amino acids, the polypeptide of SEQ ID NO:2. The nucleotide sequence encoding the polypeptide of SEQ ID NO:2 may be identical to the polypeptide encoding sequence contained in SEQ ID NO:1 or it may be a sequence other than the one contained in SEQ ID NO:1, which, as a
20 result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2. The polypeptide of SEQ ID NO:2 is structurally related to other proteins of the Polo-like Kinase family, having homology and/or structural similarity with Murine Serum Inducible Kinase (Simmons et al., Mol. Cell. Biol. 12(9):4164-4169, 1992).

Preferred polypeptides and polynucleotides of the present invention are expected to
25 have, *inter alia*, similar biological functions/properties to their homologous polypeptides and polynucleotides. Furthermore, preferred polypeptides and polynucleotides of the present invention have at least one Serum Inducible Kinase (Snk) activity.

The present invention also relates to partial or other polynucleotide and polypeptide sequences which were first identified prior to the determination of the corresponding full
30 length sequences of SEQ ID NO:1 and SEQ ID NO:2.

- 6 -

Accordingly, in a further aspect, the present invention provides for an isolated polynucleotide comprising:

- (a) a nucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% identity to SEQ ID NO:3 over the entire length of SEQ ID NO:3;
- (b) a nucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% identity, to SEQ ID NO:3 over the entire length of SEQ ID NO:3; or
- (c) the polynucleotide of SEQ ID NO:3.

The nucleotide sequence of SEQ ID NO:3 and the peptide sequence encoded thereby are derived from EST (Expressed Sequence Tag) sequences. It is recognized by those skilled in the art that there will inevitably be some nucleotide sequence reading errors in EST sequences (see Adams, M.D. *et al.*, *Nature* 377 (supp) 3, 1995). Accordingly, the nucleotide sequence of SEQ ID NO:3 and the peptide sequence encoded therefrom are therefore subject to the same inherent limitations in sequence accuracy. Furthermore, the peptide sequence encoded by SEQ ID NO:3 comprises a region of identity or close homology and/or close structural similarity (for example a conservative amino acid difference) with the closest homologous or structurally similar protein.

Polynucleotides of the present invention may be obtained, using standard cloning and screening techniques, from a cDNA library derived from mRNA in cells of human H2LAS46, colon carcinoma, using the expressed sequence tag (EST) analysis (Adams, M.D., *et al.* *Science* (1991) 252:1651-1656; Adams, M.D. *et al.*, *Nature*, (1992) 355:632-634; Adams, M.D., *et al.*, *Nature* (1995) 377 Supp:3-174). Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

When polynucleotides of the present invention are used for the recombinant production of polypeptides of the present invention, the polynucleotide may include the coding sequence for the mature polypeptide, by itself; or the coding sequence for the mature

- 7 -

polypeptide in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc Natl Acad Sci USA* (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Further embodiments of the present invention include polynucleotides encoding polypeptide variants which comprise the amino acid sequence of SEQ ID NO:2 and in which several, for instance from 5 to 10, 1 to 5, 1 to 3, 1 to 2 or 1, amino acid residues are substituted, deleted or added, in any combination.

Polynucleotides which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO:1, may be used as hybridization probes for cDNA and genomic DNA or as primers for a nucleic acid amplification (PCR) reaction, to isolate full-length cDNAs and genomic clones encoding polypeptides of the present invention and to isolate cDNA and genomic clones of other genes (including genes encoding homologs and orthologs from species other than human) that have a high sequence similarity to SEQ ID NO:1. Typically these nucleotide sequences are 70% identical, preferably 80% identical, more preferably 90% identical, most preferably 95% identical to that of the referent. The probes or primers will generally comprise at least 15 nucleotides, preferably, at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will have between 30 and 50 nucleotides.

A polynucleotide encoding a polypeptide of the present invention, including homologs and orthologs from species other than human, may be obtained by a process which comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO: 1 or a fragment thereof; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to the skilled artisan. Preferred stringent hybridization conditions include overnight incubation at 42°C in a solution

- 8 -

comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA; followed by washing the filters in 0.1x SSC at about 65°C. Thus the present invention also includes polynucleotides obtainable by screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or a fragment thereof.

The skilled artisan will appreciate that, in many cases, an isolated cDNA sequence will be incomplete, in that the region coding for the polypeptide is cut short at the 5' end of the cDNA. This is a consequence of reverse transcriptase, an enzyme with inherently low 'processivity' (a measure of the ability of the enzyme to remain attached to the template during the polymerisation reaction), failing to complete a DNA copy of the mRNA template during 1st strand cDNA synthesis.

There are several methods available and well known to those skilled in the art to obtain full-length cDNAs, or extend short cDNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman et al., PNAS USA 85, 8998-9002, 1988). Recent modifications of the technique, exemplified by the MarathonTM technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the MarathonTM technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the 'missing' 5' end of the cDNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using 'nested' primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the known gene sequence). The products of this reaction can then be analyzed by DNA sequencing and a full-length cDNA constructed either by joining the product directly to the existing cDNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

Recombinant polypeptides of the present invention may be prepared by processes well known in the art from genetically engineered host cells comprising expression systems.

Accordingly, in a further aspect, the present invention relates to expression systems which comprise a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis et al., *Basic Methods in Molecular Biology* (1986) and Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989). Preferred such methods include, for instance, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include bacterial cells, such as *streptococci*, *staphylococci*, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used, for instance, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector which is able to maintain, propagate or express a polynucleotide to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as,

- 10 -

for example, those set forth in Sambrook *et al.*, *MOLECULAR CLONING, A LABORATORY MANUAL (supra)*. Appropriate secretion signals may be incorporated into the desired polypeptide to allow secretion of the translated protein into the lumen of the endoplasmic reticulum, the periplasmic space or the extracellular environment. These
5 signals may be endogenous to the polypeptide or they may be heterologous signals.

If a polypeptide of the present invention is to be expressed for use in screening assays, it is generally preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If the polypeptide is secreted into the medium, the medium can be recovered in order to recover
10 and purify the polypeptide. If produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose
15 chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

20 This invention also relates to the use of polynucleotides of the present invention as diagnostic reagents. Detection of a mutated form of the gene characterized by the polynucleotide of SEQ ID NO:1 which is associated with a dysfunction will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, or susceptibility to a disease, which results from under-expression, over-expression or altered expression of the
25 gene. Individuals carrying mutations in the gene may be detected at the DNA level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other
30 amplification techniques prior to analysis. RNA or cDNA may also be used in similar

- 11 -

fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled Serum Inducible Kinase (Snk) nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing (e.g., Myers *et al.*, *Science* (1985) 230:1242). Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (see Cotton *et al.*, *Proc Natl Acad Sci USA* (1985) 85: 4397-4401). In another embodiment, an array of oligonucleotides probes comprising Serum Inducible Kinase (Snk) nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see for example: M.Chee *et al.*, *Science*, Vol 274, pp 610-613 (1996)).

The diagnostic assays offer a process for diagnosing or determining a susceptibility to the Diseases through detection of mutation in the Serum Inducible Kinase (Snk) gene by the methods described. In addition, such diseases may be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of polypeptide or mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, nucleic acid amplification, for instance PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as a polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit which comprises:

- 12 -

- (a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO: 1, or a fragment thereof ;
- (b) a nucleotide sequence complementary to that of (a);
- (c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2 or
5 a fragment thereof; or
- (d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO:2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility
10 to a disease, particularly proliferative diseases such as leukemia, solid tumor cancers and metastases; chronic inflammatory proliferative diseases such as psoriasis and rheumatoid arthritis; proliferative cardiovascular diseases such as restenosis; proliferative ocular disorders such as diabetic retinopathy; and benign hyperproliferative diseases such as hemangiomas, amongst others.

15 The nucleotide sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to, and can hybridize with, a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a
20 precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found in, for example, V. McKusick, Mendelian Inheritance in Man (available on-line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of
25 physically adjacent genes).

The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease. The gene of the present invention maps to human
30 chromosome 5d12.1-q13.2/D5S491-D5S427.

The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them, can also be used as immunogens to produce antibodies immunospecific for polypeptides of the present invention. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

Antibodies generated against polypeptides of the present invention may be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a non-human animal, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* (1983) 4:72) and the EBV-hybridoma technique (Cole *et al.*, MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp. 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies, such as those described in U.S. Patent No. 4,946,778, can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms, including other mammals, may be used to express humanized antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography.

Antibodies against polypeptides of the present invention may also be employed to treat the Diseases, amongst others.

In a further aspect, the present invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa. Furthermore, this invention relates to processes for the preparation of these fusion

proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

5 Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with a polypeptide of the present invention, adequate to produce antibody and/or T cell immune response to protect said animal from the Diseases hereinbefore mentioned, amongst others. Yet another aspect of the invention relates to a method of inducing immunological response in a
10 mammal which comprises, delivering a polypeptide of the present invention *via* a vector directing expression of the polynucleotide and coding for the polypeptide *in vivo* in order to induce such an immunological response to produce antibody to protect said animal from diseases.

A further aspect of the invention relates to an immunological/vaccine formulation
15 (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a polypeptide of the present invention wherein the composition comprises a polypeptide or polynucleotide of the present invention. The vaccine formulation may further comprise a suitable carrier. Since a polypeptide may be broken down in the stomach, it is preferably administered parenterally (for instance,
20 subcutaneous, intramuscular, intravenous, or intradermal injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The
25 formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend
30 on the specific activity of the vaccine and can be readily determined by routine experimentation.

- 15 -

Polypeptides of the present invention are responsible for many biological functions, including many disease states, in particular the Diseases hereinbefore mentioned. It is therefore desirous to devise screening methods to identify compounds which stimulate or which inhibit the function of the polypeptide. Accordingly, in a further aspect, the present invention provides for a method of screening compounds to identify those which stimulate or which inhibit the function of the polypeptide. In general, agonists or antagonists may be employed for therapeutic and prophylactic purposes for such Diseases as hereinbefore mentioned. Compounds may be identified from a variety of sources, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. Such agonists, antagonists or inhibitors so-identified may be natural or modified substrates, ligands, receptors, enzymes, etc., as the case may be, of the polypeptide; or may be structural or functional mimetics thereof (see Coligan *et al.*, *Current Protocols in Immunology* 1(2):Chapter 5 (1991)).

The screening method may simply measure the binding of a candidate compound to the polypeptide, or to cells or membranes bearing the polypeptide, or a fusion protein thereof by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve competition with a labeled competitor. Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells bearing the polypeptide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Constitutively active polypeptides may be employed in screening methods for inverse agonists or inhibitors, in the absence of an agonist or inhibitor, by testing whether the candidate compound results in inhibition of activation of the polypeptide. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide of the present invention, to form a mixture, measuring Serum Inducible Kinase (Snk) activity in the mixture, and comparing the Serum Inducible Kinase (Snk) activity of the mixture to a standard. Fusion proteins, such as those made from Fc portion and Serum Inducible Kinase (Snk) polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists for the polypeptide of the present invention (see

- 16 -

D. Bennett *et al.*, *J Mol Recognition*, 8:52-58 (1995); and K. Johanson *et al.*, *J Biol Chem*, 270(16):9459-9471 (1995)).

The polynucleotides, polypeptides and antibodies to the polypeptide of the present invention may also be used to configure screening methods for detecting the effect of added compounds on the production of mRNA and polypeptide in cells. For example, an ELISA assay may be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can be used to discover agents which may inhibit or enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

The polypeptide may be used to identify membrane bound or soluble receptors, if any, through standard receptor binding techniques known in the art. These include, but are not limited to, ligand binding and crosslinking assays in which the polypeptide is labeled with a radioactive isotope (for instance, ^{125}I), chemically modified (for instance, biotinylated), or fused to a peptide sequence suitable for detection or purification, and incubated with a source of the putative receptor (cells, cell membranes, cell supernatants, tissue extracts, bodily fluids). Other methods include biophysical techniques such as surface plasmon resonance and spectroscopy. These screening methods may also be used to identify agonists and antagonists of the polypeptide which compete with the binding of the polypeptide to its receptors, if any. Standard methods for conducting such assays are well understood in the art.

Examples of potential polypeptide antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligands, substrates, receptors, enzymes, etc., as the case may be, of the polypeptide, e.g., a fragment of the ligands, substrates, receptors, enzymes, etc.; or small molecules which bind to the polypeptide of the present invention but do not elicit a response, so that the activity of the polypeptide is prevented.

Thus, in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc. for

- 17 -

polypeptides of the present invention; or compounds which decrease or enhance the production of such polypeptides, which comprises:

- (a) a polypeptide of the present invention;
 - (b) a recombinant cell expressing a polypeptide of the present invention;
 - 5 (c) a cell membrane expressing a polypeptide of the present invention; or
 - (d) antibody to a polypeptide of the present invention;
- which polypeptide is preferably that of SEQ ID NO:2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

10 It will be readily appreciated by the skilled artisan that a polypeptide of the present invention may also be used in a method for the structure-based design of an agonist, antagonist or inhibitor of the polypeptide, by:

- (a) determining in the first instance the three-dimensional structure of the polypeptide;
- (b) deducing the three-dimensional structure for the likely reactive or binding site(s) of
15 an agonist, antagonist or inhibitor;
- (c) synthesizing candidate compounds that are predicted to bind to or react with the deduced binding or reactive site; and
- (d) testing whether the candidate compounds are indeed agonists, antagonists or
inhibitors.

20 It will be further appreciated that this will normally be an interactive process.

In a further aspect, the present invention provides methods of treating abnormal conditions such as, for instance, proliferative diseases such as leukemia, solid tumor cancers and metastases; chronic inflammatory proliferative diseases such as psoriasis and rheumatoid arthritis; proliferative cardiovascular diseases such as restenosis; proliferative ocular
25 disorders such as diabetic retinopathy; and benign hyperproliferative diseases such as hemangiomas, related to either an excess of, or an under-expression of, Serum Inducible Kinase (Snk) polypeptide activity.

If the activity of the polypeptide is in excess, several approaches are available. One approach comprises administering to a subject in need thereof an inhibitor compound (antagonist) as hereinabove described, optionally in combination with a pharmaceutically acceptable carrier, in an amount effective to inhibit the function of the polypeptide, such as, for example, by blocking the binding of ligands, substrates, receptors, enzymes, etc., or by inhibiting a second signal, and thereby alleviating the abnormal condition. In another approach, soluble forms of the polypeptides still capable of binding the ligand, substrate, enzymes, receptors, etc. in competition with endogenous polypeptide may be administered. Typical examples of such competitors include fragments of the Serum Inducible Kinase (Snk) polypeptide.

In still another approach, expression of the gene encoding endogenous Serum Inducible Kinase (Snk) polypeptide can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or separately administered (see, for example, O'Connor, *J Neurochem* (1991) 56:560 in Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). Alternatively, oligonucleotides which form triple helices with the gene can be supplied (see, for example, Lee *et al.*, *Nucleic Acids Res* (1979) 6:3073; Cooney *et al.*, *Science* (1988) 241:456; Dervan *et al.*, *Science* (1991) 251:1360). These oligomers can be administered *per se* or the relevant oligomers can be expressed *in vivo*.

For treating abnormal conditions related to an under-expression of Serum Inducible Kinase (Snk) and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates a polypeptide of the present invention, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of Serum Inducible Kinase (Snk) by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells

- 19 -

may be administered to a subject for engineering cells *in vivo* and expression of the polypeptide *in vivo*. For an overview of gene therapy, see Chapter 20, *Gene Therapy and other Molecular Genetic-based Therapeutic Approaches*, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd
5 (1996). Another approach is to administer a therapeutic amount of a polypeptide of the present invention in combination with a suitable pharmaceutical carrier.

In a further aspect, the present invention provides for pharmaceutical compositions comprising a therapeutically effective amount of a polypeptide, such as the soluble form of a polypeptide of the present invention, agonist/antagonist peptide or small molecule
10 compound, in combination with a pharmaceutically acceptable carrier or excipient. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention. Polypeptides and other compounds of the
15 present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The composition will be adapted to the route of administration, for instance by a systemic or an oral route. Preferred forms of systemic administration include injection, typically by intravenous injection. Other injection routes, such as subcutaneous,
20 intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if a polypeptide or other compounds of the present invention can be formulated in an enteric or an encapsulated formulation, oral administration may also be possible. Administration of these compounds
25 may also be topical and/or localized, in the form of salves, pastes, gels, and the like.

The dosage range required depends on the choice of peptide or other compounds of the present invention, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of 0.1-100 $\mu\text{g}/\text{kg}$ of subject. Wide variations in the needed
30 dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration

- 20 -

would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

Polypeptides used in treatment can also be generated endogenously in the subject, in
5 treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide *ex vivo*, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

Polynucleotide and polypeptide sequences form a valuable information resource with
10 which to identify further sequences of similar homology. This is most easily facilitated by storing the sequence in a computer readable medium and then using the stored data to search a sequence database using well known searching tools, such as GCC. Accordingly, in a further aspect, the present invention provides for a computer readable medium having stored thereon a polynucleotide comprising the sequence of SEQ ID NO:1 and/or a
15 polypeptide sequence encoded thereby.

The following definitions are provided to facilitate understanding of certain terms used frequently hereinbefore.

"Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the
20 products of an Fab or other immunoglobulin expression library.

"Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or
25 polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

"Polynucleotide" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation, single- and double-stranded DNA,
30 DNA that is a mixture of single- and double-stranded regions, single- and double-stranded

RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, “polynucleotide” refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term “polynucleotide” also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. “Modified” bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications may be made to DNA and RNA; thus, “polynucleotide” embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. “Polynucleotide” also embraces relatively short polynucleotides, often referred to as oligonucleotides.

“Polypeptide” refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. “Polypeptide” refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. “Polypeptides” include amino acid sequences modified either by natural processes, such as post-translational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications may occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present to the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from post-translation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative,

- 22 -

covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination (see, for instance, *PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993; Wold, F., *Post-translational Protein Modifications: Perspectives and Prospects*, pgs. 1-12 in *POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS*, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter *et al.*, "Analysis for protein modifications and nonprotein cofactors", *Meth Enzymol* (1990) 182:626-646 and Rattan *et al.*, "Protein Synthesis: Post-translational Modifications and Aging", *Ann NY Acad Sci* (1992) 663:48-62).

"Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

- 23 -

"Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as the case may be, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" can be readily calculated by known methods, including but not limited to those described in (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48: 1073 (1988). Methods to determine identity are designed to give the largest match between the sequences tested. Moreover, methods to determine identity are codified in publicly available computer programs. Computer program methods to determine identity between two sequences include, but are not limited to, the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN, and FASTA (Atschul, S.F. et al., *J. Molec. Biol.* 215: 403-410 (1990). The BLAST X program is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990). The well known Smith Waterman algorithm may also be used to determine identity.

Parameters for polypeptide sequence comparison include the following:

- 1) Algorithm: Needleman and Wunsch, *J. Mol Biol.* 48: 443-453 (1970)
 - 25 Comparison matrix: BLOSSUM62 from Hentikoff and Hentikoff, *Proc. Natl. Acad. Sci. USA.* 89:10915-10919 (1992)
- Gap Penalty: 12
- Gap Length Penalty: 4

- 24 -

A program useful with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison WI. The aforementioned parameters are the default parameters for peptide comparisons (along with no penalty for end gaps).

Parameters for polynucleotide comparison include the following:

- 5 1) Algorithm: Needleman and Wunsch, J. Mol Biol. 48: 443-453 (1970)

Comparison matrix: matches = +10, mismatch = 0

Gap Penalty: 50

Gap Length Penalty: 3

- 10 Available as: The "gap" program from Genetics Computer Group, Madison WI. These are the default parameters for nucleic acid comparisons.

A preferred meaning for "identity" for polynucleotides and polypeptides, as the case may be, are provided in (1) and (2) below.

- (1) Polynucleotide embodiments further include an isolated polynucleotide comprising a polynucleotide sequence having at least a 50, 60, 70, 80, 85, 90, 95, 97 or
 15 100% identity to the reference sequence of SEQ ID NO:1, wherein said polynucleotide sequence may be identical to the reference sequence of SEQ ID NO: 1 or may include up to a certain integer number of nucleotide alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion, and wherein said
 20 alterations may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of nucleotide alterations is determined by multiplying the total number of nucleotides in SEQ ID NO:1 by the integer defining the
 25 percent identity divided by 100 and then subtracting that product from said total number of nucleotides in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \cdot y),$$

- 25 -

wherein n_n is the number of nucleotide alterations, x_n is the total number of nucleotides in SEQ ID NO:1, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n . Alterations of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may create nonsense, missense or frameshift mutations in this coding sequence and thereby alter the polypeptide encoded by the polynucleotide following such alterations.

By way of example, a polynucleotide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:2, that is it may be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one nucleic acid deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference polynucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleic acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of nucleic acid alterations for a given percent identity is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_n \leq x_n - (x_n \bullet y),$$

25

wherein n_n is the number of amino acid alterations, x_n is the total number of amino acids in SEQ ID NO:2, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n .

- 26 -

(2) Polypeptide embodiments further include an isolated polypeptide comprising a polypeptide having at least a 50,60, 70, 80, 85, 90, 95, 97 or 100% identity to a polypeptide reference sequence of SEQ ID NO:2, wherein said polypeptide sequence may be identical to the reference sequence of SEQ ID NO: 2 or may include up to a certain integer number of amino acid alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of amino acid alterations is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

15

$$n_a \leq x_a - (x_a \cdot y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \cdot is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

By way of example, a polypeptide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:2, that is it may be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed

- 27 -

either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of amino acid alterations for a given % identity is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a \bullet y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., and \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

"Fusion protein" refers to a protein encoded by two, often unrelated, fused genes or fragments thereof. In one example, EP-A-0 464 discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, employing an immunoglobulin Fc region as a part of a fusion protein is advantageous for use in therapy and diagnosis resulting in, for example, improved pharmacokinetic properties [see, e.g., EP-A 0232 262]. On the other hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified.

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

SEQUENCE INFORMATION

SEQ ID NO:1

5 GGCACGAGGTTGGGTGCTATTTCGGCACCAGAGGCAAGGGTGCAGGACCACGGCCGGCTCGGACGTGTGACCGC
 GCCTAGGGGGTGGCAGCGGGCAGTGCAGGGGCGGCAAGGCGACCATGGAGCTTTTTCGGGACTATCACCTACCAGC
 CAGCCGCCAGCACCAAAATGTGCGAGCAGGCGCTGGGCAAGGGTTCGGGAGCAGACTCGAAGAAGAAGCGGGCCG
 CCGCAGCCCCCGAGGAATCGCAGCCACCTCAGTCCCAGGCGCAAGTGCCTCCCGGGCCCTCACCACCATCA
 CCACCATTTCGCACTCGGGGCCGAGATCTCGCGGATTATCGTCGACCCACGACTGGGAAGCGCTACTGCCGGG
 10 GCAAAGTGC TGGGAAAGGGTGGCTTTGCAAAATGTTACGAGATGACAGATTTGACAAATAACAAAGTCTACGCC
 GCAAAAATTATTCTCACAGCAGAGTAGCTAAACCTCATCAAAGGGAAAAGATTGACAAAGAAATAGAGCTTCA
 CAGAATTCCTCATATAAGCATGTAGTGCAGTTTTACCCTACTTTCGAGGACAAAGAAAACATTTACATTTCTCT
 TGGAATACTGCAGTAGAAGGCAATGGCTCATATTTTTGAAAGCAAGAAAGGTGTTGACAGAGCCAGAAGTTCGAT
 ACTACCTCAGGCAGATTGTGTCTGGACTGAAATACCTTCATGAACAAGAAATCTTGCACAGAGATCTCAAACCTA
 GGGAACTTTTTTTATTAATGAAGCCATGGAACATAAAAGTTGGGGACTTCGGTCTGGCAGCCAGGCTAGAACCCTT
 15 GGAACACAGAAGGAGAACGATATGTGGTACCCCAAATTATCTCTCTCTGAAAGTCTCAACAAAACAAGGACATG
 GCTGTGAATCAGACATTTGGGCCCTGGGCTGTGTAATGTATACAATGTTACTAGGGAGGCCCCCATTTGAAACT
 ACAAAATCTCAAAGAACTTATAGGTGCATAAGGGAAGCAAGGTATACAATGCCGTCTCATTGCTGGCTCTGC
 CAAGCACTTAATTGCTAGTATGTTGTCCAAAACCCAGAGGATAGGCCCTAGTTTGGATGACATCATTGACATG
 ACTTTTTTTTTG CAGGGCTTCACTCCGGACAGACTGTCTTCTAGCTGTTGTCATACAGTTCAGATTTCCACTTA
 20 TCAAGCCCAGCTAAGAATTTCTTTAAGAAAGCAGCTGCTGCTCTTTTTTGGTGGCAAAAAGACAAAGCAAGATA
 TATTGACACACATAATAGAGTGTCTAAAGAAGATGAAGACATCTACAAGCTTAGGCATGATTTGAAAAAGACTT
 CAATAACTCAGCAACCCAGCAAACACAGGACAGATGAGGAGCTCCAGCCACCTACCACCACAGTTGCCAGGTCT
 GGAACACCCG CAGTAGAAAACAAGCAGCAGATTGGGGATGCTATTTCGGATGATAGTCAGAGGGACTCTTGGCAG
 CTGTAGCAGCAGCAGTGAATGCCTTGAAGACAGTACCATGGGAAGTGTTCAGACACAGTGGCAAGGGTCTTTC
 GGGGATGTCTGGAAAACATGCCGGAAGCTGATTGCATTCCCAAAGAGCAGCTGAGCACATCATTTTCAGTGGGTC
 25 ACCAAATGGGTTGATTACTCTAACAAATATGGCTTTGGGTACCAGCTCTCAGACCACACCGTCCGGTGTCTTTTT
 CAACAATGGTGTCTCACATGAGCCTCCTTCCAGACAAAAAACAGTTCACTATTACGCAGAGCTTGGCCAATGCT
 CAGTTTTTCCAGCAACAGATGCTCCTGAGCAATTTATTAGTCAAGTGACGGTGTGAAATACTTTTCTCATTAC
 ATGGAGGAGAACCCTCATGGATGGTGGAGATCTGCCTAGTGTACTGATATTTCGAAGACCTCGGCTCTACCTCCT
 30 TCAGTGGCTAAAATCTGATAAGGCCCTAATGATGCTCTTTAATGATGGCACCTTTCAGGTGAATTTCTACCATG
 ATCATACAAAAATCATCTGTAGCCAAAATGAAGAATACCTTCTCACCTACATCAATGAGGATAGGATATCT
 ACAACTTTTCAGGCTGACAACTCTGCTGATGTCTGGCTGTTTATCATCAGAATTAATAAATCGAATGGAATATGCCCT
 GAACATGCTCTTACAAAAGATGTAAGTAAAGACTTTTTCGAATGGACCCTATGGGACTCCTCTTTTCCACTGTGA
 GATCTACAGGGAACCCAAAAGAATGATCTAGAGTATGTTGAAGAAGATGGACATGTGGTGGTACGAAAACAATT
 CCCCTGTGGCCTGTGGACTGGGTGGAACCAGAACAGGCTAAGGCATACAGTTCCTTGGACTTTGGACAATCCAAG
 35 AGTGAACCAGAATGCAGTTTTCTTGGAGATACCTGTTTTAAAAGGTTTTTTCAGACAATTTTTCAGAAAAGGTGCA
 TTGATTCTTAAATCTCTCTGTTGAGAGCATTTTCAGCCAGAGGACTTTGGAACTGTGAATATACTTCTGAAAGG
 GGAGGGAGAAGGGAGGAAGCTCCCATGTTGTTTAAAGGCTGTAATTTGGAGCAGCTTTTGGCTGCGTAACTGTGA
 ACTATGGCCATATATAATTTTTTTTTCATTAATTTTTGAAGATACTTGTGGCTGGAAAAGTGCATTCTTGTGTTAA
 40 TAAACTTTTTTATTATTACAGCCCAAAGAGCAGTATTTATTATCAAATGTCTTTUUCTTTGACCATTTTAAAC
 CGTTGGCAATAAAGAGTATGAAAACGCAAAAAAAAAAAAAAAAAAAAA

SEQ ID NO:2

45 MELLRTITYQPAASTKMCEQALGKGGADSKKKRPPQPPEESQPPQSQVPPAAPHHHHHSHSGPEISRIIV
 DPTTGKRYCRGKVLGKGGFAKCYEMTDLTNNKVYAAKIIPHSRVAKPHQREKIDKEIELHRILHKKHVQFYHY
 FEDKENIYILLEYCSRSMHILKARKVLTEPEVRYLRLQIVSGLKYLHEQEILHRDLKLGNNFFINEAMELKVG
 DFLAARLEPLEHRRRTICGTPNYLSPEVLNKQGHGCESDI WALGCVMYTMLLGRPPFETTLNKETYRCIREAR
 YTMPSSLLAPAKHLIASMLSKNPEDRPSLDDIIRHDFLQGFPTDRLSSSCCHTVPDFHLSSPAKNFFKKA
 50 LFFGKKDKARYIDTHNRVSKEDEDIYKLRHDLKKT SITQPSKHRDEELQPPTTTVARSGTPAVENKQIGDA
 IRMIVRGTGLGSCSSSECLELSTMGSVADTVARVLRGCLLENMPEADCI PKEQLSTSFQWVTKWVDYSNKYGFY
 QLSDHTVGVLFNNGAHMSLLPDKKTVHYAELGQCSVFPATDAPEQFISQVTVLKYF SHYMEENLMDGGDLPSV
 TDIRRPRLYLLQWLKSDKALMMLFNDGTFQVNFYHDHTKIIICSQNEEYLLTYINEDRISTTFRLTLLMSGCS
 SELKNRMEYALNMLLQRCN

SEQ ID NO:3

5 GGCACGAGGTTGGGTGCTATTTCGGCACCAGAGGCAAGGGTGCAGGACCACGGCCGGCTCGGACGTGTGACCGC
 GCCTAGGGGGTGGCAGCGGGCAGTGCGGGGCGGCAAGGCGACCATGGAGCTTTTTCGGACTATCACCTACCAGC
 CAGCCGCCAGCACCAAATGTGCGAGCAGGCGCTGGGCAAGGGTTGCGGAGCAGACTCGAAGAAGAAGCGGCCG
 10 CCGCAGCCCCCGAGGAATCGCAGCCACCTCAGTCCCAGGCGCAAGTGCCCCCGCGGCCCTCACCACCATCA
 CCACCATTTCGCACTCGGGGCCGAGATCTCGCGGATTATCGTCGACCCACGACTGGGAAGCGCTACTGCCGGG
 GCAAAGTGTCTGGGAAAGGGTGGCTTTGCAAAATGTTACGAGATGACAGATTTGACAAATAACAAAGTCTACGCC
 GCAAAAATTATTCCTCACAGCAGAGTAGCTAAACCTCATCAAAGGGAAAAGATTGACAAAGAAATAGAGCTTCA
 CAGAATTCCTCATCATAAGCATGTAGTGCAGTTTTACCCTACTTCGAGGACAAAGAAAACATTTACATTCTCT
 15 TGGAATACTGCAGTAGAAGGTCAATGGCTCATATTTTGAAAGCAAGAAAGGTGTTGACAGAGCCAGAAGTTCGA
 TACTACCTCAGGCAGATTGTGTCTGGACTGAAATACCTTCATGAACAAGAAATCTTGACAGAGATCTCAAACCT
 AGGGAACCTTTTTTATTAATGAAGCCATGGAACATAAAGTTGGGGACTTCGGTCTGGCAGCCAGGCTAGAACCCT
 TGGAACACAGAAGGAGAACGATATGTGGTACCCCAAATTATCTCTCTCCTGAAGTCTCAACAAAACAAGGACAT
 20 GGCTGTGAATCAGACATTTGGGCCCTGGGCTGTGTAATGTATAACAATGTTACTAGGGAGGCCCCCATTTGAAAC
 TACAAATCTCAAAGAACTTATAGGTGCATAAGGGAAGCAAGGTATAACAATGCCGTCTCATTGCTGGCTCCTG
 CCAAGCACTTAATTGCTAGTATGTTGTCCAAAGAACCCAGAGGATCGTCCAGTTTGGATGACATCATTGACAT
 GACTTTTTTTTTGCGAGGGCTTCACTCCGGACAGACTGTCTTCTAGCTGTTGTCATACAGTTCAGATTTCCACTT
 ATCAAGCCCAGCTAAGAAATTTCTTTAAGAAAGCAGCTGCTGCTCTTTTTTGGTGGCAAAAAGACAAAGCAAGAT
 25 ATATTGACACACATAATAGAGTGTCTAAAGAAAGATGAAGACATCTACAAGCTTAGGCATGATTTGAAAAAGACT
 TCAATAACTCAGCAACCCAGCAAACACAGGACAGATGAGGAGCTCCAGCCACCTACCACCACAGTTGCCAGGTC
 TGGAACACCCGCGAGTAGAAAACAAGCAGCAGATTGGGGATGCTATTTCGGATGATAGTCAGAGGGACTCTTGGA
 GCTGTAGCAGCAGCAGTGAATGCCCTTGAAAGACAGTACCATGGGAAGTGTGACAGACACAGTGGCAAGGGTCTT
 CGGGGATGTCTGGAACAACATGCCGGAAGCTGATTGCATTCCCAAAGAGCAGCTGAGCACATCATTTCAGTGGGT
 30 CACCAAATGGGTTGATTACTCTAACAAATATGGCTTTGGGTACCAGCTCTCAGACCACACCGTCCGGTGTCTTTT
 TCAACAATGGTGTCTCACATGAGCCTCCTTCCAGACAAAAAACAGTTCACTATTACGCAGAGCTTGGCCAATGC
 TCAGTTTTCCAGCAACAGATGCTCCTGAGCAATTTATTAGTCAAGTGACGGTGTGAAATACTTTTTCTCATT
 CATGGAGGAGAACCCTCATGGATGGTGGAGATCTGCCTAGTGTACTGATATTCGAAGACCTCGGCTCTACCTCC
 TTCAGTGGCTAAAATCTGATAAGGCCCTAATGATGCTCTTTAATGATGGCACCTTTCAGGTGAATTTCTACCAT
 35 GATCATACAAAATCATCATCTGTAGCCAAAATGAAGAATACCTTCTCACCTACATCAATGAGGATAGGATATC
 TACAACTTTCAGGCTGACAACTCTGCTGATGTCTGGCTGTTTCATCAGAAATTAATAAATCGAATGGAATATGCC
 TGAACATGCTCTTACAAAGATGTAAGTGAAGACTTTTTCGAATGGACCCTATGGGACTCCTCTTTTCCACTGTG
 AGATCTACAGGGAACCCAAAAGAAATGATCTAGAGTATGTTGAAGAAGATGGACATGTGGTGGTACGAAAACAAT
 TCCCCTGTGGCCTGCTGGACTGGGTGGAACCAGAACAGGCTAAGGCATACAGTTCTTGACTTTGGACAATCCAA
 40 GAGTGAACCAGAATGCAGTTTTCTTTGAGATACCTGTTTTAAAGGTTTTTTCAGACAATTTTTCAGAAAGGTGC
 ATTGATCTTAAATCTCTCTGTTGAGAGCATTTCAGCCAGAGGACTTTGGAACTGTGAATATACTTCTGAAAG
 GGGAGGGAGAAGGGAGGAAGCTCCCATGTTGTTTAAAGGCTGTAATTTGGAGCAGCTTTTTGGCTGCGTAACTGTG
 AACTATGGCCATATATAATTTTTTTTTCATTAATTTTTGAAGATACTTGTGGCTGGAAAAGTGCATTCTTGTTA
 ATAACTTTTTTATTTATTTACAGCCCAAAGAGCAGTATTTATTATCAAAATGCTTTTTTTTTTATGTTGACCATT
 TTAAACCGTTGGCAATAAAGAGTATGAAAACGCAAAAAAAAAAAAAAAAAAAAA

- 30 -

What is claimed is:

1. An isolated polypeptide selected from the group consisting of:
 - (i) an isolated polypeptide comprising an amino acid sequence selected from the group having at least:
 - 5 (a) 70% identity;
 - (b) 80% identity;
 - (c) 90% identity; or
 - (d) 95% identityto the amino acid sequence of SEQ ID NO:2 over the entire length of SEQ
10 ID NO:2;
 - (ii) an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:2
or
 - (iii) an isolated polypeptide which is the amino acid sequence of SEQ ID NO:2.
- 15 2. An isolated polynucleotide selected from the group consisting of:
 - (i) an isolated polynucleotide comprising a nucleotide sequence encoding a
polypeptide that has at least
 - (a) 70% identity;
 - (b) 80% identity;
 - 20 (c) 90% identity; or
 - (d) 95% identity;to the amino acid sequence of SEQ ID NO:2, over the entire length of SEQ ID
NO:2;
 - (ii) an isolated polynucleotide comprising a nucleotide sequence that has at least:
 - 25 (a) 70% identity

- 31 -

(b) 80% identity;

(c) 90% identity; or

(d) 95% identity;

5 over its entire length to a nucleotide sequence encoding the polypeptide of SEQ ID NO:2;

(iii) an isolated polynucleotide comprising a nucleotide sequence which has at least:

(a) 70% identity;

(b) 80% identity;

(c) 90% identity; or

10 (d) 95% identity;

to that of SEQ ID NO: 1 over the entire length of SEQ ID NO:1;

(iv) an isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2;

(vi) an isolated polynucleotide which is the polynucleotide of SEQ ID NO: 1; or

15 (vi) an isolated polynucleotide obtainable by screening an appropriate library under stringent hybridization conditions with a labelled probe having the sequence of SEQ ID NO: 1 or a fragment thereof.;

or a nucleotide sequence complementary to said isolated polynucleotide.

20 3. An antibody immunospecific for the polypeptide of claim 1.

4. A method for the treatment of a subject:

(i) in need of enhanced activity or expression of the polypeptide of claim 1 comprising:

25 (a) administering to the subject a therapeutically effective amount of an agonist to said polypeptide; and/or

SUBSTITUTE SHEET (RULE 26)

- 32 -

(b) providing to the subject an isolated polynucleotide comprising a nucleotide sequence encoding said polypeptide in a form so as to effect production of said polypeptide activity *in vivo.*; or

5 (ii) having need to inhibit activity or expression of the polypeptide of claim 1 comprising:

(a) administering to the subject a therapeutically effective amount of an antagonist to said polypeptide; and/or

(b) administering to the subject a nucleic acid molecule that inhibits the expression of a nucleotide sequence encoding said polypeptide;

10 and/or

(c) administering to the subject a therapeutically effective amount of a polypeptide that competes with said polypeptide for its ligand, substrate, or receptor.

15 5. A process for diagnosing a disease or a susceptibility to a disease in a subject related to expression or activity of the polypeptide of claim 1 in a subject comprising:

(a) determining the presence or absence of a mutation in the nucleotide sequence encoding said polypeptide in the genome of said subject; and/or

20 (b) analyzing for the presence or amount of said polypeptide expression in a sample derived from said subject.

6. A method for screening to identify compounds which stimulate or which inhibit the function of the polypeptide of claim 1 which comprises a method selected from the group consisting of:

25 (a) measuring the binding of a candidate compound to the polypeptide (or to the cells or membranes bearing the polypeptide) or a fusion protein thereof by means of a label directly or indirectly associated with the candidate compound;

- 33 -

- (b) measuring the binding of a candidate compound to the polypeptide (or to the cells or membranes bearing the polypeptide) or a fusion protein thereof in the presence of a labeled competitor;
- (c) testing whether the candidate compound results in a signal generated by
5 activation or inhibition of the polypeptide, using detection systems appropriate to the cells or cell membranes bearing the polypeptide;
- (d) mixing a candidate compound with a solution containing a polypeptide of claim 1, to form a mixture, measuring activity of the polypeptide in the mixture, and comparing the activity of the mixture to a standard; or
- 10 (e) detecting the effect of a candidate compound on the production of mRNA encoding said polypeptide and said polypeptide in cells, using for instance, an ELISA assay.
7. An agonist or an antagonist of the polypeptide of claim 1.
- 15 8. An expression system comprising a polynucleotide capable of producing a polypeptide of claim 1 when said expression system is present in a compatible host cell.
9. A process for producing a recombinant host cell comprising transforming or
20 transfecting a cell with the expression system of claim 8 such that the host cell, under appropriate culture conditions, produces a polypeptide comprising an amino acid sequence having at least 70% identity to the amino acid sequence of SEQ ID NO:2 over the entire length of SEQ ID NO:2.
- 25 10. A recombinant host cell produced by the process of claim 9.

- 34 -

11. A membrane of a recombinant host cell of claim 10 expressing a polypeptide comprising an amino acid sequence having at least 70% identity to the amino acid sequence of SEQ ID NO:2 over the entire length of SEQ ID NO:2.
- 5 12. A process for producing a polypeptide comprising culturing a host cell of claim 10 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture.
13. An isolated polynucleotide selected from the group consisting of:
- 10 (a) an isolated polynucleotide comprising a nucleotide sequence which has at least 70%, 80%, 90%, 95%, 97% identity to SEQ ID NO:3 over the entire length of SEQ ID NO:3;
- (b) an isolated polynucleotide comprising the polynucleotide of SEQ ID NO:3; or
- (c) the polynucleotide of SEQ ID NO:3.

- 1/5 -

SEQUENCE LISTING

(1) GENERAL INFORMATION

- 5 (i) APPLICANT: ANDERSON, KAREN
JACKSON, JEFFREY
HANSBURY, MICHAEL
NERURKAR, SANDHYA
10 ROSHAK, AMY
BOUZYK, MARK
- (ii) TITLE OF THE INVENTION: HUMAN SERUM INDUCIBLE KINASE (SNK)
- (iii) NUMBER OF SEQUENCES: 3
- 15 (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Ratner & Prestia
(B) STREET: P.O. Box 980
(C) CITY: Valley Forge
20 (D) STATE: PA
(E) COUNTRY: USA
(F) ZIP: 19482
- (v) COMPUTER READABLE FORM:
25 (A) MEDIUM TYPE: Diskette
(B) COMPUTER: IBM Compatible
(C) OPERATING SYSTEM: DOS
(D) SOFTWARE: FastSEQ for Windows Version 2.0
- 30 (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER: TO BE ASSIGNED
(B) FILING DATE: 20-AUG-1998
(C) CLASSIFICATION: UNKNOWN
- 35 (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 60/056,112
(B) FILING DATE: 20-AUG-1997
- 40 (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Prestia, Paul F
(B) REGISTRATION NUMBER: 23,031
45 (C) REFERENCE/DOCKET NUMBER: GH-70231
- (ix) TELECOMMUNICATION INFORMATION:
50 (A) TELEPHONE: 610-407-0700
(B) TELEFAX: 610-407-0700
(C) TELEX: 846169
- (2) INFORMATION FOR SEQ ID NO:1:
- 55 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 2783 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 60 (ii) MOLECULE TYPE: cDNA

SUBSTITUTE SHEET (RULE 26)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

	GGCACGAGGT	TGGGTGCTAT	TCGGCACCAG	AGGCAAGGGT	GCGAGGACCA	CGGCCGGCTC	60
	GGACGTGTGA	CCGCGCCTAG	GGGGTGGCAG	CGGGCAGTGC	GGGGCGGCAA	GGCGACCATG	120
5	GAGCTTTTGC	GGACTATCAC	CTACCAGCCA	GCCGCCAGCA	CCAAAATGTG	CGAGCAGGCG	180
	CTGGGCAAGG	GTTGCGGAGC	AGACTCGAAG	AAGAAGCGGC	CGCCGCAGCC	CCCCGAGGAA	240
	TCGCAGCCAC	CTCAGTCCCA	GGCGCAAGTG	CCCCCGGGCG	CCCCTCACCA	CCATCACCAC	300
	CATTCGCACT	CGGGGCCGGA	GATCTCGCGG	ATTATCGTCG	ACCCCACGAC	TGGGAAGCGC	360
	TACTGCCGGG	GCAAAGTGCT	GGGAAAGGGT	GGCTTTGCAA	AATGTTACGA	GATGACAGAT	420
10	TTGACAAATA	ACAAAGTCTA	CGCCGCAAAA	ATTATTCCCTC	ACAGCAGAGT	AGCTAAACCT	480
	CATCAAAGGG	AAAAGATTGA	CAAAGAAATA	GAGTTCACA	GAATTCCTCA	TCATAAGCAT	540
	GTAGTGCAGT	TTTACCACTA	CTTCGAGGAC	AAAGAAAACA	TTTACATTCT	CTTGGAATAC	600
	TGCAGTAGAA	GGCAATGGCT	CATATTTTGA	AAGCAAGAAA	GGTGTGACA	GAGCCAGAAG	660
	TTCGATACTA	CCTCAGGCAG	ATTGTGTCTG	GACTGAAATA	CCTTCATGAA	CAAGAAATCT	720
15	TGCACAGAGA	TCTCAAACCTA	GGGAACCTTT	TTATTAATGA	AGCCATGGAA	CTAAAAGTTG	780
	GGGACTTCGG	TCTGGCAGCC	AGGCTAGAAC	CCTTGGAACA	CAGAAGGAGA	ACGATATGTG	840
	GTACCCCAAC	TTATCTCTCT	CCTGAAGTCC	TCAACAAACA	AGGACATGGC	TGTGAATCAG	900
	ACATTTGGGG	CCTGGGCTGT	GTAATGTATA	CAATGTTACT	AGGGAGGCC	CCATTTGAAA	960
	CTACAAATCT	CAAAGAAACT	TATAGGTGCA	TAAGGGAAGC	AAGGTATACA	ATGCCGTCTT	1020
20	CATTGCTGGC	TCCTGCCAAG	CACTTAATTG	CTAGTATGTT	GTCCAAAAAC	CCAGAGGATA	1080
	GGCCTAGTTT	GGATGACATC	ATTCGACATG	ACTTTTTTTT	GCAGGGCTTC	ACTCCGGACA	1140
	GACTGTCTTC	TAGCTGTTGT	CATACAGTTC	CAGATTTCCA	CTTATCAAGC	CCAGCTAAGA	1200
	ATTTCTTTAA	GAAAGCAGCT	GCTGCTCTTT	TTGGTGGCAA	AAAAGACAAA	GCAAGATATA	1260
	TTGACACACA	TAATAGAGTG	TCTAAAGAAG	ATGAAGACAT	CTACAAGCTT	AGGCATGATT	1320
25	TGAAAAAGAC	TTCAATAACT	CAGCAACCCA	GCAAACACAG	GACAGATGAG	GAGCTCCAGC	1380
	CACCTACCAC	CACAGTTGCC	AGGTCTGGAA	CACCCGCACT	AGAAAACAAG	CAGCAGATTG	1440
	GGGATGCTAT	TCGGATGATA	GTCAGAGGGA	CTCTTGGCAG	CTGTAGCAGC	AGCATGAAAT	1500
	GCCTTGAAGA	CAGTACCATG	GGAAAGTGTG	CAGACACAGT	GGCAAGGGTT	CTTCGGGGAT	1560
	GTCTGGAAAA	CATGCCGGAA	GCTGATTGCA	TTCCCAAAGA	GCAGCTGAGC	ACATCATTTT	1620
30	AGTGGGTCAC	CAAAATGGGT	GATTACTCTA	ACAAATATGG	CTTTGGGTAC	CAGCTCTCAG	1680
	ACCACACCGT	CGGTGTCCTT	TTCAACAATG	GTGCTCACAT	GAGCCTCCTT	CCAGACAAAA	1740
	AAACAGTTCA	CTATTACGCA	GAGCTTGGCC	AATGCTCAGT	TTTCCCAGCA	ACAGATGCTC	1800
	CTGAGCAATT	TATTAGTCAA	GTGACGGTGC	TGAAATACTT	TTCTCATTAC	ATGGAGGAGA	1860
	ACCTCATGGA	TGGTGGAGAT	CTGCCTAGTG	TTACTGATAT	TCGAAGACCT	CGGCTCTACC	1920
35	TCCTTCAGTG	GCTAAAATCT	GATAAGGCC	TAATGATGCT	CTTTAATGAT	GGCACCTTTT	1980
	AGGTGAAATTT	CTACCATGAT	CATACAAAAA	TCATCATCTG	TAGCCAAAAT	GAAGAATACC	2040
	TTCTCACCTA	CATCAATGAG	GATAGGATAT	CTACAACCTT	CAGGCTGACA	ACTCTGCTGA	2100
	TGTCTGGCTG	TTCATCAGAA	TTAAAAAATC	GAATGGAATA	TGCCCTGAAC	ATGCTCTTAC	2160
	AAAGATGTAA	CTGAAAGACT	TTTCGAATGG	ACCCTATGGG	ACTCCTCTTT	TCCACTGTGA	2220
40	GATCTACAGG	GAACCCAAAA	GAATGATCTA	GAGTATGTTG	AAGAAGATGG	ACATGTGGTG	2280
	GTACGAAAAC	AATTCCCCTG	TGGCCTGCTG	GACTGGGTGG	AACCAGAACA	GGCTAAGGCA	2340
	TACAGTTCTT	GACTTTGGAC	AATCCAAGAG	TGAACCAGAA	TGCAGTTTTT	CTTGAGATAC	2400
	CTGTTTTTAA	AGGTTTTTCA	GACAATTTTG	CAGAAAGGTG	CATTGATTCT	TAAATTTCTT	2460
	CTGTTGAGAG	CATTCAGCC	AGAGGACTTT	GGAAGTGTGA	ATATACTTCC	TGAAGGGGAG	2520
45	GGAGAAGGGA	GGAAAGTCCC	ATGTTGTTTA	AAGGCTGTAA	TTGGAGCAGC	TTTTGGCTGC	2580
	GTAAGTGTGA	ACTATGGCCA	TATATAAATTT	TTTTTCATTA	ATTTTTGAAG	ATACTTGTGG	2640
	CTGAAAAAGT	GCATTCCTTG	TTAATAAACT	TTTTATTTAT	TACAGCCCAA	AGAGCAGTAT	2700
	TTATTATCAA	AATGTCTTTU	UCTTTGACCA	TTTTAAACCG	TTGGCAATAA	AGAGTATGAA	2760
50	AACGCAAAAA	AAAAAAAAAA	AAA				2783

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 685 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

	Met	Glu	Leu	Leu	Arg	Thr	Ile	Thr	Tyr	Gln	Pro	Ala	Ala	Ser	Thr	Lys
	1				5					10					15	
5	Met	Cys	Glu	Gln	Ala	Leu	Gly	Lys	Gly	Cys	Gly	Ala	Asp	Ser	Lys	Lys
			20						25				30			
	Lys	Arg	Pro	Pro	Gln	Pro	Pro	Glu	Glu	Ser	Gln	Pro	Pro	Gln	Ser	Gln
		35						40					45			
	Ala	Gln	Val	Pro	Pro	Ala	Ala	Pro	His	His	His	His	His	His	Ser	His
	50						55					60				
10	Ser	Gly	Pro	Glu	Ile	Ser	Arg	Ile	Ile	Val	Asp	Pro	Thr	Thr	Gly	Lys
	65				70						75					80
	Arg	Tyr	Cys	Arg	Gly	Lys	Val	Leu	Gly	Lys	Gly	Gly	Phe	Ala	Lys	Cys
				85						90					95	
15	Tyr	Glu	Met	Thr	Asp	Leu	Thr	Asn	Asn	Lys	Val	Tyr	Ala	Ala	Lys	Ile
				100					105						110	
	Ile	Pro	His	Ser	Arg	Val	Ala	Lys	Pro	His	Gln	Arg	Glu	Lys	Ile	Asp
			115					120					125			
	Lys	Glu	Ile	Glu	Leu	His	Arg	Ile	Leu	His	His	Lys	His	Val	Val	Gln
	130						135					140				
20	Phe	Tyr	His	Tyr	Phe	Glu	Asp	Lys	Glu	Asn	Ile	Tyr	Ile	Leu	Leu	Glu
	145					150					155					160
	Tyr	Cys	Ser	Arg	Arg	Ser	Met	Ala	His	Ile	Leu	Lys	Ala	Arg	Lys	Val
				165						170					175	
25	Leu	Thr	Glu	Pro	Glu	Val	Arg	Tyr	Tyr	Leu	Arg	Gln	Ile	Val	Ser	Gly
				180					185					190		
	Leu	Lys	Tyr	Leu	His	Glu	Gln	Glu	Ile	Leu	His	Arg	Asp	Leu	Lys	Leu
			195					200					205			
	Gly	Asn	Phe	Phe	Ile	Asn	Glu	Ala	Met	Glu	Leu	Lys	Val	Gly	Asp	Phe
	210					215						220				
30	Gly	Leu	Ala	Ala	Arg	Leu	Glu	Pro	Leu	Glu	His	Arg	Arg	Arg	Thr	Ile
	225					230						235				240
	Cys	Gly	Thr	Pro	Asn	Tyr	Leu	Ser	Pro	Glu	Val	Leu	Asn	Lys	Gln	Gly
				245						250					255	
35	His	Gly	Cys	Glu	Ser	Asp	Ile	Trp	Ala	Leu	Gly	Cys	Val	Met	Tyr	Thr
				260					265						270	
	Met	Leu	Leu	Gly	Arg	Pro	Pro	Phe	Glu	Thr	Thr	Asn	Leu	Lys	Glu	Thr
				275				280						285		
	Tyr	Arg	Cys	Ile	Arg	Glu	Ala	Arg	Tyr	Thr	Met	Pro	Ser	Ser	Leu	Leu
		290				295						300				
40	Ala	Pro	Ala	Lys	His	Leu	Ile	Ala	Ser	Met	Leu	Ser	Lys	Asn	Pro	Glu
	305					310						315				320
	Asp	Arg	Pro	Ser	Leu	Asp	Asp	Ile	Ile	Arg	His	Asp	Phe	Phe	Leu	Gln
				325						330					335	
45	Gly	Phe	Thr	Pro	Asp	Arg	Leu	Ser	Ser	Ser	Cys	Cys	His	Thr	Val	Pro
				340					345					350		
	Asp	Phe	His	Leu	Ser	Ser	Pro	Ala	Lys	Asn	Phe	Phe	Lys	Lys	Ala	Ala
			355					360					365			
	Ala	Ala	Leu	Phe	Gly	Gly	Lys	Lys	Asp	Lys	Ala	Arg	Tyr	Ile	Asp	Thr
			370				375					380				
50	His	Asn	Arg	Val	Ser	Lys	Glu	Asp	Glu	Asp	Ile	Tyr	Lys	Leu	Arg	His
	385					390					395					400
	Asp	Leu	Lys	Lys	Thr	Ser	Ile	Thr	Gln	Gln	Pro	Ser	Lys	His	Arg	Thr
				405						410					415	
55	Asp	Glu	Glu	Leu	Gln	Pro	Pro	Thr	Thr	Thr	Val	Ala	Arg	Ser	Gly	Thr
				420					425					430		
	Pro	Ala	Val	Glu	Asn	Lys	Gln	Gln	Ile	Gly	Asp	Ala	Ile	Arg	Met	Ile
			435					440					445			
	Val	Arg	Gly	Thr	Leu	Gly	Ser	Cys	Ser	Ser	Ser	Ser	Glu	Cys	Leu	Glu
		450				455						460				
60	Asp	Ser	Thr	Met	Gly	Ser	Val	Ala	Asp	Thr	Val	Ala	Arg	Val	Leu	Arg
	465					470						475				480

- 4/5 -

Gly Cys Leu Glu Asn Met Pro Glu Ala Asp Cys Ile Pro Lys Glu Gln
 485 490 495
 Leu Ser Thr Ser Phe Gln Trp Val Thr Lys Trp Val Asp Tyr Ser Asn
 500 505 510
 5 Lys Tyr Gly Phe Gly Tyr Gln Leu Ser Asp His Thr Val Gly Val Leu
 515 520 525
 Phe Asn Asn Gly Ala His Met Ser Leu Leu Pro Asp Lys Lys Thr Val
 530 535 540
 10 His Tyr Tyr Ala Glu Leu Gly Gln Cys Ser Val Phe Pro Ala Thr Asp
 545 550 555 560
 Ala Pro Glu Gln Phe Ile Ser Gln Val Thr Val Leu Lys Tyr Phe Ser
 565 570 575
 His Tyr Met Glu Glu Asn Leu Met Asp Gly Gly Asp Leu Pro Ser Val
 580 585 590
 15 Thr Asp Ile Arg Arg Pro Arg Leu Tyr Leu Leu Gln Trp Leu Lys Ser
 595 600 605
 Asp Lys Ala Leu Met Met Leu Phe Asn Asp Gly Thr Phe Gln Val Asn
 610 615 620
 20 Phe Tyr His Asp His Thr Lys Ile Ile Ile Cys Ser Gln Asn Glu Glu
 625 630 635 640
 Tyr Leu Leu Thr Tyr Ile Asn Glu Asp Arg Ile Ser Thr Thr Phe Arg
 645 650 655
 Leu Thr Thr Leu Leu Met Ser Gly Cys Ser Ser Glu Leu Lys Asn Arg
 660 665 670
 25 Met Glu Tyr Ala Leu Asn Met Leu Leu Gln Arg Cys Asn
 675 680 685

(2) INFORMATION FOR SEQ ID NO:3:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 2789 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

40 GGCACGAGGT TGGGTGCTAT TCGGCACCAG AGGCAAGGGT GCGAGGACCA CGGCCGGCTC 60
 GGACGTGTGA CCGCGCCTAG GGGGTGGCAG CGGGCAGTGC GGGGCGGCAA GGCGACCATG 120
 GAGCTTTTGC GGACTATCAC CTACCAGCCA GCCGCCAGCA CCAAATGTG CGAGCAGGCG 180
 TCGGGCAAGG GTTGC GGAGC AGACTCGAAG AAGAAGCGGC CGCCGCAGCC CCCCAGGAA 240
 TCGCAGCCAC CTCAGTCCCA GGCGCAAGTG CCCCAGGCGG CCCCACCA CCATCACCAC 300
 45 CATTTCGCACT CGGGGCCGGA GATCTCGCGG ATTATCGTCG ACCCCACGAC TGGGAAGCGC 360
 TACTGCCGGG GCAAAGTGCT GGGAAAGGGT GGCTTTGCAA AATGTTACGA GATGACAGAT 420
 TTGACAAATA ACAAAGTCTA CGCCGCAAAA ATTATTCCTC ACAGCAGAGT AGCTAAACCT 480
 CATCAAAGGG AAAAGATTGA CAAAGAAATA GAGCTTCACA GAATTCTTCA TCATAAGCAT 540
 GTAGTGCAGT TTTACCACTA CTTCGAGGAC AAAGAAAACA TTTACATTCT CTTGGAATAC 600
 50 TGCAGTAGAA GGTCAATGGC TCATATTTTG AAAGCAAGAA AGGTGTTGAC AGAGCCAGAA 660
 GTTCGATACT ACCTCAGGCA GATTGTGTCT GGACTGAAAT ACCTTCATGA ACAAGAAATC 720
 TTGCACAGAG ATCTCAAAC T AGGGAAC TTTTATTAATG AAGCCATGGA ACTAAAAGTT 780
 GGGGACTTCG GTCTGGCAGC CAGGCTAGAA CCCTTGGAAC ACAGAAGGAG AACGATATGT 840
 GGTACCCCAA ATTATCTCTC TCCTGAAGTC CTCAACAAAC AAGGACATGG CTGTGAATCA 900
 55 GACATTTGGG CCCTGGGCTG TGTAATGTAT ACAATGTTAC TAGGGAGGCC CCCATTTGAA 960
 ACTACAAATC TCAAAGAAAC TTATAGGTGC ATAAGGGAAG CAAGGTATAC AATGCCGTCC 1020
 TCATTGCTGG CTCCTGCCAA GCACTTAATT GCTAGTATGT TGTCCAAAAA CCCAGAGGAT 1080
 CGTCCAGTT TGGATGACAT CATTGACAT GACTTTTTTT TGCAGGGCTT CACTCCGGAC 1140
 AGACTGTCTT CTAGCTGTTG TCATACAGTT CCAGATTTCC ACTTATCAAG CCCAGCTAAG 1200
 60 AATTTCTTTA AGAAAGCAGC TGCTGCTCTT TTTGGTGGCA AAAAAGACAA AGCAAGATAT 1260
 ATTGACACAC ATAATAGAGT GTCTAAAGAA GATGAAGACA TCTACAAGCT TAGGCATGAT 1320

- 5/5 -

	TTGAAAAAGA	CTTCAATAAC	TCAGCAACCC	AGCAAACACA	GGACAGATGA	GGAGCTCCAG	1380
	CCACCTACCA	CCACAGTTGC	CAGGTCTGGA	ACACCCGCAG	TAGAAAACAA	GCAGCAGATT	1440
	GGGGATGCTA	TTCCGGATGAT	AGTCAGAGGG	ACTCTTGGCA	GCTGTAGCAG	CAGCAGTGAA	1500
	TGCCTTGAAG	ACAGTACCAT	GGGAAGTGTT	GCAGACACAG	TGGCAAGGGT	TCTTCGGGGGA	1560
5	TGTCTGAAA	ACATGCCGGA	AGCTGATTGC	ATTCCCAAAG	AGCAGCTGAG	CACATCATT	1620
	CAGTGGGTCA	CCAAATGGGT	TGATTACTCT	AACAAATATG	GCTTTGGGTA	CCAGCTCTCA	1680
	GACCACACCG	TCGGTGTCCCT	TTTCAACAAT	GGTGCTCACA	TGAGCCTCCT	TCCAGACAAA	1740
	AAAACAGTTC	ACTATTACGC	AGAGCTTGGC	CAATGCTCAG	TTTTCCCAGC	AACAGATGCT	1800
	CCTGAGCAAT	TTATTAGTCA	AGTGACGGTG	CTGAAATACT	TTTCTCATTA	CATGGAGGAG	1860
10	AACCTCATGG	ATGGTGGAGA	TCTGCCTAGT	GTTACTGATA	TTCGAAGACC	TCGGCTCTAC	1920
	CTCCTTCAGT	GGCTAAAATC	TGATAAGGCC	CTAATGATGC	TCTTTAATGA	TGGCACCTTT	1980
	CAGGTGAATT	TCTACCATGA	TCATACAAAA	ATCATCATCT	GTAGCCAAAA	TGAAGAATAC	2040
	CTTCTCACCT	ACATCAATGA	GGATAGGATA	TCTACAACCT	TCAGGCTGAC	AACTCTGCTG	2100
	ATGTCTGGCT	GTTCATCAGA	ATTAAAAAAT	CGAATGGAAT	ATGCCCTGAA	CATGCTCTTA	2160
15	CAAAGATGTA	ACTGAAAGAC	TTTTTCGAATG	GACCCTATGG	GACTCCTCTT	TTCCACTGTG	2220
	AGATCTACAG	GGAAACCCAAA	AGAATGATCT	AGAGTATGTT	GAAGAAGATG	GACATGTGGT	2280
	GGTACGAAAA	CAATTCCCCT	GTGGCCTGCT	GGACTGGGTG	GAACCAGAAC	AGGCTAAGGC	2340
	ATACAGTTCT	TGACTTTGGA	CAATCCAAGA	GTGAACCAGA	ATGCAGTTTT	CCTTGAGATA	2400
	CCTGTTTTAA	AAGGTTTTTC	AGACAATTTT	GCAGAAAGGT	GCATTGATTC	TTAAATTCTC	2460
20	TCTGTTGAGA	GCATTTACGC	CAGAGGACTT	TGGAAGTGTG	AATATACTTC	CTGAAGGGGA	2520
	GGGAGAAGGG	AGGAAGCTCC	CATGTTGTTT	AAAGGCTGTA	ATTGGAGCAG	CTTTTGGCTG	2580
	CGTAACTGTG	AACTATGGCC	ATATATAATT	TTTTTTTCATT	AATTTTTGAA	GATACTTGTG	2640
	GCTGGAAAAG	TGCATTCCCT	GTTAATAAAC	TTTTTATTTA	TTACAGCCCA	AAGAGCAGTA	2700
	TTTATTATCA	AAATGTCTTT	TTTTTTATGT	TGACCATTTT	AAACCGTTGG	CAATAAAGAG	2760
25	TATGAAAACG	CAAAAAAAAA	AAAAAAAAA				2789

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/17248

A. CLASSIFICATION OF SUBJECT MATTER
 IPC(6) :Please See Extra Sheet.
 US CL :Please See Extra Sheet.
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 U.S. : 435/194, 6, 15, 320.1, 440, 252.3, 325, 317.1; 536/23.2; 530/387.9, 350; 424/94.6

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 APS, STN- MEDLINE, EMBASE, BIOSIS, WPIDS, HCAPLUS, NTIS, BIOTECHDS, SCISEARCH
 SEARCH TERMS: SERUM INDUCIBLE KINASE


C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SIMMONS et al. Identification of an Early-Growth-Response Gene Encoding a Novel Putative Protein Kinase. Mol.Cell.Biol. September 1992, Vol. 12, No.9, pages 4164-4169, see figure 4 and the entire article.	1-12
X,P	Database GenBank on STN, US National Library of Medicine (Bethesda MD), No. AF059617, April 1998.	2, 13

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *G* document member of the same patent family
---	--

Date of the actual completion of the international search 28 OCTOBER 1998	Date of mailing of the international search report 19 NOV 1998
--	--

Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer  MARYAM MONSHIPOURI Telephone No. (703) 308-0196
--	--

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/17248

A. CLASSIFICATION OF SUBJECT MATTER:
IPC (6):

C12N 9/12, 15/54, 1/21, 5/10, 15/00, 15/63; C07K16/40; A01K 38/45; C12Q 1/68, 1/48; C07K 14/00

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

435/194, 6, 15, 320.1, 440, 252.3, 325, 317.1; 536/23.2; 530/387.9, 350; 424/94.6