

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 August 2011 (11.08.2011)

(10) International Publication Number
WO 2011/097345 A1

PCT

(51) International Patent Classification:
G01N 33/574 (2006.01) *G01N 35/08* (2006.01)

(21) International Application Number:
PCT/US2011/023547

(22) International Filing Date:
3 February 2011 (03.02.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/301,400 4 February 2010 (04.02.2010) US

(71) Applicant (for all designated States except US): **VIRO-STATICS SRL** [IT/IT]; Viale Umberto 1,46 (IT).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BAEV, Denis, V.** [RU/IT]; Via Don Minzoni, I-07041 Alghero (SS) (IT).
LORI, Franco [IT/US]; 4400 East West Hwy, Bethesda, MD 20814 (US).

(74) Agent: **LOOPER, Valerie, E.**; The Law Offices Of Valerie E. Looper, 11675 Cypress Canyon Rd, San Diego, CA 92131 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: KI-67 ASSAY FOR PATIENT IMMUNE SYSTEM STATUS

(57) Abstract: Novel diagnostic assay based on the detection of Ki-67 expression to anticipate the response to anti-HIV therapy.



WO 2011/097345 A1

KI-67 ASSAY FOR PATIENT IMMUNE SYSTEM STATUS**Field of the Invention**

This invention relates to a novel assay for immune system status. The expression of Ki-67 as a proliferation marker in an infected patient's T-cells can be analyzed and the results used to predict the patient's likely disease progression and responsiveness to therapies, including drug treatments and vaccines. This method can be utilized as a guide for an individual patient's therapy management (personalized therapy) over both the short and mid term.

The invention relates to a novel test that quantifies the percentage of proliferated CD3 T cells as an immune monitoring method, and to evaluate the therapy outcome in patients suffering of chronic infections, neoplastic and allergic diseases where the expression level of Ki-67 in T-cells correlates with disease progression.

The proof of principle of the utility of this marker is demonstrated in a representative chronic disease treatment scenario: where individuals who are chronically infected with HIV undergo therapy with antiviral drugs or vaccines. The number of Ki-67+ T-cells before therapy was begun correlated with the patients' reduction of an HIV viral load after therapy commenced better than the currently accepted CD4 counts and CD38 expression levels in two independent cohorts. We found that a level of the baseline Ki-67 expression in CD3, CD4 and CD8 T cell subsets but not CD4 counts or baseline values of CD38 and other activation markers (i.e. CD69, HLA-DR and PD-1), significantly correlated with changes in plasma HIV viral loads after treatment had begun.

A known flow cytometry method can be used to detect Ki-67 expression in whole CD3 cells in donor blood. In the alternative, the Ki-67 level can be detected by ELISA or bead-array methods in lysed CD3 cells sorted from donor blood samples.

This test can be used to monitor the ability of prophylactic and therapeutic approaches to produce changes in the immune system status, or competence, of individuals with HIV/AIDS, and to predict short- and long-term success or failure of treatments, including antiretroviral therapy and immune therapies based on immune modulation and targeted immune amplification, and finally to guide a HIV patient's disease management.

5

Background of the Invention

One of the great frustrations in efforts to deal with chronic infections generally in humans, and in the field of treatment of HIV/AIDS in particular is that, while drugs and treatments such as vaccines have been developed that can perform according to the design goals, the design goals, when reached, have not cured the target conditions. As a result, powerful drugs that could both reduce viral replication to undetectable levels in the blood and increase the number of CD4 cells to a more normal range, all in a relatively short period of time, could not end the disease condition. While these two factors yield some information about a patient's current immune system status, they are insufficient to support the next phase of therapy research. Other disease markers must be developed for further progress to be made. The present inventors have focused on a means to elucidate better, more predictive information about a patient's immune system, by determining a patient's present state of immune system activation. Various markers for immune system activation, namely CD 38, CD69, HLA-DR, CD4 counts and changes in plasma HIV viral loads were compared, and baseline Ki-67 expression in CD3, CD4 and CD8 T cell subsets was found to be more predictive than CD4 counts or baseline values of CD38 and other activation markers (i.e. CD69, HLA-DR).

An individual patient's degree of immune system activation plays a major role in determining the level of HIV replication and in the eventual depletion of CD4 cells that results in progression of the disease. In 1981 the very first observation of the increased expression of a protein that functions as a T cell activation marker, T10 (designated later on as CD38) in *Pneumocystis carinii* pneumonia patients was published (Masur H et al., The New England Journal of Medicine, 1981). CD38 is a glycoprotein found on the surface of many immune system cells including CD4+, CD8+, B and natural killer cells. Masur et al. connected CD 38 with the appearance of multiple cases of opportunistic infections that occur in acquired immune deficiency patients. The critical role of immune activation in HIV disease progression and in survival of HIV-infected patients was reported first in 1999 (Janis V. Giorgi et al., The Journal of Infectious Diseases 1999;179:859-70). Giorgi and colleagues reported that high levels of expression by T Cells of both CD38 and HLA-DR (cell surface receptor proteins involved in the binding

of antigens) are associated with shorter survival of HIV-infected patients and could be considered as possible predictive markers for HIV disease progression. These are said to be more likely predictors than either than plasma HIV concentrations or tests for the virus usage of CCR5 and CXCR4 chemokine receptors. Multiple current observations have reported that CD38 expression could be considered as a promising marker to evaluate progression of the disease with recommendations to use this marker as a diagnostic value (Coetzee LM, Cytometry B Clin Cytom. 2009 Nov;76(6):375-84.) in routine clinical testing.

The present inventors have explored the use of another protein. The Ki-67 protein (also known as MKI67) is a cellular marker strictly associated with cell proliferation. During interphase (collectively, phases G₁, S and G₂), Ki-67 can be exclusively detected within the cell nucleus, and during mitosis most of the protein is located on the surface of the chromosomes. Ki-67 protein is present during all active phases of the cell cycle (G₁, S, G₂, and mitosis), but is undetectable in resting cells (G₀). Maximal virus replication occurs in G₁ and S phases (Foli A. et al., J Infect Dis. 2007 Nov 1;196(9):1409-15.), when Ki-67 is also up-regulated, so that the only phase when Ki-67 is present and maximal virus replication is not occurring is the relatively short, intermediate, G₂ phase.

The persistent immune activation characteristic of chronic infections, and very particularly evident in HIV infection eventually results in the depletion of CD4 T cells because it progressively erodes the naive T cell pool. Compelling evidence of the role of Ki-67 in HIV-disease progression was reported by M. Lederman and colleagues (Sieg SF, Lederman MM et al., J Infect Dis. 2005 Jul 1;192(1):62-70.)

Thus, the inventors posited that the comparison of activation markers might yield better ways to evaluate the clinical status of HIV-infected patients and to monitor the effects of anti-HIV therapy.

Brief Description of the Drawings

Fig.1 Cohort 1: VS411 study. Ki-67 expression is dramatically up-regulated in HIV infected patients (1B) up to 2...10% of total T cells compared to 0.05...0.5% in non-infected individuals (1A) and decreases to 0.5...2% after 24 weeks of VS411 AV-HALT treatment (1C). The Ki-67 decrease is statistically significance ($p < 0.001$) by paired Student's t-test. Patient's PBMCs were stained for surface T cell markers and for Ki-67 by intracellular staining and analyzed on BD LSR-II flow cytometer. One representative dot-

plot out of 10 subjects in figure 1A and out of 32 subjects in figures 1B and 1C is presented, respectively.

Fig.2 Cohort 1: VS411 study. The expression of Ki-67 is strongly correlates with the reduction of HIV viral load. 28 subjects from Phase 2A VS-411 study were analyzed for Ki-67 expression on CD3+, CD3+CD4+ and CD3+CD8+ T cells and the Pearson's bi-variate correlation was applied between the baseline values of Ki-67 positive cells percentage and HIV viral load decrease from baseline to last day of anti-HIV treatment. The correlations were characterized by $r^2 = 0,272$ for the CD3+ subset ($p=0,004$)(2A), $r^2 = 0,19$ for CD3+CD4+ ($p=0,018$)(2B) and $r^2 = 0,21$ for CD3+CD8+ ($p=0,013$)(2C) subsets, respectively.

Fig.3 Cohort 2: RIGHT702 study. Ki-67 baseline expression in CD3 cells strongly correlates with HIV viral load decrease for short-term and long-term therapy outcomes. T cells from 7 patients from RIGHT702 study were analyzed for Ki-67 by flow cytometry analysis and correlation was found with baseline values of Ki-67 and HIV viral load changes on week 12 (A) and week 24(B) with $r^2 = 0,852$ ($p=0,003$) and $r^2 = 0,945$ ($p=0,001$), respectively

Summary of the Invention

We describe here a diagnostic assay for Ki-67 monitoring, the steps comprising:

- a. Isolation of peripheral blood mononuclear cells (PBMC)
- b. Staining of T-cells among the PBMC for surface markers
- c. Intracellular staining of T cells for Ki-67
- d. Measurement of the percentage of Ki-67 positive cells by flow cytometry.

Does this yield actively dividing immune system cells v. quiescent immune system cells?

The diagnostic assay proof of principle was demonstrated in a chronic disease, namely HIV-1/AIDS. In this assay, the cells may be frozen PBMCs derived from HIV-

infected patients and Ki-67 expression in different T cell subsets is detected by the intracellular staining with anti-Ki-67 antibodies and analyzed by multi-color flow cytometry, as demonstrated in our examples for two independent cohorts of VS411 Phase2A and RIGHT702 studies, or freshly isolated PBMCs, or sorted CD3 cells. Sorted
5 cells could be lysed and the Ki-67 expression could also be detected by ELISA or any other method. In the present application, Ki-67 was analyzed in CD3, CD4 and CD8 T cells. For HIV-1, a dramatic increase in number of circulating Ki-67 positive T cells is a particular indicator for disease progression.

This assay relies on detection of Ki-67 in patients PBMC before the anti-HIV
10 treatment is started.

To develop the assay we have started with a chronic infection that represents a world pandemic emergency: HIV infection. An advantage of this assay is that it removes a major barrier to drug development for chronic infections. In this disease, the development of effective therapy has been hampered for the past 25 years due to the absence of a rapid
15 test that can correlate immune system markers with disease progression in a case where the anti-HIV therapy is applied. Here we describe a test that yields a quantitative determination of a specific marker in T cells that can predict the therapy outcome in HIV infection.

Another advantage of the invention is that it modifies the standard flow cytometry
20 assays, such as CD4 enumeration and analysis of CD38 expression, which did not yield a good predictive potential (CD4 counts) for the therapy outcome and good correlation (CD38) with HIV-1 viral load changes

To investigate the differences between the newly developed Ki-67 assay and the standard approaches we tested the PBMC from 32 HIV-1-infected individuals from
25 Phase2A VS411 study with both assays. Surprisingly, we found that a level of the baseline Ki-67 expression in CD3, CD4 and CD8 T cell subsets but not CD4 counts or baseline values of CD38 and other activation markers (i.e. CD69, HLA-DR), significantly correlated with changes of plasma HIV viral loads. The analysis of 7 additional PBMC samples from an independent cohort of HIV-1-infected individuals from RIGHT702 study
30 confirmed the correlation between the baseline Ki-67 expression and decrease of HIV-1 viral load in all patients. These results demonstrate that the quantity of Ki-67+ T-cells detected with this test can be used as predictive tool to determine the immunological status of chronically-infected patients and to predict the therapy outcome. In addition, a certain range of baseline number of Ki-67+CD3+ cells can be correlated with a certain

range of decrease of viral load ("VL"). For example, the percentage of CD3+Ki-67+ cells below 2-2,5% were associated with VL decrease for more than 1,5 logs.

The Ki-67-based assay provides an immune monitoring tool for patients suffering from HIV/AIDS to (1) determine their specific immunological status; (2) to predict their individual disease progression; (3) for individual patient management (for example for determining when to start therapy or how intensely to monitor a patient); and (4) for the design, testing and screening of therapies based on anti-viral drugs, immune modulation, targeted immune amplification, and prophylactic and/or therapeutic vaccines.

Description of the Invention

Although several lines of evidence suggests that circulating Ki-67⁺ T-cells play a critical role as a reservoir of HIV infection in humans and SIV in the macaque model, the precise immune correlates of HIV progression and responsiveness to the therapy remain to be identified.

The flow cytometry is currently widely used to identify HIV-specific effects on T cells in humans. Current approaches include CD4 enumeration and monitoring of activation markers, like CD38 and HLA-DR, which have a predictive potential in terms of disease progression but not in the evaluation of possible anti-HIV therapy outcome. Conversely, the present invention quantifies the Ki-67 positive T cells more likely representing the HIV-related pool of T cells.

In this assay peripheral blood mononuclear cells are first collected from donors or derived from cryopreserved blood samples and stained for surface T cell markers and for Ki-67 followed by the flow cytometry analysis of the Ki-67 expression level.

In the present application, the percentage of Ki-67⁺ T cells quantified by flow cytometry was compared with other activation markers such as CD38, CD69, PD-1 and HLA-DR to discern which parameter correlates with the viral load decrease in chronically HIV-infected patients receiving the anti-HIV therapy.

We analyzed the relationships among various baseline expression levels of activation markers and plasma viral loads and viral load decreases and found that the baseline percentage of Ki-67⁺ in CD3, CD4 and CD8 subsets but not other markers have a strong and statistically significant correlation with HIV viral load decrease.

The present study is the first to demonstrate that the baseline percentages of Ki-67+ cells in CD3, CD4 and CD8 subsets inversely correlated with the decrease of plasma HIV viral load in HIV-infected individuals undergo the AV-HALT anti-viral therapy.

These results are consistent with the observation that there is a direct connection between the number of circulating S-phase (Ki-67+) T cells and the level of plasma HIV viremia as it was reported by many investigators. Results from a recent study also provides the evidence that even the baseline expression of Ki-67 in total CD3 subset could serve as a predictive marker for the changes of HIV plasma levels in patients receiving anti-viral therapy.

In the RIGHT702 study the baseline percentage of Ki-67+ T cell showed even better potential to predict both short-term (12 weeks) and long-term (24-48week) therapy outcomes in HIV-infected patients.

In conclusion, our finding represents the first evaluation of predictive potential of Ki-67 marker in HIV-infected individuals for the therapy outcome evaluation. The lower percentage of Ki-67+ T-cells is associated with higher decrease in plasma viremia and better therapy outcome.

The Ki-67 assay will be useful to evaluate the status of the immune system and the recovery of its function in patients treated with antiretroviral drugs as well as to assess the efficacy of prophylactic and therapeutic vaccines.

Examples

The following examples illustrate the practice of various aspects of the present inventions. They do not limit the inventions, or the claims, which follow them

Example 1: Cohort 1, VS411 study

Methods: Blood samples from 32 patients receiving VS411 (a novel combination of an antiviral and a cytostatic drug) during a Phase 2A, dose finding study were analyzed by 10-color flow cytometry for activation and proliferation T cell markers. Pearson correlation analysis was performed for baseline and on-treatment values of T cell markers expression versus HIV viral load (VL) and VL changes.

Results: After treatment had begun, KI67 expression in CD3+ cells level was reduced from $3,1 \pm 1,68\%$ to $2,18 \pm 0,9\%$ ($p=0.0012$). PD-1, CD38 and HLA-DR expressions

were down-regulated from $15,33 \pm 6,12\%$ to $11,32 \pm 5,01\%$, from $58,48 \pm 14,81\%$ to $52,8 \pm 14,25\%$, and from $27,5 \pm 10,01\%$ to $22,2 \pm 7,55\%$, respectively, with p values <0.005 . In CD4⁺ T cell subset the Ki-67 expression level was reduced from $3,23 \pm 1,89\%$ to $2,42 \pm 1,11\%$ ($p=0.016$). PD-1 and HLA-DR expressions were down-regulated from $15,3 \pm 7,53\%$ to $11,67 \pm 6,59\%$ and from $11,94 \pm 5,54\%$ to $9,56 \pm 4,45\%$, respectively, with p values <0.01 . In CD8⁺ T cell subset the Ki-67 expression level was reduced from $2,79 \pm 1,85\%$ to $1,89 \pm 1,09\%$ ($p=0.0025$). PD-1, CD38 and HLA-DR expressions were down-regulated from $16,75 \pm 8,11\%$ to $11,93 \pm 6,57\%$, from $59,13 \pm 15,5\%$ to $50,1 \pm 15,01\%$ and from $36,03 \pm 12,69\%$ to $30,02 \pm 10,68$, respectively, with p values <0.005 .

Most importantly, a negative correlation between baseline expression levels of Ki-67 in either CD3, CD4 or CD8 subsets and VL reduction was found. That is, higher levels of Ki-67 pre-treatment corresponded with poorer patient response. The correlations were characterized by $r^2 = 0,272$ for the CD3⁺ subset ($p=0,004$)(Fig 2A), $r^2 = 0,19$ for CD3+CD4⁺ ($p=0,018$)(Fig.2B) and $r^2 = 0,21$ for CD3+CD8⁺ ($p=0,013$)(Fig.2C) subsets, respectively. The number of Ki-67⁺ T-cell correlated with VL reduction better than the currently accepted CD4 counts and CD38 expression levels. That is, CD4 count and CD38 expression levels pre-treatment did not predict patient response as well as Ki-67 expression level.

Example 2: Cohort 2, RIGHT702 study

Methods: Blood samples from 7 patients from RIGHT 702 study were analyzed by multicolor flow cytometry for activation and proliferation T cell markers. Pearson correlation analysis was performed for baseline values of KI67 versus HIV viral load (VL) and VL changes.

Results: A negative correlation between baseline expression levels of Ki-67 in either CD3, CD4 or CD8 subsets and VL reduction was found both for short-term and long-term therapy outcomes. That is, higher levels of Ki-67 expression before treatment had begun corresponded with poorer patient response for up to 48 weeks.

For the CD3⁺ subset the correlations were characterized by $r^2 = 0,852$ for short-term ($p=0,004$)(Fig 3A) and by $r^2 = 0,945$ for the long-term therapy outcome($p=0,001$)(Fig.3B), respectively. Same negative correlation were found for the baseline Ki-67⁺ with VL in CD4 and CD8 subsets.

What is Claimed is:

1. A diagnostic assay for Ki-67 monitoring in HIV-1 infected subjects, the steps comprising:

- 5 a. Isolation of PBMS from donor blood
- b. Staining T-cells for surface markers
- c. Intracellular staining T cells for Ki-67
- d. Measuring Ki-67 level by flow cytometry.
- 10 2. The assay of Claim 1, wherein the cells are freshly derived or frozen
3. The assay of Claim 1, wherein CD3 cells were sorted from blood
4. The assay of Claim 3, wherein the Ki-67 staining is performed in cell lysates.
5. The assay of Claim 4, wherein the Ki-67 detection is performed by any other method, i.e. confocal microscopy, imaging, ELISA or multiplexed bead array (CBA, Luminex)
- 15 6. A diagnostic assay for immune system activation in humans, comprising quantification of expression of Ki-67 glycoprotein in one or more of the group consisting of CD3, CD4 and CD8 subsets of human PBMC.
7. The assay of Claim 6, wherein the subset is the total CD3 set of human PBMC.
- 20 8. The assay of Claim 6, wherein the expression of Ki-67 glycoprotein is quantified for cells in the S phase of cellular proliferation.




Fig. 1




Fig. 2




Fig. 3

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 11/23547

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - G01N 33/574, 35/08 (2011.01)

USPC - 435/7.23; 436/52

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

USPC 435/7.23; 436/52

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
USPC 435/6; 435/7.1; 435/7.4; 436/86 (see search terms below)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
PubWEST(DB=PGPB,USPT,USOC,EPAB,JPAB; PLUR=YES; OP=ADJ), Google Scholar(immune system activation Ki-67, Ki-67 HIV-1 PBMC T-cells CD3, Ki-67 ELISA)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	SACHSENBERG et al. Turnover of CD4+ and CD8+ T Lymphocytes in HIV-1 Infection as Measured by Ki-67 Antigen. J Exp Med, 1998, Vol 187, No 8, pp 1295-1303; pages 1295-1296, Study Subjects, Cell Separation and Antibody Staining, Flow Cytometry	1-3 ---- 4-5
X --- Y	LENG et al. Immune activation correlates better than HIV plasma viral load with CD4 T-cell decline during HIV infection. Journal of Acquired Immune Deficiency Syndromes, 2001, Vol 24, No 4, pp 1-11; abstract; pg 2, para 1, pg 6, para 2	6 ----- 7-8
Y	US 2005/0215479 A1 (BOYD et al.) 29 September 2005 (29.09.2005) para [0429]	4-5
Y	US 2007/0122378 A1 (FREEMAN et al.) 31 May 2007 931.05.2007) para [0184]	7
Y	SIEG et al. Peripheral S-Phase T Cells in HIV Disease Have a Central Memory Phenotype and Rarely Have Evidence of Recent T Cell Receptor Engagement. The Journal of Infectious Diseases, 2005, Vol 192, pp 62-70; abstract	8
Y	KLEIN et al. A new quantitative test method for cell proliferation based on detection of the Ki-67 protein. JOURNAL OF MATERIALS SCIENCE: MATERIALS IN MEDICINE, 2000, Vol 11, pp 125-132; abstract	5

☐ Further documents are listed in the continuation of Box C. ☐

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

18 March 2011 (18.03.2011)

Date of mailing of the international search report

19 APR 2011

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774