



US 20070292930A1

(19) **United States**

(12) **Patent Application Publication**

Shu et al.

(10) **Pub. No.: US 2007/0292930 A1**

(43) **Pub. Date: Dec. 20, 2007**

(54) **METHOD OF PRODUCING RECOMBINANT ASPERGILLUS NIGER BETA-GLUCOSIDASE AND AN AROMA SPREADING PLANT**

of international application No. PCT/IL00/00758, filed on Nov. 15, 2000, which is a continuation of application No. 09/443,338, filed on Nov. 19, 1999, now abandoned.

(75) Inventors: **Wei Shu**, Saskatoon (CA); **Daniel L. Siegel**, Rechovot (IL); **Ira Marton**, Rechovot (IL); **Ben-Ami Bravdo**, Rehovot (IL); **Mara Dekel**, Rechovot (IL); **Oded Shoseyov**, Karmei Yosef (IL)

Publication Classification

(51) **Int. Cl.**
C12P 7/10 (2006.01)
C12P 7/06 (2006.01)
(52) **U.S. Cl.** **435/165; 435/161**

Correspondence Address:

Martin D. Moynihan
PRTSI, Inc.
P.O. BOX 16446
Arlington, VA 22215 (US)

(57) **ABSTRACT**

A polypeptide having β -glucosidase enzymatic activity, a polynucleotide encoding the polypeptide, a nucleic acid constructs carrying the polynucleotide, transformed or infected cells, such as yeast cells, and transgenic organisms expressing the polynucleotide and various uses of the polypeptide, the polynucleotide, cells and/or organisms, including, producing a recombinant polypeptide having the β -glucosidase enzymatic activity, increasing the level of aroma compounds in alcoholic beverages, as well as other fermentation products of plant material, hydrolyzing cellobiose and thus increasing the level of fermentable glucose, increasing the production of alcohol, such as ethanol from plant material, increasing the aroma released from a plant or a plant product, and hydrolysis or transglycosylation of glycosides.


(73) Assignee: **Yissum Research Development Company of the Hebrew University of Jerusalem**, Jerusalem (IL)

(21) Appl. No.: **11/785,365**

(22) Filed: **Apr. 17, 2007**

Related U.S. Application Data

(63) Continuation of application No. 10/130,150, filed on May 16, 2002, now Pat. No. 7,223,902, filed as 371






Fig. 1c




Fig. 1b




Fig. 1a




Figure 2B




Figure 2A




Fig. 3




Fig. 4

2821 GCTCTGCGCAACCGATCCTGCGCTGCCGGTGGCGGTCTGGCGGCAACCCTCGCCTGTACG 2880
lySerAlaGlnProIleLeuProAlaGlyGlyGlyProGlyGlyAsnProArgLeuTyrA

2881 ACGAGCTCATCCGCGTGTGAGTACCATCAAGAACACCGGCAAGGTTGCTGGTGATGAAG 2940
 spGluLeuIleArgValSerValThrIleLysAsnThrGlyLysValAlaGlyAspGluV

2941 TTCCCAACTGGTAAGTAAACATGAGGTCCGAACGAGGTTGAACAAAGCTAATCAGTCGC 3000
 alProGlnLeu Intron#6

3001 AGTATGTTTCCCTTGGCGGTCCCAATGAGCCCAAGATCGTGTGCTGCGTCAATTCGAGCGCA 3060
 TyrValSerLeuGlyGlyProAsnGluProLysIleValLeuArgGlnPheGluArgI

3061 TCACGCTGCAGCCGTCGGAGGAGACGAAGTGGAGCAGACTCTGACGCGCCGTGACCTTG 3120
 leThrLeuGlnProSerGluGluThrLysTrpSerThrThrLeuThrArgArgAspLeuA

3121 CAAACTGGAATGTTGAGAAGCAGGACTGGGAGATTACGTGCTATCCCAAGATGGTGTGTTG 3180
 laAsnTrpAsnValGluLysGlnAspTrpGluIleThrSerTyrProLysMetValPheV

3181 TCGGAAGCTCCTCGCGGAAGCTGCCGCTCCGGGCGTCTCTGCCTACTGTTCACTAAATAG 3240
 alGlySerSerSerArgLysLeuProLeuArgAlaSerLeuProThrValHis***

3241 CTCTCAAATGGTATACCATGATGGCCGTGGTATATGAATTAATGATTTATGCCAACAGCA 3300

3301 AGACCACTGTAGATGTAGATGTAGAATGAGTATTGCGTAGTAGCGTGTAGATGATGATAC 3360

3361 AAGCGATCCGACACATGGTAGGAAGAGTGGCGCTAGTTGGGGCGGAAACCAAGCGACGTC 3420

3421 ATCCGCTGCCGACTTCGCCAGTCTTTCTTCTTTTCTTCTTCAGCCTTCTTCTCCGCTTA 3480

3481 ATCCAGCAACCATTGCCAATGCCTCTACAACAATAATTGCCATAATACTTACTCCTA 3540

3541 TTCAATATATACACCACAATCTCGACATAATCACACAAGCCTGAACACACGAGCAACCAT 3600

3601 GCCCTCTCCCGATCCTCCAGCCCCAGCGATACGACCCTTCCAACCACCATAACAGCGCT 3660

3661 CCTCATCTACCCAGCGACCCTAATCGTGGGATCACTTCTTCCGTCTCTCCCACCGC 3720

3721 ACAAGGCACACGCGACGACGGCTCCAGCACCTCCACCCACACGTCGAGCCCTAGCCCC 3780

3781 GTCCATCGCGTCAGACCTCAACCTCTECTTTCCTCCGCCGCGCCCGTCAACTACTTCGC 3840

3841 TCGCAAAGACAACATCTTCAATCTATATTCGTCAAAGTCGGC 3885

Fig. 5a (continued)




Fig. 5b




Figure 6B




Figure 6A




Fig. 7




Fig. 8




Fig. 9







Fig. 10

Fig. 11a 35SΩ + bg11

Fig. 11b 35SΩ + Ce11 + bg11

Fig. 11c 35SΩ + Ce11 + bg11 + HDEL





WT CB10 CB11 CBT3 CBT8 CBT15 B1 B34 An Glu

Fig. 14




Fig. 15

**METHOD OF PRODUCING RECOMBINANT
ASPERGILLUS NIGER BETA-GLUCOSIDASE AND
AN AROMA SPREADING PLANT**

RELATED APPLICATIONS

[0001] This application is a continuation of pending U.S. patent application Ser. No. 10/130,150 filed May 16, 2002, which is a U.S. National Phase of PCT Patent Application No. PCT/IL00/00758 filed Nov. 15, 2000, which is a continuation of U.S. patent application Ser. No. 09/443,338 filed Nov. 19, 1999, now abandoned. The contents of the above Applications are incorporated herein by reference.

FIELD AND BACKGROUND OF THE
INVENTION

[0002] The present invention relates to a polypeptide having β -glucosidase enzymatic activity, to a polynucleotide encoding the polypeptide, to nucleic acid constructs carrying the polynucleotide, to transformed or infected cells, such as yeast cells, and organisms expressing the polynucleotide and to various uses of the polypeptide, the polynucleotide, cells and/or organisms, including, but not limited to, producing a recombinant polypeptide having β -glucosidase enzymatic activity, increasing the level of aroma compounds in alcoholic beverages, as well as other fermentation products of plant material, hydrolyzing cellobiose and thus increasing the level of fermentable glucose, to increase production of alcohol, such as ethanol from plant material, increasing the aroma released from a plant or a plant product, and hydrolysis or transglycosylation of glycosides.

[0003] Abbreviations used herein include: BGL1—*Aspergillus niger* B1 β -glucosidase; bgl1—a cDNA encoding same; 2FGlcF—2-deoxy-2-fluoro β -glucosyl fluoride; DNP—2,4-dinitrophenol; DNPGlc—2,4-dinitrophenyl β -D-glucopyranoside; pNP—p-nitrophenol; pNPGlc—p-nitrophenyl β -D-glucopyranoside; MUGlc—4-methylumbelliferyl- β -D-glucopyranoside; YNB—yeast nitrogen base without amino acids; and X-glu—5-bromo-4-chloro-3-indolyl β -D-glucopyranoside.

[0004] β -Glucosidases (EC 3.2.1.21; β -D-glucoside glucohydrolase) play a number of different important roles in biology, including the degradation of cellulosic biomass by fungi and bacteria, degradation of glycolipids in mammalian lysosomes and the cleavage of glucosylated flavonoids in plants. These enzymes are therefore of considerable industrial interest, not only as constituents of cellulose-degrading systems, but also in the food industry (2, 3).

[0005] *Aspergillus* species are known as a useful source of β -glucosidases (4-6), and *Aspergillus niger* is by far the most efficient producer of β -glucosidase among the microorganisms investigated (4). Shoseyov et al. (7) have previously described a β -glucosidase from *Aspergillus niger* B1 (CMI CC 324626) which is active at low pHs, as well as in the presence of high ethanol concentrations. This enzyme effectively hydrolyzes flavor-compound glycosides in certain low-pH products, such as wine and passion fruit juice, thereby enhancing their flavor (8-12), and is particularly attractive for use in the food industry, as *A. niger* is considered non-toxic (3). In addition, β -glucosidase was found useful in enzymatic synthesis of glycosides (13-15). Other *A. niger* β -glucosidases have also been purified (16-18), however, differences in their properties have been reported,

including ranges of molecular weights (116-137 kDa), isoelectric points (pI values of 3.8-4) and pH optima (3.4-4.5). Indeed, at least two β -glucosidases, with distinct substrate specificities, have been identified in commercial *A. niger* β -glucosidase preparations (19). Attempts to clear this confusion by cloning and expression of a functional *A. niger* α -glucosidase gene in *S. cerevisiae* has been previously reported (20), however the protein was not characterized, and the sequence was not published.

[0006] Glycosidases have been assigned to families on the basis of sequence similarities, there now being some 77 different such families defined containing over 2,000 different enzymes (21, see also the CAZy (Carbohydrate Active EnZymes) website, at the Architecture of Fonction de Macromolécules Biologiques of the Centre National de la Recherche Scientifique website. With the exception of the glucosylceramidases (Family 30), all simple β -glucosidases belong to either Family 1 or 3. Family 1 contains enzymes from bacteria, plants and mammals, including also 6-phospho-glucosidases and thioglucosidases. Furthermore, most Family 1 enzymes also have significant galactosidase activity. Family 3 contains β -glucosidases and hexosaminidases of fungal, bacterial and plant origin. Enzymes of both families hydrolyze their substrates with net retention of anomeric configuration, presumably via a two-step, double-displacement mechanism, involving two key active site carboxylic acid residues (for reviews of mechanism, see 22-24). In the first step, one of the carboxylic acids (the nucleophile) attacks at the substrate anomeric center, while the other (the acid/base catalyst) protonates the glycosidic oxygen, thereby assisting the departure of the aglycone. This results in the formation of a covalent α -glycosyl-enzyme intermediate. In a second step this intermediate is hydrolyzed by general base-catalyzed attack of water at the anomeric center of the glycosyl-enzyme, to release the β -glucose product and regenerate free enzyme. Both the formation and the hydrolysis of this intermediate proceed via transition states with substantial oxocarbenium ion character.

[0007] Given that Family 3 contains fungal enzymes of similar mass, including those from other *Aspergillus* sp., it is likely that the *Aspergillus niger* β -glucosidase would be a member of this family. Mechanistic information on this family is relatively sparse: the best characterized being the glycosylated 170 kDa β -glucosidase from *Aspergillus wentii*. By labeling the active site with conduritol B-epoxide, this enzyme was shown to carry out hydrolysis, with net retention of anomeric configuration. This study has demonstrated that the labeled aspartic acid residue was the same as that derivatized by the slow substrate D-glucal (1, 25). Furthermore, it was shown that the 2-deoxyglucosyl-enzyme, trapped by use of D-glucal, was kinetically identical to that formed during the hydrolysis of PNP-2-deoxy- β -D-glucopyranoside (26). Further detailed kinetic analysis of the enzyme was performed by Legler et al. (27), including measurement of Hammett relationships, kinetic isotope effects and studies of the binding of potent reversible inhibitors, such as gluconolactone and nojirimycin.

[0008] While reducing the present invention to practice, the β -glucosidase protein was isolated from *Aspergillus niger*, purified, cloned, sequenced, expressed in yeast host cells and its enzymatic function characterized. In addition, the protein as well as signal peptide fused thereto and

optionally an endoplasmic reticulum retaining peptide fused thereto were expressed in transgenic plants and the release of aroma substances therefrom following homogenization monitored. The enzyme encoded by the isolated gene, as described above, is of known usefulness in plant and/or plant products, as well as in biotechnological processes, including the food industry. Several unexpected advantages were uncovered, including, but not limited to, pH and temperature stability of the β -glucosidase from *Aspergillus niger*, requirement for a signal peptide for obtaining catalytic activity when expressed in plants. Advantage for an endoplasmic retaining peptide or for a lack thereof when expressed in plants, depending on the application.

SUMMARY OF THE INVENTION

[0009] According to one aspect of the present invention there is provided an isolated nucleic acid comprising a genomic, complementary or composite polynucleotide preferably being derived from *Aspergillus niger*, encoding a polypeptide having a β -glucosidase catalytic activity and preferably further encoding, in frame, a signal peptide and an endoplasmic reticulum retaining peptide.

[0010] According to another aspect of the present invention there is provided a recombinant protein comprising a polypeptide having a β -glucosidase catalytic activity, the polypeptide is preferably derived from *Aspergillus niger* and it preferably fused to a signal peptide and optionally also to an endoplasmic reticulum retaining peptide.

[0011] According to yet another aspect of the present invention there is provided a nucleic acid construct comprising the isolated nucleic acid described herein.

[0012] According to still another aspect of the present invention there is provided host cell or an organism, such as a plant, comprising the nucleic acid or nucleic acid construct described herein.

[0013] According to further features in preferred embodiments of the invention described below, the polynucleotide is as set forth in SEQ ID NOs:1, 3 or a portion thereof.

[0014] According to still further features in the described preferred embodiments, the nucleic acid construct further comprising at least one cis acting control element for regulating expression of the polynucleotide.

[0015] According to still further features in the described preferred embodiments, the host cell is selected from the group consisting of a prokaryotic cell and a eukaryotic cell.

[0016] According to still further features in the described preferred embodiments the prokaryotic cell is *E. coli*.

[0017] According to still further features in the described preferred embodiments the eukaryotic cell is selected from the group consisting of a yeast cell, a fungous cell, a plant cell and an animal cell.

[0018] According to still further features in the described preferred embodiments the polypeptide is as set forth in SEQ ID NO: 2 or a portion thereof having the β -glucosidase catalytic activity.

[0019] According to an additional aspect of the present invention there is provided a method of producing recombinant β -glucosidase, the method comprising the step of introducing, in an expressible form, a nucleic acid construct

into a host cell, the nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, encoding a polypeptide having a β -glucosidase catalytic activity and preferably further encoding, in frame, a signal peptide and an endoplasmic reticulum retaining peptide.

[0020] According to further features in preferred embodiments of the invention described below, the method further comprising the step of extracting the polypeptide having the β -glucosidase catalytic activity.

[0021] According to yet an additional aspect of the present invention there is provided a method of producing a recombinant β -glucosidase overexpressing cell, the method comprising the step of introducing, in an overexpressible form, a nucleic acid construct into a host cell, the nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, encoding a polypeptide having a β -glucosidase catalytic activity and preferably further encoding, in frame, a signal peptide and an endoplasmic reticulum retaining peptide.

[0022] According to still an additional aspect of the present invention there is provided a method of increasing a level of at least one fermentation substance in a fermentation product, the method comprising the step of fermenting a glucose containing fermentation starting material by a yeast cell overexpressing a nucleic acid construct including a genomic, complementary or composite polynucleotide being preferably derived from *Aspergillus niger*, encoding a polypeptide having a β -glucosidase catalytic activity and preferably further encoding, in frame, a signal peptide and an endoplasmic reticulum retaining peptide, thereby increasing the level of the at least one fermentation substance in the fermentation product.

[0023] According to a further aspect of the present invention there is provided a method of increasing a level of at least one fermentation substance in a fermentation product, the method comprising the step of fermenting a plant derived glucose containing fermentation starting material by a yeast cell, the plant overexpressing a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, encoding a polypeptide having a β -glucosidase catalytic activity and preferably further encoding, in frame, a signal peptide and an endoplasmic reticulum retaining peptide, thereby increasing the level of the at least one fermentation substance in the fermentation product.

[0024] According to a further aspect of the present invention there is provided a method of increasing a level of at least one aroma substance in a plant derived product, the method comprising the step of incubating a glucose containing plant starting material with a yeast cell overexpressing a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, encoding a polypeptide having a β -glucosidase catalytic activity and preferably further encoding, in frame, a signal peptide and an endoplasmic reticulum retaining peptide, thereby increasing the level of the at least one aroma substance in the plant derived product.

[0025] According to yet a further aspect of the present invention there is provided a method of increasing a level of at least one aroma substance in a plant derived product, the

method comprising the step of incubating a glucose containing plant starting material with a yeast cell, said plant overexpressing a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, encoding a polypeptide having a β -glucosidase catalytic activity and preferably further encoding, in frame, a signal peptide and an endoplasmic reticulum retaining peptide, thereby increasing the level of the at least one aroma substance in the plant derived product.

[0026] According to still further features in the described preferred embodiments the plant derived product is a fermentation product, such as, but not limited to, an alcoholic beverage.

[0027] According to still a further aspect of the present invention there is provided a method of increasing a level of free glucose in a glucose containing fermentation starting material, the method comprising the step of fermenting the glucose containing fermentation starting material by a cell overexpressing a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, encoding a polypeptide having a β -glucosidase catalytic activity and preferably further encoding, in frame, a signal peptide and an endoplasmic reticulum retaining peptide, thereby increasing the level of the free glucose in the glucose containing fermentation starting material.

[0028] According to another aspect of the present invention there is provided a method of increasing a level of free glucose in a plant derived glucose containing fermentation starting material, the method comprising the step of fermenting the plant derived glucose containing fermentation starting material by a cell, the plant overexpressing a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, encoding a polypeptide having a β -glucosidase catalytic activity and preferably further encoding, in frame, a signal peptide and an endoplasmic reticulum retaining peptide, thereby increasing the level of the free glucose in the plant.

[0029] According to yet another aspect of the present invention there is provided a method of increasing a level of free glucose in a plant, the method comprising the step of overexpressing in the plant a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, encoding a polypeptide having a β -glucosidase catalytic activity and preferably further encoding, in frame, a signal peptide and an endoplasmic reticulum retaining peptide, thereby increasing the level of the free glucose in the plant.

[0030] According to still another aspect of the present invention there is provided a method of producing an alcohol, the method comprising the step of fermenting a glucose containing fermentation starting material by a cell overexpressing a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, encoding a polypeptide having a β -glucosidase catalytic activity and preferably further encoding, in frame, a signal peptide and an endoplasmic reticulum retaining peptide, and extracting the alcohol therefrom.

[0031] According to an additional aspect of the present invention there is provided a method of producing an

alcohol, the method comprising the step of fermenting a plant derived glucose containing fermentation starting material by a cell, the plant overexpressing a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, encoding a polypeptide having a β -glucosidase catalytic activity and preferably further encoding, in frame, a signal peptide and an endoplasmic reticulum retaining peptide, and extracting the alcohol therefrom.

[0032] According to an additional aspect of the present invention there is provided a method of producing an aroma spreading plant, the method comprising the step of overexpressing in the plant a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, encoding a polypeptide having a β -glucosidase catalytic activity and preferably further encoding, in frame, a signal peptide and an endoplasmic reticulum retaining peptide, thereby increasing aroma spread from the plant.

[0033] According to further features in preferred embodiments of the invention described below, overexpressing the nucleic acid construct is performed in a tissue specific manner.

[0034] According to still further features in the described preferred embodiments overexpressing the nucleic acid construct is limited to at least one tissue selected from the group consisting of flower, fruit, seed, root, stem, pollen and leaves.

[0035] The present invention successfully addresses the shortcomings of the presently known configurations by providing a polypeptide having β -glucosidase enzymatic activity, a polynucleotide encoding the polypeptide, a nucleic acid constructs carrying the polynucleotide, transformed or infected cells, such as yeast cells, and organisms expressing the polynucleotide and various uses of the polypeptide, the polynucleotide, cells and/or organisms, including, but not limited to, producing a recombinant polypeptide having β -glucosidase enzymatic activity, increasing the level of aroma compounds in alcoholic beverages, as well as other fermentation products of plant material, hydrolyzing cellobiose and thus increasing the level of fermentable and/or free glucose, to increase production of a fermentation product, such as ethanol from plant material, increasing the aroma released from a plant or a plant product, and hydrolysis or transglycosylation of glycosides.

BRIEF DESCRIPTION OF THE DRAWINGS

[0036] The invention is herein described, by way of example only, with reference to the accompanying drawings. With specific reference now to the drawings in detail, it is stressed that the particulars shown are by way of example and for purposes of illustrative discussion of the preferred embodiments of the present invention only, and are presented in the cause of providing what is believed to be the most useful and readily understood description of the principles and conceptual aspects of the invention. In this regard, no attempt is made to show structural details of the invention in more detail than is necessary for a fundamental understanding of the invention, the description taken with the drawings making apparent to those skilled in the art how the several forms of the invention may be embodied in practice.

[0037] In the drawings:

[0038] FIGS. 1a-c demonstrate plasmid maps employed as expression vectors for *bgl1* cDNA. FIG. 1a—*E. coli* expression vector containing *bgl1* cDNA, inserted into the NcoI/BamHI sites of pET3d. FIG. 1b—*S. cerevisiae* expression vector containing *bgl1* cDNA, inserted into the HindIII/BamHI sites of pYES2-*bgl1* plasmid. FIG. 1c—*P. pastoris* expression vector containing *bgl1* cDNA, inserted into the EcoRI/BamHI sites of pHIL-S1.

[0039] FIGS. 2a-b demonstrates SDS-PAGE analysis of active protein samples eluted from a MONO-Q™ (Amersham Biosciences Inc, Piscataway, N.J.) anion exchange column, stained with coomassie blue (FIG. 2a), or β -glucosidase zymogram (FIG. 2b) using MUGlc as a substrate. Lanes (for both FIGS. 2a and 2b): 1—Electroeluted band of BGL1 from preparative PAGE-SDS gel stabs; 2, 3, 4, 5—acetone precipitates from MONO-Q™ (Amersham Biosciences Inc, Piscataway, N.J.) anion exchange column separation of BGL1.

[0040] FIG. 3 demonstrates SDS-PAGE analysis of purified β -glucosidase by MONO-Q™ (Amersham Biosciences Inc, Piscataway, N.J.) anion exchange and RESOURCE-S™ (Amersham Biosciences Inc, Piscataway, N.J.) cation exchange columns. Lanes: 1—crude (27.5 μ g protein); 2—active fraction after MONO-Q™ (Amersham Biosciences Inc, Piscataway, N.J.) anion exchange (7 μ g protein); and 3—active fraction after RESOURCE-S™ (Amersham Biosciences Inc, Piscataway, N.J.) cation exchange column (10 μ g protein).

[0041] FIG. 4 demonstrates SDS-PAGE analysis of β -glucosidase deglycosylated by N-glycosidase-F. Lanes: 1—molecular weight marker; 2—native β -glucosidase; and 3—deglycosylated protein.

[0042] FIG. 5a demonstrates the DNA (SEQ ID NO: 3) and amino acid (SEQ ID NO: 2) sequences of *bgl1*. Amino acid sequences determined by Edman degradation are underlined. DNA sequences of introns are underlined. Signal peptide is indicated by italic letters.

[0043] FIG. 5b. demonstrates *bgl1* gene organization. Exons (E1-7) are indicated by filled boxes, introns by solid lines, restriction sites and the stop codon by arrows.

[0044] FIG. 6a demonstrates a Western blot analysis of recombinant BGL1 expressed in *S. cerevisiae*. Lanes: 1—native BGL1 (positive control); 2—total protein extract of *S. cerevisiae* expressing recombinant BGL1; 3—total protein extract of *S. cerevisiae* without the *bgl1* expression vector (negative control).

[0045] FIG. 6b demonstrates a Western blot analysis of recombinant BGL1 secreted from *P. pastoris*. Lanes: 1—molecular weight marker; 2—medium supernatant of *P. pastoris* expressing recombinant BGL1; 3—medium supernatant of *P. pastoris* host without the vector (negative control).

[0046] FIG. 7 demonstrates proton-NMR spectra, illustrating the stereochemical course of pNPGlc hydrolysis by *A. niger* β -glucosidase. Spectra are for the anomeric proton region of the substrate at different time intervals relative to addition of the enzyme.

[0047] FIG. 8 demonstrates inactivation of recombinant BGL1 by 2FGlcF. Pure enzyme was incubated in the pres-

ence of various concentrations of the inactivator, and residual enzyme activity was determined at different time intervals. Residual activity is presented, semilogarithmically, versus time, in the presence of the indicated concentrations of inactivator.

[0048] FIG. 9 demonstrates reactivation of 2-deoxy-2-fluoroglucosyl-recombinant BGL 1 by linamarin. Activity is plotted versus incubation time in the presence of the indicated concentrations of linamarin.

[0049] FIG. 10 demonstrates the stability of recombinant *A. niger* β -glucosidase at various temperatures. Activity is calculated as percent of a recombinant enzyme solution kept at 4° C.

[0050] FIGS. 11a-c show schematic depictions of expression cassettes used for expression of *A. niger* β -glucosidase in tobacco plants. FIG. 11a—a cassette encoding BGL1 without a signal peptide (see, SEQ ID NO:13 for the nucleotide sequence and SEQ ID NO:14 for the amino acid sequence); FIG. 11b—a cassette encoding a BGL1 fused to a Cell1 signal peptide for secretion into the apoplast (see, SEQ ID NO:15 for the nucleotide sequence and SEQ ID NO:16 for the amino acid sequence); and FIG. 11c—a cassette encoding a BGL1 fused to Cell1 signal peptide as in FIG. 11b and in addition to HDEL (SEQ ID NO:17) ER-retaining peptide at the C-terminus for accumulation in the ER (see, SEQ ID NO:18 for the nucleotide sequence and SEQ ID NO:19 for the amino acid sequence).

[0051] FIG. 12 demonstrate PCR amplification results of *bgl1* cDNA indicating the presence of *bgl1* cDNA in transgenic plants. CB10 and CB11—transgenic plants transformed with *bgl1* and Cell1 signal peptide without HDEL, SEQ ID NO:17 ER retaining peptide. CBT3, CBT8 and CBT15—different transgenic lines transformed with *bgl1*, Cell1 signal peptide and HDEL, SEQ ID NO:17. B1—a transgenic plants transformed with *bgl1*. 1 kb-1 kb DNA marker. WT—wild type non transgenic plant. pETB1-*bgl1* plasmid DNA.

[0052] FIGS. 13a-b show Western blot analyses of transgenic plants containing BGL1 without signal peptide (13a), and BGL1 with Cell1 signal peptide (13b), with and without HDEL, SEQ ID NO:17 ER retaining peptide. An glucopurified *A. niger* beta-glucosidase. WT—nontransgenic control plant. B1, B15, B16, B20, B27, B33 and B34—different transgenic lines transformed with *bgl1*. CBT1, CBT 3, CBT 7 and CBT 8—different transgenic lines transformed with *bgl1*, Cell1 signal peptide and HDEL, SEQ ID NO:17. CB10 and CB12—transgenic plants transformed with *bgl1* and Cell1 signal peptide without HDEL, SEQ ID NO:17 ER retaining peptide.

[0053] FIG. 14 show activity gel analysis of transgenic tobacco plant extracts in SDS-PAGE incubated with MUGlu. WT—non-transgenic control plant. CB10 and CB11—two independent lines of transgenic plants expressing BGL1 fused to Cell1 signal peptide (without HDEL, SEQ ID NO:17). CBT3, CBT8 and CBT15—different lines of transgenic plants expressing BGL1 fused to Cell1 signal peptide at the N terminus and HDEL, SEQ ID NO:17 ER retaining peptide at the C terminus. B1 and B34—transgenic plant expressing BGL1 without signal peptide or HDEL, SEQ ID NO:17 ER retaining peptide and which were positive for BGL1 protein in Western blot analysis. An Glu-control *A. niger* native beta-glucosidase.

[0054] FIG. 15 demonstrates level of BGL1 activity in different transgenic plants. WT—non-transgenic control plant. B1 and B21—transgenic plants expressing BGL1 without signal peptide or HDEL, SEQ ID NO:17 ER retaining peptide and which were positive for BGL1 in Western blot analysis. CBT8, CBT21, CBT0 and CBT15—-independent lines of transgenic plants expressing BGL1 fused to Cel1 signal peptide at the N terminus and HDEL, SEQ ID NO:17 ER retaining peptide at the C terminus. CB12, CB13, CB14 and CB15—four independent lines of transgenic plants expressing BGL1 fused to Cel1 signal peptide (without HDEL, SEQ ID NO:17).

DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0055] The present invention is of a polypeptide having β -glucosidase enzymatic activity, a polynucleotide encoding the polypeptide, a nucleic acid constructs carrying the polynucleotide, transformed or infected cells, such as yeast cells, and organisms expressing the polynucleotide and various uses of the polypeptide, the polynucleotide, cells and/or organisms, including, but not limited to, producing a recombinant polypeptide having the β -glucosidase enzymatic activity, increasing the level of aroma compounds in alcoholic beverages, as well as other fermentation products of plant material, hydrolyzing cellobiose and thus increasing the level of fermentable glucose, increasing the production

of alcohol, such as ethanol from plant material, increasing the aroma released from a plant or a plant product, and hydrolysis or transglycosylation of glycosides.

[0056] The principles and operation of the present invention may be better understood with reference to the drawings and accompanying descriptions.

[0057] Before explaining at least one embodiment of the invention in detail, it is to be understood that the invention is not limited in its application to the details of the components set forth in the following description or exemplified in the examples that follow. The invention is capable of other embodiments or of being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology employed herein is for the purpose of description and should not be regarded as limiting.

[0058] According to one aspect of the present invention there is provided an isolated nucleic acid comprising a genomic, complementary or composite polynucleotide encoding a polypeptide having a β -glucosidase catalytic activity. Preferably the polynucleotide is derived from *Aspergillus niger*, however other sources are applicable. These include all isolated polynucleotides encoding polypeptide having β -glucosidase catalytic activity. Such polynucleotides and polypeptides identified by their GenBank Accession Nos. are listed in Table 1 below, all of which can be used while implementing the present invention.

TABLE 1

Organism	Accession numbers of cDNA and their encoded beta-glucosidases (EC.3.2.1.21)	
	SWISS-PROT	EMBL
<i>Acetobacter xylinus</i>	O24749	AB003689; AB010645
<i>Agrobacterium</i> sp.	P12614	M19033; AAA22085.1
<i>Agrobacterium tumefaciens</i>	P27034	M59852; AAA22082.1
<i>Arabidopsis thaliana</i>	O82772, O24433, O23656	AF082157; AF082158; AC009327; U72153; U72155; AC020665; AC066691; D64088, BAA10968.1
<i>Aspergillus aculeatus</i>	P48825	AB003470
<i>Aspergillus kawachi</i>	P87076	AJ132386; CAB75696.1
<i>Aspergillus niger</i> B1	Q9P456	AF268911
<i>Aspergillus niger</i> AMS1	Q38786, Q9ZP27	X78433; AF082991
<i>Avena sativa</i>		AF090429; AAF21798.1
<i>Azospirillum irakense</i>	Q03506	M96979; AAA22266.1
<i>Bacillus circulans</i>	Q9ZNN7	AB009411; BAA36161.1; AB009410
<i>Bacillus polymyxa</i>	P22073, P22505	M60210; M60211
<i>Bacillus subtilis</i>	P40740	Z34526; CAA84287.1
<i>Bacillus subtilis</i>	P42403	D30762; BAA06429.1
<i>Bacteroides fragilis</i>	O31356	AF006658; AAB62870.1
<i>Bifidobacterium breve</i>	P94248, O08487	D84489; D88311
<i>Botryotinia fuckeliana</i>		AJ130890; CAB61489.1
<i>Brassica napus</i>	Q42618	X82577
<i>Brassica nigra</i>	O24434	U72154
<i>Butyrivibrio fibrisolvens</i>	P16084	M31120; AAA23008.1
<i>Caldocellum saccharolyticum</i>	P10482	X12575; CAA31087.1
<i>Caldicellulosiruptor</i> sp. 14B	Q9ZEN0	AJ131346
<i>Candida wickerhamii</i>	Q12601	U13672
<i>Cavia porcellus</i>	P97265	U50545
<i>Cellulomonas bizotaea</i>	O51843	AF005277; AAC38196.1
<i>Cellulomonas fimi</i>	Q46043	M94865
<i>Cellvibrio gilvus</i>	P96316	D14068; BAA03152.1
<i>Chryseobacterium meningosepticum</i>	O30713	AF015915
<i>Clostridium stercorarium</i>	O08331	Z94045
<i>Clostridium thermocellum</i>	P26208	X60268; CAA42814.1
<i>Clostridium thermocellum</i>	P14002	X15644; CAA33665.1

TABLE 1-continued

Organism	Accession numbers of cDNA and their encoded beta-glucosidases (EC.3.2.1.21)	
	SWISS-PROT	EMBL
<i>Coccidioides immitis</i>	O14424	U87805; AF022893
<i>Costus speciosus</i>	Q42707	D83177
<i>Dalbergia cochinchinensis</i>	Q9SPK3	AF163097
<i>Dictyostelium discoideum</i>	Q23892	L21014
<i>Digitalis lanata</i>	Q9ZPB6	AJ133406
<i>Erwinia chrysanthemi</i>	Q46684	U08606; AAA80156.1
<i>Erwinia herbicola</i>	Q59437	X79911; CAA56282.1
<i>Escherichia coli</i>	P33363	U15049; AAB38487.1
<i>Escherichia coli</i> K12/MG1655	E65074, Q46829	U28375; AE000373
<i>Glycine max</i>		AF000378; AAD09291.1
<i>Hansenula anomala</i>	P06835	X02903; CAA26662.1
<i>Homo sapiens</i>		AJ278964; CAC08178.1
<i>Hordeum vulgare</i>	Q40025	L41869
<i>Humicola grisea</i> var. <i>thermoidea</i>	O93784	AB003109
<i>Kluyveromyces marxianus</i>	P07337	X05918; CAA29353.1
<i>Lactobacillus plantarum</i>	O86291	Y15954; AJ250202; CAB71149.1
<i>Manihot esculenta</i>	Q40283	X94986
<i>Microbispora bispora</i>	P38645	M97265; AAA25311.1
<i>Nicotiana tabacum</i>	O82151	AB017502; BAA33065.1
<i>Orpinomyces</i> sp. PC-2		AF016864; AAD45834.1
<i>Oryza sativa</i>	Q42975	U28047
<i>Paenibacillus polymyxa</i>	P22073	M60210; AAA22263.1
<i>Paenibacillus polymyxa</i>	P22505	M60211; AAA22264.1
<i>Phaeosphaeria avenaria</i>		AJ276675; CAB82861.1
<i>Phanerochaete chrysosporium</i>	O74203	AF036872; AF036873
<i>Pichia anomala</i> (<i>Candida pelliculosa</i>)	P06835	X02903
<i>Pinus contorta</i>		AF072736; AAC696.1
<i>Polygonum tinctorium</i>		AB003089; BAA78708.1
<i>Prunus avium</i>	Q43014	U39228
<i>Prunus serotina</i>	Q43073, Q40984	U50201; U26025
<i>Prevotella albensis</i> M384		AJ276021; CAC07184.1
<i>Prevotella ruminicola</i>	Q59716	U35425
<i>Pyrococcus furiosus</i>	Q51723	AF013169; U37557
<i>Ruminococcus albus</i>	P15885 O66050	X15415; CAA33461.1 U92808
<i>Saccharomycopsis fibuligera</i>	P22506	M22475; AAA34314.1
<i>Saccharomycopsis fibuligera</i>	P22507	M22476; AAA34315.1
<i>Saccharopolyspora erythraea</i>	O70021	Y14327
<i>Salmonella typhimurium</i>	Q56078	D86507; BAA13102.1
<i>Schizophyllum commune</i>	P29091	M27313; AAA33925.1
<i>Schizosaccharomyces pombe</i>		AL355920; CAB91163.1
<i>Secale cereale</i>		AF293849; AAG00614.1
<i>Septoria lycopersici</i>	Q99324	U24701; U35462
<i>Sorghum bicolor</i>	Q41290	U33817
<i>Spodoptera frugiperda</i>	O61594	AF052729
<i>Streptomyces coelicolor</i> A3(2)		AL121596; CAB56653.1
<i>Streptomyces reticuli</i>	Q9X9R4	AJ009797
<i>Streptomyces rochei</i> A2	Q55000	X74291
<i>Streptomyces</i> sp. QM-B814	Q59976	Z29625
<i>Thermoanaerobacter brockii</i>	P96090, Q60026	Z56279; Z56279
<i>Thermobifida fusca</i> ER1		AF086819; AAF37727.1
<i>Thermococcus</i> sp.	O08324	Z70242
<i>Thermotoga maritima</i>	Q08638	X74163; CAA52276.1
<i>Thermotoga neapolitana</i>	O33843, Q60038	Z97212; Z77856; CAB10165.1
<i>Thermus</i> sp. Z-1	Q9RA58	AB034947
<i>Thermus thermophilus</i>	Q9X9D4	Y16753
<i>Trichoderma reesei</i> (<i>Hypocrea jecorina</i>)	Q12715, O93785	U09580; AAA18473.1, AB003110
<i>Trifolium repens</i>	P26204	X56734; CAA40058.1
<i>Trifolium repens</i>	P26205	X56733; CAA40057.1
<i>Tropaeolum majus</i>	O82074	AJ006501; CAA07070.1
<i>Zea mays</i>	P49235, Q41761	X74217, U25157; CAA52293.1
Unidentified bacterium	Q60055	U33816, U44087, U44773 U12011

[0059] As used herein in the specification and in the claims section that follows, the term "isolated" refers to a biological component (such as a nucleic acid or protein or organelle) that has been substantially separated or purified

away from other biological components in the cell of the organism in which the component naturally occurs, i.e., other chromosomal and extra-chromosomal DNA and RNA, proteins and organelles. Nucleic acids and proteins that have

been "isolated" include nucleic acids and proteins purified by standard purification methods. The term also embraces nucleic acids and proteins prepared by recombinant expression in a host cell as well as chemically synthesized nucleic acids.

[0060] As used herein and in the claims section that follows the terms and phrases "polynucleotide" and "polynucleotide sequence" are used interchangeably and refer to a nucleotide sequence which can be DNA or RNA of, for example, genomic or synthetic origin, which may be single- or double-stranded, and which may represent the sense or antisense strand. Similarly, the terms "polypeptide" and "polypeptide sequence" are interchangeably used herein and refer to an amino acid sequence of any length.

[0061] As used herein in the specification and in the claims section that follows, the phrase "complementary polynucleotide sequence" includes sequences, which originally result from reverse transcription of messenger RNA using a reverse transcriptase or any other RNA dependent DNA polymerase. Such sequences can be subsequently amplified in vivo or in vitro using a DNA dependent DNA polymerase.

[0062] As used herein in the specification and in the claims section that follows, the phrase "genomic polynucleotide sequence" includes sequences which originally derive from a chromosome and reflect a contiguous portion of a chromosome.

[0063] As used herein in the specification and in the claims section that follows, the phrase "composite polynucleotide sequence" includes sequences which are at least partially complementary and at least partially genomic. A composite sequence can include some exonal sequences required to encode the polypeptide having the β -glucosidase catalytic activity, as well as some intronic sequences interposing therebetween. The intronic sequences can be of any source, including of other genes, and typically will include conserved splicing signal sequences. Such intronic sequences may further include cis acting expression regulatory elements, as hereinbelow described.

[0064] As used herein in the specification and in the claims section that follows, the phrase "having a β -glucosidase catalytic activity" refers to a polypeptide sequence, protein or fragments thereof capable of serving as catalysts to a chemical reaction involving hydrolysis of the O-glycosidic bond of glucosides, the result of which is the release of a β -D-glucose residue(s), or an aglycon, in addition to the β -D-glucose residue. Specifically, hydrolysis by retaining enzymes is performed while maintaining the β -configuration of the anomeric center of the carbohydrate. A wide specificity for β -glucosides exists, thus, some examples also hydrolyze one or more of the following: β -D-galactosides, α -L-arabinosides, β -D-xylosides, and β -D-fucosides.

[0065] As used herein the term "catalyst" refers to a substance that accelerates a chemical reaction, but is not consumed or changed permanently thereby.

[0066] As used herein the term "glucoside" refers to a compound of at least two monomers, at least one of which is a glucose, including a glycoside bond. Examples of glucosides include, but are not limited to, glucose containing backbones, such as the diglucose cellobiose, and the glucose polymer, cellulose.

[0067] According to preferred embodiments, the polynucleotide according to this aspect of the present invention encodes a polypeptide as set forth in SEQ ID NO:2 or a portion thereof which retains β -glucosidase catalytic activity.

[0068] Alternatively or additionally, the polynucleotide according to this aspect of the present invention is as set forth in SEQ ID NO:1, 3 or a portion thereof, the portion encodes a polypeptide retaining β -glucosidase catalytic activity.

[0069] In a broader aspect the polynucleotides according to the present invention encode a polypeptide which is at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or more, say 95%-100% homologous to SEQ ID NO:2 as determined using the BestFit software of the Wisconsin sequence analysis package, utilizing the Smith and Waterman algorithm, where gap creation penalty equals 8 and gap extension penalty equals 2.

[0070] According to preferred embodiments, the polynucleotides according to the broader aspect of the present invention encodes a polypeptide as set forth in SEQ ID NOs:1 or 3 or a portion thereof which retains activity.

[0071] Alternatively or additionally, the polynucleotides according to this broader aspect of the present invention are hybridizable with SEQ ID NOs: 1 or 3.

[0072] Hybridization for long nucleic acids (e.g., above 200 bp in length) is effected according to preferred embodiments of the present invention by stringent or moderate hybridization, wherein stringent hybridization is effected by a hybridization solution containing 10% dextrane sulfate, 1 M NaCl, 1% SDS and 5×10^6 cpm ^{32}P labeled probe, at 65° C., with a final wash solution of 0.2 \times SSC and 0.1% SDS and final wash at 65° C.; whereas moderate hybridization is effected by a hybridization solution containing 10% dextrane sulfate, 1 M NaCl, 1% SDS and 5×10^6 cpm ^{32}P labeled probe, at 65° C., with a final wash solution of 1 \times SSC and 0.1% SDS and final wash at 50° C.

[0073] Yet alternatively or additionally, the polynucleotides according to this broad aspect of the present invention is preferably at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or more, say 95%-100%, identical with SEQ ID NOs: 1 or 3 as determined using the BestFit software of the Wisconsin sequence analysis package, utilizing the Smith and Waterman algorithm, where gap weight equals 50, length weight equals 3, average match equals 10 and average mismatch equals -9.

[0074] Thus, this broad aspect of the present invention encompasses (i) polynucleotides as set forth in SEQ ID NOs:1 or 3; (ii) fragments thereof; (iii) sequences hybridizable therewith; (iv) sequences homologous thereto; (v) sequences encoding similar polypeptides with different codon usage; (vi) altered sequences characterized by mutations, such as deletion, insertion or substitution of one or more nucleotides, either naturally occurring or man induced, either randomly or in a targeted fashion.

[0075] According to another aspect of the present invention there is provided a nucleic acid construct comprising the isolated nucleic acid described herein.

[0076] According to a preferred embodiment, the nucleic acid construct according to this aspect of the present inven-

tion further comprising at least one cis acting control (regulatory) element for regulating the expression of the isolated nucleic acid. Such cis acting regulatory elements include, for example, promoters, which are known to be sequence elements required for transcription, as they serve to bind DNA dependent RNA polymerase, which transcribes sequences present downstream thereof. Further details relating to various regulatory elements are described hereinbelow.

[0077] While the isolated nucleic acid described herein is an essential element of the invention, it is modular and can be used in different contexts. The promoter of choice that is used in conjunction with this invention is of secondary importance, and will comprise any suitable promoter. It will be appreciated by one skilled in the art, however, that it is necessary to make sure that the transcription start site(s) will be located upstream of an open reading frame. In a preferred embodiment of the present invention, the promoter that is selected comprises an element that is active in the particular host cells of interest. These elements may be selected from transcriptional regulators that activate the transcription of genes essential for the survival of these cells in conditions of stress or starvation, including the heat shock proteins.

[0078] A construct according to the present invention preferably further includes an appropriate selectable marker. In a more preferred embodiment according to the present invention the construct further includes an origin of replication. In another most preferred embodiment according to the present invention the construct is a shuttle vector, which can propagate both in *E. coli* (wherein the construct comprises an appropriate selectable marker and origin of replication) and be compatible for propagation in cells, or integration in the genome, of an organism of choice, such as a plant. The construct according to this aspect of the present invention can be, for example, a plasmid, a bacmid, a phagemid, a cosmid, a phage, a virus or an artificial chromosome.

[0079] According to an additional aspect of the present invention there is provided a recombinant protein comprising a polypeptide having a β -glucosidase catalytic activity. The polypeptide is preferably derived from an *Aspergillus niger* and preferably includes a signal peptide and optionally an endoplasmic reticulum retaining peptide.

[0080] According to preferred embodiments, the polypeptide according to this aspect of the present invention is as set forth in SEQ ID NO:2 or a portion thereof which retains β -glucosidase catalytic activity.

[0081] SEQ ID NO:2 of *A. niger* β -glucosidase is similar to the amino acid sequence of the β -glucosidase of *A. kawachii*. However, while the former is highly stable at wide range of temperatures and pH treatments, the latter is relatively unstable, and thus has certain disadvantages, rendering its use for the purpose of the present invention as is further detailed and described hereinunder, unfeasible and/or much less attractive.

[0082] Recently, Iwashita and coworkers have published the sequence of a β -glucosidase (GenBank/EMBL AB003470) obtained from *Aspergillus kawachii* strain: IF04308. Sequence comparison between *Aspergillus kawachii* β -glucosidase and *A. niger* β -glucosidase revealed that the two share 98% homology.

[0083] Enzymes of the two *Aspergillus* sp. contain seven cysteine residues and identical number of glycosylation sites, while differing in their degree of glycosylation (35).

[0084] The physical and kinetic properties of three β -glucosidases from *Aspergillus kawachii* were described (35), and the three were shown to be products of the same gene, differing solely by the degree of glycosylation. The three purified *A. kawachii* β -glucosidases were readily inactivated, even at moderate pH and temperature conditions. In sharp distinction, while examining the stability of the recombinant *A. niger* β -glucosidase according to the present invention under conditions identical to those described by Iwashita et al. and as described hereinbelow in the Examples section, revealed that the enzyme is highly stable, retaining majority of the enzymatic activity even after 1 hour incubation at 60° C. (68% activity, as defined by percent activity of an enzyme kept at 4° C.).

[0085] Thus, despite the similarity between the *A. kawachii* and *A. niger* β -glucosidases, the *A. niger* enzyme unexpectedly exhibits significantly higher thermal and pH stability.

[0086] According to yet another aspect of the present invention there is provided a host cell comprising a nucleic acid construct as described herein. The term "host cell" refers to a recipient of a heterologous nucleic acid, which host cell can be either a prokaryotic cell, such as *E. coli*, or a eukaryotic cell, such as a yeast cell, a filamentous fungus cell, a plant cell or an animal cell. Examples for a yeast cell include, but not limited to, *Pichia* sp. such as *P. pastoris*, and *Saccharomyces* sp. such as *S. cerevisiae*.

[0087] As used herein and in the claims section which follows, the term "heterologous" when used in context of a nucleic acid sequence or a protein found within a plant, plant derived tissue or plant cells, or alternatively, within a eukaryotic cell, such as yeast, or a prokaryotic cell such as bacteria, refers to nucleic acid or amino acid sequences typically not native to the plant, plant derived tissue or plant cells, or alternatively, to the eukaryotic cell, such as yeast, or the prokaryotic cell, such as bacteria. Interchangeably, nucleic acid or amino acid sequences typically not native to the plant, plant derived tissue or plant cells, or alternatively, to the eukaryotic cell, such as yeast, or the prokaryotic cell, such as bacteria, are referred to by "recombinant nucleic acid" and "recombinant protein", respectively. Thus, a recombinant nucleic acid is one that has a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques.

[0088] As used herein in the specification and in the claims section that follows, the term "eukaryotic cell" refers to a cell containing a diploid genome through at least a portion of its life cycle, having membrane-bound nucleus with chromosomes made of DNA, with cell division involving a form of mitosis in which spindles are involved. Possession of a eukaryote type of cell characterizes the four kingdoms, Protocista, Fungi, Plantae and Animalia.

[0089] As used herein in the specification and in the claims section that follows, the term "prokaryotic cell"

refers to various bacteria and blue-green algae, characterized by the absence of the nuclear organization, mitotic capacities and complex organelles that typify the eukaryote superkingdom. Examples of prokaryotic cell according to the present invention are bacteria, such as, but not limited to, *E. coli*.

[0090] According to still another aspect of the present invention there is provided an organism comprising a nucleic acid construct as described herein, such as, but not limited to, a plant. Such an organism is said to be transformed or virally infected.

[0091] As used herein the term “transformed” and its conjugations such as transformation, transforming and transform, all relate to the process of introducing heterologous nucleic acid sequences into a cell or an organism, which nucleic acid sequences are propagatable to the offspring. The term thus reads on, for example, “genetically modified”, “transgenic” and “transfected”, which may be used herein to further describe and/or claim the present invention. The term relates both to introduction of a heterologous nucleic acid sequence into the genome of an organism and/or into the genome of a nucleic acid containing organelle thereof, such as into a genome of chloroplast or a mitochondrion.

[0092] As used herein the phrase “viral infected” includes infection by a virus carrying a heterologous nucleic acid sequence. Such infection typically results in transient expression of the nucleic acid sequence, which nucleic acid sequence is typically not integrated into a genome and therefore not propagatable to offspring, unless further infection of such offspring is experienced.

[0093] There are various methods of introducing foreign genes into both monocotyledonous and dicotyledonous plants (Potrykus, I., *Annu. Rev. Plant. Physiol., Plant. Mol. Biol.* (1991) 42:205-225; Shimamoto et al., *Nature* (1989) 338:274-276). The principle methods of causing stable integration of exogenous DNA into plant genomic DNA include two main approaches:

[0094] (i) *Agrobacterium*-mediated gene transfer: Klee et al. (1987) *Annu. Rev. Plant Physiol.* 38:467-486; Klee and Rogers in *Cell Culture and Somatic Cell Genetics of Plants*, Vol. 6, *Molecular Biology of Plant Nuclear Genes*, eds. Schell, J., and Vasil, L. K., Academic Publishers, San Diego, Calif. (1989) p. 2-25; Gatenby, in *Plant Biotechnology*, eds. Kung, S, and Arntzen, C. J., Butterworth Publishers, Boston, Mass. (1989) p. 93-112.

[0095] (ii) direct DNA uptake: Paszkowski et al., in *Cell Culture and Somatic Cell Genetics of Plants*, Vol. 6, *Molecular Biology of Plant Nuclear Genes* eds. Schell, J., and Vasil, L. K., Academic Publishers, San Diego, Calif. (1989) p. 52-68; including methods for direct uptake of DNA into protoplasts, Toriyama, K. et al. (1988) *Bio/Technology* 6:1072-1074. DNA uptake induced by brief electric shock of plant cells: Zhang et al. *Plant Cell Rep.* (1988) 7:379-384. From et al. *Nature* (1986) 319:791-793. DNA injection into plant cells or tissues by particle bombardment, Klein et al. *Bio/Technology* (1988) 6:559-563; McCabe et al. *Bio/Technology* (1988) 6:923-926; Sanford, *Physiol. Plant.* (1990) 79:206-209; by the use of micropipette systems: Neuhaus et al., *Theor. Appl. Genet.* (1987) 75:30-36; Neuhaus and Spangenberg, *Physiol. Plant.* (1990) 79:213-217; or by the direct incubation of DNA with germinating pollen, DeWet et

al. in *Experimental Manipulation of Ovule Tissue*, eds. Chapman, G. P. and Mantell, S. H. and Daniels, W. Longman, London, (1985) p. 197-209; and Ohta, *Proc. Natl. Acad. Sci. USA* (1986) 83:715-719.

[0096] The *Agrobacterium* system includes the use of plasmid vectors that contain defined DNA segments that integrate into the plant genomic DNA. Methods of inoculation of the plant tissue vary depending upon the plant species and the *Agrobacterium* delivery system. A widely used approach is the leaf disc procedure, which can be performed with any tissue explant that provides a good source for initiation of whole plant differentiation. Horsch et al. in *Plant Molecular Biology Manual A5*, Kluwer Academic Publishers, Dordrecht (1988) p. 1-9. A supplementary approach employs the *Agrobacterium* delivery system in combination with vacuum infiltration. The *Agrobacterium* system is especially viable in the creation of transgenic dicotyledonous plants.

[0097] There are various methods of direct DNA transfer into plant cells. In electroporation, the protoplasts are briefly exposed to a strong electric field. In microinjection, the DNA is mechanically injected directly into the cells using very small micropipettes. In microparticle bombardment, the DNA is adsorbed on microprojectiles such as magnesium sulfate crystals or tungsten particles, and the microprojectiles are physically accelerated into cells or plant tissues.

[0098] Following transformation plant propagation is exercised. The most common method of plant propagation is by seed. Regeneration by seed propagation, however, has the deficiency that due to heterozygosity there is a lack of uniformity in the crop, since seeds are produced by plants according to the genetic variances governed by Mendelian rules. Basically, each seed is genetically different and each will grow with its own specific traits. Therefore, it is preferred that the transformed plant be produced such that the regenerated plant has the identical traits and characteristics of the parent transgenic plant. Therefore, it is preferred that the transformed plant be regenerated by micropropagation, which provides a rapid, consistent reproduction of the transformed plants.

[0099] Micropropagation is a process of growing new generation plants from a single piece of tissue that has been excised from a selected parent plant or cultivar. This process permits the mass reproduction of plants having the preferred tissue expressing the protein. The new generation plants, which are produced, are genetically identical to, and have all of the characteristics of, the original plant. Micropropagation allows mass production of quality plant material in a short period of time and offers a rapid multiplication of selected cultivars in the preservation of the characteristics of the original transgenic or transformed plant. The advantages of cloning plants are the speed of plant multiplication and the quality and uniformity of plants produced.

[0100] Micropropagation is a multi-stage procedure that requires alteration of culture medium or growth conditions between stages. Thus, the micropropagation process involves four basic stages: Stage one, initial tissue culturing; stage two, tissue culture multiplication; stage three, differentiation and plant formation; and stage four, greenhouse culturing and hardening. During stage one, initial tissue culturing, the tissue culture is established and certified contaminant-free. During stage two, the initial tissue culture

is multiplied until a sufficient number of tissue samples are produced to meet production goals. During stage three, the tissue samples grown in stage two are divided and grown into individual plantlets. At stage four, the transformed plantlets are transferred to a greenhouse for hardening where the plants' tolerance to light is gradually increased so that it can be grown in the natural environment.

[0101] Sequences suitable for permitting integration of the heterologous sequence into the plant genome are recommended. These might include transposon sequences and the like for homologous recombination as well as Ti sequences which permit random insertion of a heterologous expression cassette into a plant genome.

[0102] Suitable prokaryote selectable markers include resistance toward antibiotics such as ampicillin or tetracycline. Other DNA sequences encoding additional functions may also be present in the vector, as is known in the art.

[0103] The constructs of the subject invention will include an expression cassette for expression of the protein of interest. Usually, there will be only one expression cassette, although two or more are feasible. The recombinant expression cassette will contain in addition to the heterologous sequence one or more of the following sequence elements, a promoter region, plant 5' untranslated sequences which can include regulatory elements, initiation codon depending upon whether or not the structural gene comes equipped with one, and a transcription and translation termination sequence. Unique restriction enzyme sites at the 5' and 3' ends of the cassette allow for easy insertion into a pre-existing vector.

[0104] As used herein, the phrase "regulatory element" refers to a nucleotide sequence which are typically included within an expression cassette and function in regulating (i.e., enhancing or depressing) the expression of a coding sequence therefrom. This regulation can be effected either at the transcription or the translation stages. Examples of regulatory elements include, but are not limited to, enhancers, suppressors and transcription terminators.

[0105] As used herein the term "promoter" refers to a nucleotide sequence, which can direct gene expression in cells. Such a promoter can be derived from a plant, a plant virus, or from any other living organism including bacteria and animals.

[0106] A plant promoter can be a constitutive promoter, such as, but not limited to, CaMV35S and CaMV19S promoters, FMV34S promoter, sugarcane bacilliform badnavirus promoter, CsVMV promoter, *Arabidopsis* ACT2/ACT8 actin promoter, *Arabidopsis* ubiquitin UBQ1 promoter, barley leaf thionin BTH6 promoter, and rice actin promoter.

[0107] The promoter can alternatively be a tissue specific promoter. Examples of plant tissue specific promoters include, without being limited to, bean phaseolin storage protein promoter, DLEC promoter, PHS β promoter, zein stprotein promoter, conglutin gamma promoter from soybean, AT2S1 gene promoter, ACT11 actin promoter from *Arabidopsis*, napA promoter from *Brassica napus*, potato patatin gene promoter and the Tob promoter.

[0108] The promoter may also be a promoter which is active in a specific developmental stage of a plant's life

cycle, for example, a promoter active in late embryogenesis, such as: the LEA promoter; Endosperm-specific expression promoter (the seed storage prolamin from rice is expressed in tobacco seed at the developmental stage about 20 days after flowering) or the promoter controlling the FbL2A gene during fiber wall synthesis stages.

[0109] In case of a tissue-specific promoter, it ensures that the heterologous protein is expressed only in the desired tissue, for example, only in the flower, the fruit, the root, the seed, etc.

[0110] Both the tissue-specific and the non-specific promoters may be constitutive, i.e., may cause continuous expression of the heterologous protein.

[0111] The promoter may also be an inducible promoter, i.e., a promoter which is activated by the presence of an inducing agent, and only upon said activation, causes expression of the heterologous protein. An inducing agent can be for example, light, chemicals, drought, high salinity, osmotic shock, oxidant conditions or in case of pathogenicity and include, without being limited to, the light-inducible promoter derived from the pea *rbcS* gene, the promoter from the alfalfa *rbcS* gene, the promoters DRE, MYC and MYB active in drought; the promoters INT, INPS, prxEa, Ha hsp17.7G4 and RD21 active in high salinity and osmotic stress, the promoters hsr303J and str246C active in pathogenic stress, the copper-controllable gene expression system and the steroid-inducible gene system

[0112] Alternatively, an inducing agent may be an endogenous agent which is normally present in only certain tissues of the plant, or is produced only at certain time periods of the plant's life cycle, such as ethylene or steroids. By using such an endogenous tissue-specific inducing agent, it is possible to control the expression from such inducible promoters only in those specific tissues. By using an inducing agent produced only during a specific period of the life cycle, it is possible to control the expression from an inducible promoter to the specific phase in the life-cycle in which the inducing agent is produced.

[0113] Bacterial and yeast derived promoters are well known in the art.

[0114] Viruses are a unique class of infectious agents whose distinctive features are their simple organization and their mechanism of replication. In fact, a complete viral particle, or virion, may be regarded mainly as a block of genetic material (either DNA or RNA) capable of autonomous replication, surrounded by a protein coat and sometimes by an additional membranous envelope such as in the case of alpha viruses. The coat protects the virus from the environment and serves as a vehicle for transmission from one host cell to another.

[0115] Viruses that have been shown to be useful for the transformation of plant hosts include CaV, TMV and BV. Transformation of plants using plant viruses is described in U.S. Pat. No. 4,855,237 (BGV), EP-A 67,553 (TMV), Japanese Published Application No. 63-14693 (TMV), EPA 194,809 (BV), EPA 278,667 (BV); and Gluzman, Y. et al., Communications in Molecular Biology: Viral Vectors, Cold Spring Harbor Laboratory, New York, pp. 172-189 (1988). Pseudovirus particles for use in expressing foreign DNA in many hosts, including plants, is described in WO 87/06261.

[0116] Construction of plant RNA viruses for the introduction and expression of non-viral foreign genes in plants is demonstrated by the above references as well as by Dawson, W. O. et al., *Virology* (1989) 172:285-292; Takamatsu et al. *EMBO J.* (1987) 6:307-311; French et al. *Science* (1986) 231:1294-1297; and Takamatsu et al. *FEBS Letters* (1990) 269:73-76.

[0117] When the virus is a DNA virus, the constructions can be made to the virus itself. Alternatively, the virus can first be cloned into a bacterial plasmid for ease of constructing the desired viral vector with the foreign DNA. The virus is then excised from the plasmid. If the virus is a DNA virus, a bacterial origin of replication can be attached to the viral DNA, which is then replicated by the bacteria. Transcription and translation of this DNA will produce the coat protein which will encapsidate the viral DNA. If the virus is an RNA virus, the virus is generally cloned as a cDNA and inserted into a plasmid. The plasmid is then used to make all of the constructions. The RNA virus is then produced by transcribing the viral sequence of the plasmid and translation of the viral genes to produce the coat protein(s) which encapsidate the viral RNA.

[0118] Construction of plant RNA viruses for the introduction and expression of non-viral foreign genes in plants is demonstrated by the above references as well as in U.S. Pat. No. 5,316,931.

[0119] In one embodiment, a plant viral nucleic acid is provided in which the native coat protein coding sequence has been deleted from a viral nucleic acid, a non-native plant viral coat protein coding sequence and a non promoter, preferably the subgenomic promoter of the non-native coat protein coding sequence, capable of expression in the plant host, packaging of the recombinant plant viral nucleic acid, and ensuring a systemic infection of the host by the recombinant plant viral nucleic acid, has been inserted. Alternatively, the coat protein gene may be inactivated by insertion of the non-native nucleic acid sequence within it, such that a protein is produced. The recombinant plant viral nucleic acid may contain one or more additional non-native subgenomic promoters. Each non-native subgenomic promoter is capable of transcribing or expressing adjacent genes or nucleic acid sequences in the plant host and incapable of recombination with each other and with native subgenomic promoters. Non-native (foreign) nucleic acid sequences may be inserted adjacent the native plant viral subgenomic promoter or the native and a non-native plant viral subgenomic promoters if more than one nucleic acid sequence is included. The non-native nucleic acid sequences are transcribed or expressed in the host plant under control of the subgenomic promoter to produce the desired products.

[0120] In a second embodiment, a recombinant plant viral nucleic acid is provided as in the first embodiment except that the native coat protein coding sequence is placed adjacent one of the non-native coat protein subgenomic promoters instead of a non-native coat protein coding sequence.

[0121] In a third embodiment, a recombinant plant viral nucleic acid is provided in which the native coat protein gene is adjacent its subgenomic promoter and one or more non-native subgenomic promoters have been inserted into the viral nucleic acid. The inserted non-native subgenomic promoters are capable of transcribing or expressing adjacent

genes in a plant host and are incapable of recombination with each other and with native subgenomic promoters. Non-native nucleic acid sequences may be inserted adjacent the non-native subgenomic plant viral promoters such that said sequences are transcribed or expressed in the host plant under control of the subgenomic promoters to produce the desired product.

[0122] In a fourth embodiment, a recombinant plant viral nucleic acid is provided as in the third embodiment except that the native coat protein coding sequence is replaced by a non-native coat protein coding sequence.

[0123] The viral vectors are encapsidated by the coat proteins encoded by the recombinant plant viral nucleic acid to produce a recombinant plant virus. The recombinant plant viral nucleic acid or recombinant plant virus is used to infect appropriate host plants. The recombinant plant viral nucleic acid is capable of replication in the host, systemic spread in the host, and transcription or expression of foreign gene(s) in the host to produce the desired protein.

[0124] In many instances it is desired to target the expression of a recombinant protein. Such targeting can be into a cellular organelle or outside of the cell. This can be effected, as is well known in the art, by appropriate signal peptides, which are fused to the polypeptide to be targeted, typically at the N terminus.

[0125] Thus, as used herein and in the claims section which follows, the phrase "signal peptide" refers to a stretch of amino acids which is effective in targeting a protein expressed in a cell into a target location. Different signal peptides, which are known in the art, are effective in secreting a protein from bacteria, yeast, plant and animal cells.

[0126] It should be noted in this respect that signal peptides serve the function of translocation of produced protein across the endoplasmic reticulum membrane. Similarly, transmembrane segments halt translocation and provide anchoring of the protein to the plasma membrane, see, Johnson et al. *The Plant Cell* (1990) 2:525-532; Sauer et al. *EMBO J.* (1990) 9:3045-3050; Mueckler et al. *Science* (1985) 229:941-945. Mitochondrial, nuclear, chloroplast, or vacuolar signals target expressed protein correctly into the corresponding organelle through the secretory pathway, see, Von Heijne, *Eur. J. Biochem.* (1983) 133:17-21; Yon Heijne, *J. Mol. Biol.* (1986) 189:239-242; Iturriaga et al. *The Plant Cell* (1989) 1:381-390; McKnight et al., *Nucl. Acid Res.* (1990) 18:4939-4943; Matsuoka and Nakamura, *Proc. Natl. Acad. Sci. USA* (1991) 88:834-838. A recent book by Cunningham and Porter (*Recombinant proteins from plants*, Eds. C. Cunningham and A. J. R. Porter, 1998 Humana Press Totowa, N.J.) describe methods for the production of recombinant proteins in plants and methods for targeting the proteins to different compartments in the plant cell. In particular, two chapters therein (14 and 15) describe different methods to introduce targeting sequences that results in accumulation of recombinant proteins in compartments such as ER, vacuole, plastid, nucleus and cytoplasm. The book by Cunningham and Porter is incorporated herein by reference. Presently, the preferred site of accumulation of the fusion protein according to the present invention is the ER using signal peptide such as Cel 1 or the rice amylase signal peptide at the N-terminus and an ER retaining peptide (HDEL, SEQ ID NO:17; or KDEL, SEQ ID NO:24) at the C-terminus.

[0127] According to an additional aspect of the present invention there is provided a method of producing recombinant β -glucosidase. The method according to this aspect of the present invention is effected by introducing, in an expressible or overexpressible form, a nucleic acid construct into a host cell. The nucleic acid construct includes a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger* and encoding a polypeptide having a β -glucosidase catalytic activity. The polynucleotide preferably further encodes a signal peptide in frame with the polypeptide. Still preferably, the polynucleotide further encodes an endoplasmic reticulum retaining peptide in frame with the polypeptide.

[0128] As used herein the term "introducing" refers both to transforming and to virally infecting, as these terms are further defined hereinabove. As used herein the terms "expressible form" and "overexpressible form" refers to a recombinant form which includes the required regulatory elements to effect expression or over expression of a coding region, all as is further detailed hereinabove.

[0129] According to a preferred embodiment of this aspect of the present invention, after sufficient expression has been detected, the polypeptide having the β -glucosidase catalytic activity is extracted from the expressing host cell.

[0130] Thus host cells, expressing the polypeptide according to the present invention, provide an immediate, easy and indefinite source of the polypeptide.

[0131] Any number of well-known liquid or solid culture media may be used for appropriately culturing host cells of the present invention, although growth on liquid media is preferred as the secretion of the polypeptide into the media results in simplification of polypeptide recovery. As is further detailed hereinabove, such secretion can be effected by the incorporation of a suitable signal peptide. The β -glucosidase may be isolated or separated or purified from host cell preparations using techniques well known in the art, such as, but not limited to, centrifugation filtration, chromatography, electrophoresis and dialysis. Further concentration and/or purification of the β -glucosidase may be effected by use of conventional techniques, including, but not limited to, ultrafiltration, further dialysis, ion-exchange chromatography, HPLC, size-exclusion chromatography, cellobiose-sepharose affinity chromatography, and electrophoresis, such as polyacrylamide-gel-electrophoresis (PAGE). Using these techniques, β -glucosidase may be recovered in pure or substantially pure form.

[0132] According to an additional aspect of the present invention there is provided a method of increasing a level of at least one fermentation substance in a fermentation product. The method according to this aspect of the present invention is effected by fermenting a glucose containing fermentation starting material by a yeast cell overexpressing a nucleic acid construct which includes a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger* and which encodes a polypeptide having a β -glucosidase catalytic activity, thereby increasing the level of the at least one fermentation substance in the fermentation product. The polynucleotide preferably further encodes a signal peptide in frame with the polypeptide. Still preferably, the polynucleotide further encodes an endoplasmic reticulum retaining peptide in frame with the polypeptide.

[0133] According an alternative aspect of the present invention there is provided a method of increasing a level of at least one fermentation substance in a fermentation product. The method according to this aspect of the present invention is effected by fermenting a plant derived glucose containing fermentation starting material by a yeast cell, the plant overexpressing a nucleic acid construct which includes a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger* and which encodes a polypeptide having a β -glucosidase catalytic activity, thereby increasing the level of the at least one fermentation substance in the fermentation product. The polynucleotide preferably further encodes a signal peptide in frame with the polypeptide. Still preferably, the polynucleotide further encodes an endoplasmic reticulum retaining peptide in frame with the polypeptide.

[0134] As used herein in the specification and in the claims section that follows, the term "fermentation" refers to a chemical change induced in a complex organic compound by the action of an enzyme, whereby the substance is split into simpler compounds. Specifically, the term "fermentation" includes the anaerobic dissimilation of substrates with the production of energy and reduced compounds, the final products thereof are organic acids, alcohols, such as ethanol, isopropanol, butanol, etc., and CO₂. Such products, are typically secreted and each of which is referred to herein as a "fermentation substance", i.e., any known fermentation resultant of either microbial or yeast fermentation.

[0135] As used herein in the specification and in the claims section that follows, the phrase "fermentation product" refers to the resultant material of a fermentation process. Examples include, but are not limited to, alcohol containing fermentation medium and alcoholic beverages, such a, but not limited to, fruit-based alcohol-containing beverages, wines and beers.

[0136] When used in conjunction with, for example, a β -glucanase, the β -glucosidase is effective for hydrolyzing a variety of cellulose containing materials to glucose. The glucose produced by enzymatic hydrolysis of the cellulose and other glucose containing saccharides, may be recovered and stored, or it may be subsequently fermented to ethanol using conventional techniques. Many processes for the fermentation of glucose generated from cellulose are well known, and are suitable for use herein. Briefly, the hydrolyzate containing the glucose from the enzymatic reaction is contacted with an appropriate microorganism under conditions effective for the fermentation of the glucose to ethanol. This fermentation may be separate from and follow the enzymatic hydrolysis of the cellulose (sequentially processed), or the hydrolysis and fermentation may be concurrent and conducted in the same vessel (simultaneously processed). Details of the various fermentation techniques, conditions, and suitable microorganisms have been described, for example, by Wyman (1994, Bioresource Technol., 50:3-16) or Olsson and Hahn-Hagerdal (1996, Enzyme Microbial Technol., 18:312-331), the content of each of which is incorporated herein by reference. Following the completion of a fermentation, the alcohol may be recovered by extraction, and optionally purified e.g., by distillation.

[0137] Thus, according to still another aspect of the present invention there is provided a method of producing an

alcohol. The method according to this aspect of the present invention is effected by fermenting a glucose containing fermentation starting material by a cell overexpressing a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, encoding a polypeptide having a β -glucosidase catalytic activity, and extracting the alcohol therefrom. The polynucleotide preferably further encodes a signal peptide in frame with the polypeptide. Still preferably, the polynucleotide further encodes an endoplasmic reticulum retaining peptide in frame with the polypeptide.

[0138] According to an additional aspect of the present invention there is provided a method of producing an alcohol. The method according to this aspect of the present invention is effected by fermenting a plant derived glucose containing fermentation starting material by a cell, the plant overexpressing a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, encoding a polypeptide having a β -glucosidase catalytic activity, and extracting the alcohol therefrom. The polynucleotide preferably further encodes a signal peptide in frame with the polypeptide. Still preferably, the polynucleotide further encodes an endoplasmic reticulum retaining peptide in frame with the polypeptide.

[0139] Plants contain aroma and flavor compounds of glycosidic nature, their inherent aroma property can be released by degrading enzymes, turning a non-volatile aroma compound into its volatile form. Thus, for example, α -L-arabinofuranosidases, assist in the liberation of aroma compounds from substrates such as juices or wines, as described by Gunata et al. (European Patent Application No. 332.281, 1989; and "purification and some properties of an alpha-L-arabinofuranosidase from *A. niger* action on grape monoterpenyl arabinofuranosylglucosides. J. Agric. Food Chem. 38: 772-776, 1990). This outcome is achieved, for example, in a two step process wherein the first step comprises the use of an α -L-arabinofuranosidase, to catalyze the release of arabinose residues from monoterpenyl α -L-arabinofuranosyl glucosides contained in, for example, the fruit or vegetable juice via the cleavage of the (1 \rightarrow 6) linkage between a terminal arabinofuranosyl unit and the intermediate glucose of a monoterpenyl α -L-arabinofuranosylglucoside. The α -L-arabinofuranosidase is preferably in a purified form so as to avoid the undesirable degradation of other components of the juice which may be detrimental to its ultimate quality. In the second step, β -glucosidase is required to yield the free terpenol from the resulting des-arabinosylated monoterpenyl glucoside. If desired, both reaction steps may be performed in the same reaction vessel without the need to isolate the intermediate product (Gunata et al. (1989), supra). Thus, β -glucosidase is an essential contributor when the liberation of these aroma compounds for improving the flavor of the juice or wine is desired. Moreover, in the case of wine, the control of the liberation of aroma compounds provides wines with a more consistent flavor, thus reducing or eliminating the undesirable effect of "poor vintage years". Additional information is contained in: "Cloning and expression of DNA molecules encoding arabinan degrading enzyme of fungal origin", U.S. Pat. No. 5,863,783; Y. Gueguen, et al. "A Very Efficient β -Glucosidase Catalyst for the Hydrolysis of Flavor Precursors of

Wines and Fruit Juices", J. Agric. Food Chem. 44:2336-2340, 1996, each of which is incorporated herein by reference.

[0140] Thus, according to a further aspect of the present invention there is provided a method of increasing a level of at least one aroma substance in a plant derived product, such as, but not limited to, an alcoholic beverage. The method according to this aspect of the present invention is effected by incubating a glucose containing plant starting material with a yeast cell overexpressing a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger* which encodes a polypeptide having a β -glucosidase catalytic activity, thereby increasing the level of the at least one aroma substance in the plant derived product. The polynucleotide preferably further encodes a signal peptide in frame with the polypeptide. Still preferably, the polynucleotide further encodes an endoplasmic reticulum retaining peptide in frame with the polypeptide.

[0141] While reducing the present invention to practice it was discovered that in order to obtain activity of a β -glucosidase in a transgenic plant, the expression construct should include a signal peptide. In addition, it was found that retaining the enzyme in the endoplasmic reticulum results in higher release of aroma compounds following homogenization and incubation. It is assumed that compartmentalization of the enzyme in for example the ER prevents it from interacting with its substrates which are mainly outside the cells, limiting such interaction following homogenization. Indeed, directing the enzyme to the apoplast resulted in increased release of aroma in vivo. Thus, depending on the specific application, one can choose whether to include in the construct an endoplasmic reticulum retaining peptide or not.

[0142] According to yet a further aspect of the present invention there is provided a method of increasing a level of at least one aroma substance in a plant derived product, such as, but not limited to, an alcoholic beverage. The method according to this aspect of the present invention is effected by incubating a glucose containing plant starting material with a yeast cell, said plant overexpressing a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger* which encodes a polypeptide having a β -glucosidase catalytic activity, thereby increasing the level of the at least one aroma substance in the plant derived product. The polynucleotide preferably further encodes a signal peptide in frame with the polypeptide. Still preferably, the polynucleotide further encodes an endoplasmic reticulum retaining peptide in frame with the polypeptide.

[0143] As used herein in the specification and in the claims section that follows, the phrase "glucose containing starting material" refers to any source of energy, in the form of glucose containing compounds, other than free glucose, including, but not limited to, crushed, minced, diced or extracted plant material, plant, or portions thereof, such as fruits, examples thereof are tropical fruits and grapes.

[0144] According to an additional aspect of the present invention there is provided a method of producing an aroma spreading plant. As used herein in the specification and in the claims section that follows, the phrase "aroma spreading plant" refers to substantially any part of a plant, in which volatile compounds are generated by the catalytic activity of

the β -glucosidase polypeptide of the present invention, release of volatile compounds therefrom is perceived by the olfactory system of an organism, such as a human.

[0145] The method according to this aspect of the present invention is effected by overexpressing in the plant a nucleic acid construct including a genomic, complementary or composite polynucleotide derived from *Aspergillus niger*, which encodes a polypeptide having a β -glucosidase catalytic activity, thereby increasing aroma spread from the plant. Such overexpression is preferably performed in a tissue specific manner by, for example, employing a tissue specific promoter, as hereinabove described, to thereby overexpress a heterologous protein in a selected portion of the plant. The tissue in which such overexpression is effected is selected according to the availability of glucose containing non-volatile aroma substrates therein. Thus, such an overexpression will cause the release of a volatile and aroma constituent of the substrate. Thus, according to preferred embodiments overexpressing the nucleic acid construct is limited to at least one tissue, such as a flower, a fruit, a seed, a root, a stem, pollen and leaves.

[0146] According to still a further aspect of the present invention there is provided a method of increasing a level of free glucose in a glucose containing fermentation starting material. The method according to this aspect of the present invention is effected by fermenting the glucose containing fermentation starting material by a cell overexpressing a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, which encodes a polypeptide having a glucosidase catalytic activity, thereby increasing the level of the free glucose in the glucose containing fermentation starting material. The polynucleotide preferably further encodes a signal peptide in frame with the polypeptide. Still preferably, the polynucleotide further encodes an endoplasmic reticulum retaining peptide in frame with the polypeptide.

[0147] According to another aspect of the present invention there is provided a method of increasing a level of free glucose in a plant derived glucose containing fermentation starting material. The method according to this aspect of the present invention is effected by fermenting the plant derived glucose containing fermentation starting material by a cell, the plant overexpressing a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, which encodes a polypeptide having a β -glucosidase catalytic activity, thereby increasing the level of the free glucose in the plant. The polynucleotide preferably further encodes a signal peptide in frame with the polypeptide. Still preferably, the polynucleotide further encodes an endoplasmic reticulum retaining peptide in frame with the polypeptide.

[0148] As used herein in the specification and in the claims section that follows, the term "free glucose" refers to glucose residues in the form of a monosaccharide, the levels of which are increased by the catalytic activity of β -glucosidase.

[0149] As used herein in the specification and in the claims section that follows, the phrase "glucose containing fermentation starting material" refers to any source of energy, in the form of glucose containing compounds, other than free glucose, including, but not limited to, crushed,

minced, diced or extracted plant material, plant, or portions thereof, used in industrial fermentation processes.

[0150] According to yet another aspect of the present invention there is provided a method of increasing a level of extra- or intracellular free glucose in a plant. The method according to this aspect of the present invention is effected by overexpressing in the plant a nucleic acid construct including a genomic, complementary or composite polynucleotide preferably derived from *Aspergillus niger*, which encodes a polypeptide having a β -glucosidase catalytic activity, thereby increasing the level of the free glucose in the plant. Thus, sweeter fruits can be produced. The polynucleotide preferably further encodes a signal peptide in frame with the polypeptide. Still preferably, the polynucleotide further encodes an endoplasmic reticulum retaining peptide in frame with the polypeptide.

[0151] Glycosidases, including β -glucosidase, catalyze reactions involving the hydrolysis of O-glycosidic bond of glycosides, and synthesize oligosaccharides when the reaction is run in reverse from the normal direction, a result achieved by, for example, site directed mutagenesis, and Km reversal. As described in the Background section hereinabove, the hydrolysis reaction mechanism of glycosidases involves two catalytic steps, the second of which involves a base catalyzed H_2O attack, resulting in the regeneration of the enzyme, and the release of the saccharide residue. Thus, in addition, oligosaccharide synthesis can be achieved by adding a second saccharide to the reaction mixture, which competes with the H_2O molecule, and reacts in its place with the first saccharide in, what is known as, a transglycosylation reaction. Hence, as glycosidases are generally available and easy to handle, these enzymes have the potential to catalyze the production of many different products using inexpensive substrates. For further detail see U.S. Pat. No. 5,716,812, which is incorporated herein by reference.

[0152] Thus, according to yet an additional aspect of the present invention there is provided a method of synthesizing oligosaccharides. The method according to this aspect of the present invention is effected by mixing a polypeptide having a β -glucosidase catalytic activity with first and second saccharide molecules to thereby join the first and second saccharide molecules into an oligosaccharide.

[0153] Additional objects, advantages, and novel features of the present invention will become apparent to one ordinarily skilled in the art upon examination of the following examples, which are not intended to be limiting. Additionally, each of the various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the following examples.

EXAMPLES

[0154] Reference is now made to the following examples, which together with the above descriptions, illustrate the invention in a non limiting fashion.

[0155] Generally, the nomenclature used herein and the laboratory procedures utilized in the present invention include molecular, biochemical, microbiological and recombinant DNA techniques. Such techniques are thoroughly explained in the literature. See, for example, "Molecular Cloning: A laboratory Manual" Sambrook et al., (1989);

“Current Protocols in Molecular Biology” Volumes I-III Ausubel, R. M., ed. (1994); Ausubel et al., “Current Protocols in Molecular Biology”, John Wiley and Sons, Baltimore, Md. (1989); Perbal, “A Practical Guide to Molecular Cloning”, John Wiley & Sons, New York (1988); Watson et al., “Recombinant DNA”, Scientific American Books, New York; Birren et al. (eds) “Genome Analysis: A Laboratory Manual Series”, Vols. 1-4, Cold Spring Harbor Laboratory Press, New York (1998); methodologies as set forth in U.S. Pat. Nos. 4,666,828; 4,683,202; 4,801,531; 5,192,659 and 5,272,057; “Cell Biology: A Laboratory Handbook”, Volumes I-III Cellis, J. E., ed. (1994); “Current Protocols in” Volumes I-III Coligan J. E., ed. (1994); Stites et al. (eds), “Basic and Clinical Immunology” (8th Edition), Appleton & Lange, Norwalk, Conn. (1994); Mishell and Shiigi (eds), “Selected Methods in Cellular Immunology”, W. H. Freeman and Co., New York (1980); available immunoassays are extensively described in the patent and scientific liter, see, for example, U.S. Pat. Nos. 3,791,932; 3,839,153; 3,850,752; 3,850,578; 3,853,987; 3,867,517; 3,879,262; 3,901,654; 3,935,074; 3,984,533; 3,996,345; 4,034,074; 4,098,876; 4,879,219; 5,011,771 and 5,281,521; “Oligonucleotide Synthesis” Gait, M. J., ed. (1984); “Nucleic Acid Hybridization” Hames, B. D., and Higgins S. J., eds. (1985); “Transcription and Translation” Hames, B. D., and Higgins S. J., eds. (1984); “Animal Cell Culture” Freshney, R. I., ed. (1986); “Immobilized Cells and Enzymes” IRL Press, (1986); “A Practical Guide to Molecular Cloning” Perbal, B., (1984) and “Methods in Enzymology” Vol. 1-317, Academic Press; “PCR Protocols: A Guide To Methods And Application”, Academic Press, San Diego, Calif. (1990); Marshak et al., “Strategies for Protein Purification and Characterization—A Laboratory Course Manual” CSHL Press (1996); all of which are incorporated by reference as if fully set forth herein. Other general references are provided throughout this document. The procedures therein are believed to be well known in the art and are provided for the convenience of the reader. All the information contained therein is incorporated herein by reference.

MATERIALS AND EXPERIMENTAL METHODS

[0156] Purification of *A. niger* β -glucosidase:

[0157] A crude preparation of *A. niger* B1 (CMI CC 324626) β -glucosidase was obtained from Shaligal Ltd. (Tel-Aviv, Israel). A sample (10 ml) of the crude enzyme (140 Units/ml) was first diafiltered through a 50 kDa cut-off AMICON™ size filtration membrane (Amicon Corp., Danvers, Mass.), with 20 mM citrate buffer pH=5. The proteins were then separated on an FPLC equipped with a MONO-Q™ anion exchange RH 5/5 column (Amersham Pharmacia Biotech AB, Uppsala, Sweden), equilibrated with the same buffer. The enzyme was eluted with a linear gradient of 0 to 350 mM NaCl. Active fractions (see below, enzyme assays) were monitored and pooled (between 80-110 mM NaCl). The partially purified enzyme was dialyzed against 20 mM citrate buffer pH=3.5, applied to a RESOURCE-S™ (Amersham Biosciences Inc, Piscataway, N.J.) cation exchange column equilibrated with the same buffer, and eluted with a gradient of 0-1 M NaCl. The purified enzyme (eluted at 155 mM NaCl) was concentrated by ultrafiltration (50 kDa cut-off membrane, Amicon).

[0158] Enzyme Assays:

[0159] β -glucosidase enzyme activity was monitored using a plate assay as follows. 4-methylumbelife β -D-

glucopyranoside (MUGlc, Sigma Chemical Inc. St. Louis, Mo.) to a final concentration of 0.5 mM, was dissolved in PC buffer (50 mM phosphate, 12 mM citric acid, pH=3.4) at 45° C. The solution was mixed with 3% agar in water, previously boiled and then cooled to 45° C. The resulting solution (20 ml) was poured into a petri dish and after solidification, 10 μ l enzyme samples were spotted. The plate was incubated at 50° C. for one hour, and then illuminated with long UV. An intense fluorescence was indicative of β -glucosidase activity.

[0160] Detection of β -glucosidase in polyacrylamide gels was carried out by washing the SDS-polyacrylamide gel with 1:1 isopropanol:PC buffer to remove SDS and renature the enzyme. The gel was washed once in PC buffer and incubated in a thin layer of a solution of 0.5 mM MUGlc. After incubation at 50° C. for one hour, the active protein band was visualized by UV light.

[0161] Quantitative assays were performed using pNPGlc as a substrate according to Shoseyov (7).

[0162] Determination of Thermal Stability of *A. niger* β -Glucosidase:

[0163] Recombinant enzyme (40 μ g/ml) was dissolved in 20 mM citrate phosphate buffer, pH=5. Each tested sample (811) was covered by 1511 mineral oil. The activity was determined by the standard pNPGlc assay (7).

[0164] Deglycosylation of *A. niger* β -Glucosidase by N-Glycosidase-F:

[0165] A N-glycosidase-F (Boehringer Mannheim, Mannheim, Germany) reaction mixture, containing 0.125 μ g pure β -glucosidase (previously denatured by boiling for 3 minutes in 1% SDS and 5% β -mercaptoethanol), 0.2 units of the N-glycosidase-F, sodium phosphate buffer (50 mM, pH=7.5), EDTA (25 mM), 1% Triton X-100 and 0.02% sodium azide, in a total volume of 12.5 μ l, was incubated for 4 hours at 37° C. Reaction was stopped by addition of PAGE sample application buffer followed by 3 minutes of boiling.

[0166] Proteolysis and N-Terminal Sequences of *A. niger* B1 β -Glucosidase:

[0167] Partial enzymatic proteolysis with *Staphylococcus aureus* V8 protease was carried out as described by Cleveland (28). Briefly, FPLC-purified β -glucosidase (5 μ g), was concentrated by acetone precipitation. The protein was separated on a preparative 10% SDS-PAGE. The gel was stained with coomassie blue, destained and rinsed with cold water, and the β -glucosidase protein band was excised. The resulting gel slice was applied to a second SDS-PAGE gel (15% acrylamide) and overlaid with *Staphylococcus aureus* V8 protease. Digestion was carried out within the stacking gel by turning off the current for 30 min. As the bromophenol blue dye neared the bottom of the stacking gel, the current was restored. The electrophoresed cleavage products were electroblotted to PVDF membranes. The native protein was transferred to PVDF in parallel. The N-terminal sequence of the native protein and two of the numerous cleavage products were analyzed by Edman degradation using a gas-phase protein sequencer (Applied Biosystems model 475A microsequencer).

[0168] Cloning of bg11 cDNA and Genomic Gene:

[0169] Total RNA isolation: Total RNA was isolated from *Aspergillus niger* B1 as follows: *A. niger* B1 was grown in

liquid culture consisting of mineral media (NH₄)₂SO₄·3H₂O (0.5 g/l), KH₂PO₄ (0.2 g/l), MgSO₄ (0.2 g/l), CaCl₂·H₂O (0.1 g/l), FeSO₄·6H₂O (0.001 g/l), ZnSO₄·7H₂O (0.001 g/l), and 2 mM citric acid, at pH=3.5 with 1% w/v bran as a carbon source. The medium was autoclaved, cooled and inoculated with *A. niger* B1 (10⁶ spores/ml). Baffled flasks were used with constant shaking (200 RPM) at 37° C. The appearance of β-glucosidase activity was monitored by placing 5 μl of growth medium on 1% agar plates containing 0.5 mM MUGlc, as described above. Activity was detected following 15 hours incubation. The mycelium was harvested following 24 hours growth period, and the medium removed by filtering through GFA™ glass microfibre (Whatman Inter. Ltd., Maidstone, England). The mycelium was then frozen with liquid nitrogen and ground to fine powder with mortar and pestle. Total RNA was produced from this powder by the Guanidine thiocyanate (TRIAGENT™) method (Molecular Research Center, Inc.).

[0170] RNA reverse-transcription reaction: cDNA was obtained by reverse transcribing total RNA (10 μg) using Stratagene RT-PCR kit (Stratagene, La Jolla, Calif.). The reaction mixture (50 μl) additionally consisted of: Oligo dT18 (1 μg), RNase Block Ribonuclease Inhibitor (20 units), 1× buffer (50 mM Tris-HCl, pH=8.3, 75 mM KCl, 10 mM DTT, 3 mM MgCl₂), dNTPs (500 μM each) and reverse transcriptase (300 units). Total RNA was initially denatured at 70° C., allowed to cool to room temperature (for primers annealing), and added to the reaction mixture. The reaction mixture was incubated for 1 hour at 37° C., followed by heating (95° C., 5 minutes) and stored at -70° C. until further use.

[0171] DNA amplification: Degenerate primers for DNA amplification reaction by PCR methods were synthesized, based on part of the amino acid N-terminal sequence and an internal sequence, as determined by the Edman degradation, following V8 proteolysis (hereinbelow, experimental results). The partial sequence from β-glucosidase N-terminal derived amino acid sequence was Ser-Pro-Pro-Tyr-Tyr-Pro (SEQ ID NO:4), yielding the following primer: 5'-(C/G)(A/C/G/T)CC(A/C/G/T) CC(A/C/G/T)TA(C/T)TA(C/T)CC-3' (SEQ ID NO:5). The partial sequence from E2 internal cleavage product amino acid sequence was Gln-Pro-Ile-Leu-Pro-Ala-Gly-Gly (SEQ ID NO:6), yielding the following primer: 5'-TCCIGC(T/G/C/A)GG(TG/C/A)A(G/A) (T/G/A)AT(T/G/C/A)GG(T/C)TG-3' (SEQ ID NO: 7).

[0172] DNA amplification reaction mixture (2511) contained: reverse transcriptase reaction product (1 μl), 10×PCR buffer (2.5 μl, Promega Corp., Madison, Wis.), dNTPs (250 μM each), MgCl₂ (2.0 mM), degenerate primers (250 pmol each), DNA polymerase (3 units, Stratagene, La Jolla, Calif.) and overlaid with mineral oil (25 μl). The reaction was performed in an automated heating block (Programmable thermal controller—MJ Research, Inc.). PCR cycling conditions were 30 seconds denaturing at 94° C., 60 seconds annealing at 50° C., and 150 seconds elongation at 72° C., repeated 36 times. The resulting amplified product was electrophoresed on a 1.2% (w/v) agarose/TBE gel, resulting in a 2.2 kb cDNA gene fragment, which was further isolated using Gel Extraction Kit (QIAGEN, Hilden, Germany) and cloned directly into the single 3'-T PCR insertion site of pGEM-T cloning vector (Promega Corp., Madison, Wis.).

[0173] Probe preparation: The 2.2 kb partial cDNA was digested with PstI to produce a 1.2 kb fragment DNA probe.

A sample (25 ng) of the fragment was labeled with [³²P] dCTP, using the random sequence nononucleotide REDIPRIME™ DNA labeling system (Amersham Pharmacia Biotech AB, Buckinghamshire, England).

[0174] Preparation of genomic DNA plasmid library: An *A. niger* B1 genomic library was constructed in the pYEAUra3 yeast/*E. coli* shuttle vector (Clontech Lab. Inc. Palo Alto, Calif.). *A. niger* B1 was grown in liquid culture as described above, the mycelium harvested following 48 hours of growth, frozen in liquid nitrogen and grounded. The mycelium ground was used to produce genomic DNA by the CTAB method of Murray and Thompson (29). The library was constructed from partially digested Sau3A genomic DNA, cloned into the BamHI site of the pYEUra3 yeast shuttle vector (Clontech Lab. Inc. Palo Alto, Calif.). pYEAUra3 yeast/*E. coli* shuttle vector was digested with BamHI and dephosphorylated with CIP to prevent self ligation. The partially digested genomic DNA was cloned into the shuttle vector with T4 ligase and used to transform TOP10 *E. coli* electro-competent cells, which were then plated on LB-agar containing ampicillin (50 μg/ml). A total of 4×10⁸ colonies were grown on LB-agar plates, blotted to HYBOND-N™ membranes (Amersham Pharmacia Biotech AB, Buckinghamshire, England) and screened using the above described 1.2 kb probe. Positive clones were sub-cloned in pUC18 and sequenced (Biological Services, The Weizmann Institute of Science, Rehovot, Israel).

[0175] Expression of bgl1 cDNA in *E. coli*:

[0176] Two specific primers were designed according to the 5' and the 3' sequences, corresponding to the N-terminal and C-terminal region of the mature protein: sense primer: 5'-' (SEQ ID NO:8). Antisense primer: 5'-AAAGGATCCT-TAGTGAACAGTAGGCAGAGACGC-3' (SEQ ID NO:9). The isolated cDNA was digested with NcoI and BamHI and cloned into a pET3d expression vector (FIG. 1A, Novagen Inc., Madison, Wis.). Positive *E. coli* BL21 (DE3) pLysS colonies, containing the bgl1 cDNA, were confirmed by enzyme restriction and sequence analysis. Recombinant BGL1 was expressed according to the manufacturer's protocol.

[0177] Expression of bgl1 cDNA in *Saccharomyces cerevisiae* and *Pichia pastoris*:

[0178] The pYES2 vector (Invitrogen Inc., San Diego, Calif.) was used to successfully clone the bgl1 cDNA gene into the HindIII/BamHI of pYES2-bgl1 plasmid (FIG. 1b), and transform *Saccharomyces cerevisiae* using the lithium acetate method (30). The BGL1 was expressed by inducing the Gall promoter according to the manufacturer's protocol. *Saccharomyces cerevisiae* strain INVSc2 (MATa, his3-D200, ura3-167) was used as the host. *Pichia pastoris* strain GS115 (his4 mutant) was used as the host for shuttle and expression vector plasmid pHIL-S1 (Invitrogen Inc., San Diego, Calif.). The bgl1 cDNA was cloned into the EcoRI/BamHI sites of pHIL-S1, yielding the pHIL-S1-bgl1 expression and secretion vector (FIG. 1c). Expression in *P. pastoris* was carried out according to the manufacturer's protocol. Screening of β-glucosidase-expressing clones was facilitated by top-agar, containing 50 mg X-Glc, 30 ml methanol and 1% agar per liter. Blue color indicated a colony producing active β-glucosidase.

[0179] Western Blot Analysis:

[0180] Antibodies were produced from rabbit serum 36 days following a second injection of 100 µg purified protein and adjuvant (AniLab Biological Services, Tal-Sachar, Israel). High molecular weight ladder was from Sigma Chemical Inc. St. Louis, Mo. Western blot conditions were as described in reference 36.

[0181] Determination of the Stereochemical Course of Hydrolysis:

[0182] The method was essentially as described by Wong et al. (31). PNPgIc (10 µmols) was dissolved in 0.5 ml of 25 mM acetate buffer pH=3.5 in D₂O in an NMR tube. β-Glucosidase was lyophilized and redissolved in 100 µl D₂O (35 units/ml). The ¹H-NMR spectrum of the substrate was recorded, enzyme added (10 µl), and spectra recorded at specified time intervals on a Bruker AMX400 at 25° C.

[0183] Inactivation and Reactivation Studies:

[0184] Pure *A. niger* β-Glucosidase enzyme (0.47 mg/ml) was incubated in the presence of various concentrations of 2-deoxy-2-fluoro-β-glucosyl fluoride (2FGlcF, 0.5-6 mM) in 30 mM citrate buffer pH=4.8 at 50° C. Residual enzyme activity was determined at different time intervals by addition of an aliquot (10 µl) of the inactivation mixture, to a solution containing citrate buffer (30 mM, pH=4.8), BSA (8 µg) and 2,4-dinitrophenyl β-D-glucopyranoside (DNPgIc, 0.625 mM, 830 µl). Release of DNP was determined spectrophotometrically by measuring the absorbance at 400 nm one minute after the addition of the substrate.

[0185] Reactivation rates were determined as follows: pure *A. niger* β-glucosidase (0.34 mg/ml) was preincubated with 2FGlcF (5 mM) for 15 min, after which the excess of the inactivator was diafiltered by 20-kDa nominal molecular mass cutoff centrifugal concentrators (Sartorius Inc., Goettingen, Germany). Samples of the purified, inactivated enzyme were incubated in the presence linamarin (0-16 mM) in citrate buffer (30 mM, pH=4.8) at 50° C. for 0, 10, 20 and 30 minutes, and the activity of each sample was determined using p-nitrophenyl β-D-glucopyranoside (pNPgIc) as a substrate.

[0186] Expression of bgl1 cDNA in Tobacco Plants:

[0187] Genetic Constructs:

[0188] Bgl1 cDNA was cloned in pETB1 (37), pJD330 and pBINPlus (38) were used as an intermediate and binary vector, respectively. Cel1 signal sequence as well as 35S pLUC fragment were retrieved from pB21, modified pBLUESCRIPT® SK (39). *Nicotiana tabacum* cv. Samson was used as a model plant for gene transformation. Three gene constructs were employed (FIGS. 11a-c): (i) bgl1 without any signal peptide which served for cytoplasmic expression (FIG. 11a, plasmid pJDB1); (ii) bgl1 including a cell signal peptide at the N terminus for secretion into the apoplast (FIG. 11b, plasmid pJDCB1); and (iii) bgl1 including the cell signal peptide and the KDEL (SEQ ID NO:24) ER-retaining peptide at the C-terminus for accumulation in the ER (FIG. 11c, plasmid pJDCB1T).

[0189] To this end, bgl1 cDNA (2.5 kb) was released from pETB1 (37) with NcoI and BamHI and inserted into pJD330 between the 35S promoter Ω fragment and the nos terminator, eliminating the gus gene, resulting in plasmid pJDB1.

Endoplasmic reticulum retaining signal tetrapeptide HDEL (SEQ ID NO:17) was synthesized and fused with bgl1 at the C-terminal in pJDB1 by a fidelity PCR reaction with the following pair of primers: Forward primer (23 mer), starting from nucleotide 1248 of bgl1 cDNA 5'-(1248)-CAGTGAC-CGTGGATGCGACAATG-(1270)-3' (SEQ ID NO:20); Reverse primer (40 mer), starting at nucleotide 2506 of bgl1 cDNA encoding also for the HDEL (SEQ ID NO:17) peptide 5'-(2506)-AGAGACGGATGACAAGTACTACT-TGAAAATTGGGCCCAAAA-3' (SEQ ID NO:21). For pJDCB1T (35S Ω+Cel1+bgl1+HDEL, SEQ ID NO:17), the 35S Ω fragment of pJDB1 was replaced by a 35S Ω+Cel1 fragment digested from pB21 with BamHI and XbaI. For pJDCB1 (35S Ω+Cel1+bgl1), the fragment containing 35S Ω and Cel1 as well as part of bgl1 was cut from pJDCB1T with HindIII and NruI and ligated with the vector of pJDB1 digested with the same pair of restriction enzymes. The nucleotide sequence of all of the genetic constructs was confirmed by DNA sequencing.

[0190] Gene cassettes in the intermediate vectors of pJDB1, pJDCB1 and pJDCB1T were further isolated with HindIII and EcoRI and inserted into multiple cloning sites of the binary vector pBINPlus. Disarmed *Agrobacterium* LB4404 was transformed with pBINPlus containing bgl1 gene cassettes.

[0191] Tobacco Plant Transformation:

[0192] The young leaves of in vitro grown plantlets were excised and cut into 0.5 cm² pieces and then immersed for 5 minutes in an overnight grown culture of *Agrobacterium*. After blotted with sterile Whatman filter paper, the infected leaves were co-cultured for 2 days with *Agrobacterium* on MS medium plus 2.0 mg/L of Zeatin and 0.1 mg/L of IAA as well as 0.35% (w/v) phytigel and then transferred to the same medium but with 300 mg/L kanamycin and 300 mg/L carbenicillin. Regenerates were then transferred to the rooting media, containing only MS salts, vitamins and the same antibiotics. Rooted plants were transferred to greenhouse after PCR screening.

[0193] Screening for Transgenic Plants:

[0194] DNA and protein of plants were extracted according to Nagy et al. (40). PCR verification of gene insertion into plant genome was done with the following pairs of primers, which cover the DNA fragment from position 1248 to the end of bgl1: 5'-CAGTGACCGTGGATGCGACAATG-3' (SEQ ID NO:22) and 5'-AAAGGATCCTTAGT-GAACAGTAGGCAGAGACGC-3' (SEQ ID NO:23).

[0195] Identifying Transgenic Plants Expressing BGL1 Protein and Activity:

[0196] Western blot (40) and SDS-PAGE activity gel staining (37) were employed to screen successful transgenic lines, using the purified *A. niger* BGL1 protein as positive controls and non-transgenic plant as negative control.

[0197] SPMI-GC/MS Analysis:

[0198] The effect of bgl1 on flavor compound evolution and composition was studied. Fresh leaves of transgenic plants and of wild type control plants were excised and ground in liquid nitrogen. Ice-cold extraction buffer, containing 10 mM EDTA, 4 mM DTT in 50 mM phosphate buffer, pH 4.3, was added in a ratio of 1:3 w/w. The mixture was then shaken for 0.5 hours. 0.75 ml of supernatant from each of the centrifuged mixtures was taken into a glass vial. All manipulations were at 4° C. After 9 hours of incubation

at 37° C., the volatiles in the vial were analyzed according to Clark et al. (41) using a Saturn Varian 3800 SPMI-GC-MS apparatus, equipped with a DB-5 capillary column. The temperature of splitless injections was 250° C. and the transfer line was maintained at 280° C. Helium was used as a carrier gas. The oven was programmed as follows: 1 minute at 40° C. with gradually heating to 250° C. at a rate of 5° C./minute.

Experimental Results

[0199] Purification of Wild Type *A. niger* β -glucosidase:

[0200] *A. niger* β -glucosidase enzyme preparation was purified by MONO-Q™ (Amersham Biosciences Inc, Piscataway, N.J.) FPLC. Active protein samples eluted from the MONO-Q™ (Amersham Biosciences Inc, Piscataway, N.J.) anion exchange column were separated on a 10% SDS-PAGE gel, stained with coomassie blue, and incubated in the presence of MUGlc to demonstrate activity of the enzyme. At this stage of purification, a discrete band, having an apparent molecular mass of approximately 160 kDa and β -glucosidase activity could be detected (FIG. 2b, lanes 1-5: 1—electroeluted band of BGL1 from preparative PAGE-SDS gel stabs; 2-5—acetone precipitates from MONO-Q™ (Amersham Biosciences Inc, Piscataway, N.J.) anion exchange separation of BGL1). However, the apparent mass of the denatured enzyme (boiled for 10 min in the presence of β -mercaptoethanol), was shown to be 120 kDa on 10% SDS-PAGE (FIG. 2a). The enzyme was designated BGL1 was further purified to homogeneity on a RESOURCE-S™ (Amersham Biosciences Inc, Piscataway, N.J.) cation exchange column (FIG. 3). Deglycosylation of *A. niger* β -glucosidase was performed by N-glycosidase-F. As demonstrated in FIG. 4, SDS-PAGE analysis indicated that approximately 20 kDa of the *A. niger* β -glucosidase mass can be attributed to N-linked carbohydrates.

[0201] Proteolysis and N-Terminal Sequences of BGL1:

[0202] Partial enzymatic proteolysis with *Staphylococcus aureus* V8 protease of purified BGL1 was conducted. The undigested protein and cleavage products were separated by SDS-PAGE, followed by electroblotting onto PVDF membranes and determination of the N-terminal sequence of the native protein and two of the cleavage products. Amino acid sequences obtained were as follows:

[0203] N-terminal native protein: Asp-Glu-Leu-Ala-Tyr-Ser-Pro-Pro-Tyr-Tyr-Pro-Ser-Pro-Trp-Ala-Asn-Gly-Gln-Gly-Asp (SEQ ID NO:10). Underlined portion represents SEQ ID NO:4.

[0204] Internal cleavage product—E1 polypeptide: Val-Leu-Lys-His-Lys-Asn-Gly-Val-Phe-Thr-Ala-Thr-Asp-Asn-Trp-Ala-Ile-Asp-Gln-Ile-Glu-Ala-Leu-Ala-Lys (SEQ ID NO: 11).

[0205] Internal cleavage product—E2 polypeptide: Gly-Ala-Thr-Asp-Gly-Ser-Ala-Gln-Pro-Ile-Leu-Pro-Ala-Gly-Gly-Gly-Pro-Gly-Gly-Asn-Pro (SEQ ID NO:12). Underlined portion represents SEQ ID NO:6.

[0206] FastA analysis (32) indicated that the N-terminal sequence, as well as the internal sequences, have sequence similarity with sequences of β -glucosidase from the yeast *Saccharomyces fibuligera* which belonging to Family 3 of the glycosyl hydrolases.

[0207] Isolation and Characterization of bgl1 cDNA and Genomic DNA:

[0208] In order to clone the *A. niger* β -glucosidase gene, degenerate primers were designed according to the sequence of digest fragments of the polypeptide. These oligonucleotides were used to amplify a cDNA fragment of the β -glucosidase gene by RT-PCR. A 1.2 kb probe was excised from the resultant 2.2 kb amplification product and was used to screen a genomic library, constructed in pYEura3 yeast/*E. coli* shuttle vector. Positive clones were successfully subcloned and sequenced, resulting in full length bgl1 genomic sequence (SEQ ID NO:3, FIG. 5a). Amplification primers were then generated, according to the genomic DNA sequence, corresponding to the N- and C-terminal of the mature protein. RT-PCR was thereafter used for amplifying the full length β -glucosidase cDNA sequence (SEQ ID NO:1, FIG. 5a, GenBank Accession No. AJ132386). The cDNA sequence perfectly matched the DNA sequence of the combined exons. The open reading frame was found to encode a polypeptide with a predicted molecular weight of 92 kDa. The gene includes 7 exons intercepted by 6 introns (FIG. 5b). Analysis of the DNA sequence upstream to the sequence encoding for the mature protein revealed a putative leader sequence, intercepted by an 82 bp intron.

[0209] Production of rBGL1 in *E. coli*:

[0210] Recombinant BGL1 was overexpressed in *E. coli*. No apparent β -glucosidase activity could be detected in the *E. coli* extracts, however SDS-PAGE analysis revealed a relatively intense protein band expressed at the expected molecular weight. Western blot analysis using rabbit polyclonal anti-native BGL1 antibodies (AniLab Biological Services, Tal-Sachar, Israel), positively identified the 90 kDa protein band (not shown). Further analysis revealed that the protein was accumulated in inclusion bodies. Several refolding experiments were conducted, however, these efforts to produce active protein from *E. coli* failed (not shown).

[0211] Expression of Recombinant BGL1 in *S. cerevisiae* and *P. pastoris*:

[0212] Recombinant BGL1 was successfully expressed both in *S. cerevisiae* and *P. pastoris*. In *S. cerevisiae* a relatively low level of expression was found. The recombinant protein was detected by a Western blot analysis (FIG. 6a). The total protein extract of *S. cerevisiae* expressing bgl1 cDNA had a β -glucosidase activity of 1.9 units/mg protein. No β -glucosidase activity was detected in control *S. cerevisiae*, transformed with vector only, under the same assay conditions. However, no protein band corresponding to recombinant BGL1 could be detected by coomassie blue staining. *P. pastoris* transformed with bgl1 secreted relatively high levels of recombinant BGL1 to the medium (about 0.5 g/l) appearing as an almost pure protein in the culture supernatant (FIG. 6b). This recombinant enzyme was highly active (124 units/mg protein) and without further purification, yielded specific activity similar to that of the pure native enzyme.

[0213] ¹H-NMR Determination of Stereochemical Outcome:

[0214] ¹H-NMR spectra of a reaction mixture containing pNPGlc and BGL1 revealed that the beta anomer of glucose was formed first (H-1=4.95 ppm), with delayed appearance of the alpha anomer (H-1=5.59 ppm), the consequence of

mutarotation (FIG. 7). BGL1 is indeed, therefore, a retaining glycosidase, as has been observed for other family members (33, 34).

[0215] Inactivation and Reactivation of *A. niger* β -glucosidase:

[0216] Enzyme was incubated in the presence of various concentrations of 2FGlcF and residual enzyme activity was monitored at different time intervals. Enzyme activity decreased in a time-dependent manner, according to pseudo-first order kinetics, allowing the determination of pseudo-first order rate constants: $K_1=4.5 \text{ min}^{-1}$ and $K_2=35.4 \text{ mM}$, for inactivation at each inactivator concentration (0, 0.5, 1, 2, 4, and 6 mM, FIG. 8).

[0217] Rates of reactivation of 2-deoxy-2-fluoroglucosyl-BGL1 were determined in the presence of different concentrations of linamarin by monitoring activity regain after 0, 10, 20 and 30 min (FIG. 9). The regain of activity followed a first order process at each linamarin concentration.

[0218] Thermal stability of *A. niger* β -glucosidase:

[0219] Thermal stability of the recombinant enzyme was evaluated at different temperatures, presented as percent enzymatic activity relative to an enzyme solution kept at 4° C. Results obtained are summarized in Table 2 and illustrated in FIG. 10. The purified enzyme exhibits high thermal stability, as majority (above 50%) of the activity is maintained at a temperature ranging from 4-60° C.

TABLE 2

Temp. ° C.	% activity
4	100
50	91.5
55	83.5
60	68
65	17.8

[0220] Expression of BGL1 in Tobacco Plants:

[0221] *Agrobacterium* mediated leaf disc transformation resulted in transgenic tobacco plants as was proved by PCR (FIG. 12) for the presence of the transgene, Western blotting (FIGS. 13a-b) for presence of the protein and activity assays (FIGS. 14 and 15) for presence of protein activity. Table 3 below summarizes the results.

TABLE 3

	Gene construct		
	BGL1	Cell + BGL1 + HDEL,	Cell + BGL1
Number of Regenerates	33	14	27
PCR positive	29	9	23
Western Blot positive	4	9	18
Activity gel positive	0	9	18

[0222] Of the 29 PCR positive regenerates transformed with cDNA encoding BGL1, which fails to encode a signal peptide, only in 4 the BGL1 protein was detectable via Western blotting, however no BGL1 activity was measur-

able in any of which. The BGL1 was found smaller in molecular weight compared to wild type *A. niger* beta-glucosidase and of processed recombinant BGL1 containing a signal peptide. Its apparent size of about 95 kDa is very close to 92 kDa which is the calculated molecular weight of the un-glycosylated *A. niger* beta-glucosidase. This result coincides with the fact that a protein with no signal peptide is expected to be released from the ribosomes and remain in the cytoplasm (42) un-glycosylated, as protein glycosylation is conducted in the lumen of the endoplasmic reticulum (43).

[0223] Of the 9 PCR positive regenerates transformed with a cDNA encoding the BGL1 and a Cel1 signal peptide and in addition encodes the HDEL, ER retaining peptide, all plants expressed detectable amounts of BGL1 protein and activity.

[0224] Of the 23 PCR positive regenerates transformed with a cDNA which encodes the BGL1 protein and the Cel1 signal peptide but not the HDEL, ER retaining peptide, 18 plants expressed detectable amounts of BGL1 protein and activity.

[0225] The Effect of BGL1 on Flavor Compound Evolution and Composition in Transgenic Tobacco Plants:

[0226] Extracts of transgenic plants (CB14 and CBT21 containing similar BGL1 activity, see FIG. 15) were incubated for 9 hours at 37° C., and flavor compounds were analyzed by SPMI-GC/MS. The results, which are summarized in Table 4 below, show that with the exception of oleyl alcohol, the concentration of different flavor compounds is increased in transgenic plants expressing active BGL1 compared with the control. Furthermore, it seems that compartmentalization of BGL1 in the ER (or for that matter, any other subcellular organelle), rather than its secretion to the apoplast, results in higher release of flavor compounds. It is likely that this is resulted from the localization many flavor compounds in the apoplast, thus, secretion of BGL1 to the apoplast cause in vivo release of flavor compounds, while compartmentalization of BGL1 in the ER results in release of flavor compounds only in the event of cell disruption and decompartmentalization.

TABLE 4

Retention Time (minutes)	Scan Name	CB14	CBT 21
3.917	419 Hexanal	- ^a	-
4.749	508 3-methyl-pentanoic acid	-	-
4.863	520 2-Hexenal	-	+ ^b
5.167	552 ?	-	+
6.564	702 1-Heptanol	-	-
7.1	752 ?	+	++ ^d
8.085	865 2-ethyl-1-pexanol	-	+
8.132	870 Limonene	++	+
8.194	877 2-methyl-phenol	-	+
10.653	1139 Menthol	+	+
11.757	1258 Nerol	-	+
12.039	1288 6-Quinolinol	-	+
12.1	1294 2-butyl-1-octanol	-	+
13.0	1458 ?	-	+
13.7	1466 ?	-	+
14.091	1507 Vitispirane	-	+
14.094	1516 4-[2,6,6-trimethyl-1-cyclohexen-1-yl] 3-Buten-1-one	+	++

TABLE 4-continued

Retention Time (minutes)	Scan Name	CB14	CBT 21
15.985	1710 ?	-	-
19.327	2069 Oleyl alcohol	-- ^c	--

CB14 - transgenic plant containing Cel1 signal peptide + BGL1;
CBT 21 - transgenic plant containing Cel1 signal peptide + BGL1 + HDEL, ER retaining peptide.

^{ns}- means no significant difference in concentration compared with wild type.

^b+ means significant increase compared with the wild type.

^c- means significant decrease compared with the wild type.

^d+ means significant increase compared with a respective mark "+".

? - unknown compound.

[0227] Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications and variations that fall within the spirit and broad scope of the appended claims. All publications, patents, patent applications and sequences identified by GenBank accession numbers mentioned in this specification are herein incorporated in their entirety by reference into the specification, to the same extent as if each individual publication, patent, patent application or sequence was specifically and individually indicated to be incorporated herein by reference. In addition, citation or identification of any reference in this application shall not be construed as an admission that such reference is available as prior art to the present invention.

REFERENCES

- [0228] 1. Bause, E., and Legler, G. (1974) Hoppe-Seyler's Z. Physiol. Chem. 355, 438-442.
- [0229] 2. Beguin, P., and Aubert, J. P. (1994) FEMS Microbiol. Rev. 13, 25-58.
- [0230] 3. Rombouts, F. M. and Pilnik, W. (1978), Proc. Biochem. 8, 9-13.
- [0231] 4. Sternberg, D., Vijayakumar, P. and Reese, E. T. (1977), Can. J. Microbiol. 23, 139-147.
- [0232] 5. Woodward, J. and Wiseman, A. (1982), Enz. Microbiol. Technol. 4, 73-79.
- [0233] 6. Kems, G., Dalchow, E., Klappach, G. and Meyer, D. (1986), Acta. Biotechnol. 6(4), 355-359.
- [0234] 7. Shoseyov, O., Bravdo, B., Ikan, R. and Chet, I. (1988), Phytochem. 27(7), 1973-1976.
- [0235] 8. Shoseyov, O., Bravdo, B., Siegel, D., Goldman, A., Cohen, S., and Shoseyov, L. (1990), J. Agric. Food. Chem. 39, 1387-1390.
- [0236] 9. Dekker, R. F. H. (1986), Biotechnol. Bioengin. 26, 1438-1442.
- [0237] 10. Kitprechavanich, V. M., Hayashi, M. and Nagai, S. (1986), Agric. Biol. Chem. 50, 1703-1711.
- [0238] 11. Yeoh, H. H., Tan, T. K. and Koh, S. K. (1986), Appl. Microbiol. Biotechnol. 25, 25-28.
- [0239] 12. Crouzet, J. and Chassagne, D. (1999) in Naturally Occurring Glycosides (Ikan, R., ed.) pp. 225-274, Wiley Press.
- [0240] 13. Prade, H., L. F. Mackenzie and S. G. Withers (1998) Carbohydr. Res. 305, 371-381.
- [0241] 14. Zarevucka M., Vacek, M., Wimmer, Z., Demnerova, K. and Mackova, M. (1998) Chirality 10, 676-68.
- [0242] 15. Yi, Q., Sarney, D. B., Khan, J. A. and Vulfson, E. N. (1998) Biotechnol. Bioeng. 60, 385-390.
- [0243] 16. McCleary, B. V. and Harrington, J. (1988), Methods Enzymology, 160, 575-583.
- [0244] 17. Watanabe, T., Sato, T., Yoshioka, S., Koshijima, T. and Kuwahara, M. (1992), Eur. J. Biochem. 209, 651-659.
- [0245] 18. Unno, T., Ide, K., Yazaki, T., Tanaka, Y., Nakakuki, T. and Okada, G. (1993), Biosci. Biotech. Biochem. 57(12), 2172-2173.
- [0246] 19. Le Traon, M. M. P., and Pellerin P., (1998) Enz. Micro. Technol., 22 (5) 374-382.
- [0247] 20. Penttila, M. E., Nevalainen, H. K. M., Raynal, A., and Knowles, J. K. C., (1984), Mol. Gen. Genet. 194, 494-499.
- [0248] 21. Henrissat, B., and Bairoch, A. (1996) Biochem. J. Lett. 316, 695-696
- [0249] 22. Sinnott, M. L. (1990) Chem. Rev. 90, 1171-1202.
- [0250] 23. McCarter, J., and Withers, S. G. (1994) Curr. Opin. Struct. Biol. 4, 885-892.
- [0251] 24. Davies, G., Sinnott, M. L., and Withers, S. G. (1998) in CBiological Catalysis (Sinnott, M. L., ed.) Vol. 1, pp. 119-208, Academic Press.
- [0252] 25. Legler, G., Roeser, K. R., and Illig, H. K. (1979) Eur. J. Biochem. 101, 85-92.
- [0253] 26. Roeser, K. R., and Legler, G. (1981) Biochem. Biophys. Acta. 657, 321-333.
- [0254] 27. Legler, G., Sinnott, M. L., and Withers, S. G. (1980) J. Chem. Soc., Perkin Trans. II, 1376-1383.
- [0255] 28. Cleveland, D. W., Fischer, S. G., Kirschner, M. W. and Laemmli, U. K. (1977), J. Biol. Chem. 252(3), 1102-1106.
- [0256] 29. Murray, M. G. and Thompson, W. F. (1980), Nucl. Acids Res. 8, 4321-4325.
- [0257] 30. Ito, H., Fukuda, Y., Murata, K. and Kimura, A. (1983), J. Bacteriol. 153(1), 163-168.
- [0258] 31. Wong, A. W., He, S., Grubb, J. H., Sly, W. S., and Withers S. G. (1998), J. Biol. Chem. 273, 34057-34062.
- [0259] 32. Lipman, D. J. and Pearson, W. R. (1985) Science, 227, 1435-1441.
- [0260] 33. Legler, G. (1968) Hoppe-Zeyler's Z. Physiol. Chem. 349, 767-774.
- [0261] 34. Hrmova, M. and Fincher, G. B. (1996) J. Biol. Chem. 271, 5277-5286.

- [0262] 35. Iwashita K., Todoroki, M., Kimura, H., Shimoi, H. and Ito, K. (1998) *Biosci. Biotechnol. and Biochem.* 62 (10), 1938-1946.
- [0263] 36. Rlow E. and Lane D. (1988) *Antibodies, a laboratory manual.* Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- [0264] 37. Siegel D., Marton I., Dekel M., Bravdo B. A., He S. M., Wither S. G., Shoseyov. O., 2000, Cloning, expression, characterization, and nucleophile identification of family 3, *Aspergillus niger* beta-glucosidase. *J. Biol. Chem.* 275(7), 4973-4980.
- [0265] 38. Engelen F. A. V., Molthoff J. W., Conner A. J., Nap J. P., Pereira A and Stiekema W., 1995, pBINPLUS: an improved plant transformation vector based on pBIN19. *Transgenic Res.* 4, 288-290.
- [0266] 39. Shani Z., Dekel M., Tsabary G., Shoseyov., 1997, Cloning and characterization of elongation specific endo-1,4-beta-glucanase (Cell) from *Arabidopsis thaliana*, *Plant Molecular Biol.* 34: 837-842.
- [0267] 40. Nagy F., Kay S. A. K., Chua N. H., 1989, Analysis of gene expression in transgenic plants. In *Plant Molecular Biology: manual* edited by Gelvin et al., Kluwer Academic Publishers.
- [0268] 41. Clark T. J., Bunch J. E., 1997, Qualitative and quantitative analysis of flavor additives on tobacco products using SPME-GC-Mass Spectroscopy. *J. Agric. Food Chem.* 45 (3), 844-849.
- [0269] 42. Lewin B., 1994, The apparatus for protein localization, In *Gene V*, Oxford University Press. Pp: 279-314.
- [0270] 43. Prodi J. A., 2000, The role of N-oligosaccharide endoplasmic reticulum processing reactions in glycoprotein folding and degradation. *Biochem. J.*, 348, 1-13.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 24

<210> SEQ ID NO 1

<211> LENGTH: 2583

<212> TYPE: DNA

<213> ORGANISM: *Aspergillus niger*

<400> SEQUENCE: 1

```

atgaggttca ctttgatcga ggcggtggct ctgactgccg tctcgtggc cagcgctgat      60
gaattggcct actccccacc gtattacca tccccttggg ccaatggcca gggcgactgg      120
gcgcaggcat accagcgcgc tgttgatatt gtctcgaaa tgacattgga tgagaaggtc      180
aatctgacca caggaactgg atgggaattg gaactatgtg ttggtcagac tggcgggttt      240
ccccgattgg gagttccggg aatgtgttta caggatagcc ctctgggctg tcgcgactcc      300
gactacaact ctgctttccc tgccggcatg aacgtggctg cgacctggga caagaatctg      360
gcataccttc gcgccaaggc tatgggtcag gaatttagtg acaagggtgc cgatatccaa      420
ttgggtccag ctgcccggcc tctcggtaga agtcccgcag gtggtcgtaa ctgggagggc      480
ttctccccag accctgcctt aagtgggtgt ctctttgccg agaccatcaa gggatccaa      540
gatgctggtg tggttgcgac ggctaagcac tacattgctt acgagcaaga gcatttccgt      600
caggcgctg aagcccaagg ttttgattt aatatttccg agagtgaag tgcgaaacctc      660
gatgataaga ctatgcacga gctgtacctc tggcccttcg cggatgccat ccgtgcaggt      720
gctggcgctg tgatgtgctc ctacaaccag atcaacaaca gttatggctg ccagaacagc      780
tacactctga acaagctgct caaggccgag ctgggcttcc agggctttgt catgagtgat      840
tgggctgctc accatgctgg tgtgagtggg gctttggcag gattggatat gtctatgcca      900
ggagacgtcg actacgcagc tggtagctct tactgggta caaactgac cattagcgtg      960
ctcaacggaa cggtgcccca atggcgtggt gatgacatgg ctgtccgcat catggccgcc      1020
tactacaagg tcggccgtga ccgtctgtgg actcctocca acttcagctc atggaccaga      1080
gatgaatacg gctacaagta ctactacgtg tcggaggagc cgtacgagaa ggtcaaccag      1140
tacgtgaatg tgcaacgcaa ccacagcga ctgattgcc gcattggagc ggacagcacg      1200

```

-continued

```

gtgctcctca agaacgacgg cgctctgcct ttgactggta aggagcgctt ggtcgcgctt 1260
atcggagaag atgcggggctc caacccttat ggtgccaacg gctgcagtga cctgggatgc 1320
gacaatggaa cattggcgat gggctgggga agtgggtactg ccaacttccc atacctggtg 1380
acccccgagc aggccatctc aaacgaggtg cttaaagcaca agaatggtgt attcaccgcc 1440
accgataact gggctatcga tcaaattgag gcgcttgcta agaccgccag tgtctctctt 1500
gtctttgtca acgccgactc tggtaggggt tacatcaatg tggacggaaa cctgggtgac 1560
cgcaggaacc tgacctgtg gaggaacggc gataatgtga tcaaggctgc tgctagcaac 1620
tgcaacaaca caatcgttgt cattcactct gtcggaccag tcttggttaa cgagtggtag 1680
gacaacccca atgttaccgc tatcctctgg ggtggtttgc ccggtcagga gtctggcaac 1740
tctcttgccg acgtcctcta tggccgtgtc aaccccggtg ccaagtcgcc ctttacctgg 1800
ggcaagactc gtgaggccta ccaagactac ttggtcaccg agcccaacaa cggcaacgga 1860
gcccctcagg aagactttgt cgaggcgctc ttcattgact accgtggatt tgacaagcgc 1920
aacgagacct cgatctacga gttcggctat ggtctgagct acaccacttt caactactcg 1980
aaccttgagg tgcaggtgct gagcgcccct gcatacagac ctgcttcggg tgagaccgag 2040
gcagcgccaa ccttcggaga ggttggaat gcgtcggatt acctctacc cagcggattg 2100
ctgagaatta ccaagttcat ctaccctgg ctcaacgta ccgatctcga ggcattctcc 2160
ggggatgcta gctacgggca ggactcctcc gactatcttc ccgagggagc caccgatggc 2220
tctgcgcaac cgatcctgcc tgccggtggc ggtcctggcg gcaaccctcg cctgtacgac 2280
gagctcatcc gcgtgtcagt gaccatcaag aacaccgga aggttgctgg tgatgaagtt 2340
ccccaactgt atgtttccct tggcggtccc aatgagccca agatcgtgct gcgtcaattc 2400
gagcgcatca cgctgcagcc gtcggaggag acgaagtgga gcacgactct gacgcgccgt 2460
gaccttgcaa actggaatgt tgagaagcag gactgggaga ttacgtcgtg tcccaagatg 2520
gtgtttgtcg gaagctcctc gcggaagctg ccgctccggg cgtctctgcc tactgttcac 2580
taa 2583

```

<210> SEQ ID NO 2

<211> LENGTH: 860

<212> TYPE: PRT

<213> ORGANISM: Aspergillus niger

<400> SEQUENCE: 2

```

Met Arg Phe Thr Leu Ile Glu Ala Val Ala Leu Thr Ala Val Ser Leu
1           5           10           15
Ala Ser Ala Asp Glu Leu Ala Tyr Ser Pro Pro Tyr Tyr Pro Ser Pro
20           25           30
Trp Ala Asn Gly Gln Gly Asp Trp Ala Gln Ala Tyr Gln Arg Ala Val
35           40           45
Asp Ile Val Ser Gln Met Thr Leu Asp Glu Lys Val Asn Leu Thr Thr
50           55           60
Gly Thr Gly Trp Glu Leu Glu Leu Cys Val Gly Gln Thr Gly Gly Val
65           70           75           80
Pro Arg Leu Gly Val Pro Gly Met Cys Leu Gln Asp Ser Pro Leu Gly
85           90           95
Val Arg Asp Ser Asp Tyr Asn Ser Ala Phe Pro Ala Gly Met Asn Val

```

-continued

100					105					110					
Ala	Ala	Thr	Trp	Asp	Lys	Asn	Leu	Ala	Tyr	Leu	Arg	Gly	Lys	Ala	Met
		115					120					125			
Gly	Gln	Glu	Phe	Ser	Asp	Lys	Gly	Ala	Asp	Ile	Gln	Leu	Gly	Pro	Ala
	130					135					140				
Ala	Gly	Pro	Leu	Gly	Arg	Ser	Pro	Asp	Gly	Gly	Arg	Asn	Trp	Glu	Gly
145					150					155				160	
Phe	Ser	Pro	Asp	Pro	Ala	Leu	Ser	Gly	Val	Leu	Phe	Ala	Glu	Thr	Ile
				165					170					175	
Lys	Gly	Ile	Gln	Asp	Ala	Gly	Val	Val	Ala	Thr	Ala	Lys	His	Tyr	Ile
			180					185					190		
Ala	Tyr	Glu	Gln	Glu	His	Phe	Arg	Gln	Ala	Pro	Glu	Ala	Gln	Gly	Phe
		195					200					205			
Gly	Phe	Asn	Ile	Ser	Glu	Ser	Gly	Ser	Ala	Asn	Leu	Asp	Asp	Lys	Thr
	210					215					220				
Met	His	Glu	Leu	Tyr	Leu	Trp	Pro	Phe	Ala	Asp	Ala	Ile	Arg	Ala	Gly
225					230					235					240
Ala	Gly	Ala	Val	Met	Cys	Ser	Tyr	Asn	Gln	Ile	Asn	Asn	Ser	Tyr	Gly
				245					250					255	
Cys	Gln	Asn	Ser	Tyr	Thr	Leu	Asn	Lys	Leu	Leu	Lys	Ala	Glu	Leu	Gly
			260					265					270		
Phe	Gln	Gly	Phe	Val	Met	Ser	Asp	Trp	Ala	Ala	His	His	Ala	Gly	Val
		275					280					285			
Ser	Gly	Ala	Leu	Ala	Gly	Leu	Asp	Met	Ser	Met	Pro	Gly	Asp	Val	Asp
	290					295					300				
Tyr	Asp	Ser	Gly	Thr	Ser	Tyr	Trp	Gly	Thr	Asn	Leu	Thr	Ile	Ser	Val
305					310					315					320
Leu	Asn	Gly	Thr	Val	Pro	Gln	Trp	Arg	Val	Asp	Asp	Met	Ala	Val	Arg
				325					330					335	
Ile	Met	Ala	Ala	Tyr	Tyr	Lys	Val	Gly	Arg	Asp	Arg	Leu	Trp	Thr	Pro
			340					345					350		
Pro	Asn	Phe	Ser	Ser	Trp	Thr	Arg	Asp	Glu	Tyr	Gly	Tyr	Lys	Tyr	Tyr
		355					360					365			
Tyr	Val	Ser	Glu	Gly	Pro	Tyr	Glu	Lys	Val	Asn	Gln	Tyr	Val	Asn	Val
	370					375					380				
Gln	Arg	Asn	His	Ser	Glu	Leu	Ile	Arg	Arg	Ile	Gly	Ala	Asp	Ser	Thr
385					390					395					400
Val	Leu	Leu	Lys	Asn	Asp	Gly	Ala	Leu	Pro	Leu	Thr	Gly	Lys	Glu	Arg
				405					410					415	
Leu	Val	Ala	Leu	Ile	Gly	Glu	Asp	Ala	Gly	Ser	Asn	Pro	Tyr	Gly	Ala
			420					425					430		
Asn	Gly	Cys	Ser	Asp	Arg	Gly	Cys	Asp	Asn	Gly	Thr	Leu	Ala	Met	Gly
		435					440					445			
Trp	Gly	Ser	Gly	Thr	Ala	Asn	Phe	Pro	Tyr	Leu	Val	Thr	Pro	Glu	Gln
	450					455					460				
Ala	Ile	Ser	Asn	Glu	Val	Leu	Lys	His	Lys	Asn	Gly	Val	Phe	Thr	Ala
465					470					475					480
Thr	Asp	Asn	Trp	Ala	Ile	Asp	Gln	Ile	Glu	Ala	Leu	Ala	Lys	Thr	Ala
				485					490					495	
Ser	Val	Ser	Leu	Val	Phe	Val	Asn	Ala	Asp	Ser	Gly	Glu	Gly	Tyr	Ile
			500					505					510		

-continued

Asn Val Asp Gly Asn Leu Gly Asp Arg Arg Asn Leu Thr Leu Trp Arg
 515 520 525
 Asn Gly Asp Asn Val Ile Lys Ala Ala Ala Ser Asn Cys Asn Asn Thr
 530 535 540
 Ile Val Val Ile His Ser Val Gly Pro Val Leu Val Asn Glu Trp Tyr
 545 550 555 560
 Asp Asn Pro Asn Val Thr Ala Ile Leu Trp Gly Gly Leu Pro Gly Gln
 565 570 575
 Glu Ser Gly Asn Ser Leu Ala Asp Val Leu Tyr Gly Arg Val Asn Pro
 580 585 590
 Gly Ala Lys Ser Pro Phe Thr Trp Gly Lys Thr Arg Glu Ala Tyr Gln
 595 600 605
 Asp Tyr Leu Val Thr Glu Pro Asn Asn Gly Asn Gly Ala Pro Gln Glu
 610 615 620
 Asp Phe Val Glu Gly Val Phe Ile Asp Tyr Arg Gly Phe Asp Lys Arg
 625 630 635 640
 Asn Glu Thr Pro Ile Tyr Glu Phe Gly Tyr Gly Leu Ser Tyr Thr Thr
 645 650 655
 Phe Asn Tyr Ser Asn Leu Glu Val Gln Val Leu Ser Ala Pro Ala Tyr
 660 665 670
 Glu Pro Ala Ser Gly Glu Thr Glu Ala Ala Pro Thr Phe Gly Glu Val
 675 680 685
 Gly Asn Ala Ser Asp Tyr Leu Tyr Pro Ser Gly Leu Leu Arg Ile Thr
 690 695 700
 Lys Phe Ile Tyr Pro Trp Leu Asn Gly Thr Asp Leu Glu Ala Ser Ser
 705 710 715 720
 Gly Asp Ala Ser Tyr Gly Gln Asp Ser Ser Asp Tyr Leu Pro Glu Gly
 725 730 735
 Ala Thr Asp Gly Ser Ala Gln Pro Ile Leu Pro Ala Gly Gly Gly Pro
 740 745 750
 Gly Gly Asn Pro Arg Leu Tyr Asp Glu Leu Ile Arg Val Ser Val Thr
 755 760 765
 Ile Lys Asn Thr Gly Lys Val Ala Gly Asp Glu Val Pro Gln Leu Tyr
 770 775 780
 Val Ser Leu Gly Gly Pro Asn Glu Pro Lys Ile Val Leu Arg Gln Phe
 785 790 795 800
 Glu Arg Ile Thr Leu Gln Pro Ser Glu Glu Thr Lys Trp Ser Thr Thr
 805 810 815
 Leu Thr Arg Arg Asp Leu Ala Asn Trp Asn Val Glu Lys Gln Asp Trp
 820 825 830
 Glu Ile Thr Ser Tyr Pro Lys Met Val Phe Val Gly Ser Ser Ser Arg
 835 840 845
 Lys Leu Pro Leu Arg Ala Ser Leu Pro Thr Val His
 850 855 860

<210> SEQ ID NO 3
 <211> LENGTH: 3885
 <212> TYPE: DNA
 <213> ORGANISM: Aspergillus niger

<400> SEQUENCE: 3

tccattcgcc catgcttagc gtgtcttttc tttgaacct gcatgcgga ctgtgaattg 60

-continued

catgagtg	gg tagctttg	cg gagacag	ctg cactgg	cata catcat	cggt gggttc	ctca 120																	
attgc	catgc cgtggc	ggac ggtca	ctttg tggcg	ctcaa actat	ttaat atggcc	cage 180																	
tccc	cttct ctcg	ctgttt	tcgttt	ctgt cctcc	ctaaa cctcc	agtct ctccatt	gga 240																
cagg	gtttg	acgg	ttgctc	acct	gggg	cg acct	gccat 300																
catg	agg	ttc	aggg	ttg	ctc	gtct	cgctgg ccagc	gctgt 360															
acgt	g	cg	tt	ct	c	g	agaattgca attgtg	ctta attag	attca tttg	ttgt	g 420												
tcat	catgc	tgaca	aatg	cttt	catag	gatga	aatg	cctact	cccc	accg	tattac 480												
ccat	cccc	ctt	ggc	caatg	ccagg	gcgac	tgggc	gcagg	cata	accag	cgctgt	gat 540											
attg	tctgc	aaatg	acatt	ggatg	agaag	gtca	aatctga	ccac	agga	ac	tggg	tagggc 600											
ttac	atgg	cg	caat	ctgt	at	gctcc	ggcta	aca	actt	cta	catgg	gaatt gga	actat	gt 660									
gttg	gtc	aga	ctgt	gt	tcccc	ggtag	gtttg	aaaat	attg	t	cgaga	caggg	gacat 720										
tatt	gatt	aa	cggtg	acaga	ttggg	agttc	cgga	aatgtg	ttt	acag	gat	agcc	ctctg 780										
gcgt	t	cg	ca	ctg	taag	cca	tctg	ctgt	ttag	gct	tc	tgac	acgg	cg 840									
cagc	gacta	caact	ctg	ctg	ctg	ccg	gc	atga	acgt	ggct	gca	acc	tggg	aca	aga 900								
atct	gg	cata	ccttc	cg	gc	aag	gctat	g	tc	agga	att	tagt	gaca	ag	ggtg	ccgata 960							
tcca	att	ggg	tccag	ctg	cc	ctc	ctg	gtaga	agtcc	cgac	g	tg	ta	act	ggg	1020							
aggg	ctt	ctc	cccag	acct	gcc	ct	aatg	gtg	t	ctct	tgcc	gag	acc	atca	aggg	ta 1080							
tcca	ag	atgc	tgg	t	g	g	g	g	g	g	g	g	g	g	g	g	1140						
tccg	t	cag	gc	g	cc	ta	ag	cc	caag	gttt	g	att	ta	at	ttcc	gag	agt	gga	agt	gcga 1200			
acct	cg	atga	taag	act	atg	cacg	agct	gt	acct	t	ggcc	ctg	ctt	cg	cg	gat	gcc	at	cc	ctg 1260			
cagg	t	ctg	gc	tg	ct	ct	ac	acc	agat	caa	caac	agt	t	at	gg	t	g	cc	aga 1320				
acag	t	ac	ac	t	ctga	aca	ag	ctg	ct	caag	cg	ag	ct	ggg	ctt	cc	ag	ggc	ttg	t	catga 1380		
gtg	att	ggg	gc	tg	ct	ac	cat	gct	g	gtg	t	g	g	g	g	g	g	g	g	g	g	1440	
tgcc	ag	gaga	cg	tc	gact	ac	gac	ag	tg	ta	g	g	g	g	g	g	g	g	g	g	g	1500	
gcgt	g	ct	ca	cg	ga	ac	gg	t	g	ct	g	at	g	ct	g	ct	g	ct	g	ct	g	1560	
ccgc	ct	acta	ca	ag	g	ctg	gc	ct	g	g	act	cc	t	cc	a	act	t	ag	ct	cat	gga 1620		
ccag	ag	atga	ata	cg	gct	ac	aa	g	t	act	act	ac	g	t	g	tc	g	g	g	acc	gtac	gaga	aggtca 1680
accag	t	ac	g	ta	ctg	caa	cg	ca	acc	aca	gc	ga	act	gat	tc	g	cc	g	cat	gg	agc	ggaca 1740	
gcac	g	g	g	t	g	ct	g	ct	ct	g	ac	tg	g	ta	ag	gag	cg	ct	g	g	ct	g	1800
cgct	t	at	ctg	aga	ag	atg	cg	g	ct	cca	acc	ct	t	at	g	g	t	g	ca	ac	g	g	1860
gatg	cg	ca	aa	t	g	g	ct	g	g	g	g	g	g	g	g	g	g	g	g	g	g	g	1920
tggt	g	ac	cc	cg	ag	cagg	cc	at	ct	caa	acg	ag	gt	g	ct	ta	gc	aca	aga	aat	ggt	g	attca 1980
ccgc	ca	cc	ga	ta	act	gg	g	ct	at	ca	aaa	ttg	ag	gc	g	ct	ta	ag	acc	gcc	ag	g	taag 2040
aa	g	at	cccc	att	ct	tt	ct	g	t	g	ca	at	g	ct	g	ca	aca	at	g	ct	g	ct	ctc 2100
ttg	t	ct	tt	g	ca	ac	gc	g	ac	g	g	g	g	g	g	g	g	g	g	g	g	g	2160
accg	ca	gg	aa	cctg	ac	cctg	tgg	ag	ga	acc	gc	g	ata	at	gt	gat	ca	ag	g	ct	g	ct	agca 2220
actg	ca	ca	aa	ctg	ct	g	ct	ct	g	g	ac	ag	t	ct	g	g	tt	aac	g	ag	t	g	gt 2280
acg	ca	ac	cc	caat	g	t	at	ct	ct	ct	g	gg	g	t	g	g	tt	g	cc	cg	g	ct	agca 2340

-continued

```

actctcttgc cgacgtcctc tatggccgtg tcaaccccg tccaagtgc ccctttacct 2400
ggggcaagac tcgtgaggcc taccaagact acttggtcac cgagcccaac aacggcaacg 2460
gagcccctca ggaagacttt gtcgagggcg tcttcattga ctaccgtgga ttgacaagc 2520
gcaacgagac cccgatctac gagttcggct atggtctgag ctacaccact ttcaactact 2580
cgaaccttga ggtgcagggt ctgagcggcc ctgcatacga gcctgcttcg ggtgagaccg 2640
aggcagcgcc aaccttcgga gaggttgaa atgcgtcggg ttacctctac cccagcggat 2700
tgctgagaat taccaagttc atctaccctt ggctcaacgg taccgatctc gaggcatctt 2760
ccggggatgc tagctacggg caggactcct ccgactatct tcccagggga gccaccgatg 2820
gctctgcgca accgatcctg cctgcoggtg gcggtcctgg cggcaacct cgctgtacg 2880
acgagctcat ccgctgtgca gtgaccatca agaaccgg caaggttgcg ggtgatgaag 2940
ttcccact ggtaagtaa catgaggtcc gaacgaggtt gaacaaagct aatcagtcgc 3000
agtatgtttc ccttgcggtt cccaatgag ccaagatcgt gctgcgtcaa ttcgagcgca 3060
tcacgctgca gccgtcggag gagacgaagt ggagcagcgc tctgacgcgc cgtgacctg 3120
caactggaa tgttgagaag caggactggg agattacgct gtatccaag atggtgtttg 3180
tcggaagctc ctgcggaag ctgccgctcc gggcgtctct gcctactggt cactaaatag 3240
ctctcaaatg gtataccatg atggcoggtg tatatgaatt aatgatttat gccaacagca 3300
agaccactgt agatgtagat gtagaatgag tattgcgtag tagcgtgtag atgatgatac 3360
aagcgatccg acacatggta ggaagagtgg cgctagttgg ggcggaaacc aagcgacgtc 3420
atccgctgcc gacttcgcca gtctttcttc ttttctctt cagcctctt cctccgctta 3480
atccagcaac cattgccaat tgctctaca acaactaatt gccataatac tctactccta 3540
ttcaatatat acaccacaat ctgcacataa tcacacaagc ctgaacacac gagcaacct 3600
gccctctccc gatcctccag ccccagcgt acgacccttc caaccacca taacagcgt 3660
cctcatctac ccagcgacc taatcgtggg atcactcttc tccgtcctct ctcccaccg 3720
acaaggcaca cgcgacgac gctccagcac cctccacca cacgtcgagc ccctagcccc 3780
gtccatcgcg tcagacctca acctctcctt tctctccgcg cgcgccgca actacttcgc 3840
tcgcaagac aacatcttca atctatattc gtcaaagtgc gctgg 3885

```

```

<210> SEQ ID NO 4
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Aspergillus niger

```

```

<400> SEQUENCE: 4

```

```

Ser Pro Pro Tyr Tyr Pro
1           5

```

```

<210> SEQ ID NO 5
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Single strand DNA oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:

```

-continued

<221> NAME/KEY: misc_feature
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 5

snccncnta ytaycc 16

<210> SEQ ID NO 6
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Aspergillus niger

<400> SEQUENCE: 6

Gln Pro Ile Leu Pro Ala Gly Gly
1 5

<210> SEQ ID NO 7
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Single strand DNA oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (15)..(15)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 7

tccgcnggna rdatnggytg 20

<210> SEQ ID NO 8
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Single strand DNA oligonucleotide

<400> SEQUENCE: 8

aaaccatggc tgatgaattg gcatactccc cacc 34

<210> SEQ ID NO 9
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Single strand DNA oligonucleotide

<400> SEQUENCE: 9

aaaggatcct tagtgaacag taggcagaga cgc 33

<210> SEQ ID NO 10
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Aspergillus niger

-continued

```

<220> FEATURE:
<221> NAME/KEY: misc_feature
<223> OTHER INFORMATION: A peptide derived from partial V8 proteolysis
of purified BGL1

<400> SEQUENCE: 10

Asp Glu Leu Ala Tyr Ser Pro Pro Tyr Tyr Pro Ser Pro Trp Ala Asn
1          5          10          15

Gly Gln Gly Asp
          20

<210> SEQ ID NO 11
<211> LENGTH: 25
<212> TYPE: PRT
<213> ORGANISM: Aspergillus niger
<220> FEATURE:
<221> NAME/KEY: misc_feature
<223> OTHER INFORMATION: A peptide derived from partial V8 proteolysis
of purified BGL1

<400> SEQUENCE: 11

Val Leu Lys His Lys Asn Gly Val Phe Thr Ala Thr Asp Asn Trp Ala
1          5          10          15

Ile Asp Gln Ile Glu Ala Leu Ala Lys
          20          25

<210> SEQ ID NO 12
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Aspergillus niger
<220> FEATURE:
<221> NAME/KEY: misc_feature
<223> OTHER INFORMATION: A peptide derived from partial V8 proteolysis
of purified BGL1

<400> SEQUENCE: 12

Gly Ala Thr Asp Gly Ser Ala Gln Pro Ile Leu Pro Ala Gly Gly Gly
1          5          10          15

Pro Gly Gly Asn Pro
          20

<210> SEQ ID NO 13
<211> LENGTH: 3212
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: An expression cassettes used for expression of
A. niger
beta-glucosidase in tobacco plants

<400> SEQUENCE: 13

gaattcccga tcctatctgt cacttcatca aaaggacagt agaaaaggaa ggtggcacta    60
caaatgccat cattgcgata aaggaaaggc tatcgttcaa gatgcctctg ccgacagtgg    120
tccc aaagat ggacccccac ccacgaggag catcgtggaa aaagaagacg ttccaaccac    180
gttttcaaag caagtggatt gatgtgatat ctccactgac gtaagggatg acgcacaatc    240
ccactatcct tcgcaagacc cttcctctat ataaggaagt tcatttcatt tggagaggac    300
aggcttcttg agatccttca acaattacca acaacaacaa acaacaaaca acattacaat    360
tactatttac aattacagtc gaccatggct gatgaattgg cctactcccc accgtattac    420
ccatcccctt gggccaatgg ccagggcgac tgggcgcagg cataccagcg cgctgttgat    480

```

-continued

attgtctcgc aaatgacatt ggatgagaag gtcaatctga ccacaggaac tggatgggaa	540
ttggaactat gtgttggtca gactggcggg gttccccgat tgggagttcc gggaaatgtgt	600
ttacaggata gccctctggg cgttcgagac tccgactaca actctgcttt ccctgccggc	660
atgaacgtgg ctgcgacctg ggacaagaat ctggcatacc ttcgcgcaa ggctatgggt	720
caggaattta gtgacaaggg tgccgataac caattgggtc cagctgccgg ccctctcggg	780
agaagtcccg acggtggtcg taactgggag ggcttctccc cagaccctgc cctaagtggg	840
gtgctctttg ccgagacct caagggtatc caagatgctg gtgtggttc gacggctaag	900
cactacattg cttacgagca agagcatttc cgtcaggcgc ctgaagccca aggttttggg	960
tttaaatatt ccgagagtgg aagtgcgaac ctcgatgata agactatgca cgagctgtac	1020
ctctggccct tcgcggtatc catccgtgca ggtgctggcg ctgtgatgtg ctctacaac	1080
cagatcaaca acagtattgg ctgccagaac agctacactc tgaacaagct gctcaaggcc	1140
gagctgggct tccagggctt tgtcatgagt gattgggctg ctcaccatgc tgggtgagt	1200
ggtgctttgg caggattgga tatgtctatg ccaggagacg tcgactacga cagtgtacg	1260
tcttactggg gtacaaactt gaccattagc gtgctcaacg gaacggtgcc ccaatggcgt	1320
gttgatgaca tggctgtccg catcatggcc gcctactaca aggtcggccg tgaccgtctg	1380
tggactcctc ccaacttcag ctcatggacc agagatgaat acggctacaa gtactactac	1440
gtgtcggagg gaccgtacga gaaggtaac cagtacgtga atgtgcaacg caaccacagc	1500
gaactgattc gccgcatggg agcggacagc acggtgctcc tcaagaacga cggcgctctg	1560
cctttgactg gtaaggagcg cctggtcgcg cttatcggag aagatgcggg ctccaaccct	1620
tatggtgcca acggctgcag tgaccgtgga tgcgacaatg gaacattggc gatgggctgg	1680
ggaagtggta ctgccaactt cccatacctg gtgacccccg agcaggccat ctcaaacgag	1740
gtgcttaagc acaagaatgg tgtattcacc gccaccgata actgggctat cgatcaaatt	1800
gaggcgcttg ctaagaccgc cagtgtctct cttgtctttg tcaacgcoga ctctggtgag	1860
ggttacatca atgtggacgg aaacctgggt gaccgcagga acctgacct gtggaggaac	1920
ggcgataatg tgatcaaggc tgctgctagc aactgcaaca acacaatcgt tgcattcac	1980
tctgtcggac cagtcttggg taacgagtgg tacgacaacc ccaatgttac cgctatcctc	2040
tggggtgggt tgcccgtca ggagtctggc aactctcttg ccgacgtcct ctatggccgt	2100
gtcaaccccc gtgccaagtc gccctttacc tggggcaaga ctctgtaggc ctaccaagac	2160
tacttggtca ccgagcccaa caacggcaac ggagccccctc aggaagactt tgotgaggc	2220
gtcttcattg actaccgtgg atttgacaag cgcaacgaga ccccgatcta cgagttcggc	2280
tatggtctga gtacaccac tttcaactac tcgaacctg aggtgcagggt gctgagcgcc	2340
ctctgatacg agcctgcttc gggtagacc gaggcagcgc caacctcgg agaggttggg	2400
aatgcgtcgg attacctcta cccagcggg ttgctgagaa ttaccaagtt catctacccc	2460
tggctcaacg gtaccgatct cgaggcatct tccgggatg ctgctacgg gcaggactcc	2520
tccgactatc tccccaggg agccaocgat ggctctgcgc aaccgatoct gcctgccggg	2580
ggcgtcctg gcggcaaccc tcgcctgtac gacgagctca tccgctgtc agtgaccatc	2640
aagaacaccg gcaaggttgc tgggtatgaa gttccccaac tgtatgttcc ccttggcggg	2700
cccaatgagc ccaagatcgt gctgcgtcaa ttcgagcga tcacgctgca gccgtcggg	2760

-continued

```

gagacgaagt ggagcacgac tctgacgcgc cgtgaccttg caaactggaa tgttgagaag 2820
caggactggg agattacgtc gtatcccaag atggtgtttg tcggaagctc ctgcggaag 2880
ctgccgctcc gggcgtctct gcctactgtt cactaacccg ggcgagctcg aattgatcgt 2940
tcaaacattt ggcaataaag tttcttaaga ttgaatcctg ttgccgttct tgcgatgatt 3000
atcatataat ttctgttgaa ttacgttaag catgtaataa ttaaacaatgt aatgcatgac 3060
gttatttatg agatgggggt tttatgatta agagtccccc caattataca ttttaatacg 3120
cgatagaaaa acaaaatata gcgcccaaac taaggataaa attattcgcg ccgcgggggg 3180
gcattctatg gttactagat ctctagaatt cc 3212

```

<210> SEQ ID NO 14

<211> LENGTH: 841

<212> TYPE: PRT

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: the protein coded by An expression cassettes used for expression of A. niger beta-glucosidase in tobacco plants

<400> SEQUENCE: 14

```

Asp Glu Leu Ala Tyr Ser Pro Pro Tyr Tyr Pro Ser Pro Trp Ala Asn
1      5      10      15
Gly Gln Gly Asp Trp Ala Gln Ala Tyr Gln Arg Ala Val Asp Ile Val
20     25     30
Ser Gln Met Thr Leu Asp Glu Lys Val Asn Leu Thr Thr Gly Thr Gly
35     40     45
Trp Glu Leu Glu Leu Cys Val Gly Gln Thr Gly Gly Val Pro Arg Leu
50     55     60
Gly Val Pro Gly Met Cys Leu Gln Asp Ser Pro Leu Gly Val Arg Asp
65     70     75     80
Ser Asp Tyr Asn Ser Ala Phe Pro Ala Gly Met Asn Val Ala Ala Thr
85     90     95
Trp Asp Lys Asn Leu Ala Tyr Leu Arg Gly Lys Ala Met Gly Gln Glu
100    105   110
Phe Ser Asp Lys Gly Ala Asp Ile Gln Leu Gly Pro Ala Ala Gly Pro
115    120   125
Leu Gly Arg Ser Pro Asp Gly Gly Arg Asn Trp Glu Gly Phe Ser Pro
130    135   140
Asp Pro Ala Leu Ser Gly Val Leu Phe Ala Glu Thr Ile Lys Gly Ile
145    150   155   160
Gln Asp Ala Gly Val Val Ala Thr Ala Lys His Tyr Ile Ala Tyr Glu
165    170   175
Gln Glu His Phe Arg Gln Ala Pro Glu Ala Gln Gly Phe Gly Phe Asn
180    185   190
Ile Ser Glu Ser Gly Ser Ala Asn Leu Asp Asp Lys Thr Met His Glu
195    200   205
Leu Tyr Leu Trp Pro Phe Ala Asp Ala Ile Arg Ala Gly Ala Gly Ala
210    215   220
Val Met Cys Ser Tyr Asn Gln Ile Asn Asn Ser Tyr Gly Cys Gln Asn
225    230   235   240
Ser Tyr Thr Leu Asn Lys Leu Leu Lys Ala Glu Leu Gly Phe Gln Gly
245    250   255

```


-continued

660				665				670							
Ser	Asp	Tyr	Leu	Tyr	Pro	Ser	Gly	Leu	Leu	Arg	Ile	Thr	Lys	Phe	Ile
	675						680					685			
Tyr	Pro	Trp	Leu	Asn	Gly	Thr	Asp	Leu	Glu	Ala	Ser	Ser	Gly	Asp	Ala
	690						695				700				
Ser	Tyr	Gly	Gln	Asp	Ser	Ser	Asp	Tyr	Leu	Pro	Glu	Gly	Ala	Thr	Asp
	705				710					715					720
Gly	Ser	Ala	Gln	Pro	Ile	Leu	Pro	Ala	Gly	Gly	Gly	Pro	Gly	Gly	Asn
			725						730					735	
Pro	Arg	Leu	Tyr	Asp	Glu	Leu	Ile	Arg	Val	Ser	Val	Thr	Ile	Lys	Asn
			740						745				750		
Thr	Gly	Lys	Val	Ala	Gly	Asp	Glu	Val	Pro	Gln	Leu	Tyr	Val	Ser	Leu
		755					760					765			
Gly	Gly	Pro	Asn	Glu	Pro	Lys	Ile	Val	Leu	Arg	Gln	Phe	Glu	Arg	Ile
	770					775					780				
Thr	Leu	Gln	Pro	Ser	Glu	Glu	Thr	Lys	Trp	Ser	Thr	Thr	Leu	Thr	Arg
	785				790					795					800
Arg	Asp	Leu	Ala	Asn	Trp	Asn	Val	Glu	Lys	Gln	Asp	Trp	Glu	Ile	Thr
				805					810					815	
Ser	Tyr	Pro	Lys	Met	Val	Phe	Val	Gly	Ser	Ser	Ser	Arg	Lys	Leu	Pro
			820					825				830			
Leu	Arg	Ala	Ser	Leu	Pro	Thr	Val	His							
	835						840								

<210> SEQ ID NO 15

<211> LENGTH: 3329

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: a cassette encoding a BGL1 fused to a Cell signal peptide for secretion into the apoplast

<400> SEQUENCE: 15

```

gaattcccga tcctatctgt cacttcatca aaaggacagt agaaaaggaa ggtggcacta    60
caaatgccat cattgcgata aaggaaaggc tatcgttcaa gatgcctctg cgcacagtgg    120
tcccaaagat ggacccccac ccacgaggag catcgtggaa aaagaagacg ttccaaccac    180
gtcttcaaag caagtggatt gatgtgatat ctccactgac gtaagggatg acgcacaatc    240
ccactatcct tcgcaagacc cttcctctat ataaggaagt tcatttcatt tggagaggac    300
aggcttcttg agatccttca acaattacca acaacaacaa acaacaaaca acattacaat    360
tactatttac aattacagtc gaggggatct atggcgcgaa aatccctaat tttcccggtg    420
atthtgctcg ccgttcttct cttctctccg cogatttact ccgccggtca cgattaccgc    480
gacgctctcc gtaaactctag catggctgat gaattggcct actccccacc gtattacca    540
tccccttggg ccaatggcca gggcgactgg gcgcaggcat accagcgcgc tgttgatatt    600
gtctcgcaaa tgacattgga tgagaagtc aatctgacca caggaactgg atgggaattg    660
gaactatgtg ttggtcagac tggcgggtgt ccccgattgg gagttccggg aatgtgttta    720
caggatagcc ctctggggct tcgcgactcc gactacaact ctgctttccc tgccggcatg    780
aacgtggctg cgacctggga caagaactct gcataccttc gcggcaaggc tatgggtcag    840
gaatttagtg acaagggtgc cgatatccaa ttgggtccag ctgccggccc tctcogtaga    900

```

-continued

agtcccagcg	gtggtcgtaa	ctgggagggc	ttctcccag	accctgccct	aagtgggtg	960
ctctttgccg	agaccatcaa	gggtatccaa	gatgctggtg	tggttgcgac	ggctaagcac	1020
tacattgctt	acgagcaaga	gcatttccgt	caggcgctg	aagcccaagg	ttttggattt	1080
aatatttccg	agagtggaag	tgcaaacctc	gatgataaga	ctatgcacga	gctgtacctc	1140
tggcccttcg	cggatgccat	ccgtgcaggt	gctggcgctg	tgatgtgctc	ctacaaccag	1200
atcaacaaca	gttatggctg	ccagaacagc	tacctctga	acaagctgct	caaggccgag	1260
ctgggcttcc	agggctttgt	catgagtgat	tgggctgctc	accatgctgg	tgtgagtggt	1320
gctttggcag	gattggatat	gtctatgcca	ggagacgtcg	actacgacag	tggtagctct	1380
tactggggta	caaaacttgac	cattagcgtg	ctcaacggaa	cggtgcccca	atggcgtggt	1440
gatgacatgg	ctgtccgcac	catggccgcc	tactacaagg	tcggccgtga	ccgtctgtgg	1500
actcctccca	acttcagctc	atggaccaga	gatgaatacg	gctacaagta	ctactacgtg	1560
tcggagggac	cgtacgagaa	ggtcaaccag	tacgtgaatg	tgcaacgcaa	ccacagcgaa	1620
ctgattcgcc	gcattggagc	ggacagcacg	gtgctcctca	agaacgacgg	cgctctgcct	1680
ttgactggta	aggagcgcct	ggtcgcgctt	atcggagaag	atgctggctc	caacccttat	1740
ggtgccaaag	gctgcagtga	ccgtggatgc	gacaatggaa	cattggcgat	gggctgggga	1800
agtggtactg	ccaacttccc	atacctggtg	acccccgagc	aggccatctc	aaacgaggtg	1860
cttaagcaca	agaatggtgt	attcaccgcc	accgataact	gggctatcga	tcaaattgag	1920
gcgcttgcta	agaccgccag	tgtctctctt	gtctttgtca	acgccgactc	tggtaggggt	1980
tacatcaatg	tggacggaaa	cctgggtgac	cgcaggaacc	tgaccctgtg	gaggaacggc	2040
gataatgtga	tcaaggctgc	tgctagcaac	tgcaacaaca	caatcgttgt	cattcactct	2100
gtcggaccag	tcttggttaa	cgagtggtag	gacaacccca	atgttacgcg	tatcctctgg	2160
ggtggtttgc	ccggtcagga	gtctggcaac	tctcttgccg	acgtcctcta	tggccgtgtc	2220
aacccccggtg	ccaagtcgcc	ctttacctgg	ggcaagactc	gtgaggccta	ccaagactac	2280
ttggtcaccg	agcccaacaa	cggcaacgga	gcccctcagg	aagactttgt	cgagggcgtc	2340
ttcattgact	accgtggatt	tgacaagcgc	aacgagaccc	cgatctaoga	gttoggctat	2400
ggtctgagct	acaccacttt	caactactcg	aaccttgagg	tgacagtgct	gagcgcccct	2460
gcatacgagc	ctgcttcggg	tgagaccgag	gcagcgccaa	ccttcggaga	ggttggaat	2520
gcgtcggatt	acctctacc	cagcggattg	ctgagaatta	ccaagttcat	ctacccctgg	2580
ctcaacggta	ccgatctcga	ggcatcttcc	gggatgcta	gctacgggca	ggactcctcc	2640
gactatcttc	ccgaggggag	caccgatgac	tctgcgcaac	cgatcctgcc	tgcgggtggc	2700
ggtcctggcg	gcaaccctcg	cctgtacgac	gagctcatcc	gcgtgtcagt	gaccatcaag	2760
aacaccggca	aggttgctgg	tgatgaagtt	ccccaaactgt	atgtttccct	tggcgggtccc	2820
aatgagccca	agatcgtgct	gcgtcaattc	gagcgcatca	cgctgcagcc	gtcggaggag	2880
acgaagtgga	gcacgactct	gacgcgccgt	gaccttgcaa	actggaatgt	tgagaagcag	2940
gactgggaga	ttacgtcgta	tcccgaagtg	gtgtttgtcg	gaagctcctc	gcggaagctg	3000
ccgctccggg	cgtctctgcc	tactgttcac	taaccggggc	gagctcgaat	tgatcgttca	3060
aacatttggc	aataaagttt	cttaagattg	aatcctgttg	ccggtcttgc	gatgattatc	3120
atataatttc	tgttgaatta	cgtaagcat	gtaataatta	aacatgtaat	gcatgacggt	3180

-continued

```

atztatgaga tgggggtttt atgattaaga gtccccgcaa ttatacattt taatacgcga 3240
tagaaaaaca aaatatagcg cccaaactaa ggataaaaatt attcgcgccg cgggggggca 3300
ttctatgggtt actagatctc tagaattcc 3329

```

```

<210> SEQ ID NO 16
<211> LENGTH: 880
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: BGL1 fused to a Cell signal peptide for
secretion into the apoplast

```

```

<400> SEQUENCE: 16

```

```

Met Ala Arg Lys Ser Leu Ile Phe Pro Val Ile Leu Leu Ala Val Leu
1 5 10 15
Leu Phe Ser Pro Pro Ile Tyr Ser Ala Gly His Asp Tyr Arg Asp Ala
20 25 30
Leu Arg Lys Ser Ser Met Ala Asp Glu Leu Ala Tyr Ser Pro Pro Tyr
35 40 45
Tyr Pro Ser Pro Trp Ala Asn Gly Gln Gly Asp Trp Ala Gln Ala Tyr
50 55 60
Gln Arg Ala Val Asp Ile Val Ser Gln Met Thr Leu Asp Glu Lys Val
65 70 75 80
Asn Leu Thr Thr Gly Thr Gly Trp Glu Leu Glu Leu Cys Val Gly Gln
85 90 95
Thr Gly Gly Val Pro Arg Leu Gly Val Pro Gly Met Cys Leu Gln Asp
100 105 110
Ser Pro Leu Gly Val Arg Asp Ser Asp Tyr Asn Ser Ala Phe Pro Ala
115 120 125
Gly Met Asn Val Ala Ala Thr Trp Asp Lys Asn Leu Ala Tyr Leu Arg
130 135 140
Gly Lys Ala Met Gly Gln Glu Phe Ser Asp Lys Gly Ala Asp Ile Gln
145 150 155 160
Leu Gly Pro Ala Ala Gly Pro Leu Gly Arg Ser Pro Asp Gly Gly Arg
165 170 175
Asn Trp Glu Gly Phe Ser Pro Asp Pro Ala Leu Ser Gly Val Leu Phe
180 185 190
Ala Glu Thr Ile Lys Gly Ile Gln Asp Ala Gly Val Val Ala Thr Ala
195 200 205
Lys His Tyr Ile Ala Tyr Glu Gln Glu His Phe Arg Gln Ala Pro Glu
210 215 220
Ala Gln Gly Phe Gly Phe Asn Ile Ser Glu Ser Gly Ser Ala Asn Leu
225 230 235 240
Asp Asp Lys Thr Met His Glu Leu Tyr Leu Trp Pro Phe Ala Asp Ala
245 250 255
Ile Arg Ala Gly Ala Gly Ala Val Met Cys Ser Tyr Asn Gln Ile Asn
260 265 270
Asn Ser Tyr Gly Cys Gln Asn Ser Tyr Thr Leu Asn Lys Leu Leu Lys
275 280 285
Ala Glu Leu Gly Phe Gln Gly Phe Val Met Ser Asp Trp Ala Ala His
290 295 300
His Ala Gly Val Ser Gly Ala Leu Ala Gly Leu Asp Met Ser Met Pro
305 310 315 320

```

-continued

Gly Asp Val Asp Tyr Asp Ser Gly Thr Ser Tyr Trp Gly Thr Asn Leu
 325 330 335

Thr Ile Ser Val Leu Asn Gly Thr Val Pro Gln Trp Arg Val Asp Asp
 340 345 350

Met Ala Val Arg Ile Met Ala Ala Tyr Tyr Lys Val Gly Arg Asp Arg
 355 360 365

Leu Trp Thr Pro Pro Asn Phe Ser Ser Trp Thr Arg Asp Glu Tyr Gly
 370 375 380

Tyr Lys Tyr Tyr Tyr Val Ser Glu Gly Pro Tyr Glu Lys Val Asn Gln
 385 390 395 400

Tyr Val Asn Val Gln Arg Asn His Ser Glu Leu Ile Arg Arg Ile Gly
 405 410 415

Ala Asp Ser Thr Val Leu Leu Lys Asn Asp Gly Ala Leu Pro Leu Thr
 420 425 430

Gly Lys Glu Arg Leu Val Ala Leu Ile Gly Glu Asp Ala Gly Ser Asn
 435 440 445

Pro Tyr Gly Ala Asn Gly Cys Ser Asp Arg Gly Cys Asp Asn Gly Thr
 450 455 460

Leu Ala Met Gly Trp Gly Ser Gly Thr Ala Asn Phe Pro Tyr Leu Val
 465 470 475 480

Thr Pro Glu Gln Ala Ile Ser Asn Glu Val Leu Lys His Lys Asn Gly
 485 490 495

Val Phe Thr Ala Thr Asp Asn Trp Ala Ile Asp Gln Ile Glu Ala Leu
 500 505 510

Ala Lys Thr Ala Ser Val Ser Leu Val Phe Val Asn Ala Asp Ser Gly
 515 520 525

Glu Gly Tyr Ile Asn Val Asp Gly Asn Leu Gly Asp Arg Arg Asn Leu
 530 535 540

Thr Leu Trp Arg Asn Gly Asp Asn Val Ile Lys Ala Ala Ala Ser Asn
 545 550 555 560

Cys Asn Asn Thr Ile Val Val Ile His Ser Val Gly Pro Val Leu Val
 565 570 575

Asn Glu Trp Tyr Asp Asn Pro Asn Val Thr Ala Ile Leu Trp Gly Gly
 580 585 590

Leu Pro Gly Gln Glu Ser Gly Asn Ser Leu Ala Asp Val Leu Tyr Gly
 595 600 605

Arg Val Asn Pro Gly Ala Lys Ser Pro Phe Thr Trp Gly Lys Thr Arg
 610 615 620

Glu Ala Tyr Gln Asp Tyr Leu Val Thr Glu Pro Asn Asn Gly Asn Gly
 625 630 635 640

Ala Pro Gln Glu Asp Phe Val Glu Gly Val Phe Ile Asp Tyr Arg Gly
 645 650 655

Phe Asp Lys Arg Asn Glu Thr Pro Ile Tyr Glu Phe Gly Tyr Gly Leu
 660 665 670

Ser Tyr Thr Thr Phe Asn Tyr Ser Asn Leu Glu Val Gln Val Leu Ser
 675 680 685

Ala Pro Ala Tyr Glu Pro Ala Ser Gly Glu Thr Glu Ala Ala Pro Thr
 690 695 700

Phe Gly Glu Val Gly Asn Ala Ser Asp Tyr Leu Tyr Pro Ser Gly Leu
 705 710 715 720

-continued

gaactatgtg ttggtcagac tggcgggtgt ccccgattgg gagttccggg aatgtgttta	720
caggatagcc ctctggcgt tcgcgactcc gactacaact ctgctttccc tgcggcatg	780
aacgtggctg cgacctggga caagaatctg gcataccttc gcggcaaggc tatgggtcag	840
gaatttagtg acaagggtgc cgatatcaa ttgggtccag ctgccggccc tctcggtaga	900
agtcccagcg gtggtcgtaa ctgggagggc ttctcccag accctgcct aagtgtgtg	960
ctctttgccc agaccatcaa gggatatcaa gatgctggg tggttgcgac ggctaagcac	1020
tacattgctt acgagcaaga gcatttccgt caggcgcctg aagcccaagg ttttgattt	1080
aatatttccg agagtgaag tgcgaacctc gatgataaga ctatgcacga gctgtacctc	1140
tggcccttcg cggatgccat ccgtgcaggt gctggcgtg tgatgtgctc ctacaaccag	1200
atcaacaaca gttatggctg ccagaacagc tactactctga acaagctgct caaggccgag	1260
ctgggcttcc agggctttgt catgagtgat tgggctgctc accatgctgg tgtgagtgt	1320
gctttggcag gattggatat gtctatgcca ggagacgtcg actacgacag tggtagctt	1380
tactgggta caaactgac cattagcgtg ctcaacggaa cggtgccca atggcgtgtt	1440
gatgacatgg ctgtccgat catggccgcc tactacaagg tcggccgtga ccgtctgtg	1500
actctccca acttcagctc atggaccaga gatgaatagc gctacaagta ctactacgtg	1560
tcggaggagc cgtacgagaa ggtcaaccag tacgtgaatg tgcaacgcaa ccacagcga	1620
ctgattcgcc gattggagc ggacagcagc gtgctcctca agaacgacgg cgtctgcct	1680
ttgactggta aggagcgcct ggtcgcgctt atcggagaag atgcgggctc caacccttat	1740
gggtccaacg gctgcagtga ccgtggatgc gacaatgaa cattggcgtg gggctgggga	1800
agtgtactg ccaacttccc atacctggg acccccagc aggccatctc aaacgaggtg	1860
cttaagcaca agaatggtgt attcaccgcc accgataact gggctatcga tcaaattgag	1920
gcgcttgcta agaccgccg tgtctctctt gtctttgtca acgccgactc tggtaggggt	1980
tacatcaatg tggacgaaa cctgggtgac cgcaggaacc tgaccctgtg gaggaacggc	2040
gataatgtga tcaaggctgc tgctagcaac tgcaacaaca caatcgttgt cattactct	2100
gtcggaccag tcttggttaa cgagtgtac gacaaccca atgttacgc tctctctgg	2160
gggtggttgc ccggtcagga gtctggcaac tctcttgcg acgtccteta tggccgtgtc	2220
aaccccgggt ccaagtcgcc ctttacctgg ggcaagactc gtgaggccta ccaagactac	2280
ttggtcaccg agcccaaaa cggcaacgga gcccctcagg aagacttgt cgaggcgtc	2340
ttcattgact accgtggatt tgacaagcgc aacgagacc cgatctacga gttcggctat	2400
ggtctgagct acaccacttt caactactcg aaccttgagg tgcagggtct gagcgcctt	2460
gcatacgagc ctgcttcggg tgagaocgag gcagcgcga ccttcggaga ggttgaaat	2520
gcgtcgatt acctctacc cagcggattg ctgagaatta ccaagttcat ctaccctgg	2580
ctcaacggta ccgatctcga ggcattctcc ggggatgcta gctacgggca ggactcctc	2640
gactatcttc ccgagggagc caccgatggc tctgcgcaac cgatcctgcc tgcogtggc	2700
ggtcctggcg gcaacctcgc cctgtacgac gagctcatcc gcgtgtcagt gaccatcaag	2760
aacaccggca aggttctgtg tgatgaagtt ccccaactgt atgtttccct tggcggctcc	2820
aatgagccca agatcgtgct gcgtcaattc gagcgcacga cgtgcagcc gtcggaggag	2880
acgaagtga gcacgactct gacgcgccgt gacctgcaa actggaatgt tgagaagcag	2940

-continued

```

gactgggaga ttacgtcgta tcccaagatg gtgtttgtcg gaagctcctc gcggaagctg 3000
ccgctccggg cgtctctgcc tactgttcat gatgaacttt aaccggggcg agctcgaatt 3060
gatcgttcaa acatttggca ataaagtttc ttaagattga gttaagcatg taataattaa 3120
acatgtaatg catgacgtta tttatgagat ggggttttta tgattaagag tccccgcaat 3180
tatacathtt aatacgcgat agaaaaacaa aatatagcgc ccaaactaag gataaaatta 3240
ttcgcgccgc gggggggcat tctatgggta ctgatctct agaattcc 3288

```

<210> SEQ ID NO 19

<211> LENGTH: 883

<212> TYPE: PRT

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: BGL1 fused to Cell signal peptide and a HDEL
ER-retaining peptide

<400> SEQUENCE: 19

```

Met Ala Arg Lys Ser Leu Ile Phe Pro Val Ile Leu Leu Ala Val Leu
1           5           10          15
Leu Phe Ser Pro Pro Ile Tyr Ser Ala Gly His Asp Tyr Arg Asp Ala
20          25          30
Leu Arg Lys Ser Ser Met Ala Asp Glu Leu Ala Tyr Ser Pro Pro Tyr
35          40          45
Tyr Pro Ser Pro Trp Ala Asn Gly Gln Gly Asp Trp Ala Gln Ala Tyr
50          55          60
Gln Arg Ala Val Asp Ile Val Ser Gln Met Thr Leu Asp Glu Lys Val
65          70          75          80
Asn Leu Thr Thr Gly Thr Gly Trp Glu Leu Glu Leu Cys Val Gly Gln
85          90          95
Thr Gly Gly Val Pro Arg Leu Gly Val Pro Gly Met Cys Leu Gln Asp
100         105        110
Ser Pro Leu Gly Val Arg Asp Ser Asp Tyr Asn Ser Ala Phe Pro Ala
115        120        125
Gly Met Asn Val Ala Ala Thr Trp Asp Lys Asn Leu Ala Tyr Leu Arg
130        135        140
Gly Lys Ala Met Gly Gln Glu Phe Ser Asp Lys Gly Ala Asp Ile Gln
145        150        155        160
Leu Gly Pro Ala Ala Gly Pro Leu Gly Arg Ser Pro Asp Gly Gly Arg
165        170        175
Asn Trp Glu Gly Phe Ser Pro Asp Pro Ala Leu Ser Gly Val Leu Phe
180        185        190
Ala Glu Thr Ile Lys Gly Ile Gln Asp Ala Gly Val Val Ala Thr Ala
195        200        205
Lys His Tyr Ile Ala Tyr Glu Gln Glu His Phe Arg Gln Ala Pro Glu
210        215        220
Ala Gln Gly Phe Gly Phe Asn Ile Ser Glu Ser Gly Ser Ala Asn Leu
225        230        235        240
Asp Asp Lys Thr Met His Glu Leu Tyr Leu Trp Pro Phe Ala Asp Ala
245        250        255
Ile Arg Ala Gly Ala Gly Ala Val Met Cys Ser Tyr Asn Gln Ile Asn
260        265        270
Asn Ser Tyr Gly Cys Gln Asn Ser Tyr Thr Leu Asn Lys Leu Leu Lys
275        280        285

```

-continued

Ala Glu Leu Gly Phe Gln Gly Phe Val Met Ser Asp Trp Ala Ala His
290 295 300

His Ala Gly Val Ser Gly Ala Leu Ala Gly Leu Asp Met Ser Met Pro
305 310 315 320

Gly Asp Val Asp Tyr Asp Ser Gly Thr Ser Tyr Trp Gly Thr Asn Leu
325 330 335

Thr Ile Ser Val Leu Asn Gly Thr Val Pro Gln Trp Arg Val Asp Asp
340 345 350

Met Ala Val Arg Ile Met Ala Ala Tyr Tyr Lys Val Gly Arg Asp Arg
355 360 365

Leu Trp Thr Pro Pro Asn Phe Ser Ser Trp Thr Arg Asp Glu Tyr Gly
370 375 380

Tyr Lys Tyr Tyr Tyr Val Ser Glu Gly Pro Tyr Glu Lys Val Asn Gln
385 390 395 400

Tyr Val Asn Val Gln Arg Asn His Ser Glu Leu Ile Arg Arg Ile Gly
405 410 415

Ala Asp Ser Thr Val Leu Leu Lys Asn Asp Gly Ala Leu Pro Leu Thr
420 425 430

Gly Lys Glu Arg Leu Val Ala Leu Ile Gly Glu Asp Ala Gly Ser Asn
435 440 445

Pro Tyr Gly Ala Asn Gly Cys Ser Asp Arg Gly Cys Asp Asn Gly Thr
450 455 460

Leu Ala Met Gly Trp Gly Ser Gly Thr Ala Asn Phe Pro Tyr Leu Val
465 470 475 480

Thr Pro Glu Gln Ala Ile Ser Asn Glu Val Leu Lys His Lys Asn Gly
485 490 495

Val Phe Thr Ala Thr Asp Asn Trp Ala Ile Asp Gln Ile Glu Ala Leu
500 505 510

Ala Lys Thr Ala Ser Val Ser Leu Val Phe Val Asn Ala Asp Ser Gly
515 520 525

Glu Gly Tyr Ile Asn Val Asp Gly Asn Leu Gly Asp Arg Arg Asn Leu
530 535 540

Thr Leu Trp Arg Asn Gly Asp Asn Val Ile Lys Ala Ala Ala Ser Asn
545 550 555 560

Cys Asn Asn Thr Ile Val Val Ile His Ser Val Gly Pro Val Leu Val
565 570 575

Asn Glu Trp Tyr Asp Asn Pro Asn Val Thr Ala Ile Leu Trp Gly Gly
580 585 590

Leu Pro Gly Gln Glu Ser Gly Asn Ser Leu Ala Asp Val Leu Tyr Gly
595 600 605

Arg Val Asn Pro Gly Ala Lys Ser Pro Phe Thr Trp Gly Lys Thr Arg
610 615 620

Glu Ala Tyr Gln Asp Tyr Leu Val Thr Glu Pro Asn Asn Gly Asn Gly
625 630 635 640

Ala Pro Gln Glu Asp Phe Val Glu Gly Val Phe Ile Asp Tyr Arg Gly
645 650 655

Phe Asp Lys Arg Asn Glu Thr Pro Ile Tyr Glu Phe Gly Tyr Gly Leu
660 665 670

Ser Tyr Thr Thr Phe Asn Tyr Ser Asn Leu Glu Val Gln Val Leu Ser
675 680 685

-continued

Ala Pro Ala Tyr Glu Pro Ala Ser Gly Glu Thr Glu Ala Ala Pro Thr
690 695 700

Phe Gly Glu Val Gly Asn Ala Ser Asp Tyr Leu Tyr Pro Ser Gly Leu
705 710 715 720

Leu Arg Ile Thr Lys Phe Ile Tyr Pro Trp Leu Asn Gly Thr Asp Leu
725 730 735

Glu Ala Ser Ser Gly Asp Ala Ser Tyr Gly Gln Asp Ser Ser Asp Tyr
740 745 750

Leu Pro Glu Gly Ala Thr Asp Gly Ser Ala Gln Pro Ile Leu Pro Ala
755 760 765

Gly Gly Gly Pro Gly Gly Asn Pro Arg Leu Tyr Asp Glu Leu Ile Arg
770 775 780

Val Ser Val Thr Ile Lys Asn Thr Gly Lys Val Ala Gly Asp Glu Val
785 790 795 800

Pro Gln Leu Tyr Val Ser Leu Gly Gly Pro Asn Glu Pro Lys Ile Val
805 810 815

Leu Arg Gln Phe Glu Arg Ile Thr Leu Gln Pro Ser Glu Glu Thr Lys
820 825 830

Trp Ser Thr Thr Leu Thr Arg Arg Asp Leu Ala Asn Trp Asn Val Glu
835 840 845

Lys Gln Asp Trp Glu Ile Thr Ser Tyr Pro Lys Met Val Phe Val Gly
850 855 860

Ser Ser Ser Arg Lys Leu Pro Leu Arg Ala Ser Leu Pro Thr Val His
865 870 875 880

Asp Glu Leu

<210> SEQ ID NO 20
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Single strand DNA oligonucleotide

<400> SEQUENCE: 20

cagtgaccgt ggatgcgaca atg

23

<210> SEQ ID NO 21
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Single strand DNA oligonucleotide

<400> SEQUENCE: 21

agagacggat gacaagtact acttgaaatt gggcccaaaa

40

<210> SEQ ID NO 22
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Single strand DNA oligonucleotide

<400> SEQUENCE: 22

cagtgaccgt ggatgcgaca atg

23

<210> SEQ ID NO 23

-continued

```

<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Single strand DNA oligonucleotide

```

```

<400> SEQUENCE: 23

```

```

aaaggatcct tagtgaacag taggcagaga cgc

```

33

```

<210> SEQ ID NO 24
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: ER retaining signal peptide

```

```

<400> SEQUENCE: 24

```

```

Lys Asp Glu Leu

```

```

1

```

What is claimed is:

1. A method of increasing a level of at least one fermentation substance in a fermentation product, the method comprising the step of fermenting a plant derived glucose containing fermentation starting material by a yeast cell, said plant expressing a nucleic acid construct comprising a polynucleotide encoding a polypeptide having a β -glucosidase catalytic activity and further encoding a signal peptide being in frame with said polypeptide, said plant having greater β -glucosidase catalytic activity as compared to β -glucosidase catalytic activity of a plant not expressing said nucleic acid construct, thereby increasing the level of the at least one fermentation substance in the fermentation product.

2. The method of claim 1, wherein said polynucleotide further encodes an endoplasmic reticulum retaining peptide being in frame with said polypeptide.

3. The method of claim 1, wherein said signal peptide is an apoplast and/or vacuole targeting signal peptide.

4. The method of claim 1, wherein said signal peptide is Cell1.

5. A method of increasing a level of at least one aroma substance in a plant derived product, the method comprising the step of incubating a glucose containing plant starting material with a yeast cell, said plant expressing a nucleic acid construct comprising a polynucleotide encoding a polypeptide having a β -glucosidase catalytic activity and further encoding a signal peptide being in frame with said polypeptide, said plant having greater β -glucosidase catalytic activity as compared to β -glucosidase catalytic activity of a plant not expressing said nucleic acid construct, thereby increasing the level of the at least one aroma substance in the plant derived product.

6. The method of claim 5, wherein said polynucleotide further encodes an endoplasmic reticulum retaining peptide being in frame with said polypeptide.

7. The method of claim 5, wherein said signal peptide is an apoplast and/or vacuole targeting signal peptide.

8. The method of claim 5, wherein said signal peptide is Cell1.

9. The method of claim 5, wherein said plant derived product is a fermentation product.

10. A method of increasing a level of free glucose in a glucose containing fermentation starting material, the method comprising the step of fermenting the glucose containing fermentation starting material by a cell expressing a nucleic acid construct comprising a polynucleotide encoding a polypeptide having a β -glucosidase catalytic activity and further encoding a signal peptide being in frame with said polypeptide, said cell having greater β -glucosidase catalytic activity as compared to β -glucosidase catalytic activity of a cell not expressing said nucleic acid construct, thereby increasing the level of the free glucose in the glucose containing fermentation starting material.

11. The method of claim 10, wherein said polynucleotide further encodes an endoplasmic reticulum retaining peptide being in frame with said polypeptide.

12. The method of claim 10, wherein said signal peptide is an apoplast and/or vacuole targeting signal peptide.

13. The method of claim 10, wherein said signal peptide is Cell1.

14. A method of increasing a level of free glucose in a plant derived glucose containing fermentation starting material, the method comprising the step of fermenting the plant derived glucose containing fermentation starting material by a cell, said plant expressing a nucleic acid construct comprising a polynucleotide encoding a polypeptide having a β -glucosidase catalytic activity and further encoding a signal peptide being in frame with said polypeptide, said plant having greater β -glucosidase catalytic activity as compared to β -glucosidase catalytic activity of a plant not expressing said nucleic acid construct, thereby increasing the level of the free glucose in the plant.

15. The method of claim 14, wherein said polynucleotide further encodes an endoplasmic reticulum retaining peptide being in frame with said polypeptide.

16. The method of claim 14, wherein said signal peptide is an apoplast and/or vacuole targeting signal peptide.

17. The method of claim 14, wherein said signal peptide is Cell1.

18. A method of producing an alcohol, the method comprising the step of fermenting a glucose containing fermentation starting material by a cell expressing a nucleic acid construct comprising a polynucleotide encoding a polypeptide having a β -glucosidase catalytic activity and further encoding a signal peptide being in frame with said polypeptide, said cell having greater β -glucosidase catalytic activity as compared to β -glucosidase catalytic activity of a cell not expressing said nucleic acid construct and extracting the alcohol therefrom.

19. The method of claim 18, wherein said polynucleotide further encodes an endoplasmic reticulum retaining peptide being in frame with said polypeptide.

20. The method of claim 18, wherein said signal peptide is an apoplast and/or vacuole targeting signal peptide.

21. The method of claim 18, wherein said signal peptide is Cell1.

22. A method of producing an alcohol, the method comprising the step of fermenting a plant derived glucose containing fermentation starting material by a cell, said plant expressing a nucleic acid construct comprising a polynucleotide encoding a polypeptide having a β -glucosidase catalytic activity and further encoding a signal peptide being in frame with said polypeptide, said plant having greater β -glucosidase catalytic activity as compared to β -glucosidase catalytic activity of a plant not expressing said nucleic acid construct, and extracting the alcohol therefrom.

23. The method of claim 22, wherein said polynucleotide further encodes an endoplasmic reticulum retaining peptide being in frame with said polypeptide.

24. The method of claim 22, wherein said signal peptide is an apoplast and/or vacuole targeting signal peptide.

25. The method of claim 22, wherein said signal peptide is Cell1.

26. A method of producing a plant having increased release of flavor and/or aroma compounds in-vivo, the method comprising the step of expressing in the plant a nucleic acid construct comprising a polynucleotide encoding

a polypeptide having a β -glucosidase catalytic activity and further encoding an apoplast and/or vacuole targeting signal peptide being in frame with said polypeptide and wherein said polypeptide is secreted into the apoplast and/or vacuole, said plant having greater β -glucosidase catalytic activity in the apoplast and/or vacuole as compared to β -glucosidase catalytic activity of the apoplast and/or vacuole of a plant not expressing said nucleic acid construct, thereby increasing in-vivo release of flavor and/or aroma compounds from the plant.

27. The method of claim 26, wherein said signal peptide is Cell1.

28. A method of producing a plant having increased release of flavor and/or aroma compounds upon processing of the plant or portion thereof, the method comprising the step of expressing in the plant a nucleic acid construct comprising a polynucleotide encoding a polypeptide having a β -glucosidase catalytic activity and further encoding a signal peptide and an endoplasmic retention peptide being in frame with said polypeptide and wherein said polypeptide accumulates in the endoplasmic reticulum, said plant having greater β -glucosidase catalytic activity in said endoplasmic reticulum as compared to β -glucosidase catalytic activity of said endoplasmic reticulum of a plant not expressing said nucleic acid construct, thereby increasing release of flavor and/or aroma compounds from the processing of said plant or portion thereof, and wherein said processing comprises cell disruption and decompartmentalization.

29. The method of claim 28, wherein said signal peptide is an apoplast and/or vacuole targeting signal peptide.

30. The method of claim 28, wherein said signal peptide is Cell1.

31. The method of claim 28, wherein said endoplasmic retention peptide is selected from the group consisting of KDEL and HDEL.

* * * * *