




(86) Date de dépôt PCT/PCT Filing Date: 2014/12/08
(87) Date publication PCT/PCT Publication Date: 2015/06/18
(45) Date de délivrance/Issue Date: 2022/03/29
(85) Entrée phase nationale/National Entry: 2016/05/11
(86) N° demande PCT/PCT Application No.: EP 2014/076852
(87) N° publication PCT/PCT Publication No.: 2015/086511
(30) Priorité/Priority: 2013/12/09 (GB1321745.0)

(51) Cl.Int./Int.Cl. *C07D 487/04* (2006.01),
A61K 31/5025 (2006.01), *A61P 29/00* (2006.01)
(72) Inventeur/Inventor:
BROOKINGS, DANIEL CHRISTOPHER, GB
(73) Propriétaire/Owner:
UCB BIOPHARMA SRL, BE
(74) Agent: SMART & BIGGAR LLP

(54) Titre : DERIVES DE TRIAZOLOPYRIDAZINE A TITRE DE MODULATEURS DE L'ACTIVITE DU TNF
(54) Title: TRIAZOLOPYRIDAZINE DERIVATIVES AS MODULATORS OF TNF ACTIVITY



(I)

(57) Abrégé/Abstract:

A series of substituted [1,2,4]triazolo[4,3-b]pyridazine derivatives of formula (I), being potent modulators of human TNF α activity, are accordingly of benefit in the treatment and/or prevention of various human ailments, including autoimmune and inflammatory disorders; neurological and neurodegenerative disorders; pain and nociceptive disorders; cardiovascular disorders; metabolic disorders; ocular disorders; and oncological disorders.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(10) International Publication Number
WO 2015/086511 A1(43) International Publication Date
18 June 2015 (18.06.2015)

(51) International Patent Classification:

C07D 487/04 (2006.01) A61P 29/00 (2006.01)
A61K 31/5025 (2006.01)

(21) International Application Number:

PCT/EP2014/076852

(22) International Filing Date:

8 December 2014 (08.12.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:


1321745.0 9 December 2013 (09.12.2013) GB

(71) Applicant: UCB BIOPHARMA SPRL [BE/BE]; Allée
de la Recherche 60, B-1070 Brussels (BE).(72) Inventor: BROOKINGS, Daniel Christopher; c/o IPD,
UCB Celltech, 208 Bath Road, Slough Berkshire SL1 3WE
(GB).(74) Agent: DUCKWORTH, Timothy John; JA KEMP, 14
South Square, Gray's Inn, London, Greater London WC1R
5JJ (GB).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LI, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: TRIAZOLOPYRIDAZINE DERIVATIVES AS MODULATORS OF TNF ACTIVITY



(I)

(57) Abstract: A series of substituted
[1,2,4]triazolo[4,3-b]pyridazine derivatives of formula
(I), being potent modulators of human TNF α activity,
are accordingly of benefit in the treatment and/or pre-
vention of various human ailments, including autoim-
mune and inflammatory disorders; neurological and
neurodegenerative disorders; pain and nociceptive dis-
orders; cardiovascular disorders; metabolic disorders;
ocular disorders; and oncological disorders.

TRIAZOLOPYRIDAZINE DERIVATIVES AS MODULATORS OF TNF ACTIVITY

The present invention relates to a class of fused triazole derivatives, and to their
5 use in therapy. More particularly, this invention is concerned with pharmacologically
active substituted [1,2,4]triazolo[4,3-*b*]pyridazine derivatives. These compounds are
modulators of the signalling of TNF α , and are accordingly of benefit as pharmaceutical
agents, especially in the treatment of adverse inflammatory and autoimmune disorders,
neurological and neurodegenerative disorders, pain and nociceptive disorders,
10 cardiovascular disorders, metabolic disorders, ocular disorders, and oncological disorders.

TNF α is the prototypical member of the Tumour Necrosis Factor (TNF)
superfamily of proteins that share a primary function of regulating cell survival and cell
death. One structural feature common to all known members of the TNF superfamily is
the formation of trimeric complexes that bind to, and activate, specific TNF superfamily
15 receptors. By way of example, TNF α exists in soluble and transmembrane forms and
signals through two receptors, known as TNFR1 and TNFR2, with distinct functional
endpoints.

Various products capable of modulating TNF α activity are already commercially
available. All are approved for the treatment of inflammatory and autoimmune disorders
20 such as rheumatoid arthritis and Crohn's disease. All currently approved products are
macromolecular and act by inhibiting the binding of human TNF α to its receptor. Typical
macromolecular TNF α inhibitors include anti-TNF α antibodies; and soluble TNF α
receptor fusion proteins. Examples of commercially available anti-TNF α antibodies
include fully human antibodies such as adalimumab (Humira®) and golimumab
25 (Simponi®), chimeric antibodies such as infliximab (Remicade®), and pegylated Fab'
fragments such as certolizumab pegol (Cimzia®). An example of a commercially
available soluble TNF α receptor fusion protein is etanercept (Enbrel®).

TNF superfamily members, including TNF α itself, are implicated in a variety of
physiological and pathological functions that are believed to play a part in a range of
30 conditions of significant medical importance (see, for example, M.G. Tansey & D.E.
Szymkowski, *Drug Discovery Today*, 2009, **14**, 1082-1088; and F.S. Carneiro *et al.*, *J.*
Sexual Medicine, 2010, **7**, 3823-3834).

The compounds in accordance with the present invention, being potent modulators of human TNF α activity, are therefore beneficial in the treatment and/or prevention of various human ailments. These include autoimmune and inflammatory disorders; neurological and neurodegenerative disorders; pain and nociceptive disorders; cardiovascular disorders; metabolic disorders; ocular disorders; and oncological disorders.

In addition, the compounds in accordance with the present invention may be beneficial as pharmacological standards for use in the development of new biological tests and in the search for new pharmacological agents. Thus, in one embodiment, the compounds of this invention may be useful as radioligands in assays for detecting pharmacologically active compounds. In an alternative embodiment, certain compounds of this invention may be useful for coupling to a fluorophore to provide fluorescent conjugates that can be utilised in assays (e.g. a fluorescence polarisation assay) for detecting pharmacologically active compounds.

Co-pending international patent applications WO 2013/186229 (published 19 December 2013), WO 2014/009295 (published 16 January 2014) and WO 2014/009296 (also published 16 January 2014) describe fused imidazole derivatives which are modulators of human TNF α activity.


None of the prior art available to date, however, discloses or suggests the precise structural class of triazolopyridazine derivatives as provided by the present invention.

The compounds in accordance with the present invention potentially inhibit the binding of a fluorescence conjugate to TNF α when tested in the fluorescence polarisation assay described herein. Indeed, when tested in that assay, the compounds of the present invention exhibit an IC₅₀ value of 50 μ M or less, generally of 20 μ M or less, usually of 5 μ M or less, typically of 1 μ M or less, suitably of 500 nM or less, ideally of 100 nM or less, and preferably of 20 nM or less (the skilled person will appreciate that a *lower* IC₅₀ figure denotes a *more active* compound).

Certain compounds in accordance with the present invention potentially neutralise the activity of TNF α in a commercially available HEK-293 derived reporter cell line known as HEK-Blue™ CD40L. This is a stable HEK-293 transfected cell line expressing SEAP (secreted embryonic alkaline phosphatase) under the control of the IFN β minimal promoter fused to five NF- κ B binding sites. Secretion of SEAP by these cells is stimulated in a concentration-dependent manner by TNF α . When tested in the HEK-293 bioassay, also referred to herein as the reporter gene assay, certain compounds of the

present invention exhibit an IC_{50} value of 50 μM or less, generally of 20 μM or less, usually of 5 μM or less, typically of 1 μM or less, suitably of 500 nM or less, ideally of 100 nM or less, and preferably of 20 nM or less (as before, the skilled person will appreciate that a *lower* IC_{50} figure denotes a *more active* compound).

- 5 The present invention provides a compound of formula (I) or an *N*-oxide thereof, or a pharmaceutically acceptable salt or solvate thereof, or a glucuronide derivative thereof, or a co-crystal thereof:



(I)

10

wherein

E represents a covalent bond; or E represents -O-, -S-, -S(O)-, -S(O)₂- or -N(R⁴)-; or E represents an optionally substituted straight or branched C₁₋₄ alkylene chain;

- 15 Y represents C₃₋₇ cycloalkyl, aryl, C₃₋₇ heterocycloalkyl or heteroaryl, any of which groups may be optionally substituted by one or more substituents;

- R¹, R² and R³ independently represent hydrogen, halogen, cyano, nitro, hydroxy, trifluoromethyl, trifluoromethoxy, -OR^a, -SR^a, -SOR^a, -SO₂R^a, -SF₅, -NR^bR^c, -NR^cCOR^d, -NR^cCO₂R^d, -NHCONR^bR^c, -NR^cSO₂R^c, -N(SO₂R^c)₂, -NHSO₂NR^bR^c, -COR^d, -CO₂R^d, -CONR^bR^c, -CON(OR^a)R^b, -SO₂NR^bR^c or -SO(NR^b)R^d; or C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₇ cycloalkyl, C₄₋₇ cycloalkenyl, C₃₋₇ cycloalkyl(C₁₋₆)alkyl, aryl, aryl(C₁₋₆)-alkyl, C₃₋₇ heterocycloalkyl, C₃₋₇ heterocycloalkyl(C₁₋₆)alkyl, C₃₋₇ heterocycloalkenyl, C₄₋₉ heterobicycloalkyl, heteroaryl, heteroaryl(C₁₋₆)alkyl, (C₃₋₇)heterocycloalkyl(C₁₋₆)alkyl-aryl-, heteroaryl(C₃₋₇)heterocycloalkyl-, (C₃₋₇)cycloalkyl-heteroaryl-, (C₃₋₇)cycloalkyl-(C₁₋₆)alkyl-heteroaryl-, (C₄₋₇)cycloalkenyl-heteroaryl-, (C₄₋₉)bicycloalkyl-heteroaryl-, (C₃₋₇)heterocycloalkyl-heteroaryl-, (C₃₋₇)heterocycloalkyl(C₁₋₆)alkyl-heteroaryl-, (C₃₋₇)heterocycloalkenyl-heteroaryl-, (C₄₋₉)heterobicycloalkyl-heteroaryl- or (C₄₋₉)spiroheterocycloalkyl-heteroaryl-, any of which groups may be optionally substituted by one or more substituents;
- 20
- 25

R^d represents hydrogen or C₁₋₆ alkyl;

R^a represents C₁₋₆ alkyl, aryl, aryl(C₁₋₆)alkyl, heteroaryl or heteroaryl(C₁₋₆)alkyl, any of which groups may be optionally substituted by one or more substituents;

R^b and R^c independently represent hydrogen or trifluoromethyl; or C₁₋₆ alkyl, C₃₋₇ cycloalkyl, C₃₋₇ cycloalkyl(C₁₋₆)alkyl, aryl, aryl(C₁₋₆)alkyl, C₃₋₇ heterocycloalkyl, C₃₋₇ heterocycloalkyl(C₁₋₆)alkyl, heteroaryl or heteroaryl(C₁₋₆)alkyl, any of which groups may be optionally substituted by one or more substituents; or

R^b and R^c, when taken together with the nitrogen atom to which they are both attached, represent azetidin-1-yl, pyrrolidin-1-yl, oxazolidin-3-yl, isoxazolidin-2-yl, thiazolidin-3-yl, isothiazolidin-2-yl, piperidin-1-yl, morpholin-4-yl, thiomorpholin-4-yl, piperazin-1-yl, homopiperidin-1-yl, homomorpholin-4-yl or homopiperazin-1-yl, any of which groups may be optionally substituted by one or more substituents;

R^d represents hydrogen; or C₁₋₆ alkyl, C₃₋₇ cycloalkyl, aryl, C₃₋₇ heterocycloalkyl or heteroaryl, any of which groups may be optionally substituted by one or more substituents; and

R^e represents C₁₋₆ alkyl, aryl or heteroaryl, any of which groups may be optionally substituted by one or more substituents.

The present invention also provides a compound of formula (I) as defined above or an *N*-oxide thereof, or a pharmaceutically acceptable salt or solvate thereof, or a glucuronide derivative thereof, or a co-crystal thereof, for use in therapy.


The present invention also provides a compound of formula (I) as defined above or an *N*-oxide thereof, or a pharmaceutically acceptable salt or solvate thereof, or a glucuronide derivative thereof, or a co-crystal thereof, for use in the treatment and/or prevention of disorders for which the administration of a modulator of TNF α function is indicated.

In another aspect, the present invention provides a compound of formula (I) as defined above or an *N*-oxide thereof, or a pharmaceutically acceptable salt or solvate thereof, or a glucuronide derivative thereof, or a co-crystal thereof, for use in the treatment and/or prevention of an inflammatory or autoimmune disorder, a neurological or neurodegenerative disorder, pain or a nociceptive disorder, a cardiovascular disorder, a metabolic disorder, an ocular disorder, or an oncological disorder.

The present invention also provides a method for the treatment and/or prevention of disorders for which the administration of a modulator of TNF α function is indicated

which comprises administering to a patient in need of such treatment an effective amount of a compound of formula (I) as defined above or an *N*-oxide thereof, or a pharmaceutically acceptable salt or solvate thereof, or a glucuronide derivative thereof, or a co-crystal thereof.

- 5 In some embodiments, the compound of formula (I) is a compound of formula (IIB) or an *N*-oxide thereof, or a pharmaceutically acceptable salt or solvate thereof:



(IIB)

- 10 wherein

E represents -CH₂-;

V represents C-R²² or N;

R¹² represents hydrogen or halogen;

R¹⁵ represents difluoromethoxy;

- 15 R¹⁶ represents hydrogen, fluoro, chloro, cyano, methyl, trifluoromethyl, difluoromethoxy or amino;

R²¹ represents hydroxy(C₁₋₆)alkyl; or R²¹ represents (C₃₋₇)heterocycloalkyl, which groups may be optionally substituted by one, two or three substituents independently selected from carboxy;

- 20 R²² represents hydrogen, halogen or C₁₋₆ alkyl; and

R²³ represents hydrogen, C₁₋₆ alkyl, trifluoromethyl or C₁₋₆ alkoxy.

In another aspect, the present invention provides a method for the treatment and/or prevention of an inflammatory or autoimmune disorder, a neurological or neuro-degenerative disorder, pain or a nociceptive disorder, a cardiovascular disorder, a metabolic disorder, an ocular disorder, or an oncological disorder, which comprises administering to a
5 patient in need of such treatment an effective amount of a compound of formula (I) as defined above or an *N*-oxide thereof, or a pharmaceutically acceptable salt or solvate thereof, or a glucuronide derivative thereof, or a co-crystal thereof.

Where any of the groups in the compounds of formula (I) above is stated to be optionally substituted, this group may be unsubstituted, or substituted by one or more
10 substituents. Typically, such groups will be unsubstituted, or substituted by one or two substituents.

For use in medicine, the salts of the compounds of formula (I) will be pharmaceutically acceptable salts. Other salts may, however, be useful in the preparation of the compounds of use in the invention or of their pharmaceutically acceptable salts.
15 Standard principles underlying the selection and preparation of pharmaceutically acceptable salts are described, for example, in *Handbook of Pharmaceutical Salts: Properties, Selection and Use*, ed. P.H. Stahl & C.G. Wermuth, Wiley-VCH, 2002. Suitable pharmaceutically acceptable salts of the compounds of use in this invention include acid addition salts which may, for example, be formed by mixing a solution of the compound of
20 use in the invention with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid, methanesulphonic acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, citric acid, tartaric acid or phosphoric acid. Furthermore, where the compounds of use in the invention carry an acidic moiety, e.g. carboxy, suitable pharmaceutically acceptable salts thereof may include alkali metal salts,
25 e.g. sodium or potassium salts; alkaline earth metal salts, e.g. calcium or magnesium salts; ammonium salts; and salts formed with suitable organic ligands, e.g. quaternary ammonium salts, and meglumine salts.

The present invention includes within its scope solvates of the compounds of formula (I) above. Such solvates may be formed with common organic solvents, e.g.

hydrocarbon solvents such as benzene or toluene; chlorinated solvents such as chloroform or dichloromethane; alcoholic solvents such as methanol, ethanol or isopropanol; ethereal solvents such as diethyl ether or tetrahydrofuran; or ester solvents such as ethyl acetate. Alternatively, the solvates of the compounds of formula (I) may be formed with water, in which case they will be hydrates.

The present invention also includes co-crystals within its scope. The technical term “co-crystal” is used to describe the situation where neutral molecular components are present within a crystalline compound in a definite stoichiometric ratio. The preparation of pharmaceutical co-crystals enables modifications to be made to the crystalline form of an active pharmaceutical ingredient, which in turn can alter its physicochemical properties without compromising its intended biological activity (see *Pharmaceutical Salts and Co-crystals*, ed. J. Wouters & L. Quere, RSC Publishing, 2012). Typical examples of co-crystal formers, which may be present in the co-crystal alongside the active pharmaceutical ingredient, include *L*-ascorbic acid, citric acid, glutaric acid, urea and nicotinamide.

The present invention includes within its scope prodrugs of the compounds of formula (I) above. In general, such prodrugs will be functional derivatives of the compounds of formula (I) which are readily convertible *in vivo* into the required compound of formula (I). Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in *Design of Prodrugs*, ed. H. Bundgaard, Elsevier, 1985.

Suitable alkyl groups which may be present on the compounds of use in the invention include straight-chained and branched C₁₋₆ alkyl groups, for example C₁₋₄ alkyl groups. Typical examples include methyl and ethyl groups, and straight-chained or branched propyl, butyl and pentyl groups. Particular alkyl groups include methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *sec*-butyl, isobutyl, *tert*-butyl, 2,2-dimethylpropyl and 3-methylbutyl. Derived expressions such as “C₁₋₆ alkoxy”, “C₁₋₆ alkylthio”, “C₁₋₆ alkylsulphonyl” and “C₁₋₆ alkylamino” are to be construed accordingly.

The expression “C₁₋₄ alkylene chain” refers to a divalent straight or branched alkylene chain containing 1 to 4 carbon atoms. Typical examples include methylene, ethylene, methylenemethylene, ethylenemethylene and dimethylenemethylene.

Suitable C₂₋₆ alkenyl groups include vinyl and allyl.

Suitable C₂₋₆ alkynyl groups include ethynyl, propargyl and butynyl.

The term “C₃₋₇ cycloalkyl” as used herein refers to monovalent groups of 3 to 7 carbon atoms derived from a saturated monocyclic hydrocarbon, and may comprise benzo-fused analogues thereof. Suitable C₃₋₇ cycloalkyl groups include cyclopropyl, cyclobutyl, benzocyclobutenyl, cyclopentyl, indanyl, cyclohexyl and cycloheptyl.

5 The term “C₄₋₇ cycloalkenyl” as used herein refers to monovalent groups of 4 to 7 carbon atoms derived from a partially unsaturated monocyclic hydrocarbon. Suitable C₄₋₇ cycloalkenyl groups include cyclobutenyl, cyclopentenyl, cyclohexenyl and cycloheptenyl.

 The term “C₄₋₉ bicycloalkyl” as used herein refers to monovalent groups of 4 to 9 carbon atoms derived from a saturated bicyclic hydrocarbon. Typical bicycloalkyl groups
10 include bicyclo[3.1.0]hexanyl, bicyclo[4.1.0]heptanyl and bicyclo[2.2.2]octanyl.

 The term “aryl” as used herein refers to monovalent carbocyclic aromatic groups derived from a single aromatic ring or multiple condensed aromatic rings. Suitable aryl groups include phenyl and naphthyl, preferably phenyl.

 Suitable aryl(C₁₋₆)alkyl groups include benzyl, phenylethyl, phenylpropyl and
15 naphthylmethyl.

 The term “C₃₋₇ heterocycloalkyl” as used herein refers to saturated monocyclic rings containing 3 to 7 carbon atoms and at least one heteroatom selected from oxygen, sulphur and nitrogen, and may comprise benzo-fused analogues thereof. Suitable heterocycloalkyl groups include oxetanyl, azetidiny, tetrahydrofuranyl, dihydrobenzo-
20 furanyl, dihydrobenzothienyl, pyrrolidinyl, indolinyl, isoindolinyl, oxazolidinyl, thiazolidinyl, isothiazolidinyl, imidazolidinyl, tetrahydropyranyl, chromanyl, tetrahydrothiopyranyl, piperidinyl, 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl, piperazinyl, 1,2,3,4-tetrahydroquinoxaliny, hexahydro-[1,2,5]thiadiazolo[2,3-*a*]pyrazinyl, homopiperazinyl, morpholinyl, benzoxazinyl, thiomorpholinyl, azepanyl, oxazepanyl,
25 diazepanyl, thiadiazepanyl and azocanyl.

 The term “C₃₋₇ heterocycloalkenyl” as used herein refers to monounsaturated or polyunsaturated monocyclic rings containing 3 to 7 carbon atoms and at least one heteroatom selected from oxygen, sulphur and nitrogen, and may comprise benzo-fused analogues thereof. Suitable heterocycloalkenyl groups include thiazolinyl, isothiazolinyl,
30 imidazolinyl, dihydropyranyl, dihydrothiopyranyl and 1,2,3,6-tetrahydropyridinyl.

 The term “C₄₋₉ heterobicycloalkyl” as used herein corresponds to C₄₋₉ bicycloalkyl wherein one or more of the carbon atoms have been replaced by one or more heteroatoms selected from oxygen, sulphur and nitrogen. Typical heterobicycloalkyl groups include 3-

azabicyclo[3.1.0]hexanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 6-azabicyclo[3.2.0]heptanyl, 3-azabicyclo[3.1.1]heptanyl, 3-azabicyclo[4.1.0]heptanyl, 2-oxabicyclo[2.2.2]octanyl, quinuclidinyl, 2-oxa-5-azabicyclo[2.2.2]octanyl, 3-azabicyclo[3.2.1]octanyl, 8-azabicyclo[3.2.1]octanyl, 3-oxa-8-azabicyclo[3.2.1]octanyl, 3,8-diazabicyclo[3.2.1]octanyl, 3,6-diazabicyclo[3.2.2]nonanyl, 3-oxa-7-azabicyclo[3.3.1]nonanyl and 3,9-diazabicyclo[4.2.1]nonanyl.

The term “C₄₋₉ spiroheterocycloalkyl” as used herein refers to saturated bicyclic ring systems containing 4 to 9 carbon atoms and at least one heteroatom selected from oxygen, sulphur and nitrogen, in which the two rings are linked by a common atom.

Suitable spiroheterocycloalkyl groups include 5-azaspiro[2.3]hexanyl, 5-azaspiro[2.4]heptanyl, 2-azaspiro[3.3]heptanyl, 2-oxa-6-azaspiro[3.3]heptanyl, 2-oxa-6-azaspiro[3.4]octanyl, 2-oxa-6-azaspiro[3.5]nonanyl, 7-oxa-2-azaspiro[3.5]nonanyl, 2-oxa-7-azaspiro[3.5]nonanyl and 2,4,8-triazaspiro[4.5]decanyl.

The term “heteroaryl” as used herein refers to monovalent aromatic groups containing at least 5 atoms derived from a single ring or multiple condensed rings, wherein one or more carbon atoms have been replaced by one or more heteroatoms selected from oxygen, sulphur and nitrogen. Suitable heteroaryl groups include furyl, benzofuryl, dibenzofuryl, thienyl, benzothienyl, thieno[2,3-*c*]pyrazolyl, thieno[3,4-*b*][1,4]dioxinyl, dibenzothienyl, pyrrolyl, indolyl, pyrrolo[2,3-*b*]pyridinyl, pyrrolo[3,2-*c*]pyridinyl, pyrrolo[3,4-*b*]pyridinyl, pyrazolyl, pyrazolo[1,5-*a*]pyridinyl, pyrazolo[3,4-*d*]pyrimidinyl, indazolyl, 4,5,6,7-tetrahydroindazolyl, oxazolyl, benzoxazolyl, isoxazolyl, thiazolyl, benzothiazolyl, isothiazolyl, imidazolyl, benzimidazolyl, imidazo[2,1-*b*]thiazolyl, imidazo[1,2-*a*]pyridinyl, imidazo[4,5-*b*]pyridinyl, purinyl, imidazo[1,2-*a*]pyrimidinyl, imidazo[1,2-*a*]pyrazinyl, oxadiazolyl, thiadiazolyl, triazolyl, [1,2,4]triazolo[1,5-*a*]pyrimidinyl, benzotriazolyl, tetrazolyl, pyridinyl, quinolinyl, isoquinolinyl, naphthyridinyl, pyridazinyl, cinnolinyl, phthalazinyl, pyrimidinyl, quinazolinyl, pyrazinyl, quinoxalinyl, pteridinyl, triazinyl and chromenyl groups.

The term “halogen” as used herein is intended to include fluorine, chlorine, bromine and iodine atoms, typically fluorine, chlorine or bromine.

Where the compounds of formula (I) have one or more asymmetric centres, they may accordingly exist as enantiomers. Where the compounds of use in the invention possess two or more asymmetric centres, they may additionally exist as diastereomers. The invention is to be understood to extend to the use of all such enantiomers and

diastereomers, and to mixtures thereof in any proportion, including racemates. Formula (I) and the formulae depicted hereinafter are intended to represent all individual stereoisomers and all possible mixtures thereof, unless stated or shown otherwise. In addition, compounds of formula (I) may exist as tautomers, for example keto ($\text{CH}_2\text{C}=\text{O}$) $\leftrightarrow$ enol ($\text{CH}=\text{CHOH}$) tautomers or amide ($\text{NHC}=\text{O}$) $\leftrightarrow$ hydroxyimine ($\text{N}=\text{COH}$) tautomers.

Formula (I) and the formulae depicted hereinafter are intended to represent all individual tautomers and all possible mixtures thereof, unless stated or shown otherwise.

It is to be understood that each individual atom present in formula (I), or in the formulae depicted hereinafter, may in fact be present in the form of any of its naturally occurring isotopes, with the most abundant isotope(s) being preferred. Thus, by way of example, each individual hydrogen atom present in formula (I), or in the formulae depicted hereinafter, may be present as a ^1H , ^2H (deuterium) or ^3H (tritium) atom, preferably ^1H . Similarly, by way of example, each individual carbon atom present in formula (I), or in the formulae depicted hereinafter, may be present as a ^{12}C , ^{13}C or ^{14}C atom, preferably ^{12}C .

In a particular aspect, the present invention provides a compound of formula (I) as depicted above or an *N*-oxide thereof, or a pharmaceutically acceptable salt or solvate thereof, or a glucuronide derivative thereof, or a co-crystal thereof, wherein

R^1 represents halogen or cyano; or C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-7} cycloalkyl, C_{4-7} cycloalkenyl, C_{3-7} cycloalkyl(C_{1-6})alkyl, aryl, aryl(C_{1-6})alkyl, C_{3-7} heterocycloalkyl, C_{3-7} heterocycloalkyl(C_{1-6})alkyl, C_{3-7} heterocycloalkenyl, C_{4-9} heterobicycloalkyl, heteroaryl, heteroaryl(C_{1-6})alkyl, (C_{3-7})heterocycloalkyl(C_{1-6})alkyl-aryl-, heteroaryl(C_{3-7})heterocycloalkyl-, (C_{3-7})cycloalkyl-heteroaryl-, (C_{3-7})cycloalkyl- (C_{1-6})alkyl-heteroaryl-, (C_{4-7})cycloalkenyl-heteroaryl-, (C_{4-9})bicycloalkyl-heteroaryl-, (C_{3-7})heterocycloalkyl-heteroaryl-, (C_{3-7})heterocycloalkyl(C_{1-6})alkyl-heteroaryl-, (C_{3-7})heterocycloalkenyl-heteroaryl-, (C_{4-9})heterobicycloalkyl-heteroaryl- or (C_{4-9})spiroheterocycloalkyl-heteroaryl-, any of which groups may be optionally substituted by one or more substituents; and

E , Y , R^2 and R^3 are as defined above.

Where the compounds in accordance with the invention comprise an optionally substituted straight or branched alkylene chain, typical values thereof include methylene ($-\text{CH}_2-$), (methyl)methylene, ethylene ($-\text{CH}_2\text{CH}_2-$), (ethyl)methylene, (dimethyl)methylene, (methyl)ethylene, propylene ($-\text{CH}_2\text{CH}_2\text{CH}_2-$), (propyl)methylene and (dimethyl)ethylene, any of which chains may be optionally substituted by one or more

substituents. Suitably, such chains are unsubstituted, monosubstituted or disubstituted. Typically, such chains are unsubstituted or monosubstituted. In one embodiment, such chains are unsubstituted. In another embodiment, such chains are monosubstituted. In a further embodiment, such chains are disubstituted.

5 Examples of typical substituents on the alkylene chain which may be present in a compound in accordance with the invention include halogen, cyano, trifluoromethyl, oxo, hydroxy, C₁₋₆ alkoxy, carboxy(C₁₋₆)alkoxy, trifluoromethoxy, amino, C₁₋₆ alkylamino, di(C₁₋₆)alkylamino, C₂₋₆ alkylcarbonylamino, carboxy, benzyloxycarbonyl, tetrazolyl, aminocarbonyl, C₁₋₆ alkylaminocarbonyl and di(C₁₋₆)alkylaminocarbonyl.

10 Specific examples of suitable substituents on the alkylene chain which may be present in a compound in accordance with the invention include fluoro, cyano, trifluoromethyl, hydroxy, methoxy, carboxymethoxy, amino, acetylamino, carboxy, benzyloxycarbonyl and tetrazolyl.

 In a first embodiment, E represents a covalent bond, whereby the integer Y is
15 attached directly to the triazole ring.

 In a second embodiment, E represents -O-, -S-, -S(O)-, -S(O)₂- or -N(R⁴)-. In a first aspect of that embodiment, E represents -O-. In a second aspect of that embodiment, E represents -S-. In a third aspect of that embodiment, E represents -S(O)-. In a fourth aspect of that embodiment, E represents -S(O)₂-. In a fifth aspect of that embodiment, E
20 represents -N(R⁴)-.

 In a third embodiment, E represents an optionally substituted straight or branched C₁₋₄ alkylene chain. In a first aspect of that embodiment, E represents an optionally substituted methylene (-CH₂-) linkage. In a second aspect of that embodiment, E represents an optionally substituted (methyl)methylene linkage. In a third aspect of that
25 embodiment, E represents an optionally substituted (ethyl)methylene linkage.

 Generally, E represents a covalent bond; or E represents -N(R⁴)-; or E represents an optionally substituted straight or branched C₁₋₄ alkylene chain.

 Typically, E represents -N(R⁴)-; or E represents an optionally substituted straight or branched C₁₋₄ alkylene chain.

30 Suitably, E represents a covalent bond; or E represents -N(R⁴)-; or E represents methylene (-CH₂-), (methyl)methylene or (ethyl)methylene, any of which groups may be optionally substituted by one or more substituents.

Generally, E represents $-N(R^4)-$; or E represents methylene ($-\text{CH}_2-$) or (ethyl)methylene, either of which groups may be optionally substituted by one or more substituents.

Appositely, E represents $-N(R^4)-$, or optionally substituted methylene.

5 Selected examples of typical substituents on the linkage represented by E include halogen, trifluoromethyl, hydroxy, C_{1-6} alkoxy, carboxy(C_{1-6})alkoxy, trifluoromethoxy, amino, C_{1-6} alkylamino, di(C_{1-6})alkylamino, C_{2-6} alkylcarbonylamino, carboxy, benzyloxycarbonyl and tetrazolyl.

Specific examples of typical substituents on the linkage represented by E include
10 fluoro, trifluoromethyl, hydroxy, methoxy, carboxymethoxy, trifluoromethoxy, amino, methylamino, dimethylamino, acetylamino, carboxy, benzyloxycarbonyl and tetrazolyl.

A particular example of a typical substituent on E is hydroxy.

Typical values of E include $-N(R^4)-$, $-\text{CH}_2-$, $-\text{CH}(\text{OH})-$, $-\text{CH}(\text{OCH}_3)-$,
15 $-\text{CH}(\text{OCH}_2\text{CO}_2\text{H})-$, $-\text{CH}(\text{NH}_2)-$, $-\text{CH}(\text{NHCOCH}_3)-$, $-\text{CH}(\text{CO}_2\text{H})-$, $-\text{CH}(\text{CO}_2\text{benzyl})-$,
 $-\text{CH}(\text{CH}_3)-$, $-\text{C}(\text{CH}_3)(\text{OH})-$ and $-\text{CH}(\text{CH}_2\text{CH}_3)-$; or E may represent a covalent bond.

Suitable values of E include $-N(R^4)-$, $-\text{CH}_2-$ and $-\text{CH}(\text{OH})-$. In one embodiment, E represents $-N(R^4)-$. In another embodiment, E represents $-\text{CH}_2-$. In a further embodiment, E represents $-\text{CH}(\text{OH})-$.

In another embodiment, E represents $-\text{CH}(\text{OCH}_3)-$.

20 In another embodiment, E represents $-\text{CH}(\text{NH}_2)-$.

In an additional embodiment, E represents $-\text{CH}(\text{CH}_3)-$. In a particular aspect of that embodiment, the $-\text{CH}(\text{CH}_3)-$ linkage represented by E is in the (*S*) stereochemical configuration.

In a further embodiment, E represents $-\text{C}(\text{CH}_3)(\text{OH})-$.

25 Generally, Y represents C_{3-7} cycloalkyl, aryl or heteroaryl, any of which groups may be optionally substituted by one or more substituents.

Typically, Y represents aryl or heteroaryl, either of which groups may be optionally substituted by one or more substituents.

In a first embodiment, Y represents optionally substituted C_{3-7} cycloalkyl. In one
30 aspect of that embodiment, Y represents unsubstituted C_{3-7} cycloalkyl. In another aspect of that embodiment, Y represents monosubstituted C_{3-7} cycloalkyl. In a further aspect of that embodiment, Y represents disubstituted C_{3-7} cycloalkyl.

In a second embodiment, Y represents optionally substituted aryl. In one aspect of that embodiment, Y represents unsubstituted aryl. In another aspect of that embodiment, Y represents monosubstituted aryl. In a further aspect of that embodiment, Y represents disubstituted aryl.

5 In a third embodiment, Y represents optionally substituted C₃₋₇ heterocycloalkyl. In one aspect of that embodiment, Y represents unsubstituted C₃₋₇ heterocycloalkyl. In another aspect of that embodiment, Y represents monosubstituted C₃₋₇ heterocycloalkyl. In a further aspect of that embodiment, Y represents disubstituted C₃₋₇ heterocycloalkyl.

10 In a fourth embodiment, Y represents optionally substituted heteroaryl. In one aspect of that embodiment, Y represents unsubstituted heteroaryl. In another aspect of that embodiment, Y represents monosubstituted heteroaryl. In a further aspect of that embodiment, Y represents disubstituted heteroaryl.

Suitably, Y represents benzocyclobutenyl, phenyl, thienyl, thiazolyl or pyridinyl, any of which groups may be optionally substituted by one or more substituents.

15 Appropriately, Y represents phenyl, thienyl or thiazolyl, any of which groups may be optionally substituted by one or more substituents.

Appositely, Y represents phenyl, which may be optionally substituted by one or more substituents.

20 Examples of optional substituents which may be present on the moiety Y include one, two or three substituents independently selected from halogen, cyano, nitro, C₁₋₆ alkyl, trifluoromethyl, hydroxy, C₁₋₆ alkoxy, difluoromethoxy, trifluoromethoxy, C₁₋₆ alkylthio, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, (C₁₋₆)alkylsulfonyloxy, amino, C₁₋₆ alkyl-amino, di(C₁₋₆)alkylamino, arylamino, C₂₋₆ alkylcarbonylamino, C₁₋₆ alkylsulfonylamino, formyl, C₂₋₆ alkylcarbonyl, C₃₋₆ cycloalkylcarbonyl, C₃₋₆ heterocycloalkylcarbonyl, 25 carboxy, C₂₋₆ alkoxy carbonyl, aminocarbonyl, C₁₋₆ alkylaminocarbonyl, di(C₁₋₆)alkylaminocarbonyl, aminosulfonyl, C₁₋₆ alkylaminosulfonyl and di(C₁₋₆)alkylaminosulfonyl.

Typical examples of optional substituents on the moiety Y include halogen, cyano and difluoromethoxy.

30 Suitable examples of optional substituents on the moiety Y include difluoro-methoxy.

Examples of particular substituents on the moiety Y include fluoro, chloro, bromo, cyano, nitro, methyl, isopropyl, trifluoromethyl, hydroxy, methoxy, difluoromethoxy, trifluoromethoxy, methylthio, methylsulfinyl, methylsulfonyl, methylsulfonyloxy, amino,

methylamino, *tert*-butylamino, dimethylamino, phenylamino, acetylamino, methylsulfonylamino, formyl, acetyl, cyclopropylcarbonyl, azetidiny carbonyl, pyrrolidinylcarbonyl, piperidinylcarbonyl, piperazinylcarbonyl, morpholinylcarbonyl, carboxy, methoxycarbonyl, aminocarbonyl, methylaminocarbonyl, dimethylaminocarbonyl, aminosulfonyl, methylaminosulfonyl and dimethylaminosulfonyl.

Typical examples of particular substituents on the moiety Y include fluoro, chloro, cyano and difluoromethoxy.

Suitable examples of particular substituents on the moiety Y include difluoromethoxy.

Typical values of Y include benzocyclobutenyl, phenyl, fluorophenyl (including 2-fluorophenyl, 3-fluorophenyl and 4-fluorophenyl), chlorophenyl (including 2-chlorophenyl, 3-chlorophenyl and 4-chlorophenyl), difluorophenyl (including 2,6-difluorophenyl), (chloro)(fluoro)phenyl (including 5-chloro-2-fluorophenyl and 2-chloro-5-fluorophenyl), dichlorophenyl (including 2,5-dichlorophenyl and 2,6-dichlorophenyl), methylphenyl (including 4-methylphenyl), dimethylphenyl (including 2,5-dimethylphenyl and 2,6-dimethylphenyl), (trifluoromethyl)phenyl [including 2-(trifluoromethyl)phenyl], (chloro)(trifluoromethyl)phenyl [including 5-chloro-2-(trifluoromethyl)phenyl], (methyl)-(trifluoromethyl)phenyl [including 2-methyl-5-(trifluoromethyl)phenyl], bis(trifluoromethyl)phenyl [including 2,5-bis(trifluoromethyl)phenyl], methoxyphenyl (including 2-methoxyphenyl), (difluoromethoxy)phenyl [including 2-(difluoromethoxy)phenyl and 3-(difluoromethoxy)phenyl], (difluoromethoxy)(fluoro)phenyl [including 2-(difluoromethoxy)-5-fluorophenyl and 2-(difluoromethoxy)-6-fluorophenyl], (chloro)(difluoromethoxy)phenyl [including 5-chloro-2-(difluoromethoxy)phenyl and 6-chloro-2-(difluoromethoxy)phenyl], (cyano)(difluoromethoxy)phenyl [including 6-cyano-2-(difluoromethoxy)phenyl], (trifluoromethoxy)phenyl [including 2-(trifluoromethoxy)phenyl], methylsulfonyloxyphenyl, (amino)(chloro)phenyl (including 5-amino-2-chlorophenyl), methylthienyl (including 3-methylthien-2-yl), methylthiazolyl (including 2-methyl-1,3-thiazol-4-yl), (chloro)(methyl)thiazolyl (including 5-chloro-2-methyl-1,3-thiazol-4-yl), dimethylthiazolyl (including 2,4-dimethyl-1,3-thiazol-5-yl) and pyridinyl (including pyridin-3-yl and pyridin-4-yl).

Selected values of Y include dichlorophenyl, dimethylphenyl, (difluoromethoxy)phenyl, (difluoromethoxy)(fluoro)phenyl, methylsulfonyloxyphenyl, methylthienyl and dimethylthiazolyl.

In one embodiment, Y represents 2,5-dichlorophenyl.

In another embodiment, Y represents 2,5-dimethylphenyl.

In a particular embodiment, Y represents 2-(difluoromethoxy)phenyl.

In another embodiment, Y represents (difluoromethoxy)(fluoro)phenyl.

5 In another embodiment, Y represents 3-methylthien-2-yl.









In another embodiment, Y represents 2,4-dimethyl-1,3-thiazol-5-yl.


























Suitably, R¹, R² and R³ independently represent hydrogen, halogen, cyano, trifluoromethyl or -CO₂R^d; or C₁₋₆ alkyl, C₂₋₆ alkynyl, aryl, C₃₋₇ heterocycloalkyl, C₃₋₇ heterocycloalkenyl, heteroaryl, (C₃₋₇)heterocycloalkyl(C₁₋₆)alkyl-aryl-, heteroaryl-(C₃₋₇)heterocycloalkyl-, (C₃₋₇)cycloalkyl-heteroaryl-, (C₃₋₇)cycloalkyl(C₁₋₆)alkyl-heteroaryl-, (C₄₋₇)cycloalkenyl-heteroaryl-, (C₄₋₉)bicycloalkyl-heteroaryl-, (C₃₋₇)heterocycloalkyl-heteroaryl-, (C₃₋₇)heterocycloalkyl(C₁₋₆)alkyl-heteroaryl-, (C₃₋₇)heterocycloalkenyl-heteroaryl-, (C₄₋₉)heterobicycloalkyl-heteroaryl- or (C₄₋₉)spiroheterocycloalkyl-heteroaryl-, any of which groups may be optionally substituted by one or more substituents.











Examples of optional substituents which may be present on R¹, R² or R³ include one, two or three substituents independently selected from halogen, halo(C₁₋₆)alkyl, cyano, cyano(C₁₋₆)alkyl, nitro, nitro(C₁₋₆)alkyl, C₁₋₆ alkyl, difluoromethyl, trifluoromethyl, difluoroethyl, trifluoroethyl, C₂₋₆ alkenyl, hydroxy, hydroxy(C₁₋₆)alkyl, C₁₋₆ alkoxy, difluoromethoxy, trifluoromethoxy, trifluoroethoxy, carboxy(C₃₋₇)cycloalkyloxy, C₁₋₃ alkylenedioxy, C₁₋₆ alkoxy(C₁₋₆)alkyl, C₁₋₆ alkylthio, C₁₋₆ alkylsulphinyl, C₁₋₆ alkylsulphonyl, (C₁₋₆)alkylsulphonyl(C₁₋₆)alkyl, oxo, amino, amino(C₁₋₆)alkyl, C₁₋₆ alkylamino, di(C₁₋₆)alkylamino, hydroxy(C₁₋₆)alkylamino, C₁₋₆ alkoxyamino, (C₁₋₆)alkoxy(C₁₋₆)alkylamino, [(C₁₋₆)alkoxy](hydroxy)(C₁₋₆)alkylamino, [(C₁₋₆)alkylthio](hydroxy)(C₁₋₆)alkylamino, N-[(C₁₋₆)alkyl]-N-[hydroxy(C₁₋₆)alkyl]amino, di(C₁₋₆)alkylamino(C₁₋₆)alkylamino, N-[di(C₁₋₆)alkylamino(C₁₋₆)alkyl]-N-[hydroxy(C₁₋₆)alkyl]amino, hydroxy(C₁₋₆)alkyl-(C₃₋₇)cycloalkylamino, (hydroxy)[(C₃₋₇)cycloalkyl(C₁₋₆)alkyl]amino, (C₃₋₇)heterocycloalkyl(C₁₋₆)alkylamino, oxo(C₃₋₇)heterocycloalkyl(C₁₋₆)alkylamino, (C₁₋₆)alkylheteroarylamino, heteroaryl(C₁₋₆)alkylamino, (C₁₋₆)alkylheteroaryl(C₁₋₆)alkylamino, C₂₋₆ alkylcarbonylamino, N-[(C₁₋₆)alkyl]-N-[(C₂₋₆)alkylcarbonyl]amino, (C₂₋₆)alkylcarbonylamino(C₁₋₆)alkyl, C₃₋₆ alkenylcarbonylamino, bis[(C₃₋₆)alkenylcarbonyl]amino, N-[(C₁₋₆)alkyl]-N-[(C₃₋₇)cycloalkylcarbonyl]amino, C₂₋₆ alkoxycarbonylamino, C₂₋₆ alkoxycarbonyl(C₁₋₆)alkylamino, C₁₋₆ alkylaminocarbonylamino, C₁₋₆ alkylsulphonyl-

amino, *N*-[(C₁₋₆)alkyl]-*N*-[(C₁₋₆)alkylsulphonyl]amino, bis[(C₁₋₆)alkylsulphonyl]amino, *N*-[(C₁₋₆)alkyl]-*N*-[carboxy(C₁₋₆)alkyl]amino, carboxy(C₃₋₇)cycloalkylamino, carboxy-(C₃₋₇)cycloalkyl(C₁₋₆)alkylamino, formyl, C₂₋₆ alkylcarbonyl, (C₃₋₇)cycloalkylcarbonyl, phenylcarbonyl, (C₂₋₆)alkylcarbonyloxy(C₁₋₆)alkyl, carboxy, carboxy(C₁₋₆)alkyl, C₂₋₆ alkoxy carbonyl, C₂₋₆ alkoxy carbonyl(C₁₋₆)alkyl, morpholinyl(C₁₋₆)alkoxy carbonyl, C₂₋₆ alkoxy carbonylmethylidenyl, a carboxylic acid isostere or prodrug moiety Ω , -(C₁₋₆)alkyl- Ω , aminocarbonyl, C₁₋₆ alkylaminocarbonyl, hydroxy(C₁₋₆)alkylamino-carbonyl, di(C₁₋₆)alkylaminocarbonyl, aminocarbonyl(C₁₋₆)alkyl, aminosulphonyl, di(C₁₋₆)alkylaminosulphonyl, (C₁₋₆)alkylsulphoximinyl and [(C₁₋₆)alkyl][*N*-(C₁₋₆)alkyl]-sulphoximinyl.

By the expression “carboxylic acid isostere or prodrug moiety” is meant any functional group, structurally distinct from a carboxylic acid moiety, that will be recognised by a biological system as being similar to, and thus capable of mimicking, a carboxylic acid moiety, or will be readily convertible by a biological system *in vivo* into a carboxylic acid moiety. A synopsis of some common carboxylic acid isosteres is presented by N.A. Meanwell in *J. Med. Chem.*, 2011, **54**, 2529-2591 (cf. in particular Figures 25 and 26). An alternative carboxylic acid isostere is described by N Pemberton *et al.* in *ACS Med. Chem. Lett.*, 2012, **3**, 574-578. Typical examples of suitable carboxylic acid isostere or prodrug moieties represented by Ω include the functional groups of formula (i) to (xliii):







5 wherein

the asterisk (*) represents the site of attachment to the remainder of the molecule;

n is zero, 1 or 2;

X represents oxygen or sulphur;

R^f represents hydrogen, C₁₋₆ alkyl or -CH₂CH(OH)CH₂OH;

10 R^g represents C₁₋₆ alkyl, trifluoromethyl, -CH₂CH₂F, -CH₂CHF₂, -CH₂CF₃ or -CF₂CF₃;

R^h represents hydrogen, cyano or -CO₂R^d, in which R^d is as defined above; and

R^j represents hydrogen or halogen.

In one embodiment, n is zero. In another embodiment, n is 1. In a further

15 embodiment, n is 2.

In one embodiment, X represents oxygen. In another embodiment, X represents sulphur.

In one embodiment, R^f represents hydrogen. In another embodiment, R^f represents C₁₋₆ alkyl, especially methyl. In a further embodiment, R^f is -CH₂CH(OH)CH₂OH.

20 In one embodiment, R^g represents C₁₋₆ alkyl, especially methyl. In another embodiment, R^g represents trifluoromethyl, -CH₂CH₂F, -CH₂CHF₂, -CH₂CF₃ or -CF₂CF₃.

In a first aspect of that embodiment, R^g represents trifluoromethyl. In a second aspect of that embodiment, R^g represents $-\text{CH}_2\text{CH}_2\text{F}$. In a third aspect of that embodiment, R^g represents $-\text{CH}_2\text{CHF}_2$. In a fourth aspect of that embodiment, R^g represents $-\text{CH}_2\text{CF}_3$. In a fifth aspect of that embodiment, R^g represents $-\text{CF}_2\text{CF}_3$.

5 In one embodiment, R^h is hydrogen. In another embodiment, R^h represents cyano. In a further embodiment, R^h represents $-\text{CO}_2R^d$, especially methoxycarbonyl.

In one embodiment, R^j represents hydrogen. In another embodiment, R^j represents halogen, especially chloro.

In a selected embodiment, Ω represents tetrazolyl, especially a C-linked tetrazolyl moiety of formula (xxiv) or (xxv) as depicted above, in particular a group of formula (xxiv) as depicted above.

In another embodiment, Ω represents C_{1-6} alkylsulphonylaminocarbonyl, i.e. a moiety of formula (iii) as depicted above wherein R^g represents C_{1-6} alkyl.

In another embodiment, Ω represents C_{1-6} alkylaminosulphonyl, i.e. a moiety of formula (x) as depicted above wherein R^g represents C_{1-6} alkyl.

In a further embodiment, Ω represents $(\text{C}_{1-6})\text{alkylcarbonylaminosulphonyl}$, i.e. a moiety of formula (v) as depicted above wherein R^g represents C_{1-6} alkyl.

Typical examples of optional substituents which may be present on R^1 , R^2 or R^3 include one, two or three substituents independently selected from carboxy and amino-sulphonyl.

Examples of particular substituents on R^1 , R^2 or R^3 include fluoro, chloro, bromo, fluoromethyl, fluoroisopropyl, cyano, cyanoethyl, nitro, nitromethyl, methyl, ethyl, isopropyl, isobutyl, *tert*-butyl, difluoromethyl, trifluoromethyl, difluoroethyl, trifluoroethyl, ethenyl, hydroxy, hydroxymethyl, hydroxyisopropyl, methoxy, isopropoxy, difluoromethoxy, trifluoromethoxy, trifluoroethoxy, carboxycyclobutylloxy, methylenedioxy, ethylenedioxy, methoxymethyl, methoxyethyl, methylthio, methylsulphinyl, methylsulphonyl, methylsulphonyl ethyl, oxo, amino, aminomethyl, aminoisopropyl, methylamino, ethylamino, dimethylamino, hydroxyethylamino, hydroxypropylamino, (hydroxy)(methyl)propylamino, methoxyamino, methoxyethylamino, (hydroxy)-(methoxy)(methyl)propylamino, (hydroxy)(methylthio)butylamino, *N*-(hydroxyethyl)-*N*-(methyl)amino, dimethylaminoethylamino, (dimethylamino)(methyl)propylamino, *N*-(dimethylaminoethyl)-*N*-(hydroxyethyl)amino, hydroxymethylcyclopentylamino, hydroxycyclobutylmethylamino, (cyclopropyl)(hydroxy)propylamino, morpholinylethyl-

amino, oxopyrrolidinylmethylamino, ethyloxadiazolylamino, methylthiadiazolylamino, thiazolylmethylamino, thiazolyethylamino, pyrimidinylmethylamino, methylpyrazolylmethylamino, acetylamino, *N*-acetyl-*N*-methylamino, *N*-isopropylcarbonyl-*N*-methylamino, acetylaminomethyl, ethenylcarbonylamino, bis(ethenylcarbonyl)amino, *N*-cyclopropylcarbonyl-*N*-methylamino, methoxycarbonylamino, ethoxycarbonylamino, *tert*-butoxycarbonylamino, methoxycarbonylethylamino, ethylaminocarbonylamino, butylaminocarbonylamino, methylsulphonylamino, *N*-methyl-*N*-(methylsulphonyl)amino, bis(methylsulphonyl)amino, *N*-(carboxymethyl)-*N*-methylamino, *N*-(carboxyethyl)-*N*-methylamino, carboxycyclopentylamino, carboxycyclopropylmethylamino, formyl, acetyl, isopropylcarbonyl, cyclobutylcarbonyl, phenylcarbonyl, acetoxisopropyl, carboxy, carboxymethyl, carboxyethyl, methoxycarbonyl, ethoxycarbonyl, *n*-butoxycarbonyl, *tert*-butoxycarbonyl, methoxycarbonylmethyl, ethoxycarbonylmethyl, ethoxycarbonylethyl, morpholinylethoxycarbonyl, ethoxycarbonylmethylidenyl, methylsulphonylamino-carbonyl, acetylaminosulphonyl, methoxyaminocarbonyl, tetrazolyl, tetrazolylmethyl, hydroxyoxadiazolyl, aminocarbonyl, methylaminocarbonyl, hydroxyethylaminocarbonyl, dimethylaminocarbonyl, aminocarbonylmethyl, aminosulphonyl, methylaminosulphonyl, dimethylaminosulphonyl, methylsulphoximinyl and (methyl)(*N*-methyl)sulphoximinyl.

Typical examples of particular substituents on R¹, R² or R³ include carboxy and aminosulphonyl.

Typically, R¹ represents hydrogen, halogen, cyano or -CO₂R^d; or C₁₋₆ alkyl, C₂₋₆ alkynyl, aryl, C₃₋₇ heterocycloalkyl, C₃₋₇ heterocycloalkenyl, heteroaryl, (C₃₋₇)heterocycloalkyl(C₁₋₆)alkyl-aryl-, heteroaryl(C₃₋₇)heterocycloalkyl-, (C₃₋₇)cycloalkyl-heteroaryl-, (C₃₋₇)cycloalkyl(C₁₋₆)alkyl-heteroaryl-, (C₄₋₇)cycloalkenyl-heteroaryl-, (C₄₋₉)bicycloalkyl-heteroaryl-, (C₃₋₇)heterocycloalkyl-heteroaryl-, (C₃₋₇)heterocycloalkyl(C₁₋₆)alkyl-heteroaryl-, (C₃₋₇)heterocycloalkenyl-heteroaryl-, (C₄₋₉)heterobicycloalkyl-heteroaryl- or (C₄₋₉)spiroheterocycloalkyl-heteroaryl-, any of which groups may be optionally substituted by one or more substituents.

Suitably, R¹ represents halogen, cyano or -CO₂R^d; or C₁₋₆ alkyl, C₂₋₆ alkynyl, aryl, C₃₋₇ heterocycloalkyl, C₃₋₇ heterocycloalkenyl, heteroaryl, (C₃₋₇)heterocycloalkyl-(C₁₋₆)alkyl-aryl-, heteroaryl(C₃₋₇)heterocycloalkyl-, (C₃₋₇)cycloalkyl-heteroaryl-, (C₃₋₇)cycloalkyl(C₁₋₆)alkyl-heteroaryl-, (C₄₋₇)cycloalkenyl-heteroaryl-, (C₄₋₉)bicycloalkyl-heteroaryl-, (C₃₋₇)heterocycloalkyl-heteroaryl-, (C₃₋₇)heterocycloalkyl(C₁₋₆)alkyl-heteroaryl-, (C₃₋₇)heterocycloalkenyl-heteroaryl-, (C₄₋₉)heterobicycloalkyl-heteroaryl- or

(C₄₋₉)spiroheterocycloalkyl-heteroaryl-, any of which groups may be optionally substituted by one or more substituents.

Generally, R¹ represents halogen or cyano; or C₁₋₆ alkyl, C₂₋₆ alkynyl, aryl, C₃₋₇ heterocycloalkyl, C₃₋₇ heterocycloalkenyl, heteroaryl, (C₃₋₇)heterocycloalkyl(C₁₋₆)alkyl-
 5 aryl-, heteroaryl(C₃₋₇)heterocycloalkyl-, (C₃₋₇)cycloalkyl-heteroaryl-, (C₃₋₇)cycloalkyl-
 (C₁₋₆)alkyl-heteroaryl-, (C₄₋₇)cycloalkenyl-heteroaryl-, (C₄₋₉)bicycloalkyl-heteroaryl-,
 (C₃₋₇)heterocycloalkyl-heteroaryl-, (C₃₋₇)heterocycloalkyl(C₁₋₆)alkyl-heteroaryl-,
 (C₃₋₇)heterocycloalkenyl-heteroaryl-, (C₄₋₉)heterobicycloalkyl-heteroaryl- or
 (C₄₋₉)spiroheterocycloalkyl-heteroaryl-, any of which groups may be optionally
 10 substituted by one or more substituents.

More generally, R¹ represents aryl or (C₃₋₇)heterocycloalkyl-heteroaryl-, either of which groups may be optionally substituted by one or more substituents.

In a first embodiment, R¹ represents hydrogen.

In a second embodiment, R¹ represents halogen. In one aspect of that
 15 embodiment, R¹ represents bromo.

In a third embodiment, R¹ represents cyano.

In a fourth embodiment, R¹ represents -CO₂R^d.

In a fifth embodiment, R¹ represents optionally substituted C₁₋₆ alkyl. In one aspect of that embodiment, R¹ represents optionally substituted ethyl.

In a sixth embodiment, R¹ represents optionally substituted C₂₋₆ alkynyl. In one aspect of that embodiment, R¹ represents optionally substituted butynyl.

In a seventh embodiment, R¹ represents optionally substituted aryl. In one aspect of that embodiment, R¹ represents optionally substituted phenyl.

In an eighth embodiment, R¹ represents optionally substituted C₃₋₇
 25 heterocycloalkyl.

In a ninth embodiment, R¹ represents optionally substituted C₃₋₇ heterocycloalkenyl.

In a tenth embodiment, R¹ represents optionally substituted heteroaryl. In selected aspects of that embodiment, R¹ represents benzofuryl, thienyl, indolyl, pyrazolyl,
 30 indazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridinyl, quinolinyl, pyridazinyl, pyrimidinyl or pyrazinyl, any of which groups may be optionally substituted by one or more substituents.

In an eleventh embodiment, R^1 represents optionally substituted (C_{3-7})-heterocycloalkyl(C_{1-6})alkyl-aryl-. In a first aspect of that embodiment, R^1 represents optionally substituted pyrrolidinylmethylphenyl-. In a second aspect of that embodiment, R^1 represents optionally substituted piperazinylmethylphenyl-.

5 In a twelfth embodiment, R^1 represents optionally substituted heteroaryl(C_{3-7})-heterocycloalkyl-. In one aspect of that embodiment, R^1 represents optionally substituted pyridinylpiperazinyl-.

In a thirteenth embodiment, R^1 represents optionally substituted (C_{3-7})cycloalkyl-heteroaryl-. In a first aspect of that embodiment, R^1 represents optionally substituted
10 cyclohexylpyrazolyl-. In a second aspect of that embodiment, R^1 represents optionally substituted cyclohexylpyridinyl-. In a third aspect of that embodiment, R^1 represents optionally substituted cyclopropylpyrimidinyl-. In a fourth aspect of that embodiment, R^1 represents optionally substituted cyclobutylpyrimidinyl-. In a fifth aspect of that
15 aspect of that embodiment, R^1 represents optionally substituted cyclopentylpyrimidinyl-. In a sixth aspect of that embodiment, R^1 represents optionally substituted cyclohexylpyrimidinyl-. In a seventh aspect of that embodiment, R^1 represents optionally substituted cyclohexylpyrazinyl-.

In a fourteenth embodiment, R^1 represents optionally substituted (C_{4-7})-cycloalkenyl-heteroaryl-.

20 In a fifteenth embodiment, R^1 represents optionally substituted (C_{3-7})-heterocycloalkyl-heteroaryl-. In a first aspect of that embodiment, R^1 represents optionally substituted pyrrolidinylpyridinyl-. In a second aspect of that embodiment, R^1 represents optionally substituted tetrahydropyranylpyridinyl-. In a third aspect of that
25 embodiment, R^1 represents optionally substituted piperidinylpyridinyl-. In a fourth aspect of that embodiment, R^1 represents optionally substituted piperazinylpyridinyl-. In a fifth aspect of that embodiment, R^1 represents optionally substituted morpholinylpyridinyl-. In a sixth aspect of that embodiment, R^1 represents optionally substituted thiomorpholinylpyridinyl-. In a seventh aspect of that embodiment, R^1 represents optionally substituted diazepanylpyridinyl-. In an eighth aspect of that embodiment, R^1 represents optionally
30 substituted oxetanylpyrimidinyl-. In a ninth aspect of that embodiment, R^1 represents optionally substituted azetidinympyrimidinyl-. In a tenth aspect of that embodiment, R^1 represents optionally substituted tetrahydrofuranylpyrimidinyl-. In an eleventh aspect of that embodiment, R^1 represents optionally substituted pyrrolidinylpyrimidinyl-. In a

twelfth aspect of that embodiment, R¹ represents optionally substituted tetrahydropyranyl-pyrimidinyl-. In a thirteenth aspect of that embodiment, R¹ represents optionally substituted piperidinylpyrimidinyl-. In a fourteenth aspect of that embodiment, R¹ represents optionally substituted piperazinylpyrimidinyl-. In a fifteenth aspect of that embodiment, R¹ represents optionally substituted morpholinylpyrimidinyl-. In a sixteenth aspect of that embodiment, R¹ represents optionally substituted thiomorpholinyl-pyrimidinyl-. In a seventeenth aspect of that embodiment, R¹ represents optionally substituted azepanylpyrimidinyl-. In an eighteenth aspect of that embodiment, R¹ represents optionally substituted oxazepanylpyrimidinyl-. In a nineteenth aspect of that embodiment, R¹ represents optionally substituted diazepanylpyrimidinyl-. In a twentieth aspect of that embodiment, R¹ represents optionally substituted thiadiazepanyl-pyrimidinyl-. In a twenty-first aspect of that embodiment, R¹ represents optionally substituted oxetanylpyrazinyl-. In a twenty-second aspect of that embodiment, R¹ represents optionally substituted piperidinylpyrazinyl-.

In a sixteenth embodiment, R¹ represents optionally substituted (C₃₋₇)-heterocycloalkyl(C₁₋₆)alkyl-heteroaryl-. In a first aspect of that embodiment, R¹ represents optionally substituted morpholinylmethylthienyl-. In a second aspect of that embodiment, R¹ represents optionally substituted morpholinylethylpyrazolyl-.

In a seventeenth embodiment, R¹ represents optionally substituted (C₃₋₇)-heterocycloalkenyl-heteroaryl-.

In an eighteenth embodiment, R¹ represents optionally substituted (C₄₋₉)-heterobicycloalkyl-heteroaryl-.

In a nineteenth embodiment, R¹ represents optionally substituted (C₄₋₉)-spiroheterocycloalkyl-heteroaryl-.

In a twentieth embodiment, R¹ represents optionally substituted (C₃₋₇)cycloalkyl-(C₁₋₆)alkyl-heteroaryl-. In one aspect of that embodiment, R¹ represents optionally substituted cyclohexylmethylpyrimidinyl-.

In a twenty-first embodiment, R¹ represents optionally substituted (C₄₋₉)-bicycloalkyl-heteroaryl-.

Appositely, R¹ represents hydrogen, bromo, cyano or -CO₂R^d; or ethyl, butynyl, phenyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, 1,2,3,6-tetrahydropyridinyl, benzofuryl, thienyl, indolyl, pyrazolyl, indazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridinyl, quinolinyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrrolidinylmethylphenyl,

piperazinylmethylphenyl, pyridinylpiperazinyl, cyclohexylpyrazolyl, cyclohexylpyridinyl, cyclopropylpyrimidinyl, cyclobutylpyrimidinyl, cyclopentylpyrimidinyl, cyclohexylpyrimidinyl, cyclohexylpyrazinyl, cyclohexylmethylpyrimidinyl, cyclohexenylpyridinyl, cyclohexenylpyrimidinyl, bicyclo[3.1.0]hexanylpyridinyl, bicyclo[3.1.0]hexanylpyrimidinyl, bicyclo[4.1.0]heptanylpyrimidinyl, bicyclo[2.2.2]octanylpyrimidinyl, pyrrolidinylpyridinyl, tetrahydropyranylpyridinyl, piperidinylpyridinyl, piperazinylpyridinyl, morpholinylpyridinyl, thiomorpholinylpyridinyl, diazepanylpyridinyl, oxetanylpyrimidinyl, azetidinylypyrimidinyl, tetrahydrofuranylpyrimidinyl, pyrrolidinylpyrimidinyl, tetrahydropyranylpyrimidinyl, piperidinylpyrimidinyl, piperazinylpyrimidinyl, hexahydro-[1,2,5]thiadiazolo[2,3-*a*]pyrazinylpyrimidinyl, morpholinylpyrimidinyl, thiomorpholinylpyrimidinyl, azepanylpyrimidinyl, oxazepanylpyrimidinyl, diazepanylpyrimidinyl, thiadiazepanylpyrimidinyl, oxetanylpyrazinyl, piperidinylpyrazinyl, morpholinylmethylthienyl, morpholinylethylpyrazolyl, 3-azabicyclo[3.1.0]hexanylpyridinyl, 3-azabicyclo[3.1.0]hexanylpyridazinyl, 3-azabicyclo[3.1.0]hexanylpyrimidinyl, 2-oxa-5-azabicyclo[2.2.1]heptanylpyrimidinyl, 3-azabicyclo[3.1.1]heptanylpyrimidinyl, 3-azabicyclo[4.1.0]heptanylpyridinyl, 3-azabicyclo[4.1.0]heptanylpyrimidinyl, 2-oxabicyclo[2.2.2]octanylpyrimidinyl, 3-azabicyclo[3.2.1]octanylpyrimidinyl, 8-azabicyclo[3.2.1]octanylpyrimidinyl, 3-oxa-8-azabicyclo[3.2.1]octanylpyrimidinyl, 3,6-diazabicyclo[3.2.2]nonanylpyrimidinyl, 3-oxa-7-azabicyclo[3.3.1]nonanylpyrimidinyl, 5-azaspiro[2.3]hexanylpyrimidinyl, 5-azaspiro[2.4]heptanylpyrimidinyl, 2-azaspiro[3.3]heptanylpyrimidinyl, 2-oxa-6-azaspiro[3.3]heptanylpyrimidinyl, 2-oxa-6-azaspiro[3.4]octanylpyrimidinyl, 2-oxa-6-azaspiro[3.5]nonanylpyrimidinyl, 2-oxa-7-azaspiro[3.5]nonanylpyrimidinyl or 2,4,8-triazaspiro[4.5]decanylpyrimidinyl, any of which groups may be optionally substituted by one or more substituents.

Illustratively, R¹ represents phenyl, piperidinylpyrimidinyl or morpholinylpyrimidinyl, any of which groups may be optionally substituted by one or more substituents.

Typical examples of optional substituents on R¹ include one, two or three substituents independently selected from halogen, halo(C₁₋₆)alkyl, cyano, cyano(C₁₋₆)alkyl, nitro(C₁₋₆)alkyl, C₁₋₆ alkyl, trifluoromethyl, trifluoroethyl, C₂₋₆ alkenyl, hydroxy, hydroxy(C₁₋₆)alkyl, C₁₋₆ alkoxy, trifluoroethoxy, carboxy(C₃₋₇)cycloalkyloxy, C₁₋₆ alkylthio, C₁₋₆ alkylsulphonyl, (C₁₋₆)alkylsulphonyl(C₁₋₆)alkyl, oxo, amino, amino-

(C₁₋₆)alkyl, C₁₋₆ alkylamino, di(C₁₋₆)alkylamino, (C₁₋₆)alkoxy(C₁₋₆)alkylamino, *N*-[(C₁₋₆)alkyl]-*N*-[hydroxy(C₁₋₆)alkyl]amino, (C₂₋₆)alkylcarbonylamino(C₁₋₆)alkyl, C₁₋₆ alkylsulphonylamino, *N*-[(C₁₋₆)alkyl]-*N*-[(C₁₋₆)alkylsulphonyl]amino, bis[(C₁₋₆)alkylsulphonyl]amino, *N*-[(C₁₋₆)alkyl]-*N*-[carboxy(C₁₋₆)alkyl]amino, carboxy(C₃₋₇)cycloalkyl-amino, carboxy(C₃₋₇)cycloalkyl(C₁₋₆)alkylamino, formyl, C₂₋₆ alkylcarbonyl, (C₂₋₆)alkylcarbonyloxy(C₁₋₆)alkyl, carboxy, carboxy(C₁₋₆)alkyl, C₂₋₆ alkoxycarbonyl, C₂₋₆ alkoxycarbonyl(C₁₋₆)alkyl, morpholinyl(C₁₋₆)alkoxycarbonyl, C₂₋₆ alkoxycarbonylmethylidenyl, a carboxylic acid isostere or prodrug moiety Ω as defined herein, -(C₁₋₆)alkyl- Ω , aminocarbonyl, aminosulphonyl, (C₁₋₆)alkylsulphoximinyl and

10 [(C₁₋₆)alkyl][*N*-(C₁₋₆)alkyl]sulphoximinyl.

Suitable examples of optional substituents on R¹ include one, two or three substituents independently selected from carboxy and aminosulphonyl.

Typical examples of particular substituents on R¹ include one, two or three substituents independently selected from fluoro, chloro, fluoromethyl, fluoroisopropyl,

15 cyano, cyanoethyl, nitromethyl, methyl, ethyl, isopropyl, trifluoromethyl, trifluoroethyl, ethenyl, hydroxy, hydroxymethyl, hydroxyisopropyl, methoxy, isopropoxy, trifluoroethoxy, carboxycyclobutyl, methylthio, methylsulphonyl, methylsulphonylethyl, oxo, amino, aminomethyl, aminoisopropyl, methylamino, dimethylamino, methoxyethylamino, *N*-(hydroxyethyl)-*N*-(methyl)amino, acetaminomethyl, methylsulphonylamino, *N*-methyl-*N*-(methylsulphonyl)amino, bis(methylsulphonyl)amino, *N*-(carboxyethyl)-*N*-(methyl)amino, carboxycyclopentylamino, carboxycyclopropylmethylamino, formyl, acetyl, acetoxisopropyl, carboxy, carboxymethyl, carboxyethyl, methoxycarbonyl, ethoxycarbonyl, *n*-butoxycarbonyl, *tert*-butoxycarbonyl, methoxycarbonylmethyl, ethoxycarbonylmethyl, ethoxycarbonylethyl, morpholinylethoxycarbonyl, ethoxycarbonylmethylidenyl, methylsulphonylaminocarbonyl, acetaminosulphonyl, methoxyaminocarbonyl, tetrazolyl, tetrazolylmethyl, hydroxyoxadiazolyl, aminocarbonyl, aminosulphonyl, methylsulphoximinyl and (methyl)(*N*-methyl)sulphoximinyl.

20

25

Suitable examples of particular substituents on R¹ include one, two or three substituents independently selected from carboxy and aminosulphonyl.

30 In a particular embodiment, R¹ is substituted by hydroxy(C₁₋₆)alkyl. In one aspect of that embodiment, R¹ is substituted by hydroxyisopropyl, especially 2-hydroxyprop-2-yl.

Selected values of R¹ include hydrogen, bromo, cyano, -CO₂R^d, methoxycarbonylethyl, ethoxycarbonylethyl, hydroxybutynyl, chlorophenyl, hydroxyphenyl, methyl-

sulphonylphenyl, aminomethylphenyl, aminoisopropylphenyl, acetylaminomethylphenyl,
 acetylphenyl, methoxycarbonylphenyl, aminocarbonylphenyl, aminosulphonylphenyl,
 acetylaminosulphonylphenyl, (methoxycarbonyl)(methyl)pyrrolidinyl, oxopiperidinyl,
 ethoxycarbonylpiperidinyl, methylsulphonylpiperazinyl, morpholinyl, methylsulphonyl-
 5 1,2,3,6-tetrahydropyridinyl, acetyl-1,2,3,6-tetrahydropyridinyl, *tert*-butoxycarbonyl-
 1,2,3,6-tetrahydropyridinyl, methoxycarbonylmethyl-1,2,3,6-tetrahydropyridinyl,
 benzofuryl, thienyl, indolyl, pyrazolyl, methylpyrazolyl, dimethylpyrazolyl, (methyl)[*N*-
 methyl-*N*-(methylsulfonyl)amino]pyrazolyl, methylindazolyl, dimethylisoxazolyl,
 hydroxyisopropylthiazolyl, methylimidazolyl, dimethylimidazolyl, pyridinyl, fluoro-
 10 pyridinyl, cyanopyridinyl, methylpyridinyl, (cyano)(methyl)pyridinyl, dimethylpyridinyl,
 trifluoromethylpyridinyl, ethenylpyridinyl, hydroxyisopropylpyridinyl, methoxypyridinyl,
 (methoxy)(methyl)pyridinyl, isopropoxypyridinyl, trifluoroethoxypyridinyl, (methyl)-
 (trifluoroethoxy)pyridinyl, methylsulphonylpyridinyl, oxopyridinyl, (methyl)(oxo)-
 pyridinyl, (dimethyl)(oxo)pyridinyl, aminopyridinyl, methylaminopyridinyl, dimethyl-
 15 aminopyridinyl, methoxyethylaminopyridinyl, *N*-(hydroxyethyl)-*N*-(methyl)amino-
 pyridinyl, methylsulphonylaminopyridinyl, [bis(methylsulphonyl)amino]pyridinyl,
 carboxypyridinyl, quinolinyl, hydroxypyridazinyl, pyrimidinyl, fluoroisopropyl-
 pyrimidinyl, hydroxyisopropylpyrimidinyl, methoxypyrimidinyl, carboxycyclobutyl-
 pyrimidinyl, methylthiopyrimidinyl, methylsulphonylpyrimidinyl, oxopyrimidinyl,
 20 aminopyrimidinyl, dimethylaminopyrimidinyl, methoxyethylaminopyrimidinyl, *N*-
 (carboxyethyl)-*N*-(methyl)aminopyrimidinyl, carboxycyclopentylaminopyrimidinyl,
 carboxycyclopropylmethylaminopyrimidinyl, acetoxymethylaminopyrimidinyl,
 ethoxycarbonylmethylaminopyrimidinyl, hydroxypyrazinyl, hydroxyisopropylpyrazinyl,
 pyrrolidinylmethylphenyl, piperazinylmethylphenyl, pyridinylpiperazinyl, carboxy-
 25 cyclohexylpyrazolyl, carboxycyclohexylpyridinyl, fluoromethylcyclopropylpyrimidinyl,
 acetylaminomethylcyclopropylpyrimidinyl, hydroxycyclobutylpyrimidinyl, carboxy-
 cyclopentylpyrimidinyl, carboxycyclohexylpyrimidinyl, (carboxy)(methyl)cyclohexyl-
 pyrimidinyl, (carboxy)(hydroxy)cyclohexylpyrimidinyl, carboxymethylcyclohexyl-
 pyrimidinyl, ethoxycarbonylcyclohexylpyrimidinyl, (methoxycarbonyl)(methyl)-
 30 cyclohexylpyrimidinyl, (ethoxycarbonyl)(methyl)cyclohexylpyrimidinyl, carboxy-
 cyclohexylpyrazinyl, carboxycyclohexylmethylpyrimidinyl, carboxycyclohexenyl-
 pyridinyl, carboxycyclohexenylpyrimidinyl, ethoxycarbonylcyclohexenylpyrimidinyl,
 carboxybicyclo[3.1.0]hexanylpyridinyl, carboxybicyclo[3.1.0]hexanylpyrimidinyl,

ethoxycarbonylbicyclo[3.1.0]hexanylpyrimidinyl, carboxybicyclo[4.1.0]heptanylpyrimidinyl, carboxybicyclo[2.2.2]octanylpyrimidinyl, pyrrolidinylpyridinyl, hydroxypyrrolidinylpyridinyl, hydroxytetrahydropyranylpyridinyl, piperidinylpyridinyl, acetylpiperidinylpyridinyl, (carboxy)(methyl)piperidinylpyridinyl, [(carboxy)(methyl)piperidinyl](fluoro)pyridinyl, [(carboxy)(methyl)piperidinyl](chloro)pyridinyl, piperazinylpyridinyl, (methyl)(piperazinyl)pyridinyl, cyanoethylpiperazinylpyridinyl, trifluoroethylpiperazinylpyridinyl, methylsulphonylpiperazinylpyridinyl, methylsulphonylethylpiperazinylpyridinyl, oxopiperazinylpyridinyl, acetylpiperazinylpyridinyl, (*tert*-butoxycarbonylpiperazinyl)(methyl)pyridinyl, carboxymethylpiperazinylpyridinyl, carboxyethylpiperazinylpyridinyl, ethoxycarbonylmethylpiperazinylpyridinyl, ethoxycarbonylethylpiperazinylpyridinyl, morpholinylpyridinyl, thiomorpholinylpyridinyl, oxothiomorpholinylpyridinyl, dioxothiomorpholinylpyridinyl, oxodiazepanylpyridinyl, fluorooxetanylpyrimidinyl, hydroxyoxetanylpyrimidinyl, hydroxyazetidinyipyrimidinyl, (hydroxy)(methyl)azetidinyipyrimidinyl, carboxyazetidinyipyrimidinyl, (*tert*-butoxycarbonyl)(hydroxy)azetidinyipyrimidinyl, tetrazolylazetidinyipyrimidinyl, hydroxytetrahydrofuranylpyrimidinyl, hydroxypyrrolidinylpyrimidinyl, carboxypyrrolidinylpyrimidinyl, (carboxy)(methyl)pyrrolidinylpyrimidinyl, carboxymethylpyrrolidinylpyrimidinyl, ethoxycarbonylpyrrolidinylpyrimidinyl, fluoro-tetrahydropyranylpyrimidinyl, hydroxytetrahydropyranylpyrimidinyl, difluoropiperidinylpyrimidinyl, (cyano)(methyl)piperidinylpyrimidinyl, (hydroxy)(nitromethyl)piperidinylpyrimidinyl, (hydroxy)(methyl)piperidinylpyrimidinyl, (hydroxy)(trifluoromethyl)piperidinylpyrimidinyl, (hydroxymethyl)(methyl)piperidinylpyrimidinyl, methylsulphonylpiperidinylpyrimidinyl, oxopiperidinylpyrimidinyl, (formyl)(methyl)piperidinylpyrimidinyl, carboxypiperidinylpyrimidinyl, (carboxy)(fluoro)piperidinylpyrimidinyl, (carboxy)(methyl)piperidinylpyrimidinyl, (carboxy)(ethyl)piperidinylpyrimidinyl, (carboxy)(trifluoromethyl)piperidinylpyrimidinyl, (carboxy)(hydroxy)piperidinylpyrimidinyl, (carboxy)(hydroxymethyl)piperidinylpyrimidinyl, (carboxy)(methoxy)piperidinylpyrimidinyl, (amino)(carboxy)piperidinylpyrimidinyl, carboxymethylpiperidinylpyrimidinyl, methoxycarbonylpiperidinylpyrimidinyl, ethoxycarbonylpiperidinylpyrimidinyl, (ethoxycarbonyl)(fluoro)piperidinylpyrimidinyl, (methoxycarbonyl)(methyl)piperidinylpyrimidinyl, (ethyl)(methoxycarbonyl)piperidinylpyrimidinyl, (isopropyl)(methoxycarbonyl)piperidinylpyrimidinyl, (ethoxycarbonyl)(methyl)piperidinylpyrimidinyl, (*n*-butoxycarbonyl)(methyl)piperidinylpyrimidinyl,

(ethoxycarbonyl)(trifluoromethyl)piperidinylpyrimidinyl, (ethoxycarbonyl)-
 (hydroxymethyl)piperidinylpyrimidinyl, (methoxy)(methoxycarbonyl)piperidinyl-
 pyrimidinyl, (carboxy)(methoxycarbonyl)piperidinylpyrimidinyl, (methyl)-
 (morpholinylethoxycarbonyl)piperidinylpyrimidinyl, ethoxycarbonylmethylpiperidinyl-
 5 pyrimidinyl, methylsulphonylaminocarbonylpiperidinylpyrimidinyl, acetylamino-
 sulphonylpiperidinylpyrimidinyl, methoxyaminocarbonylpiperidinylpyrimidinyl,
 tetrazolylpiperidinylpyrimidinyl, hydroxyoxadiazolylpiperidinylpyrimidinyl, amino-
 sulphonylpiperidinylpyrimidinyl, piperazinylpyrimidinyl, methylsulphonylpiperazinyl-
 pyrimidinyl, oxopiperazinylpyrimidinyl, carboxypiperazinylpyrimidinyl, carboxyethyl-
 10 piperazinylpyrimidinyl, *tert*-butoxycarbonylpiperazinylpyrimidinyl, tetrazolylmethyl-
 piperazinylpyrimidinyl, trioxohexahydro-[1,2,5]thiadiazolo[2,3-*a*]pyrazinylpyrimidinyl,
 morpholinylpyrimidinyl, dimethylmorpholinylpyrimidinyl, hydroxymethylmorpholinyl-
 pyrimidinyl, carboxymorpholinylpyrimidinyl, (carboxy)(methyl)morpholinylpyrimidinyl,
 carboxymethylmorpholinylpyrimidinyl, thiomorpholinylpyrimidinyl, dioxo-
 15 thiomorpholinylpyrimidinyl, carboxyazepanylpyrimidinyl, carboxyoxazepanyl-
 pyrimidinyl, oxodiazepanylpyrimidinyl, (oxodiazepanyl)(trifluoromethyl)pyrimidinyl,
 (oxodiazepanyl)(methoxy)pyrimidinyl, (methyl)(oxo)diazepanylpyrimidinyl, dioxo-
 thiadiazepanylpyrimidinyl, hydroxyoxetanylpyrazinyl, (carboxy)(methyl)piperidinyl-
 pyrazinyl, (ethoxycarbonyl)(methyl)piperidinylpyrazinyl, morpholinylmethylthienyl,
 20 morpholinylethylpyrazolyl, carboxy-3-azabicyclo[3.1.0]hexanylpyridinyl, carboxy-3-
 azabicyclo[3.1.0]hexanylpyridazinyl, carboxy-3-azabicyclo[3.1.0]hexanylpyrimidinyl,
 (carboxy)(methyl)-3-azabicyclo[3.1.0]hexanylpyrimidinyl, methoxycarbonyl-3-
 azabicyclo[3.1.0]hexanylpyrimidinyl, ethoxycarbonyl-3-azabicyclo[3.1.0]hexanyl-
 pyrimidinyl, 2-oxa-5-azabicyclo[2.2.1]heptanylpyrimidinyl, carboxy-2-oxa-5-azabicyclo-
 25 [2.2.1]heptanylpyrimidinyl, carboxy-3-azabicyclo[3.1.1]heptanylpyrimidinyl, carboxy-3-
 azabicyclo[4.1.0]heptanylpyridinyl, carboxy-3-azabicyclo[4.1.0]heptanylpyrimidinyl,
 methoxycarbonyl-3-azabicyclo[4.1.0]heptanylpyrimidinyl, ethoxycarbonyl-3-azabicyclo-
 [4.1.0]heptanylpyrimidinyl, (hydroxy)(methyl)(oxo)-2-oxabicyclo[2.2.2]octanyl-
 pyrimidinyl, carboxy-3-azabicyclo[3.2.1]octanylpyrimidinyl, methoxycarbonyl-3-
 30 azabicyclo[3.2.1]octanylpyrimidinyl, oxo-8-azabicyclo[3.2.1]octanylpyrimidinyl,
 ethoxycarbonylmethylidenyl-8-azabicyclo[3.2.1]octanylpyrimidinyl, 3-oxa-8-azabicyclo-
 [3.2.1]octanylpyrimidinyl, oxo-3,6-diazabicyclo[3.2.2]nonanylpyrimidinyl, carboxy-3-
 oxa-7-azabicyclo[3.3.1]nonanylpyrimidinyl, carboxy-5-azaspiro[2.3]hexanylpyrimidinyl,

(carboxy)(methyl)-5-azaspiro[2.3]hexanylpyrimidinyl, carboxy-5-azaspiro[2.4]heptanylpyrimidinyl, carboxy-2-azaspiro[3.3]heptanylpyrimidinyl, 2-oxa-6-azaspiro[3.3]heptanylpyrimidinyl, 2-oxa-6-azaspiro[3.4]octanylpyrimidinyl, 2-oxa-6-azaspiro[3.5]nonanylpyrimidinyl, 2-oxa-7-azaspiro[3.5]nonanylpyrimidinyl and (dioxo)(methyl)-2,4,8-triazaspiro[4.5]decanylpyrimidinyl.

Illustrative values of R^1 include aminosulphonylphenyl, carboxypiperidinylpyrimidinyl and morpholinylpyrimidinyl.

Typically, R^2 represents hydrogen, halogen, trifluoromethyl or $-OR^a$; or R^2 represents optionally substituted C_{1-6} alkyl.

10 Typical examples of optional substituents on R^2 include C_{2-6} alkoxy carbonyl.

Typical examples of particular substituents on R^2 include ethoxy carbonyl.

In a first embodiment, R^2 represents hydrogen. In a second embodiment, R^2 represents halogen. In one aspect of that embodiment, R^2 represents fluoro. In another aspect of that embodiment, R^2 represents chloro. In a third embodiment, R^2 represents trifluoromethyl. In a fourth embodiment, R^2 represents $-OR^a$. In a fifth embodiment, R^2 represents optionally substituted C_{1-6} alkyl. In one aspect of that embodiment, R^2 represents unsubstituted methyl. In another aspect of that embodiment, R^2 represents unsubstituted ethyl. In a further aspect of that embodiment, R^2 represents monosubstituted methyl or monosubstituted ethyl.

20 Typical values of R^2 include hydrogen, fluoro, chloro, trifluoromethyl, $-OR^a$, methyl and ethoxy carbonyl ethyl.

Typically, R^3 represents hydrogen, halogen or C_{1-6} alkyl.

In a first embodiment, R^3 represents hydrogen. In a second embodiment, R^3 represents halogen. In one aspect of that embodiment, R^3 represents fluoro. In a third embodiment, R^3 represents C_{1-6} alkyl. In one aspect of that embodiment, R^3 represents methyl. In another aspect of that embodiment, R^3 represents ethyl.

Suitably, R^4 represents hydrogen or methyl.

In a first embodiment, R^4 represents hydrogen. In a second embodiment, R^4 represents C_{1-6} alkyl, especially methyl.

30 Typical examples of suitable substituents on R^a , R^b , R^c , R^d or R^e , or on the heterocyclic moiety $-NR^bR^c$, include halogen, C_{1-6} alkyl, C_{1-6} alkoxy, difluoromethoxy, trifluoromethoxy, C_{1-6} alkoxy(C_{1-6})alkyl, C_{1-6} alkylthio, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, hydroxy, hydroxy(C_{1-6})alkyl, amino(C_{1-6})alkyl, cyano, trifluoromethyl,

oxo, C₂₋₆ alkylcarbonyl, carboxy, C₂₋₆ alkoxy carbonyl, C₂₋₆ alkylcarbonyloxy, amino, C₁₋₆ alkylamino, di(C₁₋₆)alkylamino, phenylamino, pyridinylamino, C₂₋₆ alkylcarbonylamino, C₂₋₆ alkylcarbonylamino(C₁₋₆)alkyl, C₂₋₆ alkoxy carbonylamino, C₁₋₆ alkylsulphonylamino, aminocarbonyl, C₁₋₆ alkylaminocarbonyl and di(C₁₋₆)alkylaminocarbonyl.

- 5 Typical examples of specific substituents on R^a, R^b, R^c, R^d or R^e, or on the heterocyclic moiety -NR^bR^c, include fluoro, chloro, bromo, methyl, ethyl, isopropyl, methoxy, isopropoxy, difluoromethoxy, trifluoromethoxy, methoxymethyl, methylthio, ethylthio, methylsulphanyl, methylsulphonyl, hydroxy, hydroxymethyl, hydroxyethyl, aminomethyl, cyano, trifluoromethyl, oxo, acetyl, carboxy, methoxycarbonyl, ethoxycarbonyl, *tert*-butoxycarbonyl, acetoxymethyl, amino, methylamino, ethylamino, dimethylamino, phenylamino, pyridinylamino, acetylamino, *tert*-butoxycarbonylamino, acetylaminomethyl, methylsulphonylamino, aminocarbonyl, methylaminocarbonyl and dimethylaminocarbonyl.

- 15 Suitably, R^a represents C₁₋₆ alkyl, aryl(C₁₋₆)alkyl or heteroaryl(C₁₋₆)alkyl, any of which groups may be optionally substituted by one or more substituents.

Selected values of R^a include methyl, ethyl, benzyl and isoindolylpropyl, any of which groups may be optionally substituted by one or more substituents.

Selected examples of suitable substituents on R^a include C₁₋₆ alkoxy and oxo.

Selected examples of specific substituents on R^a include methoxy and oxo.

- 20 In one embodiment, R^a represents optionally substituted C₁₋₆ alkyl. In one aspect of that embodiment, R^a ideally represents unsubstituted C₁₋₆ alkyl, especially methyl. In another aspect of that embodiment, R^a ideally represents substituted C₁₋₆ alkyl, e.g. methoxyethyl. In another embodiment, R^a represents optionally substituted aryl. In one aspect of that embodiment, R^a represents unsubstituted aryl, especially phenyl. In another aspect of that embodiment, R^a represents monosubstituted aryl, especially methylphenyl. In another embodiment, R^a represents optionally substituted aryl(C₁₋₆)alkyl, ideally unsubstituted aryl(C₁₋₆)alkyl, especially benzyl. In a further embodiment, R^a represents optionally substituted heteroaryl. In a further embodiment, R^a represents optionally substituted heteroaryl(C₁₋₆)alkyl, e.g. dioxoisindolylpropyl.

- 30 Specific values of R^a include methyl, methoxyethyl, benzyl and dioxoisindolylpropyl.

In a particular aspect, R^b represents hydrogen or trifluoromethyl; or C₁₋₆ alkyl, C₃₋₇ cycloalkyl, C₃₋₇ cycloalkyl(C₁₋₆)alkyl, aryl, aryl(C₁₋₆)alkyl, C₃₋₇ heterocycloalkyl, C₃₋₇

heterocycloalkyl(C₁₋₆)alkyl, heteroaryl or heteroaryl(C₁₋₆)alkyl, any of which groups may be optionally substituted by one or more substituents.

Selected values of R^b include hydrogen; or C₁₋₆ alkyl, aryl(C₁₋₆)alkyl, C₃₋₇ heterocycloalkyl or C₃₋₇ heterocycloalkyl(C₁₋₆)alkyl, any of which groups may be
 5 optionally substituted by one or more substituents.

Typical values of R^b include hydrogen and C₁₋₆ alkyl.

Illustratively, R^b represents hydrogen or trifluoromethyl; or methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, 2-methylpropyl, *tert*-butyl, pentyl, hexyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl,
 10 cyclohexylmethyl, phenyl, benzyl, phenylethyl, azetidiny, tetrahydrofuryl, tetrahydrothienyl, pyrrolidiny, piperidiny, homopiperidiny, morpholiny, azetidinylmethyl, tetrahydrofurylmethyl, pyrrolidinylmethyl, pyrrolidinylethyl, pyrrolidinylpropyl, thiazolidinylmethyl, imidazolidiny, piperidinylmethyl, piperidinylethyl, tetrahydroquinoliny, piperazinylpropyl, morpholiny, morpholinyethylethyl, morpholinypropyl, pyridiny, indolylmethyl, pyrazolylmethyl,
 15 pyrazolyethylethyl, imidazolylmethyl, imidazolylethylethyl, benzimidazolylmethyl, triazolylmethyl, pyridiny, pyridinyethylethyl, any of which groups may be optionally substituted by one or more substituents.

Representative values of R^b include hydrogen; or methyl, ethyl, *n*-propyl, benzyl, pyrrolidiny or morpholinypropyl, any of which groups may be optionally substituted by
 20 one or more substituents.

Selected examples of suitable substituents on R^b include C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkylsulphiny, C₁₋₆ alkylsulphony, hydroxy, cyano, C₂₋₆ alkoxy, carbonyl, di-(C₁₋₆)alkylamino and C₂₋₆ alkoxy, carbonylamino.

Selected examples of specific substituents on R^b include methoxy, methylthio, methylsulphiny, methylsulphony, hydroxy, cyano, *tert*-butoxy, carbonyl, dimethylamino and *tert*-butoxy, carbonylamino.
 25

Specific values of R^b include hydrogen, methyl, methoxyethyl, methylthioethyl, methylsulphinyethyl, methylsulphonyethyl, hydroxyethyl, cyanoethyl, dimethylaminoethyl, *tert*-butoxy, carbonylaminoethyl, dihydroxypropyl, benzyl, pyrrolidiny, *tert*-butoxy, carbonylpyrrolidiny and morpholinypropyl.
 30

In one embodiment, R^b represents hydrogen. In another embodiment, R^b represents C₁₋₆ alkyl, especially methyl.

Selected values of R^c include hydrogen; or C₁₋₆ alkyl, C₃₋₇ cycloalkyl or C₃₋₇ heterocycloalkyl, any of which groups may be optionally substituted by one or more substituents.

In a particular aspect, R^c represents hydrogen, C₁₋₆ alkyl or C₃₋₇ cycloalkyl.

5 Representative values of R^c include hydrogen; or methyl, cyclobutyl, cyclopentyl, cyclohexyl, tetrahydropyranyl and piperidinyl, any of which groups may be optionally substituted by one or more substituents.

Selected examples of suitable substituents on R^c include C₂₋₆ alkylcarbonyl and C₂₋₆ alkoxy carbonyl.

10 Selected examples of specific substituents on R^c include acetyl and *tert*-butoxycarbonyl.

Specific values of R^c include hydrogen, methyl, cyclobutyl, cyclopentyl, cyclohexyl, tetrahydropyranyl, acetylpiperidinyl and *tert*-butoxycarbonylpiperidinyl,

Suitably, R^c represents hydrogen or C₁₋₆ alkyl. In one embodiment, R^c is hydrogen.

15 In another embodiment, R^c represents C₁₋₆ alkyl, especially methyl or ethyl, particularly methyl. In a further embodiment, R^c represents C₃₋₇ cycloalkyl, e.g. cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.

Alternatively, the moiety -NR^bR^c may suitably represent azetidin-1-yl, pyrrolidin-1-yl, oxazolidin-3-yl, isoxazolidin-2-yl, thiazolidin-3-yl, isothiazolidin-2-yl, piperidin-1-yl, morpholin-4-yl, thiomorpholin-4-yl, piperazin-1-yl, homopiperidin-1-yl, homomorpholin-4-yl or homopiperazin-1-yl, any of which groups may be optionally substituted by one or more substituents.

25 Selected examples of suitable substituents on the heterocyclic moiety -NR^bR^c include C₁₋₆ alkyl, C₁₋₆ alkylsulphonyl, hydroxy, hydroxy(C₁₋₆)alkyl, amino(C₁₋₆)alkyl, cyano, oxo, C₂₋₆ alkylcarbonyl, carboxy, C₂₋₆ alkoxy carbonyl, amino, C₂₋₆ alkylcarbonyl-amino, C₂₋₆ alkylcarbonylamino(C₁₋₆)alkyl, C₂₋₆ alkoxy carbonylamino, C₁₋₆ alkyl-sulphonylamino and aminocarbonyl.

30 Selected examples of specific substituents on the heterocyclic moiety -NR^bR^c include methyl, methylsulphonyl, hydroxy, hydroxymethyl, aminomethyl, cyano, oxo, acetyl, carboxy, ethoxycarbonyl, amino, acetylamino, acetylaminomethyl, *tert*-butoxycarbonylamino, methylsulphonylamino and aminocarbonyl.

Specific values of the moiety -NR^bR^c include azetidin-1-yl, hydroxyazetidin-1-yl, hydroxymethylazetidin-1-yl, (hydroxy)(hydroxymethyl)azetidin-1-yl, aminomethyl-

azetidin-1-yl, cyanoazetidin-1-yl, carboxyazetidin-1-yl, aminoazetidin-1-yl, aminocarbonylazetidin-1-yl, pyrrolidin-1-yl, aminomethylpyrrolidin-1-yl, oxopyrrolidin-1-yl, acetylaminoethylpyrrolidin-1-yl, *tert*-butoxycarbonylaminopyrrolidin-1-yl, oxo-oxazolidin-3-yl, hydroxyisoxazolidin-2-yl, thiazolidin-3-yl, oxothiazolidin-3-yl, dioxo-isothiazolidin-2-yl, piperidin-1-yl, hydroxypiperidin-1-yl, hydroxymethylpiperidin-1-yl, aminopiperidin-1-yl, acetylaminoethylpiperidin-1-yl, *tert*-butoxycarbonylaminopiperidin-1-yl, methylsulphonylaminopiperidin-1-yl, morpholin-4-yl, piperazin-1-yl, methylpiperazin-1-yl, methylsulphonylpiperazin-1-yl, oxopiperazin-1-yl, acetylpiperazin-1-yl, ethoxycarbonylpiperazin-1-yl and oxohomopiperazin-1-yl.

10 Suitably, R^d represents hydrogen; or C₁₋₆ alkyl, aryl or heteroaryl, any of which groups may be optionally substituted by one or more substituents.

Selected examples of suitable values for R^d include hydrogen, methyl, ethyl, isopropyl, 2-methylpropyl, *tert*-butyl, cyclopropyl, cyclobutyl, phenyl, thiazolidinyl, thienyl, imidazolyl and thiazolyl, any of which groups may be optionally substituted by
15 one or more substituents.

Selected examples of suitable substituents on R^d include halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, oxo, C₂₋₆ alkylcarbonyloxy and di(C₁₋₆)alkylamino.

Selected examples of particular substituents on R^d include fluoro, methyl, methoxy, oxo, acetoxy and dimethylamino.

20 In one embodiment, R^d represents hydrogen. In another embodiment, R^d represents optionally substituted C₁₋₆ alkyl. In one aspect of that embodiment, R^d ideally represents unsubstituted C₁₋₆ alkyl, e.g. methyl, ethyl, isopropyl, 2-methylpropyl or *tert*-butyl, especially methyl. In another aspect of that embodiment, R^d ideally represents substituted C₁₋₆ alkyl, e.g. substituted methyl or substituted ethyl, including
25 acetoxymethyl, dimethylaminomethyl and trifluoroethyl. In another embodiment, R^d represents optionally substituted aryl. In one aspect of that embodiment, R^d represents unsubstituted aryl, especially phenyl. In another aspect of that embodiment, R^d represents monosubstituted aryl, especially methylphenyl. In a further aspect of that embodiment, R^d represents disubstituted aryl, e.g. dimethoxyphenyl. In a further embodiment, R^d
30 represents optionally substituted heteroaryl, e.g. thienyl, chlorothienyl, methylthienyl, methylimidazolyl or thiazolyl. In another embodiment, R^d represents optionally substituted C₃₋₇ cycloalkyl, e.g. cyclopropyl or cyclobutyl. In a further embodiment, R^d

represents optionally substituted C₃₋₇ heterocycloalkyl, e.g. thiazolidinyl or oxo-thiazolidinyl.

Selected examples of specific values for R^d include hydrogen, methyl, acetoxy-methyl, dimethylaminomethyl, ethyl, trifluoroethyl, isopropyl, 2-methylpropyl, *tert*-butyl,
 5 cyclopropyl, cyclobutyl, phenyl, dimethoxyphenyl, thiazolidinyl, oxothiazolidinyl, thienyl, chlorothienyl, methylthienyl, methylimidazolyl and thiazolyl.


Suitably, R^e represents C₁₋₆ alkyl or aryl, either of which groups may be optionally substituted by one or more substituents.

Selected examples of suitable substituents on R^e include C₁₋₆ alkyl, especially
 10 methyl.

In one embodiment, R^e represents optionally substituted C₁₋₆ alkyl, ideally unsubstituted C₁₋₆ alkyl, e.g. methyl or propyl, especially methyl. In another embodiment, R^e represents optionally substituted aryl. In one aspect of that embodiment, R^e represents unsubstituted aryl, especially phenyl. In another aspect of that embodiment, R^e represents
 15 monosubstituted aryl, especially methylphenyl. In a further embodiment, R^e represents optionally substituted heteroaryl.

Selected values of R^e include methyl, propyl and methylphenyl.

One sub-class of compounds according to the invention is represented by the compounds of formula (IIA) and *N*-oxides thereof, and pharmaceutically acceptable salts
 20 and solvates thereof, and glucuronide derivatives thereof, and co-crystals thereof:



(IIA)

wherein

25 R¹¹ represents halogen or cyano; or R¹¹ represents C₁₋₆ alkyl, C₂₋₆ alkynyl, aryl, C₃₋₇ heterocycloalkyl, C₃₋₇ heterocycloalkenyl, heteroaryl, (C₃₋₇)heterocycloalkyl-(C₁₋₆)alkyl-aryl-, heteroaryl(C₃₋₇)heterocycloalkyl-, (C₃₋₇)cycloalkyl-heteroaryl-, (C₃₋₇)cycloalkyl(C₁₋₆)alkyl-heteroaryl-, (C₄₋₇)cycloalkenyl-heteroaryl-, (C₄₋₉)bicycloalkyl-

heteroaryl-, (C₃₋₇)heterocycloalkyl-heteroaryl-, (C₃₋₇)heterocycloalkyl(C₁₋₆)alkyl-heteroaryl-, (C₃₋₇)heterocycloalkenyl-heteroaryl-, (C₄₋₉)heterobicycloalkyl-heteroaryl- or (C₄₋₉)spiroheterocycloalkyl-heteroaryl-, any of which groups may be optionally substituted by one or more substituents;

5 R¹² represents represents hydrogen, halogen, trifluoromethyl or optionally substituted C₁₋₆ alkyl;

R¹⁵ and R¹⁶ independently represent hydrogen, halogen, cyano, nitro, C₁₋₆ alkyl, trifluoromethyl, hydroxy, C₁₋₆ alkoxy, difluoromethoxy, trifluoromethoxy, C₁₋₆ alkylthio, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, amino, C₁₋₆ alkylamino, di(C₁₋₆)alkylamino, 10 arylamino, C₂₋₆ alkylcarbonylamino, C₁₋₆ alkylsulfonylamino, formyl, C₂₋₆ alkylcarbonyl, C₃₋₆ cycloalkylcarbonyl, C₃₋₆ heterocycloalkylcarbonyl, carboxy, C₂₋₆ alkoxycarbonyl, aminocarbonyl, C₁₋₆ alkylaminocarbonyl, di(C₁₋₆)alkylaminocarbonyl, aminosulfonyl, C₁₋₆ alkylaminosulfonyl or di(C₁₋₆)alkylaminosulfonyl; and

E is as defined above.

15 Examples of optional substituents which may be present on R¹¹ include one, two or three substituents independently selected from halogen, halo(C₁₋₆)alkyl, cyano, cyano(C₁₋₆)alkyl, nitro, nitro(C₁₋₆)alkyl, C₁₋₆ alkyl, difluoromethyl, trifluoromethyl, difluoroethyl, trifluoroethyl, C₂₋₆ alkenyl, hydroxy, hydroxy(C₁₋₆)alkyl, C₁₋₆ alkoxy, difluoromethoxy, trifluoromethoxy, trifluoroethoxy, carboxy(C₃₋₇)cycloalkyloxy, C₁₋₃ 20 alkylenedioxy, C₁₋₆ alkoxy(C₁₋₆)alkyl, C₁₋₆ alkylthio, C₁₋₆ alkylsulphinyl, C₁₋₆ alkylsulphonyl, (C₁₋₆)alkylsulphonyl(C₁₋₆)alkyl, oxo, amino, amino(C₁₋₆)alkyl, C₁₋₆ alkylamino, di(C₁₋₆)alkylamino, hydroxy(C₁₋₆)alkylamino, C₁₋₆ alkoxyamino, (C₁₋₆)alkoxy(C₁₋₆)alkylamino, [(C₁₋₆)alkoxy](hydroxy)(C₁₋₆)alkylamino, [(C₁₋₆)alkylthio](hydroxy)(C₁₋₆)alkylamino, *N*-[(C₁₋₆)alkyl]-*N*-[hydroxy(C₁₋₆)alkyl]amino, di(C₁₋₆)alkylamino(C₁₋₆)alkylamino, 25 *N*-[di(C₁₋₆)alkylamino(C₁₋₆)alkyl]-*N*-[hydroxy(C₁₋₆)alkyl]amino, hydroxy(C₁₋₆)alkyl-(C₃₋₇)cycloalkylamino, (hydroxy)[(C₃₋₇)cycloalkyl(C₁₋₆)alkyl]amino, (C₃₋₇)heterocycloalkyl(C₁₋₆)alkylamino, oxo(C₃₋₇)heterocycloalkyl(C₁₋₆)alkylamino, (C₁₋₆)alkylheteroaryl-amino, heteroaryl(C₁₋₆)alkylamino, (C₁₋₆)alkylheteroaryl(C₁₋₆)alkylamino, C₂₋₆ alkylcarbonylamino, *N*-[(C₁₋₆)alkyl]-*N*-[(C₂₋₆)alkylcarbonyl]amino, (C₂₋₆)alkylcarbonylamino(C₁₋₆)alkyl, C₃₋₆ alkenylcarbonylamino, bis[(C₃₋₆)alkenylcarbonyl]amino, *N*-[(C₁₋₆)alkyl]-*N*-[(C₃₋₇)cycloalkylcarbonyl]amino, C₂₋₆ alkoxycarbonylamino, C₂₋₆ 30 alkoxycarbonyl(C₁₋₆)alkylamino, C₁₋₆ alkylaminocarbonylamino, C₁₋₆ alkylsulphonyl-amino, *N*-[(C₁₋₆)alkyl]-*N*-[(C₁₋₆)alkylsulphonyl]amino, bis[(C₁₋₆)alkylsulphonyl]amino, *N*-

- [(C₁₋₆)alkyl]-*N*-[carboxy(C₁₋₆)alkyl]amino, carboxy(C₃₋₇)cycloalkylamino, carboxy-
 (C₃₋₇)cycloalkyl(C₁₋₆)alkylamino, formyl, C₂₋₆ alkylcarbonyl, (C₃₋₇)cycloalkylcarbonyl,
 phenylcarbonyl, (C₂₋₆)alkylcarbonyloxy(C₁₋₆)alkyl, carboxy, carboxy(C₁₋₆)alkyl, C₂₋₆
 alkoxycarbonyl, C₂₋₆ alkoxycarbonyl(C₁₋₆)alkyl, morpholinyl(C₁₋₆)alkoxycarbonyl, C₂₋₆
 5 alkoxycarbonylmethylidenyl, a carboxylic acid isostere or prodrug moiety Ω as defined
 herein, -(C₁₋₆)alkyl- Ω , aminocarbonyl, C₁₋₆ alkylaminocarbonyl, hydroxy(C₁₋₆)alkylamino-
 carbonyl, di(C₁₋₆)alkylaminocarbonyl, aminocarbonyl(C₁₋₆)alkyl, aminosulphonyl,
 di(C₁₋₆)alkylaminosulphonyl, (C₁₋₆)alkylsulphoximinyl and [(C₁₋₆)alkyl][*N*-(C₁₋₆)alkyl]-
 sulphoximinyl.
- 10 Examples of particular substituents on R¹¹ include fluoro, chloro, bromo,
 fluoromethyl, fluoroisopropyl, cyano, cyanoethyl, nitro, nitromethyl, methyl, ethyl,
 isopropyl, isobutyl, *tert*-butyl, difluoromethyl, trifluoromethyl, difluoroethyl, trifluoro-
 ethyl, ethenyl, hydroxy, hydroxymethyl, hydroxyisopropyl, methoxy, isopropoxy,
 difluoromethoxy, trifluoromethoxy, trifluoroethoxy, carboxycyclobutylloxy, methylene-
 15 dioxy, ethylenedioxy, methoxymethyl, methoxyethyl, methylthio, methylsulphinyl,
 methylsulphonyl, methylsulphonylethyl, oxo, amino, aminomethyl, aminoisopropyl,
 methylamino, ethylamino, dimethylamino, hydroxyethylamino, hydroxypropylamino,
 (hydroxy)(methyl)propylamino, methoxyamino, methoxyethylamino, (hydroxy)-
 (methoxy)(methyl)propylamino, (hydroxy)(methylthio)butylamino, *N*-(hydroxyethyl)-*N*-
 20 (methyl)amino, dimethylaminoethylamino, (dimethylamino)(methyl)propylamino, *N*-
 (dimethylaminoethyl)-*N*-(hydroxyethyl)amino, hydroxymethylcyclopentylamino,
 hydroxycyclobutylmethylamino, (cyclopropyl)(hydroxy)propylamino, morpholinylethyl-
 amino, oxopyrrolidinylmethylamino, ethyloxadiazolylamino, methylthiadiazolylamino,
 thiazolylmethylamino, thiazolylethylamino, pyrimidinylmethylamino, methylpyrazolyl-
 25 methylamino, acetylaminomethyl, ethenylcarbonylamino, bis(ethenylcarbonyl)amino, *N*-
 cyclopropylcarbonyl-*N*-methylamino, methoxycarbonylamino, ethoxycarbonylamino, *tert*-
 butoxycarbonylamino, methoxycarbonylethylamino, ethylaminocarbonylamino,
 butylaminocarbonylamino, methylsulphonylamino, *N*-methyl-*N*-(methylsulphonyl)amino,
 30 bis(methylsulphonyl)amino, *N*-(carboxymethyl)-*N*-methylamino, *N*-(carboxyethyl)-*N*-
 methylamino, carboxycyclopentylamino, carboxycyclopropylmethylamino, formyl, acetyl,
 isopropylcarbonyl, cyclobutylcarbonyl, phenylcarbonyl, acetoxisopropyl, carboxy,
 carboxymethyl, carboxyethyl, methoxycarbonyl, ethoxycarbonyl, *n*-butoxycarbonyl, *tert*-

butoxycarbonyl, methoxycarbonylmethyl, ethoxycarbonylmethyl, ethoxycarbonylethyl, morpholinylethoxycarbonyl, ethoxycarbonylmethylidenyl, methylsulphonylamino-carbonyl, acetylaminosulphonyl, methoxyaminocarbonyl, tetrazolyl, tetrazolylmethyl, hydroxyoxadiazolyl, aminocarbonyl, methylaminocarbonyl, hydroxyethylaminocarbonyl, 5 dimethylaminocarbonyl, aminocarbonylmethyl, aminosulphonyl, methylaminosulphonyl, dimethylaminosulphonyl, methylsulphoximinyl and (methyl)(*N*-methyl)sulphoximinyl.

Generally, R¹¹ represents C₁₋₆ alkyl, C₂₋₆ alkynyl, aryl, C₃₋₇ heterocycloalkyl, C₃₋₇ heterocycloalkenyl, heteroaryl, (C₃₋₇)heterocycloalkyl(C₁₋₆)alkyl-aryl-, heteroaryl-(C₃₋₇)heterocycloalkyl-, (C₃₋₇)cycloalkyl-heteroaryl-, (C₃₋₇)cycloalkyl(C₁₋₆)alkyl-10 heteroaryl-, (C₄₋₇)cycloalkenyl-heteroaryl-, (C₄₋₉) bicycloalkyl-heteroaryl-, (C₃₋₇)heterocycloalkyl-heteroaryl-, (C₃₋₇)heterocycloalkyl(C₁₋₆)alkyl-heteroaryl-, (C₃₋₇)heterocycloalkenyl-heteroaryl-, (C₄₋₉)heterobicycloalkyl-heteroaryl- or (C₄₋₉)spiroheterocycloalkyl-heteroaryl-, any of which groups may be optionally substituted by one or more substituents.

15 More generally, R¹¹ represents aryl or (C₃₋₇)heterocycloalkyl-heteroaryl-, either of which groups may be optionally substituted by one or more substituents.

In a first embodiment, R¹¹ represents halogen. In one aspect of that embodiment, R¹¹ represents bromo.

In a second embodiment, R¹¹ represents cyano.

20 In a third embodiment, R¹¹ represents optionally substituted C₁₋₆ alkyl. In one aspect of that embodiment, R¹¹ represents optionally substituted ethyl.

In a fourth embodiment, R¹¹ represents optionally substituted C₂₋₆ alkynyl. In one aspect of that embodiment, R¹¹ represents optionally substituted butynyl.

25 In a fifth embodiment, R¹¹ represents optionally substituted aryl. In one aspect of that embodiment, R¹¹ represents optionally substituted phenyl.

In a sixth embodiment, R¹¹ represents optionally substituted C₃₋₇ heterocycloalkyl.

In a seventh embodiment, R¹¹ represents optionally substituted C₃₋₇ heterocycloalkenyl.

30 In an eighth embodiment, R¹¹ represents optionally substituted heteroaryl. In selected aspects of that embodiment, R¹¹ represents benzofuryl, thienyl, indolyl, pyrazolyl, indazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridinyl, quinolinyl, pyridazinyl, pyrimidinyl or pyrazinyl, any of which groups may be optionally substituted by one or more substituents.

In a ninth embodiment, R¹¹ represents optionally substituted (C₃₋₇)-heterocycloalkyl(C₁₋₆)alkyl-aryl-. In a first aspect of that embodiment, R¹¹ represents optionally substituted pyrrolidinylmethylphenyl-. In a second aspect of that embodiment, R¹¹ represents optionally substituted piperazinylmethylphenyl-.

5 In a tenth embodiment, R¹¹ represents optionally substituted heteroaryl(C₃₋₇)-heterocycloalkyl-. In one aspect of that embodiment, R¹¹ represents optionally substituted pyridinylpiperazinyl-.

In an eleventh embodiment, R¹¹ represents optionally substituted (C₃₋₇)cycloalkyl-heteroaryl-. In a first aspect of that embodiment, R¹¹ represents optionally substituted
10 cyclohexylpyrazolyl-. In a second aspect of that embodiment, R¹¹ represents optionally substituted cyclohexylpyridinyl-. In a third aspect of that embodiment, R¹¹ represents optionally substituted cyclopropylpyrimidinyl-. In a fourth aspect of that embodiment, R¹¹ represents optionally substituted cyclobutylpyrimidinyl-. In a fifth aspect of that
15 aspect of that embodiment, R¹¹ represents optionally substituted cyclohexylpyrimidinyl-. In a seventh aspect of that embodiment, R¹¹ represents optionally substituted cyclohexylpyrazinyl-.

In a twelfth embodiment, R¹¹ represents optionally substituted (C₄₋₇)cycloalkenyl-heteroaryl-.

20 In a thirteenth embodiment, R¹¹ represents optionally substituted (C₃₋₇)-heterocycloalkyl-heteroaryl-. In a first aspect of that embodiment, R¹¹ represents optionally substituted pyrrolidinylpyridinyl-. In a second aspect of that embodiment, R¹¹ represents optionally substituted tetrahydropyranylpyridinyl-. In a third aspect of that
25 embodiment, R¹¹ represents optionally substituted piperidinylpyridinyl-. In a fourth aspect of that embodiment, R¹¹ represents optionally substituted piperazinylpyridinyl-. In a fifth aspect of that embodiment, R¹¹ represents optionally substituted morpholinylpyridinyl-. In a sixth aspect of that embodiment, R¹¹ represents optionally substituted thiomorpholinylpyridinyl-. In a seventh aspect of that embodiment, R¹¹ represents
30 optionally substituted diazepanylpyridinyl-. In an eighth aspect of that embodiment, R¹¹ represents optionally substituted oxetanylpyrimidinyl-. In a ninth aspect of that embodiment, R¹¹ represents optionally substituted azetidanylpyrimidinyl-. In a tenth aspect of that embodiment, R¹¹ represents optionally substituted tetrahydrofuranylpyrimidinyl-. In an eleventh aspect of that embodiment, R¹¹ represents optionally

substituted pyrrolidinylpyrimidinyl-. In a twelfth aspect of that embodiment, R¹¹ represents optionally substituted tetrahydropyranylpyrimidinyl-. In a thirteenth aspect of that embodiment, R¹¹ represents optionally substituted piperidinylpyrimidinyl-. In a fourteenth aspect of that embodiment, R¹¹ represents optionally substituted piperazinylpyrimidinyl-. In a fifteenth aspect of that embodiment, R¹¹ represents optionally substituted morpholinylpyrimidinyl-. In a sixteenth aspect of that embodiment, R¹¹ represents optionally substituted thiomorpholinylpyrimidinyl-. In a seventeenth aspect of that embodiment, R¹¹ represents optionally substituted azepanylpyrimidinyl-. In an eighteenth aspect of that embodiment, R¹¹ represents optionally substituted oxazepanylpyrimidinyl-. In a nineteenth aspect of that embodiment, R¹¹ represents optionally substituted diazepanylpyrimidinyl-. In a twentieth aspect of that embodiment, R¹¹ represents optionally substituted thiadiazepanylpyrimidinyl-. In a twenty-first aspect of that embodiment, R¹¹ represents optionally substituted oxetanylpyrazinyl-. In a twenty-second aspect of that embodiment, R¹¹ represents optionally substituted piperidinylpyrazinyl-.

In a fourteenth embodiment, R¹¹ represents optionally substituted (C₃₋₇)-heterocycloalkyl(C₁₋₆)alkyl-heteroaryl-. In a first aspect of that embodiment, R¹¹ represents optionally substituted morpholinylmethylthienyl-. In a second aspect of that embodiment, R¹¹ represents optionally substituted morpholinylethylpyrazolyl-.

In a fifteenth embodiment, R¹¹ represents optionally substituted (C₃₋₇)-heterocycloalkenyl-heteroaryl-.

In a sixteenth embodiment, R¹¹ represents optionally substituted (C₄₋₉)-heterobicycloalkyl-heteroaryl-.

In a seventeenth embodiment, R¹¹ represents optionally substituted (C₄₋₉)-spiroheterocycloalkyl-heteroaryl-.

In an eighteenth embodiment, R¹¹ represents optionally substituted (C₃₋₇)-cycloalkyl(C₁₋₆)alkyl-heteroaryl-. In one aspect of that embodiment, R¹¹ represents optionally substituted cyclohexylmethylpyrimidinyl-.

In a nineteenth embodiment, R¹¹ represents optionally substituted (C₄₋₉)-bicycloalkyl-heteroaryl-.

Appositely, R¹¹ represents bromo or cyano; or R¹¹ represents ethyl, butynyl, phenyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, 1,2,3,6-tetrahydropyridinyl, benzofuryl, thienyl, indolyl, pyrazolyl, indazolyl, isoxazolyl, thiazolyl, imidazolyl,

pyridinyl, quinolinyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrrolidinylmethylphenyl, piperazinylmethylphenyl, pyridinylpiperazinyl, cyclohexylpyrazolyl, cyclohexylpyridinyl, cyclopropylpyrimidinyl, cyclobutylpyrimidinyl, cyclopentylpyrimidinyl, cyclohexylpyrimidinyl, cyclohexylpyrazinyl, cyclohexylmethylpyrimidinyl, cyclohexenylpyridinyl, cyclohexenylpyrimidinyl, bicyclo[3.1.0]hexanylpyridinyl, bicyclo[3.1.0]hexanylpyrimidinyl, bicyclo[4.1.0]heptanylpyrimidinyl, bicyclo[2.2.2]octanylpyrimidinyl, pyrrolidinylpyridinyl, tetrahydropyranylpyridinyl, piperidinylpyridinyl, piperazinylpyridinyl, morpholinylpyridinyl, thiomorpholinylpyridinyl, diazepanylpyridinyl, oxetanylpyrimidinyl, azetidinylpyrimidinyl, tetrahydrofuranylpyrimidinyl, pyrrolidinylpyrimidinyl, tetrahydropyranylpyrimidinyl, piperidinylpyrimidinyl, piperazinylpyrimidinyl, hexahydro-[1,2,5]thiadiazolo[2,3-*a*]pyrazinylpyrimidinyl, morpholinylpyrimidinyl, thiomorpholinylpyrimidinyl, azepanylpyrimidinyl, oxazepanylpyrimidinyl, diazepanylpyrimidinyl, thiadiazepanylpyrimidinyl, oxetanylpyrazinyl, piperidinylpyrazinyl, morpholinylmethylthienyl, morpholinylethylpyrazolyl, 3-azabicyclo[3.1.0]hexanylpyridinyl, 3-azabicyclo[3.1.0]hexanylpyridazinyl, 3-azabicyclo[3.1.0]hexanylpyrimidinyl, 2-oxa-5-azabicyclo[2.2.1]heptanylpyrimidinyl, 3-azabicyclo[3.1.1]heptanylpyrimidinyl, 3-azabicyclo[4.1.0]heptanylpyridinyl, 3-azabicyclo[4.1.0]heptanylpyrimidinyl, 2-oxabicyclo[2.2.2]octanylpyrimidinyl, 3-azabicyclo[3.2.1]octanylpyrimidinyl, 8-azabicyclo[3.2.1]octanylpyrimidinyl, 3-oxa-8-azabicyclo[3.2.1]octanylpyrimidinyl, 3,6-diazabicyclo[3.2.2]nonanylpyrimidinyl, 3-oxa-7-azabicyclo[3.3.1]nonanylpyrimidinyl, 5-azaspiro[2.3]hexanylpyrimidinyl, 5-azaspiro[2.4]heptanylpyrimidinyl, 2-azaspiro[3.3]heptanylpyrimidinyl, 2-oxa-6-azaspiro[3.3]heptanylpyrimidinyl, 2-oxa-6-azaspiro[3.4]octanylpyrimidinyl, 2-oxa-6-azaspiro[3.5]nonanylpyrimidinyl, 2-oxa-7-azaspiro[3.5]nonanylpyrimidinyl or 2,4,8-triazaspiro[4.5]decanylpyrimidinyl, any of which groups may be optionally substituted by one or more substituents.

Illustratively, R¹¹ represents phenyl, piperidinylpyrimidinyl or morpholinylpyrimidinyl, any of which groups may be optionally substituted by one or more substituents.

Typical examples of optional substituents on R¹¹ include one, two or three substituents independently selected from halogen, halo(C₁₋₆)alkyl, cyano, cyano(C₁₋₆)alkyl, nitro(C₁₋₆)alkyl, C₁₋₆ alkyl, trifluoromethyl, trifluoroethyl, C₂₋₆ alkenyl, hydroxy, hydroxy(C₁₋₆)alkyl, C₁₋₆ alkoxy, trifluoroethoxy, carboxy(C₃₋₇)cycloalkyloxy, C₁₋₆

alkylthio, C₁₋₆ alkylsulphonyl, (C₁₋₆)alkylsulphonyl(C₁₋₆)alkyl, oxo, amino, amino-(C₁₋₆)alkyl, C₁₋₆ alkylamino, di(C₁₋₆)alkylamino, (C₁₋₆)alkoxy(C₁₋₆)alkylamino, *N*-[(C₁₋₆)alkyl]-*N*-[hydroxy(C₁₋₆)alkyl]amino, (C₂₋₆)alkylcarbonylamino(C₁₋₆)alkyl, C₁₋₆ alkylsulphonylamino, *N*-[(C₁₋₆)alkyl]-*N*-[(C₁₋₆)alkylsulphonyl]amino, bis[(C₁₋₆)alkylsulphonyl]amino, *N*-[(C₁₋₆)alkyl]-*N*-[carboxy(C₁₋₆)alkyl]amino, carboxy(C₃₋₇)cycloalkylamino, carboxy(C₃₋₇)cycloalkyl(C₁₋₆)alkylamino, formyl, C₂₋₆ alkylcarbonyl, (C₂₋₆)alkylcarbonyloxy(C₁₋₆)alkyl, carboxy, carboxy(C₁₋₆)alkyl, C₂₋₆ alkoxycarbonyl, C₂₋₆ alkoxycarbonyl(C₁₋₆)alkyl, morpholinyl(C₁₋₆)alkoxycarbonyl, C₂₋₆ alkoxycarbonylmethylidenyl, a carboxylic acid isostere or prodrug moiety Ω as defined herein,

10 -(C₁₋₆)alkyl- Ω , aminocarbonyl, aminosulphonyl, (C₁₋₆)alkylsulphoximinyl and [(C₁₋₆)alkyl][*N*-(C₁₋₆)alkyl]sulphoximinyl.

Suitable examples of optional substituents on R¹¹ include one, two or three substituents independently selected from carboxy and aminosulphonyl.

Typical examples of particular substituents on R¹¹ include one, two or three

15 substituents independently selected from fluoro, chloro, fluoromethyl, fluoroisopropyl, cyano, cyanoethyl, nitromethyl, methyl, ethyl, isopropyl, trifluoromethyl, trifluoroethyl, ethenyl, hydroxy, hydroxymethyl, hydroxyisopropyl, methoxy, isopropoxy, trifluoroethoxy, carboxycyclobutyloxy, methylthio, methylsulphonyl, methylsulphonylethyl, oxo, amino, aminomethyl, aminoisopropyl, methylamino, dimethylamino, methoxyethylamino,

20 *N*-(hydroxyethyl)-*N*-(methyl)amino, acetylaminomethyl, methylsulphonylamino, *N*-methyl-*N*-(methylsulphonyl)amino, bis(methylsulphonyl)amino, *N*-(carboxyethyl)-*N*-(methyl)amino, carboxycyclopentylamino, carboxycyclopropylmethylamino, formyl, acetyl, acetoxisopropyl, carboxy, carboxymethyl, carboxyethyl, methoxycarbonyl, ethoxycarbonyl, *n*-butoxycarbonyl, *tert*-butoxycarbonyl, methoxycarbonylmethyl, ethoxycarbonylmethyl, ethoxycarbonylethyl, morpholinylethoxycarbonyl, ethoxycarbonylmethylidenyl, methylsulphonylaminocarbonyl, acetylaminosulphonyl, methoxyaminocarbonyl, tetrazolyl, tetrazolylmethyl, hydroxyoxadiazolyl, aminocarbonyl, aminosulphonyl, methylsulphoximinyl and (methyl)(*N*-methyl)sulphoximinyl.

25

Suitable examples of particular substituents on R¹¹ include one, two or three

30 substituents independently selected from carboxy and aminosulphonyl.

In a particular embodiment, R¹¹ is substituted by hydroxy(C₁₋₆)alkyl. In one aspect of that embodiment, R¹¹ is substituted by hydroxyisopropyl, especially 2-hydroxyprop-2-yl.

Selected values of R¹¹ include bromo, cyano, methoxycarbonylethyl, ethoxycarbonylethyl, hydroxybutynyl, chlorophenyl, hydroxyphenyl, methylsulphonylphenyl, aminomethylphenyl, aminoisopropylphenyl, acetylaminomethylphenyl, acetylphenyl, methoxycarbonylphenyl, aminocarbonylphenyl, aminosulphonylphenyl, 5 acetylaminosulphonylphenyl, (methoxycarbonyl)(methyl)pyrrolidinyl, oxopiperidinyl, ethoxycarbonylpiperidinyl, methylsulphonylpiperazinyl, morpholinyl, methylsulphonyl-1,2,3,6-tetrahydropyridinyl, acetyl-1,2,3,6-tetrahydropyridinyl, *tert*-butoxycarbonyl-1,2,3,6-tetrahydropyridinyl, methoxycarbonylmethyl-1,2,3,6-tetrahydropyridinyl, benzofuryl, thienyl, indolyl, pyrazolyl, methylpyrazolyl, dimethylpyrazolyl, (methyl)[*N*-10 methyl-*N*-(methylsulfonyl)amino]pyrazolyl, methylindazolyl, dimethylisoxazolyl, hydroxyisopropylthiazolyl, methylimidazolyl, dimethylimidazolyl, pyridinyl, fluoro-pyridinyl, cyanopyridinyl, methylpyridinyl, (cyano)(methyl)pyridinyl, dimethylpyridinyl, trifluoromethylpyridinyl, ethenylpyridinyl, hydroxyisopropylpyridinyl, methoxypyridinyl, (methoxy)(methyl)pyridinyl, isopropoxypyridinyl, trifluoroethoxypyridinyl, (methyl)-15 (trifluoroethoxy)pyridinyl, methylsulphonylpyridinyl, oxopyridinyl, (methyl)(oxo)-pyridinyl, (dimethyl)(oxo)pyridinyl, aminopyridinyl, methylaminopyridinyl, dimethylaminopyridinyl, methoxyethylaminopyridinyl, *N*-(hydroxyethyl)-*N*-(methyl)amino-pyridinyl, methylsulphonylaminopyridinyl, [bis(methylsulphonyl)amino]pyridinyl, carboxypyridinyl, quinolinyl, hydroxypyridazinyl, pyrimidinyl, fluoroisopropyl-20 pyrimidinyl, hydroxyisopropylpyrimidinyl, methoxypyrimidinyl, carboxycyclobutyloxy-pyrimidinyl, methylthiopyrimidinyl, methylsulphonylpyrimidinyl, oxopyrimidinyl, aminopyrimidinyl, dimethylaminopyrimidinyl, methoxyethylaminopyrimidinyl, *N*-(carboxyethyl)-*N*-(methyl)aminopyrimidinyl, carboxycyclopentylaminopyrimidinyl, carboxycyclopropylmethylaminopyrimidinyl, acetoxisopropylpyrimidinyl, 25 ethoxycarbonylethylpyrimidinyl, hydroxypyrazinyl, hydroxyisopropylpyrazinyl, pyrrolidinylmethylphenyl, piperazinylmethylphenyl, pyridinylpiperazinyl, carboxy-cyclohexylpyrazolyl, carboxycyclohexylpyridinyl, fluoromethylcyclopropylpyrimidinyl, acetylaminomethylcyclopropylpyrimidinyl, hydroxycyclobutylpyrimidinyl, carboxy-cyclopentylpyrimidinyl, carboxycyclohexylpyrimidinyl, (carboxy)(methyl)cyclohexyl-30 pyrimidinyl, (carboxy)(hydroxy)cyclohexylpyrimidinyl, carboxymethylcyclohexylpyrimidinyl, ethoxycarbonylcyclohexylpyrimidinyl, (methoxycarbonyl)(methyl)-cyclohexylpyrimidinyl, (ethoxycarbonyl)(methyl)cyclohexylpyrimidinyl, carboxy-cyclohexylpyrazinyl, carboxycyclohexylmethylpyrimidinyl, carboxycyclohexenyl-

- pyridinyl, carboxycyclohexenylpyrimidinyl, ethoxycarbonylcyclohexenylpyrimidinyl, carboxybicyclo[3.1.0]hexanylpyridinyl, carboxybicyclo[3.1.0]hexanylpyrimidinyl, ethoxycarbonylbicyclo[3.1.0]hexanylpyrimidinyl, carboxybicyclo[4.1.0]heptanylpyrimidinyl, carboxybicyclo[2.2.2]octanylpyrimidinyl, pyrrolidinylpyridinyl,
- 5 hydroxypyrrolidinylpyridinyl, hydroxytetrahydropyranylpyridinyl, piperidinylpyridinyl, acetylpiperidinylpyridinyl, (carboxy)(methyl)piperidinylpyridinyl, [(carboxy)(methyl)piperidinyl](fluoro)pyridinyl, [(carboxy)(methyl)piperidinyl](chloro)pyridinyl, piperazinylpyridinyl, (methyl)(piperazinyl)pyridinyl, cyanoethylpiperazinylpyridinyl, trifluoroethylpiperazinylpyridinyl, methylsulphonylpiperazinylpyridinyl, methyl-
- 10 sulphonylethylpiperazinylpyridinyl, oxopiperazinylpyridinyl, acetylpiperazinylpyridinyl, (*tert*-butoxycarbonylpiperazinyl)(methyl)pyridinyl, carboxymethylpiperazinylpyridinyl, carboxyethylpiperazinylpyridinyl, ethoxycarbonylmethylpiperazinylpyridinyl, ethoxycarbonylethylpiperazinylpyridinyl, morpholinylpyridinyl, thiomorpholinylpyridinyl, oxothiomorpholinylpyridinyl, dioxothiomorpholinylpyridinyl, oxodiazepanyl-
- 15 pyridinyl, fluoroxyetanylpyrimidinyl, hydroxyoxyetanylpyrimidinyl, hydroxyazetidinyipyrimidinyl, (hydroxy)(methyl)azetidinyipyrimidinyl, carboxyazetidinyipyrimidinyl, (*tert*-butoxycarbonyl)(hydroxy)azetidinyipyrimidinyl, tetrazolylazetidinyipyrimidinyl, hydroxytetrahydrofuranylpyrimidinyl, hydroxypyrrolidinylpyrimidinyl, carboxypyrrolidinylpyrimidinyl, (carboxy)(methyl)pyrrolidinylpyrimidinyl, carboxymethyl-
- 20 pyrrolidinylpyrimidinyl, ethoxycarbonylpyrrolidinylpyrimidinyl, fluoro-tetrahydropyranylpyrimidinyl, hydroxytetrahydropyranylpyrimidinyl, difluoropiperidinylpyrimidinyl, (cyano)(methyl)piperidinylpyrimidinyl, (hydroxy)(nitromethyl)piperidinylpyrimidinyl, (hydroxy)(methyl)piperidinylpyrimidinyl, (hydroxy)(trifluoromethyl)piperidinylpyrimidinyl, (hydroxymethyl)(methyl)piperidinylpyrimidinyl, methyl-
- 25 sulphonylpiperidinylpyrimidinyl, oxopiperidinylpyrimidinyl, (formyl)(methyl)piperidinylpyrimidinyl, carboxypiperidinylpyrimidinyl, (carboxy)(fluoro)piperidinylpyrimidinyl, (carboxy)(methyl)piperidinylpyrimidinyl, (carboxy)(ethyl)piperidinylpyrimidinyl, (carboxy)(trifluoromethyl)piperidinylpyrimidinyl, (carboxy)(hydroxy)piperidinylpyrimidinyl, (carboxy)(hydroxymethyl)piperidinylpyrimidinyl, (carboxy)-
- 30 (methoxy)piperidinylpyrimidinyl, (amino)(carboxy)piperidinylpyrimidinyl, carboxymethylpiperidinylpyrimidinyl, methoxycarbonylpiperidinylpyrimidinyl, ethoxycarbonylpiperidinylpyrimidinyl, (ethoxycarbonyl)(fluoro)piperidinylpyrimidinyl, (methoxycarbonyl)(methyl)piperidinylpyrimidinyl, (ethyl)(methoxycarbonyl)piperidinyl-

pyrimidinyl, (isopropyl)(methoxycarbonyl)piperidinylpyrimidinyl, (ethoxycarbonyl)-
 (methyl)piperidinylpyrimidinyl, (*n*-butoxycarbonyl)(methyl)piperidinylpyrimidinyl,
 (ethoxycarbonyl)(trifluoromethyl)piperidinylpyrimidinyl, (ethoxycarbonyl)-
 (hydroxymethyl)piperidinylpyrimidinyl, (methoxy)(methoxycarbonyl)piperidinyl-
 5 pyrimidinyl, (carboxy)(methoxycarbonyl)piperidinylpyrimidinyl, (methyl)-
 (morpholinylethoxycarbonyl)piperidinylpyrimidinyl, ethoxycarbonylmethylpiperidinyl-
 pyrimidinyl, methylsulphonylaminocarbonylpiperidinylpyrimidinyl, acetyl-amino-
 sulphonylpiperidinylpyrimidinyl, methoxyaminocarbonylpiperidinylpyrimidinyl,
 tetrazolylpiperidinylpyrimidinyl, hydroxyoxadiazolylpiperidinylpyrimidinyl, amino-
 10 sulphonylpiperidinylpyrimidinyl, piperazinylpyrimidinyl, methylsulphonylpiperazinyl-
 pyrimidinyl, oxopiperazinylpyrimidinyl, carboxypiperazinylpyrimidinyl, carboxyethyl-
 piperazinylpyrimidinyl, *tert*-butoxycarbonylpiperazinylpyrimidinyl, tetrazolylmethyl-
 piperazinylpyrimidinyl, trioxohexahydro-[1,2,5]thiadiazolo[2,3-*a*]pyrazinylpyrimidinyl,
 morpholinylpyrimidinyl, dimethylmorpholinylpyrimidinyl, hydroxymethylmorpholinyl-
 15 pyrimidinyl, carboxymorpholinylpyrimidinyl, (carboxy)(methyl)morpholinylpyrimidinyl,
 carboxymethylmorpholinylpyrimidinyl, thiomorpholinylpyrimidinyl, dioxo-
 thiomorpholinylpyrimidinyl, carboxyazepanylpyrimidinyl, carboxyoxazepanyl-
 pyrimidinyl, oxodiazepanylpyrimidinyl, (oxodiazepanyl)(trifluoromethyl)pyrimidinyl,
 (oxodiazepanyl)(methoxy)pyrimidinyl, (methyl)(oxo)diazepanylpyrimidinyl, dioxo-
 20 thiadiazepanylpyrimidinyl, hydroxyoxetanylpyrazinyl, (carboxy)(methyl)piperidinyl-
 pyrazinyl, (ethoxycarbonyl)(methyl)piperidinylpyrazinyl, morpholinylmethylthienyl,
 morpholinylethylpyrazolyl, carboxy-3-azabicyclo[3.1.0]hexanylpyridinyl, carboxy-3-
 azabicyclo[3.1.0]hexanylpyridazinyl, carboxy-3-azabicyclo[3.1.0]hexanylpyrimidinyl,
 (carboxy)(methyl)-3-azabicyclo[3.1.0]hexanylpyrimidinyl, methoxycarbonyl-3-
 25 azabicyclo[3.1.0]hexanylpyrimidinyl, ethoxycarbonyl-3-azabicyclo[3.1.0]hexanyl-
 pyrimidinyl, 2-oxa-5-azabicyclo[2.2.1]heptanylpyrimidinyl, carboxy-2-oxa-5-azabicyclo-
 [2.2.1]heptanylpyrimidinyl, carboxy-3-azabicyclo[3.1.1]heptanylpyrimidinyl, carboxy-3-
 azabicyclo[4.1.0]heptanylpyridinyl, carboxy-3-azabicyclo[4.1.0]heptanylpyrimidinyl,
 methoxycarbonyl-3-azabicyclo[4.1.0]heptanylpyrimidinyl, ethoxycarbonyl-3-azabicyclo-
 30 [4.1.0]heptanylpyrimidinyl, (hydroxy)(methyl)(oxo)-2-oxabicyclo[2.2.2]octanyl-
 pyrimidinyl, carboxy-3-azabicyclo[3.2.1]octanylpyrimidinyl, methoxycarbonyl-3-
 azabicyclo[3.2.1]octanylpyrimidinyl, oxo-8-azabicyclo[3.2.1]octanylpyrimidinyl,
 ethoxycarbonylmethylidenyl-8-azabicyclo[3.2.1]octanylpyrimidinyl, 3-oxa-8-azabicyclo-

[3.2.1]octanylpyrimidinyl, oxo-3,6-diazabicyclo[3.2.2]nonanylpyrimidinyl, carboxy-3-oxa-7-azabicyclo[3.3.1]nonanylpyrimidinyl, carboxy-5-azaspiro[2.3]hexanylpyrimidinyl, (carboxy)(methyl)-5-azaspiro[2.3]hexanylpyrimidinyl, carboxy-5-azaspiro[2.4]heptanylpyrimidinyl, carboxy-2-azaspiro[3.3]heptanylpyrimidinyl, 2-oxa-6-azaspiro[3.3]heptanylpyrimidinyl, 2-oxa-6-azaspiro[3.4]octanylpyrimidinyl, 2-oxa-6-azaspiro[3.5]nonanylpyrimidinyl, 2-oxa-7-azaspiro[3.5]nonanylpyrimidinyl and (dioxo)(methyl)-2,4,8-triazaspiro[4.5]decanylpyrimidinyl.

Illustrative values of R^{11} include aminosulphonylphenyl, carboxypiperidinylpyrimidinyl and morpholinylpyrimidinyl.

10 Typical examples of optional substituents on R^{12} include C_{2-6} alkoxy carbonyl.

Typical examples of particular substituents on R^{12} include ethoxycarbonyl.

In a first embodiment, R^{12} represents hydrogen. In a second embodiment, R^{12} represents halogen. In one aspect of that embodiment, R^{12} represents fluoro. In another aspect of that embodiment, R^{12} represents chloro. In a third embodiment, R^{12} represents trifluoromethyl. In a fourth embodiment, R^{12} represents optionally substituted C_{1-6} alkyl. In one aspect of that embodiment, R^{12} represents unsubstituted methyl. In another aspect of that embodiment, R^{12} represents unsubstituted ethyl. In a further aspect of that embodiment, R^{12} represents monosubstituted methyl or monosubstituted ethyl.

20 Typical values of R^{12} include hydrogen, fluoro, chloro, trifluoromethyl, methyl and ethoxycarbonyl ethyl.

Typically, R^{15} and R^{16} may independently represent hydrogen, fluoro, chloro, bromo, cyano, nitro, methyl, isopropyl, trifluoromethyl, hydroxy, methoxy, difluoromethoxy, trifluoromethoxy, methylthio, methylsulfinyl, methylsulfonyl, amino, methylamino, *tert*-butylamino, dimethylamino, phenylamino, acetylamino, methylsulfonylamino, 25 formyl, acetyl, cyclopropylcarbonyl, azetidylcarbonyl, pyrrolidinylcarbonyl, piperidinylcarbonyl, piperazinylcarbonyl, morpholinylcarbonyl, carboxy, methoxycarbonyl, aminocarbonyl, methylaminocarbonyl, dimethylaminocarbonyl, aminosulfonyl, methylaminosulfonyl and dimethylaminosulfonyl.

30 Typical values of R^{15} include hydrogen, halogen, C_{1-6} alkyl, trifluoromethyl, C_{1-6} alkoxy, difluoromethoxy and trifluoromethoxy.

In a first embodiment, R^{15} represents hydrogen. In a second embodiment, R^{15} represents halogen. In a first aspect of that embodiment, R^{15} represents fluoro. In a second aspect of that embodiment, R^{15} represents chloro. In a third embodiment, R^{15}

represents C₁₋₆ alkyl. In one aspect of that embodiment, R¹⁵ represents methyl. In a fourth embodiment, R¹⁵ represents trifluoromethyl. In a fifth embodiment, R¹⁵ represents C₁₋₆ alkoxy. In one aspect of that embodiment, R¹⁵ represents methoxy. In a sixth embodiment, R¹⁵ represents difluoromethoxy. In a seventh embodiment, R¹⁵ represents trifluoromethoxy.

Selected values of R¹⁵ include hydrogen, fluoro, chloro, methyl, trifluoromethyl, methoxy, difluoromethoxy and trifluoromethoxy.


Typical values of R¹⁶ include hydrogen, halogen, cyano, C₁₋₆ alkyl, trifluoromethyl, difluoromethoxy and amino.

In a first embodiment, R¹⁶ represents hydrogen. In a second embodiment, R¹⁶ represents halogen. In a first aspect of that embodiment, R¹⁶ represents fluoro. In a second aspect of that embodiment, R¹⁶ represents chloro. In a third embodiment, R¹⁶ represents cyano. In a fourth embodiment, R¹⁶ represents C₁₋₆ alkyl. In one aspect of that embodiment, R¹⁶ represents methyl. In a fifth embodiment, R¹⁶ represents trifluoromethyl. In a sixth embodiment, R¹⁶ represents difluoromethoxy. In a seventh embodiment, R¹⁶ represents amino.

Selected values of R¹⁶ include hydrogen, fluoro, chloro, cyano, methyl, trifluoromethyl, difluoromethoxy and amino.

In a particular embodiment, R¹⁶ is attached at the *para*-position of the phenyl ring relative to the integer R¹⁵.

A particular sub-group of the compounds of formula (IIA) above is represented by the compounds of formula (IIB) and *N*-oxides thereof, and pharmaceutically acceptable salts and solvates thereof, and glucuronide derivatives thereof, and co-crystals thereof:



(IIB)

wherein

V represents C-R²² or N;

R²¹ represents hydrogen, halogen, halo(C₁₋₆)alkyl, cyano, C₁₋₆ alkyl, trifluoromethyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxy, hydroxy(C₁₋₆)alkyl, C₁₋₆ alkoxy, (C₁₋₆)alkoxy-(C₁₋₆)alkyl, difluoromethoxy, trifluoromethoxy, trifluoroethoxy, carboxy(C₃₋₇)cycloalkyloxy, C₁₋₆ alkylthio, C₁₋₆ alkylsulphonyl, (C₁₋₆)alkylsulphonyl(C₁₋₆)alkyl, amino, amino-
 5 (C₁₋₆)alkyl, C₁₋₆ alkylamino, di(C₁₋₆)alkylamino, (C₁₋₆)alkoxy(C₁₋₆)alkylamino, *N*-[(C₁₋₆)alkyl]-*N*-[hydroxy(C₁₋₆)alkyl]amino, C₂₋₆ alkylcarbonylamino, (C₂₋₆)alkylcarbonylamino-(C₁₋₆)alkyl, C₂₋₆ alkoxycarbonylamino, *N*-[(C₁₋₆)alkyl]-*N*-[carboxy(C₁₋₆)alkyl]amino, carboxy(C₃₋₇)cycloalkylamino, carboxy(C₃₋₇)cycloalkyl(C₁₋₆)alkylamino, C₁₋₆ alkylsulphonylamino, C₁₋₆ alkylsulphonylamino(C₁₋₆)alkyl, formyl, C₂₋₆ alkylcarbonyl,
 10 (C₂₋₆)alkylcarbonyloxy(C₁₋₆)alkyl, carboxy, carboxy(C₁₋₆)alkyl, C₂₋₆ alkoxycarbonyl, morpholinyl(C₁₋₆)alkoxycarbonyl, C₂₋₆ alkoxycarbonyl(C₁₋₆)alkyl, C₂₋₆ alkoxycarbonylmethylidenyl, aminocarbonyl, C₁₋₆ alkylaminocarbonyl, di(C₁₋₆)alkylaminocarbonyl, aminosulphonyl, C₁₋₆ alkylaminosulphonyl, di(C₁₋₆)alkylaminosulphonyl, (C₁₋₆)alkylsulphoximinyl or [(C₁₋₆)alkyl][*N*-(C₁₋₆)alkyl]sulphoximinyl; or R²¹ represents
 15 (C₃₋₇)cycloalkyl, (C₃₋₇)cycloalkyl(C₁₋₆)alkyl, (C₄₋₇)cycloalkenyl, (C₄₋₉)bicycloalkyl, (C₃₋₇)heterocycloalkyl, (C₃₋₇)heterocycloalkenyl, (C₄₋₉)heterobicycloalkyl or (C₄₋₉)spiroheterocycloalkyl, any of which groups may be optionally substituted by one or more substituents;

R²² represents hydrogen, halogen or C₁₋₆ alkyl;

20 R²³ represents hydrogen, C₁₋₆ alkyl, trifluoromethyl or C₁₋₆ alkoxy; and
 E, R¹², R¹⁵ and R¹⁶ are as defined above.

In one embodiment, V represents C-R²². In another embodiment, V represents N.

Typically, R²¹ represents hydrogen, halogen, halo(C₁₋₆)alkyl, cyano, C₁₋₆ alkyl, trifluoromethyl, C₂₋₆ alkenyl, hydroxy, hydroxy(C₁₋₆)alkyl, C₁₋₆ alkoxy, trifluoroethoxy,
 25 carboxy(C₃₋₇)cycloalkyloxy, C₁₋₆ alkylthio, C₁₋₆ alkylsulphonyl, amino, C₁₋₆ alkylamino, di(C₁₋₆)alkylamino, (C₁₋₆)alkoxy(C₁₋₆)alkylamino, *N*-[(C₁₋₆)alkyl]-*N*-[hydroxy(C₁₋₆)alkyl]-amino, *N*-[(C₁₋₆)alkyl]-*N*-[carboxy(C₁₋₆)alkyl]amino, carboxy(C₃₋₇)cycloalkylamino, carboxy(C₃₋₇)cycloalkyl(C₁₋₆)alkylamino, C₁₋₆ alkylsulphonylamino, (C₂₋₆)alkylcarbonyloxy(C₁₋₆)alkyl, carboxy, morpholinyl(C₁₋₆)alkoxycarbonyl, C₂₋₆ alkoxycarbonyl(C₁₋₆)alkyl
 30 or C₂₋₆ alkoxycarbonylmethylidenyl; or R²¹ represents (C₃₋₇)cycloalkyl, (C₃₋₇)cycloalkyl-(C₁₋₆)alkyl, (C₄₋₇)cycloalkenyl, (C₄₋₉)bicycloalkyl, (C₃₋₇)heterocycloalkyl, (C₄₋₉)heterobicycloalkyl or (C₄₋₉)spiroheterocycloalkyl, any of which groups may be optionally substituted by one or more substituents.

Suitably, R²¹ represents (C₃₋₇)heterocycloalkyl, which group may be optionally substituted by one or more substituents.

Where R²¹ represents an optionally substituted (C₃₋₇)cycloalkyl group, typical values include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl, any of which groups may be optionally substituted by one or more substituents.

Where R²¹ represents an optionally substituted (C₃₋₇)cycloalkyl(C₁₋₆)alkyl group, a typical value is cyclohexylmethyl, which group may be optionally substituted by one or more substituents.

Where R²¹ represents an optionally substituted (C₄₋₇)cycloalkenyl group, typical values include cyclobutenyl, cyclopentenyl, cyclohexenyl and cycloheptenyl, any of which groups may be optionally substituted by one or more substituents.

Where R²¹ represents an optionally substituted (C₄₋₉)bicycloalkyl group, typical values include bicyclo[3.1.0]hexanyl, bicyclo[4.1.0]heptanyl and bicyclo[2.2.2]octanyl, any of which groups may be optionally substituted by one or more substituents.

Where R²¹ represents an optionally substituted (C₃₋₇)heterocycloalkyl group, typical values include oxetanyl, azetidiny, tetrahydrofuranyl, pyrrolidinyl, tetrahydropyranyl, piperidinyl, piperazinyl, hexahydro-[1,2,5]thiadiazolo[2,3-*a*]pyrazinyl, morpholinyl, thiomorpholinyl, azepanyl, oxazepanyl, diazepanyl and thiadiazepanyl, any of which groups may be optionally substituted by one or more substituents.

Where R²¹ represents an optionally substituted (C₃₋₇)heterocycloalkenyl group, a typical value is optionally substituted 1,2,3,6-tetrahydropyridinyl.

Where R²¹ represents an optionally substituted (C₄₋₉)heterobicycloalkyl group, typical values include 3-azabicyclo[3.1.0]hexanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 3-azabicyclo[3.1.1]heptanyl, 3-azabicyclo[4.1.0]heptanyl, 2-oxabicyclo[2.2.2]octanyl, quinuclidinyl, 2-oxa-5-azabicyclo[2.2.2]octanyl, 3-azabicyclo[3.2.1]octanyl, 8-azabicyclo[3.2.1]octanyl, 3-oxa-8-azabicyclo[3.2.1]octanyl, 3,8-diazabicyclo[3.2.1]octanyl, 3,6-diazabicyclo[3.2.2]nonanyl, 3-oxa-7-azabicyclo[3.3.1]nonanyl and 3,9-diazabicyclo[4.2.1]nonanyl, any of which groups may be optionally substituted by one or more substituents.

Where R²¹ represents an optionally substituted (C₄₋₉)spiroheterocycloalkyl group, typical values include 5-azaspiro[2.3]hexanyl, 5-azaspiro[2.4]heptanyl, 2-azaspiro[3.3]heptanyl, 2-oxa-6-azaspiro[3.3]heptanyl, 2-oxa-6-azaspiro[3.4]octanyl, 2-oxa-6-azaspiro-

[3.5]nonanyl, 2-oxa-7-azaspiro[3.5]nonanyl and 2,4,8-triazaspiro[4.5]-decanyl, any of which groups may be optionally substituted by one or more substituents.

Illustratively, R²¹ represents hydroxy, hydroxy(C₁₋₆)alkyl, methoxy, carboxy-cyclobutyloxy, methylthio, methylsulphonyl, methylamino, *N*-[carboxyethyl]-*N*-methyl-amino, carboxycyclopentylamino, carboxycyclopropylmethylamino or ethoxycarbonyl-ethyl; or R²¹ represents cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclohexyl-methyl, cyclohexenyl, bicyclo[3.1.0]hexanyl, bicyclo[4.1.0]heptanyl, bicyclo[2.2.2]-octanyl, oxetanyl, azetidiny, tetrahydrofuran, pyrrolidinyl, tetrahydropyranyl, piperidinyl, piperazinyl, hexahydro-[1,2,5]thiadiazolo[2,3-*a*]pyrazinyl, morpholinyl, thiomorpholinyl, azepanyl, oxazepanyl, diazepanyl, thiadiazepanyl, 3-azabicyclo[3.1.0]-hexanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 3-azabicyclo[3.1.1]heptanyl, 3-azabicyclo[4.1.0]heptanyl, 2-oxabicyclo[2.2.2]octanyl, 3-azabicyclo[3.2.1]octanyl, 8-azabicyclo[3.2.1]octanyl, 3-oxa-8-azabicyclo[3.2.1]octanyl, 3,6-diazabicyclo[3.2.2]nonanyl, 3-oxa-7-azabicyclo[3.3.1]nonanyl, 5-azaspiro[2.3]hexanyl, 5-azaspiro[2.4]heptanyl or 2-azaspiro-[3.3]heptanyl, any of which groups may be optionally substituted by one or more substituents.

Appositely, R²¹ represents piperidinyl or morpholinyl, either of which groups may be optionally substituted by one or more substituents.

Examples of optional substituents which may be present on R²¹ include one, two or three substituents independently selected from halogen, halo(C₁₋₆)alkyl, cyano, cyano-(C₁₋₆)alkyl, nitro, nitro(C₁₋₆)alkyl, C₁₋₆ alkyl, trifluoromethyl, trifluoroethyl, C₂₋₆ alkenyl, hydroxy, hydroxy(C₁₋₆)alkyl, C₁₋₆ alkoxy, difluoromethoxy, trifluoromethoxy, trifluoroethoxy, C₁₋₆ alkylthio, C₁₋₆ alkylsulphonyl, (C₁₋₆)alkylsulphonyl(C₁₋₆)alkyl, oxo, amino, C₁₋₆ alkylamino, di(C₁₋₆)alkylamino, C₂₋₆ alkylcarbonylamino, (C₂₋₆)alkylcarbonylamino-(C₁₋₆)alkyl, C₂₋₆ alkoxycarbonylamino, C₁₋₆ alkylsulphonylamino, formyl, C₂₋₆ alkylcarbonyl, carboxy, carboxy(C₁₋₆)alkyl, C₂₋₆ alkoxycarbonyl, morpholinyl-(C₁₋₆)alkoxycarbonyl, C₂₋₆ alkoxycarbonyl(C₁₋₆)alkyl, C₂₋₆ alkoxycarbonylmethylenyl, a carboxylic acid isostere or prodrug moiety Ω as defined herein, -(C₁₋₆)alkyl- Ω , amino-carbonyl, C₁₋₆ alkylaminocarbonyl, di(C₁₋₆)alkylaminocarbonyl, aminosulphonyl, di(C₁₋₆)alkylaminosulphonyl, (C₁₋₆)alkylsulphoximinyl and [(C₁₋₆)alkyl][*N*-(C₁₋₆)alkyl]-sulphoximinyl.

Selected examples of optional substituents on R²¹ include one, two or three substituents independently selected from carboxy.

Suitable examples of particular substituents on R²¹ include one, two or three substituents independently selected from fluoro, fluoromethyl, chloro, bromo, cyano, cyanomethyl, cyanoethyl, nitro, nitromethyl, methyl, ethyl, isopropyl, trifluoromethyl, trifluoroethyl, ethenyl, hydroxy, hydroxymethyl, methoxy, ethoxy, difluoromethoxy, trifluoromethoxy, trifluoroethoxy, methylthio, methylsulphonyl, methylsulphonylmethyl, methylsulphonylethyl, oxo, amino, methylamino, dimethylamino, acetylamino, acetylaminomethyl, methoxycarbonylamino, ethoxycarbonylamino, *tert*-butoxycarbonylamino, methylsulphonylamino, formyl, acetyl, carboxy, carboxymethyl, carboxyethyl, methoxycarbonyl, ethoxycarbonyl, *n*-butoxycarbonyl, *tert*-butoxycarbonyl, morpholinylethoxycarbonyl, methoxycarbonylmethyl, ethoxycarbonylmethyl, ethoxycarbonylethyl, ethoxycarbonylmethylidenyl, acetylaminosulphonyl, methoxyaminocarbonyl, tetrazolyl, tetrazolylmethyl, hydroxyoxadiazolyl, aminocarbonyl, methylaminocarbonyl, dimethylaminocarbonyl, methylsulphonylaminocarbonyl, aminosulphonyl, methylaminosulphonyl, dimethylaminosulphonyl, methylsulphoximinyl and (methyl)(*N*-methyl)sulphoximinyl.

Selected examples of particular substituents on R²¹ include one, two or three substituents independently selected from carboxy.

Typically, R²¹ represents hydrogen, fluoro, fluoroisopropyl, cyano, methyl, trifluoromethyl, ethenyl, hydroxy, hydroxyisopropyl, methoxy, isopropoxy, trifluoroethoxy, carboxycyclobutyl, methylthio, methylsulphonyl, amino, methylamino, dimethylamino, methoxyethylamino, *N*-(hydroxyethyl)-*N*-(methyl)amino, *N*-[carboxyethyl]-*N*-methylamino, carboxycyclopentylamino, carboxycyclopropylmethylamino, methylsulphonylamino, acetoxisopropyl, carboxy, ethoxycarbonylethyl, fluoromethylcyclopropyl, acetylaminomethylcyclopropyl, hydroxycyclobutyl, carboxycyclopentyl, carboxycyclohexyl, (carboxy)(methyl)cyclohexyl, (carboxy)(hydroxy)cyclohexyl, carboxymethylcyclohexyl, ethoxycarbonylcyclohexyl, (methoxycarbonyl)(methyl)cyclohexyl, (ethoxycarbonyl)(methyl)cyclohexyl, carboxycyclohexylmethyl, carboxycyclohexenyl, ethoxycarbonylcyclohexenyl, carboxybicyclo[3.1.0]hexanyl, ethoxycarbonylbicyclo[3.1.0]hexanyl, carboxybicyclo[4.1.0]heptanyl, carboxybicyclo[2.2.2]octanyl, fluoroxyetanyl, hydroxyoxyetanyl, hydroxyazetidiny, (hydroxy)(methyl)azetidiny, carboxyazetidiny, (*tert*-butoxycarbonyl)(hydroxy)azetidiny, tetrazolylazetidiny, hydroxytetrahydrofuranyl, pyrrolidiny, hydroxypyrrolidiny, carboxypyrrolidiny, (carboxy)(methyl)pyrrolidiny, carboxymethylpyrrolidiny, ethoxycarbonylpyrrolidiny, fluorotetrahydropyranyl, hydroxytetrahydropyranyl, piperidiny, difluoro-

piperidinyl, (cyano)(methyl)piperidinyl, (hydroxy)(nitromethyl)piperidinyl, (hydroxy)-
 (methyl)piperidinyl, (hydroxy)(trifluoromethyl)piperidinyl, (hydroxymethyl)(methyl)-
 piperidinyl, methylsulphonylpiperidinyl, oxopiperidinyl, (formyl)(methyl)piperidinyl,
 acetylpiperidinyl, carboxypiperidinyl, (carboxy)(fluoro)piperidinyl, (carboxy)(methyl)-
 5 piperidinyl, (carboxy)(ethyl)piperidinyl, (carboxy)(trifluoromethyl)piperidinyl, (carboxy)-
 (hydroxy)piperidinyl, (carboxy)(hydroxymethyl)piperidinyl, (carboxy)(methoxy)-
 piperidinyl, (amino)(carboxy)piperidinyl, carboxymethylpiperidinyl, methoxycarbonyl-
 piperidinyl, (methoxycarbonyl)(methyl)piperidinyl, (ethyl)(methoxycarbonyl)piperidinyl,
 (isopropyl)(methoxycarbonyl)piperidinyl, (methoxy)(methoxycarbonyl)piperidinyl,
 10 (carboxy)(methoxycarbonyl)piperidinyl, ethoxycarbonylpiperidinyl, (ethoxycarbonyl)-
 (fluoro)piperidinyl, (ethoxycarbonyl)(methyl)piperidinyl, (ethoxycarbonyl)(trifluoro-
 methyl)piperidinyl, (ethoxycarbonyl)(hydroxymethyl)piperidinyl, (*n*-butoxycarbonyl)-
 (methyl)piperidinyl, (methyl)(morpholinylethoxycarbonyl)piperidinyl, ethoxycarbonyl-
 methylpiperidinyl, methylsulphonylaminocarbonylpiperidinyl, acetylaminosulphonyl-
 15 piperidinyl, methoxyaminocarbonylpiperidinyl, tetrazolypiperidinyl, hydroxyoxadiazolyl-
 piperidinyl, aminosulphonylpiperidinyl, piperazinyl, cyanoethylpiperazinyl, trifluoroethyl-
 piperazinyl, methylsulphonylpiperazinyl, methylsulphonylethylpiperazinyl,
 oxopiperazinyl, acetylpiperazinyl, carboxypiperazinyl, *tert*-butoxycarbonylpiperazinyl,
 carboxymethylpiperazinyl, carboxyethylpiperazinyl, ethoxycarbonylmethylpiperazinyl,
 20 ethoxycarbonylethylpiperazinyl, tetrazolylmethylpiperazinyl, trioxohexahydro-
 [1,2,5]thiadiazolo[2,3-*a*]pyrazinyl, morpholinyl, dimethylmorpholinyl, hydroxymethyl-
 morpholinyl, carboxymorpholinyl, (carboxy)(methyl)morpholinyl, carboxymethyl-
 morpholinyl, thiomorpholinyl, oxothiomorpholinyl, dioxothiomorpholinyl, carboxy-
 azepanyl, carboxyoxazepanyl, oxodiazepanyl, (methyl)(oxo)diazepanyl, dioxo-
 25 thiadiazepanyl, carboxy-3-azabicyclo[3.1.0]hexanyl, (carboxy)(methyl)-3-azabicyclo-
 [3.1.0]hexanyl, methoxycarbonyl-3-azabicyclo[3.1.0]hexanyl, ethoxycarbonyl-3-
 azabicyclo[3.1.0]hexanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, carboxy-2-oxa-5-
 azabicyclo[2.2.1]heptanyl, carboxy-3-azabicyclo[3.1.1]heptanyl, carboxy-3-azabicyclo-
 [4.1.0]heptanyl, methoxycarbonyl-3-azabicyclo[4.1.0]heptanyl, ethoxycarbonyl-3-
 30 azabicyclo[4.1.0]heptanyl, (hydroxy)(methyl)(oxo)-2-oxabicyclo[2.2.2]octanyl, carboxy-
 3-azabicyclo[3.2.1]octanyl, methoxycarbonyl-3-azabicyclo[3.2.1]octanyl, oxo-8-
 azabicyclo[3.2.1]octanyl, ethoxycarbonylmethylidenyl-8-azabicyclo[3.2.1]octanyl, 3-oxa-
 8-azabicyclo[3.2.1]octanyl, oxo-3,6-diazabicyclo[3.2.2]nonanyl, carboxy-3-oxa-7-

azabicyclo[3.3.1]nonanyl, carboxy-5-azaspiro[2.3]hexanyl, (carboxy)(methyl)-5-azaspiro[2.3]hexanyl, carboxy-5-azaspiro[2.4]heptanyl, carboxy-2-azaspiro[3.3]heptanyl, 2-oxa-6-azaspiro[3.3]heptanyl, 2-oxa-6-azaspiro[3.4]octanyl, 2-oxa-6-azaspiro[3.5]nonanyl, 2-oxa-7-azaspiro[3.5]nonanyl or (dioxo)(methyl)-2,4,8-triazaspiro[4.5]decanyl.

5 Illustrative values of R^{21} include carboxypiperidiny and morpholinyl.

In a particular embodiment, R^{21} represents hydroxy(C_{1-6})alkyl. In one aspect of that embodiment, R^{21} represents hydroxyisopropyl, especially 2-hydroxyprop-2-yl.

Generally, R^{22} represents hydrogen or C_{1-6} alkyl.

Suitably, R^{22} represents hydrogen, chloro or methyl.

10 Typically, R^{22} represents hydrogen or methyl.

In one embodiment, R^{22} represents hydrogen. In another embodiment, R^{22} represents C_{1-6} alkyl, especially methyl. In a further embodiment, R^{22} represents halogen. In one aspect of that embodiment, R^{22} represents fluoro. In another aspect of that embodiment, R^{22} represents chloro.


15 Generally, R^{23} represents hydrogen or C_{1-6} alkyl.




Suitably, R^{23} represents hydrogen, methyl, trifluoromethyl or methoxy.

Typically, R^{23} represents hydrogen or methyl.



In one embodiment, R^{23} represents hydrogen. In another embodiment, R^{23} represents C_{1-6} alkyl, especially methyl. In a further embodiment, R^{23} represents trifluoromethyl. In an additional embodiment, R^{23} represents C_{1-6} alkoxy, especially methoxy.



25 Particular sub-groups of the compounds of formula (IIB) above are represented by the compounds of formula (IIC), (IID), (IIE), (IIF), (IIG), (IIH), (IIJ), (IIK) and (IIL), and *N*-oxides thereof, and pharmaceutically acceptable salts and solvates thereof, and glucuronide derivatives thereof, and co-crystals thereof:






5





5



wherein

T represents $-\text{CH}_2-$ or $-\text{CH}_2\text{CH}_2-$;

10 U represents $\text{C}(\text{O})$ or $\text{S}(\text{O})_2$;

W represents O, S, $\text{S}(\text{O})$, $\text{S}(\text{O})_2$, $\text{S}(\text{O})(\text{NR}^4)$, $\text{N}(\text{R}^{31})$ or $\text{C}(\text{R}^{32})(\text{R}^{33})$;

-M- represents $-\text{CH}_2-$ or $-\text{CH}_2\text{CH}_2-$;

15 R^{31} represents hydrogen, cyano(C_{1-6})alkyl, C_{1-6} alkyl, trifluoromethyl, trifluoroethyl, C_{1-6} alkylsulphonyl, (C_{1-6})alkylsulphonyl(C_{1-6})alkyl, formyl, C_{2-6} alkylcarbonyl, carboxy, carboxy(C_{1-6})alkyl, C_{2-6} alkoxy carbonyl, C_{2-6} alkoxy carbonyl(C_{1-6})alkyl, a carboxylic acid isostere or prodrug moiety Ω , $-(\text{C}_{1-6})\text{alkyl}-\Omega$, aminocarbonyl, C_{1-6} alkylaminocarbonyl, di(C_{1-6})alkylaminocarbonyl, aminosulphonyl or di(C_{1-6})alkylamino-sulphonyl;

R^{32} represents hydrogen, halogen, cyano, hydroxy, hydroxy(C_{1-6})alkyl, C_{1-6} alkylsulphonyl, formyl, C_{2-6} alkylcarbonyl, carboxy, carboxy(C_{1-6})alkyl, C_{2-6} alkoxy, C_{2-6} alkoxy(C_{1-6})alkyl, aminosulphonyl, (C_{1-6})alkylsulphoximinyl, [(C_{1-6})alkyl][N -(C_{1-6})alkyl]sulphoximinyl, a carboxylic acid isostere or
5 prodrug moiety Ω , or -(C_{1-6})alkyl- Ω ;

R^{33} represents hydrogen, halogen, C_{1-6} alkyl, trifluoromethyl, hydroxy, hydroxy-(C_{1-6})alkyl, C_{1-6} alkoxy, amino or carboxy;

R^{34} represents hydrogen, halogen, halo(C_{1-6})alkyl, hydroxy, C_{1-6} alkoxy, C_{1-6} alkylthio, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, amino, C_{1-6} alkylamino, di(C_{1-6})alkyl-
10 amino, (C_{2-6})alkylcarbonylamino, (C_{2-6})alkylcarbonylamino(C_{1-6})alkyl, (C_{1-6})alkylsulphonylamino or (C_{1-6})alkylsulphonylamino(C_{1-6})alkyl; and

V , E , R^4 , R^{12} , R^{15} , R^{16} , R^{23} and Ω are as defined above.

In a first embodiment, T represents $-CH_2-$. In a second embodiment, T represents $-CH_2CH_2-$.

15 In a first embodiment, U represents $C(O)$. In a second embodiment, U represents $S(O)_2$.

Generally, W represents O , $S(O)_2$, $N(R^{31})$ or $C(R^{32})(R^{33})$.

Typically, W represents O , $N(R^{31})$ or $C(R^{32})(R^{33})$.

Suitably, W represents O or $C(R^{32})(R^{33})$.

20 In a first embodiment, W represents O . In a second embodiment, W represents S . In a third embodiment, W represents $S(O)$. In a fourth embodiment, W represents $S(O)_2$. In a fifth embodiment, W represents $S(O)(NR^4)$. In a sixth embodiment, W represents $N(R^{31})$. In a seventh embodiment, W represents $C(R^{32})(R^{33})$.

In one embodiment, $-M-$ represents $-CH_2-$. In another embodiment, $-M-$ represents
25 $-CH_2CH_2-$.

Typically, R^{31} represents hydrogen, cyano(C_{1-6})alkyl, C_{1-6} alkyl, trifluoromethyl, trifluoroethyl, C_{1-6} alkylsulphonyl, (C_{1-6})alkylsulphonyl(C_{1-6})alkyl, formyl, C_{2-6} alkylcarbonyl, carboxy, carboxy(C_{1-6})alkyl, C_{2-6} alkoxy, C_{2-6} alkoxy(C_{1-6})alkyl, (C_{1-6})alkyl, tetrazolyl(C_{1-6})alkyl, aminocarbonyl, C_{1-6} alkylaminocarbonyl, di(C_{1-6})alkyl-
30 aminocarbonyl, aminosulphonyl, C_{1-6} alkylaminosulphonyl or di(C_{1-6})alkylaminosulphonyl.

Suitably, R^{31} represents hydrogen or C_{1-6} alkylsulphonyl.

Typical values of R^{31} include hydrogen, cyanoethyl, methyl, ethyl, isopropyl, trifluoromethyl, trifluoroethyl, methylsulphonyl, methylsulphonylethyl, formyl, acetyl, carboxy, carboxymethyl, carboxyethyl, methoxycarbonyl, ethoxycarbonyl, *tert*-butoxycarbonyl, ethoxycarbonylmethyl, ethoxycarbonylethyl, tetrazolylmethyl, aminocarbonyl, methylamino-carbonyl, dimethylaminocarbonyl, aminosulphonyl, methylaminosulphonyl and dimethylaminosulphonyl.

Particular values of R^{31} include hydrogen and methylsulphonyl.

Generally, R^{32} represents halogen, carboxy, carboxy(C_{1-6})alkyl, C_{2-6} alkoxy carbonyl, C_{2-6} alkoxy carbonyl(C_{1-6})alkyl, a carboxylic acid isostere or prodrug moiety Ω , or $-(C_{1-6})$ alkyl- Ω .

Typically, R^{32} represents hydrogen, halogen, cyano, hydroxy, hydroxy(C_{1-6})alkyl, C_{1-6} alkylsulphonyl, formyl, carboxy, carboxy(C_{1-6})alkyl, C_{2-6} alkoxy carbonyl, C_{2-6} alkoxy carbonyl(C_{1-6})alkyl, aminosulphonyl, (C_{1-6})alkylsulphoximinyl, [(C_{1-6})alkyl][*N*-(C_{1-6})alkyl]sulphoximinyl, (C_{1-6})alkylsulphonylaminocarbonyl, (C_{2-6})alkylcarbonylamino-sulphonyl, (C_{1-6})alkoxyaminocarbonyl, tetrazolyl or hydroxyoxadiazolyl.

Typical values of R^{32} include hydrogen, fluoro, cyano, hydroxy, hydroxymethyl, methylsulphonyl, formyl, carboxy, carboxymethyl, carboxyethyl, methoxycarbonyl, ethoxycarbonyl, *tert*-butoxycarbonyl, methoxycarbonylmethyl, methoxycarbonylethyl, ethoxycarbonylmethyl, ethoxycarbonylethyl, aminosulphonyl, methylsulphoximinyl, (methyl)(*N*-methyl)sulphoximinyl, methylsulphonylaminocarbonyl, acetylaminosulphonyl, methoxyaminocarbonyl, tetrazolyl and hydroxyoxadiazolyl.

In a selected embodiment, R^{32} represents carboxy.

Generally, R^{33} represents hydrogen, halogen or C_{1-6} alkyl.

Suitably, R^{33} represents hydrogen or C_{1-6} alkyl.

Selected values of R^{33} include hydrogen, fluoro, methyl, ethyl, isopropyl, trifluoromethyl, hydroxy, hydroxymethyl, methoxy, amino and carboxy.

Particular values of R^{33} include hydrogen and methyl.

In a first embodiment, R^{33} represents hydrogen. In a second embodiment, R^{33} represents halogen. In one aspect of that embodiment, R^{33} represents fluoro. In a third embodiment, R^{33} represents C_{1-6} alkyl. In a first aspect of that embodiment, R^{33} represents methyl. In a second aspect of that embodiment, R^{33} represents ethyl. In a third aspect of that embodiment, R^{33} represents isopropyl. In a fourth embodiment, R^{33} represents trifluoromethyl. In a fifth embodiment, R^{33} represents hydroxy. In a sixth embodiment,

R^{33} represents hydroxy(C_{1-6})alkyl. In one aspect of that embodiment, R^{33} represents hydroxymethyl. In a seventh embodiment, R^{33} represents C_{1-6} alkoxy. In one aspect of that embodiment, R^{33} represents methoxy. In an eighth embodiment, R^{33} represents amino. In a ninth embodiment, R^{33} represents carboxy.


- 5 In a first embodiment, R^{34} represents hydrogen. In a second embodiment, R^{34} represents halogen. In one aspect of that embodiment, R^{34} represents fluoro. In a third embodiment, R^{34} represents halo(C_{1-6})alkyl. In one aspect of that embodiment, R^{34} represents fluoromethyl. In a fourth embodiment, R^{34} represents hydroxy. In a fifth embodiment, R^{34} represents C_{1-6} alkoxy, especially methoxy. In a sixth embodiment, R^{34} represents C_{1-6} alkylthio, especially methylthio. In a seventh embodiment, R^{34} represents C_{1-6} alkylsulphinyl, especially methylsulphinyl. In an eighth embodiment, R^{34} represents C_{1-6} alkylsulphonyl, especially methylsulphonyl. In a ninth embodiment, R^{34} represents amino. In a tenth embodiment, R^{34} represents C_{1-6} alkylamino, especially methylamino. In an eleventh embodiment, R^{34} represents di(C_{1-6})alkylamino, especially dimethylamino.
- 15 In a twelfth embodiment, R^{34} represents (C_{2-6})alkylcarbonylamino, especially acetylamino. In a thirteenth embodiment, R^{34} represents (C_{2-6})alkylcarbonylamino(C_{1-6})alkyl, especially acetylaminomethyl. In a fourteenth embodiment, R^{34} represents (C_{1-6})alkylsulphonylamino, especially methylsulphonylamino. In a fifteenth embodiment, R^{34} represents (C_{1-6})alkylsulphonylamino(C_{1-6})alkyl, especially methylsulphonylaminomethyl.
- 20 Typically, R^{34} represents hydrogen, halogen, halo(C_{1-6})alkyl, hydroxy or (C_{2-6})alkylcarbonylamino(C_{1-6})alkyl.

Selected values of R^{34} include hydrogen, fluoro, fluoromethyl, hydroxy, methoxy, methylthio, methylsulphinyl, methylsulphonyl, amino, methylamino, dimethylamino and acetylaminomethyl.

- 25 Particular values of R^{34} include hydrogen, fluoro, fluoromethyl, hydroxy and acetylaminomethyl.

Suitably, R^{34} represents hydrogen or hydroxy.

- An alternative sub-class of compounds according to the invention is represented by the compounds of formula (IIM) and *N*-oxides thereof, and pharmaceutically acceptable salts and solvates thereof, and glucuronide derivatives thereof, and co-crystals thereof:
- 30



wherein

E, W, R¹², R¹⁵, R¹⁶ and R²¹ are as defined above.

5 With specific reference to formula (IIM), the integer W is suitably O, S or N-R³¹, especially S or N-R³¹.

Specific novel compounds in accordance with the present invention include each of the compounds whose preparation is described in the accompanying Examples, and pharmaceutically acceptable salts and solvates thereof, and co-crystals thereof.

10 The compounds in accordance with the present invention are beneficial in the treatment and/or prevention of various human ailments. These include autoimmune and inflammatory disorders; neurological and neurodegenerative disorders; pain and nociceptive disorders; cardiovascular disorders; metabolic disorders; ocular disorders; and oncological disorders.

15 Inflammatory and autoimmune disorders include systemic autoimmune disorders, autoimmune endocrine disorders and organ-specific autoimmune disorders. Systemic autoimmune disorders include systemic lupus erythematosus (SLE), psoriasis, psoriatic arthropathy, vasculitis, polymyositis, scleroderma, multiple sclerosis, systemic sclerosis, ankylosing spondylitis, rheumatoid arthritis, non-specific inflammatory arthritis, juvenile
20 inflammatory arthritis, juvenile idiopathic arthritis (including oligoarticular and polyarticular forms thereof), anaemia of chronic disease (ACD), Still's disease (juvenile and/or adult onset), Behçet's disease and Sjögren's syndrome. Autoimmune endocrine disorders include thyroiditis. Organ-specific autoimmune disorders include Addison's disease, haemolytic or pernicious anaemia, acute kidney injury (AKI; including cisplatin-
25 induced AKI), diabetic nephropathy (DN), obstructive uropathy (including cisplatin-induced obstructive uropathy), glomerulonephritis (including Goodpasture's syndrome, immune complex-mediated glomerulonephritis and antineutrophil cytoplasmic antibodies (ANCA)-associated glomerulonephritis), lupus nephritis (LN), minimal change disease,

- Graves' disease, idiopathic thrombocytopenic purpura, inflammatory bowel disease (including Crohn's disease, ulcerative colitis, indeterminate colitis and pouchitis), pemphigus, atopic dermatitis, autoimmune hepatitis, primary biliary cirrhosis, autoimmune pneumonitis, autoimmune carditis, myasthenia gravis, spontaneous infertility,
- 5 osteoporosis, osteopenia, erosive bone disease, chondritis, cartilage degeneration and/or destruction, fibrosing disorders (including various forms of hepatic and pulmonary fibrosis), asthma, rhinitis, chronic obstructive pulmonary disease (COPD), respiratory distress syndrome, sepsis, fever, muscular dystrophy (including Duchenne muscular dystrophy) and organ transplant rejection (including kidney allograft rejection).
- 10 Neurological and neurodegenerative disorders include Alzheimer's disease, Parkinson's disease, Huntington's disease, ischaemia, stroke, amyotrophic lateral sclerosis, spinal cord injury, head trauma, seizures and epilepsy.
- Cardiovascular disorders include thrombosis, cardiac hypertrophy, hypertension, irregular contractility of the heart (e.g. during heart failure), and sexual disorders
- 15 (including erectile dysfunction and female sexual dysfunction). Modulators of TNF α function may also be of use in the treatment and/or prevention of myocardial infarction (see J.J. Wu *et al.*, *JAMA*, 2013, **309**, 2043-2044).
- Metabolic disorders include diabetes (including insulin-dependent diabetes mellitus and juvenile diabetes), dyslipidemia and metabolic syndrome.
- 20 Ocular disorders include retinopathy (including diabetic retinopathy, proliferative retinopathy, non-proliferative retinopathy and retinopathy of prematurity), macular oedema (including diabetic macular oedema), age-related macular degeneration (ARMD), vascularisation (including corneal vascularisation and neovascularisation), retinal vein occlusion, and various forms of uveitis and keratitis.
- 25 Oncological disorders, which may be acute or chronic, include proliferative disorders, especially cancer, and cancer-associated complications (including skeletal complications, cachexia and anaemia). Particular categories of cancer include haematological malignancy (including leukaemia and lymphoma) and non-haematological malignancy (including solid tumour cancer, sarcoma, meningioma, glioblastoma
- 30 multiforme, neuroblastoma, melanoma, gastric carcinoma and renal cell carcinoma). Chronic leukaemia may be myeloid or lymphoid. Varieties of leukaemia include lymphoblastic T cell leukaemia, chronic myelogenous leukaemia (CML), chronic lymphocytic/lymphoid leukaemia (CLL), hairy-cell leukaemia, acute lymphoblastic

leukaemia (ALL), acute myelogenous leukaemia (AML), myelodysplastic syndrome, chronic neutrophilic leukaemia, acute lymphoblastic T cell leukaemia, plasmacytoma, immunoblastic large cell leukaemia, mantle cell leukaemia, multiple myeloma, acute megakaryoblastic leukaemia, acute megakaryocytic leukaemia, promyelocytic leukaemia and erythroleukaemia. Varieties of lymphoma include malignant lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, lymphoblastic T cell lymphoma, Burkitt's lymphoma, follicular lymphoma, MALT1 lymphoma and marginal zone lymphoma. Varieties of non-haematological malignancy include cancer of the prostate, lung, breast, rectum, colon, lymph node, bladder, kidney, pancreas, liver, ovary, uterus, cervix, brain, skin, bone, stomach and muscle. Modulators of TNF α function may also be used to increase the safety of the potent anticancer effect of TNF (see F.V. Hauwermeiren *et al.*, *J. Clin. Invest.*, 2013, **123**, 2590-2603).

The present invention also provides a pharmaceutical composition which comprises a compound in accordance with the invention as described above, or a pharmaceutically acceptable salt or solvate thereof, in association with one or more pharmaceutically acceptable carriers.

Pharmaceutical compositions according to the invention may take a form suitable for oral, buccal, parenteral, nasal, topical, ophthalmic or rectal administration, or a form suitable for administration by inhalation or insufflation.

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets, lozenges or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methyl cellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium hydrogenphosphate); lubricants (e.g. magnesium stearate, talc or silica); disintegrants (e.g. potato starch or sodium glycollate); or wetting agents (e.g. sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents, emulsifying agents, non-aqueous vehicles or preservatives. The preparations may also contain buffer salts, flavouring agents, colouring agents or sweetening agents, as appropriate.

Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

- 5 The compounds of formula (I) may be formulated for parenteral administration by injection, e.g. by bolus injection or infusion. Formulations for injection may be presented in unit dosage form, e.g. in glass ampoules or multi-dose containers, e.g. glass vials. The compositions for injection may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending,
10 stabilising, preserving and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use.

- In addition to the formulations described above, the compounds of formula (I) may also be formulated as a depot preparation. Such long-acting formulations may be
15 administered by implantation or by intramuscular injection.

- For nasal administration or administration by inhalation, the compounds according to the present invention may be conveniently delivered in the form of an aerosol spray presentation for pressurised packs or a nebuliser, with the use of a suitable propellant, e.g. dichlorodifluoromethane, fluorotrichloromethane, dichlorotetrafluoroethane, carbon
20 dioxide or other suitable gas or mixture of gases.

 The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack or dispensing device may be accompanied by instructions for administration.

- For topical administration the compounds of use in the present invention may be
25 conveniently formulated in a suitable ointment containing the active component suspended or dissolved in one or more pharmaceutically acceptable carriers. Particular carriers include, for example, mineral oil, liquid petroleum, propylene glycol, polyoxyethylene, polyoxypropylene, emulsifying wax and water. Alternatively, the compounds of use in the present invention may be formulated in a suitable lotion containing the active component
30 suspended or dissolved in one or more pharmaceutically acceptable carriers. Particular carriers include, for example, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, benzyl alcohol, 2-octyldodecanol and water.

For ophthalmic administration the compounds of use in the present invention may be conveniently formulated as micronized suspensions in isotonic, pH-adjusted sterile saline, either with or without a preservative such as a bactericidal or fungicidal agent, for example phenylmercuric nitrate, benzylalkonium chloride or chlorhexidine acetate.

- 5 Alternatively, for ophthalmic administration compounds may be formulated in an ointment such as petrolatum.


For rectal administration the compounds of use in the present invention may be conveniently formulated as suppositories. These can be prepared by mixing the active component with a suitable non-irritating excipient which is solid at room temperature but
10 liquid at rectal temperature and so will melt in the rectum to release the active component. Such materials include, for example, cocoa butter, beeswax and polyethylene glycols.

The quantity of a compound of use in the invention required for the prophylaxis or treatment of a particular condition will vary depending on the compound chosen and the condition of the patient to be treated. In general, however, daily dosages may range from
15 around 10 ng/kg to 1000 mg/kg, typically from 100 ng/kg to 100 mg/kg, e.g. around 0.01 mg/kg to 40 mg/kg body weight, for oral or buccal administration, from around 10 ng/kg to 50 mg/kg body weight for parenteral administration, and from around 0.05 mg to around 1000 mg, e.g. from around 0.5 mg to around 1000 mg, for nasal administration or administration by inhalation or insufflation.

- 20 If desired, a compound in accordance with the present invention may be co-administered with another pharmaceutically active agent, e.g. an anti-inflammatory molecule such as methotrexate or prednisolone.

The compounds of formula (I) above may be prepared by a process which comprises cyclising a compound of formula (III):

25




(III)

wherein E, Y, R¹, R² and R³ are as defined above.

The cyclisation may conveniently be effected by heating compound (III) in phosphorus oxychloride.

The compounds of formula (III) above may be prepared by reacting a compound of formula Y-E-CO₂H with a compound of formula (IV):

5




(IV)

wherein E, Y, R¹, R² and R³ are as defined above.

The reaction may be accomplished in the presence of a coupling reagent such as 3-(3-dimethylaminopropyl)-1-ethylcarbodiimide (EDCI), typically in the presence of a suitable base, e.g. an organic base such as triethylamine. The reaction is conveniently effected at ambient temperature in a suitable solvent, e.g. a chlorinated solvent such as dichloromethane.

The compounds of formula (IV) above may be prepared by reacting a compound of formula (V):

15



(V)

wherein R¹, R² and R³ are as defined above, and L¹ represents a suitable leaving group; with hydrazine hydrate.

20

The leaving group L¹ is typically a halogen atom, e.g. chloro.

The reaction is conveniently effected at an elevated temperature in a suitable solvent, e.g. a cyclic ether such as tetrahydrofuran.

Where they are not commercially available, the starting materials of formula (V) may be prepared by methods analogous to those described in the accompanying Examples, or by standard methods well known from the art.

It will be understood that any compound of formula (I) initially obtained from any
5 of the above processes may, where appropriate, subsequently be elaborated into a further compound of formula (I) by techniques known from the art. By way of example, a compound of formula (I) wherein E represents -C(O)- may be converted into the corresponding compound wherein E represents -CH(OH)- by treatment with a reducing agent such as sodium borohydride.

10 A compound of formula (I) wherein E represents -CH(OH)- may be converted into the corresponding compound wherein E represents -CH₂- by heating with elemental iodine and phosphinic acid in acetic acid; or by treating with triethylsilane and an acid, e.g. an organic acid such as trifluoroacetic acid, or a Lewis acid such as boron trifluoride diethyl etherate; or by treating with chlorotrimethylsilane and sodium iodide; or by a two-step
15 procedure which comprises: (i) treatment with thionyl bromide; and (ii) treatment of the product thereby obtained with a transition metal catalyst, e.g. (2,2'-bipyridine)dichlororuthenium(II) hydrate, in the presence of diethyl 1,4-dihydro-2,6-dimethyl-3,5-pyridinedicarboxylate (Hantzsch ester) and a base, e.g. an organic base such as *N,N*-diisopropylethylamine.

20 A compound of formula (I) wherein E represents -CH₂- may be converted into the corresponding compound wherein E represents -CH(CH₃)- by treatment with a methyl halide, e.g. methyl iodide, in the presence of a base such as lithium hexamethyldisilazide.

A compound of formula (I) which contains a hydroxy group may be alkylated by treatment with the appropriate alkyl halide in the presence of a base, e.g. sodium hydride,
25 or silver oxide. A compound of formula (I) which contains hydroxy may be converted into the corresponding fluoro-substituted compound by treatment with diethylaminosulfur trifluoride (DAST) or bis(2-methoxyethyl)aminosulfur trifluoride (BAST). A compound of formula (I) which contains hydroxy may be converted into the corresponding difluoro-substituted compound via a two-step procedure which comprises: (i) treatment with an
30 oxidising agent, e.g. manganese dioxide; and (ii) treatment of the carbonyl-containing compound thereby obtained with DAST.

A compound of formula (I) which contains an N-H moiety may be alkylated by treatment with the appropriate alkyl halide, typically at an elevated temperature in an

organic solvent such as acetonitrile; or at ambient temperature in the presence of a base, e.g. an alkali metal carbonate such as potassium carbonate or cesium carbonate, in a suitable solvent, e.g. a dipolar aprotic solvent such as *N,N*-dimethylformamide.

Alternatively, a compound of formula (I) which contains an N-H moiety may be alkylated
5 by treatment with the appropriate alkyl tosylate in the presence of a base, e.g. an inorganic base such as sodium hydride, or an organic base such as 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU).

A compound of formula (I) which contains an N-H moiety may be methylated by treatment with formaldehyde in the presence of a reducing agent, e.g. sodium
10 triacetoxyborohydride.

A compound of formula (I) which contains an N-H moiety may be acylated by treatment with the appropriate acid chloride, e.g. acetyl chloride, or with the appropriate carboxylic acid anhydride, e.g. acetic anhydride, typically at ambient temperature in the presence of a base, e.g. an organic base such as triethylamine.

15 A compound of formula (I) which contains an N-H moiety may be converted into the corresponding compound wherein the nitrogen atom is substituted by C₁₋₆ alkylsulphonyl, e.g. methylsulphonyl, by treatment with the appropriate C₁₋₆ alkylsulphonyl chloride, e.g. methanesulphonyl chloride, or with the appropriate C₁₋₆ alkylsulphonic acid anhydride, e.g. methanesulphonic anhydride, typically at ambient temperature in the
20 presence of a base, e.g. an organic base such as triethylamine or *N,N*-diisopropylethylamine.

A compound of formula (I) substituted by amino (-NH₂) may be converted into the corresponding compound substituted by C₁₋₆ alkylsulphonylamino, e.g. methylsulphonylamino, or bis[(C₁₋₆)alkylsulphonyl]amino, e.g. bis(methylsulphonyl)amino, by treatment
25 with the appropriate C₁₋₆ alkylsulphonyl halide, e.g. a C₁₋₆ alkylsulphonyl chloride such as methanesulphonyl chloride. Similarly, a compound of formula (I) substituted by hydroxy (-OH) may be converted into the corresponding compound substituted by C₁₋₆ alkylsulphonyloxy, e.g. methylsulphonyloxy, by treatment with the appropriate C₁₋₆ alkylsulphonyl halide, e.g. a C₁₋₆ alkylsulphonyl chloride such as methanesulphonyl chloride.

30 A compound of formula (I) containing the moiety -S- may be converted into the corresponding compound containing the moiety -S(O)- by treatment with 3-chloroperoxybenzoic acid. Likewise, a compound of formula (I) containing the moiety -S(O)- may be converted into the corresponding compound containing the moiety -S(O)₂- by treatment

with 3-chloroperoxybenzoic acid. Alternatively, a compound of formula (I) containing the moiety -S- may be converted into the corresponding compound containing the moiety -S(O)₂- by treatment with Oxone® (potassium peroxymonosulfate).

A compound of formula (I) containing an aromatic nitrogen atom may be
5 converted into the corresponding *N*-oxide derivative by treatment with 3-chloroperoxybenzoic acid.

A bromophenyl derivative of formula (I) may be converted into the corresponding optionally substituted 2-oxopyrrolidin-1-ylphenyl or 2-oxooxazolidin-3-ylphenyl derivative by treatment with pyrrolidin-2-one or oxazolidin-2-one, or an appropriately
10 substituted analogue thereof. The reaction is conveniently effected at an elevated temperature in the presence of copper(I) iodide, *trans*-*N,N'*-dimethylcyclohexane-1,2-diamine and an inorganic base such as potassium carbonate.

A compound of formula (I) wherein R¹ represents halogen, e.g. bromo, may be converted into the corresponding compound wherein R¹ represents an optionally
15 substituted aryl or heteroaryl moiety by treatment with the appropriately substituted aryl or heteroaryl boronic acid or a cyclic ester thereof formed with an organic diol, e.g. pinacol, 1,3-propanediol or neopentyl glycol. The reaction is typically effected in the presence of a transition metal catalyst, e.g. [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), tetrakis(triphenylphosphine)palladium(0), or bis[3-(diphenylphosphanyl)cyclopenta-2,4-
20 dien-1-yl]iron-dichloropalladium-dichloromethane complex, and a base, e.g. an inorganic base such as sodium carbonate or potassium carbonate, or potassium phosphate.

A compound of formula (I) wherein R¹ represents halogen, e.g. bromo, may be converted into the corresponding compound wherein R¹ represents an optionally substituted aryl, heteroaryl or heterocycloalkenyl moiety via a two-step procedure which
25 comprises: (i) reaction with bis(pinacolato)diboron or bis(neopentyl glycolato)diboron; and (ii) reaction of the compound thereby obtained with an appropriately functionalised halo- or tosyloxy-substituted aryl, heteroaryl or heterocycloalkenyl derivative. Step (i) is conveniently effected in the presence of a transition metal catalyst such as [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), or bis[3-(diphenylphosphanyl)-
30 cyclopenta-2,4-dien-1-yl]iron-dichloropalladium-dichloromethane complex. Step (ii) is conveniently effected in the presence of a transition metal catalyst such as tetrakis(triphenylphosphine)palladium(0), or bis[3-(diphenylphosphanyl)cyclopenta-2,4-dien-1-

yl]iron-dichloropalladium-dichloromethane complex, and a base, e.g. an inorganic base such as sodium carbonate or potassium carbonate.

5 A compound of formula (I) wherein R¹ represents halogen, e.g. bromo, may be converted into the corresponding compound wherein R¹ represents an optionally substituted C₂₋₆ alkynyl moiety by treatment with an appropriately substituted alkyne derivative, e.g. 2-hydroxybut-3-yne. The reaction is conveniently accomplished with the assistance of a transition metal catalyst, e.g. tetrakis(triphenylphosphine)palladium(0), typically in the presence of copper(I) iodide and a base, e.g. an organic base such as triethylamine.

10 A compound of formula (I) wherein R¹ represents halogen, e.g. bromo, may be converted into the corresponding compound wherein R¹ represents an optionally substituted imidazol-1-yl moiety by treatment with the appropriately substituted imidazole derivative, typically in the presence of copper(II) acetate and an organic base such as *N,N,N',N'*-tetramethylethylenediamine (TMEDA).

15 A compound of formula (I) wherein R¹ represents halogen, e.g. bromo, may be converted into the corresponding compound wherein R¹ represents 2-(methoxycarbonyl)-ethyl via a two-step procedure which comprises: (i) reaction with methyl acrylate; and (ii) catalytic hydrogenation of the alkenyl derivative thereby obtained, typically by treatment with a hydrogenation catalyst, e.g. palladium on charcoal, under an atmosphere of
20 hydrogen gas. Step (i) is typically effected in the presence of a transition metal catalyst, e.g. palladium(II) acetate or bis(dibenzylideneacetone)palladium(0), and a reagent such as tri(*ortho*-tolyl)phosphine.

In general, a compound of formula (I) containing a -C=C- functionality may be converted into the corresponding compound containing a -CH-CH- functionality by
25 catalytic hydrogenation, typically by treatment with a hydrogenation catalyst, e.g. palladium on charcoal, under an atmosphere of hydrogen gas, optionally in the presence of a base, e.g. an alkali metal hydroxide such as sodium hydroxide.

A compound of formula (I) wherein R¹ represents 6-methoxypyridin-3-yl may be converted into the corresponding compound wherein R¹ represents 2-oxo-1,2-dihydro-
30 pyridin-5-yl by treatment with pyridine hydrochloride; or by heating with a mineral acid such as hydrochloric acid. By utilising similar methodology, a compound of formula (I) wherein R¹ represents 6-methoxy-4-methylpyridin-3-yl may be converted into the corresponding compound wherein R¹ represents 4-methyl-2-oxo-1,2-dihydropyridin-5-yl;

and a compound of formula (I) wherein R¹ represents 6-methoxy-5-methylpyridin-3-yl may be converted into the corresponding compound wherein R¹ represents 3-methyl-2-oxo-1,2-dihydropyridin-5-yl.

5 A compound of formula (I) wherein R¹ represents 2-oxo-1,2-dihydropyridin-5-yl may be converted into the corresponding compound wherein R¹ represents 2-oxopiperidin-5-yl by catalytic hydrogenation, typically by treatment with gaseous hydrogen in the presence of a hydrogenation catalyst such as platinum(IV) oxide.

A compound of formula (I) containing an ester moiety, e.g. a C₂₋₆ alkoxy carbonyl group such as methoxycarbonyl or ethoxycarbonyl, may be converted into the
10 corresponding compound containing a carboxy (-CO₂H) moiety by treatment with an acid, e.g. a mineral acid such as hydrochloric acid.

A compound of formula (I) containing an *N*-(*tert*-butoxycarbonyl) moiety may be converted into the corresponding compound containing an N-H moiety by treatment with an acid, e.g. a mineral acid such as hydrochloric acid, or an organic acid such as
15 trifluoroacetic acid.

A compound of formula (I) containing an ester moiety, e.g. a C₂₋₆ alkoxy carbonyl group such as methoxycarbonyl or ethoxycarbonyl, may alternatively be converted into the corresponding compound containing a carboxy (-CO₂H) moiety by treatment with a base, e.g. an alkali metal hydroxide selected from lithium hydroxide, sodium hydroxide and
20 potassium hydroxide; or an organic base such as sodium methoxide or sodium ethoxide.

A compound of formula (I) containing a carboxy (-CO₂H) moiety may be converted into the corresponding compound containing an amide moiety by treatment with the appropriate amine in the presence of a condensing agent such as 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide.

25 A compound of formula (I) containing a carbonyl (C=O) moiety may be converted into the corresponding compound containing a -C(CH₃)(OH)- moiety by treatment with methylmagnesium bromide. Similarly, a compound of formula (I) containing a carbonyl (C=O) moiety may be converted into the corresponding compound containing a -C(CF₃)(OH)- moiety by treatment with (trifluoromethyl)trimethylsilane and cesium
30 fluoride. A compound of formula (I) containing a carbonyl (C=O) moiety may be converted into the corresponding compound containing a -C(CH₂NO₂)(OH)- moiety by treatment with nitromethane.

A compound of formula (I) containing a hydroxymethyl moiety may be converted into the corresponding compound containing a formyl (-CHO) moiety by treatment with an oxidising agent such as Dess-Martin periodinane. A compound of formula (I) containing a hydroxymethyl moiety may be converted into the corresponding compound
5 containing a carboxy moiety by treatment with an oxidising agent such as tetrapropylammonium perruthenate.

A compound of formula (I) wherein R¹ represents a substituent containing at least one nitrogen atom, which substituent is linked to the remainder of the molecule via a nitrogen atom, may be prepared by reacting a compound of formula (I) wherein R¹
10 represents halogen, e.g. bromo, with the appropriate compound of formula R¹-H [e.g. 1-(pyridin-3-yl)piperazine or morpholine]. The reaction is conveniently effected with the assistance of a transition metal catalyst, e.g. tris(dibenzylideneacetone)dipalladium(0), in the presence of an amination ligand such as 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl (XPhos) or 2,2'-bis(diphenylphosphino)-1,1'-binaphthalene (BINAP) and a base,
15 e.g. an inorganic base such as sodium *tert*-butoxide. Alternatively, the reaction may be effected using palladium diacetate, in the presence of a reagent such as [2',6'-bis(propan-2-yloxy)biphenyl-2-yl](dicyclohexyl)phosphane and a base, e.g. an inorganic base such as cesium carbonate.

A compound of formula (I) containing an oxo moiety can be converted into the
20 corresponding compound containing an ethoxycarbonylmethylidene moiety by treatment with triethyl phosphonoacetate in the presence of a base such as sodium hydride.

A compound of formula (IIB) wherein R²¹ represents ethenyl may be prepared by reacting a compound of formula (IIB) wherein R²¹ represents halogen, e.g. chloro, with potassium vinyl trifluoroborate. The reaction is typically effected in the presence of a
25 transition metal catalyst, e.g. [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), and a base, e.g. an organic base such as triethylamine.

A compound of formula (IIB) wherein R²¹ represents halogen, e.g. chloro, may be converted into the corresponding compound wherein R²¹ represents an optionally substituted C₄₋₇ cycloalkenyl moiety by treatment with the appropriately substituted
30 cycloalkenyl boronic acid or a cyclic ester thereof formed with an organic diol, e.g. pinacol, 1,3-propanediol or neopentyl glycol. The reaction is typically effected in the presence of a transition metal catalyst, e.g. bis[3-(diphenylphosphanyl)cyclopenta-2,4-

dien-1-yl]iron-dichloropalladium-dichloromethane complex, and a base, e.g. an inorganic base such as potassium carbonate.

A compound of formula (IIB) wherein R^{21} represents a substituent containing at least one nitrogen atom, which substituent is linked to the remainder of the molecule via a nitrogen atom, may be prepared by reacting a compound of formula (IIB) wherein R^{21} represents halogen, e.g. chloro, with the appropriate compound of formula R^{21} -H [e.g. 2-methoxyethylamine, *N*-methyl-*L*-alanine, 2-aminocyclopentanecarboxylic acid, 3-aminocyclopentanecarboxylic acid, 1-(aminomethyl)cyclopropanecarboxylic acid, methyl azetidine-3-carboxylate, pyrrolidin-3-ol, pyrrolidine-3-carboxylic acid, piperidine-2-carboxylic acid, piperidine-3-carboxylic acid, 4-(1*H*-tetrazol-5-yl)piperidine, piperazine, 1-(methylsulfonyl)piperazine, piperazin-2-one, 2-(piperazin-1-yl)propanoic acid, morpholine, morpholine-2-carboxylic acid, thiomorpholine, thiomorpholine 1,1-dioxide, 1,4-diazepan-5-one, 2-oxa-5-azabicyclo[2.2.1]heptane or an appropriately substituted azaspiroalkane], optionally in the presence of a base, e.g. an organic base such as triethylamine or *N,N*-diisopropylethylamine and/or 1-methyl-2-pyrrolidinone, or pyridine, or an inorganic base such as potassium carbonate.

Where a mixture of products is obtained from any of the processes described above for the preparation of compounds according to the invention, the desired product can be separated therefrom at an appropriate stage by conventional methods such as preparative HPLC; or column chromatography utilising, for example, silica and/or alumina in conjunction with an appropriate solvent system.

Where the above-described processes for the preparation of the compounds according to the invention give rise to mixtures of stereoisomers, these isomers may be separated by conventional techniques. In particular, where it is desired to obtain a particular enantiomer of a compound of formula (I) this may be produced from a corresponding mixture of enantiomers using any suitable conventional procedure for resolving enantiomers. Thus, for example, diastereomeric derivatives, e.g. salts, may be produced by reaction of a mixture of enantiomers of formula (I), e.g. a racemate, and an appropriate chiral compound, e.g. a chiral base. The diastereomers may then be separated by any convenient means, for example by crystallisation, and the desired enantiomer recovered, e.g. by treatment with an acid in the instance where the diastereomer is a salt. In another resolution process a racemate of formula (I) may be separated using chiral HPLC. Moreover, if desired, a particular enantiomer may be obtained by using an

appropriate chiral intermediate in one of the processes described above. Alternatively, a particular enantiomer may be obtained by performing an enantiomer-specific enzymatic biotransformation, e.g. an ester hydrolysis using an esterase, and then purifying only the enantiomerically pure hydrolysed acid from the unreacted ester antipode.

- 5 Chromatography, recrystallisation and other conventional separation procedures may also be used with intermediates or final products where it is desired to obtain a particular geometric isomer of the invention.

During any of the above synthetic sequences it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in
10 *Protective Groups in Organic Chemistry*, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons, 3rd edition, 1999. The protecting groups may be removed at any convenient subsequent stage utilising methods known from the art.

- 15 The following Examples illustrate the preparation of compounds according to the invention.

The compounds in accordance with this invention potently inhibit the binding of a fluorescence conjugate to TNF α when tested in the fluorescence polarisation assay described below. Moreover, certain compounds in accordance with this invention
20 potently inhibit TNF α -induced NF- κ B activation in the reporter gene assay described below.

Fluorescence Polarisation Assay

Preparation of Compound (A)

- 25 1-(2,5-Dimethylbenzyl)-6-[4-(piperazin-1-ylmethyl)phenyl]-2-(pyridin-4-ylmethyl)-1*H*-benzimidazole – hereinafter referred to as “*Compound (A)*” – can be prepared by the procedure described in Example 499 of WO 2013/186229 (published 19 December 2013); or by a procedure analogous thereto.

- 30 *Preparation of fluorescence conjugate*

Compound (A) (27.02 mg, 0.0538 mmol) was dissolved in DMSO (2 mL). 5 (-6) Carboxy-fluorescein succinimyl ester (24.16 mg, 0.0510 mmol) (Invitrogen catalogue number: C1311) was dissolved in DMSO (1 mL) to give a bright yellow solution. The

two solutions were mixed at room temperature, the mixture turning red in colour. The mixture was stirred at room temperature. Shortly after mixing a 20 μ L aliquot was removed and diluted in a 80:20 mixture of AcOH:H₂O for LC-MS analysis on the 1200RR-6140 LC-MS system. The chromatogram showed two closely eluting peaks at retention times of 1.42 and 1.50 minutes, both with mass $(M+H)^+ = 860.8$ amu, corresponding to the two products formed with the 5- and 6-substituted carboxyfluorescein group. A further peak at retention time 2.21 minutes had a mass of $(M+H)^+ = 502.8$ amu, corresponding to *Compound (A)*. No peak was observed for unreacted 5(-6) carboxyfluorescein succinimyl ester. The peak areas were 22.0%, 39.6% and 31.4% for the three signals, indicating a 61.6% conversion to the two isomers of the desired fluorescence conjugate at that time-point. Further 20 μ L aliquots were extracted after several hours and then after overnight stirring, diluted as before and subjected to LC-MS analysis. The percentage conversion was determined as 79.8% and 88.6% respectively at these time-points. The mixture was purified on a UV-directed preparative HPLC system. The pooled purified fractions were freeze-dried to remove excess solvent. After freeze-drying, an orange solid (23.3 mg) was recovered, equivalent to 0.027 mmol of fluorescence conjugate, corresponding to an overall yield of 53% for the reaction and preparative HPLC purification.

Inhibition of binding of fluorescence conjugate to TNF α

Compounds were tested at 10 concentrations starting from 25 μ M in a final assay concentration of 5% DMSO, by pre-incubation with TNF α for 60 minutes at ambient temperature in 20 mM Tris, 150 mM NaCl, 0.05% Tween™ 20, before addition of the fluorescence conjugate and a further incubation for 20 hours at ambient temperature. The final concentrations of TNF α and the fluorescence conjugate were 10 nM and 10 nM respectively in a total assay volume of 25 μ L. Plates were read on a plate reader capable of detecting fluorescence polarisation (e.g. an Analyst HT plate reader; or an Envision plate reader). An IC₅₀ value was calculated using XLfit™ (4 parameter logistic model) in ActivityBase.

When tested in the fluorescence polarisation assay, the compounds of the accompanying Examples were all found to exhibit IC₅₀ values of 50 μ M or better.

Reporter Gene Assay*Inhibition of TNF α -induced NF- κ B activation*

- Stimulation of HEK-293 cells by TNF α leads to activation of the NF- κ B pathway. The reporter cell line used to determine TNF α activity was purchased from InvivoGen.
- 5 HEK-Blue™ CD40L is a stable HEK-293 transfected cell line expressing SEAP (secreted embryonic alkaline phosphatase) under the control of the IFN β minimal promoter fused to five NF- κ B binding sites. Secretion of SEAP by these cells is stimulated in a dose-dependent manner by TNF α , with an EC₅₀ of 0.5 ng/mL for human TNF α . Compounds were diluted from 10 mM DMSO stocks (final assay concentration 0.3% DMSO) to
- 10 generate a 10-point 3-fold serial dilution curve (e.g. 30,000 nM to 2 nM final concentration). Diluted compound was preincubated with TNF α for 60 minutes prior to addition to a 384-well microtitre plate and incubated for 18 h. The final TNF α concentration in the assay plate was 0.5 ng/mL. SEAP activity was determined in the supernatant using a colorimetric substrate, e.g. QUANTI-Blue™ or HEK-Blue™
- 15 Detection media (InvivoGen). Percentage inhibitions for compound dilutions were calculated between a DMSO control and maximum inhibition (by excess control compound) and an IC₅₀ value calculated using XLfit™ (4 parameter logistic model) in ActivityBase.

- When tested in the reporter gene assay, certain compounds of the accompanying
- 20 Examples were found to exhibit IC₅₀ values of 50 μ M or better.

EXAMPLES**Abbreviations**

- 25 DCM: dichloromethane EtOAc: ethyl acetate
THF: tetrahydrofuran DMSO: dimethylsulfoxide
EtOH: ethanol Et₂O: diethyl ether
EDCI: 3-(3-dimethylaminopropyl)-1-ethylcarbodiimide
Pd(dppf)Cl₂: [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II)
- 30 h: hour M: mass
HPLC: High Performance Liquid Chromatography
LCMS: Liquid Chromatography Mass Spectrometry
ES+: Electrospray Positive Ionisation RT: retention time

Nomenclature

Compounds were named with the aid of ACD/Name Batch (Network) version 11.01, and/or Accelrys Draw 4.0.

5

Analytical Conditions***Analytical HPLC***

Column: Waters, X Bridge, 20 x 2.1 mm, 2.5 μ m
Mobile Phase A: 10 mM ammonium formate in water + 0.1% ammonia
10 Mobile Phase B: acetonitrile + 5% solvent A + 0.1% ammonia
Injection Volume: 5.0 μ L
Flow Rate: 1.00 mL/minute
Gradient program: 5% B to 95% B in 4 minutes; hold till 5.00 minutes; at 5.10 minutes
B conc. is 5% up to 6.5 minutes

15

GENERAL METHOD A

To a degassed solution of *Intermediate 3* (150 mg, 0.48 mmol), sodium carbonate (153 mg, 1.44 mmol) and the respective boronic acid (0.48 mmol) in 1,4-dioxane (1 mL)
20 and water (0.3 mL) was added Pd(dppf)Cl₂ (35 mg, 0.0042 mmol). The reaction mixture was heated at 125°C under microwave irradiation for 2 h, then diluted with EtOAc and filtered through Celite™. The organic layer was dried over anhydrous sodium sulphate, filtered and concentrated *in vacuo*, then purified by preparative HPLC.

25

INTERMEDIATE 1**(6-Chloropyridazin-3-yl)hydrazine**

A stirred solution of 3,6-dichloropyridazine (3.0 g, 20.3 mmol) and hydrazine hydrate (0.97 g, 20.3 mmol) in THF (15 mL) was heated at 80°C for 12 h in a sealed tube.
30 The reaction mixture was concentrated *in vacuo*. The crude residue was triturated with Et₂O to give the *title compound* (2.0 g, 68%), which was used without further purification. δ_{H} (400 MHz, DMSO-d₆) 8.25 (s, 1H), 7.42 (dd, 1H, *J* 9.5, 1.6 Hz), 7.10 (d, 1H, *J* 9.4 Hz), 4.99-4.10 (m, 2H).

INTERMEDIATE 2

N'-(6-Chloropyridazin-3-yl)-2-[2-(difluoromethoxy)phenyl]acetohydrazide

5 To a stirred solution of 2-(difluoromethoxy)phenylacetic acid (3.0 g, 14.8 mmol) in DCM (20 mL) at 0°C was added triethylamine (4.4 g, 44.5 mmol), followed by EDCI (3.3 g, 17.7 mmol). The reaction mixture was stirred at 0°C for 30 minutes. *Intermediate*
10 *1* (2.3 g, 16.3 mmol) was added and the reaction mixture was stirred at room temperature for 12 h, then diluted with DCM. The organic layer was washed with water and brine, then concentrated *in vacuo*. The crude residue was triturated with pentane and Et₂O to give the *title compound* (3.50 g, 72%), which was used without further purification. δ_{H} (400 MHz, DMSO-d₆) 10.21 (s, 1H), 9.17 (s, 1H), 7.54 (dd, 1H, *J* 9.3, 1.2 Hz), 7.40 (td, 1H, *J* 7.0, 6.6, 1.8 Hz), 7.33 (ddt, 1H, *J* 11.8, 5.9, 2.8 Hz), 7.26-7.10 (m, 3H), 7.05-6.92 (m, 1H), 3.62 (s, 2H).

15

INTERMEDIATE 3

6-Chloro-3-{{2-(difluoromethoxy)phenyl}methyl}[1,2,4]triazolo[4,3-*b*]pyridazine

A solution of *Intermediate 2* (4.0 g, 12.2 mmol) in POCl₃ (20 mL) was heated at
20 100°C for 12 h, then the reaction mixture was basified using saturated aqueous NaHCO₃ solution. The aqueous layer was extracted with DCM. The organic layer was washed with water, then dried over anhydrous sodium sulphate, filtered and concentrated *in vacuo*. The crude residue was triturated in Et₂O to give the *title compound* (7.0 g, 54%) as a solid, which was used without further purification. δ_{H} (400 MHz, CD₃OD) 8.24 (d, 1H, *J* 9.7 Hz), 7.42 (d, 1H, *J* 9.7 Hz), 7.35 (dd, 2H, *J* 10.3, 6.8 Hz), 7.20 (t, 2H, *J* 7.8 Hz),
25 6.79 (t, 1H, *J* 7.4 Hz), 4.58 (s, 2H). LCMS (ES⁺) 311 (M+H)⁺, RT 2.07 minutes.

INTERMEDIATE 4

2-(Morpholin-4-yl)pyrimidin-5-ylboronic acid

A mixture of 2-chloropyrimidin-5-ylboronic acid (3.0 g, 19.0 mmol), morpholine (1.66 mL, 19.0 mmol) and triethylamine (1.67 mL, 19.2 mmol) in EtOH (20 mL) was stirred at 80°C for 5 h. The reaction mixture was concentrated *in vacuo* and the residue

was taken up in Et₂O (approximately 5 mL). Et₂O was added, and the triethylamine hydrochloride salt that crystallised out was filtered and discarded. The filtrate was concentrated *in vacuo* and water (approximately 10 mL) was added. The mixture was placed in a refrigerator for 1 h, after which time the resulting solid was filtered off, washed with the minimum amount of water and dried by suction, to give the *title compound* (2.7 g, 68%) as an off-white solid. δ_{H} (400 MHz, DMSO-d₆) 8.64 (s, 2H), 8.08 (s, 2H), 3.73 (m, 4H), 3.65 (m, 4H). LCMS (ES⁺) 210 (M+H)⁺, RT 0.15 minutes.

INTERMEDIATE 5

1-(5-Boronopyrimidin-2-yl)piperidine-4-carboxylic acid

2-Chloropyrimidin-5-ylboronic acid (2.00 g, 12.6 mmol) and isonipecotic acid (1.63 g, 12.6 mmol) were suspended in EtOH (25 mL). Triethylamine (1.78 mL, 12.65 mmol) was added and the mixture was heated at 80°C for 16 h. The reaction mixture was cooled to room temperature and concentrated *in vacuo* to dryness. Water (30 mL) was added and the reaction mixture was swirled until the product completely dissolved. On standing, crystallisation occurred. The mixture was cooled in an ice bath for 30 minutes, then filtered. The resultant solid was washed sparingly with water and dried under suction, then freeze-dried, to give the *title compound* (1.90 g, 60%) as a white solid. δ_{H} (400 MHz, DMSO-d₆) 8.60 (s, 2H), 8.06 (br s, 2H), 4.60-4.52 (m, 2H), 3.14-3.02 (m, 2H), 2.60-2.54 (m, 1H), 1.90-1.80 (m, 2H), 1.55-1.39 (m, 2H). LCMS (ES⁺) 252 (M+H)⁺.

EXAMPLE 1

4-[5-(3-{[2-(Difluoromethoxy)phenyl]methyl}[1,2,4]triazolo[4,3-*b*]pyridazin-6-yl)-pyrimidin-2-yl]morpholine

Prepared from *Intermediate 3* (150 mg, 0.48 mmol), sodium carbonate (153 mg, 1.44 mmol), *Intermediate 4* (100 mg, 0.48 mmol) and Pd(dppf)Cl₂ (39 mg, 0.0048 mmol) in 1,4-dioxane (1 mL) and water (0.3 mL) according to *General Method A*. Purification of the crude material by preparative HPLC gave the *title compound* (25 mg, 12%) as a green solid. δ_{H} (400 MHz, CD₃OD) 9.00 (s, 2H), 8.20 (d, 1H, *J* 8.2 Hz), 7.82 (d, 1H, *J* 9.8 Hz), 7.40 (d, 1H, *J* 7.5 Hz), 7.37 (t, 1H, *J* 7.9 Hz), 7.21 (t, 2H, *J* 7.8 Hz), 6.81 (t, 1H, *J*

74.0 Hz), 4.65 (s, 2H), 3.95 (m, 4H), 3.79 (m, 4H). LCMS (ES+) 440 (M+H)⁺, RT 2.05 minutes.

EXAMPLE 2

5

1-[5-(3-{[2-(Difluoromethoxy)phenyl]methyl}[1,2,4]triazolo[4,3-*b*]pyridazin-6-yl)-pyrimidin-2-yl]piperidine-4-carboxylic acid

Prepared from *Intermediate 3* (150 mg, 0.48 mmol), sodium carbonate (153 mg, 1.44 mmol), *Intermediate 5* (121 mg, 0.48 mmol) and Pd(dppf)Cl₂ (39 mg, 0.0048 mmol) in 1,4-dioxane (1 mL) and water (0.3 mL) according to *General Method A*. Purification of the crude material by preparative HPLC gave the *title compound* (43 mg, 19%) as a grey solid. δ_{H} (400 MHz, CD₃OD) 12.40-12.20 (br s, 1H), 9.03 (s, 2H), 8.39 (d, 1H, *J* 9.8 Hz), 7.89 (d, 1H, *J* 9.8 Hz), 7.56-7.29 (m, 2H), 7.29-6.85 (m, 3H), 4.74-4.42 (m, 4H), 3.26-3.05 (m, 2H), 2.60 (m, 1H), 1.92 (dd, 2H, *J* 12.6, 4.3 Hz), 1.63-1.41 (m, 2H). LCMS (ES+) 482 (M+H)⁺, RT 1.53 minutes.

10
15

EXAMPLE 3

4-(3-{[2-(Difluoromethoxy)phenyl]methyl}[1,2,4]triazolo[4,3-*b*]pyridazin-6-yl)-benzenesulfonamide

20


Prepared from *Intermediate 3* (150 mg, 0.48 mmol), sodium carbonate (153 mg, 1.44 mmol), (4-sulfamoylphenyl)boronic acid (137 mg, 0.48 mmol) and Pd(dppf)Cl₂ (39 mg, 0.0048 mmol) in 1,4-dioxane (1 mL) and water (0.3 mL) according to *General Method A*. Purification of the crude material by preparative HPLC gave the *title compound* (54 mg, 26%) as a light brown solid. δ_{H} (400 MHz, CD₃OD) 8.31 (d, 1H, *J* 9.8 Hz), 8.24 (d, 2H, *J* 8.2 Hz), 8.07 (d, 2H, *J* 8.2 Hz), 7.98 (d, 1H, *J* 9.8 Hz), 7.43 (d, 1H, *J* 7.5 Hz), 7.34 (t, 1H, *J* 7.9 Hz), 7.21 (t, 2H, *J* 7.8 Hz), 6.81 (t, 1H, *J* 7.4 Hz), 4.72 (s, 2H). LCMS (ES+) 432 (M+H)⁺, RT 1.86 minutes.

25
30

CLAIMS:

1. A compound of formula (IIB) or an *N*-oxide thereof, or a pharmaceutically acceptable salt or solvate thereof:

5




wherein

- E represents -CH₂-;
- 10 V represents C-R²² or N;
- R¹² represents hydrogen or halogen;
- R¹⁵ represents difluoromethoxy;
- R¹⁶ represents hydrogen, fluoro, chloro, cyano, methyl, trifluoromethyl, difluoromethoxy or amino;
- 15 R²¹ represents hydroxy(C₁₋₆)alkyl; or R²¹ represents (C₃₋₇)heterocycloalkyl, which groups may be optionally substituted by one, two or three substituents independently selected from -CO₂H;
- R²² represents hydrogen, halogen or C₁₋₆ alkyl; and
- R²³ represents hydrogen, C₁₋₆ alkyl, trifluoromethyl or C₁₋₆ alkoxy.

20

2. A compound as claimed in claim 1 represented by formula (IIC) or an *N*-oxide thereof, or a pharmaceutically acceptable salt or solvate thereof:



wherein

W represents O or C(R³²)(R³³);

5 R³² represents -CO₂H;

R³³ represents hydrogen; and

E, V, R¹², R¹⁵, R¹⁶ and R²³ are as defined in claim 1.

3. A compound as claimed in claim 1 selected from the following:

- 10 4-[5-(3-{[2-(difluoromethoxy)phenyl]methyl}[1,2,4]triazolo[4,3-*b*]pyridazin-6-yl)-
pyrimidin-2-yl]morpholine; and
1-[5-(3-{[2-(difluoromethoxy)phenyl]methyl}[1,2,4]triazolo[4,3-*b*]pyridazin-6-yl)-
pyrimidin-2-yl]piperidine-4-carboxylic acid.

- 15 4. Use of a compound as defined in any one of claims 1 to 3 or an *N*-oxide thereof,
or a pharmaceutically acceptable salt or solvate thereof, for the treatment or prevention of
a disorder for which the administration of a modulator of TNF α function is indicated.


- 20 5. Use of a compound as defined in any one of claims 1 to 3 or an *N*-oxide thereof,
or a pharmaceutically acceptable salt or solvate thereof, for the treatment or prevention of
an inflammatory disorder, an autoimmune disorder, a neurological disorder, a
neurodegenerative disorder, pain, a nociceptive disorder, a cardiovascular disorder, a
metabolic disorder, an ocular disorder, or an oncological disorder.

- 25 6. A pharmaceutical composition comprising a compound as defined in any one of
claims 1 to 3 or an *N*-oxide thereof, or a pharmaceutically acceptable salt or solvate
thereof, in association with a pharmaceutically acceptable carrier.

7. A pharmaceutical composition as claimed in claim 6 further comprising an additional pharmaceutically active ingredient.

5 8. Use of a compound as defined in any one of claims 1 to 3 or an *N*-oxide thereof, or a pharmaceutically acceptable salt or solvate thereof, in the manufacture of a medicament for the treatment or prevention of a disorder for which the administration of a modulator of TNF α function is indicated.

10 9. Use of a compound as defined in any one of claims 1 to 3 or an *N*-oxide thereof, or a pharmaceutically acceptable salt or solvate thereof, in the manufacture of a medicament for the treatment or prevention of an inflammatory disorder, an autoimmune disorder, a neurological disorder, a neurodegenerative disorder, pain, a nociceptive disorder, a cardiovascular disorder, a metabolic disorder, an ocular disorder, or an oncological disorder.



(I)