

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2012208516 B2**

(54) Title
Use of glu-tubulin as a biomarker of drug response to furazanobenzimidazoles

(51) International Patent Classification(s)
G01N 33/50 (2006.01) **A61K 38/00** (2006.01)
A61K 31/00 (2006.01)

(21) Application No: **2012208516** (22) Date of Filing: **2012.01.19**

(87) WIPO No: **WO12/098203**

(30) Priority Data

(31) Number	(32) Date	(33) Country
11151681.1	2011.01.21	EP

(43) Publication Date: **2012.07.26**

(44) Accepted Journal Date: **2015.11.19**

(71) Applicant(s)
Basilea Pharmaceutica AG

(72) Inventor(s)
Lane, Heidi Alexandra; Bachmann, Felix

(74) Agent / Attorney
Spruson & Ferguson, L 35 St Martins Tower 31 Market St, Sydney, NSW, 2000

(56) Related Art
KREIS, T.E., EMBO J. 1987, Vol.6, pages 2597-2606
MELLI, G. et al., Neurobiol Dis. 2006, Vol.24, pages 525-530
WO 2004/103994
GOKMEN-POLAR, Y. et al., Cancer Res. 2005, Vol.65, pages 9406-9414

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
26 July 2012 (26.07.2012)




(10) International Publication Number
WO 2012/098203 A1

- (51) International Patent Classification:
G01N 33/50 (2006.01) *A61K 38/00* (2006.01)
A61K 31/00 (2006.01)
- (21) International Application Number:
PCT/EP2012/050814
- (22) International Filing Date:
19 January 2012 (19.01.2012)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
11151681.1 21 January 2011 (21.01.2011) EP
- (71) Applicant (for all designated States except US):
BASILEA PHARMACEUTICA AG [CH/CH]; Gren-
zacherstrasse 487, CH-4005 Basel (CH).
- (72) Inventors; and
(75) Inventors/Applicants (for US only): LANE, Heidi Alex-
andra [CH/CH]; Kaltbrunnenweg 8, CH-4106 Therwil
(CH). BACHMANN, Felix [CH/CH]; Grellingerstrasse
38, CH-4052 Basel (CH).
- (74) Agent: BOHEST AG; P.O. Box 160, CH-4003 Basel
(CH).
- (81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD,
SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


[Continued on next page]

(54) Title: USE OF GLU-TUBULIN AS A BIOMARKER OF DRUG RESPONSE TO FURAZANOBENZIMIDAZOLES

Figure 8



(57) Abstract: Use of glu-tubulin as a biomarker for predicting the response to a compound, preferably resistance of a disease such as cancer in a subject to said compound, wherein the compound is a furazanobenzimidazole compound of general formula (I).





(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (*Art. 21(3)*)
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (*Rule 48.2(h)*)
- with sequence listing part of description (*Rule 5.2(a)*)


USE OF GLU-TUBULIN AS A BIOMARKER OF DRUG RESPONSE TO FURAZANOBENZIMIDAZOLES

The present invention relates to use of glu-tubulin as a biomarker for predicting the response of a disease, such as a neoplastic or autoimmune disease, preferably cancer, to a compound of general formula I, such as 3-(4-{1-[2-(4-amino-phenyl)-2-oxo-ethyl]-1H-benzoimidazol-2-yl}-furazan-3-ylamino)-propionitrile (BAL27862). In other aspects it relates to methods and kits, as well as methods of treatment involving the use of the biomarker.

Microtubules are one of the components of the cell cytoskeleton and are composed of heterodimers of alpha and beta tubulin. Agents that target microtubules are among the most effective cytotoxic chemotherapeutic agents having a broad spectrum of activity. Microtubule destabilising agents (e.g. the vinca-alkaloids such as vincristine, vinblastine and vinorelbine) are used for example in the treatment of several types of hematologic malignancies, such as lymphoblastic leukaemia and lymphoma, as well as solid tumours, such as lung cancer. Microtubule stabilising agents (e.g. the taxanes such as paclitaxel, docetaxel) are used for example in the treatment of solid tumours, including breast, lung and prostate cancer.

However resistance to these known microtubule targeting agents can occur. The resistance can either be inherent or can be acquired after exposure to these agents. Such resistance therefore impacts patient survival rates, as well as choices of treatment regimes. Several potential mechanisms of resistance have been identified, and include defects in the microtubule targets, such as elevated levels of beta-tubulin subtype III and acquired mutations in beta-tubulin subtype I that are known to reduce taxane binding. Furthermore, defects in other cell proteins have been suggested to be associated with resistance to certain microtubule targeting agents, such as overexpression of the efflux pump P-glycoprotein (P-gp pump, also known as multi-drug resistance protein 1 or MDR1). Such factors may then be used as biomarkers of resistance to these conventional microtubule targeting agents.

A relatively recently discovered class of microtubule destabilising agents are compounds encompassed by the formula given below:



5 wherein

R represents phenyl, thienyl or pyridinyl

wherein phenyl is optionally substituted by one or two substituents independently selected from alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, acyloxy-lower alkyl, phenyl, hydroxy, lower alkoxy, hydroxy-lower alkoxy, lower

10 alkoxy-lower alkoxy, phenyl-lower alkoxy, lower alkylcarbonyloxy, amino, monoalkylamino, dialkylamino, lower alkoxycarbonylamino, lower alkylcarbonylamino, substituted amino wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, carboxy, lower alkoxycarbonyl, cyano, halogen, and nitro; and wherein two adjacent substituents are

15 methylenedioxy;

X represents a group C=Y, wherein Y stands for oxygen or nitrogen substituted by hydroxy or lower alkoxy;

20

R¹ represents hydrogen, lower alkylcarbonyl, hydroxy-lower alkyl or cyano-lower alkyl;

R², R³ and R⁶ represent hydrogen;

R⁴ and R⁵, independently of each other, represent hydrogen, lower alkyl or lower

25

alkoxy;

or R⁴ and R⁵ together represent methylenedioxy;

and pharmaceutically acceptable salts thereof;

or wherein

5 R represents phenyl or pyridinyl

wherein phenyl is optionally substituted by one or two substituents independently selected from alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, acyloxy-lower alkyl, phenyl, hydroxy, lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, phenyl-lower alkoxy, lower alkylcarbonyloxy, amino,

10 monoalkylamino, dialkylamino, lower alkoxy-carbonylamino, lower alkylcarbonylamino, substituted amino wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, carboxy, lower alkoxy-carbonyl, formyl, cyano, halogen, and nitro; and wherein two adjacent substituents are methylenedioxy;

15 and wherein pyridinyl is optionally substituted by lower alkoxy, amino or halogen;

X represents oxygen;

20 R¹ represents hydrogen, lower alkylcarbonyl, hydroxy-lower alkyl or cyano-lower alkyl;

R², R³ and R⁶ represent hydrogen;

R⁴ and R⁵, independently of each other, represent hydrogen, lower alkyl or lower alkoxy;

or R⁴ and R⁵ together represent methylenedioxy;


25 and pharmaceutically acceptable salts thereof;

and wherein the prefix lower denotes a radical having up to and including a maximum of 7, especially up to and including a maximum of 4 carbon atoms.

These compounds are disclosed in WO2004/103994 A1, which is incorporated by cross-reference herein. These compounds have been shown to
30 arrest tumour cell proliferation and induce apoptosis.

The synthesis of compounds of formula I is described in WO2004/103994 A1, in general on pages 29-35, and specifically on pages 39-55, which are incorporated herein by cross-reference. They may be prepared as disclosed or by an analogous method to the processes described therein.

- 5 One compound falling within this class, known as BAL27862, and shown in WO2004/103994 A1 as example 58, and specifically incorporated by reference herein, has the structure and chemical name given below:




Chemical name:

- 10 3-(4-{1-[2-(4-Amino-phenyl)-2-oxo-ethyl]-1H-benzoimidazol-2-yl}-furazan-3-ylamino)-propionitrile.

Or herein as Compound A


- Further compounds exemplified in WO2004/103994 A1 as examples 50 and 79 respectively, and also specifically incorporated by cross-reference herein, have
15 the structures and chemical names given below:



Chemical name: 2-[2-(4-Amino-furazan-3-yl)-benzoimidazol-1-yl]-1-(4-amino-phenyl)-ethanone

or herein as Compound B

5 and



Chemical name: 3-(4-{1-[2-(6-Amino-pyridin-3-yl)-2-oxo-ethyl]-1H-benzoimidazol-2-yl}-furazan-3-ylamino)-propionitrile

or herein as Compound C.

10 BAL27862 has activity across a broad panel of experimental, solid tumour xenograft models. Moreover, activity is retained even against tumour models which were selected for resistance to conventional microtubule targeting agents (including the vinca-alkaloid microtubule destabilisers and the microtubule stabilisers paclitaxel and epothilone B). BAL27862 activity is not affected by over-expression of the P-gp

pump in any models tested *in vitro*, nor in human mammary tumour xenografts. Additionally, BAL27862 retained its activity despite elevated levels of beta-tubulin subtype III and mutations in tubulin subtype I.

Hence, BAL27862 activity is not affected by a number of factors that confer resistance to conventional microtubule targeting agents.

Moreover, it is known that compounds of general formula I have a different effect on the phenotype of cells compared to other microtubule targeting agents, including other microtubule destabilisers. Treatment with a compound of general formula I induces a consistent microtubule phenotype in tumour cell lines derived from a variety of organs, for example lung, cervix and breast, as seen in Figure 1. Staining the microtubules in these cells with an anti-alpha-tubulin antibody shows that rather than the mitotic spindle fibres of untreated cells, only dot-like structures are visible in the treated cells. This same effect is also shown using Compounds C and B in Figures 2A and 2B respectively on the lung cancer cell line A549. It is however very distinct from that observed with the conventional microtubule targeting agents vinblastine, colchicine, paclitaxel and nocodazole as seen in Figures 3B, 3C, 3D and 4, respectively. The microtubules were stained with an anti-alpha-tubulin antibody and the cells viewed at a 1000 x magnification (Figures 3, 4). For the cells treated with BAL27862, multiple dot-like structures are visible, whereas, in stark contrast, the other conventional drugs produce filamentous microtubule structures, or dense microtubule aggregate structures. These differences at the phenotypic level, at compound doses considered optimal in terms of antiproliferative effect indicate a difference in the mode of action at the molecular level.

Furthermore, it is known that BAL27862 elicits a dominant microtubule phenotype in the presence of the other microtubule targeting agents. Treatment with vinblastine, colchicine, paclitaxel or nocodazole alone induced the microtubule phenotypes characteristic of these agents (Figure 5A, 5D, 5G, 6C-6F respectively). However, combination treatment with BAL27862 for the last 4 hours resulted in disruption of these phenotypes; despite the continued presence of vinblastine, colchicine, paclitaxel, or nocodazole (Figure 5B, 5E, 5H, 6G-6J respectively). In contrast, treating first with BAL27862 and subsequently for 4 hours in combination

with vinblastine, colchicine, paclitaxel, or nocodazole had no impact on generation of the phenotype consistent with BAL27862 treatment (Figure 5C, 5F, 5I, 6K-6N respectively).


These data all demonstrate that BAL27862 affects microtubule biology in a different manner than conventional microtubule targeting agents.

Thus, from information about conventional microtubule targeting agents, predictions cannot be made concerning if, or how, particular genes are involved in the action of compounds of formula I.

An object of the present invention is to identify factors which are associated with response to compounds of formula I or pharmaceutically acceptable derivatives thereof, for example to identify factors associated with resistance to compounds of general formula I, in particular BAL27862 or pharmaceutically acceptable derivatives thereof, as defined below.

It has surprisingly been found that glu-tubulin may be used as a biomarker of response to treatment with a compound of general formula I or pharmaceutically acceptable derivatives thereof, as defined below.

According to a first aspect of the present invention there is provided use of glu-tubulin as a biomarker for predicting the response of a neoplastic disease or an autoimmune disease to a compound, wherein the compound is a compound of general formula I



wherein

R represents phenyl, thienyl or pyridinyl

wherein phenyl is optionally substituted by one or two substituents independently selected from alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, acyloxy-lower alkyl, phenyl, hydroxy, lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, phenyl-

lower alkoxy, lower alkylcarbonyloxy, amino, monoalkylamino, dialkylamino, lower alkoxy-carbonylamino, lower alkylcarbonylamino, substituted amino wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, carboxy, lower alkoxy-carbonyl, cyano, halogen, and nitro; and wherein two adjacent substituents are methylenedioxy;

and wherein pyridinyl is optionally substituted by lower alkoxy, amino or halogen;

X represents a group C=Y, wherein Y stands for oxygen or nitrogen substituted by hydroxy or lower alkoxy;

R¹ represents hydrogen, lower alkylcarbonyl, hydroxy-lower alkyl or cyano-lower alkyl;

R², R³ and R⁶ represent hydrogen;

R⁴ and R⁵, independently of each other, represent hydrogen, lower alkyl or lower alkoxy; or R⁴ and R⁵ together represent methylenedioxy;

and pharmaceutically acceptable derivatives thereof;

or wherein

R represents phenyl or pyridinyl

wherein phenyl is optionally substituted by one or two substituents independently selected from alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, acyloxy-lower alkyl, phenyl, hydroxy, lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, phenyl-lower alkoxy, lower alkylcarbonyloxy, amino, monoalkylamino, dialkylamino, lower alkoxy-carbonylamino, lower alkylcarbonylamino, substituted amino wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, carboxy, lower alkoxy-carbonyl, formyl, cyano, halogen, and nitro; and wherein two adjacent substituents are methylenedioxy;

and wherein pyridinyl is optionally substituted by lower alkoxy, amino or halogen;

X represents oxygen;

R¹ represents hydrogen, lower alkylcarbonyl, hydroxy-lower alkyl or cyano-lower alkyl;

R², R³ and R⁶ represent hydrogen;

R⁴ and R⁵, independently of each other, represent hydrogen, lower alkyl or lower alkoxy; or R⁴ and R⁵ together represent methylenedioxy;

and pharmaceutically acceptable derivatives thereof,


and wherein the prefix lower denotes a radical having up to and including a maximum of 7, especially up to and including a maximum of 4 carbon atoms.

According to a second aspect of the present invention there is provided the use according to the first aspect above, for predicting the resistance of a disease in a subject to said compound.


According to a third aspect of the present invention there is provided a method for predicting the response of a neoplastic disease or an autoimmune disease in a subject to a compound of general formula I or a pharmaceutically acceptable derivative thereof as defined in the first aspect above, comprising the steps of:

- a) measuring a level of glu-tubulin in a sample pre-obtained from the subject to obtain a value or values representing this level; and
- b) comparing the value or values from step a) to a standard value or set of standard values.

According to a fourth aspect of the present invention there is provided a method of treating a neoplastic or autoimmune disease, preferably cancer, in a subject in need thereof, comprising measuring a level of glu-tubulin in a sample from the subject to obtain a value or values representing this level, and treating the subject with a compound of general formula I or a pharmaceutically acceptable derivative thereof wherein the compound is




or




if the level of glu-tubulin in said sample is not higher than a standard value or set of standard values.

According to a fifth aspect of the present invention there is provided a method of treating a neoplastic or autoimmune disease, preferably cancer, by first decreasing the level of glu-tubulin in a subject that has a sample with a higher level of glu-tubulin compared to a standard level or set of standard levels, then treating the subject with a compound of general formula I or a pharmaceutically acceptable derivative thereof wherein the compound is



or



According to a sixth aspect of the present invention there is provided a kit when used for predicting the response of a cancer to a compound of general formula I or a pharmaceutically acceptable derivative thereof, as defined according to the first aspect above, comprising reagents necessary for measuring a level of glu-tubulin in a sample and further comprising a comparator module which comprises a standard value or set of standard values of a level of glu-tubulin taken from samples of tumour tissue or circulating tumour cells of subjects with a cancer of the same histotype to which the level of glu-tubulin in the sample is compared.

According to a seventh aspect of the present invention there is provided a device for predicting the response of a neoplastic disease or an autoimmune disease to a compound of general formula I or a pharmaceutically acceptable derivative thereof, as defined in the first aspect above, comprising reagents necessary for measuring the level of glu-tubulin in a sample and a comparator module which comprises a standard value or set of standard values to which the level of glu-tubulin in the sample is compared.

In one preferred embodiment of the invention, relatively high glu-tubulin levels in a tumour sample are associated with inherent and acquired resistance to BAL27862, as described below.


Tubulin is subjected to a variety of posttranslational modifications, including detyrosination/tyrosination, acetylation, glutamylation, polyglycylation, phosphorylation of serine residues and phosphorylation of tyrosine residues, making it one of the most modified

proteins known. Many alpha tubulin genes described to date predict the presence of a carboxy terminal tyrosine on tubulin. It is currently thought that once tubulin polymerizes and assembles into microtubules, in some cases a carboxypeptidase (tubulin carboxypeptidase or tubulin tyrosine carboxypeptidase, TTCP) acts to remove the C-terminal tyrosine from alpha tubulin exposing the penultimate glutamate. Another enzyme, tubulin tyrosine ligase (TTL),

can reattach this tyrosine. The exact function of this detyrosination/tyrosination cycle in the cell has yet to be elucidated.

The form of alpha tubulin which does not have a tyrosine, but rather a glutamate at its C-terminal is known as glu-tubulin or glu tubulin or detyrosinated tubulin. The term glu-tubulin shall be used herein to refer to the form of alpha tubulin with a glutamate as the final amino acid at the C-terminal. The designation glu-tubulin, shall also encompass forms wherein other post-translational modifications may additionally be present. The alpha tubulin which forms the basis for the glu-tubulin is known to exist in multiple variants, subtypes and isoforms, as well as there being multiple alpha tubulin genes which give rise to these, and all these shall also be included in the designation glu-tubulin, with the proviso that a glutamate is the final amino acid at the C-terminal. Preferably it relates to human variants, subtypes and isoforms of alpha tubulin, with the proviso that a glutamate is the final amino acid at the C-terminal. In a more preferred embodiment, a glutamate, not a tyrosine, is the final amino acid at the C-terminus of alpha tubulin after post-translational modification. Subtypes of alpha tubulin include, but are not limited to, tubulin alpha 1A, tubulin alpha 1B, tubulin alpha 1C and tubulin alpha 3C/D. The protein sequences of these subtypes are accessible via the following National Center for Biotechnology Information (NCBI) Reference numbers NP_006000, NP_006073, NP_116093 and NP_005992, respectively. These are also listed in Figures 11-14 (NP_006000.2, NP_006073.2, NP_116093.1, NP_005992.1,) as SEQ ID NO. 1-4, respectively. Glu-tubulin, as described above, does not have the tyrosine at the final amino acid at the C-terminus of the sequences presented in SEQ. ID. NO. 1-4.

One aspect of the present invention relates to use of glu-tubulin as a biomarker for predicting the response to a compound, wherein the compound is a compound of general formula I,



wherein

R represents phenyl, thienyl or pyridinyl

wherein phenyl is optionally substituted by one or two substituents independently selected from alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, acyloxy-lower alkyl, phenyl, hydroxy, lower alkoxy, hydroxy-lower alkoxy, lower

alkoxy-lower alkoxy, phenyl-lower alkoxy, lower alkylcarbonyloxy, amino, monoalkylamino, dialkylamino, lower alkoxy-carbonylamino, lower alkylcarbonylamino, substituted amino wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, carboxy, lower alkoxy-carbonyl, cyano, halogen, and nitro; and wherein two adjacent substituents are methylenedioxy;

and wherein pyridinyl is optionally substituted by lower alkoxy, amino or halogen;

X represents a group C=Y, wherein Y stands for oxygen or nitrogen substituted by hydroxy or lower alkoxy;

R¹ represents hydrogen, lower alkylcarbonyl, hydroxy-lower alkyl or cyano-lower alkyl;

R², R³ and R⁶ represent hydrogen;

R⁴ and R⁵, independently of each other, represent hydrogen, lower alkyl or lower alkoxy;

or R⁴ and R⁵ together represent methylenedioxy;

and pharmaceutically acceptable derivatives thereof,

or wherein

R represents phenyl or pyridinyl

wherein phenyl is optionally substituted by one or two substituents independently selected from alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, acyloxy-lower alkyl, phenyl, hydroxy, lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, phenyl-lower alkoxy, lower alkylcarbonyloxy, amino, 5 monoalkylamino, dialkylamino, lower alkoxycarbonylamino, lower alkylcarbonylamino, substituted amino wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, carboxy, lower alkoxycarbonyl, formyl, cyano, halogen, and nitro; and wherein two adjacent substituents are methylenedioxy;

10 and wherein pyridinyl is optionally substituted by lower alkoxy, amino or halogen;

X represents oxygen;

R¹ represents hydrogen, lower alkylcarbonyl, hydroxy-lower alkyl or cyano-lower 15 alkyl;

R², R³ and R⁶ represent hydrogen;

R⁴ and R⁵, independently of each other, represent hydrogen, lower alkyl or lower alkoxy;

or R⁴ and R⁵ together represent methylenedioxy;

20 and pharmaceutically acceptable derivatives thereof;

and wherein the prefix lower denotes a radical having up to and including a maximum of 7, especially up to and including a maximum of 4 carbon atoms.

Preferably the response may be of a disease in a subject. Also preferably the response may be to treatment, i.e. to treatment with the compound of general 25 formula I or pharmaceutically acceptable derivatives thereof.


The biomarker glu-tubulin is measured *ex vivo* in a sample or samples taken from the human or animal body, preferably taken from the human body.

In a preferred embodiment, the invention relates to use of glu-tubulin as a biomarker for predicting the resistance of a disease in a subject to a compound of

general formula I or pharmaceutically acceptable derivatives thereof as defined above.

Preferably the pharmaceutically acceptable derivative is selected from the group consisting of a salt, solvate, pro-drug, salt of a pro-drug, polymorph and isomer of a compound of general formula I as defined above. Pro-drugs are preferably ester and amides of naturally occurring amino acids, small peptides or pegylated hydroxy acids. More preferably, the pro-drug is an amide formed from an amino group present within the R group of the compound of general formula I and the carboxy group of glycine, alanine or lysine.

Particularly preferably the compound is



or a pharmaceutically acceptable salt thereof, preferably a hydrochloride salt thereof, most preferably a dihydrochloride salt thereof.

Another aspect of the present invention relates to a method for predicting the response of a disease in a subject to a compound of general formula I or pharmaceutically acceptable derivatives thereof as defined above, comprising the steps of:

a) measuring a level of glu-tubulin in a sample pre-obtained from the subject

to obtain a value or values representing this level; and

- b) comparing the value or values from step a) to a standard value or set of standard values.

Further preferably the response which is predicted is resistance.

5 The measuring of a level or levels of glu-tubulin is performed ex-vivo in a sample or samples pre-obtained from the subject. Pre-obtained refers to the fact that the sample is obtained before it is subjected to any method involving measuring the level of the biomarker, and pre-obtained is not to be understood as in relation to treatment. In a preferred embodiment, a higher level of glu-tubulin in the sample from
10 the subject relative to the standard value or set of standard values predicts resistance.

Also preferably, the disease is a neoplastic or autoimmune disease. More preferably the disease is cancer. Especially preferably, the cancer is selected from the group consisting of breast cancer, prostate cancer, cervical cancer, ovarian
15 cancer, gastric cancer, colorectal cancer (i.e. including colon cancer and rectal cancer), pancreatic cancer, liver cancer, brain cancer, neuroendocrine cancer, lung cancer, kidney cancer, hematological malignancies, melanoma and sarcomas. More especially preferably the cancer is selected from the group consisting of breast cancer, cervical cancer, ovarian cancer, colorectal cancer, melanoma and lung
20 cancer. In an especially preferred embodiment the cancer is selected from the group consisting of lung cancer, melanoma, ovarian cancer and colorectal cancer. In a particularly preferred embodiment, wherein inherent resistance is determined, the cancer is selected from the group consisting of lung cancer, melanoma or colorectal cancer. In another particularly preferred embodiment, wherein acquired resistance is
25 determined, the cancer is lung or ovarian cancer.

In a further aspect, the invention relates to a method of treating a neoplastic or autoimmune disease, preferably cancer, in a subject in need thereof, comprising measuring a level of glu-tubulin in a sample from the subject to obtain a value or values representing this level, and treating the subject with a compound of general

formula I or a pharmaceutically acceptable derivative thereof as defined above, if the level of glu-tubulin in said sample is not higher than a standard value or set of standard values.


5 In yet a further aspect, the invention relates to glu-tubulin for use in the treatment of a neoplastic or autoimmune disease, preferably cancer, comprising measuring a level of glu-tubulin in a sample from the subject to obtain a value or values representing this level, and treating the subject with a compound of general formula I or a pharmaceutically acceptable derivative thereof as defined above, if the level of glu-tubulin is not higher than a standard value or set of standard values.

10 The measuring of a level of glu-tubulin is performed ex-vivo in a sample pre-obtained from the subject.

The invention also relates in another aspect to a method of treating a neoplastic or autoimmune disease, preferably cancer, by first decreasing the level of glu-tubulin in a subject that has a sample with a higher level of glu-tubulin compared to a standard level or set of standard levels, then treating the subject with a compound of general formula I or a pharmaceutically acceptable derivative thereof as defined above.

15 In yet another aspect the invention relates to a kit for predicting the response to a compound of general formula I or a pharmaceutically acceptable derivative thereof, as defined above, comprising reagents necessary for measuring the level of glu-tubulin in a sample. More preferably the kit also comprises a comparator module which comprises a standard value or set of standard values to which the level of glu-tubulin in the sample is compared.

20 More preferably the kit comprises a compound of general formula I or a pharmaceutically acceptable derivative thereof as defined above. In an especially preferred embodiment the kit comprises a compound of the following formula or a pharmaceutically acceptable salt thereof



Chemical name: S-2,6-Diamino-hexanoic acid [4-(2-{4-(2-cyano-ethylamino)-furazan-3-yl}-benzoimidazol-1-yl)-acetyl]-phenyl]-amide

5 In a particularly preferred embodiment the pharmaceutically acceptable salt is a dihydrochloride salt.

Another further aspect of the invention relates to a device for predicting the response to a compound of general formula I or a pharmaceutically acceptable derivative thereof as defined above, comprising reagents necessary for measuring a level of glu-tubulin in a sample and a comparator module which comprises a standard value or set of standard values to which the level of glu-tubulin in the sample is compared.

In a preferred embodiment, the reagents in the kit or device comprise a capture reagent comprising a detector for glu-tubulin, and a detector reagent. Especially preferably the capture reagent is an antibody. Also preferably, the disease is predicted to be resistant to treatment with said compound when glu-tubulin is higher relative to a standard value or set of standard values. In a preferred embodiment, the comparator module is included in instructions for use of the kit. In another preferred embodiment the comparator module is in the form of a display device.

Embodiments of the present invention will now be described by way of example with reference to the accompanying figures. The invention however is not to be understood as limited to these embodiments.

Brief Description of the Figures

5 Figure 1: Shows the treatment of human tumour cell lines from different histotypes with 50 nM BAL27862. The microtubules of mitotic or G2/M arrested cells were stained after 24 hours treatment with 50 nM BAL27862 or vehicle control.

Fig. 1A and 1B: A549 NSCLC cells;

Fig. 1C and 1D: HeLa cervical cancer cells;

10 Fig. 1E and 1F: SKBR3 breast cancer cells

Vehicle control treatment: Figures 1A, 1C & 1E,

BAL27862 treatment: Figures 1B, 1D & 1F.

15 Figure 2: Shows the treatment of A549 NSCLC cells with the Compounds B and C. The microtubules of mitotic or G2/M arrested A549 NSCLC cells were stained after 24 hours treatment with 80 nM or 20 nM of Compounds B and C, respectively. The white scale bar represents 10 micrometres.

Fig. 2A: treatment with 20 nM Compound C

20 Fig. 2B: treatment with 80 nM Compound B

25 Figure 3: Shows a comparison of treatment of cells with BAL27862 compared to conventional microtubule targeting agents. Microtubules of mitotic or G2/M arrested A549 NSCLC cells were stained after 24 hours of treatment with 50 nM of A: BAL27862; B: vinblastine; C: colchicine; D: paclitaxel. Stacks of images taken every 1 μ m were processed by using ImageJ software.

30 Figure 4: Shows a comparison of treatment of A549 NSCLC cells with BAL27862 compared to nocodazole. Microtubules of mitotic or G2/M arrested cells were stained after 24 h of treatment with various concentrations of nocodazole (B, C

& D) and BAL27862 (E, F & G). A: control, B: Nocodazole 50 nM, C: Nocodazole 100 nM, D: Nocodazole 200 nM, E: BAL27862 20 nM; F: BAL27862 30 nM and G: BAL27862 50 nM. The white scale bar represents 10 micrometres. Representative images of the microtubule phenotypes observed are shown.

5

Figure 5: Shows a combination of treatment with BAL27862 and conventional microtubule-targeting agents. Microtubules of mitotic or G2/M arrested A549 NSCLC cells were stained after treatment for the times indicated below. 50 nM BAL27862, 50 nM vinblastine, 50 nM colchicine and 25 nM paclitaxel were used. The white scale bar represents 10 micrometres.

10

Fig. 5A: 24 hours vinblastine treatment;

Fig. 5B: 24 hours vinblastine treatment with the final 4 hours including BAL27862;

15

Fig. 5C: 24 hours BAL27862 treatment with the final 4 hours including vinblastine.

Fig. 5D: 24 hours colchicine treatment;

Fig. 5E: 24 hours colchicine treatment with the final 4 hours including BAL27862;

20

Fig. 5F: 24 hours BAL27862 treatment with the final 4 hours including colchicine.

Fig. 5G: 24 hours paclitaxel treatment;

Fig. 5H: 24 hours paclitaxel treatment with the final 4 hours including BAL27862;

25

Fig. 5I: 24 hours BAL27862 treatment with the final 4 hours including paclitaxel.

Figure 6: Shows a combination of treatment with BAL27862 and nocodazole. Microtubules of mitotic or G2/M arrested A549 NSCLC cells were stained after treatment for the times indicated below. 25 nM BAL27862 and nocodazole at the concentrations indicated below were used. The white scale bar represents 10 micrometers.

30

Fig. 6A: 24 hours control treatment;

Fig. 6B: 24 hours of 25 nM BAL27862 treatment;

Fig. 6C: 24 hours of 50 nM nocodazole treatment

Fig. 6D: 24 hours of 100 nM nocodazole treatment

Fig. 6E: 24 hours of 150 nM nocodazole treatment

5 Fig. 6F: 24 hours of 200 nM nocodazole treatment

Fig. 6G: 24 hours of 50 nM nocodazole treatment with the final 4 hours including 25 nM BAL27862;

Fig. 6H: 24 hours of 100 nM nocodazole treatment with the final 4 hours including 25 nM BAL27862;

10 Fig. 6I: 24 hours of 150 nM nocodazole treatment with the final 4 hours including 25 nM BAL27862;

Fig. 6J: 24 hours of 200 nM nocodazole treatment with the final 4 hours including 25 nM BAL27862;

15 Fig. 6K: 24 hours of 25 nM BAL27862 treatment with the final 4 hours including 50 nM nocodazole;

Fig. 6L: 24 hours of 25 nM BAL27862 treatment with the final 4 hours including 100 nM nocodazole;

Fig. 6M: 24 hours of 25 nM BAL27862 treatment with the final 4 hours including 150 nM nocodazole;

20 Fig. 6N: 24 hours of 25 nM BAL27862 treatment with the final 4 hours including 200 nM nocodazole.

Figure 7: Shows protein extracts prepared from patient-derived colorectal cancer (Fig. 7A), lung cancer (Fig. 7B) and melanoma (Fig. 7C) tumours obtained from subcutaneous xenografted mice, and analysed by immunoblotting for glu-tubulin expression, with actin included as a loading control. Three independent tumours were analysed in each case (1 – 3). BAL27862, paclitaxel and vinblastine resistance and sensitivity is as defined in Table 1.

Figure 8: Shows by immunohistochemistry that glu-tubulin levels in tumour cells are increased in a patient-derived xenografted colorectal tumour defined as BAL27862 resistant by *ex vivo* colony outgrowth analysis. Patient-derived tumour xenografts (maintained in nude mice) were prepared, fixed and stained for glu-tubulin

protein expression using immunohistochemistry. BAL27862, paclitaxel and vinblastine resistance and sensitivity is defined in Table 1.

Figure 9: Shows tumour cell lines which were selected for resistance to BAL27862 through *in vitro* cultivation in the presence of the compound. Based on IC₅₀ determinations, BAL27862 resistance factors *versus* parental lines were: A549 (3.0 fold); SKOV3 (7.6 fold – resistant 1 line); H460 (5.3 fold) (see Table 2). Whole cell protein extracts were prepared from parental and resistant lines and analysed by immunoblotting for glu-tubulin expression. Actin levels were included as a loading control.

Figure 10: Shows that increased glu-tubulin protein levels are maintained in SKOV3 tumour lines during resistance development. SKOV3 tumour cell lines were selected for resistance to BAL27862 through *in vitro* cultivation in the presence of BAL27862 for increasing time periods. Based on IC₅₀ determinations, BAL27862 resistance factors *versus* parental lines were: SKOV3 resistant 1 (7.6 fold), SKOV3 resistant 2 (11.6 fold) (see Table 2). Whole cell protein extracts were prepared from parental and resistant lines and analysed by immunoblot for glu-tubulin expression. Actin levels act as a loading control.

Figure 11: Shows the protein sequence of tubulin alpha-1A chain [Homo sapiens] (SEQ. ID. No. 1)


Figure 12: Shows the protein sequence of tubulin alpha-1B chain [Homo sapiens] (SEQ ID No. 2)

Figure 13: Shows the protein sequence of tubulin alpha-1C chain [Homo sapiens] (SEQ. ID. NO. 3)

Figure 14: Shows the protein sequence of Tubulin alpha-3C/D chain [Homo sapiens] (SEQ. ID. NO. 4)

Detailed DescriptionCompounds of formula I

The compounds according to the invention are represented by general formula I:



5

wherein

R represents phenyl, thienyl or pyridinyl

wherein phenyl is optionally substituted by one or two substituents independently selected from alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl,

10 acyloxy-lower alkyl, phenyl, hydroxy, lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, phenyl-lower alkoxy, lower alkylcarbonyloxy, amino, monoalkylamino, dialkylamino, lower alkoxycarbonylamino, lower alkylcarbonylamino, substituted amino wherein the two substituents on nitrogen form together with the nitrogen heterocycl, lower alkylcarbonyl, carboxy, lower

15 alkoxycarbonyl, cyano, halogen, and nitro; and wherein two adjacent substituents are methylenedioxy;

and wherein pyridinyl is optionally substituted by lower alkoxy, amino or halogen;

X represents a group C=Y, wherein Y stands for oxygen or nitrogen substituted by

20 hydroxy or lower alkoxy;

R¹ represents hydrogen, lower alkylcarbonyl, hydroxy-lower alkyl or cyano-lower alkyl;

R², R³ and R⁶ represent hydrogen;

R^4 and R^5 , independently of each other, represent hydrogen, lower alkyl or lower alkoxy;

or R^4 and R^5 together represent methylenedioxy;

5 and pharmaceutically acceptable derivatives thereof,

or wherein

R represents phenyl or pyridinyl

wherein phenyl is optionally substituted by one or two substituents independently

10 selected from alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, acyloxy-lower alkyl, phenyl, hydroxy, lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, phenyl-lower alkoxy, lower alkylcarbonyloxy, amino, monoalkylamino, dialkylamino, lower alkoxycarbonylamino, lower alkylcarbonylamino, substituted amino wherein the two substituents on nitrogen form
15 together with the nitrogen heterocyclyl, lower alkylcarbonyl, carboxy, lower alkoxycarbonyl, formyl, cyano, halogen, and nitro; and wherein two adjacent substituents are methylenedioxy;
and wherein pyridinyl is optionally substituted by lower alkoxy, amino or halogen;

20 X represents oxygen;

R^1 represents hydrogen, lower alkylcarbonyl, hydroxy-lower alkyl or cyano-lower alkyl;

R^2 , R^3 and R^6 represent hydrogen;

25 R^4 and R^5 , independently of each other, represent hydrogen, lower alkyl or lower alkoxy;

or R^4 and R^5 together represent methylenedioxy;

and pharmaceutically acceptable derivatives thereof;

30 and wherein the prefix lower denotes a radical having up to and including a maximum of 7, especially up to and including a maximum of 4 carbon atoms.

Heterocyclyl designates preferably a saturated, partially saturated or

unsaturated, mono- or bicyclic ring containing 4-10 atoms comprising one, two or three heteroatoms selected from nitrogen, oxygen and sulfur, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a ring nitrogen atom may optionally be substituted by a group selected from lower alkyl, amino-lower alkyl, aryl, aryl-lower alkyl and acyl, and a ring carbon atom may be substituted by lower alkyl, amino-lower alkyl, aryl, aryl-lower alkyl, heteroaryl, lower alkoxy, hydroxy or oxo. Examples of heterocyclyl are pyrrolidinyl, oxazolidinyl, thiazolidinyl, piperidinyl, morpholinyl, piperazinyl, dioxolanyl and tetrahydropyranyl.


Acyl designates, for example, alkylcarbonyl, cyclohexylcarbonyl, arylcarbonyl, aryl-lower alkylcarbonyl, or heteroarylcarbonyl. Lower acyl is preferably lower alkylcarbonyl, in particular propionyl or acetyl.

Preferably, the compound of general formula I according to the invention is defined as wherein R¹ is selected from the group consisting of hydrogen, acetyl, CH₂CH₂CN and CH₂CH₂CH₂OH.

In one preferred embodiment, the compound of general formula I according to the invention is selected from the group consisting of:

4-(1-Phenacyl-1H-benzimidazol-2-yl)-furazan-3-ylamine,
4-[1-(4-Bromophenacyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine oxime,
N-{4-[1-(4-Chlorophenacyl)-1H-benzimidazol-2-yl]-furazan-3-yl}-acetamide,
4-[1-(4-Chlorophenacyl)-1H-benzimidazol-2-yl]-furazan-3-yl-N-(2-cyanoethyl)-amine
4-[1-(4-Chlorophenacyl)-1H-benzimidazol-2-yl]-furazan-3-yl-N-(3-hydroxypropyl)-amine,
4-[1-(3-Amino-4-chlorophenacyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine
4-[1-(3-Methoxy-4-methoxymethoxy-phenacyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine,
and pharmaceutically acceptable derivatives thereof.













In another preferred embodiment, the compound of general formula I according to the invention is selected from the group consisting of:


























wherein











5 R, Y and R¹ are defined as follows :








R	Y	R ¹
	O	H
	NOH	H
	NOMe	H
	O	H
	NOH	H
	NOH	H
	NOMe	H
	O	H

	NOH	H
	NOMe	H
	O	H
	NOH	H
	NOMe	H
	O	H
	NOMe	H
	O	H
	O	H
	NOH	H
	NOMe	H
	O	H

	NOH	H
	NOMe	H
	NOMe	H
	O	H
	O	Ac
	O	H
	O	H
	O	H
	O	CH ₂ CH ₂ CN
	O	CH ₂ CH ₂ CN
	O	H

	O	H
	O	CH ₂ CH ₂ CH ₂ OH
	O	H
	O	CH ₂ CH ₂ CN
	O	H
	O	CH ₂ CH ₂ CN
	O	CH ₂ CH ₂ CN
	O	CH ₂ CH ₂ CN
	O	H
	O	H
	O	H
	O	H

	O	H
	O	H
	O	H
	O	H
	O	CH ₂ CH ₂ CN
	O	H
	O	H
	O	H
	O	H
	O	H
	O	H


	O	H
	O	H
	O	H
	O	CH ₂ CH ₂ CN
	O	H
	O	H
	O	CH ₂ CH ₂ CN

or pharmaceutically acceptable derivatives thereof.

In yet another preferred embodiment, the compound of general formula I according to the invention is selected from the group consisting of:












- 5 4-(1-Phenoxymethyl-1H-benzimidazol-2-yl)-furazan-3-ylamine,
- 4-[1-(4-Fluorophenoxymethyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine,
- 4-[1-(3,4-Dimethylphenoxymethyl)-1H-benzimidazol-2-yl]-furazan-3-yl-N-(2-
- cyanoethyl)-amine

and compounds represented by the formula:




wherein R and R¹ are as defined below

R	R ¹
	H
	H
	H
	H
	H
	CH ₂ CH ₂ CN
	CH ₂ CH ₂ CN
	CH ₂ CH ₂ CN
	H

	H
	H
	H
	H
	H
	H
	H
	CH ₂ CH ₂ CN
	CH ₂ CH ₂ CH ₂ OH
	H
	H




and pharmaceutically acceptable derivatives thereof.

In still yet another preferred embodiment the compound of general formula I according to the invention is:




5 wherein R, R⁴ and R⁵ are as defined below

R	R ⁴	R ⁵
	Me	Me
	Me	Me
	Me	Me
	Me	Me
	Me	Me
	OMe	OMe
	OMe	OMe

	OMe	OMe
	OMe	OMe
	OMe	OMe

or pharmaceutically acceptable derivatives thereof.

More preferably, the compound according to the invention is a compound of general formula I



5

wherein

R represents phenyl or pyridinyl

wherein phenyl is optionally substituted by one or two substituents independently

10 selected from lower alkyl, lower alkoxy, amino, acetylamino, halogen and nitro;

and wherein pyridinyl is optionally substituted by amino or halogen;

X represents a group C=O;


15 R¹ represents hydrogen or cyano-lower alkyl;

R², R³, R⁴, R⁵ and R⁶ represent hydrogen;

and pharmaceutically acceptable derivatives thereof,

and wherein the prefix lower denotes a radical having up to and including a maximum of 7, especially up to and including a maximum of 4 carbon atoms.

Especially preferably, the compound according to the invention is represented by the following formula



5


wherein R, Y and R¹ are defined as follows :

R	Y	R ¹
	O	H
	O	CH ₂ CH ₂ CN
	O	H
	O	CH ₂ CH ₂ CN




or pharmaceutically acceptable derivatives thereof.

10

More especially preferably, the compound according to the invention is represented by the following formula




wherein R, Y and R1 are defined as follows:

R	Y	R1
	O	CH ₂ CH ₂ CN
	O	H
	O	CH ₂ CH ₂ CN

or pharmaceutically acceptable derivatives thereof.

Particularly preferably, the compound according to the invention is



5 or pharmaceutically acceptable derivatives thereof.

The term derivative or derivatives in the phrase “pharmaceutically acceptable derivative” or “pharmaceutically acceptable derivatives” of compounds of general formula I relates to salts, solvates and complexes thereof and to solvates and complexes of salts thereof, as well as to pro-drugs, polymorphs, and isomers thereof (including optical, geometric and tautomeric isomers) and also salts of pro-drugs thereof. In a more preferred embodiment, it relates to salts and pro-drugs, as well as to salts of pro-drugs thereof.

10


Salts are preferably acid addition salts. Salts are formed, preferably with organic or inorganic acids, from compounds of formula (I) with a basic nitrogen atom, especially the pharmaceutically acceptable salts. Suitable inorganic acids are, for example, halogen acids, such as hydrochloric acid, sulfuric acid, or phosphoric acid.


5 Suitable organic acids are, for example, carboxylic, phosphonic, sulfonic or sulfamic acids, for example acetic acid, propionic acid, octanoic acid, decanoic acid, dodecanoic acid, glycolic acid, lactic acid, fumaric acid, succinic acid, adipic acid, pimelic acid, suberic acid, azelaic acid, malic acid, tartaric acid, citric acid, amino acids, such as glutamic acid or aspartic acid, maleic acid, hydroxymaleic acid, 10 methylmaleic acid, cyclohexanecarboxylic acid, adamantanecarboxylic acid, benzoic acid, salicylic acid, 4-aminosalicylic acid, phthalic acid, phenylacetic acid, mandelic acid, cinnamic acid, methane- or ethane-sulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 2-naphthalenesulfonic acid, 1,5-naphthalene-disulfonic acid, 2-, 3- or 4-methylbenzenesulfonic acid, methylsulfuric 15 acid, ethylsulfuric acid, dodecylsulfuric acid, N-cyclohexylsulfamic acid, N-methyl-, N-ethyl- or N-propyl-sulfamic acid, or other organic protonic acids, such as ascorbic acid.

The compound according to the invention may be administered in the form of a pro-drug which is broken down in the human or animal body to give a compound 20 of the formula I. Examples of pro-drugs include in vivo hydrolysable esters and amides of a compound of the formula I. Particular pro-drugs considered are ester and amides of naturally occurring amino acids and ester or amides of small peptides, in particular small peptides consisting of up to five, preferably two or three amino acids as well as esters and amides of pegylated hydroxy acids, preferably hydroxy acetic 25 acid and lactic acid. Pro-drug esters are formed from the acid function of the amino acid or the C terminal of the peptide and suitable hydroxy group(s) in the compound of formula I. Pro-drug amides are formed from the amino function of the amino acid or the N terminal of the peptide and suitable carboxy group(s) in the compound of formula I, or from the acid function of the amino acid or the C terminal of the peptide 30 and suitable amino group(s) in the compound of formula I. Particularly preferably the pro-drug amides are formed from the amino group(s) present within the R group of formula I.


More preferably, the pro-drug is formed by the addition of glycine, alanine or lysine to the compound of formula I.

Even more preferably the compound of general formula I is in the form of a
5 pro-drug selected from the compounds of formulae:






In an especially preferred embodiment the compound of general formula I is
 5 in the form of a pro-drug which has the following formula




In a most especially preferred embodiment the compound according to the
 invention is a pharmaceutically acceptable salt, preferably a hydrochloride salt, most
 preferably a dihydrochloride salt, of a compound of the following formula



The pharmaceutically active metabolite in vivo in this case is BAL27862.


5 These pro-drugs may be prepared by processes that are known per se, in particular, a process, wherein a compound of formula (II)



wherein R^1 is defined as for formula (I) and Z is CH or N, or a derivative of such a compound comprising functional groups in protected form,

10 or a salt thereof is

(1) acylated with an amino acid of formula (III)



15 wherein

R¹⁰ is selected from hydrogen (Gly); methyl (Ala) and protected aminobutyl (Lys) and R¹¹ is a suitable amino protecting group, and

(2) any protecting groups in a protected derivative of the resulting compound are removed to yield a pro-drug as shown above, and, if so desired,

- 5 (3) said pro-drug is converted into a salt by treatment with an acid, or a salt of a compound of formula (II) is converted into the corresponding free compound of formula (II) or into another salt, and/or a mixture of isomeric product compounds is separated into the individual isomers.

10 Acylation of a compound of formula (II) with an amino acid of formula (III) is performed in a manner known per se, usually in the presence of a suitable polar or dipolar aprotic solvent, with cooling or heating as required, for example in a temperature range from approximately minus 80°C to approximately plus 150°C, more preferably from minus 30°C to plus 120°C, especially in a range from
15 approximately around 0°C to the reflux temperature of the used solvent. Optionally a suitable base is added, in particular an aromatic base like pyridine or collidine or a tertiary amine base such as triethylamine or diisopropylethylamine, or an inorganic basic salt, e.g. potassium or sodium carbonate.

20 Acylation may be accomplished under conditions used for amide formation known per se in peptide chemistry, e.g. with activating agents for the carboxy group, such as carbodiimides like N,N'-diethyl-, N,N'-dipropyl-, N,N'-diisopropyl-, N,N'-dicyclohexylcarbodiimide and N-(3-dimethylaminoisopropyl)-N'-ethylcarbodiimide-hydrochloride (EDC), or with agents such as 1-hydroxybenzotriazole (HOBt),
25 benzotriazol-1-yloxytris(dimethylamino)-phosphonium hexafluorophosphate (BOP), O-(7-aza-benzotriazol-1-yl)-N,N,N',N'-tetramethyl-uronium hexafluorophosphate (HATU), 2-(2-oxo-1-(2H)-pyridyl)-1,1,3,3-tetramethyluronium tetrafluoroborate (TPTU), optionally in the presence of suitable bases, catalysts or co-reagents. The carboxy group may also be activated as acyl halogenide, preferably as acyl chloride,
30 e.g. by reaction with thionylchloride or oxalylchloride, or as symmetrical or unsymmetrical anhydride, e.g. by reaction with halogeno formates like ethyl chloroformate, optionally in the presence of suitable bases, catalysts or co-reagents.

If one or more other functional groups, for example carboxy, hydroxy or amino, are or need to be protected in a compound of formula (II) or (III), because they should not take part in the reaction, these are such protecting groups as are usually applied in the synthesis of amides like, in particular peptide compounds, cephalosporins, penicillins, nucleic acid derivatives and sugars, which are known to the skilled persons. Suitable protecting groups for amino groups are for example t-butyl carbamate, benzyl carbamate or 9-fluorenylmethyl carbamate.

The protecting groups may already be present in precursors and should protect the functional groups concerned against unwanted secondary reactions, such as alkylations, acylations, etherifications, esterifications, oxidations, solvolysis, and similar reactions. It is a characteristic of protecting groups that they lend themselves readily, i.e. without undesired secondary reactions, to removal, typically by solvolysis, reduction, photolysis or also by enzyme activity, for example under conditions analogous to physiological conditions, and that they are not present in the end products. The specialist knows, or can easily establish, which protecting groups are suitable with the reactions mentioned hereinabove and hereinafter.

The protection of such functional groups by such protecting groups, the protecting groups themselves, and their removal reactions are described for example in standard reference books for peptide synthesis and in special books on protective groups such as

J. F. W. McOmie, "Protective Groups in Organic Chemistry", Plenum Press, London and New York 1973, in "Methoden der organischen Chemie" (Methods of organic chemistry), Houben-Weyl, 4th edition, Volume 15/I, Georg Thieme Verlag, Stuttgart 1974, and in T. W. Greene, G. M. Wuts "Protective Groups in Organic Synthesis", Wiley, New York, 2006.

Disease

The compounds of general formula I according to the invention have been shown to arrest cell proliferation and induce cell death, for example by apoptosis.

Dysregulation of cell proliferation, or lack of appropriate cell death, has wide ranging clinical implications. A number of diseases associated with such dysregulation involve hyperproliferation, inflammation, tissue remodeling and repair. Familiar indications in this category include cancers, restenosis, neointimal
5 hyperplasia, angiogenesis, endometriosis, lymphoproliferative disorders, transplantation related pathologies (graft rejection), polyposis, loss of neural function in the case of tissue remodeling and the like.

Cancer is associated with abnormal cell proliferation and cell death rates. As apoptosis is inhibited or delayed in most types of proliferative, neoplastic diseases,
10 induction of apoptosis is an option for treatment of cancer, especially in cancer types which show resistance to classic chemotherapy, radiation and immunotherapy (Apoptosis and Cancer Chemotherapy, Hickman and Dive, eds., Blackwell Publishing, 1999). Also in autoimmune and transplantation related diseases and pathologies compounds inducing apoptosis may be used to restore normal cell death
15 processes and therefore can eradicate the symptoms and might cure the diseases. Further applications of compounds inducing apoptosis may be in restenosis, i.e. accumulation of vascular smooth muscle cells in the walls of arteries, and in persistent infections caused by a failure to eradicate bacteria- and virus-infected cells. Furthermore, apoptosis can be induced or reestablished in epithelial cells, in
20 endothelial cells, in muscle cells, and in others which have lost contact with extracellular matrix.

A compound according to general formula I or pharmaceutically acceptable derivatives thereof may be used for the prophylactic or especially therapeutic treatment of the human or animal body, in particular for treating a neoplastic disease,
25 autoimmune disease, transplantation related pathology and/or degenerative disease. Examples of such neoplastic diseases include, but are not limited to, epithelial neoplasms, squamous cell neoplasms, basal cell neoplasms, transitional cell papillomas and carcinomas, adenomas und adenocarcinomas, adnexal and skin appendage neoplasms, mucoepidermoid neoplasms, cystic neoplasms, mucinous
30 and serous neoplasms, ductal-, lobular and medullary neoplasms, acinar cell neoplasms, complex epithelial neoplasms, specialized gonadal neoplasms, paragangliomas and glomus tumours, naevi and melanomas, soft tissue tumours and

sarcomas, fibromatous neoplasms, myxomatous neoplasms, lipomatous neoplasms, myomatous neoplasms, complex mixed and stromal neoplasms, fibroepithelial neoplasms, synovial like neoplasms, mesothelial neoplasms, germ cell neoplasms, trophoblastic neoplasms, mesonephromas, blood vessel tumours, lymphatic vessel
5 tumours, osseous and chondromatous neoplasms, giant cell tumours, miscellaneous bone tumours, odontogenic tumours, gliomas, neuroepitheliomatous neoplasms, meningiomas, nerve sheath tumours, granular cell tumours and alveolar soft part sarcomas, Hodgkin's and non-Hodgkin's lymphomas, other lymphoreticular neoplasms, plasma cell tumours, mast cell tumours, immunoproliferative diseases,
10 leukemias, miscellaneous myeloproliferative disorders, lymphoproliferative disorders and myelodysplastic syndromes.

The compounds of general formula I or pharmaceutically acceptable derivatives thereof may be used to treat autoimmune diseases. Examples of such autoimmune diseases include, but are not limited to, systemic, discoid or subacute
15 cutaneous lupus erythematosus, rheumatoid arthritis, antiphospholipid syndrome, CREST, progressive systemic sclerosis, mixed connective tissue disease (Sharp syndrome), Reiter's syndrome, juvenile arthritis, cold agglutinin disease, essential mixed cryoglobulinemia, rheumatic fever, ankylosing spondylitis, chronic polyarthritis, myasthenia gravis, multiple sclerosis, chronic inflammatory demyelinating
20 polyneuropathy, Guillan-Barre syndrome, dermatomyositis/ polymyositis, autoimmune hemolytic anemia, thrombocytopenic purpura, neutropenia, type I diabetes mellitus, thyroiditis (including Hashimoto's and Grave's disease), Addison's disease, polyglandular syndrome, pemphigus (vulgaris, foliaceus, sebaceous and vegetans), bullous and cicatricial pemphigoid, pemphigoid gestationis, epidermolysis
25 bullosa acquisita, linear IgA disease, lichen sclerosus et atrophicus, morbus Duhring, psoriasis vulgaris, guttate, generalized pustular and localized pustular psoriasis, vitiligo, alopecia areata, primary biliary cirrhosis, autoimmune hepatitis, all forms of glomerulonephritis, pulmonal hemorrhage (goodpasture syndrome), IgA nephropathy, pernicious anemia and autoimmune gastritis, inflammatory bowel
30 diseases (including colitis ulcerosa and morbus Crohn), Behcet's disease, Celiac-Sprue disease, autoimmune uveitis, autoimmune myocarditis, granulomatous orchitis, aspermatogenesis without orchitis, idiopathic and secondary pulmonary fibrosis, inflammatory diseases with a possibility of autoimmune pathogenesis, such

as pyoderma gangrenosum, lichen ruber, sarcoidosis (including Lofgren and cutaneous/subcutaneous type), granuloma anulare, allergic type I and type IV immunological reaction, asthma bronchiale, pollinosis, atopic, contact and airborne dermatitis, large vessel vasculitis (giant cell and Takayasu's arteritis), medium sized vessel vasculitis (polyarteritis nodosa, Kawasaki disease), small vessel vasculitis (Wegener's granulomatosis, Churg Strauss syndrome, microscopic polyangiitis, Henoch-Schoenlein purpura, essential cryoglobulinemic vasculitis, cutaneous leukocytoclastic angiitis), hypersensitivity syndromes, toxic epidermal necrolysis (Stevens-Johnson syndrome, erythema multiforme), diseases due to drug side effects, all forms of cutaneous, organ-specific and systemic effects due to type I-IV (Coombs classification) immunologic forms of reaction, transplantation related pathologies, such as acute and chronic graft versus host and host versus graft disease, involving all organs (skin, heart, kidney, bone marrow, eye, liver, spleen, lung, muscle, central and peripheral nerve system, connective tissue, bone, blood and lymphatic vessel, genito-urinary system, ear, cartilage, primary and secondary lymphatic system including bone marrow, lymph node, thymus, gastrointestinal tract, including oro-pharynx, esophageus, stomach, small intestine, colon, and rectum, including parts of above mentioned organs down to single cell level and substructures, e. g. stem cells).

Particularly preferably, the disease according to the invention is a neoplastic or autoimmune disease. In an especially preferred embodiment the disease is cancer.

Examples of cancers in terms of the organs and parts of the body affected include, but are not limited to, the breast, cervix, ovaries, colon, rectum, (including colon and rectum i.e. colorectal cancer), lung, (including small cell lung cancer, non-small cell lung cancer, large cell lung cancer and mesothelioma), bone, endocrine system, adrenal gland, thymus, liver, stomach, intestine, (including gastric cancer), pancreas, bone marrow, hematological malignancies, (such as lymphoma, leukemia, myeloma or lymphoid malignancies), bladder, urinary tract, kidneys, skin, thyroid, brain, head, neck, prostate and testis. Preferably the cancer is selected from the group consisting of breast cancer, prostate cancer, cervical cancer, ovarian cancer, gastric cancer, colorectal cancer, pancreatic cancer, liver cancer, brain cancer,

neuroendocrine cancer, lung cancer, kidney cancer, hematological malignancies, melanoma and sarcomas. Especially preferably the cancer is selected from the group consisting of breast cancer, cervical cancer, ovarian cancer, colorectal cancer, melanoma and lung cancer. More especially preferably the cancer is selected from the group consisting of lung cancer, melanoma, ovarian cancer and colorectal cancer. In another preferred embodiment, for the case when the resistance predicted is acquired resistance, the cancer is lung cancer or ovarian cancer. In yet another preferred embodiment, for the case where the resistance predicted is inherent resistance, the cancer is selected from the group consisting of colorectal cancer, lung cancer or melanoma.

Samples

The measurement of the level of glu-tubulin may be performed *in vitro*, on a sample of biological tissue derived from the subject. The sample may be any biological material separated from the body such as, for example, normal tissue, tumour tissue, cell lines, whole blood, serum, plasma, cerebrospinal fluid, lymph fluid, circulating tumour cells, cell lysate, tissue lysate, urine and aspirates. Preferably the sample is derived from normal tissue, tumour tissue, or circulating tumour cells. More preferably the sample is derived from tumour tissue or circulating tumour cells. In one particularly preferred embodiment the sample is derived from tumour tissue. For example, the level of glu-tubulin may be measured in a fresh, frozen or formalin fixed/paraffin embedded tumour tissue sample.

The sample is pre-obtained from the subject before the sample is subjected to the method steps involving measuring the level of the biomarker. The methods for removal of the sample are well known in the art, and it may for example be removed from the subject by biopsy, for example by punch biopsy, core biopsy or aspiration fine needle biopsy, endoscopic biopsy, or surface biopsy. Blood may be collected by venipuncture and further processed according to standard techniques. Circulating tumour cells may also be obtained from blood based on, for example, size (e.g. ISET - Isolation by Size of Epithelial Tumour cells) or immunomagnetic cell enrichment (e.g. Cellsearch®, Veridex, Raritan, NJ)

Sample comparison

The subject according to the invention may be human or animal. Preferably the subject is human.

The biomarker glu-tubulin is measured *ex vivo* in a sample or samples taken from the human or animal body, preferably taken from the human body. The sample
5 or samples are pre-obtained from the human or animal body, preferably pre-obtained from the human body before the sample is subjected to the method steps involving measuring the level of the biomarker.

A biomarker is in general a substance that is used as an indicator of a biological response, preferably as an indicator of the susceptibility to a given
10 treatment, which in the present application is treatment with a compound of general formula I or a pharmaceutically acceptable derivative thereof.

In a particularly preferred embodiment, higher glu-tubulin levels in the sample relative to a standard value or set of standard values predicts resistance.

As used herein, an increase or relatively high or high or higher levels relative
15 to a standard level or set of standard levels means the amount or concentration of the biomarker in a sample is detectably greater in the sample relative to the standard level or set of standard levels. This encompasses at least an increase of, or higher level of, about 1% relative to the standard, preferably at least an increase of about 5% relative to the standard. More preferably it is an increase of, or higher level of, at
20 least about 10% relative to the standard. More particularly preferably it is an increase of, or higher level of, at least about 20% relative to the standard. For example, such an increase of, or higher level of, may include, but is not limited to, at least about 1%, about 10%, about 20%, about 30%, about 50%, about 70%, about 80%, about 100%, about 150% or about 200% or more relative to the standard.

25 Preferably, higher glu-tubulin levels in a sample or samples

i) relative to a standard value or set of standard values from subjects with the same tumour histotype; or

ii) taken after treatment initiation and compared to a sample or samples taken from the same subject before treatment initiation; or

iii) relative to a standard value or set of standard values from normal cells or tissue;

5 are predictive of resistance.

The measuring of a level of glu-tubulin is performed ex-vivo in a sample pre-obtained from the subject. Further preferably the response which is to be predicted is resistance.

More preferably, higher glu-tubulin levels in a sample or samples

10 i) relative to a standard value or set of standard values from subjects with the same tumour histotype; or

ii) taken after treatment initiation and compared to a sample or samples taken from the same subject before treatment initiation;

are predictive of resistance.

15 Especially preferably, higher glu-tubulin levels in a sample or samples relative to a standard value or set of standard values from subjects with the same tumour histotype are predictive of resistance.

In one preferred embodiment, for the case i) where the measurement is compared in a sample or samples relative to a standard value or set of standard values from samples from subjects with the same tumour histotype as the sample to which it is to be compared, the standard value or set of standard values are established from samples from a population of subjects with that cancer type. The samples from these standard subjects may for example be derived from the tumour tissue, circulating tumour cells or blood, as long as the origin of the sample is consistent between the standard and the sample to be compared.

20

25

In another preferred embodiment, for the case ii) where the measurement is compared in a sample or samples taken after treatment initiation and compared to a sample or samples taken from the same subject before treatment initiation, it is measured preferably to predict acquired resistance. The samples are compared to
5 cells or tissue from the same biological origin. The prediction of acquired resistance would then indicate that the treatment with the compound should be discontinued. The biomarker is thus used to monitor whether further treatment with the compound is likely to give the required response (e.g. reduction of abnormal cells), or whether the cells have become non-responsive or resistant to such treatment.

10 In yet another preferred embodiment, for the case iii) where the measurement is compared in a sample or samples relative to a standard value or set of standard values from normal cells or tissue, the standard value or set of standard values may be established from a sample of normal (e.g. non-tumourous) cells or tissue or body fluid. Such data may be gathered from a population of subjects in
15 order to develop the standard value or set of standard values.

The standard value or set of standard values are established ex-vivo from pre-obtained samples which may be from cell lines, or preferably biological material from at least one subject and more preferably from an average of subjects (e.g., n=2 to 1000 or more). The standard value or set of standard values may then be
20 correlated with the response data of the same cell lines, or same subjects, to treatment with a compound of general formula I or a pharmaceutically acceptable derivative thereof. From this correlation a comparator module, for example in the form of a relative scale or scoring system, optionally including cut-off or threshold values, can be established which indicates the levels of biomarker associated with a
25 spectrum of response levels to the compound of formula I or a pharmaceutically acceptable derivative thereof. The spectrum of response levels may comprise relative sensitivity to the therapeutic activity of the compound, (e.g. high sensitivity to low sensitivity), as well as resistance to the therapeutic activity. In a preferred embodiment this comparator module comprises a cut-off value or set of values which
30 predicts resistance to treatment.

For example, if an immunohistochemical method is used to measure the

level of glu-tubulin in a sample, standard values may be in the form of a scoring system. Such a system might take into account the percentage of cells in which staining for glu-tubulin is present. The system may also take into account the relative intensity of staining in the individual cells. The standard values or set of standard values of the level of glu-tubulin may then be correlated with data indicating the response, especially resistance, of the subject or tissue or cell line to the therapeutic activity of a compound of formula I or a pharmaceutically acceptable derivative thereof. Such data may then form part of a comparator module.

Response is the reaction of the cell lines, or preferably of the subject, or more preferably of the disease in a subject, to the therapeutic activity of a compound of general formula I or a pharmaceutically acceptable derivative thereof. The spectrum of response levels may comprise relative sensitivity to the therapeutic activity of the compound, (e.g. high sensitivity to low sensitivity), as well as resistance to the therapeutic activity. The response data may for example be monitored in terms of: objective response rates, time to disease progression, progression free survival, and overall survival.

The response of a cancerous disease may be evaluated by using criteria well known to a person in the field of cancer treatment, for example but not restricted to,

Response Evaluation Criteria in Solid Tumors (RECIST) Guidelines, Source: Eisenhauer EA, Therasse P, Bogaerts J, Schwartz LH, Sargent D, Ford R, Dancey J, Arbuck S, Gwyther S, Mooney M, Rubinstein L, Shankar L, Dodd L, Kaplan R, Lacombe D, Verweij J. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). Eur J Cancer.2009; 45:228-47;

RANO Criteria for High-Grade Gliomas, Source: Wen PY, Macdonald DR, Reardon DA, Cloughesy TF, Sorensen AG, Galanis E, Degroot J, Wick W, Gilbert MR, Lassman AB, Tsien C, Mikkelsen T, Wong ET, Chamberlain MC, Stupp R, Lamborn

KR, Vogelbaum MA, van den Bent MJ, Chang SM. Updated response assessment criteria for high-grade gliomas: response assessment in neuro-oncology working group. J Clin Oncol. 2010;28(11):1963-72;

5

CA-125 Rustin Criteria for Ovarian Cancer Response,
Source: Rustin GJ, Quinn M, Thigpen T, du Bois A, Pujade-Lauraine E, Jakobsen A, Eisenhauer E, Sagae S, Greven K, Vergote I, Cervantes A, Vermorken J. Re: New guidelines to evaluate the response to treatment in solid tumors (ovarian cancer). J Natl Cancer Inst. 2004; 96(6):487-8;

10

and

15

PSA Working Group 2 Criteria for Prostate Cancer Response,
Source: Scher HI, Halabi S, Tannock I, Morris M, Sternberg CN, Carducci MA, Eisenberger MA, Higano C, Bubley GJ, Dreicer R, Petrylak D, Kantoff P, Basch E, Kelly WK, Figg WD, Small EJ, Beer TM, Wilding

20

G, Martin A, Hussain M; Prostate Cancer Clinical Trials Working Group. Design and end points of clinical trials for patients with progressive prostate cancer and castrate levels of testosterone: recommendations of the Prostate Cancer Clinical Trials Working Group. J Clin Oncol. 2008;26(7):1148-59.

25

Resistance is associated with there not being an observable and/or measurable reduction in, or absence of, one or more of the following: reduction in the number of abnormal cells, preferably cancerous cells; or absence of the abnormal cells, preferably cancerous cells; for cancerous diseases: reduction in tumour size; inhibition (i.e., slowed to some extent and preferably stopped) of further tumour growth; reduction in the levels of tumour markers such as PSA and CA-125, inhibition (i.e., slowed to some extent and preferably stopped) of cancer cell infiltration into other organs (including the spread of cancer into soft tissue and bone); inhibition (i.e., slowed to some extent and preferably stopped) of tumour metastasis; alleviation of

30

one or more of the symptoms associated with the specific cancer; and reduced morbidity and mortality.

In a preferred embodiment resistance means there is no observable and/or measurable reduction in, or absence of, one or more of the following criteria:

- 5 reduction in tumour size; inhibition of further tumour growth, inhibition of cancer cell infiltration into other organs; and inhibition of tumour metastasis.

In a more preferred embodiment resistance refers to one or more of the following criteria: no reduction in tumour size; no inhibition of further tumour growth, no inhibition of cancer cell infiltration into other organs; and no inhibition of tumour
10 metastasis.

Measurement of the aforementioned resistance criteria is according to clinical guidelines well known to a person in the field of cancer treatment, such as those listed above for measuring the response of a cancerous disease.

- Response may also be established in vitro by assessing cell proliferation
15 and/or cell death. For example, effects on cell death or proliferation may be assessed in vitro by one or more of the following well established assays: A) Nuclear staining with Hoechst 33342 dye providing information about nuclear morphology and DNA fragmentation which are hallmarks of apoptosis. B) AnnexinV binding assay which reflects the phosphatidylserine content of the outer lipid bilayer of the plasma
20 membrane. This event is considered an early hallmark of apoptosis. C) TUNEL assay (Terminal deoxynucleotidyl transferase mediated dUTP Nick End Labeling assay), a fluorescence method for evaluating cells undergoing apoptosis or necrosis by measuring DNA fragmentation by labeling the terminal end of nucleic acids. D) MTS proliferation assay measuring the metabolic activity of cells. Viable cells are
25 metabolically active whereas cells with a compromised respiratory chain show a reduced activity in this test. E) Crystal violet staining assay, where effects on cell number are monitored through direct staining of cellular components. F) Proliferation assay monitoring DNA synthesis through incorporation of bromodeoxyuridine (BrdU). Inhibitory effects on growth/proliferation can be directly determined. G) YO-PRO
30 assay which involves a membrane impermeable, fluorescent, monomeric cyanine,

nucleic acid stain, which permits analysis of dying (e.g. apoptotic) cells without interfering with cell viability. Overall effects on cell number can also be analysed after cell permeabilisation. H) Propidium iodide staining for cell cycle distribution which shows alterations in distribution among the different phases of the cell cycle. Cell cycle arresting points can be determined. I) Anchorage-independent growth assays, such as colony outgrowth assays which assess the ability of single cell suspensions to grow into colonies in soft agar.

In a preferred embodiment relating to determination of resistance in vitro, resistance means there is no decrease in the proliferation rate of abnormal cells and/or reduction in the number of abnormal cells. More preferably resistance means there is no decrease in the proliferation rate of cancerous cells and/or no reduction in the number of cancerous cells. The reduction in the number of abnormal, preferably cancerous, cells may occur through a variety of programmed and non-programmed cell death mechanisms. Apoptosis, caspase-independent programmed cell death and autophagic cell death are examples of programmed cell death. However the cell death criteria involved in embodiments of the invention are not to be taken as limited to any one cell death mechanism.

Glu-tubulin

Preferred examples of the protein sequence of alpha tubulin (human alpha tubulin) are listed in SEQ. ID No. 1-4, Figures 11-14. Alpha tubulin is a precursor of glu-tubulin. As described previously, the glu-tubulin itself has the C-terminal tyrosine removed. The term glu-tubulin also encompasses homologues, mutant forms, allelic variants, isoforms, splice variants and equivalents of the sequences represented by SEQ ID NO 1-4, with the proviso that a glutamate is the final amino acid at the C-terminal. More preferably it encompasses sequences having at least about 75% identity, especially preferably at least about 85% identity, particularly preferably at least about 95% identity, and more particularly preferably about 99% identity to said sequences, with in each case the proviso that a glutamate is the final amino acid at the C-terminal.

Level of glu-tubulin

The level of glu-tubulin may be assayed in the sample by protein analysis techniques well known to a skilled person. Examples of methods known in the art which are suitable to measure the level of glu-tubulin at the protein level include, but are not limited to, i) immunohistochemistry (IHC) analysis, ii) western blotting iii) immunoprecipitation iv) enzyme linked immunosorbant assay (ELISA) v) radioimmunoassay vi) Fluorescence activated cell sorting (FACS) vii) mass spectrometry, including matrix assisted laser desorption/ionisation (MALDI, e.g. MALDI-TOF) and electrospray ionisation mass-spectrometry (ESI—MS).

The antibodies involved in some of the above methods may be monoclonal or polyclonal antibodies, antibody fragments, and/or various types of synthetic antibodies, including chimeric antibodies. The antibody may be labelled to enable it to be detected or capable of detection following reaction with one or more further species, for example using a secondary antibody that is labelled or capable of producing a detectable result. Antibodies specific to the glu-tubulin form of alpha tubulin are available commercially from Milipore or can be prepared via conventional antibody generation methods well known to a skilled person.

Preferred methods of protein analysis are ELISA, mass spectrometry techniques, immunohistochemistry and western blotting, more preferably western blotting and immunohistochemistry. In western blotting, also known as immunoblotting, labelled antibodies may be used to assess levels of protein, where the intensity of the signal from the detectable label corresponds to the amount of protein, and can be quantified for example by densitometry.

Immunohistochemistry again uses labelled antibodies to detect the presence and relative amount of the biomarker. It can be used to assess the percentage of cells for which the biomarker is present. It can also be used to assess the localisation or relative amount of the biomarker in individual cells, the latter is seen as a function of the intensity of staining.

ELISA stands for enzyme linked immunosorbant assay, since it uses an enzyme linked to an antibody or antigen for the detection of a specific protein. ELISA

is typically performed as follows (although other variations in methodology exist): a solid substrate such as a 96 well plate is coated with a primary antibody, which recognises the biomarker. The bound biomarker is then recognised by a secondary antibody specific for the biomarker. This may be directly joined to an enzyme or a third anti-immunoglobulin antibody may be used which is joined to an enzyme. A substrate is added and the enzyme catalyses a reaction, yielding a specific colour. By measuring the optical density of this colour, the presence and amount of the biomarker can be determined.

10 Uses of biomarker

The biomarker may be used to predict inherent resistance of the disease in a subject to the compound of general formula I or a pharmaceutically acceptable derivative thereof as defined above.

15 The biomarker may be used to predict acquired resistance of the disease in a subject to the compound of general formula I or a pharmaceutically acceptable derivative thereof as defined above.

20 The biomarker may be used to select subjects suffering or predisposed to suffering from a disease, preferably cancer, for treatment with a compound of general formula I or a pharmaceutically acceptable derivative thereof as defined above. The levels of such a biomarker may be used to identify subjects likely to respond or to not respond or to continue to respond or to not continue to respond to treatment with such agents. Stratification of subjects may be made in order to avoid unnecessary treatment regimes. In particular the biomarker may be used to identify subjects from whom a sample or samples do not display a higher level of glu-tubulin, relative to a standard level or set of standard levels, whereupon such subjects may then be selected for treatment with the compound of formula I or a pharmaceutically acceptable derivative thereof as defined above.

25 The biomarker may also be used to assist in the determination of treatment regimes, regarding amounts and schedules of dosing. Additionally, the biomarker

may be used to assist in the selection of a combination of drugs to be given to a subject, including a compound or compounds of general formula I or a pharmaceutically acceptable derivative thereof, and another chemotherapeutic (cytotoxic) agent or agents. Furthermore, the biomarker may be used to assist in the determination of therapy strategies in a subject including whether a compound of general formula I or a pharmaceutically acceptable derivative thereof is to be administered in combination with targeted therapy, endocrine therapy, radiotherapy, immunotherapy or surgical intervention, or a combination of these.

Glu-tubulin may also be used in combination with other biomarkers to predict the response to a compound of general formula I or a pharmaceutically acceptable derivative thereof and to determine treatment regimes. It may furthermore be used in combination with chemo-sensitivity testing to predict resistance and to determine treatment regimes. Chemo-sensitivity testing involves directly applying a compound of general formula I to cells taken from the subject, for example from a subject with haematological malignancies or accessible solid tumours, for example breast, head and neck cancers or melanomas, to determine the response of the cells to the compound.

Method of treatment

The invention also involves in some aspects a method of treatment and glu-tubulin for use in a method of treatment, wherein the level of glu-tubulin is first established relative to a standard level or set of standard levels or pre-treatment initiation levels and then a compound of general formula I or a pharmaceutically acceptable derivative thereof as defined above, is administered if the level of glu-tubulin in said sample is not higher than a standard value or set of standard values or has not increased relative to pre-treatment initiation levels respectively. The compound of formula I or a pharmaceutically acceptable derivative thereof may be administered in a pharmaceutical composition, as is well known to a person skilled in the art. Suitable compositions and dosages are for example disclosed in WO 2004/103994 A1 pages 35-39, which are specifically incorporated by reference herein. Compositions for enteral administration, such as nasal, buccal, rectal or, especially, oral administration, and for parenteral administration, such as

intravenous, intramuscular or subcutaneous administration, to warm-blooded animals, especially humans, are especially preferred. More particularly, compositions for intravenous administration are preferred.

The compositions comprise the active ingredient and a pharmaceutically acceptable carrier. An example of a composition includes, but is not limited to, the following: 5000 soft gelatin capsules, each comprising as active ingredient 0.05 g of one of the compounds of general formula (I), are prepared as follows: 250 g pulverized active ingredient is suspended in 2 liter Lauroglykol® (propylene glycol laurate, Gattefossé S.A., Saint Priest, France) and ground in a wet pulverizer to produce a particle size of about 1 to 3 μm . 0.419 g portions of the mixture are then introduced into soft gelatin capsules using a capsule-filling machine.

The invention also relates in one aspect to a method of treating a neoplastic or autoimmune disease, preferably cancer, by first decreasing the level of glu-tubulin in a subject that has a sample with a higher level of glu-tubulin compared to a standard level or set of standard levels or pre-treatment initiation levels, then treating the subject with a compound of general formula I or a pharmaceutically acceptable derivative as defined above. The level of glu-tubulin may be decreased by direct or indirect chemical or genetic means. Examples of such methods are treatment with a drug that results in reduced glu-tubulin expression, targeted delivery of viral, plasmid or peptide constructs, or antibody or siRNA or antisense to downregulate the level of glu-tubulin. For example siRNA may be used to reduce the level of TTCP or delivery of a plasmid may be used to increase the expression of TTL, and thereby reduce the level of glu-tubulin in the cell. The subject may then be treated with a compound of general formula I or a pharmaceutically acceptable derivative thereof.

A compound of general formula I or a pharmaceutically acceptable derivative thereof can be administered alone or in combination with one or more other therapeutic agents. Possible combination therapy may take the form of fixed combinations, or the administration of a compound of the invention and one or more other therapeutic agents which are staggered or given independently of one another, or the combined administration of fixed combinations and one or more other therapeutic agents.

A compound of general formula I or a pharmaceutically acceptable derivative thereof can, besides or in addition, be administered especially for tumour therapy in combination with chemotherapy (cytotoxic therapy), targeted therapy, endocrine therapy, radiotherapy, immunotherapy, surgical intervention, or a combination of these. Long-term therapy is equally possible as is adjuvant therapy in the context of other treatment strategies, as described above. Other possible treatments are therapy to maintain the patient's status after tumour regression, or even chemo-preventive therapy, for example in patients at risk.

Kit and device


In one aspect the invention relates to a kit and in another aspect to a device for predicting the response, preferably of a disease in a subject, to a compound of general formula I or a pharmaceutically acceptable derivative thereof as defined above, comprising reagents necessary for measuring the level of glu-tubulin in a sample. Preferably, the reagents comprise a capture reagent comprising a detector for glu-tubulin and a detector reagent.

The kit and device may also preferably comprise a comparator module which comprises a standard value or set of standard values to which the level of glu-tubulin in the sample is compared. In a preferred embodiment, the comparator module is included in instructions for use of the kit. In another preferred embodiment the comparator module is in the form of a display device, for example a strip of colour or numerically coded material which is designed to be placed next to the readout of the sample measurement to indicate resistance levels. The standard value or set of standard values may be determined as described above.

The reagents are preferably antibodies or antibody fragments which selectively bind to glu-tubulin. These may for example be in the form of one specific primary antibody which binds to glu-tubulin and a secondary antibody which binds to the primary antibody, and which is itself labelled for detection. Alternatively, the primary antibody may also be labelled for direct detection. The kits or devices may optionally also contain a wash solution(s) that selectively allows retention of the

bound biomarker to the capture reagent as compared with other biomarkers after washing. Such kits can then be used in ELISA, western blotting, flow cytometry, immunohistochemistry or other immunochemical methods to detect the level of the biomarker.

5 More preferably the kit comprises a compound of general formula I, or a pharmaceutically acceptable derivative thereof as defined above. This compound may then be administered to the subject, in accordance with the level of the biomarker in the sample from the subject, as measured by the reagents comprised in the kit. Therefore the kit according to the invention may be used in the method of
10 treatment according to the invention, as defined above. In an especially preferred embodiment the kit comprises a compound of the following formula or a pharmaceutically acceptable salt thereof



15 In a particularly preferred embodiment of the kit the pharmaceutically acceptable salt is a dihydrochloride salt. In another aspect the invention relates to the use of such a kit as described above.

Furthermore the device may comprise imaging devices or measurement
20 devices (for example, but not restricted to, measurement of fluorescence) which

further process the measured signals and transfer them into a scale in a comparator module.

In the present specification the words “comprise” or “comprises” or “comprising” are to be understood as to imply the inclusion of a stated item or group of items, but not the exclusion of any other item or group of items.

Experimental methodology

Immunofluorescent staining of cultured cells

A549 human non-small cell lung cancer (NSCLC, ATCC reference number CCL-185) cells, HeLa cervical cancer cells (ATCC reference number CCL-2) and SKBR3 breast carcinoma cells (ATCC reference number HTB-30) were seeded at densities of 50% on round microscope coverslips and cultured for 24 hours in RPMI-1640 containing 10 % FCS (also referred to as FBS) at 37°C, 5% CO₂. Compounds to be tested were dissolved in DMSO. The cell culture medium was replaced with medium containing the diluted compound(s) (paclitaxel, vinblastine, colchicine and nocodazole were purchased from Sigma-Aldrich) or vehicle. After treatment for the times indicated in the Brief Description of the Figures, coverslips were washed and cells were fixed in methanol/acetone (1:1) for 5 minutes at room temperature and subsequently incubated in blocking buffer (0.5% BSA and 0.1% TX-100 in PBS) for 30 minutes at room temperature. Specimens were then incubated with anti-alpha-tubulin (Sigma, 1:2000) for 1 hour at room temperature in blocking buffer. After several washing steps cells were incubated with AlexaFluor-488 goat-anti-mouse IgG (Molecular Probes, 1:3000) for 1 hour at room temperature followed by several washing steps with blocking buffer. Specimens were then mounted with ProLong Gold antifade (Molecular Probes, sealed with nail polish and examined with a Leica immunofluorescence microscope. Images were captured with a cooled CCD-camera and processed by ImageJ software.

Colony Outgrowth Assay:

Single cell suspensions of patient-derived tumour xenografts (maintained in nude mice) were prepared. For colony outgrowth assays, cells were plated in soft agar in 24-well plates according to the assay introduced by Hamburger & Salmon (Primary bioassay of human tumour stem cells, *Science*, 1977, 197:461-463). 2×10^4 -
5 6×10^4 cells in 0.2 mL medium containing 0.4% agar were plated out on a bottom layer of 0.75% agar. Test compounds were applied in 0.2 mL culture medium. Every 24-well plate contained untreated controls and samples in triplicates. Cultures were incubated at 37°C and 7.5% CO₂ for 5 - 28 days. 24 hours prior to analysis, vital colonies were stained with a solution of metabolisable tetrazolium salt (Alley MC et
10 al, *Life Sci.* 1982, 31:3071-3078) and were counted with an automatic image analysis system (Omnicon 3600, Biosys GmbH).

Relative drug effects were expressed by the ratio of the mean number of colonies in the treated wells and the control wells. IC₇₀-values were determined by
15 plotting compound concentrations *versus* relative colony counts.

Generation and Crystal Violet Assay of BAL27862-Resistant Cell Lines

BAL27862-resistant sublines of human non-small cell lung cancer (H460 ATCC reference HTB-177; A549 ATCC reference CCL-185), ovarian cancer (SKOV3
20 ATCC reference HTB-77) lines were generated by long-term selection in complete cell culture medium (RPMI-1640 containing 10% FCS; Sigma-Aldrich) by stepwise increasing concentrations of BAL27862. Dependent on the cell line, the selection process was carried out for 8-12 months in order to achieve resistance factors (ratio of IC₅₀ of resistant cell line and appropriate wild-type cell line) between 3 and 11.6.
25 The resistant sublines were expanded at the highest tolerated BAL27862 concentration and subsequently frozen and stored in liquid nitrogen.

Cells were seeded in 96 well plates at the following densities: A549: 2000, H460: 1000, SKOV3: 2000 and, after 24 hours incubation, were incubated for 72 hours with DMSO, BAL27862, colchicine, nocodazole, paclitaxel or vinblastine
30 diluted in complete medium (final concentration DMSO max. 0.5 %). After medium

was removed, cells were fixed and stained by adding 50 µl Crystal Violet Staining (0.2 % Crystal Violet in 50 % Methanol) per well. Plates were incubated for 1 hour at room temperature. Subsequently the stain was decanted and plates were washed 4 times with double-distilled water. Plates were air-dried for several hours. Stain was dissolved by adding 100 µl buffer (0.1 M Tris pH 7.5, 0.2 % SDS, 20 % Ethanol) per well and shaking the plates. Absorbance at 590 nm was measured using a SpectraMax M2e plate reader (Molecular Devices). Anti-proliferative IC₅₀ values were calculated from concentration response curves using GraphPad Prism software. Resistance factors were calculated as a ratio of BAL27862 IC₅₀ in the resistant line variant *versus* the IC₅₀ in the parental line.

Protein Extraction

Tumour extraction: Tumours were extracted in ice-cold lysis buffer containing 50 mM HEPES (pH 7.5), 150 mM NaCl, 25 mM β-glycerophosphate, 25 mM NaF, 5 mM EGTA, 1 mM EDTA, 0.1% NP40, 15 mM pyrophosphate, 2 mM sodium orthovanadate, 10 mM sodium molybdate, leupeptin (10 µg/mL), aprotinin (10 µg/mL) and 1 mM phenylmethanesulphonyl fluoride (PMSF) (1 mL extraction volume per 45 mg tumour). After homogenisation by Polytron, lysates were adjusted to 1% NP40 and incubated on ice for 20 min. Lysates were clarified by centrifugation and frozen at -80°C.

Tumour cell line extraction: Cells were washed with ice-cold PBS containing 1 mM PMSF and with ice-cold lysis buffer (see above) without NP40. Cells were extracted in the same lysis buffer containing 1% NP40. After homogenisation, lysates were clarified by centrifugation and frozen at -80°C.

Immunoblotting/Western Blotting

Immunoblotting was performed using 20 µg of total protein per lane. Total protein concentration was determined with the BCA Protein Assay (Pierce). Protein

was separated on a 10% SDS-gel and transferred to a PVDF membrane using Semidry Blotting (90 min, 50 mA/gel). The primary antibodies used for immunoblotting were as follows:

5 Glu-tubulin antibody (available from Milipore, reference number AB3201), rabbit polyclonal, dilution 1:1000, buffer conditions: 3% BSA in PBS/0.1% Tween

Actin antibody (available from Chemicon, reference number MAB1501), mouse monoclonal, dilution 1:5000, buffer conditions: 3% BSA in PBS/0.1% Tween

10 The secondary antibodies used for immunoblotting were peroxidase-conjugated goat anti-rabbit or goat anti-mouse (available from Jackson ImmunoResearch Laboratories INC: reference number 111-035-144 JIR and 115-035-146 JIR), dilution 1:5000, buffer conditions: 0.5% milk in PBS/0.1% Tween. Labelled bands were revealed using a Raytest Stella 3200 High Performance Imaging System.

15 Immunohistochemistry

Fixation of patient-derived tumour xenografts (maintained in nude mice) was performed in 10 % neutral-buffered formalin containing 4 % formaldehyde for 20 – 28 hours at room temperature. Fixed specimens were kept in a solution of 70 % ethanol for a maximum of one week prior to dehydration and paraffin embedding according to
20 a standard procedure, using the conditions listed below:

Sequential Treatment	time (hours)
70% EtOH	1
80% EtOH	2
99% EtOH	1
100% Isopropanol	0.5
100% Isopropanol	1
Xylol	0.5
Xylol	1

Xylol	1
Paraffin	1
Paraffin	2
Paraffin	2

Paraffin sections of approximately 2 µm were cut and processed by using the automated immunostainer Benchmark XT® (Roche) running the standard processing steps. The visualisation of the specific antibody staining was done with DAB (3,3-diaminobenzidine) as chromogenic substrate at a concentration of 5 mg/ml.

- 5 The following primary antibody and processing conditions were used for staining:

Antibody Specification	Processing
Anti-Glu-Tubulin, Millipore, #AB3201, rabbit polyclonal	Cell conditioning 1 buffer from Roche for 90 minutes, antibody incubation at 37°C for 32 minutes at a dilution of 1:50

Detailed examples

- 10 Example 1: A Distinct Mitotic Phenotype Induced by compounds of general formula I

- 15 Treatment with compound A (BAL27862) or with compound B or compound C, induced a highly reproducible and distinct microtubule phenotype in all tumour cell lines tested (shown for compound A in A549, HeLa and SKBR3 cells in Figure 1, and for compound B and compound C in A549 cells in Figure 2). In dividing cells an apparent fragmentation of the mitotic spindle occurred, resulting in the formation of dot-like structures (Figure 1). This phenotype was shown to be distinct from that observed with conventional microtubule targeting agents, such as the microtubule stabiliser paclitaxel and the microtubule destabilisers vinblastine and colchicine (Figure 3) and nocodazole (Figure 4).

Example 2: BAL27862 Overcomes Microtubule Phenotype Induced by Conventional Microtubule-targeting Drugs in a Dominant Fashion

In order to show the uniqueness of its activity on microtubules, BAL27862 was tested in combination with vinblastine, colchicine and paclitaxel (Figure 5) and nocodazole (Figure 6) using A549 cells. Treatment with vinblastine, colchicine, paclitaxel or nocodazole alone induced the mitotic microtubule phenotypes characteristic of these agents. However, combination treatment with BAL27862 for the last 4 hours resulted in disruption of the microtubule structures; creating a phenotype consistent with treatment of BAL27862 alone, despite the continued presence of vinblastine, colchicine, paclitaxel or nocodazole. In contrast, treating first with BAL27862 and subsequently for 4 hours in combination with vinblastine, colchicine, paclitaxel or nocodazole had no impact on the observed microtubule phenotype that was consistent with treatment with BAL27862.

These data demonstrate that compounds of formula I affect microtubule biology consistently, but in a different manner than conventional microtubule targeting agents.

Detailed Examples according to the invention

Example 3: Association of high glu-tubulin expression levels with patient-derived tumour cells resistant to BAL27862 treatment.

Based on colony outgrowth assays, using tumour cells derived from 6 patient-derived tumours maintained as xenografts in mice, BAL27862-sensitive or relatively resistant tumour cells were identified from melanoma and colorectal and lung cancer (see Table 1). Concentrations at which 70% growth inhibition was observed *versus* controls (IC₇₀) are shown in Table 1. In this table, BAL27862-sensitive tumour cells have IC₇₀ values in the low nanomolar range, while BAL27862-resistant tumour cells are defined by IC₇₀ values >600 nanomolar. Paclitaxel and vinblastine data, using the same *ex vivo* assay, was available for 5 of the 6 tumour

models. Of these 5 models, all were resistant to treatment with paclitaxel, while 4 of 5 of these tumours were sensitive to treatment with vinblastine.

5 Table 1

Cancer type	name	Response to BAL27862	IC ₇₀ BAL27862 [microM]	Response to paclitaxel	Response to vinblastine
Colorectal cancer	CXF 1103	Sensitive	0.022	Resistant	Resistant
	CXF 243	Resistant	0.696	Resistant	Sensitive
Lung cancer	LXFE 211	Sensitive	0.021	Resistant	Sensitive
	LXFE 397	Resistant	> 3.5	Not known	Not known
Melanoma	MEXF 1341	Sensitive	0.025	Resistant	Sensitive
	MEXF 276	Resistant	> 3.5	Resistant	Sensitive

Immunoblotting analysis was performed in order to measure glu-tubulin levels in the same tumours maintained as xenografts, using the Milipore antibody (Figure 7). The actin levels were included on the immunoblot as a loading control.

Analysis of glu-tubulin levels indicated that glu-tubulin expression varied dramatically across all the tumours measured (Figure 7).

Based on the colony outgrowth assay and the same IC₇₀ criteria, there was no association between paclitaxel or vinblastine resistance and high glu-tubulin levels. This is evident since, for example, for the melanoma tumour type, both models were resistant to paclitaxel and yet for MEXF 1341 the glu-tubulin levels were clearly lower than in MEXF 276. The same lack of association was true for the vinca alkaloid, vinblastine in the melanoma model, since both these tumours were sensitive to vinblastine. Thus glu-tubulin levels were shown to be unsuitable as a reliable biomarker of resistance to the conventional microtubule agents paclitaxel and vinblastine in patient-derived tumour models.

Surprisingly, in contrast, when the BAL27862 resistance data is compared with the glu-tubulin level, glu-tubulin is shown to be higher only in the resistant tumours and not the sensitive tumours derived from the same tumour histotype. Increased levels were therefore consistently indicative of resistance to BAL27862.

- 5 Thus glu-tubulin levels were shown to be a biomarker of resistance for the compound according to the invention, BAL27862.

Example 4: Immunohistochemical analysis of colorectal tumour xenografts

Immunohistochemical analysis was performed on the colorectal tumour xenografts (Figure 8), revealing a high level of glu-tubulin in the tumour model CXF
10 243. Again a clear correlation was seen between high levels of glu-tubulin and resistance to BAL27862 (tumour model CXF 243 was BAL27862-resistant, while tumour model CXF 1103 was BAL27862-sensitive; as defined by the colony outgrowth assay). Thus glu-tubulin levels were again shown to be a biomarker of resistance for the compound according to the invention, BAL27862.

15

Example 5: Higher glu-tubulin expression is observed in tumour lines selected for resistance to a compound of general formula I

20

In vitro selection for resistance to BAL27862 resulted in the generation of three relatively resistant tumour cell lines, with the following resistance factors *versus* parental lines (based on IC₅₀ determinations using the Crystal Violet assay): A549 (3.0 fold); SKOV3 resistant 1 (7.6 fold); SKOV3 resistant 2 (11.6 fold); H460 (5.3
25 fold)(Table 2).

Table 2:

Treatment compound	Resistance factors (ratio of IC ₅₀ BAL27862-resistant cell line variant and IC ₅₀ parental cell line)			
	A549	H460	SKOV3 resistant 1	SKOV3 resistant 2

BAL27862	3.0	5.3	7.6	11.6
Colchicine	0.9	1.6	2.0	2.8
Nocodazole	1.6	1.3	3.6	3.9
Vinblastine	2.3	4.6	15.7	17.8
Paclitaxel	0.06	0.3	0.4	0.5

In general these BAL27862-resistant cells exhibited a different level of response to other microtubule destabilising agents, such as colchicine, nocodazole and vinblastine, as compared to BAL27862; and indeed increased sensitivity to the microtubule stabiliser paclitaxel was observed in all lines (Table 2).

- 5 Extraction and immunoblot analysis of these lines to measure the glu-tubulin levels, followed by comparison to BAL27862 resistance data, again shows that glu-tubulin is higher in the resistant lines, as compared to the parental lines (Figure 9). This was maintained throughout resistance development in the SKOV3 cells (Figure 10). These data show the association of increased glu-tubulin expression levels with
- 10 acquired resistance to BAL27862.

List of abbreviations


A549	human non-small cell lung cancer cell line
AnnexinV	phosphatidylserine-binding protein
BCA	bicinchoninic acid
Bcl-2	B-cell lymphoma 2 protein
BRCA1	breast cancer type 1 susceptibility protein
BrdU	bromodeoxyuridine
BSA	bovine serum albumin
CA-125	cancer antigen 125
cDNA	complementary deoxyribonucleic acid
CREST	limited scleroderma syndrome
CO ₂	carbon dioxide
CXF 243	patient-derived colorectal tumour
CXF 1103	patient-derived colorectal tumour

DAB	3,3-diaminobenzidine
DMSO	dimethylsulphoxide
DNA	deoxyribonucleic acid
dUTP	2'-Deoxyuridine 5'-Triphosphate
ELISA	enzyme-linked immunosorbent assay
ErbB-2	human epidermal growth factor receptor 2
ESI-MS	electrospray ionisation mass-spectrometry
EtOH	Ethanol
FACS	fluorescence activated cell scan/sorting
FCS/FBS	foetal calf / foetal bovine serum
G2/M	transition from G2 to the mitotic phase in the cell cycle
HeLa	human squamous cell cancer cell line
HEPES	4-(2-Hydroxyethyl)piperazine-1-ethanesulphonic acid
Hoe33342	2'-(4'-Ethoxyphenyl)-5-(4-methylpiperazin-1-yl)-2,5'-bis-1H-benzimidazole trihydrochloride trihydrate
H460	human non-small cell lung cancer cell line
IgG	immunoglobulin G
IHC	Immunohistochemistry
LXFE 211	Patient-derived lung tumour
LXFE 397	Patient-derived lung tumour
MALDI	matrix-assisted-laser-desorption/ionisation mass-spectrometry
MALDI-TOF	matrix-assisted-laser-desorption/ionisation-time-of-flight-mass-spectrometry
MEXF 276	patient-derived melanoma
MEXF 1341	patient-derived melanoma
mRNA	messenger ribonucleic acid
MTS	3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulphophenyl)-2H-tetrazolium
NaCl	Sodium chloride
NaF	Sodium fluoride
NCBI	National Center for Biotechnology Information

NSCLC	non-small cell lung cancer
NP40	Nonidet P40
PBS	phosphate buffered saline
PCR	polymerase chain reaction
P-gp	P-glycoprotein
PMSF	phenylmethanesulphonyl fluoride
PSA	prostate-specific antigen
PVDF	polyvinylidene fluoride
RANO	response assessment for high-grade gliomas
RECIST	response evaluation criteria in solid tumours
RNA	ribonucleic acid
RPMI-1640	cell culture medium used for culturing transformed and non-transformed eukaryotic cells and cell lines
SDS	sodium dodecyl sulphate
SEQ. ID No.	sequence identification number
siRNA	small inhibitory ribonucleic acid
SKBR3	human mammary carcinoma cell line
SKOV3	human ovarian carcinoma cell line
TTCP	tubulin tyrosine carboxypeptidase
TTL	tubulin tyrosine ligase
TUNEL	terminal deoxynucleotidyl transferase dUTP nick end labeling
Tween-20	detergent, Polyoxyethylene sorbitan monolaurate
TX-100	Triton-X100
YO-PRO	fluorescent, monomeric cyanine, nucleic acid stain

CLAIMS

1. Use of glu-tubulin as a biomarker for predicting the response of a neoplastic disease or an autoimmune disease to a compound, wherein the compound is a compound of general formula I



wherein

R represents phenyl, thienyl or pyridinyl

wherein phenyl is optionally substituted by one or two substituents independently selected from alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, acyloxy-lower alkyl, phenyl, hydroxy, lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, phenyl-lower alkoxy, lower alkylcarbonyloxy, amino, monoalkylamino, dialkylamino, lower alkoxy-carbonylamino, lower alkylcarbonylamino, substituted amino wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, carboxy, lower alkoxy-carbonyl, cyano, halogen, and nitro; and wherein two adjacent substituents are methylenedioxy;

and wherein pyridinyl is optionally substituted by lower alkoxy, amino or halogen;

X represents a group C=Y, wherein Y stands for oxygen or nitrogen substituted by hydroxy or lower alkoxy;

R¹ represents hydrogen, lower alkylcarbonyl, hydroxy-lower alkyl or cyano-lower alkyl;

R², R³ and R⁶ represent hydrogen;

R⁴ and R⁵, independently of each other, represent hydrogen, lower alkyl or lower alkoxy; or R⁴ and R⁵ together represent methylenedioxy;

and pharmaceutically acceptable derivatives thereof;

or wherein

R represents phenyl or pyridinyl

wherein phenyl is optionally substituted by one or two substituents independently selected from alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, acyloxy-lower alkyl, phenyl, hydroxy, lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy,

phenyl-lower alkoxy, lower alkylcarbonyloxy, amino, monoalkylamino, dialkylamino, lower alkoxy-carbonylamino, lower alkylcarbonylamino, substituted amino wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, carboxy, lower alkoxy-carbonyl, formyl, cyano, halogen, and nitro; and wherein two adjacent substituents are methylenedioxy;

and wherein pyridinyl is optionally substituted by lower alkoxy, amino or halogen;

X represents oxygen;

R¹ represents hydrogen, lower alkylcarbonyl, hydroxy-lower alkyl or cyano-lower alkyl;

R², R³ and R⁶ represent hydrogen;

R⁴ and R⁵, independently of each other, represent hydrogen, lower alkyl or lower alkoxy; or R⁴ and R⁵ together represent methylenedioxy;

and pharmaceutically acceptable derivatives thereof,

and wherein the prefix lower denotes a radical having up to and including a maximum of 7, especially up to and including a maximum of 4 carbon atoms.

2. The use according to claim 1, wherein the response is of a disease in a subject and the biomarker glu-tubulin is measured ex vivo in a sample or samples taken from the human or animal body, preferably taken from the human body.

3. The use according to claim 1 or claim 2, wherein the compound is a compound of general formula I wherein

R represents phenyl or pyridinyl;

wherein phenyl is optionally substituted by one or two substituents independently selected from lower alkyl, lower alkoxy, amino, acetylamino, halogen and nitro;

and wherein pyridinyl is optionally substituted by amino or halogen;

X represents a group C=O;

R¹ represents hydrogen or cyano-lower alkyl;


R², R³, R⁴, R⁵ and R⁶ represent hydrogen;

and pharmaceutically acceptable derivatives thereof,

and wherein the prefix lower denotes a radical having up to and including a maximum of 7, especially up to and including a maximum of 4 carbon atoms.

4. The use according to any one of claims 1 to 3, wherein the compound is represented by the following formula

70




wherein R, Y and R¹ are defined as follows:

R	Y	R ¹
	O	CH ₂ CH ₂ CN
	O	H
	O	CH ₂ CH ₂ CN


or pharmaceutically acceptable derivatives thereof.

5. The use according to any one of claims 1 to 4, wherein the compound is



or pharmaceutically acceptable derivatives thereof.

6. The use according to any one of claims 1 to 5, wherein the pharmaceutically acceptable derivative is selected from the group consisting of a salt, solvate, pro-drug, salt of a pro-drug, polymorph and isomer of a compound of general formula I as defined in any one of claims 1 to 5.
7. The use according to claim 6, wherein the pro-drug is an amide formed from an amino group present within the R group of the compound of general formula I as defined in any one of claims 1 to 5 and the carboxy group of glycine, alanine or lysine.
8. The use according to any one of claims 1 to 7, wherein the compound is



or a pharmaceutically acceptable salt thereof, preferably a hydrochloride salt thereof, most preferably a dihydrochloride salt thereof.

9. The use according to any one of claims 1 to 8, for predicting the resistance of a disease in a subject to said compound.
10. The use according to any one of claims 1 to 9, wherein the disease is a neoplastic disease or autoimmune disease, preferably cancer.
11. The use according to any one of claims 1 to 10, wherein the disease is selected from the group consisting of breast cancer, prostate cancer, cervical cancer, ovarian cancer gastric cancer, colorectal cancer, pancreatic cancer, liver cancer, brain cancer, neuroendocrine cancer, lung


cancer, kidney cancer, hematological malignancies, melanoma and sarcomas.

12. The use according to any one of claims 1 to 10, wherein the disease is selected from the group consisting of breast cancer, cervical cancer, ovarian cancer, colorectal cancer, lung cancer and melanoma.
13. The use according to any one of claims 1 to 10, wherein the disease is selected from the group consisting of ovarian cancer, colorectal cancer, lung cancer and melanoma.
14. The use according to any one of claims 2 to 13, wherein a higher level of glu-tubulin in the sample from the subject relative to a standard value or set of standard values predicts resistance.
15. The use according to claim 14, wherein higher glu-tubulin levels in a sample or samples relative to a standard value or set of standard values from subjects with the same tumour histotype; or
taken after treatment initiation and compared to a sample or samples taken from the same subject before treatment initiation; or
iii) relative to a standard value or set of standard values from normal cells or tissue; are predictive of resistance.
16. The use according to any one of claims 1 to 15, wherein the biomarker is used to select subjects suffering or predisposed to suffering from a disease, preferably cancer, for treatment with a compound of general formula I or pharmaceutically acceptable derivatives thereof as defined in any one of claims 1 to 8.
17. The use according to any one of claims 2 to 16, wherein the sample is derived from normal tissue, tumour tissue, or circulating tumour cells, preferably wherein it is derived from tumour tissue.
18. A method for predicting the response of a neoplastic disease or an autoimmune disease in a subject to a compound of general formula I or a pharmaceutically acceptable derivative thereof as defined in any one of claims 1 to 8, comprising the steps of:
 - a) measuring a level of glu-tubulin in a sample pre-obtained from the subject to obtain a value or values representing this level; and
 - b) comparing the value or values from step a) to a standard value or set of standard

values.

19. The method according to claim 18, wherein the response which is predicted is resistance.
20. The method according to claim 18 or claim 19, wherein a higher level of glu-tubulin in the sample from the subject relative to the standard value or set of standard values predicts resistance.
21. A method of treating a neoplastic or autoimmune disease, preferably cancer, in a subject in need thereof, comprising measuring a level of glu-tubulin in a sample from the subject to obtain a value or values representing this level, and treating the subject with a compound of general formula I or a pharmaceutically acceptable derivative thereof as defined in claim 5 or claim 8, if the level of glu-tubulin in said sample is not higher than a standard value or set of standard values.
22. A method of treating a neoplastic or autoimmune disease, preferably cancer, by first decreasing the level of glu-tubulin in a subject that has a sample with a higher level of glu-tubulin compared to a standard level or set of standard levels, then treating the subject with a compound of general formula I or a pharmaceutically acceptable derivative thereof as defined in claim 5 or claim 8.
23. A kit when used for predicting the response of a cancer to a compound of general formula I or a pharmaceutically acceptable derivative thereof, as defined in any one of claims 1 to 8, comprising reagents necessary for measuring a level of glu-tubulin in a sample and further comprising a comparator module which comprises a standard value or set of standard values of a level of glu-tubulin taken from samples of tumour tissue or circulating tumour cells of subjects with a cancer of the same histotype to which the level of glu-tubulin in the sample is compared.
24. The kit when used according to claim 23, wherein the reagents comprise a capture reagent comprising a detector for glu-tubulin and a detector reagent, preferably wherein the capture reagent is an antibody.
25. The kit when used according to claim 23 or 24, wherein the kit comprises a compound of the following formula or a pharmaceutically acceptable salt thereof, in particular the dihydrochloride salt thereof,

74



26. A device for predicting the response of a neoplastic disease or an autoimmune disease to a compound of general formula I or a pharmaceutically acceptable derivative thereof, as defined in any one of claims 1 to 8, comprising reagents necessary for measuring the level of glu-tubulin in a sample and a comparator module which comprises a standard value or set of standard values to which the level of glu-tubulin in the sample is compared.

Basilea Pharmaceutica AG
Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11

Tubulin alpha-1A chain [Homo sapiens] (SEQ. ID. No. 1)

```
1 mrecisihvg qagvqignac welyclehgi qpdgqmpsdk tigggddsfn tffsetgagk
61 hvpravfvdL eptvidevrt gtyrqlfhpe qlitgkeda nnyarghyti gkeiidlvld
121 rirkladqct glqgflvfhs fgggtgsgft sllmerlsvd ygkksklefs iypapqvsta
181 vvepynsilt thttlehsdc afmvdneaiy dicrrnldie rptytnlnrl igqivssita
241 slrfdgalnv dltefqtnlv pyprihfpla tyapvisaek ayheqlsvae itnacfeapan
301 qmvkcdprhg kymaccllyr gdvvpkdvna aiatiktkrt iqfvdwcptg fkvginypqp
361 tvvpaggdlak vgravcmlsn ttaiaeawar ldhkfdlmya krafvhwvyg egmeegefse
421 aredmaalek dyeevgvdsV egegeeegee y
```

Figure 12

Tubulin alpha-1B chain [Homo sapiens] (SEQ. ID. No. 2)

```
1 mrecisihvg qagvqignac welyclehgi qpdgqmpsdk tigggddsfn tffsetgagk
61 hvpravfvdL eptvidevrt gtyrqlfhpe qlitgkeda nnyarghyti gkeiidlvld
121 rirkladqct glqgflvfhs fgggtgsgft sllmerlsvd ygkksklefs iypapqvsta
181 vvepynsilt thttlehsdc afmvdneaiy dicrrnldie rptytnlnrl isqivssita
241 slrfdgalnv dltefqtnlv pyprihfpla tyapvisaek ayheqlsvae itnacfeapan
301 qmvkcdprhg kymaccllyr gdvvpkdvna aiatiktkrs iqfvdwcptg fkvginypqp
361 tvvpaggdlak vgravcmlsn ttaiaeawar ldhkfdlmya krafvhwvyg egmeegefse
421 aredmaalek dyeevgvdsV egegeeegee y
```

Figure 13

Tubulin alpha-1C chain [Homo sapiens] (SEQ. ID. No. 3)

```
1 mrecisihvg qagvqignac welyclehgi qpdgqmpsdk tigggddsfn tffsetgagk
61 hvpravfvdl eptvidevrt gtyrqlfhpe qlitgkedaa nnyarghyti gkeiidlvld
121 rirkladqct glqgflvfhs fgggtgsgft sllmerlsvd ygkksklefs iypapqvsta
181 vvepynsilt thttlehsdc afmvdneaiy dicrrnldie rptytnlnrl isqivssita
241 slrfdgalnv dltefqtnlv pyprihfpla tyapvisaek ayheqltvaе itnacfepan
301 qmvkcdprhg kymaccllyr gdvvpkdvna aiatiktkrt iqfvdwcptg fkvginypqp
361 tvvpaggdlak vgravcmIsn ttavaeawar ldhkfdlmya krafvhwvyg egmeegefse
421 aredmaalek dyeevgadsa dgedegeey
```

Figure 14

Tubulin alpha-3C/D chain [Homo sapiens] (SEQ. ID. No. 4)

```
1 mrecisihvg qagvqignac welyclehgi qpdgqmpsdk tigggddsfn tffsetgagk
61 hvpravfvdl eptvvdevrt gtyrqlfhpe qlitgkedaa nnyarghyti gkeivdlvld
121 rirkladlct glqgflifhs fgggtgsgfa sllmerlsvd ygkksklefa iypapqvsta
181 vvepynsilt thttlehsdc afmvdneaiy dicrrnldie rptytnlnrl igqivssita
241 slrfdgalnv dltefqtnlv pyprihfpla tyapvisaek ayheqlsvae itnacfepan
301 qmvkcdprhg kymaccmlyr gdvvpkdvna aiatiktkrt iqfvdwcptg fkvginypqp
361 tvvpaggdlak vgravcmIsn ttataeawar ldhkfdlmya krafvhwvyg egmeegefse
421 aredlaalek dyeevgvdsv eaeaegeey
```

eol f - seq
SEQUENCE LISTING

<110> Basilea Pharmaceutica AG

<120> Use of glu-tubulin as a biomarker of drug response

<130> P40620EP00

<160> 4

<170> Patent In version 3.4

<210> 1

<211> 451

<212> PRT

<213> Homo sapiens

<400> 1

Met Arg Gu Cys Ile Ser Ile His Val Gly Gln Ala Gly Val Gln Ile
1 5 10 15

Gly Asn Ala Cys Trp Gu Leu Tyr Cys Leu Gu His Gly Ile Gln Pro
20 25 30

Asp Gly Gln Met Pro Ser Asp Lys Thr Ile Gly Gly Gly Asp Asp Ser
35 40 45

Phe Asn Thr Phe Phe Ser Gu Thr Gly Ala Gly Lys His Val Pro Arg
50 55 60

Ala Val Phe Val Asp Leu Gu Pro Thr Val Ile Asp Gu Val Arg Thr
65 70 75 80

Gly Thr Tyr Arg Gln Leu Phe His Pro Gu Gln Leu Ile Thr Gly Lys
85 90 95

Gu Asp Ala Ala Asn Asn Tyr Ala Arg Gly His Tyr Thr Ile Gly Lys
100 105 110

Gu Ile Ile Asp Leu Val Leu Asp Arg Ile Arg Lys Leu Ala Asp Gln
115 120 125

Cys Thr Gly Leu Gln Gly Phe Leu Val Phe His Ser Phe Gly Gly Gly
130 135 140

Thr Gly Ser Gly Phe Thr Ser Leu Leu Met Gu Arg Leu Ser Val Asp
145 150 155 160

Tyr Gly Lys Lys Ser Lys Leu Gu Phe Ser Ile Tyr Pro Ala Pro Gln
165 170 175

Val Ser Thr Ala Val Val Gu Pro Tyr Asn Ser Ile Leu Thr Thr His
180 185 190

Thr Thr Leu Gu His Ser Asp Cys Ala Phe Met Val Asp Asn Gu Ala
Page 1

eof - seq1

195

200

205

I l e T y r A s p I l e C y s A r g A r g A s n L e u A s p I l e G l u A r g P r o T h r T y r
 210 215 220

T h r A s n L e u A s n A r g L e u I l e G l y G n I l e V a l S e r S e r I l e T h r A l a
 225 230 235 240

S e r L e u A r g P h e A s p G l y A l a L e u A s n V a l A s p L e u T h r G l u P h e G n
 245 250 255

T h r A s n L e u V a l P r o T y r P r o A r g I l e H i s P h e P r o L e u A l a T h r T y r
 260 265 270

A l a P r o V a l I l e S e r A l a G l u L y s A l a T y r H i s G l u G n L e u S e r V a l
 275 280 285

A l a G l u I l e T h r A s n A l a C y s P h e G l u P r o A l a A s n G n M e t V a l L y s
 290 295 300

C y s A s p P r o A r g H i s G l y L y s T y r M e t A l a C y s C y s L e u L e u T y r A r g
 305 310 315 320

G l y A s p V a l V a l P r o L y s A s p V a l A s n A l a A l a I l e A l a T h r I l e L y s
 325 330 335

T h r L y s A r g T h r I l e G n P h e V a l A s p T r p C y s P r o T h r G l y P h e L y s
 340 345 350

V a l G l y I l e A s n T y r G n P r o P r o T h r V a l V a l P r o G l y G l y A s p L e u
 355 360 365

A l a L y s V a l G n A r g A l a V a l C y s M e t L e u S e r A s n T h r T h r A l a I l e
 370 375 380

A l a G l u A l a T r p A l a A r g L e u A s p H i s L y s P h e A s p L e u M e t T y r A l a
 385 390 395 400

L y s A r g A l a P h e V a l H i s T r p T y r V a l G l y G l u G l y M e t G l u G l u G l y
 405 410 415

G l u P h e S e r G l u A l a A r g G l u A s p M e t A l a A l a L e u G l u L y s A s p T y r
 420 425 430

G l u G l u V a l G l y V a l A s p S e r V a l G l u G l y G l u G l y G l u G l u G l y
 435 440 445

G l u G l u T y r
 450

<210> 2

eof - seq1

<211> 451
 <212> PRT
 <213> Homo sapiens

<400> 2

```

Met Arg Gu Cys Ile Ser Ile His Val Gly Gln Ala Gly Val Gln Ile
1      5      10      15

Gly Asn Ala Cys Trp Gu Leu Tyr Cys Leu Gu His Gly Ile Gln Pro
20      25      30

Asp Gly Gln Met Pro Ser Asp Lys Thr Ile Gly Gly Gly Asp Asp Ser
35      40      45

Phe Asn Thr Phe Phe Ser Gu Thr Gly Ala Gly Lys His Val Pro Arg
50      55      60

Ala Val Phe Val Asp Leu Gu Pro Thr Val Ile Asp Gu Val Arg Thr
65      70      75

Gly Thr Tyr Arg Gln Leu Phe His Pro Gu Gln Leu Ile Thr Gly Lys
85      90      95

Gu Asp Ala Ala Asn Asn Tyr Ala Arg Gly His Tyr Thr Ile Gly Lys
100     105     110

Gu Ile Ile Asp Leu Val Leu Asp Arg Ile Arg Lys Leu Ala Asp Gln
115     120     125

Cys Thr Gly Leu Gln Gly Phe Leu Val Phe His Ser Phe Gly Gly Gly
130     135     140

Thr Gly Ser Gly Phe Thr Ser Leu Leu Met Gu Arg Leu Ser Val Asp
145     150     155

Tyr Gly Lys Lys Ser Lys Leu Gu Phe Ser Ile Tyr Pro Ala Pro Gln
165     170     175

Val Ser Thr Ala Val Val Gu Pro Tyr Asn Ser Ile Leu Thr Thr His
180     185     190

Thr Thr Leu Gu His Ser Asp Cys Ala Phe Met Val Asp Asn Gu Ala
195     200     205

Ile Tyr Asp Ile Cys Arg Arg Asn Leu Asp Ile Gu Arg Pro Thr Tyr
210     215     220

Thr Asn Leu Asn Arg Leu Ile Ser Gln Ile Val Ser Ser Ile Thr Ala
225     230     235

Ser Leu Arg Phe Asp Gly Ala Leu Asn Val Asp Leu Thr Gu Phe Gln
245     250     255
  
```

eof - seq1

Thr Asn Leu Val Pro Tyr Pro Arg Ile His Phe Pro Leu Ala Thr Tyr
260 265 270

Ala Pro Val Ile Ser Ala Gu Lys Ala Tyr His Gu Gn Leu Ser Val
275 280 285

Ala Gu Ile Thr Asn Ala Cys Phe Gu Pro Ala Asn Gn Met Val Lys
290 295 300

Cys Asp Pro Arg His Gly Lys Tyr Met Ala Cys Cys Leu Leu Tyr Arg
305 310 315 320

Gly Asp Val Val Pro Lys Asp Val Asn Ala Ala Ile Ala Thr Ile Lys
325 330 335

Thr Lys Arg Ser Ile Gn Phe Val Asp Trp Cys Pro Thr Gly Phe Lys
340 345 350

Val Gly Ile Asn Tyr Gn Pro Pro Thr Val Val Pro Gly Gly Asp Leu
355 360 365

Ala Lys Val Gn Arg Ala Val Cys Met Leu Ser Asn Thr Thr Ala Ile
370 375 380

Ala Gu Ala Trp Ala Arg Leu Asp His Lys Phe Asp Leu Met Tyr Ala
385 390 395 400

Lys Arg Ala Phe Val His Trp Tyr Val Gly Gu Gly Met Gu Gu Gly
405 410 415

Gu Phe Ser Gu Ala Arg Gu Asp Met Ala Ala Leu Gu Lys Asp Tyr
420 425 430

Gu Gu Val Gly Val Asp Ser Val Gu Gly Gu Gly Gu Gu Gu Gly
435 440 445

Gu Gu Tyr
450

<210> 3
<211> 449
<212> PRT
<213> Homo sapiens

<400> 3

Met Arg Gu Cys Ile Ser Ile His Val Gly Gn Ala Gly Val Gn Ile
1 5 10 15

Gly Asn Ala Cys Trp Gu Leu Tyr Cys Leu Gu His Gly Ile Gn Pro
20 25 30

eof - seq1

Asp Gly Gln Met Pro Ser Asp Lys Thr Ile Gly Gly Gly Asp Asp Ser
 35 40 45
 Phe Asn Thr Phe Phe Ser Glu Thr Gly Ala Gly Lys His Val Pro Arg
 50 55 60
 Ala Val Phe Val Asp Leu Glu Pro Thr Val Ile Asp Glu Val Arg Thr
 65 70 75 80
 Gly Thr Tyr Arg Gln Leu Phe His Pro Glu Gln Leu Ile Thr Gly Lys
 85 90 95
 Glu Asp Ala Ala Asn Asn Tyr Ala Arg Gly His Tyr Thr Ile Gly Lys
 100 105 110
 Glu Ile Ile Asp Leu Val Leu Asp Arg Ile Arg Lys Leu Ala Asp Gln
 115 120 125
 Cys Thr Gly Leu Gln Gly Phe Leu Val Phe His Ser Phe Gly Gly Gly
 130 135 140
 Thr Gly Ser Gly Phe Thr Ser Leu Leu Met Glu Arg Leu Ser Val Asp
 145 150 155 160
 Tyr Gly Lys Lys Ser Lys Leu Glu Phe Ser Ile Tyr Pro Ala Pro Gln
 165 170 175
 Val Ser Thr Ala Val Val Glu Pro Tyr Asn Ser Ile Leu Thr Thr His
 180 185 190
 Thr Thr Leu Glu His Ser Asp Cys Ala Phe Met Val Asp Asn Glu Ala
 195 200 205
 Ile Tyr Asp Ile Cys Arg Arg Asn Leu Asp Ile Glu Arg Pro Thr Tyr
 210 215 220
 Thr Asn Leu Asn Arg Leu Ile Ser Gln Ile Val Ser Ser Ile Thr Ala
 225 230 235 240
 Ser Leu Arg Phe Asp Gly Ala Leu Asn Val Asp Leu Thr Glu Phe Gln
 245 250 255
 Thr Asn Leu Val Pro Tyr Pro Arg Ile His Phe Pro Leu Ala Thr Tyr
 260 265 270
 Ala Pro Val Ile Ser Ala Glu Lys Ala Tyr His Glu Gln Leu Thr Val
 275 280 285
 Ala Glu Ile Thr Asn Ala Cys Phe Glu Pro Ala Asn Gln Met Val Lys
 290 295 300

eof - seq1

Cys Asp Pro Arg His Gly Lys Tyr Met Ala Cys Cys Leu Leu Tyr Arg
305 310 315 320

Gly Asp Val Val Pro Lys Asp Val Asn Ala Ala Ile Ala Thr Ile Lys
325 330 335

Thr Lys Arg Thr Ile Gln Phe Val Asp Trp Cys Pro Thr Gly Phe Lys
340 345 350

Val Gly Ile Asn Tyr Gln Pro Pro Thr Val Val Pro Gly Gly Asp Leu
355 360 365

Ala Lys Val Gln Arg Ala Val Cys Met Leu Ser Asn Thr Thr Ala Val
370 375 380

Ala Glu Ala Trp Ala Arg Leu Asp His Lys Phe Asp Leu Met Tyr Ala
385 390 395 400

Lys Arg Ala Phe Val His Trp Tyr Val Gly Glu Gly Met Glu Glu Gly
405 410 415

Glu Phe Ser Glu Ala Arg Glu Asp Met Ala Ala Leu Glu Lys Asp Tyr
420 425 430

Glu Glu Val Gly Ala Asp Ser Ala Asp Gly Glu Asp Glu Gly Glu Glu
435 440 445

Tyr

<210> 4
<211> 450
<212> PRT
<213> Homo sapiens

<400> 4

Met Arg Glu Cys Ile Ser Ile His Val Gly Gln Ala Gly Val Gln Ile
1 5 10 15

Gly Asn Ala Cys Trp Glu Leu Tyr Cys Leu Glu His Gly Ile Gln Pro
20 25 30

Asp Gly Gln Met Pro Ser Asp Lys Thr Ile Gly Gly Gly Asp Asp Ser
35 40 45

Phe Asn Thr Phe Phe Ser Glu Thr Gly Ala Gly Lys His Val Pro Arg
50 55 60

Ala Val Phe Val Asp Leu Glu Pro Thr Val Val Asp Glu Val Arg Thr
65 70 75 80

eol f - seq
 Gly Thr Tyr Arg G n Leu Phe H i s Pro G u G n Leu I l e Thr Gly Lys
 85 90 95
 G u Asp Al a Al a Asn Asn Tyr Al a Arg Gly H i s Tyr Thr I l e Gly Lys
 100 105 110
 G u I l e Val Asp Leu Val Leu Asp Arg I l e Arg Lys Leu Al a Asp Leu
 115 120 125
 Cys Thr Gly Leu G n Gly Phe Leu I l e Phe H i s Ser Phe Gly Gly Gly
 130 135 140
 Thr Gly Ser Gly Phe Al a Ser Leu Leu Met G u Arg Leu Ser Val Asp
 145 150 155 160
 Tyr Gly Lys Lys Ser Lys Leu G u Phe Al a I l e Tyr Pro Al a Pro G n
 165 170 175
 Val Ser Thr Al a Val Val G u Pro Tyr Asn Ser I l e Leu Thr Thr H i s
 180 185 190
 Thr Thr Leu G u H i s Ser Asp Cys Al a Phe Met Val Asp Asn G u Al a
 195 200 205
 I l e Tyr Asp I l e Cys Arg Arg Asn Leu Asp I l e G u Arg Pro Thr Tyr
 210 215 220
 Thr Asn Leu Asn Arg Leu I l e Gly G n I l e Val Ser Ser I l e Thr Al a
 225 230 235 240
 Ser Leu Arg Phe Asp Gly Al a Leu Asn Val Asp Leu Thr G u Phe G n
 245 250 255
 Thr Asn Leu Val Pro Tyr Pro Arg I l e H i s Phe Pro Leu Al a Thr Tyr
 260 265 270
 Al a Pro Val I l e Ser Al a G u Lys Al a Tyr H i s G u G n Leu Ser Val
 275 280 285
 Al a G u I l e Thr Asn Al a Cys Phe G u Pro Al a Asn G n Met Val Lys
 290 295 300
 Cys Asp Pro Arg H i s Gly Lys Tyr Met Al a Cys Cys Met Leu Tyr Arg
 305 310 315 320
 Gly Asp Val Val Pro Lys Asp Val Asn Al a Al a I l e Al a Thr I l e Lys
 325 330 335
 Thr Lys Arg Thr I l e G n Phe Val Asp Trp Cys Pro Thr Gly Phe Lys
 340 345 350

Val Gly Ile Asn Tyr Gln Pro Pro Thr eol f - seq
 355 360 365 Val Val Pro Gly Gly Asp Leu
 Ala Lys Val Gln Arg Ala Val Cys Met Leu Ser Asn Thr Thr Ala Ile
 370 375 380
 Ala Glu Ala Trp Ala Arg Leu Asp His Lys Phe Asp Leu Met Tyr Ala
 385 390 395 400
 Lys Arg Ala Phe Val His Trp Tyr Val Gly Glu Gly Met Glu Glu Gly
 405 410 415
 Glu Phe Ser Glu Ala Arg Glu Asp Leu Ala Ala Leu Glu Lys Asp Tyr
 420 425 430
 Glu Glu Val Gly Val Asp Ser Val Glu Ala Glu Ala Glu Glu Gly Glu
 435 440 445
 Glu Tyr
 450