



(86) Date de dépôt PCT/PCT Filing Date: 2019/12/18
(87) Date publication PCT/PCT Publication Date: 2020/06/25
(45) Date de délivrance/Issue Date: 2023/10/03
(85) Entrée phase nationale/National Entry: 2021/06/15
(86) N° demande PCT/PCT Application No.: CN 2019/126178
(87) N° publication PCT/PCT Publication No.: 2020/125659
(30) Priorités/Priorities: 2018/12/18 (CN201811550604.8);
2019/12/04 (CN201911226373.X)

(51) Cl.Int./Int.Cl. *C07D 213/00* (2006.01),
A61K 31/136 (2006.01), *A61P 27/00* (2006.01)

(72) Inventeurs/Inventors:

LI, PENG, CN;
LI, XIAOLIN, CN;
LUO, ZHI, CN;
HE, HAIYING, CN;
HU, GUOPING, CN;
LI, JIAN, CN;
CHEN, SHUHUI, CN


(73) Propriétaire/Owner:

ZHUHAI UNITED LABORATORIES CO., LTD., CN

(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : COMPOSE DESTINE A ETRE UTILISE DANS DES MALADIES RETINIENNES

(54) Title: COMPOUND FOR USE IN RETINAL DISEASES





(57) Abrégé/Abstract:

An aldehyde binder compound as represented by formula (II) or a pharmaceutically acceptable salt. T_1 , T_2 , T_3 , and T_4 are each independently N, C, or CR_1 , T_5 is C, CR_5 , or C=O. T_6 is C, CR_6 , or N. T_7 is N or CR_7 . When T_5 is C=O and T_6 is N, then ==- is a single bond. L is a single bond, -O-, -S-, -NR₂- or -(CR₃R₄)_n. Each R_1 is independently H, F, Cl, Br, I, OH, or NH₂. R_2 is H or C₁₋₃ alkyl, optionally substituted. R_3 and R_4 are each independently H, F, Cl, Br, I, OH, NH₂, CN or C₁₋₃ alkyl, optionally substituted. R_5 , R_6 and R_7 are each independently H, F, Cl, Br or I. n is 1, 2, or 3. The compound may be used for treating the xerophthalmia.

(see formula II)

Abstract

An aldehyde binder compound as represented by formula (II) or a pharmaceutically acceptable salt. T_1 , T_2 , T_3 , and T_4 are each independently N, C, or CR_1 . T_5 is C, CR_5 , or $C=O$. T_6 is C, CR_6 , or N. T_7 is N or CR_7 . When T_5 is $C=O$ and T_6 is N, then  is a single bond. L is a single bond, -O-, -S-, -NR₂- or -(CR₃R₄)_n. Each R₁ is independently H, F, Cl, Br, I, OH, or NH₂. R₂ is H or C₁₋₃ alkyl, optionally substituted. R₃ and R₄ are each independently H, F, Cl, Br, I, OH, NH₂, CN or C₁₋₃ alkyl, optionally substituted. R₅, R₆ and R₇ are each independently H, F, Cl, Br or I. n is 1, 2, or 3. The compound may be used for treating the xerophthalmia.



Compound for Use in Retinal Diseases

References to Relevant Applications

[0001] This application claims the following priority:

[0002] CN201811550604.8, filed on December 18, 2018;

[0003] CN201911226373.X, filed on December 04, 2019.

Technical Field

[0004] The present disclosure relates to an aldehyde binder, in particular to a compound represented by formula (II) or a pharmaceutically acceptable salt.

Background

[0005] Xerophthalmia, also known as conjunctival xerosis, refers to a general term of a variety of diseases characterized by decreased tear film stability due to abnormal tear quality or quantity or kinetic abnormalities for any reasons, accompanied by eye discomfort (or) ocular surface tissue lesions, with specific discomfort symptoms manifested as eye irritation, visual disturbance, and tear film instability. Some of such syndromes are caused by ocular surface inflammation, resulting in loss of lacrimal functions. In addition, the syndromes are also associated with systemic autoimmunity.


[0006] Because some toxic aldehydes, such as malondialdehyde (MDA), 4-hydroxy-2-nonenal (4HNE), etc., are produced by metabolic mechanisms of *in vivo* or ocular tissues and organs, these aldehydes highly react with proteins, carbohydrates, grease, and DNA, resulting in chemically modified biomolecules to activate modulators of inflammatory molecules such as NF-kappaB, thereby causing injury to different organs, which is one of the triggers of xerophthalmia.

[0007] After studies by the present disclosure, a small-molecule drug enters the ocular inflammatory site in the form of eye drops or oral administration, through complexation reaction with aldehydes *in vivo*, thereby reducing aldehyde toxicity and decreasing inflammation to exerting the effect of treating xerophthalmia.

Content of the invention

[0008] The present disclosure provides a compound represented by formula (I) or a

pharmaceutically acceptable salt thereof,



(I)

[0009] wherein

[0010] T₁, T₂, T₃, and T₄ are each independently selected from N, C, and CR₁;

[0011] L is selected from a single bond, -O-, -S-, -NR₂- and -(CR₃R₄)_n-;

[0012] each R₁ is independently selected from H, F, Cl, Br, I, OH, and NH₂;

[0013] R₂ is selected from H and C₁₋₃ alkyl optionally substituted with 1, 2 or 3 R_a;

[0014] R₃ and R₄ are each independently selected from H, F, Cl, Br, I, OH, NH₂, CN and C₁₋₃ alkyl optionally substituted with 1, 2 or 3 R_b;

[0015] n is selected from 1, 2, and 3;

[0016] R_a and R_b are each independently selected from H, F, Cl, Br, I, OH, NH₂, CN, and CH₃

[0017] In some embodiments of the present disclosure, R₂ is selected from H, CH₃ and CH₂CH₃, wherein the CH₃ and the CH₂CH₃ are optionally substituted with 1, 2 or 3 R_a, and other variables are as defined in the present disclosure.

[0018] In some embodiments of the present disclosure, R₂ is selected from H, CH₃ and CH₂CH₃, and other variables are as defined in the present disclosure.


[0019] In some embodiments of the present disclosure, R₃ and R₄ are each independently selected from H, F, Cl, Br, I, OH, NH₂, CN, CH₃ and CH₂CH₃, wherein the CH₃ and the CH₂CH₃ are optionally substituted with 1, 2, or 3 R_b, and other variables are as defined in the present disclosure.

[0020] In some embodiments of the present disclosure, R₃ and R₄ are each independently selected from H, F, Cl, Br, I, OH, NH₂, CN, CH₃ and CH₂CH₃, and other

variables are as defined in the present disclosure.

[0021] In some embodiments of the present disclosure, L is selected from a single bond, -O-, -S-, -NH-, -(CH₂)₂- and -CH₂-, and other variables are as defined in the present disclosure.

[0022] The present invention provides a compound represented by formula (II) or a pharmaceutically acceptable salt thereof,



[0023] wherein

[0024] is selected from a single bond and a double bond;

[0025] T₁, T₂, T₃, and T₄ are each independently selected from N, C, and CR₁;

[0026] T₅ is selected from C, CR₅, and C=O;

[0027] T₆ is selected from C, CR₆, and N;

[0028] T₇ is selected from N and CR₇;

[0029] when T₅ is selected from C=O and T₆ is selected from N, then is selected from a single bond;

[0030] L is selected from a single bond, -O-, -S-, -NR₂- and -(CR₃R₄)_n-;

[0031] each R₁ is independently selected from H, F, Cl, Br, I, OH, and NH₂;

[0032] R₂ is selected from H and C₁₋₃ alkyl optionally substituted with 1, 2 or 3 R_a;

[0033] R₃ and R₄ are each independently selected from H, F, Cl, Br, I, OH, NH₂, CN and C₁₋₃ alkyl optionally substituted with 1, 2 or 3 R_b;

[0034] R₅, R₆ and R₇ are each independently selected from H, F, Cl, Br and I;

[0035] n is selected from 1, 2, and 3;

[0036] R_a and R_b are each independently selected from H, F, Cl, Br, I, OH, NH₂, CN, and CH₃.

[0037] In some embodiments of the present disclosure, R_2 is selected from H, CH₃ and CH₂CH₃, wherein the CH₃ and the CH₂CH₃ are optionally substituted with 1, 2 or 3 R_a , and other variables are as defined in the present disclosure.

[0038] In some embodiments of the present disclosure, R_2 is selected from H, CH₃ and CH₂CH₃, and other variables are as defined in the present disclosure.


[0039] In some embodiments of the present disclosure, R_3 and R_4 are each independently selected from H, F, Cl, Br, I, OH, NH₂, CN, CH₃ and CH₂CH₃, wherein the CH₃ and the CH₂CH₃ are optionally substituted with 1, 2, or 3 R_b , and other variables are as defined in the present disclosure.

[0040] In some embodiments of the present disclosure, R_3 and R_4 are each independently selected from H, F, Cl, Br, I, OH, NH₂, CN, CH₃ and CH₂CH₃, and other variables are as defined in the present disclosure.

[0041] In some embodiments of the present disclosure, L is selected from a single bond, -O-, -S-, -NH-, -(CH₂)₂- and -CH₂-, and other variables are as defined in the present disclosure.

[0042] Some embodiments of the present disclosure are derived from any combination of the variables described above.

[0043] In some embodiments of the present disclosure, the compound described above or a pharmaceutically acceptable salt thereof is selected from




[0044] wherein

[0045] T₃ and T₄ are each independently selected from N and CR₁;

[0046] R₁ and L are as defined in the present disclosure.


[0047] In some embodiments of the present disclosure, the compound described above or a pharmaceutically acceptable salt thereof is selected from




[0048] wherein

[0049] R₁ and L are as defined in the present disclosure.

[0050] The present disclosure also provides a compound represented by the following formula or a pharmaceutically acceptable salt thereof, wherein the compound represented by the following formula or the pharmaceutically acceptable salt thereof is selected from:





[0051] The present disclosure also provides a pharmaceutical composition comprising a therapeutically effective amount of the compound, or the pharmaceutically acceptable salt thereof as an active ingredient, and a pharmaceutically acceptable carrier.

[0052] In some embodiments of the present disclosure, the compound or pharmaceutically acceptable salts thereof or the use of the composition in the manufacture of the medicament related to aldehyde binder.

[0053] In some embodiments of the present disclosure, in the use described above, the medicament related to aldehyde binder is used for the treatment of xerophthalmia.

Definition and Explanation

[0054] Unless otherwise stated, the following terms and phrases used herein are intended to have the following meanings. A particular term or phrase should not be considered uncertain or unclear if not specially defined, but should be understood in its ordinary meaning. When a trade name appears herein, it is intended to refer to its corresponding commodity or its active ingredient. The term "pharmaceutically acceptable" used herein refers to those compounds, materials, compositions and/or dosage forms that are within the scope of reliable medical judgment, suitable for use in contact with human and animal tissues, without excessive toxicity, irritation, allergic reactions or other problems or complications, and commensurate with a reasonable benefit/risk ratio.

[0055] The term "pharmaceutically acceptable salt" refers to a salt of a compound of the present disclosure that is prepared from a compound with a specific substituent found in the present disclosure with a relatively nontoxic acid or base. When the compound of the present disclosure contains a relatively acidic functional group, a base

addition salt can be obtained by a neutral contact between a sufficient amount of base and such compounds in a pure solution or a suitable inert solvent. The pharmaceutically acceptable base addition salt includes sodium, potassium, calcium, amine, organic amine or magnesium salts, or similar salts. When the compound of the present disclosure contains a relatively basic functional group, an acidic addition salt can be obtained by a neutral contact between a sufficient amount of acid and such compounds in a pure solution or a suitable inert solvent. Examples of the pharmaceutically acceptable acid addition salt include salts of inorganic acids including, for example, hydrochloric acid, hydrobromic acid, nitric acid, carbonic acid, bicarbonate, phosphoric acid, monohydrogen phosphate, dihydrogen phosphate, sulfuric acid, hydrogen sulfate, hydroiodic acid, phosphorous acid, etc., salts of organic acids including, for example, acetic acid, propionic acid, isobutyric acid, maleic acid, malonic acid, benzoic acid, succinic acid, suberic acid, fumaric acid, lactic acid, mandelic acid, phthalic acid, benzenesulfonic acid, p-toluenesulfonic acid, citric acid, tartaric acid, methanesulfonic acid, and similar acids, salts of amino acids such as arginine, etc., and salts of organic acids such as glucuronic acid, etc. Some specific compounds of the present disclosure contain basic and acidic functional groups so that the compounds can be converted to either base or acid addition salts.

[0056] The pharmaceutically acceptable salt of the present disclosure can be synthesized by parent compounds containing acid radicals or bases with conventional chemical methods. In general, such salts are prepared by reacting the compounds in forms of free acids or bases with stoichiometrically appropriate bases or acids in water or an organic solvent or a mixture thereof.

[0057] In addition to the salt form, the compound provided by the present disclosure also has a prodrug form. The prodrug of the compound described herein readily has chemical changes under physiological conditions so that the prodrug is converted into the compound of the present disclosure. In addition, the prodrug may be converted into the compound of the present disclosure in the *in vivo* environment by chemical or biochemical methods.

[0058] Some compounds of the present disclosure may be present in an unsolvated or solvated form, including a hydrate form. Generally, the solvated form is comparable to the unsolvated form, both of which are within the scope of the present disclosure.

[0059] Optically active (R)-and (S)-isomers and D and L isomers can be prepared by chiral synthesis or chiral reagents or other conventional techniques. If one enantiomer of some compound of the present disclosure is desired, the enantiomer can be prepared by asymmetric synthesis or derivatization with a chiral auxiliary, wherein the obtained diastereo-isomer mixture is separated and an auxiliary group splits to provide a pure desired enantiomer. Alternatively, when a molecule contains a basic functional group (such as amino) or an acidic functional group (such as carboxyl), a salt of the diastereo-isomer is formed by the functional group with a suitably optically active acid or base, and the diastereo-isomer is resolved by a conventional method known in the art and then recovered to obtain a pure enantiomer. In addition, the separation of the enantiomer and the diastereo-isomer is usually completed by using chromatography, wherein the chromatography employs a chiral stationary phase and is optionally combined with chemical derivatization (e.g., formation of carbamate from amine). The compound of the present disclosure may contain atomic isotopes in unnatural proportions on one or more atoms constituting the compound. For example, the compound may be labeled by radioactive isotopes, such as tritium (^3H), iodine-125 (^{125}I) or C-14 (^{14}C). For example, hydrogen can be substituted with heavy hydrogen to form a deuterated drug, and a bond composed of the tritium and carbon is stronger than that of ordinary hydrogen and carbon. Compared with a non-deuterated drug, the deuterated drug has the advantages of reduced toxic and side effects, increased drug stability, enhanced efficacy, prolonged biological half-life of the drug, etc. All transformations in all isotopic compositions of the compound of the present disclosure, whether radioactive or not, are included within the scope of the present disclosure.

[0060] The term "optional" or "optionally" refers to that an event or condition described herein below may or may not occur, and the description includes presence of the event or condition and absence of the event or condition.

[0061] The term "substituted" refers to that any one or more hydrogen atoms on a particular atom are substituted with a substituent, wherein the substituent may include variants of heavy hydrogen and hydrogen, provided that a valence state of the particular atom is normal and the substituted compound is stable. When the substituent is oxygen (i.e.=O), two hydrogen atoms are substituted. Oxygen substitution does not occur on aryl. The term "optionally substituted" means that the atom may be substituted or unsubstituted; unless otherwise specified, the type and number of substituents can be


arbitrary provided that the substituents can be chemically realized.


[0062] When any variable (e.g. R) appears more than once in the composition or structure of the compound, the definition of the variable in each case is independent. Therefore, for example, if one group is substituted with 0-2 Rs, the group may optionally be substituted with at most 2 Rs, and the R has an independent option in each case. Furthermore, combinations of the substituents and/or variants thereof are allowed only if such combinations result in stable compounds.


[0063] When the number of linking groups is 0, e.g. -(CRR)0- indicates that the linking group is a single bond.

[0064] When one of the variables is selected from the single bond, two groups linked by the single bond are directly linked, for example, when L in A-L-Z represents the single bond, the structure is actually A-Z.

[0065] When one substituent is vacant, the substituent is absent, for example, when X in A-X is vacant, the structure is actually A. When the listed substituents do not indicate by which atom the substituents are linked to the substituted group, such substituents can be bonded through any atom thereof, for example, pyridyl as the substituent can be linked to the substituted group through any carbon atom on a pyridine ring. When the listed linking group does not indicate a linking direction thereof, the linking direction

thereof is arbitrary, e.g., the linking group L in  is -M-W-, at which time the -M-W- not only can be linked to rings A and B in the same direction as the

reading order from left to right to constitute , but also can be linked to the rings A and B in the opposite direction as the reading order from left to

right to constitute . Combinations of the linking group, the substituents and/or variants thereof are allowed only if such combinations result in stable compounds.

[0066] Unless otherwise specified, the term "C₁₋₆ alkyl" is used to represent a linear or branched saturated hydrocarbon group consisting of 1 to 6 carbon atoms. The C₁₋₆ alkyl includes C₁₋₅, C₁₋₄, C₁₋₃, C₁₋₂, C₂₋₆, C₂₋₄, C₆ and C₅ alkyl, etc.; C₁₋₆ alkyl may be

monovalent (e.g. methyl), divalent (e.g., methylene), or multivalent (e.g., methyne). Examples of C₁₋₆ alkyl include, but are not limited to, methyl (Me), ethyl (Et), propyl (including n-propyl and isopropyl), butyl (including n-butyl, isobutyl, s-butyl and t-butyl), pentyl (including n-pentyl, isopentyl and neopentyl), hexyl, etc.

[0067] Unless otherwise specified, the term "C₁₋₃ alkyl" is used to represent a linear or branched saturated hydrocarbon group consisting of 1 to 6 carbon atoms. The C₁₋₃ alkyl includes C₁₋₂ and C₂₋₃ alkyl, etc.; the C₁₋₃ alkyl may be monovalent (e.g. methyl), divalent (e.g., methylene), or multivalent (e.g., methyne). Examples of C₁₋₃ alkyl include, but are not limited to, methyl (Me), ethyl (Et), propyl (including n-propyl and isopropyl), etc.

[0068] Unless otherwise specified, the term "C₁₋₃ alkoxy" represents those alkyl groups which are linked to the rest of a molecule by one oxygen atom and contains 1 to 3 carbon atoms. The C₁₋₃ alkoxy includes C₁₋₂, C₂₋₃, C₃ and C₂ alkoxy, etc. Examples of C₁₋₃ alkoxy include, but are not limited to, methoxy, ethoxy, propoxy (including n-propoxy and isopropoxy), etc.

[0069] Unless otherwise specified, C_{n-n+m} or C_{n-C_{n+m}} includes any particular case of n to n+m carbons, for example, C₁₋₁₂ includes C₁, C₂, C₃, C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, and C₁₂, and furthermore includes any one range from n to n+m, for example, C₁₋₁₂ includes C₁₋₃, C₁₋₆, C₁₋₉, C₃₋₆, C₃₋₉, C₃₋₁₂, C₆₋₉, C₆₋₁₂, and C₉₋₁₂, etc.; similarly, n membered to n+m membered indicates that the number of atoms on the ring is from n to n+m, for example, a 3-12 membered ring includes a 3 membered ring, a 4 membered ring, a 5 membered ring, a 6 membered ring, a 7 membered ring, an 8 membered ring, a 9 membered ring, a 10 membered ring, an 11 membered ring, and a 12 membered ring, and furthermore includes any one range from n to n+m, for example, a 3-12 membered ring includes a 3-6 membered ring, a 3-9 membered ring, a 5-6 membered ring, a 5-7 membered ring, a 6-7 membered ring, a 6-8 membered ring, a 6-10 membered ring, etc.

[0070] The term "leaving group" refers to a functional group or an atom that may be substituted with another functional group or atom by a substitution reaction (e.g., an affinity substitution reaction). For example, representative leaving groups include trifluoromethanesulfonate, chlorine, bromine, iodine, sulfonate such as methanesulfonate, toluenesulfonate, p-bromobenzenesulfonate, p-toluenesulfonate,

etc., acyloxy such as acetoxy, trifluoroacetoxy, etc.

[0071] The term "protective group" includes, but is not limited to, "an amino protective group", "a hydroxy protective group", or "a sulfhydryl protective group". The term "amino protective group" refers to a protective group suitable for preventing side reactions at a nitrogen position of the amino. Representative amino protective groups include, but are not limited to, formyl, acyl, such as alkanoyl (e.g., acetyl, trichloroacetyl, or trifluoroacetyl), alkoxy carbonyl, such as tert-butoxycarbonyl (Boc), arylmethoxycarbonyl, such as benzyloxycarbonyl (Cbz) and 9-fluorenylmethoxycarbonyl (Fmoc), arylmethyl, such as benzyl (Bn), triphenylmethyl (Tr), 1,1-di-(4'-methoxyphenyl) methyl, silyl such as trimethylsilyl (TMS), tert-butyl dimethylsilyl (TBS), etc. The term "hydroxyl protective group" refers to a protective group suitable for preventing side reactions of hydroxyl. Representative hydroxyl protective groups include, but are not limited to, alkyl, such as methyl, ethyl, and tert-butyl, acyl, such as alkanoyl (e.g., acetyl), arylmethyl, such as benzyl (Bn), p-methoxybenzyl (PMB), 9-fluorenylmethyl (Fm), and diphenylmethyl (DPM), silyl, such as trimethylsilyl (TMS), tert-butyl dimethylsilyl (TBS), etc.

[0072] The compound of the present disclosure can be prepared by a variety of synthetic methods known to those skilled in the art, including specific embodiments listed below, embodiments formed in combination with other chemical synthesis methods, and equivalent substitutions known to those skilled in the art. Preferred embodiments include, but are not limited to, examples of the present disclosure.

[0073] The solvent used in the present disclosure is commercially available. The following abbreviations are used in the present disclosure: aq represents water; HATU represents O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethylurea hexafluorophosphate; EDC represents N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride; m-CPBA represents 3-chloroperoxybenzoic acid; eq represents equivalent and equal amount; CDI represents carbonyl diimidazole; DCM represents dichloromethane; PE represents PE; DIAD represents diisopropyl azodicarboxylate; DMF represents N,N-dimethylformamide; DMSO represents dimethyl sulfoxide; EtOAc represents ethyl acetate; EtOH represents ethanol; MeOH represents methanol; CBz represents benzyloxycarbonyl, which is an amine protective group; BOC represents tert-butoxycarbonyl, which is an amine protective group; HOAc

represents acetic acid; NaCNBH₃ represents sodium cyanoborohydride; r.t. represents room temperature; O/N represents overnight; THF represents tetrahydrofuran; Boc₂O represents di-tert-butyl dicarbonate; TFA represents trifluoroacetic acid; DIPEA represents diisopropylethylamine; SOCl₂ represents thionyl chloride; CS₂ represents carbon disulfide; TsOH represents p-toluenesulfonic acid; NFSI represents N-fluoro-N-(benzenesulfonyl) benzenesulfonamide; NCS represents N-chlorosuccinimide; n-Bu₄NF represents tetrabutylammonium fluoride; iPrOH represents 2-propanol; mp represents melting point; LDA represents lithium diisopropylamide; LiHMDS represents lithium hexamethyldisilazide; Xantphos represents 4,5-bisdiphenylphosphine-9,9-dimethylxanthene; LiAlH₄ represents lithium aluminum hydride; Pd(dba)₂ represents tris(dibenzylideneacetone) dipalladium; mCPBA represents m-chloroperoxybenzoic acid; Pd(dppf)Cl₂ represents [1,1'-bis(diphenylphosphino) ferrocene] dichloropalladium (II); DBU represents 1,8-diazabicyclo[5.4.0] undec-7-ene.

[0074] Compounds are named according to conventional nomenclature principles in the art or using ChemDraw[®] software, and commercially available compounds are named in the vendor's catalogue.

Brief Description of the Drawings

[0075] Figure 1: Test results of *in vitro* aldehyde capture capacity;

[0076] Figure 2: Tear secretion results of an efficacy test on xerophthalmia in mice;


[0077] Figure 3: Fluorescence staining results of cornea in an efficacy test on xerophthalmia in mice.

Technical Effect

[0078] The compound of the present disclosure has excellent aldehyde complexing capacity, and is helpful to remit eye inflamed, thereby treating the xerophthalmia.

Detailed Description of the Embodiment


[0079] The following examples further illustrate the present disclosure, but the present disclosure is not adversely limited thereto. The present disclosure has been described in detail herein, wherein specific embodiments thereof are also disclosed, and it will be apparent to those skilled in the art to make various changes and improvements to




[0085] A substrate 1-2 (200 mg, 605 μmol , 1 eq) was dissolved in tetrahydrofuran (20 mL), methyl magnesium bromide (3 M, 4.04 mL, 20 eq) was slowly added dropwise at 0 °C, and the mixture was reacted under stirring at 0 °C for 2 hours. A reaction solution was quenched by adding water (20 mL), then extracted with ethyl acetate (50 mL*3), and concentrated for organic phases. The mixture was dissolved in dimethylformamide (DMF) and purified by HPLC (neutral) to obtain Compound 1.

[0086] ^1H NMR (400MHz, DMSO- d_6) δ 7.80 (d, J=8.3 Hz, 2H), 7.00 (d, J=8.3 Hz, 2H), 5.56 (s, 4H), 5.47 (s, 2H), 1.56 (s, 12H); LCMS: $[\text{M}+\text{H}]^+=302.9$


[0087] Example 2: Synthesis of Compound 2



[0088] Synthetic Route of Compound 2:




[0089] Step 1: Synthesis of Compound 2-3



[0090] Substrates 2-1 (702.07 mg, 2.55 mmol, 1.5 eq) and 2-2 (310 mg, 1.70 mmol, 1 eq) were dissolved in acetonitrile (20 mL), cesium carbonate (1.11 g, 3.40 mmol, 2 eq) was added, and the mixture was reacted under stirring at 50 °C for 3 hours. After the reaction was over, a reaction solution was filtered directly, and filtrate was spin-dried to obtain crude. The crude was purified by a rapid silica gel column (petroleum ether: ethyl acetate=100: 0 to 60: 40) to obtain Compound 2-3.

[0091] ^1H NMR (400 MHz, CDCl_3) δ 8.42 (d, $J=9.03$ Hz, 1H), 7.24 (d, $J=4.02$ Hz, 1H), 7.15-7.20 (m, 1H), 7.07-7.13 (m, 1H), 5.82 (br s, 2H), 4.39 (dq, $J=4.27, 7.11$ Hz, 4H), 1.35 (td, $J=7.15, 18.57$ Hz, 6H).


[0092] Step 2: Synthesis of Compound 2-4



[0093] The Compound 2-3 (360 mg, 956.63 μmol , 1 eq) and hydrochloric acid (12 M, 478.32 μL , 6 eq) were dissolved in ethanol (20 mL) and water (5 mL); reducing iron powders (534.23 mg, 9.57 mmol, 10 eq) were added, and stirred at 25 °C for 2 hours. After the reaction was over, a reaction solution was adjusted to pH=9 with a saturated sodium carbonate aqueous solution (30 mL), and extracted with ethyl acetate (30 mL*2). After organic phases were combined, the organic phases were dried over anhydrous sodium sulfate and spin-dried to obtain crude. The crude was purified by a rapid silica gel column (petroleum ether: ethyl acetate=100: 0 to 40: 60) to obtain Compound 2-4.

[0094] ^1H NMR (400 MHz, CDCl_3) δ 7.17-7.24 (m, 2H), 7.09-7.16 (m, 2H), 5.64 (br s, 4H), 4.37 (q, $J=7.03$ Hz, 4H), 1.37 (t, $J=7.03$ Hz, 6H).

[0095] Step 3: Synthesis of Compound 2




[0096] The Compound 2-4 (120 mg, 346.48 μmol , 1 eq) was dissolved in

tetrahydrofuran (10 mL) and cooled to 0 °C. Methyl magnesium bromide (3 M, 3.00 mL, 26 eq) was added dropwise in a 2-methyltetrahydrofuran solution at 0 °C. The solution was reacted under stirring at 0 °C for 30 min, and then stirred at 25 °C for 2 hours. After the reaction was over, a reaction solution was quenched by adding a saturated amine chloride aqueous solution (40 mL), and then extracted with ethyl acetate (100 mL). The organic phases were dried over anhydrous sodium sulfate and spin-dried to obtain the crude. The crude was purified by HPLC separation (column model: Waters Xbridge 150*25 mm 5 μm; mobile phase: [water (10 mM amine bicarbonate solution)-acetonitrile]; %B (acetonitrile): 0%-50%, 10 min) to obtain Compound 2.


[0097] LCMS: [MS+H⁺]=318.9.

[0098] ¹H NMR (400 MHz, DMSO-d₆) δ 7.03 (d, J=8.28 Hz, 2H), 6.57 (d, J=8.53 Hz, 2H), 5.37 (s, 2H), 5.27 (s, 4H), 1.38 (s, 12H).


[0099] Example 3: Synthesis of Compound 3



[0100] Synthetic Route of Compound 3:




[0101] Step 1: Synthesis of Compound 3-3



[0102] Compounds 3-1 (300 mg, 1.30 mmol, 1 eq) and 3-2 (379.73 mg, 1.69 mmol, 1.3 eq), potassium phosphate (551.24 mg, 2.60 mmol, 2 eq) and [1,1'-bis(diphenylphosphino) ferrocene] palladium (II) dichloromethane (530.18 mg, 649.22 μ mol, 0.5 eq) were dissolved in dichloroethane (15 mL), and stirred at 80 °C for 15 hours. After the reaction was over, a reaction solution was filtered directly, and filtrate was spin-dried to obtain crude. The crude was purified by a rapid silica gel column (petroleum ether: ethyl acetate=100: 0 to 50: 50) to obtain Compound 3-3.

[0103] ^1H NMR (400 MHz, CDCl_3) δ 8.29 (s, 1H), 8.08 (s, 1H), 7.88 (s, 2H), 7.25 (s, 1H), 5.95 (br s, 2H), 4.03 (s, 3H), 3.96 (s, 3H).

[0104] Step 2: Synthesis of Compound 3-4




[0105] The Compound 3-3 (250 mg, 754.65 μ mol, 1 eq) and hydrochloric acid (12 M, 1.26 mL, 20 eq) were dissolved in methanol (12 mL) and water (3 mL); reducing iron powders (421.43 mg, 7.55 mmol, 10 eq) were added, and stirred at 25 °C for 1 day and 15 hours. After the reaction was over, a reaction solution was adjusted to pH=9 with a saturated sodium carbonate aqueous solution (50 mL), and extracted with ethyl acetate (60 mL*2). After organic phases were combined, the organic phases were dried over anhydrous sodium sulfate and spin-dried to obtain crude. The crude was purified by a rapid silica gel column (petroleum ether: ethyl acetate=100: 0 to 50: 50) to obtain Compound 3-4.

[0106] ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ 8.09 (d, $J=2.01$ Hz, 1H), 7.82 (d, $J=8.28$ Hz, 1H), 7.43 (d, $J=2.01$ Hz, 1H), 7.08 (d, $J=1.51$ Hz, 1H), 6.71-6.86 (m, 5H), 3.83 (d,

$J=5.77$ Hz, 6H).

[0107] Step 3: Synthesis of Compound 3




[0108] The Compound 3-4 (80 mg, 265.52 μmol , 1 eq) was dissolved in tetrahydrofuran (10 mL) and cooled to 0 °C. Methyl magnesium bromide (3 M, 2 mL, 22.60 eq) was added dropwise in a 2-methyltetrahydrofuran solution at 0 °C. The solution was reacted under stirring at 0 °C for 30 min, and then stirred at 25 °C for 15 hours. After the reaction was over, a reaction solution was quenched by adding a saturated amine chloride aqueous solution (20 mL), and then extracted in ethyl acetate (50 mL). The organic phases were dried over anhydrous sodium sulfate and spin-dried to obtain the crude. The crude was purified by thin-layer chromatographic silica gel plate separation (mobile phase: petroleum ether: ethyl acetate=2: 5) to obtain Compound 3.


[0109] LCMS: $[\text{MS}+\text{H}^+]=301.9$.

[0110] ^1H NMR (400 MHz, CDCl_3) δ 8.12 (d, $J=1.76$ Hz, 1H), 7.20 (d, $J=8.03$ Hz, 1H), 7.07 (d, $J=2.01$ Hz, 1H), 6.85 (dd, $J=2.01, 8.03$ Hz, 1H), 6.81 (d, $J=2.01$ Hz, 1H), 4.31-5.07 (m, 4H), 3.32 (br s, 2H), 1.71 (d, $J=5.27$ Hz, 12H).


[0111] **Example 4: Synthesis of Compound 4**



[0112] **Synthetic Route of Compound 4:**




[0113] Step 1: Synthesis of Compound 4-2



[0114] The Compound 3-1 (3 g, 12.98 mmol, 1 eq) was dissolved in tetrahydrofuran (10 mL), a reaction temperature was reduced to $-78\text{ }^{\circ}\text{C}$, and lithium methyl (1 M, 64.92 mL, 5 eq) was added dropwise and stirred at $-78\text{ }^{\circ}\text{C}$ for 1 hour. TLC plate monitoring (petroleum ether: ethyl acetate=3: 1) showed a new point generated in the reaction. A reaction solution was quenched by adding water (50 mL), and then extracted with ethyl acetate (50 mL*2). Organic phases were spin-dried. Products were purified by a rapid silica gel column (petroleum ether/ethyl acetate=3: 1 to 1: 1) to obtain Compound 4-2.

[0115] $^1\text{H NMR}$ (400MHz, CDCl_3) δ 7.93 (d, $J=2.0$ Hz, 1H), 7.06 (d, $J=2.0$ Hz, 1H), 4.67 (br s, 2H), 1.64 (s, 6H)

[0116] Step 2: Synthesis of Compound 4




[0117] The Compound 4-2 (500 mg, 2.16 mmol), hexamethylditin (1.74 mmol, 360 mL), and bis (tri-tert-butylphosphine) palladium(0) (330 mg, 645.73 μmol) were added in toluene (15 mL), and stirred under nitrogen atmosphere for reaction at $80\text{ }^{\circ}\text{C}$ for 12 hours. Thin layer chromatography (dichloromethane: methanol=10: 1) showed formation of a new point with large polarity. After the reaction was over, methanol (10 mL) and the dichloromethane (100 mL) were added in the reaction; after solids were dissolved, the solids were filtered, and filtrate was concentrated under reduced pressure. The residue was subjected to column chromatography (dichloromethane: methanol=10:

1) to obtain an impure product, and the impure product was separated by high performance liquid preparative chromatography (neutral, column: Waters Xbridge 150*25 mm 5 μ m; mobile phase: [water (10 mM amine bicarbonate)-acetonitrile]; %B (acetonitrile): 10% -35%, 8.2 min) to obtain Compound 4.


[0118] LCMS (ESI): $[M+H]^+ = 303.0$

[0119] $^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ 7.91 (d, $J=2.0$ Hz, 2H), 7.12 (d, $J=2.0$ Hz, 2H), 5.68 (s, 4H), 5.48 (br s, 2H), 1.52 (m, 12H).


[0120] Example 5: Synthesis of Compound 5



[0121] Synthetic Route of Compound 5:



[0122] Step 1: Synthesis of Compound 5-3




[0123] Substrates 5-1 (3 g, 17.74 mmol, 1 eq) and 5-2 (4.03 g, 17.91 mmol, 1.01 eq) were dissolved in acetic acid (50 mL) and stirred at 25 $^{\circ}\text{C}$ for 2 hours. After the reaction

was over, a reaction solution was spin-dried directly. Crude was dissolved in ethyl acetate (130 mL), and washed once with a saturated sodium carbonate aqueous solution (100 mL), sodium thiosulfate (100 mL, 1 M) and saturated saline (100 mL), respectively. The organic phases were dried over anhydrous sodium sulfate and spin-dried to obtain the crude. The crude was purified by a rapid silica gel column (petroleum ether: ethyl acetate=100: 0 to 98: 2) to obtain Compound 5-3.

[0124] $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.59-7.67 (m, 2H), 6.22 (br s, 2H), 3.90 (s, 3H).


[0125] Step 2: Synthesis of Compound 5-5



[0126] Compounds 5-3 (1 g, 3.39 mmol, 1 eq) and 5-4 (860.66 mg, 3.39 mmol, 1 eq) and potassium acetate (332.62 mg, 3.39 mmol, 1 eq) were dissolved in toluene (15 mL), and stirred under nitrogen atmosphere at 25 °C for 10 min; [1,1'-bis(diphenylphosphino) ferrocene] palladium (II) dichloromethane [$\text{Pd}(\text{dppf})\text{Cl}_2 \cdot \text{CH}_2\text{Cl}_2$] (2.77 g, 3.39 mmol, 1 eq) was added. The solution was reacted under stirring for 15 hours after the temperature was increased to 100 °C. After the reaction was over, a reaction solution was quenched by a saturated sodium carbonate aqueous solution (60 mL), and extracted with ethyl acetate (60 mL). Organic phases were washed with saturated saline (60 mL), dried over anhydrous sodium sulfate, and spin-dried to obtain crude. The crude was purified by a rapid silica gel column (petroleum ether: ethyl acetate=100: 0 to 98.8: 1.2) to obtain Compound 5-5.

[0127] LCMS: $[\text{MS}+\text{H}^+]=295.9$.


[0128] Step 3: Synthesis of Compound 5-6



[0129] Compounds 2-1 (1.5 g, 5.45 mmol, 1 eq) and 5-5 (2.09 g, 7.09 mmol, 1.3 eq), potassium phosphate (3.47 g, 16.36 mmol, 3 eq) and Pd(dppf)Cl₂.CH₂Cl₂ (1.34 g, 1.64 mmol, 0.3 eq) were dissolved in dimethoxyethane (60 mL), and stirred at 80 °C for 15 hours. After the reaction was over, a reaction solution was filtered directly, and filtrate was spin-dried to obtain crude. The crude was purified by a rapid silica gel column (petroleum ether: ethyl acetate=100: 0 to 60: 40) to obtain Compound 5-6.

[0130] ¹H NMR (400 MHz, CDCl₃) δ 8.46 (d, *J*=8.78 Hz, 1H), 8.12 (br s, 2H), 7.88 (d, *J*=8.78 Hz, 1H), 7.81 (dd, *J*=3.01, 9.03 Hz, 1H), 7.51 (dd, *J*=3.01, 9.03 Hz, 1H), 4.53 (q, *J*=7.03 Hz, 2H), 3.87-3.94 (s, 3H), 1.44 (t, *J*=7.15 Hz, 3H).


[0131] Step 4: Synthesis of Compound 5-7



[0132] The Compound 5-6 (890 mg, 2.45 mmol, 1 eq) was dissolved in ethyl acetate (100 mL), under nitrogen atmosphere, Pd/C (500 mg, 10% purity) was added, and a hydrogen balloon (15 psi) was replaced by gas for three times. The solution was stirred at 25 °C for 15 hours, and then stirred at 65 °C for 15 hours. After the reaction was over, a reaction solution was filtered by diatomaceous earth. Filtrate was spin-dried directly to obtain Compound 5-7.

[0133] ¹H NMR (400 MHz, CDCl₃) δ 8.54 (br s, 2H), 7.60-7.68 (m, 2H), 7.45 (dd, *J*=3.14, 9.66 Hz, 1H), 7.16 (d, *J*=8.78 Hz, 1H), 5.84 (br s, 2H), 4.42 (q, *J*=7.03 Hz, 2H), 3.90 (s, 3H), 1.45 (t, *J*=7.15 Hz, 3H).

[0134] Step 5: Synthesis of Compound 5-8




[0135] The Compound 5-7 (500 mg, 1.50 mmol, 1 eq) was dissolved in

tetrahydrofuran (50 mL) and cooled to 0 °C. Methyl magnesium bromide (3 M, 7.50 mL, 15 eq) was added dropwise in a 2-methyltetrahydrofuran solution at 0 °C. The solution was reacted under stirring at 0 °C for 1.5 hours. After the reaction was over, a reaction solution was quenched by adding a saturated amine chloride aqueous solution (100 mL), and then extracted with ethyl acetate (100 mL*2). After organic phases were combined, the organic phases were dried over anhydrous sodium sulfate and spin-dried to obtain crude. The crude was purified by HPLC separation (column model: Boston Uni C18 40*150 mm*5 µm; mobile phase: [water (10 mM amine bicarbonate solution)-acetonitrile]; %B (acetonitrile): 28%-58%, 10 min) to obtain Compound 5-8.

[0136] ¹H NMR (400 MHz, CDCl₃) δ 7.30 (d, J=8.28 Hz, 1H), 6.99-7.10 (m, 2H), 6.89 (dd, J=2.89, 10.16 Hz, 1H), 1.71(d, J=1.76 Hz, 12H).


[0137] Step 6: Synthesis of Compound 5





[0138] The Compound 5-8 (100 mg, 313.11 µmol, 1 eq) was dissolved in acetonitrile (5 mL), and then purified by separation with an HPLC machine (Column Model: Phenomenex Gemini-NX 150*30 mm*5 µm; mobile phase: [Water (0.04% amine aqueous solution+10 mM amine bicarbonate solution)-acetonitrile]; %B (acetonitrile): 39%-49%, 8 min) to obtain Compound 5.

[0139] ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.36 (d, J=8.53 Hz, 1H), 7.05-7.13 (m, 2H), 6.89 (s, 2H), 6.84 (dd, J=3.01, 10.54 Hz, 1H), 5.72 (s, 2H), 5.55 (s, 1H), 5.47 (s, 1H), 1.53 (d, J=4.02 Hz, 12H).


[0140] Example 6: Synthesis of Compound 6



[0141] Synthetic Route of Compound 6:**[0142] Step 1: Synthesis of Compound 6-2**

[0143] The Compound 2-1 (10 g, 36.36 mmol, 1 eq) and benzyl alcohol (7.86 g, 72.71 mmol, 7.56 mL, 2 eq) were dissolved in acetonitrile (100 mL), and cesium carbonate (23.69 g, 72.71 mmol, 2eq) was added at 20 °C; the mixed solution was stirred at 20 °C for 12 hours. After the reaction was over, a reaction solution was quenched in water (40 mL), and extracted with ethyl acetate (80 mL*3). After organic phases were combined, the organic phases were dried over anhydrous sodium sulfate, filtered, and spin-dried to obtain crude. The crude was purified by a rapid silica gel column (SiO₂, petroleum ether/ethyl acetate=5: 1 to 3: 1) to obtain Compound 6-2.


[0144] ¹H NMR (400MHz, CDCl₃): δ 8.39-8.33 (m, 1H), 7.48-7.38 (m, 5H), 6.99-6.93 (m, 1H), 5.52-5.48 (m, 2H), 4.57-4.48 (m, 2H), 1.48-1.41 (m, 3H).

[0145] Step 2: Synthesis of Compound 6-3

[0146] The Compound 6-2 (4.9 g, 16.21 mmol, 1 eq) was dissolved in ethanol (90 mL), under nitrogen atmosphere, palladium on carbon (Pd/C, 0.5 g, 16.21 mmol, 10% purity, 1 eq) was added, and a hydrogen balloon as replaced by gas for three times. The solution was stirred at 30 °C for 12 hours. After the reaction was over, a reaction solution was filtered, and filtrate was spin-dried to obtain crude. The crude was purified by a rapid silica gel column (SiO₂, dichloromethane: methanol=10: 0 to 10: 1) to obtain Compound 6-3.

[0147] ¹H NMR (400MHz, CDCl₃): δ 9.17 (br s, 1H), 7.10 (d, J=9.8 Hz, 1H), 6.73 (d, J=9.8 Hz, 1H), 5.41 (br s, 2H), 4.36 (q, J=7.0 Hz, 2H), 1.38 (t, J=7.2 Hz, 3H).


[0148] Step 3: Synthesis of Compound 6-5



[0149] The Compounds 6-4 (1 g, 5.34 mmol, 1 eq) and 6-3 (974.4 mg, 5.35 mmol, 1 eq) were dissolved in DMSO (20 mL), potassium phosphate (1.70 g, 8.00 mmol, 1.5 eq) was added, and the solution was stirred at 100 °C for 12 hours. After the reaction was over, a reaction solution was diluted in ethyl acetate (50 mL), and washed with water (30 mL*2) and saturated saline (30 mL) sequentially. Organic phases were dried over anhydrous sodium sulfate, filtered, and spin-dried to obtain crude. The crude was purified by a rapid silica gel column (SiO₂, petroleum ether/ethyl acetate=5: 1 to 2: 1) to obtain Compound 6-5.

[0150] ¹H NMR (400 MHz, CDCl₃) δ 7.64 (d, J=11.8 Hz, 1H), 7.21-7.13 (m, 1H), 7.02-6.97 (m, 1H), 6.56(d, J=6.8 Hz, 1H), 4.44-4.31 (m, 2H), 3.89 (s, 3H), 1.40 (t, J=7.2 Hz, 3H).

[0151] Step 4: Synthesis of Compound 6




[0152] The Compound 6-5 (200 mg, 572.55 μmol , 1 eq) was dissolved in tetrahydrofuran (40 mL) and cooled to 0 °C. Methyl magnesium bromide (3 M, 5 mL, 26.20 eq) was added dropwise in a 2-methyltetrahydrofuran solution at 0 °C. The solution was reacted under stirring at 0 °C for 1 hour, and then stirred at 15 °C for 15 hours. After the reaction was over, a reaction solution was quenched by adding a saturated amine chloride aqueous solution (50 mL), and then extracted in ethyl acetate (60 mL). The organic phases were dried over anhydrous sodium sulfate and spin-dried to obtain the crude. The crude was purified by HPLC separation (column model: Waters Xbridge 150*25 mm 5 μm ; mobile phase: [water (10 mM amine bicarbonate solution)-acetonitrile]; %B (acetonitrile): 15%-45%, 10 min) to obtain Compound 2.


[0153] LCMS: $[\text{MS}+\text{H}^+]=336.0$.

[0154] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.06 (d, $J=8.28$ Hz, 1H), 6.88 (d, $J=13.05$ Hz, 1H), 6.55 (d, $J=8.28$ Hz, 1H), 6.33(d, $J=7.53$ Hz, 1H), 5.37 (s, 1H), 5.30 (br d, $J=3.01$ Hz, 4H), 5.26 (s, 1H), 1.48 (s, 6H), 1.38 (s, 6H).


[0155] Example 7: Synthesis of Compound 7



[0156] Synthetic Route of Compound 7:




[0157] Step 1: Synthesis of Compound 7-2



[0158] The Compound 5-4 (3.65 g, 14.39 mmol, 2 eq), Pd(dppf)Cl₂ (526.35 mg, 719.34 μmol, 0.1 eq), and potassium acetate (1.41 g, 14.39 mmol, 2 eq) were added in a toluene (30 mL) solution containing the Compound 7-1 (2 g, 7.19 mmol, 1 eq), and the mixed solution was reacted at 110 °C for 3 hours. After the reaction was over, a reaction solution was concentrated under reduced pressure to obtain crude 7-2.


[0159] Step 2: Synthesis of Compound 7-3



[0160] The Compound 3-1 (2.3 g, 7.07 mmol, 1 eq), cesium carbonate (4.61 g, 14.14 mmol, 2 eq), Pd(dppf)Cl₂ (517.32 mg, 707.00 μmol, 0.1 eq) were added in a toluene (50 mL) and water (10 mL) solution containing the Compound 7-2 (1.63 g, 7.07 mmol, 1 eq), and the mixed solution was reacted at 110 °C under nitrogen atmosphere for 6 hours. After the reaction was over, a reaction solution was filtered with diatomaceous earth, and filtrate was concentrated under reduced pressure to obtain crude. The crude was separated by silica gel column chromatography (SiO₂, petroleum ether/ethyl acetate=5: 1 to 0: 1) to obtain Compound 7-3.

[0161] LCMS: 350.1 [M+1]⁺.

[0162] Step 3: Synthesis of Compound 7-4




[0163] Pd/C (1.7 g, 5% purity) was added in a methanol (50 mL) and ethyl acetate (50 mL) solution containing the Compound 7-3 (1.58 g, 4.52 mmol, 1 eq), and the mixed solution was reacted at 15 psi under hydrogen atmosphere at 20 °C for 3 hours. After the reaction was over, a reaction solution was filtered with diatomaceous earth, and filtrate was concentrated under reduced pressure to obtain Compound 7-4.

[0164] LCMS: 320 [M+1]⁺.

[0165] ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 7.96 (t, J= 1.76 Hz, 1H), 7.54 (d, J= 11.84 Hz, 1H), 7.39 (s, 1H), 6.95 (d, J= 6.58 Hz, 1H), 6.82 (s, 2H), 6.66 (s, 2H), 3.83 (d, J= 3.96 Hz, 6 H).

[0166] Step 4: Synthesis of Compound 7




[0167] Methyl magnesium bromide (3M, 31.32 mL, 20 eq) was added in a tetrahydrofuran (150 mL) solution containing the Compound 7-4 (1.5 g, 4.70 mmol, 1 eq) at 0 °C, and the mixed solution was reacted at 20 °C for 3 hours. After the reaction was over, a reaction solution was poured into a saturated amine chloride solution, 50 mL of ethyl acetate was added for liquid separation, and an aqueous phase was extracted with ethyl acetate (50 mL*3); organic phases were combined, dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure to obtain crude. The crude was separated by HPLC (column model: Xtimate C18 150*25 mm*5 μm; mobile phase: [water (10 mM amine bicarbonate solution)-acetonitrile]; %B (acetonitrile): 25%-45%, 10.5 min) to obtain Compound 7.

[0168] LCMS: 320.0[M+1]⁺;


[0169] ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 7.77 (s, 1H), 7.05 (s, 1H), 6.91 (d, J=12.28 Hz, 1H), 6.67 (d, J= 7.46 Hz, 1H), 5.63 (s,2H), 5.27-5.48 (m, 4H), 1.51 (s, 12H)

[0170] Example 8: Synthesis of Compound 8




8

[0171] Synthetic Route of Compound 8:



[0172] Step 1: Synthesis of Compound 8-1




[0173] The Compound 3-4 (1 g, 3.32 mmol, 1 eq) and NCS (509.67 mg, 3.82 mmol, 1.15 eq) were dissolved in glacial acetic acid (60 mL), and stirred at 25 °C for 15 hours. After the reaction was over, a reaction solution was adjusted to pH about 9 with a saturated sodium carbonate aqueous solution (200 mL), and extracted with ethyl acetate (150 mL). The organic phase was washed once with saturated saline (50 mL), then dried over anhydrous sodium sulfate, and spin-dried to obtain crude. The crude was purified by a rapid silica gel column (petroleum ether: ethyl acetate=100: 0 to 70: 30) to obtain Compound 8-1.

[0174] ¹H NMR (400 MHz, CDCl₃) δ 8.10 (d, J=1.76 Hz, 1H), 7.97 (s, 1H), 7.12 (d,

$J=1.76$ Hz, 1H), 6.63 (s, 1H), 5.83(br s, 4H), 4.01 (s, 3H), 3.91 (s, 3H).

[0175] Step 2: Synthesis of Compound 8






[0176] The Compound 8-1 (265 mg, 789.30 μ mol, 1 eq) was dissolved in tetrahydrofuran (50 mL) and cooled to 0 °C. Methyl magnesium bromide (3 M, 4 mL, 15.20 eq) was added dropwise in a 2-methyltetrahydrofuran solution at 0 °C. The solution was reacted under stirring at 0 °C for 2 hours, and then stirred at 25 °C for 15 hours. After the reaction was over, a reaction solution was quenched by adding a saturated amine chloride aqueous solution (50 mL), and then extracted with ethyl acetate (60 mL*2). After organic phases were combined, the organic phases were dried over anhydrous sodium sulfate and spin-dried to obtain crude. The crude was purified by HPLC separation (column model: watersXbridge 150*25 mm*5 μ m; mobile phase: [water (10 mM amine bicarbonate solution)-acetonitrile]; %B (acetonitrile): 25%-48%, 7.8 min) to obtain Compound 8.

[0177] LCMS: [MS+H⁺]=336.0;

[0178] ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.66 (d, $J=2.01$ Hz, 1H), 7.07 (s, 1H), 6.96 (d, $J=2.01$ Hz, 1H), 6.61 (s, 1H), 5.61 (br d, $J=17.32$ Hz, 4H), 5.47 (s, 1H), 5.34 (s, 1H), 1.51 (s, 12H).

[0179] **Example 9: Synthesis of Compound 9**




[0180] Synthetic Route of Compound 9:**[0181] Step 1: Synthesis of Compound 9-2**

[0182] A Compound 9-1 (24.8 g, 120.37 mmol, 1 eq) and benzylamine (15.48 g, 144.45 mmol, 15.75 mL, 1.2 eq) were dissolved in DMF (200 mL), and triethylamine (36.54 g, 361.12 mmol, 50.26 mL, 3 eq) was added at 20 °C. A reaction solution was stirred at 20 °C for 21 hours. After the reaction was complete, a reaction solution was quenched by adding water (250 mL), and then extracted with ethyl acetate (500 mL *2). Organic phases were spin-dried. Products were purified by a rapid silica gel column (SiO₂, petroleum ether/ethyl acetate=5: 1 to 3: 1) to obtain Compound 9-2.

[0183] LCMS (ESI): [M+H]⁺: 277.1.

[0184] ¹H NMR (400 MHz, CDCl₃) δ 8.71 (s, 1H), 8.55 (br s, 1H), 7.36-7.43 (m, 2H), 7.29-7.35 (m, 3H), 6.56 (s, 1H), 4.44 (d, J=5.52 Hz, 2H), 3.88-3.92 (m, 3H).


[0185] Step 2: Synthesis of Compound 9-3



[0186] The Compound 9-2 (5 g, 18.07 mmol, 1 eq) and sodium methoxide (9.76 g, 180.69 mmol, 10 eq) were dissolved in MeOH (50 mL), and stirred at 90 °C under nitrogen atmosphere for 15 hours. After the reaction was over, products were purified by a rapid silica gel column (SiO₂, petroleum ether/ethyl acetate=5: 1 to 3: 1) to obtain Compound 9-3.

[0187] LCMS (ESI): [M+H]⁺: 273.

[0188] Step 3: Synthesis of Compound 9-4




[0189] The Compound 9-3 (3.26 g, 11.97 mmol, 1 eq), trimethylchlorosilane (5.20 g, 47.89 mmol, 6.08 mL, 4 eq) and NaI (7.18 g, 47.89 mmol, 4 eq) were dissolved in acetonitrile (50 mL), and then reacted under stirring at 80 °C under nitrogen atmosphere for 15 hours. After the reaction was over, a reaction solution was adjusted to PH=7 with sodium bicarbonate, then quenched with water (20 mL), and extracted twice with dichloromethane (100 mL); finally organic phases were combined, and spin-dried in a rotary evaporator. Crude was purified with a rapid silica gel column (SiO₂, petroleum ether/ethyl acetate=5/1 to 3: 1) to obtain Compound 9-4.

[0190] LCMS (ESI): [M+H]⁺: 259.1

[0191] ¹H NMR (400 MHz, CDCl₃) δ 7.47 (br s, 1H), 6.47-6.58 (m, 5H), 4.65 (s, 1H), 3.55 (br d, J=5.27 Hz, 2H), 3.00-3.08 (m, 3H)


[0192] Step 4: Synthesis of Compound 9-6



[0193] The Compound 9-4 (2.29 g, 8.87 mmol, 1 eq), the Compound 9-5 (2.77 g, 10.64 mmol, 1.2 eq), cuprous iodide (337.73 mg, 1.77 mmol, 0.2 eq), potassium carbonate (2.45 g, 17.73 mmol, 2 eq) and (1S, 2S)-N1, N2-dimethylcyclohexane-1,2-diamine (252.24 mg, 1.77 mmol, 0.2 eq) were dissolved in toluene (30 mL), and reacted under stirring at 110 °C under nitrogen atmosphere for 13 hours. After the reaction was over, the solution was filtered and spin-dried in the rotary evaporator. The crude was purified by a rapid silica gel column (SiO₂, petroleum ether/ethyl acetate=1: 1 to 0: 1) to obtain Compound 9-6.

[0194] LCMS (ESI): [M+H]⁺: 438.1

[0195] Step 5: Synthesis of Compound 9-7




[0196] The Compound 9-6 (1 g, 2.29 mmol, 1 eq) was dissolved in tetrahydrofuran (30 mL), and palladium on carbon (1.5 g, 10% purity) was added in the mixed solution under nitrogen atmosphere. The mixed solution was stirred for 12 hours at 65 °C under hydrogen atmosphere. After the reaction was over, a reaction mixture was filtered and spin-dried. The crude was purified by a rapid silica gel column (SiO₂, petroleum

ether/ethyl acetate=1: 1 to 0: 1) to obtain Compound 9-7.

[0197] LCMS (ESI): $[M+H]^+$: 318.1

[0198] ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ 8.12 (s, 1H), 7.76 (d, $J=8.53$ Hz, 1H), 7.06 (br s, 2H), 6.84 (s, 2H), 6.77(d, $J=1.76$ Hz, 1H), 6.51 (dd, $J=2.01, 8.53$ Hz, 1H), 5.42 (s, 1H), 3.81 (s, 3H), 3.74 (s, 3H)

[0199] Step 6: Synthesis of Compound 9



[0200] The Compound 9-7 (200 mg, 630.33 μmol , 1 eq) was dissolved in tetrahydrofuran (20 mL); under nitrogen atmosphere, a reaction solution was cooled to -78 $^{\circ}\text{C}$ in a dry ice-ethanol bath, and lithium methyl (1.6 M, 5.91 mL, 15 eq) was slowly added dropwise. The reaction solution was slowly heated to 0 $^{\circ}\text{C}$ thereafter and stirred for 30 min. After the reaction was over, a reaction solution was quenched in water (10 mL) at 0 $^{\circ}\text{C}$, diluted with water (5 mL) and extracted with THF (40 mL). After organic phases were combined, the organic phases were dried over anhydrous sodium sulfate, filtered, and spin-dried to obtain Compound 9.

[0201] LCMS (ESI): $[M+H]^+$: 318.

[0202] ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ 6.20 (d, $J=8.28$ Hz, 1H), 6.14 (s, 1H), 5.69 (d, $J=2.26$ Hz, 1H), 5.53 (dd, $J=2.13, 8.16$ Hz, 1H), 5.45 (s, 2H), 4.73 (s, 2H), 4.57 (s, 1H), 4.53 (s, 1H), 4.41 (s, 1H), 0.67 (s, 6H), 0.60 (s, 6H).

[0203] Experimental Example 1: *In Vitro* Aldehyde Capture Capacity Test

[0204] 1. Experiment Objective and Process

[0205] Objective: Xerophthalmia is caused by inflammation in eyes, wherein the inflammation may produce aldehydes that will accelerate the inflammation symptoms and worsen the xerophthalmia if not eliminated in time. In the experiment, a better

compound was selected according to the complexing capacity of the drug and aldehydes *in vivo* by simulating the *in vivo* environment.

[0206] Procedures: Sulfobutyl-B-cyclodextrin (310 mg) was dissolved in a phosphate buffer (1.25 mL) to prepare a solution.

[0207] Nonanal (5.0 mg, 32 μ mol, 1.0 eq) and triolein (300 mg) were added to a reaction flask at room temperature, after addition of the prepared solution described above, linoleic acid (300 mg) was added, and finally a dimethyl sulfoxide (0.15 ml) solution containing the compound (32 μ mol, 1.0 eq) of the present disclosure was added; the reaction solution was reacted at 20-23 $^{\circ}$ C.

[0208] The reaction solution was reacted under stirring for 10 min, 100 min, 200 min, and 300 min respectively, and sampled for high performance liquid chromatography after the solution was stratified after standing for 2 min.

[0209] Sampling method: 25 μ l of an upper emulsified layer and 50 μ l of a lower aqueous phase were sampled by a pipette, and samples were diluted with 1 mL of methanol.

[0210] 2. Test Results

[0211] The ultraviolet absorption of the nonanal was weak at 254 nm, so the nonanal had little overall effect on the content of complexed products. Therefore, the percentage contents of complexes at 254 nm under high performance liquid chromatography were compared to observe aldehyde capture and complexing capacities, as shown in Figure 1 and Table 1:




Table 1 Statistics of Test Results of Aldehyde Capture Capacity of the Compound

Compound	Linear Equation	Slope	Regression Coefficient R ²	Area under the curve AUC (min. Conversion)
Compound 1	y=0.0157x+1.718	0.0157	R ² =0.7521	1257
Compound 2	y=0.0103x+2.7459	0.0103	R ² =0.9712	1271
Compound 3	y=0.0652x+3.362	0.0652	R ² =0.9891	3979

Compound 4	$y=0.0589x+1.3064$	0.0589	$R^2=0.9964$	3061
Compound 5	$y=0.0271x+0.1945$	0.0271	$R^2=0.9326$	1208
Compound 6	$y=0.1083x+12.77$	0.1083	$R^2=0.9027$	8953
Compound 7	$y=0.1002x+7.7455$	0.1002	$R^2=0.9686$	6944
Compound 8	$y=0.0519x+4.9828$	0.0519	$R^2=0.9535$	3879
Compound 9	$y=0.1241x+14.964$	0.1241	$R^2=0.9299$	10284

[0212] The HPLC method is as follows: 2.5 μ m, 3.0*100mm 5-95CD_XBEH_12min_0.8.1cm or XBRIGE 2.5 μ m, 3.0*100mm 5-80CD_XBEH_12min_0.8.1cm

[0213] Specific conditions: XBRIGE 2.5 μ m, 3.0*100mm 5-80CD_XBEH_12min_0.8.1cm

Column	XBridge BEHC18 3.0*100mm, 2.5 μ m		
Detection Wavelength	220, 254nm		
Column Temperature	40 °C		
Flow Rate	0.8mL/min		
Injection Volume	1 μ L		
Mobile Phase	A: 0.02% aqueous amine solution		
	B: Acetonitrile		
Gradient elution	Time (min)	A%	B%
	0.01	95	5
	6.00	40	60
	9.00	20	80
	12.00	20	80

[0214] XBRIGE 2.5 μ m, 3.0*100mm 5-95CD_XBEH_12min_0.8.1cm

Column	XBridge BEHC18 3.0*100mm, 2.5 μ m
Detection Wavelength	220, 254nm
Column Temperature	40 °C
Flow Rate	0.8mL/min
Injection Volume	1 μ L

Mobile Phase	A: 0.02% aqueous amine solution		
	B: Acetonitrile		
Gradient elution	Time (min)	A%	B%
	0.01	95	5
	6.00	15	85
	9.00	5	95
	12.00	5	95

[0215] Taking The compound 7 as an example, after 300 min of reaction, the analytical method for HPLC was XBRIGE 2.5 μ m, 3.0*100mm 5-80CD_XBEH_12min_0.8.1cm;

[0216] Retention time 6.689 min, 6.787 min, 6.966 min, and 7.102 min were absorption peaks of products complexed with monomolecular aldehyde;

[0217] Retention time 8.905 min, 9.010 min, and 9.075 min were absorption peaks of products complexed with bimolecular aldehyde.

[0218] The percentage content of the specific complexed product at 254 nm was calculated as the sum of the percentage contents of the absorption peaks for the retention time described above, i.e. (1.839+1.715+14.993+13.029)%+(1.004+2.212+1.247)%= 36.039%

[0219] Specific HPLC data of the percentage contents of complexed products of the compounds of the present disclosure are shown in Table 2 below:

Time (min) \ Compound	10	100	200	300
1	0.917	4.909	4.383	6.224
4	1.475	7.663	13.353	18.636
2	2.658	4.097	4.721	5.796
3	3.265	10.569	17.191	22.182
5	0.882	2.935	4.328	9.139
7	6.634	20.013	29.424	36.039
6	9.738	27.850	37.847	41.686

8	4.117	12.121	15.49	19.873
9	12.268	31.427	43.086	48.761

[0220] Conclusion: Experiment results showed that the compounds of the present disclosure had far significant aldehydes complexing capability and speed.

[0221] Experimental Example 2: *In Vitro* Evaluation

[0222] Objective: To study the inhibitory effect of compounds on human liver microsomal cytochrome P450.

[0223] Experimental procedures: In the study on inhibition of the compounds on the human liver microsomal cytochrome P450 (CYP1A2, CYP2C9, CYP2C19, CYP2D6 and CYP3A4), fused human liver microsomes were selected as a CYP450 enzyme source. The compounds of different concentrations (10, 5, 1.5, 0.5, 0.15, 0.05, 0.015 mM) were incubated with probe substrates (where CYP3A4 employs two substrates) and cofactors (NADPH) of 5 CYP enzymes to determine the IC₅₀ value of each compound for inhibition of each CYP enzyme. Test results are as follows:

Table 3 Inhibition Test Results of Compounds of the Present Disclosure on Human Liver Microsomal Cytochrome P450 Isoenzymes

Compound No.	CYP1A2/CYP2C9/CYP2C19/CYP2D6/CYP3A4(μM)
Compound 3	>50/>50/>50/>50/>50
Compound 7	>50/>50/>50/>50/>50
Compound 8	>50/>50/>50/>50/>50

[0224] Conclusion: The present disclosure has high safety and low possibility of drug interaction.

[0225] Experimental Example 3: Pharmacokinetic Evaluation of Compound

[0226] I.

[0227] Objective: To study 5-day toxicology of compounds in SD rats

[0228] Experimental materials: SD rats (male, 200-300 g, 7-9 weeks old, Shanghai Lingchang)

[0229] Experimental procedures:

[0230] Pharmacokinetic profiles of the compound were tested by intravenous injection in rodents according to the standard protocol. In the study, the candidate compound was prepared as a clear solution, and was administered to SD rats at 20 mg/kg by single intravenous injection for 5 consecutive days. A vehicle for intravenous injection was a 10% hydroxypropyl β -cyclodextrin aqueous solution. Whole blood samples were collected at 0.033, 0.083, 0.25, 0.5, 1, 2, 4, 8, and 24 hours postdose on Day 1 and Day 5 of dosing, centrifuged at 3000 r for 10 min, and separated from supernatant to obtain plasma samples. 15 μ L of plasma samples were mixed with 300 μ L of acetonitrile precipitated proteins containing an internal standard, and 2 μ L of the supernatant was injected by centrifugation. Plasma concentrations were quantified by LC-MS/MS analysis, and pharmacokinetic parameters, such as clearance, half-life, and area under the concentration-time curve, were calculated.

[0231] Experiment Results:

Table 4 Pharmacokinetic Test Results

Test Sample	Clearance (mL/min/kg)		Half-time T _{1/2} (h)		Area under the concentration-time curve AUC(nM.hr)	
	Day 1	Day 5	Day 1	Day 5	Day 1	Day 5
Compound 3	7.11	9.04	2.45	2.72	155784	123277

[0232] Conclusion: The compound of the present disclosure has no risk of drug accumulation after injection for 5 consecutive days.

[0233] II.

[0234] Objective: To study pharmacokinetics of the compound in Beagle dogs

[0235] Experimental material: Beagle dogs (male, 8-11 kg, greater than or equal to 6 months, Marshall)

[0236] Experimental procedures:

[0237] Pharmacokinetic profiles of the compound were tested by intravenous administration in rodents according to the standard protocol. In the study, the candidate compound was prepared as a clear solution, and was administered to Beagle dogs at 1

mg/kg by single intravenous injection. A vehicle for intravenous injection was a 10% hydroxypropyl β -cyclodextrin aqueous solution. Whole blood samples were collected at 0.033, 0.083, 0.25, 0.5, 1, 2, 4, 8, 12, and 24 hours, respectively, centrifuged at 3000 r for 10 min, and separated from supernatant to obtain plasma samples. 20 μ L of plasma samples were mixed with 400 μ L of acetonitrile precipitated proteins containing an internal standard, and 2 μ L of the supernatant was injected by centrifugation. Plasma concentrations were quantified by LC-MS/MS analysis, and pharmacokinetic parameters, such as clearance, half-life, and area under the concentration-time curve, etc. were calculated.

[0238] Experiment Results:

Table 5 Pharmacokinetic Test Results

Test Sample	Clearance (mL/min/kg)	Half-time $T_{1/2}$ (h)	Area under the concentration-time curve AUC(nM.hr)
Compound 3	15	2.17	3709
Compound 7	11.3	2.94	4672

[0239] Conclusion: The present disclosure has high clearance, moderate half-life and good pharmacokinetic profile.

[0240] III.

[0241] Objective: To study pharmacokinetics of the compound in cynomolgus monkeys

[0242] Experimental material: Cynomolgus monkeys (male, 2.5-4 kg, greater than or equal to 2 years, Jingang Biotech)

[0243] Experimental procedures:

[0244] Pharmacokinetic profiles of the compound were tested by intravenous administration in monkeys according to the standard protocol. In the study, the candidate compound was prepared as a clear solution, and was administered to cynomolgus monkeys at 1 mg/kg by single intravenous injection. A vehicle for

intravenous injection was a 10% hydroxypropyl β -cyclodextrin aqueous solution. Whole blood samples were collected at 0.033, 0.083, 0.25, 0.5, 1, 2, 4, 8, 12, and 24 hours, respectively, centrifuged at 3000 r for 10 min, and separated from supernatant to obtain plasma samples. 20 μ L of plasma samples were mixed with 400 μ L of acetonitrile precipitated proteins containing an internal standard, and 2 μ L of the supernatant was injected by centrifugation. Plasma concentrations were quantified by LC-MS/MS analysis, and pharmacokinetic parameters, such as clearance, half-life, and area under the concentration-time curve, etc. were calculated.

[0245] Experiment Results:

Table 6 Pharmacokinetic Test Results

Test Sample	Clearance (mL/min/kg)	Half-time $T_{1/2}$ (h)	Area under the concentration-time curve AUC(nM.hr)
Compound 3	6.2	3.81	9164
Compound 7	4.6	3.91	13016

[0246] Conclusion: The present disclosure has high clearance, moderate half-life and good pharmacokinetic profile.

[0247] Experimental Example 4: *In Vivo* Pharmacokinetic Study

[0248] Objective: To study the ratio of corneal to plasma drug concentration after eye drops in rats

[0249] Experimental materials: Male SD rats, 200-300 g, 7-9 weeks old, Shanghai Lingchang

[0250] Experimental procedures: The candidate compound was prepared as a clear solution, and administered to SD rats by eye drops. The vehicle for eye drops was 10% hydroxypropyl β -cyclodextrin aqueous solution, and the drug concentration in the eye drops was 5 mg/mL. Corneal and whole blood samples were collected at 1 and 4 hours, respectively. The corneal samples was homogenized in 15 mM phosphate buffer (PBS): MeOH (2: 1, v: v) buffer, the whole blood samples were centrifuged at 3000 r for 10 min, and the supernatant plasma samples were separated. 20 μ L of the plasma samples and homogenate samples were mixed with 400 μ L of acetonitrile precipitated

proteins containing an internal standard, respectively, and 2 μL of the supernatant was injected by centrifugation. The plasma concentrations were quantified by LC-MS/MS analysis; the drug concentrations in the cornea and plasma were measured respectively according to different time points, and the ratio of corneal/plasma concentrations was calculated.

[0251] Experiment Results:

Table 7 Pharmacokinetic Test Results

Test Sample	Time	Corneal Concentration (nM)	Plasma Concentration (nM)	Corneal Concentration/Plasma Concentration C/P ratio
Compound 3	1h	1796	162	14.3
	4h	581	43.8	16.6
Compound 7	1h	2447	172	11.8
	4h	330	24.4	12.4
Compound 8	1h	1663	148	12.0
	4h	167	36.4	3.72

[0252] Conclusion: As shown in the DMPK test, test compounds 3, 7 and 8 could enter the cornea by eye drops to show the efficacy. The ratio of cornea/plasma concentrations of the Compound 3 was relatively high (above 10). The drug took effect in the cornea, so the targeting property was good.

[0253] Experimental Example 5: *In Vivo* Efficacy Evaluation

[0254] Objective:

[0255] Subcutaneous injection of scopolamine may induce the xerophthalmia in mice. Decreased tear secretion and inflammatory infiltration can be observed by tear tests and corneal fluorescence staining scores. It can predict whether a model can achieve the expected severity at the early stage of modeling.

[0256] Study Design:

[0257] Twenty animals were selected from 25 female C57BL/6J mice and randomized

into 4 groups by body weight via Provantis or Excel, 4 animals per group, and group assignment may also refer to the results of the tear test and the corneal fluorescence staining score of each animal prior to the experiment.

[0258] The mice were injected subcutaneously with scopolamine hydrobromide (3 ± 0.5 h) for 4 times daily on Day 1 to Day 12 of the study to establish xerophthalmia models in the mice at 0.1 mL/animal/time.

[0259] On Day 1 to Day 13 of the study, animals were dosed by bilateral eye drops for 4 times daily (3 ± 0.5 h) at 3 μ L/eye (on Day 13, animals in Group 1 were not dosed). Each subcutaneous injection of the scopolamine hydrobromide was given predose the eye drops (except examinations on Day 7 and Day 12).

[0260] Tear tests and corneal fluorescence staining scores were performed on all experimental animals on Day 7 and Day 12 prior to modeling.

[0261] Criteria for corneal fluorescence staining scores: The cornea of an animal was divided into 5 regions, i.e., superior, inferior, nasal, temporal and central regions, each with 0-3 scores. Each region was scored, and the monocular score was the sum of scores of the 5 regions. 0: no staining; 1: slight staining, punctate staining with less than 5 spots; 2: moderate staining, punctate staining but no plaque staining; 3: severe staining, marked fluorescent plaques.

[0262] The tear test was performed 30 min after the second dosing on Day 7 and Day 12 for all experimental animals, and the corneal fluorescence staining was scored for all experimental animals 30 min after the third dosing on Day 7 and Day 12.

Table 8 Efficacy Experiment Protocol of Scopolamine by Subcutaneous Injection in Mice

Group	Number of animals (Female)	Modeling	Therapy	Method of Administration
			Both eyes	
Vehicles	4	Subcutaneous injection of scopolamine for 4 injections/day for 12 days (5 mg/mL, 0.1	Vehicles	Eye drops 3 μ L/eye 4 injections/day (3 ± 0.5 h)
Compound 3	4		0.5%	
Compound 7	4		0.5%	
Compound 8	4		0.3%	


		mL/injection, 3±0.5 h)	
--	--	------------------------	--

[0263] Test results: See Figure 2 and Figure 3 (Note:* in the attached drawings indicates $P \leq 0.5$, ** indicates $P \leq 0.01$, and *** indicates $P \leq 0.001$).

[0264] Conclusion: Under the conditions of the experiment, it can be seen that all tested compounds can improve tear secretion and corneal inflammation, showing the efficacy of improving the xerophthalmia.

What is claimed is:

1. A compound represented by formula (II) or a pharmaceutically acceptable salt thereof,



wherein


 is selected from a single bond and a double bond;

T₁, T₂, T₃, and T₄ are each independently selected from N, C, and CR₁;

T₅ is selected from C, CR₅, and C=O;

T₆ is selected from C, CR₆, and N;

T₇ is selected from N and CR₇;

when T₅ is selected from C=O and T₆ is selected from N, then  is selected from a single bond;

L is selected from a single bond, -O-, -S-, -NR₂- and -(CR₃R₄)_n-;

each R₁ is independently selected from H, F, Cl, Br, I, OH, and NH₂;

R₂ is selected from H and C₁₋₃ alkyl optionally substituted with 1, 2 or 3 R_a;

R₃ and R₄ are each independently selected from H, F, Cl, Br, I, OH, NH₂, CN and C₁₋₃ alkyl optionally substituted with 1, 2 or 3 R_b;


R₅, R₆ and R₇ are each independently selected from H, F, Cl, Br and I;

n is selected from 1, 2, and 3;

R_a and R_b are each independently selected from H, F, Cl, Br, I, OH, NH₂, CN, and CH₃.

2. The compound or the pharmaceutically acceptable salt thereof as defined in claim 1,

wherein the compound is selected from




wherein

T₃ and T₄ are each independently selected from N and CR₁;

R₁ and L are as defined in claim 1.

3. The compound or the pharmaceutically acceptable salt thereof as defined in claim 2, wherein the compound is selected from



wherein

R₁ and L are as defined in claim 1.

4. The compound or the pharmaceutically acceptable salt thereof as defined in any one of claims 1 to 3, wherein R₂ is selected from H, CH₃ and CH₂CH₃; wherein the CH₃

and the CH_2CH_3 are optionally substituted with 1, 2, or 3 R_a .


5. The compound or the pharmaceutically acceptable salt thereof as defined in claim 4, wherein R_2 is selected from H, CH_3 , and CH_2CH_3 .

6. The compound or the pharmaceutically acceptable salt thereof as defined in any one of claims 1 to 3, wherein R_3 and R_4 are each independently selected from H, F, Cl, Br, I, OH, NH_2 , CN, CH_3 and CH_2CH_3 , wherein the CH_3 and the CH_2CH_3 are optionally substituted with 1, 2 or 3 R_b .

7. The compound or the pharmaceutically acceptable salt thereof as defined in claim 6, wherein R_3 and R_4 are each independently selected from H, F, Cl, Br, I, OH, NH_2 , CN, CH_3 and CH_2CH_3 .

8. The compound or the pharmaceutically acceptable salt thereof as defined in any one of claims 1 to 3, wherein L is selected from a single bond, -O-, -S-, -NH-, $-(\text{CH}_2)_2-$ and $-\text{CH}_2-$.

9. A compound represented by the following formula or a pharmaceutically acceptable salt thereof, wherein the compound is selected from:



10. A pharmaceutical composition, comprising the compound or the pharmaceutically acceptable salt thereof as defined in any one of claims 1 to 9, and a pharmaceutically

acceptable carrier thereof.

11. A use of the compound or the pharmaceutically acceptable salt thereof as defined in any one of claims 1 to 9 or the composition as defined in claim 10 in the manufacture of a medicament for treating xerophthalmia.




Figure 1




Figure 2




Figure 3

