



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2018/10/24
(87) Date publication PCT/PCT Publication Date: 2019/05/02
(85) Entrée phase nationale/National Entry: 2020/04/23
(86) N° demande PCT/PCT Application No.: EP 2018/079184
(87) N° publication PCT/PCT Publication No.: 2019/081590
(30) Priorité/Priority: 2017/10/25 (DE10 2017 125 019.0)

(51) Cl.Int./Int.Cl. *C12Q 1/6881* (2018.01)
(71) Demandeur/Applicant:
EPIONTIS GMBH, DE
(72) Inventeurs/Inventors:
OLEK, SVEN, DE;
BARON, UDO, DE
(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : UTILISATION DE MVD EN TANT QUE MARQUEUR EPIGENETIQUE POUR L'IDENTIFICATION DE CELLULES IMMUNITAIRES, EN PARTICULIER LES CELLULES NK CD56+
(54) Title: PD1+ AS EPIGENETIC MARKER FOR THE IDENTIFICATION OF IMMUNE CELLS, IN PARTICULAR PD1+ CELLS

(57) **Abrégé/Abstract:**

The present invention relates to a method, in particular an in vitro method, for identifying PD1+ cells, comprising analyzing the methylation status of at least one CpG position in the mammalian gene region for Programmed cell death 1 (PDCD1), wherein a demethylation or lack of methylation of said gene region is indicative for a PD1+ cell, when compared to a non-PD1+ cell. The analyses according to the invention can identify PD1+ cells on an epigenetic level and distinguish them from all other cells in complex samples, such as, for example, other blood or immune cells. The present invention furthermore provides an improved method for quantifying PD1+ cells, in particular in complex samples. The method can be performed without a step of purifying and/or enriching cells, preferably in whole blood and/or non-trypsinized tissue.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

02 May 2019 (02.05.2019)



(10) International Publication Number

WO 2019/081590 A1

(51) International Patent Classification:

C12Q 1/6881 (2018.01)

(21) International Application Number:

PCT/EP2018/079184

(22) International Filing Date:

24 October 2018 (24.10.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

10 2017 125 019.0

25 October 2017 (25.10.2017) DE

(71) Applicant: **EPIONTIS GMBH** [DE/DE]; Rudower Chaussee 29, 12489 Berlin (DE).(72) Inventors: **OLEK, Sven**; Boothstrasse 16, 12207 Berlin (DE). **BARON, Udo**; Boxhagener Strasse 88, 10245 Berlin (DE).(74) Agent: **KRAUSS, Jan**; Boehmert & Boehmert, Pettenkoferstr. 20-22, 80336 München (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: PDCD1 AS EPIGENETIC MARKER FOR THE IDENTIFICATION OF IMMUNE CELLS, IN PARTICULAR PD1+ CELLS

(57) Abstract: The present invention relates to a method, in particular an in vitro method, for identifying PD1+ cells, comprising analyzing the methylation status of at least one CpG position in the mammalian gene region for Programmed cell death 1 (PDCD1), wherein a demethylation or lack of methylation of said gene region is indicative for a PD1+ cell, when compared to a non-PD1+ cell. The analyses according to the invention can identify PD1+ cells on an epigenetic level and distinguish them from all other cells in complex samples, such as, for example, other blood or immune cells. The present invention furthermore provides an improved method for quantifying PD1+ cells, in particular in complex samples. The method can be performed without a step of purifying and/or enriching cells, preferably in whole blood and/or non-trypsinized tissue.



WO 2019/081590 A1

PDCD1 as epigenetic marker for the identification of immune cells, in particular PD1+ cells

The present invention relates to a method, in particular an *in vitro* method, for identifying PD1+ cells, comprising analyzing epigenetic modifications/ properties of (including the methylation status) of at least one CpG position in the mammalian gene region for Programmed cell death 1 (PDCD1), wherein a demethylation or lack of methylation of said gene region is indicative for a PD1+ cell, when compared to a non-PD1+ cell. The analyses according to the invention can identify PD1+ cells on an epigenetic level and distinguish them from all other cells in complex samples, such as, for example, other blood or immune cells. The present invention furthermore provides an improved method for quantifying PD1+ cells, in particular in complex samples. The method can be performed without a step of purifying and/or enriching cells, preferably in whole blood and/or non-trypsinized tissue.

Furthermore, the present invention relates to a kit for performing the above methods as well as respective uses thereof. It is one aim of this invention to provide a novel, more robust means to quantitatively detect and measure PD1+ cells of the blood within any solid organs, tissue or body fluid of a mammal.

Background of the invention

Commonly, PD1+ cells are defined as cells that actively synthesize the protein encoded by the gene programmed cell death 1. In the current application, PD1+ cells are defined as cells that are rendered capable of expressing PD1 by providing an accessible unmodified primary DNA sequence as shown by the absence of modifications in CpG motifs in the intronic region described and defined in following.

PD1+ cells are cells expressing Programmed cell death protein-1 (PDCD1, also referred to as PD-1). PDCD1 is expressed on a variety of immune cell types, such as activated Thymus-derived lymphocytes (T lymphocytes, T cells), pro-B cells, myeloid-derived dendritic cells, and natural killer (NK) cells. PDCD1 is expressed during a number of different stages of

immune development and inflammation, and PDCD1 serves as an important checkpoint receptor involved in immunity regulation and self-tolerance. PDCD1 expression slows the immune response during initial acute antigen recognition by reducing tissue residency and cytokine production, as well as by decreasing the formation of helper cells during the early immune response. Interestingly, PDCD1 promotes apoptosis (programmed cell death) in antigen specific T-cells while simultaneously prohibiting apoptosis in regulatory T cells, which are anti-inflammatory T cells. Thus, PDCD1 is an important regulator of an effective adaptive immune response.

Even though almost all cells in an individual contain the exact same complement of DNA code, higher organisms must impose and maintain different patterns of gene expression in the various types of tissue. Most gene regulation is transitory, depending on the current state of the cell and changes in external stimuli. Persistent regulation, on the other hand, is a primary role of epigenetics - heritable regulatory patterns that do not alter the basic genetic coding of the DNA. DNA methylation is the archetypical form of epigenetic regulation; it serves as the stable memory for cells and performs a crucial role in maintaining the long-term identity of various cell types. Recently, other forms of epigenetic regulation were discovered. In addition to the “fifth base” 5-methylcytosine (mC), a sixth (5-hydroxymethylcytosine, hmC), seventh (5-formylcytosine, fC) and eighth (5-carboxycytosine, cC) can be found (Michael J. Booth et al. Quantitative Sequencing of 5-Methylcytosine and 5-Hydroxymethylcytosine at Single-Base Resolution *Science* 18 May 2012, Vol. 336 no. 6083 pp. 934-937).

The primary target of mentioned DNA modifications is the two-nucleotide sequence Cytosine-Guanine (a 'CpG site'); within this context cytosine (C) can undergo a simple chemical modification to become formylated, methylated, hydroxymethylated, or carboxylated. In the human genome, the CG sequence is much rarer than expected, except in certain relatively dense clusters called 'CpG islands'. CpG islands are frequently associated with gene promoters, and it has been estimated that more than half of the human genes have CpG islands (Antequera and Bird, *Proc Natl Acad Sci USA* 90: 11995-9, 1993).

Aberrant methylation of DNA is frequently associated with the transformation from healthy to cancerous cells. Among the observed effects are genome-wide hypomethylation, increased methylation of tumor suppressor genes, and hypomethylation of many oncogenes (reviewed, for example, by Jones and Laird, *Nature Genetics* 21:163-167, 1999; Esteller, *Oncogene*

21:5427-5440, 2002; and Laird, Nature Reviews/Cancer 3:253-266, 2003). Methylation profiles have been recognized to be tumor specific (i.e., changes in the methylation pattern of particular genes or even individual CpGs are diagnostic of particular tumor types), and there is now an extensive collection of diagnostic markers for bladder, breast, colon, esophagus, stomach, liver, lung, and prostate cancers (summarized, for example, by Laird, Nature Reviews/Cancer 3:253-266, 2003).

For one of the recently described modification of cytosine, 5-hydroxymethylation, the utility of oxidative bisulfite sequencing to map and quantify 5hmC at CpG islands was shown (Michael J. Booth et al. Quantitative Sequencing of 5-Methylcytosine and 5-Hydroxymethylcytosine at Single-Base Resolution Science 18 May 2012, Vol. 336 no. 6083 pp. 934-937). High levels of 5hmC were found in CpG islands associated with transcriptional regulators and in long interspersed nuclear elements. It is suggested that these regions might undergo epigenetic reprogramming in embryonic stem cells.

WO 2012/162660 describes methods using DNA methylation arrays are provided for identifying a cell or mixture of cells and for quantification of alterations in distribution of cells in blood or in tissues, and for diagnosing, prognosing and treating disease conditions, particularly cancer. The methods use fresh and archival samples.

Youngblood et al. (in: Youngblood et al. Chronic virus infection enforces demethylation of the locus that encodes PD-1 in antigen-specific CD8⁺ T cells. 2011 Sep 23;35(3):400-12) disclosed PDCD1 expression to be dependent on methylation of a CpG rich region upstream from the transcription start. This region overlaps with previously identified conserved regions C and B (CR-C & CR-B) of the PDCD1 gene. Methylation of this PDCD1 CpG rich locus inversely correlates with PDCD1 mRNA expression, and is thus involved in regulating immune responses. For example, during differentiation of naïve to effector CD8 T cells in response to an acute infection, the PDCD1 locus is hypomethylated. This event then triggers higher expression of PDCD1 mRNA. When effector CD8 T cells further differentiate into functional memory cells, the PDCD1 locus is being remethylated. Thus, methylation of PDCD1 provides a way of regulating immune responses via PDCD1 expression.

Bally et al. (in: Bally et al. NF-κB regulates PD-1 expression in macrophages. 2015 May 1;194(9):4545-54.) further investigated the methylation state of PDCD1 upstream regions

CR-C and CR-B in macrophages. They discovered no changes of methylation of PDCD1 after LPS-stimulation of Bone Marrow-Derived Macrophages (BMDMs).

Goltz et al. (in: Goltz et al. Promoter methylation of the immune checkpoint receptor PD-1 (PDCD1) is an independent prognostic biomarker for biochemical recurrence-free survival in prostate cancer patients following radical prostatectomy. *Oncoimmunology*. 2016 Sep 2;5(10):e1221555) further demonstrated the methylation state of the PDCD1 upstream locus to be correlated with carcinomas versus normal prostatic epithelium.

In view of the above, it is an object of the present invention to provide an improved and in particular robust method based on DNA-methylation analysis as a superior tool in order to more conveniently and reliably detect, identify, discriminate, and quantify PD1+ cells.

The present invention solves the above object by providing a method for identifying PD1+ cells in a sample, comprising analyzing the methylation status (bisulfite convertibility) of at least one CpG position in the mammalian (e.g. human) gene region for Programmed cell death 1 (PDCD1), wherein preferably said gene region as analyzed is positioned based on/according to SEQ ID No. 1, wherein a demethylation of said gene region is indicative for a PD1+ cell, when compared to a non-PD1+ cell.

The Programmed cell death protein-1 (PDCD1, also referred to as PD-1) belongs in the immunoglobulin superfamily and is a 288 amino acid long cell surface receptor expressed on a variety of immune cell types. Importantly, the formation of a complex between PDCD1 and its ligand Programmed death ligand 1 (PD-L1) or Programmed death ligand 2 (PD-L2) transmits an inhibitory signal that reduces the proliferation and inflammatory activity of cells expressing PDCD1, and thereby suppressing the immune response. Thus, expression of PDCD1 enables the regulated activation and expansion of immune cells, which is necessary for an effective adaptive immune response. The gene for human PDCD1 is found on chromosome 2, 241849881-241858908 reverse strand; Ensembl-ID: ENSG00000188389.

In the context of the present invention, the gene region shall comprise all of the genomic region relating to and encoding for PDCD1. Thus, included are enhancer regions, promoter region(s), introns, exons, and non-coding regions (5'- and/or 3'-regions) that belong to PDCD1. Preferred is thus a method according to the present invention, wherein the at least

one CpG position is present in the 5' region upstream from the transcription start, promoter region, the 5' or 3' untranslated regions, exon, intron, exon/intron border and/or in the 3' region downstream of the transcriptional stop of the gene as analyzed.

The present invention is further based on the surprising identification of a region of the PDCD1 gene by the inventors, as specific epigenetic marker, allowing the identification of PD1+ cells as well as the clinical routine application of said analysis.

In the context of the present invention, the genomic region of PDCD1, in particular according to SEQ ID No. 1, more preferably SEQ ID NOs. 2 (Amp 1876), 3 (Amp 1877) or 4 (Amp 1878) allow the identification of PD1+ cells. Surprisingly, the discriminatory pattern of bisulfite convertible and non-convertible cytosine is particularly and even exclusively limited to the genomic region according to SEQ ID No. 1 for PD1+ cells as shown using the amplicon according to SEQ ID No. 1, and in particular in the bisulfite converted sequences according to SEQ ID No. 12 and/or 13 (TpG converted and CpG converted sequences for AMP 1877).

The inventors could demonstrate that in the PD1+ cells the CpG motifs as disclosed are almost completely demethylated (i.e. to more than 70%, preferably 80%, preferably, more than 90% and most preferred more than 95%), whereas the same motifs are completely methylated in PD1- cells.

The differential methylation of the CpG motifs within the aforementioned regions is a valuable tool to identify PD1+ cells, such as will be required/or at least of some value for identifying and quantifying said cells in autoimmune diseases, transplant rejections, cancer, allergy, primary and secondary immunodeficiencies, such as, for example, HIV infections and AIDS, Graft versus Host (GvH), hematologic malignancies, rheumatoid arthritis, multiple sclerosis, or a cytotoxic T cell related immune status in any envisionable diagnostic context. The assay allows measurement of PD1+ cells without purification or any staining procedures.

Another preferred aspect of the method according to the present invention then further comprises a quantification of the relative amount of PD1+ cells based on comparing relative amounts of said methylation frequency in the region as analyzed with relative amounts of the methylation frequency in a control gene, such as, for example, GAPDH. Said quantification is thus achieved based on the ratio of the bisulfite convertible DNA to non-convertible DNA in

the genetic region of PDCD1 (e.g. of SEQ ID No. 1) as described and analyzed herein. Most preferred is a quantification of the relative amount of PD1+ cells is based on an (preferably parallel or simultaneous) analysis of the relative amount of bisulfite convertible DNA of cell-specific region for PDCD1, and of the relative amount of bisulfite convertible DNA of cell-unspecific genes (preferably designated “control genes” or “control regions”, such as, for example, the gene for GAPDH).

In a further preferred embodiment of the method according to the present invention, said analysis of bisulfite convertibility comprises amplification with at least one primer of suitable primer pairs that can be suitably designed based on SEQ ID No. 1, preferably oligomers according to any of SEQ ID No. 6 to 11.

In contrast to FACS and mRNA measurements, using the methods according to the present invention, the measurement(s) and analyses can be done independent of purification, storage - and to quite some extent - also to tissue quality.

Preferably, the amplification involves a polymerase enzyme, a PCR or chemical amplification reaction, or other amplification methods as known to the person of skill as described below, e.g. in the context of MSP, HeavyMethyl, Scorpion, MS-SNUPE, MethylLight, bisulfite sequencing, methyl specific restriction assays and/or digital PCR (see, for example Kristensen and Hansen PCR-Based Methods for Detecting Single-Locus DNA Methylation Biomarkers in Cancer Diagnostics, Prognostics, and Response to Treatment *Clinical Chemistry* 55:8 1471–1483 (2009)).

With the amplification, an amplicon of the PDCD1 gene region is produced that is a particularly preferred “tool” for performing the method(s) according to the present invention. Consequently, oligomers according to any of SEQ ID No. 6 to 11 or an amplicon as amplified by a primer pair based on SEQ ID No. 6 and 7 or 9 and 10 as mentioned herein constitute preferred embodiments of the present invention. Thus, the sequences of SEQ ID No. 1 to 4 (and, if needed, the complementary sequences thereto) can be used to design primers for amplifications, i.e. serve as “beacons” in the sequence as relevant. Similarly, additional primers and probes can be designed based on the amplicon according to SEQ ID No. 1. Amplification can take place either in the genomic and/or bisulfite (i.e. “converted”) DNA sequence.

The person of skill will furthermore be able to select specific subsets of CpG positions in order to minimize the amount of sites to be analyzed, for example at least one of CpG position selected from a CpG position in an amplicon according to SEQ ID No. 1, and is preferably selected from the CpG positions 27, 47, 82, 136, 194, 197, 249, 285, 290, 303, 336, 354, and 369 in the amplicon 1876 according to SEQ ID No. 2, CpG positions 31, 60, 75, 86, 114, 138, 142, 171, 184, 210, 217, and 241 in the amplicon 1877 according to SEQ ID No. 3, CpG positions 35, 56, 74, 104, 118, 130, 150, 182, 196, and 212 in the amplicon 1878 according to SEQ ID No. 4, and is preferably selected from CpG positions 60, 75, 86, 114, 138, 142, 171, 184, 210, 217, and 241 in a fragment of the amplicon 1877 according to SEQ ID No. 3. Preferred are combinations of 3, 4, 5, 6, 7, 8, 9, or 10 positions, the analysis of which produces sufficient data and/or information in order to be informative in the context of the present invention.

The person of skill will furthermore be able to select specific subsets of CpG positions in order to minimize the amount of sites to be analyzed, for example at least one of CpG position 60, 75, 86, 114, 138, 142, 171, 184, 210, 217, and 241 in the amplicon No. 1877 of the PDCD1 specific bisulfite convertible region (SEQ ID No. 1), or all sites as present on the bisulfite convertible region according to SEQ ID No 1. One or more of positions 60, and/or 138 in AMP 1877 may be excluded.

In order to analyze the bisulfite convertibility of CpG positions, any known method to analyze DNA methylation can be used. In a preferred embodiment of the method according to the present invention, the analysis of the methylation status comprises a method selected from methylation specific enzymatic digests, bisulphite sequencing, analysis selected from promoter methylation, CpG island methylation, MSP, HeavyMethyl, MethyLight, Ms-SNuPE or other methods relying on a detection of amplified DNA. These methods are well known to the person of skill, and can be found in the respective literature.

In a preferred embodiment of the method according to the present invention, said method is suitable for routine application, for example on a DNA-chip. Based on the above information and the respective literature, the person of skill will be able to adjust the method as above to such settings.

In yet another preferred embodiment of the methods according to the present invention, said method is performed without a step of purifying and/or enriching said cells to be identified, preferably using whole blood and/or non-trypsinized tissue.

In another preferred embodiment of the method according to the present invention, the identification comprises a distinction of said PD1+ cells from all major peripheral blood cell types and/or non-blood cells, preferably, but not limited to, cytotoxic T-cells, granulocytes, monocytes, B-cells, CD56++ NK cells, T-helper cells, and NKT cells, and other cell types derived from other organs than blood.

In yet another preferred embodiment of the method according to the present invention, the sample is selected from a mammalian body fluid, including human blood samples, or a tissue, organ or a sample of leukocytes or a purified or separated fraction of such tissue, organ or leukocytes or a cell type sample. Preferably, said mammal is a mouse, goat, dog, pig, cat, cow, rat, monkey or human. The samples can be suitably pooled, if required.

Another preferred aspect of the method according to the present invention then further comprises the step of concluding on the immune status of said mammal based on said B cells. The B cells can be quantified and be used as a benchmark to relatively quantify further detailed subpopulations, or it can be used as a predictive and/or screening and/or diagnostic and/or prognostic and/or adverse events detecting factor, or it can be used to finally detect this population to determine the overall immune activity status.

In yet another preferred embodiment of the methods according to the present invention, the mammal suffers from or is likely to suffer from autoimmune diseases, transplant rejections, infection diseases, cancer, and/or allergy as but not limited to Trypanosoma cruzi-infection, Malaria and HIV infection; Hematologic Malignancies as but not limited to chronic Myelogenous Leukemia, Multiple Myeloma, Non Hodgkin's Lymphoma, Hodgkin's Disease, chronic Lymphocytic Leukemia, Graft versus Host and Host versus Graft Disease, Mycosis fungoides, Extranodal T cell lymphoma, Cutaneous T cell lymphomas, Anaplastic large cell lymphoma, Angioimmunoblastic T cell lymphoma and other T-cell, B-cell and NK cell neoplasms, T cell deficiencies such as but not limited to lymphocytopenia, severe combined immunodeficiency (SCID), Omenn syndrome, Cartilage-hair hypoplasia, acquired immune deficiency syndrome (AIDS), and hereditary conditions such as DiGeorge syndrome (DGS),

chromosomal breakage syndromes (CBSs), multiple sclerosis, rheumatoid arthritis, systemic lupus erythematosus, Sjögren's syndrome, systemic sclerosis, dermatomyositis, primary biliary cirrhosis, primary sclerosing cholangitis, ulcerative colitis, Crohn's disease, psoriasis, vitiligo, bullous pemphigoid, alopecia areata, idiopathic dilated cardiomyopathy, type 1 diabetes mellitus, Graves' disease, Hashimoto's thyroiditis, myasthenia gravis, IgA nephropathy, membranous nephropathy, and pernicious anemia; and B-cell and T-cell combined disorders such as but not limited to ataxia telangiectasia (AT) and Wiskott-Aldrich syndrome (WAS); and carcinomas such as but not limited to breast cancer, colorectal cancer, gastric cancer, pancreatic cancer, hepatocellular carcinoma, cholangiocarcinoma, melanoma, and head and neck cancer.

Another preferred aspect of the method according to the present invention then relates to a method as above, further comprising measuring and/or monitoring the amount of PD1+ cells in response to chemical and/or biological substances that are provided to said mammal, i.e. in response to a treatment of said patient. Said method comprises the steps as above, and comparing said relative amount of said cells as identified to a sample taken earlier or in parallel from the same mammal, and/or to a control sample. Based on the results as provided by the method(s) of the invention, the attending physician will be able to conclude on the immune status of the patient, and adjust a treatment of the underlying disease accordingly.

Preferably, said method is performed without a step of purifying and/or enriching cells, preferably in whole blood and/or non-trypsinized tissue, or any other biological sample potentially containing said PD1+ cells as e.g. a sample for cell transfer into a patient.

Another preferred aspect of the method according to the present invention then relates to a method as above, further comprising formulating said PD1+ cells as identified for transplantation into a patient. Pharmaceutical preparations for these purposes and methods for their production are performed according to methods known in the art of transplantation medicine.

Another preferred aspect of the method according to the present invention relates to an oligomer according to any of SEQ ID No. 6 to 11, or an amplicon according to SEQ ID No. 2 to 5.

Yet another preferred aspect of the present invention then relates to a kit for identifying, quantifying, and/or monitoring PD1+ cells in a mammal based on the analysis of the bisulfite accessibility of CpG positions in the gene region of PDCD1, comprising components for performing a method according to invention as described herein, in particular a kit comprising a) a bisulfite reagent, and b) materials for the analysis of the methylation status of CpG positions selected from the CpG positions in the region according to SEQ ID NO: 1, such as an oligomer selected from the sequences according to SEQ ID No. 6 to 11.

The present invention also encompasses the use of oligomers or amplicon or a kit according to the present invention for identifying and/or for monitoring PD1+ cells in a mammal as described herein.

As mentioned above, recently three new cytosine modifications were discovered. Therefore, it is expected that future scientific findings will correct epigenetic patterns of modification described in the past. These past patterns of cytosine modification encompass bisulfite convertible (non-methylated, non-modified) and non-convertible (methylated, modified) cytosine. Both termini need to be corrected, as described. According to the novel scientific findings (i) non-bisulfite convertible cytosine encompasses 5-methylcytosine (mC) and 5-hydroxymethylcytosine (hmC), and (ii) bisulfite convertible (i.e. the “bisulfite convertibility”) cytosine encompasses 5-formylcytosine (fC), 5-carboxycytosine (cC), as well as non-modified cytosine.

Additionally, past inventions are based on (i) the ratio of bisulfite convertible cytosine to whole amount of chromatin (cell-type independent, 100% bisulfite convertible DNA locus) or (ii) on the ratio of bisulfite convertible cytosine (fC, cC, non-modified cytosine) to non-bisulfite convertible cytosine (hmC and mC). These ratios characterize cell type, cell differentiation, cell stage as well as pathological cell stages. Therefore, new techniques will result in novel, more specific ratios and might supplement current cell specific, cell state specific as well as pathological patterns of epigenetic modifications and therefore, define potential novel biomarkers. Novel ratios to be discovered as biomarkers can be defined as:

$$\text{Biomarker Ratio} = a/b$$

$$a = \sum (\text{C and/or mC and/or hmC and/or fC and/or cC})$$

$$b = \sum (\text{C and/or mC and/or hmC and/or fC and/or cC}),$$

whereby a and b differs from each other by one to four kinds of modifications. Discovery of

novel DNA modifications will enlarge this enumeration.

For the purpose of definition for the present application, “epigenetic modifications” in the DNA sequence is referred to by the terminology of (i) bisulfite convertible cytosine (5-formylcytosine, (fC) and/or 5-carboxycytosine (cC)) and (ii) non-bisulfite convertible cytosine ((including 5-methylcytosine (mC), 5-hydroxymethylcytosine, (hmC)). As both kinds of methylation, mC and hmC, are not bisulfite convertible, it is not possible to distinguish between these two. Likewise, fC, cC as well as non-modified cytosine are bisulfite convertible and can also not be distinguished from each other as well. The term “methylated” DNA encompasses mC as well as hmC. The term “non-methylated” DNA encompasses fC, cC, and non-modified DNA. It is expected that novel variants of DNA modifications will be discovered in future. Each type of modification will be either bisulfite convertible or not. However, since the present method reliably distinguishes between the two groups, these novel modifications will also be usable as markers.

Furthermore, apart from the modifications of DNA, also histones undergo posttranslational modifications that alter their interaction with DNA and nuclear proteins. Modifications include methylation, acetylation, phosphorylation, ubiquitination, sumoylation, citrullination, and ADP-ribosylation. The core of the histones H2A, H2B, and H3 can also be modified. Histone modifications act in diverse biological processes such as gene regulation, DNA repair, chromosome condensation (mitosis) and spermatogenesis (meiosis). Also for these modifications a specific pattern of modification is specific for different cell types, cell stages, differentiation status and such a pattern can be analyzed for bisulfite convertibility or similar methods in order to identify certain cells and cell stages. The present invention also encompasses a use of these modifications.

In summary, using the PDCD1 genetic region and in particular the amplicon as described herein as a marker, the inventors very specifically identified, quantified and particularly differentiated PD1+ cells, and in their relation to other cell types in a sample, for example to other blood cells.

The invention will now be further described in the following examples and with reference to the accompanying figures and the sequence listing, without being limited thereto. For the purposes of the present invention, all references as cited herein are incorporated by reference

in their entirety.

Figure 1 shows the analysis of CpG sites on amplicons No. 1876, 1877, and 1878 (SEQ ID No. 2 to 4, respectively) according to the invention. The horizontal boxes in the table correspond to the CpG positions in the amplicon as analyzed (e.g. CpG 1, 2, etc.) with the positions indicated (AMP1876:27 corresponding to CpG 1 of Amplicon 1876 ...etc.), and the columns correspond to the cell types as analyzed.

Figure 2 shows the specificity of the TpG-specific PCR-system according to the invention using test-templates (plasmid-DNA).

Figure 3 shows the genomic region of the amplicons according to the present invention, amplicon sequences are underlined.

Figure 4 shows the positions of the amplicons of the invention in the genome.

SEQ ID No. 1 shows the genomic region of the amplicons No. 1876, 1877, 1878, and 1879 according to the present invention (see Figure 3).

SEQ ID Nos. 2 to 5 show the sequences of amplicons No. 1876, 1877, 1878, and 1879 respectively.

SEQ ID Nos. 6 to 11 show the sequences of specific oligomers (primers and probes) according to the present invention.

SEQ ID Nos. 12 to 13 show the TpG converted and CpG converted sequences, respectively, of the AMP1877 of the invention.

EXAMPLES

Example 1

In order to identify PD1+ cells, qPCR was performed on bisulphite converted samples stemming from the human genomic region according to the sequence SEQ ID No. 1 (see Figure 3), in particular the regions AMP 1876, AMP 1877, AMP 1878, and AMP 1879

(underlined)

For the actual epigenetic profiling of the amplicon region in blood cell subtypes, the immune cell populations as analyzed were as shown in Figure 1.

The bisulfite-converted target-regions of preferred qPCR-assay-system as developed were:

1877 Primers (qPCR30 FW_T)	-	GTTTAGATTAGATTTGGTATTTTTGATT
(SEQ ID NO: 6)		
qPCR30_RV_T	-	CAAATCCTCTAAAAACAACTCA (SEQ ID NO: 7)
qPCR30 Probe_T and C:	-	TCCCAACACAACCCATAAAACAATTTC (SEQ ID NO: 8)
qPCR30_FW_C	-	AGATTAGATTTCGGTATTTTTGATCG (SEQ ID NO: 9)
qPCR30_RV_C	-	CAAATCCTCTAAAAACAACTCG (SEQ ID NO: 10)
qPCR30_P_C	-	CCCAACACAACCCGTAAAACGATTTC (SEQ ID NO: 11)

1877-TpG converted (SEQ ID NO: 12)

TTaggtTTtTtagggaTaagTtTgTtgtTTtTatTTTtagTaTagTTTgtgggaTggtttTTttgtTTTtaatgggaTTaTgggTagagatgTTgggtTtggtTtgggTTagTaggttTTtTTgTTTggggTaggTagTTttTtTtgtgTgTttTtggaaagTaatgtTTtgtaatgTgggTtTtTgTgggagTaTTTTTaTTgTTaTTtTaTaggTTtgttTTaTagTTT TgggatgggTtTgtTtTTTTTTtgaTTTtgT

1877 -CpG converted (SEQ ID NO: 13)

TTaggtTTtTtagggaTaagTtCgTtgtTTtTatTTTtagTaTagTTCgtgggaCggtttTTttgtTTTtaatgggaTTaCgggTagagatgTCgggtTtggtTtgggTTagTaggttTTtTCgTTCggggTaggTagTTttTtTtgtgCgTttTtggaaagTaatgtTTtgtaatgCgggTtTtTgCgggagTaTTTTTaTCgTTaTTtTaTaggTTtgttTTaTagTTT CgggatgggTtTgtTtTTTTTTtgaTTTtgT

The specificity of the TpG-specific PCR-system was demonstrated using test-templates (plasmid-DNA) as shown in Figure 2.

The cell type specificity (as measured by qPCR) was found as follows (table 1):

Cell type	Description	Demethylation (%)
T helper cells	CD3+CD4+	4.7
Cytotoxic T cells	CD3+CD8+	0.8
NK cells	CD56+	0.1
Granulocytes	CD15+	0.7
Monocytes	CD14+	0.4
B cells	CD19+	0.3
TFH cells	CD3+CD4+CXCR5+Bcl+PD1+	76.6

CLAIMS

1. A method for identifying PD1+ cells in a sample, comprising analyzing the methylation status of at least one CpG position in the mammalian gene region for Programmed cell death 1 (PDCD1), wherein preferably said gene region as analyzed is positioned according to SEQ ID No. 1, wherein a demethylation or lack of methylation of said gene region is indicative for a PD1+ cell, when compared to a non-PD1+ cell.
2. The method according to claim 1, wherein said at least one CpG position is present in the 5' region upstream from the transcription start, promoter region, the 5' or 3' untranslated regions, exon, intron, exon/intron border and/or in the 3' region downstream of the transcriptional stop of said gene region as analyzed.
3. The method according to claim 1 or 2, wherein said at least one CpG position is selected from a CpG selected from the CpG positions 27, 47, 82, 136, 194, 197, 249, 285, 290, 303, 336, 354, and 369 in the amplicon 1876 according to SEQ ID No. 2, CpG positions 31, 60, 75, 86, 114, 138, 142, 171, 184, 210, 217, and 241 in the amplicon 1877 according to SEQ ID No. 3, CpG positions 35, 56, 74, 104, 118, 130, 150, 182, 196, and 212 in the amplicon 1878 according to SEQ ID No. 4, and is preferably selected from CpG positions 60, 75, 86, 114, 138, 142, 171, 184, 210, 217, and 241 in a fragment of the amplicon 1877 according to SEQ ID No. 3.
4. The method according to any one of claims 1 to 3, wherein said analysis of the bisulfite convertibility comprises a method selected from a methylation specific enzymatic digest, bisulfite sequencing, an analysis selected from promoter methylation, CpG island methylation, MSP, HeavyMethyl, MethyLight, Ms-SNuPE, and other methods relying on a detection of amplified DNA.
5. The method according to any one of claims 1 to 4, further comprising a quantification of the relative amount of PD1+ cells based on comparing relative amounts of said methylation frequency in the region as analyzed with relative amounts of the methylation frequency in a control gene, such as, for example, GAPDH.

6. The method according to any one of claims 1 to 5, wherein said sample is selected from a mammalian body fluid, including human blood samples, or a tissue, organ or cell type blood sample, a sample of blood lymphocytes or a fraction thereof.
7. The method according to any one of claims 1 to 6, further comprising a distinguishing of said PD1+ cells from all or at least one of the cell types selected from PD1- follicular cells, cytotoxic T-cells, granulocytes, monocytes, B-cells, CD56++ NK cells, T-helper cells, and NKT cells, and other cell types derived from other organs than blood.
8. The method according to any one of claims 1 to 7, wherein said method is performed without a step of purifying and/or enriching said cells to be identified, preferably using whole blood and/or non-trypsinized tissue.
9. The method according to any one of claims 1 to 8, further comprising the step of concluding on the immune status of said mammal based on said PD1+ cells as identified.
10. A method for monitoring the level of PD1+ cells in a mammal, comprising performing the method according to any one of claims 5 to 9, and furthermore comparing said relative amount of said cells as identified to a sample taken earlier or in parallel from the same mammal, and/or to a control sample.
11. The method according to any one of claims 1 to 10, further comprising measuring and/or monitoring the amount of said PD1+ cells in response to chemical and/or biological substances that are provided to said mammal.
12. The method according to any one of claims 1 to 11, wherein said mammal suffers from or is likely to suffer from autoimmune diseases, transplant rejections, infection diseases, cancer, and/or allergy.
13. A kit for identifying, quantifying, and/or monitoring PD1+ cells in a mammal based on the analysis of the bisulfite accessibility of CpG positions in the gene region of PDCD1, comprising components for performing a method according to any of claims 1 to 12, in particular a kit comprising a) a bisulfite reagent, and b) materials for the analysis of the

methylation status of CpG positions selected from the CpG positions in the region according to SEQ ID NO: 1, such as an oligomer selected from the sequences according to SEQ ID NOS: 6 to 11.

14. An oligomer according to any of SEQ ID No. 6 to 11, or the amplicon according to SEQ ID No. 2, 3, 4, or 5.

15. Use of the kit according to claim 13, or of the oligomer or amplicon according to claim 14 for identifying, quantifying, and/or monitoring PD1+ cells in a mammal.




Fig. 1




Fig. 1 continued





Fig. 1 continued

Fig. 2

agtagacaatgtggagagacatcaggcagccctggcctcagtgaggcgccagctccctggtccacagctgggcccac
cccaccaggtctcctctcccatggcccaggggcttcagtgaggactgagaggaggagggaaggagagtggggtgac
agggaaagaactgcaggagagagaggagaggggtgggagaaggagaagggaagggaagggttaggatggaagctgggtt
tctccctgtgcccgccttactccaggacatgtgtccaagccctggcaggtggaattttgggggcagggccttg
gtggtgaggagaccttccaggggtctgatagcatctcccatctcagagcccacctcctgggcccagcctccctc
cagcccacacagtggttcctcagtcctcagaggacagcttcgtcccacaaagctcagagccttgagggaaggccc
actgctgcccgtggaacagagacagcattcaacagaggttggaaacaaggctctacagggctgggggcagagggagg
ttctgtccagaatctgccttcaggacaagtacagccagcaggggagccttagccacttatccactgcctgggcga
ggcacagggtatggaggcacctaccaaccaacagtttctccagccccagagccccagccccctgaggcacaagggt
gggtgtgccaggagacagttgctgcggggccaccttagctgtctggcagcacagtggtgctgccaggtccctgg
gggccccccgccaagcccacctggccagctgggccccccccccacctccccaccaagccacccacacagcctcaca
tctctgagacccgggagtgggcctttgttcataaacgagagcttctcgcgctggccgcgctcgcagacatcat
ctttgatgtctcttttccactgtttcgtgtctttaatgttttcccttcagagccgggcccagtgctctcctggagc
caggcagcgcgcagctgtcaggcgggttcttagcctcgtctcgtttattttaagctgatgagcctgacgcactct
catcactaataatcagcagtttcatcttctgttttccactcgtctgtaataaaatgctcagcacagaataacaagg
agataagcaagccatttcacaaaacgcggggcgccagccagggcccagggcactggacccccctgaaccacccccccc
tggcacagtggtggctggagggcagggccccggggaagaagggtcaaggctggaaggggagggtcagcctcacagcca
gccccctgccaccgccccagcccccccgctcaggctgttgaggcatcacacggtggaaagatctggaactgtggcc
atggtgtgaggccatccacaagggtggaagctttgagggggagccgattagccatggacagttgtcattcagtagg
gtcacctgtgccccagcgaagggggatgggcccgggaaggcagaggccagggcacctgccccagcaggggagagg
ctgtgggcagccgggaggtcccagaggtccgacagaatgggagtggggttgagcccacccctcactgcagccc
aggaacctgagcccagagggggccacccaccttcccaggcaggggagggcccgccccagggagatgggggggat
gggggaggagaagggctgccccacccgcgcagcctcaggaggggcagctcgggcgggatatggaaagaggccac
agcagtgagcagagacacagaggaggaaggggccctgagctggggagacccccacggggtagggcggtgggggcca
cgggcccacctcctccccatctcctctgtctcctctgtctctctctctcctccccacacctctccccagtc
ctacccccctcctcacccctcctccccagcactgcctctgtcactctcgcccacgtggatgtggagggaagagggg
gcgggagcaaggggcgggcacccctcccttcaacctgacctgggacagtttcccttccgctcacctccgctgago
agtggagaaggcggcactctggtggggctgctccaggcatgcagatcccacaggcgccctggccagtcgtctggg
cgggtgtacaaactgggctggcgccaggatggttcttaggtagggtggggtcggcggtcaggtgtccagagccag
gggtctggagggaccttccacctcagtccttggcaggtcggggggtgctgaggcgggcctggccctggcagccc
aggggtcccggagcaggggtctggagggacctttcactctcagtccttggcaggtcgggggggtgctgtggcagg
cccagccttgccccacagctctgcccccttaccttgagctgtgtggtttgggcagctcgaactcctgggttctc
tctgggccccaaactcctccccctggcccaagtcctcttctgtctcgtggcaggcaggacctgtccccctcag
ccgttcttgggctgcgtgtttctgtagaatgacgggtcaggctggccagaacccccaaaccttggccgtgggga
gtctgcgtggcggtctgccttggccaggcatccttggctcctcactcgagttttcctaaggatgggatgagcccc
atgtgggactaaccttggctttacgacgtcaaagtttagatgagctggtgatatttttctcattatatccaaagt
gtacctgttcgagtgaggacagttcttctgtctccaggatccctcctgggtggggattgtgccgcctgggtctc
tgcccagattccagggtctctccccgagccctgttcagaccatccgtgggggaggccttggcctcactctcccga
tcgaggagagaggggagcctcttctgggtgcccgtgacctgggcccctctgtgtacactgtgaccacagccgc
tcttgacctctgtgcccggctggccctctgtgcccagccagcctgcacctggggatgccaaggcctggggagg
gtggtttcaccagggccaagcctaagacagtcctctgtggccctgtgtggtaccggggtgtgacaccactgggag
gacaagatgaggggacccctggggcgccctgacacccct_{cg}aggctcctgccc_{cg}ggggtcctggtgcccctt
cactgtggcagg_{cg}actgggggttccccacct_{cg}gccccctcc_{cg}gggctgctccccggcacctgaggcagcatc
cttgtcagggc_{cg}tgcccttctgcctcag_{cg}ccacctcttaaggttggcc_{cg}tgggtcactcaggactcagaactgg
agattctgggcaaaaggcaaaagagcaaaaggccaaaaggcatcccaggagagcagctcggggggaaccagagggc
agaggggcgctcgtcacaggggagggggagctgagcgaggcagggaggggagccgagcctctccccccgtgtcccg
gctcttcaggcacgcctcgggacgccacctccccgacccaggcgggaaagataagagcaagggtgtccgcagcc
tgacactcgtgcctcaggtgcccgcgcttgtgccggacaagactctcacaggtggcatgcctcggtttccccact
ggtaacagcacagggcactcagcaaggcgagtgggcatgactggggctctgtgggtcctgacccagatgtggcc
accccgccgcagtggtcttccattccaggatgcctcttttccctcctgatctattcactgcgttcgccattcgg
ctattggccccgtccgagggcaggagcataaataagagcctctgcttttggcggtgggaccactgtgagctccagtc
agcgtgccactgctgagctctgggcttcgacaggacttggcccccttactgactctccgtgtgctttgggtca
tgggtgaggacgctcctggcaaggctgcgtcctgaggattaaatcggggtcatctgtgaaaactaccagcccaag

Fig. 3 a

cccttgacacttttttgtttgtttcttttagtgacaggggtcttgcctctgtcaccagggctggagtgacgtggtgt
gatctcggtctcactcgacctccggggctcaagcaattctcccacctctgcctccagagtagctgggactatagggc
acgtgccacctctgccaggctaatcttctccatttttttagagacaggggtctcgctatgttccccaggctgggct
caaatggctcctcccacctcagcctcccaagtaactgggattacaggcataagccactgcattctggcctccatgac
acatatttttaaaagtctgatttttaaaagtcaaaacttttgaagtacagatttttaaacggactattttgaaaaatata
caaaaacgttttaaaaacaatgaatatccctcacctagaatcaataactaagaatattgacacatttgcctttgggg
actggggcggtggagctgccatgacaaagctccgcgcagcagagtggttttaaacagagcctgcccctctcgccga
ctgagggctggagctgcaggatggagctccgcagggtcggtctcccctgtgctctgaggggtctgctcagcctct
cccggctgtggcttaaaaaacagagcctgtcctccgcgcgtggggggctggacatgcaggaccgagggggccacagg
gtcggtctccctgtgctccgagagggctctgctcagcttctcctggctgggggggttttggggccacctctgtggt
cctgggttcagaagcatccccaggctctgcttcatctctgcacgggtgactctgtacaggaagccaggcctgc
tggtaaatggccaccagccctgtgcccctcatcttacctagtccagctgcggtcaccctattcttaataaggcc
gcctctgaggtcatgggggttaggacttccacataggaatctgtggggacaggttcggcccacagcccttccca
cctccacacacacacagactgtgaggagttggaagacctcactcctcaccctgcccaggctcctctagggacaag
ctcgctgtcctcattccagcagagcccggtgggaagggtttccttgtccctaatgggaccacggtcagagatgcggg
gtctggtctggggccagcaggttctcgcgcgggggcaggcagccttctctgtgcgcttctggaaagcaatgtcc
tgtaatgcgggtctctctgcggggagcacccccacggccacctcagaggcctgttccacagccccgggatgggctct
gtctccctcctgacctgcattagggcacagccctctctcattcaacccacgactcctacgtggatccgagaggggagc
acctggggaaacaatggaatcccatagaaacacccccaaatctaaacttgatccaggaccagccagtggtcacttct
gaatattcaccttcttagtagacactaccagccaagggagggccaggaagccttctggaggaggtggcctgagga
ctgggggtgaggcagggcctgcgtgggggtgcgccaccagcacccccacactgggtgggagccagctctctgagact
ggctgggggaggtgggagaggggggtgcttgaactgcagacaccgaggtctagccccacccccacccagccagtt
ggtggaggcaggggagggccgagggggccagctggacctgctccccgggggtggattccaaaataggggggttgggg
ggggcggaacaggagcccagggtcctggcttgaggcccagtggtgagggctggtgcaagccagacaggaaaagg
gttgagcctgtcagcgccagcagatcaagtcaggagcaggtccctccaccaatgtgtgcaataaataagcagc
taagtttccagttacaagaacaatgcacagatgggtcccaggggacattg_{cg}gtgtggacacacag_{cg}gccattgtcc
tgt_{cg}ccagcacct_{cg}ccctacagctgggggggtcccttagcacttcttagccatgcagggtcctgctcacagtac
c_{cg}tgatgacttctgttctcactgctgctgtctgtcccagacagctgcattggcagccctggcctgggagatggagac
cc_{cg}aggggggtgctg_{cg}agtggtggggccctgggtcccccagtcattccagaaacccaggggcagggcatttc
cctgctctgtgc_{cg}agtcaccccagccccagctaggccccagtaaggggtgcagcccacctgtcccaggctgccc
tcccaggagccctcttggccctgatgccagaagcccatcttctccattcaggcaggtctctgagtgccctggcc
tggtgctgctggtgcccctgagagtcacactaccccacagccctccttgggtcaaaaatccactctggagtggtgga
agattccccggggccacgcgcacacgcctatgcaggagcttcccctggccggccggcagacaaagggcggtctc
agagaggggggtcactcagcagcccttgtgtagctggccctcgccctgccacctctgggaacaccaccagga
agctgggggacaggcacgcaggtgaaggaggcgagcgcttgtcagccgggagggccatgggacagaggggaacagg
gacacctgggtggcctcaaggtcacttcaaacccctcactcgctcccctgggaggggtgcccagtgaggttggcac
taggagttggtcctggtcacatgacagaccccccacctctgggtgtccagccagcagccgctggggccagcctggc
tgcagggacacgagggcagcagcccccctcctcctctgagctgggttgccttgagtcataccacccgctgccac
ggaggccgctgtcccaggaagcagagggacgcagctgtggcaaccaggggcctggtctctgtgtcacctcgctg
gggggcgctgcccaggcctgagacggaactgagtgacagtgcaactgggtctgacagtggtggggctggcgccatgt
ttggggaaacctgtggcatgggaacctgtgggtgagccgggaaaaatcaccccggttgcatggcatctcgggcctgga
tcttaagcgctgtgttgggtgctcgcctggcggaagagccgcgacccccacggttgccatgcgggtatccccag
cctgaccttggcaggcatatgtttcaggaggtccttgtcttgggagcccagggtcgggggcccccgctgtctgtcc
acatccgagtcattggcccatctcgctctctgaagcatcttctgtgtgagctctagtcacctgtcttgcctggaa
aatgtggagggccacctgcccactgcccagggcagcaatgcccataccacgtgggtcccagctccgagcttgcct
gaaaaggggggcaaaagactggacctgagcctgccaaggggccacactcctcccagggtcggggtctccatgggca
gccccccacccacccagaccagttacactcccctgtgcccagagcagtgccagacaggaccaggccaggatgcccac
gggtcaggggtggtggatgggtagcccccaaacagcccttcttgggggaactggcctcaacgggggaaggggtga
aggctcttagtaggaaatcagggagacccaagtacagagccaggtgctgtgcagaagctgcagcctcacgtagaag
gaagaggctctgcagtgaggccagtgcccacccccgggtggcagagggcccagcagagacttctcaatgacatt
ccagctgggggtggcccttccagagcccttgcctgcccaggggatgtgagcaggtggccgggggaggtcttgtggggc
caccagcccttctcactctctccatctctcagactcccagacaggccctggaaacccccccaccttctccc
cagccctgctcggtgacccaaggggacaacgccaccttcaactgcagcttctccaacacatcgagagacttgc
tgctaaactgggtaccgcatgagccccagcaaccagacggacaagctggccgcttcccagaggaccgcagccagc
ccggccaggactgcgcttccgtgtcacacaactgcccacggggcgtgacttccacatgagcgtgggcaggggccc
ggcgcaatgacagcggcacctacctctgtggggccatctccttggcccccaaggcgagatcaaagagagcctgc
gggcagagctcaggggtgacaggtgcggcctcgaggggcccggggcaggggtgagctgagccggctcctgggggtggg

Fig. 3 b

[illegible]

Fig. 3 c




Fig. 4