

**(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. AU 2006273928 B2

(54) Title
Cell/ligand marking system, wherein the marker is of Eph type, cell material comprising said system, method for preparing same and proangiogenetic use

(51) International Patent Classification(s)
C12N 5/10 (2006.01) **A61K 38/17** (2006.01)
A61K 35/14 (2006.01) **A61P 9/00** (2006.01)
A61K 35/44 (2006.01) **C12N 5/074** (2010.01)

(21) Application No: **2006273928** (22) Date of Filing: **2006.07.27**

(87) WIPO No: **WO07/012764**

(30) Priority Data

(31) Number **0508029** (32) Date **2005.07.27** (33) Country **FR**

(43) Publication Date: **2007.02.01**
(44) Accepted Journal Date: **2012.02.16**

(71) Applicant(s)
Institut des Vaisseaux et du Sang;Universite Paris 7-Denis Diderot

(72) Inventor(s)
Foubert, Philippe;Silvestre, Jean-Sebastien;Le Ricousse-Roussanne, Sophie

(74) Agent / Attorney
Davies Collison Cave, 1 Nicholson Street, Melbourne, VIC, 3000

(56) Related Art
Maekawa et al, Arteriosclerosis Thrombosis Vascular Biology (2003), Vol 23, No 11 pages 2008-2014.

WO 2007/012764 A3

(12) DEMANDE INTERNATIONALE PUBLIÉE EN VERTU DU TRAITÉ DE COOPÉRATION
EN MATIÈRE DE BREVETS (PCT)

(19) Organisation Mondiale de la Propriété
Intellectuelle
Bureau international



(43) Date de la publication internationale
1 février 2007 (01.02.2007)

PCT

(10) Numéro de publication internationale
WO 2007/012764 A3

(51) Classification internationale des brevets :
C12N 5/06 (2006.01) A61K 35/44 (2006.01) [FR/FR]; Bureau de la Valorisation et des Recherches, 2, Place Jussieu, F-75251 Paris Cedex 05 (FR).

(21) Numéro de la demande internationale : (72) Inventeurs; et
PCT/FR2006/001837 (75) Inventeurs/Déposants (pour US seulement) : FOU-BERT, Philippe [FR/FR]; 9, rue de Conflans, F-94220 Charenton le Pont (FR). SILVESTRE, Jean-Sébastien [FR/FR]; 63, Avenue Ledru Rollin, F-75012 Paris (FR). LE RICOUSSE-ROUSSANNE, Sophie [FR/FR]; 56, rue Musselburgh-Escalier 17, F-94500 Champigny-sur-Marne (FR).

(22) Date de dépôt international : 27 juillet 2006 (27.07.2006)

(25) Langue de dépôt : français

(26) Langue de publication : français

(30) Données relatives à la priorité : 0508029 27 juillet 2005 (27.07.2005) FR

(71) Déposants (pour tous les États désignés sauf US) : INSTITUT DES VAISSEAUX ET DU SANG [FR/FR]; Hôpital Lariboisière, 2, Rue Ambroise Paré, F-75475 Paris Cedex 10 (FR). UNIVERSITE PARIS 7-DENIS DIDEROT (74) Mandataire : BERTRAND, Didier; c/o SA l'edit-Loriot & Autres Conseils en Propriété, Industrielle, 38, avenue Hoche, F-75008 Paris (FR).


(81) États désignés (sauf indication contraire, pour tout titre de protection nationale disponible) : AE, AG, AL, AM, AT,

[Suite sur la page suivante]

(54) Title: CELL/LIGAND MARKING SYSTEM, WHEREIN THE MARKER IS OF EPH TYPE, CELL MATERIAL COMPRISING SAID SYSTEM, METHOD FOR PREPARING SAME AND PROANGIOGENETIC USE

(54) Titre : SYSTÈME MARQUEUR CELLULAIRE/LIGAND, OÙ LE MARQUEUR EST DU TYPE EPH, MATÉRIAUX CELLULAIRES COMPRENANT CE SYSTÈME, PROCÉDÉ DE PRÉPARATION ET UTILISATION PROANGIOGÉNIQUE

AA
Score angiographique
(Patte ischémique / patte non ischémique)



AA ANGIOGRAPHIC SCORE (ISCHEMIC PAW/NON-ISCHEMIC PAW)

(57) Abstract: The invention concerns a cell/ligand specific marking system, said system being characterized in that it comprises: (a) an endothelial precursor cell (EPC) including a cell marker selected among the group consisting of the Eph (in particular EphB4 or EphB1), and (b) a protein material of structure L-K, which consists of a ligand (L) specific of said marker and is associated or fused with a binding protein (K), said system being capable of providing a proangiogenic cell material of structure EPC-Eph-L-K. The invention also concerns said cell material as product capable of stimulating angiogenesis, its preparation method and its therapeutic use, in particular with respect to vascular insufficiencies.

[Suite sur la page suivante]



AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **États désignés (sauf indication contraire, pour tout titre de protection régionale disponible) :** ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), eurasien (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), européen (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,

RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Publiée :

- *avec rapport de recherche internationale*
- *avant l'expiration du délai prévu pour la modification des revendications, sera republiée si des modifications sont reçues*

(88) Date de publication du rapport de recherche internationale:

31 mai 2007

En ce qui concerne les codes à deux lettres et autres abréviations, se référer aux "Notes explicatives relatives aux codes et abréviations" figurant au début de chaque numéro ordinaire de la Gazette du PCT.

(57) Abrégé : La présente invention concerne un système du type marqueur cellulaire/ligand spécifique, ledit système, étant caractérisé en ce qu'il comprend : (a) un précurseur de cellule endothéliale (EPC) comportant un marqueur cellulaire choisi parmi l'ensemble constitué par les Eph (notamment EphB4 ou EphB1), et (b) un matériau protéïnique de structure L-K, qui est constitué d'un ligand (L) spécifique dudit marqueur et est associé ou fusionné avec une protéine de liaison (K), ledit système étant capable de fournir un matériau cellulaire proangiogénique de structure EPC-Eph-L-K. Elle concerne également ce matériau cellulaire en tant que produit capable de stimuler l'angiogenèse, son procédé de préparation et son utilisation en thérapeutique, notamment vis-à-vis des insuffisances vasculaires.

Cell/ligand marking system, wherein the marker is of Eph type, cell material comprising said system, method for preparing same and proangiogenetic use

5 ***Field of the invention***

The present invention relates to a novel technical solution using a system comprising a cell marker and a ligand specific for said marker, wherein said marker is 10 selected from the group consisting of Ephs, in particular the EphB4 marker. It also relates to the cell material comprising this system and to the method for preparing it and its therapeutic use as a proangiogenic agent.

15

Prior art

It is known that the members of the ephrin family and their Eph receptor tyrosine kinases, which were 20 initially demonstrated in the nervous system for neurone guidance, are factors involved in angiogenesis. Many isoforms for Eph receptors and their ephrin ligands have been described with tissue expression specificities. In vertebrates, firstly, at least 16 Eph 25 receptors are known, namely: 10 EphA receptors (EphA1 to EphA10) and 6 EphB receptors (EphB1 to EphB6) and, secondly, at least 9 ephrin ligands are known, namely: ephrin-A1 to ephrin-A6 and ephrin-B1 to ephrin-B3. Originally, the subdivision into two classes, EphA and 30 EphB, was based on the homology of the sequence of their extracellular domain, but this subdivision also corresponds to the preferential binding of their ligands, the ephrin-A ligands being in general bound to the membrane by a glycosylphosphatidylinositol (GPI) 35 and the ephrin-B ligands being transmembrane with an intracytoplasmic domain having a PDZ-binding domain.

As represented in figure 1 hereinafter:

- 5 • the Eph receptor comprises the extracellular domain, fibronectin type III repeats, a cysteine rich region and a specific ligand-binding domain; it also has an extension in the intracellular domain comprising a domain with tyrosine kinase activity, a sterile α motif (SAM) and a PDZ-binding motif;
- 10 • the ephrin-A ligand is attached to the membrane via a GPI (GPI anchor); and
- 15 • the ephrin-B transmembrane ligand has an intracellular extension comprising a cytoplasmic tail and a PDZ-binding motif.

With regard to the interaction between the Eph receptor tyrosine kinase (in particular EphB4) and its ligand (in particular ephrin-B2), a transmembrane protein, 20 bidirectional signaling takes place, which comprises signaling induced by the Eph receptor in the cell expressing Eph on its surface (forward signaling) and reverse signaling induced by the ephrin ligand in the cell expressing ephrin at its surface. This 25 bidirectional signaling plays a role in cell/cell contacts, migration and cell adhesion.

The specificity of arteriovenous expression of the 30 EphB4/ephrin-B2 couple is known from the article by H. U. Wang et al., "Molecular distinction and angiogenic interaction between embryonic arteries and veins revealed by ephrin-B2 and its receptor EphB4", *Cell*, 1998; 93: 741-753, EphB4 being expressed only by the veins, and ephrin-B2 being expressed by the 35 arteries.

EphB receptors and their ephrin-B ligands are expressed particularly during embryonic development. According to

Z. Wang *et al.*, "Ephrin receptor, EphB4, regulates ES cell differentiation of primitive mammalian hemangioblasts, blood, cardiomyocytes and blood vessels", *Blood*, 2004; 103: 100-109, EphB4 appears to 5 be involved in the differentiation of mouse embryonic cells (ES) to endothelial cells.

It is also known that certain Eph receptors, in particular EphB4, are overexpressed in tumors. This 10 observation suggests that these Eph receptors play a role in tumor progression. The article by N. K. Noren *et al.*, "Interplay between EphB4 on tumor cells and vascular ephrin-B2 regulates tumor growth", *Proc. Natl. Acad. Sci. USA*, 101: 5583-5588, demonstrates that the 15 activation of ephrin-B2 by EphB4 (reverse signaling) stimulates tumor growth.

On the other hand, what is observed with the prior literature is (i) an increase in the expression of 20 ephrin-B2 during tumorigenesis and, secondarily, (ii) a concern, due to this increase, that ephrin-B2 would be capable of inducing tumors. This explains why several publications propose, on the one hand, using the detection of ephrin-B2 as a tumor marker or, on the 25 other hand, inhibiting ephrin-B2; see, in particular, (i) WO 02/058538 A, which recommends detection of the ephrin-B2 ligand as a marker of the tumor vasculature and describes a technique for visualizing said tumor vasculature, (ii) US 2003/0207447 A, which proposes the 30 use of an inhibitor of Eph, in particular EphB4, or of an inhibitor of ephrin, in particular ephrin-B2, for limiting angiogenesis by preventing the interaction between the receptor and the ligand, and (iii) US 6864227 A and WO 03/102144 A, which recommend using 35 either an anti-(Eph) antibody [in particular an anti-(EphB4) antibody] or an anti-(ephrin) antibody [in particular an anti-(ephrin-B2) antibody] for reducing or modulating angiogenesis.

Briefly, the published patent documents relating to the Eph/ephrin couple concern methods that relate in particular to:

5

- ephrin-B signaling (reverse signaling) by proteins binding to the PDZ domain, see WO 02/079382 A and WO 00/031124;
- 10 • modulation of the expression of ephrin-B2 or of EphB4, see in particular US 2004/0110150 A, WO 04/006846 A and WO 04/080418;
- 15 • inhibition of the ephrin-B2/EphB4 interaction with a view to treating cancers and angiogenesis-related diseases, see in particular document WO 04/080418 A mentioned above and WO 04/069264 A; and
- 20 • the use of ephrin-B2 as a marker for the tumor vasculature, see in particular document WO 02/058538 A mentioned above.

In addition, the article by Maekawa Hiromitsu et al. "Ephrin-B2 induces migration of endothelial cells through the phosphatidylinositol-3 kinase pathway and promotes angiogenesis in adult vasculature", *Arteriosclerosis thrombosis and vascular biology*, 2003, 23: 2008-2014, shows that the ephrin-B2 ligand, coupled 25 to the Fc protein (i.e. the Fc fragment of an antibody), induces the migration of endothelial cells, HUVECs.

Finally, an adenovirus having the coding sequence of sphingosine kinase and a coding sequence of an angiogenic protein is known from US 6610534 B. The objective of this patent is the expression of sphingosine kinase and of the angiogenic protein, which 30

may be ephrin-B2, after local injection of said adenovirus. Moreover, a method for reducing the abnormal proliferation of hematopoietic stem cells, which comprises the administration by injection of a 5 water-soluble inhibitor of ephrin-B2, is known from the publication US 2005/0049194 A.

Summary of the invention

10 According to the invention, it is proposed to provide a novel solution, neither described nor suggested in the abovementioned prior art, to the technical problem of the stimulation of angiogenesis. More specifically, it is proposed here to provide a novel technical solution 15 using a "cell marker/ligand specific for said marker" system which is applied to particular cells, namely endothelial precursor cells, for solving the problem of angiogenic stimulation. An aspect of the invention is the improvement of previously known cell therapies by 20 activating the cells before they are injected, so as to obtain greater effectiveness of revascularization processes, in particular post-ischemic revascularization.

25 ***Subject of the invention***

The novel solution according to the invention uses a cell marker/ligand system wherein said marker is associated with an endothelial cell precursor cell. In 30 this system, the marker is an Eph receptor and the Eph-specific ligand is an ephrin ligand.

According to a first aspect of the invention, a novel cell marker/specific ligand proangiogenic system is 35 provided, said system, in which the cell marker is present on the outer membrane of the cell, wherein said system comprises:

(a) an endothelial precursor cell (EPC) comprising a cell marker selected from the group consisting of the Ephs, in particular EphB4 or EphB1, and

5

(b) a protein material of structure:

L-K (I)

10 which consists of a ligand (L) specific for said marker and is associated or fused with a binding protein (K), in particular an antibody Fc fragment,

15 said system being capable of providing a cell material of structure:

EPC-Eph-L-K (II)

20 which stimulates angiogenesis.

In other words, in order to stimulate angiogenesis, the EPC cells comprising the Eph marker are activated with the specific ligand L belonging to the ephrin family.

25 The ligand L may also consist of a peptide fragment of an ephrin, for example of ephrin-B2, which then would have the same biological activity.

According to a second aspect of the invention, a cell 30 material capable of stimulating angiogenesis is provided, wherein said cell material has the structure:

EPC-Eph-L-K (II)

35

wherein said ligand L is associated with or fused with a binding protein K.

18 Jan 2012

2006273928

This cell material is capable (i) of being in the form of a substantially purified cell culture or in the form of a cell culture in combination with other precursor cells, in particular mononuclear cells, and (ii) of being, where appropriate, frozen.

According to a third aspect of the invention, a method for preparing said cell material is provided, said method comprising the steps consisting of:

- (i) making use of EPCs expressing on their outer membrane a marker of the Eph family, in particular EphB4 or EphB1, and
- (ii) bringing said EPCs into contact, *in vitro*, with a protein material L-K wherein L is a ligand specific for said marker, and K is a binding protein associated or fused with L.

A medicament that can be used for reconstituting damaged vessels, wherein said medicament comprises, in combination with a physiologically acceptable excipient, a therapeutically acceptable amount of the cell material according to the invention, in particular as a proangiogenic active ingredient, is also recommended.

Finally, according to another aspect of the invention, a novel use of said cell material is provided, wherein said use is made of said cell material, as proangiogenic active ingredient, in combination with a physiologically acceptable excipient, for the preparation of a composition for therapeutic use in the treatment of vascular insufficiencies, in particular in the revascularization of ischemic cardiac, brain or peripheral tissues.

Brief description of the drawings

In the attached drawings,

5 - figure 1, presented in detail above, illustrates the prior state of knowledge and represents schematically the Eph receptor tyrosine kinases and their ligands of ephrin-A type or of ephrin-B type; and

10 - figures 2 to 4 show graphically the proangiogenic properties, with regard to the tests which were carried out, of the cell material according to the invention.

15

Detailed description of the invention

The article by E. B. Pasquale, *Curr. Opin. Cell Biol.*, 1997; 9(5):608, in particular describes the 20 specificities of the components of the Eph receptor/ephrin ligand system. For convenience, table I, given below and established according to this article (as indicated in US 6579683 A), gives these 25 specificities for certain couples, the ligands being placed in the second column by decreasing affinity with respect to their receptors reported in the first column.

In the cell marker/specific ligand system according to 30 the invention, one can imagine that the two components of the couple can each be in the form of an amino acid sequence or, where appropriate, in the form of a nucleic acid sequence. It is, however, clearly rather preferable to have, at the start, an Eph receptor 35 expressed on the EPC membrane in the form of an amino acid sequence. Similarly, the ephrin ligand will be very advantageously used in the form of an amino acid sequence or else of an ephrin peptide fragment. This is

because it is in the form of amino acids that Eph and ephrin bind, the protein/protein interaction being necessary in order to observe the proangiogenic activity. Consequently, in the subsequent text, Eph and 5 ephrin each intervene exclusively in the protein form of an amino acid sequence.

As Eph marker, an EphA or an EphB can be used here. However, it is preferred, according to the invention, 10 for the Eph marker to rather be an EphB, since EphB markers are predominantly involved in angiogenesis, whereas (in the current state of knowledge) EphA markers appear to be mainly involved in the nervous system.

15

According to the invention, the EphB marker will advantageously be EphB4 or EphB1; among the EphB markers, EphB4 will be preferred to EphB1.

20 According to the invention, among the ephrin ligands, preference will be given to ephrin-B ligands, in particular ephrin-B2 or ephrin-B1. In addition, a variant of ephrin-B2, for example, which corresponds to a peptide fragment of ephrin-B2 having the same 25 biological activity, may also be appropriate.

Consequently, in the system according to the invention, use will advantageously be made, firstly, of the EphB4 or EphB1 marker and, secondly, of the ephrin-B2 or 30 ephrin-B1 ligand, the Eph/ephrin couple that is more particularly preferred according to the invention being EphB4/ephrin-B2.

35 In order for the ligand L to be able to activate EPC-Eph, it is important, in the majority of cases, for it to be associated with or fused with a binding protein K in the form of a protein material of structure:

L-K

(I)

the recombinant ephrin-B2 protein being, to the applicant's knowledge, a substance capable of 5 activating EPC-Eph alone without the introduction of said binding protein K.

Among the binding proteins K which are suitable according to the invention, mention may in particular 10 be made of the numerous antibody Fc fragments (for example, those obtained by cleavage of antibodies with pepsin, papain or any other appropriate substance). For convenience, an Fc fragment readily available on the market as a by-product of the preparation of 15 antibodies, such as Fab' and F(ab)², is recommended here.

As a variant, and given the above, said protein material L-K can be replaced with the recombinant, 20 preferably human, ephrin-B2 protein.

The EPCs which are suitable according to the invention are cells which comprise an Eph cell marker (in particular an EphB, preferably EphB1 or better still 25 EphB4) expressed at their outer membrane. Such EPC cells are obtained from mononuclear cells or cells expressing CD34 or CD133, which originate from the bone marrow, from peripheral blood or better still from umbilical cord blood.

30

The mononuclear cells are produced in the bone marrow, where they are found at a high concentration, pass into the blood stream and are found in cord blood and in peripheral blood. They constitute a material of choice, 35 in the sense that, within their make-up, a considerable number of cells possess the genetic material required to (i) express the Eph marker (and more particularly the EphB4 marker) or (ii) include on their outer

membrane said marker already expressed. The mononuclear cells that are preferred according to the invention are those which are CD34⁺ or CD133⁺, since, after differentiation, they provide a relatively large amount 5 of EPCs which substantially express or include the EphB4 marker.

The concentration of EPCs produced by differentiation, which are contained in the population of mononuclear 10 cells, varies according to the origin of the cells. Said concentration is more or less equivalent in the bone marrow and the umbilical cord blood. On the other hand, it is lower in peripheral blood.

15 The cell material according to the invention, which is represented by structure II above, in which the protein material L-K can be replaced with the recombinant ephrin-B2 protein, consists of one or more cells, each having on its outer membrane an Eph marker, in particular EphB4 or EphB1, bound to its specific ligand. As a variant, said cell material may consist of a culture containing several cells of structure II in combination, where appropriate, with other cells not activated by said protein material; such a culture may 20 25 therefore be a cell mixture of activated EPCs, of nonactivated EPCs and of mononuclear cells.

According to a specific embodiment of the invention, said cell material is obtained by incubation 30

(a) of an endothelial precursor cell (EPC) comprising a cell marker selected from the group consisting of the Ephs, in particular EphB4 or EphB1, with

35

(b) a protein material of structure:

18 Jan 2012

2006273928

which consists of a ligand (L) specific for said marker and is associated or fused with a binding protein (K), in particular an antibody Fc fragment,
5

before being brought to its site of administration.

According to an preferred embodiment, the incubation
10 is carried out *in vitro* for a period of 10 to 60 minutes, in particular for 30 minutes, this incubation period being just before the administration of the cell material.

15 As a variant, said cell material may be in the form of a cell culture which is purified or which is a mixture of cells containing (i) EPC cells activated with L-K or the recombinant ephrin-B2 protein, and (ii) nonactivated precursors, for example mononuclear cells
20 and/or EPCs, which are not activated with L-K or said recombinant ephrin-B2 protein. Furthermore, the cell material may be conserved in the frozen state.

25 The cell material that is preferred according to the invention, from the point of view of what was indicated above, is of structure II wherein said marker is EphB4 or EphB1, and wherein said ligand is ephrin-B2 or ephrin-B1.

30 According to a preferred embodiment, the method of the invention for preparing said cell material comprises the following steps:

35 (1°) making use of mononuclear cells originating from the bone marrow, peripheral blood or umbilical cord blood,

(2°) differentiating said cells of the preceding step in order to obtain EPCs having the Eph marker, and

5 (3°) activating the EPCs thus obtained in step (2°), *in vitro*, by binding of ephrin to Eph.

Said method comprises, where appropriate, an additional
10 step between steps (1°) and (2°), i.e.:

(1a°) isolating the CD34⁺ and/or CD133⁺ mononuclear cells.

15 Step (1a°) implies a ternary process (1°) + (1a°) + (2°). However, the implementation of said step (1a°) increases the production costs compared with the binary process (1°) + (2°). In practice, since the nonactivated EPC cells serve to dilute the activated
20 EPC cells in the cell medium containing them, without impairing their action, the binary process is at the current time manifestly more profitable than the ternary process.


25 According to the method of the invention, said ligand L is associated with or fused with a binding protein K in order to provide a cell material of structure:

EPC-Eph-L-K (II)

30 wherein said Eph marker is EphB4 or EphB1, and said ligand L is ephrin-B2 or ephrin-B1.

As a medicament, the cell material according to the
35 invention can be used for treating especially vascular insufficiencies, in particular in the revascularization of ischemic cardiac, brain or peripheral tissues.

The cell material according to the invention can be packaged in unit dose form, each dose containing the material of structure II. As a variant, it is possible to envision a packaging according to which the 5 components of said material II, i.e. EPC-Eph and the protein material L-K, are not in contact; the incubation according to the reaction mechanism:



10

being carried out before administration, in particular for the 10 to 60 minutes (preferably for the 20 to 30 minutes) that precede this administration.

15 In this case, a medicinal composition is recommended, which is characterized in that it comprises a therapeutically acceptable amount of the two components of the proangiogenic system according to the invention, which are packaged separately, each in a 20 physiologically acceptable excipient, said two components being incubated before administration so as to give a cell material of structure II.

As indicated above, the cell material according to the 25 invention is of use for regenerating vascular tissues which have been damaged in particular at the level of the heart or the brain or at the peripheral level. It is in particular suitable for the preparation of a composition for use in the treatment of arteritis, of 30 coronary or cardiac vascular insufficiency and of cerebral vascular insufficiency. On an animal model, it has given good results, firstly, in the treatment of "critical" ischemia of the lower limbs and, secondly, in ensuring recovery making it possible to avoid 35 amputation.

The cell material according to the invention can be administered to mammals, and in particular to humans,

according to a method known per se. For example, said material can be (i) injected into or in the vicinity of the vascular lesion, (ii) injected into the blood via the IV route, or else (iii) brought to the site of the
5 lesion by means of an appropriate vector. As a variant, it is possible to administer separately, by injection (in particular IV injection), firstly, the EPC-Eph cells and, secondly, the protein material bound to a vector known in the gene therapy field.

10

In adult humans, it is possible to administer, by IV injection, cells of structure II contained, where appropriate, in a cell mixture of nonactivated EPCs. Such a cell mixture may contain a total of
15 approximately 10^5 to 10^9 cells per injection.

Other advantages and characteristics of the invention will be understood more clearly on reading the following examples of preparation and of
20 pharmacological trials. Of course, this group of elements is not limiting but is given by way of illustration, the cell material used being: EPC-EphB4-ephrin-B2-Fc.

25 **Example 1**

Obtaining EPC-EphB4

(A) Human umbilical cord blood samples are taken (from
30 30 to 50 ml each) and placed in sterile tubes containing an anticoagulant solution of sodium heparin. The umbilical cord blood mononuclear cells are isolated by density gradient centrifugation using Pancoll (1.077 g/ml, product sold by the company Dominique
35 Dutscher S.A., Brumath, France). The mononuclear cells are then separated from the adherent cells by culturing on plastic dishes for 24 hours at 37°C. A cell mixture containing mononuclear cells expressing the EphB4

marker and mononuclear cells not expressing said marker is obtained.

(B) The cell mixture, obtained at the end of example 5 1(A), is placed in the wells of a 6-well plate coated with collagen type I (product sold by the company Sigma-Aldrich, Saint-Quentin, France) in a culture medium containing hVEGF for differentiation (as defined in the article by Le Ricousse-Roussanne S. et al., 10 *Cardiovasc. Res.*, 2004; 62: 176-184). After culturing for 15 days, a cell mixture enriched in EPC-EphB4 is collected.

Example 2

15

Obtaining EPC-EphB4

(A) Using the cell mixture obtained at the end of example 1(A), the CD34⁺ cells are isolated and purified from the nonadherent cells by a standard immunomagnetic separation technique, in particular by means of the "CD34 isolation kit" (sold by the company Miltenyi Biotech, Paris, France), which comprises an anti-CD34 monoclonal antibody. The analysis of the cells thus obtained by flow cytometry and using an anti-CD34 monoclonal antibody (preferably different from the previous one), coupled to FITC, shows that 75% (\pm 5.6%) of them have the CD34 marker.

30 **(B)** The cell mixture thus obtained, which contains 1.5×10^6 to 3.5×10^6 CD34 $^+$ cells, can be placed in the wells of a 6-well plate coated with a matrix containing fibronectin, laminin, sodium heparin sulfate and collagen type I and IV (products sold by the company 35 Sigma-Aldrich mentioned above) and in a culture medium containing hVEGF, bFGF and IGF1 (products sold by the company R&D Systems Inc., Oxford, United Kingdom).

After culturing for 15 days, a cell mixture enriched in EPC-EphB4 is recovered.

Example 3

5

Obtaining EPC-EphB4-ephrin-B2-Fc

After obtaining the cell mixture according to example 1(B), which contains EPC cells provided with the EphB4 marker, it is treated with 3 µg/ml of fusion protein ephrin-B2-Fc, EphB4-Fc or CD6-Fc (CD6-Fc being involved as the negative control for demonstrating the effect observed by the activation of the EPCs with ephrin-B2) for an incubation period of 30 minutes at 37°C. Each 10 unbound fusion protein is discarded by rinsing (at 15 least two rinses are carried out here).

Three cell mixtures are obtained, one of which contains the cell material according to the invention of 20 structure EPC-EphB4-ephrin-B2-Fc, the second contains the cell material resulting from the activation of EPCs with the EphB4-Fc fusion protein, and the third contains the cell material resulting from the activation of EPCs with the CD6-Fc fusion protein.

25

Example 4

Obtaining EPC-EphB4-ephrin-B2-Fc

30 The process is carried out as indicated in example 3 for obtaining cells of structure EPC-EphB4-ephrin-B2-Fc, with the difference that the starting cell mixture is that obtained at the end of example 2(B). The stated activated cells are obtained.

35

Trial protocol

At the time $T = 0$, the right femoral artery of 7-week-old male nude mice (one batch of 6 animals per trial and per test product, including the PBS, nonactivated EPC and HUVEC controls) is ligatured so as to induce 5 ischemia. At the time $T = + 4.5$ h, the incubation according to example 3 is carried out so as to obtain the cell material of structure EPC-EphB4-ephrin-B2-Fc according to the invention and the other two cell materials for comparison. Next, at the time $T = + 5$ h, 10 each of the three cell mixtures obtained in example 3 is injected intravenously into the retroorbital sinus (10^6 cells/mouse). At the time $T = + 12$ d, the mice are sacrificed and the gastrocnemius muscles of the limb rendered ischemic and of the nonischemic limb are 15 removed. The following are determined:

- the angiographic score;
- the capillary density (i.e. the number of capillaries/ mm^2); and
- the cutaneous blood flow.

20 In figures 2-4, the numerical values obtained are given 25 in the form: mean \pm SEM.

Trial 1

Angiographic score

30 The angiographic score (i.e. vessel density) was determined by microangiography. More specifically, the vessel density in the limb rendered ischemic, compared with the nonischemic limb, is measured by high-35 definition microangiography (see operating methods in the article by Silvestre J. S. et al., *Cir. Res.*, 2001; 89: 259-264). The results obtained, given in the form

of a ratio (ischemic limb /nonischemic limb), are reported in the form of a graph in figure 2.

It is observed that the injection of the EPCs that are
5 nonactivated or have been activated with the EphB4-Fc or CD6-Fc fusion protein slightly increase, and at comparable levels, the vessel density compared with the control group having received PBS. When the EPC cells are activated with the ephrin-B2-Fc fusion protein, it
10 is noted that the increase in vessel density is 25.5% greater than that observed after injection of the nonactivated EPCs, i.e. a 1.34 times increase in the angiographic score.

15 **Trial 2**

Capillary density (number of capillaries/mm²)

The capillary density of the muscle rendered ischemic
20 was studied by labeling sections of gastrocnemius muscle with an antibody directed against the CD31 marker, which is specific for endothelial cells, compared with sections of the same muscle of the nonischemic limb. The results obtained, presented in
25 the form of a ratio (ischemic limb/nonischemic limb) are reported in the form of a graph in figure 3.

It is observed that the capillary density is 36.7% greater when the mice were treated by injection with
30 the EPCs activated with the ephrin-B2-Fc fusion protein according to the invention, compared with the nonactivated EPCs, i.e. a 1.57 times increase in the capillary density.

35 **Trial 3**

Cutaneous blood flow

A quantitative evaluation of the blood flow, expressed by the ratio of the ischemic limb/nonischemic limb blood flow, was also carried out in order to verify that the variation in vessel number corresponds to a 5 functional adaptation and therefore to a variation in the perfusion of the limb rendered ischemic. The results obtained are reported in the form of a graph in figure 4.

10 It is observed that the injection of the EPCs activated according to the invention with the ephrin-B2-Fc fusion protein increases the ratio of the blood flow in the limb rendered ischemic to the blood flow in the nonischemic limb by 1.37 times (27.1% increase).
15

Example 5

Role of the EphB4 marker

20 In this last example, the role of the EphB4 marker was demonstrated by virtue of cell preparations in which the EphB4 marker protein synthesis was inhibited.

25 For this, use was made of "interfering RNAs", or "siRNAs" capable in particular of specifically degrading the messenger RNAs encoding a given gene, and in particular in this case encoding the gene expressing the EphB4 marker. The specific action of these interfering RNAs is called transfection.
30

These cell preparations were administered to male nude mice according to a trial protocol similar to the trial protocol mentioned above.

35 In the first step, endothelial progenitor cells, or EPCs, are cultured to 80% confluence. Solutions of EphB4 siRNA, i.e. containing the interfering RNA, capable of inhibiting EphB4 marker protein synthesis

are diluted in an M199 medium containing neither antibiotics nor serum, and incubated for five minutes at ambient temperature. Moreover, a solution of "control siRNAs" not corresponding to any particular 5 gene is diluted under the same conditions. The latter diluted solution, which thus has no biological effect on the expression of the EphB4 marker, will simply make it possible to verify that the interfering RNAs have no activity of their own independently of their role of 10 inhibitor.

Moreover, a transfection agent, Dharmafect2 (Dharmacon, Perbio) is prepared under the same conditions as the 15 abovementioned solutions, and then mixed with these solutions, which are then subsequently incubated.

These incubated solutions are then brought into contact with the endothelial progenitor cells, EPCs, so as, in a first cell preparation, to cause inhibition of the 20 EphB4 marker protein synthesis, and in a second cell preparation, where the expression of the gene encoding the EphB4 marker is not inhibited, to constitute a control.

25 These two cell preparations will, in turn, be respectively divided into two, and one of them will be stimulated with the ephrin-B2-Fc fusion protein before being injected intravenously into mice having been subjected to a ligature of the right femoral artery. In 30 this way, four different cell preparations will be administered to the mice.

A summary of the results of this example 5 will be found in table I below.

Table I

No.	PBS	1	2	3	4
Angiographic score	100	161.5 +/- 10.7	218.8 +/- 12.8	153.5 +/- 2.9	158.9 +/- 12.5
Blood flow	100	144.5 +/- 4.5	193.5 +/- 4.7	151.1 +/- 8	138.5 +/- 14.1
Capillary density	100	147.4 +/- 10.7	201.7 +/- 12.8	142.6 +/- 11	140.8 +/- 11.3

1: EPCs transfected with the control siRNA and not
5 stimulated before injection;

2: EPCs transfected with the control siRNA and
stimulated with ephrin-B2-Fc before injection;

10 3: EPCs transfected with the EphB4 siRNA and not
stimulated before injection;

4: EPCs transfected with the EphB4 siRNA and stimulated
with ephrin-B2-Fc before injection.

15 It will be noted here again that the measurements of
the angiographic score, of the blood flow and of the
capillary density for the four cell preparations
administered evolve substantially in parallel.

20 Moreover, it is immediately apparent that the result of
the measurements for cell preparations no. 1, 3 and 4
are substantially identical, whereas the result for
cell preparation no. 2 is greater by close to 40%.

25 Since the hypothesis is, of course, that the
proangiogenic activity is promoted by the association
of the EphB4 marker and the ephrin-B2-Fc protein
material, it is noted, upon reading table I, by
30 comparing the results of cell preparations no. 1 and 4,
that the effect of the endothelial progenitor cells

18 Jan 2012
2006273928

incorporating the "control siRNA" but not associated with the ephrin-B2-Fc protein material is substantially equivalent to the effect of the endothelial progenitor cells in which the expression of the EphB4 marker is inhibited but which are associated with the ephrin-B2-Fc protein material. Thus, it is shown that the interfering RNAs have no activity of their own independently of their role of inhibitor and that the endothelial progenitor cells, EPCs, expressing the EphB4 marker have no more activity than the endothelial progenitor cells, EPCs, not expressing the marker and which are associated with the ephrin-B2-Fc protein material.

Moreover, by comparing the results of the measurements of cell preparations no. 1 and 3, it is shown that the endothelial progenitor cells, EPCs, alone, expressing the EphB4 marker or not expressing it, have a comparable activity.

In addition, and this is the most important aspect, it is clearly shown, in view of the results of cell preparation no. 2 and in comparison with any of the other preparations, that the specific association of the EphB4 marker and of the ephrin-B2-Fc protein material exhibits considerable proangiogenic activity.

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.

18 Jan 2012

2006273928

- 23a -

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

Table II

Specificities for some Eph/ephrin couples

Eph receptors	Ephrin ligands (by decreasing affinity)
EphA1	ephrin-A1
EphA2	ephrin-A3, -A1, -A5, -A4
EphA3	ephrin-A5, -A2, -A3, -A1
EphA4	ephrin-A5, -A1, -A3, -A2, -B2, -B3
EphA5	ephrin-A5, -A1, -A2, -A3, -A4
EphA6	ephrin-A2, -A1, -A3, -A4, -A5
EphA7	ephrin-A2, -A3, -A1
EphA8	ephrin-A5, -A3, -A2
EphB1	ephrin-B2, -B1, -A3
EphB2	ephrin-B1, -B2, -B3
EphB3	ephrin-B1, -B2, -B3
EphB4	ephrin-B2, -B1

18 Jan 2012

2006273928

- 25 -

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A cell marker/specific ligand proangiogenic system, wherein said system, in which the cell marker is present on the outer membrane of the cell, comprises:

(a) an endothelial precursor cell (EPC) comprising a cell marker selected from the group consisting of the Ephs, and

(b) a protein material of structure:

L-K (I)

which consists of a ligand (L) specific for said marker and is associated or fused with a binding protein (K),

said system being capable of providing a cell material of structure:

EPC-Eph-L-K (II)

which stimulates angiogenesis.

2. A cell material capable of stimulating angiogenesis, wherein said cell material has the structure:

EPC-Eph-L-K (II)

wherein said ligand L is associated with or fused with a binding protein K, said cell material being capable

18 Jan 2012

2006273928

- 26 -

(i) of being in the form of a substantially purified cell culture or in the form of a cell culture in combination with other precursor cells, in particular mononuclear cells, and (ii) of being, where appropriate, frozen.

3. The cell material according to claim 2, which is obtained by incubating:

(a) an endothelial precursor cell (EPC) comprising a cell marker selected from the group consisting of the Ephs, with

(b) a protein material of structure:

L-K (I)

which consists of a ligand (L) specific for said marker, and is associated or fused with a binding protein (K),

before being brought to its site of administration.

4. The cell marker/specific ligand proangiogenic system or the cell material according to claim 2, wherein the Eph is EphB4 or EphB1.

5. The cell marker/specific ligand proangiogenic system or the cell material according to claim 2, wherein the binding protein (K) is an antibody Fc fragment A.

6. The system according to any one of claims 1, 4 or 5,

- 27 -

wherein said Eph marker is in the form of an amino acid sequence.

7. The system according to any one of claims 1 or 4 to 6, wherein said Eph marker is an EphB.
8. The system according to any one of claims 1 or 4 to 7, or the cell material according to claim 2 or claim 3, wherein said marker is EphB4 or EphB1, and said ligand is ephrin-B2 or ephrin-B1.
9. The system according to any one of claims 1 or 4 to 8, or the cell material according to any one of claims 2, 3 or 8, wherein said ligand (L) is a peptide fragment of ephrin.
10. The system according to claim 1, wherein said protein material L-K is replaced with the recombinant ephrin-B2 protein.
11. The system according to any one of claims 1 or 4 to 10, wherein the EPCs are obtained from mononuclear cells or from cells expressing CD34 or CD133, which originate from the bone marrow, from peripheral blood or from umbilical cord blood.
12. The system according to any one of claims 1 or 4 to 11, wherein the EPCs are obtained from peripheral blood or from umbilical cord blood.
13. A method for preparing a cell material as claimed in either one of claims 11 and 12, said method comprising

- 28 -

the steps consisting of:

- (i) making use of EPCs expressing on their outer membrane a marker of the Eph family, in particular EphB4 or EphB1, and
- (ii) bringing said EPCs into contact, *in vitro*, with a protein material L-K wherein L is a ligand specific for said marker and K is a binding protein associated or fused with L.

14. The method according to claim 13, wherein said method comprises the following steps:

- (1°) making use of mononuclear cells originating from the bone marrow, peripheral blood or umbilical cord blood,
- (2°) differentiating said cells of the preceding step in order to obtain EPCs having the Eph marker, and
- (3°) activating the EPCs thus obtained in step (2°), *in vitro*, by binding of ephrin to Eph.

15. The method according to claim 14, wherein said method comprises, between steps (1) and (2), an additional step:

- (1a) isolating the CD34⁺ and/or CD133⁺ mononuclear cells.

16. The method according to claim 14, wherein in the step

- 29 -

(1°), the mononuclear cells originate from peripheral blood or from umbilical cord blood.

17. The method according to any one of claims 13 to 16, wherein said ligand L is associated with or fused with a binding protein K in order to provide a cell material of structure:

EPC-Eph-L-K (II)

where K is Fc.

18. The method according to any one of claims 13 to 17, wherein the marker is EphB4 or EphB1, and in that said ligand L is ephrin-B2 or ephrin-B1.
19. The method according to any one of claims 13 to 18, wherein said ligand is a peptide fragment of ephrin.
20. A medicament that can be used for reconstituting damaged vessels, wherein said medicament comprises, in combination with a physiologically acceptable excipient, a therapeutically acceptable amount of a cell material as claimed in any one of claims 2, 3, 8 or 9.
21. The medicament according to claim 20, wherein said cell material is a proangiogenic active ingredient.
22. A medicinal composition for reconstituting damaged vessels, wherein said composition comprises a therapeutically acceptable amount of the two components

18 Jan 2012
2006273928

- 30 -

of a proangiogenic system as claimed in any one of claims 1 or 4 to 12, which are packaged separately, each in a physiologically acceptable excipient, said two components being incubated before administration so as to give a cell material of structure II.

23. The use of a cell material, wherein said use is made of a cell material as claimed in any one of claims 2, 3, 8 or 9 as proangiogenic active ingredient, in combination with a physiologically acceptable excipient, for the preparation of a composition for therapeutic use in the treatment of vascular insufficiencies.

24. The use according to claim 23, wherein said therapeutic use is the revascularization of ischemic cardiac, brain or peripheral tissues.

25. The use as claimed in claim 23 or claim 24, said composition is for use in the treatment of arteritis, of coronary insufficiency, of cardiac insufficiency or of cerebral insufficiency.

26. A cell marker/specific ligand proangiogenic system according to any one of claims 1 or 4 to 12, or the cell material according to any one of claims 3, 4, 8 or 9, or a method according to any one of claims 13 to 19, or a medicament according to claim 20 or 21, or a medicinal composition according to claim 22, or use according to any one of claims 23 to 25 substantially as hereinbefore defined.

Fig. 1





Fig. 2




Fig. 3




Fig. 4