



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION
(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2018/05/18
(87) Date publication PCT/PCT Publication Date: 2018/11/22
(85) Entrée phase nationale/National Entry: 2019/11/14
(86) N° demande PCT/PCT Application No.: US 2018/033437
(87) N° publication PCT/PCT Publication No.: 2018/213732
(30) Priorités/Priorities: 2017/05/18 (US62/508,363);
2017/05/19 (US62/508,481); 2017/10/27 (US62/578,204)

(51) Cl.Int./Int.Cl. *A61K 45/06* (2006.01),
A61K 31/4439 (2006.01), *A61K 31/454* (2006.01),
A61P 35/00 (2006.01)
(71) Demandeur/Applicant:
TESARO, INC., US
(72) Inventeurs/Inventors:
SUN, KAIMING, US;
WANG, JING YU, US;
WANG, ZEBIN, US
(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : POLYTHERAPIES POUR LE TRAITEMENT DU CANCER
(54) Title: COMBINATION THERAPIES FOR TREATING CANCER




FIG. 6A

(57) Abrégé/Abstract:
The present invention provides methods of treatment for cancer(s) through combination therapy with an agent that inhibits poly [ADP-ribose] polymerase (PARP) signaling and an agent that regulates activity within the tumor microenvironment.

- (51) International Patent Classification:

A61K 45/06 (2006.01)

A61K 31/454 (2006.01)

A61K 31/4439 (2006.01)

A61P 35/00 (2006.01)
- (21) International Application Number:

PCT/US2018/033437
- (22) International Filing Date:

18 May 2018 (18.05.2018)
- (25) Filing Language:

English
- (26) Publication Language:

English
- (30) Priority Data:

62/508,363

18 May 2017 (18.05.2017)

US

62/508,481

19 May 2017 (19.05.2017)

US

62/578,204

27 October 2017 (27.10.2017)

US
- (71) Applicant: TESARO, INC. [US/US]; 1000 Winter Street North, Suite 3300, Waltham, MA 02451 (US).
- (72) Inventors: SUN, Kaiming; c/o Tesaro, Inc., 1000 Winter Street North, Suite 3300, Waltham, MA 02451 (US). WANG, Jing, Yu; c/o Tesaro, Inc., 1000 Winter Street North, Suite 3300, Waltham, MA 02451 (US). WANG, Ze-
- bin; c/o Tesaro, Inc., 1000 Winter Street North, Suite 3300, Waltham, MA 02451 (US).
- (74) Agent: ESPINO, Christine, G. et al.; Proskauer Rose LLP, One International Place, Boston, MA 02110 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) Title: COMBINATION THERAPIES FOR TREATING CANCER




FIG. 6A

(57) Abstract: The present invention provides methods of treatment for cancer(s) through combination therapy with an agent that inhibits poly [ADP-ribose] polymerase (PARP) signaling and an agent that regulates activity within the tumor microenvironment.

WO 2018/213732 A1

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

COMBINATION THERAPIES FOR TREATING CANCER**CROSS-REFERENCE TO RELATED APPLICATIONS**

[0001] The present application claims benefit of U.S. Provisional Application No. 62/508,363, filed May 18, 2017; U.S. Provisional Application No. 62/508,481, filed May 19, 2017; and U.S. Provisional Application No. 62/578,204, filed October 27, 2017, each of which is hereby incorporated by reference in its entirety.

BACKGROUND

[0002] Cancer is a serious public health problem, with about 595,690 people in the United States of America expected to die of cancer in 2016 alone according to the American Cancer Society, Cancer Facts & Figures 2016 (<http://www.cancer.org/acs/groups/content/@research/documents/document/acspc-047079.pdf>).

SUMMARY

[0003] Provided herein are methods of treating a subject with a disease or condition comprising administering to the subject (a) a first agent that inhibits poly [ADP-ribose] polymerase (PARP); and (b) a second agent, wherein the second agent comprises a regulatory T cell (Treg) inhibitory agent, a macrophage inhibitory agent, an antigen specific immune response enhancer agent, or a combination thereof.

[0004] Provided herein are methods of enhancing an immune response or increasing the activity of an immune cell in a subject with a disease or condition comprising administering to the subject (a) a first agent that inhibits poly [ADP-ribose] polymerase (PARP); and (b) a second agent, wherein the second agent comprises a regulatory T cell (Treg) inhibitory agent, a macrophage inhibitory agent, an antigen specific immune response enhancer agent, or a combination thereof. In embodiments, the method enhances an anti-tumor response in the subject.

[0005] Provided herein are methods of inducing an immune response in a subject with a disease or condition comprising administering to the subject (a) a first agent that inhibits poly [ADP-ribose] polymerase (PARP); and (b) a second agent, wherein the second agent comprises a regulatory T cell (Treg) inhibitory agent, a macrophage inhibitory agent, an antigen specific immune response enhancer agent, or a combination thereof. In embodiments, the method induces an anti-tumor response in the subject.

[0006] In some embodiments, the first agent inhibits PARP 1 and/or 2. In some embodiments, a first agent that inhibits PARP 1 and/or 2 is selected from the group consisting of: ABT-767, AZD 2461, BGB-290, BGP 15, CEP 8983, CEP 9722, DR 2313, E7016, E7449, fluzoparib (SHR 3162), IMP 4297, INO1001, JPI 289, JPI 547, monoclonal antibody B3-LysPE40 conjugate, MP 124, niraparib (ZEJULA) (MK-4827), NU 1025, NU 1064, NU 1076, NU1085, olaparib (AZD2281), ONO2231, PD 128763, R 503, R554, rucaparib (RUBRACA) (AG-014699, PF-01367338), SBP 101, SC 101914, Simmiparib, talazoparib (BMN-673), veliparib (ABT-888), WW 46, 2-(4-(Trifluoromethyl)phenyl)-7,8-dihydro-5H-thiopyrano[4,3-d]pyrimidin-4-ol, and salts or derivatives thereof. In some embodiments, the first agent is a small organic or inorganic molecule; a saccharine; an oligosaccharide; a polysaccharide; a carbohydrate; a peptide; a protein; a peptide analog; a peptide derivative; a lipid; an antibody; an antibody fragment; a peptidomimetic; a nucleic acid; a nucleic acid analog; a nucleic acid derivative; an extract made from biological materials; a naturally occurring or synthetic composition; a metal; a toxin; or any combination thereof. In some embodiments, the first agent is a small molecule.

[0007] In some cases, the first agent is selected from the group consisting of: niraparib, olaparib, rucaparib, talazoparib, and veliparib, or salts or derivatives thereof. In some cases, the first agent comprises niraparib or a pharmaceutically acceptable salt or derivative thereof. In some instances the Treg inhibitory agent inhibits or decreases the activity, function, or migration of a Treg cell. In some cases, the Treg inhibitory agent decreases a population of Treg cells in the subject. In some embodiments, the Treg inhibitory agent substantially ablates or eliminates a population of Treg cells in the subject. In some embodiments, the macrophage inhibitory agent inhibits or decreases the activity, function, or migration of a macrophage. In some embodiments, the macrophage inhibitory agent decreases a population of macrophage cells in the subject. In some embodiments, the macrophage inhibitory agent substantially ablates or eliminates a population of macrophage cells in the subject. In some embodiments, the Treg cell is an infiltrating T cell.

[0008] In some embodiments, the macrophage comprises a tumor-associated macrophage (TAM). In some embodiments, the second agent enhances an antigen specific CD4⁺ T cell activity. In some embodiments, the second agent enhances an antigen specific CD8⁺ T cell activity. In some embodiments, the second agent is selected from the group consisting of a small organic or inorganic molecule; a saccharine; an oligosaccharide; a

polysaccharide; a carbohydrate; a peptide; a protein; a peptide analog; a peptide derivative; a lipid; an antibody; an antibody fragment; a peptidomimetic; a nucleic acid; a nucleic acid analog; a nucleic acid derivative; an extract made from biological materials; a naturally occurring or synthetic composition; a metal; a toxin; and any combination thereof.

[0009] In some embodiments, the administering comprises administering the first and second agent sequentially. In some embodiments, the administering comprises administering the first and second agent simultaneously. In some embodiments, the administering comprises administering the first agent before administering the second agent second agent. In some embodiments, the subject is a mammalian subject. In some embodiments, the subject is a human. In some embodiments, the second agent is a regulatory T cell (Treg) inhibitory agent selected from the group consisting of a Treg ablating agent, a Treg migration inhibitor agent, a Treg function inhibitor agent, and combinations thereof. In some embodiments, the Treg ablating agent is selected from the group consisting of cyclophosphamide, paclitaxel, imatinib, sunitinib, sorafenib, dasatinib, temozolomide, daclizumab, denileukin diftitox, and combinations thereof. In some embodiments, the Treg migration inhibitor agent is selected from the group consisting of AMD3100, mogamulizumab, casuarinin, fucoidan, and combinations thereof. In some embodiments, the Treg function inhibitor agent is selected from the group consisting of an anti-CTLA4 agent (e.g., ipilimumab, tremelimumab), an anti-OX40 agent, an anti-GITR agent, an adenosine receptor antagonist (e.g., caffeine, theophylline, theobromine, and 8-phenylxanthines), P60, and combinations thereof.

[0010] In some embodiments, the second agent is a macrophage inhibitory agent selected from the group consisting of a macrophage recruitment inhibitory agent, an M2 macrophage antisurvival agent, an M1 macrophage enhancing agent, an M2 to M1 polarizing agent, a macrophage activity inhibitor agent and combinations thereof. In some embodiments, the macrophage recruitment inhibitory agent is selected from the group consisting of an anti-CCL2/CCR2 agent, an anti-IL6 agent, an anti-M-CSFR agent, and combinations thereof. In some embodiments, a macrophage recruitment inhibitory agent is an anti-M-CSFR agent. In some embodiments, the macrophage recruitment inhibitory agent is selected from the group consisting of trabectedin, RS102895, PF-04136309, CNTO888, MLN1202, siltuximab, JNJ-28312141, GW2580, IMC-CS4 (LY3022855), emactuzumab, AMG820, pexidartinib, linifanib, OSI-930, CEP-32496, PLX7846, ARRY-382, JNJ-40346527, MCS110, PLX3397, PLX6134, PD-0360324, FPA008, and combinations thereof. In some embodiments, a

macrophage recruitment inhibitory agent is BLZ945, PLX7846, GW2580, ARRY-382, JNJ-40346527, emactuzumab, pexidartinib, AMG820, IMC-CS4 (LY3022855), MCS110, PLX3397, PLX6134, PD-0360324, or FPA008. In some embodiments, a macrophage recruitment inhibitory agent is BLZ945. In some embodiments, the M2 macrophage antisurvival agent is selected from the group consisting of an MMP inhibitor, clodronate, zoledronic acid, dichloromethylene bisphosphonate, trabectedin, dasatinib, retinoic acid, attenuated bacteria (e.g., *Shigella flexneri*, *Salmonella typhimurium*, *Listeria monocytogens*, *Chlamydia psittaci*, *Legionella pneumophila*), and combinations thereof.

[0011] In some embodiments, the M1 macrophage enhancing agent or the M2 to M1 polarizing agent is selected from the group consisting of an anti-CD40 agent, an anti-IL-10R agent, a CD47 antagonist (e.g., Hu5F9-G4, CC-90002, and CD47-Fc fusion protein TTI-621), PolyI:C, LPS, monophosphoryl A, imiquimod, R-848, CpG-ODN, IFN- α , IFN- β , IFN- γ , GM-CSF, IL-12, IL-2, IL-15, T α 1, ibrutinib, EF-022 and combinations thereof. In some embodiments, the macrophage activity inhibitory agent is selected from the group consisting of a STAT3 inhibitor, a STAT6 inhibitor, or an anti-tumor drug agent. In some embodiments, the macrophage activity inhibitory agent is selected from the group consisting of WP1066, sunitinib, sorafenib, STA-21, IS3 295, S3I-M2001, AS1517499, leflunomide, TMC-264, histidine-rich glycoprotein (HRG), copper chelate (CuNG), 5,6-dimethylxanthenone-4-acetic acid (MDXAA), vadimezan (ASA404), cisplatin, silibinin, proton pump inhibitor pantoprazole (PPZ), CNI-1493 and combinations thereof. In some embodiments, the macrophage inhibitor agent is an anti-IL-1 α agent (e.g., xilonix).

[0012] In some embodiments, the second agent is an antigen specific immune response enhancer agent selected from the group consisting of an anti-PD-1 agent, an anti-PD-L1 agent, a GITR (glucocorticoid-induced TNFR-related protein) stimulating agent, an anti-CTLA4 agent, an anti-TIM-3 agent, an anti-LAG-3 agent, an agent that enhances tumor antigen presentation (e.g., personalized cancer vaccine, autologous antigen presenting cell, autologous dendritic cells, artificial antigen presenting cell), a chemokine signaling agent, an anti-VEGF agent, a cytokine signal stimulating agent, and combinations thereof. In some embodiments, the anti-PD-1 agent is selected from the group consisting of pembrolizumab, nivolumab, PDR001, REGN2810 (SAR-439684), BGB-A317, BI 754091, IBI308, INCSHR-1210, JNJ-63723283, JS-001, MEDI0680 (AMP-514), MGA-012, PF-06801591, REGN-2810, TSR-042, PDR-001, camrelizumab (HR-301210), BCD-100, AGEN-2034, CS1001, Sym-021, LZM009, KN-035, AB122, genolimzumab (CBT-501), AK 104, GLS-010, and

combinations thereof. In some embodiments, the anti-PD-L1 agent is selected from the group consisting of atezolizumab, durvalumab, avelumab, LY3300054, BGB-A333, SHR-1316, CK-301, and combinations thereof. In some embodiments, the GITR stimulating agent is selected from the group consisting of DTA-1, mGITRL, pGITRL, and combinations thereof. In some embodiments, the anti-CTLA4 agent is selected from the group consisting of ipilimumab, tremelimumab, and combinations thereof. In some embodiments, the chemokine signaling agent is selected from the group consisting of CXCL16, a CXCR6 chemokine receptor (CD186) agonist, and combinations thereof. In some embodiments, the anti-VEGF agent is selected from the group consisting of bevacizumab, pazopanib, sunitinib, sorafenib, axitinib, ponatinib, regorafenib, cabozantinib, vandetanib, ramucirumab, lenvatinib, ziv-aflibercept, and combinations thereof. In some embodiments, the cytokine signal stimulating agent is an interleukin or an interferon. In some embodiments, the interleukin is selected from the group consisting of IL-2, IL-1, IL-7, IL-15, IL-12, IL-18, and combinations thereof. In some embodiments, the interferon is IFN alpha.

[0013] In some embodiments, the second agent is an antigen specific immune response enhancer agent selected from the group consisting of a flavonoid (e.g., flavonoid glycoside), lidocaine, lamotrigine, sulfamethoxazole, phenytoin, carbamazepine, sulfamethoxazole, phenytoin, allopurinol, paracetamol, mepivacaine, p-phenylenediamine, ciprofloxacin and moxifloxacin. In some embodiments, the disease or condition is cancer. In some embodiments, the cancer is selected from the group consisting of ovarian cancer, breast cancer, cervical cancer, endometrial cancer, prostate cancer, testicular cancer, pancreatic cancer, esophageal cancer, head and neck cancer, gastric cancer, bladder cancer, lung cancer, bone cancer, colon cancer, rectal cancer, thyroid cancer, brain and central nervous system cancers, glioblastoma, neuroblastoma, neuroendocrine cancer, rhabdoid cancer, keratoacanthoma, epidermoid carcinoma, seminoma, melanoma, sarcoma, bladder cancer, liver cancer, kidney cancer, myeloma, lymphoma, and combinations thereof.

[0014] In some embodiments, the administering comprises administering a composition comprising a capsule comprising the first agent. In some embodiments, the capsule comprises a formulation comprising the first agent and one or more pharmaceutically acceptable excipients. In some embodiments, the one or more pharmaceutically acceptable excipients comprises lactose monohydrate, magnesium stearate, or a combination thereof. In some embodiments, a therapeutically effective amount of the first or second agent is administered. In some embodiments, the method further comprises administering a third

agent to the subject. In some embodiments, the third agent comprises an antigen specific immune response enhancer agent, an anti-angiogenic agent, a chemotherapeutic agent, or combinations thereof. In some embodiments, the antigen specific immune response enhancer agent comprises an anti-PD-1 agent, an anti-PD-L1 agent, an anti-CTLA4 agent, an anti-TIM-3 agent, or an anti-LAG-3 agent. In some embodiments, the anti-angiogenic agent is selected from the group consisting of TNP-470, platelet factor 4, thrombospondin-1, tissue inhibitors of metalloproteases (TIMP1 and TIMP2), prolactin, angiostatin, endostatin, bFGF soluble receptor, transforming growth factor beta, interferon alpha, soluble KDR and FLT-1 receptors, placental proliferin-related protein, and combinations thereof.

[0015] In some embodiments, the chemotherapeutic agent is selected from the group consisting of aminoglutethimide, amsacrine, anastrozole, asparaginase, bcg, bicalutamide, bleomycin, buserelin, busulfan, camptothecin, capecitabine, carboplatin, carmustine, chlorambucil, cisplatin, cladribine, clodronate, colchicine, cyclophosphamide, cyproterone, cytarabine, dacarbazine, dactinomycin, daunorubicin, dienestrol, diethylstilbestrol, docetaxel, doxorubicin, epirubicin, estradiol, estramustine, etoposide, exemestane, filgrastim, fludarabine, fludrocortisone, fluorouracil, fluoxymesterone, flutamide, gemcitabine, genistein, goserelin, hydroxyurea, idarubicin, ifosfamide, imatinib, interferon, irinotecan, irinotecan, letrozole, leucovorin, leuprolide, levamisole, lomustine, mechlorethamine, medroxyprogesterone, megestrol, melphalan, mercaptopurine, mesna, methotrexate, mitomycin, mitotane, mitoxantrone, nilutamide, nocodazole, octreotide, oxaliplatin, paclitaxel, pamidronate, pentostatin, plicamycin, porfimer, procarbazine, raltitrexed, rituximab, streptozocin, suramin, tamoxifen, temozolomide, teniposide, testosterone, thioguanine, thiotepa, titanocene dichloride, topotecan, trastuzumab, tretinoin, vinblastine, vincristine, vindesine, vinorelbine, and combinations thereof.

[0016] Provided herein are pharmaceutical compositions comprising (a) a first agent that inhibits poly [ADP-ribose] polymerase (PARP); and (b) a second agent, wherein the second agent comprises a regulatory T cell (Treg) inhibitory agent, a macrophage inhibitory agent, an antigen specific immune response enhancer agent, or a combination thereof.

[0017] In some embodiments, the first agent is an agent that inhibits PARP 1 and/or 2. In some embodiments, a first agent that inhibits PARP 1 and/or 2 is selected from the group consisting of: ABT-767, AZD 2461, BGB-290, BGP 15, CEP 8983, CEP 9722, DR 2313, E7016, E7449, fluzoparib (SHR 3162), IMP 4297, INO1001, JPI 289, JPI 547, monoclonal antibody B3-LysPE40 conjugate, MP 124, niraparib (ZEJULA) (MK-4827), NU 1025, NU

1064, NU 1076, NU1085, olaparib (AZD2281), ONO2231, PD 128763, R 503, R554, rucaparib (RUBRACA) (AG-014699, PF-01367338), SBP 101, SC 101914, Simmiparib, talazoparib (BMN-673), veliparib (ABT-888), WW 46, 2-(4-(Trifluoromethyl)phenyl)-7,8-dihydro-5H-thiopyrano[4,3-d]pyrimidin-4-ol, and salts or derivatives thereof. In some embodiments, the first agent is a small organic or inorganic molecule; a saccharine; an oligosaccharide; a polysaccharide; a carbohydrate; a peptide; a protein; a peptide analog; a peptide derivative; a lipid; an antibody; an antibody fragment; a peptidomimetic; a nucleic acid; a nucleic acid analog; a nucleic acid derivative; an extract made from biological materials; a naturally occurring or synthetic composition; a metal; a toxin; or any combination thereof. In some embodiments, the first agent is a small molecule. In some cases, the first agent is selected from the group consisting of: niraparib, olaparib, rucaparib, talazoparib, and veliparib, or salts or derivatives thereof. In some cases, the first agent comprises niraparib or a pharmaceutically acceptable salt or derivative thereof.

[0018] In some instances the Treg inhibitory agent inhibits or decreases the activity, function, or migration of a Treg cell. In some cases, the Treg inhibitory agent decreases a population of Treg cells in the subject. In some embodiments, the Treg inhibitory agent substantially ablates or eliminates a population of Treg cells in the subject. In some embodiments, the macrophage inhibitory agent inhibits or decreases the activity, function, or migration of a macrophage. In some embodiments, the macrophage inhibitory agent decreases a population of macrophage cells in the subject. In some embodiments, the macrophage inhibitory agent substantially ablates or eliminates a population of macrophage cells in the subject. In some embodiments, the Treg cell is an infiltrating T cell.

[0019] In some embodiments, the macrophage comprises a tumor-associated macrophage (TAM). In some embodiments, the second agent enhances an antigen specific CD4⁺ T cell activity. In some embodiments, the second agent enhances an antigen specific CD8⁺ T cell activity. In some embodiments, the second agent is selected from the group consisting of a small organic or inorganic molecule; a saccharine; an oligosaccharide; a polysaccharide; a carbohydrate; a peptide; a protein; a peptide analog; a peptide derivative; a lipid; an antibody; an antibody fragment; a peptidomimetic; a nucleic acid; a nucleic acid analog; a nucleic acid derivative; an extract made from biological materials; a naturally occurring or synthetic composition; a metal; a toxin; and any combination thereof.

[0020] In some embodiments, the pharmaceutical composition is administered to a subject and the administering comprises administering the first and second agent sequentially.

In some embodiments, the administering comprises administering the first and second agent simultaneously. In some embodiments, the administering comprises administering the first agent before administering the second agent. In some embodiments, the subject is a mammalian subject. In some embodiments, the subject is a human. In some embodiments, the second agent is a regulatory T cell (Treg) inhibitory agent selected from the group consisting of a Treg ablating agent, a Treg migration inhibitor agent, a Treg function inhibitor agent, and combinations thereof. In some embodiments, the Treg ablating agent is selected from the group consisting of cyclophosphamide, paclitaxel, imatinib, sunitinib, sorafenib, dasatinib, temozolomide, daclizumab, denileukin diftitox, and combinations thereof. In some embodiments, the Treg migration inhibitor agent is selected from the group consisting of AMD3100, mogamulizumab, casuarinin, fucoidan, and combinations thereof. In some embodiments, the Treg function inhibitor agent is selected from the group consisting of an anti-CTLA4 agent (e.g., ipilimumab, tremelimumab), an anti-OX40 agent, an anti-GITR agent, an adenosine receptor antagonist (e.g., caffeine, theophylline, theobromine, and 8-phenylxanthines), P60, and combinations thereof.

[0021] In some embodiments, the second agent is a macrophage inhibitory agent selected from the group consisting of a macrophage recruitment inhibitory agent, an M2 macrophage antisurvival agent, an M1 macrophage enhancing agent, an M2 to M1 polarizing agent, a macrophage activity inhibitor agent and combinations thereof. In some embodiments, the macrophage recruitment inhibitory agent is selected from the group consisting of an anti-CCL2/CCR2 agent, an anti-IL6 agent, an anti-M-CSFR agent, and combinations thereof. In some embodiments, a macrophage recruitment inhibitory agent is an anti-M-CSFR agent. In some embodiments, the macrophage recruitment inhibitory agent is selected from the group consisting of trabectedin, RS102895, PF-04136309, CNTO888, MLN1202, siltuximab, JNJ-28312141, GW2580, IMC-CS4 (LY3022855), emactuzumab, AMG820, pexidartinib, linifanib, OSI-930, CEP-32496, PLX7846, ARRY-382, JNJ-40346527, MCS110, PLX3397, PLX6134, PD-0360324, FPA008, and combinations thereof. In some embodiments, a macrophage recruitment inhibitory agent is BLZ945, PLX7846, GW2580, ARRY-382, JNJ-40346527, emactuzumab, pexidartinib, AMG820, IMC-CS4 (LY3022855), MCS110, PLX3397, PLX6134, PD-0360324, or FPA008. In some embodiments, a macrophage recruitment inhibitory agent is BLZ945. In some embodiments, the M2 macrophage antisurvival agent is selected from the group consisting of an MMP inhibitor, clodronate, zoledronic acid, dichloromethylene bisphosphonate, trabectedin, dasatinib, retinoic acid,

attenuated bacteria (e.g., *Shigella flexneri*, *Salmonella typhimurium*, *Listeria monocytogens*, *Chlamydia psittaci*, *Legionella pneumophila*), and combinations thereof.

[0022] In some embodiments, the M1 macrophage enhancing agent or the M2 to M1 polarizing agent is selected from the group consisting of an anti-CD40 agent, an anti-IL-10R agent, a CD47 antagonist (e.g., Hu5F9-G4, CC-90002, and CD47-Fc fusion protein TTI-621), PolyI:C, LPS, monophosphoryl A, imiquimod, R-848, CpG-ODN, IFN- α , IFN- β , IFN- γ , GM-CSF, IL-12, IL-2, IL-15, T α 1, ibrutinib, EF-022 and combinations thereof. In some embodiments, the macrophage activity inhibitory agent is selected from the group consisting of a STAT3 inhibitor, a STAT6 inhibitor, or an anti-tumor drug agent. In some embodiments, the macrophage activity inhibitory agent is selected from the group consisting of WP1066, sunitinib, sorafenib, STA-21, IS3 295, S3I-M2001, AS1517499, leflunomide, TMC-264, histidine-rich glycoprotein (HRG), copper chelate (CuNG), 5,6-dimethylxanthenone-4-acetic acid (MDXAA), vadimezan (ASA404), cisplatin, silibinin, proton pump inhibitor pantoprazole (PPZ), CNI-1493 and combinations thereof. In some embodiments, the macrophage inhibitor agent is an anti-IL-1 α agent (e.g., xilonix).

[0023] In some embodiments, the second agent is an antigen specific immune response enhancer agent selected from the group consisting of an anti-PD-1 agent, an anti-PD-L1 agent, a GITR (glucocorticoid-induced TNFR-related protein) stimulating agent, an anti-CTLA4 agent, an anti-TIM-3 agent, an anti-LAG-3 agent, an agent that enhances tumor antigen presentation (e.g., personalized cancer vaccine, autologous antigen presenting cell, autologous dendritic cells, artificial antigen presenting cell), a chemokine signaling agent, an anti-VEGF agent, a cytokine signal stimulating agent, and combinations thereof. In some embodiments, the anti-PD-1 agent is selected from the group consisting of pembrolizumab, nivolumab, PDR001, REGN2810 (SAR-439684), BGB-A317, BI 754091, IBI308, INCSHR-1210, JNJ-63723283, JS-001, MEDI0680 (AMP-514), MGA-012, PF-06801591, REGN-2810, TSR-042, PDR-001, camrelizumab (HR-301210), BCD-100, AGEN-2034, CS1001, Sym-021, LZM009, KN-035, AB122, genolimzumab (CBT-501), AK 104, GLS-010, and combinations thereof. In some embodiments, the anti-PD-L1 agent is selected from the group consisting of atezolizumab, durvalumab, avelumab, LY3300054, BGB-A333, SHR-1316, CK-301, and combinations thereof. In some embodiments, the GITR stimulating agent is selected from the group consisting of DTA-1, mGITRL, pGITRL, and combinations thereof. In some embodiments, the anti-CTLA4 agent is selected from the group consisting of ipilimumab, tremelimumab, and combinations thereof. In some embodiments, the

chemokine signaling agent is selected from the group consisting of CXCL16, a CXCR6 chemokine receptor (CD186) agonist, and combinations thereof. In some embodiments, the anti-VEGF agent is selected from the group consisting of bevacizumab, pazopanib, sunitinib, sorafenib, axitinib, ponatinib, regorafenib, cabozantinib, vandetanib, ramucirumab, lenvatinib, ziv-aflibercept, and combinations thereof. In some embodiments, the cytokine signal stimulating agent is an interleukin or an interferon. In some embodiments, the interleukin is selected from the group consisting of IL-2, IL-1, IL-7, IL-15, IL-12, IL-18 and combinations thereof. In some embodiments, the interferon is IFN alpha.

[0024] In some embodiments, the second agent is an antigen specific immune response enhancer agent selected from the group consisting of a flavonoid (e.g., flavonoid glycoside), lidocaine, lamotrigine, sulfamethoxazole, phenytoin, carbamazepine, sulfamethoxazole, phenytoin, allopurinol, paracetamol, mepivacaine, p-phenylenediamine, ciprofloxacin and moxifloxacin.

[0025] In some embodiments, the disease or condition is cancer. In some embodiments, the cancer is selected from the group consisting of ovarian cancer, breast cancer, cervical cancer, endometrial cancer, prostate cancer, testicular cancer, pancreatic cancer, esophageal cancer, head and neck cancer, gastric cancer, bladder cancer, lung cancer, bone cancer, colon cancer, rectal cancer, thyroid cancer, brain and central nervous system cancers, glioblastoma, neuroblastoma, neuroendocrine cancer, rhabdoid cancer, keratoacanthoma, epidermoid carcinoma, seminoma, melanoma, sarcoma, bladder cancer, liver cancer, kidney cancer, myeloma, lymphoma, and combinations thereof.

[0026] In some embodiments, the pharmaceutical composition is administered to a subject and the administering comprises administering a composition comprising a capsule comprising the first agent. In some embodiments, the capsule comprises a formulation comprising the first agent and one or more pharmaceutically acceptable excipients. In some embodiments, the one or more pharmaceutically acceptable excipients comprises lactose monohydrate, magnesium stearate, or a combination thereof. In some embodiments, a therapeutically effective amount of the first or second agent is administered. In some embodiments, the method further comprises administering a third agent to the subject. In some embodiments, the third agent comprises an antigen specific immune response enhancer agent, an anti-angiogenic agent, a chemotherapeutic agent, or combinations thereof. In some embodiments, the antigen specific immune response enhancer agent comprises an anti-PD-1 agent, an anti-PD-L1 agent, an anti-CTLA4 agent, an anti-TIM-3 agent, or an anti-LAG-3

agent. In some embodiments, the anti-angiogenic agent is selected from the group consisting of TNP-470, platelet factor 4, thrombospondin-1, tissue inhibitors of metalloproteases (TIMP1 and TIMP2), prolactin, angiostatin, endostatin, bFGF soluble receptor, transforming growth factor beta, interferon alpha, soluble KDR and FLT-1 receptors, placental proliferin-related protein, and combinations thereof.

[0027] In some embodiments, the chemotherapeutic agent is selected from the group consisting of aminoglutethimide, amsacrine, anastrozole, asparaginase, bcr, bicalutamide, bleomycin, buserelin, busulfan, campothecin, capecitabine, carboplatin, carmustine, chlorambucil, cisplatin, cladribine, clodronate, colchicine, cyclophosphamide, cyproterone, cytarabine, dacarbazine, dactinomycin, daunorubicin, dienestrol, diethylstilbestrol, docetaxel, doxorubicin, epirubicin, estradiol, estramustine, etoposide, exemestane, filgrastim, fludarabine, fludrocortisone, fluorouracil, fluoxymesterone, flutamide, gemcitabine, genistein, goserelin, hydroxyurea, idarubicin, ifosfamide, imatinib, interferon, irinotecan, irinotecan, letrozole, leucovorin, leuprolide, levamisole, lomustine, mechlorethamine, medroxyprogesterone, megestrol, melphalan, mercaptopurine, mesna, methotrexate, mitomycin, mitotane, mitoxantrone, nilutamide, nocodazole, octreotide, oxaliplatin, paclitaxel, pamidronate, pentostatin, plicamycin, porfimer, procarbazine, raltitrexed, rituximab, streptozocin, suramin, tamoxifen, temozolomide, teniposide, testosterone, thioguanine, thiotepa, titanocene dichloride, topotecan, trastuzumab, tretinoin, vinblastine, vincristine, vindesine, vinorelbine, and combinations thereof.

[0028] The present disclosure encompasses the recognition that a combination therapy with an agent that regulates activity within the tumor microenvironment (e.g., activity of T cells and/or the infiltration of T cells into the tumor environment) and an agent that inhibits PARP is useful for treating certain cancers.

[0029] In some embodiments, a PARP inhibitor increases infiltration of T cells in the tumor microenvironment. In some embodiments, a PARP inhibitor increases infiltration of CD4⁺ T cells in the tumor microenvironment. In some embodiments, a PARP inhibitor increases infiltration of CD8⁺ T cells in the tumor microenvironment. In some embodiments, a PARP inhibitor increases infiltration of CD4⁺ and CD8⁺ T cells in the tumor microenvironment. In some embodiments, a PARP inhibitor increases infiltration of macrophages in the tumor microenvironment. In some embodiments, a PARP inhibitor increases infiltration of Treg cells into the tumor microenvironment.

[0030] In some embodiments, agents that inhibit PARP include agents that inhibit PARP-1 and/or PARP-2. In some embodiments, agents that inhibit PARP include ABT-767, AZD 2461, BGB-290, BGP 15, CEP 8983, CEP 9722, DR 2313, E7016, E7449, fluzoparib (SHR 3162), IMP 4297, INO1001, JPI 289, JPI 547, monoclonal antibody B3-LysPE40 conjugate, MP 124, niraparib (ZEJULA) (MK-4827), NU 1025, NU 1064, NU 1076, NU1085, olaparib (AZD2281), ONO2231, PD 128763, R 503, R554, rucaparib (RUBRACA) (AG-014699, PF-01367338), SBP 101, SC 101914, Simmiparib, talazoparib (BMN-673), veliparib (ABT-888), WW 46, 2-(4-(Trifluoromethyl)phenyl)-7,8-dihydro-5H-thiopyrano[4,3-d]pyrimidin-4-ol, and salts or derivatives thereof. In some embodiments, agents that inhibit PARP are combinations of two or more agents selected from ABT-767, AZD 2461, BGB-290, BGP 15, CEP 8983, CEP 9722, DR 2313, E7016, E7449, fluzoparib (SHR 3162), IMP 4297, INO1001, JPI 289, JPI 547, monoclonal antibody B3-LysPE40 conjugate, MP 124, niraparib (ZEJULA) (MK-4827), NU 1025, NU 1064, NU 1076, NU1085, olaparib (AZD2281), ONO2231, PD 128763, R 503, R554, rucaparib (RUBRACA) (AG-014699, PF-01367338), SBP 101, SC 101914, Simmiparib, talazoparib (BMN-673), veliparib (ABT-888), WW 46, 2-(4-(Trifluoromethyl)phenyl)-7,8-dihydro-5H-thiopyrano[4,3-d]pyrimidin-4-ol, and salts or derivatives thereof. In some embodiments, an agent that inhibits part is or comprises niraparib, olaparib, rucaparib, talazoparib, veliparib, or any combination thereof. In some certain embodiments, an agent that inhibits PARP is niraparib ((3S)-3-[4-{7-(aminocarbonyl)-2H-indazol-2-yl}phenyl]piperidine), an orally active PARP inhibitor.

[0031] In some certain embodiments, an agent that regulates T cell activity is administered to a patient who is receiving, has received or will receive treatment with niraparib. In some certain embodiments, niraparib is administered to patient who is receiving, has received or will receive treatment with an agent that regulates T cell activity.

[0032] In some embodiments, cancers for treatment with a combination therapy of the present disclosure include melanoma, renal cell carcinoma, lung cancer, bladder cancer, breast cancer, cervical cancer, colon cancer, gall bladder cancer, laryngeal cancer, liver cancer, thyroid cancer, stomach cancer, salivary gland cancer, esophageal cancer, squamous cell carcinoma of the head and neck, ovarian cancer, prostate cancer, pancreatic cancer, or Merkel cell carcinoma. In some embodiments, a cancer is a solid tumor. In some embodiments, a patient or population of patients has a hematological cancer.

[0033] In some embodiments, the method comprises administering one or both of a therapy that regulates T cell activity and a therapy that inhibits poly [ADP-ribose]

polymerase (PARP) (“anti-PARP therapy”) to a subject so that the subject receives treatment with both therapies. In some embodiments, an agent that regulates T cell activity is an agent that enhances activity of antigen-specific cells in a tumor microenvironment. In some embodiments, an agent that regulates T cell activity is an agent that enhances activity of CD8⁺ T cells in tumor microenvironments. In some embodiments, an agent that regulates T cell activity is an agent that enhances activity of CD4⁺ T cells in tumor microenvironments. In some embodiments, regulating activity of T cells (e.g., CD4⁺ and/or CD8⁺ T cells) in the tumor environment involves upregulating the anti-tumor activity of individual T cells, increasing a rate of proliferation of the T cells, or enhancing the recruitment of the T cells to the tumor microenvironment. In some embodiments, an agent that regulates. In some embodiments, an agent that enhances T cell activity is an agent that blocks macrophage recruitment. In some embodiments, an agent that regulates T cell activity is an agent that blocks recruitment of M2 macrophages. In some embodiments, an agent that regulates T cell activity is an agent that blocks Treg cell recruitment.

[0034] In some embodiments, an agent that regulates T cell activity is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, a cell, a cell preparation, or a toxin. In some embodiments, an agent that regulates T cell activity is an antibody agent. Antibody agents can include any polypeptide or polypeptide complex that includes immunoglobulin structural elements sufficient to confer specific binding. Exemplary antibody agents include, but are not limited to, monoclonal antibodies, polyclonal antibodies, antibody fragments such as Fab fragments, Fab’ fragments, F(ab’)2 fragments, Fd’ fragments, Fd fragments, and isolated CDRs or sets thereof; single chain Fvs; polypeptide-Fc fusions; single domain antibodies (e.g., shark single domain antibodies such as IgNAR or fragments thereof); cameloid antibodies; masked antibodies (e.g., Probodies[®]); Small Modular ImmunoPharmaceuticals (“SMIPsTM”); single chain or Tandem diabodies (TandAb[®]); VHHs; Anticalins[®]; Nanobodies[®] minibodies; BiTE[®]s; ankyrin repeat proteins or DARPINS[®]; Avimers[®]; DARTs; TCR-like antibodies; Adnectins[®]; Affilins[®]; Trans-bodies[®]; Affibodies[®]; TrimerX[®]; MicroProteins; Fynomers[®]; Centyrins[®]; and KALBITOR[®]s.

[0035] In some embodiments, an anti-PARP therapy comprises administration of an agent that inhibits PARP. In some embodiments, an agent that inhibits PARP is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, or a toxin. In some embodiments, an agent that inhibits PARP is a small molecule. In some embodiments, an agent that inhibits PARP is selected from the group consisting of: ABT-

767, AZD 2461, BGB-290, BGP 15, CEP 8983, CEP 9722, DR 2313, E7016, E7449, fluzoparib (SHR 3162), IMP 4297, INO1001, JPI 289, JPI 547, monoclonal antibody B3-LysPE40 conjugate, MP 124, niraparib (ZEJULA) (MK-4827), NU 1025, NU 1064, NU 1076, NU1085, olaparib (AZD2281), ONO2231, PD 128763, R 503, R554, rucaparib (RUBRACA) (AG-014699, PF-01367338), SBP 101, SC 101914, Simmiparib, talazoparib (BMN-673), veliparib (ABT-888), WW 46, 2-(4-(Trifluoromethyl)phenyl)-7,8-dihydro-5H-thiopyrano[4,3-d]pyrimidin-4-ol, and salts or derivatives thereof. In some embodiments, an agent that inhibits PARP is selected from the group consisting of: niraparib, olaparib, rucaparib, talazoparib, and veliparib or salts or derivatives thereof. In some embodiments, an agent that inhibits PARP is niraparib or a salt or derivative thereof.

[0036] In some embodiments, an agent that regulates T cell activity is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, a cell, a cell preparation, or a toxin. In some embodiments, an agent that regulates T cell activity is an anti-PD-1 antibody agent. In some embodiments, an agent that enhances T cell activity is an anti-PD-1 antibody selected from the group consisting of: BGB-A317, BI 754091, IBI308, INCSHR-1210, JNJ-63723283, JS-001, MEDI-0680, MGA-012, nivolumab, PDR001, pembrolizumab, PF-06801591, REGN-2810, TSR-042 and derivatives thereof. In some embodiments, an anti-PD-1 antibody is pembrolizumab or a derivative thereof. .

BRIEF DESCRIPTION OF THE DRAWINGS

[0037] *Figure 1A* depicts representative CD4 IHC staining on tumor samples developed from Apc^{Min}/J heterozygous background upon vehicle or niraparib treatment.

[0038] *Figure 1B* depicts quantification of CD4 IHC staining images showing CD4 positive cell numbers in vehicle or niraparib treated mice.

[0039] *Figure 2A* depicts CD8 IHC staining on tumor samples developed from Apc^{Min}/J heterozygous background upon vehicle or niraparib treatment.

[0040] *Figure 2B* depicts quantification of CD8 IHC staining images showing CD8 positive cell numbers in vehicle or niraparib treated mice.

[0041] *Figure 3A* depicts Foxp3 IHC staining on tumor samples developed from Apc^{Min}/J heterozygous background upon vehicle or niraparib treatment.

[0042] *Figure 3B* depicts quantification of Foxp3 IHC staining images showing Foxp3 positive cell numbers in vehicle or niraparib treated mice.

[0043] *Figure 4A* depicts Iba1 IHC staining on tumor samples developed from Apc^{Min}/J heterozygous background upon vehicle or niraparib treatment.

[0044] *Figure 4B* depicts quantification of Iba1 IHC staining images showing Iba1 positive cell numbers in vehicle or niraparib treated mice.

[0045] *Figure 5A* depicts percentage of Ki67-positive CD4 and CD8 positive cells among total CD3 population in control or niraparib treated mice as assessed by flow cytometry.

[0046] *Figure 5B* depicts tumor volume in MDA-MB-436 huNOG-EXL mice treated with niraparib or control.

[0047] *Figure 6A* depicts tumor volume in MDA-MB-436 huNOG-EXL mice treated with niraparib, BLZ945, niraparib and BLZ945, or control.

[0048] *Figure 6B* depicts tumor volume on day 26 in MDA-MB-436 huNOG-EXL mice treated with niraparib, BLZ945, niraparib and BLZ945, or control.

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS

Definitions

[0049] The articles “a” and “an” as used herein in the specification and in the claims, unless clearly indicated to the contrary, should be understood to include the plural referents. Claims or descriptions that include “or” between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The invention includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The invention also includes embodiments in which more than one, or the entire group members are present in, employed in, or otherwise relevant to a given product or process. Furthermore, it is to be understood that the invention encompasses all variations, combinations, and permutations in which one or more limitations, elements, clauses, descriptive terms, etc., from one or more of the listed claims is introduced into another claim dependent on the same base claim (or, as relevant, any other claim) unless otherwise indicated or unless it would be evident to one of ordinary skill in the art that a contradiction or inconsistency would arise. Where elements are presented as lists, (e.g., in Markush group or similar format) it is to be understood that each subgroup of the elements is also disclosed, and any element(s) can be removed from the group. It should be understood

that, in general, where the invention, or aspects of the invention, is/are referred to as comprising particular elements, features, etc., certain embodiments of the invention or aspects of the invention consist, or consist essentially of, such elements, features, etc. For purposes of simplicity those embodiments have not in every case been specifically set forth in so many words herein. It should also be understood that any embodiment or aspect of the invention can be explicitly excluded from the claims, regardless of whether the specific exclusion is recited in the specification. The publications, websites and other reference materials referenced herein to describe the background of the invention and to provide additional detail regarding its practice are hereby incorporated by reference.

[0050] As used herein, the term “administration” typically refers to the administration of a composition to a subject or system. Those of ordinary skill in the art will be aware of a variety of routes that may, in appropriate circumstances, be utilized for administration to a subject, for example a human subject. For example, in some embodiments, administration may be ocular, oral, parenteral, topical, etc. In some particular embodiments, administration may be bronchial (e.g., by bronchial instillation), buccal, dermal (which may be or comprise, for example, one or more of topical to the dermis, intradermal, interdermal, transdermal, etc.), enteral, intra-arterial, intradermal, intragastric, intramedullary, intramuscular, intranasal, intraperitoneal, intrathecal, intravenous, intraventricular, within a specific organ (e.g., intrahepatic), mucosal, nasal, oral, rectal, subcutaneous, sublingual, topical, tracheal (e.g., by intratracheal instillation), vaginal, vitreal, etc. In some embodiments, administration may involve dosing that is intermittent (e.g., a plurality of doses separated in time) and/or periodic (e.g., individual doses separated by a common period of time) dosing. In some embodiments, administration may involve continuous dosing (e.g., perfusion) for at least a selected period of time.

[0051] As used herein, the terms “dosage form” or “unit dosage form” refer to a physically discrete unit of an active agent (e.g., a therapeutic or diagnostic agent) for administration to a subject. Typically, each such unit contains a predetermined quantity of active agent. In some embodiments, such quantity is a unit dosage amount (or a whole fraction thereof) appropriate for administration in accordance with a regimen that has been determined to correlate with a desired or beneficial outcome when administered to a relevant population (e.g., with a therapeutic regimen). Those of ordinary skill in the art appreciate that the total amount of a therapeutic composition or agent administered to a particular subject is

determined by one or more attending physicians and may involve administration of multiple dosage forms.

[0052] As used herein, the term “regimen” refers to a set of unit doses (typically more than one) that are administered individually to a subject, typically separated by one or more periods of time. In some embodiments, a given therapeutic agent is administered according to a regimen, which may involve one or more doses. In some embodiments, a regimen comprises a plurality of doses each of which is separated in time from other doses. In some embodiments, individual doses are separated from one another by a time period of the same length; in some embodiments, a regimen comprises a plurality of doses, wherein the doses are separated by time periods of different length. In some embodiments, a regimen comprises doses of the same amount. In some embodiments, a regimen comprises doses of different amounts. In some embodiments, a regimen comprises at least one dose, wherein the dose comprises one unit dose of the therapeutic agent. In some embodiments, a regimen comprises at least one dose, wherein the dose comprises two or more unit doses of the therapeutic agent.

[0053] As used herein, the term “patient”, “subject”, or “test subject” refers to any organism, including a human or non-human, to which provided compound or compounds described herein are administered in accordance with the present invention e.g., for experimental, diagnostic, prophylactic, and/or therapeutic purposes. Typical subjects include animals (e.g., mammals such as mice, rats, rabbits, canines, felines, horses, cattle, pigs, deer, non-human primates, and humans; insects; worms; birds; reptiles; amphibians; etc.). In embodiments, the subject is a human. In some embodiments, a subject may be suffering from, and/or susceptible to a disease, disorder, and/or condition (e.g., cancer). In some embodiments, a patient is a human that has been diagnosed with a cancer. In some embodiments, a patient is a human possessing one or more female reproductive organs.

[0054] The term “cancer” includes both solid tumors and hematological malignancies. Cancers include, but are not limited to, gynecological cancers, ovarian cancer, cancer of the fallopian tube(s), peritoneal cancer, breast cancer, cervical cancer, endometrial cancer, prostate cancer, testicular cancer, pancreatic cancer, esophageal cancer, head and neck cancer, gastric cancer, bladder cancer, lung cancer (e.g., adenocarcinoma, NSCLC and SCLC), bone cancer (e.g., osteosarcoma), colon cancer, rectal cancer, thyroid cancer, brain and central nervous system cancers, glioblastoma, neuroblastoma, neuroendocrine cancer, rhabdoid cancer, keratoacanthoma, epidermoid carcinoma, seminoma, melanoma, sarcoma

(e.g., liposarcoma), bladder cancer, liver cancer (e.g., hepatocellular carcinoma), kidney cancer (e.g., renal cell carcinoma), myeloid disorders (e.g., AML, CML, myelodysplastic syndrome and promyelocytic leukemia), and lymphoid disorders (e.g., leukemia, multiple myeloma, mantle cell lymphoma, ALL, CLL, B-cell lymphoma, T-cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma). Cancers include, but are not limited to, ovarian cancer, breast cancer, cervical cancer, endometrial cancer, prostate cancer, testicular cancer, pancreatic cancer, esophageal cancer, head and neck cancer, gastric cancer, bladder cancer, lung cancer, bone cancer, colon cancer, rectal cancer, thyroid cancer, brain and central nervous system cancers, glioblastoma, neuroblastoma, neuroendocrine cancer, rhabdoid cancer, keratoacanthoma, epidermoid carcinoma, seminoma, melanoma, sarcoma, bladder cancer, liver cancer, kidney cancer, myeloma, lymphoma, and combinations thereof.

[0055] The term “composition”, as in pharmaceutical composition, is intended to encompass a drug product comprising niraparib or its pharmaceutically acceptable salts, esters, solvates, polymorphs, stereoisomers or mixtures thereof, as well as, in some embodiments, one or more additional pharmaceutically active ingredients in combination with the niraparib. The composition may also include one or more inert ingredient(s) (e.g., pharmaceutically acceptable excipients). Such pharmaceutical compositions are synonymous with “formulation” and “dosage form”. Pharmaceutical composition of the invention include, but is not limited to, granules, tablets (single layered tablets, multilayered tablets, mini tablets, bioadhesive tablets, caplets, matrix tablets, tablet within a tablet, mucoadhesive tablets, modified release tablets, orally disintegrating tablets, pulsatile release tablets, timed release tablets, delayed release, controlled release, extended release and sustained release tablets), capsules (hard and soft or liquid filled soft gelatin capsules), pills, troches, sachets, powders, microcapsules, minitables, tablets in capsules and microspheres, matrix composition and the like. In some embodiments, the pharmaceutical composition refers to capsules. In some embodiments, the pharmaceutical composition refers to hard gelatin capsules or HPMC based capsules. In some embodiments, the pharmaceutical composition refers to hard gelatin capsules.

[0056] “Diluents” increase bulk of the composition to facilitate compression or create sufficient bulk for homogenous blend for capsule filling. Such compounds include e.g., lactose, starch, mannitol, sorbitol, dextrose, microcrystalline cellulose such as Avicel[®]; dibasic calcium phosphate, dicalcium phosphate dihydrate; tricalcium phosphate, calcium phosphate; anhydrous lactose, spray-dried lactose; pregelatinized starch, compressible sugar,

such as Di-Pac[®] (Amstar); mannitol, hydroxypropylmethylcellulose, hydroxypropylmethylcellulose acetate stearate, sucrose-based diluents, confectioner's sugar; monobasic calcium sulfate monohydrate, calcium sulfate dihydrate; calcium lactate trihydrate, dextrates; hydrolyzed cereal solids, amylose; powdered cellulose, calcium carbonate; glycine, kaolin; mannitol, sodium chloride; inositol, bentonite, and the like. Combinations of one or more diluents can also be used.

[0057] As used herein, the term “effective amount” or “therapeutically effective amount” refers to an amount of one or more therapeutic agents (e.g., niraparib in combination with one or more additional pharmaceutically active ingredients) that produces the desired effect for which it is administered. In some embodiments, the term refers to an amount that is sufficient, when administered to a population suffering from or susceptible to a disease, disorder, and/or condition in accordance with a regimen, to treat the disease, disorder, and/or condition. In some embodiments, a therapeutically effective amount is one that reduces the incidence and/or severity of, and/or delays onset of, one or more symptoms of the disease, disorder, and/or condition. Those of ordinary skill in the art will appreciate that the term “therapeutically effective amount” does not in fact require successful treatment be achieved in a particular individual. Rather, a therapeutically effective amount may be that amount that provides a particular desired pharmacological response in a significant number of subjects when administered to patients in need of such treatment. In some embodiments, reference to a therapeutically effective amount may be a reference to an amount as measured in one or more specific tissues (e.g., a tissue affected by the disease, disorder or condition) or fluids (e.g., blood, saliva, serum, sweat, tears, urine, etc.). Those of ordinary skill in the art will appreciate that, in some embodiments, a therapeutically effective amount of a particular agent or therapy may be formulated and/or administered in a single dose. In some embodiments, a therapeutically effective agent may be formulated and/or administered in a plurality of doses, for example, as part of a regimen.

[0058] The terms “enhance” or “enhancing” refers to an increase or prolongation of either the potency or duration of a desired effect of a composition described herein, or a diminution of any adverse symptomatology that is consequent upon the administration of the therapeutic agent or agents. Thus, in regard to enhancing the effect of niraparib disclosed herein, the term “enhancing” refers to the ability to increase or prolong, either in potency or duration, the effect of other therapeutic agents that are used in combination with niraparib disclosed herein. An “enhancing-effective amount,” as used herein, refers to an amount of niraparib or other

therapeutic agent which is adequate to enhance the effect of another therapeutic agent or niraparib in a desired system. When used in a patient, amounts effective for this use will depend on the severity and course of the disease, disorder or condition, previous therapy, the patient's health status and response to the drugs, and the judgment of the treating physician.

[0059] The term “excipient” means a pharmacologically inactive component such as a diluent, lubricant, surfactant, carrier, or the like. Excipients that are useful in preparing a pharmaceutical composition are generally safe, non-toxic and are acceptable for human pharmaceutical use. Reference to an excipient includes both one and more than one such excipient. Co-processed excipients are also covered under the scope of present invention.

[0060] “Filling agents” or “fillers” include compounds such as lactose, lactose monohydrate, calcium carbonate, calcium phosphate, dibasic calcium phosphate, calcium sulfate, microcrystalline cellulose, cellulose powder, dextrose, dextrans, dextran, starches, pregelatinized starch, sucrose, xylitol, lactitol, mannitol, sorbitol, sodium chloride, polyethylene glycol, and the like.

[0061] “Lubricants” and “glidants” are compounds that prevent, reduce or inhibit adhesion or friction of materials. Exemplary lubricants include, e.g., stearic acid, magnesium stearate, calcium hydroxide, talc, sodium stearyl fumarate, a hydrocarbon such as mineral oil, or hydrogenated vegetable oil such as hydrogenated soybean oil (Sterotex[®]), higher fatty acids and their alkali-metal and alkaline earth metal salts, such as aluminum, calcium, magnesium, zinc, stearic acid, sodium stearates, glycerol, talc, waxes, Stearowet[®], boric acid, sodium benzoate, sodium acetate, sodium chloride, leucine, a polyethylene glycol (e.g., PEG-4000) or a methoxypolyethylene glycol such as Carbowax[™], sodium oleate, sodium benzoate, glyceryl behenate, polyethylene glycol, magnesium or sodium lauryl sulfate, colloidal silica such as Syloid[™], Cab-O-Sil[®], a starch such as corn starch, silicone oil, a surfactant, and the like.

[0062] As used herein, “CA-125” means cancer antigen 125. A CA-125 test is used to measure the amount of the protein CA-125 in the blood of a patient. A CA-125 test may be used to monitor certain cancers during and after treatment, including use to evaluate prolongation of progression free survival. In some cases, a CA-125 test may be used to look for early signs of ovarian cancer in women with a very high risk of the disease.

[0063] As used herein, a “chemotherapeutic agent” refers to a chemical agent that inhibits the proliferation, growth, life-span and/or metastatic activity of cancer cells. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and CYTOXAN[®]

cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines (e.g., altretamine, triethylenemelamine, trietylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine); acetogenins; delta-9-tetrahydrocannabinol (e.g., dronabinol, MARINOL[®]); beta-lapachone; lapachol; colchicines; betulinic acid; a camptothecin (including the synthetic analogue topotecan (HYCAMTIN[®]), CPT-11 (irinotecan, CAMPTOSAR[®]), acetylcamptothecin, scoplectin, and 9-aminocamptothecin); bryostatin; callystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); podophyllotoxin; podophyllinic acid; teniposide; cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CB1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as the enediyne antibiotics (e.g., calicheamicin); dynemicin, including dynemicin A; bisphosphonates, such as clodronate; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antiobiotic chromophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabycin, caminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, ADRIAMYCIN[®] doxorubicin (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epitiostanol, mepitiothane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestrabucil; bisantrene; edatraxate;

defofamine; demecolcine; diaziquone; elformithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; 2-ethylhydrazide; procarbazine; PSK[®] polysaccharide complex (JHS Natural Products, Eugene, Oreg.); razoxane; rhizoxin; sizofuran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine; trichothecenes (e.g., T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine (ELDISINE[®], FILDESIN[®]); dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxanes, e.g., TAXOL[®] paclitaxel (Bristol-Myers Squibb Oncology, Princeton, N.J.), ABRAXANE[™] Cremophor-free, albumin-engineered nanoparticle formulation of paclitaxel (American Pharmaceutical Partners, Schaumburg, Ill.), and TAXOTERE[®] doxetaxel (Rhone-Poulenc Rorer, Antony, France); chloranbucil; gemcitabine (GEMZAR[®]); 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine (VELBAN[®]); platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine (ONCOVIN[®]); oxaliplatin; leucovorin; vinorelbine (NAVELBINE[®]); novantrone; edatrexate; daunomycin; aminopterin; xeloda; ibandronate; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; capecitabine; pharmaceutically acceptable salts, acids or derivatives of any of the above; as well as combinations of two or more of the above such as CHOP, an abbreviation for a combined therapy of cyclophosphamide, doxorubicin, vincristine, and prednisolone, and FOLFOX, an abbreviation for a treatment regimen with oxaliplatin (ELOXATIN[™]) combined with 5-FU and leucovorin.

[0064] Also included in this definition are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens and selective estrogen receptor modulators (SERMs), including, for example, tamoxifen (including NOLVADEX[®] tamoxifen), raloxifene, droloxifene, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and FARESTON[®] toremifene; aromatase inhibitors that inhibit the enzyme aromatase, which regulates estrogen production in the adrenal glands, such as, for example, 4(5)-imidazoles, aminoglutethimide, MEGASE[®] megestrol acetate, AROMASIN[®] exemestane, formestane, fadrozole, RIVISOR[®] vorozole, FEMARA[®] letrozole, and ARIMIDEX[®] anastrozole; and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; as well as troxacitabine (a 1,3-dioxolane nucleoside cytosine

analog); antisense oligonucleotides, particularly those that inhibit expression of genes in signaling pathways implicated in aberrant cell proliferation, such as, for example, PKC-alpha, Raf, H-Ras, and epidermal growth factor receptor (EGF-R); vaccines such as gene therapy vaccines, for example, ALLOVECTIN[®] vaccine, LEUVECTIN[®] vaccine, and VAXID[®] vaccine; PROLEUKIN[®] rIL-2; LURTOTECAN[®] topoisomerase 1 inhibitor; ABARELIX[®] rmRH; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

[0065] Also included in this definition are “antimetabolite chemotherapeutic agents” that are structurally similar to a metabolite, but cannot be used by the body in a productive manner. Many antimetabolite chemotherapeutic agents interfere with the production of the nucleic acids, RNA and DNA. Examples of antimetabolite chemotherapeutic agents include gemcitabine (GEMZAR[®]), 5-fluorouracil (5-FU), capecitabine (XELODA[™]), 6-mercaptopurine, methotrexate, 6-thioguanine, pemetrexed, raltitrexed, arabinosylcytosine ARA-C cytarabine (CYTOSAR-U[®]), dacarbazine (DTIC-DOMED), azocytosine, deoxycytosine, pyridimidine, fludarabine (FLUDARA[®]), cladribine, 2-deoxy-D-glucose etc. In some embodiments, an antimetabolite chemotherapeutic agent is gemcitabine. Gemcitabine HCl is sold by Eli Lilly under the trademark GEMZAR[®].

[0066] Also included in this definition are “platinum-based chemotherapeutic agents” that comprises an organic compound which contains platinum as an integral part of the molecule. In some embodiments, a chemotherapeutic agent is a platinum agent. In some such embodiments, the platinum agent is selected from cisplatin, carboplatin, oxaliplatin, nedaplatin, triplatin tetranitrate, phenanthriplatin, picoplatin, or satraplatin.

[0067] As used herein, the term “treatment” (also “treat” or “treating”) refers to any administration of a therapy that partially or completely alleviates, ameliorates, relieves, inhibits, prevents or delays onset of, reduces severity of, and/or reduces incidence of one or more symptoms, features, and/or causes of a particular disease, disorder, and/or condition. In some embodiments, such treatment may be of a subject who does not exhibit signs of the relevant disease, disorder and/or condition and/or of a subject who exhibits only early signs of the disease, disorder, and/or condition. Alternatively or additionally, such treatment may be of a subject who exhibits one or more established signs of the relevant disease, disorder and/or condition. In some embodiments, treatment may be of a subject who has been diagnosed as suffering from the relevant disease, disorder, and/or condition. In some embodiments, treatment may be of a subject known to have one or more susceptibility factors

that are statistically correlated with increased risk of development of the relevant disease, disorder, and/or condition.

[0068] As used herein, the term “pharmaceutically acceptable salt” refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge et al., describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 1977, 66, 1–19, incorporated herein by reference. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like.

[0069] Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C_{1-4}alkyl)_4$ salts. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate.

[0070] As used herein, the term “pharmaceutical composition” refers to a composition in which an active agent is formulated together with one or more pharmaceutically acceptable carriers. In some embodiments, the active agent is present in unit dose amount appropriate for administration in a therapeutic regimen that shows a statistically significant probability of

achieving a predetermined therapeutic effect when administered to a relevant population. In some embodiments, a pharmaceutical composition may be specially formulated for administration in solid or liquid form, including those adapted for oral administration, for example, drenches (aqueous or non-aqueous solutions or suspensions), tablets, e.g., those targeted for buccal, sublingual, and systemic absorption, boluses, powders, granules, and pastes for application to the tongue. A pharmaceutical composition can also refer to a medicament.

[0071] As used herein, the term “antibody” refers to a polypeptide that includes canonical immunoglobulin sequence elements sufficient to confer specific binding to a particular target antigen. As is known in the art, intact antibodies as produced in nature are approximately 150 kD tetrameric agents comprised of two identical heavy chain polypeptides (about 50 kD each) and two identical light chain polypeptides (about 25 kD each) that associate with each other into what is commonly referred to as a “Y-shaped” structure. Each heavy chain is comprised of at least four domains (each about 110 amino acids long) – an amino-terminal variable (VH) domain (located at the tips of the Y structure), followed by three constant domains: CH1, CH2, and the carboxy-terminal CH3 (located at the base of the Y’s stem). A short region, known as the “switch”, connects the heavy chain variable and constant regions. The “hinge” connects CH2 and CH3 domains to the rest of the antibody. Two disulfide bonds in this hinge region connect the two heavy chain polypeptides to one another in an intact antibody. Each light chain is comprised of two domains – an amino-terminal variable (VL) domain, followed by a carboxy-terminal constant (CL) domain, separated from one another by another “switch”. Those skilled in the art are well familiar with antibody structure and sequence elements, recognize “variable” and “constant” regions in provided sequences, and understand that there may be some flexibility in definition of a “boundary” between such domains such that different presentations of the same antibody chain sequence may, for example, indicate such a boundary at a location that is shifted one or a few residues relative to a different presentation of the same antibody chain sequence. Intact antibody tetramers are comprised of two heavy chain-light chain dimers in which the heavy and light chains are linked to one another by a single disulfide bond; two other disulfide bonds connect the heavy chain hinge regions to one another, so that the dimers are connected to one another and the tetramer is formed. Naturally-produced antibodies are also glycosylated, typically on the CH2 domain. Each domain in a natural antibody has a structure characterized by an “immunoglobulin fold” formed from two beta sheets (e.g., 3-, 4-, or 5-stranded sheets)

packed against each other in a compressed antiparallel beta barrel. Each variable domain contains three hypervariable loops known as “complement determining regions” (CDR1, CDR2, and CDR3) and four somewhat invariant “framework” regions (FR1, FR2, FR3, and FR4). When natural antibodies fold, the FR regions form the beta sheets that provide the structural framework for the domains, and the CDR loop regions from both the heavy and light chains are brought together in three-dimensional space so that they create a single hypervariable antigen binding site located at the tip of the Y structure. The Fc region of naturally-occurring antibodies binds to elements of the complement system, and also to receptors on effector cells, including for example effector cells that mediate cytotoxicity. As is known in the art, affinity and/or other binding attributes of Fc regions for Fc receptors can be modulated through glycosylation or other modification. In some embodiments, antibodies produced and/or utilized in accordance with the present invention include glycosylated Fc domains, including Fc domains with modified or engineered such glycosylation. For purposes of the present invention, in certain embodiments, any polypeptide or complex of polypeptides that includes sufficient immunoglobulin domain sequences as found in natural antibodies can be referred to and/or used as an “antibody”, whether such polypeptide is naturally produced (e.g., generated by an organism reacting to an antigen), or produced by recombinant engineering, chemical synthesis, or other artificial system or methodology. In some embodiments, an antibody is polyclonal; in some embodiments, an antibody is monoclonal. In some embodiments, an antibody has constant region sequences that are characteristic of mouse, rabbit, primate, or human antibodies. In some embodiments, antibody sequence elements are humanized, primatized, chimeric, etc., as is known in the art. Moreover, the term “antibody” as used herein, can refer in appropriate embodiments (unless otherwise stated or clear from context) to any of the art-known or developed constructs or formats for utilizing antibody structural and functional features in alternative presentation. For example, in some embodiments, an antibody utilized in accordance with the present invention is in a format selected from, but not limited to, intact IgA, IgG, IgE or IgM antibodies; bi- or multi-specific antibodies (e.g., Zybodies[®], etc); antibody fragments such as Fab fragments, Fab’ fragments, F(ab’)2 fragments, Fd’ fragments, Fd fragments, and isolated CDRs or sets thereof; single chain Fvs; polypeptide-Fc fusions; single domain antibodies (e.g., shark single domain antibodies such as IgNAR or fragments thereof); cameloid antibodies; masked antibodies (e.g., Probodies[®]); Small Modular ImmunoPharmaceuticals (“SMIPsTM”); single chain or Tandem diabodies (TandAb[®]); VHHs; Anticalins[®]; Nanobodies[®] minibodies; BiTE[®]s;

ankyrin repeat proteins or DARPINs[®]; Avimers[®]; DARTs; TCR-like antibodies; Adnectins[®]; Affilins[®]; Trans-bodies[®]; Affibodies[®]; TrimerX[®]; MicroProteins; Fynomers[®]; Centyrins[®]; and KALBITOR[®]s. In some embodiments, an antibody may lack a covalent modification (e.g., attachment of a glycan) that it would have if produced naturally. In some embodiments, an antibody may contain a covalent modification (e.g., attachment of a glycan, a payload [e.g., a detectable moiety, a therapeutic moiety, a catalytic moiety, etc], or other pendant group [e.g., poly-ethylene glycol, etc.]

[0072] As used herein, the term “antibody agent” refers to an agent that specifically binds to a particular antigen. In some embodiments, the term encompasses any polypeptide or polypeptide complex that includes immunoglobulin structural elements sufficient to confer specific binding. Exemplary antibody agents include, but are not limited to monoclonal antibodies or polyclonal antibodies. In some embodiments, an antibody agent may include one or more constant region sequences that are characteristic of mouse, rabbit, primate, or human antibodies. In some embodiments, an antibody agent may include one or more sequence elements are humanized, primatized, chimeric, etc., as is known in the art. In many embodiments, the term “antibody agent” is used to refer to one or more of the art-known or developed constructs or formats for utilizing antibody structural and functional features in alternative presentation. For example, in some embodiments, an antibody agent utilized in accordance with the present invention is in a format selected from, but not limited to, intact IgA, IgG, IgE or IgM antibodies; bi- or multi- specific antibodies (e.g., Zybodies[®], etc); antibody fragments such as Fab fragments, Fab’ fragments, F(ab’)2 fragments, Fd’ fragments, Fd fragments, and isolated CDRs or sets thereof; single chain Fvs; polypeptide-Fc fusions; single domain antibodies (e.g., shark single domain antibodies such as IgNAR or fragments thereof); cameloid antibodies; masked antibodies (e.g., Probodies[®]); Small Modular ImmunoPharmaceuticals (“SMIPsTM”); single chain or Tandem diabodies (TandAb[®]); VHHs; Anticalins[®]; Nanobodies[®] minibodies; BiTE[®]s; ankyrin repeat proteins or DARPINs[®]; Avimers[®]; DARTs; TCR-like antibodies; Adnectins[®]; Affilins[®]; Trans-bodies[®]; Affibodies[®]; TrimerX[®]; MicroProteins; Fynomers[®]; Centyrins[®]; and KALBITOR[®]s. In some embodiments, an antibody may lack a covalent modification (e.g., attachment of a glycan) that it would have if produced naturally. In some embodiments, an antibody may contain a covalent modification (e.g., attachment of a glycan, a payload [e.g., a detectable moiety, a therapeutic moiety, a catalytic moiety, etc], or other pendant group [e.g., poly-ethylene glycol, etc.]. In many embodiments, an antibody agent is or comprises a

polypeptide whose amino acid sequence includes one or more structural elements recognized by those skilled in the art as a complementarity determining region (CDR); in some embodiments an antibody agent is or comprises a polypeptide whose amino acid sequence includes at least one CDR (e.g., at least one heavy chain CDR and/or at least one light chain CDR) that is substantially identical to one found in a reference antibody. In some embodiments an included CDR is substantially identical to a reference CDR in that it is either identical in sequence or contains between 1-5 amino acid substitutions as compared with the reference CDR. In some embodiments an included CDR is substantially identical to a reference CDR in that it shows at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity with the reference CDR. In some embodiments an included CDR is substantially identical to a reference CDR in that it shows at least 96%, 96%, 97%, 98%, 99%, or 100% sequence identity with the reference CDR. In some embodiments an included CDR is substantially identical to a reference CDR in that at least one amino acid within the included CDR is deleted, added, or substituted as compared with the reference CDR but the included CDR has an amino acid sequence that is otherwise identical with that of the reference CDR. In some embodiments an included CDR is substantially identical to a reference CDR in that 1-5 amino acids within the included CDR are deleted, added, or substituted as compared with the reference CDR but the included CDR has an amino acid sequence that is otherwise identical to the reference CDR. In some embodiments an included CDR is substantially identical to a reference CDR in that at least one amino acid within the included CDR is substituted as compared with the reference CDR but the included CDR has an amino acid sequence that is otherwise identical with that of the reference CDR. In some embodiments an included CDR is substantially identical to a reference CDR in that 1-5 amino acids within the included CDR are deleted, added, or substituted as compared with the reference CDR but the included CDR has an amino acid sequence that is otherwise identical to the reference CDR. In some embodiments, an antibody agent is or comprises a polypeptide whose amino acid sequence includes structural elements recognized by those skilled in the art as an immunoglobulin variable domain. In some embodiments, an antibody agent is a polypeptide protein having a binding domain which is homologous or largely homologous to an immunoglobulin-binding domain.

As used herein, the term “homology” refers to the overall relatedness between polymeric molecules, e.g., between nucleic acid molecules (e.g., DNA molecules and/or RNA molecules) and/or between polypeptide molecules. In some embodiments, polymeric

molecules are considered to be “homologous” to one another if their sequences are at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% identical. In some embodiments, polymeric molecules are considered to be “homologous” to one another if their sequences are at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% similar (e.g., containing residues with related chemical properties at corresponding positions). For example, as is well known by those of ordinary skill in the art, certain amino acids are typically classified as similar to one another as “hydrophobic” or “hydrophilic” amino acids, and/or as having “polar” or “non-polar” side chains. Substitution of one amino acid for another of the same type may often be considered a “homologous” substitution.

[0073] As will be understood by those skilled in the art, a variety of algorithms are available that permit comparison of sequences in order to determine their degree of homology, including by permitting gaps of designated length in one sequence relative to another when considering which residues “correspond” to one another in different sequences. Calculation of the percent homology between two nucleic acid sequences, for example, can be performed by aligning the two sequences for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second nucleic acid sequences for optimal alignment and non-corresponding sequences can be disregarded for comparison purposes). In certain embodiments, the length of a sequence aligned for comparison purposes is at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or substantially 100% of the length of the reference sequence. The nucleotides at corresponding nucleotide positions are then compared. When a position in the first sequence is occupied by the same nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position; when a position in the first sequence is occupied by a similar nucleotide as the corresponding position in the second sequence, then the molecules are similar at that position. The percent homology between the two sequences is a function of the number of identical and similar positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which needs to be introduced for optimal alignment of the two sequences. Representative algorithms and computer programs useful in determining the percent homology between two nucleotide sequences include, for example, the algorithm of Meyers and Miller (CABIOS, 1989, 4: 11-17), which has been incorporated into the ALIGN program (version 2.0) using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. The percent homology between two

nucleotide sequences can, alternatively, be determined for example using the GAP program in the GCG software package using an NWSgapdna.CMP matrix.

[0074] As used herein, the term “combination therapy” refers to a clinical intervention in which a subject is simultaneously exposed to two or more therapeutic regimens (*e.g.*, two or more therapeutic agents). In some embodiments, the two or more therapeutic regimens may be administered simultaneously. In some embodiments, the two or more therapeutic regimens may be administered sequentially (*e.g.*, a first regimen administered prior to administration of any doses of a second regimen). In some embodiments, the two or more therapeutic regimens are administered in overlapping dosing regimens. In some embodiments, administration of combination therapy may involve administration of one or more therapeutic agents or modalities to a subject receiving the other agent(s) or modality. In some embodiments, combination therapy does not necessarily require that individual agents be administered together in a single composition (or even necessarily at the same time). In some embodiments, two or more therapeutic agents or modalities of a combination therapy are administered to a subject separately, *e.g.*, in separate compositions, via separate administration routes (*e.g.*, one agent orally and another agent intravenously), and/or at different time points. In some embodiments, two or more therapeutic agents may be administered together in a combination composition, or even in a combination compound (*e.g.*, as part of a single chemical complex or covalent entity), via the same administration route, and/or at the same time.

Cancers

[0075] Cancer is an abnormal growth of cells which tend to proliferate in an uncontrolled way and, in some cases, to metastasize (spread). Cancer is not one disease. It is a group of more than 100 different and distinctive diseases. Cancer can involve any tissue of the body and have many different forms in each body area. Most cancers are named for the type of cell or organ in which they start. A tumor can be cancerous or benign. A benign tumor means the tumor can grow but does not spread. A cancerous tumor is malignant, meaning it can grow and spread to other parts of the body. If a cancer spreads (metastasizes), the new tumor bears the same name as the original (primary) tumor. The frequency of a particular cancer may depend on gender. While skin cancer is the most common type of malignancy for both men and women, the second most common type in men is prostate cancer and in women, breast cancer.

[0076] The methods of the disclosure can be used to treat any type of cancer known in the art. Non-limiting examples of cancers to be treated by the methods of the present disclosure can include melanoma (e.g., metastatic malignant melanoma), renal cancer (e.g., clear cell carcinoma), prostate cancer (e.g., hormone refractory prostate adenocarcinoma), pancreatic adenocarcinoma, breast cancer, colon cancer, lung cancer (e.g., non-small cell lung cancer), esophageal cancer, squamous cell carcinoma of the head and neck, liver cancer, ovarian cancer, cervical cancer, thyroid cancer, glioblastoma, glioma, leukemia, lymphoma, and other neoplastic malignancies. Additionally, the invention includes refractory or recurrent malignancies whose growth may be inhibited using the methods of the invention. In some embodiments, a cancer to be treated by the methods of the present disclosure include, for example, carcinoma, squamous carcinoma (for example, cervical canal, eyelid, tunica conjunctiva, vagina, lung, oral cavity, skin, urinary bladder, tongue, larynx, and gullet), and adenocarcinoma (for example, prostate, small intestine, endometrium, cervical canal, large intestine, lung, pancreas, gullet, intestinum rectum, uterus, stomach, mammary gland, and ovary). In some embodiments, a cancer to be treated by the methods of the present disclosure further include sarcomata (for example, myogenic sarcoma), leukosis, neuroma, melanoma, and lymphoma.

[0077] In some embodiments, a patient or population of patients to be treated with a combination therapy of the present disclosure have a solid tumor. In some embodiments, a solid tumor is a melanoma, renal cell carcinoma, lung cancer, bladder cancer, breast cancer, cervical cancer, colon cancer, gall bladder cancer, laryngeal cancer, liver cancer, thyroid cancer, stomach cancer, salivary gland cancer, prostate cancer, pancreatic cancer, or Merkel cell carcinoma. In some embodiments, a patient or population of patients to be treated with a combination therapy of the present disclosure have a hematological cancer. In some embodiments, the patient has a hematological cancer such as Diffuse large B cell lymphoma (“DLBCL”), Hodgkin’s lymphoma (“HL”), Non-Hodgkin’s lymphoma (“NHL”), Follicular lymphoma (“FL”), acute myeloid leukemia (“AML”), or Multiple myeloma (“MM”).

Role of poly(ADP-ribose) polymerases (PARPs)

[0078] Poly(ADP-ribose) polymerases (PARPs) are a family of enzymes that cleave NAD⁺, releasing nicotinamide, and successively add ADP-ribose units to form ADP-ribose polymers. Accordingly, activation of PARP enzymes can lead to depletion of cellular NAD⁺ levels (e.g., PARPs as NAD⁺ consumers) and mediates cellular signaling through ADP-

ribosylation of downstream targets. PARP-1 is a zinc-finger DNA-binding enzyme that is activated by binding to DNA double or single strand breaks. It was known that anti-alkylating agents could deplete the NAD⁺ content of tumor cells, and the discovery of PARPs explained this phenomena. (Parp Inhibitors and Cancer Therapy. Curtin N. *in* Poly ADP Ribosylation. ed. Alexander Burke, Lands Bioscience and Springer Bioscience, 2006: 218-233). Anti-alkylating agents induce DNA strand breaks, which activates of PARP-1, which is part of the DNA repair pathway. Poly ADP-ribosylation of nuclear proteins by PARP-1 converts DNA damage into intracellular signals that can either activate DNA repair (e.g., by the base excision repair (BER) pathway); or trigger cell death in the presence of DNA damage that is too extensive and cannot be efficiently repaired.

[0079] PARP-2 contains a catalytic domain and is capable of catalyzing a poly(ADP-ribosyl)ation reaction. PARP-2 displays auto-modification properties similar to PARP-1. The protein is localized in the nucleus *in vivo* and may account for the residual poly(ADP-ribose) synthesis observed in PARP-1-deficient cells, treated with alkylating agents or hydrogen peroxide. Some agents that inhibit PARP (e.g., agents primarily aimed at inhibiting PARP-1) may also inhibit PARP-2 (e.g., niraparib).

[0080] The role of PARP enzymes in DNA damage response (e.g., repair of DNA in response to genotoxic stress) has led to the compelling suggestion that PARP inhibitors may be useful anti-cancer agents. PARP inhibitors may be particularly effective in treating cancers resulting from germ line or sporadic deficiency in the homologous recombination DNA repair pathway, such as BRCA-1 and/or BRCA-2 deficient cancers.

[0081] Pre-clinical *ex vivo* and *in vivo* experiments suggest that PARP inhibitors are selectively cytotoxic for tumors with homozygous inactivation of BRCA-1 and/or BRCA-2 genes, which are known to be important in the homologous recombination (HR) DNA repair pathway. The biological basis for the use of PARP inhibitors as single agents in cancers with defects in BRCA-1 and/or BRCA-2 is the requirement of PARP-1 and PARP-2 for base excision repair (BER) of the damaged DNA. Upon formation of single-strand DNA breaks, PARP-1 and PARP-2 bind at sites of lesions, become activated, and catalyze the addition of long polymers of ADP-ribose (PAR chains) on several proteins associated with chromatin, including histones, PARP itself, and various DNA repair proteins. This results in chromatin relaxation and fast recruitment of DNA repair factors that access and repair DNA breaks. Normal cells repair up to 10,000 DNA defects daily and single strand breaks are the most common form of DNA damage. Cells with defects in the BER pathway enter S phase with

unrepaired single strand breaks. Pre-existing single strand breaks are converted to double strand breaks as the replication machinery passes through the break. Double strand breaks present during S phase are preferentially repaired by the error-free HR pathway. Cells with inactivation of genes required for HR, such as BRCA-1 and/or BRCA-2, accumulate stalled replication forks during S phase and may use error-prone non-homologous end joining (NHEJ) to repair damaged DNA. Both the inability to complete S phase (because of stalled replication forks) and error-prone repair by NHEJ, are thought to contribute to cell death.

[0082] Without wishing to be bound by theory, it is hypothesized that treatment with PARP inhibitors may selectively kill a subset of cancer cells with deficiencies in DNA repair pathways (e.g., inactivation of BRCA-1 and/or BRCA-2). For example, a tumor arising in a patient with a germline BRCA mutation has a defective homologous recombination DNA repair pathway and would be increasingly dependent on BER, a pathway blocked by PARP inhibitors, for maintenance of genomic integrity. This concept of inducing death by use of PARP inhibitors to block one DNA repair pathway in tumors with pre-existing deficiencies in a complementary DNA repair pathways is called synthetic lethality.

[0083] The therapeutic potential of PARP inhibitors is further expanded by the observation that PARP inhibitors not only have monotherapy activity in HR-deficient tumors, but are also effective in preclinical models in combination with other agents such as cisplatin, carboplatin, alkylating and methylating agents, radiation therapy, and topoisomerase I inhibitors. In contrast to the rationale for monotherapy in which PARP inhibition alone is sufficient for cell death in HR-deficient cancers (due to endogenous DNA damage), PARP is required for repair of DNA damage induced by standard cytotoxic chemotherapy. In some cases, the specific role of PARP is not known, but PARP is known to be required to release trapped topoisomerase I/irinotecan complexes from DNA. Temozolomide-induced DNA damage is repaired by the BER pathway, which requires PARP to recruit repair proteins. Combination therapies that enhance or synergize the cancer therapy without significantly increasing toxicity would provide substantial benefit to cancer patients, including ovarian cancer patients.

PARP inhibitors

[0084] PARP inhibitors have shown activity against tumors with existing DNA repair defects, such as BRCA1 and BRCA2. Without wishing to be bound by theory, treatment with PARP inhibitors (e.g., PARP-1/2 inhibitors) may selectively kill a subset of cancer cell types by exploiting their deficiencies in DNA repair. Human cancers exhibit genomic

instability and an increased mutation rate due to underlying defects in DNA repair. These deficiencies render cancer cells more dependent on the remaining DNA repair pathways and targeting these pathways is expected to have a much greater impact on the survival of the tumor cells than on normal cells.

[0085] In some embodiments, a PARP inhibitor increases infiltration of T cells in the tumor microenvironment. In some embodiments, a PARP inhibitor increases infiltration of CD4⁺ T cells in the tumor microenvironment. In some embodiments, a PARP inhibitor increases infiltration of CD8⁺ T cells in the tumor microenvironment. In some embodiments, a PARP inhibitor increases infiltration of CD4⁺ and CD8⁺ T cells in the tumor microenvironment. In some embodiments, a PARP inhibitor increases infiltration of macrophages in the tumor microenvironment. In some embodiments, a PARP inhibitor increases infiltration of Treg cells into the tumor microenvironment. In some embodiments, a PARP inhibitor increases infiltration of CD335⁺ T cells in the tumor microenvironment. In some embodiments, a PARP inhibitor increases infiltration of Fox3p⁺ T cells in the tumor microenvironment. In some embodiments, a PARP inhibitor increases infiltration of Iba1⁺ T cells in the tumor microenvironment.


[0086] In some embodiments, agents that inhibit PARP include agents that inhibit PARP-1 and/or PARP-2. In some embodiments, agents that inhibit PARP include ABT-767, AZD 2461, BGB-290, BGP 15, CEP 8983, CEP 9722, DR 2313, E7016, E7449, fluzoparib (SHR 3162), IMP 4297, INO1001, JPI 289, JPI 547, monoclonal antibody B3-LysPE40 conjugate, MP 124, niraparib (ZEJULA) (MK-4827), NU 1025, NU 1064, NU 1076, NU1085, olaparib (AZD2281), ONO2231, PD 128763, R 503, R554, rucaparib (RUBRACA) (AG-014699, PF-01367338), SBP 101, SC 101914, Simmiparib, talazoparib (BMN-673), veliparib (ABT-888), WW 46, 2-(4-(Trifluoromethyl)phenyl)-7,8-dihydro-5H-thiopyrano[4,3-d]pyrimidin-4-ol, and salts or derivatives thereof. In some embodiments, a PARP inhibitor is niraparib, olaparib, rucaparib, talazoparib, veliparib, or any combination thereof. In some embodiments, a PARP inhibitor can be prepared as a pharmaceutically acceptable salt. In some embodiments, an agent that inhibits PARP is niraparib or a salt or derivative thereof. One of skill in the art will appreciate that such salt forms can exist as solvated or hydrated polymorphic forms.

Niraparib

[0087] Niraparib is an orally active and potent poly (ADP-ribose) polymerase, or PARP, inhibitor. Niraparib and pharmaceutically acceptable salts thereof, are disclosed in

International Publication No. WO2007/113596 and European Patent No. EP2007733B1; International Publication No. WO2008/084261 and U.S. Patent No. 8,071,623; and International Publication No. WO2009/087381 and U.S. Patent No. 8,436,185. Methods of making niraparib and pharmaceutically acceptable salts thereof are disclosed in International Publication Nos. WO2014/088983 and WO2014/088984. Methods to treat cancer with niraparib and pharmaceutically acceptable salts thereof are disclosed in U.S. Provisional Patent Application Nos. 62/356,461 and 62/402,427. The contents of each of the foregoing references are incorporated herein by reference in their entirety.

[0088] In some embodiments, the present invention relates to use of niraparib in combination with one or more additional pharmaceutically active agents affecting activity within the tumor microenvironment. Niraparib, (3S)-3-[4-{7-(aminocarbonyl)-2H-indazol-2-yl}phenyl]piperidine, is an orally available, potent, poly (adenosine diphosphate [ADP]-ribose) polymerase (PARP)-1 and -2 inhibitor. Niraparib has the following structure:



[0089] The empirical molecular formula for niraparib is $C_{26}H_{30}N_4O_5S$ and its molecular weight is 510.61. Niraparib tosylate monohydrate drug substance is a white to off-white, non-hygroscopic crystalline solid. Niraparib solubility is pH independent below the pKa of 9.95, with an aqueous free base solubility of 0.7 mg/mL to 1.1 mg/mL across the physiological pH range. See WO 2008/084261 (published on July 17, 2008) and WO 2009/087381 (published July 16, 2009), the entirety of each of which is hereby incorporated by reference. Niraparib can be prepared according to Scheme 1 of WO 2008/084261. As used herein, the term “niraparib” means any of the free base compound ((3S)-3-[4-{7-(aminocarbonyl)-2H-indazol-2-yl}phenyl]piperidine), a salt form, including pharmaceutically acceptable salts, of (3S)-3-[4-{7-(aminocarbonyl)-2H-indazol-2-yl}phenyl]piperidine (e.g., (3S)-3-[4-{7-(aminocarbonyl)-2H-indazol-2-yl}phenyl]piperidine tosylate), or a solvated or hydrated form thereof (e.g., (3S)-3-[4-{7-(aminocarbonyl)-2H-indazol-2-yl}phenyl]piperidine tosylate monohydrate). In some embodiments, such forms may be individually referred to as “niraparib free base”, “niraparib tosylate” and “niraparib tosylate monohydrate”, respectively. Unless otherwise specified, the term “niraparib” includes all forms of the compound (3S)-3-[4-{7-(aminocarbonyl)-2H-indazol-2-yl}phenyl]piperidine.

[0090] In some embodiments, niraparib can be prepared as a pharmaceutically acceptable salt. One of skill in the art will appreciate that such salt forms can exist as solvated or hydrated polymorphic forms. In some embodiments, niraparib is prepared in the form of a hydrate.

[0091] In certain embodiments, niraparib is prepared in the form of a tosylate salt. In some embodiments, niraparib is prepared in the form of a tosylate monohydrate.

[0092] The crystalline tosylate monohydrate salt of niraparib is being developed as a monotherapy agent for tumors with defects in the homologous recombination (HR) deoxyribonucleic acid (DNA) repair pathway and as a sensitizing agent in combination with cytotoxic agents and radiotherapy.

[0093] Provided herein are compositions containing niraparib or its pharmaceutically acceptable salts. The compositions may further include one or more additional active ingredients which impact activity in the tumor microenvironment (e.g., activity of T cells and/or the infiltration of T cells into the tumor environment).

[0094] In some embodiments, the niraparib a pharmaceutically acceptable salt thereof. In some embodiments, the pharmaceutically acceptable salt is niraparib tosylate monohydrate.

[0095] The formulation can comprise one or more components, including niraparib. The components can be combined to create granules that are then compressed to form tablets.

[0096] The niraparib may be present in the formulation as a pharmaceutically acceptable salt. For example, the niraparib can be niraparib tosylate monohydrate.

[0097] The niraparib formulations described herein are administered and dosed in accordance with good medical practice, taking into account the clinical condition of the individual patient, the site and method of administration, scheduling of administration, and other factors known to medical practitioners. In human therapy, the dosage forms described herein deliver niraparib formulations that maintain a therapeutically effective amount of niraparib in plasma the while reducing the side effects associated with an elevated $C_{\max}$ blood plasma level of niraparib.

Pharmaceutically acceptable salts

[0098] In some embodiments, the niraparib used in a composition disclosed herein is the form of a free base, pharmaceutically acceptable salt, prodrug, analog or complex. In some instances, the niraparib comprises the form of a pharmaceutically acceptable salt. In some embodiments, with respect to niraparib in a composition, a pharmaceutically acceptable salt includes, but is not limited to, 4-methylbenzenesulfonate salts, sulfate salts, benzenesulfate

salts, fumarate salts, succinate salts, and stereoisomers or tautomers thereof. In some embodiments, with respect to niraparib in a composition, a pharmaceutically acceptable salt includes, but is not limited to, tosylate salts. In some embodiments, with respect to niraparib in a composition, a pharmaceutically acceptable salt includes, but is not limited to, tosylate monohydrate salts.

Additional Pharmaceutically acceptable excipients

[0099] In some aspects, the pharmaceutical composition disclosed herein further comprises one or more pharmaceutically acceptable excipients. In some embodiments, the one or more pharmaceutically acceptable excipient is present in an amount of about 0.1-99 % by weight. Exemplary pharmaceutically acceptable excipients for the purposes of pharmaceutical compositions disclosed herein include, but are not limited to, binders, disintegrants, superdisintegrants, lubricants, diluents, fillers, flavors, glidants, sorbents, solubilizers, chelating agents, emulsifiers, thickening agents, dispersants, stabilizers, suspending agents, adsorbents, granulating agents, preservatives, buffers, coloring agents and sweeteners or combinations thereof. Examples of binders include microcrystalline cellulose, hydroxypropyl methylcellulose, carboxyvinyl polymer, polyvinylpyrrolidone, polyvinylpolypyrrolidone, carboxymethylcellulose calcium, carboxymethylcellulose sodium, ceratonia, chitosan, cottonseed oil, dextrates, dextrin, ethylcellulose, gelatin, glucose, glyceryl behenate, galactomannan polysaccharide, hydroxyethyl cellulose, hydroxyethylmethyl cellulose, hydroxypropyl cellulose, hypromellose, inulin, lactose, magnesium aluminum silicate, maltodextrin, methylcellulose, poloxamer, polycarbophil, polydextrose, polyethylene glycol, polyethylene oxide, polymethacrylates, sodium alginate, sorbitol, starch, sucrose, sunflower oil, vegetable oil, tocopherols, zein, or combinations thereof. Examples of disintegrants include hydroxypropyl methylcellulose (HPMC), low substituted hydroxypropyl cellulose (L-HPC), croscarmellose sodium, sodium starch glycolate, lactose, magnesium aluminum silicate, methylcellulose, polacrillin potassium, sodium alginate, starch, or combinations thereof. Examples of a lubricant include stearic acid, sodium stearyl fumarate, glyceryl behenate, calcium stearate, glycerin monostearate, glyceryl palmitostearate, magnesium lauryl sulfate, mineral oil, palmitic acid, myristic acid, poloxamer, polyethylene glycol, sodium benzoate, sodium chloride, sodium lauryl sulfate, talc, zinc stearate, potassium benzoate, magnesium stearate or combinations thereof. Examples of diluents include talc, ammonium alginate, calcium carbonate, calcium lactate, calcium phosphate, calcium silicate, calcium sulfate, cellulose, cellulose acetate, corn starch,

dextrates, dextrin, dextrose, erythritol, ethylcellulose, fructose, fumaric acid, glyceryl palmitostearate, isomalt, kaolin, lactitol, lactose, magnesium carbonate, magnesium oxide, maltodextrin, maltose, mannitol, microcrystalline cellulose, polydextrose, polymethacrylates, simethicone, sodium alginate, sodium chloride, sorbitol, starch, sucrose, sulfobutylether β -cyclodextrin, tragacanth, trehalose, xylitol, or combinations thereof. In some embodiments, the pharmaceutically acceptable excipient is hydroxypropyl methylcellulose (HPMC). In some embodiments, the pharmaceutically acceptable excipient is low substituted hydroxypropyl cellulose (L-HPC). In some embodiments, the pharmaceutically acceptable excipient is lactose. In some embodiments, the pharmaceutically acceptable excipient is lactose monohydrate. In some embodiments, the pharmaceutically acceptable excipient is magnesium stearate. In some embodiments, the pharmaceutically acceptable excipient is lactose monohydrate and magnesium stearate.

[00100] Various useful fillers or diluents include, but are not limited to calcium carbonate (Barcroft™, MagGran™, Millicarb™, Pharma- Carb™, Precarb™, Sturcal™, Vivapres Ca™), calcium phosphate, dibasic anhydrous (Emcompress Anhydrous™, Fujicalin™), calcium phosphate, dibasic dihydrate (Calstar™, Di-Cafos™, Emcompress™), calcium phosphate tribasic (Tri-Cafos™, TRI- TAB™), calcium sulphate (Destab™, Drierite™, Snow White™, Cal-Tab™, Compactrol™), cellulose powdered (Arbocel™, Elcema™, Sanacet™), silicified microcrystalline cellulose, cellulose acetate, compressible sugar (Di-Pac™), confectioner's sugar, dextrates (Candex™, Emdex™), dextrin (Avedex™, Caloreen™, Primogran W™), dextrose (Caridex™, Dextrofin™, Tab fine D-100™), fructose (Fructofin™, Krystar™), kaolin (Lion™, Sim 90™), lactitol (Finlac DC™, Finlac MCX™), lactose (Anhydrox™, CapsuLac™, Fast-Flo™, FlowLac™, GranuLac™, InhaLac™, Lactochem™, Lactohaie™, Lactopress™, Microfme™, Microtose™, Pharmatose™, Prisma Lac™, Respiritose™, SacheLac™, SorboLac™, Super-Tab™, Tablettose™, Wyndale™, Zeparox™), lactose monohydrate, magnesium carbonate, magnesium oxide (MagGran MO™), maltodextrin (C*Dry MD™, Lycatab DSH™, Maldex™, Maitagran™, Maltrin™, Maltrin QD™, Paselli MD 10 PH™, Star-Dri™), maltose (Advantose 100™), mannitol (Mannogem™, Pearlitol™), microcrystalline cellulose (Avicel PH™, Celex™, Celphere™, Ceolus KG™, Emcocel™, Pharmacel™, Tabulose™, Vivapur™), polydextrose (Litesse™), simethicone (Dow Corning Q7- 2243 LVA™, Cow Coming Q7-2587™, Sentry Simethicone™), sodium alginate (Keltone™, Protanal™), sodium chloride (Alberger™), sorbitol (Liponec 70-NC™, Liponic 76-NCv, Meritol™,

Neosorb™, Sorbitol Instant™, Sorbogem™), starch (Fluflax W™, Instant Pure-Cote™, Melojel™, Meritena Paygel 55™, Perfectamyl D6PH™, Pure-Cote™, Pure-Dent™, Pure-Gel™, Pure-Set™, Purity 21™, Purity 826™, Tablet White™), pregelatinized starch, sucrose, trehalose and xylitol, or mixtures thereof.

[00101] Various useful disintegrants include, but are not limited to, alginic acid (Protacid™, Satialgine H8™), calcium phosphate, tribasic (TRI-TAB™), carboxymethylcellulose calcium (ECG 505™), carboxymethylcellulose sodium (Akucell™, Finnfix™, Nymcel Tylose CB™), colloidal silicon dioxide (Aerosil™, Cab-O-Sil™, Wacker HDK™), croscarmellose sodium (Ac-Di-Sol™, Pharmacel XL™, Primellose™, Solutab™, Vivasol™), crospovidone (Collison CL™, Collison CL-M™, Polyplasdone XL™), docusate sodium, guar gum (Meyprodor™, Meyprofm™, Meyproguar™), low substituted hydroxypropyl cellulose, magnesium aluminum silicate (Magnabite™, Neusilin™, Pharmsorb™, Veegum™), methylcellulose (Methocel™, Metolose™), microcrystalline cellulose (Avicel PH™, Ceolus KG™, Emcoel™, Ethispheres™, Fibrocel™, Pharmacel™, Vivapur™), povidone (Collison™, Plasdone™) sodium alginate (Kelcosol™, Ketone™, Protanal™), sodium starch glycolate, polacrilin potassium (Amberlite IRP88™), silicified microcrystalline cellulose (ProSotv™), starch (Aytex P™, Fluftex W™, Melojel™, Meritena™, Paygel 55™, Perfectamyl D6PH™, Pure-Bind™, Pure-Cote™, Pure-Dent™, Purity 21™, Purity 826™, Tablet White™) or pre-gelatinized starch (Lycatab PGST™, Merigel™, National 78-1551™, Pharma-Gel™, Prejel™, Sepistab ST 200™, Spreess B820™, Starch 1500 G™, Tablitz™, Unipure LD™), or mixtures thereof.

[00102] Various useful lubricants include, but are not limited to, calcium stearate (HyQual™), glycerine monostearate (Imwitor™ 191 and 900, Kessco GMS5™, 450 and 600, Myvaplex 600P™, Myvatex™, Rita GMS™, Stepan GMS™, Tegin™, Tegin™ 503 and 515, Tegin 4100™, Tegin M™, Unimate GMS™), glyceryl behenate (Compritol 888 ATO™), glyceryl palmitostearate (Precirol ATO 5™), hydrogenated castor oil (Castorwax MP 80™, Croduret™, Cutina HR™, Fancol™, Simulsol 1293™), hydrogenated vegetable oil 0 type I (Sterotex™, Dynasan P60™, Hydrocote™, Lipovol HS-K™, Sterotex HM™), magnesium lauryl sulphate, magnesium stearate, medium-chain triglycerides (Captex 300™, Labrafac CC™, Miglyol 810™, Neobee M5™, Nesatol™, Waglinol 3/9280™), poloxamer (Pluronic™, Synperonic™), polyethylene glycol (Carbowax Sentry™, Lipo™, Lipoxol™, Lutrol E™, Pluriol E™), sodium benzoate (Antimol™), sodium chloride, sodium lauryl sulphate (Elfan 240™, Texapon KI 2P™), sodium stearyl fumarate (Pruv™), stearic acid

(Hystrene™, industrene™, Kortacid 1895™, Pristerene™), talc (Altaic™, Luzenac™, Luzenac Pharma™, Magsil Osmanthus™, 0 Magsil Star™, Superiore™), sucrose stearate (Surfhope SE Pharma D-1803 F™) and zinc stearate (HyQual™) or mixtures thereof. Examples of suitable lubricants include, but are not limited to, magnesium stearate, calcium stearate, zinc stearate, stearic acid, talc, glyceryl behenate, polyethylene glycol, polyethylene oxide polymers, sodium lauryl sulfate, magnesium lauryl sulfate, sodium oleate, sodium stearyl fumarate, DL-leucine, colloidal silica, and others as known in the art. In some embodiments a lubricant is magnesium stearate.

[00103] Various useful glidants include, but are not limited to, tribasic calcium phosphate (TRI- TAB™), calcium silicate, cellulose, powdered (Sanacel™, Solka- Floe™), colloidal silicon dioxide (Aerosil™, Cab-O-Sil M-5P™, Wacker HDK™), magnesium silicate, magnesium trisilicate, starch (Melojel™, Meritena™, Paygel 55™, Perfectamyl D6PH™, Pure-Bind™, Pure-Cote™, Pure-Dent™, Pure-Gel™, Pure-Set™, Purity 21™, Purity 826™, Tablet White™) and talc (Luzenac Pharma™, Magsil Osmanthus™, Magsil Star™, Superiore™), or mixtures thereof.

[00104] Pharmaceutically acceptable surfactants include, but are limited to both non-ionic and ionic surfactants suitable for use in pharmaceutical dosage forms. Ionic surfactants may include one or more of anionic, cationic or zwitterionic surfactants. Various useful surfactants include, but are not limited to, sodium lauryl sulfate, monooleate, monolaurate, monopalmitate, monostearate or another ester of polyoxyethylene sorbitane, sodium dioctylsulfosuccinate (DOSS), lecithin, stearyl alcohol, cetostearyl alcohol, cholesterol, polyoxyethylene ricin oil, polyoxyethylene fatty acid glycerides, poloxamer, or any other commercially available co-processed surfactant like SEPITRAP® 80 or SEPITRAP® 4000 and mixtures thereof.

T cells

[00105] T cells can be either CD4⁺ or CD8⁺ T cells, and can be CD28 positive T cells or CD28 negative T cells. T cells can also be either memory T cells or naïve T cells. Furthermore, T cells can express CD3. T cells can be regulatory cells. In some embodiments, T regulatory cells express proteins, including, for example CD25, CTLA-4, or FoxP3, or combinations thereof. In some embodiments, T cells are Th1 cells or Th2 cells. In some embodiments, Th1 cells are capable of secreting cytokines including, for example, interferon gamma, interleukin 2, and TNF-beta. In some embodiments, Th1 cells express markers, including, for example, CD4, CD94, CD119 (IFN γ R1), CD183 (CXCR3), CD186 (CXCR6),

CD191 (CCR1), CD195 (CCR5), CD212 (IL-12R β 1&2), CD254 (RANKL), CD278 (ICOS), IL-18R, MRP1, NOTCH3, or TIM3, or combinations thereof. In some embodiments, Th2 cells are capable of secreting cytokines including, for example, IL-4, IL-5, IL-6, IL-9, IL-10, or IL-13, or combinations thereof. In some embodiments, Th2 cells express markers including, for example, CRTH2, CCR4, or CCR3, or combinations thereof.

Promoting Antigen Specific T cell Signaling

[00106] T cells are a class of lymphocytes, having specific T cell receptors (TCRs) that are produced as a result of gene rearrangement. T cells have diverse roles, which are accomplished by the differentiation of distinct subsets of T cells, recognizable by discrete patterns of gene expression. T cells capable of antigen recognition (e.g., antigen-specific T cells) are generally classified as “CD4⁺” or “CD8⁺,” depending on whether a CD4 or a CD8 molecule is displayed on the cell surface. CD4⁺ cells recognize exogenously-produced antigen which has been taken up by an antigen presenting cell (APC), processed, and displayed on the APC cell surface together with a major histocompatibility complex (MHC) class II molecule. In general, CD4⁺ T cells provide the signals to activate other cells, e.g., CD4⁺ cells activate CD8⁺ cells, to induce B cells to produce antibodies, or to activate macrophages. In contrast, CD8⁺ cells are cytotoxic, and recognize antigen produced from within a cell and displayed on the cell surface together with an MHC Class I molecule.

[00107] In some embodiments, a combination therapy comprises one or more agents that activate T cell proliferation and/or stimulate antigen-specific T cell activity. An agent that activates antigen-specific T cell proliferation and/or activity may be or comprise an agent of any chemical class including, for example, a carbohydrate, a lipid, a nucleic acid, a polypeptide, a small molecule, a metal, a cell, etc. In some embodiments, an agent that activates antigen-specific T cell proliferation and/or activity may be or comprise a polypeptide (or complex thereof). In some embodiments, an agent that activates antigen-specific T cell proliferation and/or activity may be or comprise an antibody agent, a cytokine, a ligand, a receptor, a toxin, etc.

[00108] An agent that activates or stimulates antigen-specific T cell activity typically functions to enhance the total activity of antigen-specific T cells in a tumor microenvironment in order to initiate, strengthen or maintain an anti-tumor immunogenic response. In some embodiments, activation of antigen-specific T cell activity involves the stimulation of individual antigen-specific T cells (e.g., CD4⁺ or CD8⁺ cells) from an unstimulated or partially stimulated state. In some embodiments, activation of antigen-

specific T cell activity involves inducing proliferation of antigen-specific T cells (e.g., CD4⁺ or CD8⁺ cells). In some embodiments, activation of antigen-specific T cell activity involves the recruitment of antigen-specific T cells (e.g., CD4⁺ or CD8⁺ cells) to a tumor microenvironment. Herein in the term “activated” in reference to an antigen-specific T cell refers to the activated state of the cell that results from an interaction between the T cell receptor of the antigen-specific T cell (e.g., CD4⁺ or CD8⁺ cell) and an activating signal (e.g., peptide antigen) specific for the T cell receptor presented to the antigen-specific T cell by an antigen-presenting cell. A composition can comprise a disease-specific immunogenic neoantigen peptide. A composition can comprise two or more disease-specific immunogenic neoantigen peptides. A composition may comprise a precursor to a disease-specific immunogenic peptide (such as a protein, peptide, DNA and RNA). A precursor to a disease-specific immunogenic peptide can generate or be generated to the identified disease-specific immunogenic neoantigen peptide. In some embodiments, a therapeutic composition comprises a precursor of an immunogenic peptide. The precursor to a disease-specific immunogenic peptide can be a pro-drug. In some embodiments, the composition comprising a disease-specific immunogenic neoantigen peptide may further comprise an adjuvant. For example, the neoantigen peptide can be utilized as a vaccine. In some embodiments, an immunogenic vaccine may comprise a pharmaceutically acceptable immunogenic neoantigen peptide. In some embodiments, an immunogenic vaccine may comprise a pharmaceutically acceptable precursor to an immunogenic neoantigen peptide (such as a protein, peptide, DNA and RNA). In some embodiments, the neoantigen peptide is directed to a shared antigen that can be recognized by patient T cells within a large patient group. In some embodiments, a method of treatment comprises administering to a subject an effective amount of an antibody specifically recognizing an immunogenic neoantigen peptide.

[00109] The methods described herein can be useful in the personalized medicine context, where immunogenic neoantigen peptides are used to develop therapeutics (such as vaccines or therapeutic antibodies) for the same individual. Thus, a method of treating a disease in a subject can comprise identifying an immunogenic neoantigen peptide in a subject according to the methods described herein; and synthesizing the peptide (or a precursor thereof); and administering the peptide or an antibody specifically recognizing the peptide to the subject. In some embodiments, an expression pattern of an immunogenic neoantigen can serve as the essential basis for the generation of patient specific vaccines. In some embodiments, an expression pattern of an immunogenic neoantigen can serve as the essential basis for the

generation of a vaccine for a group of patients with a particular disease. Thus, particular diseases, e.g., particular types of tumors, can be selectively treated in a patient group.

[00110] There are a variety of ways in which to produce immunogenic neoantigens. Proteins or peptides may be made by any technique known to those of skill in the art, including the expression of proteins, polypeptides or peptides through standard molecular biological techniques, the isolation of proteins or peptides from natural sources, in vitro translation, or the chemical synthesis of proteins or peptides. In general, such disease specific neoantigens may be produced either in vitro or in vivo. Immunogenic neoantigens may be produced in vitro as peptides or polypeptides, which may then be formulated into a personalized vaccine or immunogenic composition and administered to a subject. In vitro production of immunogenic neoantigens can comprise peptide synthesis or expression of a peptide/polypeptide from a DNA or RNA molecule in any of a variety of bacterial, eukaryotic, or viral recombinant expression systems, followed by purification of the expressed peptide/polypeptide. Alternatively, immunogenic neoantigens can be produced in vivo by introducing molecules (e.g., DNA, RNA, and viral expression systems) that encode an immunogenic neoantigen into a subject, whereupon the encoded immunogenic neoantigens are expressed. In some embodiments, a polynucleotide encoding an immunogenic neoantigen peptide can be used to produce the neoantigen peptide in vitro.

[00111] In some embodiments, a polynucleotide comprises a sequence with at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identity to a polynucleotide encoding an immunogenic neoantigen.

[00112] The polynucleotide may be, e.g., DNA, cDNA, PNA, CNA, RNA, single- and/or double-stranded, native or stabilized forms of polynucleotides, or combinations thereof. A nucleic acid encoding an immunogenic neoantigen peptide may or may not contain introns so long as it codes for the peptide. In some embodiments in vitro translation is used to produce the peptide.

Cytokines

[00113] In some embodiments an agent that activates an antigen-specific T cell proliferation and/or activity is an interleukin or an agent that increases the expression and/or activity of an interleukin. For example, Interleukin-2 (IL-2) is a cytokine synthesized by T-cells which was first identified in conjunction with its role in the expansion of T-cells in response to an antigen (Smith, K. A. *Science* 240:1169 (1988)). Several studies have demonstrated that IL-2 has antitumor effects (see e.g., Lotze, M. T. et al, in “*Interleukin 2*”,

ed. K. A. Smith, Academic Press, Inc., San Diego, Calif., p237 (1988); Rosenberg, S., *Ann. Surgery* 208:121 (1988)). In fact, IL-2 has been utilized to treat subjects suffering from malignant melanoma, renal cell carcinoma, and acute myelogenous leukemia. (Rosenberg, S. A., et al., *N. Eng. J. Med.* 316:889-897 (1987); Dutcher, J. P., et al., *J. Clin. Oncol.* 7:477-485 (1989); Foa, R., et al., *Br. J. Haematol.* 77:491-496 (1991)).

[00114] In some embodiments, other interleukins, such as IL-1, IL-7, IL-15, IL-12 and IL-18 can be employed to activate antigen-specific T cells in a tumor microenvironment. These interleukins have been shown to directly promote antigen-specific T cell proliferation/survival and development of cytolytic effector functions. In some cases, these interleukins act on cytokine-induced killer (CIK) cells, which are a heterogeneous population of effector CD8⁺ T cells with diverse TCR specificities, possessing non-MHC-restricted cytolytic activities against tumor cells.

[00115] Another cytokine which in some embodiments may activate an antigen-specific T cell is interferon- α (IFN- α). IFN- α is an IFN type I cytokine, has been employed to treat leukemia, myeloma, and renal cell carcinomas. IFN type I cytokines have been shown to increase class I MHC molecule expression. Because most cytolytic T-cells (CTLs) recognize foreign antigens bound to class I MHC molecules, type I IFNs may boost the effector phase of cell-mediated immune responses by enhancing the efficiency of CTL-mediated killing. At the same time, type I IFN may inhibit the cognitive phase of immune responses, by preventing the activation of class II MHC-restricted helper T-cells.

[00116] Members of the chemokine family of cytokines may also act to promote antigen-specific T cell anti-tumor activity. Chemokines are known to act as a chemoattractant to mediate chemotaxis in nearby responsive cells. In cases where responsive T cells express T cell receptors which are specific for tumor antigens, administration of chemokines can contribute to anti-tumor activity. For example, expression of the cytokine CXCL16 by tumor cells can enhance recruitment of tumor infiltrating cells such as CD4⁺ and CD8⁺ T cells to the tumor, via binding of CXCL16 to its receptor CXCR6 expressed in T cells.

Checkpoint inhibitors

[00117] A balance between co-stimulatory and inhibitory signals regulates the amplitude and the quality of T-cell responses driven by TCR signaling. T cells require CD28-mediated co-stimulation (also known as signal 2) for the full acquisition of effector functions. However, excessive T-cell activation can result in the loss of self-tolerance, underscoring the importance of immune inhibitory pathways, or immune checkpoints, that regulate T-cell

activity. The immunosuppressive tumor microenvironment directly affects the expression of immune checkpoint proteins, thereby favoring resistance to anti-tumor immune response. T cells are essential effectors for cancer immune surveillance, and inhibition of T-cell-dependent anti-tumor response can promote tumor progression. Engagement of the CD28 homologue receptor cytotoxic T-lymphocyte-associated protein 4 (CTLA-4) on T cells by co-stimulatory molecules negatively regulates T-cell activation. It has been demonstrated that administration of neutralizing CTLA-4 antibody into tumor-bearing mice resulted in tumor rejection. In addition, mice that had rejected their tumors following anti-CTLA-4 treatment were protected against subsequent tumor rechallenge, indicating the establishment of immunological memory. Additional mouse and human studies have validated these results and shown that CTLA-4 blockade triggers anticancer immune responses. For example, administration of the anti-CTLA-4 antibody tremelimumab to human patients resulted in a highly significant increase in intratumoral infiltration by CD8⁺ cells in biopsy samples taken after tremelimumab treatment (Huang *et al.*, “CTLA4 blockade induces frequent tumor infiltration by activated lymphocytes regardless of clinical response in humans,” *Clin Can Res*, 17:4101-4109 (2011)). Importantly, inhibition of CTLA-4 signaling not only enhances effector T-cell functions, but it also renders effector T cells insensitive to regulatory T-cell-driven suppression. Infusion of anti-CTLA-4 antibodies after vaccination with irradiated, autologous tumor cells secreting GM-CSF (GVAX)-induced anti-tumor immunity but no toxicity in metastatic melanoma patients. The clinical efficacy of anti-CTLA-4 therapy was further confirmed in a phase III clinical trial where ipilimumab, a human mAb against CTLA-4, was shown to enhance the overall survival of metastatic melanoma patients. The demonstrated anticancer activity of ipilimumab (Yervoy) led to its approval by the FDA for the treatment of metastatic melanoma.

[00118] Other key inhibitory checkpoints that are relevant in cancer immunotherapy include PD-1 and Tim-3. Expression of the PD-1 receptor is induced in T cells upon activation [48]. Tumor cells can drive T-cell dysfunction because of their expression of PD-1 receptor ligands, PD-L1 and PD-L2. It has been demonstrated that transgenic expression of PD-L1 in mastocytoma tumor cells prevented their elimination by CTL and enhanced their invasiveness *in vivo*. Thus, cancer tissues limit the host immune response through PD-1 ligands and their ligation to PD-1 on antigen-specific CD8 T cells, a phenomenon termed adaptive immune resistance. The molecular bases accounting for adaptive immune resistance remain elusive. However, it has been suggested that the therapeutic efficacy of PD-1

blockade is due to the restoration of CD8 T-cell effector function in the tumor microenvironment. Preclinical models have demonstrated that blockade of PD-L1/PD-1 interactions could reinforce anticancer immune responses and promote tumor control. Tim-3 is another T-cell inhibitory receptor that was initially identified on fully differentiated Th1 cells. The Tim-3 ligand, galectin-9, induces T-cell death. In the tumor microenvironment, dysfunctional CD8 T cells could be identified by the co-expression of Tim-3 and PD-1. Tim-3 and PD-1 expression are associated with tumor antigen-specific CD8⁺ T-cell dysfunction in melanoma patients and prevent the expansion of tumor antigen-specific CD8⁺ T cells induced by vaccination. Other therapies targeting immune checkpoints are currently in development such as agonist antibodies targeting molecules which activate T cells such as CD137 (BMS-663513), OX40 (MEDI6383) NCT02221960, CD40 (CP870,893) or GITR (TRAX518) NCT01239134 as well as drugs favoring DC activation such as LAG3-Fusion protein (IMP321)

Programmed Death 1 (PD-1)

[00119] Programmed Death 1 (PD-1) (also known as Programmed Cell Death 1) (encoded by the gene *Pdcd1*) is a type I transmembrane protein of 268 amino acids originally identified by subtractive hybridization of a mouse T cell line undergoing apoptosis (Ishida et al., *Embo J.*, 11: 3887-95 (1992)). The normal function of PD-1, expressed on the cell surface of activated T cells under healthy conditions, is to down-modulate unwanted or excessive immune responses, including autoimmune reactions.

[00120] PD-1 is a member of the CD28/CTLA-4 family of T-cell regulators, and is expressed on activated T-cells, B-cells, and myeloid lineage cells (Greenwald et al., *Annu. Rev. Immunol.*, 23: 515-548 (2005); and Sharpe et al., *Nat. Immunol.*, 8: 239-245 (2007)). PD-1 is an inhibitory member of the CD28 family of receptors that also includes CD28, CTLA-4, ICOS and BTLA. PD-1 is expressed on activated B cells, T cells, and myeloid cells (Agata et al., supra; Okazaki et al. (2002) *Curr. Opin. Immunol.* 14:391779-82; Bennett et al. (2003) *J. Immunol.* 170:711-8).

[00121] Two ligands for PD-1 have been identified, PD ligand 1 (PD-L1) and PD ligand 2 (PD-L2), both of which belong to the B7 protein superfamily (Greenwald et al, supra). PD-1 has been shown to negatively regulate antigen receptor signaling upon engagement of its ligands (PD-L1 and/or PD-L2).

[00122] PD-L1 is expressed in a variety of cell types, including cells of the lung, heart, thymus, spleen, and kidney (see, e.g., Freeman et al., *J. Exp. Med.*, 192(7): 1027-1034

(2000); and Yamazaki et al., *J. Immunol.*, 169(10): 5538-5545 (2002)). PD-L1 expression is upregulated on macrophages and dendritic cells (DCs) in response to lipopolysaccharide (LPS) and GM-CSF treatment, and on T-cells and B-cells upon signaling via T-cell and B-cell receptors. PD-L1 also is expressed in a variety of murine tumor cell lines (see, e.g., Iwai et al., *Proc. Natl. Acad. Sci. USA*, 99(9): 12293-12297 (2002); and Blank et al., *Cancer Res.*, 64(3): 1140-1145 (2004)). In contrast, PD-L2 exhibits a more restricted expression pattern and is expressed primarily by antigen presenting cells (e.g., dendritic cells and macrophages), and some tumor cell lines (see, e.g., Latchman et al., *Nat. Immunol.*, 2(3): 261-238 (2001)). High PD-L1 expression in tumors, whether on the tumor cell, stroma, or other cells within the tumor microenvironment, correlates with poor clinical prognosis, presumably by inhibiting effector T cells and upregulating regulatory T cells (Treg) in the tumor.

[00123] PD-1 and family members are type I transmembrane glycoproteins containing an Ig variable-type (V-type) domain responsible for ligand binding and a cytoplasmic tail, which is responsible for the binding of signaling molecules. The cytoplasmic tail of PD-1 contains 2 tyrosine-based signaling motifs, an immunoreceptor tyrosine-based inhibition motif (ITIM) and an immunoreceptor tyrosine-based switch motif (ITSM). PD-1 negatively regulates T-cell activation, and this inhibitory function is linked to an ITSM in the cytoplasmic domain (see, e.g., Greenwald et al., *supra*; and Parry et al., *Mol. Cell. Biol.*, 25: 9543-9553 (2005)). Following T cell stimulation, PD-1 recruits the tyrosine phosphatases SHP-1 and SHP-2 to the ITSM motif within its cytoplasmic tail, leading to the dephosphorylation of effector molecules, such as CD3 ζ , PKC θ and ZAP70, which are involved in the CD3 T cell signaling cascade. The mechanism by which PD-1 down-modulates T cell responses is similar to, but distinct from, that of CTLA-4. PD-1 was shown to be expressed on activated lymphocytes, including peripheral CD4⁺ and CD8⁺ T cells, B cells, T regs, and natural killer cells. Expression has also been shown during thymic development on CD4-/CD8- (double-negative) T cells, as well as subsets of macrophages and dendritic cells. The ligands for PD-1 (PD-L1 and PD-L2) are constitutively expressed or can be induced in a variety of cell types. PD-L1 is expressed at low levels on various non-hematopoietic tissues, most notably on vascular endothelium, whereas PD-L2 protein is predominantly expressed on antigen-presenting cells found in lymphoid tissue or chronic inflammatory environments. Both ligands are type I transmembrane receptors containing both IgV- and IgC-like domains in the extracellular region and short cytoplasmic regions with no known signaling motifs. Binding of either PD-1 ligand to PD-1 inhibits T cell activation triggered through the T cell receptor.

PD-L2 is thought to control immune T cell activation in lymphoid organs, whereas PD-L1 serves to dampen unwarranted T cell function in peripheral tissues. Although healthy organs express little (if any) PD-L1, a variety of cancers were demonstrated to express abundant levels of this T cell inhibitor, which, via its interaction with the PD-1 receptor on tumor-specific T cells, plays a critical role in immune evasion by tumors.

[00124] PD-1 deficiency can lead to autoimmunity. For example, C57BL/6 PD-1 knockout mice have been shown to develop a lupus-like syndrome (see, e.g., Nishimura et al., *Immunity*, 11: 141-1151 (1999)). In humans, a single nucleotide polymorphism in the PD-1 gene is associated with higher incidences of systemic lupus erythematosus, type 1 diabetes, rheumatoid arthritis, and progression of multiple sclerosis (see, e.g., Nielsen et al., *Tissue Antigens*, 62(6): 492-497 (2003); Bertsias et al., *Arthritis Rheum.*, 60(1): 207-218 (2009); Ni et al., *Hum. Genet.*, 121(2): 223-232 (2007); Tahoori et al., *Clin. Exp. Rheumatol.*, 29(5): 763-767 (2011); and Kroner et al., *Ann. Neurol.*, 58(1): 50-57 (2005)). Abnormal PD-1 expression also has been implicated in T-cell dysfunctions in several pathologies, such as tumor immune evasion and chronic viral infections (see, e.g., Barber et al., *Nature*, 439: 682-687 (2006); and Sharpe et al., *supra*). PD-1 is abnormally expressed in a variety of cancers (see, e.g., Brown et al., *J. Immunol.*, 170: 1257-1266 (2003); and Flies et. al., *Yale Journal of Biology and Medicine*, 84: 409-421 (2011)), and PD-L1 expression in some renal cell carcinoma patients correlates with tumor aggressiveness.

[00125] Recent studies demonstrate that T-cell suppression induced by PD-1 also plays a role in the suppression of anti-tumor immunity. For example, PD-L1 is expressed on a variety of human and mouse tumors, and binding of PD-1 to PD-L1 on tumors results in T-cell suppression and tumor immune evasion and protection (Dong et al., *Nat. Med.*, 8: 793-800 (2002)). Expression of PD-L1 by tumor cells has been directly associated with their resistance to lysis by anti-tumor T-cells in vitro (Dong et al., *supra*; and Blank et al., *Cancer Res.*, 64: 1140-1145 (2004)). PD-1 knockout mice are resistant to tumor challenge (Iwai et al., *Int. Immunol.*, 17: 133-144 (2005)), and T-cells from PD-1 knockout mice are highly effective in tumor rejection when adoptively transferred to tumor-bearing mice (Blank et al., *supra*). Blocking PD-1 inhibitory signals using a monoclonal antibody can potentiate host anti-tumor immunity in mice (Iwai et al., *supra*; and Hirano et al., *Cancer Res.*, 65: 1089-1096 (2005)), and high levels of PD-L1 expression in tumors are associated with poor prognosis for many human cancer types (Hamanishi et al., *Proc. Natl. Acad. Sci. USA*, 104:

3360-335 (2007), Brown *et al*, *J. Immunol.*, 170: 1257-1266 (2003); and Flies *et al.*, *Yale Journal of Biology and Medicine*, 84(4): 409-421 (2011)).

[00126] Further, several studies have shown that interaction of PD-1 with its ligands (PD-L1 and PD-L2) promotes inhibition of lymphocyte proliferation *in vitro* and *in vivo*. Blockade of the PD-1/PD-L1 interaction may accordingly lead to enhanced tumor-specific T-cell immunity and therefore be helpful in clearance of tumor cells by the immune system. For example, in a murine model of aggressive pancreatic cancer, the therapeutic efficacy of PD-1/PD-L1 blockade was demonstrated (Nomi, T., *et al.* (2007) *Clin. Cancer Res.* 13: 2151-2157). Administration of either PD-1 or PDL1 directed antibody significantly inhibited tumor growth. Antibody blockade effectively promoted tumor reactive CD8⁺ T cell infiltration into the tumor resulting in the up-regulation of anti-tumor effectors including IFN gamma, granzyme B and perforin. Additionally, the authors showed that PD-1 blockade can be effectively combined with chemotherapy to yield a synergistic effect.

[00127] In view of the foregoing, strategies for inhibiting PD-1 activity to treat various types of cancer and for immunopotentialization (e.g., to treat infectious diseases) have been developed (see, e.g., Ascierto *et al.*, *Clin. Cancer. Res.*, 19(5): 1009-1020 (2013)).

Agents that inhibit PD-1 Signaling

[00128] Agents that inhibit PD-1 signaling for use in combination therapies of the present disclosure include those that bind to and block PD-1 receptors on T cells without triggering inhibitory signal transduction, agents that bind to PD-1 ligands to prevent their binding to PD-1, agents that do both, and agents that prevent expression of genes that encode either PD-1 or natural ligands of PD-1. Compounds that bind to natural ligands of PD-1 include PD-1 itself, as well as active fragments of PD-1, and in the case of the B7-H1 ligand, B7.1 proteins and fragments. Such antagonists include proteins, antibodies, anti-sense molecules and small organics.

[00129] In some embodiments, an agent that enhances T cell activity binds to human PD-1. In some embodiments an agent that enhances T cell activity binds to human PD-L1. In some embodiments, an agent that enhances T cell activity is a monoclonal antibody, or a fragment thereof. In some embodiments, an antibody agent that enhances T cell activity is a PD-1 or PD-L1 antibody or fragment thereof. Examples of such agents that enhance T cell activity by binding to human PD-1 include BGB-A317, BI 754091, IBI308, INCSHR-1210, JNJ-63723283, JS-001, MEDI-0680, MGA-012, nivolumab, PDR001, pembrolizumab, PF-06801591, REGN-2810, TSR-042, PDR-001, camrelizumab (HR-301210), BCD-100,

AGEN-2034, CS1001, Sym-021, LZM009, KN-035, AB122, genolimzumab (CBT-501), AK 104, GLS-010, any of the antibodies disclosed in WO2014/179664, PCT/US17/59618, PCT/US18/13029, and derivatives thereof. Examples of agents that enhance T cell activity by binding to human PD-L1 include atezolizumab, avelumab, CX-072, durvalumab, FAZ053, LY3300054, PD-L1 millamolecule, BGB-A333, SHR-1316, CK-301, or derivatives thereof.

[00130] In some embodiments, an agent that enhances T cell activity for use in combination therapies of the present disclosure is an antibody agent. In some embodiments, a PD-1 antibody agent binds an epitope of PD-1 which blocks the binding of PD-1 to any one or more of its putative ligands. In some embodiments, a PD-1 antibody agent binds an epitope of PD-1 which blocks the binding of PD-1 to two or more of its putative ligands. In some embodiments, the PD-1 antibody agent binds an epitope of a PD-1 protein which blocks the binding of PD-1 to PD-L1 and/or PD-L2. PD-1 antibody agents of the present disclosure may comprise a heavy chain constant region (Fc) of any suitable class. In some embodiments, a PD-1 antibody agent comprises a heavy chain constant region that is based upon wild-type IgG1, IgG2, or IgG4 antibodies, or variants thereof.

Tim-3

[00131] The protein T Cell Immunoglobulin and Mucin Domain-3 (TIM-3), also known as Hepatitis A Virus Cellular Receptor 2 (HAVCR2), is a Th1-specific cell surface protein that regulates macrophage activation and enhances the severity of experimental autoimmune encephalomyelitis in mice. TIM-3 is highly expressed on the surface of multiple immune cell types, including, for example, Th1 IFN- γ ⁺ cells, Th17 cells, natural killer (NK) cells, monocytes, and tumor-associated dendritic cells (DCs). TIM-3 also is highly expressed on “exhausted” or impaired CD8⁺ T-cells in a variety of chronic viral infections (e.g., HIV, HCV, and HBV) and in certain cancers. Putative ligands for TIM-3 include phosphatidylserine, galectin-9, high-mobility group protein 1 (HMGB1), and carcinoembryonic antigen cell adhesion molecule 1 (CEACAM1).

[00132] TIM-3 functions to regulate various aspects of the immune response. The interaction of TIM-3 and galectin-9 (Gal-9) induces cell death and in vivo blockade of this interaction exacerbates autoimmunity and abrogates tolerance in experimental models, strongly suggesting that TIM-3 is a negative regulatory molecule. In contrast to its effect on T-cells, the TIM-3-Gal-9 interaction can exhibit antimicrobial effects by promoting macrophage clearance of intracellular pathogen. Suppression of TIM-3 has been shown to enhance the pathological severity of experimental autoimmune encephalomyelitis.

Dysregulation of the TIM-3-galectin-9 pathway may play a role in chronic autoimmune diseases, such as multiple sclerosis. TIM-3 can promote clearance of apoptotic cells by binding phosphatidyl serine through its unique binding cleft.

[00133] Anti-TIM3 antibodies can promote antitumor immunity and suppress tumor growth. The current disclosure provides compositions and methods of cancer combination therapy. In some embodiments, the compositions and methods of the cancer combination therapy provided herein can include a PARP inhibitor and a TIM-3 inhibitory agent. In some embodiments, the PARP inhibitor is niraparib or pharmaceutically acceptable salts thereof. In some embodiments, the TIM-3 inhibitory agent can be a small molecule inhibitor. In some embodiments, the TIM-3 inhibitory agent can be an anti-TIM3 antibody or fragment thereof.

LAG-3

[00134] LAG-3 is upregulated following T-cell activation, and modulates T-cell function as well as T-cell homeostasis. The LAG-3/MHC class II interaction may play a role in down-regulating antigen-dependent stimulation of CD4⁺ T lymphocytes, higher expression of activation antigens such as CD25, and higher concentrations of cytokines such as interferon-gamma and interleukin-4. CD4⁺CD25⁺ regulatory T-cells (Treg) can also express LAG-3 upon activation and antibodies to LAG-3 inhibit suppression by induced Treg cells, both *in vitro* and *in vivo*, suggesting that LAG-3 contributes to the suppressor activity of Treg cells. Furthermore, LAG-3 can negatively regulate T-cell homeostasis by regulatory T cell-dependent and -independent mechanisms.

[00135] Subsets of conventional T-cells that are anergic or display impaired functions express LAG-3, and LAG-3⁺ T-cells are enriched at tumor sites and during chronic viral infections. In a self-tolerance/tumor mouse model where transgenic CD8⁺ T-cells were rendered unresponsive/anergic *in vivo*, LAG-3 blockade enhanced T-cell proliferation, T-cell recruitment and effector functions at the tumor site (Grosso et al., *J. Clin. Invest.*, 117: 3383-92 (2007)).

[00136] In addition, the interaction between LAG-3 and its major ligand, MHC class II, may play a role in modulating dendritic cell function (Andreae et al., *J Immunol.*, 168:3874-3880, 2002). Recent preclinical studies have documented a role for LAG-3 in CD8⁺ T cell exhaustion (Blackburn et al., *Nat Immunol.*, 10: 29-37, 2009), and blockade of the LAG-3/MHC class II interaction using a LAG-3Ig fusion protein may be useful for cancer therapy.

[00137] The current disclosure provides compositions and methods of cancer combination therapy. In some embodiments, the compositions and methods of the cancer combination

therapy provided herein can include a PARP inhibitor and a LAG-3 inhibitory agent. In some embodiments, the PARP inhibitor is niraparib or pharmaceutically acceptable salts thereof. In some embodiments, the LAG-3 inhibitory agent can be a small molecule inhibitor. In some embodiments, the LAG-3 inhibitory agent can be an anti-LAG-3 antibody or fragment thereof.

Indoleamine-pyrrole 2,3-dioxygenase (IDO)

[00138] IDO is an inducible enzyme that catalyzes the rate limiting step in tryptophan catabolism. This enzyme is overexpressed in response to IFN- γ in a variety of different malignancies. IDO can cause immunosuppression through breakdown of tryptophan in the tumor microenvironment and tumor-draining lymph nodes. The depletion of tryptophan and toxic catabolites can render effector T cells inactive and dendritic cells immunosuppressive. IDO inhibition can delay tumor growth, enhance dendritic cell vaccines, and synergize with chemotherapy through immune-mediated mechanisms. IDO inhibitory agents can include, but not limited to, d-1-methyl-tryptophan (d-1-MT), norharmane, rosmarinic acid, COX-2 inhibitors, 1-methyltryptophan, epacadostat and GDC-0919. IDO inhibitory agents can be anti-IDO antibodies.

[00139] The current disclosure provides methods and compositions of cancer combination therapy. In some embodiments, the combination therapy can include a PARP inhibitor and an IDO inhibitory agent. In some embodiments, the PARP inhibitor is niraparib or pharmaceutically acceptable salts thereof. In some embodiments, the IDO inhibitory agent can be a small molecule inhibitor. In some embodiments, the IDO inhibitory agent can be an anti-IDO antibody or fragment thereof.

Glucocorticoid-induced TNFR-related protein (GITR)

[00140] Glucocorticoid-induced TNFR-related protein (GITR) is a member of the tumor necrosis factor receptor (TNFR) superfamily, is a key regulator in a multitude of immune functions. GITR is expressed in most immune cell types including T regulatory cells (Tregs), naïve T cells, natural killer cells (NKs), and at low levels in B cells, macrophages, and dendritic cells. GITR signaling is triggered by its ligand (GITRL), which is expressed in antigen-presenting cells and endothelial cells and is involved in regulating T cell receptor-mediated cell death. Upregulation of GITR signaling in CD4⁺ and CD8⁺ T cells causes enhanced T cell expansion and cytokine production.

[00141] In some embodiments, molecules which induce GITR signaling may be used to activate antigen-specific T cells. For example, administration of the GITR agonist DTA-1 in

mice increases intratumor infiltration of CD4⁺ and CD8⁺ T cells. In addition to DTA-1, administration of the GITR agonists mGITRL and pGITRL (dimeric and pentameric versions of the GITR ligand, respectively), has been effective to induce tumor regression and activate CD8⁺ cells in tumor microenvironments.

Angiogenesis Inhibitors

[00142] Tumor growth and metastasis depend on new growth in the vascular network supporting the tumor. Vascular Endothelial Growth Factor A (VEGF) is secreted by tumor cells and acts on endothelial cells to stimulate angiogenesis during tumor growth. Anti-angiogenic antibodies such as VEGF blockers (e.g., bevacizumab) can increase numbers of antigen-specific T cells in solid tumors and enhance the efficiency of immunotherapy. For example, combination treatment of bevacizumab with either atezolizumab (inhibiting PD-L1) or ipilimumab (inhibiting CTLA-4) increases the number of intratumoral CD8⁺ cells. Other examples of VEGF blockers which may activate antigen-specific T cells in tumor microenvironments include pazopanib, sunitinib, sorafenib, axitinib, ponatinib, regorafenib, cabozantinib, vandetanib, ramucirumab, lenvatinib and *ziv*-aflibercept.

Other Agents

[00143] The present invention contemplates the use of any factor capable of activating or stimulating antigen-specific T cell activity in a tumor cell microenvironment. Further examples of agents which in some embodiments may be used to activate antigen-specific T cells (e.g., CD4⁺ or CD8⁺ cells) include flavonoids (e.g., flavonoid glycoside), lidocaine, lamotrigine, sulfamethoxazole, phenytoin, carbamazepine, sulfamethoxazole, phenytoin, allopurinol, paracetamol, mepivacaine, *p*-phenylenediamine, ciprofloxacin and moxifloxacin.

Combination Therapy of PARP Inhibitors and Antigen-Specific T Cell Activators

[00144] The current disclosure provides compositions and methods of cancer combination therapy involving a PARP inhibitor and an activator of antigen-specific T cells. In some embodiments, the compositions and methods of the cancer combination therapy provided herein can include niraparib and an agent which activates antigen-specific T cells in a tumor microenvironment. In some embodiments, the compositions and methods of the cancer combination therapy provided herein can include niraparib and at least one agent which activates antigen-specific T cells in a tumor microenvironment. The at least one agent which activates antigen-specific T cells can be small molecule inhibitors/agonists, proteins or protein fragments, antibodies, antibody fragments and/or polynucleotides. In some

embodiments, the Treg-inhibiting agents can be recombinant proteins. In some embodiments, the Treg-inhibiting agents can be delivered by gene therapy.

[00145] In some embodiments, the agent that activates antigen-specific T cells can be a cytokine. In some embodiments, the agent that activates antigen-specific T cells can be an inhibitor of PD-1 signaling. In some embodiments, the agent that activates antigen-specific T cells can be an angiogenesis inhibitor such as a VEGF inhibitor. In some embodiments, the agent that activates antigen-specific T cells can be an agent that triggers GITR signaling. In some embodiments, the agent that activates antigen-specific T cells can be an agent that inhibits CTLA-4 signaling. In some embodiments, the agent that activates antigen-specific T cells can be any agent capable of activating CD4⁺ and/or CD8⁺ cells in a tumor microenvironment.

[00146] In some embodiments, the agent that activates antigen-specific T cells in a tumor microenvironment can be an agent selected from the group consisting of: pembrolizumab, nivolumab, PDR001, REGN2810 (SAR-439684), BGB-A317, BI 754091, IBI308, INCSHR-1210, JNJ-63723283, JS-001, MEDI0680 (AMP-514), MGA-012, PF-06801591, REGN-2810, TSR-042, PDR-001, camrelizumab (HR-301210), BCD-100, AGEN-2034, CS1001, Sym-021, LZM009, KN-035, AB122, genolimzumab (CBT-501), AK 104, GLS-010, atezolizumab, durvalumab, avelumab, CX-072, FAZ053, LY3300054, PD-L1 millamolecule, BGB-A333, SHR-1316, CK-301, or derivatives thereof, LY3300054, DTA-1, mGITRL, pGITRL, ipilimumab, ipilimumab, interleukin-2 (IL-2), IL-1, IL-7, IL-15, IL-12, IL-18, interferon- α (IFN- α), CXCL16, bevacizumab, pazopanib, sunitinib, sorafenib, axitinib, ponatinib, regorafenib, cabozantinib, vandetanib, ramucirumab, lenvatinib, *ziv*-aflibercept, moxifloxacin, flavonoids (e.g., flavonoid glycoside), lidocaine, lamotrigine, sulfamethoxazole, phenytoin, carbamazepine, sulfamethoxazole, phenytoin, allopurinol, paracetamol, mepivacaine, *p*-phenylenediamine, ciprofloxacin and moxifloxacin.

Tumor-Associated Macrophages (TAMs)

[00147] While a few tumors may grow as cell suspensions, for example leukemia and ascites tumor, most tumors form into solid masses of tissues that can be mainly composed of malignant cells and stroma. The tumor stroma which can be largely produced from normal host tissue includes matrix components, blood vessels as well as inflammatory cells. Malignant cells may alter the properties their stroma by cell-to-cell contact, soluble factors and/or by modification of the extra-cellular matrix (ECM) in support of their growth. On the

other hand, in response to malignant cells, the stromal cells can modify the phenotypes, invasiveness and metastatic capacity of tumor cells, typically promoting their progression. A vast diversity of molecules are produced by different cellular components within the complex tumor microenvironment such as cytokines, chemokines, growth factors and proteases, which may positively or negatively influence tumor survival and growth.

[00148] Tumor-associated macrophages (TAMs) can be the main population of inflammatory cells in tumor stroma. TAMs can be originated from peripheral blood monocytes in the blood circulation which can be recruited into the tumor mass by tumor-derived chemoattractants and then differentiated into tissue macrophages. TAMs generally may fail to express pro-inflammatory cytokines for T helper type 1 (Th1) responses but can be excellent producers of immunosuppressive cytokines for Th2 responses. As TAMs generally can exhibit low antigen-presenting and co-stimulating capacity, they ordinarily fail to activate T cell-mediated adaptive immunity. Therefore, unlike M1 macrophages, which can be highly microbicidal and tumoricidal, the M2-like TAMs can be immunosuppressive and facilitate tumor progression.

Macrophage classification and the heterogeneity of TAM phenotypes

[00149] According to the different phenotypes and distinct patterns of gene expression, macrophages can be subdivided into type I and type II macrophages. The type I macrophages (M1) have a phenotypic pattern of high interleukin-12 (IL-12), low IL-10, or low IL-4/-13. The propensity of M1 can be increased in response to opsonized ligands and toll-like receptor (TLR) engagement. The typical type I macrophages can play an indispensable role in both innate and acquired immunity, and therefore they can have a proinflammatory characteristic. They can provide an in-front defense line against different kinds of pathogens and malignant cells via phagocytosis and/or induction of antibody-dependent cellular cytotoxicity. Activated macrophages can have the capability to recognize and bind to tumorigenic cells, a process that lead to the subsequent lysis or phagocytosis of tumor cells.

[00150] The type II macrophages (M2), which have the IL-10+/IL-12- phenotype, can have an immunosuppressive characteristic. TAMs can express typical M2 markers and have defective expression of some proinflammatory cytokines, such as IL-12, tumor necrosis factor-alpha (TNF- α), CC chemokine ligand 3 (CCL3), and IL-1 but high expression of IL-10, hence the TAMs can be M2-like. IL-12, typically produced by macrophages, is a cytokine in immune resistance against pathogens. IL-12 is known as a T cell stimulating factor, which can stimulate the growth and function of T cells. It can also facilitate the

production of IFN- γ and TNF- α and alleviate IL-4-mediated suppression of IFN- γ . Therefore, lack of IL-12 production may result in dysfunction of the antitumor responses of macrophages. It was observed that the expression of IL-12 gene is partially controlled by members of nuclear factor-kappa B (NF- κ B) family and failure to produce IL-12 may be associated with defective activation of p50/p65 NF- κ B in TAMs. This defective activation of NF- κ B in TAMs can also correlate with impaired expression of NF- κ B-dependent inflammatory functions, for example, the expression of TNF- α , IL-1 and other cytotoxic mediators. Moreover, high level production of IL-10 triggered by CCL2 or other tumor-derived chemotactic factors (TDCFs) may play a role in defective IL-12 production. By interrupting the communication between tumor cells and their stroma, TAMs could be switched towards IFN- γ production and tumor rejection. Therapeutics targeted to block the production of IL-10 as well as other immunosuppressive cytokines in tumor sites may restore the antitumor functions of TAMs.

Macrophage polarization and M2-polarized TAMs

[00151] In response to different intercellular signals, macrophages can be polarized into different phenotypes. The type I macrophages which have the antitumor activity can be differentiated from monocytes exposed to certain factors such as granulocyte-macrophage colony-stimulating factor (GM-CSF), IFN- γ , LPS and some bacterial products; whereas macrophage colony-stimulating factor (M-CSF), IL-4/-13/-10 and some immunosuppressive agents trigger macrophages to differentiate toward type II macrophages. Treatment of monocytes with IFN- γ alone or in combination with TNF- α , GM-CSF or PPD, before/after co-cultured with tumor (HPC-4) cells can enhance the *de novo* production of molecular factors related to antitumor activity of macrophages. The activation of NF- κ B turns on the inflammatory repertoire of macrophages, leading to the expression of proinflammatory cytokines. Defective NF- κ B activation could inhibit the transcription of some key cytokines related to M1 phenotypes and the production of cytotoxic mediators. As an upstream component of NF- κ B activation, TLR/IL-1R signaling may also play a role in regulating macrophage polarization. Additionally, the Tie-2/Ang-2 pathway, the TRIF/TBK1/IRF3 pathway, and hypoxia-induced pathway can be possible pathways in regulating the “shift” in macrophage phenotypes.

[00152] TAMs can be triggered into M2-like phenotypes by tumor-derived factors, such as cytokines, growth factors, chemotactic molecules and proteases which may either up-regulate or down-regulate the expression of macrophage effector molecules, thus influence the

functions of macrophages. Many tumor-derived molecules can deactivate or suppress the cytotoxic activity of TAMs, including IL-4, IL-6, IL-10, macrophage-derived chemotactic factor (MDF), transforming growth factor beta1 (TGF- β 1), prostaglandin E2 (PGE2), and M-CSF. One of the players in the regulation of macrophage infiltration can be monocyte chemotactic protein-1 (MCP-1/CCL2) and related CC chemokines. Accumulation of TAMs in primary tumors can be correlated with the level of CCL2 expression. CCL2 has been shown to be produced by a variety of human tumors, such as melanoma, malignant glioma, ovarian cancer and meningioma. MCP-2 and MCP-3, also known as CCL8 and CCL7, are the other two members of the CCL2 family which were isolated from tumor cell lines. In addition, breast carcinoma at advanced stage was reported to be associated with increased expression of CCL5, and CCL5 receptor (CCR5) was detected on TAMs. CCL5 may play a role in macrophage migration, tumor progression, and protumorigenic activity.

[00153] TAMs can promote tumor angiogenesis by secreting a vast diversity of factors, including vascular endothelial growth factor (VEGF), granulocyte colony-stimulating factor (G-CSF), basic fibroblasts growth factor (bFGF), insulin-like growth factor-I (IGF-I), platelet derived growth factor (PDGF), transforming growth factor- β (TGF- β), tumor necrosis factor- α (TNF- α), IL-1, IL-6, IL-8, substance P, prostaglandins and other kinds of monokines. Proteolytic enzymes, including matrix metalloproteinases (MMPs), which may be one of the key molecular factors in angiogenesis, can be secreted by TAMs. Synthetic inhibitors against MMPs (MMPIs) have been developed and put into clinical trials as anti-cancer therapeutics. Moreover, TNF- α can be largely produced by TAMs and is also a hypoxia-inducible pro-angiogenic cytokines. TNF- α can initiate a cascade of signal transductions upon its secretion, leading to the production of other factors which in turn function positively on angiogenesis.

TAM-targeted antitumor strategies

[00154] The function of TAMs can depend on their accumulation and activation in tumor tissues, therefore, TAM-targeted anti-tumor approaches can based on the following four strategies: 1) inhibiting macrophage recruitment; 2) suppressing TAM survival; 3) enhancing M1 tumoricidal activity of TAMs; and 4) blocking M2 tumor-promoting activity of TAMs.

[00155] Chemoattractants released by tumor cells can facilitate the recruitment of macrophages into tumor tissues. These chemoattractants can include, but not limited to, CCL2, macrophage colony-stimulating factor (M-CSF, or also known as colony stimulating factor 1, CSF-1), CCL5, C-X-C motif chemokine ligand-12 (CXCL-12) and vascular endothelial growth factor (VEGF). Binding agents targeting these chemoattractants or

neutralizing their corresponding receptors can be used to inhibit the recruitment of macrophages. For example, binding agents of CCL2/CCR2 can include pharmaceutical inhibitors (e.g., trabectedin, RS102895, and PF-04136309) and antibodies (e.g., CNTO888 and MLN1202). Anti-IL6 antibody siltuximab may also inhibit macrophage infiltration in tumour tissue via declining the plasma level of some chemoattractants such as CCL2, VEGF and CXCL-12. Binding agents of M-CSF/M-CSFR can include anti-M-CSFR (or anti-CSF-1R) antibodies JNJ-28312141, GW2580, IMC-CS4 (LY3022855), emactuzumab, AMG820, MCS110, small molecule inhibitors pexidartinib, PLX7846, linifanib, OSI-930, CEP-32496, ARRY-382, and JNJ-40346527. Other anti-M-CSF/M-CSFR agents include PLX3397, PLX6134, PD-0360324, or FPA008. Inhibitors of other chemoattractants (e.g., VEGF, CXCL-12 and CCL5) and their receptors can be useful for TAM depletion and tumor rejection. In addition, inhibitors of hypoxia-inducible factors (HIFs), which are transcriptional activators for *VEGF* and *CXCR4* (encoding for CXCL-12 receptor) genes, may be anti-tumor candidates for their potential to inhibit angiogenesis and macrophage recruitment.

[00156] To suppress TAM survival, two approaches can be used. One can be to directly induce macrophage apoptosis using chemical reagents, immunotoxin-conjugated monoclonal antibodies or attenuated bacteria; the other can be to trigger the immune cells, for example T lymphocytes, to recognize and abrogate TAMs. Bisphosphonates (e.g., clodronate, zoledronic acid and dichloromethylene bisphosphonate), generally packed in liposomes, can be used for macrophage depletion by inducing apoptosis. Trabectedin can activate caspase-8-dependent apoptosis and selectively deplete monocytes including TAMs. Dasatinib, a Src kinase inhibitor, can reduce MMP9⁺ macrophage density and inhibit MMP9 expression in the tumor environment. To deplete TAMs by targeting their surface molecules with immunotoxin-conjugated agents can be another approach for tumor therapy. The surface proteins of TAMs that may be targets can include scavenger receptor-A, CD52 and folate receptor β (FR β). Bacteria can also take macrophages as targets. For example, *Shigella flexneri* infection could selectively induce the apoptosis of macrophages, and a single injection of an attenuated strain of *Shigella flexneri* to tumor-bearing mice resulted in the apoptosis of TAMs, followed by a 74% reduction in size of tumors. Other bacteria that can be used for TAM-targeted immunotherapy include, but not limited to, *Salmonella typhimurium*, *Listeria monocytogens*, *Chlamydia psittaci* and *Legionella pneumophila*. As mentioned above herein, another available approach for TAM suppression can be to evoke

acquired immune responses, in which cytotoxic T lymphocytes can act as the scavengers of TAMs because they can naturally target the membrane molecules of macrophages. In other words, up-regulating the membrane molecules that could be recognized by T cells in TAMs can be a method of TAM depletion. One such molecule is legumain, a lysosomal protease highly expressed in many human tumors and M2-like TAMs. A legumain based DNA vaccine can activate dendritic cells, which then triggered multi-step reactions including the antigen presenting, co-stimulation of cytotoxic CD8⁺ T cells and the specific abrogation of legumain-expressing TAMs. Another membrane protein involved in T-cell-mediated TAM depletion can be CD1d, a target of V α 24-invariant natural killer T (NKT) cells. Agents that can promote the expression of CD1d in TAMs may improve the tumoricidal function of NKT cells. One such agent can be retinoic acid, which can strongly up-regulate the CD1d expression in macrophages and is now used as a standard therapeutic drug for high-risk neuroblastoma in clinic.

[00157] To enhance M1 tumoricidal activity of TAMs, TAMs can be re-polarized from M2 type to M1 type. Manipulating transcription factors and their up-/down-stream regulators can contribute to targeted tumor therapy. These transcription factors can include M1-promoting modulators STAT1 and NF- κ B (p50p65 heterodimer). NF- κ B activating agents can include the agonists of Toll-like receptor (TLRs), anti-CD40 mAb and anti-IL-10R mAb. The TLR agonists can be diverse, including PolyI:C (for TLR3), lipopolysaccharide (LPS) and monophosphoryl A (for TLR4), imiquimod and R-848 (for TLR7), and CpG-oligodeoxynucleotide (CpG-ODN, for TLR9). In addition, anti-CD40 mAb may be used to promote TLR9 to respond to CpG-ODN. Agonists of STAT1 can include IFN- α (FDA approved), IFN- β (FDA approved) and IFN- γ . Agonistic anti-CD40 antibodies can include CP-870893 and RO7009789. In addition, other factors such as GM-CSF, IL-12, IL-2 and IL-15 may be used to induce M1 function. These factors may be delivered into a subject by recombinant proteins or gene therapy. A peptide drug, thymosin- α 1 (T α 1) can also be used to induce M1 function. Tumor cells can express CD47, a 'don't eat me' signal that, via interaction with macrophage surface receptor SIRP α can prevent phagocytosis by macrophages. Thus, interference with the SIRP α -CD47 pathway, for example, using antagonistic antibodies, can activate macrophage-mediated antibody-dependent cellular phagocytosis (ADCP), which subsequently results in functional skewing of macrophages in an M1 direction that is associated with antitumor activity. CD47 antagonists can include anti-CD47 antibodies Hu5F9-G4 and CC-90002, and CD47-Fc fusion protein TTI-621.

Moreover, B cell-macrophage interactions can promote PI3K γ - and Bruton tyrosine kinase (BTK)-dependent macrophage M2 polarization. Inhibitors of Bruton tyrosine kinase, such as ibrutinib, can reprogram TAM toward M1 phenotype. Vitamin-D-binding protein (macrophage activating factor), for example EF-022, can also promote tumoricidal activity of macrophage and prevent angiogenesis in tumors.

[00158] To block M2 tumor-promoting activity of TAMs, M2-promoting transcription factors STAT3 and STAT6 can be inhibited. STAT3 inhibitors can include WP1066, tyrosine kinase inhibitors sunitinib and sorafenib, STA-21, IS3 295 and S3I-M2001. STAT6 inhibitors can include AS1517499, leflunomide and TMC-264. Several up-/down-stream mediators of STAT6 can act as modulators of TAM function. These modulators include phosphatidylinositol 3-kinase (PI3K), Src homology 2-containing inositol-5'-phosphatase (SHIP), Krüppel-like factor 4 (KLF4) and c-Myc. Other proteins that may promote M2 function and thus can be targeted for cancer therapy include, but not limited to, peroxisome proliferator-activated receptor (PPARs), HIFs, Ets family member 2 (Ets2), Decoy receptor (DcR3) and mammalian target of rapamycin (mTOR). Several anti-tumour drugs that can suppress M2 macrophages can include histidine-rich glycoprotein (HRG), copper chelate (CuNG), 5,6-dimethylxanthenone-4-acetic acid (MDXAA), vadimezan (ASA404), cisplatin, silibinin, CNI-1493, and proton pump inhibitor pantoprazole (PPZ). Other agents that can repolarize M2-like TAMs include, but not limited to, anti-IL-1 α antibody xilonix.

Combination Therapy of PARP Inhibitors and TAM-Targeting Agents

[00159] A TAM-targeting agent or TAM inhibitory agent can be used interchangeably herein. A TAM inhibitory agent can be an agent that 1) can inhibit macrophage recruitment; 2) can suppress TAM survival; 3) can enhance M1 tumoricidal activity of TAMs; and 4) can block M2 tumor-promoting activity of TAMs. A TAM inhibitory agent can decrease or eliminate the population of TAM. A TAM inhibitory agent can also regulate the function of TAM.

[00160] The current disclosure provides compositions and methods of cancer combination therapy of a PARP inhibitor and a TAM-targeting agent. In some embodiments, the compositions and methods of the cancer combination therapy provided herein can include niraparib and pharmaceutically acceptable salts thereof and a TAM-targeting agent. In some embodiments, the compositions and methods of the cancer combination therapy provided herein can include niraparib and at least one TAM-targeting agent. The TAM-targeting agents can be small molecule inhibitors/agonists, proteins or protein fragments, antibodies or

antibody fragments, or bacteria. In some embodiments, the TAM-targeting agents can be recombinant proteins. In some embodiments, the TAM-targeting agents can be delivered by gene therapy.

[00161] In some embodiments, the TAM-targeting agents can be agents that inhibit recruitment of macrophages. In some embodiments, the TAM-targeting agent can be an inhibitor of a chemoattractant selected from the group consisting of CCL2, M-CSF (CSF-1), CCL5, CXCL-12 and VEGF. In some embodiments, the TAM-targeting agent can be an antibody targeting M-CSF/M-CSFR (or CSF-1/CSF-1R). In some embodiments, the compositions and methods of the cancer combination therapy provided herein can include niraparib and at least one TAM-targeting agent selected from the group consisting of trabectedin, RS102895, PF-04136309, CNTO888, MLN1202, siltuximab, JNJ-28312141, GW2580, IMC-CS4 (LY3022855), emactuzumab, AMG820, pexidartinib, linifanib, OSI-930, CEP-32496, PLX7846, ARRY-382, JNJ-40346527, MCS110, PLX3397, PLX6134, PD-0360324, FPA008, and any combinations thereof.

[00162] In some embodiments, the TAM-targeting agents can be agents that suppress the M2-like TAM survival. In some embodiments, the TAM-targeting agents can be chemical reagents, immunotoxin-conjugated monoclonal antibodies or attenuated bacteria that directly induce apoptosis of TAMs. In some embodiments, the TAM-targeting agents can be agents that can trigger immune cells to abrogate TAMs. In some embodiments, the TAM-targeting agents can be bisphosphonates. In some embodiments, the compositions and methods of the cancer combination therapy provided herein can include niraparib and at least one TAM-targeting agent selected from the group consisting of clodronate, zoledronic acid, dichloromethylene bisphosphonate, trabectedin, dasatinib, retinoic acid, and any combinations thereof. In some embodiments, the compositions and methods of the cancer combination therapy provided herein can include niraparib and at least one attenuated bacteria selected from the group consisting of *Shigella flexneri*, *Salmonella typhimurium*, *Listeria monocytogens*, *Chlamydia psittaci*, *Legionella pneumophila*, and any combinations thereof.

[00163] In some embodiments, the TAM-targeting agents can be agents that enhance the tumoricidal activity of M1 type macrophages or repolarizing the M2 type macrophages into M1 type. In some embodiments, the TAM-targeting agents can be NF- κ B activating agents. In some other embodiments, the TAM-targeting agents can be STAT1 activating agents. In some embodiments, the TAM-targeting agents can include the agonists of Toll-like receptor

(TLRs), anti-CD40 mAb and anti-IL-10R mAb. In some embodiments, the TAM-targeting agents can be PolyI:C (for TLR3), lipopolysaccharide (LPS) or monophosphoryl A (for TLR4), imiquimod or R-848 (for TLR7), or CpG-oligodeoxynucleotide (CpG-ODN, for TLR9), or any combinations thereof. In some embodiments, the TAM-targeting agents can be IFN- α , IFN- β , IFN- γ , GM-CSF, IL-12, IL-2, IL-15, or any combinations thereof. In some embodiments, the compositions and methods of the cancer combination therapy provided herein can include niraparib and at least one TAM-targeting agent selected from the group consisting of anti-CD40 mAb, anti-IL-10R mAb, CD47 antagonists (e.g., anti-CD47 antibodies Hu5F9-G4 and CC-90002, and CD47-Fc fusion protein TTI-621), PolyI:C, LPS, monophosphoryl A, imiquimod, R-848, CpG-ODN, IFN- α , IFN- β , IFN- γ , GM-CSF, IL-12, IL-2, IL-15, T α 1, ibrutinib, EF-022 and any combinations thereof.

[00164] In some embodiments, the TAM-targeting agents can be agents that block M2 tumor-promoting activity of TAMs. In some embodiments, the TAM-targeting agents can include inhibitors of STAT3 or STAT6. In some embodiments, the TAM-targeting agents can include other M2 modulators, including PI3K, SHIP, KLF4, c-Myc, PPARs, HIFs, Ets2, DcR3, and mTOR. In some embodiments, the compositions and methods of the cancer combination therapy provided herein can include niraparib and at least one TAM-targeting agent selected from the group consisting of WP1066, sunitinib, sorafenib, STA-21, IS3 295, S3I-M2001, AS1517499, leflunomide, TMC-264, histidine-rich glycoprotein (HRG), copper chelate (CuNG), 5,6-dimethylxanthenone-4-acetic acid (MDXAA), vadimezan (ASA404), cisplatin, silibinin, proton pump inhibitor pantoprazole (PPZ), CNI-1493, anti-IL-1 α antibody xilonix, EF-022 and any combinations thereof.

Regulatory T Cells

[00165] Forkhead box protein 3 (Foxp3)-expressing regulatory T cells (Treg cells) can function in the regulation of immune responses and in the maintenance of immunological self-tolerance. These cells can be therapeutic targets for autoimmune diseases and cancer. Treg cells can be characterized by the expression of the high-affinity interleukin-2 (IL-2) receptor α -chain (IL-2R α ; also known as CD25) and the X-linked gene *Foxp3*, encoding the transcription factor Foxp3, which serves as a lineage specification factor for the development and function of CD4⁺ CD25⁺ Treg cells.

[00166] Treg can infiltrate tumor tissues, and Treg cell-mediated suppression of tumor-associated antigens can be a potential mechanism to explain the failure of antitumor immunity. Tumor-induced expansion of Treg cells can be an obstacle to successful cancer

immunotherapy. Treg cells can selectively interfere with the release of cytolytic granules by cytotoxic T lymphocytes (CTLs) in a reversible and TGF β -dependent manner, thereby attenuating CTL-mediated cytotoxicity without detectably affecting CTL priming or differentiation. Local or systemic interference with suppressor pathways of CTL activity, such as cytokines, Toll-like receptor agonists, or Treg cell depletion, may be effective in tumor therapy.

[00167] One approach used to inhibit Treg cells can be Treg depletion. Some conventional chemotherapy agents can affect adaptive immune system, resulting in the inhibition of Treg function or viability. For example, these agents can include, but not limited to, cyclophosphamide and paclitaxel. Angiogenesis inhibition can overcome various immunosuppressive networks including Treg. The immunogenic effects of more specific molecularly targeted therapeutic anticancer agents can also impair Treg functions. These targeted therapeutic anticancer agents include some tyrosine kinase inhibitors, but not limited to, imatinib (Gleevec; Novartis), sunitinib (Sutent; Pfizer), sorafenib (Nexavar; Bayer/Onyx), dasatinib, and temozolomide. Some of these tyrosine inhibitors can block STAT3 and STAT5 signaling and decrease Treg cell frequency, or limit infiltration by Treg cells while inhibiting STAT3 activity. Strategies for Treg cell depletion can include agents that target IL-2R which is upregulated in Treg cells. These agents can include anti-CD25 monoclonal antibody daclizumab (Zenapax; PDL BioPharma) and denileukin diftitox (Ontak; Esai).

[00168] A second approach to inhibit Treg cells can include the targeting of molecules that are involved in Treg cell migration. Treg cells from cancer patients, as compared to healthy subjects, can be characterized by a distinct expression profile of chemokine receptors, such as CCR4, CXCR4, and CCR5, which can facilitate their migration into tumors in response to the corresponding chemokine ligands derived from tumor microenvironment. Treg cell trafficking to tumors can be triggered by a cohort of tumor-associated chemokines or hypoxia-induced factors, including CCL22, CCL17, CXCL12, CCL28, and VEGF. CC motif chemokine 22 (CCL22) blockage can reduce Treg cell-mediated tumor trafficking. Agents that block CCL22 can include, but are not limited to, casuarinin and fucoidan. CXCR4 and CXCL12 can contribute to Treg cell migration to the bone marrow. Antagonists of CXCR4, for example AMD3100, can promote antitumor immunity. Antagonists of CCR4, for example anti-CCR4 antibody mogamulizumab, can prevent the interaction of CCL22/CCL17 with their receptor. Other chemokine receptors such as CCR7 may also play a role in Treg migration.

[00169] Another approach to inhibit Treg cells can include using antibodies to target molecules constitutively expressed by Treg leading to their functional inhibition. Ipilimumab and tremelimumab, antibodies that are specific for cytotoxic T lymphocyte antigen 4 (CTLA4) can block an inhibitory signal for activated T cells, thereby bolstering T cell responses and potentiating tumor destruction. The mechanism of CTLA4-specific antibodies may be inhibiting Treg-dependent immune suppression. Adenosine A_{2A} receptor can inhibit T cell responses in part by upregulating Foxp3 expression in CD4⁺ T cells. Engagement of adenosine A_{2A} receptor by adenosine can result in Treg cell induction which may lead to self-amplifying loop within the tumor. Inhibition of adenosine A_{2A} receptor can be achieved using antibodies to block adenosine or using adenosine analogues. GITR can be constitutively expressed by Treg, but it is also detected, albeit at lower levels, on CD4⁺ and CD8⁺ effector T cells. Stimulation by agonistic antibodies to either GITR or GITR ligand can have a dual effect leading to suppression of Treg activity and enhanced proliferation of effector T cells and possible resistance to Treg-mediated suppression. OX40, a costimulatory molecule of the TNF receptor family, can be constitutively expressed on Treg and transiently expressed on activated T cells. Activation of OX40 signaling by an agonistic anti-OX40 mAb can inhibit the suppressive activity of Treg.

[00170] Treg can express various TLRs and notably high levels of TLR4, TLR5, TLR7 and TLR8. TLR 8 activation by its natural or synthetic ligands can inhibit Treg function and enhance *in vivo* tumor immunity. Appropriate TLR stimulation might therefore be an important tool for vaccination. Treg can produce adenosine via catabolism of adenine nucleotides (ATP, ADP and AMP) by extracellular ectonucleotidases, CD39 and CD73. Adenosine is a major immunosuppressive factor that may participate in the immunosuppressive activity of Foxp3⁺ T cells. Low molecular weight inhibitors and adenosine receptor antagonists can be used to block adenosine-mediated immune suppression. Exemplary adenosine receptor antagonists can include caffeine, theophylline, theobromine, and 8-phenylxanthines. Inhibition of CD39 with enzymatic inhibitors can block Treg function and improve the effects of chemotherapy. In addition, a peptide inhibitor of Foxp3 (Peptide P60) can impair Treg activity and improve vaccine efficacy.

Combination Therapy of PARP Inhibitors and Treg-Inhibiting Agents

[00171] A Treg inhibitory agent or Treg inhibitor or Treg-inhibiting agents can refer to an agent that: (1) inhibits or decreases the activity or function of a regulatory T cell, (2) decreases the population of regulatory T cells in a subject (in one embodiment, the decrease

can be temporary, for example, for a few hours, a day, a few days, a week, or a few weeks), or (3) substantially ablates or eliminates the population of regulatory T cells in a subject (in one embodiment, the ablation or elimination can be temporary, for example, for a few hours, a day, a few days, a week, or a few weeks). A Treg inhibitor can decrease the suppression of immune system activation and can decrease prevention of self-reactivity. Exemplary Treg inhibitors may include, but are not limited to, a compound, antibody, fragment of an antibody, or chemical that targets a Treg cell surface marker (such as CD25, CD4, CD28, CD38, CD62L (selectin), OX-40 ligand (OX-40L), CTLA4, CCR4, CCR8, FOXP3, LAG3, CD103, NRP-1, glucocorticoid-induced TNF receptor (GITR), galectin-1, TNFR2, or TGF- β R1). In certain embodiments, a Treg inhibitor targets a Treg cell surface marker that is involved in Treg activation such that the Treg inhibitor prevents Treg activation. A Treg inhibitor may include, but is not limited to, antibodies, fusion proteins, ONTAK, HuMax-Tac, Zenapax, or MDX-010, aptamers, siRNA, ribozymes, antisense oligonucleotides, and the like. The administration of a Treg inhibitor or derivatives thereof can block the action of its target, such as a Treg cell surface marker. A Treg inhibitor can have an attached toxic moiety such that upon internalization of the inhibitor, the attached toxic moiety can kill the T regulatory cell.

[00172] The current disclosure provides compositions and methods of cancer combination therapy of a PARP inhibitor and a Treg-inhibiting agent. In some embodiments, the compositions and methods of the cancer combination therapy provided herein can include niraparib and pharmaceutically acceptable salts thereof and a Treg-inhibiting agent. In some embodiments, the compositions and methods of the cancer combination therapy provided herein can include niraparib and at least one Treg-inhibiting agent. The Treg-inhibiting agents can be small molecule inhibitors/agonists, proteins or protein fragments, antibodies or antibody fragments. In some embodiments, the Treg-inhibiting agents can be recombinant proteins. In some embodiments, the Treg-inhibiting agents can be delivered by gene therapy.

[00173] In some embodiments, the Treg-inhibiting agents can be agents that can lead to Treg cell death. In some embodiments, the Treg-inhibiting agents can be chemotherapy agents. In some embodiments, the Treg-inhibiting agents can be angiogenesis inhibitors. In some embodiments, the Treg-inhibiting agents can be tyrosine kinase inhibitors. In some embodiments, the Treg-inhibiting agents can be STAT3 or STAT5 inhibitors. In some embodiments, the Treg-inhibiting agents can be the agents selected from the group consisting of cyclophosphamide, paclitaxel, imatinib (Gleevec; Novartis), sunitinib (Sutent; Pfizer),

sorafenib (Nexavar; Bayer/Onyx), dasatinib, temozolomide, daclizumab (Zenapax; PDL BioPharma), denileukin diftitox (Ontak; Esai), and any combinations thereof.

[00174] In some embodiments, the Treg-inhibiting agents can be agents that inhibit Treg migration. In some embodiments, the Treg-inhibiting agents can be CC motif chemokine 22 (CCL22) inhibitors. In some embodiments, the Treg-inhibiting agents can be CCR4, CCR5, or CCR7 inhibitors. In some embodiments, the Treg-inhibiting agents can be antagonists of CXCR4. In some embodiments, the Treg-inhibiting agents can be agents selected from the group consisting of AMD3100 or mogamulizumab, casuarinin, fucoidan, and any combinations thereof.

[00175] In some embodiments, the Treg-inhibiting agents can be agents that block Treg function. In some embodiments, the Treg-inhibiting agents can be antibodies that target the constitutively expressed molecules of Treg. In some embodiments, the Treg-inhibiting agents can be antibodies against cytotoxic T lymphocyte antigen 4 (CTLA4), such as ipilimumab and tremelimumab. In some embodiments, the Treg-inhibiting agents can be the agents that inhibit adenosine A_{2A} receptor. In some embodiments, the Treg-inhibiting agents can be antibodies that block adenosine or can be adenosine analogues. In some embodiments, the Treg-inhibiting agents can be immune agonists, including anti-CTLA4, anti-OX40, or anti-GITR antibodies. In some embodiments, the Treg-inhibiting agent can be a peptide inhibitor of Foxp3 Peptide P60.

Other combination therapies

[00176] Therapeutic methods of the invention can be combined with additional immunotherapies and therapies. For example, when used for treating cancer, inhibitors of the invention can be used in combination with conventional cancer therapies, such as, e.g., surgery, radiotherapy, chemotherapy or combinations thereof, depending on type of the tumor, patient condition, other health issues, and a variety of factors. In certain aspects, other therapeutic agents useful for combination cancer therapy with the inhibitors of the invention include anti-angiogenic agents. Many anti-angiogenic agents have been identified and are known in the art, including, e.g., TNP-470, platelet factor 4, thrombospondin-1, tissue inhibitors of metalloproteases (TIMP1 and TIMP2), prolactin (16-Kd fragment), angiostatin (38-Kd fragment of plasminogen), endostatin, bFGF soluble receptor, transforming growth factor beta, interferon alpha, soluble KDR and FLT-1 receptors, placental proliferin-related protein, as well as those listed by Carmeliet and Jain (2000). In one embodiment, the inhibitors of the invention can be used in combination with a VEGF antagonist or a VEGF

receptor antagonist such as anti-VEGF antibodies, VEGF variants, soluble VEGF receptor fragments, aptamers capable of blocking VEGF or VEGFR, neutralizing anti-VEGFR antibodies, inhibitors of VEGFR tyrosine kinases and any combinations thereof (e.g., anti-hVEGF antibody A4.6.1, bevacizumab or ranibizumab).

[00177] Non-limiting examples of chemotherapeutic compounds which can be used in combination treatments of the present invention include, for example, aminoglutethimide, amsacrine, anastrozole, asparaginase, bcg, bicalutamide, bleomycin, buserelin, busulfan, camptothecin, capecitabine, carboplatin, carmustine, chlorambucil, cisplatin, cladribine, clodronate, colchicine, cyclophosphamide, cyproterone, cytarabine, dacarbazine, dactinomycin, daunorubicin, dienestrol, diethylstilbestrol, docetaxel, doxorubicin, epirubicin, estradiol, estramustine, etoposide, exemestane, filgrastim, fludarabine, fludrocortisone, fluorouracil, fluoxymesterone, flutamide, gemcitabine, genistein, goserelin, hydroxyurea, idarubicin, ifosfamide, imatinib, interferon, irinotecan, irinotecan, letrozole, leucovorin, leuprolide, levamisole, lomustine, mechlorethamine, medroxyprogesterone, megestrol, melphalan, mercaptopurine, mesna, methotrexate, mitomycin, mitotane, mitoxantrone, nilutamide, nocodazole, octreotide, oxaliplatin, paclitaxel, pamidronate, pentostatin, plicamycin, porfimer, procarbazine, raltitrexed, rituximab, streptozocin, suramin, tamoxifen, temozolomide, teniposide, testosterone, thioguanine, thiotepa, titanocene dichloride, topotecan, trastuzumab, tretinoin, vinblastine, vincristine, vindesine, and vinorelbine.

[00178] These chemotherapeutic compounds may be categorized by their mechanism of action into, for example, following groups: anti-metabolites/anti-cancer agents, such as pyrimidine analogs (5-fluorouracil, floxuridine, capecitabine, gemcitabine and cytarabine) and purine analogs, folate antagonists and related inhibitors (mercaptopurine, thioguanine, pentostatin and 2-chlorodeoxyadenosine (cladribine)); antiproliferative/antimitotic agents including natural products such as vinca alkaloids (vinblastine, vincristine, and vinorelbine), microtubule disruptors such as taxane (paclitaxel, docetaxel), vincristin, vinblastin, nocodazole, epothilones and navelbine, epidipodophyllotoxins (etoposide, teniposide), DNA damaging agents (actinomycin, amsacrine, anthracyclines, bleomycin, busulfan, camptothecin, carboplatin, chlorambucil, cisplatin, cyclophosphamide, cytoxan, dactinomycin, daunorubicin, doxorubicin, epirubicin, hex amethylnelamineoxaliplatin, iphosphamide, melphalan, merchlorhtamine, mitomycin, mitoxantrone, nitrosourea, plicamycin, procarbazine, taxol, taxotere, teniposide, triethylenethiophosphoramidate and etoposide (VP16)); antibiotics such as dactinomycin (actinomycin D), daunorubicin,

doxorubicin (adriamycin), idarubicin, anthracyclines, mitoxantrone, bleomycins, plicamycin (mithramycin) and mitomycin; enzymes (L-asparaginase which systemically metabolizes L-asparagine and deprives cells which do not have the capacity to synthesize their own asparagine); antiplatelet agents; antiproliferative/antimitotic alkylating agents such as nitrogen mustards (mechlorethamine, cyclophosphamide and analogs, melphalan, chlorambucil), ethylenimines and methylmelamines (hexamethylmelamine and thiotepa), alkyl sulfonates-busulfan, nitrosoureas (carmustine (BCNU) and analogs, streptozocin), trazenes-dacarbazine (DTIC); antiproliferative/antimitotic antimetabolites such as folic acid analogs (methotrexate); platinum coordination complexes (cisplatin, carboplatin), procarbazine, hydroxyurea, mitotane, aminoglutethimide; hormones, hormone analogs (estrogen, tamoxifen, goserelin, bicalutamide, nilutamide) and aromatase inhibitors (letrozole, anastrozole); anticoagulants (heparin, synthetic heparin salts and other inhibitors of thrombin); fibrinolytic agents (such as tissue plasminogen activator, streptokinase and urokinase), aspirin, dipyridamole, ticlopidine, clopidogrel, abciximab; antimigratory agents; antisecretory agents (breveldin); immunosuppressives (cyclosporine, tacrolimus (FK-506), sirolimus (rapamycin), azathioprine, mycophenolate mofetil); anti-angiogenic compounds (e.g., TNP-470, genistein, bevacizumab) and growth factor inhibitors (e.g., fibroblast growth factor (FGF) inhibitors); angiotensin receptor blocker; nitric oxide donors; anti-sense oligonucleotides; antibodies (trastuzumab); cell cycle inhibitors and differentiation inducers (tretinoin); mTOR inhibitors, topoisomerase inhibitors (doxorubicin (adriamycin), amsacrine, camptothecin, daunorubicin, dactinomycin, eniposide, epirubicin, etoposide, idarubicin and mitoxantrone, topotecan, irinotecan), corticosteroids (cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisone, and prednisolone); growth factor signal transduction kinase inhibitors; mitochondrial dysfunction inducers and caspase activators; and chromatin disruptors.

Pharmacokinetics

[00179] In some embodiments patients may be evaluated for pharmacokinetics information. Pharmacokinetic data can provide insight regarding the fate of a given drug (e.g., a therapeutic agent) from administration to elimination from the human body.

[00180] Pharmacokinetic data can be obtained by known techniques in the art. Due to the inherent variation in pharmacokinetic and pharmacodynamic parameters of drug metabolism in human subjects, appropriate pharmacokinetic and pharmacodynamic profile components describing a particular composition can vary. Typically, pharmacokinetic and

pharmacodynamic profiles are based on the determination of the mean parameters of a group of subjects. The group of subjects includes any reasonable number of subjects suitable for determining a representative mean, for example, 5 subjects, 10 subjects, 16 subjects, 20 subjects, 25 subjects, 30 subjects, 35 subjects, or more. The mean is determined by calculating the average of all subject's measurements for each parameter measured.

[00181] In some embodiments, a patient population includes one or more subjects ("a population of subjects") suffering from metastatic disease.

[00182] In some embodiments, a patient population includes one or more subjects that is suffering from or susceptible to cancer. In some embodiments, a patient population includes one or more subjects (e.g., comprises or consists of subjects) suffering from cancer. For example, in some embodiments, a patient population suffering from cancer may have previously been treated with a prior therapy, for example, radiation and/or chemotherapy.

[00183] In some embodiments, the pharmacokinetic parameter(s) can be any parameters suitable for describing the present composition.

General Protocol for Dosing

[00184] As described herein, provided methods comprise administering a therapy that inhibits PARP and a therapy that regulates activity in the tumor microenvironment (e.g., T cell activity and/or infiltration of T cells into the tumor environment) in combination to a patient, a subject, or a population of subjects according to a regimen that achieves a therapeutic effect.

[00185] In some embodiments, administration "in combination" includes administration of one or more doses of an agent that inhibits PARP (e.g., niraparib) before, during, or after administration of one or more doses of an agent that enhances activity in the tumor microenvironment. In some embodiments, an agent that inhibits PARP (e.g., niraparib) and an agent that regulates activity in the tumor microenvironment are administered in overlapping regimens. In some embodiments, an agent that inhibits PARP (e.g., niraparib) is administered simultaneously or sequentially to an agent that enhances activity in the tumor microenvironment.

[00186] The number of times a composition is administered to an individual in need thereof depends on the discretion of a medical professional, the disorder, the severity of the disorder, and the individual's response to the formulation. In some embodiments, a composition disclosed herein is administered once to an individual in need thereof with a mild acute condition. In some embodiments, a composition disclosed herein is administered

more than once to an individual in need thereof with a moderate or severe acute condition. In the case wherein the patient's condition does not improve, upon the doctor's discretion the administration of a combination drug product described herein may be administered chronically, that is, for an extended period of time, including throughout the duration of the patient's life in order to ameliorate or otherwise control or limit the symptoms of the patient's disease or condition.

Indications Suitable for Treatment

[00187] Any subject having cancer, including breast cancer, ovarian cancer, cervical cancer, epithelial ovarian cancer, fallopian tube cancer, primary peritoneal cancer, endometrial cancer, prostate cancer, testicular cancer, pancreatic cancer, esophageal cancer, head and neck cancer, gastric cancer, bladder cancer, lung cancer (e.g., adenocarcinoma, NSCLC and SCLC), bone cancer (e.g., osteosarcoma), colon cancer, rectal cancer, thyroid cancer, brain and central nervous system cancers, glioblastoma, neuroblastoma, neuroendocrine cancer, rhabdoid cancer, keratoacanthoma, epidermoid carcinoma, seminoma, melanoma, sarcoma (e.g., liposarcoma), bladder cancer, liver cancer (e.g., hepatocellular carcinoma), kidney cancer (e.g., renal cell carcinoma), myeloid disorders (e.g., AML, CML, myelodysplastic syndrome and promyelocytic leukemia), and lymphoid disorders (e.g., leukemia, multiple myeloma, mantle cell lymphoma, ALL, CLL, B-cell lymphoma, T-cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma) may be treated with compounds and methods described herein.

[00188] In some embodiments, the methods of the invention treat subjects with a pediatric cancer. Exemplary pediatric cancers include, but are not limited to adrenocortical carcinoma, astrocytoma, atypical teratoid rhabdoid tumor, brain tumors, chondroblastoma, choroid plexus tumor, craniopharyngioma, desmoid tumor, dysembryoplastic neuroepithelial tumor (DNT), ependymoma, fibrosarcoma, germ cell tumor of the brain, glioblastoma multiforme, diffuse pontine glioma, low grade glioma, gliomatosis cerebri, hepatoblastoma, histiocytosis, kidney tumor, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), chronic myelogenous leukemia (CML), liposarcoma, liver cancer, Burkitt lymphoma, Hodgkin lymphoma, non-Hodgkin lymphoma, malignant fibrous histiocytoma, melanoma, myelodysplastic syndrome, nephroblastoma, neuroblastoma, neurofibrosarcoma, osteosarcoma, pilocytic astrocytoma, retinoblastoma, rhabdoid tumor of the kidney, rhabdomyosarcoma, Ewing sarcoma, soft tissue sarcoma, synovial sarcoma, spinal cord tumor and Wilm's tumor.

EXAMPLES

[00189] The following examples are provided to illustrate, but not limit the claimed invention.

Example 1 – Effects of an agent that inhibits PARP on the tumor microenvironment

[00190] This example describes the effects of treatment with an agent that inhibits PARP signaling on tumor microenvironment in a mouse model of colorectal cancer. 50 mg/kg of niraparib or vehicle (0.5% methylcellulose) was orally administered once daily for 21 days to C57BL/6 mice inoculated with primary Murine skin cancer model mSK6005 fragment developed from Apc^{Min}/J heterozygous background. Formalin-fixed, paraffin-embedded (FFPE) blocks were prepared from tumor samples collected from the mice. Immunohistochemistry (IHC) staining was performed on FFPE samples to assess expression of various markers for immune response in the tumor samples. Five fields in each staining, without necrosis, were randomly selected and imaged at 20x magnification.

[00191] All the images were analyzed with Image J software. For CD4, CD8, and FoxP3 IHC, positive cells were counted and the average of 5 fields positive cell numbers was taken as the score value of each case. For Iba1 IHC, the percentage of Iba1 positive expression area or mean gray was measured and taken as the score value of each case. The percentages of tumor cells at different intensity levels were evaluated according to the calculation below.

$$\text{Total Score} = (\% \text{ at } 0) \times 0 + (\% \text{ at } 1) \times 1 + (\% \text{ at } 2) \times 2 + (\% \text{ at } 3) \times 3$$

[00192] Increased expression of CD4, CD8, FoxP3, and Iba1 were observed in tumor samples obtained from mice treated with an agent that inhibits PARP signaling (e.g., niraparib) as compared to vehicle treated mice. These results can indicate an increased presence of CD4⁺ (see Figures 1A and 1B), CD8⁺ (see Figures 2A and 2B), Treg (see Figures 3A and 3B), and macrophage (see Figures 4A and 4B) cells upon administration of an agent that inhibits PARP signaling (e.g., niraparib) in an immuno-competent syngeneic mouse model. Similarly, increased percentage of CD8⁺ cells were observed in MDA-MB-436 huNOG-EXL mice treated with 35 mg/kg niraparib at QDx5/week (see Figure 5A), which resulted in a reduction of tumor growth (see Figure 5B). Thus, administration of an exemplary inhibitor of PARP signaling can enhance targeting of immune cells in the tumor microenvironment.

[00193] **Example 2 – Administration of an agent that regulates activity in the tumor microenvironment enhances the anti-tumor activity of an agent that inhibits PARP**

[00194] This example describes the effects of treatment with an agent that inhibits PARP together with an agent that regulates activity in the tumor microenvironment in a mouse model of breast cancer. HuNOG-EXL mice inoculated with MDA-MB436 breast carcinoma cell line were orally administered: i) 80 mg/kg of niraparib tosylate once daily; ii) 200 mg/kg BLZ945 (an anti-CSF-1R that binds to CSF1R and inhibits CSF1R-mediated signal transduction pathways in tumor associated macrophages) on a 5 day on and 2 day off schedule; or iii) 80 mg/kg of niraparib tosylate and 200 mg/kg BLZ945 with the same dosing schedule as the single agent groups. Control mice were administered an isotype control antibody twice weekly via intraperitoneal injection.

[00195] Tumor volumes were recorded twice in a week with a gap of 2-3 days in between two measurements until tumor volume reached 600-1000 mm³ in the control mice. The results are shown in Figure 6A, and tumor volume as measured on day 26 of the study is shown in Figure 6B. Consistent with the data described in Example 1, these results indicate that BLZ945, an agent that regulates activity in the tumor microenvironment, enhances anti-tumor activity of niraparib, an agent that inhibits PARP.

Embodiments:

1. A method of treating a subject with a disease or condition comprising administering to the subject
 - (a) a first agent that inhibits poly [ADP-ribose] polymerase (PARP); and
 - (b) a second agent, wherein the second agent comprises a regulatory T cell (Treg) inhibitory agent, a macrophage inhibitory agent, an antigen specific immune response enhancer agent, or a combination thereof.
2. A method of enhancing an immune response or increasing the activity of an immune cell in a subject with a disease or condition comprising administering to the subject
 - (a) a first agent that inhibits poly [ADP-ribose] polymerase (PARP); and
 - (b) a second agent, wherein the second agent comprises a regulatory T cell (Treg) inhibitory agent, a macrophage inhibitory agent, an antigen specific immune response enhancer agent, or a combination thereof.
3. A method of inducing an immune response in a subject with a disease or condition comprising administering to the subject
 - (a) a first agent that inhibits poly [ADP-ribose] polymerase (PARP); and

- (b) a second agent, wherein the second agent comprises a regulatory T cell (Treg) inhibitory agent, a macrophage inhibitory agent, an antigen specific immune response enhancer agent, or a combination thereof.
4. The method of any one of embodiments 1-3, wherein the first agent inhibits PARP 1 and/or 2.
 5. The method of any one of embodiments 1-4, wherein the first agent is a small organic or inorganic molecule; a saccharine; an oligosaccharide; a polysaccharide; a carbohydrate; a peptide; a protein; a peptide analog; a peptide derivative; a lipid; an antibody; an antibody fragment; a peptidomimetic; a nucleic acid; a nucleic acid analog; a nucleic acid derivative; an extract made from biological materials; a naturally occurring or synthetic composition; a metal; a toxin; or any combination thereof.
 6. The method of any one of embodiments 1-5, wherein the first agent is a small molecule.
 7. The method of any one of embodiments 1-6, wherein the first agent is selected from the group consisting of: ABT-767, AZD 2461, BGB-290, BGP 15, CEP 9722, E7016, E7449, fluzoparib, INO1001, JPI 289, MP 124, niraparib, olaparib, ONO2231, rucaparib, SC 101914, talazoparib, veliparib, WW 46, and salts or derivatives thereof.
 8. The method of any one of embodiments 1-7, wherein the first agent is selected from the group consisting of: niraparib, olaparib, rucaparib, talazoparib, and veliparib, or salts or derivatives thereof.
 9. The method of any one of embodiments 1-8, wherein the first agent comprises niraparib or a pharmaceutically acceptable salt or derivative thereof.
 10. The method of any one of embodiments 1-9, wherein the Treg inhibitory agent inhibits or decreases the activity, function, or migration of a Treg cell.
 11. The method of any one of embodiments 1-10, wherein the Treg inhibitory agent decreases a population of Treg cells in the subject.
 12. The method of any one of embodiments 1-11, wherein the Treg inhibitory agent substantially ablates or eliminates a population of Treg cells in the subject.
 13. The method of any one of embodiments 1-12, wherein the macrophage inhibitory agent inhibits or decreases the activity, function, or migration of a macrophage.
 14. The method of any one of embodiments 1-13, wherein the macrophage inhibitory agent decreases a population of macrophage cells in the subject.

15. The method of any one of embodiments 1-14, wherein the macrophage inhibitory agent substantially ablates or eliminates a population of macrophage cells in the subject.
16. The method of any one of embodiments 1-15, wherein the Treg cell is an infiltrating T cell.
17. The method of any one of embodiments 1-16, wherein the macrophage comprises a tumor-associated macrophage (TAM).
18. The method of any one of embodiments 1-17, wherein the second agent enhances an antigen specific CD4⁺ T cell activity.
19. The method of any one of embodiments 1-18, wherein the second agent enhances an antigen specific CD8⁺ T cell activity.
20. The method of any one of embodiments 1-19, wherein the second agent is selected from the group consisting of a small organic or inorganic molecule; a saccharine; an oligosaccharide; a polysaccharide; a carbohydrate; a peptide; a protein; a peptide analog; a peptide derivative; a lipid; an antibody; an antibody fragment; a peptidomimetic; a nucleic acid; a nucleic acid analog; a nucleic acid derivative; an extract made from biological materials; a naturally occurring or synthetic composition; a metal; a toxin; and any combination thereof.
21. The method of any one of embodiments 1-20, wherein the administering comprises administering the first and second agent sequentially.
22. The method of any one of embodiments 1-20, wherein the administering comprises administering the first and second agent simultaneously.
23. The method of any one of embodiments 1-20, wherein the administering comprises administering the first agent before administering the second agent.
24. The method of any one of embodiments 1-23, wherein the subject is a mammalian subject.
25. The method of any one of embodiments 1-24, wherein the subject is a human.
26. The method of any one of embodiments 1-25, wherein the second agent is a regulatory T cell (Treg) inhibitory agent selected from the group consisting of a Treg ablating agent, a Treg migration inhibitor agent, a Treg function inhibitor agent, and combinations thereof.
27. The method of embodiment 26, wherein the Treg ablating agent is selected from the group consisting of cyclophosphamide, paclitaxel, imatinib, sunitinib, sorafenib, dasatinib, temozolomide, daclizumab, denileukin diftitox, and combinations thereof.

28. The method of embodiment 26, wherein the Treg migration inhibitor agent is selected from the group consisting of AMD3100, mogamulizumab, casuarinin, fucoidan, and combinations thereof.

29. The method of embodiment 26, wherein the Treg function inhibitor agent is selected from the group consisting of an anti-CTLA4 agent (e.g., ipilimumab, tremelimumab), an anti-OX40 agent, an anti-GITR agent, an adenosine receptor antagonist (e.g., caffeine, theophylline, theobromine, and 8-phenylxanthines), P60, and combinations thereof.

30. The method of any one of embodiments 1-29, wherein the second agent is a macrophage inhibitory agent selected from the group consisting of a macrophage recruitment inhibitory agent, an M2 macrophage antisurvival agent, an M1 macrophage enhancing agent, an M2 to M1 polarizing agent, a macrophage activity inhibitor agent and combinations thereof.

31. The method of embodiment 30, wherein the macrophage recruitment inhibitory agent is selected from the group consisting of an anti-CCL2/CCR2 agent, an anti-IL6 agent, an anti-M-CSFR agent, and combinations thereof (e.g., an anti-M-CSFR agent).

32. The method of embodiment 31, wherein the macrophage recruitment inhibitory agent is selected from the group consisting of trabectedin, RS102895, PF-04136309, CNTO888, MLN1202, siltuximab, JNJ-28312141, GW2580, IMC-CS4 (LY3022855), emactuzumab, AMG820, pexidartinib, linifanib, OSI-930, CEP-32496, PLX7846, BLZ945, ARRY-382, JNJ-40346527, MCS110, PLX3397, PLX6134, PD-0360324, FPA008, and combinations thereof (e.g., BLZ945).

33. The method of embodiment 30, wherein the M2 macrophage antisurvival agent is selected from the group consisting of an MMP inhibitor, clodronate, zoledronic acid, dichloromethylene bisphosphonate, trabectedin, dasatinib, retinoic acid, attenuated bacteria (e.g., *Shigella flexneri*, *Salmonella typhimurium*, *Listeria monocytogens*, *Chlamydia psittaci*, *Legionella pneumophila*), and combinations thereof.

34. The method of embodiment 30, wherein the M1 macrophage enhancing agent or the M2 to M1 polarizing agent is selected from the group consisting of an anti-CD40 agent, an anti-IL-10R agent, a CD47 antagonist (e.g., Hu5F9-G4, CC-90002, and CD47-Fc fusion protein TTI-621), PolyI:C, LPS, monophosphoryl A, imiquimod, R-848, CpG-ODN, IFN- α , IFN- β , IFN- γ , GM-CSF, IL-12, IL-2, IL-15, T α 1, ibrutinib, EF-022 and combinations thereof.

35. The method of embodiment 30, wherein the macrophage activity inhibitory agent is selected from the group consisting of a STAT3 inhibitor, a STAT6 inhibitor, or an anti-tumor drug agent.

36. The method of embodiment 35, wherein the macrophage activity inhibitory agent is selected from the group consisting of WP1066, sunitinib, sorafenib, STA-21, IS3 295, S3I-M2001, AS1517499, leflunomide, TMC-264, histidine-rich glycoprotein (HRG), copper chelate (CuNG), 5,6-dimethylxanthenone-4-acetic acid (MDXAA), vadimezan (ASA404), cisplatin, silibinin, proton pump inhibitor pantoprazole (PPZ), CNI-1493 and combinations thereof.

37. The method of embodiment 30, wherein the macrophage inhibitor agent is an anti-IL-1 α agent (e.g., xilonix).

38. The method of any one of embodiments 1-37, wherein the second agent is an antigen specific immune response enhancer agent selected from the group consisting of an anti-PD-1 agent, an anti-PD-L1 agent, a GITR (glucocorticoid-induced TNFR-related protein) stimulating agent, an anti-CTLA4 agent, an anti-TIM-3 agent, an anti-LAG-3 agent, an anti-IDO agent, an agent that enhances tumor antigen presentation (e.g., personalized cancer vaccine, autologous antigen presenting cell, autologous dendritic cells, artificial antigen presenting cell), a chemokine signaling agent, an anti-VEGF agent, a cytokine signal stimulating agent, and combinations thereof.

39. The method of embodiment 38, wherein the anti-PD-1 agent is selected from the group consisting of pembrolizumab, nivolumab, PDR001, REGN2810 (SAR-439684), BGB-A317, BI 754091, IBI308, INCSHR-1210, JNJ-63723283, JS-001, MEDI0680 (AMP-514), MGA-012, PF-06801591, REGN-2810, TSR-042, atezolizumab, avelumab, CX-072, durvalumab, FAZ053, LY3300054, PD-L1 millamolecule, PDR-001, camrelizumab (HR-301210), BCD-100, AGEN-2034, CS1001, Sym-021, LZM009, KN-035, AB122, genolimzumab (CBT-501), AK 104, GLS-010, BGB-A333, SHR-1316, CK-301, and combinations thereof.

40. The method of embodiment 38 or 39, wherein the anti-PD-L1 agent is selected from the group consisting of atezolizumab, durvalumab, avelumab, LY3300054, BGB-A333, SHR-1316, CK-301, and combinations thereof.

41. The method of any one of embodiments 38-40, wherein the GITR stimulating agent is selected from the group consisting of DTA-1, mGITRL, pGITRL, and combinations thereof.

42. The method of any one of embodiments 38-41, wherein the anti-CTLA4 agent is selected from the group consisting of ipilimumab, tremelimumab, and combinations thereof.
43. The method of any one of embodiments 38-42, wherein the chemokine signaling agent is selected from the group consisting of CXCL16, a CXCR6 chemokine receptor (CD186) agonist, and combinations thereof.
44. The method of any one of embodiments 38-43, wherein the anti-VEGF agent is selected from the group consisting of bevacizumab, pazopanib, sunitinib, sorafenib, axitinib, ponatinib, regorafenib, cabozantinib, vandetanib, ramucirumab, lenvatinib, ziv-aflibercept, and combinations thereof.
45. The method of any one of embodiments 38-44, wherein the cytokine signal stimulating agent is an interleukin or an interferon.
46. The method of embodiment 45, wherein the interleukin is selected from the group consisting of IL-2, IL-1, IL-7, IL-15, IL-12, IL-18 and combinations thereof.
47. The method of embodiment 45, wherein the interferon is IFN alpha.
48. The method of any one of embodiments 1-47, wherein the second agent is an antigen specific immune response enhancer agent selected from the group consisting of a flavonoid (e.g., flavonoid glycoside), lidocaine, lamotrigine, sulfamethoxazole, phenytoin, carbamazepine, sulfamethoxazole, phenytoin, allopurinol, paracetamol, mepivacaine, p-phenylenediamine, ciprofloxacin and moxifloxacin.
49. The method of any one of embodiments 1-48, wherein the disease or condition is cancer.
50. The method of embodiment 49, wherein the cancer is selected from the group consisting of endometrial cancer, breast cancer, ovarian cancer, cervical cancer, fallopian tube cancer, primary peritoneal cancer, colon cancer, squamous cell carcinoma of the anogenital region, melanoma, renal cell carcinoma, lung cancer, non-small cell lung cancer, squamous cell carcinoma of the lung, stomach cancer, bladder cancer, gall bladder cancer, liver cancer, thyroid cancer, laryngeal cancer, salivary gland cancer, esophageal cancer, squamous cell carcinoma of the head and neck, prostate cancer, pancreatic cancer, mesothelioma, sarcoma, hematological cancer, and combinations thereof.
51. The method of any one of embodiments 1-50, wherein the administering comprises administering a composition comprising a capsule comprising the first agent.
52. The method of embodiment 51, wherein the capsule comprises a formulation comprising the first agent and one or more pharmaceutically acceptable excipients.

53. The method of embodiment 52, wherein the one or more pharmaceutically acceptable excipients comprises lactose monohydrate, magnesium stearate, or a combination thereof.

54. The method of any one of embodiments 1-53, wherein a therapeutically effective amount of the first or second agent is administered.

55. The method of any one of embodiments 1-54, wherein the method further comprises administering a third agent to the subject or performing a therapy on the subject selected from the group consisting of surgery, radiotherapy, and combinations thereof.

56. The method of embodiment 55, wherein the third agent comprises an antigen specific immune response enhancer agent, an anti-angiogenic agent, a chemotherapeutic agent, or combinations thereof.

57. The method of embodiment 56, wherein the antigen specific immune response enhancer agent comprises an anti-PD-1 agent, an anti-PD-L1 agent, an anti-CTLA4 agent, an anti-TIM-3 agent, or an anti-LAG-3 agent.

58. The method of embodiment 56, wherein the anti-angiogenic agent is selected from the group consisting of TNP-470, platelet factor 4, thrombospondin-1, tissue inhibitors of metalloproteases (TIMP1 and TIMP2), prolactin, angiostatin, endostatin, bFGF soluble receptor, transforming growth factor beta, interferon alpha, soluble KDR and FLT-1 receptors, placental proliferin-related protein, and combinations thereof.

59. The method of embodiment 56, wherein the chemotherapeutic agent is selected from the group consisting of aminoglutethimide, amsacrine, anastrozole, asparaginase, bcg, bicalutamide, bleomycin, buserelin, busulfan, campothecin, capecitabine, carboplatin, carmustine, chlorambucil, cisplatin, cladribine, clodronate, colchicine, cyclophosphamide, cyproterone, cytarabine, dacarbazine, dactinomycin, daunorubicin, dienestrol, diethylstilbestrol, docetaxel, doxorubicin, epirubicin, estradiol, estramustine, etoposide, exemestane, filgrastim, fludarabine, fludrocortisone, fluorouracil, fluoxymesterone, flutamide, gemcitabine, genistein, goserelin, hydroxyurea, idarubicin, ifosfamide, imatinib, interferon, irinotecan, ironotecan, letrozole, leucovorin, leuprolide, levamisole, lomustine, mechlorethamine, medroxyprogesterone, megestrol, melphalan, mercaptopurine, mesna, methotrexate, mitomycin, mitotane, mitoxantrone, nilutamide, nocodazole, octreotide, oxaliplatin, paclitaxel, pamidronate, pentostatin, plicamycin, porfimer, procarbazine, raltitrexed, rituximab, streptozocin, suramin, tamoxifen, temozolomide, teniposide, testosterone, thioguanine, thiotepa, titanocene dichloride, topotecan, trastuzumab, tretinoin, vinblastine, vincristine, vindesine, vinorelbine, and combinations thereof.

60. A pharmaceutical composition comprising
- (a) a first agent that inhibits poly [ADP-ribose] polymerase (PARP); and
 - (b) a second agent, wherein the second agent comprises a regulatory T cell (Treg) inhibitory agent, a macrophage inhibitory agent, an antigen specific immune response enhancer agent, or a combination thereof.
61. The pharmaceutical composition of embodiment 61, wherein the first agent inhibits PARP 1 and/or 2.
62. The pharmaceutical composition of embodiment 60 or 61, wherein the first agent is a small organic or inorganic molecule; a saccharine; an oligosaccharide; a polysaccharide; a carbohydrate; a peptide; a protein; a peptide analog; a peptide derivative; a lipid; an antibody; an antibody fragment; a peptidomimetic; a nucleic acid; a nucleic acid analog; a nucleic acid derivative; an extract made from biological materials; a naturally occurring or synthetic composition; a metal; a toxin; or any combination thereof.
63. The pharmaceutical composition of embodiment any one of embodiments 60-62, wherein the first agent is a small molecule.
64. The pharmaceutical composition of any one of embodiments 60-63, wherein the first agent is selected from the group consisting of: ABT-767, AZD 2461, BGB-290, BGP 15, CEP 9722, E7016, E7449, fluzoparib, INO1001, JPI 289, MP 124, niraparib, olaparib, ONO2231, rucaparib, SC 101914, talazoparib, veliparib, WW 46, and salts or derivatives thereof.
65. The pharmaceutical composition of any one of embodiments 60-64, wherein the first agent comprises niraparib or a pharmaceutically acceptable salt or derivative thereof.
66. The pharmaceutical composition of any one of embodiments 60-65, wherein the Treg inhibitory agent inhibits or decreases the activity, function, or migration of a Treg cell.
67. The pharmaceutical composition of any one of embodiments 60-66, wherein the Treg inhibitory agent decreases a population of Treg cells in the subject.
68. The pharmaceutical composition of any one of embodiments 60-67, wherein the Treg inhibitory agent substantially ablates or eliminates a population of Treg cells in the subject.
69. The pharmaceutical composition of any one of embodiments 60-68, wherein the macrophage inhibitory agent inhibits or decreases the activity, function, or migration of a macrophage.
70. The pharmaceutical composition of any one of embodiments 60-69, wherein the macrophage inhibitory agent decreases a population of macrophage cells in the subject.

71. The pharmaceutical composition of any one of embodiments 60-70, wherein the macrophage inhibitory agent substantially ablates or eliminates a population of macrophage cells in the subject.
72. The pharmaceutical composition of any one of embodiments 60-71, wherein the Treg cell is an infiltrating T cell.
73. The pharmaceutical composition of any one of embodiments 60-72, wherein the macrophage comprises a tumor-associated macrophage (TAM).
74. The pharmaceutical composition of any one of embodiments 60-73, wherein the second agent enhances an antigen specific CD4⁺ T cell activity.
75. The pharmaceutical composition of any one of embodiments 60-74, wherein the second agent enhances an antigen specific CD8⁺ T cell activity.
76. The pharmaceutical composition of any one of embodiments 60-75, wherein the second agent is selected from the group consisting of a small organic or inorganic molecule; a saccharine; an oligosaccharide; a polysaccharide; a carbohydrate; a peptide; a protein; a peptide analog; a peptide derivative; a lipid; an antibody; an antibody fragment; a peptidomimetic; a nucleic acid; a nucleic acid analog; a nucleic acid derivative; an extract made from biological materials; a naturally occurring or synthetic composition; a metal; a toxin; and any combination thereof.
77. The pharmaceutical composition of any one of embodiments 60-76, wherein the second agent is a regulatory T cell (Treg) inhibitory agent selected from the group consisting of a Treg ablating agent, a Treg migration inhibitor agent, a Treg function inhibitor agent, and combinations thereof.
78. The pharmaceutical composition of embodiment 79, wherein the Treg ablating agent is selected from the group consisting of cyclophosphamide, paclitaxel, imatinib, sunitinib, sorafenib, dasatinib, temozolomide, daclizumab, denileukin diftitox, and combinations thereof.
79. The pharmaceutical composition of embodiment 79, wherein the Treg migration inhibitor agent is selected from the group consisting of AMD3100, mogamulizumab, casuarinin, fucoidan, and combinations thereof.
80. The pharmaceutical composition of embodiment 79, wherein the Treg function inhibitor agent is selected from the group consisting of an anti-CTLA4 agent (e.g., ipilimumab, tremelimumab), an anti-OX40 agent, an anti-GITR agent, an adenosine receptor

antagonist (e.g., caffeine, theophylline, theobromine, and 8-phenylxanthines), P60, and combinations thereof.

81. The pharmaceutical composition of any one of embodiments 60-80, wherein the second agent is a macrophage inhibitory agent selected from the group consisting of a macrophage recruitment inhibitory agent, an M2 macrophage antisurvival agent, an M1 macrophage enhancing agent, an M2 to M1 polarizing agent, a macrophage activity inhibitor agent and combinations thereof.

82. The pharmaceutical composition of embodiment 81, wherein the macrophage recruitment inhibitory agent is selected from the group consisting of an anti-CCL2/CCR2 agent, an anti-IL6 agent, an anti-M-CSFR agent, and combinations thereof (e.g., an anti-M-CSFR agent).

83. The pharmaceutical composition of embodiment 82, wherein the macrophage recruitment inhibitory agent is selected from the group consisting of trabectedin, RS102895, PF-04136309, CNTO888, MLN1202, siltuximab, JNJ-28312141, GW2580, IMC-CS4 (LY3022855), emactuzumab, AMG820, pexidartinib, linifanib, OSI-930, CEP-32496, PLX7846, BLZ945, ARRY-382, JNJ-40346527, MCS110, PLX3397, PLX6134, PD-0360324, FPA008, and combinations thereof (e.g., BLZ945).

84. The pharmaceutical composition of embodiment 81, wherein the M2 macrophage antisurvival agent is selected from the group consisting of an MMP inhibitor, clodronate, zoledronic acid, dichloromethylene bisphosphonate, trabectedin, dasatinib, retinoic acid, attenuated bacteria (e.g., *Shigella flexneri*, *Salmonella typhimurium*, *Listeria monocytogens*, *Chlamydia psittaci*, *Legionella pneumophila*), and combinations thereof.

85. The pharmaceutical composition of embodiment 81, wherein the M1 macrophage enhancing agent or the M2 to M1 polarizing agent is selected from the group consisting of an anti-CD40 agent, an anti-IL-10R agent, a CD47 antagonist (e.g., Hu5F9-G4, CC-90002, and CD47-Fc fusion protein TTI-621), PolyI:C, LPS, monophosphoryl A, imiquimod, R-848, CpG-ODN, IFN- α , IFN- β , IFN- γ , GM-CSF, IL-12, IL-2, IL-15, T α 1, ibrutinib, EF-022 and combinations thereof.

86. The pharmaceutical composition of embodiment 81, wherein the macrophage activity inhibitory agent is selected from the group consisting of a STAT3 inhibitor, a STAT6 inhibitor, or an anti-tumor drug agent.

87. The pharmaceutical composition of embodiment 86, wherein the macrophage activity inhibitory agent is selected from the group consisting of WP1066, sunitinib, sorafenib, STA-

21, IS3 295, S3I-M2001, AS1517499, leflunomide, TMC-264, histidine-rich glycoprotein (HRG), copper chelate (CuNG), 5,6-dimethylxanthenone-4-acetic acid (MDXAA), vadimezan (ASA404), cisplatin, silibinin, proton pump inhibitor pantoprazole (PPZ), CNI-1493 and combinations thereof.

88. The pharmaceutical composition of embodiment 81, wherein the macrophage inhibitor agent is an anti-IL-1 α agent (e.g., xilonix).

89. The pharmaceutical composition of any one of embodiments 60-88, wherein the second agent is an antigen specific immune response enhancer agent selected from the group consisting of an anti-PD-1 agent, an anti-PD-L1 agent, a GITR (glucocorticoid-induced TNFR-related protein) stimulating agent, an anti-CTLA4 agent, an anti-TIM-3 agent, an anti-LAG-3 agent, an anti-IDO agent, an agent that enhances tumor antigen presentation (e.g., personalized cancer vaccine, autologous antigen presenting cell, autologous dendritic cells, artificial antigen presenting cell), a chemokine signaling agent, an anti-VEGF agent, a cytokine signal stimulating agent, and combinations thereof.

90. The pharmaceutical composition of embodiment 94, wherein the anti-PD-1 agent is selected from the group consisting of pembrolizumab, nivolumab, PDR001, REGN2810 (SAR-439684), BGB-A317, BI 754091, IBI308, INCSHR-1210, JNJ-63723283, JS-001, MEDI0680 (AMP-514), MGA-012, PF-06801591, REGN-2810, TSR-042, atezolizumab, avelumab, CX-072, durvalumab, FAZ053, LY3300054, PD-L1 millamolecule, PDR-001, camrelizumab (HR-301210), BCD-100, AGEN-2034, CS1001, Sym-021, LZM009, KN-035, AB122, genolimzumab (CBT-501), AK 104, GLS-010, BGB-A333, SHR-1316, CK-301, and combinations thereof.

91. The pharmaceutical composition of embodiment 94 or 95, wherein the anti-PD-L1 agent is selected from the group consisting of atezolizumab, durvalumab, avelumab, LY3300054, BGB-A333, SHR-1316, CK-301, and combinations thereof.

92. The pharmaceutical composition of any one of embodiments 89-91, wherein the GITR stimulating agent is selected from the group consisting of DTA-1, mGITRL, pGITRL, and combinations thereof.

93. The pharmaceutical composition of any one of embodiments 89-92, wherein the anti-CTLA4 agent is selected from the group consisting of ipilimumab, tremelimumab, and combinations thereof.

94. The pharmaceutical composition of any one of embodiments 89-93, wherein the chemokine signaling agent is selected from the group consisting of CXCL16, a CXCR6 chemokine receptor (CD186) agonist, and combinations thereof.

95. The pharmaceutical composition of any one of embodiments 89-94, wherein the anti-VEGF agent is selected from the group consisting of bevacizumab, pazopanib, sunitinib, sorafenib, axitinib, ponatinib, regorafenib, cabozantinib, vandetanib, ramucirumab, lenvatinib, ziv-aflibercept, and combinations thereof.

96. The pharmaceutical composition of any one of embodiments 89-95, wherein the cytokine signal stimulating agent is an interleukin or an interferon.

97. The pharmaceutical composition of embodiment 96, wherein the interleukin is selected from the group consisting of IL-2, IL-1, IL-7, IL-15, IL-12, IL-18 and combinations thereof.

98. The pharmaceutical composition of embodiment 96, wherein the interferon is IFN alpha.

99. The pharmaceutical composition of any one of embodiments 60-98, wherein the second agent is an antigen specific immune response enhancer agent selected from the group consisting of a flavonoid (e.g., flavonoid glycoside), lidocaine, lamotrigine, sulfamethoxazole, phenytoin, carbamazepine, sulfamethoxazole, phenytoin, allopurinol, paracetamol, mepivacaine, p-phenylenediamine, ciprofloxacin and moxifloxacin.

100. The pharmaceutical composition of embodiment 99, wherein the disease or condition is cancer.

101. The pharmaceutical composition of embodiment 99, wherein the cancer is selected from the group consisting of endometrial cancer, breast cancer, ovarian cancer, cervical cancer, fallopian tube cancer, primary peritoneal cancer, colon cancer, squamous cell carcinoma of the anogenital region, melanoma, renal cell carcinoma, lung cancer, non-small cell lung cancer, squamous cell carcinoma of the lung, stomach cancer, bladder cancer, gall bladder cancer, liver cancer, thyroid cancer, laryngeal cancer, salivary gland cancer, esophageal cancer, squamous cell carcinoma of the head and neck, prostate cancer, pancreatic cancer, mesothelioma, sarcoma, a hematological cancer, and combinations thereof.

102. The pharmaceutical composition of any one of embodiments 60-101, wherein the administering comprises administering a composition comprising a capsule comprising the first agent.

103. The pharmaceutical composition of embodiment 102, wherein the capsule comprises a formulation comprising the first agent and one or more pharmaceutically acceptable excipients.

104. The pharmaceutical composition of embodiment 103, wherein the one or more pharmaceutically acceptable excipients comprises lactose monohydrate, magnesium stearate, or a combination thereof.

105. The pharmaceutical composition of any one of embodiments 60-104, wherein a therapeutically effective amount of the first or second agent is administered.

106. The pharmaceutical composition of any one of embodiments 60-105, wherein the method further comprises administering a third agent to the subject.

107. The pharmaceutical composition of embodiment 106, wherein the third agent comprises an antigen specific immune response enhancer agent, an anti-angiogenic agent, a chemotherapeutic agent, or combinations thereof.

108. The pharmaceutical composition of embodiment 107, wherein the antigen specific immune response enhancer agent comprises an anti-PD-1 agent, an anti-PD-L1 agent, an anti-CTLA4 agent, an anti-TIM-3 agent, or an anti-LAG-3 agent.

109. The pharmaceutical composition of embodiment 107, wherein the anti-angiogenic agent is selected from the group consisting of TNP-470, platelet factor 4, thrombospondin-1, tissue inhibitors of metalloproteases (TIMP1 and TIMP2), prolactin, angiostatin, endostatin, bFGF soluble receptor, transforming growth factor beta, interferon alpha, soluble KDR and FLT-1 receptors, placental proliferin-related protein, and combinations thereof.

110. The pharmaceutical composition of embodiment 109, wherein the chemotherapeutic agent is selected from the group consisting of aminoglutethimide, amsacrine, anastrozole, asparaginase, bcr, bicalutamide, bleomycin, buserelin, busulfan, camptothecin, capecitabine, carboplatin, carmustine, chlorambucil, cisplatin, cladribine, clodronate, colchicine, cyclophosphamide, cyproterone, cytarabine, dacarbazine, dactinomycin, daunorubicin, dienestrol, diethylstilbestrol, docetaxel, doxorubicin, epirubicin, estradiol, estramustine, etoposide, exemestane, filgrastim, fludarabine, fludrocortisone, fluorouracil, fluoxymesterone, flutamide, gemcitabine, genistein, goserelin, hydroxyurea, idarubicin, ifosfamide, imatinib, interferon, irinotecan, irinotecan, letrozole, leucovorin, leuprolide, levamisole, lomustine, mechlorethamine, medroxyprogesterone, megestrol, melphalan, mercaptopurine, mesna, methotrexate, mitomycin, mitotane, mitoxantrone, nilutamide, nocodazole, octreotide, oxaliplatin, paclitaxel, pamidronate, pentostatin, plicamycin,

porfimer, procarbazine, raltitrexed, rituximab, streptozocin, suramin, tamoxifen, temozolomide, teniposide, testosterone, thioguanine, thiotepa, titanocene dichloride, topotecan, trastuzumab, tretinoin, vinblastine, vincristine, vindesine, vinorelbine, and combinations thereof.

CLAIMS

1. A method of treating a subject with a disease or condition comprising administering to the subject
 - (a) a first agent that inhibits poly [ADP-ribose] polymerase (PARP); and
 - (b) a second agent, wherein the second agent comprises a regulatory T cell (Treg) inhibitory agent, a macrophage inhibitory agent, an antigen specific immune response enhancer agent, or a combination thereof.
2. A method of enhancing an immune response or increasing the activity of an immune cell in a subject with a disease or condition comprising administering to the subject
 - (a) a first agent that inhibits poly [ADP-ribose] polymerase (PARP); and
 - (b) a second agent, wherein the second agent comprises a regulatory T cell (Treg) inhibitory agent, a macrophage inhibitory agent, an antigen specific immune response enhancer agent, or a combination thereof.
3. A method of inducing an immune response in a subject with a disease or condition comprising administering to the subject
 - (a) a first agent that inhibits poly [ADP-ribose] polymerase (PARP); and
 - (b) a second agent, wherein the second agent comprises a regulatory T cell (Treg) inhibitory agent, a macrophage inhibitory agent, an antigen specific immune response enhancer agent, or a combination thereof.
4. The method of any one of claims 1-3, wherein the first agent inhibits PARP 1 and/or 2.
5. The method of any one of claims 1-4, wherein the first agent is a small organic or inorganic molecule; a saccharine; an oligosaccharide; a polysaccharide; a carbohydrate; a peptide; a protein; a peptide analog; a peptide derivative; a lipid; an antibody; an antibody fragment; a peptidomimetic; a nucleic acid; a nucleic acid analog; a nucleic acid derivative; an extract made from biological materials; a naturally occurring or synthetic composition; a metal; a toxin; or any combination thereof.
6. The method of any one of claims 1-5, wherein the first agent is a small molecule.
7. The method of any one of claims 1-6, wherein the first agent is selected from the group consisting of: ABT-767, AZD 2461, BGB-290, BGP 15, CEP 8983, CEP 9722, DR 2313, E7016, E7449, fluzoparib (SHR 3162), IMP 4297, INO1001, JPI 289, JPI 547,

monoclonal antibody B3-LysPE40 conjugate, MP 124, niraparib (Zejula) (MK-4827), NU 1025, NU 1064, NU 1076, NU1085, olaparib (AZD2281), ONO2231, PD 128763, R 503, R554, rucaparib (RUBRACA) (AG-014699, PF-01367338), SBP 101, SC 101914, Simmiparib, talazoparib (BMN-673), veliparib (ABT-888), WW 46, 2-(4-(Trifluoromethyl)phenyl)-7,8-dihydro-5H-thiopyrano[4,3-d]pyrimidin-4-ol, and salts or derivatives thereof.

8. The method of any one of claims 1-7, wherein the first agent is selected from the group consisting of: niraparib, olaparib, rucaparib, talazoparib, and veliparib, or salts or derivatives thereof.
9. The method of any one of claims 1-8, wherein the first agent comprises niraparib or a pharmaceutically acceptable salt or derivative thereof.
10. The method of any one of claims 1-9, wherein the macrophage inhibitory agent inhibits or decreases the activity, function, or migration of a macrophage.
11. The method of any one of claims 1-10, wherein the macrophage inhibitory agent decreases a population of macrophage cells in the subject.
12. The method of any one of claims 1-11, wherein the macrophage inhibitory agent substantially ablates or eliminates a population of macrophage cells in the subject.
13. The method of any one of claims 1-12, wherein the macrophage comprises a tumor-associated macrophage (TAM).
14. The method of any one of claims 1-13, wherein the Treg inhibitory agent inhibits or decreases the activity, function, or migration of a Treg cell.
15. The method of any one of claims 1-14, wherein the Treg inhibitory agent decreases a population of Treg cells in the subject.
16. The method of any one of claims 1-15, wherein the Treg inhibitory agent substantially ablates or eliminates a population of Treg cells in the subject.
17. The method of any one of claims 1-16, wherein the Treg cell is an infiltrating T cell.

18. The method of any one of claims 1-17, wherein the second agent enhances an antigen specific CD4⁺ T cell activity.
19. The method of any one of claims 1-18, wherein the second agent enhances an antigen specific CD8⁺ T cell activity.
20. The method of any one of claims 1-19, wherein the second agent is selected from the group consisting of a small organic or inorganic molecule; a saccharine; an oligosaccharide; a polysaccharide; a carbohydrate; a peptide; a protein; a peptide analog; a peptide derivative; a lipid; an antibody; an antibody fragment; a peptidomimetic; a nucleic acid; a nucleic acid analog; a nucleic acid derivative; an extract made from biological materials; a naturally occurring or synthetic composition; a metal; a toxin; and any combination thereof.
21. The method of any one of claims 1-20, wherein the administering comprises administering the first and second agent sequentially.
22. The method of any one of claims 1-20, wherein the administering comprises administering the first and second agent simultaneously.
23. The method of any one of claims 1-20, wherein the administering comprises administering the first agent before administering the second agent second agent.
24. The method of any one of claims 1-23, wherein the subject is a mammalian subject.
25. The method of any one of claims 1-24, wherein the subject is a human.
26. The method of any one of claims 1-25, wherein the second agent is a macrophage inhibitory agent selected from the group consisting of a macrophage recruitment inhibitory agent, an M2 macrophage antisurvival agent, an M1 macrophage enhancing agent, an M2 to M1 polarizing agent, a macrophage activity inhibitor agent and combinations thereof.
27. The method of claim 26, wherein the macrophage recruitment inhibitory agent is selected from the group consisting of an anti-CCL2/CCR2 agent, an anti-IL6 agent, an anti-M-CSFR agent, and combinations thereof.

28. The method of claim 27, wherein the macrophage recruitment inhibitory agent is an anti-M-CSFR agent.
29. The method of claim 27, wherein the macrophage recruitment inhibitory agent is selected from the group consisting of trabectedin, RS102895, PF-04136309, CNTO888, MLN1202, siltuximab, JNJ-28312141, GW2580, IMC-CS4 (LY3022855), emactuzumab, AMG820, pexidartinib, linifanib, OSI-930, CEP-32496, PLX7846, BLZ945, ARRY-382, JNJ-40346527, MCS110, PLX3397, PLX6134, PD-0360324, FPA008, and combinations thereof.
30. The method of claim 28 or 29, wherein the macrophage recruitment inhibitory agent is BLZ945, PLX7846, GW2580, ARRY-382, JNJ-40346527, emactuzumab, pexidartinib, AMG820, IMC-CS4 (LY3022855), MCS110, PLX3397, PLX6134, PD-0360324, or FPA008.
31. The method of claim 30, wherein the macrophage recruitment inhibitory agent is BLZ945.
32. The method of claim 26, wherein the M2 macrophage antisurvival agent is selected from the group consisting of an MMP inhibitor, clodronate, zoledronic acid, dichloromethylene bisphosphonate, trabectedin, dasatinib, retinoic acid, attenuated bacteria (e.g., *Shigella flexneri*, *Salmonella typhimurium*, *Listeria monocytogens*, *Chlamydia psittaci*, *Legionella pneumophila*), and combinations thereof.
33. The method of claim 26, wherein the M1 macrophage enhancing agent or the M2 to M1 polarizing agent is selected from the group consisting of an anti-CD40 agent, an anti-IL-10R agent, a CD47 antagonist (e.g., Hu5F9-G4, CC-90002, and CD47-Fc fusion protein TTI-621), PolyI:C, LPS, monophosphoryl A, imiquimod, R-848, CpG-ODN, IFN- α , IFN- β , IFN- γ , GM-CSF, IL-12, IL-2, IL-15, T α 1, ibrutinib, EF-022 and combinations thereof.
34. The method of claim 26, wherein the macrophage activity inhibitory agent is selected from the group consisting of a STAT3 inhibitor, a STAT6 inhibitor, or an anti-tumor drug agent.

35. The method of claim 34, wherein the macrophage activity inhibitory agent is selected from the group consisting of WP1066, sunitinib, sorafenib, STA-21, IS3 295, S3I-M2001, AS1517499, leflunomide, TMC-264, histidine-rich glycoprotein (HRG), copper chelate (CuNG), 5,6-dimethylxanthenone-4-acetic acid (MDXAA), vadimezan (ASA404), cisplatin, silibinin, proton pump inhibitor pantoprazole (PPZ), CNI-1493 and combinations thereof.
36. The method of claim 35, wherein the macrophage inhibitor agent is an anti-IL-1 α agent (e.g., xilonix).
37. The method of any one of claims 1-36, wherein the second agent is a regulatory T cell (Treg) inhibitory agent selected from the group consisting of a Treg ablating agent, a Treg migration inhibitor agent, a Treg function inhibitor agent, and combinations thereof.
38. The method of claim 37, wherein the Treg ablating agent is selected from the group consisting of cyclophosphamide, paclitaxel, imatinib, sunitinib, sorafenib, dasatinib, temozolomide, daclizumab, denileukin diftitox, and combinations thereof.
39. The method of claim 37, wherein the Treg migration inhibitor agent is selected from the group consisting of AMD3100, mogamulizumab, casuarinin, fucoidan, and combinations thereof.
40. The method of claim 37, wherein the Treg function inhibitor agent is selected from the group consisting of an anti-CTLA4 agent (e.g., ipilimumab, tremelimumab), an anti-OX40 agent, an anti-GITR agent, an adenosine receptor antagonist (e.g., caffeine, theophylline, theobromine, and 8-phenylxanthines), P60, and combinations thereof.
41. The method of any one of claims 1-36, wherein the second agent is an antigen specific immune response enhancer agent selected from the group consisting of an anti-PD-1 agent, an anti-PD-L1 agent, a GITR (glucocorticoid-induced TNFR-related protein) stimulating agent, an anti-CTLA4 agent, an anti-TIM-3 agent, an anti-LAG-3 agent, an anti-IDO agent, an agent that enhances tumor antigen presentation (e.g., personalized cancer vaccine, autologous antigen presenting cell, autologous dendritic cells, artificial

antigen presenting cell), a chemokine signaling agent, an anti-VEGF agent, a cytokine signal stimulating agent, and combinations thereof.

42. The method of claim 41, wherein the anti-PD-1 agent is selected from the group consisting of pembrolizumab, nivolumab, PDR001, REGN2810 (SAR-439684), BGB-A317, BI 754091, IBI308, INCSHR-1210, JNJ-63723283, JS-001, MEDI0680 (AMP-514), MGA-012, PF-06801591, REGN-2810, TSR-042, atezolizumab, avelumab, CX-072, durvalumab, FAZ053, LY3300054, PD-L1 millamolecule, PDR-001, camrelizumab (HR-301210), BCD-100, AGEN-2034, CS1001, Sym-021, LZM009, KN-035, AB122, genolimzumab (CBT-501), AK 104, GLS-010, BGB-A333, SHR-1316, CK-301, and combinations thereof.
43. The method of claim 41 or 42, wherein the anti-PD-L1 agent is selected from the group consisting of atezolizumab, durvalumab, avelumab, LY3300054, BGB-A333, SHR-1316, CK-301, and combinations thereof.
44. The method of any one of claims 41-43, wherein the GITR stimulating agent is selected from the group consisting of DTA-1, mGITRL, pGITRL, and combinations thereof.
45. The method of any one of claims 41-44, wherein the anti-CTLA4 agent is selected from the group consisting of ipilimumab, tremelimumab, and combinations thereof.
46. The method of any one of claims 41-45, wherein the chemokine signaling agent is selected from the group consisting of CXCL16, a CXCR6 chemokine receptor (CD186) agonist, and combinations thereof.
47. The method of any one of claims 41-46, wherein the anti-VEGF agent is selected from the group consisting of bevacizumab, pazopanib, sunitinib, sorafenib, axitinib, ponatinib, regorafenib, cabozantinib, vandetanib, ramucirumab, lenvatinib, ziv-aflibercept, and combinations thereof.
48. The method of any one of claims 41-47, wherein the cytokine signal stimulating agent is an interleukin or an interferon.
49. The method of claim 48, wherein the interleukin is selected from the group consisting of IL-2, IL-1, IL-7, IL-15, IL-12, IL-18 and combinations thereof.

50. The method of claim 48, wherein the interferon is IFN alpha.
51. The method of any one of claims 1-50, wherein the second agent is an antigen specific immune response enhancer agent selected from the group consisting of a flavonoid (e.g., flavonoid glycoside), lidocaine, lamotrigine, sulfamethoxazole, phenytoin, carbamazepine, sulfamethoxazole, phenytoin, allopurinol, paracetamol, mepivacaine, p-phenylenediamine, ciprofloxacin and moxifloxacin.
52. The method of any one of claims 1-51, wherein the disease or condition is cancer.
53. The method of claim 52, wherein the cancer is selected from the group consisting of endometrial cancer, breast cancer, ovarian cancer, cervical cancer, fallopian tube cancer, primary peritoneal cancer, colon cancer, squamous cell carcinoma of the anogenital region, melanoma, renal cell carcinoma, lung cancer, non-small cell lung cancer, squamous cell carcinoma of the lung, stomach cancer, bladder cancer, gall bladder cancer, liver cancer, thyroid cancer, laryngeal cancer, salivary gland cancer, esophageal cancer, squamous cell carcinoma of the head and neck, prostate cancer, pancreatic cancer, mesothelioma, sarcoma, hematological cancer, and combinations thereof.
54. The method of any one of claims 1-53, wherein the administering comprises administering a composition comprising a capsule comprising the first agent.
55. The method of claim 54, wherein the capsule comprises a formulation comprising the first agent and one or more pharmaceutically acceptable excipients.
56. The method of claim 55, wherein the one or more pharmaceutically acceptable excipients comprises lactose monohydrate, magnesium stearate, or a combination thereof.
57. The method of any one of claims 1-56, wherein a therapeutically effective amount of the first or second agent is administered.
58. The method of any one of claims 1-57, wherein the method further comprises administering a third agent to the subject or performing a therapy on the subject selected from the group consisting of surgery, radiotherapy, and combinations thereof.

59. The method of claim 58, wherein the third agent comprises an antigen specific immune response enhancer agent, an anti-angiogenic agent, a chemotherapeutic agent, or combinations thereof.
60. The method of claim 59, wherein the antigen specific immune response enhancer agent comprises an anti-PD-1 agent, an anti-PD-L1 agent, an anti-CTLA4 agent, an anti-TIM-3 agent, or an anti-LAG-3 agent.
61. The method of claim 59, wherein the anti-angiogenic agent is selected from the group consisting of TNP-470, platelet factor 4, thrombospondin-1, tissue inhibitors of metalloproteases (TIMP1 and TIMP2), prolactin, angiostatin, endostatin, bFGF soluble receptor, transforming growth factor beta, interferon alpha, soluble KDR and FLT-1 receptors, placental proliferin-related protein, and combinations thereof.
62. The method of claim 59, wherein the chemotherapeutic agent is selected from the group consisting of aminoglutethimide, amsacrine, anastrozole, asparaginase, bcr, bicalutamide, bleomycin, buserelin, busulfan, camptothecin, capecitabine, carboplatin, carmustine, chlorambucil, cisplatin, cladribine, clodronate, colchicine, cyclophosphamide, cyproterone, cytarabine, dacarbazine, dactinomycin, daunorubicin, dienestrol, diethylstilbestrol, docetaxel, doxorubicin, epirubicin, estradiol, estramustine, etoposide, exemestane, filgrastim, fludarabine, fludrocortisone, fluorouracil, fluoxymesterone, flutamide, gemcitabine, genistein, goserelin, hydroxyurea, idarubicin, ifosfamide, imatinib, interferon, irinotecan, irinotecan, letrozole, leucovorin, leuprolide, levamisole, lomustine, mechlorethamine, medroxyprogesterone, megestrol, melphalan, mercaptopurine, mesna, methotrexate, mitomycin, mitotane, mitoxantrone, nilutamide, nocodazole, octreotide, oxaliplatin, paclitaxel, pamidronate, pentostatin, plicamycin, porfimer, procarbazine, raltitrexed, rituximab, streptozocin, suramin, tamoxifen, temozolomide, teniposide, testosterone, thioguanine, thiotepa, titanocene dichloride, topotecan, trastuzumab, tretinoin, vinblastine, vincristine, vindesine, vinorelbine, and combinations thereof.
63. A pharmaceutical composition comprising
 - (a) a first agent that inhibits poly [ADP-ribose] polymerase (PARP); and

- (b) a second agent, wherein the second agent comprises a regulatory T cell (Treg) inhibitory agent, a macrophage inhibitory agent, an antigen specific immune response enhancer agent, or a combination thereof.
64. The pharmaceutical composition of claim 63, wherein the first agent inhibits PARP 1 and/or 2.
65. The pharmaceutical composition of claim 63 or 64, wherein the first agent is a small organic or inorganic molecule; a saccharine; an oligosaccharide; a polysaccharide; a carbohydrate; a peptide; a protein; a peptide analog; a peptide derivative; a lipid; an antibody; an antibody fragment; a peptidomimetic; a nucleic acid; a nucleic acid analog; a nucleic acid derivative; an extract made from biological materials; a naturally occurring or synthetic composition; a metal; a toxin; or any combination thereof.
66. The pharmaceutical composition of claim any one of claims 63-65, wherein the first agent is a small molecule.
67. The pharmaceutical composition of any one of claims 63-66, wherein the first agent is selected from the group consisting of: ABT-767, AZD 2461, BGB-290, BGP 15, CEP 8983, CEP 9722, DR 2313, E7016, E7449, fluzoparib (SHR 3162), IMP 4297, INO1001, JPI 289, JPI 547, monoclonal antibody B3-LysPE40 conjugate, MP 124, niraparib (ZEJULA) (MK-4827), NU 1025, NU 1064, NU 1076, NU1085, olaparib (AZD2281), ONO2231, PD 128763, R 503, R554, rucaparib (RUBRACA) (AG-014699, PF-01367338), SBP 101, SC 101914, Simmiparib, talazoparib (BMN-673), veliparib (ABT-888), WW 46, 2-(4-(Trifluoromethyl)phenyl)-7,8-dihydro-5H-thiopyrano[4,3-d]pyrimidin-4-ol, and salts or derivatives thereof.
68. The pharmaceutical composition of any one of claims 63-67, wherein the first agent comprises niraparib or a pharmaceutically acceptable salt or derivative thereof.
69. The pharmaceutical composition of any one of claims 63-68, wherein the Treg inhibitory agent inhibits or decreases the activity, function, or migration of a Treg cell.
70. The pharmaceutical composition of any one of claims 63-69, wherein the Treg inhibitory agent decreases a population of Treg cells in the subject.

71. The pharmaceutical composition of any one of claims 63-70, wherein the Treg inhibitory agent substantially ablates or eliminates a population of Treg cells in the subject.
72. The pharmaceutical composition of any one of claims 63-71, wherein the macrophage inhibitory agent inhibits or decreases the activity, function, or migration of a macrophage.
73. The pharmaceutical composition of any one of claims 63-72, wherein the macrophage inhibitory agent decreases a population of macrophage cells in the subject.
74. The pharmaceutical composition of any one of claims 63-73, wherein the macrophage inhibitory agent substantially ablates or eliminates a population of macrophage cells in the subject.
75. The pharmaceutical composition of any one of claims 63-74, wherein the Treg cell is an infiltrating T cell.
76. The pharmaceutical composition of any one of claims 63-75, wherein the macrophage comprises a tumor-associated macrophage (TAM).
77. The pharmaceutical composition of any one of claims 63-76, wherein the second agent enhances an antigen specific CD4⁺ T cell activity.
78. The pharmaceutical composition of any one of claims 63-77, wherein the second agent enhances an antigen specific CD8⁺ T cell activity.
79. The pharmaceutical composition of any one of claims 63-78, wherein the second agent is selected from the group consisting of a small organic or inorganic molecule; a saccharine; an oligosaccharide; a polysaccharide; a carbohydrate; a peptide; a protein; a peptide analog; a peptide derivative; a lipid; an antibody; an antibody fragment; a peptidomimetic; a nucleic acid; a nucleic acid analog; a nucleic acid derivative; an extract made from biological materials; a naturally occurring or synthetic composition; a metal; a toxin; and any combination thereof.
80. The pharmaceutical composition of any one of claims 63-79, wherein the second agent is a macrophage inhibitory agent selected from the group consisting of a macrophage recruitment inhibitory agent, an M2 macrophage antisurvival agent, an M1 macrophage

enhancing agent, an M2 to M1 polarizing agent, a macrophage activity inhibitor agent and combinations thereof.

81. The pharmaceutical composition of claim 80, wherein the macrophage recruitment inhibitory agent is selected from the group consisting of an anti-CCL2/CCR2 agent, an anti-IL6 agent, an anti-M-CSFR agent, and combinations thereof.
82. The pharmaceutical composition of claim 81, wherein the macrophage recruitment inhibitory agent is an anti-M-CSFR agent.
83. The pharmaceutical composition of claim 81, wherein the macrophage recruitment inhibitory agent is selected from the group consisting of trabectedin, RS102895, PF-04136309, CNTO888, MLN1202, siltuximab, JNJ-28312141, GW2580, IMC-CS4 (LY3022855), emactuzumab, AMG820, pexidartinib, linifanib, OSI-930, CEP-32496, PLX7846, BLZ945, ARRY-382, JNJ-40346527, MCS110, PLX3397, PLX6134, PD-0360324, FPA008, and combinations thereof.
84. The pharmaceutical composition of claim 82 or 83, wherein the macrophage recruitment inhibitory agent is BLZ945, PLX7846, GW2580, ARRY-382, JNJ-40346527, emactuzumab, pexidartinib, AMG820, IMC-CS4 (LY3022855), MCS110, PLX3397, PLX6134, PD-0360324, or FPA008.
85. The pharmaceutical composition of claim 84, wherein the macrophage recruitment inhibitory agent is BLZ945.
86. The pharmaceutical composition of claim 80, wherein the M2 macrophage antisurvival agent is selected from the group consisting of an MMP inhibitor, clodronate, zoledronic acid, dichloromethylene bisphosphonate, trabectedin, dasatinib, retinoic acid, attenuated bacteria (e.g., *Shigella flexneri*, *Salmonella typhimurium*, *Listeria monocytogens*, *Chlamydia psittaci*, *Legionella pneumophila*), and combinations thereof.
87. The pharmaceutical composition of claim 80, wherein the M1 macrophage enhancing agent or the M2 to M1 polarizing agent is selected from the group consisting of an anti-CD40 agent, an anti-IL-10R agent, a CD47 antagonist (e.g., Hu5F9-G4, CC-90002, and CD47-Fc fusion protein TTI-621), PolyI:C, LPS, monophosphoryl A, imiquimod, R-848,

CpG-ODN, IFN- α , IFN- β , IFN- γ , GM-CSF, IL-12, IL-2, IL-15, T α 1, ibrutinib, EF-022 and combinations thereof.

88. The pharmaceutical composition of claim 80, wherein the macrophage activity inhibitory agent is selected from the group consisting of a STAT3 inhibitor, a STAT6 inhibitor, or an anti-tumor drug agent.
89. The pharmaceutical composition of claim 88, wherein the macrophage activity inhibitory agent is selected from the group consisting of WP1066, sunitinib, sorafenib, STA-21, IS3 295, S3I-M2001, AS1517499, leflunomide, TMC-264, histidine-rich glycoprotein (HRG), copper chelate (CuNG), 5,6-dimethylxanthenone-4-acetic acid (MDXAA), vadimezan (ASA404), cisplatin, silibinin, proton pump inhibitor pantoprazole (PPZ), CNI-1493 and combinations thereof.
90. The pharmaceutical composition of claim 80, wherein the macrophage inhibitor agent is an anti-IL-1 α agent (e.g., xilonix).
91. The pharmaceutical composition of any one of claims 63-90, wherein the second agent is a regulatory T cell (Treg) inhibitory agent selected from the group consisting of a Treg ablating agent, a Treg migration inhibitor agent, a Treg function inhibitor agent, and combinations thereof.
92. The pharmaceutical composition of claim 91, wherein the Treg ablating agent is selected from the group consisting of cyclophosphamide, paclitaxel, imatinib, sunitinib, sorafenib, dasatinib, temozolomide, daclizumab, denileukin diftitox, and combinations thereof.
93. The pharmaceutical composition of claim 91, wherein the Treg migration inhibitor agent is selected from the group consisting of AMD3100, mogamulizumab, casuarinin, fucoidan, and combinations thereof.
94. The pharmaceutical composition of claim 91, wherein the Treg function inhibitor agent is selected from the group consisting of an anti-CTLA4 agent (e.g., ipilimumab, tremelimumab), an anti-OX40 agent, an anti-GITR agent, an adenosine receptor antagonist (e.g., caffeine, theophylline, theobromine, and 8-phenylxanthines), P60, and combinations thereof.

95. The pharmaceutical composition of any one of claims 63-94, wherein the second agent is an antigen specific immune response enhancer agent selected from the group consisting of an anti-PD-1 agent, an anti-PD-L1 agent, a GITR (glucocorticoid-induced TNFR-related protein) stimulating agent, an anti-CTLA4 agent, an anti-TIM-3 agent, an anti-LAG-3 agent, an anti-IDO agent, an agent that enhances tumor antigen presentation (e.g., personalized cancer vaccine, autologous antigen presenting cell, autologous dendritic cells, artificial antigen presenting cell), a chemokine signaling agent, an anti-VEGF agent, a cytokine signal stimulating agent, and combinations thereof.
96. The pharmaceutical composition of claim 95, wherein the anti-PD-1 agent is selected from the group consisting of pembrolizumab, nivolumab, PDR001, REGN2810 (SAR-439684), BGB-A317, BI 754091, IBI308, INCISHR-1210, JNJ-63723283, JS-001, MEDI0680 (AMP-514), MGA-012, PF-06801591, REGN-2810, TSR-042, atezolizumab, avelumab, CX-072, durvalumab, FAZ053, LY3300054, PD-L1 millamolecule, PDR-001, camrelizumab (HR-301210), BCD-100, AGEN-2034, CS1001, Sym-021, LZM009, KN-035, AB122, genolimzumab (CBT-501), AK 104, GLS-010, BGB-A333, SHR-1316, CK-301, and combinations thereof.
97. The pharmaceutical composition of claim 95 or 96, wherein the anti-PD-L1 agent is selected from the group consisting of atezolizumab, durvalumab, avelumab, LY3300054, BGB-A333, SHR-1316, CK-301, and combinations thereof.
98. The pharmaceutical composition of any one of claims 95-97, wherein the GITR stimulating agent is selected from the group consisting of DTA-1, mGITRL, pGITRL, and combinations thereof.
99. The pharmaceutical composition of any one of claims 95-98, wherein the anti-CTLA4 agent is selected from the group consisting of ipilimumab, tremelimumab, and combinations thereof.
100. The pharmaceutical composition of any one of claims 95-99, wherein the chemokine signaling agent is selected from the group consisting of CXCL16, a CXCR6 chemokine receptor (CD186) agonist, and combinations thereof.
101. The pharmaceutical composition of any one of claims 95-100, wherein the anti-VEGF agent is selected from the group consisting of bevacizumab, pazopanib, sunitinib,

sorafenib, axitinib, ponatinib, regorafenib, cabozantinib, vandetanib, ramucirumab, lenvatinib, ziv-aflibercept, and combinations thereof.


102. The pharmaceutical composition of any one of claims 95-101, wherein the cytokine signal stimulating agent is an interleukin or an interferon.
103. The pharmaceutical composition of claim 102, wherein the interleukin is selected from the group consisting of IL-2, IL-1, IL-7, IL-15, IL-12, IL-18 and combinations thereof.
104. The pharmaceutical composition of claim 102, wherein the interferon is IFN alpha.
105. The pharmaceutical composition of any one of claims 63-104, wherein the second agent is an antigen specific immune response enhancer agent selected from the group consisting of a flavonoid (e.g., flavonoid glycoside), lidocaine, lamotrigine, sulfamethoxazole, phenytoin, carbamazepine, sulfamethoxazole, phenytoin, allopurinol, paracetamol, mepivacaine, p-phenylenediamine, ciprofloxacin and moxifloxacin.
106. The pharmaceutical composition of claim 105, wherein the disease or condition is cancer.
107. The pharmaceutical composition of claim 105, wherein the cancer is selected from the group consisting of endometrial cancer, breast cancer, ovarian cancer, cervical cancer, fallopian tube cancer, primary peritoneal cancer, colon cancer, squamous cell carcinoma of the anogenital region, melanoma, renal cell carcinoma, lung cancer, non-small cell lung cancer, squamous cell carcinoma of the lung, stomach cancer, bladder cancer, gall bladder cancer, liver cancer, thyroid cancer, laryngeal cancer, salivary gland cancer, esophageal cancer, squamous cell carcinoma of the head and neck, prostate cancer, pancreatic cancer, mesothelioma, sarcoma, a hematological cancer, and combinations thereof.
108. The pharmaceutical composition of any one of claims 63-107, wherein the administering comprises administering a composition comprising a capsule comprising the first agent.
109. The pharmaceutical composition of claim 108, wherein the capsule comprises a formulation comprising the first agent and one or more pharmaceutically acceptable excipients.

110. The pharmaceutical composition of claim 109, wherein the one or more pharmaceutically acceptable excipients comprises lactose monohydrate, magnesium stearate, or a combination thereof.
111. The pharmaceutical composition of any one of claims 63-110, wherein a therapeutically effective amount of the first or second agent is administered.
112. The pharmaceutical composition of any one of claims 63-111, wherein the method further comprises administering a third agent to the subject.
113. The pharmaceutical composition of claim 112, wherein the third agent comprises an antigen specific immune response enhancer agent, an anti-angiogenic agent, a chemotherapeutic agent, or combinations thereof.
114. The pharmaceutical composition of claim 113, wherein the antigen specific immune response enhancer agent comprises an anti-PD-1 agent, an anti-PD-L1 agent, an anti-CTLA4 agent, an anti-TIM-3 agent, or an anti-LAG-3 agent.
115. The pharmaceutical composition of claim 113, wherein the anti-angiogenic agent is selected from the group consisting of TNP-470, platelet factor 4, thrombospondin-1, tissue inhibitors of metalloproteases (TIMP1 and TIMP2), prolactin, angiostatin, endostatin, bFGF soluble receptor, transforming growth factor beta, interferon alpha, soluble KDR and FLT-1 receptors, placental proliferin-related protein, and combinations thereof.
116. The pharmaceutical composition of claim 113, wherein the chemotherapeutic agent is selected from the group consisting of aminoglutethimide, amsacrine, anastrozole, asparaginase, bcr, bicalutamide, bleomycin, buserelin, busulfan, camptothecin, capecitabine, carboplatin, carmustine, chlorambucil, cisplatin, cladribine, clodronate, colchicine, cyclophosphamide, cyproterone, cytarabine, dacarbazine, dactinomycin, daunorubicin, dienestrol, diethylstilbestrol, docetaxel, doxorubicin, epirubicin, estradiol, estramustine, etoposide, exemestane, filgrastim, fludarabine, fludrocortisone, fluorouracil, fluoxymesterone, flutamide, gemcitabine, genistein, goserelin, hydroxyurea, idarubicin, ifosfamide, imatinib, interferon, irinotecan, irinotecan, letrozole, leucovorin, leuprolide, levamisole, lomustine, mechlorethamine, medroxyprogesterone, megestrol, melphalan, mercaptopurine, mesna, methotrexate, mitomycin, mitotane, mitoxantrone,

nilutamide, nocodazole, octreotide, oxaliplatin, paclitaxel, pamidronate, pentostatin, plicamycin, porfimer, procarbazine, raltitrexed, rituximab, streptozocin, suramin, tamoxifen, temozolomide, teniposide, testosterone, thioguanine, thiotepa, titanocene dichloride, topotecan, trastuzumab, tretinoin, vinblastine, vincristine, vindesine, vinorelbine, and combinations thereof.

1/12

Niraparib



Vehicle




FIG. 1A





FIG. 1B

3/12

Niraparib



Vehicle




FIG. 2A

4/12




FIG. 2B

5/12

Niraparib



Vehicle




FIG. 3A

6/12




FIG. 3B

7/12

Niraparib



Vehicle



FIG. 4A




FIG. 4B

9/12




FIG. 5A

10/12




FIG. 5B




FIG. 6A

12/12




FIG. 6B




FIG. 6A