

(19)



(11) Publication number:

SG 185432 A1

(43) Publication date:

28.12.2012

(51) Int. Cl:

C12P 5/02, C12N 1/20;

(12)

Patent Application

(21) Application number: **2012081519**

(71) Applicant: **GENOMATICA, INC. 10520 WATERIDGE CIRCLE, SAN DIEGO, CA 92121 CA US**

(22) Date of filing: **04.05.2011**

(30) Priority: **US 61/331,821 05.05.2010**

(72) Inventor: **BURK, MARK, J. 10520 WATERIDGE CIRCLE SAN DIEGO, CA 92121 US
BURGARD, ANTHONY, P. 10520 WATERIDGE CIRCLE SAN DIEGO, CA 92121 US
SUN, JUN 10520 WATERIDGE CIRCLE SAN DIEGO, CA 92121 US
OSTERHOUT, ROBIN, E. 10520 WATERIDGE CIRCLE SAN DIEGO, CA 92121 US
PHARKYA, PRITI 10520 WATERIDGE CIRCLE SAN DIEGO, CA 92121 US**

(54) **Title:**
MICROORGANISMS AND METHODS FOR THE BIOSYNTHESIS OF BUTADIENE

(57) **Abstract:**
The invention provides non-naturally occurring microbial organisms having a butadiene pathway. The invention additionally provides methods of using such organisms to produce butadiene.

(51) International Patent Classification:
C12P 5/02 (2006.01)(21) International Application Number:
PCT/US2011/035105(22) International Filing Date:
4 May 2011 (04.05.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/331,821 5 May 2010 (05.05.2010) US(71) Applicant (for all designated States except US): **GENOMATICA, INC.** [US/US]; 10520 Wateridge Circle, San Diego, CA 92121 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BURK, Mark, J.** [US/US]; 10520 Wateridge Circle, San Diego, CA 92121 (US). **BURGARD, Anthony, P.** [US/US]; 10520 Wateridge Circle, San Diego, CA 92121 (US). **SUN, Jun** [CN/US]; 10520 Wateridge Circle, San Diego, CA 92121 (US). **OSTERHOUT, Robin, E.** [US/US]; 10520 Wateridge Circle, San Diego, CA 92121 (US). **PHARKYA, Priti** [IN/US]; 10520 Wateridge Circle, San Diego, CA 92121 (US).(74) Agents: **GAY, David, A.** et al.; Jones Day, 222 East 41st Street, New York, NY 10017-6702 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: MICROORGANISMS AND METHODS FOR THE BIOSYNTHESIS OF BUTADIENE



(57) Abstract: The invention provides non-naturally occurring microbial organisms having a butadiene pathway. The invention additionally provides methods of using such organisms to produce butadiene.

MICROORGANISMS AND METHODS FOR THE BIOSYNTHESIS OF BUTADIENE

5 BACKGROUND OF THE INVENTION

The present invention relates generally to biosynthetic processes, and more specifically to organisms having butadiene biosynthetic capability.

Over 25 billion pounds of butadiene (1,3-butadiene, BD) are produced annually and is applied in the manufacture of polymers such as synthetic rubbers and ABS resins, and
10 chemicals such as hexamethylenediamine and 1,4-butanediol. Butadiene is typically produced as a by-product of the steam cracking process for conversion of petroleum feedstocks such as naphtha, liquefied petroleum gas, ethane or natural gas to ethylene and other olefins. The ability to manufacture butadiene from alternative and/or renewable feedstocks would represent a major advance in the quest for more sustainable chemical
15 production processes

One possible way to produce butadiene renewably involves fermentation of sugars or other feedstocks to produce diols, such as 1,4-butanediol or 1,3-butanediol, which are separated, purified, and then dehydrated to butadiene in a second step involving metal-based catalysis. Direct fermentative production of butadiene from renewable feedstocks would obviate the
20 need for dehydration steps and butadiene gas (bp -4.4°C) would be continuously emitted from the fermenter and readily condensed and collected. Developing a fermentative production process would eliminate the need for fossil-based butadiene and would allow substantial savings in cost, energy, and harmful waste and emissions relative to petrochemically-derived butadiene.

25 Microbial organisms and methods for effectively producing butadiene from cheap renewable feedstocks such as molasses, sugar cane juice, and sugars derived from biomass sources, including agricultural and wood waste, as well as C1 feedstocks such as syngas and carbon dioxide, are described herein and include related advantages.

SUMMARY OF THE INVENTION

30 The invention provides non-naturally occurring microbial organisms containing butadiene pathways comprising at least one exogenous nucleic acid encoding a butadiene pathway enzyme expressed in a sufficient amount to produce butadiene. The invention additionally provides methods of using such microbial organisms to produce butadiene, by culturing a

non-naturally occurring microbial organism containing butadiene pathways as described herein under conditions and for a sufficient period of time to produce butadiene.

BRIEF DESCRIPTION OF THE DRAWINGS

- Figure 1 shows a natural pathway to isoprenoids and terpenes. Enzymes for transformation of the identified substrates to products include: A. acetyl-CoA:acetyl-CoA acyltransferase, B. hydroxymethylglutaryl-CoA synthase, C. 3-hydroxy-3-methylglutaryl-CoA reductase (alcohol forming), D. mevalonate kinase, E. phosphomevalonate kinase, F. diphosphomevalonate decarboxylase, G. isopentenyl-diphosphate isomerase, H. isoprene synthase.
- Figure 2 shows exemplary pathways for production of butadiene from acetyl-CoA, glutaconyl-CoA, glutaryl-CoA, 3-aminobutyryl-CoA or 4-hydroxybutyryl-CoA via crotyl alcohol. Enzymes for transformation of the identified substrates to products include: A. acetyl-CoA:acetyl-CoA acyltransferase, B. acetoacetyl-CoA reductase, C. 3-hydroxybutyryl-CoA dehydratase, D. crotonyl-CoA reductase (aldehyde forming), E. crotonaldehyde reductase (alcohol forming), F. crotyl alcohol kinase, G. 2-butenyl-4-phosphate kinase, H. butadiene synthase, I. crotonyl-CoA hydrolase, synthetase, transferase, J. crotonate reductase, K. crotonyl-CoA reductase (alcohol forming), L. glutaconyl-CoA decarboxylase, M., glutaryl-CoA dehydrogenase, N. 3-aminobutyryl-CoA deaminase, O. 4-hydroxybutyryl-CoA dehydratase, P. crotyl alcohol diphosphokinase.
- Figure 3 shows exemplary pathways for production of butadiene from erythrose-4-phosphate. Enzymes for transformation of the identified substrates to products include: A. Erythrose-4-phosphate reductase, B. Erythritol-4-phosphate cytidyltransferase, C. 4-(cytidine 5'-diphospho)-erythritol kinase, D. Erythritol 2,4-cyclodiphosphate synthase, E. 1-Hydroxy-2-butenyl 4-diphosphate synthase, F. 1-Hydroxy-2-butenyl 4-diphosphate reductase, G. Butenyl 4-diphosphate isomerase, H. Butadiene synthase I. Erythrose-4-phosphate kinase, J. Erythrose reductase, K. Erythritol kinase.
- Figure 4 shows an exemplary pathway for production of butadiene from malonyl-CoA plus acetyl-CoA. Enzymes for transformation of the identified substrates to products include: A. malonyl-CoA:acetyl-CoA acyltransferase, B. 3-oxoglutaryl-CoA reductase (ketone-reducing), C. 3-hydroxyglutaryl-CoA reductase (aldehyde forming), D. 3-hydroxy-5-oxopentanoate reductase, E. 3,5-dihydroxypentanoate kinase, F. 3H5PP kinase, G. 3H5PDP decarboxylase, H. butenyl 4-diphosphate isomerase, I. butadiene synthase, J. 3-

hydroxyglutaryl-CoA reductase (alcohol forming), K. 3-oxoglutaryl-CoA reductase (aldehyde forming), L. 3,5-dioxopentanoate reductase (ketone reducing), M. 3,5-dioxopentanoate reductase (aldehyde reducing), N. 5-hydroxy-3-oxopentanoate reductase, O. 3-oxo-glutaryl-CoA reductase (CoA reducing and alcohol forming). Compound
5 abbreviations include: 3H5PP = 3-Hydroxy-5-phosphonatooxypentanoate and 3H5PDP = 3-Hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to the design and production of cells and organisms having biosynthetic production capabilities for butadiene. The invention, in particular, relates to the
10 design of microbial organism capable of producing butadiene by introducing one or more nucleic acids encoding a butadiene pathway enzyme.

In one embodiment, the invention utilizes *in silico* stoichiometric models of *Escherichia coli* metabolism that identify metabolic designs for biosynthetic production of butadiene. The results described herein indicate that metabolic pathways can be designed and recombinantly
15 engineered to achieve the biosynthesis of butadiene in *Escherichia coli* and other cells or organisms. Biosynthetic production of butadiene, for example, for the *in silico* designs can be confirmed by construction of strains having the designed metabolic genotype. These metabolically engineered cells or organisms also can be subjected to adaptive evolution to further augment butadiene biosynthesis, including under conditions approaching theoretical
20 maximum growth.

In certain embodiments, the butadiene biosynthesis characteristics of the designed strains make them genetically stable and particularly useful in continuous bioprocesses. Separate strain design strategies were identified with incorporation of different non-native or heterologous reaction capabilities into *E. coli* or other host organisms leading to butadiene
25 producing metabolic pathways from acetyl-CoA, glutaconyl-CoA, glutaryl-CoA, 3-aminobutyryl-CoA, 4-hydroxybutyryl-CoA, erythrose-4-phosphate or malonyl-CoA plus acetyl-CoA. *In silico* metabolic designs were identified that resulted in the biosynthesis of butadiene in microorganisms from each of these substrates or metabolic intermediates.

Strains identified via the computational component of the platform can be put into actual
30 production by genetically engineering any of the predicted metabolic alterations, which lead to the biosynthetic production of butadiene or other intermediate and/or downstream products. In yet a further embodiment, strains exhibiting biosynthetic production of these

compounds can be further subjected to adaptive evolution to further augment product biosynthesis. The levels of product biosynthesis yield following adaptive evolution also can be predicted by the computational component of the system.

The maximum theoretical butadiene yield from glucose is 1.09 mol/mol (0.33 g/g).



The pathways presented in Figure(s) 2 and 4 achieve a yield of 1.0 moles butadiene per mole of glucose utilized. Increasing product yields to theoretical maximum value is possible if cells are capable of fixing CO₂ through pathways such as the reductive (or reverse) TCA cycle or the Wood-Ljungdahl pathway. Organisms engineered to possess the pathway depicted in
10 Figure 3 are also capable of reaching near theoretical maximum yields of butadiene.

As used herein, the term “non-naturally occurring” when used in reference to a microbial organism or microorganism of the invention is intended to mean that the microbial organism has at least one genetic alteration not normally found in a naturally occurring strain of the referenced species, including wild-type strains of the referenced species. Genetic alterations
15 include, for example, modifications introducing expressible nucleic acids encoding metabolic polypeptides, other nucleic acid additions, nucleic acid deletions and/or other functional disruption of the microbial organism’s genetic material. Such modifications include, for example, coding regions and functional fragments thereof, for heterologous, homologous or both heterologous and homologous polypeptides for the referenced species. Additional
20 modifications include, for example, non-coding regulatory regions in which the modifications alter expression of a gene or operon. Exemplary metabolic polypeptides include enzymes or proteins within a butadiene biosynthetic pathway.

A metabolic modification refers to a biochemical reaction that is altered from its naturally occurring state. Therefore, non-naturally occurring microorganisms can have genetic
25 modifications to nucleic acids encoding metabolic polypeptides, or functional fragments thereof. Exemplary metabolic modifications are disclosed herein.

As used herein, the term “butadiene,” having the molecular formula C₄H₆ and a molecular mass of 54.09 g/mol (see Figures 2-4) (IUPAC name Buta-1,3-diene) is used interchangeably throughout with 1,3-butadiene, biethylene, erythrene, divinyl, vinylethylene. Butadiene is a
30 colorless, non corrosive liquefied gas with a mild aromatic or gasoline-like odor. Butadiene is both explosive and flammable because of its low flash point.

As used herein, the term “isolated” when used in reference to a microbial organism is intended to mean an organism that is substantially free of at least one component as the referenced microbial organism is found in nature. The term includes a microbial organism that is removed from some or all components as it is found in its natural environment. The term also includes a microbial organism that is removed from some or all components as the microbial organism is found in non-naturally occurring environments. Therefore, an isolated microbial organism is partly or completely separated from other substances as it is found in nature or as it is grown, stored or subsisted in non-naturally occurring environments. Specific examples of isolated microbial organisms include partially pure microbes, substantially pure microbes and microbes cultured in a medium that is non-naturally occurring.

As used herein, the terms “microbial,” “microbial organism” or “microorganism” are intended to mean any organism that exists as a microscopic cell that is included within the domains of archaea, bacteria or eukarya. Therefore, the term is intended to encompass prokaryotic or eukaryotic cells or organisms having a microscopic size and includes bacteria, archaea and eubacteria of all species as well as eukaryotic microorganisms such as yeast and fungi. The term also includes cell cultures of any species that can be cultured for the production of a biochemical.

As used herein, the term “CoA” or “coenzyme A” is intended to mean an organic cofactor or prosthetic group (nonprotein portion of an enzyme) whose presence is required for the activity of many enzymes (the apoenzyme) to form an active enzyme system. Coenzyme A functions in certain condensing enzymes, acts in acetyl or other acyl group transfer and in fatty acid synthesis and oxidation, pyruvate oxidation and in other acetylation.

As used herein, the term “substantially anaerobic” when used in reference to a culture or growth condition is intended to mean that the amount of oxygen is less than about 10% of saturation for dissolved oxygen in liquid media. The term also is intended to include sealed chambers of liquid or solid medium maintained with an atmosphere of less than about 1% oxygen.

“Exogenous” as it is used herein is intended to mean that the referenced molecule or the referenced activity is introduced into the host microbial organism. The molecule can be introduced, for example, by introduction of an encoding nucleic acid into the host genetic material such as by integration into a host chromosome or as non-chromosomal genetic material such as a plasmid. Therefore, the term as it is used in reference to expression of an

encoding nucleic acid refers to introduction of the encoding nucleic acid in an expressible form into the microbial organism. When used in reference to a biosynthetic activity, the term refers to an activity that is introduced into the host reference organism. The source can be, for example, a homologous or heterologous encoding nucleic acid that expresses the
5 referenced activity following introduction into the host microbial organism. Therefore, the term “endogenous” refers to a referenced molecule or activity that is present in the host. Similarly, the term when used in reference to expression of an encoding nucleic acid refers to expression of an encoding nucleic acid contained within the microbial organism. The term “heterologous” refers to a molecule or activity derived from a source other than the
10 referenced species whereas “homologous” refers to a molecule or activity derived from the host microbial organism. Accordingly, exogenous expression of an encoding nucleic acid of the invention can utilize either or both a heterologous or homologous encoding nucleic acid.

It is understood that when more than one exogenous nucleic acid is included in a microbial organism that the more than one exogenous nucleic acids refers to the referenced encoding
15 nucleic acid or biosynthetic activity, as discussed above. It is further understood, as disclosed herein, that such more than one exogenous nucleic acids can be introduced into the host microbial organism on separate nucleic acid molecules, on polycistronic nucleic acid molecules, or a combination thereof, and still be considered as more than one exogenous nucleic acid. For example, as disclosed herein a microbial organism can be engineered to
20 express two or more exogenous nucleic acids encoding a desired pathway enzyme or protein. In the case where two exogenous nucleic acids encoding a desired activity are introduced into a host microbial organism, it is understood that the two exogenous nucleic acids can be introduced as a single nucleic acid, for example, on a single plasmid, on separate plasmids, can be integrated into the host chromosome at a single site or multiple sites, and still be
25 considered as two exogenous nucleic acids. Similarly, it is understood that more than two exogenous nucleic acids can be introduced into a host organism in any desired combination, for example, on a single plasmid, on separate plasmids, can be integrated into the host chromosome at a single site or multiple sites, and still be considered as two or more exogenous nucleic acids, for example three exogenous nucleic acids. Thus, the number of
30 referenced exogenous nucleic acids or biosynthetic activities refers to the number of encoding nucleic acids or the number of biosynthetic activities, not the number of separate nucleic acids introduced into the host organism.

The non-naturally occurring microbial organisms of the invention can contain stable genetic alterations, which refers to microorganisms that can be cultured for greater than five generations without loss of the alteration. Generally, stable genetic alterations include modifications that persist greater than 10 generations, particularly stable modifications will persist more than about 25 generations, and more particularly, stable genetic modifications will be greater than 50 generations, including indefinitely.

Those skilled in the art will understand that the genetic alterations, including metabolic modifications exemplified herein, are described with reference to a suitable host organism such as *E. coli* and their corresponding metabolic reactions or a suitable source organism for desired genetic material such as genes for a desired metabolic pathway. However, given the complete genome sequencing of a wide variety of organisms and the high level of skill in the area of genomics, those skilled in the art will readily be able to apply the teachings and guidance provided herein to essentially all other organisms. For example, the *E. coli* metabolic alterations exemplified herein can readily be applied to other species by incorporating the same or analogous encoding nucleic acid from species other than the referenced species. Such genetic alterations include, for example, genetic alterations of species homologs, in general, and in particular, orthologs, paralogs or nonorthologous gene displacements.

An ortholog is a gene or genes that are related by vertical descent and are responsible for substantially the same or identical functions in different organisms. For example, mouse epoxide hydrolase and human epoxide hydrolase can be considered orthologs for the biological function of hydrolysis of epoxides. Genes are related by vertical descent when, for example, they share sequence similarity of sufficient amount to indicate they are homologous, or related by evolution from a common ancestor. Genes can also be considered orthologs if they share three-dimensional structure but not necessarily sequence similarity, of a sufficient amount to indicate that they have evolved from a common ancestor to the extent that the primary sequence similarity is not identifiable. Genes that are orthologous can encode proteins with sequence similarity of about 25% to 100% amino acid sequence identity. Genes encoding proteins sharing an amino acid similarity less than 25% can also be considered to have arisen by vertical descent if their three-dimensional structure also shows similarities. Members of the serine protease family of enzymes, including tissue plasminogen activator and elastase, are considered to have arisen by vertical descent from a common ancestor.

Orthologs include genes or their encoded gene products that through, for example, evolution, have diverged in structure or overall activity. For example, where one species encodes a gene product exhibiting two functions and where such functions have been separated into distinct genes in a second species, the three genes and their corresponding products are considered to be orthologs. For the production of a biochemical product, those skilled in the art will understand that the orthologous gene harboring the metabolic activity to be introduced or disrupted is to be chosen for construction of the non-naturally occurring microorganism. An example of orthologs exhibiting separable activities is where distinct activities have been separated into distinct gene products between two or more species or within a single species. A specific example is the separation of elastase proteolysis and plasminogen proteolysis, two types of serine protease activity, into distinct molecules as plasminogen activator and elastase. A second example is the separation of mycoplasma 5'-3' exonuclease and *Drosophila* DNA polymerase III activity. The DNA polymerase from the first species can be considered an ortholog to either or both of the exonuclease or the polymerase from the second species and vice versa.

In contrast, paralogous are homologs related by, for example, duplication followed by evolutionary divergence and have similar or common, but not identical functions. Paralogous can originate or derive from, for example, the same species or from a different species. For example, microsomal epoxide hydrolase (epoxide hydrolase I) and soluble epoxide hydrolase (epoxide hydrolase II) can be considered paralogous because they represent two distinct enzymes, co-evolved from a common ancestor, that catalyze distinct reactions and have distinct functions in the same species. Paralogous are proteins from the same species with significant sequence similarity to each other suggesting that they are homologous, or related through co-evolution from a common ancestor. Groups of paralogous protein families include HipA homologs, luciferase genes, peptidases, and others.

A nonorthologous gene displacement is a nonorthologous gene from one species that can substitute for a referenced gene function in a different species. Substitution includes, for example, being able to perform substantially the same or a similar function in the species of origin compared to the referenced function in the different species. Although generally, a nonorthologous gene displacement will be identifiable as structurally related to a known gene encoding the referenced function, less structurally related but functionally similar genes and their corresponding gene products nevertheless will still fall within the meaning of the term as it is used herein. Functional similarity requires, for example, at least some structural

similarity in the active site or binding region of a nonorthologous gene product compared to a gene encoding the function sought to be substituted. Therefore, a nonorthologous gene includes, for example, a paralog or an unrelated gene.

Therefore, in identifying and constructing the non-naturally occurring microbial organisms of the invention having butadiene biosynthetic capability, those skilled in the art will understand with applying the teaching and guidance provided herein to a particular species that the identification of metabolic modifications can include identification and inclusion or inactivation of orthologs. To the extent that paralogs and/or nonorthologous gene displacements are present in the referenced microorganism that encode an enzyme catalyzing a similar or substantially similar metabolic reaction, those skilled in the art also can utilize these evolutionally related genes.

Orthologs, paralogs and nonorthologous gene displacements can be determined by methods well known to those skilled in the art. For example, inspection of nucleic acid or amino acid sequences for two polypeptides will reveal sequence identity and similarities between the compared sequences. Based on such similarities, one skilled in the art can determine if the similarity is sufficiently high to indicate the proteins are related through evolution from a common ancestor. Algorithms well known to those skilled in the art, such as Align, BLAST, Clustal W and others compare and determine a raw sequence similarity or identity, and also determine the presence or significance of gaps in the sequence which can be assigned a weight or score. Such algorithms also are known in the art and are similarly applicable for determining nucleotide sequence similarity or identity. Parameters for sufficient similarity to determine relatedness are computed based on well known methods for calculating statistical similarity, or the chance of finding a similar match in a random polypeptide, and the significance of the match determined. A computer comparison of two or more sequences can, if desired, also be optimized visually by those skilled in the art. Related gene products or proteins can be expected to have a high similarity, for example, 25% to 100% sequence identity. Proteins that are unrelated can have an identity which is essentially the same as would be expected to occur by chance, if a database of sufficient size is scanned (about 5%). Sequences between 5% and 24% may or may not represent sufficient homology to conclude that the compared sequences are related. Additional statistical analysis to determine the significance of such matches given the size of the data set can be carried out to determine the relevance of these sequences.

Exemplary parameters for determining relatedness of two or more sequences using the BLAST algorithm, for example, can be as set forth below. Briefly, amino acid sequence alignments can be performed using BLASTP version 2.0.8 (Jan-05-1999) and the following parameters: Matrix: 0 BLOSUM62; gap open: 11; gap extension: 1; x_dropoff: 50; expect: 10.0; wordsize: 3; filter: on. Nucleic acid sequence alignments can be performed using BLASTN version 2.0.6 (Sept-16-1998) and the following parameters: Match: 1; mismatch: -2; gap open: 5; gap extension: 2; x_dropoff: 50; expect: 10.0; wordsize: 11; filter: off. Those skilled in the art will know what modifications can be made to the above parameters to either increase or decrease the stringency of the comparison, for example, and determine the relatedness of two or more sequences.

In some embodiments, the invention provides a non-naturally occurring microbial organism, including a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding a butadiene pathway enzyme expressed in a sufficient amount to produce butadiene, the butadiene pathway including an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase, or transferase, a crotonate reductase, a crotonyl-CoA reductase (alcohol forming), a glutaconyl-CoA decarboxylase, a glutaryl-CoA dehydrogenase, an 3-aminobutyryl-CoA deaminase, a 4-hydroxybutyryl-CoA dehydratase or a crotyl alcohol diphosphokinase (Figure 2). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase (Figure 2, steps A-H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase and crotonyl-CoA reductase (alcohol forming) (Figure 2, steps A-C, K, F, G, H). In one aspect, the non-naturally occurring microbial

organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a butadiene synthase, a crotonyl-CoA reductase (alcohol forming) and a crotyl alcohol diphosphokinase (Figure 2, steps A-C, K, P, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase, or transferase and a crotonate reductase, (Figure 2, steps A-C, I, J, E, F, G, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase, a crotonate reductase and a crotyl alcohol diphosphokinase (Figure 2, steps A-C, I, J, E, P, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a butadiene synthase and a crotyl alcohol diphosphokinase (Figure 2, steps A-E, P, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a glutaconyl-CoA decarboxylase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase (Figure 2, steps L, D-H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway

including a glutaconyl-CoA decarboxylase, a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase and crotonyl-CoA reductase (alcohol forming) (Figure 2, steps L, K, F, G, H). In one aspect, the non-naturally occurring microbial organism includes a

microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a glutaconyl-CoA decarboxylase, a butadiene synthase, a crotonyl-CoA reductase (alcohol forming) and a crotyl alcohol diphosphokinase (Figure 2, steps L, K, P, H). In one aspect, the non-naturally occurring microbial organism includes a

microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene,

the butadiene pathway including a glutaconyl-CoA decarboxylase, a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase, or transferase and a crotonate reductase (Figure 2, steps L, I, J, E, F, G, H). In one aspect, the non-naturally occurring

microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a glutaconyl-CoA decarboxylase, a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase, a crotonate reductase and a crotyl alcohol

diphosphokinase (Figure 2, steps L, I, J, E, P, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase

(alcohol forming), a butadiene a glutaconyl-CoA decarboxylase and a crotyl alcohol diphosphokinase (Figure 2, steps L, C, D, E, P, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a glutaryl-CoA

dehydrogenase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase (Figure 2, steps M, D-H). In one aspect, the non-naturally occurring microbial

organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient

amount to produce butadiene, the butadiene pathway including a glutaryl-CoA

dehydrogenase, a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase and crotonyl-CoA reductase (alcohol forming) (Figure 2, steps M, K, F, G, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a glutaryl-CoA dehydrogenase, a butadiene synthase, a crotonyl-CoA reductase (alcohol forming) and a crotyl alcohol diphosphokinase (Figure 2, steps M, K, P, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a glutaryl-CoA dehydrogenase, a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase, or transferase and a crotonate reductase (Figure 2, steps M, I, J, E, F, G, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a glutaryl-CoA dehydrogenase, a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase, a crotonate reductase and a crotyl alcohol diphosphokinase (Figure 2, steps M, I, J, E, P, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a glutaryl-CoA dehydrogenase and a crotyl alcohol diphosphokinase (Figure 2, steps M, C, D, E, P, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including an 3-aminobutyryl-CoA deaminase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase (Figure 2, steps N, D-H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including an 3-aminobutyryl-CoA deaminase, a crotyl alcohol kinase, a 2-butenyl-4-

phosphate kinase, a butadiene synthase and crotonyl-CoA reductase (alcohol forming) (Figure 2, steps N, K, F, G, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including an 3-aminobutyryl-CoA deaminase, a butadiene synthase, a crotonyl-CoA reductase (alcohol forming) and a crotyl alcohol diphosphokinase (Figure 2, steps N, K, P, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including an 3-aminobutyryl-CoA deaminase, a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase, or transferase and a crotonate reductase (Figure 2, steps N, I, J, E, F, G, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including an 3-aminobutyryl-CoA deaminase, a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase, a crotonate reductase and a crotyl alcohol diphosphokinase (Figure 2, steps N, I, J, E, P, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a 3-aminobutyryl-CoA deaminase and a crotyl alcohol diphosphokinase (Figure 2, steps N, C, D, E, P, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a 4-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase (Figure 2, steps O, D-H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a 4-hydroxybutyryl-CoA dehydratase, a crotyl alcohol kinase, a 2-butenyl-4-

phosphate kinase, a butadiene synthase and crotonyl-CoA reductase (alcohol forming) (Figure 2, steps O, K, F, G, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a 4-hydroxybutyryl-CoA dehydratase, a butadiene synthase, a crotonyl-CoA reductase (alcohol forming) and a crotyl alcohol diphosphokinase (Figure 2, steps O, K, P, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a 4-hydroxybutyryl-CoA dehydratase, a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase, or transferase and a crotonate reductase (Figure 2, steps O, I, J, E, F, G, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a 4-hydroxybutyryl-CoA dehydratase, a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase, a crotonate reductase and a crotyl alcohol diphosphokinase (Figure 2, steps O, I, J, E, P, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a 4-hydroxybutyryl-CoA dehydratase and a crotyl alcohol diphosphokinase (Figure 2, steps L, C, D, E, P, H).

In some embodiments, the invention provides a non-naturally occurring microbial organism, including a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding a butadiene pathway enzyme expressed in a sufficient amount to produce butadiene, the butadiene pathway including an erythrose-4-phosphate reductase, an erythritol-4-phosphate cytidylyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an erythritol 2,4-cyclodiphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase, a butenyl 4-diphosphate isomerase, a butadiene synthase, an erythrose-4-phosphate kinase, an erythrose reductase or an erythritol kinase

(Figure 3). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including an erythrose-4-phosphate reductase, an erythritol-4-phosphate cytidyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an erythritol 2,4-cyclodiphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase and a butadiene synthase (Figure 3, steps A-F, and H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including an erythrose-4-phosphate reductase, an erythritol-4-phosphate cytidyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an erythritol 2,4-cyclodiphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase, a butenyl 4-diphosphate isomerase and butadiene synthase (Figure 3, steps A-H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including an erythritol-4-phosphate cytidyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an erythritol 2,4-cyclodiphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase, a butadiene synthase, an erythrose-4-phosphate kinase, an erythrose reductase and a erythritol kinase (Figure 3, steps I, J, K, B-F, H). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including an erythritol-4-phosphate cytidyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an erythritol 2,4-cyclodiphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase, a butenyl 4-diphosphate isomerase, a butadiene synthase, an erythrose-4-phosphate kinase, an erythrose reductase and an erythritol kinase (Figure 3, steps I, J, K, B-H).

In some embodiments, the invention provides a non-naturally occurring microbial organism, including a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding a butadiene pathway enzyme expressed in a sufficient amount to produce butadiene, the butadiene pathway including a malonyl-CoA:acetyl-CoA

acyltransferase, an 3-oxoglutaryl-CoA reductase (ketone-reducing), a 3-hydroxyglutaryl-CoA reductase (aldehyde forming), a 3-hydroxy-5-oxopentanoate reductase, a 3,5-dihydroxypentanoate kinase, a 3-hydroxy-5-phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase, a butadiene synthase, a 3-hydroxyglutaryl-CoA reductase (alcohol forming), an 3-oxoglutaryl-CoA reductase (aldehyde forming), a 3,5-dioxopentanoate reductase (ketone reducing), a 3,5-dioxopentanoate reductase (aldehyde reducing), a 5-hydroxy-3-oxopentanoate reductase or an 3-oxo-glutaryl-CoA reductase (CoA reducing and alcohol forming) (Figure 4). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a malonyl-CoA:acetyl-CoA acyltransferase, an 3-oxoglutaryl-CoA reductase (ketone-reducing), a 3-hydroxyglutaryl-CoA reductase (aldehyde forming), a 3-hydroxy-5-oxopentanoate reductase, a 3,5-dihydroxypentanoate kinase, a 3-hydroxy-5-phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase and a butadiene synthase (Figure 4, steps A-I). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a malonyl-CoA:acetyl-CoA acyltransferase, a 3,5-dihydroxypentanoate kinase, a 3-hydroxy-5-phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase, a butadiene synthase, an 3-oxoglutaryl-CoA reductase (aldehyde forming), a 3,5-dioxopentanoate reductase (aldehyde reducing) and a 5-hydroxy-3-oxopentanoate reductase. (Figure 4, steps A, K, M, N, E, F, G, H, I). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a malonyl-CoA:acetyl-CoA acyltransferase, a 3-hydroxy-5-oxopentanoate reductase, a 3,5-dihydroxypentanoate kinase, a 3-Hydroxy-5-phosphonatooxypentanoate kinase, a 3-Hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase, a butadiene synthase, an 3-oxoglutaryl-CoA reductase (aldehyde forming) and a 3,5-dioxopentanoate reductase (ketone reducing). (Figure 4, steps A, K, L, D, E, F, G, H, I). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway

having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a malonyl-CoA:acetyl-CoA acyltransferase, a 3,5-dihydroxypentanoate kinase, a 3-hydroxy-5-phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase, a butadiene synthase, a 5-hydroxy-3-oxopentanoate reductase and a 3-oxo-glutaryl-CoA reductase (CoA reducing and alcohol forming). (Figure 4, steps A, O, N, E, F, G, H, I). In one aspect, the non-naturally occurring microbial organism includes a microbial organism having a butadiene pathway having at least one exogenous nucleic acid encoding butadiene pathway enzymes expressed in a sufficient amount to produce butadiene, the butadiene pathway including a malonyl-CoA:acetyl-CoA acyltransferase, an 3-oxoglutaryl-CoA reductase (ketone-reducing), a 3,5-dihydroxypentanoate kinase, a 3-hydroxy-5-phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase, a butadiene synthase and a 3-hydroxyglutaryl-CoA reductase (alcohol forming). (Figure 4, steps A, B, J, E, F, G, H, I).

In an additional embodiment, the invention provides a non-naturally occurring microbial organism having a butadiene pathway, wherein the non-naturally occurring microbial organism comprises at least one exogenous nucleic acid encoding an enzyme or protein that converts a substrate to a product selected from the group consisting of acetyl-CoA to acetoacetyl-CoA, acetoacetyl-CoA to 3-hydroxybutyryl-CoA, 3-hydroxybutyryl-CoA to crotonyl-CoA, crotonyl-CoA to crotonaldehyde, crotonaldehyde to crotyl alcohol, crotyl alcohol to 2-butenyl-phosphate, 2-butenyl-phosphate to 2-butenyl-4-diphosphate, 2-butenyl-4-diphosphate to butadiene, erythrose-4-phosphate to erythritol-4-phosphate, erythritol-4-phosphate to 4-(cytidine 5'-diphospho)-erythritol, 4-(cytidine 5'-diphospho)-erythritol to 2-phospho-4-(cytidine 5'-diphospho)-erythritol, 2-phospho-4-(cytidine 5'-diphospho)-erythritol to erythritol-2,4-cyclodiphosphate, erythritol-2,4-cyclodiphosphate to 1-hydroxy-2-butenyl 4-diphosphate, 1-hydroxy-2-butenyl 4-diphosphate to butenyl 4-diphosphate, butenyl 4-diphosphate to 2-butenyl 4-diphosphate, 1-hydroxy-2-butenyl 4-diphosphate to 2-butenyl 4-diphosphate, 2-butenyl 4-diphosphate to butadiene, malonyl-CoA and acetyl-CoA to 3-oxoglutaryl-CoA, 3-oxoglutaryl-CoA to 3-hydroxyglutaryl-CoA to 3-hydroxy-5-oxopentanoate, 3-hydroxy-5-oxopentanoate to 3,5-dihydroxy pentanoate, 3,5-dihydroxy pentanoate to 3-hydroxy-5-phosphonatooxypentanoate, 3-hydroxy-5-phosphonatooxypentanoate to 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate, 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate to butenyl 4-

biphosphate, glutaconyl-CoA to crotonyl-CoA, glutaryl-CoA to crotonyl-CoA, 3-aminobutyryl-CoA to crotonyl-CoA, 4-hydroxybutyryl-CoA to crotonyl-CoA, crotonyl-CoA to crotonate, crotonate to crotonaldehyde, crotonyl-CoA to crotyl alcohol, crotyl alcohol to 2-butenyl-4-diphosphate, erythrose-4-phosphate to erythrose, erythrose to erythritol, erythritol to erythritol-4-phosphate, 3-oxoglutaryl-CoA to 3,5-dioxopentanoate, 3,5-dioxopentanoate to 5-hydroxy-3-oxopentanoate, 5-hydroxy-3-oxopentanoate to 3,5-dihydroxypentanoate, 3-oxoglutaryl-CoA to 5-hydroxy-3-oxopentanoate, 3,5-dioxopentanoate to 3-hydroxy-5-oxopentanoate and 3-hydroxyglutaryl-CoA to 3,5-dihydroxypentanoate. One skilled in the art will understand that these are merely exemplary and that any of the substrate-product pairs disclosed herein suitable to produce a desired product and for which an appropriate activity is available for the conversion of the substrate to the product can be readily determined by one skilled in the art based on the teachings herein. Thus, the invention provides a non-naturally occurring microbial organism containing at least one exogenous nucleic acid encoding an enzyme or protein, where the enzyme or protein converts the substrates and products of a butadiene pathway, such as that shown in Figures 2-4.

While generally described herein as a microbial organism that contains a butadiene pathway, it is understood that the invention additionally provides a non-naturally occurring microbial organism comprising at least one exogenous nucleic acid encoding a butadiene pathway enzyme expressed in a sufficient amount to produce an intermediate of a butadiene pathway.

For example, as disclosed herein, a butadiene pathway is exemplified in Figures 2-4.

Therefore, in addition to a microbial organism containing a butadiene pathway that produces butadiene, the invention additionally provides a non-naturally occurring microbial organism comprising at least one exogenous nucleic acid encoding a butadiene pathway enzyme, where the microbial organism produces a butadiene pathway intermediate, for example, acetoacetyl-CoA, 3-hydroxybutyryl-CoA, crotonyl-CoA, crotonaldehyde, crotyl alcohol, 2-butenyl-phosphate, 2-butenyl-4-diphosphate, erythritol-4-phosphate, 4-(cytidine 5'-diphospho)-erythritol, 2-phospho-4-(cytidine 5'-diphospho)-erythritol, erythritol-2,4-cyclodiphosphate, 1-hydroxy-2-butenyl 4-diphosphate, butenyl 4-diphosphate, 2-butenyl 4-diphosphate, 3-oxoglutaryl-CoA, 3-hydroxyglutaryl-CoA, 3-hydroxy-5-oxopentanoate, 3,5-dihydroxypentanoate, 3-hydroxy-5-phosphonatooxypentanoate, 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate, crotonate, erythrose, erythritol, 3,5-dioxopentanoate or 5-hydroxy-3-oxopentanoate.

It is understood that any of the pathways disclosed herein, as described in the Examples and exemplified in the Figures, including the pathways of Figures 2-4, can be utilized to generate a non-naturally occurring microbial organism that produces any pathway intermediate or product, as desired. As disclosed herein, such a microbial organism that produces an
5 intermediate can be used in combination with another microbial organism expressing downstream pathway enzymes to produce a desired product. However, it is understood that a non-naturally occurring microbial organism that produces a butadiene pathway intermediate can be utilized to produce the intermediate as a desired product.

The invention is described herein with general reference to the metabolic reaction, reactant or
10 product thereof, or with specific reference to one or more nucleic acids or genes encoding an enzyme associated with or catalyzing, or a protein associated with, the referenced metabolic reaction, reactant or product. Unless otherwise expressly stated herein, those skilled in the art will understand that reference to a reaction also constitutes reference to the reactants and products of the reaction. Similarly, unless otherwise expressly stated herein, reference to a
15 reactant or product also references the reaction, and reference to any of these metabolic constituents also references the gene or genes encoding the enzymes that catalyze or proteins involved in the referenced reaction, reactant or product. Likewise, given the well known fields of metabolic biochemistry, enzymology and genomics, reference herein to a gene or encoding nucleic acid also constitutes a reference to the corresponding encoded enzyme and
20 the reaction it catalyzes or a protein associated with the reaction as well as the reactants and products of the reaction.

As disclosed herein, the intermediats crotonate; 3,5-dioxopentanoate, 5-hydroxy-3-oxopentanoate, 3-hydroxy-5-oxopentanoate, 3-oxoglutaryl-CoA and 3-hydroxyglutaryl-CoA, as well as other intermediates, are carboxylic acids, which can occur in various ionized
25 forms, including fully protonated, partially protonated, and fully deprotonated forms. Accordingly, the suffix “-ate,” or the acid form, can be used interchangeably to describe both the free acid form as well as any deprotonated form, in particular since the ionized form is known to depend on the pH in which the compound is found. It is understood that carboxylate products or intermediates includes ester forms of carboxylate products or
30 pathway intermediates, such as O-carboxylate and S-carboxylate esters. O- and S-carboxylates can include lower alkyl, that is C1 to C6, branched or straight chain carboxylates. Some such O- or S-carboxylates include, without limitation, methyl, ethyl, n-propyl, n-butyl, i-propyl, sec-butyl, and tert-butyl, pentyl, hexyl O- or S-carboxylates, any of

which can further possess an unsaturation, providing for example, propenyl, butenyl, pentyl, and hexenyl O- or S-carboxylates. O-carboxylates can be the product of a biosynthetic pathway. Exemplary O-carboxylates accessed via biosynthetic pathways can include, without limitation: methyl crotonate; methyl-3,5-dioxopentanoate; methyl-5-hydroxy-3-oxopentanoate; methyl-3-hydroxy-5-oxopentanoate; 3-oxoglutaryl-CoA, methyl ester; 3-hydroxyglutaryl-CoA, methyl ester; ethyl crotonate; ethyl-3,5-dioxopentanoate; ethyl-5-hydroxy-3-oxopentanoate; ethyl-3-hydroxy-5-oxopentanoate; 3-oxoglutaryl-CoA, ethyl ester; 3-hydroxyglutaryl-CoA, ethyl ester; n-propyl crotonate; n-propyl-3,5-dioxopentanoate; n-propyl-5-hydroxy-3-oxopentanoate; n-propyl-3-hydroxy-5-oxopentanoate; 3-oxoglutaryl-CoA, n-propyl ester; and 3-hydroxyglutaryl-CoA, n-propyl ester. Other biosynthetically accessible O-carboxylates can include medium to long chain groups, that is C7-C22, O-carboxylate esters derived from fatty alcohols, such heptyl, octyl, nonyl, decyl, undecyl, lauryl, tridecyl, myristyl, pentadecyl, cetyl, palmitolyl, heptadecyl, stearyl, nonadecyl, arachidyl, heneicosyl, and behenyl alcohols, any one of which can be optionally branched and/or contain unsaturations. O-carboxylate esters can also be accessed via a biochemical or chemical process, such as esterification of a free carboxylic acid product or transesterification of an O- or S-carboxylate. S-carboxylates are exemplified by CoA S-esters, cysteinyl S-esters, alkylthioesters, and various aryl and heteroaryl thioesters.

The non-naturally occurring microbial organisms of the invention can be produced by introducing expressible nucleic acids encoding one or more of the enzymes or proteins participating in one or more butadiene biosynthetic pathways. Depending on the host microbial organism chosen for biosynthesis, nucleic acids for some or all of a particular butadiene biosynthetic pathway can be expressed. For example, if a chosen host is deficient in one or more enzymes or proteins for a desired biosynthetic pathway, then expressible nucleic acids for the deficient enzyme(s) or protein(s) are introduced into the host for subsequent exogenous expression. Alternatively, if the chosen host exhibits endogenous expression of some pathway genes, but is deficient in others, then an encoding nucleic acid is needed for the deficient enzyme(s) or protein(s) to achieve butadiene biosynthesis. Thus, a non-naturally occurring microbial organism of the invention can be produced by introducing exogenous enzyme or protein activities to obtain a desired biosynthetic pathway or a desired biosynthetic pathway can be obtained by introducing one or more exogenous enzyme or protein activities that, together with one or more endogenous enzymes or proteins, produces a desired product such as butadiene.

Host microbial organisms can be selected from, and the non-naturally occurring microbial organisms generated in, for example, bacteria, yeast, fungus or any of a variety of other microorganisms applicable to fermentation processes. Exemplary bacteria include species selected from *Escherichia coli*, *Klebsiella oxytoca*, *Anaerobiospirillum succiniciproducens*,
5 *Actinobacillus succinogenes*, *Mannheimia succiniciproducens*, *Rhizobium etli*, *Bacillus subtilis*, *Corynebacterium glutamicum*, *Gluconobacter oxydans*, *Zymomonas mobilis*, *Lactococcus lactis*, *Lactobacillus plantarum*, *Streptomyces coelicolor*, *Clostridium acetobutylicum*, *Pseudomonas fluorescens*, and *Pseudomonas putida*. Exemplary yeasts or fungi include species selected from *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*,
10 *Kluyveromyces lactis*, *Kluyveromyces marxianus*, *Aspergillus terreus*, *Aspergillus niger*, *Pichia pastoris*, *Rhizopus arrhizus*, *Rhizobus oryzae*, *Yarrowia lipolytica*, and the like. *E. coli* is a particularly useful host organism since it is a well characterized microbial organism suitable for genetic engineering. Other particularly useful host organisms include yeast such as *Saccharomyces cerevisiae*. It is understood that any suitable microbial host organism can
15 be used to introduce metabolic and/or genetic modifications to produce a desired product.

Depending on the butadiene biosynthetic pathway constituents of a selected host microbial organism, the non-naturally occurring microbial organisms of the invention will include at least one exogenously expressed butadiene pathway-encoding nucleic acid and up to all encoding nucleic acids for one or more butadiene biosynthetic pathways. For example,
20 butadiene biosynthesis can be established in a host deficient in a pathway enzyme or protein through exogenous expression of the corresponding encoding nucleic acid. In a host deficient in all enzymes or proteins of a butadiene pathway, exogenous expression of all enzyme or proteins in the pathway can be included, although it is understood that all enzymes or proteins of a pathway can be expressed even if the host contains at least one of the pathway enzymes
25 or proteins. For example, exogenous expression of all enzymes or proteins in a pathway for production of butadiene can be included, such as an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase (Figure 2, steps A-H).

30 Given the teachings and guidance provided herein, those skilled in the art will understand that the number of encoding nucleic acids to introduce in an expressible form will, at least, parallel the butadiene pathway deficiencies of the selected host microbial organism. Therefore, a non-naturally occurring microbial organism of the invention can have one, two,

three, four, five, six, seven, eight, nine or ten, up to all nucleic acids encoding the enzymes or proteins constituting a butadiene biosynthetic pathway disclosed herein. In some embodiments, the non-naturally occurring microbial organisms also can include other genetic modifications that facilitate or optimize butadiene biosynthesis or that confer other useful functions onto the host microbial organism. One such other functionality can include, for example, augmentation of the synthesis of one or more of the butadiene pathway precursors such as acetyl-CoA, glutaconyl-CoA, glutaryl-CoA, 3-aminobutyryl-CoA, 4-hydroxybutyryl-CoA, erythrose-4-phosphate or malonyl-CoA.

Generally, a host microbial organism is selected such that it produces the precursor of a butadiene pathway, either as a naturally produced molecule or as an engineered product that either provides *de novo* production of a desired precursor or increased production of a precursor naturally produced by the host microbial organism. For example, acetyl-CoA, glutaconyl-CoA, glutaryl-CoA, 3-aminobutyryl-CoA, 4-hydroxybutyryl-CoA, erythrose-4-phosphate or malonyl-CoA are produced naturally in a host organism such as *E. coli*. A host organism can be engineered to increase production of a precursor, as disclosed herein. In addition, a microbial organism that has been engineered to produce a desired precursor can be used as a host organism and further engineered to express enzymes or proteins of a butadiene pathway.

In some embodiments, a non-naturally occurring microbial organism of the invention is generated from a host that contains the enzymatic capability to synthesize butadiene. In this specific embodiment it can be useful to increase the synthesis or accumulation of a butadiene pathway product to, for example, drive butadiene pathway reactions toward butadiene production. Increased synthesis or accumulation can be accomplished by, for example, overexpression of nucleic acids encoding one or more of the above-described butadiene pathway enzymes or proteins. Overexpression the enzyme or enzymes and/or protein or proteins of the butadiene pathway can occur, for example, through exogenous expression of the endogenous gene or genes, or through exogenous expression of the heterologous gene or genes. Therefore, naturally occurring organisms can be readily generated to be non-naturally occurring microbial organisms of the invention, for example, producing butadiene, through overexpression of one, two, three, four, five, six, seven, eight, nine, or ten, that is, up to all nucleic acids encoding butadiene biosynthetic pathway enzymes or proteins. In addition, a non-naturally occurring organism can be generated by mutagenesis of an endogenous gene that results in an increase in activity of an enzyme in the butadiene biosynthetic pathway.

In particularly useful embodiments, exogenous expression of the encoding nucleic acids is employed. Exogenous expression confers the ability to custom tailor the expression and/or regulatory elements to the host and application to achieve a desired expression level that is controlled by the user. However, endogenous expression also can be utilized in other
5 embodiments such as by removing a negative regulatory effector or induction of the gene's promoter when linked to an inducible promoter or other regulatory element. Thus, an endogenous gene having a naturally occurring inducible promoter can be up-regulated by providing the appropriate inducing agent, or the regulatory region of an endogenous gene can be engineered to incorporate an inducible regulatory element, thereby allowing the regulation
10 of increased expression of an endogenous gene at a desired time. Similarly, an inducible promoter can be included as a regulatory element for an exogenous gene introduced into a non-naturally occurring microbial organism.

It is understood that, in methods of the invention, any of the one or more exogenous nucleic acids can be introduced into a microbial organism to produce a non-naturally occurring
15 microbial organism of the invention. The nucleic acids can be introduced so as to confer, for example, a butadiene biosynthetic pathway onto the microbial organism. Alternatively, encoding nucleic acids can be introduced to produce an intermediate microbial organism having the biosynthetic capability to catalyze some of the required reactions to confer butadiene biosynthetic capability. For example, a non-naturally occurring microbial
20 organism having a butadiene biosynthetic pathway can comprise at least two exogenous nucleic acids encoding desired enzymes or proteins, such as the combination of a crotyl alcohol kinase and a butadiene synthase, or alternatively a 4-(cytidine 5'-diphospho)-erythritol kinase and butadiene synthase, or alternatively a 1-hydroxy-2-butenyl 4-diphosphate synthase and a butadiene synthase, or alternatively a 3-hydroxy-5-
25 phosphonatooxypentanoate kinase and a butadiene synthase, or alternatively a crotonyl-CoA hydrolase and a crotyl alcohol diphosphokinase, or alternatively a an erythrose reductase and butadiene synthase or alternatively an 3-oxo-glutaryl-CoA reductase (CoA reducing and alcohol forming) and 3-Hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, and the like. Thus, it is understood that any combination of two or more
30 enzymes or proteins of a biosynthetic pathway can be included in a non-naturally occurring microbial organism of the invention. Similarly, it is understood that any combination of three or more enzymes or proteins of a biosynthetic pathway can be included in a non-naturally occurring microbial organism of the invention, for example, a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase, or alternatively a 1-hydroxy-2-butenyl

4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase, and butadiene synthase, or alternatively an 3-oxoglutaryl-CoA reductase, a 3-hydroxy-5-oxopentanoate reductase, and a butadiene synthase, or alternatively an acetyl-CoA:acetyl-CoA acyltransferase, a crotyl alcohol kinase and a butadiene synthase, or alternatively a
5 glutaconyl-CoA decarboxylase, a crotonyl-CoA reductase (alcohol forming), and a crotyl alcohol diphosphokinase, or alternatively a an erythrose-4-phosphate kinase, a 4-(cytidine 5'-diphospho)-erythritol kinase and a 1-hydroxy-2-butenyl 4-diphosphate synthase, or alternatively a 3,5-dioxopentanoate reductase (aldehyde reducing), a butenyl 4-diphosphate isomerase, and a butadiene synthase, and so forth, as desired, so long as the combination of
10 enzymes and/or proteins of the desired biosynthetic pathway results in production of the corresponding desired product. Similarly, any combination of four, such as a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase, or alternatively a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase, a butenyl 4-diphosphate isomerase and
15 butadiene synthase, or alternatively a 3-hydroxy-5-phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate kinase, a butenyl 4-diphosphate isomerase and a butadiene synthase, or alternatively an erythrose-4-phosphate reductase, an erythritol-4-phosphate cytidyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase and butadiene synthase, or alternatively an 3-aminobutyryl-CoA deaminase, a
20 crotonyl-CoA reductase (alcohol forming), a crotyl alcohol diphosphokinase and a butadiene synthase, or alternatively an erythrose reductase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an erythritol 2,4-cyclodiphosphate synthase and a 1-hydroxy-2-butenyl 4-diphosphate reductase, or alternatively a malonyl-CoA:acetyl-CoA acyltransferase, a 3-hydroxyglutaryl-CoA reductase (alcohol forming), a butenyl 4-diphosphate isomerase and a butadiene
25 synthase, or more enzymes or proteins of a biosynthetic pathway as disclosed herein can be included in a non-naturally occurring microbial organism of the invention, as desired, so long as the combination of enzymes and/or proteins of the desired biosynthetic pathway results in production of the corresponding desired product.

In addition to the biosynthesis of butadiene as described herein, the non-naturally occurring
30 microbial organisms and methods of the invention also can be utilized in various combinations with each other and with other microbial organisms and methods well known in the art to achieve product biosynthesis by other routes. For example, one alternative to produce butadiene other than use of the butadiene producers is through addition of another microbial organism capable of converting a butadiene pathway intermediate to butadiene.

One such procedure includes, for example, the fermentation of a microbial organism that produces a butadiene pathway intermediate. The butadiene pathway intermediate can then be used as a substrate for a second microbial organism that converts the butadiene pathway intermediate to butadiene. The butadiene pathway intermediate can be added directly to another culture of the second organism or the original culture of the butadiene pathway intermediate producers can be depleted of these microbial organisms by, for example, cell separation, and then subsequent addition of the second organism to the fermentation broth can be utilized to produce the final product without intermediate purification steps.

In other embodiments, the non-naturally occurring microbial organisms and methods of the invention can be assembled in a wide variety of subpathways to achieve biosynthesis of, for example, butadiene. In these embodiments, biosynthetic pathways for a desired product of the invention can be segregated into different microbial organisms, and the different microbial organisms can be co-cultured to produce the final product. In such a biosynthetic scheme, the product of one microbial organism is the substrate for a second microbial organism until the final product is synthesized. For example, the biosynthesis of butadiene can be accomplished by constructing a microbial organism that contains biosynthetic pathways for conversion of one pathway intermediate to another pathway intermediate or the product. Alternatively, butadiene also can be biosynthetically produced from microbial organisms through co-culture or co-fermentation using two organisms in the same vessel, where the first microbial organism produces a butadiene intermediate and the second microbial organism converts the intermediate to butadiene.

Given the teachings and guidance provided herein, those skilled in the art will understand that a wide variety of combinations and permutations exist for the non-naturally occurring microbial organisms and methods of the invention together with other microbial organisms, with the co-culture of other non-naturally occurring microbial organisms having subpathways and with combinations of other chemical and/or biochemical procedures well known in the art to produce butadiene.

Sources of encoding nucleic acids for a butadiene pathway enzyme or protein can include, for example, any species where the encoded gene product is capable of catalyzing the referenced reaction. Such species include both prokaryotic and eukaryotic organisms including, but not limited to, bacteria, including archaea and eubacteria, and eukaryotes, including yeast, plant, insect, animal, and mammal, including human. Exemplary species for such sources include, for example, *Escherichia coli*, *Acidaminococcus fermentans*, *Acinetobacter baylyi*,

- Acinetobacter calcoaceticus*, *Acinetobacter* sp. ADP1, *Acinetobacter* sp. Strain M-1, *Aquifex aeolicus*, *Arabidopsis thaliana*, *Arabidopsis thaliana* col, *Arabidopsis thaliana* col, *Archaeoglobus fulgidus* DSM 4304, *Azoarcus* sp. CIB, *Bacillus cereus*, *Bacillus subtilis*, *Bos Taurus*, *Brucella melitensis*, *Burkholderia ambifaria* AMMD, *Burkholderia phymatum*,
- 5 *Campylobacter jejuni*, *Candida albicans*, *Candida magnoliae*, *Chloroflexus aurantiacus*, *Citrobacter youngae* ATCC 29220, *Clostridium acetobutylicum*, *Clostridium aminobutyricum*, *Clostridium beijerinckii*, *Clostridium beijerinckii* NCIMB 8052, *Clostridium beijerinckii* NRRL B593, *Clostridium botulinum* C str. Eklund, *Clostridium kluyveri*, *Clostridium kluyveri* DSM 555, *Clostridium novyi* NT, *Clostridium propionicum*, *Clostridium*
- 10 *saccharoperbutylacetonicum*, *Corynebacterium glutamicum* ATCC 13032, *Cupriavidus taiwanensis*, *Cyanobium* PCC7001, *Dictyostelium discoideum* AX4, *Enterococcus faecalis*, *Erythrobacter* sp. NAPI, *Escherichia coli* K12, *Escherichia coli* str. K-12 substr. MG1655, *Eubacterium rectale* ATCC 33656, *Fusobacterium nucleatum*, *Fusobacterium nucleatum* subsp. *nucleatum* ATCC 25586, *Geobacillus thermoglucosidasius*, *Haematococcus pluvialis*,
- 15 *Haemophilus influenzae*, *Haloarcula marismortui* ATCC 43049, *Helicobacter pylori*, *Homo sapiens*, *Klebsiella pneumoniae*, *Lactobacillus plantarum*, *Leuconostoc mesenteroides*, *marine gamma proteobacterium* HTCC2080, *Metallosphaera sedula*, *Methanocaldococcus jannaschii*, *Mus musculus*, *Mycobacterium avium* subsp. *paratuberculosis* K-10, *Mycobacterium bovis* BCG, *Mycobacterium marinum* M, *Mycobacterium smegmatis* MC2
- 20 155, *Mycobacterium tuberculosis*, *Mycoplasma pneumoniae* M129, *Nocardia farcinica* IFM 10152, *Nocardia iowensis* (sp. NRRL 5646), *Oryctolagus cuniculus*, *Paracoccus denitrificans*, *Penicillium chrysogenum*, *Populus alba*, *Populus tremula* x *Populus alba*, *Porphyromonas gingivalis*, *Porphyromonas gingivalis* W83, *Pseudomonas aeruginosa*, *Pseudomonas aeruginosa* PAO1, *Pseudomonas fluorescens*, *Pseudomonas fluorescens* Pf-5,
- 25 *Pseudomonas knackmussii* (B13), *Pseudomonas putida*, *Pseudomonas putida* E23, *Pseudomonas putida* KT2440, *Pseudomonas* sp, *Pueraria Montana*, *Pyrobaculum aerophilum* str. IM2, *Pyrococcus furiosus*, *Ralstonia eutropha*, *Ralstonia eutropha* H16, *Ralstonia eutropha* H16, *Ralstonia metallidurans*, *Rattus norvegicus*, *Rhodobacter spaeroides*, *Rhodococcus rubber*, *Rhodopseudomonas palustris*, *Roseburia intestinalis* L1-82,
- 30 *Roseburia inulinivorans* DSM 16841, *Roseburia* sp. A2-183, *Roseiflexus castenholzii*, *Saccharomyces cerevisiae*, *Saccharopolyspora rythraea* NRRL 2338, *Salmonella enterica* subsp. *arizonae* serovar, *Salmonella typhimurium*, *Schizosaccharomyces pombe*, *Simmondsia chinensis*, *Sinorhizobium meliloti*, *Staphylococcus* , *ureus*, *Streptococcus pneumoniae*, *Streptomyces coelicolor*, *Streptomyces griseus* subsp. *griseus* , BRC 13350, *Streptomyces* sp.
- 35 *ACT-1*, *Sulfolobus acidocaldarius*, *Sulfolobus shibatae*, *Sulfolobus solfataricus*, *Sulfolobus*

tokodaii, *Synechocystis* sp. strain PCC6803, *Syntrophus*, *ciditrophicus*, *Thermoanaerobacter brockii* HTD4, *Thermoanaerobacter tengcongensis* MB4, *Thermosynechococcus elongates*, *Thermotoga maritime* MSB8, *Thermus thermophilus*, *Thermus, hermophilus* HB8, *Trichomonas vaginalis* G3, *Trichosporonoides megachiliensis*, *Trypanosoma brucei*,

- 5 *Tsukamurella paurometabola* DSM 20162, *Yersinia intermedia* ATCC 29909, *Zoogloea ramigera*, *Zygosaccharomyces rouxii*, *Zymomonas mobilis*, as well as other exemplary species disclosed herein are available as source organisms for corresponding genes.

However, with the complete genome sequence available for now more than 550 species (with more than half of these available on public databases such as the NCBI), including 395
10 microorganism genomes and a variety of yeast, fungi, plant, and mammalian genomes, the identification of genes encoding the requisite butadiene biosynthetic activity for one or more genes in related or distant species, including for example, homologues, orthologs, paralogs and nonorthologous gene displacements of known genes, and the interchange of genetic alterations between organisms is routine and well known in the art. Accordingly, the
15 metabolic alterations allowing biosynthesis of butadiene described herein with reference to a particular organism such as *E. coli* can be readily applied to other microorganisms, including prokaryotic and eukaryotic organisms alike. Given the teachings and guidance provided herein, those skilled in the art will know that a metabolic alteration exemplified in one organism can be applied equally to other organisms.

20 In some instances, such as when an alternative butadiene biosynthetic pathway exists in an unrelated species, butadiene biosynthesis can be conferred onto the host species by, for example, exogenous expression of a paralog or paralogs from the unrelated species that catalyzes a similar, yet non-identical metabolic reaction to replace the referenced reaction. Because certain differences among metabolic networks exist between different organisms,
25 those skilled in the art will understand that the actual gene usage between different organisms may differ. However, given the teachings and guidance provided herein, those skilled in the art also will understand that the teachings and methods of the invention can be applied to all microbial organisms using the cognate metabolic alterations to those exemplified herein to construct a microbial organism in a species of interest that will synthesize butadiene.

30 Methods for constructing and testing the expression levels of a non-naturally occurring butadiene-producing host can be performed, for example, by recombinant and detection methods well known in the art. Such methods can be found described in, for example, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Third Ed., Cold Spring Harbor

Laboratory, New York (2001); and Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley and Sons, Baltimore, MD (1999).

Exogenous nucleic acid sequences involved in a pathway for production of butadiene can be introduced stably or transiently into a host cell using techniques well known in the art including, but not limited to, conjugation, electroporation, chemical transformation, transduction, transfection, and ultrasound transformation. For exogenous expression in *E. coli* or other prokaryotic cells, some nucleic acid sequences in the genes or cDNAs of eukaryotic nucleic acids can encode targeting signals such as an N-terminal mitochondrial or other targeting signal, which can be removed before transformation into prokaryotic host cells, if desired. For example, removal of a mitochondrial leader sequence led to increased expression in *E. coli* (Hoffmeister et al., *J. Biol. Chem.* 280:4329-4338 (2005)). For exogenous expression in yeast or other eukaryotic cells, genes can be expressed in the cytosol without the addition of leader sequence, or can be targeted to mitochondrion or other organelles, or targeted for secretion, by the addition of a suitable targeting sequence such as a mitochondrial targeting or secretion signal suitable for the host cells. Thus, it is understood that appropriate modifications to a nucleic acid sequence to remove or include a targeting sequence can be incorporated into an exogenous nucleic acid sequence to impart desirable properties. Furthermore, genes can be subjected to codon optimization with techniques well known in the art to achieve optimized expression of the proteins.

An expression vector or vectors can be constructed to include one or more butadiene biosynthetic pathway encoding nucleic acids as exemplified herein operably linked to expression control sequences functional in the host organism. Expression vectors applicable for use in the microbial host organisms of the invention include, for example, plasmids, phage vectors, viral vectors, episomes and artificial chromosomes, including vectors and selection sequences or markers operable for stable integration into a host chromosome. Additionally, the expression vectors can include one or more selectable marker genes and appropriate expression control sequences. Selectable marker genes also can be included that, for example, provide resistance to antibiotics or toxins, complement auxotrophic deficiencies, or supply critical nutrients not in the culture media. Expression control sequences can include constitutive and inducible promoters, transcription enhancers, transcription terminators, and the like which are well known in the art. When two or more exogenous encoding nucleic acids are to be co-expressed, both nucleic acids can be inserted, for example, into a single expression vector or in separate expression vectors. For single vector

expression, the encoding nucleic acids can be operationally linked to one common expression control sequence or linked to different expression control sequences, such as one inducible promoter and one constitutive promoter. The transformation of exogenous nucleic acid sequences involved in a metabolic or synthetic pathway can be confirmed using methods well known in the art. Such methods include, for example, nucleic acid analysis such as Northern blots or polymerase chain reaction (PCR) amplification of mRNA, or immunoblotting for expression of gene products, or other suitable analytical methods to test the expression of an introduced nucleic acid sequence or its corresponding gene product. It is understood by those skilled in the art that the exogenous nucleic acid is expressed in a sufficient amount to produce the desired product, and it is further understood that expression levels can be optimized to obtain sufficient expression using methods well known in the art and as disclosed herein.

In some embodiments, the invention provides a method for producing butadiene that includes culturing a non-naturally occurring microbial organism, including a microbial organism having a butadiene pathway, the butadiene pathway including at least one exogenous nucleic acid encoding a butadiene pathway enzyme expressed in a sufficient amount to produce butadiene, the butadiene pathway including an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase, or transferase, a crotonate reductase, a crotonyl-CoA reductase (alcohol forming), a glutaconyl-CoA decarboxylase, a glutaryl-CoA dehydrogenase, an 3-aminobutyryl-CoA deaminase, a 4-hydroxybutyryl-CoA dehydratase or a crotyl alcohol diphosphokinase (Figure 2). In one aspect, the method includes a microbial organism having a butadiene pathway including an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase (Figure 2, steps A-H). In one aspect, the method includes a microbial organism having a butadiene pathway including an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a butadiene synthase, a

crotonyl-CoA reductase (alcohol forming) and a crotyl alcohol diphosphokinase (Figure 2, steps A-C, K, P, H). In one aspect, the method includes a microbial organism having a butadiene pathway including an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase, or transferase and a crotonate reductase, (Figure 2, steps A-C, I, J, E, F, G, H). In one aspect, the method includes a microbial organism having a butadiene pathway including an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase, a crotonate reductase and a crotyl alcohol diphosphokinase (Figure 2, steps A-C, I, J, E, P, H). In one aspect, the method includes a microbial organism having a butadiene pathway including an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a butadiene synthase and a crotyl alcohol diphosphokinase (Figure 2, steps A-E, P, H). In one aspect, the method includes a microbial organism having a butadiene pathway including a glutaconyl-CoA decarboxylase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase (Figure 2, steps L, D-H). In one aspect, the method includes a microbial organism having a butadiene pathway including a glutaconyl-CoA decarboxylase, a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase and crotonyl-CoA reductase (alcohol forming) (Figure 2, steps L, K, F, G, H). In one aspect, the method includes a microbial organism having a butadiene pathway including a glutaconyl-CoA decarboxylase, a butadiene synthase, a crotonyl-CoA reductase (alcohol forming) and a crotyl alcohol diphosphokinase (Figure 2, steps L, K, P, H). In one aspect, the method includes a microbial organism having a butadiene pathway including a glutaconyl-CoA decarboxylase, a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase, or transferase and a crotonate reductase (Figure 2, steps L, I, J, E, F, G, H). In one aspect, the method includes a microbial organism having a butadiene pathway including a glutaconyl-CoA decarboxylase, a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase, a crotonate reductase and a crotyl alcohol diphosphokinase (Figure 2, steps L, I, J, E, P, H). In one aspect, the method includes a microbial organism having a butadiene pathway including a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a

crotonaldehyde reductase (alcohol forming), a butadiene a glutaconyl-CoA decarboxylase and a crotyl alcohol diphosphokinase (Figure 2, steps L, C, D, E, P, H). In one aspect, the method includes a microbial organism having a butadiene pathway including a glutaryl-CoA dehydrogenase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase (Figure 2, steps M, D-H). In one aspect, the method includes a microbial organism having a butadiene pathway including a glutaryl-CoA dehydrogenase, a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase and crotonyl-CoA reductase (alcohol forming) (Figure 2, steps M, K, F, G, H). In one aspect, the method includes a microbial organism having a butadiene pathway including a glutaryl-CoA dehydrogenase, a butadiene synthase, a crotonyl-CoA reductase (alcohol forming) and a crotyl alcohol diphosphokinase (Figure 2, steps M, K, P, H). In one aspect, the method includes a microbial organism having a butadiene pathway including a glutaryl-CoA dehydrogenase, a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase, or transferase and a crotonate reductase (Figure 2, steps M, I, J, E, F, G, H). In one aspect, the method includes a microbial organism having a butadiene pathway including a glutaryl-CoA dehydrogenase, a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase, a crotonate reductase and a crotyl alcohol diphosphokinase (Figure 2, steps M, I, J, E, P, H). In one aspect, the method includes a microbial organism having a butadiene pathway including a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a glutaryl-CoA dehydrogenase and a crotyl alcohol diphosphokinase (Figure 2, steps M, C, D, E, P, H). In one aspect, the method includes a microbial organism having a butadiene pathway including an 3-aminobutyryl-CoA deaminase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase (Figure 2, steps N, D-H). In one aspect, the method includes a microbial organism having a butadiene pathway including an 3-aminobutyryl-CoA deaminase, a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase and crotonyl-CoA reductase (alcohol forming) (Figure 2, steps N, K, F, G, H). In one aspect, the method includes a microbial organism having a butadiene pathway including an 3-aminobutyryl-CoA deaminase, a butadiene synthase, a crotonyl-CoA reductase (alcohol forming) and a crotyl alcohol diphosphokinase (Figure 2, steps N, K, P, H). In one aspect, the method includes a microbial organism having a butadiene pathway including an 3-aminobutyryl-CoA deaminase, a crotonaldehyde reductase (alcohol forming),

a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase, or transferase and a crotonate reductase (Figure 2, steps N, I, J, E, F, G, H). In one aspect, the method includes a microbial organism having a butadiene pathway including an 3-aminobutyryl-CoA deaminase, a crotonaldehyde reductase (alcohol forming),
5 a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase, a crotonate reductase and a crotyl alcohol diphosphokinase (Figure 2, steps N, I, J, E, P, H). In one aspect, the method includes a microbial organism having a butadiene pathway including a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a 3-aminobutyryl-CoA
10 deaminase and a crotyl alcohol diphosphokinase (Figure 2, steps N, C, D, E, P, H). In one aspect, the method includes a microbial organism having a butadiene pathway including a 4-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase (Figure 2, steps O, D-H). In one aspect, the method includes
15 a microbial organism having a butadiene pathway including a 4-hydroxybutyryl-CoA dehydratase, a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase and crotonyl-CoA reductase (alcohol forming) (Figure 2, steps O, K, F, G, H). In one aspect, the method includes a microbial organism having a butadiene pathway including a 4-hydroxybutyryl-CoA dehydratase, a butadiene synthase, a crotonyl-CoA reductase (alcohol
20 forming) and a crotyl alcohol diphosphokinase (Figure 2, steps O, K, P, H). In one aspect, the method includes a microbial organism having a butadiene pathway including a 4-hydroxybutyryl-CoA dehydratase, a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase, or transferase and a crotonate reductase (Figure 2, steps O, I, J, E, F,
25 G, H). In one aspect, the method includes a microbial organism having a butadiene pathway including a 4-hydroxybutyryl-CoA dehydratase, a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase, a crotonate reductase and a crotyl alcohol diphosphokinase (Figure 2, steps O, I, J, E, P, H). In one aspect, the method includes a microbial organism having a butadiene pathway including
30 a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a 4-hydroxybutyryl-CoA dehydratase and a crotyl alcohol diphosphokinase (Figure 2, steps O, C, D, E, P, H).

In some embodiments, the invention provides a method for producing butadiene that includes culturing a non-naturally occurring microbial organism, including a microbial organism

having a butadiene pathway, the butadiene pathway including at least one exogenous nucleic acid encoding a butadiene pathway enzyme expressed in a sufficient amount to produce butadiene, the butadiene pathway including an erythrose-4-phosphate reductase, an erythritol-4-phosphate cytidylyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an erythritol 2,4-cyclodiphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase, a butenyl 4-diphosphate isomerase, a butadiene synthase, an erythrose-4-phosphate kinase, an erythrose reductase or an erythritol kinase (Figure 3). In one aspect, the method includes a microbial organism having a butadiene pathway including an erythrose-4-phosphate reductase, an erythritol-4-phosphate cytidylyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an erythritol 2,4-cyclodiphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase and a butadiene synthase (Figure 3, steps A-F, and H). In one aspect, the method includes a microbial organism having a butadiene pathway including an erythrose-4-phosphate reductase, an erythritol-4-phosphate cytidylyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an erythritol 2,4-cyclodiphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase, a butenyl 4-diphosphate isomerase and butadiene synthase (Figure 3, steps A-H). In one aspect, the method includes a microbial organism having a butadiene pathway including an erythritol-4-phosphate cytidylyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an erythritol 2,4-cyclodiphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase, a butadiene synthase, an erythrose-4-phosphate kinase, an erythrose reductase and an erythritol kinase (Figure 3, steps I, J, K, B-F, H). In one aspect, the method includes a microbial organism having a butadiene pathway including an erythritol-4-phosphate cytidylyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an erythritol 2,4-cyclodiphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase, a butenyl 4-diphosphate isomerase, a butadiene synthase, an erythrose-4-phosphate kinase, an erythrose reductase and an erythritol kinase (Figure 3, steps I, J, K, B-H).

In some embodiments, the invention provides a method for producing butadiene that includes culturing a non-naturally occurring microbial organism, including a microbial organism having a butadiene pathway, the butadiene pathway including at least one exogenous nucleic acid encoding a butadiene pathway enzyme expressed in a sufficient amount to produce butadiene, the butadiene pathway including a malonyl-CoA:acetyl-CoA acyltransferase, an 3-oxoglutaryl-CoA reductase (ketone-reducing), a 3-hydroxyglutaryl-CoA reductase (aldehyde

forming), a 3-hydroxy-5-oxopentanoate reductase, a 3,5-dihydroxypentanoate kinase, a 3-hydroxy-5-phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase, a butadiene synthase, a 3-hydroxyglutaryl-CoA reductase (alcohol forming), an 3-oxoglutaryl-CoA reductase (aldehyde forming), a 3,5-dioxopentanoate reductase (ketone reducing), a 3,5-dioxopentanoate reductase (aldehyde reducing), a 5-hydroxy-3-oxopentanoate reductase or an 3-oxo-glutaryl-CoA reductase (CoA reducing and alcohol forming) (Figure 4). In one aspect, the method includes a microbial organism having a butadiene pathway including a malonyl-CoA:acetyl-CoA acyltransferase, an 3-oxoglutaryl-CoA reductase (ketone-reducing), a 3-hydroxyglutaryl-CoA reductase (aldehyde forming), a 3-hydroxy-5-oxopentanoate reductase, a 3,5-dihydroxypentanoate kinase, a 3-hydroxy-5-phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase and a butadiene synthase (Figure 4, steps A-I). In one aspect, the method includes a microbial organism having a butadiene pathway including a malonyl-CoA:acetyl-CoA acyltransferase, a 3,5-dihydroxypentanoate kinase, a 3-hydroxy-5-phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase, a butadiene synthase, an 3-oxoglutaryl-CoA reductase (aldehyde forming), a 3,5-dioxopentanoate reductase (aldehyde reducing) and a 5-hydroxy-3-oxopentanoate reductase. (Figure 4, steps A, K, M, N, E, F, G, H, I). In one aspect, the method includes a microbial organism having a butadiene pathway including a malonyl-CoA:acetyl-CoA acyltransferase, a 3-hydroxy-5-oxopentanoate reductase, a 3,5-dihydroxypentanoate kinase, a 3-Hydroxy-5-phosphonatooxypentanoate kinase, a 3-Hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase, a butadiene synthase, an 3-oxoglutaryl-CoA reductase (aldehyde forming) and a 3,5-dioxopentanoate reductase (ketone reducing). (Figure 4, steps A, K, L, D, E, F, G, H, I). In one aspect, the method includes a microbial organism having a butadiene pathway including a malonyl-CoA:acetyl-CoA acyltransferase, a 3,5-dihydroxypentanoate kinase, a 3-hydroxy-5-phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase, a butadiene synthase, a 5-hydroxy-3-oxopentanoate reductase and a 3-oxo-glutaryl-CoA reductase (CoA reducing and alcohol forming). (Figure 4, steps A, O, N, E, F, G, H, I). In one aspect, the method includes a microbial organism having a butadiene pathway including a malonyl-CoA:acetyl-CoA acyltransferase, an 3-oxoglutaryl-CoA reductase (ketone-reducing), a 3,5-dihydroxypentanoate kinase, a 3-hydroxy-5-

phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase, a butadiene synthase and a 3-hydroxyglutaryl-CoA reductase (alcohol forming). (Figure 4, steps A, B, J, E, F, G, H, I).

- Suitable purification and/or assays to test for the production of butadiene can be performed
- 5 using well known methods. Suitable replicates such as triplicate cultures can be grown for each engineered strain to be tested. For example, product and byproduct formation in the engineered production host can be monitored. The final product and intermediates, and other organic compounds, can be analyzed by methods such as HPLC (High Performance Liquid Chromatography), GC-MS (Gas Chromatography-Mass Spectroscopy) and LC-MS (Liquid
- 10 Chromatography-Mass Spectroscopy) or other suitable analytical methods using routine procedures well known in the art. The release of product in the fermentation broth can also be tested with the culture supernatant. Byproducts and residual glucose can be quantified by HPLC using, for example, a refractive index detector for glucose and alcohols, and a UV detector for organic acids (Lin et al., *Biotechnol. Bioeng.* 90:775-779 (2005)), or other
- 15 suitable assay and detection methods well known in the art. The individual enzyme or protein activities from the exogenous DNA sequences can also be assayed using methods well known in the art. For typical Assay Methods, see Manual on Hydrocarbon Analysis (ASTM Manula Series, A.W. Drews, ed., 6th edition, 1998, American Society for Testing and Materials, Baltimore, Maryland).
- 20 The butadiene can be separated from other components in the culture using a variety of methods well known in the art. Such separation methods include, for example, extraction procedures as well as methods that include continuous liquid-liquid extraction, pervaporation, membrane filtration, membrane separation, reverse osmosis, electrodialysis, distillation, crystallization, centrifugation, extractive filtration, ion exchange chromatography, size
- 25 exclusion chromatography, adsorption chromatography, and ultrafiltration. All of the above methods are well known in the art.

Any of the non-naturally occurring microbial organisms described herein can be cultured to produce and/or secrete the biosynthetic products of the invention. For example, the butadiene producers can be cultured for the biosynthetic production of butadiene.

- 30 For the production of butadiene, the recombinant strains are cultured in a medium with carbon source and other essential nutrients. It is sometimes desirable and can be highly desirable to maintain anaerobic conditions in the fermenter to reduce the cost of the overall

process. Such conditions can be obtained, for example, by first sparging the medium with nitrogen and then sealing the flasks with a septum and crimp-cap. For strains where growth is not observed anaerobically, microaerobic or substantially anaerobic conditions can be applied by perforating the septum with a small hole for limited aeration. Exemplary

5 anaerobic conditions have been described previously and are well-known in the art. Exemplary aerobic and anaerobic conditions are described, for example, in United State publication 2009/0047719, filed August 10, 2007. Fermentations can be performed in a batch, fed-batch or continuous manner, as disclosed herein.

If desired, the pH of the medium can be maintained at a desired pH, in particular neutral pH,
10 such as a pH of around 7 by addition of a base, such as NaOH or other bases, or acid, as needed to maintain the culture medium at a desirable pH. The growth rate can be determined by measuring optical density using a spectrophotometer (600 nm), and the glucose uptake rate by monitoring carbon source depletion over time.


The growth medium can include, for example, any carbohydrate source which can supply a
15 source of carbon to the non-naturally occurring microorganism. Such sources include, for example, sugars such as glucose, xylose, arabinose, galactose, mannose, fructose, sucrose and starch. Other sources of carbohydrate include, for example, renewable feedstocks and biomass. Exemplary types of biomasses that can be used as feedstocks in the methods of the invention include cellulosic biomass, hemicellulosic biomass and lignin feedstocks or
20 portions of feedstocks. Such biomass feedstocks contain, for example, carbohydrate substrates useful as carbon sources such as glucose, xylose, arabinose, galactose, mannose, fructose and starch. Given the teachings and guidance provided herein, those skilled in the art will understand that renewable feedstocks and biomass other than those exemplified above also can be used for culturing the microbial organisms of the invention for the production of
25 butadiene.

In addition to renewable feedstocks such as those exemplified above, the butadiene microbial organisms of the invention also can be modified for growth on syngas as its source of carbon. In this specific embodiment, one or more proteins or enzymes are expressed in the butadiene producing organisms to provide a metabolic pathway for utilization of syngas or other
30 gaseous carbon source.

Synthesis gas, also known as syngas or producer gas, is the major product of gasification of coal and of carbonaceous materials such as biomass materials, including agricultural crops

and residues. Syngas is a mixture primarily of H₂ and CO and can be obtained from the gasification of any organic feedstock, including but not limited to coal, coal oil, natural gas, biomass, and waste organic matter. Gasification is generally carried out under a high fuel to oxygen ratio. Although largely H₂ and CO, syngas can also include CO₂ and other gases in smaller quantities. Thus, synthesis gas provides a cost effective source of gaseous carbon such as CO and, additionally, CO₂.

The Wood-Ljungdahl pathway catalyzes the conversion of CO and H₂ to acetyl-CoA and other products such as acetate. Organisms capable of utilizing CO and syngas also generally have the capability of utilizing CO₂ and CO₂/H₂ mixtures through the same basic set of enzymes and transformations encompassed by the Wood-Ljungdahl pathway. H₂-dependent conversion of CO₂ to acetate by microorganisms was recognized long before it was revealed that CO also could be used by the same organisms and that the same pathways were involved. Many acetogens have been shown to grow in the presence of CO₂ and produce compounds such as acetate as long as hydrogen is present to supply the necessary reducing equivalents (see for example, Drake, *Acetogenesis*, pp. 3-60 Chapman and Hall, New York, (1994)). This can be summarized by the following equation:



Hence, non-naturally occurring microorganisms possessing the Wood-Ljungdahl pathway can utilize CO₂ and H₂ mixtures as well for the production of acetyl-CoA and other desired products.

The Wood-Ljungdahl pathway is well known in the art and consists of 12 reactions which can be separated into two branches: (1) methyl branch and (2) carbonyl branch. The methyl branch converts syngas to methyl-tetrahydrofolate (methyl-THF) whereas the carbonyl branch converts methyl-THF to acetyl-CoA. The reactions in the methyl branch are catalyzed in order by the following enzymes or proteins: ferredoxin oxidoreductase, formate dehydrogenase, formyltetrahydrofolate synthetase, methenyltetrahydrofolate cyclodehydratase, methylenetetrahydrofolate dehydrogenase and methylenetetrahydrofolate reductase. The reactions in the carbonyl branch are catalyzed in order by the following enzymes or proteins: methyltetrahydrofolate:corrinoid protein methyltransferase (for example, AcsE), corrinoid iron-sulfur protein, nickel-protein assembly protein (for example, AcsF), ferredoxin, acetyl-CoA synthase, carbon monoxide dehydrogenase and nickel-protein assembly protein (for example, CooC). Following the teachings and guidance provided

herein for introducing a sufficient number of encoding nucleic acids to generate a butadiene pathway, those skilled in the art will understand that the same engineering design also can be performed with respect to introducing at least the nucleic acids encoding the Wood-Ljungdahl enzymes or proteins absent in the host organism. Therefore, introduction of one or more encoding nucleic acids into the microbial organisms of the invention such that the modified organism contains the complete Wood-Ljungdahl pathway will confer syngas utilization ability.

Additionally, the reductive (reverse) tricarboxylic acid cycle coupled with carbon monoxide dehydrogenase and/or hydrogenase activities can also be used for the conversion of CO, CO₂ and/or H₂ to acetyl-CoA and other products such as acetate. Organisms capable of fixing carbon via the reductive TCA pathway can utilize one or more of the following enzymes: ATP citrate-lyase, citrate lyase, aconitase, isocitrate dehydrogenase, alpha-ketoglutarate:ferredoxin oxidoreductase, succinyl-CoA synthetase, succinyl-CoA transferase, fumarate reductase, fumarase, malate dehydrogenase, NAD(P)H:ferredoxin oxidoreductase, carbon monoxide dehydrogenase, and hydrogenase. Specifically, the reducing equivalents extracted from CO and/or H₂ by carbon monoxide dehydrogenase and hydrogenase are utilized to fix CO₂ via the reductive TCA cycle into acetyl-CoA or acetate. Acetate can be converted to acetyl-CoA by enzymes such as acetyl-CoA transferase, acetate kinase/phosphotransacetylase, and acetyl-CoA synthetase. Acetyl-CoA can be converted to the p-toluate, terephthalate, or (2-hydroxy-3-methyl-4-oxobutoxy)phosphonate precursors, glyceraldehyde-3-phosphate, phosphoenolpyruvate, and pyruvate, by pyruvate:ferredoxin oxidoreductase and the enzymes of gluconeogenesis. Following the teachings and guidance provided herein for introducing a sufficient number of encoding nucleic acids to generate a p-toluate, terephthalate or (2-hydroxy-3-methyl-4-oxobutoxy)phosphonate pathway, those skilled in the art will understand that the same engineering design also can be performed with respect to introducing at least the nucleic acids encoding the reductive TCA pathway enzymes or proteins absent in the host organism. Therefore, introduction of one or more encoding nucleic acids into the microbial organisms of the invention such that the modified organism contains the complete reductive TCA pathway will confer syngas utilization ability.

Accordingly, given the teachings and guidance provided herein, those skilled in the art will understand that a non-naturally occurring microbial organism can be produced that secretes the biosynthesized compounds of the invention when grown on a carbon source such as a carbohydrate. Such compounds include, for example, butadiene and any of the intermediate

metabolites in the butadiene pathway. All that is required is to engineer in one or more of the required enzyme or protein activities to achieve biosynthesis of the desired compound or intermediate including, for example, inclusion of some or all of the butadiene biosynthetic pathways. Accordingly, the invention provides a non-naturally occurring microbial organism that produces and/or secretes butadiene when grown on a carbohydrate or other carbon source and produces and/or secretes any of the intermediate metabolites shown in the butadiene pathway when grown on a carbohydrate or other carbon source. The butadiene producing microbial organisms of the invention can initiate synthesis from an intermediate, for example, acetoacetyl-CoA, 3-hydroxybutyryl-CoA, crotonyl-CoA, crotonaldehyde, crotyl alcohol, 2-butenyl-phosphate, 2-butenyl-4-diphosphate, erythritol-4-phosphate, 4-(cytidine 5'-diphospho)-erythritol, 2-phospho-4-(cytidine 5'-diphospho)-erythritol, erythritol-2,4-cyclodiphosphate, 1-hydroxy-2-butenyl 4-diphosphate, butenyl 4-diphosphate, 2-butenyl 4-diphosphate, 3-oxoglutaryl-CoA, 3-hydroxyglutaryl-CoA, 3-hydroxy-5-oxopentanoate, 3,5-dihydroxy pentanoate, 3-hydroxy-5-phosphonatooxypentanoate, 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate, crotonate, erythrose, erythritol, 3,5-dioxopentanoate or 5-hydroxy-3-oxopentanoate.

The non-naturally occurring microbial organisms of the invention are constructed using methods well known in the art as exemplified herein to exogenously express at least one nucleic acid encoding a butadiene pathway enzyme or protein in sufficient amounts to produce butadiene. It is understood that the microbial organisms of the invention are cultured under conditions sufficient to produce butadiene. Following the teachings and guidance provided herein, the non-naturally occurring microbial organisms of the invention can achieve biosynthesis of butadiene resulting in intracellular concentrations between about 0.001-2000 mM or more. Generally, the intracellular concentration of butadiene is between about 3-1500 mM, particularly between about 5-1250 mM and more particularly between about 8-1000 mM, including about 10 mM, 100 mM, 200 mM, 500 mM, 800 mM, or more. Intracellular concentrations between and above each of these exemplary ranges also can be achieved from the non-naturally occurring microbial organisms of the invention.

In some embodiments, culture conditions include anaerobic or substantially anaerobic growth or maintenance conditions. Exemplary anaerobic conditions have been described previously and are well known in the art. Exemplary anaerobic conditions for fermentation processes are described herein and are described, for example, in U.S. publication 2009/0047719, filed August 10, 2007. Any of these conditions can be employed with the non-naturally occurring

microbial organisms as well as other anaerobic conditions well known in the art. Under such anaerobic or substantially anaerobic conditions, the butadiene producers can synthesize butadiene at intracellular concentrations of 5-10 mM or more as well as all other concentrations exemplified herein. It is understood that, even though the above description refers to intracellular concentrations, butadiene producing microbial organisms can produce butadiene intracellularly and/or secrete the product into the culture medium.

In addition to the culturing and fermentation conditions disclosed herein, growth condition for achieving biosynthesis of butadiene can include the addition of an osmoprotectant to the culturing conditions. In certain embodiments, the non-naturally occurring microbial organisms of the invention can be sustained, cultured or fermented as described herein in the presence of an osmoprotectant. Briefly, an osmoprotectant refers to a compound that acts as an osmolyte and helps a microbial organism as described herein survive osmotic stress.

Osmoprotectants include, but are not limited to, betaines, amino acids, and the sugar trehalose. Non-limiting examples of such are glycine betaine, praline betaine, dimethylthetin, dimethylsulfoniopropionate, 3-dimethylsulfonio-2-methylpropionate, pipercolic acid, dimethylsulfonioacetate, choline, L-carnitine and ectoine. In one aspect, the osmoprotectant is glycine betaine. It is understood to one of ordinary skill in the art that the amount and type of osmoprotectant suitable for protecting a microbial organism described herein from osmotic stress will depend on the microbial organism used. The amount of osmoprotectant in the culturing conditions can be, for example, no more than about 0.1 mM, no more than about 0.5 mM, no more than about 1.0 mM, no more than about 1.5 mM, no more than about 2.0 mM, no more than about 2.5 mM, no more than about 3.0 mM, no more than about 5.0 mM, no more than about 7.0 mM, no more than about 10 mM, no more than about 50 mM, no more than about 100 mM or no more than about 500 mM.

The culture conditions can include, for example, liquid culture procedures as well as fermentation and other large scale culture procedures. As described herein, particularly useful yields of the biosynthetic products of the invention can be obtained under anaerobic or substantially anaerobic culture conditions.

As described herein, one exemplary growth condition for achieving biosynthesis of butadiene includes anaerobic culture or fermentation conditions. In certain embodiments, the non-naturally occurring microbial organisms of the invention can be sustained, cultured or fermented under anaerobic or substantially anaerobic conditions. Briefly, anaerobic conditions refers to an environment devoid of oxygen. Substantially anaerobic conditions

include, for example, a culture, batch fermentation or continuous fermentation such that the dissolved oxygen concentration in the medium remains between 0 and 10% of saturation. Substantially anaerobic conditions also includes growing or resting cells in liquid medium or on solid agar inside a sealed chamber maintained with an atmosphere of less than 1% oxygen.

- 5 The percent of oxygen can be maintained by, for example, sparging the culture with an N₂/CO₂ mixture or other suitable non-oxygen gas or gases.

- The culture conditions described herein can be scaled up and grown continuously for manufacturing of butadiene. Exemplary growth procedures include, for example, fed-batch fermentation and batch separation; fed-batch fermentation and continuous separation, or
- 10 continuous fermentation and continuous separation. All of these processes are well known in the art. Fermentation procedures are particularly useful for the biosynthetic production of commercial quantities of butadiene. Generally, and as with non-continuous culture procedures, the continuous and/or near-continuous production of butadiene will include culturing a non-naturally occurring butadiene producing organism of the invention in
- 15 sufficient nutrients and medium to sustain and/or nearly sustain growth in an exponential phase. Continuous culture under such conditions can be include, for example, growth for 1 day, 2, 3, 4, 5, 6 or 7 days or more. Additionally, continuous culture can include longer time periods of 1 week, 2, 3, 4 or 5 or more weeks and up to several months. Alternatively, organisms of the invention can be cultured for hours, if suitable for a particular application.
- 20 It is to be understood that the continuous and/or near-continuous culture conditions also can include all time intervals in between these exemplary periods. It is further understood that the time of culturing the microbial organism of the invention is for a sufficient period of time to produce a sufficient amount of product for a desired purpose.

- Fermentation procedures are well known in the art. Briefly, fermentation for the biosynthetic
- 25 production of butadiene can be utilized in, for example, fed-batch fermentation and batch separation; fed-batch fermentation and continuous separation, or continuous fermentation and continuous separation. Examples of batch and continuous fermentation procedures are well known in the art.

- In addition to the above fermentation procedures using the butadiene producers of the
- 30 invention for continuous production of substantial quantities of butadiene, the butadiene producers also can be, for example, simultaneously subjected to chemical synthesis procedures to convert the product to other compounds or the product can be separated from

the fermentation culture and sequentially subjected to chemical or enzymatic conversion to convert the product to other compounds, if desired.

To generate better producers, metabolic modeling can be utilized to optimize growth conditions. Modeling can also be used to design gene knockouts that additionally optimize utilization of the pathway (see, for example, U.S. patent publications US 2002/0012939, US 2003/0224363, US 2004/0029149, US 2004/0072723, US 2003/0059792, US 2002/0168654 and US 2004/0009466, and U.S. Patent No. 7,127,379). Modeling analysis allows reliable predictions of the effects on cell growth of shifting the metabolism towards more efficient production of butadiene.

One computational method for identifying and designing metabolic alterations favoring biosynthesis of a desired product is the OptKnock computational framework (Burgard et al., *Biotechnol. Bioeng.* 84:647-657 (2003)). OptKnock is a metabolic modeling and simulation program that suggests gene deletion or disruption strategies that result in genetically stable microorganisms which overproduce the target product. Specifically, the framework examines the complete metabolic and/or biochemical network of a microorganism in order to suggest genetic manipulations that force the desired biochemical to become an obligatory byproduct of cell growth. By coupling biochemical production with cell growth through strategically placed gene deletions or other functional gene disruption, the growth selection pressures imposed on the engineered strains after long periods of time in a bioreactor lead to improvements in performance as a result of the compulsory growth-coupled biochemical production. Lastly, when gene deletions are constructed there is a negligible possibility of the designed strains reverting to their wild-type states because the genes selected by OptKnock are to be completely removed from the genome. Therefore, this computational methodology can be used to either identify alternative pathways that lead to biosynthesis of a desired product or used in connection with the non-naturally occurring microbial organisms for further optimization of biosynthesis of a desired product.

Briefly, OptKnock is a term used herein to refer to a computational method and system for modeling cellular metabolism. The OptKnock program relates to a framework of models and methods that incorporate particular constraints into flux balance analysis (FBA) models.

These constraints include, for example, qualitative kinetic information, qualitative regulatory information, and/or DNA microarray experimental data. OptKnock also computes solutions to various metabolic problems by, for example, tightening the flux boundaries derived through flux balance models and subsequently probing the performance limits of metabolic

networks in the presence of gene additions or deletions. OptKnock computational framework allows the construction of model formulations that allow an effective query of the performance limits of metabolic networks and provides methods for solving the resulting mixed-integer linear programming problems. The metabolic modeling and simulation methods referred to herein as OptKnock are described in, for example, U.S. publication 2002/0168654, filed January 10, 2002, in International Patent No. PCT/US02/00660, filed January 10, 2002, and U.S. publication 2009/0047719, filed August 10, 2007.

Another computational method for identifying and designing metabolic alterations favoring biosynthetic production of a product is a metabolic modeling and simulation system termed SimPheny®. This computational method and system is described in, for example, U.S. publication 2003/0233218, filed June 14, 2002, and in International Patent Application No. PCT/US03/18838, filed June 13, 2003. SimPheny® is a computational system that can be used to produce a network model *in silico* and to simulate the flux of mass, energy or charge through the chemical reactions of a biological system to define a solution space that contains any and all possible functionalities of the chemical reactions in the system, thereby determining a range of allowed activities for the biological system. This approach is referred to as constraints-based modeling because the solution space is defined by constraints such as the known stoichiometry of the included reactions as well as reaction thermodynamic and capacity constraints associated with maximum fluxes through reactions. The space defined by these constraints can be interrogated to determine the phenotypic capabilities and behavior of the biological system or of its biochemical components.

These computational approaches are consistent with biological realities because biological systems are flexible and can reach the same result in many different ways. Biological systems are designed through evolutionary mechanisms that have been restricted by fundamental constraints that all living systems must face. Therefore, constraints-based modeling strategy embraces these general realities. Further, the ability to continuously impose further restrictions on a network model via the tightening of constraints results in a reduction in the size of the solution space, thereby enhancing the precision with which physiological performance or phenotype can be predicted.

Given the teachings and guidance provided herein, those skilled in the art will be able to apply various computational frameworks for metabolic modeling and simulation to design and implement biosynthesis of a desired compound in host microbial organisms. Such metabolic modeling and simulation methods include, for example, the computational systems

exemplified above as SimPheny® and OptKnock. For illustration of the invention, some methods are described herein with reference to the OptKnock computation framework for modeling and simulation. Those skilled in the art will know how to apply the identification, design and implementation of the metabolic alterations using OptKnock to any of such other
5 metabolic modeling and simulation computational frameworks and methods well known in the art.

The methods described above will provide one set of metabolic reactions to disrupt. Elimination of each reaction within the set or metabolic modification can result in a desired product as an obligatory product during the growth phase of the organism. Because the
10 reactions are known, a solution to the bilevel OptKnock problem also will provide the associated gene or genes encoding one or more enzymes that catalyze each reaction within the set of reactions. Identification of a set of reactions and their corresponding genes encoding the enzymes participating in each reaction is generally an automated process, accomplished through correlation of the reactions with a reaction database having a
15 relationship between enzymes and encoding genes.

Once identified, the set of reactions that are to be disrupted in order to achieve production of a desired product are implemented in the target cell or organism by functional disruption of at least one gene encoding each metabolic reaction within the set. One particularly useful
20 means to achieve functional disruption of the reaction set is by deletion of each encoding gene. However, in some instances, it can be beneficial to disrupt the reaction by other genetic aberrations including, for example, mutation, deletion of regulatory regions such as promoters or cis binding sites for regulatory factors, or by truncation of the coding sequence at any of a number of locations. These latter aberrations, resulting in less than total deletion of the gene set can be useful, for example, when rapid assessments of the coupling of a
25 product are desired or when genetic reversion is less likely to occur.

To identify additional productive solutions to the above described bilevel OptKnock problem which lead to further sets of reactions to disrupt or metabolic modifications that can result in the biosynthesis, including growth-coupled biosynthesis of a desired product, an optimization method, termed integer cuts, can be implemented. This method proceeds by iteratively
30 solving the OptKnock problem exemplified above with the incorporation of an additional constraint referred to as an integer cut at each iteration. Integer cut constraints effectively prevent the solution procedure from choosing the exact same set of reactions identified in any previous iteration that obligatorily couples product biosynthesis to growth. For example, if a

previously identified growth-coupled metabolic modification specifies reactions 1, 2, and 3 for disruption, then the following constraint prevents the same reactions from being simultaneously considered in subsequent solutions. The integer cut method is well known in the art and can be found described in, for example, Burgard et al., *Biotechnol. Prog.* 17:791-797 (2001). As with all methods described herein with reference to their use in combination with the OptKnock computational framework for metabolic modeling and simulation, the integer cut method of reducing redundancy in iterative computational analysis also can be applied with other computational frameworks well known in the art including, for example, SimPheny®.

The methods exemplified herein allow the construction of cells and organisms that biosynthetically produce a desired product, including the obligatory coupling of production of a target biochemical product to growth of the cell or organism engineered to harbor the identified genetic alterations. Therefore, the computational methods described herein allow the identification and implementation of metabolic modifications that are identified by an *in silico* method selected from OptKnock or SimPheny®. The set of metabolic modifications can include, for example, addition of one or more biosynthetic pathway enzymes and/or functional disruption of one or more metabolic reactions including, for example, disruption by gene deletion.

As discussed above, the OptKnock methodology was developed on the premise that mutant microbial networks can be evolved towards their computationally predicted maximum-growth phenotypes when subjected to long periods of growth selection. In other words, the approach leverages an organism's ability to self-optimize under selective pressures. The OptKnock framework allows for the exhaustive enumeration of gene deletion combinations that force a coupling between biochemical production and cell growth based on network stoichiometry. The identification of optimal gene/reaction knockouts requires the solution of a bilevel optimization problem that chooses the set of active reactions such that an optimal growth solution for the resulting network overproduces the biochemical of interest (Burgard et al., *Biotechnol. Bioeng.* 84:647-657 (2003)).

An *in silico* stoichiometric model of *E. coli* metabolism can be employed to identify essential genes for metabolic pathways as exemplified previously and described in, for example, U.S. patent publications US 2002/0012939, US 2003/0224363, US 2004/0029149, US 2004/0072723, US 2003/0059792, US 2002/0168654 and US 2004/0009466, and in U.S. Patent No. 7,127,379. As disclosed herein, the OptKnock mathematical framework can be

applied to pinpoint gene deletions leading to the growth-coupled production of a desired product. Further, the solution of the bilevel OptKnock problem provides only one set of deletions. To enumerate all meaningful solutions, that is, all sets of knockouts leading to growth-coupled production formation, an optimization technique, termed integer cuts, can be implemented. This entails iteratively solving the OptKnock problem with the incorporation of an additional constraint referred to as an integer cut at each iteration, as discussed above.

As disclosed herein, a nucleic acid encoding a desired activity of a butadiene pathway can be introduced into a host organism. In some cases, it can be desirable to modify an activity of a butadiene pathway enzyme or protein to increase production of butadiene. For example, known mutations that increase the activity of a protein or enzyme can be introduced into an encoding nucleic acid molecule. Additionally, optimization methods can be applied to increase the activity of an enzyme or protein and/or decrease an inhibitory activity, for example, decrease the activity of a negative regulator.

One such optimization method is directed evolution. Directed evolution is a powerful approach that involves the introduction of mutations targeted to a specific gene in order to improve and/or alter the properties of an enzyme. Improved and/or altered enzymes can be identified through the development and implementation of sensitive high-throughput screening assays that allow the automated screening of many enzyme variants (for example, $>10^4$). Iterative rounds of mutagenesis and screening typically are performed to afford an enzyme with optimized properties. Computational algorithms that can help to identify areas of the gene for mutagenesis also have been developed and can significantly reduce the number of enzyme variants that need to be generated and screened. Numerous directed evolution technologies have been developed (for reviews, see Hibbert et al., *Biomol.Eng* 22:11-19 (2005); Huisman and Lalonde, *In Biocatalysis in the pharmaceutical and biotechnology industries* pgs. 717-742 (2007), Patel (ed.), CRC Press; Otten and Quax. *Biomol.Eng* 22:1-9 (2005).; and Sen et al., *Appl Biochem.Biotechnol* 143:212-223 (2007)) to be effective at creating diverse variant libraries, and these methods have been successfully applied to the improvement of a wide range of properties across many enzyme classes. Enzyme characteristics that have been improved and/or altered by directed evolution technologies include, for example: selectivity/specificity, for conversion of non-natural substrates; temperature stability, for robust high temperature processing; pH stability, for bioprocessing under lower or higher pH conditions; substrate or product tolerance, so that high product titers can be achieved; binding (K_m), including broadening substrate binding to

include non-natural substrates; inhibition (K_i), to remove inhibition by products, substrates, or key intermediates; activity (kcat), to increase enzymatic reaction rates to achieve desired flux; expression levels, to increase protein yields and overall pathway flux; oxygen stability, for operation of air sensitive enzymes under aerobic conditions; and anaerobic activity, for operation of an aerobic enzyme in the absence of oxygen.

Described below in more detail are exemplary methods that have been developed for the mutagenesis and diversification of genes to target desired properties of specific enzymes. Such methods are well known to those skilled in the art. Any of these can be used to alter and/or optimize the activity of a butadiene pathway enzyme or protein.

EpPCR (Pritchard et al., *J Theor. Biol.* 234:497-509 (2005)) introduces random point mutations by reducing the fidelity of DNA polymerase in PCR reactions by the addition of Mn^{2+} ions, by biasing dNTP concentrations, or by other conditional variations. The five step cloning process to confine the mutagenesis to the target gene of interest involves: 1) error-prone PCR amplification of the gene of interest; 2) restriction enzyme digestion; 3) gel purification of the desired DNA fragment; 4) ligation into a vector; 5) transformation of the gene variants into a suitable host and screening of the library for improved performance. This method can generate multiple mutations in a single gene simultaneously, which can be useful to screen a larger number of potential variants having a desired activity. A high number of mutants can be generated by EpPCR, so a high-throughput screening assay or a selection method, for example, using robotics, is useful to identify those with desirable characteristics.

Error-prone Rolling Circle Amplification (epRCA) (Fujii et al., *Nucleic Acids Res.* 32:e145 (2004); and Fujii et al., *Nat. Protoc.* 1:2493-2497 (2006)) has many of the same elements as epPCR except a whole circular plasmid is used as the template and random 6-mers with exonuclease resistant thiophosphate linkages on the last 2 nucleotides are used to amplify the plasmid followed by transformation into cells in which the plasmid is re-circularized at tandem repeats. Adjusting the Mn^{2+} concentration can vary the mutation rate somewhat. This technique uses a simple error-prone, single-step method to create a full copy of the plasmid with 3 - 4 mutations/kbp. No restriction enzyme digestion or specific primers are required. Additionally, this method is typically available as a commercially available kit.

DNA or Family Shuffling (Stemmer, *Proc Natl Acad Sci USA* 91:10747-10751 (1994)); and Stemmer, *Nature* 370:389-391 (1994)) typically involves digestion of two or more variant

genes with nucleases such as Dnase I or EndoV to generate a pool of random fragments that are reassembled by cycles of annealing and extension in the presence of DNA polymerase to create a library of chimeric genes. Fragments prime each other and recombination occurs when one copy primes another copy (template switch). This method can be used with >1kbp DNA sequences. In addition to mutational recombinants created by fragment reassembly, this method introduces point mutations in the extension steps at a rate similar to error-prone PCR. The method can be used to remove deleterious, random and neutral mutations.

Staggered Extension (StEP) (Zhao et al., *Nat. Biotechnol.* 16:258-261 (1998)) entails template priming followed by repeated cycles of 2 step PCR with denaturation and very short duration of annealing/extension (as short as 5 sec). Growing fragments anneal to different templates and extend further, which is repeated until full-length sequences are made. Template switching means most resulting fragments have multiple parents. Combinations of low-fidelity polymerases (Taq and Mutazyme) reduce error-prone biases because of opposite mutational spectra.

In Random Priming Recombination (RPR) random sequence primers are used to generate many short DNA fragments complementary to different segments of the template (Shao et al., *Nucleic Acids Res* 26:681-683 (1998)). Base misincorporation and mispriming via epPCR give point mutations. Short DNA fragments prime one another based on homology and are recombined and reassembled into full-length by repeated thermocycling. Removal of templates prior to this step assures low parental recombinants. This method, like most others, can be performed over multiple iterations to evolve distinct properties. This technology avoids sequence bias, is independent of gene length, and requires very little parent DNA for the application.

In Heteroduplex Recombination linearized plasmid DNA is used to form heteroduplexes that are repaired by mismatch repair (Volkov et al, *Nucleic Acids Res.* 27:e18 (1999); and Volkov et al., *Methods Enzymol.* 328:456-463 (2000)). The mismatch repair step is at least somewhat mutagenic. Heteroduplexes transform more efficiently than linear homoduplexes. This method is suitable for large genes and whole operons.

Random Chimeragenesis on Transient Templates (RACHITT) (Coco et al., *Nat. Biotechnol.* 19:354-359 (2001)) employs Dnase I fragmentation and size fractionation of single stranded DNA (ssDNA). Homologous fragments are hybridized in the absence of polymerase to a complementary ssDNA scaffold. Any overlapping unhybridized fragment ends are trimmed

down by an exonuclease. Gaps between fragments are filled in and then ligated to give a pool of full-length diverse strands hybridized to the scaffold, which contains U to preclude amplification. The scaffold then is destroyed and is replaced by a new strand complementary to the diverse strand by PCR amplification. The method involves one strand (scaffold) that is
5 from only one parent while the priming fragments derive from other genes; the parent scaffold is selected against. Thus, no reannealing with parental fragments occurs.

Overlapping fragments are trimmed with an exonuclease. Otherwise, this is conceptually similar to DNA shuffling and StEP. Therefore, there should be no siblings, few inactives, and no unshuffled parentals. This technique has advantages in that few or no parental genes
10 are created and many more crossovers can result relative to standard DNA shuffling.

Recombined Extension on Truncated templates (RETT) entails template switching of unidirectionally growing strands from primers in the presence of unidirectional ssDNA fragments used as a pool of templates (Lee et al., *J. Molec. Catalysis* 26:119-129 (2003)). No DNA endonucleases are used. Unidirectional ssDNA is made by DNA polymerase with
15 random primers or serial deletion with exonuclease. Unidirectional ssDNA are only templates and not primers. Random priming and exonucleases do not introduce sequence bias as true of enzymatic cleavage of DNA shuffling/RACHITT. RETT can be easier to optimize than StEP because it uses normal PCR conditions instead of very short extensions. Recombination occurs as a component of the PCR steps, that is, no direct shuffling. This
20 method can also be more random than StEP due to the absence of pauses.

In Degenerate Oligonucleotide Gene Shuffling (DOGS) degenerate primers are used to control recombination between molecules; (Bergquist and Gibbs, *Methods Mol.Biol* 352:191-204 (2007); Bergquist et al., *Biomol.Eng* 22:63-72 (2005); Gibbs et al., *Gene* 271:13-20 (2001)) this can be used to control the tendency of other methods such as DNA shuffling to
25 regenerate parental genes. This method can be combined with random mutagenesis (epPCR) of selected gene segments. This can be a good method to block the reformation of parental sequences. No endonucleases are needed. By adjusting input concentrations of segments made, one can bias towards a desired backbone. This method allows DNA shuffling from unrelated parents without restriction enzyme digests and allows a choice of random
30 mutagenesis methods.

Incremental Truncation for the Creation of Hybrid Enzymes (ITCHY) creates a combinatorial library with 1 base pair deletions of a gene or gene fragment of interest (Ostermeier et al., *Proc. Natl. Acad. Sci. USA* 96:3562-3567 (1999); and Ostermeier et al., *Nat. Biotechnol.*

17:1205-1209 (1999)). Truncations are introduced in opposite direction on pieces of 2 different genes. These are ligated together and the fusions are cloned. This technique does not require homology between the 2 parental genes. When ITCHY is combined with DNA shuffling, the system is called SCRATCHY (see below). A major advantage of both is no need for homology between parental genes; for example, functional fusions between an E. coli and a human gene were created via ITCHY. When ITCHY libraries are made, all possible crossovers are captured.

Thio-Incremental Truncation for the Creation of Hybrid Enzymes (THIO-ITCHY) is similar to ITCHY except that phosphothioate dNTPs are used to generate truncations (Lutz et al., Nucleic Acids Res 29:E16 (2001)). Relative to ITCHY, THIO-ITCHY can be easier to optimize, provide more reproducibility, and adjustability.

SCRATCHY combines two methods for recombining genes, ITCHY and DNA shuffling (Lutz et al., Proc. Natl. Acad. Sci. USA 98:11248-11253 (2001)). SCRATCHY combines the best features of ITCHY and DNA shuffling. First, ITCHY is used to create a comprehensive set of fusions between fragments of genes in a DNA homology-independent fashion. This artificial family is then subjected to a DNA-shuffling step to augment the number of crossovers. Computational predictions can be used in optimization. SCRATCHY is more effective than DNA shuffling when sequence identity is below 80%.

In Random Drift Mutagenesis (RNDM) mutations made via epPCR followed by screening/selection for those retaining usable activity (Bergquist et al., Biomol. Eng. 22:63-72 (2005)). Then, these are used in DOGS to generate recombinants with fusions between multiple active mutants or between active mutants and some other desirable parent. Designed to promote isolation of neutral mutations; its purpose is to screen for retained catalytic activity whether or not this activity is higher or lower than in the original gene. RNDM is usable in high throughput assays when screening is capable of detecting activity above background. RNDM has been used as a front end to DOGS in generating diversity. The technique imposes a requirement for activity prior to shuffling or other subsequent steps; neutral drift libraries are indicated to result in higher/quicker improvements in activity from smaller libraries. Though published using epPCR, this could be applied to other large-scale mutagenesis methods.

Sequence Saturation Mutagenesis (SeSaM) is a random mutagenesis method that: 1) generates a pool of random length fragments using random incorporation of a phosphothioate

nucleotide and cleavage; this pool is used as a template to 2) extend in the presence of “universal” bases such as inosine; 3) replication of an inosine-containing complement gives random base incorporation and, consequently, mutagenesis (Wong et al., *Biotechnol. J.* 3:74-82 (2008); Wong et al., *Nucleic Acids Res.* 32:e26 (2004); and Wong et al., *Anal. Biochem.* 341:187-189 (2005)). Using this technique it can be possible to generate a large library of mutants within 2 to 3 days using simple methods. This technique is non-directed in comparison to the mutational bias of DNA polymerases. Differences in this approach makes this technique complementary (or an alternative) to epPCR.

In Synthetic Shuffling, overlapping oligonucleotides are designed to encode “all genetic diversity in targets” and allow a very high diversity for the shuffled progeny (Ness et al., *Nat. Biotechnol.* 20:1251-1255 (2002)). In this technique, one can design the fragments to be shuffled. This aids in increasing the resulting diversity of the progeny. One can design sequence/codon biases to make more distantly related sequences recombine at rates approaching those observed with more closely related sequences. Additionally, the technique does not require physically possessing the template genes.

Nucleotide Exchange and Excision Technology NexT exploits a combination of dUTP incorporation followed by treatment with uracil DNA glycosylase and then piperidine to perform endpoint DNA fragmentation (Muller et al., *Nucleic Acids Res.* 33:e117 (2005)). The gene is reassembled using internal PCR primer extension with proofreading polymerase. The sizes for shuffling are directly controllable using varying dUPT::dTTP ratios. This is an end point reaction using simple methods for uracil incorporation and cleavage. Other nucleotide analogs, such as 8-oxo-guanine, can be used with this method. Additionally, the technique works well with very short fragments (86 bp) and has a low error rate. The chemical cleavage of DNA used in this technique results in very few unshuffled clones.

In Sequence Homology-Independent Protein Recombination (SHIPREC), a linker is used to facilitate fusion between two distantly related or unrelated genes. Nuclease treatment is used to generate a range of chimeras between the two genes. These fusions result in libraries of single-crossover hybrids (Sieber et al., *Nat. Biotechnol.* 19:456-460 (2001)). This produces a limited type of shuffling and a separate process is required for mutagenesis. In addition, since no homology is needed, this technique can create a library of chimeras with varying fractions of each of the two unrelated parent genes. SHIPREC was tested with a heme-binding domain of a bacterial CP450 fused to N-terminal regions of a mammalian CP450; this produced mammalian activity in a more soluble enzyme.

In Gene Site Saturation Mutagenesis™ (GSSM™) the starting materials are a supercoiled dsDNA plasmid containing an insert and two primers which are degenerate at the desired site of mutations (Kretz et al., *Methods Enzymol.* 388:3-11 (2004)). Primers carrying the mutation of interest, anneal to the same sequence on opposite strands of DNA. The mutation is typically in the middle of the primer and flanked on each side by approximately 20 nucleotides of correct sequence. The sequence in the primer is NNN or NNK (coding) and MNN (noncoding) (N = all 4, K = G, T, M = A, C). After extension, DpnI is used to digest dam-methylated DNA to eliminate the wild-type template. This technique explores all possible amino acid substitutions at a given locus (that is, one codon). The technique facilitates the generation of all possible replacements at a single-site with no nonsense codons and results in equal to near-equal representation of most possible alleles. This technique does not require prior knowledge of the structure, mechanism, or domains of the target enzyme. If followed by shuffling or Gene Reassembly, this technology creates a diverse library of recombinants containing all possible combinations of single-site up-mutations. The usefulness of this technology combination has been demonstrated for the successful evolution of over 50 different enzymes, and also for more than one property in a given enzyme.

Combinatorial Cassette Mutagenesis (CCM) involves the use of short oligonucleotide cassettes to replace limited regions with a large number of possible amino acid sequence alterations (Reidhaar-Olson et al. *Methods Enzymol.* 208:564-586 (1991); and Reidhaar-Olson et al. *Science* 241:53-57 (1988)). Simultaneous substitutions at two or three sites are possible using this technique. Additionally, the method tests a large multiplicity of possible sequence changes at a limited range of sites. This technique has been used to explore the information content of the lambda repressor DNA-binding domain.

Combinatorial Multiple Cassette Mutagenesis (CMCM) is essentially similar to CCM except it is employed as part of a larger program: 1) use of epPCR at high mutation rate to 2) identify hot spots and hot regions and then 3) extension by CMCM to cover a defined region of protein sequence space (Reetz et al., *Angew. Chem. Int. Ed Engl.* 40:3589-3591 (2001)). As with CCM, this method can test virtually all possible alterations over a target region. If used along with methods to create random mutations and shuffled genes, it provides an excellent means of generating diverse, shuffled proteins. This approach was successful in increasing, by 51-fold, the enantioselectivity of an enzyme.

In the Mutator Strains technique, conditional ts mutator plasmids allow increases of 20 to 4000-X in random and natural mutation frequency during selection and block accumulation

of deleterious mutations when selection is not required (Selifonova et al., Appl. Environ. Microbiol. 67:3645-3649 (2001)). This technology is based on a plasmid-derived mutD5 gene, which encodes a mutant subunit of DNA polymerase III. This subunit binds to endogenous DNA polymerase III and compromises the proofreading ability of polymerase III in any strain that harbors the plasmid. A broad-spectrum of base substitutions and frameshift mutations occur. In order for effective use, the mutator plasmid should be removed once the desired phenotype is achieved; this is accomplished through a temperature sensitive (ts) origin of replication, which allows for plasmid curing at 41°C. It should be noted that mutator strains have been explored for quite some time (see Low et al., J. Mol. Biol. 260:359-3680 (1996)). In this technique, very high spontaneous mutation rates are observed. The conditional property minimizes non-desired background mutations. This technology could be combined with adaptive evolution to enhance mutagenesis rates and more rapidly achieve desired phenotypes.

Look-Through Mutagenesis (LTM) is a multidimensional mutagenesis method that assesses and optimizes combinatorial mutations of selected amino acids (Rajpal et al., Proc. Natl. Acad. Sci. USA 102:8466-8471 (2005)). Rather than saturating each site with all possible amino acid changes, a set of nine is chosen to cover the range of amino acid R-group chemistry. Fewer changes per site allows multiple sites to be subjected to this type of mutagenesis. A >800-fold increase in binding affinity for an antibody from low nanomolar to picomolar has been achieved through this method. This is a rational approach to minimize the number of random combinations and can increase the ability to find improved traits by greatly decreasing the numbers of clones to be screened. This has been applied to antibody engineering, specifically to increase the binding affinity and/or reduce dissociation. The technique can be combined with either screens or selections.

Gene Reassembly is a DNA shuffling method that can be applied to multiple genes at one time or to create a large library of chimeras (multiple mutations) of a single gene (Tunable GeneReassembly™ (TGR™) Technology supplied by Verenum Corporation). Typically this technology is used in combination with ultra-high-throughput screening to query the represented sequence space for desired improvements. This technique allows multiple gene recombination independent of homology. The exact number and position of cross-over events can be pre-determined using fragments designed via bioinformatic analysis. This technology leads to a very high level of diversity with virtually no parental gene reformation and a low level of inactive genes. Combined with GSSM™, a large range of mutations can

be tested for improved activity. The method allows “blending” and “fine tuning” of DNA shuffling, for example, codon usage can be optimized.

In Silico Protein Design Automation (PDA) is an optimization algorithm that anchors the structurally defined protein backbone possessing a particular fold, and searches sequence space for amino acid substitutions that can stabilize the fold and overall protein energetics (Hayes et al., Proc. Natl. Acad. Sci. USA 99:15926-15931 (2002)). This technology uses in silico structure-based entropy predictions in order to search for structural tolerance toward protein amino acid variations. Statistical mechanics is applied to calculate coupling interactions at each position. Structural tolerance toward amino acid substitution is a measure of coupling. Ultimately, this technology is designed to yield desired modifications of protein properties while maintaining the integrity of structural characteristics. The method computationally assesses and allows filtering of a very large number of possible sequence variants (1050). The choice of sequence variants to test is related to predictions based on the most favorable thermodynamics. Ostensibly only stability or properties that are linked to stability can be effectively addressed with this technology. The method has been successfully used in some therapeutic proteins, especially in engineering immunoglobulins. In silico predictions avoid testing extraordinarily large numbers of potential variants. Predictions based on existing three-dimensional structures are more likely to succeed than predictions based on hypothetical structures. This technology can readily predict and allow targeted screening of multiple simultaneous mutations, something not possible with purely experimental technologies due to exponential increases in numbers.

Iterative Saturation Mutagenesis (ISM) involves: 1) using knowledge of structure/function to choose a likely site for enzyme improvement; 2) performing saturation mutagenesis at chosen site using a mutagenesis method such as Stratagene QuikChange (Stratagene; San Diego CA); 3) screening/selecting for desired properties; and 4) using improved clone(s), start over at another site and continue repeating until a desired activity is achieved (Reetz et al., Nat. Protoc. 2:891-903 (2007); and Reetz et al., Angew. Chem. Int. Ed Engl. 45:7745-7751 (2006)). This is a proven methodology, which assures all possible replacements at a given position are made for screening/selection.

Any of the aforementioned methods for mutagenesis can be used alone or in any combination. Additionally, any one or combination of the directed evolution methods can be used in conjunction with adaptive evolution techniques, as described herein.

It is understood that modifications which do not substantially affect the activity of the various embodiments of this invention are also provided within the definition of the invention provided herein. Accordingly, the following examples are intended to illustrate but not limit the present invention.

EXAMPLE I

Pathways for Producing Butadiene

Disclosed herein are novel processes for the direct production of butadiene using engineered non-natural microorganisms that possess the enzymes necessary for conversion of common metabolites into the four carbon diene, 1,3-butadiene. One novel route to direct production of butadiene entails reduction of the known butanol pathway metabolite crotonyl-CoA to crotyl alcohol via reduction with aldehyde and alcohol dehydrogenases, followed by phosphorylation with kinases to afford crotyl pyrophosphate and subsequent conversion to butadiene using isoprene synthases or variants thereof (see Figure 2). Another route (Figure 3) is a variant of the well-characterized DXP pathway for isoprenoid biosynthesis. In this route, the substrate lacks a 2-methyl group and provides butadiene rather than isoprene via a butadiene synthase. Such a butadiene synthase can be derived from a isoprene synthase using methods, such as directed evolution, as described herein. Finally, Figure 4 shows a pathway to butadiene involving the substrate 3-hydroxyglutaryl-CoA, which serves as a surrogate for the natural mevalonate pathway substrate 3-hydroxy-3-methyl-glutaryl-CoA (shown in Figure 1). Enzyme candidates for steps A-P of Figure 2, steps A-K of Figure 3 and steps A-O of Figure 4 are provided below.

Acetyl-CoA:acetyl-CoA acyltransferase (Figure 2, Step A)

Acetoacetyl-CoA thiolase converts two molecules of acetyl-CoA into one molecule each of acetoacetyl-CoA and CoA. Exemplary acetoacetyl-CoA thiolase enzymes include the gene products of *atoB* from *E. coli* (Martin et al., *Nat. Biotechnol* 21:796-802 (2003)), *thlA* and *thlB* from *C. acetobutylicum* (Hanai et al., *Appl Environ Microbiol* 73:7814-7818 (2007); Winzer et al., *J.Mol.Microbiol Biotechnol* 2:531-541 (2000)), and *ERG10* from *S. cerevisiae* (Hiser et al., *J.Biol.Chem.* 269:31383-31389 (1994)).

<u>Protein</u>	<u>GenBank ID</u>	<u>GI number</u>	<u>Organism</u>
<i>AtoB</i>	NP_416728	16130161	<i>Escherichia coli</i>
<i>ThlA</i>	NP_349476.1	15896127	<i>Clostridium acetobutylicum</i>
<i>ThlB</i>	NP_149242.1	15004782	<i>Clostridium acetobutylicum</i>
<i>ERG10</i>	NP_015297	6325229	<i>Saccharomyces cerevisiae</i>

Acetoacetyl-CoA reductase (Figure 2, Step B)

Acetoacetyl-CoA reductase catalyzing the reduction of acetoacetyl-CoA to 3-hydroxybutyryl-CoA participates in the acetyl-CoA fermentation pathway to butyrate in several species of *Clostridia* and has been studied in detail (Jones et al., *Microbiol Rev.* 50:484-524 (1986)).

5 The enzyme from *Clostridium acetobutylicum*, encoded by *hbd*, has been cloned and functionally expressed in *E. coli* (Youngleson et al., *J Bacteriol.* 171:6800-6807 (1989)).

Additionally, subunits of two fatty acid oxidation complexes in *E. coli*, encoded by *fadB* and *fadJ*, function as 3-hydroxyacyl-CoA dehydrogenases (Binstock et al., *Methods Enzymol.* 71

10 Pt C:403-411 (1981)). Yet other gene candidates demonstrated to reduce acetoacetyl-CoA to 3-hydroxybutyryl-CoA are *phbB* from *Zoogloea ramigera* (Ploux et al., *Eur.J Biochem.* 174:177-182 (1988)) and *phaB* from *Rhodobacter sphaeroides* (Alber et al., *Mol.Microbiol* 61:297-309 (2006)). The former gene candidate is NADPH-dependent, its nucleotide

sequence has been determined (Peoples et al., *Mol.Microbiol* 3:349-357 (1989)) and the gene

15 has been expressed in *E. coli*. Substrate specificity studies on the gene led to the conclusion that it could accept 3-oxopropionyl-CoA as a substrate besides acetoacetyl-CoA (Ploux et al., *supra*, (1988)). Additional gene candidates include *Hbd1* (C-terminal domain) and *Hbd2* (N-terminal domain) in *Clostridium kluyveri* (Hillmer and Gottschalk, *Biochim. Biophys. Acta* 3334:12-23 (1974)) and *HSD17B10* in *Bos taurus* (WAKIL et al., *J Biol.Chem.* 207:631-638

20 (1954)).

<u>Protein</u>	<u>Genbank ID</u>	<u>GI number</u>	<u>Organism</u>
<i>fadB</i>	P21177.2	119811	<i>Escherichia coli</i>
<i>fadJ</i>	P77399.1	3334437	<i>Escherichia coli</i>
<i>Hbd2</i>	EDK34807.1	146348271	<i>Clostridium kluyveri</i>
<i>Hbd1</i>	EDK32512.1	146345976	<i>Clostridium kluyveri</i>
<i>hbd</i>	P52041.2	18266893	<i>Clostridium acetobutylicum</i>
<i>HSD17B10</i>	O02691.3	3183024	<i>Bos Taurus</i>
<i>phbB</i>	P23238.1	130017	<i>Zoogloea ramigera</i>
<i>phaB</i>	YP_353825.1	77464321	<i>Rhodobacter sphaeroides</i>

A number of similar enzymes have been found in other species of *Clostridia* and in *Metallosphaera sedula* (Berg et al., *Science*. 318:1782-1786 (2007)).

<u>Protein</u>	<u>GenBank ID</u>	<u>GI number</u>	<u>Organism</u>
<i>hbd</i>	NP_349314.1	NP_349314.1	<i>Clostridium acetobutylicum</i>
<i>hbd</i>	AAM14586.1	AAM14586.1	<i>Clostridium beijerinckii</i>
<i>Msed_1423</i>	YP_001191505	YP_001191505	<i>Metallosphaera sedula</i>
<i>Msed_0399</i>	YP_001190500	YP_001190500	<i>Metallosphaera sedula</i>
<i>Msed_0389</i>	YP_001190490	YP_001190490	<i>Metallosphaera sedula</i>
<i>Msed_1993</i>	YP_001192057	YP_001192057	<i>Metallosphaera sedula</i>

5 **3-Hydroxybutyryl-CoA dehydratase (Figure 2, Step C)**

3-Hydroxybutyryl-CoA dehydratase (EC 4.2.1.55), also called crotonase, is an enoyl-CoA hydratase that reversibly dehydrates 3-hydroxybutyryl-CoA to form crotonyl-CoA.

Crotonase enzymes are required for n-butanol formation in some organisms, particularly *Clostridial* species, and also comprise one step of the 3-hydroxypropionate/4-

- 10 hydroxybutyrate cycle in thermoacidophilic Archaea of the genera *Sulfolobus*, *Acidianus*, and *Metallosphaera*. Exemplary genes encoding crotonase enzymes can be found in *C. acetobutylicum* (Atsumi et al., *Metab Eng.* 10:305-311 (2008); Boynton et al., *J Bacteriol.* 178:3015-3024 (1996)), *C. kluyveri* (Hillmer et al., *FEBS Lett.* 21:351-354 (1972)), and *Metallosphaera sedula* (Berg et al., *Science* 318:1782-1786 (2007a)) though the sequence of
- 15 the latter gene is not known. The enoyl-CoA hydratase of *Pseudomonas putida*, encoded by

ech, catalyzes the conversion of crotonyl-CoA to 3-hydroxybutyryl-CoA (Roberts et al., *Arch Microbiol.* 117:99-108 (1978)). Additional enoyl-CoA hydratase candidates are *phaA* and *phaB*, of *P. putida*, and *paaA* and *paaB* from *P. fluorescens* (Olivera et al., *Proc.Natl.Acad.Sci U.S.A* 95:6419-6424 (1998)). Lastly, a number of *Escherichia coli* genes have been shown to demonstrate enoyl-CoA hydratase functionality including *maoC* (Park et al., *J Bacteriol.* 185:5391-5397 (2003)), *paaF* (Ismail et al., *Eur.J Biochem.* 270:3047-3054 (2003); Park et al., *Appl.Biochem.Biotechnol* 113-116:335-346 (2004); Park et al., *Biotechnol Bioeng* 86:681-686 (2004)) and *paaG* (Ismail et al., *supra*, (2003); Park and Lee, *supra*, (2004); Park and Yup, *supra*, (2004)). These proteins are identified below.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>crt</i>	NP_349318.1	15895969	<i>Clostridium acetobutylicum</i>
<i>crtI</i>	YP_001393856.1	153953091	<i>Clostridium kluyveri</i>
<i>ech</i>	NP_745498.1	26990073	<i>Pseudomonas putida</i>
<i>paaA</i>	NP_745427.1	26990002	<i>Pseudomonas putida</i>
<i>paaB</i>	NP_745426.1	26990001	<i>Pseudomonas putida</i>
<i>phaA</i>	ABF82233.1	106636093	<i>Pseudomonas fluorescens</i>
<i>phaB</i>	ABF82234.1	106636094	<i>Pseudomonas fluorescens</i>
<i>maoC</i>	NP_415905.1	16129348	<i>Escherichia coli</i>
<i>paaF</i>	NP_415911.1	16129354	<i>Escherichia coli</i>
<i>paaG</i>	NP_415912.1	16129355	<i>Escherichia coli</i>

Crotonyl-CoA reductase (aldehyde forming) (Figure 2, Step D)

Several acyl-CoA dehydrogenases are capable of reducing an acyl-CoA to its corresponding aldehyde. Thus they can naturally reduce crotonyl-CoA to crotonaldehyde or can be engineered to do so. Exemplary genes that encode such enzymes include the *Acinetobacter calcoaceticus acrI* encoding a fatty acyl-CoA reductase (Reiser et al., *J. Bacteriol.* 179:2969-2975 (1997)), the *Acinetobacter sp. M-1* fatty acyl-CoA reductase (Ishige et al., *Appl.Environ.Microbiol.* 68:1192-1195 (2002)), and a CoA- and NADP- dependent succinate semialdehyde dehydrogenase encoded by the *sucD* gene in *Clostridium kluyveri* (Sohling et al., *J Bacteriol.* 178:871-880 (1996); Sohling et al., *J. Bacteriol.* 178:871-80 (1996)). *SucD* of *P. gingivalis* is another succinate semialdehyde dehydrogenase (Takahashi et al., *J.Bacteriol.* 182:4704-4710 (2000)). These succinate semialdehyde dehydrogenases were specifically shown in ref. (Burk et al., WO/2008/115840: (2008)) to convert 4-

hydroxybutyryl-CoA to 4-hydroxybutanal as part of a pathway to produce 1,4-butanediol. The enzyme acylating acetaldehyde dehydrogenase in *Pseudomonas sp*, encoded by *bphG*, is yet another capable enzyme as it has been demonstrated to oxidize and acylate acetaldehyde, propionaldehyde, butyraldehyde, isobutyraldehyde and formaldehyde (Powlowski et al., *J. Bacteriol.* 175:377-385 (1993)).

Protein	GenBank ID	GI Number	Organism
<i>acrI</i>	YP_047869.1	50086359	<i>Acinetobacter calcoaceticus</i>
<i>acrI</i>	AAC45217	1684886	<i>Acinetobacter baylyi</i>
<i>acrI</i>	BAB85476.1	18857901	<i>Acinetobacter sp.</i> Strain M-1
<i>sucD</i>	P38947.1	172046062	<i>Clostridium kluyveri</i>
<i>sucD</i>	NP_904963.1	34540484	<i>Porphyromonas gingivalis</i>
<i>bphG</i>	BAA03892.1	425213	<i>Pseudomonas sp</i>

An additional enzyme type that converts an acyl-CoA to its corresponding aldehyde is malonyl-CoA reductase which transforms malonyl-CoA to malonic semialdehyde. Malonyl-CoA reductase is a key enzyme in autotrophic carbon fixation via the 3-hydroxypropionate cycle in thermoacidophilic archaeal bacteria (Berg et al., *Science* 318:1782-1786 (2007b); Thauer, 318:1732-1733 (2007)). The enzyme utilizes NADPH as a cofactor and has been characterized in *Metallosphaera* and *Sulfolobus spp* (Alber et al., *J. Bacteriol.* 188:8551-8559 (2006); Hugler et al., *J. Bacteriol.* 184:2404-2410 (2002)). The enzyme is encoded by *Msed_0709* in *Metallosphaera sedula* (Alber et al., *supra*, (2006); Berg et al., *supra*, (2007b)). A gene encoding a malonyl-CoA reductase from *Sulfolobus tokodaii* was cloned and heterologously expressed in *E. coli* (Alber et al., *supra*, (2006)). Although the aldehyde dehydrogenase functionality of these enzymes is similar to the bifunctional dehydrogenase from *Chloroflexus aurantiacus*, there is little sequence similarity. Both malonyl-CoA reductase enzyme candidates have high sequence similarity to aspartate-semialdehyde dehydrogenase, an enzyme catalyzing the reduction and concurrent dephosphorylation of aspartyl-4-phosphate to aspartate semialdehyde. Additional gene candidates can be found by sequence homology to proteins in other organisms including *Sulfolobus solfataricus* and *Sulfolobus acidocaldarius*. Yet another candidate for CoA-acylating aldehyde dehydrogenase is the *ald* gene from *Clostridium beijerinckii* (Toth, *Appl. Environ. Microbiol.* 65:4973-4980 (1999)). This enzyme has been reported to reduce acetyl-CoA and butyryl-CoA to their corresponding aldehydes. This gene is very similar to *eutE* that encodes acetaldehyde dehydrogenase of *Salmonella typhimurium* and *E. coli* (Toth, *Appl. Environ. Microbiol.* 65:4973-4980 (1999)). These proteins are identified below.

Protein	GenBank ID	GI Number	Organism
<i>Msed_0709</i>	YP_001190808.1	146303492	<i>Metallosphaera sedula</i>
<i>Mcr</i>	NP_378167.1	15922498	<i>Sulfolobus tokodaii</i>
<i>asd-2</i>	NP_343563.1	15898958	<i>Sulfolobus solfataricus</i>
<i>Saci_2370</i>	YP_256941.1	70608071	<i>Sulfolobus acidocaldarius</i>
<i>Ald</i>	AAT66436	49473535	<i>Clostridium beijerinckii</i>
<i>eutE</i>	AAA80209	687645	<i>Salmonella typhimurium</i>
<i>eutE</i>	P77445	2498347	<i>Escherichia coli</i>

Crotonaldehyde reductase (alcohol forming) (Figure 2, Step E)

Enzymes exhibiting crotonaldehyde reductase (alcohol forming) activity are capable of forming crotyl alcohol from crotonaldehyde. The following enzymes can naturally possess this activity or can be engineered to exhibit this activity. Exemplary genes encoding enzymes that catalyze the conversion of an aldehyde to alcohol (i.e., alcohol dehydrogenase or equivalently aldehyde reductase) include *alrA* encoding a medium-chain alcohol dehydrogenase for C2-C14 (Tani et al., *Appl. Environ. Microbiol.* 66:5231-5235 (2000)), ADH2 from *Saccharomyces cerevisiae* (Atsumi et al., *Nature* 451:86-89 (2008)), *yqhD* from *E. coli* which has preference for molecules longer than C(3) (Sulzenbacher et al., *J. Mol. Biol.* 342:489-502 (2004)), and *bdh* I and *bdh* II from *C. acetobutylicum* which converts butyraldehyde into butanol (Walter et al., *J. Bacteriol.* 174:7149-7158 (1992)). ADH1 from *Zymomonas mobilis* has been demonstrated to have activity on a number of aldehydes including formaldehyde, acetaldehyde, propionaldehyde, butyraldehyde, and acrolein (Kinoshita, *Appl. Microbiol. Biotechnol.* 22:249-254 (1985)). Cbei_2181 from *Clostridium beijerinckii* NCIMB 8052 encodes yet another useful alcohol dehydrogenase capable of converting crotonaldehyde to crotyl alcohol.

Protein	GenBank ID	GI Number	Organism
<i>alrA</i>	BAB12273.1	9967138	<i>Acinetobacter</i> sp. Strain M-1
<i>ADH2</i>	NP_014032.1	6323961	<i>Saccharomyces cerevisiae</i>
<i>yqhD</i>	NP_417484.1	16130909	<i>Escherichia coli</i>
<i>bdh</i> I	NP_349892.1	15896543	<i>Clostridium acetobutylicum</i>
<i>bdh</i> II	NP_349891.1	15896542	<i>Clostridium acetobutylicum</i>
<i>adhA</i>	YP_162971.1	56552132	<i>Zymomonas mobilis</i>
Cbei_2181	YP_001309304.1	150017050	<i>Clostridium beijerinckii</i> NCIMB 8052

Enzymes exhibiting 4-hydroxybutyrate dehydrogenase activity (EC 1.1.1.61) also fall into this category. Such enzymes have been characterized in *Ralstonia eutropha* (Bravo et al., *J. Forensic Sci.* 49:379-387 (2004)), *Clostridium kluyveri* (Wolff et al., *Protein Expr. Purif.*

6:206-212 (1995)) and *Arabidopsis thaliana* (Breitkreuz et al., *J.Biol.Chem.* 278:41552-41556 (2003)).

Protein	GenBank ID	GI Number	Organism
<i>4hbd</i>	YP_726053.1	113867564	<i>Ralstonia eutropha</i> H16
<i>4hbd</i>	L21902.1	146348486	<i>Clostridium kluyveri</i> DSM 555
<i>4hbd</i>	Q94B07	75249805	<i>Arabidopsis thaliana</i>

Crotyl alcohol kinase (Figure 2, Step F)

- 5 Crotyl alcohol kinase enzymes catalyze the transfer of a phosphate group to the hydroxyl group of crotyl alcohol. The enzymes described below naturally possess such activity or can be engineered to exhibit this activity. Kinases that catalyze transfer of a phosphate group to an alcohol group are members of the EC 2.7.1 enzyme class. The table below lists several useful kinase enzymes in the EC 2.7.1 enzyme class.

Enzyme Commission Number	Enzyme Name	Enzyme Commission Number	Enzyme Name	Enzyme Commission Number	Enzyme Name
2.7.1.1	hexokinase	2.7.1.48	uridine kinase	2.7.1.94	acylglycerol kinase
2.7.1.2	glucokinase	2.7.1.49	hydroxymethylpyrimidine kinase	2.7.1.95	kanamycin kinase
2.7.1.3	ketoheokinase	2.7.1.50	hydroxyethylthiazole kinase	2.7.1.100	S-methyl-5-thioribose kinase
2.7.1.4	fructokinase	2.7.1.51	L-fuculokinase	2.7.1.101	tagatose kinase
2.7.1.5	rhamnulokinase	2.7.1.52	fucokinase	2.7.1.102	hamamelose kinase
2.7.1.6	galactokinase	2.7.1.53	L-xylulokinase	2.7.1.103	viomycin kinase
2.7.1.7	mannokinase	2.7.1.54	D-arabinokinase	2.7.1.105	6-phosphofructo-2-kinase
2.7.1.8	glucosamine kinase	2.7.1.55	allose kinase	2.7.1.106	glucose-1,6-bisphosphate synthase
2.7.1.10	phosphoglucokinase	2.7.1.56	1-phosphofructokinase	2.7.1.107	diacylglycerol kinase
2.7.1.11	6-phosphofructokinase	2.7.1.58	2-dehydro-3-deoxygalactonokinase	2.7.1.108	dolichol kinase
2.7.1.12	gluconokinase	2.7.1.59	N-acetylglucosamine kinase	2.7.1.113	deoxyguanosine kinase
2.7.1.13	dehydrogluconokinase	2.7.1.60	N-acylmannosamine kinase	2.7.1.114	AMP—thymidine kinase
2.7.1.14	sedoheptulokinase	2.7.1.61	acyl-phosphate—hexose phosphotransferase	2.7.1.118	ADP—thymidine kinase
2.7.1.15	ribokinase	2.7.1.62	phosphoramidate—hexose phosphotransferase	2.7.1.119	hygromycin-B 7"-O-kinase
2.7.1.16	ribulokinase	2.7.1.63	polyphosphate—glucose phosphotransferase	2.7.1.121	phosphoenolpyruvate—glycerone phosphotransferase
2.7.1.17	xylulokinase	2.7.1.64	inositol 3-kinase	2.7.1.122	xylitol kinase
2.7.1.18	phosphoribokinase	2.7.1.65	scyllo-inosamine 4-kinase	2.7.1.127	inositol-trisphosphate 3-kinase
2.7.1.19	phosphoribulokinase	2.7.1.66	undecaprenol kinase	2.7.1.130	tetraacyldisaccharide 4'-kinase
2.7.1.20	adenosine kinase	2.7.1.67	1-phosphatidylinositol 4-kinase	2.7.1.134	inositol-tetrakisphosphate 1-kinase
2.7.1.21	thymidine kinase	2.7.1.68	1-phosphatidylinositol-4-phosphate 5-kinase	2.7.1.136	macrolide 2'-kinase
2.7.1.22	ribosylnicotinamide kinase	2.7.1.69	protein-Np-phosphohistidine—sugar phosphotransferase	2.7.1.137	phosphatidylinositol 3-kinase
2.7.1.23	NAD ⁺ kinase	2.7.1.70	identical to EC 2.7.1.37.	2.7.1.138	ceramide kinase
2.7.1.24	dephospho-CoA kinase	2.7.1.71	shikimate kinase	2.7.1.140	inositol-tetrakisphosphate 5-kinase
2.7.1.25	adenylyl-sulfate kinase	2.7.1.72	streptomycin 6-kinase	2.7.1.142	glycerol—3-phosphate-glucose phosphotransferase

Enzyme Commission Number	Enzyme Name	Enzyme Commission Number	Enzyme Name	Enzyme Commission Number	Enzyme Name
2.7.1.26	riboflavin kinase	2.7.1.73	inosine kinase	2.7.1.143	diphosphate-purine nucleoside kinase
2.7.1.27	erythritol kinase	2.7.1.74	deoxycytidine kinase	2.7.1.144	tagatose-6-phosphate kinase
2.7.1.28	triokinase	2.7.1.76	deoxyadenosine kinase	2.7.1.145	deoxynucleoside kinase
2.7.1.29	glycerone kinase	2.7.1.77	nucleoside phosphotransferase	2.7.1.146	ADP-dependent phosphofructokinase
2.7.1.30	glycerol kinase	2.7.1.78	polynucleotide 5'-hydroxyl-kinase	2.7.1.147	ADP-dependent glucokinase
2.7.1.31	glycerate kinase	2.7.1.79	diphosphate—glycerol phosphotransferase	2.7.1.148	4-(cytidine 5'-diphospho)-2-C-methyl-D-erythritol kinase
2.7.1.32	choline kinase	2.7.1.80	diphosphate—serine phosphotransferase	2.7.1.149	1-phosphatidylinositol-5-phosphate 4-kinase
2.7.1.33	pantothenate kinase	2.7.1.81	hydroxylysine kinase	2.7.1.150	1-phosphatidylinositol-3-phosphate 5-kinase
2.7.1.34	pantetheine kinase	2.7.1.82	ethanolamine kinase	2.7.1.151	inositol-polyphosphate multikinase
2.7.1.35	pyridoxal kinase	2.7.1.83	pseudouridine kinase	2.7.1.153	phosphatidylinositol-4,5-bisphosphate 3-kinase
2.7.1.36	mevalonate kinase	2.7.1.84	alkylglycerone kinase	2.7.1.154	phosphatidylinositol-4-phosphate 3-kinase
2.7.1.39	homoserine kinase	2.7.1.85	β -glucoside kinase	2.7.1.156	adenosylcobinamide kinase
2.7.1.40	pyruvate kinase	2.7.1.86	NADH kinase	2.7.1.157	N-acetylgalactosamine kinase
2.7.1.41	glucose-1-phosphate phosphodismutase	2.7.1.87	streptomycin 3"-kinase	2.7.1.158	inositol-pentakisphosphate 2-kinase
2.7.1.42	riboflavin phosphotransferase	2.7.1.88	dihydrostreptomycin-6-phosphate 3'a-kinase	2.7.1.159	inositol-1,3,4-trisphosphate 5/6-kinase
2.7.1.43	glucuronokinase	2.7.1.89	thiamine kinase	2.7.1.160	2'-phosphotransferase
2.7.1.44	galacturonokinase	2.7.1.90	diphosphate—fructose-6-phosphate 1-phosphotransferase	2.7.1.161	CTP-dependent riboflavin kinase
2.7.1.45	2-dehydro-3-deoxygluconokinase	2.7.1.91	sphinganine kinase	2.7.1.162	N-acetylhexosamine 1-kinase
2.7.1.46	L-arabinokinase	2.7.1.92	5-dehydro-2-deoxygluconokinase	2.7.1.163	hygromycin B 4-O-kinase
2.7.1.47	D-ribulokinase	2.7.1.93	alkylglycerol kinase	2.7.1.164	O-phosphoserine-tRNA ^{Sec} kinase

A good candidate for this step is mevalonate kinase (EC 2.7.1.36) that phosphorylates the terminal hydroxyl group of the methyl analog, mevalonate, of 3,5-dihydroxypentanote. Some gene candidates for this step are *erg12* from *S. cerevisiae*, *mvk* from *Methanocaldococcus jannaschi*, *MVK* from *Homo sapiens*, and *mvk* from *Arabidopsis thaliana* col.

Protein	GenBank ID	GI Number	Organism
<i>erg12</i>	CAA39359.1	3684	<i>Saccharomyces cerevisiae</i>
<i>mvk</i>	Q58487.1	2497517	<i>Methanocaldococcus jannaschii</i>
<i>mvk</i>	AAH16140.1	16359371	<i>Homo sapiens</i>
<i>M\mvk</i>	NP_851084.1	30690651	<i>Arabidopsis thaliana</i>

Glycerol kinase also phosphorylates the terminal hydroxyl group in glycerol to form glycerol-3-phosphate. This reaction occurs in several species, including *Escherichia coli*,

Saccharomyces cerevisiae, and *Thermotoga maritima*. The *E. coli* glycerol kinase has been shown to accept alternate substrates such as dihydroxyacetone and glyceraldehyde (Hayashi et al., *J Biol.Chem.* 242:1030-1035 (1967)). *T. maritima* has two glycerol kinases (Nelson et al., *Nature* 399:323-329 (1999)). Glycerol kinases have been shown to have a wide range of substrate specificity. Crans and Whiteside studied glycerol kinases from four different organisms (*Escherichia coli*, *S. cerevisiae*, *Bacillus stearothermophilus*, and *Candida mycoderma*) (Crans et al., *J.Am.Chem.Soc.* 107:7008-7018 (2010); Nelson et al., *supra*, (1999)). They studied 66 different analogs of glycerol and concluded that the enzyme could accept a range of substituents in place of one terminal hydroxyl group and that the hydrogen atom at C2 could be replaced by a methyl group. Interestingly, the kinetic constants of the enzyme from all four organisms were very similar. The gene candidates are:

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>glpK</i>	AP_003883.1	89110103	<i>Escherichia coli K12</i>
<i>glpK1</i>	NP_228760.1	15642775	<i>Thermotoga maritima MSB8</i>
<i>glpK2</i>	NP_229230.1	15642775	<i>Thermotoga maritima MSB8</i>
<i>Gut1</i>	NP_011831.1	82795252	<i>Saccharomyces cerevisiae</i>

Homoserine kinase is another possible candidate that can lead to the phosphorylation of 3,5-dihydroxypentanoate. This enzyme is also present in a number of organisms including *E. coli*, *Streptomyces sp.*, and *S. cerevisiae*. Homoserine kinase from *E. coli* has been shown to have activity on numerous substrates, including, L-2-amino,1,4- butanediol, aspartate semialdehyde, and 2-amino-5-hydroxyvalerate (Huo et al., *Biochemistry* 35:16180-16185 (1996); Huo et al., *Arch.Biochem.Biophys.* 330:373-379 (1996)). This enzyme can act on substrates where the carboxyl group at the alpha position has been replaced by an ester or by a hydroxymethyl group. The gene candidates are:

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>thrB</i>	BAB96580.2	85674277	<i>Escherichia coli K12</i>
SACT1DRAFT_4809	ZP_06280784.1	282871792	<i>Streptomyces sp. ACT-1</i>
<i>Thr1</i>	AAA35154.1	172978	<i>Saccharomyces cerevisiae</i>

2-Butenyl-4-phosphate kinase (Figure 2, Step G)

2-Butenyl-4-phosphate kinase enzymes catalyze the transfer of a phosphate group to the phosphate group of 2-butenyl-4-phosphate. The enzymes described below naturally possess such activity or can be engineered to exhibit this activity. Kinases that catalyze transfer of a

phosphate group to another phosphate group are members of the EC 2.7.4 enzyme class. The table below lists several useful kinase enzymes in the EC 2.7.4 enzyme class.

Enzyme Commission Number	Enzyme Name
2.7.4.1	polyphosphate kinase
2.7.4.2	phosphomevalonate kinase
2.7.4.3	adenylate kinase
2.7.4.4	nucleoside-phosphate kinase
2.7.4.6	nucleoside-diphosphate kinase
2.7.4.7	phosphomethylpyrimidine kinase
2.7.4.8	guanylate kinase
2.7.4.9	dTMP kinase
2.7.4.10	nucleoside-triphosphate—adenylate kinase
2.7.4.11	(deoxy)adenylate kinase
2.7.4.12	T2-induced deoxynucleotide kinase
2.7.4.13	(deoxy)nucleoside-phosphate kinase
2.7.4.14	cytidylate kinase
2.7.4.15	thiamine-diphosphate kinase
2.7.4.16	thiamine-phosphate kinase
2.7.4.17	3-phosphoglyceroyl-phosphate—polyphosphate phosphotransferase
2.7.4.18	farnesyl-diphosphate kinase
2.7.4.19	5-methyldeoxycytidine-5'-phosphate kinase
2.7.4.20	dolichyl-diphosphate—polyphosphate phosphotransferase
2.7.4.21	inositol-hexakisphosphate kinase
2.7.4.22	UMP kinase
2.7.4.23	ribose 1,5-bisphosphate phosphokinase
2.7.4.24	diphosphoinositol-pentakisphosphate kinase

- Phosphomevalonate kinase enzymes are of particular interest. Phosphomevalonate kinase (EC 2.7.4.2) catalyzes the analogous transformation to 2-butenyl-4-phosphate kinase. This enzyme is encoded by *erg8* in *Saccharomyces cerevisiae* (Tsay et al., *Mol. Cell Biol.* 11:620-631 (1991)) and *mvaK2* in *Streptococcus pneumoniae*, *Staphylococcus aureus* and *Enterococcus faecalis* (Doun et al., *Protein Sci.* 14:1134-1139 (2005); Wilding et al., *J Bacteriol.* 182:4319-4327 (2000)). The *Streptococcus pneumoniae* and *Enterococcus faecalis* enzymes were cloned and characterized in *E. coli* (Pilloff et al., *J Biol. Chem.* 278:4510-4515 (2003); Doun et al., *Protein Sci.* 14:1134-1139 (2005)).

Protein	GenBank ID	GI Number	Organism
<i>Erg8</i>	AAA34596.1	171479	<i>Saccharomyces cerevisiae</i>
<i>mvaK2</i>	AAG02426.1	9937366	<i>Staphylococcus aureus</i>
<i>mvaK2</i>	AAG02457.1	9937409	<i>Streptococcus pneumoniae</i>
<i>mvaK2</i>	AAG02442.1	9937388	<i>Enterococcus faecalis</i>

Butadiene synthase (Figure 2, Step H)

Butadiene synthase catalyzes the conversion of 2-butenyl-4-diphosphate to 1,3-butadiene.

The enzymes described below naturally possess such activity or can be engineered to exhibit this activity. Isoprene synthase naturally catalyzes the conversion of dimethylallyl

- 5 diphosphate to isoprene, but can also catalyze the synthesis of 1,3-butadiene from 2-butenyl-4-diphosphate. Isoprene synthases can be found in several organisms including *Populus alba* (Sasaki et al., FEBS Letters, 2005, 579 (11), 2514-2518), *Pueraria montana* (Lindberg et al., Metabolic Eng, 2010, 12 (1), 70-79; Sharkey et al., Plant Physiol., 2005, 137 (2), 700-712), and *Populus tremula* x *Populus alba* (Miller et al., Planta, 2001, 213 (3), 483-487).
- 10 Additional isoprene synthase enzymes are described in (Chotani et al., WO/2010/031079, Systems Using Cell Culture for Production of Isoprene; Cervin et al., US Patent Application 20100003716, Isoprene Synthase Variants for Improved Microbial Production of Isoprene).

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>ispS</i>	BAD98243.1	63108310	<i>Populus alba</i>
<i>ispS</i>	AAQ84170.1	35187004	<i>Pueraria montana</i>
<i>ispS</i>	CAC35696.1	13539551	<i>Populus tremula</i> x <i>Populus alba</i>

Crotonyl-CoA hydrolase, synthetase, transferase (Figure 2, Step I)

- 15 Crotonyl-CoA hydrolase catalyzes the conversion of crotonyl-CoA to crotonate. The enzymes described below naturally possess such activity or can be engineered to exhibit this activity. 3-Hydroxyisobutyryl-CoA hydrolase efficiently catalyzes the conversion of 3-hydroxyisobutyryl-CoA to 3-hydroxyisobutyrate during valine degradation (Shimomura et al., *J Biol Chem.* 269:14248-14253 (1994)). Genes encoding this enzyme include *hibch* of
- 20 *Rattus norvegicus* (Shimomura et al., *supra*; Shimomura et al., *Methods Enzymol.* 324:229-240 (2000)) and *Homo sapiens* (Shimomura et al., *supra*). The *H. sapiens* enzyme also accepts 3-hydroxybutyryl-CoA and 3-hydroxypropionyl-CoA as substrates (Shimomura et al., *supra*). Candidate genes by sequence homology include *hibch* of *Saccharomyces cerevisiae* and BC_2292 of *Bacillus cereus*. These proteins are identified below.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>hibch</i>	Q5XIE6.2	146324906	<i>Rattus norvegicus</i>
<i>hibch</i>	Q6NVY1.2	146324905	<i>Homo sapiens</i>
<i>hibch</i>	P28817.2	2506374	<i>Saccharomyces cerevisiae</i>
<i>BC_2292</i>	AP09256	29895975	<i>Bacillus cereus</i>

Several eukaryotic acetyl-CoA hydrolases (EC 3.1.2.1) have broad substrate specificity and thus represent suitable candidate enzymes. For example, the enzyme from *Rattus norvegicus* brain (Robinson et al., Res. Commun. 71:959-965 (1976)) can react with butyryl-CoA, hexanoyl-CoA and malonyl-CoA. Though its sequence has not been reported, the enzyme from the mitochondrion of the pea leaf also has a broad substrate specificity, with demonstrated activity on acetyl-CoA, propionyl-CoA, butyryl-CoA, palmitoyl-CoA, oleoyl-CoA, succinyl-CoA, and crotonyl-CoA (Zeihner et al., Plant. Physiol. 94:20-27 (1990)). The acetyl-CoA hydrolase, ACH1, from *S. cerevisiae* represents another candidate hydrolase (Buu et al., J. Biol. Chem. 278:17203-17209 (2003)). These proteins are identified below.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>acot12</i>	NP_570103.1	18543355	<i>Rattus norvegicus</i>
<i>ACH1</i>	NP_009538	6319456	<i>Saccharomyces cerevisiae</i>

Another candidate hydrolase is the human dicarboxylic acid thioesterase, *acot8*, which exhibits activity on glutaryl-CoA, adipyl-CoA, suberyl-CoA, sebacyl-CoA, and dodecanedioyl-CoA (Westin et al., J Biol. Chem. 280:38125-38132 (2005)) and the closest *E. coli* homolog, *tesB*, which can also hydrolyze a broad range of CoA thioesters (Naggert et al., J Biol. Chem. 266:11044-11050 (1991)). A similar enzyme has also been characterized in the rat liver (Deana et al., Biochem. Int. 26:767-773 (1992)). Other potential *E. coli* thioester hydrolases include the gene products of *tesA* (Bonner et al., Chem. 247:3123-3133 (1972)), *ybgC* (Kuznetsova et al., FEMS Microbiol Rev 29:263-279 (2005); and (Zhuang et al., FEBS Lett. 516:161-163 (2002)), *paal* (Song et al., J Biol. Chem. 281:11028-11038 (2006)), and *ybdB* (Leduc et al., J Bacteriol. 189:7112-7126 (2007)). These proteins are identified below.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>tesB</i>	NP_414986	16128437	<i>Escherichia coli</i>
<i>acot8</i>	CAA15502	3191970	<i>Homo sapiens</i>
<i>acot8</i>	NP_570112	51036669	<i>Rattus norvegicus</i>
<i>tesA</i>	NP_415027	16128478	<i>Escherichia coli</i>
<i>ybgC</i>	NP_415264	16128711	<i>Escherichia coli</i>
<i>paal</i>	NP_415914	16129357	<i>Escherichia coli</i>
<i>ybdB</i>	NP_415129	16128580	<i>Escherichia coli</i>

Yet another candidate hydrolase is the glutaconate CoA-transferase from *Acidaminococcus fermentans*. This enzyme was transformed by site-directed mutagenesis into an acyl-CoA

hydrolase with activity on glutaryl-CoA, acetyl-CoA and 3-butenoyl-CoA (Mack et al., *FEBS.Lett.* 405:209-212 (1997)). This suggests that the enzymes encoding succinyl-CoA:3-ketoacid-CoA transferases and acetoacetyl-CoA:acetyl-CoA transferases can also serve as candidates for this reaction step but would require certain mutations to change their function.

5 These proteins are identified below.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>gctA</i>	CAA57199	559392	<i>Acidaminococcus fermentans</i>
<i>gctB</i>	CAA57200	559393	<i>Acidaminococcus fermentans</i>

Crotonyl-CoA synthetase catalyzes the conversion of crotonyl-CoA to crotonate. The enzymes described below naturally possess such activity or can be engineered to exhibit this activity. One candidate enzyme, ADP-forming acetyl-CoA synthetase (ACD, EC 6.2.1.13),
 10 couples the conversion of acyl-CoA esters to their corresponding acids with the concurrent synthesis of ATP. Several enzymes with broad substrate specificities have been described in the literature. ACD I from *Archaeoglobus fulgidus*, encoded by AF1211, was shown to operate on a variety of linear and branched-chain substrates including acetyl-CoA, propionyl-CoA, butyryl-CoA, acetate, propionate, butyrate, isobutyryate, isovalerate, succinate,
 15 fumarate, phenylacetate, indoleacetate (Musfeldt et al., *J Bacteriol* 184:636-644 (2002)). The enzyme from *Haloarcula marismortui* (annotated as a succinyl-CoA synthetase) accepts propionate, butyrate, and branched-chain acids (isovalerate and isobutyrate) as substrates, and was shown to operate in the forward and reverse directions (Brasen et al., *Arch Microbiol* 182:277-287 (2004)). The ACD encoded by PAE3250 from hyperthermophilic crenarchaeon
 20 *Pyrobaculum aerophilum* showed the broadest substrate range of all characterized ACDs, reacting with acetyl-CoA, isobutyryl-CoA (preferred substrate) and phenylacetyl-CoA (Brasen et al., *supra*). The enzymes from *A. fulgidus*, *H. marismortui* and *P. aerophilum* have all been cloned, functionally expressed, and characterized in *E. coli* (Musfeldt et al., *supra*; Brasen et al., *supra*). These proteins are identified below.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>AF1211</i>	NP_070039.1	11498810	<i>Archaeoglobus fulgidus</i> DSM 4304
<i>scs</i>	YP_135572.1	55377722	<i>Haloarcula marismortui</i> ATCC 43049
<i>PAE3250</i>	NP_560604.1	18313937	<i>Pyrobaculum aerophilum</i> str. IM2

Another candidate CoA synthetase is succinyl-CoA synthetase. The *sucCD* genes of *E. coli* form a succinyl-CoA synthetase complex which naturally catalyzes the formation of succinyl-CoA from succinate with the concomitant consumption of one ATP, a reaction which is reversible *in vivo* (Buck et al., *Biochem.* 24:6245-6252 (1985)). These proteins are identified below.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>sucC</i>	NP_415256.1	16128703	<i>Escherichia coli</i>
<i>sucD</i>	AAC73823.1	1786949	<i>Escherichia coli</i>

Additional exemplary CoA-ligases include the rat dicarboxylate-CoA ligase for which the sequence is yet uncharacterized (Vamecq et al., *Biochemical Journal* 230:683-693 (1985)), either of the two characterized phenylacetate-CoA ligases from *P. chrysogenum* (Lamas-Maceiras et al., *Biochem. J.* 395:147-155 (2005); Wang et al., *Biochem Biophys Res Commun* 360(2):453-458 (2007)), the phenylacetate-CoA ligase from *Pseudomonas putida* (Martinez-Blanco et al., *J. Biol. Chem.* 265:7084-7090 (1990)), and the 6-carboxyhexanoate-CoA ligase from *Bacillus subtilis* (Bower et al., *J. Bacteriol.* 178(14):4122-4130 (1996)). Additional candidate enzymes are acetoacetyl-CoA synthetases from *Mus musculus* (Hasegawa et al., *Biochim Biophys Acta* 1779:414-419 (2008)) and *Homo sapiens* (Ohgami et al., *Biochem Pharmacol* 65:989-994 (2003)) which naturally catalyze the ATP-dependant conversion of acetoacetate into acetoacetyl-CoA. These proteins are identified below.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>phl</i>	CAJ15517.1	77019264	<i>Penicillium chrysogenum</i>
<i>phlB</i>	ABS19624.1	152002983	<i>Penicillium chrysogenum</i>
<i>paaF</i>	AAC24333.2	22711873	<i>Pseudomonas putida</i>
<i>bioW</i>	NP_390902.2	50812281	<i>Bacillus subtilis</i>
<i>AACS</i>	NP_084486.1	21313520	<i>Mus musculus</i>
<i>AACS</i>	NP_076417.2	31982927	<i>Homo sapiens</i>

Crotonyl-CoA transferase catalyzes the conversion of crotonyl-CoA to crotonate. The enzymes described below naturally possess such activity or can be engineered to exhibit this activity. Many transferases have broad specificity and thus can utilize CoA acceptors as diverse as acetate, succinate, propionate, butyrate, 2-methylacetoacetate, 3-ketohexanoate, 3-ketopentanoate, valerate, crotonate, 3-mercaptopropionate, propionate, vinylacetate, butyrate, among others. For example, an enzyme from *Roseburia* sp. A2-183 was shown to have butyryl-CoA:acetate:CoA transferase and propionyl-CoA:acetate:CoA transferase activity

(Charrier et al., *Microbiology* 152, 179-185 (2006)). Close homologs can be found in, for example, *Roseburia intestinalis* L1-82, *Roseburia inulinivorans* DSM 16841, *Eubacterium rectale* ATCC 33656. Another enzyme with propionyl-CoA transferase activity can be found in *Clostridium propionicum* (Selmer et al., *Eur J Biochem* 269, 372-380 (2002)). This enzyme can use acetate, (R)-lactate, (S)-lactate, acrylate, and butyrate as the CoA acceptor (Selmer et al., *Eur J Biochem* 269, 372-380 (2002); Schweiger and Buckel, *FEBS Letters*, 171(1) 79-84 (1984)). Close homologs can be found in, for example, *Clostridium novyi* NT, *Clostridium beijerinckii* NCIMB 8052, and *Clostridium botulinum* C str. Eklund. *YgfH* encodes a propionyl CoA:succinate CoA transferase in *E. coli* (Haller et al., *Biochemistry*, 39(16) 4622-4629). Close homologs can be found in, for example, *Citrobacter youngae* ATCC 29220, *Salmonella enterica* subsp. *arizonae* serovar, and *Yersinia intermedia* ATCC 29909. These proteins are identified below .

Protein	GenBank ID	GI Number	Organism
<i>AchI</i>	AAX19660.1	60396828	<i>Roseburia</i> sp. A2-183
<i>ROSINTL182_07121</i>	ZP_04743841.2	257413684	<i>Roseburia intestinalis</i> L1-82
<i>ROSEINA2194_03642</i>	ZP_03755203.1	225377982	<i>Roseburia inulinivorans</i> DSM 16841
<i>EUBREC_3075</i>	YP_002938937.1	238925420	<i>Eubacterium rectale</i> ATCC 33656
<i>pct</i>	CAB77207.1	7242549	<i>Clostridium propionicum</i>
<i>NT01CX_2372</i>	YP_878445.1	118444712	<i>Clostridium novyi</i> NT
<i>Cbei_4543</i>	YP_001311608.1	150019354	<i>Clostridium beijerinckii</i> NCIMB 8052
<i>CBC_A0889</i>	ZP_02621218.1	168186583	<i>Clostridium botulinum</i> C str. Eklund
<i>ygfH</i>	NP_417395.1	16130821	<i>Escherichia coli</i> str. K-12 substr. MG1655
<i>CIT292_04485</i>	ZP_03838384.1	227334728	<i>Citrobacter youngae</i> ATCC 29220
<i>SARI_04582</i>	YP_001573497.1	161506385	<i>Salmonella enterica</i> subsp. <i>arizonae</i> serovar
<i>yinte0001_14430</i>	ZP_04635364.1	238791727	<i>Yersinia intermedia</i> ATCC 29909

An additional candidate enzyme is the two-unit enzyme encoded by *pcaI* and *pcaJ* in

Pseudomonas, which has been shown to have 3-oxoadipyl-CoA/succinate transferase activity (Kaschabek et al., *supra*). Similar enzymes based on homology exist in *Acinetobacter* sp. ADP1 (Kowalchuk et al., *Gene* 146:23-30 (1994)) and *Streptomyces coelicolor*. Additional exemplary succinyl-CoA:3-oxoacid-CoA transferases are present in *Helicobacter pylori*

(Corthesy-Theulaz et al., *J.Biol.Chem.* 272:25659-25667 (1997)) and *Bacillus subtilis* (Stols et al., *Protein.Expr.Purif.* 53:396-403 (2007)). These proteins are identified below.

Protein	GenBank ID	GI Number	Organism
<i>pcaI</i>	AAN69545.1	24985644	<i>Pseudomonas putida</i>
<i>pcaJ</i>	NP_746082.1	26990657	<i>Pseudomonas putida</i>
<i>pcaI</i>	YP_046368.1	50084858	<i>Acinetobacter sp. ADP1</i>
<i>pcaJ</i>	AAC37147.1	141776	<i>Acinetobacter sp. ADP1</i>
<i>pcaI</i>	NP_630776.1	21224997	<i>Streptomyces coelicolor</i>
<i>pcaJ</i>	NP_630775.1	21224996	<i>Streptomyces coelicolor</i>
<i>HPAGI_0676</i>	YP_627417	108563101	<i>Helicobacter pylori</i>
<i>HPAGI_0677</i>	YP_627418	108563102	<i>Helicobacter pylori</i>
<i>ScoA</i>	NP_391778	16080950	<i>Bacillus subtilis</i>
<i>ScoB</i>	NP_391777	16080949	<i>Bacillus subtilis</i>

A CoA transferase that can utilize acetate as the CoA acceptor is acetoacetyl-CoA

- 5 transferase, encoded by the *E. coli* *atoA* (alpha subunit) and *atoD* (beta subunit) genes (Vanderwinkel et al., *Biochem.Biophys.Res Commun.* 33:902-908 (1968); Korolev et al., *Acta Crystallogr.D Biol Crystallogr.* 58:2116-2121 (2002)). This enzyme has also been shown to transfer the CoA moiety to acetate from a variety of branched and linear acyl-CoA substrates, including isobutyrate (Matthies et al., *Appl Environ Microbiol* 58:1435-1439 (1992)),
- 10 valerate (Vanderwinkel et al., *supra*) and butanoate (Vanderwinkel et al., *supra*). Similar enzymes exist in *Corynebacterium glutamicum* ATCC 13032 (Duncan et al., *Appl Environ Microbiol* 68:5186-5190 (2002)), *Clostridium acetobutylicum* (Cary et al., *Appl Environ Microbiol* 56:1576-1583 (1990)), and *Clostridium saccharoperbutylacetonicum* (Kosaka et al., *Biosci.Biotechnol Biochem.* 71:58-68 (2007)). These proteins are identified below.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>atoA</i>	P76459.1	2492994	<i>Escherichia coli K12</i>
<i>atoD</i>	P76458.1	2492990	<i>Escherichia coli K12</i>
<i>actA</i>	YP_226809.1	62391407	<i>Corynebacterium glutamicum</i> <i>ATCC 13032</i>
<i>cg0592</i>	YP_224801.1	62389399	<i>Corynebacterium glutamicum</i> <i>ATCC 13032</i>
<i>ctfA</i>	NP_149326.1	15004866	<i>Clostridium acetobutylicum</i>
<i>ctfB</i>	NP_149327.1	15004867	<i>Clostridium acetobutylicum</i>
<i>ctfA</i>	AAP42564.1	31075384	<i>Clostridium</i> <i>saccharoperbutylacetonicum</i>
<i>ctfB</i>	AAP42565.1	31075385	<i>Clostridium</i> <i>saccharoperbutylacetonicum</i>

The above enzymes can also exhibit the desired activities on crotonyl-CoA. Additional exemplary transferase candidates are catalyzed by the gene products of *cat1*, *cat2*, and *cat3* of *Clostridium kluyveri* which have been shown to exhibit succinyl-CoA, 4-hydroxybutyryl-CoA, and butyryl-CoA transferase activity, respectively (Seedorf et al., *supra*; Sohling et al., *Eur.J Biochem.* 212:121-127 (1993); Sohling et al., *J Bacteriol.* 178:871-880 (1996)). Similar CoA transferase activities are also present in *Trichomonas vaginalis* (van Grinsven et al., *J.Biol.Chem.* 283:1411-1418 (2008)) and *Trypanosoma brucei* (Riviere et al., *J.Biol.Chem.* 279:45337-45346 (2004)). These proteins are identified below.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>cat1</i>	P38946.1	729048	<i>Clostridium kluyveri</i>
<i>cat2</i>	P38942.2	172046066	<i>Clostridium kluyveri</i>
<i>cat3</i>	EDK35586.1	146349050	<i>Clostridium kluyveri</i>
<i>TVAG 395550</i>	XP_001330176	123975034	<i>Trichomonas vaginalis G3</i>
<i>Tb11.02.0290</i>	XP_828352	71754875	<i>Trypanosoma brucei</i>

The glutaconate-CoA-transferase (EC 2.8.3.12) enzyme from anaerobic bacterium *Acidaminococcus fermentans* reacts with diacid glutaconyl-CoA and 3-butenoyl-CoA (Mack et al., *FEBS Lett.* 405:209-212 (1997)). The genes encoding this enzyme are *gctA* and *gctB*. This enzyme has reduced but detectable activity with other CoA derivatives including glutaryl-CoA, 2-hydroxyglutaryl-CoA, adipyl-CoA and acrylyl-CoA (Buckel et al., *Eur.J.Biochem.* 118:315-321 (1981)). The enzyme has been cloned and expressed in *E. coli* (Mack et al., *Eur.J.Biochem.* 226:41-51 (1994)). These proteins are identified below.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>gctA</i>	CAA57199.1	559392	<i>Acidaminococcus fermentans</i>
<i>gctB</i>	CAA57200.1	559393	<i>Acidaminococcus fermentans</i>

Crotonate reductase (Figure 2, Step J)

Crotonate reductase enzymes are capable of catalyzing the conversion of crotonate to crotonaldehyde. The enzymes described below naturally possess such activity or can be engineered to exhibit this activity. Carboxylic acid reductase catalyzes the magnesium, ATP and NADPH-dependent reduction of carboxylic acids to their corresponding aldehydes (Venkitasubramanian et al., *J. Biol. Chem.* 282:478-485 (2007)). This enzyme, encoded by *car*, was cloned and functionally expressed in *E. coli* (Venkitasubramanian et al., *J. Biol. Chem.* 282:478-485 (2007)). Expression of the *npt* gene product improved activity of the enzyme via post-transcriptional modification. The *npt* gene encodes a specific phosphopantetheine transferase (PPTase) that converts the inactive apo-enzyme to the active holo-enzyme. The natural substrate of this enzyme is vanillic acid, and the enzyme exhibits broad acceptance of aromatic and aliphatic substrates (Venkitasubramanian et al., in *Biocatalysis in the Pharmaceutical and Biotechnology Industries*, ed. R.N. Patel, Chapter 15, pp. 425-440, CRC Press LLC, Boca Raton, FL. (2006)).

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>Car</i>	AAR91681.1	40796035	<i>Nocardia iowensis</i> (sp. NRRL 5646)
<i>Npt</i>	ABI83656.1	114848891	<i>Nocardia iowensis</i> (sp. NRRL 5646)

Additional *car* and *npt* genes can be identified based on sequence homology.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>fadD9</i>	YP_978699.1	121638475	<i>Mycobacterium bovis</i> BCG
<i>BCG_2812c</i>	YP_978898.1	121638674	<i>Mycobacterium bovis</i> BCG
<i>nfa20150</i>	YP_118225.1	54023983	<i>Nocardia farcinica</i> IFM 10152
<i>nfa40540</i>	YP_120266.1	54026024	<i>Nocardia farcinica</i> IFM 10152
<i>SGR_6790</i>	YP_001828302.1	182440583	<i>Streptomyces griseus</i> subsp. <i>griseus</i> NBRC 13350
<i>SGR_665</i>	YP_001822177.1	182434458	<i>Streptomyces griseus</i> subsp. <i>griseus</i> NBRC 13350

Protein	GenBank ID	GI Number	Organism
<i>MSMEG_2956</i>	YP_887275.1	118473501	<i>Mycobacterium smegmatis</i> MC2 155
<i>MSMEG_5739</i>	YP_889972.1	118469671	<i>Mycobacterium smegmatis</i> MC2 155
<i>MSMEG_2648</i>	YP_886985.1	118471293	<i>Mycobacterium smegmatis</i> MC2 155
<i>MAP1040c</i>	NP_959974.1	41407138	<i>Mycobacterium avium</i> subsp. <i>paratuberculosis</i> K-10
<i>MAP2899c</i>	NP_961833.1	41408997	<i>Mycobacterium avium</i> subsp. <i>paratuberculosis</i> K-10
<i>MMAR_2117</i>	YP_001850422.1	183982131	<i>Mycobacterium marinum</i> M
<i>MMAR_2936</i>	YP_001851230.1	183982939	<i>Mycobacterium marinum</i> M
<i>MMAR_1916</i>	YP_001850220.1	183981929	<i>Mycobacterium marinum</i> M
<i>TpauDRAFT_33060</i>	ZP_04027864.1	227980601	<i>Tsukamurella paurometabola</i> DSM 20162
<i>TpauDRAFT_20920</i>	ZP_04026660.1	227979396	<i>Tsukamurella paurometabola</i> DSM 20162
<i>CPCC7001_1320</i>	ZP_05045132.1	254431429	<i>Cyanobium</i> PCC7001
<i>DDBDRAFT_0187729</i>	XP_636931.1	66806417	<i>Dictyostelium discoideum</i> AX4

An additional enzyme candidate found in *Streptomyces griseus* is encoded by the *griC* and *griD* genes. This enzyme is believed to convert 3-amino-4-hydroxybenzoic acid to 3-amino-4-hydroxybenzaldehyde as deletion of either *griC* or *griD* led to accumulation of extracellular 3-acetyl-amino-4-hydroxybenzoic acid, a shunt product of 3-amino-4-hydroxybenzoic acid metabolism (Suzuki, et al., *J. Antibiot.* 60(6):380-387 (2007)). Co-expression of *griC* and *griD* with SGR_665, an enzyme similar in sequence to the *Nocardia iowensis* *npt*, can be beneficial.

Protein	GenBank ID	GI Number	Organism
<i>griC</i>	YP_001825755.1	182438036	<i>Streptomyces griseus</i> subsp. <i>griseus</i> NBRC 13350
<i>Grid</i>	YP_001825756.1	182438037	<i>Streptomyces griseus</i> subsp. <i>griseus</i> NBRC 13350

An enzyme with similar characteristics, alpha-aminoadipate reductase (AAR, EC 1.2.1.31), participates in lysine biosynthesis pathways in some fungal species. This enzyme naturally reduces alpha-aminoadipate to alpha-aminoadipate semialdehyde. The carboxyl group is first

activated through the ATP-dependent formation of an adenylate that is then reduced by NAD(P)H to yield the aldehyde and AMP. Like CAR, this enzyme utilizes magnesium and requires activation by a PPTase. Enzyme candidates for AAR and its corresponding PPTase are found in *Saccharomyces cerevisiae* (Morris et al., *Gene* 98:141-145 (1991)), *Candida albicans* (Guo et al., *Mol. Genet. Genomics* 269:271-279 (2003)), and *Schizosaccharomyces pombe* (Ford et al., *Curr. Genet.* 28:131-137 (1995)). The AAR from *S. pombe* exhibited significant activity when expressed in *E. coli* (Guo et al., *Yeast* 21:1279-1288 (2004)). The AAR from *Penicillium chrysogenum* accepts S-carboxymethyl-L-cysteine as an alternate substrate, but did not react with adipate, L-glutamate or diaminopimelate (Hijarrubia et al., *J. Biol. Chem.* 278:8250-8256 (2003)). The gene encoding the *P. chrysogenum* PPTase has not been identified to date.

Protein	GenBank ID	GI Number	Organism
<i>LYS2</i>	AAA34747.1	171867	<i>Saccharomyces cerevisiae</i>
<i>LYS5</i>	P50113.1	1708896	<i>Saccharomyces cerevisiae</i>
<i>LYS2</i>	AAC02241.1	2853226	<i>Candida albicans</i>
<i>LYS5</i>	AAO26020.1	28136195	<i>Candida albicans</i>
<i>Lys1p</i>	P40976.3	13124791	<i>Schizosaccharomyces pombe</i>
<i>Lys7p</i>	Q10474.1	1723561	<i>Schizosaccharomyces pombe</i>
<i>Lys2</i>	CAA74300.1	3282044	<i>Penicillium chrysogenum</i>

Crotonyl-CoA reductase (alcohol forming) (Figure 2, Step K)

Crotonaldehyde reductase (alcohol forming) enzymes catalyze the 2 reduction steps required to form crotyl alcohol from crotonyl-CoA. Exemplary 2-step oxidoreductases that convert an acyl-CoA to an alcohol are provided below. Such enzymes can naturally convert crotonyl-CoA to crotyl alcohol or can be engineered to do so. These enzymes include those that transform substrates such as acetyl-CoA to ethanol (e.g., *adhE* from *E. coli* (Kessler et al., *FEBS.Lett.* 281:59-63 (1991))) and butyryl-CoA to butanol (e.g. *adhE2* from *C. acetobutylicum* (Fontaine et al., *J.Bacteriol.* 184:821-830 (2002))). The *adhE2* enzyme from *C. acetobutylicum* was specifically shown in ref. (Burk et al., *supra*, (2008)) to produce BDO from 4-hydroxybutyryl-CoA. In addition to reducing acetyl-CoA to ethanol, the enzyme encoded by *adhE* in *Leuconostoc mesenteroides* has been shown to oxide the branched chain compound isobutyraldehyde to isobutyryl-CoA (Kazahaya et al., *J.Gen.Appl.Microbiol.* 18:43-55 (1972); Koo et al., *Biotechnol. Lett.* 27:505-510 (2005)).

Protein	GenBank ID	GI Number	Organism
<i>adhE</i>	NP_415757.1	16129202	<i>Escherichia coli</i>

<i>adhE2</i>	AAK09379.1	12958626	<i>Clostridium acetobutylicum</i>
<i>adhE</i>	AAV66076.1	55818563	<i>Leuconostoc mesenteroides</i>

Another exemplary enzyme can convert malonyl-CoA to 3-HP. An NADPH-dependent enzyme with this activity has characterized in *Chloroflexus aurantiacus* where it participates in the 3-hydroxypropionate cycle (Hugler et al., *supra*, (2002); Strauss et al., 215:633-643 (1993)). This enzyme, with a mass of 300 kDa, is highly substrate-specific and shows little sequence similarity to other known oxidoreductases (Hugler et al., *supra*, (2002)). No enzymes in other organisms have been shown to catalyze this specific reaction; however there is bioinformatic evidence that other organisms can have similar pathways (Klatt et al., *Environ Microbiol.* 9:2067-2078 (2007)). Enzyme candidates in other organisms including *Roseiflexus castenholzii*, *Erythrobacter sp. NAP1* and marine gamma proteobacterium HTCC2080 can be inferred by sequence similarity.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>mcr</i>	AAS20429.1	42561982	<i>Chloroflexus aurantiacus</i>
<i>Rcas_2929</i>	YP_001433009.1	156742880	<i>Roseiflexus castenholzii</i>
<i>NAP1_02720</i>	ZP_01039179.1	85708113	<i>Erythrobacter sp. NAP1</i>
MGP2080_00535	ZP_01626393.1	119504313	marine gamma proteobacterium HTCC2080

Glutaconyl-CoA decarboxylase (Figure 2, Step L)

Glutaconyl-CoA decarboxylase enzymes, characterized in glutamate-fermenting anaerobic bacteria, are sodium-ion translocating decarboxylases that utilize biotin as a cofactor and are composed of four subunits (alpha, beta, gamma, and delta) (Boiangiu et al., *J Mol. Microbiol Biotechnol* 10:105-119 (2005); Buckel, *Biochim Biophys Acta.* 1505:15-27 (2001)). Such enzymes have been characterized in *Fusobacterium nucleatum* (Beatrix et al., *Arch Microbiol.* 154:362-369 (1990)) and *Acidaminococcus fermentans* (Braune et al., *Mol. Microbiol* 31:473-487 (1999)). Analogs to the *F. nucleatum* glutaconyl-CoA decarboxylase alpha, beta and delta subunits are found in *S. aciditrophicus*. A gene annotated as an enoyl-CoA dehydrogenase, *syn_00480*, another GCD, is located in a predicted operon between a biotin-carboxyl carrier (*syn_00479*) and a glutaconyl-CoA decarboxylase alpha subunit (*syn_00481*). The protein sequences for exemplary gene products can be found using the following GenBank accession numbers shown below.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>gcdA</i>	CAA49210	49182	<i>Acidaminococcus fermentans</i>
<i>gcdC</i>	AAC69172	3777506	<i>Acidaminococcus fermentans</i>
<i>gcdD</i>	AAC69171	3777505	<i>Acidaminococcus fermentans</i>
<i>gcdB</i>	AAC69173	3777507	<i>Acidaminococcus fermentans</i>
FN0200	AAL94406	19713641	<i>Fusobacterium nucleatum</i>
FN0201	AAL94407	19713642	<i>Fusobacterium nucleatum</i>
FN0204	AAL94410	19713645	<i>Fusobacterium nucleatum</i>
<i>syn_00479</i>	YP_462066	85859864	<i>Syntrophus aciditrophicus</i>
<i>syn_00481</i>	YP_462068	85859866	<i>Syntrophus aciditrophicus</i>
<i>syn_01431</i>	YP_460282	85858080	<i>Syntrophus aciditrophicus</i>
<i>syn_00480</i>	ABC77899	85722956	<i>Syntrophus aciditrophicus</i>

Glutaryl-CoA dehydrogenase (Figure 2 Step M)

Glutaryl-CoA dehydrogenase (GCD, EC 1.3.99.7 and EC 4.1.1.70) is a bifunctional enzyme that catalyzes the oxidative decarboxylation of glutaryl-CoA to crotonyl-CoA (Figure 3, step 3). Bifunctional GCD enzymes are homotetramers that utilize electron transfer flavoprotein as an electron acceptor (Hartel et al., *Arch Microbiol.* 159:174-181 (1993)). Such enzymes were first characterized in cell extracts of *Pseudomonas* strains KB740 and K172 during growth on aromatic compounds (Hartel et al., *supra*, (1993)), but the associated genes in these organisms is unknown. Genes encoding glutaryl-CoA dehydrogenase (*gcdH*) and its cognate transcriptional regulator (*gcdR*) were identified in *Azoarcus* sp. *CIB* (Blazquez et al., *Environ Microbiol.* 10:474-482 (2008)). An *Azoarcus* strain deficient in *gcdH* activity was used to identify the a heterologous gene *gcdH* from *Pseudomonas putida* (Blazquez et al., *supra*, (2008)). The cognate transcriptional regulator in *Pseudomonas putida* has not been identified but the locus *PP_0157* has a high sequence homology (> 69% identity) to the *Azoarcus* enzyme. Additional GCD enzymes are found in *Pseudomonas fluorescens* and *Paracoccus denitrificans* (Husain et al., *J Bacteriol.* 163:709-715 (1985)). The human GCD has been extensively studied, overexpressed in *E. coli* (Dwyer et al., *Biochemistry* 39:11488-11499 (2000)), crystallized, and the catalytic mechanism involving a conserved glutamate residue in the active site has been described (Fu et al., *Biochemistry* 43:9674-9684 (2004)). A GCD in *Syntrophus aciditrophicus* operates in the CO₂-assimilating direction during growth on crotonate (Moultaki et al., *Appl Environ Microbiol.* 73:930-938 (2007)). Two GCD genes in *S. aciditrophicus* were identified by protein sequence homology to the *Azoarcus* GcdH: *syn_00480* (31%) and *syn_01146* (31%). No significant homology was found to the

Azoarcus GcdR regulatory protein. The protein sequences for exemplary gene products can be found using the following GenBank accession numbers shown below.

Protein	GenBank ID	GI Number	Organism
<i>gcdH</i>	ABM69268.1	123187384	<i>Azoarcus</i> sp. CIB
<i>gcdR</i>	ABM69269.1	123187385	<i>Azoarcus</i> sp. CIB
<i>gcdH</i>	AAN65791.1	24981507	<i>Pseudomonas putida</i> KT2440
<i>PP_0157 (gcdR)</i>	AAN65790.1	24981506	<i>Pseudomonas putida</i> KT2440
<i>gcdH</i>	YP_257269.1	70733629	<i>Pseudomonas fluorescens</i> Pf-5
<i>gcvA (gcdR)</i>	YP_257268.1	70733628	<i>Pseudomonas fluorescens</i> Pf-5
<i>gcd</i>	YP_918172.1	119387117	<i>Paracoccus denitrificans</i>
<i>gcdR</i>	YP_918173.1	119387118	<i>Paracoccus denitrificans</i>
<i>gcd</i>	AAH02579.1	12803505	<i>Homo sapiens</i>
<i>syn_00480</i>	ABC77899	85722956	<i>Syntrophus aciditrophicus</i>
<i>syn_01146</i>	ABC76260	85721317	<i>Syntrophus aciditrophicus</i>

3-Aminobutyryl-CoA deaminase (Figure 2, Step N)

- 5 3-aminobutyryl-CoA is an intermediate in lysine fermentation. It also can be formed from acetoacetyl-CoA via a transaminase or an aminating dehydrogenase. 3-aminobutyryl-CoA deaminase (or 3-aminobutyryl-CoA ammonia lyase) catalyzes the deamination of 3-aminobutyryl-CoA to form crotonyl-CoA. This reversible enzyme is present in *Fusobacterium nucleatum*, *Porphyromonas gingivalis*, *Thermoanaerobacter tengcongensis*, and several other organisms and is co-localized with several genes involved in lysine fermentation (Kreimeyer et al., J Biol Chem, 2007, 282(10) 7191-7197).

Protein	GenBank ID	GI Number	Organism
<i>kal</i>	NP_602669.1	19705174	<i>Fusobacterium nucleatum</i> subsp. <i>nucleatum</i> ATCC 25586
<i>kal</i>	NP_905282.1	34540803	<i>Porphyromonas gingivalis</i> W83
<i>kal</i>	NP_622376.1	20807205	<i>Thermoanaerobacter tengcongensis</i> MB4

4-Hydroxybutyryl-CoA dehydratase (Figure 2, Step O)

- 15 Several enzymes naturally catalyze the dehydration of 4-hydroxybutyryl-CoA to crotonyl-CoA. This transformation is required for 4-aminobutyrate fermentation by *Clostridium aminobutyricum* (Scherf et al., Eur.J Biochem. 215:421-429 (1993)) and succinate-ethanol fermentation by *Clostridium kluyveri* (Scherf et al., Arch.Microbiol 161:239-245 (1994)). The transformation is also a key step in Archaea, for example, *Metallosphaera sedula*, as part of the 3-hydroxypropionate/4-hydroxybutyrate autotrophic carbon dioxide assimilation pathway

(Berg et al., *supra*, (2007)). The reversibility of 4-hydroxybutyryl-CoA dehydratase is well-documented (Muh et al., *Biochemistry*. 35:11710-11718 (1996); Friedrich et al., *Angew.Chem.Int.Ed.Engl.* 47:3254-3257 (2008); Muh et al., *Eur.J.Biochem.* 248:380-384 (1997)) and the equilibrium constant has been reported to be about 4 on the side of crotonoyl-CoA (Scherf and Buckel, *supra*, (1993)).

Protein	GenBank ID	GI Number	Organism
<i>AbfD</i>	CAB60035	70910046	<i>Clostridium aminobutyricum</i>
<i>AbfD</i>	YP_001396399	153955634	<i>Clostridium kluyveri</i>
<i>Msed_1321</i>	YP_001191403	146304087	<i>Metallosphaera sedula</i>
<i>Msed_1220</i>	YP_001191305	146303989	<i>Metallosphaera sedula</i>

Crotyl alcohol diphosphokinase (Figure 2, Step P)

Crotyl alcohol diphosphokinase enzymes catalyze the transfer of a diphosphate group to the hydroxyl group of crotyl alcohol. The enzymes described below naturally possess such activity or can be engineered to exhibit this activity. Kinases that catalyze transfer of a diphosphate group are members of the EC 2.7.6 enzyme class. The table below lists several useful kinase enzymes in the EC 2.7.6 enzyme class.

Enzyme Commission Number	Enzyme Name
2.7.6.1	ribose-phosphate diphosphokinase
2.7.6.2	thiamine diphosphokinase
2.7.6.3	2-amino-4-hydroxy-6-hydroxymethyldihydropteridine diphosphokinase
2.7.6.4	nucleotide diphosphokinase
2.7.6.5	GTP diphosphokinase

Of particular interest are ribose-phosphate diphosphokinase enzymes which have been identified in *Escherichia coli* (Hove-Jenson et al., J Biol Chem, 1986, 261(15);6765-71) and *Mycoplasma pneumoniae* M129 (McElwain et al, International Journal of Systematic Bacteriology, 1988, 38:417-423) as well as thiamine diphosphokinase enzymes. Exemplary thiamine diphosphokinase enzymes are found in *Arabidopsis thaliana* (Ajjawi, Plant Mol Biol, 2007, 65(1-2);151-62).

Protein	GenBank ID	GI Number	Organism
prs	NP_415725.1	16129170	<i>Escherichia coli</i>
prsA	NP_109761.1	13507812	<i>Mycoplasma pneumoniae</i> M129

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
TPK1	BAH19964.1	222424006	<i>Arabidopsis thaliana col</i>
TPK2	BAH57065.1	227204427	<i>Arabidopsis thaliana col</i>

Erythrose-4-phosphate reductase (Figure 3, Step A)

In Step A of the pathway, erythrose-4-phosphate is converted to erythritol-4-phosphate by the erythrose-4-phosphate reductase or erythritol-4-phosphate dehydrogenase. The reduction of erythrose-4-phosphate was observed in *Leuconostoc oenos* during the production of erythritol (Veiga-da-Cunha et al., *J Bacteriol.* 175:3941-3948 (1993)). NADPH was identified as the cofactor (Veiga-da-Cunha et al., *supra*, (1993)). However, gene for erythrose-4-phosphate was not identified. Thus, it is possible to identify the erythrose-4-phosphate reductase gene from *Leuconostoc oenos* and apply to this step. Additionally, enzymes catalyzing similar reactions can be utilized for this step. An example of these enzymes is 1-deoxy-D-xylulose-5-phosphate reductoisomerase (EC 1.1.1.267) catalyzing the conversion of 1-deoxy-D-xylulose 5-phosphate to 2-C-methyl-D-erythritol-4-phosphate, which has one additional methyl group comparing to Step A. The *dxr* or *ispC* genes encode the 1-deoxy-D-xylulose-5-phosphate reductoisomerase have been well studied: the Dxr proteins from *Escherichia coli* and *Mycobacterium tuberculosis* were purified and their crystal structures were determined (Yajima et al., *Acta Crystallogr.Sect.F.Struct.Biol.Cryst.Comm.* 63:466-470 (2007); Mac et al., *J Mol.Biol.* 345:115-127 (2005); Henriksson et al., *Acta Crystallogr.D.Biol.Crystallogr.* 62:807-813 (2006); Henriksson et al., *J Biol.Chem.* 282:19905-19916 (2007)); the Dxr protein from *Synechocystis sp* was studied by site-directed mutagenesis with modified activity and altered kinetics (Fernandes et al., *Biochim.Biophys.Acta* 1764:223-229 (2006); Fernandes et al., *Arch.Biochem.Biophys.* 444:159-164 (2005)). Furthermore, glyceraldehyde 3-phosphate reductase YghZ from *Escherichia coli* catalyzes the conversion between glyceraldehyde 3-phosphate and glycerol-3-phosphate (Desai et al., *Biochemistry* 47:7983-7985 (2008)) can also be applied to this step. The following genes can be used for Step A conversion:

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>dxr</i>	P45568.2	2506592	<i>Escherichia coli</i> strain K12
<i>dxr</i>	A5U6M4.1	166218269	<i>Mycobacterium tuberculosis</i>
<i>dxr</i>	Q55663.1	2496789	<i>Synechocystis sp.</i> strain PCC6803
<i>yghZ</i>	NP_417474.1	16130899	<i>Escherichia coli</i> strain K12

Erythritol-4-phosphate cytidyltransferase (Figure 3, Step B)

In Step B of the pathway, erythritol-4-phosphate is converted to 4-(cytidine 5'-diphospho)-erythritol by the erythritol-4-phosphate cytidyltransferase or 4-(cytidine 5'-diphospho)-erythritol synthase. The exact enzyme for this step has not been identified. However, enzymes catalyzing similar reactions can be applied to this step. An example is the 2-C-methyl-D-erythritol 4-phosphate cytidyltransferase or 4-(cytidine 5'-diphospho)-2-C-methyl-D-erythritol synthase (EC 2.7.7.60). The 2-C-methyl-D-erythritol 4-phosphate cytidyltransferase is in the methylerythritol phosphate pathway for the isoprenoid biosynthesis and catalyzes the conversion between 2-C-methyl-D-erythritol 4-phosphate and 4-(cytidine 5'-diphospho)-2-C-methyl-D-erythritol, with an extra methyl group comparing to Step B conversion. The 2-C-methyl-D-erythritol 4-phosphate cytidyltransferase is encoded by *ispD* gene and the crystal structure of *Escherichia coli* IspD was determined (Kemp et al., *Acta Crystallogr.D.Biol.Crystallogr.* 57:1189-1191 (2001); Kemp et al., *Acta Crystallogr.D.Biol.Crystallogr.* 59:607-610 (2003); Richard et al., *Nat.Struct.Biol.* 8:641-648 (2001)). The *ispD* gene from *Mycobacterium tuberculosis* H37Rv was cloned and expressed in *Escherichia coli*, and the recombinant proteins were purified with N-terminal His-tag (Shi et al., *J Biochem.Mol.Biol.* 40:911-920 (2007)). Additionally, the *Streptomyces coelicolor* *ispD* gene was cloned and expressed in *E. coli*, and the recombinant proteins were characterized physically and kinetically (Cane et al., *Bioorg.Med.Chem.* 9:1467-1477 (2001)). The following genes can be used for Step B conversion:

Protein	GenBank ID	GI Number	Organism
<i>ispD</i>	Q46893.3	2833415	<i>Escherichia coli</i> strain K12
<i>ispD</i>	A5U8Q7.1	166215456	<i>Mycobacterium tuberculosis</i>
<i>ispD</i>	Q9L0Q8.1	12230289	<i>Streptomyces coelicolor</i>

4-(Cytidine 5'-diphospho)-erythritol kinase (Figure 3, Step C)

In Step C of the pathway, 4-(cytidine 5'-diphospho)-erythritol is converted to 2-phospho-4-(cytidine 5'-diphospho)-erythritol by the 4-(cytidine 5'-diphospho)-erythritol kinase. The exact enzyme for this step has not been identified. However, enzymes catalyzing similar reactions can be applied to this step. An example is the 4-diphosphocytidyl-2-C-methylerythritol kinase (EC 2.7.1.148). The 4-diphosphocytidyl-2-C-methylerythritol kinase is also in the methylerythritol phosphate pathway for the isoprenoid biosynthesis and catalyzes the conversion between 4-(cytidine 5'-diphospho)-2-C-methyl-D-erythritol and 2-

phospho-4-(cytidine 5'-diphospho)-2-C-methyl-D-erythritol, with an extra methyl group comparing to Step C conversion. The 4-diphosphocytidyl-2-C-methylerythritol kinase is encoded by *ispE* gene and the crystal structures of *Escherichia coli*, *Thermus thermophilus* HB8, and *Aquifex aeolicus* IspE were determined (Sgraja et al., *FEBS J* 275:2779-2794 (2008); Miallau et al., *Proc.Natl.Acad.Sci.U.S.A* 100:9173-9178 (2003); Wada et al., *J Biol.Chem.* 278:30022-30027 (2003)). The *ispE* genes from above organism were cloned and expressed, and the recombinant proteins were purified for crystallization. The following genes can be used for Step C conversion:

Protein	GenBank ID	GI Number	Organism
<i>ispE</i>	P62615.1	50402174	<i>Escherichia coli</i> strain K12
<i>ispE</i>	P83700.1	51316201	<i>Thermus thermophilus</i> HB8
<i>ispE</i>	O67060.1	6919911	<i>Aquifex aeolicus</i>

10 **Erythritol 2,4-cyclodiphosphate synthase (Figure 3, Step D)**

In Step D of the pathway, 2-phospho-4-(cytidine 5'-diphospho)-erythritol is converted to erythritol-2,4-cyclodiphosphate by the Erythritol 2,4-cyclodiphosphate synthase. The exact enzyme for this step has not been identified. However, enzymes catalyzing similar reactions can be applied to this step. An example is the 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase (EC 4.6.1.12). The 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase is also in the methylerythritol phosphate pathway for the isoprenoid biosynthesis and catalyzes the conversion between 2-phospho-4-(cytidine 5'diphospho)-2-C-methyl-D-erythritol and 2-C-methyl-D-erythritol-2,4-cyclodiphosphate, with an extra methyl group comparing to step D conversion. The 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase is encoded by *ispF* gene and the crystal structures of *Escherichia coli*, *Thermus thermophilus*, *Haemophilus influenzae*, and *Campylobacter jejuni* IspF were determined (Richard et al., *J Biol.Chem.* 277:8667-8672 (2002); Steinbacher et al., *J Mol.Biol.* 316:79-88 (2002); Lehmann et al., *Proteins* 49:135-138 (2002); Kishida et al., *Acta Crystallogr.D.Biol.Crystallogr.* 59:23-31 (2003); Gabrielsen et al., *J Biol.Chem.* 279:52753-52761 (2004)). The *ispF* genes from above organism were cloned and expressed, and the recombinant proteins were purified for crystallization. The following genes can be used for Step D conversion:

Protein	GenBank ID	GI Number	Organism
<i>ispF</i>	P62617.1	51317402	<i>Escherichia coli</i> strain K12
<i>ispF</i>	Q8RQP5.1	51701599	<i>Thermus thermophilus</i> HB8
<i>ispF</i>	P44815.1	1176081	<i>Haemophilus influenzae</i>
<i>ispF</i>	Q9PM68.1	12230305	<i>Campylobacter jejuni</i>

1-Hydroxy-2-butenyl 4-diphosphate synthase (Figure 3, Step E)

Step E of Figure 3 entails conversion of erythritol-2,4-cyclodiphosphate to 1-hydroxy-2-butenyl 4-diphosphate by 1-hydroxy-2-butenyl 4-diphosphate synthase. An enzyme with this activity has not been characterized to date. This transformation is analogous to the reduction of 2-C-methyl-D-erythritol-2,4-cyclodiphosphate to 1-hydroxy-2-methyl-2-(E)-butenyl 4-diphosphate by (E)-4-hydroxy-3-methylbut-2-enyl-diphosphate synthase (EC 1.17.7.1). This enzyme is an iron-sulfur protein that participates in the non-mevalonate pathway for isoprenoid biosynthesis found in bacteria and plants. Most bacterial enzymes including the *E. coli* enzyme, encoded by *ispG*, utilize reduced ferredoxin or flavodoxin as an electron donor (Zepeck et al., *J Org.Chem.* 70:9168-9174 (2005)). An analogous enzyme from the thermophilic cyanobacterium *Thermosynechococcus elongatus* BP-1, encoded by *gcpE*, was heterologously expressed and characterized in *E. coli* (Okada et al., *J Biol.Chem.* 280:20672-20679 (2005)). Additional enzyme candidates from *Thermus thermophilus* and *Arabidopsis thaliana* have been characterized and expressed in *E. coli* (Seemann et al., *J Biol.Inorg.Chem.* 10:131-137 (2005); Kollas et al., *FEBS Lett.* 532:432-436 (2002)).

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>ispG</i>	NP_417010.1	16130440	<i>Escherichia coli</i>
<i>gcpE</i>	NP_681786.1	22298539	<i>Thermosynechococcus elongatus</i>
<i>gcpE</i>	AAO21364.1	27802077	<i>Thermus thermophilus</i>
<i>gcpE</i>	AAO15446.1	27462472	<i>Arabidopsis thaliana</i>

1-Hydroxy-2-butenyl 4-diphosphate reductase (Figure 3, Step F)

The concurrent dehydration and reduction of 1-hydroxy-2-butenyl 4-diphosphate is catalyzed by an enzyme with 1-hydroxy-2-butenyl 4-diphosphate reductase activity (Figure 3, Step F). Such an enzyme will form a mixture of products, butenyl 4-diphosphate or 2-butenyl 4-diphosphate. An analogous reaction is catalyzed by 4-hydroxy-3-methylbut-2-enyl diphosphate reductase (EC 1.17.1.2) in the non-mevalonate pathway for isoprenoid biosynthesis. This enzyme is an iron-sulfur protein that utilizes reduced ferredoxin or flavodoxin as an electron donor. Maximal activity of 4-hydroxy-3-methylbut-2-enyl diphosphate reductase *E. coli*, encoded by *ispH*, requires both flavodoxin and flavodoxin reductase (Wolff et al., *FEBS Lett.* 541:115-120 (2003); Grawert et al., *J Am.Chem.Soc.* 126:12847-12855 (2004)). In the characterized catalytic system, reduced flavodoxin is regenerated by the NAD(P)⁺-dependent flavodoxin reductase. The enzyme from *Aquifex aeolicus*, encoded by *lytB*, was expressed as a His-tagged enzyme in *E. coli* and characterized

(Altincicek et al., *FEBS Lett.* 532:437-440 (2002)). An analogous enzyme in plants is encoded by *hdr* of *Arabidopsis thaliana* (Botella-Pavia et al., *Plant J* 40:188-199 (2004)).

Protein	GenBank ID	GI Number	Organism
<i>ispH</i>	AAL38655.1	18652795	<i>Escherichia coli</i>
<i>lytB</i>	O67625.1	8928180	<i>Aquifex aeolicus</i>
<i>hdr</i>	NP_567965.1	18418433	<i>Arabidopsis thaliana</i>

Altering the expression level of genes involved in iron-sulfur cluster formation can have an advantageous effect on the activities of iron-sulfur proteins in the proposed pathways (for example, enzymes required in Steps E and F of Figure 3). In *E. coli*, it was demonstrated that overexpression of the iron-sulfur containing protein IspH (analogous to Step F of Figure 3) is enhanced by coexpression of genes from the *isc* region involved in assembly of iron-sulfur clusters (Grawert et al., *J Am.Chem.Soc.* 126:12847-12855 (2004)). The gene cluster is composed of the genes *icsS*, *icsU*, *icsA*, *hscB*, *hscA* and *fdx*. Overexpression of these genes was shown to improve the synthetic capability of the iron-sulfur assembly pipeline, required for functional expression of iron-sulfur proteins. A similar approach can be applicable in the current application.

Protein	GenBank ID	GI Number	Organism
<i>iscS</i>	AAT48142.1	48994898	<i>Escherichia coli</i>
<i>iscU</i>	AAC75582.1	1788878	<i>Escherichia coli</i>
<i>iscA</i>	AAC75581.1	1788877	<i>Escherichia coli</i>
<i>hscB</i>	AAC75580.1	1788876	<i>Escherichia coli</i>
<i>hscA</i>	AAC75579.1	1788875	<i>Escherichia coli</i>
<i>fdx</i>	AAC75578.1	1788874	<i>Escherichia coli</i>

15 **Butenyl 4-diphosphate isomerase (Figure 3, Step G)**

Butenyl 4-diphosphate isomerase catalyzes the reversible interconversion of 2-butenyl-4-diphosphate and butenyl-4-diphosphate. The following enzymes can naturally possess this activity or can be engineered to exhibit this activity. Useful genes include those that encode enzymes that interconvert isopenenyl diphosphate and dimethylallyl diphosphate. These include isopentenyl diphosphate isomerase enzymes from *Escherichia coli* (Rodríguez-Concepción et al., *FEBS Lett*, 473(3):328-332), *Saccharomyces cerevisiae* (Anderson et al., *J Biol Chem*, 1989, 264(32):19169-75), and *Sulfolobus shibatae* (Yamashita et al, *Eur J Biochem*, 2004, 271(6):1087-93). The reaction mechanism of isomerization, catalyzed by the Idi protein of *E. coli*, has been characterized in mechanistic detail (de Ruyck et al., *J*

Biol.Chem. 281:17864-17869 (2006)). Isopentenyl diphosphate isomerase enzymes from *Saccharomyces cerevisiae*, *Bacillus subtilis* and *Haematococcus pluvialis* have been heterologously expressed in *E. coli* (Laupitz et al., *Eur.J Biochem.* 271:2658-2669 (2004); Kajiwara et al., *Biochem.J* 324 (Pt 2):421-426 (1997)).

Protein	GenBank ID	GI Number	Organism
<i>Idi</i>	NP_417365.1	16130791	<i>Escherichia coli</i>
<i>IDII</i>	NP_015208.1	6325140	<i>Saccharomyces cerevisiae</i>
<i>Idi</i>	BAC82424.1	34327946	<i>Sulfolobus shibatae</i>
<i>Idi</i>	AAC32209.1	3421423	<i>Haematococcus pluvialis</i>
<i>Idi</i>	BAB32625.1	12862826	<i>Bacillus subtilis</i>

5

Butadiene synthase (Figure 3, Step H)

Butadiene synthase catalyzes the conversion of 2-butenyl-4-diphosphate to 1,3-butadiene.

The enzymes described below naturally possess such activity or can be engineered to exhibit this activity. Isoprene synthase naturally catalyzes the conversion of dimethylallyl

10 diphosphate to isoprene, but can also catalyze the synthesis of 1,3-butadiene from 2-butenyl-4-diphosphate. Isoprene synthases can be found in several organisms including *Populus alba* (Sasaki et al., *FEBS Letters*, 579 (11), 2514-2518 (2005)), *Pueraria montana* (Lindberg et al., *Metabolic Eng.*, 12(1):70-79 (2010); Sharkey et al., *Plant Physiol.*, 137(2):700-712 (2005)), and *Populus tremula* x *Populus alba* (Miller et al., *Planta*, 213(3):483-487 (2001)).

15 Additional isoprene synthase enzymes are described in (Chotani et al., WO/2010/031079, Systems Using Cell Culture for Production of Isoprene; Cervin et al., US Patent Application 20100003716, Isoprene Synthase Variants for Improved Microbial Production of Isoprene).

Protein	GenBank ID	GI Number	Organism
<i>ispS</i>	BAD98243.1	63108310	<i>Populus alba</i>
<i>ispS</i>	AAQ84170.1	35187004	<i>Pueraria montana</i>
<i>ispS</i>	CAC35696.1	13539551	<i>Populus tremula</i> x <i>Populus alba</i>

Erythrose-4-phosphate kinase (Figure 3, Step I)

20 In Step I of the pathway, erythrose-4-phosphate is converted to erythrose by the erythrose-4-phosphate kinase. In industrial fermentative production of erythritol by yeasts, glucose was converted to erythrose-4-phosphate through the pentose phosphate pathway and erythrose-4-phosphate was dephosphorylated and reduced to produce erythritol (Moon et al.,

Appl. Microbiol. Biotechnol. 86:1017-1025 (2010)). Thus, erythrose-4-phosphate kinase is present in many of these erythritol-producing yeasts, including *Trichosporonoides megachiliensis* SN-G42 (Sawada et al., *J. Biosci. Bioeng.* 108:385-390 (2009)), *Candida magnolia* (Kohl et al., *Biotechnol. Lett.* 25:2103-2105 (2003)), and *Torula* sp. (HAJNY et al., *Appl. Microbiol.* 12:240-246 (1964); Oh et al., *J. Ind. Microbiol. Biotechnol.* 26:248-252 (2001)). However, the erythrose-4-phosphate kinase genes were not identified yet. There are many polyol phosphotransferases with wide substrate range that can be applied to this step. An example is the triose kinase (EC 2.7.1.28) catalyzing the reversible conversion between glyceraldehydes and glyceraldehydes-3-phosphate, which are one carbon shorter comparing to Step I. Other examples include the xylulokinase (EC 2.7.1.17) or arabinokinase (EC 2.7.1.54) that catalyzes phosphorylation of 5C polyol aldehyde. The following genes can be used for Step I conversion:

Protein	GenBank ID	GI Number	Organism
<i>xylB</i>	P09099.1	139849	<i>Escherichia coli</i> strain K12
<i>xksI</i>	P42826.2	1723736	<i>Saccharomyces cerevisiae</i>
<i>xylB</i>	P29444.1	267426	<i>Klebsiella pneumoniae</i>
<i>dakI</i>	Q9HFC5	74624685	<i>Zygosaccharomyces rouxii</i>

Erythrose reductase (Figure 3, Step J)

In Step J of the pathway, erythrose is converted to erythritol by the erythrose reductase. In industrial fermentative production of erythritol by yeasts, glucose was converted to erythrose-4-phosphate through the pentose phosphate pathway and erythrose-4-phosphate was dephosphorylated and reduced to produce erythritol (Moon et al., *supra*, (2010)). Thus, erythrose reductase is present in many of these erythritol-producing yeasts, including *Trichosporonoides megachiliensis* SN-G42 (Sawada et al., *supra*, (2009)), *Candida magnolia* (Kohl et al., *supra*, (2003)), and *Torula* sp. (HAJNY et al., *supra*, (1964); Oh et al., *supra*, (2001)). Erythrose reductase was characterized and purified from *Torula corallina* (Lee et al., *Biotechnol. Prog.* 19:495-500 (2003); Lee et al., *Appl. Environ. Microbiol.* 68:4534-4538 (2002)), *Candida magnolia* (Lee et al., *Appl. Environ. Microbiol.* 69:3710-3718 (2003)) and *Trichosporonoides megachiliensis* SN-G42 (Sawada et al., *supra*, (2009)). Several erythrose reductase genes were cloned and can be applied to Step J. The following genes can be used for Step J conversion:

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>alr</i>	ACT78580.1	254679867	<i>Candida magnoliae</i>
<i>er1</i>	BAD90687.1	60458781	<i>Trichosporonoides megachiliensis</i>
<i>er2</i>	BAD90688.1	60458783	<i>Trichosporonoides megachiliensis</i>
<i>er3</i>	BAD90689.1	60458785	<i>Trichosporonoides megachiliensis</i>

Erythritol kinase (Figure 3, Step K)

In Step K of the pathway, erythritol is converted to erythritol-4-phosphate by the erythritol kinase. Erythritol kinase (EC 2.7.1.27) catalyzes the phosphorylation of erythritol. Erythritol kinase was characterized in erythritol utilizing bacteria such as *Brucella abortus* (Sperry et al., *J Bacteriol.* 121:619-630 (1975)). The *eryA* gene of *Brucella abortus* has been functionally expressed in *Escherichia coli* and the resultant EryA was shown to catalyze the ATP-dependent conversion of erythritol to erythritol-4-phosphate (Lillo et al., *Bioorg.Med.Chem.Lett.* 13:737-739 (2003)). The following genes can be used for Step K conversion:

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>eryA</i>	Q8YCU8	81850596	<i>Brucella melitensis</i>
<i>eriA</i>	Q92NH0	81774560	<i>Sinorhizobium meliloti</i>
<i>eryA</i>	YP_001108625.1	134102964	<i>Saccharopolyspora erythraea</i> NRRL 2338

Malonyl-CoA:acetyl-CoA acyltransferase (Figure 4, Step A)

In Step A of the pathway described in Figure 4, malonyl-CoA and acetyl-CoA are condensed to form 3-oxoglutaryl-CoA by malonyl-CoA:acetyl-CoA acyl transferase, a beta-keothiolase. Although no enzyme with activity on malonyl-CoA has been reported to date, a good candidate for this transformation is beta-ketoadipyl-CoA thiolase (EC 2.3.1.174), also called 3-oxoadipyl-CoA thiolase that converts beta-ketoadipyl-CoA to succinyl-CoA and acetyl-CoA, and is a key enzyme of the beta-ketoadipate pathway for aromatic compound degradation. The enzyme is widespread in soil bacteria and fungi including *Pseudomonas putida* (Harwood et al., *J Bacteriol.* 176:6479-6488 (1994)) and *Acinetobacter calcoaceticus* (Doten et al., *J Bacteriol.* 169:3168-3174 (1987)). The gene products encoded by *pcaF* in *Pseudomonas* strain B13 (Kaschabek et al., *J Bacteriol.* 184:207-215 (2002)), *phaD* in *Pseudomonas putida* U (Olivera et al., *supra*, (1998)), *paaE* in *Pseudomonas fluorescens* ST (Di Gennaro et al., *Arch Microbiol.* 88:117-125 (2007)), and *paaJ* from *E. coli* (Nogales et al., *Microbiology*, 153:357-365 (2007)) also catalyze this transformation. Several beta-

ketothiolases exhibit significant and selective activities in the oxoadipyl-CoA forming direction including *bkt* from *Pseudomonas putida*, *pcaF* and *bkt* from *Pseudomonas aeruginosa* PAO1, *bkt* from *Burkholderia ambifaria* AMMD, *paaJ* from *E. coli*, and *phaD* from *P. putida*. These enzymes can also be employed for the synthesis of 3-oxoglutaryl-CoA, a compound structurally similar to 3-oxoadipyl-CoA.

Protein	GenBank ID	GI Number	Organism
<i>paaJ</i>	NP_415915.1	16129358	<i>Escherichia coli</i>
<i>pcaF</i>	AAL02407	17736947	<i>Pseudomonas knackmussii</i> (B13)
<i>phaD</i>	AAC24332.1	3253200	<i>Pseudomonas putida</i>
<i>pcaF</i>	AAA85138.1	506695	<i>Pseudomonas putida</i>
<i>pcaF</i>	AAC37148.1	141777	<i>Acinetobacter calcoaceticus</i>
<i>paaE</i>	ABF82237.1	106636097	<i>Pseudomonas fluorescens</i>
<i>bkt</i>	YP_777652.1	115360515	<i>Burkholderia ambifaria</i> AMMD
<i>bkt</i>	AAG06977.1	9949744	<i>Pseudomonas aeruginosa</i> PAO1
<i>pcaF</i>	AAG03617.1	9946065	<i>Pseudomonas aeruginosa</i> PAO1

Another relevant beta-ketothiolase is oxopimeloyl-CoA:glutaryl-CoA acyltransferase (EC 2.3.1.16) that combines glutaryl-CoA and acetyl-CoA to form 3-oxopimeloyl-CoA. An enzyme catalyzing this transformation is found in *Ralstonia eutropha* (formerly known as *Alcaligenes eutrophus*), encoded by genes *bktB* and *bktC* (Slater et al., *J. Bacteriol.* 180:1979-1987 (1998); Haywood et al., *FEMS Microbiology Letters* 52:91-96 (1988)). The sequence of the BktB protein is known; however, the sequence of the BktC protein has not been reported. The *pim* operon of *Rhodopseudomonas palustris* also encodes a beta-ketothiolase, encoded by *pimB*, predicted to catalyze this transformation in the degradative direction during benzoyl-CoA degradation (Harrison et al., *Microbiology* 151:727-736 (2005)). A beta-ketothiolase enzyme candidate in *S. aciditrophicus* was identified by sequence homology to *bktB* (43% identity, *evalue* = 1e-93).

Protein	GenBank ID	GI Number	Organism
<i>bktB</i>	YP_725948	11386745	<i>Ralstonia eutropha</i>
<i>pimB</i>	CAE29156	39650633	<i>Rhodopseudomonas palustris</i>
<i>syn_02642</i>	YP_462685.1	85860483	<i>Syntrophus aciditrophicus</i>

Beta-ketothiolase enzymes catalyzing the formation of beta-ketovaleryl-CoA from acetyl-CoA and propionyl-CoA can also be able to catalyze the formation of 3-oxoglutaryl-CoA. *Zoogloea ramigera* possesses two ketothiolases that can form β -ketovaleryl-CoA from

propionyl-CoA and acetyl-CoA and *R. eutropha* has a β -oxidation ketothiolase that is also capable of catalyzing this transformation (Slater et al., *J. Bacteriol*, 180:1979-1987 (1998)).

The sequences of these genes or their translated proteins have not been reported, but several candidates in *R. eutropha*, *Z. ramigera*, or other organisms can be identified based on

5 sequence homology to *bktB* from *R. eutropha*. These include:

Protein	GenBank ID	GI Number	Organism
<i>phaA</i>	YP_725941.1	113867452	<i>Ralstonia eutropha</i>
<i>h16_A1713</i>	YP_726205.1	113867716	<i>Ralstonia eutropha</i>
<i>pcaF</i>	YP_728366.1	116694155	<i>Ralstonia eutropha</i>
<i>h16_B1369</i>	YP_840888.1	116695312	<i>Ralstonia eutropha</i>
<i>h16_A0170</i>	YP_724690.1	113866201	<i>Ralstonia eutropha</i>
<i>h16_A0462</i>	YP_724980.1	113866491	<i>Ralstonia eutropha</i>
<i>h16_A1528</i>	YP_726028.1	113867539	<i>Ralstonia eutropha</i>
<i>h16_B0381</i>	YP_728545.1	116694334	<i>Ralstonia eutropha</i>
<i>h16_B0662</i>	YP_728824.1	116694613	<i>Ralstonia eutropha</i>
<i>h16_B0759</i>	YP_728921.1	116694710	<i>Ralstonia eutropha</i>
<i>h16_B0668</i>	YP_728830.1	116694619	<i>Ralstonia eutropha</i>
<i>h16_A1720</i>	YP_726212.1	113867723	<i>Ralstonia eutropha</i>
<i>h16_A1887</i>	YP_726356.1	113867867	<i>Ralstonia eutropha</i>
<i>phbA</i>	P07097.4	135759	<i>Zoogloea ramigera</i>
<i>bktB</i>	YP_002005382.1	194289475	<i>Cupriavidus taiwanensis</i>
<i>Rmet_1362</i>	YP_583514.1	94310304	<i>Ralstonia metallidurans</i>
<i>Bphy_0975</i>	YP_001857210.1	186475740	<i>Burkholderia phymatum</i>

Additional candidates include beta-ketothiolases that are known to convert two molecules of acetyl-CoA into acetoacetyl-CoA (EC 2.1.3.9). Exemplary acetoacetyl-CoA thiolase enzymes include the gene products of *atoB* from *E. coli* (Martin et al., *supra*, (2003)), *thlA* and *thlB* from *C. acetobutylicum* (Hanai et al., *supra*, (2007); Winzer et al., *supra*, (2000)), and *ERG10* from *S. cerevisiae* (Hiser et al., *supra*, (1994)).

Protein	GenBank ID	GI Number	Organism
<i>toB</i>	NP_416728	16130161	<i>Escherichia coli</i>
<i>thlA</i>	NP_349476.1	15896127	<i>Clostridium acetobutylicum</i>
<i>thlB</i>	NP_149242.1	15004782	<i>Clostridium acetobutylicum</i>
<i>ERG10</i>	NP_015297	6325229	<i>Saccharomyces cerevisiae</i>

3-oxoglutaryl-CoA reductase (ketone-reducing) (Figure 4, Step B)

This enzyme catalyzes the reduction of the 3-oxo group in 3-oxoglutaryl-CoA to the 3-hydroxy group in Step B of the pathway shown in Figure 4.

3-Oxoacyl-CoA dehydrogenase enzymes convert 3-oxoacyl-CoA molecules into 3-hydroxyacyl-CoA molecules and are often involved in fatty acid beta-oxidation or phenylacetate catabolism. For example, subunits of two fatty acid oxidation complexes in *E. coli*, encoded by *fadB* and *fadJ*, function as 3-hydroxyacyl-CoA dehydrogenases (Binstock et al., *Methods Enzymol.* 71 Pt C:403-411 (1981)). Furthermore, the gene products encoded by *phaC* in *Pseudomonas putida* U (Olivera et al., *supra*, (1998)) and *paaC* in *Pseudomonas fluorescens* ST (Di et al., *supra*, (2007)) catalyze the reversible oxidation of 3-hydroxyadipyl-CoA to form 3-oxoadipyl-CoA, during the catabolism of phenylacetate or styrene. In addition, given the proximity in *E. coli* of *paaH* to other genes in the phenylacetate degradation operon (Nogales et al., *supra*, (2007)) and the fact that *paaH* mutants cannot grow on phenylacetate (Ismail et al., *supra*, (2003)), it is expected that the *E. coli* *paaH* gene encodes a 3-hydroxyacyl-CoA dehydrogenase.

Protein	GenBank ID	GI Number	Organism
<i>fadB</i>	P21177.2	119811	<i>Escherichia coli</i>
<i>fadJ</i>	P77399.1	3334437	<i>Escherichia coli</i>
<i>paaH</i>	NP_415913.1	16129356	<i>Escherichia coli</i>
<i>phaC</i>	NP_745425.1	26990000	<i>Pseudomonas putida</i>
<i>paaC</i>	ABF82235.1	106636095	<i>Pseudomonas fluorescens</i>

3-Hydroxybutyryl-CoA dehydrogenase, also called acetoacetyl-CoA reductase, catalyzes the reversible NAD(P)H-dependent conversion of acetoacetyl-CoA to 3-hydroxybutyryl-CoA. This enzyme participates in the acetyl-CoA fermentation pathway to butyrate in several species of *Clostridia* and has been studied in detail (Jones and Woods, *supra*, (1986)). Enzyme candidates include *hbd* from *C. acetobutylicum* (Boynton et al., *J. Bacteriol.* 178:3015-3024 (1996)), *hbd* from *C. beijerinckii* (Colby et al., *Appl Environ. Microbiol.* 58:3297-3302 (1992)), and a number of similar enzymes from *Metallosphaera sedula* (Berg et al., *supra*, (2007)). The enzyme from *Clostridium acetobutylicum*, encoded by *hbd*, has been cloned and functionally expressed in *E. coli* (Youngleson et al., *supra*, (1989)). Yet other genes demonstrated to reduce acetoacetyl-CoA to 3-hydroxybutyryl-CoA are *phbB* from *Zoogloea ramigera* (Ploux et al., *supra*, (1988)) and *phaB* from *Rhodobacter sphaeroides* (Alber et al., *supra*, (2006)). The former gene is NADPH-dependent, its nucleotide sequence has been determined (Peoples and Sinskey, *supra*, (1989)) and the gene has been expressed in *E. coli*. Additional genes include *hbd1* (C-terminal domain) and *hbd2* (N-terminal domain) in *Clostridium kluyveri* (Hillmer and Gottschalk, *Biochim. Biophys. Acta* 3334:12-23 (1974)) and *HSD17B10* in *Bos taurus* (WAKIL et al., *supra*, (1954)).

Protein	GenBank ID	GI Number	Organism
<i>hbd</i>	NP_349314.1	15895965	<i>Clostridium acetobutylicum</i>
<i>hbd</i>	AAM14586.1	20162442	<i>Clostridium beijerinckii</i>
<i>Msed_1423</i>	YP_001191505	146304189	<i>Metallosphaera sedula</i>
<i>Msed_0399</i>	YP_001190500	146303184	<i>Metallosphaera sedula</i>
<i>Msed_0389</i>	YP_001190490	146303174	<i>Metallosphaera sedula</i>
<i>Msed_1993</i>	YP_001192057	146304741	<i>Metallosphaera sedula</i>
<i>hbd2</i>	EDK34807.1	146348271	<i>Clostridium kluyveri</i>
<i>hbd1</i>	EDK32512.1	146345976	<i>Clostridium kluyveri</i>
<i>HSD17B10</i>	O02691.3	3183024	<i>Bos taurus</i>
<i>phaB</i>	YP_353825.1	77464321	<i>Rhodobacter sphaeroides</i>
<i>phbB</i>	P23238.1	130017	<i>Zoogloea ramigera</i>

3-hydroxyglutaryl-CoA reductase (aldehyde forming) (Figure 4, Step C)

3-hydroxyglutaryl-CoA reductase reduces 3-hydroxyglutaryl-CoA to 3-hydroxy-5-oxopentanoate. Several acyl-CoA dehydrogenases reduce an acyl-CoA to its corresponding aldehyde (EC 1.2.1). Exemplary genes that encode such enzymes include the *Acinetobacter calcoaceticus* *acr1* encoding a fatty acyl-CoA reductase (Reiser and Somerville, *supra*, (1997)), the *Acinetobacter* sp. *M-1* fatty acyl-CoA reductase (Ishige et al., *supra*, (2002)), and a CoA- and NADP- dependent succinate semialdehyde dehydrogenase encoded by the *sucD* gene in *Clostridium kluyveri* (Sohling and Gottschalk, *supra*, (1996); Sohling and Gottschalk, *supra*, (1996)). *SucD* of *P. gingivalis* is another succinate semialdehyde dehydrogenase (Takahashi et al., *supra*, (2000)). The enzyme acylating acetaldehyde dehydrogenase in *Pseudomonas* sp, encoded by *bphG*, is yet another as it has been demonstrated to oxidize and acylate acetaldehyde, propionaldehyde, butyraldehyde, isobutyraldehyde and formaldehyde (Powlowski et al., *supra*, (1993)). In addition to reducing acetyl-CoA to ethanol, the enzyme encoded by *adhE* in *Leuconostoc mesenteroides* has been shown to oxidize the branched chain compound isobutyraldehyde to isobutyryl-CoA (Koo et al., *Biotechnol Lett.* 27:505-510 (2005)). Butyraldehyde dehydrogenase catalyzes a similar reaction, conversion of butyryl-CoA to butyraldehyde, in solventogenic organisms such as *Clostridium saccharoperbutylacetonicum* (Kosaka et al., *Biosci.Biotechnol Biochem.* 71:58-68 (2007)).

Protein	GenBank ID	GI Number	Organism
<i>acr1</i>	YP_047869.1	50086359	<i>Acinetobacter calcoaceticus</i>
<i>acr1</i>	AAC45217	1684886	<i>Acinetobacter baylyi</i>
<i>acr1</i>	BAB85476.1	18857901	<i>Acinetobacter</i> sp. Strain <i>M-1</i>
<i>sucD</i>	P38947.1	172046062	<i>Clostridium kluyveri</i>
<i>sucD</i>	NP_904963.1	34540484	<i>Porphyromonas gingivalis</i>
<i>bphG</i>	BAA03892.1	425213	<i>Pseudomonas</i> sp
<i>adhE</i>	AAV66076.1	55818563	<i>Leuconostoc mesenteroides</i>

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>bld</i>	AAP42563.1	31075383	<i>Clostridium saccharoperbutylacetonicum</i>

An additional enzyme type that converts an acyl-CoA to its corresponding aldehyde is malonyl-CoA reductase which transforms malonyl-CoA to malonic semialdehyde. Malonyl-CoA reductase is a key enzyme in autotrophic carbon fixation via the 3-hydroxypropionate cycle in thermoacidophilic archaeal bacteria (Berg et al., *supra*, (2007b); Thauer, *supra*, (2007)). The enzyme utilizes NADPH as a cofactor and has been characterized in *Metallosphaera* and *Sulfolobus spp* (Alber et al., *supra*, (2006); Hugler et al., *supra*, (2002)). The enzyme is encoded by *Msed_0709* in *Metallosphaera sedula* (Alber et al., *supra*, (2006); Berg et al., *supra*, (2007b)). A gene encoding a malonyl-CoA reductase from *Sulfolobus tokodaii* was cloned and heterologously expressed in *E. coli* (Alber et al., *supra*, (2006)). This enzyme has also been shown to catalyze the conversion of methylmalonyl-CoA to its corresponding aldehyde (WO/2007/141208). Although the aldehyde dehydrogenase functionality of these enzymes is similar to the bifunctional dehydrogenase from *Chloroflexus aurantiacus*, there is little sequence similarity. Both malonyl-CoA reductase enzyme candidates have high sequence similarity to aspartate-semialdehyde dehydrogenase, an enzyme catalyzing the reduction and concurrent dephosphorylation of aspartyl-4-phosphate to aspartate semialdehyde. Additional gene candidates can be found by sequence homology to proteins in other organisms including *Sulfolobus solfataricus* and *Sulfolobus acidocaldarius*. Yet another acyl-CoA reductase (aldehyde forming) candidate is the *ald* gene from *Clostridium beijerinckii* (Toth et al., *Appl Environ. Microbiol* 65:4973-4980 (1999)). This enzyme has been reported to reduce acetyl-CoA and butyryl-CoA to their corresponding aldehydes. This gene is very similar to *eutE* that encodes acetaldehyde dehydrogenase of *Salmonella typhimurium* and *E. coli* (Toth et al., *supra*, (1999)).

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>MSED_0709</i>	YP_001190808.1	146303492	<i>Metallosphaera sedula</i>
<i>mcr</i>	NP_378167.1	15922498	<i>Sulfolobus tokodaii</i>
<i>asd-2</i>	NP_343563.1	15898958	<i>Sulfolobus solfataricus</i>
<i>Saci_2370</i>	YP_256941.1	70608071	<i>Sulfolobus acidocaldarius</i>
<i>Ald</i>	AAT66436	9473535	<i>Clostridium beijerinckii</i>
<i>eutE</i>	AAA80209	687645	<i>Salmonella typhimurium</i>
<i>eutE</i>	P77445	2498347	<i>Escherichia coli</i>

3-hydroxy-5-oxopentanoate reductase (Figure 4, Step D)

This enzyme reduces the terminal aldehyde group in 3-hydroxy-5-oxopentanoate to the alcohol group. Exemplary genes encoding enzymes that catalyze the conversion of an aldehyde to alcohol (i.e., alcohol dehydrogenase or equivalently aldehyde reductase, 1.1.1.a) include *alrA* encoding a medium-chain alcohol dehydrogenase for C2-C14 (Tani et al., *supra*, (2000)), ADH2 from *Saccharomyces cerevisiae* (Atsumi et al., *supra*, (2008)), *yqhD* from *E. coli* which has preference for molecules longer than C(3) (Sulzenbacher et al., *supra*, (2004)), and *bdh I* and *bdh II* from *C. acetobutylicum* which converts butyrylaldehyde into butanol (Walter et al., *supra*, (1992)). The gene product of *yqhD* catalyzes the reduction of acetaldehyde, malondialdehyde, propionaldehyde, butyraldehyde, and acrolein using NADPH as the cofactor (Perez et al., 283:7346-7353 (2008); Perez et al., *J Biol.Chem.* 283:7346-7353 (2008)). The *adhA* gene product from *Zymomonas mobilis* has been demonstrated to have activity on a number of aldehydes including formaldehyde, acetaldehyde, propionaldehyde, butyraldehyde, and acrolein (Kinoshita et al., *Appl Microbiol Biotechnol* 22:249-254 (1985)).

Protein	GenBank ID	GI Number	Organism
<i>alrA</i>	BAB12273.1	9967138	<i>Acinetobacter sp. Strain M-1</i>
<i>ADH2</i>	NP_014032.1	6323961	<i>Saccharomyces cerevisiae</i>
<i>yqhD</i>	NP_417484.1	16130909	<i>Escherichia coli</i>
<i>bdh I</i>	NP_349892.1	15896543	<i>Clostridium acetobutylicum</i>
<i>bdh II</i>	NP_349891.1	15896542	<i>Clostridium acetobutylicum</i>
<i>adhA</i>	YP_162971.1	56552132	<i>Zymomonas mobilis</i>

Enzymes exhibiting 4-hydroxybutyrate dehydrogenase activity (EC 1.1.1.61) also fall into this category. Such enzymes have been characterized in *Ralstonia eutropha* (Bravo et al., *supra*, (2004)), *Clostridium kluyveri* (Wolff and Kenealy, *supra*, (1995)) and *Arabidopsis thaliana* (Breitkreuz et al., *supra*, (2003)). The *A. thaliana* enzyme was cloned and characterized in yeast [12882961]. Yet another gene is the alcohol dehydrogenase *adhI* from *Geobacillus thermoglucosidasius* (Jeon et al., *J Biotechnol* 135:127-133 (2008)).

Protein	GenBank ID	GI Number	Organism
4hbd	YP_726053.1	113867564	<i>Ralstonia eutropha H16</i>
4hbd	EDK35022.1	146348486	<i>Clostridium kluyveri</i>
4hbd	Q94B07	75249805	<i>Arabidopsis thaliana</i>
adhI	AAR91477.1	40795502	<i>Geobacillus thermoglucosidasius</i>

Another exemplary *enzyme* is 3-hydroxyisobutyrate dehydrogenase (EC 1.1.1.31) which catalyzes the reversible oxidation of 3-hydroxyisobutyrate to methylmalonate semialdehyde. This enzyme participates in valine, leucine and isoleucine degradation and has been identified in bacteria, eukaryotes, and mammals. The enzyme encoded by *P84067* from *Thermus thermophilus* HB8 has been structurally characterized (Lokanath et al., *J Mol Biol* 352:905-17 (2005)). The reversibility of the human 3-hydroxyisobutyrate dehydrogenase was demonstrated using isotopically-labeled substrate (Manning et al., *Biochem J* 231:481-4 (1985)). Additional genes encoding this enzyme include *3hidh* in *Homo sapiens* (Hawes et al., *Methods Enzymol* 324:218-228 (2000)) and *Oryctolagus cuniculus* (Hawes et al., *supra*, (2000); Chowdhury et al., *Biosci.Biotechnol Biochem.* 60:2043-2047 (1996)), *mmsb* in *Pseudomonas aeruginosa*, and *dhat* in *Pseudomonas putida* (Aberhart et al., *J Chem.Soc.[Perkin 1]* 6:1404-1406 (1979); Chowdhury et al., *supra*, (1996); Chowdhury et al., *Biosci.Biotechnol Biochem.* 67:438-441 (2003)).

Protein	GenBank ID	GI Number	Organism
P84067	P84067	75345323	<i>Thermus thermophilus</i>
mmsb	P28811.1	127211	<i>Pseudomonas aeruginosa</i>
dhat	Q59477.1	2842618	<i>Pseudomonas putida</i>
3hidh	P31937.2	12643395	<i>Homo sapiens</i>
3hidh	P32185.1	416872	<i>Oryctolagus cuniculus</i>

The conversion of malonic *semialdehyde* to 3-HP can also be accomplished by two other enzymes: NADH-dependent 3-hydroxypropionate dehydrogenase and NADPH-dependent malonate semialdehyde reductase. An NADH-dependent 3-hydroxypropionate dehydrogenase is thought to participate in beta-alanine biosynthesis pathways from propionate in bacteria and plants (Rathinasabapathi B., *Journal of Plant Pathology* 159:671-674 (2002); Stadtman, *J.Am.Chem.Soc.* 77:5765-5766 (1955)). This enzyme has not been associated with a gene in any organism to date. NADPH-dependent malonate semialdehyde reductase catalyzes the reverse reaction in autotrophic CO₂-fixing bacteria. Although the enzyme activity has been detected in *Metallosphaera sedula*, the identity of the gene is not known (Alber et al., *supra*, (2006)).

3,5-dihydroxypentanoate kinase (Figure 4, Step E)

This enzyme *phosphorylates* 3,5-dihydroxypentanoate in Figure 4 (Step E) to form 3-hydroxy-5-phosphonatooxypentanoate (3H5PP). This transformation can be catalyzed by

enzymes in the EC class 2.7.1 that enable the ATP-dependent transfer of a phosphate group to an alcohol.

A good candidate for this step is mevalonate kinase (EC 2.7.1.36) that phosphorylates the terminal hydroxyl group of the methyl analog, mevalonate, of 3,5-dihydroxypentanoate. Some gene candidates for this step are *erg12* from *S. cerevisiae*, *mvk* from *Methanocaldococcus jannaschi*, *MVK* from *Homo sapiens*, and *mvk* from *Arabidopsis thaliana* col.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>erg12</i>	CAA39359.1	3684	<i>Saccharomyces cerevisiae</i>
<i>mvk</i>	Q58487.1	2497517	<i>Methanocaldococcus jannaschii</i>
<i>mvk</i>	AAH16140.1	16359371	<i>Homo sapiens</i>
<i>M\mvk</i>	NP_851084.1	30690651	<i>Arabidopsis thaliana</i>

Glycerol kinase also phosphorylates the terminal hydroxyl group in glycerol to form glycerol-3-phosphate. This reaction occurs in several species, including *Escherichia coli*,

- 10 *Saccharomyces cerevisiae*, and *Thermotoga maritima*. The *E. coli* glycerol kinase has been shown to accept alternate substrates such as dihydroxyacetone and glyceraldehyde (Hayashi and Lin, *supra*, (1967)). *T. maritima* has two glycerol kinases (Nelson et al., *supra*, (1999)). Glycerol kinases have been shown to have a wide range of substrate specificity. Crans and Whiteside studied glycerol kinases from four different organisms (*Escherichia coli*, *S.*
- 15 *cerevisiae*, *Bacillus stearothermophilus*, and *Candida mycoderma*) (Crans and Whitesides, *supra*, (2010); Nelson et al., *supra*, (1999)). They studied 66 different analogs of glycerol and concluded that the enzyme could accept a range of substituents in place of one terminal hydroxyl group and that the hydrogen atom at C2 could be replaced by a methyl group. Interestingly, the kinetic constants of the enzyme from all four organisms were very similar.
- 20 The gene candidates are:

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>glpK</i>	AP_003883.1	89110103	<i>Escherichia coli</i> K12
<i>glpK1</i>	NP_228760.1	15642775	<i>Thermotoga maritima</i> MSB8
<i>glpK2</i>	NP_229230.1	15642775	<i>Thermotoga maritima</i> MSB8
<i>Gut1</i>	NP_011831.1	82795252	<i>Saccharomyces cerevisiae</i>

Homoserine kinase is another possible candidate that can lead to the phosphorylation of 3,5-dihydroxypentanoate. This enzyme is also present in a number of organisms including *E. coli*,

Streptomyces sp., and *S. cerevisiae*. Homoserine kinase from *E. coli* has been shown to have activity on numerous substrates, including, L-2-amino,1,4- butanediol, aspartate semialdehyde, and 2-amino-5-hydroxyvalerate (Huo and Viola, *supra*, (1996); Huo and Viola, *supra*, (1996)). This enzyme can act on substrates where the carboxyl group at the alpha position has been replaced by an ester or by a hydroxymethyl group. The gene candidates are:

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>thrB</i>	BAB96580.2	85674277	<i>Escherichia coli K12</i>
<i>SACT1DRAFT 4809</i>	ZP_06280784.1	282871792	<i>Streptomyces sp. ACT-1</i>
<i>Thr1</i>	AAA35154.1	172978	<i>Saccharomyces cerevisiae</i>

3H5PP kinase (Figure 4, Step F)

- 10 Phosphorylation of 3H5PP to 3H5PDP is catalyzed by 3H5PP kinase (Figure 4, Step F). Phosphomevalonate kinase (EC 2.7.4.2) catalyzes the analogous transformation in the mevalonate pathway. This enzyme is encoded by *erg8* in *Saccharomyces cerevisiae* (Tsay et al., *Mol. Cell Biol.* 11:620-631 (1991)) and *mvaK2* in *Streptococcus pneumoniae*, *Staphylococcus aureus* and *Enterococcus faecalis* (Doun et al., *Protein Sci.* 14:1134-1139 (2005); Wilding et al., *J Bacteriol.* 182:4319-4327 (2000)). The *Streptococcus pneumoniae* and *Enterococcus faecalis* enzymes were cloned and characterized in *E. coli* (Pilloff et al., *J Biol. Chem.* 278:4510-4515 (2003); Doun et al., *Protein Sci.* 14:1134-1139 (2005)).

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>Erg8</i>	AAA34596.1	171479	<i>Saccharomyces cerevisiae</i>
<i>mvaK2</i>	AAG02426.1	9937366	<i>Staphylococcus aureus</i>
<i>mvaK2</i>	AAG02457.1	9937409	<i>Streptococcus pneumoniae</i>
<i>mvaK2</i>	AAG02442.1	9937388	<i>Enterococcus faecalis</i>

3H5PDP decarboxylase (Figure 4, Step G)

- 20 Butenyl 4-diphosphate is formed from the ATP-dependent decarboxylation of 3H5PDP by 3H5PDP decarboxylase (Figure 4, Step G). Although an enzyme with this activity has not been characterized to date a similar reaction is catalyzed by mevalonate diphosphate decarboxylase (EC 4.1.1.33), an enzyme participating in the mevalonate pathway for isoprenoid biosynthesis. This reaction is catalyzed by MVD1 in *Saccharomyces cerevisiae*,

MVD in *Homo sapiens* and MDD in *Staphylococcus aureus* and *Trypsonoma brucei* (Toth et al., *J Biol.Chem.* 271:7895-7898 (1996); Byres et al., *J Mol.Biol.* 371:540-553 (2007)).

Protein	GenBank ID	GI Number	Organism
<i>MVD1</i>	P32377.2	1706682	<i>Saccharomyces cerevisiae</i>
<i>MVD</i>	NP_002452.1	4505289	<i>Homo sapiens</i>
<i>MDD</i>	ABQ48418.1	147740120	<i>Staphylococcus aureus</i>
<i>MDD</i>	EAN78728.1	70833224	<i>Trypsonoma brucei</i>

Butenyl 4-diphosphate isomerase (Figure 4, Step H)

- 5 Butenyl 4-diphosphate isomerase catalyzes the reversible interconversion of 2-butenyl-4-diphosphate and butenyl-4-diphosphate. The following enzymes can naturally possess this activity or can be engineered to exhibit this activity. Useful genes include those that encode enzymes that interconvert isopenenyl diphosphate and dimethylallyl diphosphate. These include isopentenyl diphosphate isomerase enzymes from *Escherichia coli* (Rodríguez-Concepción et al., *FEBS Lett*, 473(3):328-332), *Saccharomyces cerevisiae* (Anderson et al., *J Biol Chem*, 1989, 264(32):19169-75), and *Sulfolobus shibatae* (Yamashita et al, *Eur J Biochem*, 2004, 271(6):1087-93). The reaction mechanism of isomerization, catalyzed by the Idi protein of *E. coli*, has been characterized in mechanistic detail (de Ruyck et al., *J Biol.Chem.* 281:17864-17869 (2006)). Isopentenyl diphosphate isomerase enzymes from
- 10
- 15 *Saccharomyces cerevisiae*, *Bacillus subtilis* and *Haematococcus pluvialis* have been heterologously expressed in *E. coli* (Laupitz et al., *Eur.J Biochem.* 271:2658-2669 (2004); Kajiwarara et al., *Biochem.J* 324 (Pt 2):421-426 (1997)).

Protein	GenBank ID	GI Number	Organism
Idi	NP_417365.1	16130791	<i>Escherichia coli</i>
IDI1	NP_015208.1	6325140	<i>Saccharomyces cerevisiae</i>
Idi	BAC82424.1	34327946	<i>Sulfolobus shibatae</i>
Idi	AAC32209.1	3421423	<i>Haematococcus pluvialis</i>
Idi	BAB32625.1	12862826	<i>Bacillus subtilis</i>

Butadiene synthase (Figure 4, Step I)

- 20 Butadiene synthase catalyzes the conversion of 2-butenyl-4-diphosphate to 1,3-butadiene. The enzymes described below naturally possess such activity or can be engineered to exhibit this activity. Isoprene synthase naturally catalyzes the conversion of dimethylallyl diphosphate to isoprene, but can also catalyze the synthesis of 1,3-butadiene from 2-butenyl-4-diphosphate. Isoprene synthases can be found in several organisms including *Populus alba*

(Sasaki et al., FEBS Letters, 2005, 579 (11), 2514-2518), *Pueraria montana* (Lindberg et al., *Metabolic Eng*, 12(1):70-79 (2010); Sharkey et al., *Plant Physiol.*, 137(2):700-712 (2005)), and *Populus tremula* x *Populus alba* (Miller et al., *Planta*, 213(3):483-487 (2001)).

Additional isoprene synthase enzymes are described in (Chotani et al., WO/2010/031079,

- 5 Systems Using Cell Culture for Production of Isoprene; Cervin et al., US Patent Application 20100003716, Isoprene Synthase Variants for Improved Microbial Production of Isoprene).

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>ispS</i>	BAD98243.1	63108310	<i>Populus alba</i>
<i>ispS</i>	AAQ84170.1	35187004	<i>Pueraria montana</i>
<i>ispS</i>	CAC35696.1	13539551	<i>Populus tremula</i> x <i>Populus alba</i>

3-Hydroxyglutaryl-CoA reductase (alcohol forming) (Figure 4, Step J)

- This step catalyzes the reduction of the acyl-CoA group in 3-hydroxyglutaryl-CoA to the alcohol group. Exemplary bifunctional oxidoreductases that convert an acyl-CoA to alcohol include those that transform substrates such as acetyl-CoA to ethanol (e.g., *adhE* from *E. coli* (Kessler et al., *supra*, (1991)) and butyryl-CoA to butanol (e.g. *adhE2* from *C. acetobutylicum* (Fontaine et al., *supra*, (2002)). In addition to reducing acetyl-CoA to ethanol, the enzyme encoded by *adhE* in *Leuconostoc mesenteroides* has been shown to oxidize the branched chain compound isobutyraldehyde to isobutyryl-CoA (Kazahaya et al., *supra*, (1972); Koo et al., *supra*, (2005)).

- Another exemplary enzyme can convert malonyl-CoA to 3-HP. An NADPH-dependent enzyme with this activity has been characterized in *Chloroflexus aurantiacus* where it participates in the 3-hydroxypropionate cycle (Hugler et al., *supra*, (2002); Strauss and Fuchs, *supra*, (1993)). This enzyme, with a mass of 300 kDa, is highly substrate-specific and shows little sequence similarity to other known oxidoreductases (Hugler et al., *supra*, (2002)). No enzymes in other organisms have been shown to catalyze this specific reaction; however there is bioinformatic evidence that other organisms can have similar pathways (Klatt et al., *supra*, (2007)). Enzyme candidates in other organisms including *Roseiflexus castenholzii*, *Erythrobacter* sp. *NAP1* and marine gamma proteobacterium HTCC2080 can be inferred by sequence similarity.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>adhE</i>	NP_415757.1	16129202	<i>Escherichia coli</i>
<i>adhE2</i>	AAK09379.1	12958626	<i>Clostridium acetobutylicum</i>
<i>adhE</i>	AAV66076.1	55818563	<i>Leuconostoc mesenteroides</i>

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>mcr</i>	<i>AAS20429.1</i>	<i>42561982</i>	<i>Chloroflexus aurantiacus</i>
<i>Rcas 2929</i>	YP_001433009.1	156742880	<i>Roseiflexus castenholzii</i>
<i>NAP1 02720</i>	ZP_01039179.1	85708113	<i>Erythrobacter sp. NAP1</i>
<i>MGP2080_00535</i>	ZP_01626393.1	119504313	<i>marine gamma proteobacterium HTCC2080</i>

Longer chain acyl-CoA molecules can be reduced to their corresponding alcohols by enzymes such as the jojoba (*Simmondsia chinensis*) FAR which encodes an alcohol-forming fatty acyl-CoA reductase. Its overexpression in *E. coli* resulted in FAR activity and the accumulation of fatty alcohol (Metz et al., *Plant Physiology* 122:635-644 (2000)).

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
FAR	AAD38039.1	5020215	<i>Simmondsia chinensis</i>

Another candidate for catalyzing this step is 3-hydroxy-3-methylglutaryl-CoA reductase (or HMG-CoA reductase). This enzyme reduces the CoA group in 3-hydroxy-3-methylglutaryl-CoA to an alcohol forming mevalonate. Gene candidates for this step include:

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>HMG1</i>	<i>CAA86503.1</i>	<i>587536</i>	<i>Saccharomyces cerevisiae</i>
HMG2	NP_013555	6323483	<i>Saccharomyces cerevisiae</i>
HMG1	CAA70691.1	1694976	<i>Arabidopsis thaliana</i>
<i>hmgA</i>	AAC45370.1	2130564	<i>Sulfolobus solfataricus</i>

The *hmgA* gene of *Sulfolobus solfataricus*, encoding 3-hydroxy-3-methylglutaryl-CoA reductase, has been cloned, sequenced, and expressed in *E. coli* (Bochar et al., *J Bacteriol.* 179:3632-3638 (1997)). *S. cerevisiae* also has two HMG-CoA reductases in it (Basson et al., *Proc.Natl.Acad.Sci.U.S.A* 83:5563-5567 (1986)). The gene has also been isolated from *Arabidopsis thaliana* and has been shown to complement the HMG-COA reductase activity in *S. cerevisiae* (Learned et al., *Proc.Natl.Acad.Sci.U.S.A* 86:2779-2783 (1989)).

3-oxoglutaryl-CoA reductase (aldehyde forming) (Figure 4, Step K)

Several acyl-CoA dehydrogenases are capable of reducing an acyl-CoA to its corresponding aldehyde. Thus they can naturally reduce 3-oxoglutaryl-CoA to 3,5-dioxopentanoate or can be engineered to do so. Exemplary genes that encode such enzymes were discussed in Figure 4, Step C.

3,5-dioxopentanoate reductase (ketone reducing) (Figure 4, Step L)

There exist several exemplary alcohol dehydrogenases that convert a ketone to a hydroxyl functional group. Two such enzymes from *E. coli* are encoded by malate dehydrogenase (*mdh*) and lactate dehydrogenase (*ldhA*). In addition, lactate dehydrogenase from *Ralstonia eutropha* has been shown to demonstrate high activities on 2-ketoacids of various chain lengths including lactate, 2-oxobutyrate, 2-oxopentanoate and 2-oxoglutarate (Steinbuchel et al., *Eur.J.Biochem.* 130:329-334 (1983)). Conversion of alpha-ketoadipate into alpha-hydroxyadipate can be catalyzed by 2-ketoadipate reductase, an enzyme reported to be found in rat and in human placenta (Suda et al., *Arch.Biochem.Biophys.* 176:610-620 (1976); Suda et al., *Biochem.Biophys.Res.Comm.* 77:586-591 (1977)). An additional candidate for this step is the mitochondrial 3-hydroxybutyrate dehydrogenase (*bdh*) from the human heart which has been cloned and characterized (Marks et al., *J.Biol.Chem.* 267:15459-15463 (1992)). This enzyme is a dehydrogenase that operates on a 3-hydroxyacid. Another exemplary alcohol dehydrogenase converts acetone to isopropanol as was shown in *C. beijerinckii* (Ismaiel et al., *J.Bacteriol.* 175:5097-5105 (1993)) and *T. brockii* (Lamed et al., *Biochem.J.* 195:183-190 (1981); Peretz et al., *Biochemistry.* 28:6549-6555 (1989)). Methyl ethyl ketone reductase, or alternatively, 2-butanol dehydrogenase, catalyzes the reduction of MEK to form 2-butanol. Exemplary enzymes can be found in *Rhodococcus ruber* (Kosjek et al., *Biotechnol Bioeng.* 86:55-62 (2004)) and *Pyrococcus furiosus* (van der et al., *Eur.J.Biochem.* 268:3062-3068 (2001)).

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>mdh</i>	AAC76268.1	1789632	<i>Escherichia coli</i>
<i>ldhA</i>	NP_415898.1	16129341	<i>Escherichia coli</i>
<i>ldh</i>	YP_725182.1	113866693	<i>Ralstonia eutropha</i>
<i>bdh</i>	AAA58352.1	177198	<i>Homo sapiens</i>
<i>adh</i>	AAA23199.2	60592974	<i>Clostridium beijerinckii</i> NRRL B593
<i>adh</i>	P14941.1	113443	<i>Thermoanaerobacter brockii</i> HTD4
<i>adhA</i>	AAC25556	3288810	<i>Pyrococcus furiosus</i>
<i>adh-A</i>	CAD36475	21615553	<i>Rhodococcus ruber</i>

A number of organisms can catalyze the reduction of 4-hydroxy-2-butanone to 1,3-butanediol, including those belonging to the genus *Bacillus*, *Brevibacterium*, *Candida*, and *Klebsiella* among others, as described by Matsuyama et al. US Patent 5,413,922. A mutated Rhodococcus phenylacetaldehyde reductase (Sar268) and a Leifonia alcohol dehydrogenase have also been shown to catalyze this transformation at high yields (Itoh et al., *Appl. Microbiol. Biotechnol.* 75(6):1249-1256).

Homoserine dehydrogenase (EC 1.1.1.13) catalyzes the NAD(P)H-dependent reduction of aspartate semialdehyde to homoserine. In many organisms, including *E. coli*, homoserine dehydrogenase is a bifunctional enzyme that also catalyzes the ATP-dependent conversion of aspartate to aspartyl-4-phosphate (Starnes et al., *Biochemistry* 11:677-687 (1972)). The functional domains are catalytically independent and connected by a linker region (Sibilli et al., *J Biol Chem* 256:10228-10230 (1981)) and both domains are subject to allosteric inhibition by threonine. The homoserine dehydrogenase domain of the *E. coli* enzyme, encoded by *thrA*, was separated from the aspartate kinase domain, characterized, and found to exhibit high catalytic activity and reduced inhibition by threonine (James et al., *Biochemistry* 41:3720-3725 (2002)). This can be applied to other bifunctional threonine kinases including, for example, *hom1* of *Lactobacillus plantarum* (Cahyanto et al., *Microbiology* 152:105-112 (2006)) and *Arabidopsis thaliana*. The monofunctional homoserine dehydrogenases encoded by *hom6* in *S. cerevisiae* (Jacques et al., *Biochim Biophys Acta* 1544:28-41 (2001)) and *hom2* in *Lactobacillus plantarum* (Cahyanto et al., *supra*, (2006)) have been functionally expressed and characterized in *E. coli*.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI number</u>	<u>Organism</u>
<i>thrA</i>	AAC73113.1	1786183	<i>Escherichia coli K12</i>
<i>akthr2</i>	O81852	75100442	<i>Arabidopsis thaliana</i>
<i>hom6</i>	CAA89671	1015880	<i>Saccharomyces cerevisiae</i>
<i>hom1</i>	CAD64819	28271914	<i>Lactobacillus plantarum</i>
<i>hom2</i>	CAD63186	28270285	<i>Lactobacillus plantarum</i>

3,5-dioxopentanoate reductase (aldehyde reducing) (Figure 4, Step M)

Several aldehyde reducing reductases are capable of reducing an aldehyde to its corresponding alcohol. Thus they can naturally reduce 3,5-dioxopentanoate to 5-hydroxy-3-oxopentanoate or can be engineered to do so. Exemplary genes that encode such enzymes were discussed in Figure 4, Step D.

5-hydroxy-3-oxopentanoate reductase (Figure 4, Step N)

Several ketone reducing reductases are capable of reducing a ketone to its corresponding hydroxyl group. Thus they can naturally reduce 5-hydroxy-3-oxopentanoate to 3,5-dihydroxypentanoate or can be engineered to do so. Exemplary genes that encode such enzymes were discussed in Figure 4, Step L.

3-oxo-glutaryl-CoA reductase (CoA reducing and alcohol forming) (Figure 4, Step O)

3-oxo-glutaryl-CoA reductase (CoA reducing and alcohol forming) enzymes catalyze the 2 reduction steps required to form 5-hydroxy-3-oxopentanoate from 3-oxo-glutaryl-CoA. Exemplary 2-step oxidoreductases that convert an acyl-CoA to an alcohol were provided for Figure 4, Step J. Such enzymes can naturally convert 3-oxo-glutaryl-CoA to 5-hydroxy-3-oxopentanoate or can be engineered to do so.

Throughout this application various publications have been referenced. The disclosures of these publications in their entireties, including GenBank and GI number publications, are hereby incorporated by reference in this application in order to more fully describe the state of the art to which this invention pertains. Although the invention has been described with reference to the examples provided above, it should be understood that various modifications can be made without departing from the spirit of the invention.

What is claimed is:

1. A non-naturally occurring microbial organism, comprising a microbial organism having a butadiene pathway comprising at least one exogenous nucleic acid encoding a butadiene pathway enzyme expressed in a sufficient amount to produce butadiene, said
5 butadiene pathway comprising a butadiene synthase, an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a crotonyl-CoA hydrolase, synthetase, or transferase, a crotonate reductase, a crotonyl-CoA reductase (alcohol forming), a
10 glutaconyl-CoA decarboxylase, a glutaryl-CoA dehydrogenase, an 3-aminobutyryl-CoA deaminase, a 4-hydroxybutyryl-CoA dehydratase or a crotyl alcohol diphosphokinase.
2. The non-naturally occurring microbial organism of claim 1, wherein said microbial organism comprises two exogenous nucleic acids each encoding a butadiene pathway enzyme.
- 15 3. The non-naturally occurring microbial organism of claim 1, wherein said microbial organism comprises three exogenous nucleic acids each encoding a butadiene pathway enzyme.
4. The non-naturally occurring microbial organism of claim 1, wherein said microbial organism comprises four exogenous nucleic acids each encoding a butadiene pathway
20 enzyme.
5. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate
25 kinase and a butadiene synthase.
6. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase and crotonyl-CoA reductase (alcohol forming).
- 30 7. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase,

a 3-hydroxybutyryl-CoA dehydratase, a butadiene synthase, a crotonyl-CoA reductase (alcohol forming) and a crotyl alcohol diphosphokinase.

8. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase,
5 a 3-hydroxybutyryl-CoA dehydratase, a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase and a crotonate reductase.

9. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA reductase,
10 a 3-hydroxybutyryl-CoA dehydratase, a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase, a crotonate reductase and a crotyl alcohol diphosphokinase.

10. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises an acetyl-CoA:acetyl-CoA acyltransferase, an acetoacetyl-CoA
15 reductase, a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a butadiene synthase and a crotyl alcohol diphosphokinase.

11. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a glutaconyl-CoA decarboxylase, a crotonyl-CoA reductase
20 (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase.

12. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a glutaconyl-CoA decarboxylase, a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase and crotonyl-CoA reductase (alcohol
25 forming).

13. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a glutaconyl-CoA decarboxylase, a butadiene synthase, a crotonyl-CoA reductase (alcohol forming) and a crotyl alcohol diphosphokinase.

14. The non-naturally occurring microbial organism of claim 1, wherein said
30 butadiene pathway comprises a glutaconyl-CoA decarboxylase, a crotonaldehyde reductase

(alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase, or transferase and a crotonate reductase.

15. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a glutaconyl-CoA decarboxylase, a crotonaldehyde reductase
5 (alcohol forming), a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase, a crotonate reductase and a crotyl alcohol diphosphokinase.

16. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a butadiene a
10 glutaconyl-CoA decarboxylase and a crotyl alcohol diphosphokinase.

17. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a glutaryl-CoA dehydrogenase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase.

15 18. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a glutaryl-CoA dehydrogenase, a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase and crotonyl-CoA reductase (alcohol forming).

19. The non-naturally occurring microbial organism of claim 1, wherein said
20 butadiene pathway comprises a glutaryl-CoA dehydrogenase, a butadiene synthase, a crotonyl-CoA reductase (alcohol forming) and a crotyl alcohol diphosphokinase.

20. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a glutaryl-CoA dehydrogenase, a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene
25 synthase, a crotonyl-CoA hydrolase, synthetase, or transferase and a crotonate reductase.

21. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a glutaryl-CoA dehydrogenase, a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase, a crotonate reductase and a crotyl alcohol diphosphokinase.

22. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a glutaryl-CoA dehydrogenase and a crotyl alcohol diphosphokinase.

5 23. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises an 3-aminobutyryl-CoA deaminase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase.

10 24. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises an 3-aminobutyryl-CoA deaminase, a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase and crotonyl-CoA reductase (alcohol forming).

15 25. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises an 3-aminobutyryl-CoA deaminase, a butadiene synthase, a crotonyl-CoA reductase (alcohol forming) and a crotyl alcohol diphosphokinase.

26. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises an 3-aminobutyryl-CoA deaminase, a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase and a crotonate reductase.

20 27. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises an 3-aminobutyryl-CoA deaminase, a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase, a crotonate reductase and a crotyl alcohol diphosphokinase.

25 28. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a 3-aminobutyryl-CoA deaminase and a crotyl alcohol diphosphokinase.

30 29. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a 4-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase and a butadiene synthase.

30. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a 4-hydroxybutyryl-CoA dehydratase, a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase and crotonyl-CoA reductase (alcohol forming).

5 31. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a 4-hydroxybutyryl-CoA dehydratase, a butadiene synthase, a crotonyl-CoA reductase (alcohol forming) and a crotyl alcohol diphosphokinase.

10 32. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a 4-hydroxybutyryl-CoA dehydratase, a crotonaldehyde reductase (alcohol forming), a crotyl alcohol kinase, a 2-butenyl-4-phosphate kinase, a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase and a crotonate reductase.

15 33. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a 4-hydroxybutyryl-CoA dehydratase, a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a crotonyl-CoA hydrolase, synthetase or transferase, a crotonate reductase and a crotyl alcohol diphosphokinase.

20 34. The non-naturally occurring microbial organism of claim 1, wherein said butadiene pathway comprises a 3-hydroxybutyryl-CoA dehydratase, a crotonyl-CoA reductase (aldehyde forming), a crotonaldehyde reductase (alcohol forming), a butadiene synthase, a 4-hydroxybutyryl-CoA dehydratase and a crotyl alcohol diphosphokinase.

35. The non-naturally occurring microbial organism of claim 1, wherein said at least one exogenous nucleic acid is a heterologous nucleic acid.

36. The non-naturally occurring microbial organism of claim 1, wherein said non-naturally occurring microbial organism is in a substantially anaerobic culture medium.

25 37. A non-naturally occurring microbial organism, comprising a microbial organism having a butadiene pathway comprising at least one exogenous nucleic acid encoding a butadiene pathway enzyme expressed in a sufficient amount to produce butadiene, said butadiene pathway comprising a butadiene synthase, an erythrose-4-phosphate reductase, an erythritol-4-phosphate cytidyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an
30 erythritol 2,4-cyclodiphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-

hydroxy-2-butenyl 4-diphosphate reductase, a butenyl 4-diphosphate isomerase, an erythrose-4-phosphate kinase, an erythrose reductase or an erythritol kinase.

38. The non-naturally occurring microbial organism of claim 37, wherein said microbial organism comprises two exogenous nucleic acids each encoding a butadiene pathway enzyme.

39. The non-naturally occurring microbial organism of claim 37, wherein said microbial organism comprises three exogenous nucleic acids each encoding a butadiene pathway enzyme.

40. The non-naturally occurring microbial organism of claim 37, wherein said microbial organism comprises four exogenous nucleic acids each encoding a butadiene pathway enzyme.

41. The non-naturally occurring microbial organism of claim 37, wherein said butadiene pathway comprises an erythrose-4-phosphate reductase, an erythritol-4-phosphate cytidyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an erythritol 2,4-cyclodiphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase and a butadiene synthase.

42. The non-naturally occurring microbial organism of claim 37, wherein said butadiene pathway comprises an erythrose-4-phosphate reductase, an erythritol-4-phosphate cytidyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an erythritol 2,4-cyclodiphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase, a butenyl 4-diphosphate isomerase and a butadiene synthase.

43. The non-naturally occurring microbial organism of claim 37, wherein said butadiene pathway comprises an erythritol-4-phosphate cytidyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an erythritol 2,4-cyclodiphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase, a butadiene synthase, an erythrose-4-phosphate kinase, an erythrose reductase and a erythritol kinase.

44. The non-naturally occurring microbial organism of claim 37, wherein said butadiene pathway comprises an erythritol-4-phosphate cytidyltransferase, a 4-(cytidine 5'-diphospho)-erythritol kinase, an erythritol 2,4-cyclodiphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate synthase, a 1-hydroxy-2-butenyl 4-diphosphate reductase, a butenyl 4-

diphosphate isomerase, a butadiene synthase, an erythrose-4-phosphate kinase, an erythrose reductase and an erythritol kinase.

45. The non-naturally occurring microbial organism of claim 37, wherein said at least one exogenous nucleic acid is a heterologous nucleic acid.

5 46. The non-naturally occurring microbial organism of claim 37, wherein said non-naturally occurring microbial organism is in a substantially anaerobic culture medium.

47. A non-naturally occurring microbial organism, comprising a microbial organism having a butadiene pathway comprising at least one exogenous nucleic acid encoding a butadiene pathway enzyme expressed in a sufficient amount to produce butadiene, said
10 butadiene pathway comprising a butadiene synthase, a malonyl-CoA:acetyl-CoA acyltransferase, an 3-oxoglutaryl-CoA reductase (ketone-reducing), a 3-hydroxyglutaryl-CoA reductase (aldehyde forming), a 3-hydroxy-5-oxopentanoate reductase, a 3,5-dihydroxypentanoate kinase, a 3-hydroxy-5-phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-
15 diphosphate isomerase, a 3-hydroxyglutaryl-CoA reductase (alcohol forming), an 3-oxoglutaryl-CoA reductase (aldehyde forming), a 3,5-dioxopentanoate reductase (ketone reducing), a 3,5-dioxopentanoate reductase (aldehyde reducing), a 5-hydroxy-3-oxopentanoate reductase or an 3-oxo-glutaryl-CoA reductase (CoA reducing and alcohol forming).

20 48. The non-naturally occurring microbial organism of claim 47, wherein said microbial organism comprises two exogenous nucleic acids each encoding a butadiene pathway enzyme.

49. The non-naturally occurring microbial organism of claim 47, wherein said microbial organism comprises three exogenous nucleic acids each encoding a butadiene
25 pathway enzyme.

50. The non-naturally occurring microbial organism of claim 47, wherein said microbial organism comprises four exogenous nucleic acids each encoding a butadiene pathway enzyme.

51. The non-naturally occurring microbial organism of claim 47, wherein said
30 butadiene pathway comprises a malonyl-CoA:acetyl-CoA acyltransferase, an 3-oxoglutaryl-CoA reductase (ketone-reducing), a 3-hydroxyglutaryl-CoA reductase (aldehyde forming), a

3-hydroxy-5-oxopentanoate reductase, a 3,5-dihydroxypentanoate kinase, a 3-hydroxy-5-phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase and a butadiene synthase.

52. The non-naturally occurring microbial organism of claim 47, wherein said
5 butadiene pathway comprises a malonyl-CoA:acetyl-CoA acyltransferase, a 3,5-dihydroxypentanoate kinase, a 3-hydroxy-5-phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase, a butadiene synthase, an 3-oxoglutaryl-CoA reductase (aldehyde forming), a 3,5-dioxopentanoate reductase (aldehyde reducing) and a 5-hydroxy-3-oxopentanoate reductase.
10

53. The non-naturally occurring microbial organism of claim 47, wherein said butadiene pathway comprises a malonyl-CoA:acetyl-CoA acyltransferase, a 3-hydroxy-5-oxopentanoate reductase, a 3,5-dihydroxypentanoate kinase, a 3-Hydroxy-5-phosphonatooxypentanoate kinase, a 3-Hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy
15 pentanoate decarboxylase, a butenyl 4-diphosphate isomerase, a butadiene synthase, an 3-oxoglutaryl-CoA reductase (aldehyde forming) and a 3,5-dioxopentanoate reductase (ketone reducing).

54. The non-naturally occurring microbial organism of claim 47, wherein said butadiene pathway comprises a malonyl-CoA:acetyl-CoA acyltransferase, a 3,5-
20 dihydroxypentanoate kinase, a 3-hydroxy-5-phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase, a butadiene synthase, a 5-hydroxy-3-oxopentanoate reductase and a 3-oxo-glutaryl-CoA reductase (CoA reducing and alcohol forming).

55. The non-naturally occurring microbial organism of claim 47, wherein said
25 butadiene pathway comprises a malonyl-CoA:acetyl-CoA acyltransferase, an 3-oxoglutaryl-CoA reductase (ketone-reducing), a 3,5-dihydroxypentanoate kinase, a 3-hydroxy-5-phosphonatooxypentanoate kinase, a 3-hydroxy-5-[hydroxy(phosphonooxy)phosphoryl]oxy pentanoate decarboxylase, a butenyl 4-diphosphate isomerase, a butadiene synthase and a 3-hydroxyglutaryl-CoA reductase (alcohol forming).

30 56. The non-naturally occurring microbial organism of claim 47, wherein said at least one exogenous nucleic acid is a heterologous nucleic acid.

57. The non-naturally occurring microbial organism of claim 47, wherein said non-naturally occurring microbial organism is in a substantially anaerobic culture medium.

58. A method for producing butadiene, comprising culturing the non-naturally occurring microbial organism of any one of claims 1-57 under conditions and for a sufficient
5 period of time to produce butadiene.

59. The method of claim 58, wherein said non-naturally occurring microbial organism is in a substantially anaerobic culture medium.