

(19)
(12)

(KR)
(A)

(51) 。 Int. Cl.⁷
A61K 35/78

(11)
(43)

2003-0081435
2003 10 17

(21) 10-2003-7010455

(22) 2003 08 08

2003 08 08

(86) PCT/US2002/02427

(87)

WO 2002/63982

(86) 2002 01 30

(87)

2002 08 22

(30) 60/267,428

2001 02 09

(US)

(71) , .

1947

22 (: 05301)

(72) , , .

63132

704

,
05301

222

(74)

:

(54)

가

가

(Burros, M., Tea? Turmeric? The Quest for Safer

- Barbecue, The New York Times on the Web, July 5, 2000)(, ') .
(marination) , , , , , 가 가 , C E
().
- (: P. Kalp
agam, T.C. Raghuram, T. P. Krishna and K. Krishnaswamy, 'Effect of Turmeric on Urinary Mutagens in Smokers', *Mutagenesis*, Vol. 7, no. 2, pp 107-109, 1992)(
).
[Krishnaswamy, K., and Raghuramulu, N., Bioactive Phytochemicals with Emphasis on Dietary Practices, *Indian J Med Res* 108, November 1998, pp. 167-181; Deshpande, S.S., Ingle, A.D., and Maru, G.B., Inhibitory Effects of Curcumin-Free Aqueous Turmeric Extract on Benzo[alpha]pyrene-Induced Forestomach Papillomas in Mice, *Cancer Letters*, 118 (1997) 79-85; Srimal R.C., Turmeric: A Brief Review of Medicinal Properties, *Fitoterapia*, Vol. LXVIII, No. 6, 1997, pp. 483-493; Arbiser, J.L., Klauber, N., Rohan, R., van Leeuwen, R., Huang, M.T., Fisher, C., Flynn, E., Byers, H.R., Curcumin is an In Vivo Inhibitor of Angiogenesis, *Mol Med* (June 1998), 4(6):376-83; Plummer, S.M., Holloway, K.A., Manson, M.M., Munks, R.J., Kaptein, A., Farrow, S., and Howells, L., Inhibition of Cyclo-Oxygenase 2 Expression in Colon Cells by the Chemopreventive Agent Curcumin Involves Inhibition of NF-kappaB activation Via the NIK/IKK Signaling Complex, *Oncogene* (Oct 28, 1999), 18(44):6013-20; Singhal, S.S., Awasthi, S., Pandya, U., Piper, J.T., Saini, M.K., Cheng, J.Z., and Awasthi, Y.C., The Effect of Curcumin on Glutathione-Linked Enzymes in K562 Human Leukemia Cells, *Toxicol Lett*, Sept. 20, 1999, 109(1-2):87-95; Kang, B.Y., Song, Y.J., Kim, K.M., Choe, Y. K., Hwang, S.Y., Kim, T.S., Curcumin Inhibits Th1 Cytokine Profile in CD4+ T Cells By Suppressing Interleukin-12 Production in Macrophages, *Br J Pharmacol*, Sept. 1999, 128(2):380-4].
- [: Ramirez-Bosca A., Soler A., Carrion-Gutierrez, M. A., Mira D. P., Zapata J. P., Diaz-Alperi J., Bernd A., Almagro E. Q. and Miquel, J., A Hydroalcoholic extract of *Curcuma Longa* Lowers the Apo B/Apo A Ratio Implications for Atherogenesis Prevention, *Mechanisms of Ageing and Development* 114 (2000) 207-210].
- (eugenol)
[: Sukumaran K. and Ramadasan K., Inhibition of Tobacco-Induced Mutagenesis by Eugenol and Plant Extracts, *Mutation Research* 343 (1995) 25-30].
[: Krishnaswamy, K., and Raghuraimulu, N., Bioactive Phytochemicals with Emphasis on Dietary Practices, *Indian J Med Res* 108, November 1998, pp. 167-181].
- 가 [: Lee I. P., Kim Y. H., Kang M. H., Roberts C., Shim J. S., and Roh J. K., Chemopreventive Effect of Green Tea (*Camellia sinensis*) Against Cigarette Patton Ref.: 9510.101 Smoke-Induced Mutations (SCE) in Humans, *Journal of Cellular Biochemistry Supplement* 27:68-75 (1997)].
[Klaunig J. E., Xu Y., Han C., Kamendulis L. M., Chen J., Heiser C., Gordon M. S., and Mohler III E. R. The Effect of Tea Consumption on Oxidative Stress in Smokers and Nonsmokers, *Antioxidant Effects of Tea*, pp. 249-254]
(anticlastogenicity activity) [Renqing Z., Zhou Y., Chen D., Shenben L., and Haug A., Effects of Soaking Temperature and Soaking Time During Preparation of Water Extract of Tea on Anticlastogenicity Against Environmental Tobacco Smoke in the Sister-Chromatid Exchange Assay, *Toxicology Letters* 115 (2000) 23-32]
(: Katiyar K. S., Agarwal R., Zaim M. T., and Mukhtar H., Protection Against N-nitrosodiethylamine and Benzo[alpha]pyrene-induced Forestomach and Lung Tumorigenesis in A/J Mice by Green Tea, *Carcinogenesis*, vol. 14. no. 5, pp. 849-855 (1993); Suganuma M., Okabe S., Kai Y., Sueoka N., Sueoka E., and Fujiki, H., Synergistic Effects of (-)-Epigallocatechin Gallate with (-)-Epicatechin, Sulindac, or Tarnoxifen on Cancer-Preventive Activity in the Human Lung Cancer Cell Line PC-9, *Cancer Research* 59, 44-47, January 1, 1999].
- [: 1. Emani, C. Rolando, M. Rojas, K. Alexandrov, H. Scherf, and H. Bartsch: A Rosmary Cigarette Filter May Reduce Tobacco-Linked Cancer, *Biosyntech Chemopreventive Filter Coresta*,

pp. 3-10, October 2000].

[: Zheng G., Kenney P. M., Zhang J., and Lam L. K. T., Inhibition of Benzo[alpha]pyrene-induced Tumorigenesis by Myristicin, a Volatile Aroma Constituent of Parsley Leaf Oil, Carcinogenesis, Vol. 13, no. 10, pp. 1921-1923 (1992)].

가

(A)

(B)

(D)

(C)

가

()

()

()

()

350mg

375

2000mg

1

()
4 1

(/), , , , ,

(sachets)

가

(turmeron)

()

가

가

가

가 (2),

가) (') (ii) , (i) (, - /) (E. Stahl, K.W. Quirin, D. Gerard: Dense Gases for Extraction and Refining, Springer Verlag, 1988)

25 33 1
 , 60 80 40 20 / (/)
 () , / 6
 , 1 1 3.0 6.0
 1 5.3
 , 가

가 , 1 4.4

0 / , / (/) , 60 80 40 2
 , 가 , 가

(i) 1.75:1 2.25:1 2:1
 ; (ii) 1.75:1 2.25:1 2:1
 ; (iii) 4:1 6:1 4.4:1
 ; (iv) 1.75:1 2.25:1 2:1 ; /
 (v) 1.75:1 2.25:1 2:1
 , 1 , 5.3 3 6 1

(i) , 80 (10 60 , (ii)
 , (iii))
 [R.Zhou et al., Toxicology Letters 115 (2000)
 23-32]
 [S.K. Katiyar et al., Carcinogenesis, vol. 14, no. 5, pp849-855 (1993)]

(i) , , , , , (ii) 1:1 2:1,
 1.5 1
 (, , (i) 45 % 55 % , 50 %)
 () ; (ii) 1.5 % 2.5 % , 2.0 %
 ; (iii) 1.5 % 2.5 % , 2.0 %
 ; (iv) 1.5 % 2.5 % , 2.0 % ()
 ; (v) 1.5 % 2.5 % , 2.0 % ; (vi) 1.5 %
 % 2.5 % , 2.0 % ; (vii) 35 % 45 %
 % , 40 %

43% 47%, 45%
 11%, 11% 15%
 28% 32% 6% 10% 1% 30% 8%
 43% 47%, 45%
 63% 67%, 65% 23% 2
 7%, 25% 33% 37%, 35%
 21% 21% 25%, 23%
 33% 23%
 21% 25%, 23%

(Handbook of Food Additives, CRC Press)()

가

(1)

[1]

1					
	30mg (45% - 13.5mg)	160mg (11% 17.6mg)	- N/A	N/A	
	N/A	N/A	N/A	150mg (45% - 67.5mg)	
	2.5mg (65% - 1.6mg)	N/A	5mg (10:1)	N/A	
	1.4mg (30% 8% - 0.1mg)	6.1mg (3% - 0.18mg)	N/A	N/A	
	2.5mg ((25% - 0.6mg)	N/A	5mg () (8:1)	N/A	/
	2.5mg (35% - 0.85mg)	N/A	5mg (4:1)	N/A	
	2.5mg (23%	N/A	5mg (23%	N/A	

;

9.

8 1 , 가 3.0 6.0

10.

8 1 , 가 4.4

11.

8 , (i) 1.75:1 2.25:1 , (ii) 1.75:1 2.25:1 , (iii) 4:1 6:1 , (iv) 1.75:1 2.25:1 , (v) 1:1 2:1 , (vi) 1:1 2:1 , (vii) 1.75:1 2.25:1

12.

8 2.5 % , (i) 45 % 55 % , (ii) 1.5 % , (iii) 1.5 % 2.5 % , (iv) 1.5 % 2.5 % , (v) 1.5 % 2.5 % , (vi) 1.5 % 2.5 % , (vii) 35% 45%

13.

14.

13 1 , 가 3.0 6.0

15.

13 , , ,

16.

13 11% 15% , 43% 47% ,