



US00PP27269P3

(12) **United States Plant Patent**
Clark et al.

(10) **Patent No.:** **US PP27,269 P3**

(45) **Date of Patent:** **Oct. 11, 2016**

(54) **COLEUS PLANT NAMED ‘UF12-22-1’**

(50) Latin Name: *Solenostemon scutellarioides*
Varietal Denomination: **UF12-22-1**

(71) Applicant: **Florida Foundation Seed Producers, Inc.**, Marianna, FL (US)

(72) Inventors: **David G. Clark**, Gainesville, FL (US);
Grayson M. Clark, Gainesville, FL (US)

(73) Assignee: **Florida Foundation Seed Producers, Inc.**, Marianna, FL (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/544,562**

(22) Filed: **Jan. 21, 2015**

(65) **Prior Publication Data**
US 2016/0212904 P1 Jul. 21, 2016

(51) **Int. Cl.**
A01H 5/12 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./373**

(58) **Field of Classification Search**

USPC Plt./373
See application file for complete search history.

(56) **References Cited**

PUBLICATIONS

BallFloraPlant website. FlameThrower™ Chipotle Coleus (http://www.ballfloraplant.com/series_info.aspx?phid=018006557030884). Accessed Jan. 14, 2016. 1 page.*
Rosy Dawn Gardens—Chipolte Coleus. (<http://www.rosydawngardens.com/coleus-a-to-d/chipolte>). Accessed Jan. 14, 2016. 1 page.*
Nguyen et al., “Genetics of growth habit and development of new coleus (*Solenostemon scutellarioides* (L.) Codd) varieties with trailing habit and bright color,” *J Heredity* 99:573-580, 2008.

* cited by examiner

Primary Examiner — Susan McCormick Ewoldt

Assistant Examiner — Karen Redden

(74) *Attorney, Agent, or Firm* — Dentons US LLP

(57) **ABSTRACT**

‘UF12-22-1’ is a new coleus plant distinguished by having consistent bright orange-bronze leaves with distinct red-magenta venation, and a vigorous but compact growth habit, as disclosed.

3 Drawing Sheets

1

Latin name of the genus and species of the plant claimed: *Solenostemon scutellarioides*.
Cultivar denomination: ‘UF12-22-1’.

BACKGROUND OF THE INVENTION

The invention relates to a new and distinct cultivar of coleus plant named ‘UF12-22-1’. ‘UF12-22-1’ originated from an open pollination conducted in May-November 2011 in Gainesville, Fla. between the female coleus plant ‘UF11-3-11’ (unpatented) and an unknown male coleus plant. A single seedling was chosen in May 2012 for further asexual propagation in Gainesville, Fla. (see FIG. 1 for pedigree).

‘UF12-22-1’ has been reproduced asexually for over one year through vegetative cuttings and has been found to retain its distinctive characteristics through successive asexual propagations. ‘UF12-22-1’ was first propagated asexually by meristem tip cuttings in May, 2012 in Gainesville, Fla., and has remained true-to-type since that time.

‘UF12-22-1’ has not been made publicly available more than one year prior to the filing date of this application.

When compared to the female parent ‘UF11-3-11’, ‘UF12-22-1’ has large, orange-bronze leaves and a compact branched habit, while ‘UF11-3-11’ has slightly smaller leaves colored deep bronze with a more vigorous and upright plant habit.

When ‘UF12-22-1’ is compared to the commercial cultivar ‘Keystone Kopper’ ‘UF09-8-87’ (commercial, unpatented), both plants have orange-bronze foliage color

2

and purple stems, although ‘UF12-22-1’ leaves are larger, and more brightly colored in both sun and shade conditions. ‘UF12-22-1’ also has a more vigorous growth habit than ‘Keystone Kopper’ ‘UF09-8-87’.

SUMMARY OF THE INVENTION

The following are the most outstanding and distinguishing characteristics of ‘UF12-22-1’ when grown under normal horticultural practices in Gainesville, Fla. ‘UF12-22-1’ has a combination of a novel, vigorous, compact upright growth habit, excellent heat tolerance, and consistent bright orange-bronze leaves that are significantly different than other coleus plants. It has superior stability in foliage color in both sun and shade conditions, maintaining stable color in all conditions. It has a vigorous but compact upright growth habit with excellent lateral branching when grown as a stock plant, thus providing ample vegetative propagules for producers. This plant is desirable for long-season performance in the landscape.

BRIEF DESCRIPTION OF THE DRAWINGS

This new coleus plant is illustrated by the accompanying photographs, which show the plant’s form and foliage. The colors shown are as true as can be reasonably obtained by

conventional photographic procedures. FIGS. 2 and 3 were taken from twelve-week-old plants pinched once at four weeks after stick and grown in March-May, 2014 in a glass greenhouse in Gainesville, Fla.

FIG. 1—shows the pedigree of the claimed plant.

FIG. 2—shows the growth habit, form, and foliage of the claimed plant.

FIG. 3—shows a close-up of the foliage.

DETAILED BOTANICAL DESCRIPTION

The following detailed description sets forth the distinctive characteristics of 'UF12-22-1'. The detailed description was obtained using twelve-week-old plants grown in March-May, 2014 in a glass greenhouse in Gainesville, Fla. The plants were pinched 4 weeks after cuttings were rooted, then grown in 1-gallon pots for approximately 8 weeks. Color references are to The R.H.S. Colour Chart of The Royal Horticultural Society of London (R.H.S.), 2007 5th Edition. Classification:

Family.—Lamiaceae.

Botanical.—*Solenostemon scutellarioides*.

Common name.—Coleus.

Cultivar name.—'UF12-22-1'.

Plant description:

Form.—Spreading.

Habit.—Upright.

Height (from top of soil).—40-45 cm.

Width (horizontal plant diameter).—65-70 cm.

Propagation:

Type cuttings.—Vegetative meristems having at least 1 node.

Time to initiate roots.—3-4 days.

Time to produce a rooted cutting.—7-10 days.

Root habit.—Fibrous.

Root description.—Callus forms in 2 to 3 days, roots initiate in 3-4 days and become a highly branched cutting in 7-10 days.

Branches:

Quantity per plant.—6 to 8 main branches per plant with numerous side branches, pinched once.

Branch color.—RHS N186C.

Texture.—Smooth.

Pubescence.—Not present.

Stem description.—Square-shaped stem, 1.5 cm in diameter at the soil line.

Branch diameter.—0.6-0.7 cm at the base of a 40-cm long branch.

Branch length.—40-42 cm.

Internode length.—6-7 cm.

Anthocyanin.—N/A.

Leaves:

Quantity of leaves per branch.—14 to 16. Arrangement: Opposite.

Fragrance.—Not fragrant.

Shape.—Ovate.

Length.—14-16 cm.

Width.—10-12 cm.

Apex.—Broadly acuminate.

Base.—Attenuate.

Margin.—Crenate.

Leaf texture (both surfaces).—Slightly pubescent upper surface; smooth lower surface.

Pubescence color (both surfaces).—Non-descript with naked eye.

Venation color.—Upper surface: RHS 79A. Lower surface: RHS N77D.

Venation pattern.—Upper surface: Reticulate. Lower surface: Reticulate.

Color.—Immature leaf: Upper surface: RHS 175A. Lower surface: RHS 59A.

Color.—Mature leaf: Upper surface: RHS 175A. Lower surface: RHS N79A.

Petiole length.—3-4 cm.

Petiole diameter.—0.3-0.4 cm.

Petiole color.—RHS N77A.

Petiole texture.—Smooth, no pubescence.

Flowers and seeds: Flowers and seeds have not been observed.

Fruit/seed set: Fruit/seed not observed.

Disease and insect resistance: Disease and insect resistance is typical of the species, thus no claims are made of any superior disease or insect resistance with this cultivar. The most common insect pests observed on this plant in Gainesville, Fla. have been long-tailed or citrus mealybugs (*Pseudococcus* sp.), which occur on older stock plant material held in the greenhouse for over 3-4 months. Impatiens Necrotic Spot Virus (*Bunyaviridae*) has also been observed in plants confined in greenhouses with mixed crops (peppers) infected with Western flower thrips (*Frankliniella occidentalis*). The most common pathogen of this species in the U.S. is downy mildew (*Peronospora lamii*). This pathogen has been observed in stock materials grown closely together in cooler growing seasons. What is claimed is:

1. A new and distinct *Solenostemon scutellarioides* plant called 'UF12-22-1' as shown and described herein.

* * * * *

UF12-22-1



UF11-3-11



UF09-8-88



UF08-29-1



UF07-8-20



UF06-4-56



'Elfers'

FIG. 1



FIG. 2




FIG. 3