

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2016252145 B2**

(54) Title
Compositions and methods for altering flowering and plant architecture to improve yield potential

(51) International Patent Classification(s)
A01H 1/00 (2006.01) **A01H 5/00** (2006.01)
A01H 1/02 (2006.01) **C12N 15/82** (2006.01)

(21) Application No: **2016252145** (22) Date of Filing: **2016.04.18**

(87) WIPO No: **WO16/172051**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/233,019	2015.09.25	US
62/150,142	2015.04.20	US

(43) Publication Date: **2016.10.27**

(44) Accepted Journal Date: **2022.01.13**

(71) Applicant(s)
Monsanto Technology LLC

(72) Inventor(s)
Brower-Toland, Brent;Caldo, Rico A.;Dai, Shunhong;Gabbert, Karen;Goldshmidt, Alexander;Howell, Miya D.;Karunanandaa, Balasulojini;Manjunath, Sivalinganna;Mcdill, Bradley W.;Ovadya, Daniel J.;Preuss, Sasha;Rice, Elena A.;Savidge, Beth;Sharma, Vijay K.

(74) Agent / Attorney
Davies Collison Cave Pty Ltd, Level 14 255 Elizabeth Street, Sydney, NSW, 2000, AU

(56) Related Art
WO 2013192081 A1
CN 102146124 A
US 6225530 B1
US 20050223428 A1
Nan H et al., "GmFT2a and GmFT5a redundantly and differentially regulate flowering through interaction with and upregulation of the bZIP transcription factor GmFDL19 in soybean", PLoS ONE



(51) International Patent Classification:

A01H 1/00 (2006.01) *A01H 5/00* (2006.01)
A01H 1/02 (2006.01)

(21) International Application Number:

PCT/US2016/028130

(22) International Filing Date:

18 April 2016 (18.04.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/150,142 20 April 2015 (20.04.2015) US
62/233,019 25 September 2015 (25.09.2015) US

(71) Applicant: **MONSANTO TECHNOLOGY LLC**
[US/US]; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US).(72) Inventors: **BROWER-TOLAND, Brent**; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US). **CALDO, Rico A.**; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US). **DAI, Shunhong**; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US). **GABBERT, Karen**; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US). **GOLDSHMIDT, Alexan-**

der; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US). **HOWELL, Miya D.**; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US). **KAR-UNANANDAA, Balasulojini**; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US). **MAN-JUNATH, Sivalinganna**; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US). **MCDILL, Bradley W.**; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US). **OVADYA, Daniel J.**; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US). **PREUSS, Sasha**; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US). **RICE, Elena A.**; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US). **SAVIDGE, Beth**; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US). **SHARMA, Vijay K.**; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US).

(74) Agents: **MARSH, David R.** et al.; 601 Massachusetts Ave., NW, Washington, District of Columbia 20001-3743 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,

[Continued on next page]

(54) Title: COMPOSITIONS AND METHODS FOR ALTERING FLOWERING AND PLANT ARCHITECTURE TO IMPROVE YIELD POTENTIAL

	SEQUENCE	1	2	3	4	5	6	7	8	9	10
1	Gm.FT2a	-	90.6 (481)	69.5 (369)	71 (377)	66.7 (354)	60.6 (322)	68.7 (365)	64.2 (341)	73.4 (390)	66.5 (353)
2	Gm.FT2b	90.6 (481)	-	67.6 (359)	69.9 (371)	64.2 (341)	58.4 (310)	65.9 (350)	62 (329)	70.4 (374)	65.9 (350)
3	Nt.FT	70.3 (369)	68.4 (359)	-	76.6 (402)	67.4 (354)	58.5 (307)	67 (352)	62.9 (330)	73.7 (387)	67.8 (356)
4	Le.FT	70.6 (377)	69.5 (371)	75.3 (402)	-	67.4 (380)	60.1 (321)	67.4 (360)	64.6 (345)	73 (390)	66.9 (357)
5	Gm.FT5A	68.2 (354)	65.7 (341)	68.2 (354)	69.4 (360)	-	61.1 (317)	65.9 (342)	62.6 (325)	70.7 (367)	65.7 (341)
6	Zm.ZCN8	61 (322)	58.7 (310)	58.1 (307)	60.8 (321)	60 (317)	-	57 (301)	59.3 (313)	61.4 (324)	56.3 (297)
7	At.FT	69.1 (365)	66.3 (350)	66.7 (352)	68.2 (360)	64.8 (342)	57 (301)	-	65.9 (348)	72.5 (383)	82.2 (434)
8	Os.HD3A	63.1 (341)	60.9 (329)	61.1 (330)	63.9 (345)	60.2 (325)	58 (313)	64.4 (348)	-	71.1 (384)	63.9 (345)
9	Pt.FT	74.3 (390)	71.2 (374)	73.7 (387)	74.3 (390)	69.9 (387)	61.7 (324)	73 (383)	73.1 (384)	-	71.8 (377)
10	At.TSF	66.9 (353)	66.3 (350)	67.4 (356)	67.6 (357)	64.6 (341)	56.3 (297)	82.2 (434)	65.3 (345)	71.4 (377)	-

FIG. 1A

(57) Abstract: The present invention provides recombinant DNA constructs, vectors and molecules comprising a polynucleotide sequence encoding a florigenic FT protein operably linked to a vegetative stage promoter, which may also be a meristem-preferred or meristem-specific promoter. Transgenic plants, plant cells and tissues, and plant parts are further provided comprising a polynucleotide sequence encoding a florigenic FT protein. Transgenic plants comprising a florigenic FT transgene may produce more bolls, siliques, fruits, nuts, or pods per node on the transgenic plant, particularly on the main stem of the plant, relative to a control or wild type plant. Methods are further provided for introducing a florigenic FT transgene into a plant, and planting transgenic FT plants in the field including at higher densities. Transgenic plants of the present invention may thus provide greater yield potential than wild type plants and may be planted at a higher density due to their altered plant architecture.



KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE,

SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(88) Date of publication of the international search report:

1 December 2016

COMPOSITIONS AND METHODS FOR ALTERING FLOWERING AND PLANT ARCHITECTURE TO IMPROVE YIELD POTENTIAL

CROSS -REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority benefit to U.S. Provisional Patent Application Nos. 62/150,142 and 62/233,019, filed on April 20, 2015 and September 25, 2015, respectively, which are incorporated herein by reference in their entirety.

INCORPORATION OF SEQUENCE LISTING

[0002] A computer readable form of a sequence listing is filed with this application by electronic submission and is incorporated into this application by reference in its entirety. The sequence listing is contained in the file named 60760_anniversary_ST25.txt, which is 61,079 bytes in size (measured in operating system MS Windows) and created on April 1, 2016.

FIELD OF THE INVENTION

[0003] The present invention relates to compositions and methods for modulating floral development and vegetative growth by genetic modification of crop plants to increase yield.

BACKGROUND

[0004] The transition from vegetative growth to flowering is a crucial process during plant development that is necessary for the production of grain yield in crop plants. There are four major pathways controlling flowering time in land plants that respond to environmental or developmental cues, including photoperiodism (i.e., day length), vernalization (i.e., response to winter cold), and plant hormones (e.g., gibberellins or GA), in addition to the autonomous (environmentally independent) pathways. Except for the GA and autonomous pathways, regulation of flowering in plants generally involves two central regulators of flowering time, *CONSTANS* (CO) and *FLOWERING LOCUS C* (FLC). The *FLC* gene is a floral repressor that regulates flowering in response to vernalization, whereas the *CO* gene is a floral activator that responds to photoperiod conditions. Under inductive photoperiodic conditions, CO activity in source leaves increases expression of *FLOWERING LOCUS T* (FT), which translocates to the meristem to trigger expression of downstream floral activating genes, including *LEAFY* (LFY), *APETALA1* (AP1) and *SUPPRESSOR OF OVEREXPRESSION OF*

CO 1 (SOC1). Other genes, such as *FLOWERING LOCUS C* (FLC) and *TERMINAL FLOWER 1* (TFL1), act to inhibit the expression or activity of these genes.

[0005] Except for day length neutral plants, most flowering plants respond to daily photoperiodic cycles and are classified as either short day (SD) or long day (LD) plants based on the photoperiod conditions required to induce flowering. The photoperiod refers to the relative length or duration of light and dark periods within a 24-hour cycle. In general, long day plants tend to flower when the day length exceeds a photoperiod threshold (e.g., as the days are getting longer in the spring), whereas short day plants tend to flower when the day length falls below a photoperiod threshold (e.g., as the days are getting shorter after the summer solstice). In other words, SD plants flower as the days are getting shorter, while LD plants flower as the days are getting longer. Soybean is an example of a short day (SD) plant in which flowering is induced when plants are exposed to shorter daylight conditions.

[0006] Plant growers are always looking for new methods to manipulate the yield of a plant, especially to enhance the seed yield of agronomically important crops. Thus, there is a continuing need in the art for improved compositions and methods for increasing yields of various crop plants. It is presently proposed that improved crop yields may be achieved by enhancing agronomic traits related to flowering and reproductive development.

SUMMARY

[0007] According to a first aspect of the present invention, a recombinant DNA construct is provided comprising a polynucleotide sequence encoding a florigenic FT protein operably linked to a vegetative stage promoter. The florigenic FT protein encoded by the polynucleotide sequence may comprise an amino acid sequence having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or least 99% identity to a sequence selected from the group consisting of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, and 20, or a functional fragment thereof. The polynucleotide sequence may also be at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or least 99% identity to a sequence selected from the group consisting of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, and 19. The vegetative stage promoter may further be a meristem-preferred or meristem-specific promoter. DNA

molecules and vectors comprising such a recombinant DNA construct are further provided.

[0008] According to a second aspect of the present invention, transgenic plants, plant cells, plant tissues and plant parts are further provided comprising an insertion of the recombinant DNA construct of the present invention into the genome of such plants, cells, tissues, and plant parts. A transgenic plant of the present invention may be homozygous or hemizygous for an insertion of the recombinant DNA construct. A transgenic plant may be a short day plant and/or a dicotyledonous plant. Depending on the plant species, transgenic plants of the present invention may produce more bolls, siliques, fruits, nuts, or pods per node of the transgenic plant, relative to a control or wild type plant not having the recombinant DNA construct. Transgenic plants of the present invention may also produce more flowers and/or floral racemes per node relative to a control or wild type plant not having the recombinant DNA construct.

[0009] According to a third aspect of the present invention, methods for producing a transgenic plant having improved yield-related traits or phenotypes are provided comprising (a) transforming at least one cell of an explant with a recombinant DNA construct comprising a polynucleotide sequence encoding a florigenic FT protein operably linked to a vegetative stage promoter; and (b) regenerating or developing the transgenic plant from the transformed explant. Such methods may further comprise (c) selecting a transgenic plant having one or more of the following traits or phenotypes: earlier flowering, longer reproductive or flowering duration, increased number of flowers per node, increased number of floral racemes per node, increased number of pods, bolls, siliques, fruits, or nuts per node, and increased number of seeds per node, as compared to a control plant not having the recombinant DNA construct.

[0010] According a fourth aspect of the present invention, methods are provided for planting a transgenic crop plant of the present invention at a normal or higher density in the field. According to some embodiments, methods are provided comprising: planting a transgenic crop plant at a higher density in the field, wherein the transgenic crop plant is transformed with a recombinant DNA construct comprising a polynucleotide sequence encoding a florigenic FT protein operably linked to a vegetative stage promoter. According to some of these embodiments, the vegetative stage promoter may be a meristem-preferred or meristem-specific promoter. For soybean, a higher density of about 150,000 to 250,000 seeds of the transgenic soybean plant may be planted per acre. For cotton, a higher density of about

48,000 to 60,000 seeds of the transgenic cotton plant may be planted per acre. For canola, a higher density of about 450,000 to 680,000 seeds of the transgenic canola plant may be planted per acre.

BRIEF DESCRIPTION OF THE DRAWINGS

[0011] **FIG. 1A** provides a matrix table showing a comparison of nucleotide sequences for each combination of the various FT genes including their percent identity.

[0012] **FIG. 1B** provides a matrix table showing a comparison of protein sequences for each combination of the various FT proteins including their percent identity.

[0013] **FIG. 1C** provides a CLUSTAL 2.0.9 multiple sequence alignment of various FT proteins.

[0014] **FIG. 2** shows the total FT transcript levels in soybean leaf and apex tissues collected at 1, 3 and 5 days after either a short day or long day light treatment.

[0015] **FIG. 3** and **FIG. 4** show the expression pattern of the *pAt.Erecta* promoter by monitoring GUS activity during early soybean development. **FIG. 3** is a set of black and white images of stained tissues, and the images in **FIG. 4** correspond to **FIG. 3** but are filtered for blue GUS staining. **FIGS. 3A to 3C** and **4A to 4C** show expression in a 3-day-old germinating seedling; **FIGS. 3D to 3I** and **4D to 4I** show expression in a 10-day-old vegetative shoot (grown in 14 hour light / 10 hour dark photoperiod); **FIGS. 3J to 3L** and **4J to 4L** show expression in a 16-day-old reproductive shoot; and **FIGS. 3M to 3O** and **4M to 4O** show expression in the 30d old mature and immature leaves of the reproductive shoot. Bars are 100µm.

[0016] **FIG. 5** and **FIG. 6** show the GUS expression pattern with the *pAT.Erecta* promoter during R1 and floral stages of development (35-40 days after germination). **FIG. 5** is a set of black and white images of stained tissues, and the images in **FIG. 6** correspond to **FIG. 5** but are filtered for blue GUS staining. **FIGS. 5A and 6A** show expression in the inflorescence stems or pedicels (arrows), and **FIGS. 5B and 6B** show expression in the floral peduncle (arrows). Expression is also shown in the vasculature and parenchyma cells (**FIGS. 5C and 6C**), in stamen filaments (**FIGS. 5D and 6D**; arrow), and un-pollinated ovules (**FIGS. 5E, 5F, 6E and 6F**; arrows). Bars are 1 mm.

[0017] **FIG. 7** shows section imaging of the shoot apical meristem (SAM) from wild type

versus GmFT2a-expressing transgenic plants at 7 days after planting using scanning electron microscopy (eSEM) analysis.

[0018] FIG. 8 shows scanning electron microscopy (eSEM) micrographs of an axillary inflorescence primordia from a wild type plant (collected at 27 days after planting), in comparison to an axillary inflorescence primordia from a transgenic event expressing Gm.FT2a (collected at 9 days after planting).

[0019] FIG. 9 shows the effects of Gm.FT2a expression driven by the *At.Erecta* promoter in soybean. FIG. 9A depicts a null segregant showing normal axillary buds, whereas FIG. 9B and FIG. 9C (corresponding to plants homozygous or hemizygous for the *Gm.FT2a* transgene, respectively) each show early flowering and increased pods per node relative to the null segregant.

[0020] FIG. 10 shows a whole plant image of a wild type null segregant next to plants hemizygous and homozygous for the *Gm.FT2a* transgene as indicated.

[0021] FIG. 11 shows images of the main stem of plants that are homozygous or hemizygous for the *pAt.Erecta-Gm.FT2a* transgene in comparison to a null segregant as indicated.

[0022] FIG. 12A shows whole plant images of a wild type null segregant and a plant homozygous for the *pEr:Zm.ZCN8* transgene as indicated.

[0023] FIG. 12B shows close up images of pods on the mainstem of a wild type null segregant and a plant homozygous for the *pEr:Zm.ZCN8* transgene as indicated.

[0024] FIG. 13A shows whole plant images of a wild type null segregant and a plant homozygous for the *pEr:Nt.FT-like* transgene as indicated.

[0025] FIG. 13B shows close up images of pods on the mainstem of a wild type null segregant and a plant homozygous for the *pEr:Nt.FT-like* transgene as indicated.

[0026] FIG. 14 shows whole plant images of a wild type null segregant and a plant homozygous for the *pEr:Gm.FT2b* transgene as indicated.

[0027] FIG. 15 shows whole plant images of a wild type null segregant and a plant homozygous for the *pEr:Le.SFT* transgene as indicated.

[0028] FIG. 16 shows whole plant images of a wild type null segregant and a plant homozygous for the *pEr:FT5a* transgene as indicated.

DETAILED DESCRIPTION

[0029] The goal of improving yield is common to all crops across agriculture. The present invention includes methods and compositions for improving yield in flowering (angiosperm) or seed-bearing plants by modification of traits associated with flowering time, reproductive development, and vegetative growth to improve one or more flowering and/or yield-related traits or phenotypes, such as the number of flowers, seeds and/or pods per plant, and/or the number of flowers, seeds and/or pods per node (and/or per main stem) of the plant. Without being bound by any theory, compositions and methods of the present invention may operate to improve yield of a plant by increasing the number of floral meristems, increasing synchronization of lateral meristem release, and/or extending the time period for pod or seed development in the plant.

[0030] Previously, it was discovered that growing short day plants, such as soybean, under long day conditions (e.g., about 14-16 hours of light per day) and then briefly subjecting those plants to short day growing conditions (e.g., about 9-11 hours of light per day for about 3-21 days) before returning the plants to long day (non-inductive) growing conditions, produced plants having increased numbers of pods/seeds per plant (and pods/seeds per node and/or per branch). See, e.g., U.S. Patent No. 8,935,880 and U.S. Patent Application Publication No. 2014/0259905, the entire contents and disclosures of which are incorporated herein by reference.

[0031] As described further below, this short day induction phenotype in soybean was used to identify genes having altered expression in these plants through transcriptional profiling. These studies identified several genes with altered expression in these treated soybean plants including an endogenous FT gene, *Gm.FT2a*, having increased expression in response to the short day induction treatment. Thus, it is presently proposed that transgenic FT expression as described herein may be used in place of short day induction to increase seed yield, alter reproductive traits or phenotypes in plants, or both. According to an aspect of the present invention, ectopic or transgenic expression of a *Gm.FT2a* gene or other FT sequence, or a functional fragment, homolog or ortholog thereof, in a flowering or seed-bearing plant may be used to increase seed yield and/or alter one or more reproductive phenotypes or traits, which may involve an increase in the number of pods/seeds per plant (and/or the number of pods/seeds per node or main stem of the plant). As explained further below and depending on the particular plant species, these yield-related or reproductive phenotypes or traits may

also apply to other botanical structures analogous to pods of leguminous plants, such as bolls, siliques, fruits, nuts, tubers, etc. Thus, a plant ectopically expressing a FT sequence may instead have an increased number of bolls, siliques, fruits, nuts, tubers, etc., per node(s), main stem, and/or branch(es) of the plant, and/or an increased number of bolls, siliques, fruits, nuts, tubers, etc., per plant.

[0032] According to embodiments of the present invention, a recombinant DNA molecule comprising an FT transgene is provided, which may be used in transformation to generate a transgenic plant expressing the FT transgene. The polynucleotide coding sequence of the FT transgene may include *Gm.FT2a* (SEQ ID NO: 1), or any polynucleotide sequence encoding the Gm.FT2a protein (SEQ ID NO: 2). The polynucleotide coding sequence of an FT transgene may also correspond to other FT genes in soybean or other plants. For example, other polynucleotide FT coding sequences from soybean that may be used according to present embodiments include: *Gm.FT5a* (SEQ ID NO: 3) or a polynucleotide encoding a Gm.FT5a protein (SEQ ID NO: 4), or *Gm.FT2b* (SEQ ID NO: 5) or a polynucleotide encoding a Gm.FT2b protein (SEQ ID NO: 6). In addition, examples of polynucleotide FT coding sequences from other plant species that may be used include: *Zm.ZCN8* (SEQ ID NO: 7) from maize or a polynucleotide encoding Zm.ZCN8 protein (SEQ ID NO: 8), *Nt.FT-like* or *Nt.FT4* (SEQ ID NO: 9) from tobacco or a polynucleotide encoding Nt.FT-like or Nt.FT4 protein (SEQ ID NO: 10), *Le.FT* or *SFT* (SEQ ID NO: 11) from tomato or a polynucleotide encoding Le.FT or SFT protein (SEQ ID NO: 12), *At.FT* (SEQ ID NO: 13) from *Arabidopsis* or a polynucleotide encoding At.FT protein (SEQ ID NO: 14), *At.TSF* (SEQ ID NO: 15) from *Arabidopsis* or a polynucleotide encoding At.TSF protein (SEQ ID NO: 16), *Os.HD3a* (SEQ ID NO: 17) from rice or a polynucleotide encoding Os.HD3a protein (SEQ ID NO: 18), or *Pt.FT* (SEQ ID NO: 19) from *Populus trichocarpa* or a polynucleotide encoding Pt.FT protein (SEQ ID NO: 20).

[0033] Polynucleotide coding sequences for FT transgenes encoding additional FT proteins from other species having known amino acid sequences may also be used according to embodiments of the present invention, which may, for example, include the following: Md.FT1 and Md.FT2 from apple (*Malus domestica*); Hv.FT2 and Hv.FT3 from barley (*Hordeum vulgare*); Cs.FTL3 from Chrysanthemum; Ls.FT from lettuce (*Lactuca sativa*); Pn.FT1 and Pn.FT2 from Lombardy poplar (*Populus nigra*); Pa.FT from London plane tree (*Platanus acerifolia*); Dl.FT1 from Longan (*Dimocarpus longan*); Ps.FTa1, Ps.FTa2, Ps.FTb1, Ps.FTb2, and Ps.FTc from pea (*Pisum sativum*); Ac.FT from pineapple (*Ananas*

comosus); Cm-FTL1 and Cm-FTL2 from pumpkin (*Cucurbita maxima*); Ro.FT from rose; Cg.FT from spring orchid (*Cymbidium*); Fv.FT1 from strawberry (*Fragaria vesca*); Bv.FT2 from sugar beet (*Beta Vulgaris*); Ha.FT4 from sunflower (*Helianthus annuus*); and Ta.FT or TaFT1 from wheat (*Triticum aestivum*). See, e.g., Wickland, DP et al., “The *Flowering Locus T/Terminal Flower 1* Gene Family: Functional Evolution and Molecular Mechanisms”, *Molecular Plant* 8: 983-997 (2015), the content and disclosure of which is incorporated herein by reference.

[0034] Unless otherwise stated, nucleic acid or polynucleotide sequences described herein are provided (left-to-right) in the 5' to 3' direction, and amino acid or protein sequences are provided (left-to-right) in the N-terminus to C-terminus direction. Additional known or later discovered FT genes and proteins from these or other species may also be used according to embodiments of the present invention. These FT genes may be known or inferred from their nucleotide and/or protein sequences, which may be determined by visual inspection or by use of a computer-based searching and identification tool or software (and database) based on a comparison algorithm with known FT sequences, structural domains, etc., and according to any known sequence alignment technique, such as BLAST, FASTA, etc.

[0035] According to embodiments of the present invention, an FT transgene of a recombinant DNA molecule may comprise a polynucleotide sequence that (when optimally aligned) is at least 60% identical, at least 65% identical, at least 70% identical, at least 75% identical, at least 80% identical, at least 85% identical, or at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or least 99% identical to one or more of the polynucleotide FT coding sequences listed above (e.g., SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, or 19), or to any other known florigenic FT coding sequence. Sequence identity percentages among polynucleotide sequences of the above listed full length FT genes are presented in FIG. 1A. Each cell in the table in FIG. 1A shows the percentage identity for the FT gene in the corresponding row (query sequence) as compared to the FT gene in the corresponding column (subject sequence) divided by the total length of the query sequence, and the number in parenthesis is the total number of identical bases between the query and subject sequences. As shown in this figure, the percentage identities among polynucleotide sequences for these sampled FT genes range from about 60% to about 90% identity. Thus, a polynucleotide sequence that is within one or more of these sequence identity ranges or has a higher sequence identity may be used according to embodiments of the present invention to induce flowering, increase yield, and/or alter one or more

reproductive traits of a plant. Similar polynucleotide coding sequences for FT may be designed or chosen based on known FT protein sequences, conserved amino acid residues and domains, the degeneracy of the genetic code, and any known codon optimizations for the particular plant species to be transformed.

[0036] As described below, an FT transgene comprising any one of the above coding sequences may further include one or more expression and/or regulatory element(s), such as leader(s), intron(s), etc. Indeed, an FT transgene may comprise a genomic sequence encoding an FT protein or amino acid sequence, or a fragment or portion thereof.

[0037] According to embodiments of the present invention, an FT transgene of a recombinant DNA molecule may comprise a polynucleotide sequence encoding an amino acid or protein sequence that (when optimally aligned) is at least 60% identical, at least 65% identical, at least 70% identical, at least 75% identical, at least 80% identical, at least 85% identical, or at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to any one or more of the FT protein or amino acid sequences listed above (e.g., SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, or 20) or any other known florigenic FT protein sequence, or a functional fragment thereof. Such a “functional fragment” is defined as a protein having a polypeptide sequence that is identical or highly similar to a full-length FT protein but lacking one or more amino acid residues, portions, protein domains, etc., of the full-length FT protein, as long as the fragment remains active in causing one or more of the phenotypic effects or changes similar to the full-length protein when transgenically expressed in a plant. Sequence identity percentages among the above listed full length FT proteins are presented in FIG. 1B. The percentages are calculated as described above in reference to FIG. 1A based on the number of identical amino acid residues (in parenthesis) between the query and subject FT protein sequences. Multiple sequence alignment of these FT proteins is also shown in FIG. 1C. As can be seen from these figures, the percentage identity among protein sequences for these FT genes ranges from about 60% to about 90% identity. Thus, a polynucleotide sequence encoding an amino acid or protein sequence that is within one or more of these sequence identity ranges or has a higher sequence identity may be used according to embodiments of the present invention to induce flowering, increase seed yield, and/or alter one or more reproductive traits of a plant. These FT protein sequences encoded by a polynucleotide sequence of the present invention may be designed or chosen based on known FT protein sequences and their conserved amino acid residues and domains.

[0038] As used herein, the term “sequence identity” refers to the extent to which two optimally aligned DNA sequences are identical. Various pair-wise or multiple sequence alignment algorithms and programs are known in the art, such as ClustalW, etc., that may be used to compare the sequence identity or similarity between two or more sequences, such as between two or more FT genes or protein sequences, or an FT gene (nucleotide) or protein sequence and another nucleotide or protein sequence. For example, the percentage identity of one sequence (query) to another sequence (subject) may be calculated as described above in reference to FIGS. 1A and 1B (i.e., with the sequences optimally aligned, divide the number of identical bases or residues by the total number of bases or residues for the query sequence, and multiply by 100%). Although other alignment and comparison methods are known in the art, the alignment and percent identity between two sequences (including the percent identity ranges described above) may be as determined by the ClustalW algorithm, see, e.g., Chenna R. et al., “Multiple sequence alignment with the Clustal series of programs,” *Nucleic Acids Research* 31: 3497-3500 (2003); Thompson JD et al., “Clustal W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice,” *Nucleic Acids Research* 22: 4673-4680 (1994); and Larkin MA et al., “Clustal W and Clustal X version 2.0,” *Bioinformatics* 23: 2947-48 (2007), the entire contents and disclosures of which are incorporated herein by reference. For purposes of the present invention, when two sequences are optimally aligned (with allowance for gaps in their alignment), then the “percent identity” for the query sequence is calculated as described above in reference to FIGS. 1A and 1B – i.e., Percent Identity = (Number of Identical Positions between query and subject sequences / Total Number of Positions in the Query Sequence) x 100%, with each sequence consisting of a series of positions (nucleotide bases or amino acid residues).

[0039] An FT protein sequence encoded by a polynucleotide sequence or transgene of the present invention may also be designed or chosen to have one or more amino acid substitution(s) known to be chemically and/or structurally conservative (e.g., replacing one amino acid with another having similar chemical or physical properties, such as hydrophobicity, polarity, charge, steric effect, acid/base chemistry, similar side chain group, such as hydroxyl, sulfhydryl, amino, etc.) to avoid or minimize structural changes to the protein that might affect its function. For example, valine is often a conservative substitute for alanine, and threonine may be a conservative substitute for serine. Additional examples of conservative amino acid substitutions in proteins include: valine/leucine, valine/isoleucine,

phenylalanine/tyrosine, lysine/arginine, aspartic acid/glutamic acid, and asparagine/glutamine. An FT protein sequence encoded by a polynucleotide sequence or transgene of the present invention may also include proteins that differ in one or more amino acids from those of a known FT protein or similar sequence as a result of deletion(s) and/or insertion(s) involving one or more amino acids.

[0040] Various FT genes and proteins from different plant species may be identified and considered FT homologs or orthologs for use in the present invention if they have a similar nucleic acid and/or protein sequence and share conserved amino acids and/or structural domain(s) with at least one known FT gene or protein. As used herein, the term “homolog” in reference to a FT gene or protein is intended to collectively include any homologs, analogs, orthologs, paralogs, etc., of the FT gene or protein, and the term “homologous” in reference to polynucleotide or protein sequences is intended to mean similar or identical sequences. Such a FT homolog may also be defined as having the same or similar biological function as known FT genes (e.g., acting to similarly influence flowering and/or other reproductive or yield-related traits or phenotypes when ectopically expressed in a plant).

[0041] Sequence analysis and alignment of FT protein sequences from different plant species further reveals a number of conserved amino acid residues and at least one conserved structural domain. By subjecting the various aligned FT protein sequences (see, e.g., FIGS. 1B and 1C) to a protein domain identification tool using a Pfam database (e.g., Pfam version 26.0, released November 2011, or later version), these FT proteins have been found to contain and share at least a portion of a putative phosphatidyl ethanolamine-binding protein (PEBP) domain (Pfam domain name: PBP_N; Accession number: PF01161). See, e.g., Banfield, MJ et al., “The structure of *Antirrhinum* centroradialis protein (CEN) suggests a role as a kinase inhibitor,” *Journal of Mol Biol.*, 297(5): 1159-1170 (2000), the entire contents and disclosure of which are incorporated herein by reference. This PEBP domain was found to correspond, for example, to amino acids 28 through 162 of the full length Gm.FT2a protein (See Table 5 below). Thus, FT proteins encompassed by embodiments of the present invention may include those identified or characterized as having or containing at least a PEBP domain (Accession number: PF01161) according to Pfam analysis. Accordingly, the present invention may further include a polynucleotide sequence(s) encoding an FT protein having at least a PEBP domain. As known in the art, the “Pfam” database is a large collection of multiple sequence alignments and hidden Markov models covering many common protein families and containing information about various protein

families and their domain structure(s). By identifying a putative Pfam structural domain(s) for a given protein sequence, the classification and function of the protein may be inferred or determined. See, e.g., Finn, RD et al., “The Pfam protein families database,” *Nucleic Acids Research (Database Issue)*, 42:D222-D230 (2014), the entire contents and disclosure of which are incorporated herein by reference.

[0042] Embodiments of the present invention may further include polynucleotide sequence(s) encoding inductive or florigenic FT proteins. An FT protein encoded by a polynucleotide sequence may be “inductive” or “florigenic” if the FT protein, when ectopically expressed in a plant, is able to cause earlier flowering and/or an increased prolificacy in the number of flowers, pods, and/or seeds per one or more node(s) of the plant. Without being bound by any theory, such increased prolificacy in the number of flowers, pods, and/or seeds per node(s) of the plant may result from an increase in the number of meristems at those node(s) that undergo a vegetative to reproductive transition and produce flowers. Such an increased prolificacy at each node due to ectopic expression of a “florigenic” FT may be due to increased synchronization of the release and floral development of early racemes and lateral meristems at each node. Although a “florigenic” FT protein may function to induce earlier flowering when ectopically expressed in a plant, a transgenically expressed “florigenic” FT protein may increase the number of flowers, pods, and/or seeds per node(s) of a plant through one or more pathways or mechanisms that are independent of, or in addition to, any florigenic effects related to flowering time and/or reproductive duration.

[0043] Florigenic FT-like genes from various plant species are generally well conserved. However, many proteins in the PEBP family have amino acid sequences that are substantially similar to florigenic FT proteins but do not behave as florigens. For example, *Terminal Flower* (TFL) genes from various plant species have similar protein sequences to florigenic FT genes but actually delay flowering. Recent work has identified specific amino acid residues that are generally not shared between florigenic FT proteins and other PEBP proteins, such as TFLs, and substitutions at many of these positions have been shown to convert florigenic FT proteins into floral repressor proteins. See, e.g., Ho and Weigel, *Plant Cell* 26: 552-564 (2014); Danilevskaya et al., *Plant Physiology* 146(1): 250-264 (2008); Harig et al., *Plant Journal* 72: 908-921 (2012); Hsu et al., *Plant Cell* 18: 1846-1861 (2006); Kojima et al., *Plant Cell Physiology* 43(10): 1096-1105 (2002); Kong et al., *Plant Physiology* 154: 1220-1231 (2010); Molinero-Rosales et al., *Planta* 218: 427-434 (2004); Zhai et al., *PLoS ONE*, 9(2): e89030 (2014), and Wickland DP et al. (2015), *supra*, the entire contents

and disclosures of which are incorporated herein by reference. Thus, these amino acid residues can serve as signatures to further define and distinguish florigenic FT proteins of the present invention.

[0044] According to embodiments of the present invention, an “inductive” or “florigenic” FT protein may be further defined or characterized as comprising one or more of the following amino acid residue(s) (amino acid positions refer to corresponding positions of the full-length *Arabidopsis* FT protein, SEQ ID NO: 14): a proline at amino acid position 21 (P21); an arginine or lysine at amino acid position 44 (R44 or K44); a glycine at amino acid position 57 (G57); a glutamic acid or an aspartic acid at amino acid position 59 (E59 or D59); a tyrosine at amino acid position 85 (Y85); a leucine at amino acid position 128 (L128); a glycine at amino acid position 129 (G129); a threonine at amino acid position 132 (T132); an alanine at amino acid position 135 (A135); a tryptophan at amino acid position 138 (W138); a glutamic acid or an aspartic acid at amino acid position 146 (E146 or D146); and/or a cysteine at amino acid position 164 (C164). Corresponding amino acid positions of other FT proteins can be determined by alignment with the *Arabidopsis* FT sequence (see, e.g., FIG. 1C). One skilled in the art would be able to identify corresponding amino acid positions of other FT proteins based on their sequence alignment. Several of these key residues fall within an external loop domain of FT-like proteins, defined as amino acids 128 through 145 of the *Arabidopsis* full-length FT sequence (SEQ ID NO: 14) and corresponding sequences of other FT proteins (see, e.g., FIG. 1C). Thus, polynucleotides of the present invention may encode florigenic FT proteins having one or more of these conserved amino acid residues.

[0045] Florigenic FT proteins of the present invention may also have one or more other amino acids at one or more of the above identified residue positions. For example, in reference to the above amino acid positions of the *Arabidopsis* FT (At.FT) protein sequence (SEQ ID NO: 14), a florigenic FT protein may alternatively have one or more of the following amino acids: an alanine (in place of proline) at the position corresponding to position 21 of the At.FT protein sequence (P21A), or possibly other small, nonpolar residues, such as glycine or valine, at this position; a histidine (in place of lysine or arginine) at the amino acid position corresponding to position 44 of the At.FT protein sequence, or possibly other polar amino acids at this position; an alanine or cysteine (in place of glycine) at the amino acid position corresponding to position 57 of the At.FT protein sequence, or possibly other small, nonpolar residues, such proline or valine, at this position; an asparagine or serine (in place of glutamic acid or aspartic acid) at the amino acid position corresponding to

position 59 of the At.FT protein sequence, or possibly other small, polar residues, such as glutamine, cysteine, or threonine, at this position; a variety of polar and nonpolar uncharged residues (other than tyrosine) at the amino acid position corresponding to position 85 of the At.FT protein sequence; a nonpolar or hydrophobic uncharged residue (other than leucine), such as isoleucine, valine, or methionine, at the amino acid position corresponding to position 128 of the At.FT protein sequence; a variety of smaller nonpolar and uncharged residues (other than glycine), such as alanine, valine, leucine, isoleucine, methionine, etc., at the amino acid position corresponding to position 129 of the At.FT protein sequence, although some polar and charged residues may be tolerated at this position; a polar uncharged residue (other than threonine) at the amino acid position corresponding to position 132 of the At.FT protein sequence; a variety of amino acids other than proline, such as threonine, at the amino acid position corresponding to position 135 of the At.FT protein sequence; a variety of other bulky nonpolar or hydrophobic amino acids (in place of tryptophan), such as methionine or phenylalanine, at the amino acid position corresponding to position 138 of the At.FT protein sequence; a variety of other polar or non-positively charged amino acids, such as asparagine or serine, at the amino acid position corresponding to position 146 of the At.FT protein sequence; and/or a variety of other polar or nonpolar amino acids (in place of cysteine, such as isoleucine, at the amino acid position corresponding to position 164 of the At.FT protein sequence. One skilled in the art would be able to identify corresponding amino acid positions and substitutions of FT proteins based on their sequence alignment to the *Arabidopsis* FT protein sequence. In addition, other chemically conservative amino acid substitutions are also contemplated within the scope of florigenic FT proteins based on the knowledge of one skilled in the art of protein biochemistry. Accordingly, polynucleotides of the present invention may further encode florigenic FT proteins having one or more conservative amino acid substitutions. Indeed, florigenic FT proteins encoded by polynucleotides of the present invention include native sequences and artificial sequences containing one or more conservative amino acid substitutions, as well as functional fragments thereof.

[0046] Florigenic FT proteins of the present invention may also be defined as excluding (i.e., not having) one or more amino acid substitutions that may be characteristic of, or associated with, TFL or other non-florigenic or anti-florigenic proteins. For example, in reference to the amino acid positions of the *Arabidopsis* FT protein sequence (SEQ ID NO: 14), a florigenic FT protein may exclude one or more of the following amino acids: a phenylalanine or serine

at the position corresponding to position 21 of the At.FT protein sequence (e.g., in place of proline or alanine); a phenylalanine at the position corresponding to position 44 of the At.FT protein sequence (e.g., in place of arginine or lysine); a histidine, glutamic acid, or aspartic acid at the position corresponding to position 57 of the At.FT protein sequence (e.g., in place of glycine); a glycine or alanine at the position corresponding to position 59 of the At.FT protein sequence (e.g., in place of glutamic acid or aspartic acid); a histidine at the position corresponding to position 85 of the At.FT protein sequence (e.g., in place of tyrosine); a lysine, arginine, alanine, or methionine at the position corresponding to position 109 of the At.FT protein sequence; a lysine or arginine at the position corresponding to position 128 of the At.FT protein sequence (e.g., in place of leucine); a glutamine or asparagine at the position corresponding to position 129 of the At.FT protein sequence (e.g., in place of glycine); a valine or cysteine at the position corresponding to position 132 of the At.FT protein sequence (e.g., in place of threonine); a lysine, arginine, or alanine at the position corresponding to position 134 of the At.FT protein sequence (e.g., in place of tyrosine); a proline at the position corresponding to position 135 of the At.FT protein sequence (e.g., in place of alanine or threonine); a serine, aspartic acid, glutamic acid, alanine, lysine, or arginine at the position corresponding to position 138 of the At.FT protein sequence (e.g., in place of tryptophan or methionine); a lysine or arginine at the position corresponding to position 140 of the At.FT protein sequence; a lysine or arginine at the position corresponding to position 146 of the At.FT protein sequence (e.g., in place of acidic or uncharged polar residues); a lysine or arginine at the position corresponding to position 152 of the At.FT protein sequence; and/or an alanine at the position corresponding to position 164 of the At.FT protein sequence (e.g., in place of cysteine or isoleucine). One skilled in the art would be able to identify corresponding amino acid positions and substitutions of other FT proteins based on their sequence alignment. Accordingly, embodiments of the present invention may exclude polynucleotides that encode FT-like proteins having one or more of the above amino acid substitutions associated with TFL or other anti-florigens. However, an FT protein may tolerate one or some of these amino acid substitutions while still maintaining florigenic activity.

[0047] A florigenic FT protein of the present invention may also be defined as being similar to a known FT protein in addition to having one or more of the above signature amino acid residues. For example, a florigenic protein may be defined as having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least

92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or least 99% identity to a sequence selected from the group consisting of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, and 20, or a functional fragment thereof, in addition to one or more of the following signature residues: a tyrosine or other uncharged polar or nonpolar residue (e.g., alanine, tryptophan, methionine, leucine, threonine, cysteine, serine, or asparagine) at the amino acid position corresponding to position 85 of the At.FT protein sequence; a leucine or other nonpolar or hydrophobic residue (e.g., isoleucine, valine, or methionine) at the amino acid position corresponding to position 128 of the At.FT protein sequence; and/or a tryptophan or other large nonpolar or hydrophobic residue (e.g., methionine or phenylalanine) at the amino acid position corresponding to position 138 of the At.FT protein sequence. Such a florigenic FT protein may be further defined as having additional signature amino acid residue(s), such as one or more of the following: a glycine or other small nonpolar and uncharged residue (e.g., alanine, valine, leucine, isoleucine, or methionine) at the amino acid position corresponding to position 129 of the At.FT protein sequence; and/or a threonine at the amino acid position corresponding to position 132 of the At.FT protein sequence.

[0048] A florigenic FT protein of the present invention may also be defined as having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or least 99% identity to a sequence selected from the group consisting of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, and 20, or a functional fragment thereof, but not having (i.e., excluding) one or more non-florigenic or anti-florigenic residues, such as one or more of the following: a histidine at the amino acid position corresponding to position 85 of the At.FT protein sequence; a lysine or arginine at the amino acid position corresponding to position 128 of the At.FT protein sequence; and/or a serine, aspartic acid, glutamic acid, lysine or arginine at the amino acid position corresponding to position 138 of the At.FT protein sequence. Such a florigenic FT protein may be further defined as not having (i.e., excluding) one or more additional residues, such as one or more of the following: a glutamine or asparagine at the amino acid position corresponding to position 129 of the At.FT protein sequence; and/or a valine or cysteine at the amino acid position corresponding to position 132 of the At.FT protein sequence.

[0049] *Flowering Locus T* (FT) genes play a key role in higher plants and function to integrate floral pathways. FT proteins have been shown to function as a mobile signal or florigen transported from leaves to the shoot apical apex where it triggers initiation of

reproductive development in diverse species. See, e.g., Jaeger, K.E. et al., “Interlocking feedback loops govern the dynamic behavior of the floral transition in *Arabidopsis*,” *The Plant Cell*, 25:820-833 (2013); Corbesier, L et al., “FT protein movement contributes to long distance signaling in floral induction of *Arabidopsis*,” *Science* 316: 1030-1033 (2007); Jaeger, KE et al., “FT protein acts as a long range signal in *Arabidopsis*,” *Curr Biol* 17: 1050-1054 (2007); and Amasino, R.M. et al., “The Timing of Flowering,” *Plant Physiology*, 154(2):516-520 (2010), the entire contents and disclosures of which are incorporated herein by reference. In *Arabidopsis*, FT protein binds to 14-3-3 and *Flowering Locus D* (FD) proteins in the meristem to form a flowering complex triggering activation of key floral meristem identity genes, such as *APETATAL1* (AP1) and *SOC1* at the shoot apex. See, e.g., Taoka, K. et al., “14-3-3 protein act as intracellular receptors for rice Hd3a florigen,” *Nature* 476:332-335 (2011). The *TERMINAL FLOWER 1* (TFL1) gene is a key repressor of FT targets that maintains the center of the shoot apical meristem (SAM) in a vegetative state. TFL1 acts by repressing the *LEAFY* (LFY) and *API* genes. Thus, the relative concentrations of FT and TFL1 in the target tissues act competitively to control the timing of the reproductive transition of meristems from a vegetative state that may terminate further vegetative growth. . See, e.g., Abe, M et al., *Science* 309:1052-1055 (2005); and McGarry, RC et al., *Plant Science* 188/189: 71-81 (2012).

[0050] FT genes have been identified from many diverse species, and ectopic FT expression has been reported to induce early flowering. See, e.g., Kong, F. et al., “Two Coordinately Regulated Homologs of *Flowering Locus T* Are Involved in the Control of Photoperiodic Flowering in Soybean,” *Plant Physiology* 154: 1220-1231 (2010); Turck, F. et al., “Regulation and identity of florigen: *Flowering Locus T* moves center stage,” *Ann Rev Plant Biol* 59: 573-594 (2008); Blackman, BK et al., “The role of recently derived FT paralogs in sunflower domestication,” *Curr Biol* 20: 629-635 (2010); Lifschitz, E. et al., “The tomato FT orthologs triggers systemic signals that regulate growth and flowering and substitute for diverse environmental stimuli,” *PNAS* 103: 6398-6403 (2006); Trankner, C. et al., “Over-expression of an FT-homologous gene of apple induces early flowering in annual and perennial plants,” *Planta* 232: 1309-1324 (2010); and Xiang, L. et al., “Functional analysis of *Flowering Locus T* orthologs from spring orchid (*Cymbidium goeringii* Rehb. f.) that regulates the vegetative to reproductive transition,” *Plant Cell & Biochem* 58: 98-105 (2012), the entire contents and disclosures of which are incorporated herein by reference. However, prior studies with expression of FT transgenes used constitutive or tissue specific promoters

that produced either very severe phenotypes, non-cell autonomous (systemic) phenotypes, or autonomous leaf specific phenotypes with plants or seedlings flowering earlier than controls and terminating at early stages of development. Given these findings, ectopic FT expression was generally not seen as a viable approach to increasing yield in plants by inducing flowers or altering flowering time.

[0051] According to embodiments of the present invention, a recombinant DNA molecule or polynucleotide is provided comprising a polynucleotide coding sequence encoding a FT protein that is operably linked to one or more promoter(s) and/or other regulatory element(s) that are operable in a plant cell to control or bias the timing and/or location of FT expression when transformed into a plant. As commonly understood in the art, the term “promoter” may generally refer to a DNA sequence that contains an RNA polymerase binding site, transcription start site, and/or TATA box and assists or promotes the transcription and expression of an associated transcribable polynucleotide sequence and/or gene (or transgene). A promoter may be synthetically produced, varied or derived from a known or naturally occurring promoter sequence or other promoter sequence (e.g., as provided herein). A promoter may also include a chimeric promoter comprising a combination of two or more heterologous sequences. A promoter of the present invention may thus include variants of promoter sequences that are similar in composition, but not identical to, other promoter sequence(s) known or provided herein. As used herein, the term “operably linked” refers to a functional linkage between a promoter or other regulatory element and an associated transcribable polynucleotide sequence or coding sequence of a gene (or transgene), such that the promoter, etc., operates to initiate, assist, affect, cause, and/or promote the transcription and expression of the associated coding or transcribable sequence, at least in particular tissue(s), developmental stage(s), and/or under certain condition(s).

[0052] A promoter may be classified according to a variety of criteria relating to the pattern of expression of a coding sequence or gene (including a transgene) operably linked to the promoter, such as constitutive, developmental, tissue-specific, inducible, etc. Promoters that initiate transcription in all or most tissues of the plant are referred to as “constitutive” promoters. Promoters that initiate transcription during certain periods or stages of development are referred to as “developmental” promoters. Promoters whose expression is enhanced in certain tissues of the plant relative to other plant tissues are referred to as “tissue-enhanced” or “tissue-preferred” promoters. Thus, a “tissue-preferred” promoter causes relatively higher or preferential expression in a specific tissue(s) of the plant, but with lower

levels of expression in other tissue(s) of the plant. Promoters that express within a specific tissue(s) of the plant, with little or no expression in other plant tissues, are referred to as “tissue-specific” promoters. A promoter that expresses in a certain cell type of the plant is referred to as a “cell type specific” promoter. An “inducible” promoter is a promoter that initiates transcription in response to an environmental stimulus such as cold, drought or light, or other stimuli, such as wounding or chemical application. A promoter may also be classified in terms of its origin, such as being heterologous, homologous, chimeric, synthetic, etc. A “heterologous” promoter is a promoter sequence having a different origin relative to its associated transcribable sequence, coding sequence, or gene (or transgene), and/or not naturally occurring in the plant species to be transformed. The term “heterologous” may refer more broadly to a combination of two or more DNA molecules or sequences when such a combination is not normally found in nature.

[0053] According to embodiments of the present invention, a recombinant DNA molecule or polynucleotide is provided comprising a florigenic FT transgene or coding sequence operably linked to a promoter that functions in a plant, which may be introduced or transformed into a plant to cause the plant to have an altered flowering, reproductive and/or yield-related trait or phenotype. Since FT proteins are believed to operate in the meristems of a plant, including the apical and/or axillary meristems, to trigger floral transitioning of those meristems and induce flowering, embodiments of the present invention provide a recombinant DNA molecule comprising a FT transgene or coding sequence operably linked to a “vegetative stage” promoter to cause, when introduced or transformed into a plant, expression of the FT transgene earlier in the development of the plant (i.e., during the vegetative growth phase of the plant) to produce an increased level of FT in target tissues than would otherwise occur in a wild type plant at the same stage of development. Timing FT transgene expression during the vegetative stage(s) of development may be important for affecting one or more reproductive, flowering and/or yield-related traits or phenotypes by providing a timely inductive signal for the production of an increased number of floral meristems and successful flowers at one or more node(s) of the plant. Without being bound by any theory, vegetative stage expression of an FT transgene in a plant may operate to synchronize and/or increase early flowering at one or more node(s) to produce more flowers per node of the plant. The promoters described below as a part of the present invention provide options for timing FT expression.

[0054] As used herein, a “vegetative stage” promoter includes any promoter that initiates, causes, drives, etc., transcription or expression of its associated gene (or transgene) during one or more vegetative stage(s) of plant development, such as during one or more of V_e, V_c, V₁, V₂, V₃, V₄, etc., and/or any or all later vegetative stages of development (e.g., up to V_n stage). Such a “vegetative stage” promoter may be further defined as a “vegetative stage preferred” promoter that initiates, causes, drives, etc., transcription or expression of its associated gene (or transgene) at least preferably or mostly, if not exclusively, during one or more vegetative stage(s) of plant development (as opposed to reproductive stages). However, a “vegetative stage” and a “vegetative stage preferred” promoter may each also permit, allow, cause, etc., transcription or expression of its associated gene (or transgene) during reproductive phase(s) of development in one or more cells or tissues of the plant. The features and characteristics of these vegetative stages for a given plant species are known in the art. For dicot plants, vegetative morphological features and characteristics of the plant during vegetative stages of development may include cotyledon form, vegetative meristems (apical, lateral/axillary, and root), leaf arrangement, leaf shape, leaf margin, leaf venation, petioles, stipules, ochrea, hypocotyl, and roots. According to embodiments of the present invention, a “vegetative stage” promoter may also be further defined by the particular vegetative stage during which observable or pronounced transcription or expression of its associated gene (or transgene) is first initiated. For example, a vegetative stage promoter may be a V_c stage promoter, a V₁ stage promoter, a V₂ stage promoter, a V₃ stage promoter, etc. As such, a “V_c stage” promoter is defined as a vegetative stage promoter that first initiates transcription of its associated gene (or transgene) during the V_c stage of plant development, a “V₁ stage” promoter is defined as a vegetative stage promoter that first initiates transcription of its associated gene (or transgene) during the V₁ stage of plant development, a “V₂ stage” promoter is defined as a vegetative stage promoter that first initiates transcription of its associated gene (or transgene) during the V₂ stage of plant development, and so on, although expression of the associated gene (or transgene) may be present continuously or discontinuously in one or more tissues during later vegetative stage(s) of development. One skilled in the art would be able to determine the timing of expression of a given gene (or transgene) during plant development using various molecular assays and techniques known in the art.

[0055] According to embodiments of the present invention, a “vegetative stage” promoter may include a constitutive, tissue-preferred, or tissue-specific promoter. For example, a

vegetative stage promoter may drive FT expression in one or more plant tissue(s), such as in one or more of the root(s), stem(s), leaf/leaves, meristem(s), etc., during a vegetative stage(s) of plant development. However, such a vegetative stage promoter may preferably drive expression of its associated FT transgene or coding sequence in one or more meristem(s) of the plant. According to many embodiments, a “vegetative stage” promoter may preferably be a “meristem-specific” or “meristem-preferred” promoter to cause expression of the FT transgene or coding sequence in meristematic tissue. FT proteins are known to operate in the meristems of a plant to help trigger the transition from vegetative to reproductive growth after translocation of the FT protein from the leaves. In contrast, embodiments of the present invention provide selective expression of an FT transgene directly in the meristem of a plant to induce flowering and cause the plant to adopt an altered reproductive and/or yield-related trait or phenotype. Thus, according to embodiments of the present invention, a recombinant DNA molecule is provided comprising an FT transgene or coding sequence operably linked to a “meristem-specific” or “meristem-preferred” promoter that drives expression of the FT transgene at least preferentially in one or more meristematic tissues of a plant when transformed into the plant. As used herein, “meristem-preferred promoter” refers to promoters that preferentially cause expression of an associated gene or transgene in at least one meristematic tissue of a plant relative to other plant tissues, whereas a “meristem-specific promoter” refers to promoters that cause expression of an associated gene or transgene exclusively (or almost exclusively) in at least one meristem of a plant.

[0056] Recent work using artificial early “short day” light treatments during vegetative stages of development revealed that flowering time could be altered in a way that alters one or more yield-related traits or phenotypes (e.g., by causing an increased number of pods or seeds per node on a plant) and that the effect of these treatments was dosage-dependent with the number of flowers, seeds and/or pods per plant (and/or per node of the plant) depending on (i) the duration of the short day exposure (i.e., floral induction signal dosage) and (ii) the length of the post-short day photoperiods under long day conditions (i.e., the dosage or length of the vegetative growth inducing signal after the short day induction signal). See, e.g., U.S. Patent No. 8,935,880 and U.S. Patent Application Publication No. 2014/0259905, introduced above. Soybean plants experiencing a lower or less prolonged early short day induction (eSDI) treatment (prior to returning to long day growing conditions) had more flowers, pods and seeds per plant with more normal plant height and maturity, whereas soybean plants exposed to a greater or more prolonged eSDI treatment produced shorter, earlier-terminating

plants with fewer pods and seeds per plant (albeit perhaps with an increased number of pods and/or seeds per node).

[0057] Without being bound by any theory, it is proposed that an early florigenic signal (e.g., short days for soybean and other SD plants) triggers an early vegetative to reproductive transition of plants and even termination of a subset of its primary meristems. However, by returning those plants to non-inductive growth conditions (e.g., long days for SD plants) after the initial SD signal, the remaining meristematic reserves of the plant may be preserved and reproductive and/or flowering duration may be extended or maintained, thus allowing for the successful development of a greater number of productive flowers, pods and/or seeds per node (and/or per plant) during the reproductive phase. With early floral induction, a greater overlap may also be created between reproductive development and vegetative growth of the plant, which may further promote or coincide with an extended reproductive and/or flowering duration. For purposes of the present invention, “reproductive duration” refers to the length of time from the initiation of flowering until the end of seed/pod development and/or filling, whereas “flowering duration” or “duration of flowering” refers the length of time from the appearance of the first open flower until the last open flower closes. By returning to non-inductive growth conditions after early floral induction, more abundant resources may be available and directed toward the production of an increased number of successful (i.e., non-aborting) flowers, pods and/or seeds per plant, unlike normal floral development in short day plants, which may coincide with declining plant resources due to termination of meristematic growth and maturation of the plant.

[0058] However, in addition to early flowering, a floral induction signal (e.g., early short day conditions) also causes early termination of the plant. Therefore, it is proposed that an optimal dosage and timing of the floral induction signal may be needed to maximize yield by balancing (i) the early vegetative to reproductive transition and/or synchronization of flowering with the early floral induction signal (leading to potential yield gains at each node of the plant) against (ii) earlier growth termination (leading to smaller plants with fewer internodes, less branching, and fewer nodes and/or flowers per plant). It is believed that lower dosages of a floral induction signal may be sufficient to induce flowering while lessening or minimizing earlier termination of the plant, such that larger plants are produced with increased numbers of flowers, pods and/or seeds per node (and/or per plant). On the other hand, higher dosages of a floral induction signal may cause early termination of the plant (in addition to early flowering) to produce smaller plants with relatively fewer numbers

of flowers, pods and/or seeds per plant due to the smaller plant size with fewer internodes and/or branches per plant, despite having perhaps a greater number of flowers, pods and/or seeds per node (and/or per plant) relative to wild-type or control plants under normal growth conditions. As stated above, these effects of ectopic FT expression may also include an increased number of bolls, siliques, fruits, nuts, tubers, etc., per node (and/or per plant), depending on the particular plant species.

[0059] As mentioned above, the short day induction phenotype in soybean was used to screen for genes having altered expression in those plants through transcriptional profiling. These studies led to the identification of an endogenous FT gene, *Gm.FT2a*, having increased expression in response to the short day induction treatment. Accordingly, it is presently proposed that expression of a florigenic FT transgene can be used as a floral induction signal to cause early flowering and increased flowers, pods and/or seeds per node (and/or per plant) relative to a wild type or control plant not having the FT transgene. According to embodiments of the present invention, appropriate control of the timing, location and dosage of florigenic FT expression during vegetative stages of development can be used to induce flowering and produce plants having increased flowers, pods and/or seeds per node relative to a wild type or control plant not having the FT transgene. Rather than attempting to transiently express FT and recapitulate the timing of the eSDI treatments, it is proposed that FT could be weakly expressed in the vegetative meristem to provide the early floral induction signal. Accordingly, a promoter from the *Erecta* gene (*pErecta* or *pEr*) having weak meristematic expression during vegetative stages of development was selected for initial testing with a *Gm.FT2a* transgene. However, given that prior studies showed that constitutive FT expression produced plants having a severe, early termination phenotype, and further that the site of action for FT produced peripherally and translocated from the leaves is in the meristem, it was possible that direct meristematic expression of FT could produce even more potent and severe phenotypes (and/or non-viable plants) relative to constitutive FT expression.

[0060] The effects of *Gm.FT2a* overexpression were immediately seen in R₀ transformed soybean plants, which had early flowering, reduced seed yield (e.g., only about 8 seeds/plant), and very early termination, suggesting that the balance between floral induction and floral repression/vegetative growth was strongly in favor of flowering and early termination. However, enough R₁ seed was salvaged from these plants to allow for additional experiments to be performed. It was proposed that growing the R₁ soybean seed

under long day (floral repressive) photoperiod conditions in the greenhouse might delay the early flowering and termination phenotypes observed in the R_0 plants. Given the theorized dosage response, it was further proposed that segregating FT2a homozygous, hemizygous and null soybean plants could be tested together in the greenhouse to evaluate the dosage response resulting from FT overexpression. In these experiments (as described further below), it was observed that segregating plants did have different phenotypes: null plants were similar to wild-type plants in terms of plant architecture and pods per node (and per plant), while homozygous plants terminated early with a severe dwarf phenotype (although possibly with an increased number of pods per node). However, hemizygous plants were almost as large as null or wild-type plants but exhibited the hyper-flowering phenotype with an increased number of pods per node (and/or per plant). These findings show that vegetative stage and/or meristematic expression of a florigenic FT transgene may be used to produce a high yielding plant (similar to the eSDI treatment), and that the effect of FT expression is dosage-dependent since soybean plants hemizygous for the FT2a transgene under the control of a weak meristematic promoter displayed the high yield phenotype of increased pods per node without the more severe early termination and short plant height phenotypes observed with homozygous FT2a plants when grown under long day (vegetative) conditions.

[0061] Interestingly, however, increased numbers of pods per node was often observed independently of reproductive (R1-R7) duration under greenhouse conditions and was not perfectly correlated with early flowering among the FT transgenes tested, although it is possible that the duration of flowering may still be prolonged in some cases (at least during one or more reproductive stages) even if the total reproductive duration is not significantly changed relative to control plants. Without being bound by any theory, it is believed that increased numbers of pods per node in transgenic FT plants may result at least in part from an increase in the number of inflorescence and floral meristems induced from vegetative shoot apical and axillary meristems at each of the affected node(s), which may give rise to a greater number of flowers and/or released floral racemes at those node(s). Such an increase in the number of floral meristems induced at each node of the plant in response to FT overexpression may operate through one or more mechanisms or pathways, which may be independent of flowering time and/or reproductive duration. However, meristematic changes may be microscopic at first, and thus not observed to cause “early flowering” at such stage by simple visual inspection even though reproductive changes to the meristem may have already begun to occur.

[0062] Without being bound by any theory, early vegetative FT expression may cause more reproductive meristems to form and develop earlier than normal at one or more node(s) of the transgenic plant. These reproductive meristems may then allow or cause a greater number of floral racemes to form and elongate with flowers at each node. On the other hand, it is further theorized that later expression of FT during reproductive stages may function to repress further floral development at each node. Thus, later developing flowers within the respective raceme may become terminated, and thus more of the plant's resources may be directed to the earlier developing flowers within the raceme to more effectively produce full-sized pods. Accordingly, it is contemplated that by (i) forming a greater number of inflorescence and floral meristems at each node by early FT induction, and then (ii) directing more plant resources to the earlier, more developed flowers within each of the racemes by termination of the later-developing flowers, increased synchronization of floral development may occur with a greater number of mature pods being formed per node of the plant.

[0063] According to embodiments of the present invention, a recombinant DNA molecule is provided comprising an FT coding sequence operatively linked to a vegetative stage promoter, which may also be a meristem-preferred and/or meristem-specific promoter. For example, the promoter may include the *pAt.Erecta* promoter from *Arabidopsis* (SEQ ID NO: 21), or a functional fragment or portion thereof. Two examples of a truncated portion of the *pAt.Erecta* promoter according to embodiments of the present invention are provided as SEQ ID NO: 22 and SEQ ID NO: 38. See, e.g., Yokoyama, R. et al., "The *Arabidopsis ERECTA* gene is expressed in the shoot apical meristem and organ primordia," *The Plant Journal* **15**(3): 301-310 (1998). *pAt.Erecta* is an example of a vegetative stage promoter that is also meristem-preferred. Other vegetative stage, meristem-preferred or meristem-specific promoters have been identified based on their characterized expression profile (see, e.g., Examples 4 and 7 below) that may also be used to drive FT expression according to embodiments of the present invention. For example, the following soybean receptor like kinase (RLK) genes were identified that could be used as vegetative stage, meristem-preferred promoters: Glyma10g38730 (SEQ ID NO: 23), Glyma09g27950 (SEQ ID NO: 24), Glyma06g05900 (SEQ ID NO: 25), and Glyma17g34380 (SEQ ID NO: 26). Vegetative stage, meristem-preferred promoters according to embodiments of the present invention may also include receptor like kinase (RLK) gene promoters from potato: PGSC0003DMP400032802 (SEQ ID NO: 27) and PGSC0003DMP400054040 (SEQ ID NO: 28). Given the characterization provided herein of the *pAt.Erecta* promoter driving FT

expression and the similar expression profiles identified for other RLK, *Erecta* or *Erecta-like* (*Erl*) genes, vegetative-stage, meristem-preferred or meristem-specific promoters of the present invention may further comprise any known or later identified promoter sequences of RLK, *Erecta* and *Erecta-like* genes from other dicotyledonous species having vegetative-stage pattern of expression in the meristems of plants.

[0064] Additional examples of vegetative stage, meristem-preferred or meristem-specific promoters may include those from the following *Arabidopsis* genes: *Pinhead* (*At.PNH*) (SEQ ID NO: 29), *Angustifolia 3* or *At.AN3* (SEQ ID NO: 30), *At.MYB17* (*At.LMI2* or *Late Meristem Identity 2*; At3g61250) (SEQ ID NO: 31), *Kinesin-like gene* (At5g55520) (SEQ ID NO: 32), AP2/B3-like genes, including *At.REM17* (SEQ ID NO: 33) or *At.REM19*, and *Erecta-like 1* and *2* genes, *At.Erl1* (SEQ ID NO: 34) and *At.Erl2* (SEQ ID NO: 35), and functional portions thereof. The polynucleotide sequence of these promoters (or a functional fragment thereof) may also have a relaxed sequence identity while still maintaining a similar or identical pattern of expression of an associated gene or transgene operably linked to the promoter. For example, the promoter may comprise a polynucleotide sequence that is at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or least 99% identical to a polynucleotide sequence selected from the above SEQ ID NOs: 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, or 35, or a functional portion thereof. A “functional portion” of a known or provided promoter sequence is defined as one or more continuous or discontinuous portion(s) of the known or provided promoter sequence that may functionally drive, cause, promote, etc., expression of its associated gene (or transgene) in a manner that is identical or similar to the known or provided promoter sequence. Based on the present disclosure, one skilled in the art would be able to determine if a promoter comprising one or more portion(s) of a known or provided promoter sequence, and/or having a shorter sequence and/or more relaxed sequence identity relative to a known or provided promoter sequence, causes a similar pattern of expression and/or similar phenotypes or effects when its associated FT transgene is expressed in a plant as compared to the known or provided promoter sequence.

[0065] As stated above, a recombinant DNA molecule of the present invention may generally comprise an FT transgene or expression cassette including a polynucleotide sequence encoding an FT protein that is operatively linked to a vegetative stage promoter, which may also be a meristem-preferred or meristem-specific promoter. The polynucleotide coding sequence of the FT transgene or expression cassette may also be operatively linked to one or

more additional regulatory element(s), such as an enhancer(s), leader, transcription start site (TSS), linker, 5' and 3' untranslated region(s), intron(s), polyadenylation signal, termination region or sequence, etc., that are suitable or necessary for regulating or allowing expression of the FT transgene or cassette to effectively produce an FT protein in a plant cell. Such additional regulatory element(s) may be optional and used to enhance or optimize expression of the transgene. For purposes of the present invention, an "enhancer" may be distinguished from a "promoter" in that an enhancer typically lacks a transcription start site, TATA box, or equivalent sequence and is thus insufficient alone to drive transcription. As used herein, a "leader" may be defined generally as the DNA sequence of the untranslated 5' region (5' UTR) of a gene (or transgene) between the transcription start site (TSS) and the protein coding sequence start site.

[0066] According to embodiments of the present invention, the term "recombinant" in reference to a DNA molecule, construct, vector, etc., refers to a DNA molecule or sequence that is not found in nature and/or is present in a context in which it is not found in nature, including a DNA molecule, construct, etc., comprising a combination of DNA sequences that would not naturally occur contiguously or in close proximity together without human intervention, and/or a DNA molecule, construct, etc., comprising at least two DNA sequences that are heterologous with respect to each other. A recombinant DNA molecule, construct, etc., may comprise DNA sequence(s) that is/are separated from other polynucleotide sequence(s) that exist in proximity to such DNA sequence(s) in nature, and/or a DNA sequence that is adjacent to (or contiguous with) other polynucleotide sequence(s) that are not naturally in proximity with each other. A recombinant DNA molecule, construct, etc., may also refer to a DNA molecule or sequence that has been genetically engineered and constructed outside of a cell. For example, a recombinant DNA molecule may comprise any suitable plasmid, vector, etc., and may include a linear or circular DNA molecule. Such plasmids, vectors, etc., may contain various maintenance elements including a prokaryotic origin of replication and selectable marker, as well as a FT expressing transgene or expression cassette perhaps in addition to a plant selectable marker gene, etc.

[0067] As introduced above, the florigenic effects of FT expression may be opposed by the activity of various other anti-florigenic (or non-florigenic) proteins, such as *Terminal Flower* (TFL) genes, in the meristem of a plant. Flowering time and duration may thus be seen as a balance between florigenic and anti-florigenic signals present within the meristem(s) of a plant. Accordingly, it is further proposed that flowering time in a plant may be altered or

induced by suppression of one or more anti-florigenic genes during vegetative stages of development to render the vegetative meristem more responsive to a florigenic signal. In soybean, for example, genes related to *Arabidopsis* TFL1 are primarily expressed in floral meristems where they oppose the functions of florigenic signals to regulate developmental decisions like stem growth habit. In particular, the TFL1-like genes, *TFL1a* and *TFL1b*, in soybean are expressed in the shoot apical meristem, and allelic diversity in *TFL1b* may be largely responsible for changes in stem growth habit in soy resulting in determinate or indeterminate growth. See, e.g., Liu, B. et al., "The soybean stem growth habit gene Dt1 is an ortholog of *Arabidopsis* TERMINAL FLOWER 1." *Plant Physiol* 153(1):198-210 (2010). Accordingly, suppressing the expression of TFL or another anti-florigenic protein in the meristem of a dicot plant may result in increased sensitivity of those meristematic tissues to florigenic signals, such as FT, resulting in early flowering and increased pods per node, similarly to overexpression of FT in these tissues.

[0068] Thus, it is contemplated that a recombinant DNA molecule of the present invention may further comprise a suppression construct having a transcribable DNA sequence operatively linked to a vegetative stage promoter, which may also be a meristem-preferred or meristem-specific promoter, wherein the transcribable DNA sequence encodes an RNA molecule that causes targeted suppression of an endogenous anti-florigenic gene, such as a *TFL* gene. Various methods for suppressing the expression of an endogenous gene are known in the art.

[0069] According to another broad aspect of the present invention, methods are provided for transforming a plant cell, tissue or explant with a recombinant DNA molecule or construct comprising an FT transgene or expression cassette to produce a transgenic plant. Numerous methods for transforming chromosomes in a plant cell with a recombinant DNA molecule are known in the art, which may be used according to methods of the present invention to produce a transgenic plant cell and plant. Any suitable method or technique for transformation of a plant cell known in the art may be used according to present methods. Effective methods for transformation of plants include bacterially mediated transformation, such as *Agrobacterium*-mediated or *Rhizobium*-mediated transformation, and microprojectile bombardment-mediated transformation. A variety of methods are known in the art for transforming explants with a transformation vector via bacterially mediated transformation or microprojectile bombardment and then subsequently culturing, etc, those explants to regenerate or develop transgenic plants. Other methods for plant transformation,

such as microinjection, electroporation, vacuum infiltration, pressure, sonication, silicon carbide fiber agitation, PEG-mediated transformation, etc., are also known in the art. Transgenic plants produced by these transformation methods may be chimeric or non-chimeric for the transformation event depending on the methods and explants used. Methods are further provided for expressing an FT transgene in one or more plant cells or tissues under the control of a vegetative-stage promoter, which may also be a meristem-preferred or meristem-specific promoter. Such methods may be used to alter flowering time of a plant and/or the number of productive or successful flowers, fruits, pods, and/or seeds per node of the plant relative to a wild type or control plant not having the FT transgene. Indeed, methods of the present invention may be used to alter reproductive or yield-related phenotype(s) or trait(s) of the transgenic plant.

[0070] Transformation of a target plant material or explant may be practiced in tissue culture on nutrient media, for example a mixture of nutrients that allow cells to grow in vitro. Recipient cell targets or explants may include, but are not limited to, meristems, shoot tips, protoplasts, hypocotyls, calli, immature or mature embryos, shoots, buds, nodal sections, leaves, gametic cells such as microspores, pollen, sperm and egg cells, etc., or any suitable portions thereof. It is contemplated that any transformable cell or tissue from which a fertile plant can be regenerated or grown/developed may be used as a target for transformation. Transformed explants, cells or tissues may be subjected to additional culturing steps, such as callus induction, selection, regeneration, etc., as known in the art. Transformed cells, tissues or explants containing a recombinant DNA insertion may be grown, developed or regenerated into transgenic plants in culture, plugs or soil according to methods known in the art. Transgenic plants may be further crossed to themselves or other plants to produce transgenic seeds and progeny. A transgenic plant may also be prepared by crossing a first plant comprising the recombinant DNA sequence or transformation event with a second plant lacking the insertion. For example, a recombinant DNA sequence may be introduced into a first plant line that is amenable to transformation, which may then be crossed with a second plant line to introgress the recombinant DNA sequence into the second plant line. Progeny of these crosses can be further back crossed into the more desirable line multiple times, such as through 6 to 8 generations or back crosses, to produce a progeny plant with substantially the same genotype as the original parental line but for the introduction of the recombinant DNA sequence.

[0071] A recombinant DNA molecule or construct of the present invention may be included within a DNA transformation vector for use in transformation of a target plant cell, tissue or explant. Such a transformation vector of the present invention may generally comprise sequences or elements necessary or beneficial for effective transformation in addition to the FT expressing transgene or expression cassette. For *Agrobacterium*-mediated transformation, the transformation vector may comprise an engineered transfer DNA (or T-DNA) segment or region having two border sequences, a left border (LB) and a right border (RB), flanking at least the FT expressing transgene or expression cassette, such that insertion of the T-DNA into the plant genome will create a transformation event for the FT transgene or expression cassette. In other words, the FT transgene or expression cassette would be located between the left and right borders of the T-DNA, perhaps along with an additional transgene(s) or expression cassette(s), such as a plant selectable marker transgene and/or other gene(s) of agronomic interest that may confer a trait or phenotype of agronomic interest to a plant. In addition to protein encoding sequences, a gene of agronomic interest may further comprise a polynucleotide sequence encoding a RNA suppression element. According to alternative embodiments, the FT transgene or expression cassette and the plant selectable marker transgene (or other gene of agronomic interest) may be present in separate T-DNA segments on the same or different recombinant DNA molecule(s), such as for co-transformation. A transformation vector or construct may further comprise prokaryotic maintenance elements, which for *Agrobacterium*-mediated transformation may be located in the vector backbone outside of the T-DNA region(s).

[0072] A plant selectable marker transgene in a transformation vector or construct of the present invention may be used to assist in the selection of transformed cells or tissue due to the presence of a selection agent, such as an antibiotic or herbicide, wherein the plant selectable marker transgene provides tolerance or resistance to the selection agent. Thus, the selection agent may bias or favor the survival, development, growth, proliferation, etc., of transformed cells expressing the plant selectable marker gene, such as to increase the proportion of transformed cells or tissues in the R₀ plant. Commonly used plant selectable marker genes include, for example, those conferring tolerance or resistance to antibiotics, such as kanamycin and paromomycin (*nptII*), hygromycin B (*aph IV*), streptomycin or spectinomycin (*aadA*) and gentamycin (*aac3* and *aacC4*), or those conferring tolerance or resistance to herbicides such as glufosinate (*bar* or *pat*), dicamba (*DMO*) and glyphosate (*aroA* or *EPSPS*). Plant screenable marker genes may also be used, which provide an ability

to visually screen for transformants, such as luciferase or green fluorescent protein (GFP), or a gene expressing a beta glucuronidase or *uidA* gene (GUS) for which various chromogenic substrates are known.

[0073] According to embodiments of the present invention, methods for transforming a plant cell, tissue or explant with a recombinant DNA molecule or construct may further include site-directed or targeted integration. According to these methods, a portion of a recombinant DNA donor template molecule (i.e., an insertion sequence) may be inserted or integrated at a desired site or locus within the plant genome. The insertion sequence of the donor template may comprise a transgene or construct, such as an FT transgene or construct comprising a polynucleotide sequence encoding a florigenic FT protein operatively linked to a vegetative-stage promoter, which may also be a meristem-preferred or meristem-specific promoter. The donor template may also have one or two homology arms flanking the insertion sequence to promote the targeted insertion event through homologous recombination and/or homology-directed repair. Thus, a recombinant DNA molecule of the present invention may further include a donor template for site-directed or targeted integration of a transgene or construct, such as an FT transgene or construct, into the genome of a plant.

[0074] Any site or locus within the genome of a plant may potentially be chosen for site-directed integration of a transgene or construct of the present invention. For site-directed integration, a double-strand break (DSB) or nick may first be made at a selected genomic locus with a site-specific nuclease, such as, for example, a zinc-finger nuclease, an engineered or native meganuclease, a TALE-endonuclease, or an RNA-guided endonuclease (e.g., Cas9 or Cpf1). Any method known in the art for site-directed integration may be used. In the presence of a donor template molecule, the DSB or nick may then be repaired by homologous recombination between the homology arm(s) of the donor template and the plant genome, or by non-homologous end joining (NHEJ), resulting in site-directed integration of the insertion sequence into the plant genome to create the targeted insertion event at the site of the DSB or nick. Thus, site-specific insertion or integration of a transgene or construct may be achieved.

[0075] According to embodiments of the present invention, a plant that may be transformed with a recombinant DNA molecule or transformation vector comprising an FT transgene may include a variety of flowering plants or angiosperms, which may be further defined as including various dicotyledonous (dicot) plant species, such as soybean, cotton, alfalfa,

canola, sugar beets, alfalfa and other leguminous plants. A dicot plant could be a member of the *Brassica* sp. (e.g., *B. napus*, *B. rapa*, *B. juncea*), particularly those *Brassica* species useful as sources of seed oil, alfalfa (*Medicago sativa*), sunflower (*Helianthus annuus*), safflower (*Carthamus tinctorius*), oil palm (*Elaeis* spp.), sesame (*Sesamum* spp.), coconut (*Cocos* spp.), soybean (*Glycine max*), tobacco (*Nicotiana tabacum*), potato (*Solanum tuberosum*), peanuts (*Arachis hypogaea*), cotton (*Gossypium barbadense*, *Gossypium hirsutum*), sweet potato (*Ipomoea batatas*), cassava (*Manihot esculenta*), coffee (*Coffea* spp.), tea (*Camellia* spp.), fruit trees, such as apple (*Malus* spp.), *Prunus* spp., such as plum, apricot, peach, cherry, etc., pear (*Pyrus* spp.), fig (*Ficus casica*), banana (*Musa* spp.), etc., citrus trees (*Citrus* spp.), cocoa (*Theobroma cacao*), avocado (*Persea americana*), olive (*Olea europaea*), almond (*Prunus amygdalus*), walnut (*Juglans* spp.), strawberry (*Fragaria* spp.), watermelon (*Citrullus lanatus*), pepper (*Capsicum* spp.), sugar beet (*Beta vulgaris*), grape (*Vitis*, *Muscadinia*), tomato (*Lycopersicon esculentum*, *Solanum lycopersicum*), and cucumber (*Cucumis sativis*). Leguminous plants include beans and peas. Beans include, for example, guar, locust bean, fenugreek, soybean, garden beans, cowpea, mungbean, lima bean, fava bean, lentils, and chickpea. Given that the present invention may apply to a broad range of plant species, the present invention further applies to other botanical structures analogous to pods of leguminous plants, such as bolls, siliques, fruits, nuts, tubers, etc. According to embodiments of the present invention and depending on the particular plant species transformed, a plant ectopically expressing a florigenic FT sequence may have an altered or greater number of bolls, siliques, fruits, nuts, tubers, etc., per node(s), main stem, and/or branch(es) of the plant, and/or an altered or greater number of bolls, siliques, fruits, nuts, tubers, etc., per plant, relative to a wild type or control plant not having the FT transgene.

[0076] According to another broad aspect of the present invention, a transgenic plant(s), plant cell(s), seed(s), and plant part(s) are provided comprising a transformation event or insertion into the genome of at least one plant cell thereof, the transformation event or insertion comprising a recombinant DNA sequence, construct or polynucleotide including a Flowering Locus T (FT) transgene or expression cassette, wherein the FT transgene or expression cassette further comprises a polynucleotide sequence encoding an FT protein operably linked to a vegetative stage promoter, which may also be a meristem-preferred or meristem-specific promoter. The FT protein encoded by the polynucleotide sequence may be native to the transgenic plant transformed with the polynucleotide sequence, or homologous or otherwise similar to a FT protein native to the transgenic plant (i.e., not native to the transgenic plant).

Such a transgenic plant may be produced by any suitable transformation method, which may be followed by selection, culturing, regeneration, development, etc., as desired or needed to produce a transgenic R₀ plant, which may then be selfed or crossed to other plants to generate R₁ seed and subsequent progeny generations and seed through additional crosses, etc. Similarly, embodiments of the present invention further include a plant cell, tissue, explant, etc., comprising one or more transgenic cells having a transformation event or genomic insertion of a recombinant DNA or polynucleotide sequence comprising an FT transgene.

[0077] Transgenic plants, plant cells, seeds, and plant parts of the present invention may be homozygous or hemizygous for a transgenic event or insertion of an FT transgene or expression cassette into the genome of at least one plant cell thereof, or may contain any number of copies of a transgenic event(s) or insertion(s) comprising an FT transgene or expression cassette. The dosage or amount of expression of an FT transgene or expression cassette may be altered by its zygosity and/or number of copies, which may affect the degree or extent of phenotypic changes in the transgenic plant, etc. According to some embodiments, a transgenic plant comprising an FT transgene may be further characterized as having one or more altered flowering or reproductive phenotypes or traits, which may include altered yield-related phenotypes or traits, such as an increase in the number of flowers, pods, etc., and/or seeds per plant (and/or per node of the plant) relative to a wild type or control plant not having the FT transgene. Such a transgenic plant may be further characterized as having an altered structure, morphology, and/or architecture due to altered plant height, branching patterns, number of floral nodes, etc., relative to a wild type or control plant. Indeed, yield-related phenotypes or traits altered by FT overexpression in a transgenic plant may include: flowering time, reproductive duration, flowering duration, amount or timing of abscission of flowers, pods, siliques, bolls, fruits, nuts, etc., number of flowers per node, number of racemes per node, number of branches, number of nodes per plant, number of nodes on the main stem, number of nodes on branches, number of pods, bolls, siliques, fruits, nuts, etc., per plant, number of pods, bolls, siliques, fruits, nuts, etc., per node, number of pods on the main stem, number of pods, bolls, siliques, fruits, nuts, etc., on branches, 1000 seed weight, number of seeds per plant, number of seeds per node, and/or altered plant architecture, as compared to a wild type or control plant not having the FT transgene.

[0078] For purposes of the present invention, a “plant” may include an explant, seedling, plantlet or whole plant at any stage of regeneration or development. As used herein, a “transgenic plant” refers to a plant whose genome has been altered by the integration or

insertion of a recombinant DNA molecule, construct or sequence. A transgenic plant includes an R_0 plant developed or regenerated from an originally transformed plant cell(s) as well as progeny transgenic plants in later generations or crosses from the R_0 transgenic plant. As used herein, a “plant part” may refer to any organ or intact tissue of a plant, such as a meristem, shoot organ/structure (e.g., leaf, stem and tuber), root, flower or floral organ/structure (e.g., bract, sepal, petal, stamen, carpel, anther and ovule), seed (e.g., embryo, endosperm, and seed coat), fruit (e.g., the mature ovary), propagule, or other plant tissues (e.g., vascular tissue, ground tissue, and the like), or any portion thereof. Plant parts of the present invention may be viable, nonviable, regenerable, and/or non-regenerable. A “propagule” may include any plant part that is capable of growing into an entire plant. For purposes of the present invention, a plant cell transformed with an FT transgene according to embodiments of the present invention may include any plant cell that is competent for transformation as understood in the art based on the method of transformation, such as a meristem cell, an embryonic cell, a callus cell, etc. As used herein, a “transgenic plant cell” simply refers to any plant cell that is transformed with a stably-integrated recombinant DNA molecule or sequence. A transgenic plant cell may include an originally-transformed plant cell, a transgenic plant cell of a regenerated or developed R_0 plant, or a transgenic plant cell from any progeny plant or offspring of the transformed R_0 plant, including cell(s) of a plant seed or embryo, or a cultured plant or callus cell, etc.

[0079] Embodiments of the present invention may further include methods for making or producing transgenic plants having altered reproductive and/or yield-related traits or phenotypes, such as by transformation, crossing, etc., wherein the method comprises introducing a recombinant DNA molecule or sequence comprising an FT transgene into a plant cell, and then regenerating or developing the transgenic plant from the transformed plant cell, which may be performed under selection pressure favoring the transgenic event. Such methods may comprise transforming a plant cell with a recombinant DNA molecule or sequence comprising an FT transgene, and selecting for a plant having one or more altered phenotypes or traits, such as one or more of the following: flowering time, reproductive duration, flowering duration, amount or timing of abscission of flowers, pods, bolls, siliques, fruits, nuts, etc., number of flowers per node, number of racemes per node, number of branches, number of nodes per plant, number of nodes on the main stem, number of nodes on branches, number of pods, bolls, siliques, fruits, or nuts per plant, number of pods, bolls, siliques, fruits, nuts, etc., per node, number of pods, bolls, siliques, fruits, nuts, etc., on the

main stem, number of pods, bolls, siliques, fruits, nuts, etc., on branches, 1000 seed weight, number of seeds per plant, number of seeds per node, and altered plant architecture, as compared to a wild type or control plant not having the FT transgene. For example, embodiments of the present invention may comprise methods for producing a transgenic plant having an increased number of flowers, pods, and/or seeds per plant (and/or an increased number of flowers, pods, and/or seeds per node of the plant), wherein the method comprises introducing a recombinant DNA molecule comprising an FT transgene into a plant cell, and then regenerating or developing the transgenic plant from the plant cell.

[0080] According to another broad aspect of the present invention, methods are provided for planting a transgenic plant(s) of the present invention at a normal or high density in field. According to some embodiments, the yield of a crop plant per acre (or per land area) may be increased by planting a transgenic plant(s) of the present invention at a higher density in the field. As described herein, transgenic plants of the present invention expressing a florigenic FT protein during vegetative stage(s) of development may exhibit increased pods per node (particularly on the main stem), but may also have an altered plant architecture with reduced branching and fewer nodes per branch. Thus, it is proposed that transgenic plants of the present invention may be planted at a higher density to increase yield per acre in the field. For row crops, higher density may be achieved by planting a greater number of seeds/plants per row length and/or by decreasing the spacing between rows. According to some embodiments, a transgenic crop plant of the present invention may be planted at a density in the field (plants per land/field area) that is at least 5%, 10%, 15%, 20%, 25%, 50%, 75%, 100%, 125%, 150%, 175%, 200%, 225%, or 250% higher than a normal planting density for that crop plant according to standard agronomic practices.

[0081] For soybean, the typical planting density is in a range from about 100,000 to 150,000 seeds per acre, and the typical row spacing is in a range from about 26 to about 40 inches, such as 30 inch or 36 inch row spacing. Within a given row, about 6-8 soybean seeds may typically be planted per foot. In contrast, high density planting for soybean may include a range of approximately 150,000 to 250,000 seeds per acre, and the row spacing may be within a range from about 10 inches or less to about 25 inches, such as 10 inch, 15 inch or 20 inch row spacing. For high density planting, approximately 9-12 soybean seeds per foot may be planted within each row, perhaps in combination with narrower row spacing. However, high crop density may be achieved by narrow row spacing without an increase in planting density within each row.

[0082] For cotton, the typical planting density is in a range from about 28,000 to 45,000 seeds per acre, and the typical row spacing is in a range from about 38 to about 40 inches, such as 38 inch or 40 inch row spacing. Within a given row, about 2-3 cotton seeds may typically be planted per foot. In contrast, high density planting for soybean may include a range of approximately 48,000 to 60,000 seeds per acre, and the row spacing may be within a range from about 30 inches or less to about 36 inches. For high density planting, approximately 3-5 cotton seeds per foot may be planted within each row, perhaps in combination with narrower row spacing. However, high crop density for cotton may be achieved by narrow row spacing without an increase in planting density within each row.

[0083] For canola, the typical planting density is in a range from about 360,000 to 550,000 seeds per acre, and the typical row spacing between openers is in a range from about 6 inches to about 16 inches. Within a given row, about 8-12 canola seeds may typically be planted per foot. In contrast, high density planting for soybean may include a range of approximately 450,000 to 680,000 seeds per acre, and the row spacing may be within a range from about 5 inches or less to about 10 inches. For high density planting, approximately 10-16 canola seeds per foot may be planted within each row, perhaps in combination with the narrower row spacing. However, high crop density for canola may be achieved by narrow row spacing without an increase in planting density within each row.

EXAMPLES

Example 1. Soybean short day induction treatment and identification of Flowering Locus T (FT) genes by transcriptional profiling.

[0084] Methods for the photoperiodic light treatment (i.e., short day induction of flowering in plants) are described in U.S. Patent No. 8,935,880 and U.S. Patent Application Publication No. 2014/0259905, which are incorporated herein by reference in their entirety. As described further therein, the early short day induction treatment produced soybean plants having altered reproductive traits including an increased number of pods/seeds per plant. Transcriptional profiling experiments were performed using gene expression microarrays to determine if particular transcripts were up-regulated in these light-induced plants to identify genes that may be responsible for mediating the short day induction phenotypes. In these experiments, an analysis of transcripts was conducted on soybean leaf and floral apex tissues collected after 1, 3 and 5 days from plants that received a short day inductive light treatment

(Short day) in comparison to tissues from plants that did not receive the inductive treatment (Long day).

[0085] As shown in FIG. 2, an increased accumulation of transcripts was observed for a particular *Flowering Locus T* gene, *Gm.FT2a* (SEQ ID NO: 1), in leaf tissue harvested at 3 and 5 days after the early short day induction (eSDI) treatment in comparison to samples taken from either (i) floral apex tissues of the same short day induction plants, or (ii) leaf tissues and floral apex tissues of soybean plants that instead received the long day treatment. These data support the conclusion that *Gm.FT2a* expression is induced in leaf tissue of plants experiencing the eSDI treatment, which was not seen in plants grown under long day conditions. *Gm.FT2a* expression was also not observed in the floral apex of eSDI treated plants, which is consistent with the model of FT protein expression being induced in peripheral leafy tissues in response to inductive photoperiod conditions and then migrating to its site of action in the meristems to induce flowering. However, additional experiments using a more sensitive RNA sequencing analysis of transcripts did show some *Gm.FT2a* induction in the shoot apex and axillary buds in response to the eSDI treatment (data not shown).

Example 2. Characterization of the *pAt.Erecta* promoter expression patterns in soybean.

[0086] Achieving desirable traits or phenotypes by transgenic approaches may require control of the temporal and spatial patterns of ectopic FT gene expression. Soy physiological experiments identifying *Gm.FT2a* expression in vegetative tissues following the short day induction treatment (see FIG. 2) indicated that achieving yield positive traits may rely on earlier FT expression during the vegetative stage. On the other hand, even though FT transcripts are not detected in the vegetative apex, FT protein has been shown to move long distance from the leaves to the apical tissue where it triggers a vegetative to reproductive transition. See, e.g., Lifschitz, E. et al., (2006), *supra*; and Corbeiser, L. et al., “FT Protein Movement Contributes to Long-Distance Signaling in Floral Induction of *Arabidopsis*”, *Science* 316: 1030-1033 (2007). Thus, in light of our own observations, we proposed using a vegetative stage promoter that is active in the meristem to control ectopic FT expression in a plant. By expressing the morphogenetic FT signal directly in the meristem at the desired developmental stage, multiple endogenous pathways and regulatory feedbacks (e.g., control

of FT transduction in the leaf and long distance translocation of the FT signal) may be bypassed or avoided. Previous experiments with the short day induction treatment (described above in Example 1) revealed up-regulation of the *Gm.Erecta* gene in the meristems of soybean plants. The *pErecta* promoter (SEQ ID NO: 21) from *Arabidopsis* had been shown to have weak expression in the meristem(s) of plants during vegetative stages of development. Accordingly, the *pAt.Erecta* promoter was selected for initial FT expression experiments.

[0087] Additional experiments were performed to further characterize the expression patterns of *pAt.Erecta* fused to a GUS reporter gene in vegetative and floral meristematic tissues. Analysis of GUS expression patterns during the development of soy seedlings indicated that the *pAt.Erecta* promoter exhibits a temporal and spatial pattern of expression, preferably in the meristematic tissues during the vegetative stage of development. FIGS. 3 and 4 and FIGS. 5 and 6 include two sets of images to show the pattern of GUS staining. FIGS. 3 and 5 provide black and white images of the stained tissues, and FIGS. 4 and 6 provide black and white images corresponding to FIGS. 3 and 5, respectively, but color filtered to show the pattern and intensity of blue GUS staining. Thus, the GUS staining pattern of expression can be viewed with these black and white images by comparing the corresponding images of FIGS. 3 and 4 or FIGS. 5 and 6. As shown in FIGS. 3 and 4, GUS staining was detected in the soy immature uni-foliate blade and petiole (FIGS. 4A and 4B) at three days after sowing/germination. *pAt.Erecta:GUS* expression was also broadly detected in the trifoliate primordia, shoot apical meristem (SAM) and axillary meristem sites at this early vegetative stage (FIG. 4C). GUS activity was not detected in the fully expanded uni-foliate and trifoliate leaves at ten days after germination or planting (FIGS. 4D and 4E). However, GUS activity was detected at the proximal part of the immature, unexpanded, but fully developed trifoliate blade, and at the adaxial side of the petiole (FIG. 4F). Detailed observation of the developing apical tissue showed that broad expression was retained in the developing immature leaf primordia, axillary meristems and shoot apical meristems (FIGS. 4G-I).

[0088] At the early reproductive stage, *pAt.Erecta* promoter activity was not detected in the mature blade and was reduced in the developing leaf primordia. The GUS signal was not detected in the indeterminate vegetative apex at the shoot apical meristem (SAM) or in the axillary meristem (AM) once these tissues started to form inflorescences (FIGS. 4J-4L). In all later stages, any additional meristematic activity could not be detected in the apex or in the axillaries or flower primordia. However, GUS expression continued in the adaxial side of the

petiole and proximal part of the immature leaf blade (FIGS. 4M and 4N), but not in the fully expanded leaf blade (FIG. 4O). GUS expression patterns with the *pAt.Erecta* promoter were also analyzed at the later R1 stages of development (35-40 days after germination). Similar to earlier stages of development, no expression was detected in the mature leaves or stems. However, strong promoter activity was detected in the inflorescence stems (FIGS. 5A and 6A; see arrow) and floral pedicels (FIGS. 5B and 6B; see arrow). In both tissues, expression was detected in vasculature and parenchyma cells (FIGS. 5C and 6C). At this stage, expression was also detected in the stamen filaments (FIG. 5D and 6D; see arrows) and in the un-pollinated ovules (FIGS. 5E, 5F, 6E and 6F; see arrows in 6F).

[0089] Previously, the *pAt.Erecta* promoter was characterized in *Arabidopsis*. Interestingly, *pAt.Erecta* expression patterns in *Arabidopsis* were comparable to the patterns observed in soy during the vegetative stage, but not during late reproductive stages. In contrast, the *pAt.Erecta* expression pattern in soybean is diminished in early reproductive tissues but reemerges in some later reproductive organs and tissues, including the inflorescence stems and floral pedicels. See, e.g., Chen, M-K et al., *FEBS Letters* 588: 3912-17 (2014); Yokoyama, R et al.; Shpak, ED et al., *Science* 309: 290-293 (2005); and Yokoyama, R et al., *Plant J* 15(3): 301-310 (1998), the entire contents and disclosures of which are incorporated herein by reference. Thus, the *pAt.Erecta* promoter provides a novel expression pattern in soybean.

Example 3. Expression of Flowering Locus T gene, *Gm.FT2a*, under control of a *pAt.Erecta* promoter alters flowering time and pods per node in soybean.

[0090] Transgenic soybean plants were produced by transforming soybean explants with a recombinant DNA molecule (i.e., a T-DNA transformation vector) comprising the *pAt.Erecta* promoter operably linked to the *Gm.FT2a* gene via *Agrobacterium*-mediated transformation to generate four *pErecta::Gm.FT2a* events that were carried forward for further testing. The effect of FT2a overexpression was immediately seen in R₀ plants, which had very early flowering and termination with reduced seed yield (e.g., only about 8 seeds/plant). These transgenic *Gm.FT2a* plants also had a short plant height and very few, if any, branches. Segregating R1 plants and their progeny were subsequently grown in the greenhouse under long day conditions for initial study and characterization. By growing these plants under long day conditions, the severe dwarf phenotypes observed with *Gm.FT2a* transgenic R₀ plants were improved. In these experiments, both homozygous and hemizygous plants grown in the greenhouse under 16-hour long day conditions (i.e., 16/8 hours of day/night photoperiods) flowered much earlier than wild type null segregants. *Gm.FT2a* transgenic plants flowered at about 19-22 days after planting or seeding). (see, e.g., FIG. 9). Under these growth conditions, transgenic soybean plants expressing *Gm.FT2a* further had an increased number of pods per node on the main stem in comparison to wild type controls (see, e.g., FIGS. 10 and 11, discussed further below).

[0091] Plants containing one of the *pEr::Gm.FT2a* transgenic events (Event 1) grown in controlled environment conditions were further analyzed via scanning electron microscopy analysis (eSEM). Analysis of the shoot apical meristem (SAM) of these transgenic plants (collected at 7 days after planting) revealed an early transition of the SAM into an inflorescence meristem (IM) and floral meristem (FM) (FIG. 7). In contrast, the SAMs of wild type soybean plants were not differentiated into IM at this growth stage. Similarly, imaging of the axillary meristem of the FT2a transformants (collected at 9 days after planting) indicated the development of dormant inflorescence meristems (dIMs) (or lateral primordial racemes) into IM and FM (FIG. 8), leading to more earlier-developing floral branches (racemes) per node in these transgenic plants. Additional phenotypic characterization revealed early flowering at the V1 stage in *Gm.FT2a* expressing soybean plants, which was well before the floral transition occurred in null segregating plants (FIG. 9). These data in combination with the *pAt.Erecta*:GUS expression pattern described above indicate that early flowering, and more particularly the formation of inflorescence and floral

meristems, were induced by ectopic expression of *Gm.FT2a* during the vegetative stage in leaf primordia and the shoot apical and axillary meristems of seedlings. The formation of a higher number of inflorescence and floral meristems is believed to further cause earlier release and elongation of the secondary and tertiary racemes, leading to a greater number of productive flowers and pods being formed per node.

[0092] Not only did *Gm.FT2a* transgenic soybean plants experience earlier flowering and produce more pods per node on the main stem (relative to segregating null plants), the effects of ectopic *Gm.FT2a* expression in transgenic plants were also found to be dosage dependent. Although both homozygous and hemizygous plants had a reduced height and less branching, plants homozygous for the *Gm.FT2a* transgene were more severely affected than hemizygous plants, presumably because homozygous plants contain two copies of the transgene (i.e., a higher dosage), as opposed to only one copy (i.e., a lower dosage) in hemizygous plants. Under long day growth conditions, homozygous plants terminated earlier and had a shorter overall height with fewer nodes and branches on the main stem in comparison to plants hemizygous for the transgene (FIG. 10). Unlike homozygous plants, which exhibited a number of sub-optimal dwarf phenotypes including very few (if any) branches on the main stem, hemizygous plants had an intermediate phenotype in terms of their vegetative growth, plant height, and the number of nodes present on the main stem relative to wild type and homozygous plants. Under 16-hour long day conditions, hemizygous plants had a more normal plant height with some degree of branching and a more extended duration of flowering, relative to homozygous plants (FIG. 10). Hemizygous plants also flowered for 40-64 days after initiation of R1, whereas homozygous plants flowered for only 16-24 days due to their earlier termination.

[0093] Additional experiments were conducted with plants transformed with the *Gm.FT2a* construct (3 events) in long day (16 hour) controlled environment conditions to further characterize the dosage response between hemizygous and homozygous plants. Differences in the number of nodes and pods on the main stem and branches, as well as the average number of pods per node and the average height per plant are shown in Table 1 for three homozygous events (Homo-Event 2, Homo-Event 3, Homo-Event 4) and three hemizygous events (Hemi-Event 2, Hemi-Event 3, Hemi-Event 4). These events are distinguished from Event 1 above.

Table 1. Event level data for homozygous and hemizygous *Gm.FT2a* transgenic plants.

Zygosity- Event #	Avg. # MS nodes per plant	Avg. # BR nodes per plant	Avg. # MS pods per plant	Avg. # BR pods per plant	Avg. Pods per Node	Avg. Height (in) per plant
Homo- Event 2	11.8	6.9	46	9	2.9	17.5
Homo- Event 3	12.3	6.5	66.4	9	4	21
Homo- Event 4	12.5	6.8	49.6	9.1	3	19.5
Hemi- Event 2	25.3	12.4	183.5	47.3	6.1	37.5
Hemi- Event 3	23.9	13.2	200.3	28.8	6.1	40
Hemi- Event 4	25.4	15.3	186.8	58	6	41.5

[0094] As shown in Table 1, hemizygous plants consistently had a higher number of nodes on the main stem (MS) and branches (BR) and a greater plant height than homozygous plants. Thus, hemizygous plants were generally less affected than homozygous plants and more like wild type plants. Hemizygous plants also had an increased number of pods per node and a higher number of pods on the main stem and branches, relative to homozygous plants. Therefore, hemizygous plants generally had a closer-to-normal plant architecture with a greater number of pods per node (and per plant), presumably due to their lower *Gm.FT2a* transgene dosage. The relative dosage level of *Gm.FT2a* based on transgene zygosity was further confirmed by additional experiments showing that *Gm.FT2a* transcript levels were higher in tissues from homozygous plants, than in tissues from hemizygous plants (data not shown).

[0095] The early induction of flowering in these *Gm.FT2a* transgenic plants was associated with more pods (and seeds) per node on the main stem in both hemizygous and homozygous plants. Homozygous and hemizygous plants containing the *Gm.FT2a* transgene each had an increased number of pods/seeds per node on the main stem of the plant in comparison to wild type segregants (FIG. 11). The distribution of pods on the main stem was also found to be different between *FT2a* transgenic and wild type null plants. Both homozygous and hemizygous plants grown under long day conditions were found to have more pods on at least the lower nodes of the main stem and more pods per node on average, in comparison to wild type null plants (data not shown). Plants hemizygous for the *Gm.FT2a* transgene contained the highest number of pods per node over the length of the main stem. However, these effects were dependent on the particular growing conditions including day length, etc. In general, experiments performed with soybean under longer day conditions tended to

produce greater differences between transgenic and null plants.

[0096] The dosage-dependent effects of transgenic *Gm.FT2a* expression were also observed in field trial experiments. In a 2014 field trial, soybean plants hemizygous for two *Gm.FT2a* events (Events 1 and 2 above) showed an average of about 2.68 pods per node on the main stem, and plants homozygous for these events had about 1.40 pods per node on average, whereas null segregating plants had about 1.63 pods per node. In a 2013 field trial, however, plants hemizygous for transgenic *Gm.FT2a* (Event 2) were found to have an average number of about 3.21 pods per node, as compared to an average of about 3.05 pods per node in homozygous plants and about 2.19 pods per node in null segregating plants. In another 2013 micro plot experiment conducted at a different field location, plants hemizygous for the *Gm.FT2a* transgene (Event 1) were found to have about 2.17 pods per node on average, as compared to an average of about 2.05 pods per node in plants homozygous for the *Gm.FT2a* transgene (Event 2) and about 1.30 pods per node in null segregating plants. Thus, the number of pods per node on plants containing the *Gm.FT2a* transgene may depend on a variety of factors including dosage of the FT transgene, environmental and field conditions, and perhaps differences in agronomic practices. However, much like transgenic *Gm.FT2a* plants grown in the greenhouse, homozygous and hemizygous *Gm.FT2a* transgenic plants grown under field conditions often had fewer nodes on the main stem, shorter overall plant height, and/or reduced branching in transgenic plants. Indeed, wild type plants typically had more branching and a greater number of total nodes per plant than hemizygous and homozygous *Gm.FT2a* plants.

[0097] Additional physiological data was collected from homozygous *Gm.FT2a* transgenic plants and wild type (WT) control plants grown in the greenhouse under 14-hour long day conditions (see Table 2). These data provide an average of measurements taken from six *Gm.FT2a* transgenic plants for each event, or from eight wild type plants. The following matrices were collected for phenotypic characterization of these plants: Days to flowering at R1 (DOFR1); Days to R7 (DOR7); reproductive duration in days from R1 to R7 (PDR1R7); number of branches per plant (BRPP); total fertile nodes on branches (FNBR); total fertile nodes per plant (FNLP); total fertile nodes on main stem (FNST); number of nodes on branches (NDBR); number of nodes on main stem (NDMS); number of nodes/plant (NDPL); percent fertile nodes on branches (PFNB); percent total fertile nodes (PFNN); percent fertile nodes on main stem (PFNS); number of pods per plant (PDPP); number of pods on main stem (PODMS); number of pods on branches (PODBR); number of pods/node; seeds per plant at

R8 (SDPPR8); and weight of 1000 seeds (SW1000). Each of these measurements was taken at harvest unless another time point is specified.

Table 2. Construct level phenotypic data for transgenic homozygous *Gm.FT2a* and WT plants.

	WT	pErecta::Gm.FT2a
DOFR1	33.5	21.3
DOR7	106.9	92.9
PDR1R7	76.5	71.6
BRPP	20.1	1
FNBR	190.6	2
FNLP	214.6	15
FNST	24.0	14.3
NDBR	211.4	3
NDMS	33.4	15.3
NDPL	244.9	16.3
PDPP	575.8	61.2
PFNB	90.4	75
PFNN	87.8	92.0
PFNS	71.4	92.9
PODBR	487.3	3
PODMS	88.4	60.2
Pods / Node	2.4	3.8
SDPPR8	1319.6	122.1
SW1000 (grams)	146	122.5

[0098] Consistent with the observations noted above, homozygous *Gm.FT2a* transgenic plants experienced earlier floral induction than WT plants (DOFR1 about 21 days after planting, instead of about 33-34 days in wild type plants). These measurements further showed that the number of branches (and other measurements related to branching, such as the number of nodes or pods on branches) was greatly reduced. Due to the transgenic plants having a shorter stature with very little branching, the total numbers of nodes or pods per plant were also greatly reduced. However, the number of pods per node on the main stem was increased in transgenic plants (e.g., about 3.8 average pods/node) relative to wild type null plants (e.g., about 2.4 pods/node).

[0099] Without being bound by any theory, the larger number of pods per node observed with transgenic soybean plants expressing FT2a in the meristem during vegetative stages of development may be caused at least in part by synchronization of early flowering with early

secondary and/or tertiary raceme release and/or better resource utilization to produce more pod-producing flowers per node. Early FT expression in the meristem (see, e.g., FIGS. 3 and 4) may cause early release of the dormant inflorescence meristems to produce a greater number of racemes per node of the plant, such that a greater number of racemes produce mature flowers and fully developed pods at each node. However, subsequent FT expression in reproductive tissues (see, e.g., FIGS. 5 and 6) may terminate floral development of later developing flowers at each node leading to more efficient resource allocation to the earlier developing racemes, flowers and pods. In wild-type soybean plants, a much lower percentage of secondary and tertiary racemes produce flowers and fully developed pods relative to primary racemes, and later developing flowers of the primary raceme typically do not produce mature flowers and/or full-sized pods prior to abscission. Thus, it is theorized that more pods per node may be generated in plants expressing FT proteins in the vegetative meristem by synchronizing early flower development with early release of the lateral racemes at one or more node(s) of the plant. With at least the *pAt.Erecta* promoter driving FT expression, later developing flowers (that may not otherwise produce fully developed or full-sized pods) may also become terminated by later reproductive-stage expression of FT to direct resources to the earlier developing flowers.

Example 4. Expression of Flowering Locus T gene, *Gm.FT2a*, under control of alternative vegetative stage promoters in soybean.

[00100] Based on the phenotypes observed in the preceding Example 3, two promoters were also proposed to drive *Gm.FT2a* transgene expression that were considered vegetative-stage, leaf-preferred promoters: *pAt.BLS* (SEQ ID NO: 36) and *pAt.ALMT6* (SEQ ID NO: 37). As used herein, a “leaf-preferred” promoter refers to a promoter that preferentially initiates transcription of its associated gene in leaf tissues relative to other plant tissues. Since FT is believed to function as a mobile florigen, early FT expression during vegetative stages in peripheral tissues, such as in the leaf with a leaf-preferred or leaf-specific promoter, may lead to phenotypes similar to the meristem-preferred *pAt.Erecta:Gm.FT2a* expression. It was further theorized that FT expression with a vegetative leaf promoter might also attenuate the floral induction signal, and thus mitigate the early termination phenotypes observed with homozygous FT expression in the meristem, and increase plant height and branching.

[00101] In these experiments, transformation vectors for *pAt.ALMT6::Gm.FT2a* and *pAt.BLS::Gm.FT2a* were constructed and used to transform a soybean line by *Agrobacterium*-mediated transformation. Expression with the *pAt.BLS* promoter has been shown to start in leaf primordia number 5 (p5) and is expressed in the source leaf veins only until transition to flowering, and the *pAt.ALMT6* promoter is also a vegetative leaf promoter with expression at later developmental stages relative to *pAt.BLS*. See, e.g., Efroni et al., “A Protracted and Dynamic Maturation Schedule Underlies *Arabidopsis* Leaf Development,” *The Plant Cell* 20(9): 2293-2306 (2008); and Shani et al., “Stage-Specific Regulation of *Solanum lycopersicum* Leaf Maturation by Class 1 KNOTTED1-LIKE HOMEODOMAIN Proteins,” *The Plant Cell* 21(10): 3078-3092 (2009). Transgenic soybean plants were produced for each of these vector constructs and characterized for phenotypes in growth chambers under 14-hour photoperiod conditions in comparison to wild type plants. For each of the *pAt.BLS* construct, six transgenic events were tested (5 plants per event), and for the *pAt.ALMT6* promoter, seven transgenic events were tested (5 plants per event). For each of these constructs, control data was collected from five wild type plants.

[00102] The following matrices were collected for phenotypic characterization of these transgenic plants (Tables 3 and 4). The individual measurements are as defined above, and phenotypic characterization was conducted on plants homozygous for the transgene.

Table 3. Construct level phenotypic data for *pALMT6::Gm.FT2a* and WT plants.

	<u>WT</u>	<u>pALMT6::FT2a</u>
DOFR1	35.2	38.8
DOR7	84.7	88.8
PDR1R7	49.5	50.0
BRPP	7.7	8.9
FNBR	57.8	73.3
FNLP	69.7	85.0
FNST	12.0	11.7
NDBR	78.9	96.0
NDMS	21.3	22.5
NDPL	100.2	118.5
PDPP	120.2	141.1
PFNB	73.2	76.8
PFNN	71.7	72.1
PFNS	57.9	51.7
PODBR	91.8	118.1
PODMS	28.3	22.9
Pods / Node	1.4	1.2

Table 4. Construct level phenotypic data for *pBLS::Gm.FT2a* and WT plants.

	<u>WT</u>	<u>pBLS::FT2a</u>
DOFR1	31.3	35.2
DOR7	78.1	82.6
PDR1R7	46.9	47.5
BRPP	7.5	8.8
FNBR	65.7	81.2
FNLP	80.5	94.0
FNST	14.9	12.7
NDBR	72.2	95.6
NDMS	21.9	22.3
NDPL	94.0	117.9
PDPP	137.0	148.1
PFNB	92.3	85.3
PFNN	87.4	80.1
PFNS	68.1	57.3
PODBR	100.9	123.4
PODMS	36.1	24.8
Pods / Node	1.7	1.3

[00103] Transgenic plants expressing *Gm.FT2a* under the control of the alternative *pAt.ALMT6* and *pAt.BLS* promoters were phenotypically more similar to wild type (WT) plants than *pAT.Erecta::Gm.FT2a* transgenic plants. Plants transformed with the

pAt.ALMT6::Gm.FT2a and *pAt.BLS::Gm.FT2a* constructs had flowering times and vegetative growth traits similar to wild type control plants, perhaps with a slightly increased number of nodes on branches as compared to wild type plants (Tables 3 and 4). These data may be interpreted to indicate that both the timing and location of transgenic FT expression are important for producing reproductive and yield-related traits or phenotypes that differ from wild-type plants. Merely expressing a FT transgene during earlier vegetative stages of development (e.g., in leaf tissues) may not be sufficient to alter the reproductive or yield-related phenotypes of a plant (e.g., pods per node). Thus, according to embodiments of the present invention, a promoter operably linked to a florigenic FT transgene may preferably be a meristem-specific or meristem-preferred promoter in addition to driving expression during the vegetative stages of plant development. However, when the expression profiles for the above two leaf-preferred promoters were tested in soybean plants, no GUS staining was observed in the developing leaf with the *pAt.BLS* promoter, and the *pAt.ALMT6* promoter did not produce detectable GUS expression in the leaf until late vegetative stages with much higher expression during early reproductive stages. Thus, it remains possible that expression of FT transgenes in peripheral (leaf) tissues during early vegetative stages using different tissue-specific promoters may be sufficient in some cases to induce early flowering and/or cause other reproductive or yield-related traits or phenotypes, which may also depend on the particular plant species tested.

Example 5. Identification of protein domains of FT homologs by Pfam analysis.

[00104] *Gm.FT2a* orthologs were identified by sequence analysis and literature review, and a few examples of these FT homologs are listed in Table 5 along with *Gm.FT2a*. These included other soybean FT genes as well as a few FT genes from other plant species. The amino acid sequences of these FT proteins were analyzed to identify any Pfam protein domains using the HMMER software and Pfam databases (version 27.0). These FT protein sequences (SEQ ID NOs: 2, 4, 6, 8, 10 and 12) were found to have the same Pfam domain identified as a phosphatidyl ethanolamine binding domain protein (PEBP) having a Pfam domain name of “PBP_N”, and a Pfam accession number of PF01161. The location of the PBP_N domains in each of these FT protein sequences are also listed in Table 5. The location of the PBP_N domain in other FT proteins can be determined by sequence alignment. It is thus contemplated that any DNA sequence encoding at least an FT protein comprising the PBP_N domain may be used in a recombinant DNA molecule of the present

invention, as long as the corresponding FT protein has florigenic activity when ectopically expressed in the meristem of a plant.

Table 5. Location of PBP_N (Pfam) domain in FT protein sequences.

PROTEIN SEQ ID NO.	Gene Name	Domain location
2	<i>Gm.FT2a</i>	28-162
4	<i>Gm.FT5a</i>	26-157
6	<i>Gm.FT2b</i>	28-162
8	<i>Zm.ZCN8</i>	26-154
10	<i>Nt.FT-like</i>	25-159
12	<i>Le.SFT</i>	29-161

Example 6. Expression of FT homologs under control of *pAt.Erecta* promoter in soybean.

[00105] Additional transformation vectors containing other FT homologs (Table 6) under control of the *pAt.Erecta* promoter were constructed and used to transform soybeans via *Agrobacterium*-mediated transformation. Transgenic plants generated from these events were characterized for their phenotypes in the greenhouse with a 14 to 14.5 hour natural daylight photoperiod. For each construct, six events were tested (6 plants per event). Six plants were also tested and averaged for wild type (WT) control plants. Different groups of experiments (A-E) were conducted as shown in Table 6 with separate wild type controls.

Table 6. List of constructs for some *Gm.FT2a* and its homologs with their protein sequences.

Construct Description	Gene Name	PROTEIN SEQ ID NO.	Testing Group
pErecta: <i>Gm.FT2a</i>	<i>Gm.FT2a</i>	2	A
pErecta: <i>Gm.FT2b</i>	<i>Gm.FT2b</i>	6	C
pErecta: <i>Gm.FT5a</i>	<i>Gm.FT5a</i>	4	E
pErecta: <i>Zm.ZCN8</i>	<i>Zm.ZCN8</i>	8	B
pErecta: <i>Nt.FT-like</i>	<i>Nt.FT-like</i>	10	B
pErecta: <i>Le.SFT</i>	<i>Le.SFT</i>	12	D

[00106] The following matrices were collected for phenotypic characterization of plants transformed with each of the constructs listed in Table 6 for expressing other FT homologs with the *pAt.Erecta* promoter, in addition to data collected for the *Gm.FT2a* construct as described above. The individual measurements are as defined above, and phenotypic characterization of transformants was conducted on plants homozygous for the transgene.

[00107] Phenotypic data was collected for plants expressing the *Zm.ZCN8* and *Nt.FT-like* transgenes under the control of the *pAt.Erecta* promoter (see Tables 7 and 8). Trait values for each Event in Tables 7 and 8 are an average of all plants tested containing the Event. A column is also provided with an average of the Event values for each trait.

Table 7. Construct and event level phenotypic data for *Zm.ZCN8* and WT plants.

	<u>WT</u>	<u>Average</u>	<u>Event 1</u>	<u>Event 2</u>	<u>Event 3</u>	<u>Event 4</u>	<u>Event 5</u>	<u>Event 6</u>
DOFR1	33.5	28.6	29	29.2	27.5	27	30.7	28
DOR7	106.9	93.5	97.3	89.2	88.2	93.5	100.3	92.8
PDR1R7	76.5	64.1	69.8	60	59	59.5	71.7	64.8
BRPP	20.1	3.2	2.8	1.3	1.5	1.3	9.5	3
FNBR	190.6	26.9	32	8	7	2.3	95.3	17
FNLP	214.6	54.9	67.5	28	40.5	20.3	132.5	40.5
FNST	24.0	28.3	35.5	22	33.5	18	37.3	23.5
NDBR	211.4	30.2	32.5	9	7.5	3.5	110.8	17.8
NDMS	33.4	30.5	36.3	24	34.3	20	44.8	24
NDPL	244.9	60.3	68.8	30.8	41.8	23.5	155.5	41.8
PDPP	575.8	317.5	498.3	144.8	319	76.3	658	208.8
PFNB	90.4	87.2	98.6	90.3	85.4	64.6	91.1	93.2
PFNN	87.8	93.1	98.1	92.3	97.0	86.5	88.5	96.4
PFNS	71.4	93.2	97.9	92.5	97.9	90.6	82.1	98.1
PODBR	487.3	105.4	162	19	18.5	3.3	384.5	45.3
PODMS	88.4	212.9	336.3	130.5	300.5	73	273.5	163.5
Pods / Node	2.4	5.5	7.2	4.6	7.7	3.2	4.9	5.2
SDPP8	1319.6	564.7	961	200.5	562	136.8	1166.3	361.5
SW1000 (grams)	146	108.9	102.9	127.4	105.3	82.9	116.6	117.9

Table 8. Construct and event level phenotypic data for *Nt.FT-like* and WT plants.

	<u>WT</u>	<u>Average</u>	<u>Event 1</u>	<u>Event 2</u>	<u>Event 3</u>	<u>Event 4</u>	<u>Event 5</u>	<u>Event 6</u>
DOFR1	33.5	31.5	39.3	27.7	25.3	29	37.2	30.7
DOR7	106.9	93.9	115.8	90.7	80.7	83.7	102.2	90.2
PDR1R7	76.5	62.3	76.4	63	55.3	54.7	65	59.5
BRPP	20.1	9.8	20	8.3	2.3	5.3	17	6
FNBR	190.6	108.7	190.5	95.3	11	54.3	223	78.3
FNLP	214.6	131.4	212.3	118	29.5	77.8	248	103
FNST	24.0	23.2	21.8	22.8	21.3	23.5	25	24.8
NDBR	211.4	128.7	281.8	97	11	54.5	247.7	80.5
NDMS	33.4	28.9	33.8	27	23.3	24.8	35.7	28.8
NDPL	244.9	157.1	315.5	124	31.5	79.3	283.3	109.3
PDPP	575.8	462.1	638	511.3	150.8	296	745	431.5
PFNB	90.4	92.5	68.0	98.6	100	99.7	91.3	97.2
PFNN	87.8	89.6	67.6	95.4	93.3	98.3	88.4	94.2
PFNS	71.4	81.9	64.7	83.3	91.3	95.2	70.6	86.2
PODBR	487.3	326.3	529	342.3	22.7	147	633.3	283.5
PODMS	88.4	136.7	109	169	133.8	149	111.7	148
Pods / Node	2.4	3.6	2.0	4.3	4.9	3.8	2.7	4.0
SDPPR8	1319.6	928.7	1359.8	965.5	382.7	591.5	1714.3	558.7
SW1000 (grams)	146	149.0	143.7	121.2	133.8	179.3	142.2	174.0

[00108] Transgenic soybean plants expressing the *Zm.ZCN8* and *Nt.FT-like* proteins flowered earlier than wild type control plants and had an increased number of pods per node (similar to plants expressing the *Gm.FT2a* transgene). Indeed, soybean plants expressing the *Zm.ZCN8* and *Nt.FT-like* transgenes had several phenotypes similar to the *Gm.FT2a* transgenic plants, including reduced number of days to flowering (DOFR1), reduced number of branches (BRPP), fewer nodes per plant (NDPL), fewer nodes on branches (NDBR), reduced number of pods per plant (PDPP), and fewer pods on branches (PODBR), along with an increase in the number of pods per node and a decrease in the number of seeds per plant (Tables 7 and 8), relative to wild type controls. However, several of the negative phenotypes observed in homozygous *Gm.FT2a* plants were less pronounced in the *Zm.ZCN8* and *Nt.FT-like* expressing transgenic plants. Overall, plants expressing the *Zm.ZCN8* transgene had shorter plant height and less branching but more pods per node on the main stem (FIGS. 12A and 12B). Similarly, plants expressing the *Nt.FT-like* transgene had shorter plant height, reduced branching and increased pods per node on the main stem (FIGS. 13A and 13B), relative to wild type control plants.

[00109] Two transgenic *Zm.ZCN8* events and four *Nt.FT-like* events from above were also tested in 2015 field trials at two different locations. Phenotypic data was collected for plants expressing *Zm.ZCN8* and *Nt.FT-like* transgenes under the control of the *pAt.Erecta* promoter (Tables 9 and 10). Events 1 and 2 in Table 9 correspond to Events 2 and 3 in Table 7, and Events 1-4 in Table 10 correspond to Events 1-4 in Table 8, respectively. Except for days to flowering at R1 (DOFR1) and reproductive duration in days from R1 to R8 (PDR1R8), all phenotypic measurements were derived based on data collected from two locations. Similar to the observations in the greenhouse, transgenic soybean plants expressing *Zm.ZCN8* and *Nt.FT-like* proteins also flowered earlier than wild-type control plants in the field. The *Zm.ZCN8* transgenic plants had an increased number of pods per node, while the *Nt.FT-like* plants did not clearly show increased pods per node in the field trial.

Table 9. Phenotypic data from 2015 field trial for *Zm.ZCN8* and WT plants.

	WT	Average	Event 1	Event 2
DOFR1*	42.4	27.9	28.0	27.7
DOR8	110.7	95.0	92.0	98.0
PDR1R8*	65.7	67.1	63.5	70.7
BRPP	2.6	0.1	0.2	0.0
NDBR	9.7	0.3	0.5	0.1
NDMS	18.3	13.6	12.5	14.7
NDPL	28.0	13.9	13.0	14.8
PDPP	44.2	35.1	30.1	40.0
TPBR	9.5	0.3	0.5	0.1
PODMS	34.7	34.7	29.5	39.9
Pods/Node	1.6	2.5	2.3	2.6
SDPPR8	99.9	67.6	54.7	80.5
SW1000 (ounces)	5.1	4.1	3.8	4.3

(* - single location data)

Table 10. Phenotypic data from 2015 field trial for *Nt.FT-like* and WT plants.

	WT	Average	Event 1	Event 2	Event 3	Event 4
DOFR1*	42.4	38.0	42.5	26.8	26.8	25.8
DOR8	110.7	93.3	111.3	88.2	86.6	87.1
PDR1R8*	65.7	63.1	66.8	62.2	60.3	63.0
BRPP	2.6	0.7	2.4	0.1	0.2	0.1
NDBR	9.7	2.7	9.2	0.5	0.8	0.3
NDMS	18.3	11.5	18.3	9.9	7.6	10.1
NDPL	28.0	14.2	27.5	10.4	8.5	10.4
PDPP	44.2	23.5	43.0	18.9	11.6	20.3
TPBR	9.5	2.6	8.4	0.5	0.8	0.5
PODMS	34.7	20.9	34.6	18.5	10.8	19.8
Pods/Node	1.6	1.6	1.6	1.8	1.4	1.7
SDPPR8	99.9	49.9	98.6	36.3	25.0	39.7
SW1000 (ounces)	5.1	4.5	5.1	4.4	4.6	4.0

(* - single location data)

[00110] Additional phenotypic data was collected for plants expressing the *Gm.FT2b* transgene under the control of the *pAt.Erecta* promoter (Table 11).

Table 11. Construct and event level phenotypic data for *Gm.FT2b* and WT plants.

	WT	Average	Event 1	Event 2	Event 3	Event 4	Event 5	Event 6
DOFR1	43.7	34.6	41.2	34.3	22.7	33.2	37.2	39.3
DOR7	105.9	100.4	100.5	100.3	99.8	100.3	98.7	102.8
PDR1R7	62.2	65.8	59.3	66	77.2	67.2	61.5	63.5
BRPP	13.4	4.7	7	5	1.7	3.3	3.7	7.7
FNBR	103.8	32.4	52	29.7	12	30.7	21.7	48.7
FNLP	125.0	46.6	68.7	41.3	24.3	45	37.3	63
FNST	21.2	14.2	16.7	11.7	12.3	14.3	15.7	14.3
NDBR	108.4	34.2	54	30.3	12.7	33.7	24.7	50
NDMS	30.2	18.0	18.3	15.3	15	19	19.3	21
NDPL	138.7	52.2	72.3	45.7	27.7	52.7	44	71
PDPP	387.4	143.0	167	140	96	145.7	108.7	200.7
PFNB	95.5	94.6	96.4	97.7	96.8	91.0	87.7	97.7
PFNN	90.1	89.1	94.9	90.3	88.0	86.0	86.1	89.4
PFNS	69.7	79.2	91.5	74.5	82.5	77.0	81.1	68.7
PODBR	284.9	90.2	109.3	96	43	94.7	55.3	143
PODMS	102.5	52.7	57.7	44	53	51	53.3	57.7
Pods / Node	2.8	2.7	2.3	3.1	3.5	2.8	2.5	2.8
SDPPR8	1159.3	322.3	411.3	292.3	195.3	346.7	245	443.3
SW1000 (grams)	174.0	154.0	170.4	156.7	154.1	155	130.2	157.8

[00111] Transgenic soybean plants expressing the *Gm.FT2b* transgene flowered earlier and had less branching than wild type control plants. *Gm.FT2b* expressing soybean plants had a reduced number of days to flowering (DOFR1), reduced number of branches (BRPP), fewer nodes per plant (NDPL), fewer nodes on branches (NDBR), reduced number of pods per plant (PDPP), and fewer pods on branches (PODBR) (Table 9). However, transgenic *Gm.FT2b* plants did not show an increase in the number of pods per node. Overall, plants expressing the *Gm.FT2b* transgene had shorter plant height and less branching relative to wild type control plants (FIG. 14). Transgenic soybean plants expressing four different events of the *Gm.FT2b* transgene were also tested in 2015 field trials. Phenotypic data was collected for plants expressing the *Gm.FT2b* transgene under the control of the *pAt.Erecta* promoter (Table 12). Events 1-4 in Table 11 correspond to Events 3, 2, 1, and 4 in Table 12, respectively. Similar to the observations in the greenhouse, *Gm.FT2b* expressing soybean plants showed a reduced number of days to flowering (DOFR1) in the field. The other phenotypic measurements also exhibited similar traits as observed in the greenhouse relative to wild-type control plants.

Table 12. Phenotypic data from 2015 field trial for *Gm.FT2b* and WT plants.

	WT	Average	Event 1	Event 2	Event 3	Event 4
DOFR1	41.9	37.3	38.3	38.3	36.3	36.2
DOR8	115.4	109.2	111.1	110.6	110.6	104.6
PDR1R8	73.5	71.8	75.0	72.1	74.1	66.1
SDPPR8	188.5	95.5	99.4	81.1	117.7	83.6
SW1000 (grams)	153.4	137.3	144.0	129.6	134.4	141.5

[00112] Additional phenotypic data was collected from plants expressing the *Le.SFT* transgene under the control of the *pAt.Erecta* promoter (Table 13).

Table 13. Construct and event level phenotypic data for *Le.SFT* and WT plants.

	WT	Average	Event 1	Event 2	Event 3	Event 4	Event 5	Event 6
DOFR1	42.9	41.4	30	44.4	30.7	28	60.2	55
DOR7	108.6	103.8	90.7	106.2	99	91.5	116.5	119
PDR1R7	65.5	64.0	60.2	71	67	65	56.4	64.2
BRPP	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
FNBR	131.5	37.1	2.7	125.2	4.7	1	47.7	41.3
FNLP	156.8	50.7	15.7	142.6	18.1	18.3	56.7	52.7
FNST	25.3	13.7	13	17.7	13.7	17.3	9	11.3
NDBR	140.2	38.9	3	129.4	5.4	1	51	43.7
NDMS	32.4	17.9	16	25.8	16.3	21	12.3	16.3
NDPL	172.4	56.7	19	154.7	21.2	22	63.3	60
PDPP	473.3	201.3	53.7	432.3	69.8	85.7	279.3	287
PFNB	94.1	94.0	100	96.9	83.3	100	92.9	90.7
PFNN	90.4	86.4	82.5	92.3	83.7	83.5	88.8	87.6
PFNS	77.1	76.5	81.2	69.5	84.0	82.8	72.7	68.6
PODBR	366.2	141.8	3.7	361.5	15.5	1.3	238.7	230.7
PODMS	114	60.3	50	73.6	57.1	84.3	40.7	56.3
Pods / Node	2.7	3.6	2.8	2.8	3.3	3.9	4.4	4.8
SDPPR8	1247.4	476.0	136.7	1036	148.5	183	655	697
SW1000 (grams)	167.7	153.2	170.0	182.8	157.5	148.5	131.5	128.8

[00113] Overall, soybean plants expressing the *Le.SFT* transgene had shorter plant height with less branching (FIG. 15) and an increased number of pods per node on average relative to wild type plants (Table 13). However, these effects were variable and event-specific. For example, Events 1, 3 and 4 displayed early flowering (DOFR1), while other events were neutral or actually had delayed flowering. In addition, some of the *Le.SFT* transgenic events showed increased pods per node on average to varying extents, while a

couple of the events were neutral in terms of the average number of pods per node. Interestingly, two of the events (Events 5 and 6) had the greatest number of pods per node on average despite having a delay in flowering.

[00114] Additional phenotypic data was collected from plants expressing the *Gm.FT5a* transgene under the control of the *pAt.Erecta* promoter (Table 14).

Table 14. Construct and event level phenotypic data for *Gm.FT5a* and WT plants.

	<u>WT</u>	<u>Average</u>	<u>Event 1</u>	<u>Event 2</u>	<u>Event 3</u>	<u>Event 4</u>	<u>Event 5</u>
DOFR1	48.2	29.9	32.2	29	28.6	29.2	30.5
DOR7	110	92.5	96.6	90.4	91	92.8	91.8
PDR1R7	61.8	62.7	64.4	61.4	62.4	63.6	61.3
BRPP	12.4	2.5	7	1.7	1	1.3	1.7
FNBR	105.6	7.3	20.3	4.7	3	4	4.3
FNLP	126.5	24.5	41.7	20	18.7	19.3	22.7
FNST	20.9	17.2	21.3	15.3	15.7	15.3	18.3
NDBR	108.6	7.5	21	5	3	4	4.3
NDMS	29	17.7	22	15.7	16.3	16	18.7
NDPL	137.6	25.2	43	20.7	19.3	20	23
PDPP	304.3	131.9	214.7	111	100.3	104.3	129.3
PFNB	97.2	98.0	97.3	93.3	100	100	100
PFNN	98.1	97.0	95.9	90	99.1	100	100
PFNS	72.1	97.0	97.1	98.1	96.1	95.8	97.9
PODBR	233.4	16.5	60.5	8	4	6	4
PODMS	75.1	108.6	159	98.5	95	92.5	98
Pods / Node	2.2	5.2	5.0	5.4	5.2	5.2	5.6
SDPPR8	778.8	271.7	516	232.7	175.3	182.3	252
SW1000 (grams)	151.6	126.0	143.7	122.4	122.2	121.8	116.8

[00115] Transgenic soybean plants expressing the *Gm.FT5a* transgene flowered significantly earlier than wild type control plants and had an increased number of pods per node (similar to plants expressing the *Gm.FT2a* transgene). Indeed, soybean plants expressing the *Gm.FT5a* transgene had several phenotypes (similar to the *Gm.FT2a* transgenic plants), including reduced number of days to flowering (DOFR1), reduced number of branches (BRPP), fewer nodes per plant (NDPL), fewer nodes on branches (NDBR), reduced number of pods per plant (PDPP), and fewer pods on branches (PODBR), along with an increase in the number of pods per node and a decrease in the number of seeds per plant (Table 14). Overall, plants expressing the *Gm.FT2a* transgene had shorter plant height and less branching, but more pods per node (particularly on the main stem) relative to wild type

control plants (FIG. 16).

[00116] Without being bound by any theory, these data support a model of FT overexpression acting in a dosage-dependent manner with the degree or extent of associated phenotypes (e.g., early flowering, increase in pods per node, and altered plant architecture) depending on (i) the level and timing of FT expression, (ii) tissue specificity of FT expression, and (iii) the relative activity and target specificity of the particular FT protein being expressed. For example, expression of the FT protein orthologs from other plant species in soybean may produce a more attenuated effect relative to overexpression of an endogenous FT protein (Gm.FT2a) in soybean, which may result from the non-native FT protein homologs having a lower activity in soybean. However, expression of some native FT proteins may not produce significant phenotypic effects if they have a different or specialized role in their native state or context. Different FT proteins may also act on different tissue targets and receptors and thus have differential effects on the various plant architecture and flowering traits and phenotypes.

[00117] Regardless of the activity level of the particular FT homolog, altered reproductive and plant architecture phenotypes appear to correlate with the timing and location of FT expression. Vegetative-stage expression of FT transgenes may be necessary to induce early flowering and/or cause increased numbers of floral meristems, flowers, pods, etc., per node of the plant. Indeed, FT expression in meristematic tissues during vegetative stages of development is shown with proper dosing of the FT transgene to cause reproductive changes in plants leading to increased numbers of flowers, pods, and/or seeds per node. In contrast, expression of a *Gm.FT2a* transgene under the control of leaf-preferred promoters produced very little, if any, phenotypic changes, relative to wild type plants. These data indicate that both the timing, and tissue specificity (or tissue preference), of FT expression are important factors that affect reproductive and/or yield-related phenotypic changes in transgenic plants.

[00118] The present data suggest that different FT proteins may have different activity levels and/or target specificities despite being expressed using the same *pErecta* promoter. While several constructs expressing *Gm.FT2a*, *Zm.ZCN8*, *Nt.FT-like*, and *Gm.FT5a* each caused early flowering and termination in addition to an increased number of pods per node, other constructs expressing *Gm.FT2b* and *Le.SFT* had different correlative effects on flowering. Expression of *Gm.FT2b* did cause early flowering and termination of plants but

without a significant increase in the number of pods per node. On the other hand, Le.SFT expression showed increased pods per node and early termination despite a delay in flowering. Interestingly, increased numbers of pods per node in transgenic FT plants did not correlate with an extended reproductive duration (PDR1R7) and was not always aligned with early flowering (DOFR1) as noted above. These data suggest that reproductive changes in response to vegetative-stage expression of FT proteins in the meristem may operate through one or more independent mechanisms or pathways. Increased numbers of pods per node in transgenic FT plants may depend on the number of inflorescent and floral meristems induced from vegetative meristems at each node, which may occur independently of flowering time and/or reproductive duration. As noted above, however, reproductive duration may not necessarily correlate with the duration of flowering.

Example 7. Identification of additional vegetative-stage meristem promoters.

[00119] Having observed phenotypic effects with expression of *Gm.FT2a* under the control of a vegetative-stage, meristem-preferred promoter, *pAt.Erecta*, it is contemplated that other vegetative-stage, meristem-preferred (or meristem-specific) promoters may be used to drive expression of FT proteins to cause reproductive or yield-related traits or phenotypes in plants, such as increased number of pods per node (and/or per plant or main stem). Using the characterized expression pattern of the *pAt.Erecta* promoter (see Example 2), other vegetative-stage, meristem-preferred (or meristem-specific) promoters were identified from soybean, potato and *Arabidopsis*. Two bioinformatic approaches were utilized to identify candidate genes from other dicotyledonous species including, for example, *Arabidopsis*, soybean, *Medicago*, potato and tomato, having similar expression profiles to *pAt.Erecta*: BAR EspressoLog and Expression Angler. See, e.g., BAR expressolog identification: expression profile similarity ranking of homologous genes in plant species,” *Plant J* 71(6): 1038-50 (2012); and Toufighi, K et al., “The Botany Array Resource: e-Northerns, Expression Angling, and promoter analyses,” *Plant J* 43(1): 153-163 (2005). The promoter sequences from these genes are thus proposed for use in expressing FT transgenes according to embodiments of the present invention.

[00120] Examples of gene promoters identified by this analysis include the following: four receptor like kinase (RLK) genes from soybean, including Glyma10g38730 (SEQ ID NO: 23), Glyma09g27950 (SEQ ID NO: 24), Glyma06g05900 (SEQ ID NO: 25), and Glyma17g34380 (SEQ ID NO: 26). Additional examples include receptor like kinase (RLK)

gene promoters from potato, PGSC0003DMP400032802 (SEQ ID NO: 27) and PGSC0003DMP400054040 (SEQ ID NO: 28). It is possible that these RLK genes may be related structurally and/or functionally to *Erecta* and *Erecta-like* genes from *Arabidopsis* and other species since they are also RLK genes. Other vegetative stage, meristem-preferred promoters from *Arabidopsis* genes include the following: *At.MYB17* (*At.LMI2*; At3g61250) (SEQ ID NO: 31), *Kinesin-like gene* (At5g55520) (SEQ ID NO: 32), AP2/B3-like genes including *At.REM17* (SEQ ID NO: 33) or *At.REM19*, and *Erecta-like* 1 and 2 genes, *At.Erl1* (SEQ ID NO: 34) and *At.Erl2* (SEQ ID NO: 35). Each of these promoters and similar functional sequences may be operatively linked to a FT gene to cause ectopic expression of FT genes in one or more meristem(s) of plants at least during vegetative stage(s) of development.

[00121] With regard to the *At.MYB17* (*At.LMI2*) gene, see Pastore, JL et al., “LATE MERISTEM IDENTITY 2 acts together with LEAFY to activate *APETALA1*,” *Development* 138: 3189-3198 (2011), the entire contents and disclosure of which are incorporated herein by reference. With regard to the *Kinesin-like* gene, see Fleury, D et al., “The *Arabidopsis thaliana* Homolog of Yeast *BRE1* Has a Function in Cell Cycle Regulation during Early Leaf and Root Growth,” *Plant Cell*, 19(2): 417-432 (2007), the entire contents and disclosure of which are incorporated herein by reference. With regard to the *REM17* and *REM19* *Arabidopsis* genes, see Mantegazza, O et al., “Analysis of the *Arabidopsis* REM gene family predicts functions during flower development,” *Ann Bot* 114(7): 1507-1515 (2014), the entire contents and disclosure of which are incorporated herein by reference. Further, with regard to the *At.Erl2* gene, see “Special Issue: Receptor-like Kinases,” *JIPB* 55(12): 1181-1286 (2013), and particularly Shpak, E., “Diverse Roles of *ERECTA* Family Genes in Plant Development,” *JIPB* 55(12): 1251-1263 (2013), the entire contents and disclosures of which are incorporated herein by reference.

Example 8. Expression of Flowering Locus T gene, *Gm.FT2a*, under control of a *pAt.Er11* promoter alters flowering time and pods per node in soybean.

[00122] A transformation vector containing *Gm.FT2a* under control of the vegetative stage, meristem-preferred *pAt.Er11* promoter (SEQ ID NO: 34) was constructed and used to transform soybeans via *Agrobacterium*-mediated transformation. Transgenic plants generated from these events were characterized for their phenotypes in the greenhouse with a 14 to 14.5 hour natural daylight photoperiod. For each *pAt.Er11:Gm.FT2a* construct, six events were tested (6 plants per event). Six plants were also tested and averaged for wild type (WT) control plants. The following matrices were collected for phenotypic characterization of these plants and expressed as an average for each Event as well as the wild type plants (see Table 15). A column providing an average for all the Events per trait is further provided.

Table 15. Phenotypic data for *pAt.Er11:Gm.FT2a* and WT plants.

	WT	Average	Event 1	Event 2	Event 3	Event 4	Event 5	Event 6
DOFR1	46.1	32.6	40.0	32.3	34.0	29.3	28.2	31.8
DOR7	115.1	99.0	109.7	99.0	99.0	93.0	91.7	101.7
PDR1R7	69.0	66.4	69.7	66.7	65.0	63.7	63.5	69.8
BRPP	23.5	7.4	16.0	6.0	9.7	1.3	4.3	7.3
NDBR	277.6	80.8	215.7	51.7	139.3	3.3	18.0	56.7
NDMS	29.8	32.3	30.7	33.7	32.7	30.7	32.7	33.3
NDPL	307.4	113.0	246.3	85.3	172.0	34.0	50.7	90.0
PDPP	605.8	346.4	447.3	349.7	493.7	240.7	194.7	352.3
PODBR	503.1	173.7	332.3	164.7	323.3	8.3	42.7	171.0
PODMS	103.0	172.7	115.0	185.0	170.3	232.3	152.0	181.3
Pods/Node	1.9	4.0	2.1	4.1	2.9	7.1	3.8	4.0
SDPPR8	1290.0	747.5	1129.0	603.5	881.0	577.0	432.3	862.3
SW1000 (grams)	157.6	157.3	187.7	142.6	173.8	144.3	144.9	150.3

[00123] Transgenic soybean plants expressing a *pAt.Er11::Gm.FT2a* construct flowered earlier than wild type control plants and had an increased number of pods per node (similar to plants expressing the *Gm.FT2a* transgene under control of the *pAt.Erecta* promoter). Indeed, soybean plants expressing *pAt.Er11:Gm.FT2a* had several phenotypes similar to the *pAt.Erecta:Gm.FT2a* transgenic plants, including reduced number of days to flowering (DOFR1), reduced number of days to R7 (DOR7), reduced number of branches (BRPP), fewer nodes per plant (NDPL), a reduced number of pods per plant (PDPP), along

with an increase in the number of pods per node (Table 15), relative to wild type control plants. However, several phenotypes observed in *pAt.Erecta::Gm.FT2a* plants such as number of pods on main stem (PODMS), number of pods on branches (PODBR), and weight of 1000 seeds (SW1000), were less pronounced in the *pAt.Er11::Gm.FT2a* expressing transgenic plants.

[00124] The expression pattern for the *Arabidopsis Erecta-like 1* promoter (*pAt.Er11*) in soybean as measured by GUS staining is more restricted than the expression pattern of *pAt.Erecta* in soybean as described above. *pAt.Er11* drives GUS expression in vegetative axillary meristems and in early floral meristems derived from axillary tissue. However, GUS staining is not observed in the shoot apical meristem at any stage where it can be distinguished from other meristematic tissues of the developing plant. Expression of the GUS reporter under the control of the *pAt.Er11* promoter is not observed in leaf tissue, stem or root at any stage (data not shown). Given that FT expression under the control of either the *pAt.Erecta* or *pAt.Er11* promoter induced early flowering and increased pods per node, vegetative expression of an FT transgene at or near the meristem(s) of a plant may generally be sufficient to induce these reproductive and yield-related phenotypes or traits.

[00125] Having described the present disclosure in detail, it will be apparent that modifications, variations, and equivalent embodiments are possible without departing from the spirit and scope of the present disclosure as described herein and in the appended claims. Furthermore, it should be appreciated that all examples in the present disclosure are provided as non-limiting examples.

[00126] While various embodiments of the present invention have been described above, it should be understood that they have been presented by way of example only, and not by way of limitation. It will be apparent to a person skilled in the relevant art that various changes in form and detail can be made therein without departing from the spirit and scope of the invention. Thus, the present invention should not be limited by any of the above described exemplary embodiments.

[00127] Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

[00128] The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.

WHAT IS CLAIMED IS:

1. A recombinant DNA construct comprising a polynucleotide sequence encoding a florigenic FT protein comprising an amino acid sequence having at least 80% identity to SEQ ID NO: 2 and operably linked to a promoter comprising a polynucleotide sequence having at least 90% identity to SEQ ID NO: 21, SEQ ID NO: 22 or SEQ ID NO: 38..
2. The recombinant DNA construct of claim 1, wherein the florigenic FT protein comprises an amino acid sequence having at least 90% identity to SEQ ID NO: 2.
3. The recombinant DNA construct of claim 2, wherein the florigenic FT protein comprises an amino acid sequence SEQ ID NO: 2 .
4. The recombinant DNA construct of claim 1, wherein the florigenic FT protein is encoded by a polynucleotide comprising at least 90%, identity to SEQ ID NO: 1.
5. The recombinant DNA construct of claim 1, wherein the promoter comprises a polynucleotide sequence that is at least 95%, identical to SEQ ID NO: 21.
6. The recombinant DNA construct of claim 1, wherein the promoter comprises SEQ ID NO: 21
7. The recombinant DNA construct of claim 1, wherein the promoter comprises a polynucleotide sequence having at least 95% identity to SEQ ID NO: 22 or SEQ ID NO: 38.
8. The recombinant DNA construct of claim 1, wherein the promoter comprises SEQ ID NO: 22, or SEQ ID NO: 38.
9. A DNA molecule or vector comprising the recombinant DNA construct of claim 1.
10. A plasmid vector for *Agrobacterium-mediated* transformation comprising the recombinant DNA construct of claim 1.
11. A donor template molecule for site-directed integration comprising the recombinant DNA construct of claim 1.
12. A transgenic plant, or plant part thereof, wherein the transgenic plant or plant part thereof comprises the recombinant DNA construct of claim 1.
13. The transgenic plant, or plant part thereof, of claim 10, wherein the transgenic plant or plant part thereof, is homozygous for the insertion of the recombinant DNA construct.

14. The transgenic plant, or plant part thereof, of claim 12, wherein the transgenic plant or plant part thereof is hemizygous for the insertion of the recombinant DNA construct.
15. The transgenic plant, or plant part thereof, of claim 12, wherein the transgenic plant or a plant grown from the plant part thereof is a short day plant.
16. The transgenic plant, or plant part thereof, of claim 12, wherein the transgenic plant or a plant grown from the plant part thereof is a dicotyledonous plant.
17. The transgenic plant, or plant part thereof, of claim 16, wherein the transgenic plant or a plant grown from the plant part thereof is a leguminous plant.
18. The transgenic plant, or plant part thereof, of claim 17, wherein the transgenic plant or a plant grown from the plant part thereof is soybean.
19. The transgenic plant, or plant part thereof, of claim 18, wherein the transgenic soybean plant or transgenic soybean plant grown from the plant part thereof produces more pods per node than a control plant not having the recombinant DNA construct.
20. The transgenic plant, or plant part thereof, of claim 12, wherein the transgenic plant or a plant grown from the plant part thereof produces more flowers per node than a control plant not having the recombinant DNA construct.
21. The transgenic plant or plant part thereof of claim 12, wherein the transgenic plant or a plant grown from the plant part thereof produces more bolls, siliques, fruits, nuts or pods per node of the transgenic plant than a control plant not having the recombinant DNA construct.
22. The transgenic plant or plant part thereof of claim 12, wherein the transgenic plant or a plant grown from the plant part thereof flowers earlier than a control plant not having the recombinant DNA construct.
23. The transgenic plant or plant part thereof of claim 12, wherein the transgenic plant or a plant grown from the plant part thereof has more floral racemes per node than a control plant not having the recombinant DNA construct.
24. A transgenic plant part comprising the recombinant DNA construct of claim 1.
25. The transgenic plant part of claim 24, wherein the transgenic plant part is one of the following: a seed, fruit, leaf, cotyledon, hypocotyl, meristem, embryo, endosperm, root, shoot,

stem, pod, flower, inflorescence, stalk, pedicel, style, stigma, receptacle, petal, sepal, pollen, anther, filament, ovary, ovule, pericarp, phloem, or vascular tissue.

26. A method for producing a transgenic plant, comprising
 - a) transforming at least one cell of an explant with the recombinant DNA construct of claim 1; and
 - b) regenerating or developing the transgenic plant from the transformed explant.
27. The method of claim 26, further comprising:
 - a) selecting a transgenic plant having one or more of the following traits or phenotypes: earlier flowering, longer reproductive or flowering duration, increased number of flowers per node, increased number of floral racemes per node, increased number of pods, bolls, siliques, fruits, or nuts per node, and increased number of seeds per node, as compared to a control plant not having the recombinant DNA construct.
28. The method of claim 26, wherein the transforming step (a) is carried out via *Agrobacterium-mediated* transformation or microprojectile bombardment of the explant.
29. The method of claim 26, wherein the transforming step (a) comprises site-directed integration of the recombinant DNA construct.
30. A method of planting a transgenic crop plant, comprising:

planting the transgenic crop plant at a higher density in the field, wherein the transgenic crop plant is transformed with a recombinant DNA construct of claim 1.
31. The recombinant DNA construct of claim 1, wherein the florigenic protein comprises the amino acid sequence SEQ ID NO: 2.
32. The recombinant DNA construct of claim 1, wherein the recombinant DNA construct comprises SEQ ID NO: 1.

1/25

	SEQUENCE	1	2	3	4	5	6	7	8	9	10
1	Gm.FT2a	-	90.6 (481)	69.5 (369)	71 (377)	66.7 (354)	60.6 (322)	68.7 (365)	64.2 (341)	73.4 (390)	66.5 (353)
2	Gm.FT2b	90.6 (481)	-	67.6 (359)	69.9 (371)	64.2 (341)	58.4 (310)	65.9 (350)	62 (329)	70.4 (374)	65.9 (350)
3	Nt.FT	70.3 (369)	68.4 (359)	-	76.6 (402)	67.4 (354)	58.5 (307)	67 (352)	62.9 (330)	73.7 (387)	67.8 (356)
4	Le.FT	70.6 (377)	69.5 (371)	75.3 (402)	-	67.4 (360)	60.1 (321)	67.4 (360)	64.6 (345)	73 (390)	66.9 (357)
5	Gm.FT5A	68.2 (354)	65.7 (341)	68.2 (354)	69.4 (360)	-	61.1 (317)	65.9 (342)	62.6 (325)	70.7 (367)	65.7 (341)
6	Zm.ZCN8	61 (322)	58.7 (310)	58.1 (307)	60.8 (321)	60 (317)	-	57 (301)	59.3 (313)	61.4 (324)	56.3 (297)
7	At.FT	69.1 (365)	66.3 (350)	66.7 (352)	68.2 (360)	64.8 (342)	57 (301)	-	65.9 (348)	72.5 (383)	82.2 (434)
8	Os.HD3A	63.1 (341)	60.9 (329)	61.1 (330)	63.9 (345)	60.2 (325)	58 (313)	64.4 (348)	-	71.1 (384)	63.9 (345)
9	Pt.FT	74.3 (390)	71.2 (374)	73.7 (387)	74.3 (390)	69.9 (367)	61.7 (324)	73 (383)	73.1 (384)	-	71.8 (377)
10	At.TSF	66.9 (353)	66.3 (350)	67.4 (356)	67.6 (357)	64.6 (341)	56.3 (297)	82.2 (434)	65.3 (345)	71.4 (377)	-

FIG. 1A

2/25

	SEQUENCE	1	2	3	4	5	6	7	8	9	10
1	Gm.FT2a	-	90.9 (160)	74.4 (131)	79.5 (140)	65.6 (116)	60.2 (106)	72.2 (127)	76.7 (135)	80.7 (142)	71.6 (126)
2	Gm.FT2b	90.9 (160)	-	72.2 (127)	75.6 (133)	61.9 (109)	57.4 (101)	71.6 (126)	71.6 (126)	76.1 (134)	71 (125)
3	Nt.FT	75.3 (131)	73 (127)	-	77.6 (135)	65.5 (114)	59.2 (103)	74.1 (129)	71.8 (125)	77.6 (135)	72.4 (126)
4	Le.FT	79.1 (140)	75.1 (133)	76.3 (135)	-	66.1 (117)	59.3 (105)	75.7 (134)	79.1 (140)	84.7 (150)	75.1 (133)
5	Gm.FT5a	67.4 (116)	63.4 (109)	66.3 (114)	68 (117)	-	58.7 (101)	65.1 (112)	65.1 (112)	69.2 (119)	67.4 (116)
6	Zm.ZCN8	60.6 (106)	57.7 (101)	58.9 (103)	60 (105)	57.7 (101)	-	56.6 (99)	58.3 (102)	62.3 (109)	56.6 (99)
7	At.FT	72.6 (127)	72 (126)	73.7 (129)	76.6 (134)	64 (112)	56.6 (99)	-	71.4 (125)	77.7 (136)	81.7 (143)
8	Os.HD3A	75.4 (135)	70.4 (126)	69.8 (125)	78.2 (140)	62.6 (112)	57 (102)	69.8 (125)	-	79.3 (142)	69.8 (125)
9	Pt.FT	81.6 (142)	77 (134)	77.6 (135)	86.2 (150)	68.4 (119)	62.6 (109)	78.2 (136)	81.6 (142)	-	75.9 (132)
10	At.TSF	72 (126)	71.4 (125)	72 (126)	76 (133)	66.3 (116)	56.6 (99)	81.7 (143)	71.4 (125)	75.4 (132)	-

FIG. 1B

Gm. FT2a	RLVFVLFRLGRQTVYAPGWRQNTKEFAELYNLGLPVAAVYFNIQRESGGRRLY--	176
Gm. FT2b	RIVFVLFRLGRQTVYAPGWRQNTITREFAELYNLGLPVAAVYFNIQRESGGRRLC--	176
Le. FT	RFVFLFRQLGRQTVYAPGWRQNTDRFAELYNLGLPVAAVYFNCQRESGGRRRSAD	177
Pt. FT	RFVFLFRQLGRQTVYAPGWRQNTDRFAEVYNLGLPVAAVYFNCQRESGGRRR---	174
Os. HD3a	RLVFVLFQQLGRQTVYAPGWRQNTKDFAEIYNLGLPVAAVYFNCQREAGSGRRVYP-	179
At. FT	RVVFILFRQLGRQTVYAPGWRQNTREFAEIYNLGLPVAAVFYNCQRESGGRRLL---	175
At. TSF	RIVLVLFRLGRQTVYAPGWRQNTREFAEIYNLGLPVAASYFNCQRENGCGGRRT---	175
Nt. FT	RFVFLFRQLGRQTVYAPGWRQNESTRDFAEVYNLGLPVSAYFNCHRESGTGRRAY--	174
Gm. FT5a	RIVFVLFQQLGRDVTITPEWRHNENSRNFAEINNLL-APVAAAAYANCQRERGCGGRRY---	172
Zm. ZCN8	RLVFVLFRLGRQTVFAPEMRHNENCRSFARQYHLSIATAT-HFNCQREGGSGGRRFREE	175
	*.***:.* ** ** :.* ** ** :.* ** ** :.* ** ** :.* ** ** :.* ** *	

FIG. 1C
CONTINUED

5/25




FIG. 2

6/25




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 3E










FIG. 3F

7/25

**FIG. 3G****FIG. 3H****FIG. 3I****FIG. 3J****FIG. 3K****FIG. 3L**

8/25




FIG. 3M




FIG. 3N














FIG. 3O

9/25

**FIG. 4A****FIG. 4B****FIG. 4C****FIG. 4D****FIG. 4E****FIG. 4F**

10/25

**FIG. 4G****FIG. 4H****FIG. 4I****FIG. 4J****FIG. 4K****FIG. 4L**

11/25



FIG. 4M




FIG. 4N



FIG. 4O

12/25




FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5D




FIG. 5E




FIG. 5F

13/25




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 6D




FIG. 6E




FIG. 6F





FIG. 7

9 DAY ER::FT



27 DAY WILDTYPE






FIG. 8

16/25




HEMI

FIG. 9C



HOMO

FIG. 9B



NULLS

FIG. 9A

17/25




FIG. 10

18/25



pAt.ERECTA-GmFT2a
HOMOZYGOUS



pAt.ERECTA-GmFT2a
HEMIZYGOUS




NULL

FIG. 11

19/25



pER:Zm:ZCN8



WT

FIG. 12A

20/25




pER:Zm.ZCN8




WT

FIG. 12B

21/25



pER:NtLFT-LIKE



WT

FIG. 13A

22/25



pER:NFT-LIKE



WT

FIG. 13B

23/25




WT




pER::FT2b

FIG. 14

24/25



WT




pER::Le.SFT

FIG. 15

25/25



WT



pER::Gm.FT5a

FIG. 16

60760_anniversary_ST25
SEQUENCE LISTING

<110> Monsanto Technology LLC
Brower-Toland, Brent
Caldo, Rico A.
Dai, Shunhong
Gabbert, Karen
Goldshmidt, Alexander
Howell, Miya D.
Karunanandaa, Balasulojini
Manjunath, Sivalinganna
McDill, Bradley W.
Ovadya, Daniel J.
Preuss, Sasha
Rice, Elena A.
Savidge, Beth
Sharma, Vijay K.

<120> COMPOSITIONS AND METHODS FOR ALTERING FLOWERING AND PLANT
ARCHITECTURE TO IMPROVE YIELD POTENTIAL

<130> P34317W000/0016518.00327

<150> US 62/150,142
<151> 2015-04-20

<150> US 62/233,019
<151> 2015-09-25

<160> 38

<170> PatentIn version 3.5

<210> 1
<211> 531
<212> DNA
<213> Glycine max

```

<400> 1
atgcctagtg gaagtaggga tcctctcggt gttgggggag taattgggga tgtattggat      60
ccttttgaat attctattcc tatgaggggt acctacaata acagagatgt cagcaatgga      120
tgtgaattca aaccctcaca agttgtcaac caaccaaggg taaatatcgg tggatgatgac      180
ctcaggaact tctatacttt gattgcgggt gatcccgatg cacctagccc aagtgacccc      240
aatttgagag aatacctcca ttggttggtg actgatatcc cagcaacaac aggggctagt      300
ttcggccatg aggttgtaac atatgaaagt ccaagaccaa tgatggggat tcatcgtttg      360
gtgttttgtgt tatttcgtca actgggtagg gagaccgtgt atgcaccagg atggcgccag      420
aatttcaaca ctaaagaatt tgctgaactt tacaaccttg gattgccagt tgctgctgtc      480
tatttcaaca ttcagaggga atctggttct ggtggaagga gggtatacta a              531

```

<210> 2
<211> 176
<212> PRT
<213> Glycine max

<400> 2

Met Pro Ser Gly Ser Arg Asp Pro Leu Val Val Gly Gly Val Ile Gly
1 5 10 15

60760_anniversary_ST25

Asp Val Leu Asp Pro Phe Glu Tyr Ser Ile Pro Met Arg Val Thr Tyr
20 25 30

Asn Asn Arg Asp Val Ser Asn Gly Cys Glu Phe Lys Pro Ser Gln Val
35 40 45

Val Asn Gln Pro Arg Val Asn Ile Gly Gly Asp Asp Leu Arg Asn Phe
50 55 60

Tyr Thr Leu Ile Ala Val Asp Pro Asp Ala Pro Ser Pro Ser Asp Pro
65 70 75 80

Asn Leu Arg Glu Tyr Leu His Trp Leu Val Thr Asp Ile Pro Ala Thr
85 90 95

Thr Gly Ala Ser Phe Gly His Glu Val Val Thr Tyr Glu Ser Pro Arg
100 105 110

Pro Met Met Gly Ile His Arg Leu Val Phe Val Leu Phe Arg Gln Leu
115 120 125

Gly Arg Glu Thr Val Tyr Ala Pro Gly Trp Arg Gln Asn Phe Asn Thr
130 135 140

Lys Glu Phe Ala Glu Leu Tyr Asn Leu Gly Leu Pro Val Ala Ala Val
145 150 155 160

Tyr Phe Asn Ile Gln Arg Glu Ser Gly Ser Gly Gly Arg Arg Leu Tyr
165 170 175

<210> 3
<211> 519
<212> DNA
<213> Glycine max

<400> 3
atggcacggg agaaccctct tgttattggt ggtgtgattg gggatgttct caaccctttt 60
acaagctccg tttctttgac tgtttcaatc aataataggg cgattagcaa tggcttggaa 120
ctcaggccct ctcaagttgt taatcgccct agggttactg ttggtggtga agacctaaagg 180
accttctaca ctctggttat ggtggatgca gatgcaccta gccctagcaa ccctgtcttg 240
agggaatacc ttactggat ggtgacagat attccagcta ccacaaatgc aagctttggg 300
agagaggttg tgttttatga gagcccgaac cttcagtag ggattcatcg aatcgtgttc 360
gtattgttcc agcaattggg cagagacact gtcacaccc cagaatggcg ccataatttc 420
aattccagaa actttgctga aattaataac cttgcacctg ttgcagcagc ttatgccaac 480
tgccaaagag agcgtggttg cggtggaagg agatattaa 519

<210> 4
<211> 172

60760_anniversary_ST25

<212> PRT
 <213> Glycine max

<400> 4

Met Ala Arg Glu Asn Pro Leu Val Ile Gly Gly Val Ile Gly Asp Val
 1 5 10 15

Leu Asn Pro Phe Thr Ser Ser Val Ser Leu Thr Val Ser Ile Asn Asn
 20 25 30

Arg Ala Ile Ser Asn Gly Leu Glu Leu Arg Pro Ser Gln Val Val Asn
 35 40 45

Arg Pro Arg Val Thr Val Gly Gly Glu Asp Leu Arg Thr Phe Tyr Thr
 50 55 60

Leu Val Met Val Asp Ala Asp Ala Pro Ser Pro Ser Asn Pro Val Leu
 65 70 75 80

Arg Glu Tyr Leu His Trp Met Val Thr Asp Ile Pro Ala Thr Thr Asn
 85 90 95

Ala Ser Phe Gly Arg Glu Val Val Phe Tyr Glu Ser Pro Asn Pro Ser
 100 105 110

Val Gly Ile His Arg Ile Val Phe Val Leu Phe Gln Gln Leu Gly Arg
 115 120 125

Asp Thr Val Ile Thr Pro Glu Trp Arg His Asn Phe Asn Ser Arg Asn
 130 135 140

Phe Ala Glu Ile Asn Asn Leu Ala Pro Val Ala Ala Ala Tyr Ala Asn
 145 150 155 160

Cys Gln Arg Glu Arg Gly Cys Gly Gly Arg Arg Tyr
 165 170

<210> 5
 <211> 531
 <212> DNA
 <213> Glycine max

<400> 5

atgcctcgtg gaagtaggga ccctctagtt gttgggcgtg tgattgggga tgtattggac	60
ccttttgaat gttctattcc tatgagggtc acctacaata acaaagatgt cagcaatgga	120
tgtgaattca aaccctcaca agttgtcaac caaccaagaa taaatatcgg tggtgatgat	180
ttcaggaact tctacacttt gatcgcggtt gatcctgatg cacctagccc aagtgatccc	240
aatttcagag aatacctcca ttggttagta actgacattc cagcaacaac ggggcctact	300
ttcggtcatg aggttgtaac atatgaaaat ccacgaccca tgatggggat ccatcgtata	360
gtctttgtgt tatttcgtca acagggtaga gagacagtgt atgcaccagg atggcgccaa	420

60760_anniversary_ST25

aatttcatta ctagagaatt tgctgaactt tacaatcttg gattgccagt tgctgctgtc 480
 tattttaaca tccagagaga atctggttgt ggtggaagaa ggctatgtta a 531

<210> 6
 <211> 176
 <212> PRT
 <213> Glycine max

<400> 6

Met Pro Arg Gly Ser Arg Asp Pro Leu Val Val Gly Arg Val Ile Gly
 1 5 10 15

Asp Val Leu Asp Pro Phe Glu Cys Ser Ile Pro Met Arg Val Thr Tyr
 20 25 30

Asn Asn Lys Asp Val Ser Asn Gly Cys Glu Phe Lys Pro Ser Gln Val
 35 40 45

Val Asn Gln Pro Arg Ile Asn Ile Gly Gly Asp Asp Phe Arg Asn Phe
 50 55 60

Tyr Thr Leu Ile Ala Val Asp Pro Asp Ala Pro Ser Pro Ser Asp Pro
 65 70 75 80

Asn Phe Arg Glu Tyr Leu His Trp Leu Val Thr Asp Ile Pro Ala Thr
 85 90 95

Thr Gly Pro Thr Phe Gly His Glu Val Val Thr Tyr Glu Asn Pro Arg
 100 105 110

Pro Met Met Gly Ile His Arg Ile Val Phe Val Leu Phe Arg Gln Gln
 115 120 125

Gly Arg Glu Thr Val Tyr Ala Pro Gly Trp Arg Gln Asn Phe Ile Thr
 130 135 140

Arg Glu Phe Ala Glu Leu Tyr Asn Leu Gly Leu Pro Val Ala Ala Val
 145 150 155 160

Tyr Phe Asn Ile Gln Arg Glu Ser Gly Cys Gly Gly Arg Arg Leu Cys
 165 170 175

<210> 7
 <211> 528
 <212> DNA
 <213> Zea mays

<400> 7

atgtcagcaa ccgatcattt gggtatggct cgtgtcatac aggatgtatt ggatcccttt 60

acaccaacca ttccactaag aataacgtac aacaataggc tacttctgcc aagtgtctgag 120

ctaaagccat ccgcggttgt aagtaaacca cgagtcgata tcggtggcag tgacatgagg 180

60760_anniversary_ST25

gctttctaca ccctggtact gattgacccg gatgcccacaa gtccaagcca tccatcacta 240
 agggagtact tgcactggat ggtgacagat attccagaaa caactagtgt caactttggc 300
 caagagctaa tattttatga gaggccggac ccaagatctg gcatccacag gctggtatatt 360
 gtgctgttcc gtcaacttgg caggggggaca gtttttgcac cagaaatgcg ccacaacttc 420
 aactgcagaa gctttgcacg gcaatatcac ctcagcattg ccaccgctac acatttcaac 480
 tgtcaaaggg aaggtggatc cggcggaaga aggtttaggg aagagtag 528

<210> 8
 <211> 175
 <212> PRT
 <213> Zea mays

<400> 8

Met Ser Ala Thr Asp His Leu Val Met Ala Arg Val Ile Gln Asp Val
 1 5 10 15

Leu Asp Pro Phe Thr Pro Thr Ile Pro Leu Arg Ile Thr Tyr Asn Asn
 20 25 30

Arg Leu Leu Leu Pro Ser Ala Glu Leu Lys Pro Ser Ala Val Val Ser
 35 40 45

Lys Pro Arg Val Asp Ile Gly Gly Ser Asp Met Arg Ala Phe Tyr Thr
 50 55 60

Leu Val Leu Ile Asp Pro Asp Ala Pro Ser Pro Ser His Pro Ser Leu
 65 70 75 80

Arg Glu Tyr Leu His Trp Met Val Thr Asp Ile Pro Glu Thr Thr Ser
 85 90 95

Val Asn Phe Gly Gln Glu Leu Ile Phe Tyr Glu Arg Pro Asp Pro Arg
 100 105 110

Ser Gly Ile His Arg Leu Val Phe Val Leu Phe Arg Gln Leu Gly Arg
 115 120 125

Gly Thr Val Phe Ala Pro Glu Met Arg His Asn Phe Asn Cys Arg Ser
 130 135 140

Phe Ala Arg Gln Tyr His Leu Ser Ile Ala Thr Ala Thr His Phe Asn
 145 150 155 160

Cys Gln Arg Glu Gly Gly Ser Gly Gly Arg Arg Phe Arg Glu Glu
 165 170 175

<210> 9
 <211> 525
 <212> DNA

60760_anniversary_ST25

<213> Nicotiana tabacum

<400> 9

```

atgccaagaa tagatccttt gatagttggt cgtgtggttag gagatgtttt agatccattc      60
acaaggctctg ttgatccttag agtggttttac aataataggg aagtcaacaa tgcattgtggc    120
ttgaaacctt ctcaaattgt tacgcaacct agggttcaaa ttggagggga tgatcttcgc      180
aactttttaca ctctgggttat ggtggatcct gatgctccaa gcccaagcaa ccctaacctg    240
agggagtatc tacactggct ggtcacagat atcccagcaa ctacagatac aagcttttga      300
aatgaagtta tatgctacga gaatccacaa ccatcattgg gaattcatcg ctttgttttc      360
gtgttggtttc gacaattggg tcgcgaaact gtgtatgcac caggttggcg tcagaatttc      420
agcacaagag actttgcaga agttttacaat cttggtttgc ccgtttctgc tgtttacttc      480
aattgccata gggagagtgg tactggtggc cgccgcgcac attaa                        525

```

<210> 10

<211> 174

<212> PRT

<213> Nicotiana tabacum

<400> 10

```

Met Pro Arg Ile Asp Pro Leu Ile Val Gly Arg Val Val Gly Asp Val
1           5           10          15

```

```

Leu Asp Pro Phe Thr Arg Ser Val Asp Leu Arg Val Val Tyr Asn Asn
          20          25          30

```

```

Arg Glu Val Asn Asn Ala Cys Gly Leu Lys Pro Ser Gln Ile Val Thr
          35          40          45

```

```

Gln Pro Arg Val Gln Ile Gly Gly Asp Asp Leu Arg Asn Phe Tyr Thr
          50          55          60

```

```

Leu Val Met Val Asp Pro Asp Ala Pro Ser Pro Ser Asn Pro Asn Leu
65          70          75          80

```

```

Arg Glu Tyr Leu His Trp Leu Val Thr Asp Ile Pro Ala Thr Thr Asp
          85          90          95

```

```

Thr Ser Phe Gly Asn Glu Val Ile Cys Tyr Glu Asn Pro Gln Pro Ser
          100          105          110

```

```

Leu Gly Ile His Arg Phe Val Phe Val Leu Phe Arg Gln Leu Gly Arg
          115          120          125

```

```

Glu Thr Val Tyr Ala Pro Gly Trp Arg Gln Asn Phe Ser Thr Arg Asp
          130          135          140

```

```

Phe Ala Glu Val Tyr Asn Leu Gly Leu Pro Val Ser Ala Val Tyr Phe
145          150          155          160

```

60760_anniversary_ST25

Asn Cys His Arg Glu Ser Gly Thr Gly Gly Arg Arg Ala Tyr
165 170

<210> 11
<211> 534
<212> DNA
<213> Solanum lycopersicum

<400> 11
atgcctagag aacgtgatcc tcttggtgtt ggctcgtgtgg taggggatgt attggaccct 60
ttcacaagaa ctattggcct aagagttata tatagagata gagaagttaa taatggatgc 120
gagcttaggc cttcccaagt tattaaccag ccaaggggtg aagttggagg agatgaccta 180
cgtacctttt tcacttttgt tatggtggac cctgatgctc caagtccgag tgatccaaat 240
ctgagagaat accttactg gttggtcacc gatattccag ctaccacagg ttcaagtttt 300
gggcaagaaa tagtgagcta tgaaagtcca agaccatcaa tgggaataca tcgatttgta 360
tttgtattat tcagacaatt aggtcggcaa acagtgtatg ctccaggatg gcgtcagaat 420
ttcaacacaa gagattttgc agaactttat aatcttggtt tacctgttgc tgctgtctat 480
tttaattgtc aaagagagag tggcagtggg ggacgtagaa gatctgctga ttga 534

<210> 12
<211> 177
<212> PRT
<213> Solanum lycopersicum

<400> 12
Met Pro Arg Glu Arg Asp Pro Leu Val Val Gly Arg Val Val Gly Asp
1 5 10 15
Val Leu Asp Pro Phe Thr Arg Thr Ile Gly Leu Arg Val Ile Tyr Arg
20 25 30
Asp Arg Glu Val Asn Asn Gly Cys Glu Leu Arg Pro Ser Gln Val Ile
35 40 45
Asn Gln Pro Arg Val Glu Val Gly Gly Asp Asp Leu Arg Thr Phe Phe
50 55 60
Thr Leu Val Met Val Asp Pro Asp Ala Pro Ser Pro Ser Asp Pro Asn
65 70 75 80
Leu Arg Glu Tyr Leu His Trp Leu Val Thr Asp Ile Pro Ala Thr Thr
85 90 95
Gly Ser Ser Phe Gly Gln Glu Ile Val Ser Tyr Glu Ser Pro Arg Pro
100 105 110
Ser Met Gly Ile His Arg Phe Val Phe Val Leu Phe Arg Gln Leu Gly
115 120 125

60760_anniversary_ST25

Arg Gln Thr Val Tyr Ala Pro Gly Trp Arg Gln Asn Phe Asn Thr Arg
130 135 140

Asp Phe Ala Glu Leu Tyr Asn Leu Gly Leu Pro Val Ala Ala Val Tyr
145 150 155 160

Phe Asn Cys Gln Arg Glu Ser Gly Ser Gly Gly Arg Arg Arg Ser Ala
165 170 175

Asp

<210> 13
<211> 528
<212> DNA
<213> Arabidopsis thaliana

<400> 13
atgtctataa atataagaga ccctcttata gtaagcagag ttgttggaga cgttcttgat 60
ccgtttaata gatcaatcac tctaaagggt acttatggcc aaagagaggt gactaatggc 120
ttggatctaa ggccttctca ggttcaaaac aagccaagag ttgagattgg tggagaagac 180
ctcaggaact tctatacttt ggttatggtg gatccagatg ttccaagtcc tagcaaccct 240
cacctccgag aatatctcca ttggttggtg actgatatcc ctgctacaac tggaacaacc 300
tttggcaatg agattgtgtg ttacgaaaat ccaagtccca ctgcaggaat tcatcgtgtc 360
gtgtttatat tgtttcgaca gcttggcagg caaacagtgt atgcaccagg gtggcgccag 420
aacttcaaca ctgcgcagtt tgctgagatc tacaatctcg gccttcccgt ggccgcagtt 480
ttctacaatt gtcagagga gagtggctgc ggaggaagaa gacttttag 528

<210> 14
<211> 175
<212> PRT
<213> Arabidopsis thaliana

<400> 14

Met Ser Ile Asn Ile Arg Asp Pro Leu Ile Val Ser Arg Val Val Gly
1 5 10 15

Asp Val Leu Asp Pro Phe Asn Arg Ser Ile Thr Leu Lys Val Thr Tyr
20 25 30

Gly Gln Arg Glu Val Thr Asn Gly Leu Asp Leu Arg Pro Ser Gln Val
35 40 45

Gln Asn Lys Pro Arg Val Glu Ile Gly Gly Glu Asp Leu Arg Asn Phe
50 55 60

Tyr Thr Leu Val Met Val Asp Pro Asp Val Pro Ser Pro Ser Asn Pro
65 70 75 80

60760_anniversary_ST25

His Leu Arg Glu Tyr Leu His Trp Leu Val Thr Asp Ile Pro Ala Thr
85 90 95

Thr Gly Thr Thr Phe Gly Asn Glu Ile Val Cys Tyr Glu Asn Pro Ser
100 105 110

Pro Thr Ala Gly Ile His Arg Val Val Phe Ile Leu Phe Arg Gln Leu
115 120 125

Gly Arg Gln Thr Val Tyr Ala Pro Gly Trp Arg Gln Asn Phe Asn Thr
130 135 140

Arg Glu Phe Ala Glu Ile Tyr Asn Leu Gly Leu Pro Val Ala Ala Val
145 150 155 160

Phe Tyr Asn Cys Gln Arg Glu Ser Gly Cys Gly Gly Arg Arg Leu
165 170 175

<210> 15
<211> 528
<212> DNA
<213> Arabidopsis thaliana

<400> 15
atgtctttaa gtcgtagaga tcctcttgtg gtcggcagtg ttgttggaga tgttcttgat 60
cctttcacga gggttggtctc tcttaaggtc acttatggcc atagagaggt tactaatggc 120
ttggatctaa ggccttctca agttctgaac aaaccaatag tggagattgg aggagacgac 180
ttcagaaatt tctacacctt ggttatggtg gatccagatg tgccgagtc aagcaaccct 240
caccaacgag aatatctcca ctggttggtg actgatatac ctgccaccac tggaaatgcc 300
tttggcaatg aggtggtgtg ctacgagagt ccacgtcccc cctcgggaat tcatcgtatt 360
gtgttggtat tgttccggca actcgggaaga caaacggttt atgcaccggg gtggcgccaa 420
cagttcaaca ctctgtagatt tgctgagatc tacaatcttg gtcttcctgt ggctgcctct 480
tacttcaact gccagaggga gaatggctgt gggggaagaa gaacgtag 528

<210> 16
<211> 175
<212> PRT
<213> Arabidopsis thaliana

<400> 16
Met Ser Leu Ser Arg Arg Asp Pro Leu Val Val Gly Ser Val Val Gly
1 5 10 15

Asp Val Leu Asp Pro Phe Thr Arg Leu Val Ser Leu Lys Val Thr Tyr
20 25 30

Gly His Arg Glu Val Thr Asn Gly Leu Asp Leu Arg Pro Ser Gln Val
35 40 45

60760_anniversary_ST25

Leu Asn Lys Pro Ile Val Glu Ile Gly Gly Asp Asp Phe Arg Asn Phe
50 55 60

Tyr Thr Leu Val Met Val Asp Pro Asp Val Pro Ser Pro Ser Asn Pro
65 70 75 80

His Gln Arg Glu Tyr Leu His Trp Leu Val Thr Asp Ile Pro Ala Thr
85 90 95

Thr Gly Asn Ala Phe Gly Asn Glu Val Val Cys Tyr Glu Ser Pro Arg
100 105 110

Pro Pro Ser Gly Ile His Arg Ile Val Leu Val Leu Phe Arg Gln Leu
115 120 125

Gly Arg Gln Thr Val Tyr Ala Pro Gly Trp Arg Gln Gln Phe Asn Thr
130 135 140

Arg Glu Phe Ala Glu Ile Tyr Asn Leu Gly Leu Pro Val Ala Ala Ser
145 150 155 160

Tyr Phe Asn Cys Gln Arg Glu Asn Gly Cys Gly Gly Arg Arg Thr
165 170 175

<210> 17
<211> 540
<212> DNA
<213> Oryza sativa

<400> 17
atggccggaa gtggcagga caggaccct cttgtggttg gtagggttgt gggatgatgtg 60
ctggacgcgt tcgtccggag caccaacctc aagggtcacct atgggtccaa gaccgtgtcc 120
aatggctgctg agtcaagcc gtccatggct acccaccagc ctaggggtcga ggctcgccggc 180
aatgacatga ggacattcta cacccttggt atggttagacc cagatgcacc aagcccaagt 240
gaccctaacc ttagggagta tctacattgg ttggctactg atattcctgg tactactgca 300
gcgtcatttg ggcaagaggt gatgtgctac gagagcccaa ggccaaccat ggggatccac 360
cggctggtgt tcgtgctgtt ccagcagctg gggcgtcaga cagtgtacgc gcccggtggtg 420
cgtcagaact tcaacaccaa ggacttcgcc gagctctaca acctcggtc gccggtcgcc 480
gccgtctact tcaactgcca gcgcgaggca ggctccggcg gcaggagggt ctaccctag 540

<210> 18
<211> 179
<212> PRT
<213> Oryza sativa

<400> 18

Met Ala Gly Ser Gly Arg Asp Arg Asp Pro Leu Val Val Gly Arg Val
1 5 10 15

60760_anniversary_ST25

Val Gly Asp Val Leu Asp Ala Phe Val Arg Ser Thr Asn Leu Lys Val
20 25 30

Thr Tyr Gly Ser Lys Thr Val Ser Asn Gly Cys Glu Leu Lys Pro Ser
35 40 45

Met Val Thr His Gln Pro Arg Val Glu Val Gly Gly Asn Asp Met Arg
50 55 60

Thr Phe Tyr Thr Leu Val Met Val Asp Pro Asp Ala Pro Ser Pro Ser
65 70 75 80

Asp Pro Asn Leu Arg Glu Tyr Leu His Trp Leu Val Thr Asp Ile Pro
85 90 95

Gly Thr Thr Ala Ala Ser Phe Gly Gln Glu Val Met Cys Tyr Glu Ser
100 105 110

Pro Arg Pro Thr Met Gly Ile His Arg Leu Val Phe Val Leu Phe Gln
115 120 125

Gln Leu Gly Arg Gln Thr Val Tyr Ala Pro Gly Trp Arg Gln Asn Phe
130 135 140

Asn Thr Lys Asp Phe Ala Glu Leu Tyr Asn Leu Gly Ser Pro Val Ala
145 150 155 160

Ala Val Tyr Phe Asn Cys Gln Arg Glu Ala Gly Ser Gly Gly Arg Arg
165 170 175

Val Tyr Pro

<210> 19
<211> 525
<212> DNA
<213> Populus trichocarpa

<400> 19
atgtcaaggg acagagatcc tctgagcggt ggccgtgtta taggggacgt gctggacccc 60
ttcacaaagt ctatctccct cagggtcact tacagctcca gagaggtcaa caatggttgc 120
gagctcaagc cctctcaggt tgccaaccag cctaggggtg atattggcgg ggaagatcta 180
aggaccttct acactctggt tatggtggac cctgatgcac ccagcccaag tgaccccagc 240
ctaagagaat atttgattg gttggtgact gatattccag caacaactgg ggcaagcttt 300
ggccatgaaa ctgtgtgcta tgagagcccg aggccgacaa tgggaattca tcggtttgtt 360
ttcgtcttgt ttcggcaact gggcaggcaa actgtgtatg cccctgggtg gcgccagaac 420
ttcaacacca gagactttgc tgaggtctac aatcttgat cgccagtggc tgctgtttat 480
ttcaactgcc agaggggagag tggctctggt ggtaggaggc gataa 525

60760_anniversary_ST25

<210> 20
 <211> 174
 <212> PRT
 <213> Populus trichocarpa

<400> 20

Met Ser Arg Asp Arg Asp Pro Leu Ser Val Gly Arg Val Ile Gly Asp
 1 5 10 15

Val Leu Asp Pro Phe Thr Lys Ser Ile Ser Leu Arg Val Thr Tyr Ser
 20 25 30

Ser Arg Glu Val Asn Asn Gly Cys Glu Leu Lys Pro Ser Gln Val Ala
 35 40 45

Asn Gln Pro Arg Val Asp Ile Gly Gly Glu Asp Leu Arg Thr Phe Tyr
 50 55 60

Thr Leu Val Met Val Asp Pro Asp Ala Pro Ser Pro Ser Asp Pro Ser
 65 70 75 80

Leu Arg Glu Tyr Leu His Trp Leu Val Thr Asp Ile Pro Ala Thr Thr
 85 90 95

Gly Ala Ser Phe Gly His Glu Thr Val Cys Tyr Glu Ser Pro Arg Pro
 100 105 110

Thr Met Gly Ile His Arg Phe Val Phe Val Leu Phe Arg Gln Leu Gly
 115 120 125

Arg Gln Thr Val Tyr Ala Pro Gly Trp Arg Gln Asn Phe Asn Thr Arg
 130 135 140

Asp Phe Ala Glu Val Tyr Asn Leu Gly Ser Pro Val Ala Ala Val Tyr
 145 150 155 160

Phe Asn Cys Gln Arg Glu Ser Gly Ser Gly Gly Arg Arg Arg
 165 170

<210> 21
 <211> 901
 <212> DNA
 <213> Arabidopsis thaliana

<400> 21

aaaccgaccg gagccaacca aaccgggttaa catcctaaaa ccaatcatat tttattaagt 60

tttgtgttga tgctaaacca aaaatcattg gcatgcatat ttctaaattt agtaataaac 120

aaaaacactt agaaatcaca cgttcactat actaaaaaac gttgacaaaa acacaacaac 180

tatactaata attaaagaag agaaaactga accaaacttt ttgtaaactc ctgaatttaa 240

attagtaatt gaagtaagaa gatgaagaag aacatgttaa gcaaacaaaa aaattacact 300

60760_anniversary_ST25

```

aaaatcatat aaaaatacat aattacaaaa gtacccataa gatggattta ttgatatggg      360
tcattctgtga aacaagccac agagagacaa agactcgtaa gtattgggca acgaaagcga      420
cctcctttat tcaccactgc cattaacatg ttcttcttct ccttcttctt ctacatttta      480
tgaccgtttt acccttcaag agagagaaac aaaatcactc cctctcactc actctatctc      540
tctcttctgc aaagcttcag aactctggca gagagataaa agatgatggg gtttttaact      600
ttatcctccc caaataattc ttcttccctt catctctctc tcttacacaa caggtcctta      660
catttgtaca atctcctctc tttaaagact ctctctcttt ctctctccat ctctatctta      720
ctctgtatth ctgtcgtctg agcactcaat gaaaccactg taaatttccg ccagaatttg      780
atgtgatgga acgataaaaa tcattttttc tcgggttaaag taaaaaaaca aaaacaaatt      840
tctgtagaaa tcataataaa agaaagaaaa aaaatctaag gtcggtacat aatacggttc      900
t                                                                                   901

```

<210> 22
 <211> 507
 <212> DNA
 <213> Arabidopsis thaliana

```

<400> 22
aaaccgaccg gagccaacca aaccggttaa catcctaaaa ccaatcatat tttattaagt      60
tttgtgttga tgctaaacca aaaatcattg gcatgcataat ttctaaattt agtaataaac      120
aaaaacactt agaaatcaca cgttcactat actaaaaaac gttgacaaaa acacaacaac      180
tatactaata attaaagaag agaaaactga accaaacttt ttgtaaactc ctgaatttaa      240
attagtaatt gcacaacagg tccctacatt tgtacaatct cctctcttta aagactctct      300
ctctttctct ctccatctct atcttactct gtatttctgt cgtctgagca ctcaatgaaa      360
ccactgtaaa tttccgccag aatttgatgt gatggaacga taaaaatcat tttttctcgg      420
ttaaagtaaa aaaacaaaaa caaatttctg tagaaatcat aataaaaagaa agaaaaaaaa      480
tctaattgtc gtacataata cggtttct                                                                                   507

```

<210> 23
 <211> 2000
 <212> DNA
 <213> Glycine max

```

<400> 23
catgagcaag taagtaaaca ttttatctct gttacactcc aaacacatac actaacttaa      60
gcagagtcct ttttcggcta ttcactccat caccaaagaa tgagctcatc cataaaaaaca      120
tatacatgtc aacatgtcga aagatgtaga acctggtaac aataataaag aatgataatt      180
tttttattht taaaacgaag cactaaaata gatttaatta tttttatcat ataaaattac      240
caattaatca tataatctagt gaaatgatca cttcaagtat aaaaaaagat tttatacttt      300
tcgtctttct gtatttagaa atattagcaa agtcttggat aaaaaagtgg tcgtagcctg      360
tccctgtagt ttccttgtha gtcttgatga acaagaagtt ctcagttctc cccaccctat      420

```

60760_anniversary_ST25

tctgattcgg	tttacatgga	agtagtaagt	aaccatacac	cattatagaa	ataatacgat	480
aaccacacgc	catgtcttac	ctcatgcgtt	tactgaagtt	gttttcttct	tttatttttc	540
tttgatggag	ttatggtatt	aatattcaat	attagattgg	aacttgcaga	tcaacttcaa	600
gaggcactct	ttgataagga	taccccaggc	attgcttttt	gacataacgc	caaaaccctc	660
ctaaaaaacc	cttcattttc	tatctcttag	ttccatttta	tgcaatgaaa	taaaacttct	720
caaatagtgt	caaagcccg	aaattgccta	ccatatattt	atcacgattc	atgaagggt	780
tcttaattcc	tttttttttt	ttttcctgaa	atgttttttt	tgaaggaatt	ttcttgaaat	840
gttttaatgc	cttttttttt	actcaagaaa	tgtcttaatg	cttgtttact	tacataaagt	900
aatatcgttt	gtcttttttt	acgcaatatt	atattctagt	actctgtctc	tccaatctta	960
ttatttttaa	aatttttctt	ccttcctatc	ctattatgca	caaaaagggtg	taattttaac	1020
atttttctac	tagtaaaaaa	cctacaaact	ttttctattg	ttaaaattaa	atataaaagt	1080
aattattttt	attttatata	aaaatacaaa	gattttatgg	aaaaatatgt	aagatataaa	1140
aatatgatta	attattttac	tttcatctta	acttagcaaa	tactttctga	tcagtgccct	1200
atctcgcaca	atccacaaac	attatctcgc	acaagccaca	aacacgcgct	tcatgatcca	1260
aaattgtacg	agcgacgctg	tccatgtctc	ctagaacgcg	cgtagtaaga	aataagtgtc	1320
cctttgattt	catgttgcat	agttaatttt	tagtttaaga	ttaatttttag	atagttttcc	1380
atatttttaa	ttttatatta	aagataaagt	caaaattaat	gtttagaatt	aattgagttt	1440
aagtcatttt	aggtaccctt	tggataaaga	aatttaaatt	aaattttaat	tcaaaattaa	1500
tataaaccaa	aattattaaa	acataaatca	tattacttta	aaattaattt	ttcgaaaaag	1560
cacattcaaa	gctccactaa	aaattgtctt	tgattaatag	ccgtggatga	gatttgattt	1620
attagtgaga	aaagacaaag	aggtttaagc	gcacgcgaag	agaggcgcgt	aagtaaatag	1680
gagaaacttt	agctgtcaaa	tatgctggga	aacggcgagt	acgaatgacg	gcggctacca	1740
cccttatatt	acagtgcacg	tctcactctc	acctatctag	cctaacgtcg	cttcaccgcc	1800
gtttcccatt	cttattctct	ctcttcataa	cactcttcct	atttacagtt	cacgccaaat	1860
gcctgcactc	tttctctact	attaccaagc	attggccaca	ccaacaccaa	cgaataacct	1920
ttgttattgt	aactaataac	cactgcattt	ttccatact	cgttgatctc	ttccactaag	1980
tgctgtgggt	ggtgaccgca					2000

<210> 24
 <211> 2479
 <212> DNA
 <213> Glycine max

<400> 24	
ttggattgtg	tagaagaaaa aaattgaaaa aaaatatgac atccatgtta atttttgaaa 60
ttttcttata	ataattttcc gtcttcaacc aaaccagact cctattaagt atagtaaata 120
ctattatatt	tttgtggact tcttaataaa tagtagtaat attttgaaaa gcttttttat 180

60760_anniversary_ST25

ttttacgaaa	gtcataataa	attataaaca	aactttataaa	aaatattaaa	tctttatagta	240
ctactttttat	ataataataa	taataataat	aataataata	ataataataa	taaaacgtgt	300
tcaaaattat	taatattcct	cttgaagttc	cgtttcatat	tctgtaaaaa	aaagttgcgt	360
ttgatattaa	aatataacag	tactaaaaaa	acaacataaa	aaagaaaatc	atgtttgatg	420
aaagaaaata	caaatatatt	ttcataaaga	gaacttcaca	attactcgca	atgctgtgtg	480
aaatagggat	ataaccttta	tccaagacac	gttcccatca	ttgaagtata	attaaatcct	540
ttacggttaa	ttataatgaa	atcatttttg	atttgctttt	gcctattatc	acttttcaca	600
cgatgatact	taattattca	tagacctttt	tgtcgagtaa	gaggggaaat	gctaaacttt	660
tctgcttaga	ttttttggca	tagttaatgg	attttagcct	ttttctttct	tattaatttt	720
tttctttcat	aagcatagtc	ccggtaaaa	tctcactttc	agttgatact	ttacctcctc	780
cgaaaagttt	cccatattag	agactcaatg	gcgtataaaa	tcactttaaa	catttactta	840
taatgaatgg	aaataaaaatc	taaaaagtta	gctactaatt	ctttcacggc	cattacgaag	900
actttgctta	aaaatggaaa	aaaagcaaaa	tataaaagag	tgtacattgt	ctatttttat	960
aattgacttg	gctctgtatg	tattatgtaa	ttaattttta	atcttatatt	ttgattattt	1020
atagagatat	aaaatgaatt	tgatagttaa	agaaagaaaa	gaagagatga	aaaattgtgt	1080
gttcgatcct	ctaataaaac	taacatttta	ataaattaat	atttatcatt	tttttaatta	1140
tttgagtttt	ggaagtgtaa	tgagtcgagt	aattttattt	gatggttgtt	tggttcactt	1200
atctatgttg	atcaagtaat	cgatcaattt	atctccatga	ataatgatga	tttttaagaa	1260
tatttaacat	ttgaccatca	attccttaaa	tcagtgaatt	atttttgtca	accatgcaac	1320
ctctataaat	ataggtccta	catatgtttg	acattcatca	tagtgtgtaa	tgtatttttt	1380
ttattaaaaa	aaacagagat	gatgaactcg	tgataaagaa	tcacctaaca	cattactgat	1440
actctctata	aatatcacat	gacaacctta	aacaaatag	cacaattcat	atatcaatat	1500
ccattacttt	gtcatattct	aatttgagtg	taaaagtctt	tattattaca	gtcttttaag	1560
ttggtttaga	gcaatttgag	ttaatatctt	tatacaaaaa	taacttaatt	ttttaatatt	1620
atttttaaga	catatttctc	ataaaaaatc	acataattta	gtttataatt	tttaatttaa	1680
tattatcttc	atttttattt	ataaaattta	attaccta	tcataaatat	taaaaaaata	1740
aattaattta	attaatatca	ataaatttat	cctaaactta	taaacattat	cattaatgct	1800
cttctctctt	aaatgtttat	ggatataact	tcttttat	aattaaaatg	tttattttaa	1860
attaaattaa	tgtaagaaat	aatacaaatt	gaatattgta	taaaggaaca	aacataat	1920
tgttttgtat	tagaccataa	gtaatactcc	atattagatt	atatatataa	cttttatttt	1980
aaaattatag	agtatacttt	ttttagagga	aattatagag	caactacat	tcatatgatt	2040
tctcttttat	aaatattgaa	aacaaaatag	ggatatgcaa	cagcaaacga	gggaggtttg	2100
aggagagagg	gagagagaga	gaatgtaggc	gcgtgtggca	cagttatgag	ttaagactta	2160
ggagaagtac	acattggcat	aggcattgtt	attggattat	gtgtagagtc	cgatagacta	2220

60760_anniversary_ST25

gaatgacggc tactagttac tactctctct cttcataaac acaccattta tgtttttccc	2280
ttcccttcac gccaaacgcc tgcactctac actctactct ctcgtgctct gtgactactg	2340
tcactctctc ataaacccaa catgccctta atccattttc catagtagtt agtggttgta	2400
ctcatctctt ccactttcaa tctctcttct ttccttattg ttgctcacca aggtgggggtt	2460
ttttgtacgt gtggtggca	2479

<210> 25
 <211> 2000
 <212> DNA
 <213> Glycine max

<400> 25	
cattaaataa ttctaaaaaa gatataaatt tttgtataca aatctatatt taagaaactt	60
ttaatctaga tgtcgatttt aaaaaatatt atttaattaa aaaatattag atggtgtaat	120
aattaatcaa aattatatca agataatctg attcctttct atacacacat aatattattt	180
catccttagt ccctaataatt ttcaattctc attttgttac cagacttgta ccgaacaaaa	240
acaaaatatac taatcatagt ttctattcaa caaaaatgat ttttaactca atttgaaaca	300
cttttcattc atttttaaaa ctaagaaaaa atttgtgatt tttttttata attttgaaaa	360
acattctacc atcattatta ttgtttctac taccatcatt attaatgta ctactatcac	420
cttcctagtt ataaccgtaa gcattatata tttttattat tattgttatt atgttatttt	480
gttaatatat ttatttttgt tctaaaaaat tattatTTTT tcatatcttt cactatTTTT	540
gttattattt tagcaagttt gattatTTTT tattttaaat atttttatgt gtcactTTTT	600
atatcacatt atttaacagt gtgaatcgat aaaaaatata ataattatct ttaatttgta	660
agaatTTTT caaaattaaa actgatttta gttcttgaaa aatgtaaaac taaaaatgaa	720
aaccacctaa cggggcctta gtaattagga catggtctcc ctgggttacc acgggatttt	780
ttcacatcaa agaagacctg gtattttcat tttcatgaga tttttgcata tcgaacaagg	840
cattaagaca ggggttgta ttgtcgtgat agtataatTT acatggtcga agtgatagaa	900
actttaacca tcatttacct tgtaccttac tataacacaa aatactacga tttccaaaca	960
ctagatcgcg cgcttatgtt ttcagacaca ttattcttct tcattcataa ataaatttgc	1020
agctagtata tgataattgt accaatttat gtaagtTTTT tacaaaggac attcttatct	1080
caataaaaaa ctaaattgtt aaaatattct ctagcacatt ttttaaatat attttgtcta	1140
attaattaaa attaaaagag gatataaaaa atatgctgct aacatcttga acatttccgc	1200
aatcaataat ttctcaatct atctgaatat ttttgcaact gtatacaaaa atctcagaac	1260
agaaaattat tgattaaact ggaagaattt aataacattt gattcacgtt tgttttagtga	1320
ttaaaaaatc ataacattac actatctaac aaatgcagca tccataacta ccaaacatta	1380
aacaagagaa acagacaaag tccaataatc acagagacac gcagtgacaa agaaaagaaa	1440
gagggaaacgg taaagagaaa ggtgtctctg tcatctcaaa tagattgcca taactccctc	1500
cttctctctc acaagctctt gcagagtgaag agcgaccact ttccgatctc aattaaaagt	1560

60760_anniversary_ST25

atggcataat	ttgcaatggc	ggaactgaac	gaataataat	aagagatacc	atagttaaga	1620
gagagaaaca	caaacatgga	aaaagctggg	cctcactccc	tgggtacaca	tagatagaga	1680
ctatggtgca	gtgttgacagg	ttgttagcaga	agctctgccca	aatagtgtta	actttattcg	1740
agaaaattat	tattattatt	attattatta	ttattctctc	tctctagtct	attatcagtg	1800
gtaattcagt	aatgtttgtg	cattatagag	agagcgtggg	ctatgtgccca	gggtgatgtg	1860
atgtcatttc	actaccttca	aagccagaaa	aatgcaacag	aaaaagcttt	catcccatca	1920
catcatttga	accatgaatc	atgaactagt	tttctaaact	aaaactataa	caacaccttc	1980
ggttggttgtt	gttggttggt					2000

<210> 26
 <211> 2000
 <212> DNA
 <213> Glycine max

<400> 26						
aacatttaag	atcttaaaga	tgccaagagc	ttcatatgaa	aatgtacaaa	agagatttta	60
aaggcaatat	caatgctgtg	acgccatatt	aaaataaaag	ggatgggtttc	tcctgtatat	120
tgagcaattt	gtattactta	tatacacaaa	atctaaattg	attcttaaca	aatatgtaaa	180
gaaattaata	atatgatcaa	gttacctgaa	gaagctaaaa	taaaatagaa	aattaagtaa	240
aagaaatgag	gagtagaata	taagataaca	tcaaaaaatt	atttcagcat	attttaagaa	300
catcaaattt	acctttcatc	aaaattaatc	ttaaaagact	aaaacattta	attaagttta	360
taaatactca	cacaaaatat	taattttatt	tgtaattatt	attttttata	tttttattta	420
ctattgcctc	aaaatttgca	ctaaacaaga	gaccctagag	atttcgttag	aacaataata	480
gacacggtat	taaataatta	aattaatacg	aggatgcata	actaccaaac	aaatgcgata	540
aataaatgag	acgacgagag	agcacaacgc	gggaatgaga	taattaagaa	aaaaaatcta	600
ataaattagg	aaaaaaaaaga	cataatatca	taagcttgaa	tccaatgtac	aaagagaggt	660
tggcaataaa	gagaaagaga	aaagacgtcc	ctgtcacctc	aaatggattg	cattactcat	720
tgaaaaggac	attattactt	ccgacttttt	atattaactt	actaattata	aaatatataa	780
aaaaatactt	caaagatgca	tatatTTTTAT	tttattacat	aattacataa	cagaataata	840
taaaataatg	taactacaca	ttaaaacatt	aaaatagtga	ttggagtagt	ggtataagag	900
gacgttgaat	tcacgcggaa	gagaaggata	tatttcatgt	ttaatttggt	gtcatgccta	960
gttcaatgta	atctaataag	taaaaataaa	atacaaaca	aataaagatt	ttggtttctt	1020
aacaaaagta	cttttacttt	aaatatatat	ttttatctgg	tttttaaaca	tgcacatatt	1080
taacataaaa	gttcatatta	aactttttcc	tacatacttg	gatcaaatag	tcacgtattg	1140
caggtaaaaa	ataatagtgt	agcttataga	aatcgtagaa	ataagtctat	aaaccagaag	1200
aaaaaaaaaca	ttaaaataat	agtatagaaa	tctatatcag	tgtccccagt	tcttacattc	1260
atgacccatt	tccccataaa	ctctttgcag	ataatgcaat	ggcaaaacca	cacagaaagt	1320

60760_anniversary_ST25

gacccctggg	aatcaaaagt	taaaaccaat	ggcacagcat	agcacagtgt	acagtgttta	1380
tttactatat	agcaaaacac	tcactggcat	aacacttttag	ggagagagag	agtgaaaaca	1440
agtgtaaaaa	gagagaaagt	taggaggggg	atagagagtg	tgtgtgtgtg	cagagtttgc	1500
aggcttgtag	cagaaatggg	ggcagatggg	tttaacttta	tgtgtgaaat	aattttcttc	1560
tatctctttt	ctcttttagtg	ttttctctct	ctctctctct	cttctttttc	ttcctgcatc	1620
ttcttgtgtt	tagggagtgt	gatgttttgt	ggcagaagaa	cgatgtgatt	ggacacagcc	1680
aaagctgtgg	acttgttctg	ttactacttt	gtaattgtaa	tcacataaaa	ggctagaggg	1740
tatgaagagt	gcacagaaaa	atactagtac	tagtttcaaa	caaaactcac	cttactacta	1800
cccttccatc	tcaagccata	gttgagttga	gtggtgcaca	gtgtcactat	acataccact	1860
aacacccttt	tttggttctt	gttctgtggc	tccttgtgct	ttgagcaaga	gctttttgag	1920
aaagagcttg	gtggtgggtg	ttgttgttga	gtggtttcat	ggttaggctg	ttgttaagtt	1980
gaagttcatc	agttgcagct					2000

<210> 27
 <211> 1399
 <212> DNA
 <213> Solanum tuberosum

<400> 27	
aacgaaaaat	ttagaaacta ttagtgatcc aaatgttcgt gattacctgc aacgagaaca 60
acaacgaata	cttgaaaaaa gaaatcgaca atcacaacca caaccataat cgcaacaatt 120
ctcagaatca	tatcctaatt ttttttcgaa tagtgctaaa ttgaaaacg acctaccgaa 180
tttctaaatt	attgtttgtga tcaattaatt attatgttat gtattgtatt ttatcttgta 240
tttaaattat	tatgttatgt attgtattgt tatcttgtat ttaaattatc atatcatgta 300
ttgtattttt	aaattaattt tttttgcata ttctttataa tgaaaattaa taataaaaca 360
attttattat	tcacgaaaaat tagaaaaaaa gttaaaatac tattaatttg aaattaaaat 420
agtatatatt	aaataatatt tttaaaaata ttatattata tttaaaaaga attatgaata 480
ttagatattt	aattaatgga attatatgta aaataatatg ttaattagaa agtaatagaa 540
aaataataaa	ataatgaaaa agtagaaata gagagtgtga atagtagaat ttggagaact 600
attcaactct	ctaaatttga agaatatagg gtgatttgga ggtgggttgg agtgtccatt 660
ctctatttta	ctctcaaaat atagagaatg gagagaaaaa tagaggtgga ttggagatgg 720
tcttagtgac	atttttgatt ccgccaatgc tcagttggcg tagtcgctgt caaacttgag 780
aaaggattac	ccctttaggc ttgcacagac agtgacttat gatgaaatga agccagagaa 840
ggcactctgt	tataacactt aaatgaaaat acatgtgtat ggactagcaa taaaaggggc 900
actagtaatt	ttagtaattg aaaagcaagt gtatagagag agataatgag agagaaagag 960
taagtacact	actactgcta ctatcccata tagctgtaat gttgcaggtc tgatttttgc 1020
agttgcagac	ccccttcttg gcacaagctc ttttaacttt tatctttctca aataattctc 1080
tctctctctc	tctctctttt ttctcttttt acattgtgag gaaactgaat acccattgta 1140

60760_anniversary_ST25

tgtattagtg	tgaggcctat	ctgccacaag	gatgtgatgg	aacactatgc	ttcctctgct	1200
aaaacccccac	aacccccaaaa	ctcttttttca	cttcacattt	aatcacaatt	cctcagtgaa	1260
attattctgt	tgctctctct	aatttcaatt	tcaatgtcgg	taagtccaag	aactggtttt	1320
tcaattcaaa	ggagctgagt	tagtgcaaac	acttgaggtt	ttgagttttg	acagagactt	1380
gagtctcaga	gaaactacc					1399

<210> 28
 <211> 2000
 <212> DNA
 <213> Solanum tuberosum

<400> 28						
accttatata	agttacaatt	tagttatgta	tataagttaa	aattaaatta	aaagacattt	60
cgaaataata	tgattatacc	atttcgaaat	taattagaga	gagaaataag	atctcgcaaa	120
attaagtgtc	ttcttgaaat	taagaaccat	ttttaggaga	taattatgta	ttttttcatt	180
tttaatttga	cacgtatgca	tatccactat	tttgttttat	tccaaagtga	cccctacttc	240
ttttggtaat	ttctttgagt	attttaaact	ctagtccccc	tttctcaagc	aaaaaggctc	300
actcgcgcac	gcgcgagag	acattgtgac	gcgctggatg	gaaaatccag	aagcgtaact	360
gtcaaaaaat	agaacaactt	tgggaaacgg	ggtgacggcc	gctgccacca	cttttttcat	420
ttccaaacac	tcattaacta	acgtcgtttc	accgccgttt	actgcttaat	gagtatgaat	480
tacactctaa	tagtctattt	ttacttattt	ttaatgtgtt	tatcaaatta	tattttttaa	540
tataatactt	taaaaatatt	atcatcaata	ataagagtaa	attaaaaaat	aaatgacaaa	600
ttgtttctta	aattgttaaa	ttaaacaatt	aaaactgaat	atttacaaaa	tacctcttaa	660
cttgctaaat	taaacaattg	aaactatatt	tatattaata	aattgaactg	acaaaaataa	720
ataaaggaac	tatatatfff	ctcaattata	tctttttact	aaaatattat	ttttctaata	780
ctagttaaac	ttttaaaaaa	catctaataa	agaaaaagaa	tttgttcaat	tatactttag	840
aagcttttat	tattattatt	attattagta	gtagtagtag	tagtaataaa	ttagattaaa	900
ttaaagagag	aagtattcaa	aactcccaa	actattgtat	tagttttatt	tcagaactat	960
tgacaatctt	aatttttttt	tttttaattt	gactagggtga	acttaaatat	acttcatttt	1020
ttgcaaaaca	agtgaagtac	actcttaa	tttcatcaag	tttagaaatg	ttttcaacaa	1080
tttactagac	tctttattaa	gaacttcatg	ttctttcaag	agtttatgag	cacttgctat	1140
gtcatgttac	agatcaagaa	tatctacaga	gtgtatctaa	atttagtact	agtaaagtag	1200
aaaatgtatt	acttatctct	caaacaatag	gtattcatta	tactatfff	agatgtccaa	1260
caattttttt	tcactttatg	aatcaatga	ataatttaac	acttagttcc	taattcccag	1320
taagcattaa	ttatagttat	ttacttatta	tatttttcaa	cacattatat	tgaaaaagtg	1380
atatagtaaa	tctatctfff	tattttatta	tttcttaaaa	tttgtacaaa	cttaataata	1440
gacaaatatt	gttgaatagg	aataataatt	tacattaaat	ccaatatatt	tttcaatagt	1500

60760_anniversary_ST25

tgtcactaaa	tgaaaatact	tcatctgttt	caatttatgt	gatagttttc	atTTTTcaaa	1560
agtcagacaa	ttatatatTT	ataaattaag	taaaaaatat	tataagtcac	actaattaac	1620
aattcgaaat	attcggtacg	gaggaactaa	cacttatgtt	tttagaccat	attagtcTTT	1680
tctctctatt	tattatataa	tattgagagg	agagtgcAAC	caccatggca	actttctctg	1740
tcttcataaa	acgcagctga	cattaaaaac	acagacacac	acttcgcatt	tcatatccct	1800
ctcactacac	gccaaatgcc	tgctcttcct	atttctcttc	ttcttctttt	tcttcttctc	1860
tctcattcac	ataacacaca	ttcttgtact	aactctgcat	cataaactct	acccactttt	1920
cttcttcttc	tccggtcata	ttgctctgaa	actccactta	ttgctctctc	ccggcattta	1980
tttttagttt	ctcagaaata					2000

<210> 29
 <211> 471
 <212> DNA
 <213> Arabidopsis thaliana

<400> 29	
taataagaga	cgaaaaaaaa
ataactaact	gatcattacc
atccataaat	aaatagttgc
	60
tgccataaac	caaacacatt
gtgcttatca	aaaagaagaa
atttgtactt	aatgaaacat
	120
tcattattag	caaagtgtaa
aaccaaagaa	aaacaaactt
tatttctcat	tttattagta
	180
aaagtgaaga	agagtaaaga
aaaagagaga	ctgagatgag
gctgagagcc	tgagtctgcg
	240
ggTggagaga	gagagaaaga
aagcctcttt	acacgtgatt
tttaaaagag	acccaaaccc
	300
caaaagcaaa	cctcttttgc
atgcgtcctt	aaaagacata
aatttctctc	aaaattttct
	360
acatcacaaa	atcaatcttt
ttctcttctt	cttcgtcttc
atcatcatca	tcatcatctt
	420
cctctttctc	ttcctactga
gatattttct	ccacattgag
aggaaagcta	t
	471

<210> 30
 <211> 1231
 <212> DNA
 <213> Arabidopsis thaliana

<400> 30	
aggataaatt	tcatctatta
agatatcagt	caattataat
gtgttacgtg	attcgataaa
	60
aaaaaaagac	caaaaaaaaa
aagaagataa	ctattggtaa
gcgtaagaaa	tgtgtttaca
	120
ttttggcatt	ttgccaaaac
acataaagat	ggttagtgat
gagacgagac	gagtcatgcg
	180
ctacttttaa	aacaaaatga
aaaacatcat	taagctaaca
aaccaaacac	acttgttttg
	240
ataacatgtt	ctaggggaact
agttatgcca	aatctaattc
gcataagaag	actaagtcac
	300
aacataattc	agtaatttgg
ttgagattaa	atcctataaa
tatgatttta	aggtataaga
	360
gagaagagac	tcttttgatc
aacacaatca	aacatctaca
aagaaaatta	tctcacatag
	420
ctacttctta	atctaatttt
ttcattaatc	cattttatTT
taaatgtgaa	gaatcgcatc
	480
tagatgtgac	ctctcatgat
aaaaaattaa	accattgtaa
aaaaaatggt	gtgtaaaact
	540
aaatataata	aattattaaa
aaaatacaaa	ttcaatccac
taggttaaaa	actcctatgt
	600

60760_anniversary_ST25

agaacatttt	tttatattaa	aatgtaaata	catgaatctt	atttttcgaa	aaactaaaga	660
catctttttt	ttatatatta	attaccaaaa	caaaataaga	cgacaaaaat	attctttgat	720
atagtaaaag	aaaactagaa	aactagaaaa	caataaatta	ccaaaacaat	ctagaaaaca	780
ataaatccta	ctttgcatta	ctttattata	aaatcccgaa	atgaatctat	aaatgtagaa	840
aatattatac	aaaagttgta	agagatttta	atatacataa	ttacatatat	atacaagtaa	900
atacaccgta	tatacatagc	aaggagtaaa	cagtattatt	tggtatatag	ttacgtctct	960
atatacgaag	ggttcaaact	tcaaagtaat	aatttaataca	acaatgtgta	catattgata	1020
agtagtagta	tatatgtaaa	ggtctcacgt	ctctataata	aagtatgact	cgtcacgtga	1080
cctcctcttc	ttcgagaga	cagagatagg	atgagacaga	aagaaaccaa	caaaaccaa	1140
ccccaaaacc	caagaaaaag	agaaaaacac	tctcttctct	tctctctctc	tctttctatt	1200
taagagactt	cactgtctct	ctcagtcttt	t			1231

<210> 31
 <211> 1548
 <212> DNA
 <213> Arabidopsis thaliana

<400> 31						
ttcacgtggt	tatttattta	tttgggttat	taaacataaa	tcatgtaa	ctgaatcctg	60
tgagatctc	tccctagttg	atgaatagat	atgatgaatt	taattctttc	atgaaataaa	120
aatatatgaa	acatatgtag	cagaaaaaga	agcatatcta	tgaacaaca	aacattcaaa	180
aaaaaaagga	aaacggaaaa	ttattaatat	gaaaactacg	gctttgactt	gtagctgact	240
acatttacga	catatatata	taaatggacc	ccactgagtg	tctgcaaggt	ctttacacaa	300
cagtatcttc	ttctgtttct	ttgactcttt	gtgatcccta	agcctacca	taatacgtgt	360
ctacatttta	ttggattggt	tcgtgactct	gtaatctttt	ttataagaaa	acaagtaata	420
gtgaaattga	agtaaatagc	tcagcacaga	aacttcgaca	aaaataactc	acagattaga	480
aaagaaaata	tatgcataaa	tagccatggt	tcatttatga	acaatttatt	cgttttttta	540
gtttataatt	tcattaaaac	atgtttgtca	catcacattt	catgtccttc	ggctcctact	600
acaacaacaa	gtcactgtca	tctccattac	ttccacttct	gctcctttct	ttattaactt	660
gttcaaaaac	aattctaaga	taaataacaa	taaatgttgg	tctctcttta	ttatttcccg	720
gctaaagaag	gaggatgtct	cgtattatcc	gccatcaatg	ctcttttggt	tcctgtttct	780
tgcaatttga	atccctgaga	atcctagccc	acttatttac	tactttgcct	tagctgtttt	840
cgacatcaaa	attttggta	tatgactcat	atcaatcttc	aaatttgata	aaatatgttc	900
ccaattcaca	aaaacaaaaa	agttttcgaa	agctcaaaaa	cctttaccat	ttcaatagta	960
gataggattc	tttttagattt	gcatttcacg	aaaagagaag	aaaaaaaatc	gaaaaatatt	1020
tgcaatcatg	attttttggt	tctgaaggag	acctgtagtt	gctgtcatga	acattaaata	1080
caaatcta	aaatgttgta	cgaattttgc	gtgtaataaa	tggtcagggc	cggtcgaag	1140
ctcgctgatc	gtcctttttt	cgtgtctcta	tagcaacaca	caatcgtatt	tatttcaaac	1200

60760_anniversary_ST25

tttttttact ttgtttccca tccatcaa	ataagtataa aaatgtaaag aatcatcata	1260
tatagatcgt aaattcattg cttcctttg	ctttttatatt catctagacg acgttaaaac	1320
cagaccagac caaatacatt tatcattttt	cccttttttc taaaattctc tctttgattc	1380
ctatcttctt ctctttatatt tcactttgtg	ctttctctgt ctctcctatt atgagtctaa	1440
aagtctacta gctgttcaat agttttgtct	ttctgtgttt cttctttctc aaaaccgaaa	1500
gaaattcaaa aagagtcttt cgctgcttgt	tagtgggggtg aggaacaa	1548

<210> 32
 <211> 440
 <212> DNA
 <213> Arabidopsis thaliana

<400> 32		
aaacccgaac ccgaaccaa cccgaacaa	aatcttaaat taccggaatg ggtcttaaat	60
ttctaaatcc gaaaaacccg aacccaaaat	accaacccg aatctgaccc gaatatccga	120
acgcctaatt tttctatgtt aatgaaatca	attatatgac atgtttataa agagaaataa	180
attacggtga gaattaagcc catttacgtt	acggaaataa aacacccatt taaaaagcc	240
caacacgtga agcccatttc cgagtgcgtc	ccacatttac tccaacggtc gaatcgactc	300
aaacattcaa aatacaaaaa cgctatcttt	atcgtcttcc tctgtctctc tctcacaaca	360
cataacgttc aaatcctctc tctctctatc	tcgtctctta tctctagatc taaaaatctc	420
ttctttcctc aatctctgtt		440

<210> 33
 <211> 317
 <212> DNA
 <213> Arabidopsis thaliana

<400> 33		
acagacattt acttatacgg ttattgaggt	tgaactggac cggagtagca ataaattatc	60
ggttcagttt gggagatcaa accgttttaa	agaaaataat ttgaaatggc cagcgagaat	120
acgaggggtc gaggattgta cctcctttct	ctgcaaaaac ttaaacgttg atttgactca	180
agcgtcaagg taaggtagtc tctcttcata	caacatttta gctttacttt ttctctttac	240
tcttctctct ctctttctct ttctctttct	ctttcactcg ttctctctca ctactctct	300
tcacacacag atccaag		317

<210> 34
 <211> 3912
 <212> DNA
 <213> Arabidopsis thaliana

<400> 34		
acaccaataa aaatacacag caataaaatc	gctacgtata tatatatata atatgtatta	60
tctattacaa gatagtaata gagtatagca	agttgtatca tctaacaaac tatgcgaata	120
aaatttgaac attgtgacat gtagatgtag	tgtaatttag ctaagtgcct atcatcagta	180

60760_anniversary_ST25

acatagaccg	acttaacttt	ttacgaaaa	aaaaaagtaa	catagaccga	aaaaatgcat	240
atcgtaaatt	taatggaaaa	cacaatttac	gataagtaaa	aaacaaaaag	aaattacgat	300
aagtcgagaa	aaatgcaaca	aattgagata	aagtattgat	aaaaccatga	aagtgtcggc	360
gtatgtaaat	gcggtgatta	atgtgatcat	tagagcgtgt	gtgttaaacg	cggcggtttt	420
agtggagatt	gatcagctga	taacactctt	accgggacga	atctaattcc	atattcatgg	480
cttgttaaaa	cctaagacat	acgcaatctc	taatttgcta	gtatagttag	ttctatatta	540
tttttcgact	aataatgtaa	acatatgatt	attaagtcgc	aaaaagagtg	cttaacaacc	600
aaaaagtgga	ttaattaact	tggtgggaaa	agttacaaaa	cctttaatga	ttactctttg	660
taccaagaat	agtggcgaag	cactataaga	gcagagaaaa	gaagctcaat	aatgtactaa	720
aagttgtaga	tttttacagc	ttaaatacac	caaaattaat	agaaaagttg	gtaatttttt	780
aattcatggc	tactgattta	gattttagaa	aacaatagta	gtatcattgt	cacatcttaa	840
acacacaata	ggtatgtttt	aaatcaaagg	ccgtagttaa	tttgtcaaaa	atgtatgcat	900
ttggtatttg	gatgtctccg	aaaggatgga	tatatggact	tgtagataa	tttcatacct	960
cagtatcaat	agtcattggg	cccaaattgc	tcaaaaacat	atttttaatt	ccaagacttt	1020
gatgaagacg	taataatgag	tccaatgggc	catcagatac	aatgttcgga	atttaacggg	1080
tttgtagttt	ataagtattg	ggcttgacct	atctggttca	atgatatgta	ggaacaaccc	1140
aatttgcaaa	gctttattaa	aagactcttt	agttgtcgtc	aaggtttaac	ttgtagtagt	1200
tggtgaagaa	ttctacgtga	aataggcaac	attacaaaaa	caaaaatcaa	ttcgaaatca	1260
tacaaaacga	aaccaagtag	taaccaacta	cactattatg	acattaatga	ttagacattc	1320
ccaaatcata	caagttcctg	tcatgaagga	aacaatggtc	cgtatttgca	aacgattaca	1380
aaaattcaaa	ccaaaaatga	aaaaacgagt	taaattattt	ggtttataaa	aatagtaatg	1440
tcaacagaag	actagattgg	gaaacctgaa	gcgaacagag	cttttaaaaa	cgagtttgaa	1500
cggctgggat	catttggtac	aatacccacc	gtaagtttgt	ttaccctagg	gatgcaagcc	1560
aaaggcccaa	atcagttact	acttactgct	acaaccatcg	tctcagcttt	ttgtctcagc	1620
tttttactaa	tgaagcatac	aatttcttgg	gcatgtcaca	tctcgacacg	tgtccactat	1680
tctcttctct	tattggctac	tcgttcgtag	gcttctgtta	atagatgata	tctctataac	1740
tctaacagtc	ttttctttct	ctttatttcg	ttttggtatt	ttaagtttca	aattgaaaat	1800
aataggagga	aaagtctagt	tttaaataat	gtttttttac	aagtgaacgt	gaaccaattt	1860
acctcttttt	ttttatatat	cctatcggct	aatctgggta	gtatcggtag	aaatgcaccg	1920
aggtgctaca	gagattaatg	ctagggatag	tcagaccgct	tgtatttctg	actatcaagt	1980
aaatctacgc	ccaactcaca	tatttcccaa	acaaatgtga	tttttttttt	tttttttttt	2040
tttttttttt	ttttgtaaca	aatgtgattt	tgttttcaag	gaaaatagaa	cttacgtttg	2100
ggaatttcac	ccttcactaa	agcttccttc	tgccattaga	ccacaaaggc	ttgggcaatt	2160
taccattttt	gtaaaagtag	aaaacaaaat	gcctaaaatg	ttcatacttc	attacatcaa	2220

60760_anniversary_ST25

caagggttatg	cccacgatat	agaggcatgt	aacattttata	tatatagtgg	aagaagccta	2280
cgagctttat	taataagtat	aaactctgat	tattaggtaa	ataaattact	taaaacgatt	2340
actcaactga	caaaaccgta	gttgaataat	aaggttacta	tgaataccga	ttgaatattg	2400
caaagccgga	attgaaaaat	atataacaga	tcaaagtgtc	aagtgtggtc	ataatttctca	2460
cataggtcac	atagctgaac	ccatgcatct	atttactagt	ctatagaaag	tactagagac	2520
gcatacagct	gaacctactc	tattctttta	ttaatttttg	ttctcgtgga	tacaaaattc	2580
ctccaacatt	tattagaacg	aataaaacca	atatgatgat	gattagttat	tggtaaacat	2640
ataaacgttg	agtaaacttc	aaaatagatt	gaagtactat	taagacttgc	atTTTTTccc	2700
cttggtttat	attcttgaat	cgtttcgaag	tattttaact	ttcaagaata	gaaggttcct	2760
caactataaa	caattacatt	aatcaaaacc	atttctatgt	aaacaacata	atTTTTgtat	2820
atTTtagtct	tcccaaaaag	tttgaccgat	agggcggttt	agaccgtata	gtacgactgt	2880
acaacaaaaa	ggactctgga	gacctaaaga	tccaaaacta	tgcaaaataa	agatacggtc	2940
ggaccaatTT	aatctaacia	aaccaaacc	ttatactaaa	ctattttaccg	atacatttcc	3000
atataacaca	gtacacacaa	ttaaatcaaa	cattattgga	agaacaagat	agaatattgg	3060
cttaatctcg	aacgattaga	gttatcctag	agcctcggag	cttttgtcac	atataatata	3120
aactatggta	tatataaaca	tgactctcat	ttgtatttat	cgcaagggtac	aattccacca	3180
atTTTTttcg	tccactcat	acagctttta	ttgtgaaatc	aatccataaa	aaaccaacat	3240
gtgacatggt	ctctataact	ataactataa	gatagtaaaa	aattcacatc	aacataaaag	3300
aaaaccaatc	atattggcta	aaaaaaacta	acggtcgaaa	aacgtataac	cacaaaacca	3360
aaccggtcca	accggtgtcc	ccaatcacta	tcaaagcatt	aactaacttt	cacaaggaaa	3420
agcatagttc	agtttctcta	catcgcttcc	catcctctta	accctgttta	ctcgaatcat	3480
ccaccgttgg	atcaaacacg	cgctacaaat	ctagcgcgtg	accgaggttt	ttacacagtg	3540
gaatattacc	atgcattgga	aagcggcgtc	tacaacaaac	ggcgggtcat	gtcaccgtca	3600
aatcaacct	ttcttaattc	ctaacgccgt	tacttatctc	cgtttactaa	aatgtttaat	3660
gcgtgtgaga	gtgaagatca	tatactaatt	agaagtggct	aatgttttaa	cgtgacatta	3720
ttatcatagt	taatggttcg	atcagagttt	taagtagtaa	atgatataag	tgtgtgtata	3780
taattgcata	catatatact	ctcacactct	gacagatttg	tcgtgggtctt	agtatttctt	3840
ttcatggcta	gttatatagg	gctctagtac	attatctctc	tctccccatt	tctctgtctc	3900
tctcttcttt	aa					3912

<210> 35

<211> 2000

<212> DNA

<213> Arabidopsis thaliana

<400> 35

atctttatgg tcaccgagtc tactgatata attttactgt cgcagtttgt ttccactact 60

taagtttcta taatttcaca gtttgaaaga aaaattactg gttattcagc taaattacaa 120

60760_anniversary_ST25

agattagttt	aattagttta	gccagtataa	tgtttttagta	aagtattaaa	cggcattttt	180
cgttgggaga	attatgttat	ggtataatct	actaatacat	actttttacac	atatatcaaa	240
aagtttgacc	atagtaggta	gtacaacata	gaagaatcaa	gatcggaacc	agcaaggaag	300
aagatacggg	cgggtccatat	taagctaaga	ggaccaacgt	aactcgatat	atatttttct	360
gttctctacg	ttaccaccat	ataaatttta	atattgaaaa	aatcatcttt	tggcattgtg	420
tttgatgtcg	gattcggaat	atggaaagag	gagagatatg	agattttggc	acaaaggaag	480
ctgccaaagc	attagggcaa	ccgagtagta	acgagatcaa	acatcgtttc	aatcggacgg	540
tcgggggtttg	accaatattt	ctcggatatc	ttttggaccc	tacgttctga	cttgaacttg	600
atcagtcact	tcagtacctt	agttttcatt	ttcaatgtga	tcatgagttt	ttttttacat	660
gtagcttca	aaacaaatac	taatattcat	taactatgga	tcggcatagt	tttcatgtaa	720
tcagctgagc	gtttatcata	ttgattgaag	ctaactgtga	aaattctcat	gatcacattg	780
acttttgcct	acaaatttta	aaagagtata	caaataattg	cttaatgaag	atagcttcca	840
tagagaaaga	gtaacagctt	tatacggagg	catagcttta	gacacgatct	ctgctcttgt	900
gttttttggt	taacactgaa	tccacagtga	aattactgct	catttttttt	cattttttat	960
tacatttttt	tttttacttt	tttattttatt	acaatctaca	gttctaccaa	cttatttcaac	1020
ctagtggtag	catatcgacc	ccaaaattaa	tcaatctaata	tacaaggtag	aaatagaaag	1080
attttatcaa	aggagacaac	tctgatcgat	aatatgttgc	aataaaaacca	tgaaaaactg	1140
taaaaaatat	tgaaagctga	agaaaaattt	tcaaatcgat	aaaaggatag	tactaaacca	1200
atccgggttg	tggcatcttt	ttcacccaat	cactatcaaa	gcattaacta	aaaatcacaa	1260
ggaagagcat	aatttgattc	tctacatcgc	agtccacgat	aggattttct	tatccaccgt	1320
tggatcaaata	ttaataatga	tgcacgcgcg	cgtcaacggg	attttaccba	gcaaaggaat	1380
gtcttttcac	cggactctta	aaagacgttc	ttcttttttc	acctttgcat	tggaacacgg	1440
cgtttcttct	tagaacgctc	gccgtcaaata	catacggcct	aataaatctc	cgtttaacgc	1500
cgttacttta	ccgttaagta	ctaaaaaac	aaaaaaaaaat	catttcgatc	actgtctcat	1560
taagatgatc	ggagatgttt	tagcagggtt	taacaagtga	tgatagtaat	gtatgtatat	1620
atgttactga	cattattttg	tcgttgtcta	ataggagggt	actaaagttt	ctctctctca	1680
tggcgtcgga	gctcagcctc	tagtaatgta	gactgtcctc	tctttctctc	tctcttcttt	1740
aaacatctct	gctctgtttt	ccttccagtt	cacgctaata	tcctgtgtcg	gtcccctctc	1800
tcttttcctt	tggtctctcc	caacaatggc	agaacgactt	tgtacccttc	ttttgctctt	1860
tgtttgaatt	tcgtttcttg	ctacaaagct	tcaaaggatc	tgacttttcc	ctaaacagaa	1920
aaagagggtct	ttaacaaaaa	aagggttgta	cttgttttct	gggtttcgtg	gtgttactct	1980
tgaggaagaa	gaagaagaag					2000

<210> 36

<211> 1470

60760_anniversary_ST25

<212> DNA
 <213> Arabidopsis thaliana

<400> 36
 ctacctaata taactagcta gggatttcta ctcttgTTTT cataatcgat ctacggacat 60
 ttctcggAAC gtggtcaaga tTcatgagtc ttctgTTTT tatgtctctg ttcaatttgg 120
 tttagagatt agtatgctta tttgtttatt tcatatatgg ttatgagagg agaggctaata 180
 ggcataact ctgatgtttg tgatggctgc taatatcggt gaggagtTat tcacgttgTt 240
 tcatgcgcaa aaatcaacag aaaaaattct gattatgagc caactctgtg aacccttata 300
 gtgcgcccag aggtttgcga ggcaaaatcc cgatgaacca gaaggaattt tagatctcta 360
 tcaacaataa ctatgatgga gctcgTttaa attcatcaca gcgacaacat cattaggctg 420
 cccaacgtct atgtctcctg gaggtgatgg tacttgatct ctcaaccaat tttcttgaaa 480
 atatcatgcc ttgtgagcgc tttcatattg cgcctaaaat acccaatacg caatgaacct 540
 acttccaaag gcatagaaaa aaaactgata atgataatga gatttgTcac tatacttattc 600
 ctatccctac ataggagccg tttgattgtt tagtccatgt tttcattttg tttagtctaa 660
 tgctatataa cttttcttta tcagtctatt gttatatgac ttatatatat ctcaagagat 720
 aaggccaata aatcttcttc ttaattatat ctgaagactc aaaacatatt ttgagtttaa 780
 taaaataaat aacgtccaaa tgctacatac aaacggacca aattcatgga ggtataaatt 840
 taaattattt tttgttccaa agtgatgca gtgatttatt gatgaatgcg atagagcggc 900
 gaaagagaat aatcgTcacc tagaagacaa attgatcggc cgtacatata tacataaata 960
 caaacctgcc acttcacatg tcaccacct ttaagcacc ccttcacata catactttct 1020
 ataacaaaaa tatcagcttc tagttcatat ttatgttaca ataactcgag tgaatcatac 1080
 taaaaaaatg taatgctttc tctaaatagg agataaaatg caccctccga cctaactaaa 1140
 gattccttat tttagctatt taagacatat tgcacatgta tagagataca taaacacata 1200
 tgcaatatgc acatcttcta tacattgaaa aaagctgac ttgcaaatat ttgtcttaca 1260
 caacacaagc gaccaaagcg atgcgtttcc caatgataag gttacgacat acttacacga 1320
 ctctctctat tgtctcgtct ctttctttcc tcatccctct ctttTgtct ctttactct 1380
 atttttcact tttcagaata cttttacgta aaaatcatgg acatgtcatt gtctccaccc 1440
 tactatactc tttttttgtt ctttttgTtt 1470

<210> 37
 <211> 1793
 <212> DNA
 <213> Arabidopsis thaliana

<400> 37
 ttggttgTct ggcacatca ttttgtaccg tttctcccaa agtaagaaac ggtacaattc 60
 tctcttatat agatttcata ccccaaaacc ctaaattcat tagggTTTT aaaaaaaaaa 120
 atcacgttta cctctaaacc aatcttctct tatatagata aatcataacg tttgtttgat 180
 ttttcagttt tctacttaac caatattaaa ctaaagtcga attgagatga gtggtagcaa 240

60760_anniversary_ST25

accacagttt	aattaagaag	ttaattatag	gccacatgat	tgagcaagcc	tttttgtttt	300
gtaacacatc	ttatcagctg	cttaaaat	tggctgcctc	ccattggcca	actggtctaa	360
acatcattgc	attggcattc	tcataatcaa	tcaatcta	gagaaacttt	gaatatttat	420
gaaaaactga	ataacaacat	aacataaacg	aacaatgtaa	aaaagaaaaa	cacaaaaaaa	480
aaaacacttt	aaaaaacaaa	aaccaaaaac	tcttaacta	taaactcatg	aacacttagt	540
gatgaggtct	gaaaggggtg	aaccaccacc	tggtgtcaat	aggtgacaac	ttcttcttgg	600
gaacattcgg	gaaagtgaag	gcttaggtga	cggttgtctt	aaaagtctct	tttagttaat	660
tcacgtatt	ttcgaatggc	attacgtttg	atgcataaag	gcccatatgg	gctatacatg	720
tactgcgttt	gagtggcttt	ctaaggattg	atgtattgtc	tctatgagag	tattcgttta	780
actcatggag	atctactctc	cacgatattt	tctgtaaact	tttctttttg	ttgattagat	840
aaatagaaaa	ttgtgtagag	cgaaactttt	aatgaattaa	aatgcggaag	cgattaaagc	900
atgaatagat	aaattggaca	agagattaaa	cgagggatca	tctagttttt	acactgatca	960
ctagtcatct	gcttgcaaaa	gaagtatatc	attaatcaag	caaaacgagg	gcataaattt	1020
cttacaataa	acttttacag	taggttaatg	attttttaat	aacttgtcca	tttcacatgc	1080
atgtgtatct	ttgtactata	catgctaagt	gtttcattaa	tcaagataaa	cgtgtctacg	1140
aataacttaa	aacagtacaa	cttcctaaa	aaattcatta	aatgaaaggt	ttttatagat	1200
tatacattgc	acgggtacgt	tcggttacca	ttcgaagtct	aaaaagagaa	tgacggttct	1260
gataatgctt	ttaatcgctt	ttgtattgta	aatcattaaa	acagtaagcc	ggataccgaa	1320
ttactaatca	gacccaaaaa	agaatctata	ggaaaaatat	caactgaaga	gcgggtaggc	1380
ttgacctga	aaggaagaat	ggtgagcgag	cggtggatag	atatgtaata	aattgtaacg	1440
ctttcaaaat	gtcaaagtca	caagtcacat	tactcacgag	ccaacactaa	ccatgcaact	1500
tttgttttga	cattttccta	aactttaggt	ataaaatacc	cgcgtaataa	ataacctctt	1560
cataattggg	tccaccact	cacaggtcca	cacataagga	accgaaaaag	gtaaaattca	1620
aaaacttaca	aagtttttta	gagatgatgt	ggtgaagtat	tgcattaatg	gaataatggg	1680
aaaagaaagt	aattgcaacg	tacgtataga	ttaatccatt	gacacaaatg	aaaagtttct	1740
ttctatttaa	tgtacacaac	aaaggttctc	ttcagagtaa	tttaggggaa	aaa	1793

<210> 38
 <211> 257
 <212> DNA
 <213> Arabidopsis thaliana

<400> 38						
acacaacagg	tccctacatt	tgtacaatct	cctctcttta	aagactctct	ctctttctct	60
ctccatctct	atcttactct	gtattttctgt	cgtctgagca	ctcaatgaaa	ccactgtaaa	120
tttccgccag	aatttgatgt	gatggaacga	taaaaatcat	tttttctcgg	ttaaagtaaa	180
aaaacaaaaa	caaatttctg	tagaaatcat	aataaaagaa	agaaaaaaa	tctaattgtcg	240

gtacataata cggttct

60760_anniversary_ST25

257