



(86) Date de dépôt PCT/PCT Filing Date: 2012/10/18
(87) Date publication PCT/PCT Publication Date: 2013/04/25
(85) Entrée phase nationale/National Entry: 2014/04/09
(86) N° demande PCT/PCT Application No.: US 2012/060741
(87) N° publication PCT/PCT Publication No.: 2013/059407
(30) Priorités/Priorities: 2011/10/18 (US61/548,518);
2012/07/09 (US61/669,480)

(51) Cl.Int./Int.Cl. *A61K 47/26* (2006.01),
A61K 47/02 (2006.01), *A61K 47/10* (2006.01),
A61K 47/48 (2006.01)
(71) Demandeur/Applicant:
COHERUS BIOSCIENCES, INC., US
(72) Inventeurs/Inventors:
MANNING, MARK, US;
MURPHY, BRIAN, US
(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : FORMULATIONS D'ETANERCEPT STABILISEES PAR DU CHLORURE DE SODIUM
(54) Title: ETANERCEPT FORMULATIONS STABILIZED WITH SODIUM CHLORIDE

(57) **Abrégé/Abstract:**

The present invention is an aqueous etanercept formulation stabilized to reduce instability, aggregation and/or fragmentation of the etanercept, said formulation comprising about 25 to about 75 mg/ml of etanercept and one or more stabilizers, wherein the stabilizers are selected from the group consisting of sodium chloride; and sodium chloride in combination with sucrose or trehalose; and a combination of sodium chloride, sucrose and trehalose. Various technical terms used in the following discussion are defined below in the section entitled "Definitions" and throughout the remainder of the specification. The stabilized etanercept formulations of the present invention elicit long term storage stability.



- (51) International Patent Classification:
A61K 39/395 (2006.01)

(21) International Application Number:
PCT/US2012/060741

(22) International Filing Date:
18 October 2012 (18.10.2012)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/548,518 18 October 2011 (18.10.2011) US
61/669,480 9 July 2012 (09.07.2012) US

(71) Applicant: COHERUS BIOSCIENCES, INC. [US/US];
201 Redwood Shores Parkway, Suite 200, Redwood City,
CA 94065 (US).

(72) Inventors: MANNING, Mark; 4630 Sorrel Lane, John-
stown, CO 80534 (US). MURPHY, Brian; 1708 West
Prospect, Fort Collins, CO 80526 (US).

(74) Agents: POLYAKOV, Mark, V. et al.; Wood, Phillips,
Katz, Clark & Mortimer, 500 West Madison Street, Suite
1130, Chicago, IL 60661 (US).
- (81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,
RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: ETANERCEPT FORMULATIONS STABILIZED WITH SODIUM CHLORIDE

(57) Abstract: The present invention is an aqueous etanercept formulation stabilized to reduce instability, aggregation and/or frag-
mentation of the etanercept, said formulation comprising about 25 to about 75 mg/ml of etanercept and one or more stabilizers,
wherein the stabilizers are selected from the group consisting of sodium chloride; and sodium chloride in combination with sucrose
or trehalose; and a combination of sodium chloride, sucrose and trehalose. Various technical terms used in the following discussion
are defined below in the section entitled "Definitions" and throughout the remainder of the specification. The stabilized etanercept
formulations of the present invention elicit long term storage stability.

Etanercept Formulations Stabilized with Sodium Chloride

Field of the Invention

The present invention relates to aqueous pharmaceutical compositions stabilized with sodium chloride for long-term storage of etanercept, methods of manufacture of the compositions, methods of their administration, and kits containing the same. The invention includes etanercept formulations that do not require arginine for stabilization.

Background of the Invention

Polypeptides must often be stored prior to their use. When stored for extended periods, polypeptides are frequently unstable in solution (Manning et al., 1989, Pharm. Res. 6:903-918). To extend their shelf life, additional processing steps have been developed, such as drying, e.g., lyophilization. However, lyophilized pharmaceutical compositions are less convenient to use.

Typical practices to improve polypeptide stability can be addressed by varying the concentration of elements with the formulation, or by adding excipients to modify the formulation (See, for example, U.S. Pat. Nos. 5,580,856 and 6,171,586). However, the use of additives can still result in inactive polypeptides. In addition, in the case of lyophilization, the rehydration step can result in inactivation of the polypeptide by, for example, aggregation or denaturation (Hora et al., 1992, Pharm. Res., 9:33-36; Liu et al., 1991, Biotechnol. Bioeng., 37:177-184). Aggregation of polypeptides is undesirable, as it may result in immunogenicity (Cleland et al., 1993, Crit. Rev. Therapeutic Drug Carrier Systems, 10:307-377; and Robbins et al., 1987, Diabetes, 36:838-845).

Another way to improve polypeptide stability is to use L-arginine at a specific concentration (U.S. Pat. No. 7,648,702).

One of the polypeptides that is stored for up to two years prior to use is etanercept (Enbrel®, Immunex Corporation), which is a dimeric fusion polypeptide consisting of the extracellular ligand-binding portion of the human 75 kilodalton (p75) tumor necrosis factor receptor (TNFR) linked to the Fc portion of human IgG1. It consists of 934 amino acids and has an apparent molecular weight of approximately

150 kilodaltons (Physicians Desk Reference, 2002, Medical Economics Company Inc.) The Fc component of etanercept contains the constant heavy 2 (CH2) domain, the constant heavy 3 (CH3) domain and hinge region, but not the constant heavy 1 (CH1) domain of human IgG1. An Fc domain can contain one or all of the domains
5 described above. Etanercept is usually produced by recombinant DNA technology in a Chinese hamster ovary (CHO) mammalian cell expression system.

The present invention provides novel stable liquid formulations of etanercept that allow its long-term storage.

Summary of the Invention

The present invention is an aqueous etanercept formulation stabilized to reduce instability, aggregation and/or fragmentation of the etanercept, said formulation comprising about 25 to about 75 mg/ml of etanercept and one or more stabilizers, wherein the stabilizers are selected from the group consisting of (i) sodium chloride; and (ii) sodium chloride in combination with sucrose or trehalose; and (iii) a combination of sodium chloride, sucrose and trehalose.

Various technical terms used in the following discussion are defined below in the section entitled "Definitions" and throughout the remainder of the specification.

The stabilized etanercept formulations of the present invention elicit long term storage stability as characterized by at least one of: (1) SEC analysis at M_3 or T_2 or T_4 of: monomer content greater than about 90%; aggregates content of less than about 3 wt%; and fragment 3 content less than about 5 wt%; and (2) HIC analysis at M_3 or T_2 , or T_4 wherein the amount of the composition represented by peak 1 of the HIC chromatogram is less than about 3 wt.%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than 80 wt.%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than about 20 wt.%.

In a related aspect the formulations elicit long term storage stability as characterized by: (1) SEC analysis at M_3 or T_2 or T_4 of greater than about 90 wt.% monomer content; less than about 3 wt.% aggregate(s) content; and less than about 5 wt% fragment 3; and (2) HIC analysis at M_3 or T_2 or T_4 wherein the amount of the

composition represented by peak 1 of the HIC chromatogram is less than about 3 wt.%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than 80 wt.%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than about 20 wt.%; and wherein the formulation is free or substantially free of arginine.

In preferred aspects of the stabilized formulations, the formulations elicit long term storage stability as characterized by: an HIC analysis at M_3 or T_2 or T_4 wherein the amount of the composition represented by peak 2 of the HIC chromatogram is greater than or equal to about 95 wt.%; and wherein, if peak 3 is present on the HIC chromatogram, the amount of the composition represented by peak 3 is less than or equal to about 3 wt.%.

The stabilized etanercept formulation as summarized above, optionally and preferably, contains no arginine, or are essentially free of arginine.

The formulations of the invention have excellent stability as determined by SEC (Size Exclusion Chromatography) and HIC (Hydrophobic Interaction Chromatography) analysis conducted after one, two or three months of storage at 5° C. These formulations are comparable to or better than a commercially available formulation of etanercept, in which arginine is a required component. Accordingly the present invention is further directed to formulations of stabilized etanercept, as summarized above, which contain no arginine, or are essentially free of arginine, and wherein the composition at M_3 or T_2 or T_4 elicits long term storage stability that meets one or both of the following criteria: (A) stability comparable to or better than commercially available etanercept marketed under the trademark Enbrel®, as measured by (i) SEC analysis of the amounts of aggregate(s), monomer and fragment 3 in the composition (as defined in the specification) and (ii) HIC analysis of amounts of material in the composition corresponding to peaks 1, 2 and 3 of the HIC chromatogram (as defined in the specification); and (B) an HIC chromatogram in which (i) peak 3 is absent, or essentially absent and (ii) peak 2 represents greater than about 95 wt% of the composition; an SEC chromatogram containing essentially no peak corresponding to aggregate(s); and an SEC chromatogram in which the monomer content represents at least about 95 wt% of the composition.

In one preferred aspect, the formulation of the invention comprises about 25 to 75 mg/ml etanercept, up to about 150 mM of sodium chloride, about 1 to about 30 mM sodium phosphate; and about 0 to 5 wt% sucrose or trehalose or combination thereof; wherein the composition has a pH of about 6.0 to about 6.6; and wherein the composition is characterized by SEC analysis at M_3 or T_2 or T_4 in which: monomer content is greater than about 80 wt%; aggregate(s) content is less than about 3 wt%, and fragment 3 content is less than about 8 wt%. Moreover, formulations meeting these criteria do not require arginine for stabilization.

In a further preferred embodiment the stabilized etanercept formulation is further characterized by (a) an SEC analysis at M_3 or T_2 or T_4 of greater than about 90 wt% monomer content; and less than about 3 wt% aggregate(s) content; and (b) an HIC analysis at M_3 or T_2 or T_4 wherein the amount of the composition represented by peak 1 of the HIC chromatogram is less than about 4 wt%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than about 80 wt%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than about 20 wt%. Formulations meeting these criteria do not require arginine for stabilization.

The etanercept compositions of the invention further afford the ability to provide formulations which contain acceptable levels of subvisible particles. Accordingly, the invention is further directed to sodium chloride stabilized etanercept formulations having, at M_3 or T_2 or T_4 no more than, on average, about 10,000 subvisible particles per mL having a size greater than 5 μ m.

The stabilized etanercept composition of the present invention are further characterized by: (a) an SEC analysis at M_3 or T_2 or T_4 of greater than about 90 wt% monomer content; and less than about 3 wt% aggregate(s) content; and (b) an HIC analysis at M_3 or T_2 or T_4 wherein the amount of the composition represented by peak 1 of the HIC chromatogram is less than about 3 wt%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than 80 wt%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than about 20 wt%. Formulations meeting these criteria do not require arginine for stabilization.

The stability of the formulations may be further characterized in that the compositions exhibit an HIC analysis at M_3 or T_2 or T_4 wherein the amount of the composition represented by peak 1 of the HIC chromatogram is less than about 1%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than about 95 wt%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than about 1 wt%. Surprisingly, formulations meeting these criteria do not require arginine for stabilization.

In a further preferred embodiment, the sodium chloride stabilized etanercept formulation of the invention comprises up to about 150 mM sodium chloride, about 1 to 30 mM sodium phosphate, and about 0-5 wt.% sucrose or trehalose, or combination of sucrose and trehalose and having a pH of about 6.0 to 6.6; and characterized by: an SEC analysis at M_3 or T_2 or T_4 of greater than about 95 wt.% monomer content and less than about 1 wt.% aggregate(s) content; and an HIC analysis at M_3 or T_2 or T_4 wherein the amount of the composition represented by peak 1 of the HIC chromatogram is less than or equal to about 3 wt%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than about 82 wt%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than about 15 wt%. Preferably, formulations meeting these analytical criteria do not contain, or are substantially free of arginine.

It is surprising and unexpected that excellent stabilization can be provided in an etanercept formulation without need for arginine as a stabilizing component. Accordingly the present invention is directed to the following sodium chloride stabilized formulations which contain no arginine or are substantially free of arginine:

An etanercept formulation free of arginine, comprising: about 50 mg/ml of etanercept; about 75 mM sodium chloride; about 10 mM sodium phosphate; about 3 wt.% sucrose.

An etanercept formulation free of arginine, comprising: about 50 mg/ml of etanercept; about 75mM sodium chloride; about 25 mM sodium phosphate; about 3 wt.% trehalose.

An etanercept formulation free of arginine, comprising: about 50 mg/ml of etanercept; about 75mM sodium chloride; about 25 mM sodium phosphate; about 3 wt.% sucrose and having pH of about 6.3 to 6.5.

An etanercept formulation free of arginine, comprising: about 50 mg/ml of etanercept; about 150 mM sodium chloride; about 25 mM sodium phosphate, in the absence of sucrose or trehalose.

An etanercept formulation free of arginine, comprising: about 50 mg/ml of etanercept; about 100 mM sodium chloride; about 15 mM sodium phosphate; and about 2 wt.% sucrose.

Preferred stabilized compositions of the invention, preferably free of arginine, exhibit an HIC analysis at M_3 or T_2 or T_4 wherein the amount of the composition represented by peak 1 of the HIC chromatogram is less than about 2% or preferably less than about 1%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than about 95 wt.% and preferably greater than about 97%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than about 1 wt%, and preferably 0 to 1%.

As differentiated from commercially available etanercept provided in an arginine-containing formulation, we found it surprising, in light of US. Pat. No. 7,648,702, that the formulation embodiments of etanercept described and exemplified herein do not require arginine for long-term stabilization, although
5 arginine may still be added if desired. The ability to provide etanercept formulations stabilized without arginine represents a potentially significant benefit to the health care system by providing patients and health care providers with alternative formulations of etanercept that may become available at lower cost compared with present commercial etanercept formulation (i.e., Enbrel®) that require arginine for
10 stabilization.

As used herein the term "instability" or like terms denotes the tendency of the etanercept monomer to undergo a variety of undesired transformations during storage. Such transformations include the formation of oligomers and high molecular weight aggregate(s) (hereinafter terms "aggregate(s)" in which multiple
15 copies of the essentially intact etanercept monomer become irreversibly associated

with one another through a variety of non-covalent attractions (e.g., electrostatic interactions.) Undesired transformations during storage may also include degradation of the etanercept monomer to smaller fragments and/or clipped species. Ideally, a formulation of etanercept should minimize, to the greatest extent possible, the tendency of the formulation to result, during storage, in the formation of aggregates, misfolded protein, oligomers and/or fragments of etanercept. An important benefit resulting from the ability to reduce formation of unwanted aggregates or fragments is a reduction in the potential toxicity and/or immunogenicity of the drug.

The etanercept formulation of the present invention which is optionally and preferably free, or essentially free of arginine. The term "essentially free of arginine" is intended to mean that arginine, even if present, is not contributing to the stabilization of the etanercept monomer in the formulation to such an extent that a person skilled in the art would judge its presence beneficial or necessary from a stabilization standpoint.

These and other aspects will become apparent from the following description although variations and modifications therein may be affected without departing from the spirit and scope of the novel concepts of the disclosure.

It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

DETAILED DESCRIPTION OF THE INVENTION

Various embodiments of the invention are now described in detail. As used in the description and throughout the claims, the meaning of "a", "an", and "the" includes plural reference unless the context clearly dictates otherwise. Also, as used in the description and throughout the claims, the meaning of "in" includes "in" and "on" unless the context clearly dictates otherwise. Additionally, some terms used in this specification are more specifically defined below.

DEFINITIONS

The terms used in this specification generally have their ordinary meanings in the art, within the context of the invention, and in the specific context where each term is used. Certain terms that are used to describe the invention are discussed
5 below, or elsewhere in the specification, to provide additional guidance to the practitioner regarding the description of the invention. Synonyms for certain terms are provided. A recital of one or more synonyms does not exclude the use of other synonyms. The use of examples anywhere in this specification including examples of any terms discussed herein is illustrative only, and in no way limits the scope and
10 meaning of the invention or of any exemplified term. The invention is not limited to the various embodiments given in this specification.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. In the case of conflict, the present document, including
15 definitions will control.

"Around," "about" or "approximately" shall generally mean within 20 percent, within 10 percent, within 5, 4, 3, 2 or 1 percent of a given value or range. Numerical quantities given are approximate, meaning that the term "around," "about" or "approximately" can be inferred if not expressly stated.

20 The term "etanercept" or "etanercept monomer" or "monomer" is synonymous with Enbrel®. It refers to a polypeptide which is a dimeric fusion polypeptide consisting of the extracellular ligand-binding portion of the human 75 kilodalton (p75) tumor necrosis factor receptor (TNFR) linked to the Fc portion of human IgG1. It consists of 934 amino acids and has an apparent molecular weight of approximately
25 150 kilodaltons. For the purposes of the present application, the term "etanercept" also encompasses etanercept with minor modifications in the amino acid structure (including deletions, additions, and/or substitutions of amino acids) which do not significantly affect the function, potency, or avidity of etanercept. The term "etanercept" encompasses all forms and formulations of Enbrel®, including but not
30 limited to concentrated formulations, injectable ready-to-use formulations; formulations reconstituted with water, alcohol, and/or other ingredients, and others.

The term "sugar" refers to monosaccharides, disaccharides, and polysaccharides. Examples of sugars include, but are not limited to, sucrose, trehalose, dextrose, and others.

The term "polyol" refers to an alcohol containing multiple hydroxyl groups. Examples of polyols include, but are not limited to, mannitol, sorbitol, and others.

The term "long-term storage" is understood to mean that the pharmaceutical composition can be stored for three months or more, for six months or more, and preferably for one year or more. Long-term storage is also understood to mean that the pharmaceutical composition is stored either as a liquid at 2-8° C, or is frozen, e.g., at -20°C, or colder. It is also contemplated that the composition can be frozen and thawed more than once.

The term "stable" or "stabilized" with respect to long-term storage is understood to mean that etanercept contained in the pharmaceutical compositions does not lose more than 20%, or more preferably 15%, or even more preferably 10%, and most preferably 5% of its activity relative to activity of the composition at the beginning of storage.

The term "mammal" includes, but is not limited to, a human.

The term "pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material, formulation auxiliary, or excipient of any conventional type. A pharmaceutically acceptable carrier is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation.

The term "composition" refers to a mixture that usually contains a carrier, such as a pharmaceutically acceptable carrier or excipient that is conventional in the art and which is suitable for administration into a subject for therapeutic, diagnostic, or prophylactic purposes. It may include a cell culture in which the polypeptide or polynucleotide is present in the cells or in the culture medium. For example, compositions for oral administration can form solutions, suspensions, tablets, pills, capsules, sustained release formulations, oral rinses or powders.

The terms "pharmaceutical composition" and "formulation" are used interchangeably.

The term "treatment" refers to any administration or application of remedies for disease in a mammal and includes inhibiting the disease, arresting its development, relieving the disease, for example, by causing regression, or restoring or repairing a lost, missing, or defective function; or stimulating an inefficient process. The term includes obtaining a desired pharmacologic and/or physiologic effect, covering any treatment of a pathological condition or disorder in a mammal. The effect may be prophylactic in terms of completely or partially preventing a disorder or symptom thereof and/or may be therapeutic in terms of a partial or complete cure for a disorder and/or adverse affect attributable to the disorder. It includes (1) preventing the disorder from occurring or recurring in a subject who may be predisposed to the disorder but is not yet symptomatic, (2) inhibiting the disorder, such as arresting its development, (3) stopping or terminating the disorder or at least its associated symptoms, so that the host no longer suffers from the disorder or its symptoms, such as causing regression of the disorder or its symptoms, for example, by restoring or repairing a lost, missing or defective function, or stimulating an inefficient process, or (4) relieving, alleviating or ameliorating the disorder, or symptoms associated therewith, where ameliorating is used in a broad sense to refer to at least a reduction in the magnitude of a parameter, such as inflammation, pain and/or tumor size.

The term "disease" refers to any condition, infection, disorder or syndrome that requires medical intervention or for which medical intervention is desirable. Such medical intervention can include treatment, diagnosis and/or prevention.

The term "therapeutically effective amount" refers to an amount which, when administered to a living subject, achieves a desired effect on the living subject. For example, an effective amount of the polypeptide of the invention for administration to the living subject is an amount that prevents and/or treats an integrin $\alpha\beta 3$ -mediated disease. The exact amount will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques. As is known in the art, adjustments for systemic versus localized delivery, age, body weight, general health, sex, diet, time of administration, drug interaction and the severity of the condition may be necessary, and will be ascertainable with routine experimentation by those skilled in the art.

The term "T₁" refers to a point in time at which an etanercept formulation has been stored for about one week at 40° C

The term "T₂" refers to a point in time at which an etanercept formulation has been stored for about two weeks at 40° C

5 The term "T₄" refers to a point in time at which an etanercept formulation has been stored for about four weeks at 40° C.

The term "M₃" refers, collectively, to three points in time, and in particular to an analytical result being observed for an etanercept formulation after duration of either about one, about two or about three months of storage at a storage
10 temperature of 5° C. For example, reference herein to an analysis being conducted at M₃ should be understood to mean that such analysis is to be done at the point in time at which etanercept formulation has been in storage for a duration selected from about one, about two, or about three months. Thus, a requirement herein that an etanercept formulation elicit a certain analytical value or measurement at M₃ is
15 satisfied if the required value is observed at a point in time corresponding to at least one of the following storage durations: at approximately one month, at approximately two months or at approximately three months of storage at 5°C.

The terms "Peak 1," "Peak 2" and "Peak 3" when used herein in connection with discussion of HIC chromatography results refers to the same peaks 1, 2 and 3
20 discussed in US Patent 7,294,481.

Embodiments of the Invention

When pharmaceutical compositions containing etanercept (Enbrel®), including aqueous and lyophilized formulations of etanercept are stored on a long term basis, the activity of etanercept can be lost or decreased due to instability of the
25 etanercept monomer via aggregation and/or chemical degradation including formation of fragments. Thus, the present invention provides several embodiments of aqueous formulations of etanercept that allow stable long-term storage of etanercept, so that etanercept is stable over the course of storage either in liquid or frozen states. The provided formulations include, but are not limited to formulations
30 that do not contain arginine and do not require any extra steps such as rehydrating.

These embodiments are explained in a greater detail below.

Etanercept

All of the compositions of the present invention comprise etanercept (Enbrel®). As explained in the Background section of this application, etanercept is a dimeric fusion polypeptide consisting of the extracellular ligand-binding portion of the human 75 kilodalton (p75) tumor necrosis factor receptor (TNFR) linked to the Fc portion of human IgG1. Etanercept consists of 934 amino acids. The Fc component of etanercept contains the constant heavy 2 (CH2) domain, the constant heavy 3 (CH3) domain and hinge region of human IgG1. An Fc domain can contain one or all of the domains described above.

Etanercept suitable for storage in the present pharmaceutical composition can be produced by living host cells that express etanercept, such as hybridomas in the case of antibodies, or host cells that have been genetically engineered to produce the polypeptide in the case of fusion polypeptides or antibodies. Methods of genetically engineering cells to produce polypeptides are well known in the art. See, e.g., Ausubel et al., eds. (1990), *Current Protocols in Molecular Biology* (Wiley, New York). Such methods include introducing nucleic acids that encode and allow expression of the polypeptide into living host cells. These host cells can be bacterial cells, fungal cells, or, preferably, animal cells grown in culture. Bacterial host cells include, but are not limited to, *Escherichia coli* cells. Examples of suitable *E. coli* strains include: HB101, DH5.alpha, GM2929, JM109, KW251, NM538, NM539, and any *E. coli* strain that fails to cleave foreign DNA. Fungal host cells that can be used include, but are not limited to, *Saccharomyces cerevisiae*, *Pichia pastoris* and *Aspergillus* cells. A few examples of animal cell lines that can be used are CHO, VERO, BHK, HeLa, Cos, MDCK, 293, 3T3, and W138. New animal cell lines can be established using methods well known by those skilled in the art (e.g., by transformation, viral infection, and/or selection). Optionally, etanercept can be secreted by the host cells into the medium.

Purification of the expressed etanercept can be performed by any standard method. When etanercept is produced intracellularly, the particulate debris is removed, for example, by centrifugation or ultrafiltration. When etanercept is secreted into the medium, supernatants from such expression systems can be first

concentrated using standard polypeptide concentration filters. Protease inhibitors can also be added to inhibit proteolysis and antibiotics can be included to prevent the growth of microorganisms.

Etanercept can be purified using, for example, hydroxyapatite
5 chromatography, gel electrophoresis, dialysis, and affinity chromatography, and any combination of known or yet to be discovered purification techniques, including but not limited to Protein A chromatography, fractionation on an ion-exchange column, ethanol precipitation, reverse phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSE®, an anion or cation exchange resin
10 chromatography (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation.

Etanercept Formulations Stabilized with NaCl

The invention provides an aqueous etanercept formulation stabilized to reduce instability, aggregation and/or fragmentation of the etanercept, said
15 formulation comprising about 25 to about 75 mg/ml of etanercept and one or more stabilizers, wherein the stabilizers are selected from the group consisting of (i) sodium chloride and (ii) sodium chloride in combination with sucrose or trehalose; and (iii) a combination of sodium chloride, sucrose and trehalose.

The pharmaceutical compositions of the invention may be prepared by
20 combining, a purified etanercept and sodium chloride, optionally with sucrose and/or trehalose. Further, a buffer, a tonicity modifier and an additional excipient and other commonly used inactive ingredients can be added as needed. For simplicity, these are discussed more fully later in the specification. A person of ordinary skill in the art will understand that the combining of the various components to be included in the
25 composition can be done in any appropriate order. For example, the buffer can be added first, middle or last, and the tonicity modifier can also be added first, middle or last. A person of ordinary skill in the art will also understand that some of these chemicals can be incompatible in certain combinations, and accordingly, are easily substituted with different chemicals that have similar properties but are compatible in
30 the relevant mixture.

In an embodiment using sodium chloride for stabilization, an etanercept formulation of the invention comprises about 25 to 75 mg/ml etanercept, up to about 150 mM of sodium chloride, about 1 to about 30 mM sodium phosphate; and about 0 to 5 wt.% sucrose or trehalose or combination thereof; wherein the composition has a pH of about 6.0 to about pH 7.0, and more preferably about 6.0 to about 6.6 and most preferably about 6.3 to about 6.5.

The sodium chloride stabilized composition is preferably characterized by SEC analysis at T_2 in which: monomer content is greater than about 80 wt.%; aggregate(s) content is less than about 3 wt.%, and fragment 3 content is less than about 8 wt.%.

The sodium chloride-stabilized etanercept composition is preferably characterized by:

(a) an SEC analysis at T_4 of greater than about 90, 91, 92, 93, 94, 95, 96, or 97 wt.% monomer content; and less than about 3, 2 or 1 wt.% aggregate(s) content; and

(b) an HIC analysis at T_2 wherein the amount of the composition represented by peak 1 of the HIC chromatogram is less than about 4, 3, 2 or 1 wt.%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than 80, 81, 82, 83, 84 85 or 86 wt.%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than about 20, 19, 18, 17, 16, 15, 14, or 13 wt.%; and

(c) an HIC analysis at T_4 wherein the amount of the composition represented by peak 1 of the HIC chromatogram is less than about 3, 2 or 1 wt.%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than 80, 81, 82, 83, 84 or 85 wt.%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than 20, 19, 18, 17, 16, 15, 14 or 13 wt.%.

In a further embodiment, preferred composition using sodium chloride for stabilization comprise up to about 150 mM sodium chloride, about 1 to 30 mM sodium phosphate, and about 0-5 wt.% sucrose or trehalose, or combination of sucrose and trehalose and having a pH of about 6.0 to 6.6; and characterized by: an

SEC analysis at T_4 of greater than about 95 wt.% monomer content and less than about 1 wt.% aggregate(s) content; an HIC analysis at T_2 wherein the amount of the composition represented by peak 1 of the HIC chromatogram is less than or equal to about 3 wt.%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than about 82 wt.%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than about 15 wt.%; and an HIC analysis at T_4 wherein the amount of the composition represented by peak 1 of the HIC chromatogram is less than about 2 wt.%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than about 84 wt.%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than or equal to about 14 wt.%.

Particularly preferred compositions in terms of reduced aggregates and fragments are those in which the sodium chloride stabilized etanercept formulations exhibit HIC analysis at T_4 or T_2 wherein the amount of the composition represented by peak 1 of the HIC chromatogram is less than about 1%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than about 95 wt.% and preferably greater than about 99 wt.%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than about 3 wt.%.

In a further embodiment of the invention, an NaCl stabilized etanercept formulation contains up to about 5 mM arginine.

In the above-referenced NaCl stabilized etanercept formulations, the terms "SEC", " T_2 ", " T_4 ", "HIC", "monomer content", "aggregate(s)" and "fragment 3", "peak 1," "peak 2," and "peak 3," are defined in the examples below.

Additional Components of the Provided Pharmaceutical Compositions

The formulations of the invention may also include buffers, tonicity modifiers, excipients, pharmaceutically acceptable carriers and other commonly used inactive ingredients of the pharmaceutical compositions. For simplicity, these are discussed more fully later in the application.

Buffers maintain pH in a desired range. Suitable buffers include histidine, potassium phosphate, sodium or potassium citrate, maleic acid, ammonium acetate,

tris-(hydroxymethyl)-aminomethane (tris), various forms of acetate and diethanolamine. The concentration of the buffer in the formulation is preferably between about 1 mM to about 1M, and more preferably about 10 mM to about 200 mM. Buffers are well known in the art and are manufactured by known methods and
5 available from commercial suppliers.

Examples of suitable buffers are phosphate, histidine, citrate, maleate, tartrate, succinate, acetate, tris-(hydroxymethyl)-aminomethane (tris), bicarbonate.

In a preferred embodiment, the buffer is sodium phosphate.

In a preferred embodiment, the pH of the pharmaceutical composition is at or
10 near physiological levels. Thus, preferably, the pH of the provided compositions is between about 5.8 and about 8.4; and even more preferably, between about 6.2 and about 7.4. A person of ordinary skill in the art will understand that the pH can be adjusted as necessary to maximize stability and solubility of etanercept in a particular formulation. Thus, etanercept formulations at a pH outside of physiological
15 ranges, yet tolerable to the patient, are also within the scope of the invention.

A tonicity modifier is a molecule that contributes to the osmolality of a solution. The osmolality of a pharmaceutical composition is preferably adjusted to maximize the active ingredient's stability and/or to minimize discomfort to the patient upon administration. It is generally preferred that a pharmaceutical composition be isotonic
20 with serum, i.e., having the same or similar osmolality, which is achieved by addition of a tonicity modifier.

In a preferred embodiment, the osmolality of the provided formulations is from about 180 to about 420 mOsm. However, it is to be understood that the osmolality can be either higher or lower as specific conditions require.

25 Examples of tonicity modifiers suitable for modifying osmolality include, but are not limited to amino acids (not including arginine) (e.g., cysteine, histidine and glycine), salts (e.g., sodium sulfate, potassium chloride and sodium citrate) and/or saccharides (e.g., sucrose, glucose and mannitol).

Preferred tonicity modifiers are glycine, alanine, sodium chloride, potassium
30 chloride, and sodium sulfate.

In a preferred embodiment, the concentration of the tonicity modifier in the formulation is preferably between about 1 mM to about 1 M, more preferably about 10 mM to about 200 mM. Tonicity modifiers are well known in the art and are manufactured by known methods and available from commercial suppliers.

5 Excipients, also referred to as chemical additives, co-solutes, or co-solvents, that stabilize the polypeptide while in solution (also in dried or frozen forms) can also be added to a pharmaceutical composition. Excipients are well known in the art and are manufactured by known methods and available from commercial suppliers.

10 Examples of suitable excipients include but are not limited to sugars/polyols such as: sucrose, lactose, glycerol, xylitol, sorbitol, mannitol, maltose, inositol, trehalose, glucose; polymers such as: serum albumin (bovine serum albumin (BSA), human SA or recombinant HA), dextran, poly(vinyl alcohol) PVA, hydroxypropyl methylcellulose (HPMC), polyethyleneimine, gelatin, polyvinylpyrrolidone (PVP), hydroxyethylcellulose (HEC); non-aqueous solvents such as: polyhydric alcohols,
15 (e.g., PEG, and glycerol) and dimethylformamide (DMF); amino acids such as: proline, L-serine, sodium glutamic acid, alanine, glycine, lysine hydrochloride, sarcosine and gamma-aminobutyric acid; surfactants such as: Tween®-80 (polysorbate 80), Tween®-20 (polysorbate 20), SDS, polysorbate, poloxamers; and
20 miscellaneous excipients such as: potassium phosphate, sodium acetate, ammonium sulfate, magnesium sulfate, sodium sulfate, trimethylamine N-oxide, betaine, metal ions (e.g., zinc, calcium, and magnesium), CHAPS, monolaurate, 2-O-beta-mannoglycerate or any combination of the above.

25 Preferred excipients are sucrose, lactose, glycerol, xylitol, sorbitol, mannitol, maltose, inositol, trehalose, glucose, bovine serum albumin (BSA), human serum albumin (HSA), recombinant albumin, dextran, PVA, hydroxypropyl methylcellulose (HPMC), polyethyleneimine, gelatin, polyvinylpyrrolidone (PVP), hydroxyethylcellulose (HEC), polyethylene glycol, ethylene glycol, glycerol, alanine, glycine, lysine hydrochloride, sarcosine, SDS, polysorbate 20, polysorbate 80, poloxamer 188, trimethylamine N-oxide, betaine, zinc ions, calcium ions, magnesium
30 ions, CHAPS, sucrose monolaurate, and 2-O-beta-mannoglycerate.

The concentration of one or more excipients in a formulation of the invention is/are preferably between about 0.001 to 5 weight percent, more preferably about 0.1 to 2 weight percent.

Methods of Treatment

5 In another embodiment, the invention provides a method of treating a mammal comprising administering a therapeutically effective amount of the pharmaceutical compositions of the invention to a mammal, wherein the mammal has a disease or disorder that can be beneficially treated with etanercept.

10 In a preferred embodiment, the etanercept is derived from the same species of mammal as is to be treated with the composition.

In a preferred embodiment, the mammal is a human.

Diseases or disorders that can be treated with the provided compositions include but are not limited to rheumatoid arthritis, psoriatic arthritis, ankylosing spondylitis, Wegener's disease (granulomatosis), Crohn's disease (or inflammatory
15 bowel disease), chronic obstructive pulmonary disease (COPD), Hepatitis C, endometriosis, asthma, cachexia, psoriasis, and atopic dermatitis. Additional diseases or disorders that can be treated with the compositions of the present invention include those described in WO 00/62790, WO 01/62272, U.S. Patent Application No. 2001/0021380, and US Pat. 7,648,702 B2, the relevant portions of
20 which are incorporated herein by reference.

The provided pharmaceutical compositions may be administered to a subject in need of treatment by injection systemically, such as by intravenous injection; or by injection or application to the relevant site, such as by direct injection, or direct application to the site when the site is exposed in surgery; or by topical application.

25 In one embodiment, the invention provides a method of treatment and/or prevention of rheumatoid arthritis comprises administering to a mammal in need thereof a therapeutically effective amount of one of the provided etanercept compositions.

30 The therapeutically effective amount of the etanercept in the provided compositions will depend on the condition to be treated, the severity of the condition,

prior therapy, and the patient's clinical history and response to the therapeutic agent. The proper dose can be adjusted according to the judgment of the attending physician such that it can be administered to the patient one time or over a series of administrations.

5 In one embodiment, the effective etanercept amount per adult dose is from about 1-500 mg/m², or from about 1-200 mg/m², or from about 1-40 mg/m² or about 5-25 mg/m².

 Alternatively, a flat dose may be administered, whose amount may range from 2-500 mg/dose, 2-100 mg/dose or from about 10-80 mg/dose.

10 If the dose is to be administered more than one time per week, an exemplary dose range is the same as the foregoing described dose ranges or lower and preferably administered two or more times per week at a per dose range of 25-100 mg/dose.

 In another embodiment, an acceptable dose for administration by injection
15 contains 80-100 mg/dose, or alternatively, containing 80 mg per dose.

 The dose can be administered weekly, biweekly, or separated by several weeks (for example 2 to 8).

 In one embodiment, etanercept is administered at 25 to 75 mg/ml by a single subcutaneous (SC) injection.

20 In some instances, an improvement in a patient's condition will be obtained by administering a dose of up to about 100 mg of the pharmaceutical composition one to three times per week over a period of at least three weeks. Treatment for longer periods may be necessary to induce the desired degree of improvement. For incurable chronic conditions the regimen may be continued indefinitely. For pediatric
25 patients (ages 4-17), a suitable regimen may involve administering a dose of 0.4 mg/kg to 5 mg/kg of etanercept, one or more times per week.

 In another embodiment, the pharmaceutical formulations of the invention may be prepared in a bulk formulation, and as such, the components of the pharmaceutical composition are adjusted to be higher than would be required for
30 administration and diluted appropriately prior to administration.

The pharmaceutical compositions can be administered as a sole therapeutic or in combination with additional therapies as needed. Thus, in one embodiment, the provided methods of treatment and/or prevention are used in combination with administering a therapeutically effective amount of another active agent. The other
5 active agent may be administered before, during, or after administering the pharmaceutical compositions of the present invention. Another active agent may be administered either as a part of the provided compositions, or alternatively, as a separate formulation.

Administration of the provided pharmaceutical compositions can be achieved
10 in various ways, including parenteral, peroral, buccal, sublingual, nasal, rectal, intraperitoneal, intradermal, transdermal, subcutaneous, intravenous, intra-arterial, intracardiac, intraventricular, intracranial, intratracheal, intrathecal administration, intramuscular injection, intravitreal injection, and topical application.

The pharmaceutical compositions of this invention are particularly useful for
15 parenteral administration, i.e., subcutaneously, intramuscularly, intravenously, intraperitoneal, intracerebrospinal, intra-articular, intrasynovial, intravitreal, and/or intrathecal. Parenteral administration can be by bolus injection or continuous infusion. Pharmaceutical compositions for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. In
20 addition, a number of recent drug delivery approaches have been developed and the pharmaceutical compositions of the present invention are suitable for administration using these new methods, e.g., Inject-ease®, Genject®, injector pens such as GenPen®, and needleless devices such as MediJector® and BioJector®. The present pharmaceutical composition can also be adapted for yet to be discovered
25 administration methods. See also Langer, 1990, Science, 249:1527-1533.

The provided pharmaceutical compositions can also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the formulations may be modified with suitable polymeric or hydrophobic
30 materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

The pharmaceutical compositions may, if desired, be presented in a vial, pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. In one embodiment the dispenser device can comprise a syringe having a single dose of the liquid formulation ready for injection. The syringe can be accompanied by instructions for administration.

In another embodiment, the present invention is directed to a kit or container, which contains an aqueous pharmaceutical composition of the invention. The concentration of the polypeptide in the aqueous pharmaceutical composition can vary over a wide range, but is generally within the range of from about 0.05 to about 20,000 micrograms per milliliter ($\mu\text{g/ml}$) of aqueous formulation. The kit can also be accompanied by instructions for use.

The present invention is more particularly described in the following examples that are intended as illustrative only, since many modifications and variations therein will be apparent to those skilled in the art.

EXAMPLE

Etanercept Stabilized with NaCl

Etanercept formulations stabilized with NaCl, alone, or NaCl in combination with sucrose, trehalose and/or arginine, may be prepared and tested using the procedures generally described below.

Each solid formulation component is weighed to the amount required for a given volume of formulation buffer. These components are combined into a beaker or vessel capable of carrying and measuring the given volume of formulation buffer. A volume of deionized water equal to approximately $\frac{3}{4}$ of the target given formulation buffer is added to the beaker, and the components are then solublized. The pH of the buffer is adjusted to the target formulation pH using 1 M sodium hydroxide and/or 1 M hydrogen chloride. The final formulation buffer volume is then raised to the target volume through the addition of deionized water. Etanercept protein solution is placed in dialysis material housing (such as Thermo Scientific Slide-A-Lyzer MINI Dialysis Unit 10,000 MWCO), which is then placed in contact with the desired formulation buffer for 12 hours at 4°C. Formulation buffer volume to protein solution volume ratio should be no less than 1000:1. The dialysis housing and protein

solution it contains is then placed in a second, equal volume of formulation buffer for an additional 12 hours at 4°C. Resulting protein solution is removed from the dialysis material housing, and the concentration of protein determined using ultraviolet spectroscopy. Protein concentration is adjusted to the desired level using centrifugation (such as Amicon Ultra 10,000 MWCO Centrifugal Concentrators) and/or dilution with formulation buffer.

The compositions can be tested for long-term stability by size exclusion chromatography (SEC), denatured SEC (dSEC), hydrophobic interaction chromatography (HIC), sodium dodecylsulfate polyacrylamide gel electrophoresis (SDS-PAGE), and for binding and bioactivity at various timepoints. The bioactivity can be measured by any number of well-known assays.

For example, the techniques of Size Exclusion Chromatography are described in Hawe et al, Pharm. Res. 2011, 28: 2302 and/or van Marrschalkerweerd et al., Eur. J. Pharm. Biopharm. 2011, 78: 213. Similarly, the techniques of Denatured Size Exclusion Chromatography, Hydrophobic Interaction Chromatography, and Sodium DodecylSulfate-PolyAcrylamide Gel Electrophoresis are also well known to persons having ordinary skill in the art.

It is believed that the composition will be stable over the term of two years or more.

(Formulation 2:8)

Ingredient	concentration
Etanercept (active ingredient)	50 mg/ml
Sodium phosphate, pH 6.32 (inactive)	25 mM
NaCl (inactive)	150 mM

(Formulation 2:6)

Ingredient	concentration
Etanercept (active ingredient)	50 mg/ml
Sodium phosphate, pH 6.3 (inactive)	15 mM

NaCl (inactive)	100 mM
Sucrose (inactive)	2 % (w/v)

(Formulation 3:10)

Ingredient	concentration
Etanercept (active ingredient)	50 mg/ml
Sodium phosphate, pH 6.57 (inactive)	10 mM
NaCl (inactive)	75 mM
Sucrose (inactive)	3 % (w/v)

(Formulation 3:11)

Ingredient	concentration
Etanercept (active ingredient)	50 mg/ml
Sodium phosphate, pH 6.30 (inactive)	25 mM
NaCl (inactive)	75 mM
Trehalose (inactive)	3 % (w/v)

5

(Formulation 3:12)

Ingredient	concentration
Etanercept (active ingredient)	50 mg/ml
Sodium phosphate, pH 6.3 (inactive)	25 mM
NaCl (inactive)	75 mM
Sucrose (inactive)	3 % (w/v)

(Formulation 3:13)

Ingredient	concentration
Etanercept (active ingredient)	50 mg/ml
Sodium phosphate, pH 6.3 (inactive)	25 mM
NaCl (inactive)	120 mM

Sucrose (inactive)	1% (w/v)
Arginine (inactive)	5 mM

EXAMPLE 2

Preparation of Etanercept

STEP 1. Cell Expansion. In a manner known in the art, cell expansion
 5 necessary to generate a sufficient number of cells for inoculation of a production
 bioreactor is performed using a clone of CHO cells expressing the etanercept fusion
 protein. The product of this expression process (a harvested cell culture fluid)
 results in a mixture of correctly folded etanercept, as well as incorrectly folded and/or
 aggregated etanercept, along with additional impurities. The harvested cell culture
 10 fluid comprising such protein mixture is subjected to detergent viral inactivation.

STEP 2. Affinity Chromatography. Affinity chromatography is
 performed on the harvested cell culture obtain in Step 1 above using a conventional
 Protein A affinity column in a well known manner. Product recovery is approximately
 85%. The product obtained is a complex protein mixture comprising correctly folded
 15 etanercept, incorrectly folded etanercept, and/or aggregates of correctly and/or
 incorrectly folded etanercept, or protein fragments. The product obtained from this
 Protein A affinity column purification step is adjusted to pH 3.5 and then subjected to
 a viral inactivation step. Following viral inactivation the product is adjusted to pH 5.5
 and then further clarified in a known manner using a commercially obtained capsule
 20 filter.

STEP 3A. Mixed-Mode Cation Exchange Chromatography. A 31.8 L
 (45 cm diameter X 20 cm bed height) packed bed GE Healthcare Capto MMC
 chromatography column is used to purify the product obtained in Step 2 above. Prior
 to use, the column is equilibrated with 2 CV of 25 mM acetate pH 5.5 and sanitized
 25 with 2 CV of 0.1 N NaOH, 1 M NaCl and neutralized with 2 CV of 25 mM acetate, 0.7
 M NaCl, pH 5.5. The column is then equilibrated with 8-10 CV of 25 mM acetate pH
 5.5 until the effluent is pH 5.5 and 3.5 mS/cm. The Protein A pool from step 2 above
 is diluted to ≤ 6 mS/cm with WFI and applied to a column loading of up to 15 g/L
 media for each cycle. The column is operated at a linear velocity of 200 cm/h to give

a 6 minute residence time. After loading, the column is washed with 2 CV of 25 mM acetate pH 5.5. The product is then eluted with an 8.5 CV, 15% to 85% gradient of 25 mM acetate pH 5.5 to 25 mM acetate, 0.7 M NaCl, pH 5.5. Product collection begins at 0.15 OD (A280, 1.0 cm path length) and collection ends at 50% of peak maximum. The eluate volume is approximately 5 CV. Residual product and contaminants are stripped from the column with 2 CV of 10 mM Tris, 1 M NaCl, pH 8.0 and discarded. The product obtained from the mixed mode column is filtered using a Millipore Opticap XL10, 0.22 μ m Durapore capsule filter, (0.69 m²). The product obtained from this step represents a recovery of about 70% of the Protein A material obtained in Step 2

STEP 3B. Mixed Mode Anion Exchange Chromatography. A 27.0 L (45 cm diameter X 17 cm bed height) packed bed GE Healthcare Canto Adhere chromatography column is used to further purify the product obtained in step 3A above. Prior to use, the column is equilibrated with 2 CV of 25 mM Tris, pH 8.0 and sanitized with 2 CV of 0.1 N NaOH, 1M NaCl and neutralized and equilibrated with 2 CV of 25 mM Tris, pH 8.0. Prior to product loading, the column is equilibrated with 3 CV of 10 mM Tris, pH 8.0. The Canto MMC pool from Step 3A above is adjusted to pH 8.1 with ~0.045 kg of 1 M Tris, pH 8.3 per kg of pool. The product from Step 3A above was diluted in-line 1:3.8 with WFI to adjust the conductivity to 12.0 mS/cm and pH 8.0. The resulting material is then applied to a column loading of up to 15 g/L media. The column is operated at a linear velocity of 170 cm/h to give a 6 minute residence time. After loading, the column is washed with 2 CV of 25 mM Tris, pH 8.0. The product is then eluted with a 10 CV gradient (20% to 90%) of 25 mM Tris, pH 8.0 to 10 mM Tris, 1 M NaCl, pH 8.0. Product collection is started at 0.15 OD (A280, 1.0 cm path length) and collection ended at 25% of peak maximum. The eluate volume is 4-6 CV. The eluted product is filtered using a commercially available capsule filter and then subjected in a known manner to viral filtration and tangential flow filtration steps. Overall product recovery from step3B (including the final viral and tangential flow filtration steps) was approximately 68%. Product recovery measured before the filtration steps was about 75%. A schematic representation of HIC data obtained on elution fractions from this step are representing in Figure 12.

Analysis: The final filtered product obtained in this example is found to have greater than about 90 wt % correctly folded etanercept as determined by HIC;

less than 5 wt% incorrectly folded etanercept species as determined by HIC; less than about 3 wt% of clipped material by HIC analysis (believed to be fragments of etanercept in which the TNFR portion thereof has been truncated) and a combined amount of correctly and incorrectly folded etanercept of greater than 95 wt% as
5 determined by size exclusion chromatography.

Analysis of Etanercept Formulations

A. Thermal Stability Storage

Following dialysis and concentration, samples of the etanercept formulations
10 exemplified above were sterile filtered in a bio safety cabinet. Using sterilized pipettes and autoclaved pipette tips, samples of the etanercept formulations were transferred to pre-labeled and autoclaved 1 mL lyophilization vials. Vials were stoppered with sterile butyl stoppers and crimped with aluminum caps. All vials were then transferred to thermal stability ovens. Samples were subject to two thermal
15 stability regimes: (1) two weeks at 40 °C and (2) four weeks at 25 °C. Throughout this specification, these two temperature regimes are denoted "T₂" and T₄," respectively.

B. Size Exclusion Chromatography (SEC)

20

Etanercept formulations disclosed herein were analyzed using the well known technique of Size Exclusion Chromatography (SEC), a high-performance liquid chromatography method in which analytes are separated by size (see Rogner, M. (2000). Size Exclusion Chromatography. Protein Liquid Chromatography. M. Kastner. Amsterdam, Elsevier. 61: 89-145.). In order to evaluate thermal stability of the Etanercept samples described above, the samples were examined by a SEC method based on the literature (van Maarschaalkerweerd, A., G. J. Wolbink, et al. (2011). "Comparison of analytical methods to detect instability of etanercept during thermal stress testing." European Journal of Pharmaceutics and Biopharmaceutics
25 78(2): 213-221.) The mobile phase buffer was prepared to contain 50 mM sodium phosphate monobasic monohydrate and 150 mM arginine. The pH was adjusted to
30

6.5 using 1 M HCl. All separations were performed using a Tosoh TSK-Gel SWxl 6 mm x 4 cm guard column (cat. no. 8543) attached linearly to a Tosoh TSK-Gel G4000 SWxl 7.8 mm x 30 cm (cat. no. 8542). To perform a separation, the columns were brought to room temperature (23°C) and equilibrated with mobile phase at a flow rate of 0.5 mL/min. 5 microliters of 50 mg/mL etanercept formulation were injected onto the column using an autosampler. The separation was accomplished over 30 minutes at a flow rate of 0.5 mL/minute. Column eluent was monitored at a wavelength of 280 nm during this time.

C. Integration of Size Exclusion Chromatography Chromatograms

All integration was performed using Chromeleon software (Dionex). Prior to integration, the SEC chromatogram for a buffer containing no etanercept was subtracted from all chromatograms. All integration was performed between retention times of 12 minutes and 26 minutes. Several parameters were used to define a peak. The minimum area for a detected peak was set to 0.05 mAu * min. The two-dimensional sensitivity for peak detection was set to 0.01 mAu and 75 seconds. Peak shoulders were added manually using a manual integration tool. All detected peaks were manually adjusted in two steps. First, peak baselines (the bottom boundary of the peak) were adjusted to horizontal. Secondly, the vertical positions of the peak baselines were adjusted to that of the chromatogram baseline. The chromatogram baseline value was defined as the signal in absence of analyte. The signal in absence of analyte was defined as the absorbance in mAu at 12 minutes retention time.

D. SEC Fractions of Etanercept Formulations

In the SEC analysis of etanercept formulations described above, three SEC chromatography fractions were identified and studied. The fractions that were analyzed were, in the order of elution from the SEC column: (1) a high molecular weight fraction representing aggregates of the intact etanercept TNFR:Fc fusion protein likely assembled via non-covalent electrostatic attraction among intact etanercept molecules (hereinafter "aggregate(s)" or aggregate(s) content); (2) monomer content, representing the intact etanercept TNFR:Fc fusion protein

(hereinafter referred to as "monomer" of "monomer content"); (3) a fraction likely representing one fragment or a population of fragments of the etanercept molecule in which one portion of the TNFR:molecule fusion protein has become cleaved from the monomer; in the loss of an arm of the Fabportion of the fusion protein at the hinge region of the molecule. The most common fragment or clipped species, as measured by SEC, is referred to as Fragment 3. In conducting the SEC analysis, it will be observed that aggregates elute first, followed by monomer, followed by fragment 3.

The following tables shows the relative amounts of Aggregates, Monomer and Fragment 3 determined by SEC analysis as described above.

TABLE 1

SEC ANALYSIS OF MONOMER

Note: Amounts reported Tables I, II and III are percentages by weight

T₀ = formulation maintained at 5 C and analyzed within 24 hours of creation.

T₁ = formulation stored for one week at 40° C

T₂ = formulation stored for two weeks at 40 C

Formulation No.	t ₀	t ₁	t ₂
Commercial Enbrel (comparative) [1:2]	98.81	92.58	87.64
xx2:6	98.09	--	87.77
xx2:8	98.07	--	88.38
xx3:10	98.16	--	89.77
xx3:11	98.32	--	89.87
xx3:12	98.33	--	90.92
xx3:13	98.18	--	90.74

TABLE II

SEC ANALYSIS OF AGGREGATES

Note: Amounts reported Tables I, II and III are percentages by weight

T₀ = formulation maintained at 5 C and analyzed within 24 hours of creation.

5 T₁ = formulation stored for one week at 40° C

T₂ = formulation stored for two weeks at 40 C

Formulation No.	t ₀	t ₁	t ₂
Commercial Enbrel (comparative)	0.09	0.59	1.02
2:6	0.30	--	1.81
2:8	0.30	--	1.42
3:10	0.27	--	1.33
3:11	0.20	--	1.24
3:12	0.21	--	0.85
3:13	0.28	--	0.86

10

15

TABLE III
ANALYSIS OF FRAGMENT 3

Formulation No	t_0	t_1	t_2
Commercial Enbrel (comparative)	0.00	3.30	6.29
2:6	0.00	--	4.73
2:8	0.00	--	6.29
3:10	0		3.76
3:11	0		3.59
3:12	0		3.68
3:13	0		3.88

5

10

TABLE IV**SEC MONOMER CONTENT**

(4 weeks/25° C)

Table IV below shows monomer (etanercept) content of etanercept formulations prepared according to the present invention, when stored for four weeks at 25 C°- denoted by the symbol T₄. In the following table T₀ represents SEC measurements conducted within 24 hours of formulation preparation, at sample temperature of 5° C; and T₄ represents etanercept formulation samples subjected to SEC analysis after 4 weeks storage at 25° C.

FORMULATION No.	T ₀ Monomer Content	T ₄ Monomer Content
Commercial Enbrel (comparative)	98.15	97.86
3:10	98.16	97.66
3:11	98.32	97.75
3:12	98.33	97.90
3:13	98.18	97.78

TABLE V**SEC AGGREGATES CONTENT****(4 weeks/25° C)**

Table V below shows aggregate(s) content of etanercept formulations prepared according to the present invention after storage for four weeks at 25 C°. In the following table T₀ represents SEC measurements conducted within 24 hours of formulation preparation, at sample temperature of 5° C; and T₄ represents etanercept formulation samples subjected to SEC analysis after 4 weeks storage at 25° C.

FORMULATION No.	T₀ Aggregate(s) Content	T₄ Aggregate(s) Content
Commercial Enbrel (comparative)	0.28	0.25
3:10	0.27	0.32
3:11	0.20	0.27
3:12	0.21	0.26
3:13	0.28	0.32

HIC ANALYSIS OF ETANERCEPT FORMULATIONS

The following tables (Tables VI and VII) show the results of hydrophobic interaction chromatography ("HIC chromatography") conducted on samples 3:5 and 3:8. HIC chromatography was carried out in a manner known in the art and generally described in U.S. Patent 7,294,481, incorporated herein by reference. Samples were evaluated at t_0 (within 24 hours of preparation at 5°C.) and again after either two weeks of storage at 25°C. (t_2) (see Table VI) or after 4 weeks of storage at 25°C. (t_4) (See Table VII). Peak 1 in the HIC chromatogram is believed to be or include "Fragment 3", which is identified and quantified using SEC, as referenced above in the discussion of SEC data; Peak 2 is etanercept monomer as referenced above in the discussion of SEC data; and Peak 3 includes "Aggregate(s)" as referenced above in the discussion of SEC data. It should further be understood that the terms "peak 1", "peak 2" and "peak 3" as used here also constitute a reference to the HIC peak 1, peak 2 and peak 3 referred to and disclosed in Figure 4 of U.S. patent 7,294,481 incorporated herein by reference.

TABLE VI

HIC Data after Two Weeks Storage at 40°C.

	PEAK 1		PEAK 2		PEAK 3	
Form. #	T ₀	T ₂	T ₀	T ₂	T ₀	T ₂
Commercial Enbrel (comparative)	0.91	3.23	86.72	83.41	12.33	13.36
3:10	0.74	3.06	87.36	84.24	11.90	12.70
3:11	0.56	3.10	86.46	83.73	12.98	13.18
3:12	0.68	3.07	86.80	83.52	12.52	13.40
3:13	0.77	2.86	86.45	84.33	12.78	12.82

TABLE VII

HIC Data after Storage at 25°C for 4 Weeks

	PEAK 1		PEAK 2		PEAK 3	
Form. #	T ₀	T ₄	T ₀	T ₄	T ₀	T ₄
Commercial Enbrel (comparative)	0.91	1.09	86.76	86.95	12.33	11.97
3:10	0.74	1.03	87.36	85.83	11.90	13.14
3:11	0.56	1.11	86.46	85.32	12.98	13.57
3:12	0.68	0.81	86.80	86.36	12.52	12.83
3:13	0.77	1.01	86.45	85.78	12.78	13.21

5. Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the invention being indicated by the following claims.

WHAT IS CLAIMED IS:

1. An aqueous etanercept formulation stabilized to reduce instability, aggregation misfolding and/or fragmentation of the etanercept, said formulation comprising about 25 to about 75 mg/ml of etanercept and one or more stabilizers, wherein the stabilizers are selected from the group consisting of (i) sodium chloride and (ii) sodium chloride in combination with sucrose or trehalose; and (iii) a combination of sodium chloride, sucrose and trehalose.
2. The formulation of claim 1, further comprising one or more additional components selected from: of a buffer, a tonicity modifier and an excipient.
3. The formulation of claim 2 wherein the stabilizer consists essentially of either: (a) sodium chloride; or (b) sodium chloride in combination with sucrose; or (c) sodium chloride in combination with trehalose; and wherein, the formulation, optionally, is free or essentially free of arginine, and the formulation elicits long term storage stability as characterized by at least one of:

SEC analysis at M_3 or T_2 or T_4 of: monomer content greater than about 90%; aggregates content of less than about 3 wt%; and fragment 3 content less than about 5 wt%; and

HIC analysis at M_3 or T_2 or T_4 wherein the amount of the composition represented by peak 1 of the HIC chromatogram is less than about 3 wt.%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than 80 wt.%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than about 20 wt. %.

4. The formulation of claim 2 comprising about 25 to 75 mg/ml etanercept, sodium chloride in an amount up to about 150 mM, about 1 to about 30 mM sodium phosphate; and about 0 to 5 wt.% sucrose or trehalose or combination thereof; wherein the composition has a pH of about 6.0 to about 6.6; and wherein the composition is characterized by SEC analysis at M_3 or T_2 or T_4 in

which: monomer content is greater than about 80 wt.%; aggregate(s) content is less than about 3 wt.%, and fragment 3 content is less than about 8 wt.%.

5. The formulation of claim 4, optionally essentially free of arginine, eliciting long term storage stability characterized by:
 - (a) an SEC analysis at M_3 or T_2 or T_4 of greater than about 90 wt.% monomer content; and less than about 3 wt.% aggregate(s) content; and
 - (b) an HIC analysis at M_3 or T_2 or T_4 wherein the amount of the composition represented by peak 1 of the HIC chromatogram is less than about 4 wt.%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than about 80 wt.%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than about 20 wt.%; and
6. The formulation of claim 5 having an HIC analysis at M_3 or T_2 or T_4 wherein the amount of the composition represented by peak 1 of the HIC chromatogram is less than about 1 wt.%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than about 95 wt.%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than about 3 wt.%.
7. The formulation of claim 2 comprising up to about 150 mM sodium chloride, about 1 to 30 mM sodium phosphate, and about 0-5 wt.% sucrose or trehalose, or combination of sucrose and trehalose, and having a pH of about 6.0 to 6.6; and characterized by: an SEC analysis at M_3 or T_2 or T_4 of greater than about 95 wt.% monomer content and less than about 1 wt.% aggregate(s) content; and an HIC analysis at M_3 or T_2 or T_4 wherein the amount of the composition represented by peak 1 of the HIC chromatogram is less than or equal to about 3 wt.%; the amount of the composition represented by peak 2 of the HIC chromatogram is greater than about 82 wt.%; and the amount of the composition represented by peak 3 of the HIC chromatogram is less than about 15 wt.%.

8. The formulation of claim 3 comprising: about 50 mg/ml of etanercept; about 75 mM sodium chloride; about 10 to 30 mM sodium phosphate; about 3 wt.% sucrose.
9. The formulation of claim 3 comprising: about 50 mg/ml of etanercept; about 75mM sodium chloride; about 10 to 30 mM sodium phosphate; about 3 wt.% trehalose.
10. The formulation of claim 3 comprising: about 50 mg/ml of etanercept; about 75mM sodium chloride; about 10 to 30 mM sodium phosphate; about 3 wt.% sucrose and having pH of about 6.3 to 6.5.
11. The formulation of claim 2 comprising: about 50 mg/ml of etanercept; about 120mM sodium chloride; about 10 to 30 mM sodium phosphate; about 1 wt.% sucrose; about 5 mM arginine, and having pH of about 6.3 to 6.5.
12. The formulation of claim 3 comprising: about 50 mg/ml of etanercept; about 150 mM sodium chloride; about 10 to 30 mM sodium phosphate, in the absence of sucrose or trehalose.
13. The formulation of claim 3 comprising: about 50 mg/ml of etanercept; about 100 mM sodium chloride; about 10 to 30 mM sodium phosphate; and about 2 wt.% sucrose.
14. The formulation of claim 3 eliciting long term storage stability as characterized by: an HIC analysis at M_3 or T_2 or T_4 wherein the amount of the composition represented by peak 2 of the HIC chromatogram is greater than or equal to about 95 wt.%; and wherein, if peak 3 is present on the HIC chromatogram, the amount of the composition represented by peak 3 is less than or equal to about 1 wt.%.
15. The formulation of claim 3 having, at M_3 or T_2 or T_4 no more than, on average, about 10,000 subvisible particles per mL having a size greater than 5 μ m.
16. The formulation of claim 1, containing no arginine, or essentially free of arginine, wherein the composition, at M_3 or T_2 or T_4 , elicits long term storage stability that meets one or both of the following criteria:

(A) stability comparable to or better than commercially available etanercept marketed under the trademark Enbrel®, as measured by (i) SEC analysis of the amounts of aggregate(s), monomer and fragment 3 in the composition (as defined in the specification) and (ii) HIC analysis of amounts of material in the composition corresponding to peaks 1, 2 and 3 of the HIC chromatogram (as defined in the specification); and

(B) an HIC chromatogram in which (i) peak 3 is absent, or essentially absent and (ii) peak 2 represents greater than about 95 wt% of the composition; an SEC chromatogram containing essentially no peak corresponding to aggregate(s); and an SEC chromatogram in which the monomer content represents at least about 95 wt% of the composition.