



(86) **Date de dépôt PCT/PCT Filing Date:** 2007/10/12
(87) **Date publication PCT/PCT Publication Date:** 2008/04/17
(45) **Date de délivrance/Issue Date:** 2017/02/28
(85) **Entrée phase nationale/National Entry:** 2009/04/07
(86) **N° demande PCT/PCT Application No.:** US 2007/021897
(87) **N° publication PCT/PCT Publication No.:** 2008/045559
(30) **Priorités/Priorities:** 2006/10/12 (US60/829,291);
2007/10/12 (US11/871,848)

(51) **Cl.Int./Int.Cl. A61K 31/439** (2006.01),
A61P 1/02 (2006.01), A61K 33/32 (2006.01)


(72) **Inventeurs/Inventors:**
ROTHSTEIN, DAVID M., US;
MURPHY, CHRIS, US;
WONG, VIVIEN, US;
KAZO, GLENN, US

(73) **Propriétaire/Owner:**
GALERA LABS, LLC, US

(74) **Agent:** SMART & BIGGAR

(54) **Titre : MIMETIQUES DE SUPEROXYDE DISMUTASE POUR LE TRAITEMENT DE LA MUCOSITE ORALE**
(54) **Title: SUPEROXIDE DISMUTASE MIMETICS FOR THE TREATMENT OF ORAL MUCOSITIS**

Study timeline.



(57) **Abbrégé/Abstract:**

Methods and kits for treating oral mucositis are disclosed. The treatment comprises administering to a patient in need thereof a Reactive Oxygen Species scavenger in a pharmaceutically acceptable formulation.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



PCT



(43) International Publication Date
17 April 2008 (17.04.2008)

(10) International Publication Number
WO 2008/045559 A3

(51) International Patent Classification:

A01N 55/02 (2006.01) A61K 31/573 (2006.01)
A61K 31/555 (2006.01)

(74) Agent: BLOSSER, Harley, G.; Sonnenschein Nath And
Rosenthal LLP, P.O. Box 061080, Wacker Drive Station,
Sears Tower, Chicago, IL 60606-1080 (US).

(21) International Application Number:

PCT/US2007/021897

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date: 12 October 2007 (12.10.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/829,291 12 October 2006 (12.10.2006) US
11/871,848 12 October 2007 (12.10.2007) US

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **ACTIVBIOTICS** [US/US]; 110 Hartwell Ave, Lexington, MA 02421 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **ROTHSTEIN, David, M.** [US/US]; 107 Cedar Street, Lexington, MA 02421 (US). **MURPHY, Chris** [US/US]; 7 Warren Street, Upton, MA 01568 (US). **WONG, Vivian** [US/US]; 423 Fort Hill Rd., Scarsdale, NY 10583 (US). **KAZO, Glen** [US/US]; 110 Hartwell Ave, Lexington, MA 02421 (US).


Published:

— with international search report

(88) Date of publication of the international search report:
31 July 2008

(54) Title: METHODS OF TREATING ORAL MUCOSITIS

Study timeline.



(57) Abstract: Methods and kits for treating oral mucositis are disclosed. The treatment comprises administering to a patient in need thereof a Reactive Oxygen Species scavenger in a pharmaceutically acceptable formulation.

WO 2008/045559 A3

75975-31

TITLE OF THE INVENTION**Superoxide Dismutase Mimetics for the Treatment of Oral Mucositis****[0001]****STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR
DEVELOPMENT****[0002]** Not Applicable.**INCORPORATION-BY-REFERENCE OF MATERIAL SUBMITTED ON A COMPACT
DISC****[0003]** Not Applicable.**BACKGROUND OF THE INVENTION****1. Field of the Invention**

[0004] This invention relates generally to the treatment of oral mucositis and, more particularly, to methods for treating oral mucositis with a Reactive Oxygen Species ("ROS") scavenger. The compositions and methods are useful in treating oral mucositis.

2. Description of Related Art

[0005] Oral ulcerative mucositis is a common, painful, dose-limiting toxicity of drug and radiation therapy for cancer (1). The disorder is characterized by breakdown of the oral mucosa that results in the formation of ulcerative lesions. In granulocytopenic patients, the ulcerations that accompany mucositis are frequent portals of entry for indigenous oral bacteria often leading to sepsis or bacteremia (2). Mucositis occurs to some degree in more than one third of patients receiving anti-neoplastic drug therapy (3). The frequency and severity are significantly greater among patients who are treated with induction therapy for leukemia or with many of

the conditioning regimens for bone marrow transplant (4). Among these individuals, moderate to severe mucositis can occur in more than three-quarters of patients. Moderate to severe mucositis occurs in virtually all patients who receive radiation therapy for tumors of the head and neck and typically begins with cumulative exposures of 15 Gy and then worsens as total doses of 60 Gy or more are reached (1- 4).

[0006] Clinically mucositis progresses through three stages:

1. Atrophic changes accompanied by painful mucosal erythema, which can respond to local anesthetics.
2. Painful ulceration with pseudomembrane formation and, in the case of myelosuppressive treatment, potentially life-threatening sepsis, requiring antimicrobial therapy. Pain is often of such intensity as to require parenteral narcotic analgesia.
3. Spontaneous healing, occurring about 2 - 3 weeks after cessation of anti-neoplastic therapy.

[0007] Standard therapy for mucositis is predominantly palliative, including application of topical analgesics such as lidocaine and/or systemic administration of narcotics and antibiotics. At present, the only approved treatment for oral mucositis is palifermin (Kepivance) which is a member of the fibroblast growth factor (FGF) superfamily of molecules. Palifermin's approval is limited to the treatment of oral mucositis in the patients undergoing conditioning regimens prior to hematopoietic stem cell transplants for the treatment of hematologic malignancies.


[0008] The complexity of mucositis as a biological process has only been recently appreciated (5-7). It has been suggested that the condition represents a sequential interaction of oral mucosal cells and tissues, reactive oxygen species, pro-inflammatory cytokines, mediators of apoptosis and local factors such as saliva and the oral microbiota. While epithelial degeneration and breakdown ultimately result in mucosal ulceration, it appears that the early changes associated with radiation-induced mucosal toxicity occur within the endothelium, and connective tissue of the submucosa. For example, electron microscopic evaluation of mucosa within 1 week of radiation shows damage to both endothelium and connective tissue, but not epithelium. It appears that the overall mechanism for mucositis development is similar for both radiation and chemotherapy (8).

[0009] Recently, a superoxide dismutase mimetic, M40403, was shown to be

effective in an animal models of inflammation (Salvemini *et al.*, Science 286:304-306, 1999) and, more specifically, in an animal model of rheumatoid arthritis (Salvemini *et al.*, Arthritis & Rheumatism 44:2909-2921, 2001). Nevertheless, treatment of oral mucositis using M40403 has neither been reported nor suggested.

BRIEF SUMMARY OF THE INVENTION

[0010] The present teachings provide methods for treating oral mucositis. A method comprises administering administering to a patient in need thereof, a superoxide dismutase mimetic. In accordance with one aspect of the invention, the superoxide dismutase mimetic can be represented by the formula:



[0011] wherein

[0012] (i) $R_1, R'_1, R_2, R'_2, R_3, R'_3, R_4, R'_4, R_5, R'_5, R_6, R'_6, R_7, R'_7, R_8, R'_8, R_9, R'_9, R_{10},$ and R'_{10} are independently:

[0013] (i^a) hydrogen; or

[0014] (i^b) a moiety independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, heterocyclyl, and aralkyl radicals and radicals attached to the α -carbon or α -amino acids; or

[0015] (i^c) a moiety independently selected from the group consisting of $-OR_{11}, -NR_{11}R_{12}, -COR_{11}, -CO_2R_{11}, -CONR_{11}R_{12}, -SR_{11}, -SOR_{11}, -SO_2R_{11}, -SO_2NR_{11}R_{12}, -N(OR_{11})(R_{12}), -P(O)(OR_{11})(OR_{12}), -P(O)(OR_{11})(R_{12}), -OP(O)(OR_{11})(OR_{12}),$ and substituents attached to the α -carbon of α -amino acids, wherein R_{11} and R_{12} are independently hydrogen or alkyl; and

[0016] (ii) optionally, one or more of R_1 or R'_1 and R_2 or R'_2, R_3 or R'_3 and R_4 or R'_4, R_5 or R'_5 and R_6 or R'_6, R_7 or R'_7 and R_8 or R'_8, R_9 or R'_9 and R_{10} or R'_{10} together

with the carbon atoms to which they are attached independently form a substituted or unsubstituted and saturated, partially saturated, or unsaturated cycle or heterocycle having 3 to 20 carbon atoms; and

[0017] (iii) optionally, one or more of R_1 and R'_1 , R_2 and R'_2 , R_3 and R'_3 , R_4 and R'_4 , R_5 and R'_5 , R_6 and R'_6 , R_7 and R'_7 , R_8 and R'_8 , R_9 and R'_9 , and R_{10} and R'_{10} , together with the carbon atom to which they are attached independently form a substituted or unsubstituted and saturated, partially saturated, or unsaturated cycle or heterocycle having 3 to 20 carbon atoms; and

[0018] (iv) optionally, one or more of R_{10} or R'_{10} and R_1 or R'_1 , R_2 or R'_2 and R_3 or R'_3 , R_4 or R'_4 and R_5 or R'_5 , R_6 or R'_6 and R_7 or R'_7 , or R_8 or R'_8 and R_9 or R'_9 together with the carbon atoms to which they are attached independently form a substituted or unsubstituted nitrogen containing heterocycle having 3 to 20 carbon atoms, which may be an aromatic heterocycle in which case the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and the R groups attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent; and

[0019] (v) optionally, one or more of R_1 , R'_1 , R_2 , R'_2 , R_3 , R'_3 , R_4 , R'_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R'_7 , R_8 , R'_8 , R_9 , R'_9 , R_{10} , and R'_{10} , together with a different one of R_1 , R'_1 , R_2 , R'_2 , R_3 , R'_3 , R_4 , R'_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R'_7 , R_8 , R'_8 , R_9 , R'_9 , R_{10} , and R'_{10} , which is attached to a different carbon atom in the macrocyclic ligand may be bound to form a strap represented by the formula:

[0020] $-(CH_2)_I-Q-(CH_2)_J-R-(CH_2)_K-S-(CH_2)_L-$

[0021] wherein

[0022] I, J, K and L independently are integers from 0 to 10 and Q, R and S are independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, and heterocyclyl, aza, amide, ammonium, oxa, thia, sulfonyl, sulfinyl, sulfonamide, phosphoryl, phosphinyl, phosphino, phosphonium, keto, ester, alcohol, carbamate, urea, thiocarbonyl, borates, boranes, boraza, silyl, siloxy, silaza, and combinations thereof; and

[0023] (vi) combinations of any of (i) through (v) above;

[0024] wherein

[0025] M is a transition metal;

[0026] X, Y and Z are independently selected from the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxy, hydroperoxy, alkylperoxy, arylperoxy, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof; or


[0027] X, Y and Z are independently selected from the group consisting of charge-neutralizing anions which are derived from any monodentate or polydentate coordinating ligand and a ligand system and the corresponding anion thereof; or

[0028] X, Y and Z are independently attached to one or more of R₁, R'₁, R₂, R'₂, R₃, R'₃, R₄, R'₄, R₅, R'₅, R₆, R'₆, R₇, R'₇, R₈, R'₈, R₉, R'₉, R₁₀, and R'₁₀; and

[0029] n is an integer from 0 to 3.

[0030] Preferably, M is selected from the group consisting of Mn²⁺, Mn³⁺, Mn⁴⁺, Mn⁶⁺, Mn⁷⁺, Fe²⁺, Fe³⁺, Fe⁴⁺, Fe⁶⁺, Ni²⁺, Ni³⁺, Cu¹⁺, Cu²⁺, V²⁺, V³⁺, V⁴⁺, and V⁵⁺.

[0031] In an alternative, the superoxide dismutase mimetic can be represented by the formula:



[0032] wherein

[0033] (i) a nitrogen of the macrocycle and two adjacent carbon atoms to which the nitrogen is attached independently form a substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing heterocycle W having 2 to 20 carbon atoms, which may be an aromatic heterocycle in which case the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and the R groups attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent; and

[0034] (ii) one or more of R_1 , R_2 , R'_2 , R_3 , R'_3 , R_4 , R'_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R'_7 , R_8 , R'_8 , R_9 , R'_9 , and R_{10} are independently:

[0035] (ii^a) hydrogen; or

[0036] (ii^b) a moiety independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, heterocyclyl, and aralkyl radicals and radicals attached to the α -carbon or α -amino acids; or

[0037] (ii^c) a moiety independently selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, $-OP(O)(OR_{11})(OR_{12})$, and substituents attached to the α -carbon of α -amino acids, wherein R_{11} and R_{12} are independently hydrogen or alkyl; and

[0038] (iii) optionally, one or more of R_1 and R_2 or R'_2 , R_3 or R'_3 and R_4 or R'_4 , R_5

or R'₅ and R₆ or R'₆, R₇ or R'₇ and R₈ or R'₈, R₉ or R'₉ and R₁₀ together with the carbon atoms to which they are attached independently form a substituted or unsubstituted and saturated, partially saturated, or unsaturated cycle or heterocycle having 3 to 20 carbon atoms; and

[0039] (iv) optionally, one or more of R₂ and R'₂, R₃ and R'₃, R₄ and R'₄, R₅ and R'₅, R₆ and R'₆, R₇ and R'₇, R₈ and R'₈, and R₉ and R'₉, together with the carbon atom to which they are attached independently form a substituted or unsubstituted and saturated, partially saturated, or unsaturated cycle or heterocycle having 3 to 20 carbon atoms; and

[0040] (v) optionally, one or more of R₂ or R'₂ and R₃ or R'₃, R₄ or R'₄ and R₅ or R'₅, R₆ or R'₆ and R₇ or R'₇, or R₈ or R'₈ and R₉ or R'₉ together with the carbon atoms to which they are attached independently form a substituted or unsubstituted nitrogen containing heterocycle having 3 to 20 carbon atoms, which may be an aromatic heterocycle in which case the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and the R groups attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent; and

[0041] (vi) optionally, one or more of R₁, R₂, R'₂, R₃, R'₃, R₄, R'₄, R₅, R'₅, R₆, R'₆, R₇, R'₇, R₈, R'₈, R₉, R'₉, and R₁₀, together with a different one of R₁, R₂, R'₂, R₃, R'₃, R₄, R'₄, R₅, R'₅, R₆, R'₆, R₇, R'₇, R₈, R'₈, R₉, R'₉, and R₁₀, which is attached to a different carbon atom in the macrocyclic ligand may be bound to form a strap represented by the formula:

[0042] $-(CH_2)_I-Q-(CH_2)_J-R-(CH_2)_K-S-(CH_2)_L-$

[0043] wherein

[0044] I, J, K and L independently are integers from 0 to 10 and Q, R and S are independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, and heterocyclyl, aza, amide, ammonium, oxa, thia, sulfonyl, sulfinyl, sulfonamide, phosphoryl, phosphinyl, phosphino, phosphonium, keto, ester, alcohol, carbamate, urea, thiocarbonyl, borates, boranes, boraza, silyl, siloxy, silaza, and combinations thereof; and

[0045] (vii) optionally, one or more of R₁, R₂, R'₂, R₃, R'₃, R₄, R'₄, R₅, R'₅, R₆, R'₆, R₇, R'₇, R₈, R'₈, R₉, R'₉, and R₁₀, may be bound to an atom of heterocycle W to form

a strap represented by the formula:

[0046] $-(CH_2)_I-Q-(CH_2)_J-R-(CH_2)_K-S-(CH_2)_L-$

[0047] wherein

[0048] I, J, K and L independently are integers from 0 to 10 and Q, R and S are independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, and heterocyclyl, aza, amide, ammonium, oxa, thia, sulfonyl, sulfinyl, sulfonamide, phosphoryl, phosphinyl, phosphino, phosphonium, keto, ester, alcohol, carbamate, urea, thiocarbonyl, borates, boranes, boraza, silyl, siloxy, silaza, and combinations thereof; and

[0049] (viii) combinations of any of (i) through (vii) above;

[0050] wherein

[0051] M is a transition metal;

[0052] X, Y and Z are independently selected from the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxo, hydroperoxo, alkylperoxo, arylperoxo, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate,

carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof; or


[0053] X, Y and Z are independently selected from the group consisting of charge-neutralizing anions which are derived from any monodentate or polydentate coordinating ligand and a ligand system and the corresponding anion thereof; or

[0054] X, Y and Z are independently attached to one or more of R_1 , R_2 , R'_2 , R_3 , R'_3 , R_4 , R'_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R'_7 , R_8 , R'_8 , R_9 , R'_9 , and R_{10} ; and

[0055] n is an integer from 0 to 3.

[0056] Preferably, M is selected from the group consisting of Mn^{2+} , Mn^{3+} , Mn^{4+} , Mn^{6+} , Mn^{7+} , Fe^{2+} , Fe^{3+} , Fe^{4+} , Fe^{6+} , Ni^{2+} , Ni^{3+} , Cu^{1+} , Cu^{2+} , V^{2+} , V^{3+} , V^{4+} , and V^{5+} , and W is a substituted or unsubstituted pyridino moiety.

[0057] In yet another alternative, the superoxide dismutase mimetic can be represented by the formula:



[0058] wherein

[0059] (i) a nitrogen of the macrocycle and two adjacent carbon atoms to which the nitrogen is attached independently form a substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing heterocycle W having 2 to 20 carbon atoms, which may be an aromatic heterocycle in which case the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and the R groups attached to the carbon atoms which are both part of the heterocycle

and the macrocycle are absent; and

[0060] (ii) two sets of two adjacent carbon atoms of the macrocycle independently form substituted or unsubstituted, saturated, partially saturated or unsaturated, cycles or heterocycles U and V having 3 to 20 carbon atoms; and

[0061] (iii) R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are independently:

[0062] (iii^a) hydrogen; or

[0063] (iii^b) a moiety independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, heterocyclyl, and aralkyl radicals and radicals attached to the α -carbon or α -amino acids; or

[0064] (iii^c) a moiety independently selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, $-OP(O)(OR_{11})(OR_{12})$, and substituents attached to the α -carbon of α -amino acids, wherein R_{11} and R_{12} are independently hydrogen or alkyl; and

[0065] (iv) optionally, one or more of R_1 and R_2 or R'_2 , R_5 or R'_5 and R_6 or R'_6 , R_9 or R'_9 and R_{10} together with the carbon atoms to which they are attached independently form a substituted or unsubstituted and saturated, partially saturated, or unsaturated cycle or heterocycle having 3 to 20 carbon atoms; and

[0066] (v) optionally, one or more of R_2 and R'_2 , R_5 and R'_5 , R_6 and R'_6 , and R_9 and R'_9 , together with the carbon atom to which they are attached independently form a substituted or unsubstituted and saturated, partially saturated, or unsaturated cycle or heterocycle having 3 to 20 carbon atoms; and

[0067] (vi) optionally, one or more of R_2 or R'_2 and R_3 , R_4 and R_5 or R'_5 , R_6 or R'_6 and R_7 , or R_8 and R_9 or R'_9 together with the carbon atoms to which they are attached independently form a substituted or unsubstituted nitrogen containing heterocycle having 3 to 20 carbon atoms, which may be an aromatic heterocycle in which case the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and the R groups attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent; and

[0068] (vii) optionally, one or more of R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} , together with a different one of R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 ,

R₈, R₉, R'₉, and R₁₀, which is attached to a different carbon atom in the macrocyclic ligand may be bound to form a strap represented by the formula:

[0069] $-(CH_2)_I-Q-(CH_2)_J-R-(CH_2)_K-S-(CH_2)_L-$

[0070] wherein

[0071] I, J, K and L independently are integers from 0 to 10 and Q, R and S are independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, and heterocyclyl, aza, amide, ammonium, oxa, thia, sulfonyl, sulfinyl, sulfonamide, phosphoryl, phosphinyl, phosphino, phosphonium, keto, ester, alcohol, carbamate, urea, thiocarbonyl, borates, boranes, boraza, silyl, siloxy, silaza, and combinations thereof; and

[0072] (viii) optionally, one or more of R₁, R₂, R'₂, R₃, R₄, R₅, R'₅, R₆, R'₆, R₇, R₈, R₉, R'₉, and R₁₀, may be individually bound to an atom of heterocycles U, V and W to form a strap represented by the formula:

[0073] $-(CH_2)_I-Q-(CH_2)_J-R-(CH_2)_K-S-(CH_2)_L-$

[0074] wherein

[0075] I, J, K and L independently are integers from 0 to 10 and Q, R and S are independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, and heterocyclyl, aza, amide, ammonium, oxa, thia, sulfonyl, sulfinyl, sulfonamide, phosphoryl, phosphinyl, phosphino, phosphonium, keto, ester, alcohol, carbamate, urea, thiocarbonyl, borates, boranes, boraza, silyl, siloxy, silaza, and combinations thereof; and

[0076] (ix) combinations of any of (i) through (viii) above;

[0077] wherein

[0078] M is a transition metal;

[0079] X, Y and Z are independently selected from the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxo, hydroperoxo, alkylperoxo, arylperoxo, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid,

alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof; or


[0080] X, Y and Z are independently selected from the group consisting of charge-neutralizing anions which are derived from any monodentate or polydentate coordinating ligand and a ligand system and the corresponding anion thereof; or

[0081] X, Y and Z are independently attached to one or more of R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} ; and

[0082] n is an integer from 0 to 3.

[0083] Preferably, M is selected from the group consisting of Mn^{2+} , Mn^{3+} , Mn^{4+} , Mn^{6+} , Mn^{7+} , Fe^{2+} , Fe^{3+} , Fe^{4+} , Fe^{6+} , Ni^{2+} , Ni^{3+} , Cu^{1+} , Cu^{2+} , V^{2+} , V^{3+} , V^{4+} , and V^{5+} . In accordance with a further aspect of the invention, U and V are saturated cycloalkyl heterocycles having 3 to 20 carbon atoms preferably saturated cycloalkyl heterocycles having 4 to 10 carbon atoms, and still more preferably U and V are trans-cyclohexanyl fused rings. In yet another aspect of the present invention, W is a substituted or unsubstituted pyridino moiety, more preferably, U and V are trans-cyclohexanyl fused rings and W is a substituted pyridino moiety. Preferably, the

superoxide dismutase mimetic can be represented by the formula:



[0084] The ROS scavenger can be administered in an amount of at most 0.015 mg/kg, or preferably at most 2 mg/kg. In yet another aspect, the pharmaceutically acceptable formulation is a pharmaceutically acceptable oral formulation and administering comprises administering orally. Preferably, the patient is a human patient and the oral mucositis is a result of chemotherapy or radiation therapy.


[0085] Another aspect of the invention can be a method of treating a cancer, the method comprising: a) administering to a subject in need of cancer treatment a pharmaceutical composition comprising a superoxide dismutase mimetic; and b) administering to the subject an effective amount of a cancer treatment, whereby the superoxide dismutase mimetic prevents or reduces oral mucositis in the subject. The cancer treatment can be comprised of radiation therapy and chemotherapy. In said method of treating a cancer, the superoxide dismutase mimetic can be a reactive oxygen species scavenger, and the pharmaceutical composition can further comprise at least one additional reactive oxygen species scavenger selected from the group consisting of amifostine and N-acetylcysteine. Additionally, the method of treating a cancer can further comprise administering a pharmaceutical composition which upregulates expression of at least one transcription factor which increases expression of one or more genes controlling at least one naturally occurring antioxidant pathway. The composition which upregulates expression of at least one transcription factor can be palifermin, and the transcription factor upregulated can be Nrf-2.

[0086] In yet another aspect of the present invention, a kit can be provided for treating oral mucositis, the kit comprising: (a) a ROS scavenger, wherein the ROS scavenger can be a superoxide dismutase; (b) one additional pharmaceutical compound selected from the group consisting of a chemotherapeutic agent; (c) a non-superoxide dismutase mimetic radical scavenger; and (d) instructions for

75975-31

administering the superoxide dismutase to a subject in need of cancer therapy. A chemotherapeutic agent can be selected from a group consisting of all-trans retinoic acid, azacitidine, azathioprine, bleomycin, carboplatin, capecitabine, cisplatin, chlorambucil, cyclophosphamide, cytarabine, daunorubicin, docetaxel, doxifluridine, doxorubicin, epirubicin, epothilone, etoposide, fluorouracil, gemcitabine, hydroxyurea, idarubicin, imatinib, mechlorethamine, mercaptopurine, methotrexate, mitoxantrone, oxaliplatin, paclitaxel, pemetrexed, teniposide, tiguanine, valrubicin, vinblastine, vincristine, vindesine, vinorelbine. A non-superoxide dismutase mimetic radical scavenger can be selected from the group of amifostine and N-acetylcysteine. A superoxide dismutase mimetic can be M40403.

[0086a] In one aspect, the present invention relates to use of a superoxide dismutase mimetic to lessen severity of an oral mucositis in a mammal, wherein the superoxide dismutase mimetic is for parenteral administration prior to or simultaneous with a cancer treatment, and the superoxide dismutase mimetic is a compound of the formula:



wherein

W is a substituted or unsubstituted pyridino moiety;

U and V are trans-cyclohexanyl fused rings;

75975-31

$R_1, R_2, R'_2, R_3, R_4, R_5, R'_5, R_6, R'_6, R_7, R_8, R_9, R'_9$, and R_{10} are each independently selected from the group consisting of hydrogen, alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, heterocyclyl, and aralkyl;

M is a cation of a transition metal;


and X, Y and Z are independently selected from the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxy, hydroperoxy, alkylperoxy, arylperoxy, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid,

75975-31

hydroxamic acid, and thiotosylate;

and n is an integer from 0 to 3.

[0086b] In one aspect, the present invention relates to use of a superoxide
dismutase mimetic to lessen severity of an oral mucositis in a mammal, wherein the
5 superoxide dismutase mimetic is for parenteral administration prior to or
simultaneous with a radiation therapy treatment, and the superoxide dismutase
mimetic is a compound of the formula:



wherein

10 W is a substituted or unsubstituted pyridino moiety;

U and V are trans-cyclohexanyl fused rings;

R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are each
independently selected from the group consisting of hydrogen, alkenyl,
alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl,
15 aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl,
cycloalkenylalkyl, heterocyclyl, and aralkyl;

M is a cation of a transition metal;

75975-31


and X, Y and Z are independently selected from the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxo, hydroperoxo, alkylperoxo, arylperoxo, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric
 5 oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl
 10 carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl
 15 phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl
 20 dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid,
 25 hydroxamic acid, and thiotosylate;

and n is an integer from 0 to 3.

[0086c] In one aspect, the present invention relates to use of a superoxide dismutase mimetic to lessen severity of an oral mucositis in a mammal, wherein the superoxide dismutase mimetic is for parenteral administration prior to or
 30 simultaneous with a chemotherapy treatment, and the superoxide dismutase mimetic

75975-31

is a compound of the formula:



wherein

W is a substituted or unsubstituted pyridino moiety;

5 U and V are trans-cyclohexanyl fused rings;

R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are each independently selected from the group consisting of hydrogen, alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, heterocyclyl, and aralkyl;

M is a cation of a transition metal;

and X, Y and Z are independently selected from the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxo, hydroperoxo, alkylperoxo, arylperoxo, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl

75975-31

- sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate,
- 5 bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite,
- 10 triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite,
- 15 hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, and thiotosylate;

and n is an integer from 0 to 3.

75975-31

[0087] These and other features, aspects and advantages of the present invention will become better understood with reference to the following description, examples and appended claims.

DETAILED DESCRIPTION OF THE INVENTION

[0088] Abbreviations and Definitions

[0089] To facilitate understanding of the invention, a number of terms and abbreviations as used herein are defined below as follows:

[0090] The term "alkenyl", alone or in combination, means an alkyl substituent having one or more double bonds. Examples of such alkenyl substituents include, but are not limited to, ethenyl, propenyl, 1-butenyl, cis-2-butenyl, trans-2-butenyl, isobutylenyl, cis-2-pentenyl, trans-2-pentenyl, 3-methyl-1-butenyl, 2,3-dimethyl-2-butenyl, 1-pentenyl, 1-hexenyl, 1-octenyl, decenyl, dodecenyl, tetradecenyl, hexadecenyl, cis- and trans-9-octadecenyl, 1,3-pentadienyl, 2,4-pentadienyl, 2,3-pentadienyl, 1,3-hexadienyl, 2,4-hexadienyl, 5,8,11,14-eicosatetraenyl, and 9,12,15-octadecatrienyl.

[0091] The term "alkyl", alone or in combination, means a straight-chain or branched-chain alkyl substituent containing from 1 to about 22 carbon atoms, preferably from about 1 to about 18 carbon atoms, and most preferably from about 1 to about 12 carbon atoms. Examples of such substituents include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isoamyl, hexyl, octyl, nonyl, decyl, dodecyl, tetradecyl, hexadecyl, octadecyl and eicosyl.

[0092] The terms "alkylcycloalkyl" and "alkenylcycloalkyl" mean a cycloalkyl

substituent as defined above which is substituted by an alkyl or alkenyl substituent as defined above. Examples of alkylcycloalkyl and alkenylcycloalkyl substituents include, but are not limited to, 2-ethylcyclobutyl, 1-methylcyclopentyl, 1-hexylcyclopentyl, 1-methylcyclohexyl, 1-(9-octadecenyl)cyclopentyl and 1-(9-octadecenyl)cyclohexyl.

[0093] The terms "alkylcycloalkenyl" and "alkenylcycloalkenyl" means a cycloalkenyl substituent as defined above which is substituted by an alkyl or alkenyl substituent as defined above. Examples of alkylcycloalkenyl and alkenylcycloalkenyl substituents include, but are not limited to, 1-methyl-2-cyclopentyl, 1-hexyl-2-cyclopentenyl, 1-ethyl-2-cyclohexenyl, 1-butyl-2-cyclohexenyl, 1-(9-octadecenyl)-2-cyclohexenyl and 1-(2-pentenyl)-2-cyclohexenyl.

[0094] The term "alkynyl", alone or in combination, means an alkyl substituent having one or more triple bonds. Examples of such alkynyl groups include, but are not limited to, ethynyl, propynyl (propargyl), 1-butylnyl, 1-octynyl, 9-octadecynyl, 1,3-pentadiynyl, 2,4-pentadiynyl, 1,3-hexadiynyl, and 2,4-hexadiynyl.

[0095] The term "aralkyl", alone or in combination, means an alkyl or cycloalkyl substituent as defined above in which one hydrogen atom is replaced by an aryl substituent as defined above, such as benzyl, 2-phenylethyl, and the like.

[0096] The term "aryl", alone or in combination, means a phenyl or naphthyl substituent which optionally carries one or more substituents selected from alkyl, cycloalkyl, cycloalkenyl, aryl, heterocycle, alkoxyaryl, alkaryl, alkoxy, halogen, hydroxy, amine, cyano, nitro, alkylthio, phenoxy, ether, trifluoromethyl and the like, such as phenyl, p-tolyl, 4-methoxyphenyl, 4-(tert-butoxy)phenyl, 4-fluorophenyl, 4-chlorophenyl, 4-hydroxyphenyl, 1-naphthyl, 2-naphthyl, and the like.

[0097] The term "cycloalkenyl", alone or in combination, means a cycloalkyl substituent having one or more double bonds. Examples of cycloalkenyl substituents include, but are not limited to, cyclopentenyl, cyclohexenyl, cyclooctenyl, cyclopentadienyl, cyclohexadienyl and cyclooctadienyl.

[0098] The terms "cyclic", "cycle" or "cyclyl" means a ring structure containing 3 to 20 carbon atoms, preferably 5 to 10 carbon atoms, which may be heterocyclic. The cyclic, cycle or cyclyl can also contain more than one ring.

[0099] The term "cycloalkenylalkyl" means an alkyl substituent as defined above which is substituted by a cycloalkenyl substituent as defined above. Examples of cycloalkenylalkyl substituents include, but are not limited to, 2-cyclohexen-1-

ylmethyl, 1-cyclopenten-1-ylmethyl, 2-(1-cyclohexen-1-yl)ethyl, 3-(1-cyclopenten-1-yl)propyl, 1-(1-cyclohexen-1-ylmethyl)pentyl, 1-(1-cyclopenten-1-yl)hexyl, 6-(1-cyclohexen-1-yl)hexyl, 1-(1-cyclopenten-1-yl)nonyl and 1-(1-cyclohexen-1-yl)nonyl.

[0100] The term "cycloalkyl", alone or in combination means a cycloalkyl radical containing from 3 to about 10, preferably from 3 to about 8, and most preferably from 3 to about 6, carbon atoms. Examples of such cycloalkyl substituents include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, and perhydronaphthyl.

[0101] The term "cycloalkylalkyl" means an alkyl substituent as defined above which is substituted by a cycloalkyl substituent as defined above. Examples of cycloalkylalkyl substituents include, but are not limited to, cyclohexylmethyl, cyclopentylmethyl, (4-isopropylcyclohexyl)methyl, (4-*t*-butyl-cyclohexyl)methyl, 3-cyclohexylpropyl, 2-cyclohexylmethylpentyl, 3-cyclopentylmethylhexyl, 1-(4-neopentylcyclohexyl)methylhexyl, and 1-(4-isopropylcyclohexyl)methylheptyl.

[0102] The term "cycloalkylcycloalkyl" means a cycloalkyl substituent as defined above which is substituted by another cycloalkyl substituent as defined above. Examples of cycloalkylcycloalkyl substituents include, but are not limited to, cyclohexylcyclopentyl and cyclohexylcyclohexyl.

[0103] The term "halide" means chloride, fluoride, iodide, or bromide.

[0104] The term "heterocyclic", "heterocycle" or "heterocyclyl" means a cyclic, cycle or cyclyl containing at least one other kind of atom, in addition to carbon, in the ring. Such atoms include, but are not limited to, nitrogen, oxygen and sulfur. The heterocyclic can also contain more than one ring. Examples of heterocyclics include, but are not limited to, pyrrolidinyl, piperidyl, imidazolidinyl, tetrahydrofuryl, tetrahydrothienyl, furyl, thienyl, pyridyl, quinolyl, isoquinolyl, pyridazinyl, pyrazinyl, indolyl, imidazolyl, oxazolyl, thiazolyl, pyrazolyl, pyridinyl, benzoxadiazolyl, benzothiadiazolyl, triazolyl and tetrazolyl groups.

[0105] The term "nitrogen containing heterocycle" means a ring structure in which 2 carbons and a nitrogen of the ring are shared with the fifteen-membered macrocyclic ligand. The nitrogen containing heterocycle can contain 2 to 20, preferably 4 to 10, carbon atoms, can be substituted or unsubstituted, saturated, partially saturated or unsaturated, and can also contain nitrogen, oxygen and/or sulfur atoms in the portion of the ring which is not also part of the fifteen-membered macrocyclic ligand.

[0106] The term "oral mucositis" shall also include stomatitis, small intestine-titis,

large intestine-titis, proctitis, and similar conditions affecting the mucosal lining of the entire gastrointestinal tract, and related conditions.

[0107] The term "R groups" means the group of variable substituents designated as "R" attached to the carbon atoms of the macrocycle, *i.e.*, R₁, R'₁, R₂, R'₂, R₃, R'₃, R₄, R'₄, R₅, R'₅, R₆, R'₆, R₇, R'₇, R₈, R'₈, R₉, R'₉, R₁₀, and R'₁₀.

[0108] The term "saturated, partially saturated or unsaturated cycle or heterocycle" means a fused ring structure in which 2 carbons of the ring are also part of the fifteen-membered macrocyclic ligand in which the ring can contain no double bonds (in the case of a saturated ring structure) or at least one double bond, which may be conjugated or unconjugated with another double bond. The ring structure can contain 3 to 20 carbon atoms, preferably 5 to 10 carbon atoms, which may be heterocyclic. The cyclic can also contain more than one ring.

[0109] In addition, the following abbreviations have the following meanings:


AUC	Area under the curve
bid	Twice daily
FGF	Fibroblast growth factor
g, mg, ml, kg	Gram, milligram, milliliter, kilogram
Gy	Gray
ip	Intraperitoneal
ma	Milliamp
mM	Millimolar
mm, cm	Millimeter, centimeter
SEM	Standard error of the mean
top	Topical

[0110] **Methods for Treating Oral Mucositis**

75975-31

[0111] The present invention involves the administration of a ROS scavenger in the treatment of oral mucositis. A ROS scavenger of the present invention is a superoxide dismutase mimetic, a non-proteinaceous molecule that catalyzes the conversion the of the superoxide radical, $O_2^{\cdot-}$, to molecular oxygen and hydrogen peroxide. Such a molecule can, like a native superoxide dismutase enzyme, reduce cell injury resulting from superoxide radical found in diseases involving oxidative stress such as inflammation (Salvemini *et al*, Arthritis & Rheumatism 44:2909-2921, 2001). The ROS scavengers of the present invention can be pentaaza-macrocyclic complexes, and more specifically, those compositions as disclosed in U.S. Patent Nos. 5,610,293, 5,637,578, 5,874,421, 5,976,498, 6,084,093, 6,180,620, 6,204,259, 6,214,817, 6,245,758, 6,395,725, and 6,525,041.

[0112] The superoxide dismutase mimetics of the present invention can be represented by the following formula:



[0113] wherein

[0114] (i) $R_1, R'_1, R_2, R'_2, R_3, R'_3, R_4, R'_4, R_5, R'_5, R_6, R'_6, R_7, R'_7, R_8, R'_8, R_9, R'_9, R_{10},$ and R'_{10} are independently:

[0115] (i^a) hydrogen; or

[0116] (i^b) a moiety independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, and aralkyl radicals and radicals attached to the α -carbon or α -amino acidsheterocyclyl; or

[0117] (i^c) a moiety independently selected from the group consisting of $-OR_{11}, -NR_{11}R_{12}, -COR_{11}, -CO_2R_{11}, -CONR_{11}R_{12}, -SR_{11}, -SOR_{11}, -SO_2R_{11}, -SO_2NR_{11}R_{12}, -N(OR_{11})(R_{12}), -P(O)(OR_{11})(OR_{12}), -P(O)(OR_{11})(R_{12}), -OP(O)(OR_{11})(OR_{12}),$ and

substituents attached to the α -carbon of α -amino acids, wherein R_{11} and R_{12} are independently hydrogen or alkyl; and

[0118] (ii) optionally, one or more of R_1 or R'_1 and R_2 or R'_2 , R_3 or R'_3 and R_4 or R'_4 , R_5 or R'_5 and R_6 or R'_6 , R_7 or R'_7 and R_8 or R'_8 , R_9 or R'_9 and R_{10} or R'_{10} together with the carbon atoms to which they are attached independently form a substituted or unsubstituted and saturated, partially saturated, or unsaturated cycle or heterocycle having 3 to 20 carbon atoms; and

[0119] (iii) optionally, one or more of R_1 and R'_1 , R_2 and R'_2 , R_3 and R'_3 , R_4 and R'_4 , R_5 and R'_5 , R_6 and R'_6 , R_7 and R'_7 , R_8 and R'_8 , R_9 and R'_9 , and R_{10} and R'_{10} , together with the carbon atom to which they are attached independently form a substituted or unsubstituted and saturated, partially saturated, or unsaturated cycle or heterocycle having 3 to 20 carbon atoms; and

[0120] (iv) optionally, one or more of R_{10} or R'_{10} and R_1 or R'_1 , R_2 or R'_2 and R_3 or R'_3 , R_4 or R'_4 and R_5 or R'_5 , R_6 or R'_6 and R_7 or R'_7 , or R_8 or R'_8 and R_9 or R'_9 together with the carbon atoms to which they are attached independently form a substituted or unsubstituted nitrogen containing heterocycle having 3 to 20 carbon atoms, which may be an aromatic heterocycle in which case the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and the R groups attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent; and

[0121] (v) optionally, one or more of R_1 , R'_1 , R_2 , R'_2 , R_3 , R'_3 , R_4 , R'_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R'_7 , R_8 , R'_8 , R_9 , R'_9 , R_{10} , and R'_{10} , together with a different one of R_1 , R'_1 , R_2 , R'_2 , R_3 , R'_3 , R_4 , R'_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R'_7 , R_8 , R'_8 , R_9 , R'_9 , R_{10} , and R'_{10} , which is attached to a different carbon atom in the macrocyclic ligand may be bound to form a strap represented by the formula:

[0122] $-(CH_2)_I-Q-(CH_2)_J-R-(CH_2)_K-S-(CH_2)_L-$

[0123] wherein

[0124] I, J, K and L independently are integers from 0 to 10 and Q, R and S are independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, and heterocyclyl, aza, amide, ammonium, oxa, thia, sulfonyl, sulfinyl, sulfonamide, phosphoryl, phosphinyl, phosphino, phosphonium, keto, ester, alcohol, carbamate, urea, thiocarbonyl, borates, boranes, boraza, silyl, siloxy, silaza, and combinations

thereof; and

[0125] (vi) combinations of any of (i) through (v) above.


[0126] Thus, the pentaaza-macrocyclic ligand compositions useful in the present invention can have any combinations of substituted or unsubstituted R groups, saturated, partially saturated or unsaturated cyclics, heterocyclics, nitrogen containing heterocycles, or straps as defined above.

[0127] M can be a transition metal, preferably Mn^{2+} , Mn^{3+} , Mn^{4+} , Mn^{6+} , Mn^{7+} , Fe^{2+} , Fe^{3+} , Fe^{4+} , Fe^{6+} , Ni^{2+} , Ni^{3+} , Cu^{1+} , Cu^{2+} , V^{2+} , V^{3+} , V^{4+} , or V^{5+} . X, Y and Z can independently be selected from the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxo, hydroperoxo, alkylperoxo, arylperoxo, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof; or

[0128] X, Y and Z are independently selected from the group consisting of charge-neutralizing anions which are derived from any monodentate or polydentate coordinating ligand and a ligand system and the corresponding anion thereof; or

[0129] X, Y and Z are independently attached to one or more of R_1 , R'_1 , R_2 , R'_2 , R_3 , R'_3 , R_4 , R'_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R'_7 , R_8 , R'_8 , R_9 , R'_9 , R_{10} , and R'_{10} . n is preferably an integer from 0 to 3.

[0130] Alternatively, the superoxide dismutase mimetic can be represented by the formula:



[0131] wherein

[0132] (i) a nitrogen of the macrocycle and two adjacent carbon atoms to which the nitrogen is attached independently form a substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing heterocycle W having 2 to 20 carbon atoms, which may be an aromatic heterocycle in which case the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and the R groups attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent; and

[0133] (ii) one or more of R_1 , R_2 , R'_2 , R_3 , R'_3 , R_4 , R'_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R'_7 , R_8 , R'_8 , R_9 , R'_9 , and R_{10} are independently:

[0134] (ii^a) hydrogen; or

[0135] (ii^b) a moiety independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, heterocyclyl, and aralkyl radicals and radicals attached to the α -carbon or α -amino acids; or

[0136] (ii^c) a moiety independently selected from the group consisting of -OR₁₁, -NR₁₁R₁₂, -COR₁₁, -CO₂R₁₁, -CONR₁₁R₁₂, -SR₁₁, -SOR₁₁, -SO₂R₁₁, -SO₂NR₁₁R₁₂, -N(OR₁₁)(R₁₂), -P(O)(OR₁₁)(OR₁₂), -P(O)(OR₁₁)(R₁₂), -OP(O)(OR₁₁)(OR₁₂), and substituents attached to the α-carbon of α-amino acids, wherein R₁₁ and R₁₂ are independently hydrogen or alkyl; and

[0137] (iii) optionally, one or more of R₁ and R₂ or R'₂, R₃ or R'₃ and R₄ or R'₄, R₅ or R'₅ and R₆ or R'₆, R₇ or R'₇ and R₈ or R'₈, R₉ or R'₉ and R₁₀ together with the carbon atoms to which they are attached independently form a substituted or unsubstituted and saturated, partially saturated, or unsaturated cycle or heterocycle having 3 to 20 carbon atoms; and

[0138] (iv) optionally, one or more of R₂ and R'₂, R₃ and R'₃, R₄ and R'₄, R₅ and R'₅, R₆ and R'₆, R₇ and R'₇, R₈ and R'₈, and R₉ and R'₉, together with the carbon atom to which they are attached independently form a substituted or unsubstituted and saturated, partially saturated, or unsaturated cycle or heterocycle having 3 to 20 carbon atoms; and

[0139] (v) optionally, one or more of R₂ or R'₂ and R₃ or R'₃, R₄ or R'₄ and R₅ or R'₅, R₆ or R'₆ and R₇ or R'₇, or R₈ or R'₈ and R₉ or R'₉ together with the carbon atoms to which they are attached independently form a substituted or unsubstituted nitrogen containing heterocycle having 3 to 20 carbon atoms, which may be an aromatic heterocycle in which case the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and the R groups attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent; and

[0140] (vi) optionally, one or more of R₁, R₂, R'₂, R₃, R'₃, R₄, R'₄, R₅, R'₅, R₆, R'₆, R₇, R'₇, R₈, R'₈, R₉, R'₉, and R₁₀, together with a different one of R₁, R₂, R'₂, R₃, R'₃, R₄, R'₄, R₅, R'₅, R₆, R'₆, R₇, R'₇, R₈, R'₈, R₉, R'₉, and R₁₀, which is attached to a different carbon atom in the macrocyclic ligand may be bound to form a strap represented by the formula:

[0141] $-(\text{CH}_2)_I - \text{Q} - (\text{CH}_2)_J - \text{R} - (\text{CH}_2)_K - \text{S} - (\text{CH}_2)_L -$

[0142] wherein

[0143] I, J, K and L independently are integers from 0 to 10 and Q, R and S are independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, and

heterocyclyl, aza, amide, ammonium, oxa, thia, sulfonyl, sulfinyl, sulfonamide, phosphoryl, phosphinyl, phosphino, phosphonium, keto, ester, alcohol, carbamate, urea, thiocarbonyl, borates, boranes, boraza, silyl, siloxy, silaza, and combinations thereof; and

[0144] (vii) optionally, one or more of R_1 , R_2 , R'_2 , R_3 , R'_3 , R_4 , R'_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R'_7 , R_8 , R'_8 , R_9 , R'_9 , and R_{10} , may be bound to an atom of heterocycle W to form a strap represented by the formula:

[0145] $-(CH_2)_I-Q-(CH_2)_J-R-(CH_2)_K-S-(CH_2)_L-$

[0146] wherein

[0147] I, J, K and L independently are integers from 0 to 10 and Q, R and S are independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, and heterocyclyl, aza, amide, ammonium, oxa, thia, sulfonyl, sulfinyl, sulfonamide, phosphoryl, phosphinyl, phosphino, phosphonium, keto, ester, alcohol, carbamate, urea, thiocarbonyl, borates, boranes, boraza, silyl, siloxy, silaza, and combinations thereof; and


[0148] (viii) combinations of any of (i) through (vii) above.

[0149] Thus, the pentaaza-macrocyclic ligand compositions useful in the present invention can have any combinations of substituted or unsubstituted R groups, saturated, partially saturated or unsaturated cyclics, heterocyclics, nitrogen containing heterocycles, or straps as defined above, which may or may not independently connect the W loop and the pentaaza macrocycle.

[0150] M can be a transition metal, preferably Mn^{2+} , Mn^{3+} , Mn^{4+} , Mn^{6+} , Mn^{7+} , Fe^{2+} , Fe^{3+} , Fe^{4+} , Fe^{6+} , Ni^{2+} , Ni^{3+} , Cu^{1+} , Cu^{2+} , V^{2+} , V^{3+} , V^{4+} , or V^{5+} . X, Y and Z can independently be selected from the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxo, hydroperoxo, alkylperoxo, arylperoxo, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic

acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof. n is preferably an integer from 0 to 3. W can be a substituted or unsubstituted pyridino moiety.

[0151] In another alternative, the superoxide dismutase mimetic can be represented by the formula:



[0152] wherein

[0153] (i) a nitrogen of the macrocycle and two adjacent carbon atoms to which the nitrogen is attached independently form a substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing heterocycle W having 2 to 20

carbon atoms, which may be an aromatic heterocycle in which case the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and the R groups attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent; and

[0154] (ii) two sets of two adjacent carbon atoms of the macrocycle independently form substituted or unsubstituted, saturated, partially saturated or unsaturated, cycles or heterocycles U and V having 3 to 20 carbon atoms; and

[0155] (iii) R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are independently:

[0156] (iii^a) hydrogen; or

[0157] (iii^b) a moiety independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, heterocyclyl, and aralkyl radicals and radicals attached to the α -carbon or α -amino acids; or

[0158] (iii^c) a moiety independently selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, $-OP(O)(OR_{11})(OR_{12})$, and substituents attached to the α -carbon of α -amino acids, wherein R_{11} and R_{12} are independently hydrogen or alkyl; and

[0159] (iv) optionally, one or more of R_1 and R_2 or R'_2 , R_5 or R'_5 and R_6 or R'_6 , R_9 or R'_9 and R_{10} together with the carbon atoms to which they are attached independently form a substituted or unsubstituted and saturated, partially saturated, or unsaturated cycle or heterocycle having 3 to 20 carbon atoms; and

[0160] (v) optionally, one or more of R_2 and R'_2 , R_5 and R'_5 , R_6 and R'_6 , and R_9 and R'_9 , together with the carbon atom to which they are attached independently form a substituted or unsubstituted and saturated, partially saturated, or unsaturated cycle or heterocycle having 3 to 20 carbon atoms; and

[0161] (vi) optionally, one or more of R_2 or R'_2 and R_3 , R_4 and R_5 or R'_5 , R_6 or R'_6 and R_7 , or R_8 and R_9 or R'_9 together with the carbon atoms to which they are attached independently form a substituted or unsubstituted nitrogen containing heterocycle having 3 to 20 carbon atoms, which may be an aromatic heterocycle in which case the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and the R groups attached to the carbon atoms

which are both part of the heterocycle and the macrocycle are absent; and

[0162] (vii) optionally, one or more of R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} , together with a different one of R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} , which is attached to a different carbon atom in the macrocyclic ligand may be bound to form a strap represented by the formula:

[0163] $-(CH_2)_I-Q-(CH_2)_J-R-(CH_2)_K-S-(CH_2)_L-$

[0164] wherein

[0165] I, J, K and L independently are integers from 0 to 10 and Q, R and S are independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, and heterocyclyl, aza, amide, ammonium, oxa, thia, sulfonyl, sulfinyl, sulfonamide, phosphoryl, phosphinyl, phosphino, phosphonium, keto, ester, alcohol, carbamate, urea, thiocarbonyl, borates, boranes, boraza, silyl, siloxy, silaza, and combinations thereof; and

[0166] (viii) optionally, one or more of R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} , may be individually bound to an atom of heterocycles U, V and W to form a strap represented by the formula:

[0167] $-(CH_2)_I-Q-(CH_2)_J-R-(CH_2)_K-S-(CH_2)_L-$

[0168] wherein


[0169] I, J, K and L independently are integers from 0 to 10 and Q, R and S are independently selected from the group consisting of alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, and heterocyclyl, aza, amide, ammonium, oxa, thia, sulfonyl, sulfinyl, sulfonamide, phosphoryl, phosphinyl, phosphino, phosphonium, keto, ester, alcohol, carbamate, urea, thiocarbonyl, borates, boranes, boraza, silyl, siloxy, silaza, and combinations thereof; and

[0170] (ix) combinations of any of (i) through (viii) above;

[0171] Thus, the pentaaza-macrocyclic ligand compositions useful in the present invention can have any combinations of substituted or unsubstituted R groups, saturated, partially saturated or unsaturated cyclics, heterocyclics, nitrogen containing heterocycles, or straps as defined above, which may or may not independently connect the W, U or V loops and the pentaaza macrocycle.

[0172] M can be a transition metal, preferably Mn^{2+} , Mn^{3+} , Mn^{4+} , Mn^{6+} , Mn^{7+} , Fe^{2+} , Fe^{3+} , Fe^{4+} , Fe^{6+} , Ni^{2+} , Ni^{3+} , Cu^{1+} , Cu^{2+} , V^{2+} , V^{3+} , V^{4+} , or V^{5+} . X, Y and Z can independently be selected from the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxo, hydroperoxo, alkylperoxo, arylperoxo, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof. n is preferably an integer from 0 to 3. W can be a substituted or unsubstituted pyridino moiety. U and V can be independently saturated cycloalkyl heterocycles having 3 to 20 carbon atoms, more preferably 4 to 10 carbon atoms, still more preferably trans-cyclohexanyl fused rings. U and V can be trans-cyclohexanyl fused rings while W is a substituted pyridino moiety.

[0173] In certain embodiments, the superoxide dismutase mimetic can be the compound identified as M40403, which can be represented by the formula:



[0174] The present invention can involve administration of a ROS scavenger, in particular, M40403 or a suitable derivative or analog thereof as described above, to a patient in need thereof. Administration of the ROS scavenger, in particular M40403, can be by numerous routes of administration well known to those of skill in the art.

[0175] Administration of the ROS scavenger can be by any suitable route of administration such as, for example, oral, buccal, sublingual, intranasal, inhalation, rectal, intravaginal, transdermal, intradermal, subcutaneous, intramuscular, intraperitoneal, intravenous, intraarterial, intrasternal, intrathecal and the like.

[0176] Pharmaceutically acceptable formulations for parenteral or nonparenteral drug delivery are known in the art such as, for example, are set forth in Remington's Pharmaceutical Sciences, 18th Edition, Mack Publishing (1990). For pharmaceutical compositions and methods of treatment disclosed herein, dosage forms and administration regimes can be determined using standard methods known to skilled artisans, for example as set forth in standard references such as Remington: the Science and Practice of Pharmacy (Alfonso R. Gennaro ed. 19th ed. 1995); Hardman, J.G., et al., Goodman & Gilman's The Pharmacological Basis of Therapeutics, Ninth Edition, McGraw-Hill, 1996; and Rowe, R.C., et al., Handbook of Pharmaceutical Excipients, Fourth Edition, Pharmaceutical Press, 2003.

[0177] Pharmaceutical compositions can be formulated to be compatible with the intended route of administration. Solutions or suspensions used for parenteral, intradermal or subcutaneous application can include: a sterile diluent, such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents, such as benzyl alcohol or methyl parabens; antioxidants, such as ascorbic acid or sodium bisulfite; chelating agents, such as ethylenediaminetetraacetic acid (EDTA); buffers such as acetates, citrates or phosphates, and agents for the adjustment of tonicity, such as sodium chloride or dextrose. Suitable carriers include physiological saline, bacteriostatic

water, Cremophor® EL (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). The compositions can be stable during manufacture and storage and preserved against contamination from microorganisms, such as bacteria and fungi. Proper fluidity can be maintained, for example, by using a coating such as lecithin; by maintaining the required particle size in the case of dispersion, and by using surfactants. Various antibacterial and antifungal agents, such as parabens, chlorobutanol, phenol, ascorbic acid, and thimerosal, can control microorganism contamination. Isotonic agents, such as sugars, polyalcohols such as manitol, sorbitol, and sodium chloride can be included in the composition. Compositions that delay absorption can be prepared by including such agents as aluminum monostearate and gelatin.

[0178] Sterile injectable solutions can be prepared by incorporating the active compound (e.g., an SCMP) in an appropriate solvent with one or more ingredient, followed by sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and any other required ingredients. Sterile powders for the preparation of sterile injectable solutions include vacuum- and freeze-drying that yield a powder containing the active ingredient and any desired ingredient from a sterile solution. The concentration of active drug, *i.e.* the ROS scavenger, can be from about 0.1% to about 90% by weight, from about 5% to about 20% by weight, from about 5% to about 17% by weight, from about 8% to about 14% by weight or, in certain embodiments, about 10% by weight.

[0179] Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included. Tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, primogel, or corn starch; a lubricant such as magnesium stearate or sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent

such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring. The concentration of active drug, *i.e.* the ROS scavenger, can be from about 0.1% to about 99% by weight, from about 5% to about 95% by weight, from about 10% to about 90% by weight, from about 15% to about 85% by weight, from about 20% to about 80% by weight, from about 25% to about 75 % by weight, from about 30% to about 70% by weight, from about 35% to about 64% by weight or from about 40% to about 60% by weight.

[0180] Administration by inhalation, can be by aerosol spray from a nebulizer or a pressurized container that contains a suitable propellant, *e.g.*, a gas such as carbon dioxide.

[0181] Systemic administration can also be transmucosal or transdermal. For transmucosal or transdermal administration, penetrants that can permeate the target barrier(s) are selected. Transmucosal penetrants include detergents, bile salts and fusidic acid derivatives. Nasal sprays or suppositories can be used for transmucosal administration. For transdermal administration, the active compounds are formulated into ointments, salves, gels or creams.

[0182] The compounds can also be prepared as suppositories (with bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

[0183] In various embodiments, the active compounds can be prepared with carriers that protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid (Alza Corporation; Mountain View, CA and Nova Pharmaceuticals, Inc.; Lake Elsinore, CA). Liposomal suspensions can also be used as pharmaceutically acceptable carriers (Eppstein, 1985).

[0184] Oral formulations or parenteral compositions in unit dosage form can be created to facilitate administration and dosage uniformity. Unit dosage form refers to physically discrete units suited as single doses for a subject to be treated, containing a therapeutically effective quantity of active compound in association with the required pharmaceutical carrier. The specification for unit dosage forms are dictated by, and directly dependent on, the unique characteristics of the active compound and the particular desired therapeutic effect, and the inherent limitations of compounding the active compound.

75975-31

[0185] A typical dose of the ROS scavenger can be from about 0.1 mg up to about 1000 mg or from about 0.001 up to about 10 mg/kg body weight. Doses of 5 mg/kg and 10 mg/kg administered intraperitoneally have shown to produce beneficial effects in rats treated with type II collagen to induced arthritis (Salvemini *et al.*, Arthritis & Rheumatism, 44:2909-2921, 2001). A dose of 2 mg/kg produced less of an effect that was nevertheless significantly different from that in placebo animals. Low doses of a ROS scavenger can be doses of less than about 5 mg/kg or doses equal to or less than about 2 mg/kg body weight, in particular, a dose of about 0.1 mg, about 0.2 mg, about 0.5 mg, about 0.8 mg, about 1 mg, about 2 mg, about 5 mg, about 8 mg, about 10 mg, about 20 mg, about 50 mg, about 80 mg, about 100 mg or about 200 mg or about 0.001 mg/kg, about 0.002 mg/kg, about 0.005 mg/kg, about 0.01 mg/kg, about 0.02 mg/kg, about 0.05 mg/kg, about 0.1 mg/kg, about 0.2 mg/kg, about 0.5 mg/kg, about 1 mg/kg, about 2 mg/kg, about 3 mg/kg or about 4 mg/kg body weight.

[0186] Total daily doses of the ROS scavenger can be administered in single or divided doses and in amounts such as, for example, from about 1 to about 2 mg/kg body weight daily and more usually about 0.05 to 1 mg/kg. Dosage unit compositions may contain such amounts of submultiples thereof to make up the total dose. However, one skilled in the art will recognize that the total dosage will vary on the particular composition the particular ROS scavenger administered.

75975-31

[0186a] According to embodiments of the invention, the superoxide dismutase mimetic may be administered: on the day before or the day of, but prior to the cancer treatment; on the day before or the day of, but prior to the chemotherapy treatment; at least 30 minutes prior to the cancer treatment; up to three days after the radiation
5 therapy treatment; or at least 30 minutes prior to the radiation therapy treatment.

[0187] Individuals receiving treatment are, typically, human patients, however, patients receiving treatment can also be animal including companion animal such as dogs and cats, farm animal such as cows, horses, swine as well as birds and exotic animal such as zoo animals.

10 **[0188]** The amount of active ingredients that may be combined with the carrier materials to produce a single dosage form can vary depending upon the host treated and the particular mode of administration. It will be appreciated that the unit content of active ingredients contained in an individual dose of each dosage form need not in
15 itself constitute an effective amount, as the necessary effective amount could be reached by administration of a number of individual doses. The selection of dosage depends upon the dosage form utilized, the condition being treated, and the particular purpose to be achieved according to the determination of those skilled in the art.

[0189] The dosage regimen for treating a disease condition with the compounds

and/or compositions of this invention can be selected in accordance with a variety of factors, including the type, age, weight, sex, diet and medical condition of the patient, the route of administration, pharmacological considerations such as the activity, efficacy, pharmacokinetic and toxicology profiles of the particular compound employed, whether a drug delivery system is utilized and whether the compound is administered as part of a drug combination. Thus, the dosage regimen actually employed can, therefore, can deviate from the preferred dosage regimen set forth above.

[0190] In various embodiments, the present invention can also involve kits. Such kits can include pharmaceutical compositions and ,in addition in certain embodiments, instructions for administration. When supplied as a kit, the different components of the composition can be packaged in separate containers and admixed immediately before use. Such packaging of the components separately can, in certain instances, permit long-term storage without losing activity of the components. In addition, if more than one route of administration is intended or more than one schedule for administration is intended, the different components can be packaged separately and not mixed prior to use. In various embodiments, the different components can be packaged in one composition for administration together.

[0191] Kits may also include reagents in separate containers such as, for example sterile water or saline to be added to a lyophilized active component packaged separately. For example, sealed glass ampules may contain lyophilized ROS scavenger, sterile water, sterile saline or sterile each of which has been packaged under a neutral non-reacting gas, such as nitrogen. Ampules may consist of any suitable material, such as glass, organic polymers, such as polycarbonate, polystyrene, etc., ceramic, metal or any other material typically employed to hold reagents. Other examples of suitable containers include bottles that may be fabricated from similar substances as ampules, and envelopes that may consist of foil-lined interiors, such as aluminum or an alloy. Other containers include test tubes, vials, flasks, bottles, syringes, etc. Containers may have a sterile access port, such as a bottle having a stopper that can be pierced by a hypodermic injection needle. Other containers may have two compartments that are separated by a readily removable membrane that upon removal permits the components to mix. Removable membranes may be glass, plastic, rubber, etc.

[0192] In certain embodiments, kits can be supplied with instructional materials. Instructions may be printed on paper or other substrate, and/or may be supplied as an electronic-readable medium, such as a floppy disc, mini-CD-ROM, CD-ROM, DVD-ROM, Zip disc, videotape, audio tape, etc. Detailed instructions may not be physically associated with the kit; instead, a user may be directed to an internet web site specified by the manufacturer or distributor of the kit, or supplied as electronic mail.

[0193] EXAMPLES

[0194] Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following specific examples are offered by way of illustration and not by way of limiting the remaining disclosure.

[0195] EXAMPLE 1

[0196] OBJECTIVE

[0197] The objective of this study is to evaluate the effect of M40403, administered by either ip or topical routes, with 2 different schedules, on the frequency, severity and duration of oral mucositis induced in hamsters by acute radiation.

[0198] SUMMARY

[0199] M40403 was given as an intraperitoneal (ip) injection at a 30mg/kg/dose twice daily (a) from day -1 to day 15, or (b) day -1 to day 3. M40403 was also given by ip injection at a dose of 3 mg/kg twice daily from day -1 to day 15. In addition, M40403 was given as a topical dose directed to the buccal mucosa in 0.2 ml doses of either 3 mg/ml or 30 mg/ml twice daily from day -1 to day 15. The greatest reduction in oral mucositis was seen in the group treated with M40403 at 30 mg/kg/dose ip twice daily from day -1 to day 3. This group had a statistically significant reduction in the number of animal days with a mucositis score of 3 or higher ($P<0.001$) and significantly lower mean mucositis scores than the control group on days 18 ($P=0.041$), 20 ($P<0.001$), 22 ($P<0.001$), 24 ($P<0.001$) and 26 ($P=0.002$). No other treatment groups in this study showed a reduction in mucositis either by long term (day-1 to day 15) ip dosing or by topical dosing. This study establishes a schedule-dependant dosing method for the treatment of oral mucositis

with M40403.

[0200] Acute radiation model

[0201] The acute radiation model in hamsters, developed by the Principal Investigator, has proven to be an accurate, efficient and cost-effective technique to provide a preliminary evaluation of anti-mucositis compounds (9). The course of mucositis in this model is well defined and results in peak scores approximately 14-16 Days following radiation. The acute model has little systemic toxicity, resulting in few hamster deaths, thus permitting the use of smaller groups (N=7-8) for initial efficacy studies. It has also been used to study specific mechanistic elements in the pathogenesis of mucositis. Molecules that show efficacy in the acute radiation model may be further evaluated in the more complex models of fractionated radiation, chemotherapy, or concomitant therapy.

[0202] In this study, an acute radiation dose of 40 Gy on day 0 was administered locally to the cheek pouch. Clinically significant mucositis was observed on days 12 through 28.

[0203] PROTOCOL SUMMARY

[0204] Forty-eight male Syrian Golden Hamsters were divided randomly into 6 groups of 8 animals and given an acute radiation dose of 40 Gy directed to their left buccal cheek pouch. This was accomplished by anesthetizing the animals and everting the left buccal pouch, while protecting the rest of the animal with a lead shield. Test materials were given topically or by ip injection twice daily as detailed in Table 1. Mucositis was evaluated clinically starting on Day 6, and continued on alternate days until day 28.

[0205] EVALUATION

[0206] Mucositis evaluation

[0207] The grade of mucositis was scored, beginning on day 6, and for every second day thereafter, through and including day 28 (for scoring scheme, see Table 2 and Figure 2). The effect on mucositis of each drug treatment compared to ip vehicle-treated controls was assessed according to the following parameters:

[0208] The difference in the number of days hamsters in each group have ulcerative (score ≥ 3) mucositis.

[0209] On each evaluation day, the number of animals with a blinded mucositis score of ≥ 3 in each drug treatment group was compared to the vehicle-treated control group. Differences were compared on a cumulative basis and statistical significance was determined by chi-square analysis. Efficacy, in this analysis, is defined as a significant reduction in the number of days that a group of animals had ulcerations (scores ≥ 3) when compared to the control group.

TABLE 2

Table 2 ACT-02: Mucositis Scoring.

Score:	Description:
0	Pouch completely healthy. No erythema or vasodilation.
1	Light to severe erythema and vasodilation. No erosion of mucosa.
2	Severe erythema and vasodilation. Erosion of superficial aspects of mucosa leaving denuded areas. Decreased stippling of mucosa.
3	Formation of off-white ulcers in one or more places. Ulcers may have a yellow/gray due to pseudomembrane. Cumulative size of ulcers should equal about $\frac{1}{4}$ of the pouch. Severe erythema and vasodilation.
4	Cumulative size of ulcers should equal about $\frac{1}{2}$ of the pouch. Loss of pliability. Severe erythema and vasodilation.
5	Virtually all of pouch is ulcerated. Loss of pliability (pouch can only partially be extracted from mouth).

[0210] Rank sum differences in daily mucositis scores.

[0211] For each evaluation day the scores of the control group were compared to those of the treated groups using non-parametric rank sum analysis. Treatment success was considered as a statistically significant lowering of scores in the treated group on 2 or more days from day 6 to day 28.

[0212] Weights and survival

[0213] All animals were weighed daily and their survival recorded, in order to assess possible differences in animal weight among treatment groups as an indication for mucositis severity and/or possible toxicity resulting from the treatments.

[0214] STUDY DESIGN

[0215] All forty-eight (48) male Syrian Golden Hamsters were given an acute radiation dose of 40 Gy directed to their left buccal cheek pouch. This was accomplished by anesthetizing the animals and everting the left buccal pouch, while protecting the rest of the animal with a lead shield. Test materials were given either

topically or by ip injection twice daily as detailed in Table 1. Mucositis was evaluated clinically starting on day 6, and continuing on alternate days until day 28. The study timeline is depicted in Figure 1. Dose levels were determined based on the results of a previous acute toxicity study (data not shown), and the levels shown in Table 1.

TABLE 1

Table 1. ACT-02. Study Design

Group Number	Number of Animals	Treatment	Treatment Schedule*	Volume (mL)
1	8 males	Vehicle, ip, bid	Day -1 to 15	Adjust per body weigh
2	8 males	M40403, ip, bid 3 mg/kg	Day -1 to 15	Adjust per body weigh
3	8 males	M40403, ip, bid 30 mg/kg	Day -1 to 15	Adjust per body weigh
4	8 males	M40403, topical, bid 3 mg/ml	Day -1 to 15	0.2 ml per dose
5	8 males	M40403, topical, bid 30 mg/ml	Day -1 to 15	0.2 ml per dose
6	8 males	M40403, ip, bid 30 mg/kg	Day -1 to 3	Adjust per body weigh

*The first dose on day 0 was performed 30 minutes prior to radiation. The second dose was given at least 4 hours after radiation.

[0216] At the end of the study on day 28, the animals in group 1 were used in a modified pK study as follows: Two animals were injected with vehicle and blood was drawn 30 minutes after the injection. Six animals were injected with 30 mg/kg of M40403. At 15, 30 and 90 minutes post injection, 2 animals from this group were sacrificed and blood was obtained by cardiac puncture. Blood was collected in lithium heparin, kept on ice for 30 minutes and the plasma was obtained after centrifugation. Plasma was transferred to labeled tubes, snap frozen in liquid nitrogen and shipped to ActivBiotics.

[0217] MATERIAL AND METHODS

[0218] Location of study performance

[0219] The study was performed at Biomodels AAALAC accredited facility in Cambridge MA. The IACUC approval number 04-0624-2 for this study was obtained from Biomodels IACUC.

[0220] Animals

[0221] Male LVG Syrian Golden Hamsters (Charles River Laboratories), aged 5 to 6 weeks, with average body weight of 84.1g at study commencement, were used. Animals were individually numbered using an ear punch and housed in small groups of approximately 8 animals per cage. Animals were acclimatized prior to study commencement. During this period of 2 days, the animals were observed daily in order to reject animals that presented in poor condition.

[0222] Housing

[0223] The study was performed in animal rooms provided with filtered air at a temperature of 70oF+/-5 oF and 50% +/-20% relative humidity. Animal rooms were set to maintain a minimum of 12 to 15 air changes per hour. The room was on an automatic timer for a light/dark cycle of 12 hours on and 12 hours off with no twilight. Bed-O-Cobs® bedding was used. Bedding was changed a minimum of once per week. Cages, tops, bottles, etc. were washed with a commercial detergent and allowed to air dry. A commercial disinfectant was used to disinfect surfaces and materials introduced into the hood. Floors were swept daily and mopped a minimum of twice weekly with a commercial detergent. Walls and cage racks were sponged a minimum of once per month with a dilute bleach solution. A cage card or label with the appropriate information necessary to identify the study, dose, animal number and treatment group marked all cages. The temperature and relative humidity was recorded during the study, and the records were retained.

[0224] Diet

[0225] Animals were fed with a Purina Labdiet® 5061 rodent diet and water was provided ad libitum. Animal randomization and allocations.

[0226] Animals were randomly and prospectively divided into eight (8) treatment groups prior to irradiation. Each animal was identified by an ear punch corresponding to an individual number. For more consistent identification, ear punch numbering was used rather than tagging, since tags may become dislodged during the course of the study. A cage card was used to identify each cage and was marked with the study number, treatment group number and animal numbers.

[0227] Dosing and drug application

[0228] Dosing solutions were made immediately prior to use. Aseptic technique was used for all preparation procedures. A 26-mM sodium bicarbonate buffer solution was prepared as the vehicle. The resultant pH was approximately 8.1 to 8.3.

[0229] Mucositis induction

[0230] Mucositis was induced using a standardized acute radiation protocol. A single dose of radiation (40 Gy/dose) was administered to all animals on day 0. Radiation was generated with a 250 kilovolt potential (15-ma) source at a focal distance of 50 cm, hardened with a 0.35 mm Cu filtration system. Irradiation targeted the left buccal pouch mucosa at a rate of 3.2 Gy/minute. Prior to irradiation, animals were anesthetized with an intra-peritoneal injection of Ketamine (160 mg/kg) and Xylazine (8 mg/kg). The left buccal pouch was everted, fixed and isolated using a lead shield.

[0231] Mucositis scoring

[0232] The mucositis score, weight change and survival were measured throughout the study as described above. For the evaluation of mucositis, the animals were anesthetized with isoflurane and the left pouch everted. Mucositis was scored visually by comparison to a validated photographic scale (Figure 2), ranging from 0 for normal, to 5 for severe ulceration (clinical scoring). In descriptive terms, this scale is defined as follows:

[0233] A score of 1-2 is considered to represent a mild stage of the disease, whereas a score of 3-5 is considered to indicate moderate to severe mucositis. Following visual scoring, a photograph was taken of each animal's mucosa using a standardized technique. At the conclusion of the experiment, all films were developed and the photographs randomly numbered. At least two independent trained observers graded the photographs in blinded fashion using the above described scale (blinded scoring).

[0234] RESULTS AND DISCUSSION

[0235] Survival

[0236] Five deaths occurred during this study. Two animals, hamsters 8 and 23, in the vehicle control group and the group treated with M40403 at 30 mg/kg twice a day from day -1 to day 15 died on day 0 from apparent anesthesia overdoses. In

addition, two animals (hamsters 45 and 46) died in the group treated with M40403 at 30 mg/kg twice daily from day -1 to day 3. Hamster 45 died during radiation and hamster 46 was found dead the following morning. Both deaths appeared to be the result of anesthesia overdose, in the case of hamster 46 the response was delayed. The occurrence of anesthesia deaths in this model is anticipated in the experimental design. Hamster 19, in the group treated ip with M40403 at 30 mg/kg twice a day from day -1 to day 15 died on day 12. This death resulted after an extended period of failure to gain weight and may reflect toxicity of this dosing regimen as all animals in this group showed lack of weight gain (see Figures 3 and 4 below).

[0237] Weight change (Figures 3 and 4).

[0238] The percent daily weight change for each group is shown in Figure 3. The groups treated by ip injection were groups 1, 2, 3 and 6. The control animals (group 1) gained an average of 64.0% of their starting weights by the end of the study. The animals treated ip with M40403 at 3 mg/kg/dose twice daily from day -1 to day 15 (group 2) gained an average of 66.0% of their starting weight during the study. The group treated ip with M40403 at 30 mg/kg/dose twice daily from day -1 to day 15 (group 3) gained an average of 33.9% of their starting weight by day 28. The group treated ip with M40403 at 30 mg/kg/dose twice daily from day 1 to day 3 gained an average of 54.7% of their starting weight during the study. In both groups treated by ip injection with 30 mg/kg M40403, there was a general lack of weight gain during dosing.

[0239] In the 2 groups treated topically with M40403 (groups 4 and 5) there was no apparent change in weight gain when compared to the control group. The animals treated topically with M40403 at 3 mg/ml twice daily from day -1 to day 15 gained an average of 65.8% of their starting weight by day 28. The group treated topically with M40403 at 30 mg/ml twice daily from day -1 to day 15 gained an average of 71.9% of their starting weight during the study.

[0240] The significance of these differences was evaluated by calculating the mean area under the curve for the percentage weight gain for each animal and comparing the groups using a OneWay ANOVA test. There was a significant difference between the control group and group 3 treated by ip injection with M40403 twice daily (30 mg/kg) on days -1 to 15 ($P < 0.001$). There was also a significant difference between the control group and the group treated by ip injection with M40403 twice daily at 30

mg/kg/dose from day -1 to day 3 ($P=0.018$). No other significant differences were seen. The topical treatment groups (groups 4 and 5) showed no significant weight differences when compared with the control group. The AUC data is shown in Figure 4. Many of the groups with a significant reduction in weight gain are the same groups in which animal deaths were observed.

TABLE 4

Group Comparison	Day											
	6	8	10	12	14	16	18	20	22	24	26	28
Control v M40403 ip 3 mg/kg day -1 to 15	0.099	0.630	0.950	0.266	0.983	0.630	0.436	0.571	0.754	0.325	0.516	0.070
Control v M40403 ip 30 mg/kg day -1 to 15	0.055	0.981	0.981	0.055	0.856	0.938	0.221	0.622	0.698	0.067	0.979	0.225
Control v M40403 top 3 mg/ml day -1 to 15	0.099	0.099	0.630	0.463	0.463	0.438	0.983	0.983	0.163	0.632	0.517	0.139
Control v M40403 top 30 mg/ml day -1 to 15	0.630	0.099	0.002	0.573	0.465	0.571	0.884	0.884	0.573	0.632	0.754	0.884
Control v M40403 ip 30 mg/kg day -1 to 3	0.127	0.059	0.546	0.938	0.221	0.312	0.041	<0.001	<0.001	<0.001	0.002	0.551

Table 4. ACT-02. The significance of group differences observed in daily mucositis scores was determined using the Mann-Whitney rank sum test. This nonparametric statistic is appropriate for the visual mucositis scoring scale. The p values for each calculation are shown.

[0241] Mucositis (Figures 5 & 6, Tables 3 & 4)

[0242] Mean daily mucositis scores for each group are shown in Figure 5. The groups treated by ip injection were groups 1, 2, 3 and 6. The control animals (group 1) had a peak mean mucositis score of 2.8 which occurred on day 16. The animals treated with M40403 at 3 mg/kg/dose ip twice daily from day -1 to day 15 (group 2) had a peak mucositis score of 2.8 on day 18, and the overall progression of mucositis severity in this group was very similar to that observed in the control group. The group treated with M40403 at 30 mg/kg/dose twice daily from day -1 to day 15 (group 3) had a peak mucositis score of 3.0 on day 18 and had a mucositis progression that was indistinguishable from that observed in groups 1 and 2. The group treated with M40403 at 30 mg/kg/dose ip twice daily from day -1 to day 3 had a peak mucositis score of 3.0 on day 14, but from day 16 to day 28, the mucositis severity in this group decreased much more rapidly than the control group suggesting that this schedule of treatment was efficacious.

[0243] In the 2 groups treated topically with M40403 (groups 4 and 5) there was no apparent effect of treatment on the course of mucositis. The animals treated topically with M40403 at 3 mg/ml twice daily from day -1 to day 15 had mucositis scores that closely paralleled those in the control group. The group treated topically with M40403 at 30 mg/ml twice daily from day -1 to day 15 showed a reduction of mucositis severity on day 10. However, from day 12 to the end of the study, the scores in this group were only slightly less than those in the control group.

[0244] The significance of the reductions in the mucositis scores seen in the groups treated with M40403 were evaluated by calculating the percentage of animal days with a score of 3 or higher. The results of this analysis are shown in Figure 6 and Table 3. In the control group, the percentage of animal days with a mucositis score of 3 or higher was 31%. Treatment by ip injection with M40403 at 3 mg/kg/dose twice daily from day -1 to day 15 resulted in a percentage of animal days with a score of 3 or higher of 34%. Treatment by ip injection with M40403 at 30 mg/kg/dose twice daily from day to -1 to day 15 resulted in 37% percent of animal days with a score of 3 or higher. Treatment by ip injection with M40403 at 30 mg/kg twice daily from day -1 to day 3 (group 6) resulted in a decrease in the number of animal days with a score of 3 or higher to 17%.

TABLE 3

Group	Days >=3	Days<3	Total Days	% Days >=3	Chi Sq v control	P Value
Vehicle ip control day -1 to 15	52	116	168	0.31	---	---
M40403 ip 3 mg/kg day -1 to 15	66	126	192	0.34	0.3440	0.563
M40403 ip 30 mg/kg day -1 to 15	56	96	152	0.37	0.9890	0.320
M40403 top 3 mg/ml day -1 to 15	73	119	192	0.38	1.6760	0.196
M40403 top 30 mg/ml day -1 to 15	64	128	192	0.33	0.136	0.712
M40403 ip 30 mg/kg day -1 to 3	24	120	144	0.17	7.83	0.005

Table 3. ACT-02. Chi-square analysis of the total number of days the animals in each group spent with a mucositis score of 3 or more. This statistic is a measure of severity of ulceration, a clinically important outcome.

[0245] In the animals receiving M40403 topically, there were no apparent reductions of mucositis severity at either dosing level. Topical treatment with M40403 at 3 mg/ml twice daily from day -1 to day 15 resulted in an increase in the number of animal days with a score of 3 or higher to 38%. Topical treatment with M40403 at 30 mg/ml twice daily from days -1 to 15 resulted in 33% of animal days with a score of 3 or higher.

[0246] When compared using a chi-squared test, the group treated by ip injection with M40403 at 30 mg/kg/dose twice daily on days -1 to day 3 had the greatest reduction in mucositis and significantly fewer days with a score of 3 or higher (P=0.005). No other study group exhibited a reduction in mucositis severity and no group demonstrated a significant difference in mucositis severity by this analysis.

[0247] Further analysis of the mucositis scores was performed by comparing the

scores for the M40403 treated groups with the control on each day of scoring using the Mann-Whitney Rank Sum test. The results of this analysis are shown in Table 4. The group treated by ip injection with M40403 at 30 mg/kg/dose twice daily on days -1 to day 3 (group 6) had the greatest reduction in mucositis and significantly lower scores than the control group on days 18 ($P=0.041$), 20 ($P<0.001$), 22 ($P<0.001$), 24 ($P<0.001$) and 26 ($P=0.002$). No other study group had more than a single day showing a significant lowering of mucositis severity. Given that group 2 received the same dose of drug for a longer duration, these results suggest that the schedule of dosing M40403 is crucial in establishing an effective treatment for mucositis.

[0248] CONCLUSIONS

[0249] 1. Based on observations of mortality and body weight, ip injections of M40403, twice daily, at 30 mg/kg appear to show some toxicity. The longer dosing in group 3 resulted in mortality and significant weight loss. When dosed for the shorter period of day -1 to day 3 (group 6), the weight loss was reversed after the cessation of dosing.

[0250] 2. Topical dosing of M40403 did not appear to have any effect on either weight or mucositis severity. It appears the topical dosing, in the formulation used here, is an ineffective method of delivery of M40403 in this model.

[0251] 3. The group treated by ip injection with M40403 at 30 mg/kg/dose twice daily on days 1 to 3 had a statistically significant reduction in the number of animal days with a mucositis score of 3 or higher ($P=0.005$) and significantly lower mucositis scores than the control group on days 18 ($P=0.041$), 20 ($P<0.001$), 22 ($P<0.001$), 24 ($P<0.001$) and 26 ($P=0.002$). This result suggests that M40403 may be an effective agent in the treatment of oral mucositis. Further studies of dose and schedule may be required to optimize the efficacy of M40403.

[0252] EXAMPLE 2 -- Repeat of day -1 to 3 of 30 mg/kg IV of Example 1 study

[0253] Acute radiation model

[0254] The acute radiation model in hamsters, developed by the Principal Investigator, has proven to be an accurate, efficient and cost-effective technique to provide a preliminary evaluation of anti-mucositis compounds (9). The course of mucositis in this model is well defined and results in peak scores approximately 14-16 Days following

radiation. The acute model has little systemic toxicity, resulting in few hamster deaths, thus permitting the use of smaller groups (N=7-8) for initial efficacy studies. It has also been used to study specific mechanistic elements in the pathogenesis of mucositis. Molecules that show efficacy in the acute radiation model may be further evaluated in the more complex models of fractionated radiation, chemotherapy, or concomitant therapy.

[0255] In this study, an acute radiation dose of 40 Gy on day 0 was administered locally to the cheek pouch. Clinically significant mucositis was observed on days 12 through 28.

[0256] PROTOCOL SUMMARY

[0257] Seventy-two male Syrian Golden Hamsters were divided randomly into 9 groups of 8 animals and given an acute radiation dose of 40 Gy directed to their left buccal cheek pouch. This was accomplished by anesthetizing the animals and everting the left buccal pouch, while protecting the rest of the animal with a lead shield. Test materials were given topically or by ip injection twice daily as detailed in Table 5. Mucositis was evaluated clinically starting on Day 6, and continued on alternate days until day 28.

Table 1. ACT-03. Study Design

Group Number	Number of Animals	Treatment	Treatment Schedule*	Volume (mL)
1	8 males	Vehicle, ip, bid	Day -1 to 3	Adjust per body weigh
2	8 males	M40403, ip, QD 30 mg/kg	Day -1 to 3	Adjust per body weigh
3	8 males	M40403, ip, bid 3 mg/kg	Day -1 to 3	Adjust per body weigh
4	8 males	M40403, ip, bid 10 mg/kg	Day -1 to 3	Adjust per body weigh
5	8 males	M40403, ip, bid 30 mg/kg	Day -1 to 3	Adjust per body weigh
6	8 males	M40403, ip, QD 30 mg/kg	Day 0 to 3	Adjust per body weigh
7	8 males	M40403, ip, bid 30 mg/kg	Day 0 to 3	Adjust per body weigh
8	8 males	M40403, ip, bid 30 mg/kg	Day 0	Adjust per body weigh
9	8 males	M40403, ip, bid 30 mg/kg	Day 0 and Day 7	Adjust per body weigh

*For BID dosing the first dose on day 0 will be performed 30 minutes prior to radiation. The second dose will be given at least 4 hours after radiation. For QD dosing, the dose will be administered on day 0 at 30 minutes prior to irradiation.

TABLE 5

[0258] EVALUATION

[0259] Mucositis evaluation

[0260] The grade of mucositis was scored, beginning on day 6, and for every second day thereafter, through and including day 28 (for scoring scheme, see Table 6 and Figure 8). The effect on mucositis of each drug treatment compared to ip vehicle-treated controls was assessed according to the following parameters:

[0261] The difference in the number of days hamsters in each group have ulcerative (score ≥ 3) mucositis.

[0262] On each evaluation day, the number of animals with a blinded mucositis score of ≥ 3 in each drug treatment group was compared to the vehicle-treated control group. Differences were compared on a cumulative basis and statistical significance was determined by chi-square analysis. Efficacy, in this analysis, is defined as a significant reduction in the number of days that a group of animals had ulcerations (scores ≥ 3) when compared to the control group.

Table 2 ACT-03: Mucositis Scoring.

Score:	Description:
0	Pouch completely healthy. No erythema or vasodilation.
1	Light to severe erythema and vasodilation. No erosion of mucosa.
2	Severe erythema and vasodilation. Erosion of superficial aspects of mucosa leaving denuded areas. Decreased stippling of mucosa.
3	Formation of off-white ulcers in one or more places. Ulcers may have a yellow/gray due to pseudomembrane. Cumulative size of ulcers should equal about $\frac{1}{4}$ of the pouch. Severe erythema and vasodilation.
4	Cumulative size of ulcers should equal about $\frac{1}{2}$ of the pouch. Loss of pliability. Severe erythema and vasodilation.
5	Virtually all of pouch is ulcerated. Loss of pliability (pouch can only partially be extracted from mouth).

TABLE 6

[0263] **Rank sum differences in daily mucositis scores.**

[0264] For each evaluation day the scores of the control group were compared to those of the treated groups using non-parametric rank sum analysis. Treatment success was considered as a statistically significant lowering of scores in the treated group on 2 or more days from day 6 to day 28.

[0265] **Weights and survival**

[0266] All animals were weighed daily and their survival recorded, in order to assess

possible differences in animal weight among treatment groups as an indication for mucositis severity and/or possible toxicity resulting from the treatments.

[0267] STUDY DESIGN

[0268] All seventy-two (72) male Syrian Golden Hamsters were given an acute radiation dose of 40 Gy directed to their left buccal cheek pouch. This was accomplished by anesthetizing the animals and everting the left buccal pouch, while protecting the rest of the animal with a lead shield. Test materials were given either topically or by ip injection twice daily as detailed in Table 5. Mucositis was evaluated clinically starting on day 6, and continuing on alternate days until day 28. The study timeline is depicted in Figure 7. Dose levels were determined based on the results of a previous acute toxicity study (data not published), and in the prior mucositis study of Example 1, shown in Table 5.

[0269] MATERIAL AND METHODS

[0270] Location of study performance

[0271] The study was performed at Biomodels AAALAC accredited facility in Cambridge MA. The IACUC approval number 04-0624-2 for this study was obtained from Biomodels IACUC.

[0272] Animals

[0273] Male LVG Syrian Golden Hamsters (Charles River Laboratories), aged 5 to 6 weeks, with average body weight of 92.7 at study commencement, were used. Animals were individually numbered using an ear punch and housed in small groups of approximately 8 animals per cage. Animals were acclimatized prior to study commencement. During this period of 3 days, the animals were observed daily in order to reject animals that presented in poor condition.

[0274] Housing

[0275] The study was performed in animal rooms provided with filtered air at a temperature of 70oF+/-5 oF and 50% +/-20% relative humidity. Animal rooms were set to maintain a minimum of 12 to 15 air changes per hour. The room was on an automatic

timer for a light/dark cycle of 12 hours on and 12 hours off with no twilight. Bed-O-Cobs® bedding was used. Bedding was changed a minimum of once per week. Cages, tops, bottles, etc. were washed with a commercial detergent and allowed to air dry. A commercial disinfectant was used to disinfect surfaces and materials introduced into the hood. Floors were swept daily and mopped a minimum of twice weekly with a commercial detergent. Walls and cage racks were sponged a minimum of once per month with a dilute bleach solution. A cage card or label with the appropriate information necessary to identify the study, dose, animal number and treatment group marked all cages. The temperature and relative humidity was recorded during the study, and the records were retained.

[0276] Diet

[0277] Animals were fed with a Purina Labdiet® 5061 rodent diet and water was provided ad libitum. Animal randomization and allocations.

[0278] Animals were randomly and prospectively divided into eight (8) treatment groups prior to irradiation. Each animal was identified by an ear punch corresponding to an individual number. For more consistent identification, ear punch numbering was used rather than tagging, since tags may become dislodged during the course of the study. A cage card was used to identify each cage and was marked with the study number (ACT-03), treatment group number and animal numbers.

[0279] Dosing and drug application

[0280] Dosing solutions were made immediately prior to use. Aseptic technique was used for all preparation procedures. A 26-mM sodium bicarbonate buffer solution was prepared as the vehicle. The resultant pH was approximately 8.1 to 8.3.

[0281] Mucositis induction

[0282] Mucositis was induced using a standardized acute radiation protocol. A single dose of radiation (40 Gy/dose) was administered to all animals on day 0. Radiation was generated with a 250 kilovolt potential (15-ma) source at a focal distance of 50 cm, hardened with a 0.35 mm Cu filtration system. Irradiation targeted the left buccal pouch mucosa at a rate of 3.2 Gy/minute. Prior to irradiation, animals were anesthetized with

an intra-peritoneal injection of Ketamine (160 mg/kg) and Xylazine (8 mg/kg). The left buccal pouch was everted, fixed and isolated using a lead shield.

[0283] Mucositis scoring

[0284] The mucositis score, weight change and survival were measured throughout the study as described above. For the evaluation of mucositis, the animals were anesthetized with isoflurane and the left pouch everted. Mucositis was scored visually by comparison to a validated photographic scale (Figure 8), ranging from 0 for normal, to 5 for severe ulceration (clinical scoring). In descriptive terms, this scale is defined as follows:

[0285] A score of 1-2 is considered to represent a mild stage of the disease, whereas a score of 3-5 is considered to indicate moderate to severe mucositis. Following visual scoring, a photograph was taken of each animal's mucosa using a standardized technique. At the conclusion of the experiment, all films were developed and the photographs randomly numbered. At least two independent trained observers graded the photographs in blinded fashion using the above-described scale (blinded scoring).

[0286] RESULTS AND DISCUSSION

[0287] Survival

[0288] Seven deaths occurred during this study, all on the day of radiation (Day 0). Two animals each died in the group treated with M40403 at 30 mg/kg once daily from day -1 to day 3, the group treated with M40403 at 30 mg/kg twice daily from day -1 to day 3 and the group treated with M40403 at 30 mg/kg once daily from day 0 to day 3. One animal died in the group treated with M40403 at 30 mg/kg once daily on day 0 and day 7

[0289] Weight change (Figures 9 and 10).

[0290] The percent daily weight change for each group is shown in Figure 9. The control animals (group 1) gained an average of 62.5% of their starting weights by the end of the study. The animals treated ip with M40403 at 30 mg/kg/dose once daily from day -1 to day 3 (group 2) gained an average of 60.7% of their starting weight during the study. The group treated ip with M40403 at 3 mg/kg/dose twice daily from day -1 to day

3 (group 3) gained an average of 59.3% of their starting weight by day 28. The group treated ip with M40403 at 10 mg/kg/dose twice daily from day -1 to day 3 (group 4) gained an average of 56.2% of their starting weight during the study. The group treated ip with M40403 at 30 mg/kg/dose twice daily from day -1 to day 3 (group 5) gained an average of 52.9% of their starting weight during the study. The group treated ip with M40403 at 30 mg/kg/dose once daily from day 0 to day 3 (group 6) gained an average of 54.8% of their starting weight during the study. The group treated ip with M40403 at 30 mg/kg/dose twice daily from day 0 to day 3 (group 7) gained an average of 58.1% of their starting weight during the study. The group treated ip with M40403 at 30 mg/kg/dose twice daily on day 0 (group 8) gained an average of 59.3% of their starting weight during the study. The group treated ip with M40403 at 30 mg/kg/dose twice daily on day 0 and on day 7 (group 9) gained an average of 57.3% of their starting weight during the study.

[0291] The significance of these differences was evaluated by calculating the mean area under the curve for the percentage weight gain for each animal and comparing the groups using a One-Way ANOVA test. There were significant differences between the vehicle control group and the group treated with M40403 at 30 mg/kg/dose once daily from day -1 to day 3 ($P=0.039$), the group treated with M40403 at 3 mg/kg/dose twice daily from day -1 to day 3 ($P=0.040$), the group treated with M40403 at 10 mg/kg/dose twice daily from day -1 to day 3 ($P=0.018$), the group treated with M40403 at 30 mg/kg/dose twice daily from day -1 to day 3 ($P<0.001$), the group treated with M40403 at 30 mg/kg/dose once daily from day 0 to day 3 ($P=0.009$) and the group treated with M40403 at 30 mg/kg/dose twice daily from day 0 to day 3 ($P=0.002$). The results of this analysis are shown in Figure 10.

[0292] Mucositis (Figures 11 & 12, Tables 7 & 8)

[0293] Mean daily mucositis scores for each group are shown in Figure 11. The control animals (group 1) had a peak mean mucositis score of 3.1 on day 16. The animals treated ip with M40403 at 30 mg/kg/dose once daily from day -1 to day 3 (group 2) had a peak mean mucositis score of 2.8 on day 16. The group treated ip with M40403 at 3 mg/kg/dose twice daily from day -1 to day 3 (group 3) had a peak mean mucositis score of 2.9, which occurred on days 16 and 18. The group treated ip with M40403 at 10

mg/kg/dose twice daily from day -1 to day 3 (group 4) had a peak mean mucositis score of 3.0 on day 16. The group treated ip with M40403 at 30 mg/kg/dose twice daily from day -1 to day 3 (group 5) had a peak mean mucositis score of 2.8 on day 16. The group treated ip with M40403 at 30 mg/kg/dose once daily from day 0 to day 3 (group 6) had a peak mean mucositis score of 3.2 on day 16. The group treated ip with M40403 at 30 mg/kg/dose twice daily from day 0 to day 3 (group 7) had a peak mean mucositis score of 2.2 on day 16. The group treated ip with M40403 at 30 mg/kg/dose twice daily on day 0 (group 8) had a peak mean mucositis score of 2.4 on day 14. The group treated ip with M40403 at 30 mg/kg/dose twice daily on day 0 and on day 7 (group 9) had a peak mean mucositis score of 2.4 which occurred on days 16 and 18.

[0294] The significance of the reductions in the mucositis scores seen in the groups treated with M40403 were evaluated by calculating the percentage of animal days with a score of 3 or higher. The results of this analysis are shown in Figure 12 and Table 7. In the control group, the percentage of animal days with a mucositis score of 3 or higher was 36.5%. Treatment by ip injection with M40403 at 30 mg/kg/dose once daily from day -1 to day 3 (group 2) or M40403 at 3 mg/kg/dose twice daily from day -1 to day 3 (group 3) resulted in a percentage of animal days with a score of 3 or higher of 25%. Treatment with M40403 at 10 mg/kg/dose twice daily from day -1 to day 3 (group 4) resulted in a percentage of animal days with a score of 3 or higher of 29.2%. Treatment with M40403 at 30 mg/kg/dose twice daily from day -1 to day 3 (group 5) resulted in a percentage of animal days with a score of 3 or higher of 22.2%. Treatment with M40403 at 30 mg/kg/dose once daily from day 0 to day 3 (group 6) resulted in a percentage of animal days with a score of 3 or higher of 30.6%. Treatment with M40403 at 30 mg/kg/dose twice daily from day 0 to day 3 (group 7) resulted in a percentage of animal days with a score of 3 or higher of 9.4%. Treatment with M40403 at 30 mg/kg/dose twice daily on day 0 (group 8) resulted in a percentage of animal days with a score of 3 or higher of 13.5%. Treatment with M40403 at 30 mg/kg/dose twice daily on day 0 and on day 7 (group 9) resulted in a percentage of animal days with a score of 3 or higher of 15.5%. When compared using a chi-squared test, significant reductions in the number of animal days with a score of 3 or higher were seen in the groups treated with M40403 at 30 mg/kg/dose once daily from day -1 to day 3 (group 2, $P=0.034$), M40403 at 3 mg/kg/dose twice daily from day -1 to day 3 (group 3, $P=0.020$), M40403 at 30

mg/kg/dose twice daily from day -1 to day 3 (group 5, $P=0.007$), M40403 at 30 mg/kg/dose twice daily from day 0 to day 3 (group 7, $P<0.001$), M40403 at 30 mg/kg/dose twice daily on day 0 (group 8, $P<0.001$) and M40403 at 30 mg/kg/dose twice daily on day 0 and on day 7 (group 9, $P<0.001$). This data is shown in Figure 12 and Table 7.

Group	Days ≥ 3	Days < 3	Total Days	% Days ≥ 3	Chi Sq v control	P Value
Vehicle ip control bid day -1 to 3	70	122	192	36.46%	---	---
M40403 ip 30 mg/kg qd day -1 to 3	36	108	144	25.00%	4.4860	0.034
M40403 ip 3 mg/kg bid day -1 to 3	48	144	192	25.00%	5.3950	0.020
M40403 ip 10 mg/kg bid day -1 to 3	56	136	192	29.17%	1.1996	0.158
M40403 ip 30 mg/kg bid day -1 to 3	32	112	144	22.22%	7.229	0.007
M40403 ip 30 mg/kg qd day 0 to 3	44	100	144	30.56%	1.029	0.310
M40403 ip 30 mg/kg bid day 0 to 3	18	174	192	9.38%	38.3440	<0.001
M40403 ip 30 mg/kg bid day 0	26	166	192	13.54%	25.681	<0.001
M40403 ip 30 mg/kg bid day 0 & 7	26	142	168	15.48%	19.113	<0.001

Table 3. ACT-03. Chi-square analysis of the total number of days the animals in each group spent with a mucositis score of three or more. This statistic is a measure of severity of ulceration, a clinically important outcome.

TABLE 7

[0295] Further analysis of the mucositis scores was performed by comparing the scores for the M40403 treated groups with the control on each day of scoring using the Mann-Whitney Rank Sum test. The results of this analysis are shown in Table 8. The groups treated with M40403 at 30 mg/kg/dose once daily from day -1 to day 3 (group 2),

M40403 at 10 mg/kg/dose twice daily from day -1 to day 3 (group 4) and M40403 at 30 mg/kg/dose once daily from day 0 to day 3 (group 6) did not show any days with significant differences relative to controls. The scores for the group treated with M40403 at 3 mg/kg/dose twice daily from day -1 to day 3 (group 3) were significant different from controls on days 14 ($P=0.013$), 22 ($P=0.047$) and 28 ($P=0.012$). The group treated with M40403 at 30 mg/kg/dose twice daily from day -1 to day 3 (group 5) was significant different from controls on days 24 ($P=0.048$). The group treated with M40403 at 30 mg/kg/dose twice daily from day 0 to day 3 (group 7) had significantly lower scores than the control group on days 14 ($P<0.001$), 16 ($P=0.001$), 18 ($P=0.007$), 20 ($P=0.001$), 22 ($P<0.001$), 24 ($P=0.006$), 26 ($P=0.001$) and 28 ($P=0.012$). The group treated with M40403 at 30 mg/kg/dose twice daily on day 0 (group 8) had significantly lower scores than the control group on days 16 ($P=0.001$), 18 ($P=0.033$), 20 ($P=0.002$), 22 ($P<0.001$), 24 ($P=0.010$), 26 ($P<0.001$) and 28 ($P=0.009$). The group treated with M40403 at 30 mg/kg/dose twice daily on day 0 and on day 7 (group 9) had significantly lower scores than the control group on days 14 ($P=0.004$), 16 ($P=0.047$), 20 ($P<0.001$), 22 ($P=0.015$) and 26 ($P=0.047$).

Group Comparison	Day											
	6	8	10	12	14	16	18	20	22	24	26	28
Control v M40403 ip 30 mg/kg qd day -1 to 3	0.467	0.655	0.944	0.499	0.097	0.231	0.233	0.315	0.235	0.294	0.273	0.253
Control v M40403 ip 3 mg/kg bid day -1 to 3	0.985	0.231	0.985	0.748	<u>0.013</u>	0.297	0.865	0.459	<u>0.047</u>	0.610	0.283	<u>0.012</u>
Control v M40403 ip 10 mg/kg bid day -1 to 3	0.230	0.985	0.985	0.865	0.506	0.955	0.437	0.414	0.354	0.595	0.463	0.584
Control v M40403 ip 30 mg/kg bid day -1 to 3	0.725	0.655	0.410	0.428	0.097	0.141	0.216	0.097	0.170	<u>0.048</u>	0.053	0.066
Control v M40403 ip 30 mg/kg qd day 0 to 3	0.981	0.944	0.798	0.834	0.361	0.870	0.592	0.981	0.437	0.388	0.362	0.798
Control v M40403 ip 30 mg/kg bid day 0 to 3	0.776	0.555	0.472	0.940	<u><0.001</u>	<u>0.001</u>	<u>0.007</u>	<u>0.001</u>	<u><0.001</u>	<u>0.006</u>	<u>0.001</u>	<u>0.012</u>
Control v M40403 ip 30 mg/kg bid day 0	0.776	0.985	0.374	0.519	0.061	<u>0.001</u>	<u>0.033</u>	<u>0.002</u>	<u><0.001</u>	<u>0.010</u>	<u><0.001</u>	<u>0.009</u>
Control v M40403 ip 30 mg/kg bid day 0 & 7	0.415	0.721	0.156	0.851	<u>0.004</u>	<u>0.047</u>	0.091	<u><0.001</u>	<u>0.015</u>	0.163	<u>0.047</u>	0.219

Table 4. ACT-03. The significance of group differences observed in daily mucositis scores was determined using the Mann-Whitney rank sum test. This nonparametric statistic is appropriate for the visual mucositis scoring scale. The p values for each calculation are shown. Significant improvements are shown underlined

TABLE 8

[0296] CONCLUSIONS

[0297] 1. Based on observations of body weight, ip injections of M40403, appear to show some toxicity. The only groups that did not exhibit some statistically significant reductions in growth rate were those dosed with M40404 on day 0 only or on day 0 and day 7. Some mortality was noted on the day of radiation, but it is not clear whether this was related to treatment with M40403, or simply accidental death during the anesthesia process

[0298] 2. The groups treated with M40403 at 30 mg/kg/dose once daily from day -1 to day 3, M40403 at 10 mg/kg/dose twice daily from day -1 to day 3 or M40403 at 30 mg/kg/dose once daily from day 0 to day 3 did not generally show any significant reduction in the course of mucositis relative to controls, although the group treated with M40403 at 30 mg/kg/dose once daily from day -1 to day 3 showed a significant reduction in the number of animals days with a score of 3 or higher ($P=0.034$).

[0299] 3. The group treated with M40403 at 3 mg/kg/dose twice daily from day -1 to day 3 had significant lower mucositis scores from controls on days 14 ($P=0.013$), 22 ($P=0.047$) and 28 ($P=0.012$), and had a significant reduction in the number of animals days with a score of 3 or higher ($P=0.020$).

[0300] 4. The group treated with M40403 at 30 mg/kg/dose twice daily from day -1 to day 3 had significantly lower scores than the control group on day 24 ($P=0.048$) and had a significant reduction in the number of animals days with a score of 3 or higher ($P=0.007$).

[0301] 5. The group treated with M40403 at 30 mg/kg/dose twice daily from day 0 to day 3 (group 7) had significantly lower scores than the control group on days 14 ($P<0.001$), 16 ($P=0.001$), 18 ($P=0.007$), 20 ($P=0.001$), 22 ($P<0.001$), 24 ($P=0.006$), 26 ($P=0.001$) and 28 ($P=0.012$) and had a significant reduction in the number of animals days with a score of 3 or higher ($P<0.001$).

[0302] 6. The group treated with M40403 at 30 mg/kg/dose twice daily on day 0 (group 8) had significantly lower scores than the control group on days 16 ($P=0.001$), 18 ($P=0.033$), 20 ($P=0.002$), 22 ($P<0.001$), 24 ($P=0.010$), 26 ($P<0.001$) and 28 ($P=0.009$) and had a significant reduction in the number of animals days with a score of 3 or higher

75975-31

(P<0.001).

[0303] 7. The group treated with M40403 at 30 mg/kg/dose twice daily on day 0 and on day 7 had significantly lower scores than the control group on days 14 (P=0.004), 16 (P=0.047), 20 (P<0.001), 22 (P=0.015) and 26 (P=0.047) and had a significant reduction in the number of animals days with a score of 3 or higher (P<0.001).

[0304] Other Embodiments

[0305] The detailed description set-forth above is provided to aid those skilled in the art in practicing the present invention. However, the invention described and claimed herein is not to be limited in scope by the specific embodiments herein disclosed because these embodiments are intended as illustration of several aspects of the invention.

[0306] References Cited

[0307] Citation of a reference herein shall not be construed as an admission that such is prior art to the present invention.

1. Knox JJ, Puodziunas AL, Feld R. Chemotherapy-induced oral mucositis. Prevention and management. *Drugs Aging* 2000;17(4):257-67.
2. Peterson DE. Research advances in oral mucositis. *Curr Opin Oncol* 1999;11(4):261-6.
3. Plevova P. Prevention and treatment of chemotherapy- and radiotherapy-induced oral mucositis: a review. *Oral Oncol* 1999;35(5):453-70.
4. Sonis ST, Oster G, Fuchs H, Bellm L, Bradford WZ, Edelsberg J, et al. Oral

mucositis and the clinical and economic outcomes of hematopoietic stem-cell transplantation. J Clin Oncol 2001;19(8):2201-5.

5. Eldor A, Fuks Z, Matzner Y, Witte LD, Vlodavsky I. Perturbation of endothelial functions by ionizing irradiation: effects on prostaglandins, chemoattractants and mitogens. Semin Thromb Hemost 1989;15(2):215-25.

6. Sonis ST, Van Vugt AG, McDonald J, Dotoli E, Schwertschlag U, Szklut P, et al. Mitigating effects of interleukin 11 on consecutive courses of 5- fluorouracil-induced ulcerative mucositis in hamsters. Cytokine 1997;9(8):605-12.

7. Sonis ST, Van Vugt AG, Brien JP, Muska AD, Bruskin AM, Rose A, et al. Transforming growth factor-beta 3 mediated modulation of cell cycling and attenuation of 5-fluorouracil induced oral mucositis. Oral Oncol 1997;33(1):47-54.

8. Sonis ST, Peterson RL, Edwards LJ, Lucey CA, Wang L, Mason L, et al. Defining mechanisms of action of interleukin-11 on the progression of radiation-induced oral mucositis in hamsters. Oral Oncol 2000;36(4):373-81.

9. Sonis ST, Tracey C, Shklar G, Jenson J, Florine D. An animal model for mucositis induced by cancer chemotherapy. Oral Surg Oral Med Oral Pathol 1990;69(4):437-448

APPENDICES

Appendix 1: Animal Weights

		DAY																							
Group	Animal	-1	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
1	1	89	92	92	96	101	102	107	110	113	116	118	119	121	118	125	126	126	130	132	133	137	138	139	143
1	2	94	101	101	108	109	109	116	118	118	122	121	125	128	123	134	134	134	138	140	144	146	151	151	156
1	3	93	98	98	99	102	105	107	104	112	115	115	118	120	118	118	119	123	124	127	130	131	129	133	135
1	4	88	92	90	93	96	98	98	101	102	106	108	109	110	132	114	115	118	117	121	124	126	134	129	132
1	5	98	98	98	102	106	109	113	108	117	123	123	127	129	130	130	133	135	138	141	142	148	147	150	153
1	6	97	102	101	107	109	111	116	118	121	123	125	127	130	131	130	131	131	136	138	141	141	142	143	146
1	7	87	95	90	94	97	98	105	104	101	105	105	105	107	110	109	111	112	116	116	120	121	122	123	127
1	8	91	96	97	100	102	103	105	109	111	115	115	117	119	126	126	129	130	130	132	137	137	140	141	145
2	9	100	dead																						
2	10	91	93	90	95	91	93	92	99	101	106	107	107	108	114	115	116	116	119	121	121	122	124	125	127
2	11	89	89	87	88	83	86	86	91	93	96	99	103	105	110	111	112	113	118	120	121	124	126	127	133
2	12	98	99	98	109	106	108	116	117	121	103	123	125	126	125	130	131	133	135	138	142	147	148	151	155
2	13	98	104	102	106	109	110	116	120	123	126	127	129	132	145	139	140	142	144	147	149	154	150	156	160
2	14	101	102	98	98	97	98	106	110	112	117	119	120	122	128	128	129	129	132	135	125	138	139	140	144
2	15	78	76	75	74	77	79	81	82	84	86	88	88	89	92	93	99	95	99	101	101	103	106	106	109
2	16	90	dead																						
3	17	93	99	96	96	99	103	105	103	106	108	107	111	113	116	116	117	118	119	121	124	124	126	127	129
3	18	94	95	98	97	98	100	103	107	108	109	107	108	108	115	116	116	118	120	123	125	126	130	132	136
3	19	95	96	100	101	105	107	109	118	114	115	113	116	119	121	121	122	125	128	131	132	135	137	138	140
3	20	99	99	102	97	106	105	113	117	120	124	123	125	127	134	134	132	134	138	140	143	145	146	149	152
3	21	91	93	95	104	99	102	105	107	110	115	113	113	115	120	123	123	125	127	129	132	132	134	136	138
3	22	105	104	95	110	112	115	120	124	128	132	130	136	138	144	145	149	150	155	158	160	164	166	169	172
3	23	92	93	93	97	99	102	103	104	106	107	110	112	115	114	117	117	118	117	120	120	126	126	128	130
3	24	93	91	93	90	96	99	101	105	107	109	107	107	110	101	105	105	110	112	115	117	116	124	126	129

Animal weights from a first study (1/3)

Appendix 1: Animal Weights (continued)

Group	Animal	DAY																													
		-1	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
4	25	96	97	95	93	98	98	102	106	107	110	111	113	115	119	118	121	124	128	120	131	132	133	134	137	138	140	142	145	145	144
	26	89	89	88	90	93	92	97	99	101	104	103	102	103	110	108	109	110	112	117	111	117	119	119	121	128	125	126	127	130	128
	27	93	88	93	96	102	105	106	107	109	116	118	119	120	123	119	123	124	128	131	130	135	138	137	143	145	145	146	150	156	152
4	28	99	100	99	96	100	103	106	110	112	115	118	121	122	122	125	126	126	133	135	137	139	140	140	144	145	148	150	152	151	152
	29	90	91	90	96	100	102	100	101	104	107	107	108	110	112	111	117	119	121	124	125	129	131	126	133	134	136	140	141	140	142
	30	93	96	95	99	102	103	106	111	111	116	118	119	121	125	122	121	126	131	134	134	136	141	141	145	148	152	154	158	157	
4	31	93	94	89	91	95	98	99	103	105	108	109	112	115	116	114	117	120	123	125	128	127	131	130	132	132	134	136	138	130	139
	32	91	92	91	95	98	100	102	102	105	108	111	112	114	114	116	116	120	125	126	127	130	133	134	137	138	141	143	145	133	146
	33	74	dead																												
5	34	85	83	84	83	88	88	87	90	94	97	98	100	102	106	108	107	108	109	111	120	118	120	120	128	129	128	130	134	134	134
	35	96	98	98	100	101	102	100	103	108	111	115	115	118	119	118	120	119	123	125	129	129	130	129	134	137	137	138	140	141	142
	36	90	92	91	94	97	98	95	96	101	104	105	107	109	114	113	116	117	119	121	120	125	127	127	131	132	133	135	139	134	140
5	37	100	dead																												
	38	98	101	100	98	101	103	104	108	110	114	117	119	120	128	129	134	134	138	140	152	144	148	144	149	152	154	156	159	161	161
	39	87	84	82	79	78	79	81	84	86	91	93	94	99	97	98	100	103	108	108	117	113	118	115	121	122	123	124	130	132	130
6	40	89	82	79	76	79	82	81	86	87	91	94	98	100	101	102	104	103	104	105	118	113	112	111	115	117	118	119	125	128	124
	41	94	92	96	97	99	102	99	98	103	108	109	110	112	116	115	114	118	117	119	129	125	127	128	136	134	136	140	142	151	144
	42	94	95	93	96	98	101	104	103	109	111	112	114	118	119	119	120	121	124	127	133	132	133	133	141	140	142	143	146	149	150
6	43	101	101	101	107	109	110	115	117	122	126	127	129	130	134	134	135	138	141	142	150	150	154	155	160	160	164	165	172	172	172
	44	90	dead																												
	45	89	88	88	90	91	96	94	92	95	98	102	102	105	105	102	101	98	100	103	105	104	108	106	111	111	111	113	115	114	115
6	46	93	94	92	103	102	102	102	104	108	110	112	114	116	119	119	122	123	125	128	132	131	134	132	138	139	141	145	146	159	144
	47	99	101	98	97	106	109	111	111	119	118	119	123	126	125	126	130	131	135	137	138	140	143	142	147	150	152	153	158	141	160
	48	91	dead																												

Animal weights from a first study, continued (2/3)

Appendix 1: Animal Weights (continued)

Group	Animal	DAY																													
		-1	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
7	49	91	94	91	95	101	104	98	102	106	107	102	113	115	119	120	127	120	131	133	135	140	141	142	148	149	150	152	158	158	156
7	50	93	95	92	91	93	93	98	104	103	105	106	109	111	115	116	118	117	124	127	120	126	129	129	135	137	136	140	143	142	146
7	51	90	87	88	87	97	98	95	95	99	101	103	107	108	113	113	116	121	118	121	125	127	133	131	137	138	138	140	144	148	145
7	52	95	99	95	99	102	104	107	108	113	115	114	113	115	120	120	124	122	128	131	132	133	137	136	143	142	143	145	148	140	149
7	53	92	93	88	89	91	93	90	93	97	99	101	102	106	109	108	110	112	117	120	118	121	124	124	128	129	130	132	135	138	137
7	54	98	101	98	102	105	108	101	106	109	112	113	116	119	121	122	125	126	129	129	135	137	138	136	142	144	146	147	151	145	152
7	55	91	93	90	93	99	101	98	101	104	109	109	111	113	117	118	120	123	125	127	130	130	132	133	139	140	142	145	148	148	150
7	56	88	87	81	83	78	81	78	80	82	85	86	92	95	94	99	100	102	105	108	111	112	113	113	116	121	123	124	127	121	130
8	57	94	97	93	99	102	105	105	108	112	113	114	117	119	120	121	123	125	128	131	138	132	137	137	139	140	143	143	147	147	148
8	58	88	90	84	89	93	95	98	104	105	106	107	111	114	116	118	119	120	120	124	128	126	131	131	134	134	135	137	139	135	142
8	59	88	92	84	91	93	97	97	100	102	105	103	105	108	110	112	111	115	118	121	122	116	122	121	123	125	127	131	129	128	127
8	60	86	89	86	96	95	96	99	100	105	106	106	109	111	114	114	117	118	122	124	125	126	131	131	136	138	139	143	145	148	146
8	61	94	94	91	97	99	103	103	106	110	113	115	117	119	125	124	126	128	128	130	138	137	139	141	142	144	144	148	151	151	153
8	62	93	94	88	93	98	101	104	108	111	108	112	118	120	122	121	122	126	128	130	134	133	137	136	138	142	145	147	147	154	153
8	63	91	94	91	96	99	100	103	105	109	110	113	115	117	118	120	123	125	127	131	129	132	135	135	137	138	139	147	143	145	146
8	64	98	101	96	101	107	107	111	113	117	118	119	125	127	124	124	125	126	127	130	131	135	137	135	139	142	142	144	147	142	151
9	65	98	101	100	104	108	110	113	115	116	114	117	119	131	124	124	127	126	133	137	135	136	138	139	141	142	144	146	149	148	151
9	66	94	98	94	98	102	105	107	110	112	113	114	117	118	125	124	127	129	130	134	137	139	141	139	143	145	148	150	152	144	152
9	67	89	93	90	95	98	101	104	106	106	105	109	112	115	115	118	119	118	123	126	125	126	130	130	132	135	136	137	140	128	143
9	68	93	96	91	95	100	103	105	104	111	109	112	113	116	117	119	120	119	121	124	129	127	130	129	131	122	133	136	134	122	136
9	69	98	99	98	102	106	109	110	114	116	119	121	124	129	131	129	133	134	138	141	142	146	152	149	151	153	155	159	182	165	166
9	70	92	dead																												
9	71	91	92	89	94	96	100	102	105	106	104	107	110	111	116	120	122	122	125	128	128	131	143	134	137	139	139	141	145	142	147
9	72	95	96	95	96	101	105	106	109	111	111	114	114	116	116	120	120	120	121	124	133	126	129	125	129	132	154	137	136	138	140

Animal weights from a first study, continued (3/3)

Appendix 1: Animal Weights

Group	Animal	DAY																													
		-1	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
1	1	68	92	91	94	96	98	102	103	103	106	108	110	114	113	116	115	115	117	119	119	114	115	116	117	118	121	122	123	124	127
1	2	69	93	92	96	100	102	103	107	109	110	114	116	119	117	120	124	124	125	129	131	131	135	136	136	138	140	141	147	149	148
1	3	96	98	96	101	105	108	110	112	115	120	122	125	126	125	123	131	132	137	142	145	144	148	152	153	155	159	161	161	163	169
1	4	95	97	97	102	103	107	108	111	114	117	118	121	125	123	126	124	123	124	129	130	131	135	138	139	140	143	147	145	149	152
1	5	68	91	88	91	94	97	98	100	103	105	106	110	112	109	111	111	112	113	117	119	119	123	127	127	128	131	132	133	136	136
1	6	90	95	93	98	101	104	107	108	111	114	115	122	122	121	124	127	125	124	128	130	130	134	136	139	141	128	147	148	149	153
1	7	66	89	88	91	94	96	92	99	103	102	104	108	110	108	111	113	112	114	116	117	118	123	123	124	125	145	128	139	133	134
1	8	92	AOD																												
2	9	65	86	86	90	93	96	96	110	104	107	109	111	113	112	113	116	115	121	123	125	127	133	135	138	139	142	144	146	149	152
2	10	66	89	86	91	94	96	97	100	104	106	107	108	112	112	114	118	117	118	121	123	125	129	130	132	135	139	142	143	147	146
2	11	88	91	91	94	96	97	98	103	104	105	107	109	112	112	115	119	116	117	119	120	121	125	126	128	129	131	134	134	138	139
2	12	100	104	102	106	105	112	113	116	121	121	124	129	132	129	132	134	135	135	138	142	140	145	144	150	150	154	155	156	162	160
2	13	77	81	80	84	87	83	94	95	97	98	101	104	107	106	108	112	111	113	117	119	117	122	125	124	122	129	130	132	136	136
2	14	92	98	96	101	103	106	108	112	111	115	116	117	118	117	120	118	120	122	123	127	128	131	134	135	138	140	144	142	144	150
2	15	88	91	89	93	95	95	97	101	102	104	106	109	111	113	115	118	120	121	123	124	126	129	131	132	135	135	137	138	140	142
2	16	94	97	97	100	102	105	106	113	112	115	116	120	123	123	126	127	126	128	129	131	131	135	136	136	140	143	145	143	113	151
3	17	96	99	93	95	97	95	95	96	99	98	97	101	100	96	99	99	97	94	98	102	102	108	113	115	117	121	124	124	127	131
3	18	68	91	86	89	92	94	94	94	98	99	96	97	93	91	93	91	86	84	84	88	91	95	99	101	104	108	111	112	115	118
3	19	90	93	90	90	92	94	82	80	84	84	85	82	87	dead																
3	20	68	91	85	89	82	84	86	79	81	80	76	80	81	80	82	83	78	77	79	81	85	89	93	97	99	103	99	108	110	115
3	21	101	102	98	100	101	103	101	101	107	108	104	107	108	108	106	103	99	99	98	106	107	112	115	119	117	127	128	129	132	135
3	22	92	96	87	93	92	95	94	88	91	86	82	83	80	78	80	79	74	72	75	79	81	85	88	91	93	97	106	107	108	109
3	23	65	dead																												
3	24	92	95	91	95	97	99	99	100	104	103	100	101	100	98	100	102	99	101	105	108	110	114	117	121	123	127	130	132	135	138

Animal weights from a second study (1/2)

Appendix 1: Animal Weights (continued)

		DAY																													
Group	Animal	-1	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
4	25	86	90	88	91	93	95	99	97	102	105	106	109	105	110	114	115	116	119	124	123	123	129	131	133	133	137	138	140	145	146
4	26	92	95	93	99	102	103	106	109	112	113	116	121	120	119	123	123	124	129	134	134	133	141	144	145	143	150	151	152	156	160
4	27	85	87	86	89	90	94	94	96	99	102	119	97	107	105	108	110	110	111	114	116	114	118	121	123	124	126	130	131	133	136
4	28	94	98	96	101	103	106	107	111	113	115	119	120	127	125	127	129	127	129	132	132	135	138	142	143	143	128	148	158	148	154
4	29	94	96	96	99	101	104	107	110	112	117	117	123	122	121	123	125	125	130	132	135	135	142	142	145	149	151	153	157	160	161
4	30	91	94	95	99	102	104	104	100	109	113	115	117	119	118	119	122	120	126	127	130	127	134	136	139	139	142	142	144	146	149
4	31	93	85	84	88	89	93	94	99	97	98	100	105	105	104	106	111	109	112	114	116	117	120	123	124	122	129	130	130	133	135
4	32	74	74	75	79	79	82	86	87	92	94	95	103	98	95	98	107	104	108	111	112	112	116	119	122	126	125	125	124	126	132
5	33	83	83	83	89	89	98	95	98	101	104	105	115	110	108	111	112	111	114	118	119	120	124	124	124	128	131	132	138	141	140
5	34	79	80	79	83	86	88	90	93	95	102	99	101	104	103	106	111	110	114	118	119	121	123	127	126	128	132	133	134	137	139
5	35	83	87	85	90	93	95	97	94	101	102	103	105	106	105	109	120	113	114	116	118	119	124	126	125	128	130	132	137	134	138
5	36	84	86	86	91	93	97	100	102	107	111	110	109	115	118	114	120	121	124	127	130	132	135	137	139	144	145	148	150	152	156
5	37	85	85	85	88	89	91	93	102	99	101	102	106	108	107	111	110	108	111	115	117	116	121	123	124	124	129	129	131	134	136
5	38	78	79	79	82	84	88	88	97	95	101	99	103	105	102	104	107	108	110	114	117	118	122	125	125	128	131	134	133	137	139
5	39	85	87	87	90	94	94	99	101	104	106	107	111	112	111	113	113	118	121	125	125	127	130	132	132	134	136	137	146	142	148
5	40	81	83	82	87	89	90	93	95	97	100	101	107	107	107	109	112	110	113	116	118	118	121	123	124	125	128	128	130	133	135
6	41	86	87	81	85	93	97	78	82	85	89	91	96	98	100	102	106	107	110	112	116	117	120	121	123	125	128	128	130	136	136
6	42	91	93	88	93	95	99	102	107	112	114	116	120	122	120	123	118	117	115	108	106	104	104	103	104	104	106	108	108	109	110
6	43	84	87	84	83	79	78	84	93	89	93	95	98	102	103	107	109	108	112	115	115	116	122	124	124	125	127	130	129	131	134
6	44	91	94	92	92	93	94	96	101	104	106	108	113	114	114	119	120	120	123	125	127	128	132	133	135	137	137	138	136	139	142
6	45	78	dead																												
6	46	86	89	dead																											
6	47	89	93	86	91	79	91	92	99	101	105	109	112	114	114	116	124	124	127	129	133	133	137	137	138	138	141	143	142	150	146
6	48	88	92	87	89	92	86	90	97	100	109	105	111	114	114	118	122	122	126	129	129	131	135	137	138	139	142	145	143	146	149

Animal weights from a second study (2/2)

WO 2008/045559

PCT/US2007/021897

Appendix 2: Mucositis Scores

Cheek pouch photographs were scored by 2 independent scorers in a blinded manner,
resulting in 2 scores for each animal at each time point.

ACT-03 Blinded Scores													
DAY													
Group	Animal	6	8	10	12	14	16	18	20	22	24	26	28
1	1	0	0	0	2	3	3	3	3	3	2	2	1
1	1	0	0	0	1	3	3	3	3	3	2	1	1
1	2	1	0	1	2	3	3	4	3	3	3	2	2
1	2	0	0	0	2	3	3	3	3	3	3	2	1
1	3	1	0	1	2	3	4	3	3	3	2	2	2
1	3	0	0	0	1	3	4	3	3	3	1	2	1
1	4	1	0	0	2	3	3	3	4	2	2	2	2
1	4	0	0	0	1	3	3	3	3	2	2	2	1
1	5	0	1	0	2	3	3	3	3	3	2	2	2
1	5	0	0	1	2	3	3	3	3	3	1	1	1
1	6	0	0	1	2	3	3	4	3	2	2	2	2
1	6	0	0	1	2	3	3	3	3	2	2	2	1
1	7	0	0	1	2	3	3	2	2	1	1	1	1
1	7	0	0	1	2	3	3	1	1	1	1	1	1
1	8	1	0	1	1	2	3	3	3	3	2	2	2
1	8	0	0	1	0	2	3	3	3	3	2	2	1
2	9	dead											
2	9												
2	10	0	0	1	2	2	3	2	2	1	1	1	1
2	10	0	0	1	1	2	3	2	2	1	1	1	1
2	11	0	0	1	2	3	3	3	3	3	2	1	1
2	11	0	0	0	1	3	3	3	3	3	2	2	2
2	12	0	0	1	1	3	3	3	3	2	2	2	1
2	12	0	0	0	1	3	3	3	3	2	1	2	1
2	13	0	0	1	2	3	3	3	3	3	2	2	1
2	13	0	1	0	2	3	3	3	3	3	2	2	1
2	14	0	0	1	2	2	3	3	3	2	2	1	0
2	14	1	1	0	1	2	3	3	3	2	2	2	1
2	15	0	0	1	2	2	2	2	2	2	1	1	1
2	15	0	0	0	1	2	2	2	2	2	1	1	2
2	16	dead											
2	16												
3	17	0	0	1	1	1	3	3	3	2	2	1	0
3	17	1	0	1	1	2	3	3	3	2	2	2	1
3	18	0	0	1	1	2	3	3	3	3	3	2	0
3	18	1	1	1	1	2	3	3	3	3	3	2	1
3	19	0	0	1	2	3	3	3	3	2	2	2	0
3	19	1	1	0	2	3	3	3	3	2	2	1	2
3	20	0	0	1	1	2	3	4	3	2	1	2	1
3	20	0	0	1	2	2	3	3	3	2	1	1	2
3	21	0	0	0	1	2	2	2	2	2	1	2	0
3	21	0	0	0	1	2	2	2	2	2	2	2	1
3	22	0	0	1	2	1	2	2	2	1	1	1	0
3	22	0	1	0	1	2	2	2	2	2	2	1	1
3	23	0	0	0	3	3	4	4	3	2	2	2	0
3	23	0	1	0	3	3	4	4	3	2	2	1	1
3	24	0	0	1	2	3	3	3	3	2	1	2	0
3	24	1	1	0	2	3	3	3	3	2	1	1	1

Mucositis scores from a first study (1/3)

WO 2008/045559

PCT/US2007/021897

Appendix 2: Mucositis Scores (continued)

Cheek pouch photographs were scored by 2 independent scorers in a blinded manner, resulting in 2 scores for each animal at each time point.

Group	Animal	DAY											
		6	8	10	12	14	16	18	20	22	24	26	28
4	25	0	0	1	2	1	1	1	1	1	1	2	1
4	25	0	0	0	2	1	1	1	1	2	2	1	2
4	26	0	0	0	0	3	3	3	3	2	1	2	1
4	26	0	0	0	0	3	3	3	3	2	2	1	2
4	27	0	0	1	2	2	3	3	3	3	2	2	0
4	27	0	1	0	2	2	4	3	3	3	2	2	1
4	28	0	0	2	2	3	3	3	3	3	2	2	1
4	28	0	0	2	2	3	3	3	3	3	2	2	2
4	29	0	0	1	2	3	3	4	3	2	2	2	0
4	29	0	0	0	1	3	4	3	3	2	2	1	1
4	30	0	0	1	2	3	3	3	3	2	2	2	0
4	30	0	0	0	1	3	4	3	3	2	2	2	2
4	31	0	0	1	2	3	3	2	2	2	2	2	1
4	31	0	0	0	2	3	3	2	2	2	2	1	2
4	32	0	0	1	2	3	3	3	3	3	2	2	1
4	32	0	0	0	2	3	4	3	3	3	2	1	2
5	33	dead											
5	33												
5	34	0	0	0	2	3	3	3	2	2	1	1	0
5	34	0	0	0	2	3	3	3	2	2	2	1	1
5	35	0	0	1	2	3	3	3	3	2	2	2	1
5	35	0	0	0	2	3	4	3	3	2	2	1	2
5	36	0	0	1	2	2	2	2	2	0	1	1	1
5	36	0	0	0	1	2	2	2	2	1	1	0	2
5	37	dead											
5	37												
5	38	0	0	0	2	2	2	2	2	2	0	2	0
5	38	1	0	0	1	2	2	1	2	1	0	1	1
5	39	0	0	0	1	3	3	3	3	3	1	1	0
5	39	0	1	0	0	3	3	3	3	3	0	1	1
5	40	1	0	2	1	2	3	3	3	3	2	2	0
5	40	0	1	1	1	2	3	3	3	3	2	2	1
6	41	0	0	1	1	3	3	3	3	3	2	2	0
6	41	1	0	1	0	3	3	3	3	3	2	2	1
6	42	0	0	1	2	3	3	3	4	2	2	2	1
6	42	0	0	1	1	3	3	3	3	2	2	1	2
6	43	0	0	0	2	3	3	2	2	2	2	2	1
6	43	1	0	0	2	3	3	2	2	2	1	2	2
6	44	dead											
6	44												
6	45	0	0	1	2	2	4	4	4	4	4	3	2
6	45	1	0	0	2	2	4	4	4	4	4	3	2
6	46	0	0	0	2	3	3	3	3	2	2	2	0
6	46	0	0	0	2	3	3	3	3	2	2	2	1
6	47	0	0	1	2	2	3	2	2	0	2	2	2
6	47	0	1	0	0	2	3	2	2	1	2	1	1
6	48	dead											
6	48												

Mucositis scores from a first study (2/3)

WO 2008/045559

PCT/US2007/021897

Appendix 2: Mucositis Scores (continued)

Cheek pouch photographs were scored by 2 independent scorers in a blinded manner, resulting in 2 scores for each animal at each time point.

		DAY											
Group	Animal	6	8	10	12	14	16	18	20	22	24	26	28
7	49	0	0	1	2	2	1	1	1	0	1	1	1
7	49	0	1	0	2	2	1	0	0	1	0	1	1
7	50	0	0	0	2	2	2	3	1	1	0	0	0
7	50	1	0	0	2	2	1	3	0	2	1	1	1
7	51	0	0	1	1	2	3	1	2	1	0	0	0
7	51	1	0	0	0	2	3	2	1	2	0	1	1
7	52	0	0	0	2	2	2	2	2	1	0	0	0
7	52	1	1	0	2	2	2	1	0	2	1	1	1
7	53	0	0	1	1	2	3	3	3	2	1	1	0
7	53	0	1	0	0	2	3	3	3	2	2	1	1
7	54	0	0	2	2	2	3	3	3	2	0	1	0
7	54	0	0	0	2	2	3	3	3	1	2	1	2
7	55	0	0	0	2	3	2	2	2	2	1	2	0
7	55	0	0	0	2	3	2	2	1	2	2	1	2
7	56	0	0	1	2	2	2	1	1	1	2	2	0
7	56	0	0	1	1	2	2	1	1	1	2	1	1
8	57	0	0	0	2	3	3	3	3	1	1	0	1
8	57	0	1	0	2	3	3	3	3	1	2	0	2
8	58	1	0	0	2	2	2	2	2	1	1	1	1
8	58	1	0	0	2	2	2	2	2	1	2	0	1
8	59	0	0	1	2	1	1	1	1	1	1	2	0
8	59	1	0	0	2	1	2	1	2	1	1	2	1
8	60	0	0	1	1	3	3	3	1	1	0	0	0
8	60	0	0	0	1	3	3	3	2	1	2	0	1
8	61	0	0	1	2	2	2	3	2	1	1	1	0
8	61	0	0	0	2	2	2	3	1	1	1	1	1
8	62	0	0	0	2	3	2	1	2	2	0	1	0
8	62	0	0	0	1	3	2	2	1	2	2	0	1
8	63	0	0	1	2	2	2	1	2	2	0	1	0
8	63	1	0	1	2	2	2	1	1	2	2	0	1
8	64	0	0	1	2	3	3	3	3	2	0	2	0
8	64	1	0	0	2	3	3	3	3	2	1	0	1
9	65	0	0	1	2	2	3	3	2	2	0	2	1
9	65	1	0	0	2	2	3	3	2	2	1	1	2
9	66	0	0	0	2	2	2	2	2	0	1	1	0
9	66	1	0	0	1	1	1	1	1	1	2	0	1
9	67	0	0	0	2	1	1	2	1	0	0	1	0
9	67	1	0	0	1	1	1	1	1	1	1	0	1
9	68	1	1	2	2	2	3	3	2	1	1	2	0
9	68	1	0	0	1	2	3	3	1	2	2	1	1
9	69	0	0	0	2	3	3	3	3	3	3	2	1
9	69	1	1	0	1	3	3	3	3	3	3	1	2
9	70												
9	70												
9	71	0	0	1	2	2	2	2	2	2	1	2	0
9	71	0	0	0	2	2	2	2	2	2	2	1	2
9	72	0	0	0	2	3	4	3	1	2	2	2	1
9	72	0	0	0	2	3	3	3	2	2	1	1	2

Mucositis scores from a first study (3/3)

Mucositis scores from a second study (1/2)

Appendix 2: Mucositis Scores (continued)
Cheek pouch photographs were scored by 2 independent scorers in a blinded manner, resulting in 2 scores for each animal at each time point.


ACT-02 Blinded Scores													
Group	Animal	6	8	10	12	14	16	18	20	22	24	26	28
4	25	1	0	1	2	3	3	3	3	3	3	3	3
4	25	1	0	1	2	3	3	3	3	3	3	3	3
4	26	1	1	2	3	3	3	3	3	3	2	2	2
4	26	1	1	2	3	3	3	3	3	3	2	2	2
4	27	0	1	1	2	3	3	3	3	3	3	2	2
4	27	0	1	1	2	3	3	3	3	3	3	2	2
4	28	0	0	1	1	3	3	3	3	3	1	1	1
4	28	0	0	1	1	3	3	3	3	3	1	1	1
4	29	0	0	1	3	3	3	3	3	2	1	1	1
4	29	0	0	1	3	3	3	3	3	2	1	1	1
4	30	1	1	1	2	3	2	3	3	2	2	2	1
4	30	1	1	1	2	3	3	3	3	2	2	2	1
4	31	1	0	1	2	3	2	2	2	2	2	2	1
4	31	1	0	1	2	3	2	2	2	2	2	2	1
4	32	0	1	2	2	2	2	1	1	2	3	3	2
4	32	0	1	2	2	2	2	1	1	2	3	3	2
5	33	0	1	1	2	3	3	3	3	3	2	1	1
5	33	0	0	1	2	3	3	3	3	3	2	1	1
5	34	0	0	0	1	1	1	1	1	3	2	2	2
5	34	0	1	0	1	1	1	1	1	3	2	2	2
5	35	1	0	0	2	3	3	3	3	3	3	2	1
5	35	1	0	0	2	3	3	3	3	3	3	2	1
5	36	1	1	0	2	3	3	3	3	3	3	3	1
5	36	1	1	1	2	3	3	3	3	3	3	3	1
5	37	0	1	1	2	3	3	3	3	2	3	2	1
5	37	0	1	0	2	3	3	3	3	2	3	2	1
5	38	0	0	0	3	3	3	3	3	2	2	2	1
5	38	0	0	0	3	3	3	3	3	2	2	2	1
5	39	0	0	0	1	1	2	1	1	2	1	1	1
5	39	0	0	0	1	1	2	1	1	2	1	1	1
5	40	0	1	1	1	2	3	3	3	1	1	1	1
5	40	0	1	1	1	2	3	3	3	1	1	1	1
6	41	1	1	1	2	3	3	3	2	2	1	1	1
6	41	1	1	1	2	3	3	3	2	2	1	1	1
6	42	1	0	1	1	3	3	3	2	1	1	1	1
6	42	1	0	1	1	3	3	3	2	2	1	1	1
6	43	0	1	1	2	3	3	2	2	2	1	1	1
6	43	0	0	1	2	3	3	2	2	1	1	1	1
6	44	0	0	1	2	3	3	2	1	1	1	1	1
6	44	0	0	1	2	3	3	2	1	1	1	1	1
6	45												
6	45												
6	46												
6	46												
6	47	0	0	1	2	3	1	1	1	1	1	1	1
6	47	0	0	1	2	3	1	1	1	1	1	1	1
6	48	1	1	1	2	3	1	1	1	1	1	1	1
6	48	1	1	1	2	3	1	1	1	1	1	1	1

Mucositis scores from a second study (2/2)

75975-31

CLAIMS:

1. Use of a superoxide dismutase mimetic to lessen severity of an oral mucositis in a mammal, wherein the superoxide dismutase mimetic is for parenteral administration prior to or simultaneous with a cancer treatment, and the superoxide
- 5 dismutase mimetic is a compound of the formula:



wherein

W is a substituted or unsubstituted pyridino moiety;

U and V are trans-cyclohexanyl fused rings;

- 10 R₁, R₂, R'₂, R₃, R₄, R₅, R'₅, R₆, R'₆, R₇, R₈, R₉, R'₉, and R₁₀ are each independently selected from the group consisting of hydrogen, alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, heterocyclyl, and aralkyl;

- 15 M is a cation of a transition metal;

and X, Y and Z are independently selected from the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxo, hydroperoxo, alkylperoxo, arylperoxo, ammonia, alkylamino, arylamino, heterocycloalkyl amino,

75975-31


heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, and thiotosylate;

and n is an integer from 0 to 3.

2. The use in accordance with claim 1, wherein M is selected from the group consisting of Mn^{2+} , Mn^{3+} , Fe^{2+} , and Fe^{3+} .
3. The use in accordance with claim 1, wherein M is Mn^{2+} or Mn^{3+} .
4. The use in accordance with claim 1, where W is an unsubstituted pyridino moiety.

75975-31

5. The use in accordance with claim 1, wherein the superoxide dismutase mimetic is a compound represented by the formula:




6. The use in accordance with claim 1, wherein the mammal is a human
5 patient.

7. The use in accordance with claim 6, wherein R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are each hydrogen.

8. The use in accordance with claim 6, wherein R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are each hydrogen and W is an unsubstituted pyridino
10 moiety.

9. Use of a superoxide dismutase mimetic to lessen severity of an oral mucositis in a mammal, wherein the superoxide dismutase mimetic is for parenteral administration prior to or simultaneous with a radiation therapy treatment, and the superoxide dismutase mimetic is a compound of the formula:

75975-31



wherein

W is a substituted or unsubstituted pyridino moiety;

U and V are trans-cyclohexanyl fused rings;

5 $R_1, R_2, R'_2, R_3, R_4, R_5, R'_5, R_6, R'_6, R_7, R_8, R_9, R'_9,$ and R_{10} are each independently selected from the group consisting of hydrogen, alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, heterocyclyl, and aralkyl;

10 M is a cation of a transition metal;

and X, Y and Z are independently selected from the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxo, hydroperoxo, alkylperoxo, arylperoxo, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric
 15 oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl

75975-31

carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, and thiotosylate;

and n is an integer from 0 to 3.

10. The use in accordance with claim 9, wherein the superoxide dismutase mimetic is for administration on the day before or the day of, but prior to the radiation therapy treatment.


11. The use in accordance with claim 9, wherein M is selected from the group consisting of Mn^{2+} , Mn^{3+} , Fe^{2+} , and Fe^{3+} .

12. The use in accordance with claim 9, wherein M is Mn^{2+} or Mn^{3+} .

13. The use in accordance with claim 9, where W is an unsubstituted pyridino moiety.

14. The use in accordance with claim 9, wherein the superoxide dismutase mimetic is a compound represented by the formula:

75975-31




15. The use in accordance with claim 9, wherein the mammal is a human patient.

16. The use in accordance with claim 9, wherein R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are each hydrogen.

17. The use in accordance with claim 15, wherein R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are each hydrogen and W is an unsubstituted pyridino moiety.

18. Use of a superoxide dismutase mimetic to lessen severity of an oral mucositis in a mammal, wherein the superoxide dismutase mimetic is for parenteral administration prior to or simultaneous with a chemotherapy treatment, and the superoxide dismutase mimetic is a compound of the formula:



75975-31

wherein

W is a substituted or unsubstituted pyridino moiety;

U and V are trans-cyclohexanyl fused rings;

R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are each
 5 independently selected from the group consisting of hydrogen, alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, heterocyclyl, and aralkyl;

M is a cation of a transition metal;

10 and X, Y and Z are independently selected from the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxo, hydroperoxo, alkylperoxo, arylperoxo, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl
 15 nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea,
 20 thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid,
 25 aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl

75975-31

dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate,
 5 tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, and thiotosylate;

and n is an integer from 0 to 3.

19. The use in accordance with claim 18, wherein the superoxide dismutase mimetic is for administration prior to the chemotherapy treatment.

10 20. The use in accordance with claim 18, wherein the superoxide dismutase mimetic is for administration of less than 3 mg/kg body weight of the superoxide dismutase mimetic to the mammal.

21. The use in accordance with claim 18, wherein the superoxide dismutase mimetic is for administration prior to the chemotherapy treatment, and is
 15 for administration of less than 3 mg/kg body weight of the superoxide dismutase mimetic to the mammal.


22. The use in accordance with claim 18, wherein M is selected from the group consisting of Mn^{2+} , Mn^{3+} , Fe^{2+} , and Fe^{3+} .

23. The use in accordance with claim 18, wherein M is Mn^{2+} or Mn^{3+} .

20 24. The use in accordance with claim 18, where W is an unsubstituted pyridino moiety.

25. The use in accordance with claim 18, wherein the superoxide dismutase mimetic is a compound represented by the formula:

75975-31



26. The use in accordance with claim 18, wherein the mammal is a human patient.

27. The use in accordance with claim 18, wherein R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 ,
5 R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are each hydrogen.

28. The use in accordance with claim 26, wherein R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 ,
 R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are each hydrogen and W is an unsubstituted pyridino moiety.

29. The use in accordance with claim 1, wherein the superoxide dismutase
10 mimetic is for administration on the day before or the day of, but prior to the cancer treatment.

30. The use in accordance with claim 18, wherein the superoxide dismutase mimetic is for administration on the day before or the day of, but prior to the chemotherapy treatment.

15 31. The use in accordance with claim 1, wherein the superoxide dismutase mimetic is for further administration for up to three days after the cancer therapy treatment.


32. The use in accordance with claim 1, wherein the superoxide dismutase mimetic is for administration at least 30 minutes prior to the cancer therapy treatment.

75975-31

33. The use in accordance with claim 9, wherein the superoxide dismutase mimetic is for further administration for up to three days after the radiation therapy treatment.

34. The use in accordance with claim 9, wherein the superoxide dismutase
5 mimetic is for administration at least 30 minutes prior to the radiation therapy treatment.

FIG. 1

Figure 1. Study timeline.

2/12
FIG. 2

A. Score of 0. The pouch is completely healthy with no erythema or vasodilation.



D. Score of 3. Formation of ulcer occupying $\frac{1}{4}$ or less of the pouch.



B. Score of 1. Light to severe erythema and vasodilation with no erosion of the mucosa.



E. Score of 4. Total size of ulcer occupies $\frac{1}{2}$ of the pouch with loss of pliability.





C. Score of 2. Severe erythema and vasodilation with erosion and decreased stippling of superficial mucosa leaving denuded areas.



F. Score of 5. Ulceration to virtually all of the pouch. Pouch can only be partially everted.


Figure 2. ACT-02. Mucositis Score Scale.

3/12
FIG. 3

4/12
FIG. 4

5/12

A. Intraperitoneal Dosing



B. Topical Dosing




FIG. 5



6/12
FIG. 6


FIG. 7

Study Timeline






A. Score of 0. The pouch is completely healthy with no erythema or vasodilation.



D. Score of 3. Formation of ulcer occupying $\frac{1}{4}$ or less of the pouch.



B. Score of 1. Light to severe erythema and vasodilation with no erosion of the mucosa.



E. Score of 4. Total size of ulcer occupies $\frac{1}{2}$ of the pouch with loss of pliability.



C. Score of 2. Severe erythema and vasodilation with erosion and decreased stippling of superficial mucosa leaving denuded areas.



F. Score of 5. Ulceration to virtually all of the pouch. Pouch can only be partially everted.

Figure 2. ACT-03. Mucositis Score Scale.

FIG. 8







FIG. 9

Note: Y axes are mean % weight gain (-20 to 80%)

10/12
FIG. 10

11/12

A



B




FIG. 11

12/12




FIG. 12

Study timeline.

