

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 June 2011 (03.06.2011)

(10) International Publication Number
WO 2011/063947 A1

PCT

(51) International Patent Classification:

A01N 37/42 (2006.01) *A01N 43/828* (2006.01)
A01N 43/653 (2006.01) *A01P 21/00* (2006.01)

(21) International Application Number:

PCT/EP2010/007129

(22) International Filing Date:

24 November 2010 (24.11.2010)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0920892.7 27 November 2009 (27.11.2009) GB

(71) Applicant (for all designated States except US): **SYNGENTA PARTICIPATIONS AG** [CH/CH];
Schwarzwaldallee 215, CH-4058 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HAAS, Ulrich Johannes** [DE/DE]; Syngenta Crop Protection AG, Werk Rosental, Schwarzwaldallee 215, CH-4058 Basel (CH).
HARP, Tyler L. [US/CH]; Syngenta Crop Protection AG, Schaffhauserstrasse, CH-4332 Stein (CH).

(74) Agents: **SYNGENTA LIMITED** et al.; Intellectual Property Department, PO Box 3538, Jealott's Hill Research centre, Bracknell RG42 6YA (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: PLANT GROWTH REGULATION

(57) Abstract: The present invention relates to a method for improving the plant growth regulation of and/or enhancing crop plants, by applying to the plants a mixture of a plant growth regulator, and acibenzolar-S-methyl, and to a composition comprising the same.



WO 2011/063947 A1

PLANT GROWTH REGULATION

The present invention relates to a method for improving the plant growth regulation of crop
5 plants, by applying to the crop plants a mixture of a plant growth regulator and acibenzolar-S-
methyl, and to a composition comprising the same.

Plant growth regulators are often used to regulate the growth and development of crop plants.
For example, plant growth regulators are used to slow the development of a crop (such as oil
10 seed rape) so that it flowers at a desired time, reduce the height of a crop (such as in cereals)
so that it is less susceptible to lodging, increase nitrogen efficiency, regulate flowering and
fruit set of a crop (such as fruit trees), and slow turfgrass growth rate to reduce mowing
frequency.

15 There are several different classes of plant growth regulator. Known classes include azoles
(such as uniconazole, and paclobutrazol), cyclohexane carboxylates (such as trinexapac-ethyl,
and prohexadione-calcium), pyrimidinyl carbinols (such as flurprimidol, and ancymidol),
quarternary ammoniums (such as chlormequat-chloride, and mepiquat-chloride), and
sulphonyl-amino phenyl-acetamides (such as mefluidide).

20 Plant growth regulators operate by various modes of action. For example, onium-type plant
growth retardants such as chlormequat-chloride and mepiquat-chloride, that possess a
positively charged ammonium, phosphonium or sulphonium group, function by blocking the
synthesis of gibberellin early in the biosynthetic pathway. Growth retardants comprising a
25 nitrogen-containing heterocycle, such as flurprimidol, paclobutrazol and uniconazole-P, act
as inhibitors of monooxygenases that catalyse oxidative steps in gibberellin biosynthesis.
Structural mimics of 2-oxoglutaric acid, such as the acylcyclohexanediones trinexapac-ethyl
and prohexadione-calcium, interfere with the late steps of gibberellin biosynthesis. Other
plant growth regulators, such as mefluidide, inhibit cell division and differentiation.

30 Plant growth regulators such as trinexapac-ethyl are commonly used on crops to reduce the
risk of lodging through stem thickening and shortening, and improved rooting..

In some cases, active ingredients have been shown to be more effective when mixed with other active ingredients compared to when applied individually, and this is referred to as “synergism”, since the combination demonstrates a potency or activity level exceeding that which it would be expected to have based on knowledge of the individual potencies of the components.

International patent publication number WO2008/020872 relates to mixtures of plant growth activators and plant activators for plant stress tolerance, manipulating seed germination or controlling disease in plants, primarily via seed treatment. There is no mention in WO2008/020872 that such mixtures could have a synergistic plant growth regulation effect.

The present invention resides in the discovery that plant growth regulators exhibit an improved plant growth regulation effect when applied in combination with acibenzolar-S-methyl.

The present invention is also useful in providing a plant growth regulation effect in crop plants that is as good as, or better than, that achieved using existing plant growth regulating products, using a lower concentration of any one plant growth regulator. This enables the composition to be applied at an earlier growth stage of the crop plants without causing phytotoxicity.

According to the present invention, there is provided a method for regulating the growth of crop plants, comprising applying to the plants, plant parts, plant propagation material, or a plant growing locus, a plant growth regulator and acibenzolar-S-methyl in a synergistically effective amount.

The term ‘regulating the growth’ includes restricting shoot growth, promoting root growth, stunting, and the like.

The term ‘plants’ refers to all physical parts of a plant, including seeds, seedlings, saplings, roots, tubers, stems, stalks, foliage and fruits.

The term 'plant propagation material' denotes generative parts of the plant, such as seeds, which can be used for the multiplication of the latter, and vegetative material, such as cuttings or tubers, for example potatoes. In particular, it includes seeds (in the strict sense), roots, fruits, tubers, bulbs, rhizomes and parts of plants. Germinated plants and young plants
5 which are to be transplanted after germination or after emergence from the soil, may also be mentioned - these young plants may be protected before transplantation by a total or partial treatment by immersion. Suitably "plant propagation material" is understood to denote seeds.

The term 'plant growing locus' is intended to embrace the place on which the plants are
10 growing, where the plant propagation materials of the plants are sown or where the plant propagation materials of the useful plants will be placed into the soil. An example for such a locus is a field, on which crop plants are growing.

The present invention also provides a method for enhancing crop plants, comprising applying
15 to the plants, plant parts, plant propagation material, or a plant growing locus, a plant growth regulator and acibenzolar-S-methyl in a synergistically effective amount.

According to the present invention, 'enhancing crop plants' means improving plant quality and/or plant vigour and/or tolerance to stress factors, any of which may lead to increased
20 yield. In one embodiment, the present invention relates to a method for improving plant yield, comprising applying to the plant, plant part, plant propagation material, or a plant growing locus, a plant growth regulator and acibenzolar-S-methyl. Such improved yield may be as a result of improved root growth. In a further embodiment, the present invention relates to a method for improving plant vigour and/or plant quality, and/or plant tolerance to
25 stress factors, comprising applying to the plant, plant part, plant propagation material, or a plant growing locus, a plant growth regulator and acibenzolar-S-methyl.

According to the present invention, an improvement in plant vigour means that certain traits are improved qualitatively or quantitatively when compared with the same trait in a control
30 plant which has been grown under the same conditions in the absence of the method of the invention. Such traits include, but are not limited to, early and/or improved germination, improved emergence, the ability to use less seeds, increased root growth, a more developed root system, increased shoot growth, increased tillering, stronger tillers, more productive

tillers, increased or improved plant stand, less plant verse (lodging), an increase and/or improvement in plant height, an increase in plant weight (fresh or dry), bigger leaf blades, greener leaf colour, increased pigment content, increased photosynthetic activity, earlier flowering, homogenous flowering, longer panicles, early grain maturity, increased seed, fruit
5 or pod size, increased pod or ear number, increased seed number per pod or ear, increased seed mass, enhanced seed filling, less dead basal leaves, delay of senescence, improved vitality of the plant and/or less inputs needed (e.g. less fertiliser, water and/or labour needed). A plant with improved vigour may have an increase in any of the aforementioned traits or any combination or two or more of the aforementioned traits. Suitably, the method of the present
10 invention increases plant height, plant weight and/or provides enhanced germination.

According to the present invention, an improvement in plant quality means also that certain traits are improved qualitatively or quantitatively when compared with the same trait in a control plant which has been grown under the same conditions in the absence of the method
15 of the invention. Such traits include, but are not limited to, improved visual appearance of the plant (e.g. improved colour, density, uniformity, compactness), reduced ethylene (reduced production and/or inhibition of reception), improved quality of harvested material, e.g. seeds, fruits, leaves, vegetables (such improved quality may manifest as improved visual appearance of the harvested material, improved carbohydrate content (e.g. increased
20 quantities of sugar and/or starch, improved sugar acid ratio, reduction of reducing sugars, increased rate of development of sugar), improved protein content, improved oil content and composition, improved nutritional value, reduction in anti-nutritional compounds, improved organoleptic properties (e.g. improved taste) and/or improved consumer health benefits (e.g. increased levels of vitamins and anti-oxidants)), improved post-harvest characteristics (e.g.
25 enhanced shelf-life and/or storage stability, easier processability, easier extraction of compounds) and/or improved seed quality (e.g. for use in following seasons). A plant with improved quality may have an increase in any of the aforementioned traits or any combination or two or more of the aforementioned traits.

30 According to the present invention, an improved tolerance to stress factors means that certain traits are improved qualitatively or quantitatively when compared with the same trait in a control plant which has been grown under the same conditions in the absence of the method of the invention. Such traits include, but are not limited to, an increased tolerance and/or

resistance to abiotic stress factors which cause sub-optimal growing conditions such as drought (e.g. any stress which leads to a lack of water content in plants, a lack of water uptake potential or a reduction in the water supply to plants), cold exposure, heat exposure, osmotic stress, UV stress, flooding, increased salinity (e.g. in the soil), increased mineral exposure, ozone exposure, high light exposure and/or limited availability of nutrients (e.g. nitrogen and/or phosphorus nutrients). A plant with improved tolerance to stress factors may have an increase in any of the aforementioned traits or any combination or two or more of the aforementioned traits. In the case of drought and nutrient stress, such improved tolerances may be due to, for example, more efficient uptake, use or retention of water and nutrients. Suitably, the method of the present invention increases tolerance of plants to drought.

Any or all of the above crop enhancements may lead to an improved yield by improving e.g. plant physiology, plant growth and development and/or plant architecture. In the context of the present invention 'yield' includes, but is not limited to, (i) an increase in biomass production, grain yield (e.g. grain size, grain number, grain density), starch content, oil content and/or protein content, which may result from (a) an increase in the amount produced by the plant *per se* or (b) an improved ability to harvest plant matter, (ii) an improvement in the composition of the harvested material (e.g. improved sugar acid ratios, improved oil composition, increased nutritional value, reduction of anti-nutritional compounds, increased consumer health benefits) and/or (iii) an increased/facilitated ability to harvest the crop, improved processability of the crop and/or better storage stability/shelf life. Increased yield of an agricultural plant means that, where it is possible to take a quantitative measurement, the yield of a product of the respective plant is increased by a measurable amount over the yield of the same product of the plant produced under the same conditions, but without application of the present invention. According to the present invention, it is preferred that the yield be increased by at least 0.5%, more preferred at least 1%, even more preferred at least 2%, still more preferred at least 4% , preferably 5% or even more.

Any or all of the above crop enhancements may also lead to an improved utilisation of land, i.e. land which was previously unavailable or sub-optimal for cultivation may become available. For example, plants which show an increased ability to survive in drought conditions, may be able to be cultivated in areas of sub-optimal rainfall, e.g. perhaps on the fringe of a desert or even the desert itself.

The term 'synergistically effective amount' indicates the quantity of such compounds which is capable of modifying the effect on the growth of plants, where said effect is greater than the sum of the effects obtained by applying each of the compounds individually.

5

Acibenzolar-S-methyl is a plant activator that stimulates a systemic acquired resistance (SAR) response in plants, so that they are better able to fight biotic and abiotic stress.

Any plant growth regulator may be used in accordance with the present invention. A complete list of all commercially available plant growth regulators may be obtained from the Pesticide Manual (14th edition, published by the British Crop Protection Council). In one embodiment, the plant growth regulator is selected from the group consisting of trinexapac-ethyl, prohexadione-calcium, paclobutrazol, uniconazole, flurprimidol, mefluidide, mepiquat-chloride, chlormequat-chloride, and a mixture thereof.

15

Suitably, the plant growth regulator is a gibberellin biosynthesis inhibitor. Suitably, the plant growth regulator is a class A gibberellin biosynthesis inhibitor. Suitably, the plant growth regulator is a class B gibberellin biosynthesis inhibitor. In a preferred embodiment the plant growth regulator is trinexapac-ethyl, prohexadione-calcium or chlormequat-chloride. In one embodiment, the plant growth regulator is trinexapac-ethyl. In one embodiment, the plant growth regulator is prohexadione-calcium. In one embodiment, the plant growth regulator is chlormequat-chloride. In one embodiment, the plant growth regulator is paclobutrazol. In one embodiment, the plant growth regulator is flurprimidol.

25 If desired, it is possible to use more than one plant growth regulator in combination, in accordance with the present invention, such as mixtures of trinexapac-ethyl and paclobutrazol.

In the present invention, the mixture ratio of plant growth regulator to acibenzolar-S-methyl at which the growth regulation effect is synergistic lies within the range from about 1:1000 to about 1000:1 by weight. Suitably, the mixture ratio of plant growth regulator to acibenzolar-S-methyl is from about 1:100 to about 100:1 by weight. More suitably, the mixture ratio of plant growth regulator to acibenzolar-S-methyl is from about 10:1 to about 1:1 by weight.

30

For example, when the plant growth regulator is trinexapac-ethyl, a mixture ratio of trinexapac-ethyl to acibenzolar-S-methyl from about 5:1 to about 2:1: by weight is preferred.

5 The rate of application of the compounds of the present invention may vary within wide limits and depends upon the nature of the soil, the method of application, the target insect pest to be controlled, the prevailing climatic conditions, and other factors governed by the method of application and the time of application. The compounds of the present invention are generally applied at a rate of 0.001 to 4 kg/ha, especially from 0.005 to 1 kg/ha, in particular of 0.01 to 0.5 kg/ha. Suitably, the plant growth regulator is applied at a rate from 10 about 50 to about 100 g ai/ha, and acibenzolar-S-methyl is applied at a rate from about 5 to about 25 g ai/ha. When the plant growth regulator is trinexapac-ethyl, a particularly preferred rate is 100 g ai/ha.

The method of the present invention may be applied to any crop plants, in particular 15 monocotyledons such as cereals (wheat, millet, sorghum, rye, triticale, oats, barley, teff, spelt, buckwheat, fonio and quinoa), rice, maize (corn), and/or sugar cane; or dicotyledon crops such as beet (such as sugar beet or fodder beet); fruits (such as pomes, stone fruits or soft fruits, for example apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries or blackberries); leguminous plants (such as beans, lentils, peas or soybeans); oil plants (such 20 as rape, mustard, poppy, olives, sunflowers, coconut, castor oil plants, cocoa beans or groundnuts); cucumber plants (such as marrows, cucumbers or melons); fibre plants (such as cotton, flax, hemp or jute); citrus fruit (such as oranges, lemons, grapefruit or mandarins); vegetables (such as spinach, lettuce, cabbages, carrots, tomatoes, cucurbits or paprika); lauraceae (such as avocados, cinnamon or camphor); tobacco; nuts; coffee; tea; 25 vines; hops; durian; bananas; natural rubber plants; and ornamentals (such as flowers, shrubs, broad-leaved trees or evergreens, for example conifers). This list does not represent any limitation.

Suitably the crop plants are monocotyledonous plants. More suitably, the crop plants are 30 cereals, in particular wheat or barley. In one embodiment, the cereal crop is wheat. In a further embodiment, the cereal crop is barley. In a further embodiment, the crop plants are rice plants. In a further embodiment, the crop plants are sugar cane plants. In further embodiment, the crop plants are corn plants.

Suitably the crop plants are dicotyledonous plants. In one embodiment, the crop plants are oil seed rape plants.

- 5 Crops include those that have been rendered tolerant to herbicides like bromoxynil or classes of herbicides (such as HPPD inhibitors, ALS inhibitors (for example primisulfuron, prosulfuron and trifloxysulfuron), EPSPS (5-enol-pyrovyl-shikimate-3-phosphate-synthase) inhibitors, GS (glutamine synthetase) inhibitors or PPO (protoporphyrinogen-oxidase) inhibitors) as a result of conventional methods of breeding or genetic engineering. An
- 10 example of a crop that has been rendered tolerant to imidazolinones, e.g. imazamox, by conventional methods of breeding (mutagenesis) is Clearfield® summer rape (Canola). Examples of crops that have been rendered tolerant to herbicides or classes of herbicides by genetic engineering methods include glyphosate- and glufosinate-resistant maize varieties commercially available under the trade names RoundupReady® , Herculex IO and
- 15 LibertyLink®. Crops also includes plants that have been transformed by the use of recombinant DNA techniques so that they are capable of synthesising one or more selectively acting toxins, such as are known, for example, from toxin-producing bacteria, especially those of the genus *Bacillus*. Crops also includes plants which have been transformed by the use of recombinant DNA techniques so that they are capable of synthesising antipathogenic
- 20 substances having a selective action, such as, for example, the so-called "pathogenesis-related proteins". Examples of such antipathogenic substances and transgenic plants capable of synthesising such antipathogenic substances are known, for example, from EP-A-0 392 225, WO 95/33818, and EP-A-0 353 191. The methods of producing such transgenic plants are generally known to the person skilled in the art and are described, for example, in the
- 25 publications mentioned above.

The plant growth regulator and acibenzolar-S-methyl of the present invention may be applied either simultaneously or sequentially in any order. If administered sequentially, the components may be administered in any order in a suitable timescale, for example, with no

30 longer than 1 month, no longer than 1 week, or no longer than 24 hours between the time of administering the first component and the time of administering the last component. Suitably, the components are administered within a timescale of a few hours, such as one hour. If the plant growth regulator and acibenzolar-S-methyl components are administered

simultaneously, they may be administered separately or as a tank mix or as a pre-formulated mixture. In one embodiment the mixture or composition of the present invention may be applied to the crop plants as a seed treatment prior to planting.

- 5 When the method of the present invention refers to the application to crop plants of a co-formulated composition, the composition comprises both plant growth regulator and acibenzolar-S-methyl. The compounds may be homogeneously mixed together with other formulation components necessary to make the desired formulation type, as is known to those skilled in the art.

10 In one embodiment of the present invention, the plant growth regulator and acibenzolar-S-methyl, are applied in the form of a composition comprising an agriculturally acceptable carrier.

- 15 The compounds of the present invention may be used in unmodified form, but are generally formulated into compositions using formulation adjuvants, such as carriers, solvents and surface-active substances. The formulations can be in various physical forms, for example dusting powders, gels, wettable powders, water-dispersible granules, water-dispersible tablets, effervescent compressed tablets, emulsifiable concentrates, microemulsifiable concentrates, oil-in-water emulsions, oil flowables, aqueous dispersions, oil dispersions, 20 suspoemulsions, capsule suspensions, emulsifiable granules, soluble liquids, water-soluble concentrates (with water or a water-miscible organic solvent as carrier), or impregnated polymer films. Such formulations can either be used directly or are diluted prior to use. Diluted formulations can be prepared, for example, with water, liquid fertilizers, micro- 25 nutrients, biological organisms, oil or solvents. These formulations may contain as little as about 0.5% to as much as about 95% or more by weight of active ingredient. The optimum amount for any given compound will depend on formulation, application equipment and nature of the plants to be controlled.

- 30 Wettable powders are in the form of finely divided particles which disperse readily in water or other liquid carriers. The particles contain the active ingredient retained in a solid matrix. Typical solid matrices include fuller's earth, kaolin clays, silicas and other readily wet

organic or inorganic solids. Wettable powders normally contain about 5% to about 95% of the active ingredient plus a small amount of wetting, dispersing or emulsifying agent.

Emulsifiable concentrates are homogeneous liquid compositions dispersible in water or other liquid and may consist entirely of the active compound with a liquid or solid emulsifying agent, or may also contain a liquid carrier, such as xylene, heavy aromatic naphthas, isophorone and other non-volatile organic solvents. In use, these concentrates are dispersed in water or other liquid and normally applied as a spray to the area to be treated. The amount of active ingredient may range from about 0.5% to about 95% of the concentrate.

Granular formulations include both extrudates and relatively coarse particles and are usually applied without dilution to the area in which suppression of vegetation is desired. Typical carriers for granular formulations include fertiliser, sand, fuller's earth, attapulgite clay, bentonite clays, montmorillonite clay, vermiculite, perlite, calcium carbonate, brick, pumice, pyrophyllite, kaolin, dolomite, plaster, wood flour, ground corn cobs, ground peanut hulls, sugars, sodium chloride, sodium sulphate, sodium silicate, sodium borate, magnesia, mica, iron oxide, zinc oxide, titanium oxide, antimony oxide, cryolite, gypsum, diatomaceous earth, calcium sulphate and other organic or inorganic materials which absorb or which can be coated with the active compound. Particularly suitable is a fertiliser granule carrier.

Granular formulations normally contain about 5% to about 25% active ingredients which may include surface-active agents such as heavy aromatic naphthas, kerosene and other petroleum fractions, or vegetable oils; and/or stickers such as dextrans, glue or synthetic resins. The granular substrate material can be one of the typical carriers mentioned above and/or can be a fertiliser material e.g. urea/formaldehyde fertilisers, ammonium, liquid nitrogen, urea, potassium chloride, ammonium compounds, phosphorus compounds, sulphur, similar plant nutrients and micronutrients and mixtures or combinations thereof. The plant growth regulator and acibenzolar-S-methyl may be homogeneously distributed throughout the granule or may be spray impregnated or absorbed onto the granule substrate after the granules are formed.

Encapsulated granules are generally porous granules with porous membranes sealing the granule pore openings, retaining the active species in liquid form inside the granule pores. Granules typically range from 1 millimetre to 1 centimetre, preferably 1 to 2 millimetres in

diameter. Granules are formed by extrusion, agglomeration or prilling, or are naturally occurring. Examples of such materials are vermiculite, sintered clay, kaolin, attapulgite clay, sawdust and granular carbon. Shell or membrane materials include natural and synthetic rubbers, cellulosic materials, styrene-butadiene copolymers, polyacrylonitriles, polyacrylates, polyesters, polyamides, polyureas, polyurethanes and starch xanthates.

Dusts are free-flowing admixtures of the active ingredient with finely divided solids such as talc, clays, flours and other organic and inorganic solids which act as dispersants and carriers.

Microcapsules are typically droplets or granules of the active material enclosed in an inert porous shell which allows escape of the enclosed material to the surroundings at controlled rates. Encapsulated droplets are typically about 1 to 50 microns in diameter. The enclosed liquid typically constitutes about 50 to 95% of the weight of the capsule and may include solvent in addition to the active compound.

Other useful formulations for plant growth regulation applications include simple solutions of the active ingredients in a solvent in which it is completely soluble at the desired concentration, such as acetone, alkylated naphthalenes, xylene and other organic solvents. Pressurised sprayers, wherein the active ingredient is dispersed in finely-divided form as a result of vaporisation of a low boiling dispersant solvent carrier, may also be used.

Many of the formulations described above include wetting, dispersing or emulsifying agents. Examples are alkyl and alkylaryl sulphonates and sulphates and their salts, polyhydric alcohols; polyethoxylated alcohols, esters and fatty amines. These agents, when used, normally comprise from 0.1% to 15% by weight of the formulation.

Suitable agricultural adjuvants and carriers, either formulated together and/or added separately, that are useful in formulating the compositions of the invention in the formulation types described above are well known to those skilled in the art. Suitable examples of the different classes are found in the non-limiting list below.

Liquid carriers that can be employed include water, toluene, xylene, petroleum naphtha, crop oils, AMS; acetone, methyl ethyl ketone, cyclohexanone, acetic anhydride, acetonitrile,

acetophenone, amyl acetate, 2-butanone, chlorobenzene, cyclohexane, cyclohexanol, alkyl acetates, diacetonol, 1,2-dichloropropane, diethanolamine, p-diethylbenzene, diethylene glycol, diethylene glycol abietate, diethylene glycol butyl ether, diethylene glycol ethyl ether, diethylene glycol methyl ether, N,N-dimethyl formamide, dimethyl sulfoxide, 1,4-dioxane, 5 dipropylene glycol, dipropylene glycol methyl ether, dipropylene glycol dibenzoate, diproxitol, alkyl pyrrolidinone, ethyl acetate, 2-ethyl hexanol, ethylene carbonate, 1,1,1-trichloroethane, 2-heptanone, alpha pinene, d-limonene, ethylene glycol, ethylene glycol butyl ether, ethylene glycol methyl ether, gamma-butyrolactone, glycerol, glycerol diacetate, glycerol monoacetate, glycerol triacetate, hexadecane, hexylene glycol, isoamyl acetate, 10 isobornyl acetate, isooctane, isophorone, isopropyl benzene, isopropyl myristate, lactic acid, laurylamine, mesityl oxide, methoxy-propanol, methyl isoamyl ketone, methyl isobutyl ketone, methyl laurate, methyl octanoate, methyl oleate, methylene chloride, m-xylene, n-hexane, n-octylamine, octadecanoic acid, octyl amine acetate, oleic acid, oleylamine, o-xylene, phenol, polyethylene glycol (PEG400), propionic acid, propylene glycol, propylene glycol monomethyl ether, p-xylene, toluene, triethyl phosphate, triethylene glycol, xylene 15 sulfonic acid, paraffin, mineral oil, trichloroethylene, perchloroethylene, ethyl acetate, amyl acetate, butyl acetate, methanol, ethanol, isopropanol, and higher molecular weight alcohols such as amyl alcohol, tetrahydrofurfuryl alcohol, hexanol, octanol, etc. ethylene glycol, propylene glycol, glycerine, N-methyl-2-pyrrolidinone, and the like. Water is generally the 20 carrier of choice for the dilution of concentrates.

Suitable solid carriers include talc, titanium dioxide, pyrophyllite clay, silica, attapulgite clay, kieselguhr, chalk, diatomaceous earth, lime, calcium carbonate, bentonite clay, fuller's earth, fertiliser, cotton seed hulls, wheat flour, soybean flour, pumice, wood flour, walnut shell 25 flour, lignin and the like.

A broad range of surface-active agents are advantageously employed in both said liquid and solid compositions, especially those designed to be diluted with carrier before application. The surface-active agents can be anionic, cationic, non-ionic or polymeric in character and 30 can be employed as emulsifying agents, wetting agents, suspending agents or for other purposes. Typical surface active agents include salts of alkyl sulfates, such as diethanolammonium lauryl sulphate; alkylarylsulfonate salts, such as calcium dodecylbenzenesulfonate; alkylphenol-alkylene oxide addition products, such as

nonylphenol-C.sub. 18 ethoxylate; alcohol-alkylene oxide addition products, such as tridecyl alcohol-C.sub. 16 ethoxylate; soaps, such as sodium stearate; alkylnaphthalenesulfonate salts, such as sodium dibutylnaphthalenesulfonate; dialkyl esters of sulfosuccinate salts, such as sodium di(2-ethylhexyl) sulfosuccinate; sorbitol esters, such as sorbitol oleate; quaternary
5 amines, such as lauryl trimethylammonium chloride; polyethylene glycol esters of fatty acids, such as polyethylene glycol stearate; block copolymers of ethylene oxide and propylene oxide; and salts of mono and dialkyl phosphate esters.

Other adjuvants commonly utilized in agricultural compositions include crystallisation
10 inhibitors, viscosity modifiers, suspending agents, spray droplet modifiers, pigments, antioxidants, foaming agents, light-blocking agents, compatibilizing agents, antifoam agents, sequestering agents, neutralising agents and buffers, corrosion inhibitors, dyes, odorants, spreading agents, penetration aids, micronutrients, emollients, lubricants, sticking agents, and the like. The compositions can also be formulated with liquid fertilizers or solid, particulate
15 fertiliser carriers such as ammonium nitrate, urea and the like.

Also, the present invention may optionally include one or more additional pesticides such as insecticides, nematicides, fungicides or herbicides or additional plant growth regulators. Co-application of pesticides with the present invention has the added benefit of minimising
20 farmer time spent applying products to crops, since only a single application may be required to both provide growth regulation and control pests.

According to the present invention, there is provided the use of a composition comprising a synergistically effective amount of a plant growth regulator and acibenzolar-S-methyl, for
25 regulating the growth of and/or enhancing crop plants, as described above.

According to the present invention, there is provided a plant growth regulating composition, comprising a plant growth regulator and acibenzolar-S-methyl at a weight ratio from about 10:1 to about 1:1. In one embodiment, the weight ratio is about 5:1. In a further
30 embodiment, the plant growth regulator and acibenzolar-S-methyl are present in a synergistically effective amount. In an alternative embodiment, the plant growth regulator is trinexapac-ethyl.

Compositions of the present invention may contain from about 0.001% to about 99% by weight active ingredients. Suitably, the composition contains from about 0.001% to about 50% by weight active ingredients. More suitably, the composition contains from about 0.001% to about 10% by weight active ingredients. More suitably, the composition contains
5 from about 0.001% to about 1% by weight active ingredients. If the formulation is in the form of a concentrate, requiring dilution with water before use, it will contain a higher amount of active ingredients than a composition that is ready to use without dilution.

The following examples further exemplify the present invention. Although the invention has
10 been described with reference to preferred embodiments and examples thereof, the scope of the present invention is not limited only to those described embodiments. As will be apparent to persons skilled in the art, modifications and adaptations to the above-described invention can be made without departing from the spirit and scope of the invention, which is defined and circumscribed by the attached claims.

EXAMPLES

Example 1

A glasshouse trial was setup to compare growth regulation effects of trinexapac-ethyl, acibenzolar-S-methyl, and mixtures of trinexapac-ethyl and acibenzolar-S-methyl at various rates, on both winter and summer varieties of barley and wheat (summer barley Pasadena, winter barley Hasso, winter wheat Arina, and summer wheat Lona).

Table 1 describes the treatments made. Each treatment was applied as a spray to the leaves of the plant at plant growth stage 30 (start of stem elongation). Assessments of growth were made two weeks after application, and the results are expressed as percentage growth reduction in table 2.

TABLE 1: Treatment list

Treatment Number	Treatment description	Treatment details	Rate (g AI / ha)
1	Trinexapac-ethyl (TXP) only	Moddus®	100
2	TXP	Moddus®	50
3	Acibenzolar-S-methyl (ASM) only	Bion®	20
4	ASM	Bion®	10
5	ASM	Bion®	5
6	TXP + ASM	Moddus®+Bion®	100+20
7	TXP + ASM	Moddus®+Bion®	50+20

TABLE 2: Results

	Summer-Barley Pasadena		Winter-Barley Hasso		Winter-Wheat Arina		Summer-Wheat Lona	
Treatment	% growth reduction (actual)	Colby (expected)	% growth reduction (actual)	Colby (expected)	% growth reduction (actual)	Colby (expected)	% growth reduction (actual)	Colby (expected)
1	0	n/a	8	n/a	10	n/a	5	n/a
2	0	n/a	3	n/a	5	n/a	3	n/a
3	3	n/a	8	n/a	3	n/a	3	n/a
4	0	n/a	0	n/a	0	n/a	0	n/a
5	0	n/a	0	n/a	0	n/a	0	n/a
6	18	3	33	14	20	12	13	7
7	8	3	23	10	8	7	13	5

Shading / bold type indicates synergy (i.e. where the actual percentage growth reduction observed is greater than the 'Colby' expected growth reduction).

5

A synergistic growth reduction effect was observed when applying mixtures of trinexapac-ethyl and acibenzolar-S-methyl, at all rates tested, and against all 4 species.

CLAIMS

1. A method for regulating the growth of and/or enhancing crop plants, comprising applying to the plants a plant growth regulator and acibenzolar-S-methyl.
- 5 2. A method according to claim 1, wherein the plant growth regulator and acibenzolar-S_methyl are applied in a synergistically effective amount.
3. A method according to claim 1 or 2, wherein the plant growth regulator is selected from
10 the group consisting of trinexapac-ethyl, prohexadione-calcium, paclobutrazol, uniconazole, flurprimidol, mefluidide, mepiquat-chloride, chlormequat-chloride, and a mixture thereof.
4. A method according to claim 3, wherein the plant growth regulator is trinexapac-ethyl.
- 15 5. A method according to claim 3, wherein the plant growth regulator is paclobutrazol.
6. A method according to claim 1, wherein the crop plants are monocotyledonous plants.
7. A method according to claim 6, wherein the crop plants are selected from the group
20 consisting of cereals, rice, maize and sugar cane.
8. A method according to claim 1, wherein the plant growth regulator and acibenzolar-S-methyl are applied to the plant in a weight ratio from about 5:1 to about 1:1.
- 25 9. A method according to claim 1, wherein the plant growth regulator is applied at a rate from about 50 to about 100 g ai/ ha.
10. A method according to claim 1, wherein the acibenzolar-S-methyl is applied at a rate
30 from about 5 to about 25 g ai/ ha.
11. A method according to claim 1, wherein the enhancement is improved plant yield.

12. A method according to claim 1, wherein the enhancement is improved plant vigour and/or plant quality, and/or plant tolerance to stress factors.
13. A method according to claim 12, wherein the enhancement is improved tolerance to drought conditions.
14. A plant growth regulating composition, comprising trinexapac-ethyl and acibenzolar-S-methyl in a weight ratio from about 10:1 to about 1:1.
15. A plant growth regulating composition according to claim 14, comprising a plant growth regulator and acibenzolar-S-methyl in a weight ratio of about 5:1.
16. Use of a composition comprising a synergistically effective amount of a plant growth regulator and acibenzolar-S-methyl, for regulating the growth of and/or enhancing crop plants.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2010/007129

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

4, 14(completely); 1-3, 6-13, 15, 16(partially)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2010/007129

A. CLASSIFICATION OF SUBJECT MATTER
INV. A01N37/42 A01N43/653 A01N43/828 A01P21/00
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MAHESANIYA A ET AL: "Efficacy of acibenzolar-S-methyl and paclobutrazol for the control of bacterial speck (Pseudomonas syringae pv tomato) in tomato plug transplants", CANADIAN JOURNAL OF PLANT PATHOLOGY, GUELPH, ONT, CA, vol. 27, 1 January 2002 (2002-01-01), pages 377-378, XP009144261, ISSN: 0706-0661 abstract ----- -/--	1-3,5, 12,16



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

28 February 2011

Date of mailing of the international search report

26/04/2011

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Habermann, Jörg

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2010/007129

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MAHESANIYA A; PITBLADO R E; SOUZA MACHADO V: "Ontario Tomato Research Institute - Research Summary Results - 2002: Paclobutrazol and Acibenzolar-S-Methyl induced tomato seeding growth response and resistance to bacterial speck (<i>Pseudomonas syringae</i> pv. <i>tomato</i>)", University of Guelph, Ridgetown Campus, Ontario, Canada 1 January 2002 (2002-01-01), XP002621840, Retrieved from the Internet: URL: http://www.opvg.org/2002%20TRI4.pdf [retrieved on 2011-02-07] the whole document	1-3,5, 12,16
X	----- WO 2008/020872 A2 (SYNGENTA PARTICIPATIONS AG [CH]; HASKELL JACK [US]; SHETTY KIRAN [US];) 21 February 2008 (2008-02-21) cited in the application	1-4,6-16
A	paragraphs [0001], [0006], [0019], [0021], [0028], [0033], [0034], [0066] paragraphs [0069] - [0071], [0075] - [0083], [0091], [0128], [0131] paragraphs [0132], [0144] - [0146], [0148], [0151], [0152], [0155] examples 1-12 -----	5

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2010/007129

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2008020872 A2	21-02-2008	AR 058622 A1	13-02-2008
		AU 2006347294 A1	21-02-2008
		CA 2633987 A1	21-02-2008
		EP 1962588 A2	03-09-2008
		JP 2009524593 T	02-07-2009
		UY 30053 A1	31-07-2007

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 4, 14(completely); 1-3, 6-13, 15, 16(partially)

A method for regulating the growth of and/or enhancing crop plants, comprising applying to the plants trinexapac-ethyl and acibenzolar-S-methyl.

2. claims: 1-3, 6-13, 15, 16(all partially)

A method for regulating the growth of and/or enhancing crop plants, comprising applying to the plants prohexadione-calcium and acibenzolar-S-methyl.

3. claims: 5(completely); 1-3, 6-13, 15, 16(partially)

A method for regulating the growth of and/or enhancing crop plants, comprising applying to the plants paclobutrazol and acibenzolar-S-methyl.

4. claims: 1-3, 6-13, 15, 16(all partially)

A method for regulating the growth of and/or enhancing crop plants, comprising applying to the plants uniconazole and acibenzolar-S-methyl.

5. claims: 1-3, 6-13, 15, 16(all partially)

A method for regulating the growth of and/or enhancing crop plants, comprising applying to the plants flurprimidol and acibenzolar-S-methyl.

6. claims: 1-3, 6-13, 15, 16(all partially)

A method for regulating the growth of and/or enhancing crop plants, comprising applying to the plants mefluidide and acibenzolar-S-methyl.

7. claims: 1-3, 6-13, 15, 16(all partially)

A method for regulating the growth of and/or enhancing crop plants, comprising applying to the plants mepiquat-chloride and acibenzolar-S-methyl.

8. claims: 1-3, 6-13, 15, 16(all partially)

A method for regulating the growth of and/or enhancing crop

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

plants, comprising applying to the plants
chlormequat-chloride and acibenzolar-S-methyl.
